

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.2-Cosine/92-4.2.3.1-a+b-cos^m-c+d-cosⁿ-
A+B-cos-

Nasser M. Abbasi

December 8, 2023

Compiled on December 8, 2023 at 11:21pm

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	212
4	Appendix	5894

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [644]. This is test number [92].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (644)	0.00 (0)
Mathematica	98.60 (635)	1.40 (9)
Maple	97.67 (629)	2.33 (15)
Fricas	72.98 (470)	27.02 (174)
Mupad	35.87 (231)	64.13 (413)
Giac	32.61 (210)	67.39 (434)
Maxima	32.45 (209)	67.55 (435)
Sympy	10.71 (69)	89.29 (575)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	98.447	0.000	0.000	1.553
Mathematica	62.267	9.627	25.311	2.795
Maple	46.118	49.224	0.776	3.882
Fricas	41.460	6.522	23.447	28.571
Giac	21.739	8.696	0.621	68.944
Maxima	15.683	14.441	0.776	69.099
Sympy	2.019	7.298	0.000	90.683
Mupad	0.000	34.317	0.000	65.683

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	9	88.89	11.11	0.00
Maple	15	66.67	33.33	0.00
Fricas	174	56.90	43.10	0.00
Mupad	413	0.00	100.00	0.00
Giac	434	83.18	14.98	1.84
Maxima	435	80.46	7.36	12.18
Sympy	575	35.30	64.70	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	1.19
Rubi	1.27
Maxima	2.36
Mupad	2.70
Giac	4.13
Mathematica	4.40
Sympy	15.96
Maple	23.19

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	226.23	1.02	194.00	1.01
Fricas	290.10	1.62	197.50	1.24
Mathematica	461.19	1.86	207.00	1.07
Sympy	643.57	5.18	252.00	2.50
Maple	878.21	2.99	355.00	2.13
Mupad	883.30	4.10	202.00	1.43
Giac	2408.05	13.67	188.50	1.45
Maxima	14869.09	85.36	269.00	1.75

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{449, 455, 456, 457, 458, 635, 641, 642, 643, 644}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {123, 124, 125, 126, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 163, 164, 165, 195, 196, 200, 201, 206, 207, 208, 209, 213, 214, 339, 340, 382, 395, 402, 410, 411, 412, 418, 426, 431, 432, 435, 436, 454, 521, 522, 523, 524, 529, 530, 531, 532, 537, 538, 543, 544, 572, 573, 577, 578, 579, 580, 582, 583, 590, 591, 597, 598, 599, 600, 605, 606, 607, 608, 609, 612, 614, 620, 621, 624, 625, 626, 627, 628, 630, 634, 640}

Maple {106, 122, 164, 165, 209, 339, 340, 382, 395, 398, 402, 403, 404, 405, 406, 410, 411, 412, 413, 414, 415, 418, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 487, 541, 548, 583, 593, 597, 600, 601, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```


1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v0.6

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	192

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	22
2.1.3	Maple	23
2.1.4	Fricas	24
2.1.5	Maxima	25
2.1.6	Giac	26
2.1.7	Mupad	27
2.1.8	Sympy	28

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 450, 451, 452, 453, 454, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549,

550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 636, 637, 638, 639, 640 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 35, 36, 37, 41, 49, 51, 60, 61, 62, 68, 69, 70, 71, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 197, 198, 199, 202, 203, 204, 205, 208, 209, 211, 212, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 237, 238, 239, 240, 241, 242, 243, 244, 245, 247, 248, 249, 250, 251, 252, 253, 254, 255, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 275, 276, 277, 278, 279, 280, 281, 282, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 304, 305, 306, 311, 312, 313, 319, 320, 321, 322, 323, 326, 327, 328, 329, 333, 334, 335, 336, 337, 341, 342, 343, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 394, 397, 398, 399, 419, 421, 422, 423, 424, 437, 438, 439, 440, 450, 451, 452, 453, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 525, 528, 533, 539, 540, 545, 546, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 597, 598, 599, 601, 606, 607, 613, 614, 615, 616, 617, 618, 620, 621, 622, 626, 627, 628, 631, 632, 633, 634, 636, 637, 638, 639 }

B grade { 23, 32, 33, 34, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 50, 52, 53, 54, 55, 56, 57, 58, 59, 63, 64, 65, 66, 67, 72, 73, 210, 236, 246, 256, 257, 273, 274, 283, 369, 393, 454, 547, 573, 595, 596, 600, 602, 603, 604, 605, 608, 609, 610, 611, 612, 619, 623, 624, 625, 629, 630, 640 }

C grade { 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160,

161, 162, 163, 164, 165, 194, 195, 196, 200, 201, 206, 207, 213, 214, 301, 302, 303, 307, 308, 309, 310, 314, 315, 316, 317, 318, 324, 325, 330, 331, 332, 338, 339, 340, 344, 395, 396, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 420, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 520, 521, 522, 523, 524, 526, 527, 529, 530, 531, 532, 534, 535, 536, 537, 538, 541, 542, 543, 544, 548, 549 }

F normal fail { 441, 442, 443, 444, 445, 446, 447, 448 }

F(-1) timedout fail { 641 }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 82, 83, 84, 85, 91, 92, 93, 100, 101, 102, 107, 130, 131, 133, 137, 138, 139, 140, 145, 146, 147, 148, 149, 151, 158, 159, 166, 167, 168, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 196, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 300, 322, 323, 329, 330, 342, 343, 344, 367, 368, 370, 390, 391, 422, 438, 439, 461, 464, 467, 468, 469, 473, 474, 475, 477, 478, 479, 480, 481, 483, 484, 485, 486, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 525, 526, 527, 528, 529, 530, 534, 535, 552, 555, 567, 568, 569, 570, 617, 632, 633, 634 }

B grade { 78, 79, 80, 81, 86, 87, 88, 89, 90, 94, 95, 96, 97, 98, 99, 103, 104, 105, 106, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 132, 134, 135, 136, 141, 142, 143, 144, 150, 152, 153, 154, 155, 156, 157, 160, 161, 162, 163, 164, 165, 169, 194, 195, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 296, 297, 298, 299, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 324, 325, 326, 327, 328, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 358, 359, 360, 361, 362, 365, 366, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 387, 388, 389, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 440, 441, 442, 443, 444, 445, 446, 447, 448, 459, 460, 462, 463, 465, 466, 470, 471, 472, 476, 482, 487, 506, 523, 524, 531, 532, 533, 536, 537, 538, 539, 540, 541, 542, 543, 544,

545, 546, 547, 548, 549, 550, 551, 553, 554, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 610, 611, 612, 613, 614, 615, 616, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631 }

C grade { 341, 357, 363, 364, 386 }

F normal fail { 450, 451, 452, 453, 454, 636, 637, 638, 639, 640 }

F(-1) timedout fail { 605, 606, 607, 608, 609 }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 73, 74, 75, 76, 77, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 107, 108, 109, 110, 114, 115, 116, 117, 122, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 257, 258, 261, 281, 282, 283, 284, 286, 287, 288, 289, 290, 291, 292, 293, 295, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549 }

B grade { 6, 53, 72, 78, 79, 103, 104, 105, 106, 111, 112, 113, 118, 119, 120, 121, 219, 255, 256, 259, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 285, 294 }

C grade { 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 296, 297, 298, 299, 304, 305, 306, 311, 312, 313, 319, 320, 321, 322, 326, 327, 328, 329, 333, 334, 335, 336, 337, 341, 343, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 383, 384, 385, 386, 387, 388, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 584, 585, 586, 587, 588, 589 }

F normal fail { 300, 307, 308, 314, 365, 396, 398, 399, 400, 401, 403, 404, 405, 406, 407, 408, 409, 411, 412, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 450, 451, 452, 453, 454, 590 }

591, 592, 593, 594, 595, 597, 598, 599, 600, 601, 602, 603, 605, 606, 607, 608, 611, 612, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 631, 632, 633, 634, 636, 637, 638, 639, 640 }

F(-1) timeout fail { 301, 302, 303, 309, 310, 315, 316, 317, 318, 323, 324, 325, 330, 331, 332, 338, 339, 340, 342, 344, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 389, 390, 391, 392, 393, 394, 395, 397, 402, 410, 413, 414, 431, 432, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 596, 604, 609, 610, 613, 629, 630 }

F(-2) exception fail { }

2.1.5 Maxima

A grade { 1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 50, 51, 52, 57, 58, 59, 60, 61, 62, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 82, 83, 84, 85, 86, 91, 92, 93, 94, 104, 215, 216, 217, 218, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249 }

B grade { 6, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 53, 54, 55, 56, 63, 64, 79, 80, 81, 87, 88, 89, 90, 95, 96, 97, 100, 101, 102, 103, 105, 106, 111, 112, 113, 120, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 219, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 517, 518, 519 }

C grade { 193, 194, 195, 196, 525 }

F normal fail { 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 191, 197, 198, 199, 200, 201, 202, 203, 204, 205, 207, 208, 209, 210, 211, 212, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 333, 334, 335, 336, 337, 338, 339, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 376, 377, 378, 379, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 450, 451, 452, 453, 454, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 488, 489, 490, 491, 492, 520, 521, 522, 523, 524, 527, 528, 529, 530, 531, 532, 533, 534, 535, 539, 540, 541, 542, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 574, 575, 576, 577, 580, 581, 582, 583, 584, 585, 586,

587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 636, 637, 638, 639, 640 }

F(-1) timeout fail { 98, 99, 107, 108, 109, 110, 115, 116, 117, 118, 119, 163, 165, 206, 213, 214, 332, 340, 375, 380, 382, 487, 516, 536, 537, 538, 543, 544, 572, 573, 578, 579 }

F(-2) exception fail { 114, 121, 122, 164, 192, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 381, 526 }

2.1.6 Giac

A grade { 1, 2, 3, 4, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 114, 115, 116, 117, 118, 119, 120, 121, 215, 216, 217, 223, 224, 225, 231, 232, 233, 235, 240, 241, 242, 244, 251, 252, 254, 255, 258, 260, 261, 262, 264, 281, 282, 283, 286, 287, 288, 289, 290, 292, 293, 294 }

B grade { 5, 6, 7, 15, 79, 104, 105, 106, 111, 112, 218, 219, 220, 221, 222, 226, 227, 228, 229, 230, 234, 236, 237, 238, 239, 243, 245, 246, 247, 248, 249, 250, 253, 256, 257, 259, 263, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 285, 291, 295 }

C grade { 173, 180, 188, 284 }

F normal fail { 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 170, 174, 175, 182, 183, 194, 195, 196, 200, 201, 206, 207, 213, 214, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 450, 451, 452, 453, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 497, 499, 500, 508, 509, 518, 519, 520, 521, 522, 523, 524, 529, 530, 531, 532, 536, 537, 538, 543, 544, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614,

615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 636, 637, 638, 639 }

F(-1) timeout fail { 169, 171, 172, 176, 177, 178, 179, 181, 184, 185, 186, 187, 189, 190, 191, 192, 193, 197, 198, 199, 202, 203, 204, 205, 208, 209, 210, 211, 212, 493, 494, 495, 496, 498, 501, 502, 503, 504, 505, 506, 507, 510, 511, 512, 513, 514, 515, 516, 517, 525, 526, 527, 528, 533, 534, 535, 539, 540, 541, 542, 545, 546, 547, 548, 549 }

F(-2) exception fail { 107, 108, 109, 110, 113, 122, 454, 640 }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 102, 103, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 171, 172, 173, 179, 180, 181, 188, 189, 190, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 321, 322, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 493, 494, 495, 496, 501, 502, 503, 504, 510, 511, 512, 513 }

C grade { }

F normal fail { }

F(-1) timeout fail { 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 174, 175, 176, 177, 178, 182, 183, 184, 185, 186, 187, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 450, 451, 452, 453, 454, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 497, 498, 499,

500, 505, 506, 507, 508, 509, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 636, 637, 638, 639, 640 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { 42, 50, 51, 59, 60, 61, 67, 68, 69, 70, 225, 284, 288 }

B grade { 1, 2, 3, 4, 10, 11, 12, 13, 19, 20, 21, 28, 29, 30, 38, 39, 40, 41, 47, 48, 49, 56, 57, 58, 65, 66, 215, 216, 217, 223, 224, 231, 232, 233, 240, 241, 242, 251, 252, 253, 261, 281, 282, 283, 285, 286, 292 }

C grade { }

F normal fail { 5, 6, 7, 8, 9, 14, 15, 16, 17, 22, 23, 24, 31, 32, 43, 44, 45, 46, 52, 53, 54, 55, 62, 63, 64, 71, 72, 73, 75, 76, 77, 78, 79, 80, 85, 101, 102, 103, 104, 105, 106, 110, 111, 112, 113, 114, 119, 120, 121, 126, 148, 168, 169, 170, 171, 176, 177, 178, 192, 193, 194, 195, 198, 199, 200, 204, 205, 218, 219, 220, 221, 222, 226, 227, 228, 229, 234, 235, 236, 243, 244, 254, 255, 256, 257, 262, 263, 264, 270, 271, 272, 278, 279, 280, 287, 293, 294, 295, 297, 298, 299, 300, 301, 302, 306, 320, 321, 322, 323, 324, 325, 330, 331, 332, 338, 339, 340, 341, 342, 343, 344, 348, 396, 397, 398, 399, 404, 405, 406, 421, 422, 423, 424, 427, 428, 429, 433, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 453, 462, 463, 464, 468, 469, 474, 475, 478, 479, 480, 483, 484, 488, 498, 499, 500, 525, 526, 527, 528, 533, 534, 553, 554, 555, 559, 560, 565, 566, 569, 570, 571, 574, 575, 579, 585, 586, 587, 588, 594, 595, 596, 617, 618, 619, 622, 623, 632, 633, 637, 638, 639, 640 }

F(-1) timedout fail { 18, 25, 26, 27, 33, 34, 35, 36, 37, 74, 81, 82, 83, 84, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 107, 108, 109, 115, 116, 117, 118, 122, 123, 124, 125, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 172, 173, 174, 175, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 196, 197, 201, 202, 203, 206, 207, 208, 209, 210, 211, 212, 213, 214, 230, 237, 238, 239, 245, 246, 247, 248, 249, 250, 258, 259, 260, 265, 266, 267, 268, 269, 273, 274, 275, 276, 277, 289, 290, 291, 296, 303, 304, 305, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 326, 327, 328, 329, 333, 334, 335, 336, 337, 345, 346, 347, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 400, 401, 402, 403, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 425, 426, 430, 431, 432, 434, 435, 436, 450, 451, 452,

454, 459, 460, 461, 465, 466, 467, 470, 471, 472, 473, 476, 477, 481, 482, 485, 486, 487, 489, 490, 491, 492, 493, 494, 495, 496, 497, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 529, 530, 531, 532, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 556, 557, 558, 561, 562, 563, 564, 567, 568, 572, 573, 576, 577, 578, 580, 581, 582, 583, 584, 589, 590, 591, 592, 593, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 620, 621, 624, 625, 626, 627, 628, 629, 630, 631, 634, 636, 641 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	120	77	87	124	88	333	112	236
N.S.	1	0.96	0.62	0.70	0.99	0.70	2.66	0.90	1.89
time (sec)	N/A	0.692	0.248	3.455	0.207	0.327	0.270	0.323	0.975

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	93	75	67	101	74	252	89	212
N.S.	1	0.96	0.77	0.69	1.04	0.76	2.60	0.92	2.19
time (sec)	N/A	0.565	0.188	2.516	0.202	0.349	0.184	0.319	0.768

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	79	65	54	79	56	168	68	84
N.S.	1	1.03	0.84	0.70	1.03	0.73	2.18	0.88	1.09
time (sec)	N/A	0.391	0.129	1.806	0.198	0.312	0.129	0.303	0.143

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	44	37	55	38	94	45	50
N.S.	1	1.00	0.94	0.79	1.17	0.81	2.00	0.96	1.06
time (sec)	N/A	0.213	0.077	0.946	0.196	0.368	0.094	0.292	0.120

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	46	48	47	51	0	79	100
N.S.	1	1.00	1.44	1.50	1.47	1.59	0.00	2.47	3.12
time (sec)	N/A	0.444	0.018	1.482	0.211	0.335	0.000	0.304	0.177

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	43	50	73	79	0	84	100
N.S.	1	1.00	1.34	1.56	2.28	2.47	0.00	2.62	3.12
time (sec)	N/A	0.444	0.018	2.326	0.207	0.395	0.000	0.306	0.179

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	59	75	75	95	89	0	124	94
N.S.	1	1.05	1.34	1.34	1.70	1.59	0.00	2.21	1.68
time (sec)	N/A	0.556	0.019	2.917	0.235	0.321	0.000	0.315	0.505

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	87	56	81	127	105	0	154	126
N.S.	1	1.01	0.65	0.94	1.48	1.22	0.00	1.79	1.47
time (sec)	N/A	0.669	0.222	2.976	0.237	0.282	0.000	0.307	1.280

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	102	77	119	163	127	0	188	166
N.S.	1	0.96	0.73	1.12	1.54	1.20	0.00	1.77	1.57
time (sec)	N/A	0.698	0.267	3.055	0.242	0.302	0.000	0.312	1.668

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	191	177	134	111	216	130	600	166	315
N.S.	1	0.93	0.70	0.58	1.13	0.68	3.14	0.87	1.65
time (sec)	N/A	1.027	0.399	4.450	0.455	0.320	0.394	0.323	0.991

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	151	108	93	178	110	459	137	277
N.S.	1	0.94	0.68	0.58	1.11	0.69	2.87	0.86	1.73
time (sec)	N/A	0.900	0.237	3.520	0.265	0.315	0.296	0.316	0.967

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	129	128	86	74	144	90	338	110	134
N.S.	1	0.99	0.67	0.57	1.12	0.70	2.62	0.85	1.04
time (sec)	N/A	0.579	0.196	2.840	0.248	0.292	0.191	0.319	0.191

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	83	84	61	110	70	199	85	98
N.S.	1	0.88	0.89	0.65	1.17	0.74	2.12	0.90	1.04
time (sec)	N/A	0.308	0.322	1.932	0.251	0.279	0.135	0.312	0.138

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	82	96	74	94	79	0	145	141
N.S.	1	1.00	1.17	0.90	1.15	0.96	0.00	1.77	1.72
time (sec)	N/A	0.701	1.025	1.526	0.229	0.292	0.000	0.302	0.215

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	143	81	105	108	0	155	161
N.S.	1	1.00	1.93	1.09	1.42	1.46	0.00	2.09	2.18
time (sec)	N/A	0.707	1.341	2.470	0.205	0.292	0.000	0.307	0.192

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	53	106	142	119	0	154	162
N.S.	1	1.00	0.60	1.20	1.61	1.35	0.00	1.75	1.84
time (sec)	N/A	0.736	0.538	3.292	0.310	0.295	0.000	0.310	0.187

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	119	63	120	174	125	0	178	145
N.S.	1	1.05	0.56	1.06	1.54	1.11	0.00	1.58	1.28
time (sec)	N/A	0.879	0.591	4.119	0.210	0.296	0.000	0.319	1.301

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	145	81	150	230	145	0	212	183
N.S.	1	1.01	0.56	1.04	1.60	1.01	0.00	1.47	1.27
time (sec)	N/A	1.027	0.816	4.477	0.243	0.311	0.000	0.329	1.943

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	197	134	112	262	130	695	166	315
N.S.	1	0.98	0.67	0.56	1.30	0.65	3.46	0.83	1.57
time (sec)	N/A	1.211	0.294	4.290	0.229	0.310	0.408	0.318	1.632

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	146	108	93	213	110	530	136	277
N.S.	1	0.95	0.70	0.60	1.38	0.71	3.44	0.88	1.80
time (sec)	N/A	0.628	0.234	3.355	0.212	0.301	0.299	0.310	1.510

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	101	116	79	167	90	371	112	134
N.S.	1	0.87	1.00	0.68	1.44	0.78	3.20	0.97	1.16
time (sec)	N/A	0.352	0.390	2.516	0.220	0.285	0.222	0.319	0.303

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	117	113	93	141	102	0	180	178
N.S.	1	1.05	1.02	0.84	1.27	0.92	0.00	1.62	1.60
time (sec)	N/A	0.945	1.175	2.550	0.219	0.323	0.000	0.345	0.452

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	111	272	120	140	127	0	192	197
N.S.	1	1.01	2.47	1.09	1.27	1.15	0.00	1.75	1.79
time (sec)	N/A	0.944	2.638	2.520	0.366	0.318	0.000	0.316	0.385

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	114	116	208	129	165	137	0	192	207
N.S.	1	1.02	1.82	1.13	1.45	1.20	0.00	1.68	1.82
time (sec)	N/A	0.991	3.261	3.426	0.235	0.303	0.000	0.333	0.397

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	131	70	142	212	141	0	189	209
N.S.	1	1.05	0.56	1.14	1.70	1.13	0.00	1.51	1.67
time (sec)	N/A	1.009	1.103	4.017	0.220	0.333	0.000	0.330	0.361

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	165	173	178	269	145	0	212	185
N.S.	1	1.07	1.12	1.16	1.75	0.94	0.00	1.38	1.20
time (sec)	N/A	1.204	0.982	4.707	0.220	0.281	0.000	0.351	2.895

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	191	101	196	337	165	0	246	224
N.S.	1	1.03	0.55	1.06	1.82	0.89	0.00	1.33	1.21
time (sec)	N/A	1.364	1.761	5.860	0.222	0.312	0.000	0.327	2.964

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	241	242	156	131	356	150	960	193	353
N.S.	1	1.00	0.65	0.54	1.48	0.62	3.98	0.80	1.46
time (sec)	N/A	1.582	0.425	5.701	0.223	0.314	0.592	0.336	1.794

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	170	134	111	297	130	765	166	316
N.S.	1	0.92	0.72	0.60	1.61	0.70	4.14	0.90	1.71
time (sec)	N/A	0.664	0.273	4.098	0.238	0.311	0.427	0.345	1.697

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	125	133	94	236	110	544	139	278
N.S.	1	0.83	0.89	0.63	1.57	0.73	3.63	0.93	1.85
time (sec)	N/A	0.387	0.578	3.672	0.218	0.310	0.310	0.306	1.642

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	162	138	108	198	118	0	214	188
N.S.	1	1.07	0.91	0.72	1.31	0.78	0.00	1.42	1.25
time (sec)	N/A	1.232	1.738	3.459	0.215	0.326	0.000	0.312	0.738

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	156	312	140	187	150	0	226	242
N.S.	1	1.04	2.08	0.93	1.25	1.00	0.00	1.51	1.61
time (sec)	N/A	1.293	4.165	3.506	0.229	0.341	0.000	0.340	0.495

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	162	154	343	168	199	156	0	230	243
N.S.	1	0.95	2.12	1.04	1.23	0.96	0.00	1.42	1.50
time (sec)	N/A	1.301	7.850	3.288	0.231	0.321	0.000	0.338	0.488

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	165	172	380	165	235	159	0	227	254
N.S.	1	1.04	2.30	1.00	1.42	0.96	0.00	1.38	1.54
time (sec)	N/A	1.399	9.285	4.061	0.214	0.324	0.000	0.357	0.507

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	173	184	178	203	307	157	0	223	255
N.S.	1	1.06	1.03	1.17	1.77	0.91	0.00	1.29	1.47
time (sec)	N/A	1.384	3.682	4.777	0.233	0.339	0.000	0.367	0.466

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	214	217	217	376	165	0	246	224
N.S.	1	1.08	1.10	1.10	1.90	0.83	0.00	1.24	1.13
time (sec)	N/A	1.566	4.345	5.591	0.221	0.309	0.000	0.361	2.938

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	240	259	251	464	185	0	280	262
N.S.	1	1.05	1.13	1.10	2.03	0.81	0.00	1.22	1.14
time (sec)	N/A	1.784	5.109	6.185	0.296	0.301	0.000	0.371	2.954

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	134	311	96	394	120	1794	181	170
N.S.	1	0.88	2.03	0.63	2.58	0.78	11.73	1.18	1.11
time (sec)	N/A	0.682	1.442	0.934	0.313	0.329	1.981	0.326	0.404

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	107	249	78	310	98	1161	151	138
N.S.	1	0.88	2.04	0.64	2.54	0.80	9.52	1.24	1.13
time (sec)	N/A	0.556	1.228	0.951	0.298	0.294	1.268	0.312	1.496

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	98	197	61	225	83	665	124	107
N.S.	1	1.09	2.19	0.68	2.50	0.92	7.39	1.38	1.19
time (sec)	N/A	0.400	1.026	0.875	0.303	0.301	0.873	0.293	0.573

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	47	93	42	143	61	264	78	65
N.S.	1	0.87	1.72	0.78	2.65	1.13	4.89	1.44	1.20
time (sec)	N/A	0.457	0.373	0.835	0.301	0.293	0.573	0.292	0.323

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	76	28	73	43	49	43	30
N.S.	1	1.00	2.24	0.82	2.15	1.26	1.44	1.26	0.88
time (sec)	N/A	0.271	0.247	0.926	0.289	0.321	0.356	0.296	0.239

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	109	54	99	74	0	71	42
N.S.	1	1.00	2.48	1.23	2.25	1.68	0.00	1.61	0.95
time (sec)	N/A	0.328	0.468	1.269	0.215	0.321	0.000	0.314	0.263

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	70	201	93	196	127	0	110	78
N.S.	1	1.01	2.91	1.35	2.84	1.84	0.00	1.59	1.13
time (sec)	N/A	0.529	1.374	1.209	0.226	0.301	0.000	0.331	0.352

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	100	289	126	282	156	0	157	119
N.S.	1	0.93	2.70	1.18	2.64	1.46	0.00	1.47	1.11
time (sec)	N/A	0.659	2.937	1.215	0.212	0.309	0.000	0.318	0.428

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	118	490	170	368	168	0	182	152
N.S.	1	0.90	3.74	1.30	2.81	1.28	0.00	1.39	1.16
time (sec)	N/A	0.695	3.697	1.404	0.219	0.301	0.000	0.320	0.707

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	160	369	108	372	154	1425	192	189
N.S.	1	0.94	2.17	0.64	2.19	0.91	8.38	1.13	1.11
time (sec)	N/A	0.874	1.809	0.973	0.328	0.305	3.177	0.310	0.347

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	152	315	88	283	138	843	164	152
N.S.	1	1.03	2.14	0.60	1.93	0.94	5.73	1.12	1.03
time (sec)	N/A	0.655	1.539	0.964	0.299	0.295	1.813	0.310	0.299

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	100	137	73	191	117	411	119	105
N.S.	1	1.01	1.38	0.74	1.93	1.18	4.15	1.20	1.06
time (sec)	N/A	0.769	1.285	1.085	0.320	0.325	1.091	0.322	0.277

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	73	153	49	120	91	105	86	65
N.S.	1	1.04	2.19	0.70	1.71	1.30	1.50	1.23	0.93
time (sec)	N/A	0.521	0.750	0.873	0.406	0.286	0.680	0.302	0.242

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	64	43	42	93	58	94	60	45
N.S.	1	0.98	0.66	0.65	1.43	0.89	1.45	0.92	0.69
time (sec)	N/A	0.303	0.100	0.898	0.208	0.292	0.503	0.307	0.195

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	80	170	75	145	131	0	113	74
N.S.	1	1.01	2.15	0.95	1.84	1.66	0.00	1.43	0.94
time (sec)	N/A	0.515	0.658	1.239	0.222	0.324	0.000	0.331	0.262

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	115	264	126	244	207	0	155	123
N.S.	1	1.07	2.47	1.18	2.28	1.93	0.00	1.45	1.15
time (sec)	N/A	0.792	1.792	1.441	0.261	0.290	0.000	0.345	0.299

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	151	496	158	336	228	0	198	165
N.S.	1	0.99	3.26	1.04	2.21	1.50	0.00	1.30	1.09
time (sec)	N/A	0.962	2.940	1.447	0.257	0.297	0.000	0.332	0.308

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	171	609	196	425	247	0	226	203
N.S.	1	0.96	3.40	1.09	2.37	1.38	0.00	1.26	1.13
time (sec)	N/A	1.018	4.023	1.397	0.228	0.321	0.000	0.352	0.354

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	218	216	491	124	412	205	1584	228	238
N.S.	1	0.99	2.25	0.57	1.89	0.94	7.27	1.05	1.09
time (sec)	N/A	1.209	2.553	0.990	0.305	0.314	5.949	0.333	0.323

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	193	207	435	106	322	190	966	200	203
N.S.	1	1.07	2.25	0.55	1.67	0.98	5.01	1.04	1.05
time (sec)	N/A	0.972	2.170	0.979	0.290	0.297	3.904	0.329	0.286

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	154	361	89	231	165	496	155	152
N.S.	1	1.05	2.46	0.61	1.57	1.12	3.37	1.05	1.03
time (sec)	N/A	1.114	1.863	1.111	0.294	0.322	2.371	0.337	0.282

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	125	241	69	160	137	148	120	134
N.S.	1	1.08	2.08	0.59	1.38	1.18	1.28	1.03	1.16
time (sec)	N/A	0.821	1.207	0.882	0.308	0.308	1.342	0.310	0.420

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	104	135	56	115	93	117	75	66
N.S.	1	1.02	1.32	0.55	1.13	0.91	1.15	0.74	0.65
time (sec)	N/A	0.578	0.818	0.906	0.207	0.310	1.006	0.322	0.204

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	99	63	56	115	93	114	75	66
N.S.	1	0.97	0.62	0.55	1.13	0.91	1.12	0.74	0.65
time (sec)	N/A	0.411	0.140	1.125	0.364	0.304	0.779	0.301	0.221

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	127	197	95	187	185	0	148	130
N.S.	1	1.09	1.68	0.81	1.60	1.58	0.00	1.26	1.11
time (sec)	N/A	0.765	0.936	1.122	0.232	0.303	0.000	0.339	0.260

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	161	482	143	286	272	0	190	168
N.S.	1	1.11	3.32	0.99	1.97	1.88	0.00	1.31	1.16
time (sec)	N/A	1.139	2.776	1.448	0.219	0.318	0.000	0.322	0.294

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	203	610	175	377	295	0	233	216
N.S.	1	1.04	3.11	0.89	1.92	1.51	0.00	1.19	1.10
time (sec)	N/A	1.365	4.025	1.344	0.246	0.336	0.000	0.353	0.301

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	251	555	123	364	238	1085	233	259
N.S.	1	1.10	2.42	0.54	1.59	1.04	4.74	1.02	1.13
time (sec)	N/A	1.268	5.026	1.159	0.299	0.319	7.966	0.348	0.337

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	202	481	106	271	213	578	188	201
N.S.	1	1.09	2.60	0.57	1.46	1.15	3.12	1.02	1.09
time (sec)	N/A	1.451	4.767	0.969	0.300	0.307	5.071	0.332	0.427

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	173	329	89	201	180	192	155	162
N.S.	1	1.12	2.14	0.58	1.31	1.17	1.25	1.01	1.05
time (sec)	N/A	1.113	4.365	0.858	0.305	0.297	3.240	0.324	0.371

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	147	193	82	175	124	182	117	86
N.S.	1	1.08	1.42	0.60	1.29	0.91	1.34	0.86	0.63
time (sec)	N/A	0.879	3.090	0.975	0.214	0.287	2.197	0.395	0.278

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	137	163	76	174	124	178	117	84
N.S.	1	0.99	1.18	0.55	1.26	0.90	1.29	0.85	0.61
time (sec)	N/A	0.678	1.085	0.970	0.214	0.272	1.628	0.348	0.294

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	132	81	80	175	125	177	117	87
N.S.	1	0.96	0.59	0.58	1.27	0.91	1.28	0.85	0.63
time (sec)	N/A	0.516	0.341	1.280	0.221	0.312	1.431	0.329	0.299

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	167	239	113	228	236	0	182	199
N.S.	1	1.14	1.63	0.77	1.55	1.61	0.00	1.24	1.35
time (sec)	N/A	1.033	1.444	1.172	0.236	0.302	0.000	0.360	0.428

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	199	595	160	326	337	0	224	236
N.S.	1	1.14	3.40	0.91	1.86	1.93	0.00	1.28	1.35
time (sec)	N/A	1.442	4.604	1.473	0.225	0.311	0.000	0.361	0.353

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	232	247	798	192	419	360	0	267	273
N.S.	1	1.06	3.44	0.83	1.81	1.55	0.00	1.15	1.18
time (sec)	N/A	1.736	8.170	1.536	0.213	0.323	0.000	0.385	0.362

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	187	187	103	121	145	99	0	182	0
N.S.	1	1.00	0.55	0.65	0.78	0.53	0.00	0.97	0.00
time (sec)	N/A	0.893	0.395	2.710	0.406	0.305	0.000	0.689	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	146	80	102	118	82	0	147	0
N.S.	1	1.01	0.56	0.71	0.82	0.57	0.00	1.02	0.00
time (sec)	N/A	0.660	0.184	2.756	0.416	0.297	0.000	0.438	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	112	64	83	88	64	0	107	0
N.S.	1	1.11	0.63	0.82	0.87	0.63	0.00	1.06	0.00
time (sec)	N/A	0.601	0.109	2.714	0.372	0.300	0.000	0.332	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	46	62	57	47	0	70	0
N.S.	1	1.00	0.74	1.00	0.92	0.76	0.00	1.13	0.00
time (sec)	N/A	0.305	0.057	2.239	0.377	0.282	0.000	0.331	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	66	219	21	127	0	89	0
N.S.	1	1.00	1.00	3.32	0.32	1.92	0.00	1.35	0.00
time (sec)	N/A	0.389	0.069	3.710	0.345	0.317	0.000	0.330	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	85	567	710	153	0	121	0
N.S.	1	1.00	1.25	8.34	10.44	2.25	0.00	1.78	0.00
time (sec)	N/A	0.422	0.135	4.707	0.421	0.320	0.000	0.339	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	109	101	936	3352	178	0	195	0
N.S.	1	0.93	0.86	8.00	28.65	1.52	0.00	1.67	0.00
time (sec)	N/A	0.555	0.568	5.240	3.241	0.302	0.000	0.346	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	150	129	1282	5021	197	0	244	0
N.S.	1	0.94	0.81	8.01	31.38	1.23	0.00	1.52	0.00
time (sec)	N/A	0.732	1.188	5.806	3.229	0.322	0.000	0.439	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	234	238	125	142	185	125	0	235	0
N.S.	1	1.02	0.53	0.61	0.79	0.53	0.00	1.00	0.00
time (sec)	N/A	1.251	0.569	2.552	0.395	0.276	0.000	2.301	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	197	103	123	154	107	0	191	0
N.S.	1	1.04	0.54	0.65	0.81	0.57	0.00	1.01	0.00
time (sec)	N/A	0.987	0.301	2.589	0.393	0.279	0.000	1.058	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	146	81	104	123	88	0	155	0
N.S.	1	1.06	0.59	0.75	0.89	0.64	0.00	1.12	0.00
time (sec)	N/A	0.756	0.208	2.603	0.368	0.271	0.000	0.511	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	99	65	85	93	69	0	115	0
N.S.	1	0.98	0.64	0.84	0.92	0.68	0.00	1.14	0.00
time (sec)	N/A	0.412	0.131	2.503	0.368	0.309	0.000	0.359	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	108	85	269	39	149	0	140	0
N.S.	1	1.03	0.81	2.56	0.37	1.42	0.00	1.33	0.00
time (sec)	N/A	0.630	0.150	4.250	0.344	0.288	0.000	0.402	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	109	98	603	1315	172	0	149	0
N.S.	1	1.06	0.95	5.85	12.77	1.67	0.00	1.45	0.00
time (sec)	N/A	0.651	0.208	4.811	0.374	0.324	0.000	0.407	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	118	109	938	3339	182	0	201	0
N.S.	1	0.99	0.92	7.88	28.06	1.53	0.00	1.69	0.00
time (sec)	N/A	0.700	0.354	5.389	0.496	0.335	0.000	0.408	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	160	132	1283	7567	202	0	252	0
N.S.	1	0.98	0.80	7.82	46.14	1.23	0.00	1.54	0.00
time (sec)	N/A	0.879	0.628	5.879	154.990	0.342	0.000	0.444	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	201	151	1617	10504	220	0	302	0
N.S.	1	0.96	0.72	7.74	50.26	1.05	0.00	1.44	0.00
time (sec)	N/A	1.103	0.986	6.362	154.623	0.331	0.000	0.484	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	249	127	142	207	137	0	257	0
N.S.	1	1.05	0.54	0.60	0.87	0.58	0.00	1.08	0.00
time (sec)	N/A	1.374	0.630	11.388	0.398	0.308	0.000	3.285	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	180	105	123	172	116	0	213	0
N.S.	1	1.03	0.60	0.70	0.98	0.66	0.00	1.22	0.00
time (sec)	N/A	0.884	0.377	4.783	0.390	0.285	0.000	1.131	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	133	83	104	139	95	0	169	0
N.S.	1	0.96	0.60	0.75	1.01	0.69	0.00	1.22	0.00
time (sec)	N/A	0.509	0.220	3.231	0.361	0.303	0.000	0.569	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	150	104	310	61	177	0	202	0
N.S.	1	1.06	0.73	2.18	0.43	1.25	0.00	1.42	0.00
time (sec)	N/A	0.862	0.271	6.214	0.336	0.327	0.000	0.692	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	153	120	683	8114	202	0	209	0
N.S.	1	1.06	0.83	4.74	56.35	1.40	0.00	1.45	0.00
time (sec)	N/A	0.920	0.351	15.191	0.647	0.394	0.000	0.657	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	159	126	995	11782	204	0	239	0
N.S.	1	1.02	0.81	6.38	75.53	1.31	0.00	1.53	0.00
time (sec)	N/A	0.963	0.431	54.749	3.425	0.325	0.000	0.687	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	169	131	1282	7994	212	0	268	0
N.S.	1	1.03	0.80	7.82	48.74	1.29	0.00	1.63	0.00
time (sec)	N/A	1.010	0.713	163.576	3.619	0.340	0.000	0.686	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	212	152	1650	0	232	0	322	0
N.S.	1	1.01	0.73	7.89	0.00	1.11	0.00	1.54	0.00
time (sec)	N/A	1.207	1.118	2.423	0.000	0.357	0.000	0.732	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	254	253	176	1975	0	252	0	376	0
N.S.	1	1.00	0.69	7.78	0.00	0.99	0.00	1.48	0.00
time (sec)	N/A	1.419	1.422	2.584	0.000	0.348	0.000	0.794	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	226	111	281	1604723	184	0	217	0
N.S.	1	1.12	0.55	1.39	7944.17	0.91	0.00	1.07	0.00
time (sec)	N/A	1.303	0.433	4.207	32.549	0.317	0.000	0.727	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	174	94	240	927957	166	0	183	0
N.S.	1	1.09	0.59	1.51	5836.21	1.04	0.00	1.15	0.00
time (sec)	N/A	0.948	0.241	4.263	17.267	0.331	0.000	0.464	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	125	78	194	38386	149	0	149	160
N.S.	1	1.06	0.66	1.64	325.31	1.26	0.00	1.26	1.36
time (sec)	N/A	0.616	0.106	3.727	0.900	0.320	0.000	0.326	0.397

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	60	160	19040	135	0	125	112
N.S.	1	1.00	0.77	2.05	244.10	1.73	0.00	1.60	1.44
time (sec)	N/A	0.322	0.049	2.799	0.593	0.311	0.000	0.320	0.367

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	72	272	91	171	0	165	0
N.S.	1	1.00	0.79	2.99	1.00	1.88	0.00	1.81	0.00
time (sec)	N/A	0.484	0.050	4.608	0.340	0.336	0.000	0.346	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	128	95	703	18436	259	0	219	0
N.S.	1	1.08	0.80	5.91	154.92	2.18	0.00	1.84	0.00
time (sec)	N/A	0.751	0.241	5.452	0.528	0.332	0.000	0.354	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	165	177	114	1152	76209	284	0	287	0
N.S.	1	1.07	0.69	6.98	461.87	1.72	0.00	1.74	0.00
time (sec)	N/A	1.044	0.538	5.878	2.970	0.339	0.000	0.356	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	287	167	448	0	241	0	0	0
N.S.	1	1.10	0.64	1.72	0.00	0.92	0.00	0.00	0.00
time (sec)	N/A	1.762	0.737	4.919	0.000	0.326	0.000	0.000	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	232	142	445	0	224	0	0	0
N.S.	1	1.07	0.66	2.06	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	1.342	0.605	4.571	0.000	0.319	0.000	0.000	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	180	97	327	0	205	0	0	0
N.S.	1	1.05	0.57	1.91	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	0.952	0.497	4.217	0.000	0.288	0.000	0.000	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	121	104	256	0	189	0	0	0
N.S.	1	1.03	0.88	2.17	0.00	1.60	0.00	0.00	0.00
time (sec)	N/A	0.608	0.278	4.095	0.000	0.319	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	87	63	220	62254	172	0	160	0
N.S.	1	1.00	0.72	2.53	715.56	1.98	0.00	1.84	0.00
time (sec)	N/A	0.340	0.143	4.064	2.480	0.295	0.000	0.394	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	133	131	364	15722	281	0	220	0
N.S.	1	1.05	1.03	2.87	123.80	2.21	0.00	1.73	0.00
time (sec)	N/A	0.745	0.417	5.186	0.841	0.345	0.000	0.863	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	178	141	860	47933	339	0	0	0
N.S.	1	1.05	0.83	5.06	281.96	1.99	0.00	0.00	0.00
time (sec)	N/A	1.093	0.770	5.953	1.751	0.364	0.000	0.000	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	232	205	1372	0	361	0	354	0
N.S.	1	1.05	0.93	6.21	0.00	1.63	0.00	1.60	0.00
time (sec)	N/A	1.480	0.994	6.819	0.000	0.373	0.000	0.487	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	287	139	467	0	270	0	309	0
N.S.	1	1.10	0.53	1.79	0.00	1.03	0.00	1.18	0.00
time (sec)	N/A	1.778	1.008	5.100	0.000	0.298	0.000	10.413	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	232	117	397	0	254	0	263	0
N.S.	1	1.07	0.54	1.84	0.00	1.18	0.00	1.22	0.00
time (sec)	N/A	1.341	0.709	4.461	0.000	0.332	0.000	5.930	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	178	100	327	0	237	0	236	0
N.S.	1	1.05	0.59	1.93	0.00	1.40	0.00	1.40	0.00
time (sec)	N/A	0.981	0.486	4.474	0.000	0.343	0.000	3.200	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	132	87	292	0	223	0	206	0
N.S.	1	1.05	0.69	2.32	0.00	1.77	0.00	1.63	0.00
time (sec)	N/A	0.675	0.382	4.318	0.000	0.313	0.000	1.380	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	124	80	292	0	223	0	194	0
N.S.	1	0.98	0.63	2.32	0.00	1.77	0.00	1.54	0.00
time (sec)	N/A	0.448	0.321	4.235	0.000	0.315	0.000	0.752	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	176	126	430	84333	339	0	252	0
N.S.	1	1.07	0.77	2.62	514.23	2.07	0.00	1.54	0.00
time (sec)	N/A	1.050	1.040	5.787	15.383	0.366	0.000	3.036	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	207	223	142	925	0	404	0	335	0
N.S.	1	1.08	0.69	4.47	0.00	1.95	0.00	1.62	0.00
time (sec)	N/A	1.458	2.256	6.608	0.000	0.375	0.000	0.813	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	264	282	178	1438	0	428	0	0	0
N.S.	1	1.07	0.67	5.45	0.00	1.62	0.00	0.00	0.00
time (sec)	N/A	1.909	4.018	7.575	0.000	0.407	0.000	0.000	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	159	158	914	411	0	193	0	0	177
N.S.	1	0.99	5.75	2.58	0.00	1.21	0.00	0.00	1.11
time (sec)	N/A	0.833	6.998	13.836	0.000	0.141	0.000	0.000	1.146

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	132	130	872	383	0	179	0	0	166
N.S.	1	0.98	6.61	2.90	0.00	1.36	0.00	0.00	1.26
time (sec)	N/A	0.692	6.853	10.972	0.000	0.125	0.000	0.000	0.669

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	101	103	363	355	0	161	0	0	128
N.S.	1	1.02	3.59	3.51	0.00	1.59	0.00	0.00	1.27
time (sec)	N/A	0.643	5.462	8.569	0.000	0.107	0.000	0.000	0.574

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	70	73	309	321	0	142	0	0	79
N.S.	1	1.04	4.41	4.59	0.00	2.03	0.00	0.00	1.13
time (sec)	N/A	0.546	4.624	6.461	0.000	0.113	0.000	0.000	0.665

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	256	242	0	173	0	0	90
N.S.	1	1.00	3.88	3.67	0.00	2.62	0.00	0.00	1.36
time (sec)	N/A	0.533	4.324	5.780	0.000	0.122	0.000	0.000	1.161

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	95	97	293	399	0	196	0	0	150
N.S.	1	1.02	3.08	4.20	0.00	2.06	0.00	0.00	1.58
time (sec)	N/A	0.629	6.205	7.309	0.000	0.106	0.000	0.000	1.427

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	132	126	865	634	0	219	0	0	177
N.S.	1	0.95	6.55	4.80	0.00	1.66	0.00	0.00	1.34
time (sec)	N/A	0.680	6.805	10.376	0.000	0.130	0.000	0.000	1.863

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	194	190	944	413	0	223	0	0	266
N.S.	1	0.98	4.87	2.13	0.00	1.15	0.00	0.00	1.37
time (sec)	N/A	1.057	6.565	13.625	0.000	0.122	0.000	0.000	1.206

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	161	162	898	385	0	203	0	0	231
N.S.	1	1.01	5.58	2.39	0.00	1.26	0.00	0.00	1.43
time (sec)	N/A	0.989	6.544	11.960	0.000	0.124	0.000	0.000	1.110

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	126	132	369	357	0	179	0	0	153
N.S.	1	1.05	2.93	2.83	0.00	1.42	0.00	0.00	1.21
time (sec)	N/A	0.838	6.144	8.463	0.000	0.123	0.000	0.000	1.108

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	118	121	623	244	0	198	0	0	134
N.S.	1	1.03	5.28	2.07	0.00	1.68	0.00	0.00	1.14
time (sec)	N/A	0.828	7.365	7.375	0.000	0.123	0.000	0.000	1.241

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	120	124	311	513	0	210	0	0	196
N.S.	1	1.03	2.59	4.28	0.00	1.75	0.00	0.00	1.63
time (sec)	N/A	0.880	6.310	7.327	0.000	0.115	0.000	0.000	1.936

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	159	156	883	714	0	239	0	0	229
N.S.	1	0.98	5.55	4.49	0.00	1.50	0.00	0.00	1.44
time (sec)	N/A	0.990	7.893	10.220	0.000	0.123	0.000	0.000	2.138

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	194	186	925	824	0	263	0	0	235
N.S.	1	0.96	4.77	4.25	0.00	1.36	0.00	0.00	1.21
time (sec)	N/A	1.037	8.182	13.635	0.000	0.113	0.000	0.000	2.594

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	237	238	990	441	0	243	0	0	360
N.S.	1	1.00	4.18	1.86	0.00	1.03	0.00	0.00	1.52
time (sec)	N/A	1.381	6.680	16.941	0.000	0.123	0.000	0.000	1.380

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	204	210	944	413	0	223	0	0	323
N.S.	1	1.03	4.63	2.02	0.00	1.09	0.00	0.00	1.58
time (sec)	N/A	1.311	6.619	13.125	0.000	0.108	0.000	0.000	1.094

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	171	182	898	385	0	203	0	0	255
N.S.	1	1.06	5.25	2.25	0.00	1.19	0.00	0.00	1.49
time (sec)	N/A	1.184	7.630	11.546	0.000	0.116	0.000	0.000	1.079

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	169	175	888	337	0	229	0	0	229
N.S.	1	1.04	5.25	1.99	0.00	1.36	0.00	0.00	1.36
time (sec)	N/A	1.154	7.743	9.076	0.000	0.130	0.000	0.000	1.091

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	161	170	879	654	0	223	0	0	251
N.S.	1	1.06	5.46	4.06	0.00	1.39	0.00	0.00	1.56
time (sec)	N/A	1.152	7.953	9.469	0.000	0.128	0.000	0.000	1.691

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	171	175	890	916	0	243	0	0	287
N.S.	1	1.02	5.20	5.36	0.00	1.42	0.00	0.00	1.68
time (sec)	N/A	1.179	8.450	11.998	0.000	0.131	0.000	0.000	2.732

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	204	206	925	902	0	263	0	0	307
N.S.	1	1.01	4.53	4.42	0.00	1.29	0.00	0.00	1.50
time (sec)	N/A	1.330	8.883	13.761	0.000	0.126	0.000	0.000	2.770

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	237	234	967	1151	0	283	0	0	552
N.S.	1	0.99	4.08	4.86	0.00	1.19	0.00	0.00	2.33
time (sec)	N/A	1.372	9.712	18.177	0.000	0.115	0.000	0.000	3.290

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	156	148	946	281	0	269	0	0	0
N.S.	1	0.95	6.06	1.80	0.00	1.72	0.00	0.00	0.00
time (sec)	N/A	0.658	7.418	6.043	0.000	0.126	0.000	0.000	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	123	120	893	262	0	250	0	0	0
N.S.	1	0.98	7.26	2.13	0.00	2.03	0.00	0.00	0.00
time (sec)	N/A	0.620	7.064	5.016	0.000	0.111	0.000	0.000	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	85	90	862	244	0	237	0	0	0
N.S.	1	1.06	10.14	2.87	0.00	2.79	0.00	0.00	0.00
time (sec)	N/A	0.506	6.910	4.190	0.000	0.121	0.000	0.000	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	83	89	858	243	0	241	0	0	0
N.S.	1	1.07	10.34	2.93	0.00	2.90	0.00	0.00	0.00
time (sec)	N/A	0.508	7.274	3.110	0.000	0.119	0.000	0.000	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	119	116	894	319	0	290	0	0	0
N.S.	1	0.97	7.51	2.68	0.00	2.44	0.00	0.00	0.00
time (sec)	N/A	0.629	7.619	4.175	0.000	0.120	0.000	0.000	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	153	144	931	466	0	320	0	0	0
N.S.	1	0.94	6.08	3.05	0.00	2.09	0.00	0.00	0.00
time (sec)	N/A	0.664	8.029	5.486	0.000	0.125	0.000	0.000	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	203	200	1024	465	0	383	0	0	0
N.S.	1	0.99	5.04	2.29	0.00	1.89	0.00	0.00	0.00
time (sec)	N/A	0.991	7.799	8.276	0.000	0.128	0.000	0.000	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	166	171	980	435	0	367	0	0	0
N.S.	1	1.03	5.90	2.62	0.00	2.21	0.00	0.00	0.00
time (sec)	N/A	0.968	7.527	7.706	0.000	0.126	0.000	0.000	0.000

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	136	142	945	421	0	352	0	0	0
N.S.	1	1.04	6.95	3.10	0.00	2.59	0.00	0.00	0.00
time (sec)	N/A	0.780	7.264	5.200	0.000	0.121	0.000	0.000	0.000

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	121	130	694	350	0	314	0	0	0
N.S.	1	1.07	5.74	2.89	0.00	2.60	0.00	0.00	0.00
time (sec)	N/A	0.756	7.046	4.584	0.000	0.106	0.000	0.000	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	121	130	695	350	0	318	0	0	0
N.S.	1	1.07	5.74	2.89	0.00	2.63	0.00	0.00	0.00
time (sec)	N/A	0.764	7.660	3.987	0.000	0.115	0.000	0.000	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	168	168	979	494	0	407	0	0	0
N.S.	1	1.00	5.83	2.94	0.00	2.42	0.00	0.00	0.00
time (sec)	N/A	0.930	7.748	4.276	0.000	0.131	0.000	0.000	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	201	197	1020	723	0	436	0	0	0
N.S.	1	0.98	5.07	3.60	0.00	2.17	0.00	0.00	0.00
time (sec)	N/A	0.971	8.102	6.472	0.000	0.110	0.000	0.000	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	254	259	482	493	0	495	0	0	0
N.S.	1	1.02	1.90	1.94	0.00	1.95	0.00	0.00	0.00
time (sec)	N/A	1.371	5.265	15.764	0.000	0.158	0.000	0.000	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	219	229	441	465	0	478	0	0	0
N.S.	1	1.05	2.01	2.12	0.00	2.18	0.00	0.00	0.00
time (sec)	N/A	1.297	4.100	15.413	0.000	0.126	0.000	0.000	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	188	203	400	451	0	467	0	0	0
N.S.	1	1.08	2.13	2.40	0.00	2.48	0.00	0.00	0.00
time (sec)	N/A	1.119	3.052	15.039	0.000	0.122	0.000	0.000	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	195	845	451	0	465	0	0	0
N.S.	1	1.08	4.69	2.51	0.00	2.58	0.00	0.00	0.00
time (sec)	N/A	1.103	6.475	5.330	0.000	0.112	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	193	664	451	0	465	0	0	0
N.S.	1	1.08	3.73	2.53	0.00	2.61	0.00	0.00	0.00
time (sec)	N/A	1.073	6.346	5.433	0.000	0.112	0.000	0.000	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	182	197	1029	451	0	465	0	0	0
N.S.	1	1.08	5.65	2.48	0.00	2.55	0.00	0.00	0.00
time (sec)	N/A	1.129	7.866	5.203	0.000	0.118	0.000	0.000	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	221	227	1069	685	0	521	0	0	0
N.S.	1	1.03	4.84	3.10	0.00	2.36	0.00	0.00	0.00
time (sec)	N/A	1.315	7.921	5.097	0.000	0.133	0.000	0.000	0.000

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	C	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	254	255	1110	876	0	548	0	0	0
N.S.	1	1.00	4.37	3.45	0.00	2.16	0.00	0.00	0.00
time (sec)	N/A	1.350	8.361	6.065	0.000	0.131	0.000	0.000	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	209	135	348	8220	151	0	0	0
N.S.	1	0.95	0.61	1.57	37.19	0.68	0.00	0.00	0.00
time (sec)	N/A	0.922	0.677	15.005	0.825	0.390	0.000	0.000	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	176	166	118	276	2981	134	0	0	0
N.S.	1	0.94	0.67	1.57	16.94	0.76	0.00	0.00	0.00
time (sec)	N/A	0.755	0.364	14.315	0.649	0.349	0.000	0.000	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	123	100	206	1851	117	0	0	0
N.S.	1	0.94	0.76	1.57	14.13	0.89	0.00	0.00	0.00
time (sec)	N/A	0.582	0.201	15.518	0.538	0.400	0.000	0.000	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	83	141	939	97	0	0	0
N.S.	1	1.00	1.06	1.81	12.04	1.24	0.00	0.00	0.00
time (sec)	N/A	0.425	0.109	18.239	0.462	0.356	0.000	0.000	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	86	116	245	109	0	0	0
N.S.	1	1.00	1.13	1.53	3.22	1.43	0.00	0.00	0.00
time (sec)	N/A	0.408	0.119	7.563	0.392	0.342	0.000	0.000	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	57	62	289	67	0	0	112
N.S.	1	1.00	0.67	0.73	3.40	0.79	0.00	0.00	1.32
time (sec)	N/A	0.433	0.100	7.331	0.347	0.288	0.000	0.000	1.643

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	128	78	86	428	86	0	0	194
N.S.	1	0.98	0.60	0.66	3.29	0.66	0.00	0.00	1.49
time (sec)	N/A	0.585	0.175	7.384	0.342	0.303	0.000	0.000	3.441

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	171	102	108	522	104	0	136951	479
N.S.	1	0.98	0.58	0.62	2.98	0.59	0.00	782.58	2.74
time (sec)	N/A	0.753	0.270	7.376	0.353	0.291	0.000	237.662	6.435

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	219	136	349	8904	162	0	0	0
N.S.	1	0.96	0.60	1.54	39.22	0.71	0.00	0.00	0.00
time (sec)	N/A	1.083	0.741	15.596	0.932	0.392	0.000	0.000	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	176	119	277	3023	144	0	0	0
N.S.	1	0.98	0.66	1.54	16.79	0.80	0.00	0.00	0.00
time (sec)	N/A	0.864	0.433	15.192	0.658	0.334	0.000	0.000	0.000

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	132	101	207	1884	125	0	0	0
N.S.	1	0.99	0.76	1.56	14.17	0.94	0.00	0.00	0.00
time (sec)	N/A	0.675	0.243	18.612	0.550	0.367	0.000	0.000	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	107	189	1801	135	0	0	0
N.S.	1	1.00	0.85	1.50	14.29	1.07	0.00	0.00	0.00
time (sec)	N/A	0.689	0.223	18.348	0.537	0.355	0.000	0.000	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	128	106	188	1124	133	0	0	0
N.S.	1	1.02	0.85	1.50	8.99	1.06	0.00	0.00	0.00
time (sec)	N/A	0.675	0.259	7.799	0.448	0.321	0.000	0.000	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	139	80	87	344	88	0	0	195
N.S.	1	1.04	0.60	0.65	2.57	0.66	0.00	0.00	1.46
time (sec)	N/A	0.702	0.209	7.443	0.338	0.337	0.000	0.000	3.226

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	181	102	109	481	107	0	147871	236
N.S.	1	1.00	0.56	0.60	2.66	0.59	0.00	816.97	1.30
time (sec)	N/A	0.893	0.311	7.655	0.337	0.288	0.000	260.783	7.065

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	224	124	131	573	126	0	0	289
N.S.	1	0.98	0.54	0.57	2.51	0.55	0.00	0.00	1.27
time (sec)	N/A	1.097	0.395	7.777	0.353	0.296	0.000	0.000	7.453

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	274	273	159	423	10042	194	0	0	0
N.S.	1	1.00	0.58	1.54	36.65	0.71	0.00	0.00	0.00
time (sec)	N/A	1.450	1.265	16.231	0.965	0.377	0.000	0.000	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	230	137	351	9415	174	0	0	0
N.S.	1	1.01	0.60	1.55	41.48	0.77	0.00	0.00	0.00
time (sec)	N/A	1.188	0.770	15.850	0.925	0.431	0.000	0.000	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	185	121	279	3071	154	0	0	0
N.S.	1	1.03	0.67	1.55	17.06	0.86	0.00	0.00	0.00
time (sec)	N/A	1.007	0.482	20.252	0.658	0.371	0.000	0.000	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	178	126	263	2080	164	0	0	0
N.S.	1	1.00	0.71	1.48	11.69	0.92	0.00	0.00	0.00
time (sec)	N/A	1.007	0.453	18.326	0.540	0.342	0.000	0.000	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	173	177	130	267	2370	169	0	0	0
N.S.	1	1.02	0.75	1.54	13.70	0.98	0.00	0.00	0.00
time (sec)	N/A	1.024	0.464	18.648	0.557	0.345	0.000	0.000	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	172	180	130	227	1548	161	0	0	0
N.S.	1	1.05	0.76	1.32	9.00	0.94	0.00	0.00	0.00
time (sec)	N/A	0.966	0.522	7.944	0.469	0.333	0.000	0.000	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	191	104	111	396	114	0	169051	551
N.S.	1	1.06	0.57	0.61	2.19	0.63	0.00	933.98	3.04
time (sec)	N/A	1.044	0.397	8.281	0.342	0.321	0.000	244.268	7.192

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	235	126	133	533	135	0	0	647
N.S.	1	1.03	0.55	0.58	2.34	0.59	0.00	0.00	2.84
time (sec)	N/A	1.238	0.497	7.871	0.346	0.286	0.000	0.000	8.703

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	275	278	147	155	626	156	0	0	773
N.S.	1	1.01	0.53	0.56	2.28	0.57	0.00	0.00	2.81
time (sec)	N/A	1.481	0.602	7.415	0.363	0.290	0.000	0.000	7.565

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	190	203	228	253	0	184	0	0	0
N.S.	1	1.07	1.20	1.33	0.00	0.97	0.00	0.00	0.00
time (sec)	N/A	1.178	0.390	16.231	0.000	2.437	0.000	0.000	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	150	167	188	0	168	0	0	0
N.S.	1	1.06	1.18	1.33	0.00	1.19	0.00	0.00	0.00
time (sec)	N/A	0.809	0.262	16.632	0.000	1.360	0.000	0.000	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	82	132	1221	96	0	0	0
N.S.	1	1.00	0.82	1.32	12.21	0.96	0.00	0.00	0.00
time (sec)	N/A	0.549	0.099	8.207	0.972	1.100	0.000	0.000	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	C	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	203	199	1188	143	0	0	0
N.S.	1	1.00	2.05	2.01	12.00	1.44	0.00	0.00	0.00
time (sec)	N/A	0.437	1.116	7.980	0.958	0.351	0.000	0.000	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	C	A	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	142	150	627	271	1485	163	0	0	0
N.S.	1	1.06	4.42	1.91	10.46	1.15	0.00	0.00	0.00
time (sec)	N/A	0.717	6.666	7.802	0.942	0.327	0.000	0.000	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	C	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	187	205	1759	309	1826	180	0	0	0
N.S.	1	1.10	9.41	1.65	9.76	0.96	0.00	0.00	0.00
time (sec)	N/A	0.994	7.242	7.205	0.966	0.327	0.000	0.000	0.000

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	205	255	381	0	237	0	0	0
N.S.	1	1.04	1.29	1.93	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	1.177	0.878	16.219	0.000	4.055	0.000	0.000	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	150	171	279	0	203	0	0	0
N.S.	1	1.03	1.18	1.92	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.815	0.808	5.455	0.000	3.122	0.000	0.000	0.000

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	184	213	0	164	0	0	0
N.S.	1	1.00	1.72	1.99	0.00	1.53	0.00	0.00	0.00
time (sec)	N/A	0.453	0.773	7.895	0.000	0.334	0.000	0.000	0.000

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	156	158	423	353	0	201	0	0	0
N.S.	1	1.01	2.71	2.26	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.738	2.593	7.906	0.000	0.325	0.000	0.000	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	203	213	1054	403	0	221	0	0	0
N.S.	1	1.05	5.19	1.99	0.00	1.09	0.00	0.00	0.00
time (sec)	N/A	1.086	6.575	8.150	0.000	0.323	0.000	0.000	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	262	311	587	0	302	0	0	0
N.S.	1	1.07	1.26	2.39	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	1.589	2.186	17.798	0.000	8.570	0.000	0.000	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	205	252	447	0	267	0	0	0
N.S.	1	1.06	1.30	2.30	0.00	1.38	0.00	0.00	0.00
time (sec)	N/A	1.148	1.174	6.058	0.000	6.873	0.000	0.000	0.000

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	160	168	343	0	215	0	0	0
N.S.	1	1.04	1.09	2.23	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.718	4.631	5.956	0.000	0.326	0.000	0.000	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	162	150	343	0	217	0	0	0
N.S.	1	1.04	0.96	2.20	0.00	1.39	0.00	0.00	0.00
time (sec)	N/A	0.748	1.607	8.030	0.000	0.307	0.000	0.000	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	203	213	728	485	0	248	0	0	0
N.S.	1	1.05	3.59	2.39	0.00	1.22	0.00	0.00	0.00
time (sec)	N/A	1.075	6.661	8.185	0.000	0.322	0.000	0.000	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	250	268	1148	535	0	270	0	0	0
N.S.	1	1.07	4.59	2.14	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	1.426	8.072	7.977	0.000	0.309	0.000	0.000	0.000

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	293	317	378	795	0	368	0	0	0
N.S.	1	1.08	1.29	2.71	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	1.987	2.940	17.277	0.000	14.790	0.000	0.000	0.000

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	241	260	319	640	0	327	0	0	0
N.S.	1	1.08	1.32	2.66	0.00	1.36	0.00	0.00	0.00
time (sec)	N/A	1.518	2.168	5.945	0.000	8.464	0.000	0.000	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	215	613	475	0	266	0	0	0
N.S.	1	1.07	3.05	2.36	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	1.077	6.318	5.967	0.000	0.348	0.000	0.000	0.000

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	215	167	475	0	264	0	0	0
N.S.	1	1.07	0.83	2.36	0.00	1.31	0.00	0.00	0.00
time (sec)	N/A	1.069	1.669	5.885	0.000	0.306	0.000	0.000	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	203	217	168	475	0	266	0	0	0
N.S.	1	1.07	0.83	2.34	0.00	1.31	0.00	0.00	0.00
time (sec)	N/A	1.111	1.586	8.149	0.000	0.318	0.000	0.000	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	250	268	798	603	0	298	0	0	0
N.S.	1	1.07	3.19	2.41	0.00	1.19	0.00	0.00	0.00
time (sec)	N/A	1.413	6.914	8.032	0.000	0.333	0.000	0.000	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	297	323	1256	667	0	319	0	0	0
N.S.	1	1.09	4.23	2.25	0.00	1.07	0.00	0.00	0.00
time (sec)	N/A	1.802	8.436	7.896	0.000	0.358	0.000	0.000	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	97	91	86	101	81	252	89	117
N.S.	1	0.92	0.87	0.82	0.96	0.77	2.40	0.85	1.11
time (sec)	N/A	0.571	0.208	2.962	0.216	0.300	0.185	0.295	0.511

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	86	75	65	79	60	168	68	84
N.S.	1	1.02	0.89	0.77	0.94	0.71	2.00	0.81	1.00
time (sec)	N/A	0.400	0.151	2.049	0.202	0.287	0.138	0.308	0.425

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	52	51	51	55	42	94	45	50
N.S.	1	1.00	0.98	0.98	1.06	0.81	1.81	0.87	0.96
time (sec)	N/A	0.215	0.095	0.915	0.192	0.276	0.090	0.269	0.385

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	46	48	47	54	0	79	100
N.S.	1	1.00	1.31	1.37	1.34	1.54	0.00	2.26	2.86
time (sec)	N/A	0.435	0.029	1.621	0.202	0.294	0.000	0.295	0.517

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	43	50	73	85	0	84	114
N.S.	1	1.00	1.23	1.43	2.09	2.43	0.00	2.40	3.26
time (sec)	N/A	0.449	0.010	2.590	0.206	0.314	0.000	0.294	0.551

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	64	75	75	95	96	0	151	104
N.S.	1	1.05	1.23	1.23	1.56	1.57	0.00	2.48	1.70
time (sec)	N/A	0.583	0.013	3.249	0.212	0.292	0.000	0.305	1.408

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	91	67	81	127	115	0	210	145
N.S.	1	0.98	0.72	0.87	1.37	1.24	0.00	2.26	1.56
time (sec)	N/A	0.673	0.198	4.176	0.267	0.298	0.000	0.325	2.687

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	114	106	85	119	163	136	0	304	194
N.S.	1	0.93	0.75	1.04	1.43	1.19	0.00	2.67	1.70
time (sec)	N/A	0.698	0.439	4.341	0.200	0.325	0.000	0.304	4.119

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	164	146	147	176	142	459	156	307
N.S.	1	0.87	0.77	0.78	0.93	0.75	2.43	0.83	1.62
time (sec)	N/A	0.820	1.154	3.999	0.191	0.284	0.292	0.301	4.085

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	177	118	118	142	114	338	124	169
N.S.	1	1.04	0.69	0.69	0.84	0.67	1.99	0.73	0.99
time (sec)	N/A	0.639	1.639	2.762	0.193	0.296	0.198	0.289	0.532

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	110	90	88	108	85	199	93	115
N.S.	1	1.03	0.84	0.82	1.01	0.79	1.86	0.87	1.07
time (sec)	N/A	0.356	0.574	2.078	0.192	0.300	0.138	0.303	0.469

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	120	94	92	87	0	178	169
N.S.	1	1.00	1.40	1.09	1.07	1.01	0.00	2.07	1.97
time (sec)	N/A	0.618	0.940	1.532	0.223	0.291	0.000	0.291	0.759

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	109	79	103	117	0	152	169
N.S.	1	1.00	1.82	1.32	1.72	1.95	0.00	2.53	2.82
time (sec)	N/A	0.577	1.212	2.716	0.196	0.280	0.000	0.304	1.027

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	84	67	104	140	136	0	190	176
N.S.	1	1.05	0.84	1.30	1.75	1.70	0.00	2.38	2.20
time (sec)	N/A	0.642	0.186	3.717	0.208	0.285	0.000	0.313	1.166

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	122	92	118	172	150	0	294	227
N.S.	1	1.05	0.79	1.02	1.48	1.29	0.00	2.53	1.96
time (sec)	N/A	0.809	0.313	4.518	0.383	0.300	0.000	0.325	3.850

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	149	120	148	228	180	0	478	314
N.S.	1	0.96	0.77	0.95	1.46	1.15	0.00	3.06	2.01
time (sec)	N/A	0.929	0.481	5.230	0.198	0.292	0.000	0.320	4.014

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	269	231	289	213	266	211	721	230	352
N.S.	1	0.86	1.07	0.79	0.99	0.78	2.68	0.86	1.31
time (sec)	N/A	1.211	1.675	5.089	0.226	0.318	0.429	0.316	1.251

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	252	176	176	217	174	551	188	277
N.S.	1	1.04	0.72	0.72	0.89	0.72	2.27	0.77	1.14
time (sec)	N/A	0.880	2.309	3.947	0.205	0.285	0.297	0.321	0.898

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	179	140	142	171	136	386	148	202
N.S.	1	1.05	0.82	0.83	1.00	0.80	2.26	0.87	1.18
time (sec)	N/A	0.557	1.361	3.165	0.211	0.280	0.204	0.283	0.604

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	146	159	135	145	131	0	314	1924
N.S.	1	1.07	1.16	0.99	1.06	0.96	0.00	2.29	14.04
time (sec)	N/A	0.949	1.698	2.923	0.213	0.301	0.000	0.330	2.038

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	134	217	124	144	152	0	234	236
N.S.	1	1.02	1.66	0.95	1.10	1.16	0.00	1.79	1.80
time (sec)	N/A	0.922	1.963	3.241	0.201	0.306	0.000	0.310	1.459

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	124	126	277	133	169	167	0	239	249
N.S.	1	1.02	2.23	1.07	1.36	1.35	0.00	1.93	2.01
time (sec)	N/A	0.970	3.286	3.727	0.217	0.292	0.000	0.318	1.668

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	154	108	146	216	189	0	336	526
N.S.	1	1.06	0.74	1.01	1.49	1.30	0.00	2.32	3.63
time (sec)	N/A	1.028	0.401	4.875	0.201	0.314	0.000	0.319	2.169

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	188	199	140	185	273	211	0	586	395
N.S.	1	1.06	0.74	0.98	1.45	1.12	0.00	3.12	2.10
time (sec)	N/A	1.236	0.561	5.534	0.225	0.319	0.000	0.328	4.129

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	225	181	200	341	249	0	722	470
N.S.	1	0.95	0.77	0.85	1.44	1.06	0.00	3.06	1.99
time (sec)	N/A	1.447	2.095	6.201	0.209	0.330	0.000	0.361	4.445

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	366	315	408	273	366	289	1017	313	436
N.S.	1	0.86	1.11	0.75	1.00	0.79	2.78	0.86	1.19
time (sec)	N/A	1.806	1.893	6.263	0.204	0.320	0.594	0.334	2.909

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	325	336	333	242	307	243	811	263	403
N.S.	1	1.03	1.02	0.74	0.94	0.75	2.50	0.81	1.24
time (sec)	N/A	1.191	2.881	4.755	0.205	0.305	0.432	0.314	1.470

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	241	254	263	197	246	197	580	212	307
N.S.	1	1.05	1.09	0.82	1.02	0.82	2.41	0.88	1.27
time (sec)	N/A	0.798	1.527	3.922	0.204	0.282	0.312	0.333	0.963

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	214	210	191	208	183	0	603	369
N.S.	1	1.07	1.05	0.96	1.04	0.92	0.00	3.02	1.84
time (sec)	N/A	1.451	2.027	3.084	0.198	0.304	0.000	0.327	1.558

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	195	204	257	164	197	196	0	371	2522
N.S.	1	1.05	1.32	0.84	1.01	1.01	0.00	1.90	12.93
time (sec)	N/A	1.422	2.606	4.008	0.260	0.296	0.000	0.334	2.462

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	201	310	178	209	202	0	526	330
N.S.	1	0.96	1.48	0.85	1.00	0.97	0.00	2.52	1.58
time (sec)	N/A	1.471	3.245	3.854	0.198	0.300	0.000	0.383	2.654

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	205	415	175	245	219	0	387	636
N.S.	1	1.04	2.10	0.88	1.24	1.11	0.00	1.95	3.21
time (sec)	N/A	1.467	6.804	5.034	0.202	0.303	0.000	0.349	3.014

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	230	160	213	317	250	0	635	1969
N.S.	1	1.06	0.74	0.99	1.47	1.16	0.00	2.94	9.12
time (sec)	N/A	1.493	0.722	5.474	0.205	0.307	0.000	0.351	3.172

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	267	283	198	237	386	281	0	850	555
N.S.	1	1.06	0.74	0.89	1.45	1.05	0.00	3.18	2.08
time (sec)	N/A	1.732	2.677	6.398	0.262	0.290	0.000	0.380	4.167

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	324	302	244	277	474	327	0	1186	706
N.S.	1	0.93	0.75	0.85	1.46	1.01	0.00	3.66	2.18
time (sec)	N/A	1.967	1.956	7.462	0.230	0.309	0.000	0.371	4.181

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	195	152	240	0	541	0	360	4568
N.S.	1	1.10	0.85	1.35	0.00	3.04	0.00	2.02	25.66
time (sec)	N/A	1.046	0.942	1.212	0.000	0.361	0.000	0.294	5.321

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	146	121	169	0	426	10409	227	3761
N.S.	1	1.09	0.90	1.26	0.00	3.18	77.68	1.69	28.07
time (sec)	N/A	0.692	0.679	1.082	0.000	0.317	173.247	0.331	4.289

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	86	85	110	0	322	3225	142	541
N.S.	1	0.97	0.96	1.24	0.00	3.62	36.24	1.60	6.08
time (sec)	N/A	0.500	0.466	0.952	0.000	0.312	56.388	0.298	1.228

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	68	73	0	242	524	296	344
N.S.	1	1.00	1.01	1.09	0.00	3.61	7.82	4.42	5.13
time (sec)	N/A	0.300	0.135	1.010	0.000	0.323	11.239	0.297	1.871

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	112	92	0	304	0	127	342
N.S.	1	1.00	1.47	1.21	0.00	4.00	0.00	1.67	4.50
time (sec)	N/A	0.387	0.344	1.383	0.000	0.669	0.000	0.308	1.821

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	96	129	144	0	460	0	175	675
N.S.	1	0.97	1.30	1.45	0.00	4.65	0.00	1.77	6.82
time (sec)	N/A	0.521	0.737	1.645	0.000	0.395	0.000	0.315	2.301

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	143	154	300	229	0	589	0	269	4051
N.S.	1	1.08	2.10	1.60	0.00	4.12	0.00	1.88	28.33
time (sec)	N/A	0.990	1.675	1.561	0.000	2.785	0.000	0.357	4.442

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	187	204	422	335	0	729	0	412	4696
N.S.	1	1.09	2.26	1.79	0.00	3.90	0.00	2.20	25.11
time (sec)	N/A	1.401	1.995	1.860	0.000	0.732	0.000	0.324	5.410

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	263	277	184	267	0	965	0	338	6744
N.S.	1	1.05	0.70	1.02	0.00	3.67	0.00	1.29	25.64
time (sec)	N/A	1.302	1.496	1.399	0.000	0.355	0.000	0.295	9.932

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	185	147	205	0	788	0	1116	3276
N.S.	1	1.19	0.95	1.32	0.00	5.08	0.00	7.20	21.14
time (sec)	N/A	0.868	1.144	1.271	0.000	0.336	0.000	0.389	5.501

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	149	119	161	0	552	0	199	3775
N.S.	1	1.22	0.98	1.32	0.00	4.52	0.00	1.63	30.94
time (sec)	N/A	0.637	0.749	1.038	0.000	0.356	0.000	0.282	8.385

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	111	97	128	0	379	4974	159	113
N.S.	1	1.11	0.97	1.28	0.00	3.79	49.74	1.59	1.13
time (sec)	N/A	0.358	0.374	1.049	0.000	0.311	157.097	0.333	0.922

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	158	191	182	0	684	0	223	3763
N.S.	1	1.19	1.44	1.37	0.00	5.14	0.00	1.68	28.29
time (sec)	N/A	0.738	0.796	1.569	0.000	2.990	0.000	0.310	8.390

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	208	240	241	0	1088	0	404	5464
N.S.	1	1.10	1.27	1.28	0.00	5.76	0.00	2.14	28.91
time (sec)	N/A	1.211	1.695	2.107	0.000	8.040	0.000	0.305	13.252

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	284	438	326	0	1329	0	378	6692
N.S.	1	1.05	1.62	1.21	0.00	4.92	0.00	1.40	24.79
time (sec)	N/A	1.744	6.902	2.059	0.000	12.981	0.000	0.324	9.654

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	398	424	734	402	0	1812	0	2712	10598
N.S.	1	1.07	1.84	1.01	0.00	4.55	0.00	6.81	26.63
time (sec)	N/A	2.131	3.664	1.932	0.000	0.501	0.000	0.543	12.391

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	280	317	232	341	0	1561	0	543	5542
N.S.	1	1.13	0.83	1.22	0.00	5.58	0.00	1.94	19.79
time (sec)	N/A	1.407	2.257	1.703	0.000	0.414	0.000	0.361	7.989

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	247	204	282	0	1152	0	455	6923
N.S.	1	1.17	0.97	1.34	0.00	5.46	0.00	2.16	32.81
time (sec)	N/A	0.992	1.598	1.295	0.000	0.369	0.000	0.331	10.658

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	206	172	234	0	740	0	391	248
N.S.	1	1.14	0.96	1.30	0.00	4.11	0.00	2.17	1.38
time (sec)	N/A	0.702	1.006	1.155	0.000	0.362	0.000	0.324	3.763

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	190	157	232	0	742	0	390	248
N.S.	1	1.16	0.96	1.41	0.00	4.52	0.00	2.38	1.51
time (sec)	N/A	0.570	0.601	1.261	0.000	0.340	0.000	0.314	3.657

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	258	269	302	0	1400	0	481	6913
N.S.	1	1.21	1.26	1.41	0.00	6.54	0.00	2.25	32.30
time (sec)	N/A	1.200	1.250	2.221	0.000	13.011	0.000	0.339	9.896

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	299	335	352	376	0	2100	0	574	9312
N.S.	1	1.12	1.18	1.26	0.00	7.02	0.00	1.92	31.14
time (sec)	N/A	2.018	4.339	2.536	0.000	27.862	0.000	0.359	13.448

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	402	426	507	460	0	2416	0	1395	10547
N.S.	1	1.06	1.26	1.14	0.00	6.01	0.00	3.47	26.24
time (sec)	N/A	2.804	3.001	3.043	0.000	43.605	0.000	0.371	12.723

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	409	466	1155	552	0	2567	0	966	7823
N.S.	1	1.14	2.82	1.35	0.00	6.28	0.00	2.36	19.13
time (sec)	N/A	2.177	7.668	2.483	0.000	0.575	0.000	0.387	12.379

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	301	356	717	459	0	1857	0	813	9733
N.S.	1	1.18	2.38	1.52	0.00	6.17	0.00	2.70	32.34
time (sec)	N/A	1.446	3.630	1.730	0.000	0.494	0.000	0.350	12.761

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	274	319	251	371	0	1220	0	689	440
N.S.	1	1.16	0.92	1.35	0.00	4.45	0.00	2.51	1.61
time (sec)	N/A	1.095	1.636	1.580	0.000	0.381	0.000	0.350	6.715

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	263	302	252	384	0	1232	0	722	451
N.S.	1	1.15	0.96	1.46	0.00	4.68	0.00	2.75	1.71
time (sec)	N/A	1.029	1.290	1.447	0.000	0.382	0.000	0.335	6.626

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	279	227	372	0	1228	0	691	440
N.S.	1	1.18	0.96	1.57	0.00	5.18	0.00	2.92	1.86
time (sec)	N/A	0.860	1.676	1.540	0.000	0.401	0.000	0.323	4.000

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	301	365	368	479	0	2269	0	837	9727
N.S.	1	1.21	1.22	1.59	0.00	7.54	0.00	2.78	32.32
time (sec)	N/A	1.859	1.611	2.578	0.000	37.890	0.000	0.364	19.058

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	420	476	549	587	0	3393	0	996	13119
N.S.	1	1.13	1.31	1.40	0.00	8.08	0.00	2.37	31.24
time (sec)	N/A	3.048	2.969	3.304	0.000	76.714	0.000	0.366	25.786

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	547	591	781	672	0	3819	0	1090	14398
N.S.	1	1.08	1.43	1.23	0.00	6.98	0.00	1.99	26.32
time (sec)	N/A	3.913	4.639	5.062	0.000	113.158	0.000	0.372	14.276

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	27	28	23	0	25	56	25	24
N.S.	1	0.96	1.00	0.82	0.00	0.89	2.00	0.89	0.86
time (sec)	N/A	0.211	0.008	1.726	0.000	0.292	0.422	0.270	0.518

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	24	21	0	24	68	33	50
N.S.	1	1.00	0.89	0.78	0.00	0.89	2.52	1.22	1.85
time (sec)	N/A	0.204	0.012	0.811	0.000	0.305	0.388	0.282	0.912

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	23	12	0	11	31	11	11
N.S.	1	1.00	2.09	1.09	0.00	1.00	2.82	1.00	1.00
time (sec)	N/A	0.185	0.003	0.645	0.000	0.283	0.339	0.286	0.563

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	0	3	2	10	3
N.S.	1	1.00	1.00	1.33	0.00	1.00	0.67	3.33	1.00
time (sec)	N/A	0.136	0.000	0.302	0.000	0.259	0.065	0.274	0.522

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	20	0	31	39	47	16
N.S.	1	1.00	1.00	1.67	0.00	2.58	3.25	3.92	1.33
time (sec)	N/A	0.190	0.003	1.067	0.000	0.295	1.841	0.284	0.530

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	0	19	32	11	30
N.S.	1	1.00	1.00	1.09	0.00	1.73	2.91	1.00	2.73
time (sec)	N/A	0.204	0.002	1.830	0.000	0.283	1.440	0.289	0.511

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	36	37	0	64	0	52	73
N.S.	1	1.00	1.00	1.03	0.00	1.78	0.00	1.44	2.03
time (sec)	N/A	0.268	0.007	1.577	0.000	0.321	0.000	0.304	0.988

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	27	24	25	0	32	42	25	39
N.S.	1	0.96	0.86	0.89	0.00	1.14	1.50	0.89	1.39
time (sec)	N/A	0.222	0.030	1.767	0.000	0.256	8.067	0.292	0.693

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	114	122	98	139	0	350	0	185	173
N.S.	1	1.07	0.86	1.22	0.00	3.07	0.00	1.62	1.52
time (sec)	N/A	0.617	0.131	1.075	0.000	0.313	0.000	0.321	1.237

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	81	73	98	0	281	0	128	193
N.S.	1	1.03	0.92	1.24	0.00	3.56	0.00	1.62	2.44
time (sec)	N/A	0.423	0.068	0.977	0.000	0.301	0.000	0.301	0.897

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	59	66	0	231	0	245	101
N.S.	1	1.00	0.97	1.08	0.00	3.79	0.00	4.02	1.66
time (sec)	N/A	0.303	0.033	0.856	0.000	0.291	0.000	0.341	0.828

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	50	49	45	0	177	190	78	44
N.S.	1	1.00	0.98	0.90	0.00	3.54	3.80	1.56	0.88
time (sec)	N/A	0.222	0.017	0.803	0.000	0.297	155.041	0.286	0.535

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	103	86	0	292	0	122	101
N.S.	1	1.00	1.47	1.23	0.00	4.17	0.00	1.74	1.44
time (sec)	N/A	0.360	0.043	1.208	0.000	0.322	0.000	0.357	0.817

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	90	116	125	0	398	0	155	326
N.S.	1	1.02	1.32	1.42	0.00	4.52	0.00	1.76	3.70
time (sec)	N/A	0.491	0.239	1.324	0.000	0.366	0.000	0.308	1.119

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	132	239	194	0	487	0	221	1099
N.S.	1	1.07	1.94	1.58	0.00	3.96	0.00	1.80	8.93
time (sec)	N/A	0.821	0.757	1.423	0.000	0.474	0.000	0.309	1.903

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	386	408	292	1635	0	639	0	0	0
N.S.	1	1.06	0.76	4.24	0.00	1.66	0.00	0.00	0.00
time (sec)	N/A	2.182	1.410	17.280	0.000	0.194	0.000	0.000	0.000

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	303	317	232	1305	0	561	0	0	0
N.S.	1	1.05	0.77	4.31	0.00	1.85	0.00	0.00	0.00
time (sec)	N/A	1.583	0.975	15.457	0.000	0.135	0.000	0.000	0.000

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	231	237	179	993	0	492	0	0	0
N.S.	1	1.03	0.77	4.30	0.00	2.13	0.00	0.00	0.00
time (sec)	N/A	1.218	2.081	13.495	0.000	0.146	0.000	0.000	0.000

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	172	146	600	0	435	0	0	0
N.S.	1	1.01	0.85	3.51	0.00	2.54	0.00	0.00	0.00
time (sec)	N/A	0.829	0.536	10.721	0.000	0.134	0.000	0.000	0.000

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	179	107	247	0	0	0	0	0
N.S.	1	1.01	0.60	1.39	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.218	26.060	7.423	0.000	0.000	0.000	0.000	0.000

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	227	372	746	0	0	0	0	0
N.S.	1	1.07	1.75	3.50	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.783	15.819	10.244	0.000	0.000	0.000	0.000	0.000

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	292	298	420	1290	0	0	0	0	0
N.S.	1	1.02	1.44	4.42	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.504	3.020	12.902	0.000	0.000	0.000	0.000	0.000

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	378	389	480	2213	0	0	0	0	0
N.S.	1	1.03	1.27	5.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.249	5.125	18.684	0.000	0.000	0.000	0.000	0.000

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	378	394	291	1635	0	639	0	0	0
N.S.	1	1.04	0.77	4.33	0.00	1.69	0.00	0.00	0.00
time (sec)	N/A	2.070	1.827	17.544	0.000	0.168	0.000	0.000	0.000

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	305	233	1305	0	562	0	0	0
N.S.	1	1.03	0.78	4.39	0.00	1.89	0.00	0.00	0.00
time (sec)	N/A	1.582	3.092	15.979	0.000	0.146	0.000	0.000	0.000

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	225	231	203	993	0	493	0	0	0
N.S.	1	1.03	0.90	4.41	0.00	2.19	0.00	0.00	0.00
time (sec)	N/A	1.131	0.931	13.116	0.000	0.126	0.000	0.000	0.000

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	244	406	701	0	0	0	0	0
N.S.	1	1.03	1.72	2.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.967	2.198	10.703	0.000	0.000	0.000	0.000	0.000

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	232	246	398	991	0	0	0	0	0
N.S.	1	1.06	1.72	4.27	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.977	1.919	12.200	0.000	0.000	0.000	0.000	0.000

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	295	309	422	1403	0	0	0	0	0
N.S.	1	1.05	1.43	4.76	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.595	3.596	14.721	0.000	0.000	0.000	0.000	0.000

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	375	390	483	2327	0	0	0	0	0
N.S.	1	1.04	1.29	6.21	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.349	5.117	18.458	0.000	0.000	0.000	0.000	0.000

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	462	480	357	1983	0	726	0	0	0
N.S.	1	1.04	0.77	4.29	0.00	1.57	0.00	0.00	0.00
time (sec)	N/A	2.574	2.632	27.776	0.000	0.183	0.000	0.000	0.000

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	372	382	291	1635	0	639	0	0	0
N.S.	1	1.03	0.78	4.40	0.00	1.72	0.00	0.00	0.00
time (sec)	N/A	1.984	3.977	19.431	0.000	0.164	0.000	0.000	0.000

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	288	299	254	1305	0	562	0	0	0
N.S.	1	1.04	0.88	4.53	0.00	1.95	0.00	0.00	0.00
time (sec)	N/A	1.541	1.776	15.981	0.000	0.128	0.000	0.000	0.000

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	292	305	453	1067	0	0	0	0	0
N.S.	1	1.04	1.55	3.65	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.489	2.530	14.267	0.000	0.000	0.000	0.000	0.000

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	296	310	442	1490	0	0	0	0	0
N.S.	1	1.05	1.49	5.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.591	3.776	22.722	0.000	0.000	0.000	0.000	0.000

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	315	323	451	1742	0	0	0	0	0
N.S.	1	1.03	1.43	5.53	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.579	4.163	64.484	0.000	0.000	0.000	0.000	0.000

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	376	395	486	2438	0	0	0	0	0
N.S.	1	1.05	1.29	6.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.302	4.124	174.976	0.000	0.000	0.000	0.000	0.000

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	465	485	549	3548	0	0	0	0	0
N.S.	1	1.04	1.18	7.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.166	5.831	458.616	0.000	0.000	0.000	0.000	0.000

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	320	340	230	1305	0	562	0	0	0
N.S.	1	1.06	0.72	4.08	0.00	1.76	0.00	0.00	0.00
time (sec)	N/A	1.720	1.237	13.143	0.000	0.178	0.000	0.000	0.000

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	258	180	993	0	493	0	0	0
N.S.	1	1.05	0.73	4.04	0.00	2.00	0.00	0.00	0.00
time (sec)	N/A	1.230	1.042	11.777	0.000	0.141	0.000	0.000	0.000

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	187	154	671	0	435	0	0	199
N.S.	1	1.02	0.84	3.67	0.00	2.38	0.00	0.00	1.09
time (sec)	N/A	0.948	1.308	10.131	0.000	0.116	0.000	0.000	0.898

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	130	93	249	0	371	0	0	135
N.S.	1	1.00	0.72	1.92	0.00	2.85	0.00	0.00	1.04
time (sec)	N/A	0.604	2.993	6.027	0.000	0.110	0.000	0.000	1.015

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	118	81	194	0	0	0	0	0
N.S.	1	1.00	0.69	1.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.814	0.390	5.776	0.000	0.000	0.000	0.000	0.000

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	227	320	639	0	0	0	0	0
N.S.	1	1.05	1.48	2.96	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.808	13.549	8.256	0.000	0.000	0.000	0.000	0.000

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	299	305	420	1182	0	0	0	0	0
N.S.	1	1.02	1.40	3.95	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.501	4.190	10.812	0.000	0.000	0.000	0.000	0.000

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	387	391	304	1312	0	910	0	0	0
N.S.	1	1.01	0.79	3.39	0.00	2.35	0.00	0.00	0.00
time (sec)	N/A	2.039	1.796	14.767	0.000	0.206	0.000	0.000	0.000

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	262	280	189	1275	0	789	0	0	0
N.S.	1	1.07	0.72	4.87	0.00	3.01	0.00	0.00	0.00
time (sec)	N/A	1.435	1.472	14.819	0.000	0.174	0.000	0.000	0.000

Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	204	219	170	519	0	683	0	0	0
N.S.	1	1.07	0.83	2.54	0.00	3.35	0.00	0.00	0.00
time (sec)	N/A	1.097	0.905	12.082	0.000	0.149	0.000	0.000	0.000

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	196	151	432	0	610	0	0	0
N.S.	1	1.06	0.82	2.34	0.00	3.30	0.00	0.00	0.00
time (sec)	N/A	0.891	0.497	8.425	0.000	0.129	0.000	0.000	0.000

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	190	198	460	433	0	0	0	0	0
N.S.	1	1.04	2.42	2.28	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.256	2.778	9.937	0.000	0.000	0.000	0.000	0.000

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	303	335	482	912	0	0	0	0	0
N.S.	1	1.11	1.59	3.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.695	4.038	14.114	0.000	0.000	0.000	0.000	0.000

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	398	425	546	1568	0	0	0	0	0
N.S.	1	1.07	1.37	3.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.508	5.720	19.364	0.000	0.000	0.000	0.000	0.000

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	550	555	372	1750	0	1538	0	0	0
N.S.	1	1.01	0.68	3.18	0.00	2.80	0.00	0.00	0.00
time (sec)	N/A	3.019	3.189	30.716	0.000	0.347	0.000	0.000	0.000

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	413	421	334	1412	0	1348	0	0	0
N.S.	1	1.02	0.81	3.42	0.00	3.26	0.00	0.00	0.00
time (sec)	N/A	2.232	2.483	27.650	0.000	0.228	0.000	0.000	0.000

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	331	338	274	954	0	1193	0	0	0
N.S.	1	1.02	0.83	2.88	0.00	3.60	0.00	0.00	0.00
time (sec)	N/A	1.619	2.009	24.432	0.000	0.211	0.000	0.000	0.000

Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	307	317	224	864	0	1076	0	0	0
N.S.	1	1.03	0.73	2.81	0.00	3.50	0.00	0.00	0.00
time (sec)	N/A	1.528	1.690	21.465	0.000	0.158	0.000	0.000	0.000

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	275	285	193	754	0	956	0	0	0
N.S.	1	1.04	0.70	2.74	0.00	3.48	0.00	0.00	0.00
time (sec)	N/A	1.315	1.162	19.202	0.000	0.158	0.000	0.000	0.000

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	349	375	553	858	0	0	0	0	0
N.S.	1	1.07	1.58	2.46	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.887	4.917	19.819	0.000	0.000	0.000	0.000	0.000

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	437	471	750	1345	0	0	0	0	0
N.S.	1	1.08	1.72	3.08	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.751	6.882	29.578	0.000	0.000	0.000	0.000	0.000

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	532	569	820	2004	0	0	0	0	0
N.S.	1	1.07	1.54	3.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.688	7.191	40.589	0.000	0.000	0.000	0.000	0.000

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	58	79	0	148	0	0	0
N.S.	1	1.00	1.00	1.36	0.00	2.55	0.00	0.00	0.00
time (sec)	N/A	0.299	0.039	6.006	0.000	0.093	0.000	0.000	0.000

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	59	167	0	0	0	0	0
N.S.	1	1.00	1.00	2.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.398	0.033	7.436	0.000	0.000	0.000	0.000	0.000

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	108	84	218	0	491	0	0	0
N.S.	1	1.00	0.78	2.02	0.00	4.55	0.00	0.00	0.00
time (sec)	N/A	0.434	0.147	13.902	0.000	0.112	0.000	0.000	0.000

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	186	403	377	0	0	0	0	0
N.S.	1	1.04	2.25	2.11	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.195	3.415	16.666	0.000	0.000	0.000	0.000	0.000

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	162	125	451	0	211	0	0	177
N.S.	1	0.95	0.74	2.65	0.00	1.24	0.00	0.00	1.04
time (sec)	N/A	0.774	1.724	17.602	0.000	0.118	0.000	0.000	1.509

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	134	103	413	0	192	0	0	166
N.S.	1	0.96	0.74	2.95	0.00	1.37	0.00	0.00	1.19
time (sec)	N/A	0.669	1.320	11.266	0.000	0.112	0.000	0.000	1.271

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	107	86	371	0	175	0	0	128
N.S.	1	0.99	0.80	3.44	0.00	1.62	0.00	0.00	1.19
time (sec)	N/A	0.621	0.994	9.451	0.000	0.106	0.000	0.000	1.135

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	78	67	326	0	156	0	0	85
N.S.	1	1.04	0.89	4.35	0.00	2.08	0.00	0.00	1.13
time (sec)	N/A	0.518	0.669	7.774	0.000	0.100	0.000	0.000	1.123

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	64	246	0	185	0	0	96
N.S.	1	1.00	0.90	3.46	0.00	2.61	0.00	0.00	1.35
time (sec)	N/A	0.519	0.441	7.123	0.000	0.105	0.000	0.000	1.627

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	102	107	401	0	213	0	0	150
N.S.	1	0.99	1.04	3.89	0.00	2.07	0.00	0.00	1.46
time (sec)	N/A	0.625	0.545	9.123	0.000	0.104	0.000	0.000	2.383

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	130	134	636	0	235	0	0	177
N.S.	1	0.93	0.96	4.54	0.00	1.68	0.00	0.00	1.26
time (sec)	N/A	0.663	0.746	12.938	0.000	0.108	0.000	0.000	2.674

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	264	231	196	666	0	299	0	0	275
N.S.	1	0.88	0.74	2.52	0.00	1.13	0.00	0.00	1.04
time (sec)	N/A	1.057	2.332	42.909	0.000	0.133	0.000	0.000	1.700

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	223	203	167	610	0	271	0	0	264
N.S.	1	0.91	0.75	2.74	0.00	1.22	0.00	0.00	1.18
time (sec)	N/A	0.940	2.048	14.207	0.000	0.133	0.000	0.000	1.516

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	182	175	139	548	0	243	0	0	229
N.S.	1	0.96	0.76	3.01	0.00	1.34	0.00	0.00	1.26
time (sec)	N/A	0.913	1.860	12.631	0.000	0.110	0.000	0.000	1.488

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	146	106	487	0	216	0	0	177
N.S.	1	1.04	0.76	3.48	0.00	1.54	0.00	0.00	1.26
time (sec)	N/A	0.792	1.171	9.590	0.000	0.106	0.000	0.000	1.674

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	124	102	405	0	240	0	0	158
N.S.	1	1.02	0.84	3.35	0.00	1.98	0.00	0.00	1.31
time (sec)	N/A	0.729	1.008	9.836	0.000	0.104	0.000	0.000	1.849

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	129	105	574	0	255	0	0	194
N.S.	1	1.02	0.83	4.56	0.00	2.02	0.00	0.00	1.54
time (sec)	N/A	0.758	1.309	10.721	0.000	0.113	0.000	0.000	2.598

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	172	161	175	723	0	286	0	0	227
N.S.	1	0.94	1.02	4.20	0.00	1.66	0.00	0.00	1.32
time (sec)	N/A	0.878	1.352	13.069	0.000	0.115	0.000	0.000	2.889

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	305	272	235	825	0	358	0	0	364
N.S.	1	0.89	0.77	2.70	0.00	1.17	0.00	0.00	1.19
time (sec)	N/A	1.343	2.541	18.436	0.000	0.144	0.000	0.000	2.084

Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	255	244	197	745	0	321	0	0	328
N.S.	1	0.96	0.77	2.92	0.00	1.26	0.00	0.00	1.29
time (sec)	N/A	1.302	1.967	16.220	0.000	0.131	0.000	0.000	1.885

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	216	158	664	0	284	0	0	275
N.S.	1	1.05	0.77	3.24	0.00	1.39	0.00	0.00	1.34
time (sec)	N/A	1.159	2.379	13.337	0.000	0.122	0.000	0.000	1.587

Problem 362	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	208	150	641	0	302	0	0	248
N.S.	1	1.03	0.74	3.17	0.00	1.50	0.00	0.00	1.23
time (sec)	N/A	1.167	2.131	11.409	0.000	0.128	0.000	0.000	1.640

Problem 363	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	191	165	771	0	306	0	0	255
N.S.	1	0.99	0.86	4.02	0.00	1.59	0.00	0.00	1.33
time (sec)	N/A	1.137	2.129	12.889	0.000	0.129	0.000	0.000	2.655

Problem 364	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	204	208	176	950	0	326	0	0	291
N.S.	1	1.02	0.86	4.66	0.00	1.60	0.00	0.00	1.43
time (sec)	N/A	1.130	3.020	14.865	0.000	0.138	0.000	0.000	4.239

Problem 365	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	182	200	260	1074	0	0	0	0	0
N.S.	1	1.10	1.43	5.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.484	1.718	7.506	0.000	0.000	0.000	0.000	0.000

Problem 366	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	148	207	822	0	0	0	0	0
N.S.	1	1.08	1.51	6.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.030	1.100	5.879	0.000	0.000	0.000	0.000	0.000

Problem 367	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	86	128	295	0	0	0	0	0
N.S.	1	0.97	1.44	3.31	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.582	0.735	4.993	0.000	0.000	0.000	0.000	0.000

Problem 368	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	58	217	0	0	0	0	0
N.S.	1	1.00	0.95	3.56	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.434	0.351	3.301	0.000	0.000	0.000	0.000	0.000

Problem 369	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	86	206	300	0	0	0	0	0
N.S.	1	1.00	2.40	3.49	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.684	1.678	4.599	0.000	0.000	0.000	0.000	0.000

Problem 370	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	162	260	441	0	0	0	0	0
N.S.	1	1.08	1.73	2.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.440	1.570	6.415	0.000	0.000	0.000	0.000	0.000

Problem 371	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	303	295	318	1066	0	0	0	0	0
N.S.	1	0.97	1.05	3.52	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.871	2.375	19.240	0.000	0.000	0.000	0.000	0.000

Problem 372	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	224	220	280	849	0	0	0	0	0
N.S.	1	0.98	1.25	3.79	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.309	1.954	7.792	0.000	0.000	0.000	0.000	0.000

Problem 373	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	192	260	808	0	0	0	0	0
N.S.	1	0.97	1.31	4.08	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.194	1.828	6.708	0.000	0.000	0.000	0.000	0.000

Problem 374	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	185	274	721	0	0	0	0	0
N.S.	1	0.92	1.37	3.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.315	1.924	6.464	0.000	0.000	0.000	0.000	0.000

Problem 375	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	256	242	316	856	0	0	0	0	0
N.S.	1	0.95	1.23	3.34	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.887	2.904	7.465	0.000	0.000	0.000	0.000	0.000

Problem 376	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	345	327	363	1004	0	0	0	0	0
N.S.	1	0.95	1.05	2.91	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.593	4.619	11.490	0.000	0.000	0.000	0.000	0.000

Problem 377	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	367	372	390	1977	0	0	0	0	0
N.S.	1	1.01	1.06	5.39	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.171	3.427	83.352	0.000	0.000	0.000	0.000	0.000

Problem 378	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	344	349	360	1937	0	0	0	0	0
N.S.	1	1.01	1.05	5.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.061	2.540	11.477	0.000	0.000	0.000	0.000	0.000

Problem 379	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	337	334	365	1850	0	0	0	0	0
N.S.	1	0.99	1.08	5.49	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.012	3.119	11.049	0.000	0.000	0.000	0.000	0.000

Problem 380	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	345	336	383	1744	0	0	0	0	0
N.S.	1	0.97	1.11	5.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.238	3.341	10.655	0.000	0.000	0.000	0.000	0.000

Problem 381	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	420	412	458	1975	0	0	0	0	0
N.S.	1	0.98	1.09	4.70	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.071	3.813	12.740	0.000	0.000	0.000	0.000	0.000

Problem 382	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	523	513	570	2131	0	0	0	0	0
N.S.	1	0.98	1.09	4.07	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.002	7.133	20.434	0.000	0.000	0.000	0.000	0.000

Problem 383	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	41	203	0	77	0	0	0
N.S.	1	1.00	0.93	4.61	0.00	1.75	0.00	0.00	0.00
time (sec)	N/A	0.259	0.031	4.679	0.000	0.099	0.000	0.000	0.000

Problem 384	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	37	180	0	71	0	0	0
N.S.	1	1.00	0.84	4.09	0.00	1.61	0.00	0.00	0.00
time (sec)	N/A	0.265	0.025	3.364	0.000	0.097	0.000	0.000	0.000

Problem 385	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	134	0	59	0	0	0
N.S.	1	1.00	1.00	7.88	0.00	3.47	0.00	0.00	0.00
time (sec)	N/A	0.197	0.012	2.359	0.000	0.122	0.000	0.000	0.000

Problem 386	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	19	0	53	0	0	0
N.S.	1	1.00	1.00	1.12	0.00	3.12	0.00	0.00	0.00
time (sec)	N/A	0.196	0.013	0.491	0.000	0.088	0.000	0.000	0.000

Problem 387	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	40	183	0	96	0	0	0
N.S.	1	1.00	1.00	4.58	0.00	2.40	0.00	0.00	0.00
time (sec)	N/A	0.254	0.036	2.083	0.000	0.088	0.000	0.000	0.000

Problem 388	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	37	214	0	95	0	0	0
N.S.	1	1.00	0.84	4.86	0.00	2.16	0.00	0.00	0.00
time (sec)	N/A	0.253	0.043	2.045	0.000	0.094	0.000	0.000	0.000

Problem 389	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	125	159	553	0	0	0	0	0
N.S.	1	1.08	1.37	4.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.928	0.955	6.426	0.000	0.000	0.000	0.000	0.000

Problem 390	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	81	82	228	0	0	0	0	0
N.S.	1	1.04	1.05	2.92	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.572	0.055	3.870	0.000	0.000	0.000	0.000	0.000

Problem 391	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	49	189	0	0	0	0	0
N.S.	1	1.00	0.89	3.44	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.402	0.018	3.091	0.000	0.000	0.000	0.000	0.000

Problem 392	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	30	151	0	0	0	0	0
N.S.	1	1.00	1.00	5.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.243	0.025	1.999	0.000	0.000	0.000	0.000	0.000

Problem 393	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	79	196	355	0	0	0	0	0
N.S.	1	0.99	2.45	4.44	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.629	1.770	4.023	0.000	0.000	0.000	0.000	0.000

Problem 394	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	140	211	425	0	0	0	0	0
N.S.	1	1.05	1.59	3.20	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.284	2.606	6.313	0.000	0.000	0.000	0.000	0.000

Problem 395	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	560	563	1224	3997	0	0	0	0	0
N.S.	1	1.01	2.19	7.14	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.784	6.556	16.716	0.000	0.000	0.000	0.000	0.000

Problem 396	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	473	471	1175	2772	0	0	0	0	0
N.S.	1	1.00	2.48	5.86	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.062	19.790	9.717	0.000	0.000	0.000	0.000	0.000

Problem 397	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	385	391	408	1268	0	0	0	0	0
N.S.	1	1.02	1.06	3.29	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.573	7.635	11.872	0.000	0.000	0.000	0.000	0.000

Problem 398	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	351	351	273	953	0	0	0	0	0
N.S.	1	1.00	0.78	2.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.196	8.402	14.281	0.000	0.000	0.000	0.000	0.000

Problem 399	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	284	285	407	1955	0	0	0	0	0
N.S.	1	1.00	1.43	6.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.991	10.990	18.363	0.000	0.000	0.000	0.000	0.000

Problem 400	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	350	359	1315	3286	0	0	0	0	0
N.S.	1	1.03	3.76	9.39	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.507	6.553	22.892	0.000	0.000	0.000	0.000	0.000

Problem 401	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	433	450	1408	4604	0	0	0	0	0
N.S.	1	1.04	3.25	10.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.005	6.559	26.957	0.000	0.000	0.000	0.000	0.000

Problem 402	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	670	675	1284	5533	0	0	0	0	0
N.S.	1	1.01	1.92	8.26	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.448	7.007	18.241	0.000	0.000	0.000	0.000	0.000

Problem 403	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	566	568	1227	4259	0	0	0	0	0
N.S.	1	1.00	2.17	7.52	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.785	6.899	16.061	0.000	0.000	0.000	0.000	0.000

Problem 404	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	472	475	1198	3310	0	0	0	0	0
N.S.	1	1.01	2.54	7.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.181	7.023	17.602	0.000	0.000	0.000	0.000	0.000

Problem 405	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	449	461	1196	2380	0	0	0	0	0
N.S.	1	1.03	2.66	5.30	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.167	6.577	17.902	0.000	0.000	0.000	0.000	0.000

Problem 406	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	419	420	1236	2468	0	0	0	0	0
N.S.	1	1.00	2.95	5.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.671	6.539	19.027	0.000	0.000	0.000	0.000	0.000

Problem 407	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	353	359	1314	3550	0	0	0	0	0
N.S.	1	1.02	3.72	10.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.534	6.642	23.428	0.000	0.000	0.000	0.000	0.000

Problem 408	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	433	444	1407	4605	0	0	0	0	0
N.S.	1	1.03	3.25	10.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.063	6.682	27.508	0.000	0.000	0.000	0.000	0.000

Problem 409	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	522	533	1515	5953	0	0	0	0	0
N.S.	1	1.02	2.90	11.40	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.673	6.868	30.803	0.000	0.000	0.000	0.000	0.000

Problem 410	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	779	786	1353	7110	0	0	0	0	0
N.S.	1	1.01	1.74	9.13	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.335	7.572	20.893	0.000	0.000	0.000	0.000	0.000

Problem 411	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	664	672	1287	5791	0	0	0	0	0
N.S.	1	1.01	1.94	8.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.527	7.496	18.363	0.000	0.000	0.000	0.000	0.000

Problem 412	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	564	572	1251	4802	0	0	0	0	0
N.S.	1	1.01	2.22	8.51	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.934	7.475	19.204	0.000	0.000	0.000	0.000	0.000

Problem 413	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	547	555	1241	4453	0	0	0	0	0
N.S.	1	1.01	2.27	8.14	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.884	6.671	17.706	0.000	0.000	0.000	0.000	0.000

Problem 414	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	536	547	1269	4719	0	0	0	0	0
N.S.	1	1.02	2.37	8.80	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.845	6.691	20.898	0.000	0.000	0.000	0.000	0.000

Problem 415	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	493	499	1319	4433	0	0	0	0	0
N.S.	1	1.01	2.68	8.99	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.244	6.742	22.237	0.000	0.000	0.000	0.000	0.000

Problem 416	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	434	445	1409	4874	0	0	0	0	0
N.S.	1	1.03	3.25	11.23	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.035	6.867	26.956	0.000	0.000	0.000	0.000	0.000

Problem 417	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	522	533	1517	5955	0	0	0	0	0
N.S.	1	1.02	2.91	11.41	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.656	6.994	30.972	0.000	0.000	0.000	0.000	0.000

Problem 418	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	622	638	1640	7347	0	0	0	0	0
N.S.	1	1.03	2.64	11.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.386	7.131	37.176	0.000	0.000	0.000	0.000	0.000

Problem 419	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	418	423	458	3179	0	0	0	0	0
N.S.	1	1.01	1.10	7.61	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.830	12.687	21.444	0.000	0.000	0.000	0.000	0.000

Problem 420	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	479	480	1175	2525	0	0	0	0	0
N.S.	1	1.00	2.45	5.27	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.073	21.256	15.454	0.000	0.000	0.000	0.000	0.000

Problem 421	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	427	436	402	981	0	0	0	0	0
N.S.	1	1.02	0.94	2.30	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.791	7.041	11.545	0.000	0.000	0.000	0.000	0.000

Problem 422	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	228	144	181	0	0	0	0	0
N.S.	1	1.00	0.63	0.79	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.623	2.258	14.268	0.000	0.000	0.000	0.000	0.000

Problem 423	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	230	230	299	771	0	0	0	0	0
N.S.	1	1.00	1.30	3.35	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.704	9.684	15.736	0.000	0.000	0.000	0.000	0.000

Problem 424	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	290	294	416	1844	0	0	0	0	0
N.S.	1	1.01	1.43	6.36	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.012	12.138	19.463	0.000	0.000	0.000	0.000	0.000

Problem 425	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	363	375	1319	3290	0	0	0	0	0
N.S.	1	1.03	3.63	9.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.473	6.591	22.404	0.000	0.000	0.000	0.000	0.000

Problem 426	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	500	527	1234	3562	0	0	0	0	0
N.S.	1	1.05	2.47	7.12	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.432	6.832	16.509	0.000	0.000	0.000	0.000	0.000

Problem 427	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	416	420	1012	1819	0	0	0	0	0
N.S.	1	1.01	2.43	4.37	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.488	16.322	12.144	0.000	0.000	0.000	0.000	0.000

Problem 428	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	284	307	1223	1533	0	0	0	0	0
N.S.	1	1.08	4.31	5.40	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.014	6.666	13.703	0.000	0.000	0.000	0.000	0.000

Problem 429	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	305	331	1281	1978	0	0	0	0	0
N.S.	1	1.09	4.20	6.49	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.157	6.657	18.714	0.000	0.000	0.000	0.000	0.000

Problem 430	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	393	412	1357	3846	0	0	0	0	0
N.S.	1	1.05	3.45	9.79	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.764	6.775	22.205	0.000	0.000	0.000	0.000	0.000

Problem 431	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	674	698	1396	10417	0	0	0	0	0
N.S.	1	1.04	2.07	15.46	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.430	7.188	18.246	0.000	0.000	0.000	0.000	0.000

Problem 432	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	545	569	1342	6574	0	0	0	0	0
N.S.	1	1.04	2.46	12.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.390	6.947	16.556	0.000	0.000	0.000	0.000	0.000

Problem 433	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	391	410	1335	4743	0	0	0	0	0
N.S.	1	1.05	3.41	12.13	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.526	6.907	14.549	0.000	0.000	0.000	0.000	0.000

Problem 434	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	429	444	1384	5455	0	0	0	0	0
N.S.	1	1.03	3.23	12.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.709	6.616	18.115	0.000	0.000	0.000	0.000	0.000

Problem 435	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	456	477	1431	6516	0	0	0	0	0
N.S.	1	1.05	3.14	14.29	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.975	6.758	20.066	0.000	0.000	0.000	0.000	0.000

Problem 436	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	567	581	1499	8901	0	0	0	0	0
N.S.	1	1.02	2.64	15.70	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.621	7.056	24.345	0.000	0.000	0.000	0.000	0.000

Problem 437	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	419	426	225	835	0	0	0	0	0
N.S.	1	1.02	0.54	1.99	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.687	0.802	14.515	0.000	0.000	0.000	0.000	0.000

Problem 438	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	131	146	0	0	0	0	0
N.S.	1	1.00	1.12	1.25	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.302	0.088	11.976	0.000	0.000	0.000	0.000	0.000

Problem 439	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	171	112	0	0	0	0	0
N.S.	1	1.00	1.55	1.02	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.305	0.648	12.300	0.000	0.000	0.000	0.000	0.000

Problem 440	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	226	226	212	659	0	0	0	0	0
N.S.	1	1.00	0.94	2.92	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.664	1.410	17.055	0.000	0.000	0.000	0.000	0.000

Problem 441	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	0	475	0	0	0	0	0
N.S.	1	1.00	0.00	6.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.278	0.000	11.352	0.000	0.000	0.000	0.000	0.000

Problem 442	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	0	462	0	0	0	0	0
N.S.	1	1.00	0.00	6.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.274	0.000	9.832	0.000	0.000	0.000	0.000	0.000

Problem 443	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	93	93	0	474	0	0	0	0	0
N.S.	1	1.00	0.00	5.10	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.456	0.000	11.336	0.000	0.000	0.000	0.000	0.000

Problem 444	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	0	316	0	0	0	0	0
N.S.	1	1.00	0.00	3.33	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.445	0.000	12.191	0.000	0.000	0.000	0.000	0.000

Problem 445	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	0	477	0	0	0	0	0
N.S.	1	1.00	0.00	6.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.280	0.000	12.358	0.000	0.000	0.000	0.000	0.000

Problem 446	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	0	485	0	0	0	0	0
N.S.	1	1.00	0.00	6.55	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.295	0.000	11.587	0.000	0.000	0.000	0.000	0.000

Problem 447	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	0	313	0	0	0	0	0
N.S.	1	1.00	0.00	3.19	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.469	0.000	12.997	0.000	0.000	0.000	0.000	0.000

Problem 448	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	B	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	96	96	0	331	0	0	0	0	0
N.S.	1	1.00	0.00	3.45	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.448	0.000	12.615	0.000	0.000	0.000	0.000	0.000

Problem 449	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	33	33	35	33	35	35	32	35	35
N.S.	1	1.00	1.06	1.00	1.06	1.06	0.97	1.06	1.06
time (sec)	N/A	0.260	16.143	0.866	3.954	0.302	123.044	1.010	2.727

Problem 450	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	595	614	319	0	0	0	0	0	0
N.S.	1	1.03	0.54	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.942	4.729	0.000	0.000	0.000	0.000	0.000	0.000

Problem 451	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	406	408	263	0	0	0	0	0	0
N.S.	1	1.00	0.65	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.762	1.794	0.000	0.000	0.000	0.000	0.000	0.000

Problem 452	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	287	291	211	0	0	0	0	0	0
N.S.	1	1.01	0.74	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.069	1.084	0.000	0.000	0.000	0.000	0.000	0.000

Problem 453	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	196	193	162	0	0	0	0	0	0
N.S.	1	0.98	0.83	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.648	0.541	0.000	0.000	0.000	0.000	0.000	0.000

Problem 454	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	286	281	10482	0	0	0	0	0	0
N.S.	1	0.98	36.65	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.876	30.990	0.000	0.000	0.000	0.000	0.000	0.000

Problem 455	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	54	34	35	35
N.S.	1	1.00	1.06	0.94	1.00	1.54	0.97	1.00	1.00
time (sec)	N/A	0.840	79.327	0.786	2.595	0.300	147.502	16.257	4.403

Problem 456	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	35	34	35	35
N.S.	1	1.00	1.06	0.94	1.00	1.00	0.97	1.00	1.00
time (sec)	N/A	0.300	28.846	0.798	2.591	0.272	7.322	0.956	2.716

Problem 457	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	35	34	35	35
N.S.	1	1.00	1.06	0.94	1.00	1.00	0.97	1.00	1.00
time (sec)	N/A	0.298	34.238	0.923	2.418	0.276	3.757	0.736	4.305

Problem 458	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	63	34	35	35
N.S.	1	1.00	1.06	0.94	1.00	1.80	0.97	1.00	1.00
time (sec)	N/A	0.846	32.201	0.896	2.590	0.283	9.252	1.288	6.389

Problem 459	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	172	166	292	634	0	219	0	0	0
N.S.	1	0.97	1.70	3.69	0.00	1.27	0.00	0.00	0.00
time (sec)	N/A	0.857	1.726	37.108	0.000	0.099	0.000	0.000	0.000

Problem 460	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	137	225	399	0	188	0	0	0
N.S.	1	1.01	1.67	2.96	0.00	1.39	0.00	0.00	0.00
time (sec)	N/A	0.799	1.103	34.895	0.000	0.098	0.000	0.000	0.000

Problem 461	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	106	157	242	0	141	0	0	0
N.S.	1	1.00	1.48	2.28	0.00	1.33	0.00	0.00	0.00
time (sec)	N/A	0.687	1.394	7.657	0.000	0.093	0.000	0.000	0.000

Problem 462	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	113	148	321	0	142	0	0	0
N.S.	1	1.03	1.35	2.92	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	0.709	1.498	8.608	0.000	0.091	0.000	0.000	0.000

Problem 463	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	143	148	355	0	169	0	0	0
N.S.	1	1.01	1.05	2.52	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	0.809	1.600	10.193	0.000	0.102	0.000	0.000	0.000

Problem 464	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	172	170	182	383	0	187	0	0	0
N.S.	1	0.99	1.06	2.23	0.00	1.09	0.00	0.00	0.00
time (sec)	N/A	0.891	2.161	11.717	0.000	0.102	0.000	0.000	0.000

Problem 465	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	199	196	299	714	0	239	0	0	0
N.S.	1	0.98	1.50	3.59	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	1.128	2.555	63.487	0.000	0.099	0.000	0.000	0.000

Problem 466	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	164	188	513	0	202	0	0	0
N.S.	1	1.02	1.18	3.21	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	1.005	2.299	60.966	0.000	0.100	0.000	0.000	0.000

Problem 467	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	164	152	244	0	166	0	0	0
N.S.	1	1.02	0.95	1.52	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.981	2.329	9.708	0.000	0.097	0.000	0.000	0.000

Problem 468	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	172	153	357	0	187	0	0	0
N.S.	1	1.04	0.92	2.15	0.00	1.13	0.00	0.00	0.00
time (sec)	N/A	1.017	2.573	11.113	0.000	0.097	0.000	0.000	0.000

Problem 469	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	202	193	385	0	211	0	0	0
N.S.	1	1.00	0.96	1.92	0.00	1.05	0.00	0.00	0.00
time (sec)	N/A	1.168	2.769	12.975	0.000	0.102	0.000	0.000	0.000

Problem 470	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	244	246	435	902	0	263	0	0	0
N.S.	1	1.01	1.78	3.70	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	1.518	3.523	179.701	0.000	0.100	0.000	0.000	0.000

Problem 471	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	215	268	916	0	243	0	0	0
N.S.	1	1.02	1.27	4.34	0.00	1.15	0.00	0.00	0.00
time (sec)	N/A	1.333	3.446	181.489	0.000	0.106	0.000	0.000	0.000

Problem 472	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	199	201	202	654	0	215	0	0	0
N.S.	1	1.01	1.02	3.29	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	1.301	2.573	187.012	0.000	0.097	0.000	0.000	0.000

Problem 473	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	215	207	337	0	197	0	0	0
N.S.	1	1.02	0.98	1.60	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	1.346	2.136	11.495	0.000	0.095	0.000	0.000	0.000

Problem 474	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	222	194	385	0	211	0	0	0
N.S.	1	1.05	0.92	1.82	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	1.367	3.222	13.997	0.000	0.096	0.000	0.000	0.000

Problem 475	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	244	250	196	413	0	231	0	0	0
N.S.	1	1.02	0.80	1.69	0.00	0.95	0.00	0.00	0.00
time (sec)	N/A	1.593	3.131	16.787	0.000	0.103	0.000	0.000	0.000

Problem 476	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	193	185	409	466	0	308	0	0	0
N.S.	1	0.96	2.12	2.41	0.00	1.60	0.00	0.00	0.00
time (sec)	N/A	0.981	6.693	8.747	0.000	0.102	0.000	0.000	0.000

Problem 477	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	157	400	319	0	250	0	0	0
N.S.	1	0.99	2.52	2.01	0.00	1.57	0.00	0.00	0.00
time (sec)	N/A	0.912	4.202	4.785	0.000	0.098	0.000	0.000	0.000

Problem 478	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	129	200	243	0	241	0	0	0
N.S.	1	1.05	1.63	1.98	0.00	1.96	0.00	0.00	0.00
time (sec)	N/A	0.763	2.072	3.641	0.000	0.091	0.000	0.000	0.000

Problem 479	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	130	422	244	0	237	0	0	0
N.S.	1	1.04	3.38	1.95	0.00	1.90	0.00	0.00	0.00
time (sec)	N/A	0.765	2.941	4.711	0.000	0.094	0.000	0.000	0.000

Problem 480	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	159	444	262	0	261	0	0	0
N.S.	1	0.98	2.72	1.61	0.00	1.60	0.00	0.00	0.00
time (sec)	N/A	0.906	4.739	5.448	0.000	0.099	0.000	0.000	0.000

Problem 481	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	187	518	281	0	278	0	0	0
N.S.	1	0.95	2.64	1.43	0.00	1.42	0.00	0.00	0.00
time (sec)	N/A	0.948	3.631	5.897	0.000	0.107	0.000	0.000	0.000

Problem 482	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	208	208	303	494	0	367	0	0	0
N.S.	1	1.00	1.46	2.38	0.00	1.76	0.00	0.00	0.00
time (sec)	N/A	1.247	2.801	5.185	0.000	0.099	0.000	0.000	0.000

Problem 483	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	161	171	256	350	0	326	0	0	0
N.S.	1	1.06	1.59	2.17	0.00	2.02	0.00	0.00	0.00
time (sec)	N/A	1.080	2.164	4.714	0.000	0.096	0.000	0.000	0.000

Problem 484	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	175	256	350	0	324	0	0	0
N.S.	1	1.04	1.52	2.08	0.00	1.93	0.00	0.00	0.00
time (sec)	N/A	1.091	2.334	5.218	0.000	0.096	0.000	0.000	0.000

Problem 485	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	176	182	475	421	0	362	0	0	0
N.S.	1	1.03	2.70	2.39	0.00	2.06	0.00	0.00	0.00
time (sec)	N/A	1.107	5.476	5.667	0.000	0.102	0.000	0.000	0.000

Problem 486	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	206	211	525	435	0	376	0	0	0
N.S.	1	1.02	2.55	2.11	0.00	1.83	0.00	0.00	0.00
time (sec)	N/A	1.291	6.761	6.122	0.000	0.106	0.000	0.000	0.000

Problem 487	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	261	267	358	685	0	481	0	0	0
N.S.	1	1.02	1.37	2.62	0.00	1.84	0.00	0.00	0.00
time (sec)	N/A	1.731	4.353	6.000	0.000	0.107	0.000	0.000	0.000

Problem 488	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	222	237	793	451	0	476	0	0	0
N.S.	1	1.07	3.57	2.03	0.00	2.14	0.00	0.00	0.00
time (sec)	N/A	1.520	7.606	5.521	0.000	0.101	0.000	0.000	0.000

Problem 489	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	231	792	451	0	472	0	0	0
N.S.	1	1.07	3.67	2.09	0.00	2.19	0.00	0.00	0.00
time (sec)	N/A	1.490	7.497	5.766	0.000	0.103	0.000	0.000	0.000

Problem 490	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	222	237	793	451	0	474	0	0	0
N.S.	1	1.07	3.57	2.03	0.00	2.14	0.00	0.00	0.00
time (sec)	N/A	1.471	7.979	6.267	0.000	0.105	0.000	0.000	0.000

Problem 491	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	243	817	451	0	478	0	0	0
N.S.	1	1.07	3.58	1.98	0.00	2.10	0.00	0.00	0.00
time (sec)	N/A	1.522	8.447	6.152	0.000	0.117	0.000	0.000	0.000

Problem 492	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	259	269	589	465	0	489	0	0	0
N.S.	1	1.04	2.27	1.80	0.00	1.89	0.00	0.00	0.00
time (sec)	N/A	1.683	5.182	6.607	0.000	0.123	0.000	0.000	0.000

Problem 493	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	220	235	124	121	659	121	0	0	479
N.S.	1	1.07	0.56	0.55	3.00	0.55	0.00	0.00	2.18
time (sec)	N/A	1.168	0.373	9.799	0.368	0.257	0.000	0.000	6.195

Problem 494	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	192	102	101	568	104	0	0	441
N.S.	1	1.10	0.58	0.58	3.25	0.59	0.00	0.00	2.52
time (sec)	N/A	0.939	0.299	9.616	0.374	0.255	0.000	0.000	4.824

Problem 495	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	149	78	84	475	86	0	0	196
N.S.	1	1.15	0.60	0.65	3.65	0.66	0.00	0.00	1.51
time (sec)	N/A	0.754	0.193	9.730	0.352	0.277	0.000	0.000	2.824

Problem 496	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	106	57	60	380	65	0	0	114
N.S.	1	1.25	0.67	0.71	4.47	0.76	0.00	0.00	1.34
time (sec)	N/A	0.608	0.116	9.581	0.352	0.280	0.000	0.000	1.165

Problem 497	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	96	97	86	120	906	91	0	0	0
N.S.	1	1.01	0.90	1.25	9.44	0.95	0.00	0.00	0.00
time (sec)	N/A	0.607	0.124	9.934	0.470	0.300	0.000	0.000	0.000

Problem 498	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	99	103	147	939	97	0	0	0
N.S.	1	1.01	1.05	1.50	9.58	0.99	0.00	0.00	0.00
time (sec)	N/A	0.585	0.148	22.685	0.485	0.298	0.000	0.000	0.000

Problem 499	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	144	120	206	1851	127	0	0	0
N.S.	1	0.95	0.79	1.36	12.26	0.84	0.00	0.00	0.00
time (sec)	N/A	0.740	0.269	20.259	0.546	0.304	0.000	0.000	0.000

Problem 500	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	187	138	274	2981	146	0	0	0
N.S.	1	0.95	0.70	1.40	15.21	0.74	0.00	0.00	0.00
time (sec)	N/A	0.973	0.467	20.121	0.677	0.307	0.000	0.000	0.000

Problem 501	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	275	288	146	142	712	144	0	0	348
N.S.	1	1.05	0.53	0.52	2.59	0.52	0.00	0.00	1.27
time (sec)	N/A	1.512	0.469	10.151	0.349	0.257	0.000	0.000	5.298

Problem 502	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	245	124	122	619	126	0	0	316
N.S.	1	1.07	0.54	0.54	2.71	0.55	0.00	0.00	1.39
time (sec)	N/A	1.298	0.463	10.160	0.371	0.290	0.000	0.000	5.302

Problem 503	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	202	102	102	527	107	0	0	259
N.S.	1	1.12	0.56	0.56	2.91	0.59	0.00	0.00	1.43
time (sec)	N/A	1.083	0.368	10.201	0.370	0.289	0.000	0.000	5.152

Problem 504	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	160	80	85	436	88	0	0	197
N.S.	1	1.19	0.60	0.63	3.25	0.66	0.00	0.00	1.47
time (sec)	N/A	0.913	0.242	10.145	0.364	0.278	0.000	0.000	2.651

Problem 505	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	149	106	200	1462	130	0	0	0
N.S.	1	1.03	0.73	1.38	10.08	0.90	0.00	0.00	0.00
time (sec)	N/A	0.884	0.275	10.323	0.498	0.306	0.000	0.000	0.000

Problem 506	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	147	107	274	1801	119	0	0	0
N.S.	1	1.01	0.73	1.88	12.34	0.82	0.00	0.00	0.00
time (sec)	N/A	0.908	0.222	22.793	0.542	0.330	0.000	0.000	0.000

Problem 507	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	153	121	213	1884	133	0	0	0
N.S.	1	1.00	0.79	1.39	12.31	0.87	0.00	0.00	0.00
time (sec)	N/A	0.902	0.300	22.879	0.570	0.326	0.000	0.000	0.000

Problem 508	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	197	139	277	3023	153	0	0	0
N.S.	1	0.98	0.70	1.38	15.12	0.76	0.00	0.00	0.00
time (sec)	N/A	1.109	0.555	20.701	0.707	0.341	0.000	0.000	0.000

Problem 509	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	247	240	156	345	8901	171	0	0	0
N.S.	1	0.97	0.63	1.40	36.04	0.69	0.00	0.00	0.00
time (sec)	N/A	1.308	0.935	20.376	0.978	0.375	0.000	0.000	0.000

Problem 510	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	322	342	171	164	763	176	0	0	789
N.S.	1	1.06	0.53	0.51	2.37	0.55	0.00	0.00	2.45
time (sec)	N/A	1.945	0.867	2.205	0.369	0.300	0.000	0.000	6.741

Problem 511	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	275	299	147	144	672	156	0	0	751
N.S.	1	1.09	0.53	0.52	2.44	0.57	0.00	0.00	2.73
time (sec)	N/A	1.692	0.782	2.208	0.371	0.283	0.000	0.000	5.441

Problem 512	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	256	126	124	579	135	0	0	617
N.S.	1	1.12	0.55	0.54	2.54	0.59	0.00	0.00	2.71
time (sec)	N/A	1.424	0.557	2.680	0.384	0.296	0.000	0.000	5.597

Problem 513	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	212	104	104	488	114	0	0	579
N.S.	1	1.17	0.57	0.57	2.70	0.63	0.00	0.00	3.20
time (sec)	N/A	1.246	0.491	2.086	0.358	0.303	0.000	0.000	5.254

Problem 514	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	201	130	237	1713	162	0	0	0
N.S.	1	1.05	0.68	1.23	8.92	0.84	0.00	0.00	0.00
time (sec)	N/A	1.177	0.572	2.440	0.529	0.305	0.000	0.000	0.000

Problem 515	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	193	198	130	334	2780	166	0	0	0
N.S.	1	1.03	0.67	1.73	14.40	0.86	0.00	0.00	0.00
time (sec)	N/A	1.218	0.480	2.511	0.617	0.353	0.000	0.000	0.000

Problem 516	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	199	126	310	0	147	0	0	0
N.S.	1	1.01	0.64	1.57	0.00	0.74	0.00	0.00	0.00
time (sec)	N/A	1.215	0.414	22.910	0.000	0.337	0.000	0.000	0.000

Problem 517	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	206	141	285	3071	163	0	0	0
N.S.	1	1.03	0.70	1.42	15.36	0.82	0.00	0.00	0.00
time (sec)	N/A	1.220	0.670	22.477	0.696	0.334	0.000	0.000	0.000

Problem 518	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	247	251	157	351	9390	183	0	0	0
N.S.	1	1.02	0.64	1.42	38.02	0.74	0.00	0.00	0.00
time (sec)	N/A	1.447	1.131	21.158	0.957	0.364	0.000	0.000	0.000

Problem 519	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	294	294	179	419	10042	203	0	0	0
N.S.	1	1.00	0.61	1.43	34.16	0.69	0.00	0.00	0.00
time (sec)	N/A	1.705	1.889	20.834	0.946	0.380	0.000	0.000	0.000

Problem 520	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	295	336	272	405	0	198	0	0	0
N.S.	1	1.14	0.92	1.37	0.00	0.67	0.00	0.00	0.00
time (sec)	N/A	1.971	8.750	9.421	0.000	0.328	0.000	0.000	0.000

Problem 521	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	250	281	2442	365	0	181	0	0	0
N.S.	1	1.12	9.77	1.46	0.00	0.72	0.00	0.00	0.00
time (sec)	N/A	1.559	10.904	9.897	0.000	0.323	0.000	0.000	0.000

Problem 522	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	207	226	1719	325	0	164	0	0	0
N.S.	1	1.09	8.30	1.57	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	1.187	7.210	9.838	0.000	0.279	0.000	0.000	0.000

Problem 523	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	162	171	617	284	0	143	0	0	0
N.S.	1	1.06	3.81	1.75	0.00	0.88	0.00	0.00	0.00
time (sec)	N/A	0.883	6.592	9.943	0.000	0.295	0.000	0.000	0.000

Problem 524	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	119	120	203	211	0	110	0	0	0
N.S.	1	1.01	1.71	1.77	0.00	0.92	0.00	0.00	0.00
time (sec)	N/A	0.615	1.059	10.183	0.000	0.295	0.000	0.000	0.000

Problem 525	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	121	102	138	1221	96	0	0	0
N.S.	1	0.86	0.73	0.99	8.72	0.69	0.00	0.00	0.00
time (sec)	N/A	0.744	0.157	10.306	1.078	1.045	0.000	0.000	0.000

Problem 526	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	171	467	196	0	168	0	0	0
N.S.	1	0.94	2.58	1.08	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	0.997	1.562	20.338	0.000	1.252	0.000	0.000	0.000

Problem 527	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	230	224	412	264	0	194	0	0	0
N.S.	1	0.97	1.79	1.15	0.00	0.84	0.00	0.00	0.00
time (sec)	N/A	1.355	1.652	19.926	0.000	2.256	0.000	0.000	0.000

Problem 528	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	183	143	281	0	208	0	0	0
N.S.	1	0.95	0.74	1.46	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	1.186	0.320	24.582	0.000	13.774	0.000	0.000	0.000

Problem 529	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	317	344	2966	497	0	237	0	0	0
N.S.	1	1.09	9.36	1.57	0.00	0.75	0.00	0.00	0.00
time (sec)	N/A	2.029	8.377	9.577	0.000	0.293	0.000	0.000	0.000

Problem 530	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	270	289	2166	457	0	220	0	0	0
N.S.	1	1.07	8.02	1.69	0.00	0.81	0.00	0.00	0.00
time (sec)	N/A	1.629	7.226	10.385	0.000	0.288	0.000	0.000	0.000

Problem 531	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	223	234	981	417	0	197	0	0	0
N.S.	1	1.05	4.40	1.87	0.00	0.88	0.00	0.00	0.00
time (sec)	N/A	1.255	6.588	10.332	0.000	0.279	0.000	0.000	0.000

Problem 532	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	176	179	443	353	0	163	0	0	0
N.S.	1	1.02	2.52	2.01	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	0.925	2.901	10.227	0.000	0.280	0.000	0.000	0.000

Problem 533	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	128	191	215	0	144	0	0	0
N.S.	1	1.01	1.50	1.69	0.00	1.13	0.00	0.00	0.00
time (sec)	N/A	0.640	1.319	10.543	0.000	0.274	0.000	0.000	0.000

Problem 534	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	171	243	283	0	203	0	0	0
N.S.	1	0.92	1.31	1.53	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	1.067	2.158	7.445	0.000	2.892	0.000	0.000	0.000

Problem 535	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	226	836	387	0	246	0	0	0
N.S.	1	0.95	3.53	1.63	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	1.421	7.444	20.863	0.000	3.748	0.000	0.000	0.000

Problem 536	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	317	344	261	589	0	266	0	0	0
N.S.	1	1.09	0.82	1.86	0.00	0.84	0.00	0.00	0.00
time (sec)	N/A	1.999	10.236	10.839	0.000	0.297	0.000	0.000	0.000

Problem 537	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	270	289	1152	549	0	246	0	0	0
N.S.	1	1.07	4.27	2.03	0.00	0.91	0.00	0.00	0.00
time (sec)	N/A	1.638	8.070	11.249	0.000	0.283	0.000	0.000	0.000

Problem 538	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	223	234	732	485	0	210	0	0	0
N.S.	1	1.05	3.28	2.17	0.00	0.94	0.00	0.00	0.00
time (sec)	N/A	1.290	6.776	11.084	0.000	0.279	0.000	0.000	0.000

Problem 539	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	176	183	162	367	0	207	0	0	0
N.S.	1	1.04	0.92	2.09	0.00	1.18	0.00	0.00	0.00
time (sec)	N/A	0.954	1.888	11.460	0.000	0.285	0.000	0.000	0.000

Problem 540	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	174	181	182	340	0	205	0	0	0
N.S.	1	1.04	1.05	1.95	0.00	1.18	0.00	0.00	0.00
time (sec)	N/A	0.951	3.676	8.016	0.000	0.273	0.000	0.000	0.000

Problem 541	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	234	226	264	469	0	277	0	0	0
N.S.	1	0.97	1.13	2.00	0.00	1.18	0.00	0.00	0.00
time (sec)	N/A	1.407	2.855	8.392	0.000	6.027	0.000	0.000	0.000

Problem 542	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	286	283	929	597	0	313	0	0	0
N.S.	1	0.99	3.25	2.09	0.00	1.09	0.00	0.00	0.00
time (sec)	N/A	1.825	7.899	20.986	0.000	7.925	0.000	0.000	0.000

Problem 543	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	317	344	1260	681	0	295	0	0	0
N.S.	1	1.09	3.97	2.15	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	2.026	8.876	10.717	0.000	0.322	0.000	0.000	0.000

Problem 544	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	A	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	270	289	802	643	0	260	0	0	0
N.S.	1	1.07	2.97	2.38	0.00	0.96	0.00	0.00	0.00
time (sec)	N/A	1.664	7.016	10.804	0.000	0.301	0.000	0.000	0.000

Problem 545	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	223	238	180	465	0	257	0	0	0
N.S.	1	1.07	0.81	2.09	0.00	1.15	0.00	0.00	0.00
time (sec)	N/A	1.297	1.856	11.718	0.000	0.296	0.000	0.000	0.000

Problem 546	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	236	179	488	0	255	0	0	0
N.S.	1	1.07	0.81	2.21	0.00	1.15	0.00	0.00	0.00
time (sec)	N/A	1.288	1.804	8.293	0.000	0.285	0.000	0.000	0.000

Problem 547	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	236	561	490	0	257	0	0	0
N.S.	1	1.07	2.54	2.22	0.00	1.16	0.00	0.00	0.00
time (sec)	N/A	1.337	6.426	8.270	0.000	0.292	0.000	0.000	0.000

Problem 548	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	281	281	281	567	0	338	0	0	0
N.S.	1	1.00	1.00	2.02	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	1.773	4.071	8.177	0.000	7.949	0.000	0.000	0.000

Problem 549	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	333	338	1017	811	0	379	0	0	0
N.S.	1	1.02	3.05	2.44	0.00	1.14	0.00	0.00	0.00
time (sec)	N/A	2.238	8.347	20.985	0.000	13.730	0.000	0.000	0.000

Problem 550	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	170	132	636	0	235	0	0	0
N.S.	1	0.94	0.73	3.53	0.00	1.31	0.00	0.00	0.00
time (sec)	N/A	0.849	1.746	102.974	0.000	0.099	0.000	0.000	0.000

Problem 551	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	143	142	104	401	0	205	0	0	0
N.S.	1	0.99	0.73	2.80	0.00	1.43	0.00	0.00	0.00
time (sec)	N/A	0.800	0.972	100.541	0.000	0.097	0.000	0.000	0.000

Problem 552	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	111	85	246	0	153	0	0	0
N.S.	1	1.00	0.77	2.22	0.00	1.38	0.00	0.00	0.00
time (sec)	N/A	0.712	1.382	9.325	0.000	0.114	0.000	0.000	0.000

Problem 553	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	118	90	326	0	156	0	0	0
N.S.	1	1.03	0.78	2.83	0.00	1.36	0.00	0.00	0.00
time (sec)	N/A	0.687	0.811	9.842	0.000	0.101	0.000	0.000	0.000

Problem 554	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	148	147	108	371	0	184	0	0	0
N.S.	1	0.99	0.73	2.51	0.00	1.24	0.00	0.00	0.00
time (sec)	N/A	0.834	1.000	10.470	0.000	0.104	0.000	0.000	0.000

Problem 555	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	174	125	413	0	203	0	0	0
N.S.	1	0.97	0.69	2.29	0.00	1.13	0.00	0.00	0.00
time (sec)	N/A	0.912	1.391	12.427	0.000	0.106	0.000	0.000	0.000

Problem 556	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	206	171	723	0	286	0	0	0
N.S.	1	0.93	0.77	3.27	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	1.409	2.937	490.579	0.000	0.102	0.000	0.000	0.000

Problem 557	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	177	178	125	650	0	247	0	0	0
N.S.	1	1.01	0.71	3.67	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	1.198	4.398	495.723	0.000	0.102	0.000	0.000	0.000

Problem 558	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	161	164	124	405	0	208	0	0	0
N.S.	1	1.02	0.77	2.52	0.00	1.29	0.00	0.00	0.00
time (sec)	N/A	1.135	4.130	11.533	0.000	0.099	0.000	0.000	0.000

Problem 559	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	174	128	487	0	226	0	0	0
N.S.	1	1.02	0.75	2.85	0.00	1.32	0.00	0.00	0.00
time (sec)	N/A	1.137	4.194	12.227	0.000	0.111	0.000	0.000	0.000

Problem 560	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	202	161	548	0	254	0	0	0
N.S.	1	0.95	0.76	2.57	0.00	1.19	0.00	0.00	0.00
time (sec)	N/A	1.409	4.471	15.016	0.000	0.107	0.000	0.000	0.000

Problem 561	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	295	277	225	917	0	364	0	0	0
N.S.	1	0.94	0.76	3.11	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	1.937	3.823	1876.703	0.000	0.119	0.000	0.000	0.000

Problem 562	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	244	248	192	970	0	326	0	0	0
N.S.	1	1.02	0.79	3.98	0.00	1.34	0.00	0.00	0.00
time (sec)	N/A	1.751	10.302	1955.723	0.000	0.126	0.000	0.000	0.000

Problem 563	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	239	238	166	886	0	298	0	0	0
N.S.	1	1.00	0.69	3.71	0.00	1.25	0.00	0.00	0.00
time (sec)	N/A	1.657	7.282	2002.346	0.000	0.116	0.000	0.000	0.000

Problem 564	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	241	172	641	0	270	0	0	0
N.S.	1	1.02	0.73	2.70	0.00	1.14	0.00	0.00	0.00
time (sec)	N/A	1.672	6.702	15.570	0.000	0.114	0.000	0.000	0.000

Problem 565	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	245	253	180	664	0	295	0	0	0
N.S.	1	1.03	0.73	2.71	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	1.670	7.095	16.213	0.000	0.121	0.000	0.000	0.000

Problem 566	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	295	281	219	745	0	332	0	0	0
N.S.	1	0.95	0.74	2.53	0.00	1.13	0.00	0.00	0.00
time (sec)	N/A	1.928	7.445	18.822	0.000	0.129	0.000	0.000	0.000

Problem 567	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	210	224	225	441	0	0	0	0	0
N.S.	1	1.07	1.07	2.10	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.970	9.764	20.235	0.000	0.000	0.000	0.000	0.000

Problem 568	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	125	300	0	0	0	0	0
N.S.	1	1.00	0.99	2.38	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.108	2.023	5.453	0.000	0.000	0.000	0.000	0.000

Problem 569	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	101	76	217	0	0	0	0	0
N.S.	1	1.00	0.75	2.15	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.803	0.916	3.765	0.000	0.000	0.000	0.000	0.000

Problem 570	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	149	126	220	295	0	0	0	0	0
N.S.	1	0.85	1.48	1.98	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.001	19.911	5.425	0.000	0.000	0.000	0.000	0.000

Problem 571	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	203	278	822	0	0	0	0	0
N.S.	1	1.03	1.41	4.17	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.518	5.013	6.477	0.000	0.000	0.000	0.000	0.000

Problem 572	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	405	389	735	1004	0	0	0	0	0
N.S.	1	0.96	1.81	2.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.150	6.982	44.153	0.000	0.000	0.000	0.000	0.000

Problem 573	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	316	306	681	856	0	0	0	0	0
N.S.	1	0.97	2.16	2.71	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.399	6.872	8.246	0.000	0.000	0.000	0.000	0.000

Problem 574	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	260	252	443	721	0	0	0	0	0
N.S.	1	0.97	1.70	2.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.829	6.010	7.460	0.000	0.000	0.000	0.000	0.000

Problem 575	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	258	247	441	808	0	0	0	0	0
N.S.	1	0.96	1.71	3.13	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.731	5.286	7.943	0.000	0.000	0.000	0.000	0.000

Problem 576	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	284	275	461	849	0	0	0	0	0
N.S.	1	0.97	1.62	2.99	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.911	6.522	9.533	0.000	0.000	0.000	0.000	0.000

Problem 577	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	363	350	701	1066	0	0	0	0	0
N.S.	1	0.96	1.93	2.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.578	6.841	11.300	0.000	0.000	0.000	0.000	0.000

Problem 578	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	480	476	844	1975	0	0	0	0	0
N.S.	1	0.99	1.76	4.11	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.672	7.136	15.520	0.000	0.000	0.000	0.000	0.000

Problem 579	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	405	403	797	1744	0	0	0	0	0
N.S.	1	1.00	1.97	4.31	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.831	6.975	12.596	0.000	0.000	0.000	0.000	0.000

Problem 580	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	402	397	784	1850	0	0	0	0	0
N.S.	1	0.99	1.95	4.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.743	6.954	13.375	0.000	0.000	0.000	0.000	0.000

Problem 581	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	400	397	551	1937	0	0	0	0	0
N.S.	1	0.99	1.38	4.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.705	6.010	13.431	0.000	0.000	0.000	0.000	0.000

Problem 582	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	427	428	820	1977	0	0	0	0	0
N.S.	1	1.00	1.92	4.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.027	7.143	15.438	0.000	0.000	0.000	0.000	0.000

Problem 583	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	521	519	865	2195	0	0	0	0	0
N.S.	1	1.00	1.66	4.21	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.860	7.352	17.786	0.000	0.000	0.000	0.000	0.000

Problem 584	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	64	47	214	0	91	0	0	0
N.S.	1	1.00	0.73	3.34	0.00	1.42	0.00	0.00	0.00
time (sec)	N/A	0.339	0.065	3.661	0.000	0.119	0.000	0.000	0.000

Problem 585	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	46	183	0	76	0	0	0
N.S.	1	1.00	0.77	3.05	0.00	1.27	0.00	0.00	0.00
time (sec)	N/A	0.343	0.041	2.946	0.000	0.118	0.000	0.000	0.000

Problem 586	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	37	134	0	53	0	0	0
N.S.	1	1.00	1.00	3.62	0.00	1.43	0.00	0.00	0.00
time (sec)	N/A	0.260	0.029	2.074	0.000	0.097	0.000	0.000	0.000

Problem 587	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	37	134	0	59	0	0	0
N.S.	1	1.00	1.00	3.62	0.00	1.59	0.00	0.00	0.00
time (sec)	N/A	0.272	0.031	3.259	0.000	0.100	0.000	0.000	0.000

Problem 588	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	64	50	180	0	71	0	0	0
N.S.	1	1.00	0.78	2.81	0.00	1.11	0.00	0.00	0.00
time (sec)	N/A	0.349	0.042	4.025	0.000	0.102	0.000	0.000	0.000

Problem 589	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	64	56	203	0	77	0	0	0
N.S.	1	1.00	0.88	3.17	0.00	1.20	0.00	0.00	0.00
time (sec)	N/A	0.352	0.057	5.029	0.000	0.106	0.000	0.000	0.000

Problem 590	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	473	471	519	4628	0	0	0	0	0
N.S.	1	1.00	1.10	9.78	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.339	15.833	29.527	0.000	0.000	0.000	0.000	0.000

Problem 591	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	390	380	373	3320	0	0	0	0	0
N.S.	1	0.97	0.96	8.51	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.606	11.880	25.135	0.000	0.000	0.000	0.000	0.000

Problem 592	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	324	306	318	1975	0	0	0	0	0
N.S.	1	0.94	0.98	6.10	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.174	11.004	18.414	0.000	0.000	0.000	0.000	0.000

Problem 593	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	411	372	358	960	0	0	0	0	0
N.S.	1	0.91	0.87	2.34	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.378	7.173	18.247	0.000	0.000	0.000	0.000	0.000

Problem 594	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	445	412	787	1268	0	0	0	0	0
N.S.	1	0.93	1.77	2.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.728	16.630	15.048	0.000	0.000	0.000	0.000	0.000

Problem 595	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	533	492	1121	2728	0	0	0	0	0
N.S.	1	0.92	2.10	5.12	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.276	16.836	14.563	0.000	0.000	0.000	0.000	0.000

Problem 596	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	620	584	1533	3922	0	0	0	0	0
N.S.	1	0.94	2.47	6.33	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.911	13.942	17.608	0.000	0.000	0.000	0.000	0.000

Problem 597	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	562	554	625	5977	0	0	0	0	0
N.S.	1	0.99	1.11	10.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.862	18.972	32.824	0.000	0.000	0.000	0.000	0.000

Problem 598	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	473	465	518	4627	0	0	0	0	0
N.S.	1	0.98	1.10	9.78	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.252	16.284	28.852	0.000	0.000	0.000	0.000	0.000

Problem 599	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	393	380	379	3582	0	0	0	0	0
N.S.	1	0.97	0.96	9.11	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.710	12.706	24.059	0.000	0.000	0.000	0.000	0.000

Problem 600	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	479	441	5981	2489	0	0	0	0	0
N.S.	1	0.92	12.49	5.20	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.900	22.339	17.873	0.000	0.000	0.000	0.000	0.000

Problem 601	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	509	482	927	2386	0	0	0	0	0
N.S.	1	0.95	1.82	4.69	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.377	14.770	19.430	0.000	0.000	0.000	0.000	0.000

Problem 602	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	532	496	1134	3310	0	0	0	0	0
N.S.	1	0.93	2.13	6.22	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.407	15.120	18.057	0.000	0.000	0.000	0.000	0.000

Problem 603	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	626	589	1489	4189	0	0	0	0	0
N.S.	1	0.94	2.38	6.69	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.053	15.363	17.608	0.000	0.000	0.000	0.000	0.000

Problem 604	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	730	696	1888	5425	0	0	0	0	0
N.S.	1	0.95	2.59	7.43	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.663	19.454	21.102	0.000	0.000	0.000	0.000	0.000

Problem 605	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F(-1)	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	662	659	4198	0	0	0	0	0	0
N.S.	1	1.00	6.34	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.674	24.740	180.000	0.000	0.000	0.000	0.000	0.000

Problem 606	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F(-1)	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	562	554	629	0	0	0	0	0	0
N.S.	1	0.99	1.12	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.869	19.362	180.000	0.000	0.000	0.000	0.000	0.000

Problem 607	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F(-1)	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	474	466	524	0	0	0	0	0	0
N.S.	1	0.98	1.11	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.303	17.200	180.000	0.000	0.000	0.000	0.000	0.000

Problem 608	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F(-1)	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	553	520	7032	0	0	0	0	0	0
N.S.	1	0.94	12.72	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.476	24.853	180.000	0.000	0.000	0.000	0.000	0.000

Problem 609	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F(-1)	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	596	568	7700	0	0	0	0	0	0
N.S.	1	0.95	12.92	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.079	23.262	180.000	0.000	0.000	0.000	0.000	0.000

Problem 610	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	607	576	1278	4459	0	0	0	0	0
N.S.	1	0.95	2.11	7.35	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.048	16.118	21.309	0.000	0.000	0.000	0.000	0.000

Problem 611	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	624	593	1504	4802	0	0	0	0	0
N.S.	1	0.95	2.41	7.70	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.085	16.972	20.147	0.000	0.000	0.000	0.000	0.000

Problem 612	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	724	693	1857	5695	0	0	0	0	0
N.S.	1	0.96	2.56	7.87	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.765	16.903	21.539	0.000	0.000	0.000	0.000	0.000

Problem 613	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	839	807	703	6970	0	0	0	0	0
N.S.	1	0.96	0.84	8.31	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.584	13.174	23.115	0.000	0.000	0.000	0.000	0.000

Problem 614	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	403	396	385	3320	0	0	0	0	0
N.S.	1	0.98	0.96	8.24	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.673	13.679	25.691	0.000	0.000	0.000	0.000	0.000

Problem 615	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	330	315	324	1864	0	0	0	0	0
N.S.	1	0.95	0.98	5.65	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.175	11.153	21.358	0.000	0.000	0.000	0.000	0.000

Problem 616	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	251	279	778	0	0	0	0	0
N.S.	1	0.93	1.03	2.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.871	9.699	18.685	0.000	0.000	0.000	0.000	0.000

Problem 617	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	268	249	157	181	0	0	0	0	0
N.S.	1	0.93	0.59	0.68	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.795	3.021	16.179	0.000	0.000	0.000	0.000	0.000

Problem 618	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	487	457	610	958	0	0	0	0	0
N.S.	1	0.94	1.25	1.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.962	9.363	16.820	0.000	0.000	0.000	0.000	0.000

Problem 619	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	539	501	1157	2480	0	0	0	0	0
N.S.	1	0.93	2.15	4.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.254	15.453	17.493	0.000	0.000	0.000	0.000	0.000

Problem 620	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	433	433	498	3850	0	0	0	0	0
N.S.	1	1.00	1.15	8.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.930	16.322	25.286	0.000	0.000	0.000	0.000	0.000

Problem 621	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	345	352	375	1811	0	0	0	0	0
N.S.	1	1.02	1.09	5.25	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.378	13.777	22.102	0.000	0.000	0.000	0.000	0.000

Problem 622	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	324	328	305	1316	0	0	0	0	0
N.S.	1	1.01	0.94	4.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.236	9.266	18.912	0.000	0.000	0.000	0.000	0.000

Problem 623	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	476	441	1050	1602	0	0	0	0	0
N.S.	1	0.93	2.21	3.37	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.745	12.798	17.634	0.000	0.000	0.000	0.000	0.000

Problem 624	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	560	548	1551	3270	0	0	0	0	0
N.S.	1	0.98	2.77	5.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.604	15.813	18.009	0.000	0.000	0.000	0.000	0.000

Problem 625	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	607	602	4316	9528	0	0	0	0	0
N.S.	1	0.99	7.11	15.70	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.896	24.300	25.711	0.000	0.000	0.000	0.000	0.000

Problem 626	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	496	498	610	6408	0	0	0	0	0
N.S.	1	1.00	1.23	12.92	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.165	18.731	22.023	0.000	0.000	0.000	0.000	0.000

Problem 627	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	469	465	557	5287	0	0	0	0	0
N.S.	1	0.99	1.19	11.27	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.960	17.147	18.875	0.000	0.000	0.000	0.000	0.000

Problem 628	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	431	431	528	4567	0	0	0	0	0
N.S.	1	1.00	1.23	10.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.792	14.532	17.006	0.000	0.000	0.000	0.000	0.000

Problem 629	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	602	590	1496	6418	0	0	0	0	0
N.S.	1	0.98	2.49	10.66	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.707	14.197	19.327	0.000	0.000	0.000	0.000	0.000

Problem 630	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	F(-1)	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	733	719	2318	10353	0	0	0	0	0
N.S.	1	0.98	3.16	14.12	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.653	20.137	21.290	0.000	0.000	0.000	0.000	0.000

Problem 631	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	266	246	298	659	0	0	0	0	0
N.S.	1	0.92	1.12	2.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.807	6.055	20.427	0.000	0.000	0.000	0.000	0.000

Problem 632	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	130	104	112	0	0	0	0	0
N.S.	1	1.00	0.80	0.86	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.420	0.076	15.870	0.000	0.000	0.000	0.000	0.000

Problem 633	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	137	147	138	0	0	0	0	0
N.S.	1	1.00	1.07	1.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.429	0.099	15.263	0.000	0.000	0.000	0.000	0.000

Problem 634	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	479	446	236	820	0	0	0	0	0
N.S.	1	0.93	0.49	1.71	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.858	0.976	16.896	0.000	0.000	0.000	0.000	0.000

Problem 635	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	33	33	35	33	35	35	32	35	37
N.S.	1	1.00	1.06	1.00	1.06	1.06	0.97	1.06	1.12
time (sec)	N/A	0.436	20.031	1.883	5.085	0.286	165.508	0.658	2.848

Problem 636	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	644	617	317	0	0	0	0	0	0
N.S.	1	0.96	0.49	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.275	2.661	0.000	0.000	0.000	0.000	0.000	0.000

Problem 637	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	455	450	259	0	0	0	0	0	0
N.S.	1	0.99	0.57	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.826	1.425	0.000	0.000	0.000	0.000	0.000	0.000

Problem 638	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	327	315	205	0	0	0	0	0	0
N.S.	1	0.96	0.63	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.797	0.621	0.000	0.000	0.000	0.000	0.000	0.000

Problem 639	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	217	220	163	0	0	0	0	0	0
N.S.	1	1.01	0.75	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.912	0.250	0.000	0.000	0.000	0.000	0.000	0.000

Problem 640	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	299	305	10630	0	0	0	0	0	0
N.S.	1	1.02	35.55	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.372	32.020	0.000	0.000	0.000	0.000	0.000	0.000

Problem 641	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	F(-1)	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	0	33	35	54	0	35	37
N.S.	1	1.00	0.00	0.94	1.00	1.54	0.00	1.00	1.06
time (sec)	N/A	1.062	0.000	2.019	2.686	0.312	0.000	1.159	5.760

Problem 642	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	35	34	35	37
N.S.	1	1.00	1.06	0.94	1.00	1.00	0.97	1.00	1.06
time (sec)	N/A	0.477	33.021	1.774	2.710	0.281	8.463	0.694	3.271

Problem 643	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	35	34	35	37
N.S.	1	1.00	1.06	0.94	1.00	1.00	0.97	1.00	1.06
time (sec)	N/A	0.501	39.765	1.580	2.614	0.278	3.118	0.730	4.944

Problem 644	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	35	35	37	33	35	63	34	35	37
N.S.	1	1.00	1.06	0.94	1.00	1.80	0.97	1.00	1.06
time (sec)	N/A	1.101	31.590	1.713	2.316	0.304	11.397	1.261	8.631

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [318] had the largest ratio of [.818181999999999965]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	14	13	0.96	29	0.448
2	A	12	11	0.96	29	0.379
3	A	6	6	1.03	27	0.222
4	A	2	2	1.00	21	0.095
5	A	8	8	1.00	27	0.296
6	A	8	8	1.00	29	0.276
7	A	11	10	1.05	29	0.345
8	A	13	12	1.01	29	0.414
9	A	13	12	0.96	29	0.414
10	A	16	15	0.93	31	0.484
11	A	14	13	0.94	31	0.419
12	A	8	8	0.99	29	0.276
13	A	4	4	0.88	23	0.174
14	A	10	10	1.00	29	0.345
15	A	10	10	1.00	31	0.323
16	A	10	10	1.00	31	0.323
17	A	13	12	1.05	31	0.387
18	A	15	14	1.01	31	0.452
19	A	17	16	0.98	31	0.516
20	A	9	9	0.95	29	0.310
21	A	5	5	0.87	23	0.217
22	A	13	13	1.05	29	0.448

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	12	12	1.01	31	0.387
24	A	12	12	1.02	31	0.387
25	A	13	13	1.05	31	0.419
26	A	15	14	1.07	31	0.452
27	A	17	16	1.03	31	0.516
28	A	19	18	1.00	31	0.581
29	A	9	9	0.92	29	0.310
30	A	5	5	0.83	23	0.217
31	A	15	15	1.07	29	0.517
32	A	15	15	1.04	31	0.484
33	A	15	15	0.95	31	0.484
34	A	15	15	1.04	31	0.484
35	A	15	15	1.06	31	0.484
36	A	17	16	1.08	31	0.516
37	A	20	19	1.05	31	0.613
38	A	12	11	0.88	31	0.355
39	A	10	9	0.88	31	0.290
40	A	4	4	1.09	31	0.129
41	A	9	9	0.87	29	0.310
42	A	4	4	1.00	23	0.174
43	A	5	5	1.00	29	0.172
44	A	9	8	1.01	31	0.258
45	A	11	10	0.93	31	0.323
46	A	11	10	0.90	31	0.323
47	A	13	12	0.94	31	0.387
48	A	6	6	1.03	31	0.194
49	A	11	11	1.01	31	0.355
50	A	9	9	1.04	29	0.310
51	A	4	4	0.98	23	0.174
52	A	7	7	1.01	29	0.241
53	A	11	10	1.07	31	0.323
54	A	13	12	0.99	31	0.387
55	A	14	13	0.96	31	0.419
56	A	15	14	0.99	31	0.452

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	8	8	1.07	31	0.258
58	A	13	13	1.05	31	0.419
59	A	11	11	1.08	31	0.355
60	A	9	9	1.02	29	0.310
61	A	6	6	0.97	23	0.261
62	A	9	9	1.09	29	0.310
63	A	13	12	1.11	31	0.387
64	A	15	14	1.04	31	0.452
65	A	10	10	1.10	31	0.323
66	A	15	15	1.09	31	0.484
67	A	13	13	1.12	31	0.419
68	A	11	11	1.08	31	0.355
69	A	11	11	0.99	29	0.379
70	A	8	8	0.96	23	0.348
71	A	11	11	1.14	29	0.379
72	A	15	14	1.14	31	0.452
73	A	17	16	1.06	31	0.516
74	A	11	11	1.00	33	0.333
75	A	9	9	1.01	33	0.273
76	A	9	9	1.11	31	0.290
77	A	4	4	1.00	25	0.160
78	A	6	5	1.00	31	0.161
79	A	6	5	1.00	33	0.152
80	A	8	7	0.93	33	0.212
81	A	10	9	0.94	33	0.273
82	A	14	14	1.02	33	0.424
83	A	12	12	1.04	33	0.364
84	A	11	11	1.06	31	0.355
85	A	6	6	0.98	25	0.240
86	A	9	8	1.03	31	0.258
87	A	9	8	1.06	33	0.242
88	A	9	8	0.99	33	0.242
89	A	11	10	0.98	33	0.303
90	A	13	12	0.96	33	0.364

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
91	A	15	15	1.05	33	0.455
92	A	13	13	1.03	31	0.419
93	A	8	8	0.96	25	0.320
94	A	12	11	1.06	31	0.355
95	A	12	11	1.06	33	0.333
96	A	12	11	1.02	33	0.333
97	A	12	11	1.03	33	0.333
98	A	14	13	1.01	33	0.394
99	A	16	15	1.00	33	0.455
100	A	17	16	1.12	33	0.485
101	A	14	13	1.09	33	0.394
102	A	11	10	1.06	31	0.323
103	A	6	5	1.00	25	0.200
104	A	8	7	1.00	31	0.226
105	A	11	10	1.08	33	0.303
106	A	14	13	1.07	33	0.394
107	A	20	19	1.10	33	0.576
108	A	17	16	1.07	33	0.485
109	A	14	13	1.05	33	0.394
110	A	11	10	1.03	31	0.323
111	A	6	5	1.00	25	0.200
112	A	11	10	1.05	31	0.323
113	A	14	13	1.05	33	0.394
114	A	17	16	1.05	33	0.485
115	A	20	19	1.10	33	0.576
116	A	17	16	1.07	33	0.485
117	A	14	13	1.05	33	0.394
118	A	11	10	1.05	31	0.323
119	A	8	7	0.98	25	0.280
120	A	14	13	1.07	31	0.419
121	A	17	16	1.08	33	0.485
122	A	20	19	1.07	33	0.576
123	A	14	14	0.99	31	0.452
124	A	12	12	0.98	31	0.387

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
125	A	12	12	1.02	31	0.387
126	A	10	10	1.04	31	0.323
127	A	10	10	1.00	31	0.323
128	A	12	12	1.02	31	0.387
129	A	12	12	0.95	31	0.387
130	A	14	14	0.98	33	0.424
131	A	14	14	1.01	33	0.424
132	A	12	12	1.05	33	0.364
133	A	12	12	1.03	33	0.364
134	A	12	12	1.03	33	0.364
135	A	14	14	0.98	33	0.424
136	A	14	14	0.96	33	0.424
137	A	17	17	1.00	33	0.515
138	A	18	18	1.03	33	0.545
139	A	15	15	1.06	33	0.455
140	A	15	15	1.04	33	0.455
141	A	16	16	1.06	33	0.485
142	A	15	15	1.02	33	0.455
143	A	17	17	1.01	33	0.515
144	A	18	18	0.99	33	0.545
145	A	10	10	0.95	33	0.303
146	A	10	10	0.98	33	0.303
147	A	8	8	1.06	33	0.242
148	A	8	8	1.07	33	0.242
149	A	10	10	0.97	33	0.303
150	A	10	10	0.94	33	0.303
151	A	12	12	0.99	33	0.364
152	A	13	13	1.03	33	0.394
153	A	10	10	1.04	33	0.303
154	A	10	10	1.07	33	0.303
155	A	10	10	1.07	33	0.303
156	A	12	12	1.00	33	0.364
157	A	13	13	0.98	33	0.394
158	A	15	15	1.02	33	0.455

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
159	A	15	15	1.05	33	0.455
160	A	13	13	1.08	33	0.394
161	A	13	13	1.08	33	0.394
162	A	13	13	1.08	33	0.394
163	A	13	13	1.08	33	0.394
164	A	15	15	1.03	33	0.455
165	A	15	15	1.00	33	0.455
166	A	12	11	0.95	35	0.314
167	A	10	9	0.94	35	0.257
168	A	8	7	0.94	35	0.200
169	A	6	5	1.00	35	0.143
170	A	6	5	1.00	35	0.143
171	A	4	4	1.00	35	0.114
172	A	6	6	0.98	35	0.171
173	A	8	8	0.98	35	0.229
174	A	13	12	0.96	35	0.343
175	A	11	10	0.98	35	0.286
176	A	9	8	0.99	35	0.229
177	A	9	8	1.00	35	0.229
178	A	9	8	1.02	35	0.229
179	A	7	7	1.04	35	0.200
180	A	9	9	1.00	35	0.257
181	A	11	11	0.98	35	0.314
182	A	16	15	1.00	35	0.429
183	A	14	13	1.01	35	0.371
184	A	12	11	1.03	35	0.314
185	A	12	11	1.00	35	0.314
186	A	12	11	1.02	35	0.314
187	A	12	11	1.05	35	0.314
188	A	10	10	1.06	35	0.286
189	A	12	12	1.03	35	0.343
190	A	14	14	1.01	35	0.400
191	A	14	13	1.07	35	0.371
192	A	11	10	1.06	35	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
193	A	8	7	1.00	35	0.200
194	A	7	6	1.00	35	0.171
195	A	10	9	1.06	35	0.257
196	A	13	12	1.10	35	0.343
197	A	14	13	1.04	35	0.371
198	A	11	10	1.03	35	0.286
199	A	7	6	1.00	35	0.171
200	A	10	9	1.01	35	0.257
201	A	13	12	1.05	35	0.343
202	A	17	16	1.07	35	0.457
203	A	14	13	1.06	35	0.371
204	A	10	9	1.04	35	0.257
205	A	10	9	1.04	35	0.257
206	A	13	12	1.05	35	0.343
207	A	16	15	1.07	35	0.429
208	A	20	19	1.08	35	0.543
209	A	17	16	1.08	35	0.457
210	A	13	12	1.07	35	0.343
211	A	13	12	1.07	35	0.343
212	A	13	12	1.07	35	0.343
213	A	16	15	1.07	35	0.429
214	A	19	18	1.09	35	0.514
215	A	12	11	0.92	29	0.379
216	A	6	6	1.02	27	0.222
217	A	2	2	1.00	21	0.095
218	A	8	8	1.00	27	0.296
219	A	8	8	1.00	29	0.276
220	A	11	10	1.05	29	0.345
221	A	13	12	0.98	29	0.414
222	A	13	12	0.93	29	0.414
223	A	12	11	0.87	31	0.355
224	A	8	8	1.04	29	0.276
225	A	4	4	1.03	23	0.174
226	A	8	8	1.00	29	0.276

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
227	A	9	9	1.00	31	0.290
228	A	9	9	1.05	31	0.290
229	A	12	11	1.05	31	0.355
230	A	14	13	0.96	31	0.419
231	A	15	14	0.86	31	0.452
232	A	10	10	1.04	29	0.345
233	A	6	6	1.05	23	0.261
234	A	11	11	1.07	29	0.379
235	A	10	10	1.02	31	0.323
236	A	11	11	1.02	31	0.355
237	A	12	12	1.06	31	0.387
238	A	14	13	1.06	31	0.419
239	A	16	15	0.95	31	0.484
240	A	17	16	0.86	31	0.516
241	A	13	13	1.03	29	0.448
242	A	8	8	1.05	23	0.348
243	A	13	13	1.07	29	0.448
244	A	13	13	1.05	31	0.419
245	A	13	13	0.96	31	0.419
246	A	14	14	1.04	31	0.452
247	A	14	14	1.06	31	0.452
248	A	16	15	1.06	31	0.484
249	A	19	18	0.93	31	0.581
250	A	13	12	1.10	31	0.387
251	A	10	9	1.09	31	0.290
252	A	11	10	0.97	29	0.345
253	A	6	5	1.00	23	0.217
254	A	7	6	1.00	29	0.207
255	A	11	10	0.97	31	0.323
256	A	13	12	1.08	31	0.387
257	A	16	15	1.09	31	0.484
258	A	14	13	1.05	31	0.419
259	A	10	9	1.19	31	0.290
260	A	10	9	1.22	29	0.310

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
261	A	8	7	1.11	23	0.304
262	A	9	8	1.19	29	0.276
263	A	12	11	1.10	31	0.355
264	A	15	14	1.05	31	0.452
265	A	17	16	1.07	31	0.516
266	A	13	12	1.13	31	0.387
267	A	11	10	1.17	31	0.323
268	A	12	11	1.14	29	0.379
269	A	11	10	1.16	23	0.435
270	A	11	10	1.21	29	0.345
271	A	14	13	1.12	31	0.419
272	A	17	16	1.06	31	0.516
273	A	17	16	1.14	31	0.516
274	A	15	14	1.18	31	0.452
275	A	12	11	1.16	31	0.355
276	A	14	13	1.15	29	0.448
277	A	13	12	1.18	23	0.522
278	A	14	13	1.21	29	0.448
279	A	16	15	1.13	31	0.484
280	A	19	18	1.08	31	0.581
281	A	5	4	0.96	34	0.118
282	A	4	4	1.00	34	0.118
283	A	3	3	1.00	32	0.094
284	A	2	2	1.00	26	0.077
285	A	3	3	1.00	32	0.094
286	A	5	4	1.00	34	0.118
287	A	5	5	1.00	34	0.147
288	A	5	4	0.96	34	0.118
289	A	11	10	1.07	34	0.294
290	A	11	10	1.03	34	0.294
291	A	7	6	1.00	32	0.188
292	A	5	4	1.00	26	0.154
293	A	8	7	1.00	32	0.219
294	A	12	11	1.02	34	0.324

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
295	A	14	13	1.07	34	0.382
296	A	21	21	1.06	33	0.636
297	A	18	18	1.05	33	0.545
298	A	17	17	1.03	31	0.548
299	A	12	12	1.01	25	0.480
300	A	14	14	1.01	31	0.452
301	A	18	18	1.07	33	0.545
302	A	21	21	1.02	33	0.636
303	A	24	24	1.03	33	0.727
304	A	21	21	1.04	33	0.636
305	A	20	20	1.03	31	0.645
306	A	15	15	1.03	25	0.600
307	A	18	18	1.03	31	0.581
308	A	18	18	1.06	33	0.545
309	A	21	21	1.05	33	0.636
310	A	24	24	1.04	33	0.727
311	A	24	24	1.04	33	0.727
312	A	23	23	1.03	31	0.742
313	A	18	18	1.04	25	0.720
314	A	21	21	1.04	31	0.677
315	A	21	21	1.05	33	0.636
316	A	21	21	1.03	33	0.636
317	A	24	24	1.05	33	0.727
318	A	27	27	1.04	33	0.818
319	A	18	18	1.06	33	0.545
320	A	15	15	1.05	33	0.455
321	A	14	14	1.02	31	0.452
322	A	9	9	1.00	25	0.360
323	A	9	9	1.00	31	0.290
324	A	18	18	1.05	33	0.545
325	A	21	21	1.02	33	0.636
326	A	18	18	1.01	33	0.545
327	A	15	15	1.07	33	0.455
328	A	14	14	1.07	31	0.452

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
329	A	12	12	1.06	25	0.480
330	A	14	14	1.04	31	0.452
331	A	21	21	1.11	33	0.636
332	A	24	24	1.07	33	0.727
333	A	21	21	1.01	33	0.636
334	A	18	18	1.02	33	0.545
335	A	15	15	1.02	33	0.455
336	A	17	17	1.03	31	0.548
337	A	15	15	1.04	25	0.600
338	A	21	21	1.07	31	0.677
339	A	24	24	1.08	33	0.727
340	A	27	27	1.07	33	0.818
341	A	5	5	1.00	28	0.179
342	A	5	5	1.00	34	0.147
343	A	8	8	1.00	28	0.286
344	A	15	15	1.04	34	0.441
345	A	14	14	0.95	31	0.452
346	A	12	12	0.96	31	0.387
347	A	12	12	0.99	31	0.387
348	A	10	10	1.04	31	0.323
349	A	10	10	1.00	31	0.323
350	A	12	12	0.99	31	0.387
351	A	12	12	0.93	31	0.387
352	A	15	15	0.88	33	0.455
353	A	13	13	0.91	33	0.394
354	A	13	13	0.96	33	0.394
355	A	11	11	1.04	33	0.333
356	A	11	11	1.02	33	0.333
357	A	11	11	1.02	33	0.333
358	A	13	13	0.94	33	0.394
359	A	16	16	0.89	33	0.485
360	A	16	16	0.96	33	0.485
361	A	14	14	1.05	33	0.424
362	A	14	14	1.03	33	0.424

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
363	A	14	14	0.99	33	0.424
364	A	14	14	1.02	33	0.424
365	A	15	15	1.10	33	0.455
366	A	12	12	1.08	33	0.364
367	A	8	8	0.97	33	0.242
368	A	5	5	1.00	33	0.152
369	A	10	10	1.00	33	0.303
370	A	15	15	1.08	33	0.455
371	A	15	15	0.97	33	0.455
372	A	12	12	0.98	33	0.364
373	A	12	12	0.97	33	0.364
374	A	12	12	0.92	33	0.364
375	A	15	15	0.95	33	0.455
376	A	18	18	0.95	33	0.545
377	A	15	15	1.01	33	0.455
378	A	15	15	1.01	33	0.455
379	A	15	15	0.99	33	0.455
380	A	15	15	0.97	33	0.455
381	A	18	18	0.98	33	0.545
382	A	21	21	0.98	33	0.636
383	A	5	5	1.00	36	0.139
384	A	5	5	1.00	36	0.139
385	A	3	3	1.00	36	0.083
386	A	3	3	1.00	36	0.083
387	A	5	5	1.00	36	0.139
388	A	5	5	1.00	36	0.139
389	A	13	13	1.08	36	0.361
390	A	9	9	1.04	36	0.250
391	A	6	6	1.00	36	0.167
392	A	3	3	1.00	36	0.083
393	A	11	11	0.99	36	0.306
394	A	16	16	1.05	36	0.444
395	A	17	17	1.01	35	0.486
396	A	13	13	1.00	35	0.371

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
397	A	11	11	1.02	35	0.314
398	A	8	8	1.00	35	0.229
399	A	8	8	1.00	35	0.229
400	A	11	11	1.03	35	0.314
401	A	14	14	1.04	35	0.400
402	A	20	20	1.01	35	0.571
403	A	17	17	1.00	35	0.486
404	A	14	14	1.01	35	0.400
405	A	13	13	1.03	35	0.371
406	A	11	11	1.00	35	0.314
407	A	11	11	1.02	35	0.314
408	A	14	14	1.03	35	0.400
409	A	17	17	1.02	35	0.486
410	A	23	23	1.01	35	0.657
411	A	20	20	1.01	35	0.571
412	A	17	17	1.01	35	0.486
413	A	16	16	1.01	35	0.457
414	A	16	16	1.02	35	0.457
415	A	14	14	1.01	35	0.400
416	A	14	14	1.03	35	0.400
417	A	17	17	1.02	35	0.486
418	A	20	20	1.03	35	0.571
419	A	11	11	1.01	43	0.256
420	A	14	14	1.00	35	0.400
421	A	14	14	1.02	35	0.400
422	A	5	5	1.00	35	0.143
423	A	5	5	1.00	35	0.143
424	A	8	8	1.01	35	0.229
425	A	11	11	1.03	35	0.314
426	A	13	13	1.05	35	0.371
427	A	10	10	1.01	35	0.286
428	A	7	7	1.08	35	0.200
429	A	8	8	1.09	35	0.229
430	A	11	11	1.05	35	0.314

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
431	A	16	16	1.04	35	0.457
432	A	13	13	1.04	35	0.371
433	A	10	10	1.05	35	0.286
434	A	10	10	1.03	35	0.286
435	A	11	11	1.05	35	0.314
436	A	14	14	1.02	35	0.400
437	A	16	16	1.02	38	0.421
438	A	3	3	1.00	38	0.079
439	A	3	3	1.00	38	0.079
440	A	6	6	1.00	38	0.158
441	A	2	2	1.00	33	0.061
442	A	2	2	1.00	33	0.061
443	A	4	4	1.00	33	0.121
444	A	4	4	1.00	33	0.121
445	A	2	2	1.00	33	0.061
446	A	2	2	1.00	33	0.061
447	A	4	4	1.00	33	0.121
448	A	4	4	1.00	33	0.121
449	N/A	2	0	1.00	33	0.000
450	A	12	12	1.03	33	0.364
451	A	10	10	1.00	33	0.303
452	A	8	8	1.01	33	0.242
453	A	8	8	0.98	31	0.258
454	A	9	8	0.98	33	0.242
455	N/A	5	0	1.00	35	0.000
456	N/A	2	0	1.00	35	0.000
457	N/A	2	0	1.00	35	0.000
458	N/A	5	0	1.00	35	0.000
459	A	14	14	0.97	31	0.452
460	A	14	14	1.01	31	0.452
461	A	12	12	1.00	31	0.387
462	A	12	12	1.03	31	0.387
463	A	14	14	1.01	31	0.452

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
464	A	14	14	0.99	31	0.452
465	A	16	16	0.98	33	0.485
466	A	14	14	1.02	33	0.424
467	A	14	14	1.02	33	0.424
468	A	15	15	1.04	33	0.455
469	A	17	17	1.00	33	0.515
470	A	19	19	1.01	33	0.576
471	A	17	17	1.02	33	0.515
472	A	18	18	1.01	33	0.545
473	A	17	17	1.02	33	0.515
474	A	17	17	1.05	33	0.515
475	A	20	20	1.02	33	0.606
476	A	14	14	0.96	33	0.424
477	A	14	14	0.99	33	0.424
478	A	12	12	1.05	33	0.364
479	A	12	12	1.04	33	0.364
480	A	14	14	0.98	33	0.424
481	A	14	14	0.95	33	0.424
482	A	16	16	1.00	33	0.485
483	A	15	15	1.06	33	0.455
484	A	15	15	1.04	33	0.455
485	A	14	14	1.03	33	0.424
486	A	17	17	1.02	33	0.515
487	A	19	19	1.02	33	0.576
488	A	17	17	1.07	33	0.515
489	A	18	18	1.07	33	0.545
490	A	17	17	1.07	33	0.515
491	A	17	17	1.07	33	0.515
492	A	19	19	1.04	33	0.576
493	A	12	12	1.07	35	0.343
494	A	10	10	1.10	35	0.286
495	A	8	8	1.15	35	0.229
496	A	6	6	1.25	35	0.171
497	A	8	7	1.01	35	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
498	A	8	7	1.01	35	0.200
499	A	10	9	0.95	35	0.257
500	A	12	11	0.95	35	0.314
501	A	15	15	1.05	35	0.429
502	A	13	13	1.07	35	0.371
503	A	11	11	1.12	35	0.314
504	A	9	9	1.19	35	0.257
505	A	11	10	1.03	35	0.286
506	A	11	10	1.01	35	0.286
507	A	11	10	1.00	35	0.286
508	A	13	12	0.98	35	0.343
509	A	15	14	0.97	35	0.400
510	A	18	18	1.06	35	0.514
511	A	16	16	1.09	35	0.457
512	A	14	14	1.12	35	0.400
513	A	12	12	1.17	35	0.343
514	A	14	13	1.05	35	0.371
515	A	14	13	1.03	35	0.371
516	A	14	13	1.01	35	0.371
517	A	14	13	1.03	35	0.371
518	A	16	15	1.02	35	0.429
519	A	18	17	1.00	35	0.486
520	A	21	20	1.14	35	0.571
521	A	18	17	1.12	35	0.486
522	A	15	14	1.09	35	0.400
523	A	12	11	1.06	35	0.314
524	A	9	8	1.01	35	0.229
525	A	10	9	0.86	35	0.257
526	A	13	12	0.94	35	0.343
527	A	16	15	0.97	35	0.429
528	A	13	12	0.95	54	0.222
529	A	21	20	1.09	35	0.571
530	A	18	17	1.07	35	0.486
531	A	15	14	1.05	35	0.400

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
532	A	12	11	1.02	35	0.314
533	A	9	8	1.01	35	0.229
534	A	13	12	0.92	35	0.343
535	A	16	15	0.95	35	0.429
536	A	21	20	1.09	35	0.571
537	A	18	17	1.07	35	0.486
538	A	15	14	1.05	35	0.400
539	A	12	11	1.04	35	0.314
540	A	12	11	1.04	35	0.314
541	A	16	15	0.97	35	0.429
542	A	19	18	0.99	35	0.514
543	A	21	20	1.09	35	0.571
544	A	18	17	1.07	35	0.486
545	A	15	14	1.07	35	0.400
546	A	15	14	1.07	35	0.400
547	A	15	14	1.07	35	0.400
548	A	19	18	1.00	35	0.514
549	A	22	21	1.02	35	0.600
550	A	14	14	0.94	31	0.452
551	A	14	14	0.99	31	0.452
552	A	12	12	1.00	31	0.387
553	A	12	12	1.03	31	0.387
554	A	14	14	0.99	31	0.452
555	A	14	14	0.97	31	0.452
556	A	18	18	0.93	33	0.545
557	A	16	16	1.01	33	0.485
558	A	16	16	1.02	33	0.485
559	A	16	16	1.02	33	0.485
560	A	18	18	0.95	33	0.545
561	A	21	21	0.94	33	0.636
562	A	19	19	1.02	33	0.576
563	A	19	19	1.00	33	0.576
564	A	19	19	1.02	33	0.576
565	A	19	19	1.03	33	0.576

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
566	A	21	21	0.95	33	0.636
567	A	20	20	1.07	33	0.606
568	A	15	15	1.00	33	0.455
569	A	11	11	1.00	33	0.333
570	A	14	14	0.85	33	0.424
571	A	17	17	1.03	33	0.515
572	A	23	23	0.96	33	0.697
573	A	20	20	0.97	33	0.606
574	A	17	17	0.97	33	0.515
575	A	17	17	0.96	33	0.515
576	A	17	17	0.97	33	0.515
577	A	20	20	0.96	33	0.606
578	A	23	23	0.99	33	0.697
579	A	20	20	1.00	33	0.606
580	A	20	20	0.99	33	0.606
581	A	20	20	0.99	33	0.606
582	A	20	20	1.00	33	0.606
583	A	23	23	1.00	33	0.697
584	A	7	7	1.00	36	0.194
585	A	7	7	1.00	36	0.194
586	A	5	5	1.00	36	0.139
587	A	5	5	1.00	36	0.139
588	A	7	7	1.00	36	0.194
589	A	7	7	1.00	36	0.194
590	A	16	16	1.00	35	0.457
591	A	13	13	0.97	35	0.371
592	A	10	10	0.94	35	0.286
593	A	10	10	0.91	35	0.286
594	A	13	13	0.93	35	0.371
595	A	15	15	0.92	35	0.429
596	A	19	19	0.94	35	0.543
597	A	19	19	0.99	35	0.543
598	A	16	16	0.98	35	0.457
599	A	13	13	0.97	35	0.371

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
600	A	13	13	0.92	35	0.371
601	A	15	15	0.95	35	0.429
602	A	16	16	0.93	35	0.457
603	A	19	19	0.94	35	0.543
604	A	22	22	0.95	35	0.629
605	A	22	22	1.00	35	0.629
606	A	19	19	0.99	35	0.543
607	A	16	16	0.98	35	0.457
608	A	16	16	0.94	35	0.457
609	A	18	18	0.95	35	0.514
610	A	18	18	0.95	35	0.514
611	A	19	19	0.95	35	0.543
612	A	22	22	0.96	35	0.629
613	A	25	25	0.96	35	0.714
614	A	13	13	0.98	35	0.371
615	A	10	10	0.95	35	0.286
616	A	7	7	0.93	35	0.200
617	A	7	7	0.93	35	0.200
618	A	16	16	0.94	35	0.457
619	A	16	16	0.93	35	0.457
620	A	13	13	1.00	35	0.371
621	A	10	10	1.02	35	0.286
622	A	9	9	1.01	35	0.257
623	A	12	12	0.93	35	0.343
624	A	15	15	0.98	35	0.429
625	A	16	16	0.99	35	0.457
626	A	13	13	1.00	35	0.371
627	A	12	12	0.99	35	0.343
628	A	12	12	1.00	35	0.343
629	A	15	15	0.98	35	0.429
630	A	18	18	0.98	35	0.514
631	A	8	8	0.92	38	0.211
632	A	5	5	1.00	38	0.132
633	A	5	5	1.00	38	0.132

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
634	A	18	18	0.93	38	0.474
635	N/A	4	0	1.00	33	0.000
636	A	21	21	0.96	33	0.636
637	A	19	19	0.99	33	0.576
638	A	16	16	0.96	33	0.485
639	A	11	11	1.01	31	0.355
640	A	15	14	1.02	33	0.424
641	N/A	7	0	1.00	35	0.000
642	N/A	4	0	1.00	35	0.000
643	N/A	4	0	1.00	35	0.000
644	N/A	7	0	1.00	35	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \dots$	232
3.2	$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \dots$	240
3.3	$\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \dots$	247
3.4	$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx \dots$	253
3.5	$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx \dots$	258
3.6	$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx \dots$	264
3.7	$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx \dots$	270
3.8	$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx \dots$	277
3.9	$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx \dots$	284
3.10	$\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \dots$	292
3.11	$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \dots$	302
3.12	$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \dots$	311
3.13	$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \dots$	318
3.14	$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec(c + dx) dx \dots$	324
3.15	$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^2(c + dx) dx \dots$	331
3.16	$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^3(c + dx) dx \dots$	338
3.17	$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^4(c + dx) dx \dots$	345
3.18	$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^5(c + dx) dx \dots$	354
3.19	$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \dots$	363
3.20	$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \dots$	373
3.21	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \dots$	381
3.22	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) \sec(c + dx) dx \dots$	387
3.23	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) \sec^2(c + dx) dx \dots$	395
3.24	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) \sec^3(c + dx) dx \dots$	404
3.25	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) \sec^4(c + dx) dx \dots$	413
3.26	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) \sec^5(c + dx) dx \dots$	422
3.27	$\int (a + a \cos(c + dx))^3(A + B \cos(c + dx)) \sec^6(c + dx) dx \dots$	431
3.28	$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx \dots$	441

3.29	$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$	452
3.30	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$	461
3.31	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec(c + dx) dx$	468
3.32	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^2(c + dx) dx$	478
3.33	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^3(c + dx) dx$	488
3.34	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^4(c + dx) dx$	498
3.35	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^5(c + dx) dx$	508
3.36	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^6(c + dx) dx$	517
3.37	$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^7(c + dx) dx$	527
3.38	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	537
3.39	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	546
3.40	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	554
3.41	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	561
3.42	$\int \frac{A+B \cos(c+dx)}{a+a \cos(c+dx)} dx$	567
3.43	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{a+a \cos(c+dx)} dx$	572
3.44	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{a+a \cos(c+dx)} dx$	578
3.45	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{a+a \cos(c+dx)} dx$	585
3.46	$\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+a \cos(c+dx)} dx$	592
3.47	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	600
3.48	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	609
3.49	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	617
3.50	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	625
3.51	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2} dx$	632
3.52	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^2} dx$	637
3.53	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^2} dx$	643
3.54	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^2} dx$	651
3.55	$\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{(a+a \cos(c+dx))^2} dx$	660
3.56	$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	669
3.57	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	679
3.58	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	687
3.59	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	696
3.60	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	703
3.61	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3} dx$	710
3.62	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^3} dx$	716
3.63	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^3} dx$	723

3.64	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^3} dx$	732
3.65	$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$	742
3.66	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$	751
3.67	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$	761
3.68	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$	770
3.69	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$	777
3.70	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^4} dx$	784
3.71	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^4} dx$	790
3.72	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^4} dx$	798
3.73	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^4} dx$	808
3.74	$\int \cos^3(c+dx) \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	819
3.75	$\int \cos^2(c+dx) \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	827
3.76	$\int \cos(c+dx) \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	834
3.77	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	840
3.78	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec(c+dx) dx$	845
3.79	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^2(c+dx) dx$	851
3.80	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^3(c+dx) dx$	858
3.81	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^4(c+dx) dx$	866
3.82	$\int \cos^3(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$	875
3.83	$\int \cos^2(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$	884
3.84	$\int \cos(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$	892
3.85	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$	899
3.86	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec(c+dx) dx$	905
3.87	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^2(c+dx) dx$	912
3.88	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^3(c+dx) dx$	920
3.89	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^4(c+dx) dx$	928
3.90	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^5(c+dx) dx$	937
3.91	$\int \cos^2(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$	947
3.92	$\int \cos(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$	956
3.93	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$	964
3.94	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec(c+dx) dx$	970
3.95	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^2(c+dx) dx$	978
3.96	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^3(c+dx) dx$	987
3.97	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^4(c+dx) dx$	996
3.98	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^5(c+dx) dx$	1005
3.99	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^6(c+dx) dx$	1014
3.100	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$	1023

3.101	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$	1033
3.102	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$	1042
3.103	$\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	1049
3.104	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	1055
3.105	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	1062
3.106	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	1070
3.107	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$	1079
3.108	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$	1089
3.109	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$	1098
3.110	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$	1106
3.111	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$	1112
3.112	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$	1118
3.113	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$	1126
3.114	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$	1135
3.115	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$	1145
3.116	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$	1156
3.117	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$	1165
3.118	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$	1173
3.119	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$	1180
3.120	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$	1186
3.121	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$	1195
3.122	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$	1205
3.123	$\int \cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$	1215
3.124	$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$	1224
3.125	$\int \sqrt{\cos(c+dx)}(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$	1233
3.126	$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	1241
3.127	$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	1248
3.128	$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	1255
3.129	$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	1263
3.130	$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$	1273
3.131	$\int \sqrt{\cos(c+dx)}(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$	1283
3.132	$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	1293

3.133	$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	1302
3.134	$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	1311
3.135	$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	1320
3.136	$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$	1330
3.137	$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$	1340
3.138	$\int \sqrt{\cos(c+dx)}(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$	1352
3.139	$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	1363
3.140	$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	1373
3.141	$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	1384
3.142	$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	1395
3.143	$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$	1406
3.144	$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$	1418
3.145	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	1430
3.146	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	1438
3.147	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$	1446
3.148	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))} dx$	1454
3.149	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))} dx$	1462
3.150	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))} dx$	1470
3.151	$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	1479
3.152	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	1487
3.153	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	1496
3.154	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$	1504
3.155	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^2} dx$	1513
3.156	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$	1521
3.157	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$	1530
3.158	$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	1539
3.159	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	1548
3.160	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	1557
3.161	$\int \frac{\cos^{\frac{1}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	1565

3.162	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$	1574
3.163	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^3} dx$	1583
3.164	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$	1591
3.165	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$	1600
3.166	$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	1610
3.167	$\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	1619
3.168	$\int \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) dx$	1627
3.169	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	1634
3.170	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	1640
3.171	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	1646
3.172	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	1652
3.173	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$	1659
3.174	$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$	1667
3.175	$\int \sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$	1676
3.176	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	1685
3.177	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	1693
3.178	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	1701
3.179	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	1708
3.180	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$	1715
3.181	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$	1724
3.182	$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$	1733
3.183	$\int \sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$	1743
3.184	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	1752
3.185	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	1761
3.186	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	1770
3.187	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	1778
3.188	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$	1786
3.189	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$	1795
3.190	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$	1804
3.191	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$	1813

3.192	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$	1821
3.193	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}} dx$	1828
3.194	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+a \cos(c+dx)}} dx$	1835
3.195	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+a \cos(c+dx)}} dx$	1842
3.196	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+a \cos(c+dx)}} dx$	1850
3.197	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	1859
3.198	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	1868
3.199	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	1875
3.200	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	1881
3.201	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	1888
3.202	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	1896
3.203	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	1906
3.204	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	1915
3.205	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	1922
3.206	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	1929
3.207	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	1938
3.208	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	1947
3.209	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	1958
3.210	$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	1968
3.211	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	1976
3.212	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	1984
3.213	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	1992
3.214	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	2002
3.215	$\int \cos^2(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$	2013
3.216	$\int \cos(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$	2020
3.217	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) dx$	2026
3.218	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec(c+dx) dx$	2031
3.219	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^2(c+dx) dx$	2037
3.220	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^3(c+dx) dx$	2043
3.221	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^4(c+dx) dx$	2050
3.222	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^5(c+dx) dx$	2057

3.223	$\int \cos^2(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx)) dx$	2065
3.224	$\int \cos(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx)) dx$	2074
3.225	$\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) dx$	2082
3.226	$\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) \sec(c+dx) dx$	2088
3.227	$\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) \sec^2(c+dx) dx$	2095
3.228	$\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) \sec^3(c+dx) dx$	2102
3.229	$\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) \sec^4(c+dx) dx$	2109
3.230	$\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) \sec^5(c+dx) dx$	2117
3.231	$\int \cos^2(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx$	2126
3.232	$\int \cos(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx$	2136
3.233	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx$	2145
3.234	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) \sec(c+dx) dx$	2152
3.235	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) \sec^2(c+dx) dx$	2161
3.236	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) \sec^3(c+dx) dx$	2170
3.237	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) \sec^4(c+dx) dx$	2179
3.238	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) \sec^5(c+dx) dx$	2188
3.239	$\int (a+b\cos(c+dx))^3(A+B\cos(c+dx)) \sec^6(c+dx) dx$	2198
3.240	$\int \cos^2(c+dx)(a+b\cos(c+dx))^4(A+B\cos(c+dx)) dx$	2209
3.241	$\int \cos(c+dx)(a+b\cos(c+dx))^4(A+B\cos(c+dx)) dx$	2221
3.242	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) dx$	2231
3.243	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec(c+dx) dx$	2239
3.244	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec^2(c+dx) dx$	2249
3.245	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec^3(c+dx) dx$	2259
3.246	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec^4(c+dx) dx$	2269
3.247	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec^5(c+dx) dx$	2280
3.248	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec^6(c+dx) dx$	2291
3.249	$\int (a+b\cos(c+dx))^4(A+B\cos(c+dx)) \sec^7(c+dx) dx$	2302
3.250	$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$	2314
3.251	$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$	2323
3.252	$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$	2332
3.253	$\int \frac{A+B\cos(c+dx)}{a+b\cos(c+dx)} dx$	2340
3.254	$\int \frac{(A+B\cos(c+dx)) \sec(c+dx)}{a+b\cos(c+dx)} dx$	2347
3.255	$\int \frac{(A+B\cos(c+dx)) \sec^2(c+dx)}{a+b\cos(c+dx)} dx$	2353
3.256	$\int \frac{(A+B\cos(c+dx)) \sec^3(c+dx)}{a+b\cos(c+dx)} dx$	2361
3.257	$\int \frac{(A+B\cos(c+dx)) \sec^4(c+dx)}{a+b\cos(c+dx)} dx$	2370
3.258	$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$	2380
3.259	$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$	2390
3.260	$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$	2399

3.261	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2} dx$	2406
3.262	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx$	2413
3.263	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$	2421
3.264	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$	2431
3.265	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	2442
3.266	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	2454
3.267	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	2465
3.268	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	2474
3.269	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3} dx$	2482
3.270	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^3} dx$	2490
3.271	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^3} dx$	2500
3.272	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^3} dx$	2511
3.273	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$	2524
3.274	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$	2537
3.275	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$	2549
3.276	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$	2559
3.277	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^4} dx$	2569
3.278	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$	2579
3.279	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^4} dx$	2591
3.280	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$	2603
3.281	$\int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$	2617
3.282	$\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$	2622
3.283	$\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$	2627
3.284	$\int \frac{aB+bB \cos(c+dx)}{a+b \cos(c+dx)} dx$	2632
3.285	$\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx$	2636
3.286	$\int \frac{(aB+bB \cos(c+dx)) \sec^2(c+dx)}{a+b \cos(c+dx)} dx$	2641
3.287	$\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$	2646
3.288	$\int \frac{(aB+bB \cos(c+dx)) \sec^4(c+dx)}{a+b \cos(c+dx)} dx$	2651
3.289	$\int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	2656
3.290	$\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	2664
3.291	$\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	2671
3.292	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^2} dx$	2677
3.293	$\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx$	2683

3.294	$\int \frac{(aB+bB \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$	2689
3.295	$\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$	2697
3.296	$\int \cos^3(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$	2706
3.297	$\int \cos^2(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$	2718
3.298	$\int \cos(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$	2729
3.299	$\int \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$	2738
3.300	$\int \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) \sec(c+dx) dx$	2746
3.301	$\int \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) \sec^2(c+dx) dx$	2755
3.302	$\int \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) \sec^3(c+dx) dx$	2766
3.303	$\int \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) \sec^4(c+dx) dx$	2779
3.304	$\int \cos^2(c+dx) (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) dx$	2793
3.305	$\int \cos(c+dx) (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) dx$	2805
3.306	$\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) dx$	2815
3.307	$\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec(c+dx) dx$	2824
3.308	$\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^2(c+dx) dx$	2835
3.309	$\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^3(c+dx) dx$	2846
3.310	$\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^4(c+dx) dx$	2859
3.311	$\int \cos^2(c+dx) (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) dx$	2873
3.312	$\int \cos(c+dx) (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) dx$	2885
3.313	$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) dx$	2897
3.314	$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec(c+dx) dx$	2907
3.315	$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^2(c+dx) dx$	2919
3.316	$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^3(c+dx) dx$	2931
3.317	$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^4(c+dx) dx$	2943
3.318	$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^5(c+dx) dx$	2957
3.319	$\int \frac{\cos^3(c+dx) (A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$	2972
3.320	$\int \frac{\cos^2(c+dx) (A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$	2983
3.321	$\int \frac{\cos(c+dx) (A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$	2992
3.322	$\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	3001
3.323	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	3008
3.324	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	3014
3.325	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	3024
3.326	$\int \frac{\cos^3(c+dx) (A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$	3036
3.327	$\int \frac{\cos^2(c+dx) (A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$	3047
3.328	$\int \frac{\cos(c+dx) (A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$	3057
3.329	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$	3066
3.330	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$	3074

3.331	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$	3083
3.332	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$	3095
3.333	$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$	3109
3.334	$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$	3122
3.335	$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$	3133
3.336	$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$	3143
3.337	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$	3153
3.338	$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$	3162
3.339	$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$	3175
3.340	$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$	3189
3.341	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$	3204
3.342	$\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$	3209
3.343	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$	3214
3.344	$\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$	3220
3.345	$\int \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$	3229
3.346	$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$	3237
3.347	$\int \sqrt{\cos(c+dx)}(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$	3245
3.348	$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	3253
3.349	$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	3260
3.350	$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	3267
3.351	$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	3275
3.352	$\int \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$	3284
3.353	$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$	3294
3.354	$\int \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$	3304
3.355	$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	3313
3.356	$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	3321
3.357	$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	3329
3.358	$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$	3338
3.359	$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx$	3347
3.360	$\int \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx$	3359
3.361	$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$	3369
3.362	$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$	3379
3.363	$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$	3389

3.364	$\int \frac{(a+b \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^2(c+dx)} dx$	3399
3.365	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$	3409
3.366	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$	3418
3.367	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$	3426
3.368	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx$	3432
3.369	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$	3437
3.370	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$	3444
3.371	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	3453
3.372	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	3463
3.373	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	3472
3.374	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} dx$	3481
3.375	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$	3490
3.376	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$	3500
3.377	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	3511
3.378	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	3521
3.379	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$	3531
3.380	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^3} dx$	3541
3.381	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$	3551
3.382	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$	3562
3.383	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$	3575
3.384	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$	3580
3.385	$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$	3585
3.386	$\int \frac{aB+bB \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx$	3590
3.387	$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$	3595
3.388	$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$	3600
3.389	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	3605
3.390	$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	3613
3.391	$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$	3620
3.392	$\int \frac{aB+bB \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} dx$	3626

3.393	$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx \dots \dots \dots$	3631
3.394	$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx \dots \dots \dots$	3638
3.395	$\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) dx \dots \dots \dots$	3647
3.396	$\int \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) dx \dots \dots \dots$	3659
3.397	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \dots \dots \dots$	3670
3.398	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx \dots \dots \dots$	3679
3.399	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx \dots \dots \dots$	3687
3.400	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx \dots \dots \dots$	3695
3.401	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx \dots \dots \dots$	3704
3.402	$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx)) dx \dots \dots \dots$	3714
3.403	$\int \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx)) dx \dots \dots \dots$	3726
3.404	$\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \dots \dots \dots$	3739
3.405	$\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx \dots \dots \dots$	3750
3.406	$\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx \dots \dots \dots$	3760
3.407	$\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx \dots \dots \dots$	3769
3.408	$\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx \dots \dots \dots$	3779
3.409	$\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx \dots \dots \dots$	3790
3.410	$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx)) dx \dots \dots \dots$	3802
3.411	$\int \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx)) dx \dots \dots \dots$	3815
3.412	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \dots \dots \dots$	3828
3.413	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx \dots \dots \dots$	3840
3.414	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx \dots \dots \dots$	3852
3.415	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx \dots \dots \dots$	3864
3.416	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx \dots \dots \dots$	3874
3.417	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx \dots \dots \dots$	3884
3.418	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx \dots \dots \dots$	3895
3.419	$\int \frac{(a+b \cos(c+dx))^{\frac{5}{2}} \left(\frac{3bB}{2a} + B \cos(c+dx) \right)}{\cos^{\frac{5}{2}}(c+dx)} dx \dots \dots \dots$	3908
3.420	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx \dots \dots \dots$	3917
3.421	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx \dots \dots \dots$	3928

3.422	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx$	3938
3.423	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$	3944
3.424	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$	3951
3.425	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$	3959
3.426	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	3968
3.427	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	3979
3.428	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	3989
3.429	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	3997
3.430	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	4005
3.431	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	4014
3.432	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	4025
3.433	$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	4034
3.434	$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	4043
3.435	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	4051
3.436	$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	4060
3.437	$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	4070
3.438	$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	4080
3.439	$\int \frac{aB+bB \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	4085
3.440	$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	4090
3.441	$\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{2+3 \cos(c+dx)}} dx$	4096
3.442	$\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{-2+3 \cos(c+dx)}} dx$	4101
3.443	$\int \frac{1+\cos(c+dx)}{\sqrt{2-3 \cos(c+dx)} \cos^{\frac{3}{2}}(c+dx)} dx$	4106
3.444	$\int \frac{1+\cos(c+dx)}{\sqrt{-2-3 \cos(c+dx)} \cos^{\frac{3}{2}}(c+dx)} dx$	4111
3.445	$\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{3+2 \cos(c+dx)}} dx$	4116
3.446	$\int \frac{1+\cos(c+dx)}{\sqrt{3-2 \cos(c+dx)} \cos^{\frac{3}{2}}(c+dx)} dx$	4121
3.447	$\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{-3+2 \cos(c+dx)}} dx$	4126
3.448	$\int \frac{1+\cos(c+dx)}{\sqrt{-3-2 \cos(c+dx)} \cos^{\frac{3}{2}}(c+dx)} dx$	4131
3.449	$\int (c \cos(e+fx))^m (a+b \cos(e+fx))^n (A+B \cos(e+fx)) dx$	4136
3.450	$\int (c \cos(e+fx))^m (a+b \cos(e+fx))^4 (A+B \cos(e+fx)) dx$	4141
3.451	$\int (c \cos(e+fx))^m (a+b \cos(e+fx))^3 (A+B \cos(e+fx)) dx$	4150

3.452	$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx$	4158
3.453	$\int (c \cos(e + fx))^m (a + b \cos(e + fx)) (A + B \cos(e + fx)) dx$	4165
3.454	$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx$	4171
3.455	$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx$	4177
3.456	$\int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx$	4183
3.457	$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$	4188
3.458	$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx$	4193
3.459	$\int (a + a \cos(c + dx)) (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$	4199
3.460	$\int (a + a \cos(c + dx)) (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$	4208
3.461	$\int (a + a \cos(c + dx)) (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$	4216
3.462	$\int (a + a \cos(c + dx)) (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$	4223
3.463	$\int \frac{(a + a \cos(c + dx)) (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$	4230
3.464	$\int \frac{(a + a \cos(c + dx)) (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx$	4238
3.465	$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$	4246
3.466	$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$	4255
3.467	$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$	4264
3.468	$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$	4272
3.469	$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$	4281
3.470	$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$	4291
3.471	$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$	4301
3.472	$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$	4310
3.473	$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$	4320
3.474	$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$	4329
3.475	$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$	4339
3.476	$\int \frac{(A + B \cos(c + dx)) \sec^{5/2}(c + dx)}{a + a \cos(c + dx)} dx$	4349
3.477	$\int \frac{(A + B \cos(c + dx)) \sec^{3/2}(c + dx)}{a + a \cos(c + dx)} dx$	4358
3.478	$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx$	4366
3.479	$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sqrt{\sec(c + dx)}} dx$	4374
3.480	$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{3/2}(c + dx)} dx$	4382
3.481	$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{5/2}(c + dx)} dx$	4390
3.482	$\int \frac{(A + B \cos(c + dx)) \sec^{3/2}(c + dx)}{(a + a \cos(c + dx))^2} dx$	4398
3.483	$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx$	4407
3.484	$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$	4415
3.485	$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{3/2}(c + dx)} dx$	4424

3.486	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$	4432
3.487	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$	4441
3.488	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^3} dx$	4451
3.489	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$	4461
3.490	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$	4472
3.491	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$	4482
3.492	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$	4492
3.493	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$	4502
3.494	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$	4510
3.495	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$	4518
3.496	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$	4525
3.497	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$	4531
3.498	$\int \sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	4538
3.499	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	4545
3.500	$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$	4553
3.501	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{13}{2}}(c+dx) dx$	4561
3.502	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$	4570
3.503	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$	4579
3.504	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$	4587
3.505	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$	4595
3.506	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$	4603
3.507	$\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	4611
3.508	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	4619
3.509	$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$	4628
3.510	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{15}{2}}(c+dx) dx$	4637
3.511	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{13}{2}}(c+dx) dx$	4647
3.512	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$	4656
3.513	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$	4665
3.514	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$	4673
3.515	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$	4681
3.516	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$	4689
3.517	$\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	4697
3.518	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	4706
3.519	$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$	4715

3.520	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	4725
3.521	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	4736
3.522	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	4745
3.523	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	4753
3.524	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$	4761
3.525	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{\sqrt{a+a \cos(c+dx)}} dx$	4767
3.526	$\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}} dx$	4775
3.527	$\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$	4783
3.528	$\int \frac{(aA+(Ab+aB) \cos(c+dx)+bB \cos^2(c+dx)) \sqrt{\sec(c+dx)}}{\sqrt{a+a \cos(c+dx)}} dx$	4792
3.529	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	4800
3.530	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	4811
3.531	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	4821
3.532	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	4830
3.533	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$	4838
3.534	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)}} dx$	4844
3.535	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{3}{2}} \sec^{\frac{3}{2}}(c+dx)} dx$	4852
3.536	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	4862
3.537	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	4873
3.538	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	4883
3.539	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$	4892
3.540	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}} \sqrt{\sec(c+dx)}} dx$	4899
3.541	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}} \sec^{\frac{3}{2}}(c+dx)} dx$	4906
3.542	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}} \sec^{\frac{5}{2}}(c+dx)} dx$	4915
3.543	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	4927
3.544	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	4939
3.545	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$	4949
3.546	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{7}{2}} \sqrt{\sec(c+dx)}} dx$	4957
3.547	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{\frac{7}{2}} \sec^{\frac{3}{2}}(c+dx)} dx$	4966

3.548	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{5/2}(c+dx)} dx$	4975
3.549	$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{7/2}(c+dx)} dx$	4986
3.550	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$	4999
3.551	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$	5008
3.552	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$	5016
3.553	$\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	5023
3.554	$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	5030
3.555	$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sec^{3/2}(c+dx)} dx$	5038
3.556	$\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$	5046
3.557	$\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$	5055
3.558	$\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$	5064
3.559	$\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	5073
3.560	$\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	5082
3.561	$\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{9/2}(c+dx) dx$	5092
3.562	$\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$	5103
3.563	$\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$	5113
3.564	$\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$	5123
3.565	$\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	5133
3.566	$\int \frac{(a+b \cos(c+dx))^3 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	5143
3.567	$\int \frac{(A+B \cos(c+dx)) \sec^{5/2}(c+dx)}{a+b \cos(c+dx)} dx$	5154
3.568	$\int \frac{(A+B \cos(c+dx)) \sec^{3/2}(c+dx)}{a+b \cos(c+dx)} dx$	5164
3.569	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{a+b \cos(c+dx)} dx$	5172
3.570	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sqrt{\sec(c+dx)}} dx$	5179
3.571	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{3/2}(c+dx)} dx$	5187
3.572	$\int \frac{(A+B \cos(c+dx)) \sec^{5/2}(c+dx)}{(a+b \cos(c+dx))^2} dx$	5197
3.573	$\int \frac{(A+B \cos(c+dx)) \sec^{3/2}(c+dx)}{(a+b \cos(c+dx))^2} dx$	5209
3.574	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx$	5220
3.575	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$	5230
3.576	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{3/2}(c+dx)} dx$	5240
3.577	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{5/2}(c+dx)} dx$	5250
3.578	$\int \frac{(A+B \cos(c+dx)) \sec^{3/2}(c+dx)}{(a+b \cos(c+dx))^3} dx$	5261
3.579	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$	5274
3.580	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$	5286

3.581	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$	5298
3.582	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$	5309
3.583	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$	5321
3.584	$\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$	5334
3.585	$\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$	5340
3.586	$\int \frac{(aB+bB \cos(c+dx)) \sqrt{\sec(c+dx)}}{a+b \cos(c+dx)} dx$	5346
3.587	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx)) \sqrt{\sec(c+dx)}} dx$	5351
3.588	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$	5356
3.589	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)} dx$	5362
3.590	$\int \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$	5368
3.591	$\int \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$	5379
3.592	$\int \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$	5388
3.593	$\int \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$	5396
3.594	$\int \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	5404
3.595	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	5414
3.596	$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$	5425
3.597	$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$	5438
3.598	$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$	5449
3.599	$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$	5460
3.600	$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$	5469
3.601	$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$	5478
3.602	$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	5489
3.603	$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	5500
3.604	$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$	5512
3.605	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{13}{2}}(c+dx) dx$	5525
3.606	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$	5537
3.607	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$	5548
3.608	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$	5558
3.609	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$	5567
3.610	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$	5577
3.611	$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$	5589
3.612	$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$	5601
3.613	$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$	5614
3.614	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	5627

3.615	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	5637
3.616	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$	5646
3.617	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx$	5653
3.618	$\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sqrt{\sec(c+dx)}} dx$	5660
3.619	$\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$	5671
3.620	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	5683
3.621	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	5693
3.622	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	5702
3.623	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)}} dx$	5710
3.624	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}} \sec^{\frac{3}{2}}(c+dx)} dx$	5721
3.625	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	5733
3.626	$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	5744
3.627	$\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$	5753
3.628	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}} \sqrt{\sec(c+dx)}} dx$	5761
3.629	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}} \sec^{\frac{3}{2}}(c+dx)} dx$	5770
3.630	$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}} \sec^{\frac{5}{2}}(c+dx)} dx$	5781
3.631	$\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	5793
3.632	$\int \frac{(aB+bB \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$	5800
3.633	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}} \sqrt{\sec(c+dx)}} dx$	5805
3.634	$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}} \sec^{\frac{3}{2}}(c+dx)} dx$	5810
3.635	$\int (a+b \cos(e+fx))^n (A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5821
3.636	$\int (a+b \cos(e+fx))^4 (A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5827
3.637	$\int (a+b \cos(e+fx))^3 (A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5838
3.638	$\int (a+b \cos(e+fx))^2 (A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5848
3.639	$\int (a+b \cos(e+fx))(A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5856
3.640	$\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{a+b \cos(e+fx)} dx$	5863
3.641	$\int (a+b \cos(e+fx))^{\frac{3}{2}} (A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5871
3.642	$\int \sqrt{a+b \cos(e+fx)} (A+B \cos(e+fx))(c \sec(e+fx))^m dx$	5877
3.643	$\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{\sqrt{a+b \cos(e+fx)}} dx$	5883
3.644	$\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{(a+b \cos(e+fx))^{\frac{3}{2}}} dx$	5888

3.1 $\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.1.1	Optimal result	232
3.1.2	Mathematica [A] (verified)	233
3.1.3	Rubi [A] (verified)	233
3.1.4	Maple [A] (verified)	236
3.1.5	Fricas [A] (verification not implemented)	237
3.1.6	Sympy [B] (verification not implemented)	237
3.1.7	Maxima [A] (verification not implemented)	238
3.1.8	Giac [A] (verification not implemented)	238
3.1.9	Mupad [B] (verification not implemented)	239

3.1.1 Optimal result

Integrand size = 29, antiderivative size = 125

$$\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3}{8}a(A + B)x + \frac{a(5A + 4B) \sin(c + dx)}{5d}$$

$$+ \frac{3a(A + B) \cos(c + dx) \sin(c + dx)}{8d} + \frac{a(A + B) \cos^3(c + dx) \sin(c + dx)}{4d}$$

$$+ \frac{aB \cos^4(c + dx) \sin(c + dx)}{5d} - \frac{a(5A + 4B) \sin^3(c + dx)}{15d}$$

output $\frac{3}{8}a*(A+B)*x+1/5*a*(5*A+4*B)*\sin(d*x+c)/d+3/8*a*(A+B)*\cos(d*x+c)*\sin(d*x+c)/d+1/4*a*(A+B)*\cos(d*x+c)^3*\sin(d*x+c)/d+1/5*a*B*\cos(d*x+c)^4*\sin(d*x+c)/d-1/15*a*(5*A+4*B)*\sin(d*x+c)^3/d$

3.1.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.62

$$\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{a(480(A + B) \sin(c + dx) - 160(A + 2B) \sin^3(c + dx) + 96B \sin^5(c + dx) + 15(A + B)(12(c + dx) + 8 \sin^2(c + dx) + \sin[4(c + dx)]))}{480d}$$

input `Integrate[Cos[c + d*x]^3*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(a*(480*(A + B)*Sin[c + d*x] - 160*(A + 2*B)*Sin[c + d*x]^3 + 96*B*SIN[c + d*x]^5 + 15*(A + B)*(12*(c + d*x) + 8*SIN[2*(c + d*x)] + SIN[4*(c + d*x)])))/(480*d)`

3.1.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.96, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$, Rules used = {3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^3 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int \cos^3(c + dx) \left((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)\right) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^3 \left((aA + aB) \sin\left(c + dx + \frac{\pi}{2}\right) + aA + aB \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{3502}$$

$$\frac{1}{5} \int \cos^3(c + dx) (a(5A + 4B) + 5a(A + B) \cos(c + dx)) dx + \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d}$$

3.1. $\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{1}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 \left(a(5A + 4B) + 5a(A + B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \\
& \quad \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{3227} \\
& \frac{1}{5} \left(5a(A + B) \int \cos^4(c + dx) dx + a(5A + 4B) \int \cos^3(c + dx) dx \right) + \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \left(a(5A + 4B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 dx + 5a(A + B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^4 dx \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{3113} \\
& \frac{1}{5} \left(5a(A + B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^4 dx - \frac{a(5A + 4B) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{2009} \\
& \frac{1}{5} \left(5a(A + B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^4 dx - \frac{a(5A + 4B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{3115} \\
& \frac{1}{5} \left(5a(A + B) \left(\frac{3}{4} \int \cos^2(c + dx) dx + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) - \frac{a(5A + 4B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \left(5a(A + B) \left(\frac{3}{4} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) - \frac{a(5A + 4B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^4(c + dx)}{5d} \\
& \downarrow \text{3115}
\end{aligned}$$

3.1. $\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\frac{1}{5} \left(5a(A+B) \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) - \frac{a(5A+4B) \left(\frac{1}{3} \sin^3(c+dx) - \frac{aB \sin(c+dx) \cos^4(c+dx)}{5d} \right)}{d} \right)$$

↓ 24

$$\frac{1}{5} \left(5a(A+B) \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right) - \frac{a(5A+4B) \left(\frac{1}{3} \sin^3(c+dx) - \frac{aB \sin(c+dx) \cos^4(c+dx)}{5d} \right)}{d} \right)$$

input `Int[Cos[c + d*x]^3*(a + a*cos[c + d*x])*(A + B*cos[c + d*x]),x]`

output `(a*B*cos[c + d*x]^4*sin[c + d*x])/(5*d) + (-((a*(5*A + 4*B)*(-sin[c + d*x] + sin[c + d*x]^3/3))/d) + 5*a*(A + B)*((cos[c + d*x]^3*sin[c + d*x])/(4*d) + (3*(x/2 + (cos[c + d*x]*sin[c + d*x])/(2*d)))/4))/5`

3.1.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.1. $\int \cos^3(c+dx)(a + a \cos(c+dx))(A + B \cos(c+dx)) dx$

```
rule 3227 Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]
```

3.1.4 Maple [A] (verified)

Time = 3.46 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.70

method	result
parallelrisch	$\frac{(8(A+B) \sin(2dx+2c) + \frac{2(4A+5B) \sin(3dx+3c)}{3} + (A+B) \sin(4dx+4c) + \frac{2B \sin(5dx+5c)}{5} + 4(6A+5B) \sin(dx+c) + 12(A+B))}{32d}$
parts	$\frac{(aA+Ba) \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(\frac{dx+c}{2})}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right)}{d} + \frac{aA(2+\cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{Ba \left(\frac{8}{3} + \cos^4(dx+c) \right)}{d}$
derivativedivides	$\frac{Ba \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + aA \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(\frac{dx+c}{2})}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + Ba \left(\frac{\cos^3(dx+c)}{d} \right)$
default	$\frac{Ba \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + aA \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(\frac{dx+c}{2})}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + Ba \left(\frac{\cos^3(dx+c)}{d} \right)$
risch	$\frac{3axA}{8} + \frac{3aBx}{8} + \frac{3 \sin(dx+c)aA}{4d} + \frac{5aB \sin(dx+c)}{8d} + \frac{Ba \sin(5dx+5c)}{80d} + \frac{\sin(4dx+4c)aA}{32d} + \frac{\sin(4dx+4c)Ba}{32d} +$
norman	$\frac{3a(A+B)x}{8} + \frac{13a(A+B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4d} + \frac{3a(A+B) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{4d} + \frac{15a(A+B)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{8} + \frac{15a(A+B)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{4} +$

3.1. $\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

input `int(cos(d*x+c)^3*(a+cos(d*x+c)*a)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/32*(8*(A+B)*sin(2*d*x+2*c))+2/3*(4*A+5*B)*sin(3*d*x+3*c)+(A+B)*sin(4*d*x+4*c)+2/5*B*sin(5*d*x+5*c)+4*(6*A+5*B)*sin(d*x+c)+12*(A+B)*x*d*a/d`

3.1.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.70

$$\int \cos^3(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{45(A+B)adx + (24Ba\cos(dx+c)^4 + 30(A+B)a\cos(dx+c)^3 + 8(5A+4B)a\cos(dx+c)^2 + 45(A+B)a\cos(dx+c) + 16(5A+4B)a)\sin(dx+c)}{120d}$$

input `integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/120*(45*(A+B)*a*d*x + (24*B*a*cos(d*x+c)^4 + 30*(A+B)*a*cos(d*x+c)^3 + 8*(5*A+4*B)*a*cos(d*x+c)^2 + 45*(A+B)*a*cos(d*x+c) + 16*(5*A+4*B)*a)*sin(d*x+c))/d`

3.1.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. $2(117) = 234$.

Time = 0.27 (sec) , antiderivative size = 333, normalized size of antiderivative = 2.66

$$\int \cos^3(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \begin{cases} \frac{3Aax\sin^4(c+dx)}{8} + \frac{3Aax\sin^2(c+dx)\cos^2(c+dx)}{4} + \frac{3Aax\cos^4(c+dx)}{8} + \frac{3Aa\sin^3(c+dx)\cos(c+dx)}{8d} + \frac{2Aa\sin^3(c+dx)}{3d} + \frac{5Aa\sin(c+dx)}{3d} \\ x(A+B\cos(c))(a\cos(c)+a)\cos^3(c) \end{cases}$$

input `integrate(cos(d*x+c)**3*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

3.1. $\int \cos^3(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$

output `Piecewise((3*A*a*x*sin(c + d*x)**4/8 + 3*A*a*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*A*a*x*cos(c + d*x)**4/8 + 3*A*a*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 2*A*a*sin(c + d*x)**3/(3*d) + 5*A*a*sin(c + d*x)*cos(c + d*x)**3/(8*d) + A*a*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*a*x*sin(c + d*x)**4/8 + 3*B*a*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*B*a*x*cos(c + d*x)**4/8 + 8*B*a*sin(c + d*x)**5/(15*d) + 4*B*a*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 3*B*a*sin(c + d*x)**3*cos(c + d*x)/(8*d) + B*a*sin(c + d*x)*cos(c + d*x)**4/d + 5*B*a*sin(c + d*x)*cos(c + d*x)**3/(8*d), Ne(d, 0)), (x*(A + B*cos(c))* (a*cos(c) + a)*cos(c)**3, True))`

3.1.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.99

$$\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx =$$

$$\frac{160 (\sin(dx + c)^3 - 3 \sin(dx + c))Aa - 15(12dx + 12c + \sin(4dx + 4c) + 8 \sin(2dx + 2c))Aa - 32(3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c))B^2a - 15(12dx + 12c + \sin(4dx + 4c) + 8 \sin(2dx + 2c))B^2a}{d}$$

input `integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `-1/480*(160*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a - 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a - 32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a - 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a)/d`

3.1.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.90

$$\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3}{8} (Aa + Ba)x + \frac{Ba \sin(5dx + 5c)}{80d}$$

$$+ \frac{(Aa + Ba) \sin(4dx + 4c)}{32d} + \frac{(4Aa + 5Ba) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(Aa + Ba) \sin(2dx + 2c)}{4d} + \frac{(6Aa + 5Ba) \sin(dx + c)}{8d}$$

3.1. $\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

input `integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `3/8*(A*a + B*a)*x + 1/80*B*a*sin(5*d*x + 5*c)/d + 1/32*(A*a + B*a)*sin(4*d*x + 4*c)/d + 1/48*(4*A*a + 5*B*a)*sin(3*d*x + 3*c)/d + 1/4*(A*a + B*a)*sin(2*d*x + 2*c)/d + 1/8*(6*A*a + 5*B*a)*sin(d*x + c)/d`

3.1.9 Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.89

$$\int \cos^3(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{3Aa}{4} + \frac{3Ba}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{29Aa}{6} + \frac{13Ba}{6}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{20Aa}{3} + \frac{116Ba}{15}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{35Aa}{6} + \frac{19Ba}{6}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{3Aa}{4} + \frac{3Ba}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \frac{3a \operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}(A + B)}{4d} + \frac{3a \operatorname{atan}\left(\frac{3a \tan\left(\frac{c}{2} + \frac{dx}{2}\right)(A + B)}{4\left(\frac{3Aa}{4} + \frac{3Ba}{4}\right)}\right)(A + B)}{4d}$$

input `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`

output `(tan(c/2 + (d*x)/2)*((13*A*a)/4 + (13*B*a)/4) + tan(c/2 + (d*x)/2)^9*((3*A*a)/4 + (3*B*a)/4) + tan(c/2 + (d*x)/2)^7*((29*A*a)/6 + (13*B*a)/6) + tan(c/2 + (d*x)/2)^3*((35*A*a)/6 + (19*B*a)/6) + tan(c/2 + (d*x)/2)^5*((20*A*a)/3 + (116*B*a)/15))/(d*(5*tan(c/2 + (d*x)/2)^2 + 10*tan(c/2 + (d*x)/2)^4 + 10*tan(c/2 + (d*x)/2)^6 + 5*tan(c/2 + (d*x)/2)^8 + tan(c/2 + (d*x)/2)^10 + 1)) - (3*a*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2)*(A + B))/(4*d) + (3*a*a*tan((3*a*tan(c/2 + (d*x)/2)*(A + B))/(4*((3*A*a)/4 + (3*B*a)/4)))*(A + B))/(4*d)`

3.2 $\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.2.1	Optimal result	240
3.2.2	Mathematica [A] (verified)	240
3.2.3	Rubi [A] (verified)	241
3.2.4	Maple [A] (verified)	244
3.2.5	Fricas [A] (verification not implemented)	244
3.2.6	Sympy [B] (verification not implemented)	245
3.2.7	Maxima [A] (verification not implemented)	245
3.2.8	Giac [A] (verification not implemented)	246
3.2.9	Mupad [B] (verification not implemented)	246

3.2.1 Optimal result

Integrand size = 29, antiderivative size = 97

$$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{1}{8}a(4A + 3B)x + \frac{a(A + B) \sin(c + dx)}{d} + \frac{a(4A + 3B) \cos(c + dx) \sin(c + dx)}{8d}$$

$$+ \frac{aB \cos^3(c + dx) \sin(c + dx)}{4d} - \frac{a(A + B) \sin^3(c + dx)}{3d}$$

output $\frac{1}{8}a*(4*A+3*B)*x+a*(A+B)*\sin(d*x+c)/d+\frac{1}{8}a*(4*A+3*B)*\cos(d*x+c)*\sin(d*x+c)/d+\frac{1}{4}a*B*\cos(d*x+c)^3*\sin(d*x+c)/d-\frac{1}{3}a*(A+B)*\sin(d*x+c)^3/d$

3.2.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{a(48Ac + 36Bc + 48Adx + 36Bdx + 96(A + B) \sin(c + dx) - 32(A + B) \sin^3(c + dx) + 24(A + B) \sin(2c + 2dx))}{96d}$$

input `Integrate[Cos[c + d*x]^2*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output $(a*(48*A*c + 36*B*c + 48*A*d*x + 36*B*d*x + 96*(A + B)*\text{Sin}[c + d*x] - 32*(A + B)*\text{Sin}[c + d*x]^3 + 24*(A + B)*\text{Sin}[2*(c + d*x)] + 3*B*\text{Sin}[4*(c + d*x)])/(96*d)$

3.2.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.96, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \cos^2(c + dx) \left((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)\right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left((aA + aB) \sin\left(c + dx + \frac{\pi}{2}\right) + aA + aB \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{1}{4} \int \cos^2(c + dx) (a(4A + 3B) + 4a(A + B) \cos(c + dx)) dx + \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a(4A + 3B) + 4a(A + B) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\
 & \quad \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
 & \quad \downarrow \text{3227} \\
 & \frac{1}{4} \left(4a(A + B) \int \cos^3(c + dx) dx + a(4A + 3B) \int \cos^2(c + dx) dx\right) + \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.2. $\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \frac{1}{4} \left(a(4A + 3B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + 4a(A + B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
& \quad \downarrow \text{3113} \\
& \frac{1}{4} \left(a(4A + 3B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4a(A + B) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
& \quad \downarrow \text{2009} \\
& \frac{1}{4} \left(a(4A + 3B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4a(A + B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
& \quad \downarrow \text{3115} \\
& \frac{1}{4} \left(a(4A + 3B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{4a(A + B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d} \\
& \quad \downarrow \text{24} \\
& \frac{1}{4} \left(a(4A + 3B) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) - \frac{4a(A + B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^3(c + dx)}{4d}
\end{aligned}$$

input `Int[Cos[c + d*x]^2*(a + a*cos[c + d*x])*(A + B*cos[c + d*x]),x]`

output `(a*B*cos[c + d*x]^3*sin[c + d*x])/(4*d) + (a*(4*A + 3*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (4*a*(A + B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3)/d)/4`

3.2. $\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.2.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.2.4 Maple [A] (verified)

Time = 2.52 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.69

method	result
parallelrisch	$\frac{\left(\frac{(A+B)\sin(2dx+2c)}{2} + \frac{(A+B)\sin(3dx+3c)}{6} + \frac{\sin(4dx+4c)B}{16} + \frac{3(A+B)\sin(dx+c)}{2} + dx\left(A + \frac{3B}{4}\right)\right)a}{2d}$
parts	$\frac{(aA+Ba)(2+\cos^2(dx+c))\sin(dx+c)}{3d} + \frac{aA\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx+c}{2}\right)}{d} + \frac{Ba\left(\frac{\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}}{4}\right)\sin(dx+c)}{d}$
derivativedivides	$\frac{Ba\left(\frac{\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}}{4}\right)\sin(dx+c) + \frac{3dx + \frac{3c}{8}}{8}}{d} + \frac{aA(2+\cos^2(dx+c))\sin(dx+c)}{3} + \frac{Ba(2+\cos^2(dx+c))\sin(dx+c)}{3} + aA\left(\frac{\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}}{4}\right)\sin(dx+c)$
default	$\frac{Ba\left(\frac{\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}}{4}\right)\sin(dx+c) + \frac{3dx + \frac{3c}{8}}{8}}{d} + \frac{aA(2+\cos^2(dx+c))\sin(dx+c)}{3} + \frac{Ba(2+\cos^2(dx+c))\sin(dx+c)}{3} + aA\left(\frac{\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}}{4}\right)\sin(dx+c)$
risch	$\frac{axA}{2} + \frac{3aBx}{8} + \frac{3\sin(dx+c)aA}{4d} + \frac{3aB\sin(dx+c)}{4d} + \frac{\sin(4dx+4c)Ba}{32d} + \frac{\sin(3dx+3c)aA}{12d} + \frac{\sin(3dx+3c)Ba}{12d} +$
norman	$\frac{\frac{a(4A+3B)x}{8} + \frac{a(4A+3B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{4d} + \frac{a(4A+3B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2} + \frac{3a(4A+3B)x\left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{4} + \frac{a(4A+3B)x\left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2}}{(1+\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))}$

input `int(cos(d*x+c)^2*(a+cos(d*x+c))*a*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2*(1/2*(A+B)*sin(2*d*x+2*c)+1/6*(A+B)*sin(3*d*x+3*c)+1/16*sin(4*d*x+4*c)*B+3/2*(A+B)*sin(d*x+c)+d*x*(A+3/4*B))*a/d`

3.2.5 Fricas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.76

$$\int \cos^2(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{3(4A+3B)adx + (6Ba\cos(dx+c))^3 + 8(A+B)a\cos(dx+c)^2 + 3(4A+3B)a\cos(dx+c) + 16(A+B)a\sin(dx+c)}{24d}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))*a*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/24*(3*(4*A + 3*B)*a*d*x + (6*B*a*cos(d*x + c))^3 + 8*(A + B)*a*cos(d*x + c)^2 + 3*(4*A + 3*B)*a*cos(d*x + c) + 16*(A + B)*a)*sin(d*x + c)/d`

3.2. $\int \cos^2(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$

3.2.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(88) = 176.

Time = 0.18 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.60

$$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aax \sin^2(c+dx)}{2} + \frac{Aax \cos^2(c+dx)}{2} + \frac{2Aa \sin^3(c+dx)}{3d} + \frac{Aa \sin(c+dx) \cos^2(c+dx)}{d} + \frac{Aa \sin(c+dx) \cos(c+dx)}{2d} + \frac{3Bax \sin^4(c+dx)}{8} \\ x(A + B \cos(c))(a \cos(c) + a) \cos^2(c) \end{cases}$$

input `integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a*x*sin(c + d*x)**2/2 + A*a*x*cos(c + d*x)**2/2 + 2*A*a*sin(c + d*x)**3/(3*d) + A*a*sin(c + d*x)*cos(c + d*x)**2/d + A*a*sin(c + d*x)*cos(c + d*x)/(2*d) + 3*B*a*x*sin(c + d*x)**4/8 + 3*B*a*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*B*a*x*cos(c + d*x)**4/8 + 3*B*a*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 2*B*a*sin(c + d*x)**3/(3*d) + 5*B*a*sin(c + d*x)*cos(c + d*x)**3/(8*d) + B*a*sin(c + d*x)*cos(c + d*x)**2/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)*cos(c)**2, True))`

3.2.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.04

$$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx =$$

$$\frac{32 (\sin(dx + c)^3 - 3 \sin(dx + c))Aa - 24 (2dx + 2c + \sin(2dx + 2c))Aa + 32 (\sin(dx + c)^3 - 3 \sin(dx + c))Ba - 24 (2dx + 2c + \sin(2dx + 2c))Ba + 32 (\sin(dx + c)^3 - 3 \sin(dx + c))Aa - 24 (2dx + 2c + \sin(2dx + 2c))Aa + 32 (\sin(dx + c)^3 - 3 \sin(dx + c))Ba - 24 (2dx + 2c + \sin(2dx + 2c))Ba}{96d}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `-1/96*(32*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a - 24*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a + 32*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a - 3*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a)/d`

3.2.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.92

$$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{1}{8} (4Aa + 3Ba)x + \frac{Ba \sin(4dx + 4c)}{32d} + \frac{(Aa + Ba) \sin(3dx + 3c)}{12d}$$

$$+ \frac{(Aa + Ba) \sin(2dx + 2c)}{4d} + \frac{3(Aa + Ba) \sin(dx + c)}{4d}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/8*(4*A*a + 3*B*a)*x + 1/32*B*a*sin(4*d*x + 4*c)/d + 1/12*(A*a + B*a)*sin(3*d*x + 3*c)/d + 1/4*(A*a + B*a)*sin(2*d*x + 2*c)/d + 3/4*(A*a + B*a)*sin(d*x + c)/d`

3.2.9 Mupad [B] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 212, normalized size of antiderivative = 2.19

$$\int \cos^2(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{(Aa + \frac{3Ba}{4}) \tan(\frac{c}{2} + \frac{dx}{2})^7 + (\frac{7Aa}{3} + \frac{49Ba}{12}) \tan(\frac{c}{2} + \frac{dx}{2})^5 + (\frac{13Aa}{3} + \frac{31Ba}{12}) \tan(\frac{c}{2} + \frac{dx}{2})^3 + (3Aa + \frac{13Ba}{4}) \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(\tan(\frac{c}{2} + \frac{dx}{2})^8 + 4 \tan(\frac{c}{2} + \frac{dx}{2})^6 + 6 \tan(\frac{c}{2} + \frac{dx}{2})^4 + 4 \tan(\frac{c}{2} + \frac{dx}{2})^2 + 1 \right)}$$

$$+ \frac{a \operatorname{atan}\left(\frac{a \tan(\frac{c}{2} + \frac{dx}{2}) (4A + 3B)}{4(Aa + \frac{3Ba}{4})}\right) (4A + 3B)}{4d} - \frac{a(4A + 3B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{4d}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`

output `(tan(c/2 + (d*x)/2)*(3*A*a + (13*B*a)/4) + tan(c/2 + (d*x)/2)^7*(A*a + (3*B*a)/4) + tan(c/2 + (d*x)/2)^3*((13*A*a)/3 + (31*B*a)/12) + tan(c/2 + (d*x)/2)^5*((7*A*a)/3 + (49*B*a)/12))/(d*(4*tan(c/2 + (d*x)/2)^2 + 6*tan(c/2 + (d*x)/2)^4 + 4*tan(c/2 + (d*x)/2)^6 + tan(c/2 + (d*x)/2)^8 + 1)) + (a*atan((a*tan(c/2 + (d*x)/2)*(4*A + 3*B))/(4*(A*a + (3*B*a)/4)))*(4*A + 3*B))/(4*d) - (a*(4*A + 3*B)*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(4*d)`

3.3 $\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.3.1	Optimal result	247
3.3.2	Mathematica [A] (verified)	247
3.3.3	Rubi [A] (verified)	248
3.3.4	Maple [A] (verified)	250
3.3.5	Fricas [A] (verification not implemented)	250
3.3.6	Sympy [B] (verification not implemented)	251
3.3.7	Maxima [A] (verification not implemented)	251
3.3.8	Giac [A] (verification not implemented)	252
3.3.9	Mupad [B] (verification not implemented)	252

3.3.1 Optimal result

Integrand size = 27, antiderivative size = 77

$$\begin{aligned} & \int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \frac{1}{2}a(A + B)x + \frac{a(3A + 2B) \sin(c + dx)}{3d} \\ & \quad + \frac{a(A + B) \cos(c + dx) \sin(c + dx)}{2d} + \frac{aB \cos^2(c + dx) \sin(c + dx)}{3d} \end{aligned}$$

output `1/2*a*(A+B)*x+1/3*a*(3*A+2*B)*sin(d*x+c)/d+1/2*a*(A+B)*cos(d*x+c)*sin(d*x+c)/d+1/3*a*B*cos(d*x+c)^2*sin(d*x+c)/d`

3.3.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.84

$$\begin{aligned} & \int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \frac{a(6Ac + 6Bc + 6Adx + 6Bdx + 3(4A + 3B) \sin(c + dx) + 3(A + B) \sin(2(c + dx)) + B \sin(3(c + dx)))}{12d} \end{aligned}$$

input `Integrate[Cos[c + d*x]*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output $(a*(6*A*c + 6*B*c + 6*A*d*x + 6*B*d*x + 3*(4*A + 3*B)*\text{Sin}[c + d*x] + 3*(A + B)*\text{Sin}[2*(c + d*x)] + B*\text{Sin}[3*(c + d*x)])/(12*d)$

3.3.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 3447, 3042, 3502, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \cos(c + dx) \left((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)\right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left((aA + aB) \sin\left(c + dx + \frac{\pi}{2}\right) + aA + aB \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{1}{3} \int \cos(c + dx) \left(a(3A + 2B) + 3a(A + B) \cos(c + dx)\right) dx + \frac{aB \sin(c + dx) \cos^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a(3A + 2B) + 3a(A + B) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\
 & \quad \frac{aB \sin(c + dx) \cos^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3213} \\
 & \frac{1}{3} \left(\frac{a(3A + 2B) \sin(c + dx)}{d} + \frac{3a(A + B) \sin(c + dx) \cos(c + dx)}{2d} + \frac{3}{2} ax(A + B) \right) + \\
 & \quad \frac{aB \sin(c + dx) \cos^2(c + dx)}{3d}
 \end{aligned}$$

3.3. $\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

input `Int[Cos[c + d*x]*(a + a*cos[c + d*x])*(A + B*cos[c + d*x]),x]`

output `(a*B*cos[c + d*x]^2*sin[c + d*x])/(3*d) + ((3*a*(A + B)*x)/2 + (a*(3*A + 2*B)*sin[c + d*x])/d + (3*a*(A + B)*cos[c + d*x]*sin[c + d*x])/(2*d))/3`

3.3.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Int[(a + b*sin[e + f*x])^m*(A*c + (B*c + A*d)*sin[e + f*x] + B*d*sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.3.4 Maple [A] (verified)

Time = 1.81 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.70

method	result
parallelrisch	$\frac{a\left(\frac{(A+B)\sin(2dx+2c)}{2} + \frac{B\sin(3dx+3c)}{6}\right) + \left(2A + \frac{3B}{2}\right)\sin(dx+c) + (A+B)xd}{2d}$
parts	$\frac{(aA+Ba)\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx+c}{2}\right)}{d} + \frac{\sin(dx+c)aA}{d} + \frac{Ba(2+\cos^2(dx+c))\sin(dx+c)}{3d}$
derivativedivides	$\frac{\frac{Ba(2+\cos^2(dx+c))\sin(dx+c)}{3} + aA\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx+c}{2}\right) + Ba\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx+c}{2}\right) + aA\sin(dx+c)}{d}$
default	$\frac{\frac{Ba(2+\cos^2(dx+c))\sin(dx+c)}{3} + aA\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx+c}{2}\right) + Ba\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx+c}{2}\right) + aA\sin(dx+c)}{d}$
risch	$\frac{aAx}{2} + \frac{aBx}{2} + \frac{\sin(dx+c)aA}{d} + \frac{3aB\sin(dx+c)}{4d} + \frac{\sin(3dx+3c)Ba}{12d} + \frac{\sin(2dx+2c)aA}{4d} + \frac{\sin(2dx+2c)Ba}{4d}$
norman	$\frac{\frac{a(A+B)\left(\tan^5\left(\frac{dx+c}{2}\right)\right)}{d} + \frac{a(A+B)x}{2} + \frac{3a(A+B)\tan\left(\frac{dx+c}{2}\right)}{d} + \frac{3a(A+B)x\left(\tan^2\left(\frac{dx+c}{2}\right)\right)}{2} + \frac{3a(A+B)x\left(\tan^4\left(\frac{dx+c}{2}\right)\right)}{2} + \frac{a(A+B)}{\left(1+\tan^2\left(\frac{dx+c}{2}\right)\right)^3}}{d}$

input `int(cos(d*x+c)*(a+cos(d*x+c))*a*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2*a*(1/2*(A+B)*sin(2*d*x+2*c)+1/6*B*sin(3*d*x+3*c)+(2*A+3/2*B)*sin(d*x+c)+(A+B)*x*d)/d`

3.3.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.73

$$\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3(A + B)adx + (2Ba \cos(dx + c))^2 + 3(A + B)a \cos(dx + c) + 2(3A + 2B)a \sin(dx + c)}{6d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))*a*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/6*(3*(A + B)*a*d*x + (2*B*a*cos(d*x + c))^2 + 3*(A + B)*a*cos(d*x + c) + 2*(3*A + 2*B)*a)*sin(d*x + c)/d`

3.3.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(70) = 140$.

Time = 0.13 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.18

$$\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aax \sin^2(c+dx)}{2} + \frac{Aax \cos^2(c+dx)}{2} + \frac{Aa \sin(c+dx) \cos(c+dx)}{2d} + \frac{Aa \sin(c+dx)}{d} + \frac{Bax \sin^2(c+dx)}{2} + \frac{Bax \cos^2(c+dx)}{2} + \frac{2Ba \sin^3(c+dx)}{3} \\ x(A + B \cos(c))(a \cos(c) + a) \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a*x*sin(c + d*x)**2/2 + A*a*x*cos(c + d*x)**2/2 + A*a*sin(c + d*x)*cos(c + d*x)/(2*d) + A*a*sin(c + d*x)/d + B*a*x*sin(c + d*x)**2/2 + B*a*x*cos(c + d*x)**2/2 + 2*B*a*sin(c + d*x)**3/(3*d) + B*a*sin(c + d*x)*cos(c + d*x)**2/d + B*a*sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)*cos(c), True))`

3.3.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.03

$$\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3(2dx + 2c + \sin(2dx + 2c))Aa - 4(\sin(dx + c)^3 - 3\sin(dx + c))Ba + 3(2dx + 2c + \sin(2dx + 2c))Aa}{12d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/12*(3*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a - 4*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a + 3*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a + 12*A*a*sin(d*x + c))/d`

3.3.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.88

$$\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{1}{2} (Aa + Ba)x + \frac{Ba \sin(3dx + 3c)}{12d}$$

$$+ \frac{(Aa + Ba) \sin(2dx + 2c)}{4d} + \frac{(4Aa + 3Ba) \sin(dx + c)}{4d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/2*(A*a + B*a)*x + 1/12*B*a*sin(3*d*x + 3*c)/d + 1/4*(A*a + B*a)*sin(2*d*x + 2*c)/d + 1/4*(4*A*a + 3*B*a)*sin(d*x + c)/d`

3.3.9 Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.09

$$\int \cos(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{Aax}{2} + \frac{Bax}{2} + \frac{Aa \sin(c + dx)}{d} + \frac{3Ba \sin(c + dx)}{4d}$$

$$+ \frac{Aa \sin(2c + 2dx)}{4d} + \frac{Ba \sin(2c + 2dx)}{4d} + \frac{Ba \sin(3c + 3dx)}{12d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`

output `(A*a*x)/2 + (B*a*x)/2 + (A*a*sin(c + d*x))/d + (3*B*a*sin(c + d*x))/(4*d) + (A*a*sin(2*c + 2*d*x))/(4*d) + (B*a*sin(2*c + 2*d*x))/(4*d) + (B*a*sin(3*c + 3*d*x))/(12*d)`

3.4 $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.4.1	Optimal result	253
3.4.2	Mathematica [A] (verified)	253
3.4.3	Rubi [A] (verified)	254
3.4.4	Maple [A] (verified)	255
3.4.5	Fricas [A] (verification not implemented)	255
3.4.6	Sympy [B] (verification not implemented)	256
3.4.7	Maxima [A] (verification not implemented)	256
3.4.8	Giac [A] (verification not implemented)	257
3.4.9	Mupad [B] (verification not implemented)	257

3.4.1 Optimal result

Integrand size = 21, antiderivative size = 47

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx = \frac{1}{2}a(2A + B)x + \frac{a(A + B) \sin(c + dx)}{d} + \frac{aB \cos(c + dx) \sin(c + dx)}{2d}$$

```
output 1/2*a*(2*A+B)*x+a*(A+B)*sin(d*x+c)/d+1/2*a*B*cos(d*x+c)*sin(d*x+c)/d
```

3.4.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.94

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx = \frac{a(2Bc + 4Adx + 2Bdx + 4(A + B) \sin(c + dx) + B \sin(2(c + dx)))}{4d}$$

```
input Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]
```

```
output (a*(2*B*c + 4*A*d*x + 2*B*d*x + 4*(A + B)*Sin[c + d*x] + B*Sine[2*(c + d*x)]))/(4*d)
```

3.4.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \cos(c + dx) + a)(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx$$

$$\downarrow \text{3213}$$

$$\frac{a(A + B) \sin(c + dx)}{d} + \frac{1}{2}ax(2A + B) + \frac{aB \sin(c + dx) \cos(c + dx)}{2d}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(a*(2*A + B)*x)/2 + (a*(A + B)*Sin[c + d*x])/d + (a*B*Cos[c + d*x]*Sin[c + d*x])/(2*d)`

3.4.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

3.4.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.79

method	result
parallelrisc	$\frac{\left(\frac{\sin(2dx+2c)B}{4} + (A+B)\sin(dx+c) + dx\left(A + \frac{B}{2}\right)\right)a}{d}$
risc	$axA + \frac{aBx}{2} + \frac{\sin(dx+c)aA}{d} + \frac{aB\sin(dx+c)}{d} + \frac{\sin(2dx+2c)Ba}{4d}$
parts	$axA + \frac{(aA+Ba)\sin(dx+c)}{d} + \frac{Ba\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
derivativedivides	$\frac{Ba\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + aA\sin(dx+c) + Ba\sin(dx+c) + aA(dx+c)}{d}$
default	$\frac{Ba\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + aA\sin(dx+c) + Ba\sin(dx+c) + aA(dx+c)}{d}$
norman	$\frac{\frac{a(2A+B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d} + a(2A+B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \frac{a(2A+3B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{a(2A+B)x}{2} + \frac{a(2A+B)x\left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$

input `int((a+cos(d*x+c))*a)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `(1/4*sin(2*d*x+2*c)*B+(A+B)*sin(d*x+c)+d*x*(A+1/2*B))*a/d`

3.4.5 Fracas [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{(2A + B)adx + (Ba \cos(dx + c) + 2(A + B)a) \sin(dx + c)}{2d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/2*((2*A + B)*a*d*x + (B*a*cos(d*x + c) + 2*(A + B)*a)*sin(d*x + c))/d`

3.4.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(42) = 84$.

Time = 0.09 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.00

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \begin{cases} Aax + \frac{Aa \sin(c+dx)}{d} + \frac{Bax \sin^2(c+dx)}{2} + \frac{Bax \cos^2(c+dx)}{2} + \frac{Ba \sin(c+dx) \cos(c+dx)}{2d} + \frac{Ba \sin(c+dx)}{d} & \text{for } d \neq 0 \\ x(A + B \cos(c))(a \cos(c) + a) & \text{otherwise} \end{cases}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a*x + A*a*sin(c + d*x)/d + B*a*x*sin(c + d*x)**2/2 + B*a*x*cos(c + d*x)**2/2 + B*a*sin(c + d*x)*cos(c + d*x)/(2*d) + B*a*sin(c + d*x)/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a), True))`

3.4.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.17

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{4(dx + c)Aa + (2dx + 2c + \sin(2dx + 2c))Ba + 4Aa \sin(dx + c) + 4Ba \sin(dx + c)}{4d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/4*(4*(d*x + c)*A*a + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*a + 4*A*a*sin(d*x + c) + 4*B*a*sin(d*x + c))/d`

3.4.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx = \frac{1}{2} (2 A a + B a) x + \frac{B a \sin(2 d x + 2 c)}{4 d} + \frac{(A a + B a) \sin(d x + c)}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/2*(2*A*a + B*a)*x + 1/4*B*a*sin(2*d*x + 2*c)/d + (A*a + B*a)*sin(d*x + c)/d`

3.4.9 Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.06

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) dx = A a x + \frac{B a x}{2} + \frac{A a \sin(c + d x)}{d} + \frac{B a \sin(c + d x)}{d} + \frac{B a \sin(2 c + 2 d x)}{4 d}$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`

output `A*a*x + (B*a*x)/2 + (A*a*sin(c + d*x))/d + (B*a*sin(c + d*x))/d + (B*a*sin(2*c + 2*d*x))/(4*d)`

3.5 $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$

3.5.1	Optimal result	258
3.5.2	Mathematica [A] (verified)	258
3.5.3	Rubi [A] (verified)	259
3.5.4	Maple [A] (verified)	261
3.5.5	Fricas [A] (verification not implemented)	261
3.5.6	Sympy [F]	262
3.5.7	Maxima [A] (verification not implemented)	262
3.5.8	Giac [B] (verification not implemented)	263
3.5.9	Mupad [B] (verification not implemented)	263

3.5.1 Optimal result

Integrand size = 27, antiderivative size = 32

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx \\ &= a(A + B)x + \frac{aA \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{aB \sin(c + dx)}{d} \end{aligned}$$

output `a*(A+B)*x+a*A*arctanh(sin(d*x+c))/d+a*B*sin(d*x+c)/d`

3.5.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.44

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx \\ &= aAx + aBx + \frac{aA \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{aB \cos(dx) \sin(c)}{d} + \frac{aB \cos(c) \sin(dx)}{d} \end{aligned}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `a*A*x + a*B*x + (a*A*ArcTanh[Sin[c + d*x]])/d + (a*B*Cos[d*x]*Sin[c])/d + (a*B*Cos[c]*Sin[d*x])/d`

3.5. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$

3.5.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c+dx)(a \cos(c+dx) + a)(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c+dx + \frac{\pi}{2}) + a)(A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec(c+dx)((aA + aB) \cos(c+dx) + aA + aB \cos^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aA + aB) \sin(c+dx + \frac{\pi}{2}) + aA + aB \sin(c+dx + \frac{\pi}{2})^2}{\sin(c+dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3502} \\
 & \int (aA + a(A+B) \cos(c+dx)) \sec(c+dx) dx + \frac{aB \sin(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA + a(A+B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})} dx + \frac{aB \sin(c+dx)}{d} \\
 & \quad \downarrow \text{3214} \\
 & aA \int \sec(c+dx) dx + ax(A+B) + \frac{aB \sin(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & aA \int \csc(c+dx + \frac{\pi}{2}) dx + ax(A+B) + \frac{aB \sin(c+dx)}{d} \\
 & \quad \downarrow \text{4257} \\
 & \frac{aA \operatorname{Arctanh}(\sin(c+dx))}{d} + ax(A+B) + \frac{aB \sin(c+dx)}{d}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `a*(A + B)*x + (a*A*ArcTanh[Sin[c + d*x]])/d + (a*B*Sin[c + d*x])/d`

3.5.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.5.4 Maple [A] (verified)

Time = 1.48 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.50

method	result
derivativdivides	$\frac{aA(dx+c)+Ba \sin(dx+c)+aA \ln(\sec(dx+c)+\tan(dx+c))+Ba(dx+c)}{d}$
default	$\frac{aA(dx+c)+Ba \sin(dx+c)+aA \ln(\sec(dx+c)+\tan(dx+c))+Ba(dx+c)}{d}$
parallelrisc	$\frac{(-A \ln(\tan(\frac{dx}{2}+\frac{c}{2}))-1)+A \ln(\tan(\frac{dx}{2}+\frac{c}{2}))+1)+B \sin(dx+c)+(A+B)xd}{d}a$
parts	$\frac{aA \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{(aA+Ba)(dx+c)}{d} + \frac{aB \sin(dx+c)}{d}$
risc	$aAx + aBx - \frac{iBa e^{i(dx+c)}}{2d} + \frac{iBa e^{-i(dx+c)}}{2d} + \frac{aA \ln(e^{i(dx+c)}+i)}{d} - \frac{aA \ln(e^{i(dx+c)}-i)}{d}$
norman	$\frac{(aA+Ba)x+(aA+Ba)x(\tan^4(\frac{dx}{2}+\frac{c}{2}))+(2aA+2Ba)x(\tan^2(\frac{dx}{2}+\frac{c}{2}))+\frac{2Ba \tan(\frac{dx}{2}+\frac{c}{2})}{d}+\frac{2Ba(\tan^3(\frac{dx}{2}+\frac{c}{2}))}{d}}{(1+\tan^2(\frac{dx}{2}+\frac{c}{2}))^2} + \dots$

input `int((a+cos(d*x+c))*a)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `1/d*(a*A*(d*x+c)+B*a*sin(d*x+c)+a*A*ln(sec(d*x+c)+tan(d*x+c))+B*a*(d*x+c))`

3.5.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2(A + B)adx + Aa \log(\sin(dx + c) + 1) - Aa \log(-\sin(dx + c) + 1) + 2Ba \sin(dx + c)}{2d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fracas")`

output `1/2*(2*(A + B)*a*d*x + A*a*log(sin(d*x + c) + 1) - A*a*log(-sin(d*x + c) + 1) + 2*B*a*sin(d*x + c))/d`

3.5.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx \\ &= a \left(\int A \sec(c + dx) dx + \int A \cos(c + dx) \sec(c + dx) dx \right. \\ & \quad \left. + \int B \cos(c + dx) \sec(c + dx) dx + \int B \cos^2(c + dx) \sec(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `a*(Integral(A*sec(c + d*x), x) + Integral(A*cos(c + d*x)*sec(c + d*x), x) + Integral(B*cos(c + d*x)*sec(c + d*x), x) + Integral(B*cos(c + d*x)**2*sec(c + d*x), x))`

3.5.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.47

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{(dx + c)Aa + (dx + c)Ba + Aa \log(\sec(dx + c) + \tan(dx + c)) + Ba \sin(dx + c)}{d} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `((d*x + c)*A*a + (d*x + c)*B*a + A*a*log(sec(d*x + c) + tan(d*x + c)) + B*a*sin(d*x + c))/d`

3.5.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(32) = 64$.

Time = 0.30 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.47

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{Aa \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - Aa \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) + (Aa + Ba)(dx + c) + \frac{2Ba \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1}}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `(A*a*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - A*a*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + (A*a + B*a)*(d*x + c) + 2*B*a*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 1))/d`

3.5.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 100, normalized size of antiderivative = 3.12

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{Ba \sin(c + dx)}{d} + \frac{2Aa \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{2Aa \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2Ba \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x),x)`

output `(B*a*sin(c + d*x))/d + (2*A*a*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*A*a*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d`

3.6 $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.6.1	Optimal result	264
3.6.2	Mathematica [A] (verified)	264
3.6.3	Rubi [A] (verified)	265
3.6.4	Maple [A] (verified)	267
3.6.5	Fricas [B] (verification not implemented)	267
3.6.6	Sympy [F]	268
3.6.7	Maxima [B] (verification not implemented)	268
3.6.8	Giac [B] (verification not implemented)	269
3.6.9	Mupad [B] (verification not implemented)	269

3.6.1 Optimal result

Integrand size = 29, antiderivative size = 32

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= aBx + \frac{a(A + B)\operatorname{arctanh}(\sin(c + dx))}{d} + \frac{aA \tan(c + dx)}{d}$$

output `a*B*x+a*(A+B)*arctanh(sin(d*x+c))/d+a*A*tan(d*x+c)/d`

3.6.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.34

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= aBx + \frac{aA \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{aB \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{aA \tan(c + dx)}{d}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `a*B*x + (a*A*ArcTanh[Sin[c + d*x]])/d + (a*B*ArcTanh[Sin[c + d*x]])/d + (a*A*Tan[c + d*x])/d`

3.6. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.6.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 3447, 3042, 3500, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx)(a \cos(c+dx) + a)(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c+dx + \frac{\pi}{2}) + a)(A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec^2(c+dx)((aA + aB) \cos(c+dx) + aA + aB \cos^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aA + aB) \sin(c+dx + \frac{\pi}{2}) + aA + aB \sin(c+dx + \frac{\pi}{2})^2}{\sin(c+dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3500} \\
 & \int (a(A+B) + aB \cos(c+dx)) \sec(c+dx) dx + \frac{aA \tan(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a(A+B) + aB \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})} dx + \frac{aA \tan(c+dx)}{d} \\
 & \quad \downarrow \text{3214} \\
 & a(A+B) \int \sec(c+dx) dx + \frac{aA \tan(c+dx)}{d} + aBx \\
 & \quad \downarrow \text{3042} \\
 & a(A+B) \int \csc(c+dx + \frac{\pi}{2}) dx + \frac{aA \tan(c+dx)}{d} + aBx \\
 & \quad \downarrow \text{4257} \\
 & \frac{a(A+B) \operatorname{arctanh}(\sin(c+dx))}{d} + \frac{aA \tan(c+dx)}{d} + aBx
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `a*B*x + (a*(A + B)*ArcTanh[Sin[c + d*x]])/d + (a*A*Tan[c + d*x])/d`

3.6.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.6.4 Maple [A] (verified)

Time = 2.33 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.56

method	result
parts	$\frac{aA \tan(dx+c)}{d} + \frac{(aA+Ba) \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{Ba(dx+c)}{d}$
derivativedivides	$\frac{aA \ln(\sec(dx+c)+\tan(dx+c))+Ba(dx+c)+aA \tan(dx+c)+Ba \ln(\sec(dx+c)+\tan(dx+c))}{d}$
default	$\frac{aA \ln(\sec(dx+c)+\tan(dx+c))+Ba(dx+c)+aA \tan(dx+c)+Ba \ln(\sec(dx+c)+\tan(dx+c))}{d}$
parallelrisc	$-\frac{\left((A+B) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)-(A+B) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)-dx B \cos(dx+c)-A \sin(dx+c)\right) a}{d \cos(dx+c)}$
risc	$aBx + \frac{2iaA}{d(e^{2i(dx+c)}+1)} + \frac{aA \ln(e^{i(dx+c)}+i)}{d} + \frac{a \ln(e^{i(dx+c)}+i)B}{d} - \frac{aA \ln(e^{i(dx+c)}-i)}{d} - \frac{a \ln(e^{i(dx+c)}-i)B}{d}$
norman	$\frac{aBx \left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+aBx \left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-aBx - \frac{2aA \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} - \frac{4aA \left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d} - \frac{2aA \left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d} - aBx \left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2 \left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}$

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBOSE)
```

```
output a*A*tan(d*x+c)/d+(A*a+B*a)/d*ln(sec(d*x+c)+tan(d*x+c))+B*a/d*(d*x+c)
```

3.6.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. 2(32) = 64.

Time = 0.39 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.47

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2 B a d x \cos(dx + c) + (A + B) a \cos(dx + c) \log(\sin(dx + c) + 1) - (A + B) a \cos(dx + c) \log(-\sin(dx + c))}{2 d \cos(dx + c)}$$

```
input integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fracas")
```

```
output 1/2*(2*B*a*d*x*cos(d*x + c) + (A + B)*a*cos(d*x + c)*log(sin(d*x + c) + 1) - (A + B)*a*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*A*a*sin(d*x + c))/(d*cos(d*x + c))
```

3.6.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= a \left(\int A \sec^2(c + dx) dx + \int A \cos(c + dx) \sec^2(c + dx) dx \right. \\ & \quad \left. + \int B \cos(c + dx) \sec^2(c + dx) dx + \int B \cos^2(c + dx) \sec^2(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `a*(Integral(A*sec(c + d*x)**2, x) + Integral(A*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)**2*sec(c + d*x)**2, x))`

3.6.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(32) = 64$.

Time = 0.21 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.28

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= \frac{2(dx + c)Ba + Aa(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + Ba(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1))}{2d} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*B*a + A*a*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + B*a*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*A*a*tan(d*x + c))/d`

3.6.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(32) = 64$.

Time = 0.31 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.62

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{(dx + c)Ba + (Aa + Ba) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - (Aa + Ba) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2Aa \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1}}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `((d*x + c)*B*a + (A*a + B*a)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (A*a + B*a)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*A*a*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 1))/d`

3.6.9 Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 100, normalized size of antiderivative = 3.12

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{Aa \tan(c + dx)}{d} + \frac{2Aa \operatorname{atanh} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d}$$

$$+ \frac{2Ba \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} + \frac{2Ba \operatorname{atanh} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^2,x)`

output `(A*a*tan(c + d*x))/d + (2*A*a*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d`

3.7 $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.7.1	Optimal result	270
3.7.2	Mathematica [A] (verified)	270
3.7.3	Rubi [A] (verified)	271
3.7.4	Maple [A] (verified)	273
3.7.5	Fricas [A] (verification not implemented)	274
3.7.6	Sympy [F]	274
3.7.7	Maxima [A] (verification not implemented)	275
3.7.8	Giac [B] (verification not implemented)	275
3.7.9	Mupad [B] (verification not implemented)	276

3.7.1 Optimal result

Integrand size = 29, antiderivative size = 56

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{a(A + 2B)\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{a(A + B)\tan(c + dx)}{d} + \frac{aA \sec(c + dx)\tan(c + dx)}{2d}$$

output `1/2*a*(A+2*B)*arctanh(sin(d*x+c))/d+a*(A+B)*tan(d*x+c)/d+1/2*a*A*sec(d*x+c)*tan(d*x+c)/d`

3.7.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.34

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{aA\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{aB\operatorname{arctanh}(\sin(c + dx))}{d}$$

$$+ \frac{aA \tan(c + dx)}{d} + \frac{aB \tan(c + dx)}{d} + \frac{aA \sec(c + dx)\tan(c + dx)}{2d}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output $(a*A*ArcTanh[Sin[c + d*x]])/(2*d) + (a*B*ArcTanh[Sin[c + d*x]])/d + (a*A*Tan[c + d*x])/d + (a*B*Tan[c + d*x])/d + (a*A*Sec[c + d*x]*Tan[c + d*x])/(2*d)$

3.7.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec^3(c + dx)((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{1}{2} \int (2a(A + B) + a(A + 2B) \cos(c + dx)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{2a(A + B) + a(A + 2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow \text{3227} \\
 & \frac{1}{2} \left(2a(A + B) \int \sec^2(c + dx) dx + a(A + 2B) \int \sec(c + dx) dx \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(a(A+2B) \int \csc \left(c+dx+\frac{\pi}{2} \right) dx + 2a(A+B) \int \csc \left(c+dx+\frac{\pi}{2} \right)^2 dx \right) + \\
& \quad \frac{aA \tan(c+dx) \sec(c+dx)}{2d} \\
& \quad \downarrow \text{4254} \\
& \frac{1}{2} \left(a(A+2B) \int \csc \left(c+dx+\frac{\pi}{2} \right) dx - \frac{2a(A+B) \int 1d(-\tan(c+dx))}{d} \right) + \\
& \quad \frac{aA \tan(c+dx) \sec(c+dx)}{2d} \\
& \quad \downarrow \text{24} \\
& \frac{1}{2} \left(a(A+2B) \int \csc \left(c+dx+\frac{\pi}{2} \right) dx + \frac{2a(A+B) \tan(c+dx)}{d} \right) + \frac{aA \tan(c+dx) \sec(c+dx)}{2d} \\
& \quad \downarrow \text{4257} \\
& \frac{1}{2} \left(\frac{a(A+2B) \operatorname{arctanh}(\sin(c+dx))}{d} + \frac{2a(A+B) \tan(c+dx)}{d} \right) + \frac{aA \tan(c+dx) \sec(c+dx)}{2d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((a*(A + 2*B)*ArcTanh[Sin[c + d*x]])/d + (2*a*(A + B)*Tan[c + d*x])/d)/2`

3.7.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.7.4 Maple [A] (verified)

Time = 2.92 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.34

method	result
derivativedivides	$\frac{aA \tan(dx+c) + Ba \ln(\sec(dx+c) + \tan(dx+c)) + aA \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + Ba \tan(dx+c)}{d}$
default	$\frac{aA \tan(dx+c) + Ba \ln(\sec(dx+c) + \tan(dx+c)) + aA \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + Ba \tan(dx+c)}{d}$
parts	$\frac{aA \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d} + \frac{(aA + Ba) \tan(dx+c)}{d} + \frac{Ba \ln(\sec(dx+c) + \tan(dx+c))}{d}$
parallelrisch	$-\frac{\left((1 + \cos(2dx+2c))(A+2B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - (1 + \cos(2dx+2c))(A+2B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + (-2B - 2A) \sin(2c) \right)}{2d(1 + \cos(2dx+2c))}$
risch	$-\frac{ia(Ae^{3i(dx+c)} - 2Ae^{2i(dx+c)} - 2Be^{2i(dx+c)} - Ae^{i(dx+c)} - 2A - 2B)}{d(e^{2i(dx+c)} + 1)^2} + \frac{aA \ln(e^{i(dx+c)} + i)}{2d} + \frac{a \ln(e^{i(dx+c)} + i)B}{d}$
norman	$\frac{a(A-2B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{a(3A+2B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{a(5A+2B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{d} - \frac{a(A+2B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{d}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{a(A+2B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d}$

3.7. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$


```
input int((a*cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(a*A*tan(d*x+c)+B*a*ln(sec(d*x+c)+tan(d*x+c))+a*A*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c))))+B*a*tan(d*x+c)
```

3.7.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.59

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(A + 2B)a \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (A + 2B)a \cos(dx + c)^2 \log(-\sin(dx + c) + 1) + 2(2(A + B)a \cos(dx + c) + Aa) \sin(dx + c)}{4d \cos(dx + c)^2}$$

```
input integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fricas")
```

```
output 1/4*((A + 2*B)*a*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A + 2*B)*a*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(2*(A + B)*a*cos(d*x + c) + A*a)*sin(d*x + c))/(d*cos(d*x + c)^2)
```

3.7.6 Sympy [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= a \left(\int A \sec^3(c + dx) dx + \int A \cos(c + dx) \sec^3(c + dx) dx \right. \\ \left. + \int B \cos(c + dx) \sec^3(c + dx) dx + \int B \cos^2(c + dx) \sec^3(c + dx) dx \right)$$

```
input integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)
```

```
output a*(Integral(A*sec(c + d*x)**3, x) + Integral(A*cos(c + d*x)*sec(c + d*x)**3, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**3, x) + Integral(B*cos(c + d*x)**2*sec(c + d*x)**3, x))
```

3.7. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.7.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.70

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx =$$

$$\frac{Aa \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1) \right) - 2Ba(\log(\sin(dx+c) + 1) - \log(\sin(dx+c) - 1)) - 4Aa \tan(dx+c) - 4Ba \tan(dx+c)}{4d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `-1/4*(A*a*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 2*B*a*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) - 4*A*a*tan(d*x + c) - 4*B*a*tan(d*x + c))/d`

3.7.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(52) = 104.

Time = 0.31 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.21

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(Aa + 2Ba) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - (Aa + 2Ba) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2 \left(Aa \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right)^3 + 2Ba \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{2d}}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output `1/2*((A*a + 2*B*a)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (A*a + 2*B*a)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(A*a*tan(1/2*d*x + 1/2*c)^3 + 2*B*a*tan(1/2*d*x + 1/2*c)^3 - 3*A*a*tan(1/2*d*x + 1/2*c) - 2*B*a*tan(1/2*d*x + 1/2*c)))/(tan(1/2*d*x + 1/2*c)^2 - 1)^2/d`

3.7.9 Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.68

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (3Aa + 2Ba) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (Aa + 2Ba)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)} + \frac{a \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (A + 2B)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^3,x)`output `(tan(c/2 + (d*x)/2)*(3*A*a + 2*B*a) - tan(c/2 + (d*x)/2)^3*(A*a + 2*B*a))/
(d*(tan(c/2 + (d*x)/2)^4 - 2*tan(c/2 + (d*x)/2)^2 + 1)) + (a*atanh(tan(c/2
+ (d*x)/2))*(A + 2*B))/d`

3.8 $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$

3.8.1	Optimal result	277
3.8.2	Mathematica [A] (verified)	277
3.8.3	Rubi [A] (verified)	278
3.8.4	Maple [A] (verified)	281
3.8.5	Fricas [A] (verification not implemented)	281
3.8.6	Sympy [F]	282
3.8.7	Maxima [A] (verification not implemented)	282
3.8.8	Giac [A] (verification not implemented)	283
3.8.9	Mupad [B] (verification not implemented)	283

3.8.1 Optimal result

Integrand size = 29, antiderivative size = 86

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= \frac{a(A + B) \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{a(2A + 3B) \tan(c + dx)}{3d} \\ &+ \frac{a(A + B) \sec(c + dx) \tan(c + dx)}{2d} + \frac{aA \sec^2(c + dx) \tan(c + dx)}{3d} \end{aligned}$$

```
output 1/2*a*(A+B)*arctanh(sin(d*x+c))/d+1/3*a*(2*A+3*B)*tan(d*x+c)/d+1/2*a*(A+B)
*sec(d*x+c)*tan(d*x+c)/d+1/3*a*A*sec(d*x+c)^2*tan(d*x+c)/d
```

3.8.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.65

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= \frac{a(3(A + B) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (6(A + B) + 3(A + B) \sec(c + dx) + 2A \tan^2(c + dx)))}{6d} \end{aligned}$$

```
input Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]
```

output $(a*(3*(A + B)*\text{ArcTanh}[\text{Sin}[c + d*x]] + \text{Tan}[c + d*x]*(6*(A + B) + 3*(A + B)*\text{Sec}[c + d*x] + 2*A*\text{Tan}[c + d*x]^2)))/(6*d)$

3.8.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.01, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec^4(c + dx)((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{1}{3} \int (3a(A + B) + a(2A + 3B) \cos(c + dx)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{3a(A + B) + a(2A + 3B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3227} \\
 & \frac{1}{3} \left(3a(A + B) \int \sec^3(c + dx) dx + a(2A + 3B) \int \sec^2(c + dx) dx \right) + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3} \left(a(2A + 3B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^2 dx + 3a(A + B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{4254} \\
& \frac{1}{3} \left(3a(A + B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx - \frac{a(2A + 3B) \int 1d(-\tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{24} \\
& \frac{1}{3} \left(3a(A + B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx + \frac{a(2A + 3B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{4255} \\
& \frac{1}{3} \left(3a(A + B) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{a(2A + 3B) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(3a(A + B) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{a(2A + 3B) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow \text{4257} \\
& \frac{1}{3} \left(3a(A + B) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{a(2A + 3B) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((a*(2*A + 3*B)*Tan[c + d*x])/d + 3*a*(A + B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/3`

3.8.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.8.4 Maple [A] (verified)

Time = 2.98 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.94

method	result
parts	$-\frac{aA\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c)}{d} + \frac{(aA+Ba)\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d} + \frac{Ba\tan(dx+c)}{d}$
derivativedivides	$\frac{aA\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) + Ba\tan(dx+c) - aA\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c) + Ba\left(\frac{\sec(dx+c)}{2}\right)}{d}$
default	$\frac{aA\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) + Ba\tan(dx+c) - aA\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c) + Ba\left(\frac{\sec(dx+c)}{2}\right)}{d}$
parallelrisch	$\left(\frac{3\left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c)\right)(A+B)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2} + \frac{3\left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c)\right)(A+B)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2}\right) + (A+B) \frac{d(\cos(3dx+3c)+3\cos(dx+c))}{d}$
norman	$\frac{4a(A-3B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3d} - \frac{2a(A-3B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3d} - \frac{3a(A+B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} - \frac{a(A+B)\left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d} - \frac{2a(7A+3B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3d}$
risch	$\frac{ia(3Ae^{5i(dx+c)} + 3Be^{5i(dx+c)} - 6Be^{4i(dx+c)} - 12Ae^{2i(dx+c)} - 12Be^{2i(dx+c)} - 3Ae^{i(dx+c)} - 3Be^{i(dx+c)} - 4A - 6B)}{3d(e^{2i(dx+c)} + 1)^3} +$

input `int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBOSE)`

output `-a*A/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(A*a+B*a)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+B*a/d*tan(d*x+c)`

3.8.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.22

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(A + B)a \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3(A + B)a \cos(dx + c)^3 \log(-\sin(dx + c) + 1) + 2(2(A + B)a \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 2(A + B)a \cos(dx + c)^3 \log(-\sin(dx + c) + 1))}{12d \cos(dx + c)^3}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")`

output `1/12*(3*(A + B)*a*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*(A + B)*a*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(2*(2*A + 3*B)*a*cos(d*x + c)^2 + 3*(A + B)*a*cos(d*x + c) + 2*A*a)*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.8.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= a \left(\int A \sec^4(c + dx) dx + \int A \cos(c + dx) \sec^4(c + dx) dx \right. \\ & \quad \left. + \int B \cos(c + dx) \sec^4(c + dx) dx + \int B \cos^2(c + dx) \sec^4(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `a*(Integral(A*sec(c + d*x)**4, x) + Integral(A*cos(c + d*x)*sec(c + d*x)**4, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**4, x) + Integral(B*cos(c + d*x)**2*sec(c + d*x)**4, x))`

3.8.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.48

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= \frac{4 \left(\tan(dx + c)^3 + 3 \tan(dx + c) \right) Aa - 3Aa \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) + 12B*a*\tan(dx + c)}{12d} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output `1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a - 3*A*a*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 3*B*a*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 12*B*a*tan(d*x + c))/d`

3.8.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.79

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(Aa + Ba) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - 3(Aa + Ba) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right) - \frac{2\left(3Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 3Ba \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 4Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 4Ba\right)}{6d}}{6d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")`

output `1/6*(3*(A*a + B*a)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*(A*a + B*a)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(3*A*a*tan(1/2*d*x + 1/2*c)^5 + 3*B*a*tan(1/2*d*x + 1/2*c)^3 - 4*A*a*tan(1/2*d*x + 1/2*c) + 4*B*a))/(tan(1/2*d*x + 1/2*c)^2 - 1)^3/d`

3.8.9 Mupad [B] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.47

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{a \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (A + B)}{d} - \frac{(Aa + Ba) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{4Aa}{3} - 4Ba\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (3Aa + 3Ba) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^4,x)`

output `(a*atanh(tan(c/2 + (d*x)/2))*(A + B))/d - (tan(c/2 + (d*x)/2)*(3*A*a + 3*B*a) + tan(c/2 + (d*x)/2)^5*(A*a + B*a) - tan(c/2 + (d*x)/2)^3*((4*A*a)/3 + 4*B*a))/(d*(3*tan(c/2 + (d*x)/2)^2 - 3*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 - 1))`

3.9 $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.9.1	Optimal result	284
3.9.2	Mathematica [A] (verified)	285
3.9.3	Rubi [A] (verified)	285
3.9.4	Maple [A] (verified)	288
3.9.5	Fricas [A] (verification not implemented)	289
3.9.6	Sympy [F]	289
3.9.7	Maxima [A] (verification not implemented)	290
3.9.8	Giac [A] (verification not implemented)	290
3.9.9	Mupad [B] (verification not implemented)	291

3.9.1 Optimal result

Integrand size = 29, antiderivative size = 106

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{a(3A + 4B)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{a(A + B)\tan(c + dx)}{d}$$

$$+ \frac{a(3A + 4B)\sec(c + dx)\tan(c + dx)}{8d}$$

$$+ \frac{aA\sec^3(c + dx)\tan(c + dx)}{4d} + \frac{a(A + B)\tan^3(c + dx)}{3d}$$

```
output 1/8*a*(3*A+4*B)*arctanh(sin(d*x+c))/d+a*(A+B)*tan(d*x+c)/d+1/8*a*(3*A+4*B)
*sec(d*x+c)*tan(d*x+c)/d+1/4*a*A*sec(d*x+c)^3*tan(d*x+c)/d+1/3*a*(A+B)*tan
(d*x+c)^3/d
```

3.9.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.73

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{a(3(3A + 4B)\operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx)(9A + 12B + 8(A + B)(2 + \cos(2(c + dx)))) \sec(c + dx)}{24d}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*(3*(3*A + 4*B)*ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*(9*A + 12*B + 8*(A + B)*(2 + Cos[2*(c + d*x)])*Sec[c + d*x] + 6*A*Sec[c + d*x]^2)*Tan[c + d*x])/(24*d)`

3.9.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.96, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 2009, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow 3447$$

$$\int \sec^5(c + dx)((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow 3500$$

$$\begin{aligned}
& \frac{1}{4} \int (4a(A+B) + a(3A+4B) \cos(c+dx)) \sec^4(c+dx) dx + \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \int \frac{4a(A+B) + a(3A+4B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^4} dx + \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{4} \left(4a(A+B) \int \sec^4(c+dx) dx + a(3A+4B) \int \sec^3(c+dx) dx \right) + \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left(a(3A+4B) \int \csc\left(c+dx + \frac{\pi}{2}\right)^3 dx + 4a(A+B) \int \csc\left(c+dx + \frac{\pi}{2}\right)^4 dx \right) + \\
& \quad \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{4254} \\
& \frac{1}{4} \left(a(3A+4B) \int \csc\left(c+dx + \frac{\pi}{2}\right)^3 dx - \frac{4a(A+B) \int (\tan^2(c+dx) + 1) d(-\tan(c+dx))}{d} \right) + \\
& \quad \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{2009} \\
& \frac{1}{4} \left(a(3A+4B) \int \csc\left(c+dx + \frac{\pi}{2}\right)^3 dx - \frac{4a(A+B) \left(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)\right)}{d} \right) + \\
& \quad \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{4255} \\
& \frac{1}{4} \left(a(3A+4B) \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{4a(A+B) \left(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)\right)}{d} \right) + \\
& \quad \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left(a(3A+4B) \left(\frac{1}{2} \int \csc\left(c+dx + \frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{4a(A+B) \left(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx)\right)}{d} \right) + \\
& \quad \frac{aA \tan(c+dx) \sec^3(c+dx)}{4d} \\
& \quad \downarrow \text{4257}
\end{aligned}$$

$$\frac{1}{4} \left(a(3A + 4B) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{4a(A + B) \left(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx) \right)}{d} \right. \\ \left. \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \right)$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*A*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (a*(3*A + 4*B)*(ArcTanh[Sin[c + d*x]])/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)) - (4*a*(A + B)*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d)/4`

3.9.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.9.4 Maple [A] (verified)

Time = 3.06 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.12

method	result
parts	$aA \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right) - \frac{(aA+Bd) \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d}$
derivativedivides	$-aA \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + Ba \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + aA \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)$
default	$-aA \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + Ba \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + aA \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)$
parallelrisch	$8 \left(- \frac{9(A + \frac{4B}{3}) \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{16} + \frac{9(A + \frac{4B}{3}) \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) \ln(\tan(\frac{dx}{2} + \frac{c}{2}))}{16} \right)$
norman	$\frac{3d(\cos(4dx+4c) + 4 \cos(2dx+2c))}{(1 + \tan^2(\frac{dx}{2} + \frac{c}{2}))^2 (\tan^2(\frac{dx}{2} + \frac{c}{2}) - 1)^4} - \frac{a(3A+4B) \tan^{11}(\frac{dx}{2} + \frac{c}{2})}{4d} + \frac{a(13A-20B) \tan^5(\frac{dx}{2} + \frac{c}{2})}{6d} + \frac{a(13A+12B) \tan(\frac{dx}{2} + \frac{c}{2})}{4d} + \frac{a(29A-4B) \tan^7(\frac{dx}{2} + \frac{c}{2})}{6d} + \frac{a(31A+12B) \tan^9(\frac{dx}{2} + \frac{c}{2})}{6d}$
risch	$- \frac{ia(9Ae^{7i(dx+c)} + 12Be^{7i(dx+c)} + 33Ae^{5i(dx+c)} + 12Be^{5i(dx+c)} - 48Ae^{4i(dx+c)} - 48Be^{4i(dx+c)} - 33Ae^{3i(dx+c)} - 12Be^{3i(dx+c)} - 12Ae^{2i(dx+c)} - 12Be^{2i(dx+c)} - 9Ae^{i(dx+c)} - 9Be^{i(dx+c)} - 9A - 9B)}{12d(e^{2i(dx+c)} + 1)^4}$

input `int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBOSE)`

output `a*A/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))-(A*a+B*a)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+B*a/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))`

3.9.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.20

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3A + 4B)a \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(3A + 4B)a \cos(dx + c)^4 \log(-\sin(dx + c) + 1) + 48 d \cos^5(dx + c)}{48 d \cos^5(dx + c)}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")`

output `1/48*(3*(3*A + 4*B)*a*cos(d*x + c)^4*log(sin(d*x + c) + 1) - 3*(3*A + 4*B)*a*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*(16*(A + B)*a*cos(d*x + c)^3 + 3*(3*A + 4*B)*a*cos(d*x + c)^2 + 8*(A + B)*a*cos(d*x + c) + 6*A*a)*sin(d*x + c))/(d*cos(d*x + c)^4)`

3.9.6 Sympy [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= a \left(\int A \sec^5(c + dx) dx + \int A \cos(c + dx) \sec^5(c + dx) dx \right. \\ \left. + \int B \cos(c + dx) \sec^5(c + dx) dx + \int B \cos^2(c + dx) \sec^5(c + dx) dx \right)$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output `a*(Integral(A*sec(c + d*x)**5, x) + Integral(A*cos(c + d*x)*sec(c + d*x)**5, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**5, x) + Integral(B*cos(c + d*x)**2*sec(c + d*x)**5, x))`

3.9.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.54

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa + 16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ba - 3 Aa \left(\frac{2 (3 \sin(dx+c)^3 - 5 \sin(dx+c))}{\sin(dx+c)^4 - 2 \sin(dx+c)^2 + 1} \right) - 12 B a (2 \sin(dx+c) / (\sin(dx+c)^2 - 1) - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1))}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output `1/48*(16*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a + 16*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a - 3*A*a*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 12*B*a*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)))/d`

3.9.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.77

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3Aa + 4Ba) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 3(3Aa + 4Ba) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) - \frac{2(9Aa \tan(\frac{1}{2}dx + \frac{1}{2}c) + 12B a \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 49Aa \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 28Ba \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 31Aa \tan(\frac{1}{2}dx + \frac{1}{2}c)^9 + 52Ba \tan(\frac{1}{2}dx + \frac{1}{2}c)^{11} - 39Aa \tan(\frac{1}{2}dx + \frac{1}{2}c)^{13} - 36Ba \tan(\frac{1}{2}dx + \frac{1}{2}c)^{15})}{\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1}}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")`

output `1/24*(3*(3*A*a + 4*B*a)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*(3*A*a + 4*B*a)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(9*A*a*tan(1/2*d*x + 1/2*c)^7 + 12*B*a*tan(1/2*d*x + 1/2*c)^9 - 49*A*a*tan(1/2*d*x + 1/2*c)^11 - 28*B*a*tan(1/2*d*x + 1/2*c)^13 + 31*A*a*tan(1/2*d*x + 1/2*c)^15 + 52*B*a*tan(1/2*d*x + 1/2*c)^17 - 39*A*a*tan(1/2*d*x + 1/2*c)^19 - 36*B*a*tan(1/2*d*x + 1/2*c)^21))/(tan(1/2*d*x + 1/2*c)^2 - 1)^4/d`

3.9.9 Mupad [B] (verification not implemented)

Time = 1.67 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.57

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{\left(-\frac{3Aa}{4} - Ba\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{49Aa}{12} + \frac{7Ba}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{31Aa}{12} - \frac{13Ba}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{13Aa}{4} + Ba\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)} + \frac{a \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (3A + 4B)}{4d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^5,x)`

output `(tan(c/2 + (d*x)/2)*((13*A*a)/4 + 3*B*a) - tan(c/2 + (d*x)/2)^7*((3*A*a)/4 + B*a) - tan(c/2 + (d*x)/2)^3*((31*A*a)/12 + (13*B*a)/3) + tan(c/2 + (d*x)/2)^5*((49*A*a)/12 + (7*B*a)/3))/(d*(6*tan(c/2 + (d*x)/2)^4 - 4*tan(c/2 + (d*x)/2)^2 - 4*tan(c/2 + (d*x)/2)^6 + tan(c/2 + (d*x)/2)^8 + 1)) + (a*atanh(tan(c/2 + (d*x)/2))*(3*A + 4*B))/(4*d)`

3.10 $\int \cos^3(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.10.1	Optimal result	292
3.10.2	Mathematica [A] (verified)	293
3.10.3	Rubi [A] (verified)	293
3.10.4	Maple [A] (verified)	297
3.10.5	Fricas [A] (verification not implemented)	298
3.10.6	Sympy [B] (verification not implemented)	298
3.10.7	Maxima [A] (verification not implemented)	299
3.10.8	Giac [A] (verification not implemented)	300
3.10.9	Mupad [B] (verification not implemented)	300

3.10.1 Optimal result

Integrand size = 31, antiderivative size = 191

$$\int \cos^3(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$$

$$= \frac{1}{16}a^2(12A+11B)x + \frac{a^2(9A+8B) \sin(c+dx)}{5d} + \frac{a^2(12A+11B) \cos(c+dx) \sin(c+dx)}{16d}$$

$$+ \frac{a^2(12A+11B) \cos^3(c+dx) \sin(c+dx)}{24d} + \frac{a^2(6A+7B) \cos^4(c+dx) \sin(c+dx)}{30d}$$

$$+ \frac{B \cos^4(c+dx) (a^2+a^2 \cos(c+dx)) \sin(c+dx)}{6d} - \frac{a^2(9A+8B) \sin^3(c+dx)}{15d}$$

```
output 1/16*a^2*(12*A+11*B)*x+1/5*a^2*(9*A+8*B)*sin(d*x+c)/d+1/16*a^2*(12*A+11*B)
*cos(d*x+c)*sin(d*x+c)/d+1/24*a^2*(12*A+11*B)*cos(d*x+c)^3*sin(d*x+c)/d+1/
30*a^2*(6*A+7*B)*cos(d*x+c)^4*sin(d*x+c)/d+1/6*B*cos(d*x+c)^4*(a^2+a^2*cos
(d*x+c))*sin(d*x+c)/d-1/15*a^2*(9*A+8*B)*sin(d*x+c)^3/d
```

3.10.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.70

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{a^2(660Bc + 720Adx + 660Bdx + 120(11A + 10B) \sin(c + dx) + 15(32A + 31B) \sin(2(c + dx)) + 180A \sin(3(c + dx)) + 200B \sin(3(c + dx)) + 60A \sin(4(c + dx)) + 75B \sin(4(c + dx)) + 12A \sin(5(c + dx)) + 24B \sin(5(c + dx)) + 5B \sin(6(c + dx)))}{960d}$$

input `Integrate[Cos[c + d*x]^3*(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]),x]`

output $(a^2(660Bc + 720A*d*x + 660B*d*x + 120(11A + 10B)*\text{Sin}[c + d*x] + 15(32A + 31B)*\text{Sin}[2*(c + d*x)] + 180A*\text{Sin}[3*(c + d*x)] + 200B*\text{Sin}[3*(c + d*x)] + 60A*\text{Sin}[4*(c + d*x)] + 75B*\text{Sin}[4*(c + d*x)] + 12A*\text{Sin}[5*(c + d*x)] + 24B*\text{Sin}[5*(c + d*x)] + 5B*\text{Sin}[6*(c + d*x)])/(960*d)$

3.10.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.93, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3455, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^3 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3455}$$

$$\frac{1}{6} \int \cos^3(c + dx)(\cos(c + dx)a + a)(2a(3A + 2B) + a(6A + 7B) \cos(c + dx)) dx + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

$$\downarrow \text{3042}$$

3.10. $\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\frac{1}{6} \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right) \left(2a(3A + 2B) + a(6A + 7B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3447

$$\frac{1}{6} \int \cos^3(c + dx) \left((6A + 7B) \cos^2(c + dx) a^2 + 2(3A + 2B) a^2 + (2(3A + 2B) a^2 + (6A + 7B) a^2) \cos(c + dx) \right) dx + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3042

$$\frac{1}{6} \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 \left((6A + 7B) \sin \left(c + dx + \frac{\pi}{2} \right)^2 a^2 + 2(3A + 2B) a^2 + (2(3A + 2B) a^2 + (6A + 7B) a^2) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3502

$$\frac{1}{6} \left(\frac{1}{5} \int \cos^3(c + dx) (6(9A + 8B) a^2 + 5(12A + 11B) \cos(c + dx) a^2) dx + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx)}{5d} \right) + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 (6(9A + 8B) a^2 + 5(12A + 11B) \sin \left(c + dx + \frac{\pi}{2} \right) a^2) dx + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx)}{5d} \right) + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3227

$$\frac{1}{6} \left(\frac{1}{5} \left(5a^2(12A + 11B) \int \cos^4(c + dx) dx + 6a^2(9A + 8B) \int \cos^3(c + dx) dx \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx)}{5d} \right) + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(6a^2(9A + 8B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 dx + 5a^2(12A + 11B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^4 dx \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx)}{5d} \right) + \frac{B \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d}$$

↓ 3113

$$\frac{1}{6} \left(\frac{1}{5} \left(5a^2(12A + 11B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^4 dx - \frac{6a^2(9A + 8B) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d} \right)$$

↓ 2009

$$\frac{1}{6} \left(\frac{1}{5} \left(5a^2(12A + 11B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^4 dx - \frac{6a^2(9A + 8B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d} \right)$$

↓ 3115

$$\frac{1}{6} \left(\frac{1}{5} \left(5a^2(12A + 11B) \left(\frac{3}{4} \int \cos^2(c + dx) dx + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) - \frac{6a^2(9A + 8B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(5a^2(12A + 11B) \left(\frac{3}{4} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) - \frac{6a^2(9A + 8B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d} \right)$$

↓ 3115

$$\frac{1}{6} \left(\frac{1}{5} \left(5a^2(12A + 11B) \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) + \frac{\sin(c + dx) \cos^3(c + dx)}{4d} \right) - \frac{6a^2(9A + 8B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d} \right)$$

↓ 24

$$\frac{1}{6} \left(\frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx)}{5d} + \frac{1}{5} \left(5a^2(12A + 11B) \left(\frac{\sin(c + dx) \cos^3(c + dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} \right) \right) + \frac{a^2(6A + 7B) \sin(c + dx) \cos^4(c + dx) (a^2 \cos(c + dx) + a^2)}{6d} \right)$$

input `Int[Cos[c + d*x]^3*(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]),x]`

output `(B*cos[c + d*x]^4*(a^2 + a^2*cos[c + d*x])*sin[c + d*x])/(6*d) + ((a^2*(6*A + 7*B)*cos[c + d*x]^4*sin[c + d*x])/(5*d) + ((-6*a^2*(9*A + 8*B)*(-sin[c + d*x] + sin[c + d*x]^3/3))/d + 5*a^2*(12*A + 11*B)*((cos[c + d*x]^3*sin[c + d*x])/(4*d) + (3*(x/2 + (cos[c + d*x]*sin[c + d*x])/(2*d))/4))/5)/6`

3.10.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*sin[e + f*x])^m*(A*c + (B*c + A*d)*sin[e + f*x] + B*d*sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)^2], x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.10.4 Maple [A] (verified)

Time = 4.45 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.58

method	result
parallelrisch	$\frac{\left(\left(8A + \frac{31B}{4}\right) \sin(2dx+2c) + \left(3A + \frac{10B}{3}\right) \sin(3dx+3c) + \left(A + \frac{5B}{4}\right) \sin(4dx+4c) + \frac{(A+2B) \sin(5dx+5c)}{5} + \frac{B \sin(6dx+6c)}{12}\right) + 2(1}{16d}$
parts	$\frac{(A a^2 + 2B a^2) \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right) \sin(dx+c)}{5d} + \frac{(2A a^2 + B a^2) \left(\frac{\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}}{4}\right) \sin(dx+c)}{d}$
risch	$\frac{3a^2xA}{4} + \frac{11a^2Bx}{16} + \frac{11 \sin(dx+c)Aa^2}{8d} + \frac{5 \sin(dx+c)Ba^2}{4d} + \frac{Ba^2 \sin(6dx+6c)}{192d} + \frac{\sin(5dx+5c)Aa^2}{80d} + \frac{\sin(5dx+5c)Ba^2}{80d}$
derivativedivides	$\frac{A a^2 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right) \sin(dx+c)}{5} + B a^2 \left(\frac{\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8}}{6}\right) \sin(dx+c) + \frac{5dx}{16} + \frac{5c}{16}$
default	$\frac{A a^2 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right) \sin(dx+c)}{5} + B a^2 \left(\frac{\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8}}{6}\right) \sin(dx+c) + \frac{5dx}{16} + \frac{5c}{16}$
norman	$\frac{a^2(12A+11B)x}{16} + \frac{17a^2(12A+11B)\left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{24d} + \frac{a^2(12A+11B)\left(\tan^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{8d} + \frac{3a^2(12A+11B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{8} + \frac{15a^2(12A+11B)x}{16}$

3.10. $\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$


```
input int(cos(d*x+c)^3*(a+cos(d*x+c))*a^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/16*((8*A+31/4*B)*sin(2*d*x+2*c)+(3*A+10/3*B)*sin(3*d*x+3*c)+(A+5/4*B)*sin(4*d*x+4*c)+1/5*(A+2*B)*sin(5*d*x+5*c)+1/12*B*sin(6*d*x+6*c)+2*(11*A+10*B)*sin(d*x+c)+12*(11/12*B+A)*x*d)*a^2/d
```

3.10.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.68

$$\int \cos^3(c+dx)(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{15(12A+11B)a^2dx + (40Ba^2\cos(dx+c))^5 + 48(A+2B)a^2\cos(dx+c)^4 + 10(12A+11B)a^2\cos(dx+c)^3 + 16(9A+8B)a^2\cos(dx+c)^2 + 15(12A+11B)a^2\cos(dx+c) + 32(9A+8B)a^2\sin(dx+c)}{24d}$$

```
input integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/240*(15*(12*A + 11*B)*a^2*d*x + (40*B*a^2*cos(d*x + c)^5 + 48*(A + 2*B)*a^2*cos(d*x + c)^4 + 10*(12*A + 11*B)*a^2*cos(d*x + c)^3 + 16*(9*A + 8*B)*a^2*cos(d*x + c)^2 + 15*(12*A + 11*B)*a^2*cos(d*x + c) + 32*(9*A + 8*B)*a^2*sin(d*x + c))/d
```

3.10.6 SymPy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 600 vs. 2(173) = 346.

Time = 0.39 (sec) , antiderivative size = 600, normalized size of antiderivative = 3.14

$$\int \cos^3(c+dx)(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \left\{ \begin{array}{l} \frac{3Aa^2x\sin^4(c+dx)}{4} + \frac{3Aa^2x\sin^2(c+dx)\cos^2(c+dx)}{2} + \frac{3Aa^2x\cos^4(c+dx)}{4} + \frac{8Aa^2\sin^5(c+dx)}{15d} + \frac{4Aa^2\sin^3(c+dx)\cos^2(c+dx)}{3d} + \frac{3Aa^2\sin(c+dx)\cos^4(c+dx)}{3d} \\ x(A+B\cos(c))(a\cos(c)+a)^2\cos^3(c) \end{array} \right.$$

```
input integrate(cos(d*x+c)**3*(a+a*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)
```

3.10. $\int \cos^3(c+dx)(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx$

output `Piecewise((3*A*a**2*x*sin(c + d*x)**4/4 + 3*A*a**2*x*sin(c + d*x)**2*cos(c + d*x)**2/2 + 3*A*a**2*x*cos(c + d*x)**4/4 + 8*A*a**2*sin(c + d*x)**5/(15*d) + 4*A*a**2*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 3*A*a**2*sin(c + d*x)**3*cos(c + d*x)/(4*d) + 2*A*a**2*sin(c + d*x)**3/(3*d) + A*a**2*sin(c + d*x)*cos(c + d*x)**4/d + 5*A*a**2*sin(c + d*x)*cos(c + d*x)**3/(4*d) + A*a**2*sin(c + d*x)*cos(c + d*x)**2/d + 5*B*a**2*x*sin(c + d*x)**6/16 + 15*B*a**2*x*sin(c + d*x)**4*cos(c + d*x)**2/16 + 3*B*a**2*x*sin(c + d*x)**4/8 + 15*B*a**2*x*sin(c + d*x)**2*cos(c + d*x)**4/16 + 3*B*a**2*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 5*B*a**2*x*cos(c + d*x)**6/16 + 3*B*a**2*x*cos(c + d*x)**4/8 + 5*B*a**2*sin(c + d*x)**5*cos(c + d*x)/(16*d) + 16*B*a**2*sin(c + d*x)**5/(15*d) + 5*B*a**2*sin(c + d*x)**3*cos(c + d*x)**3/(6*d) + 8*B*a**2*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 3*B*a**2*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 11*B*a**2*sin(c + d*x)*cos(c + d*x)**5/(16*d) + 2*B*a**2*sin(c + d*x)*cos(c + d*x)**4/d + 5*B*a**2*sin(c + d*x)*cos(c + d*x)**3/(8*d), Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**2*cos(c)**3, True))`

3.10.7 Maxima [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.13

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{64 (3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c)) Aa^2 - 320 (\sin(dx + c)^3 - 3 \sin(dx + c)) Aa^2 + \dots}{\dots}$$

input `integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/960*(64*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*A*a^2 - 320*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^2 + 60*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^2 + 128*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a^2 - 5*(4*sin(2*d*x + 2*c)^3 - 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*sin(2*d*x + 2*c))*B*a^2 + 30*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^2)/d`

3.10.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.87

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^2 \sin(6dx + 6c)}{192d} + \frac{1}{16} (12Aa^2 + 11Ba^2)x + \frac{(Aa^2 + 2Ba^2) \sin(5dx + 5c)}{80d}$$

$$+ \frac{(4Aa^2 + 5Ba^2) \sin(4dx + 4c)}{64d} + \frac{(9Aa^2 + 10Ba^2) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(32Aa^2 + 31Ba^2) \sin(2dx + 2c)}{64d} + \frac{(11Aa^2 + 10Ba^2) \sin(dx + c)}{8d}$$

```
input integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/192*B*a^2*sin(6*d*x + 6*c)/d + 1/16*(12*A*a^2 + 11*B*a^2)*x + 1/80*(A*a^2 + 2*B*a^2)*sin(5*d*x + 5*c)/d + 1/64*(4*A*a^2 + 5*B*a^2)*sin(4*d*x + 4*c)/d + 1/48*(9*A*a^2 + 10*B*a^2)*sin(3*d*x + 3*c)/d + 1/64*(32*A*a^2 + 31*B*a^2)*sin(2*d*x + 2*c)/d + 1/8*(11*A*a^2 + 10*B*a^2)*sin(d*x + c)/d
```

3.10.9 Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.65

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{3Aa^2}{2} + \frac{11Ba^2}{8}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{11} + \left(\frac{17Aa^2}{2} + \frac{187Ba^2}{24}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{107Aa^2}{5} + \frac{331Ba^2}{20}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{12} + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 15 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)}$$

$$- \frac{a^2 (12A + 11B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{8d}$$

$$+ \frac{a^2 \operatorname{atan}\left(\frac{a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (12A + 11B)}{8 \left(\frac{3Aa^2}{2} + \frac{11Ba^2}{8}\right)}\right) (12A + 11B)}{8d}$$

```
input int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2,x)
```

output

$$\begin{aligned} & (\tan(c/2 + (d*x)/2)*((13*A*a^2)/2 + (53*B*a^2)/8) + \tan(c/2 + (d*x)/2)^{11}* \\ & ((3*A*a^2)/2 + (11*B*a^2)/8) + \tan(c/2 + (d*x)/2)^3*((31*A*a^2)/2 + (87*B* \\ & a^2)/8) + \tan(c/2 + (d*x)/2)^9*((17*A*a^2)/2 + (187*B*a^2)/24) + \tan(c/2 + \\ & (d*x)/2)^7*((107*A*a^2)/5 + (331*B*a^2)/20) + \tan(c/2 + (d*x)/2)^5*((117* \\ & A*a^2)/5 + (501*B*a^2)/20))/(d*(6*\tan(c/2 + (d*x)/2)^2 + 15*\tan(c/2 + (d*x) \\ &)/2)^4 + 20*\tan(c/2 + (d*x)/2)^6 + 15*\tan(c/2 + (d*x)/2)^8 + 6*\tan(c/2 + (\\ & d*x)/2)^{10} + \tan(c/2 + (d*x)/2)^{12} + 1)) - (a^2*(12*A + 11*B)*(atan(\tan(c/ \\ & 2 + (d*x)/2)) - (d*x)/2))/(8*d) + (a^2*atan((a^2*\tan(c/2 + (d*x)/2)*(12*A \\ & + 11*B))/(8*((3*A*a^2)/2 + (11*B*a^2)/8)))*(12*A + 11*B))/(8*d) \end{aligned}$$

3.11 $\int \cos^2(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.11.1	Optimal result	302
3.11.2	Mathematica [A] (verified)	303
3.11.3	Rubi [A] (verified)	303
3.11.4	Maple [A] (verified)	307
3.11.5	Fricas [A] (verification not implemented)	308
3.11.6	Sympy [B] (verification not implemented)	308
3.11.7	Maxima [A] (verification not implemented)	309
3.11.8	Giac [A] (verification not implemented)	309
3.11.9	Mupad [B] (verification not implemented)	310

3.11.1 Optimal result

Integrand size = 31, antiderivative size = 160

$$\int \cos^2(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$$

$$= \frac{1}{8}a^2(7A+6B)x + \frac{a^2(10A+9B) \sin(c+dx)}{5d}$$

$$+ \frac{a^2(7A+6B) \cos(c+dx) \sin(c+dx)}{8d} + \frac{a^2(5A+6B) \cos^3(c+dx) \sin(c+dx)}{20d}$$

$$+ \frac{B \cos^3(c+dx) (a^2+a^2 \cos(c+dx)) \sin(c+dx)}{5d} - \frac{a^2(10A+9B) \sin^3(c+dx)}{15d}$$

```
output 1/8*a^2*(7*A+6*B)*x+1/5*a^2*(10*A+9*B)*sin(d*x+c)/d+1/8*a^2*(7*A+6*B)*cos(
d*x+c)*sin(d*x+c)/d+1/20*a^2*(5*A+6*B)*cos(d*x+c)^3*sin(d*x+c)/d+1/5*B*cos
(d*x+c)^3*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d-1/15*a^2*(10*A+9*B)*sin(d*x+c)
^3/d
```

3.11.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.68

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{a^2(360Bc + 420Adx + 360Bdx + 60(12A + 11B) \sin(c + dx) + 240(A + B) \sin(2(c + dx)) + 80A \sin(3(c + dx)))}{480d}$$

input `Integrate[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(a^2*(360*B*c + 420*A*d*x + 360*B*d*x + 60*(12*A + 11*B)*Sin[c + d*x] + 240*(A + B)*Sin[2*(c + d*x)] + 80*A*Ssin[3*(c + d*x)] + 90*B*Ssin[3*(c + d*x)] + 15*A*Ssin[4*(c + d*x)] + 30*B*Ssin[4*(c + d*x)] + 6*B*Ssin[5*(c + d*x)])/(480*d)`

3.11.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.94, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3455, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3455}$$

$$\frac{1}{5} \int \cos^2(c + dx)(\cos(c + dx)a + a)(a(5A + 3B) + a(5A + 6B) \cos(c + dx)) dx + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

$$\downarrow \text{3042}$$

3.11. $\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\frac{1}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right) \left(a(5A + 3B) + a(5A + 6B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

↓ 3447

$$\frac{1}{5} \int \cos^2(c + dx) \left((5A + 6B) \cos^2(c + dx) a^2 + (5A + 3B) a^2 + ((5A + 3B) a^2 + (5A + 6B) a^2) \cos(c + dx) \right) dx + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left((5A + 6B) \sin \left(c + dx + \frac{\pi}{2} \right)^2 a^2 + (5A + 3B) a^2 + ((5A + 3B) a^2 + (5A + 6B) a^2) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

↓ 3502

$$\frac{1}{5} \left(\frac{1}{4} \int \cos^2(c + dx) (5(7A + 6B) a^2 + 4(10A + 9B) \cos(c + dx) a^2) dx + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx)}{4d} \right) + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 (5(7A + 6B) a^2 + 4(10A + 9B) \sin \left(c + dx + \frac{\pi}{2} \right) a^2) dx + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx)}{4d} \right) + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

↓ 3227

$$\frac{1}{5} \left(\frac{1}{4} \left(4a^2(10A + 9B) \int \cos^3(c + dx) dx + 5a^2(7A + 6B) \int \cos^2(c + dx) dx \right) + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx)}{4d} \right) + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(5a^2(7A + 6B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + 4a^2(10A + 9B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx)}{4d} \right) + \frac{B \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d}$$

3.11. $\int \cos^2(c + dx) (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$

↓ 3113

$$\frac{1}{5} \left(\frac{1}{4} \left(5a^2(7A + 6B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4a^2(10A + 9B) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right) + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d} \right)$$

↓ 2009

$$\frac{1}{5} \left(\frac{1}{4} \left(5a^2(7A + 6B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4a^2(10A + 9B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d} \right)$$

↓ 3115

$$\frac{1}{5} \left(\frac{1}{4} \left(5a^2(7A + 6B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{4a^2(10A + 9B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d} \right)$$

↓ 24

$$\frac{1}{5} \left(\frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx)}{4d} + \frac{1}{4} \left(5a^2(7A + 6B) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) - \frac{4a^2(10A + 9B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{a^2(5A + 6B) \sin(c + dx) \cos^3(c + dx) (a^2 \cos(c + dx) + a^2)}{5d} \right)$$

input `Int[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(B*Cos[c + d*x]^3*(a^2 + a^2*Cos[c + d*x])*Sin[c + d*x])/(5*d) + ((a^2*(5*A + 6*B)*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (5*a^2*(7*A + 6*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (4*a^2*(10*A + 9*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/4)/5`

3.11.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3455 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

$$3.11. \int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.11.4 Maple [A] (verified)

Time = 3.52 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.58

method	result
parallelrisch	$\frac{(16(A+B) \sin(2dx+2c)+2\left(\frac{8A}{3}+3B\right) \sin(3dx+3c)+(A+2B) \sin(4dx+4c)+\frac{2B \sin(5dx+5c)}{5}+4(12A+11B) \sin(dx+c)+28(A+6/7B)x^2/d}{32d}$
parts	$\frac{(Aa^2+2Ba^2) \left(\frac{(\cos^3(dx+c)+\frac{3\cos(\frac{dx+c}{2})) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right)}{d} + \frac{(2Aa^2+Ba^2)(2+\cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{Aa^2(2+\cos^2(dx+c))}{16d}$
risch	$\frac{7a^2xA}{8} + \frac{3a^2Bx}{4} + \frac{3 \sin(dx+c)Aa^2}{2d} + \frac{11 \sin(dx+c)Ba^2}{8d} + \frac{\sin(5dx+5c)Ba^2}{80d} + \frac{\sin(4dx+4c)Aa^2}{32d} + \frac{\sin(4dx+4c)Ba^2}{16d}$
derivativedivides	$Aa^2 \left(\frac{(\cos^3(dx+c)+\frac{3\cos(\frac{dx+c}{2})) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{Ba^2 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + \frac{2Aa^2(2+\cos^2(dx+c))}{16d}$
default	$Aa^2 \left(\frac{(\cos^3(dx+c)+\frac{3\cos(\frac{dx+c}{2})) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{Ba^2 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + \frac{2Aa^2(2+\cos^2(dx+c))}{16d}$
norman	$\frac{a^2(7A+6B)x}{8} + \frac{7a^2(7A+6B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{6d} + \frac{a^2(7A+6B) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{4d} + \frac{5a^2(7A+6B)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{8} + \frac{5a^2(7A+6B)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{4d}$

```
input int(cos(d*x+c)^2*(a+cos(d*x+c))*a^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/32*(16*(A+B)*sin(2*d*x+2*c)+2*(8/3*A+3*B)*sin(3*d*x+3*c)+(A+2*B)*sin(4*d
*x+4*c)+2/5*B*sin(5*d*x+5*c)+4*(12*A+11*B)*sin(d*x+c)+28*(A+6/7*B)*x^2/d
```

$$3.11. \int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

3.11.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.69

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{15(7A + 6B)a^2 dx + (24Ba^2 \cos(dx + c))^4 + 30(A + 2B)a^2 \cos(dx + c)^3 + 8(10A + 9B)a^2 \cos(dx + c)^2 + 16(10A + 9B)a^2 \sin(dx + c)}{120d}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `1/120*(15*(7*A + 6*B)*a^2*d*x + (24*B*a^2*cos(d*x + c)^4 + 30*(A + 2*B)*a^2*cos(d*x + c)^3 + 8*(10*A + 9*B)*a^2*cos(d*x + c)^2 + 15*(7*A + 6*B)*a^2*cos(d*x + c) + 16*(10*A + 9*B)*a^2*sin(d*x + c))/d`

3.11.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 459 vs. $2(144) = 288$.

Time = 0.30 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.87

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \left\{ \begin{array}{l} \frac{3Aa^2 x \sin^4(c+dx)}{8} + \frac{3Aa^2 x \sin^2(c+dx) \cos^2(c+dx)}{4} + \frac{Aa^2 x \sin^2(c+dx)}{2} + \frac{3Aa^2 x \cos^4(c+dx)}{8} + \frac{Aa^2 x \cos^2(c+dx)}{2} + \frac{3Aa^2 \sin^3(c+dx)}{8} \\ x(A + B \cos(c)) (a \cos(c) + a)^2 \cos^2(c) \end{array} \right.$$

input `integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Piecewise((3*A*a**2*x**sin(c + d*x)**4/8 + 3*A*a**2*x**sin(c + d*x)**2*cos(c + d*x)**2/4 + A*a**2*x**sin(c + d*x)**2/2 + 3*A*a**2*x*cos(c + d*x)**4/8 + A*a**2*x*cos(c + d*x)**2/2 + 3*A*a**2*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 4*A*a**2*sin(c + d*x)**3/(3*d) + 5*A*a**2*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 2*A*a**2*sin(c + d*x)*cos(c + d*x)**2/d + A*a**2*sin(c + d*x)*cos(c + d*x)/(2*d) + 3*B*a**2*x**sin(c + d*x)**4/4 + 3*B*a**2*x**sin(c + d*x)**2*cos(c + d*x)**2/2 + 3*B*a**2*x*cos(c + d*x)**4/4 + 8*B*a**2*sin(c + d*x)**5/(15*d) + 4*B*a**2*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 3*B*a**2*sin(c + d*x)**3*cos(c + d*x)/(4*d) + 2*B*a**2*sin(c + d*x)**3/(3*d) + B*a**2*sin(c + d*x)*cos(c + d*x)**4/d + 5*B*a**2*sin(c + d*x)*cos(c + d*x)**3/(4*d) + B*a**2*sin(c + d*x)*cos(c + d*x)**2/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**2*cos(c)**2, True))`

3.11. $\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

3.11.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.11

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx =$$

$$\frac{320 (\sin(dx + c)^3 - 3 \sin(dx + c))Aa^2 - 15 (12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(2 dx + 2 c))Aa^2 -$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `-1/480*(320*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^2 - 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^2 - 120*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^2 - 32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a^2 + 160*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^2 - 30*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^2)/d`

3.11.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.86

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^2 \sin(5 dx + 5 c)}{80 d} + \frac{1}{8} (7 Aa^2 + 6 Ba^2) x$$

$$+ \frac{(Aa^2 + 2 Ba^2) \sin(4 dx + 4 c)}{32 d} + \frac{(8 Aa^2 + 9 Ba^2) \sin(3 dx + 3 c)}{48 d}$$

$$+ \frac{(Aa^2 + Ba^2) \sin(2 dx + 2 c)}{2 d} + \frac{(12 Aa^2 + 11 Ba^2) \sin(dx + c)}{8 d}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/80*B*a^2*sin(5*d*x + 5*c)/d + 1/8*(7*A*a^2 + 6*B*a^2)*x + 1/32*(A*a^2 + 2*B*a^2)*sin(4*d*x + 4*c)/d + 1/48*(8*A*a^2 + 9*B*a^2)*sin(3*d*x + 3*c)/d + 1/2*(A*a^2 + B*a^2)*sin(2*d*x + 2*c)/d + 1/8*(12*A*a^2 + 11*B*a^2)*sin(d*x + c)/d`

3.11. $\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

3.11.9 Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.73

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{7Aa^2}{4} + \frac{3Ba^2}{2}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{49Aa^2}{6} + 7Ba^2\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{40Aa^2}{3} + \frac{72Ba^2}{5}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{7Aa^2}{4} + \frac{3Ba^2}{2}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{49Aa^2}{6} + 7Ba^2\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \left(\frac{40Aa^2}{3} + \frac{72Ba^2}{5}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)} - \frac{a^2(7A + 6B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{4d} + \frac{a^2 \operatorname{atan}\left(\frac{a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (7A + 6B)}{4 \left(\frac{7Aa^2}{4} + \frac{3Ba^2}{2}\right)}\right) (7A + 6B)}{4d}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2,x)`

output

```
(tan(c/2 + (d*x)/2)*((25*A*a^2)/4 + (13*B*a^2)/2) + tan(c/2 + (d*x)/2)^9*(
(7*A*a^2)/4 + (3*B*a^2)/2) + tan(c/2 + (d*x)/2)^7*((49*A*a^2)/6 + 7*B*a^2)
+ tan(c/2 + (d*x)/2)^3*((79*A*a^2)/6 + 9*B*a^2) + tan(c/2 + (d*x)/2)^5*((
40*A*a^2)/3 + (72*B*a^2)/5))/(d*(5*tan(c/2 + (d*x)/2)^2 + 10*tan(c/2 + (d*
x)/2)^4 + 10*tan(c/2 + (d*x)/2)^6 + 5*tan(c/2 + (d*x)/2)^8 + tan(c/2 + (d*
x)/2)^10 + 1)) - (a^2*(7*A + 6*B)*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(4
*d) + (a^2*atan((a^2*tan(c/2 + (d*x)/2)*(7*A + 6*B))/(4*((7*A*a^2)/4 + (3*
B*a^2)/2)))*(7*A + 6*B))/(4*d)
```

3.12 $\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

3.12.1	Optimal result	311
3.12.2	Mathematica [A] (verified)	311
3.12.3	Rubi [A] (verified)	312
3.12.4	Maple [A] (verified)	314
3.12.5	Fricas [A] (verification not implemented)	315
3.12.6	Sympy [B] (verification not implemented)	315
3.12.7	Maxima [A] (verification not implemented)	316
3.12.8	Giac [A] (verification not implemented)	316
3.12.9	Mupad [B] (verification not implemented)	317

3.12.1 Optimal result

Integrand size = 29, antiderivative size = 129

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{1}{8}a^2(8A + 7B)x + \frac{a^2(8A + 7B) \sin(c + dx)}{6d} + \frac{a^2(8A + 7B) \cos(c + dx) \sin(c + dx)}{24d}$$

$$+ \frac{(4A - B)(a + a \cos(c + dx))^2 \sin(c + dx)}{12d} + \frac{B(a + a \cos(c + dx))^3 \sin(c + dx)}{4ad}$$

```
output 1/8*a^2*(8*A+7*B)*x+1/6*a^2*(8*A+7*B)*sin(d*x+c)/d+1/24*a^2*(8*A+7*B)*cos(
d*x+c)*sin(d*x+c)/d+1/12*(4*A-B)*(a+a*cos(d*x+c))^2*sin(d*x+c)/d+1/4*B*(a+
a*cos(d*x+c))^3*sin(d*x+c)/a/d
```

3.12.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.67

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{a^2(84Bc + 96Adx + 84Bdx + 24(7A + 6B) \sin(c + dx) + 48(A + B) \sin(2(c + dx)) + 8A \sin(3(c + dx)))}{96d}$$

input `Integrate[Cos[c + d*x]*(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]),x]`

output $(a^2*(84*B*c + 96*A*d*x + 84*B*d*x + 24*(7*A + 6*B)*\sin[c + d*x] + 48*(A + B)*\sin[2*(c + d*x)] + 8*A*\sin[3*(c + d*x)] + 16*B*\sin[3*(c + d*x)] + 3*B*\sin[4*(c + d*x)])/(96*d)$

3.12.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.99, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 3447, 3042, 3502, 3042, 3230, 3042, 3123}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int (a \cos(c + dx) + a)^2 (A \cos(c + dx) + B \cos^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{\int (\cos(c + dx)a + a)^2 (3aB + a(4A - B) \cos(c + dx)) dx}{4a} + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^3}{4ad} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (\sin(c + dx + \frac{\pi}{2})a + a)^2 (3aB + a(4A - B) \sin(c + dx + \frac{\pi}{2})) dx}{4a} + \\
 & \quad \frac{B \sin(c + dx)(a \cos(c + dx) + a)^3}{4ad} \\
 & \quad \downarrow \text{3230}
 \end{aligned}$$

3.12. $\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\frac{\frac{1}{3}a(8A + 7B) \int (\cos(c + dx)a + a)^2 dx + \frac{a(4A-B) \sin(c+dx)(a \cos(c+dx)+a)^2}{3d}}{\frac{4a}{4ad} B \sin(c + dx)(a \cos(c + dx) + a)^3} +$$

↓ 3042

$$\frac{\frac{1}{3}a(8A + 7B) \int (\sin(c + dx + \frac{\pi}{2})a + a)^2 dx + \frac{a(4A-B) \sin(c+dx)(a \cos(c+dx)+a)^2}{3d}}{\frac{4a}{4ad} B \sin(c + dx)(a \cos(c + dx) + a)^3} +$$

↓ 3123

$$\frac{\frac{1}{3}a(8A + 7B) \left(\frac{2a^2 \sin(c+dx)}{d} + \frac{a^2 \sin(c+dx) \cos(c+dx)}{2d} + \frac{3a^2 x}{2} \right) + \frac{a(4A-B) \sin(c+dx)(a \cos(c+dx)+a)^2}{3d}}{\frac{4a}{4ad} B \sin(c + dx)(a \cos(c + dx) + a)^3} +$$

```
input Int[Cos[c + d*x]*(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]),x]
```

```
output (B*(a + a*cos[c + d*x])^3*sin[c + d*x])/(4*a*d) + ((a*(4*A - B)*(a + a*cos[c + d*x])^2*sin[c + d*x])/(3*d) + (a*(8*A + 7*B)*((3*a^2*x)/2 + (2*a^2*sin[c + d*x])/d + (a^2*cos[c + d*x]*sin[c + d*x])/(2*d)))/3)/(4*a)
```

3.12.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3123 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^2, x_Symbol] := Simp[(2*a^2 + b^2)*(x/2), x] + (-Simp[2*a*b*(Cos[c + d*x]/d), x] - Simp[b^2*cos[c + d*x]*(Sin[c + d*x]/(2*d)), x]) /; FreeQ[{a, b, c, d}, x]
```

```
rule 3230 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

3.12. $\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$


```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.12.4 Maple [A] (verified)

Time = 2.84 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.57

method	result
parallelrisch	$\frac{\left(\frac{(A+B)\sin(2dx+2c)}{2} + \frac{\left(\frac{A}{2}+B\right)\sin(3dx+3c)}{6} + \frac{\sin(4dx+4c)B}{32} + \frac{\left(\frac{7A}{2}+3B\right)\sin(dx+c)}{2} + dx\left(A+\frac{7B}{8}\right) \right) a^2}{d}$
parts	$\frac{(A a^2+2B a^2)(2+\cos^2(dx+c))\sin(dx+c)}{3d} + \frac{(2A a^2+B a^2)\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{\sin(dx+c)A a^2}{d} + \frac{B a^2}{d}$
risch	$a^2 x A + \frac{7a^2 B x}{8} + \frac{7\sin(dx+c)A a^2}{4d} + \frac{3\sin(dx+c)B a^2}{2d} + \frac{\sin(4dx+4c)B a^2}{32d} + \frac{\sin(3dx+3c)A a^2}{12d} + \frac{\sin(3dx+3c)B a^2}{6d}$
derivativedivides	$\frac{A a^2(2+\cos^2(dx+c))\sin(dx+c)}{3} + B a^2 \left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2})\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 2A a^2 \left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
default	$\frac{A a^2(2+\cos^2(dx+c))\sin(dx+c)}{3} + B a^2 \left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2})\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 2A a^2 \left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
norman	$\frac{a^2(8A+7B)x}{8} + \frac{11a^2(8A+7B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{12d} + \frac{a^2(8A+7B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{4d} + \frac{a^2(8A+7B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2} + \frac{3a^2(8A+7B)x\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{4} + \frac{B a^2}{d}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (1/2*(A+B)*sin(2*d*x+2*c)+1/6*(1/2*A+B)*sin(3*d*x+3*c)+1/32*sin(4*d*x+4*c)
*B+1/2*(7/2*A+3*B)*sin(d*x+c)+d*x*(A+7/8*B))*a^2/d
```

$$3.12. \int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

3.12.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.70

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{3(8A + 7B)a^2 dx + (6Ba^2 \cos(dx + c))^3 + 8(A + 2B)a^2 \cos(dx + c)^2 + 3(8A + 7B)a^2 \cos(dx + c) + 8Aa^2}{24d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/24*(3*(8*A + 7*B)*a^2*d*x + (6*B*a^2*cos(d*x + c)^3 + 8*(A + 2*B)*a^2*cos(d*x + c)^2 + 3*(8*A + 7*B)*a^2*cos(d*x + c) + 8*(5*A + 4*B)*a^2)*sin(d*x + c))/d`

3.12.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 338 vs. 2(112) = 224.

Time = 0.19 (sec) , antiderivative size = 338, normalized size of antiderivative = 2.62

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \begin{cases} Aa^2x \sin^2(c + dx) + Aa^2x \cos^2(c + dx) + \frac{2Aa^2 \sin^3(c + dx)}{3d} + \frac{Aa^2 \sin(c + dx) \cos^2(c + dx)}{d} + \frac{Aa^2 \sin(c + dx) \cos(c + dx)}{d} \\ x(A + B \cos(c)) (a \cos(c) + a)^2 \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a**2*x*sin(c + d*x)**2 + A*a**2*x*cos(c + d*x)**2 + 2*A*a**2*sin(c + d*x)**3/(3*d) + A*a**2*sin(c + d*x)*cos(c + d*x)**2/d + A*a**2*sin(c + d*x)*cos(c + d*x)/d + A*a**2*sin(c + d*x)/d + 3*B*a**2*x*sin(c + d*x)**4/8 + 3*B*a**2*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + B*a**2*x*sin(c + d*x)**2/2 + 3*B*a**2*x*cos(c + d*x)**4/8 + B*a**2*x*cos(c + d*x)**2/2 + 3*B*a**2*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 4*B*a**2*sin(c + d*x)**3/(3*d) + 5*B*a**2*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 2*B*a**2*sin(c + d*x)*cos(c + d*x)**2/d + B*a**2*sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**2*cos(c), True))`

3.12.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.12

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx =$$

$$\frac{32 (\sin(dx + c)^3 - 3 \sin(dx + c))Aa^2 - 48(2dx + 2c + \sin(2dx + 2c))Aa^2 + 64 (\sin(dx + c)^3 - 3 \sin(dx + c))Ba^2 - 32(2dx + 2c + \sin(2dx + 2c))Ba^2 - 96Aa^2 \sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `-1/96*(32*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^2 - 48*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^2 + 64*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^2 - 3*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^2 - 24*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^2 - 96*A*a^2*sin(d*x + c))/d`

3.12.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.85

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^2 \sin(4dx + 4c)}{32d} + \frac{1}{8}(8Aa^2 + 7Ba^2)x + \frac{(Aa^2 + 2Ba^2) \sin(3dx + 3c)}{12d}$$

$$+ \frac{(Aa^2 + Ba^2) \sin(2dx + 2c)}{2d} + \frac{(7Aa^2 + 6Ba^2) \sin(dx + c)}{4d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/32*B*a^2*sin(4*d*x + 4*c)/d + 1/8*(8*A*a^2 + 7*B*a^2)*x + 1/12*(A*a^2 + 2*B*a^2)*sin(3*d*x + 3*c)/d + 1/2*(A*a^2 + B*a^2)*sin(2*d*x + 2*c)/d + 1/4*(7*A*a^2 + 6*B*a^2)*sin(d*x + c)/d`

3.12.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.04

$$\int \cos(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= A a^2 x + \frac{7 B a^2 x}{8} + \frac{7 A a^2 \sin(c + dx)}{4 d} + \frac{3 B a^2 \sin(c + dx)}{2 d}$$

$$+ \frac{A a^2 \sin(2 c + 2 d x)}{2 d} + \frac{A a^2 \sin(3 c + 3 d x)}{12 d} + \frac{B a^2 \sin(2 c + 2 d x)}{2 d}$$

$$+ \frac{B a^2 \sin(3 c + 3 d x)}{6 d} + \frac{B a^2 \sin(4 c + 4 d x)}{32 d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2,x)`output `A*a^2*x + (7*B*a^2*x)/8 + (7*A*a^2*sin(c + d*x))/(4*d) + (3*B*a^2*sin(c + d*x))/(2*d) + (A*a^2*sin(2*c + 2*d*x))/(2*d) + (A*a^2*sin(3*c + 3*d*x))/(12*d) + (B*a^2*sin(2*c + 2*d*x))/(2*d) + (B*a^2*sin(3*c + 3*d*x))/(6*d) + (B*a^2*sin(4*c + 4*d*x))/(32*d)`

3.13 $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$

3.13.1	Optimal result	318
3.13.2	Mathematica [A] (verified)	318
3.13.3	Rubi [A] (verified)	319
3.13.4	Maple [A] (verified)	320
3.13.5	Fricas [A] (verification not implemented)	321
3.13.6	Sympy [B] (verification not implemented)	321
3.13.7	Maxima [A] (verification not implemented)	322
3.13.8	Giac [A] (verification not implemented)	322
3.13.9	Mupad [B] (verification not implemented)	323

3.13.1 Optimal result

Integrand size = 23, antiderivative size = 94

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{1}{2} a^2 (3A + 2B)x + \frac{2a^2(3A + 2B) \sin(c + dx)}{3d}$$

$$+ \frac{a^2(3A + 2B) \cos(c + dx) \sin(c + dx)}{6d} + \frac{B(a + a \cos(c + dx))^2 \sin(c + dx)}{3d}$$

output `1/2*a^2*(3*A+2*B)*x+2/3*a^2*(3*A+2*B)*sin(d*x+c)/d+1/6*a^2*(3*A+2*B)*cos(d*x+c)*sin(d*x+c)/d+1/3*B*(a+a*cos(d*x+c))^2*sin(d*x+c)/d`

3.13.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.89

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{a^2 \sin(c + dx) \left(12A + 11B + 3(A + 2B) \cos(c + dx) + B \cos(2(c + dx)) \right) + \frac{6(3A+2B) \arcsin\left(\sqrt{\sin^2\left(\frac{1}{2}(c+dx)\right)}\right)}{\sqrt{\sin^2(c+dx)}}}{6d}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output $(a^2 \sin[c + dx] * (12A + 11B + 3(A + 2B) \cos[c + dx] + B \cos[2(c + dx)]) + (6(3A + 2B) \operatorname{ArcSin}[\sqrt{\sin[(c + dx)/2]^2}]) / \sqrt{\sin[c + dx]^2}) / (6d)$

3.13.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.88, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {3042, 3230, 3042, 3123}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \cos(c + dx) + a)^2 (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx \\ & \quad \downarrow \text{3230} \\ & \frac{1}{3}(3A + 2B) \int (\cos(c + dx)a + a)^2 dx + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^2}{3d} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{3}(3A + 2B) \int \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^2 dx + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^2}{3d} \\ & \quad \downarrow \text{3123} \\ & \frac{1}{3}(3A + 2B) \left(\frac{2a^2 \sin(c + dx)}{d} + \frac{a^2 \sin(c + dx) \cos(c + dx)}{2d} + \frac{3a^2 x}{2} \right) + \\ & \quad \frac{B \sin(c + dx)(a \cos(c + dx) + a)^2}{3d} \end{aligned}$$

input $\text{Int}[(a + a \cos[c + dx])^2 (A + B \cos[c + dx]), x]$

output $(B(a + a \cos[c + dx])^2 \sin[c + dx]) / (3d) + ((3A + 2B) * ((3a^2 x) / 2 + (2a^2 \sin[c + dx]) / d + (a^2 \cos[c + dx] * \sin[c + dx]) / (2d))) / 3$

3.13.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3123 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^2, x_Symbol] := Simp[(2*a^2 + b^
2)*(x/2), x] + (-Simp[2*a*b*(Cos[c + d*x]/d), x] - Simp[b^2*Cos[c + d*x]*(S
in[c + d*x]/(2*d)), x]) /; FreeQ[{a, b, c, d}, x]
```

```
rule 3230 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

3.13.4 Maple [A] (verified)

Time = 1.93 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.65

method	result
parallelrisch	$\frac{3\left(\left(\frac{A}{6} + \frac{B}{3}\right)\sin(2dx+2c) + \frac{B\sin(3dx+3c)}{18} + \left(\frac{4A}{3} + \frac{7B}{6}\right)\sin(dx+c) + dx\left(A + \frac{2B}{3}\right)\right)a^2}{2d}$
parts	$a^2xA + \frac{(Aa^2+2Ba^2)\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{(2Aa^2+Ba^2)\sin(dx+c)}{d} + \frac{Ba^2(2+\cos^2(dx+c))\sin(dx+c)}{3d}$
risch	$\frac{3a^2xA}{2} + a^2Bx + \frac{2\sin(dx+c)Aa^2}{d} + \frac{7\sin(dx+c)Ba^2}{4d} + \frac{\sin(3dx+3c)Ba^2}{12d} + \frac{\sin(2dx+2c)Aa^2}{4d} + \frac{\sin(2dx+2c)Ba^2}{2d}$
derivativdivides	$\frac{Aa^2\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + \frac{Ba^2(2+\cos^2(dx+c))\sin(dx+c)}{3} + 2Aa^2\sin(dx+c) + 2Ba^2\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
default	$\frac{Aa^2\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + \frac{Ba^2(2+\cos^2(dx+c))\sin(dx+c)}{3} + 2Aa^2\sin(dx+c) + 2Ba^2\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
norman	$\frac{\frac{a^2(3A+2B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d} + \frac{a^2(5A+6B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{a^2(3A+2B)x}{2} + \frac{8a^2(3A+2B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3d} + \frac{3a^2(3A+2B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^3}$

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 3/2*((1/6*A+1/3*B)*sin(2*d*x+2*c)+1/18*B*sin(3*d*x+3*c)+(4/3*A+7/6*B)*sin(
d*x+c)+d*x*(A+2/3*B))*a^2/d
```

3.13. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$

3.13.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.74

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{3(3A + 2B)a^2 dx + (2Ba^2 \cos(dx + c))^2 + 3(A + 2B)a^2 \cos(dx + c) + 2(6A + 5B)a^2 \sin(dx + c)}{6d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `1/6*(3*(3*A + 2*B)*a^2*d*x + (2*B*a^2*cos(d*x + c))^2 + 3*(A + 2*B)*a^2*cos(d*x + c) + 2*(6*A + 5*B)*a^2)*sin(d*x + c))/d`

3.13.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 199 vs. $2(85) = 170$.

Time = 0.14 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.12

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aa^2 x \sin^2(c+dx)}{2} + \frac{Aa^2 x \cos^2(c+dx)}{2} + Aa^2 x + \frac{Aa^2 \sin(c+dx) \cos(c+dx)}{2d} + \frac{2Aa^2 \sin(c+dx)}{d} + Ba^2 x \sin^2(c + dx) + Ba^2 x \cos^2(c + dx) \\ x(A + B \cos(c)) (a \cos(c) + a)^2 \end{cases}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a**2*x*sin(c + d*x)**2/2 + A*a**2*x*cos(c + d*x)**2/2 + A*a**2*x + A*a**2*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*A*a**2*sin(c + d*x)/d + B*a**2*x*sin(c + d*x)**2 + B*a**2*x*cos(c + d*x)**2 + 2*B*a**2*sin(c + d*x)**3/(3*d) + B*a**2*sin(c + d*x)*cos(c + d*x)**2/d + B*a**2*sin(c + d*x)*cos(c + d*x)/d + B*a**2*sin(c + d*x)/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**2, True))`

3.13.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.17

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{3(2dx + 2c + \sin(2dx + 2c))Aa^2 + 12(dx + c)Aa^2 - 4(\sin(dx + c)^3 - 3\sin(dx + c))Ba^2 + 6(2dx + 2c + \sin(2dx + 2c))Ba^2 + 24Aa^2 \sin(dx + c) + 12Ba^2 \sin(dx + c)}{12d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")`output `1/12*(3*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^2 + 12*(d*x + c)*A*a^2 - 4*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^2 + 6*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^2 + 24*A*a^2*sin(d*x + c) + 12*B*a^2*sin(d*x + c))/d`**3.13.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.90

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{Ba^2 \sin(3dx + 3c)}{12d} + \frac{1}{2} (3Aa^2 + 2Ba^2)x$$

$$+ \frac{(Aa^2 + 2Ba^2) \sin(2dx + 2c)}{4d} + \frac{(8Aa^2 + 7Ba^2) \sin(dx + c)}{4d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/12*B*a^2*sin(3*d*x + 3*c)/d + 1/2*(3*A*a^2 + 2*B*a^2)*x + 1/4*(A*a^2 + 2*B*a^2)*sin(2*d*x + 2*c)/d + 1/4*(8*A*a^2 + 7*B*a^2)*sin(d*x + c)/d`

3.13.9 Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.04

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{3 A a^2 x}{2} + B a^2 x + \frac{2 A a^2 \sin(c + dx)}{d} + \frac{7 B a^2 \sin(c + dx)}{4 d}$$

$$+ \frac{A a^2 \sin(2c + 2dx)}{4 d} + \frac{B a^2 \sin(2c + 2dx)}{2 d} + \frac{B a^2 \sin(3c + 3dx)}{12 d}$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2,x)`output `(3*A*a^2*x)/2 + B*a^2*x + (2*A*a^2*sin(c + d*x))/d + (7*B*a^2*sin(c + d*x))/
(4*d) + (A*a^2*sin(2*c + 2*d*x))/(4*d) + (B*a^2*sin(2*c + 2*d*x))/(2*d)
+ (B*a^2*sin(3*c + 3*d*x))/(12*d)`

3.14 $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$

3.14.1	Optimal result	324
3.14.2	Mathematica [A] (verified)	324
3.14.3	Rubi [A] (verified)	325
3.14.4	Maple [A] (verified)	328
3.14.5	Fricas [A] (verification not implemented)	328
3.14.6	Sympy [F]	329
3.14.7	Maxima [A] (verification not implemented)	329
3.14.8	Giac [A] (verification not implemented)	330
3.14.9	Mupad [B] (verification not implemented)	330

3.14.1 Optimal result

Integrand size = 29, antiderivative size = 82

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{1}{2} a^2 (4A + 3B)x + \frac{a^2 A \operatorname{arctanh}(\sin(c + dx))}{d} \\ & \quad + \frac{a^2 (2A + 3B) \sin(c + dx)}{2d} + \frac{B(a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{2d} \end{aligned}$$

```
output 1/2*a^2*(4*A+3*B)*x+a^2*A*arctanh(sin(d*x+c))/d+1/2*a^2*(2*A+3*B)*sin(d*x+c)/d+1/2*B*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d
```

3.14.2 Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.17

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{a^2 (8A dx + 6B dx - 4A \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 4A \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{4d} \end{aligned}$$

```
input Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x],x]
```

output $(a^2(8Adx + 6Bdx - 4A \log[\cos[(c + dx)/2] - \sin[(c + dx)/2]] + 4A \log[\cos[(c + dx)/2] + \sin[(c + dx)/2]] + 4(A + 2B)\sin[c + dx] + B \sin[2(c + dx)])/(4d)$

3.14.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 3455, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{3455} \\ & \frac{1}{2} \int (\cos(c + dx)a + a)(2aA + a(2A + 3B) \cos(c + dx)) \sec(c + dx) dx + \\ & \quad \frac{B \sin(c + dx)(a^2 \cos(c + dx) + a^2)}{2d} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)(2aA + a(2A + 3B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \\ & \quad \frac{B \sin(c + dx)(a^2 \cos(c + dx) + a^2)}{2d} \\ & \quad \downarrow \text{3447} \\ & \frac{1}{2} \int ((2A + 3B) \cos^2(c + dx)a^2 + 2Aa^2 + (2Aa^2 + (2A + 3B)a^2) \cos(c + dx)) \sec(c + dx) dx + \\ & \quad \frac{B \sin(c + dx)(a^2 \cos(c + dx) + a^2)}{2d} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2} \int \frac{(2A + 3B) \sin(c + dx + \frac{\pi}{2})^2 a^2 + 2Aa^2 + (2Aa^2 + (2A + 3B)a^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \\ & \quad \frac{B \sin(c + dx)(a^2 \cos(c + dx) + a^2)}{2d} \end{aligned}$$

3.14. $\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec(c + dx) dx$

$$\begin{aligned}
& \downarrow \text{3502} \\
& \frac{1}{2} \left(\int (2Aa^2 + (4A + 3B) \cos(c + dx)a^2) \sec(c + dx) dx + \frac{a^2(2A + 3B) \sin(c + dx)}{d} \right) + \\
& \quad \frac{B \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \downarrow \text{3042} \\
& \frac{1}{2} \left(\int \frac{2Aa^2 + (4A + 3B) \sin(c + dx + \frac{\pi}{2}) a^2}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a^2(2A + 3B) \sin(c + dx)}{d} \right) + \\
& \quad \frac{B \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \downarrow \text{3214} \\
& \frac{1}{2} \left(2a^2 A \int \sec(c + dx) dx + \frac{a^2(2A + 3B) \sin(c + dx)}{d} + a^2 x(4A + 3B) \right) + \\
& \quad \frac{B \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \downarrow \text{3042} \\
& \frac{1}{2} \left(2a^2 A \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{a^2(2A + 3B) \sin(c + dx)}{d} + a^2 x(4A + 3B) \right) + \\
& \quad \frac{B \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \downarrow \text{4257} \\
& \frac{1}{2} \left(\frac{2a^2 A \operatorname{Arctanh}(\sin(c + dx))}{d} + \frac{a^2(2A + 3B) \sin(c + dx)}{d} + a^2 x(4A + 3B) \right) + \\
& \quad \frac{B \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{2d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(B*(a^2 + a^2*Cos[c + d*x])*Sin[c + d*x])/(2*d) + (a^2*(4*A + 3*B)*x + (2*a^2*A*ArcTanh[Sin[c + d*x]]))/d + (a^2*(2*A + 3*B)*Sin[c + d*x])/d/2`

3.14.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3455 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.14.4 Maple [A] (verified)

Time = 1.53 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.90

method	result
parallelrisc	$-\frac{\left(A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{\sin(2dx+2c)B}{4} + (-A-2B) \sin(dx+c) - 2dx\left(A + \frac{3B}{4}\right)\right) a^2}{d}$
derivativedivides	$\frac{A a^2 \sin(dx+c) + B a^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + 2A a^2(dx+c) + 2B a^2 \sin(dx+c) + A a^2 \ln(\sec(dx+c) + \tan(dx+c))}{d}$
default	$\frac{A a^2 \sin(dx+c) + B a^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + 2A a^2(dx+c) + 2B a^2 \sin(dx+c) + A a^2 \ln(\sec(dx+c) + \tan(dx+c))}{d}$
parts	$\frac{A a^2 \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(A a^2 + 2B a^2) \sin(dx+c)}{d} + \frac{(2A a^2 + B a^2)(dx+c)}{d} + \frac{B a^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
risc	$2a^2 x A + \frac{3a^2 B x}{2} - \frac{ie^{i(dx+c)} A a^2}{2d} - \frac{ie^{i(dx+c)} B a^2}{d} + \frac{ie^{-i(dx+c)} A a^2}{2d} + \frac{ie^{-i(dx+c)} B a^2}{d} + \frac{A a^2 \ln(e^{i(dx+c)} + 1)}{d}$
norman	$\frac{(2A a^2 + \frac{3}{2} B a^2) x + (2A a^2 + \frac{3}{2} B a^2) x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (6A a^2 + \frac{9}{2} B a^2) x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (6A a^2 + \frac{9}{2} B a^2) x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^3}$

input `int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output $-(A*\ln(\tan(1/2*d*x+1/2*c)-1)-A*\ln(\tan(1/2*d*x+1/2*c)+1)-1/4*\sin(2*d*x+2*c)*B+(-A-2*B)*\sin(d*x+c)-2*d*x*(A+3/4*B))*a^2/d$

3.14.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.96

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{(4A + 3B)a^2 dx + Aa^2 \log(\sin(dx + c) + 1) - Aa^2 \log(-\sin(dx + c) + 1) + (Ba^2 \cos(dx + c) + 2(A + B)a^2 \sin(dx + c))}{2d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fracas")`

output $1/2*((4*A + 3*B)*a^2*d*x + A*a^2*\log(\sin(d*x + c) + 1) - A*a^2*\log(-\sin(d*x + c) + 1) + (B*a^2*\cos(d*x + c) + 2*(A + 2*B)*a^2)*\sin(d*x + c))/d$

3.14. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$

3.14.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= a^2 \left(\int A \sec(c + dx) dx + \int 2A \cos(c + dx) \sec(c + dx) dx \right. \\ &\quad \left. + \int A \cos^2(c + dx) \sec(c + dx) dx + \int B \cos(c + dx) \sec(c + dx) dx \right. \\ &\quad \left. + \int 2B \cos^2(c + dx) \sec(c + dx) dx + \int B \cos^3(c + dx) \sec(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `a**2*(Integral(A*sec(c + d*x), x) + Integral(2*A*cos(c + d*x)*sec(c + d*x), x) + Integral(A*cos(c + d*x)**2*sec(c + d*x), x) + Integral(B*cos(c + d*x)*sec(c + d*x), x) + Integral(2*B*cos(c + d*x)**2*sec(c + d*x), x) + Integral(B*cos(c + d*x)**3*sec(c + d*x), x))`

3.14.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.15

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{8(dx + c)Aa^2 + (2dx + 2c + \sin(2dx + 2c))Ba^2 + 4(dx + c)Ba^2 + 4Aa^2 \log(\sec(dx + c) + \tan(dx + c)) + 8Ba^2 \sin(dx + c)}{4d} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `1/4*(8*(d*x + c)*A*a^2 + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^2 + 4*(d*x + c)*B*a^2 + 4*A*a^2*log(sec(d*x + c) + tan(d*x + c)) + 4*A*a^2*sin(d*x + c) + 8*B*a^2*sin(d*x + c))/d`

3.14.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.77

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2 A a^2 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 2 A a^2 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + (4 A a^2 + 3 B a^2)(dx + c) + \frac{2(2 A a^2 + 3 B a^2) \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{2 d}}{2 d}$$

```
input integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")
```

```
output 1/2*(2*A*a^2*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 2*A*a^2*log(abs(tan(1/2*d*x + 1/2*c) - 1))) + (4*A*a^2 + 3*B*a^2)*(d*x + c) + 2*(2*A*a^2*tan(1/2*d*x + 1/2*c)^3 + 3*B*a^2*tan(1/2*d*x + 1/2*c)^3 + 2*A*a^2*tan(1/2*d*x + 1/2*c) + 5*B*a^2*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^2/d
```

3.14.9 Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.72

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{A a^2 \sin(c + dx)}{d} + \frac{2 B a^2 \sin(c + dx)}{d} + \frac{4 A a^2 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d}$$

$$+ \frac{2 A a^2 \operatorname{atanh} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} + \frac{3 B a^2 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} + \frac{B a^2 \sin(2c + 2dx)}{4d}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x),x)
```

```
output (A*a^2*sin(c + d*x))/d + (2*B*a^2*sin(c + d*x))/d + (4*A*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*A*a^2*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (3*B*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (B*a^2*sin(2*c + 2*d*x))/(4*d)
```

3.15 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.15.1	Optimal result	331
3.15.2	Mathematica [A] (verified)	331
3.15.3	Rubi [A] (verified)	332
3.15.4	Maple [A] (verified)	335
3.15.5	Fricas [A] (verification not implemented)	335
3.15.6	Sympy [F]	336
3.15.7	Maxima [A] (verification not implemented)	336
3.15.8	Giac [B] (verification not implemented)	337
3.15.9	Mupad [B] (verification not implemented)	337

3.15.1 Optimal result

Integrand size = 31, antiderivative size = 74

$$\begin{aligned} & \int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= a^2(A + 2B)x + \frac{a^2(2A + B)\operatorname{arctanh}(\sin(c + dx))}{d} \\ & \quad - \frac{a^2(A - B)\sin(c + dx)}{d} + \frac{A(a^2 + a^2 \cos(c + dx)) \tan(c + dx)}{d} \end{aligned}$$

```
output a^2*(A+2*B)*x+a^2*(2*A+B)*arctanh(sin(d*x+c))/d-a^2*(A-B)*sin(d*x+c)/d+A*(
a^2+a^2*cos(d*x+c))*tan(d*x+c)/d
```

3.15.2 Mathematica [A] (verified)

Time = 1.34 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.93

$$\begin{aligned} & \int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= \frac{a^2(Ac + 2Bc + Adx + 2Bdx - 2A \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) - B \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{d} \end{aligned}$$

```
input Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]
```

output $(a^2(Ac + 2Bc + Adx + 2Bdx - 2A\text{Log}[\text{Cos}[(c + dx)/2] - \text{Sin}[(c + dx)/2]] - B\text{Log}[\text{Cos}[(c + dx)/2] - \text{Sin}[(c + dx)/2]] + 2A\text{Log}[\text{Cos}[(c + dx)/2] + \text{Sin}[(c + dx)/2]] + B\text{Log}[\text{Cos}[(c + dx)/2] + \text{Sin}[(c + dx)/2]] + B\text{Sin}[c + dx] + A\text{Tan}[c + dx])/d$

3.15.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3454, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx$$

$$\downarrow \text{3454}$$

$$\int (\cos(c + dx)a + a)(a(2A + B) - a(A - B) \cos(c + dx)) \sec(c + dx) dx + \frac{A \tan(c + dx) (a^2 \cos(c + dx) + a^2)}{d}$$

$$\downarrow \text{3042}$$

$$\int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)(a(2A + B) - a(A - B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{A \tan(c + dx) (a^2 \cos(c + dx) + a^2)}{d}$$

$$\downarrow \text{3447}$$

$$\int (-((A - B) \cos^2(c + dx)a^2) + (2A + B)a^2 + (a^2(2A + B) - a^2(A - B)) \cos(c + dx)) \sec(c + dx) dx + \frac{A \tan(c + dx) (a^2 \cos(c + dx) + a^2)}{d}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \frac{-\left((A-B)\sin\left(c+dx+\frac{\pi}{2}\right)^2 a^2\right) + (2A+B)a^2 + (a^2(2A+B) - a^2(A-B))\sin\left(c+dx+\frac{\pi}{2}\right)}{\frac{\sin\left(c+dx+\frac{\pi}{2}\right)}{A \tan(c+dx) (a^2 \cos(c+dx) + a^2)}} dx + \\
& \qquad \qquad \qquad \downarrow \text{3502} \\
& \int \left(\frac{(2A+B)a^2 + (A+2B)\cos(c+dx)a^2}{A \tan(c+dx) (a^2 \cos(c+dx) + a^2)}\right) \sec(c+dx) dx - \frac{a^2(A-B)\sin(c+dx)}{d} + \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \int \frac{(2A+B)a^2 + (A+2B)\sin\left(c+dx+\frac{\pi}{2}\right) a^2}{\sin\left(c+dx+\frac{\pi}{2}\right) A \tan(c+dx) (a^2 \cos(c+dx) + a^2)} dx - \frac{a^2(A-B)\sin(c+dx)}{d} + \\
& \qquad \qquad \qquad \downarrow \text{3214} \\
& a^2(2A+B) \int \sec(c+dx) dx - \frac{a^2(A-B)\sin(c+dx)}{d} + a^2x(A+2B) + \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& a^2(2A+B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx - \frac{a^2(A-B)\sin(c+dx)}{d} + a^2x(A+2B) + \\
& \qquad \qquad \qquad \downarrow \text{4257} \\
& \frac{a^2(2A+B)\operatorname{arctanh}(\sin(c+dx))}{d} - \frac{a^2(A-B)\sin(c+dx)}{d} + a^2x(A+2B) + \\
& \qquad \qquad \qquad \frac{A \tan(c+dx) (a^2 \cos(c+dx) + a^2)}{d}
\end{aligned}$$

input `Int[(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x])*Sec[c + d*x]^2,x]`

output `a^2*(A + 2*B)*x + (a^2*(2*A + B)*ArcTanh[Sin[c + d*x]])/d - (a^2*(A - B)*Sin[c + d*x])/d + (A*(a^2 + a^2*cos[c + d*x])*Tan[c + d*x])/d`

3.15.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.15.4 Maple [A] (verified)

Time = 2.47 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.09

method	result
parts	$\frac{a^2 A \tan(dx+c)}{d} + \frac{(A a^2+2B a^2)(dx+c)}{d} + \frac{(2A a^2+B a^2) \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{\sin(dx+c)B a^2}{d}$
derivativedivides	$\frac{A a^2(dx+c)+B a^2 \sin(dx+c)+2A a^2 \ln(\sec(dx+c)+\tan(dx+c))+2B a^2(dx+c)+A a^2 \tan(dx+c)+B a^2 \ln(\sec(dx+c)+\tan(dx+c))}{d}$
default	$\frac{A a^2(dx+c)+B a^2 \sin(dx+c)+2A a^2 \ln(\sec(dx+c)+\tan(dx+c))+2B a^2(dx+c)+A a^2 \tan(dx+c)+B a^2 \ln(\sec(dx+c)+\tan(dx+c))}{d}$
parallelrisch	$-\frac{2a^2 \left(\cos(dx+c) \left(A + \frac{B}{2} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) - \cos(dx+c) \left(A + \frac{B}{2} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \frac{\sin(2dx+2c)B}{4} - \frac{dx(A+2B)\cos(dx+c)}{2}}{d \cos(dx+c)}$
risch	$a^2 x A + 2a^2 B x - \frac{ie^{i(dx+c)} B a^2}{2d} + \frac{ie^{-i(dx+c)} B a^2}{2d} + \frac{2iA a^2}{d(e^{2i(dx+c)}+1)} - \frac{2A a^2 \ln(e^{i(dx+c)}-i)}{d} - \frac{a^2 \ln(e^{i(dx+c)}+i)}{d}$
norman	$\frac{(-A a^2-2B a^2)x+(-2A a^2-4B a^2)x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (A a^2+2B a^2)x \left(\tan^8 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (2A a^2+4B a^2)x \left(\tan^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{\left(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^3}$

```
input int((a+cos(d*x+c))*a^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBOSE)
```

```
output a^2*A*tan(d*x+c)/d+(A*a^2+2*B*a^2)/d*(d*x+c)+(2*A*a^2+B*a^2)/d*ln(sec(d*x+c)+tan(d*x+c))+1/d*sin(d*x+c)*B*a^2
```

3.15.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.46

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2(A + 2B)a^2 dx \cos(dx + c) + (2A + B)a^2 \cos(dx + c) \log(\sin(dx + c) + 1) - (2A + B)a^2 \cos(dx + c)}{2d \cos(dx + c)}$$

```
input integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fracas")
```

```
output 1/2*(2*(A + 2*B)*a^2*d*x*cos(d*x + c) + (2*A + B)*a^2*cos(d*x + c)*log(sin(d*x + c) + 1) - (2*A + B)*a^2*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*(B*a^2*cos(d*x + c) + A*a^2)*sin(d*x + c))/(d*cos(d*x + c))
```

3.15. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.15.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= a^2 \left(\int A \sec^2(c + dx) dx + \int 2A \cos(c + dx) \sec^2(c + dx) dx \right. \\ &\quad \left. + \int A \cos^2(c + dx) \sec^2(c + dx) dx + \int B \cos(c + dx) \sec^2(c + dx) dx \right. \\ &\quad \left. + \int 2B \cos^2(c + dx) \sec^2(c + dx) dx + \int B \cos^3(c + dx) \sec^2(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `a**2*(Integral(A*sec(c + d*x)**2, x) + Integral(2*A*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(A*cos(c + d*x)**2*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(2*B*cos(c + d*x)**2*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)**3*sec(c + d*x)**2, x))`

3.15.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.42

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= \frac{2(dx + c)Aa^2 + 4(dx + c)Ba^2 + 2Aa^2(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + Ba^2(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 2Ba^2 \sin(dx + c) + 2Aa^2 \tan(dx + c)}{2d} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*A*a^2 + 4*(d*x + c)*B*a^2 + 2*A*a^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + B*a^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*B*a^2*sin(d*x + c) + 2*A*a^2*tan(d*x + c))/d`

3.15.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 155 vs. 2(74) = 148.

Time = 0.31 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.09

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{(Aa^2 + 2Ba^2)(dx + c) + (2Aa^2 + Ba^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - (2Aa^2 + Ba^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `((A*a^2 + 2*B*a^2)*(d*x + c) + (2*A*a^2 + B*a^2)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (2*A*a^2 + B*a^2)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(A*a^2*tan(1/2*d*x + 1/2*c)^3 - B*a^2*tan(1/2*d*x + 1/2*c)^3 + A*a^2*tan(1/2*d*x + 1/2*c) + B*a^2*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^4 - 1))/d`

3.15.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.18

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{B a^2 \sin(c + dx)}{d} + \frac{2 A a^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{4 A a^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{4 B a^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2 B a^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{A a^2 \sin(c + dx)}{d \cos(c + dx)}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^2,x)`

output `(B*a^2*sin(c + d*x))/d + (2*A*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2))/d + (4*A*a^2*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2))/d + (4*B*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a^2*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (A*a^2*sin(c + d*x))/(d*cos(c + d*x))`

3.16 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.16.1	Optimal result	338
3.16.2	Mathematica [A] (verified)	338
3.16.3	Rubi [A] (verified)	339
3.16.4	Maple [A] (verified)	342
3.16.5	Fricas [A] (verification not implemented)	342
3.16.6	Sympy [F]	343
3.16.7	Maxima [A] (verification not implemented)	343
3.16.8	Giac [A] (verification not implemented)	344
3.16.9	Mupad [B] (verification not implemented)	344

3.16.1 Optimal result

Integrand size = 31, antiderivative size = 88

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= a^2 Bx + \frac{a^2(3A + 4B)\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{a^2(3A + 2B) \tan(c + dx)}{2d}$$

$$+ \frac{A(a^2 + a^2 \cos(c + dx)) \sec(c + dx) \tan(c + dx)}{2d}$$

```
output a^2*B*x+1/2*a^2*(3*A+4*B)*arctanh(sin(d*x+c))/d+1/2*a^2*(3*A+2*B)*tan(d*x+c)/d+1/2*A*(a^2+a^2*cos(d*x+c))*sec(d*x+c)*tan(d*x+c)/d
```

3.16.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.60

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{a^2(2Bdx + (3A + 4B)\operatorname{arctanh}(\sin(c + dx)) + (4A + 2B + A \sec(c + dx)) \tan(c + dx))}{2d}$$

```
input Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]
```

output $(a^2(2Bdx + (3A + 4B)\text{ArcTanh}[\text{Sin}[c + dx]] + (4A + 2B + A\text{Sec}[c + dx])\text{Tan}[c + dx]))/(2d)$

3.16.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3454, 3042, 3447, 3042, 3500, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3454} \\
 & \frac{1}{2} \int (\cos(c + dx)a + a)(a(3A + 2B) + 2aB \cos(c + dx)) \sec^2(c + dx) dx + \\
 & \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)(a(3A + 2B) + 2aB \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx + \\
 & \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
 & \quad \downarrow \text{3447} \\
 & \frac{1}{2} \int (2B \cos^2(c + dx)a^2 + (3A + 2B)a^2 + (2Ba^2 + (3A + 2B)a^2) \cos(c + dx)) \sec^2(c + dx) dx + \\
 & \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{2B \sin(c + dx + \frac{\pi}{2})^2 a^2 + (3A + 2B)a^2 + (2Ba^2 + (3A + 2B)a^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx + \\
 & \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d}
 \end{aligned}$$

3.16. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

$$\begin{aligned}
& \downarrow \text{3500} \\
& \frac{1}{2} \left(\int \left((3A + 4B)a^2 + 2B \cos(c + dx)a^2 \right) \sec(c + dx) dx + \frac{a^2(3A + 2B) \tan(c + dx)}{d} \right) + \\
& \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\int \frac{(3A + 4B)a^2 + 2B \sin(c + dx + \frac{\pi}{2}) a^2}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a^2(3A + 2B) \tan(c + dx)}{d} \right) + \\
& \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \quad \downarrow \text{3214} \\
& \frac{1}{2} \left(a^2(3A + 4B) \int \sec(c + dx) dx + \frac{a^2(3A + 2B) \tan(c + dx)}{d} + 2a^2 Bx \right) + \\
& \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(a^2(3A + 4B) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{a^2(3A + 2B) \tan(c + dx)}{d} + 2a^2 Bx \right) + \\
& \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d} \\
& \quad \downarrow \text{4257} \\
& \frac{1}{2} \left(\frac{a^2(3A + 4B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{a^2(3A + 2B) \tan(c + dx)}{d} + 2a^2 Bx \right) + \\
& \quad \frac{A \tan(c + dx) \sec(c + dx) (a^2 \cos(c + dx) + a^2)}{2d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(A*(a^2 + a^2*Cos[c + d*x])*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (2*a^2*B*x + (a^2*(3*A + 4*B)*ArcTanh[Sin[c + d*x]])/d + (a^2*(3*A + 2*B)*Tan[c + d*x])/d)/2`

3.16.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.16.4 Maple [A] (verified)

Time = 3.29 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.20

method	result
parts	$\frac{A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d} + \frac{(A a^2 + 2B a^2) \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(2A a^2 + B a^2) \tan(dx+c)}{d}$
derivativedivides	$\frac{A a^2 \ln(\sec(dx+c) + \tan(dx+c)) + B a^2 (dx+c) + 2A a^2 \tan(dx+c) + 2B a^2 \ln(\sec(dx+c) + \tan(dx+c)) + A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
default	$\frac{A a^2 \ln(\sec(dx+c) + \tan(dx+c)) + B a^2 (dx+c) + 2A a^2 \tan(dx+c) + 2B a^2 \ln(\sec(dx+c) + \tan(dx+c)) + A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
parallelrisch	$-\frac{3 \left(\left(A + \frac{4B}{3} \right) (1 + \cos(2dx+2c)) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) - \left(A + \frac{4B}{3} \right) (1 + \cos(2dx+2c)) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \frac{2dx B \cos(2dx+2c)}{3} \right)}{2d(1 + \cos(2dx+2c))}$
risch	$a^2 B x - \frac{ia^2 (A e^{3i(dx+c)} - 4A e^{2i(dx+c)} - 2B e^{2i(dx+c)} - A e^{i(dx+c)} - 4A - 2B)}{d(e^{2i(dx+c)} + 1)^2} - \frac{3A a^2 \ln(e^{i(dx+c)} - i)}{2d} - \frac{2a^2 \ln(e^{i(dx+c)} + i)}{2d}$
norman	$\frac{a^2 B x + a^2 B x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + a^2 B x \left(\tan^8 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + a^2 B x \left(\tan^{10} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \frac{a^2 (5A + 2B) \tan \left(\frac{dx}{2} + \frac{c}{2} \right)}{d} + \frac{6A a^2 \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d}}{(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right))^2}$

input `int((a+cos(d*x+c))*a^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `A*a^2/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(A*a^2+2*B*a^2)/d*ln(sec(d*x+c)+tan(d*x+c))+(2*A*a^2+B*a^2)/d*tan(d*x+c)+B*a^2/d*(d*x+c)`

3.16.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.35

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4 B a^2 dx \cos(dx + c)^2 + (3 A + 4 B) a^2 \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (3 A + 4 B) a^2 \cos(dx + c)^2 \log(\sin(dx + c) - 1)}{4 d \cos(dx + c)^2}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fracas")`

output `1/4*(4*B*a^2*d*x*cos(d*x + c)^2 + (3*A + 4*B)*a^2*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (3*A + 4*B)*a^2*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(2*(2*A + B)*a^2*cos(d*x + c) + A*a^2)*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.16. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.16.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= a^2 \left(\int A \sec^3(c + dx) dx + \int 2A \cos(c + dx) \sec^3(c + dx) dx \right. \\ & \quad \left. + \int A \cos^2(c + dx) \sec^3(c + dx) dx + \int B \cos(c + dx) \sec^3(c + dx) dx \right. \\ & \quad \left. + \int 2B \cos^2(c + dx) \sec^3(c + dx) dx + \int B \cos^3(c + dx) \sec^3(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `a**2*(Integral(A*sec(c + d*x)**3, x) + Integral(2*A*cos(c + d*x)*sec(c + d*x)**3, x) + Integral(A*cos(c + d*x)**2*sec(c + d*x)**3, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**3, x) + Integral(2*B*cos(c + d*x)**2*sec(c + d*x)**3, x) + Integral(B*cos(c + d*x)**3*sec(c + d*x)**3, x))`

3.16.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.61

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= \frac{4(dx+c)Ba^2 - Aa^2 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c)+1) + \log(\sin(dx+c)-1) \right) + 2Aa^2(\log(\sin(dx+c)+1) - \log(\sin(dx+c)-1)) + 4B*a^2*(\log(\sin(dx+c)+1) - \log(\sin(dx+c)-1)) + 8*A*a^2*\tan(dx+c) + 4*B*a^2*\tan(dx+c))/d}{1} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `1/4*(4*(d*x + c)*B*a^2 - A*a^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 2*A*a^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 4*B*a^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 8*A*a^2*tan(d*x + c) + 4*B*a^2*tan(d*x + c))/d`

3.16.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.75

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2(dx + c)Ba^2 + (3Aa^2 + 4Ba^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - (3Aa^2 + 4Ba^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right)}{2d}$$

```
input integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")
```

```
output 1/2*(2*(d*x + c)*B*a^2 + (3*A*a^2 + 4*B*a^2)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (3*A*a^2 + 4*B*a^2)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(3*A*a^2*tan(1/2*d*x + 1/2*c)^3 + 2*B*a^2*tan(1/2*d*x + 1/2*c)^3 - 5*A*a^2*tan(1/2*d*x + 1/2*c) - 2*B*a^2*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^2)/d
```

3.16.9 Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.84

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{3Aa^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2Ba^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{4Ba^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{2Aa^2 \sin(c + dx)}{d \cos(c + dx)} + \frac{Aa^2 \sin(c + dx)}{2d \cos(c + dx)^2} + \frac{Ba^2 \sin(c + dx)}{d \cos(c + dx)}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^3,x)
```

```
output (3*A*a^2*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (4*B*a^2*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*A*a^2*sin(c + d*x))/(d*cos(c + d*x)) + (A*a^2*sin(c + d*x))/(2*d*cos(c + d*x)^2) + (B*a^2*sin(c + d*x))/(d*cos(c + d*x))
```

3.17 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.17.1	Optimal result	345
3.17.2	Mathematica [A] (verified)	346
3.17.3	Rubi [A] (verified)	346
3.17.4	Maple [A] (verified)	350
3.17.5	Fricas [A] (verification not implemented)	350
3.17.6	Sympy [F]	351
3.17.7	Maxima [A] (verification not implemented)	351
3.17.8	Giac [A] (verification not implemented)	352
3.17.9	Mupad [B] (verification not implemented)	352

3.17.1 Optimal result

Integrand size = 31, antiderivative size = 113

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{a^2(2A + 3B)\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{a^2(5A + 6B) \tan(c + dx)}{3d}$$

$$+ \frac{a^2(4A + 3B) \sec(c + dx) \tan(c + dx)}{6d}$$

$$+ \frac{A(a^2 + a^2 \cos(c + dx)) \sec^2(c + dx) \tan(c + dx)}{3d}$$

```
output 1/2*a^2*(2*A+3*B)*arctanh(sin(d*x+c))/d+1/3*a^2*(5*A+6*B)*tan(d*x+c)/d+1/6
*a^2*(4*A+3*B)*sec(d*x+c)*tan(d*x+c)/d+1/3*A*(a^2+a^2*cos(d*x+c))*sec(d*x+
c)^2*tan(d*x+c)/d
```


3.17.2 Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.56

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{a^2((6A + 9B)\operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (12(A + B) + 3(2A + B) \sec(c + dx) + 2A \tan^2(c + dx)))}{6d}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a^2*((6*A + 9*B)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(12*(A + B) + 3*(2*A + B)*Sec[c + d*x] + 2*A*Tan[c + d*x]^2)))/(6*d)`

3.17.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3454, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c + dx)(a \cos(c + dx) + a)^2 (A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx$$

$$\downarrow 3454$$

$$\frac{1}{3} \int (\cos(c + dx)a + a)(a(4A + 3B) + a(A + 3B) \cos(c + dx)) \sec^3(c + dx) dx + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow 3042$$

$$\frac{1}{3} \int \frac{(\sin(c + dx + \frac{\pi}{2}) a + a) (a(4A + 3B) + a(A + 3B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

3.17. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

$$\downarrow \text{3447}$$

$$\frac{1}{3} \int ((A + 3B) \cos^2(c + dx)a^2 + (4A + 3B)a^2 + ((A + 3B)a^2 + (4A + 3B)a^2) \cos(c + dx)) \sec^3(c + dx) dx + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \int \frac{(A + 3B) \sin(c + dx + \frac{\pi}{2})^2 a^2 + (4A + 3B)a^2 + ((A + 3B)a^2 + (4A + 3B)a^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow \text{3500}$$

$$\frac{1}{3} \left(\frac{1}{2} \int (2(5A + 6B)a^2 + 3(2A + 3B) \cos(c + dx)a^2) \sec^2(c + dx) dx + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \left(\frac{1}{2} \int \frac{2(5A + 6B)a^2 + 3(2A + 3B) \sin(c + dx + \frac{\pi}{2}) a^2}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow \text{3227}$$

$$\frac{1}{3} \left(\frac{1}{2} \left(2a^2(5A + 6B) \int \sec^2(c + dx) dx + 3a^2(2A + 3B) \int \sec(c + dx) dx \right) + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3a^2(2A + 3B) \int \csc(c + dx + \frac{\pi}{2}) dx + 2a^2(5A + 6B) \int \csc(c + dx + \frac{\pi}{2})^2 dx \right) + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

$$\downarrow \text{4254}$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3a^2(2A + 3B) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx - \frac{2a^2(5A + 6B) \int 1d(-\tan(c + dx))}{d} \right) + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right)$$

$$\frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

↓ 24

$$\frac{1}{3} \left(\frac{1}{2} \left(3a^2(2A + 3B) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{2a^2(5A + 6B) \tan(c + dx)}{d} \right) + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right)$$

$$\frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{3a^2(2A + 3B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{2a^2(5A + 6B) \tan(c + dx)}{d} \right) + \frac{a^2(4A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right)$$

$$\frac{A \tan(c + dx) \sec^2(c + dx) (a^2 \cos(c + dx) + a^2)}{3d}$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(A*(a^2 + a^2*Cos[c + d*x])*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((a^2*(4*A + 3*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((3*a^2*(2*A + 3*B)*ArcTanh[Sin[c + d*x]])/d + (2*a^2*(5*A + 6*B)*Tan[c + d*x])/d)/2)/3`

3.17.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.17.4 Maple [A] (verified)

Time = 4.12 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.06

method	result
parts	$-\frac{A a^2 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d} + \frac{(A a^2 + 2B a^2) \tan(dx+c)}{d} + \frac{(2A a^2 + B a^2) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c))}{2} \right)}{d}$
derivativedivides	$\frac{A a^2 \tan(dx+c) + B a^2 \ln(\sec(dx+c) + \tan(dx+c)) + 2A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 2B a^2 \tan(dx+c)}{d}$
default	$\frac{A a^2 \tan(dx+c) + B a^2 \ln(\sec(dx+c) + \tan(dx+c)) + 2A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 2B a^2 \tan(dx+c)}{d}$
parallelrisc	$2 \left(-\frac{3 \left(\frac{3B}{2} + A \right) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)}{2} + \frac{3 \left(\frac{3B}{2} + A \right) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{2} \right) + \frac{A}{d}$
risc	$-\frac{ia^2(6Ae^{5i(dx+c)} + 3Be^{5i(dx+c)} - 6Ae^{4i(dx+c)} - 12Be^{4i(dx+c)} - 24Ae^{2i(dx+c)} - 24Be^{2i(dx+c)} - 6Ae^{i(dx+c)} - 3Be^{i(dx+c)})}{3d(e^{2i(dx+c)} + 1)^3}$
norman	$\frac{-\frac{2a^2(2A-3B) \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} - \frac{a^2(2A+3B) \left(\tan^9 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{3d} - \frac{a^2(2A+3B) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} + \frac{2a^2(2A+5B) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} - \frac{a^2}{d}}{\left(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^3 \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)^3}$

input `int((a+cos(d*x+c))*a^2*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBOSE)`

output `-A*a^2/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(A*a^2+2*B*a^2)/d*tan(d*x+c)+(2*A*a^2+B*a^2)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+1/d*B*ln(sec(d*x+c)+tan(d*x+c))*a^2`

3.17.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.11

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(2A + 3B)a^2 \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3(2A + 3B)a^2 \cos(dx + c)^3 \log(-\sin(dx + c) + 1)}{12d \cos(dx + c)^3}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fracas")`

3.17. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

output `1/12*(3*(2*A + 3*B)*a^2*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*(2*A + 3*B)*a^2*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(2*(5*A + 6*B)*a^2*cos(d*x + c)^2 + 3*(2*A + B)*a^2*cos(d*x + c) + 2*A*a^2)*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.17.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= a^2 \left(\int A \sec^4(c + dx) dx + \int 2A \cos(c + dx) \sec^4(c + dx) dx \right. \\ &\quad \left. + \int A \cos^2(c + dx) \sec^4(c + dx) dx + \int B \cos(c + dx) \sec^4(c + dx) dx \right. \\ &\quad \left. + \int 2B \cos^2(c + dx) \sec^4(c + dx) dx + \int B \cos^3(c + dx) \sec^4(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `a**2*(Integral(A*sec(c + d*x)**4, x) + Integral(2*A*cos(c + d*x)*sec(c + d*x)**4, x) + Integral(A*cos(c + d*x)**2*sec(c + d*x)**4, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**4, x) + Integral(2*B*cos(c + d*x)**2*sec(c + d*x)**4, x) + Integral(B*cos(c + d*x)**3*sec(c + d*x)**4, x))`

3.17.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.54

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= \frac{4 (\tan(dx + c))^3 + 3 \tan(dx + c) A a^2 - 6 A a^2 \left(\frac{2 \sin(dx + c)}{\sin(dx + c)^2 - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right)}{d} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output $1/12*(4*(\tan(dx + c)^3 + 3*\tan(dx + c))*A*a^2 - 6*A*a^2*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) - 3*B*a^2*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) + 6*B*a^2*(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 12*A*a^2*\tan(dx + c) + 24*B*a^2*\tan(dx + c))/d$

3.17.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.58

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(2Aa^2 + 3Ba^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 3(2Aa^2 + 3Ba^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) - \frac{2(6Aa^2 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 3Aa^2 + 3Ba^2)}{6d}}{6d}$$

input `integrate((a+a*cos(dx+c))^2*(A+B*cos(dx+c))*sec(dx+c)^4,x, algorithm="giac")`

output $1/6*(3*(2*A*a^2 + 3*B*a^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(2*A*a^2 + 3*B*a^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*A*a^2*\tan(1/2*d*x + 1/2*c)^5 + 9*B*a^2*\tan(1/2*d*x + 1/2*c)^5 - 16*A*a^2*\tan(1/2*d*x + 1/2*c)^3 - 24*B*a^2*\tan(1/2*d*x + 1/2*c)^3 + 18*A*a^2*\tan(1/2*d*x + 1/2*c) + 15*B*a^2*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^3/d$

3.17.9 Mupad [B] (verification not implemented)

Time = 1.30 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.28

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{2a^2 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) \left(A + \frac{3B}{2}\right) + (2Aa^2 + 3Ba^2) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{16Aa^2}{3} - 8Ba^2\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (6Aa^2 + 5Ba^2) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^4,x)`

3.17. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

output $(2*a^2*atanh(\tan(c/2 + (d*x)/2))*(A + (3*B)/2))/d - (\tan(c/2 + (d*x)/2)*(6*A*a^2 + 5*B*a^2) + \tan(c/2 + (d*x)/2)^5*(2*A*a^2 + 3*B*a^2) - \tan(c/2 + (d*x)/2)^3*((16*A*a^2)/3 + 8*B*a^2))/(d*(3*\tan(c/2 + (d*x)/2)^2 - 3*\tan(c/2 + (d*x)/2)^4 + \tan(c/2 + (d*x)/2)^6 - 1))$

3.18 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.18.1	Optimal result	354
3.18.2	Mathematica [A] (verified)	355
3.18.3	Rubi [A] (verified)	355
3.18.4	Maple [A] (verified)	359
3.18.5	Fricas [A] (verification not implemented)	360
3.18.6	Sympy [F(-1)]	360
3.18.7	Maxima [A] (verification not implemented)	361
3.18.8	Giac [A] (verification not implemented)	361
3.18.9	Mupad [B] (verification not implemented)	362

3.18.1 Optimal result

Integrand size = 31, antiderivative size = 144

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{a^2(7A + 8B)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{a^2(4A + 5B) \tan(c + dx)}{3d}$$

$$+ \frac{a^2(7A + 8B) \sec(c + dx) \tan(c + dx)}{8d} + \frac{a^2(5A + 4B) \sec^2(c + dx) \tan(c + dx)}{12d}$$

$$+ \frac{A(a^2 + a^2 \cos(c + dx)) \sec^3(c + dx) \tan(c + dx)}{4d}$$

output `1/8*a^2*(7*A+8*B)*arctanh(sin(d*x+c))/d+1/3*a^2*(4*A+5*B)*tan(d*x+c)/d+1/8*a^2*(7*A+8*B)*sec(d*x+c)*tan(d*x+c)/d+1/12*a^2*(5*A+4*B)*sec(d*x+c)^2*tan(d*x+c)/d+1/4*A*(a^2+a^2*cos(d*x+c))*sec(d*x+c)^3*tan(d*x+c)/d`

3.18.2 Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.56

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{a^2(3(7A + 8B)\operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx)(48(A + B) + 3(7A + 8B)\sec(c + dx) + 6A\sec^3(c + dx))}{24d}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a^2*(3*(7*A + 8*B)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(48*(A + B) + 3*(7*A + 8*B)*Sec[c + d*x] + 6*A*Sec[c + d*x]^3 + 8*(2*A + B)*Tan[c + d*x]^2))/ (24*d)`

3.18.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3454, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow 3454$$

$$\frac{1}{4} \int (\cos(c + dx)a + a)(a(5A + 4B) + 2a(A + 2B)\cos(c + dx)) \sec^4(c + dx) dx + \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \int \frac{(\sin(c + dx + \frac{\pi}{2}) a + a) (a(5A + 4B) + 2a(A + 2B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx + \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

3.18. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

$$\downarrow 3447$$

$$\frac{1}{4} \int (2(A+2B) \cos^2(c+dx)a^2 + (5A+4B)a^2 + (2(A+2B)a^2 + (5A+4B)a^2) \cos(c+dx)) \sec^4(c+dx) dx + \frac{A \tan(c+dx) \sec^3(c+dx) (a^2 \cos(c+dx) + a^2)}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \int \frac{2(A+2B) \sin(c+dx + \frac{\pi}{2})^2 a^2 + (5A+4B)a^2 + (2(A+2B)a^2 + (5A+4B)a^2) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^4} dx + \frac{A \tan(c+dx) \sec^3(c+dx) (a^2 \cos(c+dx) + a^2)}{4d}$$

$$\downarrow 3500$$

$$\frac{1}{4} \left(\frac{1}{3} \int (3(7A+8B)a^2 + 4(4A+5B) \cos(c+dx)a^2) \sec^3(c+dx) dx + \frac{a^2(5A+4B) \tan(c+dx) \sec^2(c+dx)}{3d} \right) + \frac{A \tan(c+dx) \sec^3(c+dx) (a^2 \cos(c+dx) + a^2)}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \left(\frac{1}{3} \int \frac{3(7A+8B)a^2 + 4(4A+5B) \sin(c+dx + \frac{\pi}{2}) a^2}{\sin(c+dx + \frac{\pi}{2})^3} dx + \frac{a^2(5A+4B) \tan(c+dx) \sec^2(c+dx)}{3d} \right) + \frac{A \tan(c+dx) \sec^3(c+dx) (a^2 \cos(c+dx) + a^2)}{4d}$$

$$\downarrow 3227$$

$$\frac{1}{4} \left(\frac{1}{3} \left(3a^2(7A+8B) \int \sec^3(c+dx) dx + 4a^2(4A+5B) \int \sec^2(c+dx) dx \right) + \frac{a^2(5A+4B) \tan(c+dx) \sec^2(c+dx)}{3d} \right) + \frac{A \tan(c+dx) \sec^3(c+dx) (a^2 \cos(c+dx) + a^2)}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \left(\frac{1}{3} \left(4a^2(4A+5B) \int \csc(c+dx + \frac{\pi}{2})^2 dx + 3a^2(7A+8B) \int \csc(c+dx + \frac{\pi}{2})^3 dx \right) + \frac{a^2(5A+4B) \tan(c+dx) \sec^2(c+dx)}{3d} \right) + \frac{A \tan(c+dx) \sec^3(c+dx) (a^2 \cos(c+dx) + a^2)}{4d}$$

$$\downarrow 4254$$

$$\frac{1}{4} \left(\frac{1}{3} \left(3a^2(7A + 8B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx - \frac{4a^2(4A + 5B) \int 1d(-\tan(c + dx))}{d} \right) + \frac{a^2(5A + 4B) \tan(c + dx)}{3d} \right) \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

↓ 24

$$\frac{1}{4} \left(\frac{1}{3} \left(3a^2(7A + 8B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx + \frac{4a^2(4A + 5B) \tan(c + dx)}{d} \right) + \frac{a^2(5A + 4B) \tan(c + dx) \sec^2(c + dx)}{3d} \right) \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

↓ 4255

$$\frac{1}{4} \left(\frac{1}{3} \left(3a^2(7A + 8B) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4a^2(4A + 5B) \tan(c + dx)}{d} \right) + \frac{a^2(5A + 4B) \tan(c + dx) \sec^2(c + dx)}{3d} \right) \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(3a^2(7A + 8B) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4a^2(4A + 5B) \tan(c + dx)}{d} \right) + \frac{a^2(5A + 4B) \tan(c + dx) \sec^2(c + dx)}{3d} \right) \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

↓ 4257

$$\frac{1}{4} \left(\frac{1}{3} \left(3a^2(7A + 8B) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4a^2(4A + 5B) \tan(c + dx)}{d} \right) + \frac{a^2(5A + 4B) \tan(c + dx) \sec^2(c + dx)}{3d} \right) \frac{A \tan(c + dx) \sec^3(c + dx) (a^2 \cos(c + dx) + a^2)}{4d}$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(A*(a^2 + a^2*Cos[c + d*x])*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((a^2*(5*A + 4*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((4*a^2*(4*A + 5*B)*Tan[c + d*x])/d + 3*a^2*(7*A + 8*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/3)/4`

3.18.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

output $A*a^2/d*(-(-1/4*\sec(d*x+c)^3-3/8*\sec(d*x+c))*\tan(d*x+c)+3/8*\ln(\sec(d*x+c)+\tan(d*x+c)))+(A*a^2+2*B*a^2)/d*(1/2*\sec(d*x+c)*\tan(d*x+c)+1/2*\ln(\sec(d*x+c)+\tan(d*x+c)))-(2*A*a^2+B*a^2)/d*(-2/3-1/3*\sec(d*x+c)^2)*\tan(d*x+c)+B*a^2/d*\tan(d*x+c)$

3.18.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.01

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(7A + 8B)a^2 \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(7A + 8B)a^2 \cos(dx + c)^4 \log(-\sin(dx + c) + 1)}{48}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fracas")`

output $1/48*(3*(7*A + 8*B)*a^2*\cos(d*x + c)^4*\log(\sin(d*x + c) + 1) - 3*(7*A + 8*B)*a^2*\cos(d*x + c)^4*\log(-\sin(d*x + c) + 1) + 2*(8*(4*A + 5*B)*a^2*\cos(d*x + c)^3 + 3*(7*A + 8*B)*a^2*\cos(d*x + c)^2 + 8*(2*A + B)*a^2*\cos(d*x + c) + 6*A*a^2)*\sin(d*x + c))/(d*\cos(d*x + c)^4)$

3.18.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output Timed out

3.18. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.18.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.60

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{32 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^2 + 16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ba^2 - 3 Aa^2 \left(\frac{2(3 \sin(dx+c)^3 - 5 \sin(dx+c))}{\sin(dx+c)^4 - 2 \sin(dx+c)^2 + 1} - 3 \log(\sin(dx+c) + 1) + 3 \log(\sin(dx+c) - 1) - 12 Aa^2 \frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1) - 24 Ba^2 \frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1) + 48 Ba^2 \tan(dx+c) \right)}{d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output `1/48*(32*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^2 + 16*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^2 - 3*A*a^2*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 12*A*a^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 24*B*a^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 48*B*a^2*tan(d*x + c))/d`

3.18.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.47

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(7Aa^2 + 8Ba^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 3(7Aa^2 + 8Ba^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) - \frac{2(21Aa^2 \tan(1/2 dx + 1/2 c) + 7Aa^2 + 8Ba^2)}{\tan(1/2 dx + 1/2 c)^2 - 1}}{d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")`

output `1/24*(3*(7*A*a^2 + 8*B*a^2)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*(7*A*a^2 + 8*B*a^2)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(21*A*a^2*tan(1/2*d*x + 1/2*c)^7 + 24*B*a^2*tan(1/2*d*x + 1/2*c)^7 - 77*A*a^2*tan(1/2*d*x + 1/2*c)^5 - 88*B*a^2*tan(1/2*d*x + 1/2*c)^5 + 83*A*a^2*tan(1/2*d*x + 1/2*c)^3 + 136*B*a^2*tan(1/2*d*x + 1/2*c)^3 - 75*A*a^2*tan(1/2*d*x + 1/2*c) - 72*B*a^2*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^4/d`

3.18. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.18.9 Mupad [B] (verification not implemented)

Time = 1.94 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.27

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{\left(-\frac{7Aa^2}{4} - 2Ba^2\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{77Aa^2}{12} + \frac{22Ba^2}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{83Aa^2}{12} - \frac{34Ba^2}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)} + \frac{2a^2 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) \left(\frac{7A}{8} + B\right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^5,x)`output `(tan(c/2 + (d*x)/2)*((25*A*a^2)/4 + 6*B*a^2) - tan(c/2 + (d*x)/2)^7*((7*A*a^2)/4 + 2*B*a^2) + tan(c/2 + (d*x)/2)^5*((77*A*a^2)/12 + (22*B*a^2)/3) - tan(c/2 + (d*x)/2)^3*((83*A*a^2)/12 + (34*B*a^2)/3))/(d*(6*tan(c/2 + (d*x)/2)^4 - 4*tan(c/2 + (d*x)/2)^2 - 4*tan(c/2 + (d*x)/2)^6 + tan(c/2 + (d*x)/2)^8 + 1)) + (2*a^2*atanh(tan(c/2 + (d*x)/2))*((7*A)/8 + B))/d`

3.19 $\int \cos^2(c+dx)(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$

3.19.1	Optimal result	363
3.19.2	Mathematica [A] (verified)	364
3.19.3	Rubi [A] (verified)	364
3.19.4	Maple [A] (verified)	369
3.19.5	Fricas [A] (verification not implemented)	369
3.19.6	Sympy [B] (verification not implemented)	370
3.19.7	Maxima [A] (verification not implemented)	371
3.19.8	Giac [A] (verification not implemented)	371
3.19.9	Mupad [B] (verification not implemented)	372

3.19.1 Optimal result

Integrand size = 31, antiderivative size = 201

$$\int \cos^2(c+dx)(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$$

$$= \frac{1}{16}a^3(26A+23B)x + \frac{a^3(19A+17B) \sin(c+dx)}{5d}$$

$$+ \frac{a^3(26A+23B) \cos(c+dx) \sin(c+dx)}{16d} + \frac{a^3(22A+21B) \cos^3(c+dx) \sin(c+dx)}{40d}$$

$$+ \frac{aB \cos^3(c+dx)(a+a \cos(c+dx))^2 \sin(c+dx)}{6d}$$

$$+ \frac{(3A+4B) \cos^3(c+dx) (a^3+a^3 \cos(c+dx)) \sin(c+dx)}{15d}$$

$$- \frac{a^3(19A+17B) \sin^3(c+dx)}{15d}$$

```
output 1/16*a^3*(26*A+23*B)*x+1/5*a^3*(19*A+17*B)*sin(d*x+c)/d+1/16*a^3*(26*A+23*
B)*cos(d*x+c)*sin(d*x+c)/d+1/40*a^3*(22*A+21*B)*cos(d*x+c)^3*sin(d*x+c)/d+
1/6*a*B*cos(d*x+c)^3*(a+a*cos(d*x+c))^2*sin(d*x+c)/d+1/15*(3*A+4*B)*cos(d*
x+c)^3*(a^3+a^3*cos(d*x+c))*sin(d*x+c)/d-1/15*a^3*(19*A+17*B)*sin(d*x+c)^3
/d
```

3.19.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.67

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{a^3(1380Bc + 1560Adx + 1380Bdx + 120(23A + 21B) \sin(c + dx) + 15(64A + 63B) \sin(2(c + dx)) + 340A \sin(3(c + dx)) + 380B \sin(3(c + dx)) + 90A \sin(4(c + dx)) + 135B \sin(4(c + dx)) + 12A \sin(5(c + dx)) + 36B \sin(5(c + dx)) + 5B \sin(6(c + dx)))}{960d}$$

input `Integrate[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output $(a^3(1380Bc + 1560A*d*x + 1380B*d*x + 120*(23A + 21B)*\text{Sin}[c + d*x] + 15*(64A + 63B)*\text{Sin}[2*(c + d*x)] + 340A*\text{Sin}[3*(c + d*x)] + 380B*\text{Sin}[3*(c + d*x)] + 90A*\text{Sin}[4*(c + d*x)] + 135B*\text{Sin}[4*(c + d*x)] + 12A*\text{Sin}[5*(c + d*x)] + 36B*\text{Sin}[5*(c + d*x)] + 5B*\text{Sin}[6*(c + d*x)])/(960*d)$

3.19.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.98, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3455, 3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx)(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3455}$$

$$\frac{1}{6} \int \cos^2(c + dx)(\cos(c + dx)a + a)^2(3a(2A + B) + 2a(3A + 4B) \cos(c + dx)) dx + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^2}{6d}$$

$$\downarrow \text{3042}$$

3.19. $\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

$$\frac{1}{6} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right)^2 \left(3a(2A + B) + 2a(3A + 4B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^2}{6d}$$

↓ 3455

$$\frac{1}{6} \left(\frac{1}{5} \int 3 \cos^2(c + dx) (\cos(c + dx)a + a) ((16A + 13B)a^2 + (22A + 21B) \cos(c + dx)a^2) dx + \frac{2(3A + 4B) \sin(c + dx) (a \cos(c + dx) + a)^2}{6d} \right)$$

↓ 27

$$\frac{1}{6} \left(\frac{3}{5} \int \cos^2(c + dx) (\cos(c + dx)a + a) ((16A + 13B)a^2 + (22A + 21B) \cos(c + dx)a^2) dx + \frac{2(3A + 4B) \sin(c + dx) (a \cos(c + dx) + a)^2}{6d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{3}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right) \left((16A + 13B)a^2 + (22A + 21B) \sin \left(c + dx + \frac{\pi}{2} \right) a^2 \right) dx + \frac{aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^2}{6d} \right)$$

↓ 3447

$$\frac{1}{6} \left(\frac{3}{5} \int \cos^2(c + dx) ((22A + 21B) \cos^2(c + dx)a^3 + (16A + 13B)a^3 + ((16A + 13B)a^3 + (22A + 21B)a^3) \cos(c + dx)) dx + \frac{aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^2}{6d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{3}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left((22A + 21B) \sin \left(c + dx + \frac{\pi}{2} \right)^2 a^3 + (16A + 13B)a^3 + ((16A + 13B)a^3 + (22A + 21B)a^3) \sin \left(c + dx + \frac{\pi}{2} \right) a^2 \right) dx + \frac{aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^2}{6d} \right)$$

↓ 3502

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \int \cos^2(c+dx) (5(26A+23B)a^3 + 8(19A+17B)\cos(c+dx)a^3) dx + \frac{a^3(22A+21B)\sin(c+dx)\cos^3(c+dx)}{4d} \right) \right. \\ \left. \frac{aB \sin(c+dx)\cos^3(c+dx)(a\cos(c+dx)+a)^2}{6d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 (5(26A+23B)a^3 + 8(19A+17B)\sin\left(c+dx+\frac{\pi}{2}\right)a^3) dx + \frac{a^3(22A+21B)\sin(c+dx)\cos^3(c+dx)}{4d} \right) \right. \\ \left. \frac{aB \sin(c+dx)\cos^3(c+dx)(a\cos(c+dx)+a)^2}{6d} \right) \\ \downarrow \text{3227}$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(8a^3(19A+17B) \int \cos^3(c+dx) dx + 5a^3(26A+23B) \int \cos^2(c+dx) dx \right) + \frac{a^3(22A+21B)\sin(c+dx)\cos^3(c+dx)}{4d} \right) \right. \\ \left. \frac{aB \sin(c+dx)\cos^3(c+dx)(a\cos(c+dx)+a)^2}{6d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(5a^3(26A+23B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx + 8a^3(19A+17B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 dx \right) + \frac{a^3(22A+21B)\sin(c+dx)\cos^3(c+dx)}{4d} \right) \right. \\ \left. \frac{aB \sin(c+dx)\cos^3(c+dx)(a\cos(c+dx)+a)^2}{6d} \right) \\ \downarrow \text{3113}$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(5a^3(26A+23B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx - \frac{8a^3(19A+17B) \int (1-\sin^2(c+dx)) d(-\sin(c+dx))}{d} \right) + \frac{a^3(22A+21B)\sin(c+dx)\cos^3(c+dx)}{4d} \right) \right. \\ \left. \frac{aB \sin(c+dx)\cos^3(c+dx)(a\cos(c+dx)+a)^2}{6d} \right) \\ \downarrow \text{2009}$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(5a^3(26A+23B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx - \frac{8a^3(19A+17B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d} \right) + \frac{a^3(22A+21B)\sin(c+dx)\cos^3(c+dx)}{4d} \right) \right. \\ \left. \frac{aB \sin(c+dx)\cos^3(c+dx)(a\cos(c+dx)+a)^2}{6d} \right) \\ \downarrow \text{3115}$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(5a^3(26A + 23B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{8a^3(19A + 17B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^2}{6d} \right) \right) \right. \\ \left. \downarrow 24 \right.$$

$$\frac{1}{6} \left(\frac{3}{5} \left(\frac{a^3(22A + 21B) \sin(c + dx) \cos^3(c + dx)}{4d} + \frac{1}{4} \left(5a^3(26A + 23B) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) - \frac{8a^3(19A + 17B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) \right. \right. \\ \left. \left. \frac{aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^2}{6d} \right) \right)$$

input `Int[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^3*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(6*d) + ((2*(3*A + 4*B)*Cos[c + d*x]^3*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(5*d) + (3*((a^3*(22*A + 21*B)*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (5*a^3*(26*A + 23*B))*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (8*a^3*(19*A + 17*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/4))/5)/6`

3.19.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

$$3.19. \quad \int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*SIN[e + f*x])^m*(A*c + (B*c + A*d)*SIN[e + f*x] + B*d*SIN[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b)*B*COS[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*SIN[e + f*x])^(m - 1)*(c + d*SIN[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*SIN[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*COS[e + f*x]*((a + b*SIN[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*SIN[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*SIN[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.19.4 Maple [A] (verified)

Time = 4.29 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.56

method	result
parallelrisc	$3 \left(\left(\frac{32A}{3} + \frac{21B}{2} \right) \sin(2dx+2c) + \frac{2(17A+19B) \sin(3dx+3c)}{9} + \left(\frac{3B}{2} + A \right) \sin(4dx+4c) + \frac{2 \left(\frac{4}{3} + B \right) \sin(5dx+5c)}{5} + \frac{B \sin(6dx+6c)}{18} + \dots \right)$
risc	$\frac{13a^3 Ax}{8} + \frac{23a^3 Bx}{16} + \frac{23a^3 A \sin(dx+c)}{8d} + \frac{21a^3 B \sin(dx+c)}{8d} + \frac{B a^3 \sin(6dx+6c)}{192d} + \frac{\sin(5dx+5c) A a^3}{80d} + \frac{3 \sin(5dx+5c) B a^3}{80d}$
parts	$\frac{(A a^3 + 3B a^3) \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5d} + \frac{(3A a^3 + B a^3) (2 + \cos^2(dx+c)) \sin(dx+c)}{3d} + \dots$
derivativedivides	$\frac{A a^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + B a^3 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right)$
default	$\frac{A a^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + B a^3 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right)$
norman	$\frac{a^3(26A+23B)x}{16} + \frac{33a^3(26A+23B) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{20d} + \frac{17a^3(26A+23B) \left(\tan^9 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{24d} + \frac{a^3(26A+23B) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{8d} + \frac{3a^3(26A+23B)}{16}$

input `int(cos(d*x+c)^2*(a+cos(d*x+c))*a^3*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `3/32*((32/3*A+21/2*B)*sin(2*d*x+2*c)+2/9*(17*A+19*B)*sin(3*d*x+3*c)+(3/2*B+A)*sin(4*d*x+4*c)+2/5*(1/3*A+B)*sin(5*d*x+5*c)+1/18*B*sin(6*d*x+6*c)+4*(2/3/3*A+7*B)*sin(d*x+c)+52/3*(23/26*B+A)*x*d)*a^3/d`

3.19.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.65

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{15(26A + 23B)a^3 dx + (40Ba^3 \cos(dx + c))^5 + 48(A + 3B)a^3 \cos(dx + c)^4 + 10(18A + 23B)a^3 \cos(dx + c)^3 + \dots}{16}$$

3.19. $\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$


```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="f
ricas")
```

```
output 1/240*(15*(26*A + 23*B)*a^3*d*x + (40*B*a^3*cos(d*x + c)^5 + 48*(A + 3*B)*
a^3*cos(d*x + c)^4 + 10*(18*A + 23*B)*a^3*cos(d*x + c)^3 + 16*(19*A + 17*B
)*a^3*cos(d*x + c)^2 + 15*(26*A + 23*B)*a^3*cos(d*x + c) + 32*(19*A + 17*B
)*a^3)*sin(d*x + c))/d
```

3.19.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 695 vs. $2(184) = 368$.

Time = 0.41 (sec) , antiderivative size = 695, normalized size of antiderivative = 3.46

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{9Aa^3x \sin^4(c+dx)}{8} + \frac{9Aa^3x \sin^2(c+dx) \cos^2(c+dx)}{4} + \frac{Aa^3x \sin^2(c+dx)}{2} + \frac{9Aa^3x \cos^4(c+dx)}{8} + \frac{Aa^3x \cos^2(c+dx)}{2} + \frac{8Aa^3 \sin^5(c+dx)}{15d} \\ x(A + B \cos(c))(a \cos(c) + a)^3 \cos^2(c) \end{cases}$$

```
input integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)
```

```
output Piecewise((9*A*a**3*x*sin(c + d*x)**4/8 + 9*A*a**3*x*sin(c + d*x)**2*cos(c
+ d*x)**2/4 + A*a**3*x*sin(c + d*x)**2/2 + 9*A*a**3*x*cos(c + d*x)**4/8 +
A*a**3*x*cos(c + d*x)**2/2 + 8*A*a**3*sin(c + d*x)**5/(15*d) + 4*A*a**3*s
in(c + d*x)**3*cos(c + d*x)**2/(3*d) + 9*A*a**3*sin(c + d*x)**3*cos(c + d*
x)/(8*d) + 2*A*a**3*sin(c + d*x)**3/d + A*a**3*sin(c + d*x)*cos(c + d*x)**
4/d + 15*A*a**3*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 3*A*a**3*sin(c + d*x)
*cos(c + d*x)**2/d + A*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + 5*B*a**3*x*s
in(c + d*x)**6/16 + 15*B*a**3*x*sin(c + d*x)**4*cos(c + d*x)**2/16 + 9*B*a
**3*x*sin(c + d*x)**4/8 + 15*B*a**3*x*sin(c + d*x)**2*cos(c + d*x)**4/16 +
9*B*a**3*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 5*B*a**3*x*cos(c + d*x)**6
/16 + 9*B*a**3*x*cos(c + d*x)**4/8 + 5*B*a**3*sin(c + d*x)**5*cos(c + d*x)
/(16*d) + 8*B*a**3*sin(c + d*x)**5/(5*d) + 5*B*a**3*sin(c + d*x)**3*cos(c
+ d*x)**3/(6*d) + 4*B*a**3*sin(c + d*x)**3*cos(c + d*x)**2/d + 9*B*a**3*si
n(c + d*x)**3*cos(c + d*x)/(8*d) + 2*B*a**3*sin(c + d*x)**3/(3*d) + 11*B*a
**3*sin(c + d*x)*cos(c + d*x)**5/(16*d) + 3*B*a**3*sin(c + d*x)*cos(c + d*
x)**4/d + 15*B*a**3*sin(c + d*x)*cos(c + d*x)**3/(8*d) + B*a**3*sin(c + d*
x)*cos(c + d*x)**2/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**3*cos(c
)**2, True))
```

3.19. $\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.19.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.30

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{64(3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c))Aa^3 - 960(\sin(dx + c)^3 - 3 \sin(dx + c))Aa^3 +$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output 1/960*(64*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*A*a^3 -
960*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^3 + 90*(12*d*x + 12*c + sin(4*d
*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^3 + 240*(2*d*x + 2*c + sin(2*d*x + 2*c
))*A*a^3 + 192*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*
a^3 - 5*(4*sin(2*d*x + 2*c)^3 - 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*si
n(2*d*x + 2*c))*B*a^3 - 320*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^3 + 90*(
12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^3)/d
```

3.19.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.83

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^3 \sin(6dx + 6c)}{192d} + \frac{1}{16}(26Aa^3 + 23Ba^3)x + \frac{(Aa^3 + 3Ba^3) \sin(5dx + 5c)}{80d}$$

$$+ \frac{3(2Aa^3 + 3Ba^3) \sin(4dx + 4c)}{64d} + \frac{(17Aa^3 + 19Ba^3) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(64Aa^3 + 63Ba^3) \sin(2dx + 2c)}{64d} + \frac{(23Aa^3 + 21Ba^3) \sin(dx + c)}{8d}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/192*B*a^3*sin(6*d*x + 6*c)/d + 1/16*(26*A*a^3 + 23*B*a^3)*x + 1/80*(A*a^
3 + 3*B*a^3)*sin(5*d*x + 5*c)/d + 3/64*(2*A*a^3 + 3*B*a^3)*sin(4*d*x + 4*c
)/d + 1/48*(17*A*a^3 + 19*B*a^3)*sin(3*d*x + 3*c)/d + 1/64*(64*A*a^3 + 63*
B*a^3)*sin(2*d*x + 2*c)/d + 1/8*(23*A*a^3 + 21*B*a^3)*sin(d*x + c)/d
```

3.19. $\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.19.9 Mupad [B] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.57

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{13Aa^3}{4} + \frac{23Ba^3}{8}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{11} + \left(\frac{221Aa^3}{12} + \frac{391Ba^3}{24}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{429Aa^3}{10} + \frac{759Ba^3}{20}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{12} + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 15 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + a^3(26A + 23B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)} + \frac{a^3 \operatorname{atan}\left(\frac{a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (26A + 23B)}{8 \left(\frac{13Aa^3}{4} + \frac{23Ba^3}{8}\right)}\right) (26A + 23B)}{8d}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3,x)`

```
output (tan(c/2 + (d*x)/2)*((51*A*a^3)/4 + (105*B*a^3)/8) + tan(c/2 + (d*x)/2)^11
*((13*A*a^3)/4 + (23*B*a^3)/8) + tan(c/2 + (d*x)/2)^3*((419*A*a^3)/12 + (2
11*B*a^3)/8) + tan(c/2 + (d*x)/2)^9*((221*A*a^3)/12 + (391*B*a^3)/24) + ta
n(c/2 + (d*x)/2)^7*((429*A*a^3)/10 + (759*B*a^3)/20) + tan(c/2 + (d*x)/2)^
5*((499*A*a^3)/10 + (969*B*a^3)/20))/(d*(6*tan(c/2 + (d*x)/2)^2 + 15*tan(c
/2 + (d*x)/2)^4 + 20*tan(c/2 + (d*x)/2)^6 + 15*tan(c/2 + (d*x)/2)^8 + 6*ta
n(c/2 + (d*x)/2)^10 + tan(c/2 + (d*x)/2)^12 + 1)) - (a^3*(26*A + 23*B)*(at
an(tan(c/2 + (d*x)/2)) - (d*x)/2))/(8*d) + (a^3*atan((a^3*tan(c/2 + (d*x)/
2)*(26*A + 23*B))/(8*((13*A*a^3)/4 + (23*B*a^3)/8)))*(26*A + 23*B))/(8*d)
```

3.20 $\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.20.1	Optimal result	373
3.20.2	Mathematica [A] (verified)	374
3.20.3	Rubi [A] (verified)	374
3.20.4	Maple [A] (verified)	377
3.20.5	Fricas [A] (verification not implemented)	377
3.20.6	Sympy [B] (verification not implemented)	378
3.20.7	Maxima [A] (verification not implemented)	379
3.20.8	Giac [A] (verification not implemented)	379
3.20.9	Mupad [B] (verification not implemented)	380

3.20.1 Optimal result

Integrand size = 29, antiderivative size = 154

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{1}{8}a^3(15A + 13B)x + \frac{a^3(15A + 13B) \sin(c + dx)}{5d}$$

$$+ \frac{3a^3(15A + 13B) \cos(c + dx) \sin(c + dx)}{40d} + \frac{(5A - B)(a + a \cos(c + dx))^3 \sin(c + dx)}{20d}$$

$$+ \frac{B(a + a \cos(c + dx))^4 \sin(c + dx)}{5ad} - \frac{a^3(15A + 13B) \sin^3(c + dx)}{60d}$$

```
output 1/8*a^3*(15*A+13*B)*x+1/5*a^3*(15*A+13*B)*sin(d*x+c)/d+3/40*a^3*(15*A+13*B)
)*cos(d*x+c)*sin(d*x+c)/d+1/20*(5*A-B)*(a+a*cos(d*x+c))^3*sin(d*x+c)/d+1/5
*B*(a+a*cos(d*x+c))^4*sin(d*x+c)/a/d-1/60*a^3*(15*A+13*B)*sin(d*x+c)^3/d
```

3.20.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.70

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{a^3(780Bc + 900Adx + 780Bdx + 60(26A + 23B) \sin(c + dx) + 480(A + B) \sin(2(c + dx)) + 120A \sin(3(c + dx)))}{480d}$$

input `Integrate[Cos[c + d*x]*(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(a^3*(780*B*c + 900*A*d*x + 780*B*d*x + 60*(26*A + 23*B)*Sin[c + d*x] + 480*(A + B)*Sin[2*(c + d*x)] + 120*A*SIN[3*(c + d*x)] + 170*B*SIN[3*(c + d*x)]) + 15*A*SIN[4*(c + d*x)] + 45*B*SIN[4*(c + d*x)] + 6*B*SIN[5*(c + d*x)])/(480*d)`

3.20.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.95, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3447, 3042, 3502, 3042, 3230, 3042, 3124, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(c + dx)(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int (a \cos(c + dx) + a)^3 (A \cos(c + dx) + B \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{3502}$$

$$\begin{aligned}
& \frac{\int (\cos(c+dx)a+a)^3(4aB+a(5A-B)\cos(c+dx))dx}{5a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^4}{5ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\int (\sin(c+dx+\frac{\pi}{2})a+a)^3(4aB+a(5A-B)\sin(c+dx+\frac{\pi}{2}))dx}{5a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^4}{5ad} \\
& \quad \downarrow \text{3230} \\
& \frac{\frac{1}{4}a(15A+13B)\int (\cos(c+dx)a+a)^3dx + \frac{a(5A-B)\sin(c+dx)(a\cos(c+dx)+a)^3}{4d}}{5a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^4}{5ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{4}a(15A+13B)\int (\sin(c+dx+\frac{\pi}{2})a+a)^3dx + \frac{a(5A-B)\sin(c+dx)(a\cos(c+dx)+a)^3}{4d}}{5a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^4}{5ad} \\
& \quad \downarrow \text{3124} \\
& \frac{\frac{1}{4}a(15A+13B)\int (\cos^3(c+dx)a^3+3\cos^2(c+dx)a^3+3\cos(c+dx)a^3+a^3)dx + \frac{a(5A-B)\sin(c+dx)(a\cos(c+dx)+a)^3}{4d}}{5a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^4}{5ad} \\
& \quad \downarrow \text{2009} \\
& \frac{\frac{1}{4}a(15A+13B)\left(-\frac{a^3\sin^3(c+dx)}{3d} + \frac{4a^3\sin(c+dx)}{d} + \frac{3a^3\sin(c+dx)\cos(c+dx)}{2d} + \frac{5a^3x}{2}\right) + \frac{a(5A-B)\sin(c+dx)(a\cos(c+dx)+a)^3}{4d}}{5a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^4}{5ad}
\end{aligned}$$

input `Int[Cos[c + d*x]*(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(B*(a + a*Cos[c + d*x])^4*Sin[c + d*x])/(5*a*d) + ((a*(5*A - B)*(a + a*Cos[c + d*x])^3*Sin[c + d*x])/(4*d) + (a*(15*A + 13*B)*((5*a^3*x)/2 + (4*a^3*Sin[c + d*x])/d + (3*a^3*Cos[c + d*x]*Sin[c + d*x])/(2*d) - (a^3*Sin[c + d*x]^3)/(3*d)))/4)/(5*a)`

3.20.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3124 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(a + b*sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.20.4 Maple [A] (verified)

Time = 3.36 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.60

method	result
parallelrisc	$(32(A+B) \sin(2dx+2c)+2(4A+\frac{17B}{3}) \sin(3dx+3c)+(A+3B) \sin(4dx+4c)+\frac{2B \sin(5dx+5c)}{5}+4(26A+23B) \sin(dx+c)+\frac{60A^3}{32d})$
risc	$\frac{15a^3Ax}{8} + \frac{13a^3Bx}{8} + \frac{13a^3A \sin(dx+c)}{4d} + \frac{23a^3B \sin(dx+c)}{8d} + \frac{\sin(5dx+5c)B a^3}{80d} + \frac{\sin(4dx+4c)A a^3}{32d} + \frac{3 \sin(4dx+4c)A a^3}{32d}$
parts	$(A a^3+3B a^3) \left(\frac{\left(\cos^3(dx+c)+\frac{3 \cos(\frac{dx+c}{2})}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{(3A a^3+B a^3) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d}$
derivatividevides	$A a^3 \left(\frac{\left(\cos^3(dx+c)+\frac{3 \cos(\frac{dx+c}{2})}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{B a^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + A a^3 (2+\cos^2(dx+c))$
default	$A a^3 \left(\frac{\left(\cos^3(dx+c)+\frac{3 \cos(\frac{dx+c}{2})}{2} \right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + \frac{B a^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + A a^3 (2+\cos^2(dx+c))$
norman	$\frac{a^3(15A+13B)x}{8} + \frac{32a^3(15A+13B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{15d} + \frac{7a^3(15A+13B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{6d} + \frac{a^3(15A+13B) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{4d} + \frac{5a^3(15A+13B)}{4d}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/32*(32*(A+B)*sin(2*d*x+2*c)+2*(4*A+17/3*B)*sin(3*d*x+3*c)+(A+3*B)*sin(4*d*x+4*c)+2/5*B*sin(5*d*x+5*c)+4*(26*A+23*B)*sin(d*x+c)+60*(A+13/15*B)*x*d)*a^3/d
```

3.20.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.71

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{15(15A + 13B)a^3 dx + (24Ba^3 \cos(dx + c))^4 + 30(A + 3B)a^3 \cos(dx + c)^3 + 8(15A + 19B)a^3 \cos(dx + c)}{120d}$$

```
input integrate(cos(d*x+c)*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

3.20. $\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

output $1/120*(15*(15*A + 13*B)*a^3*d*x + (24*B*a^3*\cos(d*x + c)^4 + 30*(A + 3*B)*a^3*\cos(d*x + c)^3 + 8*(15*A + 19*B)*a^3*\cos(d*x + c)^2 + 15*(15*A + 13*B)*a^3*\cos(d*x + c) + 8*(45*A + 38*B)*a^3)*\sin(d*x + c))/d$

3.20.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 530 vs. $2(136) = 272$.

Time = 0.30 (sec) , antiderivative size = 530, normalized size of antiderivative = 3.44

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{3Aa^3x \sin^4(c+dx)}{8} + \frac{3Aa^3x \sin^2(c+dx) \cos^2(c+dx)}{4} + \frac{3Aa^3x \sin^2(c+dx)}{2} + \frac{3Aa^3x \cos^4(c+dx)}{8} + \frac{3Aa^3x \cos^2(c+dx)}{2} + \frac{3Aa^3 \sin^3(c)}{2} \\ x(A + B \cos(c))(a \cos(c) + a)^3 \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)`

output `Piecewise((3*A*a**3*x*sin(c + d*x)**4/8 + 3*A*a**3*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*A*a**3*x*sin(c + d*x)**2/2 + 3*A*a**3*x*cos(c + d*x)**4/8 + 3*A*a**3*x*cos(c + d*x)**2/2 + 3*A*a**3*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 2*A*a**3*sin(c + d*x)**3/d + 5*A*a**3*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 3*A*a**3*sin(c + d*x)*cos(c + d*x)**2/d + 3*A*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + A*a**3*sin(c + d*x)/d + 9*B*a**3*x*sin(c + d*x)**4/8 + 9*B*a**3*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + B*a**3*x*sin(c + d*x)**2/2 + 9*B*a**3*x*cos(c + d*x)**4/8 + B*a**3*x*cos(c + d*x)**2/2 + 8*B*a**3*sin(c + d*x)**5/(15*d) + 4*B*a**3*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 9*B*a**3*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 2*B*a**3*sin(c + d*x)**3/d + B*a**3*sin(c + d*x)*cos(c + d*x)**4/d + 15*B*a**3*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 3*B*a**3*sin(c + d*x)*cos(c + d*x)**2/d + B*a**3*sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**3*cos(c), True))`

3.20.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.38

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx =$$

$$\frac{480 (\sin(dx + c)^3 - 3 \sin(dx + c))Aa^3 - 15 (12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(2 dx + 2 c))Aa^3 -$$

```
input integrate(cos(d*x+c)*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output -1/480*(480*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^3 - 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^3 - 360*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^3 - 32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a^3 + 480*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^3 - 45*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^3 - 120*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^3 - 480*A*a^3*sin(d*x + c))/d
```

3.20.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.88

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^3 \sin(5 dx + 5 c)}{80 d} + \frac{1}{8} (15 Aa^3 + 13 Ba^3)x$$

$$+ \frac{(Aa^3 + 3 Ba^3) \sin(4 dx + 4 c)}{32 d} + \frac{(12 Aa^3 + 17 Ba^3) \sin(3 dx + 3 c)}{48 d}$$

$$+ \frac{(Aa^3 + Ba^3) \sin(2 dx + 2 c)}{d} + \frac{(26 Aa^3 + 23 Ba^3) \sin(dx + c)}{8 d}$$

```
input integrate(cos(d*x+c)*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/80*B*a^3*sin(5*d*x + 5*c)/d + 1/8*(15*A*a^3 + 13*B*a^3)*x + 1/32*(A*a^3 + 3*B*a^3)*sin(4*d*x + 4*c)/d + 1/48*(12*A*a^3 + 17*B*a^3)*sin(3*d*x + 3*c)/d + (A*a^3 + B*a^3)*sin(2*d*x + 2*c)/d + 1/8*(26*A*a^3 + 23*B*a^3)*sin(d*x + c)/d
```

3.20. $\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.20.9 Mupad [B] (verification not implemented)

Time = 1.51 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.80

$$\int \cos(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{15Aa^3}{4} + \frac{13Ba^3}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{35Aa^3}{2} + \frac{91Ba^3}{6}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(32Aa^3 + \frac{416Ba^3}{15}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right) - a^3(15A + 13B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{4d}$$

$$+ \frac{a^3 \operatorname{atan}\left(\frac{a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)(15A + 13B)}{4\left(\frac{15Aa^3}{4} + \frac{13Ba^3}{4}\right)}\right)(15A + 13B)}{4d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3,x)`output `(tan(c/2 + (d*x)/2)*((49*A*a^3)/4 + (51*B*a^3)/4) + tan(c/2 + (d*x)/2)^9*(
(15*A*a^3)/4 + (13*B*a^3)/4) + tan(c/2 + (d*x)/2)^7*((35*A*a^3)/2 + (91*B*
a^3)/6) + tan(c/2 + (d*x)/2)^5*((61*A*a^3)/2 + (133*B*a^3)/6) + tan(c/2 +
(d*x)/2)^3*((32*A*a^3 + (416*B*a^3)/15))/(d*(5*tan(c/2 + (d*x)/2)^2 + 10*ta
n(c/2 + (d*x)/2)^4 + 10*tan(c/2 + (d*x)/2)^6 + 5*tan(c/2 + (d*x)/2)^8 + ta
n(c/2 + (d*x)/2)^10 + 1)) - (a^3*(15*A + 13*B)*(atan(tan(c/2 + (d*x)/2)) -
(d*x)/2))/(4*d) + (a^3*atan((a^3*tan(c/2 + (d*x)/2)*(15*A + 13*B))/(4*((1
5*A*a^3)/4 + (13*B*a^3)/4)))*(15*A + 13*B))/(4*d)`

3.21 $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$

3.21.1	Optimal result	381
3.21.2	Mathematica [A] (verified)	381
3.21.3	Rubi [A] (verified)	382
3.21.4	Maple [A] (verified)	383
3.21.5	Fricas [A] (verification not implemented)	384
3.21.6	Sympy [B] (verification not implemented)	385
3.21.7	Maxima [A] (verification not implemented)	385
3.21.8	Giac [A] (verification not implemented)	386
3.21.9	Mupad [B] (verification not implemented)	386

3.21.1 Optimal result

Integrand size = 23, antiderivative size = 116

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{5}{8}a^3(4A + 3B)x + \frac{a^3(4A + 3B) \sin(c + dx)}{d} + \frac{3a^3(4A + 3B) \cos(c + dx) \sin(c + dx)}{8d}$$

$$+ \frac{B(a + a \cos(c + dx))^3 \sin(c + dx)}{4d} - \frac{a^3(4A + 3B) \sin^3(c + dx)}{12d}$$

```
output 5/8*a^3*(4*A+3*B)*x+a^3*(4*A+3*B)*sin(d*x+c)/d+3/8*a^3*(4*A+3*B)*cos(d*x+c)
)*sin(d*x+c)/d+1/4*B*(a+a*cos(d*x+c))^3*sin(d*x+c)/d-1/12*a^3*(4*A+3*B)*si
n(d*x+c)^3/d
```

3.21.2 Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{a^3 \sin(c + dx) \left(30(4A + 3B) \arcsin \left(\sqrt{\sin^2 \left(\frac{1}{2}(c + dx) \right)} \right) + (88A + 72B + 9(4A + 5B) \cos(c + dx) + 8(A + B \cos(c + dx))) \right)}{24d \sqrt{\sin^2(c + dx)}}$$

```
input Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]
```

output $(a^3 \sin[c + d*x] * (30*(4*A + 3*B) * \text{ArcSin}[\text{Sqrt}[\text{Sin}[(c + d*x)/2]^2]] + (88*A + 72*B + 9*(4*A + 5*B) * \text{Cos}[c + d*x] + 8*(A + 3*B) * \text{Cos}[c + d*x]^2 + 6*B * \text{Cos}[c + d*x]^3) * \text{Sqrt}[\text{Sin}[c + d*x]^2])) / (24*d * \text{Sqrt}[\text{Sin}[c + d*x]^2])$

3.21.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.87, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {3042, 3230, 3042, 3124, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx \\
 & \quad \downarrow \text{3230} \\
 & \frac{1}{4}(4A + 3B) \int (\cos(c + dx)a + a)^3 dx + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(4A + 3B) \int \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^3 dx + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\
 & \quad \downarrow \text{3124} \\
 & \frac{1}{4}(4A + 3B) \int (\cos^3(c + dx)a^3 + 3 \cos^2(c + dx)a^3 + 3 \cos(c + dx)a^3 + a^3) dx + \\
 & \quad \frac{B \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{4}(4A + 3B) \left(-\frac{a^3 \sin^3(c + dx)}{3d} + \frac{4a^3 \sin(c + dx)}{d} + \frac{3a^3 \sin(c + dx) \cos(c + dx)}{2d} + \frac{5a^3 x}{2} \right) + \\
 & \quad \frac{B \sin(c + dx)(a \cos(c + dx) + a)^3}{4d}
 \end{aligned}$$

input $\text{Int}[(a + a \text{Cos}[c + d*x])^3 * (A + B * \text{Cos}[c + d*x]), x]$

output $(B*(a + a*\cos[c + d*x])^3*\sin[c + d*x]/(4*d) + ((4*A + 3*B)*((5*a^3*x)/2 + (4*a^3*\sin[c + d*x])/d + (3*a^3*\cos[c + d*x]*\sin[c + d*x])/(2*d) - (a^3*\sin[c + d*x]^3)/(3*d)))/4$

3.21.3.1 Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3124 $\text{Int}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrig}[(a + b*\sin[c + d*x])^n, x], x] \text{ ; FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{IGtQ}[n, 0]$

rule 3230 $\text{Int}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]^{(m_)}*((c_) + (d_)*\sin[(e_) + (f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(-d)*\cos[e + f*x]*((a + b*\sin[e + f*x])^m/(f*(m + 1))), x] + \text{Simp}[(a*d*m + b*c*(m + 1))/(b*(m + 1)) \ \text{Int}[(a + b*\sin[e + f*x])^m, x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, m\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{!LtQ}[m, -2^{(-1)}]$

3.21.4 Maple [A] (verified)

Time = 2.52 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.68

method	result
parallelrisch	$5 \left(\frac{\left(\frac{3A}{2} + 2B\right) \sin(2dx+2c)}{5} + \frac{\left(\frac{A}{3} + B\right) \sin(3dx+3c)}{10} + \frac{\sin(4dx+4c)B}{80} + \frac{\left(3A + \frac{13B}{5}\right) \sin(dx+c)}{2} + dx \left(A + \frac{3B}{4}\right) \right) a^3$
risch	$\frac{5a^3 Ax}{2} + \frac{15a^3 Bx}{8} + \frac{15a^3 A \sin(dx+c)}{4d} + \frac{13a^3 B \sin(dx+c)}{4d} + \frac{\sin(4dx+4c)B a^3}{32d} + \frac{\sin(3dx+3c)A a^3}{12d} + \frac{\sin(3dx+3c)A a^3}{12d} + \frac{\sin(3dx+3c)A a^3}{12d}$
parts	$a^3 Ax + \frac{(A a^3 + 3B a^3)(2 + \cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{(3A a^3 + B a^3) \sin(dx+c)}{d} + \frac{(3A a^3 + 3B a^3) \left(\frac{\cos(dx+c) \sin(dx+c)}{2}\right)}{d}$
derivativedivides	$\frac{A a^3 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + B a^3 \left(\frac{\left(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}\right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 3A a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
default	$\frac{A a^3 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + B a^3 \left(\frac{\left(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}\right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 3A a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
norman	$\frac{5a^3(4A+3B)x}{8} + \frac{73a^3(4A+3B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{12d} + \frac{55a^3(4A+3B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{12d} + \frac{5a^3(4A+3B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{4d} + \frac{5a^3(4A+3B)x\left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2d} + \dots$

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 5/2*(1/5*(3/2*A+2*B)*sin(2*d*x+2*c)+1/10*(1/3*A+B)*sin(3*d*x+3*c)+1/80*sin(4*d*x+4*c)*B+1/2*(3*A+13/5*B)*sin(d*x+c)+d*x*(A+3/4*B))*a^3/d
```

3.21.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.78

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{15(4A + 3B)a^3 dx + (6Ba^3 \cos(dx + c))^3 + 8(A + 3B)a^3 \cos(dx + c)^2 + 9(4A + 5B)a^3 \cos(dx + c) + 8Aa^3}{24d}$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/24*(15*(4*A + 3*B)*a^3*d*x + (6*B*a^3*cos(d*x + c))^3 + 8*(A + 3*B)*a^3*cos(d*x + c)^2 + 9*(4*A + 5*B)*a^3*cos(d*x + c) + 8*(11*A + 9*B)*a^3*sin(d*x + c))/d
```

3.21. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$

3.21.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 371 vs. $2(107) = 214$.

Time = 0.22 (sec) , antiderivative size = 371, normalized size of antiderivative = 3.20

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{3Aa^3 x \sin^2(c+dx)}{2} + \frac{3Aa^3 x \cos^2(c+dx)}{2} + Aa^3 x + \frac{2Aa^3 \sin^3(c+dx)}{3d} + \frac{Aa^3 \sin(c+dx) \cos^2(c+dx)}{d} + \frac{3Aa^3 \sin(c+dx) \cos(c+dx)}{2d} \\ x(A + B \cos(c)) (a \cos(c) + a)^3 \end{cases}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)`

output `Piecewise((3*A*a**3*x*sin(c + d*x)**2/2 + 3*A*a**3*x*cos(c + d*x)**2/2 + A*a**3*x + 2*A*a**3*sin(c + d*x)**3/(3*d) + A*a**3*sin(c + d*x)*cos(c + d*x)**2/d + 3*A*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + 3*A*a**3*sin(c + d*x)/d + 3*B*a**3*x*sin(c + d*x)**4/8 + 3*B*a**3*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*B*a**3*x*sin(c + d*x)**2/2 + 3*B*a**3*x*cos(c + d*x)**4/8 + 3*B*a**3*x*cos(c + d*x)**2/2 + 3*B*a**3*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 2*B*a**3*sin(c + d*x)**3/d + 5*B*a**3*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 3*B*a**3*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + B*a**3*sin(c + d*x)/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**3, True))`

3.21.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.44

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx =$$

$$\frac{32 (\sin(dx + c)^3 - 3 \sin(dx + c)) Aa^3 - 72 (2 dx + 2c + \sin(2 dx + 2c)) Aa^3 - 96 (dx + c) Aa^3 + 96 (s$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `-1/96*(32*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^3 - 72*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^3 - 96*(d*x + c)*A*a^3 + 96*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^3 - 3*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^3 - 72*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^3 - 288*A*a^3*sin(d*x + c) - 96*B*a^3*sin(d*x + c))/d`

3.21. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$

3.21.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.97

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{Ba^3 \sin(4dx + 4c)}{32d} + \frac{5}{8} (4Aa^3 + 3Ba^3)x + \frac{(Aa^3 + 3Ba^3) \sin(3dx + 3c)}{12d}$$

$$+ \frac{(3Aa^3 + 4Ba^3) \sin(2dx + 2c)}{4d} + \frac{(15Aa^3 + 13Ba^3) \sin(dx + c)}{4d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/32*B*a^3*sin(4*d*x + 4*c)/d + 5/8*(4*A*a^3 + 3*B*a^3)*x + 1/12*(A*a^3 + 3*B*a^3)*sin(3*d*x + 3*c)/d + 1/4*(3*A*a^3 + 4*B*a^3)*sin(2*d*x + 2*c)/d + 1/4*(15*A*a^3 + 13*B*a^3)*sin(d*x + c)/d`**3.21.9 Mupad [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.16

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{5Aa^3x}{2} + \frac{15Ba^3x}{8} + \frac{15Aa^3 \sin(c + dx)}{4d} + \frac{13Ba^3 \sin(c + dx)}{4d}$$

$$+ \frac{3Aa^3 \sin(2c + 2dx)}{4d} + \frac{Aa^3 \sin(3c + 3dx)}{12d}$$

$$+ \frac{Ba^3 \sin(2c + 2dx)}{d} + \frac{Ba^3 \sin(3c + 3dx)}{4d} + \frac{Ba^3 \sin(4c + 4dx)}{32d}$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3,x)`output `(5*A*a^3*x)/2 + (15*B*a^3*x)/8 + (15*A*a^3*sin(c + d*x))/(4*d) + (13*B*a^3*sin(c + d*x))/(4*d) + (3*A*a^3*sin(2*c + 2*d*x))/(4*d) + (A*a^3*sin(3*c + 3*d*x))/(12*d) + (B*a^3*sin(2*c + 2*d*x))/d + (B*a^3*sin(3*c + 3*d*x))/(4*d) + (B*a^3*sin(4*c + 4*d*x))/(32*d)`

3.22 $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$

3.22.1	Optimal result	387
3.22.2	Mathematica [A] (verified)	387
3.22.3	Rubi [A] (verified)	388
3.22.4	Maple [A] (verified)	391
3.22.5	Fricas [A] (verification not implemented)	392
3.22.6	Sympy [F]	393
3.22.7	Maxima [A] (verification not implemented)	393
3.22.8	Giac [A] (verification not implemented)	394
3.22.9	Mupad [B] (verification not implemented)	394

3.22.1 Optimal result

Integrand size = 29, antiderivative size = 111

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{1}{2}a^3(7A + 5B)x + \frac{a^3 A \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{5a^3(A + B) \sin(c + dx)}{2d}$$

$$+ \frac{aB(a + a \cos(c + dx))^2 \sin(c + dx)}{3d} + \frac{(3A + 5B)(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{6d}$$

```
output 1/2*a^3*(7*A+5*B)*x+a^3*A*arctanh(sin(d*x+c))/d+5/2*a^3*(A+B)*sin(d*x+c)/d
+1/3*a*B*(a+a*cos(d*x+c))^2*sin(d*x+c)/d+1/6*(3*A+5*B)*(a^3+a^3*cos(d*x+c)
)*sin(d*x+c)/d
```

3.22.2 Mathematica [A] (verified)

Time = 1.18 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.02

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{a^3(42Adx + 30Bdx - 12A \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 12A \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{12d}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(a^3*(42*A*d*x + 30*B*d*x - 12*A*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 12*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 9*(4*A + 5*B)*Sin[c + d*x] + 3*(A + 3*B)*Sin[2*(c + d*x)] + B*Sin[3*(c + d*x)])/(12*d)`

3.22.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$, Rules used = {3042, 3455, 3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c + dx)(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3455} \\
 & \frac{1}{3} \int (\cos(c + dx)a + a)^2 (3aA + a(3A + 5B) \cos(c + dx)) \sec(c + dx) dx + \\
 & \quad \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^2}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 (3aA + a(3A + 5B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \\
 & \quad \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^2}{3d} \\
 & \quad \downarrow \text{3455} \\
 & \frac{1}{3} \left(\frac{1}{2} \int 3(\cos(c + dx)a + a) (2Aa^2 + 5(A + B) \cos(c + dx)a^2) \sec(c + dx) dx + \frac{(3A + 5B) \sin(c + dx) (a^3 \cos(c + dx) + a^2 \sin(c + dx))}{2d} \right) \\
 & \quad \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^2}{3d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.22. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$

$$\frac{1}{3} \left(\frac{3}{2} \int (\cos(c+dx)a+a)(2Aa^2+5(A+B)\cos(c+dx)a^2)\sec(c+dx)dx + \frac{(3A+5B)\sin(c+dx)(a^3\cos(c+dx)+a^2)}{2d} \right. \\ \left. \frac{aB\sin(c+dx)(a\cos(c+dx)+a)^2}{3d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(2Aa^2+5(A+B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{(3A+5B)\sin(c+dx)(a^3\cos(c+dx)+a^2)}{2d} \right. \\ \left. \frac{aB\sin(c+dx)(a\cos(c+dx)+a)^2}{3d} \right) \\ \downarrow \text{3447}$$

$$\frac{1}{3} \left(\frac{3}{2} \int (5(A+B)\cos^2(c+dx)a^3+2Aa^3+(2Aa^3+5(A+B)a^3)\cos(c+dx))\sec(c+dx)dx + \frac{(3A+5B)\sin(c+dx)(a^3\cos(c+dx)+a^2)}{2d} \right. \\ \left. \frac{aB\sin(c+dx)(a\cos(c+dx)+a)^2}{3d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{5(A+B)\sin(c+dx+\frac{\pi}{2})^2 a^3+2Aa^3+(2Aa^3+5(A+B)a^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{(3A+5B)\sin(c+dx)(a^3\cos(c+dx)+a^2)}{2d} \right. \\ \left. \frac{aB\sin(c+dx)(a\cos(c+dx)+a)^2}{3d} \right) \\ \downarrow \text{3502}$$

$$\frac{1}{3} \left(\frac{3}{2} \left(\int (2Aa^3+(7A+5B)\cos(c+dx)a^3)\sec(c+dx)dx + \frac{5a^3(A+B)\sin(c+dx)}{d} \right) + \frac{(3A+5B)\sin(c+dx)(a^3\cos(c+dx)+a^2)}{2d} \right. \\ \left. \frac{aB\sin(c+dx)(a\cos(c+dx)+a)^2}{3d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(\frac{3}{2} \left(\int \frac{2Aa^3+(7A+5B)\sin(c+dx+\frac{\pi}{2})a^3}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{5a^3(A+B)\sin(c+dx)}{d} \right) + \frac{(3A+5B)\sin(c+dx)(a^3\cos(c+dx)+a^2)}{2d} \right. \\ \left. \frac{aB\sin(c+dx)(a\cos(c+dx)+a)^2}{3d} \right) \\ \downarrow \text{3214}$$

$$\frac{1}{3} \left(\frac{3}{2} \left(2a^3 A \int \sec(c+dx) dx + \frac{5a^3(A+B)\sin(c+dx)}{d} + a^3 x(7A+5B) \right) + \frac{(3A+5B)\sin(c+dx)(a^3 \cos(c+dx) + aB \sin(c+dx)(a \cos(c+dx) + a)^2)}{2d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \left(2a^3 A \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{5a^3(A+B)\sin(c+dx)}{d} + a^3 x(7A+5B) \right) + \frac{(3A+5B)\sin(c+dx)(a^3 \cos(c+dx) + aB \sin(c+dx)(a \cos(c+dx) + a)^2)}{2d} \right)$$

↓ 4257

$$\frac{1}{3} \left(\frac{3}{2} \left(\frac{2a^3 A \operatorname{arctanh}(\sin(c+dx))}{d} + \frac{5a^3(A+B)\sin(c+dx)}{d} + a^3 x(7A+5B) \right) + \frac{(3A+5B)\sin(c+dx)(a^3 \cos(c+dx) + aB \sin(c+dx)(a \cos(c+dx) + a)^2)}{2d} \right)$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x], x]`

output `(a*B*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + (((3*A + 5*B)*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(2*d) + (3*(a^3*(7*A + 5*B)*x + (2*a^3*A*ArcTanh[Sin[c + d*x]])/d + (5*a^3*(A + B)*Sin[c + d*x])/d))/2)/3`

3.22.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.22.4 Maple [A] (verified)

Time = 2.55 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.84

method	result
parallelrisch	$\frac{a^3 \left(A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + \frac{(-A - 3B) \sin(2dx + 2c)}{4} - \frac{B \sin(3dx + 3c)}{12} + 3\left(-A - \frac{5B}{4}\right) \sin(dx + c) \right)}{d}$
parts	$\frac{A a^3 \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(A a^3 + 3B a^3) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d} + \frac{(3A a^3 + B a^3)(dx+c)}{d} + \frac{(3A a^3 - B a^3) \sin(dx+c)}{d}$
derivativedivides	$\frac{A a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + \frac{B a^3 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + 3A a^3 \sin(dx+c) + 3B a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d}$
default	$\frac{A a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + \frac{B a^3 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + 3A a^3 \sin(dx+c) + 3B a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d}$
risch	$\frac{7a^3 Ax}{2} + \frac{5a^3 Bx}{2} - \frac{3ie^{i(dx+c)} A a^3}{2d} - \frac{15ie^{i(dx+c)} B a^3}{8d} + \frac{3ie^{-i(dx+c)} A a^3}{2d} + \frac{15ie^{-i(dx+c)} B a^3}{8d} + \frac{A a^3 \ln(e^{i(dx+c)} + 1)}{d}$
norman	$\frac{\left(\frac{7}{2} A a^3 + \frac{5}{2} B a^3\right) x + \left(\frac{7}{2} A a^3 + \frac{5}{2} B a^3\right) x \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (14A a^3 + 10B a^3) x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (14A a^3 + 10B a^3) x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6d}$

```
input int((a+cos(d*x+c))*a)^3*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)
```

```
output -a^3*(A*ln(tan(1/2*d*x+1/2*c)-1)-A*ln(tan(1/2*d*x+1/2*c)+1)+1/4*(-A-3*B)*sin(2*d*x+2*c)-1/12*B*sin(3*d*x+3*c)+3*(-A-5/4*B)*sin(d*x+c)-7/2*(A+5/7*B)*x*d)/d
```

3.22.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.92

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{3(7A + 5B)a^3 dx + 3Aa^3 \log(\sin(dx + c) + 1) - 3Aa^3 \log(-\sin(dx + c) + 1) + (2Ba^3 \cos(dx + c))^2 + 2(9A + 11B)a^3 \sin(dx + c)}{6d}$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fracas")
```

```
output 1/6*(3*(7*A + 5*B)*a^3*d*x + 3*A*a^3*log(sin(d*x + c) + 1) - 3*A*a^3*log(-sin(d*x + c) + 1) + (2*B*a^3*cos(d*x + c))^2 + 3*(A + 3*B)*a^3*cos(d*x + c) + 2*(9*A + 11*B)*a^3*sin(d*x + c))/d
```

3.22. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$

3.22.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= a^3 \left(\int A \sec(c + dx) dx + \int 3A \cos(c + dx) \sec(c + dx) dx \right. \\ &\quad + \int 3A \cos^2(c + dx) \sec(c + dx) dx + \int A \cos^3(c + dx) \sec(c + dx) dx \\ &\quad + \int B \cos(c + dx) \sec(c + dx) dx + \int 3B \cos^2(c + dx) \sec(c + dx) dx \\ &\quad \left. + \int 3B \cos^3(c + dx) \sec(c + dx) dx + \int B \cos^4(c + dx) \sec(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `a**3*(Integral(A*sec(c + d*x), x) + Integral(3*A*cos(c + d*x)*sec(c + d*x), x) + Integral(3*A*cos(c + d*x)**2*sec(c + d*x), x) + Integral(A*cos(c + d*x)**3*sec(c + d*x), x) + Integral(B*cos(c + d*x)*sec(c + d*x), x) + Integral(3*B*cos(c + d*x)**2*sec(c + d*x), x) + Integral(3*B*cos(c + d*x)**3*sec(c + d*x), x) + Integral(B*cos(c + d*x)**4*sec(c + d*x), x))`

3.22.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.27

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{3(2dx + 2c + \sin(2dx + 2c))Aa^3 + 36(dx + c)Aa^3 - 4(\sin(dx + c)^3 - 3\sin(dx + c))Ba^3 + 9(2dx + \dots}{\dots} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `1/12*(3*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^3 + 36*(d*x + c)*A*a^3 - 4*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^3 + 9*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^3 + 12*(d*x + c)*B*a^3 + 12*A*a^3*log(sec(d*x + c) + tan(d*x + c)) + 36*A*a^3*sin(d*x + c) + 36*B*a^3*sin(d*x + c))/d`

3.22.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.62

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{6 A a^3 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 6 A a^3 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + 3 (7 A a^3 + 5 B a^3) (dx + c) + \frac{2 (15 A^2 a^3 \tan^2 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 15 B a^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 36 A^2 a^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 + 40 B a^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 + 21 A a^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 33 B a^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right))}{(\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 + 1)^3} / d$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="gias")
```

```
output 1/6*(6*A*a^3*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 6*A*a^3*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + 3*(7*A*a^3 + 5*B*a^3)*(d*x + c) + 2*(15*A*a^3*tan(1/2*d*x + 1/2*c)^5 + 15*B*a^3*tan(1/2*d*x + 1/2*c)^5 + 36*A*a^3*tan(1/2*d*x + 1/2*c)^3 + 40*B*a^3*tan(1/2*d*x + 1/2*c)^3 + 21*A*a^3*tan(1/2*d*x + 1/2*c) + 33*B*a^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^3/d
```

3.22.9 Mupad [B] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.60

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{3 A a^3 \sin(c + dx)}{d} + \frac{15 B a^3 \sin(c + dx)}{4 d} + \frac{7 A a^3 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d}$$

$$+ \frac{2 A a^3 \operatorname{atanh} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} + \frac{5 B a^3 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d}$$

$$+ \frac{A a^3 \sin(2c + 2dx)}{4 d} + \frac{3 B a^3 \sin(2c + 2dx)}{4 d} + \frac{B a^3 \sin(3c + 3dx)}{12 d}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x),x)
```

```
output (3*A*a^3*sin(c + d*x))/d + (15*B*a^3*sin(c + d*x))/(4*d) + (7*A*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*A*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (5*B*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (A*a^3*sin(2*c + 2*d*x))/(4*d) + (3*B*a^3*sin(2*c + 2*d*x))/(4*d) + (B*a^3*sin(3*c + 3*d*x))/(12*d)
```

3.23 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.23.1	Optimal result	395
3.23.2	Mathematica [B] (verified)	396
3.23.3	Rubi [A] (verified)	396
3.23.4	Maple [A] (verified)	400
3.23.5	Fricas [A] (verification not implemented)	401
3.23.6	Sympy [F]	401
3.23.7	Maxima [A] (verification not implemented)	402
3.23.8	Giac [A] (verification not implemented)	402
3.23.9	Mupad [B] (verification not implemented)	403

3.23.1 Optimal result

Integrand size = 31, antiderivative size = 110

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{1}{2}a^3(6A + 7B)x + \frac{a^3(3A + B)\operatorname{arctanh}(\sin(c + dx))}{d} + \frac{5a^3B \sin(c + dx)}{2d}$$

$$- \frac{(2A - B)(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{2d} + \frac{aA(a + a \cos(c + dx))^2 \tan(c + dx)}{d}$$

```
output 1/2*a^3*(6*A+7*B)*x+a^3*(3*A+B)*arctanh(sin(d*x+c))/d+5/2*a^3*B*sin(d*x+c)
/d-1/2*(2*A-B)*(a^3+a^3*cos(d*x+c))*sin(d*x+c)/d+a*A*(a+a*cos(d*x+c))^2*tan(d*x+c)/d
```

3.23.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 272 vs. $2(110) = 220$.

Time = 2.64 (sec) , antiderivative size = 272, normalized size of antiderivative = 2.47

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{1}{32} a^3 (1 + \cos(c + dx))^3 \sec^6\left(\frac{1}{2}(c + dx)\right) \left(2(6A + 7B)x \right. \\ \left. - \frac{4(3A + B) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)}{d} \right. \\ \left. + \frac{4(3A + B) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)}{d} + \frac{4(A + 3B) \cos(dx) \sin(c)}{d} \right. \\ \left. + \frac{B \cos(2dx) \sin(2c)}{d} + \frac{4(A + 3B) \cos(c) \sin(dx)}{d} + \frac{B \cos(2c) \sin(2dx)}{d} \right. \\ \left. + \frac{4A \sin\left(\frac{dx}{2}\right)}{d \left(\cos\left(\frac{c}{2}\right) - \sin\left(\frac{c}{2}\right)\right) \left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)} \right. \\ \left. + \frac{4A \sin\left(\frac{dx}{2}\right)}{d \left(\cos\left(\frac{c}{2}\right) + \sin\left(\frac{c}{2}\right)\right) \left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)} \right)$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `(a^3*(1 + Cos[c + d*x])^3*Sec[(c + d*x)/2]^6*(2*(6*A + 7*B)*x - (4*(3*A + B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]])/d + (4*(3*A + B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/d + (4*(A + 3*B)*Cos[d*x]*Sin[c])/d + (B*Cos[2*d*x]*Sin[2*c])/d + (4*(A + 3*B)*Cos[c]*Sin[d*x])/d + (B*Cos[2*c]*Sin[2*d*x])/d + (4*A*Sin[(d*x)/2])/(d*(Cos[c/2] - Sin[c/2])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) + (4*A*Sin[(d*x)/2])/(d*(Cos[c/2] + Sin[c/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])))`

3.23.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3454, 3042, 3455, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.23. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$

$$\begin{aligned}
& \int \sec^2(c+dx)(a \cos(c+dx) + a)^3(A + B \cos(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a \sin(c+dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^2} dx \\
& \quad \downarrow \text{3454} \\
& \int (\cos(c+dx)a + a)^2(a(3A + B) - a(2A - B) \cos(c+dx)) \sec(c+dx) dx + \\
& \quad \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^2 (a(3A + B) - a(2A - B) \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})} dx + \\
& \quad \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3455} \\
& \frac{1}{2} \int (\cos(c+dx)a + a) (2(3A + B)a^2 + 5B \cos(c+dx)a^2) \sec(c+dx) dx - \\
& \frac{(2A - B) \sin(c+dx) (a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a) (2(3A + B)a^2 + 5B \sin(c+dx + \frac{\pi}{2})a^2)}{\sin(c+dx + \frac{\pi}{2})} dx - \\
& \frac{(2A - B) \sin(c+dx) (a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3447} \\
& \frac{1}{2} \int (5B \cos^2(c+dx)a^3 + 2(3A + B)a^3 + (5Ba^3 + 2(3A + B)a^3) \cos(c+dx)) \sec(c+dx) dx - \\
& \frac{(2A - B) \sin(c+dx) (a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{5B \sin(c+dx + \frac{\pi}{2})^2 a^3 + 2(3A + B)a^3 + (5Ba^3 + 2(3A + B)a^3) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})} dx - \\
& \frac{(2A - B) \sin(c+dx) (a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3502}
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(\int \frac{(2(3A+B)a^3 + (6A+7B)\cos(c+dx)a^3) \sec(c+dx) dx + \frac{5a^3 B \sin(c+dx)}{d}}{(2A-B)\sin(c+dx)(a^3 \cos(c+dx) + a^3)} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \right) - \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\int \frac{2(3A+B)a^3 + (6A+7B)\sin(c+dx + \frac{\pi}{2})a^3}{\sin(c+dx + \frac{\pi}{2})} dx + \frac{5a^3 B \sin(c+dx)}{d} \right) - \\
& \frac{(2A-B)\sin(c+dx)(a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3214} \\
& \frac{1}{2} \left(2a^3(3A+B) \int \sec(c+dx) dx + a^3 x(6A+7B) + \frac{5a^3 B \sin(c+dx)}{d} \right) - \\
& \frac{(2A-B)\sin(c+dx)(a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(2a^3(3A+B) \int \csc\left(c+dx + \frac{\pi}{2}\right) dx + a^3 x(6A+7B) + \frac{5a^3 B \sin(c+dx)}{d} \right) - \\
& \frac{(2A-B)\sin(c+dx)(a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d} \\
& \quad \downarrow \text{4257} \\
& \frac{1}{2} \left(\frac{2a^3(3A+B)\operatorname{arctanh}(\sin(c+dx))}{d} + a^3 x(6A+7B) + \frac{5a^3 B \sin(c+dx)}{d} \right) - \\
& \frac{(2A-B)\sin(c+dx)(a^3 \cos(c+dx) + a^3)}{2d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^2}{d}
\end{aligned}$$

input `Int[(a + a*cos[c + d*x])^3*(A + B*cos[c + d*x])*Sec[c + d*x]^2,x]`

output `-1/2*((2*A - B)*(a^3 + a^3*cos[c + d*x])*Sin[c + d*x])/d + (a^3*(6*A + 7*B)*x + (2*a^3*(3*A + B)*ArcTanh[Sin[c + d*x]])/d + (5*a^3*B*SIN[c + d*x])/d)/2 + (a*A*(a + a*cos[c + d*x])^2*Tan[c + d*x])/d`

3.23.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.23.4 Maple [A] (verified)

Time = 2.52 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.09

method	result
parts	$\frac{A a^3 \tan(dx+c)}{d} + \frac{(A a^3 + 3B a^3) \sin(dx+c)}{d} + \frac{(3A a^3 + B a^3) \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(3A a^3 + 3B a^3)(dx+c)}{d}$
parallelrisch	$\frac{3 \left(\cos(dx+c) \left(A + \frac{B}{3} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) - \cos(dx+c) \left(A + \frac{B}{3} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) + \left(-\frac{A}{6} - \frac{B}{2} \right) \sin(2dx+2c) - \frac{B \sin(dx+c)}{2} \right)}{d \cos(dx+c)}$
derivativedivides	$\frac{A a^3 \sin(dx+c) + B a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 3A a^3 (dx+c) + 3B a^3 \sin(dx+c) + 3A a^3 \ln(\sec(dx+c) + \tan(dx+c))}{d}$
default	$\frac{A a^3 \sin(dx+c) + B a^3 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 3A a^3 (dx+c) + 3B a^3 \sin(dx+c) + 3A a^3 \ln(\sec(dx+c) + \tan(dx+c))}{d}$
risch	$3a^3 Ax + \frac{7a^3 Bx}{2} - \frac{ib a^3 e^{2i(dx+c)}}{8d} - \frac{ie^{i(dx+c)} A a^3}{2d} - \frac{3ie^{i(dx+c)} B a^3}{2d} + \frac{ie^{-i(dx+c)} A a^3}{2d} + \frac{3ie^{-i(dx+c)} B a^3}{2d}$
norman	$\frac{\left(-\frac{7}{2} B a^3 - 3A a^3 \right) x + \left(-\frac{21}{2} B a^3 - 9A a^3 \right) x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \left(\frac{7}{2} B a^3 + 3A a^3 \right) x \left(\tan^{10} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \left(\frac{21}{2} B a^3 + 9A a^3 \right) x \left(\tan^{14} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d}$

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBO
SE)
```

```
output A*a^3/d*tan(d*x+c)+(A*a^3+3*B*a^3)/d*sin(d*x+c)+(3*A*a^3+B*a^3)/d*ln(sec(d
*x+c)+tan(d*x+c))+(3*A*a^3+3*B*a^3)/d*(d*x+c)+B*a^3/d*(1/2*cos(d*x+c)*sin(
d*x+c)+1/2*d*x+1/2*c)
```

3.23. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.23.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.15

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{(6A + 7B)a^3 dx \cos(dx + c) + (3A + B)a^3 \cos(dx + c) \log(\sin(dx + c) + 1) - (3A + B)a^3 \cos(dx + c)}{2d \cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/2*((6*A + 7*B)*a^3*d*x*cos(d*x + c) + (3*A + B)*a^3*cos(d*x + c)*log(sin(d*x + c) + 1) - (3*A + B)*a^3*cos(d*x + c)*log(-sin(d*x + c) + 1) + (B*a^3*cos(d*x + c)^2 + 2*(A + 3*B)*a^3*cos(d*x + c) + 2*A*a^3)*sin(d*x + c))/(d*cos(d*x + c))`

3.23.6 Sympy [F]

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= a^3 \left(\int A \sec^2(c + dx) dx + \int 3A \cos(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int 3A \cos^2(c + dx) \sec^2(c + dx) dx + \int A \cos^3(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int B \cos(c + dx) \sec^2(c + dx) dx + \int 3B \cos^2(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int 3B \cos^3(c + dx) \sec^2(c + dx) dx + \int B \cos^4(c + dx) \sec^2(c + dx) dx \right)$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `a**3*(Integral(A*sec(c + d*x)**2, x) + Integral(3*A*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(3*A*cos(c + d*x)**2*sec(c + d*x)**2, x) + Integral(A*cos(c + d*x)**3*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(3*B*cos(c + d*x)**2*sec(c + d*x)**2, x) + Integral(3*B*cos(c + d*x)**3*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)**4*sec(c + d*x)**2, x))`

3.23.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.27

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{12(dx + c)Aa^3 + (2dx + 2c + \sin(2dx + 2c))Ba^3 + 12(dx + c)Ba^3 + 6Aa^3(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 4Aa^3 \sin(dx + c) + 12Ba^3 \sin(dx + c) + 4Aa^3 \tan(dx + c)}{d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `1/4*(12*(d*x + c)*A*a^3 + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^3 + 12*(d*x + c)*B*a^3 + 6*A*a^3*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*B*a^3*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 4*A*a^3*sin(d*x + c) + 12*B*a^3*sin(d*x + c) + 4*A*a^3*tan(d*x + c))/d`

3.23.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.75

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$\frac{4Aa^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)}{\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1} - (6Aa^3 + 7Ba^3)(dx + c) - 2(3Aa^3 + Ba^3) \log(|\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1|) + 2(3Aa^3 + Ba^3) \log(|\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1|) + 4Aa^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 12Ba^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)}{d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `-1/2*(4*A*a^3*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 1) - (6*A*a^3 + 7*B*a^3)*(d*x + c) - 2*(3*A*a^3 + B*a^3)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) + 2*(3*A*a^3 + B*a^3)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(2*A*a^3*tan(1/2*d*x + 1/2*c)^3 + 5*B*a^3*tan(1/2*d*x + 1/2*c)^3 + 2*A*a^3*tan(1/2*d*x + 1/2*c) + 7*B*a^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)/d`

3.23.9 Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.79

$$\begin{aligned}
& \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx \\
&= \frac{A a^3 \sin(c + dx)}{d} + \frac{3 B a^3 \sin(c + dx)}{d} + \frac{6 A a^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{6 A a^3 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{7 B a^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{2 B a^3 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{A a^3 \sin(c + dx)}{d \cos(c + dx)} + \frac{B a^3 \cos(c + dx) \sin(c + dx)}{2d}
\end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^2,x)`output `(A*a^3*sin(c + d*x))/d + (3*B*a^3*sin(c + d*x))/d + (6*A*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (6*A*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (7*B*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (A*a^3*sin(c + d*x))/(d*cos(c + d*x)) + (B*a^3*cos(c + d*x)*sin(c + d*x))/(2*d)`

3.24 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.24.1	Optimal result	404
3.24.2	Mathematica [A] (verified)	405
3.24.3	Rubi [A] (verified)	405
3.24.4	Maple [A] (verified)	409
3.24.5	Fricas [A] (verification not implemented)	409
3.24.6	Sympy [F]	410
3.24.7	Maxima [A] (verification not implemented)	410
3.24.8	Giac [A] (verification not implemented)	411
3.24.9	Mupad [B] (verification not implemented)	412

3.24.1 Optimal result

Integrand size = 31, antiderivative size = 114

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= a^3(A + 3B)x + \frac{a^3(7A + 6B)\operatorname{arctanh}(\sin(c + dx))}{2d}$$

$$- \frac{5a^3 A \sin(c + dx)}{2d} + \frac{(2A + B)(a^3 + a^3 \cos(c + dx)) \tan(c + dx)}{d}$$

$$+ \frac{aA(a + a \cos(c + dx))^2 \sec(c + dx) \tan(c + dx)}{2d}$$

```
output a^3*(A+3*B)*x+1/2*a^3*(7*A+6*B)*arctanh(sin(d*x+c))/d-5/2*a^3*A*sin(d*x+c)
/d+(2*A+B)*(a^3+a^3*cos(d*x+c))*tan(d*x+c)/d+1/2*a*A*(a+a*cos(d*x+c))^2*se
c(d*x+c)*tan(d*x+c)/d
```

3.24.2 Mathematica [A] (verified)

Time = 3.26 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.82

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= a^3 \left(4Ac + 12Bc + 4Adx + 12Bdx - 14A \log \left(\cos \left(\frac{1}{2}(c + dx) \right) - \sin \left(\frac{1}{2}(c + dx) \right) \right) - 12B \log \left(\cos \left(\frac{1}{2}(c + dx) \right) \right) \right)$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output $(a^3(4Ac + 12Bc + 4Adx + 12Bdx - 14A \log[\cos[(c + dx)/2] - \sin[(c + dx)/2]] - 12B \log[\cos[(c + dx)/2] - \sin[(c + dx)/2]] + 14A \log[\cos[(c + dx)/2] + \sin[(c + dx)/2]] + 12B \log[\cos[(c + dx)/2] + \sin[(c + dx)/2]] + A/(\cos[(c + dx)/2] - \sin[(c + dx)/2])^2 - A/(\cos[(c + dx)/2] + \sin[(c + dx)/2])^2 + 4B \sin[c + dx] + 4(3A + B) \tan[c + dx]))/(4d)$

3.24.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3454, 3042, 3454, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^3(c + dx)(a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx$$

$$\downarrow \text{3454}$$

$$\frac{1}{2} \int (\cos(c + dx)a + a)^2 (2a(2A + B) - a(A - 2B) \cos(c + dx)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

$$\downarrow \text{3042}$$

3.24. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

$$\frac{1}{2} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 (2a(2A + B) - a(A - 2B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

↓ 3454

$$\frac{1}{2} \left(\int (\cos(c + dx)a + a) (a^2(7A + 6B) - 5a^2A \cos(c + dx)) \sec(c + dx) dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{(\sin(c + dx + \frac{\pi}{2})a + a) (a^2(7A + 6B) - 5a^2A \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

↓ 3447

$$\frac{1}{2} \left(\int (-5A \cos^2(c + dx)a^3 + (7A + 6B)a^3 + (a^3(7A + 6B) - 5a^3A) \cos(c + dx)) \sec(c + dx) dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{-5A \sin(c + dx + \frac{\pi}{2})^2 a^3 + (7A + 6B)a^3 + (a^3(7A + 6B) - 5a^3A) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

↓ 3502

$$\frac{1}{2} \left(\int ((7A + 6B)a^3 + 2(A + 3B) \cos(c + dx)a^3) \sec(c + dx) dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^2}{2d}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{(7A + 6B)a^3 + 2(A + 3B) \sin(c + dx + \frac{\pi}{2}) a^3}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} - \frac{5a^3 A \sin(c + dx)}{d} - \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^2}{2d} \right)$$

↓ 3214

$$\frac{1}{2} \left(a^3(7A + 6B) \int \sec(c + dx) dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} + 2a^3 x(A + 3B) - \frac{5a^3 A \sin(c + dx)}{d} - \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^2}{2d} \right)$$

↓ 3042

$$\frac{1}{2} \left(a^3(7A + 6B) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} + 2a^3 x(A + 3B) - \frac{5a^3 A \sin(c + dx)}{d} - \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^2}{2d} \right)$$

↓ 4257

$$\frac{1}{2} \left(\frac{a^3(7A + 6B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{2(2A + B) \tan(c + dx) (a^3 \cos(c + dx) + a^3)}{d} + 2a^3 x(A + 3B) - \frac{5a^3 A \sin(c + dx)}{d} - \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^2}{2d} \right)$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*(a + a*Cos[c + d*x])^2*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (2*a^3*(A + 3*B)*x + (a^3*(7*A + 6*B)*ArcTanh[Sin[c + d*x]])/d - (5*a^3*A*Sin[c + d*x])/d + (2*(2*A + B)*(a^3 + a^3*Cos[c + d*x])*Tan[c + d*x])/d)/2`

3.24.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.24.4 Maple [A] (verified)

Time = 3.43 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.13

method	result
parts	$\frac{A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d} + \frac{(A a^3 + 3B a^3)(dx+c)}{d} + \frac{(3A a^3 + B a^3) \tan(dx+c)}{d} + \frac{(3A a^3 + B a^3) \ln(\sec(dx+c) + \tan(dx+c))}{d}$
derivativedivides	$\frac{A a^3(dx+c) + B a^3 \sin(dx+c) + 3A a^3 \ln(\sec(dx+c) + \tan(dx+c)) + 3B a^3(dx+c) + 3A a^3 \tan(dx+c) + 3B a^3 \ln(\sec(dx+c) + \tan(dx+c))}{d}$
default	$\frac{A a^3(dx+c) + B a^3 \sin(dx+c) + 3A a^3 \ln(\sec(dx+c) + \tan(dx+c)) + 3B a^3(dx+c) + 3A a^3 \tan(dx+c) + 3B a^3 \ln(\sec(dx+c) + \tan(dx+c))}{d}$
parallelrisch	$-\frac{7 \left(\left(A + \frac{6B}{7} \right) (1 + \cos(2dx+2c)) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) - \left(A + \frac{6B}{7} \right) (1 + \cos(2dx+2c)) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \frac{2dx(A+3B) \cos(dx+c)}{7}}{2d(1 + \cos(2dx+2c))}$
risch	$a^3 A x + 3a^3 B x - \frac{i e^{i(dx+c)} B a^3}{2d} + \frac{i e^{-i(dx+c)} B a^3}{2d} - \frac{i a^3 (A e^{3i(dx+c)} - 6A e^{2i(dx+c)} - 2B e^{2i(dx+c)} - A e^{i(dx+c)})}{d(e^{2i(dx+c)} + 1)^2}$
norman	$\frac{(A a^3 + 3B a^3)x + (-4A a^3 - 12B a^3)x \left(\tan^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (-A a^3 - 3B a^3)x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (-A a^3 - 3B a^3)x \left(\tan^8 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{4d \cos(dx+c)}$

input `int((a+cos(d*x+c))*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `A*a^3/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(A*a^3+3*B*a^3)/d*(d*x+c)+(3*A*a^3+B*a^3)/d*tan(d*x+c)+(3*A*a^3+3*B*a^3)/d*ln(sec(d*x+c)+tan(d*x+c))+a^3*B*sin(d*x+c)/d`

3.24.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.20

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4(A + 3B)a^3 dx \cos(dx + c)^2 + (7A + 6B)a^3 \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (7A + 6B)a^3 \cos(dx + c)}{4d \cos(dx+c)}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fracas")`

output $\frac{1}{4}(4*(A + 3*B)*a^3*d*x*\cos(d*x + c)^2 + (7*A + 6*B)*a^3*\cos(d*x + c)^2*\log(\sin(d*x + c) + 1) - (7*A + 6*B)*a^3*\cos(d*x + c)^2*\log(-\sin(d*x + c) + 1) + 2*(2*B*a^3*\cos(d*x + c)^2 + 2*(3*A + B)*a^3*\cos(d*x + c) + A*a^3)*\sin(d*x + c))/(d*\cos(d*x + c)^2)$

3.24.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= a^3 \left(\int A \sec^3(c + dx) dx + \int 3A \cos(c + dx) \sec^3(c + dx) dx \right. \\ & \quad + \int 3A \cos^2(c + dx) \sec^3(c + dx) dx + \int A \cos^3(c + dx) \sec^3(c + dx) dx \\ & \quad + \int B \cos(c + dx) \sec^3(c + dx) dx + \int 3B \cos^2(c + dx) \sec^3(c + dx) dx \\ & \quad \left. + \int 3B \cos^3(c + dx) \sec^3(c + dx) dx + \int B \cos^4(c + dx) \sec^3(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `a**3*(Integral(A*sec(c + d*x)**3, x) + Integral(3*A*cos(c + d*x)*sec(c + d*x)**3, x) + Integral(3*A*cos(c + d*x)**2*sec(c + d*x)**3, x) + Integral(A*cos(c + d*x)**3*sec(c + d*x)**3, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**3, x) + Integral(3*B*cos(c + d*x)**2*sec(c + d*x)**3, x) + Integral(3*B*cos(c + d*x)**3*sec(c + d*x)**3, x) + Integral(B*cos(c + d*x)**4*sec(c + d*x)**3, x))`

3.24.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.45

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= \frac{4(dx + c)Aa^3 + 12(dx + c)Ba^3 - Aa^3 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) + 6}{1} \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output $\frac{1}{4}*(4*(d*x + c)*A*a^3 + 12*(d*x + c)*B*a^3 - A*a^3*(2*\sin(d*x + c)/(\sin(d*x + c)^2 - 1) - \log(\sin(d*x + c) + 1) + \log(\sin(d*x + c) - 1)) + 6*A*a^3*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 6*B*a^3*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 4*B*a^3*\sin(d*x + c) + 12*A*a^3*\tan(d*x + c) + 4*B*a^3*\tan(d*x + c))/d$

3.24.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.68

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4Ba^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)}{\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + 1} + 2(Aa^3 + 3Ba^3)(dx + c) + (7Aa^3 + 6Ba^3) \log(|\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1|) - (7Aa^3 + 6Ba^3) \log(|\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1|)$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output $\frac{1}{2}*(4*B*a^3*\tan(1/2*d*x + 1/2*c)/(\tan(1/2*d*x + 1/2*c)^2 + 1) + 2*(A*a^3 + 3*B*a^3)*(d*x + c) + (7*A*a^3 + 6*B*a^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - (7*A*a^3 + 6*B*a^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(5*A*a^3*\tan(1/2*d*x + 1/2*c)^3 + 2*B*a^3*\tan(1/2*d*x + 1/2*c)^3 - 7*A*a^3*\tan(1/2*d*x + 1/2*c) - 2*B*a^3*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^2)/d$

3.24.9 Mupad [B] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.82

$$\begin{aligned}
& \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx \\
&= \frac{B a^3 \sin(c + dx)}{d} + \frac{2 A a^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{7 A a^3 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{6 B a^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{6 B a^3 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{3 A a^3 \sin(c + dx)}{d \cos(c + dx)} + \frac{A a^3 \sin(c + dx)}{2 d \cos(c + dx)^2} + \frac{B a^3 \sin(c + dx)}{d \cos(c + dx)}
\end{aligned}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^3,x)
```

```
output (B*a^3*sin(c + d*x))/d + (2*A*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (7*A*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (6*B*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (6*B*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (3*A*a^3*sin(c + d*x))/(d*cos(c + d*x)) + (A*a^3*sin(c + d*x))/(2*d*cos(c + d*x)^2) + (B*a^3*sin(c + d*x))/(d*cos(c + d*x))
```

3.25 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.25.1	Optimal result	413
3.25.2	Mathematica [A] (verified)	414
3.25.3	Rubi [A] (verified)	414
3.25.4	Maple [A] (verified)	418
3.25.5	Fricas [A] (verification not implemented)	419
3.25.6	Sympy [F(-1)]	419
3.25.7	Maxima [A] (verification not implemented)	419
3.25.8	Giac [A] (verification not implemented)	420
3.25.9	Mupad [B] (verification not implemented)	421

3.25.1 Optimal result

Integrand size = 31, antiderivative size = 125

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= a^3 B x + \frac{a^3 (5A + 7B) \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{5a^3 (A + B) \tan(c + dx)}{2d}$$

$$+ \frac{(5A + 3B) (a^3 + a^3 \cos(c + dx)) \sec(c + dx) \tan(c + dx)}{6d}$$

$$+ \frac{aA (a + a \cos(c + dx))^2 \sec^2(c + dx) \tan(c + dx)}{3d}$$

output

```
a^3*B*x+1/2*a^3*(5*A+7*B)*arctanh(sin(d*x+c))/d+5/2*a^3*(A+B)*tan(d*x+c)/d
+1/6*(5*A+3*B)*(a^3+a^3*cos(d*x+c))*sec(d*x+c)*tan(d*x+c)/d+1/3*a*A*(a+a*cos(d*x+c))^2*sec(d*x+c)^2*tan(d*x+c)/d
```

3.25.2 Mathematica [A] (verified)

Time = 1.10 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.56

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{a^3(6Bdx + 3(5A + 7B)\operatorname{arctanh}(\sin(c + dx)) + 3(8A + 6B + (3A + B)\sec(c + dx))\tan(c + dx) + 2A \tan(c + dx))}{6d}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a^3*(6*B*d*x + 3*(5*A + 7*B)*ArcTanh[Sin[c + d*x]] + 3*(8*A + 6*B + (3*A + B)*Sec[c + d*x])*Tan[c + d*x] + 2*A*Tan[c + d*x]^3)/(6*d)`

3.25.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3454, 3042, 3454, 27, 3042, 3447, 3042, 3500, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c + dx)(a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx$$

$$\downarrow \text{3454}$$

$$\frac{1}{3} \int (\cos(c + dx)a + a)^2 (a(5A + 3B) + 3aB \cos(c + dx)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a \cos(c + dx) + a)^2}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 (a(5A + 3B) + 3aB \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a \cos(c + dx) + a)^2}{3d}$$

3.25. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

↓ 3454

$$\frac{1}{3} \left(\frac{1}{2} \int 3(\cos(c+dx)a+a)(5(A+B)a^2+2B\cos(c+dx)a^2)\sec^2(c+dx)dx + \frac{(5A+3B)\tan(c+dx)\sec(c+dx)}{2d} \right. \\ \left. \frac{aA\tan(c+dx)\sec^2(c+dx)(a\cos(c+dx)+a)^2}{3d} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{3}{2} \int (\cos(c+dx)a+a)(5(A+B)a^2+2B\cos(c+dx)a^2)\sec^2(c+dx)dx + \frac{(5A+3B)\tan(c+dx)\sec(c+dx)}{2d} \right. \\ \left. \frac{aA\tan(c+dx)\sec^2(c+dx)(a\cos(c+dx)+a)^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(5(A+B)a^2+2B\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{(5A+3B)\tan(c+dx)\sec(c+dx)}{2d} \right. \\ \left. \frac{aA\tan(c+dx)\sec^2(c+dx)(a\cos(c+dx)+a)^2}{3d} \right)$$

↓ 3447

$$\frac{1}{3} \left(\frac{3}{2} \int (2B\cos^2(c+dx)a^3+5(A+B)a^3+(2Ba^3+5(A+B)a^3)\cos(c+dx))\sec^2(c+dx)dx + \frac{(5A+3B)\tan(c+dx)\sec(c+dx)}{2d} \right. \\ \left. \frac{aA\tan(c+dx)\sec^2(c+dx)(a\cos(c+dx)+a)^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{2B\sin(c+dx+\frac{\pi}{2})^2a^3+5(A+B)a^3+(2Ba^3+5(A+B)a^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{(5A+3B)\tan(c+dx)\sec(c+dx)}{2d} \right. \\ \left. \frac{aA\tan(c+dx)\sec^2(c+dx)(a\cos(c+dx)+a)^2}{3d} \right)$$

↓ 3500

$$\frac{1}{3} \left(\frac{3}{2} \left(\int ((5A+7B)a^3+2B\cos(c+dx)a^3)\sec(c+dx)dx + \frac{5a^3(A+B)\tan(c+dx)}{d} \right) + \frac{(5A+3B)\tan(c+dx)\sec(c+dx)}{2d} \right. \\ \left. \frac{aA\tan(c+dx)\sec^2(c+dx)(a\cos(c+dx)+a)^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \left(\int \frac{(5A + 7B)a^3 + 2B \sin(c + dx + \frac{\pi}{2}) a^3}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{5a^3(A + B) \tan(c + dx)}{d} \right) + \frac{(5A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^2}{3d}$$

↓ 3214

$$\frac{1}{3} \left(\frac{3}{2} \left(a^3(5A + 7B) \int \sec(c + dx) dx + \frac{5a^3(A + B) \tan(c + dx)}{d} + 2a^3 Bx \right) + \frac{(5A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^2}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \left(a^3(5A + 7B) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{5a^3(A + B) \tan(c + dx)}{d} + 2a^3 Bx \right) + \frac{(5A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^2}{3d}$$

↓ 4257

$$\frac{1}{3} \left(\frac{3}{2} \left(\frac{a^3(5A + 7B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{5a^3(A + B) \tan(c + dx)}{d} + 2a^3 Bx \right) + \frac{(5A + 3B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^2}{3d}$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*(a + a*Cos[c + d*x])^2*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + (((5*A + 3*B)*(a^3 + a^3*Cos[c + d*x])*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (3*(2*a^3*B*x + (a^3*(5*A + 7*B)*ArcTanh[Sin[c + d*x]]))/d + (5*a^3*(A + B)*Tan[c + d*x])/d))/2)/3`

3.25.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.25.4 Maple [A] (verified)

Time = 4.02 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.14

method	result
parts	$-\frac{A a^3 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d} + \frac{(A a^3 + 3B a^3) \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(3A a^3 + B a^3) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
parallelrisc	$3 \left(-\frac{5 \left(A + \frac{7B}{5} \right) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)}{2} + \frac{5 \left(A + \frac{7B}{5} \right) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{2} \right) + \frac{dx}{d(\cos(3dx+3c) + 3 \cos(dx+c))}$
derivativedivides	$\frac{A a^3 \ln(\sec(dx+c) + \tan(dx+c)) + B a^3 (dx+c) + 3A a^3 \tan(dx+c) + 3B a^3 \ln(\sec(dx+c) + \tan(dx+c)) + 3A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
default	$\frac{A a^3 \ln(\sec(dx+c) + \tan(dx+c)) + B a^3 (dx+c) + 3A a^3 \tan(dx+c) + 3B a^3 \ln(\sec(dx+c) + \tan(dx+c)) + 3A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
risc	$a^3 B x - \frac{ia^3 (9A e^{5i(dx+c)} + 3B e^{5i(dx+c)} - 18A e^{4i(dx+c)} - 18B e^{4i(dx+c)} - 48A e^{2i(dx+c)} - 36B e^{2i(dx+c)} - 9A e^{i(dx+c)} - 9B e^{i(dx+c)})}{3d(e^{2i(dx+c)} + 1)^3}$
norman	$\frac{a^3 B x \left(\tan^{12} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + a^3 B x \left(\tan^{14} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - a^3 B x - a^3 B x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 3a^3 B x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 3a^3 B x \left(\tan^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d}$

input `int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBOSE)`

output `-A*a^3/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(A*a^3+3*B*a^3)/d*ln(sec(d*x+c)+tan(d*x+c))+(3*A*a^3+B*a^3)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(3*A*a^3+3*B*a^3)/d*tan(d*x+c)+B*a^3/d*(d*x+c)`

3.25.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.13

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12 B a^3 dx \cos(dx + c)^3 + 3(5A + 7B)a^3 \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3(5A + 7B)a^3 \cos(dx + c)}{12 d c}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")`

output `1/12*(12*B*a^3*d*x*cos(d*x + c)^3 + 3*(5*A + 7*B)*a^3*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*(5*A + 7*B)*a^3*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(2*(11*A + 9*B)*a^3*cos(d*x + c)^2 + 3*(3*A + B)*a^3*cos(d*x + c) + 2*A*a^3)*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.25.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.25.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.70

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{4(\tan(dx + c)^3 + 3 \tan(dx + c))Aa^3 + 12(dx + c)Ba^3 - 9Aa^3 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right)}{12 d c}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output $\frac{1}{12}*(4*(\tan(dx + c)^3 + 3*\tan(dx + c))*A*a^3 + 12*(dx + c)*B*a^3 - 9*A*a^3*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) - 3*B*a^3*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) + 6*A*a^3*(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 18*B*a^3*(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 36*A*a^3*\tan(dx + c) + 36*B*a^3*\tan(dx + c))/d$

3.25.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.51

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{6(dx + c)Ba^3 + 3(5Aa^3 + 7Ba^3) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 3(5Aa^3 + 7Ba^3) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)\right|\right)}{d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")`

output $\frac{1}{6}*(6*(dx + c)*B*a^3 + 3*(5*A*a^3 + 7*B*a^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(5*A*a^3 + 7*B*a^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(15*A*a^3*\tan(1/2*d*x + 1/2*c)^5 + 15*B*a^3*\tan(1/2*d*x + 1/2*c)^5 - 40*A*a^3*\tan(1/2*d*x + 1/2*c)^3 - 36*B*a^3*\tan(1/2*d*x + 1/2*c)^3 + 33*A*a^3*\tan(1/2*d*x + 1/2*c) + 21*B*a^3*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^3/d$

3.25.9 Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.67

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{5 A a^3 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2 B a^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{7 B a^3 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{11 A a^3 \sin(c + dx)}{3 d \cos(c + dx)} + \frac{3 A a^3 \sin(c + dx)}{2 d \cos(c + dx)^2}$$

$$+ \frac{A a^3 \sin(c + dx)}{3 d \cos(c + dx)^3} + \frac{3 B a^3 \sin(c + dx)}{d \cos(c + dx)} + \frac{B a^3 \sin(c + dx)}{2 d \cos(c + dx)^2}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^4,x)`output `(5*A*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2))/d + (2*B*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (7*B*a^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (11*A*a^3*sin(c + d*x))/(3*d*cos(c + d*x)) + (3*A*a^3*sin(c + d*x))/(2*d*cos(c + d*x)^2) + (A*a^3*sin(c + d*x))/(3*d*cos(c + d*x)^3) + (3*B*a^3*sin(c + d*x))/(d*cos(c + d*x)) + (B*a^3*sin(c + d*x))/(2*d*cos(c + d*x)^2)`

3.26 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.26.1	Optimal result	422
3.26.2	Mathematica [A] (verified)	423
3.26.3	Rubi [A] (verified)	423
3.26.4	Maple [A] (verified)	427
3.26.5	Fricas [A] (verification not implemented)	428
3.26.6	Sympy [F(-1)]	428
3.26.7	Maxima [A] (verification not implemented)	429
3.26.8	Giac [A] (verification not implemented)	429
3.26.9	Mupad [B] (verification not implemented)	430

3.26.1 Optimal result

Integrand size = 31, antiderivative size = 154

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx \\ &= \frac{5a^3(3A + 4B)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{a^3(9A + 11B) \tan(c + dx)}{3d} \\ &+ \frac{a^3(27A + 28B) \sec(c + dx) \tan(c + dx)}{24d} \\ &+ \frac{(3A + 2B) (a^3 + a^3 \cos(c + dx)) \sec^2(c + dx) \tan(c + dx)}{6d} \\ &+ \frac{aA(a + a \cos(c + dx))^2 \sec^3(c + dx) \tan(c + dx)}{4d} \end{aligned}$$

```
output 5/8*a^3*(3*A+4*B)*arctanh(sin(d*x+c))/d+1/3*a^3*(9*A+11*B)*tan(d*x+c)/d+1/
24*a^3*(27*A+28*B)*sec(d*x+c)*tan(d*x+c)/d+1/6*(3*A+2*B)*(a^3+a^3*cos(d*x+
c))*sec(d*x+c)^2*tan(d*x+c)/d+1/4*a*A*(a+a*cos(d*x+c))^2*sec(d*x+c)^3*tan(
d*x+c)/d
```

3.26.2 Mathematica [A] (verified)

Time = 0.98 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.12

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{15a^3 A \operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{5a^3 B \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{4a^3 A \tan(c + dx)}{d}$$

$$+ \frac{4a^3 B \tan(c + dx)}{d} + \frac{15a^3 A \sec(c + dx) \tan(c + dx)}{8d} + \frac{3a^3 B \sec(c + dx) \tan(c + dx)}{2d}$$

$$+ \frac{a^3 A \sec^3(c + dx) \tan(c + dx)}{4d} + \frac{a^3 A \tan^3(c + dx)}{d} + \frac{a^3 B \tan^3(c + dx)}{3d}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output $(15*a^3*A*\operatorname{ArcTanh}[\operatorname{Sin}[c + d*x]])/(8*d) + (5*a^3*B*\operatorname{ArcTanh}[\operatorname{Sin}[c + d*x]])/(2*d) + (4*a^3*A*\operatorname{Tan}[c + d*x])/d + (4*a^3*B*\operatorname{Tan}[c + d*x])/d + (15*a^3*A*\operatorname{Sec}[c + d*x]*\operatorname{Tan}[c + d*x])/(8*d) + (3*a^3*B*\operatorname{Sec}[c + d*x]*\operatorname{Tan}[c + d*x])/(2*d) + (a^3*A*\operatorname{Sec}[c + d*x]^3*\operatorname{Tan}[c + d*x])/(4*d) + (a^3*A*\operatorname{Tan}[c + d*x]^3)/d + (a^3*B*\operatorname{Tan}[c + d*x]^3)/(3*d)$

3.26.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3454, 3042, 3454, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3454}$$

$$\frac{1}{4} \int (\cos(c + dx)a + a)^2 (2a(3A + 2B) + a(A + 4B) \cos(c + dx)) \sec^4(c + dx) dx + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d}$$

3.26. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

$$\begin{aligned} & \downarrow \text{3042} \\ & \frac{1}{4} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2 (2a(3A+2B) + a(A+4B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^4} dx + \\ & \quad \frac{aA \tan(c+dx) \sec^3(c+dx)(a \cos(c+dx) + a)^2}{4d} \\ & \downarrow \text{3454} \\ & \frac{1}{4} \left(\frac{1}{3} \int (\cos(c+dx)a+a) ((27A+28B)a^2 + (9A+16B)\cos(c+dx)a^2) \sec^3(c+dx) dx + \frac{2(3A+2B)\tan(c+dx)}{4d} \right. \\ & \quad \left. \frac{aA \tan(c+dx) \sec^3(c+dx)(a \cos(c+dx) + a)^2}{4d} \right) \\ & \downarrow \text{3042} \\ & \frac{1}{4} \left(\frac{1}{3} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a) ((27A+28B)a^2 + (9A+16B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^3} dx + \frac{2(3A+2B)\tan(c+dx)}{4d} \right. \\ & \quad \left. \frac{aA \tan(c+dx) \sec^3(c+dx)(a \cos(c+dx) + a)^2}{4d} \right) \\ & \downarrow \text{3447} \\ & \frac{1}{4} \left(\frac{1}{3} \int ((9A+16B)\cos^2(c+dx)a^3 + (27A+28B)a^3 + ((9A+16B)a^3 + (27A+28B)a^3)\cos(c+dx)) \sec^3(c+dx) dx + \right. \\ & \quad \left. \frac{aA \tan(c+dx) \sec^3(c+dx)(a \cos(c+dx) + a)^2}{4d} \right) \\ & \downarrow \text{3042} \\ & \frac{1}{4} \left(\frac{1}{3} \int \frac{(9A+16B)\sin(c+dx+\frac{\pi}{2})^2 a^3 + (27A+28B)a^3 + ((9A+16B)a^3 + (27A+28B)a^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3} dx + \right. \\ & \quad \left. \frac{aA \tan(c+dx) \sec^3(c+dx)(a \cos(c+dx) + a)^2}{4d} \right) \\ & \downarrow \text{3500} \\ & \frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (8(9A+11B)a^3 + 15(3A+4B)\cos(c+dx)a^3) \sec^2(c+dx) dx + \frac{a^3(27A+28B)\tan(c+dx)\sec(c+dx)}{2d} \right) \right. \\ & \quad \left. \frac{aA \tan(c+dx) \sec^3(c+dx)(a \cos(c+dx) + a)^2}{4d} \right) \\ & \downarrow \text{3042} \end{aligned}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{8(9A + 11B)a^3 + 15(3A + 4B) \sin(c + dx + \frac{\pi}{2}) a^3}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{a^3(27A + 28B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d} \right) \downarrow \text{3227}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(8a^3(9A + 11B) \int \sec^2(c + dx) dx + 15a^3(3A + 4B) \int \sec(c + dx) dx \right) + \frac{a^3(27A + 28B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d} \right) \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15a^3(3A + 4B) \int \csc(c + dx + \frac{\pi}{2}) dx + 8a^3(9A + 11B) \int \csc(c + dx + \frac{\pi}{2})^2 dx \right) + \frac{a^3(27A + 28B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d} \right) \downarrow \text{4254}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15a^3(3A + 4B) \int \csc(c + dx + \frac{\pi}{2}) dx - \frac{8a^3(9A + 11B) \int 1d(-\tan(c + dx))}{d} \right) + \frac{a^3(27A + 28B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d} \right) \downarrow \text{24}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15a^3(3A + 4B) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{8a^3(9A + 11B) \tan(c + dx)}{d} \right) + \frac{a^3(27A + 28B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d} \right) \downarrow \text{4257}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\frac{15a^3(3A + 4B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{8a^3(9A + 11B) \tan(c + dx)}{d} \right) + \frac{a^3(27A + 28B) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)(a \cos(c + dx) + a)^2}{4d} \right)$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output $(aA*(a + a*\cos[c + d*x])^2*\sec[c + d*x]^3*\tan[c + d*x])/(4*d) + ((2*(3*A + 2*B)*(a^3 + a^3*\cos[c + d*x])*sec[c + d*x]^2*\tan[c + d*x])/(3*d) + ((a^3*(27*A + 28*B)*sec[c + d*x]*\tan[c + d*x])/(2*d) + ((15*a^3*(3*A + 4*B)*ArcTanh[\sin[c + d*x]])/d + (8*a^3*(9*A + 11*B)*\tan[c + d*x])/d)/2)/3)/4$

3.26.3.1 Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3227 $\text{Int}[(b_*\sin[e_*] + f_*(x_*))^m * (c_* + d_*\sin[e_*] + f_*(x_*))], x_Symbol] \rightarrow \text{Simp}[c \text{ Int}[(b*\sin[e + f*x])^m, x], x] + \text{Simp}[d/b \text{ Int}[(b*\sin[e + f*x])^{m+1}, x], x] \text{ ; FreeQ}[\{b, c, d, e, f, m\}, x]$

rule 3447 $\text{Int}[(a_* + b_*\sin[e_*] + f_*(x_*))^m * (A_* + B_*\sin[e_*] + f_*(x_*)) * (c_* + d_*\sin[e_*] + f_*(x_*))], x_Symbol] \rightarrow \text{Int}[(a + b*\sin[e + f*x])^m * (A*c + (B*c + A*d)*\sin[e + f*x] + B*d*\sin[e + f*x]^2), x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0]$

rule 3454 $\text{Int}[(a_* + b_*\sin[e_*] + f_*(x_*))^m * (A_* + B_*\sin[e_*] + f_*(x_*))^n * (c_* + d_*\sin[e_*] + f_*(x_*))^n], x_Symbol] \rightarrow \text{Simp}[(-b^2)*(B*c - A*d)*\cos[e + f*x]*(a + b*\sin[e + f*x])^{m-1} * ((c + d*\sin[e + f*x])^{n+1}/(d*f*(n+1)*(b*c + a*d))), x] - \text{Simp}[b/(d*(n+1)*(b*c + a*d)) \text{ Int}[(a + b*\sin[e + f*x])^{m-1} * (c + d*\sin[e + f*x])^{n+1} * \text{Simp}[a*A*d*(m-n-2) - B*(a*c*(m-1) + b*d*(n+1)) - (A*b*d*(m+n+1) - B*(b*c*m - a*d*(n+1)))*\sin[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{GtQ}[m, 1/2] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*m] \ \&\& \ (\text{IntegerQ}[2*n] \ || \ \text{EqQ}[c, 0])]$

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-A*b^2
- a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.26.4 Maple [A] (verified)

Time = 4.71 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.16

method	result
parallelrisch	$10 \left(-\frac{3(A + \frac{4B}{3}) \left(\frac{3}{4} + \frac{\cos(4dx+4c) + \cos(2dx+2c)}{4} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 3 \left(\frac{3}{4} + \frac{\cos(4dx+4c) + \cos(2dx+2c)}{4} \right) \left(A + \frac{4B}{3} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{d(\cos(4dx+4c)+4c)} \right)$
parts	$A a^3 \left(-\left(-\frac{(\sec^3(dx+c))}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c)+\tan(dx+c))}{8} \right) + \frac{(A a^3 + 3B a^3) \tan(dx+c)}{d} - \frac{(3A a^3 + 3B a^3) \ln(\sec(dx+c)+\tan(dx+c))}{d}$
derivativedivides	$A a^3 \tan(dx+c) + B a^3 \ln(\sec(dx+c)+\tan(dx+c)) + 3A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) + 3B a^3 \tan(dx+c)$
default	$A a^3 \tan(dx+c) + B a^3 \ln(\sec(dx+c)+\tan(dx+c)) + 3A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) + 3B a^3 \tan(dx+c)$
risch	$-\frac{ia^3(45A e^{7i(dx+c)} + 36B e^{7i(dx+c)} - 24A e^{6i(dx+c)} - 72B e^{6i(dx+c)} + 69A e^{5i(dx+c)} + 36B e^{5i(dx+c)} - 216A e^{4i(dx+c)} - 216B e^{4i(dx+c)} + 12d(e^{2i(dx+c)} + e^{-2i(dx+c)}))}{12d(e^{2i(dx+c)} + e^{-2i(dx+c)})}$
norman	$\frac{19a^3(3A+4B) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 5a^3(3A+4B) \left(\tan^{13} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 5a^3(3A+4B) \left(\tan^{15} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + a^3(49A+44B) \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + a^3(49A+44B) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{(1+\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right))}$

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBO
SE)
```

3.26. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

output $10*(-3/4*(A+4/3*B)*(3/4+1/4*\cos(4*d*x+4*c)+\cos(2*d*x+2*c))*\ln(\tan(1/2*d*x+1/2*c)-1)+3/4*(3/4+1/4*\cos(4*d*x+4*c)+\cos(2*d*x+2*c))*(A+4/3*B)*\ln(\tan(1/2*d*x+1/2*c)+1)+(A+13/15*B)*\sin(2*d*x+2*c)+(3/8*A+3/10*B)*\sin(3*d*x+3*c)+(3/10*A+11/30*B)*\sin(4*d*x+4*c)+23/40*(A+12/23*B)*\sin(d*x+c))*a^3/d/(\cos(4*d*x+4*c)+4*\cos(2*d*x+2*c)+3)$

3.26.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.94

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{15(3A + 4B)a^3 \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 15(3A + 4B)a^3 \cos(dx + c)^4 \log(-\sin(dx + c) + 1)}{4}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")`

output $1/48*(15*(3*A + 4*B)*a^3*\cos(d*x + c)^4*\log(\sin(d*x + c) + 1) - 15*(3*A + 4*B)*a^3*\cos(d*x + c)^4*\log(-\sin(d*x + c) + 1) + 2*(8*(9*A + 11*B)*a^3*\cos(d*x + c)^3 + 9*(5*A + 4*B)*a^3*\cos(d*x + c)^2 + 8*(3*A + B)*a^3*\cos(d*x + c) + 6*A*a^3)*\sin(d*x + c))/(d*\cos(d*x + c)^4)$

3.26.6 SymPy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output Timed out

3.26.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.75

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{48 (\tan(dx + c)^3 + 3 \tan(dx + c)) A a^3 + 16 (\tan(dx + c)^3 + 3 \tan(dx + c)) B a^3 - 3 A a^3 \left(\frac{2(3 \sin(dx+c)^3 - 5 \sin(dx+c))}{\sin(dx+c)^4 - 2 \sin(dx+c)^2 + 1} - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1) \right) - 36 A a^3 (2 \sin(dx+c) / (\sin(dx+c)^2 - 1) - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1)) - 36 B a^3 (2 \sin(dx+c) / (\sin(dx+c)^2 - 1) - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1)) + 24 B a^3 (\log(\sin(dx+c) + 1) - \log(\sin(dx+c) - 1)) + 48 A a^3 \tan(dx+c) + 144 B a^3 \tan(dx+c)) / d}{}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output `1/48*(48*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^3 + 16*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^3 - 3*A*a^3*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 36*A*a^3*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 36*B*a^3*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 24*B*a^3*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 48*A*a^3*tan(d*x + c) + 144*B*a^3*tan(d*x + c))/d`

3.26.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.38

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{15 (3 A a^3 + 4 B a^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 15 (3 A a^3 + 4 B a^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2(45 A a^3 \tan(1/2 dx + 1/2 c)^7 + 60 B a^3 \tan(1/2 dx + 1/2 c)^7 - 165 A a^3 \tan(1/2 dx + 1/2 c)^5 - 220 B a^3 \tan(1/2 dx + 1/2 c)^5 + 219 A a^3 \tan(1/2 dx + 1/2 c)^3 + 292 B a^3 \tan(1/2 dx + 1/2 c)^3 - 147 A a^3 \tan(1/2 dx + 1/2 c) - 132 B a^3 \tan(1/2 dx + 1/2 c))}{(\tan(1/2 dx + 1/2 c)^2 - 1)^4} / d}{}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")`

output `1/24*(15*(3*A*a^3 + 4*B*a^3)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 15*(3*A*a^3 + 4*B*a^3)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(45*A*a^3*tan(1/2*d*x + 1/2*c)^7 + 60*B*a^3*tan(1/2*d*x + 1/2*c)^7 - 165*A*a^3*tan(1/2*d*x + 1/2*c)^5 - 220*B*a^3*tan(1/2*d*x + 1/2*c)^5 + 219*A*a^3*tan(1/2*d*x + 1/2*c)^3 + 292*B*a^3*tan(1/2*d*x + 1/2*c)^3 - 147*A*a^3*tan(1/2*d*x + 1/2*c) - 132*B*a^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^4/d`

3.26. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.26.9 Mupad [B] (verification not implemented)

Time = 2.90 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.20

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{\left(-\frac{15Aa^3}{4} - 5Ba^3\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{55Aa^3}{4} + \frac{55Ba^3}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{73Aa^3}{4} - \frac{73Ba^3}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)} + \frac{5a^3 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (3A + 4B)}{4d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^5,x)`output `(tan(c/2 + (d*x)/2)*((49*A*a^3)/4 + 11*B*a^3) - tan(c/2 + (d*x)/2)^7*((15*A*a^3)/4 + 5*B*a^3) + tan(c/2 + (d*x)/2)^5*((55*A*a^3)/4 + (55*B*a^3)/3) - tan(c/2 + (d*x)/2)^3*((73*A*a^3)/4 + (73*B*a^3)/3))/(d*(6*tan(c/2 + (d*x)/2)^4 - 4*tan(c/2 + (d*x)/2)^2 - 4*tan(c/2 + (d*x)/2)^6 + tan(c/2 + (d*x)/2)^8 + 1)) + (5*a^3*atanh(tan(c/2 + (d*x)/2))*(3*A + 4*B))/(4*d)`

3.27 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^6(c+dx) dx$

3.27.1	Optimal result	431
3.27.2	Mathematica [A] (verified)	432
3.27.3	Rubi [A] (verified)	432
3.27.4	Maple [A] (verified)	437
3.27.5	Fricas [A] (verification not implemented)	437
3.27.6	Sympy [F(-1)]	438
3.27.7	Maxima [A] (verification not implemented)	438
3.27.8	Giac [A] (verification not implemented)	439
3.27.9	Mupad [B] (verification not implemented)	439

3.27.1 Optimal result

Integrand size = 31, antiderivative size = 185

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{a^3(13A + 15B)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{a^3(38A + 45B) \tan(c + dx)}{15d}$$

$$+ \frac{a^3(13A + 15B) \sec(c + dx) \tan(c + dx)}{8d} + \frac{a^3(43A + 45B) \sec^2(c + dx) \tan(c + dx)}{60d}$$

$$+ \frac{(7A + 5B) (a^3 + a^3 \cos(c + dx)) \sec^3(c + dx) \tan(c + dx)}{20d}$$

$$+ \frac{aA(a + a \cos(c + dx))^2 \sec^4(c + dx) \tan(c + dx)}{5d}$$

output

```
1/8*a^3*(13*A+15*B)*arctanh(sin(d*x+c))/d+1/15*a^3*(38*A+45*B)*tan(d*x+c)/
d+1/8*a^3*(13*A+15*B)*sec(d*x+c)*tan(d*x+c)/d+1/60*a^3*(43*A+45*B)*sec(d*x
+c)^2*tan(d*x+c)/d+1/20*(7*A+5*B)*(a^3+a^3*cos(d*x+c))*sec(d*x+c)^3*tan(d*
x+c)/d+1/5*a*A*(a+a*cos(d*x+c))^2*sec(d*x+c)^4*tan(d*x+c)/d
```

3.27.2 Mathematica [A] (verified)

Time = 1.76 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.55

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{a^3(15(13A + 15B)\operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx)(15(13A + 15B)\sec(c + dx) + 30(3A + B)\sec^3(c + dx))}{120d}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a^3*(15*(13*A + 15*B)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(15*(13*A + 15*B)*Sec[c + d*x] + 30*(3*A + B)*Sec[c + d*x]^3 + 8*(60*(A + B) + 5*(5*A + 3*B)*Tan[c + d*x]^2 + 3*A*Tan[c + d*x]^4)))/(120*d)`

3.27.3 Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.03, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3454, 3042, 3454, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^6(c + dx)(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^6} dx$$

$$\downarrow \text{3454}$$

$$\frac{1}{5} \int (\cos(c + dx)a + a)^2 (a(7A + 5B) + a(2A + 5B)\cos(c + dx)) \sec^5(c + dx) dx + \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{5} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 (a(7A + 5B) + a(2A + 5B)\sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx + \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d}$$

↓ 3454

$$\frac{1}{5} \left(\frac{1}{4} \int (\cos(c + dx)a + a) ((43A + 45B)a^2 + 2(11A + 15B)\cos(c + dx)a^2) \sec^4(c + dx) dx + \frac{(7A + 5B)\tan(c + dx)}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a) ((43A + 45B)a^2 + 2(11A + 15B)\sin(c + dx + \frac{\pi}{2})a^2)}{\sin(c + dx + \frac{\pi}{2})^4} dx + \frac{(7A + 5B)\tan(c + dx)}{5d} \right)$$

↓ 3447

$$\frac{1}{5} \left(\frac{1}{4} \int (2(11A + 15B)\cos^2(c + dx)a^3 + (43A + 45B)a^3 + (2(11A + 15B)a^3 + (43A + 45B)a^3)\cos(c + dx)) \sec^4(c + dx) dx + \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \frac{2(11A + 15B)\sin(c + dx + \frac{\pi}{2})^2 a^3 + (43A + 45B)a^3 + (2(11A + 15B)a^3 + (43A + 45B)a^3)\sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^4} dx + \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \right)$$

↓ 3500

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int (15(13A + 15B)a^3 + 4(38A + 45B)\cos(c + dx)a^3) \sec^3(c + dx) dx + \frac{a^3(43A + 45B)\tan(c + dx)\sec^2(c + dx)}{3d} \right) + \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \frac{15(13A + 15B)a^3 + 4(38A + 45B) \sin(c + dx + \frac{\pi}{2}) a^3}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{a^3(43A + 45B) \tan(c + dx) \sec^2(c + dx)}{3d} \right) \right) \\ \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \\ \downarrow \text{3227}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15a^3(13A + 15B) \int \sec^3(c + dx) dx + 4a^3(38A + 45B) \int \sec^2(c + dx) dx \right) \right) + \frac{a^3(43A + 45B) \tan(c + dx)}{3d} \right) \\ \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \\ \downarrow \text{3042}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(4a^3(38A + 45B) \int \csc(c + dx + \frac{\pi}{2})^2 dx + 15a^3(13A + 15B) \int \csc(c + dx + \frac{\pi}{2})^3 dx \right) \right) + \frac{a^3(43A + 45B)}{3d} \right) \\ \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \\ \downarrow \text{4254}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15a^3(13A + 15B) \int \csc(c + dx + \frac{\pi}{2})^3 dx - \frac{4a^3(38A + 45B) \int 1d(-\tan(c + dx))}{d} \right) \right) + \frac{a^3(43A + 45B)}{3d} \right) \\ \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \\ \downarrow \text{24}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15a^3(13A + 15B) \int \csc(c + dx + \frac{\pi}{2})^3 dx + \frac{4a^3(38A + 45B) \tan(c + dx)}{d} \right) \right) + \frac{a^3(43A + 45B) \tan(c + dx)}{3d} \right) \\ \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \\ \downarrow \text{4255}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15a^3(13A + 15B) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) \right) + \frac{4a^3(38A + 45B) \tan(c + dx)}{d} \right) \right) + \\ \frac{aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^2}{5d} \\ \downarrow \text{3042}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15a^3(13A + 15B) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) \right) + \frac{4a^3(38A + 45B) \tan(c + dx)}{d} \right) \right) + \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^2}{5d}$$

↓ 4257

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15a^3(13A + 15B) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) \right) + \frac{4a^3(38A + 45B) \tan(c + dx)}{d} \right) \right) + \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^2}{5d}$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a*A*(a + a*Cos[c + d*x])^2*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) + (((7*A + 5*B)*(a^3 + a^3*Cos[c + d*x])*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((a^3*(43*A + 45*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((4*a^3*(38*A + 45*B)*Tan[c + d*x])/d + 15*a^3*(13*A + 15*B)*(ArcTanh[Sin[c + d*x]])/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/3)/4)/5`

3.27.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3500 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.27.4 Maple [A] (verified)

Time = 5.86 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.06

method	result
parts	$-\frac{A a^3 \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c}}{d} + \frac{(A a^3 + 3B a^3) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d}$
parallelrisch	$40 \left(-\frac{39 \left(\frac{\cos(5dx+5c)}{10} + \frac{\cos(3dx+3c)}{2} + \cos(dx+c) \right) \left(A + \frac{15B}{13} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)}{32} + \frac{39 \left(\frac{\cos(5dx+5c)}{10} + \frac{\cos(3dx+3c)}{2} + \cos(dx+c) \right)}{32} \right)$
derivativedivides	$A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + B a^3 \tan(dx+c) - 3A a^3 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + 3B a^3$
default	$A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + B a^3 \tan(dx+c) - 3A a^3 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + 3B a^3$
risch	$-\frac{i a^3 (195 A e^{9i(dx+c)} + 225 B e^{9i(dx+c)} - 120 B e^{8i(dx+c)} + 750 A e^{7i(dx+c)} + 570 B e^{7i(dx+c)} - 720 A e^{6i(dx+c)} - 1200 B e^{6i(dx+c)} - 1200 A e^{5i(dx+c)} - 1200 B e^{5i(dx+c)} - 1200 A e^{4i(dx+c)} - 1200 B e^{4i(dx+c)} - 1200 A e^{3i(dx+c)} - 1200 B e^{3i(dx+c)} - 1200 A e^{2i(dx+c)} - 1200 B e^{2i(dx+c)} - 1200 A e^{i(dx+c)} - 1200 B e^{i(dx+c)})}{32}$

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x,method=_RETURNVERBOSE)
```

```
output -A*a^3/d*(-8/15-1/5*sec(d*x+c)^4-4/15*sec(d*x+c)^2)*tan(d*x+c)+(A*a^3+3*B*a^3)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(3*A*a^3+B*a^3)/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))-(3*A*a^3+3*B*a^3)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+B*a^3/d*tan(d*x+c)
```

3.27.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.89

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{15 (13 A + 15 B) a^3 \cos(dx + c)^5 \log(\sin(dx + c) + 1) - 15 (13 A + 15 B) a^3 \cos(dx + c)^5 \log(-\sin(dx + c) + 1)}{1}$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="fracas")
```

output $1/240*(15*(13*A + 15*B)*a^3*\cos(d*x + c)^5*\log(\sin(d*x + c) + 1) - 15*(13*A + 15*B)*a^3*\cos(d*x + c)^5*\log(-\sin(d*x + c) + 1) + 2*(8*(38*A + 45*B)*a^3*\cos(d*x + c)^4 + 15*(13*A + 15*B)*a^3*\cos(d*x + c)^3 + 8*(19*A + 15*B)*a^3*\cos(d*x + c)^2 + 30*(3*A + B)*a^3*\cos(d*x + c) + 24*A*a^3)*\sin(d*x + c))/(d*\cos(d*x + c)^5)$

3.27.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**6,x)`

output Timed out

3.27.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.82

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{16 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c)) A a^3 + 240 (\tan(dx + c)^3 + 3 \tan(dx + c)) A a^3}{d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="maxima")`

output $1/240*(16*(3*\tan(d*x + c)^5 + 10*\tan(d*x + c)^3 + 15*\tan(d*x + c))*A*a^3 + 240*(\tan(d*x + c)^3 + 3*\tan(d*x + c))*A*a^3 + 240*(\tan(d*x + c)^3 + 3*\tan(d*x + c))*B*a^3 - 45*A*a^3*(2*(3*\sin(d*x + c)^3 - 5*\sin(d*x + c)))/(\sin(d*x + c)^4 - 2*\sin(d*x + c)^2 + 1) - 3*\log(\sin(d*x + c) + 1) + 3*\log(\sin(d*x + c) - 1)) - 15*B*a^3*(2*(3*\sin(d*x + c)^3 - 5*\sin(d*x + c)))/(\sin(d*x + c)^4 - 2*\sin(d*x + c)^2 + 1) - 3*\log(\sin(d*x + c) + 1) + 3*\log(\sin(d*x + c) - 1)) - 60*A*a^3*(2*\sin(d*x + c))/(\sin(d*x + c)^2 - 1) - \log(\sin(d*x + c) + 1) + \log(\sin(d*x + c) - 1)) - 180*B*a^3*(2*\sin(d*x + c))/(\sin(d*x + c)^2 - 1) - \log(\sin(d*x + c) + 1) + \log(\sin(d*x + c) - 1)) + 240*B*a^3*\tan(d*x + c))/d$

3.27. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$

3.27.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.33

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{15(13Aa^3 + 15Ba^3) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 15(13Aa^3 + 15Ba^3) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) - \frac{2}{d} \left(\frac{13Aa^3 + 15Ba^3}{4} \tan^9\left(\frac{c}{2} + \frac{dx}{2}\right) + \left(-\frac{91Aa^3}{6} - \frac{35Ba^3}{2}\right) \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right) + \left(\frac{416Aa^3}{15} + 32Ba^3\right) \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right) - \left(\frac{13Aa^3}{4} + \frac{15Ba^3}{4}\right) \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right) + 735Ba^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \right)}{d \left(\tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 1 \right)^5}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="giac")`

output `1/120*(15*(13*A*a^3 + 15*B*a^3)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 15*(13*A*a^3 + 15*B*a^3)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(195*A*a^3*tan(1/2*d*x + 1/2*c)^9 + 225*B*a^3*tan(1/2*d*x + 1/2*c)^9 - 910*A*a^3*tan(1/2*d*x + 1/2*c)^7 - 1050*B*a^3*tan(1/2*d*x + 1/2*c)^7 + 1664*A*a^3*tan(1/2*d*x + 1/2*c)^5 + 1920*B*a^3*tan(1/2*d*x + 1/2*c)^5 - 1330*A*a^3*tan(1/2*d*x + 1/2*c)^3 - 1830*B*a^3*tan(1/2*d*x + 1/2*c)^3 + 765*A*a^3*tan(1/2*d*x + 1/2*c) + 735*B*a^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^5/d`

3.27.9 Mupad [B] (verification not implemented)

Time = 2.96 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.21

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{a^3 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (13A + 15B)}{4d} - \frac{\left(\frac{13Aa^3}{4} + \frac{15Ba^3}{4}\right) \tan^9\left(\frac{c}{2} + \frac{dx}{2}\right) + \left(-\frac{91Aa^3}{6} - \frac{35Ba^3}{2}\right) \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right) + \left(\frac{416Aa^3}{15} + 32Ba^3\right) \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right) - \left(\frac{13Aa^3}{4} + \frac{15Ba^3}{4}\right) \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right) + 735Ba^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) - 1 \right)^5}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^6,x)`

output $(a^3 \operatorname{atanh}(\tan(c/2 + (d*x)/2)) * (13*A + 15*B)) / (4*d) - (\tan(c/2 + (d*x)/2) * ((51*A*a^3)/4 + (49*B*a^3)/4) + \tan(c/2 + (d*x)/2)^9 * ((13*A*a^3)/4 + (15*B*a^3)/4) - \tan(c/2 + (d*x)/2)^7 * ((91*A*a^3)/6 + (35*B*a^3)/2) - \tan(c/2 + (d*x)/2)^3 * ((133*A*a^3)/6 + (61*B*a^3)/2) + \tan(c/2 + (d*x)/2)^5 * ((416*A*a^3)/15 + 32*B*a^3)) / (d * (5 * \tan(c/2 + (d*x)/2)^2 - 10 * \tan(c/2 + (d*x)/2)^4 + 10 * \tan(c/2 + (d*x)/2)^6 - 5 * \tan(c/2 + (d*x)/2)^8 + \tan(c/2 + (d*x)/2)^{10} - 1))$

3.28 $\int \cos^2(c+dx)(a+a \cos(c+dx))^4(A+B \cos(c+dx)) dx$

3.28.1	Optimal result	441
3.28.2	Mathematica [A] (verified)	442
3.28.3	Rubi [A] (verified)	442
3.28.4	Maple [A] (verified)	447
3.28.5	Fricas [A] (verification not implemented)	448
3.28.6	Sympy [B] (verification not implemented)	448
3.28.7	Maxima [A] (verification not implemented)	449
3.28.8	Giac [A] (verification not implemented)	450
3.28.9	Mupad [B] (verification not implemented)	451

3.28.1 Optimal result

Integrand size = 31, antiderivative size = 241

$$\int \cos^2(c+dx)(a+a \cos(c+dx))^4(A+B \cos(c+dx)) dx$$

$$= \frac{1}{16}a^4(49A+44B)x + \frac{a^4(252A+227B) \sin(c+dx)}{35d}$$

$$+ \frac{a^4(49A+44B) \cos(c+dx) \sin(c+dx)}{16d} + \frac{a^4(301A+276B) \cos^3(c+dx) \sin(c+dx)}{280d}$$

$$+ \frac{aB \cos^3(c+dx)(a+a \cos(c+dx))^3 \sin(c+dx)}{7d}$$

$$+ \frac{(7A+10B) \cos^3(c+dx) (a^2+a^2 \cos(c+dx))^2 \sin(c+dx)}{42d}$$

$$+ \frac{7(A+B) \cos^3(c+dx) (a^4+a^4 \cos(c+dx)) \sin(c+dx)}{15d}$$

$$- \frac{a^4(252A+227B) \sin^3(c+dx)}{105d}$$

```
output 1/16*a^4*(49*A+44*B)*x+1/35*a^4*(252*A+227*B)*sin(d*x+c)/d+1/16*a^4*(49*A+
44*B)*cos(d*x+c)*sin(d*x+c)/d+1/280*a^4*(301*A+276*B)*cos(d*x+c)^3*sin(d*x
+c)/d+1/7*a*B*cos(d*x+c)^3*(a+a*cos(d*x+c))^3*sin(d*x+c)/d+1/42*(7*A+10*B)
*cos(d*x+c)^3*(a^2+a^2*cos(d*x+c))^2*sin(d*x+c)/d+7/15*(A+B)*cos(d*x+c)^3*
(a^4+a^4*cos(d*x+c))*sin(d*x+c)/d-1/105*a^4*(252*A+227*B)*sin(d*x+c)^3/d
```


3.28.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.65

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{a^4(18480Bc + 20580Adx + 18480Bdx + 105(352A + 323B) \sin(c + dx) + 105(127A + 124B) \sin(2(c + dx) + \dots)}{720d}$$

input `Integrate[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output $(a^4(18480Bc + 20580A*d*x + 18480B*d*x + 105(352A + 323B)*\text{Sin}[c + d*x] + 105(127A + 124B)*\text{Sin}[2(c + d*x)] + 5040A*\text{Sin}[3(c + d*x)] + 5495B*\text{Sin}[3(c + d*x)] + 1575A*\text{Sin}[4(c + d*x)] + 2100B*\text{Sin}[4(c + d*x)] + 336A*\text{Sin}[5(c + d*x)] + 651B*\text{Sin}[5(c + d*x)] + 35A*\text{Sin}[6(c + d*x)] + 140B*\text{Sin}[6(c + d*x)] + 15B*\text{Sin}[7(c + d*x)]))/(6720*d)$

3.28.3 Rubi [A] (verified)

Time = 1.58 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 3455, 3042, 3455, 3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx)(a \cos(c + dx) + a)^4(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^4 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3455}$$

$$\frac{1}{7} \int \cos^2(c + dx)(\cos(c + dx)a + a)^3(a(7A + 3B) + a(7A + 10B) \cos(c + dx)) dx + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{7} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right)^3 \left(a(7A + 3B) + a(7A + 10B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d}$$

↓ 3455

$$\frac{1}{7} \left(\frac{1}{6} \int \cos^2(c + dx)(\cos(c + dx)a + a)^2 (3(21A + 16B)a^2 + 98(A + B) \cos(c + dx)a^2) dx + \frac{(7A + 10B) \sin(c + dx) a^2}{7d} \right) + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{6} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right)^2 \left(3(21A + 16B)a^2 + 98(A + B) \sin \left(c + dx + \frac{\pi}{2} \right) a^2 \right) dx + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right)$$

↓ 3455

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \int 3 \cos^2(c + dx)(\cos(c + dx)a + a) ((203A + 178B)a^3 + (301A + 276B) \cos(c + dx)a^3) dx + \frac{98(A + B) \sin(c + dx) a^2}{7d} \right) + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \int \cos^2(c + dx)(\cos(c + dx)a + a) ((203A + 178B)a^3 + (301A + 276B) \cos(c + dx)a^3) dx + \frac{98(A + B) \sin(c + dx) a^2}{7d} \right) + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\sin \left(c + dx + \frac{\pi}{2} \right) a + a \right) \left((203A + 178B)a^3 + (301A + 276B) \sin \left(c + dx + \frac{\pi}{2} \right) a^3 \right) dx + \frac{98(A + B) \cos(c + dx) a^2}{7d} \right) + \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right)$$

↓ 3447

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \int \cos^2(c+dx) \left((301A+276B) \cos^2(c+dx)a^4 + (203A+178B)a^4 + ((203A+178B)a^4 + (301A+276B)a^4 \right) \right. \right. \\ \left. \left. \frac{aB \sin(c+dx) \cos^3(c+dx)(a \cos(c+dx) + a)^3}{7d} \right) \right) \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \int \sin \left(c+dx + \frac{\pi}{2} \right)^2 \left((301A+276B) \sin \left(c+dx + \frac{\pi}{2} \right)^2 a^4 + (203A+178B)a^4 + ((203A+178B)a^4 + (301A+276B)a^4 \right) \right. \right. \\ \left. \left. \frac{aB \sin(c+dx) \cos^3(c+dx)(a \cos(c+dx) + a)^3}{7d} \right) \right) \\ \downarrow \text{3502}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \int \cos^2(c+dx) (35(49A+44B)a^4 + 8(252A+227B) \cos(c+dx)a^4) dx + \frac{a^4(301A+276B) \sin(c+dx)}{4d} \right) \right. \right. \\ \left. \left. \frac{aB \sin(c+dx) \cos^3(c+dx)(a \cos(c+dx) + a)^3}{7d} \right) \right) \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \int \sin \left(c+dx + \frac{\pi}{2} \right)^2 (35(49A+44B)a^4 + 8(252A+227B) \sin \left(c+dx + \frac{\pi}{2} \right) a^4) dx + \frac{a^4(301A+276B) \cos(c+dx)}{4d} \right) \right. \right. \\ \left. \left. \frac{aB \sin(c+dx) \cos^3(c+dx)(a \cos(c+dx) + a)^3}{7d} \right) \right) \\ \downarrow \text{3227}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(8a^4(252A+227B) \int \cos^3(c+dx) dx + 35a^4(49A+44B) \int \cos^2(c+dx) dx \right) \right) \right. \right. \\ \left. \left. \frac{aB \sin(c+dx) \cos^3(c+dx)(a \cos(c+dx) + a)^3}{7d} \right) \right) \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(35a^4(49A+44B) \int \sin \left(c+dx + \frac{\pi}{2} \right)^2 dx + 8a^4(252A+227B) \int \sin \left(c+dx + \frac{\pi}{2} \right)^3 dx \right) \right) \right. \right. \\ \left. \left. \frac{aB \sin(c+dx) \cos^3(c+dx)(a \cos(c+dx) + a)^3}{7d} \right) \right) \\ \downarrow \text{3113}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(35a^4(49A + 44B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{8a^4(252A + 227B) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right. \right. \right. \right. \\ \left. \left. \left. \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right) \right) \right) \downarrow \text{2009}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(35a^4(49A + 44B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{8a^4(252A + 227B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) \right) \right) \right) + \\ \left. \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right) \downarrow \text{3115}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{1}{4} \left(35a^4(49A + 44B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{8a^4(252A + 227B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right. \right. \right. \right. \\ \left. \left. \left. \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right) \right) \right) \downarrow \text{24}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{3}{5} \left(\frac{a^4(301A + 276B) \sin(c + dx) \cos^3(c + dx)}{4d} + \frac{1}{4} \left(35a^4(49A + 44B) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) - \frac{8a^4(252A + 227B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) \right) \right) \right) + \\ \left. \frac{aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^3}{7d} \right)$$

input `Int[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^3*(a + a*Cos[c + d*x])^3*Sin[c + d*x])/(7*d) + (((7*A + 10*B)*Cos[c + d*x]^3*(a^2 + a^2*Cos[c + d*x])^2*Sin[c + d*x])/(6*d) + ((98*(A + B)*Cos[c + d*x]^3*(a^4 + a^4*Cos[c + d*x])*Sin[c + d*x])/(5*d) + (3*((a^4*(301*A + 276*B)*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (35*a^4*(49*A + 44*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (8*a^4*(252*A + 227*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/4))/5)/6)/7`

3.28.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.28.4 Maple [A] (verified)

Time = 5.70 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.54

method	result
parallelrisch	$15 \left(\frac{(127A+124B)\sin(2dx+2c)}{15} + \frac{(16A+\frac{157B}{9})\sin(3dx+3c)}{5} + \left(A+\frac{4B}{3}\right)\sin(4dx+4c) + \frac{(16A+31B)\sin(5dx+5c)}{75} + \frac{(A+4B)\sin(6dx+6c)}{45} \right)$
risch	$\frac{49a^4xA}{16} + \frac{11a^4Bx}{4} + \frac{11\sin(dx+c)a^4A}{2d} + \frac{323\sin(dx+c)Ba^4}{64d} + \frac{Ba^4\sin(7dx+7c)}{448d} + \frac{\sin(6dx+6c)a^4A}{192d} + \frac{\sin(6dx+6c)a^4B}{192d}$
parts	$(a^4A+4Ba^4) \left(\frac{\left(\cos^5(dx+c) + \frac{5\cos^3(dx+c)}{4} + \frac{15\cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{(4a^4A+B a^4)(2+\cos^2(dx+c))\sin(dx+c)}{3d}$
derivativedivides	$a^4A \left(\frac{\left(\cos^5(dx+c) + \frac{5\cos^3(dx+c)}{4} + \frac{15\cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{Ba^4 \left(\frac{16}{5} + \cos^6(dx+c) + \frac{6\cos^4(dx+c)}{5} + \frac{8\cos^2(dx+c)}{5} \right)}{7}$
default	$a^4A \left(\frac{\left(\cos^5(dx+c) + \frac{5\cos^3(dx+c)}{4} + \frac{15\cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{Ba^4 \left(\frac{16}{5} + \cos^6(dx+c) + \frac{6\cos^4(dx+c)}{5} + \frac{8\cos^2(dx+c)}{5} \right)}{7}$
norman	$\frac{a^4(49A+44B)x}{16} + \frac{128a^4(49A+44B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{35d} + \frac{283a^4(49A+44B)\left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{120d} + \frac{5a^4(49A+44B)\left(\tan^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6d} + \frac{a^4(49A+44B)\left(\tan^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{11d}$

3.28. $\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$

```
input int(cos(d*x+c)^2*(a+cos(d*x+c))*a^4*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 15/64*(1/15*(127*A+124*B)*sin(2*d*x+2*c)+1/5*(16*A+157/9*B)*sin(3*d*x+3*c)
+(A+4/3*B)*sin(4*d*x+4*c)+1/75*(16*A+31*B)*sin(5*d*x+5*c)+1/45*(A+4*B)*sin
(6*d*x+6*c)+1/105*B*sin(7*d*x+7*c)+1/15*(352*A+323*B)*sin(d*x+c)+196/15*(A
+44/49*B)*x*d)*a^4/d
```

3.28.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.62

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{105(49A + 44B)a^4 dx + (240Ba^4 \cos(dx + c))^6 + 280(A + 4B)a^4 \cos(dx + c)^5 + 192(7A + 12B)a^4 \cos(dx + c)^4 + 70(41A + 44B)a^4 \cos(dx + c)^3 + 16(252A + 227B)a^4 \cos(dx + c)^2 + 105(49A + 44B)a^4 \cos(dx + c) + 32(252A + 227B)a^4 \sin(dx + c)}{d}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/1680*(105*(49*A + 44*B))*a^4*d*x + (240*B*a^4*cos(d*x + c)^6 + 280*(A + 4
*B)*a^4*cos(d*x + c)^5 + 192*(7*A + 12*B)*a^4*cos(d*x + c)^4 + 70*(41*A +
44*B)*a^4*cos(d*x + c)^3 + 16*(252*A + 227*B)*a^4*cos(d*x + c)^2 + 105*(49
*A + 44*B)*a^4*cos(d*x + c) + 32*(252*A + 227*B)*a^4*sin(d*x + c))/d
```

3.28.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 960 vs. 2(226) = 452.

Time = 0.59 (sec) , antiderivative size = 960, normalized size of antiderivative = 3.98

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))**4*(A+B*cos(d*x+c)),x)
```

output `Piecewise((5*A*a**4*x*sin(c + d*x)**6/16 + 15*A*a**4*x*sin(c + d*x)**4*cos(c + d*x)**2/16 + 9*A*a**4*x*sin(c + d*x)**4/4 + 15*A*a**4*x*sin(c + d*x)**2*cos(c + d*x)**4/16 + 9*A*a**4*x*sin(c + d*x)**2*cos(c + d*x)**2/2 + A*a**4*x*sin(c + d*x)**2/2 + 5*A*a**4*x*cos(c + d*x)**6/16 + 9*A*a**4*x*cos(c + d*x)**4/4 + A*a**4*x*cos(c + d*x)**2/2 + 5*A*a**4*sin(c + d*x)**5*cos(c + d*x)/(16*d) + 32*A*a**4*sin(c + d*x)**5/(15*d) + 5*A*a**4*sin(c + d*x)**3*cos(c + d*x)**3/(6*d) + 16*A*a**4*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 9*A*a**4*sin(c + d*x)**3*cos(c + d*x)/(4*d) + 8*A*a**4*sin(c + d*x)**3/(3*d) + 11*A*a**4*sin(c + d*x)*cos(c + d*x)**5/(16*d) + 4*A*a**4*sin(c + d*x)*cos(c + d*x)**4/d + 15*A*a**4*sin(c + d*x)*cos(c + d*x)**3/(4*d) + 4*A*a**4*sin(c + d*x)*cos(c + d*x)**2/d + A*a**4*sin(c + d*x)*cos(c + d*x)/(2*d) + 5*B*a**4*x*sin(c + d*x)**6/4 + 15*B*a**4*x*sin(c + d*x)**4*cos(c + d*x)**2/4 + 3*B*a**4*x*sin(c + d*x)**4/2 + 15*B*a**4*x*sin(c + d*x)**2*cos(c + d*x)**4/4 + 3*B*a**4*x*sin(c + d*x)**2*cos(c + d*x)**2 + 5*B*a**4*x*cos(c + d*x)**6/4 + 3*B*a**4*x*cos(c + d*x)**4/2 + 16*B*a**4*sin(c + d*x)**7/(35*d) + 8*B*a**4*sin(c + d*x)**5*cos(c + d*x)**2/(5*d) + 5*B*a**4*sin(c + d*x)**5*cos(c + d*x)/(4*d) + 16*B*a**4*sin(c + d*x)**5/(5*d) + 2*B*a**4*sin(c + d*x)**3*cos(c + d*x)**4/d + 10*B*a**4*sin(c + d*x)**3*cos(c + d*x)**3/(3*d) + 8*B*a**4*sin(c + d*x)**3*cos(c + d*x)**2/d + 3*B*a**4*sin(c + d*x)**3*cos(c + d*x)/(2*d) + 2*B*a**4*sin(c + d*x)**3/(3*d) + B*a**4*si...`

3.28.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.48

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{1792 (3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c)) Aa^4 - 35 (4 \sin(2dx + 2c)^3 - 60 dx - 60c - 9)}{105}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="maxima")`


```
output 1/6720*(1792*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*A*a^
4 - 35*(4*sin(2*d*x + 2*c)^3 - 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*sin
(2*d*x + 2*c))*A*a^4 - 8960*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^4 + 1260
*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^4 + 1680*(2*d
*x + 2*c + sin(2*d*x + 2*c))*A*a^4 - 192*(5*sin(d*x + c)^7 - 21*sin(d*x +
c)^5 + 35*sin(d*x + c)^3 - 35*sin(d*x + c))*B*a^4 + 2688*(3*sin(d*x + c)^5
- 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a^4 - 140*(4*sin(2*d*x + 2*c)^3
- 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*sin(2*d*x + 2*c))*B*a^4 - 2240*(
sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^4 + 840*(12*d*x + 12*c + sin(4*d*x +
4*c) + 8*sin(2*d*x + 2*c))*B*a^4)/d
```

3.28.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.80

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^4 \sin(7dx + 7c)}{448d} + \frac{1}{16} (49Aa^4 + 44Ba^4)x$$

$$+ \frac{(Aa^4 + 4Ba^4) \sin(6dx + 6c)}{192d} + \frac{(16Aa^4 + 31Ba^4) \sin(5dx + 5c)}{320d}$$

$$+ \frac{5(3Aa^4 + 4Ba^4) \sin(4dx + 4c)}{64d} + \frac{(144Aa^4 + 157Ba^4) \sin(3dx + 3c)}{192d}$$

$$+ \frac{(127Aa^4 + 124Ba^4) \sin(2dx + 2c)}{64d} + \frac{(352Aa^4 + 323Ba^4) \sin(dx + c)}{64d}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="g
iac")
```

```
output 1/448*B*a^4*sin(7*d*x + 7*c)/d + 1/16*(49*A*a^4 + 44*B*a^4)*x + 1/192*(A*a
^4 + 4*B*a^4)*sin(6*d*x + 6*c)/d + 1/320*(16*A*a^4 + 31*B*a^4)*sin(5*d*x +
5*c)/d + 5/64*(3*A*a^4 + 4*B*a^4)*sin(4*d*x + 4*c)/d + 1/192*(144*A*a^4 +
157*B*a^4)*sin(3*d*x + 3*c)/d + 1/64*(127*A*a^4 + 124*B*a^4)*sin(2*d*x +
2*c)/d + 1/64*(352*A*a^4 + 323*B*a^4)*sin(d*x + c)/d
```

3.28.9 Mupad [B] (verification not implemented)

Time = 1.79 (sec) , antiderivative size = 353, normalized size of antiderivative = 1.46

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{49Aa^4}{8} + \frac{11Ba^4}{2}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{13} + \left(\frac{245Aa^4}{6} + \frac{110Ba^4}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{11} + \left(\frac{13867Aa^4}{120} + \frac{3113Ba^4}{30}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{896Aa^4}{5} + \frac{5632Ba^4}{35}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{67Aa^4}{120} + \frac{3113Ba^4}{30}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{19157Aa^4}{120} + \frac{1501Ba^4}{10}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{207Aa^4}{8} + \frac{53Ba^4}{2}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \left(\frac{49Aa^4}{8} + \frac{11Ba^4}{2}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{14} + 7 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{12} + 21 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 35 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + 21 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 7 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 1 \right) - \frac{a^4(49A + 44B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{8d} + \frac{a^4 \operatorname{atan}\left(\frac{a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (49A + 44B)}{8 \left(\frac{49Aa^4}{8} + \frac{11Ba^4}{2}\right)}\right) (49A + 44B)}{8d}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4,x)`output `(tan(c/2 + (d*x)/2)*((207*A*a^4)/8 + (53*B*a^4)/2) + tan(c/2 + (d*x)/2)^13 * ((49*A*a^4)/8 + (11*B*a^4)/2) + tan(c/2 + (d*x)/2)^11 * ((245*A*a^4)/6 + (110*B*a^4)/3) + tan(c/2 + (d*x)/2)^9 * ((13867*A*a^4)/120 + (3113*B*a^4)/30) + tan(c/2 + (d*x)/2)^7 * ((896*A*a^4)/5 + (5632*B*a^4)/35) + tan(c/2 + (d*x)/2)^5 * ((19157*A*a^4)/120 + (1501*B*a^4)/10) / (d*(7*tan(c/2 + (d*x)/2)^2 + 21*tan(c/2 + (d*x)/2)^4 + 35*tan(c/2 + (d*x)/2)^6 + 21*tan(c/2 + (d*x)/2)^8 + 7*tan(c/2 + (d*x)/2)^10 + 1) - (a^4*(49*A + 44*B)*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2)) / (8*d) + (a^4*atan((a^4*tan(c/2 + (d*x)/2)*(49*A + 44*B)) / (8*((49*A*a^4)/8 + (11*B*a^4)/2)))) * (49*A + 44*B)) / (8*d)`

3.29 $\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$

3.29.1	Optimal result	452
3.29.2	Mathematica [A] (verified)	453
3.29.3	Rubi [A] (verified)	453
3.29.4	Maple [A] (verified)	456
3.29.5	Fricas [A] (verification not implemented)	456
3.29.6	Sympy [B] (verification not implemented)	457
3.29.7	Maxima [A] (verification not implemented)	458
3.29.8	Giac [A] (verification not implemented)	459
3.29.9	Mupad [B] (verification not implemented)	459

3.29.1 Optimal result

Integrand size = 29, antiderivative size = 185

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{7}{16}a^4(8A + 7B)x + \frac{4a^4(8A + 7B) \sin(c + dx)}{5d} + \frac{27a^4(8A + 7B) \cos(c + dx) \sin(c + dx)}{80d}$$

$$+ \frac{a^4(8A + 7B) \cos^3(c + dx) \sin(c + dx)}{40d} + \frac{(6A - B)(a + a \cos(c + dx))^4 \sin(c + dx)}{30d}$$

$$+ \frac{B(a + a \cos(c + dx))^5 \sin(c + dx)}{6ad} - \frac{2a^4(8A + 7B) \sin^3(c + dx)}{15d}$$

output `7/16*a^4*(8*A+7*B)*x+4/5*a^4*(8*A+7*B)*sin(d*x+c)/d+27/80*a^4*(8*A+7*B)*cos(d*x+c)*sin(d*x+c)/d+1/40*a^4*(8*A+7*B)*cos(d*x+c)^3*sin(d*x+c)/d+1/30*(6*A-B)*(a+a*cos(d*x+c))^4*sin(d*x+c)/d+1/6*B*(a+a*cos(d*x+c))^5*sin(d*x+c)/a/d-2/15*a^4*(8*A+7*B)*sin(d*x+c)^3/d`

3.29.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.72

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{a^4(2940Bc + 3360Adx + 2940Bdx + 120(49A + 44B) \sin(c + dx) + 15(128A + 127B) \sin(2(c + dx)) + 580A \sin(3(c + dx)) + 720B \sin(3(c + dx)) + 120A \sin(4(c + dx)) + 225B \sin(4(c + dx)) + 12A \sin(5(c + dx)) + 48B \sin(5(c + dx)) + 5B \sin(6(c + dx)))}{960d}$$

input `Integrate[Cos[c + d*x]*(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output `(a^4*(2940*B*c + 3360*A*d*x + 2940*B*d*x + 120*(49*A + 44*B)*Sin[c + d*x] + 15*(128*A + 127*B)*Sin[2*(c + d*x)] + 580*A*Ssin[3*(c + d*x)] + 720*B*Ssin[3*(c + d*x)] + 120*A*Ssin[4*(c + d*x)] + 225*B*Ssin[4*(c + d*x)] + 12*A*Ssin[5*(c + d*x)] + 48*B*Ssin[5*(c + d*x)] + 5*B*Ssin[6*(c + d*x)])/(960*d)`

3.29.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.92, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3447, 3042, 3502, 3042, 3230, 3042, 3124, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(c + dx)(a \cos(c + dx) + a)^4(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^4 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int (a \cos(c + dx) + a)^4 (A \cos(c + dx) + B \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^4 \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{3502}$$

$$\begin{aligned}
& \frac{\int (\cos(c+dx)a+a)^4(5aB+a(6A-B)\cos(c+dx))dx}{6a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^5}{6ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\int (\sin(c+dx+\frac{\pi}{2})a+a)^4(5aB+a(6A-B)\sin(c+dx+\frac{\pi}{2}))dx}{6a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^5}{6ad} \\
& \quad \downarrow \text{3230} \\
& \frac{\frac{3}{5}a(8A+7B)\int (\cos(c+dx)a+a)^4dx + \frac{a(6A-B)\sin(c+dx)(a\cos(c+dx)+a)^4}{5d}}{6a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^5}{6ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{3}{5}a(8A+7B)\int (\sin(c+dx+\frac{\pi}{2})a+a)^4dx + \frac{a(6A-B)\sin(c+dx)(a\cos(c+dx)+a)^4}{5d}}{6a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^5}{6ad} \\
& \quad \downarrow \text{3124} \\
& \frac{\frac{3}{5}a(8A+7B)\int (\cos^4(c+dx)a^4 + 4\cos^3(c+dx)a^4 + 6\cos^2(c+dx)a^4 + 4\cos(c+dx)a^4 + a^4)dx + \frac{a(6A-B)\sin(c+dx)(a\cos(c+dx)+a)^4}{5d}}{6a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^5}{6ad} \\
& \quad \downarrow \text{2009} \\
& \frac{\frac{3}{5}a(8A+7B)\left(-\frac{4a^4\sin^3(c+dx)}{3d} + \frac{8a^4\sin(c+dx)}{d} + \frac{a^4\sin(c+dx)\cos^3(c+dx)}{4d} + \frac{27a^4\sin(c+dx)\cos(c+dx)}{8d} + \frac{35a^4x}{8}\right) + \frac{a(6A-B)\sin(c+dx)(a\cos(c+dx)+a)^4}{5d}}{6a} + \frac{B\sin(c+dx)(a\cos(c+dx)+a)^5}{6ad}
\end{aligned}$$

input `Int[Cos[c + d*x]*(a + a*cos[c + d*x])^4*(A + B*cos[c + d*x]),x]`

output `(B*(a + a*cos[c + d*x])^5*sin[c + d*x])/(6*a*d) + ((a*(6*A - B)*(a + a*cos[c + d*x])^4*sin[c + d*x])/(5*d) + (3*a*(8*A + 7*B)*((35*a^4*x)/8 + (8*a^4*sin[c + d*x])/d) + (27*a^4*cos[c + d*x]*sin[c + d*x])/(8*d) + (a^4*cos[c + d*x]^3*sin[c + d*x])/(4*d) - (4*a^4*sin[c + d*x]^3)/(3*d))/5)/(6*a)`

3.29.3.1 Defintions of rubi rules used

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3124 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(a + b*sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`
- rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.29.4 Maple [A] (verified)

Time = 4.10 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.60

method	result
parallelrisc	$\left(\left(16A + \frac{127B}{8}\right) \sin(2dx+2c) + \left(\frac{29A}{6} + 6B\right) \sin(3dx+3c) + \left(A + \frac{15B}{8}\right) \sin(4dx+4c) + \left(\frac{A}{10} + \frac{2B}{5}\right) \sin(5dx+5c) + \frac{B \sin(6dx+6c)}{24}\right)$
risc	$\frac{7a^4xA}{2} + \frac{49a^4Bx}{16} + \frac{49 \sin(dx+c)a^4A}{8d} + \frac{11 \sin(dx+c)B a^4}{2d} + \frac{\sin(6dx+6c)B a^4}{192d} + \frac{\sin(5dx+5c)a^4A}{80d} + \frac{\sin(5dx+5c)a^4A}{80d}$
parts	$\frac{(a^4A+4B a^4) \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right) \sin(dx+c)}{5d} + \frac{(4a^4A+B a^4) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{(4a^4A)}{2}$
derivativedivides	$\frac{a^4A \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right) \sin(dx+c)}{5} + B a^4 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8}\right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16}\right)$
default	$\frac{a^4A \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right) \sin(dx+c)}{5} + B a^4 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8}\right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16}\right)$
norman	$\frac{7a^4(8A+7B)x}{16} + \frac{281a^4(8A+7B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{20d} + \frac{231a^4(8A+7B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{20d} + \frac{119a^4(8A+7B)\left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{24d} + \frac{7a^4(8A+7B)}{24d}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*a)^4*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/8*((16*A+127/8*B)*sin(2*d*x+2*c)+(29/6*A+6*B)*sin(3*d*x+3*c)+(A+15/8*B)*sin(4*d*x+4*c)+(1/10*A+2/5*B)*sin(5*d*x+5*c)+1/24*B*sin(6*d*x+6*c)+(49*A+44*B)*sin(d*x+c)+28*d*x*(A+7/8*B))*a^4/d
```

3.29.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.70

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{105(8A + 7B)a^4dx + (40Ba^4 \cos(dx + c))^5 + 48(A + 4B)a^4 \cos(dx + c)^4 + 10(24A + 41B)a^4 \cos(dx + c)^3 + \dots}{2}$$

3.29. $\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/240*(105*(8*A + 7*B)*a^4*d*x + (40*B*a^4*cos(d*x + c)^5 + 48*(A + 4*B)*a^4*cos(d*x + c)^4 + 10*(24*A + 41*B)*a^4*cos(d*x + c)^3 + 32*(17*A + 18*B)*a^4*cos(d*x + c)^2 + 105*(8*A + 7*B)*a^4*cos(d*x + c) + 16*(83*A + 72*B)*a^4*sin(d*x + c))/d`

3.29.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 765 vs. $2(170) = 340$.

Time = 0.43 (sec) , antiderivative size = 765, normalized size of antiderivative = 4.14

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{3Aa^4x \sin^4(c+dx)}{2} + 3Aa^4x \sin^2(c + dx) \cos^2(c + dx) + 2Aa^4x \sin^2(c + dx) + \frac{3Aa^4x \cos^4(c+dx)}{2} + 2Aa^4x \cos^2(c+dx) \\ x(A + B \cos(c)) (a \cos(c) + a)^4 \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))**4*(A+B*cos(d*x+c)),x)`

output `Piecewise((3*A*a**4*x*sin(c + d*x)**4/2 + 3*A*a**4*x*sin(c + d*x)**2*cos(c + d*x)**2 + 2*A*a**4*x*sin(c + d*x)**2 + 3*A*a**4*x*cos(c + d*x)**4/2 + 2*A*a**4*x*cos(c + d*x)**2 + 8*A*a**4*sin(c + d*x)**5/(15*d) + 4*A*a**4*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 3*A*a**4*sin(c + d*x)**3*cos(c + d*x)/(2*d) + 4*A*a**4*sin(c + d*x)**3/d + A*a**4*sin(c + d*x)*cos(c + d*x)**4/d + 5*A*a**4*sin(c + d*x)*cos(c + d*x)**3/(2*d) + 6*A*a**4*sin(c + d*x)*cos(c + d*x)**2/d + 2*A*a**4*sin(c + d*x)*cos(c + d*x)/d + A*a**4*sin(c + d*x)/d + 5*B*a**4*x*sin(c + d*x)**6/16 + 15*B*a**4*x*sin(c + d*x)**4*cos(c + d*x)**2/16 + 9*B*a**4*x*sin(c + d*x)**4/4 + 15*B*a**4*x*sin(c + d*x)**2*cos(c + d*x)**4/16 + 9*B*a**4*x*sin(c + d*x)**2*cos(c + d*x)**2/2 + B*a**4*x*sin(c + d*x)**2/2 + 5*B*a**4*x*cos(c + d*x)**6/16 + 9*B*a**4*x*cos(c + d*x)**4/4 + B*a**4*x*cos(c + d*x)**2/2 + 5*B*a**4*sin(c + d*x)**5*cos(c + d*x)/(16*d) + 32*B*a**4*sin(c + d*x)**5/(15*d) + 5*B*a**4*sin(c + d*x)**3*cos(c + d*x)**3/(6*d) + 16*B*a**4*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 9*B*a**4*sin(c + d*x)**3*cos(c + d*x)/(4*d) + 8*B*a**4*sin(c + d*x)**3/(3*d) + 11*B*a**4*sin(c + d*x)*cos(c + d*x)**5/(16*d) + 4*B*a**4*sin(c + d*x)*cos(c + d*x)**4/d + 15*B*a**4*sin(c + d*x)*cos(c + d*x)**3/(4*d) + 4*B*a**4*sin(c + d*x)*cos(c + d*x)**2/d + B*a**4*sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**4*cos(c), True))`

3.29.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.61

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{64(3 \sin(dx + c)^5 - 10 \sin(dx + c)^3 + 15 \sin(dx + c))Aa^4 - 1920(\sin(dx + c)^3 - 3 \sin(dx + c))Aa^4 + \dots}{d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/960*(64*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*A*a^4 - 1920*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^4 + 120*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^4 + 960*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^4 + 256*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a^4 - 5*(4*sin(2*d*x + 2*c)^3 - 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*sin(2*d*x + 2*c))*B*a^4 - 1280*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^4 + 180*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^4 + 240*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^4 + 960*A*a^4*sin(d*x + c))/d`

3.29. $\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$

3.29.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.90

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{Ba^4 \sin(6dx + 6c)}{192d} + \frac{7}{16}(8Aa^4 + 7Ba^4)x + \frac{(Aa^4 + 4Ba^4) \sin(5dx + 5c)}{80d}$$

$$+ \frac{(8Aa^4 + 15Ba^4) \sin(4dx + 4c)}{64d} + \frac{(29Aa^4 + 36Ba^4) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(128Aa^4 + 127Ba^4) \sin(2dx + 2c)}{64d} + \frac{(49Aa^4 + 44Ba^4) \sin(dx + c)}{8d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/192*B*a^4*sin(6*d*x + 6*c)/d + 7/16*(8*A*a^4 + 7*B*a^4)*x + 1/80*(A*a^4 + 4*B*a^4)*sin(5*d*x + 5*c)/d + 1/64*(8*A*a^4 + 15*B*a^4)*sin(4*d*x + 4*c)/d + 1/48*(29*A*a^4 + 36*B*a^4)*sin(3*d*x + 3*c)/d + 1/64*(128*A*a^4 + 127*B*a^4)*sin(2*d*x + 2*c)/d + 1/8*(49*A*a^4 + 44*B*a^4)*sin(d*x + c)/d`

3.29.9 Mupad [B] (verification not implemented)

Time = 1.70 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.71

$$\int \cos(c + dx)(a + a \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{\left(7Aa^4 + \frac{49Ba^4}{8}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{11} + \left(\frac{119Aa^4}{3} + \frac{833Ba^4}{24}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{462Aa^4}{5} + \frac{1617Ba^4}{20}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{12} + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 15 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8\right)}$$

$$- \frac{7a^4(8A + 7B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{8d}$$

$$+ \frac{7a^4 \operatorname{atan}\left(\frac{7a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)(8A + 7B)}{8\left(7Aa^4 + \frac{49Ba^4}{8}\right)}\right) (8A + 7B)}{8d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4,x)`

output $(\tan(c/2 + (d*x)/2)*(25*A*a^4 + (207*B*a^4)/8) + \tan(c/2 + (d*x)/2)^{11}*(7*A*a^4 + (49*B*a^4)/8) + \tan(c/2 + (d*x)/2)^9*((119*A*a^4)/3 + (833*B*a^4)/24) + \tan(c/2 + (d*x)/2)^3*((233*A*a^4)/3 + (1471*B*a^4)/24) + \tan(c/2 + (d*x)/2)^7*((462*A*a^4)/5 + (1617*B*a^4)/20) + \tan(c/2 + (d*x)/2)^5*((562*A*a^4)/5 + (1967*B*a^4)/20))/(d*(6*\tan(c/2 + (d*x)/2)^2 + 15*\tan(c/2 + (d*x)/2)^4 + 20*\tan(c/2 + (d*x)/2)^6 + 15*\tan(c/2 + (d*x)/2)^8 + 6*\tan(c/2 + (d*x)/2)^{10} + \tan(c/2 + (d*x)/2)^{12} + 1)) - (7*a^4*(8*A + 7*B)*(atan(\tan(c/2 + (d*x)/2)) - (d*x)/2))/(8*d) + (7*a^4*atan((7*a^4*\tan(c/2 + (d*x)/2)*(8*A + 7*B))/(8*(7*A*a^4 + (49*B*a^4)/8)))*(8*A + 7*B))/(8*d)$

3.30 $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$

3.30.1	Optimal result	461
3.30.2	Mathematica [A] (verified)	461
3.30.3	Rubi [A] (verified)	462
3.30.4	Maple [A] (verified)	464
3.30.5	Fricas [A] (verification not implemented)	464
3.30.6	Sympy [B] (verification not implemented)	465
3.30.7	Maxima [A] (verification not implemented)	466
3.30.8	Giac [A] (verification not implemented)	466
3.30.9	Mupad [B] (verification not implemented)	467

3.30.1 Optimal result

Integrand size = 23, antiderivative size = 150

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \frac{7}{8} a^4 (5A + 4B)x + \frac{8a^4(5A + 4B) \sin(c + dx)}{5d}$$

$$+ \frac{27a^4(5A + 4B) \cos(c + dx) \sin(c + dx)}{40d} + \frac{a^4(5A + 4B) \cos^3(c + dx) \sin(c + dx)}{20d}$$

$$+ \frac{B(a + a \cos(c + dx))^4 \sin(c + dx)}{5d} - \frac{4a^4(5A + 4B) \sin^3(c + dx)}{15d}$$

```
output 7/8*a^4*(5*A+4*B)*x+8/5*a^4*(5*A+4*B)*sin(d*x+c)/d+27/40*a^4*(5*A+4*B)*cos
(d*x+c)*sin(d*x+c)/d+1/20*a^4*(5*A+4*B)*cos(d*x+c)^3*sin(d*x+c)/d+1/5*B*(a
+a*cos(d*x+c))^4*sin(d*x+c)/d-4/15*a^4*(5*A+4*B)*sin(d*x+c)^3/d
```

3.30.2 Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.89

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \frac{a^4 \sin(c + dx) \left(210(5A + 4B) \arcsin \left(\sqrt{\sin^2 \left(\frac{1}{2}(c + dx) \right)} \right) + (800A + 664B + 15(27A + 28B) \cos(c + dx) \right)}{120d \sqrt{\sin^2(c + dx)}}$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output `(a^4*Sin[c + d*x]*(210*(5*A + 4*B)*ArcSin[Sqrt[Sin[(c + d*x)/2]^2]] + (800*A + 664*B + 15*(27*A + 28*B)*Cos[c + d*x] + 16*(10*A + 17*B)*Cos[c + d*x]^2 + 30*(A + 4*B)*Cos[c + d*x]^3 + 24*B*Cos[c + d*x]^4)*Sqrt[Sin[c + d*x]^2])/(120*d*Sqrt[Sin[c + d*x]^2])`

3.30.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.83, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {3042, 3230, 3042, 3124, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^4 (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^4 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx \\
 & \quad \downarrow \text{3230} \\
 & \frac{1}{5}(5A + 4B) \int (\cos(c + dx)a + a)^4 dx + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^4}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5}(5A + 4B) \int \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^4 dx + \frac{B \sin(c + dx)(a \cos(c + dx) + a)^4}{5d} \\
 & \quad \downarrow \text{3124} \\
 & \frac{1}{5}(5A + 4B) \int (\cos^4(c + dx)a^4 + 4 \cos^3(c + dx)a^4 + 6 \cos^2(c + dx)a^4 + 4 \cos(c + dx)a^4 + a^4) dx + \\
 & \quad \frac{B \sin(c + dx)(a \cos(c + dx) + a)^4}{5d} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$4B) \left(-\frac{4a^4 \sin^3(c+dx)}{3d} + \frac{8a^4 \sin(c+dx)}{d} + \frac{\frac{1}{5}(5A + a^4 \sin(c+dx) \cos^3(c+dx))}{4d} + \frac{27a^4 \sin(c+dx) \cos(c+dx)}{8d} + \frac{35a^4 x}{8} \right. \\ \left. \frac{B \sin(c+dx)(a \cos(c+dx) + a)^4}{5d} \right)$$

input `Int[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output `(B*(a + a*Cos[c + d*x])^4*Sin[c + d*x])/(5*d) + ((5*A + 4*B)*((35*a^4*x)/8 + (8*a^4*Sin[c + d*x])/d + (27*a^4*Cos[c + d*x]*Sin[c + d*x])/(8*d) + (a^4*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) - (4*a^4*Sin[c + d*x]^3)/(3*d)))/5`

3.30.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3124 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(a + b*sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

3.30.4 Maple [A] (verified)

Time = 3.67 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.63

method	result
parallelrisc	$\frac{((56A+64B)\sin(2dx+2c) + \left(\frac{32A}{3} + \frac{58B}{3}\right)\sin(3dx+3c) + (A+4B)\sin(4dx+4c) + \frac{2B\sin(5dx+5c)}{5} + (224A+196B)\sin(dx+c) + 140(A+4/5B)x)d}{32d}$
risc	$\frac{35a^4xA}{8} + \frac{7a^4Bx}{2} + \frac{7\sin(dx+c)a^4A}{d} + \frac{49\sin(dx+c)Ba^4}{8d} + \frac{\sin(5dx+5c)Ba^4}{80d} + \frac{\sin(4dx+4c)a^4A}{32d} + \frac{\sin(4dx+4c)a^4A}{80d}$
parts	$a^4xA + \frac{(a^4A+4Ba^4)\left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2})\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right)}{d} + \frac{(4a^4A+Ba^4)\sin(dx+c)}{d} + \frac{(4a^4A+6Ba^4)\sin(dx+c)}{d}$
derivativdivides	$a^4A\left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2})\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ba^4\left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right)\sin(dx+c)}{5} + \frac{4a^4A(2+\cos^2(dx+c))}{5}$
default	$a^4A\left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2})\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ba^4\left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3}\right)\sin(dx+c)}{5} + \frac{4a^4A(2+\cos^2(dx+c))}{5}$
norman	$\frac{7a^4(5A+4B)x}{8} + \frac{79a^4(5A+4B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6d} + \frac{224a^4(5A+4B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{15d} + \frac{49a^4(5A+4B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6d} + \frac{7a^4(5A+4B)\sin(dx+c)}{120d}$

input `int((a+cos(d*x+c))*a)^4*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/32*((56*A+64*B)*sin(2*d*x+2*c)+(32/3*A+58/3*B)*sin(3*d*x+3*c)+(A+4*B)*sin(4*d*x+4*c)+2/5*B*sin(5*d*x+5*c)+(224*A+196*B)*sin(d*x+c)+140*(A+4/5*B)*x*d)*a^4/d`

3.30.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.73

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \frac{105(5A + 4B)a^4 dx + (24Ba^4 \cos(dx + c))^4 + 30(A + 4B)a^4 \cos(dx + c)^3 + 16(10A + 17B)a^4 \cos(dx + c)^2 + 8(5A + 4B)a^4 \cos(dx + c) + 7a^4}{120d}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output $1/120*(105*(5*A + 4*B)*a^4*d*x + (24*B*a^4*\cos(d*x + c)^4 + 30*(A + 4*B)*a^4*\cos(d*x + c)^3 + 16*(10*A + 17*B)*a^4*\cos(d*x + c)^2 + 15*(27*A + 28*B)*a^4*\cos(d*x + c) + 8*(100*A + 83*B)*a^4)*\sin(d*x + c))/d$

3.30.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 544 vs. $2(141) = 282$.

Time = 0.31 (sec) , antiderivative size = 544, normalized size of antiderivative = 3.63

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{3Aa^4 x \sin^4(c+dx)}{8} + \frac{3Aa^4 x \sin^2(c+dx) \cos^2(c+dx)}{4} + 3Aa^4 x \sin^2(c+dx) + \frac{3Aa^4 x \cos^4(c+dx)}{8} + 3Aa^4 x \cos^2(c+dx) \\ x(A + B \cos(c)) (a \cos(c) + a)^4 \end{cases}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c)),x)`

output `Piecewise((3*A*a**4*x*sin(c + d*x)**4/8 + 3*A*a**4*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*A*a**4*x*sin(c + d*x)**2 + 3*A*a**4*x*cos(c + d*x)**4/8 + 3*A*a**4*x*cos(c + d*x)**2 + A*a**4*x + 3*A*a**4*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 8*A*a**4*sin(c + d*x)**3/(3*d) + 5*A*a**4*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 4*A*a**4*sin(c + d*x)*cos(c + d*x)**2/d + 3*A*a**4*sin(c + d*x)*cos(c + d*x)/d + 4*A*a**4*sin(c + d*x)/d + 3*B*a**4*x*sin(c + d*x)**4/2 + 3*B*a**4*x*sin(c + d*x)**2*cos(c + d*x)**2 + 2*B*a**4*x*sin(c + d*x)**2 + 3*B*a**4*x*cos(c + d*x)**4/2 + 2*B*a**4*x*cos(c + d*x)**2 + 8*B*a**4*sin(c + d*x)**5/(15*d) + 4*B*a**4*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 3*B*a**4*sin(c + d*x)**3*cos(c + d*x)/(2*d) + 4*B*a**4*sin(c + d*x)**3/d + B*a**4*sin(c + d*x)*cos(c + d*x)**4/d + 5*B*a**4*sin(c + d*x)*cos(c + d*x)**3/(2*d) + 6*B*a**4*sin(c + d*x)*cos(c + d*x)**2/d + 2*B*a**4*sin(c + d*x)*cos(c + d*x)/d + B*a**4*sin(c + d*x)/d, Ne(d, 0)), (x*(A + B*cos(c))*(a*cos(c) + a)**4, True))`

3.30.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.57

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx =$$

$$\frac{640 (\sin(dx + c)^3 - 3 \sin(dx + c)) Aa^4 - 15 (12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(2 dx + 2 c)) Aa^4 -$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="maxima")`output `-1/480*(640*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^4 - 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^4 - 720*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^4 - 480*(d*x + c)*A*a^4 - 32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*a^4 + 960*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^4 - 60*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a^4 - 480*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^4 - 1920*A*a^4*sin(d*x + c) - 480*B*a^4*sin(d*x + c))/d`**3.30.8 Giac [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.93

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \frac{Ba^4 \sin(5 dx + 5 c)}{80 d} + \frac{7}{8} (5 Aa^4 + 4 Ba^4) x$$

$$+ \frac{(Aa^4 + 4 Ba^4) \sin(4 dx + 4 c)}{32 d} + \frac{(16 Aa^4 + 29 Ba^4) \sin(3 dx + 3 c)}{48 d}$$

$$+ \frac{(7 Aa^4 + 8 Ba^4) \sin(2 dx + 2 c)}{4 d} + \frac{7(8 Aa^4 + 7 Ba^4) \sin(dx + c)}{8 d}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/80*B*a^4*sin(5*d*x + 5*c)/d + 7/8*(5*A*a^4 + 4*B*a^4)*x + 1/32*(A*a^4 + 4*B*a^4)*sin(4*d*x + 4*c)/d + 1/48*(16*A*a^4 + 29*B*a^4)*sin(3*d*x + 3*c)/d + 1/4*(7*A*a^4 + 8*B*a^4)*sin(2*d*x + 2*c)/d + 7/8*(8*A*a^4 + 7*B*a^4)*sin(d*x + c)/d`

3.30.9 Mupad [B] (verification not implemented)

Time = 1.64 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.85

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \frac{\left(\frac{35Aa^4}{4} + 7Ba^4\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(\frac{245Aa^4}{6} + \frac{98Ba^4}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{224Aa^4}{3} + \frac{896Ba^4}{15}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)} - \frac{7a^4(5A + 4B) \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{4d} + \frac{7a^4 \operatorname{atan}\left(\frac{7a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (5A + 4B)}{4 \left(\frac{35Aa^4}{4} + 7Ba^4\right)}\right) (5A + 4B)}{4d}$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4,x)`output `(tan(c/2 + (d*x)/2)*((93*A*a^4)/4 + 25*B*a^4) + tan(c/2 + (d*x)/2)^9*((35*A*a^4)/4 + 7*B*a^4) + tan(c/2 + (d*x)/2)^7*((245*A*a^4)/6 + (98*B*a^4)/3) + tan(c/2 + (d*x)/2)^5*((395*A*a^4)/6 + (158*B*a^4)/3) + tan(c/2 + (d*x)/2)^3*((224*A*a^4)/3 + (896*B*a^4)/15))/(d*(5*tan(c/2 + (d*x)/2)^2 + 10*tan(c/2 + (d*x)/2)^4 + 10*tan(c/2 + (d*x)/2)^6 + 5*tan(c/2 + (d*x)/2)^8 + tan(c/2 + (d*x)/2)^10 + 1)) - (7*a^4*(5*A + 4*B)*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(4*d) + (7*a^4*atan((7*a^4*tan(c/2 + (d*x)/2)*(5*A + 4*B))/(4*((35*A*a^4)/4 + 7*B*a^4)))*(5*A + 4*B))/(4*d)`

3.31 $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$

3.31.1	Optimal result	468
3.31.2	Mathematica [A] (verified)	469
3.31.3	Rubi [A] (verified)	469
3.31.4	Maple [A] (verified)	473
3.31.5	Fricas [A] (verification not implemented)	474
3.31.6	Sympy [F]	475
3.31.7	Maxima [A] (verification not implemented)	475
3.31.8	Giac [A] (verification not implemented)	476
3.31.9	Mupad [B] (verification not implemented)	476

3.31.1 Optimal result

Integrand size = 29, antiderivative size = 151

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{1}{8} a^4 (48A + 35B)x + \frac{a^4 A \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{5a^4 (8A + 7B) \sin(c + dx)}{8d}$$

$$+ \frac{aB(a + a \cos(c + dx))^3 \sin(c + dx)}{4d} + \frac{(4A + 7B)(a^2 + a^2 \cos(c + dx))^2 \sin(c + dx)}{12d}$$

$$+ \frac{(32A + 35B)(a^4 + a^4 \cos(c + dx)) \sin(c + dx)}{24d}$$

output `1/8*a^4*(48*A+35*B)*x+a^4*A*arctanh(sin(d*x+c))/d+5/8*a^4*(8*A+7*B)*sin(d*x+c)/d+1/4*a*B*(a+a*cos(d*x+c))^3*sin(d*x+c)/d+1/12*(4*A+7*B)*(a^2+a^2*cos(d*x+c))^2*sin(d*x+c)/d+1/24*(32*A+35*B)*(a^4+a^4*cos(d*x+c))*sin(d*x+c)/d`

3.31.2 Mathematica [A] (verified)

Time = 1.74 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.91

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{a^4 (576Adx + 420Bdx - 96A \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 96A \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) + 24(27A + 28B) \sin(c + dx) + 24(4A + 7B) \sin[2(c + dx)] + 8A \sin[3(c + dx)] + 32B \sin[3(c + dx)] + 3B \sin[4(c + dx)])}{96d}$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(a^4*(576*A*d*x + 420*B*d*x - 96*A*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 96*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 24*(27*A + 28*B)*Sin[c + d*x] + 24*(4*A + 7*B)*Sin[2*(c + d*x)] + 8*A*Sin[3*(c + d*x)] + 32*B*Sin[3*(c + d*x)] + 3*B*Sin[4*(c + d*x)]))/(96*d)`

3.31.3 Rubi [A] (verified)

Time = 1.23 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.517$, Rules used = {3042, 3455, 3042, 3455, 3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(c + dx) (a \cos(c + dx) + a)^4 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx$$

$$\downarrow \text{3455}$$

$$\frac{1}{4} \int (\cos(c + dx)a + a)^3 (4aA + a(4A + 7B) \cos(c + dx)) \sec(c + dx) dx + \frac{aB \sin(c + dx) (a \cos(c + dx) + a)^3}{4d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^3 (4aA + a(4A + 7B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \downarrow \text{3455}$$

$$\frac{1}{4} \left(\frac{1}{3} \int (\cos(c + dx)a + a)^2 (12Aa^2 + (32A + 35B) \cos(c + dx)a^2) \sec(c + dx) dx + \frac{(4A + 7B) \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d} \right) \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 (12Aa^2 + (32A + 35B) \sin(c + dx + \frac{\pi}{2})a^2)}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{(4A + 7B) \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d} \right) \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \downarrow \text{3455}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int 3(\cos(c + dx)a + a) (8Aa^3 + 5(8A + 7B) \cos(c + dx)a^3) \sec(c + dx) dx + \frac{(32A + 35B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{2d} \right) \right) \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \downarrow \text{27}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int (\cos(c + dx)a + a) (8Aa^3 + 5(8A + 7B) \cos(c + dx)a^3) \sec(c + dx) dx + \frac{(32A + 35B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{2d} \right) \right) \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a) (8Aa^3 + 5(8A + 7B) \sin(c + dx + \frac{\pi}{2})a^3)}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{(32A + 35B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{2d} \right) \right) \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \downarrow \text{3447}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int (5(8A + 7B) \cos^2(c + dx)a^4 + 8Aa^4 + (8Aa^4 + 5(8A + 7B)a^4) \cos(c + dx)) \sec(c + dx) dx + \frac{(32A + 35B) \sin(c + dx)}{2d} \right) \right) \\ \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\ \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{5(8A + 7B) \sin(c + dx + \frac{\pi}{2})^2 a^4 + 8Aa^4 + (8Aa^4 + 5(8A + 7B)a^4) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{(32A + 35B) \sin(c + dx)}{2d} \right) \right) \\ \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\ \downarrow \text{3502}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(\int (8Aa^4 + (48A + 35B) \cos(c + dx)a^4) \sec(c + dx) dx + \frac{5a^4(8A + 7B) \sin(c + dx)}{d} \right) \right) + \frac{(32A + 35B) \sin(c + dx)}{2d} \right) \\ \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\ \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(\int \frac{8Aa^4 + (48A + 35B) \sin(c + dx + \frac{\pi}{2}) a^4}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{5a^4(8A + 7B) \sin(c + dx)}{d} \right) \right) + \frac{(32A + 35B) \sin(c + dx)}{2d} \right) \\ \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\ \downarrow \text{3214}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(8a^4 A \int \sec(c + dx) dx + \frac{5a^4(8A + 7B) \sin(c + dx)}{d} + a^4 x(48A + 35B) \right) \right) + \frac{(32A + 35B) \sin(c + dx)}{2d} \right) \\ \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\ \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(8a^4 A \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{5a^4(8A + 7B) \sin(c + dx)}{d} + a^4 x(48A + 35B) \right) \right) + \frac{(32A + 35B) \sin(c + dx)}{2d} \right) \\ \frac{aB \sin(c + dx)(a \cos(c + dx) + a)^3}{4d} \\ \downarrow \text{4257}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(\frac{8a^4 A \operatorname{arctanh}(\sin(c+dx))}{d} + \frac{5a^4(8A+7B)\sin(c+dx)}{d} + a^4 x(48A+35B) \right) + \frac{(32A+35B)\sin(c+dx)}{2} \right) + \frac{aB \sin(c+dx)(a \cos(c+dx) + a)^3}{4d} \right)$$

input `Int[(a + a*cos[c + d*x])^4*(A + B*cos[c + d*x])*Sec[c + d*x], x]`

output `(a*B*(a + a*cos[c + d*x])^3*sin[c + d*x])/(4*d) + (((4*A + 7*B)*(a^2 + a^2*cos[c + d*x])^2*sin[c + d*x])/(3*d) + (((32*A + 35*B)*(a^4 + a^4*cos[c + d*x])*sin[c + d*x])/(2*d) + (3*(a^4*(48*A + 35*B)*x + (8*a^4*A*ArcTanh[Sin[c + d*x]]))/d + (5*a^4*(8*A + 7*B)*sin[c + d*x])/d))/2)/3)/4`

3.31.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*sin[e + f*x])^m*(A*c + (B*c + A*d)*sin[e + f*x] + B*d*sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)^2], x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.31.4 Maple [A] (verified)

Time = 3.46 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.72

method	result
parallelrisch	$-\frac{\left(A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + \left(-A - \frac{7B}{4}\right) \sin(2dx + 2c) + \left(-\frac{A}{12} - \frac{B}{3}\right) \sin(3dx + 3c) - \frac{\sin(4dx + 4c)B}{32}\right)}{d}$
parts	$\frac{a^4 A \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(a^4 A + 4B a^4)(2 + \cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{(4a^4 A + B a^4)(dx+c)}{d} + \frac{(4a^4 A + 6B a^4)}{3d}$
derivativdivides	$\frac{a^4 A (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + B a^4 \left(\frac{\left(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}\right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 4a^4 A \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
default	$\frac{a^4 A (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + B a^4 \left(\frac{\left(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}\right) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 4a^4 A \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)$
risch	$6a^4 x A + \frac{35a^4 B x}{8} - \frac{27ie^{i(dx+c)} a^4 A}{8d} - \frac{7ie^{i(dx+c)} B a^4}{2d} + \frac{27ie^{-i(dx+c)} a^4 A}{8d} + \frac{7ie^{-i(dx+c)} B a^4}{2d} + \frac{a^4 A \ln(e^{i(dx+c)} + 1)}{d}$
norman	$\frac{(6a^4 A + \frac{35}{8} B a^4)x + (6a^4 A + \frac{35}{8} B a^4)x \left(\tan^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (30a^4 A + \frac{175}{8} B a^4)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (30a^4 A + \frac{175}{8} B a^4)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{24d}$

```
input int((a+cos(d*x+c))*a)^4*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)
```

```
output -(A*ln(tan(1/2*d*x+1/2*c)-1)-A*ln(tan(1/2*d*x+1/2*c)+1)+(-A-7/4*B)*sin(2*d*x+2*c)+(-1/12*A-1/3*B)*sin(3*d*x+3*c)-1/32*sin(4*d*x+4*c)*B+(-27/4*A-7*B)*sin(d*x+c)-6*x*(A+35/48*B)*d)*a^4/d
```

3.31.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.78

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{3(48A + 35B)a^4 dx + 12Aa^4 \log(\sin(dx + c) + 1) - 12Aa^4 \log(-\sin(dx + c) + 1) + (6Ba^4 \cos(dx + c) + 24d)}{24d}$$

```
input integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")
```

```
output 1/24*(3*(48*A + 35*B)*a^4*d*x + 12*A*a^4*log(sin(d*x + c) + 1) - 12*A*a^4*log(-sin(d*x + c) + 1) + (6*B*a^4*cos(d*x + c)^3 + 8*(A + 4*B)*a^4*cos(d*x + c)^2 + 3*(16*A + 27*B)*a^4*cos(d*x + c) + 160*(A + B)*a^4)*sin(d*x + c)/d
```

3.31. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$

3.31.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= a^4 \left(\int A \sec(c + dx) dx + \int 4A \cos(c + dx) \sec(c + dx) dx \right. \\ &\quad + \int 6A \cos^2(c + dx) \sec(c + dx) dx + \int 4A \cos^3(c + dx) \sec(c + dx) dx \\ &\quad + \int A \cos^4(c + dx) \sec(c + dx) dx + \int B \cos(c + dx) \sec(c + dx) dx \\ &\quad + \int 4B \cos^2(c + dx) \sec(c + dx) dx + \int 6B \cos^3(c + dx) \sec(c + dx) dx \\ &\quad \left. + \int 4B \cos^4(c + dx) \sec(c + dx) dx + \int B \cos^5(c + dx) \sec(c + dx) dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `a**4*(Integral(A*sec(c + d*x), x) + Integral(4*A*cos(c + d*x)*sec(c + d*x), x) + Integral(6*A*cos(c + d*x)**2*sec(c + d*x), x) + Integral(4*A*cos(c + d*x)**3*sec(c + d*x), x) + Integral(A*cos(c + d*x)**4*sec(c + d*x), x) + Integral(B*cos(c + d*x)*sec(c + d*x), x) + Integral(4*B*cos(c + d*x)**2*sec(c + d*x), x) + Integral(6*B*cos(c + d*x)**3*sec(c + d*x), x) + Integral(4*B*cos(c + d*x)**4*sec(c + d*x), x) + Integral(B*cos(c + d*x)**5*sec(c + d*x), x))`

3.31.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.31

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{32 (\sin(dx + c)^3 - 3 \sin(dx + c)) Aa^4 - 96 (2 dx + 2c + \sin(2 dx + 2c)) Aa^4 - 384 (dx + c) Aa^4 + 128 B a^4 \cos^2(dx + c) \sec(dx + c)}{1}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output
$$\frac{-1/96*(32*(\sin(dx + c))^3 - 3*\sin(dx + c))*A*a^4 - 96*(2*dx + 2*c + \sin(2*dx + 2*c))*A*a^4 - 384*(dx + c)*A*a^4 + 128*(\sin(dx + c))^3 - 3*\sin(dx + c))*B*a^4 - 3*(12*dx + 12*c + \sin(4*dx + 4*c) + 8*\sin(2*dx + 2*c))*B*a^4 - 144*(2*dx + 2*c + \sin(2*dx + 2*c))*B*a^4 - 96*(dx + c)*B*a^4 - 96*A*a^4*\log(\sec(dx + c) + \tan(dx + c)) - 576*A*a^4*\sin(dx + c) - 384*B*a^4*\sin(dx + c))/d$$

3.31.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.42

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{24 A a^4 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 24 A a^4 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + 3 (48 A a^4 + 35 B a^4) (dx + c) + \dots}{\dots}$$

input `integrate((a+a*cos(dx+c))^4*(A+B*cos(dx+c))*sec(dx+c),x, algorithm="giac")`

output
$$\frac{1/24*(24*A*a^4*\log(\text{abs}(\tan(1/2*dx + 1/2*c) + 1)) - 24*A*a^4*\log(\text{abs}(\tan(1/2*dx + 1/2*c) - 1)) + 3*(48*A*a^4 + 35*B*a^4)*(dx + c) + 2*(120*A*a^4*\tan(1/2*dx + 1/2*c)^7 + 105*B*a^4*\tan(1/2*dx + 1/2*c)^7 + 424*A*a^4*\tan(1/2*dx + 1/2*c)^5 + 385*B*a^4*\tan(1/2*dx + 1/2*c)^5 + 520*A*a^4*\tan(1/2*dx + 1/2*c)^3 + 511*B*a^4*\tan(1/2*dx + 1/2*c)^3 + 216*A*a^4*\tan(1/2*dx + 1/2*c) + 279*B*a^4*\tan(1/2*dx + 1/2*c))/(\tan(1/2*dx + 1/2*c)^2 + 1)^4}{d}$$

3.31.9 Mupad [B] (verification not implemented)

Time = 0.74 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.25

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{144 A a^4 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right) + 24 A a^4 \operatorname{atanh} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right) + 105 B a^4 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right) + 12 A a^4 \sin(2c + \dots)}{\dots}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x),x)`

output `(144*A*a^4*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)) + 24*A*a^4*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)) + 105*B*a^4*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)) + 12*A*a^4*sin(2*c + 2*d*x) + A*a^4*sin(3*c + 3*d*x) + 21*B*a^4*sin(2*c + 2*d*x) + 4*B*a^4*sin(3*c + 3*d*x) + (3*B*a^4*sin(4*c + 4*d*x))/8 + 81*A*a^4*sin(c + d*x) + 84*B*a^4*sin(c + d*x))/(12*d)`

3.32 $\int (a+a \cos(c+dx))^4(A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.32.1	Optimal result	478
3.32.2	Mathematica [B] (verified)	479
3.32.3	Rubi [A] (verified)	480
3.32.4	Maple [A] (verified)	484
3.32.5	Fricas [A] (verification not implemented)	485
3.32.6	Sympy [F]	485
3.32.7	Maxima [A] (verification not implemented)	486
3.32.8	Giac [A] (verification not implemented)	486
3.32.9	Mupad [B] (verification not implemented)	487

3.32.1 Optimal result

Integrand size = 31, antiderivative size = 150

$$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{1}{2}a^4(13A + 12B)x + \frac{a^4(4A + B)\operatorname{arctanh}(\sin(c + dx))}{d}$$

$$+ \frac{5a^4(A + 2B) \sin(c + dx)}{2d} - \frac{(3A - B)(a^2 + a^2 \cos(c + dx))^2 \sin(c + dx)}{3d}$$

$$- \frac{(3A - 8B)(a^4 + a^4 \cos(c + dx)) \sin(c + dx)}{6d} + \frac{aA(a + a \cos(c + dx))^3 \tan(c + dx)}{d}$$

```
output 1/2*a^4*(13*A+12*B)*x+a^4*(4*A+B)*arctanh(sin(d*x+c))/d+5/2*a^4*(A+2*B)*si
n(d*x+c)/d-1/3*(3*A-B)*(a^2+a^2*cos(d*x+c))^2*sin(d*x+c)/d-1/6*(3*A-8*B)*
(a^4+a^4*cos(d*x+c))*sin(d*x+c)/d+a*A*(a+a*cos(d*x+c))^3*tan(d*x+c)/d
```

3.32.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 312 vs. $2(150) = 300$.

Time = 4.16 (sec) , antiderivative size = 312, normalized size of antiderivative = 2.08

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{1}{192} a^4 (1 + \cos(c + dx))^4 \sec^8\left(\frac{1}{2}(c + dx)\right) \left(78Ax + 72Bx \right. \\ \left. - \frac{12(4A + B) \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{d} \right. \\ \left. + \frac{12(4A + B) \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{d} + \frac{3(16A + 27B) \cos(dx) \sin(c)}{d} \right. \\ \left. + \frac{3(A + 4B) \cos(2dx) \sin(2c)}{d} + \frac{B \cos(3dx) \sin(3c)}{d} + \frac{3(16A + 27B) \cos(c) \sin(dx)}{d} \right. \\ \left. + \frac{3(A + 4B) \cos(2c) \sin(2dx)}{d} + \frac{B \cos(3c) \sin(3dx)}{d} \right. \\ \left. + \frac{12A \sin(\frac{dx}{2})}{d (\cos(\frac{c}{2}) - \sin(\frac{c}{2})) (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))} \right. \\ \left. + \frac{12A \sin(\frac{dx}{2})}{d (\cos(\frac{c}{2}) + \sin(\frac{c}{2})) (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))} \right)$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `(a^4*(1 + Cos[c + d*x])^4*Sec[(c + d*x)/2]^8*(78*A*x + 72*B*x - (12*(4*A + B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]])/d + (12*(4*A + B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/d + (3*(16*A + 27*B)*Cos[d*x]*Sin[c])/d + (3*(A + 4*B)*Cos[2*d*x]*Sin[2*c])/d + (B*Cos[3*d*x]*Sin[3*c])/d + (3*(16*A + 27*B)*Cos[c]*Sin[d*x])/d + (3*(A + 4*B)*Cos[2*c]*Sin[2*d*x])/d + (B*Cos[3*c]*Sin[3*d*x])/d + (12*A*Sin[(d*x)/2])/(d*(Cos[c/2] - Sin[c/2])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) + (12*A*Sin[(d*x)/2])/(d*(Cos[c/2] + Sin[c/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])))`

3.32.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3454, 3042, 3455, 3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx)(a \cos(c+dx) + a)^4(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c+dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3454} \\
 & \int (\cos(c+dx)a + a)^3 (a(4A + B) - a(3A - B) \cos(c+dx)) \sec(c+dx) dx + \\
 & \quad \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^3}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^3 (a(4A + B) - a(3A - B) \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})} dx + \\
 & \quad \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^3}{d} \\
 & \quad \downarrow \text{3455} \\
 & \frac{1}{3} \int (\cos(c+dx)a + a)^2 (3a^2(4A + B) - a^2(3A - 8B) \cos(c+dx)) \sec(c+dx) dx - \\
 & \quad \frac{(3A - B) \sin(c+dx) (a^2 \cos(c+dx) + a^2)^2}{3d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^3}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^2 (3a^2(4A + B) - a^2(3A - 8B) \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})} dx - \\
 & \quad \frac{(3A - B) \sin(c+dx) (a^2 \cos(c+dx) + a^2)^2}{3d} + \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^3}{d} \\
 & \quad \downarrow \text{3455}
 \end{aligned}$$

$$\frac{1}{3} \left(\frac{1}{2} \int 3(\cos(c+dx)a+a)(2(4A+B)a^3+5(A+2B)\cos(c+dx)a^3)\sec(c+dx)dx - \frac{(3A-8B)\sin(c+dx)(a^4)}{2d} \right. \\ \left. \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{3}{2} \int (\cos(c+dx)a+a)(2(4A+B)a^3+5(A+2B)\cos(c+dx)a^3)\sec(c+dx)dx - \frac{(3A-8B)\sin(c+dx)(a^4)}{2d} \right. \\ \left. \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(2(4A+B)a^3+5(A+2B)\sin(c+dx+\frac{\pi}{2})a^3)}{\sin(c+dx+\frac{\pi}{2})} dx - \frac{(3A-8B)\sin(c+dx)(a^4)}{2d} \right. \\ \left. \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d} \right)$$

↓ 3447

$$\frac{1}{3} \left(\frac{3}{2} \int (5(A+2B)\cos^2(c+dx)a^4+2(4A+B)a^4+(2(4A+B)a^4+5(A+2B)a^4)\cos(c+dx))\sec(c+dx)dx - \right. \\ \left. \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{5(A+2B)\sin(c+dx+\frac{\pi}{2})^2a^4+2(4A+B)a^4+(2(4A+B)a^4+5(A+2B)a^4)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx - \right. \\ \left. \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d} \right)$$

↓ 3502

$$\frac{1}{3} \left(\frac{3}{2} \left(\int (2(4A+B)a^4+(13A+12B)\cos(c+dx)a^4)\sec(c+dx)dx + \frac{5a^4(A+2B)\sin(c+dx)}{d} \right) - \frac{(3A-8B)\sin(c+dx)(a^4)}{2d} \right. \\ \left. \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \left(\int \frac{2(4A+B)a^4 + (13A+12B)\sin(c+dx+\frac{\pi}{2})a^4}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{5a^4(A+2B)\sin(c+dx)}{d} \right) - \frac{(3A-8B)\sin(c+dx)}{2d} \right) + \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d}$$

↓ 3214

$$\frac{1}{3} \left(\frac{3}{2} \left(2a^4(4A+B) \int \sec(c+dx) dx + \frac{5a^4(A+2B)\sin(c+dx)}{d} + a^4x(13A+12B) \right) - \frac{(3A-8B)\sin(c+dx)}{2d} \right) + \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} \left(2a^4(4A+B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{5a^4(A+2B)\sin(c+dx)}{d} + a^4x(13A+12B) \right) - \frac{(3A-8B)\sin(c+dx)}{2d} \right) + \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d}$$

↓ 4257

$$\frac{1}{3} \left(\frac{3}{2} \left(\frac{2a^4(4A+B)\operatorname{arctanh}(\sin(c+dx))}{d} + \frac{5a^4(A+2B)\sin(c+dx)}{d} + a^4x(13A+12B) \right) - \frac{(3A-8B)\sin(c+dx)}{2d} \right) + \frac{(3A-B)\sin(c+dx)(a^2\cos(c+dx)+a^2)^2}{3d} + \frac{aA\tan(c+dx)(a\cos(c+dx)+a)^3}{d}$$

input `Int[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `-1/3*((3*A - B)*(a^2 + a^2*Cos[c + d*x])^2*Sin[c + d*x])/d + (-1/2*((3*A - 8*B)*(a^4 + a^4*Cos[c + d*x])*Sin[c + d*x])/d + (3*(a^4*(13*A + 12*B)*x + (2*a^4*(4*A + B)*ArcTanh[Sin[c + d*x]]))/d + (5*a^4*(A + 2*B)*Sin[c + d*x])/d))/2)/3 + (a*A*(a + a*Cos[c + d*x])^3*Tan[c + d*x])/d`

3.32.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos
[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.32.4 Maple [A] (verified)

Time = 3.51 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.93

method	result
parallelrisch	$\frac{(-32 \cos(dx+c) \left(A + \frac{B}{4}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 32 \cos(dx+c) \left(A + \frac{B}{4}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + \left(16A + \frac{82B}{3}\right) \sin(2dx+2c) + \dots}{8d \cos(dx+c)}$
parts	$\frac{a^4 A \tan(dx+c)}{d} + \frac{(a^4 A + 4B a^4) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{(4a^4 A + B a^4) \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(4a^4 A + B a^4) \sin(3dx+3c)}{3d}$
derivativedivides	$a^4 A \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + \frac{B a^4 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + 4a^4 A \sin(dx+c) + 4B a^4 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)$
default	$a^4 A \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + \frac{B a^4 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + 4a^4 A \sin(dx+c) + 4B a^4 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)$
risch	$\frac{13a^4 x A}{2} + 6a^4 B x + \frac{27ie^{-i(dx+c)} B a^4}{8d} + \frac{ie^{-2i(dx+c)} B a^4}{2d} - \frac{ie^{2i(dx+c)} a^4 A}{8d} - \frac{27ie^{i(dx+c)} B a^4}{8d} + \frac{2ie^{-i(dx+c)} a^4 A}{d}$
norman	$\frac{\left(-\frac{13}{2} a^4 A - 6B a^4\right) x + \left(-\frac{65}{2} a^4 A - 30B a^4\right) x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(\frac{13}{2} a^4 A + 6B a^4\right) x \left(\tan^{12}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(\frac{65}{2} a^4 A + 30B a^4\right) x \left(\tan^{20}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d \cos(dx+c)}$

```
input int((a+cos(d*x+c)*a)^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBO
SE)
```

```
output 1/8*(-32*cos(d*x+c)*(A+1/4*B)*ln(tan(1/2*d*x+1/2*c)-1)+32*cos(d*x+c)*(A+1/
4*B)*ln(tan(1/2*d*x+1/2*c)+1)+(16*A+82/3*B)*sin(2*d*x+2*c)+(A+4*B)*sin(3*d
*x+3*c)+1/3*sin(4*d*x+4*c)*B+52*(A+12/13*B)*x*d*cos(d*x+c)+9*(A+4/9*B)*sin
(d*x+c))*a^4/d/cos(d*x+c)
```

3.32. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.32.5 Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.00

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{3(13A + 12B)a^4 dx \cos(dx + c) + 3(4A + B)a^4 \cos(dx + c) \log(\sin(dx + c) + 1) - 3(4A + B)a^4 \cos(dx + c) \log(\sin(dx + c) - 1) + (2Ba^4 \cos(dx + c)^3 + 3(A + 4B)a^4 \cos(dx + c)^2 + 8(3A + 5B)a^4 \cos(dx + c) + 6Aa^4) \sin(dx + c)}{(d \cos(dx + c))}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/6*(3*(13*A + 12*B)*a^4*d*x*cos(d*x + c) + 3*(4*A + B)*a^4*cos(d*x + c)*log(sin(d*x + c) + 1) - 3*(4*A + B)*a^4*cos(d*x + c)*log(-sin(d*x + c) + 1) + (2*B*a^4*cos(d*x + c)^3 + 3*(A + 4*B)*a^4*cos(d*x + c)^2 + 8*(3*A + 5*B)*a^4*cos(d*x + c) + 6*A*a^4)*sin(d*x + c))/(d*cos(d*x + c))`

3.32.6 Sympy [F]

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= a^4 \left(\int A \sec^2(c + dx) dx + \int 4A \cos(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int 6A \cos^2(c + dx) \sec^2(c + dx) dx + \int 4A \cos^3(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int A \cos^4(c + dx) \sec^2(c + dx) dx + \int B \cos(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int 4B \cos^2(c + dx) \sec^2(c + dx) dx + \int 6B \cos^3(c + dx) \sec^2(c + dx) dx \right. \\ \left. + \int 4B \cos^4(c + dx) \sec^2(c + dx) dx + \int B \cos^5(c + dx) \sec^2(c + dx) dx \right)$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `a**4*(Integral(A*sec(c + d*x)**2, x) + Integral(4*A*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(6*A*cos(c + d*x)**2*sec(c + d*x)**2, x) + Integral(4*A*cos(c + d*x)**3*sec(c + d*x)**2, x) + Integral(A*cos(c + d*x)**4*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2, x) + Integral(4*B*cos(c + d*x)**2*sec(c + d*x)**2, x) + Integral(6*B*cos(c + d*x)**3*sec(c + d*x)**2, x) + Integral(4*B*cos(c + d*x)**4*sec(c + d*x)**2, x) + Integral(B*cos(c + d*x)**5*sec(c + d*x)**2, x))`

3.32.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.25

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{3(2dx + 2c + \sin(2dx + 2c))Aa^4 + 72(dx + c)Aa^4 - 4(\sin(dx + c)^3 - 3\sin(dx + c))Ba^4 + 12(2dx + 2c + \sin(2dx + 2c))Ba^4}{1}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `1/12*(3*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^4 + 72*(d*x + c)*A*a^4 - 4*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^4 + 12*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^4 + 48*(d*x + c)*B*a^4 + 24*A*a^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 6*B*a^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 48*A*a^4*sin(d*x + c) + 72*B*a^4*sin(d*x + c) + 12*A*a^4*tan(d*x + c))/d`

3.32.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.51

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$\frac{12Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)}{\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1} - 3(13Aa^4 + 12Ba^4)(dx + c) - 6(4Aa^4 + Ba^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) + 6(4Aa^4 + Ba^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right)$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

3.32. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$

output
$$\begin{aligned} & -1/6*(12*A*a^4*\tan(1/2*d*x + 1/2*c)/(\tan(1/2*d*x + 1/2*c)^2 - 1) - 3*(13*A \\ & *a^4 + 12*B*a^4)*(d*x + c) - 6*(4*A*a^4 + B*a^4)*\log(\text{abs}(\tan(1/2*d*x + 1/2 \\ & *c) + 1)) + 6*(4*A*a^4 + B*a^4)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(21 \\ & *A*a^4*\tan(1/2*d*x + 1/2*c)^5 + 30*B*a^4*\tan(1/2*d*x + 1/2*c)^5 + 48*A*a^4 \\ & *\tan(1/2*d*x + 1/2*c)^3 + 76*B*a^4*\tan(1/2*d*x + 1/2*c)^3 + 27*A*a^4*\tan(1 \\ & /2*d*x + 1/2*c) + 54*B*a^4*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 + \\ & 1)^3)/d \end{aligned}$$

3.32.9 Mupad [B] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.61

$$\begin{aligned} & \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx \\ & = \frac{4 A a^4 \sin(c + dx)}{d} + \frac{20 B a^4 \sin(c + dx)}{3 d} + \frac{13 A a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\ & + \frac{8 A a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{12 B a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\ & + \frac{2 B a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{A a^4 \sin(c + dx)}{d \cos(c + dx)} + \frac{B a^4 \cos(c + dx)^2 \sin(c + dx)}{3 d} \\ & + \frac{A a^4 \cos(c + dx) \sin(c + dx)}{2 d} + \frac{2 B a^4 \cos(c + dx) \sin(c + dx)}{d} \end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x)^2,x)`

output
$$\begin{aligned} & (4*A*a^4*\sin(c + d*x))/d + (20*B*a^4*\sin(c + d*x))/(3*d) + (13*A*a^4*\operatorname{atan} \\ & (\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (8*A*a^4*\operatorname{atanh}(\sin(c/2 + (d*x) \\ & /2)/\cos(c/2 + (d*x)/2)))/d + (12*B*a^4*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (\\ & d*x)/2)))/d + (2*B*a^4*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (\\ & A*a^4*\sin(c + d*x))/(d*\cos(c + d*x)) + (B*a^4*\cos(c + d*x)^2*\sin(c + d*x)) \\ & / (3*d) + (A*a^4*\cos(c + d*x)*\sin(c + d*x))/(2*d) + (2*B*a^4*\cos(c + d*x)*\sin \\ & (c + d*x))/d \end{aligned}$$

3.33 $\int (a+a \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.33.1	Optimal result	488
3.33.2	Mathematica [B] (verified)	489
3.33.3	Rubi [A] (verified)	490
3.33.4	Maple [A] (verified)	494
3.33.5	Fricas [A] (verification not implemented)	495
3.33.6	Sympy [F(-1)]	495
3.33.7	Maxima [A] (verification not implemented)	495
3.33.8	Giac [A] (verification not implemented)	496
3.33.9	Mupad [B] (verification not implemented)	497

3.33.1 Optimal result

Integrand size = 31, antiderivative size = 162

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{1}{2} a^4 (8A + 13B)x + \frac{a^4 (13A + 8B) \operatorname{arctanh}(\sin(c + dx))}{2d}$$

$$- \frac{5a^4 (A - B) \sin(c + dx)}{2d} - \frac{(6A + B) (a^4 + a^4 \cos(c + dx)) \sin(c + dx)}{2d}$$

$$+ \frac{(5A + 2B) (a^2 + a^2 \cos(c + dx))^2 \tan(c + dx)}{2d}$$

$$+ \frac{aA (a + a \cos(c + dx))^3 \sec(c + dx) \tan(c + dx)}{2d}$$

output

```
1/2*a^4*(8*A+13*B)*x+1/2*a^4*(13*A+8*B)*arctanh(sin(d*x+c))/d-5/2*a^4*(A-B)
)*sin(d*x+c)/d-1/2*(6*A+B)*(a^4+a^4*cos(d*x+c))*sin(d*x+c)/d+1/2*(5*A+2*B)
*(a^2+a^2*cos(d*x+c))^2*tan(d*x+c)/d+1/2*a*A*(a+a*cos(d*x+c))^3*sec(d*x+c)
*tan(d*x+c)/d
```

3.33.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 343 vs. $2(162) = 324$.

Time = 7.85 (sec) , antiderivative size = 343, normalized size of antiderivative = 2.12

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{1}{64} a^4 (1 + \cos(c + dx))^4 \sec^8\left(\frac{1}{2}(c + dx)\right) \left(2(8A + 13B)x \right. \\ \left. - \frac{2(13A + 8B) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)}{d} \right. \\ \left. + \frac{2(13A + 8B) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)}{d} + \frac{4(A + 4B) \cos(dx) \sin(c)}{d} \right. \\ \left. + \frac{B \cos(2dx) \sin(2c)}{d} + \frac{4(A + 4B) \cos(c) \sin(dx)}{d} + \frac{B \cos(2c) \sin(2dx)}{d} \right. \\ \left. + \frac{A}{d \left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)^2} \right. \\ \left. + \frac{4(4A + B) \sin\left(\frac{dx}{2}\right)}{d \left(\cos\left(\frac{c}{2}\right) - \sin\left(\frac{c}{2}\right)\right) \left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)} \right. \\ \left. - \frac{A}{d \left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)^2} \right. \\ \left. + \frac{4(4A + B) \sin\left(\frac{dx}{2}\right)}{d \left(\cos\left(\frac{c}{2}\right) + \sin\left(\frac{c}{2}\right)\right) \left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)} \right)$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a^4*(1 + Cos[c + d*x])^4*Sec[(c + d*x)/2]^8*(2*(8*A + 13*B)*x - (2*(13*A + 8*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]])/d + (2*(13*A + 8*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/d + (4*(A + 4*B)*Cos[d*x]*Sin[c])/d + (B*Cos[2*d*x]*Sin[2*c])/d + (4*(A + 4*B)*Cos[c]*Sin[d*x])/d + (B*Cos[2*c]*Sin[2*d*x])/d + A/(d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2) + (4*(4*A + B)*Sin[(d*x)/2])/(d*(Cos[c/2] - Sin[c/2])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) - A/(d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2) + (4*(4*A + B)*Sin[(d*x)/2])/(d*(Cos[c/2] + Sin[c/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])))/64`

3.33.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.95, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3454, 3042, 3454, 3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c+dx)(a \cos(c+dx) + a)^4(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c+dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3454} \\
 & \frac{1}{2} \int (\cos(c+dx)a + a)^3 (a(5A + 2B) - 2a(A - B) \cos(c+dx)) \sec^2(c+dx) dx + \\
 & \quad \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^3 (a(5A + 2B) - 2a(A - B) \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^2} dx + \\
 & \quad \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \\
 & \quad \downarrow \text{3454} \\
 & \frac{1}{2} \left(\int (\cos(c+dx)a + a)^2 (a^2(13A + 8B) - 2a^2(6A + B) \cos(c+dx)) \sec(c+dx) dx + \frac{(5A + 2B) \tan(c+dx) (a^2)}{d} \right. \\
 & \quad \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(\int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^2 (a^2(13A + 8B) - 2a^2(6A + B) \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})} dx + \frac{(5A + 2B) \tan(c+dx) (a^2)}{d} \right. \\
 & \quad \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right) \\
 & \quad \downarrow \text{3455}
 \end{aligned}$$

3.33. $\int (a + a \cos(c+dx))^4 (A + B \cos(c+dx)) \sec^3(c+dx) dx$

$$\frac{1}{2} \left(\frac{1}{2} \int 2(\cos(c+dx)a+a) (a^3(13A+8B) - 5a^3(A-B)\cos(c+dx)) \sec(c+dx) dx - \frac{(6A+B)\sin(c+dx)(a^4)}{d} \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right. \\ \left. \downarrow 27 \right.$$

$$\frac{1}{2} \left(\int (\cos(c+dx)a+a) (a^3(13A+8B) - 5a^3(A-B)\cos(c+dx)) \sec(c+dx) dx - \frac{(6A+B)\sin(c+dx)(a^4 \cos)}{d} \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{2} \left(\int \frac{(\sin(c+dx+\frac{\pi}{2})a+a) (a^3(13A+8B) - 5a^3(A-B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})} dx - \frac{(6A+B)\sin(c+dx)(a^4 \cos)}{d} \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right. \\ \left. \downarrow 3447 \right.$$

$$\frac{1}{2} \left(\int (-5(A-B)\cos^2(c+dx)a^4 + (13A+8B)a^4 + (a^4(13A+8B) - 5a^4(A-B))\cos(c+dx)) \sec(c+dx) dx - \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{2} \left(\int \frac{-5(A-B)\sin(c+dx+\frac{\pi}{2})^2 a^4 + (13A+8B)a^4 + (a^4(13A+8B) - 5a^4(A-B))\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx - \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right. \\ \left. \downarrow 3502 \right.$$

$$\frac{1}{2} \left(\int ((13A+8B)a^4 + (8A+13B)\cos(c+dx)a^4) \sec(c+dx) dx - \frac{5a^4(A-B)\sin(c+dx)}{d} - \frac{(6A+B)\sin(c+dx)}{d} \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a \cos(c+dx) + a)^3}{2d} \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{2} \left(\int \frac{(13A + 8B)a^4 + (8A + 13B) \sin(c + dx + \frac{\pi}{2}) a^4}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{5a^4(A - B) \sin(c + dx)}{d} - \frac{(6A + B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{d} \right) \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^3}{2d}$$

↓ 3214

$$\frac{1}{2} \left(a^4(13A + 8B) \int \sec(c + dx) dx - \frac{5a^4(A - B) \sin(c + dx)}{d} - \frac{(6A + B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{d} + a^4 \right) \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^3}{2d}$$

↓ 3042

$$\frac{1}{2} \left(a^4(13A + 8B) \int \csc(c + dx + \frac{\pi}{2}) dx - \frac{5a^4(A - B) \sin(c + dx)}{d} - \frac{(6A + B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{d} \right) \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^3}{2d}$$

↓ 4257

$$\frac{1}{2} \left(\frac{a^4(13A + 8B) \operatorname{arctanh}(\sin(c + dx))}{d} - \frac{5a^4(A - B) \sin(c + dx)}{d} - \frac{(6A + B) \sin(c + dx) (a^4 \cos(c + dx) + a^4)}{d} \right) \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^3}{2d}$$

input `Int[(a + a*cos[c + d*x])^4*(A + B*cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*(a + a*cos[c + d*x])^3*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (a^4*(8*A + 13*B)*x + (a^4*(13*A + 8*B)*ArcTanh[Sin[c + d*x]])/d - (5*a^4*(A - B)*Sin[c + d*x])/d - ((6*A + B)*(a^4 + a^4*cos[c + d*x])*Sin[c + d*x])/d + ((5*A + 2*B)*(a^2 + a^2*cos[c + d*x])^2*Tan[c + d*x])/d)/2`

3.33.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.33.4 Maple [A] (verified)

Time = 3.29 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.04

method	result
parallelrisch	$\frac{(-13(A + \frac{8B}{13})(1 + \cos(2dx + 2c)) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 1) + 13(A + \frac{8B}{13})(1 + \cos(2dx + 2c)) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 1) + 8(A + \frac{13B}{8}) \cos(2dx + 2c))}{2d(1 + \cos(2dx + 2c))}$
parts	$\frac{a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d} + \frac{(a^4 A + 4B a^4) \sin(dx+c)}{d} + \frac{(4a^4 A + B a^4) \tan(dx+c)}{d} + \frac{(4a^4 A + 13B a^4) \cos(2dx+2c)}{2d}$
derivativedivides	$a^4 A \sin(dx+c) + B a^4 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 4a^4 A(dx+c) + 4B a^4 \sin(dx+c) + 6a^4 A \ln(\sec(dx+c) + \tan(dx+c)) + \frac{(4a^4 A + 13B a^4) \cos(2dx+2c)}{2d}$
default	$a^4 A \sin(dx+c) + B a^4 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 4a^4 A(dx+c) + 4B a^4 \sin(dx+c) + 6a^4 A \ln(\sec(dx+c) + \tan(dx+c)) + \frac{(4a^4 A + 13B a^4) \cos(2dx+2c)}{2d}$
risch	$4a^4 x A + \frac{13a^4 B x}{2} - \frac{ie^{2i(dx+c)} B a^4}{8d} - \frac{ie^{i(dx+c)} a^4 A}{2d} - \frac{2ie^{i(dx+c)} B a^4}{d} + \frac{ie^{-i(dx+c)} a^4 A}{2d} + \frac{2ie^{-i(dx+c)} B a^4}{d}$
norman	$\frac{(4a^4 A + \frac{13}{2} B a^4) x + (-20a^4 A - \frac{65}{2} B a^4) x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + (-20a^4 A - \frac{65}{2} B a^4) x \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + (4a^4 A + \frac{13}{2} B a^4) x \left(\tan^{10}\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{2d(1 + \cos(2dx + 2c))}$

```
input int((a+cos(d*x+c)*a)^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBO
SE)
```

```
output 1/2*(-13*(A+8/13*B)*(1+cos(2*d*x+2*c))*ln(tan(1/2*d*x+1/2*c)-1)+13*(A+8/13
*B)*(1+cos(2*d*x+2*c))*ln(tan(1/2*d*x+1/2*c)+1)+8*(A+13/8*B)*x*d*cos(2*d*x
+2*c)+(8*A+5/2*B)*sin(2*d*x+2*c)+(A+4*B)*sin(3*d*x+3*c)+1/4*sin(4*d*x+4*c)
*B+(3*A+4*B)*sin(d*x+c)+8*(A+13/8*B)*x*d)*a^4/d/(1+cos(2*d*x+2*c))
```

3.33. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.33.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.96

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2(8A + 13B)a^4 dx \cos(dx + c)^2 + (13A + 8B)a^4 \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (13A + 8B)a^4 \cos(dx + c)^2 \log(-\sin(dx + c) + 1) + 2(Ba^4 \cos(dx + c)^3 + 2(A + 4B)a^4 \cos(dx + c)^2 + 2(4A + B)a^4 \cos(dx + c) + Aa^4) \sin(dx + c)}{(d \cos(dx + c))^2}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fracas")`

output `1/4*(2*(8*A + 13*B)*a^4*d*x*cos(d*x + c)^2 + (13*A + 8*B)*a^4*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (13*A + 8*B)*a^4*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(B*a^4*cos(d*x + c)^3 + 2*(A + 4*B)*a^4*cos(d*x + c)^2 + 2*(4*A + B)*a^4*cos(d*x + c) + A*a^4)*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.33.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Timed out`

3.33.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.23

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{16(dx + c)Aa^4 + (2dx + 2c + \sin(2dx + 2c))Ba^4 + 24(dx + c)Ba^4 - Aa^4 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx + c) + 1) - \log(-\sin(dx + c) + 1) \right)}{(d \cos(dx + c))^2}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `1/4*(16*(d*x + c)*A*a^4 + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^4 + 24*(d*x + c)*B*a^4 - A*a^4*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 12*A*a^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 8*B*a^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 4*A*a^4*sin(d*x + c) + 16*B*a^4*sin(d*x + c) + 16*A*a^4*tan(d*x + c) + 4*B*a^4*tan(d*x + c))/d`

3.33.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.42

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(8Aa^4 + 13Ba^4)(dx + c) + (13Aa^4 + 8Ba^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - (13Aa^4 + 8Ba^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right)}{d}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output `1/2*((8*A*a^4 + 13*B*a^4)*(d*x + c) + (13*A*a^4 + 8*B*a^4)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (13*A*a^4 + 8*B*a^4)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(5*A*a^4*tan(1/2*d*x + 1/2*c)^7 - 5*B*a^4*tan(1/2*d*x + 1/2*c)^7 + 7*A*a^4*tan(1/2*d*x + 1/2*c)^5 + 7*B*a^4*tan(1/2*d*x + 1/2*c)^5 - 9*A*a^4*tan(1/2*d*x + 1/2*c)^3 + 9*B*a^4*tan(1/2*d*x + 1/2*c)^3 - 11*A*a^4*tan(1/2*d*x + 1/2*c) - 11*B*a^4*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^4 - 1)^2)/d`

3.33.9 Mupad [B] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.50

$$\begin{aligned}
& \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx \\
&= \frac{A a^4 \sin(c + dx)}{d} + \frac{4 B a^4 \sin(c + dx)}{d} + \frac{8 A a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{13 A a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{13 B a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{8 B a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{4 A a^4 \sin(c + dx)}{d \cos(c + dx)} + \frac{A a^4 \sin(c + dx)}{2 d \cos(c + dx)^2} \\
&+ \frac{B a^4 \sin(c + dx)}{d \cos(c + dx)} + \frac{B a^4 \cos(c + dx) \sin(c + dx)}{2 d}
\end{aligned}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x)^3,x)
```

```
output (A*a^4*sin(c + d*x))/d + (4*B*a^4*sin(c + d*x))/d + (8*A*a^4*atan(sin(c/2 +
+ (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (13*A*a^4*atanh(sin(c/2 + (d*x)/2)/cos
(c/2 + (d*x)/2)))/d + (13*B*a^4*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)
))/d + (8*B*a^4*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (4*A*a^4
*sin(c + d*x))/(d*cos(c + d*x)) + (A*a^4*sin(c + d*x))/(2*d*cos(c + d*x)^2
) + (B*a^4*sin(c + d*x))/(d*cos(c + d*x)) + (B*a^4*cos(c + d*x)*sin(c + d*
x))/(2*d)
```


3.34 $\int (a+a \cos(c+dx))^4(A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.34.1	Optimal result	498
3.34.2	Mathematica [B] (verified)	499
3.34.3	Rubi [A] (verified)	500
3.34.4	Maple [A] (verified)	504
3.34.5	Fricas [A] (verification not implemented)	504
3.34.6	Sympy [F(-1)]	505
3.34.7	Maxima [A] (verification not implemented)	505
3.34.8	Giac [A] (verification not implemented)	506
3.34.9	Mupad [B] (verification not implemented)	506

3.34.1 Optimal result

Integrand size = 31, antiderivative size = 165

$$\begin{aligned} & \int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= a^4(A + 4B)x + \frac{a^4(12A + 13B)\operatorname{arctanh}(\sin(c + dx))}{2d} \\ & \quad - \frac{5a^4(2A + B) \sin(c + dx)}{2d} + \frac{(11A + 9B)(a^4 + a^4 \cos(c + dx)) \tan(c + dx)}{3d} \\ & \quad + \frac{(2A + B)(a^2 + a^2 \cos(c + dx))^2 \sec(c + dx) \tan(c + dx)}{2d} \\ & \quad + \frac{aA(a + a \cos(c + dx))^3 \sec^2(c + dx) \tan(c + dx)}{3d} \end{aligned}$$

output

```
a^4*(A+4*B)*x+1/2*a^4*(12*A+13*B)*arctanh(sin(d*x+c))/d-5/2*a^4*(2*A+B)*sin(d*x+c)/d+1/3*(11*A+9*B)*(a^4+a^4*cos(d*x+c))*tan(d*x+c)/d+1/2*(2*A+B)*(a^2+a^2*cos(d*x+c))^2*sec(d*x+c)*tan(d*x+c)/d+1/3*a*A*(a+a*cos(d*x+c))^3*sec(d*x+c)^2*tan(d*x+c)/d
```

3.34.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 380 vs. $2(165) = 330$.

Time = 9.28 (sec) , antiderivative size = 380, normalized size of antiderivative = 2.30

$$\begin{aligned} & \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= a^4 \left(\frac{(A + 4B)(c + dx)}{d} + \frac{(-12A - 13B) \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{2d} \right. \\ & \quad \left. + \frac{(12A + 13B) \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{2d} \right. \\ & \quad + \frac{13A + 3B}{12d (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^2} + \frac{A \sin(\frac{1}{2}(c + dx))}{6d (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^3} \\ & \quad + \frac{A \sin(\frac{1}{2}(c + dx))}{6d (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^3} + \frac{-13A - 3B}{12d (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^2} \\ & \quad \left. + \frac{4(5A \sin(\frac{1}{2}(c + dx)) + 3B \sin(\frac{1}{2}(c + dx)))}{3d (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))} \right. \\ & \quad \left. + \frac{4(5A \sin(\frac{1}{2}(c + dx)) + 3B \sin(\frac{1}{2}(c + dx)))}{3d (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))} + \frac{B \sin(c + dx)}{d} \right) \end{aligned}$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `a^4*(((A + 4*B)*(c + d*x))/d + ((-12*A - 13*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]])/(2*d) + ((12*A + 13*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/(2*d) + (13*A + 3*B)/(12*d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2) + (A*Sin[(c + d*x)/2])/(6*d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^3) + (A*Sin[(c + d*x)/2])/(6*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^3) + (-13*A - 3*B)/(12*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2) + (4*(5*A*Sin[(c + d*x)/2] + 3*B*Sin[(c + d*x)/2]))/(3*d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) + (4*(5*A*Sin[(c + d*x)/2] + 3*B*Sin[(c + d*x)/2]))/(3*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])) + (B*Sin[c + d*x])/d)`

3.34.3 Rubi [A] (verified)

Time = 1.40 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3454, 3042, 3454, 3042, 3454, 27, 3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c+dx)(a \cos(c+dx) + a)^4(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c+dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3454} \\
 & \frac{1}{3} \int (\cos(c+dx)a + a)^3 (3a(2A+B) - a(A-3B)\cos(c+dx)) \sec^3(c+dx) dx + \\
 & \quad \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^3 (3a(2A+B) - a(A-3B)\sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^3} dx + \\
 & \quad \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \\
 & \quad \downarrow \text{3454} \\
 & \frac{1}{3} \left(\frac{1}{2} \int (\cos(c+dx)a + a)^2 (2a^2(11A+9B) - a^2(8A-3B)\cos(c+dx)) \sec^2(c+dx) dx + \frac{3(2A+B)\tan(c+dx)}{3d} \right. \\
 & \quad \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{(\sin(c+dx + \frac{\pi}{2})a + a)^2 (2a^2(11A+9B) - a^2(8A-3B)\sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})^2} dx + \frac{3(2A+B)\tan(c+dx)}{3d} \right. \\
 & \quad \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\
 & \quad \downarrow \text{3454}
 \end{aligned}$$

3.34. $\int (a + a \cos(c+dx))^4 (A + B \cos(c+dx)) \sec^4(c+dx) dx$

$$\frac{1}{3} \left(\frac{1}{2} \left(\int 3(\cos(c+dx)a+a)(a^3(12A+13B)-5a^3(2A+B)\cos(c+dx))\sec(c+dx)dx + \frac{2(11A+9B)\tan(c+dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\ \downarrow 27$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int (\cos(c+dx)a+a)(a^3(12A+13B)-5a^3(2A+B)\cos(c+dx))\sec(c+dx)dx + \frac{2(11A+9B)\tan(c+dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\ \downarrow 3042$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a^3(12A+13B)-5a^3(2A+B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2(11A+9B)\tan(c+dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\ \downarrow 3447$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int (-5(2A+B)\cos^2(c+dx)a^4 + (12A+13B)a^4 + (a^4(12A+13B)-5a^4(2A+B))\cos(c+dx))\sec(c+dx)dx + \frac{2(11A+9B)\tan(c+dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\ \downarrow 3042$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{-5(2A+B)\sin(c+dx+\frac{\pi}{2})^2 a^4 + (12A+13B)a^4 + (a^4(12A+13B)-5a^4(2A+B))\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2(11A+9B)\tan(c+dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\ \downarrow 3502$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(\int ((12A+13B)a^4 + 2(A+4B)\cos(c+dx)a^4)\sec(c+dx)dx - \frac{5a^4(2A+B)\sin(c+dx)}{d} \right) + \frac{2(11A+9B)\tan(c+dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec^2(c+dx)(a \cos(c+dx) + a)^3}{3d} \right) \\ \downarrow 3042$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(\int \frac{(12A + 13B)a^4 + 2(A + 4B) \sin(c + dx + \frac{\pi}{2}) a^4}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{5a^4(2A + B) \sin(c + dx)}{d} \right) + \frac{2(11A + 9B) \tan(c + dx)}{aA \tan(c + dx) \sec^2(c + dx)(a \cos(c + dx) + a)^3} \right) \right) + \frac{2(11A + 9B) \tan(c + dx)}{3d}$$

↓ 3214

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(a^4(12A + 13B) \int \sec(c + dx) dx - \frac{5a^4(2A + B) \sin(c + dx)}{d} + 2a^4x(A + 4B) \right) + \frac{2(11A + 9B) \tan(c + dx)}{aA \tan(c + dx) \sec^2(c + dx)(a \cos(c + dx) + a)^3} \right) \right) + \frac{2(11A + 9B) \tan(c + dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(a^4(12A + 13B) \int \csc(c + dx + \frac{\pi}{2}) dx - \frac{5a^4(2A + B) \sin(c + dx)}{d} + 2a^4x(A + 4B) \right) + \frac{2(11A + 9B) \tan(c + dx)}{aA \tan(c + dx) \sec^2(c + dx)(a \cos(c + dx) + a)^3} \right) \right) + \frac{2(11A + 9B) \tan(c + dx)}{3d}$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(\frac{a^4(12A + 13B) \operatorname{arctanh}(\sin(c + dx))}{d} - \frac{5a^4(2A + B) \sin(c + dx)}{d} + 2a^4x(A + 4B) \right) + \frac{2(11A + 9B) \tan(c + dx)}{aA \tan(c + dx) \sec^2(c + dx)(a \cos(c + dx) + a)^3} \right) \right) + \frac{2(11A + 9B) \tan(c + dx)}{3d}$$

input `Int[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*(a + a*Cos[c + d*x])^3*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((3*(2*A + B)*(a^2 + a^2*Cos[c + d*x])^2*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (3*(2*a^4*(A + 4*B)*x + (a^4*(12*A + 13*B)*ArcTanh[Sin[c + d*x]])/d - (5*a^4*(2*A + B)*Sin[c + d*x])/d) + (2*(11*A + 9*B)*(a^4 + a^4*Cos[c + d*x])*Tan[c + d*x])/d)/2)/3`

3.34.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]) / ((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.) * ((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) * ((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m * (A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_) * ((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) * ((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1) * ((c + d*Sin[e + f*x])^(n + 1) / (d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1) * (c + d*Sin[e + f*x])^(n + 1) * Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.) * ((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x] * ((a + b*Sin[e + f*x])^(m + 1) / (b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m * Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.34.4 Maple [A] (verified)

Time = 4.06 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.00

method	result
parts	$-\frac{a^4 A \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3}\right) \tan(dx+c)}{d} + \frac{(a^4 A + 4B a^4)(dx+c)}{d} + \frac{(4a^4 A + B a^4) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \ln(\sec(dx+c))\right)}{d}$
parallelrisch	$4 \left(-\frac{9 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c)\right) \left(A + \frac{13B}{12}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2} + \frac{9 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c)\right) \left(A + \frac{13B}{12}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2} \right) \frac{1}{d(\cos(3dx+c))}$
derivativedivides	$a^4 A(dx+c) + B a^4 \sin(dx+c) + 4a^4 A \ln(\sec(dx+c) + \tan(dx+c)) + 4B a^4(dx+c) + 6a^4 A \tan(dx+c) + 6B a^4 \ln(\sec(dx+c) + \tan(dx+c))$
default	$a^4 A(dx+c) + B a^4 \sin(dx+c) + 4a^4 A \ln(\sec(dx+c) + \tan(dx+c)) + 4B a^4(dx+c) + 6a^4 A \tan(dx+c) + 6B a^4 \ln(\sec(dx+c) + \tan(dx+c))$
risch	$a^4 x A + 4a^4 B x - \frac{ie^{i(dx+c)} B a^4}{2d} + \frac{ie^{-i(dx+c)} B a^4}{2d} - \frac{ia^4 (12A e^{5i(dx+c)} + 3B e^{5i(dx+c)} - 36A e^{4i(dx+c)} - 24B e^{3i(dx+c)})}{2d}$
norman	$\frac{(-a^4 A - 4B a^4)x + (-6a^4 A - 24B a^4)x \left(\tan^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-2a^4 A - 8B a^4)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-2a^4 A - 8B a^4)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d}$

```
input int((a+cos(d*x+c)*a)^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBOSE)
```

```
output -a^4*A/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(A*a^4+4*B*a^4)/d*(d*x+c)+(4*A*a^4+B*a^4)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(4*A*a^4+6*B*a^4)/d*ln(sec(d*x+c)+tan(d*x+c))+(6*A*a^4+4*B*a^4)/d*tan(d*x+c)+B*a^4/d*sin(d*x+c)
```

3.34.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.96

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12 (A + 4 B) a^4 dx \cos(dx + c)^3 + 3 (12 A + 13 B) a^4 \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3 (12 A + 13 B) a^4 \cos(dx + c)^3 \log(\sin(dx + c) - 1)}{d}$$

```
input integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")
```

3.34. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

output $1/12*(12*(A + 4*B)*a^4*d*x*cos(d*x + c)^3 + 3*(12*A + 13*B)*a^4*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*(12*A + 13*B)*a^4*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(6*B*a^4*cos(d*x + c)^3 + 8*(5*A + 3*B)*a^4*cos(d*x + c)^2 + 3*(4*A + B)*a^4*cos(d*x + c) + 2*A*a^4)*sin(d*x + c))/(d*cos(d*x + c)^3)$

3.34.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output Timed out

3.34.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.42

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{4 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^4 + 12 (dx + c) Aa^4 + 48 (dx + c) Ba^4 - 12 Aa^4 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c)) \right)}{d}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output $1/12*(4*(\tan(d*x + c)^3 + 3*\tan(d*x + c))*A*a^4 + 12*(d*x + c)*A*a^4 + 48*(d*x + c)*B*a^4 - 12*A*a^4*(2*\sin(d*x + c)/(\sin(d*x + c)^2 - 1) - \log(\sin(d*x + c) + 1) + \log(\sin(d*x + c) - 1)) - 3*B*a^4*(2*\sin(d*x + c)/(\sin(d*x + c)^2 - 1) - \log(\sin(d*x + c) + 1) + \log(\sin(d*x + c) - 1)) + 24*A*a^4*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 36*B*a^4*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 12*B*a^4*\sin(d*x + c) + 72*A*a^4*\tan(d*x + c) + 48*B*a^4*\tan(d*x + c))/d$

3.34.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.38

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12 B a^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1} + 6 (A a^4 + 4 B a^4) (dx + c) + 3 (12 A a^4 + 13 B a^4) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - 3 (12 A a^4 + 13 B a^4) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|) + \frac{20 A a^4 \sin(c + dx)}{3 d \cos(c + dx)} + \frac{2 A a^4 \sin(c + dx)}{d \cos(c + dx)^2} + \frac{A a^4 \sin(c + dx)}{3 d \cos(c + dx)^3} + \frac{4 B a^4 \sin(c + dx)}{d \cos(c + dx)} + \frac{B a^4 \sin(c + dx)}{2 d \cos(c + dx)^2}$$

```
input integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")
```

```
output 1/6*(12*B*a^4*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 1) + 6*(A*a^4 + 4*B*a^4)*(d*x + c) + 3*(12*A*a^4 + 13*B*a^4)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*(12*A*a^4 + 13*B*a^4)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(30*A*a^4*tan(1/2*d*x + 1/2*c)^5 + 21*B*a^4*tan(1/2*d*x + 1/2*c)^5 - 76*A*a^4*tan(1/2*d*x + 1/2*c)^3 - 48*B*a^4*tan(1/2*d*x + 1/2*c)^3 + 54*A*a^4*tan(1/2*d*x + 1/2*c) + 27*B*a^4*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^3)/d
```

3.34.9 Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.54

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{B a^4 \sin(c + dx)}{d} + \frac{2 A a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{12 A a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{8 B a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{13 B a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{20 A a^4 \sin(c + dx)}{3 d \cos(c + dx)}$$

$$+ \frac{2 A a^4 \sin(c + dx)}{d \cos(c + dx)^2} + \frac{A a^4 \sin(c + dx)}{3 d \cos(c + dx)^3} + \frac{4 B a^4 \sin(c + dx)}{d \cos(c + dx)} + \frac{B a^4 \sin(c + dx)}{2 d \cos(c + dx)^2}$$

```
input int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x)^4,x
```

output $(B*a^4*\sin(c + d*x))/d + (2*A*a^4*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (12*A*a^4*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (8*B*a^4*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (13*B*a^4*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (20*A*a^4*\sin(c + d*x))/(3*d*\cos(c + d*x)) + (2*A*a^4*\sin(c + d*x))/(d*\cos(c + d*x)^2) + (A*a^4*\sin(c + d*x))/(3*d*\cos(c + d*x)^3) + (4*B*a^4*\sin(c + d*x))/(d*\cos(c + d*x)) + (B*a^4*\sin(c + d*x))/(2*d*\cos(c + d*x)^2)$

3.35 $\int (a+a \cos(c+dx))^4(A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.35.1	Optimal result	508
3.35.2	Mathematica [A] (verified)	509
3.35.3	Rubi [A] (verified)	509
3.35.4	Maple [A] (verified)	513
3.35.5	Fricas [A] (verification not implemented)	514
3.35.6	Sympy [F(-1)]	514
3.35.7	Maxima [A] (verification not implemented)	515
3.35.8	Giac [A] (verification not implemented)	515
3.35.9	Mupad [B] (verification not implemented)	516

3.35.1 Optimal result

Integrand size = 31, antiderivative size = 173

$$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= a^4 Bx + \frac{a^4(35A + 48B)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{5a^4(7A + 8B) \tan(c + dx)}{8d}$$

$$+ \frac{(35A + 32B)(a^4 + a^4 \cos(c + dx)) \sec(c + dx) \tan(c + dx)}{24d}$$

$$+ \frac{(7A + 4B)(a^2 + a^2 \cos(c + dx))^2 \sec^2(c + dx) \tan(c + dx)}{12d}$$

$$+ \frac{aA(a + a \cos(c + dx))^3 \sec^3(c + dx) \tan(c + dx)}{4d}$$

output

```
a^4*B*x+1/8*a^4*(35*A+48*B)*arctanh(sin(d*x+c))/d+5/8*a^4*(7*A+8*B)*tan(d*x+c)/d+1/24*(35*A+32*B)*(a^4+a^4*cos(d*x+c))*sec(d*x+c)*tan(d*x+c)/d+1/12*(7*A+4*B)*(a^2+a^2*cos(d*x+c))^2*sec(d*x+c)^2*tan(d*x+c)/d+1/4*a*A*(a+a*cos(d*x+c))^3*sec(d*x+c)^3*tan(d*x+c)/d
```

3.35.2 Mathematica [A] (verified)

Time = 3.68 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.03

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= a^4 B x + \frac{35a^4 A \operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{6a^4 B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{8a^4 A \tan(c + dx)}{d}$$

$$+ \frac{7a^4 B \tan(c + dx)}{d} + \frac{27a^4 A \sec(c + dx) \tan(c + dx)}{8d} + \frac{2a^4 B \sec(c + dx) \tan(c + dx)}{d}$$

$$+ \frac{a^4 A \sec^3(c + dx) \tan(c + dx)}{4d} + \frac{4a^4 A \tan^3(c + dx)}{3d} + \frac{a^4 B \tan^3(c + dx)}{3d}$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `a^4*B*x + (35*a^4*A*ArcTanh[Sin[c + d*x]])/(8*d) + (6*a^4*B*ArcTanh[Sin[c + d*x]])/d + (8*a^4*A*Tan[c + d*x])/d + (7*a^4*B*Tan[c + d*x])/d + (27*a^4*A*Sec[c + d*x]*Tan[c + d*x])/(8*d) + (2*a^4*B*Sec[c + d*x]*Tan[c + d*x])/d + (a^4*A*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (4*a^4*A*Tan[c + d*x]^3)/(3*d) + (a^4*B*Tan[c + d*x]^3)/(3*d)`

3.35.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3454, 3042, 3454, 3042, 3454, 27, 3042, 3447, 3042, 3500, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a \cos(c + dx) + a)^4 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3454}$$

$$\frac{1}{4} \int (\cos(c + dx)a + a)^3 (a(7A + 4B) + 4aB \cos(c + dx)) \sec^4(c + dx) dx + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^3}{4d}$$

↓ 3042

$$\frac{1}{4} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^3 (a(7A + 4B) + 4aB \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^3}{4d}$$

↓ 3454

$$\frac{1}{4} \left(\frac{1}{3} \int (\cos(c + dx)a + a)^2 ((35A + 32B)a^2 + 12B \cos(c + dx)a^2) \sec^3(c + dx) dx + \frac{(7A + 4B) \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^3}{4d} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 ((35A + 32B)a^2 + 12B \sin(c + dx + \frac{\pi}{2})a^2)}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{(7A + 4B) \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^3}{4d} \right)$$

↓ 3454

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int 3(\cos(c + dx)a + a) (5(7A + 8B)a^3 + 8B \cos(c + dx)a^3) \sec^2(c + dx) dx + \frac{(35A + 32B) \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^3}{4d} \right) \right)$$

↓ 27

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int (\cos(c + dx)a + a) (5(7A + 8B)a^3 + 8B \cos(c + dx)a^3) \sec^2(c + dx) dx + \frac{(35A + 32B) \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^3}{4d} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(5(7A+8B)a^3+8B\sin(c+dx+\frac{\pi}{2})a^3)}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{(35A+32B)\tan(c+dx)\sec(c+dx)}{aA\tan(c+dx)\sec^3(c+dx)(a\cos(c+dx)+a)^3} \right) \right)$$

↓ 3447

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int (8B\cos^2(c+dx)a^4+5(7A+8B)a^4+(8Ba^4+5(7A+8B)a^4)\cos(c+dx))\sec^2(c+dx)dx + \frac{(35A+32B)\tan(c+dx)\sec(c+dx)}{aA\tan(c+dx)\sec^3(c+dx)(a\cos(c+dx)+a)^3} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \int \frac{8B\sin(c+dx+\frac{\pi}{2})^2a^4+5(7A+8B)a^4+(8Ba^4+5(7A+8B)a^4)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{(35A+32B)\tan(c+dx)\sec(c+dx)}{aA\tan(c+dx)\sec^3(c+dx)(a\cos(c+dx)+a)^3} \right) \right)$$

↓ 3500

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(\int ((35A+48B)a^4+8B\cos(c+dx)a^4)\sec(c+dx)dx + \frac{5a^4(7A+8B)\tan(c+dx)}{d} \right) + \frac{(35A+32B)\tan(c+dx)\sec(c+dx)}{aA\tan(c+dx)\sec^3(c+dx)(a\cos(c+dx)+a)^3} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(\int \frac{(35A+48B)a^4+8B\sin(c+dx+\frac{\pi}{2})a^4}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{5a^4(7A+8B)\tan(c+dx)}{d} \right) + \frac{(35A+32B)\tan(c+dx)\sec(c+dx)}{aA\tan(c+dx)\sec^3(c+dx)(a\cos(c+dx)+a)^3} \right) \right)$$

↓ 3214

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(a^4(35A+48B) \int \sec(c+dx)dx + \frac{5a^4(7A+8B)\tan(c+dx)}{d} + 8a^4Bx \right) + \frac{(35A+32B)\tan(c+dx)\sec(c+dx)}{aA\tan(c+dx)\sec^3(c+dx)(a\cos(c+dx)+a)^3} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(a^4(35A + 48B) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{5a^4(7A + 8B) \tan(c + dx)}{d} + 8a^4Bx \right) + \frac{(35A + 32B) \tan(c + dx)}{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^3} \right) \right)$$

↓ 4257

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{3}{2} \left(\frac{a^4(35A + 48B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{5a^4(7A + 8B) \tan(c + dx)}{d} + 8a^4Bx \right) + \frac{(35A + 32B) \tan(c + dx)}{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^3} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*A*(a + a*Cos[c + d*x])^3*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (((7*A + 4*B)*(a^2 + a^2*Cos[c + d*x])^2*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + (((35*A + 32*B)*(a^4 + a^4*Cos[c + d*x])*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (3*(8*a^4*B*x + (a^4*(35*A + 48*B)*ArcTanh[Sin[c + d*x]]))/d + (5*a^4*(7*A + 8*B)*Tan[c + d*x])/d))/2)/3)/4`

3.35.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3500 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2
- a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.35.4 Maple [A] (verified)

Time = 4.78 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.17

method	result
parts	$a^4 A \left(- \left(\frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right) + \frac{(a^4 A + 4B a^4) \ln(\sec(dx+c) + \tan(dx+c))}{d}$
parallelrisch	$56a^4 \left(- \frac{15 \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) \left(A + \frac{48B}{35} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + \frac{15 \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) \left(A + \frac{48B}{35} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{16} \right)$
derivativedivides	$a^4 A \ln(\sec(dx+c) + \tan(dx+c)) + B a^4 (dx+c) + 4a^4 A \tan(dx+c) + 4B a^4 \ln(\sec(dx+c) + \tan(dx+c)) + 6a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)$
default	$a^4 A \ln(\sec(dx+c) + \tan(dx+c)) + B a^4 (dx+c) + 4a^4 A \tan(dx+c) + 4B a^4 \ln(\sec(dx+c) + \tan(dx+c)) + 6a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)$
risch	$a^4 B x - \frac{ia^4 (81A e^{7i(dx+c)} + 48B e^{7i(dx+c)} - 96A e^{6i(dx+c)} - 144B e^{6i(dx+c)} + 105A e^{5i(dx+c)} + 48B e^{5i(dx+c)} - 480A e^{4i(dx+c)} - 144B e^{4i(dx+c)} + 105A e^{3i(dx+c)} + 48B e^{3i(dx+c)} - 96A e^{2i(dx+c)} - 144B e^{2i(dx+c)} + 105A e^{i(dx+c)} + 48B e^{i(dx+c)} - 480A)}{16}$

3.35. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

input `int((a+cos(d*x+c))*a^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBOSE)`

output `a^4*A/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))+(A*a^4+4*B*a^4)/d*ln(sec(d*x+c)+tan(d*x+c))-(4*A*a^4+B*a^4)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(4*A*a^4+6*B*a^4)/d*tan(d*x+c)+(6*A*a^4+4*B*a^4)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+B*a^4/d*(d*x+c)`

3.35.5 Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.91

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{48 B a^4 dx \cos(dx + c)^4 + 3(35 A + 48 B) a^4 \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(35 A + 48 B) a^4 \cos(dx + c)^4 \log(\sin(dx + c) - 1)}{1}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")`

output `1/48*(48*B*a^4*d*x*cos(d*x + c)^4 + 3*(35*A + 48*B)*a^4*cos(d*x + c)^4*log(sin(d*x + c) + 1) - 3*(35*A + 48*B)*a^4*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*(160*(A + B)*a^4*cos(d*x + c)^3 + 3*(27*A + 16*B)*a^4*cos(d*x + c)^2 + 8*(4*A + B)*a^4*cos(d*x + c) + 6*A*a^4)*sin(d*x + c))/(d*cos(d*x + c)^4)`

3.35.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output Timed out

3.35. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.35.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.77

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{64 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^4 + 16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ba^4 + 48 (dx + c) Ba^4 - 3 Aa^4 \log\left(\frac{\sin(dx + c) + 1}{\sin(dx + c) - 1}\right) + 3 Aa^4 \log(\sin(dx + c) - 1) - 72 Aa^4 \frac{2 \sin(dx + c)}{\sin^2(dx + c) - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) - 48 Ba^4 \frac{2 \sin(dx + c)}{\sin^2(dx + c) - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) + 24 Aa^4 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 96 Ba^4 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 192 Aa^4 \tan(dx + c) + 288 Ba^4 \tan(dx + c)}{d}$$

```
input integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")
```

```
output 1/48*(64*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^4 + 16*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^4 + 48*(d*x + c)*B*a^4 - 3*A*a^4*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 72*A*a^4*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 48*B*a^4*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 24*A*a^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 96*B*a^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 192*A*a^4*tan(d*x + c) + 288*B*a^4*tan(d*x + c))/d
```

3.35.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.29

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{24 (dx + c) Ba^4 + 3 (35 Aa^4 + 48 Ba^4) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - 3 (35 Aa^4 + 48 Ba^4) \log\left(\left|\tan\left(\frac{1}{2} dx - \frac{1}{2} c\right) + 1\right|\right)}{d}$$

```
input integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")
```

output $\frac{1}{24}*(24*(d*x + c)*B*a^4 + 3*(35*A*a^4 + 48*B*a^4)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(35*A*a^4 + 48*B*a^4)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(105*A*a^4*\tan(1/2*d*x + 1/2*c)^7 + 120*B*a^4*\tan(1/2*d*x + 1/2*c)^7 - 385*A*a^4*\tan(1/2*d*x + 1/2*c)^5 - 424*B*a^4*\tan(1/2*d*x + 1/2*c)^5 + 511*A*a^4*\tan(1/2*d*x + 1/2*c)^3 + 520*B*a^4*\tan(1/2*d*x + 1/2*c)^3 - 279*A*a^4*\tan(1/2*d*x + 1/2*c) - 216*B*a^4*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^4)/d$

3.35.9 Mupad [B] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.47

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{35 A a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{4d} + \frac{2 B a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{12 B a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{20 A a^4 \sin(c + dx)}{3d \cos(c + dx)} + \frac{27 A a^4 \sin(c + dx)}{8d \cos(c + dx)^2} + \frac{4 A a^4 \sin(c + dx)}{3d \cos(c + dx)^3}$$

$$+ \frac{A a^4 \sin(c + dx)}{4d \cos(c + dx)^4} + \frac{20 B a^4 \sin(c + dx)}{3d \cos(c + dx)} + \frac{2 B a^4 \sin(c + dx)}{d \cos(c + dx)^2} + \frac{B a^4 \sin(c + dx)}{3d \cos(c + dx)^3}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x)^5,x)`

output $(35*A*a^4*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/(4*d) + (2*B*a^4*a*\tan(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (12*B*a^4*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + (20*A*a^4*\sin(c + d*x))/(3*d*\cos(c + d*x)) + (27*A*a^4*\sin(c + d*x))/(8*d*\cos(c + d*x)^2) + (4*A*a^4*\sin(c + d*x))/(3*d*\cos(c + d*x)^3) + (A*a^4*\sin(c + d*x))/(4*d*\cos(c + d*x)^4) + (20*B*a^4*\sin(c + d*x))/(3*d*\cos(c + d*x)) + (2*B*a^4*\sin(c + d*x))/(d*\cos(c + d*x)^2) + (B*a^4*\sin(c + d*x))/(3*d*\cos(c + d*x)^3)$

3.36 $\int (a+a \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^6(c+dx) dx$

3.36.1	Optimal result	517
3.36.2	Mathematica [A] (verified)	518
3.36.3	Rubi [A] (verified)	518
3.36.4	Maple [A] (verified)	523
3.36.5	Fricas [A] (verification not implemented)	523
3.36.6	Sympy [F(-1)]	524
3.36.7	Maxima [B] (verification not implemented)	524
3.36.8	Giac [A] (verification not implemented)	525
3.36.9	Mupad [B] (verification not implemented)	526

3.36.1 Optimal result

Integrand size = 31, antiderivative size = 198

$$\begin{aligned} & \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx \\ &= \frac{7a^4(4A + 5B)\operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{a^4(83A + 100B) \tan(c + dx)}{15d} \\ &+ \frac{a^4(244A + 275B) \sec(c + dx) \tan(c + dx)}{120d} \\ &+ \frac{(26A + 25B) (a^4 + a^4 \cos(c + dx)) \sec^2(c + dx) \tan(c + dx)}{30d} \\ &+ \frac{(8A + 5B) (a^2 + a^2 \cos(c + dx))^2 \sec^3(c + dx) \tan(c + dx)}{20d} \\ &+ \frac{aA(a + a \cos(c + dx))^3 \sec^4(c + dx) \tan(c + dx)}{5d} \end{aligned}$$

```
output 7/8*a^4*(4*A+5*B)*arctanh(sin(d*x+c))/d+1/15*a^4*(83*A+100*B)*tan(d*x+c)/d
+1/120*a^4*(244*A+275*B)*sec(d*x+c)*tan(d*x+c)/d+1/30*(26*A+25*B)*(a^4+a^4
*cos(d*x+c))*sec(d*x+c)^2*tan(d*x+c)/d+1/20*(8*A+5*B)*(a^2+a^2*cos(d*x+c))
^2*sec(d*x+c)^3*tan(d*x+c)/d+1/5*a*A*(a+a*cos(d*x+c))^3*sec(d*x+c)^4*tan(d
*x+c)/d
```

3.36.2 Mathematica [A] (verified)

Time = 4.34 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.10

$$\begin{aligned} & \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx \\ &= \frac{7a^4 A \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{35a^4 B \operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{8a^4 A \tan(c + dx)}{d} \\ &+ \frac{8a^4 B \tan(c + dx)}{d} + \frac{7a^4 A \sec(c + dx) \tan(c + dx)}{2d} + \frac{27a^4 B \sec(c + dx) \tan(c + dx)}{8d} \\ &+ \frac{a^4 A \sec^3(c + dx) \tan(c + dx)}{d} + \frac{a^4 B \sec^3(c + dx) \tan(c + dx)}{d} \\ &+ \frac{8a^4 A \tan^3(c + dx)}{3d} + \frac{4a^4 B \tan^3(c + dx)}{3d} + \frac{a^4 A \tan^5(c + dx)}{5d} \end{aligned}$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(7*a^4*A*ArcTanh[Sin[c + d*x]])/(2*d) + (35*a^4*B*ArcTanh[Sin[c + d*x]])/(8*d) + (8*a^4*A*Tan[c + d*x])/d + (8*a^4*B*Tan[c + d*x])/d + (7*a^4*A*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (27*a^4*B*Sec[c + d*x]*Tan[c + d*x])/(8*d) + (a^4*A*Sec[c + d*x]^3*Tan[c + d*x])/d + (a^4*B*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + (8*a^4*A*Tan[c + d*x]^3)/(3*d) + (4*a^4*B*Tan[c + d*x]^3)/(3*d) + (a^4*A*Tan[c + d*x]^5)/(5*d)`

3.36.3 Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.08, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3454, 3042, 3454, 3042, 3454, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^6(c + dx) (a \cos(c + dx) + a)^4 (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^6} dx \end{aligned}$$

$$\begin{aligned}
& \downarrow 3454 \\
& \frac{1}{5} \int (\cos(c+dx)a+a)^3 (a(8A+5B) + a(A+5B)\cos(c+dx)) \sec^5(c+dx) dx + \\
& \quad \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^3}{5d} \\
& \downarrow 3042 \\
& \frac{1}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^3 (a(8A+5B) + a(A+5B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^5} dx + \\
& \quad \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^3}{5d} \\
& \downarrow 3454 \\
& \frac{1}{5} \left(\frac{1}{4} \int (\cos(c+dx)a+a)^2 (2(26A+25B)a^2 + (12A+25B)\cos(c+dx)a^2) \sec^4(c+dx) dx + \frac{(8A+5B)\tan(c+dx)}{5d} \right. \\
& \quad \left. \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^3}{5d} \right) \\
& \downarrow 3042 \\
& \frac{1}{5} \left(\frac{1}{4} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2 (2(26A+25B)a^2 + (12A+25B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^4} dx + \frac{(8A+5B)\tan(c+dx)}{5d} \right. \\
& \quad \left. \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^3}{5d} \right) \\
& \downarrow 3454 \\
& \frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int (\cos(c+dx)a+a) ((244A+275B)a^3 + (88A+125B)\cos(c+dx)a^3) \sec^3(c+dx) dx + \frac{2(26A+25B)\tan(c+dx)}{5d} \right. \right. \\
& \quad \left. \left. \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^3}{5d} \right) \right) \\
& \downarrow 3042 \\
& \frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a) ((244A+275B)a^3 + (88A+125B)\sin(c+dx+\frac{\pi}{2})a^3)}{\sin(c+dx+\frac{\pi}{2})^3} dx + \frac{2(26A+25B)\tan(c+dx)}{5d} \right. \right. \\
& \quad \left. \left. \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^3}{5d} \right) \right) \\
& \downarrow 3447
\end{aligned}$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \frac{((88A + 125B) \cos^2(c + dx)a^4 + (244A + 275B)a^4 + ((88A + 125B)a^4 + (244A + 275B)a^4) \cos(c + dx) + aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^3}{5d} dx \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \frac{(88A + 125B) \sin(c + dx + \frac{\pi}{2})^2 a^4 + (244A + 275B)a^4 + ((88A + 125B)a^4 + (244A + 275B)a^4) \sin(c + dx + \frac{\pi}{2}) + aA \tan(c + dx) \sec^4(c + dx)(a \cos(c + dx) + a)^3}{5d} dx \right) \right)$$

↓ 3500

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (8(83A + 100B)a^4 + 105(4A + 5B) \cos(c + dx)a^4) \sec^2(c + dx) dx + \frac{a^4(244A + 275B) \tan(c + dx)}{2d} \right) \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{8(83A + 100B)a^4 + 105(4A + 5B) \sin(c + dx + \frac{\pi}{2}) a^4}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{a^4(244A + 275B) \tan(c + dx) \sec(c + dx)}{2d} \right) \right) \right)$$

↓ 3227

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(8a^4(83A + 100B) \int \sec^2(c + dx) dx + 105a^4(4A + 5B) \int \sec(c + dx) dx \right) \right) \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(105a^4(4A + 5B) \int \csc(c + dx + \frac{\pi}{2}) dx + 8a^4(83A + 100B) \int \csc(c + dx + \frac{\pi}{2})^2 dx \right) \right) \right) \right)$$

↓ 4254

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(105a^4(4A + 5B) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx - \frac{8a^4(83A + 100B) \int 1d(-\tan(c + dx))}{d} \right) \right) + \frac{a^4(244A + 275B) \tan(c + dx)}{2d} \right) \right) + \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^3}{5d}$$

↓ 24

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(105a^4(4A + 5B) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{8a^4(83A + 100B) \tan(c + dx)}{d} \right) \right) + \frac{a^4(244A + 275B) \tan(c + dx)}{2d} \right) \right) + \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^3}{5d}$$

↓ 4257

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\frac{105a^4(4A + 5B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{8a^4(83A + 100B) \tan(c + dx)}{d} \right) \right) + \frac{a^4(244A + 275B) \tan(c + dx)}{2d} \right) \right) + \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^3}{5d}$$

input `Int[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a*A*(a + a*Cos[c + d*x])^3*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) + (((8*A + 5*B)*(a^2 + a^2*Cos[c + d*x])^2*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((2*(26*A + 25*B)*(a^4 + a^4*Cos[c + d*x])*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((a^4*(244*A + 275*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((105*a^4*(4*A + 5*B)*ArcTanh[Sin[c + d*x]])/d + (8*a^4*(83*A + 100*B)*Tan[c + d*x])/d)/2)/3)/4)/5`

3.36.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.36.4 Maple [A] (verified)

Time = 5.59 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.10

method	result
parallelrisch	$70 \left(-\frac{3 \left(\frac{\cos(5dx+5c)}{10} + \frac{\cos(3dx+3c)}{2} + \cos(dx+c) \right) \left(A + \frac{5B}{4} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)}{2} + \frac{3 \left(\frac{\cos(5dx+5c)}{10} + \frac{\cos(3dx+3c)}{2} + \cos(dx+c) \right) \left(A + \frac{5B}{4} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{2} \right)$
parts	$\frac{a^4 A \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c)}{d} + \frac{(a^4 A + 4B a^4) \tan(dx+c)}{d} + \frac{(4a^4 A + B a^4) \left(-\left(-\frac{\sec^3(dx+c)}{3} \right) \right)}{d}$
derivativedivides	$\frac{a^4 A \tan(dx+c) + B a^4 \ln(\sec(dx+c) + \tan(dx+c)) + 4a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 4B a^4 \tan(dx+c)}{d}$
default	$\frac{a^4 A \tan(dx+c) + B a^4 \ln(\sec(dx+c) + \tan(dx+c)) + 4a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 4B a^4 \tan(dx+c)}{d}$
risch	$-\frac{ia^4 (420A e^{9i(dx+c)} + 405B e^{9i(dx+c)} - 120A e^{8i(dx+c)} - 480B e^{8i(dx+c)} + 1320A e^{7i(dx+c)} + 930B e^{7i(dx+c)} - 1920A e^{6i(dx+c)} + 1440B e^{6i(dx+c)} - 360A e^{5i(dx+c)} - 360B e^{5i(dx+c)} + 360A e^{4i(dx+c)} + 360B e^{4i(dx+c)} - 360A e^{3i(dx+c)} - 360B e^{3i(dx+c)} + 360A e^{2i(dx+c)} + 360B e^{2i(dx+c)} - 360A e^{i(dx+c)} - 360B e^{i(dx+c)} + 360A + 360B)}{d}$

```
input int((a+cos(d*x+c)*a)^4*(A+B*cos(d*x+c))*sec(d*x+c)^6,x,method=_RETURNVERBOSE)
```

```
output 70/3*(-3/2*(1/10*cos(5*d*x+5*c)+1/2*cos(3*d*x+3*c)+cos(d*x+c))*(A+5/4*B)*ln(tan(1/2*d*x+1/2*c)-1)+3/2*(1/10*cos(5*d*x+5*c)+1/2*cos(3*d*x+3*c)+cos(d*x+c))*(A+5/4*B)*ln(tan(1/2*d*x+1/2*c)+1)+(33/35*A+93/140*B)*sin(2*d*x+2*c)+(11/10*A+38/35*B)*sin(3*d*x+3*c)+(3/10*A+81/280*B)*sin(4*d*x+4*c)+(83/350*A+2/7*B)*sin(5*d*x+5*c)+sin(d*x+c)*(A+4/5*B))*a^4/d/(cos(5*d*x+5*c)+5*cos(3*d*x+3*c)+10*cos(d*x+c))
```

3.36.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.83

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{105 (4 A + 5 B) a^4 \cos(dx + c)^5 \log(\sin(dx + c) + 1) - 105 (4 A + 5 B) a^4 \cos(dx + c)^5 \log(-\sin(dx + c))}{d}$$

```
input integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="fracas")
```

3.36. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$

output $1/240*(105*(4*A + 5*B)*a^4*\cos(d*x + c)^5*\log(\sin(d*x + c) + 1) - 105*(4*A + 5*B)*a^4*\cos(d*x + c)^5*\log(-\sin(d*x + c) + 1) + 2*(8*(83*A + 100*B)*a^4*\cos(d*x + c)^4 + 15*(28*A + 27*B)*a^4*\cos(d*x + c)^3 + 16*(17*A + 10*B)*a^4*\cos(d*x + c)^2 + 30*(4*A + B)*a^4*\cos(d*x + c) + 24*A*a^4)*\sin(d*x + c))/(d*\cos(d*x + c)^5)$

3.36.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**6,x)`

output Timed out

3.36.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 376 vs. $2(186) = 372$.

Time = 0.22 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.90

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{16 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c)) Aa^4 + 480 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^4}{\dots}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="maxima")`

output $\frac{1}{240}(16(3\tan(dx+c)^5 + 10\tan(dx+c)^3 + 15\tan(dx+c))Aa^4 + 480(\tan(dx+c)^3 + 3\tan(dx+c))Aa^4 + 320(\tan(dx+c)^3 + 3\tan(dx+c))Ba^4 - 60Aa^4(2(3\sin(dx+c)^3 - 5\sin(dx+c))/(\sin(dx+c)^4 - 2\sin(dx+c)^2 + 1) - 3\log(\sin(dx+c) + 1) + 3\log(\sin(dx+c) - 1)) - 15Ba^4(2(3\sin(dx+c)^3 - 5\sin(dx+c))/(\sin(dx+c)^4 - 2\sin(dx+c)^2 + 1) - 3\log(\sin(dx+c) + 1) + 3\log(\sin(dx+c) - 1)) - 240Aa^4(2\sin(dx+c)/(\sin(dx+c)^2 - 1) - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1)) - 360Ba^4(2\sin(dx+c)/(\sin(dx+c)^2 - 1) - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1)) + 120Ba^4(\log(\sin(dx+c) + 1) - \log(\sin(dx+c) - 1)) + 240Aa^4\tan(dx+c) + 960Ba^4\tan(dx+c))/d$

3.36.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.24

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{105(4Aa^4 + 5Ba^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 105(4Aa^4 + 5Ba^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) - \frac{2(420Aa^4 \tan^9(1/2dx + 1/2c) + 525Ba^4 \tan^9(1/2dx + 1/2c) - 1960Aa^4 \tan^7(1/2dx + 1/2c) - 2450Ba^4 \tan^7(1/2dx + 1/2c) + 3584Aa^4 \tan^5(1/2dx + 1/2c) + 4480Ba^4 \tan^5(1/2dx + 1/2c) - 3160Aa^4 \tan^3(1/2dx + 1/2c) - 3950Ba^4 \tan^3(1/2dx + 1/2c) + 1500Aa^4 \tan(1/2dx + 1/2c) + 1395Ba^4 \tan(1/2dx + 1/2c))}{(\tan(1/2dx + 1/2c)^2 - 1)^5}}{d}$$

input `integrate((a+a*cos(dx+c))^4*(A+B*cos(dx+c))*sec(dx+c)^6,x, algorithm="giac")`

output $\frac{1}{120}(105(4Aa^4 + 5Ba^4)\log(\tan(1/2dx + 1/2c) + 1) - 105(4Aa^4 + 5Ba^4)\log(\tan(1/2dx + 1/2c) - 1) - 2(420Aa^4 \tan^9(1/2dx + 1/2c) + 525Ba^4 \tan^9(1/2dx + 1/2c) - 1960Aa^4 \tan^7(1/2dx + 1/2c) - 2450Ba^4 \tan^7(1/2dx + 1/2c) + 3584Aa^4 \tan^5(1/2dx + 1/2c) + 4480Ba^4 \tan^5(1/2dx + 1/2c) - 3160Aa^4 \tan^3(1/2dx + 1/2c) - 3950Ba^4 \tan^3(1/2dx + 1/2c) + 1500Aa^4 \tan(1/2dx + 1/2c) + 1395Ba^4 \tan(1/2dx + 1/2c))/(\tan(1/2dx + 1/2c)^2 - 1)^5)/d$

3.36.9 Mupad [B] (verification not implemented)

Time = 2.94 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.13

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{7a^4 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (4A + 5B)}{4d} - \frac{\left(7Aa^4 + \frac{35Ba^4}{4}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(-\frac{98Aa^4}{3} - \frac{245Ba^4}{6}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{896Aa^4}{15} + \frac{224Ba^4}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{158Aa^4}{3} + \frac{395Ba^4}{6}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{224Ba^4}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} - 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1 \right)}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x)^6,x)`output `(7*a^4*atanh(tan(c/2 + (d*x)/2))*(4*A + 5*B))/(4*d) - (tan(c/2 + (d*x)/2)*(25*A*a^4 + (93*B*a^4)/4) + tan(c/2 + (d*x)/2)^9*(7*A*a^4 + (35*B*a^4)/4) - tan(c/2 + (d*x)/2)^7*((98*A*a^4)/3 + (245*B*a^4)/6) - tan(c/2 + (d*x)/2)^5*((158*A*a^4)/3 + (395*B*a^4)/6) + tan(c/2 + (d*x)/2)^3*((158*A*a^4)/3 + (395*B*a^4)/6) + tan(c/2 + (d*x)/2)^1*((224*B*a^4)/3))/(d*(5*tan(c/2 + (d*x)/2)^2 - 10*tan(c/2 + (d*x)/2)^4 + 10*tan(c/2 + (d*x)/2)^6 - 5*tan(c/2 + (d*x)/2)^8 + tan(c/2 + (d*x)/2)^10 - 1))`

3.37 $\int (a+a \cos(c+dx))^4(A+B \cos(c+dx)) \sec^7(c+dx) dx$

3.37.1	Optimal result	527
3.37.2	Mathematica [A] (verified)	528
3.37.3	Rubi [A] (verified)	528
3.37.4	Maple [A] (verified)	533
3.37.5	Fricas [A] (verification not implemented)	534
3.37.6	Sympy [F(-1)]	534
3.37.7	Maxima [B] (verification not implemented)	535
3.37.8	Giac [A] (verification not implemented)	535
3.37.9	Mupad [B] (verification not implemented)	536

3.37.1 Optimal result

Integrand size = 31, antiderivative size = 229

$$\int (a + a \cos(c + dx))^4(A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{7a^4(7A + 8B)\operatorname{arctanh}(\sin(c + dx))}{16d} + \frac{a^4(72A + 83B) \tan(c + dx)}{15d}$$

$$+ \frac{7a^4(7A + 8B) \sec(c + dx) \tan(c + dx)}{16d} + \frac{a^4(159A + 176B) \sec^2(c + dx) \tan(c + dx)}{120d}$$

$$+ \frac{(73A + 72B) (a^4 + a^4 \cos(c + dx)) \sec^3(c + dx) \tan(c + dx)}{120d}$$

$$+ \frac{(3A + 2B) (a^2 + a^2 \cos(c + dx))^2 \sec^4(c + dx) \tan(c + dx)}{120d}$$

$$+ \frac{aA(a + a \cos(c + dx))^3 \sec^5(c + dx) \tan(c + dx)}{6d}$$

output

```
7/16*a^4*(7*A+8*B)*arctanh(sin(d*x+c))/d+1/15*a^4*(72*A+83*B)*tan(d*x+c)/d
+7/16*a^4*(7*A+8*B)*sec(d*x+c)*tan(d*x+c)/d+1/120*a^4*(159*A+176*B)*sec(d*
x+c)^2*tan(d*x+c)/d+1/120*(73*A+72*B)*(a^4+a^4*cos(d*x+c))*sec(d*x+c)^3*ta
n(d*x+c)/d+1/10*(3*A+2*B)*(a^2+a^2*cos(d*x+c))^2*sec(d*x+c)^4*tan(d*x+c)/d
+1/6*a*A*(a+a*cos(d*x+c))^3*sec(d*x+c)^5*tan(d*x+c)/d
```

3.37.2 Mathematica [A] (verified)

Time = 5.11 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.13

$$\begin{aligned} & \int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx \\ &= \frac{49a^4 A \operatorname{Arctanh}(\sin(c + dx))}{16d} + \frac{7a^4 B \operatorname{Arctanh}(\sin(c + dx))}{2d} \\ &+ \frac{8a^4 A \tan(c + dx)}{d} + \frac{8a^4 B \tan(c + dx)}{d} + \frac{49a^4 A \sec(c + dx) \tan(c + dx)}{16d} \\ &+ \frac{7a^4 B \sec(c + dx) \tan(c + dx)}{2d} + \frac{41a^4 A \sec^3(c + dx) \tan(c + dx)}{24d} \\ &+ \frac{a^4 B \sec^3(c + dx) \tan(c + dx)}{d} + \frac{a^4 A \sec^5(c + dx) \tan(c + dx)}{6d} \\ &+ \frac{4a^4 A \tan^3(c + dx)}{d} + \frac{8a^4 B \tan^3(c + dx)}{3d} + \frac{4a^4 A \tan^5(c + dx)}{5d} + \frac{a^4 B \tan^5(c + dx)}{5d} \end{aligned}$$

input `Integrate[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^7,x]`

output `(49*a^4*A*ArcTanh[Sin[c + d*x]])/(16*d) + (7*a^4*B*ArcTanh[Sin[c + d*x]])/(2*d) + (8*a^4*A*Tan[c + d*x])/d + (8*a^4*B*Tan[c + d*x])/d + (49*a^4*A*Sec[c + d*x]*Tan[c + d*x])/(16*d) + (7*a^4*B*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (41*a^4*A*Sec[c + d*x]^3*Tan[c + d*x])/(24*d) + (a^4*B*Sec[c + d*x]^3*Tan[c + d*x])/d + (a^4*A*Sec[c + d*x]^5*Tan[c + d*x])/(6*d) + (4*a^4*A*Tan[c + d*x]^3)/d + (8*a^4*B*Tan[c + d*x]^3)/(3*d) + (4*a^4*A*Tan[c + d*x]^5)/(5*d) + (a^4*B*Tan[c + d*x]^5)/(5*d)`

3.37.3 Rubi [A] (verified)

Time = 1.78 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.05, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.613$, Rules used = {3042, 3454, 3042, 3454, 3042, 3454, 27, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^7(c + dx)(a \cos(c + dx) + a)^4 (A + B \cos(c + dx)) dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^7} dx \\
& \quad \downarrow \text{3454} \\
& \frac{1}{6} \int (\cos(c + dx)a + a)^3 (3a(3A + 2B) + 2a(A + 3B) \cos(c + dx)) \sec^6(c + dx) dx + \\
& \quad \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^3 (3a(3A + 2B) + 2a(A + 3B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^6} dx + \\
& \quad \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d} \\
& \quad \downarrow \text{3454} \\
& \frac{1}{6} \left(\frac{1}{5} \int (\cos(c + dx)a + a)^2 ((73A + 72B)a^2 + 14(2A + 3B) \cos(c + dx)a^2) \sec^5(c + dx) dx + \frac{3(3A + 2B) \tan(c + dx)}{5} \right. \\
& \quad \left. \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6} \left(\frac{1}{5} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^2 ((73A + 72B)a^2 + 14(2A + 3B) \sin(c + dx + \frac{\pi}{2})a^2)}{\sin(c + dx + \frac{\pi}{2})^5} dx + \frac{3(3A + 2B) \tan(c + dx)}{5} \right. \\
& \quad \left. \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d} \right) \\
& \quad \downarrow \text{3454} \\
& \frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \int 3(\cos(c + dx)a + a) ((159A + 176B)a^3 + 2(43A + 52B) \cos(c + dx)a^3) \sec^4(c + dx) dx + \frac{(73A + 72B) \tan(c + dx)}{4} \right. \right. \\
& \quad \left. \left. \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d} \right) \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int (\cos(c + dx)a + a) ((159A + 176B)a^3 + 2(43A + 52B) \cos(c + dx)a^3) \sec^4(c + dx) dx + \frac{(73A + 72B) \tan(c + dx)}{4} \right. \right. \\
& \quad \left. \left. \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d} \right) \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.37. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)((159A+176B)a^3+2(43A+52B)\sin(c+dx+\frac{\pi}{2})a^3)}{\sin(c+dx+\frac{\pi}{2})^4} dx + \frac{(73A+72B)\tan(c+dx)}{aA \tan(c+dx) \sec^5(c+dx)(a \cos(c+dx)+a)^3} \right) \right)$$

$$\frac{1}{6d} \downarrow 3447$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int (2(43A+52B)\cos^2(c+dx)a^4+(159A+176B)a^4+(2(43A+52B)a^4+(159A+176B)a^4)\cos(c+dx)) \right) \right)$$

$$\frac{1}{6d} \downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int \frac{2(43A+52B)\sin(c+dx+\frac{\pi}{2})^2 a^4+(159A+176B)a^4+(2(43A+52B)a^4+(159A+176B)a^4)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^4} \right) \right)$$

$$\frac{1}{6d} \downarrow 3500$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \int (105(7A+8B)a^4+8(72A+83B)\cos(c+dx)a^4)\sec^3(c+dx)dx + \frac{a^4(159A+176B)\tan(c+dx)}{3d} \right) \right) \right)$$

$$\frac{1}{6d} \downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \int \frac{105(7A+8B)a^4+8(72A+83B)\sin(c+dx+\frac{\pi}{2})a^4}{\sin(c+dx+\frac{\pi}{2})^3} dx + \frac{a^4(159A+176B)\tan(c+dx)\sec^2(c+dx)}{3d} \right) \right) \right)$$

$$\frac{1}{6d} \downarrow 3227$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(105a^4(7A+8B) \int \sec^3(c+dx)dx + 8a^4(72A+83B) \int \sec^2(c+dx)dx \right) \right) \right) \right) + \frac{a^4(159A+176B)\tan(c+dx)}{3d}$$

$$\frac{1}{6d} \downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(8a^4(72A + 83B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^2 dx + 105a^4(7A + 8B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \frac{a^4(159A - 176B) \tan(c + dx)}{d} \right) \right) \right) + \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d}$$

↓ 4254

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(105a^4(7A + 8B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx - \frac{8a^4(72A + 83B) \int 1d(-\tan(c + dx))}{d} \right) \right) \right) + \frac{a^4(159A + 176B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d}$$

↓ 24

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(105a^4(7A + 8B) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx + \frac{8a^4(72A + 83B) \tan(c + dx)}{d} \right) \right) \right) + \frac{a^4(159A + 176B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d}$$

↓ 4255

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(105a^4(7A + 8B) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) \right) \right) \right) + \frac{8a^4(72A + 83B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(105a^4(7A + 8B) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) \right) \right) \right) + \frac{8a^4(72A + 83B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d}$$

↓ 4257

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \left(105a^4(7A + 8B) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) \right) \right) \right) + \frac{8a^4(72A + 83B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec^5(c + dx)(a \cos(c + dx) + a)^3}{6d}$$

input `Int[(a + a*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^7,x]`

```
output (a*A*(a + a*cos[c + d*x])^3*Sec[c + d*x]^5*Tan[c + d*x]/(6*d) + ((3*(3*A
+ 2*B)*(a^2 + a^2*cos[c + d*x])^2*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) + (((
73*A + 72*B)*(a^4 + a^4*cos[c + d*x])*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) +
(3*((a^4*(159*A + 176*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((8*a^4*(72
*A + 83*B)*Tan[c + d*x])/d + 105*a^4*(7*A + 8*B)*(ArcTanh[Sin[c + d*x])/(2
*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))))/3)/4)/5)/6
```

3.37.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_))*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_))*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-A*b^2
- a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.37.4 Maple [A] (verified)

Time = 6.18 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.10

method	result
parallelrisch	$125 \left(-\frac{147 \left(\frac{\cos(6dx+6c)}{15} + \frac{2 \cos(4dx+4c)}{5} + \cos(2dx+2c) + \frac{2}{3} \right) \left(A + \frac{8B}{7} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)}{100} + \frac{147 \left(\frac{\cos(6dx+6c)}{15} + \frac{2 \cos(4dx+4c)}{5} + \cos(2dx+2c) + \frac{2}{3} \right) \left(A + \frac{8B}{7} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{100} \right)$
parts	$a^4 A \left(-\left(-\frac{\sec^5(dx+c)}{6} - \frac{5 \sec^3(dx+c)}{24} - \frac{5 \sec(dx+c)}{16} \right) \tan(dx+c) + \frac{5 \ln(\sec(dx+c) + \tan(dx+c))}{16} \right) + \frac{(a^4 A + 4B a^4) \left(\frac{\sec(dx+c)}{2} \right)}{d}$
derivativedivides	$a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + B a^4 \tan(dx+c) - 4a^4 A \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + 4B a^4 \left(\frac{\sec(dx+c)}{2} \right)$
default	$a^4 A \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + B a^4 \tan(dx+c) - 4a^4 A \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + 4B a^4 \left(\frac{\sec(dx+c)}{2} \right)$
risch	$-\frac{ia^4 (735A e^{11i(dx+c)} + 840B e^{11i(dx+c)} - 240B e^{10i(dx+c)} + 3845A e^{9i(dx+c)} + 3480B e^{9i(dx+c)} - 1920A e^{8i(dx+c)} - 4080B e^{7i(dx+c)} + 1920A e^{6i(dx+c)} + 1920B e^{6i(dx+c)} - 1920A e^{5i(dx+c)} - 1920B e^{5i(dx+c)} + 1920A e^{4i(dx+c)} + 1920B e^{4i(dx+c)} - 1920A e^{3i(dx+c)} - 1920B e^{3i(dx+c)} + 1920A e^{2i(dx+c)} + 1920B e^{2i(dx+c)} - 1920A e^{i(dx+c)} - 1920B e^{i(dx+c)} + 1920A + 1920B)}{100}$

3.37. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$

input `int((a+cos(d*x+c))*a^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x,method=_RETURNVERBOSE)`

output `125/4*(-147/100*(1/15*cos(6*d*x+6*c)+2/5*cos(4*d*x+4*c)+cos(2*d*x+2*c)+2/3)*(A+8/7*B)*ln(tan(1/2*d*x+1/2*c)-1)+147/100*(1/15*cos(6*d*x+6*c)+2/5*cos(4*d*x+4*c)+cos(2*d*x+2*c)+2/3)*(A+8/7*B)*ln(tan(1/2*d*x+1/2*c)+1)+28/125*(8*A+7*B)*sin(2*d*x+2*c)+1/125*(116*B+769/6*A)*sin(3*d*x+3*c)+48/625*(12*A+13*B)*sin(4*d*x+4*c)+7/125*(7/2*A+4*B)*sin(5*d*x+5*c)+4/625*(24*A+83/3*B)*sin(6*d*x+6*c)+sin(d*x+c)*(88/125*B+A))*a^4/d/(cos(6*d*x+6*c)+6*cos(4*d*x+4*c)+15*cos(2*d*x+2*c)+10)`

3.37.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.81

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{105(7A + 8B)a^4 \cos(dx + c)^6 \log(\sin(dx + c) + 1) - 105(7A + 8B)a^4 \cos(dx + c)^6 \log(-\sin(dx + c))}{1}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x, algorithm="fracas")`

output `1/480*(105*(7*A + 8*B))*a^4*cos(d*x + c)^6*log(sin(d*x + c) + 1) - 105*(7*A + 8*B))*a^4*cos(d*x + c)^6*log(-sin(d*x + c) + 1) + 2*(16*(72*A + 83*B))*a^4*cos(d*x + c)^5 + 105*(7*A + 8*B))*a^4*cos(d*x + c)^4 + 32*(18*A + 17*B))*a^4*cos(d*x + c)^3 + 10*(41*A + 24*B))*a^4*cos(d*x + c)^2 + 48*(4*A + B))*a^4*cos(d*x + c) + 40*A*a^4)*sin(d*x + c))/(d*cos(d*x + c)^6)`

3.37.6 SymPy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**7,x)`

output Timed out

3.37. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$

3.37.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 464 vs. $2(215) = 430$.

Time = 0.30 (sec) , antiderivative size = 464, normalized size of antiderivative = 2.03

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{128 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c)) Aa^4 + 640 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^4}{\dots}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x, algorithm="maxima")`

output `1/480*(128*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*A*a^4 + 640*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^4 + 32*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*B*a^4 + 960*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^4 - 5*A*a^4*(2*(15*sin(d*x + c)^5 - 40*sin(d*x + c)^3 + 33*sin(d*x + c))/(sin(d*x + c)^6 - 3*sin(d*x + c)^4 + 3*sin(d*x + c)^2 - 1) - 15*log(sin(d*x + c) + 1) + 15*log(sin(d*x + c) - 1)) - 180*A*a^4*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 120*B*a^4*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 120*A*a^4*(2*sin(d*x + c))/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 480*B*a^4*(2*sin(d*x + c))/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 480*B*a^4*tan(d*x + c))/d`

3.37.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.22

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{105 (7 Aa^4 + 8 Ba^4) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 105 (7 Aa^4 + 8 Ba^4) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) - \frac{2(73}{\dots}$$

input `integrate((a+a*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x, algorithm="giac")`

3.37. $\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$

output $\frac{1}{240} \cdot (105 \cdot (7Aa^4 + 8Ba^4) \cdot \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1)) - 105 \cdot (7Aa^4 + 8Ba^4) \cdot \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) - 2 \cdot (735Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^{11} + 840Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^{11} - 4165Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^9 - 4760Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^9 + 9702Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 11088Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 11802Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 13488Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 7355Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 9320Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 3105Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 3000Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)) / (\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1)^6) / d$

3.37.9 Mupad [B] (verification not implemented)

Time = 2.95 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.14

$$\int (a + a \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{\left(-\frac{49Aa^4}{8} - 7Ba^4\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{11} + \left(\frac{833Aa^4}{24} + \frac{119Ba^4}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(-\frac{1617Aa^4}{20} - \frac{462Ba^4}{5}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{1967Aa^4}{20} + \frac{562Ba^4}{5}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{1471Aa^4}{24} + \frac{233Ba^4}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{735Aa^4}{20} + \frac{11088Ba^4}{5}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 15 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{12} - 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^{10} + 15 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 1\right)} + \frac{7a^4 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (7A + 8B)}{8d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^4)/cos(c + d*x)^7,x)`

output $(\tan(c/2 + (d*x)/2) \cdot ((207Aa^4)/8 + 25Ba^4) - \tan(c/2 + (d*x)/2)^{11} \cdot ((49Aa^4)/8 + 7Ba^4) + \tan(c/2 + (d*x)/2)^9 \cdot ((833Aa^4)/24 + (119Ba^4)/3) - \tan(c/2 + (d*x)/2)^7 \cdot ((1471Aa^4)/24 + (233Ba^4)/3) - \tan(c/2 + (d*x)/2)^5 \cdot ((1617Aa^4)/20 + (462Ba^4)/5) + \tan(c/2 + (d*x)/2)^3 \cdot ((1967Aa^4)/20 + (562Ba^4)/5) + \tan(c/2 + (d*x)/2) \cdot ((735Aa^4)/20 + (11088Ba^4)/5) + 15 \tan(c/2 + (d*x)/2) + 1) / (d \cdot (15 \tan(c/2 + (d*x)/2)^4 - 6 \tan(c/2 + (d*x)/2)^2 - 20 \tan(c/2 + (d*x)/2)^6 + 15 \tan(c/2 + (d*x)/2)^8 - 6 \tan(c/2 + (d*x)/2)^{10} + \tan(c/2 + (d*x)/2)^{12} + 1)) + (7a^4 \operatorname{atanh}(\tan(c/2 + (d*x)/2)) \cdot (7A + 8B)) / (8d)$

3.38 $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.38.1 Optimal result 537
 3.38.2 Mathematica [B] (verified) 538
 3.38.3 Rubi [A] (verified) 538
 3.38.4 Maple [A] (verified) 541
 3.38.5 Fricas [A] (verification not implemented) 542
 3.38.6 Sympy [B] (verification not implemented) 542
 3.38.7 Maxima [B] (verification not implemented) 543
 3.38.8 Giac [A] (verification not implemented) 544
 3.38.9 Mupad [B] (verification not implemented) 545

3.38.1 Optimal result

Integrand size = 31, antiderivative size = 153

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = -\frac{3(4A-5B)x}{8a} + \frac{4(A-B) \sin(c+dx)}{ad} - \frac{3(4A-5B) \cos(c+dx) \sin(c+dx)}{8ad} - \frac{(4A-5B) \cos^3(c+dx) \sin(c+dx)}{4ad} + \frac{(A-B) \cos^4(c+dx) \sin(c+dx)}{d(a+a \cos(c+dx))} - \frac{4(A-B) \sin^3(c+dx)}{3ad}$$

output

```
-3/8*(4*A-5*B)*x/a+4*(A-B)*sin(d*x+c)/a/d-3/8*(4*A-5*B)*cos(d*x+c)*sin(d*x+c)/a/d-1/4*(4*A-5*B)*cos(d*x+c)^3*sin(d*x+c)/a/d+(A-B)*cos(d*x+c)^4*sin(d*x+c)/d/(a+a*cos(d*x+c))-4/3*(A-B)*sin(d*x+c)^3/a/d
```


3.38.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 311 vs. $2(153) = 306$.

Time = 1.44 (sec) , antiderivative size = 311, normalized size of antiderivative = 2.03

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(-72(4A-5B)dx \cos\left(\frac{dx}{2}\right) - 72(4A-5B)dx \cos\left(c+\frac{dx}{2}\right) + 552A \sin\left(\frac{dx}{2}\right) - 552B \sin\left(c+\frac{dx}{2}\right)\right)}{(192a^2d(1+\cos(c+dx)))}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(-72*(4*A - 5*B)*d*x*Cos[(d*x)/2] - 72*(4*A - 5*B)*d*x*Cos[c + (d*x)/2] + 552*A*Sin[(d*x)/2] - 552*B*Sin[(d*x)/2] + 168*A*Sin[c + (d*x)/2] - 168*B*Sin[c + (d*x)/2] + 144*A*Sin[c + (3*d*x)/2] - 120*B*Sin[c + (3*d*x)/2] + 144*A*Sin[2*c + (3*d*x)/2] - 120*B*Sin[2*c + (3*d*x)/2] - 16*A*Sin[2*c + (5*d*x)/2] + 40*B*Sin[2*c + (5*d*x)/2] - 16*A*Sin[3*c + (5*d*x)/2] + 40*B*Sin[3*c + (5*d*x)/2] + 8*A*Sin[3*c + (7*d*x)/2] - 5*B*Sin[3*c + (7*d*x)/2] + 8*A*Sin[4*c + (7*d*x)/2] - 5*B*Sin[4*c + (7*d*x)/2] + 3*B*Sin[4*c + (9*d*x)/2] + 3*B*Sin[5*c + (9*d*x)/2]))/(192*a*d*(1 + Cos[c + d*x]))`

3.38.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.88, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3456, 3042, 3227, 3042, 3113, 2009, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a\cos(c+dx)+a} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^4 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{a\sin\left(c+dx+\frac{\pi}{2}\right)+a} dx$$

$$\downarrow \text{3456}$$

$$\begin{aligned}
& \frac{\int \cos^3(c+dx)(4a(A-B) - a(4A-5B)\cos(c+dx))dx}{a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sin(c+dx+\frac{\pi}{2})^3(4a(A-B) - a(4A-5B)\sin(c+dx+\frac{\pi}{2}))dx}{a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3227} \\
& \frac{4a(A-B)\int \cos^3(c+dx)dx - a(4A-5B)\int \cos^4(c+dx)dx}{a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{4a(A-B)\int \sin(c+dx+\frac{\pi}{2})^3dx - a(4A-5B)\int \sin(c+dx+\frac{\pi}{2})^4dx}{a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3113} \\
& \frac{-\frac{4a(A-B)\int(1-\sin^2(c+dx))d(-\sin(c+dx))}{d} - a(4A-5B)\int \sin(c+dx+\frac{\pi}{2})^4dx}{a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{2009} \\
& \frac{-a(4A-5B)\int \sin(c+dx+\frac{\pi}{2})^4dx - \frac{4a(A-B)(\frac{1}{3}\sin^3(c+dx)-\sin(c+dx))}{d}}{a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3115} \\
& \frac{-a(4A-5B)\left(\frac{3}{4}\int \cos^2(c+dx)dx + \frac{\sin(c+dx)\cos^3(c+dx)}{4d}\right) - \frac{4a(A-B)(\frac{1}{3}\sin^3(c+dx)-\sin(c+dx))}{d}}{a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{-a(4A-5B)\left(\frac{3}{4}\int \sin(c+dx+\frac{\pi}{2})^2dx + \frac{\sin(c+dx)\cos^3(c+dx)}{4d}\right) - \frac{4a(A-B)(\frac{1}{3}\sin^3(c+dx)-\sin(c+dx))}{d}}{a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{d(a\cos(c+dx)+a)}
\end{aligned}$$

3.38. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3115} \\
 & \frac{-a(4A - 5B) \left(\frac{3}{4} \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{\sin(c+dx) \cos^3(c+dx)}{4d} \right) - \frac{4a(A-B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d}}{a^2} + \\
 & \quad \frac{(A - B) \sin(c + dx) \cos^4(c + dx)}{d(a \cos(c + dx) + a)} \\
 & \quad \downarrow \text{24} \\
 & \frac{-\frac{4a(A-B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d} - a(4A - 5B) \left(\frac{\sin(c+dx) \cos^3(c+dx)}{4d} + \frac{3}{4} \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right)}{a^2} + \\
 & \quad \frac{(A - B) \sin(c + dx) \cos^4(c + dx)}{d(a \cos(c + dx) + a)}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*cos[c + d*x]),x]`

output `((A - B)*Cos[c + d*x]^4*Sin[c + d*x])/(d*(a + a*cos[c + d*x])) + ((-4*a*(A - B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d - a*(4*A - 5*B)*((Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (3*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d))))/4) /a^2`

3.38.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.38.4 Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.63

method	result
parallelrisch	$\frac{((-8A+38B)\cos(2dx+2c)+(8A-2B)\cos(3dx+3c)+3B\cos(4dx+4c)+(136A-82B)\cos(dx+c)+248A-221B)\tan\left(\frac{dx}{2}\right)}{96ad}$
derivativedivides	$A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{2\left(\left(\frac{25B}{8} - \frac{5A}{2}\right)\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(\frac{115B}{24} - \frac{31A}{6}\right)\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(\frac{109B}{24} - \frac{25A}{6}\right)\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(\frac{11B}{24} - \frac{A}{6}\right)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^4}$
default	$A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{2\left(\left(\frac{25B}{8} - \frac{5A}{2}\right)\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(\frac{115B}{24} - \frac{31A}{6}\right)\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(\frac{109B}{24} - \frac{25A}{6}\right)\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(\frac{11B}{24} - \frac{A}{6}\right)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^4}$
risch	$-\frac{3xA}{2a} + \frac{15Bx}{8a} - \frac{7ie^{i(dx+c)}A}{8ad} + \frac{7ie^{i(dx+c)}B}{8ad} + \frac{7ie^{-i(dx+c)}A}{8ad} - \frac{7ie^{-i(dx+c)}B}{8ad} + \frac{2iA}{da(e^{i(dx+c)}+1)} - \frac{2iB}{da(e^{i(dx+c)}-1)}$
norman	$\frac{(A-B)\tan^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} - \frac{3(4A-5B)x}{8a} + \frac{86(A-B)\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{3ad} - \frac{15(4A-5B)x\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)}{8a} - \frac{15(4A-5B)x\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)}{4a}$

input `int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a), x, method=_RETURNVERBOSE)`

3.38. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

output $1/96*(((-8*A+38*B)*\cos(2*d*x+2*c)+(8*A-2*B)*\cos(3*d*x+3*c)+3*B*\cos(4*d*x+4*c)+(136*A-82*B)*\cos(d*x+c)+248*A-221*B)*\tan(1/2*d*x+1/2*c)-144*(A-5/4*B)*x*d)/a/d$

3.38.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.78

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \frac{9(4A-5B)dx\cos(dx+c)+9(4A-5B)dx-(6B\cos(dx+c))^4+2(4A-B)\cos(dx+c)^3-(4A-19B)\cos(dx+c)+64A-64B*\sin(dx+c)}{24(ad\cos(dx+c)+ad)}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output $-1/24*(9*(4*A - 5*B)*d*x*\cos(d*x + c) + 9*(4*A - 5*B)*d*x - (6*B*\cos(d*x + c))^4 + 2*(4*A - B)*\cos(d*x + c)^3 - (4*A - 13*B)*\cos(d*x + c)^2 + (28*A - 19*B)*\cos(d*x + c) + 64*A - 64*B)*\sin(d*x + c))/(a*d*\cos(d*x + c) + a*d)$

3.38.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1794 vs. $2(134) = 268$.

Time = 1.98 (sec) , antiderivative size = 1794, normalized size of antiderivative = 11.73

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x)`

output `Piecewise((-36*A*d*x*tan(c/2 + d*x/2)**8/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) - 144*A*d*x*tan(c/2 + d*x/2)**6/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) - 216*A*d*x*tan(c/2 + d*x/2)**4/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) - 144*A*d*x*tan(c/2 + d*x/2)**2/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) - 36*A*d*x/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) + 24*A*tan(c/2 + d*x/2)**9/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) + 216*A*tan(c/2 + d*x/2)**7/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) + 392*A*tan(c/2 + d*x/2)**5/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) + 296*A*tan(c/2 + d*x/2)**3/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 144*a*d*tan(c/2 + d*x/2)**4 + 96*a*d*tan(c/2 + d*x/2)**2 + 24*a*d) + 96*A*tan(c/2 + d*x/2)/(24*a*d*tan(c/2 + d*x/2)**8 + 96*a*d*tan(c/2 + d*x/2)**6 + 1...`

3.38.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 394 vs. $2(145) = 290$.

Time = 0.31 (sec) , antiderivative size = 394, normalized size of antiderivative = 2.58

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx =$$

$$\frac{B \left(\frac{21 \sin(dx+c)}{\cos(dx+c)+1} + \frac{109 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{115 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{75 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} - \frac{45 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} + \frac{12 \sin(dx+c)}{a(\cos(dx+c)+1)} \right) - 4A \left(\frac{1}{a} \right)}{12d}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output
$$-1/12*(B*((21*\sin(dx + c)/(\cos(dx + c) + 1) + 109*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 115*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 + 75*\sin(dx + c)^7/(\cos(dx + c) + 1)^7)/(a + 4*a*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 6*a*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + 4*a*\sin(dx + c)^6/(\cos(dx + c) + 1)^6 + a*\sin(dx + c)^8/(\cos(dx + c) + 1)^8) - 45*\arctan(\sin(dx + c)/(\cos(dx + c) + 1))/a + 12*\sin(dx + c)/(a*(\cos(dx + c) + 1))) - 4*A*((9*\sin(dx + c)/(\cos(dx + c) + 1) + 16*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 15*\sin(dx + c)^5/(\cos(dx + c) + 1)^5)/(a + 3*a*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 3*a*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + a*\sin(dx + c)^6/(\cos(dx + c) + 1)^6) - 9*\arctan(\sin(dx + c)/(\cos(dx + c) + 1))/a + 3*\sin(dx + c)/(a*(\cos(dx + c) + 1))))/d$$

3.38.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.18

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx = \frac{9(dx+c)(4A-5B)}{a} - \frac{24(A \tan(\frac{1}{2} dx + \frac{1}{2} c) - B \tan(\frac{1}{2} dx + \frac{1}{2} c))}{a} - \frac{2(60A \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 - 75B \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 + 124A \tan(\frac{1}{2} dx + \frac{1}{2} c)^5)}{24d}$$

24d

input `integrate(cos(dx+c)^4*(A+B*cos(dx+c))/(a+a*cos(dx+c)),x, algorithm="giac")`

output
$$-1/24*(9*(dx + c)*(4*A - 5*B)/a - 24*(A*\tan(1/2*dx + 1/2*c) - B*\tan(1/2*dx + 1/2*c))/a - 2*(60*A*\tan(1/2*dx + 1/2*c)^7 - 75*B*\tan(1/2*dx + 1/2*c)^7 + 124*A*\tan(1/2*dx + 1/2*c)^5 - 115*B*\tan(1/2*dx + 1/2*c)^5 + 100*A*\tan(1/2*dx + 1/2*c)^3 - 109*B*\tan(1/2*dx + 1/2*c)^3 + 36*A*\tan(1/2*dx + 1/2*c) - 21*B*\tan(1/2*dx + 1/2*c))/((\tan(1/2*dx + 1/2*c)^2 + 1)^4*a))/d$$

3.38.9 Mupad [B] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.11

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \frac{15Bx}{8a} - \frac{3Ax}{2a} + \frac{7A\sin(c+dx)}{4ad} - \frac{7B\sin(c+dx)}{4ad} - \frac{A\sin(2c+2dx)}{4ad} + \frac{A\sin(3c+3dx)}{12ad} + \frac{A\tan(\frac{c}{2} + \frac{dx}{2})}{ad} + \frac{B\sin(2c+2dx)}{2ad} - \frac{B\sin(3c+3dx)}{12ad} + \frac{B\sin(4c+4dx)}{32ad} - \frac{B\tan(\frac{c}{2} + \frac{dx}{2})}{ad}$$

input `int((cos(c + d*x))^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`

output `(15*B*x)/(8*a) - (3*A*x)/(2*a) + (7*A*sin(c + d*x))/(4*a*d) - (7*B*sin(c + d*x))/(4*a*d) - (A*sin(2*c + 2*d*x))/(4*a*d) + (A*sin(3*c + 3*d*x))/(12*a*d) + (A*tan(c/2 + (d*x)/2))/(a*d) + (B*sin(2*c + 2*d*x))/(2*a*d) - (B*sin(3*c + 3*d*x))/(12*a*d) + (B*sin(4*c + 4*d*x))/(32*a*d) - (B*tan(c/2 + (d*x)/2))/(a*d)`

3.39 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.39.1	Optimal result	546
3.39.2	Mathematica [B] (verified)	546
3.39.3	Rubi [A] (verified)	547
3.39.4	Maple [A] (verified)	550
3.39.5	Fricas [A] (verification not implemented)	550
3.39.6	Sympy [B] (verification not implemented)	551
3.39.7	Maxima [B] (verification not implemented)	551
3.39.8	Giac [A] (verification not implemented)	552
3.39.9	Mupad [B] (verification not implemented)	553

3.39.1 Optimal result

Integrand size = 31, antiderivative size = 122

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = \frac{3(A-B)x}{2a} - \frac{(3A-4B) \sin(c+dx)}{ad} + \frac{3(A-B) \cos(c+dx) \sin(c+dx)}{2ad} + \frac{(A-B) \cos^3(c+dx) \sin(c+dx)}{d(a+a \cos(c+dx))} + \frac{(3A-4B) \sin^3(c+dx)}{3ad}$$

```
output 3/2*(A-B)*x/a-(3*A-4*B)*sin(d*x+c)/a/d+3/2*(A-B)*cos(d*x+c)*sin(d*x+c)/a/d
+(A-B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))+1/3*(3*A-4*B)*sin(d*x+c)
^3/a/d
```

3.39.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 249 vs. 2(122) = 244.

Time = 1.23 (sec) , antiderivative size = 249, normalized size of antiderivative = 2.04

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(36(A-B)dx \cos\left(\frac{dx}{2}\right) + 36(A-B)dx \cos\left(c+\frac{dx}{2}\right) - 60A \sin\left(\frac{dx}{2}\right) + 69B \sin\left(\frac{dx}{2}\right)\right)}{2(a+a \cos(c+dx))}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*cos[c + d*x]),x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(36*(A - B)*d*x*cos[(d*x)/2] + 36*(A - B)*d*x*Cos[c + (d*x)/2] - 60*A*Sin[(d*x)/2] + 69*B*Sin[(d*x)/2] - 12*A*Sin[c + (d*x)/2] + 21*B*Sin[c + (d*x)/2] - 9*A*Sin[c + (3*d*x)/2] + 18*B*Sin[c + (3*d*x)/2] - 9*A*Sin[2*c + (3*d*x)/2] + 18*B*Sin[2*c + (3*d*x)/2] + 3*A*Sin[2*c + (5*d*x)/2] - 2*B*Sin[2*c + (5*d*x)/2] + 3*A*Sin[3*c + (5*d*x)/2] - 2*B*Sin[3*c + (5*d*x)/2] + B*Sin[3*c + (7*d*x)/2] + B*Sin[4*c + (7*d*x)/2]))/(24*a*d*(1 + Cos[c + d*x]))`

3.39.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.88, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3456, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{a \sin(c + dx + \frac{\pi}{2}) + a} dx$$

↓ 3456

$$\frac{\int \cos^2(c + dx)(3a(A - B) - a(3A - 4B) \cos(c + dx)) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^3(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3042

$$\frac{\int \sin(c + dx + \frac{\pi}{2})^2 (3a(A - B) - a(3A - 4B) \sin(c + dx + \frac{\pi}{2})) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^3(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3227

$$\frac{3a(A - B) \int \cos^2(c + dx) dx - a(3A - 4B) \int \cos^3(c + dx) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^3(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3042

3.39. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \frac{3a(A-B) \int \sin(c+dx + \frac{\pi}{2})^2 dx - a(3A-4B) \int \sin(c+dx + \frac{\pi}{2})^3 dx}{a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx) + a)} \\
& \quad \downarrow \text{3113} \\
& \frac{\frac{a(3A-4B) \int (1-\sin^2(c+dx)) d(-\sin(c+dx))}{d} + 3a(A-B) \int \sin(c+dx + \frac{\pi}{2})^2 dx}{a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx) + a)} \\
& \quad \downarrow \text{2009} \\
& \frac{3a(A-B) \int \sin(c+dx + \frac{\pi}{2})^2 dx + \frac{a(3A-4B)(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d}}{a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx) + a)} \\
& \quad \downarrow \text{3115} \\
& \frac{3a(A-B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{a(3A-4B)(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d}}{a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx) + a)} \\
& \quad \downarrow \text{24} \\
& \frac{\frac{a(3A-4B)(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx))}{d} + 3a(A-B) \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right)}{a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx) + a)}
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `((A - B)*Cos[c + d*x]^3*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])) + (3*a*(A - B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) + (a*(3*A - 4*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/a^2`

3.39.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.39.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.64

method	result
parallelrisch	$\frac{((3A-B)\cos(2dx+2c)+B\cos(3dx+3c)+(-6A+17B)\cos(dx+c)-21A+31B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+18dx(A-B)}{12ad}$
derivativedivides	$-A\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+\frac{2\left(-\frac{3A}{2}+\frac{5B}{2}\right)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2\left(\frac{8B}{3}-2A\right)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2\left(-\frac{A}{2}+\frac{3B}{2}\right)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^3}+3$
default	$-A\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+\frac{2\left(-\frac{3A}{2}+\frac{5B}{2}\right)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2\left(\frac{8B}{3}-2A\right)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2\left(-\frac{A}{2}+\frac{3B}{2}\right)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^3}+3$
risch	$\frac{3xA}{2a}-\frac{3Bx}{2a}+\frac{ie^{i(dx+c)}A}{2ad}-\frac{7ie^{i(dx+c)}B}{8ad}-\frac{ie^{-i(dx+c)}A}{2ad}+\frac{7ie^{-i(dx+c)}B}{8ad}-\frac{2iA}{da(e^{i(dx+c)}+1)}+\frac{2iB}{da(e^{i(dx+c)}-1)}$
norman	$\frac{3(A-B)x}{2a}-\frac{2(A-2B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{ad}-\frac{(A-B)\left(\tan^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{ad}+\frac{6(A-B)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{9(A-B)x\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{6(A-B)x}{a(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right))}$

input `int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)`

output `1/12*((3*A-B)*cos(2*d*x+2*c)+B*cos(3*d*x+3*c)+(-6*A+17*B)*cos(d*x+c)-21*A+31*B)*tan(1/2*d*x+1/2*c)+18*d*x*(A-B))/a/d`

3.39.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.80

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{9(A-B)dx\cos(dx+c)+9(A-B)dx+(2B\cos(dx+c))^3+(3A-B)\cos(dx+c)^2-(3A-7B)\cos(dx+c)}{6(ad\cos(dx+c)+ad)}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `1/6*(9*(A-B)*d*x*cos(d*x+c)+9*(A-B)*d*x+(2*B*cos(d*x+c))^3+(3*A-B)*cos(d*x+c)^2-(3*A-7*B)*cos(d*x+c)-12*A+16*B)*sin(d*x+c))/(a*d*cos(d*x+c)+a*d)`

3.39. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

3.39.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1161 vs. $2(105) = 210$.

Time = 1.27 (sec) , antiderivative size = 1161, normalized size of antiderivative = 9.52

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)), x)
```

```
output Piecewise((9*A*d*x*tan(c/2 + d*x/2)**6/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d
*tan(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) + 27*A*d*x*tan(
c/2 + d*x/2)**4/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 +
18*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) + 27*A*d*x*tan(c/2 + d*x/2)**2/(6*a*d*
tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/2)
**2 + 6*a*d) + 9*A*d*x/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)
)**4 + 18*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) - 6*A*tan(c/2 + d*x/2)**7/(6*a*
d*tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/
2)**2 + 6*a*d) - 36*A*tan(c/2 + d*x/2)**5/(6*a*d*tan(c/2 + d*x/2)**6 + 18*
a*d*tan(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) - 42*A*tan(c
/2 + d*x/2)**3/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 + 1
8*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) - 12*A*tan(c/2 + d*x/2)/(6*a*d*tan(c/2
+ d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/2)**2 + 6*
a*d) - 9*B*d*x*tan(c/2 + d*x/2)**6/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d*tan
(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) - 27*B*d*x*tan(c/2
+ d*x/2)**4/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 + 18*a
*d*tan(c/2 + d*x/2)**2 + 6*a*d) - 27*B*d*x*tan(c/2 + d*x/2)**2/(6*a*d*tan(
c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4 + 18*a*d*tan(c/2 + d*x/2)**2
+ 6*a*d) - 9*B*d*x/(6*a*d*tan(c/2 + d*x/2)**6 + 18*a*d*tan(c/2 + d*x/2)**4
+ 18*a*d*tan(c/2 + d*x/2)**2 + 6*a*d) + 6*B*tan(c/2 + d*x/2)**7/(6*a*d...
```

3.39.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 310 vs. $2(116) = 232$.

Time = 0.30 (sec) , antiderivative size = 310, normalized size of antiderivative = 2.54

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{B \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} + \frac{16 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{15 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{9 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} + \frac{3 \sin(dx+c)}{a(\cos(dx+c)+1)} \right) - 3A \left(\frac{\sin(dx+c)}{a + \frac{2a \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a}{\cos(dx+c)}} \right)}{3d}$$

3.39. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `1/3*(B*((9*sin(d*x + c)/(cos(d*x + c) + 1) + 16*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 15*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(a + 3*a*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*a*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + a*sin(d*x + c)^6/(cos(d*x + c) + 1)^6) - 9*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a + 3*sin(d*x + c)/(a*(cos(d*x + c) + 1))) - 3*A*((sin(d*x + c)/(cos(d*x + c) + 1) + 3*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(a + 2*a*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a*sin(d*x + c)^4/(cos(d*x + c) + 1)^4) - 3*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a + sin(d*x + c)/(a*(cos(d*x + c) + 1)))/d`

3.39.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.24

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{\frac{9(dx+c)(A-B)}{a} - \frac{6(A\tan(\frac{1}{2}dx+\frac{1}{2}c)-B\tan(\frac{1}{2}dx+\frac{1}{2}c))}{a} - \frac{2(9A\tan(\frac{1}{2}dx+\frac{1}{2}c)^5-15B\tan(\frac{1}{2}dx+\frac{1}{2}c)^5+12A\tan(\frac{1}{2}dx+\frac{1}{2}c)^3-16B\tan(\frac{1}{2}dx+\frac{1}{2}c))}{(\tan(\frac{1}{2}dx+\frac{1}{2}c)^2+1)^3}}{6d}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `1/6*(9*(d*x + c)*(A - B)/a - 6*(A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a - 2*(9*A*tan(1/2*d*x + 1/2*c)^5 - 15*B*tan(1/2*d*x + 1/2*c)^5 + 12*A*tan(1/2*d*x + 1/2*c)^3 - 16*B*tan(1/2*d*x + 1/2*c)^3 + 3*A*tan(1/2*d*x + 1/2*c) - 9*B*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 + 1)^3*a))/d`

3.39.9 Mupad [B] (verification not implemented)

Time = 1.50 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.13

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{3x(A-B)}{2a}$$

$$- \frac{(3A-5B)\tan\left(\frac{c}{2}+\frac{dx}{2}\right)^5 + \left(4A-\frac{16B}{3}\right)\tan\left(\frac{c}{2}+\frac{dx}{2}\right)^3 + (A-3B)\tan\left(\frac{c}{2}+\frac{dx}{2}\right)}{d\left(a\tan\left(\frac{c}{2}+\frac{dx}{2}\right)^6 + 3a\tan\left(\frac{c}{2}+\frac{dx}{2}\right)^4 + 3a\tan\left(\frac{c}{2}+\frac{dx}{2}\right)^2 + a\right)}$$

$$- \frac{\tan\left(\frac{c}{2}+\frac{dx}{2}\right)(A-B)}{ad}$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`output `(3*x*(A - B))/(2*a) - (tan(c/2 + (d*x)/2)^5*(3*A - 5*B) + tan(c/2 + (d*x)/2)^3*(4*A - (16*B)/3) + tan(c/2 + (d*x)/2)*(A - 3*B))/(d*(a + 3*a*tan(c/2 + (d*x)/2)^2 + 3*a*tan(c/2 + (d*x)/2)^4 + a*tan(c/2 + (d*x)/2)^6)) - (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.40 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.40.1 Optimal result 554
 3.40.2 Mathematica [B] (verified) 554
 3.40.3 Rubi [A] (verified) 555
 3.40.4 Maple [A] (verified) 556
 3.40.5 Fricas [A] (verification not implemented) 557
 3.40.6 Sympy [B] (verification not implemented) 558
 3.40.7 Maxima [B] (verification not implemented) 558
 3.40.8 Giac [A] (verification not implemented) 559
 3.40.9 Mupad [B] (verification not implemented) 559

3.40.1 Optimal result

Integrand size = 31, antiderivative size = 90

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = -\frac{(A-B)x}{a} + \frac{Bx}{2a} + \frac{(A-B) \sin(c+dx)}{ad} + \frac{B \cos(c+dx) \sin(c+dx)}{2ad} + \frac{(A-B) \sin(c+dx)}{ad(1+\cos(c+dx))}$$

output `-(A-B)*x/a+1/2*B*x/a+(A-B)*sin(d*x+c)/a/d+1/2*B*cos(d*x+c)*sin(d*x+c)/a/d+(A-B)*sin(d*x+c)/a/d/(1+cos(d*x+c))`

3.40.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 197 vs. 2(90) = 180.

Time = 1.03 (sec) , antiderivative size = 197, normalized size of antiderivative = 2.19

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(-4(2A-3B)dx \cos\left(\frac{dx}{2}\right) - 4(2A-3B)dx \cos\left(c+\frac{dx}{2}\right) + 20A \sin\left(\frac{dx}{2}\right) - 20B \sin\left(c+\frac{dx}{2}\right)\right)}{2a}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output $(\text{Cos}[(c + d*x)/2]*\text{Sec}[c/2]*(-4*(2*A - 3*B)*d*x*\text{Cos}[(d*x)/2] - 4*(2*A - 3*B)*d*x*\text{Cos}[c + (d*x)/2] + 20*A*\text{Sin}[(d*x)/2] - 20*B*\text{Sin}[(d*x)/2] + 4*A*\text{Sin}[c + (d*x)/2] - 4*B*\text{Sin}[c + (d*x)/2] + 4*A*\text{Sin}[c + (3*d*x)/2] - 3*B*\text{Sin}[c + (3*d*x)/2] + 4*A*\text{Sin}[2*c + (3*d*x)/2] - 3*B*\text{Sin}[2*c + (3*d*x)/2] + B*\text{Sin}[2*c + (5*d*x)/2] + B*\text{Sin}[3*c + (5*d*x)/2]))/(8*a*d*(1 + \text{Cos}[c + d*x]))$

3.40.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {3042, 3456, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{a \sin(c + dx + \frac{\pi}{2}) + a} dx$$

↓ 3456

$$\frac{\int \cos(c + dx)(2a(A - B) - a(2A - 3B) \cos(c + dx)) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3042

$$\frac{\int \sin(c + dx + \frac{\pi}{2})(2a(A - B) - a(2A - 3B) \sin(c + dx + \frac{\pi}{2})) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3213

$$\frac{\frac{2a(A - B) \sin(c + dx)}{d} - \frac{a(2A - 3B) \sin(c + dx) \cos(c + dx)}{2d} - \frac{1}{2}ax(2A - 3B)}{a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{d(a \cos(c + dx) + a)}$$

input $\text{Int}[(\text{Cos}[c + d*x]^2*(A + B*\text{Cos}[c + d*x]))/(a + a*\text{Cos}[c + d*x]),x]$

```
output ((A - B)*Cos[c + d*x]^2*Sin[c + d*x]/(d*(a + a*cos[c + d*x])) + (-1/2*(a*(2*A - 3*B)*x) + (2*a*(A - B)*Sin[c + d*x])/d - (a*(2*A - 3*B)*Cos[c + d*x]*Sin[c + d*x]/(2*d))/a^2
```

3.40.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3213 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.40.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.68

method	result
parallelrisch	$\frac{(B \cos(2dx+2c)+(4A-2B) \cos(dx+c)+8A-7B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 4x\left(-\frac{3B}{2} + A\right) d}{4ad}$
derivativdivides	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{2\left(\left(\frac{3B}{2} - A\right) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(-A + \frac{B}{2}\right) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2} - (2A-3B) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$
default	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{2\left(\left(\frac{3B}{2} - A\right) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(-A + \frac{B}{2}\right) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2} - (2A-3B) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$
risch	$-\frac{x A}{a} + \frac{3 B x}{2 a} - \frac{i e^{i(dx+c)} A}{2 a d} + \frac{i e^{i(dx+c)} B}{2 a d} + \frac{i e^{-i(dx+c)} A}{2 a d} - \frac{i e^{-i(dx+c)} B}{2 a d} + \frac{2 i A}{d a \left(e^{i(dx+c)} + 1\right)} - \frac{2 i B}{d a \left(e^{i(dx+c)} + 1\right)}$
norman	$\frac{(A-B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a d} + \frac{(3A-2B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{a d} + \frac{(5A-6B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a d} - \frac{(2A-3B)x}{2a} + \frac{7(A-B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a d} - \frac{3(2A-3B)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^3}$

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)
```

```
output 1/4*((B*cos(2*d*x+2*c)+(4*A-2*B)*cos(d*x+c)+8*A-7*B)*tan(1/2*d*x+1/2*c)-4*x*(-3/2*B+A)*d)/a/d
```

3.40.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.92

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = \frac{(2A-3B)dx \cos(dx+c) + (2A-3B)dx - (B \cos(dx+c))^2 + (2A-B) \cos(dx+c) + 4A-4B}{2(ad \cos(dx+c) + ad)}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="fricas")
```

```
output -1/2*((2*A - 3*B)*d*x*cos(d*x + c) + (2*A - 3*B)*d*x - (B*cos(d*x + c))^2 + (2*A - B)*cos(d*x + c) + 4*A - 4*B)*sin(d*x + c)/(a*d*cos(d*x + c) + a*d)
```

3.40. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.40.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 665 vs. 2(68) = 136.

Time = 0.87 (sec) , antiderivative size = 665, normalized size of antiderivative = 7.39

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \left\{ \begin{array}{l} \frac{2Adx \tan^4\left(\frac{c}{2}+\frac{dx}{2}\right)}{2ad \tan^4\left(\frac{c}{2}+\frac{dx}{2}\right)+4ad \tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)+2ad} - \frac{4Adx \tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)}{2ad \tan^4\left(\frac{c}{2}+\frac{dx}{2}\right)+4ad \tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)+2ad} - \frac{2Adx}{2ad \tan^4\left(\frac{c}{2}+\frac{dx}{2}\right)+4ad \tan^2\left(\frac{c}{2}+\frac{dx}{2}\right)+2ad} \\ \frac{x(A+B\cos(c))\cos^2(c)}{a\cos(c)+a} \end{array} \right.$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)), x)`

output `Piecewise((-2*A*d*x*tan(c/2 + d*x/2)**4/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) - 4*A*d*x*tan(c/2 + d*x/2)**2/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) - 2*A*d*x/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) + 2*A*tan(c/2 + d*x/2)**5/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) + 8*A*tan(c/2 + d*x/2)**3/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) + 6*A*tan(c/2 + d*x/2)/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) + 3*B*d*x*tan(c/2 + d*x/2)**4/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) + 6*B*d*x*tan(c/2 + d*x/2)**2/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) + 3*B*d*x/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) - 2*B*tan(c/2 + d*x/2)**5/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) - 10*B*tan(c/2 + d*x/2)**3/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d) - 4*B*tan(c/2 + d*x/2)/(2*a*d*tan(c/2 + d*x/2)**4 + 4*a*d*tan(c/2 + d*x/2)**2 + 2*a*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**2/(a*cos(c) + a), True))`

3.40.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 225 vs. 2(86) = 172.

Time = 0.30 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.50

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \frac{B \left(\frac{\frac{\sin(dx+c)}{\cos(dx+c)+1} + \frac{3\sin(dx+c)^3}{(\cos(dx+c)+1)^3}}{a + \frac{2a\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a\sin(dx+c)^4}{(\cos(dx+c)+1)^4}} - \frac{3\arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} + \frac{\sin(dx+c)}{a(\cos(dx+c)+1)} \right) + A \left(\frac{2\arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} - \frac{1}{a + \frac{a\sin(dx+c)}{\cos(dx+c)+1}} \right)}{d}$$

3.40. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output
$$\begin{aligned} & -(B*((\sin(dx + c)/(\cos(dx + c) + 1) + 3*\sin(dx + c)^3/(\cos(dx + c) + 1)^3)/ \\ & (a + 2*a*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + a*\sin(dx + c)^4/(\cos(dx + c) + 1)^4) - \\ & 3*\arctan(\sin(dx + c)/(\cos(dx + c) + 1))/a + \sin(dx + c)/(a*(\cos(dx + c) + 1))) + \\ & A*(2*\arctan(\sin(dx + c)/(\cos(dx + c) + 1))/a - 2*\sin(dx + c)/((a + a*\sin(dx + c)^2/(\cos(dx + c) + 1)^2)* \\ & (\cos(dx + c) + 1)) - \sin(dx + c)/(a*(\cos(dx + c) + 1))))/d \end{aligned}$$

3.40.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.38

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx = \frac{\frac{(dx+c)(2A-3B)}{a} - \frac{2(A \tan(\frac{1}{2} dx + \frac{1}{2} c) - B \tan(\frac{1}{2} dx + \frac{1}{2} c))}{a} - \frac{2(2A \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 3B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 2A \tan(\frac{1}{2} dx + \frac{1}{2} c) - B \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^2}{a}}{2d}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="giac")`

output
$$\begin{aligned} & -1/2*((dx + c)*(2A - 3*B)/a - 2*(A*\tan(1/2*d*x + 1/2*c) - B*\tan(1/2*d*x \\ & + 1/2*c))/a - 2*(2*A*\tan(1/2*d*x + 1/2*c)^3 - 3*B*\tan(1/2*d*x + 1/2*c)^3 + \\ & 2*A*\tan(1/2*d*x + 1/2*c) - B*\tan(1/2*d*x + 1/2*c))/((\tan(1/2*d*x + 1/2*c) \\ & ^2 + 1)^2*a))/d \end{aligned}$$

3.40.9 Mupad [B] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.19

$$\begin{aligned} & \int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx \\ & = \frac{(2A - 3B) \tan(\frac{c}{2} + \frac{dx}{2})^3 + (2A - B) \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(a \tan(\frac{c}{2} + \frac{dx}{2})^4 + 2a \tan(\frac{c}{2} + \frac{dx}{2})^2 + a \right)} \\ & \quad - \frac{x(2A - 3B)}{2a} + \frac{\tan(\frac{c}{2} + \frac{dx}{2})(A - B)}{ad} \end{aligned}$$

3.40. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`

output `(tan(c/2 + (d*x)/2)^3*(2*A - 3*B) + tan(c/2 + (d*x)/2)*(2*A - B))/(d*(a + 2*a*tan(c/2 + (d*x)/2)^2 + a*tan(c/2 + (d*x)/2)^4) - (x*(2*A - 3*B))/(2*a) + (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.41 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.41.1	Optimal result	561
3.41.2	Mathematica [A] (verified)	561
3.41.3	Rubi [A] (verified)	562
3.41.4	Maple [A] (verified)	564
3.41.5	Fricas [A] (verification not implemented)	564
3.41.6	Sympy [B] (verification not implemented)	565
3.41.7	Maxima [B] (verification not implemented)	565
3.41.8	Giac [A] (verification not implemented)	566
3.41.9	Mupad [B] (verification not implemented)	566

3.41.1 Optimal result

Integrand size = 29, antiderivative size = 54

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx = \frac{(A - B)x}{a} + \frac{B \sin(c + dx)}{ad} - \frac{(A - B) \sin(c + dx)}{ad(1 + \cos(c + dx))}$$

output `(A-B)*x/a+B*sin(d*x+c)/a/d-(A-B)*sin(d*x+c)/a/d/(1+cos(d*x+c))`

3.41.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.72

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx = \frac{B \sin(c + dx)}{ad} + (A - B) \left(-\frac{\sin(c + dx)}{ad(1 + \cos(c + dx))} - \frac{\arcsin(\cos(c + dx)) \sin(c + dx)}{ad\sqrt{1 - \cos(c + dx)}\sqrt{1 + \cos(c + dx)}} \right)$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `(B*Sin[c + d*x])/(a*d) + (A - B)*(-(Sin[c + d*x]/(a*d*(1 + Cos[c + d*x]))) - (ArcSin[Cos[c + d*x]]*Sin[c + d*x])/(a*d*Sqrt[1 - Cos[c + d*x]]*Sqrt[1 + Cos[c + d*x]]))`

3.41.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.87, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a\cos(c+dx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{a\sin(c+dx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{a\cos(c+dx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{a\sin(c+dx+\frac{\pi}{2})+a} dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{\int \frac{(A-B)\cos(c+dx)}{\cos(c+dx)+1} dx}{a} + \frac{B\sin(c+dx)}{ad} \\
 & \quad \downarrow \text{27} \\
 & \frac{(A-B)\int \frac{\cos(c+dx)}{\cos(c+dx)+1} dx}{a} + \frac{B\sin(c+dx)}{ad} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A-B)\int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})+1} dx}{a} + \frac{B\sin(c+dx)}{ad} \\
 & \quad \downarrow \text{3214} \\
 & \frac{(A-B)\left(x - \int \frac{1}{\cos(c+dx)+1} dx\right)}{a} + \frac{B\sin(c+dx)}{ad} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{(A - B) \left(x - \int \frac{1}{\sin(c+dx+\frac{\pi}{2})+1} dx \right)}{a} + \frac{B \sin(c + dx)}{ad}$$

↓ 3127

$$\frac{(A - B) \left(x - \frac{\sin(c+dx)}{d(\cos(c+dx)+1)} \right)}{a} + \frac{B \sin(c + dx)}{ad}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `(B*Sin[c + d*x])/(a*d) + ((A - B)*(x - Sin[c + d*x]/(d*(1 + Cos[c + d*x])))`
`)/a`

3.41.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.41.4 Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

method	result
parallelrisc	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)(B \cos(dx+c) - A + 2B) + dx(A-B)}{ad}$
derivativedivides	$\frac{-A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + 2(A-B) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$
default	$\frac{-A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + 2(A-B) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$
risc	$\frac{x A}{a} - \frac{B x}{a} - \frac{ie^{i(dx+c)} B}{2ad} + \frac{ie^{-i(dx+c)} B}{2ad} - \frac{2i A}{da(e^{i(dx+c)} + 1)} + \frac{2i B}{da(e^{i(dx+c)} + 1)}$
norman	$\frac{\frac{(A-B)x}{a} + \frac{(A-B)x \tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)}{a} - \frac{(A-3B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} - \frac{2(A-2B) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} - \frac{(A-B) \tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} + \frac{2(A-B)x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{a}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)
```

```
output (tan(1/2*d*x+1/2*c)*(B*cos(d*x+c)-A+2*B)+d*x*(A-B))/a/d
```

3.41.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.13

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx$$

$$= \frac{(A - B)dx \cos(dx + c) + (A - B)dx + (B \cos(dx + c) - A + 2B) \sin(dx + c)}{ad \cos(dx + c) + ad}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="fracas")
```

3.41.
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$$

output $((A - B)*d*x*cos(d*x + c) + (A - B)*d*x + (B*cos(d*x + c) - A + 2*B)*sin(d*x + c))/(a*d*cos(d*x + c) + a*d)$

3.41.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 264 vs. 2(39) = 78.

Time = 0.57 (sec) , antiderivative size = 264, normalized size of antiderivative = 4.89

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx$$

$$= \left\{ \begin{array}{l} \frac{Adx \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{ad \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} + \frac{Adx}{ad \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} - \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{ad \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} - \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{ad \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} - \frac{Bdx \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{ad \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} - \frac{Bdx}{ad \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} \\ \frac{x(A+B \cos(c)) \cos(c)}{a \cos(c) + a} \end{array} \right.$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)), x)`

output `Piecewise((A*d*x*tan(c/2 + d*x/2)**2/(a*d*tan(c/2 + d*x/2)**2 + a*d) + A*d*x/(a*d*tan(c/2 + d*x/2)**2 + a*d) - A*tan(c/2 + d*x/2)**3/(a*d*tan(c/2 + d*x/2)**2 + a*d) - A*tan(c/2 + d*x/2)/(a*d*tan(c/2 + d*x/2)**2 + a*d) - B*d*x*tan(c/2 + d*x/2)**2/(a*d*tan(c/2 + d*x/2)**2 + a*d) - B*d*x/(a*d*tan(c/2 + d*x/2)**2 + a*d) + B*tan(c/2 + d*x/2)**3/(a*d*tan(c/2 + d*x/2)**2 + a*d) + 3*B*tan(c/2 + d*x/2)/(a*d*tan(c/2 + d*x/2)**2 + a*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)/(a*cos(c) + a), True))`

3.41.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(54) = 108.

Time = 0.30 (sec) , antiderivative size = 143, normalized size of antiderivative = 2.65

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx =$$

$$\frac{B \left(\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} - \frac{2 \sin(dx+c)}{\left(a + \frac{a \sin(dx+c)^2}{(\cos(dx+c)+1)^2}\right)(\cos(dx+c)+1)} - \frac{\sin(dx+c)}{a(\cos(dx+c)+1)} \right) - A \left(\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} - \frac{\sin(dx+c)}{a(\cos(dx+c)+1)} \right)}{d}$$

3.41. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `-(B*(2*arctan(sin(d*x + c)/(cos(d*x + c) + 1)))/a - 2*sin(d*x + c)/((a + a*sin(d*x + c)^2/(cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) - sin(d*x + c)/(a*(cos(d*x + c) + 1))) - A*(2*arctan(sin(d*x + c)/(cos(d*x + c) + 1)))/a - sin(d*x + c)/(a*(cos(d*x + c) + 1)))/d`

3.41.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx$$

$$= \frac{\frac{(dx+c)(A-B)}{a} - \frac{A \tan(\frac{1}{2} dx + \frac{1}{2} c) - B \tan(\frac{1}{2} dx + \frac{1}{2} c)}{a} + \frac{2 B \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)a}}{d}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `((d*x + c)*(A - B)/a - (A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a + 2*B*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 + 1)*a))/d`

3.41.9 Mupad [B] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.20

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{a + a \cos(c + dx)} dx = \frac{x(A - B)}{a} + \frac{2 B \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(a \tan^2(\frac{c}{2} + \frac{dx}{2}) + a \right)}$$

$$- \frac{\tan(\frac{c}{2} + \frac{dx}{2})(A - B)}{a d}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`

output `(x*(A - B))/a + (2*B*tan(c/2 + (d*x)/2))/(d*(a + a*tan(c/2 + (d*x)/2)^2)) - (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.41. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.42 $\int \frac{A+B \cos(c+dx)}{a+a \cos(c+dx)} dx$

3.42.1	Optimal result	567
3.42.2	Mathematica [B] (verified)	567
3.42.3	Rubi [A] (verified)	568
3.42.4	Maple [A] (verified)	569
3.42.5	Fricas [A] (verification not implemented)	569
3.42.6	Sympy [A] (verification not implemented)	570
3.42.7	Maxima [B] (verification not implemented)	570
3.42.8	Giac [A] (verification not implemented)	570
3.42.9	Mupad [B] (verification not implemented)	571

3.42.1 Optimal result

Integrand size = 23, antiderivative size = 34

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = \frac{Bx}{a} + \frac{(A - B) \sin(c + dx)}{d(a + a \cos(c + dx))}$$

output `B*x/a+(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))`

3.42.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 76 vs. 2(34) = 68.

Time = 0.25 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.24

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = -\frac{\sin(c + dx) \left(B \arcsin(\cos(c + dx))(1 + \cos(c + dx)) + (-A + B) \sqrt{\sin^2(c + dx)} \right)}{ad \sqrt{1 - \cos(c + dx)} (1 + \cos(c + dx))^{3/2}}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x]),x]`

output `-((Sin[c + d*x]*(B*ArcSin[Cos[c + d*x]]*(1 + Cos[c + d*x]) + (-A + B)*Sqrt[Sin[c + d*x]^2]))/(a*d*Sqrt[1 - Cos[c + d*x]]*(1 + Cos[c + d*x])^(3/2))`

3.42.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{a \cos(c + dx) + a} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} dx \\ & \quad \downarrow \text{3214} \\ & (A - B) \int \frac{1}{\cos(c + dx)a + a} dx + \frac{Bx}{a} \\ & \quad \downarrow \text{3042} \\ & (A - B) \int \frac{1}{\sin\left(c + dx + \frac{\pi}{2}\right)a + a} dx + \frac{Bx}{a} \\ & \quad \downarrow \text{3127} \\ & \frac{(A - B) \sin(c + dx)}{d(a \cos(c + dx) + a)} + \frac{Bx}{a} \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x]),x]`

output `(B*x)/a + ((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))`

3.42.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3127 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*SIN[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

```
rule 3214 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)
)*(x_)], x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d
*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

3.42.4 Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

method	result	size
parallelrisch	$\frac{dx B + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)(A - B)}{ad}$	28
derivativedivides	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2B \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$	45
default	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 2B \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$	45
risch	$\frac{Bx}{a} + \frac{2iA}{da(e^{i(dx+c)}+1)} - \frac{2iB}{da(e^{i(dx+c)}+1)}$	54
norman	$\frac{\frac{Bx}{a} + \frac{Bx \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} + \frac{(A - B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} + \frac{(A - B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{ad}}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)}$	85

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a), x, method=_RETURNVERBOSE)
```

```
output (d*x*B+tan(1/2*d*x+1/2*c)*(A-B))/a/d
```

3.42.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.26

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = \frac{B dx \cos(dx + c) + B dx + (A - B) \sin(dx + c)}{ad \cos(dx + c) + ad}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c)), x, algorithm="fricas")
```

```
output (B*d*x*cos(d*x + c) + B*d*x + (A - B)*sin(d*x + c))/(a*d*cos(d*x + c) + a*d)
```


3.42.6 Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.44

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = \begin{cases} \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{ad} + \frac{Bx}{a} - \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{ad} & \text{for } d \neq 0 \\ \frac{x(A+B \cos(c))}{a \cos(c)+a} & \text{otherwise} \end{cases}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x)`

output `Piecewise((A*tan(c/2 + d*x/2)/(a*d) + B*x/a - B*tan(c/2 + d*x/2)/(a*d), Ne(d, 0)), (x*(A + B*cos(c))/(a*cos(c) + a), True))`

3.42.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(34) = 68.

Time = 0.29 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.15

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = \frac{B \left(\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a} - \frac{\sin(dx+c)}{a(\cos(dx+c)+1)} \right) + \frac{A \sin(dx+c)}{a(\cos(dx+c)+1)}}{d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `(B*(2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a - sin(d*x + c)/(a*(cos(d*x + c) + 1))) + A*sin(d*x + c)/(a*(cos(d*x + c) + 1)))/d`

3.42.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.26

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = \frac{\frac{(dx+c)B}{a} + \frac{A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{a}}{d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `((d*x + c)*B/a + (A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a)/d`

3.42.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \frac{A + B \cos(c + dx)}{a + a \cos(c + dx)} dx = \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)(A-B)}{a} + \frac{B dx}{a}$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x)),x)`

output `((tan(c/2 + (d*x)/2)*(A - B))/a + (B*d*x)/a)/d`

3.43 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{a+a \cos(c+dx)} dx$

3.43.1 Optimal result	572
3.43.2 Mathematica [B] (verified)	572
3.43.3 Rubi [A] (verified)	573
3.43.4 Maple [A] (verified)	574
3.43.5 Fricas [A] (verification not implemented)	575
3.43.6 Sympy [F]	575
3.43.7 Maxima [B] (verification not implemented)	576
3.43.8 Giac [A] (verification not implemented)	576
3.43.9 Mupad [B] (verification not implemented)	577

3.43.1 Optimal result

Integrand size = 29, antiderivative size = 44

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx = \frac{A \operatorname{arctanh}(\sin(c + dx))}{ad} - \frac{(A - B) \sin(c + dx)}{d(a + a \cos(c + dx))}$$

output `A*arctanh(sin(d*x+c))/a/d-(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))`

3.43.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 109 vs. 2(44) = 88.

Time = 0.47 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.48

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx = \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) \left(A \cos\left(\frac{1}{2}(c + dx)\right) \left(-\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) + (-A + B) \operatorname{Sec}\left[\frac{c}{2}\right] \operatorname{Sin}\left[\frac{d x}{2}\right]}{ad(1 + \cos(c + dx))}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x]),x]`

output `(2*Cos[(c + d*x)/2]*(A*Cos[(c + d*x)/2]*(-Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + (-A + B)*Sec[c/2]*Sin[(d*x)/2))/(a*d*(1 + Cos[c + d*x]))`

3.43.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.172$, Rules used = {3042, 3457, 27, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(A+B\cos(c+dx))}{a\cos(c+dx)+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a\sin(c+dx+\frac{\pi}{2})+a)} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int aA\sec(c+dx)dx}{a^2} - \frac{(A-B)\sin(c+dx)}{d(a\cos(c+dx)+a)} \\
 & \quad \downarrow \text{27} \\
 & \frac{A\int\sec(c+dx)dx}{a} - \frac{(A-B)\sin(c+dx)}{d(a\cos(c+dx)+a)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A\int\csc(c+dx+\frac{\pi}{2})dx}{a} - \frac{(A-B)\sin(c+dx)}{d(a\cos(c+dx)+a)} \\
 & \quad \downarrow \text{4257} \\
 & \frac{A\operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{(A-B)\sin(c+dx)}{d(a\cos(c+dx)+a)}
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x]),x]`

output `(A*ArcTanh[Sin[c + d*x]])/(a*d) - ((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))`

3.43.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

3.43.4 Maple [A] (verified)

Time = 1.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.23

method	result	size
parallelrisch	$\frac{-A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \tan\left(\frac{dx}{2} + \frac{c}{2}\right)(A - B)}{ad}$	54
derivativedivides	$\frac{-A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{da}$	61
default	$\frac{-A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{da}$	61
risch	$-\frac{2iA}{da(e^{i(dx+c)}+1)} + \frac{2iB}{da(e^{i(dx+c)}+1)} + \frac{A \ln(e^{i(dx+c)}+i)}{ad} - \frac{A \ln(e^{i(dx+c)}-i)}{ad}$	91
norman	$\frac{\frac{(A-B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} - \frac{(A-B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{ad}}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{ad} - \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{ad}$	106

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)
```

3.43.
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{a+a \cos(c+dx)} dx$$

output $(-A \ln(\tan(1/2 dx + 1/2 c) - 1) + A \ln(\tan(1/2 dx + 1/2 c) + 1) - \tan(1/2 dx + 1/2 c) * (A - B)) / a / d$

3.43.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.68

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{(A \cos(dx + c) + A) \log(\sin(dx + c) + 1) - (A \cos(dx + c) + A) \log(-\sin(dx + c) + 1) - 2(A - B) \sin(dx + c)}{2(ad \cos(dx + c) + ad)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output $1/2*((A \cos(dx + c) + A) \log(\sin(dx + c) + 1) - (A \cos(dx + c) + A) \log(-\sin(dx + c) + 1) - 2*(A - B) \sin(dx + c)) / (a*d \cos(dx + c) + a*d)$

3.43.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx = \frac{\int \frac{A \sec(c + dx)}{\cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec(c + dx)}{\cos(c + dx) + 1} dx}{a}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c)),x)`

output $(\text{Integral}(A \sec(c + d*x) / (\cos(c + d*x) + 1), x) + \text{Integral}(B \cos(c + d*x) * \sec(c + d*x) / (\cos(c + d*x) + 1), x)) / a$

3.43.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(44) = 88$.

Time = 0.22 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.25

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{A \left(\frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a} - \frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a} - \frac{\sin(dx+c)}{a(\cos(dx+c)+1)} \right) + \frac{B \sin(dx+c)}{a(\cos(dx+c)+1)}}{d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `(A*(log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a - log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a - sin(d*x + c)/(a*(cos(d*x + c) + 1))) + B*sin(d*x + c)/(a*(cos(d*x + c) + 1)))/d`

3.43.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.61

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{\frac{A \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right)}{a} - \frac{A \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right)}{a} - \frac{A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{a}}{d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `(A*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a - A*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a - (A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a)/d`

3.43.9 Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.95

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + a \cos(c + dx)} dx = \frac{2 A \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{a d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A - B)}{a d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))),x)`

output `(2*A*atanh(tan(c/2 + (d*x)/2)))/(a*d) - (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.44 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{a+a \cos(c+dx)} dx$

3.44.1 Optimal result 578
 3.44.2 Mathematica [B] (verified) 578
 3.44.3 Rubi [A] (verified) 579
 3.44.4 Maple [A] (verified) 581
 3.44.5 Fricas [A] (verification not implemented) 582
 3.44.6 Sympy [F] 582
 3.44.7 Maxima [B] (verification not implemented) 582
 3.44.8 Giac [A] (verification not implemented) 583
 3.44.9 Mupad [B] (verification not implemented) 583

3.44.1 Optimal result

Integrand size = 31, antiderivative size = 69

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx = -\frac{(A - B) \operatorname{arctanh}(\sin(c + dx))}{ad} + \frac{(2A - B) \tan(c + dx)}{ad} - \frac{(A - B) \tan(c + dx)}{d(a + a \cos(c + dx))}$$

output `-(A-B)*arctanh(sin(d*x+c))/a/d+(2*A-B)*tan(d*x+c)/a/d-(A-B)*tan(d*x+c)/d/(a+a*cos(d*x+c))`

3.44.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 201 vs. 2(69) = 138.

Time = 1.37 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.91

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx = \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) \left((A - B) \sec\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right) + \cos\left(\frac{1}{2}(c + dx)\right) \left((A - B) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) - \sin\left(\frac{1}{2}(c - dx)\right) \right) \right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x]),x]`

output $(2*\text{Cos}[(c + d*x)/2]*((A - B)*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] + \text{Cos}[(c + d*x)/2]*((A - B)*(\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] - \text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2])) + (A*\text{Sin}[d*x])/((\text{Cos}[c/2] - \text{Sin}[c/2])*(\text{Cos}[c/2] + \text{Sin}[c/2]))*(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2])*(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2])))/((a*d*(1 + \text{Cos}[c + d*x])))$

3.44.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 3457, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2 (a \sin(c + dx + \frac{\pi}{2}) + a)} dx$$

↓ 3457

$$\frac{\int (a(2A - B) - a(A - B) \cos(c + dx)) \sec^2(c + dx) dx}{a^2} - \frac{(A - B) \tan(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3042

$$\frac{\int \frac{a(2A - B) - a(A - B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx}{a^2} - \frac{(A - B) \tan(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3227

$$\frac{a(2A - B) \int \sec^2(c + dx) dx - a(A - B) \int \sec(c + dx) dx}{a^2} - \frac{(A - B) \tan(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 3042

$$\frac{a(2A - B) \int \csc(c + dx + \frac{\pi}{2})^2 dx - a(A - B) \int \csc(c + dx + \frac{\pi}{2}) dx}{a^2} - \frac{(A - B) \tan(c + dx)}{d(a \cos(c + dx) + a)}$$

↓ 4254

$$\frac{-\frac{a(2A - B) \int 1 d(-\tan(c + dx))}{d} - a(A - B) \int \csc(c + dx + \frac{\pi}{2}) dx}{a^2} - \frac{(A - B) \tan(c + dx)}{d(a \cos(c + dx) + a)}$$

3.44. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx$

$$\begin{array}{c} \downarrow 24 \\ \frac{\frac{a(2A-B)\tan(c+dx)}{d} - a(A-B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx}{a^2} - \frac{(A-B)\tan(c+dx)}{d(a\cos(c+dx)+a)} \\ \downarrow 4257 \\ \frac{\frac{a(2A-B)\tan(c+dx)}{d} - \frac{a(A-B)\operatorname{arctanh}(\sin(c+dx))}{d}}{a^2} - \frac{(A-B)\tan(c+dx)}{d(a\cos(c+dx)+a)} \end{array}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x]),x]`

output `-(((A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x]))) + (-((a*(A - B)*ArcTanh[Sin[c + d*x]])/d) + (a*(2*A - B)*Tan[c + d*x])/d)/a^2`

3.44.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.44.4 Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.35

method	result
parallelrisch	$\frac{(A-B) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - (A-B) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 2\left(\left(A - \frac{B}{2}\right) \cos(dx+c) + \frac{A}{2}\right) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad \cos(dx+c)}$
derivativedivides	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1} + (-A+B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1} + (A-B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$
default	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1} + (-A+B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1} + (A-B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{da}$
norman	$\frac{(A-B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad} - \frac{2A \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{ad} - \frac{(3A-B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad}\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(A-B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{ad} - \frac{(A-B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{ad}$
risch	$\frac{2i(A e^{2i(dx+c)} - B e^{2i(dx+c)} + A e^{i(dx+c)} + 2A - B)}{da(e^{2i(dx+c)} + 1)(e^{i(dx+c)} + 1)} - \frac{A \ln(e^{i(dx+c)} + i)}{ad} + \frac{\ln(e^{i(dx+c)} + i) B}{ad} + \frac{A \ln(e^{i(dx+c)} - i)}{ad} -$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a), x, method=_RETURNVERBOSE)`

output `((A-B)*cos(d*x+c)*ln(tan(1/2*d*x+1/2*c)-1)-(A-B)*cos(d*x+c)*ln(tan(1/2*d*x+1/2*c)+1)+2*((A-1/2*B)*cos(d*x+c)+1/2*A)*tan(1/2*d*x+1/2*c)/a/d/cos(d*x+c)`

3.44.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.84

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx =$$

$$\frac{((A - B) \cos(dx + c))^2 + (A - B) \cos(dx + c)) \log(\sin(dx + c) + 1) - ((A - B) \cos(dx + c))^2 + (A - B) \cos(dx + c)}{2(ad \cos(dx + c))^2 + ad \cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `-1/2*(((A - B)*cos(d*x + c)^2 + (A - B)*cos(d*x + c))*log(sin(d*x + c) + 1) - ((A - B)*cos(d*x + c)^2 + (A - B)*cos(d*x + c))*log(-sin(d*x + c) + 1) - 2*((2*A - B)*cos(d*x + c) + A)*sin(d*x + c))/(a*d*cos(d*x + c)^2 + a*d*cos(d*x + c))`

3.44.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx = \frac{\int \frac{A \sec^2(c + dx)}{\cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec^2(c + dx)}{\cos(c + dx) + 1} dx}{a}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c)),x)`

output `(Integral(A*sec(c + d*x)**2/(cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2/(cos(c + d*x) + 1), x))/a`

3.44.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(69) = 138.

Time = 0.23 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.84

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx =$$

$$\frac{A \left(\frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a} - \frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a} - \frac{2 \sin(dx+c)}{\left(a - \frac{a \sin(dx+c)^2}{(\cos(dx+c)+1)^2}\right) (\cos(dx+c)+1)} - \frac{\sin(dx+c)}{a(\cos(dx+c)+1)} \right) - B \left(\frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)}\right)}{a} \right)}{d}$$

3.44. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{a+a \cos(c+dx)} dx$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `-(A*(log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a - log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a - 2*sin(d*x + c)/((a - a*sin(d*x + c)^2/(cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) - sin(d*x + c)/(a*(cos(d*x + c) + 1))) - B*(log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a - log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a - sin(d*x + c)/(a*(cos(d*x + c) + 1))))/d`

3.44.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.59

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx = \frac{\frac{(A-B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a} - \frac{(A-B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a} - \frac{A \tan(\frac{1}{2} dx + \frac{1}{2} c) - B \tan(\frac{1}{2} dx + \frac{1}{2} c)}{a} + \frac{2 A \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1)a}}{d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `-((A - B)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a - (A - B)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a - (A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a + 2*A*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 - 1)*a))/d`

3.44.9 Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.13

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + a \cos(c + dx)} dx = \frac{2 A \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(a - a \tan(\frac{c}{2} + \frac{dx}{2})^2 \right)} - \frac{2 a \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (A - B)}{a d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A - B)}{a d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))),x)`

output `(2*A*tan(c/2 + (d*x)/2))/(d*(a - a*tan(c/2 + (d*x)/2)^2)) - (2*atanh(tan(c/2 + (d*x)/2))*(A - B))/(a*d) + (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.45 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{a+a \cos(c+dx)} dx$

3.45.1	Optimal result	585
3.45.2	Mathematica [B] (verified)	585
3.45.3	Rubi [A] (verified)	586
3.45.4	Maple [A] (verified)	589
3.45.5	Fricas [A] (verification not implemented)	589
3.45.6	Sympy [F]	590
3.45.7	Maxima [B] (verification not implemented)	590
3.45.8	Giac [A] (verification not implemented)	591
3.45.9	Mupad [B] (verification not implemented)	591

3.45.1 Optimal result

Integrand size = 31, antiderivative size = 107

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx = \frac{(3A - 2B) \operatorname{arctanh}(\sin(c + dx))}{2ad} - \frac{2(A - B) \tan(c + dx)}{ad} + \frac{(3A - 2B) \sec(c + dx) \tan(c + dx)}{2ad} - \frac{(A - B) \sec(c + dx) \tan(c + dx)}{d(a + a \cos(c + dx))}$$

```
output 1/2*(3*A-2*B)*arctanh(sin(d*x+c))/a/d-2*(A-B)*tan(d*x+c)/a/d+1/2*(3*A-2*B)*sec(d*x+c)*tan(d*x+c)/a/d-(A-B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))
```

3.45.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 289 vs. 2(107) = 214.

Time = 2.94 (sec) , antiderivative size = 289, normalized size of antiderivative = 2.70

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx = \frac{\cos\left(\frac{1}{2}(c + dx)\right) \left(4(-A + B) \sec\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right) + \cos\left(\frac{1}{2}(c + dx)\right) \left((-6A + 4B) \log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) - \sin\right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x]),x]`

output `(Cos[(c + d*x)/2]*(4*(-A + B)*Sec[c/2]*Sin[(d*x)/2] + Cos[(c + d*x)/2]*((-6*A + 4*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 6*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - 4*B*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + A/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2 - A/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 - (4*(A - B)*Sin[d*x])/((Cos[c/2] - Sin[c/2])*(Cos[c/2] + Sin[c/2])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))))/(2*a*d*(1 + Cos[c + d*x]))`

3.45.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.93, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3457, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3 (a \sin(c + dx + \frac{\pi}{2}) + a)} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int (a(3A - 2B) - 2a(A - B) \cos(c + dx)) \sec^3(c + dx) dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(3A - 2B) - 2a(A - B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
 & \quad \downarrow \text{3227} \\
 & \frac{a(3A - 2B) \int \sec^3(c + dx) dx - 2a(A - B) \int \sec^2(c + dx) dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.45. $\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx$

$$\begin{aligned}
& \frac{a(3A - 2B) \int \csc(c + dx + \frac{\pi}{2})^3 dx - 2a(A - B) \int \csc(c + dx + \frac{\pi}{2})^2 dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow 4254 \\
& \frac{\frac{2a(A - B) \int \frac{1d(-\tan(c + dx))}{d} + a(3A - 2B) \int \csc(c + dx + \frac{\pi}{2})^3 dx}{a^2} - (A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow 24 \\
& \frac{a(3A - 2B) \int \csc(c + dx + \frac{\pi}{2})^3 dx - \frac{2a(A - B) \tan(c + dx)}{d}}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow 4255 \\
& \frac{a(3A - 2B) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{2a(A - B) \tan(c + dx)}{d}}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow 3042 \\
& \frac{a(3A - 2B) \left(\frac{1}{2} \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{2a(A - B) \tan(c + dx)}{d}}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow 4257 \\
& \frac{a(3A - 2B) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{2a(A - B) \tan(c + dx)}{d}}{a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{d(a \cos(c + dx) + a)}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x]),x]`

output `-(((A - B)*Sec[c + d*x]*Tan[c + d*x])/(d*(a + a*Cos[c + d*x]))) + ((-2*a*(A - B)*Tan[c + d*x])/d + a*(3*A - 2*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/a^2`

3.45.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.45.4 Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.18

method	result
parallelrisc	$\frac{-3(1+\cos(2dx+2c))(A-\frac{2B}{3})\ln(\tan(\frac{dx}{2}+\frac{c}{2})-1)+3(1+\cos(2dx+2c))(A-\frac{2B}{3})\ln(\tan(\frac{dx}{2}+\frac{c}{2})+1)-2\tan(\frac{dx}{2}+\frac{c}{2})}{2ad(1+\cos(2dx+2c))}$
derivativedivides	$\frac{-A\tan(\frac{dx}{2}+\frac{c}{2})+B\tan(\frac{dx}{2}+\frac{c}{2})-\frac{A}{2(\tan(\frac{dx}{2}+\frac{c}{2})+1)^2}-\frac{-\frac{3A}{2}+B}{\tan(\frac{dx}{2}+\frac{c}{2})+1}+(\frac{3A}{2}-B)\ln(\tan(\frac{dx}{2}+\frac{c}{2})+1)+\frac{A}{2(\tan(\frac{dx}{2}+\frac{c}{2}))}}{da}$
default	$\frac{-A\tan(\frac{dx}{2}+\frac{c}{2})+B\tan(\frac{dx}{2}+\frac{c}{2})-\frac{A}{2(\tan(\frac{dx}{2}+\frac{c}{2})+1)^2}-\frac{-\frac{3A}{2}+B}{\tan(\frac{dx}{2}+\frac{c}{2})+1}+(\frac{3A}{2}-B)\ln(\tan(\frac{dx}{2}+\frac{c}{2})+1)+\frac{A}{2(\tan(\frac{dx}{2}+\frac{c}{2}))}}{da}$
norman	$\frac{\frac{(3A-B)(\tan^3(\frac{dx}{2}+\frac{c}{2}))}{ad}+\frac{(4A-3B)(\tan^5(\frac{dx}{2}+\frac{c}{2}))}{ad}-\frac{(A-B)(\tan^7(\frac{dx}{2}+\frac{c}{2}))}{ad}-\frac{(2A-3B)\tan(\frac{dx}{2}+\frac{c}{2})}{ad}}{(1+\tan^2(\frac{dx}{2}+\frac{c}{2}))(\tan^2(\frac{dx}{2}+\frac{c}{2})-1)^2}-\frac{(3A-2B)\ln(\tan(\frac{dx}{2}+\frac{c}{2}))}{2ad}}$
risc	$\frac{i(3Ae^{4i(dx+c)}-2Be^{4i(dx+c)}+3Ae^{3i(dx+c)}-2Be^{3i(dx+c)}+5Ae^{2i(dx+c)}-6Be^{2i(dx+c)}+Ae^{i(dx+c)}-2Be^{i(dx+c)}+4)}{da(e^{i(dx+c)}+1)(e^{2i(dx+c)}+1)^2}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)
```

```
output 1/2*(-3*(1+cos(2*d*x+2*c))*(A-2/3*B)*ln(tan(1/2*d*x+1/2*c)-1)+3*(1+cos(2*d*x+2*c))*(A-2/3*B)*ln(tan(1/2*d*x+1/2*c)+1)-2*tan(1/2*d*x+1/2*c)*((-2*B+2*A)*cos(2*d*x+2*c)+(1+cos(d*x+c))*(A-2*B))/a/d/(1+cos(2*d*x+2*c))
```

3.45.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.46

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{((3A - 2B) \cos(dx + c))^3 + (3A - 2B) \cos(dx + c)^2 \log(\sin(dx + c) + 1) - ((3A - 2B) \cos(dx + c))^3}{4(ad \cos$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c)),x, algorithm="fricas")
```

output $1/4*((3*A - 2*B)*\cos(d*x + c)^3 + (3*A - 2*B)*\cos(d*x + c)^2)*\log(\sin(d*x + c) + 1) - ((3*A - 2*B)*\cos(d*x + c)^3 + (3*A - 2*B)*\cos(d*x + c)^2)*\log(-\sin(d*x + c) + 1) - 2*(4*(A - B)*\cos(d*x + c)^2 + (A - 2*B)*\cos(d*x + c) - A)*\sin(d*x + c)/(a*d*\cos(d*x + c)^3 + a*d*\cos(d*x + c)^2)$

3.45.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx = \frac{\int \frac{A \sec^3(c + dx)}{\cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec^3(c + dx)}{\cos(c + dx) + 1} dx}{a}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c)),x)`

output `(Integral(A*sec(c + d*x)**3/(cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**3/(cos(c + d*x) + 1), x))/a`

3.45.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 282 vs. $2(103) = 206$.

Time = 0.21 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.64

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx =$$

$$A \left(\frac{2 \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - \frac{3 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a - \frac{2a \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{a \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} - \frac{3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a} + \frac{3 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a} + \frac{2 \sin(dx+c)}{a(\cos(dx+c)+1)} \right) + 2B \left(\frac{\log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{2d} \right)$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output $-1/2*(A*(2*(\sin(d*x + c)/(\cos(d*x + c) + 1) - 3*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(a - 2*a*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + a*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4) - 3*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)/a + 3*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) - 1)/a + 2*\sin(d*x + c)/(a*(\cos(d*x + c) + 1))) + 2*B*(\log(\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)/a - \log(\sin(d*x + c)/(\cos(d*x + c) + 1) - 1)/a - 2*\sin(d*x + c)/((a - a*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2)*(\cos(d*x + c) + 1)) - \sin(d*x + c)/(a*(\cos(d*x + c) + 1))))/d$

3.45. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{a+a \cos(c+dx)} dx$

3.45.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.47

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{(3A-2B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a} - \frac{(3A-2B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a} - \frac{2(A \tan(\frac{1}{2} dx + \frac{1}{2} c) - B \tan(\frac{1}{2} dx + \frac{1}{2} c))}{a} + \frac{2(3A \tan(\frac{1}{2} dx + \frac{1}{2} c) - A \tan(\frac{1}{2} dx + \frac{1}{2} c) + 2B \tan(\frac{1}{2} dx + \frac{1}{2} c))}{2d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `1/2*((3*A - 2*B)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a - (3*A - 2*B)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a - 2*(A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a + 2*(3*A*tan(1/2*d*x + 1/2*c)^3 - 2*B*tan(1/2*d*x + 1/2*c)^3 - A*tan(1/2*d*x + 1/2*c) + 2*B*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^2*a))/d`

3.45.9 Mupad [B] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.11

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{\tan(\frac{c}{2} + \frac{dx}{2})^3 (3A - 2B) - \tan(\frac{c}{2} + \frac{dx}{2}) (A - 2B)}{d \left(a \tan(\frac{c}{2} + \frac{dx}{2})^4 - 2a \tan(\frac{c}{2} + \frac{dx}{2})^2 + a \right)} + \frac{2 \operatorname{atanh}(\tan(\frac{c}{2} + \frac{dx}{2})) \left(\frac{3A}{2} - B \right) - \tan(\frac{c}{2} + \frac{dx}{2}) (A - B)}{ad}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))),x)`

output `(tan(c/2 + (d*x)/2)^3*(3*A - 2*B) - tan(c/2 + (d*x)/2)*(A - 2*B))/(d*(a - 2*a*tan(c/2 + (d*x)/2)^2 + a*tan(c/2 + (d*x)/2)^4) + (2*atanh(tan(c/2 + (d*x)/2))*(3*A/2 - B))/(a*d) - (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.46
$$\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+a \cos(c+dx)} dx$$

3.46.1 Optimal result 592
 3.46.2 Mathematica [B] (verified) 593
 3.46.3 Rubi [A] (verified) 593
 3.46.4 Maple [A] (verified) 596
 3.46.5 Fricas [A] (verification not implemented) 597
 3.46.6 Sympy [F] 597
 3.46.7 Maxima [B] (verification not implemented) 597
 3.46.8 Giac [A] (verification not implemented) 598
 3.46.9 Mupad [B] (verification not implemented) 599

3.46.1 Optimal result

Integrand size = 31, antiderivative size = 131

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx = -\frac{3(A - B) \operatorname{arctanh}(\sin(c + dx))}{2ad} + \frac{(4A - 3B) \tan(c + dx)}{ad} - \frac{3(A - B) \sec(c + dx) \tan(c + dx)}{2ad} - \frac{(A - B) \sec^2(c + dx) \tan(c + dx)}{d(a + a \cos(c + dx))} + \frac{(4A - 3B) \tan^3(c + dx)}{3ad}$$

output

```
-3/2*(A-B)*arctanh(sin(d*x+c))/a/d+(4*A-3*B)*tan(d*x+c)/a/d-3/2*(A-B)*sec(d*x+c)*tan(d*x+c)/a/d-(A-B)*sec(d*x+c)^2*tan(d*x+c)/d/(a+a*cos(d*x+c))+1/3*(4*A-3*B)*tan(d*x+c)^3/a/d
```

3.46.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 490 vs. $2(131) = 262$.

Time = 3.70 (sec) , antiderivative size = 490, normalized size of antiderivative = 3.74

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c + dx)\right) \left(144(A - B) \cos\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) + \sec\left(\frac{c}{2}\right) \sec^3\left(\frac{c + dx}{2}\right) \left(6(A + B) \sin\left(\frac{dx}{2}\right) + 3(13A - 9B) \sin\left(\frac{3dx}{2}\right) - 24A \sin\left[c - \frac{dx}{2}\right] + 12B \sin\left[c - \frac{dx}{2}\right] - 6A \sin\left[c + \frac{dx}{2}\right] + 6B \sin\left[c + \frac{dx}{2}\right] - 24A \sin\left[2c + \frac{dx}{2}\right] + 24B \sin\left[2c + \frac{dx}{2}\right] + 21A \sin\left[c + \frac{3dx}{2}\right] - 9B \sin\left[c + \frac{3dx}{2}\right] + 9A \sin\left[2c + \frac{3dx}{2}\right] - 9B \sin\left[2c + \frac{3dx}{2}\right] - 9A \sin\left[3c + \frac{3dx}{2}\right] + 9B \sin\left[3c + \frac{3dx}{2}\right] + 7A \sin\left[c + \frac{5dx}{2}\right] - 3B \sin\left[c + \frac{5dx}{2}\right] + A \sin\left[2c + \frac{5dx}{2}\right] + 3B \sin\left[2c + \frac{5dx}{2}\right] - 3A \sin\left[3c + \frac{5dx}{2}\right] + 3B \sin\left[3c + \frac{5dx}{2}\right] - 9A \sin\left[4c + \frac{5dx}{2}\right] + 9B \sin\left[4c + \frac{5dx}{2}\right] + 16A \sin\left[2c + \frac{7dx}{2}\right] - 12B \sin\left[2c + \frac{7dx}{2}\right] + 10A \sin\left[3c + \frac{7dx}{2}\right] - 6B \sin\left[3c + \frac{7dx}{2}\right] + 6A \sin\left[4c + \frac{7dx}{2}\right] - 6B \sin\left[4c + \frac{7dx}{2}\right])}{48ad(1 + \cos(c + dx))}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + a*Cos[c + d*x]),x]`

output `(Cos[(c + d*x)/2]*(144*(A - B)*Cos[(c + d*x)/2]*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Sec[c/2]*Sec[c]*Sec[c + d*x]^3*(6*(A + B)*Sin[(d*x)/2] + 3*(13*A - 9*B)*Sin[(3*d*x)/2] - 24*A*Sin[c - (d*x)/2] + 12*B*Sin[c - (d*x)/2] - 6*A*Sin[c + (d*x)/2] + 6*B*Sin[c + (d*x)/2] - 24*A*Sin[2*c + (d*x)/2] + 24*B*Sin[2*c + (d*x)/2] + 21*A*Sin[c + (3*d*x)/2] - 9*B*Sin[c + (3*d*x)/2] + 9*A*Sin[2*c + (3*d*x)/2] - 9*B*Sin[2*c + (3*d*x)/2] - 9*A*Sin[3*c + (3*d*x)/2] + 9*B*Sin[3*c + (3*d*x)/2] + 7*A*Sin[c + (5*d*x)/2] - 3*B*Sin[c + (5*d*x)/2] + A*Sin[2*c + (5*d*x)/2] + 3*B*Sin[2*c + (5*d*x)/2] - 3*A*Sin[3*c + (5*d*x)/2] + 3*B*Sin[3*c + (5*d*x)/2] - 9*A*Sin[4*c + (5*d*x)/2] + 9*B*Sin[4*c + (5*d*x)/2] + 16*A*Sin[2*c + (7*d*x)/2] - 12*B*Sin[2*c + (7*d*x)/2] + 10*A*Sin[3*c + (7*d*x)/2] - 6*B*Sin[3*c + (7*d*x)/2] + 6*A*Sin[4*c + (7*d*x)/2] - 6*B*Sin[4*c + (7*d*x)/2]))/(48*a*d*(1 + Cos[c + d*x]))`

3.46.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.90, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3457, 3042, 3227, 3042, 4254, 2009, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^4(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

3.46. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^4 (a \sin(c + dx + \frac{\pi}{2}) + a)} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int (a(4A - 3B) - 3a(A - B) \cos(c + dx)) \sec^4(c + dx) dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(4A - 3B) - 3a(A - B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^4} dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{3227} \\
& \frac{a(4A - 3B) \int \sec^4(c + dx) dx - 3a(A - B) \int \sec^3(c + dx) dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(4A - 3B) \int \csc(c + dx + \frac{\pi}{2})^4 dx - 3a(A - B) \int \csc(c + dx + \frac{\pi}{2})^3 dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{4254} \\
& \frac{-\frac{a(4A - 3B) \int (\tan^2(c + dx) + 1) d(-\tan(c + dx))}{d} - 3a(A - B) \int \csc(c + dx + \frac{\pi}{2})^3 dx}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{2009} \\
& \frac{-3a(A - B) \int \csc(c + dx + \frac{\pi}{2})^3 dx - \frac{a(4A - 3B)(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d}}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{4255} \\
& \frac{-3a(A - B) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{a(4A - 3B)(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d}}{a^2} - \frac{(A - B) \tan(c + dx) \sec^2(c + dx)}{d(a \cos(c + dx) + a)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.46. $\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx$

$$\frac{-3a(A-B) \left(\frac{1}{2} \int \csc(c+dx + \frac{\pi}{2}) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{a(4A-3B)(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d}}{a^2 \frac{(A-B) \tan(c+dx) \sec^2(c+dx)}{d(a \cos(c+dx) + a)}}{-}$$

↓ 4257

$$\frac{-3a(A-B) \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{a(4A-3B)(-\frac{1}{3} \tan^3(c+dx) - \tan(c+dx))}{d}}{a^2 \frac{(A-B) \tan(c+dx) \sec^2(c+dx)}{d(a \cos(c+dx) + a)}}{-}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + a*Cos[c + d*x]),x]`

output `-(((A - B)*Sec[c + d*x]^2*Tan[c + d*x]/(d*(a + a*Cos[c + d*x]))) + (-3*a*(A - B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)) - (a*(4*A - 3*B)*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d)/a^2`

3.46.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.46. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+a \cos(c+dx)} dx$

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.46.4 Maple [A] (verified)

Time = 1.40 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.30

method	result
parallelrisch	$\frac{27\left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c)\right)(A-B)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - 27\left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c)\right)(A-B)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 44\left(\frac{1}{11} * (4 * A - 3 * B) * \cos(3 * dx + 3 * c) + 1\right) / (22 * (7 * A - 3 * B) * \cos(2 * dx + 2 * c) + (A - 6 / 11 * B) * \cos(dx + c) + 1) * \tan(1/2 * dx + 1/2 * c)}{6ad(\cos(3dx+3c)+3\cos(dx+c))}$
derivativedivides	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{A}{3\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} - \frac{2A - B}{2\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} + \left(\frac{3A}{2} - \frac{3B}{2}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - \frac{\frac{5A}{2} - \frac{3B}{2}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}}{da}$
default	$\frac{A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{A}{3\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3} - \frac{2A - B}{2\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} + \left(\frac{3A}{2} - \frac{3B}{2}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - \frac{\frac{5A}{2} - \frac{3B}{2}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}}{da}$
norman	$\frac{(A-B)\left(\frac{\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad}\right) + \frac{(A-3B)\left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{3ad}\right) - 2(2A-B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - (7A-5B)\left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad}\right) + \frac{(13A-15B)\left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3ad}\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3}}{ad}$
risch	$\frac{i(9Ae^{6i(dx+c)} - 9Be^{6i(dx+c)} + 9Ae^{5i(dx+c)} - 9Be^{5i(dx+c)} + 24Ae^{4i(dx+c)} - 24Be^{4i(dx+c)} + 24Ae^{3i(dx+c)} - 12Be^{3i(dx+c)} - 12Ae^{2i(dx+c)} + 12Be^{2i(dx+c)} - 12Ae^{i(dx+c)} + 12Be^{i(dx+c)} - 12A + 12B)}{3da(e^{2i(dx+c)} + 1)^3(e^{i(dx+c)} + 1)}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+cos(d*x+c)*a), x, method=_RETURNVERBOSE)`

output `1/6*(27*(1/3*cos(3*d*x+3*c)+cos(d*x+c))*(A-B)*ln(tan(1/2*d*x+1/2*c)-1)-27*(1/3*cos(3*d*x+3*c)+cos(d*x+c))*(A-B)*ln(tan(1/2*d*x+1/2*c)+1)+44*(1/11*(4*A-3*B)*cos(3*d*x+3*c)+1/22*(7*A-3*B)*cos(2*d*x+2*c)+(A-6/11*B)*cos(d*x+c)+1/2*A-3/22*B)*tan(1/2*d*x+1/2*c))/a/d/(cos(3*d*x+3*c)+3*cos(d*x+c))`

$$3.46. \int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+a \cos(c+dx)} dx$$

3.46.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.28

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx =$$

$$\frac{9((A - B) \cos(dx + c)^4 + (A - B) \cos(dx + c)^3) \log(\sin(dx + c) + 1) - 9((A - B) \cos(dx + c)^4 + (A - B) \cos(dx + c)^3) \log(-\sin(dx + c) + 1) - 2(4(4A - 3B) \cos(dx + c)^3 + (7A - 3B) \cos(dx + c)^2 - (A - 3B) \cos(dx + c) + 2A) \sin(dx + c)}{a^2 d \cos^4(dx + c) + a^2 d \cos^3(dx + c)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `-1/12*(9*((A - B)*cos(d*x + c)^4 + (A - B)*cos(d*x + c)^3)*log(sin(d*x + c) + 1) - 9*((A - B)*cos(d*x + c)^4 + (A - B)*cos(d*x + c)^3)*log(-sin(d*x + c) + 1) - 2*(4*(4*A - 3*B)*cos(d*x + c)^3 + (7*A - 3*B)*cos(d*x + c)^2 - (A - 3*B)*cos(d*x + c) + 2*A)*sin(d*x + c))/(a*d*cos(d*x + c)^4 + a*d*cos(d*x + c)^3)`

3.46.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx = \frac{\int \frac{A \sec^4(c + dx)}{\cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec^4(c + dx)}{\cos(c + dx) + 1} dx}{a}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**4/(a+a*cos(d*x+c)),x)`

output `(Integral(A*sec(c + d*x)**4/(cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**4/(cos(c + d*x) + 1), x))/a`

3.46.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 368 vs. 2(125) = 250.

Time = 0.22 (sec) , antiderivative size = 368, normalized size of antiderivative = 2.81

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{A \left(\frac{2 \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} - \frac{16 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{15 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a - \frac{3a \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3a \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - \frac{a \sin(dx+c)^6}{(\cos(dx+c)+1)^6}} - \frac{9 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a} + \frac{9 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a} + \frac{6 \sin(dx+c)}{a(\cos(dx+c)+1)} \right) - 3B}{6d}$$

3.46. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+a \cos(c+dx)} dx$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `1/6*(A*(2*(9*sin(d*x + c)/(cos(d*x + c) + 1) - 16*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 15*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(a - 3*a*sin(d*x + c))^2/(cos(d*x + c) + 1)^2 + 3*a*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - a*sin(d*x + c)^6/(cos(d*x + c) + 1)^6) - 9*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a + 9*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a + 6*sin(d*x + c)/(a*(cos(d*x + c) + 1))) - 3*B*(2*(sin(d*x + c)/(cos(d*x + c) + 1) - 3*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(a - 2*a*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a*sin(d*x + c)^4/(cos(d*x + c) + 1)^4) - 3*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a + 3*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a + 2*sin(d*x + c)/(a*(cos(d*x + c) + 1))))/d`

3.46.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.39

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx =$$

$$\frac{9(A-B) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right)}{a} - \frac{9(A-B) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right)}{a} - \frac{6(A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right))}{a} + \frac{2\left(15A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 15A \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15B \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 15A \tan^5\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15B \tan^5\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 15A \tan^7\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15B \tan^7\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 15A \tan^9\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15B \tan^9\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)}{6d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `-1/6*(9*(A - B)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a - 9*(A - B)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a - 6*(A*tan(1/2*d*x + 1/2*c) - B*tan(1/2*d*x + 1/2*c))/a + 2*(15*A*tan(1/2*d*x + 1/2*c)^5 - 9*B*tan(1/2*d*x + 1/2*c)^5 - 16*A*tan(1/2*d*x + 1/2*c)^3 + 12*B*tan(1/2*d*x + 1/2*c)^3 + 9*A*tan(1/2*d*x + 1/2*c) - 3*B*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 - 1)^3*a))/d`

3.46.9 Mupad [B] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.16

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{(5A - 3B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(4B - \frac{16A}{3}\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (3A - B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(-a \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 3a \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 3a \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a\right)}$$

$$- \frac{3a \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (A - B)}{ad} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A - B)}{ad}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^4*(a + a*cos(c + d*x))),x)`output `(tan(c/2 + (d*x)/2)^5*(5*A - 3*B) - tan(c/2 + (d*x)/2)^3*((16*A)/3 - 4*B) + tan(c/2 + (d*x)/2)*(3*A - B))/(d*(a - 3*a*tan(c/2 + (d*x)/2)^2 + 3*a*tan(c/2 + (d*x)/2)^4 - a*tan(c/2 + (d*x)/2)^6) - (3*atanh(tan(c/2 + (d*x)/2))*(A - B))/(a*d) + (tan(c/2 + (d*x)/2)*(A - B))/(a*d)`

3.47
$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.47.1 Optimal result 600
 3.47.2 Mathematica [B] (verified) 601
 3.47.3 Rubi [A] (verified) 601
 3.47.4 Maple [A] (verified) 605
 3.47.5 Fricas [A] (verification not implemented) 605
 3.47.6 Sympy [B] (verification not implemented) 606
 3.47.7 Maxima [B] (verification not implemented) 607
 3.47.8 Giac [A] (verification not implemented) 607
 3.47.9 Mupad [B] (verification not implemented) 608

3.47.1 Optimal result

Integrand size = 31, antiderivative size = 170

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = \frac{(7A-10B)x}{2a^2} - \frac{4(2A-3B) \sin(c+dx)}{a^2d} + \frac{(7A-10B) \cos(c+dx) \sin(c+dx)}{2a^2d} + \frac{(7A-10B) \cos^3(c+dx) \sin(c+dx)}{3a^2d(1+\cos(c+dx))} + \frac{(A-B) \cos^4(c+dx) \sin(c+dx)}{3d(a+a \cos(c+dx))^2} + \frac{4(2A-3B) \sin^3(c+dx)}{3a^2d}$$

```
output 1/2*(7*A-10*B)*x/a^2-4*(2*A-3*B)*sin(d*x+c)/a^2/d+1/2*(7*A-10*B)*cos(d*x+c)*sin(d*x+c)/a^2/d+1/3*(7*A-10*B)*cos(d*x+c)^3*sin(d*x+c)/a^2/d/(1+cos(d*x+c))+1/3*(A-B)*cos(d*x+c)^4*sin(d*x+c)/d/(a+a*cos(d*x+c))^2+4/3*(2*A-3*B)*sin(d*x+c)^3/a^2/d
```

3.47.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 369 vs. $2(170) = 340$.

Time = 1.81 (sec) , antiderivative size = 369, normalized size of antiderivative = 2.17

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) (36(7A-10B)dx \cos\left(\frac{dx}{2}\right) + 36(7A-10B)dx \cos\left(c+\frac{dx}{2}\right) + 84Adx \cos\left(c+\frac{3dx}{2}\right))}{(48a^2d(1+\cos(c+dx))^2)}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(36*(7*A - 10*B)*d*x*Cos[(d*x)/2] + 36*(7*A - 10*B)*d*x*Cos[c + (d*x)/2] + 84*A*d*x*Cos[c + (3*d*x)/2] - 120*B*d*x*Cos[c + (3*d*x)/2] + 84*A*d*x*Cos[2*c + (3*d*x)/2] - 120*B*d*x*Cos[2*c + (3*d*x)/2] - 381*A*Sin[(d*x)/2] + 516*B*Sin[(d*x)/2] + 147*A*Sin[c + (d*x)/2] - 156*B*Sin[c + (d*x)/2] - 239*A*Sin[c + (3*d*x)/2] + 342*B*Sin[c + (3*d*x)/2] - 63*A*Sin[2*c + (3*d*x)/2] + 118*B*Sin[2*c + (3*d*x)/2] - 15*A*Sin[2*c + (5*d*x)/2] + 30*B*Sin[2*c + (5*d*x)/2] - 15*A*Sin[3*c + (5*d*x)/2] + 30*B*Sin[3*c + (5*d*x)/2] + 3*A*Sin[3*c + (7*d*x)/2] - 3*B*Sin[3*c + (7*d*x)/2] + 3*A*Sin[4*c + (7*d*x)/2] - 3*B*Sin[4*c + (7*d*x)/2] + B*Sin[4*c + (9*d*x)/2] + B*Sin[5*c + (9*d*x)/2]))/(48*a^2*d*(1 + Cos[c + d*x])^2)`

3.47.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.94, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3456, 3042, 3456, 27, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^4 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a\sin\left(c+dx+\frac{\pi}{2}\right)+a)^2} dx$$

$$\downarrow \text{3456}$$

3.47. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{\int \frac{\cos^3(c+dx)(4a(A-B)-3a(A-2B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^3(4a(A-B)-3a(A-2B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3456} \\
& \frac{\int 3\cos^2(c+dx)(a^2(7A-10B)-4a^2(2A-3B)\cos(c+dx)) dx}{a^2} + \frac{(7A-10B)\sin(c+dx)\cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{3\int \cos^2(c+dx)(a^2(7A-10B)-4a^2(2A-3B)\cos(c+dx)) dx}{a^2} + \frac{(7A-10B)\sin(c+dx)\cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3\int \sin(c+dx+\frac{\pi}{2})^2(a^2(7A-10B)-4a^2(2A-3B)\sin(c+dx+\frac{\pi}{2})) dx}{a^2} + \frac{(7A-10B)\sin(c+dx)\cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3227} \\
& \frac{3(a^2(7A-10B)\int \cos^2(c+dx)dx-4a^2(2A-3B)\int \cos^3(c+dx)dx)}{a^2} + \frac{(7A-10B)\sin(c+dx)\cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3(a^2(7A-10B)\int \sin(c+dx+\frac{\pi}{2})^2 dx-4a^2(2A-3B)\int \sin(c+dx+\frac{\pi}{2})^3 dx)}{a^2} + \frac{(7A-10B)\sin(c+dx)\cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3113}
\end{aligned}$$

3.47. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{3 \left(\frac{4a^2(2A-3B) \int (1-\sin^2(c+dx)) d(-\sin(c+dx))}{d} + a^2(7A-10B) \int \sin(c+dx+\frac{\pi}{2})^2 dx \right)}{a^2} + \frac{(7A-10B) \sin(c+dx) \cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
 & \frac{3a^2}{(A-B) \sin(c+dx) \cos^4(c+dx)} \\
 & \frac{3d(a \cos(c+dx) + a)^2}{3d(a \cos(c+dx) + a)^2} \\
 & \quad \downarrow \text{2009} \\
 & \frac{3 \left(a^2(7A-10B) \int \sin(c+dx+\frac{\pi}{2})^2 dx + \frac{4a^2(2A-3B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d} \right)}{a^2} + \frac{(7A-10B) \sin(c+dx) \cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
 & \frac{3a^2}{(A-B) \sin(c+dx) \cos^4(c+dx)} \\
 & \frac{3d(a \cos(c+dx) + a)^2}{3d(a \cos(c+dx) + a)^2} \\
 & \quad \downarrow \text{3115} \\
 & \frac{3 \left(a^2(7A-10B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx) \cos(c+dx)}{2d} \right) + \frac{4a^2(2A-3B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d} \right)}{a^2} + \frac{(7A-10B) \sin(c+dx) \cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
 & \frac{3a^2}{(A-B) \sin(c+dx) \cos^4(c+dx)} \\
 & \frac{3d(a \cos(c+dx) + a)^2}{3d(a \cos(c+dx) + a)^2} \\
 & \quad \downarrow \text{24} \\
 & \frac{3 \left(\frac{4a^2(2A-3B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d} + a^2(7A-10B) \left(\frac{\sin(c+dx) \cos(c+dx)}{2d} + \frac{x}{2} \right) \right)}{a^2} + \frac{(7A-10B) \sin(c+dx) \cos^3(c+dx)}{d(\cos(c+dx)+1)} + \\
 & \frac{3a^2}{(A-B) \sin(c+dx) \cos^4(c+dx)} \\
 & \frac{3d(a \cos(c+dx) + a)^2}{3d(a \cos(c+dx) + a)^2}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Cos[c + d*x]^4*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (((7*A - 10*B)*Cos[c + d*x]^3*Sin[c + d*x])/(d*(1 + Cos[c + d*x])) + (3*(a^2*(7*A - 10*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) + (4*a^2*(2*A - 3*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d))/a^2)/(3*a^2)`

3.47. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

3.47.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.47.4 Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.64

method	result
parallelrisch	$\frac{-163\left(\frac{4(3A-7B)\cos(2dx+2c)}{163} + \frac{(-3A+2B)\cos(3dx+3c)}{163} - \frac{B\cos(4dx+4c)}{163}\right) + \left(A - \frac{258B}{163}\right)\cos(dx+c) + \frac{140A}{163} - \frac{219B}{163}}{48a^2d} \tan\left(\frac{dx}{2} + \frac{c}{2}\right)$
derivativedivides	$\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A - \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B - 7A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 9B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{8\left(-\frac{5A}{4} + \frac{5B}{2}\right)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(-2A + \frac{10B}{3}\right)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$
default	$\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A - \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B - 7A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 9B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{8\left(-\frac{5A}{4} + \frac{5B}{2}\right)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(-2A + \frac{10B}{3}\right)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$
risch	$\frac{7xA}{2a^2} - \frac{5Bx}{a^2} - \frac{ie^{2i(dx+c)}A}{8a^2d} + \frac{ie^{2i(dx+c)}B}{4a^2d} + \frac{ie^{i(dx+c)}A}{a^2d} - \frac{15ie^{i(dx+c)}B}{8a^2d} - \frac{ie^{-i(dx+c)}A}{a^2d} + \frac{15ie^{-i(dx+c)}B}{8a^2d}$
norman	$\frac{(7A-10B)x}{2a} + \frac{(A-B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} + \frac{5(7A-10B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2a} + \frac{5(7A-10B)x\left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} + \frac{5(7A-10B)x\left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a}$

input `int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)`

output `1/48*(-163*(4/163*(3*A-7*B)*cos(2*d*x+2*c)+1/163*(-3*A+2*B)*cos(3*d*x+3*c)-1/163*B*cos(4*d*x+4*c)+(A-258/163*B)*cos(d*x+c)+140/163*A-219/163*B)*tan(1/2*d*x+1/2*c)*sec(1/2*d*x+1/2*c)^2+168*(A-10/7*B)*x*d)/a^2/d`

3.47.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.91

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{3(7A-10B)dx\cos(dx+c)^2 + 6(7A-10B)dx\cos(dx+c) + 3(7A-10B)dx + (2B\cos(dx+c))^4}{6(a^2d\cos(dx+c))^2} -$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="fricas")`

output $\frac{1}{6}*(3*(7*A - 10*B)*d*x*cos(d*x + c)^2 + 6*(7*A - 10*B)*d*x*cos(d*x + c) + 3*(7*A - 10*B)*d*x + (2*B*cos(d*x + c))^4 + (3*A - 2*B)*cos(d*x + c)^3 - 6*(A - 2*B)*cos(d*x + c)^2 - (43*A - 66*B)*cos(d*x + c) - 32*A + 48*B)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)$

3.47.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1425 vs. $2(155) = 310$.

Time = 3.18 (sec) , antiderivative size = 1425, normalized size of antiderivative = 8.38

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)`

output `Piecewise((21*A*d*x*tan(c/2 + d*x/2)**6/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 63*A*d*x*tan(c/2 + d*x/2)**4/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 63*A*d*x*tan(c/2 + d*x/2)**2/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 21*A*d*x/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + A*tan(c/2 + d*x/2)**9/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 18*A*tan(c/2 + d*x/2)**7/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 90*A*tan(c/2 + d*x/2)**5/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 110*A*tan(c/2 + d*x/2)**3/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 39*A*tan(c/2 + d*x/2)/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 30*B*d*x*tan(c/2 + d*x/2)**6/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 90*B*d*x*tan(c/2 + d*x/2)**4/(6*a**2*d*tan(c/2 + d*x/2)**6 + 18*a**2*d*tan(c/2 + d*x/2)**4 + 18*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 90*B...`

3.47.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 372 vs. $2(160) = 320$.

Time = 0.33 (sec) , antiderivative size = 372, normalized size of antiderivative = 2.19

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{B \left(\frac{4 \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} + \frac{20 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{15 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a^2 + \frac{3a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3a^2 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{a^2 \sin(dx+c)^6}{(\cos(dx+c)+1)^6}} + \frac{27 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{60 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^2} \right) - A \left(\frac{6 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{2a^2 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^2 + \frac{2a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2}} \right)}{6d}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `1/6*(B*(4*(9*sin(d*x + c)/(cos(d*x + c) + 1) + 20*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 15*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(a^2 + 3*a^2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*a^2*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + a^2*sin(d*x + c)^6/(cos(d*x + c) + 1)^6) + (27*sin(d*x + c)/(cos(d*x + c) + 1) - sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2 - 60*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^2) - A*(6*(3*sin(d*x + c)/(cos(d*x + c) + 1) + 5*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(a^2 + 2*a^2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a^2*sin(d*x + c)^4/(cos(d*x + c) + 1)^4) + (21*sin(d*x + c)/(cos(d*x + c) + 1) - sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2 - 42*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^2)/d`

3.47.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.13

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{3(dx+c)(7A-10B)}{a^2} - \frac{2 \left(15A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 30B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 + 24A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 40B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 9A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 18B \right)}{\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1 \right)^3 a^2}$$

6d

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

3.47. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

output $\frac{1}{6}(3(dx + c)(7A - 10B)/a^2 - 2(15A \tan(1/2 dx + 1/2 c)^5 - 30B \tan(1/2 dx + 1/2 c)^5 + 24A \tan(1/2 dx + 1/2 c)^3 - 40B \tan(1/2 dx + 1/2 c)^3 + 9A \tan(1/2 dx + 1/2 c) - 18B \tan(1/2 dx + 1/2 c))/((\tan(1/2 dx + 1/2 c)^2 + 1)^3 a^2) + (A a^4 \tan(1/2 dx + 1/2 c)^3 - B a^4 \tan(1/2 dx + 1/2 c)^3 - 21A a^4 \tan(1/2 dx + 1/2 c) + 27B a^4 \tan(1/2 dx + 1/2 c))/a^6)/d$

3.47.9 Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.11

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{x(7A - 10B)}{2a^2}$$

$$- \frac{(5A - 10B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + (8A - \frac{40B}{3}) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (3A - 6B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 3a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^2 \right)}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{2(A-B)}{a^2} + \frac{3A-5B}{2a^2} \right)}{d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A - B)}{6a^2 d}$$

input $\text{int}((\cos(c + d*x))^4*(A + B*\cos(c + d*x)))/(a + a*\cos(c + d*x))^2,x)$

output $(x*(7*A - 10*B))/(2*a^2) - (\tan(c/2 + (d*x)/2)^5*(5*A - 10*B) + \tan(c/2 + (d*x)/2)^3*(8*A - (40*B)/3) + \tan(c/2 + (d*x)/2)*(3*A - 6*B))/(d*(3*a^2*\tan(c/2 + (d*x)/2)^2 + 3*a^2*\tan(c/2 + (d*x)/2)^4 + a^2*\tan(c/2 + (d*x)/2)^6 + a^2)) - (\tan(c/2 + (d*x)/2)*((2*(A - B))/a^2 + (3*A - 5*B)/(2*a^2)))/d + (\tan(c/2 + (d*x)/2)^3*(A - B))/(6*a^2*d)$

3.48
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.48.1	Optimal result	609
3.48.2	Mathematica [B] (verified)	609
3.48.3	Rubi [A] (verified)	610
3.48.4	Maple [A] (verified)	612
3.48.5	Fricas [A] (verification not implemented)	613
3.48.6	Sympy [B] (verification not implemented)	613
3.48.7	Maxima [B] (verification not implemented)	614
3.48.8	Giac [A] (verification not implemented)	615
3.48.9	Mupad [B] (verification not implemented)	615

3.48.1 Optimal result

Integrand size = 31, antiderivative size = 147

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = -\frac{(4A-7B)x}{2a^2} + \frac{2(5A-8B) \sin(c+dx)}{3a^2d} - \frac{(4A-7B) \cos(c+dx) \sin(c+dx)}{2a^2d} + \frac{(5A-8B) \cos^2(c+dx) \sin(c+dx)}{3a^2d(1+\cos(c+dx))} + \frac{(A-B) \cos^3(c+dx) \sin(c+dx)}{3d(a+a \cos(c+dx))^2}$$

output

```
-1/2*(4*A-7*B)*x/a^2+2/3*(5*A-8*B)*sin(d*x+c)/a^2/d-1/2*(4*A-7*B)*cos(d*x+c)*sin(d*x+c)/a^2/d+1/3*(5*A-8*B)*cos(d*x+c)^2*sin(d*x+c)/a^2/d/(1+cos(d*x+c))+1/3*(A-B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^2
```

3.48.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 315 vs. 2(147) = 294.

Time = 1.54 (sec) , antiderivative size = 315, normalized size of antiderivative = 2.14

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) (-36(4A-7B)dx \cos\left(\frac{dx}{2}\right) - 36(4A-7B)dx \cos\left(c+\frac{dx}{2}\right) - 48Adx \cos\left(c+\frac{3dx}{2}\right)}{\dots}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output
$$\frac{(\text{Cos}[(c + d*x)/2] * \text{Sec}[c/2] * (-36*(4*A - 7*B)*d*x*\text{Cos}[(d*x)/2] - 36*(4*A - 7*B)*d*x*\text{Cos}[c + (d*x)/2] - 48*A*d*x*\text{Cos}[c + (3*d*x)/2] + 84*B*d*x*\text{Cos}[c + (3*d*x)/2] - 48*A*d*x*\text{Cos}[2*c + (3*d*x)/2] + 84*B*d*x*\text{Cos}[2*c + (3*d*x)/2] + 264*A*\text{Sin}[(d*x)/2] - 381*B*\text{Sin}[(d*x)/2] - 120*A*\text{Sin}[c + (d*x)/2] + 147*B*\text{Sin}[c + (d*x)/2] + 164*A*\text{Sin}[c + (3*d*x)/2] - 239*B*\text{Sin}[c + (3*d*x)/2] + 36*A*\text{Sin}[2*c + (3*d*x)/2] - 63*B*\text{Sin}[2*c + (3*d*x)/2] + 12*A*\text{Sin}[2*c + (5*d*x)/2] - 15*B*\text{Sin}[2*c + (5*d*x)/2] + 12*A*\text{Sin}[3*c + (5*d*x)/2] - 15*B*\text{Sin}[3*c + (5*d*x)/2] + 3*B*\text{Sin}[3*c + (7*d*x)/2] + 3*B*\text{Sin}[4*c + (7*d*x)/2]))}{(48*a^2*d*(1 + \text{Cos}[c + d*x])^2)}$$

3.48.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 3456, 3042, 3456, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c + dx + \frac{\pi}{2})^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{(a \sin(c + dx + \frac{\pi}{2}) + a)^2} dx \\ & \quad \downarrow \text{3456} \\ & \frac{\int \frac{\cos^2(c+dx)(3a(A-B)-a(2A-5B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(3a(A-B)-a(2A-5B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2} \\ & \quad \downarrow \text{3456} \end{aligned}$$

3.48. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{\int \cos(c+dx) \frac{(2a^2(5A-8B)-3a^2(4A-7B) \cos(c+dx)) dx}{a^2} + \frac{(5A-8B) \sin(c+dx) \cos^2(c+dx)}{d(\cos(c+dx)+1)}}{3a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sin(c+dx+\frac{\pi}{2}) \frac{(2a^2(5A-8B)-3a^2(4A-7B) \sin(c+dx+\frac{\pi}{2})) dx}{a^2} + \frac{(5A-8B) \sin(c+dx) \cos^2(c+dx)}{d(\cos(c+dx)+1)}}{3a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3213} \\
& \frac{\frac{2a^2(5A-8B) \sin(c+dx)}{d} - \frac{3a^2(4A-7B) \sin(c+dx) \cos(c+dx)}{2d} - \frac{3}{2}a^2x(4A-7B)}{a^2} + \frac{(5A-8B) \sin(c+dx) \cos^2(c+dx)}{d(\cos(c+dx)+1)}}{3a^2} + \\
& \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{3d(a \cos(c+dx) + a)^2}
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Cos[c + d*x]^3*Sin[c + d*x]/(3*d*(a + a*Cos[c + d*x])^2) + (((5*A - 8*B)*Cos[c + d*x]^2*Sin[c + d*x])/(d*(1 + Cos[c + d*x])) + ((-3*a^2*(4*A - 7*B)*x)/2 + (2*a^2*(5*A - 8*B)*Sin[c + d*x])/d - (3*a^2*(4*A - 7*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/a^2)/(3*a^2)`

3.48.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]) , x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

3.48.4 Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.60

method	result
parallelrisch	$\frac{7 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\frac{3 \cos(2dx+2c)(A-B)}{28} + \frac{3B \cos(3dx+3c)}{112} + \left(A - \frac{163B}{112}\right) \cos(dx+c) + \frac{23A}{28} - \frac{5B}{4} \right) \left(\sec^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 6 \left(A - \frac{7B}{4} \right) x}{3a^2d}$
derivativedivides	$-\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} + \frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} + 5A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 7B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{4\left(\left(\frac{5B}{2} - A\right)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(\frac{3B}{2} - A\right)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$
default	$-\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} + \frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} + 5A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 7B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{4\left(\left(\frac{5B}{2} - A\right)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(\frac{3B}{2} - A\right)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$
risch	$-\frac{2xA}{a^2} + \frac{7Bx}{2a^2} - \frac{ie^{2i(dx+c)}B}{8a^2d} + \frac{ie^{i(dx+c)}B}{a^2d} - \frac{ie^{i(dx+c)}A}{2a^2d} - \frac{ie^{-i(dx+c)}B}{a^2d} + \frac{ie^{-i(dx+c)}A}{2a^2d} + \frac{ie^{-2i(dx+c)}B}{8a^2d} + \frac{ie^{-2i(dx+c)}A}{8a^2d}$
norman	$\frac{(11A-18B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{ad} - \frac{(4A-7B)x}{2a} - \frac{(A-B)\left(\tan^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} - \frac{2(4A-7B)x\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} - \frac{3(4A-7B)x\left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a}$

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBO
SE)
```

```
output 1/3*(7*tan(1/2*d*x+1/2*c)*(3/28*cos(2*d*x+2*c)*(A-B)+3/112*B*cos(3*d*x+3*c)
)+(A-163/112*B)*cos(d*x+c)+23/28*A-5/4*B)*sec(1/2*d*x+1/2*c)^2-6*(A-7/4*B)
*x*d)/a^2/d
```

$$3.48. \int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.48.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.94

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx =$$

$$\frac{3(4A-7B)dx\cos(dx+c)^2 + 6(4A-7B)dx\cos(dx+c) + 3(4A-7B)dx - (3B\cos(dx+c))^3 +}{6(a^2d\cos(dx+c))^2 + 2a^2d\cos(dx+c)}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="fracas")`

output `-1/6*(3*(4*A - 7*B)*d*x*cos(d*x + c)^2 + 6*(4*A - 7*B)*d*x*cos(d*x + c) + 3*(4*A - 7*B)*d*x - (3*B*cos(d*x + c)^3 + 6*(A - B)*cos(d*x + c)^2 + (28*A - 43*B)*cos(d*x + c) + 20*A - 32*B)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.48.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 843 vs. $2(136) = 272$.

Time = 1.81 (sec) , antiderivative size = 843, normalized size of antiderivative = 5.73

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \left\{ \begin{array}{l} -\frac{12Adx \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 12a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} - \frac{24Adx \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 12a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} - \frac{12Adx}{6a^2d \tan^4\left(\frac{c}{2} + \frac{dx}{2}\right) + 12a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} \\ \frac{x(A+B\cos(c))\cos^3(c)}{(a\cos(c)+a)^2} \end{array} \right.$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)`

output `Piecewise((-12*A*d*x*tan(c/2 + d*x/2)**4/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 24*A*d*x*tan(c/2 + d*x/2)**2/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 12*A*d*x/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - A*tan(c/2 + d*x/2)**7/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 13*A*tan(c/2 + d*x/2)**5/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 41*A*tan(c/2 + d*x/2)**3/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 27*A*tan(c/2 + d*x/2)/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 21*B*d*x*tan(c/2 + d*x/2)**4/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 42*B*d*x*tan(c/2 + d*x/2)**2/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 21*B*d*x/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + B*tan(c/2 + d*x/2)**7/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 19*B*tan(c/2 + d*x/2)**5/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 71*B*tan(c/2 + d*x/2)**3/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 39*B*tan(c/2 + d*x/2)/(6*a**2*d*tan(c/2 + d*x/2)**4 + 12*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**3/(a*cos(c) + a)**2, True))`

3.48.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. $2(137) = 274$.

Time = 0.30 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.93

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx =$$

$$\frac{B \left(\frac{6 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{5 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^2 + \frac{2a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^2 \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} + \frac{21 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{42 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^2} \right) - A \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)}{a^2} \right)}{6d}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output
$$\frac{-1/6*(B*(6*(3*\sin(d*x + c)/(\cos(d*x + c) + 1) + 5*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(a^2 + 2*a^2*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + a^2*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4) + (21*\sin(d*x + c)/(\cos(d*x + c) + 1) - \sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/a^2 - 42*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1))/a^2) - A*((15*\sin(d*x + c)/(\cos(d*x + c) + 1) - \sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/a^2 - 24*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1))/a^2 + 12*\sin(d*x + c)/((a^2 + a^2*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2)*(\cos(d*x + c) + 1)))}{d}$$

3.48.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.12

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \frac{3(dx+c)(4A-7B)}{a^2} - \frac{6\left(2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 5B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 3B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)\right)}{\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1\right)^2 a^2} + \frac{Aa^4 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - B}{6d}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

output
$$\frac{-1/6*(3*(d*x + c)*(4*A - 7*B)/a^2 - 6*(2*A*\tan(1/2*d*x + 1/2*c)^3 - 5*B*\tan(1/2*d*x + 1/2*c)^3 + 2*A*\tan(1/2*d*x + 1/2*c) - 3*B*\tan(1/2*d*x + 1/2*c))/((\tan(1/2*d*x + 1/2*c)^2 + 1)^2*a^2) + (A*a^4*\tan(1/2*d*x + 1/2*c)^3 - B*a^4*\tan(1/2*d*x + 1/2*c)^3 - 15*A*a^4*\tan(1/2*d*x + 1/2*c) + 21*B*a^4*\tan(1/2*d*x + 1/2*c))/a^6}{d}$$

3.48.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.03

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3(A-B)}{2a^2} + \frac{2A-4B}{2a^2}\right) - x(4A-7B)}{d} - \frac{(2A-5B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (2A-3B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A-B)}{d \left(a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 2a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^2\right)} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A-B)}{6a^2 d}$$

3.48.
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2,x)`

output `(tan(c/2 + (d*x)/2)*((3*(A - B))/(2*a^2) + (2*A - 4*B)/(2*a^2)))/d - (x*(4*A - 7*B))/(2*a^2) + (tan(c/2 + (d*x)/2)^3*(2*A - 5*B) + tan(c/2 + (d*x)/2)*(2*A - 3*B))/(d*(2*a^2*tan(c/2 + (d*x)/2)^2 + a^2*tan(c/2 + (d*x)/2)^4 + a^2)) - (tan(c/2 + (d*x)/2)^3*(A - B))/(6*a^2*d)`

3.48. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

3.49 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

3.49.1 Optimal result 617
 3.49.2 Mathematica [A] (verified) 617
 3.49.3 Rubi [A] (verified) 618
 3.49.4 Maple [A] (verified) 621
 3.49.5 Fricas [A] (verification not implemented) 621
 3.49.6 Sympy [B] (verification not implemented) 622
 3.49.7 Maxima [B] (verification not implemented) 622
 3.49.8 Giac [A] (verification not implemented) 623
 3.49.9 Mupad [B] (verification not implemented) 623

3.49.1 Optimal result

Integrand size = 31, antiderivative size = 99

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = \frac{(A-2B)x}{a^2} - \frac{(A-4B) \sin(c+dx)}{3a^2d} - \frac{(A-2B) \sin(c+dx)}{a^2d(1+\cos(c+dx))} + \frac{(A-B) \cos^2(c+dx) \sin(c+dx)}{3d(a+a \cos(c+dx))^2}$$

output `(A-2*B)*x/a^2-1/3*(A-4*B)*sin(d*x+c)/a^2/d-(A-2*B)*sin(d*x+c)/a^2/d/(1+cos(d*x+c))+1/3*(A-B)*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^2`

3.49.2 Mathematica [A] (verified)

Time = 1.28 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.38

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = \frac{2 \cos\left(\frac{1}{2}(c+dx)\right) \left((A-B) \sec\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right) - 2(5A-8B) \cos^2\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right) + 6 \cos^3\left(\frac{1}{2}(c+dx)\right) \right)}{3a^2d(1+\cos(c+dx))^2}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output $(2*\text{Cos}[(c + d*x)/2]*((A - B)*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] - 2*(5*A - 8*B)*\text{Cos}[(c + d*x)/2]^2*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] + 6*\text{Cos}[(c + d*x)/2]^3*((A - 2*B)*d*x + B*\text{Sin}[c + d*x]) + (A - B)*\text{Cos}[(c + d*x)/2]*\text{Tan}[c/2]))/(3*a^2*d*(1 + \text{Cos}[c + d*x])^2)$

3.49.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3456, 3042, 3447, 3042, 3502, 27, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{(a \sin(c + dx + \frac{\pi}{2}) + a)^2} dx$$

↓ 3456

$$\frac{\int \frac{\cos(c+dx)(2a(A-B)-a(A-4B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{3d(a \cos(c + dx) + a)^2}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(2a(A-B)-a(A-4B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{3d(a \cos(c + dx) + a)^2}$$

↓ 3447

$$\frac{\int \frac{2a(A-B)\cos(c+dx)-a(A-4B)\cos^2(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{3d(a \cos(c + dx) + a)^2}$$

↓ 3042

$$\frac{\int \frac{2a(A-B)\sin(c+dx+\frac{\pi}{2})-a(A-4B)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{(A - B) \sin(c + dx) \cos^2(c + dx)}{3d(a \cos(c + dx) + a)^2}$$

↓ 3502

3.49. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{\int \frac{3a^2(A-2B)\cos(c+dx)}{\cos(c+dx)a+a} dx - \frac{(A-4B)\sin(c+dx)}{d}}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{3a(A-2B)\int \frac{\cos(c+dx)}{\cos(c+dx)a+a} dx - \frac{(A-4B)\sin(c+dx)}{d}}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3a(A-2B)\int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})a+a} dx - \frac{(A-4B)\sin(c+dx)}{d}}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3214} \\
& \frac{3a(A-2B)\left(\frac{x}{a} - \int \frac{1}{\cos(c+dx)a+a} dx\right) - \frac{(A-4B)\sin(c+dx)}{d}}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3a(A-2B)\left(\frac{x}{a} - \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx\right) - \frac{(A-4B)\sin(c+dx)}{d}}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3127} \\
& \frac{3a(A-2B)\left(\frac{x}{a} - \frac{\sin(c+dx)}{d(a\cos(c+dx)+a)}\right) - \frac{(A-4B)\sin(c+dx)}{d}}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}
\end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Cos[c + d*x]^2*Sin[c + d*x]/(3*d*(a + a*Cos[c + d*x])^2) + (-(((A - 4*B)*Sin[c + d*x])/d) + 3*a*(A - 2*B)*(x/a - Sin[c + d*x]/(d*(a + a*Cos[c + d*x]))))/((3*a^2)`

3.49.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.49. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.49.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.74

method	result
parallelrisc	$\frac{(-20A + \left(\sec^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)(3B \cos(2dx+2c) + 28B \cos(dx+c) + 2A + 23B)) \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 12dx(A-2B)}{12a^2d}$
derivativedivides	$\frac{\left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)A - \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)B - 3A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{4B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + 4(A-2B) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2}$
default	$\frac{\left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)A - \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)B - 3A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{4B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + 4(A-2B) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2}$
risc	$\frac{x A}{a^2} - \frac{2 B x}{a^2} - \frac{i e^{i(dx+c)} B}{2 a^2 d} + \frac{i e^{-i(dx+c)} B}{2 a^2 d} - \frac{2 i (6 A e^{2 i(dx+c)} - 9 B e^{2 i(dx+c)} + 9 A e^{i(dx+c)} - 15 B e^{i(dx+c)} + 5 A - 8 B)}{3 d a^2 (e^{i(dx+c)} + 1)^3}$
norman	$\frac{(A-2B)x}{a} + \frac{(A-2B)x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} - \frac{3(A-3B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2ad} - \frac{(A-2B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{ad} + \frac{3(A-2B)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} + \frac{3(A-2B)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^3 a}$

input `int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)`

output `1/12*((-20*A+sec(1/2*d*x+1/2*c)^2*(3*B*cos(2*d*x+2*c)+28*B*cos(d*x+c)+2*A+23*B))*tan(1/2*d*x+1/2*c)+12*d*x*(A-2*B))/a^2/d`

3.49.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.18

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

$$= \frac{3(A-2B)dx \cos(dx+c)^2 + 6(A-2B)dx \cos(dx+c) + 3(A-2B)dx + (3B \cos(dx+c))^2 - (5A - 3(a^2d \cos(dx+c)^2 + 2a^2d \cos(dx+c) + a^2d))}{3(a^2d \cos(dx+c)^2 + 2a^2d \cos(dx+c) + a^2d)}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="fracas")`

output `1/3*(3*(A - 2*B)*d*x*cos(d*x + c)^2 + 6*(A - 2*B)*d*x*cos(d*x + c) + 3*(A - 2*B)*d*x + (3*B*cos(d*x + c)^2 - (5*A - 14*B)*cos(d*x + c) - 4*A + 10*B)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.49. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

3.49.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 411 vs. $2(90) = 180$.

Time = 1.09 (sec) , antiderivative size = 411, normalized size of antiderivative = 4.15

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \begin{cases} \frac{6Adx \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} + \frac{6Adx}{6a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} + \frac{A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} - \frac{8A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} - \frac{9A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 6a^2d} \\ \frac{x(A+B\cos(c))\cos^2(c)}{(a\cos(c)+a)^2} \end{cases}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)`

output `Piecewise((6*A*d*x*tan(c/2 + d*x/2)**2/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 6*A*d*x/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + A*tan(c/2 + d*x/2)**5/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 8*A*tan(c/2 + d*x/2)**3/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 9*A*tan(c/2 + d*x/2)/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 12*B*d*x*tan(c/2 + d*x/2)**2/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - 12*B*d*x/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) - B*tan(c/2 + d*x/2)**5/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 14*B*tan(c/2 + d*x/2)**3/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d) + 27*B*tan(c/2 + d*x/2)/(6*a**2*d*tan(c/2 + d*x/2)**2 + 6*a**2*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**2/(a*cos(c) + a)**2, True))`

3.49.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(95) = 190$.

Time = 0.32 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.93

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{B \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{24 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^2} + \frac{12 \sin(dx+c)}{\left(a^2 + \frac{a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2}\right)(\cos(dx+c)+1)} \right) - A \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{\sin(dx+c)}{a^2} \right)}{6d}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

3.49. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

output $\frac{1}{6} * (B * ((15 * \sin(dx + c)) / (\cos(dx + c) + 1) - \sin(dx + c)^3 / (\cos(dx + c) + 1)^3) / a^2 - 24 * \arctan(\sin(dx + c) / (\cos(dx + c) + 1)) / a^2 + 12 * \sin(dx + c) / ((a^2 + a^2 * \sin(dx + c)^2 / (\cos(dx + c) + 1)^2) * (\cos(dx + c) + 1))) - A * ((9 * \sin(dx + c) / (\cos(dx + c) + 1) - \sin(dx + c)^3 / (\cos(dx + c) + 1)^3) / a^2 - 12 * \arctan(\sin(dx + c) / (\cos(dx + c) + 1)) / a^2) / d$

3.49.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.20

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \frac{6(dx+c)(A-2B)}{a^2} + \frac{12B \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)a^2} + \frac{Aa^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - Ba^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 9Aa^4 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 15Ba^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{a^6}$$

$6d$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

output $\frac{1}{6} * (6 * (dx + c) * (A - 2 * B) / a^2 + 12 * B * \tan(1/2 * dx + 1/2 * c) / ((\tan(1/2 * dx + 1/2 * c)^2 + 1) * a^2) + (A * a^4 * \tan(1/2 * dx + 1/2 * c)^3 - B * a^4 * \tan(1/2 * dx + 1/2 * c)^3 - 9 * A * a^4 * \tan(1/2 * dx + 1/2 * c) + 15 * B * a^4 * \tan(1/2 * dx + 1/2 * c)) / a^6) / d$

3.49.9 Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.06

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \frac{x(A - 2B)}{a^2} - \frac{\tan(\frac{c}{2} + \frac{dx}{2})}{d} \left(\frac{A-B}{a^2} + \frac{A-3B}{2a^2} \right) + \frac{2B \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(a^2 \tan^2(\frac{c}{2} + \frac{dx}{2}) + a^2 \right)} + \frac{\tan(\frac{c}{2} + \frac{dx}{2})^3 (A - B)}{6a^2 d}$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2,x)`

3.49. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

output $(x*(A - 2*B))/a^2 - (\tan(c/2 + (d*x)/2)*((A - B)/a^2 + (A - 3*B)/(2*a^2))$
 $/d + (2*B*\tan(c/2 + (d*x)/2))/(d*(a^2*\tan(c/2 + (d*x)/2)^2 + a^2)) + (\tan(c/2 + (d*x)/2)^3*(A - B))/(6*a^2*d)$

3.50
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.50.1	Optimal result	625
3.50.2	Mathematica [B] (verified)	625
3.50.3	Rubi [A] (verified)	626
3.50.4	Maple [A] (verified)	628
3.50.5	Fricas [A] (verification not implemented)	629
3.50.6	Sympy [A] (verification not implemented)	629
3.50.7	Maxima [A] (verification not implemented)	630
3.50.8	Giac [A] (verification not implemented)	630
3.50.9	Mupad [B] (verification not implemented)	631

3.50.1 Optimal result

Integrand size = 29, antiderivative size = 70

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{Bx}{a^2} + \frac{(2A - 5B) \sin(c + dx)}{3a^2d(1 + \cos(c + dx))} - \frac{(A - B) \sin(c + dx)}{3d(a + a \cos(c + dx))^2}$$

output `B*x/a^2+1/3*(2*A-5*B)*sin(d*x+c)/a^2/d/(1+cos(d*x+c))-1/3*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2`

3.50.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 153 vs. 2(70) = 140.

Time = 0.75 (sec) , antiderivative size = 153, normalized size of antiderivative = 2.19

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\sec\left(\frac{c}{2}\right) \sec^3\left(\frac{1}{2}(c + dx)\right) \left(9Bdx \cos\left(\frac{dx}{2}\right) + 9Bdx \cos\left(c + \frac{dx}{2}\right) + 3Bdx \cos\left(c + \frac{3dx}{2}\right) + 3Bdx \cos\left(2c + \frac{3dx}{2}\right)\right)}{2a^2d}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output $(\text{Sec}[c/2]*\text{Sec}[(c + d*x)/2]^3*(9*B*d*x*\text{Cos}[(d*x)/2] + 9*B*d*x*\text{Cos}[c + (d*x)/2] + 3*B*d*x*\text{Cos}[c + (3*d*x)/2] + 3*B*d*x*\text{Cos}[2*c + (3*d*x)/2] + 6*A*\text{Sin}[(d*x)/2] - 18*B*\text{Sin}[(d*x)/2] - 6*A*\text{Sin}[c + (d*x)/2] + 12*B*\text{Sin}[c + (d*x)/2] + 4*A*\text{Sin}[c + (3*d*x)/2] - 10*B*\text{Sin}[c + (3*d*x)/2]))/(24*a^2*d)$

3.50.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3447, 3042, 3498, 25, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^2} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a\cos(c+dx)+a)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{(a\sin(c+dx+\frac{\pi}{2})+a)^2} dx \\ & \quad \downarrow \text{3498} \\ & -\frac{\int -\frac{2a(A-B)+3aB\cos(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} - \frac{(A-B)\sin(c+dx)}{3d(a\cos(c+dx)+a)^2} \\ & \quad \downarrow \text{25} \\ & \frac{\int \frac{2a(A-B)+3aB\cos(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} - \frac{(A-B)\sin(c+dx)}{3d(a\cos(c+dx)+a)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{2a(A-B)+3aB\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} - \frac{(A-B)\sin(c+dx)}{3d(a\cos(c+dx)+a)^2} \end{aligned}$$

3.50. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{array}{c}
 \downarrow \text{3214} \\
 \frac{a(2A - 5B) \int \frac{1}{\cos(c+dx)a+a} dx + 3Bx}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\
 \downarrow \text{3042} \\
 \frac{a(2A - 5B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx + 3Bx}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\
 \downarrow \text{3127} \\
 \frac{\frac{a(2A-5B) \sin(c+dx)}{d(a \cos(c+dx)+a)} + 3Bx}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2}
 \end{array}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `-1/3*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + (3*B*x + (a*(2*A - 5*B)*Sin[c + d*x]))/(d*(a + a*Cos[c + d*x]))/(3*a^2)`

3.50.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*SIN[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3498 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b - a*B + b*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[1/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]`

3.50.4 Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.70

method	result
parallelrisch	$\frac{(-A+B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (3A-9B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 6dxB}{6a^2d}$
derivativedivides	$-\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} + \frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} + A\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3B\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 4B\arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2}$
default	$-\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} + \frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} + A\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3B\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 4B\arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2da^2}$
risch	$\frac{Bx}{a^2} + \frac{2i(3Ae^{2i(dx+c)} - 6Be^{2i(dx+c)} + 3Ae^{i(dx+c)} - 9Be^{i(dx+c)} + 2A - 5B)}{3da^2(e^{i(dx+c)} + 1)^3}$
norman	$\frac{Bx}{a} + \frac{Bx\left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} + \frac{2Bx\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} + \frac{(A-7B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} + \frac{(A-3B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2ad} - \frac{(A-B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} + \frac{5}{a\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$

input `int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)`

output `1/6*((-A+B)*tan(1/2*d*x+1/2*c)^3+(3*A-9*B)*tan(1/2*d*x+1/2*c)+6*d*x*B)/a^2/d`

3.50.
$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

3.50.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.30

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{3Bdx\cos(dx+c)^2 + 6Bdx\cos(dx+c) + 3Bdx + ((2A-5B)\cos(dx+c) + A-4B)\sin(dx+c)}{3(a^2d\cos(dx+c)^2 + 2a^2d\cos(dx+c) + a^2d)}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="fricas")
```

```
output 1/3*(3*B*d*x*cos(d*x + c)^2 + 6*B*d*x*cos(d*x + c) + 3*B*d*x + ((2*A - 5*B)*cos(d*x + c) + A - 4*B)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)
```

3.50.6 Sympy [A] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.50

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \begin{cases} -\frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2a^2d} + \frac{Bx}{a^2} + \frac{B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2d} - \frac{3B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2a^2d} & \text{for } d \neq 0 \\ \frac{x(A+B\cos(c))\cos(c)}{(a\cos(c)+a)^2} & \text{otherwise} \end{cases}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)
```

```
output Piecewise((-A*tan(c/2 + d*x/2)**3/(6*a**2*d) + A*tan(c/2 + d*x/2)/(2*a**2*d) + B*x/a**2 + B*tan(c/2 + d*x/2)**3/(6*a**2*d) - 3*B*tan(c/2 + d*x/2)/(2*a**2*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)/(a*cos(c) + a)**2, True))
```

3.50.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.71

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{B \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{12 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^2} \right) - A \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{6d}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/6*(B*((9*sin(d*x + c))/(cos(d*x + c) + 1) - sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2 - 12*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^2) - A*(3*sin(d*x + c)/(cos(d*x + c) + 1) - sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2)/d`

3.50.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.23

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{\frac{6(dx+c)B}{a^2} - \frac{Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 3Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 9Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)}{a^6}}{6d}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

output `1/6*(6*(d*x + c)*B/a^2 - (A*a^4*tan(1/2*d*x + 1/2*c)^3 - B*a^4*tan(1/2*d*x + 1/2*c)^3 - 3*A*a^4*tan(1/2*d*x + 1/2*c) + 9*B*a^4*tan(1/2*d*x + 1/2*c))/a^6)/d`

3.50.9 Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.93

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{3A \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 9B \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + 6B dx}{6a^2 d}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2,x)`output `(3*A*tan(c/2 + (d*x)/2) - 9*B*tan(c/2 + (d*x)/2) - A*tan(c/2 + (d*x)/2)^3 + B*tan(c/2 + (d*x)/2)^3 + 6*B*d*x)/(6*a^2*d)`

3.51 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2} dx$

3.51.1	Optimal result	632
3.51.2	Mathematica [A] (verified)	632
3.51.3	Rubi [A] (verified)	633
3.51.4	Maple [A] (verified)	634
3.51.5	Fricas [A] (verification not implemented)	635
3.51.6	Sympy [A] (verification not implemented)	635
3.51.7	Maxima [A] (verification not implemented)	635
3.51.8	Giac [A] (verification not implemented)	636
3.51.9	Mupad [B] (verification not implemented)	636

3.51.1 Optimal result

Integrand size = 23, antiderivative size = 65

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{(A - B) \sin(c + dx)}{3d(a + a \cos(c + dx))^2} + \frac{(A + 2B) \sin(c + dx)}{3d(a^2 + a^2 \cos(c + dx))}$$

output `1/3*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2+1/3*(A+2*B)*sin(d*x+c)/d/(a^2+a^2*cos(d*x+c))`

3.51.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.66

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{(2A + B + (A + 2B) \cos(c + dx)) \sin(c + dx)}{3a^2d(1 + \cos(c + dx))^2}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^2,x]`

output `((2*A + B + (A + 2*B)*Cos[c + d*x])*Sin[c + d*x])/(3*a^2*d*(1 + Cos[c + d*x])^2)`

3.51.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {3042, 3229, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a \cos(c + dx) + a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2} dx \\
 & \quad \downarrow \text{3229} \\
 & \frac{(A + 2B) \int \frac{1}{\cos(c+dx)a+a} dx}{3a} + \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A + 2B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a} + \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\
 & \quad \downarrow \text{3127} \\
 & \frac{(A + 2B) \sin(c + dx)}{3ad(a \cos(c + dx) + a)} + \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((A + 2*B)*Sin[c + d*x])/(3*a*d*(a + a*Cos[c + d*x]))`

3.51.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3229 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.51.4 Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

method	result	size
parallelrisc	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(3A + 3B + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) (A - B)}{6a^2d}$	42
derivativedivides	$\frac{\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} - \frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2da^2}$	60
default	$\frac{\frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} - \frac{\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2da^2}$	60
risc	$\frac{2i(3B e^{2i(dx+c)} + 3A e^{i(dx+c)} + 3B e^{i(dx+c)} + A + 2B)}{3da^2(e^{i(dx+c)} + 1)^3}$	64
norman	$\frac{\frac{(A-B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} + \frac{(A+B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2ad} + \frac{(2A+B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a}$	89

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)`

output `1/6*tan(1/2*d*x+1/2*c)*(3*A+3*B+tan(1/2*d*x+1/2*c)^2*(A-B))/a^2/d`

3.51.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.89

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{((A + 2B) \cos(dx + c) + 2A + B) \sin(dx + c)}{3(a^2 d \cos(dx + c)^2 + 2a^2 d \cos(dx + c) + a^2 d)}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="fracas")`output `1/3*((A + 2*B)*cos(d*x + c) + 2*A + B)*sin(d*x + c)/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`**3.51.6 Sympy [A] (verification not implemented)**

Time = 0.50 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.45

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \begin{cases} \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2 d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2a^2 d} - \frac{B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^2 d} + \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2a^2 d} & \text{for } d \neq 0 \\ \frac{x(A+B \cos(c))}{(a \cos(c)+a)^2} & \text{otherwise} \end{cases}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)`output `Piecewise((A*tan(c/2 + d*x/2)**3/(6*a**2*d) + A*tan(c/2 + d*x/2)/(2*a**2*d) - B*tan(c/2 + d*x/2)**3/(6*a**2*d) + B*tan(c/2 + d*x/2)/(2*a**2*d), Ne(d, 0)), (x*(A + B*cos(c))/(a*cos(c) + a)**2, True))`**3.51.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.43

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{A \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^2} + \frac{B \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^2}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output $\frac{1}{6} * (A * (3 * \sin(d * x + c) / (\cos(d * x + c) + 1) + \sin(d * x + c)^3 / (\cos(d * x + c) + 1)^3) / a^2 + B * (3 * \sin(d * x + c) / (\cos(d * x + c) + 1) - \sin(d * x + c)^3 / (\cos(d * x + c) + 1)^3) / a^2) / d$

3.51.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.92

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 3 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{6 a^2 d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

output $\frac{1}{6} * (A * \tan(1/2 * d * x + 1/2 * c)^3 - B * \tan(1/2 * d * x + 1/2 * c)^3 + 3 * A * \tan(1/2 * d * x + 1/2 * c) + 3 * B * \tan(1/2 * d * x + 1/2 * c)) / (a^2 * d)$

3.51.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.69

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A - B)}{6 a^2 d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A + B)}{2 a^2 d}$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^2,x)`

output $(\tan(c/2 + (d*x)/2)^3 * (A - B)) / (6 * a^2 * d) + (\tan(c/2 + (d*x)/2) * (A + B)) / (2 * a^2 * d)$

3.52 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^2} dx$

3.52.1 Optimal result 637
 3.52.2 Mathematica [B] (verified) 637
 3.52.3 Rubi [A] (verified) 638
 3.52.4 Maple [A] (verified) 640
 3.52.5 Fricas [A] (verification not implemented) 640
 3.52.6 Sympy [F] 641
 3.52.7 Maxima [A] (verification not implemented) 641
 3.52.8 Giac [A] (verification not implemented) 642
 3.52.9 Mupad [B] (verification not implemented) 642

3.52.1 Optimal result

Integrand size = 29, antiderivative size = 79

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{A \operatorname{arctanh}(\sin(c + dx))}{a^2 d} - \frac{(4A - B) \sin(c + dx)}{3a^2 d(1 + \cos(c + dx))} - \frac{(A - B) \sin(c + dx)}{3d(a + a \cos(c + dx))^2}$$

output `A*arctanh(sin(d*x+c))/a^2/d-1/3*(4*A-B)*sin(d*x+c)/a^2/d/(1+cos(d*x+c))-1/3*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2`

3.52.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 170 vs. 2(79) = 158.

Time = 0.66 (sec) , antiderivative size = 170, normalized size of antiderivative = 2.15

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) (6A \cos^3\left(\frac{1}{2}(c + dx)\right) (\log(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right))) - \log(\cos\left(\frac{1}{2}(c + dx)\right))}{(a + a \cos(c + dx))^2}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^2,x]`

output $(-2*\text{Cos}[(c + d*x)/2]*(6*A*\text{Cos}[(c + d*x)/2]^3*(\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] - \text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]]) + (A - B)*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] + 2*(4*A - B)*\text{Cos}[(c + d*x)/2]^2*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] + (A - B)*\text{Cos}[(c + d*x)/2]*\text{Tan}[c/2]))/(3*a^2*d*(1 + \text{Cos}[c + d*x])^2)$

3.52.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.241$, Rules used = {3042, 3457, 3042, 3457, 27, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})(a \sin(c + dx + \frac{\pi}{2}) + a)^2} dx \\ & \quad \downarrow \text{3457} \\ & \frac{\int \frac{(3aA - a(A - B) \cos(c + dx)) \sec(c + dx)}{\cos(c + dx)a + a} dx}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{3aA - a(A - B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})(\sin(c + dx + \frac{\pi}{2})a + a)} dx}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{3457} \\ & \frac{\int \frac{3a^2 A \sec(c + dx) dx}{a^2} - \frac{(4A - B) \sin(c + dx)}{d(\cos(c + dx) + 1)}}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{27} \\ & \frac{3A \int \sec(c + dx) dx - \frac{(4A - B) \sin(c + dx)}{d(\cos(c + dx) + 1)}}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{3A \int \csc(c + dx + \frac{\pi}{2}) dx - \frac{(4A - B) \sin(c + dx)}{d(\cos(c + dx) + 1)}}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d(a \cos(c + dx) + a)^2} \end{aligned}$$

3.52. $\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx$

$$\frac{\frac{3A \operatorname{arctanh}(\sin(c+dx))}{d} - \frac{(4A-B) \sin(c+dx)}{d(\cos(c+dx)+1)}}{3a^2} - \frac{(A-B) \sin(c+dx)}{3d(a \cos(c+dx) + a)^2}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^2,x]`

output `-1/3*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + ((3*A*ArcTanh[Sin[c + d*x]])/d - ((4*A - B)*Sin[c + d*x])/(d*(1 + Cos[c + d*x]))) / (3*a^2)`

3.52.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.52.4 Maple [A] (verified)

Time = 1.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.95

method	result
parallelrisch	$\frac{-6A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 6A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) (A - B) + 9A - 3B}{6a^2d}$
derivativedivides	$\frac{-\frac{(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right))A}{3} + \frac{(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right))B}{3} - 3A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 2A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2da^2}$
default	$\frac{-\frac{(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right))A}{3} + \frac{(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right))B}{3} - 3A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 2A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 2A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2da^2}$
risch	$\frac{2i(3A e^{2i(dx+c)} + 9A e^{i(dx+c)} - 3B e^{i(dx+c)} + 4A - B)}{3da^2(e^{i(dx+c)} + 1)^3} + \frac{A \ln(e^{i(dx+c)} + i)}{a^2d} - \frac{A \ln(e^{i(dx+c)} - i)}{a^2d}$
norman	$\frac{\frac{(A-B)(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right))}{6ad} - \frac{(3A-B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2ad} - \frac{(5A-2B)(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right))}{3ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a} + \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^2d} - \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^2d}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)`

output `1/6*(-6*A*ln(tan(1/2*d*x+1/2*c)-1)+6*A*ln(tan(1/2*d*x+1/2*c)+1)-tan(1/2*d*x+1/2*c)*(tan(1/2*d*x+1/2*c)^2*(A-B)+9*A-3*B))/a^2/d`

3.52.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.66

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{3(A \cos(dx + c)^2 + 2A \cos(dx + c) + A) \log(\sin(dx + c) + 1) - 3(A \cos(dx + c)^2 + 2A \cos(dx + c) + A) \log(-\sin(dx + c) + 1) - 2((4A - B) \cos(dx + c) + 5A - 2B) \sin(dx + c)}{6(a^2d \cos(dx + c))^2 + 2a^2d \cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^2,x, algorithm="fricas")`

output `1/6*(3*(A*cos(d*x + c)^2 + 2*A*cos(d*x + c) + A)*log(sin(d*x + c) + 1) - 3*(A*cos(d*x + c)^2 + 2*A*cos(d*x + c) + A)*log(-sin(d*x + c) + 1) - 2*((4*A - B)*cos(d*x + c) + 5*A - 2*B)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.52.
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^2} dx$$

3.52.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\int \frac{A \sec(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sec(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))**2,x)`

output `(Integral(A*sec(c + d*x)/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x))/a**2`

3.52.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.84

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx =$$

$$\frac{A \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{6 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^2} + \frac{6 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^2} \right) - \frac{B \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^2}}{6d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `-1/6*(A*((9*sin(d*x + c)/(cos(d*x + c) + 1) + sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2 - 6*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a^2 + 6*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a^2) - B*(3*sin(d*x + c)/(cos(d*x + c) + 1) + sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2/d`

3.52.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.43

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\frac{6 A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^2} - \frac{6 A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^2} - \frac{A a^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - B a^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 9 A a^4 \tan(\frac{1}{2} dx + \frac{1}{2} c) - 3 B a^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{a^6}}{6 d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

output `1/6*(6*A*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^2 - 6*A*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^2 - (A*a^4*tan(1/2*d*x + 1/2*c)^3 - B*a^4*tan(1/2*d*x + 1/2*c)^3 + 9*A*a^4*tan(1/2*d*x + 1/2*c) - 3*B*a^4*tan(1/2*d*x + 1/2*c))/a^6)/d`

3.52.9 Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.94

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{2 A \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{a^2 d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A - B)}{6 a^2 d}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{A}{a^2} + \frac{A - B}{2 a^2}\right)}{d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^2),x)`

output `(2*A*atanh(tan(c/2 + (d*x)/2)))/(a^2*d) - (tan(c/2 + (d*x)/2)^3*(A - B))/(6*a^2*d) - (tan(c/2 + (d*x)/2)*(A/a^2 + (A - B)/(2*a^2)))/d`

3.53 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^2} dx$

3.53.1 Optimal result 643
 3.53.2 Mathematica [B] (verified) 643
 3.53.3 Rubi [A] (verified) 644
 3.53.4 Maple [A] (verified) 647
 3.53.5 Fricas [B] (verification not implemented) 647
 3.53.6 Sympy [F] 648
 3.53.7 Maxima [B] (verification not implemented) 648
 3.53.8 Giac [A] (verification not implemented) 649
 3.53.9 Mupad [B] (verification not implemented) 649

3.53.1 Optimal result

Integrand size = 31, antiderivative size = 107

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx = -\frac{(2A - B)\operatorname{arctanh}(\sin(c + dx))}{a^2 d} + \frac{2(5A - 2B) \tan(c + dx)}{3a^2 d} - \frac{(2A - B) \tan(c + dx)}{a^2 d(1 + \cos(c + dx))} - \frac{(A - B) \tan(c + dx)}{3d(a + a \cos(c + dx))^2}$$

```
output - (2*A-B)*arctanh(sin(d*x+c))/a^2/d+2/3*(5*A-2*B)*tan(d*x+c)/a^2/d-(2*A-B)*tan(d*x+c)/a^2/d/(1+cos(d*x+c))-1/3*(A-B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^2
```

3.53.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 264 vs. 2(107) = 214.

Time = 1.79 (sec) , antiderivative size = 264, normalized size of antiderivative = 2.47

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) \left((A - B) \sec\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right) + 2(7A - 4B) \cos^2\left(\frac{1}{2}(c + dx)\right) \sec\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right) + 6 \cos^3\left(\frac{1}{2}(c + dx)\right) \right)}{(a + a \cos(c + dx))^2}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^2,x]`

output $(2*\text{Cos}[(c + d*x)/2]*((A - B)*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] + 2*(7*A - 4*B)*\text{Cos}[(c + d*x)/2]^2*\text{Sec}[c/2]*\text{Sin}[(d*x)/2] + 6*\text{Cos}[(c + d*x)/2]^3*((2*A - B)*(\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] - \text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]])) + (A*\text{Sin}[d*x])/((\text{Cos}[c/2] - \text{Sin}[c/2])*(\text{Cos}[c/2] + \text{Sin}[c/2])*(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2])*(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]))) + (A - B)*\text{Cos}[(c + d*x)/2]*\text{Tan}[c/2))/(3*a^2*d*(1 + \text{Cos}[c + d*x])^2)$

3.53.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3457, 3042, 3457, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2 (a \sin(c + dx + \frac{\pi}{2}) + a)^2} dx \\ & \quad \downarrow \text{3457} \\ & \frac{\int \frac{(a(4A - B) - 2a(A - B) \cos(c + dx)) \sec^2(c + dx)}{\cos(c + dx) a + a} dx}{3a^2} - \frac{(A - B) \tan(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{a(4A - B) - 2a(A - B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2 (\sin(c + dx + \frac{\pi}{2}) a + a)} dx}{3a^2} - \frac{(A - B) \tan(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{3457} \\ & \frac{\int (2a^2(5A - 2B) - 3a^2(2A - B) \cos(c + dx)) \sec^2(c + dx) dx}{3a^2} - \frac{3(2A - B) \tan(c + dx)}{d(\cos(c + dx) + 1)} - \frac{(A - B) \tan(c + dx)}{3d(a \cos(c + dx) + a)^2} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.53. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx$

$$\begin{aligned}
& \frac{\int \frac{2a^2(5A-2B) - 3a^2(2A-B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2} dx}{3a^2} - \frac{3(2A-B) \tan(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B) \tan(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3227} \\
& \frac{2a^2(5A-2B) \int \sec^2(c+dx) dx - 3a^2(2A-B) \int \sec(c+dx) dx}{3a^2} - \frac{3(2A-B) \tan(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B) \tan(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{2a^2(5A-2B) \int \csc(c+dx+\frac{\pi}{2})^2 dx - 3a^2(2A-B) \int \csc(c+dx+\frac{\pi}{2}) dx}{3a^2} - \frac{3(2A-B) \tan(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B) \tan(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{4254} \\
& \frac{-\frac{2a^2(5A-2B)}{d} \int \frac{1}{d} (-\tan(c+dx)) - 3a^2(2A-B) \int \csc(c+dx+\frac{\pi}{2}) dx}{3a^2} - \frac{3(2A-B) \tan(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B) \tan(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{24} \\
& \frac{\frac{2a^2(5A-2B) \tan(c+dx)}{d} - 3a^2(2A-B) \int \csc(c+dx+\frac{\pi}{2}) dx}{3a^2} - \frac{3(2A-B) \tan(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B) \tan(c+dx)}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{4257} \\
& \frac{\frac{2a^2(5A-2B) \tan(c+dx)}{d} - \frac{3a^2(2A-B) \operatorname{arctanh}(\sin(c+dx))}{d}}{3a^2} - \frac{3(2A-B) \tan(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B) \tan(c+dx)}{3d(a \cos(c+dx) + a)^2}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^2,x]`

output `-1/3*((A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + ((-3*(2*A - B)*Tan[c + d*x])/(d*(1 + Cos[c + d*x])) + ((-3*a^2*(2*A - B)*ArcTanh[Sin[c + d*x]])/d + (2*a^2*(5*A - 2*B)*Tan[c + d*x])/d)/a^2)/(3*a^2)`

3.53.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.53.4 Maple [A] (verified)

Time = 1.44 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.18

method	result
parallelrisch	$\frac{12 \cos(dx+c) \left(A - \frac{B}{2}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - 12 \cos(dx+c) \left(A - \frac{B}{2}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 14 \left(\left(\frac{5A}{14} - \frac{B}{7}\right) \cos(2dx+2c) + \left(\frac{5A}{14} - \frac{B}{7}\right) \cos(2dx+c)\right)}{6d a^2 \cos(dx+c)}$
derivativedivides	$\frac{\left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)A - \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)B + 5A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (2B - 4A) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{2A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}}{2d a^2}$
default	$\frac{\left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)A - \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}\right)B + 5A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (2B - 4A) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{2A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}}{2d a^2}$
norman	$\frac{\frac{(A-B)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} - \frac{3(3A-B)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2ad} + \frac{(5A-3B)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2ad} - \frac{(13A-B)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)a} + \frac{(2A-B)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^2d}$
risch	$\frac{2i(6A e^{4i(dx+c)} - 3B e^{4i(dx+c)} + 18A e^{3i(dx+c)} - 9B e^{3i(dx+c)} + 22A e^{2i(dx+c)} - 7B e^{2i(dx+c)} + 24A e^{i(dx+c)} - 9B e^{i(dx+c)} - 1)}{3d a^2 (e^{i(dx+c)} + 1)^3 (e^{2i(dx+c)} + 1)}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)
```

```
output 1/6*(12*cos(d*x+c)*(A-1/2*B)*ln(tan(1/2*d*x+1/2*c)-1)-12*cos(d*x+c)*(A-1/2*B)*ln(tan(1/2*d*x+1/2*c)+1)+14*((5/14*A-1/7*B)*cos(2*d*x+2*c)+(A-5/14*B)*cos(d*x+c)+4/7*A-1/7*B)*tan(1/2*d*x+1/2*c)*sec(1/2*d*x+1/2*c)^2/d/a^2/cos(d*x+c)
```

3.53.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. 2(103) = 206.

Time = 0.29 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.93

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{3 \left((2A - B) \cos(dx + c)^3 + 2(2A - B) \cos(dx + c)^2 + (2A - B) \cos(dx + c) \right) \log(\sin(dx + c) + 1) - 1}{3d a^2 (e^{i(dx+c)} + 1)^3 (e^{2i(dx+c)} + 1)}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^2,x, algorithm="fracas")
```

output
$$\frac{-1/6*(3*((2*A - B)*\cos(d*x + c))^3 + 2*(2*A - B)*\cos(d*x + c)^2 + (2*A - B)*\cos(d*x + c))*\log(\sin(d*x + c) + 1) - 3*((2*A - B)*\cos(d*x + c))^3 + 2*(2*A - B)*\cos(d*x + c)^2 + (2*A - B)*\cos(d*x + c))*\log(-\sin(d*x + c) + 1) - 2*(2*(5*A - 2*B)*\cos(d*x + c)^2 + (14*A - 5*B)*\cos(d*x + c) + 3*A)*\sin(d*x + c))/(a^2*d*\cos(d*x + c)^3 + 2*a^2*d*\cos(d*x + c)^2 + a^2*d*\cos(d*x + c))$$

3.53.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\int \frac{A \sec^2(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sec^2(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c))**2,x)`

output `(Integral(A*sec(c + d*x)**2/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x))/a**2`

3.53.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. $2(103) = 206$.

Time = 0.26 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.28

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{A \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{12 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^2} + \frac{12 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^2} + \frac{12 \sin(dx+c)}{\left(a^2 - \frac{a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2}\right)(\cos(dx+c)+1)} \right) - B}{6d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output $\frac{1}{6} \left(A \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right) / a^2 - 12 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right) / a^2 + 12 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right) / a^2 + 12 \sin(dx+c) / ((a^2 - a^2 \sin(dx+c))^2 / (\cos(dx+c)+1)^2 * (\cos(dx+c)+1)) \right) - B \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right) / a^2 - 6 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right) / a^2 + 6 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right) / a^2 \right) / d$

3.53.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.45

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{6(2A-B) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right)}{a^2} - \frac{6(2A-B) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right)}{a^2} + \frac{12A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 1\right) a^2} - \frac{Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{6d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

output $\frac{-1}{6} \left(6 \left(2A - B \right) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) / a^2 - 6 \left(2A - B \right) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right) / a^2 + 12A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) / \left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 1\right) a^2 - \left(Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 15Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 9Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right) / a^6 \right) / d$

3.53.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.15

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{A-B}{a^2} + \frac{3A-B}{2a^2}\right)}{d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A-B)}{6a^2 d} - \frac{2A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - a^2\right)} - \frac{2a \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (2A-B)}{a^2 d}$$

3.53. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^2} dx$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^2),x)`

output `(tan(c/2 + (d*x)/2)*((A - B)/a^2 + (3*A - B)/(2*a^2)))/d + (tan(c/2 + (d*x)/2)^3*(A - B))/(6*a^2*d) - (2*A*tan(c/2 + (d*x)/2))/(d*(a^2*tan(c/2 + (d*x)/2)^2 - a^2)) - (2*atanh(tan(c/2 + (d*x)/2))*(2*A - B))/(a^2*d)`

3.54
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^2} dx$$

3.54.1 Optimal result 651
 3.54.2 Mathematica [B] (verified) 652
 3.54.3 Rubi [A] (verified) 652
 3.54.4 Maple [A] (verified) 656
 3.54.5 Fricas [A] (verification not implemented) 656
 3.54.6 Sympy [F] 657
 3.54.7 Maxima [B] (verification not implemented) 657
 3.54.8 Giac [A] (verification not implemented) 658
 3.54.9 Mupad [B] (verification not implemented) 659

3.54.1 Optimal result

Integrand size = 31, antiderivative size = 152

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^2} dx = \frac{(7A - 4B) \operatorname{arctanh}(\sin(c + dx))}{2a^2d} - \frac{2(8A - 5B) \tan(c + dx)}{3a^2d} + \frac{(7A - 4B) \sec(c + dx) \tan(c + dx)}{2a^2d} - \frac{(8A - 5B) \sec(c + dx) \tan(c + dx)}{3a^2d(1 + \cos(c + dx))} - \frac{(A - B) \sec(c + dx) \tan(c + dx)}{3d(a + a \cos(c + dx))^2}$$

output

```
1/2*(7*A-4*B)*arctanh(sin(d*x+c))/a^2/d-2/3*(8*A-5*B)*tan(d*x+c)/a^2/d+1/2
*(7*A-4*B)*sec(d*x+c)*tan(d*x+c)/a^2/d-1/3*(8*A-5*B)*sec(d*x+c)*tan(d*x+c)
/a^2/d/(1+cos(d*x+c))-1/3*(A-B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^2
```

3.54.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 496 vs. $2(152) = 304$.

Time = 2.94 (sec) , antiderivative size = 496, normalized size of antiderivative = 3.26

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^2} dx =$$

$$\frac{96(7A - 4B) \cos^4\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) + \sin\left(\frac{1}{2}(c + dx)\right)}{a^2 d (1 + \cos(c + dx))^2}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^2,x]`

output

```
-1/48*(96*(7*A - 4*B)*Cos[(c + d*x)/2]^4*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Cos[(c + d*x)/2]*Sec[c/2]*Sec[c]*Sec[c + d*x]^2*(-14*(A - B)*Sin[(d*x)/2] + (97*A - 64*B)*Sin[(3*d*x)/2] - 126*A*Sin[c - (d*x)/2] + 84*B*Sin[c - (d*x)/2] + 42*A*Sin[c + (d*x)/2] - 42*B*Sin[c + (d*x)/2] - 98*A*Sin[2*c + (d*x)/2] + 56*B*Sin[2*c + (d*x)/2] - 3*A*Sin[c + (3*d*x)/2] + 6*B*Sin[c + (3*d*x)/2] + 37*A*Sin[2*c + (3*d*x)/2] - 34*B*Sin[2*c + (3*d*x)/2] - 63*A*Sin[3*c + (3*d*x)/2] + 36*B*Sin[3*c + (3*d*x)/2] + 75*A*Sin[c + (5*d*x)/2] - 48*B*Sin[c + (5*d*x)/2] + 15*A*Sin[2*c + (5*d*x)/2] - 6*B*Sin[2*c + (5*d*x)/2] + 39*A*Sin[3*c + (5*d*x)/2] - 30*B*Sin[3*c + (5*d*x)/2] - 21*A*Sin[4*c + (5*d*x)/2] + 12*B*Sin[4*c + (5*d*x)/2] + 32*A*Sin[2*c + (7*d*x)/2] - 20*B*Sin[2*c + (7*d*x)/2] + 12*A*Sin[3*c + (7*d*x)/2] - 6*B*Sin[3*c + (7*d*x)/2] + 20*A*Sin[4*c + (7*d*x)/2] - 14*B*Sin[4*c + (7*d*x)/2]))/(a^2*d*(1 + Cos[c + d*x])^2)
```

3.54.3 Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.99, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3457, 3042, 3457, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx$$

↓ 3042

3.54. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^3 (a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^2} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(a(5A-2B) - 3a(A-B)\cos(c+dx)) \sec^3(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(5A-2B) - 3a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3 (\sin(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3457} \\
& \frac{\int (3a^2(7A-4B) - 2a^2(8A-5B)\cos(c+dx)) \sec^3(c+dx) dx}{a^2} - \frac{(8A-5B)\tan(c+dx)\sec(c+dx)}{d(\cos(c+dx)+1)} \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3a^2(7A-4B) - 2a^2(8A-5B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3} dx}{a^2} - \frac{(8A-5B)\tan(c+dx)\sec(c+dx)}{d(\cos(c+dx)+1)} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3227} \\
& \frac{3a^2(7A-4B) \int \sec^3(c+dx) dx - 2a^2(8A-5B) \int \sec^2(c+dx) dx}{a^2} - \frac{(8A-5B)\tan(c+dx)\sec(c+dx)}{d(\cos(c+dx)+1)} \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3a^2(7A-4B) \int \csc(c+dx+\frac{\pi}{2})^3 dx - 2a^2(8A-5B) \int \csc(c+dx+\frac{\pi}{2})^2 dx}{a^2} - \frac{(8A-5B)\tan(c+dx)\sec(c+dx)}{d(\cos(c+dx)+1)} \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{4254} \\
& \frac{2a^2(8A-5B) \int \frac{1d(-\tan(c+dx))}{d} + 3a^2(7A-4B) \int \csc(c+dx+\frac{\pi}{2})^3 dx}{a^2} - \frac{(8A-5B)\tan(c+dx)\sec(c+dx)}{d(\cos(c+dx)+1)} \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{3a^2}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{24}
\end{aligned}$$

3.54. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^2} dx$

$$\frac{3a^2(7A-4B) \int \csc(c+dx+\frac{\pi}{2})^3 dx - \frac{2a^2(8A-5B) \tan(c+dx)}{d} - \frac{(8A-5B) \tan(c+dx) \sec(c+dx)}{d(\cos(c+dx)+1)}}{a^2} - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{3d(a \cos(c+dx) + a)^2}$$

↓ 4255

$$\frac{3a^2(7A-4B) \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{2a^2(8A-5B) \tan(c+dx)}{d} - \frac{(8A-5B) \tan(c+dx) \sec(c+dx)}{d(\cos(c+dx)+1)}}{a^2} - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{3d(a \cos(c+dx) + a)^2}$$

↓ 3042

$$\frac{3a^2(7A-4B) \left(\frac{1}{2} \int \csc(c+dx+\frac{\pi}{2}) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{2a^2(8A-5B) \tan(c+dx)}{d} - \frac{(8A-5B) \tan(c+dx) \sec(c+dx)}{d(\cos(c+dx)+1)}}{a^2} - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{3d(a \cos(c+dx) + a)^2}$$

↓ 4257

$$\frac{3a^2(7A-4B) \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{2a^2(8A-5B) \tan(c+dx)}{d} - \frac{(8A-5B) \tan(c+dx) \sec(c+dx)}{d(\cos(c+dx)+1)}}{a^2} - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{3d(a \cos(c+dx) + a)^2}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^2,x]`

output `-1/3*((A - B)*Sec[c + d*x]*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + (-((8*A - 5*B)*Sec[c + d*x]*Tan[c + d*x])/(d*(1 + Cos[c + d*x]))) + ((-2*a^2*(8*A - 5*B)*Tan[c + d*x])/d + 3*a^2*(7*A - 4*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/a^2)/(3*a^2)`

3.54.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

3.54. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^2} dx$

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d)), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.54.4 Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.04

method	result
parallelrisc	$\frac{-42(1+\cos(2dx+2c))(A-\frac{4B}{7})\ln(\tan(\frac{dx}{2}+\frac{c}{2})-1)+42(1+\cos(2dx+2c))(A-\frac{4B}{7})\ln(\tan(\frac{dx}{2}+\frac{c}{2})+1)-60\tan(\frac{dx}{2}+\frac{c}{2})}{12da^2(1+\cos(2dx+2c))}$
derivativedivides	$\frac{-\frac{(\tan^3(\frac{dx}{2}+\frac{c}{2}))^A}{3}+\frac{(\tan^3(\frac{dx}{2}+\frac{c}{2}))^B}{3}-7A\tan(\frac{dx}{2}+\frac{c}{2})+5B\tan(\frac{dx}{2}+\frac{c}{2})-\frac{-5A+2B}{\tan(\frac{dx}{2}+\frac{c}{2})+1}+(7A-4B)\ln(\tan(\frac{dx}{2}+\frac{c}{2}))}{2da^2}$
default	$\frac{-\frac{(\tan^3(\frac{dx}{2}+\frac{c}{2}))^A}{3}+\frac{(\tan^3(\frac{dx}{2}+\frac{c}{2}))^B}{3}-7A\tan(\frac{dx}{2}+\frac{c}{2})+5B\tan(\frac{dx}{2}+\frac{c}{2})-\frac{-5A+2B}{\tan(\frac{dx}{2}+\frac{c}{2})+1}+(7A-4B)\ln(\tan(\frac{dx}{2}+\frac{c}{2}))}{2da^2}$
norman	$\frac{\frac{(A-B)(\tan^9(\frac{dx}{2}+\frac{c}{2}))}{6ad}-\frac{(10A-7B)(\tan^7(\frac{dx}{2}+\frac{c}{2}))}{3ad}-\frac{(13A-9B)\tan(\frac{dx}{2}+\frac{c}{2})}{2ad}+\frac{2(13A-7B)(\tan^5(\frac{dx}{2}+\frac{c}{2}))}{3ad}+\frac{(16A-7B)(\tan^3(\frac{dx}{2}+\frac{c}{2}))}{3ad}}{(1+\tan^2(\frac{dx}{2}+\frac{c}{2}))(\tan^2(\frac{dx}{2}+\frac{c}{2})-1)^2 a}$
risc	$\frac{-i(21Ae^{6i(dx+c)}-12Be^{6i(dx+c)}+63Ae^{5i(dx+c)}-36Be^{5i(dx+c)}+98Ae^{4i(dx+c)}-56Be^{4i(dx+c)}+126Ae^{3i(dx+c)}-84Ae^{2i(dx+c)}+21Ae^{i(dx+c)}-12Be^{i(dx+c)}+6A-6)}{3da^2(e^{i(dx+c)}+1)^3(e^{2i(dx+c)}+1)}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{12}*(-42*(1+\cos(2*d*x+2*c))*(A-4/7*B)*\ln(\tan(1/2*d*x+1/2*c)-1)+42*(1+\cos(2*d*x+2*c))*(A-4/7*B)*\ln(\tan(1/2*d*x+1/2*c)+1)-60*\tan(1/2*d*x+1/2*c)*((43/60*A-7/15*B)*\cos(2*d*x+2*c)+(4/15*A-1/6*B)*\cos(3*d*x+3*c)+(A-7/10*B)*\cos(d*x+c)+37/60*A-7/15*B)*\sec(1/2*d*x+1/2*c)^2)/d/a^2/(1+\cos(2*d*x+2*c))$$

3.54.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.50

$$\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{3((7A-4B)\cos(dx+c)^4+2(7A-4B)\cos(dx+c)^3+(7A-4B)\cos(dx+c)^2)\log(\sin(dx+c))+3(7A-4B)\cos(dx+c)}{(a+a\cos(c+dx))^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^2,x, algorithm="fracas")`

3.54.
$$\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^2} dx$$

output $1/12*(3*((7*A - 4*B)*\cos(d*x + c)^4 + 2*(7*A - 4*B)*\cos(d*x + c)^3 + (7*A - 4*B)*\cos(d*x + c)^2)*\log(\sin(d*x + c) + 1) - 3*((7*A - 4*B)*\cos(d*x + c)^4 + 2*(7*A - 4*B)*\cos(d*x + c)^3 + (7*A - 4*B)*\cos(d*x + c)^2)*\log(-\sin(d*x + c) + 1) - 2*(4*(8*A - 5*B)*\cos(d*x + c)^3 + (43*A - 28*B)*\cos(d*x + c)^2 + 6*(A - B)*\cos(d*x + c) - 3*A)*\sin(d*x + c))/(a^2*d*\cos(d*x + c)^4 + 2*a^2*d*\cos(d*x + c)^3 + a^2*d*\cos(d*x + c)^2)$

3.54.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\int \frac{A \sec^3(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sec^3(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c))**2,x)`

output `(Integral(A*sec(c + d*x)**3/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**3/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x))/a**2`

3.54.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 336 vs. $2(142) = 284$.

Time = 0.26 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.21

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^2} dx =$$

$$A \left(\frac{6 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - \frac{5 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^2 - \frac{2 a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^2 \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} + \frac{21 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{21 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^2} + \frac{21 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^2} \right) - B d$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output
$$\begin{aligned} & -1/6*(A*(6*(3*\sin(dx + c)/(\cos(dx + c) + 1) - 5*\sin(dx + c)^3/(\cos(dx + c) + 1)^3)/(a^2 - 2*a^2*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + a^2*\sin(dx + c)^4/(\cos(dx + c) + 1)^4) + (21*\sin(dx + c)/(\cos(dx + c) + 1) + \sin(dx + c)^3/(\cos(dx + c) + 1)^3)/a^2 - 21*\log(\sin(dx + c)/(\cos(dx + c) + 1) + 1)/a^2 + 21*\log(\sin(dx + c)/(\cos(dx + c) + 1) - 1)/a^2) - B*((15*\sin(dx + c)/(\cos(dx + c) + 1) + \sin(dx + c)^3/(\cos(dx + c) + 1)^3)/a^2 - 12*\log(\sin(dx + c)/(\cos(dx + c) + 1) + 1)/a^2 + 12*\log(\sin(dx + c)/(\cos(dx + c) + 1) - 1)/a^2 + 12*\sin(dx + c)/((a^2 - a^2*\sin(dx + c)^2/(\cos(dx + c) + 1)^2)*(cos(dx + c) + 1))))/d \end{aligned}$$

3.54.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.30

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{3(7A-4B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^2} - \frac{3(7A-4B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^2} + \frac{6(5A \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 2B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 3A \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1)^2 a^2}$$

$6d$

input `integrate((A+B*cos(dx+c))*sec(dx+c)^3/(a+a*cos(dx+c))^2,x, algorithm="giac")`

output
$$\begin{aligned} & 1/6*(3*(7*A - 4*B)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1))/a^2 - 3*(7*A - 4*B)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1))/a^2 + 6*(5*A*\tan(1/2*d*x + 1/2*c)^3 - 2*B*\tan(1/2*d*x + 1/2*c)^3 - 3*A*\tan(1/2*d*x + 1/2*c) + 2*B*\tan(1/2*d*x + 1/2*c))/((\tan(1/2*d*x + 1/2*c)^2 - 1)^2*a^2) - (A*a^4*\tan(1/2*d*x + 1/2*c)^3 - B*a^4*\tan(1/2*d*x + 1/2*c)^3 + 21*A*a^4*\tan(1/2*d*x + 1/2*c) - 15*B*a^4*\tan(1/2*d*x + 1/2*c))/a^6)/d \end{aligned}$$

3.54.9 Mupad [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.09

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (5A - 2B) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (3A - 2B)}{d \left(a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^2 \right)}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3(A-B)}{2a^2} + \frac{4A-2B}{2a^2} \right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A - B)}{6a^2 d}$$

$$+ \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (7A - 4B)}{a^2 d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^2),x)`output `(tan(c/2 + (d*x)/2)^3*(5*A - 2*B) - tan(c/2 + (d*x)/2)*(3*A - 2*B))/(d*(a^2*tan(c/2 + (d*x)/2)^4 - 2*a^2*tan(c/2 + (d*x)/2)^2 + a^2)) - (tan(c/2 + (d*x)/2)*((3*(A - B))/(2*a^2) + (4*A - 2*B)/(2*a^2)))/d - (tan(c/2 + (d*x)/2)^3*(A - B))/(6*a^2*d) + (atanh(tan(c/2 + (d*x)/2))*(7*A - 4*B))/(a^2*d)`

3.55 $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{(a+a \cos(c+dx))^2} dx$

3.55.1	Optimal result	660
3.55.2	Mathematica [B] (verified)	661
3.55.3	Rubi [A] (verified)	661
3.55.4	Maple [A] (verified)	665
3.55.5	Fricas [A] (verification not implemented)	666
3.55.6	Sympy [F]	666
3.55.7	Maxima [B] (verification not implemented)	667
3.55.8	Giac [A] (verification not implemented)	667
3.55.9	Mupad [B] (verification not implemented)	668

3.55.1 Optimal result

Integrand size = 31, antiderivative size = 179

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx = -\frac{(10A - 7B) \operatorname{arctanh}(\sin(c + dx))}{2a^2d} + \frac{4(3A - 2B) \tan(c + dx)}{a^2d} - \frac{(10A - 7B) \sec(c + dx) \tan(c + dx)}{2a^2d} - \frac{(10A - 7B) \sec^2(c + dx) \tan(c + dx)}{3a^2d(1 + \cos(c + dx))} - \frac{(A - B) \sec^2(c + dx) \tan(c + dx)}{3d(a + a \cos(c + dx))^2} + \frac{4(3A - 2B) \tan^3(c + dx)}{3a^2d}$$

output

```
-1/2*(10*A-7*B)*arctanh(sin(d*x+c))/a^2/d+4*(3*A-2*B)*tan(d*x+c)/a^2/d-1/2
*(10*A-7*B)*sec(d*x+c)*tan(d*x+c)/a^2/d-1/3*(10*A-7*B)*sec(d*x+c)^2*tan(d*
x+c)/a^2/d/(1+cos(d*x+c))-1/3*(A-B)*sec(d*x+c)^2*tan(d*x+c)/d/(a+a*cos(d*x
+c))^2+4/3*(3*A-2*B)*tan(d*x+c)^3/a^2/d
```

3.55.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 609 vs. $2(179) = 358$.

Time = 4.02 (sec) , antiderivative size = 609, normalized size of antiderivative = 3.40

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{192(10A - 7B) \cos^4\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{a^2}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + a*Cos[c + d*x])^2,x]`

output

```
(192*(10*A - 7*B)*Cos[(c + d*x)/2]^4*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Cos[(c + d*x)/2]*Sec[c/2]*Sec[c + d*x]^3*((-6*A + 45*B)*Sin[(d*x)/2] + (310*A - 201*B)*Sin[(3*d*x)/2] - 306*A*Sin[c - (d*x)/2] + 195*B*Sin[c - (d*x)/2] + 42*A*Sin[c + (d*x)/2] - 51*B*Sin[c + (d*x)/2] - 270*A*Sin[2*c + (d*x)/2] + 189*B*Sin[2*c + (d*x)/2] + 50*A*Sin[c + (3*d*x)/2] - B*Sin[c + (3*d*x)/2] + 90*A*Sin[2*c + (3*d*x)/2] - 81*B*Sin[2*c + (3*d*x)/2] - 170*A*Sin[3*c + (3*d*x)/2] + 119*B*Sin[3*c + (3*d*x)/2] + 198*A*Sin[c + (5*d*x)/2] - 129*B*Sin[c + (5*d*x)/2] + 42*A*Sin[2*c + (5*d*x)/2] - 9*B*Sin[2*c + (5*d*x)/2] + 66*A*Sin[3*c + (5*d*x)/2] - 57*B*Sin[3*c + (5*d*x)/2] - 90*A*Sin[4*c + (5*d*x)/2] + 63*B*Sin[4*c + (5*d*x)/2] + 114*A*Sin[2*c + (7*d*x)/2] - 75*B*Sin[2*c + (7*d*x)/2] + 36*A*Sin[3*c + (7*d*x)/2] - 15*B*Sin[3*c + (7*d*x)/2] + 48*A*Sin[4*c + (7*d*x)/2] - 39*B*Sin[4*c + (7*d*x)/2] - 30*A*Sin[5*c + (7*d*x)/2] + 21*B*Sin[5*c + (7*d*x)/2] + 48*A*Sin[3*c + (9*d*x)/2] - 32*B*Sin[3*c + (9*d*x)/2] + 22*A*Sin[4*c + (9*d*x)/2] - 12*B*Sin[4*c + (9*d*x)/2] + 26*A*Sin[5*c + (9*d*x)/2] - 20*B*Sin[5*c + (9*d*x)/2]))/(96*a^2*d*(1 + Cos[c + d*x])^2)
```

3.55.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.96, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3457, 3042, 3457, 27, 3042, 3227, 3042, 4254, 2009, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.55. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{\sec^4(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A+B\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^4 (a\sin\left(c+dx+\frac{\pi}{2}\right)+a)^2} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(3a(2A-B)-4a(A-B)\cos(c+dx))\sec^4(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} - \frac{(A-B)\tan(c+dx)\sec^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3a(2A-B)-4a(A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^4 (\sin\left(c+dx+\frac{\pi}{2}\right)a+a)} dx}{3a^2} - \frac{(A-B)\tan(c+dx)\sec^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3457} \\
& \frac{\int 3(4a^2(3A-2B)-a^2(10A-7B)\cos(c+dx))\sec^4(c+dx) dx}{a^2} - \frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)} - \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)} \\
& \quad \frac{(A-B)\tan(c+dx)\sec^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{3\int (4a^2(3A-2B)-a^2(10A-7B)\cos(c+dx))\sec^4(c+dx) dx}{a^2} - \frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)} - \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)} \\
& \quad \frac{(A-B)\tan(c+dx)\sec^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3\int \frac{4a^2(3A-2B)-a^2(10A-7B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^4} dx}{a^2} - \frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)} - \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)} \\
& \quad \frac{(A-B)\tan(c+dx)\sec^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3227} \\
& \frac{3(4a^2(3A-2B)\int \sec^4(c+dx) dx - a^2(10A-7B)\int \sec^3(c+dx) dx)}{a^2} - \frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)} - \\
& \quad \frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)} \\
& \quad \frac{(A-B)\tan(c+dx)\sec^2(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.55. $\int \frac{(A+B\cos(c+dx))\sec^4(c+dx)}{(a+a\cos(c+dx))^2} dx$

$$\frac{3\left(4a^2(3A-2B)\int\csc(c+dx+\frac{\pi}{2})^4dx-a^2(10A-7B)\int\csc(c+dx+\frac{\pi}{2})^3dx\right)}{a^2}-\frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)}-\frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)}\frac{3a^2}{3d(a\cos(c+dx)+a)^2}$$

↓ 4254

$$\frac{3\left(-\frac{4a^2(3A-2B)\int(\tan^2(c+dx)+1)d(-\tan(c+dx))}{d}-\left(a^2(10A-7B)\int\csc(c+dx+\frac{\pi}{2})^3dx\right)\right)}{a^2}-\frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)}-\frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)}\frac{3a^2}{3d(a\cos(c+dx)+a)^2}$$

↓ 2009

$$\frac{3\left(-a^2(10A-7B)\int\csc(c+dx+\frac{\pi}{2})^3dx-\frac{4a^2(3A-2B)\left(-\frac{1}{3}\tan^3(c+dx)-\tan(c+dx)\right)}{d}\right)}{a^2}-\frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)}-\frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)}\frac{3a^2}{3d(a\cos(c+dx)+a)^2}$$

↓ 4255

$$\frac{3\left(-a^2(10A-7B)\left(\frac{1}{2}\int\sec(c+dx)dx+\frac{\tan(c+dx)\sec(c+dx)}{2d}\right)-\frac{4a^2(3A-2B)\left(-\frac{1}{3}\tan^3(c+dx)-\tan(c+dx)\right)}{d}\right)}{a^2}-\frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)}-\frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)}\frac{3a^2}{3d(a\cos(c+dx)+a)^2}$$

↓ 3042

$$\frac{3\left(-a^2(10A-7B)\left(\frac{1}{2}\int\csc(c+dx+\frac{\pi}{2})dx+\frac{\tan(c+dx)\sec(c+dx)}{2d}\right)-\frac{4a^2(3A-2B)\left(-\frac{1}{3}\tan^3(c+dx)-\tan(c+dx)\right)}{d}\right)}{a^2}-\frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)}-\frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)}\frac{3a^2}{3d(a\cos(c+dx)+a)^2}$$

↓ 4257

$$\frac{3\left(-\left(a^2(10A-7B)\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d}+\frac{\tan(c+dx)\sec(c+dx)}{2d}\right)\right)-\frac{4a^2(3A-2B)\left(-\frac{1}{3}\tan^3(c+dx)-\tan(c+dx)\right)}{d}\right)}{a^2}-\frac{(10A-7B)\tan(c+dx)\sec^2(c+dx)}{d(\cos(c+dx)+1)}-\frac{3a^2}{(A-B)\tan(c+dx)\sec^2(c+dx)}\frac{3a^2}{3d(a\cos(c+dx)+a)^2}$$

3.55. $\int \frac{(A+B\cos(c+dx))\sec^4(c+dx)}{(a+a\cos(c+dx))^2} dx$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + a*Cos[c + d*x])^2,x]`

output `-1/3*((A - B)*Sec[c + d*x]^2*Tan[c + d*x]/(d*(a + a*Cos[c + d*x])^2) + (-
(((10*A - 7*B)*Sec[c + d*x]^2*Tan[c + d*x]/(d*(1 + Cos[c + d*x]))) + (3*(
-(a^2*(10*A - 7*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*
x])/(2*d))) - (4*a^2*(3*A - 2*B)*(-Tan[c + d*x] - Tan[c + d*x]^3/3)/d))/a
^2)/(3*a^2)`

3.55.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_))*((A_) + (B_)*sin[(e_) +
(f_)*(x_)]*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d)), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
d(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 4254 `Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]`

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
  && IntegerQ[2*n]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
  /; FreeQ[{c, d}, x]
```

3.55.4 Maple [A] (verified)

Time = 1.40 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.09

method	result
parallelrisch	$\frac{180 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \left(A - \frac{7B}{10} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) - 180 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \left(A - \frac{7B}{10} \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{\left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^A - \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^B} + 9A \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 7B \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (-10A+7B) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \frac{-6A}{2 \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}$
derivativedivides	$\frac{\left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^A - \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^B}{3} + 9A \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 7B \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (-10A+7B) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \frac{-6A}{2 \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}$
default	$\frac{\left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^A - \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^B}{3} + 9A \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 7B \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (-10A+7B) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \frac{-6A}{2 \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}$
norman	$\frac{(A-B) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (11A-10B) \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - (19A-12B) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - (21A-13B) \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (25A-19B) \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{6ad} + \frac{(11A-10B) \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - (19A-12B) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - (21A-13B) \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (25A-19B) \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{3ad} - \frac{(19A-12B) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - (21A-13B) \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (25A-19B) \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{ad} - \frac{(21A-13B) \tan \left(\frac{dx}{2} + \frac{c}{2} \right) + (25A-19B) \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{2ad} + \frac{(25A-19B) \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{6ad}$
risch	$\frac{i(30A e^{8i(dx+c)} - 21B e^{8i(dx+c)} + 90A e^{7i(dx+c)} - 63B e^{7i(dx+c)} + 170A e^{6i(dx+c)} - 119B e^{6i(dx+c)} + 270A e^{5i(dx+c)} - 189B e^{5i(dx+c)} + 126A e^{4i(dx+c)} - 105B e^{4i(dx+c)} + 63A e^{3i(dx+c)} - 52B e^{3i(dx+c)} + 27A e^{2i(dx+c)} - 21B e^{2i(dx+c)} + 9A e^{i(dx+c)} - 7B e^{i(dx+c)} + 3A - 2B)}{(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right)) \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)^3 a}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+cos(d*x+c)*a)^2,x,method=_RETURNVERBO
SE)
```

```
output 1/12*(180*(1/3*cos(3*d*x+3*c)+cos(d*x+c))*(A-7/10*B)*ln(tan(1/2*d*x+1/2*c)
-1)-180*(1/3*cos(3*d*x+3*c)+cos(d*x+c))*(A-7/10*B)*ln(tan(1/2*d*x+1/2*c)+1
)+24*sec(1/2*d*x+1/2*c)^2*((11/4*A-43/24*B)*cos(3*d*x+3*c)+(5*A-19/6*B)*co
s(2*d*x+2*c)+(A-2/3*B)*cos(4*d*x+4*c)+(95/12*A-39/8*B)*cos(d*x+c)+13/3*A-5
/2*B)*tan(1/2*d*x+1/2*c))/d/a^2/(cos(3*d*x+3*c)+3*cos(d*x+c))
```

$$3.55. \int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{(a+a \cos(c+dx))^2} dx$$

3.55.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.38

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx =$$

$$\frac{3((10A - 7B) \cos(dx + c)^5 + 2(10A - 7B) \cos(dx + c)^4 + (10A - 7B) \cos(dx + c)^3) \log(\sin(dx + c) + 1) - 3((10A - 7B) \cos(dx + c)^5 + 2(10A - 7B) \cos(dx + c)^4 + (10A - 7B) \cos(dx + c)^3) \log(-\sin(dx + c) + 1) - 2(16(3A - 2B) \cos(dx + c)^4 + (66A - 43B) \cos(dx + c)^3 + 6(2A - B) \cos(dx + c)^2 - (2A - 3B) \cos(dx + c) + 2A) \sin(dx + c)}{a^2 d \cos(dx + c)^5 + 2a^2 d \cos(dx + c)^4 + a^2 d \cos(dx + c)^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+a*cos(d*x+c))^2,x, algorithm="fricas")`

output `-1/12*(3*((10*A - 7*B)*cos(d*x + c)^5 + 2*(10*A - 7*B)*cos(d*x + c)^4 + (10*A - 7*B)*cos(d*x + c)^3)*log(sin(d*x + c) + 1) - 3*((10*A - 7*B)*cos(d*x + c)^5 + 2*(10*A - 7*B)*cos(d*x + c)^4 + (10*A - 7*B)*cos(d*x + c)^3)*log(-sin(d*x + c) + 1) - 2*(16*(3*A - 2*B)*cos(d*x + c)^4 + (66*A - 43*B)*cos(d*x + c)^3 + 6*(2*A - B)*cos(d*x + c)^2 - (2*A - 3*B)*cos(d*x + c) + 2*A)*sin(d*x + c))/(a^2*d*cos(d*x + c)^5 + 2*a^2*d*cos(d*x + c)^4 + a^2*d*cos(d*x + c)^3)`

3.55.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\int \frac{A \sec^4(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sec^4(c+dx)}{\cos^2(c+dx)+2 \cos(c+dx)+1} dx}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**4/(a+a*cos(d*x+c))**2,x)`

output `(Integral(A*sec(c + d*x)**4/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**4/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x))/a**2`

3.55.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 425 vs. $2(169) = 338$.

Time = 0.23 (sec) , antiderivative size = 425, normalized size of antiderivative = 2.37

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= A \left(\frac{4 \left(\frac{9 \sin(dx+c)}{\cos(dx+c)+1} - \frac{20 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{15 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a^2 - \frac{3a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3a^2 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - \frac{a^2 \sin(dx+c)^6}{(\cos(dx+c)+1)^6}} + \frac{27 \sin(dx+c)}{\cos(dx+c)+1} + \frac{\sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{30 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^2} + \frac{30 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^2} \right)$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `1/6*(A*(4*(9*sin(d*x + c)/(cos(d*x + c) + 1) - 20*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 15*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(a^2 - 3*a^2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*a^2*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - a^2*sin(d*x + c)^6/(cos(d*x + c) + 1)^6) + (27*sin(d*x + c)/(cos(d*x + c) + 1) + sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2 - 30*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a^2 + 30*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a^2) - B*(6*(3*sin(d*x + c)/(cos(d*x + c) + 1) - 5*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(a^2 - 2*a^2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a^2*sin(d*x + c)^4/(cos(d*x + c) + 1)^4) + (21*sin(d*x + c)/(cos(d*x + c) + 1) + sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/a^2 - 21*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a^2 + 21*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a^2))/d`

3.55.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.26

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx =$$

$$\frac{3(10A - 7B) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right)}{a^2} - \frac{3(10A - 7B) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right)}{a^2} + \frac{2\left(30A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 15B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 40A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 20B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3\right)}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+a*cos(d*x+c))^2,x, algorithm="giac")`

3.55. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{(a+a \cos(c+dx))^2} dx$

```
output -1/6*(3*(10*A - 7*B)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^2 - 3*(10*A - 7*
B)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^2 + 2*(30*A*tan(1/2*d*x + 1/2*c)^5
- 15*B*tan(1/2*d*x + 1/2*c)^5 - 40*A*tan(1/2*d*x + 1/2*c)^3 + 24*B*tan(1/
2*d*x + 1/2*c)^3 + 18*A*tan(1/2*d*x + 1/2*c) - 9*B*tan(1/2*d*x + 1/2*c))/(
(tan(1/2*d*x + 1/2*c)^2 - 1)^3*a^2) - (A*a^4*tan(1/2*d*x + 1/2*c)^3 - B*a^
4*tan(1/2*d*x + 1/2*c)^3 + 27*A*a^4*tan(1/2*d*x + 1/2*c) - 21*B*a^4*tan(1/
2*d*x + 1/2*c))/a^6)/d
```

3.55.9 Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.13

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{2(A-B)}{a^2} + \frac{5A-3B}{2a^2}\right)}{d}$$

$$- \frac{(10A - 5B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + (8B - \frac{40A}{3}) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (6A - 3B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - a^2\right)}$$

$$+ \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A - B)}{6a^2 d} - \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (10A - 7B)}{a^2 d}$$

```
input int((A + B*cos(c + d*x))/(cos(c + d*x)^4*(a + a*cos(c + d*x))^2),x)
```

```
output (tan(c/2 + (d*x)/2)*((2*(A - B))/a^2 + (5*A - 3*B)/(2*a^2)))/d - (tan(c/2
+ (d*x)/2)^5*(10*A - 5*B) - tan(c/2 + (d*x)/2)^3*((40*A)/3 - 8*B) + tan(c/
2 + (d*x)/2)*(6*A - 3*B))/(d*(3*a^2*tan(c/2 + (d*x)/2)^2 - 3*a^2*tan(c/2 +
(d*x)/2)^4 + a^2*tan(c/2 + (d*x)/2)^6 - a^2)) + (tan(c/2 + (d*x)/2)^3*(A
- B))/(6*a^2*d) - (atanh(tan(c/2 + (d*x)/2))*(10*A - 7*B))/(a^2*d)
```

3.56
$$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.56.1 Optimal result 669
 3.56.2 Mathematica [B] (verified) 670
 3.56.3 Rubi [A] (verified) 670
 3.56.4 Maple [A] (verified) 674
 3.56.5 Fricas [A] (verification not implemented) 675
 3.56.6 Sympy [B] (verification not implemented) 676
 3.56.7 Maxima [B] (verification not implemented) 677
 3.56.8 Giac [A] (verification not implemented) 677
 3.56.9 Mupad [B] (verification not implemented) 678

3.56.1 Optimal result

Integrand size = 31, antiderivative size = 218

$$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{(13A-23B)x}{2a^3} - \frac{4(19A-34B) \sin(c+dx)}{5a^3d} + \frac{(13A-23B) \cos(c+dx) \sin(c+dx)}{2a^3d} + \frac{(A-B) \cos^5(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(8A-13B) \cos^4(c+dx) \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} + \frac{(13A-23B) \cos^3(c+dx) \sin(c+dx)}{3d(a^3+a^3 \cos(c+dx))} + \frac{4(19A-34B) \sin^3(c+dx)}{15a^3d}$$

```
output 1/2*(13*A-23*B)*x/a^3-4/5*(19*A-34*B)*sin(d*x+c)/a^3/d+1/2*(13*A-23*B)*cos
(d*x+c)*sin(d*x+c)/a^3/d+1/5*(A-B)*cos(d*x+c)^5*sin(d*x+c)/d/(a+a*cos(d*x+
c))^3+1/15*(8*A-13*B)*cos(d*x+c)^4*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+1/3*(
13*A-23*B)*cos(d*x+c)^3*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))+4/15*(19*A-34*B)
*sin(d*x+c)^3/a^3/d
```

3.56.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 491 vs. $2(218) = 436$.

Time = 2.55 (sec) , antiderivative size = 491, normalized size of antiderivative = 2.25

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) (600(13A-23B)dx \cos\left(\frac{dx}{2}\right) + 600(13A-23B)dx \cos\left(c+\frac{dx}{2}\right) + 3900Adx \cos(c$$

input `Integrate[(Cos[c + d*x]^5*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output

$$\frac{(\cos\left(\frac{c+dx}{2}\right) \sec\left(\frac{c}{2}\right) (600(13A-23B)dx \cos\left(\frac{dx}{2}\right) + 600(13A-23B)dx \cos\left(c+\frac{dx}{2}\right) + 3900A dx \cos\left[c+\frac{3dx}{2}\right] - 6900B dx \cos\left[c+\frac{3dx}{2}\right] + 3900A dx \cos\left[2c+\frac{3dx}{2}\right] - 6900B dx \cos\left[2c+\frac{3dx}{2}\right] + 780A dx \cos\left[2c+\frac{5dx}{2}\right] - 1380B dx \cos\left[2c+\frac{5dx}{2}\right] + 780A dx \cos\left[3c+\frac{5dx}{2}\right] - 1380B dx \cos\left[3c+\frac{5dx}{2}\right] - 12760A \sin\left(\frac{dx}{2}\right) + 20410B \sin\left(\frac{dx}{2}\right) + 7560A \sin\left[c+\frac{dx}{2}\right] - 11110B \sin\left[c+\frac{dx}{2}\right] - 9230A \sin\left[c+\frac{3dx}{2}\right] + 15380B \sin\left[c+\frac{3dx}{2}\right] + 930A \sin\left[2c+\frac{3dx}{2}\right] - 380B \sin\left[2c+\frac{3dx}{2}\right] - 2782A \sin\left[2c+\frac{5dx}{2}\right] + 4777B \sin\left[2c+\frac{5dx}{2}\right] - 750A \sin\left[3c+\frac{5dx}{2}\right] + 1625B \sin\left[3c+\frac{5dx}{2}\right] - 105A \sin\left[3c+\frac{7dx}{2}\right] + 230B \sin\left[3c+\frac{7dx}{2}\right] - 105A \sin\left[4c+\frac{7dx}{2}\right] + 230B \sin\left[4c+\frac{7dx}{2}\right] + 15A \sin\left[4c+\frac{9dx}{2}\right] - 20B \sin\left[4c+\frac{9dx}{2}\right] + 15A \sin\left[5c+\frac{9dx}{2}\right] - 20B \sin\left[5c+\frac{9dx}{2}\right] + 5B \sin\left[5c+\frac{11dx}{2}\right] + 5B \sin\left[6c+\frac{11dx}{2}\right])}{(480a^3d(1+\cos[c+dx])^3)}$$

3.56.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.99, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3456, 3042, 3456, 3042, 3456, 27, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

3.56. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^5 \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^4(c+dx)(5a(A-B)-a(3A-8B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^4(5a(A-B)-a(3A-8B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^3(c+dx)(4a^2(8A-13B)-3a^2(11A-21B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{a(8A-13B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^5(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^3(4a^2(8A-13B)-3a^2(11A-21B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{a(8A-13B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^5(c+dx) \\
& \quad \downarrow \text{3456} \\
& \frac{\int 3\cos^2(c+dx)(5a^3(13A-23B)-4a^3(19A-34B)\cos(c+dx)) dx}{a^2} + \frac{5a^2(13A-23B)\sin(c+dx)\cos^3(c+dx)}{d(a\cos(c+dx)+a)} + \frac{a(8A-13B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^5(c+dx) \\
& \quad \downarrow \text{27} \\
& \frac{3\int \cos^2(c+dx)(5a^3(13A-23B)-4a^3(19A-34B)\cos(c+dx)) dx}{a^2} + \frac{5a^2(13A-23B)\sin(c+dx)\cos^3(c+dx)}{d(a\cos(c+dx)+a)} + \frac{a(8A-13B)\sin(c+dx)\cos^4(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^5(c+dx) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.56. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{3 \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(5a^3(13A-23B)-4a^3(19A-34B) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx}{a^2} + \frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{a(8A-13B) \sin(c+dx) \cos^4(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{5a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3227

$$\frac{3 \left(5a^3(13A-23B) \int \cos^2(c+dx) dx - 4a^3(19A-34B) \int \cos^3(c+dx) dx\right)}{a^2} + \frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{a(8A-13B) \sin(c+dx) \cos^4(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{5a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{3 \left(5a^3(13A-23B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx - 4a^3(19A-34B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 dx\right)}{a^2} + \frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{a(8A-13B) \sin(c+dx) \cos^4(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{5a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3113

$$\frac{3 \left(\frac{4a^3(19A-34B) \int (1-\sin^2(c+dx)) d(-\sin(c+dx))}{d} + 5a^3(13A-23B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx\right)}{a^2} + \frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{a(8A-13B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{5a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 2009

$$\frac{3 \left(5a^3(13A-23B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx + \frac{4a^3(19A-34B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx)\right)}{d}\right)}{a^2} + \frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{a(8A-13B) \sin(c+dx) \cos^4(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{5a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3115

3.56. $\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

$$\frac{3 \left(\frac{5a^3(13A-23B) \left(\frac{\int \frac{1dx}{2} + \frac{\sin(c+dx)\cos(c+dx)}{2d} \right) + \frac{4a^3(19A-34B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d}}{a^2} \right)}{3a^2} + \frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{a(8A-13B) \sin(c+dx)}{3d(a \cos(c+dx)+a)}$$

$$\frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 24

$$\frac{\frac{5a^2(13A-23B) \sin(c+dx) \cos^3(c+dx)}{d(a \cos(c+dx)+a)} + \frac{3 \left(\frac{4a^3(19A-34B) \left(\frac{1}{3} \sin^3(c+dx) - \sin(c+dx) \right)}{d} + \frac{5a^3(13A-23B) \left(\frac{\sin(c+dx)\cos(c+dx)}{2d} + \frac{x}{2} \right)}{a^2} \right)}{3a^2}}{3a^2} + \frac{a(8A-13B) \sin(c+dx)}{3d(a \cos(c+dx)+a)}$$

$$\frac{(A-B) \sin(c+dx) \cos^5(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

```
input Int[(Cos[c + d*x]^5*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]
```

```
output ((A - B)*Cos[c + d*x]^5*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((a*(8*A - 13*B)*Cos[c + d*x]^4*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((5*a^2*(13*A - 23*B)*Cos[c + d*x]^3*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])) + (3*(5*a^3*(13*A - 23*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) + (4*a^3*(19*A - 34*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/a^2)/(3*a^2))/(5*a^2)
```

3.56.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.56. $\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.56.4 Maple [A] (verified)

Time = 0.99 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.57

3.56.
$$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

method	result
parallelrisch	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\frac{8(-232A+427B)\cos(2dx+2c)}{15} + \left(-6A + \frac{43B}{3}\right) \cos(3dx+3c) + (A-B) \cos(4dx+4c) + \frac{B \cos(5dx+5c)}{3} + \frac{2(-1001A+545B)\cos(dx+c)}{15} \right)}{64a^3d}$
derivativedivides	$-\frac{A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5} + \frac{\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{5} + \frac{8 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} - \frac{10 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} - 31A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 49B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) +$
default	$-\frac{A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5} + \frac{\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{5} + \frac{8 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} - \frac{10 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} - 31A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 49B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) +$
risch	$\frac{13Ax}{2a^3} - \frac{23Bx}{2a^3} - \frac{iB e^{3i(dx+c)}}{24a^3d} - \frac{ie^{2i(dx+c)}A}{8a^3d} + \frac{3ie^{2i(dx+c)}B}{8a^3d} + \frac{3ie^{i(dx+c)}A}{2a^3d} - \frac{27ie^{i(dx+c)}B}{8a^3d} - \frac{3ie^{-i(dx+c)}}{2a^3d}$
norman	$\frac{(13A-23B)x}{2a} - \frac{(A-B)\left(\tan^{17}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{20ad} - \frac{(9A-16B)\left(\tan^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2ad} + \frac{(11A-16B)\left(\tan^{15}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{30ad} + \frac{3(13A-23B)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a}$

input `int(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE)`

output `1/64*(tan(1/2*d*x+1/2*c)*(8/15*(-232*A+427*B)*cos(2*d*x+2*c)+(-6*A+43/3*B)*cos(3*d*x+3*c)+(A-B)*cos(4*d*x+4*c)+1/3*B*cos(5*d*x+5*c)+2/5*(-1001*A+545B/3)*cos(dx+c)-4303/15*A+7783/15*B)*sec(1/2*d*x+1/2*c)^4+416*(A-23/13*B)*x*d)/a^3/d`

3.56.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.94

$$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

$$= \frac{15(13A-23B)dx \cos(dx+c)^3 + 45(13A-23B)dx \cos(dx+c)^2 + 45(13A-23B)dx \cos(dx+c) + \dots}{64a^3d}$$

input `integrate(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fracas")`

3.56. $\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.56.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 412 vs. $2(204) = 408$.

Time = 0.30 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.89

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{B \left(\frac{20 \left(\frac{33 \sin(dx+c)}{\cos(dx+c)+1} + \frac{76 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{51 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a^3 + \frac{3a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3a^3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{a^3 \sin(dx+c)^6}{(\cos(dx+c)+1)^6}} + \frac{735 \sin(dx+c)}{\cos(dx+c)+1} - \frac{50 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{1380 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^3} \right)}{60d}$$

input `integrate(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `1/60*(B*(20*(33*sin(d*x + c)/(cos(d*x + c) + 1) + 76*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 51*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/(a^3 + 3*a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*a^3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + a^3*sin(d*x + c)^6/(cos(d*x + c) + 1)^6) + (735*sin(d*x + c)/(cos(d*x + c) + 1) - 50*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 1380*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3) - A*(60*(5*sin(d*x + c)/(cos(d*x + c) + 1) + 7*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(a^3 + 2*a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a^3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4) + (465*sin(d*x + c)/(cos(d*x + c) + 1) - 40*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 780*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3))/d`

3.56.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.05

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{30(dx+c)(13A-23B)}{a^3} - \frac{20 \left(21A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 51B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 + 36A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 76B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 15A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) \right)}{\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1 \right)^3 a^3}$$

input `integrate(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="giac")`

3.56. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

output $\frac{1}{60} \cdot (30 \cdot (d \cdot x + c) \cdot (13 \cdot A - 23 \cdot B) / a^3 - 20 \cdot (21 \cdot A \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 - 5 \cdot 1 \cdot B \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 + 36 \cdot A \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 - 76 \cdot B \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 + 15 \cdot A \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) - 33 \cdot B \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)) / ((\tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 1)^3 \cdot a^3) - (3 \cdot A \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 - 3 \cdot B \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 - 40 \cdot A \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 + 50 \cdot B \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 + 465 \cdot A \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) - 735 \cdot B \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)) / a^{15}) / d$

3.56.9 Mupad [B] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.09

$$\int \frac{\cos^5(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{x(13A - 23B)}{2a^3} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{5(A-B)}{2a^3} + \frac{4A-6B}{a^3} + \frac{5A-15B}{4a^3}\right)}{d}$$

$$- \frac{(7A - 17B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + (12A - \frac{76B}{3}) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (5A - 11B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 3a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^3\right)}$$

$$+ \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{3a^3} + \frac{4A-6B}{12a^3}\right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A - B)}{20a^3 d}$$

input `int((cos(c + d*x))^5*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output $(x \cdot (13 \cdot A - 23 \cdot B)) / (2 \cdot a^3) - (\tan(c/2 + (d \cdot x)/2) \cdot ((5 \cdot (A - B)) / (2 \cdot a^3) + (4 \cdot A - 6 \cdot B) / a^3 + (5 \cdot A - 15 \cdot B) / (4 \cdot a^3))) / d - (\tan(c/2 + (d \cdot x)/2)^5 \cdot (7 \cdot A - 17 \cdot B) + \tan(c/2 + (d \cdot x)/2)^3 \cdot (12 \cdot A - (76 \cdot B) / 3) + \tan(c/2 + (d \cdot x)/2) \cdot (5 \cdot A - 11 \cdot B)) / (d \cdot (3 \cdot a^3 \cdot \tan(c/2 + (d \cdot x)/2)^2 + 3 \cdot a^3 \cdot \tan(c/2 + (d \cdot x)/2)^4 + a^3 \cdot \tan(c/2 + (d \cdot x)/2)^6 + a^3)) + (\tan(c/2 + (d \cdot x)/2)^3 \cdot ((A - B) / (3 \cdot a^3) + (4 \cdot A - 6 \cdot B) / (12 \cdot a^3))) / d - (\tan(c/2 + (d \cdot x)/2)^5 \cdot (A - B)) / (20 \cdot a^3 \cdot d)$

3.57 $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.57.1	Optimal result	679
3.57.2	Mathematica [B] (verified)	680
3.57.3	Rubi [A] (verified)	680
3.57.4	Maple [A] (verified)	683
3.57.5	Fricas [A] (verification not implemented)	683
3.57.6	Sympy [B] (verification not implemented)	684
3.57.7	Maxima [A] (verification not implemented)	685
3.57.8	Giac [A] (verification not implemented)	685
3.57.9	Mupad [B] (verification not implemented)	686

3.57.1 Optimal result

Integrand size = 31, antiderivative size = 193

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = -\frac{(6A-13B)x}{2a^3} + \frac{8(9A-19B) \sin(c+dx)}{15a^3d} - \frac{(6A-13B) \cos(c+dx) \sin(c+dx)}{2a^3d} + \frac{(A-B) \cos^4(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(6A-11B) \cos^3(c+dx) \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} + \frac{4(9A-19B) \cos^2(c+dx) \sin(c+dx)}{15d(a^3+a^3 \cos(c+dx))}$$

output

```
-1/2*(6*A-13*B)*x/a^3+8/15*(9*A-19*B)*sin(d*x+c)/a^3/d-1/2*(6*A-13*B)*cos(d*x+c)*sin(d*x+c)/a^3/d+1/5*(A-B)*cos(d*x+c)^4*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(6*A-11*B)*cos(d*x+c)^3*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+4/15*(9*A-19*B)*cos(d*x+c)^2*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))
```

3.57.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 435 vs. $2(193) = 386$.

Time = 2.17 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.25

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(-600(6A-13B)dx \cos\left(\frac{dx}{2}\right) - 600(6A-13B)dx \cos\left(c+\frac{dx}{2}\right) - 1800Adx \cos\left(c\right) \right)}{(480a^3d(1+\cos(c+dx)))^3}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(-600*(6*A - 13*B)*d*x*Cos[(d*x)/2] - 600*(6*A - 13*B)*d*x*Cos[c + (d*x)/2] - 1800*A*d*x*Cos[c + (3*d*x)/2] + 3900*B*d*x*Cos[c + (3*d*x)/2] - 1800*A*d*x*Cos[2*c + (3*d*x)/2] + 3900*B*d*x*Cos[2*c + (3*d*x)/2] - 360*A*d*x*Cos[2*c + (5*d*x)/2] + 780*B*d*x*Cos[2*c + (5*d*x)/2] - 360*A*d*x*Cos[3*c + (5*d*x)/2] + 780*B*d*x*Cos[3*c + (5*d*x)/2] + 7020*A*Sin[(d*x)/2] - 12760*B*Sin[(d*x)/2] - 4500*A*Sin[c + (d*x)/2] + 7560*B*Sin[c + (d*x)/2] + 4860*A*Sin[c + (3*d*x)/2] - 9230*B*Sin[c + (3*d*x)/2] - 900*A*Sin[2*c + (3*d*x)/2] + 930*B*Sin[2*c + (3*d*x)/2] + 1452*A*Sin[2*c + (5*d*x)/2] - 2782*B*Sin[2*c + (5*d*x)/2] + 300*A*Sin[3*c + (5*d*x)/2] - 750*B*Sin[3*c + (5*d*x)/2] + 60*A*Sin[3*c + (7*d*x)/2] - 105*B*Sin[3*c + (7*d*x)/2] + 60*A*Sin[4*c + (7*d*x)/2] - 105*B*Sin[4*c + (7*d*x)/2] + 15*B*Sin[4*c + (9*d*x)/2] + 15*B*Sin[5*c + (9*d*x)/2]))/(480*a^3*d*(1 + Cos[c + d*x])^3)`

3.57.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 3456, 3042, 3456, 3042, 3456, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

3.57. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^4 \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^3(c+dx)(4a(A-B)-a(2A-7B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^3(4a(A-B)-a(2A-7B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^2(c+dx)(3a^2(6A-11B)-a^2(18A-43B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{a(6A-11B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(3a^2(6A-11B)-a^2(18A-43B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{a(6A-11B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos(c+dx)(8a^3(9A-19B)-15a^3(6A-13B)\cos(c+dx))}{a^2} dx}{3a^2} + \frac{4a^2(9A-19B)\sin(c+dx)\cos^2(c+dx)}{d(a\cos(c+dx)+a)} + \frac{a(6A-11B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(8a^3(9A-19B)-15a^3(6A-13B)\sin(c+dx+\frac{\pi}{2}))}{a^2} dx}{3a^2} + \frac{4a^2(9A-19B)\sin(c+dx)\cos^2(c+dx)}{d(a\cos(c+dx)+a)} + \frac{a(6A-11B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad \downarrow \text{3213}
\end{aligned}$$

3.57. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{\frac{4a^2(9A-19B)\sin(c+dx)\cos^2(c+dx)}{d(a\cos(c+dx)+a)} + \frac{\frac{8a^3(9A-19B)\sin(c+dx)}{d} - \frac{15a^3(6A-13B)\sin(c+dx)\cos(c+dx)}{2d} - \frac{15}{2}a^3x(6A-13B)}{3a^2} + \frac{a(6A-11B)\sin(c+dx)\cos^3(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^4(c+dx)$$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^4*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((a*(6*A - 11*B)*Cos[c + d*x]^3*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((4*a^2*(9*A - 19*B)*Cos[c + d*x]^2*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])) + ((-15*a^3*(6*A - 13*B)*x)/2 + (8*a^3*(9*A - 19*B)*Sin[c + d*x])/d - (15*a^3*(6*A - 13*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/a^2)/(3*a^2)/(5*a^2)`

3.57.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; Free Q[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; Free Q[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.57.4 Maple [A] (verified)

Time = 0.98 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.55

method	result
parallelrisch	$\frac{2916 \left(\left(\frac{26A}{81} - \frac{464B}{729} \right) \cos(2dx+2c) + \left(\frac{5A}{243} - \frac{5B}{162} \right) \cos(3dx+3c) + \frac{5B \cos(4dx+4c)}{972} + \left(A - \frac{1001B}{486} \right) \cos(dx+c) + \frac{58A}{81} - \frac{4303B}{2916} \right)}{960a^3d}$
derivativedivides	$\frac{A \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) B - 2 \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) A + \frac{8 \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) B}{3} + 17A \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 31B \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 10}{4da^3}$
default	$\frac{A \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) B - 2 \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) A + \frac{8 \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) B}{3} + 17A \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 31B \tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 10}{4da^3}$
risch	$-\frac{3Ax}{a^3} + \frac{13Bx}{2a^3} - \frac{ie^{2i(dx+c)}B}{8a^3d} - \frac{ie^{i(dx+c)}A}{2a^3d} + \frac{3ie^{i(dx+c)}B}{2a^3d} + \frac{ie^{-i(dx+c)}A}{2a^3d} - \frac{3ie^{-i(dx+c)}B}{2a^3d} + \frac{ie^{-2i(dx+c)}B}{8a^3d}$
norman	$-\frac{(6A-13B)x}{2a} + \frac{(A-B) \left(\tan^{15} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{20ad} - \frac{(3A-5B) \left(\tan^{13} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{12ad} - \frac{5(6A-13B)x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{2a} - \frac{5(6A-13B)x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{a}$

input `int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE)`

output `1/960*(2916*((26/81*A-464/729*B)*cos(2*d*x+2*c)+(5/243*A-5/162*B)*cos(3*d*x+3*c)+5/972*B*cos(4*d*x+4*c)+(A-1001/486*B)*cos(d*x+c)+58/81*A-4303/2916*B)*tan(1/2*d*x+1/2*c)*sec(1/2*d*x+1/2*c)^4-2880*(A-13/6*B)*x*d)/a^3/d`

3.57.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.98

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{15(6A-13B)dx \cos(dx+c)^3 + 45(6A-13B)dx \cos(dx+c)^2 + 45(6A-13B)dx \cos(dx+c) + 15(6A-13B)dx}{30}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fracas")`

output
$$\frac{-1/30*(15*(6*A - 13*B)*d*x*cos(d*x + c)^3 + 45*(6*A - 13*B)*d*x*cos(d*x + c)^2 + 45*(6*A - 13*B)*d*x*cos(d*x + c) + 15*(6*A - 13*B)*d*x - (15*B*cos(d*x + c)^4 + 15*(2*A - 3*B)*cos(d*x + c)^3 + (234*A - 479*B)*cos(d*x + c)^2 + 3*(114*A - 239*B)*cos(d*x + c) + 144*A - 304*B)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)}$$

3.57.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 966 vs. $2(178) = 356$.

Time = 3.90 (sec) , antiderivative size = 966, normalized size of antiderivative = 5.01

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)`

output
$$\text{Piecewise}\left(\frac{-180*A*d*x*\tan(c/2 + d*x/2)**4}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{360*A*d*x*\tan(c/2 + d*x/2)**2}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{180*A*d*x}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{3*A*\tan(c/2 + d*x/2)**9}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{24*A*\tan(c/2 + d*x/2)**7}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{198*A*\tan(c/2 + d*x/2)**5}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{600*A*\tan(c/2 + d*x/2)**3}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{375*A*\tan(c/2 + d*x/2)}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{390*B*d*x*\tan(c/2 + d*x/2)**4}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{780*B*d*x*\tan(c/2 + d*x/2)**2}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{390*B*d*x}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{3*B*\tan(c/2 + d*x/2)**9}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} + \frac{34*B*\tan(c/2 + d*x/2)**7}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{388*B*\tan(c/2 + d*x/2)**5}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{1310*B*\tan(c/2 + d*x/2)**3}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{1310*B*\tan(c/2 + d*x/2)}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)} - \frac{1310*B}{(60*a**3*d*\tan(c/2 + d*x/2)**4 + 120*a**3*d*\tan(c/2 + d*x/2)**2 + 60*a**3*d)}\right)$$

3.57.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.67

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{B \left(\frac{60 \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} + \frac{7 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^3 + \frac{2a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} + \frac{465 \sin(dx+c)}{\cos(dx+c)+1} - \frac{40 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{780 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^3} \right)}{60d} - 3A \left(\frac{1}{(a+a\cos(c+dx))^3} \right)$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `-1/60*(B*(60*(5*sin(d*x + c)/(cos(d*x + c) + 1) + 7*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/(a^3 + 2*a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a^3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4) + (465*sin(d*x + c)/(cos(d*x + c) + 1) - 40*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 780*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3) - 3*A*(40*sin(d*x + c)/((a^3 + a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) + (85*sin(d*x + c)/(cos(d*x + c) + 1) - 10*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 120*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3))/d`

3.57.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.04

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{30(dx+c)(6A-13B)}{a^3} - \frac{60 \left(2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 7B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 5B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) \right)}{\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1\right)^2 a^3} - \frac{3Aa^{12} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{60d}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="giac")`

output
$$\frac{-1/60*(30*(d*x + c)*(6*A - 13*B)/a^3 - 60*(2*A*\tan(1/2*d*x + 1/2*c)^3 - 7*B*\tan(1/2*d*x + 1/2*c)^3 + 2*A*\tan(1/2*d*x + 1/2*c) - 5*B*\tan(1/2*d*x + 1/2*c)))/((\tan(1/2*d*x + 1/2*c)^2 + 1)^2*a^3) - (3*A*a^{12}*\tan(1/2*d*x + 1/2*c)^5 - 3*B*a^{12}*\tan(1/2*d*x + 1/2*c)^5 - 30*A*a^{12}*\tan(1/2*d*x + 1/2*c)^3 + 40*B*a^{12}*\tan(1/2*d*x + 1/2*c)^3 + 255*A*a^{12}*\tan(1/2*d*x + 1/2*c) - 465*B*a^{12}*\tan(1/2*d*x + 1/2*c))/a^{15}}{d}$$

3.57.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.05

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3(A-B)}{2a^3} + \frac{3(3A-5B)}{4a^3} + \frac{2A-10B}{4a^3}\right)}{d} - \frac{x(6A - 13B)}{2a^3}$$

$$+ \frac{(2A - 7B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (2A - 5B) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 2a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^3\right)}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{4a^3} + \frac{3A-5B}{12a^3}\right)}{d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A - B)}{20a^3 d}$$

input `int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output
$$\frac{(\tan(c/2 + (d*x)/2)*((3*(A - B))/(2*a^3) + (3*(3*A - 5*B))/(4*a^3) + (2*A - 10*B)/(4*a^3)))/d - (x*(6*A - 13*B))/(2*a^3) + (\tan(c/2 + (d*x)/2)^3*(2*A - 7*B) + \tan(c/2 + (d*x)/2)*(2*A - 5*B))/(d*(2*a^3*\tan(c/2 + (d*x)/2)^2 + a^3*\tan(c/2 + (d*x)/2)^4 + a^3)) - (\tan(c/2 + (d*x)/2)^3*((A - B)/(4*a^3) + (3*A - 5*B)/(12*a^3)))/d + (\tan(c/2 + (d*x)/2)^5*(A - B))/(20*a^3*d)}$$

3.58 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.58.1 Optimal result 687
 3.58.2 Mathematica [B] (verified) 687
 3.58.3 Rubi [A] (verified) 688
 3.58.4 Maple [A] (verified) 692
 3.58.5 Fricas [A] (verification not implemented) 692
 3.58.6 Sympy [B] (verification not implemented) 693
 3.58.7 Maxima [A] (verification not implemented) 694
 3.58.8 Giac [A] (verification not implemented) 694
 3.58.9 Mupad [B] (verification not implemented) 695

3.58.1 Optimal result

Integrand size = 31, antiderivative size = 147

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{(A-3B)x}{a^3} - \frac{(7A-27B) \sin(c+dx)}{15a^3d} + \frac{(A-B) \cos^3(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(4A-9B) \cos^2(c+dx) \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} - \frac{(A-3B) \sin(c+dx)}{d(a^3+a^3 \cos(c+dx))}$$

output `(A-3*B)*x/a^3-1/15*(7*A-27*B)*sin(d*x+c)/a^3/d+1/5*(A-B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(4*A-9*B)*cos(d*x+c)^2*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2-(A-3*B)*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))`

3.58.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 361 vs. 2(147) = 294.

Time = 1.86 (sec) , antiderivative size = 361, normalized size of antiderivative = 2.46

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{\cos(\frac{1}{2}(c+dx)) \sec(\frac{c}{2}) (300(A-3B)dx \cos(\frac{dx}{2}) + 300(A-3B)dx \cos(c+\frac{dx}{2}) + 150Adx \cos(c+\frac{3dx}{2}))}{(a+a \cos(c+dx))^3}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(300*(A - 3*B)*d*x*Cos[(d*x)/2] + 300*(A - 3*B)*d*x*Cos[c + (d*x)/2] + 150*A*d*x*Cos[c + (3*d*x)/2] - 450*B*d*x*Cos[c + (3*d*x)/2] + 150*A*d*x*Cos[2*c + (3*d*x)/2] - 450*B*d*x*Cos[2*c + (3*d*x)/2] + 30*A*d*x*Cos[2*c + (5*d*x)/2] - 90*B*d*x*Cos[2*c + (5*d*x)/2] + 30*A*d*x*Cos[3*c + (5*d*x)/2] - 90*B*d*x*Cos[3*c + (5*d*x)/2] - 740*A*Sin[(d*x)/2] + 1755*B*Sin[(d*x)/2] + 540*A*Sin[c + (d*x)/2] - 1125*B*Sin[c + (d*x)/2] - 460*A*Sin[c + (3*d*x)/2] + 1215*B*Sin[c + (3*d*x)/2] + 180*A*Sin[2*c + (3*d*x)/2] - 225*B*Sin[2*c + (3*d*x)/2] - 128*A*Sin[2*c + (5*d*x)/2] + 363*B*Sin[2*c + (5*d*x)/2] + 75*B*Sin[3*c + (5*d*x)/2] + 15*B*Sin[3*c + (7*d*x)/2] + 15*B*Sin[4*c + (7*d*x)/2]))/(120*a^3*d*(1 + Cos[c + d*x])^3)`

3.58.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3456, 3042, 3456, 3042, 3447, 3042, 3502, 27, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx$$

↓ 3456

$$\frac{\int \frac{\cos^2(c+dx)(3a(A-B)-a(A-6B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(3a(A-B)-a(A-6B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3456

3.58. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{\cos(c+dx)(2a^2(4A-9B)-a^2(7A-27B)\cos(c+dx))}{3a^2} dx + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}}{5a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(2a^2(4A-9B)-a^2(7A-27B)\sin(c+dx+\frac{\pi}{2}))}{3a^2} dx + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}}{5a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3447} \\
& \frac{\int \frac{2a^2(4A-9B)\cos(c+dx)-a^2(7A-27B)\cos^2(c+dx)}{3a^2} dx + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}}{5a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2a^2(4A-9B)\sin(c+dx+\frac{\pi}{2})-a^2(7A-27B)\sin(c+dx+\frac{\pi}{2})^2}{3a^2} dx + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}}{5a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3502} \\
& \frac{\int \frac{15a^3(A-3B)\cos(c+dx)}{\cos(c+dx)a+a} dx - \frac{a(7A-27B)\sin(c+dx)}{d} + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}}{3a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{15a^2(A-3B)\int \frac{\cos(c+dx)}{\cos(c+dx)a+a} dx - \frac{a(7A-27B)\sin(c+dx)}{d} + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2}}{3a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.58. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \frac{15a^2(A-3B) \int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})a+a} dx - \frac{a(7A-27B)\sin(c+dx)}{d}}{3a^2} + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \frac{5a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \frac{5d(a\cos(c+dx)+a)^3}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow \text{3214} \\
& \frac{15a^2(A-3B)\left(\frac{x}{a} - \int \frac{1}{\cos(c+dx)a+a} dx\right) - \frac{a(7A-27B)\sin(c+dx)}{d}}{3a^2} + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \frac{5a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \frac{5d(a\cos(c+dx)+a)^3}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow \text{3042} \\
& \frac{15a^2(A-3B)\left(\frac{x}{a} - \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx\right) - \frac{a(7A-27B)\sin(c+dx)}{d}}{3a^2} + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \frac{5a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \frac{5d(a\cos(c+dx)+a)^3}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow \text{3127} \\
& \frac{15a^2(A-3B)\left(\frac{x}{a} - \frac{\sin(c+dx)}{d(a\cos(c+dx)+a)}\right) - \frac{a(7A-27B)\sin(c+dx)}{d}}{3a^2} + \frac{a(4A-9B)\sin(c+dx)\cos^2(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \frac{5a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \frac{5d(a\cos(c+dx)+a)^3}{5d(a\cos(c+dx)+a)^3}
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^3*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((a*(4*A - 9*B)*Cos[c + d*x]^2*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (-((a*(7*A - 27*B)*Sin[c + d*x])/d) + 15*a^2*(A - 3*B)*(x/a - Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))) / (3*a^2) / (5*a^2)`

3.58. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

3.58.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/(c_ + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_ + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_ + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)^2], x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.58.
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.58.4 Maple [A] (verified)

Time = 1.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.61

method	result
parallelrisch	$\frac{-204 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\left(\frac{16A}{51} - \frac{39B}{34} \right) \cos(2dx+2c) - \frac{5B \cos(3dx+3c)}{68} + \left(A - \frac{243B}{68} \right) \cos(dx+c) + \frac{38A}{51} - \frac{87B}{34} \right) \left(\sec^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \dots}{240a^3d}$
derivativedivides	$\frac{-\frac{A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{5} + \frac{4 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{3} - 2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B - 7A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 17B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots}{4da^3}$
default	$\frac{-\frac{A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{5} + \frac{4 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{3} - 2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B - 7A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 17B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots}{4da^3}$
risch	$\frac{Ax}{a^3} - \frac{3Bx}{a^3} - \frac{ie^{i(dx+c)}B}{2a^3d} + \frac{ie^{-i(dx+c)}B}{2a^3d} - \frac{2i(45Ae^{4i(dx+c)} - 90Be^{4i(dx+c)} + 135Ae^{3i(dx+c)} - 300Be^{3i(dx+c)} + 15d a^3)}{15d a^3}$
norman	$\frac{(A-3B)x}{a} + \frac{(A-3B)x \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a} + \frac{4(A-3B)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a} + \frac{6(A-3B)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a} + \frac{4(A-3B)x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a} - \dots$

input `int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE)`

output `1/240*(-204*tan(1/2*d*x+1/2*c)*((16/51*A-39/34*B)*cos(2*d*x+2*c)-5/68*B*cos(3*d*x+3*c)+(A-243/68*B)*cos(d*x+c)+38/51*A-87/34*B)*sec(1/2*d*x+1/2*c)^4+240*d*x*(A-3*B))/a^3/d`

3.58.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.12

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

$$= \frac{15(A-3B)dx \cos(dx+c)^3 + 45(A-3B)dx \cos(dx+c)^2 + 45(A-3B)dx \cos(dx+c) + 15(A-3B)dx}{15(a^3d \cos(dx+c)^3 + 3a^3d)}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fricas")`

```
output 1/15*(15*(A - 3*B)*d*x*cos(d*x + c)^3 + 45*(A - 3*B)*d*x*cos(d*x + c)^2 +
45*(A - 3*B)*d*x*cos(d*x + c) + 15*(A - 3*B)*d*x + (15*B*cos(d*x + c)^3 -
(32*A - 117*B)*cos(d*x + c)^2 - 3*(17*A - 57*B)*cos(d*x + c) - 22*A + 72*B
)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*c
os(d*x + c) + a^3*d)
```

3.58.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 496 vs. $2(134) = 268$.

Time = 2.37 (sec) , antiderivative size = 496, normalized size of antiderivative = 3.37

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \begin{cases} \frac{60Adx \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{60a^3d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 60a^3d} + \frac{60Adx}{60a^3d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 60a^3d} - \frac{3A \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{60a^3d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 60a^3d} + \frac{17A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{60a^3d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 60a^3d} - \frac{85A}{60a^3d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 60a^3d} \\ \frac{x(A+B\cos(c))\cos^3(c)}{(a\cos(c)+a)^3} \end{cases}$$

```
input integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)
```

```
output Piecewise((60*A*d*x*tan(c/2 + d*x/2)**2/(60*a**3*d*tan(c/2 + d*x/2)**2 + 6
0*a**3*d) + 60*A*d*x/(60*a**3*d*tan(c/2 + d*x/2)**2 + 60*a**3*d) - 3*A*tan
(c/2 + d*x/2)**7/(60*a**3*d*tan(c/2 + d*x/2)**2 + 60*a**3*d) + 17*A*tan(c/
2 + d*x/2)**5/(60*a**3*d*tan(c/2 + d*x/2)**2 + 60*a**3*d) - 85*A*tan(c/2 +
d*x/2)**3/(60*a**3*d*tan(c/2 + d*x/2)**2 + 60*a**3*d) - 105*A*tan(c/2 + d
*x/2)/(60*a**3*d*tan(c/2 + d*x/2)**2 + 60*a**3*d) - 180*B*d*x*tan(c/2 + d*
x/2)**2/(60*a**3*d*tan(c/2 + d*x/2)**2 + 60*a**3*d) - 180*B*d*x/(60*a**3*d
*tan(c/2 + d*x/2)**2 + 60*a**3*d) + 3*B*tan(c/2 + d*x/2)**7/(60*a**3*d*tan
(c/2 + d*x/2)**2 + 60*a**3*d) - 27*B*tan(c/2 + d*x/2)**5/(60*a**3*d*tan(c/
2 + d*x/2)**2 + 60*a**3*d) + 225*B*tan(c/2 + d*x/2)**3/(60*a**3*d*tan(c/2
+ d*x/2)**2 + 60*a**3*d) + 375*B*tan(c/2 + d*x/2)/(60*a**3*d*tan(c/2 + d*x
/2)**2 + 60*a**3*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**3/(a*cos(c) + a)
**3, True))
```

3.58.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.57

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{3B \left(\frac{40 \sin(dx+c)}{\left(a^3 + \frac{a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2}\right) (\cos(dx+c)+1)} + \frac{85 \sin(dx+c)}{\cos(dx+c)+1} - \frac{10 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{\sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{120 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^3} \right) - A \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} \right)}{60d}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")
```

```
output 1/60*(3*B*(40*sin(d*x + c)/((a^3 + a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) + (85*sin(d*x + c)/(cos(d*x + c) + 1) - 10*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 120*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3) - A*((105*sin(d*x + c)/(cos(d*x + c) + 1) - 20*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 120*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3)/d
```

3.58.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.05

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{60(dx+c)(A-3B)}{a^3} + \frac{120B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1\right)a^3} - \frac{3Aa^{12} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 3Ba^{12} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 20Aa^{12} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 30Ba^{12} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{a^{15}}}{60d}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="giac")
```

```
output 1/60*(60*(d*x + c)*(A - 3*B)/a^3 + 120*B*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 + 1)*a^3) - (3*A*a^12*tan(1/2*d*x + 1/2*c)^5 - 3*B*a^12*tan(1/2*d*x + 1/2*c)^5 - 20*A*a^12*tan(1/2*d*x + 1/2*c)^3 + 30*B*a^12*tan(1/2*d*x + 1/2*c)^3 + 105*A*a^12*tan(1/2*d*x + 1/2*c) - 255*B*a^12*tan(1/2*d*x + 1/2*c))/a^15)/d
```

3.58. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

3.58.9 Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.03

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{6a^3} + \frac{2A-4B}{12a^3}\right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3(A-B)}{4a^3} - \frac{3B}{2a^3} + \frac{2A-4B}{2a^3}\right)}{d} + \frac{x(A-3B)}{a^3} + \frac{2B\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d\left(a^3\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^3\right)} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5(A-B)}{20a^3d}$$

input `int((cos(c + d*x))^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`output `(tan(c/2 + (d*x)/2)^3*((A - B)/(6*a^3) + (2*A - 4*B)/(12*a^3)))/d - (tan(c/2 + (d*x)/2)*((3*(A - B))/(4*a^3) - (3*B)/(2*a^3) + (2*A - 4*B)/(2*a^3)))/d + (x*(A - 3*B))/a^3 + (2*B*tan(c/2 + (d*x)/2))/(d*(a^3*tan(c/2 + (d*x)/2)^2 + a^3)) - (tan(c/2 + (d*x)/2)^5*(A - B))/(20*a^3*d)`

3.59 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.59.1	Optimal result	696
3.59.2	Mathematica [B] (verified)	696
3.59.3	Rubi [A] (verified)	697
3.59.4	Maple [A] (verified)	700
3.59.5	Fricas [A] (verification not implemented)	700
3.59.6	Sympy [A] (verification not implemented)	701
3.59.7	Maxima [A] (verification not implemented)	701
3.59.8	Giac [A] (verification not implemented)	702
3.59.9	Mupad [B] (verification not implemented)	702

3.59.1 Optimal result

Integrand size = 31, antiderivative size = 116

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{Bx}{a^3} + \frac{(A-B) \cos^2(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} - \frac{(2A-7B) \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} + \frac{(4A-29B) \sin(c+dx)}{15d(a^3+a^3 \cos(c+dx))}$$

```
output B*x/a^3+1/5*(A-B)*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^3-1/15*(2*A-7
*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+1/15*(4*A-29*B)*sin(d*x+c)/d/(a^3+a^
3*cos(d*x+c))
```

3.59.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 241 vs. 2(116) = 232.

Time = 1.21 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.08

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{\sec\left(\frac{c}{2}\right) \sec^5\left(\frac{1}{2}(c+dx)\right) \left(150Bdx \cos\left(\frac{dx}{2}\right) + 150Bdx \cos\left(c+\frac{dx}{2}\right) + 75Bdx \cos\left(c+\frac{3dx}{2}\right) + 75Bdx \cos\left(2c+dx\right)\right)}{\dots}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*cos[c + d*x])^3,x]`

output `(Sec[c/2]*Sec[(c + d*x)/2]^5*(150*B*d*x*cos[(d*x)/2] + 150*B*d*x*cos[c + (d*x)/2] + 75*B*d*x*cos[c + (3*d*x)/2] + 75*B*d*x*cos[2*c + (3*d*x)/2] + 15*B*d*x*cos[2*c + (5*d*x)/2] + 15*B*d*x*cos[3*c + (5*d*x)/2] + 80*A*sin[(d*x)/2] - 370*B*sin[(d*x)/2] - 60*A*sin[c + (d*x)/2] + 270*B*sin[c + (d*x)/2] + 40*A*sin[c + (3*d*x)/2] - 230*B*sin[c + (3*d*x)/2] - 30*A*sin[2*c + (3*d*x)/2] + 90*B*sin[2*c + (3*d*x)/2] + 14*A*sin[2*c + (5*d*x)/2] - 64*B*sin[2*c + (5*d*x)/2]))/(480*a^3*d)`

3.59.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3456, 3042, 3447, 3042, 3498, 25, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})^2(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx \\
 & \quad \downarrow \text{3456} \\
 & \frac{\int \frac{\cos(c+dx)(2a(A-B)+5aB\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(2a(A-B)+5aB\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
 & \quad \downarrow \text{3447} \\
 & \frac{\int \frac{5aB\cos^2(c+dx)+2a(A-B)\cos(c+dx)}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.59. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{5aB \sin(c+dx+\frac{\pi}{2})^2 + 2a(A-B) \sin(c+dx+\frac{\pi}{2})}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3498} \\
& \frac{-\int \frac{2(2A-7B)a^2 + 15B \cos(c+dx)a^2}{3a^2} dx - \frac{a(2A-7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2}}{5a^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{2(2A-7B)a^2 + 15B \cos(c+dx)a^2}{3a^2} dx - \frac{a(2A-7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2}}{5a^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2(2A-7B)a^2 + 15B \sin(c+dx+\frac{\pi}{2})a^2}{3a^2} dx - \frac{a(2A-7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2}}{5a^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3214} \\
& \frac{a^2(4A-29B) \int \frac{1}{\cos(c+dx)a+a} dx + 15aBx}{5a^2} - \frac{a(2A-7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{a^2(4A-29B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx + 15aBx}{5a^2} - \frac{a(2A-7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3127} \\
& \frac{\frac{a^2(4A-29B) \sin(c+dx)}{d(a \cos(c+dx)+a)} + 15aBx}{5a^2} - \frac{a(2A-7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx) + a)^3}
\end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^2*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + (-1/3*(a*(2*A - 7*B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + (15*a*B*x + (a^2*(4*A - 29*B)*Sin[c + d*x]))/(d*(a + a*Cos[c + d*x])))/(3*a^2)/(5*a^2)`

3.59. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.59.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3498 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b - a*B + b*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m/(a*f*(2*m + 1)), x] + Simp[1/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]`

3.59.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.59

method	result
parallelrisch	$\frac{3(A-B)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+10(2B-A)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+15(A-7B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+60dxB}{60a^3d}$
derivativedivides	$\frac{\frac{A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}-\frac{\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{5}-\frac{2\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{3}+\frac{4\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{3}+A\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-7B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+8B\arcsin\left(\frac{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2}\right)}{4da^3}$
default	$\frac{\frac{A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}-\frac{\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{5}-\frac{2\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{3}+\frac{4\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{3}+A\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-7B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+8B\arcsin\left(\frac{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2}\right)}{4da^3}$
risch	$\frac{Bx}{a^3}+\frac{2i(15Ae^{4i(dx+c)}-45Be^{4i(dx+c)}+30Ae^{3i(dx+c)}-135Be^{3i(dx+c)}+40Ae^{2i(dx+c)}-185Be^{2i(dx+c)}+20Ae^{i(dx+c)}-15d)}{15da^3(e^{i(dx+c)}+1)^5}$
norman	$\frac{\frac{Bx}{a}+\frac{Bx\left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{3Bx\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{3Bx\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}-\frac{(A-11B)\left(\tan^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{60ad}+\frac{(A-7B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{4ad}+\frac{(A-11B)\arcsin\left(\frac{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2}\right)}{4ad}}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^3a^2}$

input `int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE)`

output `1/60*(3*(A-B)*tan(1/2*d*x+1/2*c)^5+10*(2*B-A)*tan(1/2*d*x+1/2*c)^3+15*(A-7*B)*tan(1/2*d*x+1/2*c)+60*d*x*B)/a^3/d`

3.59.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.18

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{15Bdx\cos(dx+c)^3+45Bdx\cos(dx+c)^2+45Bdx\cos(dx+c)+15Bdx+((7A-32B)\cos(dx+c)+15(a^3d\cos(dx+c)^3+3a^3d\cos(dx+c)^2+3a^3d\cos(dx+c))}{15(a^3d\cos(dx+c)^3+3a^3d\cos(dx+c)^2+3a^3d\cos(dx+c))}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fricas")`

output `1/15*(15*B*d*x*cos(d*x+c)^3+45*B*d*x*cos(d*x+c)^2+45*B*d*x*cos(d*x+c)+15*B*d*x+((7*A-32*B)*cos(d*x+c)^2+3*(2*A-17*B)*cos(d*x+c)+2*A-22*B)*sin(d*x+c))/(a^3*d*cos(d*x+c)^3+3*a^3*d*cos(d*x+c)^2+3*a^3*d*cos(d*x+c)+a^3*d)`

3.59.
$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

3.59.6 Sympy [A] (verification not implemented)

Time = 1.34 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.28

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \begin{cases} \frac{A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{20a^3d} - \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^3d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4a^3d} + \frac{Bx}{a^3} - \frac{B \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{20a^3d} + \frac{B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{3a^3d} - \frac{7B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4a^3d} \\ \frac{x(A+B\cos(c))\cos^2(c)}{(a\cos(c)+a)^3} \end{cases} \quad \text{for } d \neq 0$$

otherwise

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)`

output `Piecewise((A*tan(c/2 + d*x/2)**5/(20*a**3*d) - A*tan(c/2 + d*x/2)**3/(6*a**3*d) + A*tan(c/2 + d*x/2)/(4*a**3*d) + B*x/a**3 - B*tan(c/2 + d*x/2)**5/(20*a**3*d) + B*tan(c/2 + d*x/2)**3/(3*a**3*d) - 7*B*tan(c/2 + d*x/2)/(4*a**3*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**2/(a*cos(c) + a)**3, True))`

3.59.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.38

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{B \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} - \frac{20 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) - \frac{120 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^3} - \frac{A \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} - \frac{10 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a^3}}{60d}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `-1/60*(B*((105*sin(d*x + c)/(cos(d*x + c) + 1) - 20*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 120*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^3 - A*(15*sin(d*x + c)/(cos(d*x + c) + 1) - 10*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3)/d`

3.59.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.03

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{60(dx+c)B}{a^3} + \frac{3Aa^{12}\tan(\frac{1}{2}dx+\frac{1}{2}c)^5 - 3Ba^{12}\tan(\frac{1}{2}dx+\frac{1}{2}c)^5 - 10Aa^{12}\tan(\frac{1}{2}dx+\frac{1}{2}c)^3 + 20Ba^{12}\tan(\frac{1}{2}dx+\frac{1}{2}c)^3 + 15Aa^{12}\tan(\frac{1}{2}dx+\frac{1}{2}c)}{a^{15}}$$

$$60d$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="giac")`

output `1/60*(60*(d*x + c)*B/a^3 + (3*A*a^12*tan(1/2*d*x + 1/2*c)^5 - 3*B*a^12*tan(1/2*d*x + 1/2*c)^5 - 10*A*a^12*tan(1/2*d*x + 1/2*c)^3 + 20*B*a^12*tan(1/2*d*x + 1/2*c)^3 + 15*A*a^12*tan(1/2*d*x + 1/2*c) - 105*B*a^12*tan(1/2*d*x + 1/2*c))/a^15)/d`

3.59.9 Mupad [B] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.16

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \frac{Bx}{a^3}$$

$$\frac{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \left(\frac{A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{6} - \frac{B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{3} \right) - \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^4 \left(\frac{A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{7B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} \right) - \frac{A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{20}}{a^3 d \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^5}$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output `(B*x)/a^3 - (cos(c/2 + (d*x)/2)^2*((A*sin(c/2 + (d*x)/2)^3)/6 - (B*sin(c/2 + (d*x)/2)^3)/3) - cos(c/2 + (d*x)/2)^4*((A*sin(c/2 + (d*x)/2))/4 - (7*B*sin(c/2 + (d*x)/2))/4) - (A*sin(c/2 + (d*x)/2)^5)/20 + (B*sin(c/2 + (d*x)/2)^5)/20)/(a^3*d*cos(c/2 + (d*x)/2)^5)`

3.60 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.60.1 Optimal result 703
 3.60.2 Mathematica [A] (verified) 703
 3.60.3 Rubi [A] (verified) 704
 3.60.4 Maple [A] (verified) 706
 3.60.5 Fricas [A] (verification not implemented) 707
 3.60.6 Sympy [A] (verification not implemented) 707
 3.60.7 Maxima [A] (verification not implemented) 708
 3.60.8 Giac [A] (verification not implemented) 708
 3.60.9 Mupad [B] (verification not implemented) 709

3.60.1 Optimal result

Integrand size = 29, antiderivative size = 102

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = -\frac{(A - B) \sin(c + dx)}{5d(a + a \cos(c + dx))^3} + \frac{(3A - 8B) \sin(c + dx)}{15ad(a + a \cos(c + dx))^2} + \frac{(3A + 7B) \sin(c + dx)}{15d(a^3 + a^3 \cos(c + dx))}$$

output `-1/5*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(3*A-8*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+1/15*(3*A+7*B)*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))`

3.60.2 Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.32

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \frac{\cos\left(\frac{1}{2}(c + dx)\right) \sec\left(\frac{c}{2}\right) (5(3A + 8B) \sin\left(\frac{dx}{2}\right) - 15(A + 2B) \sin\left(c + \frac{dx}{2}\right) + 15A \sin\left(c + \frac{3dx}{2}\right) + 20B \sin(c + dx))}{30a^3d(1 + \cos(c + dx))^3}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output $(\text{Cos}[(c + d*x)/2]*\text{Sec}[c/2]*(5*(3*A + 8*B)*\text{Sin}[(d*x)/2] - 15*(A + 2*B)*\text{Sin}[c + (d*x)/2] + 15*A*\text{Sin}[c + (3*d*x)/2] + 20*B*\text{Sin}[c + (3*d*x)/2] - 15*B*\text{Sin}[2*c + (3*d*x)/2] + 3*A*\text{Sin}[2*c + (5*d*x)/2] + 7*B*\text{Sin}[2*c + (5*d*x)/2]))/(30*a^3*d*(1 + \text{Cos}[c + d*x])^3)$

3.60.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.02, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3447, 3042, 3498, 25, 3042, 3229, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a\cos(c+dx)+a)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx \\ & \quad \downarrow \text{3498} \\ & -\frac{\int -\frac{3a(A-B)+5aB\cos(c+dx)}{(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} \\ & \quad \downarrow \text{25} \\ & \frac{\int \frac{3a(A-B)+5aB\cos(c+dx)}{(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{3a(A-B)+5aB\sin(c+dx+\frac{\pi}{2})}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} \end{aligned}$$

3.60. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{array}{c}
 \downarrow \text{3229} \\
 \frac{\frac{1}{3}(3A + 7B) \int \frac{1}{\cos(c+dx)a+a} dx + \frac{a(3A-8B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2}}{5a^2} - \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} \\
 \downarrow \text{3042} \\
 \frac{\frac{1}{3}(3A + 7B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx + \frac{a(3A-8B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2}}{5a^2} - \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} \\
 \downarrow \text{3127} \\
 \frac{\frac{(3A+7B) \sin(c+dx)}{3d(a \cos(c+dx)+a)} + \frac{a(3A-8B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2}}{5a^2} - \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3}
 \end{array}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `-1/5*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^3) + ((a*(3*A - 8*B)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((3*A + 7*B)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x]))) / (5*a^2)`

3.60.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*SIN[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3229 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*SIN[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`


```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3498 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b - a*
B + b*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[1
/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b
*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]
```

3.60.4 Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.55

method	result
parallelrisch	$-\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{10B \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} - 5A - 5B \right)}{20a^3d}$
derivativedivides	$\frac{(-A+B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} \right) - \frac{2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4da^3}$
default	$\frac{(-A+B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} \right) - \frac{2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4da^3}$
risch	$\frac{2i(15B e^{4i(dx+c)} + 15A e^{3i(dx+c)} + 30B e^{3i(dx+c)} + 15A e^{2i(dx+c)} + 40B e^{2i(dx+c)} + 15A e^{i(dx+c)} + 20B e^{i(dx+c)} + 3A + 7B)}{15da^3(e^{i(dx+c)} + 1)^5}$
norman	$-\frac{(A-B) \left(\frac{\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)}{20ad} \right) + \frac{(A+B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4ad} + \frac{(3A+2B) \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{6ad} \right) - (3A+2B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{30ad} \right) + \frac{(6A-B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{30ad} \right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^2 a^2}$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE
)
```

```
output -1/20*tan(1/2*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^4+10/3*B*tan(1/2*d*x+1/
2*c)^2-5*A-5*B)/a^3/d
```

3.60.
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.60.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.91

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{((3A+7B)\cos(dx+c)^2 + 3(3A+2B)\cos(dx+c) + 3A+2B)\sin(dx+c)}{15(a^3d\cos(dx+c)^3 + 3a^3d\cos(dx+c)^2 + 3a^3d\cos(dx+c) + a^3d)}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fricas")
```

```
output 1/15*((3*A + 7*B)*cos(d*x + c)^2 + 3*(3*A + 2*B)*cos(d*x + c) + 3*A + 2*B)*sin(d*x + c)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)
```

3.60.6 Sympy [A] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.15

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \begin{cases} -\frac{A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{20a^3d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4a^3d} + \frac{B \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{20a^3d} - \frac{B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^3d} + \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4a^3d} & \text{for } d \neq 0 \\ \frac{x(A+B\cos(c))\cos(c)}{(a\cos(c)+a)^3} & \text{otherwise} \end{cases}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)
```

```
output Piecewise((-A*tan(c/2 + d*x/2)**5/(20*a**3*d) + A*tan(c/2 + d*x/2)/(4*a**3*d) + B*tan(c/2 + d*x/2)**5/(20*a**3*d) - B*tan(c/2 + d*x/2)**3/(6*a**3*d) + B*tan(c/2 + d*x/2)/(4*a**3*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)/(a*cos(c) + a)**3, True))
```

3.60.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.13

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{B\left(\frac{15\sin(dx+c)}{\cos(dx+c)+1} - \frac{10\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3\sin(dx+c)^5}{(\cos(dx+c)+1)^5}\right)}{a^3} + \frac{3A\left(\frac{5\sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^5}{(\cos(dx+c)+1)^5}\right)}{a^3}$$

$$60d$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `1/60*(B*(15*sin(d*x + c)/(cos(d*x + c) + 1) - 10*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 + 3*A*(5*sin(d*x + c)/(cos(d*x + c) + 1) - sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3)/d`

3.60.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.74

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{3A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 3B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 + 10B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 15A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 15B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{60a^3d}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="giac")`

output `-1/60*(3*A*tan(1/2*d*x + 1/2*c)^5 - 3*B*tan(1/2*d*x + 1/2*c)^5 + 10*B*tan(1/2*d*x + 1/2*c)^3 - 15*A*tan(1/2*d*x + 1/2*c) - 15*B*tan(1/2*d*x + 1/2*c))/(a^3*d)`

3.60.9 Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.65

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(15A + 15B - 3A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 10B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 3B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4\right)}{60a^3d}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`output `(tan(c/2 + (d*x)/2)*(15*A + 15*B - 3*A*tan(c/2 + (d*x)/2)^4 - 10*B*tan(c/2 + (d*x)/2)^2 + 3*B*tan(c/2 + (d*x)/2)^4))/(60*a^3*d)`

3.61 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3} dx$

3.61.1	Optimal result	710
3.61.2	Mathematica [A] (verified)	710
3.61.3	Rubi [A] (verified)	711
3.61.4	Maple [A] (verified)	713
3.61.5	Fricas [A] (verification not implemented)	713
3.61.6	Sympy [A] (verification not implemented)	714
3.61.7	Maxima [A] (verification not implemented)	714
3.61.8	Giac [A] (verification not implemented)	715
3.61.9	Mupad [B] (verification not implemented)	715

3.61.1 Optimal result

Integrand size = 23, antiderivative size = 102

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{(A - B) \sin(c + dx)}{5d(a + a \cos(c + dx))^3} + \frac{(2A + 3B) \sin(c + dx)}{15ad(a + a \cos(c + dx))^2} + \frac{(2A + 3B) \sin(c + dx)}{15d(a^3 + a^3 \cos(c + dx))}$$

output `1/5*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(2*A+3*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+1/15*(2*A+3*B)*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))`

3.61.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.62

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{(7A + 3B + (6A + 9B) \cos(c + dx) + (2A + 3B) \cos^2(c + dx)) \sin(c + dx)}{15a^3d(1 + \cos(c + dx))^3}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^3,x]`

output `((7*A + 3*B + (6*A + 9*B)*Cos[c + d*x] + (2*A + 3*B)*Cos[c + d*x]^2)*Sin[c + d*x])/(15*a^3*d*(1 + Cos[c + d*x])^3)`

3.61.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3042, 3229, 3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a \cos(c + dx) + a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
 & \quad \downarrow \text{3229} \\
 & \frac{(2A + 3B) \int \frac{1}{(\cos(c+dx)a+a)^2} dx}{5a} + \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(2A + 3B) \int \frac{1}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a} + \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} \\
 & \quad \downarrow \text{3129} \\
 & \frac{(2A + 3B) \left(\int \frac{\frac{1}{\cos(c+dx)a+a} dx}{3a} + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right)}{5a} + \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(2A + 3B) \left(\int \frac{\frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a} + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right)}{5a} + \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} \\
 & \quad \downarrow \text{3127} \\
 & \frac{(A - B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} + \frac{(2A + 3B) \left(\frac{\sin(c+dx)}{3ad(a \cos(c+dx)+a)} + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right)}{5a}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^3,x]`

```
output ((A - B)*Sin[c + d*x])/(5*d*(a + a*cos[c + d*x])^3) + ((2*A + 3*B)*(Sin[c
+ d*x]/(3*d*(a + a*cos[c + d*x])^2) + Sin[c + d*x]/(3*a*d*(a + a*cos[c + d
*x])))))/(5*a)
```

3.61.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3127 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c +
d*x]/(d*(b + a*sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b
^2, 0]
```

```
rule 3129 Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*cos[c
+ d*x]*((a + b*sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n
+ 1)) Int[(a + b*sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x]
&& EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 3229 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*sin[e + f*
x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) I
nt[(a + b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &&
NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]
```

3.61.4 Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.55

method	result	size
parallelrisch	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{10A \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} + 5A + 5B \right)}{20a^3d}$	56
derivativedivides	$\frac{(A-B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} \right) + \frac{2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4da^3}$	64
default	$\frac{(A-B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} \right) + \frac{2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4da^3}$	64
risch	$\frac{2i(15B e^{3i(dx+c)} + 20A e^{2i(dx+c)} + 15B e^{i(dx+c)} + 10A e^{i(dx+c)} + 15B e^{i(dx+c)} + 2A + 3B)}{15da^3 (e^{i(dx+c)} + 1)^5}$	90
norman	$\frac{(A-B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{20ad} \right) + \frac{(A+B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4ad} + \frac{(5A+3B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{12ad} + \frac{(13A-3B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{60ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a^2}$	117

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c))*a^3,x,method=_RETURNVERBOSE)`

output `1/20*tan(1/2*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^4+10/3*A*tan(1/2*d*x+1/2*c)^2+5*A+5*B)/a^3/d`

3.61.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.91

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{((2A + 3B) \cos(dx + c))^2 + 3(2A + 3B) \cos(dx + c) + 7A + 3B) \sin(dx + c)}{15(a^3d \cos(dx + c))^3 + 3a^3d \cos(dx + c)^2 + 3a^3d \cos(dx + c) + a^3d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fricas")`

output `1/15*((2*A + 3*B)*cos(d*x + c)^2 + 3*(2*A + 3*B)*cos(d*x + c) + 7*A + 3*B)*sin(d*x + c)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.61.6 Sympy [A] (verification not implemented)

Time = 0.78 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.12

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \begin{cases} \frac{A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{20a^3d} + \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{6a^3d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4a^3d} - \frac{B \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{20a^3d} + \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4a^3d} & \text{for } d \neq 0 \\ \frac{x(A+B \cos(c))}{(a \cos(c)+a)^3} & \text{otherwise} \end{cases}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)`

output `Piecewise((A*tan(c/2 + d*x/2)**5/(20*a**3*d) + A*tan(c/2 + d*x/2)**3/(6*a**3*d) + A*tan(c/2 + d*x/2)/(4*a**3*d) - B*tan(c/2 + d*x/2)**5/(20*a**3*d) + B*tan(c/2 + d*x/2)/(4*a**3*d), Ne(d, 0)), (x*(A + B*cos(c))/(a*cos(c) + a)**3, True))`

3.61.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.13

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{A \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} + \frac{10 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a^3} + \frac{3B \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{a^3}$$

$$60d$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `1/60*(A*(15*sin(d*x + c)/(cos(d*x + c) + 1) + 10*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 + 3*B*(5*sin(d*x + c)/(cos(d*x + c) + 1) - sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3/d`

3.61.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.74

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{3 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 3 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 10 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 15 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 15 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{60 a^3 d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="giac")`output `1/60*(3*A*tan(1/2*d*x + 1/2*c)^5 - 3*B*tan(1/2*d*x + 1/2*c)^5 + 10*A*tan(1/2*d*x + 1/2*c)^3 + 15*A*tan(1/2*d*x + 1/2*c) + 15*B*tan(1/2*d*x + 1/2*c))/(a^3*d)`**3.61.9 Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.65

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(15 A + 15 B + 10 A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 3 A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 3 B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4\right)}{60 a^3 d}$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^3,x)`output `(tan(c/2 + (d*x)/2)*(15*A + 15*B + 10*A*tan(c/2 + (d*x)/2)^2 + 3*A*tan(c/2 + (d*x)/2)^4 - 3*B*tan(c/2 + (d*x)/2)^4)/(60*a^3*d)`

$$3.62 \quad \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^3} dx$$

3.62.1	Optimal result	716
3.62.2	Mathematica [A] (verified)	716
3.62.3	Rubi [A] (verified)	717
3.62.4	Maple [A] (verified)	719
3.62.5	Fricas [A] (verification not implemented)	720
3.62.6	Sympy [F]	720
3.62.7	Maxima [A] (verification not implemented)	721
3.62.8	Giac [A] (verification not implemented)	721
3.62.9	Mupad [B] (verification not implemented)	722

3.62.1 Optimal result

Integrand size = 29, antiderivative size = 117

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{A \operatorname{arctanh}(\sin(c + dx))}{a^3 d} - \frac{(A - B) \sin(c + dx)}{5d(a + a \cos(c + dx))^3} - \frac{(7A - 2B) \sin(c + dx)}{15ad(a + a \cos(c + dx))^2} - \frac{2(11A - B) \sin(c + dx)}{15d(a^3 + a^3 \cos(c + dx))}$$

output `A*arctanh(sin(d*x+c))/a^3/d-1/5*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^3-1/15*(7*A-2*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2-2/15*(11*A-B)*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))`

3.62.2 Mathematica [A] (verified)

Time = 0.94 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.68

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{-240A \cos^6\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^3,x]`

3.62. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^3} dx$

output $(-240*A*\text{Cos}[(c + d*x)/2]^6*(\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] - \text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]]) + \text{Cos}[(c + d*x)/2]*\text{Sec}[c/2]*(-5*(29*A - 4*B)*\text{Sin}[(d*x)/2] + 75*A*\text{Sin}[c + (d*x)/2] - 95*A*\text{Sin}[c + (3*d*x)/2] + 10*B*\text{Sin}[c + (3*d*x)/2] + 15*A*\text{Sin}[2*c + (3*d*x)/2] - 22*A*\text{Sin}[2*c + (5*d*x)/2] + 2*B*\text{Sin}[2*c + (5*d*x)/2]))/(30*a^3*d*(1 + \text{Cos}[c + d*x])^3)$

3.62.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3457, 3042, 3457, 3042, 3457, 27, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx$$

↓ 3457

$$\frac{\int \frac{(5aA-2a(A-B)\cos(c+dx))\sec(c+dx)}{(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{\int \frac{5aA-2a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3457

$$\frac{\int \frac{(15a^2A-a^2(7A-2B)\cos(c+dx))\sec(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} - \frac{a(7A-2B)\sin(c+dx)}{3d(a\cos(c+dx)+a)^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{\int \frac{15a^2A-a^2(7A-2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(\sin(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} - \frac{a(7A-2B)\sin(c+dx)}{3d(a\cos(c+dx)+a)^2} - \frac{(A-B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3457

3.62. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+a\cos(c+dx))^3} dx$

$$\frac{\frac{\int 15a^3 A \sec(c+dx) dx - \frac{2a^2(11A-B) \sin(c+dx)}{d(a \cos(c+dx)+a)}}{3a^2} - \frac{a(7A-2B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} - \frac{(A-B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2}$$

↓ 27

$$\frac{15aA \int \sec(c+dx) dx - \frac{2a^2(11A-B) \sin(c+dx)}{d(a \cos(c+dx)+a)}}{3a^2} - \frac{a(7A-2B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} - \frac{(A-B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2}$$

↓ 3042

$$\frac{15aA \int \csc(c+dx+\frac{\pi}{2}) dx - \frac{2a^2(11A-B) \sin(c+dx)}{d(a \cos(c+dx)+a)}}{3a^2} - \frac{a(7A-2B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} - \frac{(A-B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2}$$

↓ 4257

$$\frac{\frac{15aA \operatorname{arctanh}(\sin(c+dx))}{d} - \frac{2a^2(11A-B) \sin(c+dx)}{d(a \cos(c+dx)+a)}}{3a^2} - \frac{a(7A-2B) \sin(c+dx)}{3d(a \cos(c+dx)+a)^2} - \frac{(A-B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^3,x]`

output `-1/5*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^3) + (-1/3*(a*(7*A - 2*B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + ((15*a*A*ArcTanh[Sin[c + d*x]])/d - (2*a^2*(11*A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])))/(3*a^2))/(5*a^2)`

3.62.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.62.4 Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.81

method	result
parallelrisc	$\frac{-20A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 20A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \frac{10(2A-B) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3} \right)}{20a^3d}$
derivativedivides	$\frac{-4A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 4A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + \frac{\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{5} + \frac{2\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} - \frac{4\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} - \frac{A\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3}}{4da^3}$
default	$\frac{-4A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 4A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + \frac{\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{5} + \frac{2\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)B}{3} - \frac{4\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)A}{3} - \frac{A\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3}}{4da^3}$
risc	$-\frac{2i\left(15A e^{4i(dx+c)} + 75A e^{3i(dx+c)} + 145A e^{2i(dx+c)} - 20B e^{2i(dx+c)} + 95A e^{i(dx+c)} - 10B e^{i(dx+c)} + 22A - 2B\right)}{15da^3\left(e^{i(dx+c)} + 1\right)^5} + \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^3d}$
norman	$\frac{-\frac{(A-B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{20ad} - \frac{5(5A-B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{12ad} - \frac{(7A-B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4ad} - \frac{(23A-13B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{60ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a^2} + \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^3d}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE
)
```

```
output 1/20*(-20*A*ln(tan(1/2*d*x+1/2*c)-1)+20*A*ln(tan(1/2*d*x+1/2*c)+1)-tan(1/2
*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^4+10/3*(2*A-B)*tan(1/2*d*x+1/2*c)^2+
35*A-5*B))/a^3/d
```

$$3.62. \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^3} dx$$

3.62.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.58

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{15 (A \cos(dx + c)^3 + 3 A \cos(dx + c)^2 + 3 A \cos(dx + c) + A) \log(\sin(dx + c) + 1) - 15 (A \cos(dx + c)^3 + 3 A \cos(dx + c)^2 + 3 A \cos(dx + c) + A) \log(-\sin(dx + c) + 1) - 2 * (2 * (11 * A - B) * \cos(dx + c)^2 + 3 * (17 * A - 2 * B) * \cos(dx + c) + 32 * A - 7 * B) * \sin(dx + c)}{30 (a^3 d \cos(dx + c)^3 + 3 * a^3 d \cos(dx + c)^2 + 3 * a^3 d \cos(dx + c) + a^3 d)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^3,x, algorithm="fricas")`

output `1/30*(15*(A*cos(d*x + c)^3 + 3*A*cos(d*x + c)^2 + 3*A*cos(d*x + c) + A)*log(sin(d*x + c) + 1) - 15*(A*cos(d*x + c)^3 + 3*A*cos(d*x + c)^2 + 3*A*cos(d*x + c) + A)*log(-sin(d*x + c) + 1) - 2*(2*(11*A - B)*cos(d*x + c)^2 + 3*(17*A - 2*B)*cos(d*x + c) + 32*A - 7*B)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.62.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{\int \frac{A \sec(c+dx)}{\cos^3(c+dx)+3 \cos^2(c+dx)+3 \cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sec(c+dx)}{\cos^3(c+dx)+3 \cos^2(c+dx)+3 \cos(c+dx)+1} dx}{a^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))**3,x)`

output `(Integral(A*sec(c + d*x)/(cos(c + d*x)**3 + 3*cos(c + d*x)**2 + 3*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)/(cos(c + d*x)**3 + 3*cos(c + d*x)**2 + 3*cos(c + d*x) + 1), x))/a**3`

3.62.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.60

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{A \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} + \frac{20 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{60 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^3} + \frac{60 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^3} \right) - \frac{B \left(\frac{15 \sin(dx+c)}{\cos(dx+c)+1} + \frac{10 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{60 d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `-1/60*(A*((105*sin(d*x + c)/(cos(d*x + c) + 1) + 20*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3 - 60*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a^3 + 60*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a^3) - B*(15*sin(d*x + c)/(cos(d*x + c) + 1) + 10*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)/a^3/d`

3.62.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.26

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{60 A \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - 60 A \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right) - 3 A a^{12} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 3 B a^{12} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 20 A a^{12} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3}{60 d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^3,x, algorithm="giac")`

output `1/60*(60*A*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^3 - 60*A*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^3 - (3*A*a^12*tan(1/2*d*x + 1/2*c)^5 - 3*B*a^12*tan(1/2*d*x + 1/2*c)^5 + 20*A*a^12*tan(1/2*d*x + 1/2*c)^3 - 10*B*a^12*tan(1/2*d*x + 1/2*c)^3 + 105*A*a^12*tan(1/2*d*x + 1/2*c) - 15*B*a^12*tan(1/2*d*x + 1/2*c))/a^15)/d`

3.62.9 Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.11

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{2 A \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{a^3 d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{A-B}{4a^3} + \frac{3A+B}{4a^3} + \frac{3A-B}{4a^3}\right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A - B)}{20 a^3 d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{12a^3} + \frac{3A-B}{12a^3}\right)}{d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^3),x)`output `(2*A*atanh(tan(c/2 + (d*x)/2)))/(a^3*d) - (tan(c/2 + (d*x)/2)*((A - B)/(4*a^3) + (3*A + B)/(4*a^3) + (3*A - B)/(4*a^3)))/d - (tan(c/2 + (d*x)/2)^5*(A - B))/(20*a^3*d) - (tan(c/2 + (d*x)/2)^3*((A - B)/(12*a^3) + (3*A - B)/(12*a^3)))/d`

3.63
$$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^3} dx$$

3.63.1 Optimal result 723
 3.63.2 Mathematica [B] (verified) 724
 3.63.3 Rubi [A] (verified) 724
 3.63.4 Maple [A] (verified) 728
 3.63.5 Fricas [A] (verification not implemented) 728
 3.63.6 Sympy [F] 729
 3.63.7 Maxima [B] (verification not implemented) 729
 3.63.8 Giac [A] (verification not implemented) 730
 3.63.9 Mupad [B] (verification not implemented) 731

3.63.1 Optimal result

Integrand size = 31, antiderivative size = 145

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx = -\frac{(3A - B) \operatorname{arctanh}(\sin(c + dx))}{a^3 d} + \frac{2(36A - 11B) \tan(c + dx)}{15a^3 d} - \frac{(A - B) \tan(c + dx)}{5d(a + a \cos(c + dx))^3} - \frac{(9A - 4B) \tan(c + dx)}{15ad(a + a \cos(c + dx))^2} - \frac{(3A - B) \tan(c + dx)}{d(a^3 + a^3 \cos(c + dx))}$$

```
output -(3*A-B)*arctanh(sin(d*x+c))/a^3/d+2/15*(36*A-11*B)*tan(d*x+c)/a^3/d-1/5*(
A-B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^3-1/15*(9*A-4*B)*tan(d*x+c)/a/d/(a+a*co
s(d*x+c))^2-(3*A-B)*tan(d*x+c)/d/(a^3+a^3*cos(d*x+c))
```

3.63.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 482 vs. $2(145) = 290$.

Time = 2.78 (sec) , antiderivative size = 482, normalized size of antiderivative = 3.32

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{960(3A - B) \cos^6\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{(120a^3d(1 + \cos(c + dx))^3)}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^3,x]`

output `(960*(3*A - B)*Cos[(c + d*x)/2]^6*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Cos[(c + d*x)/2]*Sec[c/2]*Sec[c]*Sec[c + d*x]*(-5*(51*A - 32*B)*Sin[(d*x)/2] + (567*A - 167*B)*Sin[(3*d*x)/2] - 600*A*Sin[c - (d*x)/2] + 170*B*Sin[c - (d*x)/2] + 375*A*Sin[c + (d*x)/2] - 170*B*Sin[c + (d*x)/2] - 480*A*Sin[2*c + (d*x)/2] + 160*B*Sin[2*c + (d*x)/2] - 60*A*Sin[c + (3*d*x)/2] + 75*B*Sin[c + (3*d*x)/2] + 402*A*Sin[2*c + (3*d*x)/2] - 167*B*Sin[2*c + (3*d*x)/2] - 225*A*Sin[3*c + (3*d*x)/2] + 75*B*Sin[3*c + (3*d*x)/2] + 315*A*Sin[c + (5*d*x)/2] - 95*B*Sin[c + (5*d*x)/2] + 30*A*Sin[2*c + (5*d*x)/2] + 15*B*Sin[2*c + (5*d*x)/2] + 240*A*Sin[3*c + (5*d*x)/2] - 95*B*Sin[3*c + (5*d*x)/2] - 45*A*Sin[4*c + (5*d*x)/2] + 15*B*Sin[4*c + (5*d*x)/2] + 72*A*Sin[2*c + (7*d*x)/2] - 22*B*Sin[2*c + (7*d*x)/2] + 15*A*Sin[3*c + (7*d*x)/2] + 57*A*Sin[4*c + (7*d*x)/2] - 22*B*Sin[4*c + (7*d*x)/2]))/(120*a^3*d*(1 + Cos[c + d*x])^3)`

3.63.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.11, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3457, 3042, 3457, 3042, 3457, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^3} dx$$

↓ 3042

3.63. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
& \quad \downarrow \text{3457} \\
& \int \frac{\frac{a(6A-B) - 3a(A-B) \cos(c+dx)}{(\cos(c+dx)a+a)^2} \sec^2(c+dx)}{5a^2} dx - \frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\frac{a(6A-B) - 3a(A-B) \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2} dx}{5a^2} - \frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3457} \\
& \int \frac{\frac{a^2(27A-7B) - 2a^2(9A-4B) \cos(c+dx)}{\cos(c+dx)a+a} \sec^2(c+dx)}{3a^2} dx - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} - \frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\frac{a^2(27A-7B) - 2a^2(9A-4B) \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)} dx}{3a^2} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} - \frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3457} \\
& \int \frac{\frac{2a^3(36A-11B) - 15a^3(3A-B) \cos(c+dx)}{a^2} \sec^2(c+dx)}{3a^2} dx - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\frac{2a^3(36A-11B) - 15a^3(3A-B) \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2} dx}{a^2} - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3227} \\
& \frac{2a^3(36A-11B) \int \sec^2(c+dx) dx - 15a^3(3A-B) \int \sec(c+dx) dx}{3a^2} - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
& \quad \downarrow \\
& \frac{5a^2}{3a^2} \frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx) + a)^3}
\end{aligned}$$

3.63. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^3} dx$

$$\begin{array}{c}
\downarrow 3042 \\
\frac{2a^3(36A-11B) \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx - 15a^3(3A-B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx}{a^2} - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
\hline
\frac{5a^2}{(A-B) \tan(c+dx)} \\
\frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3} \\
\downarrow 4254 \\
-\frac{2a^3(36A-11B) \int \frac{1d(-\tan(c+dx))}{d} - 15a^3(3A-B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx}{a^2} - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
\hline
\frac{5a^2}{(A-B) \tan(c+dx)} \\
\frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3} \\
\downarrow 24 \\
\frac{2a^3(36A-11B) \tan(c+dx)}{d} - \frac{15a^3(3A-B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx}{a^2} - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
\hline
\frac{5a^2}{(A-B) \tan(c+dx)} \\
\frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3} \\
\downarrow 4257 \\
\frac{2a^3(36A-11B) \tan(c+dx)}{d} - \frac{15a^3(3A-B) \arctan\left(\frac{\sin(c+dx)}{d}\right)}{a^2} - \frac{15a^2(3A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{a(9A-4B) \tan(c+dx)}{3d(a \cos(c+dx)+a)^2} \\
\hline
\frac{5a^2}{(A-B) \tan(c+dx)} \\
\frac{(A-B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3}
\end{array}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^3,x]`

output `-1/5*((A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^3) + (-1/3*(a*(9*A - 4*B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + ((-15*a^2*(3*A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x]))) + ((-15*a^3*(3*A - B)*ArcTanh[Sin[c + d*x]])/d + (2*a^3*(36*A - 11*B)*Tan[c + d*x])/d)/a^2)/(3*a^2))/(5*a^2)`

3.63.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.63.4 Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.99

method	result
parallelrisc	$\frac{3 \cos(dx+c) \left(A - \frac{B}{3}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - 3 \cos(dx+c) \left(A - \frac{B}{3}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + \frac{57 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\frac{A - 17B}{57}\right) \cos(2dx)}{a^3 d \cos(dx+c)}$
derivativedivides	$\frac{A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) B + 2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) A - \frac{4 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) B}{3} + 17 A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 7 B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (12)}{4d a^3}$
default	$\frac{A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) B + 2 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) A - \frac{4 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) B}{3} + 17 A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 7 B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (12)}{4d a^3}$
norman	$\frac{(A-B) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{20ad} + \frac{(3A-2B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} - \frac{(15A-2B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{6ad} - \frac{(25A-7B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4ad} + \frac{(42A-17B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{10ad}$
risc	$\frac{2i(45A e^{6i(dx+c)} - 15B e^{6i(dx+c)} + 225A e^{5i(dx+c)} - 75B e^{5i(dx+c)} + 480A e^{4i(dx+c)} - 160B e^{4i(dx+c)} + 600A e^{3i(dx+c)} - 120B e^{2i(dx+c)} + 60A e^{2i(dx+c)} - 12A e^{i(dx+c)} + 6A)}{15d a^3 (e^{i(dx+c)} + 1)^5 (e^{2i(dx+c)} - 1)}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE)`

output `3/20*(20*cos(d*x+c)*(A-1/3*B)*ln(tan(1/2*d*x+1/2*c)-1)-20*cos(d*x+c)*(A-1/3*B)*ln(tan(1/2*d*x+1/2*c)+1)+19*tan(1/2*d*x+1/2*c)*(1/2*(A-17/57*B)*cos(2*d*x+2*c)+1/19*(2*A-11/18*B)*cos(3*d*x+3*c)+(A-97/342*B)*cos(d*x+c)+67/114*A-17/114*B)*sec(1/2*d*x+1/2*c)^4)/d/a^3/cos(d*x+c)`

3.63.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.88

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{15((3A - B) \cos(dx + c))^4 + 3(3A - B) \cos(dx + c)^3 + 3(3A - B) \cos(dx + c)^2 + (3A - B) \cos(dx + c)}{15d a^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^3,x, algorithm="fracas")`

output
$$\begin{aligned} & -1/30*(15*((3*A - B)*\cos(d*x + c))^4 + 3*(3*A - B)*\cos(d*x + c)^3 + 3*(3*A \\ & - B)*\cos(d*x + c)^2 + (3*A - B)*\cos(d*x + c))*\log(\sin(d*x + c) + 1) - 15*(\\ & (3*A - B)*\cos(d*x + c)^4 + 3*(3*A - B)*\cos(d*x + c)^3 + 3*(3*A - B)*\cos(d* \\ & x + c)^2 + (3*A - B)*\cos(d*x + c))*\log(-\sin(d*x + c) + 1) - 2*(2*(36*A - 1 \\ & 1*B)*\cos(d*x + c)^3 + 3*(57*A - 17*B)*\cos(d*x + c)^2 + (117*A - 32*B)*\cos(\\ & d*x + c) + 15*A)*\sin(d*x + c))/(a^3*d*\cos(d*x + c)^4 + 3*a^3*d*\cos(d*x + c \\ &)^3 + 3*a^3*d*\cos(d*x + c)^2 + a^3*d*\cos(d*x + c)) \end{aligned}$$

3.63.6 Sympy [F]

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx \\ & = \frac{\int \frac{A \sec^2(c + dx)}{\cos^3(c + dx) + 3 \cos^2(c + dx) + 3 \cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec^2(c + dx)}{\cos^3(c + dx) + 3 \cos^2(c + dx) + 3 \cos(c + dx) + 1} dx}{a^3} \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c))**3,x)`

output
$$\begin{aligned} & (\text{Integral}(A*\sec(c + d*x)**2/(\cos(c + d*x)**3 + 3*\cos(c + d*x)**2 + 3*\cos(c \\ & + d*x) + 1), x) + \text{Integral}(B*\cos(c + d*x)*\sec(c + d*x)**2/(\cos(c + d*x)** \\ & 3 + 3*\cos(c + d*x)**2 + 3*\cos(c + d*x) + 1), x))/a**3 \end{aligned}$$

3.63.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 286 vs. $2(139) = 278$.

Time = 0.22 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.97

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx \\ & = \frac{3A \left(\frac{40 \sin(dx+c)}{\left(a^3 - \frac{a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2}\right) (\cos(dx+c)+1)} + \frac{\frac{85 \sin(dx+c)}{\cos(dx+c)+1} + \frac{10 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{\sin(dx+c)^5}{(\cos(dx+c)+1)^5}}{a^3} - \frac{60 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^3} + \frac{60 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^3} \right)}{60d} \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output $\frac{1}{60} \cdot (3A \cdot (40 \sin(dx + c) / ((a^3 - a^3 \sin(dx + c))^2 / (\cos(dx + c) + 1)^2) \cdot (\cos(dx + c) + 1)) + (85 \sin(dx + c) / (\cos(dx + c) + 1) + 10 \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + \sin(dx + c)^5 / (\cos(dx + c) + 1)^5) / a^3 - 60 \log(\sin(dx + c) / (\cos(dx + c) + 1) + 1) / a^3 + 60 \log(\sin(dx + c) / (\cos(dx + c) + 1) - 1) / a^3) - B \cdot ((105 \sin(dx + c) / (\cos(dx + c) + 1) + 20 \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + 3 \sin(dx + c)^5 / (\cos(dx + c) + 1)^5) / a^3 - 60 \log(\sin(dx + c) / (\cos(dx + c) + 1) + 1) / a^3 + 60 \log(\sin(dx + c) / (\cos(dx + c) + 1) - 1) / a^3) / d$

3.63.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.31

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{60(3A - B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^3} - \frac{60(3A - B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^3} + \frac{120 A \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1) a^3} - \frac{3 A a^{12} \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 3 B}{60 d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^3,x, algorithm="giac")`

output $\frac{-1}{60} \cdot (60 \cdot (3A - B) \cdot \log(\text{abs}(\tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + 1)) / a^3 - 60 \cdot (3A - B) \cdot \log(\text{abs}(\tan(1/2 \cdot d \cdot x + 1/2 \cdot c) - 1)) / a^3 + 120 \cdot A \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) / ((\tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 - 1) \cdot a^3) - (3 \cdot A \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 - 3 \cdot B \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 + 30 \cdot A \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 - 20 \cdot B \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 + 255 \cdot A \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) - 105 \cdot B \cdot a^{12} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)) / a^{15}) / d$

3.63.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.16

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{6a^3} + \frac{4A-2B}{12a^3}\right)}{d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3A}{2a^3} + \frac{3(A-B)}{4a^3} + \frac{4A-2B}{2a^3}\right)}{d} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A-B)}{20a^3d} - \frac{2A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - a^3\right)} - \frac{2 \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (3A - B)}{a^3d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^3),x)`output `(tan(c/2 + (d*x)/2)^3*((A - B)/(6*a^3) + (4*A - 2*B)/(12*a^3))/d + (tan(c/2 + (d*x)/2)*((3*A)/(2*a^3) + (3*(A - B))/(4*a^3) + (4*A - 2*B)/(2*a^3)))/d + (tan(c/2 + (d*x)/2)^5*(A - B))/(20*a^3*d) - (2*A*tan(c/2 + (d*x)/2))/(d*(a^3*tan(c/2 + (d*x)/2)^2 - a^3)) - (2*atanh(tan(c/2 + (d*x)/2))*(3*A - B))/(a^3*d)`

3.64 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^3} dx$

3.64.1 Optimal result 732
 3.64.2 Mathematica [B] (verified) 733
 3.64.3 Rubi [A] (verified) 733
 3.64.4 Maple [A] (verified) 737
 3.64.5 Fricas [A] (verification not implemented) 738
 3.64.6 Sympy [F] 739
 3.64.7 Maxima [B] (verification not implemented) 739
 3.64.8 Giac [A] (verification not implemented) 740
 3.64.9 Mupad [B] (verification not implemented) 741

3.64.1 Optimal result

Integrand size = 31, antiderivative size = 196

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx = \frac{(13A - 6B) \operatorname{arctanh}(\sin(c + dx))}{2a^3d} - \frac{8(19A - 9B) \tan(c + dx)}{15a^3d} + \frac{(13A - 6B) \sec(c + dx) \tan(c + dx)}{2a^3d} - \frac{(A - B) \sec(c + dx) \tan(c + dx)}{5d(a + a \cos(c + dx))^3} - \frac{(11A - 6B) \sec(c + dx) \tan(c + dx)}{15ad(a + a \cos(c + dx))^2} - \frac{4(19A - 9B) \sec(c + dx) \tan(c + dx)}{15d(a^3 + a^3 \cos(c + dx))}$$

```
output 1/2*(13*A-6*B)*arctanh(sin(d*x+c))/a^3/d-8/15*(19*A-9*B)*tan(d*x+c)/a^3/d+
1/2*(13*A-6*B)*sec(d*x+c)*tan(d*x+c)/a^3/d-1/5*(A-B)*sec(d*x+c)*tan(d*x+c)
/d/(a+a*cos(d*x+c))^3-1/15*(11*A-6*B)*sec(d*x+c)*tan(d*x+c)/a/d/(a+a*cos(d
*x+c))^2-4/15*(19*A-9*B)*sec(d*x+c)*tan(d*x+c)/d/(a^3+a^3*cos(d*x+c))
```

3.64.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 610 vs. $2(196) = 392$.

Time = 4.02 (sec) , antiderivative size = 610, normalized size of antiderivative = 3.11

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx =$$

$$\frac{1920(13A - 6B) \cos^6\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{a^3 d (1 + \cos(c + dx))^3}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^3,x]`

output

```
-1/480*(1920*(13*A - 6*B)*Cos[(c + d*x)/2]^6*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Cos[(c + d*x)/2]*Sec[c/2]*Sec[c]*Sec[c + d*x]^2*((-1235*A + 870*B)*Sin[(d*x)/2] + 5*(761*A - 366*B)*Sin[(3*d*x)/2] - 4329*A*Sin[c - (d*x)/2] + 2094*B*Sin[c - (d*x)/2] + 1989*A*Sin[c + (d*x)/2] - 1314*B*Sin[c + (d*x)/2] - 3575*A*Sin[2*c + (d*x)/2] + 1650*B*Sin[2*c + (d*x)/2] - 475*A*Sin[c + (3*d*x)/2] + 450*B*Sin[c + (3*d*x)/2] + 2005*A*Sin[2*c + (3*d*x)/2] - 1230*B*Sin[2*c + (3*d*x)/2] - 2275*A*Sin[3*c + (3*d*x)/2] + 1050*B*Sin[3*c + (3*d*x)/2] + 2673*A*Sin[c + (5*d*x)/2] - 1278*B*Sin[c + (5*d*x)/2] + 105*A*Sin[2*c + (5*d*x)/2] + 90*B*Sin[2*c + (5*d*x)/2] + 1593*A*Sin[3*c + (5*d*x)/2] - 918*B*Sin[3*c + (5*d*x)/2] - 975*A*Sin[4*c + (5*d*x)/2] + 450*B*Sin[4*c + (5*d*x)/2] + 1325*A*Sin[2*c + (7*d*x)/2] - 630*B*Sin[2*c + (7*d*x)/2] + 255*A*Sin[3*c + (7*d*x)/2] - 60*B*Sin[3*c + (7*d*x)/2] + 875*A*Sin[4*c + (7*d*x)/2] - 480*B*Sin[4*c + (7*d*x)/2] - 195*A*Sin[5*c + (7*d*x)/2] + 90*B*Sin[5*c + (7*d*x)/2] + 304*A*Sin[3*c + (9*d*x)/2] - 144*B*Sin[3*c + (9*d*x)/2] + 90*A*Sin[4*c + (9*d*x)/2] - 30*B*Sin[4*c + (9*d*x)/2] + 214*A*Sin[5*c + (9*d*x)/2] - 114*B*Sin[5*c + (9*d*x)/2]))/(a^3*d*(1 + Cos[c + d*x])^3)
```

3.64.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3457, 3042, 3457, 3042, 3457, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.64. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(a(7A-2B)-4a(A-B)\cos(c+dx))\sec^3(c+dx)}{(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(7A-2B)-4a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(a^2(43A-18B)-3a^2(11A-6B)\cos(c+dx))\sec^3(c+dx)}{\cos(c+dx)a+a} dx}{3a^2} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(43A-18B)-3a^2(11A-6B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3457} \\
& \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(15a^3(13A-6B)-8a^3(19A-9B)\cos(c+dx))\sec^3(c+dx)dx}{a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)}}{3a^2} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}
\end{aligned}$$

3.64. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^3} dx$

$$\frac{\int \frac{15a^3(13A-6B)-8a^3(19A-9B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^3} dx}{3a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3227

$$\frac{15a^3(13A-6B)\int \sec^3(c+dx)dx - 8a^3(19A-9B)\int \sec^2(c+dx)dx}{3a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{15a^3(13A-6B)\int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx - 8a^3(19A-9B)\int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx}{3a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 4254

$$\frac{8a^3(19A-9B)\int \frac{1d(-\tan(c+dx))}{d} + 15a^3(13A-6B)\int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx}{3a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 24

$$\frac{15a^3(13A-6B)\int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx - 8a^3(19A-9B)\frac{\tan(c+dx)}{d}}{3a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 4255

$$\frac{15a^3(13A-6B)\left(\frac{1}{2}\int \sec(c+dx)dx + \frac{\tan(c+dx)\sec(c+dx)}{2d}\right) - 8a^3(19A-9B)\frac{\tan(c+dx)}{d}}{3a^2} - \frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

3.64. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^3} dx$

↓ 3042

$$\frac{15a^3(13A-6B)\left(\frac{1}{2}\int \csc\left(c+dx+\frac{\pi}{2}\right)dx+\frac{\tan(c+dx)\sec(c+dx)}{2d}\right)-\frac{8a^3(19A-9B)\tan(c+dx)}{d}-\frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)}-\frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)}}{a^2} \frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

↓ 4257

$$\frac{15a^3(13A-6B)\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d}+\frac{\tan(c+dx)\sec(c+dx)}{2d}\right)-\frac{8a^3(19A-9B)\tan(c+dx)}{d}-\frac{4a^2(19A-9B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)}-\frac{a(11A-6B)\tan(c+dx)\sec(c+dx)}{3d(a\cos(c+dx)+a)}}{a^2} \frac{5a^2}{3a^2} \frac{(A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^3,x]`

output `-1/5*((A - B)*Sec[c + d*x]*Tan[c + d*x]/(d*(a + a*Cos[c + d*x])^3) + (-1/3*(a*(11*A - 6*B)*Sec[c + d*x]*Tan[c + d*x]/(d*(a + a*Cos[c + d*x])^2) + ((-4*a^2*(19*A - 9*B)*Sec[c + d*x]*Tan[c + d*x]/(d*(a + a*Cos[c + d*x]))) + ((-8*a^3*(19*A - 9*B)*Tan[c + d*x])/d + 15*a^3*(13*A - 6*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/a^2)/(3*a^2))/(5*a^2)`

3.64.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
  Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 4254 Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4255 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
  /; FreeQ[{c, d}, x]
```

3.64.4 Maple [A] (verified)

Time = 1.34 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.89

method	result
parallelrisch	$-1560(1+\cos(2dx+2c))\left(A-\frac{6B}{13}\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+1560(1+\cos(2dx+2c))\left(A-\frac{6B}{13}\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)-152\left(\sec\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^4$
derivativedivides	$-\frac{A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}+\frac{\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{5}-\frac{8\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{3}+2\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B-31A\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+17B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)$
default	$-\frac{A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}+\frac{\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{5}-\frac{8\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{3}+2\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B-31A\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+17B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)$
norman	$\frac{(A-B)\left(\tan^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{20ad}-\frac{(37A-27B)\left(\tan^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{60ad}-\frac{(51A-25B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{4ad}+\frac{(109A-45B)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{12ad}-\frac{(211A-111B)\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{4ad}$
risch	$-\frac{i\left(195Ae^{8i(dx+c)}-90Be^{8i(dx+c)}+975Ae^{7i(dx+c)}-450Be^{7i(dx+c)}+2275Ae^{6i(dx+c)}-1050Be^{6i(dx+c)}+3575Ae^{5i(dx+c)}-1050Be^{4i(dx+c)}+1050Be^{3i(dx+c)}-3575Ae^{2i(dx+c)}+90Be^{2i(dx+c)}-195Ae^{i(dx+c)}\right)}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2a^2}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a)^3,x,method=_RETURNVERBOSE)`

output `1/240*(-1560*(1+cos(2*d*x+2*c))*(A-6/13*B)*ln(tan(1/2*d*x+1/2*c)-1)+1560*(1+cos(2*d*x+2*c))*(A-6/13*B)*ln(tan(1/2*d*x+1/2*c)+1)-152*sec(1/2*d*x+1/2*c)^4*((783/76*A-189/38*B)*cos(2*d*x+2*c)+(717/152*A-9/4*B)*cos(3*d*x+3*c)+(A-9/19*B)*cos(4*d*x+4*c)+(2331/152*A-573/76*B)*cos(d*x+c)+677/76*A-9/2*B)*tan(1/2*d*x+1/2*c))/d/a^3/(1+cos(2*d*x+2*c))`

3.64.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.51

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{15 \left((13A - 6B) \cos(dx + c)^5 + 3(13A - 6B) \cos(dx + c)^4 + 3(13A - 6B) \cos(dx + c)^3 + (13A - 6B) \cos(dx + c)^2 + 3(13A - 6B) \cos(dx + c) + 15A \right)}{a^3 (1 + \cos(2dx + 2c))}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^3,x, algorithm="fracas")`

```
output 1/60*(15*((13*A - 6*B)*cos(d*x + c)^5 + 3*(13*A - 6*B)*cos(d*x + c)^4 + 3*
(13*A - 6*B)*cos(d*x + c)^3 + (13*A - 6*B)*cos(d*x + c)^2)*log(sin(d*x + c
) + 1) - 15*((13*A - 6*B)*cos(d*x + c)^5 + 3*(13*A - 6*B)*cos(d*x + c)^4 +
3*(13*A - 6*B)*cos(d*x + c)^3 + (13*A - 6*B)*cos(d*x + c)^2)*log(-sin(d*x
+ c) + 1) - 2*(16*(19*A - 9*B)*cos(d*x + c)^4 + 3*(239*A - 114*B)*cos(d*x
+ c)^3 + (479*A - 234*B)*cos(d*x + c)^2 + 15*(3*A - 2*B)*cos(d*x + c) - 1
5*A)*sin(d*x + c))/(a^3*d*cos(d*x + c)^5 + 3*a^3*d*cos(d*x + c)^4 + 3*a^3*
d*cos(d*x + c)^3 + a^3*d*cos(d*x + c)^2)
```

3.64.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{\int \frac{A \sec^3(c + dx)}{\cos^3(c + dx) + 3 \cos^2(c + dx) + 3 \cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec^3(c + dx)}{\cos^3(c + dx) + 3 \cos^2(c + dx) + 3 \cos(c + dx) + 1} dx}{a^3}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c))**3,x)
```

```
output (Integral(A*sec(c + d*x)**3/(cos(c + d*x)**3 + 3*cos(c + d*x)**2 + 3*cos(c
+ d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**3/(cos(c + d*x)**
3 + 3*cos(c + d*x)**2 + 3*cos(c + d*x) + 1), x))/a**3
```

3.64.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 377 vs. $2(184) = 368$.

Time = 0.25 (sec) , antiderivative size = 377, normalized size of antiderivative = 1.92

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx =$$

$$A \left(\frac{60 \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} - \frac{7 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^3 - \frac{2a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} + \frac{465 \sin(dx+c) + 40 \sin(dx+c)^3 + 3 \sin(dx+c)^5}{a^3 \cos(dx+c)+1} - \frac{390 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^3} + \frac{390 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^3} \right)$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^3,x, algorithm="m
axima")
```

3.64. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^3} dx$

output
$$\begin{aligned} & -1/60*(A*(60*(5*\sin(dx + c)/(\cos(dx + c) + 1) - 7*\sin(dx + c)^3/(\cos(dx + c) + 1)^3)/(a^3 - 2*a^3*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + a^3*\sin(dx + c)^4/(\cos(dx + c) + 1)^4) + (465*\sin(dx + c)/(\cos(dx + c) + 1) + 40*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 3*\sin(dx + c)^5/(\cos(dx + c) + 1)^5)/a^3 - 390*\log(\sin(dx + c)/(\cos(dx + c) + 1) + 1)/a^3 + 390*\log(\sin(dx + c)/(\cos(dx + c) + 1) - 1)/a^3) - 3*B*(40*\sin(dx + c)/((a^3 - a^3*\sin(dx + c)^2/(\cos(dx + c) + 1)^2)*(cos(dx + c) + 1)) + (85*\sin(dx + c))/(\cos(dx + c) + 1) + 10*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + \sin(dx + c)^5/(\cos(dx + c) + 1)^5)/a^3 - 60*\log(\sin(dx + c)/(\cos(dx + c) + 1) + 1)/a^3 + 60*\log(\sin(dx + c)/(\cos(dx + c) + 1) - 1)/a^3))/d \end{aligned}$$

3.64.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.19

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{30(13A - 6B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^3} - \frac{30(13A - 6B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^3} + \frac{60(7A \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 2B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 5A \tan(\frac{1}{2} dx + \frac{1}{2} c))}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1)^2 a}$$

input `integrate((A+B*cos(dx+c))*sec(dx+c)^3/(a+a*cos(dx+c))^3,x, algorithm="giac")`

output
$$\begin{aligned} & 1/60*(30*(13*A - 6*B)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1))/a^3 - 30*(13*A - 6*B)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1))/a^3 + 60*(7*A*\tan(1/2*d*x + 1/2*c)^3 - 2*B*\tan(1/2*d*x + 1/2*c)^3 - 5*A*\tan(1/2*d*x + 1/2*c) + 2*B*\tan(1/2*d*x + 1/2*c))/((\tan(1/2*d*x + 1/2*c)^2 - 1)^2*a^3) - (3*A*a^12*\tan(1/2*d*x + 1/2*c)^5 - 3*B*a^12*\tan(1/2*d*x + 1/2*c)^5 + 40*A*a^12*\tan(1/2*d*x + 1/2*c)^3 - 30*B*a^12*\tan(1/2*d*x + 1/2*c)^3 + 465*A*a^12*\tan(1/2*d*x + 1/2*c) - 255*B*a^12*\tan(1/2*d*x + 1/2*c))/a^15)/d \end{aligned}$$

3.64.9 Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.10

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (7A - 2B) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (5A - 2B)}{d \left(a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^3\right)}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3(A-B)}{2a^3} + \frac{3(5A-3B)}{4a^3} + \frac{10A-2B}{4a^3}\right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{4a^3} + \frac{5A-3B}{12a^3}\right)}{d}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A - B)}{20a^3 d} + \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (13A - 6B)}{a^3 d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^3),x)`output `(tan(c/2 + (d*x)/2)^3*(7*A - 2*B) - tan(c/2 + (d*x)/2)*(5*A - 2*B))/(d*(a^3*tan(c/2 + (d*x)/2)^4 - 2*a^3*tan(c/2 + (d*x)/2)^2 + a^3)) - (tan(c/2 + (d*x)/2)*((3*(A - B))/(2*a^3) + (3*(5*A - 3*B))/(4*a^3) + (10*A - 2*B)/(4*a^3)))/d - (tan(c/2 + (d*x)/2)^3*((A - B)/(4*a^3) + (5*A - 3*B)/(12*a^3)))/d - (tan(c/2 + (d*x)/2)^5*(A - B))/(20*a^3*d) + (atanh(tan(c/2 + (d*x)/2))*(13*A - 6*B))/(a^3*d)`

3.65 $\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

3.65.1	Optimal result	742
3.65.2	Mathematica [B] (verified)	743
3.65.3	Rubi [A] (verified)	743
3.65.4	Maple [A] (verified)	746
3.65.5	Fricas [A] (verification not implemented)	747
3.65.6	Sympy [B] (verification not implemented)	747
3.65.7	Maxima [A] (verification not implemented)	748
3.65.8	Giac [A] (verification not implemented)	749
3.65.9	Mupad [B] (verification not implemented)	750

3.65.1 Optimal result

Integrand size = 31, antiderivative size = 229

$$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = -\frac{(8A-21B)x}{2a^4} + \frac{8(83A-216B) \sin(c+dx)}{105a^4d} - \frac{(8A-21B) \cos(c+dx) \sin(c+dx)}{2a^4d} + \frac{(52A-129B) \cos^3(c+dx) \sin(c+dx)}{105a^4d(1+\cos(c+dx))^2} + \frac{4(83A-216B) \cos^2(c+dx) \sin(c+dx)}{105a^4d(1+\cos(c+dx))} + \frac{(A-B) \cos^5(c+dx) \sin(c+dx)}{7d(a+a \cos(c+dx))^4} + \frac{(A-2B) \cos^4(c+dx) \sin(c+dx)}{5ad(a+a \cos(c+dx))^3}$$

output

```
-1/2*(8*A-21*B)*x/a^4+8/105*(83*A-216*B)*sin(d*x+c)/a^4/d-1/2*(8*A-21*B)*c
os(d*x+c)*sin(d*x+c)/a^4/d+1/105*(52*A-129*B)*cos(d*x+c)^3*sin(d*x+c)/a^4/
d/(1+cos(d*x+c))^2+4/105*(83*A-216*B)*cos(d*x+c)^2*sin(d*x+c)/a^4/d/(1+cos
(d*x+c))+1/7*(A-B)*cos(d*x+c)^5*sin(d*x+c)/d/(a+a*cos(d*x+c))^4+1/5*(A-2*B
)*cos(d*x+c)^4*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```

3.65.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 555 vs. $2(229) = 458$.

Time = 5.03 (sec) , antiderivative size = 555, normalized size of antiderivative = 2.42

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(-14700(8A-21B)dx \cos\left(\frac{dx}{2}\right) - 14700(8A-21B)dx \cos\left(c+\frac{dx}{2}\right) - 70560Adx\right)}{(a+a\cos(c+dx))^4}$$

input `Integrate[(Cos[c + d*x]^5*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(-14700*(8*A - 21*B)*d*x*Cos[(d*x)/2] - 14700*(8*A - 21*B)*d*x*Cos[c + (d*x)/2] - 70560*A*d*x*Cos[c + (3*d*x)/2] + 185220*B*d*x*Cos[c + (3*d*x)/2] - 70560*A*d*x*Cos[2*c + (3*d*x)/2] + 185220*B*d*x*Cos[2*c + (3*d*x)/2] - 23520*A*d*x*Cos[2*c + (5*d*x)/2] + 61740*B*d*x*Cos[2*c + (5*d*x)/2] - 23520*A*d*x*Cos[3*c + (5*d*x)/2] + 61740*B*d*x*Cos[3*c + (5*d*x)/2] - 3360*A*d*x*Cos[3*c + (7*d*x)/2] + 8820*B*d*x*Cos[3*c + (7*d*x)/2] - 3360*A*d*x*Cos[4*c + (7*d*x)/2] + 8820*B*d*x*Cos[4*c + (7*d*x)/2] + 243320*A*Sin[(d*x)/2] - 539490*B*Sin[(d*x)/2] - 184520*A*Sin[c + (d*x)/2] + 386190*B*Sin[c + (d*x)/2] + 184464*A*Sin[c + (3*d*x)/2] - 422478*B*Sin[c + (3*d*x)/2] - 72240*A*Sin[2*c + (3*d*x)/2] + 132930*B*Sin[2*c + (3*d*x)/2] + 77168*A*Sin[2*c + (5*d*x)/2] - 181461*B*Sin[2*c + (5*d*x)/2] - 8400*A*Sin[3*c + (5*d*x)/2] + 3675*B*Sin[3*c + (5*d*x)/2] + 15164*A*Sin[3*c + (7*d*x)/2] - 36003*B*Sin[3*c + (7*d*x)/2] + 2940*A*Sin[4*c + (7*d*x)/2] - 9555*B*Sin[4*c + (7*d*x)/2] + 420*A*Sin[4*c + (9*d*x)/2] - 945*B*Sin[4*c + (9*d*x)/2] + 420*A*Sin[5*c + (9*d*x)/2] - 945*B*Sin[5*c + (9*d*x)/2] + 105*B*Sin[5*c + (11*d*x)/2] + 105*B*Sin[6*c + (11*d*x)/2]))/(6720*a^4*d*(1 + Cos[c + d*x])^4)`

3.65.3 Rubi [A] (verified)

Time = 1.27 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3456, 3042, 3456, 3042, 3456, 3042, 3456, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.65. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

$$\begin{aligned}
& \int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^5(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^4} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^4(c+dx)(5a(A-B)-a(2A-9B)\cos(c+dx))}{(\cos(c+dx)a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^4(5a(A-B)-a(2A-9B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^3(c+dx)(28a^2(A-2B)-a^2(24A-73B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{7a(A-2B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^3(28a^2(A-2B)-a^2(24A-73B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{7a(A-2B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^2(c+dx)(3a^3(52A-129B)-a^3(176A-477B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{(52A-129B)\sin(c+dx)\cos^3(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{7a(A-2B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{5a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.65. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

$$\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(3a^3(52A-129B)-a^3(176A-477B)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx}{\frac{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}{3a^2}} + \frac{(52A-129B)\sin(c+dx)\cos^3(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{7a(A-2B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{7a^2}{5a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3456

$$\frac{\int \frac{\cos(c+dx)\left(8a^4(83A-216B)-105a^4(8A-21B)\cos(c+dx)\right) dx}{a^2} + \frac{4a^3(83A-216B)\sin(c+dx)\cos^2(c+dx)}{d(a\cos(c+dx)+a)} + \frac{(52A-129B)\sin(c+dx)\cos^3(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{7a(A-2B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{7a^2}{5a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3042

$$\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)\left(8a^4(83A-216B)-105a^4(8A-21B)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx}{a^2} + \frac{4a^3(83A-216B)\sin(c+dx)\cos^2(c+dx)}{d(a\cos(c+dx)+a)} + \frac{(52A-129B)\sin(c+dx)\cos^3(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{7a(A-2B)\sin(c+dx)\cos^4(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{7a^2}{5a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3213

$$\frac{\frac{4a^3(83A-216B)\sin(c+dx)\cos^2(c+dx)}{d(a\cos(c+dx)+a)} + \frac{8a^4(83A-216B)\sin(c+dx)}{d} - \frac{105a^4(8A-21B)\sin(c+dx)\cos(c+dx)}{a^2} - \frac{105}{2}a^4x(8A-21B)}{3a^2} + \frac{(52A-129B)\sin(c+dx)\cos^3(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{7a^2}{5a^2} \frac{(A-B)\sin(c+dx)\cos^5(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

```
input Int[(Cos[c + d*x]^5*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]
```

```
output ((A - B)*Cos[c + d*x]^5*Sin[c + d*x])/(7*d*(a + a*Cos[c + d*x])^4) + ((7*a
*(A - 2*B)*Cos[c + d*x]^4*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + (((
52*A - 129*B)*Cos[c + d*x]^3*Sin[c + d*x])/(3*d*(1 + Cos[c + d*x])^2) + ((
4*a^3*(83*A - 216*B)*Cos[c + d*x]^2*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))
+ ((-105*a^4*(8*A - 21*B)*x)/2 + (8*a^4*(83*A - 216*B)*Sin[c + d*x])/d -
(105*a^4*(8*A - 21*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/a^2)/(3*a^2))/(5*a
^2))/(7*a^2)
```

3.65. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

3.65.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.65.4 Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.54

method	result
parallelrisch	$420 \left(\left(\frac{10964A}{105} - \frac{9376B}{35} \right) \cos(2dx+2c) + \left(\frac{2368A}{105} - \frac{7873B}{140} \right) \cos(3dx+3c) + (A-2B) \cos(4dx+4c) + \frac{B \cos(5dx+5c)}{4} + \left(\frac{24992A}{105} - \frac{26880A^4d}{105} \right) \right)$
derivativedivides	$-\frac{(\tan^7(\frac{dx}{2} + \frac{c}{2}))A}{7} + \frac{(\tan^7(\frac{dx}{2} + \frac{c}{2}))B}{7} + \frac{7A(\tan^5(\frac{dx}{2} + \frac{c}{2}))}{5} - \frac{9(\tan^5(\frac{dx}{2} + \frac{c}{2}))B}{5} - \frac{23(\tan^3(\frac{dx}{2} + \frac{c}{2}))A}{3} + 13(\tan^3(\frac{dx}{2} + \frac{c}{2}))$
default	$-\frac{(\tan^7(\frac{dx}{2} + \frac{c}{2}))A}{7} + \frac{(\tan^7(\frac{dx}{2} + \frac{c}{2}))B}{7} + \frac{7A(\tan^5(\frac{dx}{2} + \frac{c}{2}))}{5} - \frac{9(\tan^5(\frac{dx}{2} + \frac{c}{2}))B}{5} - \frac{23(\tan^3(\frac{dx}{2} + \frac{c}{2}))A}{3} + 13(\tan^3(\frac{dx}{2} + \frac{c}{2}))$
risch	$-\frac{4xA}{a^4} + \frac{21Bx}{2a^4} - \frac{iB e^{2i(dx+c)}}{8a^4d} - \frac{i e^{i(dx+c)}A}{2a^4d} + \frac{2i e^{i(dx+c)}B}{a^4d} + \frac{i e^{-i(dx+c)}A}{2a^4d} - \frac{2i e^{-i(dx+c)}B}{a^4d} + \frac{iB e^{-2i(dx+c)}}{8a^4d}$

input `int(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBOSE)`

3.65.
$$\int \frac{\cos^5(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$$

output $1/26880*(420*((10964/105*A-9376/35*B)*\cos(2*d*x+2*c)+(2368/105*A-7873/140*B)*\cos(3*d*x+3*c)+(A-2*B)*\cos(4*d*x+4*c)+1/4*B*\cos(5*d*x+5*c)+(24992/105*A-42881/70*B)*\cos(d*x+c)+16171/105*A-13914/35*B)*\tan(1/2*d*x+1/2*c)*\sec(1/2*d*x+1/2*c)^6-107520*(A-21/8*B)*x*d)/a^4/d$

3.65.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.04

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx = \frac{105(8A-21B)dx\cos(dx+c)^4 + 420(8A-21B)dx\cos(dx+c)^3 + 630(8A-21B)dx\cos(dx+c)^2 + 420(8A-21B)d^2x\cos(dx+c)^2 + 630(8A-21B)d^2x\cos(dx+c) + 420(8A-21B)d^2x + 105(8A-21B)d^2x - (105B\cos(dx+c)^5 + 210(A-2B)\cos(dx+c)^4 + 4(592A-1509B)\cos(dx+c)^3 + 4(1318A-3411B)\cos(dx+c)^2 + (4472A-11619B)\cos(dx+c) + 1328A-3456B)\sin(dx+c)}{(a^4d\cos(dx+c)^4 + 4a^4d\cos(dx+c)^3 + 6a^4d\cos(dx+c)^2 + 4a^4d\cos(dx+c) + a^4d)}$$

input `integrate(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="fricas")`

output $-1/210*(105*(8*A - 21*B)*d*x*\cos(d*x + c)^4 + 420*(8*A - 21*B)*d*x*\cos(d*x + c)^3 + 630*(8*A - 21*B)*d*x*\cos(d*x + c)^2 + 420*(8*A - 21*B)*d*x*\cos(d*x + c) + 105*(8*A - 21*B)*d*x - (105*B*\cos(d*x + c)^5 + 210*(A - 2*B)*\cos(d*x + c)^4 + 4*(592*A - 1509*B)*\cos(d*x + c)^3 + 4*(1318*A - 3411*B)*\cos(d*x + c)^2 + (4472*A - 11619*B)*\cos(d*x + c) + 1328*A - 3456*B)*\sin(d*x + c))/(a^4*d*\cos(d*x + c)^4 + 4*a^4*d*\cos(d*x + c)^3 + 6*a^4*d*\cos(d*x + c)^2 + 4*a^4*d*\cos(d*x + c) + a^4*d)$

3.65.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1085 vs. $2(216) = 432$.

Time = 7.97 (sec) , antiderivative size = 1085, normalized size of antiderivative = 4.74

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)**5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**4,x)`

output

```
Piecewise((-3360*A*d*x*tan(c/2 + d*x/2)**4/(840*a**4*d*tan(c/2 + d*x/2)**4
+ 1680*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 6720*A*d*x*tan(c/2 + d*
x/2)**2/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**2
+ 840*a**4*d) - 3360*A*d*x/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*t
an(c/2 + d*x/2)**2 + 840*a**4*d) - 15*A*tan(c/2 + d*x/2)**11/(840*a**4*d*t
an(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 117*A
*tan(c/2 + d*x/2)**9/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2
+ d*x/2)**2 + 840*a**4*d) - 526*A*tan(c/2 + d*x/2)**7/(840*a**4*d*tan(c/2
+ d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 3682*A*tan(
c/2 + d*x/2)**5/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*
x/2)**2 + 840*a**4*d) + 11165*A*tan(c/2 + d*x/2)**3/(840*a**4*d*tan(c/2 +
d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 6825*A*tan(c/2
+ d*x/2)/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**
2 + 840*a**4*d) + 8820*B*d*x*tan(c/2 + d*x/2)**4/(840*a**4*d*tan(c/2 + d*x
/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 17640*B*d*x*tan(c
/2 + d*x/2)**2/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x
/2)**2 + 840*a**4*d) + 8820*B*d*x/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a
**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 15*B*tan(c/2 + d*x/2)**11/(840*a
**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d)
- 159*B*tan(c/2 + d*x/2)**9/(840*a**4*d*tan(c/2 + d*x/2)**4 + 1680*a**4...
```

3.65.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.59

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx =$$

$$3B \left(\frac{280 \left(\frac{7 \sin(dx+c)}{\cos(dx+c)+1} + \frac{9 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^4 + \frac{2a^4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^4 \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} + \frac{3885 \sin(dx+c)}{\cos(dx+c)+1} - \frac{455 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{63 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{5 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} - \frac{5880 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)}\right)}{a^4} \right)$$

input

```
integrate(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="m
axima")
```

3.65. $\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

output
$$\begin{aligned} & -1/840*(3*B*(280*(7*\sin(d*x + c))/(\cos(d*x + c) + 1) + 9*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(a^4 + 2*a^4*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + a^4*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4) + (3885*\sin(d*x + c))/(\cos(d*x + c) + 1) - 455*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 63*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 - 5*\sin(d*x + c)^7/(\cos(d*x + c) + 1)^7/a^4 - 5880*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1))/a^4 - A*(1680*\sin(d*x + c)/((a^4 + a^4*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2*(\cos(d*x + c) + 1)) + (5145*\sin(d*x + c)/(\cos(d*x + c) + 1) - 805*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 147*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 - 15*\sin(d*x + c)^7/(\cos(d*x + c) + 1)^7)/a^4 - 6720*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1))/a^4)/d \end{aligned}$$

3.65.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.02

$$\int \frac{\cos^5(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx = \frac{420(dx+c)(8A-21B)}{a^4} - \frac{840\left(2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 9B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + 2A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 7B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)\right)}{\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 1\right)^2 a^4} + \frac{15Aa^{24} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{a^4}$$

input `integrate(cos(d*x+c)^5*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="giac")`

output
$$\begin{aligned} & -1/840*(420*(d*x + c)*(8*A - 21*B)/a^4 - 840*(2*A*\tan(1/2*d*x + 1/2*c)^3 - 9*B*\tan(1/2*d*x + 1/2*c)^3 + 2*A*\tan(1/2*d*x + 1/2*c) - 7*B*\tan(1/2*d*x + 1/2*c))/((\tan(1/2*d*x + 1/2*c)^2 + 1)^2*a^4) + (15*A*a^24*\tan(1/2*d*x + 1/2*c)^7 - 15*B*a^24*\tan(1/2*d*x + 1/2*c)^7 - 147*A*a^24*\tan(1/2*d*x + 1/2*c)^5 + 189*B*a^24*\tan(1/2*d*x + 1/2*c)^5 + 805*A*a^24*\tan(1/2*d*x + 1/2*c)^3 - 1365*B*a^24*\tan(1/2*d*x + 1/2*c)^3 - 5145*A*a^24*\tan(1/2*d*x + 1/2*c) + 11655*B*a^24*\tan(1/2*d*x + 1/2*c))/a^28)/d \end{aligned}$$

3.65.9 Mupad [B] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.13

$$\int \frac{\cos^5(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{5(A-B)}{4a^4} - \frac{5B}{2a^4} + \frac{3(4A-6B)}{4a^4} + \frac{3(5A-15B)}{8a^4}\right)}{d}$$

$$- \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{4a^4} + \frac{4A-6B}{8a^4} + \frac{5A-15B}{24a^4}\right)}{d} - \frac{x(8A-21B)}{2a^4}$$

$$+ \frac{(2A-9B)\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (2A-7B)\tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 2a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^4\right)}$$

$$+ \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 \left(\frac{3(A-B)}{40a^4} + \frac{4A-6B}{40a^4}\right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 (A-B)}{56a^4 d}$$

input `int((cos(c + d*x))^5*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^4,x)`output `(tan(c/2 + (d*x)/2)*((5*(A - B))/(4*a^4) - (5*B)/(2*a^4) + (3*(4*A - 6*B))/(4*a^4) + (3*(5*A - 15*B))/(8*a^4)))/d - (tan(c/2 + (d*x)/2)^3*((A - B)/(4*a^4) + (4*A - 6*B)/(8*a^4) + (5*A - 15*B)/(24*a^4)))/d - (x*(8*A - 21*B))/(2*a^4) + (tan(c/2 + (d*x)/2)^3*(2*A - 9*B) + tan(c/2 + (d*x)/2)*(2*A - 7*B))/(d*(2*a^4*tan(c/2 + (d*x)/2)^2 + a^4*tan(c/2 + (d*x)/2)^4 + a^4)) + (tan(c/2 + (d*x)/2)^5*((3*(A - B))/(40*a^4) + (4*A - 6*B)/(40*a^4)))/d - (tan(c/2 + (d*x)/2)^7*(A - B))/(56*a^4*d)`

3.66
$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$$

3.66.1 Optimal result 751
 3.66.2 Mathematica [B] (verified) 752
 3.66.3 Rubi [A] (verified) 752
 3.66.4 Maple [A] (verified) 757
 3.66.5 Fricas [A] (verification not implemented) 757
 3.66.6 Sympy [B] (verification not implemented) 758
 3.66.7 Maxima [A] (verification not implemented) 759
 3.66.8 Giac [A] (verification not implemented) 759
 3.66.9 Mupad [B] (verification not implemented) 760

3.66.1 Optimal result

Integrand size = 31, antiderivative size = 185

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = \frac{(A-4B)x}{a^4} - \frac{(55A-244B) \sin(c+dx)}{105a^4d} + \frac{(25A-88B) \cos^2(c+dx) \sin(c+dx)}{105a^4d(1+\cos(c+dx))^2} - \frac{(A-4B) \sin(c+dx)}{a^4d(1+\cos(c+dx))} + \frac{(A-B) \cos^4(c+dx) \sin(c+dx)}{7d(a+a \cos(c+dx))^4} + \frac{(5A-12B) \cos^3(c+dx) \sin(c+dx)}{35ad(a+a \cos(c+dx))^3}$$

```
output (A-4*B)*x/a^4-1/105*(55*A-244*B)*sin(d*x+c)/a^4/d+1/105*(25*A-88*B)*cos(d*
x+c)^2*sin(d*x+c)/a^4/d/(1+cos(d*x+c))^2-(A-4*B)*sin(d*x+c)/a^4/d/(1+cos(d
*x+c))+1/7*(A-B)*cos(d*x+c)^4*sin(d*x+c)/d/(a+a*cos(d*x+c))^4+1/35*(5*A-12
*B)*cos(d*x+c)^3*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```

3.66.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 481 vs. $2(185) = 370$.

Time = 4.77 (sec) , antiderivative size = 481, normalized size of antiderivative = 2.60

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(7350(A-4B)dx \cos\left(\frac{dx}{2}\right) + 7350(A-4B)dx \cos\left(c+\frac{dx}{2}\right) + 4410Adx \cos\left(c+\frac{3d}{2}\right)\right)}{(1680a^4d(1+\cos(c+dx))^4)}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(7350*(A - 4*B)*d*x*Cos[(d*x)/2] + 7350*(A - 4*B)*d*x*Cos[c + (d*x)/2] + 4410*A*d*x*Cos[c + (3*d*x)/2] - 17640*B*d*x*Cos[c + (3*d*x)/2] + 4410*A*d*x*Cos[2*c + (3*d*x)/2] - 17640*B*d*x*Cos[2*c + (3*d*x)/2] + 1470*A*d*x*Cos[2*c + (5*d*x)/2] - 5880*B*d*x*Cos[2*c + (5*d*x)/2] + 1470*A*d*x*Cos[3*c + (5*d*x)/2] - 5880*B*d*x*Cos[3*c + (5*d*x)/2] + 210*A*d*x*Cos[3*c + (7*d*x)/2] - 840*B*d*x*Cos[3*c + (7*d*x)/2] + 210*A*d*x*Cos[4*c + (7*d*x)/2] - 840*B*d*x*Cos[4*c + (7*d*x)/2] - 19880*A*Sin[(d*x)/2] + 60830*B*Sin[(d*x)/2] + 16520*A*Sin[c + (d*x)/2] - 46130*B*Sin[c + (d*x)/2] - 14280*A*Sin[c + (3*d*x)/2] + 46116*B*Sin[c + (3*d*x)/2] + 7560*A*Sin[2*c + (3*d*x)/2] - 18060*B*Sin[2*c + (3*d*x)/2] - 5600*A*Sin[2*c + (5*d*x)/2] + 19292*B*Sin[2*c + (5*d*x)/2] + 1680*A*Sin[3*c + (5*d*x)/2] - 21000*B*Sin[3*c + (5*d*x)/2] - 1040*A*Sin[3*c + (7*d*x)/2] + 3791*B*Sin[3*c + (7*d*x)/2] + 735*B*Sin[4*c + (7*d*x)/2] + 105*B*Sin[4*c + (9*d*x)/2] + 105*B*Sin[5*c + (9*d*x)/2]))/(1680*a^4*d*(1 + Cos[c + d*x])^4)`

3.66.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.09, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3456, 3042, 3456, 3042, 3456, 3042, 3447, 3042, 3502, 27, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx$$

3.66. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^4 (A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^4} dx \\
& \downarrow 3456 \\
& \frac{\int \frac{\cos^3(c+dx)(4a(A-B)-a(A-8B)\cos(c+dx))}{(\cos(c+dx)a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^3 (4a(A-B)-a(A-8B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \downarrow 3456 \\
& \frac{\int \frac{\cos^2(c+dx)(3a^2(5A-12B)-2a^2(5A-26B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2 (3a^2(5A-12B)-2a^2(5A-26B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \downarrow 3456 \\
& \frac{\int \frac{\cos(c+dx)(2a^3(25A-88B)-a^3(55A-244B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{(25A-88B)\sin(c+dx)\cos^2(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(2a^3(25A-88B)-a^3(55A-244B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{3a^2} + \frac{(25A-88B)\sin(c+dx)\cos^2(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx)
\end{aligned}$$

3.66. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

↓ 3447

$$\frac{\int \frac{2a^3(25A-88B) \cos(c+dx) - a^3(55A-244B) \cos^2(c+dx)}{3a^2} dx + \frac{(25A-88B) \sin(c+dx) \cos^2(c+dx)}{3d(\cos(c+dx)+1)^2}}{5a^2} + \frac{a(5A-12B) \sin(c+dx) \cos^3(c+dx)}{5d(a \cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a \cos(c+dx)+a)^4} (A-B) \sin(c+dx) \cos^4(c+dx)$$

↓ 3042

$$\frac{\int \frac{2a^3(25A-88B) \sin(c+dx+\frac{\pi}{2}) - a^3(55A-244B) \sin(c+dx+\frac{\pi}{2})^2}{3a^2} dx + \frac{(25A-88B) \sin(c+dx) \cos^2(c+dx)}{3d(\cos(c+dx)+1)^2}}{5a^2} + \frac{a(5A-12B) \sin(c+dx) \cos^3(c+dx)}{5d(a \cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a \cos(c+dx)+a)^4} (A-B) \sin(c+dx) \cos^4(c+dx)$$

↓ 3502

$$\frac{\int \frac{105a^4(A-4B) \cos(c+dx)}{3a^2} dx - \frac{a^2(55A-244B) \sin(c+dx)}{d} + \frac{(25A-88B) \sin(c+dx) \cos^2(c+dx)}{3d(\cos(c+dx)+1)^2}}{5a^2} + \frac{a(5A-12B) \sin(c+dx) \cos^3(c+dx)}{5d(a \cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a \cos(c+dx)+a)^4} (A-B) \sin(c+dx) \cos^4(c+dx)$$

↓ 27

$$\frac{105a^3(A-4B) \int \frac{\cos(c+dx)}{3a^2} dx - \frac{a^2(55A-244B) \sin(c+dx)}{d} + \frac{(25A-88B) \sin(c+dx) \cos^2(c+dx)}{3d(\cos(c+dx)+1)^2}}{5a^2} + \frac{a(5A-12B) \sin(c+dx) \cos^3(c+dx)}{5d(a \cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a \cos(c+dx)+a)^4} (A-B) \sin(c+dx) \cos^4(c+dx)$$

↓ 3042

$$\frac{105a^3(A-4B) \int \frac{\sin(c+dx+\frac{\pi}{2})}{3a^2} dx - \frac{a^2(55A-244B) \sin(c+dx)}{d} + \frac{(25A-88B) \sin(c+dx) \cos^2(c+dx)}{3d(\cos(c+dx)+1)^2}}{5a^2} + \frac{a(5A-12B) \sin(c+dx) \cos^3(c+dx)}{5d(a \cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a \cos(c+dx)+a)^4} (A-B) \sin(c+dx) \cos^4(c+dx)$$

↓ 3214

3.66. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

$$\frac{\frac{105a^3(A-4B)\left(\frac{x}{a} - \int \frac{1}{\cos(c+dx)a+a} dx\right) - \frac{a^2(55A-244B)\sin(c+dx)}{d}}{3a^2}}{5a^2} + \frac{(25A-88B)\sin(c+dx)\cos^2(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx)$$

↓ 3042

$$\frac{105a^3(A-4B)\left(\frac{x}{a} - \int \frac{1}{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} dx\right) - \frac{a^2(55A-244B)\sin(c+dx)}{d}}{3a^2}}{5a^2} + \frac{(25A-88B)\sin(c+dx)\cos^2(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx)$$

↓ 3127

$$\frac{105a^3(A-4B)\left(\frac{x}{a} - \frac{\sin(c+dx)}{d(a\cos(c+dx)+a)}\right) - \frac{a^2(55A-244B)\sin(c+dx)}{d}}{3a^2}}{5a^2} + \frac{(25A-88B)\sin(c+dx)\cos^2(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(5A-12B)\sin(c+dx)\cos^3(c+dx)}{5d(a\cos(c+dx)+a)^3} +$$

$$\frac{7a^2}{7d(a\cos(c+dx)+a)^4} (A-B)\sin(c+dx)\cos^4(c+dx)$$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `((A - B)*Cos[c + d*x]^4*Sin[c + d*x])/(7*d*(a + a*Cos[c + d*x])^4) + ((a*(5*A - 12*B)*Cos[c + d*x]^3*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((25*A - 88*B)*Cos[c + d*x]^2*Sin[c + d*x])/(3*d*(1 + Cos[c + d*x])^2) + (-((a^2*(55*A - 244*B)*Sin[c + d*x])/d) + 105*a^3*(A - 4*B)*(x/a - Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))) / (3*a^2) / (5*a^2) / (7*a^2)`

3.66.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.66. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.66.4 Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.57

method	result
parallelrisc	$-5840 \left(\left(\frac{31A}{73} - \frac{2741B}{1460} \right) \cos(2dx+2c) + \left(\frac{13A}{146} - \frac{148B}{365} \right) \cos(3dx+3c) - \frac{21B \cos(4dx+4c)}{1168} + \left(A - \frac{1562B}{365} \right) \cos(dx+c) + \frac{47A}{73} - \frac{16B}{5} \right)$
derivativedivides	$\frac{(\tan^7(\frac{dx}{2} + \frac{c}{2}))A - (\tan^7(\frac{dx}{2} + \frac{c}{2}))B}{7} - A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{7(\tan^5(\frac{dx}{2} + \frac{c}{2}))B}{5} + \frac{11(\tan^3(\frac{dx}{2} + \frac{c}{2}))A}{3} - \frac{23(\tan^3(\frac{dx}{2} + \frac{c}{2}))B}{3}$
default	$\frac{(\tan^7(\frac{dx}{2} + \frac{c}{2}))A - (\tan^7(\frac{dx}{2} + \frac{c}{2}))B}{7} - A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{7(\tan^5(\frac{dx}{2} + \frac{c}{2}))B}{5} + \frac{11(\tan^3(\frac{dx}{2} + \frac{c}{2}))A}{3} - \frac{23(\tan^3(\frac{dx}{2} + \frac{c}{2}))B}{3}$
risc	$\frac{x A}{a^4} - \frac{4 B x}{a^4} - \frac{i e^{i(dx+c)} B}{2 a^4 d} + \frac{i e^{-i(dx+c)} B}{2 a^4 d} - \frac{4 i (210 A e^{6 i(dx+c)} - 525 B e^{6 i(dx+c)} + 945 A e^{5 i(dx+c)} - 2625 B e^{5 i(dx+c)})}{8 d a^4}$
norman	$\frac{(A-4B)x}{a} + \frac{(A-4B)x \left(\tan^{10}\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a} + \frac{(A-22B) \left(\tan^{13}\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{84 a d} + \frac{5(A-4B)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a} + \frac{10(A-4B)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a}$

input `int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBOSE)`

output `1/6720*(-5840*((31/73*A-2741/1460*B)*cos(2*d*x+2*c)+(13/146*A-148/365*B)*cos(3*d*x+3*c)-21/1168*B*cos(4*d*x+4*c)+(A-1562/365*B)*cos(d*x+c)+47/73*A-16171/5840*B)*tan(1/2*d*x+1/2*c)*sec(1/2*d*x+1/2*c)^6+6720*d*x*(A-4*B))/a^4/d`

3.66.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.15

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$$

$$= \frac{105(A-4B)dx \cos(dx+c)^4 + 420(A-4B)dx \cos(dx+c)^3 + 630(A-4B)dx \cos(dx+c)^2 + 420(A-4B)dx \cos(dx+c) + 105(A-4B)dx}{(a+a \cos(dx+c))^4}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="fricas")`

```
output 1/105*(105*(A - 4*B)*d*x*cos(d*x + c)^4 + 420*(A - 4*B)*d*x*cos(d*x + c)^3
+ 630*(A - 4*B)*d*x*cos(d*x + c)^2 + 420*(A - 4*B)*d*x*cos(d*x + c) + 105
*(A - 4*B)*d*x + (105*B*cos(d*x + c)^4 - 4*(65*A - 296*B)*cos(d*x + c)^3 -
4*(155*A - 659*B)*cos(d*x + c)^2 - (535*A - 2236*B)*cos(d*x + c) - 160*A
+ 664*B)*sin(d*x + c))/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*
a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)
```

3.66.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 578 vs. $2(172) = 344$.

Time = 5.07 (sec) , antiderivative size = 578, normalized size of antiderivative = 3.12

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \left\{ \begin{array}{l} \frac{840Adx \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{840a^4d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 840a^4d} + \frac{840Adx}{840a^4d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 840a^4d} + \frac{15A \tan^9\left(\frac{c}{2} + \frac{dx}{2}\right)}{840a^4d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 840a^4d} - \frac{90A \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{840a^4d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 840a^4d} + \dots \\ \frac{x(A+B\cos(c))\cos^4(c)}{(a\cos(c)+a)^4} \end{array} \right.$$

```
input integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**4,x)
```

```
output Piecewise((840*A*d*x*tan(c/2 + d*x/2)**2/(840*a**4*d*tan(c/2 + d*x/2)**2 +
840*a**4*d) + 840*A*d*x/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 1
5*A*tan(c/2 + d*x/2)**9/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 90
*A*tan(c/2 + d*x/2)**7/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 280
*A*tan(c/2 + d*x/2)**5/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 119
0*A*tan(c/2 + d*x/2)**3/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 15
75*A*tan(c/2 + d*x/2)/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 3360
*B*d*x*tan(c/2 + d*x/2)**2/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) -
3360*B*d*x/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 15*B*tan(c/2 +
d*x/2)**9/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 132*B*tan(c/2 +
d*x/2)**7/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) - 658*B*tan(c/2 +
d*x/2)**5/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 4340*B*tan(c/2
+ d*x/2)**3/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d) + 6825*B*tan(c/2
+ d*x/2)/(840*a**4*d*tan(c/2 + d*x/2)**2 + 840*a**4*d), Ne(d, 0)), (x*(A
+ B*cos(c))*cos(c)**4/(a*cos(c) + a)**4, True))
```

3.66.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.46

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{B \left(\frac{1680 \sin(dx+c)}{\left(a^4 + \frac{a^4 \sin^2(dx+c)}{(\cos(dx+c)+1)^2}\right) (\cos(dx+c)+1)} + \frac{5145 \sin(dx+c)}{\cos(dx+c)+1} - \frac{805 \sin^3(dx+c)}{(\cos(dx+c)+1)^3} + \frac{147 \sin^5(dx+c)}{(\cos(dx+c)+1)^5} - \frac{15 \sin^7(dx+c)}{(\cos(dx+c)+1)^7} - \frac{6720 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^4} \right)}{840 d}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="maxima")`

output `1/840*(B*(1680*sin(d*x + c)/((a^4 + a^4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) + (5145*sin(d*x + c)/(cos(d*x + c) + 1) - 805*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 147*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 15*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4 - 6720*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^4 - 5*A*((315*sin(d*x + c)/(cos(d*x + c) + 1) - 77*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 21*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 3*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4 - 336*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/a^4)/d`

3.66.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.02

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{840(dx+c)(A-4B)}{a^4} + \frac{1680B \tan(\frac{1}{2}dx + \frac{1}{2}c)}{(\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + 1)a^4} + \frac{15Aa^{24} \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 15Ba^{24} \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 105Aa^{24} \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 147Ba^{24} \tan(\frac{1}{2}dx + \frac{1}{2}c)^3}{840 d}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="giac")`

output $\frac{1}{840} \cdot (840 \cdot (d \cdot x + c) \cdot (A - 4 \cdot B) / a^4 + 1680 \cdot B \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) / ((\tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 1) \cdot a^4) + (15 \cdot A \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^7 - 15 \cdot B \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^7 - 105 \cdot A \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 + 147 \cdot B \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^5 + 385 \cdot A \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 - 805 \cdot B \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 - 1575 \cdot A \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + 5145 \cdot B \cdot a^{24} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)) / a^{28}) / d$

3.66.9 Mupad [B] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.09

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx = \frac{A dx - 4 B dx}{a^4 d}$$

$$- \frac{\left(\frac{52 A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{21} - \frac{764 B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{105} \right) \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + \left(\frac{143 B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{105} - \frac{16 A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{21} \right) \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + a^4 d \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^7}{a^4 d \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^7}$$

$$+ \frac{2 B \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{a^4 d}$$

input `int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^4,x)`

output $(A \cdot d \cdot x - 4 \cdot B \cdot d \cdot x) / (a^4 \cdot d) - ((B \cdot \sin(c/2 + (d \cdot x)/2)) / 56 - (A \cdot \sin(c/2 + (d \cdot x)/2)) / 56 + \cos(c/2 + (d \cdot x)/2)^2 \cdot ((5 \cdot A \cdot \sin(c/2 + (d \cdot x)/2)) / 28 - (8 \cdot B \cdot \sin(c/2 + (d \cdot x)/2)) / 35) - \cos(c/2 + (d \cdot x)/2)^4 \cdot ((16 \cdot A \cdot \sin(c/2 + (d \cdot x)/2)) / 21 - (143 \cdot B \cdot \sin(c/2 + (d \cdot x)/2)) / 105) + \cos(c/2 + (d \cdot x)/2)^6 \cdot ((52 \cdot A \cdot \sin(c/2 + (d \cdot x)/2)) / 21 - (764 \cdot B \cdot \sin(c/2 + (d \cdot x)/2)) / 105)) / (a^4 \cdot d \cdot \cos(c/2 + (d \cdot x)/2)^7) + (2 \cdot B \cdot \cos(c/2 + (d \cdot x)/2) \cdot \sin(c/2 + (d \cdot x)/2)) / (a^4 \cdot d)$

3.67 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

3.67.1	Optimal result	761
3.67.2	Mathematica [B] (verified)	762
3.67.3	Rubi [A] (verified)	762
3.67.4	Maple [A] (verified)	766
3.67.5	Fricas [A] (verification not implemented)	766
3.67.6	Sympy [A] (verification not implemented)	767
3.67.7	Maxima [A] (verification not implemented)	767
3.67.8	Giac [A] (verification not implemented)	768
3.67.9	Mupad [B] (verification not implemented)	768

3.67.1 Optimal result

Integrand size = 31, antiderivative size = 154

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = \frac{Bx}{a^4} - \frac{(6A-55B) \sin(c+dx)}{105a^4d(1+\cos(c+dx))^2} + \frac{(12A-215B) \sin(c+dx)}{105a^4d(1+\cos(c+dx))} + \frac{(A-B) \cos^3(c+dx) \sin(c+dx)}{7d(a+a \cos(c+dx))^4} + \frac{(3A-10B) \cos^2(c+dx) \sin(c+dx)}{35ad(a+a \cos(c+dx))^3}$$

```
output B*x/a^4-1/105*(6*A-55*B)*sin(d*x+c)/a^4/d/(1+cos(d*x+c))^2+1/105*(12*A-215*B)*sin(d*x+c)/a^4/d/(1+cos(d*x+c))+1/7*(A-B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^4+1/35*(3*A-10*B)*cos(d*x+c)^2*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```


3.67.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 329 vs. $2(154) = 308$.

Time = 4.37 (sec) , antiderivative size = 329, normalized size of antiderivative = 2.14

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{\sec\left(\frac{c}{2}\right) \sec^7\left(\frac{1}{2}(c+dx)\right) \left(3675Bdx \cos\left(\frac{dx}{2}\right) + 3675Bdx \cos\left(c+\frac{dx}{2}\right) + 2205Bdx \cos\left(c+\frac{3dx}{2}\right) + 2205Bdx\right)}{(a+a\cos(c+dx))^4}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `(Sec[c/2]*Sec[(c + d*x)/2]^7*(3675*B*d*x*Cos[(d*x)/2] + 3675*B*d*x*Cos[c + (d*x)/2] + 2205*B*d*x*Cos[c + (3*d*x)/2] + 2205*B*d*x*Cos[2*c + (3*d*x)/2] + 735*B*d*x*Cos[2*c + (5*d*x)/2] + 735*B*d*x*Cos[3*c + (5*d*x)/2] + 105*B*d*x*Cos[3*c + (7*d*x)/2] + 105*B*d*x*Cos[4*c + (7*d*x)/2] + 1260*A*Sin[(d*x)/2] - 9940*B*Sin[(d*x)/2] - 1260*A*Sin[c + (d*x)/2] + 8260*B*Sin[c + (d*x)/2] + 882*A*Sin[c + (3*d*x)/2] - 7140*B*Sin[c + (3*d*x)/2] - 630*A*Sin[2*c + (3*d*x)/2] + 3780*B*Sin[2*c + (3*d*x)/2] + 294*A*Sin[2*c + (5*d*x)/2] - 2800*B*Sin[2*c + (5*d*x)/2] - 210*A*Sin[3*c + (5*d*x)/2] + 840*B*Sin[3*c + (5*d*x)/2] + 72*A*Sin[3*c + (7*d*x)/2] - 520*B*Sin[3*c + (7*d*x)/2])/((13440*a^4*d)`

3.67.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.12, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3456, 3042, 3456, 3042, 3447, 3042, 3498, 25, 3042, 3214, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^3 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a\sin\left(c+dx+\frac{\pi}{2}\right)+a)^4} dx$$

$$\downarrow \text{3456}$$

3.67. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

$$\begin{aligned}
& \frac{\int \frac{\cos^2(c+dx)(3a(A-B)+7aB \cos(c+dx))}{(\cos(c+dx)a+a)^3} dx}{7a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(3a(A-B)+7aB \sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos(c+dx)(2(3A-10B)a^2+35B \cos(c+dx)a^2)}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a \cos(c+dx) + a)^4} (A-B) \sin(c+dx) \cos^3(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(2(3A-10B)a^2+35B \sin(c+dx+\frac{\pi}{2})a^2)}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a \cos(c+dx) + a)^4} (A-B) \sin(c+dx) \cos^3(c+dx) \\
& \quad \downarrow \text{3447} \\
& \frac{\int \frac{35B \cos^2(c+dx)a^2+2(3A-10B) \cos(c+dx)a^2}{(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a \cos(c+dx) + a)^4} (A-B) \sin(c+dx) \cos^3(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{35B \sin(c+dx+\frac{\pi}{2})^2 a^2+2(3A-10B) \sin(c+dx+\frac{\pi}{2})a^2}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a \cos(c+dx) + a)^4} (A-B) \sin(c+dx) \cos^3(c+dx) \\
& \quad \downarrow \text{3498} \\
& \frac{\int -\frac{2(6A-55B)a^3+105B \cos(c+dx)a^3}{\cos(c+dx)a+a} dx}{3a^2} - \frac{(6A-55B) \sin(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3} + \\
& \quad \frac{7a^2}{7d(a \cos(c+dx) + a)^4} (A-B) \sin(c+dx) \cos^3(c+dx) \\
& \quad \downarrow \text{25}
\end{aligned}$$

3.67. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

$$\begin{aligned}
& \frac{\int \frac{2(6A-55B)a^3+105B \cos(c+dx)a^3}{\cos(c+dx)a+a} dx - \frac{(6A-55B) \sin(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2} \\
& \quad + \frac{7a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
& \quad \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2(6A-55B)a^3+105B \sin(c+dx+\frac{\pi}{2})a^3}{\sin(c+dx+\frac{\pi}{2})a+a} dx - \frac{(6A-55B) \sin(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2} \\
& \quad + \frac{7a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
& \quad \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3214} \\
& \frac{a^3(12A-215B) \int \frac{1}{\cos(c+dx)a+a} dx + 105a^2 Bx - \frac{(6A-55B) \sin(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2} \\
& \quad + \frac{7a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
& \quad \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{a^3(12A-215B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx + 105a^2 Bx - \frac{(6A-55B) \sin(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2} \\
& \quad + \frac{7a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
& \quad \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3127} \\
& \frac{\frac{a^3(12A-215B) \sin(c+dx)}{d(a \cos(c+dx)+a)} + 105a^2 Bx - \frac{(6A-55B) \sin(c+dx)}{3d(\cos(c+dx)+1)^2} + \frac{a(3A-10B) \sin(c+dx) \cos^2(c+dx)}{5d(a \cos(c+dx)+a)^3}}{5a^2} \\
& \quad + \frac{7a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
& \quad \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{7d(a \cos(c+dx)+a)^4}
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `((A - B)*Cos[c + d*x]^3*Sin[c + d*x])/(7*d*(a + a*Cos[c + d*x])^4) + ((a*(3*A - 10*B)*Cos[c + d*x]^2*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + (-1/3*((6*A - 55*B)*Sin[c + d*x])/(d*(1 + Cos[c + d*x])^2) + (105*a^2*B*x + (a^3*(12*A - 215*B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])))/(3*a^2))/(5*a^2)))/(7*a^2)`

$$3.67. \quad \int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$$

3.67.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3498 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b - a*B + b*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m/(a*f*(2*m + 1)), x] + Simp[1/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]`

3.67.4 Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.58

method	result
parallelrisc	$\frac{(-15A+15B)\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(63A-105B)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(-105A+385B)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(105A-1575B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{840a^4d}$
derivativedivides	$-\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{7}+\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{7}+\frac{3A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}-\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B-\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A+\frac{11\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{3}$
default	$-\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{7}+\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{7}+\frac{3A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}-\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B-\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A+\frac{11\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{3}$
risc	$\frac{Bx}{a^4}+\frac{2i(105Ae^{6i(dx+c)}-420Be^{6i(dx+c)}+315Ae^{5i(dx+c)}-1890Be^{5i(dx+c)}+630Ae^{4i(dx+c)}-4130Be^{4i(dx+c)}+630Ae^{3i(dx+c)}-105d^4(e^{i(dx+c)}-1))}{105d^4(e^{i(dx+c)}-1)}$
norman	$\frac{Bx}{a}+\frac{Bx\left(\tan^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{4Bx\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{6Bx\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{4Bx\left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}+\frac{(A-15B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{8ad}+\frac{(A-15B)\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)}{8ad}$

input `int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBOSE)`

output `1/840*((-15*A+15*B)*tan(1/2*d*x+1/2*c)^7+(63*A-105*B)*tan(1/2*d*x+1/2*c)^5+(-105*A+385*B)*tan(1/2*d*x+1/2*c)^3+(105*A-1575*B)*tan(1/2*d*x+1/2*c)+840*d*x*B)/a^4/d`

3.67.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.17

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{105 Bdx \cos(dx+c)^4 + 420 Bdx \cos(dx+c)^3 + 630 Bdx \cos(dx+c)^2 + 420 Bdx \cos(dx+c) + 105 Bdx}{105 (a^4d \cos(dx+c))^4 + 4 a^4d \cos(dx+c)}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="fracas")`

output $1/105*(105*B*d*x*cos(d*x + c)^4 + 420*B*d*x*cos(d*x + c)^3 + 630*B*d*x*cos(d*x + c)^2 + 420*B*d*x*cos(d*x + c) + 105*B*d*x + (4*(9*A - 65*B)*cos(d*x + c)^3 + (39*A - 620*B)*cos(d*x + c)^2 + (24*A - 535*B)*cos(d*x + c) + 6*A - 160*B)*sin(d*x + c))/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)$

3.67.6 Sympy [A] (verification not implemented)

Time = 3.24 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.25

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx$$

$$= \begin{cases} -\frac{A \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{56a^4d} + \frac{3A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{40a^4d} - \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} + \frac{Bx}{a^4} + \frac{B \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{56a^4d} - \frac{B \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} + \frac{11B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} - \frac{11B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} \\ \frac{x(A+B \cos(c)) \cos^3(c)}{(a \cos(c)+a)^4} \end{cases}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**4,x)`

output `Piecewise((-A*tan(c/2 + d*x/2)**7/(56*a**4*d) + 3*A*tan(c/2 + d*x/2)**5/(40*a**4*d) - A*tan(c/2 + d*x/2)**3/(8*a**4*d) + A*tan(c/2 + d*x/2)/(8*a**4*d) + B*x/a**4 + B*tan(c/2 + d*x/2)**7/(56*a**4*d) - B*tan(c/2 + d*x/2)**5/(8*a**4*d) + 11*B*tan(c/2 + d*x/2)**3/(24*a**4*d) - 15*B*tan(c/2 + d*x/2)/(8*a**4*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**3/(a*cos(c) + a)**4, True))`

3.67.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.31

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx =$$

$$\frac{5B \left(\frac{315 \sin(dx+c)}{\cos(dx+c)+1} - \frac{77 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{3 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) - \frac{336 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^4} - 3A \left(\frac{35 \sin(dx+c)}{\cos(dx+c)+1} - \frac{35 \sin(dx+c)}{(\cos(dx+c)+1)} \right)}{840d}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="maxima")`

3.67. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

output
$$-1/840*(5*B*((315*\sin(dx + c)/(\cos(dx + c) + 1) - 77*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 21*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 3*\sin(dx + c)^7/(\cos(dx + c) + 1)^7)/a^4 - 336*\arctan(\sin(dx + c)/(\cos(dx + c) + 1))/a^4 - 3*A*(35*\sin(dx + c)/(\cos(dx + c) + 1) - 35*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 21*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 5*\sin(dx + c)^7/(\cos(dx + c) + 1)^7)/a^4)/d$$

3.67.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.01

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{840(dx+c)B}{a^4} - \frac{15Aa^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)^7 - 15Ba^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)^7 - 63Aa^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)^5 + 105Ba^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)^5 + 105Aa^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)^3 - 385Ba^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)^3 - 105Aa^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c) + 1575Ba^{24}\tan(\frac{1}{2}dx+\frac{1}{2}c)}{a^{28}}$$

$840 d$

input `integrate(cos(dx+c)^3*(A+B*cos(dx+c))/(a+a*cos(dx+c))^4,x, algorithm="giac")`

output
$$1/840*(840*(dx + c)*B/a^4 - (15*A*a^{24}*\tan(1/2*dx + 1/2*c)^7 - 15*B*a^{24}*\tan(1/2*dx + 1/2*c)^7 - 63*A*a^{24}*\tan(1/2*dx + 1/2*c)^5 + 105*B*a^{24}*\tan(1/2*dx + 1/2*c)^5 + 105*A*a^{24}*\tan(1/2*dx + 1/2*c)^3 - 385*B*a^{24}*\tan(1/2*dx + 1/2*c)^3 - 105*A*a^{24}*\tan(1/2*dx + 1/2*c) + 1575*B*a^{24}*\tan(1/2*dx + 1/2*c))/a^{28})/d$$

3.67.9 Mupad [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.05

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx = \frac{Bx}{a^4}$$

$$+ \frac{\left(\frac{12A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{35} - \frac{52B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{21}\right) \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + \left(\frac{16B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{21} - \frac{23A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{70}\right) \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + \left(\frac{12A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{35} - \frac{52B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{21}\right) \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \left(\frac{16B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{21} - \frac{23A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{70}\right) \cos\left(\frac{c}{2} + \frac{dx}{2}\right)}{a^4 d \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^7}$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^4,x)`

output $(B*x)/a^4 + ((B*\sin(c/2 + (d*x)/2))/56 - (A*\sin(c/2 + (d*x)/2))/56 + \cos(c/2 + (d*x)/2)^2*((9*A*\sin(c/2 + (d*x)/2))/70 - (5*B*\sin(c/2 + (d*x)/2))/28) + \cos(c/2 + (d*x)/2)^6*((12*A*\sin(c/2 + (d*x)/2))/35 - (52*B*\sin(c/2 + (d*x)/2))/21) - \cos(c/2 + (d*x)/2)^4*((23*A*\sin(c/2 + (d*x)/2))/70 - (16*B*\sin(c/2 + (d*x)/2))/21))/(a^4*d*\cos(c/2 + (d*x)/2)^7)$

3.68
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$$

3.68.1	Optimal result	770
3.68.2	Mathematica [A] (verified)	770
3.68.3	Rubi [A] (verified)	771
3.68.4	Maple [A] (verified)	774
3.68.5	Fricas [A] (verification not implemented)	774
3.68.6	Sympy [A] (verification not implemented)	775
3.68.7	Maxima [A] (verification not implemented)	775
3.68.8	Giac [A] (verification not implemented)	776
3.68.9	Mupad [B] (verification not implemented)	776

3.68.1 Optimal result

Integrand size = 31, antiderivative size = 136

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = -\frac{2(A+27B) \sin(c+dx)}{105a^4d(1+\cos(c+dx))^2} + \frac{(13A+36B) \sin(c+dx)}{105a^4d(1+\cos(c+dx))} + \frac{(A-B) \cos^2(c+dx) \sin(c+dx)}{7d(a+a \cos(c+dx))^4} - \frac{(A-8B) \sin(c+dx)}{35ad(a+a \cos(c+dx))^3}$$

```
output -2/105*(A+27*B)*sin(d*x+c)/a^4/d/(1+cos(d*x+c))^2+1/105*(13*A+36*B)*sin(d*x+c)/a^4/d/(1+cos(d*x+c))+1/7*(A-B)*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^4-1/35*(A-8*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```

3.68.2 Mathematica [A] (verified)

Time = 3.09 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.42

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = \frac{\cos\left(\frac{1}{2}(c+dx)\right) \sec\left(\frac{c}{2}\right) \left(70(4A+9B) \sin\left(\frac{dx}{2}\right) - 35(5A+18B) \sin\left(c+\frac{dx}{2}\right) + 168A \sin\left(c+\frac{3dx}{2}\right) + 441B\right)}{(a+a \cos(c+dx))^4}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output $(\text{Cos}[(c + d*x)/2] * \text{Sec}[c/2] * (70*(4*A + 9*B) * \text{Sin}[(d*x)/2] - 35*(5*A + 18*B) * \text{Sin}[c + (d*x)/2] + 168*A * \text{Sin}[c + (3*d*x)/2] + 441*B * \text{Sin}[c + (3*d*x)/2] - 105*A * \text{Sin}[2*c + (3*d*x)/2] - 315*B * \text{Sin}[2*c + (3*d*x)/2] + 91*A * \text{Sin}[2*c + (5*d*x)/2] + 147*B * \text{Sin}[2*c + (5*d*x)/2] - 105*B * \text{Sin}[3*c + (5*d*x)/2] + 13*A * \text{Sin}[3*c + (7*d*x)/2] + 36*B * \text{Sin}[3*c + (7*d*x)/2])) / (420*a^4*d*(1 + \text{Cos}[c + d*x])^4)$

3.68.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3456, 3042, 3447, 3042, 3498, 25, 3042, 3229, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^4} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{(a \sin(c + dx + \frac{\pi}{2}) + a)^4} dx$$

↓ 3456

$$\frac{\int \frac{\cos(c+dx)(2a(A-B)+a(A+6B)\cos(c+dx))}{(\cos(c+dx)a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(2a(A-B)+a(A+6B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3447

$$\frac{\int \frac{a(A+6B)\cos^2(c+dx)+2a(A-B)\cos(c+dx)}{(\cos(c+dx)a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3042

3.68. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

$$\begin{aligned}
& \frac{\int \frac{a(A+6B)\sin(c+dx+\frac{\pi}{2})^2+2a(A-B)\sin(c+dx+\frac{\pi}{2})}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow 3498 \\
& -\frac{\int \frac{3(A-8B)a^2+5(A+6B)\cos(c+dx)a^2}{5a^2} dx}{7a^2} - \frac{a(A-8B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow 25 \\
& \frac{\int \frac{3(A-8B)a^2+5(A+6B)\cos(c+dx)a^2}{5a^2} dx}{7a^2} - \frac{a(A-8B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{3(A-8B)a^2+5(A+6B)\sin(c+dx+\frac{\pi}{2})a^2}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{7a^2} - \frac{a(A-8B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow 3229 \\
& \frac{\frac{1}{3}a(13A+36B)\int \frac{1}{\cos(c+dx)a+a} dx - \frac{2(A+27B)\sin(c+dx)}{3d(\cos(c+dx)+1)^2}}{7a^2} - \frac{a(A-8B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow 3042 \\
& \frac{\frac{1}{3}a(13A+36B)\int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx - \frac{2(A+27B)\sin(c+dx)}{3d(\cos(c+dx)+1)^2}}{7a^2} - \frac{a(A-8B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow 3127 \\
& \frac{\frac{a(13A+36B)\sin(c+dx)}{3d(a\cos(c+dx)+a)} - \frac{2(A+27B)\sin(c+dx)}{3d(\cos(c+dx)+1)^2}}{7a^2} - \frac{a(A-8B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{7d(a\cos(c+dx)+a)^4}
\end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `((A - B)*Cos[c + d*x]^2*Sin[c + d*x])/(7*d*(a + a*Cos[c + d*x])^4) + (-1/5 * (a*(A - 8*B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^3) + ((-2*(A + 27*B)*Sin[c + d*x])/(3*d*(1 + Cos[c + d*x])^2) + (a*(13*A + 36*B)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x]))) / (5*a^2) / (7*a^2)`

3.68. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

3.68.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3229 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3498 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b - a*
B + b*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[1
/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b
*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]
```

3.68.4 Maple [A] (verified)

Time = 0.98 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.60

method	result
parallelrisch	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{7(-A+3B) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + 7\left(-\frac{A}{3} - B\right) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 7A + 7B \right)}{56a^4d}$
derivativedivides	$\frac{(A-B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{7} \right) + \frac{(-A+3B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} \right) + \frac{(-A-3B) \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3} \right) + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8da^4}}$
default	$\frac{(A-B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{7} \right) + \frac{(-A+3B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} \right) + \frac{(-A-3B) \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{3} \right) + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8da^4}}$
risch	$\frac{2i(105B e^{6i(dx+c)} + 105A e^{5i(dx+c)} + 315B e^{5i(dx+c)} + 175A e^{4i(dx+c)} + 630B e^{4i(dx+c)} + 280A e^{3i(dx+c)} + 630B e^{3i(dx+c)})}{105da^4(e^{i(dx+c)} + 1)^7}$
norman	$\frac{(A-B) \left(\frac{\tan^{13}\left(\frac{dx}{2} + \frac{c}{2}\right)}{56ad} \right) + \frac{(A+B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8ad} + \frac{3(3A+B) \left(\frac{\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)}{40ad} \right) + \frac{(4A+3B) \left(\frac{\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{12ad} \right) - \frac{(4A+3B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{70ad} \right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^3 a^3}}$

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBO
SE)
```

```
output 1/56*tan(1/2*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^6+7/5*(-A+3*B)*tan(1/2*d
*x+1/2*c)^4+7*(-1/3*A-B)*tan(1/2*d*x+1/2*c)^2+7*A+7*B)/a^4/d
```

3.68.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.91

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{((13A + 36B) \cos(dx + c))^3 + 13(4A + 3B) \cos(dx + c)^2 + 8(4A + 3B) \cos(dx + c) + 8A + 6B) \sin(dx + c)}{105(a^4d \cos(dx + c))^4 + 4a^4d \cos(dx + c)^3 + 6a^4d \cos(dx + c)^2 + 4a^4d \cos(dx + c) + a^4d}$$

3.68. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="fricas")`

output `1/105*((13*A + 36*B)*cos(d*x + c)^3 + 13*(4*A + 3*B)*cos(d*x + c)^2 + 8*(4*A + 3*B)*cos(d*x + c) + 8*A + 6*B)*sin(d*x + c)/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)`

3.68.6 Sympy [A] (verification not implemented)

Time = 2.20 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.34

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \begin{cases} \frac{A \tan^7\left(\frac{c+dx}{2}\right)}{56a^4d} - \frac{A \tan^5\left(\frac{c+dx}{2}\right)}{40a^4d} - \frac{A \tan^3\left(\frac{c+dx}{2}\right)}{24a^4d} + \frac{A \tan\left(\frac{c+dx}{2}\right)}{8a^4d} - \frac{B \tan^7\left(\frac{c+dx}{2}\right)}{56a^4d} + \frac{3B \tan^5\left(\frac{c+dx}{2}\right)}{40a^4d} - \frac{B \tan^3\left(\frac{c+dx}{2}\right)}{8a^4d} \\ \frac{x(A+B\cos(c))\cos^2(c)}{(a\cos(c)+a)^4} \end{cases}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**4,x)`

output `Piecewise((A*tan(c/2 + d*x/2)**7/(56*a**4*d) - A*tan(c/2 + d*x/2)**5/(40*a**4*d) - A*tan(c/2 + d*x/2)**3/(24*a**4*d) + A*tan(c/2 + d*x/2)/(8*a**4*d) - B*tan(c/2 + d*x/2)**7/(56*a**4*d) + 3*B*tan(c/2 + d*x/2)**5/(40*a**4*d) - B*tan(c/2 + d*x/2)**3/(8*a**4*d) + B*tan(c/2 + d*x/2)/(8*a**4*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)**2/(a*cos(c) + a)**4, True))`

3.68.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.29

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{A \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} - \frac{35 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{15 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) + 3B \left(\frac{35 \sin(dx+c)}{\cos(dx+c)+1} - \frac{35 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{5 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right)}{840d}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="maxima")`

3.68. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$

output $\frac{1}{840} * (A * (105 * \sin(dx + c) / (\cos(dx + c) + 1) - 35 * \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 - 21 * \sin(dx + c)^5 / (\cos(dx + c) + 1)^5 + 15 * \sin(dx + c)^7 / (\cos(dx + c) + 1)^7) / a^4 + 3 * B * (35 * \sin(dx + c) / (\cos(dx + c) + 1) - 35 * \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + 21 * \sin(dx + c)^5 / (\cos(dx + c) + 1)^5 - 5 * \sin(dx + c)^7 / (\cos(dx + c) + 1)^7) / a^4) / d$

3.68.8 Giac [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.86

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{15 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 - 15 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 - 21 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 63 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 35 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 105 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 105 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 105 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{840 a^4 d}$$

input `integrate(cos(dx+c)^2*(A+B*cos(dx+c))/(a+a*cos(dx+c))^4,x, algorithm="giac")`

output $\frac{1}{840} * (15 * A * \tan(1/2 * dx + 1/2 * c)^7 - 15 * B * \tan(1/2 * dx + 1/2 * c)^7 - 21 * A * \tan(1/2 * dx + 1/2 * c)^5 + 63 * B * \tan(1/2 * dx + 1/2 * c)^5 - 35 * A * \tan(1/2 * dx + 1/2 * c)^3 - 105 * B * \tan(1/2 * dx + 1/2 * c)^3 + 105 * A * \tan(1/2 * dx + 1/2 * c) + 105 * B * \tan(1/2 * dx + 1/2 * c)) / (a^4 * d)$

3.68.9 Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A + 3B)}{24 a^4} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A - 3B)}{40 a^4} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 (A - B)}{56 a^4} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A + B)}{8 a^4}$$

input `int((cos(c + dx)^2*(A + B*cos(c + dx)))/(a + a*cos(c + dx))^4,x)`

output $-\left(\frac{\tan(c/2 + (dx)/2)^3 * (A + 3 * B)}{(24 * a^4)} + \frac{\tan(c/2 + (dx)/2)^5 * (A - 3 * B)}{(40 * a^4)} - \frac{\tan(c/2 + (dx)/2)^7 * (A - B)}{(56 * a^4)} - \frac{\tan(c/2 + (dx)/2) * (A + B)}{(8 * a^4)}\right) / d$

3.68. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

3.69
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$$

3.69.1 Optimal result 777
 3.69.2 Mathematica [A] (verified) 777
 3.69.3 Rubi [A] (verified) 778
 3.69.4 Maple [A] (verified) 781
 3.69.5 Fricas [A] (verification not implemented) 781
 3.69.6 Sympy [A] (verification not implemented) 782
 3.69.7 Maxima [A] (verification not implemented) 782
 3.69.8 Giac [A] (verification not implemented) 783
 3.69.9 Mupad [B] (verification not implemented) 783

3.69.1 Optimal result

Integrand size = 29, antiderivative size = 138

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = -\frac{(A-B) \sin(c+dx)}{7d(a+a \cos(c+dx))^4} + \frac{(4A-11B) \sin(c+dx)}{35ad(a+a \cos(c+dx))^3} + \frac{(8A+13B) \sin(c+dx)}{105d(a^2+a^2 \cos(c+dx))^2} + \frac{(8A+13B) \sin(c+dx)}{105d(a^4+a^4 \cos(c+dx))}$$

output `-1/7*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^4+1/35*(4*A-11*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3+1/105*(8*A+13*B)*sin(d*x+c)/d/(a^2+a^2*cos(d*x+c))^2+1/105*(8*A+13*B)*sin(d*x+c)/d/(a^4+a^4*cos(d*x+c))`

3.69.2 Mathematica [A] (verified)

Time = 1.09 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.18

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx = \frac{\cos(\frac{1}{2}(c+dx)) \sec(\frac{c}{2}) (140(A+2B) \sin(\frac{dx}{2}) - 35(4A+5B) \sin(c+\frac{dx}{2}) + 168A \sin(c+\frac{3dx}{2}) + 168B \sin(c+\frac{5dx}{2}))}{420a^4}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `(Cos[(c + d*x)/2]*Sec[c/2]*(140*(A + 2*B)*Sin[(d*x)/2] - 35*(4*A + 5*B)*Sin[c + (d*x)/2] + 168*A*Sin[c + (3*d*x)/2] + 168*B*Sin[c + (3*d*x)/2] - 105*B*Sin[2*c + (3*d*x)/2] + 56*A*Sin[2*c + (5*d*x)/2] + 91*B*Sin[2*c + (5*d*x)/2] + 8*A*Sin[3*c + (7*d*x)/2] + 13*B*Sin[3*c + (7*d*x)/2]))/(420*a^4*d*(1 + Cos[c + d*x])^4)`

3.69.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 3447, 3042, 3498, 25, 3042, 3229, 3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^4} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a\cos(c+dx)+a)^4} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{(a\sin(c+dx+\frac{\pi}{2})+a)^4} dx \\ & \quad \downarrow \text{3498} \\ & -\frac{\int -\frac{4a(A-B)+7aB\cos(c+dx)}{(\cos(c+dx)a+a)^3} dx}{7a^2} - \frac{(A-B)\sin(c+dx)}{7d(a\cos(c+dx)+a)^4} \\ & \quad \downarrow \text{25} \\ & \frac{\int \frac{4a(A-B)+7aB\cos(c+dx)}{(\cos(c+dx)a+a)^3} dx}{7a^2} - \frac{(A-B)\sin(c+dx)}{7d(a\cos(c+dx)+a)^4} \end{aligned}$$

$$\begin{aligned}
& \int \frac{4a(A-B)+7aB \sin(c+dx+\frac{\pi}{2})}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx && \downarrow \text{3042} \\
& \frac{\int \frac{4a(A-B)+7aB \sin(c+dx+\frac{\pi}{2})}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} - \frac{(A-B) \sin(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \downarrow \text{3229} \\
& \frac{\frac{1}{5}(8A+13B) \int \frac{1}{(\cos(c+dx)a+a)^2} dx + \frac{a(4A-11B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{7a^2} - \frac{(A-B) \sin(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \downarrow \text{3042} \\
& \frac{\frac{1}{5}(8A+13B) \int \frac{1}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx + \frac{a(4A-11B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{7a^2} - \frac{(A-B) \sin(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \downarrow \text{3129} \\
& \frac{\frac{1}{5}(8A+13B) \left(\int \frac{1}{\cos(c+dx)a+a} dx + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right) + \frac{a(4A-11B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{7a^2} - \frac{(A-B) \sin(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \downarrow \text{3042} \\
& \frac{\frac{1}{5}(8A+13B) \left(\int \frac{1}{\sin(c+dx+\frac{\pi}{2})a+a} dx + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right) + \frac{a(4A-11B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3}}{7a^2} - \frac{(A-B) \sin(c+dx)}{7d(a \cos(c+dx) + a)^4} \\
& \downarrow \text{3127} \\
& \frac{\frac{a(4A-11B) \sin(c+dx)}{5d(a \cos(c+dx)+a)^3} + \frac{1}{5}(8A+13B) \left(\frac{\sin(c+dx)}{3ad(a \cos(c+dx)+a)} + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right)}{7a^2} - \frac{(A-B) \sin(c+dx)}{7d(a \cos(c+dx) + a)^4}
\end{aligned}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^4,x]`

output `-1/7*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^4) + ((a*(4*A - 11*B)*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((8*A + 13*B)*(Sin[c + d*x]/(3*d*(a + a*Cos[c + d*x])^2) + Sin[c + d*x]/(3*a*d*(a + a*Cos[c + d*x]))))/5)/(7*a^2)`

3.69.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3129 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3229 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3498 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(A*b - a*B + b*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[1/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]`

3.69.4 Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.55

method	result
parallelrisc	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{7(A+B) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{7(-A+B) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} - 7A - 7B \right)}{56a^4d}$
derivativedivides	$\frac{\frac{(-A+B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{7} + \frac{(-A-B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{(A-B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8da^4}$
default	$\frac{\frac{(-A+B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{7} + \frac{(-A-B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{(A-B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8da^4}$
risc	$\frac{2i(105B e^{5i(dx+c)} + 140A e^{4i(dx+c)} + 175B e^{4i(dx+c)} + 140A e^{3i(dx+c)} + 280B e^{3i(dx+c)} + 168A e^{2i(dx+c)} + 168B e^{2i(dx+c)})}{105da^4(e^{i(dx+c)} + 1)^7}$
norman	$-\frac{(A-B) \left(\tan^{11}\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{56ad} + \frac{(A+B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8ad} + \frac{(7A+5B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{24ad} + \frac{(11A+B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{60ad} - \frac{(11A+31B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{420ad} - \frac{1}{(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^2 a^3}$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBOSE)
```

```
output -1/56*tan(1/2*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^6+7/5*(A+B)*tan(1/2*d*x+1/2*c)^4+7/3*(-A+B)*tan(1/2*d*x+1/2*c)^2-7*A-7*B)/a^4/d
```

3.69.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.90

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{((8A + 13B) \cos(dx + c))^3 + 4(8A + 13B) \cos(dx + c)^2 + 4(13A + 8B) \cos(dx + c) + 13A + 8B) \sin(dx + c)}{105(a^4d \cos(dx + c))^4 + 4a^4d \cos(dx + c)^3 + 6a^4d \cos(dx + c)^2 + 4a^4d \cos(dx + c) + a^4d}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="fricas")
```

```
output 1/105*((8*A + 13*B)*cos(d*x + c)^3 + 4*(8*A + 13*B)*cos(d*x + c)^2 + 4*(13*A + 8*B)*cos(d*x + c) + 13*A + 8*B)*sin(d*x + c)/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)
```

3.69. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^4} dx$

3.69.6 Sympy [A] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.29

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \begin{cases} -\frac{A \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{56a^4d} - \frac{A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{40a^4d} + \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{24a^4d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} + \frac{B \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{56a^4d} - \frac{B \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{40a^4d} - \frac{B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{24a^4d} \\ \frac{x(A+B\cos(c))\cos(c)}{(a\cos(c)+a)^4} \end{cases}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**4,x)`

output `Piecewise((-A*tan(c/2 + d*x/2)**7/(56*a**4*d) - A*tan(c/2 + d*x/2)**5/(40*a**4*d) + A*tan(c/2 + d*x/2)**3/(24*a**4*d) + A*tan(c/2 + d*x/2)/(8*a**4*d) + B*tan(c/2 + d*x/2)**7/(56*a**4*d) - B*tan(c/2 + d*x/2)**5/(40*a**4*d) - B*tan(c/2 + d*x/2)**3/(24*a**4*d) + B*tan(c/2 + d*x/2)/(8*a**4*d), Ne(d, 0)), (x*(A + B*cos(c))*cos(c)/(a*cos(c) + a)**4, True))`

3.69.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.26

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{A \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} + \frac{35 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{15 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right)}{a^4} + \frac{B \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} - \frac{35 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{15 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right)}{a^4}$$

$$840 d$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="maxima")`

output `1/840*(A*(105*sin(d*x + c)/(cos(d*x + c) + 1) + 35*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 21*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 15*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4 + B*(105*sin(d*x + c)/(cos(d*x + c) + 1) - 35*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 21*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 15*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4/d`

3.69.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.85

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx =$$

$$\frac{15A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^7 - 15B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^7 + 21A \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 + 21B \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 35}{840 a^4 d}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="giac")`

output `-1/840*(15*A*tan(1/2*d*x + 1/2*c)^7 - 15*B*tan(1/2*d*x + 1/2*c)^7 + 21*A*tan(1/2*d*x + 1/2*c)^5 + 21*B*tan(1/2*d*x + 1/2*c)^5 - 35*A*tan(1/2*d*x + 1/2*c)^3 + 35*B*tan(1/2*d*x + 1/2*c)^3 - 105*A*tan(1/2*d*x + 1/2*c) - 105*B*tan(1/2*d*x + 1/2*c))/(a^4*d)`

3.69.9 Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.61

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^4} dx$$

$$= \frac{\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (A+B)}{40 a^4} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (A-B)}{24 a^4} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 (A-B)}{56 a^4} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A+B)}{8 a^4}}{d}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^4,x)`

output `-((tan(c/2 + (d*x)/2)^5*(A + B))/(40*a^4) - (tan(c/2 + (d*x)/2)^3*(A - B))/(24*a^4) + (tan(c/2 + (d*x)/2)^7*(A - B))/(56*a^4) - (tan(c/2 + (d*x)/2)*(A + B))/(8*a^4))/d`

3.70 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^4} dx$

3.70.1	Optimal result	784
3.70.2	Mathematica [A] (verified)	784
3.70.3	Rubi [A] (verified)	785
3.70.4	Maple [A] (verified)	787
3.70.5	Fricas [A] (verification not implemented)	787
3.70.6	Sympy [A] (verification not implemented)	788
3.70.7	Maxima [A] (verification not implemented)	788
3.70.8	Giac [A] (verification not implemented)	789
3.70.9	Mupad [B] (verification not implemented)	789

3.70.1 Optimal result

Integrand size = 23, antiderivative size = 138

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{(A - B) \sin(c + dx)}{7d(a + a \cos(c + dx))^4} + \frac{(3A + 4B) \sin(c + dx)}{35ad(a + a \cos(c + dx))^3} + \frac{2(3A + 4B) \sin(c + dx)}{105d(a^2 + a^2 \cos(c + dx))^2} + \frac{2(3A + 4B) \sin(c + dx)}{105d(a^4 + a^4 \cos(c + dx))}$$

```
output 1/7*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^4+1/35*(3*A+4*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3+2/105*(3*A+4*B)*sin(d*x+c)/d/(a^2+a^2*cos(d*x+c))^2+2/105*(3*A+4*B)*sin(d*x+c)/d/(a^4+a^4*cos(d*x+c))
```

3.70.2 Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.59

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{(36A + 13B + 13(3A + 4B) \cos(c + dx) + 8(3A + 4B) \cos^2(c + dx) + (6A + 8B) \cos^3(c + dx)) \sin(c + dx)}{105a^4d(1 + \cos(c + dx))^4}$$

```
input Integrate[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^4,x]
```

```
output ((36*A + 13*B + 13*(3*A + 4*B)*Cos[c + d*x] + 8*(3*A + 4*B)*Cos[c + d*x]^2 + (6*A + 8*B)*Cos[c + d*x]^3)*Sin[c + d*x]/(105*a^4*d*(1 + Cos[c + d*x])^4)
```

3.70.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$, Rules used = {3042, 3229, 3042, 3129, 3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a \cos(c + dx) + a)^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^4} dx \\
 & \quad \downarrow \text{3229} \\
 & \frac{(3A + 4B) \int \frac{1}{(\cos(c+dx)a+a)^3} dx}{7a} + \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(3A + 4B) \int \frac{1}{(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a} + \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
 & \quad \downarrow \text{3129} \\
 & \frac{(3A + 4B) \left(\frac{2 \int \frac{1}{(\cos(c+dx)a+a)^2} dx}{5a} + \frac{\sin(c+dx)}{5d(a \cos(c+dx)+a)^3} \right)}{7a} + \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(3A + 4B) \left(\frac{2 \int \frac{1}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a} + \frac{\sin(c+dx)}{5d(a \cos(c+dx)+a)^3} \right)}{7a} + \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
 & \quad \downarrow \text{3129} \\
 & \frac{(3A + 4B) \left(\frac{2 \left(\frac{\int \frac{1}{\cos(c+dx)a+a} dx}{3a} + \frac{\sin(c+dx)}{3d(a \cos(c+dx)+a)^2} \right)}{5a} + \frac{\sin(c+dx)}{5d(a \cos(c+dx)+a)^3} \right)}{7a} + \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.70. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^4} dx$

$$(3A + 4B) \left(\frac{2 \left(\frac{\int \frac{1}{\sin(c+dx + \frac{\pi}{2})} a + a \, dx}{3a} + \frac{\sin(c+dx)}{3d(a \cos(c+dx) + a)^2} \right)}{5a} + \frac{\sin(c+dx)}{5d(a \cos(c+dx) + a)^3} \right) + \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4}$$

$$\frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} + \frac{(3A + 4B) \left(\frac{\sin(c+dx)}{5d(a \cos(c+dx) + a)^3} + \frac{2 \left(\frac{\sin(c+dx)}{3ad(a \cos(c+dx) + a)} + \frac{\sin(c+dx)}{3d(a \cos(c+dx) + a)^2} \right)}{5a} \right)}{7a}$$

input `Int[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^4,x]`

output `((A - B)*Sin[c + d*x])/(7*d*(a + a*Cos[c + d*x])^4) + ((3*A + 4*B)*(Sin[c + d*x]/(5*d*(a + a*Cos[c + d*x])^3) + (2*(Sin[c + d*x]/(3*d*(a + a*Cos[c + d*x])^2) + Sin[c + d*x]/(3*a*d*(a + a*Cos[c + d*x]))))/(5*a))/(7*a)`

3.70.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3229 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.70.4 Maple [A] (verified)

Time = 1.28 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.58

method	result
parallelrisc	$\frac{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{7(3A-B) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + 7\left(A + \frac{B}{3}\right) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 7A + 7B \right)}{56a^4d}$
derivativedivides	$\frac{(A-B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{7} \right) + \frac{(3A-B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{(3A+B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8da^4}$
default	$\frac{(A-B) \left(\frac{\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)}{7} \right) + \frac{(3A-B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{5} + \frac{(3A+B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3} + A \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8da^4}$
risc	$\frac{4i(70B e^{4i(dx+c)} + 105A e^{3i(dx+c)} + 70B e^{3i(dx+c)} + 63A e^{2i(dx+c)} + 84B e^{2i(dx+c)} + 21A e^{i(dx+c)} + 28B e^{i(dx+c)} + 3A + 4B)}{105da^4(e^{i(dx+c)} + 1)^7}$
norman	$\frac{(A-B) \left(\frac{\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)}{56ad} \right) + \frac{(A+B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8ad} + \frac{(3A+2B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{12ad} + \frac{(12A+B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{60ad} + \frac{(13A-6B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{140ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a^3}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c))*a^4,x,method=_RETURNVERBOSE)`

output `1/56*tan(1/2*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^6+7/5*(3*A-B)*tan(1/2*d*x+1/2*c)^4+7*(A+1/3*B)*tan(1/2*d*x+1/2*c)^2+7*A+7*B)/a^4/d`

3.70.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.91

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{(2(3A + 4B) \cos(dx + c)^3 + 8(3A + 4B) \cos(dx + c)^2 + 13(3A + 4B) \cos(dx + c) + 36A + 13B) \sin(dx + c)}{105(a^4d \cos(dx + c)^4 + 4a^4d \cos(dx + c)^3 + 6a^4d \cos(dx + c)^2 + 4a^4d \cos(dx + c) + a^4d)}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="fricas")`

output `1/105*(2*(3*A + 4*B)*cos(d*x + c)^3 + 8*(3*A + 4*B)*cos(d*x + c)^2 + 13*(3*A + 4*B)*cos(d*x + c) + 36*A + 13*B)*sin(d*x + c)/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)`

3.70.6 Sympy [A] (verification not implemented)

Time = 1.43 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.28

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \begin{cases} \frac{A \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{56a^4d} + \frac{3A \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{40a^4d} + \frac{A \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} + \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{8a^4d} - \frac{B \tan^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{56a^4d} - \frac{B \tan^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{40a^4d} + \frac{B \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{24a^4d} \\ \frac{x(A+B \cos(c))}{(a \cos(c)+a)^4} \end{cases}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**4,x)`

output `Piecewise((A*tan(c/2 + d*x/2)**7/(56*a**4*d) + 3*A*tan(c/2 + d*x/2)**5/(40*a**4*d) + A*tan(c/2 + d*x/2)**3/(8*a**4*d) + A*tan(c/2 + d*x/2)/(8*a**4*d) - B*tan(c/2 + d*x/2)**7/(56*a**4*d) - B*tan(c/2 + d*x/2)**5/(40*a**4*d) + B*tan(c/2 + d*x/2)**3/(24*a**4*d) + B*tan(c/2 + d*x/2)/(8*a**4*d), Ne(d, 0)), (x*(A + B*cos(c))/(a*cos(c) + a)**4, True))`

3.70.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.27

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{B \left(\frac{105 \sin(dx+c)}{\cos(dx+c)+1} + \frac{35 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{15 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right)}{a^4} + \frac{3A \left(\frac{35 \sin(dx+c)}{\cos(dx+c)+1} + \frac{35 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{5 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right)}{840d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="maxima")`

output `1/840*(B*(105*sin(d*x + c)/(cos(d*x + c) + 1) + 35*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 21*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 15*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4 + 3*A*(35*sin(d*x + c)/(cos(d*x + c) + 1) + 35*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 21*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 5*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4/d`

3.70.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.85

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{15 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 - 15 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^7 + 63 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 21 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 105 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 35 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 105 A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 105 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{840 a^4 d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^4,x, algorithm="giac")`output `1/840*(15*A*tan(1/2*d*x + 1/2*c)^7 - 15*B*tan(1/2*d*x + 1/2*c)^7 + 63*A*tan(1/2*d*x + 1/2*c)^5 - 21*B*tan(1/2*d*x + 1/2*c)^5 + 105*A*tan(1/2*d*x + 1/2*c)^3 + 35*B*tan(1/2*d*x + 1/2*c)^3 + 105*A*tan(1/2*d*x + 1/2*c) + 105*B*tan(1/2*d*x + 1/2*c))/(a^4*d)`**3.70.9 Mupad [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.63

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (3A+B)}{24 a^4} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 (A-B)}{56 a^4} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (A+B)}{8 a^4} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 (3A-B)}{40 a^4}$$

$$d$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^4,x)`output `((tan(c/2 + (d*x)/2)^3*(3*A + B))/(24*a^4) + (tan(c/2 + (d*x)/2)^7*(A - B))/(56*a^4) + (tan(c/2 + (d*x)/2)*(A + B))/(8*a^4) + (tan(c/2 + (d*x)/2)^5*(3*A - B))/(40*a^4))/d`

3.71 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^4} dx$

3.71.1 Optimal result	790
3.71.2 Mathematica [A] (verified)	790
3.71.3 Rubi [A] (verified)	791
3.71.4 Maple [A] (verified)	794
3.71.5 Fricas [A] (verification not implemented)	794
3.71.6 Sympy [F]	795
3.71.7 Maxima [A] (verification not implemented)	795
3.71.8 Giac [A] (verification not implemented)	796
3.71.9 Mupad [B] (verification not implemented)	796

3.71.1 Optimal result

Integrand size = 29, antiderivative size = 147

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{\text{Aarctanh}(\sin(c + dx))}{a^4 d} - \frac{(55A - 6B) \sin(c + dx)}{105a^4 d (1 + \cos(c + dx))^2} - \frac{2(80A - 3B) \sin(c + dx)}{105a^4 d (1 + \cos(c + dx))} - \frac{(A - B) \sin(c + dx)}{7d(a + a \cos(c + dx))^4} - \frac{(10A - 3B) \sin(c + dx)}{35ad(a + a \cos(c + dx))^3}$$

```
output A*arctanh(sin(d*x+c))/a^4/d-1/105*(55*A-6*B)*sin(d*x+c)/a^4/d/(1+cos(d*x+c))^2-2/105*(80*A-3*B)*sin(d*x+c)/a^4/d/(1+cos(d*x+c))-1/7*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^4-1/35*(10*A-3*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```

3.71.2 Mathematica [A] (verified)

Time = 1.44 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.63

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{-6720A \cos^8\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^4,x]`

output `(-6720*A*Cos[(c + d*x)/2]^8*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Cos[(c + d*x)/2]*Sec[c/2]*(-70*(49*A - 3*B)*Sin[(d*x)/2] + 2170*A*Sin[c + (d*x)/2] - 2625*A*Sin[c + (3*d*x)/2] + 126*B*Sin[c + (3*d*x)/2] + 735*A*Sin[2*c + (3*d*x)/2] - 1015*A*Sin[2*c + (5*d*x)/2] + 42*B*Sin[2*c + (5*d*x)/2] + 105*A*Sin[3*c + (5*d*x)/2] - 160*A*Sin[3*c + (7*d*x)/2] + 6*B*Sin[3*c + (7*d*x)/2]))/(420*a^4*d*(1 + Cos[c + d*x])^4)`

3.71.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 3457, 3042, 3457, 3042, 3457, 3042, 3457, 27, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a\sin(c+dx+\frac{\pi}{2})+a)^4} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(7aA-3a(A-B)\cos(c+dx))\sec(c+dx)}{(\cos(c+dx)a+a)^3} dx}{7a^2} - \frac{(A-B)\sin(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{7aA-3a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} - \frac{(A-B)\sin(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(35a^2A-2a^2(10A-3B)\cos(c+dx))\sec(c+dx)}{(\cos(c+dx)a+a)^2} dx}{7a^2} - \frac{a(10A-3B)\sin(c+dx)}{5d(a\cos(c+dx)+a)^3} - \frac{(A-B)\sin(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.71. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+a\cos(c+dx))^4} dx$

$$\begin{aligned}
& \frac{\int \frac{35a^2 A - 2a^2(10A - 3B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a\right)^2} dx}{5a^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
& \quad \downarrow 3457 \\
& \frac{\int \frac{(105a^3 A - a^3(55A - 6B) \cos(c + dx)) \sec(c + dx)}{\cos(c + dx) a + a} dx}{3a^2} - \frac{(55A - 6B) \sin(c + dx)}{3d(\cos(c + dx) + 1)^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{105a^3 A - a^3(55A - 6B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a\right)} dx}{3a^2} - \frac{(55A - 6B) \sin(c + dx)}{3d(\cos(c + dx) + 1)^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
& \quad \downarrow 3457 \\
& \frac{\int 105a^4 A \sec(c + dx) dx}{a^2} - \frac{2a^3(80A - 3B) \sin(c + dx)}{3a^2 d(a \cos(c + dx) + a)} - \frac{(55A - 6B) \sin(c + dx)}{3d(\cos(c + dx) + 1)^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
& \quad \downarrow 27 \\
& \frac{105a^2 A \int \sec(c + dx) dx}{3a^2} - \frac{2a^3(80A - 3B) \sin(c + dx)}{d(a \cos(c + dx) + a)} - \frac{(55A - 6B) \sin(c + dx)}{3d(\cos(c + dx) + 1)^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \frac{(A - B) \sin(c + dx)}{7d(a \cos(c + dx) + a)^4} \\
& \quad \downarrow 3042 \\
& \frac{105a^2 A \int \csc\left(c + dx + \frac{\pi}{2}\right) dx}{3a^2} - \frac{2a^3(80A - 3B) \sin(c + dx)}{d(a \cos(c + dx) + a)} - \frac{(55A - 6B) \sin(c + dx)}{3d(\cos(c + dx) + 1)^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \\
& \quad \frac{7a^2}{7d(a \cos(c + dx) + a)^4} \\
& \quad \downarrow 4257 \\
& \frac{105a^2 A \operatorname{arctanh}(\sin(c + dx))}{d} - \frac{2a^3(80A - 3B) \sin(c + dx)}{3a^2 d(a \cos(c + dx) + a)} - \frac{(55A - 6B) \sin(c + dx)}{3d(\cos(c + dx) + 1)^2} - \frac{a(10A - 3B) \sin(c + dx)}{5d(a \cos(c + dx) + a)^3} - \\
& \quad \frac{7a^2}{7d(a \cos(c + dx) + a)^4}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^4,x]`

$$3.71. \quad \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx$$

```
output -1/7*((A - B)*Sin[c + d*x])/(d*(a + a*cos[c + d*x])^4) + (-1/5*(a*(10*A -
3*B)*Sin[c + d*x])/(d*(a + a*cos[c + d*x])^3) + (-1/3*((55*A - 6*B)*Sin[c
+ d*x])/(d*(1 + Cos[c + d*x])^2) + ((105*a^2*A*ArcTanh[Sin[c + d*x]])/d -
(2*a^3*(80*A - 3*B)*Sin[c + d*x])/(d*(a + a*cos[c + d*x])))/(3*a^2)/(5*a^
2))/(7*a^2)
```

3.71.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```


3.71.4 Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.77

method	result
parallelrisch	$-840A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 840A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - 15 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) \left((A-B) \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \left(7A - \frac{21B}{5}\right) \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \frac{840a^4d}{840a^4d} \right)$
derivativedivides	$-\frac{11 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{3} + \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B - \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{7} + \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{7} + \frac{3 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{5} - A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)$
default	$-\frac{11 \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{3} + \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B - \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right) A}{7} + \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{7} + \frac{3 \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right) B}{5} - A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)$
risch	$-\frac{2i \left(105A e^{6i(dx+c)} + 735A e^{5i(dx+c)} + 2170A e^{4i(dx+c)} + 3430A e^{3i(dx+c)} - 210B e^{3i(dx+c)} + 2625A e^{2i(dx+c)} - 126B e^{2i(dx+c)} \right)}{105d a^4 \left(e^{i(dx+c)} + 1 \right)^7}$
norman	$-\frac{(A-B) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{56ad} - \frac{(15A-B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8ad} - \frac{(20A-13B) \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{140ad} - \frac{(28A-3B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{12ad} - \frac{(35A-12B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{60ad}$
	$\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a^3$

input `int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBOSE)`

output `1/840*(-840*A*ln(tan(1/2*d*x+1/2*c))-1)+840*A*ln(tan(1/2*d*x+1/2*c)+1)-15*tan(1/2*d*x+1/2*c)*((A-B)*tan(1/2*d*x+1/2*c)^6+(7*A-21/5*B)*tan(1/2*d*x+1/2*c)^4+(77/3*A-7*B)*tan(1/2*d*x+1/2*c)^2+105*A-7*B)/a^4/d`

3.71.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.61

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{105 (A \cos(dx + c)^4 + 4 A \cos(dx + c)^3 + 6 A \cos(dx + c)^2 + 4 A \cos(dx + c) + A) \log(\sin(dx + c) + 1)}{105d a^4 (e^{i(dx+c)} + 1)^7}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^4,x, algorithm="fricas")`

```
output 1/210*(105*(A*cos(d*x + c)^4 + 4*A*cos(d*x + c)^3 + 6*A*cos(d*x + c)^2 + 4
*A*cos(d*x + c) + A)*log(sin(d*x + c) + 1) - 105*(A*cos(d*x + c)^4 + 4*A*cos
os(d*x + c)^3 + 6*A*cos(d*x + c)^2 + 4*A*cos(d*x + c) + A)*log(-sin(d*x +
c) + 1) - 2*(2*(80*A - 3*B)*cos(d*x + c)^3 + (535*A - 24*B)*cos(d*x + c)^2
+ (620*A - 39*B)*cos(d*x + c) + 260*A - 36*B)*sin(d*x + c))/(a^4*d*cos(d*
x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d
*x + c) + a^4*d)
```

3.71.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{\int \frac{A \sec(c + dx)}{\cos^4(c + dx) + 4 \cos^3(c + dx) + 6 \cos^2(c + dx) + 4 \cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec(c + dx)}{\cos^4(c + dx) + 4 \cos^3(c + dx) + 6 \cos^2(c + dx) + 4 \cos(c + dx) + 1} dx}{a^4}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))**4,x)
```

```
output (Integral(A*sec(c + d*x)/(cos(c + d*x)**4 + 4*cos(c + d*x)**3 + 6*cos(c +
d*x)**2 + 4*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)/(
cos(c + d*x)**4 + 4*cos(c + d*x)**3 + 6*cos(c + d*x)**2 + 4*cos(c + d*x) +
1), x))/a**4
```

3.71.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.55

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx =$$

$$\frac{5A \left(\frac{315 \sin(dx+c)}{\cos(dx+c)+1} + \frac{77 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{21 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{3 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} - \frac{168 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^4} + \frac{168 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1\right)}{a^4} \right) - 3B}{840d}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^4,x, algorithm="max
ima")
```

3.71. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^4} dx$

output
$$\frac{-1/840*(5*A*((315*\sin(d*x + c)/(\cos(d*x + c) + 1) + 77*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 21*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 + 3*\sin(d*x + c)^7/(\cos(d*x + c) + 1)^7)/a^4 - 168*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)/a^4 + 168*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) - 1)/a^4 - 3*B*(35*\sin(d*x + c)/(\cos(d*x + c) + 1) + 35*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 21*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 + 5*\sin(d*x + c)^7/(\cos(d*x + c) + 1)^7)/a^4)/d$$

3.71.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.24

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{840 A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^4} - \frac{840 A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^4} - \frac{15 A a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 - 15 B a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^7 + 105 A a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 105 B a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 + 385 A a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 105 B a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 1575 A a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c) - 105 B a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)}{a^{28} d}$$

840

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^4,x, algorithm="giac")`

output
$$\frac{1/840*(840*A*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1))/a^4 - 840*A*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1))/a^4 - (15*A*a^{24}*\tan(1/2*d*x + 1/2*c)^7 - 15*B*a^{24}*\tan(1/2*d*x + 1/2*c)^7 + 105*A*a^{24}*\tan(1/2*d*x + 1/2*c)^5 - 63*B*a^{24}*\tan(1/2*d*x + 1/2*c)^5 + 385*A*a^{24}*\tan(1/2*d*x + 1/2*c)^3 - 105*B*a^{24}*\tan(1/2*d*x + 1/2*c)^3 + 1575*A*a^{24}*\tan(1/2*d*x + 1/2*c) - 105*B*a^{24}*\tan(1/2*d*x + 1/2*c))/a^{28})/d$$

3.71.9 Mupad [B] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.35

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{2 A \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{a^4 d}$$

$$- \frac{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)^4 \left(\frac{11 A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{24} - \frac{B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{8}\right) + \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \left(\frac{A \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{8} - \frac{3 B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{40}\right) + \cos\left(\frac{c}{2} + \frac{dx}{2}\right)}{a^4 d \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^7}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^4),x)`

output `(2*A*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(a^4*d) - (cos(c/2 + (d*x)/2)^4*((11*A*sin(c/2 + (d*x)/2)^3)/24 - (B*sin(c/2 + (d*x)/2)^3)/8) + cos(c/2 + (d*x)/2)^2*((A*sin(c/2 + (d*x)/2)^5)/8 - (3*B*sin(c/2 + (d*x)/2)^5)/40) + cos(c/2 + (d*x)/2)^6*((15*A*sin(c/2 + (d*x)/2))/8 - (B*sin(c/2 + (d*x)/2))/8) + (A*sin(c/2 + (d*x)/2)^7)/56 - (B*sin(c/2 + (d*x)/2)^7)/56)/(a^4*d*cos(c/2 + (d*x)/2)^7)`

3.72 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^4} dx$

3.72.1	Optimal result	798
3.72.2	Mathematica [B] (verified)	799
3.72.3	Rubi [A] (verified)	799
3.72.4	Maple [A] (verified)	803
3.72.5	Fricas [B] (verification not implemented)	804
3.72.6	Sympy [F]	804
3.72.7	Maxima [A] (verification not implemented)	805
3.72.8	Giac [A] (verification not implemented)	806
3.72.9	Mupad [B] (verification not implemented)	806

3.72.1 Optimal result

Integrand size = 31, antiderivative size = 175

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx = -\frac{(4A - B)\operatorname{arctanh}(\sin(c + dx))}{a^4 d} + \frac{8(83A - 20B) \tan(c + dx)}{105a^4 d} - \frac{(88A - 25B) \tan(c + dx)}{105a^4 d(1 + \cos(c + dx))^2} - \frac{(4A - B) \tan(c + dx)}{a^4 d(1 + \cos(c + dx))} - \frac{(A - B) \tan(c + dx)}{7d(a + a \cos(c + dx))^4} - \frac{(12A - 5B) \tan(c + dx)}{35ad(a + a \cos(c + dx))^3}$$

```
output - (4*A-B)*arctanh(sin(d*x+c))/a^4/d+8/105*(83*A-20*B)*tan(d*x+c)/a^4/d-1/105*(88*A-25*B)*tan(d*x+c)/a^4/d/(1+cos(d*x+c))^2-(4*A-B)*tan(d*x+c)/a^4/d/(1+cos(d*x+c))-1/7*(A-B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^4-1/35*(12*A-5*B)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```

3.72.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 595 vs. $2(175) = 350$.

Time = 4.60 (sec) , antiderivative size = 595, normalized size of antiderivative = 3.40

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{26880(4A - B) \cos^8\left(\frac{1}{2}(c + dx)\right) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{(1680a^4d(1 + \cos(c + dx))^4)}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^4,x]`

output `(26880*(4*A - B)*Cos[(c + d*x)/2]^8*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + Cos[(c + d*x)/2]*Sec[c/2]*Sec[c]*Sec[c + d*x]*(-245*(44*A - 17*B)*Sin[(d*x)/2] + 7*(2684*A - 635*B)*Sin[(3*d*x)/2] - 20524*A*Sin[c - (d*x)/2] + 4795*B*Sin[c - (d*x)/2] + 14644*A*Sin[c + (d*x)/2] - 4795*B*Sin[c + (d*x)/2] - 16660*A*Sin[2*c + (d*x)/2] + 4165*B*Sin[2*c + (d*x)/2] - 4690*A*Sin[c + (3*d*x)/2] + 2275*B*Sin[c + (3*d*x)/2] + 14378*A*Sin[2*c + (3*d*x)/2] - 4445*B*Sin[2*c + (3*d*x)/2] - 9100*A*Sin[3*c + (3*d*x)/2] + 2275*B*Sin[3*c + (3*d*x)/2] + 11668*A*Sin[c + (5*d*x)/2] - 2785*B*Sin[c + (5*d*x)/2] - 630*A*Sin[2*c + (5*d*x)/2] + 735*B*Sin[2*c + (5*d*x)/2] + 9358*A*Sin[3*c + (5*d*x)/2] - 2785*B*Sin[3*c + (5*d*x)/2] - 2940*A*Sin[4*c + (5*d*x)/2] + 735*B*Sin[4*c + (5*d*x)/2] + 4228*A*Sin[2*c + (7*d*x)/2] - 1015*B*Sin[2*c + (7*d*x)/2] + 315*A*Sin[3*c + (7*d*x)/2] + 105*B*Sin[3*c + (7*d*x)/2] + 3493*A*Sin[4*c + (7*d*x)/2] - 1015*B*Sin[4*c + (7*d*x)/2] - 420*A*Sin[5*c + (7*d*x)/2] + 105*B*Sin[5*c + (7*d*x)/2] + 664*A*Sin[3*c + (9*d*x)/2] - 160*B*Sin[3*c + (9*d*x)/2] + 105*A*Sin[4*c + (9*d*x)/2] + 559*A*Sin[5*c + (9*d*x)/2] - 160*B*Sin[5*c + (9*d*x)/2]))/(1680*a^4*d*(1 + Cos[c + d*x])^4)`

3.72.3 Rubi [A] (verified)

Time = 1.44 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.14, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3457, 3042, 3457, 3042, 3457, 3042, 3457, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.72. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^4} dx$

$$\begin{aligned}
& \int \frac{\sec^2(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A+B\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 (a\sin\left(c+dx+\frac{\pi}{2}\right)+a)^4} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(a(8A-B)-4a(A-B)\cos(c+dx))\sec^2(c+dx)}{(\cos(c+dx)a+a)^3} dx}{7a^2} - \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(8A-B)-4a(A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 (\sin\left(c+dx+\frac{\pi}{2}\right)a+a)^3} dx}{7a^2} - \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(2a^2(26A-5B)-3a^2(12A-5B)\cos(c+dx))\sec^2(c+dx)}{(\cos(c+dx)a+a)^2} dx}{7a^2} - \frac{a(12A-5B)\tan(c+dx)}{5d(a\cos(c+dx)+a)^3} - \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2a^2(26A-5B)-3a^2(12A-5B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 (\sin\left(c+dx+\frac{\pi}{2}\right)a+a)^2} dx}{7a^2} - \frac{a(12A-5B)\tan(c+dx)}{5d(a\cos(c+dx)+a)^3} - \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(a^3(244A-55B)-2a^3(88A-25B)\cos(c+dx))\sec^2(c+dx)}{\cos(c+dx)a+a} dx}{5a^2} - \frac{(88A-25B)\tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B)\tan(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a^3(244A-55B)-2a^3(88A-25B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 (\sin\left(c+dx+\frac{\pi}{2}\right)a+a)} dx}{5a^2} - \frac{(88A-25B)\tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B)\tan(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4}
\end{aligned}$$

3.72. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{(a+a\cos(c+dx))^4} dx$

$$\frac{\int \frac{(8a^4(83A-20B) - 105a^4(4A-B) \cos(c+dx)) \sec^2(c+dx) dx}{a^2} - \frac{105a^3(4A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(88A-25B) \tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3}}{3a^2} - \frac{5a^2}{5a^2}$$

$$\frac{7a^2}{(A-B) \tan(c+dx)} \frac{1}{7d(a \cos(c+dx)+a)^4}$$

↓ 3042

$$\frac{\int \frac{8a^4(83A-20B) - 105a^4(4A-B) \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2} dx}{a^2} - \frac{105a^3(4A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(88A-25B) \tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

$$\frac{7a^2}{(A-B) \tan(c+dx)} \frac{1}{7d(a \cos(c+dx)+a)^4}$$

↓ 3227

$$\frac{8a^4(83A-20B) \int \sec^2(c+dx) dx - 105a^4(4A-B) \int \sec(c+dx) dx - \frac{105a^3(4A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(88A-25B) \tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3}}{a^2} - \frac{3a^2}{3a^2} - \frac{5a^2}{5a^2}$$

$$\frac{7a^2}{(A-B) \tan(c+dx)} \frac{1}{7d(a \cos(c+dx)+a)^4}$$

↓ 3042

$$\frac{8a^4(83A-20B) \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx - 105a^4(4A-B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx - \frac{105a^3(4A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(88A-25B) \tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3}}{a^2} - \frac{3a^2}{3a^2} - \frac{5a^2}{5a^2}$$

$$\frac{7a^2}{(A-B) \tan(c+dx)} \frac{1}{7d(a \cos(c+dx)+a)^4}$$

↓ 4254

$$\frac{-\frac{8a^4(83A-20B) \int \frac{1d(-\tan(c+dx))}{d} - 105a^4(4A-B) \int \csc\left(c+dx+\frac{\pi}{2}\right) dx - \frac{105a^3(4A-B) \tan(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(88A-25B) \tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B) \tan(c+dx)}{5d(a \cos(c+dx)+a)^3}}{a^2} - \frac{3a^2}{3a^2} - \frac{5a^2}{5a^2}}$$

$$\frac{7a^2}{(A-B) \tan(c+dx)} \frac{1}{7d(a \cos(c+dx)+a)^4}$$

↓ 24

3.72. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^4} dx$

$$\frac{\frac{8a^4(83A-20B)\tan(c+dx) - 105a^4(4A-B)\int \csc\left(c+dx+\frac{\pi}{2}\right)dx - 105a^3(4A-B)\tan(c+dx)}{a^2} - \frac{(88A-25B)\tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B)\tan(c+dx)}{5d(a\cos(c+dx)+a)^3}}{3a^2} - \frac{7a^2}{5a^2} \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 4257

$$\frac{\frac{8a^4(83A-20B)\tan(c+dx) - 105a^4(4A-B)\operatorname{arctanh}(\sin(c+dx)) - 105a^3(4A-B)\tan(c+dx)}{a^2} - \frac{(88A-25B)\tan(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{a(12A-5B)\tan(c+dx)}{5d(a\cos(c+dx)+a)^3}}{3a^2} - \frac{7a^2}{5a^2} \frac{(A-B)\tan(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^4,x]`

output `-1/7*((A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^4) + (-1/5*(a*(12*A - 5*B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^3) + (-1/3*((88*A - 25*B)*Tan[c + d*x])/(d*(1 + Cos[c + d*x])^2) + ((-105*a^3*(4*A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x]))) + ((-105*a^4*(4*A - B)*ArcTanh[Sin[c + d*x]])/d + (8*a^4*(83*A - 20*B)*Tan[c + d*x])/d)/a^2)/(3*a^2))/(5*a^2))/(7*a^2)`

3.72.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 4254 Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.72.4 Maple [A] (verified)

Time = 1.47 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.91

method	result
parallelrisc	$\frac{13440 \cos(dx+c) \left(A - \frac{B}{4}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) - 13440 \cos(dx+c) \left(A - \frac{B}{4}\right) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 332 \left(\sec^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^{11} - 1\right)}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^{11} - 1}$
derivativedivides	$(-32A+8B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{8A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1} + \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^A}{7} - \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^B}{7} + \frac{7A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5} - \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)$
default	$(-32A+8B) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) - \frac{8A}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1} + \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^A}{7} - \frac{\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^B}{7} + \frac{7A \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5} - \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)$
norman	$\frac{(A-B) \left(\tan^{11}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (7A-5B) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \frac{5(13A-3B) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{8ad} + \frac{7(17A-5B) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{20ad} - \frac{(71A-11B) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{24ad}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) a^3}$
risc	$2i \left(420A e^{8i(dx+c)} - 105B e^{8i(dx+c)} + 2940A e^{7i(dx+c)} - 735B e^{7i(dx+c)} + 9100A e^{6i(dx+c)} - 2275B e^{6i(dx+c)} + 16660A e^{5i(dx+c)} - 4200B e^{5i(dx+c)} + 10500A e^{4i(dx+c)} - 1050B e^{4i(dx+c)} + 10500A e^{3i(dx+c)} - 1050B e^{3i(dx+c)} + 10500A e^{2i(dx+c)} - 1050B e^{2i(dx+c)} + 10500A e^{i(dx+c)} - 1050B e^{i(dx+c)}\right)$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBO
SE)
```

$$3.72. \int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^4} dx$$

output $\frac{1}{3360} \cdot (13440 \cdot \cos(dx+c) \cdot (A-1/4 \cdot B) \cdot \ln(\tan(1/2 \cdot dx+1/2 \cdot c)-1) - 13440 \cdot \cos(dx+c) \cdot (A-1/4 \cdot B) \cdot \ln(\tan(1/2 \cdot dx+1/2 \cdot c)+1) + 332 \cdot \sec(1/2 \cdot dx+1/2 \cdot c)^6 \cdot ((1650/83 \cdot A - 390/83 \cdot B) \cdot \cos(2 \cdot dx+2 \cdot c) + (559/83 \cdot A - 535/332 \cdot B) \cdot \cos(3 \cdot dx+3 \cdot c) + (A-20/83 \cdot B) \cdot \cos(4 \cdot dx+4 \cdot c) + (2861/83 \cdot A - 2645/332 \cdot B) \cdot \cos(dx+c) + 1672/83 \cdot A - 370/83 \cdot B) \cdot \tan(1/2 \cdot dx+1/2 \cdot c)) / d / a^4 / \cos(dx+c)$

3.72.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 337 vs. $2(167) = 334$.

Time = 0.31 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.93

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{105 \left((4A - B) \cos(dx + c)^5 + 4(4A - B) \cos(dx + c)^4 + 6(4A - B) \cos(dx + c)^3 + 4(4A - B) \cos(dx + c)^2 + a^4 d \cos(dx + c) \right)}{a^4 \cos^4(c + dx) + 4a^4 \cos^3(c + dx) + 6a^4 \cos^2(c + dx) + 4a^4 \cos(c + dx) + 1}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^4,x, algorithm="fricas")`

output $-1/210 \cdot (105 \cdot ((4A - B) \cdot \cos(dx + c)^5 + 4 \cdot (4A - B) \cdot \cos(dx + c)^4 + 6 \cdot (4A - B) \cdot \cos(dx + c)^3 + 4 \cdot (4A - B) \cdot \cos(dx + c)^2 + (4A - B) \cdot \cos(dx + c)) \cdot \log(\sin(dx + c) + 1) - 105 \cdot ((4A - B) \cdot \cos(dx + c)^5 + 4 \cdot (4A - B) \cdot \cos(dx + c)^4 + 6 \cdot (4A - B) \cdot \cos(dx + c)^3 + 4 \cdot (4A - B) \cdot \cos(dx + c)^2 + (4A - B) \cdot \cos(dx + c)) \cdot \log(-\sin(dx + c) + 1) - 2 \cdot (8 \cdot (83A - 20B) \cdot \cos(dx + c)^4 + (2236A - 535B) \cdot \cos(dx + c)^3 + 4 \cdot (659A - 155B) \cdot \cos(dx + c)^2 + 4 \cdot (296A - 65B) \cdot \cos(dx + c) + 105A) \cdot \sin(dx + c)) / (a^4 \cdot d \cdot \cos(dx + c)^5 + 4 \cdot a^4 \cdot d \cdot \cos(dx + c)^4 + 6 \cdot a^4 \cdot d \cdot \cos(dx + c)^3 + 4 \cdot a^4 \cdot d \cdot \cos(dx + c)^2 + a^4 \cdot d \cdot \cos(dx + c))$

3.72.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{\int \frac{A \sec^2(c + dx)}{\cos^4(c + dx) + 4 \cos^3(c + dx) + 6 \cos^2(c + dx) + 4 \cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sec^2(c + dx)}{\cos^4(c + dx) + 4 \cos^3(c + dx) + 6 \cos^2(c + dx) + 4 \cos(c + dx) + 1} dx}{a^4}$$

3.72. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^4} dx$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c))**4,x)`

output `(Integral(A*sec(c + d*x)**2/(cos(c + d*x)**4 + 4*cos(c + d*x)**3 + 6*cos(c + d*x)**2 + 4*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)**2/(cos(c + d*x)**4 + 4*cos(c + d*x)**3 + 6*cos(c + d*x)**2 + 4*cos(c + d*x) + 1), x))/a**4`

3.72.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.86

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= A \left(\frac{1680 \sin(dx+c)}{\left(a^4 - \frac{a^4 \sin^2(dx+c)}{(\cos(dx+c)+1)^2}\right) (\cos(dx+c)+1)} + \frac{5145 \sin(dx+c)}{\cos(dx+c)+1} + \frac{805 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{147 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{15 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) - \frac{3360 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1\right)}{a^4}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^4,x, algorithm="maxima")`

output `1/840*(A*(1680*sin(d*x + c)/((a^4 - a^4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) + (5145*sin(d*x + c)/(cos(d*x + c) + 1) + 805*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 147*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 15*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4 - 3360*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a^4 + 3360*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a^4) - 5*B*((315*sin(d*x + c)/(cos(d*x + c) + 1) + 77*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 21*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 3*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)/a^4 - 168*log(sin(d*x + c)/(cos(d*x + c) + 1) + 1)/a^4 + 168*log(sin(d*x + c)/(cos(d*x + c) + 1) - 1)/a^4)/d`

3.72.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.28

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx =$$

$$\frac{840(4A - B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^4} - \frac{840(4A - B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^4} + \frac{1680 A \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1) a^4} - \frac{15 A a^{24} \tan(\frac{1}{2} dx + \frac{1}{2} c)^7}{a^4} -$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^4,x, algorithm="giac")`

output `-1/840*(840*(4*A - B)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^4 - 840*(4*A - B)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^4 + 1680*A*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 - 1)*a^4) - (15*A*a^24*tan(1/2*d*x + 1/2*c)^7 - 15*B*a^24*tan(1/2*d*x + 1/2*c)^7 + 147*A*a^24*tan(1/2*d*x + 1/2*c)^5 - 105*B*a^24*tan(1/2*d*x + 1/2*c)^5 + 805*A*a^24*tan(1/2*d*x + 1/2*c)^3 - 385*B*a^24*tan(1/2*d*x + 1/2*c)^3 + 5145*A*a^24*tan(1/2*d*x + 1/2*c) - 1575*B*a^24*tan(1/2*d*x + 1/2*c))/a^28)/d`

3.72.9 Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.35

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{\tan(\frac{c}{2} + \frac{dx}{2})^3 \left(\frac{A-B}{8a^4} + \frac{5A-3B}{12a^4} + \frac{10A-2B}{24a^4} \right)}{d} + \frac{\tan(\frac{c}{2} + \frac{dx}{2})^5 \left(\frac{A-B}{20a^4} + \frac{5A-3B}{40a^4} \right)}{d}$$

$$+ \frac{\tan(\frac{c}{2} + \frac{dx}{2}) \left(\frac{A-B}{2a^4} + \frac{3(5A-3B)}{8a^4} + \frac{10A-2B}{4a^4} + \frac{10A+2B}{8a^4} \right)}{d} + \frac{\tan(\frac{c}{2} + \frac{dx}{2})^7 (A - B)}{56 a^4 d}$$

$$- \frac{2A \tan(\frac{c}{2} + \frac{dx}{2})}{d \left(a^4 \tan(\frac{c}{2} + \frac{dx}{2})^2 - a^4 \right)} - \frac{2 \operatorname{atanh}(\tan(\frac{c}{2} + \frac{dx}{2})) (4A - B)}{a^4 d}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^4),x)`

output $(\tan(c/2 + (d*x)/2)^3((A - B)/(8*a^4) + (5*A - 3*B)/(12*a^4) + (10*A - 2*B)/(24*a^4)))/d + (\tan(c/2 + (d*x)/2)^5((A - B)/(20*a^4) + (5*A - 3*B)/(40*a^4)))/d + (\tan(c/2 + (d*x)/2)*((A - B)/(2*a^4) + (3*(5*A - 3*B))/(8*a^4) + (10*A - 2*B)/(4*a^4) + (10*A + 2*B)/(8*a^4)))/d + (\tan(c/2 + (d*x)/2)^7*(A - B))/(56*a^4*d) - (2*A*\tan(c/2 + (d*x)/2))/(d*(a^4*\tan(c/2 + (d*x)/2)^2 - a^4)) - (2*atanh(\tan(c/2 + (d*x)/2))*(4*A - B))/(a^4*d)$

3.73 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^4} dx$

3.73.1	Optimal result	808
3.73.2	Mathematica [B] (verified)	809
3.73.3	Rubi [A] (verified)	810
3.73.4	Maple [A] (verified)	814
3.73.5	Fricas [A] (verification not implemented)	815
3.73.6	Sympy [F]	816
3.73.7	Maxima [A] (verification not implemented)	816
3.73.8	Giac [A] (verification not implemented)	817
3.73.9	Mupad [B] (verification not implemented)	818

3.73.1 Optimal result

Integrand size = 31, antiderivative size = 232

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx = \frac{(21A - 8B) \arctanh(\sin(c + dx))}{2a^4d} - \frac{8(216A - 83B) \tan(c + dx)}{105a^4d} + \frac{(21A - 8B) \sec(c + dx) \tan(c + dx)}{2a^4d} - \frac{(129A - 52B) \sec(c + dx) \tan(c + dx)}{105a^4d(1 + \cos(c + dx))^2} - \frac{4(216A - 83B) \sec(c + dx) \tan(c + dx)}{105a^4d(1 + \cos(c + dx))} - \frac{(A - B) \sec(c + dx) \tan(c + dx)}{7d(a + a \cos(c + dx))^4} - \frac{(2A - B) \sec(c + dx) \tan(c + dx)}{5ad(a + a \cos(c + dx))^3}$$

output

```
1/2*(21*A-8*B)*arctanh(sin(d*x+c))/a^4/d-8/105*(216*A-83*B)*tan(d*x+c)/a^4/d+1/2*(21*A-8*B)*sec(d*x+c)*tan(d*x+c)/a^4/d-1/105*(129*A-52*B)*sec(d*x+c)*tan(d*x+c)/a^4/d/(1+cos(d*x+c))^2-4/105*(216*A-83*B)*sec(d*x+c)*tan(d*x+c)/a^4/d/(1+cos(d*x+c))-1/7*(A-B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^4-1/5*(2*A-B)*sec(d*x+c)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^3
```

3.73.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 798 vs. $2(232) = 464$.

Time = 8.17 (sec) , antiderivative size = 798, normalized size of antiderivative = 3.44

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= -\frac{8(21A - 8B) \cos^8\left(\frac{c}{2} + \frac{dx}{2}\right) \log\left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right) - \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d(a + a \cos(c + dx))^4}$$

$$+ \frac{8(21A - 8B) \cos^8\left(\frac{c}{2} + \frac{dx}{2}\right) \log\left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right) + \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d(a + a \cos(c + dx))^4}$$

$$+ \frac{\cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sec\left(\frac{c}{2}\right) \sec(c) \sec^2(c + dx) (73206A \sin\left(\frac{dx}{2}\right) - 38668B \sin\left(\frac{dx}{2}\right) - 166668A \sin\left(\frac{3dx}{2}\right) + 64384B \sin\left(\frac{3dx}{2}\right))}{d(a + a \cos(c + dx))^4}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^4,x]`

output

```
(-8*(21*A - 8*B)*Cos[c/2 + (d*x)/2]^8*Log[Cos[c/2 + (d*x)/2] - Sin[c/2 + (d*x)/2]]/(d*(a + a*Cos[c + d*x])^4) + (8*(21*A - 8*B)*Cos[c/2 + (d*x)/2]^8*Log[Cos[c/2 + (d*x)/2] + Sin[c/2 + (d*x)/2]]/(d*(a + a*Cos[c + d*x])^4) + (Cos[c/2 + (d*x)/2]*Sec[c/2]*Sec[c]*Sec[c + d*x]^2*(73206*A*Sin[(d*x)/2] - 38668*B*Sin[(d*x)/2] - 166668*A*Sin[(3*d*x)/2] + 64384*B*Sin[(3*d*x)/2] + 183162*A*Sin[c - (d*x)/2] - 70896*B*Sin[c - (d*x)/2] - 100842*A*Sin[c + (d*x)/2] + 50316*B*Sin[c + (d*x)/2] + 155526*A*Sin[2*c + (d*x)/2] - 59248*B*Sin[2*c + (d*x)/2] + 37380*A*Sin[c + (3*d*x)/2] - 22820*B*Sin[c + (3*d*x)/2] - 101148*A*Sin[2*c + (3*d*x)/2] + 48004*B*Sin[2*c + (3*d*x)/2] + 102900*A*Sin[3*c + (3*d*x)/2] - 39200*B*Sin[3*c + (3*d*x)/2] - 119364*A*Sin[c + (5*d*x)/2] + 46032*B*Sin[c + (5*d*x)/2] + 8820*A*Sin[2*c + (5*d*x)/2] - 8750*B*Sin[2*c + (5*d*x)/2] - 78204*A*Sin[3*c + (5*d*x)/2] + 35742*B*Sin[3*c + (5*d*x)/2] + 49980*A*Sin[4*c + (5*d*x)/2] - 19040*B*Sin[4*c + (5*d*x)/2] - 64053*A*Sin[2*c + (7*d*x)/2] + 24664*B*Sin[2*c + (7*d*x)/2] - 3885*A*Sin[3*c + (7*d*x)/2] - 1050*B*Sin[3*c + (7*d*x)/2] - 44733*A*Sin[4*c + (7*d*x)/2] + 19834*B*Sin[4*c + (7*d*x)/2] + 15435*A*Sin[5*c + (7*d*x)/2] - 5880*B*Sin[5*c + (7*d*x)/2] - 21987*A*Sin[3*c + (9*d*x)/2] + 8456*B*Sin[3*c + (9*d*x)/2] - 3675*A*Sin[4*c + (9*d*x)/2] + 630*B*Sin[4*c + (9*d*x)/2] - 16107*A*Sin[5*c + (9*d*x)/2] + 6986*B*Sin[5*c + (9*d*x)/2] + 2205*A*Sin[6*c + (9*d*x)/2] - 840*B*Sin[6*c + (9*d*x)/2] - 3456*A*Sin[4*c + (11*d...
```


3.73.3 Rubi [A] (verified)

Time = 1.74 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.06, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3457, 3042, 3457, 3042, 3457, 3042, 3457, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a\sin(c+dx+\frac{\pi}{2})+a)^4} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(a(9A-2B)-5a(A-B)\cos(c+dx))\sec^3(c+dx)}{(\cos(c+dx)a+a)^3} dx}{7a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(9A-2B)-5a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)^3} dx}{7a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(a^2(73A-24B)-28a^2(2A-B)\cos(c+dx))\sec^3(c+dx)}{(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a^2(73A-24B)-28a^2(2A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{a^2(73A-24B)-28a^2(2A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{a^2(73A-24B)-28a^2(2A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{5a^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4}
 \end{aligned}$$

3.73. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^4} dx$

$$\frac{\int \frac{a^3(477A-176B)-3a^3(129A-52B)\cos(c+dx)}{\cos(c+dx)a+a} \sec^3(c+dx) dx - \frac{(129A-52B)\tan(c+dx)\sec(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}}{5a^2} = \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3042

$$\frac{\int \frac{a^3(477A-176B)-3a^3(129A-52B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(\sin(c+dx+\frac{\pi}{2})a+a)} dx - \frac{(129A-52B)\tan(c+dx)\sec(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}}{5a^2} = \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3457

$$\frac{\int \frac{(105a^4(21A-8B)-8a^4(216A-83B)\cos(c+dx))\sec^3(c+dx) dx - \frac{4a^3(216A-83B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{(129A-52B)\tan(c+dx)\sec(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}}{3a^2} = \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3042

$$\frac{\int \frac{105a^4(21A-8B)-8a^4(216A-83B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3} dx - \frac{4a^3(216A-83B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{(129A-52B)\tan(c+dx)\sec(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}}{3a^2} = \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3227

$$\frac{\frac{105a^4(21A-8B)}{a^2} \int \sec^3(c+dx) dx - \frac{8a^4(216A-83B)}{3a^2} \int \sec^2(c+dx) dx - \frac{4a^3(216A-83B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{(129A-52B)\tan(c+dx)\sec(c+dx)}{3d(\cos(c+dx)+1)^2} - \frac{7a(2A-B)\tan(c+dx)\sec(c+dx)}{5d(a\cos(c+dx)+a)^3}}{5a^2} = \frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4}$$

↓ 3042

3.73. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^4} dx$

$$\frac{105a^4(21A-8B) \int \csc(c+dx+\frac{\pi}{2})^3 dx - 8a^4(216A-83B) \int \csc(c+dx+\frac{\pi}{2})^2 dx - \frac{4a^3(216A-83B) \tan(c+dx) \sec(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}}{a^2 \quad 3a^2 \quad 5a^2} - 7a$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{7d(a \cos(c+dx)+a)^4} \quad 7a^2$$

↓ 4254

$$\frac{8a^4(216A-83B) \int \frac{1d(-\tan(c+dx))}{d} + 105a^4(21A-8B) \int \csc(c+dx+\frac{\pi}{2})^3 dx - \frac{8a^4(216A-83B) \tan(c+dx)}{d} - \frac{4a^3(216A-83B) \tan(c+dx) \sec(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}}{a^2 \quad 3a^2 \quad 5a^2} - 7a$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{7d(a \cos(c+dx)+a)^4} \quad 7a^2$$

↓ 24

$$\frac{105a^4(21A-8B) \int \csc(c+dx+\frac{\pi}{2})^3 dx - \frac{8a^4(216A-83B) \tan(c+dx)}{d} - \frac{4a^3(216A-83B) \tan(c+dx) \sec(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}}{a^2 \quad 3a^2 \quad 5a^2} - \frac{7a(2A-B)}{5d(a \cos(c+dx)+a)}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{7d(a \cos(c+dx)+a)^4} \quad 7a^2$$

↓ 4255

$$\frac{105a^4(21A-8B) \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{8a^4(216A-83B) \tan(c+dx)}{d} - \frac{4a^3(216A-83B) \tan(c+dx) \sec(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}}{a^2 \quad 3a^2 \quad 5a^2} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{7d(a \cos(c+dx)+a)^4} \quad 7a^2$$

↓ 3042

$$\frac{105a^4(21A-8B) \left(\frac{1}{2} \int \csc(c+dx+\frac{\pi}{2}) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) - \frac{8a^4(216A-83B) \tan(c+dx)}{d} - \frac{4a^3(216A-83B) \tan(c+dx) \sec(c+dx)}{d(a \cos(c+dx)+a)} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}}{a^2 \quad 3a^2 \quad 5a^2} - \frac{(129A-52B) \tan(c+dx) \sec(c+dx)}{3d(\cos(c+dx)+1)^2}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{7d(a \cos(c+dx)+a)^4} \quad 7a^2$$

↓ 4257

3.73. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^4} dx$

$$\frac{105a^4(21A-8B)\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx)\sec(c+dx)}{2d}\right) - \frac{8a^4(216A-83B)\tan(c+dx)}{d}}{a^2} - \frac{4a^3(216A-83B)\tan(c+dx)\sec(c+dx)}{d(a\cos(c+dx)+a)} - \frac{(129A-52B)\tan(c+dx)}{3d(\cos(c+dx)+1)}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{7d(a\cos(c+dx)+a)^4} \quad 7a^2$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^4,x]`

output `-1/7*((A - B)*Sec[c + d*x]*Tan[c + d*x]/(d*(a + a*Cos[c + d*x])^4) + ((-7 *a*(2*A - B)*Sec[c + d*x]*Tan[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + (-1 /3*((129*A - 52*B)*Sec[c + d*x]*Tan[c + d*x])/(d*(1 + Cos[c + d*x])^2) + ((-4*a^3*(216*A - 83*B)*Sec[c + d*x]*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])) + ((-8*a^4*(216*A - 83*B)*Tan[c + d*x])/d + 105*a^4*(21*A - 8*B)*(ArcTanh [Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/a^2)/(3*a^2))/(5*a^2))/(7*a^2)`

3.73.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int [(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
  Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 4254 Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4255 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
  /; FreeQ[{c, d}, x]
```

3.73.4 Maple [A] (verified)

Time = 1.54 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.83

$$3.73. \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^4} dx$$

method	result
parallelrisch	$-70560(1+\cos(2dx+2c))\left(A-\frac{8B}{21}\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+70560(1+\cos(2dx+2c))\left(A-\frac{8B}{21}\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)-11619$
derivativedivides	$-\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{7}+\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{7}-\frac{9A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}+\frac{7\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{5}-13\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A+\frac{23\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3}$
default	$-\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A}{7}+\frac{\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{7}-\frac{9A\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5}+\frac{7\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B}{5}-13\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)A+\frac{23\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3}$
norman	$\frac{(A-B)\left(\tan^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{56ad}-\frac{(29A-22B)\left(\tan^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{140ad}-\frac{(167A-65B)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{8ad}+\frac{(171A-62B)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{12ad}-\frac{(1161A-6$
risch	$\frac{i\left(2205Ae^{10i(dx+c)}-840Be^{10i(dx+c)}+15435Ae^{9i(dx+c)}-5880Be^{9i(dx+c)}+49980Ae^{8i(dx+c)}-19040Be^{8i(dx+c)}+10$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a)^4,x,method=_RETURNVERBOSE)
```

```
output 1/6720*(-70560*(1+cos(2*d*x+2*c))*(A-8/21*B)*ln(tan(1/2*d*x+1/2*c)-1)+70560*(1+cos(2*d*x+2*c))*(A-8/21*B)*ln(tan(1/2*d*x+1/2*c)+1)-11619*tan(1/2*d*x+1/2*c)*((23540/3873*A-3040/1291*B)*cos(2*d*x+2*c)+(3992/1291*A-13864/11619*B)*cos(3*d*x+3*c)+(A-4472/11619*B)*cos(4*d*x+4*c)+(192/1291*A-664/11619*B)*cos(5*d*x+5*c)+(34168/3873*A-39952/11619*B)*cos(d*x+c)+19387/3873*A-22888/11619*B)*sec(1/2*d*x+1/2*c)^6)/d/a^4/(1+cos(2*d*x+2*c))
```

3.73.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.55

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{105 \left((21A - 8B) \cos(dx + c)^6 + 4(21A - 8B) \cos(dx + c)^5 + 6(21A - 8B) \cos(dx + c)^4 + 4(21A - 8B) \cos(dx + c)^3 + 4(21A - 8B) \cos(dx + c)^2 + 4(21A - 8B) \cos(dx + c) + 4(21A - 8B) \right)}{(a + a \cos(dx + c))^4}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^4,x, algorithm="fracas")
```

3.73. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^4} dx$

```
output 1/420*(105*((21*A - 8*B)*cos(d*x + c)^6 + 4*(21*A - 8*B)*cos(d*x + c)^5 +
6*(21*A - 8*B)*cos(d*x + c)^4 + 4*(21*A - 8*B)*cos(d*x + c)^3 + (21*A - 8*
B)*cos(d*x + c)^2)*log(sin(d*x + c) + 1) - 105*((21*A - 8*B)*cos(d*x + c)^
6 + 4*(21*A - 8*B)*cos(d*x + c)^5 + 6*(21*A - 8*B)*cos(d*x + c)^4 + 4*(21*
A - 8*B)*cos(d*x + c)^3 + (21*A - 8*B)*cos(d*x + c)^2)*log(-sin(d*x + c) +
1) - 2*(16*(216*A - 83*B)*cos(d*x + c)^5 + (11619*A - 4472*B)*cos(d*x + c
)^4 + 4*(3411*A - 1318*B)*cos(d*x + c)^3 + 4*(1509*A - 592*B)*cos(d*x + c)
^2 + 210*(2*A - B)*cos(d*x + c) - 105*A)*sin(d*x + c))/(a^4*d*cos(d*x + c)
^6 + 4*a^4*d*cos(d*x + c)^5 + 6*a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c
)^3 + a^4*d*cos(d*x + c)^2)
```

3.73.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{\int \frac{A \sec^3(c+dx)}{\cos^4(c+dx)+4 \cos^3(c+dx)+6 \cos^2(c+dx)+4 \cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sec^3(c+dx)}{\cos^4(c+dx)+4 \cos^3(c+dx)+6 \cos^2(c+dx)+4 \cos(c+dx)+1} dx}{a^4}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c))**4,x)
```

```
output (Integral(A*sec(c + d*x)**3/(cos(c + d*x)**4 + 4*cos(c + d*x)**3 + 6*cos(c
+ d*x)**2 + 4*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sec(c + d*x)
)**3/(cos(c + d*x)**4 + 4*cos(c + d*x)**3 + 6*cos(c + d*x)**2 + 4*cos(c +
d*x) + 1), x))/a**4
```

3.73.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 419, normalized size of antiderivative = 1.81

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx =$$

$$3A \left(\frac{280 \left(\frac{7 \sin(dx+c)}{\cos(dx+c)+1} - \frac{9 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{a^4 - \frac{2a^4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^4 \sin(dx+c)^4}{(\cos(dx+c)+1)^4}} + \frac{3885 \sin(dx+c)}{\cos(dx+c)+1} + \frac{455 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{63 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + \frac{5 \sin(dx+c)^7}{(\cos(dx+c)+1)^7} - \frac{2940 \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{a^4} \right)$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^4,x, algorithm="m
axima")
```

3.73. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^4} dx$

output
$$\begin{aligned} & -1/840*(3*A*(280*(7*\sin(d*x + c))/(\cos(d*x + c) + 1) - 9*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3)/(a^4 - 2*a^4*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + a^4*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4) + (3885*\sin(d*x + c)/(\cos(d*x + c) + 1) + 455*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 63*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 + 5*\sin(d*x + c)^7/(\cos(d*x + c) + 1)^7)/a^4 - 2940*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)/a^4 + 2940*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) - 1)/a^4 - B*(1680*\sin(d*x + c)/((a^4 - a^4*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2)*(cos(d*x + c) + 1)) + (5145*\sin(d*x + c)/(\cos(d*x + c) + 1) + 805*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 147*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 + 15*\sin(d*x + c)^7/(\cos(d*x + c) + 1)^7)/a^4 - 3360*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)/a^4 + 3360*\log(\sin(d*x + c)/(\cos(d*x + c) + 1) - 1)/a^4))/d \end{aligned}$$

3.73.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.15

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx$$

$$= \frac{420(21A - 8B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^4} - \frac{420(21A - 8B) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^4} + \frac{840(9A \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 2B \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 7A \tan(\frac{1}{2} dx + \frac{1}{2} c) + 2B)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1)^2 a^4}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^4,x, algorithm="giac")`

output
$$\begin{aligned} & 1/840*(420*(21*A - 8*B)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1))/a^4 - 420*(21*A - 8*B)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1))/a^4 + 840*(9*A*\tan(1/2*d*x + 1/2*c)^3 - 2*B*\tan(1/2*d*x + 1/2*c)^3 - 7*A*\tan(1/2*d*x + 1/2*c) + 2*B*\tan(1/2*d*x + 1/2*c))/((\tan(1/2*d*x + 1/2*c)^2 - 1)^2*a^4) - (15*A*a^24*\tan(1/2*d*x + 1/2*c)^7 - 15*B*a^24*\tan(1/2*d*x + 1/2*c)^7 + 189*A*a^24*\tan(1/2*d*x + 1/2*c)^5 - 147*B*a^24*\tan(1/2*d*x + 1/2*c)^5 + 1365*A*a^24*\tan(1/2*d*x + 1/2*c)^3 - 805*B*a^24*\tan(1/2*d*x + 1/2*c)^3 + 11655*A*a^24*\tan(1/2*d*x + 1/2*c) - 5145*B*a^24*\tan(1/2*d*x + 1/2*c))/a^28)/d \end{aligned}$$

3.73.9 Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.18

$$\begin{aligned}
& \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^4} dx \\
&= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (9A - 2B) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (7A - 2B)}{d \left(a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2a^4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a^4 \right)} \\
&\quad - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{5A}{2a^4} + \frac{5(A-B)}{4a^4} + \frac{3(6A-4B)}{4a^4} + \frac{3(15A-5B)}{8a^4} \right)}{d} \\
&\quad - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{A-B}{4a^4} + \frac{6A-4B}{8a^4} + \frac{15A-5B}{24a^4} \right)}{d} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 \left(\frac{3(A-B)}{40a^4} + \frac{6A-4B}{40a^4} \right)}{d} \\
&\quad - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 (A - B)}{56a^4 d} + \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (21A - 8B)}{a^4 d}
\end{aligned}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^4),x)`output `(tan(c/2 + (d*x)/2)^3*(9*A - 2*B) - tan(c/2 + (d*x)/2)*(7*A - 2*B))/(d*(a^4*tan(c/2 + (d*x)/2)^4 - 2*a^4*tan(c/2 + (d*x)/2)^2 + a^4) - (tan(c/2 + (d*x)/2)*((5*A)/(2*a^4) + (5*(A - B))/(4*a^4) + (3*(6*A - 4*B))/(4*a^4) + (3*(15*A - 5*B))/(8*a^4)))/d - (tan(c/2 + (d*x)/2)^3*((A - B)/(4*a^4) + (6*A - 4*B)/(8*a^4) + (15*A - 5*B)/(24*a^4)))/d - (tan(c/2 + (d*x)/2)^5*((3*(A - B))/(40*a^4) + (6*A - 4*B)/(40*a^4)))/d - (tan(c/2 + (d*x)/2)^7*(A - B))/(56*a^4*d) + (atanh(tan(c/2 + (d*x)/2))*(21*A - 8*B))/(a^4*d)`

3.74 $\int \cos^3(c+dx) \sqrt{a + a \cos(c + dx)}(A+B \cos(c+dx)) dx$

3.74.1	Optimal result	819
3.74.2	Mathematica [A] (verified)	820
3.74.3	Rubi [A] (verified)	820
3.74.4	Maple [A] (verified)	823
3.74.5	Fricas [A] (verification not implemented)	824
3.74.6	Sympy [F(-1)]	824
3.74.7	Maxima [A] (verification not implemented)	825
3.74.8	Giac [A] (verification not implemented)	825
3.74.9	Mupad [F(-1)]	826

3.74.1 Optimal result

Integrand size = 33, antiderivative size = 187

$$\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{4a(9A + 8B) \sin(c + dx)}{45d \sqrt{a + a \cos(c + dx)}} + \frac{2a(9A + 8B) \cos^3(c + dx) \sin(c + dx)}{63d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{2aB \cos^4(c + dx) \sin(c + dx)}{9d \sqrt{a + a \cos(c + dx)}} - \frac{8(9A + 8B) \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{315d}$$

$$+ \frac{4(9A + 8B)(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{105ad}$$

output $\frac{4}{105}*(9*A+8*B)*(a+a*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/a/d+4/45*a*(9*A+8*B)*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/63*a*(9*A+8*B)*\cos(d*x+c)^3*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/9*a*B*\cos(d*x+c)^4*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}-8/315*(9*A+8*B)*\sin(d*x+c)*(a+a*\cos(d*x+c))^{(1/2)}/d$

3.74.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.55

$$\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} (1368A + 1321B + 94(9A + 8B) \cos(c + dx) + 4(54A + 83B) \cos(2(c + dx)) + 90A)}{1260d}$$

input `Integrate[Cos[c + d*x]^3*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])]*(1368*A + 1321*B + 94*(9*A + 8*B)*Cos[c + d*x] + 4*(54*A + 83*B)*Cos[2*(c + d*x)] + 90*A*Cos[3*(c + d*x)] + 80*B*Cos[3*(c + d*x)] + 35*B*Cos[4*(c + d*x)])*Tan[(c + d*x)/2])/(1260*d)`

3.74.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3460, 3042, 3249, 3042, 3238, 27, 3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^3 \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} (A + B \sin\left(c + dx + \frac{\pi}{2}\right)) dx$$

$$\downarrow \text{3460}$$

$$\frac{1}{9}(9A + 8B) \int \cos^3(c + dx) \sqrt{\cos(c + dx)a + a} dx + \frac{2aB \sin(c + dx) \cos^4(c + dx)}{9d \sqrt{a \cos(c + dx) + a}}$$

$$\downarrow \text{3042}$$

$$\frac{1}{9}(9A + 8B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^3 \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a} dx + \frac{2aB \sin(c + dx) \cos^4(c + dx)}{9d \sqrt{a \cos(c + dx) + a}}$$

$$\downarrow \text{3249}$$

3.74. $\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \frac{1}{9}(9A+8B) \left(\frac{6}{7} \int \cos^2(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9}(9A+8B) \left(\frac{6}{7} \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3238} \\
& 8B) \left(\frac{6}{7} \left(\frac{2 \int \frac{1}{2}(3a-2a \cos(c+dx)) \sqrt{\cos(c+dx)a+adx}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{27} \\
& 8B) \left(\frac{6}{7} \left(\frac{\int (3a-2a \cos(c+dx)) \sqrt{\cos(c+dx)a+adx}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& 8B) \left(\frac{6}{7} \left(\frac{\int (3a-2a \sin\left(c+dx+\frac{\pi}{2}\right)) \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3230} \\
& 8B) \left(\frac{6}{7} \left(\frac{\frac{7}{3}a \int \sqrt{\cos(c+dx)a+adx} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d\sqrt{a \cos(c+dx)+a}}
\end{aligned}$$

3.74. $\int \cos^3(c+dx) \sqrt{a+a \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 8B) & \left(\frac{6}{7} \left(\frac{\frac{7}{3}a \int \sqrt{\sin(c+dx+\frac{\pi}{2})} a + adx - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) + \frac{2aB \sin(c+dx) \cos^4(c+dx)}{9d \sqrt{a \cos(c+dx)+a}} \right) \\
 & \downarrow 3125 \\
 8B) & \left(\frac{6}{7} \left(\frac{\frac{14a^2 \sin(c+dx)}{3d \sqrt{a \cos(c+dx)+a}} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) + \frac{2a \sin(c+dx) \cos^4(c+dx)}{7d \sqrt{a \cos(c+dx)+a}} \right)
 \end{aligned}$$

input `Int[Cos[c + d*x]^3*sqrt[a + a*cos[c + d*x]]*(A + B*cos[c + d*x]),x]`

output `(2*a*B*cos[c + d*x]^4*sin[c + d*x])/(9*d*sqrt[a + a*cos[c + d*x]]) + ((9*A + 8*B)*((2*a*cos[c + d*x]^3*sin[c + d*x])/(7*d*sqrt[a + a*cos[c + d*x]])) + (6*((2*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(5*a*d) + ((14*a^2*sin[c + d*x])/(3*d*sqrt[a + a*cos[c + d*x]]) - (4*a*sqrt[a + a*cos[c + d*x]]*sin[c + d*x])/(3*d))/(5*a)))/7)/9`

3.74.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*sqrt[a + b*sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

```
rule 3230 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

```
rule 3238 Int[sin[(e_) + (f_)*(x_)]^2*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_),
x_Symbol] := Simp[(-Cos[e + f*x])*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2
))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*(b*(m + 1) - a*Si
n[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !
LtQ[m, -2^(-1)]
```

```
rule 3249 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])
^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(
2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]
```

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.74.4 Maple [A] (verified)

Time = 2.71 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.65

method	result
default	$\frac{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(560B \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-360A - 1440B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (756A + 1512B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-630A - 840B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 315B}{315 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$
parts	$\frac{2A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(40 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 36 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 22 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 9\right) \sqrt{2}}{35 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d + \frac{2B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$

3.74. $\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

input `int(cos(d*x+c)^3*(a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `2/315*cos(1/2*d*x+1/2*c)*a*sin(1/2*d*x+1/2*c)*(560*B*sin(1/2*d*x+1/2*c)^8+
(-360*A-1440*B)*sin(1/2*d*x+1/2*c)^6+(756*A+1512*B)*sin(1/2*d*x+1/2*c)^4+(-
630*A-840*B)*sin(1/2*d*x+1/2*c)^2+315*A+315*B)*2^(1/2)/(a*cos(1/2*d*x+1/2
*c)^2)^(1/2)/d`

3.74.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.53

$$\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{2(35B \cos(dx + c)^4 + 5(9A + 8B) \cos(dx + c)^3 + 6(9A + 8B) \cos(dx + c)^2 + 8(9A + 8B) \cos(dx + c) + 144A + 128B) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{315(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm
m="fricas")`

output `2/315*(35*B*cos(d*x + c)^4 + 5*(9*A + 8*B)*cos(d*x + c)^3 + 6*(9*A + 8*B)*
cos(d*x + c)^2 + 8*(9*A + 8*B)*cos(d*x + c) + 144*A + 128*B)*sqrt(a*cos(d*
x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

3.74.6 Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

$$3.74. \quad \int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

3.74.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.78

$$\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{18 \left(5 \sqrt{2} \sin \left(\frac{7}{2} dx + \frac{7}{2} c \right) + 7 \sqrt{2} \sin \left(\frac{5}{2} dx + \frac{5}{2} c \right) + 35 \sqrt{2} \sin \left(\frac{3}{2} dx + \frac{3}{2} c \right) + 105 \sqrt{2} \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) A \sqrt{a}}{d}$$

```
input integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm
m="maxima")
```

```
output 1/2520*(18*(5*sqrt(2)*sin(7/2*d*x + 7/2*c) + 7*sqrt(2)*sin(5/2*d*x + 5/2*c)
) + 35*sqrt(2)*sin(3/2*d*x + 3/2*c) + 105*sqrt(2)*sin(1/2*d*x + 1/2*c))*A*
sqrt(a) + (35*sqrt(2)*sin(9/2*d*x + 9/2*c) + 45*sqrt(2)*sin(7/2*d*x + 7/2*
c) + 252*sqrt(2)*sin(5/2*d*x + 5/2*c) + 420*sqrt(2)*sin(3/2*d*x + 3/2*c) +
1890*sqrt(2)*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d
```

3.74.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.97

$$\int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2} \left(35 B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \sin \left(\frac{9}{2} dx + \frac{9}{2} c \right) + 45 \left(2 A \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) + B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \right) \right)}{d}$$

```
input integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm
m="giac")
```

```
output 1/2520*sqrt(2)*(35*B*sgn(cos(1/2*d*x + 1/2*c))*sin(9/2*d*x + 9/2*c) + 45*(
2*A*sgn(cos(1/2*d*x + 1/2*c)) + B*sgn(cos(1/2*d*x + 1/2*c)))*sin(7/2*d*x +
7/2*c) + 126*(A*sgn(cos(1/2*d*x + 1/2*c)) + 2*B*sgn(cos(1/2*d*x + 1/2*c))
)*sin(5/2*d*x + 5/2*c) + 210*(3*A*sgn(cos(1/2*d*x + 1/2*c)) + 2*B*sgn(cos(
1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 1890*(A*sgn(cos(1/2*d*x + 1/2*c)
) + B*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d
```


3.74.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cos^3(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int \cos(c + dx)^3 (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx \end{aligned}$$

input `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)`output `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)`

3.75 $\int \cos^2(c+dx) \sqrt{a + a \cos(c + dx)}(A+B \cos(c+dx)) dx$

3.75.1 Optimal result	827
3.75.2 Mathematica [A] (verified)	828
3.75.3 Rubi [A] (verified)	828
3.75.4 Maple [A] (verified)	831
3.75.5 Fricas [A] (verification not implemented)	831
3.75.6 Sympy [F]	832
3.75.7 Maxima [A] (verification not implemented)	832
3.75.8 Giac [A] (verification not implemented)	833
3.75.9 Mupad [F(-1)]	833

3.75.1 Optimal result

Integrand size = 33, antiderivative size = 144

$$\begin{aligned} & \int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx \\ &= \frac{2a(7A + 6B) \sin(c + dx)}{15d \sqrt{a + a \cos(c + dx)}} + \frac{2aB \cos^3(c + dx) \sin(c + dx)}{7d \sqrt{a + a \cos(c + dx)}} \\ & \quad - \frac{4(7A + 6B) \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{105d} \\ & \quad + \frac{2(7A + 6B)(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{35ad} \end{aligned}$$

```
output 2/35*(7*A+6*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/a/d+2/15*a*(7*A+6*B)*sin(
d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/7*a*B*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos
(d*x+c))^(1/2)-4/105*(7*A+6*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.75.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.56

$$\int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} (266A + 228B + (112A + 141B) \cos(c + dx) + 6(7A + 6B) \cos(2(c + dx)) + 15B \cos(3(c + dx))) \tan((c + dx)/2)}{210d}$$

input `Integrate[Cos[c + d*x]^2*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])]*(266*A + 228*B + (112*A + 141*B)*Cos[c + d*x] + 6*(7*A + 6*B)*Cos[2*(c + d*x)] + 15*B*Cos[3*(c + d*x)])*Tan[(c + d*x)/2])/(210*d)`

3.75.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 3460, 3042, 3238, 27, 3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3460}$$

$$\frac{1}{7}(7A + 6B) \int \cos^2(c + dx) \sqrt{\cos(c + dx)a + a} dx + \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d \sqrt{a \cos(c + dx) + a}}$$

$$\downarrow \text{3042}$$

$$\frac{1}{7}(7A + 6B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a} dx + \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d \sqrt{a \cos(c + dx) + a}}$$

$$\downarrow \text{3238}$$

3.75. $\int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \frac{1}{7}(7A + \\
6B) & \left(\frac{2 \int \frac{1}{2}(3a - 2a \cos(c + dx)) \sqrt{\cos(c + dx)a + adx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) + \\
& \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d\sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow 27 \\
& \frac{1}{7}(7A + \\
6B) & \left(\frac{\int (3a - 2a \cos(c + dx)) \sqrt{\cos(c + dx)a + adx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) + \\
& \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d\sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow 3042 \\
& \frac{1}{7}(7A + \\
6B) & \left(\frac{\int (3a - 2a \sin(c + dx + \frac{\pi}{2})) \sqrt{\sin(c + dx + \frac{\pi}{2})a + adx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) + \\
& \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d\sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow 3230 \\
& \frac{1}{7}(7A + \\
6B) & \left(\frac{\frac{7}{3}a \int \sqrt{\cos(c + dx)a + adx} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) + \\
& \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d\sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow 3042 \\
& \frac{1}{7}(7A + \\
6B) & \left(\frac{\frac{7}{3}a \int \sqrt{\sin(c + dx + \frac{\pi}{2})a + adx} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) + \\
& \frac{2aB \sin(c + dx) \cos^3(c + dx)}{7d\sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow 3125
\end{aligned}$$

$$\frac{1}{7}(7A+6B) \left(\frac{\frac{14a^2 \sin(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} - \frac{4a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{5ad} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}}$$

input `Int[Cos[c + d*x]^2*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(2*a*B*Cos[c + d*x]^3*Sin[c + d*x])/(7*d*Sqrt[a + a*Cos[c + d*x]]) + ((7*A + 6*B)*((2*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*a*d) + ((14*a^2*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) - (4*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(5*a)))/7`

3.75.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3238 `Int[sin[(e_) + (f_)*(x_)]^2*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*(b*(m + 1) - a*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.75.4 Maple [A] (verified)

Time = 2.76 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.71

method	result
default	$\frac{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(-120B \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (84A + 252B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-140A - 210B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 105A + 105B}{105 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{2A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(12 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 4 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 7\right) \sqrt{2}}{15 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{2B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(40 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 35 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{35 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

```
input int(cos(d*x+c)^2*(a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 2/105*cos(1/2*d*x+1/2*c)*a*sin(1/2*d*x+1/2*c)*(-120*B*sin(1/2*d*x+1/2*c)^6
+(84*A+252*B)*sin(1/2*d*x+1/2*c)^4+(-140*A-210*B)*sin(1/2*d*x+1/2*c)^2+105
*A+105*B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.75.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.57

$$\int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{2 (15 B \cos(dx + c)^3 + 3 (7 A + 6 B) \cos(dx + c)^2 + 4 (7 A + 6 B) \cos(dx + c) + 56 A + 48 B) \sqrt{a \cos(dx + c)}}{105 (d \cos(dx + c) + d)}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorith
m="fracas")
```

3.75. $\int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

output $2/105*(15*B*\cos(d*x + c)^3 + 3*(7*A + 6*B)*\cos(d*x + c)^2 + 4*(7*A + 6*B)*\cos(d*x + c) + 56*A + 48*B)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/(d*\cos(d*x + c) + d)$

3.75.6 Sympy [F]

$$\begin{aligned} & \int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int \sqrt{a (\cos(c + dx) + 1)} (A + B \cos(c + dx)) \cos^2(c + dx) dx \end{aligned}$$

input `integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))*cos(c + d*x)**2, x)`

3.75.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.82

$$\begin{aligned} & \int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \frac{14 \left(3 \sqrt{2} \sin\left(\frac{5}{2} dx + \frac{5}{2} c\right) + 5 \sqrt{2} \sin\left(\frac{3}{2} dx + \frac{3}{2} c\right) + 30 \sqrt{2} \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right) \right) A \sqrt{a} + 3 \left(5 \sqrt{2} \sin\left(\frac{7}{2} dx + \frac{7}{2} c\right) \right.}{420 d} \end{aligned}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/420*(14*(3*sqrt(2)*sin(5/2*d*x + 5/2*c) + 5*sqrt(2)*sin(3/2*d*x + 3/2*c) + 30*sqrt(2)*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + 3*(5*sqrt(2)*sin(7/2*d*x + 7/2*c) + 7*sqrt(2)*sin(5/2*d*x + 5/2*c) + 35*sqrt(2)*sin(3/2*d*x + 3/2*c) + 105*sqrt(2)*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d`

3.75.8 Giac [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.02

$$\int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2} (15 B \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{7}{2} dx + \frac{7}{2} c) + 21 (2 A \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + B \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))))}{2}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm
m="giac")
```

```
output 1/420*sqrt(2)*(15*B*sgn(cos(1/2*d*x + 1/2*c))*sin(7/2*d*x + 7/2*c) + 21*(2
*A*sgn(cos(1/2*d*x + 1/2*c)) + B*sgn(cos(1/2*d*x + 1/2*c)))*sin(5/2*d*x +
5/2*c) + 35*(2*A*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*sgn(cos(1/2*d*x + 1/2*c))
)*sin(3/2*d*x + 3/2*c) + 105*(4*A*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*sgn(cos(
1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d
```

3.75.9 Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \cos(c + dx)^2 (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx$$

```
input int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)
```

```
output int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)
```


3.76 $\int \cos(c+dx) \sqrt{a + a \cos(c + dx)} (A+B \cos(c+dx)) dx$

3.76.1	Optimal result	834
3.76.2	Mathematica [A] (verified)	834
3.76.3	Rubi [A] (verified)	835
3.76.4	Maple [A] (verified)	837
3.76.5	Fricas [A] (verification not implemented)	838
3.76.6	Sympy [F]	838
3.76.7	Maxima [A] (verification not implemented)	838
3.76.8	Giac [A] (verification not implemented)	839
3.76.9	Mupad [F(-1)]	839

3.76.1 Optimal result

Integrand size = 31, antiderivative size = 101

$$\begin{aligned} & \int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \frac{2a(5A + 7B) \sin(c + dx)}{15d \sqrt{a + a \cos(c + dx)}} + \frac{2(5A - 2B) \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{15d} \\ &+ \frac{2B(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{5ad} \end{aligned}$$

```
output 2/5*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/a/d+2/15*a*(5*A+7*B)*sin(d*x+c)/d/
(a+a*cos(d*x+c))^(1/2)+2/15*(5*A-2*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.76.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\begin{aligned} & \int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \frac{\sqrt{a(1 + \cos(c + dx))} (20A + 19B + 2(5A + 4B) \cos(c + dx) + 3B \cos(2(c + dx))) \tan\left(\frac{1}{2}(c + dx)\right)}{15d} \end{aligned}$$

```
input Integrate[Cos[c + d*x]*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]
```

output $(\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*(20*A + 19*B + 2*(5*A + 4*B)*\text{Cos}[c + d*x] + 3*B*\text{Cos}[2*(c + d*x)])*\text{Tan}[(c + d*x)/2])/(15*d)$

3.76.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sqrt{a \cos(c + dx) + a} (A \cos(c + dx) + B \cos^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{2 \int \frac{1}{2} \sqrt{\cos(c + dx)a + a} (3aB + a(5A - 2B) \cos(c + dx)) dx}{5a} + \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \sqrt{\cos(c + dx)a + a} (3aB + a(5A - 2B) \cos(c + dx)) dx}{5a} + \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a} (3aB + a(5A - 2B) \sin\left(c + dx + \frac{\pi}{2}\right)) dx}{5a} + \\
 & \quad \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \\
 & \quad \downarrow \text{3230}
 \end{aligned}$$

3.76. $\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \frac{\frac{1}{3}a(5A+7B) \int \sqrt{\cos(c+dx)a+adx} + \frac{2a(5A-2B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \\
& \frac{2B\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{5ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{3}a(5A+7B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})a+adx} + \frac{2a(5A-2B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \\
& \frac{2B\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{5ad} \\
& \quad \downarrow \text{3125} \\
& \frac{\frac{2a^2(5A+7B)\sin(c+dx)}{3d\sqrt{a\cos(c+dx)+a}} + \frac{2a(5A-2B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \frac{2B\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{5ad}
\end{aligned}$$

input `Int[Cos[c + d*x]*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(2*B*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*a*d) + ((2*a^2*(5*A + 7*B)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (2*a*(5*A - 2*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(5*a)`

3.76.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

```
rule 3230 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.76.4 Maple [A] (verified)

Time = 2.71 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.82

method	result
default	$\frac{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(12B \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-10A - 20B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 15A + 15B\right) \sqrt{2}}{15 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{2A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(1 + 2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \sqrt{2}}{3 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{2B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(12 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 4 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 3\right) \sqrt{2}}{15 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 2/15*cos(1/2*d*x+1/2*c)*a*sin(1/2*d*x+1/2*c)*(12*B*sin(1/2*d*x+1/2*c)^4+(-
10*A-20*B)*sin(1/2*d*x+1/2*c)^2+15*A+15*B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2
)^(1/2)/d
```

3.76. $\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.76.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{2(3B \cos(dx + c)^2 + (5A + 4B) \cos(dx + c) + 10A + 8B) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{15(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `2/15*(3*B*cos(d*x + c)^2 + (5*A + 4*B)*cos(d*x + c) + 10*A + 8*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

3.76.6 Sympy [F]

$$\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \sqrt{a(\cos(c + dx) + 1)} (A + B \cos(c + dx)) \cos(c + dx) dx$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))*cos(c + d*x), x)`

3.76.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.87

$$\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{10(\sqrt{2} \sin(\frac{3}{2} dx + \frac{3}{2} c) + 3\sqrt{2} \sin(\frac{1}{2} dx + \frac{1}{2} c)) A \sqrt{a} + (3\sqrt{2} \sin(\frac{5}{2} dx + \frac{5}{2} c) + 5\sqrt{2} \sin(\frac{3}{2} dx + \frac{3}{2} c) + 3\sqrt{2} \sin(\frac{1}{2} dx + \frac{1}{2} c)) B \sqrt{a}}{30d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/30*(10*(sqrt(2)*sin(3/2*d*x + 3/2*c) + 3*sqrt(2)*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + (3*sqrt(2)*sin(5/2*d*x + 5/2*c) + 5*sqrt(2)*sin(3/2*d*x + 3/2*c) + 30*sqrt(2)*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d`

3.76.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.06

$$\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2} \left(3 B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \sin \left(\frac{5}{2} dx + \frac{5}{2} c \right) + 5 \left(2 A \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) + B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \right) \sin \left(\frac{3}{2} dx + \frac{3}{2} c \right) \right)}{30 d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/30*sqrt(2)*(3*B*sgn(cos(1/2*d*x + 1/2*c))*sin(5/2*d*x + 5/2*c) + 5*(2*A*sgn(cos(1/2*d*x + 1/2*c)) + B*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 30*(A*sgn(cos(1/2*d*x + 1/2*c)) + B*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

3.76.9 Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \cos(c + dx) (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)`

3.76. $\int \cos(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.77 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$

3.77.1	Optimal result	840
3.77.2	Mathematica [A] (verified)	840
3.77.3	Rubi [A] (verified)	841
3.77.4	Maple [A] (verified)	842
3.77.5	Fricas [A] (verification not implemented)	843
3.77.6	Sympy [F]	843
3.77.7	Maxima [A] (verification not implemented)	843
3.77.8	Giac [A] (verification not implemented)	844
3.77.9	Mupad [F(-1)]	844

3.77.1 Optimal result

Integrand size = 25, antiderivative size = 62

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{2a(3A + B) \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}} + \frac{2B\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{3d}$$

output `2/3*a*(3*A+B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/3*B*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d`

3.77.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{2\sqrt{a(1 + \cos(c + dx))}(3A + 2B + B \cos(c + dx)) \tan\left(\frac{1}{2}(c + dx)\right)}{3d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*(3*A + 2*B + B*Cos[c + d*x])*Tan[(c + d*x)/2])/ (3*d)`

3.77.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3230} \\
 & \frac{1}{3}(3A + B) \int \sqrt{\cos(c + dx)a + adx} + \frac{2B \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3}(3A + B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + adx} + \frac{2B \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{3125} \\
 & \frac{2a(3A + B) \sin(c + dx)}{3d \sqrt{a \cos(c + dx) + a}} + \frac{2B \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}
 \end{aligned}$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(2*a*(3*A + B)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (2*B*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d)`

3.77.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

3.77.4 Maple [A] (verified)

Time = 2.24 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(2B \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 3A + B\right) \sqrt{2}}{3 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	62
parts	$\frac{2Aa \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2}}{\sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{2B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(1 + 2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \sqrt{2}}{3 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$	103

input `int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `2/3*cos(1/2*d*x+1/2*c)*a*sin(1/2*d*x+1/2*c)*(2*B*cos(1/2*d*x+1/2*c)^2+3*A+B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.77.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.76

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{2(B \cos(dx + c) + 3A + 2B)\sqrt{a \cos(dx + c) + a} \sin(dx + c)}{3(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`output `2/3*(B*cos(d*x + c) + 3*A + 2*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`**3.77.6 Sympy [F]**

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \int \sqrt{a (\cos(c + dx) + 1)}(A + B \cos(c + dx)) dx$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x)), x)`**3.77.7 Maxima [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.92

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{6\sqrt{2}A\sqrt{a} \sin\left(\frac{1}{2}dx + \frac{1}{2}c\right) + \left(\sqrt{2} \sin\left(\frac{3}{2}dx + \frac{3}{2}c\right) + 3\sqrt{2} \sin\left(\frac{1}{2}dx + \frac{1}{2}c\right)\right)B\sqrt{a}}{3d}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`output `1/3*(6*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c) + (sqrt(2)*sin(3/2*d*x + 3/2*c) + 3*sqrt(2)*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d`

3.77. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$

3.77.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.13

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2}(B \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{3}{2} dx + \frac{3}{2} c) + 3(2 A \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + B \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin}{3 d}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/3*sqrt(2)*(B*sgn(cos(1/2*d*x + 1/2*c))*sin(3/2*d*x + 3/2*c) + 3*(2*A*sgn(cos(1/2*d*x + 1/2*c)) + B*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`**3.77.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)`output `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)`

3.78 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$

3.78.1 Optimal result	845
3.78.2 Mathematica [A] (verified)	845
3.78.3 Rubi [A] (verified)	846
3.78.4 Maple [B] (verified)	847
3.78.5 Fricas [B] (verification not implemented)	848
3.78.6 Sympy [F]	848
3.78.7 Maxima [A] (verification not implemented)	849
3.78.8 Giac [A] (verification not implemented)	849
3.78.9 Mupad [F(-1)]	850

3.78.1 Optimal result

Integrand size = 31, antiderivative size = 66

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2\sqrt{a}A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{d} + \frac{2aB \sin(c + dx)}{d\sqrt{a + a \cos(c + dx)}}$$

output `2*A*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d+2*a*B*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)`

3.78.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(\sqrt{2}A \operatorname{arctanh}\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2B \sin\left(\frac{1}{2}(c + dx)\right)\right)}{d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(Sqrt[a*(1 + Cos[c + d*x]])*Sec[(c + d*x)/2]*(Sqrt[2]*A*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]] + 2*B*Sin[(c + d*x)/2]))/d`

3.78. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$

3.78.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3042, 3460, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c+dx) \sqrt{a \cos(c+dx) + a} (A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a \sin(c+dx + \frac{\pi}{2}) + a} (A + B \sin(c+dx + \frac{\pi}{2}))}{\sin(c+dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3460} \\
 & A \int \sqrt{\cos(c+dx)a + a} \sec(c+dx) dx + \frac{2aB \sin(c+dx)}{d \sqrt{a \cos(c+dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & A \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2}) a + a}}{\sin(c+dx + \frac{\pi}{2})} dx + \frac{2aB \sin(c+dx)}{d \sqrt{a \cos(c+dx) + a}} \\
 & \quad \downarrow \text{3252} \\
 & \frac{2aB \sin(c+dx)}{d \sqrt{a \cos(c+dx) + a}} - \frac{2aA \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{2\sqrt{a}A \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx) + a}} \right)}{d} + \frac{2aB \sin(c+dx)}{d \sqrt{a \cos(c+dx) + a}}
 \end{aligned}$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(2*Sqrt[a]*A*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d + (2*a*B*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])`

3.78.3.1 Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3252 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (
f_.)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2),
x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.78.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 218 vs. 2(58) = 116.

Time = 3.71 (sec) , antiderivative size = 219, normalized size of antiderivative = 3.32

method	result
default	$\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(A\sqrt{2} \ln\left(\frac{2\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+4a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}}\right) + A\sqrt{2} \ln\left(-\frac{2\left(\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \right. \\ \left. + \frac{2\sqrt{a} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{d} \right)$
parts	$A\sqrt{a} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\ln\left(\frac{4\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+8a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}}\right) + \ln\left(-\frac{4\left(\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \right. \\ \left. + \frac{\sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{d} \right)$

3.78. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$

input `int((a+cos(d*x+c))*a^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `1/2/a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(A*2^(1/2)*ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+A*2^(1/2)*ln(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+4*B*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/sin(1/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.78.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs. $2(58) = 116$.

Time = 0.32 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.92

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{(A \cos(dx + c) + A)\sqrt{a} \log\left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a \cos(dx+c) + a}\sqrt{a}(\cos(dx+c) - 2) \sin(dx+c) + 8a}{\cos(dx+c)^3 + \cos(dx+c)^2}\right) + 4\sqrt{a \cos(dx+c)}}{2(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")`

output `1/2*((A*cos(d*x + c) + A)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*sqrt(a*cos(d*x + c) + a)*B*sin(d*x + c))/(d*cos(d*x + c) + d)`

3.78.6 Sympy [F]

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int \sqrt{a (\cos(c + dx) + 1)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))*sec(c + d*x), x)`

3.78.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.32

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx = \frac{2\sqrt{2}B\sqrt{a} \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{d}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `2*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c)/d`

3.78.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.35

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx = \frac{\sqrt{2} \left(\sqrt{2}A \log \left(\frac{|-2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|} \right) \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) - 4B \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{1}{2} dx + \frac{1}{2} c) \right)}{2d}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `-1/2*sqrt(2)*(sqrt(2)*A*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c)))*sgn(cos(1/2*d*x + 1/2*c)) - 4*B*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

3.78.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x),x)`output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x), x)`

3.79 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.79.1	Optimal result	851
3.79.2	Mathematica [A] (verified)	851
3.79.3	Rubi [A] (verified)	852
3.79.4	Maple [B] (verified)	854
3.79.5	Fricas [B] (verification not implemented)	855
3.79.6	Sympy [F]	855
3.79.7	Maxima [B] (verification not implemented)	856
3.79.8	Giac [B] (verification not implemented)	856
3.79.9	Mupad [F(-1)]	857

3.79.1 Optimal result

Integrand size = 33, antiderivative size = 68

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{\sqrt{a}(A + 2B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{d} + \frac{aA \tan(c + dx)}{d\sqrt{a + a \cos(c + dx)}}$$

```
output (A+2*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d+a*A*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
```

3.79.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.25

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec(c + dx) (\sqrt{2}(A + 2B)\operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right))) \cos(c + dx)}{2d}$$

```
input Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]
```

output $(\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])] * \text{Sec}[(c + d*x)/2] * \text{Sec}[c + d*x] * (\text{Sqrt}[2] * (A + 2*B) * \text{ArcTanh}[\text{Sqrt}[2] * \text{Sin}[(c + d*x)/2]] * \text{Cos}[c + d*x] + 2*A * \text{Sin}[(c + d*x)/2])) / (2*d)$

3.79.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {3042, 3459, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3459} \\
 & \frac{1}{2} (A + 2B) \int \sqrt{\cos(c + dx) a + a} \sec(c + dx) dx + \frac{aA \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} (A + 2B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{aA \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3252} \\
 & \frac{aA \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} - \frac{a(A + 2B) \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx) a + a}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right)}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{\sqrt{a} (A + 2B) \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} + \frac{aA \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}}
 \end{aligned}$$

input $\text{Int}[\text{Sqrt}[a + a*\text{Cos}[c + d*x]]*(A + B*\text{Cos}[c + d*x])* \text{Sec}[c + d*x]^2, x]$

3.79. $\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^2(c + dx) dx$

output $(\sqrt{a}*(A + 2*B)*\text{ArcTanh}[(\sqrt{a}*\sin[c + d*x])/(\sqrt{a + a*\cos[c + d*x]})]/d + (a*A*\tan[c + d*x])/(d*\sqrt{a + a*\cos[c + d*x]})$

3.79.3.1 Defintions of rubi rules used

rule 219 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3252 $\text{Int}[\sqrt{(a_ + (b_)*\sin[(e_.) + (f_)*(x_)])/(c_.) + (d_)*\sin[(e_.) + (f_)*(x_)]}], x_Symbol] \rightarrow \text{Simp}[-2*(b/f) \ \text{Subst}[\text{Int}[1/(b*c + a*d - d*x^2), x], x, b*(\cos[e + f*x]/\sqrt{a + b*\sin[e + f*x]})], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0]$

rule 3459 $\text{Int}[\sqrt{(a_ + (b_)*\sin[(e_.) + (f_)*(x_)])*(A_.) + (B_)*\sin[(e_.) + (f_)*(x_)])*(c_.) + (d_)*\sin[(e_.) + (f_)*(x_)]^n}, x_Symbol] \rightarrow \text{Simp}[(-b^2)*(B*c - A*d)*\cos[e + f*x]*(c + d*\sin[e + f*x])^{n+1}/(d*f*(n+1)*(b*c + a*d)*\sqrt{a + b*\sin[e + f*x]}), x] + \text{Simp}[(A*b*d*(2*n+3) - B*(b*c - 2*a*d*(n+1)))/(2*d*(n+1)*(b*c + a*d)) \ \text{Int}[\sqrt{a + b*\sin[e + f*x]}*(c + d*\sin[e + f*x])^{n+1}, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, A, B\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{LtQ}[n, -1]$

3.79.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 566 vs. 2(60) = 120.

Time = 4.71 (sec) , antiderivative size = 567, normalized size of antiderivative = 8.34

method	result
parts	$A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-2a \left(\ln \left(\frac{4\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+8a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}} \right) + \ln \left(-\frac{4 \left(\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)} \right) \right)$
default	$\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-2a \left(A \ln \left(-\frac{4 \left(\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right) \sqrt{a-2a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}} \right) + A \ln \left(\frac{4\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)} \right) \right)$

```
input int((a*cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNV
ERBOSE)
```

```
output A*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*a*(ln(4/(2*cos(1/2
*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+
1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2
)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a
))*sin(1/2*d*x+1/2*c)^2+ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos
(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+ln(
-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a
*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+2*2^(1/2)*(a*sin(1/2*d*x+1/2*
c)^2)^(1/2)*a^(1/2))/a^(1/2)/(2*cos(1/2*d*x+1/2*c)+2^(1/2))/(2*cos(1/2*d*x
+1/2*c)-2^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+B*a^(
1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(ln(2/(2*cos(1/2*d*
x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2
*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a
*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a)))/
sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.79. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.79.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(60) = 120.

Time = 0.32 (sec) , antiderivative size = 153, normalized size of antiderivative = 2.25

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{((A + 2B) \cos(dx + c))^2 + (A + 2B) \cos(dx + c)) \sqrt{a} \log\left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a} \cos(dx+c) + a\sqrt{a}(\cos(dx+c)^3 + \cos(dx+c)^2)}{\cos(dx+c)^3 + \cos(dx+c)^2}\right)}{4(d \cos(dx + c))^2 + d \cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/4*(((A + 2*B)*cos(d*x + c)^2 + (A + 2*B)*cos(d*x + c))*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*sqrt(a*cos(d*x + c) + a)*A*sin(d*x + c))/(d*cos(d*x + c)^2 + d*cos(d*x + c))`

3.79.6 Sympy [F]

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int \sqrt{a(\cos(c + dx) + 1)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))*sec(c + d*x)**2, x)`

3.79.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 710 vs. $2(60) = 120$.

Time = 0.42 (sec) , antiderivative size = 710, normalized size of antiderivative = 10.44

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `-1/4*(4*sqrt(2)*cos(5/2*d*x + 5/2*c)*sin(2*d*x + 2*c) + 4*sqrt(2)*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) - 4*sqrt(2)*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) - (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*log(2*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sqrt(2)*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2*sqrt(2)*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2) + (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*log(2*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sqrt(2)*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c))) - 2*sqrt(2)*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2) - (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*log(2*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 - 2*sqrt(2)*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2*sqrt(2)*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2) + (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*log(2*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 - 2*sqrt(2)*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c))) - 2*sqrt(2)*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2) - 4*(sqrt(2)*cos(2*d*x + 2*c) + sqrt(2))*sin(5/2*d*x + 5/2*c) + 4*(sqrt(2)*cos(2*d*x + 2*c)^2 + sqrt(2)*sin(2*d*x + 2*c)^2 + 2*sqrt(2)*cos(2*d*x + 2*c) + s...`

3.79.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(60) = 120$.

Time = 0.34 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.78

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$\frac{\sqrt{2} \left(\sqrt{2} \left(\operatorname{Asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) + 2 B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \right) \log \left(\frac{-2\sqrt{2} + 4 \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{\left| 2\sqrt{2} + 4 \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right|} \right) + \frac{4 \operatorname{Asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right)}{2 \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)} \right)}{4d}$$

3.79. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm
m="giac")`

output `-1/4*sqrt(2)*(sqrt(2)*(A*sgn(cos(1/2*d*x + 1/2*c)) + 2*B*sgn(cos(1/2*d*x +
1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*s
in(1/2*d*x + 1/2*c))) + 4*A*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)
/(2*sin(1/2*d*x + 1/2*c)^2 - 1))*sqrt(a)/d`

3.79.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\cos(c + dx)^2} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^2,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^2, x)`

3.80 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.80.1 Optimal result	858
3.80.2 Mathematica [A] (verified)	858
3.80.3 Rubi [A] (verified)	859
3.80.4 Maple [B] (verified)	861
3.80.5 Fricas [A] (verification not implemented)	862
3.80.6 Sympy [F]	863
3.80.7 Maxima [B] (verification not implemented)	863
3.80.8 Giac [A] (verification not implemented)	864
3.80.9 Mupad [F(-1)]	865

3.80.1 Optimal result

Integrand size = 33, antiderivative size = 117

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{\sqrt{a}(3A + 4B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{4d} + \frac{a(3A + 4B) \tan(c + dx)}{4d \sqrt{a + a \cos(c + dx)}} + \frac{aA \sec(c + dx) \tan(c + dx)}{2d \sqrt{a + a \cos(c + dx)}}$$

```
output 1/4*(3*A+4*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d
+1/4*a*(3*A+4*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/2*a*A*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
```

3.80.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.86

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))}(3\sqrt{2}(3A + 4B) \operatorname{arctanh}(\sqrt{2} \sin(\frac{1}{2}(c + dx)))) \sec(\frac{1}{2}(c + dx)) + 6(2A + (3A + 4B) \cos(\frac{1}{2}(c + dx))) \sqrt{a} \tan(\frac{1}{2}(c + dx))}{24d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(Sqrt[a*(1 + Cos[c + d*x])]*(3*Sqrt[2]*(3*A + 4*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Sec[(c + d*x)/2] + 6*(2*A + (3*A + 4*B)*Cos[c + d*x])*Sec[c + d*x]^2*Tan[(c + d*x)/2]))/(24*d)`

3.80.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.93, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3042, 3459, 3042, 3251, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3459} \\
 & \frac{1}{4} (3A + 4B) \int \sqrt{\cos(c + dx) a + a} \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} (3A + 4B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{aA \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3251} \\
 & \frac{1}{4} (3A + 4B) \left(\frac{1}{2} \int \sqrt{\cos(c + dx) a + a} \sec(c + dx) dx + \frac{a \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) + \\
 & \quad \frac{aA \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{1}{4}(3A + 4B) \left(\frac{1}{2} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}}$$

↓ 3252

$$\frac{1}{4}(3A + 4B) \left(\frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{a \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}}$$

↓ 219

$$\frac{1}{4}(3A + 4B) \left(\frac{\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} + \frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}}$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*Sec[c + d*x]*Tan[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + ((3*A + 4*B)*((Sqrt[a]*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4`

3.80.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3251 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e
+ f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Sim
p[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e
+ f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3252 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (
f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2),
x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.80.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 935 vs. $2(101) = 202$.

Time = 5.24 (sec) , antiderivative size = 936, normalized size of antiderivative = 8.00

method	result	size
parts	Expression too large to display	936
default	Expression too large to display	1003

```
input int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/2*A*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*a*(ln(4/(2*cos
(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*
d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^
(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-
2*a))*sin(1/2*d*x+1/2*c)^4+(-12*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^
(1/2)-12*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)
+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a-12*ln(-4/(2*cos(1/
2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x
+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^2+10*2^(1/2)*(a*sin(1
/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+3*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1
/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*
a))*a+3*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)
-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)/a^(1/2)/(2*cos(1/
2*d*x+1/2*c)-2^(1/2))^2/(2*cos(1/2*d*x+1/2*c)+2^(1/2))^2/sin(1/2*d*x+1/2*c
)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+B*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2
*c)^2)^(1/2)*(-2*a*(ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2
*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-2/(2*
cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1
/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*sin(1/2*d*x+1/2*c)^2+2*2^(1/2)*(a*si
n(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+ln(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*...
```

3.80.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.52

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{((3A + 4B) \cos(dx + c)^3 + (3A + 4B) \cos(dx + c)^2) \sqrt{a} \log \left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a \cos(dx+c)} + a\sqrt{a}(\cos(dx+c)^3 + \cos(dx+c)^2)}{\cos(dx+c)^3 + \cos(dx+c)^2} \right)}{16 (d \cos(dx + c))^3 + d \cos(dx + c)}$$

```
input integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorith
m="fricas")
```

```
output 1/16*(((3*A + 4*B)*cos(d*x + c)^3 + (3*A + 4*B)*cos(d*x + c)^2)*sqrt(a)*lo
g((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt
(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^
2)) + 4*((3*A + 4*B)*cos(d*x + c) + 2*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x
+ c))/(d*cos(d*x + c)^3 + d*cos(d*x + c)^2)
```

3.80. $\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.80.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= \int \sqrt{a(\cos(c + dx) + 1)}(A + B \cos(c + dx)) \sec^3(c + dx) dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))*sec(c + d*x)**3, x)`

3.80.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3352 vs. 2(101) = 202.

Time = 3.24 (sec) , antiderivative size = 3352, normalized size of antiderivative = 28.65

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

```
output 1/16*((3*(log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt
(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - log(2*cos
(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1
/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + log(2*cos(1/2*d*x + 1/2*c)^2
+ 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*s
in(1/2*d*x + 1/2*c) + 2) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x +
1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c)
+ 2))*cos(4*d*x + 4*c)^2 + 12*(log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d
*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1
/2*c) + 2) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*s
qrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + log(2*
cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x
+ 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - log(2*cos(1/2*d*x + 1/2*c
)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2
)*sin(1/2*d*x + 1/2*c) + 2))*cos(2*d*x + 2*c)^2 + 3*(log(2*cos(1/2*d*x + 1
/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sq
rt(2)*sin(1/2*d*x + 1/2*c) + 2) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2
*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x +
1/2*c) + 2) + log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2
*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - 1...
```

3.80.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.67

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx =$$

$$\sqrt{2} \left(\sqrt{2} \left(3 \operatorname{Asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) + 4 B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \right) \log \left(\frac{-2\sqrt{2} + 4 \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{2\sqrt{2} + 4 \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)} \right) \right) + \frac{4 \left(6 \operatorname{Asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \right)}{\dots}$$

```
input integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorith
m="giac")
```

```
output -1/16*sqrt(2)*(sqrt(2)*(3*A*sgn(cos(1/2*d*x + 1/2*c)) + 4*B*sgn(cos(1/2*d*
x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) +
4*sin(1/2*d*x + 1/2*c))) + 4*(6*A*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x +
1/2*c)^3 + 8*B*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 5*A*sgn(
cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) - 4*B*sgn(cos(1/2*d*x + 1/2*c))
*sin(1/2*d*x + 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^2)*sqrt(a)/d
```

3.80. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.80.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\cos(c + dx)^3} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^3,x)`output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^3, x)`

3.81 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$

3.81.1	Optimal result	866
3.81.2	Mathematica [A] (verified)	867
3.81.3	Rubi [A] (verified)	867
3.81.4	Maple [B] (verified)	870
3.81.5	Fricas [A] (verification not implemented)	871
3.81.6	Sympy [F(-1)]	872
3.81.7	Maxima [B] (verification not implemented)	872
3.81.8	Giac [A] (verification not implemented)	873
3.81.9	Mupad [F(-1)]	874

3.81.1 Optimal result

Integrand size = 33, antiderivative size = 160

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{\sqrt{a}(5A + 6B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{8d} + \frac{a(5A + 6B) \tan(c + dx)}{8d \sqrt{a + a \cos(c + dx)}} + \frac{a(5A + 6B) \sec(c + dx) \tan(c + dx)}{12d \sqrt{a + a \cos(c + dx)}} + \frac{aA \sec^2(c + dx) \tan(c + dx)}{3d \sqrt{a + a \cos(c + dx)}}$$

```
output 1/8*(5*A+6*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d
+1/8*a*(5*A+6*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/12*a*(5*A+6*B)*sec(
d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/3*a*A*sec(d*x+c)^2*tan(d*x+c)
/d/(a+a*cos(d*x+c))^(1/2)
```

3.81.2 Mathematica [A] (verified)

Time = 1.19 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.81

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} \sec^3(c + dx) (3\sqrt{2}(5A + 6B) \operatorname{arctanh}(\sqrt{2} \sin(\frac{1}{2}(c + dx))) \cos^3(c + dx) \sec(\frac{1}{2}(c + dx))}{48d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(Sqrt[a*(1 + Cos[c + d*x]])*Sec[c + d*x]^3*(3*Sqrt[2]*(5*A + 6*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^3*Sec[(c + d*x)/2] + (31*A + 18*B + 4*(5*A + 6*B)*Cos[c + d*x] + 3*(5*A + 6*B)*Cos[2*(c + d*x)])*Tan[(c + d*x)/2])/(48*d)`

3.81.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.94, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 3459, 3042, 3251, 3042, 3251, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx$$

$$\downarrow \text{3459}$$

$$\frac{1}{6}(5A + 6B) \int \sqrt{\cos(c + dx)a + a} \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d \sqrt{a \cos(c + dx) + a}}$$

$$\downarrow \text{3042}$$

$$\frac{1}{6}(5A + 6B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d \sqrt{a \cos(c + dx) + a}}$$

$$\begin{aligned}
& \downarrow \text{3251} \\
& \frac{1}{6}(5A + 6B) \left(\frac{3}{4} \int \sqrt{\cos(c + dx)a + a} \sec^2(c + dx) dx + \frac{a \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3042} \\
& \frac{1}{6}(5A + 6B) \left(\frac{3}{4} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{a \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3251} \\
& \frac{1}{6}(5A + \\
6B) & \left(\frac{3}{4} \left(\frac{1}{2} \int \sqrt{\cos(c + dx)a + a} \sec(c + dx) dx + \frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3042} \\
& \frac{1}{6}(5A + \\
6B) & \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3252} \\
& \frac{1}{6}(5A + \\
6B) & \left(\frac{3}{4} \left(\frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{a \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}} \right)}{d} \right) + \frac{a \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{219}
\end{aligned}$$

$$6B) \left(\frac{3}{4} \left(\frac{\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a \tan(c+dx)}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^2(c+dx)}{3d \sqrt{a \cos(c+dx)+a}} + \frac{1}{6}(5A +$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*Sec[c + d*x]^2*Tan[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + ((5*A + 6*B)*((a*Sec[c + d*x]*Tan[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*(Sqrt[a]*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (a*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)/6`

3.81.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3251 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3252 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.81.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1281 vs. $2(140) = 280$.

Time = 5.81 (sec) , antiderivative size = 1282, normalized size of antiderivative = 8.01

method	result	size
parts	Expression too large to display	1282
default	Expression too large to display	1327

```
input int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNV
ERBOSE)
```

output `1/6*A*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-120*a*(ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*sin(1/2*d*x+1/2*c)^6+60*(2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+3*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+3*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^4+(-90*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a-90*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a-160*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))*sin(1/2*d*x+1/2*c)^2+15*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+15*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+66*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/a^(1/2)/(2*cos(1/2*d*x+1/2*c)+2^(1/2))^3/(2*cos(1/2*d*x+1/2*c)-2^(1/2))^3/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+1/4*B*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*2^(1/2)*a*(ln(2/(2*...`

3.81.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.23

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3 \left((5A + 6B) \cos(dx + c)^4 + (5A + 6B) \cos(dx + c)^3 \right) \sqrt{a} \log \left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4 \sqrt{a \cos(dx+c) + a} \sqrt{a} \cos(dx+c)}{\cos(dx+c)^3 + \cos(dx+c)^2} \right)}{96 (d \cos(c + dx))^3}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")`

output `1/96*(3*((5*A + 6*B)*cos(d*x + c)^4 + (5*A + 6*B)*cos(d*x + c)^3)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*(3*(5*A + 6*B)*cos(d*x + c)^2 + 2*(5*A + 6*B)*cos(d*x + c) + 8*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c)^4 + d*cos(d*x + c)^3)`

3.81. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$

3.81.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.81.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5021 vs. $2(140) = 280$.

Time = 3.23 (sec) , antiderivative size = 5021, normalized size of antiderivative = 31.38

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="maxima")`

output

```
-1/96*((120*(sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 3*sin(2*d*x + 2*c))*cos(13/2*d*x + 13/2*c) - 8*(15*sin(11/2*d*x + 11/2*c) + 50*sin(9/2*d*x + 9/2*c) + 42*sin(7/2*d*x + 7/2*c) + 3*sin(5/2*d*x + 5/2*c) - 5*sin(3/2*d*x + 3/2*c))*cos(6*d*x + 6*c) + 360*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*cos(11/2*d*x + 11/2*c) + 1200*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*cos(9/2*d*x + 9/2*c) - 24*(42*sin(7/2*d*x + 7/2*c) + 3*sin(5/2*d*x + 5/2*c) - 5*sin(3/2*d*x + 3/2*c))*cos(4*d*x + 4*c) - 15*(sqrt(2)*cos(6*d*x + 6*c)^2 + 9*sqrt(2)*cos(4*d*x + 4*c)^2 + 9*sqrt(2)*cos(2*d*x + 2*c)^2 + sqrt(2)*sin(6*d*x + 6*c)^2 + 9*sqrt(2)*sin(4*d*x + 4*c)^2 + 18*sqrt(2)*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sqrt(2)*sin(2*d*x + 2*c)^2 + 2*(3*sqrt(2)*cos(4*d*x + 4*c) + 3*sqrt(2)*cos(2*d*x + 2*c) + sqrt(2))*cos(6*d*x + 6*c) + 6*(3*sqrt(2)*cos(2*d*x + 2*c) + sqrt(2))*cos(4*d*x + 4*c) + 6*(sqrt(2)*sin(4*d*x + 4*c) + sqrt(2)*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 6*sqrt(2)*cos(2*d*x + 2*c) + sqrt(2))*log(2*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c)))^2 + 2*sqrt(2)*cos(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2*sqrt(2)*sin(1/2*arctan2(sin(d*x + c), cos(d*x + c))) + 2) + 15*(sqrt(2)*cos(6*d*x + 6*c)^2 + 9*sqrt(2)*cos(4*d*x + 4*c)^2 + 9*sqrt(2)*cos(2*d*x + 2*c)^2 + sqrt(2)*sin(6*d*x + 6*c)^2 + 9*sqrt(2)*sin(4*d*x + 4*c)^2 + 18*sqrt(2)*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sqrt(2)*sin(2*d*x + 2*c)^2 + 2*(3*sqrt(2)*cos(4*d*x + 4*c) + 3*sqrt(2)*co...
```

3.81.8 Giac [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.52

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^4(c + dx) dx =$$

$$\frac{\sqrt{2} \left(3\sqrt{2} \left(5 \operatorname{Asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) + 6 B \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \right) \log \left(\frac{|-2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|} \right) + \frac{4}{60} \operatorname{Asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \right)}{1}$$

input

```
integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")
```


output `-1/96*sqrt(2)*(3*sqrt(2)*(5*A*sgn(cos(1/2*d*x + 1/2*c)) + 6*B*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))) + 4*(60*A*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 + 72*B*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 - 80*A*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 96*B*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 33*A*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) + 30*B*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^3)*sqrt(a)/d`

3.81.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\cos(c + dx)^4} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^4,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^4, x)`

3.82 $\int \cos^3(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

3.82.1	Optimal result	875
3.82.2	Mathematica [A] (verified)	876
3.82.3	Rubi [A] (verified)	876
3.82.4	Maple [A] (verified)	880
3.82.5	Fricas [A] (verification not implemented)	881
3.82.6	Sympy [F(-1)]	881
3.82.7	Maxima [A] (verification not implemented)	882
3.82.8	Giac [A] (verification not implemented)	882
3.82.9	Mupad [F(-1)]	883

3.82.1 Optimal result

Integrand size = 33, antiderivative size = 234

$$\int \cos^3(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx = \frac{4a^2(187A+168B) \sin(c+dx)}{495d\sqrt{a+a \cos(c+dx)}} + \frac{2a^2(187A+168B) \cos^3(c+dx) \sin(c+dx)}{693d\sqrt{a+a \cos(c+dx)}} + \frac{2a^2(11A+12B) \cos^4(c+dx) \sin(c+dx)}{99d\sqrt{a+a \cos(c+dx)}} - \frac{8a(187A+168B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{3465d} + \frac{2aB \cos^4(c+dx)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{11d} + \frac{4(187A+168B)(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{1155d}$$

```
output 4/1155*(187*A+168*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+4/495*a^2*(187*A+
168*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/693*a^2*(187*A+168*B)*cos(d*x
+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/99*a^2*(11*A+12*B)*cos(d*x+c)^
4*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-8/3465*a*(187*A+168*B)*sin(d*x+c)*(a
+a*cos(d*x+c))^(1/2)/d+2/11*a*B*cos(d*x+c)^4*sin(d*x+c)*(a+a*cos(d*x+c))^(
1/2)/d
```

3.82.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.53

$$\int \cos^3(c+dx)(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \frac{a\sqrt{a(1+\cos(c+dx))}(59158A+55482B+(35156A+34734B)\cos(c+dx)+8(1507A+1743B)\cos(2(c+dx))+3740A\cos(3(c+dx))+4935B\cos(3(c+dx))+770A\cos(4(c+dx))+1470B\cos(4(c+dx))+315B\cos(5(c+dx)))\tan((c+dx)/2)}{27720d}$$

input `Integrate[Cos[c + d*x]^3*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*sqrt[a*(1 + Cos[c + d*x])]*(59158*A + 55482*B + (35156*A + 34734*B)*Cos[c + d*x] + 8*(1507*A + 1743*B)*Cos[2*(c + d*x)] + 3740*A*Cos[3*(c + d*x)] + 4935*B*Cos[3*(c + d*x)] + 770*A*Cos[4*(c + d*x)] + 1470*B*Cos[4*(c + d*x)] + 315*B*Cos[5*(c + d*x)])*Tan[(c + d*x)/2])/(27720*d)`

3.82.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3238, 27, 3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^3(c+dx)(a\cos(c+dx)+a)^{3/2}(A+B\cos(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 \left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^{3/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3455} \\ & \frac{2}{11} \int \frac{1}{2} \cos^3(c+dx) \sqrt{\cos(c+dx)a+a(a(11A+8B)+a(11A+12B)\cos(c+dx))} dx + \\ & \quad \frac{2aB\sin(c+dx)\cos^4(c+dx)\sqrt{a\cos(c+dx)+a}}{11d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{11} \int \cos^3(c+dx) \sqrt{\cos(c+dx)a+a} (a(11A+8B) + a(11A+12B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^4(c+dx) \sqrt{a \cos(c+dx) + a}}{11d} \downarrow 3042$$

$$\frac{1}{11} \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} (a(11A+8B) + a(11A+12B) \sin\left(c+dx+\frac{\pi}{2}\right)) dx + \frac{2aB \sin(c+dx) \cos^4(c+dx) \sqrt{a \cos(c+dx) + a}}{11d} \downarrow 3460$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A+168B) \int \cos^3(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{2a^2(11A+12B) \sin(c+dx) \cos^4(c+dx)}{9d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aB \sin(c+dx) \cos^4(c+dx) \sqrt{a \cos(c+dx) + a}}{11d} \downarrow 3042$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A+168B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{2a^2(11A+12B) \sin(c+dx) \cos^4(c+dx)}{9d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aB \sin(c+dx) \cos^4(c+dx) \sqrt{a \cos(c+dx) + a}}{11d} \downarrow 3249$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A+168B) \left(\frac{6}{7} \int \cos^2(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^2(11A+12B) \sin(c+dx) \cos^4(c+dx)}{9d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aB \sin(c+dx) \cos^4(c+dx) \sqrt{a \cos(c+dx) + a}}{11d} \downarrow 3042$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A+168B) \left(\frac{6}{7} \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{2a \sin(c+dx) \cos^3(c+dx)}{7d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^2(11A+12B) \sin(c+dx) \cos^4(c+dx)}{9d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aB \sin(c+dx) \cos^4(c+dx) \sqrt{a \cos(c+dx) + a}}{11d} \downarrow 3238$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A + 168B) \left(\frac{6}{7} \left(\frac{2 \int \frac{1}{2} (3a - 2a \cos(c + dx)) \sqrt{\cos(c + dx)a + adx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^3}{5ad} \right) \right. \right. \\ \left. \left. \frac{2aB \sin(c + dx) \cos^4(c + dx) \sqrt{a \cos(c + dx) + a}}{11d} \right) \right. \\ \left. \downarrow 27 \right.$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A + 168B) \left(\frac{6}{7} \left(\frac{\int (3a - 2a \cos(c + dx)) \sqrt{\cos(c + dx)a + adx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \right. \\ \left. \left. \frac{2aB \sin(c + dx) \cos^4(c + dx) \sqrt{a \cos(c + dx) + a}}{11d} \right) \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A + 168B) \left(\frac{6}{7} \left(\frac{\int (3a - 2a \sin(c + dx + \frac{\pi}{2})) \sqrt{\sin(c + dx + \frac{\pi}{2})a + adx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \right. \\ \left. \left. \frac{2aB \sin(c + dx) \cos^4(c + dx) \sqrt{a \cos(c + dx) + a}}{11d} \right) \right. \\ \left. \downarrow 3230 \right.$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A + 168B) \left(\frac{6}{7} \left(\frac{\frac{7}{3} a \int \sqrt{\cos(c + dx)a + adx} - \frac{4a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \right. \\ \left. \left. \frac{2aB \sin(c + dx) \cos^4(c + dx) \sqrt{a \cos(c + dx) + a}}{11d} \right) \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{11} \left(\frac{1}{9} a(187A + 168B) \left(\frac{6}{7} \left(\frac{\frac{7}{3} a \int \sqrt{\sin(c + dx + \frac{\pi}{2})a + adx} - \frac{4a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \right. \\ \left. \left. \frac{2aB \sin(c + dx) \cos^4(c + dx) \sqrt{a \cos(c + dx) + a}}{11d} \right) \right. \\ \left. \downarrow 3125 \right.$$

$$\frac{1}{11} \left(\frac{2a^2(11A + 12B) \sin(c + dx) \cos^4(c + dx)}{9d \sqrt{a \cos(c + dx) + a}} + \frac{1}{9} a(187A + 168B) \left(\frac{6}{7} \left(\frac{\frac{14a^2 \sin(c + dx)}{3d \sqrt{a \cos(c + dx) + a}} - \frac{4a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}}{5a} \right) \right. \right. \\ \left. \left. \frac{2aB \sin(c + dx) \cos^4(c + dx) \sqrt{a \cos(c + dx) + a}}{11d} \right) \right.$$

input `Int[Cos[c + d*x]^3*(a + a*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x]),x]`

output `(2*a*B*cos[c + d*x]^4*Sqrt[a + a*cos[c + d*x]]*Sin[c + d*x])/(11*d) + ((2*a^2*(11*A + 12*B)*cos[c + d*x]^4*sin[c + d*x])/(9*d*Sqrt[a + a*cos[c + d*x]]) + (a*(187*A + 168*B)*((2*a*cos[c + d*x]^3*sin[c + d*x])/(7*d*Sqrt[a + a*cos[c + d*x]])) + (6*((2*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(5*a*d) + ((14*a^2*sin[c + d*x])/(3*d*Sqrt[a + a*cos[c + d*x]])) - (4*a*Sqrt[a + a*cos[c + d*x]]*sin[c + d*x])/(3*d))/(5*a))/7)/9)/11`

3.82.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3238 `Int[sin[(e_) + (f_)*(x_)]^2*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-Cos[e + f*x])*((a + b*sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*sin[e + f*x])^m*(b*(m + 1) - a*sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

```
rule 3249 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])
^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(
2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]
```

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.82.4 Maple [A] (verified)

Time = 2.55 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.61

method	result
default	$\frac{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(-5040B \left(\sin^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (3080A + 18480B) \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-9900A - 27720B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (3465A + 18480B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (114A + 1140B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 94A + 940B}{3465 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$
parts	$\frac{4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(280 \left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 220 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 114 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 47 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 94\right) \sqrt{2}}{315 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d + \frac{4B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(280 \left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 220 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 114 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 47 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 94\right) \sqrt{2}}{315 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$

```
input int(cos(d*x+c)^3*(a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.82. \int \cos^3(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

output $\frac{4/3465*\cos(1/2*d*x+1/2*c)*a^2*\sin(1/2*d*x+1/2*c)*(-5040*B*\sin(1/2*d*x+1/2*c)^{10}+(3080*A+18480*B)*\sin(1/2*d*x+1/2*c)^8+(-9900*A-27720*B)*\sin(1/2*d*x+1/2*c)^6+(12474*A+22176*B)*\sin(1/2*d*x+1/2*c)^4+(-8085*A-10395*B)*\sin(1/2*d*x+1/2*c)^2+3465*A+3465*B)*2^{(1/2)}}{(a*\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/d}$

3.82.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.53

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{2(315Ba \cos(dx + c)^5 + 35(11A + 21B)a \cos(dx + c)^4 + 5(187A + 168B)a \cos(dx + c)^3 + 6(187A + 168B)a \cos(dx + c)^2 + 8(187A + 168B)a \cos(dx + c) + 16(187A + 168B)a) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output $\frac{2/3465*(315*B*a*\cos(d*x + c)^5 + 35*(11*A + 21*B)*a*\cos(d*x + c)^4 + 5*(187*A + 168*B)*a*\cos(d*x + c)^3 + 6*(187*A + 168*B)*a*\cos(d*x + c)^2 + 8*(187*A + 168*B)*a*\cos(d*x + c) + 16*(187*A + 168*B)*a)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)}{(d*\cos(d*x + c) + d)}$

3.82.6 Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output Timed out

3.82.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.79

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{22(35\sqrt{2}a \sin(\frac{9}{2}dx + \frac{9}{2}c) + 135\sqrt{2}a \sin(\frac{7}{2}dx + \frac{7}{2}c) + 378\sqrt{2}a \sin(\frac{5}{2}dx + \frac{5}{2}c) + 1050\sqrt{2}a \sin(\frac{3}{2}dx + \frac{3}{2}c) + 3780\sqrt{2}a \sin(\frac{1}{2}dx + \frac{1}{2}c))A\sqrt{a} + 21(15\sqrt{2}(2)a \sin(\frac{11}{2}dx + \frac{11}{2}c) + 55\sqrt{2}(2)a \sin(\frac{9}{2}dx + \frac{9}{2}c) + 165\sqrt{2}(2)a \sin(\frac{7}{2}dx + \frac{7}{2}c) + 429\sqrt{2}(2)a \sin(\frac{5}{2}dx + \frac{5}{2}c) + 990\sqrt{2}(2)a \sin(\frac{3}{2}dx + \frac{3}{2}c) + 3630\sqrt{2}(2)a \sin(\frac{1}{2}dx + \frac{1}{2}c))B\sqrt{a}}{d}$$

```
input integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output 1/55440*(22*(35*sqrt(2)*a*sin(9/2*d*x + 9/2*c) + 135*sqrt(2)*a*sin(7/2*d*x + 7/2*c) + 378*sqrt(2)*a*sin(5/2*d*x + 5/2*c) + 1050*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 3780*sqrt(2)*a*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + 21*(15*sqrt(2)*a*sin(11/2*d*x + 11/2*c) + 55*sqrt(2)*a*sin(9/2*d*x + 9/2*c) + 165*sqrt(2)*a*sin(7/2*d*x + 7/2*c) + 429*sqrt(2)*a*sin(5/2*d*x + 5/2*c) + 990*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 3630*sqrt(2)*a*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d
```

3.82.8 Giac [A] (verification not implemented)

Time = 2.30 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.00

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(315 B a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{11}{2}dx + \frac{11}{2}c) + 385(2 A a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 3 B a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(9/2dx + 9/2c) + 495(6 A a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 7 B a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(7/2dx + 7/2c) + 693(12 A a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 13 B a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(5/2dx + 5/2c) + 2310(10 A a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 9 B a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(3/2dx + 3/2c) + 6930(12 A a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 11 B a \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(1/2dx + 1/2c)) \sqrt{a}}{d}$$

```
input integrate(cos(d*x+c)^3*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/55440*sqrt(2)*(315*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(11/2*d*x + 11/2*c) + 385*(2*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(9/2*d*x + 9/2*c) + 495*(6*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 7*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(7/2*d*x + 7/2*c) + 693*(12*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 13*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(5/2*d*x + 5/2*c) + 2310*(10*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 9*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 6930*(12*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 11*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d
```

3.82.9 Mupad [F(-1)]

Timed out.

$$\int \cos^3(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^3 (A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)`

3.83 $\int \cos^2(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

3.83.1	Optimal result	884
3.83.2	Mathematica [A] (verified)	885
3.83.3	Rubi [A] (verified)	885
3.83.4	Maple [A] (verified)	889
3.83.5	Fricas [A] (verification not implemented)	889
3.83.6	Sympy [F(-1)]	890
3.83.7	Maxima [A] (verification not implemented)	890
3.83.8	Giac [A] (verification not implemented)	891
3.83.9	Mupad [F(-1)]	891

3.83.1 Optimal result

Integrand size = 33, antiderivative size = 189

$$\int \cos^2(c+dx)(a + a \cos(c+dx))^{3/2}(A + B \cos(c+dx)) dx = \frac{2a^2(39A + 34B) \sin(c+dx)}{45d\sqrt{a + a \cos(c+dx)}} + \frac{2a^2(9A + 10B) \cos^3(c+dx) \sin(c+dx)}{63d\sqrt{a + a \cos(c+dx)}} - \frac{4a(39A + 34B) \sqrt{a + a \cos(c+dx)} \sin(c+dx)}{315d} + \frac{2aB \cos^3(c+dx) \sqrt{a + a \cos(c+dx)} \sin(c+dx)}{9d} + \frac{2(39A + 34B)(a + a \cos(c+dx))^{3/2} \sin(c+dx)}{105d}$$

```
output 2/105*(39*A+34*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/45*a^2*(39*A+34*B)
*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/63*a^2*(9*A+10*B)*cos(d*x+c)^3*sin(
d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-4/315*a*(39*A+34*B)*sin(d*x+c)*(a+a*cos(d*
x+c))^(1/2)/d+2/9*a*B*cos(d*x+c)^3*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.83.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.54

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{a\sqrt{a(1 + \cos(c + dx))}(2964A + 2689B + 2(759A + 799B) \cos(c + dx) + (468A + 548B) \cos^2(c + dx))}{1260d}$$

input `Integrate[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*(2964*A + 2689*B + 2*(759*A + 799*B)*Cos[c + d*x] + (468*A + 548*B)*Cos[2*(c + d*x)] + 90*A*Cos[3*(c + d*x)] + 170*B*Cos[3*(c + d*x)] + 35*B*Cos[4*(c + d*x)])*Tan[(c + d*x)/2])/(1260*d)`

3.83.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3455, 27, 3042, 3460, 3042, 3238, 27, 3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3455} \\ & \frac{2}{9} \int \frac{1}{2} \cos^2(c + dx) \sqrt{\cos(c + dx)a + a(3a(3A + 2B) + a(9A + 10B) \cos(c + dx))} dx + \\ & \quad \frac{2aB \sin(c + dx) \cos^3(c + dx) \sqrt{a \cos(c + dx) + a}}{9d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{9} \int \cos^2(c+dx) \sqrt{\cos(c+dx)a+a(3a(3A+2B)+a(9A+10B)\cos(c+dx))} dx + \frac{2aB \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{9d}$$

↓ 3042

$$\frac{1}{9} \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a(3a(3A+2B)+a(9A+10B)\sin\left(c+dx+\frac{\pi}{2}\right))} dx + \frac{2aB \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{9d}$$

↓ 3460

$$\frac{1}{9} \left(\frac{3}{7}a(39A+34B) \int \cos^2(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{2a^2(9A+10B) \sin(c+dx) \cos^3(c+dx)}{7d \sqrt{a \cos(c+dx)+a}} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{9d}$$

↓ 3042

$$\frac{1}{9} \left(\frac{3}{7}a(39A+34B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{2a^2(9A+10B) \sin(c+dx) \cos^3(c+dx)}{7d \sqrt{a \cos(c+dx)+a}} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{9d}$$

↓ 3238

$$\frac{1}{9} \left(\frac{3}{7}a(39A+34B) \left(\frac{2 \int \frac{1}{2}(3a-2a \cos(c+dx)) \sqrt{\cos(c+dx)a+adx}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{9d}$$

↓ 27

$$\frac{1}{9} \left(\frac{3}{7}a(39A+34B) \left(\frac{\int (3a-2a \cos(c+dx)) \sqrt{\cos(c+dx)a+adx}}{5a} + \frac{2 \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5ad} \right) \right) + \frac{2a^2}{9d} + \frac{2aB \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{9d}$$

↓ 3042

$$\frac{1}{9} \left(\frac{3}{7} a(39A + 34B) \left(\frac{\int (3a - 2a \sin(c + dx + \frac{\pi}{2})) \sqrt{\sin(c + dx + \frac{\pi}{2}) a + a dx}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)}{5ad} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) \sqrt{a \cos(c + dx) + a}}{9d} \right) \\ \downarrow \text{3230}$$

$$\frac{1}{9} \left(\frac{3}{7} a(39A + 34B) \left(\frac{\frac{7}{3} a \int \sqrt{\cos(c + dx) a + a dx} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) \sqrt{a \cos(c + dx) + a}}{9d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{9} \left(\frac{3}{7} a(39A + 34B) \left(\frac{\frac{7}{3} a \int \sqrt{\sin(c + dx + \frac{\pi}{2}) a + a dx} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) \sqrt{a \cos(c + dx) + a}}{9d} \right) \\ \downarrow \text{3125}$$

$$\frac{1}{9} \left(\frac{2a^2(9A + 10B) \sin(c + dx) \cos^3(c + dx)}{7d \sqrt{a \cos(c + dx) + a}} + \frac{3}{7} a(39A + 34B) \left(\frac{\frac{14a^2 \sin(c+dx)}{3d \sqrt{a \cos(c+dx)+a}} - \frac{4a \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{5a} + \frac{2 \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) \sqrt{a \cos(c + dx) + a}}{9d} \right)$$

input `Int[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(2*a*B*Cos[c + d*x]^3*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(9*d) + ((2*a^2*(9*A + 10*B)*Cos[c + d*x]^3*Sin[c + d*x])/(7*d*Sqrt[a + a*Cos[c + d*x]]) + (3*a*(39*A + 34*B)*((2*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*a*d) + ((14*a^2*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) - (4*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(5*a)))/7)/9`

3.83.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`
- rule 3238 `Int[sin[(e_) + (f_)*(x_)]^2*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-Cos[e + f*x])*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*(b*(m + 1) - a*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`
- rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.83.4 Maple [A] (verified)

Time = 2.59 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.65

method	result
default	$\frac{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(280B \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-180A - 900B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (504A + 1134B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-525A - 735B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 315A + 315B}{315 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$
parts	$\frac{4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(60 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 12 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 19 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 38\right) \sqrt{2}}{105 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d + \frac{4B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right)}{105 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$

```
input int(cos(d*x+c)^2*(a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 4/315*cos(1/2*d*x+1/2*c)*a^2*sin(1/2*d*x+1/2*c)*(280*B*sin(1/2*d*x+1/2*c)^
8+(-180*A-900*B)*sin(1/2*d*x+1/2*c)^6+(504*A+1134*B)*sin(1/2*d*x+1/2*c)^4+
(-525*A-735*B)*sin(1/2*d*x+1/2*c)^2+315*A+315*B)*2^(1/2)/(a*cos(1/2*d*x+1/
2*c)^2)^(1/2)/d
```

3.83.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.57

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{2(35Ba \cos(dx + c)^4 + 5(9A + 17B)a \cos(dx + c)^3 + 3(39A + 34B)a \cos(dx + c)^2 + 315A + 315B)}{315(d \cos(dx + c))^{3/2}}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorith
m="fracas")
```

3.83. $\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

output $2/315*(35*B*a*\cos(d*x + c)^4 + 5*(9*A + 17*B)*a*\cos(d*x + c)^3 + 3*(39*A + 34*B)*a*\cos(d*x + c)^2 + 4*(39*A + 34*B)*a*\cos(d*x + c) + 8*(39*A + 34*B)*a*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/(d*\cos(d*x + c) + d)$

3.83.6 Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output Timed out

3.83.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.81

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{6(15\sqrt{2}a \sin(\frac{7}{2}dx + \frac{7}{2}c) + 63\sqrt{2}a \sin(\frac{5}{2}dx + \frac{5}{2}c) + 175\sqrt{2}a \sin(\frac{3}{2}dx + \frac{3}{2}c) + 735\sqrt{2}a \sin(\frac{1}{2}dx + \frac{1}{2}c))A\sqrt{a} + (35\sqrt{2}a \sin(\frac{9}{2}dx + \frac{9}{2}c) + 135\sqrt{2}a \sin(\frac{7}{2}dx + \frac{7}{2}c) + 378\sqrt{2}a \sin(\frac{5}{2}dx + \frac{5}{2}c) + 1050\sqrt{2}a \sin(\frac{3}{2}dx + \frac{3}{2}c) + 3780\sqrt{2}a \sin(\frac{1}{2}dx + \frac{1}{2}c))B\sqrt{a}}{d}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output $1/2520*(6*(15*\sqrt{2}*a*\sin(7/2*d*x + 7/2*c) + 63*\sqrt{2}*a*\sin(5/2*d*x + 5/2*c) + 175*\sqrt{2}*a*\sin(3/2*d*x + 3/2*c) + 735*\sqrt{2}*a*\sin(1/2*d*x + 1/2*c))*A*\sqrt{a} + (35*\sqrt{2}*a*\sin(9/2*d*x + 9/2*c) + 135*\sqrt{2}*a*\sin(7/2*d*x + 7/2*c) + 378*\sqrt{2}*a*\sin(5/2*d*x + 5/2*c) + 1050*\sqrt{2}*a*\sin(3/2*d*x + 3/2*c) + 3780*\sqrt{2}*a*\sin(1/2*d*x + 1/2*c))*B*\sqrt{a})/d$

3.83.8 Giac [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.01

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(35 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{9}{2} dx + \frac{9}{2} c) + 45 (2 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 3 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(\frac{7}{2} dx + \frac{7}{2} c) + 378 (A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(\frac{5}{2} dx + \frac{5}{2} c) + 1050 (A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(\frac{3}{2} dx + \frac{3}{2} c) + 630 (7 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 6 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(\frac{1}{2} dx + \frac{1}{2} c)) \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))}{\sqrt{2}}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm
m="giac")
```

```
output 1/2520*sqrt(2)*(35*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(9/2*d*x + 9/2*c) + 45
*(2*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(7
/2*d*x + 7/2*c) + 378*(A*a*sgn(cos(1/2*d*x + 1/2*c)) + B*a*sgn(cos(1/2*d*x
+ 1/2*c)))*sin(5/2*d*x + 5/2*c) + 1050*(A*a*sgn(cos(1/2*d*x + 1/2*c)) + B
*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 630*(7*A*a*sgn(cos(1/
2*d*x + 1/2*c)) + 6*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*s
qrt(a)/d
```

3.83.9 Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^2 (A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2} dx$$

```
input int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2),x)
```

```
output int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)
```

3.84 $\int \cos(c+dx)(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

3.84.1	Optimal result	892
3.84.2	Mathematica [A] (verified)	893
3.84.3	Rubi [A] (verified)	893
3.84.4	Maple [A] (verified)	896
3.84.5	Fricas [A] (verification not implemented)	897
3.84.6	Sympy [F(-1)]	897
3.84.7	Maxima [A] (verification not implemented)	897
3.84.8	Giac [A] (verification not implemented)	898
3.84.9	Mupad [F(-1)]	898

3.84.1 Optimal result

Integrand size = 31, antiderivative size = 138

$$\int \cos(c+dx)(a + a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx = \frac{8a^2(21A+19B) \sin(c+dx)}{105d \sqrt{a+a \cos(c+dx)}} + \frac{2a(21A+19B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{105d} + \frac{2(7A-2B)(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{35d} + \frac{2B(a+a \cos(c+dx))^{5/2} \sin(c+dx)}{7ad}$$

```
output 2/35*(7*A-2*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/7*B*(a+a*cos(d*x+c))^(5/2)*sin(d*x+c)/a/d+8/105*a^2*(21*A+19*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/105*a*(21*A+19*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.84.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.59

$$\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{a\sqrt{a(1 + \cos(c + dx))}(546A + 494B + (252A + 253B)\cos(c + dx) + 6(7A + 13B)\cos(2(c + dx)) + 15B\cos(3(c + dx)))\tan((c + dx)/2)}{210d}$$

input `Integrate[Cos[c + d*x]*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*(546*A + 494*B + (252*A + 253*B)*Cos[c + d*x] + 6*(7*A + 13*B)*Cos[2*(c + d*x)] + 15*B*Cos[3*(c + d*x)])*Tan[(c + d*x)/2])/(210*d)`

3.84.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3447} \\ & \int (a \cos(c + dx) + a)^{3/2} (A \cos(c + dx) + B \cos^2(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\ & \quad \downarrow \text{3502} \end{aligned}$$

$$\begin{aligned}
& \frac{2 \int \frac{1}{2} (\cos(c+dx)a+a)^{3/2} (5aB+a(7A-2B)\cos(c+dx)) dx}{7a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{27} \\
& \frac{\int (\cos(c+dx)a+a)^{3/2} (5aB+a(7A-2B)\cos(c+dx)) dx}{7a} + \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\int (\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (5aB+a(7A-2B)\sin(c+dx+\frac{\pi}{2})) dx}{7a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{3230} \\
& \frac{\frac{1}{5}a(21A+19B) \int (\cos(c+dx)a+a)^{3/2} dx + \frac{2a(7A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d}}{7a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{5}a(21A+19B) \int (\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} dx + \frac{2a(7A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d}}{7a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{3126} \\
& \frac{\frac{1}{5}a(21A+19B) \left(\frac{4}{3}a \int \sqrt{\cos(c+dx)a+adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a(7A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d}}{7a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{5}a(21A+19B) \left(\frac{4}{3}a \int \sqrt{\sin(c+dx+\frac{\pi}{2})a+adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a(7A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d}}{7a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7ad} \\
& \quad \downarrow \text{3125}
\end{aligned}$$

$$\frac{\frac{1}{5}a(21A + 19B) \left(\frac{8a^2 \sin(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a(7A-2B) \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d}}{\frac{7a}{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}} + 7ad}$$

input `Int[Cos[c + d*x]*(a + a*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x]),x]`

output `(2*B*(a + a*cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*a*d) + ((2*a*(7*A - 2*B)*(a + a*cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (a*(21*A + 19*B)*((8*a^2*Ssin[c + d*x])/(3*d*Sqrt[a + a*cos[c + d*x]]) + (2*a*Sqrt[a + a*cos[c + d*x]])*Sin[c + d*x])/(3*d)))/5)/(7*a)`

3.84.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Ssin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Ssin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Ssin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Ssin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Ssin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.84.4 Maple [A] (verified)

Time = 2.60 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.75

method	result
default	$\frac{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(-60B \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (42A + 168B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-105A - 175B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 105A + 105B}{105 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(2 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right) \sqrt{2}}{5 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{4B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(60 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120 \cos^4\left(\frac{dx}{2} + \frac{c}{2}\right) + 60 \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 105A + 105B\right)}{105 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 4/105*cos(1/2*d*x+1/2*c)*a^2*sin(1/2*d*x+1/2*c)*(-60*B*sin(1/2*d*x+1/2*c)^
6+(42*A+168*B)*sin(1/2*d*x+1/2*c)^4+(-105*A-175*B)*sin(1/2*d*x+1/2*c)^2+10
5*A+105*B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.84. $\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

3.84.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.64

$$\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{2(15Ba \cos(dx + c)^3 + 3(7A + 13B)a \cos(dx + c)^2 + (63A + 52B)a \cos(dx + c) + 2(63A + 52B)a) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{105(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `2/105*(15*B*a*cos(d*x + c)^3 + 3*(7*A + 13*B)*a*cos(d*x + c)^2 + (63*A + 52*B)*a*cos(d*x + c) + 2*(63*A + 52*B)*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

3.84.6 Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.84.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.89

$$\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{42(\sqrt{2}a \sin(\frac{5}{2}dx + \frac{5}{2}c) + 5\sqrt{2}a \sin(\frac{3}{2}dx + \frac{3}{2}c) + 20\sqrt{2}a \sin(\frac{1}{2}dx + \frac{1}{2}c))A\sqrt{a} + (15Ba \cos(dx + c)^3 + 3(7A + 13B)a \cos(dx + c)^2 + (63A + 52B)a \cos(dx + c) + 2(63A + 52B)a) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{105(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/420*(42*(sqrt(2)*a*sin(5/2*d*x + 5/2*c) + 5*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 20*sqrt(2)*a*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + (15*sqrt(2)*a*sin(7/2*d*x + 7/2*c) + 63*sqrt(2)*a*sin(5/2*d*x + 5/2*c) + 175*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 735*sqrt(2)*a*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d`

3.84.8 Giac [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.12

$$\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(15 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{7}{2} dx + \frac{7}{2} c) + 21(2 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 3 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(5/2 dx + 5/2 c) + 35(6 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 5 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(3/2 dx + 3/2 c) + 105(8 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 7 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(1/2 dx + 1/2 c)) \sqrt{a}}{d}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/420*sqrt(2)*(15*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(7/2*d*x + 7/2*c) + 21*(2*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(5/2*d*x + 5/2*c) + 35*(6*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 5*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 105*(8*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 7*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

3.84.9 Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx) (A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)`

3.84. $\int \cos(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

3.85 $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx$

3.85.1	Optimal result	899
3.85.2	Mathematica [A] (verified)	899
3.85.3	Rubi [A] (verified)	900
3.85.4	Maple [A] (verified)	901
3.85.5	Fricas [A] (verification not implemented)	902
3.85.6	Sympy [F]	902
3.85.7	Maxima [A] (verification not implemented)	903
3.85.8	Giac [A] (verification not implemented)	903
3.85.9	Mupad [F(-1)]	904

3.85.1 Optimal result

Integrand size = 25, antiderivative size = 101

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{8a^2(5A + 3B) \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{2a(5A + 3B)\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{15d} + \frac{2B(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{5d}$$

```
output 2/5*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+8/15*a^2*(5*A+3*B)*sin(d*x+c)/d/
(a+a*cos(d*x+c))^(1/2)+2/15*a*(5*A+3*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/
d
```

3.85.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.64

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{a\sqrt{a(1 + \cos(c + dx))}(50A + 39B + 2(5A + 9B) \cos(c + dx) + 3B \cos(2(c + dx))) \tan((c + dx)/2)}{15d}$$

```
input Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]
```

```
output (a*Sqrt[a*(1 + Cos[c + d*x])]*(50*A + 39*B + 2*(5*A + 9*B)*Cos[c + d*x] +
3*B*Cos[2*(c + d*x)])*Tan[(c + d*x)/2])/(15*d)
```

3.85.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {3042, 3230, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx \\
 & \quad \downarrow \text{3230} \\
 & \frac{1}{5}(5A + 3B) \int (\cos(c + dx)a + a)^{3/2} dx + \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5}(5A + 3B) \int \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^{3/2} dx + \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3126} \\
 & \frac{1}{5}(5A + 3B) \left(\frac{4}{3} a \int \sqrt{\cos(c + dx)a + a} dx + \frac{2a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \\
 & \quad \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5}(5A + 3B) \left(\frac{4}{3} a \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a} dx + \frac{2a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \\
 & \quad \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3125} \\
 & \frac{1}{5}(5A + 3B) \left(\frac{8a^2 \sin(c + dx)}{3d \sqrt{a \cos(c + dx) + a}} + \frac{2a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \\
 & \quad \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}
 \end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

```
output (2*B*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(5*d) + ((5*A + 3*B)*((8*a^2
*Sin[c + d*x])/(3*d*Sqrt[a + a*cos[c + d*x])) + (2*a*Sqrt[a + a*cos[c + d*
x]])*Sin[c + d*x])/(3*d))/5
```

3.85.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3125 Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos
[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq
Q[a^2 - b^2, 0]
```

```
rule 3126 Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n)
Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[
a^2 - b^2, 0] && IGtQ[n - 1/2, 0]
```

```
rule 3230 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[
e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

3.85.4 Maple [A] (verified)

Time = 2.50 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.84

method	result
default	$\frac{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(6B \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-5A - 15B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 15A + 15B\right) \sqrt{2}}{15 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{4A a^2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right) \sqrt{2}}{3 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{4B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^2 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(2 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\right) \sqrt{2}}{5 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

3.85. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx$

input `int((a+cos(d*x+c))*a^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `4/15*cos(1/2*d*x+1/2*c)*a^2*sin(1/2*d*x+1/2*c)*(6*B*sin(1/2*d*x+1/2*c)^4+(-5*A-15*B)*sin(1/2*d*x+1/2*c)^2+15*A+15*B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.85.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.68

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{2(3Ba \cos(dx + c)^2 + (5A + 9B)a \cos(dx + c) + (25A + 18B)a) \sqrt{a \cos(dx + c) + a}}{15(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `2/15*(3*B*a*cos(d*x + c)^2 + (5*A + 9*B)*a*cos(d*x + c) + (25*A + 18*B)*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

3.85.6 Sympy [F]

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \int (a(\cos(c + dx) + 1))^{3/2} (A + B \cos(c + dx)) dx$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Integral((a*(cos(c + d*x) + 1))**(3/2)*(A + B*cos(c + d*x)), x)`

3.85.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.92

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{10 (\sqrt{2} a \sin(\frac{3}{2} dx + \frac{3}{2} c) + 9 \sqrt{2} a \sin(\frac{1}{2} dx + \frac{1}{2} c)) A \sqrt{a} + 3 (\sqrt{2} a \sin(\frac{5}{2} dx + \frac{5}{2} c) + 5 \sqrt{2} a \sin(\frac{3}{2} dx + \frac{3}{2} c)) B \sqrt{a}}{30 d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`output `1/30*(10*(sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 9*sqrt(2)*a*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + 3*(sqrt(2)*a*sin(5/2*d*x + 5/2*c) + 5*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 20*sqrt(2)*a*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d`**3.85.8 Giac [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.14

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{\sqrt{2} (3 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{5}{2} dx + \frac{5}{2} c) + 5 (2 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 3 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(\frac{3}{2} dx + \frac{3}{2} c) + 30 (3 A a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 2 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \sin(\frac{1}{2} dx + \frac{1}{2} c)) \sqrt{a}}{30 d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/30*sqrt(2)*(3*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(5/2*d*x + 5/2*c) + 5*(2*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 30*(3*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 2*B*a*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

3.85.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)`output `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)`

3.86 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec(c+dx) dx$

3.86.1	Optimal result	905
3.86.2	Mathematica [A] (verified)	905
3.86.3	Rubi [A] (verified)	906
3.86.4	Maple [B] (verified)	909
3.86.5	Fricas [A] (verification not implemented)	909
3.86.6	Sympy [F(-1)]	910
3.86.7	Maxima [A] (verification not implemented)	910
3.86.8	Giac [A] (verification not implemented)	911
3.86.9	Mupad [F(-1)]	911

3.86.1 Optimal result

Integrand size = 31, antiderivative size = 105

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{2a^{3/2} A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} + \frac{2a^2(3A + 4B) \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}} + \frac{2aB\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{3d}$$

output

```
2*a^(3/2)*A*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+2/3*a^2*(3*A+4*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/3*a*B*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.86.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.81

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) (3\sqrt{2}A \operatorname{arctanh}\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2(3A + 5B + B \cos(c + dx)))}{3d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(3*Sqrt[2]*A*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]] + 2*(3*A + 5*B + B*Cos[c + d*x])*Sin[(c + d*x)/2]))/(3*d)`

3.86.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 3455, 27, 3042, 3460, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3455} \\
 & \frac{2}{3} \int \frac{1}{2} \sqrt{\cos(c + dx)a + a(3aA + a(3A + 4B) \cos(c + dx))} \sec(c + dx) dx + \\
 & \quad \frac{2aB \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \int \sqrt{\cos(c + dx)a + a(3aA + a(3A + 4B) \cos(c + dx))} \sec(c + dx) dx + \\
 & \quad \frac{2aB \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a(3aA + a(3A + 4B) \sin(c + dx + \frac{\pi}{2}))}}{\sin(c + dx + \frac{\pi}{2})} dx + \\
 & \quad \frac{2aB \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\
 & \quad \downarrow \text{3460}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3} \left(3aA \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx + \frac{2a^2(3A+4B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(3aA \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2a^2(3A+4B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} \\
& \quad \downarrow \text{3252} \\
& \frac{1}{3} \left(\frac{2a^2(3A+4B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{6a^2A \int \frac{1}{a-\frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} \\
& \quad \downarrow \text{219} \\
& \frac{1}{3} \left(\frac{6a^{3/2} A \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}} \right)}{d} + \frac{2a^2(3A+4B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(2*a*B*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((6*a^(3/2)*A*ArcTan
h[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (2*a^2*(3*A + 4*B)
*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/3`

3.86.3.1 Defintions of rubi rules used

- rule 277 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.86.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. 2(91) = 182.

Time = 4.25 (sec) , antiderivative size = 269, normalized size of antiderivative = 2.56

method	result
parts	$\frac{A\sqrt{a} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(2\sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} + \ln\left(\frac{4\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} + 8a}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}}\right) a + \ln\left(\frac{\sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}{\sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}\right)}{\sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
default	$\frac{\sqrt{a} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-8B \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 3A\sqrt{2} \ln\left(-\frac{2\left(\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}}\right)}{6 \sin\left(\frac{dx}{2} + \frac{c}{2}\right)}$

input `int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `A*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+4/3*B*a^2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)*(cos(1/2*d*x+1/2*c)^2+2)^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.86.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.42

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{3(Aa \cos(dx + c) + Aa)\sqrt{a} \log\left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a \cos(dx+c)} + a\sqrt{a}(\cos(dx+c)-2) \sin(dx+c) + 8a}{\cos(dx+c)^3 + \cos(dx+c)^2}\right)}{6(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fracas")`

output $1/6*(3*(A*a*\cos(d*x + c) + A*a)*\sqrt{a}*\log((a*\cos(d*x + c))^3 - 7*a*\cos(d*x + c)^2 - 4*\sqrt{a*\cos(d*x + c) + a}*\sqrt{a}*(\cos(d*x + c) - 2)*\sin(d*x + c) + 8*a)/(\cos(d*x + c)^3 + \cos(d*x + c)^2)) + 4*(B*a*\cos(d*x + c) + (3*A + 5*B)*a)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c))/(d*\cos(d*x + c) + d)$

3.86.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `Timed out`

3.86.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.37

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{(\sqrt{2}a \sin(\frac{3}{2} dx + \frac{3}{2} c) + 9 \sqrt{2}a \sin(\frac{1}{2} dx + \frac{1}{2} c)) B \sqrt{a}}{3d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output $1/3*(\sqrt{2}*a*\sin(3/2*d*x + 3/2*c) + 9*\sqrt{2}*a*\sin(1/2*d*x + 1/2*c))*B*\sqrt{a}/d$

3.86.8 Giac [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.33

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx =$$

$$\sqrt{2} \left(8 B a \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{1}{2} dx + \frac{1}{2} c)^3 + 3 \sqrt{2} A a \log \left(\frac{|-2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|} \right) \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \right) \frac{1}{d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `-1/6*sqrt(2)*(8*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 3*sqrt(2)*A*a*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c)))*sgn(cos(1/2*d*x + 1/2*c)) - 12*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) - 24*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`

3.86.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x), x)`

3.87 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.87.1	Optimal result	912
3.87.2	Mathematica [A] (verified)	912
3.87.3	Rubi [A] (verified)	913
3.87.4	Maple [B] (verified)	916
3.87.5	Fricas [A] (verification not implemented)	917
3.87.6	Sympy [F(-1)]	917
3.87.7	Maxima [B] (verification not implemented)	917
3.87.8	Giac [A] (verification not implemented)	918
3.87.9	Mupad [F(-1)]	919

3.87.1 Optimal result

Integrand size = 33, antiderivative size = 103

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{a^{3/2}(3A + 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} - \frac{a^2(A - 2B) \sin(c + dx)}{d\sqrt{a + a \cos(c + dx)}} + \frac{aA\sqrt{a + a \cos(c + dx)} \tan(c + dx)}{d}$$

output

```
a^(3/2)*(3*A+2*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d-a^2*(A-2*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+a*A*(a+a*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.87.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.95

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec(c + dx) (\sqrt{2}(3A + 2B) \operatorname{arctanh}\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) \cos\left(\frac{1}{2}(c + dx)\right) + 2A \sin(c + dx))}{2d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x
]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]*(Sqrt[2]*(3*A
+ 2*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x] + 2*(A + 2*B*Cos[c +
d*x])*Sin[(c + d*x)/2]))/(2*d)`

3.87.3 Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3454, 27, 3042, 3460, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3454} \\
 & \int \frac{1}{2} \sqrt{\cos(c + dx)a + a}(a(3A + 2B) - a(A - 2B) \cos(c + dx)) \sec(c + dx) dx + \\
 & \quad \frac{aA \tan(c + dx) \sqrt{a \cos(c + dx) + a}}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \sqrt{\cos(c + dx)a + a}(a(3A + 2B) - a(A - 2B) \cos(c + dx)) \sec(c + dx) dx + \\
 & \quad \frac{aA \tan(c + dx) \sqrt{a \cos(c + dx) + a}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}(a(3A + 2B) - a(A - 2B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \\
 & \quad \frac{aA \tan(c + dx) \sqrt{a \cos(c + dx) + a}}{d} \\
 & \quad \downarrow \text{3460}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(a(3A + 2B) \int \frac{\sqrt{\cos(c + dx)a + a} \sec(c + dx) dx - \frac{2a^2(A - 2B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}}}{\frac{aA \tan(c + dx) \sqrt{a \cos(c + dx) + a}}{d}} \right) + \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{1}{2} \left(a(3A + 2B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{2a^2(A - 2B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \qquad \qquad \qquad \downarrow \text{3252} \\
& \frac{1}{2} \left(-\frac{2a^2(3A + 2B) \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} - \frac{2a^2(A - 2B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \qquad \qquad \qquad \downarrow \text{219} \\
& \frac{1}{2} \left(\frac{2a^{3/2}(3A + 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} - \frac{2a^2(A - 2B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \qquad \qquad \qquad \frac{aA \tan(c + dx) \sqrt{a \cos(c + dx) + a}}{d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `((2*a^(3/2)*(3*A + 2*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d - (2*a^2*(A - 2*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]))/2 + (a*A*Sqrt[a + a*Cos[c + d*x])*Tan[c + d*x])/d`

3.87.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.87.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 602 vs. 2(93) = 186.

Time = 4.81 (sec) , antiderivative size = 603, normalized size of antiderivative = 5.85

method	result
parts	$A\sqrt{a} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-6a \left(\ln \left(\frac{4\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+8a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}} \right) + \ln \left(-\frac{4\left(\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)} \right) \right)$
default	$\sqrt{a} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\left(-6A \ln \left(-\frac{4\left(\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right) \sqrt{a-2a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}} \right) a - 6A \ln \left(\frac{4\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+8a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}} \right) \right)$

input `int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNV
ERBOSE)`

output

```
A*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-6*a*(ln(4/(2
*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))
*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1
/2)-2*a)))*sin(1/2*d*x+1/2*c)^2+2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a
^(1/2)+3*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)
+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+3*ln(-4/(2*cos(1/2
*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+
1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)/(2*cos(1/2*d*x+1/2*c)+2^(1/2))/(2*cos(1/2
*d*x+1/2*c)-2^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+1
/2*B*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2^(1/2)*ln
(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+2^(1/2)*ln(2/(2*cos(1/2*d*x+
1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)
)^2)^(1/2)*a^(1/2)+2*a))*a+4*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2))/sin(1
/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.87.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.67

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{((3A + 2B)a \cos(dx + c))^2 + (3A + 2B)a \cos(dx + c) \sqrt{a} \log\left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a} \cos(dx+c)}{\cos(dx+c)^3 + a}\right)}{4(d \cos(dx + c))^2 + \dots}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="fricas")`

output `1/4*(((3*A + 2*B)*a*cos(d*x + c)^2 + (3*A + 2*B)*a*cos(d*x + c))*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*(2*B*a*cos(d*x + c) + A*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c)^2 + d*cos(d*x + c))`

3.87.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Timed out`

3.87.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1315 vs. 2(93) = 186.

Time = 0.37 (sec) , antiderivative size = 1315, normalized size of antiderivative = 12.77

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `-1/4*(2*sqrt(2)*a*cos(7/2*d*x + 7/2*c)*sin(2*d*x + 2*c) + 6*sqrt(2)*a*cos(5/2*d*x + 5/2*c)*sin(2*d*x + 2*c) + (2*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 6*sqrt(2)*a*sin(1/2*d*x + 1/2*c) - 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2))*cos(2*d*x + 2*c)^2 + (2*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 6*sqrt(2)*a*sin(1/2*d*x + 1/2*c) - 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 3*a*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2))*sin(2*d*x + 2*c)^2 - 4*sqrt(2)*a*sin(3/2*d*x + 3/2*c) + 4*sqrt(2)*a*sin(1/2*d*x + 1/2*c) - 2*(sqrt(2)*a*sin(3/2*d*x + 3/2*c) - 5*sqrt(2)*a*sin...`

3.87.8 Giac [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.45

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{\sqrt{2} \left(8 B \operatorname{asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right) - \frac{4 A \operatorname{asgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{2 \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 - 1} - \sqrt{2} (3 A \operatorname{asgn} (c + dx) \right)}{4 d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output $1/4*\sqrt{2}*(8*B*a*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c) - 4*A*a*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)/(2*\sin(1/2*d*x + 1/2*c)^2 - 1) - \sqrt{2}*(3*A*a*\text{sgn}(\cos(1/2*d*x + 1/2*c)) + 2*B*a*\text{sgn}(\cos(1/2*d*x + 1/2*c)))*\log(\text{abs}(-2*\sqrt{2} + 4*\sin(1/2*d*x + 1/2*c))/\text{abs}(2*\sqrt{2} + 4*\sin(1/2*d*x + 1/2*c))))*\sqrt{a}/d$

3.87.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)^2} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^2,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^2, x)`

3.88 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.88.1	Optimal result	920
3.88.2	Mathematica [A] (verified)	920
3.88.3	Rubi [A] (verified)	921
3.88.4	Maple [B] (verified)	924
3.88.5	Fricas [A] (verification not implemented)	925
3.88.6	Sympy [F(-1)]	925
3.88.7	Maxima [B] (verification not implemented)	925
3.88.8	Giac [A] (verification not implemented)	926
3.88.9	Mupad [F(-1)]	927

3.88.1 Optimal result

Integrand size = 33, antiderivative size = 119

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{a^{3/2}(7A + 12B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4d} + \frac{a^2(5A + 4B) \tan(c + dx)}{4d\sqrt{a + a \cos(c + dx)}} + \frac{aA\sqrt{a + a \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{2d}$$

```
output 1/4*a^(3/2)*(7*A+12*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+1/4*a^2*(5*A+4*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/2*a*A*sec(d*x+c)*(a+a*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.88.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.92

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^2(c + dx) (\sqrt{2}(7A + 12B)\operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)))}{8d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x
]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^2*(Sqrt[2]*(7*
A + 12*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^2 + 2*(2*A + (7*A
+ 4*B)*Cos[c + d*x])*Sin[(c + d*x)/2]))/(8*d)`

3.88.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3454} \\
 & \frac{1}{2} \int \frac{1}{2} \sqrt{\cos(c + dx)a + a(a(5A + 4B) + a(A + 4B) \cos(c + dx))} \sec^2(c + dx) dx + \\
 & \quad \frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4} \int \sqrt{\cos(c + dx)a + a(a(5A + 4B) + a(A + 4B) \cos(c + dx))} \sec^2(c + dx) dx + \\
 & \quad \frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a(a(5A + 4B) + a(A + 4B) \sin(c + dx + \frac{\pi}{2}))}}{\sin(c + dx + \frac{\pi}{2})^2} dx + \\
 & \quad \frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d} \\
 & \quad \downarrow \text{3459}
 \end{aligned}$$

3.88. $\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^3(c + dx) dx$

$$\begin{aligned}
& \frac{1}{4} \left(\frac{1}{2} a(7A + 12B) \int \frac{\sqrt{\cos(c + dx)a + a} \sec(c + dx) dx + \frac{a^2(5A + 4B) \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}}}{\frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d}} \right) + \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{1}{4} \left(\frac{1}{2} a(7A + 12B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a^2(5A + 4B) \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \qquad \qquad \qquad \frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d} \\
& \qquad \qquad \qquad \downarrow \text{3252} \\
& \frac{1}{4} \left(\frac{a^2(5A + 4B) \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{a^2(7A + 12B) \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} \right) + \\
& \qquad \qquad \qquad \frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d} \\
& \qquad \qquad \qquad \downarrow \text{219} \\
& \frac{1}{4} \left(\frac{a^{3/2}(7A + 12B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} + \frac{a^2(5A + 4B) \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \qquad \qquad \qquad \frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*sqrt[a + a*cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((a^(3/2)*(7*A + 12*B)*ArcTanh[(sqrt[a]*sin[c + d*x])/sqrt[a + a*cos[c + d*x]]])/d + (a^2*(5*A + 4*B)*Tan[c + d*x])/(d*sqrt[a + a*cos[c + d*x]])/4`

3.88.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.88.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 937 vs. $2(103) = 206$.

Time = 5.39 (sec) , antiderivative size = 938, normalized size of antiderivative = 7.88

method	result	size
parts	Expression too large to display	938
default	Expression too large to display	1003

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/2*A*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(28*a*(ln(
4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*
sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1
/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*
a^(1/2)-2*a)))*sin(1/2*d*x+1/2*c)^4+(-28*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2
)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^
(1/2)+2*a))*a-28*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d
*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a-28*2^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))*sin(1/2*d*x+1/2*c)^2+7*ln(4/(2*c
os(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/
2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+7*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2
)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^
(1/2)-2*a))*a+18*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/(2*cos(1/
2*d*x+1/2*c)+2^(1/2))^2/(2*cos(1/2*d*x+1/2*c)-2^(1/2))^2/sin(1/2*d*x+1/2*c
)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+B*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-6*a*(ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a
*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+l
n(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*
(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a)))*sin(1/2*d*x+1/2*c)^2+2*2^(1/
2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+3*ln(2/(2*cos(1/2*d*x+1/2*c)+...
```

3.88.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.53

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{((7A + 12B)a \cos(dx + c))^3 + (7A + 12B)a \cos(dx + c)^2 \sqrt{a} \log\left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a} \cos(dx+c) + a}{\cos(dx+c)}\right) + 4((7A + 4B)a \cos(dx + c) + 2Aa) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{16 (d \cos(dx + c))^3 + d \cos(dx + c)^2}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fricas")`

output `1/16*(((7*A + 12*B)*a*cos(d*x + c)^3 + (7*A + 12*B)*a*cos(d*x + c)^2)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*((7*A + 4*B)*a*cos(d*x + c) + 2*A*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(d*cos(d*x + c)^3 + d*cos(d*x + c)^2)`

3.88.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Timed out`

3.88.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3339 vs. 2(103) = 206.

Time = 0.50 (sec) , antiderivative size = 3339, normalized size of antiderivative = 28.06

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `-1/16*((12*a*cos(4*d*x + 4*c)^2*sin(3/2*d*x + 3/2*c) + 48*a*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 12*a*sin(4*d*x + 4*c)^2*sin(3/2*d*x + 3/2*c) + 48*a*sin(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 160*a*cos(7/2*d*x + 7/2*c)*sin(2*d*x + 2*c) + 168*a*cos(5/2*d*x + 5/2*c)*sin(2*d*x + 2*c) + 72*a*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) - 24*a*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) - 4*(a*sin(4*d*x + 4*c) + 2*a*sin(2*d*x + 2*c))*cos(13/2*d*x + 13/2*c) + 12*(a*sin(4*d*x + 4*c) + 2*a*sin(2*d*x + 2*c))*cos(11/2*d*x + 11/2*c) + 48*(a*sin(4*d*x + 4*c) + 2*a*sin(2*d*x + 2*c))*cos(9/2*d*x + 9/2*c) + 4*(12*a*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) - 20*a*sin(7/2*d*x + 7/2*c) - 21*a*sin(5/2*d*x + 5/2*c) - 3*a*sin(3/2*d*x + 3/2*c))*cos(4*d*x + 4*c) - 7*(sqrt(2)*a*cos(4*d*x + 4*c)^2 + 4*sqrt(2)*a*cos(2*d*x + 2*c)^2 + sqrt(2)*a*sin(4*d*x + 4*c)^2 + 4*sqrt(2)*a*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sqrt(2)*a*sin(2*d*x + 2*c)^2 + 4*sqrt(2)*a*cos(2*d*x + 2*c) + 2*(2*sqrt(2)*a*cos(2*d*x + 2*c) + sqrt(2)*a)*cos(4*d*x + 4*c) + sqrt(2)*a*log(2*cos(1/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c)))^2 + 2*sin(1/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c)))^2 + 2*sqrt(2)*cos(1/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c))) + 2*sqrt(2)*sin(1/3*arctan2(sin(3/2*d*x + 3/2*c), cos(3/2*d*x + 3/2*c))) + 2) + 7*(sqrt(2)*a*cos(4*d*x + 4*c)^2 + 4*sqrt(2)*a*cos(2*d*x + 2*c)^2 + sqrt(2)*a*sin(4*d*x + 4*c)^2 + 4*sqrt(2)*a*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sqrt(2)*a*sin(2...)`

3.88.8 Giac [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.69

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx =$$

$$\sqrt{2} \left(\sqrt{2} (7 A \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 12 B \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \log \left(\frac{-2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)}{2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)} \right) + \frac{4 (14 A \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 12 B \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)))}{\sqrt{2}} \right)$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output `-1/16*sqrt(2)*(sqrt(2)*(7*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 12*B*a*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c)))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))) + 4*(14*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 8*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 9*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) - 4*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^2)*sqrt(a)/d`

3.88.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)^3} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^3,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^3, x)`

3.89 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.89.1 Optimal result	928
3.89.2 Mathematica [A] (verified)	929
3.89.3 Rubi [A] (verified)	929
3.89.4 Maple [B] (verified)	933
3.89.5 Fricas [A] (verification not implemented)	934
3.89.6 Sympy [F(-1)]	934
3.89.7 Maxima [B] (verification not implemented)	934
3.89.8 Giac [A] (verification not implemented)	935
3.89.9 Mupad [F(-1)]	936

3.89.1 Optimal result

Integrand size = 33, antiderivative size = 164

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{a^{3/2}(11A + 14B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{8d} + \frac{a^2(11A + 14B) \tan(c + dx)}{8d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(7A + 6B) \sec(c + dx) \tan(c + dx)}{12d\sqrt{a + a \cos(c + dx)}} + \frac{aA\sqrt{a + a \cos(c + dx)} \sec^2(c + dx) \tan(c + dx)}{3d}$$

```
output 1/8*a^(3/2)*(11*A+14*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))
/d+1/8*a^2*(11*A+14*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/12*a^2*(7*A+6
*B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/3*a*A*sec(d*x+c)^2*(a
+a*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.89.2 Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.80

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{a \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^3(c + dx) (3\sqrt{2}(11A + 14B) \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right))$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^3*(3*Sqrt[2]*(11*A + 14*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^3 + (7*(7*A + 6*B) + 4*(11*A + 6*B)*Cos[c + d*x] + (33*A + 42*B)*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.89.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^4(c + dx) (a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx \\ & \quad \downarrow \text{3454} \\ & \frac{1}{3} \int \frac{1}{2} \sqrt{\cos(c + dx)a + a(a(7A + 6B) + 3a(A + 2B) \cos(c + dx))} \sec^3(c + dx) dx + \\ & \quad \frac{aA \tan(c + dx) \sec^2(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{6} \int \frac{\sqrt{\cos(c+dx)a+a}(a(7A+6B)+3a(A+2B)\cos(c+dx))\sec^3(c+dx)dx + aA \tan(c+dx) \sec^2(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} dx +$$

↓ 3042

$$\frac{1}{6} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a(7A+6B)+3a(A+2B)\sin(c+dx+\frac{\pi}{2}))\sin(c+dx+\frac{\pi}{2})^3 dx + aA \tan(c+dx) \sec^2(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} dx +$$

↓ 3459

$$\frac{1}{6} \left(\frac{3}{4} a(11A+14B) \int \frac{\sqrt{\cos(c+dx)a+a} \sec^2(c+dx) dx + \frac{a^2(7A+6B) \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a \cos(c+dx)+a}} \right) +$$

↓ 3042

$$\frac{1}{6} \left(\frac{3}{4} a(11A+14B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} dx + \frac{a^2(7A+6B) \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a \cos(c+dx)+a}} \right) +$$

↓ 3251

$$\frac{1}{6} \left(\frac{3}{4} a(11A+14B) \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a} \sec(c+dx) dx + \frac{a \tan(c+dx)}{d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(7A+6B) \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) +$$

↓ 3042

$$\frac{1}{6} \left(\frac{3}{4} a(11A+14B) \left(\frac{1}{2} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} dx + \frac{a \tan(c+dx)}{d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(7A+6B) \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) +$$

↓ 3252

$$\frac{1}{6} \left(\frac{3}{4} a(11A + 14B) \left(\frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{a \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \right) + \frac{a^2(7A + 6B) \tan(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right)$$

$$\frac{aA \tan(c + dx) \sec^2(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}$$

↓ 219

$$\frac{1}{6} \left(\frac{a^2(7A + 6B) \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} + \frac{3}{4} a(11A + 14B) \left(\frac{\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} + \frac{a \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right)$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*Sqrt[a + a*Cos[c + d*x]]*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((a^2*(7*A + 6*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*a*(11*A + 14*B)*((Sqrt[a]*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)/6`

3.89.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*SIN[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*SIN[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*SIN[e + f*x]]*(c + d*SIN[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*SIN[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*SIN[e + f*x])^(m - 1)*(c + d*SIN[e + f*x])^(n + 1)*Simp[A*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*SIN[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*SIN[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*SIN[e + f*x]]*(c + d*SIN[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.89.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1282 vs. $2(144) = 288$.

Time = 5.88 (sec) , antiderivative size = 1283, normalized size of antiderivative = 7.82

method	result	size
parts	Expression too large to display	1283
default	Expression too large to display	1326

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNV
ERBOSE)
```

```
output 1/6*A*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-264*a*(1
n(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(
1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)*a^(1/2)-2*a)))*sin(1/2*d*x+1/2*c)^6+132*(2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)
^2)^(1/2)*a^(1/2)+3*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2
*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+3*ln(-4
/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*s
in(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^4-22*(16*2^(
1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+9*ln(4/(2*cos(1/2*d*x+1/2*c)+
2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1
/2)*a^(1/2)+2*a))*a+9*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(
1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin
(1/2*d*x+1/2*c)^2+33*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/
2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+33*ln(
-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a
*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+126*2^(1/2)*(a*sin(1/2*d*x+1/
2*c)^2)^(1/2)*a^(1/2))/(2*cos(1/2*d*x+1/2*c)+2^(1/2))^3/(2*cos(1/2*d*x+1/2
*c)-2^(1/2))^3/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+1/4*B*a
^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(28*2^(1/2)*a*...
```

3.89.5 Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.23

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{3 \left((11A + 14B)a \cos(dx + c)^4 + (11A + 14B)a \cos(dx + c)^3 \right) \sqrt{a} \log \left(\frac{a \cos(dx + c)^3 - 7a \cos(dx + c)^2 - 4a \cos(dx + c) + a}{\cos(dx + c)} \right)}{\cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")`

output `1/96*(3*((11*A + 14*B)*a*cos(d*x + c)^4 + (11*A + 14*B)*a*cos(d*x + c)^3)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*(3*(11*A + 14*B)*a*cos(d*x + c)^2 + 2*(11*A + 6*B)*a*cos(d*x + c) + 8*A*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(d*cos(d*x + c)^4 + d*cos(d*x + c)^3)`

3.89.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.89.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7567 vs. 2(144) = 288.

Time = 154.99 (sec) , antiderivative size = 7567, normalized size of antiderivative = 46.14

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Too large to display}$$

output `-1/96*sqrt(2)*(3*sqrt(2)*(11*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 14*B*a*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))) + 4*(132*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 + 168*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 - 176*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 192*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 63*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) + 54*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^3)*sqrt(a)/d`

3.89.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)^4} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^4,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^4, x)`

3.90 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.90.1	Optimal result	937
3.90.2	Mathematica [A] (verified)	938
3.90.3	Rubi [A] (verified)	938
3.90.4	Maple [B] (verified)	942
3.90.5	Fricas [A] (verification not implemented)	943
3.90.6	Sympy [F(-1)]	944
3.90.7	Maxima [B] (verification not implemented)	944
3.90.8	Giac [A] (verification not implemented)	945
3.90.9	Mupad [F(-1)]	946

3.90.1 Optimal result

Integrand size = 33, antiderivative size = 209

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^5(c + dx) dx = \frac{a^{3/2}(75A + 88B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{64d} + \frac{a^2(75A + 88B) \tan(c + dx)}{64d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(75A + 88B) \sec(c + dx) \tan(c + dx)}{96d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(9A + 8B) \sec^2(c + dx) \tan(c + dx)}{24d\sqrt{a + a \cos(c + dx)}} + \frac{aA\sqrt{a + a \cos(c + dx)} \sec^3(c + dx) \tan(c + dx)}{4d}$$

output

```
1/64*a^(3/2)*(75*A+88*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2)
)/d+1/64*a^2*(75*A+88*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/96*a^2*(75*
A+88*B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/24*a^2*(9*A+8*B)*
sec(d*x+c)^2*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/4*a*A*sec(d*x+c)^3*(a+a
*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```


3.90.2 Mathematica [A] (verified)

Time = 0.99 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.72

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \frac{a \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^4(c + dx) (6\sqrt{2}(75A + 88B) \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right))}{768d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^4*(6*sqrt[2]*(75*A + 88*B)*ArcTanh[sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^4 + (492*A + 352*B + (1155*A + 1048*B)*Cos[c + d*x] + 4*(75*A + 88*B)*Cos[2*(c + d*x)] + 225*A*cos[3*(c + d*x)] + 264*B*cos[3*(c + d*x)])*Sin[(c + d*x)/2]))/(768*d)`

3.90.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 201, normalized size of antiderivative = 0.96, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^5(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx \\ & \quad \downarrow \text{3454} \\ & \frac{1}{4} \int \frac{1}{2} \sqrt{\cos(c + dx)a + a(a(9A + 8B) + a(5A + 8B) \cos(c + dx))} \sec^4(c + dx) dx + \\ & \quad \frac{aA \tan(c + dx) \sec^3(c + dx) \sqrt{a \cos(c + dx) + a}}{4d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{8} \int \frac{\sqrt{\cos(c+dx)a+a}(a(9A+8B)+a(5A+8B)\cos(c+dx))\sec^4(c+dx)dx + aA \tan(c+dx) \sec^3(c+dx) \sqrt{a \cos(c+dx)+a}}{4d}$$

↓ 3042

$$\frac{1}{8} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a(9A+8B)+a(5A+8B)\sin(c+dx+\frac{\pi}{2}))\sin^4(c+dx+\frac{\pi}{2})dx + aA \tan(c+dx) \sec^3(c+dx) \sqrt{a \cos(c+dx)+a}}{4d}$$

↓ 3459

$$\frac{1}{8} \left(\frac{1}{6} a(75A+88B) \int \frac{\sqrt{\cos(c+dx)a+a} \sec^3(c+dx) dx + \frac{a^2(9A+8B) \tan(c+dx) \sec^2(c+dx)}{3d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a \cos(c+dx)+a}} \right) +$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} a(75A+88B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} \sin^3(c+dx+\frac{\pi}{2}) dx + \frac{a^2(9A+8B) \tan(c+dx) \sec^2(c+dx)}{3d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a \cos(c+dx)+a}} \right) +$$

↓ 3251

$$\frac{1}{8} \left(\frac{1}{6} a(75A+88B) \left(\frac{3}{4} \int \frac{\sqrt{\cos(c+dx)a+a} \sec^2(c+dx) dx + \frac{a \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(9A+8B) \tan(c+dx)}{3d \sqrt{a \cos(c+dx)+a}} \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} a(75A+88B) \left(\frac{3}{4} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} \sin^2(c+dx+\frac{\pi}{2}) dx + \frac{a \tan(c+dx) \sec(c+dx)}{2d \sqrt{a \cos(c+dx)+a}}}{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(9A+8B) \tan(c+dx)}{3d \sqrt{a \cos(c+dx)+a}} \right)$$

↓ 3251

$$\frac{1}{8} \left(\frac{1}{6} a(75A + 88B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a} \sec(c+dx) dx}{d\sqrt{a\cos(c+dx)+a}} + \frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a\cos(c+dx)+a}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} a(75A + 88B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a\cos(c+dx)+a}}{4d}$$

↓ 3252

$$\frac{1}{8} \left(\frac{1}{6} a(75A + 88B) \left(\frac{3}{4} \left(\frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a\cos(c+dx)+a}}{4d}$$

↓ 219

$$\frac{1}{8} \left(\frac{a^2(9A + 8B) \tan(c+dx) \sec^2(c+dx)}{3d\sqrt{a\cos(c+dx)+a}} + \frac{1}{6} a(75A + 88B) \left(\frac{3}{4} \left(\frac{\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a\cos(c+dx)+a}} \right)}{d} + \frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) \sqrt{a\cos(c+dx)+a}}{4d}$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*A*sqrt[a + a*cos[c + d*x]]*sec[c + d*x]^3*tan[c + d*x])/(4*d) + ((a^2*(9*A + 8*B)*sec[c + d*x]^2*tan[c + d*x])/(3*d*sqrt[a + a*cos[c + d*x]]) + (a*(75*A + 88*B)*((a*sec[c + d*x]*tan[c + d*x])/(2*d*sqrt[a + a*cos[c + d*x]])) + (3*((sqrt[a]*ArcTanh[(sqrt[a]*sin[c + d*x])/sqrt[a + a*cos[c + d*x]])]/d + (a*tan[c + d*x])/(d*sqrt[a + a*cos[c + d*x]]))))/4)/6)/8`

3.90.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`
- rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3459 Int[Sqrt[(a_) + (b_.)*sin[(e_) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_) + (
f_.)*(x_)])*((c_) + (d_.)*sin[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.90.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1616 vs. $2(185) = 370$.

Time = 6.36 (sec) , antiderivative size = 1617, normalized size of antiderivative = 7.74

method	result	size
parts	Expression too large to display	1617
default	Expression too large to display	1651

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNV
ERBOSE)
```

output

```

1/8*A*a^(1/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(1200*a*(1
n(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(
1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)*a^(1/2)-2*a)))*sin(1/2*d*x+1/2*c)^8-1200*(2^(1/2)*(a*sin(1/2*d*x+1/2*c)^
2)^(1/2)*a^(1/2)+2*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*
d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+2*ln(-4/
(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*si
n(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^6+200*(11*2^(
1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+9*ln(4/(2*cos(1/2*d*x+1/2*c)+
2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1
/2)*a^(1/2)+2*a))*a+9*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(
1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin
(1/2*d*x+1/2*c)^4+(-1460*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-60
0*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a-600*ln(-4/(2*cos(1/2*d*x+
1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c
)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^2+362*2^(1/2)*(a*sin(1/2*d*
x+1/2*c)^2)^(1/2)*a^(1/2)+75*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*
a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a...

```

3.90.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.05

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \frac{3 \left((75A + 88B)a \cos(dx + c)^5 + (75A + 88B)a \cos(dx + c)^4 \right) \sqrt{a} \log \left(\frac{a \cos(dx + c)^3 - 7a \cos(dx + c)^2 - 4 \sqrt{a} \cos(dx + c) + a}{\cos(dx + c)} \right) + \dots}{\cos(dx + c)}$$

input

```

integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorith
m="fricas")

```

output

```

1/768*(3*((75*A + 88*B)*a*cos(d*x + c)^5 + (75*A + 88*B)*a*cos(d*x + c)^4)
*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c
) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + co
s(d*x + c)^2)) + 4*(3*(75*A + 88*B)*a*cos(d*x + c)^3 + 2*(75*A + 88*B)*a*c
os(d*x + c)^2 + 8*(15*A + 8*B)*a*cos(d*x + c) + 48*A*a)*sqrt(a*cos(d*x + c
) + a)*sin(d*x + c))/(d*cos(d*x + c)^5 + d*cos(d*x + c)^4)

```

3.90. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.90.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output `Timed out`

3.90.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10504 vs. $2(185) = 370$.

Time = 154.62 (sec) , antiderivative size = 10504, normalized size of antiderivative = 50.26

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm m="maxima")`

output

```

-1/768*(3*(140*a*cos(8*d*x + 8*c)^2*sin(3/2*d*x + 3/2*c) + 2240*a*cos(6*d*
x + 6*c)^2*sin(3/2*d*x + 3/2*c) + 5040*a*cos(4*d*x + 4*c)^2*sin(3/2*d*x +
3/2*c) + 2240*a*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 140*a*sin(8*d*x
+ 8*c)^2*sin(3/2*d*x + 3/2*c) + 2240*a*sin(6*d*x + 6*c)^2*sin(3/2*d*x + 3/
2*c) + 5040*a*sin(4*d*x + 4*c)^2*sin(3/2*d*x + 3/2*c) + 2240*a*sin(2*d*x +
2*c)^2*sin(3/2*d*x + 3/2*c) + 4064*a*cos(7/2*d*x + 7/2*c)*sin(2*d*x + 2*c
) + 336*a*cos(5/2*d*x + 5/2*c)*sin(2*d*x + 2*c) - 240*a*cos(3/2*d*x + 3/2*
c)*sin(2*d*x + 2*c) + 1360*a*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) - 36*(a
*sin(8*d*x + 8*c) + 4*a*sin(6*d*x + 6*c) + 6*a*sin(4*d*x + 4*c) + 4*a*sin(
2*d*x + 2*c))*cos(21/2*d*x + 21/2*c) + 140*(a*sin(8*d*x + 8*c) + 4*a*sin(6
*d*x + 6*c) + 6*a*sin(4*d*x + 4*c) + 4*a*sin(2*d*x + 2*c))*cos(19/2*d*x +
19/2*c) + 456*(a*sin(8*d*x + 8*c) + 4*a*sin(6*d*x + 6*c) + 6*a*sin(4*d*x +
4*c) + 4*a*sin(2*d*x + 2*c))*cos(17/2*d*x + 17/2*c) + 4*(280*a*cos(6*d*x
+ 6*c)*sin(3/2*d*x + 3/2*c) + 420*a*cos(4*d*x + 4*c)*sin(3/2*d*x + 3/2*c)
+ 280*a*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) - 290*a*sin(15/2*d*x + 15/2*
c) - 596*a*sin(13/2*d*x + 13/2*c) - 780*a*sin(11/2*d*x + 11/2*c) - 750*a*s
in(9/2*d*x + 9/2*c) - 254*a*sin(7/2*d*x + 7/2*c) - 21*a*sin(5/2*d*x + 5/2*
c) + 85*a*sin(3/2*d*x + 3/2*c))*cos(8*d*x + 8*c) + 2320*(2*a*sin(6*d*x + 6
*c) + 3*a*sin(4*d*x + 4*c) + 2*a*sin(2*d*x + 2*c))*cos(15/2*d*x + 15/2*c)
+ 4768*(2*a*sin(6*d*x + 6*c) + 3*a*sin(4*d*x + 4*c) + 2*a*sin(2*d*x + 2...

```

3.90.8 Giac [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.44

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx =$$

$$\sqrt{2} \left(3 \sqrt{2} (75 A \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 88 B \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \log \left(\frac{-2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)}{2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)} \right) + \frac{4}{1800} \right)$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")`

output `-1/768*sqrt(2)*(3*sqrt(2)*(75*A*a*sgn(cos(1/2*d*x + 1/2*c)) + 88*B*a*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))) + 4*(1800*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^7 + 2112*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^7 - 3300*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 - 3872*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 + 2190*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 2416*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 543*A*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) - 504*B*a*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^4)*sqrt(a)/d`

3.90.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)^5} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^5,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^5, x)`

3.91 $\int \cos^2(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.91.1	Optimal result	947
3.91.2	Mathematica [A] (verified)	948
3.91.3	Rubi [A] (verified)	948
3.91.4	Maple [A] (verified)	952
3.91.5	Fricas [A] (verification not implemented)	953
3.91.6	Sympy [F(-1)]	953
3.91.7	Maxima [A] (verification not implemented)	954
3.91.8	Giac [A] (verification not implemented)	954
3.91.9	Mupad [F(-1)]	955

3.91.1 Optimal result

Integrand size = 33, antiderivative size = 237

$$\int \cos^2(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx = \frac{2a^3(803A+710B) \sin(c+dx)}{495d\sqrt{a+a \cos(c+dx)}} + \frac{2a^3(209A+194B) \cos^3(c+dx) \sin(c+dx)}{693d\sqrt{a+a \cos(c+dx)}} - \frac{4a^2(803A+710B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{3465d} + \frac{2a^2(11A+14B) \cos^3(c+dx)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{99d} + \frac{2a(803A+710B)(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{1155d} + \frac{2aB \cos^3(c+dx)(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{11d}$$

```
output 2/1155*a*(803*A+710*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/11*a*B*cos(d*x+c)^3*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/495*a^3*(803*A+710*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/693*a^3*(209*A+194*B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-4/3465*a^2*(803*A+710*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+2/99*a^2*(11*A+14*B)*cos(d*x+c)^3*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.91.2 Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.54

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))}(124366A + 114640B + (68552A + 69890B) \cos(c + dx) + 16($$

input `Integrate[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x]])*(124366*A + 114640*B + (68552*A + 69890*B)*Cos[c + d*x] + 16*(1397*A + 1625*B)*Cos[2*(c + d*x)] + 5720*A*Cos[3*(c + d*x)] + 8675*B*Cos[3*(c + d*x)] + 770*A*Cos[4*(c + d*x)] + 2240*B*Cos[4*(c + d*x)] + 315*B*Cos[5*(c + d*x)])*Tan[(c + d*x)/2])/(27720*d)`

3.91.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3238, 27, 3042, 3230, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3455} \\ & \frac{2}{11} \int \frac{1}{2} \cos^2(c + dx)(\cos(c + dx)a + a)^{3/2}(a(11A + 6B) + a(11A + 14B) \cos(c + dx)) dx + \\ & \quad \frac{2aB \sin(c + dx) \cos^3(c + dx)(a \cos(c + dx) + a)^{3/2}}{11d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{11} \int \cos^2(c+dx) (\cos(c+dx)a+a)^{3/2} (a(11A+6B) + a(11A+14B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{11d}$$

↓ 3042

$$\frac{1}{11} \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{3/2} \left(a(11A+6B) + a(11A+14B) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2aB \sin(c+dx) \cos^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{11d}$$

↓ 3455

$$\frac{1}{11} \left(\frac{2}{9} \int \frac{1}{2} \cos^2(c+dx) \sqrt{\cos(c+dx)a+a} (3(55A+46B)a^2 + (209A+194B) \cos(c+dx)a^2) dx + \frac{2a^2(11A+14B)}{11d} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{11d}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \int \cos^2(c+dx) \sqrt{\cos(c+dx)a+a} (3(55A+46B)a^2 + (209A+194B) \cos(c+dx)a^2) dx + \frac{2a^2(11A+14B)}{11d} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{11d}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} (3(55A+46B)a^2 + (209A+194B) \sin\left(c+dx+\frac{\pi}{2}\right)a^2) dx + \frac{2a^2(11A+14B)}{11d} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{11d}$$

↓ 3460

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A+710B) \int \cos^2(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{2a^3(209A+194B) \sin(c+dx) \cos^3(c+dx)}{7d \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^2(11A+14B)}{11d} \right) + \frac{2aB \sin(c+dx) \cos^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{11d}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A + 710B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right) a + adx} + \frac{2a^3 (209A + 194B) \sin(c + dx) \cos(c + dx)}{7d \sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \\ \downarrow \text{3238}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A + 710B) \left(\frac{2 \int \frac{1}{2} (3a - 2a \cos(c + dx)) \sqrt{\cos(c + dx) a + adx}}{5a} + \frac{2 \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \\ \downarrow \text{27}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A + 710B) \left(\frac{\int (3a - 2a \cos(c + dx)) \sqrt{\cos(c + dx) a + adx}}{5a} + \frac{2 \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A + 710B) \left(\frac{\int (3a - 2a \sin \left(c + dx + \frac{\pi}{2} \right)) \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right) a + adx}}{5a} + \frac{2 \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \\ \downarrow \text{3230}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A + 710B) \left(\frac{\frac{7}{3} a \int \sqrt{\cos(c + dx) a + adx} - \frac{4a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}}{5a} + \frac{2 \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (803A + 710B) \left(\frac{\frac{7}{3} a \int \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right) a + adx} - \frac{4a \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d}}{5a} + \frac{2 \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{5ad} \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \\ \downarrow \text{3125}$$

$$\frac{1}{11} \left(\frac{2a^2(11A + 14B) \sin(c + dx) \cos^3(c + dx) \sqrt{a \cos(c + dx) + a}}{9d} + \frac{1}{9} \left(\frac{2a^3(209A + 194B) \sin(c + dx) \cos^3(c + dx)}{7d \sqrt{a \cos(c + dx) + a}} \right. \right. \\ \left. \left. + \frac{2aB \sin(c + dx) \cos^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{11d} \right) \right)$$

input `Int[Cos[c + d*x]^2*(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(2*a*B*Cos[c + d*x]^3*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(11*d) + ((2*a^2*(11*A + 14*B)*Cos[c + d*x]^3*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(9*d) + ((2*a^3*(209*A + 194*B)*Cos[c + d*x]^3*Sqrt[a + a*Cos[c + d*x]])/(7*d*Sqrt[a + a*Cos[c + d*x]]) + (3*a^2*(803*A + 710*B)*((2*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*a*d) + ((14*a^2*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]])) - (4*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(5*a))/7)/9)/11`

3.91.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

```
rule 3238 Int[sin[(e_.) + (f_.)*(x_)]^2*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_),
x_Symbol] := Simp[(-Cos[e + f*x])*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2
))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*(b*(m + 1) - a*Si
n[e + f*x]), x], x] /; FreeQ[{a, b, e, f, m}, x] && EqQ[a^2 - b^2, 0] && !
LtQ[m, -2^(-1)]
```

```
rule 3455 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.91.4 Maple [A] (verified)

Time = 11.39 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.60

method	result
default	$\frac{8 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(-2520B \left(\sin^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (1540A + 10780B) \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-5940A - 18810B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (1540A + 10780B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-5940A - 18810B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8B}{3465 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$
parts	$\frac{8A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(140 \left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 20 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 39 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 52 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 104\right) \sqrt{2} + 8B}{315 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} d$

```
input int(cos(d*x+c)^2*(a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

3.91. $\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

output $8/3465*\cos(1/2*d*x+1/2*c)*a^3*\sin(1/2*d*x+1/2*c)*(-2520*B*\sin(1/2*d*x+1/2*c)^{10}+(1540*A+10780*B)*\sin(1/2*d*x+1/2*c)^8+(-5940*A-18810*B)*\sin(1/2*d*x+1/2*c)^6+(9009*A+17325*B)*\sin(1/2*d*x+1/2*c)^4+(-6930*A-9240*B)*\sin(1/2*d*x+1/2*c)^2+3465*A+3465*B)*2^{(1/2)}/(a*\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/d$

3.91.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.58

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{2(315Ba^2 \cos(dx + c)^5 + 35(11A + 32B)a^2 \cos(dx + c)^4 + 5(286A + 355B)a^2 \cos(dx + c)^3 + 3(803A + 710B)a^2 \cos(dx + c)^2 + 4(803A + 710B)a^2 \cos(dx + c) + 8(803A + 710B)a^2)\sqrt{a \cos(dx + c) + a} \sin(dx + c)}{(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output $2/3465*(315*B*a^2*\cos(d*x + c)^5 + 35*(11*A + 32*B)*a^2*\cos(d*x + c)^4 + 5*(286*A + 355*B)*a^2*\cos(d*x + c)^3 + 3*(803*A + 710*B)*a^2*\cos(d*x + c)^2 + 4*(803*A + 710*B)*a^2*\cos(d*x + c) + 8*(803*A + 710*B)*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/(d*\cos(d*x + c) + d)$

3.91.6 Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output Timed out

3.91.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.87

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{22(35\sqrt{2}a^2 \sin(\frac{9}{2}dx + \frac{9}{2}c) + 225\sqrt{2}a^2 \sin(\frac{7}{2}dx + \frac{7}{2}c) + 756\sqrt{2}a^2 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 2100\sqrt{2}a^2 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 8190\sqrt{2}a^2 \sin(\frac{1}{2}dx + \frac{1}{2}c))A\sqrt{a} + 5(63\sqrt{2}a^2 \sin(\frac{11}{2}dx + \frac{11}{2}c) + 385\sqrt{2}a^2 \sin(\frac{9}{2}dx + \frac{9}{2}c) + 1287\sqrt{2}a^2 \sin(\frac{7}{2}dx + \frac{7}{2}c) + 3465\sqrt{2}a^2 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 8778\sqrt{2}a^2 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 31878\sqrt{2}a^2 \sin(\frac{1}{2}dx + \frac{1}{2}c))B\sqrt{a}}{d}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output 1/55440*(22*(35*sqrt(2)*a^2*sin(9/2*d*x + 9/2*c) + 225*sqrt(2)*a^2*sin(7/2*d*x + 7/2*c) + 756*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) + 2100*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 8190*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + 5*(63*sqrt(2)*a^2*sin(11/2*d*x + 11/2*c) + 385*sqrt(2)*a^2*sin(9/2*d*x + 9/2*c) + 1287*sqrt(2)*a^2*sin(7/2*d*x + 7/2*c) + 3465*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) + 8778*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 31878*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d
```

3.91.8 Giac [A] (verification not implemented)

Time = 3.28 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.08

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(315Ba^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{11}{2}dx + \frac{11}{2}c) + 385(2Aa^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)))}{d}$$

```
input integrate(cos(d*x+c)^2*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/55440*sqrt(2)*(315*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(11/2*d*x + 11/2*c) + 385*(2*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 5*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(9/2*d*x + 9/2*c) + 495*(10*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 13*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(7/2*d*x + 7/2*c) + 693*(24*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 25*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(5/2*d*x + 5/2*c) + 2310*(20*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 19*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 6930*(26*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 23*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d
```

3.91.9 Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^2 (A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2), x)`

3.92 $\int \cos(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.92.1	Optimal result	956
3.92.2	Mathematica [A] (verified)	957
3.92.3	Rubi [A] (verified)	957
3.92.4	Maple [A] (verified)	960
3.92.5	Fricas [A] (verification not implemented)	961
3.92.6	Sympy [F(-1)]	961
3.92.7	Maxima [A] (verification not implemented)	962
3.92.8	Giac [A] (verification not implemented)	962
3.92.9	Mupad [F(-1)]	963

3.92.1 Optimal result

Integrand size = 31, antiderivative size = 175

$$\int \cos(c+dx)(a + a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx = \frac{64a^3(15A+13B) \sin(c+dx)}{315d\sqrt{a+a \cos(c+dx)}} + \frac{16a^2(15A+13B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{315d} + \frac{2a(15A+13B)(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{105d} + \frac{2(9A-2B)(a+a \cos(c+dx))^{5/2} \sin(c+dx)}{63d} + \frac{2B(a+a \cos(c+dx))^{7/2} \sin(c+dx)}{9ad}$$

output

```
2/105*a*(15*A+13*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/63*(9*A-2*B)*(a+a*cos(d*x+c))^(5/2)*sin(d*x+c)/d+2/9*B*(a+a*cos(d*x+c))^(7/2)*sin(d*x+c)/a/d+64/315*a^3*(15*A+13*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+16/315*a^2*(15*A+13*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.92.2 Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.60

$$\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))(6240A + 5653B + (3030A + 3116B) \cos(c + dx) + 8(90A + 1$$

input `Integrate[Cos[c + d*x]*(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*(6240*A + 5653*B + (3030*A + 3116*B)*Cos[c + d*x] + 8*(90*A + 127*B)*Cos[2*(c + d*x)] + 90*A*Cos[3*(c + d*x)] + 260*B*Cos[3*(c + d*x)] + 35*B*Cos[4*(c + d*x)])*Tan[(c + d*x)/2])/(1260*d)`

3.92.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3447} \\ & \int (a \cos(c + dx) + a)^{5/2} (A \cos(c + dx) + B \cos^2(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\ & \quad \downarrow \text{3502} \end{aligned}$$

$$\begin{aligned}
& \frac{2 \int \frac{1}{2} (\cos(c+dx)a+a)^{5/2} (7aB+a(9A-2B)\cos(c+dx)) dx}{9a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 27 \\
& \frac{\int (\cos(c+dx)a+a)^{5/2} (7aB+a(9A-2B)\cos(c+dx)) dx}{9a} + \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 3042 \\
& \frac{\int (\sin(c+dx+\frac{\pi}{2})a+a)^{5/2} (7aB+a(9A-2B)\sin(c+dx+\frac{\pi}{2})) dx}{9a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 3230 \\
& \frac{\frac{3}{7}a(15A+13B) \int (\cos(c+dx)a+a)^{5/2} dx + \frac{2a(9A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d}}{9a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 3042 \\
& \frac{\frac{3}{7}a(15A+13B) \int (\sin(c+dx+\frac{\pi}{2})a+a)^{5/2} dx + \frac{2a(9A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d}}{9a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 3126 \\
& \frac{\frac{3}{7}a(15A+13B) \left(\frac{8}{5}a \int (\cos(c+dx)a+a)^{3/2} dx + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) + \frac{2a(9A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d}}{9a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 3042 \\
& \frac{\frac{3}{7}a(15A+13B) \left(\frac{8}{5}a \int (\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} dx + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) + \frac{2a(9A-2B)\sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d}}{9a} + \\
& \quad \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{7/2}}{9ad} \\
& \quad \downarrow 3126
\end{aligned}$$

$$\frac{\frac{3}{7}a(15A + 13B) \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{\cos(c + dx)a + adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) + \frac{2a(9A-2B) \sin(c+dx)}{5d}}{9a} \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{7/2}}{9ad}$$

↓ 3042

$$\frac{\frac{3}{7}a(15A + 13B) \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{\sin(c + dx + \frac{\pi}{2})a + adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) + \frac{2a(9A-2B) \sin(c+dx)}{5d}}{9a} \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{7/2}}{9ad}$$

↓ 3125

$$\frac{\frac{3}{7}a(15A + 13B) \left(\frac{8}{5}a \left(\frac{8a^2 \sin(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) + \frac{2a(9A-2B) \sin(c+dx)}{5d}}{9a} \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{7/2}}{9ad}$$

input `Int[Cos[c + d*x]*(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*(a + a*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*a*d) + ((2*a*(9*A - 2*B)*(a + a*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*d) + (3*a*(15*A + 13*B))*((2*a*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (8*a*((8*a^2*SIN[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (2*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/5)/7)/(9*a)`

3.92.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[-2*b*(Cos [c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq Q[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos [c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos [e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.92.4 Maple [A] (verified)

Time = 4.78 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.70

method	result
default	$\frac{8 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(140B \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-90A - 540B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (315A + 819B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-420A - 630B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8B}{315 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{8A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(6 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 3 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\right) \sqrt{2}}{21 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{8B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right)}{21 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

3.92. $\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

input `int(cos(d*x+c)*(a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `8/315*cos(1/2*d*x+1/2*c)*a^3*sin(1/2*d*x+1/2*c)*(140*B*sin(1/2*d*x+1/2*c)^8+(-90*A-540*B)*sin(1/2*d*x+1/2*c)^6+(315*A+819*B)*sin(1/2*d*x+1/2*c)^4+(-420*A-630*B)*sin(1/2*d*x+1/2*c)^2+315*A+315*B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.92.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.66

$$\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{2(35Ba^2 \cos(dx + c)^4 + 5(9A + 26B)a^2 \cos(dx + c)^3 + 3(60A + 73B)a^2 \cos(dx + c)^2 + (345A + 292B)a^2 \cos(dx + c) + 2(345A + 292B)a^2) \sin(dx + c)}{315(d \cos(dx + c) + d)}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `2/315*(35*B*a^2*cos(d*x + c)^4 + 5*(9*A + 26*B)*a^2*cos(d*x + c)^3 + 3*(60*A + 73*B)*a^2*cos(d*x + c)^2 + (345*A + 292*B)*a^2*cos(d*x + c) + 2*(345*A + 292*B)*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c) + d)`

3.92.6 Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.92.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.98

$$\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{30(3\sqrt{2}a^2 \sin(\frac{7}{2}dx + \frac{7}{2}c) + 21\sqrt{2}a^2 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 77\sqrt{2}a^2 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 315\sqrt{2}a^2 \sin(\frac{1}{2}dx + \frac{1}{2}c))A\sqrt{a} + (35\sqrt{2}a^2 \sin(\frac{9}{2}dx + \frac{9}{2}c) + 225\sqrt{2}a^2 \sin(\frac{7}{2}dx + \frac{7}{2}c) + 756\sqrt{2}a^2 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 2100\sqrt{2}a^2 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 8190\sqrt{2}a^2 \sin(\frac{1}{2}dx + \frac{1}{2}c))B\sqrt{a}}{d}$$

```
input integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output 1/2520*(30*(3*sqrt(2)*a^2*sin(7/2*d*x + 7/2*c) + 21*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) + 77*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 315*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*A*sqrt(a) + (35*sqrt(2)*a^2*sin(9/2*d*x + 9/2*c) + 225*sqrt(2)*a^2*sin(7/2*d*x + 7/2*c) + 756*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) + 2100*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 8190*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*B*sqrt(a))/d
```

3.92.8 Giac [A] (verification not implemented)

Time = 1.13 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.22

$$\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(35Ba^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{9}{2}dx + \frac{9}{2}c) + 45(2Aa^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 5Ba^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(\frac{7}{2}dx + \frac{7}{2}c) + 126(5Aa^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 6Ba^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(\frac{5}{2}dx + \frac{5}{2}c) + 210(11Aa^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 10Ba^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(\frac{3}{2}dx + \frac{3}{2}c) + 630(15Aa^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 13Ba^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) \sin(\frac{1}{2}dx + \frac{1}{2}c)) \sqrt{a}}{d}$$

```
input integrate(cos(d*x+c)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/2520*sqrt(2)*(35*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(9/2*d*x + 9/2*c) + 45*(2*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 5*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(7/2*d*x + 7/2*c) + 126*(5*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 6*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(5/2*d*x + 5/2*c) + 210*(11*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 10*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 630*(15*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 13*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d
```

3.92.9 Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)(A + B \cos(c + dx))(a + a \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2), x)`

3.93 $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$

3.93.1	Optimal result	964
3.93.2	Mathematica [A] (verified)	964
3.93.3	Rubi [A] (verified)	965
3.93.4	Maple [A] (verified)	967
3.93.5	Fricas [A] (verification not implemented)	967
3.93.6	Sympy [F(-1)]	968
3.93.7	Maxima [A] (verification not implemented)	968
3.93.8	Giac [A] (verification not implemented)	969
3.93.9	Mupad [F(-1)]	969

3.93.1 Optimal result

Integrand size = 25, antiderivative size = 138

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{64a^3(7A + 5B) \sin(c + dx)}{105d\sqrt{a + a \cos(c + dx)}} + \frac{16a^2(7A + 5B)\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{105d} + \frac{2a(7A + 5B)(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{35d} + \frac{2B(a + a \cos(c + dx))^{5/2} \sin(c + dx)}{7d}$$

```
output 2/35*a*(7*A+5*B)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/7*B*(a+a*cos(d*x+c))^(5/2)*sin(d*x+c)/d+64/105*a^3*(7*A+5*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+16/105*a^2*(7*A+5*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.93.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.60

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (1246A + 1040B + (392A + 505B) \cos(c + dx) + 6(7A + 20B))}{210d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*(1246*A + 1040*B + (392*A + 505*B)*Cos[c + d*x] + 6*(7*A + 20*B)*Cos[2*(c + d*x)] + 15*B*Cos[3*(c + d*x)])*Tan[(c + d*x)/2])/(210*d)`

3.93.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {3042, 3230, 3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a \right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx \\
 & \quad \downarrow \text{3230} \\
 & \frac{1}{7}(7A + 5B) \int (\cos(c + dx)a + a)^{5/2} dx + \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7}(7A + 5B) \int \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^{5/2} dx + \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3126} \\
 & \frac{1}{7}(7A + 5B) \left(\frac{8}{5} a \int (\cos(c + dx)a + a)^{3/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) + \\
 & \quad \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7}(7A + 5B) \left(\frac{8}{5} a \int \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a \right)^{3/2} dx + \frac{2a \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) + \\
 & \quad \frac{2B \sin(c + dx)(a \cos(c + dx) + a)^{5/2}}{7d}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3126} \\
& \frac{1}{7}(7A + \\
5B) & \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{\cos(c+dx)a+adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)}{5d} \right. \\
& \left. \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d} \right) \\
& \downarrow \text{3042} \\
& \frac{1}{7}(7A + \\
5B) & \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)}{5d} \right. \\
& \left. \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d} \right) \\
& \downarrow \text{3125} \\
& \frac{1}{7}(7A + \\
5B) & \left(\frac{8}{5}a \left(\frac{8a^2 \sin(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} + \frac{2a \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} \right) + \frac{2a \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d} \right) \\
& \left. \frac{2B \sin(c+dx)(a \cos(c+dx)+a)^{5/2}}{7d} \right)
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*(a + a*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*d) + ((7*A + 5*B)*((2*a*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (8*a*((8*a^2*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (2*a*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/5)/7`

3.93.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq Q[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

3.93.4 Maple [A] (verified)

Time = 3.23 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.75

method	result
default	$\frac{8 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(-30B \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (21A + 105B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-70A - 140B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 105A + 105B\right)}{105 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{8A a^3 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(3 \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\right) \sqrt{2}}{15 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{8B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \left(6 \left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \dots\right)}{21 \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

input `int((a+cos(d*x+c))*a^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `8/105*cos(1/2*d*x+1/2*c)*a^3*sin(1/2*d*x+1/2*c)*(-30*B*sin(1/2*d*x+1/2*c)^6+(21*A+105*B)*sin(1/2*d*x+1/2*c)^4+(-70*A-140*B)*sin(1/2*d*x+1/2*c)^2+105*A+105*B)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.93.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.69

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{2 (15 B a^2 \cos(dx + c)^3 + 3 (7 A + 20 B) a^2 \cos(dx + c)^2 + (98 A + 115 B) a^2 \cos(dx + c) + 105 A + 105 B)}{105 (d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output
$$\frac{2}{105}*(15*B*a^2*\cos(d*x + c)^3 + 3*(7*A + 20*B)*a^2*\cos(d*x + c)^2 + (98*A + 115*B)*a^2*\cos(d*x + c) + (301*A + 230*B)*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/(d*\cos(d*x + c) + d)$$

3.93.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.93.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.01

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{14 (3 \sqrt{2} a^2 \sin(\frac{5}{2} dx + \frac{5}{2} c) + 25 \sqrt{2} a^2 \sin(\frac{3}{2} dx + \frac{3}{2} c) + 150 \sqrt{2} a^2 \sin(\frac{1}{2} dx + \frac{1}{2} c)) A \sqrt{a + a \cos(c + dx)}}{d}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output
$$\frac{1}{420}*(14*(3*\sqrt{2}*a^2*\sin(5/2*d*x + 5/2*c) + 25*\sqrt{2}*a^2*\sin(3/2*d*x + 3/2*c) + 150*\sqrt{2}*a^2*\sin(1/2*d*x + 1/2*c))*A*\sqrt{a} + 5*(3*\sqrt{2}*a^2*\sin(7/2*d*x + 7/2*c) + 21*\sqrt{2}*a^2*\sin(5/2*d*x + 5/2*c) + 77*\sqrt{2}*a^2*\sin(3/2*d*x + 3/2*c) + 315*\sqrt{2}*a^2*\sin(1/2*d*x + 1/2*c))*B*\sqrt{a})/d$$

3.93.8 Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.22

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{\sqrt{2} (15 B a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{7}{2} dx + \frac{7}{2} c) + 21 (2 A a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 5$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/420*sqrt(2)*(15*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(7/2*d*x + 7/2*c) + 21*(2*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 5*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(5/2*d*x + 5/2*c) + 35*(10*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 11*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(3/2*d*x + 3/2*c) + 525*(4*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 3*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*sin(1/2*d*x + 1/2*c))*sqrt(a)/d`**3.93.9 Mupad [F(-1)]**

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2),x)`output `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2), x)`

3.94 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec(c+dx) dx$

3.94.1	Optimal result	970
3.94.2	Mathematica [A] (verified)	971
3.94.3	Rubi [A] (verified)	971
3.94.4	Maple [B] (verified)	974
3.94.5	Fricas [A] (verification not implemented)	975
3.94.6	Sympy [F(-1)]	976
3.94.7	Maxima [A] (verification not implemented)	976
3.94.8	Giac [A] (verification not implemented)	976
3.94.9	Mupad [F(-1)]	977

3.94.1 Optimal result

Integrand size = 31, antiderivative size = 142

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec(c + dx) dx = \frac{2a^{5/2}A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} + \frac{2a^3(35A + 32B) \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{2a^2(5A + 8B)\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{15d} + \frac{2aB(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{5d}$$

```
output 2*a^(5/2)*A*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+2/5*a*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/15*a^3*(35*A+32*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/15*a^2*(5*A+8*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.94.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.73

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) (15\sqrt{2}A \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right))) + (80A + 89B + 2(5A + 14B) \cos(c + dx) + 3B \cos(2(c + dx))) \sin\left(\frac{1}{2}(c + dx)\right)}{15d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(15*Sqrt[2]*A*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]] + (80*A + 89*B + 2*(5*A + 14*B)*Cos[c + d*x] + 3*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(15*d)`

3.94.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{3455} \\ & \frac{2}{5} \int \frac{1}{2} (\cos(c + dx)a + a)^{3/2} (5aA + a(5A + 8B) \cos(c + dx)) \sec(c + dx) dx + \\ & \quad \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \\ & \quad \downarrow \text{27} \\ & \frac{1}{5} \int (\cos(c + dx)a + a)^{3/2} (5aA + a(5A + 8B) \cos(c + dx)) \sec(c + dx) dx + \\ & \quad \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \end{aligned}$$

3.94. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx$

↓ 3042

$$\frac{1}{5} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (5aA + a(5A + 8B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 3455

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{1}{2} \sqrt{\cos(c + dx)a + a} (15Aa^2 + (35A + 32B) \cos(c + dx)a^2) \sec(c + dx) dx + \frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \sqrt{\cos(c + dx)a + a} (15Aa^2 + (35A + 32B) \cos(c + dx)a^2) \sec(c + dx) dx + \frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a} (15Aa^2 + (35A + 32B) \sin(c + dx + \frac{\pi}{2})a^2)}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 3460

$$\frac{1}{5} \left(\frac{1}{3} \left(15a^2A \int \sqrt{\cos(c + dx)a + a} \sec(c + dx) dx + \frac{2a^3(35A + 32B) \sin(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(15a^2A \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a^3(35A + 32B) \sin(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 3252

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2a^3(35A + 32B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{30a^3 A \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} \right) + \frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 219

$$\frac{1}{5} \left(\frac{2a^2(5A + 8B) \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} + \frac{1}{3} \left(\frac{30a^{5/2} A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{2a^3(35A + 32B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{2aB \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x], x]`

output `(2*a*B*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + ((2*a^2*(5*A + 8*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((30*a^(5/2)*A*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (2*a^3*(35*A + 32*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/3)/5`

3.94.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.94.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 309 vs. 2(124) = 248.

Time = 6.21 (sec) , antiderivative size = 310, normalized size of antiderivative = 2.18

method	result
default	$a^{\frac{3}{2}} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(48B \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 40 \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} (A+4B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right)$
parts	$A a^{\frac{3}{2}} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-4\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 18\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} + 3 \ln\left(\frac{4\sqrt{2} a c \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{3 \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}\right) \right)$

3.94. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `1/30*a^(3/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(48*B*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^4-40*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*(A+4*B)*sin(1/2*d*x+1/2*c)^2+15*A*2^(1/2)*ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+15*A*2^(1/2)*ln(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+180*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+240*B*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/sin(1/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.94.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.25

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{15 (Aa^2 \cos(dx + c) + Aa^2) \sqrt{a} \log \left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4 \sqrt{a} \cos(dx+c) + a \sqrt{a} (\cos(dx+c) - 2) \sin(dx+c) + 8a}{\cos(dx+c)^3 + \cos(dx+c)^2} \right)}{30(d \cos(dx+c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")`

output `1/30*(15*(A*a^2*cos(d*x + c) + A*a^2)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*(3*B*a^2*cos(d*x + c)^2 + (5*A + 14*B)*a^2*cos(d*x + c) + (40*A + 43*B)*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(d*cos(d*x + c) + d)`

3.94.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `Timed out`

3.94.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.43

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{(3\sqrt{2}a^2 \sin(\frac{5}{2}dx + \frac{5}{2}c) + 25\sqrt{2}a^2 \sin(\frac{3}{2}dx + \frac{3}{2}c) + 150\sqrt{2}a^2 \sin(\frac{1}{2}dx + \frac{1}{2}c))B\sqrt{a}}{30d}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `1/30*(3*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) + 25*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 150*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*B*sqrt(a)/d`

3.94.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.42

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{\sqrt{2} \left(48 B a^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{1}{2}dx + \frac{1}{2}c)^5 - 40 A a^2 \operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) \sin(\frac{1}{2}dx + \frac{1}{2}c) \right)}{30d}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output $\frac{1}{30}\sqrt{2}(48B^2a^2\operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))\sin(\frac{1}{2}dx + \frac{1}{2}c)^5 - 40A^2a^2\operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))\sin(\frac{1}{2}dx + \frac{1}{2}c)^3 - 160B^2a^2\operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))\sin(\frac{1}{2}dx + \frac{1}{2}c)^3 - 15\sqrt{2}A^2a^2\log(\operatorname{abs}(-2\sqrt{2} + 4\sin(\frac{1}{2}dx + \frac{1}{2}c))\operatorname{abs}(2\sqrt{2} + 4\sin(\frac{1}{2}dx + \frac{1}{2}c)))\operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)) + 180A^2a^2\operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))\sin(\frac{1}{2}dx + \frac{1}{2}c) + 240B^2a^2\operatorname{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))\sin(\frac{1}{2}dx + \frac{1}{2}c))\sqrt[3]{a}/d$

3.94.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x), x)`

3.95 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.95.1	Optimal result	978
3.95.2	Mathematica [A] (verified)	978
3.95.3	Rubi [A] (verified)	979
3.95.4	Maple [B] (verified)	983
3.95.5	Fricas [A] (verification not implemented)	984
3.95.6	Sympy [F(-1)]	984
3.95.7	Maxima [B] (verification not implemented)	984
3.95.8	Giac [A] (verification not implemented)	985
3.95.9	Mupad [F(-1)]	986

3.95.1 Optimal result

Integrand size = 33, antiderivative size = 144

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{a^{5/2}(5A + 2B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} + \frac{a^3(3A + 14B) \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}} - \frac{a^2(3A - 2B)\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{3d} + \frac{aA(a + a \cos(c + dx))^{3/2} \tan(c + dx)}{d}$$

output

```
a^(5/2)*(5*A+2*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+1/3
*a^3*(3*A+14*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-1/3*a^2*(3*A-2*B)*sin(
d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+a*A*(a+a*cos(d*x+c))^(3/2)*tan(d*x+c)/d
```

3.95.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.83

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec(c + dx) (3\sqrt{2}(5A + 2B)\operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)))}{6d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x
]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]*(3*Sqrt[2]*(
5*A + 2*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x] + 2*(3*A + B + 2
*(3*A + 8*B)*Cos[c + d*x] + B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(6*d)`

3.95.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3454, 27, 3042, 3455, 27, 3042, 3460, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3454} \\
 & \int \frac{1}{2}(\cos(c + dx)a + a)^{3/2}(a(5A + 2B) - a(3A - 2B) \cos(c + dx)) \sec(c + dx) dx + \\
 & \quad \frac{aA \tan(c + dx)(a \cos(c + dx) + a)^{3/2}}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int (\cos(c + dx)a + a)^{3/2}(a(5A + 2B) - a(3A - 2B) \cos(c + dx)) \sec(c + dx) dx + \\
 & \quad \frac{aA \tan(c + dx)(a \cos(c + dx) + a)^{3/2}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (a(5A + 2B) - a(3A - 2B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx + \\
 & \quad \frac{aA \tan(c + dx)(a \cos(c + dx) + a)^{3/2}}{d} \\
 & \quad \downarrow \text{3455}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{2}{3} \int \frac{1}{2} \sqrt{\cos(c+dx)a+a} (3(5A+2B)a^2 + (3A+14B)\cos(c+dx)a^2) \sec(c+dx) dx - \frac{2a^2(3A-2B)\sin(c+dx)}{3d} \right) - \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^{3/2}}{d} \downarrow 27$$

$$\frac{1}{2} \left(\frac{1}{3} \int \sqrt{\cos(c+dx)a+a} (3(5A+2B)a^2 + (3A+14B)\cos(c+dx)a^2) \sec(c+dx) dx - \frac{2a^2(3A-2B)\sin(c+dx)}{3d} \right) - \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^{3/2}}{d} \downarrow 3042$$

$$\frac{1}{2} \left(\frac{1}{3} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} (3(5A+2B)a^2 + (3A+14B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})} dx - \frac{2a^2(3A-2B)\sin(c+dx)}{3d} \right) - \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^{3/2}}{d} \downarrow 3460$$

$$\frac{1}{2} \left(\frac{1}{3} \left(3a^2(5A+2B) \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx + \frac{2a^3(3A+14B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \right) - \frac{2a^2(3A-2B)\sin(c+dx)}{3d} \right) - \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^{3/2}}{d} \downarrow 3042$$

$$\frac{1}{2} \left(\frac{1}{3} \left(3a^2(5A+2B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2a^3(3A+14B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \right) - \frac{2a^2(3A-2B)\sin(c+dx)}{3d} \right) - \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^{3/2}}{d} \downarrow 3252$$

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{2a^3(3A+14B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{6a^3(5A+2B) \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} \right) - \frac{2a^2(3A-2B)\sin(c+dx)}{3d} \right) - \frac{aA \tan(c+dx)(a \cos(c+dx) + a)^{3/2}}{d} \downarrow 219$$

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{6a^{5/2}(5A + 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{2a^3(3A + 14B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \right) - \frac{2a^2(3A - 2B) \sin(c+dx) \sqrt{a}}{3d} \right) + \frac{aA \tan(c+dx)(a \cos(c+dx)+a)^{3/2}}{d}$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `((-2*a^2*(3*A - 2*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((6*a^(5/2)*(5*A + 2*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d + (2*a^3*(3*A + 14*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/3)/2 + (a*A*(a + a*Cos[c + d*x])^(3/2)*Tan[c + d*x])/d`

3.95.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3252 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.95.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 682 vs. $2(128) = 256$.

Time = 15.19 (sec) , antiderivative size = 683, normalized size of antiderivative = 4.74

method	result
parts	$A a^{\frac{3}{2}} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-8\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 10 \ln \left(-\frac{4 \left(\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}} \right) \right)$
default	Expression too large to display

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNV
ERBOSE)
```

```
output A*a^(3/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-8*2^(1/2)*(a
*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^2-10*ln(-4/(2*cos(
1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d
*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*sin(1/2*d*x+1/2*c)^2*a-10*ln(4/(2*cos(1/2
*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+
1/2*c)^2)^(1/2)*a^(1/2)+2*a))*sin(1/2*d*x+1/2*c)^2*a+6*2^(1/2)*(a*sin(1/2*
d*x+1/2*c)^2)^(1/2)*a^(1/2)+5*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2
)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a
)*a+5*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(
1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a)/(2*cos(1/2*d*x+1/2*c
)-2^(1/2))/(2*cos(1/2*d*x+1/2*c)+2^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*
x+1/2*c)^2)^(1/2)/d+1/6*B*a^(3/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)
^2)^(1/2)*(-8*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^2+
3*2^(1/2)*ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c
)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+3*2^(1/2)*ln(-2/(
2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin
(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+36*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^
2)^(1/2))/sin(1/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.95.5 Fracas [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.40

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{3 \left((5A + 2B)a^2 \cos(dx + c)^2 + (5A + 2B)a^2 \cos(dx + c) \right) \sqrt{a} \log \left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a} \cos(dx+c) + a}{\cos(dx+c)} \right)}{1}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/12*(3*((5*A + 2*B)*a^2*cos(d*x + c)^2 + (5*A + 2*B)*a^2*cos(d*x + c))*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*(2*B*a^2*cos(d*x + c)^2 + 2*(3*A + 8*B)*a^2*cos(d*x + c) + 3*A*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(d*cos(d*x + c)^2 + d*cos(d*x + c))`

3.95.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Timed out`

3.95.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8114 vs. 2(128) = 256.

Time = 0.65 (sec) , antiderivative size = 8114, normalized size of antiderivative = 56.35

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `-1/252*(1449*sqrt(2)*a^2*cos(5/2*d*x + 5/2*c)^3*sin(2*d*x + 2*c) - 1260*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c)^3 - 1449*(sqrt(2)*a^2*cos(2*d*x + 2*c) + sqrt(2)*a^2)*sin(5/2*d*x + 5/2*c)^3 + 21*(25*sqrt(2)*a^2*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 25*sqrt(2)*a^2*sin(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) - 60*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) + 5*(5*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) - 12*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*cos(2*d*x + 2*c) + (25*sqrt(2)*a^2*cos(3/2*d*x + 3/2*c) + 198*sqrt(2)*a^2*cos(1/2*d*x + 1/2*c))*sin(2*d*x + 2*c))*cos(5/2*d*x + 5/2*c)^2 - 21*(12*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) - 25*(sqrt(2)*a^2*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*a^2*sin(1/2*d*x + 1/2*c)^2)*sin(3/2*d*x + 3/2*c))*cos(2*d*x + 2*c)^2 + 21*(25*sqrt(2)*a^2*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 25*sqrt(2)*a^2*sin(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 69*sqrt(2)*a^2*cos(5/2*d*x + 5/2*c)*sin(2*d*x + 2*c) - 198*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) + (25*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) - 198*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c))*cos(2*d*x + 2*c) + 5*(5*sqrt(2)*a^2*cos(3/2*d*x + 3/2*c) + 12*sqrt(2)*a^2*cos(1/2*d*x + 1/2*c))*sin(2*d*x + 2*c))*sin(5/2*d*x + 5/2*c)^2 - 21*(12*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) - 25*(sqrt(2)*a^2*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*a^2*sin(1/2*d*x + 1/2*c)^2)*sin(3/2*d*x + 3/2*c))*sin(2*d*x + 2*c)^2 - 35*(sqrt(2)*a^2*cos(5/2*d*x + 5/2*c)^2*sin(2*d*x + 2*c) + 2*sqrt(2)*a^2*cos(5/2*d*x + 5/2*c)*cos(1/2*d*x + 1/2*c)*sin(2*d*x + 2*c) + sqrt(2)*a^2*sin(5/2*d*x + 5/2*c)^2*s...`

3.95.8 Giac [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.45

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$\sqrt{2} \left(16 B a^2 \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 - 24 A a^2 \operatorname{sgn} \left(\cos \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right) \sin \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 72 B a \right)$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `-1/12*sqrt(2)*(16*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 24*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) - 72*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) + 12*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)/(2*sin(1/2*d*x + 1/2*c)^2 - 1) + 3*sqrt(2)*(5*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 2*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))))*sqrt(a)/d`

3.95.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^2} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^2,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^2, x)`

3.96 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.96.1	Optimal result	987
3.96.2	Mathematica [A] (verified)	988
3.96.3	Rubi [A] (verified)	988
3.96.4	Maple [B] (verified)	991
3.96.5	Fricas [A] (verification not implemented)	992
3.96.6	Sympy [F(-1)]	993
3.96.7	Maxima [B] (verification not implemented)	993
3.96.8	Giac [A] (verification not implemented)	994
3.96.9	Mupad [F(-1)]	995

3.96.1 Optimal result

Integrand size = 33, antiderivative size = 156

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{a^{5/2}(19A + 20B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4d} - \frac{a^3(9A - 4B) \sin(c + dx)}{4d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(7A + 4B)\sqrt{a + a \cos(c + dx)} \tan(c + dx)}{4d} + \frac{aA(a + a \cos(c + dx))^{3/2} \sec(c + dx) \tan(c + dx)}{2d}$$

```
output 1/4*a^(5/2)*(19*A+20*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))
/d-1/4*a^3*(9*A-4*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/2*a*A*(a+a*cos(
d*x+c))^(3/2)*sec(d*x+c)*tan(d*x+c)/d+1/4*a^2*(7*A+4*B)*(a+a*cos(d*x+c))^(
1/2)*tan(d*x+c)/d
```

3.96.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.81

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^2(c + dx) (\sqrt{2}(19A + 20B) \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)) + 8C)}{8d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x]])*Sec[(c + d*x)/2]*Sec[c + d*x]^2*(sqrt[2]*(19*A + 20*B)*ArcTanh[sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^2 + 2*((11*A + 4*B)*Cos[c + d*x] + 2*(A + 2*B + 2*B*Cos[2*(c + d*x)]))*Sin[(c + d*x)/2])/ (8*d)`

3.96.3 Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3460, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^3(c + dx) (a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\ & \quad \downarrow \text{3454} \\ & \frac{1}{2} \int \frac{1}{2} (\cos(c + dx) a + a)^{3/2} (a(7A + 4B) - a(A - 4B) \cos(c + dx)) \sec^2(c + dx) dx + \\ & \quad \frac{aA \tan(c + dx) \sec(c + dx) (a \cos(c + dx) + a)^{3/2}}{2d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{4} \int (\cos(c+dx)a+a)^{3/2} (a(7A+4B) - a(A-4B)\cos(c+dx)) \sec^2(c+dx) dx + \frac{aA \tan(c+dx) \sec(c+dx) (a \cos(c+dx) + a)^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (a(7A+4B) - a(A-4B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{aA \tan(c+dx) \sec(c+dx) (a \cos(c+dx) + a)^{3/2}}{2d}$$

↓ 3454

$$\frac{1}{4} \left(\int \frac{1}{2} \sqrt{\cos(c+dx)a+a} (a^2(19A+20B) - a^2(9A-4B)\cos(c+dx)) \sec(c+dx) dx + \frac{a^2(7A+4B)\tan(c+dx)}{d} \right) + \frac{aA \tan(c+dx) \sec(c+dx) (a \cos(c+dx) + a)^{3/2}}{2d}$$

↓ 27

$$\frac{1}{4} \left(\frac{1}{2} \int \sqrt{\cos(c+dx)a+a} (a^2(19A+20B) - a^2(9A-4B)\cos(c+dx)) \sec(c+dx) dx + \frac{a^2(7A+4B)\tan(c+dx)}{d} \right) + \frac{aA \tan(c+dx) \sec(c+dx) (a \cos(c+dx) + a)^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} (a^2(19A+20B) - a^2(9A-4B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{a^2(7A+4B)\tan(c+dx)}{d} \right) + \frac{aA \tan(c+dx) \sec(c+dx) (a \cos(c+dx) + a)^{3/2}}{2d}$$

↓ 3460

$$\frac{1}{4} \left(\frac{1}{2} \left(a^2(19A+20B) \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx - \frac{2a^3(9A-4B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(7A+4B)\tan(c+dx)}{d} \right) + \frac{aA \tan(c+dx) \sec(c+dx) (a \cos(c+dx) + a)^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \left(a^2(19A + 20B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{2a^3(9A - 4B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a^2(7A + 4B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^{3/2}}{2d}$$

↓ 3252

$$\frac{1}{4} \left(\frac{1}{2} \left(-\frac{2a^3(19A + 20B) \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}} \right) - \frac{2a^3(9A - 4B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a^2(7A + 4B) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^{3/2}}{2d}$$

↓ 219

$$\frac{1}{4} \left(\frac{a^2(7A + 4B) \tan(c + dx) \sqrt{a \cos(c + dx) + a}}{d} + \frac{1}{2} \left(\frac{2a^{5/2}(19A + 20B) \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} - \frac{2a^3(9A - 4B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) \right) + \frac{aA \tan(c + dx) \sec(c + dx)(a \cos(c + dx) + a)^{3/2}}{2d}$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*(a + a*Cos[c + d*x])^(3/2)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (((2*a^(5/2)*(19*A + 20*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d - (2*a^3*(9*A - 4*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/2 + (a^2*(7*A + 4*B)*Sqrt[a + a*Cos[c + d*x])*Tan[c + d*x])/d)/4`

3.96.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.96.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 994 vs. $2(136) = 272$.

Time = 54.75 (sec) , antiderivative size = 995, normalized size of antiderivative = 6.38

method	result	size
parts	Expression too large to display	995
default	Expression too large to display	1028

output $1/16*((19*A + 20*B)*a^2*\cos(d*x + c)^3 + (19*A + 20*B)*a^2*\cos(d*x + c)^2)*\sqrt{a}*\log((a*\cos(d*x + c)^3 - 7*a*\cos(d*x + c)^2 - 4*\sqrt{a*\cos(d*x + c) + a})*\sqrt{a}*(\cos(d*x + c) - 2)*\sin(d*x + c) + 8*a)/(\cos(d*x + c)^3 + \cos(d*x + c)^2)) + 4*(8*B*a^2*\cos(d*x + c)^2 + (11*A + 4*B)*a^2*\cos(d*x + c) + 2*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c))/(d*\cos(d*x + c)^3 + d*\cos(d*x + c)^2)$

3.96.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Timed out`

3.96.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11782 vs. $2(136) = 272$.

Time = 3.42 (sec) , antiderivative size = 11782, normalized size of antiderivative = 75.53

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output

```

-1/1008*(63*(150*sqrt(2)*a^2*cos(7/2*d*x + 7/2*c)*sin(2*d*x + 2*c) + 154*sqrt(2)*a^2*cos(5/2*d*x + 5/2*c)*sin(2*d*x + 2*c) - 28*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 44*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) - (3*sqrt(2)*a^2*sin(7/2*d*x + 7/2*c) + 5*sqrt(2)*a^2*sin(5/2*d*x + 5/2*c) - 17*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) - 55*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) + 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2))*cos(4*d*x + 4*c)^2 + 4*(17*sqrt(2)*a^2*sin(3/2*d*x + 3/2*c) + 55*sqrt(2)*a^2*sin(1/2*d*x + 1/2*c) - 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + 19*a^2*log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)...

```

3.96.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.53

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{\sqrt{2} \left(32 B a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) \sin(\frac{1}{2} dx + \frac{1}{2} c) - \sqrt{2} (19 A a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 20 B a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \right)}{\dots}$$

input

```

integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")

```

output $\frac{1}{16}\sqrt{2}*(32*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c) - \sqrt{2}*(19*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c)) + 20*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c)))*\log(\text{abs}(-2*\sqrt{2} + 4*\sin(1/2*d*x + 1/2*c))/\text{abs}(2*\sqrt{2} + 4*\sin(1/2*d*x + 1/2*c))) - 4*(22*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)^3 + 8*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)^3 - 13*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c) - 4*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c))/(2*\sin(1/2*d*x + 1/2*c)^2 - 1)^2)*\sqrt{a})/d$

3.96.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^3} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^3,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^3, x)`

3.97 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.97.1	Optimal result	996
3.97.2	Mathematica [A] (verified)	997
3.97.3	Rubi [A] (verified)	997
3.97.4	Maple [B] (verified)	1000
3.97.5	Fricas [A] (verification not implemented)	1001
3.97.6	Sympy [F(-1)]	1002
3.97.7	Maxima [B] (verification not implemented)	1002
3.97.8	Giac [A] (verification not implemented)	1003
3.97.9	Mupad [F(-1)]	1004

3.97.1 Optimal result

Integrand size = 33, antiderivative size = 164

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{a^{5/2}(25A + 38B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{8d} + \frac{a^3(49A + 54B) \tan(c + dx)}{24d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(3A + 2B)\sqrt{a + a \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{4d} + \frac{aA(a + a \cos(c + dx))^{3/2} \sec^2(c + dx) \tan(c + dx)}{3d}$$

```
output 1/8*a^(5/2)*(25*A+38*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))
/d+1/3*a*A*(a+a*cos(d*x+c))^(3/2)*sec(d*x+c)^2*tan(d*x+c)/d+1/24*a^3*(49*A
+54*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/4*a^2*(3*A+2*B)*sec(d*x+c)*(a
+a*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.97.2 Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.80

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^3(c + dx) (3\sqrt{2}(25A + 38B) \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)) + (91A + 66B + 4(17A + 6B)\cos(c + dx) + (75A + 66B)\cos[2(c + dx)]) \sin\left(\frac{c + dx}{2}\right))}{48d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^3*(3*Sqrt[2]*(25*A + 38*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^3 + (91*A + 66*B + 4*(17*A + 6*B)*Cos[c + d*x] + (75*A + 66*B)*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.97.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^4(c + dx) (a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx \\ & \quad \downarrow \text{3454} \\ & \frac{1}{3} \int \frac{1}{2} (\cos(c + dx)a + a)^{3/2} (3a(3A + 2B) + a(A + 6B) \cos(c + dx)) \sec^3(c + dx) dx + \\ & \quad \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^{3/2}}{3d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{6} \int (\cos(c+dx)a+a)^{3/2} (3a(3A+2B) + a(A+6B)\cos(c+dx)) \sec^3(c+dx) dx + \frac{aA \tan(c+dx) \sec^2(c+dx) (a \cos(c+dx) + a)^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{6} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (3a(3A+2B) + a(A+6B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^3} dx + \frac{aA \tan(c+dx) \sec^2(c+dx) (a \cos(c+dx) + a)^{3/2}}{3d}$$

↓ 3454

$$\frac{1}{6} \left(\frac{1}{2} \int \frac{1}{2} \sqrt{\cos(c+dx)a+a} ((49A+54B)a^2 + (13A+30B)\cos(c+dx)a^2) \sec^2(c+dx) dx + \frac{3a^2(3A+2B)\tan(c+dx)}{3d} \right) + \frac{aA \tan(c+dx) \sec^2(c+dx) (a \cos(c+dx) + a)^{3/2}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{1}{4} \int \sqrt{\cos(c+dx)a+a} ((49A+54B)a^2 + (13A+30B)\cos(c+dx)a^2) \sec^2(c+dx) dx + \frac{3a^2(3A+2B)\tan(c+dx)}{3d} \right) + \frac{aA \tan(c+dx) \sec^2(c+dx) (a \cos(c+dx) + a)^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} ((49A+54B)a^2 + (13A+30B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{3a^2(3A+2B)\tan(c+dx)}{3d} \right) + \frac{aA \tan(c+dx) \sec^2(c+dx) (a \cos(c+dx) + a)^{3/2}}{3d}$$

↓ 3459

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3}{2} a^2 (25A+38B) \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx + \frac{a^3(49A+54B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{3a^2(3A+2B)\tan(c+dx)}{3d} \right) + \frac{aA \tan(c+dx) \sec^2(c+dx) (a \cos(c+dx) + a)^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3}{2} a^2 (25A + 38B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a^3 (49A + 54B) \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{3a^2 (3A + 2B) \tan(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^{3/2}}{3d}$$

↓ 3252

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{a^3 (49A + 54B) \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} - \frac{3a^3 (25A + 38B) \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx) a + a}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right)}{d} \right) + \frac{3a^2 (3A + 2B) \tan(c + dx)}{2d} \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^{3/2}}{3d}$$

↓ 219

$$\frac{1}{6} \left(\frac{3a^2 (3A + 2B) \tan(c + dx) \sec(c + dx) \sqrt{a \cos(c + dx) + a}}{2d} + \frac{1}{4} \left(\frac{3a^{5/2} (25A + 38B) \operatorname{arctanh} \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} + \frac{a^3 (49A + 54B) \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) \right) + \frac{aA \tan(c + dx) \sec^2(c + dx) (a \cos(c + dx) + a)^{3/2}}{3d}$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*(a + a*Cos[c + d*x])^(3/2)*Sec[c + d*x]^2*Tan[c + d*x]/(3*d) + ((3*a^2*(3*A + 2*B)*Sqrt[a + a*Cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((3*a^(5/2)*(25*A + 38*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (a^3*(49*A + 54*B)*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4)/6`

3.97.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.97. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.97.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1281 vs. $2(144) = 288$.

Time = 163.58 (sec) , antiderivative size = 1282, normalized size of antiderivative = 7.82

method	result	size
parts	Expression too large to display	1282
default	Expression too large to display	1326

3.97. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNV
ERBOSE)`

output `1/6*A*a^(3/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-600*a*(1
n(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(
1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)*a^(1/2)-2*a)))*sin(1/2*d*x+1/2*c)^6+300*(2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)
^2)^(1/2)*a^(1/2)+3*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2
*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+3*ln(-4
/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*s
in(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^4+(-736*2^(
1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-450*ln(-4/(2*cos(1/2*d*x+1/2*c
)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)
(1/2)*a^(1/2)-2*a))*a-450*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*c
os(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a)*
sin(1/2*d*x+1/2*c)^2+234*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+75
*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+75*ln(4/(2*cos(1/2*d*x+1/
2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)
^2)^(1/2)*a^(1/2)+2*a))*a)/(2*cos(1/2*d*x+1/2*c)-2^(1/2))^3/(2*cos(1/2*d*x+
1/2*c)+2^(1/2))^3/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+1/4*A
B*a^(3/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(76*2^(1/2)...`

3.97.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.29

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{3 \left((25A + 38B)a^2 \cos^4(dx + c) + (25A + 38B)a^2 \cos^3(dx + c) \right) \sqrt{a} \log \left(\frac{a \cos(dx + c)^3 - 7a \cos(dx + c)^2 - c}{c} \right)}{c}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorith
m="fricas")`

output $1/96*(3*((25*A + 38*B)*a^2*\cos(d*x + c)^4 + (25*A + 38*B)*a^2*\cos(d*x + c)^3)*\sqrt{a}*\log((a*\cos(d*x + c)^3 - 7*a*\cos(d*x + c)^2 - 4*\sqrt{a*\cos(d*x + c) + a})*\sqrt{a}*(\cos(d*x + c) - 2)*\sin(d*x + c) + 8*a)/(\cos(d*x + c)^3 + \cos(d*x + c)^2)) + 4*(3*(25*A + 22*B)*a^2*\cos(d*x + c)^2 + 2*(17*A + 6*B)*a^2*\cos(d*x + c) + 8*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c))/(d*\cos(d*x + c)^4 + d*\cos(d*x + c)^3)$

3.97.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.97.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7994 vs. $2(144) = 288$.

Time = 3.62 (sec) , antiderivative size = 7994, normalized size of antiderivative = 48.74

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output

```

-1/96*((1530*a^2*cos(4*d*x + 4*c)^2*sin(3/2*d*x + 3/2*c) + 1530*a^2*cos(2*
d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 1530*a^2*sin(4*d*x + 4*c)^2*sin(3/2*d*
x + 3/2*c) + 1530*a^2*sin(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 4176*a^2*c
os(7/2*d*x + 7/2*c)*sin(2*d*x + 2*c) + 2430*a^2*cos(5/2*d*x + 5/2*c)*sin(2
*d*x + 2*c) + 678*a^2*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) + 342*a^2*cos(
2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + 10*(a^2*sin(9/2*d*x + 9/2*c) + 17*a^2*
sin(3/2*d*x + 3/2*c))*cos(6*d*x + 6*c)^2 + 10*(a^2*sin(9/2*d*x + 9/2*c) +
17*a^2*sin(3/2*d*x + 3/2*c))*sin(6*d*x + 6*c)^2 - 56*a^2*sin(3/2*d*x + 3/2
*c) + 10*(a^2*sin(6*d*x + 6*c) + 3*a^2*sin(4*d*x + 4*c) + 3*a^2*sin(2*d*x
+ 2*c))*cos(21/2*d*x + 21/2*c) - 30*(a^2*sin(6*d*x + 6*c) + 3*a^2*sin(4*d*
x + 4*c) + 3*a^2*sin(2*d*x + 2*c))*cos(19/2*d*x + 19/2*c) - 48*(a^2*sin(6*
d*x + 6*c) + 3*a^2*sin(4*d*x + 4*c) + 3*a^2*sin(2*d*x + 2*c))*cos(17/2*d*x
+ 17/2*c) + 80*(a^2*sin(6*d*x + 6*c) + 3*a^2*sin(4*d*x + 4*c) + 3*a^2*sin
(2*d*x + 2*c))*cos(15/2*d*x + 15/2*c) + 396*(a^2*sin(6*d*x + 6*c) + 3*a^2*
sin(4*d*x + 4*c) + 3*a^2*sin(2*d*x + 2*c))*cos(13/2*d*x + 13/2*c) + 6*(170
*a^2*cos(4*d*x + 4*c)*sin(3/2*d*x + 3/2*c) + 170*a^2*cos(2*d*x + 2*c)*sin(
3/2*d*x + 3/2*c) - 170*a^2*sin(11/2*d*x + 11/2*c) - 232*a^2*sin(7/2*d*x +
7/2*c) - 135*a^2*sin(5/2*d*x + 5/2*c) + 19*a^2*sin(3/2*d*x + 3/2*c) + 10*(
a^2*cos(4*d*x + 4*c) + a^2*cos(2*d*x + 2*c) - 25*a^2)*sin(9/2*d*x + 9/2*c)
)*cos(6*d*x + 6*c) + 3060*(a^2*sin(4*d*x + 4*c) + a^2*sin(2*d*x + 2*c))...

```

3.97.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.63

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx =$$

$$\frac{\sqrt{2} \left(3 \sqrt{2} (25 A a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 38 B a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \log \left(\frac{|-2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2} + 4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|} \right) + \frac{4}{30} \right)}{1}$$

input

```

integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorith
m="giac")

```

output
$$\begin{aligned} & -1/96*\sqrt{2}*(3*\sqrt{2}*(25*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c)) + 38*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))))*\log(\text{abs}(-2*\sqrt{2} + 4*\sin(1/2*d*x + 1/2*c))/\text{abs}(2*\sqrt{2} + 4*\sin(1/2*d*x + 1/2*c))) + 4*(300*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)^5 + 264*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)^5 - 368*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)^3 - 288*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c)^3 + 117*A*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c) + 78*B*a^2*\text{sgn}(\cos(1/2*d*x + 1/2*c))*\sin(1/2*d*x + 1/2*c))/(2*\sin(1/2*d*x + 1/2*c)^2 - 1)^3*\sqrt{a}/d \end{aligned}$$

3.97.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^4} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^4,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^4, x)`

3.98 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.98.1	Optimal result	1005
3.98.2	Mathematica [A] (verified)	1006
3.98.3	Rubi [A] (verified)	1006
3.98.4	Maple [B] (verified)	1010
3.98.5	Fricas [A] (verification not implemented)	1011
3.98.6	Sympy [F(-1)]	1012
3.98.7	Maxima [F(-1)]	1012
3.98.8	Giac [A] (verification not implemented)	1012
3.98.9	Mupad [F(-1)]	1013

3.98.1 Optimal result

Integrand size = 33, antiderivative size = 209

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^5(c + dx) dx = \frac{a^{5/2}(163A + 200B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{64d} + \frac{a^3(163A + 200B) \tan(c + dx)}{64d\sqrt{a + a \cos(c + dx)}} + \frac{a^3(95A + 104B) \sec(c + dx) \tan(c + dx)}{96d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(11A + 8B)\sqrt{a + a \cos(c + dx)} \sec^2(c + dx) \tan(c + dx)}{24d} + \frac{aA(a + a \cos(c + dx))^{3/2} \sec^3(c + dx) \tan(c + dx)}{4d}$$

```
output 1/64*a^(5/2)*(163*A+200*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+1/4*a*A*(a+a*cos(d*x+c))^(3/2)*sec(d*x+c)^3*tan(d*x+c)/d+1/64*a^3*(163*A+200*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/96*a^3*(95*A+104*B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/24*a^2*(11*A+8*B)*sec(d*x+c)^2*(a+a*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.98.2 Mathematica [A] (verified)

Time = 1.12 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.73

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^4(c + dx) (6\sqrt{2}(163A + 200B) \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)) + \dots}{768d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^4*(6*sqrt[2]*(163*A + 200*B)*ArcTanh[sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^4 + (844*A + 544*B + (2203*A + 2056*B)*Cos[c + d*x] + (652*A + 544*B)*Cos[2*(c + d*x)]) + 489*A*cos[3*(c + d*x)] + 600*B*cos[3*(c + d*x)])*Sin[(c + d*x)/2])/768*d)`

3.98.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^5(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx \\ & \quad \downarrow \text{3454} \\ & \frac{1}{4} \int \frac{1}{2} (\cos(c + dx)a + a)^{3/2} (a(11A + 8B) + a(3A + 8B) \cos(c + dx)) \sec^4(c + dx) dx + \\ & \quad \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{8} \int (\cos(c+dx)a+a)^{3/2} (a(11A+8B) + a(3A+8B)\cos(c+dx)) \sec^4(c+dx) dx + \frac{aA \tan(c+dx) \sec^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (a(11A+8B) + a(3A+8B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^4} dx + \frac{aA \tan(c+dx) \sec^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{4d}$$

↓ 3454

$$\frac{1}{8} \left(\frac{1}{3} \int \frac{1}{2} \sqrt{\cos(c+dx)a+a} ((95A+104B)a^2 + 3(17A+24B)\cos(c+dx)a^2) \sec^3(c+dx) dx + \frac{a^2(11A+8B)\tan(c+dx)}{4d} \right) + \frac{aA \tan(c+dx) \sec^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \int \sqrt{\cos(c+dx)a+a} ((95A+104B)a^2 + 3(17A+24B)\cos(c+dx)a^2) \sec^3(c+dx) dx + \frac{a^2(11A+8B)\tan(c+dx)}{4d} \right) + \frac{aA \tan(c+dx) \sec^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} ((95A+104B)a^2 + 3(17A+24B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^3} dx + \frac{a^2(11A+8B)\tan(c+dx)}{4d} \right) + \frac{aA \tan(c+dx) \sec^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{4d}$$

↓ 3459

$$\frac{1}{8} \left(\frac{1}{4} \left(\frac{3}{4} a^2 (163A + 200B) \int \sqrt{\cos(c+dx)a+a} \sec^2(c+dx) dx + \frac{a^3(95A+104B)\tan(c+dx)\sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(11A+8B)\tan(c+dx)}{4d} \right) + \frac{aA \tan(c+dx) \sec^3(c+dx) (a \cos(c+dx) + a)^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (163A + 200B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{a^3 (95A + 104B) \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) + \frac{a^2 (11A + 8B) \tan(c + dx) \sec^2(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d}$$

↓ 3251

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (163A + 200B) \left(\frac{1}{2} \int \sqrt{\cos(c + dx) a + a} \sec(c + dx) dx + \frac{a \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{a^3 (95A + 104B) \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (163A + 200B) \left(\frac{1}{2} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{a \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{a^3 (95A + 104B) \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right)$$

↓ 3252

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (163A + 200B) \left(\frac{a \tan(c + dx)}{d \sqrt{a \cos(c + dx) + a}} - \frac{a \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx) a + a}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right)}{d} \right) + \frac{a^3 (95A + 104B) \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right)$$

↓ 219

$$\frac{1}{8} \left(\frac{a^2 (11A + 8B) \tan(c + dx) \sec^2(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} + \frac{1}{6} \left(\frac{a^3 (95A + 104B) \tan(c + dx) \sec(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} + \frac{aA \tan(c + dx) \sec^3(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

```
output (a*A*(a + a*cos[c + d*x])^(3/2)*Sec[c + d*x]^3*Tan[c + d*x]/(4*d) + ((a^2
*(11*A + 8*B)*Sqrt[a + a*cos[c + d*x]]*Sec[c + d*x]^2*Tan[c + d*x])/(3*d)
+ ((a^3*(95*A + 104*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d*Sqrt[a + a*cos[c +
d*x]])) + (3*a^2*(163*A + 200*B)*((Sqrt[a]*ArcTanh[(Sqrt[a]*Sin[c + d*x])/S
qrt[a + a*cos[c + d*x]]])/d + (a*Tan[c + d*x])/(d*Sqrt[a + a*cos[c + d*x]]
))) / 4) / 6) / 8
```

3.98.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3251 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (
f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*SIN[e
+ f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*SIN[e + f*x]])), x] + Sim
p[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*SIN[e
+ f*x]]*(c + d*SIN[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3252 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (
f_.)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2),
x], x, b*(Cos[e + f*x]/Sqrt[a + b*SIN[e + f*x])], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```



```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.98.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1649 vs. $2(185) = 370$.

Time = 2.42 (sec) , antiderivative size = 1650, normalized size of antiderivative = 7.89

Expression too large to display

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x)
```

```
output 1/24*a^(3/2)*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(48*a*(163*
A*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/
2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))+163*A*ln(4/(2*cos(1/2*d*x+
1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c
)^2)^(1/2)*a^(1/2)+2*a))+200*B*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/
2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a
))+200*B*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)
+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a)))*sin(1/2*d*x+1/2*c)^
8-48*(163*A*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+200*B*2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+326*A*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^
(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)*a^(1/2)-2*a))*a+326*A*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos
(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+400
*B*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1
/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+400*B*ln(4/(2*cos(1/2*d
*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/
2*c)^2)^(1/2)*a^(1/2)+2*a))*a)*sin(1/2*d*x+1/2*c)^6+8*(1793*A*a^(1/2)*2^(1
/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2072*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2
)^(1/2)*a^(1/2)+1467*A*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(
1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a+1...
```

3.98.5 Fracas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.11

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \frac{3 \left((163A + 200B)a^2 \cos(dx + c)^5 + (163A + 200B)a^2 \cos(dx + c)^4 \right) \sqrt{a} \log \left(\frac{a \cos(dx + c)^3 - 7a \cos(dx + c)^2 - 4\sqrt{a} \cos(dx + c) + a}{\cos(dx + c)^3 + \cos(dx + c)^2} \right) + 4 \left(3(163A + 200B)a^2 \cos(dx + c)^3 + 2(163A + 136B)a^2 \cos(dx + c)^2 + 8(23A + 8B)a^2 \cos(dx + c) + 48Aa^2 \right) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{(d \cos(dx + c))^5 + d \cos(dx + c)^4}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorith
m="fracas")
```

```
output 1/768*(3*((163*A + 200*B)*a^2*cos(d*x + c)^5 + (163*A + 200*B)*a^2*cos(d*x
+ c)^4)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos
(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c
)^3 + cos(d*x + c)^2)) + 4*(3*(163*A + 200*B)*a^2*cos(d*x + c)^3 + 2*(163*
A + 136*B)*a^2*cos(d*x + c)^2 + 8*(23*A + 8*B)*a^2*cos(d*x + c) + 48*A*a^2
)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(d*cos(d*x + c)^5 + d*cos(d*x + c
)^4)
```

3.98. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.98.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`output `Timed out`**3.98.7 Maxima [F(-1)]**

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm m="maxima")`output `Timed out`**3.98.8 Giac [A] (verification not implemented)**

Time = 0.73 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.54

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx =$$

$$\frac{\sqrt{2} \left(3 \sqrt{2} (163 A a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c)) + 200 B a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \log \left(\frac{|-2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|} \right) + \frac{4}{\dots} \right)}{\dots}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm m="giac")`

3.98. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx$

output `-1/768*sqrt(2)*(3*sqrt(2)*(163*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 200*B*a^2*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c)))/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c)) + 4*(3912*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^7 + 4800*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^7 - 7172*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 - 8288*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 + 4606*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 4816*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 1047*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c) - 936*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^4)*sqrt(a)/d`

3.98.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^5} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^5,x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^5, x)`

3.99 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^6(c+dx) dx$

3.99.1	Optimal result	1014
3.99.2	Mathematica [A] (verified)	1015
3.99.3	Rubi [A] (verified)	1015
3.99.4	Maple [B] (verified)	1020
3.99.5	Fricas [A] (verification not implemented)	1020
3.99.6	Sympy [F(-1)]	1021
3.99.7	Maxima [F(-1)]	1021
3.99.8	Giac [A] (verification not implemented)	1022
3.99.9	Mupad [F(-1)]	1022

3.99.1 Optimal result

Integrand size = 33, antiderivative size = 254

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c + dx) dx = \frac{a^{5/2}(283A + 326B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{128d} + \frac{a^3(283A + 326B) \tan(c + dx)}{128d\sqrt{a + a \cos(c + dx)}} + \frac{a^3(283A + 326B) \sec(c + dx) \tan(c + dx)}{192d\sqrt{a + a \cos(c + dx)}} + \frac{a^3(157A + 170B) \sec^2(c + dx) \tan(c + dx)}{240d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(13A + 10B) \sqrt{a + a \cos(c + dx)} \sec^3(c + dx) \tan(c + dx)}{40d} + \frac{aA(a + a \cos(c + dx))^{3/2} \sec^4(c + dx) \tan(c + dx)}{5d}$$

output $\frac{1}{128}a^{5/2}(283A+326B)\operatorname{arctanh}\left(\frac{\sin(d*x+c)*a^{1/2}}{(a+a*\cos(d*x+c))^{1/2}}\right)/d+1/5*a*A*(a+a*\cos(d*x+c))^{3/2}*\sec(d*x+c)^4*\tan(d*x+c)/d+1/128*a^3*(283A+326B)*\tan(d*x+c)/d/(a+a*\cos(d*x+c))^{1/2}+1/192*a^3*(283A+326B)*\sec(d*x+c)*\tan(d*x+c)/d/(a+a*\cos(d*x+c))^{1/2}+1/240*a^3*(157A+170B)*\sec(d*x+c)^2*\tan(d*x+c)/d/(a+a*\cos(d*x+c))^{1/2}+1/40*a^2*(13A+10B)*\sec(d*x+c)^3*(a+a*\cos(d*x+c))^{1/2}*\tan(d*x+c)/d$

3.99.2 Mathematica [A] (verified)

Time = 1.42 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.69

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^5(c + dx) (60\sqrt{2}(283A + 326B) \operatorname{arctanh}(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)) + (24863A + 22030B + 36(781A + 650B) \cos(c + dx) + 4(6509A + 6730B) \cos(2(c + dx)) + 5660A \cos(3(c + dx)) + 6520B \cos(3(c + dx)) + 4245A \cos(4(c + dx)) + 4890B \cos(4(c + dx))) \sin\left(\frac{c + dx}{2}\right))}{15360d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^5*(60*sqrt[2]*(283*A + 326*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^5 + (24863*A + 22030*B + 36*(781*A + 650*B)*Cos[c + d*x] + 4*(6509*A + 6730*B)*Cos[2*(c + d*x)] + 5660*A*cos[3*(c + d*x)] + 6520*B*cos[3*(c + d*x)] + 4245*A*cos[4*(c + d*x)] + 4890*B*cos[4*(c + d*x)])*Sin[(c + d*x)/2])/15360*d)`

3.99.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^6(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^6} dx \\ & \quad \downarrow \text{3454} \\ & \frac{1}{5} \int \frac{1}{2} (\cos(c + dx)a + a)^{3/2} (a(13A + 10B) + 5a(A + 2B) \cos(c + dx)) \sec^5(c + dx) dx + \\ & \quad \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{10} \int (\cos(c+dx)a+a)^{3/2} (a(13A+10B) + 5a(A+2B)\cos(c+dx)) \sec^5(c+dx) dx + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{10} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (a(13A+10B) + 5a(A+2B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^5} dx + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 3454

$$\frac{1}{10} \left(\frac{1}{4} \int \frac{1}{2} \sqrt{\cos(c+dx)a+a} ((157A+170B)a^2 + 5(21A+26B)\cos(c+dx)a^2) \sec^4(c+dx) dx + \frac{a^2(13A+10B)}{5d} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 27

$$\frac{1}{10} \left(\frac{1}{8} \int \sqrt{\cos(c+dx)a+a} ((157A+170B)a^2 + 5(21A+26B)\cos(c+dx)a^2) \sec^4(c+dx) dx + \frac{a^2(13A+10B)}{5d} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{10} \left(\frac{1}{8} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} ((157A+170B)a^2 + 5(21A+26B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^4} dx + \frac{a^2(13A+10B)}{5d} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 3459

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \int \sqrt{\cos(c+dx)a+a} \sec^3(c+dx) dx + \frac{a^3(157A+170B) \tan(c+dx) \sec^2(c+dx)}{3d \sqrt{a \cos(c+dx) + a}} \right) + \frac{a^2(13A+10B)}{5d} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^3} dx + \frac{a^3(157A+170B)\tan(c+dx)\sec^2(c+dx)}{3d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(157A+170B)}{3d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d}$$

↓ 3251

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \left(\frac{3}{4} \int \sqrt{\cos(c+dx)a+a} \sec^2(c+dx) dx + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^3(157A+170B)}{3d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d} \right)$$

↓ 3042

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \left(\frac{3}{4} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^2} dx + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^3(157A+170B)}{3d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d} \right)$$

↓ 3251

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \left(\frac{3}{4} \left(\frac{1}{2} \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx + \frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d} \right)$$

↓ 3042

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d} \right)$$

↓ 3252

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (283A + 326B) \left(\frac{3}{4} \left(\frac{a \tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \right) + \frac{a \tan(c+dx) \sec(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aA \tan(c+dx) \sec^4(c+dx) (a \cos(c+dx) + a)^{3/2}}{5d} \right)$$

↓ 219

$$\frac{1}{10} \left(\frac{a^2(13A + 10B) \tan(c + dx) \sec^3(c + dx) \sqrt{a \cos(c + dx) + a}}{4d} + \frac{1}{8} \left(\frac{a^3(157A + 170B) \tan(c + dx) \sec^2(c + dx)}{3d \sqrt{a \cos(c + dx) + a}} + \frac{aA \tan(c + dx) \sec^4(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a*A*(a + a*Cos[c + d*x])^(3/2)*Sec[c + d*x]^4*Tan[c + d*x]/(5*d) + ((a^2*(13*A + 10*B)*Sqrt[a + a*Cos[c + d*x])*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((a^3*(157*A + 170*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]])) + (5*a^2*(283*A + 326*B)*((a*Sec[c + d*x]*Tan[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]])) + (3*((Sqrt[a]*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (a*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))) /4)/6)/8)/10`

3.99.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

output $1/7680*(15*((283*A + 326*B)*a^2*\cos(d*x + c)^6 + (283*A + 326*B)*a^2*\cos(d*x + c)^5)*\sqrt{a}*\log((a*\cos(d*x + c)^3 - 7*a*\cos(d*x + c)^2 - 4*\sqrt{a*\cos(d*x + c) + a}*\sqrt{a}*(\cos(d*x + c) - 2)*\sin(d*x + c) + 8*a)/(\cos(d*x + c)^3 + \cos(d*x + c)^2)) + 4*(15*(283*A + 326*B)*a^2*\cos(d*x + c)^4 + 10*(283*A + 326*B)*a^2*\cos(d*x + c)^3 + 8*(283*A + 230*B)*a^2*\cos(d*x + c)^2 + 48*(29*A + 10*B)*a^2*\cos(d*x + c) + 384*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c))/(d*\cos(d*x + c)^6 + d*\cos(d*x + c)^5)$

3.99.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**6,x)`

output Timed out

3.99.7 Maxima [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="maxima")`

output Timed out

3.99.8 Giac [A] (verification not implemented)

Time = 0.79 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.48

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c + dx) dx =$$

$$\sqrt{2} \left(15 \sqrt{2} (283 A a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) + 326 B a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))) \log \left(\frac{|-2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|} \right) \right) +$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm
m="giac")
```

```
output -1/7680*sqrt(2)*(15*sqrt(2)*(283*A*a^2*sgn(cos(1/2*d*x + 1/2*c)) + 326*B*a
^2*sgn(cos(1/2*d*x + 1/2*c)))*log(abs(-2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))
/abs(2*sqrt(2) + 4*sin(1/2*d*x + 1/2*c))) + 4*(67920*A*a^2*sgn(cos(1/2*d*x
+ 1/2*c))*sin(1/2*d*x + 1/2*c)^9 + 78240*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*
sin(1/2*d*x + 1/2*c)^9 - 158480*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*
x + 1/2*c)^7 - 182560*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)
^7 + 144896*A*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 + 16384
0*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^5 - 62780*A*a^2*sgn
(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 - 67000*B*a^2*sgn(cos(1/2*d*
x + 1/2*c))*sin(1/2*d*x + 1/2*c)^3 + 11115*A*a^2*sgn(cos(1/2*d*x + 1/2*c))
*sin(1/2*d*x + 1/2*c) + 10470*B*a^2*sgn(cos(1/2*d*x + 1/2*c))*sin(1/2*d*x
+ 1/2*c))/(2*sin(1/2*d*x + 1/2*c)^2 - 1)^5)*sqrt(a)/d
```

3.99.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c$$

$$+ dx) dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^6} dx$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^6,x)
```

```
output int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^6, x)
```

3.99. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^6(c + dx) dx$

3.100
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$$

3.100.1 Optimal result 1023
 3.100.2 Mathematica [A] (verified) 1024
 3.100.3 Rubi [A] (verified) 1024
 3.100.4 Maple [A] (verified) 1029
 3.100.5 Fricas [A] (verification not implemented) 1029
 3.100.6 Sympy [F(-1)] 1030
 3.100.7 Maxima [B] (verification not implemented) 1030
 3.100.8 Giac [A] (verification not implemented) 1031
 3.100.9 Mupad [F(-1)] 1032

3.100.1 Optimal result

Integrand size = 33, antiderivative size = 202

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx = -\frac{\sqrt{2}(A-B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{4(49A-37B) \sin(c+dx)}{105d\sqrt{a+a \cos(c+dx)}} + \frac{2(7A-B) \cos^2(c+dx) \sin(c+dx)}{35d\sqrt{a+a \cos(c+dx)}} + \frac{2B \cos^3(c+dx) \sin(c+dx)}{7d\sqrt{a+a \cos(c+dx)}} - \frac{2(7A-31B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{105ad}$$

```
output -(A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+4/105*(49*A-37*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/35*(7*A-B)*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/7*B*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-2/105*(7*A-31*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a/d
```

3.100.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.55

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c+dx)\right) \left(-420(A-B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) + 2(406A-178B + (-28A+169B)\cos(c+dx))\right)}{210d\sqrt{a(1+\cos(c+dx))}}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(Cos[(c + d*x)/2]*(-420*(A - B)*ArcTanh[Sin[(c + d*x)/2]] + 2*(406*A - 178*B + (-28*A + 169*B)*Cos[c + d*x] + 6*(7*A - B)*Cos[2*(c + d*x)] + 15*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2])/(210*d*Sqrt[a*(1 + Cos[c + d*x])])`

3.100.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.12, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3462, 27, 3042, 3462, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^3 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\sqrt{a\sin\left(c+dx+\frac{\pi}{2}\right)+a}} dx$$

$$\downarrow \text{3462}$$

$$\frac{2 \int \frac{\cos^2(c+dx)(6aB+a(7A-B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{7a} + \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \frac{\int \frac{\cos^2(c+dx)(6aB+a(7A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{7a} + \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(6aB+a(7A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{7a} + \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3462} \\
& \frac{2 \int \frac{\cos(c+dx)(4a^2(7A-B)-a^2(7A-31B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \quad \frac{7a}{7d\sqrt{a \cos(c+dx)+a}} \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cos(c+dx)(4a^2(7A-B)-a^2(7A-31B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4a^2(7A-B)-a^2(7A-31B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \quad \frac{7a}{7d\sqrt{a \cos(c+dx)+a}} \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3447} \\
& \frac{\int \frac{4a^2(7A-B)\cos(c+dx)-a^2(7A-31B)\cos^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{4a^2(7A-B)\sin(c+dx+\frac{\pi}{2})-a^2(7A-31B)\sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \quad \frac{7a}{7d\sqrt{a \cos(c+dx)+a}} \frac{2B \sin(c+dx) \cos^3(c+dx)}{7d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3502}
\end{aligned}$$

3.100. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

$$\frac{2 \int \frac{a^3(7A-31B)-2a^3(49A-37B)\cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx - \frac{2a(7A-31B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{7a}{2B\sin(c+dx)\cos^3(c+dx)} \frac{7a}{7d\sqrt{a\cos(c+dx)+a}}$$

27

$$\frac{\int \frac{a^3(7A-31B)-2a^3(49A-37B)\cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx - \frac{2a(7A-31B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{7a}{2B\sin(c+dx)\cos^3(c+dx)} \frac{7a}{7d\sqrt{a\cos(c+dx)+a}}$$

3042

$$\frac{\int \frac{a^3(7A-31B)-2a^3(49A-37B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2a(7A-31B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{7a}{2B\sin(c+dx)\cos^3(c+dx)} \frac{7a}{7d\sqrt{a\cos(c+dx)+a}}$$

3230

$$\frac{105a^3(A-B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^3(49A-37B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-31B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{7a}{2B\sin(c+dx)\cos^3(c+dx)} \frac{7a}{7d\sqrt{a\cos(c+dx)+a}}$$

3042

$$\frac{105a^3(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^3(49A-37B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-31B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}}{5a} + \frac{2a(7A-B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{7a}{2B\sin(c+dx)\cos^3(c+dx)} \frac{7a}{7d\sqrt{a\cos(c+dx)+a}}$$

3128

3.100. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

$$\begin{aligned}
& \frac{210a^3(A-B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{3a} - \frac{4a^3(49A-37B) \sin(c+dx)}{5a d \sqrt{a \cos(c+dx)+a}} - \frac{2a(7A-31B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2a(7A-B) \sin(c+dx)}{5d \sqrt{a \cos(c+dx)+a}} \\
& \frac{2B \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{7d \sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{219} \\
& \frac{105\sqrt{2}a^{5/2}(A-B) \arctanh\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{3a d} - \frac{4a^3(49A-37B) \sin(c+dx)}{5a d \sqrt{a \cos(c+dx)+a}} - \frac{2a(7A-31B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2a(7A-B) \sin(c+dx) \cos^2(c+dx)}{5d \sqrt{a \cos(c+dx)+a}} \\
& \frac{2B \sin(c+dx) \cos^3(c+dx) \sqrt{a \cos(c+dx)+a}}{7d \sqrt{a \cos(c+dx)+a}}
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*B*Cos[c + d*x]^3*Sin[c + d*x])/(7*d*Sqrt[a + a*Cos[c + d*x]]) + ((2*a*(7*A - B)*Cos[c + d*x]^2*Sin[c + d*x])/(5*d*Sqrt[a + a*Cos[c + d*x]]) + ((-2*a*(7*A - 31*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) - ((105*Sqrt[2]*a^(5/2)*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d - (4*a^3*(49*A - 37*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/(3*a)/(5*a))/(7*a)`

3.100.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3462 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]`

3.100.4 Maple [A] (verified)

Time = 4.21 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.39

method	result
default	$\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-240B\sqrt{a} \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 168\sqrt{a} \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} (A+2B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right)$
parts	$A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(24 \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 20 \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 30 \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right) + 15a^{\frac{3}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d$

input `int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNV ERBOSE)`

output `1/105*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-240*B*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^6+168*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(A+2*B)*sin(1/2*d*x+1/2*c)^4-140*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(A+2*B)*sin(1/2*d*x+1/2*c)^2-105*2^(1/2)*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a*A+105*2^(1/2)*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a*B+210*A*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)/a^(3/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.100.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.91

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{4(15B\cos(dx+c)^3 + 3(7A-B)\cos(dx+c)^2 - (7A-31B)\cos(dx+c) + 91A - 43B)\sqrt{a\cos(dx+c)}}{210(ad\cos(dx+c))}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x,algorithm="fracas")`

output `1/210*(4*(15*B*cos(d*x + c)^3 + 3*(7*A - B)*cos(d*x + c)^2 - (7*A - 31*B)*cos(d*x + c) + 91*A - 43*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c) - 105*sqrt(2)*((A - B)*a*cos(d*x + c) + (A - B)*a)*log(-(cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(a) - 2*cos(d*x + c) - 3)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1))/sqrt(a))/(a*d*cos(d*x + c) + a*d)`

3.100.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.100.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1604723 vs. 2(177) = 354.

Time = 32.55 (sec) , antiderivative size = 1604723, normalized size of antiderivative = 7944.17

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `-1/5040*(3*(84*(sqrt(2)*cos(3/2*d*x + 3/2*c)^2*sin(d*x + c) + 2*sqrt(2)*cos(3/2*d*x + 3/2*c)*cos(1/2*d*x + 1/2*c)*sin(d*x + c) + sqrt(2)*sin(3/2*d*x + 3/2*c)^2*sin(d*x + c) + 2*sqrt(2)*sin(3/2*d*x + 3/2*c)*sin(d*x + c)*sin(1/2*d*x + 1/2*c) + (sqrt(2)*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*sin(1/2*d*x + 1/2*c)^2)*sin(d*x + c))*cos(7/2*d*x + 7/2*c)^3 - 84*((sqrt(2)*cos(d*x + c) + sqrt(2))*cos(3/2*d*x + 3/2*c)^2 + (sqrt(2)*cos(d*x + c) + sqrt(2))*sin(3/2*d*x + 3/2*c)^2 + sqrt(2)*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*sin(1/2*d*x + 1/2*c)^2 + 2*(sqrt(2)*cos(d*x + c)*cos(1/2*d*x + 1/2*c) + sqrt(2)*cos(1/2*d*x + 1/2*c))*cos(3/2*d*x + 3/2*c) + (sqrt(2)*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*sin(1/2*d*x + 1/2*c)^2)*cos(d*x + c) + 2*(sqrt(2)*cos(d*x + c)*sin(1/2*d*x + 1/2*c) + sqrt(2)*sin(1/2*d*x + 1/2*c))*sin(3/2*d*x + 3/2*c))*sin(7/2*d*x + 7/2*c)^3 - 24*((sqrt(2)*cos(d*x + c)^2 + sqrt(2)*sin(d*x + c)^2 + 2*sqrt(2)*cos(d*x + c) + sqrt(2))*cos(3/2*d*x + 3/2*c)^2 + (sqrt(2)*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*sin(1/2*d*x + 1/2*c)^2)*cos(d*x + c)^2 + (sqrt(2)*cos(d*x + c)^2 + sqrt(2)*sin(d*x + c)^2 + 2*sqrt(2)*cos(d*x + c) + sqrt(2))*sin(3/2*d*x + 3/2*c)^2 + (sqrt(2)*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*sin(1/2*d*x + 1/2*c)^2)*sin(d*x + c)^2 + sqrt(2)*cos(1/2*d*x + 1/2*c)^2 + sqrt(2)*sin(1/2*d*x + 1/2*c)^2 + 2*(sqrt(2)*cos(d*x + c)^2*cos(1/2*d*x + 1/2*c) + sqrt(2)*cos(1/2*d*x + 1/2*c)*sin(d*x + c)^2 + 2*sqrt(2)*cos(d*x + c)*cos(1/2*d*x + 1/2*c) + sqrt(2)*cos(1/2*d*x + 1/2*c))*cos(3/2*d*x ...`

3.100.8 Giac [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.07

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx = \frac{105 \sqrt{2}(A\sqrt{a} - B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{105 \sqrt{2}(A\sqrt{a} - B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \frac{4 \sqrt{2}(120 B a^{\frac{13}{2}} \sin(\frac{1}{2} dx + \frac{1}{2} c)^7 - 84 A a^{\frac{13}{2}})}{210 d}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output
$$\frac{-1/210*(105*\sqrt{2}*(A*\sqrt{a} - B*\sqrt{a})*\log(\sin(1/2*d*x + 1/2*c) + 1)/(a*\text{sgn}(\cos(1/2*d*x + 1/2*c))) - 105*\sqrt{2}*(A*\sqrt{a} - B*\sqrt{a})*\log(-\sin(1/2*d*x + 1/2*c))/(a*\text{sgn}(\cos(1/2*d*x + 1/2*c))) + 4*\sqrt{2}*(120*B*a^{13/2}*\sin(1/2*d*x + 1/2*c)^7 - 84*A*a^{13/2}*\sin(1/2*d*x + 1/2*c)^5 - 168*B*a^{13/2}*\sin(1/2*d*x + 1/2*c)^3 + 70*A*a^{13/2}*\sin(1/2*d*x + 1/2*c)^3 + 140*B*a^{13/2}*\sin(1/2*d*x + 1/2*c)^3 - 105*A*a^{13/2}*\sin(1/2*d*x + 1/2*c))/(a^7*\text{sgn}(\cos(1/2*d*x + 1/2*c))))/d}$$

3.100.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^3 (A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2),x)`

output `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2), x)`

3.101 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

3.101.1 Optimal result 1033
 3.101.2 Mathematica [A] (verified) 1034
 3.101.3 Rubi [A] (verified) 1034
 3.101.4 Maple [A] (verified) 1038
 3.101.5 Fricas [A] (verification not implemented) 1038
 3.101.6 Sympy [F] 1039
 3.101.7 Maxima [B] (verification not implemented) 1039
 3.101.8 Giac [A] (verification not implemented) 1040
 3.101.9 Mupad [F(-1)] 1041

3.101.1 Optimal result

Integrand size = 33, antiderivative size = 159

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx = \frac{\sqrt{2}(A-B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} - \frac{4(5A-7B) \sin(c+dx)}{15d\sqrt{a+a \cos(c+dx)}} + \frac{2B \cos^2(c+dx) \sin(c+dx)}{5d\sqrt{a+a \cos(c+dx)}} + \frac{2(5A-B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{15ad}$$

```
output (A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)-4/15*(5*A-7*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/5*B*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/15*(5*A-B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a/d
```


3.101.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.59

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{2\cos\left(\frac{1}{2}(c+dx)\right)\left(15(A-B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right)\right) + (-10A+29B+2(5A-B)\cos(c+dx)+3B\cos(2(c+dx)))\sin\left(\frac{1}{2}(c+dx)\right)}{15d\sqrt{a(1+\cos(c+dx))}}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Cos[(c + d*x)/2]*(15*(A - B)*ArcTanh[Sin[(c + d*x)/2]] + (-10*A + 29*B + 2*(5*A - B)*Cos[c + d*x] + 3*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(15*d*Sqrt[a*(1 + Cos[c + d*x])])`

3.101.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.09, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3462, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^2(A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\sqrt{a\sin\left(c+dx+\frac{\pi}{2}\right)+a}} dx$$

$$\downarrow \text{3462}$$

$$\frac{2\int \frac{\cos(c+dx)(4aB+a(5A-B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2B\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}$$

$$\downarrow \text{27}$$

$$\frac{\int \frac{\cos(c+dx)(4aB+a(5A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2B\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}$$

3.101. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

$$\begin{aligned}
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(4aB+a(5A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx && \downarrow \text{3042} \\
 & \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4aB+a(5A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \int \frac{a(5A-B) \cos^2(c+dx)+4aB \cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx && \downarrow \text{3447} \\
 & \frac{\int \frac{a(5A-B) \cos^2(c+dx)+4aB \cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \int \frac{a(5A-B) \sin(c+dx+\frac{\pi}{2})^2+4aB \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx && \downarrow \text{3042} \\
 & \frac{\int \frac{a(5A-B) \sin(c+dx+\frac{\pi}{2})^2+4aB \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{2 \int \frac{a^2(5A-B)-2a^2(5A-7B) \cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} && \downarrow \text{3502} \\
 & \frac{\int \frac{a^2(5A-B)-2a^2(5A-7B) \cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} && \downarrow \text{27} \\
 & \frac{\int \frac{a^2(5A-B)-2a^2(5A-7B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} && \downarrow \text{3042} \\
 & \frac{\int \frac{a^2(5A-B)-2a^2(5A-7B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} && \downarrow \text{3230} \\
 & \frac{15a^2(A-B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^2(5A-7B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{3a} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} && \downarrow \text{3042} \\
 & \frac{15a^2(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^2(5A-7B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{3a} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \frac{2B \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}}
 \end{aligned}$$

3.101. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 3128 \\
 & \frac{30a^2(A-B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{3a} - \frac{4a^2(5A-7B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
 & \frac{5a}{2B \sin(c+dx) \cos^2(c+dx)} \\
 & \frac{5d\sqrt{a \cos(c+dx)+a}}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \downarrow 219 \\
 & \frac{15\sqrt{2}a^{3/2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{4a^2(5A-7B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} + \frac{2(5A-B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
 & \frac{5a}{2B \sin(c+dx) \cos^2(c+dx)} \\
 & \frac{5d\sqrt{a \cos(c+dx)+a}}{5d\sqrt{a \cos(c+dx)+a}}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*B*Cos[c + d*x]^2*Sin[c + d*x])/(5*d*Sqrt[a + a*Cos[c + d*x]]) + ((2*(5*A - B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((15*Sqrt[2]*a^(3/2)*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/d - (4*a^2*(5*A - 7*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/(3*a)))/(5*a)`

3.101.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3462 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]`

3.101.4 Maple [A] (verified)

Time = 4.26 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.51

method	result
default	$\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(24B\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 20\sqrt{a} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{2} (A+B) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right) + 15a^{\frac{3}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$
parts	$A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-4\sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 3\sqrt{2} \ln\left(\frac{4\sqrt{a} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 4a}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a \right) + 3a^{\frac{3}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d$

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/15*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(24*B*2^(1/2)*(a*si
n(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^4-20*a^(1/2)*(a*sin(1
/2*d*x+1/2*c)^2)^(1/2)*2^(1/2)*(A+B)*sin(1/2*d*x+1/2*c)^2+15*2^(1/2)*ln(4*
(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a*A-15*2^(1
/2)*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a*
B+30*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/a^(3/2)/sin(1/2*d*x
+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.101.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.04

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{4(3B\cos(dx+c)^2 + (5A-B)\cos(dx+c) - 5A + 13B)\sqrt{a\cos(dx+c)+a\sin(dx+c)} - \frac{15\sqrt{2}(A-B)}{30(ad\cos(dx+c)+ad)}}{30(ad\cos(dx+c)+ad)}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x,algorith
m="fracas")
```

output $1/30*(4*(3*B*\cos(d*x + c)^2 + (5*A - B)*\cos(d*x + c) - 5*A + 13*B)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c) - 15*\sqrt{2}*((A - B)*a*\cos(d*x + c) + (A - B)*a)*\log(-(\cos(d*x + c)^2 + 2*\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c))/\sqrt{a} - 2*\cos(d*x + c) - 3)/(\cos(d*x + c)^2 + 2*\cos(d*x + c) + 1))/\sqrt{a})/(a*d*\cos(d*x + c) + a*d)$

3.101.6 Sympy [F]

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \cos^2(c + dx)}{\sqrt{a (\cos(c + dx) + 1)}} dx$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*cos(c + d*x)**2/sqrt(a*(cos(c + d*x) + 1)), x)`

3.101.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 927957 vs. 2(138) = 276.

Time = 17.27 (sec) , antiderivative size = 927957, normalized size of antiderivative = 5836.21

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

3.101.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \int \frac{\cos(c+dx)^2(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2),x)`output `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2), x)`

3.102 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

3.102.1 Optimal result 1042
 3.102.2 Mathematica [A] (verified) 1042
 3.102.3 Rubi [A] (verified) 1043
 3.102.4 Maple [A] (verified) 1045
 3.102.5 Fricas [A] (verification not implemented) 1046
 3.102.6 Sympy [F] 1046
 3.102.7 Maxima [B] (verification not implemented) 1047
 3.102.8 Giac [A] (verification not implemented) 1047
 3.102.9 Mupad [B] (verification not implemented) 1048

3.102.1 Optimal result

Integrand size = 31, antiderivative size = 118

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx = -\frac{\sqrt{2}(A-B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{2(3A-2B) \sin(c+dx)}{3d\sqrt{a+a \cos(c+dx)}} + \frac{2B\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{3ad}$$

```
output -(A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+2/3*(3*A-2*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/3*B*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a/d
```

3.102.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.66

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx = \frac{2 \cos\left(\frac{1}{2}(c+dx)\right) \left(-3(A-B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) + 6A \sin\left(\frac{1}{2}(c+dx)\right) - 4B \sin^3\left(\frac{1}{2}(c+dx)\right)\right)}{3d\sqrt{a(1+\cos(c+dx))}}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Cos[(c + d*x)/2]*(-3*(A - B)*ArcTanh[Sin[(c + d*x)/2]] + 6*A*Sin[(c + d*x)/2] - 4*B*Sin[(c + d*x)/2]^3))/(3*d*Sqrt[a*(1 + Cos[c + d*x])])`

3.102.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{\sqrt{a\cos(c+dx)+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{2 \int \frac{aB+a(3A-2B)\cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2B\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3ad} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{aB+a(3A-2B)\cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2B\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3ad} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\int \frac{aB+a(3A-2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} + \frac{2B \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3ad} \\
& \quad \downarrow \text{3230} \\
& \frac{\frac{2a(3A-2B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - 3a(A-B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2B \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{2a(3A-2B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - 3a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} + \frac{2B \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3ad} \\
& \quad \downarrow \text{3128} \\
& \frac{6a(A-B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{3a} + \frac{2a(3A-2B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} + \frac{2B \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3ad} \\
& \quad \downarrow \text{219} \\
& \frac{\frac{2a(3A-2B)\sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{3\sqrt{2}\sqrt{a}(A-B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{3a} + \frac{2B \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3ad}
\end{aligned}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*B*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d) + ((-3*Sqrt[2]*Sqrt[a]*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a*(3*A - 2*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(3*a)`

3.102.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.102.4 Maple [A] (verified)

Time = 3.73 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.64

method	result
default	$\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(-4B\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 6A\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{a} - 3A\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)}{3a^{\frac{3}{2}}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}d$
parts	$\frac{A\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(2\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 4a}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)a\right)}{a^{\frac{3}{2}}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}d + \frac{B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{a^{\frac{3}{2}}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}d$

3.102.
$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

input `int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/3*cos(1/2*d*x+1/2*c)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^2+6*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-3*A*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a/cos(1/2*d*x+1/2*c))*a+3*B*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a/cos(1/2*d*x+1/2*c))*a)/a^(3/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.102.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.26

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{4(B\cos(dx+c)+3A-B)\sqrt{a\cos(dx+c)+a}\sin(dx+c) - \frac{3\sqrt{2}((A-B)a\cos(dx+c)+(A-B)a)\log\left(-\frac{\cos(dx+c)^2-1}{\sqrt{a}}\right)}{6(ad\cos(dx+c)+ad)}}{6(ad\cos(dx+c)+ad)}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/6*(4*(B*cos(d*x+c)+3*A-B)*sqrt(a*cos(d*x+c)+a)*sin(d*x+c)-3*sqrt(2)*((A-B)*a*cos(d*x+c)+(A-B)*a)*log(-(cos(d*x+c)^2-2*sqrt(2)*sqrt(a*cos(d*x+c)+a)*sin(d*x+c)/sqrt(a)-2*cos(d*x+c)-3)/(cos(d*x+c)^2+2*cos(d*x+c)+1))/sqrt(a))/(a*d*cos(d*x+c)+a*d)`

3.102.6 Sympy [F]

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \int \frac{(A+B\cos(c+dx))\cos(c+dx)}{\sqrt{a(\cos(c+dx)+1)}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A+B*cos(c+d*x))*cos(c+d*x)/sqrt(a*(cos(c+d*x)+1)),x)`

3.102. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

3.102.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38386 vs. $2(101) = 202$.

Time = 0.90 (sec) , antiderivative size = 38386, normalized size of antiderivative = 325.31

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output

```
1/60*((20*(cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c)^3 + 8*(cos(d*x + c)^2 +
sin(d*x + c)^2 + 2*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c)^3 - 20*cos(5/2*d
*x + 5/2*c)^3*sin(d*x + c) + 2*(15*(log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d
*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - log(cos(1/2*d*x + 1/2*c)^2 +
sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*cos(d*x + c)^2 + 15
*(log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/
2*c) + 1) - log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/
2*d*x + 1/2*c) + 1))*sin(d*x + c)^2 + 30*(log(cos(1/2*d*x + 1/2*c)^2 + sin
(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - log(cos(1/2*d*x + 1/2*
c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*cos(d*x + c)
+ 4*(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*cos(d*x + c) + 1)*sin(3/2*d*x + 3
/2*c) - 20*cos(3/2*d*x + 3/2*c)*sin(d*x + c) + 15*log(cos(1/2*d*x + 1/2*c)
^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - 15*log(cos(1/2
*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*co
s(5/2*d*x + 5/2*c)^2 + 30*((log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2
*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2
*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*cos(d*x + c)^2 + (log(cos(1
/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) -
log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2
*c) + 1))*sin(d*x + c)^2 + 2*(log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x ...
```

3.102.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.26

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx =$$

$$\frac{3\sqrt{2}(A\sqrt{a}-B\sqrt{a})\log(\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{\operatorname{asgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} - \frac{3\sqrt{2}(A\sqrt{a}-B\sqrt{a})\log(-\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{\operatorname{asgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} + \frac{4\sqrt{2}(2Ba^{\frac{5}{2}}\sin(\frac{1}{2}dx+\frac{1}{2}c)^3-3Aa^{\frac{5}{2}}\sin(\frac{1}{2}dx+\frac{1}{2}c))}{a^3\operatorname{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))}$$

$6d$

3.102. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `-1/6*(3*sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - 3*sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) + 4*sqrt(2)*(2*B*a^(5/2)*sin(1/2*d*x + 1/2*c)^3 - 3*A*a^(5/2)*sin(1/2*d*x + 1/2*c))/(a^3*sgn(cos(1/2*d*x + 1/2*c))))/d`

3.102.9 Mupad [B] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.36

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{2A\left(2E\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right) - F\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right)\right) \sqrt{\frac{a+a\cos(c+dx)}{2a}}}{d\sqrt{a+a\cos(c+dx)}} + \frac{2B\sin(c+dx)\sqrt{a+a\cos(c+dx)}}{3ad} - \frac{2B\left(4a^2E\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right) - 3a^2F\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right)\right) \sqrt{\frac{a+a\cos(c+dx)}{2a}}}{3a^2d\sqrt{a+a\cos(c+dx)}}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2),x)`

output `(2*A*(2*ellipticE(c/2 + (d*x)/2, 1) - ellipticF(c/2 + (d*x)/2, 1))*((a + a*cos(c + d*x))/(2*a))^(1/2))/(d*(a + a*cos(c + d*x))^(1/2)) + (2*B*sin(c + d*x)*(a + a*cos(c + d*x))^(1/2))/(3*a*d) - (2*B*(4*a^2*ellipticE(c/2 + (d*x)/2, 1) - 3*a^2*ellipticF(c/2 + (d*x)/2, 1))*((a + a*cos(c + d*x))/(2*a))^(1/2))/(3*a^2*d*(a + a*cos(c + d*x))^(1/2))`

3.103 $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.103.1 Optimal result	1049
3.103.2 Mathematica [A] (verified)	1049
3.103.3 Rubi [A] (verified)	1050
3.103.4 Maple [B] (verified)	1051
3.103.5 Fracas [B] (verification not implemented)	1052
3.103.6 Sympy [F]	1052
3.103.7 Maxima [B] (verification not implemented)	1053
3.103.8 Giac [A] (verification not implemented)	1053
3.103.9 Mupad [B] (verification not implemented)	1054

3.103.1 Optimal result

Integrand size = 25, antiderivative size = 78

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \frac{\sqrt{2}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2\sqrt{a+a \cos(c+dx)}}}\right)}{\sqrt{ad}} + \frac{2B \sin(c + dx)}{d\sqrt{a + a \cos(c + dx)}}$$

```
output (A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+2*B*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
```

3.103.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.77

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) \left((A - B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) + 2B \sin\left(\frac{1}{2}(c + dx)\right) \right)}{d\sqrt{a(1 + \cos(c + dx))}}$$

```
input Integrate[(A + B*Cos[c + d*x])/Sqrt[a + a*Cos[c + d*x]],x]
```

```
output (2*Cos[(c + d*x)/2]*((A - B)*ArcTanh[Sin[(c + d*x)/2]] + 2*B*Sin[(c + d*x)/2]))/(d*Sqrt[a*(1 + Cos[c + d*x])])
```


3.103.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sqrt{a \cos(c + dx) + a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx \\
 & \quad \downarrow \text{3230} \\
 & (A - B) \int \frac{1}{\sqrt{\cos(c + dx)a + a}} dx + \frac{2B \sin(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & (A - B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}} dx + \frac{2B \sin(c + dx)}{d \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3128} \\
 & \frac{2B \sin(c + dx)}{d \sqrt{a \cos(c + dx) + a}} - \frac{2(A - B) \int \frac{1}{2a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{\sqrt{2}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2} \sqrt{a \cos(c + dx) + a}}\right)}{\sqrt{ad}} + \frac{2B \sin(c + dx)}{d \sqrt{a \cos(c + dx) + a}}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/Sqrt[a + a*Cos[c + d*x]],x]`

output `(Sqrt[2]*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/(Sqrt[a]*d) + (2*B*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])`

3.103.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

3.103.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 159 vs. 2(67) = 134.

Time = 2.80 (sec) , antiderivative size = 160, normalized size of antiderivative = 2.05

method	result
default	$\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(A\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)+4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)a+2B\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{a}-B\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)}{a^{\frac{3}{2}}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}d}$
parts	$\frac{A\sqrt{2}\operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2}\middle 1\right)}{d\sec\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\operatorname{csgn}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + \frac{B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(2\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}-\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)}{a^{\frac{3}{2}}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}d}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output $\cos(1/2*d*x+1/2*c)*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(A*\ln(4*(a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+a)/\cos(1/2*d*x+1/2*c))*a+2*B*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}-B*\ln(4*(a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+a)/\cos(1/2*d*x+1/2*c))*a/a^{(3/2)}/\sin(1/2*d*x+1/2*c)/(a*\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/d$

3.103.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 135 vs. 2(67) = 134.

Time = 0.31 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.73

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{4 \sqrt{a \cos(dx + c) + a} B \sin(dx + c) - \frac{\sqrt{2}((A - B)a \cos(dx + c) + (A - B)a) \log\left(-\frac{\cos(dx + c)^2 + 2 \sqrt{2} \sqrt{a \cos(dx + c) + a} \sin(dx + c) - 2 \cos(dx + c)}{\sqrt{a} \cos(dx + c)^2 + 2 \cos(dx + c) + 1}\right)}{\sqrt{a}}}{2(ad \cos(dx + c) + ad)}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output $1/2*(4*\sqrt{a*\cos(d*x + c) + a}*B*\sin(d*x + c) - \sqrt{2}*((A - B)*a*\cos(d*x + c) + (A - B)*a)*\log(-(\cos(d*x + c))^2 + 2*\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/\sqrt{a} - 2*\cos(d*x + c) - 3)/(\cos(d*x + c)^2 + 2*\cos(d*x + c) + 1))/\sqrt{a})/(a*d*\cos(d*x + c) + a*d)$

3.103.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a (\cos(c + dx) + 1)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/sqrt(a*(cos(c + d*x) + 1)), x)`

3.103.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19040 vs. 2(67) = 134.

Time = 0.59 (sec) , antiderivative size = 19040, normalized size of antiderivative = 244.10

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output 1/12*(6*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*
sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*
x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*A/sqrt(a) - (12*sqrt(2)*cos(3/
2*d*x + 3/2*c)^3*sin(d*x + c) - 12*(sqrt(2)*cos(d*x + c) + sqrt(2))*sin(3/
2*d*x + 3/2*c)^3 - 8*sqrt(2)*sin(1/2*d*x + 1/2*c)^3 + ((3*sqrt(2)*log(cos(
1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1)
- 3*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/
2*d*x + 1/2*c) + 1) - 8*sqrt(2)*sin(1/2*d*x + 1/2*c))*cos(d*x + c)^2 + (3*
sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*
x + 1/2*c) + 1) - 3*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2
*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1) - 8*sqrt(2)*sin(1/2*d*x + 1/2*c))*sin(
d*x + c)^2 + 24*sqrt(2)*cos(1/2*d*x + 1/2*c)*sin(d*x + c) + 2*(3*sqrt(2)*l
og(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c
) + 1) - 3*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2
*sin(1/2*d*x + 1/2*c) + 1) - 8*sqrt(2)*sin(1/2*d*x + 1/2*c))*cos(d*x + c)
+ 3*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/
2*d*x + 1/2*c) + 1) - 3*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x +
1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1) - 8*sqrt(2)*sin(1/2*d*x + 1/2*c))*
cos(3/2*d*x + 3/2*c)^2 - (8*sqrt(2)*sin(1/2*d*x + 1/2*c)^3 - 3*(sqrt(2)*lo
g(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2...
```

3.103.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.60

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{4\sqrt{2}B \sin(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a} \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \frac{\sqrt{2}(A\sqrt{a} - B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(A\sqrt{a} - B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))}$$

$2d$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `1/2*(4*sqrt(2)*B*sin(1/2*d*x + 1/2*c)/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) + sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))))/d`

3.103.9 Mupad [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.44

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{A F\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right) \sqrt{\frac{2(a+a \cos(c+dx))}{a}} + 2 B E\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right) \sqrt{\frac{2(a+a \cos(c+dx))}{a}} - B F\left(\frac{c}{2} + \frac{dx}{2} \mid 1\right) \sqrt{\frac{2(a+a \cos(c+dx))}{a}}}{d \sqrt{a + a \cos(c + dx)}}$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^(1/2),x)`

output `(A*ellipticF(c/2 + (d*x)/2, 1)*((2*(a + a*cos(c + d*x)))/a)^(1/2) + 2*B*ellipticE(c/2 + (d*x)/2, 1)*((2*(a + a*cos(c + d*x)))/a)^(1/2) - B*ellipticF(c/2 + (d*x)/2, 1)*((2*(a + a*cos(c + d*x)))/a)^(1/2))/(d*(a + a*cos(c + d*x))^(1/2))`

3.104 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.104.1 Optimal result 1055
 3.104.2 Mathematica [A] (verified) 1055
 3.104.3 Rubi [A] (verified) 1056
 3.104.4 Maple [B] (verified) 1058
 3.104.5 Fricas [B] (verification not implemented) 1059
 3.104.6 Sympy [F] 1059
 3.104.7 Maxima [A] (verification not implemented) 1060
 3.104.8 Giac [B] (verification not implemented) 1060
 3.104.9 Mupad [F(-1)] 1061

3.104.1 Optimal result

Integrand size = 31, antiderivative size = 91

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{\sqrt{ad}} - \frac{\sqrt{2}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2} \sqrt{a + a \cos(c + dx)}}\right)}{\sqrt{ad}}$$

output

```
2*A*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d/a^(1/2)-(A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)
```

3.104.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.79

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \frac{2((A - B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) - \sqrt{2}A \operatorname{arctanh}\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right)) \cos\left(\frac{1}{2}(c + dx)\right)}{d \sqrt{a(1 + \cos(c + dx))}}$$

input

```
Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/Sqrt[a + a*Cos[c + d*x]],x]
```

output $(-2*((A - B)*\text{ArcTanh}[\text{Sin}[(c + d*x)/2]] - \text{Sqrt}[2]*A*\text{ArcTanh}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]])*\text{Cos}[(c + d*x)/2])/(d*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])])$

3.104.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{3464} \\
 & \frac{A \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx}{a} - (A-B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx}{a} - (A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
 & \quad \downarrow \text{3128} \\
 & \frac{2(A-B) \int \frac{1}{2a-\frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{A \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx}{a} \\
 & \quad \downarrow \text{219} \\
 & \frac{A \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx}{a} - \frac{\sqrt{2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{\sqrt{ad}} \\
 & \quad \downarrow \text{3252}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{2A \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{\sqrt{2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}}}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} - \frac{\sqrt{2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}}
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*A*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/(Sqrt[a]*d) - (Sqrt[2]*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])/(Sqrt[a]*d)`

3.104.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3252 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`


```
rule 3464 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.104.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(76) = 152.

Time = 4.61 (sec) , antiderivative size = 272, normalized size of antiderivative = 2.99

method	result
default	$\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(A\sqrt{2} \ln\left(\frac{2\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 2\sqrt{2}\sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+4a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}}\right) + A\sqrt{2} \ln\left(-\frac{2\left(\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}\sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \right)$
parts	$\frac{A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\sqrt{2} \ln\left(\frac{4\sqrt{a} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 4a}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) - \ln\left(\frac{4\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2}\sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a+8a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}}\right) \right)}{\sqrt{a} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a)^(1/2), x, method=_RETURNVERBOSE)
```

```
output 1/2*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(A*2^(1/2)*ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))+(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a)+A*2^(1/2)*ln(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))+(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a)-2*ln(2*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*A+2*ln(2*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*B/a^(1/2)/sin(1/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.104.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 171 vs. 2(76) = 152.

Time = 0.34 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.88

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{\sqrt{2}(A - B)\sqrt{a} \log\left(-\frac{\cos(dx+c)^2 - \frac{2\sqrt{2}\sqrt{a}\cos(dx+c)+a\sin(dx+c)}{\sqrt{a}} - 2\cos(dx+c) - 3}{\cos(dx+c)^2 + 2\cos(dx+c) + 1}\right) - A\sqrt{a} \log\left(\frac{a\cos(dx+c)^3 - 7a\cos(dx+c)^2 - 4\sqrt{a}\cos(dx+c) + a}{\cos(dx+c)^3 + \cos(dx+c)^2}\right)}{2ad}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output -1/2*(sqrt(2)*(A - B)*sqrt(a)*log(-(cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(a) - 2*cos(d*x + c) - 3)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - A*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)))/(a*d)
```

3.104.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a(\cos(c + dx) + 1)}} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))**(1/2),x)
```

```
output Integral((A + B*cos(c + d*x))*sec(c + d*x)/sqrt(a*(cos(c + d*x) + 1)), x)
```

3.104.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\left(\sqrt{2} \log\left(\cos\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 2 \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right) - \sqrt{2} \log\left(\cos\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 2 \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right)\right) * B / (\sqrt{a} * d)}{2 \sqrt{ad}}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(1/2),x, algorithm=
"maxima")
```

```
output 1/2*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1
/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x +
1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*B/(sqrt(a)*d)
```

3.104.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 165 vs. 2(76) = 152.

Time = 0.35 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.81

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{\frac{\sqrt{2}(A\sqrt{a}-B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(A\sqrt{a}-B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{2A \log\left(\left|\frac{1}{2} \sqrt{2} + \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right|\right)}{\sqrt{a} \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \frac{2A \log\left(\left|-\frac{1}{2} \sqrt{2} + \sin\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right|\right)}{\sqrt{a} \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))}}{2d}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(1/2),x, algorithm=
"giac")
```

```
output -1/2*(sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a*sgn
(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(-sin(1/2*d*x
+ 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - 2*A*log(abs(1/2*sqrt(2) + s
in(1/2*d*x + 1/2*c)))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) + 2*A*log(abs(-1
/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))))/d
```

3.104.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx) \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^(1/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^(1/2)), x)`

3.105 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.105.1 Optimal result	1062
3.105.2 Mathematica [A] (verified)	1062
3.105.3 Rubi [A] (verified)	1063
3.105.4 Maple [B] (verified)	1066
3.105.5 Fricas [B] (verification not implemented)	1067
3.105.6 Sympy [F]	1067
3.105.7 Maxima [B] (verification not implemented)	1068
3.105.8 Giac [B] (verification not implemented)	1068
3.105.9 Mupad [F(-1)]	1069

3.105.1 Optimal result

Integrand size = 33, antiderivative size = 119

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = -\frac{(A - 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{\sqrt{2}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{A \tan(c + dx)}{d\sqrt{a + a \cos(c + dx)}}$$

output `-(A-2*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d/a^(1/2)+(A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+A*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)`

3.105.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.80

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \frac{\cos\left(\frac{1}{2}(c + dx)\right) \left(2(A - B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) - \sqrt{2}(A - 2B) \operatorname{arctanh}\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2A \sec\left(\frac{1}{2}(c + dx)\right)\right)}{d\sqrt{a(1 + \cos(c + dx))}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/Sqrt[a + a*Cos[c + d*x]],x
]`

output `(Cos[(c + d*x)/2]*(2*(A - B)*ArcTanh[Sin[(c + d*x)/2]] - Sqrt[2]*(A - 2*B)
*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]] + 2*A*Sec[c + d*x]*Sin[(c + d*x)/2]))/(
d*Sqrt[a*(1 + Cos[c + d*x])])`

3.105.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3463, 27, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx$$

↓ 3463

$$\frac{\int -\frac{(a(A-2B)-aA\cos(c+dx))\sec(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{A\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}}$$

↓ 27

$$\frac{A\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{(a(A-2B)-aA\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{2a}$$

↓ 3042

$$\frac{A\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-2B)-aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{2a}$$

↓ 3464

$$\begin{aligned}
 & \frac{A \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{(A - 2B) \int \sqrt{\cos(c + dx)a + a} \sec(c + dx) dx - 2a(A - B) \int \frac{1}{\sqrt{\cos(c + dx)a + a}} dx}{2a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{(A - 2B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx - 2a(A - B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}} dx}{2a} \\
 & \quad \downarrow \text{3128} \\
 & \frac{A \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{4a(A - B) \int \frac{1}{2a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right) + (A - 2B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx}{2a} \\
 & \quad \downarrow \text{219} \\
 & \frac{A \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{(A - 2B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{2\sqrt{2}\sqrt{a}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2}\sqrt{a \cos(c + dx) + a}}\right)}{d}}{2a} \\
 & \quad \downarrow \text{3252} \\
 & \frac{A \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{2a(A - 2B) \int \frac{1}{a - \frac{a^2 \sin^2(c + dx)}{\cos(c + dx)a + a}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right) - \frac{2\sqrt{2}\sqrt{a}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2}\sqrt{a \cos(c + dx) + a}}\right)}{d}}{2a} \\
 & \quad \downarrow \text{219} \\
 & \frac{A \tan(c + dx)}{d\sqrt{a \cos(c + dx) + a}} - \frac{\frac{2\sqrt{a}(A - 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} - \frac{2\sqrt{2}\sqrt{a}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2}\sqrt{a \cos(c + dx) + a}}\right)}{d}}{2a}
 \end{aligned}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/Sqrt[a + a*Cos[c + d*x]],x]
```

```
output -1/2*((2*Sqrt[a]*(A - 2*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d - (2*Sqrt[2]*Sqrt[a]*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/d)/a + (A*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])
```

3.105. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.105.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`
- rule 3464 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.105.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 702 vs. $2(102) = 204$.

Time = 5.45 (sec) , antiderivative size = 703, normalized size of antiderivative = 5.91

method	result
parts	$A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-2a \left(2\sqrt{2} \ln\left(\frac{4\sqrt{a} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 4a}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) - \ln\left(\frac{4 \left(\sqrt{2} a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2} \sqrt{a \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right) \sqrt{a}}{2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}}\right) \right)$
default	Expression too large to display

input `int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNV
ERBOSE)`

output `A*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*a*(2*2^(1/2)*ln(4*(
a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))-ln(-4/(2*co
s(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2
*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))-ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(
1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+
2*a))*sin(1/2*d*x+1/2*c)^2+2*2^(1/2)*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^
2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a+2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)
*a^(1/2)-ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c
)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a-ln(4/(2*cos(1/2*d
*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/
2*c)^2)^(1/2)*a^(1/2)+2*a))*a/a^(3/2)/(2*cos(1/2*d*x+1/2*c)-2^(1/2))/(2*c
os(1/2*d*x+1/2*c)+2^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/
2)/d+1/2*B*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2^(1/2)*ln(2
/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*s
in(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))+2^(1/2)*ln(-2/(2*cos(1/2*d*x+1/2*
c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)
^(1/2)*a^(1/2)-2*a))-2*ln(2*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos
(1/2*d*x+1/2*c)))/a^(1/2)/sin(1/2*d*x+1/2*c)*2^(1/2)/(a*cos(1/2*d*x+1/2*c
^2)^(1/2)/d`

3.105.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. $2(102) = 204$.

Time = 0.33 (sec) , antiderivative size = 259, normalized size of antiderivative = 2.18

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{((A - 2B) \cos(dx + c))^2 + (A - 2B) \cos(dx + c)}{\sqrt{a}} \log \left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 - 4\sqrt{a} \cos(dx+c) + a\sqrt{a}(\cos(dx+c)^3 + \cos(dx+c)^2)}{\cos(dx+c)^3 + \cos(dx+c)^2} \right)$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `-1/4*(((A - 2*B)*cos(d*x + c)^2 + (A - 2*B)*cos(d*x + c))*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) - 4*sqrt(a*cos(d*x + c) + a)*A*sin(d*x + c) + 2*sqrt(2)*((A - B)*a*cos(d*x + c)^2 + (A - B)*a*cos(d*x + c))*log(-cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(a) - 2*cos(d*x + c) - 3)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1))/sqrt(a)/(a*d*cos(d*x + c)^2 + a*d*cos(d*x + c))`

3.105.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a} (\cos(c + dx) + 1)} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/sqrt(a*(cos(c + d*x) + 1)), x)`

3.105.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18436 vs. $2(102) = 204$.

Time = 0.53 (sec) , antiderivative size = 18436, normalized size of antiderivative = 154.92

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output 1/4*((2*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - 2*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2))*cos(d*x + c)^4 + (2*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - 2*sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 + 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) - log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) + 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2) + log(2*cos(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c)^2 - 2*sqrt(2)*cos(1/2*d*x + 1/2*c) - 2*sqrt(2)*sin(1/2*d*x + 1/2*c) + 2))*sin(d*x + c)^4 + 4*sqrt(2)*cos(1/2*d*x + 1/2*c)...
```

3.105.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. $2(102) = 204$.

Time = 0.35 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.84

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\sqrt{2}(A\sqrt{a}-B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(A\sqrt{a}-B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{(A-2B) \log\left(\left|\frac{1}{2} \sqrt{2} + \sin(\frac{1}{2} dx + \frac{1}{2} c)\right|\right)}{\sqrt{a} \operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \frac{(A-2B)}{2d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `1/2*(sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - (A - 2*B)*log(abs(1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) + (A - 2*B)*log(abs(-1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) - 2*sqrt(2)*A*sin(1/2*d*x + 1/2*c)/((2*sin(1/2*d*x + 1/2*c))^2 - 1)*sqrt(a)*sgn(cos(1/2*d*x + 1/2*c)))/d`

3.105.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^2 \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^(1/2)), x)`

3.106
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.106.1 Optimal result 1070
 3.106.2 Mathematica [A] (verified) 1071
 3.106.3 Rubi [A] (verified) 1071
 3.106.4 Maple [B] (warning: unable to verify) 1074
 3.106.5 Fricas [B] (verification not implemented) 1075
 3.106.6 Sympy [F] 1076
 3.106.7 Maxima [B] (verification not implemented) 1076
 3.106.8 Giac [B] (verification not implemented) 1077
 3.106.9 Mupad [F(-1)] 1078

3.106.1 Optimal result

Integrand size = 33, antiderivative size = 165

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \frac{(7A - 4B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{4\sqrt{ad}} - \frac{\sqrt{2}(A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2}\sqrt{a + a \cos(c + dx)}}\right)}{\sqrt{ad}} - \frac{(A - 4B) \tan(c + dx)}{4d\sqrt{a + a \cos(c + dx)}} + \frac{A \sec(c + dx) \tan(c + dx)}{2d\sqrt{a + a \cos(c + dx)}}$$

```
output 1/4*(7*A-4*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d/a^(1/2)
-(A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)-1/4*(A-4*B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/2*A*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
```

3.106.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.69

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c + dx)\right) \left(-8(A - B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) + \sqrt{2}(7A - 4B) \operatorname{arctanh}\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2s\right)}{4d\sqrt{a(1 + \cos(c + dx))}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/Sqrt[a + a*Cos[c + d*x]],x]`

output `(Cos[(c + d*x)/2]*(-8*(A - B)*ArcTanh[Sin[(c + d*x)/2]] + Sqrt[2]*(7*A - 4*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sec[c + d*x]*(-A + 4*B + 2*A*Sec[c + d*x])*Sin[(c + d*x)/2))/(4*d*Sqrt[a*(1 + Cos[c + d*x])])`

3.106.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3463, 27, 3042, 3463, 27, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a \cos(c + dx) + a}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^3 \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx$$

$$\downarrow \text{3463}$$

$$\frac{\int -\frac{(a(A-4B)-3aA \cos(c+dx)) \sec^2(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{2a} + \frac{A \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}}$$

$$\downarrow \text{27}$$

$$\frac{A \tan(c + dx) \sec(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{(a(A-4B)-3aA \cos(c+dx)) \sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a}$$

3.106. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\int \frac{a(A-4B) - 3aA \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^2 \sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx}{4a} \\
& \downarrow \text{3463} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\int \frac{(a^2(7A-4B) - a^2(A-4B) \cos(c+dx)) \sec(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)} + a} \\
& \downarrow \text{27} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)} + a} - \int \frac{(a^2(7A-4B) - a^2(A-4B) \cos(c+dx)) \sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{2a}}{4a} \\
& \downarrow \text{3042} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)} + a} - \int \frac{a^2(7A-4B) - a^2(A-4B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx}{2a}}{4a} \\
& \downarrow \text{3464} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)} + a} - \frac{a(7A-4B) \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx - 8a^2(A-B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{2a}}{4a} \\
& \downarrow \text{3042} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)} + a} - \frac{a(7A-4B) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})} dx - 8a^2(A-B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx}{2a}}{4a} \\
& \downarrow \text{3128} \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)} + a} - \frac{\frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)} + a} - \frac{16a^2(A-B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) + a(7A-4B) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})} dx}{2a}}{4a}
\end{aligned}$$

3.106. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 219 \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} - \\
 & \frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a(7A-4B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} a+a}{\sin(c+dx+\frac{\pi}{2})} dx - \frac{8\sqrt{2}a^{3/2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a} \\
 & \frac{4a}{4a} \\
 & \downarrow 3252 \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} - \\
 & \frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a^2(7A-4B) \int \frac{1}{a-\frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{8\sqrt{2}a^{3/2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a} \\
 & \frac{4a}{4a} \\
 & \downarrow 219 \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} - \\
 & \frac{a(A-4B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a^{3/2}(7A-4B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right) - \frac{8\sqrt{2}a^{3/2}(A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a} \\
 & \frac{4a}{4a}
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/Sqrt[a + a*Cos[c + d*x]],x]`

output `(A*Sec[c + d*x]*Tan[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) - (-1/2*((2*a^(3/2)*(7*A - 4*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (8*Sqrt[2]*a^(3/2)*(A - B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d)/a + (a*(A - 4*B)*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])/(4*a)`

3.106.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

$$3.106. \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3464 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.106.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1151 vs. 2(140) = 280.

Time = 5.88 (sec) , antiderivative size = 1152, normalized size of antiderivative = 6.98

method	result	size
parts	Expression too large to display	1152
default	Expression too large to display	1252

input `int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNV
ERBOSE)`

output `-1/2*A*cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*a*(8*2^(1/2)*1
n(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))-7*ln(-4
/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*s
in(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))-7*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1
/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*
a^(1/2)+2*a))*sin(1/2*d*x+1/2*c)^4+(-32*2^(1/2)*ln(4*(a^(1/2)*(a*sin(1/2*
d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a-4*2^(1/2)*(a*sin(1/2*d*x+1/2*
c)^2)^(1/2)*a^(1/2)+28*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(
1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a+28*1
n(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*
(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a)*sin(1/2*d*x+1/2*c)^2+8*2^(
1/2)*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a
-7*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/
2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a-7*ln(-4/(2*cos(1/2*d*x+1
/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)
^2)^(1/2)*a^(1/2)-2*a))*a-2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)
)/a^(3/2)/(2*cos(1/2*d*x+1/2*c)-2^(1/2))^2/(2*cos(1/2*d*x+1/2*c)+2^(1/2))^
2/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d-1/2*B*cos(1/2*d*x+1/
2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*a*(2^(1/2)*ln(2/(2*cos(1/2*d*x+1/2
c)+2^(1/2))(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c...`

3.106.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 284 vs. 2(140) = 280.

Time = 0.34 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.72

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{((7A - 4B) \cos(dx + c)^3 + (7A - 4B) \cos(dx + c)^2) \sqrt{a} \log \left(\frac{a \cos(dx+c)^3 - 7a \cos(dx+c)^2 + 4 \sqrt{a \cos(dx+c) + a} \sqrt{a}}{\cos(dx+c)^3 + \cos(dx+c)} \right)}{...}$$

3.106. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `-1/16*(((7*A - 4*B)*cos(d*x + c)^3 + (7*A - 4*B)*cos(d*x + c)^2)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 + 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*((A - 4*B)*cos(d*x + c) - 2*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c) + 8*sqrt(2)*((A - B)*a*cos(d*x + c)^3 + (A - B)*a*cos(d*x + c)^2)*log(-(cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(a) - 2*cos(d*x + c) - 3)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1))/sqrt(a))/(a*d*cos(d*x + c)^3 + a*d*cos(d*x + c)^2)`

3.106.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a (\cos(c + dx) + 1)}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/sqrt(a*(cos(c + d*x) + 1)), x)`

3.106.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76209 vs. $2(140) = 280$.

Time = 2.97 (sec) , antiderivative size = 76209, normalized size of antiderivative = 461.87

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output

```

-1/16*((4*sqrt(2)*cos(6*d*x + 6*c)^2*sin(3/2*d*x + 3/2*c) + 16*sqrt(2)*cos
(5*d*x + 5*c)^2*sin(3/2*d*x + 3/2*c) + 36*sqrt(2)*cos(4*d*x + 4*c)^2*sin(3
/2*d*x + 3/2*c) + 64*sqrt(2)*cos(3*d*x + 3*c)^2*sin(3/2*d*x + 3/2*c) + 36*
sqrt(2)*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + 4*sqrt(2)*sin(6*d*x + 6*
c)^2*sin(3/2*d*x + 3/2*c) + 16*sqrt(2)*sin(5*d*x + 5*c)^2*sin(3/2*d*x + 3/
2*c) + 36*sqrt(2)*sin(4*d*x + 4*c)^2*sin(3/2*d*x + 3/2*c) + 64*sqrt(2)*sin
(3*d*x + 3*c)^2*sin(3/2*d*x + 3/2*c) + 36*sqrt(2)*sin(2*d*x + 2*c)^2*sin(3
/2*d*x + 3/2*c) - 8*(3*sqrt(2)*cos(3/2*d*x + 3/2*c)*sin(2*d*x + 2*c) - 3*s
qrt(2)*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + 2*sqrt(2)*cos(3/2*d*x + 3/2
*c)*sin(d*x + c) + (sqrt(2)*sin(9/2*d*x + 9/2*c) + 3*sqrt(2)*sin(7/2*d*x +
7/2*c) - 3*sqrt(2)*sin(5/2*d*x + 5/2*c) - sqrt(2)*sin(3/2*d*x + 3/2*c))*c
os(6*d*x + 6*c) + 2*(sqrt(2)*sin(9/2*d*x + 9/2*c) + 3*sqrt(2)*sin(7/2*d*x
+ 7/2*c) - 3*sqrt(2)*sin(5/2*d*x + 5/2*c) - sqrt(2)*sin(3/2*d*x + 3/2*c))*
cos(5*d*x + 5*c) - (3*sqrt(2)*sin(4*d*x + 4*c) + 4*sqrt(2)*sin(3*d*x + 3*c
) + 3*sqrt(2)*sin(2*d*x + 2*c) + 2*sqrt(2)*sin(d*x + c))*cos(9/2*d*x + 9/2
*c) + 3*(3*sqrt(2)*sin(7/2*d*x + 7/2*c) - 3*sqrt(2)*sin(5/2*d*x + 5/2*c) -
sqrt(2)*sin(3/2*d*x + 3/2*c))*cos(4*d*x + 4*c) - 3*(4*sqrt(2)*sin(3*d*x +
3*c) + 3*sqrt(2)*sin(2*d*x + 2*c) + 2*sqrt(2)*sin(d*x + c))*cos(7/2*d*x +
7/2*c) - 4*(3*sqrt(2)*sin(5/2*d*x + 5/2*c) + sqrt(2)*sin(3/2*d*x + 3/2*c)
)*cos(3*d*x + 3*c) + 3*(3*sqrt(2)*sin(2*d*x + 2*c) + 2*sqrt(2)*sin(d*x ...

```

3.106.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(140) = 280$.

Time = 0.36 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.74

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{4\sqrt{2}(A\sqrt{a}-B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{4\sqrt{2}(A\sqrt{a}-B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{(7A-4B) \log\left(\left|\frac{1}{2}\sqrt{2} + \sin(\frac{1}{2} dx + \frac{1}{2} c)\right|\right)}{\sqrt{a}\operatorname{asgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \dots$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `-1/8*(4*sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - 4*sqrt(2)*(A*sqrt(a) - B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(1/2*d*x + 1/2*c))) - (7*A - 4*B)*log(abs(1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) + (7*A - 4*B)*log(abs(-1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(sqrt(a)*sgn(cos(1/2*d*x + 1/2*c))) - 2*(2*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 8*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 + sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c) + 4*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c))/((2*sin(1/2*d*x + 1/2*c)^2 - 1)^2*a*sgn(cos(1/2*d*x + 1/2*c)))/d`

3.106.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^3 \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^(1/2)), x)`

$$3.107 \quad \int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$$

3.107.1 Optimal result	1079
3.107.2 Mathematica [A] (verified)	1080
3.107.3 Rubi [A] (verified)	1080
3.107.4 Maple [A] (verified)	1086
3.107.5 Fricas [A] (verification not implemented)	1086
3.107.6 Sympy [F(-1)]	1087
3.107.7 Maxima [F(-1)]	1087
3.107.8 Giac [F(-2)]	1088
3.107.9 Mupad [F(-1)]	1088

3.107.1 Optimal result

Integrand size = 33, antiderivative size = 261

$$\begin{aligned} \int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = & -\frac{(15A-19B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} \\ & + \frac{(A-B) \cos^4(c+dx) \sin(c+dx)}{2d(a+a \cos(c+dx))^{3/2}} + \frac{(651A-799B) \sin(c+dx)}{105ad\sqrt{a+a \cos(c+dx)}} \\ & + \frac{(63A-67B) \cos^2(c+dx) \sin(c+dx)}{70ad\sqrt{a+a \cos(c+dx)}} - \frac{(7A-11B) \cos^3(c+dx) \sin(c+dx)}{14ad\sqrt{a+a \cos(c+dx)}} \\ & - \frac{(273A-397B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{210a^2d} \end{aligned}$$

output

```
1/2*(A-B)*cos(d*x+c)^4*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)-1/4*(15*A-19*B)
*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*
2^(1/2)+1/105*(651*A-799*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/70*(63
*A-67*B)*cos(d*x+c)^2*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)-1/14*(7*A-11*B
)*cos(d*x+c)^3*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)-1/210*(273*A-397*B)*s
in(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a^2/d
```

3.107.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.64

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \frac{105(15A-19B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right)\cos^5\left(\frac{1}{2}(c+dx)\right) - \frac{1}{2}\cos^5\left(\frac{1}{2}(c+dx)\right)}{(a+a\cos(c+dx))^{3/2}}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2), x]`

output `(105*(15*A - 19*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 - (Cos[(c + d*x)/2]^3*(1974*A - 2161*B + 6*(273*A - 277*B)*Cos[c + d*x] + (-84*A + 256*B)*Cos[2*(c + d*x)] + 42*A*Cos[3*(c + d*x)] - 18*B*Cos[3*(c + d*x)] + 15*B*Cos[4*(c + d*x)])*Sin[(c + d*x)/2])/2)/(105*d*(a*(1 + Cos[c + d*x]))^(3/2)*(-1 + Sin[(c + d*x)/2]^2))`

3.107.3 Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.10, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3456, 27, 3042, 3462, 27, 3042, 3462, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^4 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a\sin\left(c+dx+\frac{\pi}{2}\right)+a)^{3/2}} dx \\ & \quad \downarrow \text{3456} \\ & \frac{\int \frac{\cos^3(c+dx)(8a(A-B)-a(7A-11B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.107. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{\cos^3(c+dx)(8a(A-B)-a(7A-11B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^3(8a(A-B)-a(7A-11B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3462} \\
 & \frac{2\int -\frac{\cos^2(c+dx)(6a^2(7A-11B)-a^2(63A-67B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{7a} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} + \\
 & \quad \frac{4a^2}{2d(a\cos(c+dx)+a)^{3/2}} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & -\frac{\int \frac{\cos^2(c+dx)(6a^2(7A-11B)-a^2(63A-67B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{7a} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} + \\
 & \quad \frac{4a^2}{2d(a\cos(c+dx)+a)^{3/2}} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(6a^2(7A-11B)-a^2(63A-67B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{7a} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} + \\
 & \quad \frac{4a^2}{2d(a\cos(c+dx)+a)^{3/2}} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3462} \\
 & -\frac{2\int -\frac{\cos(c+dx)(4a^3(63A-67B)-a^3(273A-397B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} + \\
 & \quad \frac{4a^2}{2d(a\cos(c+dx)+a)^{3/2}} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.107. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\frac{\int \frac{\cos(c+dx)(4a^3(63A-67B)-a^3(273A-397B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{4a^2}{7a} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4a^3(63A-67B)-a^3(273A-397B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{4a^2}{7a} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3447

$$\frac{\int \frac{4a^3(63A-67B)\cos(c+dx)-a^3(273A-397B)\cos^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{4a^2}{7a} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3042

$$\frac{\int \frac{4a^3(63A-67B)\sin(c+dx+\frac{\pi}{2})-a^3(273A-397B)\sin^2(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{4a^2}{7a} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3502

$$\frac{2 \int \frac{a^4(273A-397B)-2a^4(651A-799B)\cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a(7A-11B)\sin(c+dx)\cos^3(c+dx)}{7d\sqrt{a\cos(c+dx)+a}} +$$

$$\frac{4a^2}{7a} \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 27

3.107. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\frac{\int \frac{a^4(273A-397B)-2a^4(651A-799B)\cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{5a} - \frac{2a(7A-11)}{7d}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \quad 4a^2$$

↓ 3042

$$\frac{\int \frac{a^4(273A-397B)-2a^4(651A-799B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{5a} - \frac{2a(7A-11)}{7d}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \quad 4a^2$$

↓ 3230

$$\frac{105a^4(15A-19B)\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^4(651A-799B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{5a} - \frac{2a(7A-11)}{7d}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \quad 4a^2$$

↓ 3042

$$\frac{105a^4(15A-19B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^4(651A-799B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{5a} - \frac{2a(7A-11)}{7d}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \quad 4a^2$$

↓ 3128

$$\frac{210a^4(15A-19B)\int \frac{1}{2a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{4a^4(651A-799B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{3a} - \frac{2a(7A-11)}{7d}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \quad 4a^2$$

3.107. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

↓ 219

$$\frac{-\frac{2a^2(63A-67B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(273A-397B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{105\sqrt{2}a^{7/2}(15A-19B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{4a^4(651A-799B)}{d\sqrt{a\cos(c+dx)+a}}}{7a} = \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

```
input Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]
```

```
output ((A - B)*Cos[c + d*x]^4*Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2)) + (-2*a*(7*A - 11*B)*Cos[c + d*x]^3*Sin[c + d*x])/(7*d*Sqrt[a + a*Cos[c + d*x]]) - ((-2*a^2*(63*A - 67*B)*Cos[c + d*x]^2*Sin[c + d*x])/(5*d*Sqrt[a + a*Cos[c + d*x]]) - ((-2*a^2*(273*A - 397*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) - ((105*Sqrt[2]*a^(7/2)*(15*A - 19*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d - (4*a^4*(651*A - 799*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(5*a))/(7*a))/(4*a^2)
```

3.107.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3128 Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]
```

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3462 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m + n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.107.4 Maple [A] (verified)

Time = 4.92 (sec) , antiderivative size = 448, normalized size of antiderivative = 1.72

method	result
default	$\sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(960B\sqrt{a} \sqrt{2} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(\sin^8 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 96\sqrt{a} \sqrt{2} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} (7A+17B) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 2 \right)$
parts	$\frac{A\sqrt{2} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(32\sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{a \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} - 32\sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{a \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} + 80\sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \right)}{20 \cos \left(\frac{dx}{2} + \frac{c}{2} \right) a}$

```
input int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/420/cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(960*B*a^(1/2)*2^(
1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^8-96*a^(1/2)*2^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(7*A+17*B)*sin(1/2*d*x+1/2*c)^6+224*a^(1/
2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(3*A+8*B)*sin(1/2*d*x+1/2*c)^4-3
5*2^(1/2)*(48*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-45*A*ln(4*(a^(1/2)*
(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a-16*B*(a*sin(1/2*d*
x+1/2*c)^2)^(1/2)*a^(1/2)+57*B*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)+a)/cos(1/2*d*x+1/2*c))*a)*sin(1/2*d*x+1/2*c)^2-1575*2^(1/2)*ln(4*(a^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a*A+1995*2^(1/2)*l
n(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a*B+178
5*A*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-1785*B*2^(1/2)*(a*sin(1
/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/a^(5/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+
1/2*c)^2)^(1/2)/d
```

3.107.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.92

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx =$$

$$105\sqrt{2}((15A-19B)\cos(dx+c)^2+2(15A-19B)\cos(dx+c)+15A-19B)\sqrt{a}\log\left(-\frac{a\cos(dx+c)^2-2}{a+a\cos(dx+c)}\right)$$

3.107. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/840*(105*sqrt(2)*((15*A - 19*B)*cos(d*x + c)^2 + 2*(15*A - 19*B)*cos(d*x + c) + 15*A - 19*B)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - 4*(60*B*cos(d*x + c)^4 + 12*(7*A - 3*B)*cos(d*x + c)^3 - 28*(3*A - 7*B)*cos(d*x + c)^2 + 12*(63*A - 67*B)*cos(d*x + c) + 1029*A - 1201*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.107.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.107.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `Timed out`

3.107.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{[-27222589353675077077069968594541456916480,0]:[1,0,-2]%
```

3.107.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^4 (A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx$$

```
input int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)
```

```
output int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2), x)
```

3.108
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$$

3.108.1 Optimal result 1089
 3.108.2 Mathematica [A] (verified) 1090
 3.108.3 Rubi [A] (verified) 1090
 3.108.4 Maple [B] (verified) 1095
 3.108.5 Fricas [A] (verification not implemented) 1096
 3.108.6 Sympy [F(-1)] 1096
 3.108.7 Maxima [F(-1)] 1096
 3.108.8 Giac [F(-2)] 1097
 3.108.9 Mupad [F(-1)] 1097

3.108.1 Optimal result

Integrand size = 33, antiderivative size = 216

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = \frac{(11A-15B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A-B) \cos^3(c+dx) \sin(c+dx)}{2d(a+a \cos(c+dx))^{3/2}} - \frac{(65A-93B) \sin(c+dx)}{15ad\sqrt{a+a \cos(c+dx)}} - \frac{(5A-9B) \cos^2(c+dx) \sin(c+dx)}{10ad\sqrt{a+a \cos(c+dx)}} + \frac{(35A-39B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{30a^2d}$$

```
output 1/2*(A-B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)+1/4*(11*A-15*B)
*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*
2^(1/2)-1/15*(65*A-93*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)-1/10*(5*A-9
*B)*cos(d*x+c)^2*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/30*(35*A-39*B)*si
n(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a^2/d
```


3.108.2 Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.66

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \frac{-15(11A-15B)\operatorname{arctanh}(\sin(\frac{1}{2}(c+dx)))\cos^5(\frac{1}{2}(c+dx)) + \cos^5(\frac{1}{2}(c+dx))}{(a+a\cos(c+dx))^{3/2}}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2), x]`

output `(-15*(11*A - 15*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 + Cos[(c + d*x)/2]^3*(85*A - 141*B + 3*(20*A - 39*B)*Cos[c + d*x] + (-10*A + 6*B)*Cos[2*(c + d*x)] - 3*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2]/(15*d*(a*(1 + Cos[c + d*x]))^(3/2)*(-1 + Sin[(c + d*x)/2]^2))`

3.108.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3456, 27, 3042, 3462, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\ & \quad \downarrow \text{3456} \\ & \frac{\int \frac{\cos^2(c+dx)(6a(A-B)-a(5A-9B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\cos^2(c+dx)(6a(A-B)-a(5A-9B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \end{aligned}$$

3.108. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{\sin(c+dx+\frac{\pi}{2})^2 (6a(A-B) - a(5A-9B) \sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{3042} \\
& \frac{2 \int -\frac{\cos(c+dx)(4a^2(5A-9B) - a^2(35A-39B) \cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{3462} \\
& - \frac{\int \frac{\cos(c+dx)(4a^2(5A-9B) - a^2(35A-39B) \cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{27} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4a^2(5A-9B) - a^2(35A-39B) \sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{3042} \\
& - \frac{\int \frac{\cos(c+dx)(4a^2(5A-9B) - a^2(35A-39B) \cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{3447} \\
& - \frac{\int \frac{4a^2(5A-9B) \cos(c+dx) - a^2(35A-39B) \cos^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{3042} \\
& - \frac{\int \frac{4a^2(5A-9B) \sin(c+dx+\frac{\pi}{2}) - a^2(35A-39B) \sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} + \\
& \frac{4a^2}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \downarrow \text{3502}
\end{aligned}$$

3.108. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & - \frac{2 \int -\frac{a^3(35A-39B)-2a^3(65A-93B) \cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{4a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
 & \frac{2d(a \cos(c+dx) + a)^{3/2}}{27} \\
 & - \frac{\int \frac{a^3(35A-39B)-2a^3(65A-93B) \cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{4a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
 & \frac{2d(a \cos(c+dx) + a)^{3/2}}{3042} \\
 & - \frac{\int \frac{a^3(35A-39B)-2a^3(65A-93B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{4a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
 & \frac{2d(a \cos(c+dx) + a)^{3/2}}{3230} \\
 & - \frac{15a^3(11A-15B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^3(65A-93B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{3a} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{4a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
 & \frac{2d(a \cos(c+dx) + a)^{3/2}}{3042} \\
 & - \frac{15a^3(11A-15B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^3(65A-93B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{3a} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx) \cos^2(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{4a^2}{(A-B) \sin(c+dx) \cos^3(c+dx)} \\
 & \frac{2d(a \cos(c+dx) + a)^{3/2}}{3128}
 \end{aligned}$$

3.108. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{30a^3(11A-15B)}{2a-\frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} \int \frac{1}{d} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) \\
 & - \frac{4a^3(65A-93B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \frac{4a^2}{3a} \\
 & \quad \downarrow \text{219} \\
 & \frac{15\sqrt{2}a^{5/2}(11A-15B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{4a^3(65A-93B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(35A-39B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} - \frac{2a(5A-9B) \sin(c+dx)}{5d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \frac{4a^2}{3a}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output `((A - B)*Cos[c + d*x]^3*Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2)) + (-2*a*(5*A - 9*B)*Cos[c + d*x]^2*Sin[c + d*x]/(5*d*Sqrt[a + a*Cos[c + d*x]])) - ((-2*a*(35*A - 39*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x]/(3*d) - ((15*Sqrt[2]*a^(5/2)*(11*A - 15*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]))/d - (4*a^3*(65*A - 93*B)*Sin[c + d*x]/(d*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(5*a))/(4*a^2)`

3.108.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.108. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &&
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])`

rule 3462 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.108.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 444 vs. $2(189) = 378$.

Time = 4.57 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.06

method	result
default	$\sqrt{2} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(-96B \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{a \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} + 80 \sqrt{a} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) A + 96B \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \right)$
parts	$\frac{A \sqrt{2} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(16 \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{a \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} + 8 \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \right) - 33 \ln \left(\frac{4 \sqrt{a} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}}{\cos \left(\frac{dx}{2} + \frac{c}{2} \right)} \right)}{12 \cos \left(\frac{dx}{2} + \frac{c}{2} \right) a^{\frac{5}{2}} \sin \left(\frac{dx}{2} + \frac{c}{2} \right) \sqrt{a \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}}$

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/60/cos(1/2*d*x+1/2*c)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-96*B*(a*s
in(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^6+80*a^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4+40*A*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^
2)^(1/2)*sin(1/2*d*x+1/2*c)^2-165*A*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(
1/2)+a)/cos(1/2*d*x+1/2*c))*sin(1/2*d*x+1/2*c)^2*a-240*B*(a*sin(1/2*d*x+1
/2*c)^2)^(1/2)*a^(1/2)*sin(1/2*d*x+1/2*c)^2+225*B*ln(4*(a^(1/2)*(a*sin(1/2
*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*sin(1/2*d*x+1/2*c)^2*a-135*A*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+165*A*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1
/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*a+255*B*(a*sin(1/2*d*x+1/2*c)^2)^(1/
2)*a^(1/2)-225*B*ln(4*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d
*x+1/2*c))*a/a^(5/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.108.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.04

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx =$$

$$\frac{15\sqrt{2}((11A-15B)\cos(dx+c)^2 + 2(11A-15B)\cos(dx+c) + 11A-15B)\sqrt{a}\log\left(-\frac{a\cos(dx+c)^2+2\sqrt{2}\sqrt{a}\cos(dx+c)+a}{a\cos(dx+c)+a}\right)}{2(a+a\cos(dx+c))^{3/2}}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/120*(15*sqrt(2)*((11*A - 15*B)*cos(d*x + c)^2 + 2*(11*A - 15*B)*cos(d*x + c) + 11*A - 15*B)*sqrt(a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - 4*(12*B*cos(d*x + c)^3 + 4*(5*A - 3*B)*cos(d*x + c)^2 - 12*(5*A - 9*B)*cos(d*x + c) - 95*A + 147*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.108.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.108.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm m="maxima")`

output Timed out

3.108.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm m="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{[%%{%%{[2475880078570760549798248448,0]:[1,0,-2]%%},[10]%%},0]

3.108.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^3(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$$

input `int((cos(c+d*x)^3*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(3/2),x)`

output `int((cos(c+d*x)^3*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(3/2),x)`

3.109
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$$

3.109.1 Optimal result 1098
 3.109.2 Mathematica [A] (verified) 1098
 3.109.3 Rubi [A] (verified) 1099
 3.109.4 Maple [B] (verified) 1103
 3.109.5 Fricas [A] (verification not implemented) 1103
 3.109.6 Sympy [F(-1)] 1104
 3.109.7 Maxima [F(-1)] 1104
 3.109.8 Giac [F(-2)] 1105
 3.109.9 Mupad [F(-1)] 1105

3.109.1 Optimal result

Integrand size = 33, antiderivative size = 171

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = -\frac{(7A-11B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A-B) \cos^2(c+dx) \sin(c+dx)}{2d(a+a \cos(c+dx))^{3/2}} + \frac{(9A-13B) \sin(c+dx)}{3ad\sqrt{a+a \cos(c+dx)}} - \frac{(3A-7B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{6a^2d}$$

output `1/2*(A-B)*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)-1/4*(7*A-11*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)+1/3*(9*A-13*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)-1/6*(3*A-7*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a^2/d`

3.109.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.57

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = \frac{-3(7A-11B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) \cos\left(\frac{1}{2}(c+dx)\right) + (15A-11B)}{6ad\sqrt{a(1-\cos(c+dx))}}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output $(-3*(7*A - 11*B)*\text{ArcTanh}[\text{Sin}[(c + d*x)/2]]*\text{Cos}[(c + d*x)/2] + (15*A - 17*B + 12*(A - B)*\text{Cos}[c + d*x] + 2*B*\text{Cos}[2*(c + d*x)])*\text{Tan}[(c + d*x)/2])/(6*a*d*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])])$

3.109.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3456, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^2(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx$$

↓ 3456

$$\frac{\int \frac{\cos(c+dx)(4a(A-B)-a(3A-7B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 27

$$\frac{\int \frac{\cos(c+dx)(4a(A-B)-a(3A-7B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4a(A-B)-a(3A-7B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3447

$$\frac{\int \frac{4a(A-B)\cos(c+dx)-a(3A-7B)\cos^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3042

$$\begin{aligned}
& \frac{\int \frac{4a(A-B) \sin(c+dx+\frac{\pi}{2}) - a(3A-7B) \sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^2(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{3502} \\
& \frac{2 \int -\frac{a^2(3A-7B) - 2a^2(9A-13B) \cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
& \quad \frac{4a^2}{(A-B) \sin(c+dx) \cos^2(c+dx)} \\
& \quad \frac{2d(a \cos(c+dx) + a)^{3/2}}{27} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a^2(3A-7B) - 2a^2(9A-13B) \cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
& \quad \frac{4a^2}{(A-B) \sin(c+dx) \cos^2(c+dx)} \\
& \quad \frac{2d(a \cos(c+dx) + a)^{3/2}}{3042} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(3A-7B) - 2a^2(9A-13B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
& \quad \frac{4a^2}{(A-B) \sin(c+dx) \cos^2(c+dx)} \\
& \quad \frac{2d(a \cos(c+dx) + a)^{3/2}}{3230} \\
& \quad \downarrow \text{3230} \\
& \frac{3a^2(7A-11B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^2(9A-13B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{3a} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
& \quad \frac{4a^2}{(A-B) \sin(c+dx) \cos^2(c+dx)} \\
& \quad \frac{2d(a \cos(c+dx) + a)^{3/2}}{3042} \\
& \quad \downarrow \text{3042} \\
& \frac{3a^2(7A-11B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^2(9A-13B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{3a} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
& \quad \frac{4a^2}{(A-B) \sin(c+dx) \cos^2(c+dx)} \\
& \quad \frac{2d(a \cos(c+dx) + a)^{3/2}}{3128} \\
& \quad \downarrow \text{3128}
\end{aligned}$$

3.109. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & -\frac{6a^2(7A-11B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{3a} - \frac{4a^2(9A-13B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
 & \frac{4a^2}{2d(a \cos(c+dx) + a)^{3/2}} (A - B) \sin(c+dx) \cos^2(c+dx) \\
 & \quad \downarrow \text{219} \\
 & -\frac{3\sqrt{2}a^{3/2}(7A-11B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{4a^2(9A-13B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2(3A-7B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d} + \\
 & \frac{4a^2}{2d(a \cos(c+dx) + a)^{3/2}} (A - B) \sin(c+dx) \cos^2(c+dx)
 \end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output `((A - B)*Cos[c + d*x]^2*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)) + (-2*(3*A - 7*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) - ((3*Sqrt[2]*a^(3/2)*(7*A - 11*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d - (4*a^2*(9*A - 13*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(4*a^2)`

3.109.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.109.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(148) = 296.

Time = 4.22 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.91

method	result
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(16B\left(\cos^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{2}\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-21A\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+4a}}{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)\sqrt{2}\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a+3\right)}{4\cos\left(\frac{dx}{2}+\frac{c}{2}\right)a^{\frac{5}{2}}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}d}$
parts	$\frac{A\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(-7\sqrt{2}\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+4a}}{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)a\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{2}\sqrt{a}+\sqrt{2}\right)}{4\cos\left(\frac{dx}{2}+\frac{c}{2}\right)a^{\frac{5}{2}}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}d}$

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/12/cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(16*B*cos(1/2*d*x+1
/2*c)^4*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-21*A*ln(2*(2*a^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*2^(1/2)*cos(1/2*
d*x+1/2*c)^2*a+33*B*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/co
s(1/2*d*x+1/2*c))*2^(1/2)*cos(1/2*d*x+1/2*c)^2*a+24*A*cos(1/2*d*x+1/2*c)^2
*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-40*B*cos(1/2*d*x+1/2*c)^2*
2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+3*A*a^(1/2)*2^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)-3*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)
)/a^(5/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.109.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.20

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx =$$

$$\frac{3\sqrt{2}((7A-11B)\cos(dx+c)^2+2(7A-11B)\cos(dx+c)+7A-11B)\sqrt{a}\log\left(-\frac{a\cos(dx+c)^2-2\sqrt{2}\sqrt{a}\cos(dx+c)}{\cos(dx+c)}\right)}{24(a^2d\cos(dx+c))^{3/2}}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm
m="fracas")
```

3.109. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

output
$$\begin{aligned} & -1/24*(3*\sqrt{2})*((7*A - 11*B)*\cos(d*x + c)^2 + 2*(7*A - 11*B)*\cos(d*x + c) \\ & + 7*A - 11*B)*\sqrt{a}*\log(-a*\cos(d*x + c)^2 - 2*\sqrt{2}*\sqrt{a*\cos(d*x + c) + a} \\ & *\sqrt{a}*\sin(d*x + c) - 2*a*\cos(d*x + c) - 3*a)/(\cos(d*x + c)^2 + 2*\cos(d*x + c) + 1)) \\ & - 4*(4*B*\cos(d*x + c)^2 + 12*(A - B)*\cos(d*x + c) + 15*A - 19*B)*\sqrt{a*\cos(d*x + c) + a} \\ & *\sin(d*x + c))/(a^2*d*\cos(d*x + c)^2 + 2*a^2*d*\cos(d*x + c) + a^2*d) \end{aligned}$$

3.109.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.109.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `Timed out`

3.109.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{%%{[-54043195528445952,0]:[1,0,-2]%%},[6]%%},0):[1,0,%%{-
```

3.109.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^2 (A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx$$

```
input int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)
```

```
output int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2), x)
```


$$3.110 \quad \int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$$

3.110.1 Optimal result 1106
 3.110.2 Mathematica [A] (verified) 1106
 3.110.3 Rubi [A] (verified) 1107
 3.110.4 Maple [B] (verified) 1109
 3.110.5 Fricas [A] (verification not implemented) 1110
 3.110.6 Sympy [F] 1110
 3.110.7 Maxima [F(-1)] 1111
 3.110.8 Giac [F(-2)] 1111
 3.110.9 Mupad [F(-1)] 1111

3.110.1 Optimal result

Integrand size = 31, antiderivative size = 118

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = \frac{(3A-7B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{(A-B) \sin(c+dx)}{2d(a+a \cos(c+dx))^{3/2}} + \frac{2B \sin(c+dx)}{ad\sqrt{a+a \cos(c+dx)}}$$

```
output -1/2*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)+1/4*(3*A-7*B)*arctanh(1/2*
sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)+2*B*si
n(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.110.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.88

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = \frac{-((3A-7B)\operatorname{arctanh}(\sin(\frac{1}{2}(c+dx))) \cos^5(\frac{1}{2}(c+dx))) + \cos^3(\frac{1}{2}(c+dx))}{d(a(1+\cos(c+dx)))^{3/2}(-1+\sin(\frac{1}{2}(c+dx)))}$$

```
input Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x
]
```

```
output (-((3*A - 7*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5) + Cos[(c + d*
x)/2]^3*(A - 5*B - 4*B*Cos[c + d*x])*Sin[(c + d*x)/2])/(d*(a*(1 + Cos[c +
d*x]))^(3/2)*(-1 + Sin[(c + d*x)/2]^2))
```

3.110. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

3.110.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3498, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a\cos(c+dx)+a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\
 & \quad \downarrow \text{3498} \\
 & -\frac{\int -\frac{3a(A-B)+4aB\cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3a(A-B)+4aB\cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3a(A-B)+4aB\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3230} \\
 & \frac{a(3A-7B)\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx + \frac{8aB\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.110. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\frac{a(3A - 7B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx + \frac{8aB \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{4a^2} - \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}}$$

↓ 3128

$$\frac{\frac{8aB \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(3A-7B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{4a^2}}{d} - \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}}$$

↓ 219

$$\frac{\frac{\sqrt{2}\sqrt{a}(3A-7B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{8aB \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}}}{4a^2} - \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output `-1/2*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^(3/2)) + ((Sqrt[2]*Sqrt[a]*(3*A - 7*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d + (8*a*B*SIN[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])/(4*a^2)`

3.110.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*SIN[c + d*x]])], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

```
rule 3230 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3498 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b - a*
B + b*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[1
/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b
*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]
```

3.110.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 255 vs. 2(101) = 202.

Time = 4.10 (sec) , antiderivative size = 256, normalized size of antiderivative = 2.17

method	result
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(3A \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \sqrt{2}\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a - 7B \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \sqrt{2}\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^{\frac{5}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a} \right)}{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^{\frac{5}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$
parts	$\frac{A \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(3\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \right)}{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^{\frac{5}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \frac{B \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^{\frac{5}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2), x, method=_RETURNVER
BOSE)
```

output $\frac{1}{4}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(3*A*\ln(2*(2*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*a)/\cos(1/2*d*x+1/2*c))*2^{(1/2)}*\cos(1/2*d*x+1/2*c)^2*a-7*B*\ln(2*(2*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*a)/\cos(1/2*d*x+1/2*c))*2^{(1/2)}*\cos(1/2*d*x+1/2*c)^2*a+8*B*\cos(1/2*d*x+1/2*c)^2*2^{(1/2)}*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}-A*a^{(1/2)}*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+B*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)})/\cos(1/2*d*x+1/2*c)/a^{(5/2)}/\sin(1/2*d*x+1/2*c)/(a*\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/d$

3.110.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.60

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \frac{\sqrt{2}((3A-7B)\cos(dx+c)^2+2(3A-7B)\cos(dx+c)+3A-7B)\sqrt{a}\log\left(-\frac{a\cos(dx+c)^2+2\sqrt{2}\sqrt{a\cos(dx+c)}}{\cos(dx+c)^2}\right)}{8(a^2d\cos(dx+c))^2+2a^2d}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output $-1/8*(\sqrt{2})*((3*A-7*B)*\cos(d*x+c)^2+2*(3*A-7*B)*\cos(d*x+c)+3*A-7*B)*\sqrt{a}*\log(-a*\cos(d*x+c)^2+2*\sqrt{2}*\sqrt{a*\cos(d*x+c)+a}*\sqrt{a}*\sin(d*x+c)-2*a*\cos(d*x+c)-3*a)/(\cos(d*x+c)^2+2*\cos(d*x+c)+1))-4*(4*B*\cos(d*x+c)-A+5*B)*\sqrt{a*\cos(d*x+c)+a}*\sin(d*x+c)/(a^2*d*\cos(d*x+c)^2+2*a^2*d*\cos(d*x+c)+a^2*d)$

3.110.6 Sympy [F]

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \int \frac{(A+B\cos(c+dx))\cos(c+dx)}{(a(\cos(c+dx)+1))^{3/2}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A+B*cos(c+d*x))*cos(c+d*x)/(a*(cos(c+d*x)+1))**(3/2),x)`

3.110. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

3.110.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
output Timed out
```

3.110.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{[%%{%%{[268435456,0]:[1,0,-2]%%},[2]%%},0]:[1,0,%%{-1,[1]%%}}
```

3.110.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{3/2}} dx$$

```
input int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)
```

```
output int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2), x)
```

3.111 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.111.1 Optimal result 1112
 3.111.2 Mathematica [A] (verified) 1112
 3.111.3 Rubi [A] (verified) 1113
 3.111.4 Maple [B] (verified) 1114
 3.111.5 Fricas [B] (verification not implemented) 1115
 3.111.6 Sympy [F] 1115
 3.111.7 Maxima [B] (verification not implemented) 1116
 3.111.8 Giac [B] (verification not implemented) 1116
 3.111.9 Mupad [F(-1)] 1117

3.111.1 Optimal result

Integrand size = 25, antiderivative size = 87

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(A + 3B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A - B) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}}$$

output $1/2*(A-B)*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^(3/2)+1/4*(A+3*B)*\operatorname{arctanh}(1/2*\sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*\cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)$

3.111.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.72

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(A + 3B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) \cos^3\left(\frac{1}{2}(c + dx)\right) + \frac{1}{2}(A - B) \sin(c + dx)}{d(a(1 + \cos(c + dx)))^{3/2}}$$

input $\operatorname{Integrate}[(A + B*\operatorname{Cos}[c + d*x])/(a + a*\operatorname{Cos}[c + d*x])^(3/2),x]$

output $((A + 3*B)*\operatorname{ArcTanh}[\operatorname{Sin}[(c + d*x)/2]]*\operatorname{Cos}[(c + d*x)/2]^3 + ((A - B)*\operatorname{Sin}[c + d*x])/2)/(d*(a*(1 + \operatorname{Cos}[c + d*x]))^(3/2))$

3.111.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3229, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a \cos(c + dx) + a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2}} dx \\
 & \quad \downarrow \text{3229} \\
 & \frac{(A + 3B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A + 3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3128} \\
 & \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}} - \frac{(A + 3B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{2ad} \\
 & \quad \downarrow \text{219} \\
 & \frac{(A + 3B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A - B) \sin(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^(3/2),x]`

output `((A + 3*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) + ((A - B)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2))`

3.111.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3229 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.111.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(72) = 144.

Time = 4.06 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.53

method	result
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(A \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \sqrt{2} \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a + 3B \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \sqrt{2} \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a}{4 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) a^{\frac{5}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d \right)}{A \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \right) + B \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{4a^{\frac{5}{2}} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \dots$
parts	$\frac{A \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{2} \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{a} \right) + B \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{4a^{\frac{5}{2}} \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d} + \dots$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVERBOSE)`

output $\frac{1}{4} \frac{\cos(1/2 dx + 1/2 c) (a \sin(1/2 dx + 1/2 c)^2)^{1/2} (A \ln(2 (2 a^{1/2}) (a \sin(1/2 dx + 1/2 c)^2)^{1/2} + 2 a) / \cos(1/2 dx + 1/2 c))^2 + 2 (A + 3 B) \cos(dx + c) + A + 3 B}{(a + a \cos(c + dx))^{3/2}} \sqrt{a} \log \left(\frac{\sqrt{2} ((A + 3 B) \cos(dx + c))^2 + 2 (A + 3 B) \cos(dx + c) + A + 3 B}{8 (a^2 d \cos(dx + c) + a^2 d)} \right) dx$

3.111.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 172 vs. $2(72) = 144$.

Time = 0.29 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.98

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2} ((A + 3 B) \cos(dx + c))^2 + 2 (A + 3 B) \cos(dx + c) + A + 3 B}{8 (a^2 d \cos(dx + c) + a^2 d)} \sqrt{a} \log \left(\frac{\sqrt{2} ((A + 3 B) \cos(dx + c))^2 + 2 (A + 3 B) \cos(dx + c) + A + 3 B}{8 (a^2 d \cos(dx + c) + a^2 d)} \right) dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output $\frac{1}{8} \sqrt{2} ((A + 3 B) \cos(dx + c))^2 + 2 (A + 3 B) \cos(dx + c) + A + 3 B \sqrt{a} \log \left(\frac{\sqrt{2} ((A + 3 B) \cos(dx + c))^2 + 2 (A + 3 B) \cos(dx + c) + A + 3 B}{8 (a^2 d \cos(dx + c) + a^2 d)} \right) + 4 \sqrt{a} \cos(dx + c) - 2 a \cos(dx + c) - 3 a}{(a + a \cos(c + dx))^{3/2}} dx$

3.111.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a (\cos(c + dx) + 1))^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))/(a*(cos(c + d*x) + 1))**(3/2), x)`

3.111.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 62254 vs. $2(72) = 144$.

Time = 2.48 (sec) , antiderivative size = 62254, normalized size of antiderivative = 715.56

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output

```
1/16*((3*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*
sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d
*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1) + 4*sqrt(2)*sin(5/2*d*x + 5/2*
c))*cos(3*d*x + 3*c)^2 + 27*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*
d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1
/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*cos(2*d*x
+ 2*c)^2 + 27*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2
+ 2*sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(
1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*cos(d*x + c)^2 + (3*(sqr
t(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x +
1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2
- 2*sin(1/2*d*x + 1/2*c) + 1))*cos(3*d*x + 3*c)^2 + 27*(sqrt(2)*log(cos(1
/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) -
sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d
*x + 1/2*c) + 1))*cos(2*d*x + 2*c)^2 + 27*(sqrt(2)*log(cos(1/2*d*x + 1/2*c
)^2 + sin(1/2*d*x + 1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(c
os(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) +
1))*cos(d*x + c)^2 + 3*(sqrt(2)*log(cos(1/2*d*x + 1/2*c)^2 + sin(1/2*d*x +
1/2*c)^2 + 2*sin(1/2*d*x + 1/2*c) + 1) - sqrt(2)*log(cos(1/2*d*x + 1/2*c)
^2 + sin(1/2*d*x + 1/2*c)^2 - 2*sin(1/2*d*x + 1/2*c) + 1))*sin(3*d*x + ...
```

3.111.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(72) = 144$.

Time = 0.39 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.84

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2}(A\sqrt{a} + 3B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(A\sqrt{a} + 3B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^2 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{2\sqrt{2}(A\sqrt{a} + 3B\sqrt{a})}{(\sin(\frac{1}{2} dx + \frac{1}{2} c))}$$

3.111. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `1/8*(sqrt(2)*(A*sqrt(a) + 3*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(A*sqrt(a) + 3*B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(1/2*d*x + 1/2*c))) - 2*sqrt(2)*(A*sqrt(a)*sin(1/2*d*x + 1/2*c) - B*sqrt(a)*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c)^2 - 1)*a^2*sgn(cos(1/2*d*x + 1/2*c)))/d`

3.111.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^(3/2), x)`

3.112 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.112.1 Optimal result 1118
 3.112.2 Mathematica [A] (verified) 1118
 3.112.3 Rubi [A] (verified) 1119
 3.112.4 Maple [B] (verified) 1122
 3.112.5 Fricas [B] (verification not implemented) 1122
 3.112.6 Sympy [F] 1123
 3.112.7 Maxima [B] (verification not implemented) 1123
 3.112.8 Giac [B] (verification not implemented) 1124
 3.112.9 Mupad [F(-1)] 1125

3.112.1 Optimal result

Integrand size = 31, antiderivative size = 127

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{3/2} d} - \frac{(5A - B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2} a^{3/2} d} - \frac{(A - B) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}}$$

output `2*A*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d-1/2*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)-1/4*(5*A-B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)`

3.112.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.03

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(5A - B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) \cos^5\left(\frac{1}{2}(c + dx)\right) - 4\sqrt{2}A \operatorname{arctan}\left(\frac{\sin\left(\frac{1}{2}(c + dx)\right)}{\cos\left(\frac{1}{2}(c + dx)\right)}\right)}{d(a(1 + \cos(c + dx)))^{3/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^(3/2),x]`

```
output ((5*A - B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 - 4*Sqrt[2]*A*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 + (A - B)*Cos[(c + d*x)/2]^3*Sin[(c + d*x)/2])/(d*(a*(1 + Cos[c + d*x]))^(3/2)*(-1 + Sin[(c + d*x)/2]^2))
```

3.112.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3457, 27, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx$$

↓ 3457

$$\frac{\int \frac{(4aA-a(A-B)\cos(c+dx))\sec(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 27

$$\frac{\int \frac{(4aA-a(A-B)\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3042

$$\frac{\int \frac{4aA-a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3464

$$\frac{4A \int \sqrt{\cos(c+dx)a+a} \sec(c+dx) dx - a(5A-B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 3042

3.112. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{4A \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx - a(5A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{3128} \\
& \frac{2a(5A-B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) + 4A \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{4A \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})} dx - \frac{\sqrt{2}\sqrt{a}(5A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{3252} \\
& \frac{8aA \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{\sqrt{2}\sqrt{a}(5A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{8\sqrt{a}A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right) - \frac{\sqrt{2}\sqrt{a}(5A-B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^(3/2),x]`

output `((8*sqrt[a]*A*ArcTanh[(sqrt[a]*Sin[c + d*x])/sqrt[a + a*Cos[c + d*x]])]/d - (sqrt[2]*sqrt[a]*(5*A - B)*ArcTanh[(sqrt[a]*Sin[c + d*x])/(sqrt[2]*sqrt[a + a*Cos[c + d*x]])])/d)/(4*a^2) - ((A - B)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2))`

3.112.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`
- rule 3252 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3464 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.112.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 363 vs. 2(106) = 212.

Time = 5.19 (sec) , antiderivative size = 364, normalized size of antiderivative = 2.87

method	result
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(2A\sqrt{2}\ln\left(\frac{2\sqrt{2}a\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+2\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a+4a}}{2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{2}}\right)a\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2A\sqrt{2}\ln\left(-\frac{2\left(\sqrt{2}a\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{2}\right)}{\dots}$
parts	$\frac{A\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(5\sqrt{2}\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+4a}}{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)a\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-4\ln\left(\frac{4\sqrt{2}a\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+4\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a}}{2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{2}}\right)}{4a^{\frac{5}{2}}\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sin\left(\frac{dx}{2}\right)}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVER
BOSE)
```

```
output 1/4/a^(5/2)/cos(1/2*d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*2^(1/2)
*ln(2/(2*cos(1/2*d*x+1/2*c)+2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)
*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*a*cos(1/2*d*x+1/2*c)^2+2*A*2
^(1/2)*ln(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2)))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-
2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*a*cos(1/2*d*x+1/2*c)^
2-5*A*ln(2*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/cos(1/2*d*x+1/2*c))*
a*cos(1/2*d*x+1/2*c)^2+B*ln(2*(a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+a)/c
os(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^2*a-A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)
*a^(1/2)+B*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/sin(1/2*d*x+1/2*c)*2^(1
/2)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.112.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 281 vs. 2(106) = 212.

Time = 0.34 (sec) , antiderivative size = 281, normalized size of antiderivative = 2.21

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{\sqrt{2}((5A - B) \cos(dx + c)^2 + 2(5A - B) \cos(dx + c) + 5A - B) \sqrt{a} \log\left(-\frac{a \cos(dx+c)^2 - 2\sqrt{2}\sqrt{a \cos(dx+c)+a}\sqrt{a}}{\cos(dx+c)^2 + 2\cos(dx+c) + 2}\right)}{\dots}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/8*(sqrt(2)*((5*A - B)*cos(d*x + c)^2 + 2*(5*A - B)*cos(d*x + c) + 5*A - B)*sqrt(a)*log(-(a*cos(d*x + c))^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - 4*(A*cos(d*x + c)^2 + 2*A*cos(d*x + c) + A)*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) + 4*sqrt(a*cos(d*x + c) + a)*(A - B)*sin(d*x + c)/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.112.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a (\cos(c + dx) + 1))^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a*(cos(c + d*x) + 1))**(3/2), x)`

3.112.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15722 vs. $2(106) = 212$.

Time = 0.84 (sec) , antiderivative size = 15722, normalized size of antiderivative = 123.80

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output $\frac{1}{8} \cdot (8A \cdot \log(\frac{-2\sqrt{2} - 4\sin(\frac{1}{2}dx + \frac{1}{2}c)}{\sqrt{2} - 4\sin(\frac{1}{2}dx + \frac{1}{2}c)}) / (a^{3/2} \cdot \text{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))) - \sqrt{2} \cdot (5A\sqrt{a} - B\sqrt{a}) \cdot \log(\frac{\sin(\frac{1}{2}dx + \frac{1}{2}c) + 1}{a^2 \cdot \text{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))}) + \sqrt{2} \cdot (5A\sqrt{a} - B\sqrt{a}) \cdot \log(\frac{-\sin(\frac{1}{2}dx + \frac{1}{2}c) + 1}{a^2 \cdot \text{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c))}) + 2\sqrt{2} \cdot (A\sqrt{a} \cdot \sin(\frac{1}{2}dx + \frac{1}{2}c) - B\sqrt{a} \cdot \sin(\frac{1}{2}dx + \frac{1}{2}c)) / ((\sin(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1) \cdot a^2 \cdot \text{sgn}(\cos(\frac{1}{2}dx + \frac{1}{2}c)))) / d$

3.112.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx) (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^(3/2)), x)`

3.113
$$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.113.1 Optimal result 1126
 3.113.2 Mathematica [A] (verified) 1126
 3.113.3 Rubi [A] (verified) 1127
 3.113.4 Maple [B] (verified) 1131
 3.113.5 Fricas [B] (verification not implemented) 1132
 3.113.6 Sympy [F] 1132
 3.113.7 Maxima [B] (verification not implemented) 1133
 3.113.8 Giac [F(-2)] 1133
 3.113.9 Mupad [F(-1)] 1134

3.113.1 Optimal result

Integrand size = 33, antiderivative size = 170

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$-\frac{(3A - 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{3/2}d} + \frac{(9A - 5B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d}$$

$$-\frac{(A - B) \tan(c + dx)}{2d(a + a \cos(c + dx))^{3/2}} + \frac{(3A - B) \tan(c + dx)}{2ad\sqrt{a + a \cos(c + dx)}}$$

output

```
-(3*A-2*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d+1/4*(9*A-5*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d-1/2*(A-B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)+1/2*(3*A-B)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.113.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.83

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\cos^3\left(\frac{1}{2}(c + dx)\right) \left(2(9A - 5B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right)\right) + \frac{4\sqrt{2}(3A - B) \tan\left(\frac{1}{2}(c + dx)\right)}{2d(a(1 + \cos(c + dx)))^{3/2}}}{2d(a(1 + \cos(c + dx)))^{3/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^(3/2),x]`

output `(Cos[(c + d*x)/2]^3*(2*(9*A - 5*B)*ArcTanh[Sin[(c + d*x)/2]] + (4*Sqrt[2]*(3*A - 2*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^2 + 2*(-3*A + B - 2*A*Sec[c + d*x])*Sin[(c + d*x)/2])/(-1 + Sin[(c + d*x)/2]^2))/(2*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.113.3 Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3457, 27, 3042, 3463, 25, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(2a(3A-B)-3a(A-B)\cos(c+dx))\sec^2(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(2a(3A-B)-3a(A-B)\cos(c+dx))\sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2a(3A-B)-3a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3463}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\int -\frac{(2a^2(3A-2B)-a^2(3A-B)\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}}dx}{4a^2} + \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{25} \\
& \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{(2a^2(3A-2B)-a^2(3A-B)\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}}dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{2a^2(3A-2B)-a^2(3A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3464} \\
& \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a(3A-2B)\int\sqrt{\cos(c+dx)a+a}\sec(c+dx)dx - a^2(9A-5B)\int\frac{1}{\sqrt{\cos(c+dx)a+a}}dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a(3A-2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})}dx - a^2(9A-5B)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3128} \\
& \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(9A-5B)\int\frac{1}{2a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) + 2a(3A-2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})}dx}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a(3A-2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})}dx - \sqrt{2}a^{3/2}(9A-5B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{4a^2} - \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}
\end{aligned}$$

3.113. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{array}{c}
 \downarrow \text{3252} \\
 \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{4a^2(3A-2B)\int \frac{1}{a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{\sqrt{2}a^{3/2}(9A-5B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{a}}{a} \\
 \frac{4a^2}{(A-B)\tan(c+dx)} \\
 \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
 \downarrow \text{219} \\
 \frac{2a(3A-B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{4a^{3/2}(3A-2B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right) - \frac{\sqrt{2}a^{3/2}(9A-5B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}}{a} \\
 \frac{4a^2}{(A-B)\tan(c+dx)} \\
 \frac{(A-B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}
 \end{array}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^(3/2),x]`

output `-1/2*((A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^(3/2)) + (-(((4*a^(3/2)*(3*A - 2*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d - (Sqrt[2]*a^(3/2)*(9*A - 5*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d)/a) + (2*a*(3*A - B)*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]))/(4*a^2)`

3.113.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.113. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

rule 3128 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)\sin[(c_) + (d_)(x_)]]], x_Symbol] \rightarrow \text{Simp}[-2/d \text{ Subst}[\text{Int}[1/(2*a - x^2)], x], x, b*(\text{Cos}[c + d*x]/\text{Sqrt}[a + b*\text{Sin}[c + d*x]])], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

rule 3252 $\text{Int}[\text{Sqrt}[(a_) + (b_)\sin[(e_) + (f_)(x_)]]/((c_) + (d_)\sin[(e_) + (f_)(x_)])], x_Symbol] \rightarrow \text{Simp}[-2*(b/f) \text{ Subst}[\text{Int}[1/(b*c + a*d - d*x^2)], x], x, b*(\text{Cos}[e + f*x]/\text{Sqrt}[a + b*\text{Sin}[e + f*x]])], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0]$

rule 3457 $\text{Int}(((a_) + (b_)\sin[(e_) + (f_)(x_)])^m * ((A_) + (B_)\sin[(e_) + (f_)(x_)]) * ((c_) + (d_)\sin[(e_) + (f_)(x_)])^n), x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*\text{Cos}[e + f*x]*(a + b*\text{Sin}[e + f*x])^m * ((c + d*\text{Sin}[e + f*x])^{n+1}) / (a*f*(2*m + 1)*(b*c - a*d)), x] + \text{Simp}[1/(a*(2*m + 1)*(b*c - a*d)) \text{ Int}[(a + b*\text{Sin}[e + f*x])^{m+1} * (c + d*\text{Sin}[e + f*x])^n * \text{Simp}[B*(a*c*m + b*d*(n+1)) + A*(b*c*(m+1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{LtQ}[m, -2^{(-1)}] \ \&\& \ !\text{GtQ}[n, 0] \ \&\& \ \text{IntegerQ}[2*m] \ \&\& \ (\text{IntegerQ}[2*n] \ || \ \text{EqQ}[c, 0])]$

rule 3463 $\text{Int}(((a_) + (b_)\sin[(e_) + (f_)(x_)])^m * ((A_) + (B_)\sin[(e_) + (f_)(x_)]) * ((c_) + (d_)\sin[(e_) + (f_)(x_)])^n), x_Symbol] \rightarrow \text{Simp}[(B*c - A*d)*\text{Cos}[e + f*x]*(a + b*\text{Sin}[e + f*x])^m * ((c + d*\text{Sin}[e + f*x])^{n+1}) / (f*(n+1)*(c^2 - d^2)), x] + \text{Simp}[1/(b*(n+1)*(c^2 - d^2)) \text{ Int}[(a + b*\text{Sin}[e + f*x])^m * (c + d*\text{Sin}[e + f*x])^{n+1} * \text{Simp}[A*(a*d*m + b*c*(n+1)) - B*(a*c*m + b*d*(n+1)) + b*(B*c - A*d)*(m + n + 2)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{IntegerQ}[n] \ || \ \text{EqQ}[m + 1/2, 0])]$

rule 3464 $\text{Int}(((A_) + (B_)\sin[(e_) + (f_)(x_)]) / (\text{Sqrt}[(a_) + (b_)\sin[(e_) + (f_)(x_)]] * ((c_) + (d_)\sin[(e_) + (f_)(x_)])), x_Symbol] \rightarrow \text{Simp}[(A*b - a*B)/(b*c - a*d) \text{ Int}[1/\text{Sqrt}[a + b*\text{Sin}[e + f*x]], x], x] + \text{Simp}[(B*c - A*d)/(b*c - a*d) \text{ Int}[\text{Sqrt}[a + b*\text{Sin}[e + f*x]] / (c + d*\text{Sin}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0]$

3.113.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 859 vs. $2(145) = 290$.

Time = 5.95 (sec) , antiderivative size = 860, normalized size of antiderivative = 5.06

method	result	size
parts	Expression too large to display	860
default	Expression too large to display	1051

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/2*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(18*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/
2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*a*cos(1/2*d*x+1/2*c)^4-12*ln
n(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^4*a-12*ln(-
4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*
sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^4*a-9*2^(1/2)
*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*a
*cos(1/2*d*x+1/2*c)^2+6*cos(1/2*d*x+1/2*c)^2*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)
)*2^(1/2)*a^(1/2)+6*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2
*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d
*x+1/2*c)^2*a+6*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*
x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+
1/2*c)^2*a-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/a^(5/2)/cos(1/2
*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)-2^(1/2))/(2*cos(1/2*d*x+1/2*c)+2^(1/2))/
sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+1/4*B/a^(5/2)/cos(1/2*
d*x+1/2*c)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*2^(1/2)*ln(-2/(2*cos(1/2*d*x+
1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)
)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^2*a+2*2^(1/2)*ln(2/(2*cos(1/2*
d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1
/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^2*a-5*ln(2*(a^(1/2)*(a...
```

3.113.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. $2(145) = 290$.

Time = 0.36 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.99

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{\sqrt{2}((9A - 5B) \cos(dx + c))^3 + 2(9A - 5B) \cos(dx + c)^2 + (9A - 5B) \cos(dx + c)}{\sqrt{a}} \log\left(-\frac{a \cos(dx + c)}{\dots}\right)$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/8*(sqrt(2)*((9*A - 5*B)*cos(d*x + c)^3 + 2*(9*A - 5*B)*cos(d*x + c)^2 + (9*A - 5*B)*cos(d*x + c))*sqrt(a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) + 2*((3*A - 2*B)*cos(d*x + c)^3 + 2*(3*A - 2*B)*cos(d*x + c)^2 + (3*A - 2*B)*cos(d*x + c))*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) - 4*((3*A - B)*cos(d*x + c) + 2*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(a^2*d*cos(d*x + c)^3 + 2*a^2*d*cos(d*x + c)^2 + a^2*d*cos(d*x + c))`

3.113.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a (\cos(c + dx) + 1))^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a*(cos(c + d*x) + 1))**(3/2), x)`

3.113.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47933 vs. $2(145) = 290$.

Time = 1.75 (sec) , antiderivative size = 47933, normalized size of antiderivative = 281.96

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(3/2),x, algorithm
m="maxima")
```

```
output 1/4*(128*cos(3/2*d*x + 3/2*c)*sin(3*d*x + 3*c)^3 + 1152*cos(3/2*d*x + 3/2*
c)*sin(2*d*x + 2*c)^3 - 128*cos(3*d*x + 3*c)^3*sin(3/2*d*x + 3/2*c) - 1152
*cos(2*d*x + 2*c)^3*sin(3/2*d*x + 3/2*c) + 32*(4*cos(3/2*d*x + 3/2*c)*sin(
2*d*x + 2*c) - 9*(3*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c) - 28*cos(2*d*x
+ 2*c)*sin(3/2*d*x + 3/2*c) + 3*cos(3/2*d*x + 3/2*c)*sin(d*x + c))*cos(3*d
*x + 3*c)^2 - 96*(11*(3*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c) - 9*cos(3/2
*d*x + 3/2*c)*sin(d*x + c))*cos(2*d*x + 2*c)^2 + 32*(28*cos(3/2*d*x + 3/2*
c)*sin(2*d*x + 2*c) - (3*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c) - 4*cos(3*
d*x + 3*c)*sin(3/2*d*x + 3/2*c) - 4*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c)
+ 27*cos(3/2*d*x + 3/2*c)*sin(d*x + c))*sin(3*d*x + 3*c)^2 - 288*((3*cos(d
*x + c) + 1)*sin(3/2*d*x + 3/2*c) + 4*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c
) - 11*cos(3/2*d*x + 3/2*c)*sin(d*x + c))*sin(2*d*x + 2*c)^2 - 32*(cos(3*d
*x + 3*c)^2*sin(3/2*d*x + 3/2*c) + 6*(3*cos(d*x + c) + 1)*cos(2*d*x + 2*c)
*sin(3/2*d*x + 3/2*c) + 9*cos(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2*c) + sin(3*
d*x + 3*c)^2*sin(3/2*d*x + 3/2*c) + 9*sin(2*d*x + 2*c)^2*sin(3/2*d*x + 3/2
*c) + 18*sin(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c)*sin(d*x + c) + 2*((3*cos(d*
x + c) + 1)*sin(3/2*d*x + 3/2*c) + 3*cos(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c)
)*cos(3*d*x + 3*c) + 6*(sin(2*d*x + 2*c)*sin(3/2*d*x + 3/2*c) + sin(3/2*d*
x + 3/2*c)*sin(d*x + c))*sin(3*d*x + 3*c) + (9*cos(d*x + c)^2 + 9*sin(d*x
+ c)^2 + 6*cos(d*x + c) + 1)*sin(3/2*d*x + 3/2*c))*cos(5*d*x + 5*c) - 9...
```

3.113.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(3/2),x, algorithm
m="giac")
```

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:index.cc index_m i_lex_is_greater Error: Bad Argument Value

3.113.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^2 (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^(3/2)), x)`

3.114
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.114.1 Optimal result 1135
 3.114.2 Mathematica [A] (verified) 1136
 3.114.3 Rubi [A] (verified) 1136
 3.114.4 Maple [B] (verified) 1141
 3.114.5 Fricas [A] (verification not implemented) 1142
 3.114.6 Sympy [F] 1142
 3.114.7 Maxima [F(-2)] 1143
 3.114.8 Giac [A] (verification not implemented) 1143
 3.114.9 Mupad [F(-1)] 1144

3.114.1 Optimal result

Integrand size = 33, antiderivative size = 221

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(19A - 12B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4a^{3/2}d} - \frac{(13A - 9B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{(7A - 6B) \tan(c + dx)}{4ad\sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec(c + dx) \tan(c + dx)}{2d(a + a \cos(c + dx))^{3/2}} + \frac{(2A - B) \sec(c + dx) \tan(c + dx)}{2ad\sqrt{a + a \cos(c + dx)}}$$

```
output 1/4*(19*A-12*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)
/d-1/4*(13*A-9*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(
1/2))/a^(3/2)/d*2^(1/2)-1/2*(A-B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*x+c)
)^(3/2)-1/4*(7*A-6*B)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/2*(2*A-B)*se
c(d*x+c)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.114.2 Mathematica [A] (verified)

Time = 0.99 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.93

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\cos^3\left(\frac{1}{2}(c + dx)\right) \sec^2(c + dx) \left(4(13A - 9B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{(a + a \cos(c + dx))^{3/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^(3/2),x]`

output `(Cos[(c + d*x)/2]^3*Sec[c + d*x]^2*(4*(13*A - 9*B)*ArcTanh[Sin[(c + d*x)/2]]*(Cos[(c + d*x)/2] + Cos[(3*(c + d*x))/2])^2 - 2*Sqrt[2]*(19*A - 12*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*(Cos[(c + d*x)/2] + Cos[(3*(c + d*x))/2])^2 + 4*(3*(A - 2*B) + (6*A - 8*B)*Cos[c + d*x] + (7*A - 6*B)*Cos[2*(c + d*x)])*Sin[(c + d*x)/2))/(16*d*(a*(1 + Cos[c + d*x]))^(3/2)*(-1 + Sin[(c + d*x)/2]^2))`

3.114.3 Rubi [A] (verified)Time = 1.48 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.05, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^3 (a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{3/2}} dx$$

↓ 3457

$$\frac{\int \frac{(4a(2A - B) - 5a(A - B) \cos(c + dx)) \sec^3(c + dx)}{2\sqrt{\cos(c + dx)a + a}} dx}{2a^2} - \frac{(A - B) \tan(c + dx) \sec(c + dx)}{2d(a \cos(c + dx) + a)^{3/2}}$$

↓ 27

3.114. $\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{(4a(2A-B)-5a(A-B)\cos(c+dx))\sec^3(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{4a(2A-B)-5a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3463} \\
& \frac{\int -\frac{2(a^2(7A-6B)-3a^2(2A-B)\cos(c+dx))\sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{2a} + \frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \\
& \quad \frac{4a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{2d(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{27}} \\
& \frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{(a^2(7A-6B)-3a^2(2A-B)\cos(c+dx))\sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{a} \\
& \quad \frac{4a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{2d(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{3042}} \\
& \frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(7A-6B)-3a^2(2A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{a} \\
& \quad \frac{4a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{2d(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{3463}} \\
& \frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int -\frac{(a^3(19A-12B)-a^3(7A-6B)\cos(c+dx))\sec(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{a^2(7A-6B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \\
& \quad \frac{4a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{2d(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{27}} \\
& \frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a^2(7A-6B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{(a^3(19A-12B)-a^3(7A-6B)\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{2a} \\
& \quad \frac{4a^2}{(A-B)\tan(c+dx)\sec(c+dx)} \\
& \quad \frac{2d(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{3042}}
\end{aligned}$$

3.114. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\frac{2a(2A-B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\frac{a^2(7A-6B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^3(19A-12B) - a^3(7A-6B) \sin\left(c+dx+\frac{\pi}{2}\right) dx}{\sin\left(c+dx+\frac{\pi}{2}\right) \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a+a}}{2a}}{a}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 3464

$$\frac{2a(2A-B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\frac{a^2(7A-6B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a^2(19A-12B) \int \sqrt{\cos(c+dx)+a} \sec(c+dx) dx - 2a^3(13A-9B) \int \frac{1}{\sqrt{\cos(c+dx)+a}} dx}{2a}}{a}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 3042

$$\frac{2a(2A-B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\frac{a^2(7A-6B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a^2(19A-12B) \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a+a}}{\sin\left(c+dx+\frac{\pi}{2}\right)} dx - 2a^3(13A-9B) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a+a}} dx}{2a}}{a}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 3128

$$\frac{2a(2A-B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\frac{a^2(7A-6B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a^2(19A-12B) \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a+a}}{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{4a^3(13A-9B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx) a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}}\right)}{2a}}{a}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 219

$$\frac{2a(2A-B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\frac{a^2(7A-6B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a^2(19A-12B) \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a+a}}{\sin\left(c+dx+\frac{\pi}{2}\right)} dx - \frac{2\sqrt{2}a^{5/2}(13A-9B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a}}{a}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 3252

3.114. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

$$\frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a^2(7A-6B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(19A-12B)\int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} dx \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{a} - \frac{2\sqrt{2}a^{5/2}(13A-9B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \frac{4a^2}{}$$

↓ 219

$$\frac{2a(2A-B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a^2(7A-6B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^{5/2}(19A-12B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{2\sqrt{2}a^{5/2}(13A-9B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \frac{4a^2}{}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^(3/2),x]`

output `-1/2*((A - B)*Sec[c + d*x]*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^(3/2)) + ((2*a*(2*A - B)*Sec[c + d*x]*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]) - (-1/2*((2*a^(5/2)*(19*A - 12*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (2*Sqrt[2]*a^(5/2)*(13*A - 9*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d)/a + (a^2*(7*A - 6*B)*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]))/a)/(4*a^2)`

3.114.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.114. \quad \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

rule 3128 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)\sin[(c_) + (d_)(x_)]]], x_Symbol] \rightarrow \text{Simp}[-2/d \text{Subst}[\text{Int}[1/(2*a - x^2)], x], x, b*(\text{Cos}[c + d*x]/\text{Sqrt}[a + b*\text{Sin}[c + d*x]])], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{EqQ}[a^2 - b^2, 0]$

rule 3252 $\text{Int}[\text{Sqrt}[(a_) + (b_)\sin[(e_) + (f_)(x_)]]/((c_) + (d_)\sin[(e_) + (f_)(x_)])], x_Symbol] \rightarrow \text{Simp}[-2*(b/f) \text{Subst}[\text{Int}[1/(b*c + a*d - d*x^2)], x], x, b*(\text{Cos}[e + f*x]/\text{Sqrt}[a + b*\text{Sin}[e + f*x]])], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$

rule 3457 $\text{Int}(((a_) + (b_)\sin[(e_) + (f_)(x_)])^m * ((A_) + (B_)\sin[(e_) + (f_)(x_)]) * ((c_) + (d_)\sin[(e_) + (f_)(x_)])^n), x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*\text{Cos}[e + f*x]*(a + b*\text{Sin}[e + f*x])^m * ((c + d*\text{Sin}[e + f*x])^{n+1}) / (a*f*(2*m + 1)*(b*c - a*d)), x] + \text{Simp}[1/(a*(2*m + 1)*(b*c - a*d)) \text{Int}[(a + b*\text{Sin}[e + f*x])^{m+1} * (c + d*\text{Sin}[e + f*x])^n * \text{Simp}[B*(a*c*m + b*d*(n+1)) + A*(b*c*(m+1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{LtQ}[m, -2^{(-1)}] \&\& !\text{GtQ}[n, 0] \&\& \text{IntegerQ}[2*m] \&\& (\text{IntegerQ}[2*n] || \text{EqQ}[c, 0])]$

rule 3463 $\text{Int}(((a_) + (b_)\sin[(e_) + (f_)(x_)])^m * ((A_) + (B_)\sin[(e_) + (f_)(x_)]) * ((c_) + (d_)\sin[(e_) + (f_)(x_)])^n), x_Symbol] \rightarrow \text{Simp}[(B*c - A*d)*\text{Cos}[e + f*x]*(a + b*\text{Sin}[e + f*x])^m * ((c + d*\text{Sin}[e + f*x])^{n+1}) / (f*(n+1)*(c^2 - d^2)), x] + \text{Simp}[1/(b*(n+1)*(c^2 - d^2)) \text{Int}[(a + b*\text{Sin}[e + f*x])^m * (c + d*\text{Sin}[e + f*x])^{n+1} * \text{Simp}[A*(a*d*m + b*c*(n+1)) - B*(a*c*m + b*d*(n+1)) + b*(B*c - A*d)*(m + n + 2)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegerQ}[n] || \text{EqQ}[m + 1/2, 0])]$

rule 3464 $\text{Int}(((A_) + (B_)\sin[(e_) + (f_)(x_)]) / (\text{Sqrt}[(a_) + (b_)\sin[(e_) + (f_)(x_)]] * ((c_) + (d_)\sin[(e_) + (f_)(x_)])), x_Symbol] \rightarrow \text{Simp}[(A*b - a*B)/(b*c - a*d) \text{Int}[1/\text{Sqrt}[a + b*\text{Sin}[e + f*x]], x], x] + \text{Simp}[(B*c - A*d)/(b*c - a*d) \text{Int}[\text{Sqrt}[a + b*\text{Sin}[e + f*x]] / (c + d*\text{Sin}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$

3.114.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1371 vs. $2(190) = 380$.

Time = 6.82 (sec) , antiderivative size = 1372, normalized size of antiderivative = 6.21

method	result	size
parts	Expression too large to display	1372
default	Expression too large to display	1540

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/2*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(104*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^6*a-76
*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2
)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^6*a-76*ln
n(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(
a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^6*a-104*2^(
1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c
))*a*cos(1/2*d*x+1/2*c)^4+28*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2
)*cos(1/2*d*x+1/2*c)^4+76*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a
cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*co
s(1/2*d*x+1/2*c)^4*a+76*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos
(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1
/2*d*x+1/2*c)^4*a+26*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)+2*a)/cos(1/2*d*x+1/2*c))*a*cos(1/2*d*x+1/2*c)^2-22*cos(1/2*d*x+1/2*c)^2*
(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*2^(1/2)*a^(1/2)-19*ln(-4/(2*cos(1/2*d*x+1/2
*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2
)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^2*a-19*ln(4/(2*cos(1/2*d*x+1/2*c)
+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(
1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^2*a+2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^
2)^(1/2)*a^(1/2))/a^(5/2)/cos(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)-2^(1...
```

3.114.5 Fricas [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.63

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{2\sqrt{2}((13A - 9B) \cos(dx + c)^4 + 2(13A - 9B) \cos(dx + c)^3 + (13A - 9B) \cos(dx + c)^2) \sqrt{a} \log\left(-\frac{a \cos(dx + c) + a}{2\sqrt{2}((13A - 9B) \cos(dx + c)^4 + 2(13A - 9B) \cos(dx + c)^3 + (13A - 9B) \cos(dx + c)^2)}\right)}{2\sqrt{2}((13A - 9B) \cos(dx + c)^4 + 2(13A - 9B) \cos(dx + c)^3 + (13A - 9B) \cos(dx + c)^2)}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(3/2),x, algorithm
m="fricas")
```

```
output -1/16*(2*sqrt(2)*((13*A - 9*B)*cos(d*x + c)^4 + 2*(13*A - 9*B)*cos(d*x + c)
)^3 + (13*A - 9*B)*cos(d*x + c)^2)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt
(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a
)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) + ((19*A - 12*B)*cos(d*x + c)^4 +
2*(19*A - 12*B)*cos(d*x + c)^3 + (19*A - 12*B)*cos(d*x + c)^2)*sqrt(a)*lo
g((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 + 4*sqrt(a*cos(d*x + c) + a)*sqrt
(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^
2)) + 4*((7*A - 6*B)*cos(d*x + c)^2 + (3*A - 4*B)*cos(d*x + c) - 2*A)*sqrt
(a*cos(d*x + c) + a)*sin(d*x + c))/(a^2*d*cos(d*x + c)^4 + 2*a^2*d*cos(d*x
+ c)^3 + a^2*d*cos(d*x + c)^2)
```

3.114.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a(\cos(c + dx) + 1))^{\frac{3}{2}}} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c))**(3/2),x)
```

```
output Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a*(cos(c + d*x) + 1))**(3/2
), x)
```

3.114.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(3/2),x, algorithm
m="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: Memory limit reached. Please j
ump to an outer pointer, quit program and enlarge thememory limits before
executing the program again.
```

3.114.8 Giac [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 354, normalized size of antiderivative = 1.60

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{\sqrt{2}(13A\sqrt{a}-9B\sqrt{a})\log(\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^2\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} - \frac{\sqrt{2}(13A\sqrt{a}-9B\sqrt{a})\log(-\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^2\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} - \frac{(19A-12B)\log\left(\left|\frac{1}{2}\sqrt{2}+\sin(\frac{1}{2}dx+\frac{1}{2}c)\right|\right)}{a^{\frac{3}{2}}\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(3/2),x, algorithm
m="giac")
```

```
output -1/8*(sqrt(2)*(13*A*sqrt(a) - 9*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(
a^2*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(13*A*sqrt(a) - 9*B*sqrt(a))*log(
-sin(1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(1/2*d*x + 1/2*c))) - (19*A - 12*B)
*log(abs(1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(a^(3/2)*sgn(cos(1/2*d*x + 1
/2*c))) + (19*A - 12*B)*log(abs(-1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(a^(
3/2)*sgn(cos(1/2*d*x + 1/2*c))) - 2*sqrt(2)*(A*sqrt(a)*sin(1/2*d*x + 1/2*c
) - B*sqrt(a)*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c)^2 - 1)*a^2*sgn(
cos(1/2*d*x + 1/2*c))) - 2*(10*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 -
8*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 3*sqrt(2)*A*sqrt(a)*sin(1/2*d
*x + 1/2*c) + 4*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c))/((2*sin(1/2*d*x +
1/2*c)^2 - 1)^2*a^2*sgn(cos(1/2*d*x + 1/2*c)))/d
```

3.114. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.114.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^3 (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^(3/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^(3/2)), x)`

3.115
$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

3.115.1 Optimal result 1145
 3.115.2 Mathematica [A] (verified) 1146
 3.115.3 Rubi [A] (verified) 1146
 3.115.4 Maple [B] (verified) 1152
 3.115.5 Fracas [A] (verification not implemented) 1153
 3.115.6 Sympy [F(-1)] 1153
 3.115.7 Maxima [F(-1)] 1154
 3.115.8 Giac [A] (verification not implemented) 1154
 3.115.9 Mupad [F(-1)] 1155

3.115.1 Optimal result

Integrand size = 33, antiderivative size = 261

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{(163A-283B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A-B) \cos^4(c+dx) \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{(13A-21B) \cos^3(c+dx) \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}} - \frac{(985A-1729B) \sin(c+dx)}{120a^2d\sqrt{a+a \cos(c+dx)}} - \frac{(85A-157B) \cos^2(c+dx) \sin(c+dx)}{80a^2d\sqrt{a+a \cos(c+dx)}} + \frac{(475A-787B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{240a^3d}$$

```
output 1/4*(A-B)*cos(d*x+c)^4*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+1/16*(13*A-21*B)
*cos(d*x+c)^3*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+1/32*(163*A-283*B)*ar
ctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(
1/2)-1/120*(985*A-1729*B)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)-1/80*(85
*A-157*B)*cos(d*x+c)^2*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)+1/240*(475*
A-787*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a^3/d
```


3.115.2 Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.53

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{30(163A-283B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right)\cos^3\left(\frac{1}{2}(c+dx)\right) + (-1895A+3491B-5(479A-887B)\cos(c+dx)+(-400A+832B)\cos[2(c+dx)]+40A\cos[3(c+dx)]-40B\cos[3(c+dx)]+12B\cos[4(c+dx)])\tan\left(\frac{c+dx}{2}\right)}{(240a^2(a(1+\cos(c+dx)))^{3/2}}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `(30*(163*A - 283*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + (-1895*A + 3491*B - 5*(479*A - 887*B)*Cos[c + d*x] + (-400*A + 832*B)*Cos[2*(c + d*x)] + 40*A*Cos[3*(c + d*x)] - 40*B*Cos[3*(c + d*x)] + 12*B*Cos[4*(c + d*x)])*Tan[(c + d*x)/2])/(240*a*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.115.3 Rubi [A] (verified)

Time = 1.78 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.10, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3462, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^4 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^{5/2}} dx \\ & \quad \downarrow \text{3456} \\ & \frac{\int \frac{\cos^3(c+dx)(8a(A-B)-a(5A-13B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\cos^3(c+dx)(8a(A-B)-a(5A-13B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \end{aligned}$$

3.115. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\sin(c+dx+\frac{\pi}{2})^3 (8a(A-B) - a(5A-13B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx \\
& \quad + \frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \quad + \frac{\int \frac{\cos^2(c+dx)(6a^2(13A-21B) - a^2(85A-157B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad + \frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad + \frac{\int \frac{\cos^2(c+dx)(6a^2(13A-21B) - a^2(85A-157B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad + \frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad + \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2 (6a^2(13A-21B) - a^2(85A-157B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad + \frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad + \frac{2 \int -\frac{\cos(c+dx)(4a^3(85A-157B) - a^3(475A-787B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad + \frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}} (A-B)\sin(c+dx)\cos^4(c+dx) \\
& \quad + \frac{\int \frac{\cos(c+dx)(4a^3(85A-157B) - a^3(475A-787B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad + \frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}} (A-B)\sin(c+dx)\cos^4(c+dx)
\end{aligned}$$

3.115. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

↓ 3042

$$\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)\left(4a^3(85A-157B)-a^3(475A-787B)\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{5a} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3447

$$\frac{\int \frac{4a^3(85A-157B)\cos(c+dx)-a^3(475A-787B)\cos^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{5a} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{4a^3(85A-157B)\sin\left(c+dx+\frac{\pi}{2}\right)-a^3(475A-787B)\sin\left(c+dx+\frac{\pi}{2}\right)^2}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{5a} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3502

$$\frac{2\int \frac{a^4(475A-787B)-2a^4(985A-1729B)\cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 27

$$\frac{\int \frac{a^4(475A-787B)-2a^4(985A-1729B)\cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-21B)\sin(c+dx)\cos^3(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

3.115. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\int \frac{a^4(475A-787B)-2a^4(985A-1729B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} + \frac{a(13A-283B)}{3a}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \quad 8a^2$$

↓ 3230

$$\frac{15a^4(163A-283B)\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^4(985A-1729B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{4a^2}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \quad 8a^2$$

↓ 3042

$$\frac{15a^4(163A-283B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^4(985A-1729B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{4a^2}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \quad 8a^2$$

↓ 3128

$$\frac{30a^4(163A-283B)\int \frac{1}{2a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{4a^4(985A-1729B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}}}{4a^2}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \quad 8a^2$$

↓ 219

$$\frac{2a^2(85A-157B)\sin(c+dx)\cos^2(c+dx)}{5d\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(475A-787B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} - \frac{15\sqrt{2}a^{7/2}(163A-283B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{5a} - \frac{4a^4(985A-1729B)\sin(c+dx)}{3a}$$

$$\frac{(A-B)\sin(c+dx)\cos^4(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \quad 8a^2$$

3.115. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `((A - B)*Cos[c + d*x]^4*Sin[c + d*x]/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (a*(13*A - 21*B)*Cos[c + d*x]^3*Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2)) + ((-2*a^2*(85*A - 157*B)*Cos[c + d*x]^2*Sin[c + d*x]/(5*d*Sqrt[a + a*Cos[c + d*x]]) - ((-2*a^2*(475*A - 787*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x]/(3*d) - ((15*Sqrt[2]*a^(7/2)*(163*A - 283*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]))/d - (4*a^4*(985*A - 1729*B)*Sin[c + d*x]/(d*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(5*a))/(4*a^2))/(8*a^2)`

3.115.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3462 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m + n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.115.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 466 vs. $2(230) = 460$.

Time = 5.10 (sec) , antiderivative size = 467, normalized size of antiderivative = 1.79

method	result
default	$\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(768B\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}+640A\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a\left(\cos^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-2176B\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a\left(\cos^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)$
parts	$A\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(128\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a\left(\cos^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}+489\sqrt{2}\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}+4a}{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)a\left(\cos^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-96\cos\left(\frac{dx}{2}+\frac{c}{2}\right)^3a^{\frac{7}{2}}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\right)$

input `int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNV
ERBOSE)`

output
$$\begin{aligned} & 1/480*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(768*B*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}*\cos(1/2*d*x+1/2*c)^8+640*A*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}*\cos(1/2*d*x+1/2*c)^6-2176*B*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}*\cos(1/2*d*x+1/2*c)^6+2445*A*2^{(1/2)}*\ln(2*(2*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*a)/\cos(1/2*d*x+1/2*c))*\cos(1/2*d*x+1/2*c)^4*a \\ & -4245*B*2^{(1/2)}*\ln(2*(2*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*a)/\cos(1/2*d*x+1/2*c))*\cos(1/2*d*x+1/2*c)^4*a-2560*A*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}*\cos(1/2*d*x+1/2*c)^4+5248*B*\cos(1/2*d*x+1/2*c)^4*2^{(1/2)}* \\ & a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}-435*A*\cos(1/2*d*x+1/2*c)^2*2^{(1/2)}* \\ & a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+555*B*\cos(1/2*d*x+1/2*c)^2*2^{(1/2)}* \\ & a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+30*A*a^{(1/2)}*2^{(1/2)}*(a*\sin(1/2*d*x \\ & +1/2*c)^2)^{(1/2)}-30*B*2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)})/\cos(\\ & 1/2*d*x+1/2*c)^3/a^{(7/2)}/\sin(1/2*d*x+1/2*c)/(a*\cos(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & /d \end{aligned}$$

3.115.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.03

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx =$$

$$\frac{15\sqrt{2}((163A-283B)\cos(dx+c)^3 + 3(163A-283B)\cos(dx+c)^2 + 3(163A-283B)\cos(dx+c) + 163A-283B)\sqrt{a}\log\left(\frac{-a\cos(dx+c)^2 + 2\sqrt{2}\sqrt{a\cos(dx+c)+a}\sqrt{a}\sin(dx+c) - 2a\cos(dx+c) - 3a}{(\cos(dx+c)^2 + 2\cos(dx+c) + 1)}\right) - 4(96B\cos(dx+c)^4 + 160(A-B)\cos(dx+c)^3 - 32(25A-49B)\cos(dx+c)^2 - 5(503A-911B)\cos(dx+c) - 1495A + 2671B)\sqrt{a\cos(dx+c)+a}\sin(dx+c)}{a^3d\cos(dx+c)^3 + 3a^3d\cos(dx+c)^2 + 3a^3d\cos(dx+c) + a^3d}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm m="fricas")`

output `-1/960*(15*sqrt(2)*((163*A - 283*B)*cos(d*x + c)^3 + 3*(163*A - 283*B)*cos(d*x + c)^2 + 3*(163*A - 283*B)*cos(d*x + c) + 163*A - 283*B)*sqrt(a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - 4*(96*B*cos(d*x + c)^4 + 160*(A - B)*cos(d*x + c)^3 - 32*(25*A - 49*B)*cos(d*x + c)^2 - 5*(503*A - 911*B)*cos(d*x + c) - 1495*A + 2671*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.115.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.115.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm m="maxima")`

output `Timed out`

3.115.8 Giac [A] (verification not implemented)

Time = 10.41 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.18

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{15\sqrt{2}(163A\sqrt{a}-283B\sqrt{a})\log(\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^3\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} - \frac{15\sqrt{2}(163A\sqrt{a}-283B\sqrt{a})\log(-\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^3\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm m="giac")`

output `1/960*(15*sqrt(2)*(163*A*sqrt(a) - 283*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - 15*sqrt(2)*(163*A*sqrt(a) - 283*B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) + 30*(29*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 37*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 27*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c) + 35*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c)^2 - 1)^2*a^3*sgn(cos(1/2*d*x + 1/2*c))) + 256*sqrt(2)*(6*B*a^(25/2)*sin(1/2*d*x + 1/2*c)^5 - 5*A*a^(25/2)*sin(1/2*d*x + 1/2*c)^3 + 5*B*a^(25/2)*sin(1/2*d*x + 1/2*c)^3 - 15*A*a^(25/2)*sin(1/2*d*x + 1/2*c) + 30*B*a^(25/2)*sin(1/2*d*x + 1/2*c))/(a^15*sgn(cos(1/2*d*x + 1/2*c))))/d`

3.115.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^4(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2), x)`output `int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2), x)`

3.116
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

3.116.1 Optimal result 1156
 3.116.2 Mathematica [A] (verified) 1157
 3.116.3 Rubi [A] (verified) 1157
 3.116.4 Maple [B] (verified) 1162
 3.116.5 Fracas [A] (verification not implemented) 1162
 3.116.6 Sympy [F(-1)] 1163
 3.116.7 Maxima [F(-1)] 1163
 3.116.8 Giac [A] (verification not implemented) 1164
 3.116.9 Mupad [F(-1)] 1164

3.116.1 Optimal result

Integrand size = 33, antiderivative size = 216

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = -\frac{(75A-163B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A-B) \cos^3(c+dx) \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{(9A-17B) \cos^2(c+dx) \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}} + \frac{(93A-197B) \sin(c+dx)}{24a^2d\sqrt{a+a \cos(c+dx)}} - \frac{(39A-95B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{48a^3d}$$

```
output 1/4*(A-B)*cos(d*x+c)^3*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+1/16*(9*A-17*B)
*cos(d*x+c)^2*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)-1/32*(75*A-163*B)*arct
anh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/
2)+1/24*(93*A-197*B)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)-1/48*(39*A-95
*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/a^3/d
```

3.116.2 Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.54

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{-6(75A-163B)\operatorname{arctanh}(\sin(\frac{1}{2}(c+dx)))\cos^3(\frac{1}{2}(c+dx)) + (19$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `(-6*(75*A - 163*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + (195*A - 379*B + (255*A - 479*B)*Cos[c + d*x] + 16*(3*A - 5*B)*Cos[2*(c + d*x)] + 8*B*Cos[3*(c + d*x)])*Tan[(c + d*x)/2]/(48*a*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.116.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3447, 3042, 3502, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx \\ & \quad \downarrow \text{3456} \\ & \frac{\int \frac{\cos^2(c+dx)(6a(A-B)-a(3A-11B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\cos^2(c+dx)(6a(A-B)-a(3A-11B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \end{aligned}$$

3.116. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2 (6a(A-B) - a(3A-11B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^3(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \downarrow \text{3456} \\
& \frac{\int \frac{\cos(c+dx)(4a^2(9A-17B) - a^2(39A-95B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a(9A-17B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{27} \\
& \downarrow \text{27} \\
& \frac{\int \frac{\cos(c+dx)(4a^2(9A-17B) - a^2(39A-95B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a(9A-17B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{3042} \\
& \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4a^2(9A-17B) - a^2(39A-95B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a(9A-17B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{3447} \\
& \downarrow \text{3447} \\
& \frac{\int \frac{4a^2(9A-17B)\cos(c+dx) - a^2(39A-95B)\cos^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a(9A-17B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{3042} \\
& \downarrow \text{3042} \\
& \frac{\int \frac{4a^2(9A-17B)\sin(c+dx+\frac{\pi}{2}) - a^2(39A-95B)\sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a(9A-17B)\sin(c+dx)\cos^2(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^3(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{3502} \\
& \downarrow \text{3502}
\end{aligned}$$

3.116. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{2 \int \frac{a^3(39A-95B) - 2a^3(93A-197B) \cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx - \frac{2a(39A-95B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{4a^2} + \frac{a(9A-17B) \sin(c+dx) \cos^2(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{4d(a \cos(c+dx)+a)^{5/2}} (A-B) \sin(c+dx) \cos^3(c+dx)$$

↓ 27

$$\frac{\int \frac{a^3(39A-95B) - 2a^3(93A-197B) \cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx - \frac{2a(39A-95B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{4a^2} + \frac{a(9A-17B) \sin(c+dx) \cos^2(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{4d(a \cos(c+dx)+a)^{5/2}} (A-B) \sin(c+dx) \cos^3(c+dx)$$

↓ 3042

$$\frac{\int \frac{a^3(39A-95B) - 2a^3(93A-197B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx - \frac{2a(39A-95B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{4a^2} + \frac{a(9A-17B) \sin(c+dx) \cos^2(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{4d(a \cos(c+dx)+a)^{5/2}} (A-B) \sin(c+dx) \cos^3(c+dx)$$

↓ 3230

$$\frac{3a^3(75A-163B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{4a^3(93A-197B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(39A-95B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{4a^2} + \frac{a(9A-17B) \sin(c+dx) \cos^2(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{4d(a \cos(c+dx)+a)^{5/2}} (A-B) \sin(c+dx) \cos^3(c+dx)$$

↓ 3042

$$\frac{3a^3(75A-163B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx - \frac{4a^3(93A-197B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(39A-95B) \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{3d}}{4a^2} + \frac{a(9A-17B) \sin(c+dx) \cos^2(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{4d(a \cos(c+dx)+a)^{5/2}} (A-B) \sin(c+dx) \cos^3(c+dx)$$

↓ 3128

3.116. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & -\frac{6a^3(75A-163B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{3a} - \frac{4a^3(93A-197B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(39A-95B) \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} + \frac{a(9A-17B) \sin(c+dx)}{2d(a \cos(c+dx)+a)} \\
 & \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \frac{8a^2}{5/2} \\
 & \quad \downarrow \text{219} \\
 & -\frac{3\sqrt{2}a^{5/2}(75A-163B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{4a^3(93A-197B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{2a(39A-95B) \sin(c+dx)\sqrt{a \cos(c+dx)+a}}{3d} + \frac{a(9A-17B) \sin(c+dx)}{2d(a \cos(c+dx)+a)} \\
 & \frac{(A-B) \sin(c+dx) \cos^3(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \frac{8a^2}{5/2}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `((A - B)*Cos[c + d*x]^3*Sin[c + d*x]/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (a*(9*A - 17*B)*Cos[c + d*x]^2*Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2)) + ((-2*a*(39*A - 95*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) - ((3*Sqrt[2]*a^(5/2)*(75*A - 163*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d - (4*a^3*(93*A - 197*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(4*a^2))/(8*a^2)`

3.116.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.116. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e
+ f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]`

3.116.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 396 vs. 2(189) = 378.

Time = 4.46 (sec) , antiderivative size = 397, normalized size of antiderivative = 1.84

method	result
default	$\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(128B\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a\left(\cos^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-225A\sqrt{2}\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+4a}}{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)\left(\cos^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a\right)$
parts	$\frac{A\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(-75\sqrt{2}\ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+4a}}{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)a\left(\cos^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+64\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{a\left(\cos^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)+2}{32\cos\left(\frac{dx}{2}+\frac{c}{2}\right)^3a^{\frac{7}{2}}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{a\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}d$

input `int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNV
ERBOSE)`

output `1/96/cos(1/2*d*x+1/2*c)^3*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(128*B*2^(1/2)*(a
*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*cos(1/2*d*x+1/2*c)^6-225*A*2^(1/2)*ln
(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(
1/2*d*x+1/2*c)^4*a+489*B*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(
1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^4*a+192*A*2^(1/2)*(a*sin
1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*cos(1/2*d*x+1/2*c)^4-512*B*cos(1/2*d*x+1/
2*c)^4*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+63*A*cos(1/2*d*x+1/
2*c)^2*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-87*B*cos(1/2*d*x+1/2*
c)^2*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-6*A*a^(1/2)*2^(1/2)*(a
*sin(1/2*d*x+1/2*c)^2)^(1/2)+6*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(
1/2))/a^(7/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.116.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.18

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx =$$

$$3\sqrt{2}((75A-163B)\cos(dx+c)^3+3(75A-163B)\cos(dx+c)^2+3(75A-163B)\cos(dx+c)+75)$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `-1/192*(3*sqrt(2)*((75*A - 163*B)*cos(d*x + c)^3 + 3*(75*A - 163*B)*cos(d*x + c)^2 + 3*(75*A - 163*B)*cos(d*x + c) + 75*A - 163*B)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - 4*(32*B*cos(d*x + c)^3 + 32*(3*A - 5*B)*cos(d*x + c)^2 + (255*A - 503*B)*cos(d*x + c) + 147*A - 299*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.116.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.116.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.116.8 Giac [A] (verification not implemented)

Time = 5.93 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.22

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{3\sqrt{2}(75A\sqrt{a}-163B\sqrt{a})\log(\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^3\operatorname{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} - \frac{3\sqrt{2}(75A\sqrt{a}-163B\sqrt{a})\log(-\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^3\operatorname{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} + \frac{6(21\sqrt{2}A\sin(\frac{1}{2}dx+\frac{1}{2}c)^3-29\sqrt{2}B\sin(\frac{1}{2}dx+\frac{1}{2}c))}{(\sin(\frac{1}{2}dx+\frac{1}{2}c))^2-1} a^{5/2} \operatorname{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c)) + 128\sqrt{2}(2B^2a^{13/2}\sin(\frac{1}{2}dx+\frac{1}{2}c)^3-3Aa^{13/2}\sin(\frac{1}{2}dx+\frac{1}{2}c)+6B^2a^{13/2}\sin(\frac{1}{2}dx+\frac{1}{2}c))/(a^9\operatorname{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c)))$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `-1/192*(3*sqrt(2)*(75*A*sqrt(a) - 163*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - 3*sqrt(2)*(75*A*sqrt(a) - 163*B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) + 6*(21*sqrt(2)*A*sin(1/2*d*x + 1/2*c)^3 - 29*sqrt(2)*B*sin(1/2*d*x + 1/2*c)^3 - 19*sqrt(2)*A*sin(1/2*d*x + 1/2*c) + 27*sqrt(2)*B*sin(1/2*d*x + 1/2*c))/(sin(1/2*d*x + 1/2*c)^2 - 1)^2*a^(5/2)*sgn(cos(1/2*d*x + 1/2*c))) + 128*sqrt(2)*(2*B*a^(13/2)*sin(1/2*d*x + 1/2*c)^3 - 3*A*a^(13/2)*sin(1/2*d*x + 1/2*c) + 6*B*a^(13/2)*sin(1/2*d*x + 1/2*c))/(a^9*sgn(cos(1/2*d*x + 1/2*c)))`
/d

3.116.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^3(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2), x)`

$$3.117 \quad \int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

3.117.1 Optimal result 1165
 3.117.2 Mathematica [A] (verified) 1165
 3.117.3 Rubi [A] (verified) 1166
 3.117.4 Maple [B] (verified) 1169
 3.117.5 Fricas [A] (verification not implemented) 1170
 3.117.6 Sympy [F(-1)] 1171
 3.117.7 Maxima [F(-1)] 1171
 3.117.8 Giac [A] (verification not implemented) 1171
 3.117.9 Mupad [F(-1)] 1172

3.117.1 Optimal result

Integrand size = 33, antiderivative size = 169

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{(19A-75B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A-B) \cos^2(c+dx) \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} - \frac{(5A-13B) \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}} - \frac{(A-9B) \sin(c+dx)}{4a^2d\sqrt{a+a \cos(c+dx)}}$$

output `1/4*(A-B)*cos(d*x+c)^2*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)-1/16*(5*A-13*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+1/32*(19*A-75*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)-1/4*(A-9*B)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)`

3.117.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.59

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{2(19A-75B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) \cos^3\left(\frac{1}{2}(c+dx)\right) + (-9A)}{16ad(a(1$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

3.117. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

output $(2*(19*A - 75*B)*\text{ArcTanh}[\text{Sin}[(c + d*x)/2]]*\text{Cos}[(c + d*x)/2]^3 + (-9*A + 65*B + (-13*A + 85*B)*\text{Cos}[c + d*x] + 16*B*\text{Cos}[2*(c + d*x)])*\text{Tan}[(c + d*x)/2])/ (16*a*d*(a*(1 + \text{Cos}[c + d*x]))^(3/2))$

3.117.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3456, 27, 3042, 3447, 3042, 3498, 27, 3042, 3230, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^2(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 3456

$$\frac{\int \frac{\cos(c+dx)(4a(A-B)-a(A-9B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 27

$$\frac{\int \frac{\cos(c+dx)(4a(A-B)-a(A-9B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(4a(A-B)-a(A-9B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3447

$$\frac{\int \frac{4a(A-B)\cos(c+dx)-a(A-9B)\cos^2(c+dx)}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{4a(A-B)\sin(c+dx+\frac{\pi}{2})-a(A-9B)\sin(c+dx+\frac{\pi}{2})^2}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

3.117. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{array}{c}
\downarrow \text{3498} \\
\frac{\int \frac{3a^2(5A-13B)-4a^2(A-9B)\cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{8a^2} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
\downarrow \text{27} \\
\frac{\int \frac{3a^2(5A-13B)-4a^2(A-9B)\cos(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{8a^2} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
\downarrow \text{3042} \\
\frac{\int \frac{3a^2(5A-13B)-4a^2(A-9B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{8a^2} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\cos^2(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
\downarrow \text{3230} \\
\frac{a^2(19A-75B)\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx - \frac{8a^2(A-9B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
\frac{8a^2}{(A-B)\sin(c+dx)\cos^2(c+dx)} \\
\frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow \text{3042}} \\
\frac{a^2(19A-75B)\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx - \frac{8a^2(A-9B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
\frac{8a^2}{(A-B)\sin(c+dx)\cos^2(c+dx)} \\
\frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow \text{3128}} \\
\frac{2a^2(19A-75B)\int \frac{1}{2a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} - \frac{8a^2(A-9B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
\frac{8a^2}{(A-B)\sin(c+dx)\cos^2(c+dx)} \\
\frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow \text{219}}
\end{array}$$

3.117. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{\frac{\sqrt{2}a^{3/2}(19A-75B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{8a^2(A-9B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}}{4a^2} + \frac{8a^2}{(A-B)\sin(c+dx)\cos^2(c+dx)} \frac{1}{4d(a\cos(c+dx)+a)^{5/2}}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `((A - B)*Cos[c + d*x]^2*Sin[c + d*x]/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(5*A - 13*B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^(3/2)) + ((Sqrt[2]*a^(3/2)*(19*A - 75*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d - (8*a^2*(A - 9*B)*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2))/(8*a^2)`

3.117.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3230 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(b*(m + 1)) Int[(a + b*Sin[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3498 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b - a*B + b*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m/(a*f*(2*m + 1)), x] + Simp[1/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]`

3.117.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(146) = 292.

Time = 4.47 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.93

method	result
default	$\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(19A\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^a - 75B\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^a \right)$
parts	$\frac{A\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(19\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 13 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{2}\sqrt{a+2\sqrt{a}} \right)}{32 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)^3 a^{\frac{7}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d}$

3.117.
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

input `int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNV
ERBOSE)`

output `1/32*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(19*A*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^4*a-75*B*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^4*a+64*B*cos(1/2*d*x+1/2*c)^4*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-13*A*cos(1/2*d*x+1/2*c)^2*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+21*B*cos(1/2*d*x+1/2*c)^2*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*A*a^(1/2)*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-2*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/cos(1/2*d*x+1/2*c)^3/a^(7/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d`

3.117.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.40

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx =$$

$$\frac{\sqrt{2}((19A-75B)\cos(dx+c)^3+3(19A-75B)\cos(dx+c)^2+3(19A-75B)\cos(dx+c)+19A-75B)}{(a+a\cos(dx+c))^3}$$

64

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm
m="fricas")`

output `-1/64*(sqrt(2)*((19*A - 75*B)*cos(d*x + c)^3 + 3*(19*A - 75*B)*cos(d*x + c)^2 + 3*(19*A - 75*B)*cos(d*x + c) + 19*A - 75*B)*sqrt(a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) - 4*(32*B*cos(d*x + c)^2 - (13*A - 85*B)*cos(d*x + c) - 9*A + 49*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.117.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.117.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.117.8 Giac [A] (verification not implemented)

Time = 3.20 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.40

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{5/2}} dx = \frac{128 \sqrt{2} B \sin(\frac{1}{2} dx + \frac{1}{2} c)}{a^{\frac{5}{2}} \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \frac{\sqrt{2}(19 A \sqrt{a} - 75 B \sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^3 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(19 A \sqrt{a} + 75 B \sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) - 1)}{a^3 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output $\frac{1}{64} * (128 * \sqrt{2} * B * \sin(1/2 * d * x + 1/2 * c) / (a^{5/2} * \operatorname{sgn}(\cos(1/2 * d * x + 1/2 * c))) + \sqrt{2} * (19 * A * \sqrt{a} - 75 * B * \sqrt{a}) * \log(\sin(1/2 * d * x + 1/2 * c) + 1) / (a^3 * \operatorname{sgn}(\cos(1/2 * d * x + 1/2 * c))) - \sqrt{2} * (19 * A * \sqrt{a} - 75 * B * \sqrt{a}) * \log(-\sin(1/2 * d * x + 1/2 * c) + 1) / (a^3 * \operatorname{sgn}(\cos(1/2 * d * x + 1/2 * c))) + 2 * (13 * \sqrt{2} * A * \sqrt{a} * \sin(1/2 * d * x + 1/2 * c)^3 - 21 * \sqrt{2} * B * \sqrt{a} * \sin(1/2 * d * x + 1/2 * c)^3 - 11 * \sqrt{2} * A * \sqrt{a} * \sin(1/2 * d * x + 1/2 * c) + 19 * \sqrt{2} * B * \sqrt{a} * \sin(1/2 * d * x + 1/2 * c)) / ((\sin(1/2 * d * x + 1/2 * c)^2 - 1)^2 * a^3 * \operatorname{sgn}(\cos(1/2 * d * x + 1/2 * c)))) / d$

3.117.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^2 (A + B \cos(c + dx))}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2), x)`

3.118 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

3.118.1 Optimal result 1173
 3.118.2 Mathematica [A] (verified) 1173
 3.118.3 Rubi [A] (verified) 1174
 3.118.4 Maple [B] (verified) 1176
 3.118.5 Fricas [B] (verification not implemented) 1177
 3.118.6 Sympy [F(-1)] 1177
 3.118.7 Maxima [F(-1)] 1178
 3.118.8 Giac [A] (verification not implemented) 1178
 3.118.9 Mupad [F(-1)] 1179

3.118.1 Optimal result

Integrand size = 31, antiderivative size = 126

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{(5A+19B)\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} - \frac{(A-B) \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{(5A-13B) \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}}$$

output `-1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+1/16*(5*A-13*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+1/32*(5*A+19*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)`

3.118.2 Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.69

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{2(5A+19B)\operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c+dx)\right)\right) \cos^3\left(\frac{1}{2}(c+dx)\right) + (A-9B)}{16ad(a(1+\cos(c+dx)))}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `(2*(5*A + 19*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + (A - 9*B + (5*A - 13*B)*Cos[c + d*x])*Tan[(c + d*x)/2])/(16*a*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.118. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

3.118.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3498, 27, 3042, 3229, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a\cos(c+dx)+a)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx \\
 & \quad \downarrow \text{3498} \\
 & -\frac{\int -\frac{5a(A-B)+8aB\cos(c+dx)}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{5a(A-B)+8aB\cos(c+dx)}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{5a(A-B)+8aB\sin(c+dx+\frac{\pi}{2})}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{3229} \\
 & \frac{\frac{1}{4}(5A+19B) \int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx + \frac{a(5A-13B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.118. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{\frac{1}{4}(5A + 19B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{a(5A-13B) \sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}}$$

↓ 3128

$$\frac{\frac{a(5A-13B) \sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{(5A+19B) \int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{8a^2}}{8a^2} - \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}}$$

↓ 219

$$\frac{\frac{(5A+19B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}\sqrt{ad}} + \frac{a(5A-13B) \sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `-1/4*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^(5/2)) + (((5*A + 19*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*Sqrt[a]*d) + (a*(5*A - 13*B)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2)`

3.118.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

```
rule 3229 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*
x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) I
nt[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] &&
NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3498 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b - a*
B + b*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[1
/(a^2*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[a*A*(m + 1) + m*(b
*B - a*C) + b*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && EqQ[a^2 - b^2, 0]
```

3.118.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 291 vs. 2(107) = 214.

Time = 4.32 (sec) , antiderivative size = 292, normalized size of antiderivative = 2.32

method	result
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(5A\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^a + 19B\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^a \right)}{32 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}$
parts	$\frac{A\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(5\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4a}}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right) a \left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 5 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \sqrt{2}\sqrt{a} - 2\sqrt{2} \right)}{32 \cos\left(\frac{dx}{2} + \frac{c}{2}\right)^3 a^{\frac{7}{2}} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{a\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} d$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2), x, method=_RETURNVER
BOSE)
```

```
output 1/32/cos(1/2*d*x+1/2*c)^3*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(5*A*2^(1/2)*ln(2
*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/
2*d*x+1/2*c)^4*a+19*B*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/
2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^4*a+5*A*cos(1/2*d*x+1/2*c)^
2*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-13*B*cos(1/2*d*x+1/2*c)^2
*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)-2*A*a^(1/2)*2^(1/2)*(a*sin
(1/2*d*x+1/2*c)^2)^(1/2)+2*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2
))/a^(7/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.118.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. $2(107) = 214$.

Time = 0.31 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.77

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{\sqrt{2}((5A+19B)\cos(dx+c)^3 + 3(5A+19B)\cos(dx+c)^2 + 3(5A+19B)\cos(dx+c) + 5A+19B)\sqrt{a}\log(-a\cos(dx+c)^2 - 2\sqrt{2}\sqrt{a\cos(dx+c)+a}\sqrt{a}\sin(dx+c) - 2a\cos(dx+c) - 3a)/(\cos(dx+c)^2 + 2\cos(dx+c) + 1) + 4((5A-13B)\cos(dx+c) + A - 9B)\sqrt{a\cos(dx+c)+a}\sin(dx+c))/(a^3d\cos(dx+c)^3 + 3a^3d\cos(dx+c)^2 + 3a^3d\cos(dx+c) + a^3d)}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm=
"fracas")
```

```
output 1/64*(sqrt(2)*((5*A + 19*B)*cos(d*x + c)^3 + 3*(5*A + 19*B)*cos(d*x + c)^2
+ 3*(5*A + 19*B)*cos(d*x + c) + 5*A + 19*B)*sqrt(a)*log(-(a*cos(d*x + c)^
2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x
+ c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) + 4*((5*A - 13*B)*cos(d
*x + c) + A - 9*B)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c))/(a^3*d*cos(d*x +
c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)
```

3.118.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)
```

```
output Timed out
```

3.118. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

3.118.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.118.8 Giac [A] (verification not implemented)

Time = 1.38 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.63

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{\sqrt{2}(5A\sqrt{a}+19B\sqrt{a})\log(\sin(\frac{1}{2}dx+\frac{1}{2}c)+1)}{a^3\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))} - \frac{\sqrt{2}(5A\sqrt{a}+19B\sqrt{a})\log(-\sin(\frac{1}{2}dx+\frac{1}{2}c))}{a^3\text{sgn}(\cos(\frac{1}{2}dx+\frac{1}{2}c))}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `1/64*(sqrt(2)*(5*A*sqrt(a) + 19*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(5*A*sqrt(a) + 19*B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - 2*(5*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 13*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 3*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c) + 11*sqrt(2)*B*sqrt(a)*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c)^2 - 1)^2*a^3*sgn(cos(1/2*d*x + 1/2*c)))/d`

3.118.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`output `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2), x)`

$$3.119 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

3.119.1 Optimal result 1180
 3.119.2 Mathematica [A] (verified) 1180
 3.119.3 Rubi [A] (verified) 1181
 3.119.4 Maple [B] (verified) 1183
 3.119.5 Fracas [B] (verification not implemented) 1183
 3.119.6 Sympy [F] 1184
 3.119.7 Maxima [F(-1)] 1184
 3.119.8 Giac [A] (verification not implemented) 1184
 3.119.9 Mupad [F(-1)] 1185

3.119.1 Optimal result

Integrand size = 25, antiderivative size = 126

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{(3A + 5B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A - B) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}} + \frac{(3A + 5B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}}$$

output `1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+1/16*(3*A+5*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+1/32*(3*A+5*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)`

3.119.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.63

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{4(3A + 5B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) \cos^5\left(\frac{1}{2}(c + dx)\right) + (7A + B + (3A + 5B) \cos(c + dx)) \sin(c + dx)}{16d(a(1 + \cos(c + dx)))^{5/2}}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^(5/2),x]`

output `(4*(3*A + 5*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 + (7*A + B + (3*A + 5*B)*Cos[c + d*x])*Sin[c + d*x])/(16*d*(a*(1 + Cos[c + d*x]))^(5/2))`

3.119. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.119.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {3042, 3229, 3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a \cos(c + dx) + a)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{5/2}} dx \\
 & \quad \downarrow \text{3229} \\
 & \frac{(3A + 5B) \int \frac{1}{(\cos(c+dx)a+a)^{3/2}} dx}{8a} + \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(3A + 5B) \int \frac{1}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a} + \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3129} \\
 & \frac{(3A + 5B) \left(\frac{\int \frac{1}{\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a} + \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(3A + 5B) \left(\frac{\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a} + \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \downarrow \text{3128} \\
 & \frac{(3A + 5B) \left(\frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{\int \frac{1}{2a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{2ad} \right)}{8a} + \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

3.119. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

$$\frac{(3A + 5B) \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a} + \frac{(A - B) \sin(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}}$$

input `Int[(A + B*Cos[c + d*x])/(a + a*Cos[c + d*x])^(5/2), x]`

output `((A - B)*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + ((3*A + 5*B)*(ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) + Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2))))/(8*a)`

3.119.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3229 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(a*f*(2*m + 1))), x] + Simp[(a*d*m + b*c*(m + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)]`

3.119.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 291 vs. 2(107) = 214.

Time = 4.24 (sec) , antiderivative size = 292, normalized size of antiderivative = 2.32

method	result
default	$\frac{\sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(3A\sqrt{2} \ln \left(\frac{4\sqrt{a} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 4a}}{\cos \left(\frac{dx}{2} + \frac{c}{2} \right)} \right) \left(\cos^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) a + 5B\sqrt{2} \ln \left(\frac{4\sqrt{a} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 4a}}{\cos \left(\frac{dx}{2} + \frac{c}{2} \right)} \right) \left(\cos^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 32 \cos \left(\frac{dx}{2} + \frac{c}{2} \right)}{\dots}$
parts	$\frac{A \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \left(3\sqrt{2} \ln \left(\frac{4\sqrt{a} \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 4a}}{\cos \left(\frac{dx}{2} + \frac{c}{2} \right)} \right) a \left(\cos^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 3 \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \sqrt{a \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} \sqrt{2} \sqrt{a} + 2\sqrt{2} \right)}{32a^{\frac{7}{2}} \cos \left(\frac{dx}{2} + \frac{c}{2} \right)^3 \sin \left(\frac{dx}{2} + \frac{c}{2} \right) \sqrt{a \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)} d$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/32/cos(1/2*d*x+1/2*c)^3*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(3*A*2^(1/2)*ln(2
*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/
2*d*x+1/2*c)^4*a+5*B*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2
)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^4*a+3*A*cos(1/2*d*x+1/2*c)^2
*2^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+5*B*cos(1/2*d*x+1/2*c)^2*2
^(1/2)*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*A*a^(1/2)*2^(1/2)*(a*sin(1
/2*d*x+1/2*c)^2)^(1/2)-2*B*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))
/a^(7/2)/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d
```

3.119.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. 2(107) = 214.

Time = 0.31 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.77

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2}((3A + 5B) \cos(dx + c)^3 + 3(3A + 5B) \cos(dx + c)^2 + 3(3A + 5B) \cos(dx + c) + 3A)}{(a + a \cos(c + dx))^{5/2}}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="fricas")
```

output $1/64*(\text{sqrt}(2)*((3*A + 5*B)*\cos(d*x + c)^3 + 3*(3*A + 5*B)*\cos(d*x + c)^2 + 3*(3*A + 5*B)*\cos(d*x + c) + 3*A + 5*B)*\text{sqrt}(a)*\log(-(a*\cos(d*x + c))^2 - 2*\text{sqrt}(2)*\text{sqrt}(a*\cos(d*x + c) + a)*\text{sqrt}(a)*\sin(d*x + c) - 2*a*\cos(d*x + c) - 3*a)/(\cos(d*x + c)^2 + 2*\cos(d*x + c) + 1)) + 4*((3*A + 5*B)*\cos(d*x + c) + 7*A + B)*\text{sqrt}(a*\cos(d*x + c) + a)*\sin(d*x + c)/(a^3*d*\cos(d*x + c)^3 + 3*a^3*d*\cos(d*x + c)^2 + 3*a^3*d*\cos(d*x + c) + a^3*d)$

3.119.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{(a(\cos(c + dx) + 1))^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))/(a*(cos(c + d*x) + 1))**(5/2), x)`

3.119.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.119.8 Giac [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.54

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2}(3A\sqrt{a} + 5B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^3 \text{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(3A\sqrt{a} + 5B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^3 \text{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{2(3\sqrt{a})}{a^3} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `1/64*(sqrt(2)*(3*A*sqrt(a) + 5*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(3*A*sqrt(a) + 5*B*sqrt(a))*log(-sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - 2*(3*sqrt(2)*A*sin(1/2*d*x + 1/2*c)^3 + 5*sqrt(2)*B*sin(1/2*d*x + 1/2*c)^3 - 5*sqrt(2)*A*sin(1/2*d*x + 1/2*c) - 3*sqrt(2)*B*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c))^2 - 1)^2*a^(5/2)*sgn(cos(1/2*d*x + 1/2*c)))/d`

3.119.9 Mupad [**F(-1)**]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))/(a + a*cos(c + d*x))^(5/2), x)`

3.120 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.120.1 Optimal result 1186
 3.120.2 Mathematica [A] (verified) 1186
 3.120.3 Rubi [A] (verified) 1187
 3.120.4 Maple [B] (verified) 1190
 3.120.5 Fricas [B] (verification not implemented) 1191
 3.120.6 Sympy [F] 1192
 3.120.7 Maxima [B] (verification not implemented) 1192
 3.120.8 Giac [A] (verification not implemented) 1193
 3.120.9 Mupad [F(-1)] 1194

3.120.1 Optimal result

Integrand size = 31, antiderivative size = 164

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{5/2}d}$$

$$- \frac{(43A - 3B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d}$$

$$- \frac{(A - B) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}} - \frac{(11A - 3B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}}$$

output `2*A*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d-1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)-1/16*(11*A-3*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)-1/32*(43*A-3*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)`

3.120.2 Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.77

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{-2(43A - 3B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) \cos^3\left(\frac{1}{2}(c + dx)\right) + 64\sqrt{2}a^{5/2}d}{(a + a \cos(c + dx))^{5/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^(5/2),x]`

```
output (-2*(43*A - 3*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + 64*Sqrt[2]
*A*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + (-15*A + 7*B + (
-11*A + 3*B)*Cos[c + d*x])*Tan[(c + d*x)/2])/(16*a*d*(a*(1 + Cos[c + d*x])
)^(3/2))
```

3.120.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(8aA-3a(A-B)\cos(c+dx))\sec(c+dx)}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(8aA-3a(A-B)\cos(c+dx))\sec(c+dx)}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{8aA-3a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(32a^2A-a^2(11A-3B)\cos(c+dx))\sec(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx}{8a^2} - \frac{a(11A-3B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.120. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{64a^2 A \int \frac{1}{a - \frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right) - \frac{\sqrt{2}a^{3/2}(43A-3B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{a(11A-3B) \sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}}{4a^2} = \frac{(A-B) \sin(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

↓ 219

$$\frac{64a^{3/2} A \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right) - \frac{\sqrt{2}a^{3/2}(43A-3B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{a(11A-3B) \sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}}{4a^2} = \frac{(A-B) \sin(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + a*Cos[c + d*x])^(5/2), x]`

output `-1/4*((A - B)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^(5/2)) + (((64*a^(3/2)*A*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d - (Sqrt[2]*a^(3/2)*(43*A - 3*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])]/d)/(4*a^2) - (a*(11*A - 3*B)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2)`

3.120.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.120. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3464 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.120.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 429 vs. 2(139) = 278.

Time = 5.79 (sec) , antiderivative size = 430, normalized size of antiderivative = 2.62

method	result
default	$\frac{\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(16A\sqrt{2} \ln\left(-\frac{2\left(\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{a-2a}\right)}{2\cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}} \right) a\left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 16A\sqrt{2} \ln\left(\frac{2\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}{2\cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{2}} \right)}{\dots}$
parts	$\frac{A\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(43\sqrt{2} \ln\left(\frac{4\sqrt{a}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 4a}{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)} \right) a\left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 32 \ln\left(\frac{4\sqrt{2}a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + 4\sqrt{2}\sqrt{a\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{2\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{2}} \right)}{\dots}$

3.120.
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

input `int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{32}a^{-7/2}/\cos(1/2*d*x+1/2*c)^3*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(16*A*2^{(1/2)}*\ln(-2/(2*\cos(1/2*d*x+1/2*c)-2^{(1/2)})*(2^{(1/2)}*a*\cos(1/2*d*x+1/2*c)-2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}-2*a))*a*\cos(1/2*d*x+1/2*c)^4+16*A*2^{(1/2)}*\ln(2/(2*\cos(1/2*d*x+1/2*c)+2^{(1/2)})*(2^{(1/2)}*a*\cos(1/2*d*x+1/2*c)+2^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}+2*a))*a*\cos(1/2*d*x+1/2*c)^4-43*A*\ln(2*(a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+a)/\cos(1/2*d*x+1/2*c))*a*\cos(1/2*d*x+1/2*c)^4+3*B*\ln(2*(a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+a)/\cos(1/2*d*x+1/2*c))*\cos(1/2*d*x+1/2*c)^4*a-11*A*\cos(1/2*d*x+1/2*c)^2*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+3*B*\cos(1/2*d*x+1/2*c)^2*a^{(1/2)}*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}-2*A*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)}+2*B*(a*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^{(1/2)})/\sin(1/2*d*x+1/2*c)*2^{(1/2)}/(a*\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/d$$

3.120.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. $2(139) = 278$.

Time = 0.37 (sec) , antiderivative size = 339, normalized size of antiderivative = 2.07

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2}((43A - 3B) \cos(dx + c)^3 + 3(43A - 3B) \cos(dx + c)^2 + 3(43A - 3B) \cos(dx + c) + 43A - 3B)}{\dots}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output
$$-1/64*(\sqrt{2})*((43*A - 3*B)*\cos(d*x + c)^3 + 3*(43*A - 3*B)*\cos(d*x + c)^2 + 3*(43*A - 3*B)*\cos(d*x + c) + 43*A - 3*B)*\sqrt{a}*\log(-(a*\cos(d*x + c)^2 - 2*\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sqrt{a}*\sin(d*x + c) - 2*a*\cos(d*x + c) - 3*a)/(\cos(d*x + c)^2 + 2*\cos(d*x + c) + 1)) - 32*(A*\cos(d*x + c)^3 + 3*A*\cos(d*x + c)^2 + 3*A*\cos(d*x + c) + A)*\sqrt{a}*\log((a*\cos(d*x + c)^3 - 7*a*\cos(d*x + c)^2 - 4*\sqrt{a*\cos(d*x + c) + a}*\sqrt{a}*(\cos(d*x + c) - 2)*\sin(d*x + c) + 8*a)/(\cos(d*x + c)^3 + \cos(d*x + c)^2)) + 4*((11*A - 3*B)*\cos(d*x + c) + 15*A - 7*B)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/(a^3*d*\cos(d*x + c)^3 + 3*a^3*d*\cos(d*x + c)^2 + 3*a^3*d*\cos(d*x + c) + a^3*d)$$

$$3.120. \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

3.120.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a (\cos(c + dx) + 1))^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a*(cos(c + d*x) + 1))**(5/2), x)`

3.120.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84333 vs. $2(139) = 278$.

Time = 15.38 (sec) , antiderivative size = 84333, normalized size of antiderivative = 514.23

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output

```

1/32*(512*((2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d*x + 5/2*c) + cos(
5/2*d*x + 5/2*c)*sin(4*d*x + 4*c) + 2*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c
) + (2*cos(2*d*x + 2*c) + cos(d*x + c))*sin(5/2*d*x + 5/2*c) + cos(4*d*x +
4*c)*sin(5/2*d*x + 5/2*c) + 2*cos(3*d*x + 3*c)*sin(5/2*d*x + 5/2*c))*cos(
5*d*x + 5*c)^2 + 2560*(5*(2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d*x +
5/2*c) + cos(5/2*d*x + 5/2*c)*sin(5*d*x + 5*c) + 5*cos(5/2*d*x + 5/2*c)*s
in(4*d*x + 4*c) + 10*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) - (10*cos(2*d*x
+ 2*c) + 5*cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c) - cos(5*d*x + 5*c)*sin(
5/2*d*x + 5/2*c) - 5*cos(4*d*x + 4*c)*sin(5/2*d*x + 5/2*c) - 10*cos(3*d*x
+ 3*c)*sin(5/2*d*x + 5/2*c))*cos(8/5*arctan2(sin(5/2*d*x + 5/2*c), cos(5/2
*d*x + 5/2*c)))^2 + 10240*(5*(2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5/2*d
*x + 5/2*c) + cos(5/2*d*x + 5/2*c)*sin(5*d*x + 5*c) + 5*cos(5/2*d*x + 5/2*
c)*sin(4*d*x + 4*c) + 10*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) - (10*cos(2
*d*x + 2*c) + 5*cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c) - cos(5*d*x + 5*c)*
sin(5/2*d*x + 5/2*c) - 5*cos(4*d*x + 4*c)*sin(5/2*d*x + 5/2*c) - 10*cos(3*
d*x + 3*c)*sin(5/2*d*x + 5/2*c))*cos(6/5*arctan2(sin(5/2*d*x + 5/2*c), cos
(5/2*d*x + 5/2*c)))^2 + 10240*(5*(2*sin(2*d*x + 2*c) + sin(d*x + c))*cos(5
/2*d*x + 5/2*c) + cos(5/2*d*x + 5/2*c)*sin(5*d*x + 5*c) + 5*cos(5/2*d*x +
5/2*c)*sin(4*d*x + 4*c) + 10*cos(5/2*d*x + 5/2*c)*sin(3*d*x + 3*c) - (10*c
os(2*d*x + 2*c) + 5*cos(d*x + c) + 1)*sin(5/2*d*x + 5/2*c) - cos(5*d*x ...
    
```

3.120.8 Giac [A] (verification not implemented)

Time = 3.04 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.54

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{64 A \log\left(\frac{|-2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}{|2\sqrt{2}+4 \sin(\frac{1}{2} dx + \frac{1}{2} c)|}\right)}{a^{5/2} \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} + \frac{\sqrt{2}(43 A\sqrt{a}-3 B\sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c)+1)}{a^3 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(43 A\sqrt{a}-3 B\sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c)+1)}{a^3 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{2}{64d}$$

input

```

integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+a*cos(d*x+c))^(5/2),x, algorithm=
"giac")
    
```


output
$$\begin{aligned} & -1/64*(64*A*\log(\text{abs}(-2*\text{sqrt}(2) + 4*\sin(1/2*d*x + 1/2*c))/\text{abs}(2*\text{sqrt}(2) + 4 \\ & * \sin(1/2*d*x + 1/2*c)))/(a^{5/2}*\text{sgn}(\cos(1/2*d*x + 1/2*c))) + \text{sqrt}(2)*(43* \\ & A*\text{sqrt}(a) - 3*B*\text{sqrt}(a))*\log(\sin(1/2*d*x + 1/2*c) + 1)/(a^3*\text{sgn}(\cos(1/2*d* \\ & x + 1/2*c))) - \text{sqrt}(2)*(43*A*\text{sqrt}(a) - 3*B*\text{sqrt}(a))*\log(-\sin(1/2*d*x + 1/2 \\ & *c) + 1)/(a^3*\text{sgn}(\cos(1/2*d*x + 1/2*c))) - 2*(11*\text{sqrt}(2)*A*\sin(1/2*d*x + 1 \\ & /2*c)^3 - 3*\text{sqrt}(2)*B*\sin(1/2*d*x + 1/2*c)^3 - 13*\text{sqrt}(2)*A*\sin(1/2*d*x + \\ & 1/2*c) + 5*\text{sqrt}(2)*B*\sin(1/2*d*x + 1/2*c))/((\sin(1/2*d*x + 1/2*c)^2 - 1)^2 \\ & *a^{5/2}*\text{sgn}(\cos(1/2*d*x + 1/2*c)))/d \end{aligned}$$

3.120.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx) (a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + a*cos(c + d*x))^(5/2)), x)`

3.121 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.121.1 Optimal result 1195
 3.121.2 Mathematica [A] (verified) 1196
 3.121.3 Rubi [A] (verified) 1196
 3.121.4 Maple [B] (verified) 1201
 3.121.5 Fracas [B] (verification not implemented) 1202
 3.121.6 Sympy [F] 1202
 3.121.7 Maxima [F(-2)] 1203
 3.121.8 Giac [A] (verification not implemented) 1203
 3.121.9 Mupad [F(-1)] 1204

3.121.1 Optimal result

Integrand size = 33, antiderivative size = 207

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = -\frac{(5A - 2B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{5/2}d}$$

$$+ \frac{(115A - 43B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} - \frac{(A - B) \tan(c + dx)}{4d(a + a \cos(c + dx))^{5/2}}$$

$$- \frac{(15A - 7B) \tan(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}} + \frac{(35A - 11B) \tan(c + dx)}{16a^2d\sqrt{a + a \cos(c + dx)}}$$

```
output - (5*A-2*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d+1/
32*(115*A-43*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+c))^(1
/2))/a^(5/2)/d*2^(1/2)-1/4*(A-B)*tan(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)-1/16*
(15*A-7*B)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+1/16*(35*A-11*B)*tan(d*x+
c)/a^2/d/(a+a*cos(d*x+c))^(1/2)
```

3.121.2 Mathematica [A] (verified)

Time = 2.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.69

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{8(115A - 43B) \operatorname{arctanh}\left(\sin\left(\frac{1}{2}(c + dx)\right)\right) \cos^5\left(\frac{1}{2}(c + dx)\right) - 128\sqrt{a} \sin^5\left(\frac{1}{2}(c + dx)\right)}{(a + a \cos(c + dx))^{5/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^(5/2), x]`

output `(8*(115*A - 43*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 - 128*Sqrt[2]*(5*A - 2*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^5 + (67*A - 11*B + 10*(11*A - 3*B)*Cos[c + d*x] + (35*A - 11*B)*Cos[2*(c + d*x)])*Tan[c + d*x]/(32*d*(a*(1 + Cos[c + d*x]))^(5/2))`

3.121.3 Rubi [A] (verified)

Time = 1.46 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.08, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 25, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^2 (a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{5/2}} dx \\ & \quad \downarrow \text{3457} \\ & \frac{\int \frac{(2a(5A - B) - 5a(A - B) \cos(c + dx)) \sec^2(c + dx)}{2(\cos(c + dx)a + a)^{3/2}} dx}{4a^2} - \frac{(A - B) \tan(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(2a(5A - B) - 5a(A - B) \cos(c + dx)) \sec^2(c + dx)}{(\cos(c + dx)a + a)^{3/2}} dx}{8a^2} - \frac{(A - B) \tan(c + dx)}{4d(a \cos(c + dx) + a)^{5/2}} \end{aligned}$$

3.121. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{2a(5A-B) - 5a(A-B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2 (\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx - \frac{(A-B) \tan(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2a^2(35A-11B) - 3a^2(15A-7B) \cos(c+dx)}{2\sqrt{\cos(c+dx)a+a}} \sec^2(c+dx) dx}{8a^2} - \frac{a(15A-7B) \tan(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{(A-B) \tan(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(2a^2(35A-11B) - 3a^2(15A-7B) \cos(c+dx)) \sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(15A-7B) \tan(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{(A-B) \tan(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{2a^2(35A-11B) - 3a^2(15A-7B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2 \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{8a^2} - \frac{a(15A-7B) \tan(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{(A-B) \tan(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(16a^3(5A-2B) - a^3(35A-11B) \cos(c+dx)) \sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{2a^2(35A-11B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a(15A-7B) \tan(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3463} \\
& \frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} \frac{(A-B) \tan(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{25} \\
& \frac{2a^2(35A-11B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{(16a^3(5A-2B) - a^3(35A-11B) \cos(c+dx)) \sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(15A-7B) \tan(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{16a^3(5A-2B) - a^3(35A-11B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2a^2(35A-11B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{a(15A-7B) \tan(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} \frac{(A-B) \tan(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}}
\end{aligned}$$

3.121. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

↓ 3464

$$\frac{\frac{2a^2(35A-11B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{16a^2(5A-2B)\int\sqrt{\cos(c+dx)a+a}\sec(c+dx)dx - a^3(115A-43B)\int\frac{1}{\sqrt{\cos(c+dx)a+a}}dx}{4a^2} - \frac{a(15A-7B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}}(A-B)\tan(c+dx)$$

↓ 3042

$$\frac{\frac{2a^2(35A-11B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{16a^2(5A-2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})}dx - a^3(115A-43B)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}dx}{4a^2} - \frac{a(15A-7B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}}(A-B)\tan(c+dx)$$

↓ 3128

$$\frac{\frac{2a^2(35A-11B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{16a^2(5A-2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})}dx + \frac{2a^3(115A-43B)\int\frac{1}{2a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} - \frac{a(15A-7B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^3}}$$

$$\frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}}(A-B)\tan(c+dx)$$

↓ 219

$$\frac{\frac{2a^2(35A-11B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{16a^2(5A-2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})}dx - \frac{\sqrt{2}a^{5/2}(115A-43B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{a(15A-7B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}}(A-B)\tan(c+dx)$$

↓ 3252

$$\frac{\frac{2a^2(35A-11B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{32a^3(5A-2B)\int\frac{1}{a-\frac{a^2\sin^2(c+dx)}{\cos(c+dx)a+a}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \frac{\sqrt{2}a^{5/2}(115A-43B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{a(15A-7B)\tan(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{8a^2}{4d(a\cos(c+dx)+a)^{5/2}}(A-B)\tan(c+dx)$$

3.121. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

↓ 219

$$\frac{\frac{2a^2(35A-11B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{32a^{5/2}(5A-2B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{\sqrt{2}a^{5/2}(115A-43B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{a(15A-7B)\tan(c+dx)}{2d(a\cos(c+dx)+a)}$$

$$\frac{8a^2(A-B)\tan(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + a*Cos[c + d*x])^(5/2),x]`

output `-1/4*((A - B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(15*A - 7*B)*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^(3/2)) + (-(((32*a^(5/2)*(5*A - 2*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (Sqrt[2]*a^(5/2)*(115*A - 43*B)*ArcTanh[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[a + a*Cos[c + d*x]])])/d)/a) + (2*a^2*(35*A - 11*B)*Tan[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]))/(4*a^2))/(8*a^2)`

3.121.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.121. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3464 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.121.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 924 vs. $2(178) = 356$.

Time = 6.61 (sec) , antiderivative size = 925, normalized size of antiderivative = 4.47

method	result	size
parts	Expression too large to display	925
default	Expression too large to display	1122

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/16*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(230*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^6*a-16
0*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/
2))*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^6*a-160*
ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)
*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^6*a-115*2
^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2
*c))*a*cos(1/2*d*x+1/2*c)^4+70*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1
/2)*cos(1/2*d*x+1/2*c)^4+80*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a
*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*c
os(1/2*d*x+1/2*c)^4*a+80*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*c
os(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos
(1/2*d*x+1/2*c)^4*a-15*cos(1/2*d*x+1/2*c)^2*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)
*2^(1/2)*a^(1/2)-2*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2))/a^(7/2)
/cos(1/2*d*x+1/2*c)^3/(2*cos(1/2*d*x+1/2*c)+2^(1/2))/(2*cos(1/2*d*x+1/2*c)
-2^(1/2))/sin(1/2*d*x+1/2*c)/(a*cos(1/2*d*x+1/2*c)^2)^(1/2)/d+1/32*B/a^(7/
2)/cos(1/2*d*x+1/2*c)^3*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(16*2^(1/2)*ln(2/(2
*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^4*a+16*2^(1/2)*ln
(-2/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)...
```


3.121.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 404 vs. $2(178) = 356$.

Time = 0.37 (sec) , antiderivative size = 404, normalized size of antiderivative = 1.95

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\sqrt{2}((115A - 43B) \cos(dx + c)^4 + 3(115A - 43B) \cos(dx + c)^3 + 3(115A - 43B) \cos(dx + c)^2 + (115A - 43B) \cos(dx + c) + 115A - 43B)$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `-1/64*(sqrt(2)*((115*A - 43*B)*cos(d*x + c)^4 + 3*(115*A - 43*B)*cos(d*x + c)^3 + 3*(115*A - 43*B)*cos(d*x + c)^2 + (115*A - 43*B)*cos(d*x + c))*sqrt(a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x + c) + 1)) + 16*((5*A - 2*B)*cos(d*x + c)^4 + 3*(5*A - 2*B)*cos(d*x + c)^3 + 3*(5*A - 2*B)*cos(d*x + c)^2 + (5*A - 2*B)*cos(d*x + c))*sqrt(a)*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 - 4*sqrt(a*cos(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x + c)^2)) - 4*((35*A - 11*B)*cos(d*x + c)^2 + 5*(11*A - 3*B)*cos(d*x + c) + 16*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/(a^3*d*cos(d*x + c)^4 + 3*a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + a^3*d*cos(d*x + c))`

3.121.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a(\cos(c + dx) + 1))^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+a*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a*(cos(c + d*x) + 1))**(5/2), x)`

3.121.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(5/2),x, algorithm
m="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.121.8 Giac [A] (verification not implemented)

Time = 0.81 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.62

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2}(115 A \sqrt{a} - 43 B \sqrt{a}) \log(\sin(\frac{1}{2} dx + \frac{1}{2} c) + 1)}{a^3 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))} - \frac{\sqrt{2}(115 A \sqrt{a} - 43 B \sqrt{a}) \log(-\sin(\frac{1}{2} dx + \frac{1}{2} c))}{a^3 \operatorname{sgn}(\cos(\frac{1}{2} dx + \frac{1}{2} c))}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+a*cos(d*x+c))^(5/2),x, algorithm
m="giac")
```

```
output 1/64*(sqrt(2)*(115*A*sqrt(a) - 43*B*sqrt(a))*log(sin(1/2*d*x + 1/2*c) + 1)
/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - sqrt(2)*(115*A*sqrt(a) - 43*B*sqrt(a))*
log(-sin(1/2*d*x + 1/2*c) + 1)/(a^3*sgn(cos(1/2*d*x + 1/2*c))) - 32*(5*A -
2*B)*log(abs(1/2*sqrt(2) + sin(1/2*d*x + 1/2*c)))/(a^(5/2)*sgn(cos(1/2*d*
x + 1/2*c))) + 32*(5*A - 2*B)*log(abs(-1/2*sqrt(2) + sin(1/2*d*x + 1/2*c))
)/(a^(5/2)*sgn(cos(1/2*d*x + 1/2*c))) - 64*sqrt(2)*A*sin(1/2*d*x + 1/2*c)/
((2*sin(1/2*d*x + 1/2*c)^2 - 1)*a^(5/2)*sgn(cos(1/2*d*x + 1/2*c))) - 2*(19
*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c)^3 - 11*sqrt(2)*B*sqrt(a)*sin(1/2*d
*x + 1/2*c)^3 - 21*sqrt(2)*A*sqrt(a)*sin(1/2*d*x + 1/2*c) + 13*sqrt(2)*B*s
qrt(a)*sin(1/2*d*x + 1/2*c))/((sin(1/2*d*x + 1/2*c)^2 - 1)^2*a^3*sgn(cos(1
/2*d*x + 1/2*c))))/d
```

3.121.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^2 (a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^(5/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + a*cos(c + d*x))^(5/2)), x)`

3.122
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

3.122.1 Optimal result 1205
 3.122.2 Mathematica [A] (verified) 1206
 3.122.3 Rubi [A] (verified) 1206
 3.122.4 Maple [B] (warning: unable to verify) 1212
 3.122.5 Fricas [A] (verification not implemented) 1213
 3.122.6 Sympy [F(-1)] 1213
 3.122.7 Maxima [F(-2)] 1214
 3.122.8 Giac [F(-2)] 1214
 3.122.9 Mupad [F(-1)] 1214

3.122.1 Optimal result

Integrand size = 33, antiderivative size = 264

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{(39A - 20B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4a^{5/2}d} - \frac{(219A - 115B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} - \frac{7(9A - 5B) \tan(c + dx)}{16a^2d\sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec(c + dx) \tan(c + dx)}{4d(a + a \cos(c + dx))^{5/2}} - \frac{(19A - 11B) \sec(c + dx) \tan(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}} + \frac{(31A - 15B) \sec(c + dx) \tan(c + dx)}{16a^2d\sqrt{a + a \cos(c + dx)}}$$

```
output 1/4*(39*A-20*B)*arctanh(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)
/d-1/32*(219*A-115*B)*arctanh(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/(a+a*cos(d*x+
c))^(1/2))/a^(5/2)/d*2^(1/2)-1/4*(A-B)*sec(d*x+c)*tan(d*x+c)/d/(a+a*cos(d*
x+c))^(5/2)-1/16*(19*A-11*B)*sec(d*x+c)*tan(d*x+c)/a/d/(a+a*cos(d*x+c))^(3
/2)-7/16*(9*A-5*B)*tan(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)+1/16*(31*A-15*B
)*sec(d*x+c)*tan(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)
```

3.122.2 Mathematica [A] (verified)

Time = 4.02 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.67

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{-8(219A - 115B) \operatorname{arctanh}(\sin(\frac{1}{2}(c + dx))) \cos^3(\frac{1}{2}(c + dx)) + 32 \sqrt{2} (39A - 20B) \operatorname{arctanh}(\sqrt{2} \sin(\frac{1}{2}(c + dx))) \cos^3(\frac{1}{2}(c + dx)) + (-158A + 110B + (-269A + 169B) \cos(c + dx) + (-190A + 110B) \cos[2(c + dx)] - 63A \cos[3(c + dx)] + 35B \cos[3(c + dx)]) \sec^2(c + dx) \operatorname{arctanh}(\frac{1}{2}(c + dx))}{(64ad(a(1 + \cos(c + dx)))^{3/2})}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^(5/2),x]`

output `(-8*(219*A - 115*B)*ArcTanh[Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + 32*sqrt[2]*(39*A - 20*B)*ArcTanh[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[(c + d*x)/2]^3 + (-158*A + 110*B + (-269*A + 169*B)*Cos[c + d*x] + (-190*A + 110*B)*Cos[2*(c + d*x)] - 63*A*Cos[3*(c + d*x)] + 35*B*Cos[3*(c + d*x)])*Sec[c + d*x]^2*ArcTanh[(c + d*x)/2]/(64*a*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.122.3 Rubi [A] (verified)Time = 1.91 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.07, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 25, 3042, 3464, 3042, 3128, 219, 3252, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^3(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3 (a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2}} dx \\ & \quad \downarrow \text{3457} \\ & \frac{\int \frac{(4a(3A-B) - 7a(A-B) \cos(c+dx)) \sec^3(c+dx)}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(4a(3A-B) - 7a(A-B) \cos(c+dx)) \sec^3(c+dx)}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \end{aligned}$$

3.122. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

$$\begin{array}{c}
\downarrow \text{3042} \\
\int \frac{4a(3A-B) - 7a(A-B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^3 (\sin(c+dx + \frac{\pi}{2})a + a)^{3/2}} dx - \frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
\downarrow \text{3457} \\
\frac{\int \frac{(4a^2(31A-15B) - 5a^2(19A-11B) \cos(c+dx)) \sec^3(c+dx)}{2\sqrt{\cos(c+dx)a+a}} dx - \frac{a(19A-11B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}}{\frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} (A-B) \tan(c+dx) \sec(c+dx)} \\
\downarrow \text{27} \\
\frac{\int \frac{(4a^2(31A-15B) - 5a^2(19A-11B) \cos(c+dx)) \sec^3(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx - \frac{a(19A-11B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}}{\frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} (A-B) \tan(c+dx) \sec(c+dx)} \\
\downarrow \text{3042} \\
\frac{\int \frac{4a^2(31A-15B) - 5a^2(19A-11B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^3 \sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx - \frac{a(19A-11B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}}{\frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} (A-B) \tan(c+dx) \sec(c+dx)} \\
\downarrow \text{3463} \\
\frac{\int -\frac{2(14a^3(9A-5B) - 3a^3(31A-15B) \cos(c+dx)) \sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx + \frac{2a^2(31A-15B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx) + a}} - \frac{a(19A-11B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}}{\frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} (A-B) \tan(c+dx) \sec(c+dx)} \\
\downarrow \text{27} \\
\frac{\frac{2a^2(31A-15B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx) + a}} - \frac{\int \frac{(14a^3(9A-5B) - 3a^3(31A-15B) \cos(c+dx)) \sec^2(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{a}}{4a^2} - \frac{a(19A-11B) \tan(c+dx) \sec(c+dx)}{2d(a \cos(c+dx) + a)^{3/2}}}{\frac{8a^2}{4d(a \cos(c+dx) + a)^{5/2}} (A-B) \tan(c+dx) \sec(c+dx)} \\
\downarrow \text{3042}
\end{array}$$

3.122. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

$$\frac{2a^2(31A-15B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{14a^3(9A-5B)-3a^3(31A-15B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^2\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2} - \frac{a(19A-11B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3463

$$\frac{2a^2(31A-15B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int -\frac{(4a^4(39A-20B)-7a^4(9A-5B)\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{14a^3(9A-5B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{a(19A-11B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 25

$$\frac{2a^2(31A-15B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{14a^3(9A-5B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{(4a^4(39A-20B)-7a^4(9A-5B)\cos(c+dx))\sec(c+dx)}{\sqrt{\cos(c+dx)a+a}} dx}{a} - \frac{a(19A-11B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{2a^2(31A-15B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{14a^3(9A-5B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{4a^4(39A-20B)-7a^4(9A-5B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{a} - \frac{a(19A-11B)\tan(c+dx)\sec(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3464

$$\frac{2a^2(31A-15B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{14a^3(9A-5B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{4a^3(39A-20B)\int\sqrt{\cos(c+dx)a+a}\sec(c+dx)dx - a^4(219A-115B)\int\frac{1}{\sqrt{\cos(c+dx)a+a}}dx}{a} - a(\dots)$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

3.122. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{2a^2(31A-15B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{14a^3(9A-5B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{4a^3(39A-20B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} a+a}{\sin(c+dx+\frac{\pi}{2})} dx - a^4(219A-115B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} a+a} dx}{4a^2} - a(1)$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \quad 8a^2$$

↓ 3128

$$\frac{2a^2(31A-15B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{14a^3(9A-5B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{4a^3(39A-20B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} a+a}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2a^4(219A-115B) \int \frac{1}{2a-\frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{a}}{4a^2}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \quad 8a^2$$

↓ 219

$$\frac{2a^2(31A-15B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{14a^3(9A-5B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{4a^3(39A-20B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} a+a}{\sin(c+dx+\frac{\pi}{2})} dx - \frac{\sqrt{2}a^{7/2}(219A-115B) \operatorname{arctanh}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \quad 8a^2$$

↓ 3252

$$\frac{2a^2(31A-15B) \tan(c+dx) \sec(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{14a^3(9A-5B) \tan(c+dx)}{d\sqrt{a \cos(c+dx)+a}} - \frac{8a^4(39A-20B) \int \frac{1}{a-\frac{a^2 \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right) - \sqrt{2}a^{7/2}(219A-115B) \operatorname{arctan}\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{a \cos(c+dx)+a}}\right)}{a}}{4a^2}$$

$$\frac{(A-B) \tan(c+dx) \sec(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \quad 8a^2$$

↓ 219

3.122. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

$$\frac{2a^2(31A-15B)\tan(c+dx)\sec(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{14a^3(9A-5B)\tan(c+dx)}{d\sqrt{a\cos(c+dx)+a}} - \frac{8a^{7/2}(39A-20B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{\sqrt{2}a^{7/2}(219A-115B)\operatorname{arctanh}\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{a\cos(c+dx)+a}}\right)}{d}$$

$$\frac{(A-B)\tan(c+dx)\sec(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + a*Cos[c + d*x])^(5/2), x]`

output `-1/4*((A - B)*Sec[c + d*x]*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(19*A - 11*B)*Sec[c + d*x]*Tan[c + d*x])/(d*(a + a*Cos[c + d*x])^(3/2)) + ((2*a^2*(31*A - 15*B)*Sec[c + d*x]*Tan[c + d*x])/(d*sqrt[a + a*Cos[c + d*x]]) - (((8*a^(7/2)*(39*A - 20*B)*ArcTanh[(sqrt[a]*Sin[c + d*x])/sqrt[a + a*Cos[c + d*x]])]/d - (sqrt[2]*a^(7/2)*(219*A - 115*B)*ArcTanh[(sqrt[a]*Sin[c + d*x])/(sqrt[2]*sqrt[a + a*Cos[c + d*x]])]/d)/a) + (14*a^3*(9*A - 5*B)*Tan[c + d*x])/(d*sqrt[a + a*Cos[c + d*x]))/a)/(4*a^2)/(8*a^2)`

3.122.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/sqrt[a + b*Sin[c + d*x]])], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

3.122. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

rule 3252 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-2*(b/f) Subst[Int[1/(b*c + a*d - d*x^2), x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

rule 3464 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.122.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1437 vs. $2(229) = 458$.

Time = 7.58 (sec) , antiderivative size = 1438, normalized size of antiderivative = 5.45

method	result	size
parts	Expression too large to display	1438
default	Expression too large to display	1610

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/8*A*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*(876*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(
1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*cos(1/2*d*x+1/2*c)^8*a-62
4*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/
2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^8*a-624
*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)
*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^8*a-876*2
^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a)/cos(1/2*d*x+1/2
*c))*cos(1/2*d*x+1/2*c)^6*a+252*2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(
1/2)*cos(1/2*d*x+1/2*c)^6+624*ln(-4/(2*cos(1/2*d*x+1/2*c)-2^(1/2))*(2^(1/2)
)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)-2*a)
)*cos(1/2*d*x+1/2*c)^6*a+624*ln(4/(2*cos(1/2*d*x+1/2*c)+2^(1/2))*(2^(1/2)*
a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)+2*a))*
cos(1/2*d*x+1/2*c)^6*a+219*2^(1/2)*ln(2*(2*a^(1/2)*(a*sin(1/2*d*x+1/2*c)^2)
)^(1/2)+2*a)/cos(1/2*d*x+1/2*c))*a*cos(1/2*d*x+1/2*c)^4-188*2^(1/2)*(a*sin
(1/2*d*x+1/2*c)^2)^(1/2)*a^(1/2)*cos(1/2*d*x+1/2*c)^4-156*ln(-4/(2*cos(1/2
*d*x+1/2*c)-2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)-2^(1/2)*(a*sin(1/2*d*x+
1/2*c)^2)^(1/2)*a^(1/2)-2*a))*cos(1/2*d*x+1/2*c)^4*a-156*ln(4/(2*cos(1/2*d
*x+1/2*c)+2^(1/2))*(2^(1/2)*a*cos(1/2*d*x+1/2*c)+2^(1/2)*(a*sin(1/2*d*x+1/
2*c)^2)^(1/2)*a^(1/2)+2*a))*cos(1/2*d*x+1/2*c)^4*a+19*cos(1/2*d*x+1/2*c)^2
*(a*sin(1/2*d*x+1/2*c)^2)^(1/2)*2^(1/2)*a^(1/2)+2*2^(1/2)*(a*sin(1/2*d*...
```

3.122.5 Fricas [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 428, normalized size of antiderivative = 1.62

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\sqrt{2}((219A - 115B) \cos(dx + c)^5 + 3(219A - 115B) \cos(dx + c)^4 + 3(219A - 115B) \cos(dx + c)^3 +$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(5/2),x, algorithm
m="fricas")
```

```
output -1/64*(sqrt(2)*((219*A - 115*B)*cos(d*x + c)^5 + 3*(219*A - 115*B)*cos(d*x
+ c)^4 + 3*(219*A - 115*B)*cos(d*x + c)^3 + (219*A - 115*B)*cos(d*x + c)^
2)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqr
t(a)*sin(d*x + c) - 2*a*cos(d*x + c) - 3*a)/(cos(d*x + c)^2 + 2*cos(d*x +
c) + 1)) + 4*((39*A - 20*B)*cos(d*x + c)^5 + 3*(39*A - 20*B)*cos(d*x + c)^
4 + 3*(39*A - 20*B)*cos(d*x + c)^3 + (39*A - 20*B)*cos(d*x + c)^2)*sqrt(a)
*log((a*cos(d*x + c)^3 - 7*a*cos(d*x + c)^2 + 4*sqrt(a*cos(d*x + c) + a)*s
qrt(a)*(cos(d*x + c) - 2)*sin(d*x + c) + 8*a)/(cos(d*x + c)^3 + cos(d*x +
c)^2)) + 4*(7*(9*A - 5*B)*cos(d*x + c)^3 + 5*(19*A - 11*B)*cos(d*x + c)^2
+ 4*(5*A - 4*B)*cos(d*x + c) - 8*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)
)/(a^3*d*cos(d*x + c)^5 + 3*a^3*d*cos(d*x + c)^4 + 3*a^3*d*cos(d*x + c)^3 +
a^3*d*cos(d*x + c)^2)
```

3.122.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+a*cos(d*x+c))**(5/2),x)
```

```
output Timed out
```

3.122.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(5/2),x, algorithm
m="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.122.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+a*cos(d*x+c))^(5/2),x, algorithm
m="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:index.cc index_m i_lex_is_greater E
rror: Bad Argument Value
```

3.122.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos^3(c + dx) (a + a \cos(c + dx))^{5/2}} dx$$

```
input int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^(5/2)),x)
```

```
output int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + a*cos(c + d*x))^(5/2)), x)
```

3.122. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.123 $\int \cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$

3.123.1 Optimal result	1215
3.123.2 Mathematica [C] (warning: unable to verify)	1216
3.123.3 Rubi [A] (verified)	1217
3.123.4 Maple [B] (verified)	1221
3.123.5 Fricas [C] (verification not implemented)	1221
3.123.6 Sympy [F(-1)]	1222
3.123.7 Maxima [F]	1222
3.123.8 Giac [F]	1223
3.123.9 Mupad [B] (verification not implemented)	1223

3.123.1 Optimal result

Integrand size = 31, antiderivative size = 159

$$\int \cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{2a(9A+7B)E(\frac{1}{2}(c+dx)|2)}{15d} + \frac{10a(A+B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{21d}$$

$$+ \frac{10a(A+B)\sqrt{\cos(c+dx)}\sin(c+dx)}{21d} + \frac{2a(9A+7B)\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{45d}$$

$$+ \frac{2a(A+B)\cos^{\frac{5}{2}}(c+dx)\sin(c+dx)}{7d} + \frac{2aB\cos^{\frac{7}{2}}(c+dx)\sin(c+dx)}{9d}$$

```
output 2/15*a*(9*A+7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE
(sin(1/2*d*x+1/2*c),2^(1/2))/d+10/21*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/45*a*(9*A+7*B
)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/7*a*(A+B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d+
2/9*a*B*cos(d*x+c)^(7/2)*sin(d*x+c)/d+10/21*a*(A+B)*sin(d*x+c)*cos(d*x+c)^(
1/2)/d
```

3.123.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.00 (sec) , antiderivative size = 914, normalized size of antiderivative = 5.75

$$\int \cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= a \left(\sqrt{\cos(c + dx)}(1 + \cos(c + dx)) \sec^2 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{(9A + 7B) \cot(c)}{15d} \right. \right.$$

$$+ \frac{23(A + B) \cos(dx) \sin(c)}{84d} + \frac{(18A + 19B) \cos(2dx) \sin(2c)}{180d} + \frac{(A + B) \cos(3dx) \sin(3c)}{28d}$$

$$+ \frac{B \cos(4dx) \sin(4c)}{72d} + \frac{23(A + B) \cos(c) \sin(dx)}{84d} + \frac{(18A + 19B) \cos(2c) \sin(2dx)}{180d}$$

$$\left. + \frac{(A + B) \cos(3c) \sin(3dx)}{28d} + \frac{B \cos(4c) \sin(4dx)}{72d} \right)$$

$$\frac{5A(1 + \cos(c + dx)) \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{21d}$$

$$\frac{5B(1 + \cos(c + dx)) \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{21d}$$

$$\frac{3A(1 + \cos(c + dx)) \csc(c) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d}$$

$$\frac{7B(1 + \cos(c + dx)) \csc(c) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{30d}$$

input `Integrate[Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output

```

a*(Sqrt[Cos[c + d*x]]*(1 + Cos[c + d*x])*Sec[c/2 + (d*x)/2]^2*(-1/15*((9*A
+ 7*B)*Cot[c])/d + (23*(A + B)*Cos[d*x]*Sin[c])/(84*d) + ((18*A + 19*B)*C
os[2*d*x]*Sin[2*c])/(180*d) + ((A + B)*Cos[3*d*x]*Sin[3*c])/(28*d) + (B*Co
s[4*d*x]*Sin[4*c])/(72*d) + (23*(A + B)*Cos[c]*Sin[d*x])/(84*d) + ((18*A +
19*B)*Cos[2*c]*Sin[2*d*x])/(180*d) + ((A + B)*Cos[3*c]*Sin[3*d*x])/(28*d)
+ (B*Cos[4*c]*Sin[4*d*x])/(72*d)) - (5*A*(1 + Cos[c + d*x])*Csc[c]*Hyperg
eometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x
)/2]^2*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[
-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x -
ArcTan[Cot[c]]])]/(21*d*Sqrt[1 + Cot[c]^2]) - (5*B*(1 + Cos[c + d*x])*Csc
[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[
c/2 + (d*x)/2]^2*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c
]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 +
Sin[d*x - ArcTan[Cot[c]]])]/(21*d*Sqrt[1 + Cot[c]^2]) - (3*A*(1 + Cos[c +
d*x])*Csc[c]*Sec[c/2 + (d*x)/2]^2*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}
, Cos[d*x + ArcTan[Tan[c]]]^2]*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 -
Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]*Sqrt[Cos[c
]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Si
n[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x +
ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]...

```

3.123.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.99, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \cos^{\frac{5}{2}}(c + dx) \left((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)\right) dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.123. $\int \cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} \left((aA + aB) \sin \left(c + dx + \frac{\pi}{2} \right) + aA + aB \sin \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx \\
& \quad \downarrow \text{3502} \\
& \frac{2}{9} \int \frac{1}{2} \cos^{5/2}(c + dx)(a(9A + 7B) + 9a(A + B) \cos(c + dx))dx + \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{9} \int \cos^{5/2}(c + dx)(a(9A + 7B) + 9a(A + B) \cos(c + dx))dx + \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} \left(a(9A + 7B) + 9a(A + B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \\
& \quad \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{9} \left(a(9A + 7B) \int \cos^{5/2}(c + dx)dx + 9a(A + B) \int \cos^{7/2}(c + dx)dx \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \left(a(9A + 7B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} dx + 9a(A + B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{7/2} dx \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{3115} \\
& \frac{1}{9} \left(a(9A + 7B) \left(\frac{3}{5} \int \sqrt{\cos(c + dx)}dx + \frac{2 \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + 9a(A + B) \left(\frac{5}{7} \int \cos^{3/2}(c + dx)dx + \frac{2 \sin(c + dx) \cos^{5/2}(c + dx)}{7d} \right) \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \left(a(9A + 7B) \left(\frac{3}{5} \int \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)}dx + \frac{2 \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + 9a(A + B) \left(\frac{5}{7} \int \sin \left(c + dx + \frac{\pi}{2} \right) \cos^{3/2}(c + dx)dx + \frac{2 \sin(c + dx) \cos^{5/2}(c + dx)}{7d} \right) \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{7/2}(c + dx)}{9d} \\
& \quad \downarrow \text{3115}
\end{aligned}$$

3.123. $\int \cos^{5/2}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\frac{1}{9} \left(9a(A+B) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) + a(9A + \frac{2aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) \downarrow 3042$$

$$\frac{1}{9} \left(9a(A+B) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) + a(9A + \frac{2aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) \downarrow 3119$$

$$\frac{1}{9} \left(9a(A+B) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) + a(9A + \frac{2aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) \downarrow 3120$$

$$\frac{1}{9} \left(9a(A+B) \left(\frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} + \frac{5}{7} \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right) + a(9A + \frac{2aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right)$$

input `Int[Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(2*a*B*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(9*d) + (a*(9*A + 7*B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)) + 9*a*(A + B)*((2*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(7*d) + (5*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/7)/9`

3.123.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.123.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. 2(191) = 382.

Time = 13.84 (sec) , antiderivative size = 411, normalized size of antiderivative = 2.58

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}\left(-1120B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(720A+2960B)\left(\sin^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)$
parts	$\frac{2(aA+Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{21\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}$

```
input int(cos(d*x+c)^(5/2)*(a+cos(d*x+c)*a)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output -2/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(-1120*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10+(720*A+2960*B)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1584*A-3152*B)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(1344*A+1792*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-366*A-408*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+75*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-189*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+75*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-147*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.123.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.21

$$\int \cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{-75i\sqrt{2}(A+B)aweierstrassPInverse(-4,0,\cos(dx+c)+i\sin(dx+c))+75i\sqrt{2}(A+B)aweierstrassPInverse(-4,0,\cos(dx+c)-i\sin(dx+c))}{2}$$

3.123. $\int \cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$

input `integrate(cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/315*(-75*I*sqrt(2)*(A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 75*I*sqrt(2)*(A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*I*sqrt(2)*(9*A + 7*B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*(9*A + 7*B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(35*B*a*cos(d*x + c)^3 + 45*(A + B)*a*cos(d*x + c)^2 + 7*(9*A + 7*B)*a*cos(d*x + c) + 75*(A + B)*a)*sqrt(cos(d*x + c))*sin(d*x + c))/d`

3.123.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.123.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*cos(d*x + c)^(5/2), x)`

3.123. $\int \cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.123.8 Giac [F]

$$\int \cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \int (B\cos(dx+c)+A)(a\cos(dx+c)+a)\cos(dx+c)^{\frac{5}{2}}dx$$

input `integrate(cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*cos(d*x + c)^(5/2), x)`

3.123.9 Mupad [B] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.11

$$\int \cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= -\frac{2Aa\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}}$$

$$-\frac{2Aa\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}}$$

$$-\frac{2Ba\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}}$$

$$-\frac{2Ba\cos(c+dx)^{11/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)^2}}$$

input `int(cos(c + d*x)^(5/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`

output `-(2*A*a*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*A*a*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a*cos(c + d*x)^(11/2)*sin(c + d*x)*hypergeom([1/2, 11/4], 15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2))`

3.124 $\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$

3.124.1 Optimal result	1224
3.124.2 Mathematica [C] (warning: unable to verify)	1225
3.124.3 Rubi [A] (verified)	1226
3.124.4 Maple [B] (verified)	1229
3.124.5 Fricas [C] (verification not implemented)	1230
3.124.6 Sympy [F(-1)]	1231
3.124.7 Maxima [F]	1231
3.124.8 Giac [F]	1231
3.124.9 Mupad [B] (verification not implemented)	1232

3.124.1 Optimal result

Integrand size = 31, antiderivative size = 132

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{6a(A+B)E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2a(7A+5B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{21d}$$

$$+ \frac{2a(7A+5B)\sqrt{\cos(c+dx)} \sin(c+dx)}{21d}$$

$$+ \frac{2a(A+B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{5d} + \frac{2aB \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{7d}$$

```
output 6/5*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(
1/2*d*x+1/2*c),2^(1/2))/d+2/21*a*(7*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/co
s(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*a*(A+B)*cos(d
*x+c)^(3/2)*sin(d*x+c)/d+2/7*a*B*cos(d*x+c)^(5/2)*sin(d*x+c)/d+2/21*a*(7*A
+5*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.124.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.85 (sec) , antiderivative size = 872, normalized size of antiderivative = 6.61

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= a \left(\sqrt{\cos(c + dx)}(1 + \cos(c + dx)) \sec^2 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{3(A + B) \cot(c)}{5d} \right. \right.$$

$$\left. \left. + \frac{(28A + 23B) \cos(dx) \sin(c)}{84d} + \frac{(A + B) \cos(2dx) \sin(2c)}{10d} + \frac{B \cos(3dx) \sin(3c)}{28d} \right. \right.$$

$$\left. \left. + \frac{(28A + 23B) \cos(c) \sin(dx)}{84d} + \frac{(A + B) \cos(2c) \sin(2dx)}{10d} + \frac{B \cos(3c) \sin(3dx)}{28d} \right) \right.$$

$$\frac{A(1 + \cos(c + dx)) \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{3d}$$

$$\frac{5B(1 + \cos(c + dx)) \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{21d}$$

$$\frac{3A(1 + \cos(c + dx)) \csc(c) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d}$$

$$\frac{3B(1 + \cos(c + dx)) \csc(c) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output

```

a*(Sqrt[Cos[c + d*x]]*(1 + Cos[c + d*x])*Sec[c/2 + (d*x)/2]^2*((-3*(A + B)
*Cot[c])/(5*d) + ((28*A + 23*B)*Cos[d*x]*Sin[c])/(84*d) + ((A + B)*Cos[2*d
*x]*Sin[2*c])/(10*d) + (B*Cos[3*d*x]*Sin[3*c])/(28*d) + ((28*A + 23*B)*Cos
[c]*Sin[d*x])/(84*d) + ((A + B)*Cos[2*c]*Sin[2*d*x])/(10*d) + (B*Cos[3*c]*
Sin[3*d*x])/(28*d)) - (A*(1 + Cos[c + d*x])*Csc[c]*HypergeometricPFQ[{1/4,
1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^2*Sec[d*x -
ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]
^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]
]/(3*d*Sqrt[1 + Cot[c]^2]) - (5*B*(1 + Cos[c + d*x])*Csc[c]*Hypergeometric
PFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^2*S
ec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1
+ Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[
Cot[c]]])]/(21*d*Sqrt[1 + Cot[c]^2]) - (3*A*(1 + Cos[c + d*x])*Csc[c]*Sec[
c/2 + (d*x)/2]^2*(HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[
Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[
Tan[c]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan
[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan
[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c]]]*S
qrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan
[c]]]*Sqrt[1 + Tan[c]^2])))/(10*d) - (3*B*(1 + Cos[c + d*x])*Csc[c]*Sec...

```

3.124.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.98, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \cos^{\frac{3}{2}}(c + dx) \left((aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)\right) dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.124. $\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} \left((aA + aB) \sin \left(c + dx + \frac{\pi}{2} \right) + aA + aB \sin \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx \\
& \quad \downarrow \text{3502} \\
& \frac{2}{7} \int \frac{1}{2} \cos^{\frac{3}{2}}(c + dx) (a(7A + 5B) + 7a(A + B) \cos(c + dx)) dx + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \cos^{\frac{3}{2}}(c + dx) (a(7A + 5B) + 7a(A + B) \cos(c + dx)) dx + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} \left(a(7A + 5B) + 7a(A + B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \\
& \quad \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{7} \left(a(7A + 5B) \int \cos^{\frac{3}{2}}(c + dx) dx + 7a(A + B) \int \cos^{\frac{5}{2}}(c + dx) dx \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \left(a(7A + 5B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} dx + 7a(A + B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} dx \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{3115} \\
& \frac{1}{7} \left(7a(A + B) \left(\frac{3}{5} \int \sqrt{\cos(c + dx)} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + a(7A + 5B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d} \right) \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \left(7a(A + B) \left(\frac{3}{5} \int \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + a(7A + 5B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d} \right) \right) + \\
& \quad \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \\
& \quad \downarrow \text{3119}
\end{aligned}$$

3.124. $\int \cos^{\frac{3}{2}}(c + dx) (a + a \cos(c + dx)) (A + B \cos(c + dx)) dx$

$$\frac{1}{7} \left(a(7A + 5B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + 7a(A + B) \left(\frac{6E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \right) \right) + \frac{1}{7} \left(7a(A + B) \left(\frac{6E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + a(7A + 5B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \right) \right)$$

↓ 3120

input `Int[Cos[c + d*x]^(3/2)*(a + a*cos[c + d*x])*(A + B*cos[c + d*x]),x]`

output `(2*a*B*cos[c + d*x]^(5/2)*sin[c + d*x]/(7*d) + (a*(7*A + 5*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*sin[c + d*x])/(3*d)) + 7*a*(A + B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*cos[c + d*x]^(3/2)*sin[c + d*x])/(5*d)))/7`

3.124.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.124.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 382 vs. 2(168) = 336.

Time = 10.97 (sec) , antiderivative size = 383, normalized size of antiderivative = 2.90

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} \left(240B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-168A - 528B)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots\right)$
parts	$-\frac{2(aA + Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \left(-8\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 8\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots\right)$

input `int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*a)*(A+B*cos(d*x+c)), x, method=_RETURNVER BOSE)`

$$3.124. \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

output `-2/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(240*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8+(-168*A-528*B)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(308*A+448*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-112*A-122*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+35*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+25*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-63*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.124.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.36

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{-5i \sqrt{2}(7A + 5B)a \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2}(7A + 5B)a \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/105*(-5*I*sqrt(2)*(7*A + 5*B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*(7*A + 5*B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 63*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 63*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(15*B*a*cos(d*x + c)^2 + 21*(A + B)*a*cos(d*x + c) + 5*(7*A + 5*B)*a)*sqrt(cos(d*x + c))*sin(d*x + c)/d`

3.124.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.124.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*cos(d*x + c)^(3/2), x)`

3.124.8 Giac [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*cos(d*x + c)^(3/2), x)`

3.124.9 Mupad [B] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.26

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{2 A a \left(\sqrt{\cos(c + dx)} \sin(c + dx) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) \right)}{3 d} - \frac{2 A a \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}} - \frac{2 B a \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}} - \frac{2 B a \cos(c + dx)^{9/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c + dx)^2\right)}{9 d \sqrt{\sin(c + dx)^2}}$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`output `(2*A*a*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) - (2*A*a*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2))`

3.125 $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.125.1 Optimal result	1233
3.125.2 Mathematica [C] (warning: unable to verify)	1233
3.125.3 Rubi [A] (verified)	1234
3.125.4 Maple [B] (verified)	1237
3.125.5 Fricas [C] (verification not implemented)	1238
3.125.6 Sympy [F(-1)]	1238
3.125.7 Maxima [F]	1239
3.125.8 Giac [F]	1239
3.125.9 Mupad [B] (verification not implemented)	1239

3.125.1 Optimal result

Integrand size = 31, antiderivative size = 101

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{2a(5A + 3B)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d} + \frac{2a(A + B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d}$$

$$+ \frac{2a(A + B) \sqrt{\cos(c + dx)} \sin(c + dx)}{3d} + \frac{2aB \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{5d}$$

```
output 2/5*a*(5*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(
sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos
(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*a*B*cos(d*x+c)
^(3/2)*sin(d*x+c)/d+2/3*a*(A+B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.125.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.46 (sec) , antiderivative size = 363, normalized size of antiderivative = 3.59

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{a(1 + \cos(c + dx)) \sec^2\left(\frac{1}{2}(c + dx)\right) \left(-6(5A + 3B) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right)\right) \sec(c) \sin(c)}{\dots}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(a*(1 + Cos[c + d*x])*Sec[(c + d*x)/2]^2*(-6*(5*A + 3*B)*HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sec[c]*Sin[d*x + ArcTan[Tan[c]]] + (9*(5*A + 3*B)*Cos[c - d*x - ArcTan[Tan[c]]]*Csc[c]*Sec[c] + 15*A*Cos[c + d*x + ArcTan[Tan[c]]]*Csc[c]*Sec[c] + 9*B*Cos[c + d*x + ArcTan[Tan[c]]]*Csc[c]*Sec[c] - 60*A*Cos[c + d*x]*Cot[c]*Sqrt[Sec[c]^2] - 36*B*Cos[c + d*x]*Cot[c]*Sqrt[Sec[c]^2] - 20*(A + B)*Cos[c + d*x]*Sqrt[Cos[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Csc[c]^2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Sec[c]^2]*Sec[d*x - ArcTan[Cot[c]]]*Sin[c] + 20*A*Cos[c + d*x]*Sqrt[Sec[c]^2]*Sin[c + d*x] + 20*B*Cos[c + d*x]*Sqrt[Sec[c]^2]*Sin[c + d*x] + 12*B*Cos[c + d*x]^2*Sqrt[Sec[c]^2]*Sin[c + d*x])*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2]))/(60*d*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c]^2]*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2])`

3.125.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a \sin\left(c+dx+\frac{\pi}{2}\right) + a\right)\left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sqrt{\cos(c+dx)}((aA + aB) \cos(c+dx) + aA + aB \cos^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left((aA + aB) \sin\left(c+dx+\frac{\pi}{2}\right) + aA + aB \sin\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{2}{5} \int \frac{1}{2} \sqrt{\cos(c+dx)}(a(5A + 3B) + 5a(A + B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d}
 \end{aligned}$$

3.125. $\int \sqrt{\cos(c+dx)}(a + a \cos(c+dx))(A + B \cos(c+dx)) dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{5} \int \sqrt{\cos(c+dx)}(a(5A+3B)+5a(A+B)\cos(c+dx))dx + \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
& \downarrow 3042 \\
& \frac{1}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a(5A+3B)+5a(A+B)\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx + \\
& \quad \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
& \downarrow 3227 \\
& \frac{1}{5} \left(5a(A+B) \int \cos^{\frac{3}{2}}(c+dx)dx + a(5A+3B) \int \sqrt{\cos(c+dx)}dx \right) + \\
& \quad \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(a(5A+3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx + 5a(A+B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}dx \right) + \\
& \quad \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
& \downarrow 3115 \\
& \frac{1}{5} \left(a(5A+3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx + 5a(A+B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) \right) + \\
& \quad \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(a(5A+3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx + 5a(A+B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) \right) + \\
& \quad \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
& \downarrow 3119 \\
& \frac{1}{5} \left(5a(A+B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2a(5A+3B)E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) + \\
& \quad \frac{2aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d}
\end{aligned}$$

↓ 3120

$$\frac{1}{5} \left(\frac{2a(5A + 3B)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + 5a(A + B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) + \frac{2aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d}$$

input `Int[Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])*(A + B*cos[c + d*x]),x]`

output `(2*a*B*cos[c + d*x]^(3/2)*sin[c + d*x])/(5*d) + ((2*a*(5*A + 3*B)*EllipticE[(c + d*x)/2, 2])/d + 5*a*(A + B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*sin[c + d*x])/(3*d)))/5`

3.125.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.125.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. $2(141) = 282$.

Time = 8.57 (sec) , antiderivative size = 355, normalized size of antiderivative = 3.51

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} \left(-24B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (20A + 44B)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + (-1\right)$
parts	$-\frac{2(aA + Ba)\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} - d$

input `int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(-24*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+(20*A+44*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A-16*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-9*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.125. $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))(A + B \cos(c + dx)) dx$

3.125.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.59

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{-5i\sqrt{2}(A+B)a\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+5i\sqrt{2}(A+B)a\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `1/15*(-5*I*sqrt(2)*(A+B)*a*weierstrassPInverse(-4,0,cos(d*x+c)+I*sin(d*x+c))+5*I*sqrt(2)*(A+B)*a*weierstrassPInverse(-4,0,cos(d*x+c)-I*sin(d*x+c))+3*I*sqrt(2)*(5*A+3*B)*a*weierstrassZeta(-4,0,weierstrassPInverse(-4,0,cos(d*x+c)+I*sin(d*x+c)))-3*I*sqrt(2)*(5*A+3*B)*a*weierstrassZeta(-4,0,weierstrassPInverse(-4,0,cos(d*x+c)-I*sin(d*x+c)))+2*(3*B*a*cos(d*x+c)+5*(A+B)*a)*sqrt(cos(d*x+c))*sin(d*x+c))/d`

3.125.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))(A+B\cos(c+dx))dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*cos(d*x+c)**(1/2),x)`

output `Timed out`

3.125.7 Maxima [F]

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \int (B\cos(dx+c)+A)(a\cos(dx+c)+a)\sqrt{\cos(dx+c)}dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.125.8 Giac [F]

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \int (B\cos(dx+c)+A)(a\cos(dx+c)+a)\sqrt{\cos(dx+c)}dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.125.9 Mupad [B] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.27

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{2Aa\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)\right)}{3d}$$

$$+ \frac{2Ba\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)\right)}{3d} + \frac{2AaE\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)}{d}$$

$$- \frac{2Ba\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}}$$

3.125. $\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))(A+B\cos(c+dx))dx$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x)),x)`

output `(2*A*a*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*B*a*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*A*a*ellipticE(c/2 + (d*x)/2, 2))/d - (2*B*a*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))`

3.126
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.126.1 Optimal result 1241
 3.126.2 Mathematica [C] (warning: unable to verify) 1241
 3.126.3 Rubi [A] (verified) 1242
 3.126.4 Maple [B] (verified) 1245
 3.126.5 Fricas [C] (verification not implemented) 1245
 3.126.6 Sympy [F] 1246
 3.126.7 Maxima [F] 1246
 3.126.8 Giac [F] 1247
 3.126.9 Mupad [B] (verification not implemented) 1247

3.126.1 Optimal result

Integrand size = 31, antiderivative size = 70

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{2a(A + B)E(\frac{1}{2}(c + dx) | 2)}{d} + \frac{2a(3A + B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2aB \sqrt{\cos(c + dx)} \sin(c + dx)}{3d}$$

```
output 2*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*(3*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*B*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.126.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.62 (sec) , antiderivative size = 309, normalized size of antiderivative = 4.41

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{a(1 + \cos(c + dx)) \sec^2(\frac{1}{2}(c + dx)) \left(-6(A + B) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right)\right) \sec(c) \sin(dx)}{\dots}$$

input `Integrate[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x
]`

output `(a*(1 + Cos[c + d*x])*Sec[(c + d*x)/2]^2*(-6*(A + B)*HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sec[c]*Sin[d*x + ArcTan[Tan[c]]) + (9*(A + B)*Cos[c - d*x - ArcTan[Tan[c]]]*Csc[c]*Sec[c] + 3*A*Cos[c + d*x + ArcTan[Tan[c]]]*Csc[c]*Sec[c] + 3*B*Cos[c + d*x + ArcTan[Tan[c]]]*Csc[c]*Sec[c] - 12*A*Cos[c + d*x]*Cot[c]*Sqrt[Sec[c]^2] - 12*B*Cos[c + d*x]*Cot[c]*Sqrt[Sec[c]^2] - 4*(3*A + B)*Cos[c + d*x]*Sqrt[Cos[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Csc[c]^2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Sec[c]^2]*Sec[d*x - ArcTan[Cot[c]]]*Sin[c] + 4*B*Cos[c + d*x]*Sqrt[Sec[c]^2]*Sin[c + d*x])*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2])/((12*d*Sqrt[Cos[c + d*x])*Sqrt[Sec[c]^2]*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2])`

3.126.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3447

$$\int \frac{(aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3502

3.126. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\begin{aligned}
& \frac{2}{3} \int \frac{a(3A+B) + 3a(A+B) \cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx + \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \\
& \quad \downarrow 27 \\
& \frac{1}{3} \int \frac{a(3A+B) + 3a(A+B) \cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \int \frac{a(3A+B) + 3a(A+B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \\
& \quad \downarrow 3227 \\
& \frac{1}{3} \left(a(3A+B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a(A+B) \int \sqrt{\cos(c+dx)} dx \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left(a(3A+B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + 3a(A+B) \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \\
& \quad \downarrow 3119 \\
& \frac{1}{3} \left(a(3A+B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{6a(A+B)E(\frac{1}{2}(c+dx)|2)}{d} \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \\
& \quad \downarrow 3120 \\
& \frac{1}{3} \left(\frac{2a(3A+B) \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6a(A+B)E(\frac{1}{2}(c+dx)|2)}{d} \right) + \\
& \quad \frac{2aB \sin(c+dx) \sqrt{\cos(c+dx)}}{3d}
\end{aligned}$$

input `Int[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output $((6*a*(A + B)*\text{EllipticE}[(c + d*x)/2, 2])/d + (2*a*(3*A + B)*\text{EllipticF}[(c + d*x)/2, 2])/d)/3 + (2*a*B*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(3*d)$

3.126.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3119 $\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3227 $\text{Int}[(b_.)*\text{sin}[(e_.) + (f_.)*(x_)]^{(m_)*((c_.) + (d_.)*\text{sin}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[c \quad \text{Int}[(b*\text{Sin}[e + f*x])^m, x], x] + \text{Simp}[d/b \quad \text{Int}[(b*\text{Sin}[e + f*x])^{(m + 1)}, x], x] /; \text{FreeQ}[\{b, c, d, e, f, m\}, x]$

rule 3447 $\text{Int}[(a_.) + (b_.)*\text{sin}[(e_.) + (f_.)*(x_)]^{(m_.)*((A_.) + (B_.)*\text{sin}[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*\text{sin}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Int}[(a + b*\text{Sin}[e + f*x])^m*(A*c + (B*c + A*d)*\text{Sin}[e + f*x] + B*d*\text{Sin}[e + f*x]^2), x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0]$

rule 3502 $\text{Int}[(a_.) + (b_.)*\text{sin}[(e_.) + (f_.)*(x_)]^{(m_.)*((A_.) + (B_.)*\text{sin}[(e_.) + (f_.)*(x_)] + (C_.)*\text{sin}[(e_.) + (f_.)*(x_)]^2), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cos}[e + f*x]*((a + b*\text{Sin}[e + f*x])^{(m + 1)}/(b*f*(m + 2))), x] + \text{Simp}[1/(b*(m + 2)) \quad \text{Int}[(a + b*\text{Sin}[e + f*x])^m*\text{Simp}[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, e, f, A, B, C, m\}, x] \ \&\& \ !\text{LtQ}[m, -1]$

3.126.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. 2(116) = 232.

Time = 6.46 (sec) , antiderivative size = 321, normalized size of antiderivative = 4.59

method	result
default	$- \frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a\left(4B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+3A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\right)}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)$
parts	$\frac{2(aA+Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}}\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right) - \frac{2Ba\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}}{\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}d$

input `int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(4*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2+B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.126.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 142, normalized size of antiderivative = 2.03

$$\int \frac{(a+a\cos(c+dx))(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

$$= \frac{2Ba\sqrt{\cos(dx+c)}\sin(dx+c) - i\sqrt{2}(3A+B)\text{aweierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))}{\sin(dx+c)}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,algorithm="fracas")`

3.126. $\int \frac{(a+a\cos(c+dx))(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

output `1/3*(2*B*a*sqrt(cos(d*x + c))*sin(d*x + c) - I*sqrt(2)*(3*A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*(3*A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.126.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = a \left(\int \frac{A}{\sqrt{\cos(c + dx)}} dx + \int A \sqrt{\cos(c + dx)} dx + \int B \sqrt{\cos(c + dx)} dx + \int B \cos^{\frac{3}{2}}(c + dx) dx \right)$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2), x)`

output `a*(Integral(A/sqrt(cos(c + d*x)), x) + Integral(A*sqrt(cos(c + d*x)), x) + Integral(B*sqrt(cos(c + d*x)), x) + Integral(B*cos(c + d*x)**(3/2), x))`

3.126.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.126. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

3.126.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.126.9 Mupad [B] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.13

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2 B a \left(\sqrt{\cos(c + dx)} \sin(c + dx) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) \right)}{3 d}$$

$$+ \frac{2 B a E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 A a \left(E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) \right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^(1/2),x)`

output `(2*B*a*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*B*a*ellipticE(c/2 + (d*x)/2, 2))/d + (2*A*a*(ellipticE(c/2 + (d*x)/2, 2) + ellipticF(c/2 + (d*x)/2, 2)))/d`

$$3.127 \quad \int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.127.1 Optimal result	1248
3.127.2 Mathematica [C] (verified)	1248
3.127.3 Rubi [A] (verified)	1249
3.127.4 Maple [B] (verified)	1251
3.127.5 Fracas [C] (verification not implemented)	1252
3.127.6 Sympy [F(-1)]	1253
3.127.7 Maxima [F]	1253
3.127.8 Giac [F]	1253
3.127.9 Mupad [B] (verification not implemented)	1254

3.127.1 Optimal result

Integrand size = 31, antiderivative size = 66

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = -\frac{2a(A - B)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + \frac{2a(A + B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}}$$

output

```
-2*a*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*a*A*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.127.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.32 (sec) , antiderivative size = 256, normalized size of antiderivative = 3.88

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \frac{a(1 + \cos(c + dx)) \sec^2\left(\frac{1}{2}(c + dx)\right) \left(\frac{\csc(c) \left(-3(A - B) \cos(c - dx - \arctan(\tan(c))) \sec(c) - (A - B) \cos(c + dx + \arctan(\tan(c))) \sec(c)\right)}{\sqrt{\sec^2(c)}} \right)}{\dots}$$

3.127. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

input `Integrate[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x
]`

output `(a*(1 + Cos[c + d*x])*Sec[(c + d*x)/2]^2*((Csc[c]*(-3*(A - B)*Cos[c - d*x
- ArcTan[Tan[c]])*Sec[c] - (A - B)*Cos[c + d*x + ArcTan[Tan[c]])*Sec[c] +
2*((2*A - B)*Cos[d*x] - B*Cos[2*c + d*x])*Sqrt[Sec[c]^2])/Sqrt[Sec[c]^2
- 4*(A + B)*Cos[c + d*x])*Sqrt[Cos[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Csc[c]^2]*
HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[d*x
- ArcTan[Cot[c]]]*Sin[c] + (2*(A - B)*HypergeometricPFQ[{-1/2, -1/4}, {3/4
}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sec[c]*Sin[d*x + ArcTan[Tan[c]]])/(Sqrt[Se
c[c]^2]*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2]))/(4*d*Sqrt[Cos[c + d*x]))`

3.127.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{(aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)}{\cos^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3500} \\ & 2 \int \frac{a(A + B) - a(A - B) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \end{aligned}$$

3.127. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{a(A+B) - a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{27} \\
& \int \frac{a(A+B) - a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& a(A+B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx - a(A-B) \int \sqrt{\cos(c+dx)} dx + \frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3227} \\
& a(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - a(A-B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& a(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(A-B)E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3119} \\
& \frac{2a(A+B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d} - \frac{2a(A-B)E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3120}
\end{aligned}$$

input `Int[((a + a*cos[c + d*x])*(A + B*cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(-2*a*(A - B)*EllipticE[(c + d*x)/2, 2])/d + (2*a*(A + B)*EllipticF[(c + d*x)/2, 2])/d + (2*a*A*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])`

3.127.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.127. $\int \frac{(a+a\cos(c+dx))(A+B\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.127.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 241 vs. 2(116) = 232.

Time = 5.78 (sec) , antiderivative size = 242, normalized size of antiderivative = 3.67

method	result
default	$\frac{2a \left(2A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)} - 1 F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)} \right)}{\dots}$
parts	$\frac{2(aA+Ba) a m^{-1} \left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2}}{d} + \frac{2Ba \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$

3.127.
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

input `int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `2*a*(2*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2-A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))-A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.127.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 173, normalized size of antiderivative = 2.62

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{-i \sqrt{2}(A + B)a \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2}(A + B)a \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - i \sqrt{2}(A - B)a \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2}(A - B)a \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2Aa \sqrt{\cos(dx + c)} \sin(dx + c)}{d \cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="fracas")`

output `(-I*sqrt(2)*(A + B)*a*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*(A + B)*a*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - I*sqrt(2)*(A - B)*a*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*(A - B)*a*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*A*a*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c))`

3.127.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Timed out`

3.127.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\cos(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`

3.127.8 Giac [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\cos(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`

3.127.9 Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.36

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 A a F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 B a \left(E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)\right)}{d}$$

$$+ \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^(3/2),x)`output `(2*A*a*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*a*(ellipticE(c/2 + (d*x)/2, 2) + ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.128
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.128.1 Optimal result 1255
 3.128.2 Mathematica [C] (warning: unable to verify) 1255
 3.128.3 Rubi [A] (verified) 1256
 3.128.4 Maple [B] (verified) 1259
 3.128.5 Fracas [C] (verification not implemented) 1260
 3.128.6 Sympy [F(-1)] 1261
 3.128.7 Maxima [F] 1261
 3.128.8 Giac [F] 1261
 3.128.9 Mupad [B] (verification not implemented) 1262

3.128.1 Optimal result

Integrand size = 31, antiderivative size = 95

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = -\frac{2a(A + B)E(\frac{1}{2}(c + dx)|2)}{d} + \frac{2a(A + 3B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2a(A + B) \sin(c + dx)}{d \sqrt{\cos(c + dx)}}$$

output `-2*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*(A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*A*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2*a*(A+B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)`

3.128.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.21 (sec) , antiderivative size = 293, normalized size of antiderivative = 3.08

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \frac{a(1 + \cos(c + dx)) \sec^2(\frac{1}{2}(c + dx)) \left(6(A + B) \cos(c + dx) {}_2F_1(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c))))\right)}{\dots}$$

input `Integrate[((a + a*cos[c + d*x])*(A + B*cos[c + d*x]))/Cos[c + d*x]^(5/2),x
]`

output `(a*(1 + Cos[c + d*x])*Sec[(c + d*x)/2]^2*(6*(A + B)*Cos[c + d*x]*HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sec[c]*Sin[d*x + ArcTan[Tan[c]]] + (-3*(A + B)*Cos[c + d*x]*(3*Cos[c - d*x - ArcTan[Tan[c]]] + Cos[c + d*x + ArcTan[Tan[c]]])*Csc[c]*Sec[c] + 2*(3*(A + B)*Cot[c] + (A*cos[d*x] - A*cos[2*c + d*x] + 3*(A + B)*Cos[c + 2*d*x])*Csc[c])*Sqrt[Sec[c]^2] - 4*(A + 3*B)*Cos[c + d*x]^2*Sqrt[Cos[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Csc[c]^2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Sec[c]^2]*Sec[d*x - ArcTan[Cot[c]]]*Sin[c])*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2))/((12*d*cos[c + d*x]^(3/2)*Sqrt[Sec[c]^2]*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2))`

3.128.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{(aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)}{\cos^{\frac{5}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx \\ & \quad \downarrow \text{3500} \end{aligned}$$

3.128. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \frac{2}{3} \int \frac{3a(A+B) + a(A+3B) \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{3a(A+B) + a(A+3B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3a(A+B) + a(A+3B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{3} \left(3a(A+B) \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx + a(A+3B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx \right) + \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(3a(A+B) \int \frac{1}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + a(A+3B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx \right) + \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3116} \\
& \frac{1}{3} \left(a(A+3B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + 3a(A+B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx \right) \right) + \\
& \quad \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(a(A+3B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + 3a(A+B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx \right) \right) + \\
& \quad \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3119} \\
& \frac{1}{3} \left(a(A+3B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + 3a(A+B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) \right) + \\
& \quad \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3120}
\end{aligned}$$

3.128. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(\frac{2a(A+3B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} + 3a(A+B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) \right) + \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])/Cos[c + d*x]^(5/2),x]`

output `(2*a*A*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((2*a*(A + 3*B)*EllipticF[(c + d*x)/2, 2])/d + 3*a*(A + B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/3`

3.128.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-A*b^2
- a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.128.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 398 vs. 2(139) = 278.

Time = 7.31 (sec) , antiderivative size = 399, normalized size of antiderivative = 4.20

method	result
default	$4\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^a \left(\frac{B\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{2\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}} + \frac{A\left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}}{6}\right)}{6} \right)$
parts	$\frac{2(aA + Ba)\left(-2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2), x, method=_RETURNVER
BOSE)
```

$$3.128. \int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

output $-4*(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a*(1/2*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+1/2*A*(-1/6*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(\cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})))+(1/2*A+1/2*B)/\sin(1/2*d*x+1/2*c)^2/(2*\sin(1/2*d*x+1/2*c)^2-1)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$

3.128.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.06

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= -i\sqrt{2}(A + 3B)a \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i\sqrt{2}(A + 3B)$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fracas")`

output $1/3*(-I*\sqrt{2}*(A + 3*B)*a*\cos(d*x + c)^2*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*(A + 3*B)*a*\cos(d*x + c)^2*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*I*\sqrt{2}*(A + B)*a*\cos(d*x + c)^2*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + 3*I*\sqrt{2}*(A + B)*a*\cos(d*x + c)^2*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*(3*(A + B)*a*\cos(d*x + c) + A*a)*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^2)$

3.128.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`output `Timed out`**3.128.7 Maxima [F]**

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\cos(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`**3.128.8 Giac [F]**

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\cos(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`

3.128.9 Mupad [B] (verification not implemented)

Time = 1.43 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.58

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 B a F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^(5/2),x)`output `(2*B*a*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) + (2*A*a*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2)) + (2*B*a*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.129
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.129.1 Optimal result 1263
 3.129.2 Mathematica [C] (warning: unable to verify) 1264
 3.129.3 Rubi [A] (verified) 1265
 3.129.4 Maple [B] (verified) 1268
 3.129.5 Fricas [C] (verification not implemented) 1269
 3.129.6 Sympy [F(-1)] 1270
 3.129.7 Maxima [F] 1270
 3.129.8 Giac [F] 1271
 3.129.9 Mupad [B] (verification not implemented) 1271

3.129.1 Optimal result

Integrand size = 31, antiderivative size = 132

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = -\frac{2a(3A + 5B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{2a(A + B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} + \frac{2a(A + B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2a(3A + 5B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

```
output -2/5*a*(3*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE
(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/co
s(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*a*A*sin(d*x+c
)/d/cos(d*x+c)^(5/2)+2/3*a*(A+B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/5*a*(3*A+
5*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.129.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.81 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.55

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = a \left(\sqrt{\cos(c + dx)}(1 + \cos(c + dx)) \sec^2 \left(\frac{c}{2} + \frac{dx}{2} \right) \right. \\ \left. + \frac{dx}{2} \right) \left(\frac{(3A + 5B) \csc(c) \sec(c)}{5d} + \frac{A \sec(c) \sec^3(c + dx) \sin(dx)}{5d} \right. \\ \left. + \frac{\sec(c) \sec^2(c + dx)(3A \sin(c) + 5A \sin(dx) + 5B \sin(dx))}{15d} \right. \\ \left. + \frac{\sec(c) \sec(c + dx)(5A \sin(c) + 5B \sin(c) + 9A \sin(dx) + 15B \sin(dx))}{15d} \right) \\ - \frac{A(1 + \cos(c + dx)) \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{3d} \\ - \frac{B(1 + \cos(c + dx)) \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{3d} \\ + \frac{3A(1 + \cos(c + dx)) \csc(c) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d} \\ + \frac{B(1 + \cos(c + dx)) \csc(c) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{2d}$$

input `Integrate[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2), x]`

$$3.129. \int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

output

```

a*(Sqrt[Cos[c + d*x]]*(1 + Cos[c + d*x])*Sec[c/2 + (d*x)/2]^2*(((3*A + 5*B)
)*Csc[c]*Sec[c])/(5*d) + (A*Sec[c]*Sec[c + d*x]^3*Sin[d*x])/(5*d) + (Sec[c
]*Sec[c + d*x]^2*(3*A*Sin[c] + 5*A*Sin[d*x] + 5*B*Sin[d*x]))/(15*d) + (Sec
[c]*Sec[c + d*x]*(5*A*Sin[c] + 5*B*Sin[c] + 9*A*Sin[d*x] + 15*B*Sin[d*x]))
/(15*d) - (A*(1 + Cos[c + d*x])*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4
}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^2*Sec[d*x - ArcTan[Cot[
c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*
Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3*d*Sqrt
[1 + Cot[c]^2]) - (B*(1 + Cos[c + d*x])*Csc[c]*HypergeometricPFQ[{1/4, 1/2
}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^2*Sec[d*x - ArcT
an[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*
Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3
*d*Sqrt[1 + Cot[c]^2]) + (3*A*(1 + Cos[c + d*x])*Csc[c]*Sec[c/2 + (d*x)/2
]^2*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*S
in[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]*Sqrt
[1 + Cos[d*x + ArcTan[Tan[c]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt
[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/
Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]
^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 +
Tan[c]^2]))/(10*d) + (B*(1 + Cos[c + d*x])*Csc[c]*Sec[c/2 + (d*x)/2]^...

```

3.129.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.95, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \cos(c + dx) + a)(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{(aA + aB) \cos(c + dx) + aA + aB \cos^2(c + dx)}{\cos^{\frac{7}{2}}(c + dx)} dx
 \end{aligned}$$

3.129. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{(aA + aB) \sin(c + dx + \frac{\pi}{2}) + aA + aB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \downarrow 3500 \\
& \frac{2}{5} \int \frac{5a(A + B) + a(3A + 5B) \cos(c + dx)}{2 \cos^{\frac{5}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow 27 \\
& \frac{1}{5} \int \frac{5a(A + B) + a(3A + 5B) \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \int \frac{5a(A + B) + a(3A + 5B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3227 \\
& \frac{1}{5} \left(5a(A + B) \int \frac{1}{\cos^{\frac{5}{2}}(c + dx)} dx + a(3A + 5B) \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx \right) + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(5a(A + B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + a(3A + 5B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \\
& \quad \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3116 \\
& \frac{1}{5} \left(5a(A + B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + a(3A + 5B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right) \right) \\
& \quad \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(5a(A + B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + a(3A + 5B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\sin(c + dx)} dx \right) \right) \\
& \quad \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)}
\end{aligned}$$

3.129. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

↓ 3119

$$\frac{1}{5} \left(5a(A+B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + a(3A+5B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx))}{d} \right) \right. \\ \left. \frac{2aA \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(5a(A+B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + a(3A+5B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx))}{d} \right) \right. \\ \left. \frac{2aA \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (5*a*(A + B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))) + a*(3*A + 5*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/5`

3.129.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.129.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 633 vs. 2(168) = 336.

Time = 10.38 (sec) , antiderivative size = 634, normalized size of antiderivative = 4.80

method	result
default	$- \frac{4 \sqrt{-2 \left(\cos^2 \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{a \left(\frac{B \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \left(2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \cos \left(\frac{dx}{2} + \frac{c}{2} \right) - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)}{2 \sin \left(\frac{dx}{2} + \frac{c}{2} \right)^2 \left(2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1 \right)} \right)}$
parts	Expression too large to display

3.129.
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$$

input `int((a+cos(d*x+c))*a)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RETURNVERBOSE)`

output `-4*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(1/2*B/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))+1/10*A/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+(1/2*A+1/2*B)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.129.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.66

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{-5i \sqrt{2}(A + B)a \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2}(A + B)a}{\dots}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="fricas")`

output $1/15*(-5*I*\sqrt{2}*(A + B)*a*\cos(d*x + c)^3*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + 5*I*\sqrt{2}*(A + B)*a*\cos(d*x + c)^3*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*I*\sqrt{2}*(3*A + 5*B)*a*\cos(d*x + c)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + 3*I*\sqrt{2}*(3*A + 5*B)*a*\cos(d*x + c)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*(3*(3*A + 5*B)*a*\cos(d*x + c)^2 + 5*(A + B)*a*\cos(d*x + c) + 3*A*a)*\sqrt{\cos(d*x + c)}*\sin(d*x + c))/(d*\cos(d*x + c)^3)$

3.129.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.129.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\cos(dx + c)^{\frac{7}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/cos(d*x + c)^(7/2), x)`

3.129.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/cos(d*x + c)^(7/2), x)`

3.129.9 Mupad [B] (verification not implemented)

Time = 1.86 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.34

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right)}{5 d \cos(c + dx)^{5/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/cos(c + d*x)^(7/2),x)`

output $(2Aa\sin(c + dx)\operatorname{hypergeom}([-3/4, 1/2], 1/4, \cos(c + dx)^2))/(3d\cos(c + dx)^{3/2}(\sin(c + dx)^2)^{1/2}) + (2Aa\sin(c + dx)\operatorname{hypergeom}([-5/4, 1/2], -1/4, \cos(c + dx)^2))/(5d\cos(c + dx)^{5/2}(\sin(c + dx)^2)^{1/2}) + (2Ba\sin(c + dx)\operatorname{hypergeom}([-1/4, 1/2], 3/4, \cos(c + dx)^2))/(d\cos(c + dx)^{1/2}(\sin(c + dx)^2)^{1/2}) + (2Ba\sin(c + dx)\operatorname{hypergeom}([-3/4, 1/2], 1/4, \cos(c + dx)^2))/(3d\cos(c + dx)^{3/2}(\sin(c + dx)^2)^{1/2})$

3.130 $\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.130.1 Optimal result	1273
3.130.2 Mathematica [C] (warning: unable to verify)	1274
3.130.3 Rubi [A] (verified)	1275
3.130.4 Maple [A] (verified)	1279
3.130.5 Fricas [C] (verification not implemented)	1280
3.130.6 Sympy [F(-1)]	1281
3.130.7 Maxima [F]	1281
3.130.8 Giac [F]	1281
3.130.9 Mupad [B] (verification not implemented)	1282

3.130.1 Optimal result

Integrand size = 33, antiderivative size = 194

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2(A+B \cos(c+dx)) dx$$

$$= \frac{4a^2(9A+8B)E(\frac{1}{2}(c+dx)|2)}{15d} + \frac{4a^2(6A+5B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{21d}$$

$$+ \frac{4a^2(6A+5B)\sqrt{\cos(c+dx)}\sin(c+dx)}{21d} + \frac{4a^2(9A+8B)\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{45d}$$

$$+ \frac{2a^2(9A+11B)\cos^{\frac{5}{2}}(c+dx)\sin(c+dx)}{63d}$$

$$+ \frac{2B\cos^{\frac{5}{2}}(c+dx)(a^2+a^2\cos(c+dx))\sin(c+dx)}{9d}$$

```
output 4/15*a^2*(9*A+8*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellipti
cE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/21*a^2*(6*A+5*B)*(cos(1/2*d*x+1/2*c)^2)
^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/45*a^2
*(9*A+8*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/63*a^2*(9*A+11*B)*cos(d*x+c)^(5
/2)*sin(d*x+c)/d+2/9*B*cos(d*x+c)^(5/2)*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d
+4/21*a^2*(6*A+5*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```


3.130.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.56 (sec) , antiderivative size = 944, normalized size of antiderivative = 4.87

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2 \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(-\frac{(9A + 8B) \cot(c)}{15d} \right. \\ \left. + \frac{(51A + 46B) \cos(dx) \sin(c)}{168d} + \frac{(36A + 37B) \cos(2dx) \sin(2c)}{360d} \right. \\ \left. + \frac{(A + 2B) \cos(3dx) \sin(3c)}{56d} + \frac{B \cos(4dx) \sin(4c)}{144d} + \frac{(51A + 46B) \cos(c) \sin(dx)}{360d} \right. \\ \left. + \frac{(36A + 37B) \cos(2c) \sin(2dx)}{360d} + \frac{(A + 2B) \cos(3c) \sin(3dx)}{56d} + \frac{B \cos(4c) \sin(4dx)}{144d} \right) \\ - \frac{2A(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{7d} \\ - \frac{5B(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{21d} \\ - \frac{3A(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d} \\ - \frac{4B(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{15d}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output $\text{Sqrt}[\text{Cos}[c + d*x]]*(a + a*\text{Cos}[c + d*x])^2*\text{Sec}[c/2 + (d*x)/2]^4*(-1/15*((9*A + 8*B)*\text{Cot}[c])/d + ((51*A + 46*B)*\text{Cos}[d*x]*\text{Sin}[c])/(168*d) + ((36*A + 37*B)*\text{Cos}[2*d*x]*\text{Sin}[2*c])/(360*d) + ((A + 2*B)*\text{Cos}[3*d*x]*\text{Sin}[3*c])/(56*d) + (B*\text{Cos}[4*d*x]*\text{Sin}[4*c])/(144*d) + ((51*A + 46*B)*\text{Cos}[c]*\text{Sin}[d*x])/(168*d) + ((36*A + 37*B)*\text{Cos}[2*c]*\text{Sin}[2*d*x])/(360*d) + ((A + 2*B)*\text{Cos}[3*c]*\text{Sin}[3*d*x])/(56*d) + (B*\text{Cos}[4*c]*\text{Sin}[4*d*x])/(144*d)) - (2*A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(7*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (5*B*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(21*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (3*A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{Sec}[c/2 + (d*x)/2]^4*((\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]]*\text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\text{Cos}[c]^2*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])/(\text{Cos}[c]^...$

3.130.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.98, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow 3455$$

$$\frac{2}{9} \int \frac{1}{2} \cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)(a(9A + 5B) + a(9A + 11B) \cos(c + dx)) dx + \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d}$$

3.130. $\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\downarrow 27$$

$$\frac{1}{9} \int \cos^{\frac{3}{2}}(c+dx) (\cos(c+dx)a+a) (a(9A+5B)+a(9A+11B)\cos(c+dx)) dx + \frac{2B \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{9d}$$

$$\downarrow 3042$$

$$\frac{1}{9} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right) \left(a(9A+5B)+a(9A+11B)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2B \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{9d}$$

$$\downarrow 3447$$

$$\frac{1}{9} \int \cos^{\frac{3}{2}}(c+dx) \left((9A+11B) \cos^2(c+dx)a^2 + (9A+5B)a^2 + ((9A+5B)a^2 + (9A+11B)a^2) \cos(c+dx) \right) dx + \frac{2B \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{9d}$$

$$\downarrow 3042$$

$$\frac{1}{9} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left((9A+11B) \sin\left(c+dx+\frac{\pi}{2}\right)^2 a^2 + (9A+5B)a^2 + ((9A+5B)a^2 + (9A+11B)a^2) \sin\left(c+dx+\frac{\pi}{2}\right) \right) dx + \frac{2B \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{9d}$$

$$\downarrow 3502$$

$$\frac{1}{9} \left(\frac{2}{7} \int \cos^{\frac{3}{2}}(c+dx) (9(6A+5B)a^2 + 7(9A+8B)\cos(c+dx)a^2) dx + \frac{2a^2(9A+11B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{7d} \right) + \frac{2B \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{9d}$$

$$\downarrow 3042$$

$$\frac{1}{9} \left(\frac{2}{7} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} (9(6A+5B)a^2 + 7(9A+8B)\sin\left(c+dx+\frac{\pi}{2}\right)a^2) dx + \frac{2a^2(9A+11B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{7d} \right) + \frac{2B \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{9d}$$

$$\downarrow 3227$$

$$\frac{1}{9} \left(\frac{2}{7} \left(9a^2(6A + 5B) \int \cos^{\frac{3}{2}}(c + dx) dx + 7a^2(9A + 8B) \int \cos^{\frac{5}{2}}(c + dx) dx \right) + \frac{2a^2(9A + 11B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d} \\ \downarrow \text{3042}$$

$$\frac{1}{9} \left(\frac{2}{7} \left(9a^2(6A + 5B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} dx + 7a^2(9A + 8B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} dx \right) + \frac{2a^2(9A + 11B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d} \\ \downarrow \text{3115}$$

$$\frac{1}{9} \left(\frac{2}{7} \left(7a^2(9A + 8B) \left(\frac{3}{5} \int \sqrt{\cos(c + dx)} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + 9a^2(6A + 5B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d} \\ \downarrow \text{3042}$$

$$\frac{1}{9} \left(\frac{2}{7} \left(7a^2(9A + 8B) \left(\frac{3}{5} \int \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + 9a^2(6A + 5B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)}} dx \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d} \\ \downarrow \text{3119}$$

$$\frac{1}{9} \left(\frac{2}{7} \left(9a^2(6A + 5B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + 7a^2(9A + 8B) \left(\frac{6E \left(\frac{1}{2}(c + dx) \mid 2 \right)}{5d} \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d} \\ \downarrow \text{3120}$$

$$\frac{1}{9} \left(\frac{2a^2(9A + 11B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} + \frac{2}{7} \left(7a^2(9A + 8B) \left(\frac{6E \left(\frac{1}{2}(c + dx) \mid 2 \right)}{5d} \right) + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{9d}$$

input `Int[Cos[c + d*x]^(3/2)*(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]),x]`

output `(2*B*cos[c + d*x]^(5/2)*(a^2 + a^2*cos[c + d*x])*Sin[c + d*x])/(9*d) + ((2*a^2*(9*A + 11*B)*cos[c + d*x]^(5/2)*Sin[c + d*x])/(7*d) + (2*(9*a^2*(6*A + 5*B))*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)) + 7*a^2*(9*A + 8*B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d))))/7)/9`

3.130.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.130.4 Maple [A] (verified)

Time = 13.62 (sec) , antiderivative size = 413, normalized size of antiderivative = 2.13

method	result
default	$-\frac{4\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^2} \left(-560B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (360A + 1840B)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)$
parts	Expression too large to display

input `int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c)), x, method=_RETURNV ERBOSE)`

$$3.130. \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

output `-4/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(-560*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10+(360*A+1840*B)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1044*A-2368*B)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(1134*A+1568*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-351*A-387*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+90*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-189*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+75*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-168*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.130.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.15

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx =$$

$$\frac{2\left(15i\sqrt{2}(6A+5B)a^2\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))-15i\sqrt{2}(6A+5B)a^2\right)}{\dots}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `-2/315*(15*I*sqrt(2)*(6*A + 5*B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 15*I*sqrt(2)*(6*A + 5*B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*I*sqrt(2)*(9*A + 8*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*(9*A + 8*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (35*B*a^2*cos(d*x + c)^3 + 45*(A + 2*B)*a^2*cos(d*x + c)^2 + 14*(9*A + 8*B)*a^2*cos(d*x + c) + 30*(6*A + 5*B)*a^2)*sqrt(cos(d*x + c))*sin(d*x + c))/d`

3.130.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+a*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.130.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2), x)`

3.130.8 Giac [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2), x)`

3.130.9 Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.37

$$\begin{aligned}
& \int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx \\
&= \frac{2Aa^2\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)\right)}{3d} \\
&\quad - \frac{4Aa^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Aa^2\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Ba^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{4Ba^2\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Ba^2\cos(c+dx)^{11/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)^2}}
\end{aligned}$$

```
input int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2,x)
```

```
output (2*A*a^2*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2))/
(3*d) - (4*A*a^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/
4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*A*a^2*cos(c + d*x)^(
9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c
+ d*x)^2)^(1/2)) - (2*B*a^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/
2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (4*B*a^2*co
s(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))
/(9*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a^2*cos(c + d*x)^(11/2)*sin(c + d*x)*
hypergeom([1/2, 11/4], 15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2)
)
```

3.131 $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

3.131.1 Optimal result	1283
3.131.2 Mathematica [C] (warning: unable to verify)	1284
3.131.3 Rubi [A] (verified)	1285
3.131.4 Maple [A] (verified)	1289
3.131.5 Fricas [C] (verification not implemented)	1290
3.131.6 Sympy [F(-1)]	1291
3.131.7 Maxima [F]	1291
3.131.8 Giac [F]	1291
3.131.9 Mupad [B] (verification not implemented)	1292

3.131.1 Optimal result

Integrand size = 33, antiderivative size = 161

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{4a^2(4A + 3B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{4a^2(7A + 6B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d}$$

$$+ \frac{4a^2(7A + 6B)\sqrt{\cos(c + dx)} \sin(c + dx)}{21d} + \frac{2a^2(7A + 9B) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{35d}$$

$$+ \frac{2B \cos^{\frac{3}{2}}(c + dx) (a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{7d}$$

```
output 4/5*a^2*(4*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/21*a^2*(7*A+6*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/35*a^2*
(7*A+9*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/7*B*cos(d*x+c)^(3/2)*(a^2+a^2*co
s(d*x+c))*sin(d*x+c)/d+4/21*a^2*(7*A+6*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.131.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.54 (sec) , antiderivative size = 898, normalized size of antiderivative = 5.58

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2 \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(-\frac{(4A + 3B) \cot(c)}{5d} \right.$$

$$+ \frac{(56A + 51B) \cos(dx) \sin(c)}{168d} + \frac{(A + 2B) \cos(2dx) \sin(2c)}{20d} + \frac{B \cos(3dx) \sin(3c)}{56d}$$

$$\left. + \frac{(56A + 51B) \cos(c) \sin(dx)}{168d} + \frac{(A + 2B) \cos(2c) \sin(2dx)}{20d} + \frac{B \cos(3c) \sin(3dx)}{56d} \right)$$

$$\frac{A(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{3a}$$

$$\frac{2B(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{7}$$

$$\frac{2A(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{5d}$$

$$\frac{3B(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d}$$

```
input Integrate[Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]
]
```

output $\text{Sqrt}[\text{Cos}[c + d*x]]*(a + a*\text{Cos}[c + d*x])^2*\text{Sec}[c/2 + (d*x)/2]^4*(-1/5*((4*A + 3*B)*\text{Cot}[c])/d + ((56*A + 51*B)*\text{Cos}[d*x]*\text{Sin}[c])/(168*d) + ((A + 2*B)*\text{Cos}[2*d*x]*\text{Sin}[2*c])/(20*d) + (B*\text{Cos}[3*d*x]*\text{Sin}[3*c])/(56*d) + ((56*A + 51*B)*\text{Cos}[c]*\text{Sin}[d*x])/(168*d) + ((A + 2*B)*\text{Cos}[2*c]*\text{Sin}[2*d*x])/(20*d) + (B*\text{Cos}[3*c]*\text{Sin}[3*d*x])/(56*d)) - (A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(3*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (2*B*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(7*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (2*A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{Sec}[c/2 + (d*x)/2]^4*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]]*\text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\text{Cos}[c]^2*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]) / (\text{Cos}[c]^2 + \text{Sin}[c]^2)) / \text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]]) / (5*d) - (3*B*(a + a*\text{Cos}...$

3.131.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow 3455$$

$$\frac{2}{7} \int \frac{1}{2} \sqrt{\cos(c + dx)}(\cos(c + dx)a + a)(a(7A + 3B) + a(7A + 9B) \cos(c + dx)) dx + \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d}$$

3.131. $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\downarrow 27$$

$$\frac{1}{7} \int \sqrt{\cos(c+dx)} (\cos(c+dx)a+a) (a(7A+3B) + a(7A+9B) \cos(c+dx)) dx + \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{7d}$$

$$\downarrow 3042$$

$$\frac{1}{7} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right) \left(a(7A+3B) + a(7A+9B) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{7d}$$

$$\downarrow 3447$$

$$\frac{1}{7} \int \sqrt{\cos(c+dx)} ((7A+9B) \cos^2(c+dx)a^2 + (7A+3B)a^2 + ((7A+3B)a^2 + (7A+9B)a^2) \cos(c+dx)) dx + \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{7d}$$

$$\downarrow 3042$$

$$\frac{1}{7} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left((7A+9B) \sin\left(c+dx+\frac{\pi}{2}\right)^2 a^2 + (7A+3B)a^2 + ((7A+3B)a^2 + (7A+9B)a^2) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{7d}$$

$$\downarrow 3502$$

$$\frac{1}{7} \left(\frac{2}{5} \int \sqrt{\cos(c+dx)} (7(4A+3B)a^2 + 5(7A+6B) \cos(c+dx)a^2) dx + \frac{2a^2(7A+9B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{7d}$$

$$\downarrow 3042$$

$$\frac{1}{7} \left(\frac{2}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} (7(4A+3B)a^2 + 5(7A+6B) \sin\left(c+dx+\frac{\pi}{2}\right)a^2) dx + \frac{2a^2(7A+9B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a^2 \cos(c+dx) + a^2)}{7d}$$

$$\downarrow 3227$$

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(7A + 6B) \int \cos^{\frac{3}{2}}(c + dx) dx + 7a^2(4A + 3B) \int \sqrt{\cos(c + dx)} dx \right) + \frac{2a^2(7A + 9B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d} \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(7a^2(4A + 3B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5a^2(7A + 6B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx \right) + \frac{2a^2(7A + 9B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d} \\ \downarrow \text{3115}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(7a^2(4A + 3B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5a^2(7A + 6B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d} \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(7a^2(4A + 3B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5a^2(7A + 6B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d} \\ \downarrow \text{3119}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(7A + 6B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + \frac{14a^2(4A + 3B) E\left(\frac{1}{2}(c + dx)\right)}{d} \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d} \\ \downarrow \text{3120}$$

$$\frac{1}{7} \left(\frac{2a^2(7A + 9B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2}{5} \left(\frac{14a^2(4A + 3B) E\left(\frac{1}{2}(c + dx)\right)}{d} + 5a^2(7A + 6B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx)\right)}{3} \right) \right) \right) \\ \frac{2B \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a^2 \cos(c + dx) + a^2)}{7d}$$

input `Int[Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]),x]`

output `(2*B*cos[c + d*x]^(3/2)*(a^2 + a^2*cos[c + d*x])*Sin[c + d*x])/(7*d) + ((2*a^2*(7*A + 9*B)*cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d) + (2*((14*a^2*(4*A + 3*B)*EllipticE[(c + d*x)/2, 2])/d + 5*a^2*(7*A + 6*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/5)/7`

3.131.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Ssin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Ssin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Ssin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Ssin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*Ssin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Ssin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.131.4 Maple [A] (verified)

Time = 11.96 (sec) , antiderivative size = 385, normalized size of antiderivative = 2.39

method	result
default	$- \frac{4 \sqrt{\left(2 \cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a^2 \left(120B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-84A - 348B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots}{\dots}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```


output
$$\begin{aligned} & -4/105*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a^2*(120*B* \\ & \cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^8+(-84*A-348*B)*\sin(1/2*d*x+1/2*c)^6 \\ & * \cos(1/2*d*x+1/2*c)+(224*A+378*B)*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+ \\ & (-91*A-117*B)*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+35*A*(\sin(1/2*d*x+1/ \\ & 2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c) \\ &),2^{(1/2)})-84*A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1 \\ & /2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})+30*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2} \\ &)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-6 \\ & 3*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*Elliptic \\ & E(\cos(1/2*d*x+1/2*c),2^{(1/2)}))/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c) \\ & ^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.131.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.26

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx = \frac{2\left(5i\sqrt{2}(7A+6B)a^2\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))-5i\sqrt{2}(7A+6B)a^2\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))\right)}{\sin(dx+c)\sqrt{\cos(dx+c)}}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm="fracas")`

output
$$\begin{aligned} & -2/105*(5*I*\sqrt{2}*(7*A+6*B)*a^2*\text{weierstrassPInverse}(-4,0,\cos(d*x+c) \\ &)+I*\sin(d*x+c))-5*I*\sqrt{2}*(7*A+6*B)*a^2*\text{weierstrassPInverse}(-4, \\ & 0,\cos(d*x+c)-I*\sin(d*x+c))-21*I*\sqrt{2}*(4*A+3*B)*a^2*\text{weierstra} \\ & \text{ssZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))) + \\ & 21*I*\sqrt{2}*(4*A+3*B)*a^2*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(- \\ & 4,0,\cos(d*x+c)-I*\sin(d*x+c))) - (15*B*a^2*\cos(d*x+c)^2+21*(A+ \\ & 2*B)*a^2*\cos(d*x+c)+10*(7*A+6*B)*a^2)*\sqrt{\cos(d*x+c)}*\sin(d*x+c))/d \end{aligned}$$

3.131.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*cos(d*x+c)**(1/2),x)`

output `Timed out`

3.131.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sqrt(cos(d*x + c)), x)`

3.131.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sqrt(cos(d*x + c)), x)`

3.131.9 Mupad [B] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.43

$$\begin{aligned}
& \int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^2(A+B\cos(c+dx))dx \\
&= \frac{2Aa^2\left(\frac{2\sqrt{\cos(c+dx)}\sin(c+dx)}{3} + \frac{2F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3}\right)}{d} \\
&+ \frac{2Ba^2\left(\sqrt{\cos(c+dx)}\sin(c+dx) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)\right)}{3d} + \frac{2Aa^2E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} \\
&- \frac{2Aa^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)}^2} \\
&- \frac{4Ba^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)}^2} \\
&- \frac{2Ba^2\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)}^2}
\end{aligned}$$

```
input int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2,x)
```

```
output (2*A*a^2*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)
/2, 2))/3))/d + (2*B*a^2*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2
+ (d*x)/2, 2)))/(3*d) + (2*A*a^2*ellipticE(c/2 + (d*x)/2, 2))/d - (2*A*a^2
*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^
2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (4*B*a^2*cos(c + d*x)^(7/2)*sin(c + d*x
)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)
) - (2*B*a^2*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, c
os(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2))
```

3.132
$$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.132.1 Optimal result 1293
 3.132.2 Mathematica [C] (warning: unable to verify) 1294
 3.132.3 Rubi [A] (verified) 1294
 3.132.4 Maple [B] (verified) 1298
 3.132.5 Fricas [C] (verification not implemented) 1299
 3.132.6 Sympy [F(-1)] 1299
 3.132.7 Maxima [F] 1300
 3.132.8 Giac [F] 1300
 3.132.9 Mupad [B] (verification not implemented) 1301

3.132.1 Optimal result

Integrand size = 33, antiderivative size = 126

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^2(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{4a^2(5A + 4B)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d} + \frac{4a^2(2A + B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} \\ &+ \frac{2a^2(5A + 7B)\sqrt{\cos(c + dx)} \sin(c + dx)}{15d} \\ &+ \frac{2B\sqrt{\cos(c + dx)}(a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{5d} \end{aligned}$$

```
output 4/5*a^2*(5*A+4*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/3*a^2*(2*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/
2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/15*a^2*(5*
A+7*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d+2/5*B*(a^2+a^2*cos(d*x+c))*sin(d*x+c)
*cos(d*x+c)^(1/2)/d
```

3.132.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.14 (sec) , antiderivative size = 369, normalized size of antiderivative = 2.93

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{a^2(1 + \cos(c + dx))^2 \sec^4\left(\frac{1}{2}(c + dx)\right) \left(-6(5A + 4B) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right)\right) \sec(c) \sin(c)}{\dots}$$

input `Integrate[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(a^2*(1 + Cos[c + d*x])^2*Sec[(c + d*x)/2]^4*(-6*(5*A + 4*B)*HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sec[c]*Sin[d*x + ArcTan[Tan[c]]] + (9*(5*A + 4*B)*Cos[c - d*x - ArcTan[Tan[c]]]*Csc[c]*Sec[c] + 15*A*Cos[c + d*x + ArcTan[Tan[c]]]*Csc[c]*Sec[c] + 12*B*Cos[c + d*x + ArcTan[Tan[c]]]*Csc[c]*Sec[c] - 60*A*Cos[c + d*x]*Cot[c]*Sqrt[Sec[c]^2] - 4*8*B*Cos[c + d*x]*Cot[c]*Sqrt[Sec[c]^2] - 20*(2*A + B)*Cos[c + d*x]*Sqrt[Cos[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Csc[c]^2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Sec[c]^2]*Sec[d*x - ArcTan[Cot[c]]]*Sin[c] + 10*A*Cos[c + d*x]*Sqrt[Sec[c]^2]*Sin[c + d*x] + 20*B*Cos[c + d*x]*Sqrt[Sec[c]^2]*Sin[c + d*x] + 6*B*Cos[c + d*x]^2*Sqrt[Sec[c]^2]*Sin[c + d*x])*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2]))/(60*d*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c]^2]*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2])`

3.132.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3455, 27, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

3.132. $\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3455} \\
& \frac{2}{5} \int \frac{(\cos(c + dx)a + a)(a(5A + B) + a(5A + 7B) \cos(c + dx))}{2\sqrt{\cos(c + dx)}} dx + \\
& \quad \frac{2B \sin(c + dx) \sqrt{\cos(c + dx)} (a^2 \cos(c + dx) + a^2)}{5d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{(\cos(c + dx)a + a)(a(5A + B) + a(5A + 7B) \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx + \\
& \quad \frac{2B \sin(c + dx) \sqrt{\cos(c + dx)} (a^2 \cos(c + dx) + a^2)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)(a(5A + B) + a(5A + 7B) \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2B \sin(c + dx) \sqrt{\cos(c + dx)} (a^2 \cos(c + dx) + a^2)}{5d} \\
& \quad \downarrow \text{3447} \\
& \frac{1}{5} \int \frac{(5A + 7B) \cos^2(c + dx)a^2 + (5A + B)a^2 + ((5A + B)a^2 + (5A + 7B)a^2) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \\
& \quad \frac{2B \sin(c + dx) \sqrt{\cos(c + dx)} (a^2 \cos(c + dx) + a^2)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{(5A + 7B) \sin(c + dx + \frac{\pi}{2})^2 a^2 + (5A + B)a^2 + ((5A + B)a^2 + (5A + 7B)a^2) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2B \sin(c + dx) \sqrt{\cos(c + dx)} (a^2 \cos(c + dx) + a^2)}{5d} \\
& \quad \downarrow \text{3502} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{5(2A + B)a^2 + 3(5A + 4B) \cos(c + dx)a^2}{\sqrt{\cos(c + dx)}} dx + \frac{2a^2(5A + 7B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + \\
& \quad \frac{2B \sin(c + dx) \sqrt{\cos(c + dx)} (a^2 \cos(c + dx) + a^2)}{5d} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{5(2A+B)a^2 + 3(5A+4B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a^2(5A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}(a^2\cos(c+dx)+a^2)}{5d}$$

↓ 3227

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(2A+B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^2(5A+4B) \int \sqrt{\cos(c+dx)} dx \right) + \frac{2a^2(5A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}(a^2\cos(c+dx)+a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(2A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a^2(5A+4B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) + \frac{2a^2(5A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}(a^2\cos(c+dx)+a^2)}{5d}$$

↓ 3119

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(2A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^2(5A+4B)E(\frac{1}{2}(c+dx)|2)}{d} \right) + \frac{2a^2(5A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}(a^2\cos(c+dx)+a^2)}{5d}$$

↓ 3120

$$\frac{1}{5} \left(\frac{2a^2(5A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} + \frac{2}{3} \left(\frac{10a^2(2A+B)\text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6a^2(5A+4B)E(\frac{1}{2}(c+dx)|2)}{d} \right) \right) + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}(a^2\cos(c+dx)+a^2)}{5d}$$

input `Int[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(2*B*Sqrt[Cos[c + d*x]]*(a^2 + a^2*Cos[c + d*x])*Sin[c + d*x])/(5*d) + ((2*((6*a^2*(5*A + 4*B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^2*(2*A + B)*EllipticF[(c + d*x)/2, 2])/d))/3 + (2*a^2*(5*A + 7*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

3.132.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`


```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.132.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs. 2(166) = 332.

Time = 8.46 (sec) , antiderivative size = 357, normalized size of antiderivative = 2.83

method	result
default	$-\frac{4\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}a^2\left(-12B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\left(10A+32B\right)\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\left(\frac{1}{2}-\frac{1}{2}\right)}$
parts	$-\frac{2\left(Aa^2+2Ba^2\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{1}{2}}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(-12*B*c
os(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+(10*A+32*B)*sin(1/2*d*x+1/2*c)^4*co
s(1/2*d*x+1/2*c)+(-5*A-13*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+10*A*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))-15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x
+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*B*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2
*c),2^(1/2))-12*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(
1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2
)/d
```

$$3.132. \int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.132.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.42

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx =$$

$$\frac{2 \left(5i \sqrt{2} (2A + B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (2A + B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - 3i \sqrt{2} (5A + 4B) a^2 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 3i \sqrt{2} (5A + 4B) a^2 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - (3B a^2 \cos(dx + c) + 5(A + 2B) a^2) \sqrt{\cos(dx + c)} \sin(dx + c) \right)}{d}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `-2/15*(5*I*sqrt(2)*(2*A + B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(2*A + B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(5*A + 4*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(5*A + 4*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (3*B*a^2*cos(d*x + c) + 5*(A + 2*B)*a^2)*sqrt(cos(d*x + c))*sin(d*x + c))/d`

3.132.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Timed out`

3.132.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/sqrt(cos(d*x + c)), x)`

3.132.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/sqrt(cos(d*x + c)), x)`

3.132.9 Mupad [B] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.21

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2 B a^2 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3} \right)}{d}$$

$$+ \frac{2 A a^2 \left(\sqrt{\cos(c + dx)} \sin(c + dx) + 6 E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + 4 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) \right)}{3 d}$$

$$+ \frac{2 B a^2 E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} - \frac{2 B a^2 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^(1/2),x)
```

```
output (2*B*a^2*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (2*A*a^2*(cos(c + d*x)^(1/2)*sin(c + d*x) + 6*ellipticE(c/2 + (d*x)/2, 2) + 4*ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*B*a^2*ellipticE(c/2 + (d*x)/2, 2))/d - (2*B*a^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))
```

3.133
$$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.133.1 Optimal result 1302
 3.133.2 Mathematica [C] (warning: unable to verify) 1302
 3.133.3 Rubi [A] (verified) 1304
 3.133.4 Maple [A] (verified) 1307
 3.133.5 Fricas [C] (verification not implemented) 1308
 3.133.6 Sympy [F(-1)] 1308
 3.133.7 Maxima [F] 1309
 3.133.8 Giac [F] 1309
 3.133.9 Mupad [B] (verification not implemented) 1310

3.133.1 Optimal result

Integrand size = 33, antiderivative size = 118

$$\int \frac{(a + a \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{4a^2BE(\frac{1}{2}(c + dx)|2)}{d} + \frac{4a^2(3A + 2B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$- \frac{2a^2(3A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{3d} + \frac{2A(a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{d\sqrt{\cos(c + dx)}}$$

output

```
4*a^2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/3*a^2*(3*A+2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*A*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(1/2)-2/3*a^2*(3*A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.133.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.37 (sec) , antiderivative size = 623, normalized size of antiderivative = 5.28

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^2 \sec^4 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{(-A + 2B + A \cos(2c) + 2B \cos(2c)) \csc(c) \sec(c)}{4d} + \frac{B \cos(dx) \sin(c)}{6d} + \frac{B \cos(c) \sin(dx)}{6d} + \frac{A \sec(c) \sec(c + dx) \sin(dx)}{2d} \right)$$

$$\frac{A(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{d}$$

$$\frac{2B(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{3d}$$

$$\frac{B(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \right)}{2d}$$

input `Integrate[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2), x]`

output `Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2*Sec[c/2 + (d*x)/2]^4*(-1/4*((-A + 2*B + A*Cos[2*c] + 2*B*Cos[2*c])*Csc[c]*Sec[c])/d + (B*Cos[d*x]*Sin[c])/(6*d) + (B*Cos[c]*Sin[d*x])/(6*d) + (A*Sec[c]*Sec[c + d*x]*Sin[d*x])/(2*d) - (A*(a + a*Cos[c + d*x])^2*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^4*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]])*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]])]/(d*Sqrt[1 + Cot[c]^2]) - (2*B*(a + a*Cos[c + d*x])^2*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^4*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]])*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]])]/(3*d*Sqrt[1 + Cot[c]^2]) - (B*(a + a*Cos[c + d*x])^2*Csc[c]*Sec[c/2 + (d*x)/2]^4*(HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]])*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]])*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2])*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]))/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]))/(2*d)`

3.133. $\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$

3.133.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3454, 27, 3042, 3447, 3042, 3502, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a \cos(c + dx) + a)^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\
 & \quad \downarrow \text{3454} \\
 & 2 \int \frac{(\cos(c + dx)a + a)(a(3A + B) - a(3A - B) \cos(c + dx))}{2\sqrt{\cos(c + dx)}} dx + \\
 & \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(\cos(c + dx)a + a)(a(3A + B) - a(3A - B) \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx + \\
 & \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(\sin(c + dx + \frac{\pi}{2}) a + a) (a(3A + B) - a(3A - B) \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \\
 & \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{-((3A - B) \cos^2(c + dx)a^2) + (3A + B)a^2 + (a^2(3A + B) - a^2(3A - B)) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \\
 & \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{-\left((3A - B) \sin\left(c + dx + \frac{\pi}{2}\right)^2 a^2\right) + (3A + B)a^2 + (a^2(3A + B) - a^2(3A - B)) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \\
& \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3502} \\
& \frac{2}{3} \int \frac{(3A + 2B)a^2 + 3B \cos(c + dx)a^2}{\sqrt{\cos(c + dx)}} dx - \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \\
& \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2}{3} \int \frac{(3A + 2B)a^2 + 3B \sin\left(c + dx + \frac{\pi}{2}\right) a^2}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx - \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \\
& \quad \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3227} \\
& \frac{2}{3} \left(a^2(3A + 2B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3a^2 B \int \sqrt{\cos(c + dx)} dx \right) - \\
& \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2}{3} \left(a^2(3A + 2B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + 3a^2 B \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx \right) - \\
& \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3119} \\
& \frac{2}{3} \left(a^2(3A + 2B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{6a^2 B E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} \right) - \\
& \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3120}
\end{aligned}$$

3.133. $\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$

$$\frac{2a^2(3A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2}{3} \left(\frac{2a^2(3A + 2B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{6a^2 B E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{d \sqrt{\cos(c + dx)}}$$

input `Int[((a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(2*((6*a^2*B*EllipticE[(c + d*x)/2, 2])/d + (2*a^2*(3*A + 2*B)*EllipticF[(c + d*x)/2, 2])/d))/3 - (2*a^2*(3*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d) + (2*A*(a^2 + a^2*cos[c + d*x])*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])`

3.133.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.133.4 Maple [A] (verified)

Time = 7.38 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.07

method	result
default	$4a^2 \left(-2B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 3A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 3A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right)$
parts	$\frac{2(Aa^2 + 2Ba^2) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + 2(2Aa^2)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2), x, method=_RETURNV
ERBOSE)
```

3.133.
$$\int \frac{(a+a \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

output
$$\frac{4/3a^2(-2B\cos(1/2dx+1/2c)\sin(1/2dx+1/2c)^4+3A\cos(1/2dx+1/2c)\sin(1/2dx+1/2c)^2-3A(\sin(1/2dx+1/2c)^2)^{1/2}(2\sin(1/2dx+1/2c)^2-1)^{1/2}\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})+B\cos(1/2dx+1/2c)\sin(1/2dx+1/2c)^2-2B(\sin(1/2dx+1/2c)^2)^{1/2}(2\sin(1/2dx+1/2c)^2-1)^{1/2}\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})+3B(\sin(1/2dx+1/2c)^2)^{1/2}(2\sin(1/2dx+1/2c)^2-1)^{1/2}\text{EllipticE}(\cos(1/2dx+1/2c),2^{1/2}))}{\sin(1/2dx+1/2c)(2\cos(1/2dx+1/2c)^2-1)^{1/2}d}$$

3.133.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.68

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx =$$

$$2 \left(i \sqrt{2} (3A + 2B) a^2 \cos(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - i \sqrt{2} (3A + 2B) a^2 \sin(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) \right) - 3i \sqrt{2} B a^2 \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 3i \sqrt{2} B a^2 \sin(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - (B a^2 \cos(dx + c) + 3A a^2) \sqrt{\cos(dx + c)} \sin(dx + c) / (d \cos(dx + c))$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="fricas")`

output
$$-2/3(I\sqrt{2}(3A + 2B)a^2\cos(dx + c)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I\sin(dx + c)) - I\sqrt{2}(3A + 2B)a^2\cos(dx + c)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I\sin(dx + c)) - 3I\sqrt{2}B a^2\cos(dx + c)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I\sin(dx + c))) + 3I\sqrt{2}B a^2\cos(dx + c)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I\sin(dx + c))) - (B a^2\cos(dx + c) + 3A a^2)\sqrt{\cos(dx + c)}\sin(dx + c))/(d\cos(dx + c))$$

3.133.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

3.133.
$$\int \frac{(a+a \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

output Timed out

3.133.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(3/2), x)`

3.133.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(3/2), x)`

3.133.9 Mupad [B] (verification not implemented)

Time = 1.24 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.14

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 B a^2 \left(\sqrt{\cos(c + dx)} \sin(c + dx) + 6 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) + 4 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) \right)}{3 d}$$

$$+ \frac{2 A a^2 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 A a^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$+ \frac{2 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^(3/2),x)`output `(2*B*a^2*(cos(c + d*x)^(1/2)*sin(c + d*x) + 6*ellipticE(c/2 + (d*x)/2, 2) + 4*ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*A*a^2*ellipticE(c/2 + (d*x)/2, 2))/d + (4*A*a^2*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.134
$$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.134.1 Optimal result 1311
 3.134.2 Mathematica [C] (warning: unable to verify) 1311
 3.134.3 Rubi [A] (verified) 1312
 3.134.4 Maple [B] (verified) 1316
 3.134.5 Fricas [C] (verification not implemented) 1317
 3.134.6 Sympy [F(-1)] 1317
 3.134.7 Maxima [F] 1318
 3.134.8 Giac [F] 1318
 3.134.9 Mupad [B] (verification not implemented) 1319

3.134.1 Optimal result

Integrand size = 33, antiderivative size = 120

$$\int \frac{(a + a \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{4a^2AE(\frac{1}{2}(c + dx)|2)}{d} + \frac{4a^2(2A + 3B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2a^2(5A + 3B) \sin(c + dx)}{3d\sqrt{\cos(c + dx)}} + \frac{2A(a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

output

```
-4*a^2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/3*a^2*(2*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*A*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/3*a^2*(5*A+3*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.134.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.31 (sec) , antiderivative size = 311, normalized size of antiderivative = 2.59

$$\int \frac{(a + a \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{a^2(1 + \cos(c + dx))^2 \sec^4(\frac{1}{2}(c + dx)) \left(12A \cos(c + dx) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right)\right) \sec(c)}{\dots}$$

input `Integrate[((a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(a^2*(1 + Cos[c + d*x])^2*Sec[(c + d*x)/2]^4*(12*A*cos[c + d*x]*HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sec[c]*Sin[d*x + ArcTan[Tan[c]]) + (Csc[c]*(-6*A*cos[c + d*x]*(3*cos[c - d*x - ArcTan[Tan[c]]) + Cos[c + d*x + ArcTan[Tan[c]])]*Sec[c] + (12*A*cos[c] + 2*A*cos[d*x] - 2*A*cos[2*c + d*x] + 12*A*cos[c + 2*d*x] + 3*B*cos[c + 2*d*x] - 3*B*cos[3*c + 2*d*x])*Sqrt[Sec[c]^2)) - 8*(2*A + 3*B)*Cos[c + d*x]^2*Sqrt[Cos[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Csc[c]^2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sqrt[Sec[c]^2]*Sec[d*x - ArcTan[Cot[c]]]*Sin[c])*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2)))/(24*d*cos[c + d*x]^(3/2)*Sqrt[Sec[c]^2]*Sqrt[Sin[d*x + ArcTan[Tan[c]]]^2))`

3.134.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3454, 27, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

$$\downarrow \text{3454}$$

$$\frac{2}{3} \int \frac{(\cos(c + dx)a + a)(a(5A + 3B) - a(A - 3B) \cos(c + dx))}{2 \cos^{\frac{3}{2}}(c + dx) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d \cos^{\frac{3}{2}}(c + dx)}} dx +$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \frac{1}{3} \int \frac{(\cos(c+dx)a+a)(a(5A+3B)-a(A-3B)\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx + \\
& \quad \frac{2A \sin(c+dx)(a^2 \cos(c+dx)+a^2)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a(5A+3B)-a(A-3B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \\
& \quad \frac{2A \sin(c+dx)(a^2 \cos(c+dx)+a^2)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3447} \\
& \frac{1}{3} \int \frac{-((A-3B)\cos^2(c+dx)a^2)+(5A+3B)a^2+(a^2(5A+3B)-a^2(A-3B))\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx + \\
& \quad \frac{2A \sin(c+dx)(a^2 \cos(c+dx)+a^2)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{-((A-3B)\sin(c+dx+\frac{\pi}{2})^2 a^2)+(5A+3B)a^2+(a^2(5A+3B)-a^2(A-3B))\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \\
& \quad \frac{2A \sin(c+dx)(a^2 \cos(c+dx)+a^2)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3500} \\
& \frac{1}{3} \left(2 \int \frac{a^2(2A+3B)-3a^2A \cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{2a^2(5A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) + \\
& \quad \frac{2A \sin(c+dx)(a^2 \cos(c+dx)+a^2)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(2 \int \frac{a^2(2A+3B)-3a^2A \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a^2(5A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) + \\
& \quad \frac{2A \sin(c+dx)(a^2 \cos(c+dx)+a^2)}{3d \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3227}
\end{aligned}$$

$$\frac{1}{3} \left(2 \left(a^2(2A + 3B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - 3a^2 A \int \sqrt{\cos(c + dx)} dx \right) + \frac{2a^2(5A + 3B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left(2 \left(a^2(2A + 3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - 3a^2 A \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) + \frac{2a^2(5A + 3B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3119

$$\frac{1}{3} \left(2 \left(a^2(2A + 3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{6a^2 AE(\frac{1}{2}(c + dx) | 2)}{d} \right) + \frac{2a^2(5A + 3B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + 2 \left(\frac{2a^2(2A + 3B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} - \frac{6a^2 AE(\frac{1}{2}(c + dx) | 2)}{d} \right) \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

input `Int[((a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*A*(a^2 + a^2*cos[c + d*x])*Sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)) + (2*((-6*a^2*A*EllipticE[(c + d*x)/2, 2])/d + (2*a^2*(2*A + 3*B)*EllipticF[(c + d*x)/2, 2])/d) + (2*a^2*(5*A + 3*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])/3`

3.134.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3454 `Int[((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[(-(A*b^2
- a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.134.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 512 vs. 2(162) = 324.

Time = 7.33 (sec) , antiderivative size = 513, normalized size of antiderivative = 4.28

method	result
default	$\frac{4 \left(6 \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right)} (2A+B) \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right)} (7A+3B) \right)}{\dots}$
parts	$\frac{2(Aa^2+2Ba^2) \operatorname{am}^{-1} \left(\frac{dx}{2} + \frac{c}{2} \mid \sqrt{2} \right)}{d} - \frac{2(2Aa^2+Ba^2) \left(-2 \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right)} \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sqrt{\frac{1}{2}} \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right)} \right)}{\dots}$

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/3*(6*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A+B)*cos(1
/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4-(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/
2*c)^2)^(1/2)*(7*A+3*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-2*(2*sin(1
/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*sin(1/2*d*x+1/2*
c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)
)+3*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*EllipticF(cos(1/2*d*x+1/2*
c),2^(1/2)))*sin(1/2*d*x+1/2*c)^2+2*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(
1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(-2*sin(1/
2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+3*A*(-2*sin(1/2*d*x+1/2*c)^4+si
n(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*
c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*(sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2
^(1/2))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2))*a^2/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3
/2)/sin(1/2*d*x+1/2*c)/d
```

3.134.
$$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.134.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.75

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx =$$

$$2 \left(i \sqrt{2} (2A + 3B) a^2 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - i \sqrt{2} (2A + 3B) a^2 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 3 i \sqrt{2} A a^2 \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 3 i \sqrt{2} A a^2 \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - (3(2A + B) a^2 \cos(dx + c) + A a^2) \sqrt{\cos(dx + c)} \sin(dx + c) \right) / (d \cos(dx + c)^2)$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fricas")`

output `-2/3*(I*sqrt(2)*(2*A + 3*B)*a^2*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - I*sqrt(2)*(2*A + 3*B)*a^2*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*A*a^2*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*A*a^2*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (3*(2*A + B)*a^2*cos(d*x + c) + A*a^2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.134.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output `Timed out`

3.134.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(5/2), x)`

3.134.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(5/2), x)`

3.134.9 Mupad [B] (verification not implemented)

Time = 1.94 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.63

$$\begin{aligned}
& \int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\
&= \frac{2 A a^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{2 B a^2 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 B a^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} \\
&+ \frac{4 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{2 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{2 B a^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}
\end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^(5/2),x)`

output `(2*A*a^2*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*a^2*ellipticE(c/2 + (d*x)/2, 2))/d + (4*B*a^2*ellipticF(c/2 + (d*x)/2, 2))/d + (4*A*a^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) + (2*A*a^2*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2)) + (2*B*a^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.135
$$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.135.1 Optimal result 1320
 3.135.2 Mathematica [C] (warning: unable to verify) 1321
 3.135.3 Rubi [A] (verified) 1322
 3.135.4 Maple [B] (verified) 1326
 3.135.5 Fracas [C] (verification not implemented) 1327
 3.135.6 Sympy [F(-1)] 1328
 3.135.7 Maxima [F] 1328
 3.135.8 Giac [F] 1329
 3.135.9 Mupad [B] (verification not implemented) 1329

3.135.1 Optimal result

Integrand size = 33, antiderivative size = 159

$$\int \frac{(a + a \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= -\frac{4a^2(4A + 5B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{4a^2(A + 2B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2a^2(7A + 5B) \sin(c + dx)}{15d \cos^{\frac{3}{2}}(c + dx)} + \frac{4a^2(4A + 5B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

$$+ \frac{2A(a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

```
output -4/5*a^2*(4*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/3*a^2*(A+2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/15*a^2*(7*A+5*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/5*A*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(5/2)+4/5*a^2*(4*A+5*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.135.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.89 (sec) , antiderivative size = 883, normalized size of antiderivative = 5.55

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^2 \sec^4 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(\frac{(4A + 5B) \csc(c) \sec(c)}{5d} + \frac{A \sec(c) \sec^3(c + dx) \sin(dx)}{10d} + \frac{\sec(c) \sec^2(c + dx) (3A \sin(c) + 10A \sin(dx) + 5B \sin(dx))}{30d} + \frac{\sec(c) \sec(c + dx) (10A \sin(c) + 5B \sin(c) + 24A \sin(dx) + 30B \sin(dx))}{30d} \right)$$

$$\frac{A(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{30d}$$

$$\frac{2B(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{30d}$$

$$+ \frac{2A(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{5d}$$

$$+ \frac{B(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{2d}$$

input `Integrate[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output $\text{Sqrt}[\text{Cos}[c + d*x]]*(a + a*\text{Cos}[c + d*x])^2*\text{Sec}[c/2 + (d*x)/2]^4*((4*A + 5*B)*\text{Csc}[c]*\text{Sec}[c])/(5*d) + (A*\text{Sec}[c]*\text{Sec}[c + d*x]^3*\text{Sin}[d*x])/(10*d) + (\text{Sec}[c]*\text{Sec}[c + d*x]^2*(3*A*\text{Sin}[c] + 10*A*\text{Sin}[d*x] + 5*B*\text{Sin}[d*x]))/(30*d) + (\text{Sec}[c]*\text{Sec}[c + d*x]*(10*A*\text{Sin}[c] + 5*B*\text{Sin}[c] + 24*A*\text{Sin}[d*x] + 30*B*\text{Sin}[d*x]))/(30*d) - (A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(3*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (2*B*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(3*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) + (2*A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{Sec}[c/2 + (d*x)/2]^4*((\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])*\text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\text{Cos}[c]^2*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])/(\text{Cos}[c]^2 + \text{Sin}[c]^2))/ \text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]))/(5*d) + (B*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[\dots$

3.135.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.98, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3454, 27, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

↓ 3454

3.135. $\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{2}{5} \int \frac{(\cos(c+dx)a+a)(a(7A+5B)+a(A+5B)\cos(c+dx))}{2\cos^{\frac{5}{2}}(c+dx)} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{(\cos(c+dx)a+a)(a(7A+5B)+a(A+5B)\cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a(7A+5B)+a(A+5B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3447} \\
& \frac{1}{5} \int \frac{(A+5B)\cos^2(c+dx)a^2+(7A+5B)a^2+((A+5B)a^2+(7A+5B)a^2)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{(A+5B)\sin(c+dx+\frac{\pi}{2})^2a^2+(7A+5B)a^2+((A+5B)a^2+(7A+5B)a^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3500} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{3(4A+5B)a^2+5(A+2B)\cos(c+dx)a^2}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a^2(7A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)} \right) + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{3(4A+5B)a^2+5(A+2B)\sin(c+dx+\frac{\pi}{2})a^2}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2a^2(7A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)} \right) + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{5d\cos^{\frac{5}{2}}(c+dx)}
\end{aligned}$$

3.135. $\int \frac{(a+a\cos(c+dx))^2(A+B\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

↓ 3227

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^2(4A + 5B) \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx + 5a^2(A + 2B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \right) + \frac{2a^2(7A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^2(4A + 5B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}}} dx + 5a^2(A + 2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2a^2(7A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3116

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right) \right) + \frac{2a^2(7A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) \right) + \frac{2a^2(7A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3119

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx)|2)}{d} \right) \right) + \frac{2a^2(7A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3120

$$\frac{1}{5} \left(\frac{2a^2(7A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2}{3} \left(\frac{10a^2(A + 2B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2}{d} \right) \right) \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

input `Int[((a + a*cos[c + d*x])^2*(A + B*cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*A*(a^2 + a^2*cos[c + d*x])*Sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)) + ((2*a^2*(7*A + 5*B)*Sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)) + (2*((10*a^2*(A + 2*B)*EllipticF[(c + d*x)/2, 2])/d + 3*a^2*(4*A + 5*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*SIN[c + d*x])/(d*Sqrt[Cos[c + d*x]))))/3)/5`

3.135.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*SIN[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*SIN[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.135.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 713 vs. $2(195) = 390$.

Time = 10.22 (sec) , antiderivative size = 714, normalized size of antiderivative = 4.49

method	result	size
default	Expression too large to display	714
parts	Expression too large to display	801

input `int((a+cos(d*x+c))*a^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)`

$$3.135. \quad \int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$$

output

```

-8*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(1/4*B*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d
*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/
2))+1/20*A/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1
/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^
6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^
4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*s
in(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*s
in(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*si
n(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+(1/2*A+1/4*B)*(-1/6*cos(1/2
*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*
d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c
)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Elliptic
F(cos(1/2*d*x+1/2*c),2^(1/2)))+(1/4*A+1/2*B)/sin(1/2*d*x+1/2*c)^2/(2*sin(1
/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2
*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*s
in(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))/sin(1
/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d

```

3.135.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.50

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{2 \left(5i \sqrt{2} (A + 2B) a^2 \cos(dx + c)^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (A \right.}{\dots}$$

input

```

integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorith
m="fricas")

```

output
$$-2/15*(5*I*sqrt(2)*(A + 2*B)*a^2*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(A + 2*B)*a^2*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(4*A + 5*B)*a^2*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(4*A + 5*B)*a^2*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (6*(4*A + 5*B)*a^2*cos(d*x + c)^2 + 5*(2*A + B)*a^2*cos(d*x + c) + 3*A*a^2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)$$

3.135.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output Timed out

3.135.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{7}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(7/2), x)`

3.135.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(7/2), x)`

3.135.9 Mupad [B] (verification not implemented)

Time = 2.14 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.44

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{6 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right) + 20 A a^2 \cos(c + dx) \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right) + 15 d \cos(c + dx)^{5/2} \sqrt{1 - \cos(c + dx)}}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 B a^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^2 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^(7/2),x)`

output `(6*A*a^2*sin(c + d*x)*hypergeom([-5/4, 1/2], -1/4, cos(c + d*x)^2) + 20*A*a^2*cos(c + d*x)*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2) + 30*A*a^2*cos(c + d*x)^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(15*d*cos(c + d*x)^(5/2)*(1 - cos(c + d*x)^2)^(1/2)) + (2*B*a^2*ellipticF(c/2 + (d*x)/2, 2))/d + (4*B*a^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) + (2*B*a^2*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2))`

3.135. $\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

3.136
$$\int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.136.1 Optimal result 1330
 3.136.2 Mathematica [C] (warning: unable to verify) 1331
 3.136.3 Rubi [A] (verified) 1332
 3.136.4 Maple [B] (verified) 1336
 3.136.5 Fracas [C] (verification not implemented) 1337
 3.136.6 Sympy [F(-1)] 1338
 3.136.7 Maxima [F] 1338
 3.136.8 Giac [F] 1339
 3.136.9 Mupad [B] (verification not implemented) 1339

3.136.1 Optimal result

Integrand size = 33, antiderivative size = 194

$$\int \frac{(a + a \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= -\frac{4a^2(3A + 4B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{4a^2(6A + 7B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d}$$

$$+ \frac{2a^2(9A + 7B) \sin(c + dx)}{35d \cos^{\frac{5}{2}}(c + dx)} + \frac{4a^2(6A + 7B) \sin(c + dx)}{21d \cos^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{4a^2(3A + 4B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}} + \frac{2A(a^2 + a^2 \cos(c + dx)) \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx)}$$

output

```
-4/5*a^2*(3*A+4*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/21*a^2*(6*A+7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/35*a^2*(9*A+7*B)*sin(d*x+c)/d/cos(d*x+c)^(5/2)+4/21*a^2*(6*A+7*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/7*A*(a^2+a^2*cos(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(7/2)+4/5*a^2*(3*A+4*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.136.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.18 (sec) , antiderivative size = 925, normalized size of antiderivative = 4.77

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^2 \sec^4 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(\frac{(3A + 4B) \csc(c) \sec(c)}{5d} + \frac{A \sec(c) \sec^4(c + dx) \sin(dx)}{14d} + \frac{\sec(c) \sec^3(c + dx) (5A \sin(c) + 14A \sin(dx) + 7B \sin(dx))}{70d} + \frac{\sec(c) \sec^2(c + dx) (42A \sin(c) + 21B \sin(c) + 60A \sin(dx) + 70B \sin(dx))}{210d} + \frac{\sec(c) \sec(c + dx) (30A \sin(c) + 35B \sin(c) + 63A \sin(dx) + 84B \sin(dx))}{105d} \right)$$

$$- \frac{2A(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{7d}$$

$$- \frac{B(a + a \cos(c + dx))^2 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{3d}$$

$$+ \frac{3A(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{10d}$$

$$+ \frac{2B(a + a \cos(c + dx))^2 \csc(c) \sec^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{5d}$$

input `Integrate[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output $\text{Sqrt}[\text{Cos}[c + d*x]]*(a + a*\text{Cos}[c + d*x])^2*\text{Sec}[c/2 + (d*x)/2]^4*((3*A + 4*B)*\text{Csc}[c]*\text{Sec}[c])/(5*d) + (A*\text{Sec}[c]*\text{Sec}[c + d*x]^4*\text{Sin}[d*x])/(14*d) + (\text{Sec}[c]*\text{Sec}[c + d*x]^3*(5*A*\text{Sin}[c] + 14*A*\text{Sin}[d*x] + 7*B*\text{Sin}[d*x]))/(70*d) + (\text{Sec}[c]*\text{Sec}[c + d*x]^2*(42*A*\text{Sin}[c] + 21*B*\text{Sin}[c] + 60*A*\text{Sin}[d*x] + 70*B*\text{Sin}[d*x]))/(210*d) + (\text{Sec}[c]*\text{Sec}[c + d*x]*(30*A*\text{Sin}[c] + 35*B*\text{Sin}[c] + 63*A*\text{Sin}[d*x] + 84*B*\text{Sin}[d*x]))/(105*d) - (2*A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(7*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (B*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^4*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(3*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) + (3*A*(a + a*\text{Cos}[c + d*x])^2*\text{Csc}[c]*\text{Sec}[c/2 + (d*x)/2]^4*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]*\text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\text{Cos}[c]^2*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])/(\text{Cos}[c]^2 + \text{Sin}[c]^2))/\text{Sq}...$

3.136.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.96, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3454, 27, 3042, 3447, 3042, 3500, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{\frac{9}{2}}} dx$$

↓ 3454

3.136. $\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{2}{7} \int \frac{(\cos(c+dx)a+a)(a(9A+7B)+a(3A+7B)\cos(c+dx))}{2\cos^{\frac{7}{2}}(c+dx)} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 27 \\
& \frac{1}{7} \int \frac{(\cos(c+dx)a+a)(a(9A+7B)+a(3A+7B)\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a(9A+7B)+a(3A+7B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3447 \\
& \frac{1}{7} \int \frac{(3A+7B)\cos^2(c+dx)a^2+(9A+7B)a^2+((3A+7B)a^2+(9A+7B)a^2)\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \int \frac{(3A+7B)\sin(c+dx+\frac{\pi}{2})^2 a^2+(9A+7B)a^2+((3A+7B)a^2+(9A+7B)a^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}} dx + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3500 \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{5(6A+7B)a^2+7(3A+4B)\cos(c+dx)a^2}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^2(9A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)} \right) + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{5(6A+7B)a^2+7(3A+4B)\sin(c+dx+\frac{\pi}{2})a^2}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2a^2(9A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)} \right) + \\
& \quad \frac{2A\sin(c+dx)(a^2\cos(c+dx)+a^2)}{7d\cos^{\frac{7}{2}}(c+dx)}
\end{aligned}$$

3.136. $\int \frac{(a+a\cos(c+dx))^2(A+B\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

↓ 3227

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(6A + 7B) \int \frac{1}{\cos^{\frac{5}{2}}(c + dx)} dx + 7a^2(3A + 4B) \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx \right) + \frac{2a^2(9A + 7B) \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(6A + 7B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + 7a^2(3A + 4B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \frac{2a^2(9A + 7B) \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3116

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(6A + 7B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + 7a^2(3A + 4B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right) \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(6A + 7B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + 7a^2(3A + 4B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right) \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3119

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(6A + 7B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + 7a^2(3A + 4B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E}{d} \right) \right) + \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3120

$$\frac{1}{7} \left(\frac{2a^2(9A + 7B) \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} + \frac{2}{5} \left(5a^2(6A + 7B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + 7a^2(3A + 4B) \frac{2A \sin(c + dx) (a^2 \cos(c + dx) + a^2)}{7d \cos^{\frac{7}{2}}(c + dx)} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])/Cos[c + d*x]^(9/2),x]`

output `(2*A*(a^2 + a^2*Cos[c + d*x])*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*a^2*(9*A + 7*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (2*(5*a^2*(6*A + 7*B))*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))) + 7*a^2*(3*A + 4*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/5/7`

3.136.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 $\text{Int}[(b \cdot \sin(e) + f \cdot x)^m \cdot (c + d \cdot \sin(e) + f \cdot x)], x_{\text{Symbol}}] \rightarrow \text{Simp}[c \cdot \text{Int}[(b \cdot \sin(e) + f \cdot x)^m, x], x] + \text{Simp}[d/b \cdot \text{Int}[(b \cdot \sin(e) + f \cdot x)^{m+1}, x], x] /;$ $\text{FreeQ}\{b, c, d, e, f, m, x\}$

rule 3447 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) \cdot (c + d \cdot \sin(e) + f \cdot x)], x_{\text{Symbol}}] \rightarrow \text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A \cdot c + (B \cdot c + A \cdot d) \cdot \sin(e) + f \cdot x + B \cdot d \cdot \sin(e) + f \cdot x)^2], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, A, B, m, x\}$ && $\text{NeQ}[b \cdot c - a \cdot d, 0]$

rule 3454 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) \cdot (c + d \cdot \sin(e) + f \cdot x)^n], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-b^2) \cdot (B \cdot c - A \cdot d) \cdot \cos(e) + f \cdot x \cdot (a + b \cdot \sin(e) + f \cdot x)^{m-1} \cdot (c + d \cdot \sin(e) + f \cdot x)^{n+1} / (d \cdot f \cdot (n+1) \cdot (b \cdot c + a \cdot d)), x] - \text{Simp}[b / (d \cdot (n+1) \cdot (b \cdot c + a \cdot d)) \cdot \text{Int}[(a + b \cdot \sin(e) + f \cdot x)^{m-1} \cdot (c + d \cdot \sin(e) + f \cdot x)^{n+1} \cdot \text{Simp}[a \cdot A \cdot d \cdot (m - n - 2) - B \cdot (a \cdot c \cdot (m - 1) + b \cdot d \cdot (n + 1)) - (A \cdot b \cdot d \cdot (m + n + 1) - B \cdot (b \cdot c \cdot m - a \cdot d \cdot (n + 1))) \cdot \sin(e) + f \cdot x], x], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, A, B, x\}$ && $\text{NeQ}[b \cdot c - a \cdot d, 0]$ && $\text{EqQ}[a^2 - b^2, 0]$ && $\text{NeQ}[c^2 - d^2, 0]$ && $\text{GtQ}[m, 1/2]$ && $\text{LtQ}[n, -1]$ && $\text{IntegerQ}[2 \cdot m]$ && $(\text{IntegerQ}[2 \cdot n] \parallel \text{EqQ}[c, 0])$

rule 3500 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) \cdot (C + d \cdot \sin(e) + f \cdot x)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C) \cdot \cos(e) + f \cdot x \cdot (a + b \cdot \sin(e) + f \cdot x)^{m+1} / (b \cdot f \cdot (m+1) \cdot (a^2 - b^2)), x] + \text{Simp}[1 / (b \cdot (m+1) \cdot (a^2 - b^2)) \cdot \text{Int}[(a + b \cdot \sin(e) + f \cdot x)^{m+1} \cdot \text{Simp}[b \cdot (a \cdot A - b \cdot B + a \cdot C) \cdot (m+1) - (A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C + b \cdot (A \cdot b - a \cdot B + b \cdot C) \cdot (m+1)) \cdot \sin(e) + f \cdot x], x], x], x] /;$ $\text{FreeQ}\{a, b, e, f, A, B, C, x\}$ && $\text{LtQ}[m, -1]$ && $\text{NeQ}[a^2 - b^2, 0]$

3.136.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 823 vs. $2(226) = 452$.

Time = 13.64 (sec) , antiderivative size = 824, normalized size of antiderivative = 4.25

method	result	size
default	Expression too large to display	824
parts	Expression too large to display	1026

3.136.
$$\int \frac{(a + a \cos(c+dx))^2 (A + B \cos(c+dx))}{\cos^2(c+dx)} dx$$

input `int((a+cos(d*x+c))*a)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RETURNV
ERBOSE)`

output `-8*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(1/4*A*(-
1/56*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/
2))/(cos(1/2*d*x+1/2*c)^2-1/2)^4-5/42*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+/
2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+5/21*(sin(
1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+/
1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))
) + 1/4*B/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+/
2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/
2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elliptic
E(cos(1/2*d*x+1/2*c),2^(1/2)))+(1/4*A+1/2*B)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*
sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2
)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2)))+1/5*(1/2*A+1/4*B)/sin(1/2*d*x+1/2*c)^2/(8*sin(1/2*d*x+1/2*c
)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)*(24*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-
24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*si
n(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2...`

3.136.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.36

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx =$$

$$\frac{2 \left(5i \sqrt{2} (6A + 7B) a^2 \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (6 \right)}{\dots}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm
m="fracas")`

output `-2/105*(5*I*sqrt(2)*(6*A + 7*B)*a^2*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(6*A + 7*B)*a^2*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*I*sqrt(2)*(3*A + 4*B)*a^2*cos(d*x + c)^4*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*(3*A + 4*B)*a^2*cos(d*x + c)^4*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (42*(3*A + 4*B)*a^2*cos(d*x + c)^3 + 10*(6*A + 7*B)*a^2*cos(d*x + c)^2 + 21*(2*A + B)*a^2*cos(d*x + c) + 15*A*a^2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^4)`

3.136.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output `Timed out`

3.136.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{9}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(9/2), x)`

3.136.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{9}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/cos(d*x + c)^(9/2), x)`

3.136.9 Mupad [B] (verification not implemented)

Time = 2.59 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.21

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{30 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{7}{4}, \frac{1}{2}; -\frac{3}{4}; \cos(c + dx)^2\right) + 84 A a^2 \cos(c + dx) \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right)}{105 d \cos(c + dx)^{7/2} \sqrt{1 - \cos(c + dx)}} + \frac{6 B a^2 \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right) + 20 B a^2 \cos(c + dx) \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{15 d \cos(c + dx)^{5/2} \sqrt{1 - \cos(c + dx)}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/cos(c + d*x)^(9/2),x)`

output `(30*A*a^2*sin(c + d*x)*hypergeom([-7/4, 1/2], -3/4, cos(c + d*x)^2) + 84*A*a^2*cos(c + d*x)*sin(c + d*x)*hypergeom([-5/4, 1/2], -1/4, cos(c + d*x)^2) + 70*A*a^2*cos(c + d*x)^2*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(105*d*cos(c + d*x)^(7/2)*(1 - cos(c + d*x)^2)^(1/2)) + (6*B*a^2*sin(c + d*x)*hypergeom([-5/4, 1/2], -1/4, cos(c + d*x)^2) + 20*B*a^2*cos(c + d*x)*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2) + 30*B*a^2*cos(c + d*x)^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(15*d*cos(c + d*x)^(5/2)*(1 - cos(c + d*x)^2)^(1/2))`

3.137 $\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$

3.137.1 Optimal result	1340
3.137.2 Mathematica [C] (warning: unable to verify)	1341
3.137.3 Rubi [A] (verified)	1342
3.137.4 Maple [A] (verified)	1347
3.137.5 Fricas [C] (verification not implemented)	1348
3.137.6 Sympy [F(-1)]	1349
3.137.7 Maxima [F]	1349
3.137.8 Giac [F]	1349
3.137.9 Mupad [B] (verification not implemented)	1350

3.137.1 Optimal result

Integrand size = 33, antiderivative size = 237

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$$

$$= \frac{4a^3(17A+15B)E(\frac{1}{2}(c+dx)|2)}{15d} + \frac{4a^3(121A+105B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{231d}$$

$$+ \frac{4a^3(121A+105B)\sqrt{\cos(c+dx)}\sin(c+dx)}{231d}$$

$$+ \frac{4a^3(17A+15B)\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{45d}$$

$$+ \frac{20a^3(22A+21B)\cos^{\frac{5}{2}}(c+dx)\sin(c+dx)}{693d}$$

$$+ \frac{2aB\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2\sin(c+dx)}{11d}$$

$$+ \frac{2(11A+15B)\cos^{\frac{5}{2}}(c+dx)(a^3+a^3 \cos(c+dx))\sin(c+dx)}{99d}$$

```
output 4/15*a^3*(17*A+15*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellip
ticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/231*a^3*(121*A+105*B)*(cos(1/2*d*x+1/
2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+4
/45*a^3*(17*A+15*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+20/693*a^3*(22*A+21*B)*c
os(d*x+c)^(5/2)*sin(d*x+c)/d+2/11*a*B*cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))^2*
sin(d*x+c)/d+2/99*(11*A+15*B)*cos(d*x+c)^(5/2)*(a^3+a^3*cos(d*x+c))*sin(d*
x+c)/d+4/231*a^3*(121*A+105*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.137. $\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3(A+B \cos(c+dx)) dx$

3.137.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.68 (sec) , antiderivative size = 990, normalized size of antiderivative = 4.18

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3 \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(-\frac{(17A + 15B) \cot(c)}{30d} \right.$$

$$+ \frac{(2134A + 1953B) \cos(dx) \sin(c)}{7392d} + \frac{(73A + 75B) \cos(2dx) \sin(2c)}{720d}$$

$$+ \frac{3(44A + 63B) \cos(3dx) \sin(3c)}{4928d} + \frac{(A + 3B) \cos(4dx) \sin(4c)}{288d} + \frac{B \cos(5dx) \sin(5c)}{704d}$$

$$+ \frac{(2134A + 1953B) \cos(c) \sin(dx)}{7392d} + \frac{(73A + 75B) \cos(2c) \sin(2dx)}{720d}$$

$$\left. + \frac{3(44A + 63B) \cos(3c) \sin(3dx)}{4928d} + \frac{(A + 3B) \cos(4c) \sin(4dx)}{288d} + \frac{B \cos(5c) \sin(5dx)}{704d} \right)$$

$$\frac{11A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{4d}$$

$$\frac{5B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{22d}$$

$$\frac{17A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{60d}$$

$$\frac{B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{4d}$$

```
input Integrate[Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x
]
```

output `Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^3*Sec[c/2 + (d*x)/2]^6*(-1/30*((17 *A + 15*B)*Cot[c])/d + ((2134*A + 1953*B)*Cos[d*x]*Sin[c])/(7392*d) + ((73 *A + 75*B)*Cos[2*d*x]*Sin[2*c])/(720*d) + (3*(44*A + 63*B)*Cos[3*d*x]*Sin[3*c])/(4928*d) + ((A + 3*B)*Cos[4*d*x]*Sin[4*c])/(288*d) + (B*cos[5*d*x]*S in[5*c])/(704*d) + ((2134*A + 1953*B)*Cos[c]*Sin[d*x])/(7392*d) + ((73*A + 75*B)*Cos[2*c]*Sin[2*d*x])/(720*d) + (3*(44*A + 63*B)*Cos[3*c]*Sin[3*d*x])/(4928*d) + ((A + 3*B)*Cos[4*c]*Sin[4*d*x])/(288*d) + (B*cos[5*c]*Sin[5*d *x])/(704*d)) - (11*A*(a + a*cos[c + d*x])^3*Csc[c]*HypergeometricPFQ[{1/4 , 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(42*d*Sqrt[1 + Cot[c]^2]) - (5*B*(a + a*cos[c + d*x])^3*Csc[c]*Hypergeo metricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/ 2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - A rcTan[Cot[c]]])]/(22*d*Sqrt[1 + Cot[c]^2]) - (17*A*(a + a*cos[c + d*x])^3* Csc[c]*Sec[c/2 + (d*x)/2]^6*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d *x + ArcTan[Tan[c]]]^2]*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d* x + ArcTan[Tan[c]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]*Sqrt[Cos[c]*Cos[d *x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*...`

3.137.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3455, 27, 3042, 3455, 3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3455}$$

3.137. $\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

$$\frac{2}{11} \int \frac{1}{2} \cos^{\frac{3}{2}}(c+dx) (\cos(c+dx)a+a)^2 (a(11A+5B) + a(11A+15B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^2}{11d}$$

↓ 27

$$\frac{1}{11} \int \cos^{\frac{3}{2}}(c+dx) (\cos(c+dx)a+a)^2 (a(11A+5B) + a(11A+15B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^2}{11d}$$

↓ 3042

$$\frac{1}{11} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2 \left(a(11A+5B) + a(11A+15B) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^2}{11d}$$

↓ 3455

$$\frac{1}{11} \left(\frac{2}{9} \int \cos^{\frac{3}{2}}(c+dx) (\cos(c+dx)a+a) ((77A+60B)a^2 + 5(22A+21B) \cos(c+dx)a^2) dx + \frac{2(11A+15B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^2}{11d} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{2}{9} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right) ((77A+60B)a^2 + 5(22A+21B) \sin\left(c+dx+\frac{\pi}{2}\right)a^2) dx + \frac{2aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^2}{11d} \right)$$

↓ 3447

$$\frac{1}{11} \left(\frac{2}{9} \int \cos^{\frac{3}{2}}(c+dx) (5(22A+21B) \cos^2(c+dx)a^3 + (77A+60B)a^3 + (5(22A+21B)a^3 + (77A+60B)a^3) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^2}{11d} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{2}{9} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} \left(5(22A + 21B) \sin \left(c + dx + \frac{\pi}{2} \right)^2 a^3 + (77A + 60B)a^3 + (5(22A + 21B)a^3 + (77A + 60B)a^3 \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2}{11d} \right) \\ \downarrow \text{3502}$$

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{2}{7} \int \frac{1}{2} \cos^{\frac{3}{2}}(c + dx) (9(121A + 105B)a^3 + 77(17A + 15B) \cos(c + dx)a^3) dx + \frac{10a^3(22A + 21B) \sin(c + dx)}{7d} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2}{11d} \right) \\ \downarrow \text{27}$$

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \int \cos^{\frac{3}{2}}(c + dx) (9(121A + 105B)a^3 + 77(17A + 15B) \cos(c + dx)a^3) dx + \frac{10a^3(22A + 21B) \sin(c + dx)}{7d} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2}{11d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} \left(9(121A + 105B)a^3 + 77(17A + 15B) \sin \left(c + dx + \frac{\pi}{2} \right) a^3 \right) dx + \frac{10a^3(22A + 21B) \sin(c + dx)}{7d} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2}{11d} \right) \\ \downarrow \text{3227}$$

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \left(9a^3(121A + 105B) \int \cos^{\frac{3}{2}}(c + dx) dx + 77a^3(17A + 15B) \int \cos^{\frac{5}{2}}(c + dx) dx \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2}{11d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \left(9a^3(121A + 105B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} dx + 77a^3(17A + 15B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} dx \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2}{11d} \right) \\ \downarrow \text{3115}$$

3.137. $\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \left(77a^3(17A + 15B) \left(\frac{3}{5} \int \sqrt{\cos(c + dx)} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + 9a^3(121A + 105B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) \right) \right) + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^2}{11d} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \left(77a^3(17A + 15B) \left(\frac{3}{5} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + 9a^3(121A + 105B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) \right) \right) + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^2}{11d} \right)$$

↓ 3119

$$\frac{1}{11} \left(\frac{2}{9} \left(\frac{1}{7} \left(9a^3(121A + 105B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) + 77a^3(17A + 15B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) \right) \right) + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^2}{11d}$$

↓ 3120

$$\frac{1}{11} \left(\frac{2(11A + 15B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a^3 \cos(c + dx) + a^3)}{9d} + \frac{2}{9} \left(\frac{10a^3(22A + 21B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} + \frac{2aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^2}{11d} \right) \right)$$

input `Int[Cos[c + d*x]^(3/2)*(a + a*cos[c + d*x])^3*(A + B*cos[c + d*x]),x]`

output `(2*a*B*cos[c + d*x]^(5/2)*(a + a*cos[c + d*x])^2*sin[c + d*x])/(11*d) + ((2*(11*A + 15*B)*cos[c + d*x]^(5/2)*(a^3 + a^3*cos[c + d*x])*sin[c + d*x])/(9*d) + (2*((10*a^3*(22*A + 21*B)*cos[c + d*x]^(5/2)*sin[c + d*x])/(7*d) + (9*a^3*(121*A + 105*B))*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x])*sin[c + d*x])/(3*d)) + 77*a^3*(17*A + 15*B))*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*cos[c + d*x]^(3/2)*sin[c + d*x])/(5*d)))/7)/9)/11`

3.137.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)^2], x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.137.4 Maple [A] (verified)

Time = 16.94 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.86

method	result
default	$-4\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} a^3 \left(10080B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^{12}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-6160A - 43680B)\left(\sin^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right)$
parts	Expression too large to display

```
input int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```
-4/3465*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(10080
*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^12+(-6160*A-43680*B)*sin(1/2*d*x+
1/2*c)^10*cos(1/2*d*x+1/2*c)+(24200*A+77280*B)*sin(1/2*d*x+1/2*c)^8*cos(1/
2*d*x+1/2*c)+(-37532*A-72240*B)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(2
9722*A+39270*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-8118*A-8820*B)*s
in(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+1815*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-39
27*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2))+1575*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(
1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3465*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1
/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/
2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.137.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.03

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^3(A+B\cos(c+dx))dx =$$

$$\frac{2\left(15i\sqrt{2}(121A+105B)a^3\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))-15i\sqrt{2}(121A+\right.}{-}$$

input

```
integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorith
m="fricas")
```

output

```
-2/3465*(15*I*sqrt(2)*(121*A + 105*B)*a^3*weierstrassPInverse(-4, 0, cos(d
*x + c) + I*sin(d*x + c)) - 15*I*sqrt(2)*(121*A + 105*B)*a^3*weierstrassPI
nverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 231*I*sqrt(2)*(17*A + 15*B)
*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*si
n(d*x + c))) + 231*I*sqrt(2)*(17*A + 15*B)*a^3*weierstrassZeta(-4, 0, weie
rstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (315*B*a^3*cos(d*
x + c)^4 + 385*(A + 3*B)*a^3*cos(d*x + c)^3 + 135*(11*A + 14*B)*a^3*cos(d*
x + c)^2 + 154*(17*A + 15*B)*a^3*cos(d*x + c) + 30*(121*A + 105*B)*a^3)*sq
rt(cos(d*x + c))*sin(d*x + c))/d
```

3.137. $\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^3(A+B\cos(c+dx))dx$

3.137.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+a*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.137.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*cos(d*x + c)^(3/2), x)`

3.137.8 Giac [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*cos(d*x + c)^(3/2), x)`

3.137. $\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.137.9 Mupad [B] (verification not implemented)

Time = 1.38 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.52

$$\begin{aligned}
& \int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^3(A+B\cos(c+dx))dx \\
&= \frac{Aa^3 \left(\frac{2\sqrt{\cos(c+dx)}\sin(c+dx)}{3} + \frac{2F\left(\frac{c+dx}{2}\middle|2\right)}{3} \right)}{d} \\
&\quad - \frac{6Aa^3 \cos(c+dx)^{7/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Aa^3 \cos(c+dx)^{9/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{3d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Aa^3 \cos(c+dx)^{11/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Ba^3 \cos(c+dx)^{7/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Ba^3 \cos(c+dx)^{9/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{3d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{6Ba^3 \cos(c+dx)^{11/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2Ba^3 \cos(c+dx)^{13/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{13}{4}; \frac{17}{4}; \cos(c+dx)^2\right)}{13d\sqrt{\sin(c+dx)^2}}
\end{aligned}$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3,x)`

output $(Aa^3((2\cos(c + dx)^{1/2}\sin(c + dx))/3 + (2\text{ellipticF}(c/2 + (dx)/2, 2))/3))/d - (6Aa^3\cos(c + dx)^{7/2}\sin(c + dx)\text{hypergeom}([1/2, 7/4], 11/4, \cos(c + dx)^2))/(7d(\sin(c + dx)^2)^{1/2}) - (2Aa^3\cos(c + dx)^{9/2}\sin(c + dx)\text{hypergeom}([1/2, 9/4], 13/4, \cos(c + dx)^2))/(3d(\sin(c + dx)^2)^{1/2}) - (2Aa^3\cos(c + dx)^{11/2}\sin(c + dx)\text{hypergeom}([1/2, 11/4], 15/4, \cos(c + dx)^2))/(11d(\sin(c + dx)^2)^{1/2}) - (2Ba^3\cos(c + dx)^{7/2}\sin(c + dx)\text{hypergeom}([1/2, 7/4], 11/4, \cos(c + dx)^2))/(7d(\sin(c + dx)^2)^{1/2}) - (2Ba^3\cos(c + dx)^{9/2}\sin(c + dx)\text{hypergeom}([1/2, 9/4], 13/4, \cos(c + dx)^2))/(3d(\sin(c + dx)^2)^{1/2}) - (6Ba^3\cos(c + dx)^{11/2}\sin(c + dx)\text{hypergeom}([1/2, 11/4], 15/4, \cos(c + dx)^2))/(11d(\sin(c + dx)^2)^{1/2}) - (2Ba^3\cos(c + dx)^{13/2}\sin(c + dx)\text{hypergeom}([1/2, 13/4], 17/4, \cos(c + dx)^2))/(13d(\sin(c + dx)^2)^{1/2})$

3.138 $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.138.1 Optimal result	1352
3.138.2 Mathematica [C] (warning: unable to verify)	1353
3.138.3 Rubi [A] (verified)	1354
3.138.4 Maple [A] (verified)	1359
3.138.5 Fricas [C] (verification not implemented)	1360
3.138.6 Sympy [F(-1)]	1361
3.138.7 Maxima [F]	1361
3.138.8 Giac [F]	1361
3.138.9 Mupad [B] (verification not implemented)	1362

3.138.1 Optimal result

Integrand size = 33, antiderivative size = 204

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{4a^3(21A + 17B)E(\frac{1}{2}(c + dx)|2)}{15d} + \frac{4a^3(13A + 11B)\text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d}$$

$$+ \frac{4a^3(13A + 11B)\sqrt{\cos(c + dx)}\sin(c + dx)}{21d} + \frac{4a^3(24A + 23B)\cos^{\frac{3}{2}}(c + dx)\sin(c + dx)}{105d}$$

$$+ \frac{2aB\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2\sin(c + dx)}{9d}$$

$$+ \frac{2(9A + 13B)\cos^{\frac{3}{2}}(c + dx)(a^3 + a^3 \cos(c + dx))\sin(c + dx)}{63d}$$

```
output 4/15*a^3*(21*A+17*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellip
ticE(sin(1/2*d*x+1/2*c), 2^(1/2))/d+4/21*a^3*(13*A+11*B)*(cos(1/2*d*x+1/2*c
)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/d+4/10
5*a^3*(24*A+23*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/9*a*B*cos(d*x+c)^(3/2)*
(a+a*cos(d*x+c))^2*sin(d*x+c)/d+2/63*(9*A+13*B)*cos(d*x+c)^(3/2)*(a^3+a^3*c
os(d*x+c))*sin(d*x+c)/d+4/21*a^3*(13*A+11*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.138.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.62 (sec) , antiderivative size = 944, normalized size of antiderivative = 4.63

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3 \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(-\frac{(21A + 17B) \cot(c)}{30d} \right.$$

$$+ \frac{(107A + 97B) \cos(dx) \sin(c)}{336d} + \frac{(54A + 73B) \cos(2dx) \sin(2c)}{720d}$$

$$+ \frac{(A + 3B) \cos(3dx) \sin(3c)}{112d} + \frac{B \cos(4dx) \sin(4c)}{288d} + \frac{(107A + 97B) \cos(c) \sin(dx)}{720d}$$

$$+ \frac{(54A + 73B) \cos(2c) \sin(2dx)}{720d} + \frac{(A + 3B) \cos(3c) \sin(3dx)}{112d} + \left. \frac{B \cos(4c) \sin(4dx)}{288d} \right)$$

$$- \frac{13A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{4d}$$

$$- \frac{11B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{4d}$$

$$- \frac{7A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d}$$

$$- \frac{17B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{60d}$$

```
input Integrate[Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x
]
```


output `Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^3*Sec[c/2 + (d*x)/2]^6*(-1/30*((21*A + 17*B)*Cot[c])/d + ((107*A + 97*B)*Cos[d*x]*Sin[c])/(336*d) + ((54*A + 73*B)*Cos[2*d*x]*Sin[2*c])/(720*d) + ((A + 3*B)*Cos[3*d*x]*Sin[3*c])/(112*d) + (B*Cos[4*d*x]*Sin[4*c])/(288*d) + ((107*A + 97*B)*Cos[c]*Sin[d*x])/(336*d) + ((54*A + 73*B)*Cos[2*c]*Sin[2*d*x])/(720*d) + ((A + 3*B)*Cos[3*c]*Sin[3*d*x])/(112*d) + (B*Cos[4*c]*Sin[4*d*x])/(288*d)) - (13*A*(a + a*cos[c + d*x])^3*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]])]/(42*d*Sqrt[1 + Cot[c]^2]) - (11*B*(a + a*cos[c + d*x])^3*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]])]/(42*d*Sqrt[1 + Cot[c]^2]) - (7*A*(a + a*cos[c + d*x])^3*Csc[c]*Sec[c/2 + (d*x)/2]^6*(HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2])...`

3.138.3 Rubi [A] (verified)

Time = 1.31 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.03, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3455, 27, 3042, 3455, 27, 3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3455}$$

3.138. $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx$

$$\frac{2}{9} \int \frac{1}{2} \sqrt{\cos(c+dx)} (\cos(c+dx)a+a)^2 (3a(3A+B) + a(9A+13B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx) + a)^2}{9d}$$

↓ 27

$$\frac{1}{9} \int \sqrt{\cos(c+dx)} (\cos(c+dx)a+a)^2 (3a(3A+B) + a(9A+13B) \cos(c+dx)) dx + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx) + a)^2}{9d}$$

↓ 3042

$$\frac{1}{9} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2 \left(3a(3A+B) + a(9A+13B) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx) + a)^2}{9d}$$

↓ 3455

$$\frac{1}{9} \left(\frac{2}{7} \int 3\sqrt{\cos(c+dx)} (\cos(c+dx)a+a) (5(3A+2B)a^2 + (24A+23B) \cos(c+dx)a^2) dx + \frac{2(9A+13B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx) + a)^2}{9d} \right)$$

↓ 27

$$\frac{1}{9} \left(\frac{6}{7} \int \sqrt{\cos(c+dx)} (\cos(c+dx)a+a) (5(3A+2B)a^2 + (24A+23B) \cos(c+dx)a^2) dx + \frac{2(9A+13B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx) + a)^2}{9d} \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right) \left(5(3A+2B)a^2 + (24A+23B) \sin\left(c+dx+\frac{\pi}{2}\right)a^2\right) dx + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx) + a)^2}{9d} \right)$$

↓ 3447

3.138. $\int \sqrt{\cos(c+dx)} (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) dx$

$$\frac{1}{9} \left(\frac{6}{7} \int \sqrt{\cos(c+dx)} ((24A+23B) \cos^2(c+dx)a^3 + 5(3A+2B)a^3 + (5(3A+2B)a^3 + (24A+23B)a^3) \cos(c+dx) + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{9d} \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left((24A+23B) \sin\left(c+dx+\frac{\pi}{2}\right)^2 a^3 + 5(3A+2B)a^3 + (5(3A+2B)a^3 + (24A+23B)a^3) \cos(c+dx) + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{9d} \right) \right)$$

↓ 3502

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{2}{5} \int \frac{1}{2} \sqrt{\cos(c+dx)} (7(21A+17B)a^3 + 15(13A+11B) \cos(c+dx)a^3) dx + \frac{2a^3(24A+23B) \sin(c+dx) \cos(c+dx)}{5d} + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{9d} \right) \right)$$

↓ 27

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \int \sqrt{\cos(c+dx)} (7(21A+17B)a^3 + 15(13A+11B) \cos(c+dx)a^3) dx + \frac{2a^3(24A+23B) \sin(c+dx) \cos(c+dx)}{5d} + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{9d} \right) \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(7(21A+17B)a^3 + 15(13A+11B) \sin\left(c+dx+\frac{\pi}{2}\right) a^3 \right) dx + \frac{2a^3(24A+23B) \sin(c+dx) \cos(c+dx)}{5d} + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{9d} \right) \right)$$

↓ 3227

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(13A+11B) \int \cos^{\frac{3}{2}}(c+dx) dx + 7a^3(21A+17B) \int \sqrt{\cos(c+dx)} dx \right) + \frac{2a^3(24A+23B) \sin(c+dx) \cos(c+dx)}{5d} + \frac{2aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{9d} \right) \right)$$

↓ 3042

3.138. $\int \sqrt{\cos(c+dx)} (a + a \cos(c+dx))^3 (A + B \cos(c+dx)) dx$

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(7a^3(21A + 17B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 15a^3(13A + 11B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx \right) + \frac{2a^3(24A + 23B) \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + \frac{2aB \sin(c + dx) \cos^{3/2}(c + dx)(a \cos(c + dx) + a)^2}{9d} \right)$$

↓ 3115

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(7a^3(21A + 17B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 15a^3(13A + 11B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d} \right) \right) + \frac{2a^3(24A + 23B) \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + \frac{2aB \sin(c + dx) \cos^{3/2}(c + dx)(a \cos(c + dx) + a)^2}{9d} \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(7a^3(21A + 17B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 15a^3(13A + 11B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx)}{3d} \right) \right) + \frac{2a^3(24A + 23B) \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + \frac{2aB \sin(c + dx) \cos^{3/2}(c + dx)(a \cos(c + dx) + a)^2}{9d} \right)$$

↓ 3119

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(13A + 11B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) + \frac{14a^3(21A + 17B) \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + \frac{2aB \sin(c + dx) \cos^{3/2}(c + dx)(a \cos(c + dx) + a)^2}{9d} \right)$$

↓ 3120

$$\frac{1}{9} \left(\frac{2(9A + 13B) \sin(c + dx) \cos^{3/2}(c + dx) (a^3 \cos(c + dx) + a^3)}{7d} + \frac{6 \left(\frac{2a^3(24A + 23B) \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + \frac{2aB \sin(c + dx) \cos^{3/2}(c + dx)(a \cos(c + dx) + a)^2}{9d} \right)$$

input `Int[Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^3*(A + B*cos[c + d*x]),x]`

```
output (2*a*B*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^2*Sin[c + d*x]/(9*d) + ((2
*(9*A + 13*B)*Cos[c + d*x]^(3/2)*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(7
*d) + (6*((2*a^3*(24*A + 23*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d) + ((
14*a^3*(21*A + 17*B)*EllipticE[(c + d*x)/2, 2])/d + 15*a^3*(13*A + 11*B)*(
(2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/
(3*d))))/5)/7)/9
```

3.138.3.1 Defintions of rubi rules used

```
rule 277 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3115 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*Sin
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)^2], x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.138.4 Maple [A] (verified)

Time = 13.12 (sec) , antiderivative size = 413, normalized size of antiderivative = 2.02

method	result
default	$-\frac{4\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^3} \left(-560B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (360A + 2200B)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right)$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

output `-4/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(-560*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10+(360*A+2200*B)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1296*A-3412*B)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(1806*A+2702*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-624*A-738*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+195*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-441*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+165*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-357*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.138.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.09

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^3(A+B\cos(c+dx))dx =$$

$$\frac{2\left(15i\sqrt{2}(13A+11B)a^3\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))-15i\sqrt{2}(13A+11B)a^3\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))-21i\sqrt{2}(21A+17B)a^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c)))+21i\sqrt{2}(21A+17B)a^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c)))-(35B*a^3*\cos(dx+c)^3+45*(A+3B)*a^3*\cos(dx+c)^2+7*(27A+34B)*a^3*\cos(dx+c)+30*(13A+11B)*a^3)*\sqrt{\cos(dx+c)}*\sin(dx+c)}{d}\right)}{d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `-2/315*(15*I*sqrt(2)*(13*A + 11*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 15*I*sqrt(2)*(13*A + 11*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*I*sqrt(2)*(21*A + 17*B)*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*(21*A + 17*B)*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (35*B*a^3*cos(d*x + c)^3 + 45*(A + 3*B)*a^3*cos(d*x + c)^2 + 7*(27*A + 34*B)*a^3*cos(d*x + c) + 30*(13*A + 11*B)*a^3)*sqrt(cos(d*x + c))*sin(d*x + c)/d`

3.138.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*cos(d*x+c)**(1/2),x)`

output `Timed out`

3.138.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))3*(A+B*cos(d*x+c))*cos(d*x+c)(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)3*sqrt(cos(d*x + c)), x)`

3.138.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))3*(A+B*cos(d*x+c))*cos(d*x+c)(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)3*sqrt(cos(d*x + c)), x)`

3.138.9 Mupad [B] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.58

$$\begin{aligned}
& \int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^3(A+B\cos(c+dx))dx \\
&= \frac{2\left(Aa^3E\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)+Aa^3F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)+Aa^3\sqrt{\cos(c+dx)}\sin(c+dx)\right)}{Ba^3\left(\frac{2\sqrt{\cos(c+dx)}\sin(c+dx)}{3}+\frac{2F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)}{3}\right)} \\
&+ \frac{d}{6Aa^3\cos(c+dx)^{7/2}\sin(c+dx)} {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right) \\
&- \frac{7d\sqrt{\sin(c+dx)^2}}{2Aa^3\cos(c+dx)^{9/2}\sin(c+dx)} {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right) \\
&- \frac{9d\sqrt{\sin(c+dx)^2}}{6Ba^3\cos(c+dx)^{7/2}\sin(c+dx)} {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right) \\
&- \frac{7d\sqrt{\sin(c+dx)^2}}{2Ba^3\cos(c+dx)^{9/2}\sin(c+dx)} {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right) \\
&- \frac{3d\sqrt{\sin(c+dx)^2}}{2Ba^3\cos(c+dx)^{11/2}\sin(c+dx)} {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right) \\
&- \frac{11d\sqrt{\sin(c+dx)^2}}{11d\sqrt{\sin(c+dx)^2}}
\end{aligned}$$

```
input int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3,x)
```

```
output (2*(A*a^3*ellipticE(c/2 + (d*x)/2, 2) + A*a^3*ellipticF(c/2 + (d*x)/2, 2)
+ A*a^3*cos(c + d*x)^(1/2)*sin(c + d*x))/d + (B*a^3*((2*cos(c + d*x)^(1/2)
)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d - (6*A*a^3*cos(c
+ d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7
*d*(sin(c + d*x)^2)^(1/2)) - (2*A*a^3*cos(c + d*x)^(9/2)*sin(c + d*x)*hype
rgeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) - (6
*B*a^3*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c +
d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a^3*cos(c + d*x)^(9/2)*sin(c
+ d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(3*d*(sin(c + d*x)^2)
^(1/2)) - (2*B*a^3*cos(c + d*x)^(11/2)*sin(c + d*x)*hypergeom([1/2, 11/4],
15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2))
```

3.139 $\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

3.139.1 Optimal result 1363
 3.139.2 Mathematica [C] (warning: unable to verify) 1364
 3.139.3 Rubi [A] (verified) 1365
 3.139.4 Maple [A] (verified) 1369
 3.139.5 Fricas [C] (verification not implemented) 1370
 3.139.6 Sympy [F(-1)] 1371
 3.139.7 Maxima [F] 1371
 3.139.8 Giac [F] 1371
 3.139.9 Mupad [B] (verification not implemented) 1372

3.139.1 Optimal result

Integrand size = 33, antiderivative size = 171

$$\int \frac{(a + a \cos(c + dx))^3(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{4a^3(9A + 7B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{4a^3(21A + 13B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d}$$

$$+ \frac{4a^3(42A + 41B)\sqrt{\cos(c + dx)} \sin(c + dx)}{105d}$$

$$+ \frac{2aB\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2 \sin(c + dx)}{7d}$$

$$+ \frac{2(7A + 11B)\sqrt{\cos(c + dx)}(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{35d}$$

```
output 4/5*a^3*(9*A+7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/21*a^3*(21*A+13*B)*(cos(1/2*d*x+1/2*c)^2
)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/105*a
^3*(42*A+41*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d+2/7*a*B*(a+a*cos(d*x+c))^2*si
n(d*x+c)*cos(d*x+c)^(1/2)/d+2/35*(7*A+11*B)*(a^3+a^3*cos(d*x+c))*sin(d*x+c
)*cos(d*x+c)^(1/2)/d
```

3.139.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.63 (sec) , antiderivative size = 898, normalized size of antiderivative = 5.25

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^3 \sec^6 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{(9A + 7B) \cot(c)}{10d} \right. \\ \left. + \frac{(84A + 107B) \cos(dx) \sin(c)}{336d} + \frac{(A + 3B) \cos(2dx) \sin(2c)}{40d} + \frac{B \cos(3dx) \sin(3c)}{112d} \right. \\ \left. + \frac{(84A + 107B) \cos(c) \sin(dx)}{336d} + \frac{(A + 3B) \cos(2c) \sin(2dx)}{40d} + \frac{B \cos(3c) \sin(3dx)}{112d} \right) \\ - \frac{A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{20d} \\ - \frac{13B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{40d} \\ - \frac{9A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d} \\ - \frac{7B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d}$$

input `Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output $\text{Sqrt}[\text{Cos}[c + d*x]]*(a + a*\text{Cos}[c + d*x])^3*\text{Sec}[c/2 + (d*x)/2]^6*(-1/10*((9*A + 7*B)*\text{Cot}[c])/d + ((84*A + 107*B)*\text{Cos}[d*x]*\text{Sin}[c])/(336*d) + ((A + 3*B)*\text{Cos}[2*d*x]*\text{Sin}[2*c])/(40*d) + (B*\text{Cos}[3*d*x]*\text{Sin}[3*c])/(112*d) + ((84*A + 107*B)*\text{Cos}[c]*\text{Sin}[d*x])/(336*d) + ((A + 3*B)*\text{Cos}[2*c]*\text{Sin}[2*d*x])/(40*d) + (B*\text{Cos}[3*c]*\text{Sin}[3*d*x])/(112*d)) - (A*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^6*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(2*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (13*B*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^6*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])]*\text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(42*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (9*A*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{Sec}[c/2 + (d*x)/2]^6*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]]*\text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\text{Cos}[c]^2*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])/(\text{Cos}[c]^2 + \text{Sin}[c]^2))/ \text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]])/(20*d) - (7*B*(a...$

3.139.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3455, 27, 3042, 3455, 3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3455

3.139. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\frac{2}{7} \int \frac{(\cos(c+dx)a+a)^2(a(7A+B)+a(7A+11B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx + \frac{2aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}{7d}$$

↓ 27

$$\frac{1}{7} \int \frac{(\cos(c+dx)a+a)^2(a(7A+B)+a(7A+11B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx + \frac{2aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}{7d}$$

↓ 3042

$$\frac{1}{7} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2(a(7A+B)+a(7A+11B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}{7d}$$

↓ 3455

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{(\cos(c+dx)a+a)((21A+8B)a^2+(42A+41B)\cos(c+dx)a^2)}{\sqrt{\cos(c+dx)}} dx + \frac{2(7A+11B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d} \right) + \frac{2aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)((21A+8B)a^2+(42A+41B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(7A+11B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d} \right) + \frac{2aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}{7d}$$

↓ 3447

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{(42A+41B)\cos^2(c+dx)a^3+(21A+8B)a^3+((21A+8B)a^3+(42A+41B)a^3)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{2(7A+11B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d} \right) + \frac{2aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{(42A + 41B) \sin(c + dx + \frac{\pi}{2})^2 a^3 + (21A + 8B)a^3 + ((21A + 8B)a^3 + (42A + 41B)a^3) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} \right. \\ \left. \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) \\ \downarrow \text{3502}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{2}{3} \int \frac{5(21A + 13B)a^3 + 21(9A + 7B) \cos(c + dx)a^3}{2\sqrt{\cos(c + dx)}} dx + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\ \downarrow \text{27}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \frac{5(21A + 13B)a^3 + 21(9A + 7B) \cos(c + dx)a^3}{\sqrt{\cos(c + dx)}} dx + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \frac{5(21A + 13B)a^3 + 21(9A + 7B) \sin(c + dx + \frac{\pi}{2}) a^3}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right. \\ \left. \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) \\ \downarrow \text{3227}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(21A + 13B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 21a^3(9A + 7B) \int \sqrt{\cos(c + dx)} dx \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(21A + 13B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(9A + 7B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) \right) \right. \\ \left. \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) + \frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\ \downarrow \text{3119}$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(21A + 13B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{42a^3(9A + 7B)E(\frac{1}{2}(c + dx)|2)}{d} \right) \right) + \frac{2a^3(42A + 41B) \sin(c + dx)}{3d} \right) + \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d}$$

↓ 3120

$$\frac{1}{7} \left(\frac{2(7A + 11B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2}{5} \left(\frac{2a^3(42A + 41B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2}{7d} \right) \right)$$

input `Int[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(2*a*B*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(7*d) + ((2*(7*A + 11*B)*Sqrt[Cos[c + d*x]]*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(5*d) + (2*(((42*a^3*(9*A + 7*B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(21*A + 13*B)*EllipticF[(c + d*x)/2, 2])/d)/3 + (2*a^3*(42*A + 41*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)))/5)/7`

3.139.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3455 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.139.4 Maple [A] (verified)

Time = 11.55 (sec) , antiderivative size = 385, normalized size of antiderivative = 2.25

method	result
default	$-\frac{4\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^3} \left(120B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-84A - 432B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots\right)$
parts	Expression too large to display

3.139.
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

input `int((a+cos(d*x+c))*a^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)`

output `-4/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(120*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8+(-84*A-432*B)*sin(1/2*d*x+1/2*c)^6
*cos(1/2*d*x+1/2*c)+(294*A+602*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+
(-126*A-208*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+105*A*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2
*c),2^(1/2))-189*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+65*B*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2
)-147*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elli
pticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/
2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.139.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.19

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx =$$

$$\frac{2 \left(5i \sqrt{2} (21A + 13B) a^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (21A + 13B) \right)}{-}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorith
m="fricas")`

output `-2/105*(5*I*sqrt(2)*(21*A + 13*B))*a^3*weierstrassPInverse(-4, 0, cos(d*x +
c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(21*A + 13*B))*a^3*weierstrassPInverse(
-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*I*sqrt(2)*(9*A + 7*B))*a^3*weier
strassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)
) + 21*I*sqrt(2)*(9*A + 7*B))*a^3*weierstrassZeta(-4, 0, weierstrassPInver
se(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (15*B*a^3*cos(d*x + c)^2 + 21*
(A + 3*B))*a^3*cos(d*x + c) + 5*(21*A + 26*B))*a^3)*sqrt(cos(d*x + c))*sin(d
*x + c))/d`

3.139. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

3.139.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

```
input integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)
```

```
output Timed out
```

3.139.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\sqrt{\cos(dx + c)}} dx \end{aligned}$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm
m="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/sqrt(cos(d*x + c)),
x)
```

3.139.8 Giac [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\sqrt{\cos(dx + c)}} dx \end{aligned}$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm
m="giac")
```

```
output integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/sqrt(cos(d*x + c)),
x)
```

3.139.9 Mupad [B] (verification not implemented)

Time = 1.08 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.49

$$\begin{aligned}
& \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\
&= \frac{2 \left(B a^3 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) + B a^3 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) + B a^3 \sqrt{\cos(c + dx)} \sin(c + dx) \right)}{d} \\
&+ \frac{6 A a^3 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 A a^3 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{2 A a^3 \sqrt{\cos(c + dx)} \sin(c + dx)}{d} \\
&- \frac{2 A a^3 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}} \\
&- \frac{6 B a^3 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}} \\
&- \frac{2 B a^3 \cos(c + dx)^{9/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c + dx)^2\right)}{9 d \sqrt{\sin(c + dx)^2}}
\end{aligned}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^(1/2),x)
```

```
output (2*(B*a^3*ellipticE(c/2 + (d*x)/2, 2) + B*a^3*ellipticF(c/2 + (d*x)/2, 2)
+ B*a^3*cos(c + d*x)^(1/2)*sin(c + d*x))/d + (6*A*a^3*ellipticE(c/2 + (d*
x)/2, 2))/d + (4*A*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a^3*cos(c + d
*x)^(1/2)*sin(c + d*x))/d - (2*A*a^3*cos(c + d*x)^(7/2)*sin(c + d*x)*hyper
geom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (6*
B*a^3*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c +
d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a^3*cos(c + d*x)^(9/2)*sin(c
+ d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(
1/2))
```

3.140
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.140.1 Optimal result 1373
 3.140.2 Mathematica [C] (warning: unable to verify) 1374
 3.140.3 Rubi [A] (verified) 1375
 3.140.4 Maple [A] (verified) 1380
 3.140.5 Fracas [C] (verification not implemented) 1381
 3.140.6 Sympy [F(-1)] 1381
 3.140.7 Maxima [F] 1382
 3.140.8 Giac [F] 1382
 3.140.9 Mupad [B] (verification not implemented) 1383

3.140.1 Optimal result

Integrand size = 33, antiderivative size = 169

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \frac{4a^3(5A + 9B)E(\frac{1}{2}(c + dx) | 2)}{5d} + \frac{4a^3(5A + 3B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} \\ & - \frac{4a^3(5A - 6B)\sqrt{\cos(c + dx)} \sin(c + dx)}{15d} + \frac{2aA(a + a \cos(c + dx))^2 \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\ & - \frac{2(5A - B)\sqrt{\cos(c + dx)}(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{5d} \end{aligned}$$

```
output 4/5*a^3*(5*A+9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/3*a^3*(5*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*a*A*(a+
*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+c)^(1/2)-4/15*a^3*(5*A-6*B)*sin(d*x+c)
*cos(d*x+c)^(1/2)/d-2/5*(5*A-B)*(a^3+a^3*cos(d*x+c))*sin(d*x+c)*cos(d*x+c)
^(1/2)/d
```

3.140.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.74 (sec) , antiderivative size = 888, normalized size of antiderivative = 5.25

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^3 \sec^6 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{(5A + 18B + 15A \cos(2c) + 18B \cos(2c)) \csc(c) \sec(c)}{40d} + \frac{(A + 3B) \cos(dx) \sin(c)}{12d} + \frac{B \cos(2dx) \sin(2c)}{40d} + \frac{(A + 3B) \cos(c) \sin(dx)}{12d} + \frac{A \sec(c) \sec(c + dx) \sin(dx)}{4d} + \frac{B \cos(2c) \sin(2dx)}{40d} \right) - \frac{5A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{6d} - \frac{B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{20d} - \frac{A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{4d} - \frac{9B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d}$$

input `Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

```

output Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^3*Sec[c/2 + (d*x)/2]^6*(-1/40*((5*
A + 18*B + 15*A*cos[2*c] + 18*B*cos[2*c])*Csc[c]*Sec[c])/d + ((A + 3*B)*Co
s[d*x]*Sin[c])/(12*d) + (B*cos[2*d*x]*Sin[2*c])/(40*d) + ((A + 3*B)*Cos[c]
*sin[d*x])/(12*d) + (A*Sec[c]*Sec[c + d*x]*Sin[d*x])/(4*d) + (B*cos[2*c]*S
in[2*d*x])/(40*d) - (5*A*(a + a*cos[c + d*x])^3*Csc[c]*HypergeometricPFQ[
{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^6*Sec[d
*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + C
ot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[
c]]]])/(6*d*Sqrt[1 + Cot[c]^2]) - (B*(a + a*cos[c + d*x])^3*Csc[c]*Hyperge
ometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)
/2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-
(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x -
ArcTan[Cot[c]]]])/(2*d*Sqrt[1 + Cot[c]^2]) - (A*(a + a*cos[c + d*x])^3*Csc
[c]*Sec[c/2 + (d*x)/2]^6*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x
+ ArcTan[Tan[c]]]^2]*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x +
ArcTan[Tan[c]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]*Sqrt[Cos[c]*Cos[d*x
+ ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + Ar
cTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*cos[c]^2*cos[d*x + ArcTan[Ta
n[c]]]*Sqrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + Ar
cTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]))/(4*d) - (9*B*(a + a*cos[c + d*x])^3...
    
```

3.140.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3454, 27, 3042, 3455, 3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}}} dx$$

↓ 3454

3.140. $\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
& 2 \int \frac{(\cos(c+dx)a+a)^2(a(5A+B)-a(5A-B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx + \\
& \quad \frac{2aA \sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{27} \\
& \int \frac{(\cos(c+dx)a+a)^2(a(5A+B)-a(5A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx + \\
& \quad \frac{2aA \sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2(a(5A+B)-a(5A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \\
& \quad \frac{2aA \sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3455} \\
& \frac{2}{5} \int \frac{(\cos(c+dx)a+a)(a^2(10A+3B)-a^2(5A-6B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx - \\
& \frac{2(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}(a^3\cos(c+dx)+a^3)}{5d} + \frac{2aA \sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{2}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a^2(10A+3B)-a^2(5A-6B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \\
& \frac{2(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}(a^3\cos(c+dx)+a^3)}{5d} + \frac{2aA \sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3447} \\
& \frac{2}{5} \int \frac{-((5A-6B)\cos^2(c+dx)a^3)+(10A+3B)a^3+(a^3(10A+3B)-a^3(5A-6B))\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx - \\
& \frac{2(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}(a^3\cos(c+dx)+a^3)}{5d} + \frac{2aA \sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.140. $\int \frac{(a+a\cos(c+dx))^3(A+B\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\frac{2}{5} \int \frac{-\left((5A - 6B) \sin(c + dx + \frac{\pi}{2})^2 a^3\right) + (10A + 3B)a^3 + (a^3(10A + 3B) - a^3(5A - 6B)) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx -$$

$$\frac{2(5A - B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^2}{d \sqrt{\cos(c + dx)}}$$

↓ 3502

$$\frac{2}{5} \left(\frac{2}{3} \int \frac{5(5A + 3B)a^3 + 3(5A + 9B) \cos(c + dx)a^3}{2\sqrt{\cos(c + dx)}} dx - \frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) -$$

$$\frac{2(5A - B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^2}{d \sqrt{\cos(c + dx)}}$$

↓ 27

$$\frac{2}{5} \left(\frac{1}{3} \int \frac{5(5A + 3B)a^3 + 3(5A + 9B) \cos(c + dx)a^3}{\sqrt{\cos(c + dx)}} dx - \frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) -$$

$$\frac{2(5A - B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^2}{d \sqrt{\cos(c + dx)}}$$

↓ 3042

$$\frac{2}{5} \left(\frac{1}{3} \int \frac{5(5A + 3B)a^3 + 3(5A + 9B) \sin(c + dx + \frac{\pi}{2}) a^3}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) -$$

$$\frac{2(5A - B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^2}{d \sqrt{\cos(c + dx)}}$$

↓ 3227

$$\frac{2}{5} \left(\frac{1}{3} \left(5a^3(5A + 3B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3a^3(5A + 9B) \int \sqrt{\cos(c + dx)} dx \right) - \frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) -$$

$$\frac{2(5A - B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^2}{d \sqrt{\cos(c + dx)}}$$

↓ 3042

$$\frac{2}{5} \left(\frac{1}{3} \left(5a^3(5A + 3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a^3(5A + 9B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) - \frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) -$$

$$\frac{2(5A - B) \sin(c + dx) \sqrt{\cos(c + dx)} (a^3 \cos(c + dx) + a^3)}{5d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^2}{d \sqrt{\cos(c + dx)}}$$

3.140. $\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$

↓ 3119

$$\frac{2}{5} \left(\frac{1}{3} \left(5a^3(5A+3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^3(5A+9B)E(\frac{1}{2}(c+dx)|2)}{d} \right) - \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}(a^3\cos(c+dx)+a^3)}{5d} + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}}$$

↓ 3120

$$- \frac{2(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}(a^3\cos(c+dx)+a^3)}{5d} + \frac{2}{5} \left(\frac{1}{3} \left(\frac{10a^3(5A+3B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d} + \frac{6a^3(5A+9B)E(\frac{1}{2}(c+dx)|2)}{d} \right) - \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{d\sqrt{\cos(c+dx)}}$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])/Cos[c + d*x]^(3/2),x]`

output `(2*a*A*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - (2*(5*A - B)*Sqrt[Cos[c + d*x]]*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(5*d) + (2*(((6*a^3*(5*A + 9*B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(5*A + 3*B)*EllipticF[(c + d*x)/2, 2])/d)/3 - (2*a^3*(5*A - 6*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

3.140.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.140.4 Maple [A] (verified)

Time = 9.08 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.99

method	result
default	$\frac{4a^3 \left(-12B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 10A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 42B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 20A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{\dots}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/15*a^3*(-12*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+10*A*cos(1/2*d*x+
1/2*c)*sin(1/2*d*x+1/2*c)^4+42*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4-2
0*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2+25*A*(sin(1/2*d*x+1/2*c)^2)^(1
/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))
-15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellipt
icE(cos(1/2*d*x+1/2*c),2^(1/2))-18*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)
^2+15*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elli
pticF(cos(1/2*d*x+1/2*c),2^(1/2))-27*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin
(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*
d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

$$3.140. \int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.140.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.36

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{2 \left(5i \sqrt{2} (5A + 3B) a^3 \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (5A + 3B) a^3 \sin(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - 3i \sqrt{2} (5A + 9B) a^3 \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 3i \sqrt{2} (5A + 9B) a^3 \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - (3B a^3 \cos(dx + c)^2 + 5(A + 3B) a^3 \cos(dx + c) + 15 A a^3) \sqrt{\cos(dx + c)} \sin(dx + c) / (d \cos(dx + c)) \right)}{d}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="fricas")`

output `-2/15*(5*I*sqrt(2)*(5*A + 3*B)*a^3*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(5*A + 3*B)*a^3*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(5*A + 9*B)*a^3*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(5*A + 9*B)*a^3*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (3*B*a^3*cos(d*x + c)^2 + 5*(A + 3*B)*a^3*cos(d*x + c) + 15*A*a^3)*sqrt(cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c))`

3.140.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Timed out`

3.140.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(3/2), x)`

3.140.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(3/2), x)`

3.140.9 Mupad [B] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.36

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{A a^3 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} | 2\right)}{3} \right)}{d} + \frac{6 A a^3 E\left(\frac{c}{2} + \frac{dx}{2} | 2\right)}{d}$$

$$+ \frac{6 A a^3 F\left(\frac{c}{2} + \frac{dx}{2} | 2\right)}{d} + \frac{6 B a^3 E\left(\frac{c}{2} + \frac{dx}{2} | 2\right)}{d} + \frac{4 B a^3 F\left(\frac{c}{2} + \frac{dx}{2} | 2\right)}{d}$$

$$+ \frac{2 B a^3 \sqrt{\cos(c + dx)} \sin(c + dx)}{d} + \frac{2 A a^3 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 B a^3 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^(3/2),x)
```

```
output (A*a^3*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (6*A*a^3*ellipticE(c/2 + (d*x)/2, 2))/d + (6*A*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (6*B*a^3*ellipticE(c/2 + (d*x)/2, 2))/d + (4*B*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*a^3*cos(c + d*x)^(1/2)*sin(c + d*x))/d + (2*A*a^3*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) - (2*B*a^3*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))
```

3.141
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.141.1 Optimal result 1384
 3.141.2 Mathematica [C] (warning: unable to verify) 1385
 3.141.3 Rubi [A] (verified) 1386
 3.141.4 Maple [B] (verified) 1391
 3.141.5 Fricas [C] (verification not implemented) 1392
 3.141.6 Sympy [F(-1)] 1392
 3.141.7 Maxima [F] 1393
 3.141.8 Giac [F] 1393
 3.141.9 Mupad [B] (verification not implemented) 1394

3.141.1 Optimal result

Integrand size = 33, antiderivative size = 161

$$\begin{aligned} & \int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx \\ &= -\frac{4a^3(A-B)E(\frac{1}{2}(c+dx)|2)}{d} + \frac{20a^3(A+B) \operatorname{EllipticF}(\frac{1}{2}(c+dx),2)}{3d} \\ & \quad - \frac{4a^3(4A+B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3d} + \frac{2aA(a+a \cos(c+dx))^2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \\ & \quad + \frac{2(7A+3B)(a^3+a^3 \cos(c+dx)) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}} \end{aligned}$$

output

```
-4*a^3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+20/3*a^3*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*A*(a+a*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/3*(7*A+3*B)*(a^3+a^3*cos(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(1/2)-4/3*a^3*(4*A+B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.141.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.95 (sec) , antiderivative size = 879, normalized size of antiderivative = 5.46

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^3 \sec^6 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{(-5A + B + A \cos(2c) + 3B \cos(2c)) \csc(c) \sec(c)}{8d} + \frac{B \cos(dx) \sin(c)}{12d} + \frac{B \cos(c) \sin(dx)}{12d} + \frac{A \sec(c) \sec^2(c + dx) \sin(dx)}{12d} + \frac{\sec(c) \sec(c + dx) (A \sin(c) + 9A \sin(dx) + 3B \sin(dx))}{12d} \right) - \frac{5A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{6d} - \frac{5B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{6d} + \frac{A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{4d} + \frac{B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{4d}$$

input `Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2), x]`

output $\text{Sqrt}[\text{Cos}[c + d*x]]*(a + a*\text{Cos}[c + d*x])^3*\text{Sec}[c/2 + (d*x)/2]^6*(-1/8*((-5*A + B + A*\text{Cos}[2*c] + 3*B*\text{Cos}[2*c])* \text{Csc}[c]*\text{Sec}[c])/d + (B*\text{Cos}[d*x]*\text{Sin}[c])/ (12*d) + (B*\text{Cos}[c]*\text{Sin}[d*x])/ (12*d) + (A*\text{Sec}[c]*\text{Sec}[c + d*x]^2*\text{Sin}[d*x])/ (12*d) + (\text{Sec}[c]*\text{Sec}[c + d*x]*(A*\text{Sin}[c] + 9*A*\text{Sin}[d*x] + 3*B*\text{Sin}[d*x]))/ (12*d)) - (5*A*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^6*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])] * \text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/ (6*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) - (5*B*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\text{Sec}[c/2 + (d*x)/2]^6*\text{Sec}[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]])] * \text{Sqrt}[1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]])/ (6*d*\text{Sqrt}[1 + \text{Cot}[c]^2]) + (A*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{Sec}[c/2 + (d*x)/2]^6*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]] * \text{Sqrt}[1 + \text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]] * \text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]] * \text{Sqrt}[1 + \text{Tan}[c]^2]) * \text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\text{Sin}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\text{Cos}[c]^2*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])/(\text{Cos}[c]^2 + \text{Sin}[c]^2))/ \text{Sqrt}[\text{Cos}[c]*\text{Cos}[d*x + \text{ArcTan}[\text{Tan}[c]]]] * \text{Sqrt}[1 + \text{Tan}[c]^2]))/ (4*d) - (B*(a + a*\text{Cos}[c + d*x])^3*\text{Csc}[c]*\text{Sec}[c/...$

3.141.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.06, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3454

3.141. $\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{2}{3} \int \frac{(\cos(c+dx)a+a)^2(a(7A+3B)-3a(A-B)\cos(c+dx))}{\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{3d\cos^{\frac{3}{2}}(c+dx)}} dx + \\
& \quad \downarrow 27 \\
& \frac{1}{3} \int \frac{(\cos(c+dx)a+a)^2(a(7A+3B)-3a(A-B)\cos(c+dx))}{\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{3d\cos^{\frac{3}{2}}(c+dx)}} dx + \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2(a(7A+3B)-3a(A-B)\sin(c+dx+\frac{\pi}{2}))}{\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{3d\cos^{\frac{3}{2}}(c+dx)}} dx + \\
& \quad \downarrow 3454 \\
& \frac{1}{3} \left(2 \int \frac{3(\cos(c+dx)a+a)(a^2(3A+2B)-a^2(4A+B)\cos(c+dx))}{\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{3d\cos^{\frac{3}{2}}(c+dx)}} dx + \frac{2(7A+3B)\sin(c+dx)(a^3\cos(c+dx))}{d\sqrt{\cos(c+dx)}} \right) \\
& \quad \downarrow 27 \\
& \frac{1}{3} \left(6 \int \frac{(\cos(c+dx)a+a)(a^2(3A+2B)-a^2(4A+B)\cos(c+dx))}{\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{3d\cos^{\frac{3}{2}}(c+dx)}} dx + \frac{2(7A+3B)\sin(c+dx)(a^3\cos(c+dx))}{d\sqrt{\cos(c+dx)}} \right) \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left(6 \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a^2(3A+2B)-a^2(4A+B)\sin(c+dx+\frac{\pi}{2}))}{\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{3d\cos^{\frac{3}{2}}(c+dx)}} dx + \frac{2(7A+3B)\sin(c+dx)(a^3\cos(c+dx))}{d\sqrt{\cos(c+dx)}} \right) \\
& \quad \downarrow 3447
\end{aligned}$$

3.141. $\int \frac{(a+a\cos(c+dx))^3(A+B\cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(6 \int \frac{-((4A+B)\cos^2(c+dx)a^3) + (3A+2B)a^3 + (a^3(3A+2B) - a^3(4A+B))\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{2(7A+3B)}{3d} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(6 \int \frac{-((4A+B)\sin(c+dx+\frac{\pi}{2})^2 a^3) + (3A+2B)a^3 + (a^3(3A+2B) - a^3(4A+B))\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(7A+3B)}{3d} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3502

$$\frac{1}{3} \left(6 \left(\frac{2}{3} \int \frac{5a^3(A+B) - 3a^3(A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx - \frac{2a^3(4A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2(7A+3B)}{3d} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{3} \left(6 \left(\frac{1}{3} \int \frac{5a^3(A+B) - 3a^3(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx - \frac{2a^3(4A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2(7A+3B)}{3d} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(6 \left(\frac{1}{3} \int \frac{5a^3(A+B) - 3a^3(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a^3(4A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2(7A+3B)}{3d} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3227

$$\frac{1}{3} \left(6 \left(\frac{1}{3} \left(5a^3(A+B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^3(A-B) \int \sqrt{\cos(c+dx)} dx \right) - \frac{2a^3(4A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)} \right)$$

3.141. $\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

↓ 3042

$$\frac{1}{3} \left(6 \left(\frac{1}{3} \left(5a^3(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^3(A-B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) - \frac{2a^3(4A+B)\sin(c+dx)}{3d} \right) \right) - \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3119

$$\frac{1}{3} \left(6 \left(\frac{1}{3} \left(5a^3(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6a^3(A-B)E(\frac{1}{2}(c+dx)|2)}{d} \right) - \frac{2a^3(4A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) \right) - \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3120

$$\frac{1}{3} \left(\frac{(2(7A+3B)\sin(c+dx)(a^3 \cos(c+dx) + a^3))}{d \sqrt{\cos(c+dx)}} + 6 \left(\frac{1}{3} \left(\frac{10a^3(A+B)\text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} - \frac{6a^3(A-B)E(\frac{1}{2}(c+dx)|2)}{d} \right) \right) - \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{3d \cos^{\frac{3}{2}}(c+dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*a*A*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((2*(7*A + 3*B)*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) + 6*(((-6*a^3*(A - B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(A + B)*EllipticF[(c + d*x)/2, 2])/d)/3 - (2*a^3*(4*A + B)*Sqrt[Cos[c + d*x])*Sin[c + d*x])/(3*d)))/3`

3.141.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.141. $\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.141.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 653 vs. $2(199) = 398$.

Time = 9.47 (sec) , antiderivative size = 654, normalized size of antiderivative = 4.06

method	result
default	$-\frac{4\left(-4B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)(9A+5B)\left(\sin^4\right)$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/3*(-4*B*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^
2)^(1/2)*sin(1/2*d*x+1/2*c)^6+2*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*(9*A+5*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(5*A+2*B)*sin(1/2*d*x+1/2*c)^2*c
os(1/2*d*x+1/2*c)-2*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*
(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(5*A*Elliptic
F(cos(1/2*d*x+1/2*c),2^(1/2))+3*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*
B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*B*EllipticE(cos(1/2*d*x+1/2*c),2
^(1/2)))*sin(1/2*d*x+1/2*c)^2+5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*
d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(-2*sin(1/2*d*
x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+3*A*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/
2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2
-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*B*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/
2))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)-3*B*(-2*sin(1/2*d
*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*si
n(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*a^3/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-
1)^(3/2)/sin(1/2*d*x+1/2*c)/d
```

$$3.141. \int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.141.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.39

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{2 \left(5i \sqrt{2} (A + B) a^3 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (A + B) a^3 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 3i \sqrt{2} (A - B) a^3 \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 3i \sqrt{2} (A - B) a^3 \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - (B a^3 \cos(dx + c)^2 + 3(3A + B) a^3 \cos(dx + c) + A a^3) \sqrt{\cos(dx + c)} \sin(dx + c) \right)}{(d \cos(dx + c))^2}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fricas")`

output `-2/3*(5*I*sqrt(2)*(A + B)*a^3*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(A + B)*a^3*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(A - B)*a^3*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(A - B)*a^3*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (B*a^3*cos(d*x + c)^2 + 3*(3*A + B)*a^3*cos(d*x + c) + A*a^3)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.141.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output `Timed out`

3.141.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(5/2), x)`

3.141.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(5/2), x)`

3.141.9 Mupad [B] (verification not implemented)

Time = 1.69 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.56

$$\begin{aligned}
& \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\
&= \frac{2(A a^3 E(\frac{c}{2} + \frac{dx}{2}|2) + 3 A a^3 F(\frac{c}{2} + \frac{dx}{2}|2))}{d} \\
&+ \frac{B a^3 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F(\frac{c}{2} + \frac{dx}{2}|2)}{3} \right)}{d} + \frac{6 B a^3 E(\frac{c}{2} + \frac{dx}{2}|2)}{d} \\
&+ \frac{6 B a^3 F(\frac{c}{2} + \frac{dx}{2}|2)}{d} + \frac{6 A a^3 \sin(c + dx) {}_2F_1(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{2 A a^3 \sin(c + dx) {}_2F_1(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{2 B a^3 \sin(c + dx) {}_2F_1(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}
\end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^(5/2),x)`output `(2*(A*a^3*ellipticE(c/2 + (d*x)/2, 2) + 3*A*a^3*ellipticF(c/2 + (d*x)/2, 2)))/d + (B*a^3*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (6*B*a^3*ellipticE(c/2 + (d*x)/2, 2))/d + (6*B*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (6*A*a^3*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) + (2*A*a^3*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2)) + (2*B*a^3*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.142
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.142.1 Optimal result 1395
 3.142.2 Mathematica [C] (warning: unable to verify) 1396
 3.142.3 Rubi [A] (verified) 1397
 3.142.4 Maple [B] (verified) 1401
 3.142.5 Fracas [C] (verification not implemented) 1402
 3.142.6 Sympy [F(-1)] 1403
 3.142.7 Maxima [F] 1403
 3.142.8 Giac [F] 1404
 3.142.9 Mupad [B] (verification not implemented) 1404

3.142.1 Optimal result

Integrand size = 33, antiderivative size = 171

$$\int \frac{(a + a \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= -\frac{4a^3(9A + 5B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{4a^3(3A + 5B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{4a^3(21A + 20B) \sin(c + dx)}{15d\sqrt{\cos(c + dx)}} + \frac{2aA(a + a \cos(c + dx))^2 \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{2(9A + 5B)(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{15d \cos^{\frac{3}{2}}(c + dx)}$$

```
output -4/5*a^3*(9*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellipti
cE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/3*a^3*(3*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*a*A*(
a+a*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+c)^(5/2)+2/15*(9*A+5*B)*(a^3+a^3*co
s(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(3/2)+4/15*a^3*(21*A+20*B)*sin(d*x+c)/d/
cos(d*x+c)^(1/2)
```

3.142.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.45 (sec) , antiderivative size = 890, normalized size of antiderivative = 5.20

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^3 \sec^6 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(-\frac{(-36A - 25B + 5B \cos(2c)) \csc(c) \sec(c)}{40d} + \frac{A \sec(c) \sec^3(c + dx) \sin(dx)}{20d} + \frac{\sec(c) \sec^2(c + dx) (3A \sin(c) + 15A \sin(dx) + 5B \sin(dx))}{60d} + \frac{\sec(c) \sec(c + dx) (15A \sin(c) + 5B \sin(c) + 54A \sin(dx) + 45B \sin(dx))}{60d} \right) - \frac{A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{20d} - \frac{5B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{60d} + \frac{9A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d} + \frac{B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}, \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{4d}$$

```
input Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]
```

output $\sqrt{\cos[c + dx]}(a + a\cos[c + dx])^3 \sec[c/2 + (dx)/2]^6 (-1/40 * ((-36A - 25B + 5B\cos[2c]) \csc[c] \sec[c])/d + (A \sec[c] \sec[c + dx]^3 \sin[dx])/(20d) + (\sec[c] \sec[c + dx]^2 (3A \sin[c] + 15A \sin[dx] + 5B \sin[dx]))/(60d) + (\sec[c] \sec[c + dx] (15A \sin[c] + 5B \sin[c] + 54A \sin[dx] + 45B \sin[dx]))/(60d) - (A(a + a\cos[c + dx])^3 \csc[c] \text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[dx - \text{ArcTan}[\text{Cot}[c]]]^2] \sec[c/2 + (dx)/2]^6 \sec[dx - \text{ArcTan}[\text{Cot}[c]]] \sqrt{1 - \sin[dx - \text{ArcTan}[\text{Cot}[c]]]} \sqrt{-(\sqrt{1 + \text{Cot}[c]^2} \sin[c] \sin[dx - \text{ArcTan}[\text{Cot}[c]])} \sqrt{1 + \sin[dx - \text{ArcTan}[\text{Cot}[c]]]})/(2d \sqrt{1 + \text{Cot}[c]^2}) - (5B(a + a\cos[c + dx])^3 \csc[c] \text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[dx - \text{ArcTan}[\text{Cot}[c]]]^2] \sec[c/2 + (dx)/2]^6 \sec[dx - \text{ArcTan}[\text{Cot}[c]]] \sqrt{1 - \sin[dx - \text{ArcTan}[\text{Cot}[c]]]} \sqrt{-(\sqrt{1 + \text{Cot}[c]^2} \sin[c] \sin[dx - \text{ArcTan}[\text{Cot}[c]])} \sqrt{1 + \sin[dx - \text{ArcTan}[\text{Cot}[c]]]})/(6d \sqrt{1 + \text{Cot}[c]^2}) + (9A(a + a\cos[c + dx])^3 \csc[c] \sec[c/2 + (dx)/2]^6 (\text{HypergeometricPFQ}[-1/2, -1/4], \{3/4\}, \cos[dx + \text{ArcTan}[\text{Tan}[c]]]^2] \sin[dx + \text{ArcTan}[\text{Tan}[c]]] \tan[c]) / (\sqrt{1 - \cos[dx + \text{ArcTan}[\text{Tan}[c]]]} \sqrt{1 + \cos[dx + \text{ArcTan}[\text{Tan}[c]]]} \sqrt{\cos[c] \cos[dx + \text{ArcTan}[\text{Tan}[c]]] \sqrt{1 + \tan[c]^2}} \sqrt{1 + \tan[c]^2}) - ((\sin[dx + \text{ArcTan}[\text{Tan}[c]]] \tan[c]) / \sqrt{1 + \tan[c]^2} + (2 \cos[c]^2 \cos[dx + \text{ArcTan}[\text{Tan}[c]]] \sqrt{1 + \tan[c]^2}) / (\cos[c]^2 + \sin[c]^2)) / \sqrt{\cos[c] \cos[dx + \text{ArcTan}[\text{Tan}[c]]] \sqrt{1 + \tan[c]^2}}) / (20d) + (B(a + a...$

3.142.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3454, 27, 3042, 3454, 3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

↓ 3454

3.142. $\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$

$$\frac{2}{5} \int \frac{(\cos(c+dx)a+a)^2(a(9A+5B)-a(A-5B)\cos(c+dx))}{2\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{5d\cos^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{(\cos(c+dx)a+a)^2(a(9A+5B)-a(A-5B)\cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{5d\cos^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2(a(9A+5B)-a(A-5B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{5d\cos^{\frac{5}{2}}(c+dx)}$$

↓ 3454

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{(\cos(c+dx)a+a)(a^2(21A+20B)-a^2(6A-5B)\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2(9A+5B)\sin(c+dx)(a^3\cos(c+dx))}{3d\cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{5d\cos^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)(a^2(21A+20B)-a^2(6A-5B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2(9A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{5d\cos^{\frac{5}{2}}(c+dx)}$$

↓ 3447

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{-((6A-5B)\cos^2(c+dx)a^3)+(21A+20B)a^3+(a^3(21A+20B)-a^3(6A-5B))\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2(9A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}{5d\cos^{\frac{5}{2}}(c+dx)}$$

↓ 3042

3.142. $\int \frac{(a+a\cos(c+dx))^3(A+B\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{-\left((6A - 5B) \sin\left(c + dx + \frac{\pi}{2}\right)^2 a^3\right) + (21A + 20B)a^3 + (a^3(21A + 20B) - a^3(6A - 5B)) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2}} dx + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3500

$$\frac{1}{5} \left(\frac{2}{3} \left(2 \int \frac{5a^3(3A + 5B) - 3a^3(9A + 5B) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2a^3(21A + 20B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2(9A + 5B) \sin(c + dx)}{3d \cos^{\frac{5}{2}}(c + dx)} + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{2}{3} \left(\int \frac{5a^3(3A + 5B) - 3a^3(9A + 5B) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2a^3(21A + 20B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2(9A + 5B) \sin(c + dx)}{3d \cos^{\frac{5}{2}}(c + dx)} + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(\int \frac{5a^3(3A + 5B) - 3a^3(9A + 5B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2a^3(21A + 20B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2(9A + 5B) \sin(c + dx)}{3d \cos^{\frac{5}{2}}(c + dx)} + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3227

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^3(3A + 5B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - 3a^3(9A + 5B) \int \sqrt{\cos(c + dx)} dx + \frac{2a^3(21A + 20B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2(9A + 5B) \sin(c + dx)}{3d \cos^{\frac{5}{2}}(c + dx)} + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^3(3A + 5B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx - 3a^3(9A + 5B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2a^3(21A + 20B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) + \frac{2(9A + 5B) \sin(c + dx)}{3d \cos^{\frac{5}{2}}(c + dx)} + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3119

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^3(3A + 5B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{6a^3(9A + 5B)E(\frac{1}{2}(c + dx)|2)}{d} + \frac{2a^3(21A + 20B)\sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) (a^3 \cos(c + dx) + a^3)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2}{3} \left(\frac{10a^3(3A + 5B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} - \frac{6a^3(9A + 5B)E(\frac{1}{2}(c + dx)|2)}{d} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

```
input Int[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]
```

```
output (2*a*A*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*(9*A + 5*B)*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2))) + (2*((-6*a^3*(9*A + 5*B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(3*A + 5*B)*EllipticF[(c + d*x)/2, 2])/d + (2*a^3*(21*A + 20*B)*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]])))/3/5
```

3.142.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

3.142. $\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

rule 3227 $\text{Int}[(b \cdot \sin(e) + f \cdot x)^m \cdot (c + d \cdot \sin(e) + f \cdot x)], x_{\text{Symbol}}] \rightarrow \text{Simp}[c \cdot \text{Int}[(b \cdot \sin[e + f \cdot x])^m, x], x] + \text{Simp}[d/b \cdot \text{Int}[(b \cdot \sin[e + f \cdot x])^{m+1}, x], x] /;$ $\text{FreeQ}[\{b, c, d, e, f, m\}, x]$

rule 3447 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) \cdot (c + d \cdot \sin(e) + f \cdot x)], x_{\text{Symbol}}] \rightarrow \text{Int}[(a + b \cdot \sin[e + f \cdot x])^m \cdot (A \cdot c + (B \cdot c + A \cdot d) \cdot \sin[e + f \cdot x] + B \cdot d \cdot \sin[e + f \cdot x]^2), x] /;$ $\text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0]$

rule 3454 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) \cdot (c + d \cdot \sin(e) + f \cdot x)^n], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-b^2) \cdot (B \cdot c - A \cdot d) \cdot \cos[e + f \cdot x] \cdot (a + b \cdot \sin[e + f \cdot x])^{m-1} \cdot (c + d \cdot \sin[e + f \cdot x])^{n+1} / (d \cdot f \cdot (n+1) \cdot (b \cdot c + a \cdot d)), x] - \text{Simp}[b / (d \cdot (n+1) \cdot (b \cdot c + a \cdot d)) \cdot \text{Int}[(a + b \cdot \sin[e + f \cdot x])^{m-1} \cdot (c + d \cdot \sin[e + f \cdot x])^{n+1} \cdot \text{Simp}[a \cdot A \cdot d \cdot (m - n - 2) - B \cdot (a \cdot c \cdot (m - 1) + b \cdot d \cdot (n + 1)) - (A \cdot b \cdot d \cdot (m + n + 1) - B \cdot (b \cdot c \cdot m - a \cdot d \cdot (n + 1))) \cdot \sin[e + f \cdot x], x], x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{GtQ}[m, 1/2] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2 \cdot m] \ \&\& \ (\text{IntegerQ}[2 \cdot n] \ || \ \text{EqQ}[c, 0])]$

rule 3500 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) + (C \cdot \sin(e) + f \cdot x)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C) \cdot \cos[e + f \cdot x] \cdot (a + b \cdot \sin[e + f \cdot x])^{m+1} / (b \cdot f \cdot (m+1) \cdot (a^2 - b^2)), x] + \text{Simp}[1 / (b \cdot (m+1) \cdot (a^2 - b^2)) \cdot \text{Int}[(a + b \cdot \sin[e + f \cdot x])^{m+1} \cdot \text{Simp}[b \cdot (a \cdot A - b \cdot B + a \cdot C) \cdot (m+1) - (A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C + b \cdot (A \cdot b - a \cdot B + b \cdot C) \cdot (m+1)) \cdot \sin[e + f \cdot x], x], x], x] /;$ $\text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

3.142.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 915 vs. $2(207) = 414$.

Time = 12.00 (sec) , antiderivative size = 916, normalized size of antiderivative = 5.36

method	result	size
default	Expression too large to display	916
parts	Expression too large to display	946

3.142.
$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^2(c + dx)} dx$$


```
input int((a+cos(d*x+c))*a)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/15*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3/(8*sin
(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/
2*d*x+1/2*c)^3*(216*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-60*A*Ellipti
cF(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4-108*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*si
n(1/2*d*x+1/2*c)^4+180*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-100*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-60*B*EllipticE(cos(1/2*d*x+1/2
*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*sin(1/2*d*x+1/2*c)^4-246*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+60*A*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1
/2*d*x+1/2*c)^2-1)^(1/2)*sin(1/2*d*x+1/2*c)^2+108*A*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/
2))*sin(1/2*d*x+1/2*c)^2-190*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+100
*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF
(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+60*B*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)
^(1/2)*sin(1/2*d*x+1/2*c)^2+72*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2-1
5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellip...
```

3.142.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.42

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx =$$

$$2 \left(5i \sqrt{2} (3A + 5B) a^3 \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (3A + 5B) a^3 \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) \right)$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,algorith
m="fracas")
```

3.142. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

output `-2/15*(5*I*sqrt(2)*(3*A + 5*B)*a^3*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(3*A + 5*B)*a^3*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(9*A + 5*B)*a^3*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(9*A + 5*B)*a^3*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (9*(6*A + 5*B)*a^3*cos(d*x + c)^2 + 5*(3*A + B)*a^3*cos(d*x + c) + 3*A*a^3)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.142.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.142.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{7}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(7/2), x)`

3.142.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(7/2), x)`

3.142.9 Mupad [B] (verification not implemented)

Time = 2.73 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.68

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2 (B a^3 E(\frac{c}{2} + \frac{dx}{2} | 2) + 3 B a^3 F(\frac{c}{2} + \frac{dx}{2} | 2))}{d} + \frac{2 A a^3 F(\frac{c}{2} + \frac{dx}{2} | 2)}{d}$$

$$+ \frac{6 A a^3 \sin(c + dx) {}_2F_1(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 A a^3 \sin(c + dx) {}_2F_1(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2)}{d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 A a^3 \sin(c + dx) {}_2F_1(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2)}{5 d \cos(c + dx)^{5/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{6 B a^3 \sin(c + dx) {}_2F_1(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^3 \sin(c + dx) {}_2F_1(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^(7/2),x)`

3.142. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

output $(2*(B*a^3*ellipticE(c/2 + (d*x)/2, 2) + 3*B*a^3*ellipticF(c/2 + (d*x)/2, 2)))/d + (2*A*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (6*A*a^3*\sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, \cos(c + d*x)^2))/(d*\cos(c + d*x)^{(1/2)}*(\sin(c + d*x)^2)^{(1/2)}) + (2*A*a^3*\sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, \cos(c + d*x)^2))/(d*\cos(c + d*x)^{(3/2)}*(\sin(c + d*x)^2)^{(1/2)}) + (2*A*a^3*\sin(c + d*x)*hypergeom([-5/4, 1/2], -1/4, \cos(c + d*x)^2))/(5*d*\cos(c + d*x)^{(5/2)}*(\sin(c + d*x)^2)^{(1/2)}) + (6*B*a^3*\sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, \cos(c + d*x)^2))/(d*\cos(c + d*x)^{(1/2)}*(\sin(c + d*x)^2)^{(1/2)}) + (2*B*a^3*\sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, \cos(c + d*x)^2))/(3*d*\cos(c + d*x)^{(3/2)}*(\sin(c + d*x)^2)^{(1/2)})$

3.143
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.143.1 Optimal result 1406
 3.143.2 Mathematica [C] (warning: unable to verify) 1407
 3.143.3 Rubi [A] (verified) 1408
 3.143.4 Maple [B] (verified) 1413
 3.143.5 Fracas [C] (verification not implemented) 1414
 3.143.6 Sympy [F(-1)] 1415
 3.143.7 Maxima [F] 1415
 3.143.8 Giac [F] 1416
 3.143.9 Mupad [B] (verification not implemented) 1416

3.143.1 Optimal result

Integrand size = 33, antiderivative size = 204

$$\int \frac{(a + a \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= -\frac{4a^3(7A + 9B)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{4a^3(13A + 21B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d}$$

$$+ \frac{4a^3(41A + 42B) \sin(c + dx)}{105d \cos^{\frac{3}{2}}(c + dx)} + \frac{4a^3(7A + 9B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

$$+ \frac{2aA(a + a \cos(c + dx))^2 \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx)} + \frac{2(11A + 7B)(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{35d \cos^{\frac{5}{2}}(c + dx)}$$

```
output -4/5*a^3*(7*A+9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellipti
cE(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/21*a^3*(13*A+21*B)*(cos(1/2*d*x+1/2*c)^
2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+4/105*
a^3*(41*A+42*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/7*a*A*(a+a*cos(d*x+c))^2*s
in(d*x+c)/d/cos(d*x+c)^(7/2)+2/35*(11*A+7*B)*(a^3+a^3*cos(d*x+c))*sin(d*x+
c)/d/cos(d*x+c)^(5/2)+4/5*a^3*(7*A+9*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.143.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.88 (sec) , antiderivative size = 925, normalized size of antiderivative = 4.53

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^3 \sec^6 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(\frac{(7A + 9B) \csc(c) \sec(c)}{10d} + \frac{A \sec(c) \sec^4(c + dx) \sin(dx)}{28d} + \frac{\sec(c) \sec^3(c + dx) (5A \sin(c) + 21A \sin(dx) + 7B \sin(dx))}{140d} + \frac{\sec(c) \sec^2(c + dx) (63A \sin(c) + 21B \sin(c) + 130A \sin(dx) + 105B \sin(dx))}{420d} + \frac{\sec(c) \sec(c + dx) (130A \sin(c) + 105B \sin(c) + 294A \sin(dx) + 378B \sin(dx))}{420d} \right) \\ - \frac{13A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{4d} \\ - \frac{B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{2d} \\ + \frac{7A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d} \\ + \frac{9B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d}$$

input `Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2), x]`

output $\sqrt{\cos[c + dx]}(a + a\cos[c + dx])^3 \sec[c/2 + (dx)/2]^6 ((7A + 9B) \operatorname{Csc}[c] \operatorname{Sec}[c]) / (10d) + (A \operatorname{Sec}[c] \operatorname{Sec}[c + dx]^4 \sin[dx]) / (28d) + (\operatorname{Sec}[c] \operatorname{Sec}[c + dx]^3 (5A \sin[c] + 21A \sin[dx] + 7B \sin[dx])) / (140d) + (\operatorname{Sec}[c] \operatorname{Sec}[c + dx]^2 (63A \sin[c] + 21B \sin[c] + 130A \sin[dx] + 105B \sin[dx])) / (420d) + (\operatorname{Sec}[c] \operatorname{Sec}[c + dx] (130A \sin[c] + 105B \sin[c] + 294A \sin[dx] + 378B \sin[dx])) / (420d) - (13A (a + a\cos[c + dx])^3 \operatorname{Csc}[c] \operatorname{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]^2] \operatorname{Sec}[c/2 + (dx)/2]^6 \operatorname{Sec}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]] \sqrt{1 - \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]}] \sqrt{-(\sqrt{1 + \operatorname{Cot}[c]^2} \sin[c] \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]])} \sqrt{1 + \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]}) / (42d \sqrt{1 + \operatorname{Cot}[c]^2}) - (B (a + a\cos[c + dx])^3 \operatorname{Csc}[c] \operatorname{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]^2] \operatorname{Sec}[c/2 + (dx)/2]^6 \operatorname{Sec}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]] \sqrt{1 - \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]}] \sqrt{-(\sqrt{1 + \operatorname{Cot}[c]^2} \sin[c] \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]])} \sqrt{1 + \sin[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]}) / (2d \sqrt{1 + \operatorname{Cot}[c]^2}) + (7A (a + a\cos[c + dx])^3 \operatorname{Csc}[c] \operatorname{Sec}[c/2 + (dx)/2]^6 (\operatorname{HypergeometricPFQ}[-1/2, -1/4, \{3/4\}, \cos[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]]^2] \sin[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Tan}[c]) / (\sqrt{1 - \cos[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]}] \sqrt{1 + \cos[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]}] \sqrt{\cos[c] \cos[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \sqrt{1 + \operatorname{Tan}[c]^2}} \sqrt{1 + \operatorname{Tan}[c]^2}) - ((\sin[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Tan}[c]) / \sqrt{1 + \operatorname{Tan}[c]^2} + (2 \operatorname{Cos}[c]^2 \cos[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \sqrt{1 + \operatorname{Tan}[c]^2}) / (\operatorname{Cos}[c]^2 + \sin...$

3.143.3 Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.01, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3454, 27, 3042, 3454, 3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{\frac{9}{2}}} dx$$

↓ 3454

3.143. $\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{2}{7} \int \frac{(\cos(c+dx)a+a)^2(a(11A+7B)+a(A+7B)\cos(c+dx))}{\frac{2\cos^{\frac{7}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \\
& \qquad \qquad \qquad \frac{7d\cos^{\frac{7}{2}}(c+dx)}{27} \\
& \frac{1}{7} \int \frac{(\cos(c+dx)a+a)^2(a(11A+7B)+a(A+7B)\cos(c+dx))}{\frac{\cos^{\frac{7}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \\
& \qquad \qquad \qquad \frac{7d\cos^{\frac{7}{2}}(c+dx)}{3042} \\
& \frac{1}{7} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2(a(11A+7B)+a(A+7B)\sin(c+dx+\frac{\pi}{2}))}{\frac{\sin(c+dx+\frac{\pi}{2})^{7/2}}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \\
& \qquad \qquad \qquad \frac{7d\cos^{\frac{7}{2}}(c+dx)}{3454} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{(\cos(c+dx)a+a)((41A+42B)a^2+(8A+21B)\cos(c+dx)a^2)}{\frac{\cos^{\frac{5}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \frac{2(11A+7B)\sin(c+dx)(a^3\cos(c+dx))}{5d\cos^{\frac{5}{2}}(c+dx)} \right) \\
& \qquad \qquad \qquad \frac{7d\cos^{\frac{7}{2}}(c+dx)}{3042} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)((41A+42B)a^2+(8A+21B)\sin(c+dx+\frac{\pi}{2})a^2)}{\frac{\sin(c+dx+\frac{\pi}{2})^{5/2}}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \frac{2(11A+7B)\sin(c+dx)(a^3\cos(c+dx))}{5d\cos^{\frac{5}{2}}(c+dx)} \right) \\
& \qquad \qquad \qquad \frac{7d\cos^{\frac{7}{2}}(c+dx)}{3447} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{(8A+21B)\cos^2(c+dx)a^3+(41A+42B)a^3+((8A+21B)a^3+(41A+42B)a^3)\cos(c+dx)}{\frac{\cos^{\frac{5}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \frac{2(11A+7B)\sin(c+dx)(a^3\cos(c+dx))}{5d\cos^{\frac{5}{2}}(c+dx)} \right) \\
& \qquad \qquad \qquad \frac{7d\cos^{\frac{7}{2}}(c+dx)}{3042}
\end{aligned}$$

3.143. $\int \frac{(a+a\cos(c+dx))^3(A+B\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{(8A + 21B) \sin(c + dx + \frac{\pi}{2})^2 a^3 + (41A + 42B)a^3 + ((8A + 21B)a^3 + (41A + 42B)a^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{5/2} \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)}} dx \right)$$

↓ 3500

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{2}{3} \int \frac{21(7A + 9B)a^3 + 5(13A + 21B) \cos(c + dx)a^3}{2 \cos^{\frac{3}{2}}(c + dx)} dx + \frac{2a^3(41A + 42B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2(11A + 7B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \frac{21(7A + 9B)a^3 + 5(13A + 21B) \cos(c + dx)a^3}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2a^3(41A + 42B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2(11A + 7B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \frac{21(7A + 9B)a^3 + 5(13A + 21B) \sin(c + dx + \frac{\pi}{2})a^3}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a^3(41A + 42B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2(11A + 7B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)}$$

↓ 3227

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(21a^3(7A + 9B) \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx + 5a^3(13A + 21B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \right) + \frac{2a^3(41A + 42B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(21a^3(7A + 9B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + 5a^3(13A + 21B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2a^3(41A + 42B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3116

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right) \right) \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) \right) \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)} \right) \right)$$

↓ 3119

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx)|2)}{d} \right) \right) \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{7} \left(\frac{2(11A + 7B) \sin(c + dx) (a^3 \cos(c + dx) + a^3)}{5d \cos^{\frac{5}{2}}(c + dx)} + \frac{2}{5} \left(\frac{2a^3(41A + 42B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{1}{3} \left(\frac{10a^3(13A + 21B) \text{EllipticE}[(c + dx)/2, 2]}{d} \right) \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `(2*a*A*(a + a*Cos[c + d*x])^2*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(11*A + 7*B)*(a^3 + a^3*Cos[c + d*x])*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (2*((2*a^3*(41*A + 42*B))*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((10*a^3*(13*A + 21*B)*EllipticF[(c + d*x)/2, 2])/d + 21*a^3*(7*A + 9*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x])))))/3)/5)/7`

3.143.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3116 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m - 1)*((c + d*Ssin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Ssin[e + f*x])^(m - 1)*(c + d*Ssin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3500 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] :> Simp[(-A*b^2
- a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.143.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 901 vs. $2(236) = 472$.

Time = 13.76 (sec) , antiderivative size = 902, normalized size of antiderivative = 4.42

method	result	size
default	Expression too large to display	902
parts	Expression too large to display	1056

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RETURNV
ERBOSE)
```

output

```

-16*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(1/8*B*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*
d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1
/2))+1/8*A*(-1/56*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^4-5/42*cos(1/2*d*x+1/2*c)*(-2*s
in(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)
^2+5/21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2)))+1/5*(1/8*B+3/8*A)/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/
2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-
24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*si
n(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*
x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1
/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+(3/8*
A+3/8*B)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x...

```

3.143.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.29

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx =$$

$$\frac{2 \left(5i \sqrt{2} (13A + 21B) a^3 \cos(dx + c)^4 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} \right)}{\dots}$$

input

```

integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorith
m="fracas")

```

output `-2/105*(5*I*sqrt(2)*(13*A + 21*B)*a^3*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(13*A + 21*B)*a^3*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*I*sqrt(2)*(7*A + 9*B)*a^3*cos(d*x + c)^4*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*(7*A + 9*B)*a^3*cos(d*x + c)^4*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (42*(7*A + 9*B)*a^3*cos(d*x + c)^3 + 5*(26*A + 21*B)*a^3*cos(d*x + c)^2 + 21*(3*A + B)*a^3*cos(d*x + c) + 15*A*a^3)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^4)`

3.143.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output `Timed out`

3.143.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{9}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(9/2), x)`

3.143.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{9}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(9/2), x)`

3.143.9 Mupad [B] (verification not implemented)

Time = 2.77 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.50

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \frac{2 B a^3 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d}$$

$$+ \frac{2 A a^3 \sin(c+dx) {}_2F_1\left(-\frac{7}{4}, \frac{1}{2}; -\frac{3}{4}; \cos(c+dx)^2\right)}{7} + \frac{6 A a^3 \cos(c+dx) \sin(c+dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c+dx)^2\right)}{5} + 2 A a^3 \cos(c + dx)$$

$$+ \frac{6 B a^3 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^3 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^3 \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right)}{5 d \cos(c + dx)^{5/2} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^(9/2),x)`

output $(2*B*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + ((2*A*a^3*\sin(c + d*x)*hypergeom$
 $([-7/4, 1/2], -3/4, \cos(c + d*x)^2))/7 + (6*A*a^3*\cos(c + d*x)*\sin(c + d*x$
 $)*hypergeom([-5/4, 1/2], -1/4, \cos(c + d*x)^2))/5 + 2*A*a^3*\cos(c + d*x)^2$
 $*\sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, \cos(c + d*x)^2) + 2*A*a^3*\cos(c$
 $+ d*x)^3*\sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, \cos(c + d*x)^2))/(d*\cos(c$
 $+ d*x)^(7/2)*(1 - \cos(c + d*x)^2)^(1/2)) + (6*B*a^3*\sin(c + d*x)*hyperge$
 $om([-1/4, 1/2], 3/4, \cos(c + d*x)^2))/(d*\cos(c + d*x)^(1/2)*(\sin(c + d*x)^$
 $2)^(1/2)) + (2*B*a^3*\sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, \cos(c + d*x)$
 $^2))/(d*\cos(c + d*x)^(3/2)*(\sin(c + d*x)^2)^(1/2)) + (2*B*a^3*\sin(c + d*x)$
 $*hypergeom([-5/4, 1/2], -1/4, \cos(c + d*x)^2))/(5*d*\cos(c + d*x)^(5/2)*(si$
 $n(c + d*x)^2)^(1/2))$

3.144
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

3.144.1 Optimal result 1418
 3.144.2 Mathematica [C] (warning: unable to verify) 1419
 3.144.3 Rubi [A] (verified) 1420
 3.144.4 Maple [B] (verified) 1425
 3.144.5 Fricas [C] (verification not implemented) 1426
 3.144.6 Sympy [F(-1)] 1427
 3.144.7 Maxima [F] 1427
 3.144.8 Giac [F] 1428
 3.144.9 Mupad [B] (verification not implemented) 1428

3.144.1 Optimal result

Integrand size = 33, antiderivative size = 237

$$\int \frac{(a + a \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$= -\frac{4a^3(17A + 21B)E(\frac{1}{2}(c + dx) | 2)}{15d} + \frac{4a^3(11A + 13B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d}$$

$$+ \frac{4a^3(23A + 24B) \sin(c + dx)}{105d \cos^{\frac{5}{2}}(c + dx)} + \frac{4a^3(11A + 13B) \sin(c + dx)}{21d \cos^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{4a^3(17A + 21B) \sin(c + dx)}{15d \sqrt{\cos(c + dx)}} + \frac{2aA(a + a \cos(c + dx))^2 \sin(c + dx)}{9d \cos^{\frac{9}{2}}(c + dx)}$$

$$+ \frac{2(13A + 9B)(a^3 + a^3 \cos(c + dx)) \sin(c + dx)}{63d \cos^{\frac{7}{2}}(c + dx)}$$

```
output -4/15*a^3*(17*A+21*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elli
pticE(sin(1/2*d*x+1/2*c), 2^(1/2))/d+4/21*a^3*(11*A+13*B)*(cos(1/2*d*x+1/2*
c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/d+4/1
05*a^3*(23*A+24*B)*sin(d*x+c)/d/cos(d*x+c)^(5/2)+4/21*a^3*(11*A+13*B)*sin(
d*x+c)/d/cos(d*x+c)^(3/2)+2/9*a*A*(a+a*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+
c)^(9/2)+2/63*(13*A+9*B)*(a^3+a^3*cos(d*x+c))*sin(d*x+c)/d/cos(d*x+c)^(7/2
)+4/15*a^3*(17*A+21*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.144.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 9.71 (sec) , antiderivative size = 967, normalized size of antiderivative = 4.08

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx = \sqrt{\cos(c + dx)} (a + a \cos(c + dx))^3 \sec^6 \left(\frac{c}{2} + \frac{dx}{2} \right) \left(\frac{(17A + 21B) \csc(c) \sec(c)}{30d} + \frac{A \sec(c) \sec^5(c + dx) \sin(dx)}{36d} + \frac{\sec(c) \sec^4(c + dx) (7A \sin(c) + 27A \sin(dx) + 9B \sin(dx))}{252d} + \frac{\sec(c) \sec(c + dx) (55A \sin(c) + 65B \sin(c) + 119A \sin(dx) + 147B \sin(dx))}{210d} + \frac{\sec(c) \sec^3(c + dx) (135A \sin(c) + 45B \sin(c) + 238A \sin(dx) + 189B \sin(dx))}{1260d} + \frac{\sec(c) \sec^2(c + dx) (238A \sin(c) + 189B \sin(c) + 330A \sin(dx) + 390B \sin(dx))}{1260d} \right)$$

$$- \frac{11A(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{4d}$$

$$- \frac{13B(a + a \cos(c + dx))^3 \csc(c) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sec(dx - \arctan(\cot(c)))}{4d}$$

$$+ \frac{17A(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{60d}$$

$$+ \frac{7B(a + a \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c)))}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))}} \right)}{20d}$$

input `Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2),x]`

output `Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^3*Sec[c/2 + (d*x)/2]^6*(((17*A + 21*B)*Csc[c]*Sec[c])/(30*d) + (A*Sec[c]*Sec[c + d*x]^5*Sin[d*x])/(36*d) + (Sec[c]*Sec[c + d*x]^4*(7*A*Sin[c] + 27*A*Sin[d*x] + 9*B*Sin[d*x]))/(252*d) + (Sec[c]*Sec[c + d*x]*(55*A*Sin[c] + 65*B*Sin[c] + 119*A*Sin[d*x] + 147*B*Sin[d*x]))/(210*d) + (Sec[c]*Sec[c + d*x]^3*(135*A*Sin[c] + 45*B*Sin[c] + 238*A*Sin[d*x] + 189*B*Sin[d*x]))/(1260*d) + (Sec[c]*Sec[c + d*x]^2*(238*A*Sin[c] + 189*B*Sin[c] + 330*A*Sin[d*x] + 390*B*Sin[d*x]))/(1260*d)) - (11*A*(a + a*cos[c + d*x])^3*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(42*d*Sqrt[1 + Cot[c]^2]) - (13*B*(a + a*cos[c + d*x])^3*Csc[c]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2 + (d*x)/2]^6*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(42*d*Sqrt[1 + Cot[c]^2]) + (17*A*(a + a*cos[c + d*x])^3*Csc[c]*Sec[c/2 + (d*x)/2]^6*(HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Tan[c]^2])*Sqrt[1 + Tan[c]^2)) - ((Sin[d*x + ArcTan[Tan[c]...`

3.144.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 234, normalized size of antiderivative = 0.99, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx$$

↓ 3454

3.144. $\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{2}{9} \int \frac{(\cos(c+dx)a+a)^2(a(13A+9B)+3a(A+3B)\cos(c+dx))}{\frac{2\cos^{\frac{9}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \\
& \qquad \qquad \qquad \frac{9d\cos^{\frac{9}{2}}(c+dx)}{27} \\
& \frac{1}{9} \int \frac{(\cos(c+dx)a+a)^2(a(13A+9B)+3a(A+3B)\cos(c+dx))}{\frac{\cos^{\frac{9}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \\
& \qquad \qquad \qquad \frac{9d\cos^{\frac{9}{2}}(c+dx)}{3042} \\
& \frac{1}{9} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^2(a(13A+9B)+3a(A+3B)\sin(c+dx+\frac{\pi}{2}))}{\frac{\sin(c+dx+\frac{\pi}{2})^{9/2}}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \\
& \qquad \qquad \qquad \frac{9d\cos^{\frac{9}{2}}(c+dx)}{3454} \\
& \frac{1}{9} \left(\frac{2}{7} \int \frac{3(\cos(c+dx)a+a)((23A+24B)a^2+5(2A+3B)\cos(c+dx)a^2)}{\frac{\cos^{\frac{7}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \frac{2(13A+9B)\sin(c+dx)(a^3\cos(c+dx))}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \\
& \qquad \qquad \qquad \frac{9d\cos^{\frac{9}{2}}(c+dx)}{27} \\
& \frac{1}{9} \left(\frac{6}{7} \int \frac{(\cos(c+dx)a+a)((23A+24B)a^2+5(2A+3B)\cos(c+dx)a^2)}{\frac{\cos^{\frac{7}{2}}(c+dx)}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \frac{2(13A+9B)\sin(c+dx)(a^3\cos(c+dx))}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \\
& \qquad \qquad \qquad \frac{9d\cos^{\frac{9}{2}}(c+dx)}{3042} \\
& \frac{1}{9} \left(\frac{6}{7} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)((23A+24B)a^2+5(2A+3B)\sin(c+dx+\frac{\pi}{2})a^2)}{\frac{\sin(c+dx+\frac{\pi}{2})^{7/2}}{2aA\sin(c+dx)(a\cos(c+dx)+a)^2}} dx + \frac{2(13A+9B)\sin(c+dx)(a^3\cos(c+dx))}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \\
& \qquad \qquad \qquad \frac{9d\cos^{\frac{9}{2}}(c+dx)}{3447}
\end{aligned}$$

3.144. $\int \frac{(a+a\cos(c+dx))^3(A+B\cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{6}{7} \int \frac{5(2A+3B) \cos^2(c+dx) a^3 + (23A+24B) a^3 + (5(2A+3B) a^3 + (23A+24B) a^3) \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)} dx + \frac{2(13A+9B)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \int \frac{5(2A+3B) \sin(c+dx + \frac{\pi}{2})^2 a^3 + (23A+24B) a^3 + (5(2A+3B) a^3 + (23A+24B) a^3) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{7/2}} dx + \frac{2(13A+9B)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3500

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{2}{5} \int \frac{15(11A+13B) a^3 + 7(17A+21B) \cos(c+dx) a^3}{2 \cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^3(23A+24B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right) + \frac{2(13A+9B)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \int \frac{15(11A+13B) a^3 + 7(17A+21B) \cos(c+dx) a^3}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^3(23A+24B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right) + \frac{2(13A+9B)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \int \frac{15(11A+13B) a^3 + 7(17A+21B) \sin(c+dx + \frac{\pi}{2}) a^3}{\sin(c+dx + \frac{\pi}{2})^{5/2}} dx + \frac{2a^3(23A+24B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right) + \frac{2(13A+9B)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3227

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(11A+13B) \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)} dx + 7a^3(17A+21B) \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx \right) + \frac{2a^3(23A+24B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^2}{9d \cos^{\frac{9}{2}}(c+dx)} \right)$$

↓ 3042

3.144. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(11A + 13B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + 7a^3(17A + 21B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \frac{2a^3(23A + 24B) \sin(c + dx)}{5d \cos^{5/2}(c + dx)} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{9d \cos^{9/2}(c + dx)} \right) \\ \downarrow \text{3116}$$

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(11A + 13B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d \cos^{3/2}(c + dx)} \right) + 7a^3(17A + 21B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} \right) \right) \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{9d \cos^{9/2}(c + dx)} \right) \right) \\ \downarrow \text{3042}$$

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(11A + 13B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{3/2}(c + dx)} \right) + 7a^3(17A + 21B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} \right) \right) \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{9d \cos^{9/2}(c + dx)} \right) \right) \\ \downarrow \text{3119}$$

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(11A + 13B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{3/2}(c + dx)} \right) + 7a^3(17A + 21B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} \right) \right) \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{9d \cos^{9/2}(c + dx)} \right) \right) \\ \downarrow \text{3120}$$

$$\frac{1}{9} \left(\frac{2(13A + 9B) \sin(c + dx) (a^3 \cos(c + dx) + a^3)}{7d \cos^{7/2}(c + dx)} + \frac{6}{7} \left(\frac{2a^3(23A + 24B) \sin(c + dx)}{5d \cos^{5/2}(c + dx)} + \frac{1}{5} \left(15a^3(11A + 13B) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} \right) \right) \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^2}{9d \cos^{9/2}(c + dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2),x]`

```
output (2*a*A*(a + a*cos[c + d*x])^2*sin[c + d*x])/(9*d*cos[c + d*x]^(9/2)) + ((2
*(13*A + 9*B)*(a^3 + a^3*cos[c + d*x])*sin[c + d*x])/(7*d*cos[c + d*x]^(7/
2)) + (6*((2*a^3*(23*A + 24*B)*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)) + (1
5*a^3*(11*A + 13*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*sin[c + d*x]
)/(3*d*cos[c + d*x]^(3/2))) + 7*a^3*(17*A + 21*B)*((-2*EllipticE[(c + d*x)
/2, 2])/d + (2*sin[c + d*x])/(d*Sqrt[Cos[c + d*x]))))/5))/7)/9
```

3.144.3.1 Defintions of rubi rules used

```
rule 277 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3116 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) I
nt[(b*sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*sin[e + f*x])^m*(A*c + (B*c + A*d)*sin[e + f*x] + B*d*sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3500 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] :> Simp[(-A*b^2
- a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.144.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1150 vs. $2(265) = 530$.

Time = 18.18 (sec) , antiderivative size = 1151, normalized size of antiderivative = 4.86

method	result	size
default	Expression too large to display	1151
parts	Expression too large to display	1421

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x,method=_RETURN
VERBOSE)
```


output

```

-16*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(1/8*B/s
in(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(
1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2
*d*x+1/2*c),2^(1/2)))+1/8*A*(-1/144*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2
*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^5-7/180*cos(1
/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/
2*d*x+1/2*c)^2-1/2)^3-14/15*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)/(-(-2*
cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)+7/15*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin
(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-7/15*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1
/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))
-EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))+(1/8*B+3/8*A)*(-1/56*cos(1/2*d*x+
1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1
/2*c)^2-1/2)^4-5/42*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*
x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+5/21*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*
d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+1/5*(3/8*A+3/8*
B)/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)...

```

3.144.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.19

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx =$$

$$\frac{2 \left(15i \sqrt{2} (11A + 13B) a^3 \cos(dx + c)^5 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 15i \right)}{\dots}$$

input

```

integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorit
hm="fricas")

```

output
$$\begin{aligned} & -2/315*(15*I*\sqrt{2}*(11*A + 13*B)*a^3*\cos(dx + c)^5*\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I*\sin(dx + c)) - 15*I*\sqrt{2}*(11*A + 13*B)*a^3*\cos(dx + c)^5*\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I*\sin(dx + c)) + 21*I*\sqrt{2}*(17*A + 21*B)*a^3*\cos(dx + c)^5*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I*\sin(dx + c))) - 21*I*\sqrt{2}*(17*A + 21*B)*a^3*\cos(dx + c)^5*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I*\sin(dx + c))) - (42*(17*A + 21*B)*a^3*\cos(dx + c)^4 + 30*(11*A + 13*B)*a^3*\cos(dx + c)^3 + 7*(34*A + 27*B)*a^3*\cos(dx + c)^2 + 45*(3*A + B)*a^3*\cos(dx + c) + 35*A*a^3)*\sqrt{\cos(dx + c)}*\sin(dx + c))/(d*\cos(dx + c)^5) \end{aligned}$$

3.144.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(11/2),x)`

output Timed out

3.144.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx \\ & = \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{11}{2}}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))3*(A+B*cos(d*x+c))/cos(d*x+c)(11/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)3/cos(d*x + c)(11/2), x)`

3.144.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{11}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/cos(d*x + c)^(11/2), x)`

3.144.9 Mupad [B] (verification not implemented)

Time = 3.29 (sec) , antiderivative size = 552, normalized size of antiderivative = 2.33

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$= \frac{2 {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right) \left(\frac{19 A a^3 \sin(c+dx)}{\cos(c+dx)^{3/2} \sqrt{1-\cos(c+dx)^2}} + \frac{9 A a^3 \sin(c+dx)}{\cos(c+dx)^{7/2} \sqrt{1-\cos(c+dx)^2}} + \frac{25 B a^3 \sin(c+dx)}{\cos(c+dx)^{3/2} \sqrt{1-\cos(c+dx)^2}} \right)}{21 d}$$

$$- \frac{8 {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{7}{4}; \cos(c + dx)^2\right) \left(\frac{34 A a^3 \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{1-\cos(c+dx)^2}} + \frac{5 A a^3 \sin(c+dx)}{\cos(c+dx)^{5/2} \sqrt{1-\cos(c+dx)^2}} + \frac{27 B a^3 \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{1-\cos(c+dx)^2}} \right)}{135 d}$$

$$+ \frac{8 \left(\frac{3 A a^3 \sin(c+dx)}{\cos(c+dx)^{3/2} \sqrt{1-\cos(c+dx)^2}} + \frac{B a^3 \sin(c+dx)}{\cos(c+dx)^{3/2} \sqrt{1-\cos(c+dx)^2}} \right) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{5}{4}; \cos(c + dx)^2\right)}{21 d}$$

$$+ \frac{2 {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right) \left(\frac{136 A a^3 \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{1-\cos(c+dx)^2}} + \frac{39 A a^3 \sin(c+dx)}{\cos(c+dx)^{5/2} \sqrt{1-\cos(c+dx)^2}} + \frac{5 A a^3 \sin(c+dx)}{\cos(c+dx)^{9/2} \sqrt{1-\cos(c+dx)^2}} \right)}{45 d}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/cos(c + d*x)^(11/2),x)`

output

```
(2*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2)*((19*A*a^3*sin(c + d*x))/(cos(c + d*x)^(3/2)*(1 - cos(c + d*x)^2)^(1/2)) + (9*A*a^3*sin(c + d*x))/(cos(c + d*x)^(7/2)*(1 - cos(c + d*x)^2)^(1/2)) + (25*B*a^3*sin(c + d*x))/(cos(c + d*x)^(3/2)*(1 - cos(c + d*x)^2)^(1/2)) + (3*B*a^3*sin(c + d*x))/(cos(c + d*x)^(7/2)*(1 - cos(c + d*x)^2)^(1/2))))/(21*d) - (8*hypergeom([-1/4, 1/2], 7/4, cos(c + d*x)^2)*((34*A*a^3*sin(c + d*x))/(cos(c + d*x)^(1/2)*(1 - cos(c + d*x)^2)^(1/2)) + (5*A*a^3*sin(c + d*x))/(cos(c + d*x)^(5/2)*(1 - cos(c + d*x)^2)^(1/2)) + (27*B*a^3*sin(c + d*x))/(cos(c + d*x)^(1/2)*(1 - cos(c + d*x)^2)^(1/2))))/(135*d) + (8*((3*A*a^3*sin(c + d*x))/(cos(c + d*x)^(3/2)*(1 - cos(c + d*x)^2)^(1/2)) + (B*a^3*sin(c + d*x))/(cos(c + d*x)^(3/2)*(1 - cos(c + d*x)^2)^(1/2))))*hypergeom([-3/4, 1/2], 5/4, cos(c + d*x)^2))/(21*d) + (2*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2)*((136*A*a^3*sin(c + d*x))/(cos(c + d*x)^(1/2)*(1 - cos(c + d*x)^2)^(1/2)) + (39*A*a^3*sin(c + d*x))/(cos(c + d*x)^(5/2)*(1 - cos(c + d*x)^2)^(1/2)) + (5*A*a^3*sin(c + d*x))/(cos(c + d*x)^(9/2)*(1 - cos(c + d*x)^2)^(1/2)) + (153*B*a^3*sin(c + d*x))/(cos(c + d*x)^(1/2)*(1 - cos(c + d*x)^2)^(1/2)) + (27*B*a^3*sin(c + d*x))/(cos(c + d*x)^(5/2)*(1 - cos(c + d*x)^2)^(1/2))))/(45*d)
```

3.145
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$$

3.145.1 Optimal result 1430
 3.145.2 Mathematica [C] (warning: unable to verify) 1431
 3.145.3 Rubi [A] (verified) 1432
 3.145.4 Maple [A] (verified) 1435
 3.145.5 Fricas [C] (verification not implemented) 1436
 3.145.6 Sympy [F(-1)] 1436
 3.145.7 Maxima [F] 1437
 3.145.8 Giac [F] 1437
 3.145.9 Mupad [F(-1)] 1437

3.145.1 Optimal result

Integrand size = 33, antiderivative size = 156

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = -\frac{3(5A-7B)E(\frac{1}{2}(c+dx)|2)}{5ad} + \frac{5(A-B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{3ad} + \frac{5(A-B)\sqrt{\cos(c+dx)}\sin(c+dx)}{3ad} - \frac{(5A-7B)\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{5ad} + \frac{(A-B)\cos^{\frac{5}{2}}(c+dx)\sin(c+dx)}{d(a+a \cos(c+dx))}$$

output

```
-3/5*(5*A-7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a/d+5/3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a/d-1/5*(5*A-7*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/a/d+(A-B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))+5/3*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d
```

3.145.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.42 (sec) , antiderivative size = 946, normalized size of antiderivative = 6.06

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \left(\frac{2(5A-5B+10A\cos(c)-16B\cos(c))\csc(c)}{5d} + \frac{4(A-B)\cos(dx)\sin(c)}{3d} + \frac{2B\cos(2dx)\sin(2c)}{5d} + \frac{2\sec(c)}{5d} \right)}{a+a\cos(c+dx)}$$

$$- \frac{5A\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))}}{3d(a+a\cos(c+dx))}$$

$$+ \frac{5B\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))}}{3d(a+a\cos(c+dx))}$$

$$+ \frac{3A\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos^2(dx + \arctan(\tan(c)))}} \right)}{2d(a+a\cos(c+dx))}$$

$$- \frac{21B\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos^2(dx + \arctan(\tan(c)))}} \right)}{10d(a+a\cos(c+dx))}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output $(\cos[c/2 + (d*x)/2]^2*\sqrt{\cos[c + d*x]}*((2*(5*A - 5*B + 10*A*\cos[c] - 16*B*\cos[c])*Csc[c])/(5*d) + (4*(A - B)*\cos[d*x]*\sin[c])/(3*d) + (2*B*\cos[2*d*x]*\sin[2*c])/(5*d) + (2*\sec[c/2]*\sec[c/2 + (d*x)/2]*(A*\sin[(d*x)/2] - B*\sin[(d*x)/2]))/d + (4*(A - B)*\cos[c]*\sin[d*x])/(3*d) + (2*B*\cos[2*c]*\sin[2*d*x])/(5*d)))/(a + a*\cos[c + d*x]) - (5*A*\cos[c/2 + (d*x)/2]^2*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\sqrt{1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}]*\sqrt{-(\sqrt{1 + \text{Cot}[c]^2}*\sin[c]*\sin[d*x - \text{ArcTan}[\text{Cot}[c]]])})*\sqrt{1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}])/(3*d*(a + a*\cos[c + d*x])*\sqrt{1 + \text{Cot}[c]^2}) + (5*B*\cos[c/2 + (d*x)/2]^2*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\sqrt{1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}]*\sqrt{-(\sqrt{1 + \text{Cot}[c]^2}*\sin[c]*\sin[d*x - \text{ArcTan}[\text{Cot}[c]]])})*\sqrt{1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}])/(3*d*(a + a*\cos[c + d*x])*\sqrt{1 + \text{Cot}[c]^2}) + (3*A*\cos[c/2 + (d*x)/2]^2*Csc[c/2]*\sec[c/2]*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}, \cos[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\sin[d*x + \text{ArcTan}[\text{Tan}[c]]]*\tan[c])/(sqrt{1 - \cos[d*x + \text{ArcTan}[\text{Tan}[c]]}]*\sqrt{1 + \cos[d*x + \text{ArcTan}[\text{Tan}[c]]}]*\sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\text{Tan}[c]]}]*\sqrt{1 + \tan[c]^2})*\sqrt{1 + \tan[c]^2}) - ((\sin[d*x + \text{ArcTan}[\text{Tan}[c]]]*\tan[c])/sqrt{1 + \tan[c]^2} + (2*\cos[c]^2*\cos[d*x + \text{ArcTan}[\text{Tan}[c]]]*\sqrt{1 + \tan[c]^2})/(\cos[c]^2 + \sin[c]^2))/sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\text{Tan}[c]]})*\sqrt{1 + \tan[c]^2}))/((2*d*(a ...$

3.145.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3456, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{a \sin(c + dx + \frac{\pi}{2}) + a} dx$$

↓ 3456

$$\frac{\int \frac{1}{2} \cos^{\frac{3}{2}}(c + dx)(5a(A - B) - a(5A - 7B) \cos(c + dx)) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{d(a \cos(c + dx) + a)}$$

3.145. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \cos^{\frac{3}{2}}(c+dx)(5a(A-B) - a(5A-7B)\cos(c+dx))dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3042 \\
& \frac{\int \sin(c+dx+\frac{\pi}{2})^{3/2}(5a(A-B) - a(5A-7B)\sin(c+dx+\frac{\pi}{2}))dx}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3227 \\
& \frac{5a(A-B)\int \cos^{\frac{3}{2}}(c+dx)dx - a(5A-7B)\int \cos^{\frac{5}{2}}(c+dx)dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3042 \\
& \frac{5a(A-B)\int \sin(c+dx+\frac{\pi}{2})^{3/2}dx - a(5A-7B)\int \sin(c+dx+\frac{\pi}{2})^{5/2}dx}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3115 \\
& \frac{5a(A-B)\left(\frac{1}{3}\int \frac{1}{\sqrt{\cos(c+dx)}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d}\right) - a(5A-7B)\left(\frac{3}{5}\int \sqrt{\cos(c+dx)}dx + \frac{2\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d}\right)}{2a^2} \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3042 \\
& \frac{5a(A-B)\left(\frac{1}{3}\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d}\right) - a(5A-7B)\left(\frac{3}{5}\int \sqrt{\sin(c+dx+\frac{\pi}{2})}dx + \frac{2\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d}\right)}{2a^2} \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3119 \\
& \frac{5a(A-B)\left(\frac{1}{3}\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d}\right) - a(5A-7B)\left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d}\right)}{2a^2} \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(a\cos(c+dx)+a)}
\end{aligned}$$

3.145. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

↓ 3120

$$\frac{5a(A - B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) - a(5A - 7B) \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right)}{2a^2 \frac{(A - B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{d(a \cos(c + dx) + a)}} +$$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `((A - B)*Cos[c + d*x]^(5/2)*Sin[c + d*x]/(d*(a + a*Cos[c + d*x])) + (5*a*(A - B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)) - a*(5*A - 7*B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)))/(2*a^2)`

3.145.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.145. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.145.4 Maple [A] (verified)

Time = 6.04 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.80

method	result
default	$-\frac{\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(25AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + 45AE\right)}{15a \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}$

input `int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a), x, method=_RETURNVERBOSE)`

output `-1/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(cos(1/2*d*x+1/2*c)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(25*A*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+45*A*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))-25*B*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-63*B*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))+48*B*sin(1/2*d*x+1/2*c)^8+(-40*A-56*B)*sin(1/2*d*x+1/2*c)^6+(90*A-30*B)*sin(1/2*d*x+1/2*c)^4+(-35*A+23*B)*sin(1/2*d*x+1/2*c)^2)/a/cos(1/2*d*x+1/2*c)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

$$3.145. \int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$$

3.145.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.72

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{2(6B\cos(dx+c)^2 + 2(5A-2B)\cos(dx+c) + 25A-25B)\sqrt{\cos(dx+c)}\sin(dx+c) - 25(\sqrt{2}(iA$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `1/30*(2*(6*B*cos(d*x + c)^2 + 2*(5*A - 2*B)*cos(d*x + c) + 25*A - 25*B)*sqrt(cos(d*x + c))*sin(d*x + c) - 25*(sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 25*(sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 9*(sqrt(2)*(5*I*A - 7*I*B)*cos(d*x + c) + sqrt(2)*(5*I*A - 7*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 9*(sqrt(2)*(-5*I*A + 7*I*B)*cos(d*x + c) + sqrt(2)*(-5*I*A + 7*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a*d*cos(d*x + c) + a*d)`

3.145.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x)`

output `Timed out`

3.145.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{a\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a), x)`

3.145.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{a\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a), x)`

3.145.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)), x)`

3.146 $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

3.146.1 Optimal result 1438
 3.146.2 Mathematica [C] (warning: unable to verify) 1439
 3.146.3 Rubi [A] (verified) 1440
 3.146.4 Maple [A] (verified) 1443
 3.146.5 Fricas [C] (verification not implemented) 1444
 3.146.6 Sympy [F(-1)] 1444
 3.146.7 Maxima [F] 1445
 3.146.8 Giac [F] 1445
 3.146.9 Mupad [F(-1)] 1445

3.146.1 Optimal result

Integrand size = 33, antiderivative size = 123

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = \frac{3(A-B)E(\frac{1}{2}(c+dx)|2)}{ad} - \frac{(3A-5B) \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3ad} - \frac{(3A-5B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3ad} + \frac{(A-B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{d(a+a \cos(c+dx))}$$

output

```
3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d-1/3*(3*A-5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d+(A-B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))-1/3*(3*A-5*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d
```

3.146.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.06 (sec) , antiderivative size = 893, normalized size of antiderivative = 7.26

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \left(-\frac{2(A-B)(1+2\cos(c))\csc(c)}{d} + \frac{4B\cos(dx)\sin(c)}{3d} - \frac{2\sec\left(\frac{c}{2}\right)\sec\left(\frac{c}{2} + \frac{dx}{2}\right)(A\sin\left(\frac{dx}{2}\right) - B\sin\left(\frac{dx}{2}\right))}{d} \right)}{a+a\cos(c+dx)}$$

$$+ \frac{A\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right)\csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{d(a+a\cos(c+dx))}}$$

$$- \frac{5B\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right)\csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))}}$$

$$- \frac{3A\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right)\csc\left(\frac{c}{2}\right)\sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos^2(dx + \arctan(\tan(c)))}} \right)}{2d(a+a\cos(c+dx))}}$$

$$+ \frac{3B\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right)\csc\left(\frac{c}{2}\right)\sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos^2(dx + \arctan(\tan(c)))}} \right)}{2d(a+a\cos(c+dx))}}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output $(\cos[c/2 + (d*x)/2]^2 \sqrt{\cos[c + d*x]} * ((-2*(A - B)*(1 + 2*\cos[c])*Csc[c])/d + (4*B*\cos[d*x]*\sin[c])/(3*d) - (2*\sec[c/2]*\sec[c/2 + (d*x)/2]*(A*\sin[(d*x)/2] - B*\sin[(d*x)/2]))/d + (4*B*\cos[c]*\sin[d*x])/(3*d)))/(a + a*\cos[c + d*x]) + (A*\cos[c/2 + (d*x)/2]^2 * Csc[c/2] * HypergeometricPFQ[{1/4, 1/2}, {5/4}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2] * \sec[c/2] * \sec[d*x - \text{ArcTan}[\text{Cot}[c]]] * \sqrt{1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}] * \sqrt{-(\sqrt{1 + \text{Cot}[c]^2} * \sin[c] * \sin[d*x - \text{ArcTan}[\text{Cot}[c]]])} * \sqrt{1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]})/(d*(a + a*\cos[c + d*x]) * \sqrt{1 + \text{Cot}[c]^2}) - (5*B*\cos[c/2 + (d*x)/2]^2 * Csc[c/2] * HypergeometricPFQ[{1/4, 1/2}, {5/4}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2] * \sec[c/2] * \sec[d*x - \text{ArcTan}[\text{Cot}[c]]] * \sqrt{1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}] * \sqrt{-(\sqrt{1 + \text{Cot}[c]^2} * \sin[c] * \sin[d*x - \text{ArcTan}[\text{Cot}[c]]])} * \sqrt{1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]})/(3*d*(a + a*\cos[c + d*x]) * \sqrt{1 + \text{Cot}[c]^2}) - (3*A*\cos[c/2 + (d*x)/2]^2 * Csc[c/2] * \sec[c/2] * ((HypergeometricPFQ[-1/2, -1/4], {3/4}, \cos[d*x + \text{ArcTan}[\text{Tan}[c]]]^2] * \sin[d*x + \text{ArcTan}[\text{Tan}[c]]] * \tan[c]) / (\sqrt{1 - \cos[d*x + \text{ArcTan}[\text{Tan}[c]]}] * \sqrt{1 + \cos[d*x + \text{ArcTan}[\text{Tan}[c]]}] * \sqrt{\cos[c] * \cos[d*x + \text{ArcTan}[\text{Tan}[c]]}] * \sqrt{1 + \tan[c]^2}] * \sqrt{1 + \tan[c]^2}) - ((\sin[d*x + \text{ArcTan}[\text{Tan}[c]]] * \tan[c]) / \sqrt{1 + \tan[c]^2} + (2*\cos[c]^2 * \cos[d*x + \text{ArcTan}[\text{Tan}[c]]] * \sqrt{1 + \tan[c]^2}) / (\cos[c]^2 + \sin[c]^2)) / \sqrt{\cos[c] * \cos[d*x + \text{ArcTan}[\text{Tan}[c]]] * \sqrt{1 + \tan[c]^2}})) / (2*d*(a + a*\cos[c + d*x])) + (3*B*\cos[c/2 + (d*x)/2]^2 * Csc[c/2] * \sec[c/2] * ((HypergeometricPFQ[-1/2, -1/4], {3/4}...$

3.146.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3456, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{a \sin(c + dx + \frac{\pi}{2}) + a} dx$$

↓ 3456

$$\frac{\int \frac{1}{2} \sqrt{\cos(c + dx)} (3a(A - B) - a(3A - 5B) \cos(c + dx)) dx}{a^2} + \frac{(A - B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{d(a \cos(c + dx) + a)}$$

3.146. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \sqrt{\cos(c+dx)}(3a(A-B) - a(3A-5B)\cos(c+dx))dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3042 \\
& \frac{\int \sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B) - a(3A-5B)\sin(c+dx+\frac{\pi}{2}))dx}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3227 \\
& \frac{3a(A-B)\int \sqrt{\cos(c+dx)}dx - a(3A-5B)\int \cos^{\frac{3}{2}}(c+dx)dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3042 \\
& \frac{3a(A-B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})}dx - a(3A-5B)\int \sin(c+dx+\frac{\pi}{2})^{3/2}dx}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3115 \\
& \frac{3a(A-B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})}dx - a(3A-5B)\left(\frac{1}{3}\int \frac{1}{\sqrt{\cos(c+dx)}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d}\right)}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3042 \\
& \frac{3a(A-B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})}dx - a(3A-5B)\left(\frac{1}{3}\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d}\right)}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3119 \\
& \frac{\frac{6a(A-B)E(\frac{1}{2}(c+dx)|2)}{d} - a(3A-5B)\left(\frac{1}{3}\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d}\right)}{2a^2} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(a\cos(c+dx)+a)} \\
& \downarrow 3120
\end{aligned}$$

3.146. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

$$\frac{\frac{6a(A-B)E\left(\frac{1}{2}(c+dx)|2\right)}{d} - a(3A - 5B) \left(\frac{2\text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + \frac{2\sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right)}{2a^2} + \frac{(A - B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{d(a \cos(c + dx) + a)}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])) + ((6*a*(A - B)*EllipticE[(c + d*x)/2, 2])/d - a*(3*A - 5*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)))/(2*a^2)`

3.146.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

3.146. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

3.146.4 Maple [A] (verified)

Time = 5.02 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.13

method	result
default	$\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\left(3AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + 9AE\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)\right)}{3a \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}$

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a), x, method=_RETURNVER
BOSE)
```

```
output 1/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(cos(1/2*d*x+1
/2*c)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(3*A*E
llipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+9*A*EllipticE(cos(1/2*d*x+1/2*c), 2^(1
/2))-5*B*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-9*B*EllipticE(cos(1/2*d*x+1
/2*c), 2^(1/2)))+8*B*sin(1/2*d*x+1/2*c)^6+(6*A-18*B)*sin(1/2*d*x+1/2*c)^4+(
-3*A+7*B)*sin(1/2*d*x+1/2*c)^2)/a/cos(1/2*d*x+1/2*c)/(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2
-1)^(1/2)/d
```

3.146.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 250, normalized size of antiderivative = 2.03

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{2(2B\cos(dx+c) - 3A + 5B)\sqrt{\cos(dx+c)}\sin(dx+c) + (\sqrt{2}(3iA - 5iB)\cos(dx+c) + \sqrt{2}(3iA -$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `1/6*(2*(2*B*cos(d*x + c) - 3*A + 5*B)*sqrt(cos(d*x + c))*sin(d*x + c) + (sqrt(2)*(3*I*A - 5*I*B)*cos(d*x + c) + sqrt(2)*(3*I*A - 5*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(-3*I*A + 5*I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A + 5*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 9*(sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 9*(sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a*d*cos(d*x + c) + a*d)`

3.146.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x)`

output `Timed out`

3.146.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{a\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a), x)`

3.146.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{a\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a), x)`

3.146.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)), x)`

3.147
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$$

3.147.1 Optimal result 1446
 3.147.2 Mathematica [C] (warning: unable to verify) 1446
 3.147.3 Rubi [A] (verified) 1448
 3.147.4 Maple [A] (verified) 1450
 3.147.5 Fricas [C] (verification not implemented) 1451
 3.147.6 Sympy [F(-1)] 1452
 3.147.7 Maxima [F] 1452
 3.147.8 Giac [F] 1452
 3.147.9 Mupad [F(-1)] 1453

3.147.1 Optimal result

Integrand size = 33, antiderivative size = 85

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx = -\frac{(A-3B)E(\frac{1}{2}(c+dx)|2)}{ad} + \frac{(A-B) \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{ad} + \frac{(A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{d(a+a \cos(c+dx))}$$

output `-(A-3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d+(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d+(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))`

3.147.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.91 (sec) , antiderivative size = 862, normalized size of antiderivative = 10.14

$$\begin{aligned}
 & \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx \\
 &= \frac{\cos^2\left(\frac{c}{2}+\frac{dx}{2}\right)\sqrt{\cos(c+dx)}\left(-\frac{2(-A+B+2B\cos(c))\csc(c)}{d}+\frac{2\sec\left(\frac{c}{2}\right)\sec\left(\frac{c}{2}+\frac{dx}{2}\right)\left(A\sin\left(\frac{dx}{2}\right)-B\sin\left(\frac{dx}{2}\right)\right)}{d}\right)}{a+a\cos(c+dx)} \\
 & \quad - \frac{A\cos^2\left(\frac{c}{2}+\frac{dx}{2}\right)\csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4},\frac{1}{2};\frac{5}{4};\sin^2(dx-\arctan(\cot(c)))\right)\sec\left(\frac{c}{2}\right)\sec(dx-\arctan(\cot(c)))\sqrt{1-\sin^2(dx-\arctan(\cot(c)))}}{d(a+a\cos(c+dx))}}{d(a+a\cos(c+dx))} \\
 & \quad + \frac{B\cos^2\left(\frac{c}{2}+\frac{dx}{2}\right)\csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4},\frac{1}{2};\frac{5}{4};\sin^2(dx-\arctan(\cot(c)))\right)\sec\left(\frac{c}{2}\right)\sec(dx-\arctan(\cot(c)))\sqrt{1-\sin^2(dx-\arctan(\cot(c)))}}{d(a+a\cos(c+dx))}}{d(a+a\cos(c+dx))} \\
 & \quad + \frac{A\cos^2\left(\frac{c}{2}+\frac{dx}{2}\right)\csc\left(\frac{c}{2}\right)\sec\left(\frac{c}{2}\right)\left(\frac{{}_2F_1\left(-\frac{1}{2},-\frac{1}{4};\frac{3}{4};\cos^2(dx+\arctan(\tan(c)))\right)\sin(dx+\arctan(\tan(c)))\tan(c)}{\sqrt{1-\cos(dx+\arctan(\tan(c)))}\sqrt{1+\cos(dx+\arctan(\tan(c)))}\sqrt{\cos(c)\cos(dx+\arctan(\tan(c)))}\sqrt{1+\cos(dx+\arctan(\tan(c)))}}}\right)}{2d(a+a\cos(c+dx))}}{2d(a+a\cos(c+dx))} \\
 & \quad - \frac{3B\cos^2\left(\frac{c}{2}+\frac{dx}{2}\right)\csc\left(\frac{c}{2}\right)\sec\left(\frac{c}{2}\right)\left(\frac{{}_2F_1\left(-\frac{1}{2},-\frac{1}{4};\frac{3}{4};\cos^2(dx+\arctan(\tan(c)))\right)\sin(dx+\arctan(\tan(c)))\tan(c)}{\sqrt{1-\cos(dx+\arctan(\tan(c)))}\sqrt{1+\cos(dx+\arctan(\tan(c)))}\sqrt{\cos(c)\cos(dx+\arctan(\tan(c)))}\sqrt{1+\cos(dx+\arctan(\tan(c)))}}}\right)}{2d(a+a\cos(c+dx))}}{2d(a+a\cos(c+dx))}
 \end{aligned}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output

```
(Cos[c/2 + (d*x)/2]^2*Sqrt[Cos[c + d*x]]*((-2*(-A + B + 2*B*Cos[c])*Csc[c]
)/d + (2*Sec[c/2]*Sec[c/2 + (d*x)/2]*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/d
)/(a + a*Cos[c + d*x]) - (A*Cos[c/2 + (d*x)/2]^2*Csc[c/2]*HypergeometricPF
Q[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTa
n[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*S
in[c]*Sin[d*x - ArcTan[Cot[c]]])] *Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(d*
(a + a*Cos[c + d*x])*Sqrt[1 + Cot[c]^2]) + (B*Cos[c/2 + (d*x)/2]^2*Csc[c/2
]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/
2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sq
rt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])] *Sqrt[1 + Sin[d*x - Arc
Tan[Cot[c]]]])/(d*(a + a*Cos[c + d*x])*Sqrt[1 + Cot[c]^2]) + (A*Cos[c/2 +
(d*x)/2]^2*Csc[c/2]*Sec[c/2]*((HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[
d*x + ArcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d
*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[
d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x
+ ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTa
n[Tan[c]]]*Sqrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x
+ ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]))/(2*d*(a + a*Cos[c + d*x])) - (3*B*
Cos[c/2 + (d*x)/2]^2*Csc[c/2]*Sec[c/2]*((HypergeometricPFQ[{-1/2, -1/4}, {
3/4}, Cos[d*x + ArcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(S...
```

3.147.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3456, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a\cos(c+dx)+a} dx$$

↓ 3042

$$\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{a\sin(c+dx+\frac{\pi}{2})+a} dx$$

↓ 3456

$$\frac{\int \frac{a(A-B)-a(A-3B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx}{a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}$$

3.147. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{a(A-B) - a(A-3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow 27 \\
& \int \frac{a(A-B) - a(A-3B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow 3042 \\
& \frac{a(A-B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx - a(A-3B) \int \sqrt{\cos(c+dx)} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow 3042 \\
& \frac{a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - a(A-3B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow 3119 \\
& \frac{a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(A-3B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow 3120 \\
& \frac{\frac{2a(A-B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d} - \frac{2a(A-3B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}
\end{aligned}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x]),x]`

output `((-2*a*(A - 3*B)*EllipticE[(c + d*x)/2, 2])/d + (2*a*(A - B)*EllipticF[(c + d*x)/2, 2])/d)/(2*a^2) + ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))`

3.147.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.147.4 Maple [A] (verified)

Time = 4.19 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.87

method	result
default	$-\frac{\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \left(AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + AE\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right)}{a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}$

3.147. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+a \cos(c+dx)} dx$

input `int((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)`

output `-((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(cos(1/2*d*x+1/2*c)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))+(-2*B+2*A)*sin(1/2*d*x+1/2*c)^4+(-A+B)*sin(1/2*d*x+1/2*c)^2)/a/cos(1/2*d*x+1/2*c)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.147.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 237, normalized size of antiderivative = 2.79

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

$$= \frac{2(A-B)\sqrt{\cos(dx+c)}\sin(dx+c) + (\sqrt{2}(-iA+iB)\cos(dx+c) + \sqrt{2}(-iA+iB))\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{a+d}$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `1/2*(2*(A - B)*sqrt(cos(d*x + c))*sin(d*x + c) + (sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + (sqrt(2)*(-I*A + 3*I*B)*cos(d*x + c) + sqrt(2)*(-I*A + 3*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + (sqrt(2)*(I*A - 3*I*B)*cos(d*x + c) + sqrt(2)*(I*A - 3*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a*d*cos(d*x + c) + a*d)`

3.147.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)**(1/2)/(a+a*cos(d*x+c)),x)`

output `Timed out`

3.147.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{a\cos(dx+c)+a} dx$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(a*cos(d*x + c) + a), x)`

3.147.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{a\cos(dx+c)+a} dx$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(a*cos(d*x + c) + a), x)`

3.147.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+a\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)),x)`output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x)), x)`

3.148 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))} dx$

3.148.1 Optimal result 1454
 3.148.2 Mathematica [C] (warning: unable to verify) 1454
 3.148.3 Rubi [A] (verified) 1456
 3.148.4 Maple [A] (verified) 1458
 3.148.5 Fricas [C] (verification not implemented) 1459
 3.148.6 Sympy [F] 1460
 3.148.7 Maxima [F] 1460
 3.148.8 Giac [F] 1460
 3.148.9 Mupad [F(-1)] 1461

3.148.1 Optimal result

Integrand size = 33, antiderivative size = 83

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx = \frac{(A - B)E(\frac{1}{2}(c + dx) | 2)}{ad} + \frac{(A + B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{ad} - \frac{(A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{d(a + a \cos(c + dx))}$$

output `(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a/d+(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a/d-(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))`

3.148.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.27 (sec) , antiderivative size = 858, normalized size of antiderivative = 10.34

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx$$

$$= \frac{\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \left(-\frac{2(A-B) \csc(c)}{d} - \frac{2 \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) \left(A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right) \right)}{d} \right)}{a + a \cos(c + dx)}$$

$$- \frac{A \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{d(a + a \cos(c + dx))}}{d(a + a \cos(c + dx))}$$

$$- \frac{B \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{d(a + a \cos(c + dx))}}{d(a + a \cos(c + dx))}$$

$$- \frac{A \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{2d(a + a \cos(c + dx))}$$

$$+ \frac{B \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{2d(a + a \cos(c + dx))}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])),x]`

output $(\cos[c/2 + (d*x)/2]^2 \sqrt{\cos[c + d*x]} * ((-2*(A - B)*\csc[c])/d - (2*\sec[c/2]*\sec[c/2 + (d*x)/2]*(A*\sin[(d*x)/2] - B*\sin[(d*x)/2]))/d)/(a + a*\cos[c + d*x]) - (A*\cos[c/2 + (d*x)/2]^2*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\sin[c]*\sin[d*x - \text{ArcTan}[\text{Cot}[c]]])] * \text{Sqrt}[1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(d*(a + a*\cos[c + d*x])* \text{Sqrt}[1 + \text{Cot}[c]^2]) - (B*\cos[c/2 + (d*x)/2]^2*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\text{Sqrt}[1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]]*\text{Sqrt}[-(\text{Sqrt}[1 + \text{Cot}[c]^2]*\sin[c]*\sin[d*x - \text{ArcTan}[\text{Cot}[c]]])] * \text{Sqrt}[1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]])/(d*(a + a*\cos[c + d*x])* \text{Sqrt}[1 + \text{Cot}[c]^2]) - (A*\cos[c/2 + (d*x)/2]^2*\csc[c/2]*\sec[c/2]*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \cos[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\sin[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \cos[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[1 + \cos[d*x + \text{ArcTan}[\text{Tan}[c]]]]*\text{Sqrt}[\cos[c]*\cos[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]]*\text{Sqrt}[1 + \text{Tan}[c]^2]) - ((\sin[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/ \text{Sqrt}[1 + \text{Tan}[c]^2] + (2*\cos[c]^2*\cos[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2])/(\cos[c]^2 + \sin[c]^2))/\text{Sqrt}[\cos[c]*\cos[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Sqrt}[1 + \text{Tan}[c]^2]])/(2*d*(a + a*\cos[c + d*x])) + (B*\cos[c/2 + (d*x)/2]^2*\csc[c/2]*\sec[c/2]*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \cos[d*x + \text{ArcTan}[\text{Tan}[c]]]^2]*\sin[d*x + \text{ArcTan}[\text{Tan}[c]]]*\text{Tan}[c])/(\text{Sqrt}[1 - \cos[d*x ...$

3.148.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3457, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a \cos(c + dx) + a)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)} dx$$

$$\downarrow \text{3457}$$

$$\frac{\int \frac{a(A+B) + a(A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx}{a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx) + a)}$$

3.148. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))} dx$

$$\begin{aligned}
& \int \frac{a(A+B)+a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx \quad \downarrow \text{27} \\
& \frac{\int \frac{a(A+B)+a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \int \frac{a(A+B)+a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \\
& \frac{\int \frac{a(A+B)+a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3227} \\
& \frac{a(A+B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx + a(A-B)\int \sqrt{\cos(c+dx)} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(A+B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + a(A-B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3119} \\
& \frac{a(A+B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a(A-B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} \\
& \quad \downarrow \text{3120} \\
& \frac{\frac{2a(A+B)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d} + \frac{2a(A-B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])),x]`

output `((2*a*(A - B)*EllipticE[(c + d*x)/2, 2])/d + (2*a*(A + B)*EllipticF[(c + d*x)/2, 2])/d)/(2*a^2) - ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))`

3.148.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.148.4 Maple [A] (verified)

Time = 3.11 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.93

method	result
default	$\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\left(AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - AE\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{a \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}$

3.148.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+a \cos(c+dx))}} dx$$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)/cos(d*x+c)^(1/2),x,method=_RETURNVER
BOSE)
```

```
output ((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-cos(1/2*d*x+1/2*
c)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(A*Ellipt
icF(cos(1/2*d*x+1/2*c),2^(1/2))-A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+B*
EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/
2)))+(-2*B+2*A)*sin(1/2*d*x+1/2*c)^4+(-A+B)*sin(1/2*d*x+1/2*c)^2)/a/cos(1/
2*d*x+1/2*c)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*
d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.148.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.90

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + a \cos(c + dx))}} dx =$$

$$\frac{2(A - B)\sqrt{\cos(dx + c)}\sin(dx + c) - (\sqrt{2}(-iA - iB)\cos(dx + c) + \sqrt{2}(-iA - iB))\text{weierstrassPInverse}(\dots)}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm=
"fricas")
```

```
output -1/2*(2*(A - B)*sqrt(cos(d*x + c))*sin(d*x + c) - (sqrt(2)*(-I*A - I*B)*co
s(d*x + c) + sqrt(2)*(-I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c)
+ I*sin(d*x + c)) - (sqrt(2)*(I*A + I*B)*cos(d*x + c) + sqrt(2)*(I*A + I*
B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - (sqrt(2)*
(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassZeta(-4, 0, weie
rstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - (sqrt(2)*(-I*A +
I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassZeta(-4, 0, weierstra
ssPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a*d*cos(d*x + c) + a*d
)
```

3.148.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx$$

$$= \frac{\int \frac{A}{\cos^{\frac{3}{2}}(c+dx) + \sqrt{\cos(c+dx)}} dx + \int \frac{B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) + \sqrt{\cos(c+dx)}} dx}{a}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `(Integral(A/(cos(c + d*x)**(3/2) + sqrt(cos(c + d*x))), x) + Integral(B*cos(c + d*x)/(cos(c + d*x)**(3/2) + sqrt(cos(c + d*x))), x))/a`

3.148.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)\sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.148.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)\sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.148.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))), x)`

3.149
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))} dx$$

3.149.1 Optimal result 1462
 3.149.2 Mathematica [C] (warning: unable to verify) 1463
 3.149.3 Rubi [A] (verified) 1464
 3.149.4 Maple [A] (verified) 1467
 3.149.5 Fricas [C] (verification not implemented) 1468
 3.149.6 Sympy [F(-1)] 1468
 3.149.7 Maxima [F] 1469
 3.149.8 Giac [F] 1469
 3.149.9 Mupad [F(-1)] 1469

3.149.1 Optimal result

Integrand size = 33, antiderivative size = 119

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx = -\frac{(3A - B)E(\frac{1}{2}(c + dx) | 2)}{ad} - \frac{(A - B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{ad} + \frac{(3A - B) \sin(c + dx)}{ad\sqrt{\cos(c + dx)}} - \frac{(A - B) \sin(c + dx)}{d\sqrt{\cos(c + dx)(a + a \cos(c + dx))}}$$

```
output -(3*A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a/d-(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a/d+(3*A-B)*sin(d*x+c)/a/d/cos(d*x+c)^(1/2)-(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))/cos(d*x+c)^(1/2)
```

3.149.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.62 (sec) , antiderivative size = 894, normalized size of antiderivative = 7.51

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx$$

$$= \frac{\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \left(\frac{(2A + A \cos(c) - B \cos(c)) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \sec(c)}{d} + \frac{2 \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) \left(A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right)\right)}{d} \right) + 4 \dots}{a + a \cos(c + dx)}$$

$$+ \frac{A \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{d(a + a \cos(c + dx))}}{d(a + a \cos(c + dx))}$$

$$- \frac{B \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{d(a + a \cos(c + dx))}}{d(a + a \cos(c + dx))}$$

$$+ \frac{3A \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{2d(a + a \cos(c + dx))}$$

$$- \frac{B \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{2d(a + a \cos(c + dx))}$$

```
input Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])),x
]
```

```
output (Cos[c/2 + (d*x)/2]^2*Sqrt[Cos[c + d*x]]*(((2*A + A*Cos[c] - B*Cos[c])*Csc
[c/2]*Sec[c/2]*Sec[c])/d + (2*Sec[c/2]*Sec[c/2 + (d*x)/2]*(A*Sin[(d*x)/2]
- B*Sin[(d*x)/2]))/d + (4*A*Sec[c]*Sec[c + d*x]*Sin[d*x])/d)/(a + a*Cos[c
+ d*x]) + (A*Cos[c/2 + (d*x)/2]^2*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2},
{5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqr
t[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x
- ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(d*(a + a*Cos[c +
d*x])*Sqrt[1 + Cot[c]^2]) - (B*Cos[c/2 + (d*x)/2]^2*Csc[c/2]*Hypergeometr
icPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - A
rcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^
2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])
/(d*(a + a*Cos[c + d*x])*Sqrt[1 + Cot[c]^2]) + (3*A*Cos[c/2 + (d*x)/2]^2*C
sc[c/2]*Sec[c/2]*((HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan
[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[
Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan
[Tan[c]]]*Sqrt[1 + Tan[c]^2])*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan
[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c]]]*S
qrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan
[c]]]*Sqrt[1 + Tan[c]^2]))/(2*d*(a + a*Cos[c + d*x])) - (B*Cos[c/2 + (d*x
)/2]^2*Csc[c/2]*Sec[c/2]*((HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d...
```

3.149.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.97, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3457, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}}(a \sin(c + dx + \frac{\pi}{2}) + a)} dx$$

↓ 3457

$$\frac{\int \frac{a(3A - B) - a(A - B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)} dx}{a^2} - \frac{(A - B) \sin(c + dx)}{d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)}$$

3.149. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx$

$$\begin{aligned}
& \int \frac{a(3A-B) - a(A-B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx \quad \downarrow \quad 27 \\
& \frac{2a^2}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \\
& \int \frac{a(3A-B) - a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}} dx \quad \downarrow \quad 3042 \\
& \frac{2a^2}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \\
& \frac{a(3A-B) \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx - a(A-B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \quad \downarrow \quad 3227 \\
& \frac{a(3A-B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}} dx - a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \quad \downarrow \quad 3042 \\
& \frac{a(3A-B) \left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx \right) - a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \quad \downarrow \quad 3116 \\
& \frac{a(3A-B) \left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) - a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \quad \downarrow \quad 3042 \\
& \frac{a(3A-B) \left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) - a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \quad \downarrow \quad 3119 \\
& \frac{a(3A-B) \left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) - a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \quad \downarrow \quad 3120 \\
& \frac{a(3A-B) \left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) - a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)}
\end{aligned}$$

3.149. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))} dx$

$$\frac{a(3A - B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) - \frac{2a(A-B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{d}}{\frac{2a^2}{(A-B) \sin(c+dx)}} - \frac{2a^2}{d\sqrt{\cos(c+dx)}(a \cos(c+dx) + a)}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])),x]`

output `-(((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x]))) + ((-2*a*(A - B)*EllipticF[(c + d*x)/2, 2])/d + a*(3*A - B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/(2*a^2)`

3.149.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.149.4 Maple [A] (verified)

Time = 4.18 (sec) , antiderivative size = 319, normalized size of antiderivative = 2.68

method	result
default	$-\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a),x,method=_RETURNVER
BOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/a*(cos(1/2*d*x+
1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(A*EllipticF(cos(1/2*d*x+1
/2*c),2^(1/2))-3*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-B*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))+B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))-2*(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(3*A-B)*sin(1/2*d*x+1/2*c)^4+
(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(5*A-B)*sin(1/2*d*x+1
/2*c)^2)/cos(1/2*d*x+1/2*c)/sin(1/2*d*x+1/2*c)^3/(2*sin(1/2*d*x+1/2*c)^2-1
)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.149.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))} dx$$

3.149.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 290, normalized size of antiderivative = 2.44

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx$$

$$= \frac{2((3A - B) \cos(dx + c) + 2A) \sqrt{\cos(dx + c)} \sin(dx + c) + (\sqrt{2}(iA - iB) \cos(dx + c))^2 + \sqrt{2}(iA - iB) \cos(dx + c)}{a^2 \cos^2(dx + c) + a d \cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output `1/2*(2*((3*A - B)*cos(d*x + c) + 2*A)*sqrt(cos(d*x + c))*sin(d*x + c) + (sqrt(2)*(I*A - I*B)*cos(d*x + c)^2 + sqrt(2)*(I*A - I*B)*cos(d*x + c))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(-I*A + I*B)*cos(d*x + c)^2 + sqrt(2)*(-I*A + I*B)*cos(d*x + c))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + (sqrt(2)*(-3*I*A + I*B)*cos(d*x + c)^2 + sqrt(2)*(-3*I*A + I*B)*cos(d*x + c))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + (sqrt(2)*(3*I*A - I*B)*cos(d*x + c)^2 + sqrt(2)*(3*I*A - I*B)*cos(d*x + c))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a*d*cos(d*x + c)^2 + a*d*cos(d*x + c))`

3.149.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c)),x)`

output `Timed out`

3.149.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.149.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.149.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{3}{2}}(a + a \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))), x)`

3.150
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))} dx$$

3.150.1 Optimal result 1470
 3.150.2 Mathematica [C] (warning: unable to verify) 1471
 3.150.3 Rubi [A] (verified) 1472
 3.150.4 Maple [B] (verified) 1475
 3.150.5 Fricas [C] (verification not implemented) 1476
 3.150.6 Sympy [F(-1)] 1477
 3.150.7 Maxima [F] 1477
 3.150.8 Giac [F] 1477
 3.150.9 Mupad [F(-1)] 1478

3.150.1 Optimal result

Integrand size = 33, antiderivative size = 153

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx = \frac{3(A - B)E(\frac{1}{2}(c + dx) | 2)}{ad} + \frac{(5A - 3B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3ad} + \frac{(5A - 3B) \sin(c + dx)}{3ad \cos^{\frac{3}{2}}(c + dx)} - \frac{3(A - B) \sin(c + dx)}{ad \sqrt{\cos(c + dx)}} - \frac{(A - B) \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))}$$

output `3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a/d+1/3*(5*A-3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a/d+1/3*(5*A-3*B)*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)-(A-B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))-3*(A-B)*sin(d*x+c)/a/d/cos(d*x+c)^(1/2)`

3.150.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.03 (sec) , antiderivative size = 931, normalized size of antiderivative = 6.08

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx$$

$$= \frac{\cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \left(-\frac{(A-B)(2+\cos(c)) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \sec(c)}{d} - \frac{2 \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) \left(A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right)\right)}{d} + \frac{4A \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right)}{d} \right)}{a + a \cos(c + dx)}$$

$$- \frac{5A \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))}}{3d(a + a \cos(c + dx))}}$$

$$+ \frac{B \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{d(a + a \cos(c + dx))}}{d(a + a \cos(c + dx))}}$$

$$- \frac{3A \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \sin^2(dx + \arctan(\tan(c)))}} \right)}{2d(a + a \cos(c + dx))}}$$

$$+ \frac{3B \cos^2\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \sin^2(dx + \arctan(\tan(c)))}} \right)}{2d(a + a \cos(c + dx))}}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])),x]`

output

```
(Cos[c/2 + (d*x)/2]^2*Sqrt[Cos[c + d*x]]*(-(((A - B)*(2 + Cos[c])*Csc[c/2]
*Sec[c/2]*Sec[c])/d) - (2*Sec[c/2]*Sec[c/2 + (d*x)/2]*(A*Sin[(d*x)/2] - B*
Sin[(d*x)/2]))/d + (4*A*Sec[c]*Sec[c + d*x]^2*Sin[d*x])/(3*d) + (4*Sec[c]*
Sec[c + d*x]*(A*Sin[c] - 3*A*Sin[d*x] + 3*B*Sin[d*x]))/(3*d)))/(a + a*Cos[
c + d*x]) - (5*A*Cos[c/2 + (d*x)/2]^2*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2
}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*
Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d
*x - ArcTan[Cot[c]])])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3*d*(a + a*Co
s[c + d*x])*Sqrt[1 + Cot[c]^2]) + (B*Cos[c/2 + (d*x)/2]^2*Csc[c/2]*Hyperge
ometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*
x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Co
t[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c
]]]])/(d*(a + a*Cos[c + d*x])*Sqrt[1 + Cot[c]^2]) - (3*A*Cos[c/2 + (d*x)/2
]^2*Csc[c/2]*Sec[c/2]*((HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + A
rcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + Ar
cTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + A
rcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2])*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTa
n[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c
]]]*Sqrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + ArcTa
n[Tan[c]]]*Sqrt[1 + Tan[c]^2]))/(2*d*(a + a*Cos[c + d*x])) + (3*B*Cos[...
```

3.150.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.94, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3457, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}}(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)} dx$$

↓ 3457

$$\frac{\int \frac{a(5A - 3B) - 3a(A - B) \cos(c + dx)}{2 \cos^{\frac{5}{2}}(c + dx)} dx}{a^2} - \frac{(A - B) \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)}$$

3.150. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx$

$$\begin{aligned}
& \int \frac{a(5A-3B)-3a(A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)} dx \quad \downarrow \text{27} \\
& \frac{2a^2}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} (A-B) \sin(c+dx) \\
& \int \frac{a(5A-3B)-3a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}}} dx \quad \downarrow \text{3042} \\
& \frac{2a^2}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} (A-B) \sin(c+dx) \\
& \frac{a(5A-3B) \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)} dx - 3a(A-B) \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx}{2a^2} - \frac{(A-B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} \quad \downarrow \text{3227} \\
& \frac{a(5A-3B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}}} dx - 3a(A-B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}} dx}{2a^2} - \frac{(A-B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} \quad \downarrow \text{3042} \\
& \frac{a(5A-3B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} - 3a(A-B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx \right)}{2a^2} - \frac{(A-B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} \quad \downarrow \text{3116} \\
& \frac{a(5A-3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 3a(A-B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right)}{2a^2} - \frac{(A-B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} \quad \downarrow \text{3042} \\
& \frac{a(5A-3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 3a(A-B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right)}{2a^2} - \frac{(A-B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)} \quad \downarrow \text{3119} \\
& \frac{a(5A-3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 3a(A-B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right)}{2a^2} - \frac{(A-B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)}
\end{aligned}$$

3.150. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))} dx$

$$\frac{a(5A - 3B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 3a(A - B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right)}{\frac{2a^2}{(A - B) \sin(c + dx)} d \cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])),x]`

output `-(((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x]))) + (a*(5*A - 3*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))) - 3*a*(A - B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/(2*a^2)`

3.150.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d)), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.150.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(195) = 390.

Time = 5.49 (sec) , antiderivative size = 466, normalized size of antiderivative = 3.05

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2})}} \left(\frac{(A-B)\left(\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right) (F(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2}) - E(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2}))}{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2})}} \right)$

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)`

$$3.150. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))} dx$$

output
$$\begin{aligned} & -(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/a*((A-B)*(\cos(1/2*d*x+1/2*c)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ &)*(\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})) \\ &)-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)/\cos(1/2*d*x+1/2*c)/(-2* \\ & \sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*A*(-1/6*\cos(1/2*d*x+1/2 \\ & *c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(\cos(1/2*d*x+1/2* \\ & c)^2-1/2)^2+1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)} \\ &)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c), \\ & 2^{(1/2)}))+(2*B-2*A)/\sin(1/2*d*x+1/2*c)^2/(2*\sin(1/2*d*x+1/2*c)^2-1)*(-2*\sin(1/2*d*x+1/2*c)^4 \\ & +\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & *(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c) \\ &)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.150.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 320, normalized size of antiderivative = 2.09

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx = \frac{2(9(A - B) \cos(dx + c)^2 + 2(2A - 3B) \cos(dx + c) - 2A) \sqrt{\cos(dx + c)} \sin(dx + c) - (\sqrt{2}(-5iA$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & -1/6*(2*(9*(A - B)*\cos(d*x + c)^2 + 2*(2*A - 3*B)*\cos(d*x + c) - 2*A)*\text{sqrt} \\ & (\cos(d*x + c))*\sin(d*x + c) - (\text{sqrt}(2)*(-5*I*A + 3*I*B)*\cos(d*x + c)^3 + \text{sqrt}(2)*(-5*I*A + 3*I*B)*\cos(d*x + c)^2)*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) - (\text{sqrt}(2)*(5*I*A - 3*I*B)*\cos(d*x + c)^3 + \text{sqrt}(2)*(5*I*A - 3*I*B)*\cos(d*x + c)^2)*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + 9*(\text{sqrt}(2)*(-I*A + I*B)*\cos(d*x + c)^3 + \text{sqrt}(2)*(-I*A + I*B)*\cos(d*x + c)^2)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + 9*(\text{sqrt}(2)*(I*A - I*B)*\cos(d*x + c)^3 + \text{sqrt}(2)*(I*A - I*B)*\cos(d*x + c)^2)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/(a*d*\cos(d*x + c)^3 + a*d*\cos(d*x + c)^2) \end{aligned}$$

3.150.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))} dx$$

3.150.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c)),x)`

output `Timed out`

3.150.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*cos(d*x + c)^(5/2)),x)`

3.150.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*cos(d*x + c)^(5/2)),x)`

3.150.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + a \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))), x)`

3.151
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.151.1 Optimal result 1479
 3.151.2 Mathematica [C] (warning: unable to verify) 1480
 3.151.3 Rubi [A] (verified) 1480
 3.151.4 Maple [A] (verified) 1484
 3.151.5 Fricas [C] (verification not implemented) 1485
 3.151.6 Sympy [F(-1)] 1485
 3.151.7 Maxima [F] 1486
 3.151.8 Giac [F] 1486
 3.151.9 Mupad [F(-1)] 1486

3.151.1 Optimal result

Integrand size = 33, antiderivative size = 203

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = -\frac{7(5A-8B)E(\frac{1}{2}(c+dx)|2)}{5a^2d} + \frac{5(2A-3B) \operatorname{EllipticF}(\frac{1}{2}(c+dx),2)}{3a^2d} + \frac{5(2A-3B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3a^2d} - \frac{7(5A-8B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{15a^2d} + \frac{(2A-3B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{a^2d(1+\cos(c+dx))} + \frac{(A-B) \cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{3d(a+a \cos(c+dx))^2}$$

output `-7/5*(5*A-8*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d+5/3*(2*A-3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d-7/15*(5*A-8*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/a^2/d+(2*A-3*B)*cos(d*x+c)^(5/2)*sin(d*x+c)/a^2/d/(1+cos(d*x+c))+1/3*(A-B)*cos(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2+5/3*(2*A-3*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d`

3.151.
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.151.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.80 (sec) , antiderivative size = 1024, normalized size of antiderivative = 5.04

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \text{Too large to display}$$

input `Integrate[(Cos[c + d*x]^(7/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2, x]`

output `(-20*A*Cos[c/2 + (d*x)/2]^4*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3*d*(a + a*Cos[c + d*x])^2*Sqrt[1 + Cot[c]^2]) + (10*B*Cos[c/2 + (d*x)/2]^4*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(d*(a + a*Cos[c + d*x])^2*Sqrt[1 + Cot[c]^2]) + (Cos[c/2 + (d*x)/2]^4*Sqrt[Cos[c + d*x]]*((4*(15*A - 20*B + 20*A*Cos[c] - 36*B*Cos[c])*Csc[c])/(5*d) + (8*(A - 2*B)*Cos[d*x]*Sin[c])/(3*d) + (4*B*Cos[2*d*x]*Sin[2*c])/(5*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(3*A*Sin[(d*x)/2] - 4*B*Sin[(d*x)/2]))/d - (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(3*d) + (8*(A - 2*B)*Cos[c]*Sin[d*x])/(3*d) + (4*B*Cos[2*c]*Sin[2*d*x])/(5*d) - (2*(A - B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(3*d)))/(a + a*Cos[c + d*x])^2 + (7*A*Cos[c/2 + (d*x)/2]^4*Csc[c/2]*Sec[c/2]*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos...`

3.151.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.99, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3456, 27, 3042, 3456, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.151. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^{7/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^2} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^{\frac{5}{2}}(c+dx)(7a(A-B)-a(5A-11B)\cos(c+dx))}{2(\cos(c+dx)a+a)} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cos^{\frac{5}{2}}(c+dx)(7a(A-B)-a(5A-11B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}(7a(A-B)-a(5A-11B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \cos^{\frac{3}{2}}(c+dx)(15a^2(2A-3B)-7a^2(5A-8B)\cos(c+dx)) dx}{a^2} + \frac{6(2A-3B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sin(c+dx+\frac{\pi}{2})^{3/2}(15a^2(2A-3B)-7a^2(5A-8B)\sin(c+dx+\frac{\pi}{2})) dx}{a^2} + \frac{6(2A-3B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3227} \\
& \frac{15a^2(2A-3B)\int \cos^{\frac{3}{2}}(c+dx) dx - 7a^2(5A-8B)\int \cos^{\frac{5}{2}}(c+dx) dx}{a^2} + \frac{6(2A-3B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.151. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\frac{15a^2(2A-3B) \int \sin(c+dx+\frac{\pi}{2})^{3/2} dx - 7a^2(5A-8B) \int \sin(c+dx+\frac{\pi}{2})^{5/2} dx}{a^2} + \frac{6(2A-3B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{d(\cos(c+dx)+1)} +$$

$$\frac{6a^2}{3d(a \cos(c+dx) + a)^2} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3115

$$\frac{15a^2(2A-3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) - 7a^2(5A-8B) \left(\frac{3}{5} \int \sqrt{\cos(c+dx)} dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right)}{a^2} + \frac{6(2A-3B) \sin(c+dx)}{d(\cos(c+dx)+1)}$$

$$\frac{6a^2}{3d(a \cos(c+dx) + a)^2} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3042

$$\frac{15a^2(2A-3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) - 7a^2(5A-8B) \left(\frac{3}{5} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right)}{a^2} + \frac{6(2A-3B) \sin(c+dx)}{d(\cos(c+dx)+1)}$$

$$\frac{6a^2}{3d(a \cos(c+dx) + a)^2} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3119

$$\frac{15a^2(2A-3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) - 7a^2(5A-8B) \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right)}{a^2} + \frac{6(2A-3B) \sin(c+dx)}{d(\cos(c+dx)+1)}$$

$$\frac{6a^2}{3d(a \cos(c+dx) + a)^2} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3120

$$\frac{15a^2(2A-3B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) - 7a^2(5A-8B) \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right)}{a^2} + \frac{6(2A-3B) \sin(c+dx)}{d(\cos(c+dx)+1)}$$

$$\frac{6a^2}{3d(a \cos(c+dx) + a)^2} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

input `Int[(Cos[c + d*x]^(7/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

3.151. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

```
output ((A - B)*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (
(6*(2*A - 3*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(d*(1 + Cos[c + d*x])) + (
15*a^2*(2*A - 3*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c +
d*x]]*Sin[c + d*x])/(3*d)) - 7*a^2*(5*A - 8*B)*((6*EllipticE[(c + d*x)/2,
2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)))/a^2/(6*a^2)
```

3.151.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3115 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*SIN[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*SIN
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int
[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Ssin[e + f*x])^(m
+ 1)*(c + d*Ssin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

3.151.4 Maple [A] (verified)

Time = 8.28 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.29

method	result
default	$-\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(96B\left(\cos^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 80A\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 352B\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 60A\left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 352B\left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 80A\left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 352B\left(\cos^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 60A\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 352B\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 60A\right)$

```
input int(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)
```

```
output -1/30*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(96*B*cos(1/
2*d*x+1/2*c)^10+80*A*cos(1/2*d*x+1/2*c)^8-352*B*cos(1/2*d*x+1/2*c)^8+60*A*
cos(1/2*d*x+1/2*c)^6+100*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/
2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3
+210*A*cos(1/2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1
/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+120*B*cos(1/2*d*x+1
/2*c)^6-150*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/
2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3-336*B*cos(1/
2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1
/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-240*A*cos(1/2*d*x+1/2*c)^4+266*B
*cos(1/2*d*x+1/2*c)^4+105*A*cos(1/2*d*x+1/2*c)^2-135*B*cos(1/2*d*x+1/2*c)^
2-5*A+5*B)/a^2/cos(1/2*d*x+1/2*c)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1
/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

$$3.151. \quad \int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.151.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 383, normalized size of antiderivative = 1.89

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{2(6B\cos(dx+c)^3 + 2(5A-4B)\cos(dx+c)^2 + (65A-94B)\cos(dx+c) + 50A-75B)\sqrt{\cos(dx+c)}}{a^2d\cos(dx+c)^2 + 2a^2d\cos(dx+c) + a^2d}$$

```
input integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm
m="fricas")
```

```
output 1/30*(2*(6*B*cos(d*x + c)^3 + 2*(5*A - 4*B)*cos(d*x + c)^2 + (65*A - 94*B)
*cos(d*x + c) + 50*A - 75*B)*sqrt(cos(d*x + c))*sin(d*x + c) - 25*(sqrt(2)
*(2*I*A - 3*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(2*I*A - 3*I*B)*cos(d*x + c) +
sqrt(2)*(2*I*A - 3*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(
d*x + c)) - 25*(sqrt(2)*(-2*I*A + 3*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(-2*I*
A + 3*I*B)*cos(d*x + c) + sqrt(2)*(-2*I*A + 3*I*B))*weierstrassPInverse(-4
, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*(sqrt(2)*(5*I*A - 8*I*B)*cos(d*x
+ c)^2 + 2*sqrt(2)*(5*I*A - 8*I*B)*cos(d*x + c) + sqrt(2)*(5*I*A - 8*I*B))
*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*
x + c))) - 21*(sqrt(2)*(-5*I*A + 8*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(-5*I*A
+ 8*I*B)*cos(d*x + c) + sqrt(2)*(-5*I*A + 8*I*B))*weierstrassZeta(-4, 0,
weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a^2*d*cos(d*x
+ c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)
```

3.151.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)
```

```
output Timed out
```

3.151. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

3.151.7 Maxima [F]

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{7}{2}}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^2, x)`

3.151.8 Giac [F]

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{7}{2}}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^2, x)`

3.151.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{7/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(7/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2,x)`

output `int((cos(c + d*x)^(7/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2, x)`

3.151. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

3.152
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.152.1 Optimal result 1487
 3.152.2 Mathematica [C] (warning: unable to verify) 1488
 3.152.3 Rubi [A] (verified) 1489
 3.152.4 Maple [B] (verified) 1493
 3.152.5 Fricas [C] (verification not implemented) 1493
 3.152.6 Sympy [F(-1)] 1494
 3.152.7 Maxima [F] 1494
 3.152.8 Giac [F] 1495
 3.152.9 Mupad [F(-1)] 1495

3.152.1 Optimal result

Integrand size = 33, antiderivative size = 166

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = \frac{(4A-7B)E(\frac{1}{2}(c+dx)|2)}{a^2d} - \frac{5(A-2B) \operatorname{EllipticF}(\frac{1}{2}(c+dx),2)}{3a^2d} - \frac{5(A-2B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3a^2d} + \frac{(4A-7B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{3a^2d(1+\cos(c+dx))} + \frac{(A-B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{3d(a+a \cos(c+dx))^2}$$

output

```
(4*A-7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d-5/3*(A-2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d+1/3*(4*A-7*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/a^2/d/(1+cos(d*x+c))+1/3*(A-B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2-5/3*(A-2*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d
```

3.152.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.53 (sec) , antiderivative size = 980, normalized size of antiderivative = 5.90

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{10A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))^2} - \frac{20B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))^2} + \frac{\cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \left(-\frac{4(2A-3B+2A\cos(c)-4B\cos(c))\csc(c)}{d} + \frac{8B\cos(dx)\sin(c)}{3d} - \frac{4\sec\left(\frac{c}{2}\right)\sec\left(\frac{c}{2} + \frac{dx}{2}\right)(2A\sin\left(\frac{dx}{2}\right) - 2B\cos\left(\frac{dx}{2}\right))}{d}\right)}{(a+a\cos(c+dx))^2} - \frac{4A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos^2(dx + \arctan(\tan(c)))}}\right)}{d(a+a\cos(c+dx))^2} - \frac{7B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos^2(dx + \arctan(\tan(c)))}}\right)}{d(a+a\cos(c+dx))^2}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2, x]`

output $(10*A*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\sqrt{1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}]*\sqrt{-(\sqrt{1 + \text{Cot}[c]^2}*\sin[c]*\sin[d*x - \text{ArcTan}[\text{Cot}[c]])})*\sqrt{1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}]/(3*d*(a + a*\cos[c + d*x])^2*\sqrt{1 + \text{Cot}[c]^2}) - (20*B*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\sqrt{1 - \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}]*\sqrt{-(\sqrt{1 + \text{Cot}[c]^2}*\sin[c]*\sin[d*x - \text{ArcTan}[\text{Cot}[c]])})*\sqrt{1 + \sin[d*x - \text{ArcTan}[\text{Cot}[c]]}]/(3*d*(a + a*\cos[c + d*x])^2*\sqrt{1 + \text{Cot}[c]^2}) + (\cos[c/2 + (d*x)/2]^4*\sqrt{\cos[c + d*x]}*((-4*(2*A - 3*B + 2*A*\cos[c] - 4*B*\cos[c])*Csc[c])/d + (8*B*\cos[d*x]*\sin[c])/(3*d) - (4*\sec[c/2]*\sec[c/2 + (d*x)/2]*(2*A*\sin[(d*x)/2] - 3*B*\sin[(d*x)/2]))/d + (2*\sec[c/2]*\sec[c/2 + (d*x)/2]^3*(A*\sin[(d*x)/2] - B*\sin[(d*x)/2]))/(3*d) + (8*B*\cos[c]*\sin[d*x])/(3*d) + (2*(A - B)*\sec[c/2 + (d*x)/2]^2*\tan[c/2])/(3*d)))/(a + a*\cos[c + d*x])^2 - (4*A*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\sec[c/2]*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \cos[d*x + \text{ArcTan}[\tan[c]]]^2]*\sin[d*x + \text{ArcTan}[\tan[c]]]*\tan[c])/(\sqrt{1 - \cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{1 + \cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{1 + \tan[c]^2})*\sqrt{1 + \tan[c]^2}) - ((\sin[d*x + \text{ArcTan}[\tan[c]]]*\tan[c])/sqrt{1 + \tan[c]^2} + (2*\cos[c]^2*\cos[d*x + \text{ArcTan}[\tan[c]]]*sqrt{1 + \tan[c]^2}))/(\cos[c]^2 + \sin[c]^2))/sqrt{\cos[c]*\cos...$

3.152.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^2} dx$$

$$\downarrow \text{3456}$$

$$\frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)-3a(A-3B)\cos(c+dx))}{2(\cos(c+dx)a+a)} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

3.152. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)-3a(A-3B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(5a(A-B)-3a(A-3B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3456 \\
& \frac{\int 3\sqrt{\cos(c+dx)}(a^2(4A-7B)-5a^2(A-2B)\cos(c+dx)) dx}{a^2} + \frac{2(4A-7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 27 \\
& \frac{3\int \sqrt{\cos(c+dx)}(a^2(4A-7B)-5a^2(A-2B)\cos(c+dx)) dx}{a^2} + \frac{2(4A-7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3042 \\
& \frac{3\int \sqrt{\sin(c+dx+\frac{\pi}{2})}(a^2(4A-7B)-5a^2(A-2B)\sin(c+dx+\frac{\pi}{2})) dx}{a^2} + \frac{2(4A-7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3227 \\
& \frac{3(a^2(4A-7B)\int \sqrt{\cos(c+dx)} dx - 5a^2(A-2B)\int \cos^{\frac{3}{2}}(c+dx) dx)}{a^2} + \frac{2(4A-7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3042 \\
& \frac{3(a^2(4A-7B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^2(A-2B)\int \sin(c+dx+\frac{\pi}{2})^{3/2} dx)}{a^2} + \frac{2(4A-7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} + \\
& \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3115
\end{aligned}$$

3.152. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\frac{3 \left(a^2(4A-7B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^2(A-2B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right)}{a^2} + \frac{2(4A-7B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} +$$

$$\frac{6a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \frac{1}{3d(a \cos(c+dx) + a)^2}$$

↓ 3042

$$\frac{3 \left(a^2(4A-7B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^2(A-2B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right)}{a^2} + \frac{2(4A-7B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} +$$

$$\frac{6a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \frac{1}{3d(a \cos(c+dx) + a)^2}$$

↓ 3119

$$\frac{3 \left(\frac{2a^2(4A-7B)E(\frac{1}{2}(c+dx)|2)}{d} - 5a^2(A-2B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right)}{a^2} + \frac{2(4A-7B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} +$$

$$\frac{6a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \frac{1}{3d(a \cos(c+dx) + a)^2}$$

↓ 3120

$$\frac{3 \left(\frac{2a^2(4A-7B)E(\frac{1}{2}(c+dx)|2)}{d} - 5a^2(A-2B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right)}{a^2} + \frac{2(4A-7B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(\cos(c+dx)+1)} +$$

$$\frac{6a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \frac{1}{3d(a \cos(c+dx) + a)^2}$$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Cos[c + d*x]^(5/2)*Sin[c + d*x]/(3*d*(a + a*Cos[c + d*x])^2) + (2*(4*A - 7*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x]/(d*(1 + Cos[c + d*x])) + (3*((2*a^2*(4*A - 7*B)*EllipticE[(c + d*x)/2, 2])/d - 5*a^2*(A - 2*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/a^2)/(6*a^2)`

3.152. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$

3.152.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.152.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(204) = 408$.

Time = 7.71 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.62

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(-16B\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+24A\left(\cos^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+10A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)$

input `int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)`

output
$$\frac{1}{6}\left(\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2-1\right)\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)^{\frac{1}{2}}\left(-16B\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^8+24A\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^6+10A\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2\right)^{\frac{1}{2}}\left(-2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2+1\right)^{\frac{1}{2}}\text{EllipticF}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^3+24A\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^3\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2\right)^{\frac{1}{2}}\left(-2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2+1\right)^{\frac{1}{2}}\text{EllipticE}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)-12B\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^6-20B\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2\right)^{\frac{1}{2}}\left(-2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2+1\right)^{\frac{1}{2}}\text{EllipticF}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^3-42B\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^3\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2\right)^{\frac{1}{2}}\left(-2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2+1\right)^{\frac{1}{2}}\text{EllipticE}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)-38A\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^4+48B\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^4+15A\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-21B\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-A+B}{a^2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^3\left(-2\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^4+\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)^{\frac{1}{2}}\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2-1\right)^{\frac{1}{2}}/d$$

3.152.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 367, normalized size of antiderivative = 2.21

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{2(2B\cos(dx+c)^2 - (6A - 13B)\cos(dx+c) - 5A + 10B)\sqrt{\cos(dx+c)}\sin(dx+c) - 5(\sqrt{2}(-iA +$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm
m="fracas")`

3.152.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

output $1/6*(2*(2*B*\cos(d*x + c)^2 - (6*A - 13*B)*\cos(d*x + c) - 5*A + 10*B)*\sqrt{\cos(d*x + c)}*\sin(d*x + c) - 5*(\sqrt{2})*(-I*A + 2*I*B)*\cos(d*x + c)^2 + 2*\sqrt{2}*(-I*A + 2*I*B)*\cos(d*x + c) + \sqrt{2}*(-I*A + 2*I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) - 5*(\sqrt{2}*(I*A - 2*I*B)*\cos(d*x + c)^2 + 2*\sqrt{2}*(I*A - 2*I*B)*\cos(d*x + c) + \sqrt{2}*(I*A - 2*I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*(\sqrt{2}*(-4*I*A + 7*I*B)*\cos(d*x + c)^2 + 2*\sqrt{2}*(-4*I*A + 7*I*B)*\cos(d*x + c) + \sqrt{2}*(-4*I*A + 7*I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - 3*(\sqrt{2}*(4*I*A - 7*I*B)*\cos(d*x + c)^2 + 2*\sqrt{2}*(4*I*A - 7*I*B)*\cos(d*x + c) + \sqrt{2}*(4*I*A - 7*I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/(a^2*d*\cos(d*x + c)^2 + 2*a^2*d*\cos(d*x + c) + a^2*d)$

3.152.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)`

output Timed out

3.152.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{5}{2}}}{(a \cos(dx + c) + a)^2} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^2, x)`

3.152.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^2, x)`

3.152.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2,x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2, x)`

3.153
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.153.1 Optimal result 1496
 3.153.2 Mathematica [C] (warning: unable to verify) 1497
 3.153.3 Rubi [A] (verified) 1498
 3.153.4 Maple [B] (verified) 1501
 3.153.5 Fricas [C] (verification not implemented) 1502
 3.153.6 Sympy [F(-1)] 1502
 3.153.7 Maxima [F] 1503
 3.153.8 Giac [F] 1503
 3.153.9 Mupad [F(-1)] 1503

3.153.1 Optimal result

Integrand size = 33, antiderivative size = 136

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = -\frac{(A-4B)E(\frac{1}{2}(c+dx)|2)}{a^2d} + \frac{(2A-5B) \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3a^2d} + \frac{(2A-5B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3a^2d(1+\cos(c+dx))} + \frac{(A-B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{3d(a+a \cos(c+dx))^2}$$

```
output -(A-4*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2
*d*x+1/2*c), 2^(1/2))/a^2/d+1/3*(2*A-5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(
1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a^2/d+1/3*(A-B)*cos(d
*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2+1/3*(2*A-5*B)*sin(d*x+c)*cos(d
*x+c)^(1/2)/a^2/d/(1+cos(d*x+c))
```

3.153.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.26 (sec) , antiderivative size = 945, normalized size of antiderivative = 6.95

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx =$$

$$\frac{4A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \cos(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))^2} +$$

$$\frac{10B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \cos(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))^2} +$$

$$\frac{\cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \left(-\frac{4(-A+2B+2B\cos(c))\csc(c)}{d} + \frac{4\sec\left(\frac{c}{2}\right)\sec\left(\frac{c}{2} + \frac{dx}{2}\right)\left(A\sin\left(\frac{dx}{2}\right) - 2B\sin\left(\frac{dx}{2}\right)\right)}{d} - \frac{2\sec\left(\frac{c}{2}\right)\sec\left(\frac{c}{2} + \frac{dx}{2}\right)}{d} \right)}{(a + a \cos(c + dx))^2} +$$

$$\frac{A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{d(a + a \cos(c + dx))^2} +$$

$$\frac{4B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{d(a + a \cos(c + dx))^2}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2, x]`

output $(-4*A*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\sqrt{1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]}]*\sqrt{-(\sqrt{1 + \text{Cot}[c]^2}*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]])})*\sqrt{1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]}]/(3*d*(a + a*\cos[c + d*x])^2*\sqrt{1 + \text{Cot}[c]^2}) + (10*B*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\text{Cot}[c]]]*\sqrt{1 - \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]}]*\sqrt{-(\sqrt{1 + \text{Cot}[c]^2}*\text{Sin}[c]*\text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]])})*\sqrt{1 + \text{Sin}[d*x - \text{ArcTan}[\text{Cot}[c]]}]/(3*d*(a + a*\cos[c + d*x])^2*\sqrt{1 + \text{Cot}[c]^2}) + (\cos[c/2 + (d*x)/2]^4*\sqrt{\cos[c + d*x]}*((-4*(-A + 2*B + 2*B*\cos[c])*\csc[c])/d + (4*\sec[c/2]*\sec[c/2 + (d*x)/2]*(A*\sin[(d*x)/2] - 2*B*\sin[(d*x)/2]))/d - (2*\sec[c/2]*\sec[c/2 + (d*x)/2]^3*(A*\sin[(d*x)/2] - B*\sin[(d*x)/2]))/(3*d) - (2*(A - B)*\sec[c/2 + (d*x)/2]^2*\tan[c/2])/(3*d)))/(a + a*\cos[c + d*x])^2 + (A*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\sec[c/2]*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \cos[d*x + \text{ArcTan}[\tan[c]]]^2]*\sin[d*x + \text{ArcTan}[\tan[c]]]*\tan[c])/(\sqrt{1 - \cos[d*x + \text{ArcTan}[\tan[c]]})*\sqrt{1 + \cos[d*x + \text{ArcTan}[\tan[c]]})*\sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\tan[c]]})*\sqrt{1 + \tan[c]^2})*\sqrt{1 + \tan[c]^2}) - ((\sin[d*x + \text{ArcTan}[\tan[c]]]*\tan[c])/(\sqrt{1 + \tan[c]^2}) + (2*\cos[c]^2*\cos[d*x + \text{ArcTan}[\tan[c]]]*\sqrt{1 + \tan[c]^2}))/(\cos[c]^2 + \sin[c]^2))/(\sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\tan[c]]})*\sqrt{1 + \tan[c]^2}))/((d*(a + a*\cos[c + d*x])^2) - (4*B...$

3.153.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3456, 27, 3042, 3456, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^2} dx$$

$$\downarrow \text{3456}$$

$$\frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)-a(A-7B)\cos(c+dx))}{2(\cos(c+dx)a+a)} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2}$$

3.153. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)-a(A-7B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)-a(A-7B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3456 \\
& \frac{\int \frac{a^2(2A-5B)-3a^2(A-4B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx}{6a^2} + \frac{2(2A-5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{a^2(2A-5B)-3a^2(A-4B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{6a^2} + \frac{2(2A-5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3227 \\
& \frac{a^2(2A-5B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^2(A-4B)\int \sqrt{\cos(c+dx)} dx}{6a^2} + \frac{2(2A-5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3042 \\
& \frac{a^2(2A-5B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^2(A-4B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{6a^2} + \frac{2(2A-5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3119 \\
& \frac{a^2(2A-5B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6a^2(A-4B)E(\frac{1}{2}(c+dx)|2)}{d}}{6a^2} + \frac{2(2A-5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} \\
& \downarrow 3120
\end{aligned}$$

3.153. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\frac{\frac{2a^2(2A-5B)\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} - \frac{6a^2(A-4B)E\left(\frac{1}{2}(c+dx)\right)}{d}}{a^2} + \frac{2(2A-5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \frac{6a^2}{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)} \frac{1}{3d(a\cos(c+dx)+a)^2}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((-6*a^2*(A - 4*B)*EllipticE[(c + d*x)/2, 2])/d + (2*a^2*(2*A - 5*B)*EllipticF[(c + d*x)/2, 2])/d)/a^2 + (2*(2*A - 5*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(1 + Cos[c + d*x]))/(6*a^2)`

3.153.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

3.153.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 420 vs. 2(178) = 356.

Time = 5.20 (sec) , antiderivative size = 421, normalized size of antiderivative = 3.10

method	result
default	$-\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(12A\left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 4A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right)$

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)
```

```
output -1/6*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*A*cos(1/2
*d*x+1/2*c)^6+4*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)
^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3+6*A*cos(
1/2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-24*B*cos(1/2*d*x+1/2*c)^6-10*B
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(
cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3-24*B*cos(1/2*d*x+1/2*c)^3
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(
cos(1/2*d*x+1/2*c),2^(1/2))-20*A*cos(1/2*d*x+1/2*c)^4+38*B*cos(1/2*d*x+1/2
*c)^4+9*A*cos(1/2*d*x+1/2*c)^2-15*B*cos(1/2*d*x+1/2*c)^2-A+B)/a^2/cos(1/2*
d*x+1/2*c)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*
d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.153.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.153.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 352, normalized size of antiderivative = 2.59

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{2(3(A-2B)\cos(dx+c) + 2A - 5B)\sqrt{\cos(dx+c)}\sin(dx+c) + (\sqrt{2}(-2iA+5iB)\cos(dx+c))^2 - \dots}{\dots}$$

```
input integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm
m="fricas")
```

```
output 1/6*(2*(3*(A - 2*B)*cos(d*x + c) + 2*A - 5*B)*sqrt(cos(d*x + c))*sin(d*x +
c) + (sqrt(2)*(-2*I*A + 5*I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(2*I*A - 5*I*B)
*cos(d*x + c) + sqrt(2)*(-2*I*A + 5*I*B))*weierstrassPInverse(-4, 0, cos(d
*x + c) + I*sin(d*x + c)) + (sqrt(2)*(2*I*A - 5*I*B)*cos(d*x + c)^2 - 2*sq
rt(2)*(-2*I*A + 5*I*B)*cos(d*x + c) + sqrt(2)*(2*I*A - 5*I*B))*weierstrass
PInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*(sqrt(2)*(I*A - 4*I*B)*
cos(d*x + c)^2 + 2*sqrt(2)*(I*A - 4*I*B)*cos(d*x + c) + sqrt(2)*(I*A - 4*I
*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*si
n(d*x + c))) - 3*(sqrt(2)*(-I*A + 4*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(-I*A
+ 4*I*B)*cos(d*x + c) + sqrt(2)*(-I*A + 4*I*B))*weierstrassZeta(-4, 0, wei
erstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a^2*d*cos(d*x +
c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)
```

3.153.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2,x)
```

```
output Timed out
```

3.153.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^2, x)`

3.153.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^2, x)`

3.153.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2,x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2, x)`

3.153. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

3.154
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx$$

3.154.1 Optimal result 1504
 3.154.2 Mathematica [C] (warning: unable to verify) 1505
 3.154.3 Rubi [A] (verified) 1506
 3.154.4 Maple [B] (verified) 1509
 3.154.5 Fricas [C] (verification not implemented) 1510
 3.154.6 Sympy [F(-1)] 1511
 3.154.7 Maxima [F] 1511
 3.154.8 Giac [F] 1511
 3.154.9 Mupad [F(-1)] 1512

3.154.1 Optimal result

Integrand size = 33, antiderivative size = 121

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^2} dx = -\frac{BE(\frac{1}{2}(c+dx)|2)}{a^2d} + \frac{(A+2B) \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3a^2d} + \frac{B\sqrt{\cos(c+dx)} \sin(c+dx)}{a^2d(1+\cos(c+dx))} + \frac{(A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3d(a+a \cos(c+dx))^2}$$

output

```
-B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/a^2/d+1/3*(A+2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a^2/d+B*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d/(1+cos(d*x+c))+1/3*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^2
```

3.154.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.05 (sec) , antiderivative size = 694, normalized size of antiderivative = 5.74

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx =$$

$$\frac{2A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \cos(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))^2} +$$

$$\frac{4B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \cos(dx - \arctan(\cot(c)))}}{3d(a+a\cos(c+dx))^2} +$$

$$\frac{\cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \left(\frac{4B \csc(c)}{d} + \frac{4B \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) \sin\left(\frac{dx}{2}\right)}{d} + \frac{2 \sec\left(\frac{c}{2}\right) \sec^3\left(\frac{c}{2} + \frac{dx}{2}\right) (A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right))}{3d} \right)}{(a+a\cos(c+dx))^2} +$$

$$\frac{B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{d(a+a\cos(c+dx))^2}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2, x]`

output $(-2*A*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[d*x - \text{ArcTan}[\cot[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\cot[c]]]*\sqrt{1 - \sin[d*x - \text{ArcTan}[\cot[c]]}]*\sqrt{-(\sqrt{1 + \cot[c]^2}*\sin[c]*\sin[d*x - \text{ArcTan}[\cot[c]]])}]*\sqrt{1 + \sin[d*x - \text{ArcTan}[\cot[c]]}]/(3*d*(a + a*\cos[c + d*x])^2*\sqrt{1 + \cot[c]^2}) - (4*B*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\text{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \sin[d*x - \text{ArcTan}[\cot[c]]]^2]*\sec[c/2]*\sec[d*x - \text{ArcTan}[\cot[c]]]*\sqrt{1 - \sin[d*x - \text{ArcTan}[\cot[c]]}]*\sqrt{-(\sqrt{1 + \cot[c]^2}*\sin[c]*\sin[d*x - \text{ArcTan}[\cot[c]]])}]*\sqrt{1 + \sin[d*x - \text{ArcTan}[\cot[c]]}]/(3*d*(a + a*\cos[c + d*x])^2*\sqrt{1 + \cot[c]^2}) + (\cos[c/2 + (d*x)/2]^4*\sqrt{\cos[c + d*x]}*((4*B*\csc[c])/d + (4*B*\sec[c/2]*\sec[c/2 + (d*x)/2]*\sin[(d*x)/2])/d + (2*\sec[c/2]*\sec[c/2 + (d*x)/2]^3*(A*\sin[(d*x)/2] - B*\sin[(d*x)/2]))/(3*d) + (2*(A - B)*\sec[c/2 + (d*x)/2]^2*\tan[c/2]/(3*d)))/(a + a*\cos[c + d*x])^2 + (B*\cos[c/2 + (d*x)/2]^4*\csc[c/2]*\sec[c/2]*(\text{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \cos[d*x + \text{ArcTan}[\tan[c]]]^2]*\sin[d*x + \text{ArcTan}[\tan[c]]]*\tan[c])/(\sqrt{1 - \cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{1 + \cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{1 + \tan[c]^2}]*\sqrt{1 + \tan[c]^2}) - ((\sin[d*x + \text{ArcTan}[\tan[c]]]*\tan[c])/(\sqrt{1 + \tan[c]^2}) + (2*\cos[c]^2*\cos[d*x + \text{ArcTan}[\tan[c]]]*\sqrt{1 + \tan[c]^2})/(\cos[c]^2 + \sin[c]^2))/\sqrt{\cos[c]*\cos[d*x + \text{ArcTan}[\tan[c]]}]*\sqrt{1 + \tan[c]^2})))/(d*(a + a*\cos[c + d*x])^2)$

3.154.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.07, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3456, 27, 3042, 3457, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^2} dx$$

↓ 3042

$$\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^2} dx$$

↓ 3456

$$\frac{\int \frac{a(A-B)+a(A+5B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx}{3a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

3.154. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{a(A-B)+a(A+5B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 27 \\
& \int \frac{a(A-B)+a(A+5B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)} dx + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a^2(A+2B)-3a^2B\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx}{6a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3457 \\
& \frac{\int \frac{a^2(A+2B)-3a^2B\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{6a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3042 \\
& \frac{a^2(A+2B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^2B\int \sqrt{\cos(c+dx)} dx}{6a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3227 \\
& \frac{a^2(A+2B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^2B\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{6a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3042 \\
& \frac{a^2(A+2B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6a^2BE(\frac{1}{2}(c+dx)|_2)}{d}}{6a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3119 \\
& \frac{a^2(A+2B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6a^2BE(\frac{1}{2}(c+dx)|_2)}{d}}{6a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \\
& \quad \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow 3120
\end{aligned}$$

3.154. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$

$$\frac{\frac{2a^2(A+2B)\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} - \frac{6a^2BE\left(\frac{1}{2}(c+dx), 2\right)}{d}}{a^2} + \frac{6B\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} + \frac{6a^2}{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{3d(a\cos(c+dx)+a)^2}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^2,x]`

output `((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((-6*a^2*B*EllipticE[(c + d*x)/2, 2])/d + (2*a^2*(A + 2*B)*EllipticF[(c + d*x)/2, 2])/d)/a^2 + (6*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(1 + Cos[c + d*x]))/(6*a^2)`

3.154.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.154.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 349 vs. $2(165) = 330$.

Time = 4.58 (sec) , antiderivative size = 350, normalized size of antiderivative = 2.89

method	result
default	$-\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(2A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 1F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 12\right)}$

```
input int((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)
```

output
$$-1/6*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))*\cos(1/2*d*x+1/2*c)^3+12*B*\cos(1/2*d*x+1/2*c)^6+4*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))*\cos(1/2*d*x+1/2*c)^3+6*B*\cos(1/2*d*x+1/2*c)^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+2*A*\cos(1/2*d*x+1/2*c)^4-20*B*\cos(1/2*d*x+1/2*c)^4-3*A*\cos(1/2*d*x+1/2*c)^2+9*B*\cos(1/2*d*x+1/2*c)^2+A-B)/a^2/\cos(1/2*d*x+1/2*c)^3/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$$

3.154.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.60

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

$$= \frac{2(3B\cos(dx+c)+A+2B)\sqrt{\cos(dx+c)}\sin(dx+c)+(\sqrt{2}(-iA-2iB)\cos(dx+c))^2-2\sqrt{2}(iA+2iB)\cos(dx+c)}{(a+a\cos(c+dx))^2}$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^2,x, algorithm="fracas")`

output
$$1/6*(2*(3*B*\cos(d*x+c)+A+2*B)*\text{sqrt}(\cos(d*x+c))*\sin(d*x+c)+(\text{sqrt}(2)*(-I*A-2*I*B)*\cos(d*x+c)^2-2*\text{sqrt}(2)*(I*A+2*I*B)*\cos(d*x+c)+\text{sqrt}(2)*(-I*A-2*I*B))*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+(\text{sqrt}(2)*(I*A+2*I*B)*\cos(d*x+c)^2-2*\text{sqrt}(2)*(-I*A-2*I*B)*\cos(d*x+c)+\text{sqrt}(2)*(I*A+2*I*B))*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))-3*(I*\text{sqrt}(2)*B*\cos(d*x+c)^2+2*I*\text{sqrt}(2)*B*\cos(d*x+c)+I*\text{sqrt}(2)*B)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c)))-3*(-I*\text{sqrt}(2)*B*\cos(d*x+c)^2-2*I*\text{sqrt}(2)*B*\cos(d*x+c)-I*\text{sqrt}(2)*B)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))))/(a^2*d*\cos(d*x+c)^2+2*a^2*d*\cos(d*x+c)+a^2*d)$$

3.154.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)**(1/2)/(a+a*cos(d*x+c))**2,x)`

output `Timed out`

3.154.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x+c)+A)*sqrt(cos(d*x+c))/(a*cos(d*x+c)+a)^2,x)`

3.154.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(a\cos(dx+c)+a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x+c)+A)*sqrt(cos(d*x+c))/(a*cos(d*x+c)+a)^2,x)`

3.154.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2, x)`output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^2, x)`

3.155 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+a \cos(c+dx))^2}} dx$

3.155.1 Optimal result 1513
 3.155.2 Mathematica [C] (warning: unable to verify) 1514
 3.155.3 Rubi [A] (verified) 1515
 3.155.4 Maple [B] (verified) 1518
 3.155.5 Fricas [C] (verification not implemented) 1519
 3.155.6 Sympy [F(-1)] 1519
 3.155.7 Maxima [F] 1520
 3.155.8 Giac [F] 1520
 3.155.9 Mupad [F(-1)] 1520

3.155.1 Optimal result

Integrand size = 33, antiderivative size = 121

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + a \cos(c + dx))^2}} dx = \frac{AE(\frac{1}{2}(c + dx) | 2)}{a^2 d} + \frac{(2A + B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3a^2 d} - \frac{A \sqrt{\cos(c + dx)} \sin(c + dx)}{a^2 d(1 + \cos(c + dx))} - \frac{(A - B) \sqrt{\cos(c + dx)} \sin(c + dx)}{3d(a + a \cos(c + dx))^2}$$

output

```
A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d+1/3*(2*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d-A*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d/(1+cos(d*x+c))-1/3*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^2
```


3.155.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.66 (sec) , antiderivative size = 695, normalized size of antiderivative = 5.74

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx =$$

$$\frac{4A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \cos(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))^2} +$$

$$\frac{2B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \cos(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))^2} +$$

$$\frac{\cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \left(-\frac{4A \csc(c)}{d} - \frac{4A \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) \sin\left(\frac{dx}{2}\right)}{d} - \frac{2 \sec\left(\frac{c}{2}\right) \sec^3\left(\frac{c}{2} + \frac{dx}{2}\right) \left(A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right)\right)}{3d} \right)}{(a + a \cos(c + dx))^2} +$$

$$\frac{A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))}} \right)}{d(a + a \cos(c + dx))^2}$$

```
input Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2),x]
```

output

$$\begin{aligned}
& (-4A \cos[c/2 + (dx)/2]^4 \operatorname{Csc}[c/2] \operatorname{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]^2] \operatorname{Sec}[c/2] \operatorname{Sec}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]] \operatorname{Sqrt}[1 - \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]] \operatorname{Sqrt}[-(\operatorname{Sqrt}[1 + \operatorname{Cot}[c]^2] \operatorname{Sin}[c] \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]])] \operatorname{Sqrt}[1 + \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]]) / (3d(a + a \cos[c + dx])^2 \operatorname{Sqrt}[1 + \operatorname{Cot}[c]^2]) - (2B \cos[c/2 + (dx)/2]^4 \operatorname{Csc}[c/2] \operatorname{HypergeometricPFQ}[\{1/4, 1/2\}, \{5/4\}, \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]^2] \operatorname{Sec}[c/2] \operatorname{Sec}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]] \operatorname{Sqrt}[1 - \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]] \operatorname{Sqrt}[-(\operatorname{Sqrt}[1 + \operatorname{Cot}[c]^2] \operatorname{Sin}[c] \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]])] \operatorname{Sqrt}[1 + \operatorname{Sin}[dx - \operatorname{ArcTan}[\operatorname{Cot}[c]]]]) / (3d(a + a \cos[c + dx])^2 \operatorname{Sqrt}[1 + \operatorname{Cot}[c]^2]) + (\cos[c/2 + (dx)/2]^4 \operatorname{Sqrt}[\cos[c + dx]] * ((-4A \operatorname{Csc}[c]) / d - (4A \operatorname{Sec}[c/2] \operatorname{Sec}[c/2 + (dx)/2] \operatorname{Sin}[(dx)/2]) / d - (2 \operatorname{Sec}[c/2] \operatorname{Sec}[c/2 + (dx)/2]^3 (A \operatorname{Sin}[(dx)/2] - B \operatorname{Sin}[(dx)/2])) / (3d) - (2(A - B) \operatorname{Sec}[c/2 + (dx)/2]^2 \operatorname{Tan}[c/2]) / (3d)) / (a + a \cos[c + dx])^2 - (A \cos[c/2 + (dx)/2]^4 \operatorname{Csc}[c/2] \operatorname{Sec}[c/2] * (\operatorname{HypergeometricPFQ}[\{-1/2, -1/4\}, \{3/4\}, \operatorname{Cos}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]]^2] \operatorname{Sin}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Tan}[c]) / (\operatorname{Sqrt}[1 - \operatorname{Cos}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]]] \operatorname{Sqrt}[1 + \operatorname{Cos}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]]] \operatorname{Sqrt}[\operatorname{Cos}[c] \operatorname{Cos}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Sqrt}[1 + \operatorname{Tan}[c]^2] \operatorname{Sqrt}[1 + \operatorname{Tan}[c]^2]) - ((\operatorname{Sin}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Tan}[c]) / \operatorname{Sqrt}[1 + \operatorname{Tan}[c]^2] + (2 \operatorname{Cos}[c]^2 \operatorname{Cos}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Sqrt}[1 + \operatorname{Tan}[c]^2]) / (\operatorname{Cos}[c]^2 + \operatorname{Sin}[c]^2)) / \operatorname{Sqrt}[\operatorname{Cos}[c] \operatorname{Cos}[dx + \operatorname{ArcTan}[\operatorname{Tan}[c]]] \operatorname{Sqrt}[1 + \operatorname{Tan}[c]^2])) / (d(a + a \cos[c + dx])^2)
\end{aligned}$$

3.155.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.07, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3457, 27, 3042, 3457, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a(5A+B) - a(A-B) \cos(c+dx)}{2\sqrt{\cos(c+dx)} (\cos(c+dx)a+a)} dx}{3a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx) + a)^2}
\end{aligned}$$

3.155. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)} (a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{a(5A+B) - a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx \quad \downarrow \quad 27 \\
& \frac{\int \frac{a(5A+B) - a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \quad 3042 \\
& \int \frac{a(5A+B) - a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)} dx \quad \downarrow \quad 3457 \\
& \frac{\int \frac{(2A+B)a^2 + 3A\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}} dx}{6a^2} - \frac{6A\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \quad 3042 \\
& \int \frac{(2A+B)a^2 + 3A\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \quad \downarrow \quad 3227 \\
& \frac{\int \frac{(2A+B)a^2 + 3A\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{6a^2} - \frac{6A\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \quad 3042 \\
& \frac{a^2(2A+B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^2 A \int \sqrt{\cos(c+dx)} dx}{6a^2} - \frac{6A\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \quad 3042 \\
& \frac{a^2(2A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a^2 A \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{6a^2} - \frac{6A\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \quad 3119 \\
& \frac{a^2(2A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^2 AE(\frac{1}{2}(c+dx)|2)}{d}}{6a^2} - \frac{6A\sin(c+dx)\sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} \\
& \quad \downarrow \quad 3120
\end{aligned}$$

3.155. $\int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a\cos(c+dx))^2} dx$

$$\frac{\frac{2a^2(2A+B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} + \frac{6a^2AE\left(\frac{1}{2}(c+dx)\right)}{d}}{a^2} - \frac{6A \sin(c+dx) \sqrt{\cos(c+dx)}}{d(\cos(c+dx)+1)} - \frac{6a^2}{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}} - \frac{6a^2}{3d(a \cos(c+dx) + a)^2}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2),x]`

output `-1/3*((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^2) + (((6*a^2*A*EllipticE[(c + d*x)/2, 2])/d + (2*a^2*(2*A + B)*EllipticF[(c + d*x)/2, 2])/d)/a^2 - (6*A*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(1 + Cos[c + d*x])))/(6*a^2)`

3.155.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.155.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 349 vs. 2(165) = 330.

Time = 3.99 (sec) , antiderivative size = 350, normalized size of antiderivative = 2.89

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(12A\left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 4A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right)$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2/cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/6*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*A*cos(1/2*
d*x+1/2*c)^6-4*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^
(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3+6*A*cos(1
/2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^
(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*B*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*
cos(1/2*d*x+1/2*c)^3-16*A*cos(1/2*d*x+1/2*c)^4-2*B*cos(1/2*d*x+1/2*c)^4+3*
A*cos(1/2*d*x+1/2*c)^2+3*B*cos(1/2*d*x+1/2*c)^2+A-B)/a^2/cos(1/2*d*x+1/2*c
)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c
)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.155.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+a \cos(c+dx))^2}} dx$$

3.155.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 318, normalized size of antiderivative = 2.63

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx =$$

$$\frac{2(3A \cos(dx + c) + 4A - B)\sqrt{\cos(dx + c)} \sin(dx + c) - (\sqrt{2}(-2iA - iB) \cos(dx + c))^2 - 2\sqrt{2}(2iA + iB) \cos(dx + c)}{a^2 d \cos(dx + c)^2 + 2a^2 d \cos(dx + c) + a^2 d}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/6*(2*(3*A*cos(d*x + c) + 4*A - B)*sqrt(cos(d*x + c))*sin(d*x + c) - (sqrt(2)*(-2*I*A - I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(2*I*A + I*B)*cos(d*x + c) + sqrt(2)*(-2*I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - (sqrt(2)*(2*I*A + I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(-2*I*A - I*B)*cos(d*x + c) + sqrt(2)*(2*I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(-I*sqrt(2)*A*cos(d*x + c)^2 - 2*I*sqrt(2)*A*cos(d*x + c) - I*sqrt(2)*A)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(I*sqrt(2)*A*cos(d*x + c)^2 + 2*I*sqrt(2)*A*cos(d*x + c) + I*sqrt(2)*A)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/((a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.155.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2/cos(d*x+c)**(1/2),x)`

output `Timed out`

3.155.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/cos(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sqrt(cos(d*x + c))), x)`

3.155.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/cos(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sqrt(cos(d*x + c))), x)`

3.155.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^2), x)`

3.156 $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$

3.156.1 Optimal result 1521
 3.156.2 Mathematica [C] (warning: unable to verify) 1522
 3.156.3 Rubi [A] (verified) 1523
 3.156.4 Maple [B] (verified) 1526
 3.156.5 Fricas [C] (verification not implemented) 1527
 3.156.6 Sympy [F(-1)] 1528
 3.156.7 Maxima [F] 1528
 3.156.8 Giac [F] 1529
 3.156.9 Mupad [F(-1)] 1529

3.156.1 Optimal result

Integrand size = 33, antiderivative size = 168

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = -\frac{(4A - B)E(\frac{1}{2}(c + dx)|2)}{a^2d} - \frac{(5A - 2B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3a^2d} + \frac{(4A - B) \sin(c + dx)}{a^2d\sqrt{\cos(c + dx)}} - \frac{(5A - 2B) \sin(c + dx)}{3a^2d\sqrt{\cos(c + dx)}(1 + \cos(c + dx))} - \frac{(A - B) \sin(c + dx)}{3d\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2}$$

output

```
-(4*A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d-1/3*(5*A-2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d+(4*A-B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(1/2)-1/3*(5*A-2*B)*sin(d*x+c)/a^2/d/(1+cos(d*x+c))/cos(d*x+c)^(1/2)-1/3*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^2/cos(d*x+c)^(1/2)
```


3.156.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.75 (sec) , antiderivative size = 979, normalized size of antiderivative = 5.83

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx$$

$$= \frac{10A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))^2} + \frac{4B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; \sin^2(dx - \arctan(\cot(c)))\right) \sec\left(\frac{c}{2}\right) \sec(dx - \arctan(\cot(c))) \sqrt{1 - \sin^2(dx - \arctan(\cot(c)))}}{3d(a + a \cos(c + dx))^2} + \frac{\cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \left(\frac{2(2A + 2A \cos(c) - B \cos(c)) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \sec(c)}{d} + \frac{2 \sec\left(\frac{c}{2}\right) \sec^3\left(\frac{c}{2} + \frac{dx}{2}\right) (A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right))}{3d} \right)}{(a + a \cos(c + dx))^2} + \frac{4A \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos(dx + \arctan(\tan(c)))}} \right)}{d(a + a \cos(c + dx))^2} + \frac{B \cos^4\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right) \left(\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{3}{4}; \cos^2(dx + \arctan(\tan(c)))\right) \sin(dx + \arctan(\tan(c))) \tan(c)}{\sqrt{1 - \cos(dx + \arctan(\tan(c)))} \sqrt{1 + \cos(dx + \arctan(\tan(c)))} \sqrt{\cos(c) \cos(dx + \arctan(\tan(c)))} \sqrt{1 - \cos(dx + \arctan(\tan(c)))}} \right)}{d(a + a \cos(c + dx))^2}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^2),x]`

output

```
(10*A*cos[c/2 + (d*x)/2]^4*csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])]/(3*d*(a + a*cos[c + d*x])^2*Sqrt[1 + Cot[c]^2]) - (4*B*cos[c/2 + (d*x)/2]^4*csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])]/(3*d*(a + a*cos[c + d*x])^2*Sqrt[1 + Cot[c]^2]) + (Cos[c/2 + (d*x)/2]^4*Sqrt[Cos[c + d*x]]*((2*(2*A + 2*A*cos[c] - B*cos[c])*Csc[c/2]*Sec[c/2]*Sec[c])/d + (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(A*sin[(d*x)/2] - B*sin[(d*x)/2]))/(3*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(2*A*sin[(d*x)/2] - B*sin[(d*x)/2])/d + (8*A*Sec[c]*Sec[c + d*x]*Sin[d*x])/d + (2*(A - B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(3*d)))/(a + a*cos[c + d*x])^2 + (4*A*cos[c/2 + (d*x)/2]^4*csc[c/2]*Sec[c/2]*((HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*cos[c]^2*cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2])/(Cos[c]^2 + Sin[c]^2))/Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]])
```

3.156.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3457, 27, 3042, 3457, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^2} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}}(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^2} dx$$

↓ 3457

$$\frac{\int \frac{a(7A - B) - 3a(A - B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)} dx}{3a^2} - \frac{(A - B) \sin(c + dx)}{3d \sqrt{\cos(c + dx)(a \cos(c + dx) + a)^2}}$$

3.156. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx$

$$\begin{aligned}
 & \int \frac{a(7A-B)-3a(A-B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)} dx \quad \downarrow \quad 27 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{a(7A-B)-3a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(\sin(c+dx+\frac{\pi}{2})a+a)} dx \quad \downarrow \quad 3042 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{3a^2(4A-B)-a^2(5A-2B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx \quad \downarrow \quad 3457 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{2(5A-2B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{3a^2(4A-B)-a^2(5A-2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx \quad \downarrow \quad 3042 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{2(5A-2B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{3a^2(4A-B)}{\cos^{\frac{3}{2}}(c+dx)} dx - a^2(5A-2B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx \quad \downarrow \quad 3227 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{3a^2(4A-B)}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx - a^2(5A-2B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \quad \downarrow \quad 3042 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{3a^2(4A-B)\left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx\right) - a^2(5A-2B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2} dx \quad \downarrow \quad 3116 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \int \frac{3a^2(4A-B)\left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx\right) - a^2(5A-2B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2} dx \quad \downarrow \quad 3042 \\
 & \frac{6a^2}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \frac{(A-B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2}
 \end{aligned}$$

3.156. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{3a^2(4A-B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) - a^2(5A-2B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2} - \frac{2(5A-2B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}(\cos(c+dx)+1)} \\
& \frac{6a^2 (A-B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3119} \\
& \frac{3a^2(4A-B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) - a^2(5A-2B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2} - \frac{2(5A-2B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}(\cos(c+dx)+1)} \\
& \frac{6a^2 (A-B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3120} \\
& \frac{3a^2(4A-B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) - \frac{2a^2(5A-2B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{d}}{a^2} - \frac{2(5A-2B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}(\cos(c+dx)+1)} \\
& \frac{6a^2 (A-B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^2}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^2),x]`

output `-1/3*((A - B)*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2) + ((-2*(5*A - 2*B)*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]]*(1 + Cos[c + d*x])) + ((-2*a^2*(5*A - 2*B)*EllipticF[(c + d*x)/2, 2])/d + 3*a^2*(4*A - B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]]))) /a^2)/(6*a^2)`

3.156.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.156. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$

rule 3116 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.156.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. 2(208) = 416.

Time = 4.28 (sec) , antiderivative size = 494, normalized size of antiderivative = 2.94

method	result
default	$-\frac{2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(5AF\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)-12AE\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\right)}{\dots}$

3.156.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$$

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)`

output `-1/6*(2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(5*A*EllipticF(cos(1/2*d*
x+1/2*c),2^(1/2))-12*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*B*EllipticF
(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*co
s(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*s
in(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
^(1/2)*(5*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-12*A*EllipticE(cos(1/2*d
*x+1/2*c),2^(1/2))-2*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*EllipticE
(cos(1/2*d*x+1/2*c),2^(1/2)))*cos(1/2*d*x+1/2*c)-12*(-2*sin(1/2*d*x+1/2*c)
^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*A-B)*sin(1/2*d*x+1/2*c)^6+2*(-2*sin(1/2*
d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(43*A-10*B)*sin(1/2*d*x+1/2*c)^4-
(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(37*A-7*B)*sin(1/2*d*
x+1/2*c)^2)/a^2/cos(1/2*d*x+1/2*c)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.156.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 407, normalized size of antiderivative = 2.42

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx$$

$$= \frac{2(3(4A - B) \cos(dx + c)^2 + (19A - 4B) \cos(dx + c) + 6A) \sqrt{\cos(dx + c)} \sin(dx + c) + (\sqrt{2}(5iA - 2B) \cos(dx + c) + (2A - B) \sin(dx + c)) \sqrt{\cos(dx + c)}}{d}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2,x, algorithm
m="fricas")`

```
output 1/6*(2*(3*(4*A - B)*cos(d*x + c)^2 + (19*A - 4*B)*cos(d*x + c) + 6*A)*sqrt
(cos(d*x + c))*sin(d*x + c) + (sqrt(2)*(5*I*A - 2*I*B)*cos(d*x + c)^3 - 2*
sqrt(2)*(-5*I*A + 2*I*B)*cos(d*x + c)^2 + sqrt(2)*(5*I*A - 2*I*B)*cos(d*x
+ c))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)
*(-5*I*A + 2*I*B)*cos(d*x + c)^3 - 2*sqrt(2)*(5*I*A - 2*I*B)*cos(d*x + c)^
2 + sqrt(2)*(-5*I*A + 2*I*B)*cos(d*x + c))*weierstrassPInverse(-4, 0, cos(
d*x + c) - I*sin(d*x + c)) - 3*(sqrt(2)*(4*I*A - I*B)*cos(d*x + c)^3 + 2*s
qrt(2)*(4*I*A - I*B)*cos(d*x + c)^2 + sqrt(2)*(4*I*A - I*B)*cos(d*x + c))*
weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x
+ c))) - 3*(sqrt(2)*(-4*I*A + I*B)*cos(d*x + c)^3 + 2*sqrt(2)*(-4*I*A + I
*B)*cos(d*x + c)^2 + sqrt(2)*(-4*I*A + I*B)*cos(d*x + c))*weierstrassZeta(
-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a^2*d*
cos(d*x + c)^3 + 2*a^2*d*cos(d*x + c)^2 + a^2*d*cos(d*x + c))
```

3.156.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c))**2,x)
```

```
output Timed out
```

3.156.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}}} dx$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2,x, algorithm
m="maxima")
```

```
output integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2))
, x)
```

3.156.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2)), x)`

3.156.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} (a + a \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^2), x)`

$$3.157 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$$

3.157.1 Optimal result 1530
 3.157.2 Mathematica [C] (warning: unable to verify) 1531
 3.157.3 Rubi [A] (verified) 1531
 3.157.4 Maple [B] (verified) 1535
 3.157.5 Fricas [C] (verification not implemented) 1536
 3.157.6 Sympy [F(-1)] 1537
 3.157.7 Maxima [F] 1537
 3.157.8 Giac [F] 1538
 3.157.9 Mupad [F(-1)] 1538

3.157.1 Optimal result

Integrand size = 33, antiderivative size = 201

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \frac{(7A - 4B)E(\frac{1}{2}(c + dx) | 2)}{a^2 d} + \frac{5(2A - B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3a^2 d} + \frac{5(2A - B) \sin(c + dx)}{3a^2 d \cos^{\frac{3}{2}}(c + dx)} - \frac{(7A - 4B) \sin(c + dx)}{a^2 d \sqrt{\cos(c + dx)}} - \frac{(7A - 4B) \sin(c + dx)}{3a^2 d \cos^{\frac{3}{2}}(c + dx)(1 + \cos(c + dx))} - \frac{(A - B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2}$$

output

```
(7*A-4*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d+5/3*(2*A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/d+5/3*(2*A-B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(3/2)-1/3*(7*A-4*B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(3/2)/(1+cos(d*x+c))-1/3*(A-B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2-(7*A-4*B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(1/2)
```

3.157.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.10 (sec) , antiderivative size = 1020, normalized size of antiderivative = 5.07

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \text{Too large to display}$$

```
input Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^2),x]
```

```
output (-20*A*Cos[c/2 + (d*x)/2]^4*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]]]/(3*d*(a + a*Cos[c + d*x])^2*Sqrt[1 + Cot[c]^2]) + (10*B*Cos[c/2 + (d*x)/2]^4*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]]]/(3*d*(a + a*Cos[c + d*x])^2*Sqrt[1 + Cot[c]^2]) + (Cos[c/2 + (d*x)/2]^4*Sqrt[Cos[c + d*x]]*((-2*(4*A - 2*B + 3*A*Cos[c] - 2*B*Cos[c])*Csc[c/2]*Sec[c/2]*Sec[c])/d - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(3*A*Sin[(d*x)/2] - 2*B*Sin[(d*x)/2]))/d - (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(3*d) + (8*A*Sec[c]*Sec[c + d*x]^2*Sin[d*x])/(3*d) + (8*Sec[c]*Sec[c + d*x]*(A*Sin[c] - 6*A*Sin[d*x] + 3*B*Sin[d*x]))/(3*d) - (2*(A - B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(3*d)))/(a + a*Cos[c + d*x])^2 - (7*A*Cos[c/2 + (d*x)/2]^4*Csc[c/2]*Sec[c/2]*((HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Tan[c]^2]*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x...
```

3.157.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.98, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.157. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{3a(3A-B) - 5a(A-B) \cos(c+dx)}{2 \cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)} dx}{3a^2} - \frac{(A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{3a(3A-B) - 5a(A-B) \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)} dx}{6a^2} - \frac{(A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3a(3A-B) - 5a(A-B) \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)} dx}{6a^2} - \frac{(A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{3(5a^2(2A-B) - a^2(7A-4B) \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx}{6a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} - \frac{(A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{3 \int \frac{5a^2(2A-B) - a^2(7A-4B) \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)} dx}{6a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} - \frac{(A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3 \int \frac{5a^2(2A-B) - a^2(7A-4B) \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2}} dx}{6a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} - \frac{(A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{3227}
\end{aligned}$$

3.157. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{3 \left(5a^2(2A-B) \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)} dx - a^2(7A-4B) \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx \right)}{a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} \\
& \frac{6a^2}{(A-B) \sin(c+dx)} \\
& \frac{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{\downarrow 3042} \\
& \frac{3 \left(5a^2(2A-B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}}} dx - a^2(7A-4B) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}} dx \right)}{a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} \\
& \frac{6a^2}{(A-B) \sin(c+dx)} \\
& \frac{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{\downarrow 3116} \\
& \frac{3 \left(5a^2(2A-B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - a^2(7A-4B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx \right) \right)}{a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} \\
& \frac{6a^2}{(A-B) \sin(c+dx)} \\
& \frac{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{\downarrow 3042} \\
& \frac{3 \left(5a^2(2A-B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - a^2(7A-4B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) \right)}{a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} \\
& \frac{6a^2}{(A-B) \sin(c+dx)} \\
& \frac{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{\downarrow 3119} \\
& \frac{3 \left(5a^2(2A-B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - a^2(7A-4B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E(\frac{1}{2}(c+dx)|2)}{d} \right) \right)}{a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} \\
& \frac{6a^2}{(A-B) \sin(c+dx)} \\
& \frac{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}{\downarrow 3120}
\end{aligned}$$

3.157. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$

$$\frac{3 \left(5a^2(2A-B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - a^2(7A-4B) \left(\frac{2 \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) \right)}{a^2} - \frac{2(7A-4B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)+1)} - \frac{6a^2 (A-B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx) + a)^2}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^2), x]`

output `-1/3*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^2) + ((-2*(7*A - 4*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(1 + Cos[c + d*x])) + (3*(5*a^2*(2*A - B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))) - a^2*(7*A - 4*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]]))))/a^2)/(6*a^2)`

3.157.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 3227 Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3457 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d)), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.157.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. $2(237) = 474$.

Time = 6.47 (sec) , antiderivative size = 723, normalized size of antiderivative = 3.60

method	result	size
default	Expression too large to display	723

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)
```

output

```
-1/2*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/a^2*((4*A-2
*B)*(cos(1/2*d*x+1/2*c)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-EllipticE(cos(1/2*d*x+1
/2*c),2^(1/2)))-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)/cos(1/2*d*x+1
/2*c)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+1/3*(A-B)*(2*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(2*EllipticF(c
os(1/2*d*x+1/2*c),2^(1/2))-3*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*sin(1/
2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/
2*d*x+1/2*c)^2-1)^(1/2)*(2*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2)))*cos(1/2*d*x+1/2*c)-12*sin(1/2*d*x+1/2*c)^6
+20*sin(1/2*d*x+1/2*c)^4-7*sin(1/2*d*x+1/2*c)^2)/cos(1/2*d*x+1/2*c)/(-2*si
n(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(-1+sin(1/2*d*x+1/2*c)^2)+4
*A*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)
^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*c
os(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+(-8*A+4*B)/sin(1/2*d*x+1/2*
c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^
2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(
1/2))))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.157.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 436, normalized size of antiderivative = 2.17

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx =$$

$$\frac{2(3(7A - 4B) \cos(dx + c)^3 + (32A - 19B) \cos(dx + c)^2 + 2(4A - 3B) \cos(dx + c) - 2A) \sqrt{\cos(dx + c)}}{\dots}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^2,x, algorithm="fracas")`

output
$$\begin{aligned} & -1/6*(2*(3*(7*A - 4*B)*\cos(d*x + c)^3 + (32*A - 19*B)*\cos(d*x + c)^2 + 2*(4*A - 3*B)*\cos(d*x + c) - 2*A)*\sqrt{\cos(d*x + c)}*\sin(d*x + c) + 5*(\sqrt{2} \\ &)*(2*I*A - I*B)*\cos(d*x + c)^4 + 2*\sqrt{2}*(2*I*A - I*B)*\cos(d*x + c)^3 + \\ & \sqrt{2}*(2*I*A - I*B)*\cos(d*x + c)^2)*\text{weierstrassPInverse}(-4, 0, \cos(d*x + \\ & c) + I*\sin(d*x + c)) + 5*(\sqrt{2}*(-2*I*A + I*B)*\cos(d*x + c)^4 + 2*\sqrt{2} \\ &)*(-2*I*A + I*B)*\cos(d*x + c)^3 + \sqrt{2}*(-2*I*A + I*B)*\cos(d*x + c)^2)* \\ & \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + 3*(\sqrt{2}*(-7 \\ & *I*A + 4*I*B)*\cos(d*x + c)^4 + 2*\sqrt{2}*(-7*I*A + 4*I*B)*\cos(d*x + c)^3 + \\ & \sqrt{2}*(-7*I*A + 4*I*B)*\cos(d*x + c)^2)*\text{weierstrassZeta}(-4, 0, \text{weierstra} \\ & \text{ssPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + 3*(\sqrt{2}*(7*I*A - 4* \\ & I*B)*\cos(d*x + c)^4 + 2*\sqrt{2}*(7*I*A - 4*I*B)*\cos(d*x + c)^3 + \sqrt{2}*(\\ & 7*I*A - 4*I*B)*\cos(d*x + c)^2)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse} \\ & (-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/(a^2*d*\cos(d*x + c)^4 + 2*a^2*d*co \\ & s(d*x + c)^3 + a^2*d*\cos(d*x + c)^2) \end{aligned}$$

3.157.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c))**2,x)`

output Timed out

3.157.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2)), x)`

3.157.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^2} dx$$

3.157.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2)), x)`

3.157.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^2} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + a \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^2), x)`

3.158 $\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.158.1 Optimal result 1539
 3.158.2 Mathematica [C] (verified) 1540
 3.158.3 Rubi [A] (verified) 1540
 3.158.4 Maple [A] (verified) 1545
 3.158.5 Fricas [C] (verification not implemented) 1545
 3.158.6 Sympy [F(-1)] 1546
 3.158.7 Maxima [F] 1546
 3.158.8 Giac [F] 1547
 3.158.9 Mupad [F(-1)] 1547

3.158.1 Optimal result

Integrand size = 33, antiderivative size = 254

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = -\frac{7(17A-33B)E(\frac{1}{2}(c+dx)|2)}{10a^3d} + \frac{(11A-21B) \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{2a^3d} + \frac{(11A-21B)\sqrt{\cos(c+dx)} \sin(c+dx)}{2a^3d} - \frac{7(17A-33B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{30a^3d} + \frac{(A-B) \cos^{\frac{9}{2}}(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(7A-12B) \cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} + \frac{3(11A-21B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{10d(a^3+a^3 \cos(c+dx))}$$

output

```
-7/10*(17*A-33*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c), 2^(1/2))/a^3/d+1/2*(11*A-21*B)*(cos(1/2*d*x+1/2*c)^2)
^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a^3/d-7/30
*(17*A-33*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/a^3/d+1/5*(A-B)*cos(d*x+c)^(9/2)*
sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(7*A-12*B)*cos(d*x+c)^(7/2)*sin(d*x+c
)/a/d/(a+a*cos(d*x+c))^2+3/10*(11*A-21*B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a
^3+a^3*cos(d*x+c))+1/2*(11*A-21*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^3/d
```

3.158. $\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.158.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.26 (sec) , antiderivative size = 482, normalized size of antiderivative = 1.90

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{\sqrt{\cos(c+dx)} \csc^5(c+dx) \left(-396A + 396B - 680A \cos(c+dx) + 680B \cos(c+dx) + 792A \cos^2(c+dx) \right)}{a^3 d}$$

input `Integrate[(Cos[c + d*x]^(9/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3, x]`

output

```
-1/30*(Sqrt[Cos[c + d*x]]*Csc[c + d*x]^5*(-396*A + 396*B - 680*A*Cos[c + d*x] + 680*B*Cos[c + d*x] + 792*A*Cos[c + d*x]^2 - 792*B*Cos[c + d*x]^2 + 680*A*Cos[c + d*x]^3 - 680*B*Cos[c + d*x]^3 - 440*A*Cos[c + d*x]^4 + 440*B*Cos[c + d*x]^4 - 180*A*Cos[c + d*x]^5 + 180*B*Cos[c + d*x]^5 + 20*A*Cos[c + d*x]^6 - 20*B*Cos[c + d*x]^6 + 66*A*Sin[c + d*x]^2 + 234*B*Sin[c + d*x]^2 + 448*B*Cos[c + d*x]*Sin[c + d*x]^2 + 40*B*Cos[c + d*x]^4*Sin[c + d*x]^2 - 12*B*Cos[c + d*x]^5*Sin[c + d*x]^2 + 165*A*Sin[c + d*x]^4 - 315*B*Sin[c + d*x]^4 + 15*(11*A - 21*B)*Hypergeometric2F1[1/4, 1/2, 5/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] - 448*B*Cos[c + d*x]*Hypergeometric2F1[3/4, 5/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] + 680*A*Cos[c + d*x]*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] - 680*B*Cos[c + d*x]*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] - 90*B*Sin[2*(c + d*x)]^2 - 24*B*Csc[c + d*x]*Sin[2*(c + d*x)]^3))/(a^3*d)
```

3.158.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3456, 27, 3042, 3456, 3042, 3456, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

3.158. $\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sin(c+dx+\frac{\pi}{2})^{9/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{(a \sin(c+dx+\frac{\pi}{2})+a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\cos^{7/2}(c+dx)(9a(A-B)-5a(A-3B)\cos(c+dx))}{2(\cos(c+dx)a+a)^2} dx + \frac{(A-B)\sin(c+dx)\cos^{9/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \int \frac{\cos^{7/2}(c+dx)(9a(A-B)-5a(A-3B)\cos(c+dx))}{10a^2} dx + \frac{(A-B)\sin(c+dx)\cos^{9/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \int \frac{\cos^{7/2}(c+dx)(9a(A-B)-5a(A-3B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx + \frac{(A-B)\sin(c+dx)\cos^{9/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^{7/2} (9a(A-B)-5a(A-3B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx + \frac{(A-B)\sin(c+dx)\cos^{9/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \int \frac{\cos^{5/2}(c+dx)(7a^2(7A-12B)-5a^2(10A-21B)\cos(c+dx))}{\cos(c+dx)a+a} dx + \frac{2a(7A-12B)\sin(c+dx)\cos^{7/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^{9/2}(c+dx) \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2} (7a^2(7A-12B)-5a^2(10A-21B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})a+a} dx + \frac{2a(7A-12B)\sin(c+dx)\cos^{7/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^{9/2}(c+dx) \\
& \quad \downarrow \text{3456} \\
& \int \frac{\cos^{3/2}(c+dx)(9a^3(11A-21B)-7a^3(17A-33B)\cos(c+dx))}{a^2} dx + \frac{9a^2(11A-21B)\sin(c+dx)\cos^{5/2}(c+dx)}{d(a\cos(c+dx)+a)} + \frac{2a(7A-12B)\sin(c+dx)\cos^{7/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\cos^{9/2}(c+dx) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.158. $\int \frac{\cos^{9/2}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{\frac{5 \int \cos^{\frac{3}{2}}(c+dx) \left(9a^3(11A-21B) - 7a^3(17A-33B) \cos(c+dx)\right) dx}{2a^2}}{3a^2} + \frac{9a^2(11A-21B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{2a(7A-12B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{5d(a \cos(c+dx)+a)^3} (A-B) \sin(c+dx) \cos^{\frac{9}{2}}(c+dx)$$

↓ 3042

$$\frac{\frac{5 \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(9a^3(11A-21B) - 7a^3(17A-33B) \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx}{2a^2}}{3a^2} + \frac{9a^2(11A-21B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{2a(7A-12B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{5d(a \cos(c+dx)+a)^3} (A-B) \sin(c+dx) \cos^{\frac{9}{2}}(c+dx)$$

↓ 3227

$$\frac{5 \left(9a^3(11A-21B) \int \cos^{\frac{3}{2}}(c+dx) dx - 7a^3(17A-33B) \int \cos^{\frac{5}{2}}(c+dx) dx\right)}{2a^2} + \frac{9a^2(11A-21B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{2a(7A-12B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{5d(a \cos(c+dx)+a)^3} (A-B) \sin(c+dx) \cos^{\frac{9}{2}}(c+dx)$$

↓ 3042

$$\frac{5 \left(9a^3(11A-21B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} dx - 7a^3(17A-33B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} dx\right)}{2a^2} + \frac{9a^2(11A-21B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{2a(7A-12B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{5d(a \cos(c+dx)+a)^3} (A-B) \sin(c+dx) \cos^{\frac{9}{2}}(c+dx)$$

↓ 3115

$$\frac{5 \left(9a^3(11A-21B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d}\right) - 7a^3(17A-33B) \left(\frac{3}{5} \int \sqrt{\cos(c+dx)} dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d}\right)\right)}{2a^2}}{3a^2} + \frac{9a^2(11A-21B) \sin(c+dx)}{d(a \cos(c+dx)+a)}$$

$$\frac{10a^2}{5d(a \cos(c+dx)+a)^3} (A-B) \sin(c+dx) \cos^{\frac{9}{2}}(c+dx)$$

↓ 3042

3.158. $\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

$$\frac{5 \left(9a^3(11A-21B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) - 7a^3(17A-33B) \left(\frac{3}{5} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) \right)}{2a^2} + \frac{9a^2(11A-21B)}{d(a \cos(c+dx) + a)}$$

$$\frac{(A - B) \sin(c + dx) \cos^{\frac{9}{2}}(c + dx)}{5d(a \cos(c + dx) + a)^3}$$

↓ 3119

$$\frac{5 \left(9a^3(11A-21B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) - 7a^3(17A-33B) \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) \right)}{2a^2} + \frac{9a^2(11A-21B) \sin(c+dx)}{d(a \cos(c+dx) + a)}$$

$$\frac{(A - B) \sin(c + dx) \cos^{\frac{9}{2}}(c + dx)}{5d(a \cos(c + dx) + a)^3}$$

↓ 3120

$$\frac{9a^2(11A-21B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{d(a \cos(c+dx) + a)} + \frac{5 \left(9a^3(11A-21B) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) - 7a^3(17A-33B) \left(\frac{6E(\frac{1}{2}(c+dx)|2)}{5d} + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) \right)}{2a^2}$$

$$\frac{(A - B) \sin(c + dx) \cos^{\frac{9}{2}}(c + dx)}{5d(a \cos(c + dx) + a)^3}$$

input `Int[(Cos[c + d*x]^(9/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^(9/2)*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((2*a*(7*A - 12*B)*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((9*a^2*(11*A - 21*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))) + (5*(9*a^3*(11*A - 21*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)) - 7*a^3*(17*A - 33*B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d))))/(2*a^2)/(3*a^2)/(10*a^2)`

3.158. $\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.158.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*SIN[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.158.4 Maple [A] (verified)

Time = 15.76 (sec) , antiderivative size = 493, normalized size of antiderivative = 1.94

method	result
default	$-\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(192B\left(\cos^{12}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+160A\left(\cos^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-864B\left(\cos^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+468A\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\dots\right)$

```
input int(cos(d*x+c)^(9/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

```
output -1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(192*B*cos(1
/2*d*x+1/2*c)^12+160*A*cos(1/2*d*x+1/2*c)^10-864*B*cos(1/2*d*x+1/2*c)^10+4
68*A*cos(1/2*d*x+1/2*c)^8+330*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d
*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2
*c)^5+714*A*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*
d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-228*B*cos(1/2*
d*x+1/2*c)^8-630*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5-1386*B*
cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2
+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-1058*A*cos(1/2*d*x+1/2*c)^
6+1590*B*cos(1/2*d*x+1/2*c)^6+474*A*cos(1/2*d*x+1/2*c)^4-744*B*cos(1/2*d*x
+1/2*c)^4-47*A*cos(1/2*d*x+1/2*c)^2+57*B*cos(1/2*d*x+1/2*c)^2+3*A-3*B)/a^3
/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.158.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.16 (sec) , antiderivative size = 495, normalized size of antiderivative = 1.95

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{2(12B\cos(dx+c)^4+4(5A-6B)\cos(dx+c)^3+3(79A-147B)\cos(dx+c)^2+2(188A-357B)\cos(dx+c)+\dots)}{(a+a\cos(c+dx))^3}$$

```
input integrate(cos(d*x+c)^(9/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorith
m="fracas")
```

$$3.158. \int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

output `1/60*(2*(12*B*cos(d*x + c)^4 + 4*(5*A - 6*B)*cos(d*x + c)^3 + 3*(79*A - 14
7*B)*cos(d*x + c)^2 + 2*(188*A - 357*B)*cos(d*x + c) + 165*A - 315*B)*sqrt
(cos(d*x + c))*sin(d*x + c) - 15*(sqrt(2)*(11*I*A - 21*I*B)*cos(d*x + c)^3
+ 3*sqrt(2)*(11*I*A - 21*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(11*I*A - 21*I*B
) *cos(d*x + c) + sqrt(2)*(11*I*A - 21*I*B))*weierstrassPInverse(-4, 0, cos
(d*x + c) + I*sin(d*x + c)) - 15*(sqrt(2)*(-11*I*A + 21*I*B)*cos(d*x + c)^
3 + 3*sqrt(2)*(-11*I*A + 21*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-11*I*A + 21*
I*B)*cos(d*x + c) + sqrt(2)*(-11*I*A + 21*I*B))*weierstrassPInverse(-4, 0,
cos(d*x + c) - I*sin(d*x + c)) - 21*(sqrt(2)*(17*I*A - 33*I*B)*cos(d*x +
c)^3 + 3*sqrt(2)*(17*I*A - 33*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(17*I*A - 33
*I*B)*cos(d*x + c) + sqrt(2)*(17*I*A - 33*I*B))*weierstrassZeta(-4, 0, wei
erstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*(sqrt(2)*(-17
*I*A + 33*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-17*I*A + 33*I*B)*cos(d*x + c)^
2 + 3*sqrt(2)*(-17*I*A + 33*I*B)*cos(d*x + c) + sqrt(2)*(-17*I*A + 33*I*B
) *weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d
*x + c))))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*
x + c) + a^3*d)`

3.158.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(9/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)`

output `Timed out`

3.158.7 Maxima [F]

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{9}{2}}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate(cos(d*x+c)^(9/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

3.158. $\int \frac{\cos^{\frac{9}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(9/2)/(a*cos(d*x + c) + a)^3, x)`

3.158.8 Giac [F]

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{9}{2}}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate(cos(d*x+c)^(9/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(9/2)/(a*cos(d*x + c) + a)^3, x)`

3.158.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \int \frac{\cos(c + dx)^{9/2} (A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx$$

input `int((cos(c + d*x)^(9/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output `int((cos(c + d*x)^(9/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3, x)`

3.159
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.159.1 Optimal result 1548
 3.159.2 Mathematica [C] (verified) 1549
 3.159.3 Rubi [A] (verified) 1549
 3.159.4 Maple [A] (verified) 1554
 3.159.5 Fricas [C] (verification not implemented) 1554
 3.159.6 Sympy [F(-1)] 1555
 3.159.7 Maxima [F] 1555
 3.159.8 Giac [F] 1556
 3.159.9 Mupad [F(-1)] 1556

3.159.1 Optimal result

Integrand size = 33, antiderivative size = 219

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{7(7A-17B)E(\frac{1}{2}(c+dx)|2)}{10a^3d} - \frac{(13A-33B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{6a^3d} - \frac{(13A-33B)\sqrt{\cos(c+dx)} \sin(c+dx)}{6a^3d} + \frac{(A-B) \cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(A-2B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{3ad(a+a \cos(c+dx))^2} + \frac{7(7A-17B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{30d(a^3+a^3 \cos(c+dx))}$$

output

```
7/10*(7*A-17*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(
sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d-1/6*(13*A-33*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/5*(A
-B)*cos(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/3*(A-2*B)*cos(d*x+c
)^(5/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+7/30*(7*A-17*B)*cos(d*x+c)^(3/2)
*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))-1/6*(13*A-33*B)*sin(d*x+c)*cos(d*x+c)^(
1/2)/a^3/d
```

3.159.
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.159.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.10 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.01

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{\sqrt{\cos(c+dx)} \csc^5(c+dx) (156A - 156B + 280A \cos(c+dx) - 280B \cos(c+dx) - 312A \cos^2(c+dx) - 312B \cos^2(c+dx))}{(a+a\cos(c+dx))^3}$$

input `Integrate[(Cos[c + d*x]^(7/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3, x]`

output

```
-1/30*(Sqrt[Cos[c + d*x]]*Csc[c + d*x]^5*(156*A - 156*B + 280*A*Cos[c + d*x] - 280*B*Cos[c + d*x] - 312*A*Cos[c + d*x]^2 + 312*B*Cos[c + d*x]^2 - 280*A*Cos[c + d*x]^3 + 280*B*Cos[c + d*x]^3 + 180*A*Cos[c + d*x]^4 - 180*B*Cos[c + d*x]^4 + 60*A*Cos[c + d*x]^5 - 60*B*Cos[c + d*x]^5 - 26*A*Sin[c + d*x]^2 - 174*B*Sin[c + d*x]^2 - 280*B*Cos[c + d*x]*Sin[c + d*x]^2 - 20*B*Cos[c + d*x]^4*Sin[c + d*x]^2 - 65*A*Sin[c + d*x]^4 + 165*B*Sin[c + d*x]^4 - 5*(13*A - 33*B)*Hypergeometric2F1[1/4, 1/2, 5/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] + 280*B*Cos[c + d*x]*Hypergeometric2F1[3/4, 5/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] - 280*A*Cos[c + d*x]*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] + 280*B*Cos[c + d*x]*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] + 60*B*Sin[2*(c + d*x)]^2 + 15*B*Csc[c + d*x]*Sin[2*(c + d*x)]^3))/(a^3*d)
```

3.159.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3456, 27, 3042, 3456, 3042, 3456, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

3.159. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{7/2} \left(A+B \sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a \sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^{5/2}(c+dx)(7a(A-B)-a(3A-13B)\cos(c+dx))}{2(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cos^{5/2}(c+dx)(7a(A-B)-a(3A-13B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} (7a(A-B)-a(3A-13B)\sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right)a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^{3/2}(c+dx)(25a^2(A-2B)-3a^2(8A-23B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{10a(A-2B)\sin(c+dx)\cos^{5/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)} \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} (25a^2(A-2B)-3a^2(8A-23B)\sin\left(c+dx+\frac{\pi}{2}\right))}{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} dx}{3a^2} + \frac{10a(A-2B)\sin(c+dx)\cos^{5/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)} \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(7a^3(7A-17B)-5a^3(13A-33B)\cos(c+dx))}{a^2} dx}{3a^2} + \frac{7a^2(7A-17B)\sin(c+dx)\cos^{3/2}(c+dx)}{d(a\cos(c+dx)+a)} + \frac{10a(A-2B)\sin(c+dx)\cos^{5/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)} \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.159. $\int \frac{\cos^{7/2}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{3 \int \frac{\sqrt{\cos(c+dx)} (7a^3(7A-17B) - 5a^3(13A-33B) \cos(c+dx)) dx}{2a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)}}{3a^2} + \frac{10a(A-2B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} (7a^3(7A-17B) - 5a^3(13A-33B) \sin(c+dx+\frac{\pi}{2})) dx}{2a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)}}{3a^2} + \frac{10a(A-2B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3227

$$\frac{3 \left(\frac{7a^3(7A-17B) \int \sqrt{\cos(c+dx)} dx - 5a^3(13A-33B) \int \cos^{\frac{3}{2}}(c+dx) dx}{2a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)} \right)}{3a^2} + \frac{10a(A-2B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{3 \left(\frac{7a^3(7A-17B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^3(13A-33B) \int \sin(c+dx+\frac{\pi}{2})^{3/2} dx}{2a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)} \right)}{3a^2} + \frac{10a(A-2B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3115

$$\frac{3 \left(\frac{7a^3(7A-17B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^3(13A-33B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right)}{2a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)} \right)}{3a^2} + \frac{10a(A-2B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3042

3.159. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

$$\frac{\frac{3 \left(7a^3(7A-17B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx - 5a^3(13A-33B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{2 \sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) \right)}{2a^2}}{3a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{10a(A-2B)}{3d}$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3119

$$\frac{\frac{3 \left(\frac{14a^3(7A-17B)E\left(\frac{1}{2}(c+dx)|2\right)}{d} - 5a^3(13A-33B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{2 \sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) \right)}{2a^2}}{3a^2} + \frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{10a(A-2B)}{3d}$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3120

$$\frac{\frac{7a^2(7A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{d(a \cos(c+dx)+a)} + \frac{3 \left(\frac{14a^3(7A-17B)E\left(\frac{1}{2}(c+dx)|2\right)}{d} - 5a^3(13A-33B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + \frac{2 \sin(c+dx)\sqrt{\cos(c+dx)}}{3d} \right) \right)}{2a^2}}{3a^2} + \frac{10a(A-2B)}{3d}$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

input `Int[(Cos[c + d*x]^(7/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + (10*a*(A - 2*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + ((7*a^2*(7*A - 17*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])) + (3*((14*a^3*(7*A - 17*B)*EllipticE[(c + d*x)/2, 2])/d - 5*a^3*(13*A - 33*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/(2*a^2))/(3*a^2))/(10*a^2)`

3.159. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.159.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.159.
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.159.4 Maple [A] (verified)

Time = 15.41 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.12

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(-160B\left(\cos^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+348A\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+130A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)$

```
input int(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-160*B*cos(1
/2*d*x+1/2*c)^10+348*A*cos(1/2*d*x+1/2*c)^8+130*A*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2
))*cos(1/2*d*x+1/2*c)^5+294*A*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/
2))-468*B*cos(1/2*d*x+1/2*c)^8-330*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(
1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*
x+1/2*c)^5-714*B*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos
(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-578*A*cos
(1/2*d*x+1/2*c)^6+1058*B*cos(1/2*d*x+1/2*c)^6+264*A*cos(1/2*d*x+1/2*c)^4-4
74*B*cos(1/2*d*x+1/2*c)^4-37*A*cos(1/2*d*x+1/2*c)^2+47*B*cos(1/2*d*x+1/2*c
)^2+3*A-3*B)/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x
+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.159.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 478, normalized size of antiderivative = 2.18

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{2(20B\cos(dx+c)^3-3(29A-79B)\cos(dx+c)^2-2(73A-188B)\cos(dx+c)-65A+165B)\sqrt{\cos(dx+c)}}{(a+a\cos(c+dx))^3}$$

```
input integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm
m="fricas")
```

3.159. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

output `1/60*(2*(20*B*cos(d*x + c)^3 - 3*(29*A - 79*B)*cos(d*x + c)^2 - 2*(73*A - 188*B)*cos(d*x + c) - 65*A + 165*B)*sqrt(cos(d*x + c))*sin(d*x + c) - 5*(sqrt(2)*(-13*I*A + 33*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-13*I*A + 33*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-13*I*A + 33*I*B)*cos(d*x + c) + sqrt(2)*(-13*I*A + 33*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*(sqrt(2)*(13*I*A - 33*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(13*I*A - 33*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(13*I*A - 33*I*B)*cos(d*x + c) + sqrt(2)*(13*I*A - 33*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*(sqrt(2)*(-7*I*A + 17*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-7*I*A + 17*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-7*I*A + 17*I*B)*cos(d*x + c) + sqrt(2)*(-7*I*A + 17*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*(sqrt(2)*(7*I*A - 17*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(7*I*A - 17*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(7*I*A - 17*I*B)*cos(d*x + c) + sqrt(2)*(7*I*A - 17*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/((a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.159.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)`

output `Timed out`

3.159.7 Maxima [F]

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{7}{2}}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^3, x)`

3.159. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.159.8 Giac [F]

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{7}{2}}}{(a\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^3, x)`

3.159.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{\cos(c+dx)^{7/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(7/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output `int((cos(c + d*x)^(7/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3, x)`

3.160
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.160.1 Optimal result 1557
 3.160.2 Mathematica [C] (verified) 1558
 3.160.3 Rubi [A] (verified) 1558
 3.160.4 Maple [B] (verified) 1562
 3.160.5 Fricas [C] (verification not implemented) 1563
 3.160.6 Sympy [F(-1)] 1563
 3.160.7 Maxima [F] 1564
 3.160.8 Giac [F] 1564
 3.160.9 Mupad [F(-1)] 1564

3.160.1 Optimal result

Integrand size = 33, antiderivative size = 188

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = -\frac{(9A-49B)E(\frac{1}{2}(c+dx)|2)}{10a^3d} + \frac{(3A-13B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{6a^3d} + \frac{(A-B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(3A-8B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} + \frac{(3A-13B) \sqrt{\cos(c+dx)} \sin(c+dx)}{6d(a^3+a^3 \cos(c+dx))}$$

output

```
-1/10*(9*A-49*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE
(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/6*(3*A-13*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/5*(A
-B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(3*A-8*B)*cos(d*
x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^2+1/6*(3*A-13*B)*sin(d*x+c)*cos
(d*x+c)^(1/2)/d/(a^3+a^3*cos(d*x+c))
```

3.160.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.05 (sec) , antiderivative size = 400, normalized size of antiderivative = 2.13

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{\sqrt{\cos(c+dx)} \operatorname{csc}^5(c+dx) \left(-252A + 252B - 360A\cos(c+dx) + 360B\cos(c+dx) + 504A\cos^2(c+dx) \right)}{a^3}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3, x]`

output

```
-1/210*(Sqrt[Cos[c + d*x]]*Csc[c + d*x]^5*(-252*A + 252*B - 360*A*Cos[c + d*x] + 360*B*Cos[c + d*x] + 504*A*Cos[c + d*x]^2 - 504*B*Cos[c + d*x]^2 + 420*A*Cos[c + d*x]^3 - 420*B*Cos[c + d*x]^3 - 420*A*Cos[c + d*x]^4 + 420*B*Cos[c + d*x]^4 + 42*A*Sin[c + d*x]^2 + 658*B*Sin[c + d*x]^2 - 420*B*Cos[c + d*x]^3*Sin[c + d*x]^2 + 105*A*Sin[c + d*x]^4 - 455*B*Sin[c + d*x]^4 + 35*(3*A - 13*B)*Hypergeometric2F1[1/4, 1/2, 5/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] - 1120*B*Cos[c + d*x]*Hypergeometric2F1[3/4, 5/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] + 360*A*Cos[c + d*x]*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] - 360*B*Cos[c + d*x]*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]^4*Sqrt[Sin[c + d*x]^2] + 560*B*Sin[c + d*x]*Sin[2*(c + d*x)] - 210*B*Sin[2*(c + d*x)]^2))/(a^3*d)
```

3.160.3 Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3456, 27, 3042, 3456, 3042, 3456, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

3.160. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\cos^{3/2}(c+dx)(5a(A-B)-a(A-11B)\cos(c+dx))}{2(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cos^{3/2}(c+dx)(5a(A-B)-a(A-11B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} (5a(A-B)-a(A-11B)\sin\left(c+dx+\frac{\pi}{2}\right))}{\left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a^2(3A-8B)-a^2(6A-41B)\cos(c+dx))}{\cos(c+dx)a+a} dx}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{3/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)} \\
& \quad \frac{5d(a\cos(c+dx)+a)^3}{} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}(3a^2(3A-8B)-a^2(6A-41B)\sin\left(c+dx+\frac{\pi}{2}\right))}{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} dx}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{3/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)} \\
& \quad \frac{5d(a\cos(c+dx)+a)^3}{} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{5a^3(3A-13B)-3a^3(9A-49B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx}{a^2} + \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} + \frac{2a(3A-8B)\sin(c+dx)\cos^{3/2}(c+dx)}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)} \\
& \quad \frac{5d(a\cos(c+dx)+a)^3}{} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.160. $\int \frac{\cos^{5/2}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{\int \frac{5a^3(3A-13B)-3a^3(9A-49B)\cos(c+dx)}{2a^2} dx + \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{5d(a\cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{\int \frac{5a^3(3A-13B)-3a^3(9A-49B)\sin(c+dx+\frac{\pi}{2})}{2a^2} dx + \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{5d(a\cos(c+dx)+a)^3}$$

↓ 3227

$$\frac{5a^3(3A-13B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^3(9A-49B)\int \sqrt{\cos(c+dx)} dx + \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{5d(a\cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{5a^3(3A-13B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^3(9A-49B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{5d(a\cos(c+dx)+a)^3}$$

↓ 3119

$$\frac{5a^3(3A-13B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6a^3(9A-49B)E(\frac{1}{2}(c+dx)|2)}{d} + \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{5d(a\cos(c+dx)+a)^3}$$

↓ 3120

3.160. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{\frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} + \frac{10a^3(3A-13B)\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - 6a^3(9A-49B)E\left(\frac{1}{2}(c+dx)|2\right)}{3a^2} + \frac{2a(3A-8B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3d(a\cos(c+dx)+a)^2}}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3}$$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((2*a*(3*A - 8*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (((-6*a^3*(9*A - 49*B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(3*A - 13*B)*EllipticF[(c + d*x)/2, 2])/d)/(2*a^2) + (5*a^2*(3*A - 13*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))/(3*a^2))/(10*a^2)`

3.160.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`


```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &&
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

3.160.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(224) = 448$.

Time = 15.04 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.40

method	result
default	$-\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(108A\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 30A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 1\right)F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$

```
input int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

```
output -1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(108*A*cos(1
/2*d*x+1/2*c)^8+30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2
+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+54*A*
cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2
+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-348*B*cos(1/2*d*x+1/2*c)^8
-130*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*Elli
pticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5-294*B*cos(1/2*d*x+1
/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*Elli
pticE(cos(1/2*d*x+1/2*c),2^(1/2))-198*A*cos(1/2*d*x+1/2*c)^6+578*B*cos(1/
2*d*x+1/2*c)^6+114*A*cos(1/2*d*x+1/2*c)^4-264*B*cos(1/2*d*x+1/2*c)^4-27*A*
cos(1/2*d*x+1/2*c)^2+37*B*cos(1/2*d*x+1/2*c)^2+3*A-3*B)/a^3/cos(1/2*d*x+1/
2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/
2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

$$3.160. \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.160.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 467, normalized size of antiderivative = 2.48

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{2(3(9A-29B)\cos(dx+c)^2 + 2(18A-73B)\cos(dx+c) + 15A-65B)\sqrt{\cos(dx+c)}\sin(dx+c) -$$

```
input integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="fricas")
```

```
output 1/60*(2*(3*(9*A - 29*B)*cos(d*x + c)^2 + 2*(18*A - 73*B)*cos(d*x + c) + 15
*A - 65*B)*sqrt(cos(d*x + c))*sin(d*x + c) - 5*(sqrt(2)*(3*I*A - 13*I*B)*c
os(d*x + c)^3 + 3*sqrt(2)*(3*I*A - 13*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(3*I
*A - 13*I*B)*cos(d*x + c) + sqrt(2)*(3*I*A - 13*I*B))*weierstrassPInverse(
-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*(sqrt(2)*(-3*I*A + 13*I*B)*cos(d
*x + c)^3 + 3*sqrt(2)*(-3*I*A + 13*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-3*I*A
+ 13*I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A + 13*I*B))*weierstrassPInverse(-
4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*(sqrt(2)*(9*I*A - 49*I*B)*cos(d*x
+ c)^3 + 3*sqrt(2)*(9*I*A - 49*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(9*I*A - 4
9*I*B)*cos(d*x + c) + sqrt(2)*(9*I*A - 49*I*B))*weierstrassZeta(-4, 0, wei
erstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*(sqrt(2)*(-9*I
*A + 49*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-9*I*A + 49*I*B)*cos(d*x + c)^2 +
3*sqrt(2)*(-9*I*A + 49*I*B)*cos(d*x + c) + sqrt(2)*(-9*I*A + 49*I*B))*wei
erstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x +
c))))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c
) + a^3*d)
```

3.160.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)
```

```
output Timed out
```

3.160. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

3.160.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(a\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^3, x)`

3.160.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(a\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^3, x)`

3.160.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3, x)`

3.160. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

3.161 $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.161.1 Optimal result 1565
 3.161.2 Mathematica [C] (verified) 1566
 3.161.3 Rubi [A] (verified) 1567
 3.161.4 Maple [B] (verified) 1571
 3.161.5 Fricas [C] (verification not implemented) 1571
 3.161.6 Sympy [F(-1)] 1572
 3.161.7 Maxima [F] 1572
 3.161.8 Giac [F] 1573
 3.161.9 Mupad [F(-1)] 1573

3.161.1 Optimal result

Integrand size = 33, antiderivative size = 180

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = -\frac{(A+9B)E(\frac{1}{2}(c+dx)|2)}{10a^3d} + \frac{(A+3B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{6a^3d} + \frac{(A-B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(A-6B) \sqrt{\cos(c+dx)} \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} + \frac{(A+9B) \sqrt{\cos(c+dx)} \sin(c+dx)}{10d(a^3+a^3 \cos(c+dx))}$$

output

```
-1/10*(A+9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/6*(A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/5*(A-B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^3+1/15*(A-6*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^2+1/10*(A+9*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a^3+a^3*cos(d*x+c))
```

3.161.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.47 (sec) , antiderivative size = 845, normalized size of antiderivative = 4.69

$$\begin{aligned}
& \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx \\
&= -\frac{(A-B)\sqrt{\cos(c+dx)}\csc(c+dx)}{15a^3d(1-\cos(c+dx))(1+\cos(c+dx))} + \frac{4B\sqrt{\cos(c+dx)}\csc(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&+ \frac{4B\cos^{\frac{3}{2}}(c+dx)\csc(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} - \frac{2B\cos^{\frac{5}{2}}(c+dx)\csc(c+dx)}{a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&+ \frac{2(A-B)\sqrt{\cos(c+dx)}\csc^3(c+dx)}{5a^3d(1-\cos(c+dx))(1+\cos(c+dx))} + \frac{4(A-B)\cos^{\frac{3}{2}}(c+dx)\csc^3(c+dx)}{21a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&- \frac{6(A-B)\cos^{\frac{5}{2}}(c+dx)\csc^3(c+dx)}{5a^3d(1-\cos(c+dx))(1+\cos(c+dx))} + \frac{2(A-B)\cos^{\frac{7}{2}}(c+dx)\csc^3(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&- \frac{(A-B)\sqrt{\cos(c+dx)}\sin(c+dx)}{6a^3d(1-\cos(c+dx))(1+\cos(c+dx))} - \frac{2B\sqrt{\cos(c+dx)}\sin(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&- \frac{(A-B)\sqrt{\cos(c+dx)}\operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{6a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&- \frac{2B\sqrt{\cos(c+dx)}\operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&- \frac{4B\cos^{\frac{3}{2}}(c+dx)\operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{5}{2}, \frac{7}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} \\
&- \frac{4(A-B)\cos^{\frac{3}{2}}(c+dx)\operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{7}{2}, \frac{7}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{21a^3d(1-\cos(c+dx))(1+\cos(c+dx))}
\end{aligned}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3, x]`

output

```

-1/15*((A - B)*Sqrt[Cos[c + d*x]]*Csc[c + d*x])/(a^3*d*(1 - Cos[c + d*x])*
(1 + Cos[c + d*x])) + (4*B*Sqrt[Cos[c + d*x]]*Csc[c + d*x])/(3*a^3*d*(1 -
Cos[c + d*x])*(1 + Cos[c + d*x])) + (4*B*Cos[c + d*x]^(3/2)*Csc[c + d*x])/
(3*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x])) - (2*B*Cos[c + d*x]^(5/2)*
Csc[c + d*x])/(a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x])) + (2*(A - B)*S
qrt[Cos[c + d*x]]*Csc[c + d*x]^3)/(5*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c +
d*x])) + (4*(A - B)*Cos[c + d*x]^(3/2)*Csc[c + d*x]^3)/(21*a^3*d*(1 - Cos
[c + d*x])*(1 + Cos[c + d*x])) - (6*(A - B)*Cos[c + d*x]^(5/2)*Csc[c + d*x
]^3)/(5*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x])) + (2*(A - B)*Cos[c +
d*x]^(7/2)*Csc[c + d*x]^3)/(3*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x]))
- ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(6*a^3*d*(1 - Cos[c + d*x])*(
1 + Cos[c + d*x])) - (2*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*a^3*d*(1 - C
os[c + d*x])*(1 + Cos[c + d*x])) - ((A - B)*Sqrt[Cos[c + d*x]]*Hypergeomet
ric2F1[1/4, 1/2, 5/4, Cos[c + d*x]^2]*Sin[c + d*x]*Sqrt[Sin[c + d*x]^2])/(
6*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x])) - (2*B*Sqrt[Cos[c + d*x]]*H
ypergeometric2F1[1/4, 1/2, 5/4, Cos[c + d*x]^2]*Sin[c + d*x]*Sqrt[Sin[c +
d*x]^2])/(3*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x])) - (4*B*Cos[c + d*
x]^(3/2)*Hypergeometric2F1[3/4, 5/2, 7/4, Cos[c + d*x]^2]*Sin[c + d*x]*Sqr
t[Sin[c + d*x]^2])/(3*a^3*d*(1 - Cos[c + d*x])*(1 + Cos[c + d*x])) - (4*(A
- B)*Cos[c + d*x]^(3/2)*Hypergeometric2F1[3/4, 7/2, 7/4, Cos[c + d*x]^...

```

3.161.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3456, 27, 3042, 3456, 3042, 3457, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx \\
 & \quad \downarrow \text{3456} \\
 & \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+a(A+9B)\cos(c+dx))}{2(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3}
 \end{aligned}$$

3.161. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+a(A+9B)\cos(c+dx))}{(\cos(c+dx)a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 3042 \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)+a(A+9B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 3456 \\
& \frac{\frac{\int \frac{(A-6B)a^2+(4A+21B)\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx}{3a^2} + \frac{2a(A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 3042 \\
& \frac{\frac{\int \frac{(A-6B)a^2+(4A+21B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} + \frac{2a(A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 3457 \\
& \frac{\frac{\frac{\int \frac{5a^3(A+3B)-3a^3(A+9B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx}{a^2} + \frac{3a^2(A+9B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 27 \\
& \frac{\frac{\frac{\int \frac{5a^3(A+3B)-3a^3(A+9B)\cos(c+dx)}{\sqrt{\cos(c+dx)}} dx}{2a^2} + \frac{3a^2(A+9B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 3042 \\
& \frac{\frac{\int \frac{5a^3(A+3B)-3a^3(A+9B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} + \frac{3a^2(A+9B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{3a^2} + \frac{2a(A-6B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}}{10a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d(a\cos(c+dx)+a)^3} \\
& \downarrow 3227
\end{aligned}$$

3.161. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{\frac{5a^3(A+3B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^3(A+9B) \int \sqrt{\cos(c+dx)} dx}{2a^2} + \frac{3a^2(A+9B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A-6B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2}}{3a^2} + \frac{10a^2}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{\frac{5a^3(A+3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^3(A+9B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{2a^2} + \frac{3a^2(A+9B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A-6B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2}}{3a^2} + \frac{10a^2}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3119

$$\frac{\frac{5a^3(A+3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6a^3(A+9B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} + \frac{3a^2(A+9B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A-6B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2}}{3a^2} + \frac{10a^2}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

↓ 3120

$$\frac{\frac{3a^2(A+9B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{\frac{10a^3(A+3B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} - \frac{6a^3(A+9B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2}}{3a^2} + \frac{2a(A-6B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2}}{3a^2} + \frac{10a^2}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d(a \cos(c+dx)+a)^3}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + ((2*a*(A - 6*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (((-6*a^3*(A + 9*B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(A + 3*B)*EllipticF[(c + d*x)/2, 2])/d)/(2*a^2) + (3*a^2*(A + 9*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))/(3*a^2))/(10*a^2)`

3.161. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.161.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.161.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.161.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(216) = 432$.

Time = 5.33 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.51

method	result
default	$-\frac{\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(12A\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+10A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)\right)}{\dots}$

input `int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)`

output
$$\begin{aligned} & -1/60*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(12*A*\cos(1/2*d*x+1/2*c)^8+10*A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\cos(1/2*d*x+1/2*c)^5+6*A*\cos(1/2*d*x+1/2*c)^5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})+108*B*\cos(1/2*d*x+1/2*c)^8+30*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\cos(1/2*d*x+1/2*c)^5+54*B*\cos(1/2*d*x+1/2*c)^5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})-2*A*\cos(1/2*d*x+1/2*c)^6-198*B*\cos(1/2*d*x+1/2*c)^6-24*A*\cos(1/2*d*x+1/2*c)^4+114*B*\cos(1/2*d*x+1/2*c)^4+17*A*\cos(1/2*d*x+1/2*c)^2-27*B*\cos(1/2*d*x+1/2*c)^2-3*A+3*B)/a^3/\cos(1/2*d*x+1/2*c)^5/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.161.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.58

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= \frac{2(3(A+9B)\cos(dx+c)^2+2(7A+18B)\cos(dx+c)+5A+15B)\sqrt{\cos(dx+c)}\sin(dx+c)-5(\sqrt{\cos(dx+c)}\sin(dx+c))^2}{(a+a\cos(c+dx))^3}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm
m="fricas")`

3.161.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

output `1/60*(2*(3*(A + 9*B)*cos(d*x + c)^2 + 2*(7*A + 18*B)*cos(d*x + c) + 5*A + 15*B)*sqrt(cos(d*x + c))*sin(d*x + c) - 5*(sqrt(2)*(I*A + 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(I*A + 3*I*B)*cos(d*x + c) + sqrt(2)*(I*A + 3*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*(sqrt(2)*(-I*A - 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-I*A - 3*I*B)*cos(d*x + c) + sqrt(2)*(-I*A - 3*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*(sqrt(2)*(I*A + 9*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(I*A + 9*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(I*A + 9*I*B)*cos(d*x + c) + sqrt(2)*(I*A + 9*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*(sqrt(2)*(-I*A - 9*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-I*A - 9*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-I*A - 9*I*B)*cos(d*x + c) + sqrt(2)*(-I*A - 9*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.161.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3,x)`

output `Timed out`

3.161.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^3, x)`

3.161. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$

3.161.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(a\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^3, x)`

3.161.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3, x)`

3.162
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx$$

3.162.1 Optimal result 1574
 3.162.2 Mathematica [C] (verified) 1575
 3.162.3 Rubi [A] (verified) 1576
 3.162.4 Maple [B] (verified) 1580
 3.162.5 Fracas [C] (verification not implemented) 1580
 3.162.6 Sympy [F(-1)] 1581
 3.162.7 Maxima [F] 1581
 3.162.8 Giac [F] 1582
 3.162.9 Mupad [F(-1)] 1582

3.162.1 Optimal result

Integrand size = 33, antiderivative size = 178

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^3} dx = \frac{(A-B)E(\frac{1}{2}(c+dx)|2)}{10a^3d} + \frac{(A+B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{6a^3d} + \frac{(A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{5d(a+a \cos(c+dx))^3} + \frac{(A+4B)\sqrt{\cos(c+dx)} \sin(c+dx)}{15ad(a+a \cos(c+dx))^2} - \frac{(A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{10d(a^3+a^3 \cos(c+dx))}$$

```
output 1/10*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/6*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/5*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^3+1/15*(A+4*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^2-1/10*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a^3+a^3*cos(d*x+c))
```

3.162.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.35 (sec) , antiderivative size = 664, normalized size of antiderivative = 3.73

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

$$= -\frac{(A-B)\sqrt{\cos(c+dx)}\csc(c+dx)}{15a^3d(1-\cos(c+dx))(1+\cos(c+dx))} + \frac{2B\sqrt{\cos(c+dx)}\csc(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

$$- \frac{2B\cos^{\frac{3}{2}}(c+dx)\csc(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))} + \frac{2(A-B)\sqrt{\cos(c+dx)}\csc^3(c+dx)}{5a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

$$- \frac{6(A-B)\cos^{\frac{3}{2}}(c+dx)\csc^3(c+dx)}{7a^3d(1-\cos(c+dx))(1+\cos(c+dx))} + \frac{2(A-B)\cos^{\frac{5}{2}}(c+dx)\csc^3(c+dx)}{5a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

$$- \frac{(A-B)\sqrt{\cos(c+dx)}\sin(c+dx)}{6a^3d(1-\cos(c+dx))(1+\cos(c+dx))} - \frac{B\sqrt{\cos(c+dx)}\sin(c+dx)}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

$$- \frac{(A-B)\sqrt{\cos(c+dx)}\operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{6a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

$$- \frac{B\sqrt{\cos(c+dx)}\operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{3a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

$$+ \frac{4(A-B)\cos^{\frac{3}{2}}(c+dx)\operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{7}{2}, \frac{7}{4}, \cos^2(c+dx)\right)\sin(c+dx)\sqrt{\sin^2(c+dx)}}{21a^3d(1-\cos(c+dx))(1+\cos(c+dx))}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3, x]`

output

$$\begin{aligned}
& -1/15*((A - B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Csc}[c + d*x])/(a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) + (2*B*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Csc}[c + d*x])/(3*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) - (2*B*\text{Cos}[c + d*x]^{3/2}*\text{Csc}[c + d*x])/(3*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) + (2*(A - B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Csc}[c + d*x]^3)/(5*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) - (6*(A - B)*\text{Cos}[c + d*x]^{3/2}*\text{Csc}[c + d*x]^3)/(7*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) + (2*(A - B)*\text{Cos}[c + d*x]^{5/2}*\text{Csc}[c + d*x]^3)/(5*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) - ((A - B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(6*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) - (B*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(3*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) - ((A - B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Hypergeometric2F1}[1/4, 1/2, 5/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(6*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) - (B*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Hypergeometric2F1}[1/4, 1/2, 5/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(3*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x])) + (4*(A - B)*\text{Cos}[c + d*x]^{3/2}*\text{Hypergeometric2F1}[3/4, 7/2, 7/4, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x]*\text{Sqrt}[\text{Sin}[c + d*x]^2])/(21*a^3*d*(1 - \text{Cos}[c + d*x])*(1 + \text{Cos}[c + d*x]))
\end{aligned}$$

3.162.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3456, 27, 3042, 3457, 3042, 3457, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{a(A-B)+a(3A+7B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.162. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{a(A-B)+a(3A+7B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(A-B)+a(3A+7B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^2} dx}{10a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{(4A+B)a^2+(A+4B)\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx}{3a^2} + \frac{2a(A+4B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(4A+B)a^2+(A+4B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} + \frac{2a(A+4B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{5(A+B)a^3+3(A-B)\cos(c+dx)a^3}{2\sqrt{\cos(c+dx)}} dx}{a^2} - \frac{3a^2(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} + \frac{2a(A+4B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{5(A+B)a^3+3(A-B)\cos(c+dx)a^3}{\sqrt{\cos(c+dx)}} dx}{2a^2} - \frac{3a^2(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} + \frac{2a(A+4B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{5(A+B)a^3+3(A-B)\sin(c+dx+\frac{\pi}{2})a^3}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{3a^2(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} + \frac{2a(A+4B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2} + \\
& \quad \frac{10a^2}{5d(a\cos(c+dx)+a)^3} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{5d(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3227}
\end{aligned}$$

3.162. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$

$$\frac{5a^3(A+B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^3(A-B) \int \sqrt{\cos(c+dx)} dx}{2a^2} - \frac{3a^2(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A+4B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx)+a)^3}$$

↓ 3042

$$\frac{5a^3(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a^3(A-B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{3a^2(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A+4B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx)+a)^3}$$

↓ 3119

$$\frac{5a^3(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^3(A-B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} - \frac{3a^2(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A+4B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx)+a)^3}$$

↓ 3120

$$\frac{\frac{10a^3(A+B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6a^3(A-B)E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2} - \frac{3a^2(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx)+a)} + \frac{2a(A+4B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx)+a)^2} +$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx)+a)^3}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^3,x]`

output `((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(5*d*(a + a*Cos[c + d*x])^3) + (2*a*(A + 4*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (((6*a^3*(A - B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(A + B)*EllipticF[(c + d*x)/2, 2])/d)/(2*a^2) - (3*a^2*(A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))/(3*a^2))/(10*a^2)`

3.162.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.162.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(214) = 428$.

Time = 5.43 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.53

method	result
default	$\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(12A\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-10A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\right)}{\dots}$

```
input int((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*A*cos(1/2
*d*x+1/2*c)^8-10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+6*A*cos
(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-12*B*cos(1/2*d*x+1/2*c)^8-10*
B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF
(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5-6*B*cos(1/2*d*x+1/2*c)^5
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(
cos(1/2*d*x+1/2*c),2^(1/2))-22*A*cos(1/2*d*x+1/2*c)^6+2*B*cos(1/2*d*x+1/2*
c)^6+6*A*cos(1/2*d*x+1/2*c)^4+24*B*cos(1/2*d*x+1/2*c)^4+7*A*cos(1/2*d*x+1/
2*c)^2-17*B*cos(1/2*d*x+1/2*c)^2-3*A+3*B)/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin
(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/
2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.162.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.61

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx =$$

$$\frac{2(3(A-B)\cos(dx+c)^2+2(2A-7B)\cos(dx+c)-5A-5B)\sqrt{\cos(dx+c)}\sin(dx+c)+5(\sqrt{\cos(dx+c)}\sin(dx+c))^2}{\dots}$$

```
input integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^3,x,algorith
m="fracas")
```

$$3.162. \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

output
$$\begin{aligned} & -1/60*(2*(3*(A - B)*\cos(d*x + c)^2 + 2*(2*A - 7*B)*\cos(d*x + c) - 5*A - 5*B)*\sqrt{\cos(d*x + c)}*\sin(d*x + c) + 5*(\sqrt{2}*(I*A + I*B)*\cos(d*x + c)^3 \\ & + 3*\sqrt{2}*(I*A + I*B)*\cos(d*x + c)^2 + 3*\sqrt{2}*(I*A + I*B)*\cos(d*x + c) + \sqrt{2}*(I*A + I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) \\ & + 5*(\sqrt{2}*(-I*A - I*B)*\cos(d*x + c)^3 + 3*\sqrt{2}*(-I*A - I*B)*\cos(d*x + c)^2 + 3*\sqrt{2}*(-I*A - I*B)*\cos(d*x + c) + \sqrt{2}*(-I*A - I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) \\ & + 3*(\sqrt{2}*(-I*A + I*B)*\cos(d*x + c)^3 + 3*\sqrt{2}*(-I*A + I*B)*\cos(d*x + c)^2 + 3*\sqrt{2}*(-I*A + I*B)*\cos(d*x + c) + \sqrt{2}*(-I*A + I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) \\ & + 3*(\sqrt{2}*(I*A - I*B)*\cos(d*x + c)^3 + 3*\sqrt{2}*(I*A - I*B)*\cos(d*x + c)^2 + 3*\sqrt{2}*(I*A - I*B)*\cos(d*x + c) + \sqrt{2}*(I*A - I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) \\ &)/(a^3*d*\cos(d*x + c)^3 + 3*a^3*d*\cos(d*x + c)^2 + 3*a^3*d*\cos(d*x + c) + a^3*d) \end{aligned}$$

3.162.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)**(1/2)/(a+a*cos(d*x+c))**3,x)`

output Timed out

3.162.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\cos(dx + c)}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(a*cos(d*x + c) + a)^3, x)`

3.162.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(a\cos(dx+c)+a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(a*cos(d*x + c) + a)^3, x)`

3.162.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3,x)`

output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^3, x)`

3.163
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^3} dx$$

3.163.1 Optimal result 1583
 3.163.2 Mathematica [C] (warning: unable to verify) 1584
 3.163.3 Rubi [A] (verified) 1584
 3.163.4 Maple [B] (verified) 1588
 3.163.5 Fricas [C] (verification not implemented) 1589
 3.163.6 Sympy [F(-1)] 1589
 3.163.7 Maxima [F(-1)] 1590
 3.163.8 Giac [F] 1590
 3.163.9 Mupad [F(-1)] 1590

3.163.1 Optimal result

Integrand size = 33, antiderivative size = 182

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \frac{(9A + B)E(\frac{1}{2}(c + dx) | 2)}{10a^3d} + \frac{(3A + B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{6a^3d} - \frac{(A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{5d(a + a \cos(c + dx))^3} - \frac{(6A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{15ad(a + a \cos(c + dx))^2} - \frac{(9A + B)\sqrt{\cos(c + dx)} \sin(c + dx)}{10d(a^3 + a^3 \cos(c + dx))}$$

```
output 1/10*(9*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/6*(3*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d-1/5*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^3-1/15*(6*A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^2-1/10*(9*A+B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a^3+a^3*cos(d*x+c))
```

3.163.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.87 (sec) , antiderivative size = 1029, normalized size of antiderivative = 5.65

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^3),x]`

output `(-2*A*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(d*(a + a*Cos[c + d*x])^3*Sqrt[1 + Cot[c]^2]) - (2*B*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3*d*(a + a*Cos[c + d*x])^3*Sqrt[1 + Cot[c]^2]) + (Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*((-4*(9*A + B)*Csc[c])/(5*d) - (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^5*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(5*d) - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(6*A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(15*d) - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(9*A*Sin[(d*x)/2] + B*Sin[(d*x)/2]))/(5*d) - (4*(6*A - B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(15*d) - (2*(A - B)*Sec[c/2 + (d*x)/2]^4*Tan[c/2])/(5*d)))/(a + a*Cos[c + d*x])^3 - (9*A*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*Sec[c/2]*(HypergeometricPFQ[{-1/2, -1/4}, {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2]*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Tan[c]^2])*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/Sqrt[1 + Tan[c]^2] + (2*Cos[c]^2*Cos[d*x + ArcTan[Tan[c]]]*Sqrt[1 + Ta...`

3.163.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3457, 27, 3042, 3457, 3042, 3457, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.163. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a \sin(c + dx + \frac{\pi}{2}) + a)^3} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a(9A+B) - 3a(A-B) \cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(9A+B) - 3a(A-B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^2} dx}{10a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(9A+B) - 3a(A-B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(\sin(c+dx + \frac{\pi}{2})a+a)^2} dx}{10a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a^2(21A+4B) - a^2(6A-B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)} dx}{3a^2} - \frac{2a(6A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx) + a)^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(21A+4B) - a^2(6A-B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(\sin(c+dx + \frac{\pi}{2})a+a)} dx}{3a^2} - \frac{2a(6A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx) + a)^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx) + a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{5(3A+B)a^3 + 3(9A+B) \cos(c+dx)a^3}{2\sqrt{\cos(c+dx)}} dx}{a^2} - \frac{3a^2(9A+B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d(a \cos(c+dx) + a)} - \frac{2a(6A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d(a \cos(c+dx) + a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{10a^2}{5d(a \cos(c+dx) + a)^3} \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{5d(a \cos(c+dx) + a)^3}
\end{aligned}$$

3.163. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^3} dx$

$$\frac{\int \frac{5(3A+B)a^3 + 3(9A+B)\cos(c+dx)a^3}{\sqrt{\cos(c+dx)}} dx}{3a^2} - \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} - \frac{2a(6A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\sqrt{\cos(c+dx)}$$

↓ 3042

$$\frac{\int \frac{5(3A+B)a^3 + 3(9A+B)\sin(c+dx+\frac{\pi}{2})a^3}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3a^2} - \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} - \frac{2a(6A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\sqrt{\cos(c+dx)}$$

↓ 3227

$$\frac{5a^3(3A+B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^3(9A+B)\int \sqrt{\cos(c+dx)} dx}{3a^2} - \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} - \frac{2a(6A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\sqrt{\cos(c+dx)}$$

↓ 3042

$$\frac{5a^3(3A+B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a^3(9A+B)\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{3a^2} - \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} - \frac{2a(6A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\sqrt{\cos(c+dx)}$$

↓ 3119

$$\frac{5a^3(3A+B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^3(9A+B)E(\frac{1}{2}(c+dx)|2)}{d}}{3a^2} - \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)} - \frac{2a(6A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\sqrt{\cos(c+dx)}$$

↓ 3120

3.163. $\int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a\cos(c+dx))^3} dx$

$$\frac{\frac{10a^3(3A+B)\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} + \frac{6a^3(9A+B)E\left(\frac{1}{2}(c+dx), 2\right)}{d} - \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a\cos(c+dx)+a)}}{\frac{2a^2}{3a^2}} - \frac{2a(6A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2}{5d(a\cos(c+dx)+a)^3} (A-B)\sin(c+dx)\sqrt{\cos(c+dx)}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^3), x]`

output `-1/5*((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^3) + ((-2*a*(6*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Cos[c + d*x])^2) + (((6*a^3*(9*A + B)*EllipticE[(c + d*x)/2, 2])/d + (10*a^3*(3*A + B)*EllipticF[(c + d*x)/2, 2])/d)/(2*a^2) - (3*a^2*(9*A + B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x]))/(3*a^2))/(10*a^2)`

3.163.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.163.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(218) = 436$.

Time = 5.20 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.48

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(108A\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 30A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 1F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right)$

input `int((A+B*cos(d*x+c))/(a*cos(d*x+c)*a)^3/cos(d*x+c)^(1/2),x,method=_RETURNV ERBOSE)`

output `1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(108*A*cos(1/2*d*x+1/2*c)^8-30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+54*A*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+12*B*cos(1/2*d*x+1/2*c)^8-10*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+6*B*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-138*A*cos(1/2*d*x+1/2*c)^6-22*B*cos(1/2*d*x+1/2*c)^6+24*A*cos(1/2*d*x+1/2*c)^4+6*B*cos(1/2*d*x+1/2*c)^4+3*A*cos(1/2*d*x+1/2*c)^2+7*B*cos(1/2*d*x+1/2*c)^2+3*A-3*B)/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.163.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.55

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \frac{2(3(9A + B) \cos(dx + c)^2 + 2(33A + 2B) \cos(dx + c) + 45A - 5B) \sqrt{\cos(dx + c)} \sin(dx + c) + 5$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/60*(2*(3*(9*A + B)*cos(d*x + c)^2 + 2*(33*A + 2*B)*cos(d*x + c) + 45*A - 5*B)*sqrt(cos(d*x + c))*sin(d*x + c) + 5*(sqrt(2)*(3*I*A + I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(3*I*A + I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(3*I*A + I*B)*cos(d*x + c) + sqrt(2)*(3*I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(-3*I*A - I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-3*I*A - I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-3*I*A - I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(sqrt(2)*(-9*I*A - I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-9*I*A - I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-9*I*A - I*B)*cos(d*x + c) + sqrt(2)*(-9*I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(sqrt(2)*(9*I*A + I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(9*I*A + I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(9*I*A + I*B)*cos(d*x + c) + sqrt(2)*(9*I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.163.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3/cos(d*x+c)**(1/2),x)`

output `Timed out`

3.163. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^3} dx$

3.163.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/cos(d*x+c)^(1/2),x, algorithm m="maxima")`

output `Timed out`

3.163.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/cos(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sqrt(cos(d*x + c))), x)`

3.163.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^3), x)`

3.164
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$$

3.164.1 Optimal result 1591
 3.164.2 Mathematica [C] (warning: unable to verify) 1592
 3.164.3 Rubi [A] (verified) 1592
 3.164.4 Maple [B] (warning: unable to verify) 1597
 3.164.5 Fricas [C] (verification not implemented) 1598
 3.164.6 Sympy [F(-1)] 1598
 3.164.7 Maxima [F(-2)] 1599
 3.164.8 Giac [F] 1599
 3.164.9 Mupad [F(-1)] 1599

3.164.1 Optimal result

Integrand size = 33, antiderivative size = 221

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = -\frac{(49A - 9B)E(\frac{1}{2}(c + dx)|2)}{10a^3d} - \frac{(13A - 3B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{6a^3d} + \frac{(49A - 9B) \sin(c + dx)}{10a^3d\sqrt{\cos(c + dx)}} - \frac{(A - B) \sin(c + dx)}{5d\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^3} - \frac{(8A - 3B) \sin(c + dx)}{15ad\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^2} - \frac{(13A - 3B) \sin(c + dx)}{6d\sqrt{\cos(c + dx)}(a^3 + a^3 \cos(c + dx))}$$

output

```
-1/10*(49*A-9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE
(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d-1/6*(13*A-3*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/10*(
49*A-9*B)*sin(d*x+c)/a^3/d/cos(d*x+c)^(1/2)-1/5*(A-B)*sin(d*x+c)/d/(a+a*co
s(d*x+c))^3/cos(d*x+c)^(1/2)-1/15*(8*A-3*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c)
)^2/cos(d*x+c)^(1/2)-1/6*(13*A-3*B)*sin(d*x+c)/d/(a^3+a^3*cos(d*x+c))/cos(
d*x+c)^(1/2)
```

3.164.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.92 (sec) , antiderivative size = 1069, normalized size of antiderivative = 4.84

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^3),x]`

output `(26*A*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3*d*(a + a*Cos[c + d*x])^3*Sqrt[1 + Cot[c]^2]) - (2*B*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2]*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(d*(a + a*Cos[c + d*x])^3*Sqrt[1 + Cot[c]^2]) + (Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*((2*(20*A + 29*A*Cos[c] - 9*B*Cos[c])*Csc[c/2]*Sec[c/2]*Sec[c])/(5*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(29*A*Sin[(d*x)/2] - 9*B*Sin[(d*x)/2]))/(5*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(11*A*Sin[(d*x)/2] - 6*B*Sin[(d*x)/2]))/(15*d) + (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^5*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(5*d) + (16*A*Sec[c]*Sec[c + d*x]*Sin[d*x])/d + (4*(11*A - 6*B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(15*d) + (2*(A - B)*Sec[c/2 + (d*x)/2]^4*Tan[c/2])/(5*d)))/(a + a*Cos[c + d*x])^3 + (49*A*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*Sec[c/2]*(HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2)*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[Cos[c]*Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Tan[c]^2])*Sqrt[1 + Tan[c]^2]) - ((Sin[d*x + Arc...`

3.164.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3457, 27, 3042, 3457, 3042, 3457, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.164. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a(11A - B) - 5a(A - B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)^2} dx}{5a^2} - \frac{(A - B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(11A - B) - 5a(A - B) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)^2} dx}{10a^2} - \frac{(A - B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(11A - B) - 5a(A - B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(\sin\left(c + dx + \frac{\pi}{2}\right)a + a\right)^2} dx}{10a^2} - \frac{(A - B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a^2(41A - 6B) - 3a^2(8A - 3B) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)} dx}{3a^2} - \frac{2a(8A - 3B) \sin(c + dx)}{3d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^2} \\
& \quad \frac{10a^2}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \frac{(A - B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(41A - 6B) - 3a^2(8A - 3B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(\sin\left(c + dx + \frac{\pi}{2}\right)a + a\right)} dx}{3a^2} - \frac{2a(8A - 3B) \sin(c + dx)}{3d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^2} \\
& \quad \frac{10a^2}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \frac{(A - B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{3a^3(49A - 9B) - 5a^3(13A - 3B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)} dx}{3a^2} - \frac{5a^2(13A - 3B) \sin(c + dx)}{d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)} - \frac{2a(8A - 3B) \sin(c + dx)}{3d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^2} \\
& \quad \frac{10a^2}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3} \frac{(A - B) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^3}
\end{aligned}$$

3.164. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{\int \frac{3a^3(49A-9B)-5a^3(13A-3B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx}{3a^2} - \frac{5a^2(13A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} - \frac{2a(8A-3B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \frac{10a^2}{(A-B)\sin(c+dx)} \\
 & \frac{(A-B)\sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^3} \\
 & \downarrow 3042 \\
 & \frac{\int \frac{3a^3(49A-9B)-5a^3(13A-3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx}{3a^2} - \frac{5a^2(13A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} - \frac{2a(8A-3B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \frac{10a^2}{(A-B)\sin(c+dx)} \\
 & \frac{(A-B)\sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^3} \\
 & \downarrow 3227 \\
 & \frac{3a^3(49A-9B)\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx - 5a^3(13A-3B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3a^2} - \frac{5a^2(13A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} - \frac{2a(8A-3B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \frac{10a^2}{(A-B)\sin(c+dx)} \\
 & \frac{(A-B)\sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^3} \\
 & \downarrow 3042 \\
 & \frac{3a^3(49A-9B)\int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx - 5a^3(13A-3B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3a^2} - \frac{5a^2(13A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} - \frac{2a(8A-3B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \frac{10a^2}{(A-B)\sin(c+dx)} \\
 & \frac{(A-B)\sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^3} \\
 & \downarrow 3116 \\
 & \frac{3a^3(49A-9B)\left(\frac{2\sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx\right) - 5a^3(13A-3B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3a^2} - \frac{5a^2(13A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} - \frac{2a(8A-3B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^2} \\
 & \frac{10a^2}{(A-B)\sin(c+dx)} \\
 & \frac{(A-B)\sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^3} \\
 & \downarrow 3042
 \end{aligned}$$

3.164. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^3} dx$

$$\frac{3a^3(49A-9B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx \right) - 5a^3(13A-3B) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{2a^2} - \frac{5a^2(13A-3B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)} - \frac{2a(8A-3B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)}$$

$$\frac{(A-B) \sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)^3}$$

↓ 3119

$$\frac{3a^3(49A-9B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) - 5a^3(13A-3B) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{2a^2} - \frac{5a^2(13A-3B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)} - \frac{2a(8A-3B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)}$$

$$\frac{(A-B) \sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)^3}$$

↓ 3120

$$\frac{3a^3(49A-9B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) - \frac{10a^3(13A-3B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d}}{2a^2} - \frac{5a^2(13A-3B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)} - \frac{2a(8A-3B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)}$$

$$\frac{(A-B) \sin(c+dx)}{5d\sqrt{\cos(c+dx)}(a \cos(c+dx)+a)^3}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^3),x]`

output `-1/5*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^3) + ((-2*a*(8*A - 3*B)*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^2) + ((-5*a^2*(13*A - 3*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x]))) + ((-10*a^3*(13*A - 3*B)*EllipticF[(c + d*x)/2, 2])/d + 3*a^3*(49*A - 9*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/(2*a^2))/(3*a^2)/(10*a^2)`

3.164. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

3.164.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3116 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.164.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 684 vs. $2(253) = 506$.

Time = 5.10 (sec) , antiderivative size = 685, normalized size of antiderivative = 3.10

method	result	size
default	Expression too large to display	685

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

```
output -1/60*(-2*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/
2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(65*A*EllipticF(cos(1/2
*d*x+1/2*c),2^(1/2))-147*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-15*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+27*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+4*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(65*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-147*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-15*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+27*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-2*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(65*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-147*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-15*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+27*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))*cos(1/2*d*x+1/2*c)+12*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(49*A-9*B)*sin(1/2*d*x+1/2*c)^8-2*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(817*A-147*B)*sin(1/2*d*x+1/2*c)^6+6*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(248*A-43*B)*sin(1/2*d*x+1/2*c)^4-(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(439*A-69*B)*sin(1/2*d*x+1/2*c)^2/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.164.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 521, normalized size of antiderivative = 2.36

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx$$

$$= \frac{2(3(49A - 9B)\cos(dx + c)^3 + 2(188A - 33B)\cos(dx + c)^2 + 5(59A - 9B)\cos(dx + c) + 60A)\sqrt{\cos(dx + c)}}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^3,x, algorithm
m="fricas")
```

```
output 1/60*(2*(3*(49*A - 9*B)*cos(d*x + c)^3 + 2*(188*A - 33*B)*cos(d*x + c)^2 +
5*(59*A - 9*B)*cos(d*x + c) + 60*A)*sqrt(cos(d*x + c))*sin(d*x + c) - 5*(
sqrt(2)*(-13*I*A + 3*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(-13*I*A + 3*I*B)*cos
(d*x + c)^3 + 3*sqrt(2)*(-13*I*A + 3*I*B)*cos(d*x + c)^2 + sqrt(2)*(-13*I*
A + 3*I*B)*cos(d*x + c))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d
*x + c)) - 5*(sqrt(2)*(13*I*A - 3*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(13*I*A
- 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(13*I*A - 3*I*B)*cos(d*x + c)^2 + sqrt
(2)*(13*I*A - 3*I*B)*cos(d*x + c))*weierstrassPInverse(-4, 0, cos(d*x + c)
- I*sin(d*x + c)) - 3*(sqrt(2)*(49*I*A - 9*I*B)*cos(d*x + c)^4 + 3*sqrt(2
)*(49*I*A - 9*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(49*I*A - 9*I*B)*cos(d*x + c
)^2 + sqrt(2)*(49*I*A - 9*I*B)*cos(d*x + c))*weierstrassZeta(-4, 0, weiers
trassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*(sqrt(2)*(-49*I*A
+ 9*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(-49*I*A + 9*I*B)*cos(d*x + c)^3 + 3*
sqrt(2)*(-49*I*A + 9*I*B)*cos(d*x + c)^2 + sqrt(2)*(-49*I*A + 9*I*B)*cos(d
*x + c))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) -
I*sin(d*x + c))))/(a^3*d*cos(d*x + c)^4 + 3*a^3*d*cos(d*x + c)^3 + 3*a^3*d
*cos(d*x + c)^2 + a^3*d*cos(d*x + c))
```

3.164.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c))**3,x)
```

3.164. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

output Timed out

3.164.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^3,x, algorithm m="maxima")`

output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.

3.164.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*cos(d*x + c)^(3/2)), x)`

3.164.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{3}{2}}(a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^3), x)`

3.164. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

3.165
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$$

3.165.1 Optimal result 1600
 3.165.2 Mathematica [C] (warning: unable to verify) 1601
 3.165.3 Rubi [A] (verified) 1601
 3.165.4 Maple [B] (warning: unable to verify) 1606
 3.165.5 Fricas [C] (verification not implemented) 1607
 3.165.6 Sympy [F(-1)] 1608
 3.165.7 Maxima [F(-1)] 1608
 3.165.8 Giac [F] 1609
 3.165.9 Mupad [F(-1)] 1609

3.165.1 Optimal result

Integrand size = 33, antiderivative size = 254

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \frac{7(17A - 7B)E(\frac{1}{2}(c + dx)|2)}{10a^3d} + \frac{(33A - 13B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{6a^3d} + \frac{(33A - 13B) \sin(c + dx)}{6a^3d \cos^{\frac{3}{2}}(c + dx)} - \frac{7(17A - 7B) \sin(c + dx)}{10a^3d \sqrt{\cos(c + dx)}} - \frac{(A - B) \sin(c + dx)}{5d \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^3} - \frac{(2A - B) \sin(c + dx)}{3ad \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^2} - \frac{7(17A - 7B) \sin(c + dx)}{30d \cos^{\frac{3}{2}}(c + dx)(a^3 + a^3 \cos(c + dx))}$$

output

```
7/10*(17*A-7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(
sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/6*(33*A-13*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/d+1/6*(3
3*A-13*B)*sin(d*x+c)/a^3/d/cos(d*x+c)^(3/2)-1/5*(A-B)*sin(d*x+c)/d/cos(d*x
+c)^(3/2)/(a+a*cos(d*x+c))^3-1/3*(2*A-B)*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)/(
a+a*cos(d*x+c))^2-7/30*(17*A-7*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a^3+a^3*c
os(d*x+c))-7/10*(17*A-7*B)*sin(d*x+c)/a^3/d/cos(d*x+c)^(1/2)
```

3.165.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$$

3.165.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.36 (sec) , antiderivative size = 1110, normalized size of antiderivative = 4.37

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \text{Too large to display}$$

```
input Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^3),x]
```

```
output (-22*A*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(d*(a + a*Cos[c + d*x])^3*Sqrt[1 + Cot[c]^2]) + (26*B*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*HypergeometricPFQ[{1/4, 1/2}, {5/4}, Sin[d*x - ArcTan[Cot[c]]]^2*Sec[c/2]*Sec[d*x - ArcTan[Cot[c]]]*Sqrt[1 - Sin[d*x - ArcTan[Cot[c]]]]*Sqrt[-(Sqrt[1 + Cot[c]^2]*Sin[c]*Sin[d*x - ArcTan[Cot[c]]])]*Sqrt[1 + Sin[d*x - ArcTan[Cot[c]]]])/(3*d*(a + a*Cos[c + d*x])^3*Sqrt[1 + Cot[c]^2]) + (Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*((-2*(60*A - 20*B + 59*A*Cos[c] - 29*B*Cos[c])*Csc[c/2]*Sec[c/2]*Sec[c]/(5*d) - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(59*A*Sin[(d*x)/2] - 29*B*Sin[(d*x)/2]))/(5*d) - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(16*A*Sin[(d*x)/2] - 11*B*Sin[(d*x)/2]))/(15*d) - (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^5*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(5*d) + (16*A*Sec[c]*Sec[c + d*x]^2*Sin[d*x])/(3*d) + (16*Sec[c]*Sec[c + d*x]*(A*Sin[c] - 9*A*Sin[d*x] + 3*B*Sin[d*x]))/(3*d) - (4*(16*A - 11*B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(15*d) - (2*(A - B)*Sec[c/2 + (d*x)/2]^4*Tan[c/2])/(5*d)))/(a + a*Cos[c + d*x])^3 - (119*A*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*Sec[c/2]*((HypergeometricPFQ[-1/2, -1/4], {3/4}, Cos[d*x + ArcTan[Tan[c]]]^2*Sin[d*x + ArcTan[Tan[c]]]*Tan[c])/(Sqrt[1 - Cos[d*x + ArcTan[Tan[c]]]]*Sqrt[1 + Cos[d*x + ArcTan[Tan[c]]]])*...
```

3.165.3 Rubi [A] (verified)

Time = 1.35 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3457, 27, 3042, 3457, 3042, 3457, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.165. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a(13A-3B)-7a(A-B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(13A-3B)-7a(A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(13A-3B)-7a(A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{3a^2(23A-8B)-25a^2(2A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)} dx}{3a^2} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2} \\
& \quad \frac{10a^2}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3} \frac{(A-B)\sin(c+dx)}{1} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3a^2(23A-8B)-25a^2(2A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)} dx}{3a^2} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2} \\
& \quad \frac{10a^2}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3} \frac{(A-B)\sin(c+dx)}{1} \\
& \quad \downarrow \text{3457}
\end{aligned}$$

3.165. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

$$\frac{\int \frac{3(5a^3(33A-13B)-7a^3(17A-7B)\cos(c+dx))}{2\cos^{\frac{5}{2}}(c+dx)} dx}{3a^2} - \frac{7a^2(17A-7B)\sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3}$$

↓ 27

$$\frac{3\int \frac{5a^3(33A-13B)-7a^3(17A-7B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)} dx}{3a^2} - \frac{7a^2(17A-7B)\sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3}$$

↓ 3042

$$3\int \frac{5a^3(33A-13B)-7a^3(17A-7B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx}{3a^2} - \frac{7a^2(17A-7B)\sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3}$$

↓ 3227

$$3\left(\frac{5a^3(33A-13B)\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)} dx - 7a^3(17A-7B)\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx}{2a^2}\right) - \frac{7a^2(17A-7B)\sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3}$$

↓ 3042

$$3\left(\frac{5a^3(33A-13B)\int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx - 7a^3(17A-7B)\int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx}{2a^2}\right) - \frac{7a^2(17A-7B)\sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} - \frac{10a(2A-B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^2}$$

$$\frac{10a^2(A-B)\sin(c+dx)}{5d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^3}$$

↓ 3116

3.165. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))^3} dx$

$$\frac{3 \left(5a^3(33A-13B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 7a^3(17A-7B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\cos(c+dx)} dx \right) \right)}{2a^2} - \frac{7a^2(17A-7B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} - \frac{10a^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{(A-B) \sin(c+dx)}{5d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^3} \frac{10a^2}{3a^2}$$

↓ 3042

$$\frac{3 \left(5a^3(33A-13B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 7a^3(17A-7B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) \right)}{2a^2} - \frac{7a^2(17A-7B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} - \frac{10a^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{(A-B) \sin(c+dx)}{5d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^3} \frac{10a^2}{3a^2}$$

↓ 3119

$$\frac{3 \left(5a^3(33A-13B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 7a^3(17A-7B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) \right)}{2a^2} - \frac{7a^2(17A-7B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} - \frac{10a^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{(A-B) \sin(c+dx)}{5d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^3} \frac{10a^2}{3a^2}$$

↓ 3120

$$\frac{3 \left(5a^3(33A-13B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + \frac{2 \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) - 7a^3(17A-7B) \left(\frac{2 \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) \right)}{2a^2} - \frac{7a^2(17A-7B) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} - \frac{10a^2}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{(A-B) \sin(c+dx)}{5d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^3} \frac{10a^2}{3a^2}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*cos[c + d*x])^3),x]`

3.165. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

```
output -1/5*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^3)
+ ((-10*a*(2*A - B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d
*x])^2) + ((-7*a^2*(17*A - 7*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a
*Cos[c + d*x])) + (3*(5*a^3*(33*A - 13*B)*((2*EllipticF[(c + d*x)/2, 2])/(
3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))) - 7*a^3*(17*A - 7*B)*((-
2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]))))
)/(2*a^2)/(3*a^2)/(10*a^2)
```

3.165.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3116 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) I
nt[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.165.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 875 vs. $2(282) = 564$.

Time = 6.06 (sec) , antiderivative size = 876, normalized size of antiderivative = 3.45

method	result	size
default	Expression too large to display	876

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

output

```

1/60*(4*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(165*A*EllipticF(cos(1/2*
d*x+1/2*c),2^(1/2))-357*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-65*B*Ellip
ticF(cos(1/2*d*x+1/2*c),2^(1/2))+147*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2
)))*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-10*(-2*sin(1/2*d*x+1/2*c)^4+si
n(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*
c)^2-1)^(1/2)*(165*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-357*A*EllipticE
(cos(1/2*d*x+1/2*c),2^(1/2))-65*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+14
7*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*
x+1/2*c)+8*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(165*A*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))-357*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-65*B*El
lipticF(cos(1/2*d*x+1/2*c),2^(1/2))+147*B*EllipticE(cos(1/2*d*x+1/2*c),2^(
1/2)))*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-2*(-2*sin(1/2*d*x+1/2*c)^4+
sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/
2*c)^2-1)^(1/2)*(165*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-357*A*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2))-65*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+
147*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*cos(1/2*d*x+1/2*c)-168*(-2*si
n(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(17*A-7*B)*sin(1/2*d*x+1/2*
c)^10+8*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(1167*A-48...

```

3.165.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 548, normalized size of antiderivative = 2.16

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx =$$

$$\frac{2(21(17A - 7B) \cos(dx + c)^4 + 2(453A - 188B) \cos(dx + c)^3 + 5(139A - 59B) \cos(dx + c)^2 + 6}$$

input

```

integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^3,x, algorith
m="fracas")

```

output

```
-1/60*(2*(21*(17*A - 7*B)*cos(d*x + c)^4 + 2*(453*A - 188*B)*cos(d*x + c)^3 + 5*(139*A - 59*B)*cos(d*x + c)^2 + 60*(2*A - B)*cos(d*x + c) - 20*A)*sqrt(cos(d*x + c))*sin(d*x + c) + 5*(sqrt(2)*(33*I*A - 13*I*B)*cos(d*x + c)^5 + 3*sqrt(2)*(33*I*A - 13*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(33*I*A - 13*I*B)*cos(d*x + c)^3 + sqrt(2)*(33*I*A - 13*I*B)*cos(d*x + c)^2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(-33*I*A + 13*I*B)*cos(d*x + c)^5 + 3*sqrt(2)*(-33*I*A + 13*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(-33*I*A + 13*I*B)*cos(d*x + c)^3 + sqrt(2)*(-33*I*A + 13*I*B)*cos(d*x + c)^2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*(sqrt(2)*(-17*I*A + 7*I*B)*cos(d*x + c)^5 + 3*sqrt(2)*(-17*I*A + 7*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(-17*I*A + 7*I*B)*cos(d*x + c)^3 + sqrt(2)*(-17*I*A + 7*I*B)*cos(d*x + c)^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*(sqrt(2)*(17*I*A - 7*I*B)*cos(d*x + c)^5 + 3*sqrt(2)*(17*I*A - 7*I*B)*cos(d*x + c)^4 + 3*sqrt(2)*(17*I*A - 7*I*B)*cos(d*x + c)^3 + sqrt(2)*(17*I*A - 7*I*B)*cos(d*x + c)^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(a^3*d*cos(d*x + c)^5 + 3*a^3*d*cos(d*x + c)^4 + 3*a^3*d*cos(d*x + c)^3 + a^3*d*cos(d*x + c)^2)
```

3.165.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c))**3,x)`

output `Timed out`

3.165.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

3.165. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^3} dx$

output Timed out

3.165.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*cos(d*x + c)^(5/2)), x)`

3.165.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^3} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{5}{2}}(a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^3), x)`

3.166 $\int \cos^{\frac{5}{2}}(c+dx) \sqrt{a + a \cos(c + dx)} (A+B \cos(c+dx)) dx$

3.166.1 Optimal result	1610
3.166.2 Mathematica [A] (verified)	1611
3.166.3 Rubi [A] (verified)	1611
3.166.4 Maple [A] (verified)	1615
3.166.5 Fricas [A] (verification not implemented)	1615
3.166.6 Sympy [F(-1)]	1616
3.166.7 Maxima [B] (verification not implemented)	1616
3.166.8 Giac [F]	1617
3.166.9 Mupad [F(-1)]	1618

3.166.1 Optimal result

Integrand size = 35, antiderivative size = 221

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{5\sqrt{a}(8A + 7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{64d} + \frac{5a(8A + 7B) \sqrt{\cos(c + dx)} \sin(c + dx)}{64d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{5a(8A + 7B) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{96d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{a(8A + 7B) \cos^{\frac{5}{2}}(c + dx) \sin(c + dx)}{24d \sqrt{a + a \cos(c + dx)}} + \frac{aB \cos^{\frac{7}{2}}(c + dx) \sin(c + dx)}{4d \sqrt{a + a \cos(c + dx)}}$$

```
output 5/64*(8*A+7*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d
+5/96*a*(8*A+7*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/2
4*a*(8*A+7*B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/4*a*B
*cos(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+5/64*a*(8*A+7*B)*sin
(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.166.2 Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.61

$$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} (A+B \cos(c+dx)) dx$$

$$= \frac{\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(15\sqrt{2}(8A+7B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c+dx)\right)\right) + 2\sqrt{\cos(c+dx)}(152A + 133B + 2(40A + 53B)\cos(c+dx) + 4(8A+7B)\cos[2(c+dx)] + 12B\cos[3(c+dx)])\right)}{(384d)}$$

input `Integrate[Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(15*Sqrt[2]*(8*A + 7*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(152*A + 133*B + 2*(40*A + 53*B)*Cos[c + d*x] + 4*(8*A + 7*B)*Cos[2*(c + d*x)] + 12*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2])/(384*d)`

3.166.3 Rubi [A] (verified)Time = 0.92 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.95, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3460, 3042, 3249, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a} (A+B \cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{5}{2}} \sqrt{a \sin\left(c+dx+\frac{\pi}{2}\right) + a} (A+B \sin\left(c+dx+\frac{\pi}{2}\right)) dx$$

$$\downarrow \text{3460}$$

$$\frac{1}{8}(8A+7B) \int \cos^{\frac{5}{2}}(c+dx) \sqrt{\cos(c+dx)a + adx} + \frac{aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4d\sqrt{a \cos(c+dx) + a}}$$

$$\downarrow \text{3042}$$

$$\frac{1}{8}(8A+7B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{5}{2}} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a + adx} + \frac{aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4d\sqrt{a \cos(c+dx) + a}}$$

3.166. $\int \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\begin{aligned}
& \downarrow \text{3249} \\
& \frac{1}{8}(8A + 7B) \left(\frac{5}{6} \int \cos^{\frac{3}{2}}(c + dx) \sqrt{\cos(c + dx)a + adx} + \frac{a \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{4d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3042} \\
& \frac{1}{8}(8A + \\
7B) & \left(\frac{5}{6} \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + adx} + \frac{a \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{4d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3249} \\
& \frac{1}{8}(8A + \\
7B) & \left(\frac{5}{6} \left(\frac{3}{4} \int \sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + adx} + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{4d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3042} \\
& \frac{1}{8}(8A + \\
7B) & \left(\frac{5}{6} \left(\frac{3}{4} \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + adx} + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{4d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3249} \\
& \frac{1}{8}(8A + \\
7B) & \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) + \frac{a \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{4d\sqrt{a \cos(c + dx) + a}} \\
& \downarrow \text{3042}
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{8}(8A + \\
7B) & \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} a+a}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4d \sqrt{a \cos(c+dx)+a}} \right) \\
& \quad \downarrow \text{3253} \\
7B) & \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4d \sqrt{a \cos(c+dx)+a}} \right) \\
& \quad \downarrow \text{223} \\
7B) & \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{\sqrt{a} \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx)+a}}
\end{aligned}$$

input `Int[Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(4*d*Sqrt[a + a*Cos[c + d*x]]) + ((8*A + 7*B)*((a*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]])) + (5*((a*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]])) + (3*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)/6)/8`

3.166.3.1 Defintions of rubi rules used

- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3249 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.166.4 Maple [A] (verified)

Time = 15.00 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.57

method	result
default	$\frac{48B(\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 64A(\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 56B(\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}{24d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$A \left(8 \sin(dx+c) (\cos^2(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 10 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 15 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 15 \arctan\left(\tan(dx+c)\right) \right)$

input `int(cos(d*x+c)^(5/2)*(a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{192d} \left(48B \cos^3(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 64A \cos^2(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 56B \cos^2(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 80A \cos(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 70B \cos(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 120A \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \sin(dx+c) + 105B \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 120A \arctan\left(\tan(dx+c)\right) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 105B \arctan\left(\tan(dx+c)\right) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \right) \cos(dx+c)^{1/2} \left(a(1+\cos(dx+c)) \right)^{1/2} / (1+\cos(dx+c)) / \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2}$$

3.166.5 Fracas [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.68

$$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} (A+B \cos(c+dx)) dx$$

$$= \frac{(48B \cos(dx+c))^3 + 8(8A+7B) \cos(dx+c)^2 + 10(8A+7B) \cos(dx+c) + 120A + 105B}{192d} \sqrt{a \cos(c+dx)}$$

input `integrate(cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output $1/192*((48*B*\cos(d*x + c)^3 + 8*(8*A + 7*B)*\cos(d*x + c)^2 + 10*(8*A + 7*B)*\cos(d*x + c) + 120*A + 105*B)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c) - 15*((8*A + 7*B)*\cos(d*x + c) + 8*A + 7*B)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c)))/(d*\cos(d*x + c) + d)$

3.166.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output Timed out

3.166.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8220 vs. $2(189) = 378$.

Time = 0.82 (sec) , antiderivative size = 8220, normalized size of antiderivative = 37.19

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/768*(8*(4*(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(3/4)*(cos(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1))*sin(3*d*x + 3*c) - (cos(3*d*x + 3*c) - 1)*sin(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)))*sqrt(a) + 6*(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(1/4)*((sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 5*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)) - (cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 3*cos(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) - 4)*sin(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)))*sqrt(a) + 15*sqrt(a)*(arctan2(-(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(1/4)*(cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(...`

3.166.8 Giac [F]

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a} \cos(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)*cos(d*x + c)^(5/2), x)`

3.166.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \cos(c + dx)^{5/2} (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx$$

input `int(cos(c + d*x)^(5/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)`output `int(cos(c + d*x)^(5/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)`

3.167 $\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a + a \cos(c + dx)} (A+B \cos(c+dx)) dx$

3.167.1 Optimal result	1619
3.167.2 Mathematica [A] (verified)	1620
3.167.3 Rubi [A] (verified)	1620
3.167.4 Maple [A] (verified)	1623
3.167.5 Fricas [A] (verification not implemented)	1623
3.167.6 Sympy [F(-1)]	1624
3.167.7 Maxima [B] (verification not implemented)	1624
3.167.8 Giac [F]	1625
3.167.9 Mupad [F(-1)]	1626

3.167.1 Optimal result

Integrand size = 35, antiderivative size = 176

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{a}(6A + 5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{8d} + \frac{a(6A + 5B) \sqrt{\cos(c + dx)} \sin(c + dx)}{8d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{a(6A + 5B) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{12d \sqrt{a + a \cos(c + dx)}} + \frac{aB \cos^{\frac{5}{2}}(c + dx) \sin(c + dx)}{3d \sqrt{a + a \cos(c + dx)}}$$

```
output 1/8*(6*A+5*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d+
1/12*a*(6*A+5*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/3*
a*B*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/8*a*(6*A+5*B)*s
in(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.167.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.67

$$\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} (A+B \cos(c+dx)) dx$$

$$= \frac{\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(3\sqrt{2}(6A+5B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c+dx)\right)\right) + 2\sqrt{\cos(c+dx)}(18A+19B)\right)}{48d}$$

input `Integrate[Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])] * Sec[(c + d*x)/2] * (3*Sqrt[2]*(6*A + 5*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(18*A + 19*B + 2*(6*A + 5*B)*Cos[c + d*x] + 4*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.167.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.94, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3460, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a} (A+B \cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{a \sin\left(c+dx+\frac{\pi}{2}\right) + a} \left(A+B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3460}$$

$$\frac{1}{6}(6A+5B) \int \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a + adx} + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx) + a}}$$

$$\downarrow \text{3042}$$

$$\frac{1}{6}(6A+5B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a + adx} + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx) + a}}$$

$$\downarrow \text{3249}$$

3.167. $\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\begin{aligned}
& \frac{1}{6}(6A+5B) \left(\frac{3}{4} \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+adx} + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6}(6A+5B) \left(\frac{3}{4} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3249} \\
& \frac{1}{6}(6A+ \\
& 5B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6}(6A+ \\
& 5B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3253} \\
& \frac{1}{6}(6A+ \\
& 5B) \left(\frac{3}{4} \left(\frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{223}
\end{aligned}$$

$$5B) \left(\frac{3}{4} \left(\frac{\sqrt{a} \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx)+a}}$$

input `Int[Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + ((6 *A + 5*B)*((a*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]))/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))) /4))/6`

3.167.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt [a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3249 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x]) ^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.) *(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Co s[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && E qQ[a^2 - b^2, 0] && EqQ[d, a/b]`

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.167.4 Maple [A] (verified)

Time = 14.32 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.57

method	result
default	$\left(8B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 10B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 18A \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \sqrt{\cos(dx+c)} \sqrt{a(1+\cos(dx+c))}$
parts	$\frac{A \left(2 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3 \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)\right) (\sqrt{\cos(dx+c)}) \sqrt{a(1+\cos(dx+c))}}{4d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)), x, method=_RET
URNVERBOSE)
```

```
output 1/24/d*(8*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+12*A
*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+10*B*cos(d*x+c)*s
in(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+18*A*(cos(d*x+c)/(1+cos(d*x+c)
))^(1/2)*sin(d*x+c)+15*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+18*A
*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+15*B*arctan(tan(d*x+
c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c))
^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

3.167.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.76

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{(8B \cos(dx + c)^2 + 2(6A + 5B) \cos(dx + c) + 18A + 15B) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c) + 24(d \cos(dx + c) + d)}{24(d \cos(dx + c) + d)}$$

3.167. $\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algo
rithm="fricas")`

output `1/24*((8*B*cos(d*x + c)^2 + 2*(6*A + 5*B)*cos(d*x + c) + 18*A + 15*B)*sqrt
(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - 3*((6*A + 5*B)*cos(
d*x + c) + 6*A + 5*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x
+ c))/(sqrt(a)*sin(d*x + c))))/(d*cos(d*x + c) + d)`

3.167.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.167.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2981 vs. 2(150) = 300.

Time = 0.65 (sec) , antiderivative size = 2981, normalized size of antiderivative = 16.94

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algo
rithm="maxima")`

output `1/96*(6*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*((cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(2*d*x + 2*c) - (cos(2*d*x + 2*c) - 2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + ((cos(2*d*x + 2*c) - 2)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + sin(2*d*x + 2*c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) - cos(2*d*x + 2*c) + 2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(a) + 3*sqrt(a)*(arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))) + 1 - arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))...`

3.167.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)*cos(d*x + c)^(3/2), x)`

3.167.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \cos(c + dx)^{3/2} (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)`output `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)`

3.168 $\int \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.168.1 Optimal result	1627
3.168.2 Mathematica [A] (verified)	1627
3.168.3 Rubi [A] (verified)	1628
3.168.4 Maple [A] (verified)	1630
3.168.5 Fricas [A] (verification not implemented)	1631
3.168.6 Sympy [F]	1631
3.168.7 Maxima [B] (verification not implemented)	1632
3.168.8 Giac [F]	1632
3.168.9 Mupad [F(-1)]	1633

3.168.1 Optimal result

Integrand size = 35, antiderivative size = 131

$$\int \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{a}(4A + 3B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right)}{4d}$$

$$+ \frac{a(4A + 3B) \sqrt{\cos(c + dx)} \sin(c + dx)}{4d \sqrt{a + a \cos(c + dx)}} + \frac{aB \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2d \sqrt{a + a \cos(c + dx)}}$$

output $1/4*(4*A+3*B)*\arcsin(\sin(d*x+c)*a^{(1/2)}/(a+a*\cos(d*x+c))^{(1/2)})*a^{(1/2)}/d+1/2*a*B*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+1/4*a*(4*A+3*B)*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(a+a*\cos(d*x+c))^{(1/2)}$

3.168.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.76

$$\int \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(\sqrt{2}(4A + 3B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) + 2\sqrt{\cos(c + dx)}(4A + B \cos(c + dx))}{8d}$$

3.168. $\int \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$

input `Integrate[Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])] * Sec[(c + d*x)/2] * (Sqrt[2]*(4*A + 3*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(4*A + 3*B + 2*B*Cos[c + d*x])*Sin[(c + d*x)/2]))/(8*d)`

3.168.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.94, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3460, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx) + a} (A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{a \sin\left(c+dx+\frac{\pi}{2}\right) + a} \left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3460} \\
 & \frac{1}{4}(4A + 3B) \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a + a} dx + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4}(4A + 3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right) a + a} dx + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx) + a}} \\
 & \quad \downarrow \text{3249} \\
 & \frac{1}{4}(4A + 3B) \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a + a}}{\sqrt{\cos(c+dx)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx) + a}} \right) + \\
 & \quad \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx) + a}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{4}(4A + 3B) \left(\frac{1}{2} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow \text{3253} \\
& \frac{1}{4}(4A + 3B) \left(\frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx)a + a}}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}} \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow \text{223} \\
& \frac{1}{4}(4A + 3B) \left(\frac{\sqrt{a} \arcsin \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}}
\end{aligned}$$

input `Int[Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + ((4*A + 3*B)*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4`

3.168.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3249 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]
```

```
rule 3253 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.168.4 Maple [A] (verified)

Time = 15.52 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.57

method	result
default	$\frac{(2B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 3B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)) \sqrt{\cos(dx+c)} \sqrt{a(1+\cos(dx+c))}}{4d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \left(\sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right) (\sqrt{\cos(dx+c)}) \sqrt{a(1+\cos(dx+c))}}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}} + \frac{B (2 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c)),x,method=_RET URNVERBOSE)
```

output $1/4/d*(2*B*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+4*A*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\sin(d*x+c)+3*B*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+4*A*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2})+3*B*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}))*\cos(d*x+c)^{1/2}*(a*(1+\cos(d*x+c)))^{1/2}/(1+\cos(d*x+c))/(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}$

3.168.5 Fricas [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.89

$$\int \sqrt{\cos(c+dx)}\sqrt{a+a\cos(c+dx)}(A+B\cos(c+dx))dx$$

$$= \frac{(2B\cos(dx+c)+4A+3B)\sqrt{a\cos(dx+c)+a}\sqrt{\cos(dx+c)}\sin(dx+c)-((4A+3B)\cos(dx+c))}{4(d\cos(dx+c)+d)}$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output $1/4*((2*B*\cos(d*x+c)+4*A+3*B)*\sqrt{a*\cos(d*x+c)+a}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-((4*A+3*B)*\cos(d*x+c)+4*A+3*B)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x+c)+a}*\sqrt{\cos(d*x+c)})/(\sqrt{a}*\sin(d*x+c)))/(\cos(d*x+c)+d)$

3.168.6 Sympy [F]

$$\int \sqrt{\cos(c+dx)}\sqrt{a+a\cos(c+dx)}(A+B\cos(c+dx))dx$$

$$= \int \sqrt{a(\cos(c+dx)+1)}(A+B\cos(c+dx))\sqrt{\cos(c+dx)}dx$$

input `integrate(cos(d*x+c)**(1/2)*(a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Integral(sqrt(a*(cos(c+d*x)+1))*(A+B*cos(c+d*x))*sqrt(cos(c+d*x)),x)`

3.168.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1851 vs. $2(111) = 222$.

Time = 0.54 (sec) , antiderivative size = 1851, normalized size of antiderivative = 14.13

$$\int \sqrt{\cos(c+dx)} \sqrt{a+a\cos(c+dx)} (A+B\cos(c+dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algo
rithm="maxima")`

output `1/16*(4*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) +
1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*
x + c) - (cos(d*x + c) - 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c) + 1)))*sqrt(a) + sqrt(a)*(arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x +
2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c),
cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*
*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)
^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) + 1))) + 1) - arctan2(-(cos(2*d*x + 2*c)^2 + sin(2
*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(
sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x
+ 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(si
n(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(
2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) - 1) - arctan2((cos(2*d*x + 2*c)^2 +
sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*
d*x + 2*c), cos(2*d*x + 2*c) + 1)), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)
^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2
*d*x + 2*c) + 1)) + 1) + arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)...`

3.168.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c+dx)} \sqrt{a+a\cos(c+dx)} (A+B\cos(c+dx)) dx \\ &= \int (B\cos(dx+c) + A) \sqrt{a\cos(dx+c) + a} \sqrt{\cos(dx+c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))
, x)`

3.168.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \sqrt{\cos(c + dx)} (A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)} dx$$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2), x)`

3.169
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.169.1 Optimal result 1634
 3.169.2 Mathematica [A] (verified) 1634
 3.169.3 Rubi [A] (verified) 1635
 3.169.4 Maple [B] (verified) 1636
 3.169.5 Fricas [A] (verification not implemented) 1637
 3.169.6 Sympy [F] 1637
 3.169.7 Maxima [B] (verification not implemented) 1638
 3.169.8 Giac [F(-1)] 1638
 3.169.9 Mupad [F(-1)] 1639

3.169.1 Optimal result

Integrand size = 35, antiderivative size = 78

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

$$= \frac{\sqrt{a}(2A+B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} + \frac{aB \sqrt{\cos(c+dx)} \sin(c+dx)}{d \sqrt{a+a \cos(c+dx)}}$$

output `(2*A+B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d+a*B*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.169.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

$$= \frac{\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(\sqrt{2}(2A+B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c+dx)\right)\right) + 2B \sqrt{\cos(c+dx)} \sin\left(\frac{1}{2}(c+dx)\right)\right)}{2d}$$

input `Integrate[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

3.169.
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

output $(\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])] * \text{Sec}[(c + d*x)/2] * (\text{Sqrt}[2]*(2*A + B) * \text{ArcSin}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]] + 2*B*\text{Sqrt}[\text{Cos}[c + d*x]] * \text{Sin}[(c + d*x)/2])) / (2*d)$

3.169.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{3460} \\ & \frac{1}{2}(2A + B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2}(2A + B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \\ & \quad \downarrow \text{3253} \\ & \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} - \frac{(2A + B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx)a + a}}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} \\ & \quad \downarrow \text{223} \\ & \frac{\sqrt{a}(2A + B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} + \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \end{aligned}$$

input $\text{Int}[(\text{Sqrt}[a + a*\text{Cos}[c + d*x]]*(A + B*\text{Cos}[c + d*x]))/\text{Sqrt}[\text{Cos}[c + d*x]],x]$

3.169. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

output $(\sqrt{a}*(2*A + B)*\text{ArcSin}[(\sqrt{a}*\text{Sin}[c + d*x])/\sqrt{a + a*\text{Cos}[c + d*x]})]/d + (a*B*\sqrt{\text{Cos}[c + d*x]}*\text{Sin}[c + d*x])/(d*\sqrt{a + a*\text{Cos}[c + d*x]})$

3.169.3.1 Defintions of rubi rules used

rule 223 $\text{Int}[1/\sqrt{(a_+) + (b_+)*(x_+)^2}, x_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\sqrt{a})]/\text{Rt}[-b, 2], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{GtQ}\{a, 0\} \ \&\& \ \text{NegQ}\{b\}$

rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}\{u, x\}$

rule 3253 $\text{Int}[\sqrt{(a_+) + (b_+)*\text{sin}[(e_+) + (f_+)*(x_+)]}/\sqrt{(d_+)*\text{sin}[(e_+) + (f_+)*(x_+)]}, x_Symbol] \rightarrow \text{Simp}[-2/f \ \text{Subst}[\text{Int}[1/\sqrt{1 - x^2/a}, x], x, b*(\text{Cos}[e + f*x]/\sqrt{a + b*\text{Sin}[e + f*x]})], x] /;$ $\text{FreeQ}\{a, b, d, e, f, x\} \ \&\& \ \text{EqQ}\{a^2 - b^2, 0\} \ \&\& \ \text{EqQ}\{d, a/b\}$

rule 3460 $\text{Int}[\sqrt{(a_+) + (b_+)*\text{sin}[(e_+) + (f_+)*(x_+)]}*((A_+) + (B_+)*\text{sin}[(e_+) + (f_+)*(x_+)])*((c_+) + (d_+)*\text{sin}[(e_+) + (f_+)*(x_+)])^{n_+}, x_Symbol] \rightarrow \text{Simp}[-2*b*B*\text{Cos}[e + f*x]*((c + d*\text{Sin}[e + f*x])^{(n + 1)})/(d*f*(2*n + 3)*\sqrt{a + b*\text{Sin}[e + f*x]}), x] + \text{Simp}[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) \ \text{Int}[\sqrt{a + b*\text{Sin}[e + f*x]}*(c + d*\text{Sin}[e + f*x])^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, A, B, n, x\} \ \&\& \ \text{NeQ}\{b*c - a*d, 0\} \ \&\& \ \text{EqQ}\{a^2 - b^2, 0\} \ \&\& \ \text{NeQ}\{c^2 - d^2, 0\} \ \&\& \ !\text{LtQ}\{n, -1\}$

3.169.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 140 vs. $2(68) = 136$.

Time = 18.24 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.81

method	result
default	$\frac{\sqrt{a(1+\cos(dx+c))} (\sqrt{\cos(dx+c)}) (B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 2A \arctan(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}) + B \arctan(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}))}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{2A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sqrt{a(1+\cos(dx+c))} \arctan(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})}{d \sqrt{\cos(dx+c)}} + \frac{B (\sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \arctan(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}))}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

3.169.
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

input `int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^(1/2)*(B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)`

3.169.5 Fricas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.24

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{\sqrt{a \cos(dx + c) + aB} \sqrt{\cos(dx + c)} \sin(dx + c) - ((2A + B) \cos(dx + c) + 2A + B) \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx + c)}}{\sqrt{\cos(dx + c)}}\right)}{d \cos(dx + c) + d}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `(sqrt(a*cos(d*x + c) + a)*B*sqrt(cos(d*x + c))*sin(d*x + c) - ((2*A + B)*cos(d*x + c) + 2*A + B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))))/(d*cos(d*x + c) + d)`

3.169.6 Sympy [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{\sqrt{a (\cos(c + dx) + 1)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))/sqrt(cos(c + d*x)), x)`

3.169. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

3.169.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 939 vs. $2(68) = 136$.

Time = 0.46 (sec) , antiderivative size = 939, normalized size of antiderivative = 12.04

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `1/4*(4*A*sqrt(a)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + cos(d*x + c)) + (2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (cos(d*x + c) - 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))))*sqrt(a) + sqrt(a)*(arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) + 1) - arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) - 1) - arctan2((co...`

3.169.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algo
rithm="giac")`

3.169. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

output Timed out

3.169.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2),x)`

$$3.170 \quad \int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.170.1 Optimal result	1640
3.170.2 Mathematica [A] (verified)	1640
3.170.3 Rubi [A] (verified)	1641
3.170.4 Maple [A] (verified)	1642
3.170.5 Fricas [A] (verification not implemented)	1643
3.170.6 Sympy [F]	1643
3.170.7 Maxima [B] (verification not implemented)	1644
3.170.8 Giac [F]	1644
3.170.9 Mupad [F(-1)]	1645

3.170.1 Optimal result

Integrand size = 35, antiderivative size = 76

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{2\sqrt{a}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} + \frac{2aA \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}$$

output `2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)/d+2*a*A*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)`

3.170.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.13

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(\sqrt{2}B \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c+dx)\right)\right)\right) \sqrt{\cos(c+dx)} + 2A \sin\left(\frac{1}{2}(c+dx)\right)}{d\sqrt{\cos(c+dx)}}$$

input `Integrate[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

3.170. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

output $(\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])] * \text{Sec}[(c + d*x)/2] * (\text{Sqrt}[2]*B*\text{ArcSin}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]] * \text{Sqrt}[\text{Cos}[c + d*x]] + 2*A*\text{Sin}[(c + d*x)/2])) / (d*\text{Sqrt}[\text{Cos}[c + d*x]])$

3.170.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 3459, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

↓ 3459

$$B \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}$$

↓ 3042

$$B \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2aA \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}$$

↓ 3253

$$\frac{2aA \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} - \frac{2B \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx)a + a}}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d}$$

↓ 223

$$\frac{2aA \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} + \frac{2\sqrt{a}B \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d}$$

input $\text{Int}[(\text{Sqrt}[a + a*\text{Cos}[c + d*x]]*(A + B*\text{Cos}[c + d*x]))/\text{Cos}[c + d*x]^{(3/2)}, x]$

$$3.170. \quad \int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

output $(2\sqrt{a}B\text{ArcSin}[\frac{\sqrt{a}\sin[c+dx]}{\sqrt{a+a\cos[c+dx]}}])/d + (2aA\sin[c+dx])/(d\sqrt{\cos[c+dx]}\sqrt{a+a\cos[c+dx]})$

3.170.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3459 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x])], x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.170.4 Maple [A] (verified)

Time = 7.56 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.53

method	result
parts	$\frac{2A\sqrt{a(1+\cos(dx+c))} \sin(dx+c)}{d(1+\cos(dx+c))\sqrt{\cos(dx+c)}} + \frac{2B\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sqrt{a(1+\cos(dx+c))} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{d\sqrt{\cos(dx+c)}}$
default	$\frac{2\left(B\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \cos(dx+c) + B\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + A \sin(dx+c)\right)}{d(1+\cos(dx+c))\sqrt{\cos(dx+c)}}$

3.170. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

```
input int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 2*A/d*(a*(1+cos(d*x+c)))^(1/2)*sin(d*x+c)/(1+cos(d*x+c))/cos(d*x+c)^(1/2)+
2*B/d/cos(d*x+c)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(a*(1+cos(d*x+c))
)^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))
```

3.170.5 Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.43

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 \left(\sqrt{a \cos(dx + c) + a} A \sqrt{\cos(dx + c)} \sin(dx + c) - (B \cos(dx + c)^2 + B \cos(dx + c)) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx + c) + a}}{\cos(dx + c)} \right) \right)}{d \cos(dx + c)^2 + d \cos(dx + c)}$$

```
input integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algo
rithm="fricas")
```

```
output 2*(sqrt(a*cos(d*x + c) + a)*A*sqrt(cos(d*x + c))*sin(d*x + c) - (B*cos(d*x
+ c)^2 + B*cos(d*x + c))*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos
(d*x + c))/(sqrt(a)*sin(d*x + c))))/(d*cos(d*x + c)^2 + d*cos(d*x + c))
```

3.170.6 Sympy [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{\sqrt{a (\cos(c + dx) + 1)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

```
input integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)
```

```
output Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))/cos(c + d*x)**(3/
2), x)
```

3.170. $\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$

3.170.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. 2(66) = 132.

Time = 0.39 (sec) , antiderivative size = 245, normalized size of antiderivative = 3.22

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$B\sqrt{a} \arctan \left((\cos(2dx + 2c)^2 + \sin(2dx + 2c)^2 + 2 \cos(2dx + 2c) + 1)^{\frac{1}{4}} \sin \left(\frac{1}{2} \arctan(\sin(2dx + 2c) / (\cos(2dx + 2c) + 1)) \right) \right) + A \sqrt{a} \arctan \left(\frac{\sin(dx + c)}{\cos(dx + c) + 1} \right) - \sqrt{2} \sqrt{a} \sin(dx + c) / (\cos(dx + c) + 1) + \sqrt{2} \sqrt{a} \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 / ((\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{\frac{3}{2}}) / d$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`

output `(B*sqrt(a)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) + cos(d*x + c)) + 2*A*(sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(3/2))*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(3/2))/d`

3.170.8 Giac [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`

3.170. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

3.170.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\cos(c + dx)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^(3/2),x)`

3.171
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.171.1 Optimal result 1646
 3.171.2 Mathematica [A] (verified) 1646
 3.171.3 Rubi [A] (verified) 1647
 3.171.4 Maple [A] (verified) 1648
 3.171.5 Fricas [A] (verification not implemented) 1649
 3.171.6 Sympy [F] 1649
 3.171.7 Maxima [B] (verification not implemented) 1649
 3.171.8 Giac [F(-1)] 1650
 3.171.9 Mupad [B] (verification not implemented) 1650

3.171.1 Optimal result

Integrand size = 35, antiderivative size = 85

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{2aA \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a(2A+3B) \sin(c+dx)}{3d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}$$

output $2/3*a*A*\sin(d*x+c)/d/\cos(d*x+c)^{(3/2)}/(a+a*\cos(d*x+c))^{(1/2)}+2/3*a*(2*A+3*B)*\sin(d*x+c)/d/\cos(d*x+c)^{(1/2)}/(a+a*\cos(d*x+c))^{(1/2)}$

3.171.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{2\sqrt{a(1+\cos(c+dx))}(A+(2A+3B)\cos(c+dx))\tan\left(\frac{1}{2}(c+dx)\right)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

input $\text{Integrate}[(\text{Sqrt}[a+a*\text{Cos}[c+d*x]])*(A+B*\text{Cos}[c+d*x]))/\text{Cos}[c+d*x]^{(5/2)},x]$

output $(2*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*(A + (2*A + 3*B)*\text{Cos}[c + d*x])* \text{Tan}[(c + d*x)/2])/(3*d*\text{Cos}[c + d*x]^{(3/2)})$

3.171.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.114$, Rules used = {3042, 3459, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3459

$$\frac{1}{3}(2A + 3B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

↓ 3042

$$\frac{1}{3}(2A + 3B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

↓ 3250

$$\frac{2a(2A + 3B) \sin(c + dx)}{3d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

input $\text{Int}[(\text{Sqrt}[a + a*\text{Cos}[c + d*x]]*(A + B*\text{Cos}[c + d*x]))/\text{Cos}[c + d*x]^{(5/2)}, x]$

output $(2*a*A*\text{Sin}[c + d*x])/(3*d*\text{Cos}[c + d*x]^{(3/2)}*\text{Sqrt}[a + a*\text{Cos}[c + d*x]]) + (2*a*(2*A + 3*B)*\text{Sin}[c + d*x])/(3*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + a*\text{Cos}[c + d*x]])$

3.171. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

3.171.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.171.4 Maple [A] (verified)

Time = 7.33 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{2 \sin(dx+c)(2A \cos(dx+c)+3B \cos(dx+c)+A) \sqrt{a(1+\cos(dx+c))}}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}}$	62
parts	$\frac{2A \sin(dx+c)(2 \cos(dx+c)+1) \sqrt{a(1+\cos(dx+c))}}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}} + \frac{2B \sqrt{a(1+\cos(dx+c))} \sin(dx+c)}{d(1+\cos(dx+c)) \sqrt{\cos(dx+c)}}$	96

input `int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RETURNVERBOSE)`

output `2/3/d*sin(d*x+c)*(2*A*cos(d*x+c)+3*B*cos(d*x+c)+A)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(3/2)`

3.171.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2((2A + 3B) \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c)}{3(d \cos(dx + c))^3 + d \cos(dx + c)^2}$$

```
input integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algo
rithm="fricas")
```

```
output 2/3*((2*A + 3*B)*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x +
c))*sin(d*x + c)/(d*cos(d*x + c)^3 + d*cos(d*x + c)^2)
```

3.171.6 Sympy [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{\sqrt{a (\cos(c + dx) + 1)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

```
input integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)
```

```
output Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))/cos(c + d*x)**(5/
2), x)
```

3.171.7 Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 289 vs. $2(73) = 146$.

Time = 0.35 (sec) , antiderivative size = 289, normalized size of antiderivative = 3.40

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 \left(\frac{3B \left(\frac{\sqrt{2}\sqrt{a} \sin(dx+c)}{\cos(dx+c)+1} - \frac{\sqrt{2}\sqrt{a} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{3}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{3}{2}}} + \frac{A \left(\frac{3\sqrt{2}\sqrt{a} \sin(dx+c)}{\cos(dx+c)+1} - \frac{4\sqrt{2}\sqrt{a} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{\sqrt{2}\sqrt{a} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^2}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{5}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{5}{2}} \left(\frac{2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)} \right)}{3d}$$

3.171. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorith="maxima")`

output `2/3*(3*B*(sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(3/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(3/2)) + A*(3*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 4*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + sqrt(2)*sqrt(a)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^2/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(5/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(5/2)*(2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 1)))/d`

3.171.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorith="giac")`

output Timed out

3.171.9 Mupad [B] (verification not implemented)

Time = 1.64 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.32

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 \sqrt{a} (\cos(c + dx) + 1) (2 A \sin(c + dx) + 3 B \sin(c + dx) + 2 A \sin(2c + 2dx) + 2 A \sin(3c + 3dx))}{3 d \sqrt{\cos(c + dx)} (3 \cos(c + dx) + 2 \cos(2c + 2dx) + \cos(3c + 3dx) + 2)}$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^(5/2),x)`

3.171. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

output $(2*(a*(\cos(c + d*x) + 1))^{(1/2)}*(2*A*\sin(c + d*x) + 3*B*\sin(c + d*x) + 2*A*\sin(2*c + 2*d*x) + 2*A*\sin(3*c + 3*d*x) + 3*B*\sin(3*c + 3*d*x)))/(3*d*\cos(c + d*x)^{(1/2)}*(3*\cos(c + d*x) + 2*\cos(2*c + 2*d*x) + \cos(3*c + 3*d*x) + 2))$

3.171. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

3.172
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.172.1 Optimal result 1652
 3.172.2 Mathematica [A] (verified) 1652
 3.172.3 Rubi [A] (verified) 1653
 3.172.4 Maple [A] (verified) 1655
 3.172.5 Fricas [A] (verification not implemented) 1655
 3.172.6 Sympy [F(-1)] 1656
 3.172.7 Maxima [B] (verification not implemented) 1656
 3.172.8 Giac [F(-1)] 1657
 3.172.9 Mupad [B] (verification not implemented) 1657

3.172.1 Optimal result

Integrand size = 35, antiderivative size = 130

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{2aA \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a(4A+5B) \sin(c+dx)}{15d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}}$$

$$+ \frac{4a(4A+5B) \sin(c+dx)}{15d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}$$

output `2/5*a*A*sin(d*x+c)/d/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+2/15*a*(4*A+5*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+4/15*a*(4*A+5*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)`

3.172.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{2\sqrt{a(1+\cos(c+dx))}(7A+5B+(4A+5B)\cos(c+dx)+(4A+5B)\cos(2(c+dx))) \tan(\frac{1}{2}(c+dx))}{15d \cos^{\frac{5}{2}}(c+dx)}$$

input `Integrate[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*(7*A + 5*B + (4*A + 5*B)*Cos[c + d*x] + (4*A + 5*B)*Cos[2*(c + d*x)])*Tan[(c + d*x)/2])/(15*d*Cos[c + d*x]^(5/2))`

3.172.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3042, 3459, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

↓ 3459

$$\frac{1}{5}(4A + 5B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{5}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

↓ 3042

$$\frac{1}{5}(4A + 5B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

↓ 3251

$$\frac{1}{5}(4A + 5B) \left(\frac{2}{3} \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2a \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

↓ 3042

3.172. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5}(4A + 5B) \left(\frac{2}{3} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

↓ 3250

$$\frac{1}{5}(4A + 5B) \left(\frac{2a \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} + \frac{4a \sin(c + dx)}{3d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}$$

input `Int[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) + ((4*A + 5*B)*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]])) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/5`

3.172.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp [(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1) * (b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

3.172.4 Maple [A] (verified)

Time = 7.38 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.66

method	result	size
default	$\frac{2 \sin(dx+c)(8A(\cos^2(dx+c))+10B(\cos^2(dx+c))+4A \cos(dx+c)+5B \cos(dx+c)+3A)\sqrt{a(1+\cos(dx+c))}}{15d(1+\cos(dx+c)) \cos(dx+c)^{\frac{5}{2}}}$	86
parts	$\frac{2A \sin(dx+c)(8(\cos^2(dx+c))+4 \cos(dx+c)+3)\sqrt{a(1+\cos(dx+c))}}{15d(1+\cos(dx+c)) \cos(dx+c)^{\frac{5}{2}}} + \frac{2B \sin(dx+c)(2 \cos(dx+c)+1)\sqrt{a(1+\cos(dx+c))}}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}}$	116

```
input int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RETURNVERBOSE)
```

```
output 2/15/d*sin(d*x+c)*(8*A*cos(d*x+c)^2+10*B*cos(d*x+c)^2+4*A*cos(d*x+c)+5*B*cos(d*x+c)+3*A)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(5/2)
```

3.172.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.66

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{2(2(4A+5B) \cos(dx+c)^2+(4A+5B) \cos(dx+c)+3A)\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)} \sin(dx+c)}{15(d \cos(dx+c))^4+d \cos(dx+c)^3}$$

```
input integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="fracas")
```

output $2/15*(2*(4*A + 5*B)*\cos(d*x + c)^2 + (4*A + 5*B)*\cos(d*x + c) + 3*A)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^4 + d*\cos(d*x + c)^3)$

3.172.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output Timed out

3.172.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 428 vs. $2(112) = 224$.

Time = 0.34 (sec) , antiderivative size = 428, normalized size of antiderivative = 3.29

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2 \left(\frac{5B \left(\frac{3\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{4\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^2}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{5}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{5}{2}} \left(\frac{2\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)} + \frac{A \left(\frac{15\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{25\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{17\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^2}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(\frac{2\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)} \right)}{15d}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo rithm="maxima")`

output $\frac{2}{15} * (5 * B * (3 * \sqrt{2} * \sqrt{a} * \sin(dx + c) / (\cos(dx + c) + 1) - 4 * \sqrt{2} * \sqrt{a} * \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + \sqrt{2} * \sqrt{a} * \sin(dx + c)^5 / (\cos(dx + c) + 1)^5) * (\sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + 1)^2 / ((\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{5/2} * (-\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{5/2} * (2 * \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 + 1)) + A * (15 * \sqrt{2} * \sqrt{a} * \sin(dx + c) / (\cos(dx + c) + 1) - 25 * \sqrt{2} * \sqrt{a} * \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + 17 * \sqrt{2} * \sqrt{a} * \sin(dx + c)^5 / (\cos(dx + c) + 1)^5 - 7 * \sqrt{2} * \sqrt{a} * \sin(dx + c)^7 / (\cos(dx + c) + 1)^7) * (\sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + 1)^3 / ((\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{7/2} * (-\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{7/2} * (3 * \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + 3 * \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 + \sin(dx + c)^6 / (\cos(dx + c) + 1)^6 + 1))) / d$

3.172.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(dx+c))^(1/2)*(A+B*cos(dx+c))/cos(dx+c)^(7/2),x, algorithm="giac")`

output Timed out

3.172.9 Mupad [B] (verification not implemented)

Time = 3.44 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.49

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$$

$$= \frac{4 \sqrt{a (\cos(c + dx) + 1)} (14 A \sin(c + dx) + 10 B \sin(c + dx) + 8 A \sin(2c + 2dx) + 18 A \sin(3c + 3dx))}{15 d \sqrt{\cos(c + dx)} (10 \cos(c + dx) + 8 \cos(2c + 2dx) + 6 \cos(3c + 3dx) + 1)}$$

input `int(((A + B*cos(c + dx))*(a + a*cos(c + dx))^(1/2))/cos(c + dx)^(7/2),x)`

3.172. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

output $(4*(a*(\cos(c + d*x) + 1))^{1/2}*(14*A*\sin(c + d*x) + 10*B*\sin(c + d*x) + 8*A*\sin(2*c + 2*d*x) + 18*A*\sin(3*c + 3*d*x) + 4*A*\sin(4*c + 4*d*x) + 4*A*\sin(5*c + 5*d*x) + 10*B*\sin(2*c + 2*d*x) + 15*B*\sin(3*c + 3*d*x) + 5*B*\sin(4*c + 4*d*x) + 5*B*\sin(5*c + 5*d*x)))/(15*d*\cos(c + d*x)^{1/2}*(10*\cos(c + d*x) + 8*\cos(2*c + 2*d*x) + 5*\cos(3*c + 3*d*x) + 2*\cos(4*c + 4*d*x) + \cos(5*c + 5*d*x) + 6))$

3.172. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

3.173
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.173.1 Optimal result 1659
 3.173.2 Mathematica [A] (verified) 1659
 3.173.3 Rubi [A] (verified) 1660
 3.173.4 Maple [A] (verified) 1662
 3.173.5 Fricas [A] (verification not implemented) 1663
 3.173.6 Sympy [F(-1)] 1663
 3.173.7 Maxima [B] (verification not implemented) 1664
 3.173.8 Giac [C] (verification not implemented) 1664
 3.173.9 Mupad [B] (verification not implemented) 1665

3.173.1 Optimal result

Integrand size = 35, antiderivative size = 175

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2aA \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a(6A+7B) \sin(c+dx)}{35d \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}}$$

$$+ \frac{8a(6A+7B) \sin(c+dx)}{105d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{16a(6A+7B) \sin(c+dx)}{105d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}$$

```
output 2/7*a*A*sin(d*x+c)/d/cos(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2)+2/35*a*(6*A+7
*B)*sin(d*x+c)/d/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+8/105*a*(6*A+7*B)
*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+16/105*a*(6*A+7*B)*s
in(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)
```

3.173.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.58

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2\sqrt{a(1+\cos(c+dx))}(27A+14B+9(6A+7B)\cos(c+dx)+2(6A+7B)\cos(2(c+dx))+12A\cos(3(c+dx)))}{105d \cos^{\frac{7}{2}}(c+dx)}$$

3.173.
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

input `Integrate[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*(27*A + 14*B + 9*(6*A + 7*B)*Cos[c + d*x] + 2*(6*A + 7*B)*Cos[2*(c + d*x)] + 12*A*Cos[3*(c + d*x)] + 14*B*Cos[3*(c + d*x)])*Tan[(c + d*x)/2])/(105*d*Cos[c + d*x]^(7/2))`

3.173.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3459, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx \\
 & \quad \downarrow \text{3459} \\
 & \frac{1}{7}(6A + 7B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{7}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7}(6A + 7B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx + \frac{2aA \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3251} \\
 & \frac{1}{7}(6A + 7B) \left(\frac{4}{5} \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{5}{2}}(c + dx)} dx + \frac{2a \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \\
 & \quad \frac{2aA \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.173. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned}
& \frac{1}{7}(6A + 7B) \left(\frac{4}{5} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + \frac{2a \sin(c + dx)}{5d \cos^{5/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{2aA \sin(c + dx)}{7d \cos^{7/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow \text{3251} \\
& \frac{1}{7}(6A + \\
7B) & \left(\frac{4}{5} \left(\frac{2}{3} \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{3/2}(c + dx)} dx + \frac{2a \sin(c + dx)}{3d \cos^{3/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \frac{2a \sin(c + dx)}{5d \cos^{5/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{2aA \sin(c + dx)}{7d \cos^{7/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7}(6A + \\
7B) & \left(\frac{4}{5} \left(\frac{2}{3} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a \sin(c + dx)}{3d \cos^{3/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \frac{2a \sin(c + dx)}{5d \cos^{5/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{2aA \sin(c + dx)}{7d \cos^{7/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow \text{3250} \\
& \frac{1}{7}(6A + \\
7B) & \left(\frac{2a \sin(c + dx)}{5d \cos^{5/2}(c + dx) \sqrt{a \cos(c + dx) + a}} + \frac{4}{5} \left(\frac{2a \sin(c + dx)}{3d \cos^{3/2}(c + dx) \sqrt{a \cos(c + dx) + a}} + \frac{4a \sin(c + dx)}{3d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx)}{7d \cos^{7/2}(c + dx) \sqrt{a \cos(c + dx) + a}} \right)
\end{aligned}$$

input `Int[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `(2*a*A*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) + ((6*A + 7*B)*((2*a*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]])) + (4*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]])) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/5)/7`

3.173.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.173.4 Maple [A] (verified)

Time = 7.38 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.62

method	result
default	$\frac{2 \sin(dx+c)(48A(\cos^3(dx+c))+56B(\cos^3(dx+c))+24A(\cos^2(dx+c))+28B(\cos^2(dx+c))+18A \cos(dx+c)+21B \cos(dx+c)+15A)}{105d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}}$
parts	$\frac{2A \sin(dx+c)(16(\cos^3(dx+c))+8(\cos^2(dx+c))+6 \cos(dx+c)+5) \sqrt{a(1+\cos(dx+c))}}{35d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}} + \frac{2B \sin(dx+c)(8(\cos^2(dx+c))+4 \cos(dx+c))}{15d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}}$

3.173.
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

input `int((a+cos(d*x+c)*a)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RETURNVERBOSE)`

output $\frac{2}{105}d\sin(dx+c)\cdot(48A\cos(dx+c)^3+56B\cos(dx+c)^3+24A\cos(dx+c)^2+28B\cos(dx+c)^2+18A\cos(dx+c)+21B\cos(dx+c)+15A)\cdot(a\cdot(1+\cos(dx+c)))^{1/2}/(1+\cos(dx+c))/\cos(dx+c)^{7/2}$

3.173.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{a+a\cos(c+dx)}(A+B\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2(8(6A+7B)\cos(dx+c)^3+4(6A+7B)\cos(dx+c)^2+3(6A+7B)\cos(dx+c)+15A)\sqrt{a\cos(dx+c)}}{105(d\cos(dx+c)^5+d\cos(dx+c)^4)}$$

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,algorithm="fricas")`

output $\frac{2}{105}\cdot(8\cdot(6A+7B)\cdot\cos(dx+c)^3+4\cdot(6A+7B)\cdot\cos(dx+c)^2+3\cdot(6A+7B)\cdot\cos(dx+c)+15A)\cdot\sqrt{a\cdot\cos(dx+c)+a}\cdot\sqrt{\cos(dx+c)}\cdot\sin(dx+c)/(d\cdot\cos(dx+c)^5+d\cdot\cos(dx+c)^4)$

3.173.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a+a\cos(c+dx)}(A+B\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output `Timed out`

3.173.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 522 vs. $2(151) = 302$.

Time = 0.35 (sec) , antiderivative size = 522, normalized size of antiderivative = 2.98

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{2 \left(\frac{7B \left(\frac{15\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{25\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{17\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{7\sqrt{2}\sqrt{a}\sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^3}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(\frac{3\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + 1 \right)} + \frac{3A \left(\frac{35\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{70\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{84\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{58\sqrt{2}\sqrt{a}\sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{9\sqrt{2}\sqrt{a}\sin(dx+c)^9}{(\cos(dx+c)+1)^9} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^{\frac{9}{2}}}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{9}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{9}{2}} \left(\frac{4\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{6\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + \frac{\sin(dx+c)^8}{(\cos(dx+c)+1)^8} + 1 \right)} \right) d$$

105 d

input `integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="maxima")`

output `2/105*(7*B*(15*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 25*sqrt(2)
)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 17*sqrt(2)*sqrt(a)*sin(d*x
+ c)^5/(cos(d*x + c) + 1)^5 - 7*sqrt(2)*sqrt(a)*sin(d*x + c)^7/(cos(d*x +
c) + 1)^7)*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^3/((sin(d*x + c)/(co
s(d*x + c) + 1) + 1)^(7/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(3
*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*sin(d*x + c)^4/(cos(d*x + c) + 1)
^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1)) + 3*A*(35*sqrt(2)*sqrt(a)*s
in(d*x + c)/(cos(d*x + c) + 1) - 70*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*
x + c) + 1)^3 + 84*sqrt(2)*sqrt(a)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 5
8*sqrt(2)*sqrt(a)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 9*sqrt(2)*sqrt(a)*
sin(d*x + c)^9/(cos(d*x + c) + 1)^9)*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2
+ 1)^4/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(9/2)*(-sin(d*x + c)/(cos(d*
x + c) + 1) + 1)^(9/2)*(4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 6*sin(d*x
+ c)^4/(cos(d*x + c) + 1)^4 + 4*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + sin(
d*x + c)^8/(cos(d*x + c) + 1)^8 + 1)))/d`

3.173.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 237.66 (sec) , antiderivative size = 136951, normalized size of antiderivative = 782.58

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Too large to display}$$

3.173. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

```
input integrate((a+a*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="giac")
```

```
output -134217728/105*sqrt(2)*sqrt(-tan(1/4*d*x + c)^4*tan(1/2*c)^8 + 14*tan(1/4*
d*x + c)^4*tan(1/2*c)^6 - 24*tan(1/4*d*x + c)^3*tan(1/2*c)^7 + 6*tan(1/4*d
*x + c)^2*tan(1/2*c)^8 + 56*tan(1/4*d*x + c)^3*tan(1/2*c)^5 - 84*tan(1/4*d
*x + c)^2*tan(1/2*c)^6 + 24*tan(1/4*d*x + c)*tan(1/2*c)^7 - tan(1/2*c)^8 -
14*tan(1/4*d*x + c)^4*tan(1/2*c)^2 + 56*tan(1/4*d*x + c)^3*tan(1/2*c)^3 -
56*tan(1/4*d*x + c)*tan(1/2*c)^5 + 14*tan(1/2*c)^6 + tan(1/4*d*x + c)^4 -
24*tan(1/4*d*x + c)^3*tan(1/2*c) + 84*tan(1/4*d*x + c)^2*tan(1/2*c)^2 - 5
6*tan(1/4*d*x + c)*tan(1/2*c)^3 - 6*tan(1/4*d*x + c)^2 + 24*tan(1/4*d*x +
c)*tan(1/2*c) - 14*tan(1/2*c)^2 + 1)*(((((((((((((((((((12*I*A*e^(1071/2*I*c)
+ 14*I*B*e^(1071/2*I*c) + 4944*I*A*e^(1069/2*I*c) + 5768*I*B*e^(1069/2*I*c)
) + 1015992*I*A*e^(1067/2*I*c) + 1185324*I*B*e^(1067/2*I*c) + 138852240*I*
A*e^(1065/2*I*c) + 161994280*I*B*e^(1065/2*I*c) + 14197641540*I*A*e^(1063/
2*I*c) + 16563915130*I*B*e^(1063/2*I*c) + 1158527549664*I*A*e^(1061/2*I*c)
+ 1351615474608*I*B*e^(1061/2*I*c) + 78586785452286*I*A*e^(1059/2*I*c) +
91684583027667*I*B*e^(1059/2*I*c) + 4558033556260200*I*A*e^(1057/2*I*c) +
5317705815636900*I*B*e^(1057/2*I*c) + 230750448790649688*I*A*e^(1055/2*I*c)
) + 269208856922424636*I*B*e^(1055/2*I*c) + 10358131257430815000*I*A*e^(10
53/2*I*c) + 12084486467002617465*I*B*e^(1053/2*I*c) + 41743268973037417024
2*I*A*e^(1051/2*I*c) + 487004804685436517529*I*B*e^(1051/2*I*c) + 15255267
392486063535840*I*A*e^(1049/2*I*c) + 17797811957900404495170*I*B*e^(104...
```

3.173.9 Mupad [B] (verification not implemented)

Time = 6.44 (sec) , antiderivative size = 479, normalized size of antiderivative = 2.74

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{a + a \left(\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2} \right)} \left(\frac{96A + 112B}{105d} \right)}{\sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}} + e^{c \operatorname{li} + dx \operatorname{li}} \sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}} + 3e^{c 2i + dx 2i} \sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}} + 3e^{c 3i + dx 3i}}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/cos(c + d*x)^(9/2),x
)
```


output $((a + a*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2))^{(1/2)*(((96*A + 112*B)*1i)/(105*d) - (B*\exp(c*3i + d*x*3i)*8i)/(3*d) + (B*\exp(c*4i + d*x*4i)*8i)/(3*d) - (\exp(c*7i + d*x*7i)*(96*A + 112*B)*1i)/(105*d) + (\exp(c*2i + d*x*2i)*(336*A + 392*B)*1i)/(105*d) - (\exp(c*5i + d*x*5i)*(336*A + 392*B)*1i)/(105*d)))/((\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + \exp(c*1i + d*x*1i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*2i + d*x*2i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*3i + d*x*3i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*4i + d*x*4i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*5i + d*x*5i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + \exp(c*6i + d*x*6i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + \exp(c*7i + d*x*7i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2}))$

3.173. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

3.174 $\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx)) dx$

3.174.1 Optimal result	1667
3.174.2 Mathematica [A] (verified)	1668
3.174.3 Rubi [A] (verified)	1668
3.174.4 Maple [A] (verified)	1672
3.174.5 Fricas [A] (verification not implemented)	1673
3.174.6 Sympy [F(-1)]	1673
3.174.7 Maxima [B] (verification not implemented)	1673
3.174.8 Giac [F]	1674
3.174.9 Mupad [F(-1)]	1675

3.174.1 Optimal result

Integrand size = 35, antiderivative size = 227

$$\int \cos^{\frac{3}{2}}(c+dx)(a + a \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx)) dx = \frac{a^{3/2}(88A+75B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{64d} + \frac{a^2(88A+75B) \sqrt{\cos(c+dx)} \sin(c+dx)}{64d \sqrt{a+a \cos(c+dx)}} + \frac{a^2(88A+75B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{96d \sqrt{a+a \cos(c+dx)}} + \frac{a^2(8A+9B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{24d \sqrt{a+a \cos(c+dx)}} + \frac{aB \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{4d}$$

output

```
1/64*a^(3/2)*(88*A+75*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))
/d+1/96*a^2*(88*A+75*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
+1/24*a^2*(8*A+9*B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
+1/64*a^2*(88*A+75*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
+1/4*a*B*cos(d*x+c)^(5/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.174.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.60

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx = \frac{a\sqrt{a(1+\cos(c+dx))}\sec\left(\frac{1}{2}(c+dx)\right)\left(3\sqrt{2}(88A+75B)\arcsin\left(\sqrt{2}\sin\left(\frac{1}{2}(c+dx)\right)\right)+2\sqrt{2}(88A+93B)\cos\left(\frac{1}{2}(c+dx)\right)+4(8A+15B)\cos(c+dx)+12B\cos\left(\frac{3}{2}(c+dx)\right)\right)}{384d}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(3*Sqrt[2]*(88*A + 75*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(296*A + 285*B + 2*(88*A + 93*B)*Cos[c + d*x] + 4*(8*A + 15*B)*Cos[2*(c + d*x)] + 12*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2])/ (384*d)`

3.174.3 Rubi [A] (verified)Time = 1.08 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.96, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}(A+B\cos(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^{3/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\ & \quad \downarrow \text{3455} \\ & \frac{1}{4}\int\frac{1}{2}\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a(a(8A+5B)+a(8A+9B)\cos(c+dx))}dx+ \\ & \quad \frac{aB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}{4d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.174. $\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx$

$$\frac{1}{8} \int \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a} (a(8A+5B) + a(8A+9B) \cos(c+dx)) dx + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{4d}$$

↓ 3042

$$\frac{1}{8} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} (a(8A+5B) + a(8A+9B) \sin\left(c+dx+\frac{\pi}{2}\right)) dx + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{4d}$$

↓ 3460

$$\frac{1}{8} \left(\frac{1}{6} a(88A+75B) \int \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{a^2(8A+9B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx) + a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} a(88A+75B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{a^2(8A+9B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx) + a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{4d}$$

↓ 3249

$$\frac{1}{8} \left(\frac{1}{6} a(88A+75B) \left(\frac{3}{4} \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+adx} + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx) + a}} \right) + \frac{a^2(8A+9B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx) + a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} a(88A+75B) \left(\frac{3}{4} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx) + a}} \right) + \frac{a^2(8A+9B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx) + a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{4d}$$

↓ 3249

3.174. $\int \cos^{\frac{3}{2}}(c+dx) (a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx)) dx$

$$\frac{1}{8} \left(\frac{1}{6} a(88A + 75B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) \right. \\ \left. \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}}{4d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{8} \left(\frac{1}{6} a(88A + 75B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) \right. \\ \left. \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}}{4d} \right) \\ \downarrow \text{3253}$$

$$\frac{1}{8} \left(\frac{1}{6} a(88A + 75B) \left(\frac{3}{4} \left(\frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{d} \right) + \frac{a \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d \sqrt{a \cos(c+dx)+a}} \right) \right. \\ \left. \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}}{4d} \right) \\ \downarrow \text{223}$$

$$\frac{1}{8} \left(\frac{a^2(8A + 9B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{3d \sqrt{a \cos(c+dx)+a}} + \frac{1}{6} a(88A + 75B) \left(\frac{3}{4} \left(\frac{\sqrt{a} \arcsin \left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}} \right)}{d} + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) \right) \right. \\ \left. \frac{aB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}}{4d} \right)$$

input `Int[Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(4*d) + ((a^2*(8*A + 9*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (a*(88*A + 75*B)*((a*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]])) + (3*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))))/4)/6)/8`

3.174.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3249 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`
- rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.174.4 Maple [A] (verified)

Time = 15.60 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.54

method	result
default	$a \left(48B(\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 64A(\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 120B(\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)$
parts	$\frac{A \left(8 \sin(dx+c) (\cos^2(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 22 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 33 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 33 \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right)}{24d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c)), x, method=_RET
URNVERBOSE)
```

```
output 1/192*a/d*(48*B*cos(d*x+c)^3*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+
64*A*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+120*B*cos(d
*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+176*A*cos(d*x+c)*sin(
d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+150*B*cos(d*x+c)*sin(d*x+c)*(cos(
d*x+c)/(1+cos(d*x+c)))^(1/2)+264*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d
*x+c)+225*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+264*A*arctan(tan(
d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+225*B*arctan(tan(d*x+c)*(cos(d*x
+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+c
os(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

3.174. $\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

3.174.5 Fricas [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.71

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \frac{(48Ba\cos(dx+c))^3 + 8(8A+15B)a\cos(dx+c)^2 + 2(88A+75B)a\cos(dx+c) + (48Ba\cos(dx+c))^3 + 8(8A+15B)a\cos(dx+c)^2 + 2(88A+75B)a\cos(dx+c) + (48Ba\cos(dx+c))^3 + 8(8A+15B)a\cos(dx+c)^2 + 2(88A+75B)a\cos(dx+c) + \dots}{(48Ba\cos(dx+c))^3 + 8(8A+15B)a\cos(dx+c)^2 + 2(88A+75B)a\cos(dx+c) + \dots}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algo rithm="fricas")`

output `1/192*((48*B*a*cos(d*x + c)^3 + 8*(8*A + 15*B)*a*cos(d*x + c)^2 + 2*(88*A + 75*B)*a*cos(d*x + c) + 3*(88*A + 75*B)*a)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - 3*((88*A + 75*B)*a*cos(d*x + c) + (88*A + 75*B)*a)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c)))/(d*cos(d*x + c) + d)`

3.174.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.174.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8904 vs. 2(195) = 390.

Time = 0.93 (sec) , antiderivative size = 8904, normalized size of antiderivative = 39.22

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algo
rithm="maxima")`

output `1/768*(8*(4*(a*cos(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x
+ 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1))*sin(
3*d*x + 3*c) - (a*cos(3*d*x + 3*c) - a)*sin(3/2*arctan2(sin(2/3*arctan2(si
n(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(
3*d*x + 3*c))) + 1))*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))
)^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*a
rctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(3/4)*sqrt(a) + 6*(cos(2/
3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d
*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(
3*d*x + 3*c))) + 1)^(1/4)*((3*a*sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*
x + 3*c))) + 11*a*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*co
s(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/
3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1) - (3*a*cos(2/3*arctan
2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 5*a*cos(1/3*arctan2(sin(3*d*x + 3
*c), cos(3*d*x + 3*c))) - 8*a)*sin(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x +
3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3
*c))) + 1))*sqrt(a) + 33*(a*arctan2(-(cos(2/3*arctan2(sin(3*d*x + 3*c), c
os(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))
^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(1/4)*(co
s(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos...`

3.174.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3/
2), x)`

3.174.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^{3/2}(A + B \cos(c + dx))(a + a \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)`

3.175 $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

3.175.1 Optimal result	1676
3.175.2 Mathematica [A] (verified)	1677
3.175.3 Rubi [A] (verified)	1677
3.175.4 Maple [A] (verified)	1681
3.175.5 Fricas [A] (verification not implemented)	1681
3.175.6 Sympy [F(-1)]	1682
3.175.7 Maxima [B] (verification not implemented)	1682
3.175.8 Giac [F]	1683
3.175.9 Mupad [F(-1)]	1684

3.175.1 Optimal result

Integrand size = 35, antiderivative size = 180

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{a^{3/2}(14A + 11B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{8d} + \frac{a^2(14A + 11B)\sqrt{\cos(c + dx)} \sin(c + dx)}{8d\sqrt{a + a \cos(c + dx)}} + \frac{a^2(6A + 7B) \cos^{3/2}(c + dx) \sin(c + dx)}{12d\sqrt{a + a \cos(c + dx)}} + \frac{aB \cos^{3/2}(c + dx)\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{3d}$$

output

```
1/8*a^(3/2)*(14*A+11*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/
d+1/12*a^2*(6*A+7*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+
1/8*a^2*(14*A+11*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)+1
/3*a*B*cos(d*x+c)^(3/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.175.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.66

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx = \frac{a\sqrt{a(1+\cos(c+dx))}\sec\left(\frac{1}{2}(c+dx)\right)\left(3\sqrt{2}(14A+11B)\arcsin\left(\sqrt{2}\sin\left(\frac{1}{2}(c+dx)\right)\right)+2\sqrt{2}(42A+37B+2(6A+11B)\cos(c+dx))+4B\cos(2(c+dx))\sin\left(\frac{1}{2}(c+dx)\right)\right)}{48d}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(3*Sqrt[2]*(14*A + 11*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(42*A + 37*B + 2*(6*A + 11*B)*Cos[c + d*x] + 4*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.175.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}(A+B\cos(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^{3/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\ & \quad \downarrow \text{3455} \\ & \frac{1}{3}\int\frac{1}{2}\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}\left(3a(2A+B)+a(6A+7B)\cos(c+dx)\right)dx+ \\ & \quad \frac{aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}{3d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.175. $\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx$

$$\frac{1}{6} \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a + a(3a(2A+B) + a(6A+7B)\cos(c+dx))} dx + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{3d}$$

↓ 3042

$$\frac{1}{6} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a + a(3a(2A+B) + a(6A+7B)\sin\left(c+dx+\frac{\pi}{2}\right))} dx + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{3d}$$

↓ 3460

$$\frac{1}{6} \left(\frac{3}{4} a(14A+11B) \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a + adx} + \frac{a^2(6A+7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3}{4} a(14A+11B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a + adx} + \frac{a^2(6A+7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{3d}$$

↓ 3249

$$\frac{1}{6} \left(\frac{3}{4} a(14A+11B) \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(6A+7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3}{4} a(14A+11B) \left(\frac{1}{2} \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(6A+7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}{3d}$$

↓ 3253

$$\frac{1}{6} \left(\frac{3}{4} a(14A + 11B) \left(\frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx) a + a}}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right)}{d} \right) + \frac{a^2(6A + 7B) \sin(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right. \\ \left. + \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right)$$

↓ 223

$$\frac{1}{6} \left(\frac{a^2(6A + 7B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} + \frac{3}{4} a(14A + 11B) \left(\frac{\sqrt{a} \arcsin \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. + \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} \right)$$

input `Int[Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(a*B*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((a^2*(6*A + 7*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*a*(14*A + 11*B)*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x])))/4)/6`

3.175.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3249 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.175.4 Maple [A] (verified)

Time = 15.19 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.54

method	result
default	$a \left(8B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 22B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 42A \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 7 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 7 \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right) \sqrt{a(1+\cos(dx+c))} (\sqrt{\cos(dx+c)})$
parts	$\frac{A \left(2 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 7 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 7 \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right) \sqrt{a(1+\cos(dx+c))} (\sqrt{\cos(dx+c)})}{4d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

input `int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output $1/24*a/d*(8*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+12*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+22*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+42*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+33*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+42*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+33*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)$

3.175.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.80

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx = \frac{(8Ba\cos(dx+c)^2+2(6A+11B)a\cos(dx+c)+3(14A+11B)a)\sqrt{a\cos(dx+c)}}{d\cos(dx+c)+d}$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x,algorithm="fracas")`

output $1/24*((8*B*a*cos(d*x+c)^2+2*(6*A+11*B)*a*cos(d*x+c)+3*(14*A+11*B)*a)*sqrt(a*cos(d*x+c)+a)*sqrt(cos(d*x+c))*sin(d*x+c)-3*((14*A+11*B)*a*cos(d*x+c)+(14*A+11*B)*a)*sqrt(a)*arctan(sqrt(a*cos(d*x+c)+a)*sqrt(cos(d*x+c))/(sqrt(a)*sin(d*x+c)))/(d*cos(d*x+c)+d)$

3.175. $\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx$

3.175.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.175.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3023 vs. 2(154) = 308.

Time = 0.66 (sec) , antiderivative size = 3023, normalized size of antiderivative = 16.79

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output

```

1/96*(6*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) +
1)^(1/4))*((a*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))*sin(2*d
*x + 2*c) + a*sin(2*d*x + 2*c) - (a*cos(2*d*x + 2*c) - 6*a)*sin(1/2*arctan
2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*cos(1/2*arctan2(sin(2*d*x + 2*c),
cos(2*d*x + 2*c) + 1)) + (a*sin(2*d*x + 2*c)*sin(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c)))) - a*cos(2*d*x + 2*c) + (a*cos(2*d*x + 2*c) - 6*a)*
cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 6*a)*sin(1/2*arctan
2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(a) + 7*(a*arctan2((cos(2*
d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2
*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))*sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(
2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1
/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*
d*x + 2*c), cos(2*d*x + 2*c))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d
*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1)
- a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c)
+ 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))*sin(1/2*
arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d
*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos...

```

3.175.8 Giac [F]

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^{3/2} \sqrt{\cos(dx + c)} dx$$

input

```

integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algo
rithm="giac")

```

output

```

integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(3/2)*sqrt(cos(d*x + c
)), x)

```

3.175.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx = \int \sqrt{\cos(c+dx)}(A+B\cos(c+dx))(a+a\cos(c+dx))^{3/2}dx$$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2), x)`

3.176
$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.176.1 Optimal result 1685
 3.176.2 Mathematica [A] (verified) 1685
 3.176.3 Rubi [A] (verified) 1686
 3.176.4 Maple [A] (verified) 1689
 3.176.5 Fricas [A] (verification not implemented) 1689
 3.176.6 Sympy [F] 1690
 3.176.7 Maxima [B] (verification not implemented) 1690
 3.176.8 Giac [F(-1)] 1691
 3.176.9 Mupad [F(-1)] 1692

3.176.1 Optimal result

Integrand size = 35, antiderivative size = 133

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx = \frac{a^{3/2}(12A+7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4d} + \frac{a^2(4A+5B)\sqrt{\cos(c+dx)} \sin(c+dx)}{4d\sqrt{a+a \cos(c+dx)}} + \frac{aB\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{2d}$$

output `1/4*a^(3/2)*(12*A+7*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d + 1/4*a^2*(4*A+5*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)+1/2*a*B*sin(d*x+c)*cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(1/2)/d`

3.176.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.76

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx = \frac{a\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(\sqrt{2}(12A+7B) a\right)}{\dots}$$

input `Integrate[((a+a*Cos[c+d*x])^(3/2)*(A+B*Cos[c+d*x]))/Sqrt[Cos[c+d*x]],x]`

output $(a*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])] * \text{Sec}[(c + d*x)/2] * (\text{Sqrt}[2]*(12*A + 7*B)*\text{ArcSin}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]] + 2*\text{Sqrt}[\text{Cos}[c + d*x]]*(4*A + 7*B + 2*B*\text{Cos}[c + d*x])*\text{Sin}[(c + d*x)/2]))/(8*d)$

3.176.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3455, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3455

$$\frac{1}{2} \int \frac{\sqrt{\cos(c + dx)a + a(a(4A + B) + a(4A + 5B) \cos(c + dx))}}{2\sqrt{\cos(c + dx)}} dx + \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d}$$

↓ 27

$$\frac{1}{4} \int \frac{\sqrt{\cos(c + dx)a + a(a(4A + B) + a(4A + 5B) \cos(c + dx))}}{\sqrt{\cos(c + dx)}} dx + \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d}$$

↓ 3042

$$\frac{1}{4} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a(a(4A + B) + a(4A + 5B) \sin(c + dx + \frac{\pi}{2}))}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d}$$

↓ 3460

$$\begin{aligned}
& \frac{1}{4} \left(\frac{1}{2} a(12A + 7B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{a^2(4A + 5B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left(\frac{1}{2} a(12A + 7B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{a^2(4A + 5B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} \\
& \quad \downarrow \text{3253} \\
& \frac{1}{4} \left(\frac{a^2(4A + 5B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} - \frac{a(12A + 7B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx)a + a}}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}} \right)}{d} \right) + \\
& \quad \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} \\
& \quad \downarrow \text{223} \\
& \frac{1}{4} \left(\frac{a^{3/2}(12A + 7B) \arcsin \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} + \frac{a^2(4A + 5B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} \right) + \\
& \quad \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d}
\end{aligned}$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(a*B*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + ((a^(3/2)*(12*A + 7*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a^2*(4*A + 5*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4`

3.176.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.176.4 Maple [A] (verified)

Time = 18.61 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.56

method	result
default	$\frac{a \left(2B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 7B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{4d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \left(3 \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + \cos(dx+c) \sin(dx+c) + 3 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{d \sqrt{\cos(dx+c)} (1+\cos(dx+c))}$

input `int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `1/4*a/d*(2*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+7*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+12*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))+7*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)`

3.176.5 Fracas [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.94

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{(2Ba \cos(dx + c) + (4A + 7B)a) \sqrt{a \cos(dx + c) + a}}{\dots}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,algorithm="fricas")`

output `1/4*((2*B*a*cos(d*x + c) + (4*A + 7*B)*a)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - ((12*A + 7*B)*a*cos(d*x + c) + (12*A + 7*B)*a)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/(d*cos(d*x + c) + d)`

3.176.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(a(\cos(c + dx) + 1))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Integral((a*(cos(c + d*x) + 1))**(3/2)*(A + B*cos(c + d*x))/sqrt(cos(c + d*x)), x)`

3.176.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1884 vs. $2(113) = 226$.

Time = 0.55 (sec) , antiderivative size = 1884, normalized size of antiderivative = 14.17

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/16*(4*(2*(a*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (a*cos(d*x + c) - a)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sqrt(a) + 3*(a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) + 1) - a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + 1) + a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x ...`

3.176.8 Giac [**F(-1)**]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.176.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(1/2),x)`

$$3.177 \quad \int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.177.1 Optimal result 1693
 3.177.2 Mathematica [A] (verified) 1693
 3.177.3 Rubi [A] (verified) 1694
 3.177.4 Maple [A] (verified) 1697
 3.177.5 Fricas [A] (verification not implemented) 1697
 3.177.6 Sympy [F] 1698
 3.177.7 Maxima [B] (verification not implemented) 1698
 3.177.8 Giac [F(-1)] 1699
 3.177.9 Mupad [F(-1)] 1700

3.177.1 Optimal result

Integrand size = 35, antiderivative size = 126

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{a^{3/2}(2A+3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} - \frac{a^2(2A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{a+a \cos(c+dx)}} + \frac{2aA\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}}$$

```
output a^(3/2)*(2*A+3*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d-a^2*(2*A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)+2*a*A*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)
```

3.177.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.85

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{a\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(\sqrt{2}(2A+3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)\right)}{d}$$

```
input Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]
```

output $(a*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])] * \text{Sec}[(c + d*x)/2] * (\text{Sqrt}[2]*(2*A + 3*B) * \text{ArcSin}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]] * \text{Sqrt}[\text{Cos}[c + d*x]] + 2*(2*A + B*\text{Cos}[c + d*x]) * \text{Sin}[(c + d*x)/2])) / (2*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

3.177.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3454, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

↓ 3454

$$2 \int \frac{\sqrt{\cos(c + dx)a + a(a(2A + B) - a(2A - B) \cos(c + dx))}}{2\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}}$$

↓ 27

$$\int \frac{\sqrt{\cos(c + dx)a + a(a(2A + B) - a(2A - B) \cos(c + dx))}}{\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}}$$

↓ 3042

$$\int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a(a(2A + B) - a(2A - B) \sin(c + dx + \frac{\pi}{2}))}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}}$$

↓ 3460

3.177. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

$$\begin{aligned}
& \frac{1}{2}a(2A + 3B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx - \frac{a^2(2A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2}a(2A + 3B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{a^2(2A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3253} \\
& \frac{a(2A + 3B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx)a + a}}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} - \frac{a^2(2A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{223} \\
& \frac{a^{3/2}(2A + 3B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} - \frac{a^2(2A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{d\sqrt{\cos(c + dx)}}
\end{aligned}$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2), x]`

output `(a^(3/2)*(2*A + 3*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d - (a^2*(2*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]) + (2*a*A*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])`

3.177.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.177.4 Maple [A] (verified)

Time = 18.35 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.50

method	result
default	$\frac{a \left(B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 2A \cos(dx+c) \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + 2A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 3B \cos(dx+c) \right)}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sqrt{\cos(dx+c)}}$
parts	$\frac{2A \left(\cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + \sin(dx+c) \right)}{d(1+\cos(dx+c)) \sqrt{\cos(dx+c)}}$

input `int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `a/d*(B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2*A*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+2*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+3*B*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/cos(d*x+c)^(1/2)`

3.177.5 Fracas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.07

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \frac{(Ba \cos(dx + c) + 2Aa) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}}{\cos^{3/2}(c + dx)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,algorithm="fracas")`

output `((B*a*cos(d*x + c) + 2*A*a)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - ((2*A + 3*B)*a*cos(d*x + c)^2 + (2*A + 3*B)*a*cos(d*x + c))*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))))/(d*cos(d*x + c)^2 + d*cos(d*x + c))`

3.177.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(a(\cos(c + dx) + 1))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Integral((a*(cos(c + d*x) + 1))**(3/2)*(A + B*cos(c + d*x))/cos(c + d*x)**(3/2), x)`

3.177.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1801 vs. $2(112) = 224$.

Time = 0.54 (sec) , antiderivative size = 1801, normalized size of antiderivative = 14.29

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/4*((2*(a*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (a*cos(d*x + c) - a)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sqrt(a) + 3*(a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) + 1) - a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + 1) + a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2...`

3.177.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorith="giac")`

output `Timed out`

3.177.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(3/2),x)`

$$3.178 \quad \int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$$

3.178.1 Optimal result 1701
 3.178.2 Mathematica [A] (verified) 1701
 3.178.3 Rubi [A] (verified) 1702
 3.178.4 Maple [A] (verified) 1704
 3.178.5 Fricas [A] (verification not implemented) 1705
 3.178.6 Sympy [F] 1705
 3.178.7 Maxima [B] (verification not implemented) 1706
 3.178.8 Giac [F(-1)] 1706
 3.178.9 Mupad [F(-1)] 1707

3.178.1 Optimal result

Integrand size = 35, antiderivative size = 125

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx = \frac{2a^{3/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} + \frac{2a^2(4A+3B) \sin(c+dx)}{3d\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}} + \frac{2aA\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{3d \cos^{3/2}(c+dx)}$$

output `2*a^(3/2)*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+2/3*a^2*(4*A+3*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)+2/3*a*A*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)`

3.178.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.85

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx = \frac{a\sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(3\sqrt{2}B \arcsin\left(\sqrt{\frac{1-\cos(c+dx)}{2}}\right) + (A+B \cos(c+dx))\right)}{\cos^{5/2}(c+dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output $(a*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*\text{Sec}[(c + d*x)/2]*(3*\text{Sqrt}[2]*B*\text{ArcSin}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]]*\text{Cos}[c + d*x]^{(3/2)} + 2*(A + (5*A + 3*B)*\text{Cos}[c + d*x]))*\text{Sin}[(c + d*x)/2])/(3*d*\text{Cos}[c + d*x]^{(3/2)})$

3.178.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.02, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3454

$$\frac{2}{3} \int \frac{\sqrt{\cos(c + dx)a + a(a(4A + 3B) + 3aB \cos(c + dx))}}{2 \cos^{3/2}(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d \cos^{3/2}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{\sqrt{\cos(c + dx)a + a(a(4A + 3B) + 3aB \cos(c + dx))}}{\cos^{3/2}(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d \cos^{3/2}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a(a(4A + 3B) + 3aB \sin(c + dx + \frac{\pi}{2}))}}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d \cos^{3/2}(c + dx)}$$

↓ 3459

$$\frac{1}{3} \left(3aB \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{2a^2(4A + 3B) \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{3d \cos^{3/2}(c + dx)}$$

3.178. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{1}{3} \left(3aB \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right) + \\
& \quad \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d\cos^{\frac{3}{2}}(c+dx)} \\
& \downarrow \text{3253} \\
& \frac{1}{3} \left(\frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{6aB \int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} \right) + \\
& \quad \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d\cos^{\frac{3}{2}}(c+dx)} \\
& \downarrow \text{223} \\
& \frac{1}{3} \left(\frac{6a^{3/2}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} + \frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right) + \\
& \quad \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d\cos^{\frac{3}{2}}(c+dx)}
\end{aligned}$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*a*A*sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((6*a^(3/2)*B*ArcSin[(sqrt[a]*Sin[c + d*x])/sqrt[a + a*Cos[c + d*x]]])/d + (2*a^2*(4*A + 3*B)*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]]*sqrt[a + a*Cos[c + d*x]]))/3`

3.178.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.178. $\int \frac{(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.178.4 Maple [A] (verified)

Time = 7.80 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.50

method	result
default	$\frac{2a \left(3B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) (\cos^2(dx+c)) + 3B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \cos(dx+c) \right)}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}}$
parts	$\frac{2A \sin(dx+c)(5 \cos(dx+c)+1) \sqrt{a(1+\cos(dx+c))} a}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}} + \frac{2B \left(\cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{d(1+\cos(dx+c))}$

3.178.
$$\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 2/3*a/d*(3*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arctan(tan(d*x+c)*(cos(d*x+
c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^2+3*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)+5*A*sin
(d*x+c)*cos(d*x+c)+3*B*sin(d*x+c)*cos(d*x+c)+A*sin(d*x+c))*(a*(1+cos(d*x+c
)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(3/2)
```

3.178.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.06

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \frac{2 \left(((5A + 3B)a \cos(dx + c) + Aa) \sqrt{a \cos(dx + c) + a} \right)}{\cos^{5/2}(c + dx)}$$

```
input integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algo
rithm="fricas")
```

```
output 2/3*(((5*A + 3*B)*a*cos(d*x + c) + A*a)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(
d*x + c))*sin(d*x + c) - 3*(B*a*cos(d*x + c)^3 + B*a*cos(d*x + c)^2)*sqrt(
a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c
))))/(d*cos(d*x + c)^3 + d*cos(d*x + c)^2)
```

3.178.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(a(\cos(c + dx) + 1))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx$$

```
input integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)
```

```
output Integral((a*(cos(c + d*x) + 1))**(3/2)*(A + B*cos(c + d*x))/cos(c + d*x)**
(5/2), x)
```

3.178. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$

3.178.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1124 vs. $2(107) = 214$.

Time = 0.45 (sec) , antiderivative size = 1124, normalized size of antiderivative = 8.99

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/6*(3*((a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*...`

3.178.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`

3.178. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$

output Timed out

3.178.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\cos(c + dx)^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(5/2),x)`

3.179
$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

3.179.1 Optimal result 1708
 3.179.2 Mathematica [A] (verified) 1708
 3.179.3 Rubi [A] (verified) 1709
 3.179.4 Maple [A] (verified) 1711
 3.179.5 Fricas [A] (verification not implemented) 1712
 3.179.6 Sympy [F(-1)] 1712
 3.179.7 Maxima [B] (verification not implemented) 1712
 3.179.8 Giac [F(-1)] 1713
 3.179.9 Mupad [B] (verification not implemented) 1713

3.179.1 Optimal result

Integrand size = 35, antiderivative size = 134

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \frac{2a^2(6A + 5B) \sin(c + dx)}{15d \cos^{3/2}(c + dx) \sqrt{a + a \cos(c + dx)}} + \frac{2a^2(18A + 25B) \sin(c + dx)}{15d \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} + \frac{2aA \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{5d \cos^{5/2}(c + dx)}$$

output `2/15*a^2*(6*A+5*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+2/15*a^2*(18*A+25*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)+2/5*a*A*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)`

3.179.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.60

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \frac{a \sqrt{a(1 + \cos(c + dx))(24A + 25B + 2(9A + 5B) \cos(c + dx))}}{15d \cos^{5/2}(c + dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

```
output (a*Sqrt[a*(1 + Cos[c + d*x])]*(24*A + 25*B + 2*(9*A + 5*B)*Cos[c + d*x] +
(18*A + 25*B)*Cos[2*(c + d*x)]*Tan[(c + d*x)/2])/(15*d*Cos[c + d*x]^(5/2)
)
```

3.179.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

↓ 3454

$$\frac{2}{5} \int \frac{\sqrt{\cos(c + dx)a + a}(a(6A + 5B) + a(2A + 5B)\cos(c + dx))}{2 \cos^{5/2}(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{5d \cos^{5/2}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{\sqrt{\cos(c + dx)a + a}(a(6A + 5B) + a(2A + 5B)\cos(c + dx))}{\cos^{5/2}(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{5d \cos^{5/2}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}(a(6A + 5B) + a(2A + 5B)\sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{5d \cos^{5/2}(c + dx)}$$

↓ 3459

3.179. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{1}{3} a(18A + 25B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2a^2(6A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} a(18A + 25B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}}} dx + \frac{2a^2(6A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3250

$$\frac{1}{5} \left(\frac{2a^2(6A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} + \frac{2a^2(18A + 25B) \sin(c + dx)}{3d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{5d \cos^{\frac{5}{2}}(c + dx)}$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*a^2*(6*A + 5*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (2*a^2*(18*A + 25*B)*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/5`

3.179.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.179.4 Maple [A] (verified)

Time = 7.44 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{2a \sin(dx+c)(18A(\cos^2(dx+c))+25B(\cos^2(dx+c))+9A \cos(dx+c)+5B \cos(dx+c)+3A)\sqrt{a(1+\cos(dx+c))}}{15d(1+\cos(dx+c)) \cos(dx+c)^{\frac{5}{2}}}$	87
parts	$\frac{2A \sin(dx+c)(6(\cos^2(dx+c))+3 \cos(dx+c)+1)\sqrt{a(1+\cos(dx+c))} a}{5d(1+\cos(dx+c)) \cos(dx+c)^{\frac{5}{2}}} + \frac{2B \sin(dx+c)(5 \cos(dx+c)+1)\sqrt{a(1+\cos(dx+c))} a}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}}$	118

input `int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RET URNVERBOSE)`

$$3.179. \int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

output $2/15*a/d*\sin(d*x+c)*(18*A*\cos(d*x+c)^2+25*B*\cos(d*x+c)^2+9*A*\cos(d*x+c)+5*B*\cos(d*x+c)+3*A)*(a*(1+\cos(d*x+c)))^{1/2}/(1+\cos(d*x+c))/\cos(d*x+c)^{5/2}$

3.179.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.66

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \frac{2((18A + 25B)a \cos(dx + c)^2 + (9A + 5B)a \cos(dx + c) + 3Aa)}{15(d \cos(dx + c))^{3/2}}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="fracas")`

output $2/15*((18*A + 25*B)*a*\cos(d*x + c)^2 + (9*A + 5*B)*a*\cos(d*x + c) + 3*A*a)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^{4 + d*\cos(d*x + c)^3}$

3.179.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output Timed out

3.179.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 344 vs. 2(116) = 232.

Time = 0.34 (sec) , antiderivative size = 344, normalized size of antiderivative = 2.57

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \frac{4 \left(\frac{5 \left(\frac{3\sqrt{2}a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{5\sqrt{2}a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{2\sqrt{2}a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) B}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{5/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{5/2}} + \frac{3 \left(\frac{3\sqrt{2}a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{5\sqrt{2}a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{2\sqrt{2}a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{5/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{5/2}} \right)}{15(d \cos(dx + c))^{3/2}}$$

3.179. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="maxima")`

output `4/15*(5*(3*sqrt(2)*a^(3/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 5*sqrt(2)*a^(
3/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 2*sqrt(2)*a^(3/2)*sin(d*x + c)^
5/(cos(d*x + c) + 1)^5)*B/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(5/2)*(-s
in(d*x + c)/(cos(d*x + c) + 1) + 1)^(5/2)) + 3*(5*sqrt(2)*a^(3/2)*sin(d*x
+ c)/(cos(d*x + c) + 1) - 10*sqrt(2)*a^(3/2)*sin(d*x + c)^3/(cos(d*x + c)
+ 1)^3 + 7*sqrt(2)*a^(3/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 2*sqrt(2)
*a^(3/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)*A*(sin(d*x + c)^2/(cos(d*x +
c) + 1)^2 + 1)^2/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(-sin(d*x +
c)/(cos(d*x + c) + 1) + 1)^(7/2)*(2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 +
sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 1)))/d`

3.179.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="giac")`

output Timed out

3.179.9 Mupad [B] (verification not implemented)

Time = 3.23 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.46

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \frac{2a \sqrt{a (\cos(c + dx) + 1)} (48A \sin(c + dx) + 50B \sin$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(7/2),x
)`

output $(2*a*(a*(\cos(c + d*x) + 1))^{1/2}*(48*A*\sin(c + d*x) + 50*B*\sin(c + d*x) + 36*A*\sin(2*c + 2*d*x) + 66*A*\sin(3*c + 3*d*x) + 18*A*\sin(4*c + 4*d*x) + 18*A*\sin(5*c + 5*d*x) + 20*B*\sin(2*c + 2*d*x) + 75*B*\sin(3*c + 3*d*x) + 10*B*\sin(4*c + 4*d*x) + 25*B*\sin(5*c + 5*d*x)))/(15*d*\cos(c + d*x)^{1/2}*(10*\cos(c + d*x) + 8*\cos(2*c + 2*d*x) + 5*\cos(3*c + 3*d*x) + 2*\cos(4*c + 4*d*x) + \cos(5*c + 5*d*x) + 6))$

3.179. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

$$3.180 \quad \int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.180.1 Optimal result	1715
3.180.2 Mathematica [A] (verified)	1715
3.180.3 Rubi [A] (verified)	1716
3.180.4 Maple [A] (verified)	1719
3.180.5 Fricas [A] (verification not implemented)	1719
3.180.6 Sympy [F(-1)]	1720
3.180.7 Maxima [B] (verification not implemented)	1720
3.180.8 Giac [C] (verification not implemented)	1721
3.180.9 Mupad [B] (verification not implemented)	1722

3.180.1 Optimal result

Integrand size = 35, antiderivative size = 181

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{2a^2(8A+7B) \sin(c+dx)}{35d \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a^2(52A+63B) \sin(c+dx)}{105d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{4a^2(52A+63B) \sin(c+dx)}{105d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} + \frac{2aA \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)}$$

output $2/35*a^2*(8*A+7*B)*\sin(d*x+c)/d/\cos(d*x+c)^{(5/2)}/(a+a*\cos(d*x+c))^{(1/2)}+2/105*a^2*(52*A+63*B)*\sin(d*x+c)/d/\cos(d*x+c)^{(3/2)}/(a+a*\cos(d*x+c))^{(1/2)}+4/105*a^2*(52*A+63*B)*\sin(d*x+c)/d/\cos(d*x+c)^{(1/2)}/(a+a*\cos(d*x+c))^{(1/2)}+2/7*a*A*\sin(d*x+c)*(a+a*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(7/2)}$

3.180.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.56

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{a \sqrt{a(1+\cos(c+dx))}(82A+63B+3(78A+77B) \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

3.180. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

output $(a*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*(82*A + 63*B + 3*(78*A + 77*B)*\text{Cos}[c + d*x] + (52*A + 63*B)*\text{Cos}[2*(c + d*x)] + 52*A*\text{Cos}[3*(c + d*x)] + 63*B*\text{Cos}[3*(c + d*x)])*\text{Tan}[(c + d*x)/2])/(105*d*\text{Cos}[c + d*x]^(7/2))$

3.180.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx$$

↓ 3454

$$\frac{2}{7} \int \frac{\sqrt{\cos(c + dx)a + a}(a(8A + 7B) + a(4A + 7B) \cos(c + dx))}{2 \cos^{7/2}(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{7d \cos^{7/2}(c + dx)}$$

↓ 27

$$\frac{1}{7} \int \frac{\sqrt{\cos(c + dx)a + a}(a(8A + 7B) + a(4A + 7B) \cos(c + dx))}{\cos^{7/2}(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{7d \cos^{7/2}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}(a(8A + 7B) + a(4A + 7B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx + \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{7d \cos^{7/2}(c + dx)}$$

↓ 3459

3.180. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{9/2}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{1}{5} a(52A + 63B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{7d\cos^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} a(52A + 63B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{7d\cos^{\frac{7}{2}}(c+dx)}$$

↓ 3251

$$\frac{1}{7} \left(\frac{1}{5} a(52A + 63B) \left(\frac{2}{3} \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{7d\cos^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} a(52A + 63B) \left(\frac{2}{3} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{7d\cos^{\frac{7}{2}}(c+dx)}$$

↓ 3250

$$\frac{1}{7} \left(\frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} + \frac{1}{5} a(52A + 63B) \left(\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} + \frac{4a\sin(c+dx)}{3d\sqrt{\cos(c+dx)+a}} \right) + \frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{7d\cos^{\frac{7}{2}}(c+dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2), x]`

```
output (2*a*A*Sqrt[a + a*cos[c + d*x]]*sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)) + (
(2*a^2*(8*A + 7*B)*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)*Sqrt[a + a*cos[c
+ d*x]]) + (a*(52*A + 63*B)*((2*a*sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)*Sq
rt[a + a*cos[c + d*x]])) + (4*a*sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[
a + a*cos[c + d*x]])))/5)/7
```

3.180.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3250 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sq
rt[a + b*sin[e + f*x]]*Sqrt[c + d*sin[e + f*x]])), x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3251 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*sin[e
+ f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*sin[e + f*x]])), x] + Sim
p[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*sin[e
+ f*x]]*(c + d*sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*sin[e + f*x])^(m - 1)*((c + d*sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*sin[e + f*x])^(m - 1)*(c + d*sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.180.4 Maple [A] (verified)

Time = 7.66 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.60

method	result
default	$\frac{2a \sin(dx+c)(104A(\cos^3(dx+c))+126B(\cos^3(dx+c))+52A(\cos^2(dx+c))+63B(\cos^2(dx+c))+39A \cos(dx+c)+21B \cos(dx+c)+15A)}{105d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}}$
parts	$\frac{2A \sin(dx+c)(104(\cos^3(dx+c))+52(\cos^2(dx+c))+39 \cos(dx+c)+15) \sqrt{a(1+\cos(dx+c))} a}{105d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}} + \frac{2B \sin(dx+c)(6(\cos^2(dx+c))+3 \cos(dx+c)+15) \sqrt{a(1+\cos(dx+c))} a}{5d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RET
URNVERBOSE)
```

```
output 2/105*a/d*sin(d*x+c)*(104*A*cos(d*x+c)^3+126*B*cos(d*x+c)^3+52*A*cos(d*x+c)
)^2+63*B*cos(d*x+c)^2+39*A*cos(d*x+c)+21*B*cos(d*x+c)+15*A)*(a*(1+cos(d*x+
c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(7/2)
```

3.180.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.59

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \frac{2 (2 (52 A + 63 B) a \cos(dx + c)^3 + (52 A + 63 B) a \cos(dx + c)) \sqrt{a(1 + \cos(dx + c))}}{105 d (1 + \cos(dx + c)) \cos(dx + c)^{\frac{7}{2}}} + \frac{2 B \sin(dx + c) (6 (\cos^2(dx + c)) + 3 \cos(dx + c) + 15) \sqrt{a(1 + \cos(dx + c))} a}{5 d (1 + \cos(dx + c)) \cos(dx + c)^{\frac{7}{2}}}$$

```
input integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="fracas")
```

3.180.
$$\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

output $2/105*(2*(52*A + 63*B)*a*\cos(d*x + c)^3 + (52*A + 63*B)*a*\cos(d*x + c)^2 + 3*(13*A + 7*B)*a*\cos(d*x + c) + 15*A*a)*\text{sqrt}(a*\cos(d*x + c) + a)*\text{sqrt}(\cos(d*x + c))*\sin(d*x + c)/(d*\cos(d*x + c)^5 + d*\cos(d*x + c)^4)$

3.180.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output Timed out

3.180.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 481 vs. $2(157) = 314$.

Time = 0.34 (sec) , antiderivative size = 481, normalized size of antiderivative = 2.66

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \frac{4 \left(21 \left(\frac{5 \sqrt{2} a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{10 \sqrt{2} a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{7 \sqrt{2} a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{2 \sqrt{2} a^{3/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{7/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{7/2} \left(\frac{2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} \right)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo rithm="maxima")`

output $4/105*(21*(5*\sqrt{2})*a^{(3/2)}*\sin(d*x + c)/(\cos(d*x + c) + 1) - 10*\sqrt{2})*$
 $a^{(3/2)}*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 7*\sqrt{2})*a^{(3/2)}*\sin(d*x +$
 $c)^5/(\cos(d*x + c) + 1)^5 - 2*\sqrt{2})*a^{(3/2)}*\sin(d*x + c)^7/(\cos(d*x + c)$
 $+ 1)^7)*B*(\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 1)^2/((\sin(d*x + c)/(\cos$
 $(d*x + c) + 1) + 1)^{(7/2)}*(-\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)^{(7/2)}*(2*$
 $\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + \sin(d*x + c)^4/(\cos(d*x + c) + 1)^4$
 $+ 1)) + (105*\sqrt{2})*a^{(3/2)}*\sin(d*x + c)/(\cos(d*x + c) + 1) - 245*\sqrt{2})*$
 $*a^{(3/2)}*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 273*\sqrt{2})*a^{(3/2)}*\sin(d*x$
 $+ c)^5/(\cos(d*x + c) + 1)^5 - 171*\sqrt{2})*a^{(3/2)}*\sin(d*x + c)^7/(\cos(d*x$
 $+ c) + 1)^7 + 38*\sqrt{2})*a^{(3/2)}*\sin(d*x + c)^9/(\cos(d*x + c) + 1)^9)*A*($
 $\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 1)^3/((\sin(d*x + c)/(\cos(d*x + c) +$
 $1) + 1)^{(9/2)}*(-\sin(d*x + c)/(\cos(d*x + c) + 1) + 1)^{(9/2)}*(3*\sin(d*x + c)$
 $^2/(\cos(d*x + c) + 1)^2 + 3*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 + \sin(d*x$
 $+ c)^6/(\cos(d*x + c) + 1)^6 + 1))))/d$

3.180.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 260.78 (sec) , antiderivative size = 147871, normalized size of antiderivative = 816.97

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="giac")`

output

```

67108864/105*sqrt(2)*sqrt(-tan(1/4*d*x + c)^4*tan(1/2*c)^8 + 14*tan(1/4*d*
x + c)^4*tan(1/2*c)^6 - 24*tan(1/4*d*x + c)^3*tan(1/2*c)^7 + 6*tan(1/4*d*x
+ c)^2*tan(1/2*c)^8 + 56*tan(1/4*d*x + c)^3*tan(1/2*c)^5 - 84*tan(1/4*d*x
+ c)^2*tan(1/2*c)^6 + 24*tan(1/4*d*x + c)*tan(1/2*c)^7 - tan(1/2*c)^8 - 1
4*tan(1/4*d*x + c)^4*tan(1/2*c)^2 + 56*tan(1/4*d*x + c)^3*tan(1/2*c)^3 - 5
6*tan(1/4*d*x + c)*tan(1/2*c)^5 + 14*tan(1/2*c)^6 + tan(1/4*d*x + c)^4 - 2
4*tan(1/4*d*x + c)^3*tan(1/2*c) + 84*tan(1/4*d*x + c)^2*tan(1/2*c)^2 - 56*
tan(1/4*d*x + c)*tan(1/2*c)^3 - 6*tan(1/4*d*x + c)^2 + 24*tan(1/4*d*x + c)
*tan(1/2*c) - 14*tan(1/2*c)^2 + 1)*(((((((((((((((((-52*I*A*a*e^(1055/2*I*c)
- 63*I*B*a*e^(1055/2*I*c) - 21008*I*A*a*e^(1053/2*I*c) - 25452*I*B*a*e^(1
053/2*I*c) - 4233112*I*A*a*e^(1051/2*I*c) - 5128578*I*B*a*e^(1051/2*I*c) -
567237008*I*A*a*e^(1049/2*I*c) - 687229452*I*B*a*e^(1049/2*I*c) - 5686551
0052*I*A*a*e^(1047/2*I*c) - 68894752563*I*B*a*e^(1047/2*I*c) - 45492408041
60*I*A*a*e^(1045/2*I*c) - 5511580205040*I*B*a*e^(1045/2*I*c) - 30252451347
6978*I*A*a*e^(1043/2*I*c) - 366520083635517*I*B*a*e^(1043/2*I*c) - 1720067
9480665512*I*A*a*e^(1041/2*I*c) - 20839284755400468*I*B*a*e^(1041/2*I*c) -
853583719248764868*I*A*a*e^(1039/2*I*c) - 1034149506008652852*I*B*a*e^(10
39/2*I*c) - 37557683649422024642*I*A*a*e^(1037/2*I*c) - 455025782669963001
63*I*B*a*e^(1037/2*I*c) - 1483528504376157661378*I*A*a*e^(1035/2*I*c) - 17
97351841782932557017*I*B*a*e^(1035/2*I*c) - 53137293718531292117900*I*A...

```

3.180.9 Mupad [B] (verification not implemented)

Time = 7.07 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.30

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \frac{\sqrt{a + a \cos(c + dx)} \left(-\frac{8 a e^{\frac{c}{2} + \frac{dx}{2}}}{6 \sqrt{\cos(c + dx)} e^{\frac{c}{2} + \frac{dx}{2}} \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + 6 \sqrt{\cos(c + dx)} \right)}{6 \sqrt{\cos(c + dx)} e^{\frac{c}{2} + \frac{dx}{2}} \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + 6 \sqrt{\cos(c + dx)}}$$

input

```

int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(9/2),x
)

```

output $((a + a*\cos(c + d*x))^{1/2}*((16*a*\exp((c*7i)/2 + (d*x*7i)/2)*\sin((3*c)/2 + (3*d*x)/2)*(13*A + 12*B))/(15*d) - (8*a*\exp((c*7i)/2 + (d*x*7i)/2)*\sin(c/2 + (d*x)/2)*(2*A + 3*B))/(3*d) + (8*a*\exp((c*7i)/2 + (d*x*7i)/2)*\sin((7*c)/2 + (7*d*x)/2)*(52*A + 63*B))/(105*d)))/(6*\cos(c + d*x)^{1/2}*\exp((c*7i)/2 + (d*x*7i)/2)*\cos(c/2 + (d*x)/2) + 6*\cos(c + d*x)^{1/2}*\exp((c*7i)/2 + (d*x*7i)/2)*\cos((3*c)/2 + (3*d*x)/2) + 2*\cos(c + d*x)^{1/2}*\exp((c*7i)/2 + (d*x*7i)/2)*\cos((5*c)/2 + (5*d*x)/2) + 2*\cos(c + d*x)^{1/2}*\exp((c*7i)/2 + (d*x*7i)/2)*\cos((7*c)/2 + (7*d*x)/2))$

3.180. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

3.181
$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

3.181.1 Optimal result 1724
 3.181.2 Mathematica [A] (verified) 1725
 3.181.3 Rubi [A] (verified) 1725
 3.181.4 Maple [A] (verified) 1729
 3.181.5 Fricas [A] (verification not implemented) 1729
 3.181.6 Sympy [F(-1)] 1730
 3.181.7 Maxima [B] (verification not implemented) 1730
 3.181.8 Giac [F(-1)] 1731
 3.181.9 Mupad [B] (verification not implemented) 1731

3.181.1 Optimal result

Integrand size = 35, antiderivative size = 228

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{2a^2(10A+9B) \sin(c+dx)}{63d \cos^{\frac{7}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a^2(34A+39B) \sin(c+dx)}{105d \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{8a^2(34A+39B) \sin(c+dx)}{315d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{16a^2(34A+39B) \sin(c+dx)}{315d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} + \frac{2aA \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)}$$

```
output 2/63*a^2*(10*A+9*B)*sin(d*x+c)/d/cos(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2)+
/105*a^2*(34*A+39*B)*sin(d*x+c)/d/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+
8/315*a^2*(34*A+39*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)
+16/315*a^2*(34*A+39*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/
2)+2/9*a*A*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(9/2)
```

3.181.2 Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.54

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \frac{a \sqrt{a(1 + \cos(c + dx))} (376A + 351B + (374A + 324B) \cos(c + dx) + 11(34A + 39B) \cos(2(c + dx)) + 68A \cos(3(c + dx)) + 78B \cos(3(c + dx)) + 68A \cos(4(c + dx)) + 78B \cos(4(c + dx))) \tan[(c + dx)/2]}{315d \cos^{9/2}(c + dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2),x]`

output `(a*sqrt[a*(1 + Cos[c + d*x])]*(376*A + 351*B + (374*A + 324*B)*Cos[c + d*x] + 11*(34*A + 39*B)*Cos[2*(c + d*x)] + 68*A*cos[3*(c + d*x)] + 78*B*cos[3*(c + d*x)] + 68*A*cos[4*(c + d*x)] + 78*B*cos[4*(c + d*x)])*Tan[(c + d*x)/2])/(315*d*cos[c + d*x]^(9/2))`

3.181.3 Rubi [A] (verified)Time = 1.10 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx \\ & \quad \downarrow \text{3454} \\ & \frac{2}{9} \int \frac{\sqrt{\cos(c + dx)a + a(a(10A + 9B) + 3a(2A + 3B) \cos(c + dx))}}{2 \cos^{9/2}(c + dx)} dx + \\ & \quad \frac{2aA \sin(c + dx) \sqrt{a \cos(c + dx) + a}}{9d \cos^{9/2}(c + dx)} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.181. $\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

$$\begin{aligned}
& \frac{1}{9} \int \frac{\sqrt{\cos(c+dx)a+a}(a(10A+9B)+3a(2A+3B)\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx + \\
& \quad \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a(10A+9B)+3a(2A+3B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{9/2}} dx + \\
& \quad \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \\
& \quad \downarrow \text{3459} \\
& \frac{1}{9} \left(\frac{3}{7} a(34A+39B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{7}{2}}(c+dx)} dx + \frac{2a^2(10A+9B)\sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \left(\frac{3}{7} a(34A+39B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}} dx + \frac{2a^2(10A+9B)\sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \\
& \quad \downarrow \text{3251} \\
& \frac{1}{9} \left(\frac{3}{7} a(34A+39B) \left(\frac{4}{5} \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{2a^2(10A+9B)\sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \left(\frac{3}{7} a(34A+39B) \left(\frac{4}{5} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{2a^2(10A+9B)\sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \\
& \quad \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \\
& \quad \downarrow \text{3251}
\end{aligned}$$

3.181. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{3}{7} a(34A + 39B) \left(\frac{4}{5} \left(\frac{2}{3} \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{3}{7} a(34A + 39B) \left(\frac{4}{5} \left(\frac{2}{3} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}} dx + \frac{2a \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) + \frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \right)$$

↓ 3250

$$\frac{1}{9} \left(\frac{2a^2(10A + 9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} + \frac{3}{7} a(34A + 39B) \left(\frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} + \frac{4}{5} \left(\frac{2a \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{a \cos(c+dx)+a}}{9d \cos^{\frac{9}{2}}(c+dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2), x]`

output `(2*a*A*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((2*a^2*(10*A + 9*B)*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) + (3*a*(34*A + 39*B)*((2*a*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x])))/5)/7)/9`

3.181.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.181. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.181.4 Maple [A] (verified)

Time = 7.78 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.57

method	result
default	$\frac{2a \sin(dx+c)(272A(\cos^4(dx+c))+312B(\cos^4(dx+c))+136A(\cos^3(dx+c))+156B(\cos^3(dx+c))+102A(\cos^2(dx+c))+117B(\cos^2(dx+c)))}{315d(1+\cos(dx+c)) \cos(dx+c)^{\frac{9}{2}}}$
parts	$\frac{2A \sin(dx+c)(272(\cos^4(dx+c))+136(\cos^3(dx+c))+102(\cos^2(dx+c))+85 \cos(dx+c)+35) \sqrt{a(1+\cos(dx+c))} a}{315d(1+\cos(dx+c)) \cos(dx+c)^{\frac{9}{2}}} + \frac{2B \sin(dx+c)(102(\cos^3(dx+c))+117(\cos^2(dx+c))+85 \cos(dx+c)+35) \sqrt{a(1+\cos(dx+c))} a}{315d(1+\cos(dx+c)) \cos(dx+c)^{\frac{9}{2}}}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x,method=_RE
TURNVERBOSE)
```

```
output 2/315*a/d*sin(d*x+c)*(272*A*cos(d*x+c)^4+312*B*cos(d*x+c)^4+136*A*cos(d*x+
c)^3+156*B*cos(d*x+c)^3+102*A*cos(d*x+c)^2+117*B*cos(d*x+c)^2+85*A*cos(d*x
+c)+45*B*cos(d*x+c)+35*A)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+
c)^(9/2)
```

3.181.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.55

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx = \frac{2(8(34A + 39B)a \cos(dx + c)^4 + 4(34A + 39B)a \cos(dx + c)^3 + 3(34A + 39B)a \cos(dx + c)^2 + 5(17A + 9B)a \cos(dx + c) + 35Aa) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c)}{d \cos(dx + c)^6 + d \cos(dx + c)^5}$$

```
input integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, alg
orithm="fricas")
```

```
output 2/315*(8*(34*A + 39*B)*a*cos(d*x + c)^4 + 4*(34*A + 39*B)*a*cos(d*x + c)^3
+ 3*(34*A + 39*B)*a*cos(d*x + c)^2 + 5*(17*A + 9*B)*a*cos(d*x + c) + 35*A
*a)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(d*cos(d*x +
c)^6 + d*cos(d*x + c)^5)
```


3.181.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(11/2),x)
```

```
output Timed out
```

3.181.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 573 vs. 2(198) = 396.

Time = 0.35 (sec) , antiderivative size = 573, normalized size of antiderivative = 2.51

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \frac{4 \left(3 \left(\frac{105 \sqrt{2} a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{245 \sqrt{2} a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{273 \sqrt{2} a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{171 \sqrt{2} a^{3/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \right)}{\dots}$$

```
input integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="maxima")
```

output
$$\frac{4/315*(3*(105*\sqrt{2})*a^{(3/2)}*\sin(dx + c)/(\cos(dx + c) + 1) - 245*\sqrt{2})*a^{(3/2)}*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 273*\sqrt{2}*a^{(3/2)}*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 171*\sqrt{2}*a^{(3/2)}*\sin(dx + c)^7/(\cos(dx + c) + 1)^7 + 38*\sqrt{2}*a^{(3/2)}*\sin(dx + c)^9/(\cos(dx + c) + 1)^9*B*(\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 1)^3/((\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(9/2)}*(-\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(9/2)}*(3*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 3*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + \sin(dx + c)^6/(\cos(dx + c) + 1)^6 + 1)) + (315*\sqrt{2})*a^{(3/2)}*\sin(dx + c)/(\cos(dx + c) + 1) - 840*\sqrt{2})*a^{(3/2)}*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 1344*\sqrt{2})*a^{(3/2)}*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 1242*\sqrt{2})*a^{(3/2)}*\sin(dx + c)^7/(\cos(dx + c) + 1)^7 + 517*\sqrt{2})*a^{(3/2)}*\sin(dx + c)^9/(\cos(dx + c) + 1)^9 - 94*\sqrt{2})*a^{(3/2)}*\sin(dx + c)^{11}/(\cos(dx + c) + 1)^{11}*A*(\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 1)^4/((\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(11/2)}*(-\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(11/2)}*(4*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 6*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + 4*\sin(dx + c)^6/(\cos(dx + c) + 1)^6 + \sin(dx + c)^8/(\cos(dx + c) + 1)^8 + 1)))}{d}$$

3.181.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(dx+c))^(3/2)*(A+B*cos(dx+c))/cos(dx+c)^(11/2),x, algorithm="giac")`

output Timed out

3.181.9 Mupad [B] (verification not implemented)

Time = 7.45 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.27

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \frac{\sqrt{a + a \cos(c + dx)} \left(- \right)}{12 \sqrt{\cos(c + dx)} e^{\frac{c}{2} + \frac{dx}{2}} \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + 8 \sqrt{\cos(c + a}}$$

3.181.
$$\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/cos(c + d*x)^(11/2),
x)`

output `((a + a*cos(c + d*x))^(1/2)*((16*a*exp((c*9i)/2 + (d*x*9i)/2)*sin((5*c)/2 + (5*d*x)/2)*(34*A + 39*B))/(35*d) - (16*B*a*exp((c*9i)/2 + (d*x*9i)/2)*sin((3*c)/2 + (3*d*x)/2))/(3*d) + (32*a*exp((c*9i)/2 + (d*x*9i)/2)*sin((9*c)/2 + (9*d*x)/2)*(34*A + 39*B))/(315*d) + (96*a*exp((c*9i)/2 + (d*x*9i)/2)*sin(c/2 + (d*x)/2)*(A + B))/(5*d))/(12*cos(c + d*x)^(1/2)*exp((c*9i)/2 + (d*x*9i)/2)*cos(c/2 + (d*x)/2) + 8*cos(c + d*x)^(1/2)*exp((c*9i)/2 + (d*x*9i)/2)*cos((3*c)/2 + (3*d*x)/2) + 8*cos(c + d*x)^(1/2)*exp((c*9i)/2 + (d*x*9i)/2)*cos((5*c)/2 + (5*d*x)/2) + 2*cos(c + d*x)^(1/2)*exp((c*9i)/2 + (d*x*9i)/2)*cos((7*c)/2 + (7*d*x)/2) + 2*cos(c + d*x)^(1/2)*exp((c*9i)/2 + (d*x*9i)/2)*cos((9*c)/2 + (9*d*x)/2))`

3.181. $\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

3.182 $\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.182.1 Optimal result 1733
 3.182.2 Mathematica [A] (verified) 1734
 3.182.3 Rubi [A] (verified) 1734
 3.182.4 Maple [A] (verified) 1739
 3.182.5 Fricas [A] (verification not implemented) 1739
 3.182.6 Sympy [F(-1)] 1740
 3.182.7 Maxima [B] (verification not implemented) 1740
 3.182.8 Giac [F] 1741
 3.182.9 Mupad [F(-1)] 1742

3.182.1 Optimal result

Integrand size = 35, antiderivative size = 274

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx = \frac{a^{5/2}(326A+283B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{128d} + \frac{a^3(326A+283B) \sqrt{\cos(c+dx)} \sin(c+dx)}{128d \sqrt{a+a \cos(c+dx)}} + \frac{a^3(326A+283B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{192d \sqrt{a+a \cos(c+dx)}} + \frac{a^3(170A+157B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{240d \sqrt{a+a \cos(c+dx)}} + \frac{a^2(10A+13B) \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{40d} + \frac{aB \cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{5d}$$

output $\frac{1}{128}a^{5/2}(326A+283B)\arcsin(\sin(dx+c)a^{1/2}/(a+a\cos(dx+c))^{1/2})/d+1/5a^3B\cos(dx+c)^{5/2}(a+a\cos(dx+c))^{3/2}\sin(dx+c)/d+1/192a^3(326A+283B)\cos(dx+c)^{3/2}\sin(dx+c)/d/(a+a\cos(dx+c))^{1/2}+1/240a^3(170A+157B)\cos(dx+c)^{5/2}\sin(dx+c)/d/(a+a\cos(dx+c))^{1/2}+1/128a^3(326A+283B)\sin(dx+c)\cos(dx+c)^{1/2}/d/(a+a\cos(dx+c))^{1/2}+1/40a^2(10A+13B)\cos(dx+c)^{5/2}\sin(dx+c)(a+a\cos(dx+c))^{1/2}/d$

3.182.2 Mathematica [A] (verified)

Time = 1.27 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.58

$$\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \frac{a^2\sqrt{a(1+\cos(c+dx))}\sec\left(\frac{1}{2}(c+dx)\right)\left(15\sqrt{2}(326A+283B)\arcsin\left(\sqrt{2}\sin\left(\frac{1}{2}(c+dx)\right)\right)+15\sqrt{2}(326A+283B)\right)}{(3840d)}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output $(a^2\sqrt{a(1+\cos(c+dx))}\sec((c+dx)/2)*(15\sqrt{2}(326A+283B)\arcsin(\sqrt{2}\sin((c+dx)/2))+15\sqrt{2}(326A+283B))+2\sqrt{\cos(c+dx)}*(5810A+5521B+(3620A+3874B)\cos(c+dx)+4*(230A+331B)\cos(2*(c+dx))+120A\cos(3*(c+dx))+348B\cos(3*(c+dx))+48B\cos(4*(c+dx))}\sin((c+dx)/2))/(3840d)$

3.182.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}(A+B\cos(c+dx))dx$$

↓ 3042

3.182. $\int \cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx$

$$\begin{aligned}
& \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3455} \\
& \frac{1}{5} \int \frac{1}{2} \cos^{3/2}(c + dx) (\cos(c + dx)a + a)^{3/2} (5a(2A + B) + a(10A + 13B) \cos(c + dx)) dx + \\
& \quad \frac{aB \sin(c + dx) \cos^{5/2}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{10} \int \cos^{3/2}(c + dx) (\cos(c + dx)a + a)^{3/2} (5a(2A + B) + a(10A + 13B) \cos(c + dx)) dx + \\
& \quad \frac{aB \sin(c + dx) \cos^{5/2}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{10} \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(\sin\left(c + dx + \frac{\pi}{2}\right) a + a\right)^{3/2} \left(5a(2A + B) + a(10A + 13B) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\
& \quad \frac{aB \sin(c + dx) \cos^{5/2}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\
& \quad \downarrow \text{3455} \\
& \frac{1}{10} \left(\frac{1}{4} \int \frac{1}{2} \cos^{3/2}(c + dx) \sqrt{\cos(c + dx)a + a} (5(26A + 21B)a^2 + (170A + 157B) \cos(c + dx)a^2) dx + \frac{a^2(10A + 13B)}{5d} \right) \\
& \quad \frac{aB \sin(c + dx) \cos^{5/2}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{10} \left(\frac{1}{8} \int \cos^{3/2}(c + dx) \sqrt{\cos(c + dx)a + a} (5(26A + 21B)a^2 + (170A + 157B) \cos(c + dx)a^2) dx + \frac{a^2(10A + 13B)}{5d} \right) \\
& \quad \frac{aB \sin(c + dx) \cos^{5/2}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{10} \left(\frac{1}{8} \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a} (5(26A + 21B)a^2 + (170A + 157B) \sin\left(c + dx + \frac{\pi}{2}\right) a^2) dx + \frac{a^2(10A + 13B)}{5d} \right) \\
& \quad \frac{aB \sin(c + dx) \cos^{5/2}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d} \\
& \quad \downarrow \text{3460}
\end{aligned}$$

3.182. $\int \cos^{3/2}(c + dx) (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \int \cos^{\frac{3}{2}}(c + dx) \sqrt{\cos(c + dx)a + adx} + \frac{a^3(170A + 157B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + adx} + \frac{a^3(170A + 157B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) \\ \downarrow \text{3249}$$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \left(\frac{3}{4} \int \sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + adx} + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \right) + \frac{a^3(170A + 157B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \left(\frac{3}{4} \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + adx} + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \right) + \frac{a^3(170A + 157B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d\sqrt{a \cos(c + dx) + a}} \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) \\ \downarrow \text{3249}$$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} \right) \right) + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d\sqrt{a \cos(c + dx) + a}} \right) \right) + \frac{a \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d} \right) \\ \downarrow \text{3253}$$

$$\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6} a^2 (326A + 283B) \left(\frac{3}{4} \left(\frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx) a + a}}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right) \right) + \frac{a \sin(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) \right) \right)$$

$$\frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d}$$

↓ 223

$$\frac{1}{10} \left(\frac{a^2 (10A + 13B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}{4d} + \frac{1}{8} \left(\frac{a^3 (170A + 157B) \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{3d \sqrt{a \cos(c + dx) + a}} \right) \right)$$

$$\frac{aB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{5d}$$

input `Int[Cos[c + d*x]^(3/2)*(a + a*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]),x]`

output `(a*B*cos[c + d*x]^(5/2)*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(5*d) + (a^2*(10*A + 13*B)*cos[c + d*x]^(5/2)*sqrt[a + a*cos[c + d*x]]*sin[c + d*x])/(4*d) + ((a^3*(170*A + 157*B)*cos[c + d*x]^(5/2)*sin[c + d*x])/(3*d*sqrt[a + a*cos[c + d*x]]) + (5*a^2*(326*A + 283*B)*((a*cos[c + d*x]^(3/2)*sin[c + d*x])/(2*d*sqrt[a + a*cos[c + d*x]]) + (3*((sqrt[a]*arcsin[(sqrt[a]*sin[c + d*x])/sqrt[a + a*cos[c + d*x]])]/d + (a*sqrt[cos[c + d*x]]*sin[c + d*x])/(d*sqrt[a + a*cos[c + d*x]]))))/4)/6)/8)/10`

3.182.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.182. $\int \cos^{\frac{3}{2}}(c + dx) (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$

rule 3249 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.182.4 Maple [A] (verified)

Time = 16.23 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.54

method	result
default	$a^2 \left(384B \sin(dx+c) (\cos^4(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 480A \sin(dx+c) (\cos^3(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 1392B (\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \dots \right)$
parts	$\frac{A \left(48 \sin(dx+c) (\cos^3(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 184 \sin(dx+c) (\cos^2(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 326 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \dots \right)}{192d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

input `int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{1920} a^2 / d \left(384 B \sin(dx+c) \cos(dx+c)^4 \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 480 A \sin(dx+c) \cos(dx+c)^3 \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 1392 B \cos(dx+c)^3 \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 1840 A \cos(dx+c)^2 \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 2264 B \cos(dx+c)^2 \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 3260 A \cos(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 2830 B \cos(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 4890 A \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \sin(dx+c) + 4245 B \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 4890 A \arctan\left(\frac{\tan(dx+c)}{\cos(dx+c)}\right) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 4245 B \arctan\left(\frac{\tan(dx+c)}{\cos(dx+c)}\right) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \right) \cos(dx+c)^{1/2} (a(1+\cos(dx+c)))^{1/2} / (1+\cos(dx+c))^{1/2}$$

3.182.5 Fracas [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.71

$$\int \cos^{\frac{3}{2}}(c+dx) (a+a\cos(c+dx))^{5/2} (A + B \cos(c+dx)) dx = \frac{(384 B a^2 \cos(dx+c)^4 + 48 (10 A + 29 B) a^2 \cos(dx+c)^3 + 8 (230 A + 283 B) a^2 \cos(dx+c)^2 + \dots}{192 d (1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x,algorithm="fracas")`

$$3.182. \quad \int \cos^{\frac{3}{2}}(c+dx) (a+a\cos(c+dx))^{5/2} (A + B \cos(c+dx)) dx$$

output `1/1920*((384*B*a^2*cos(d*x + c)^4 + 48*(10*A + 29*B)*a^2*cos(d*x + c)^3 + 8*(230*A + 283*B)*a^2*cos(d*x + c)^2 + 10*(326*A + 283*B)*a^2*cos(d*x + c) + 15*(326*A + 283*B)*a^2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - 15*((326*A + 283*B)*a^2*cos(d*x + c) + (326*A + 283*B)*a^2)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/(d*cos(d*x + c) + d)`

3.182.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.182.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10042 vs. $2(236) = 472$.

Time = 0.97 (sec) , antiderivative size = 10042, normalized size of antiderivative = 36.65

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algo rithm="maxima")`

output `1/7680*((10*(cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + 2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)^(3/4))*((135*a^2*sin(4/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 88*a^2*sin(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 135*a^2*sin(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))))*cos(3/2*arctan2(sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))), cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)) - (135*a^2*cos(4/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 88*a^2*cos(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) - 135*a^2*cos(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) - 88*a^2)*sin(3/2*arctan2(sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))), cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)))*sqrt(a) + 6*(cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + 2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)^(1/4)*(8*(a^2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2*sin(5*d*x + 5*c) + a^2*sin(5*d*x + 5*c)*sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + 2*a^2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))*sin(5*d*x + 5*c) + a^2*sin(5*d*x + 5*c))*cos(5/2*arctan2(sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))), cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)) - 5*(35*a^2*sin(4/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x...`

3.182.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(3/2), x)`

3.182.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^{3/2}(A + B \cos(c + dx))(a + a \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2), x)`output `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2), x)`

3.183 $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

3.183.1 Optimal result	1743
3.183.2 Mathematica [A] (verified)	1744
3.183.3 Rubi [A] (verified)	1744
3.183.4 Maple [A] (verified)	1748
3.183.5 Fricas [A] (verification not implemented)	1749
3.183.6 Sympy [F(-1)]	1749
3.183.7 Maxima [B] (verification not implemented)	1750
3.183.8 Giac [F]	1750
3.183.9 Mupad [F(-1)]	1751

3.183.1 Optimal result

Integrand size = 35, antiderivative size = 227

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{a^{5/2}(200A + 163B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{64d} + \frac{a^3(200A + 163B) \sqrt{\cos(c + dx)} \sin(c + dx)}{64d \sqrt{a + a \cos(c + dx)}} + \frac{a^3(104A + 95B) \cos^{3/2}(c + dx) \sin(c + dx)}{96d \sqrt{a + a \cos(c + dx)}} + \frac{a^2(8A + 11B) \cos^{3/2}(c + dx) \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{24d} + \frac{aB \cos^{3/2}(c + dx)(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{4d}$$

output

```
1/64*a^(5/2)*(200*A+163*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2)))/d+1/4*a*B*cos(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d+1/96*a^3*(104*A+95*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+1/64*a^3*(200*A+163*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)+1/24*a^2*(8*A+11*B)*cos(d*x+c)^(3/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d
```

3.183.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.60

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \frac{a^2\sqrt{a(1+\cos(c+dx))}\sec\left(\frac{1}{2}(c+dx)\right)\left(3\sqrt{2}(200A+163B)\arcsin\left(\sqrt{2}\sin\left(\frac{1}{2}(c+dx)\right)\right)+2\sqrt{2}\sin\left(\frac{1}{2}(c+dx)\right)\right)}{384d}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(3*Sqrt[2]*(200*A + 163*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[2]*Sin[(c + d*x)/2]) + 2*Sqrt[Cos[c + d*x]]*(632*A + 581*B + (272*A + 362*B)*Cos[c + d*x] + 4*(8*A + 23*B)*Cos[2*(c + d*x)] + 12*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2])/384*d`

3.183.3 Rubi [A] (verified)Time = 1.19 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}(A+B\cos(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^{5/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\ & \quad \downarrow \text{3455} \\ & \frac{1}{4}\int\frac{1}{2}\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}(a(8A+3B)+a(8A+11B)\cos(c+dx))dx + \\ & \quad \frac{aB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}{4d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.183. $\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx$

$$\frac{1}{8} \int \sqrt{\cos(c+dx)} (\cos(c+dx)a+a)^{3/2} (a(8A+3B)+a(8A+11B)\cos(c+dx)) dx + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx)+a)^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{3/2} \left(a(8A+3B)+a(8A+11B)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx)+a)^{3/2}}{4d}$$

↓ 3455

$$\frac{1}{8} \left(\frac{1}{3} \int \frac{1}{2} \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a} (3(24A+17B)a^2 + (104A+95B)\cos(c+dx)a^2) dx + \frac{a^2(8A+11B)\sin(c+dx)}{4d} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx)+a)^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a} (3(24A+17B)a^2 + (104A+95B)\cos(c+dx)a^2) dx + \frac{a^2(8A+11B)\sin(c+dx)}{4d} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx)+a)^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} (3(24A+17B)a^2 + (104A+95B)\sin\left(c+dx+\frac{\pi}{2}\right)a^2) dx + \frac{a^2(8A+11B)\cos(c+dx)}{4d} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx)+a)^{3/2}}{4d}$$

↓ 3460

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (200A+163B) \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+adx} + \frac{a^3(104A+95B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) + \frac{aB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a \cos(c+dx)+a)^{3/2}}{4d} \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (200A + 163B) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} a + dx + \frac{a^3 (104A + 95B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right) \\ \downarrow \text{3249}$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (200A + 163B) \left(\frac{1}{2} \int \frac{\sqrt{\cos(c + dx) a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{a^3 (104A + 95B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (200A + 163B) \left(\frac{1}{2} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} a + a}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{a^3 (104A + 95B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right) \\ \downarrow \text{3253}$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3}{4} a^2 (200A + 163B) \left(\frac{a \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx) a + a}}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right)}{d} \right) + \frac{a^3 (104A + 95B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right) \right. \\ \left. \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right) \\ \downarrow \text{223}$$

$$\frac{1}{8} \left(\frac{a^2 (8A + 11B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}}{3d} + \frac{1}{6} \left(\frac{a^3 (104A + 95B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} + \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2}}{4d} \right) \right)$$

input `Int[Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]),x]`

```
output (a*B*cos[c + d*x]^(3/2)*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(4*d) + (
(a^2*(8*A + 11*B)*cos[c + d*x]^(3/2)*sqrt[a + a*cos[c + d*x]]*sin[c + d*x]
)/(3*d) + ((a^3*(104*A + 95*B)*cos[c + d*x]^(3/2)*sin[c + d*x])/(2*d*sqrt[
a + a*cos[c + d*x]]) + (3*a^2*(200*A + 163*B)*((sqrt[a]*ArcSin[(sqrt[a]*Si
n[c + d*x])/sqrt[a + a*cos[c + d*x]]))/d + (a*sqrt[cos[c + d*x]]*sin[c + d
*x])/(d*sqrt[a + a*cos[c + d*x]])))/4)/6)/8
```

3.183.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 223 Int[1/sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3249 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (
f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*cos[e + f*x]*((c + d*sin[e + f*x])
^n/(f*(2*n + 1)*sqrt[a + b*sin[e + f*x]])), x] + Simp[2*n*((b*c + a*d)/(b*(
2*n + 1))) Int[Sqrt[a + b*sin[e + f*x]]*(c + d*sin[e + f*x])^(n - 1), x],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]
```

```
rule 3253 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/sqrt[(d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/sqrt[1 - x^2/a], x], x, b*(Co
s[e + f*x]/sqrt[a + b*sin[e + f*x]])], x] /; FreeQ[{a, b, d, e, f}, x] && E
qQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.183.4 Maple [A] (verified)

Time = 15.85 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.55

method	result
default	$\frac{a^2 \left(48B (\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 64A (\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 184B (\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{24d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \left(8 \sin(dx+c) (\cos^2(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 34 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 75 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 75 \arctan \left(\tan(dx+c) \right) \right)}{24d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c)),x,method=_RET
URNVERBOSE)
```

3.183. $\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

```
output 1/192*a^2/d*(48*B*cos(d*x+c)^3*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
)+64*A*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+184*B*cos
(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+272*A*cos(d*x+c)*si
n(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+326*B*cos(d*x+c)*sin(d*x+c)*(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2)+600*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin
(d*x+c)+489*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+600*A*arctan(ta
n(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+489*B*arctan(tan(d*x+c)*(cos(d
*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(1
+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

3.183.5 Fracas [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.77

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx = \frac{(48Ba^2\cos(dx+c)^3 + 8(8A+23B)a^2\cos(dx+c)^2 + 2(136A+163B)a^2\cos(dx+c) + B\cos(c+dx))}{d}$$

```
input integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algo
rithm="fricas")
```

```
output 1/192*((48*B*a^2*cos(d*x + c)^3 + 8*(8*A + 23*B)*a^2*cos(d*x + c)^2 + 2*(1
36*A + 163*B)*a^2*cos(d*x + c) + 3*(200*A + 163*B)*a^2)*sqrt(a*cos(d*x + c
) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - 3*((200*A + 163*B)*a^2*cos(d*x +
c) + (200*A + 163*B)*a^2)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos
(d*x + c))/(sqrt(a)*sin(d*x + c))))/(d*cos(d*x + c) + d)
```

3.183.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(1/2)*(a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.183. $\int \sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx$

3.183.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9415 vs. $2(195) = 390$.

Time = 0.93 (sec) , antiderivative size = 9415, normalized size of antiderivative = 41.48

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algo
rithm="maxima")`

output `1/768*(8*(4*(a^2*cos(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d
*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1))*si
n(3*d*x + 3*c) - (a^2*cos(3*d*x + 3*c) - a^2)*sin(3/2*arctan2(sin(2/3*arct
an2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c)
, cos(3*d*x + 3*c))) + 1))*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x +
3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos
(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(3/4)*sqrt(a) + 30*
(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(
sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c
, cos(3*d*x + 3*c))) + 1)^(1/4)*((a^2*sin(2/3*arctan2(sin(3*d*x + 3*c), c
os(3*d*x + 3*c))) + 5*a^2*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*
c))))*cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))
, cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)) - (a^2*cos(2/
3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 3*a^2*cos(1/3*arctan2(sin
(3*d*x + 3*c), cos(3*d*x + 3*c))) - 4*a^2)*sin(1/2*arctan2(sin(2/3*arctan2
(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), c
os(3*d*x + 3*c))) + 1))*sqrt(a) + 75*(a^2*arctan2(-(cos(2/3*arctan2(sin(3
*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(
3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))
+ 1)^(1/4)*cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x...`

3.183.8 Giac [F]

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^{\frac{5}{2}} \sqrt{\cos(dx + c)} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(5/2)*sqrt(cos(d*x + c
)), x)`

3.183.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \sqrt{\cos(c + dx)}(A + B \cos(c + dx))(a + a \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2),x)`

output `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2), x)`

3.184
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.184.1 Optimal result 1752
 3.184.2 Mathematica [A] (verified) 1753
 3.184.3 Rubi [A] (verified) 1753
 3.184.4 Maple [A] (verified) 1757
 3.184.5 Fricas [A] (verification not implemented) 1757
 3.184.6 Sympy [F(-1)] 1758
 3.184.7 Maxima [B] (verification not implemented) 1758
 3.184.8 Giac [F(-1)] 1759
 3.184.9 Mupad [F(-1)] 1760

3.184.1 Optimal result

Integrand size = 35, antiderivative size = 180

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx = \frac{a^{5/2}(38A+25B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{8d} + \frac{a^3(54A+49B)\sqrt{\cos(c+dx)} \sin(c+dx)}{24d\sqrt{a+a \cos(c+dx)}} + \frac{a^2(2A+3B)\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{4d} + \frac{aB\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{3d}$$

```
output 1/8*a^(5/2)*(38*A+25*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/
d+1/3*a*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)*cos(d*x+c)^(1/2)/d+1/24*a^3*(5
4*A+49*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)+1/4*a^2*(2*
A+3*B)*sin(d*x+c)*cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(1/2)/d
```

3.184.2 Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.67

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(3\sqrt{2}(38A + 25B) \right)}{48d}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(3*Sqrt[2]*(38*A + 25*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(66*A + 79*B + 2*(6*A + 17*B)*Cos[c + d*x] + 4*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.184.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{3455} \\ & \frac{1}{3} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(6A + B) + 3a(2A + 3B) \cos(c + dx))}{2\sqrt{\cos(c + dx)}} dx + \\ & \quad \frac{aB \sin(c + dx) \sqrt{\cos(c + dx)} (a \cos(c + dx) + a)^{3/2}}{3d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.184. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\frac{1}{6} \int \frac{(\cos(c+dx)a+a)^{3/2}(a(6A+B)+3a(2A+3B)\cos(c+dx))}{\sqrt{\cos(c+dx)} \frac{aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}}{3d}} dx +$$

↓ 3042

$$\frac{1}{6} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(6A+B)+3a(2A+3B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \frac{aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}}{3d}} dx +$$

↓ 3455

$$\frac{1}{6} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a}((30A+13B)a^2+(54A+49B)\cos(c+dx)a^2)}{2\sqrt{\cos(c+dx)} \frac{aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}}{3d}} dx + \frac{3a^2(2A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} \right)$$

↓ 27

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{\sqrt{\cos(c+dx)a+a}((30A+13B)a^2+(54A+49B)\cos(c+dx)a^2)}{\sqrt{\cos(c+dx)} \frac{aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}}{3d}} dx + \frac{3a^2(2A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((30A+13B)a^2+(54A+49B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \frac{aB \sin(c+dx)\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}}{3d}} dx + \frac{3a^2(2A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} \right)$$

↓ 3460

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3}{2} a^2(38A+25B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx + \frac{a^3(54A+49B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{3a^2(2A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3}{2} a^2 (38A + 25B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})} a + a}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{a^3 (54A + 49B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{3a^2 (2A + 3B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} \right) + \frac{3a^5/2 (38A + 25B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{4d}$$

↓ 3253

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{a^3 (54A + 49B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} - \frac{3a^2 (38A + 25B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx) a + a}}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}}\right)}{d} \right) + \frac{3a^2 (2A + 3B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} \right) + \frac{3a^5/2 (38A + 25B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{4d}$$

↓ 223

$$\frac{1}{6} \left(\frac{3a^2 (2A + 3B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} + \frac{1}{4} \left(\frac{3a^{5/2} (38A + 25B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{d} \right) \right) + \frac{3a^5/2 (38A + 25B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right)}{4d}$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(a*B*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + ((3*a^2*(2*A + 3*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + ((3*a^(5/2)*(38*A + 25*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a^3*(54*A + 49*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4)/6`

3.184.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.184.4 Maple [A] (verified)

Time = 20.25 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.55

method	result
default	$a^2 \left(8B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 34B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \dots \right)$
parts	$\frac{A \left(2(\cos^2(dx+c) \sin(dx+c) + 19 \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}) \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + 11 \cos(dx+c) \sin(dx+c) + 19 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{4d \sqrt{\cos(dx+c)} (1+\cos(dx+c))}$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{24}a^2/d * (8*B*\cos(d*x+c)^2*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2) + 12*A*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2) + 34*B*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2) + 66*A*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\sin(d*x+c) + 75*B*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2) + 114*A*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)) + 75*B*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)))*\cos(d*x+c)^(1/2)*(a*(1+\cos(d*x+c)))^(1/2)/(1+\cos(d*x+c))/(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)$$

3.184.5 Fracas [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.86

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{(8 B a^2 \cos(dx + c)^2 + 2(6 A + 17 B) a^2 \cos(dx + c) + 3(22 A + 25 B) a^2) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c) - 3((38 A + 25 B) a^2 \cos(dx + c) + (38 A + 25 B) a^2) \sqrt{a} \arctan(\sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)})}{d \cos(dx + c) + d}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,algorithm="fracas")`

output
$$\frac{1}{24} * ((8*B*a^2*\cos(d*x + c)^2 + 2*(6*A + 17*B)*a^2*\cos(d*x + c) + 3*(22*A + 25*B)*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c) - 3*((38*A + 25*B)*a^2*\cos(d*x + c) + (38*A + 25*B)*a^2)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(d*\cos(d*x + c) + d)$$

3.184.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Timed out`

3.184.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3071 vs. 2(154) = 308.

Time = 0.66 (sec) , antiderivative size = 3071, normalized size of antiderivative = 17.06

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

3.184.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(1/2),x)`

3.185
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.185.1 Optimal result 1761
 3.185.2 Mathematica [A] (verified) 1762
 3.185.3 Rubi [A] (verified) 1762
 3.185.4 Maple [A] (verified) 1766
 3.185.5 Fricas [A] (verification not implemented) 1766
 3.185.6 Sympy [F(-1)] 1767
 3.185.7 Maxima [B] (verification not implemented) 1767
 3.185.8 Giac [F(-1)] 1768
 3.185.9 Mupad [F(-1)] 1769

3.185.1 Optimal result

Integrand size = 35, antiderivative size = 178

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{a^{5/2}(20A+19B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4d}$$

$$- \frac{a^3(4A-9B)\sqrt{\cos(c+dx)} \sin(c+dx)}{4d\sqrt{a+a \cos(c+dx)}}$$

$$- \frac{a^2(4A-B)\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{2d}$$

$$+ \frac{2aA(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{d\sqrt{\cos(c+dx)}}$$

```
output 1/4*a^(5/2)*(20*A+19*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/
d+2*a*A*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(1/2)-1/4*a^3*(4*A-
9*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)-1/2*a^2*(4*A-B)*
sin(d*x+c)*cos(d*x+c)^(1/2)*(a+a*cos(d*x+c))^(1/2)/d
```


3.185.2 Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.71

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \left(\sqrt{2}(20A + 19B)\right.}{\cos^{3/2}(c + dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])] * Sec[(c + d*x)/2] * (Sqrt[2]*(20*A + 19*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] * Sqrt[Cos[c + d*x]] + 2*(8*A + B + (4*A + 11*B)*Cos[c + d*x] + B*Cos[2*(c + d*x)]) * Sin[(c + d*x)/2]) / (8*d*Sqrt[Cos[c + d*x]])`

3.185.3 Rubi [A] (verified)Time = 1.01 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3454, 27, 3042, 3455, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3454} \\ & 2 \int \frac{(\cos(c + dx)a + a)^{3/2} (a(4A + B) - a(4A - B) \cos(c + dx))}{2\sqrt{\cos(c + dx)}} dx + \\ & \quad \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{d\sqrt{\cos(c + dx)}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.185. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(\cos(c+dx)a+a)^{3/2}(a(4A+B)-a(4A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(4A+B)-a(4A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3455} \\
& \frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+a}(a^2(12A+5B)-a^2(4A-9B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx - \\
& \frac{a^2(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}{2d} + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{1}{4} \int \frac{\sqrt{\cos(c+dx)a+a}(a^2(12A+5B)-a^2(4A-9B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx - \\
& \frac{a^2(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}{2d} + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a^2(12A+5B)-a^2(4A-9B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \\
& \frac{a^2(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}{2d} + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3460} \\
& \frac{1}{4} \left(\frac{1}{2} a^2(20A+19B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx - \frac{a^3(4A-9B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) - \\
& \frac{a^2(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}{2d} + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{d\sqrt{\cos(c+dx)}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.185. $\int \frac{(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\frac{1}{4} \left(\frac{1}{2} a^2 (20A + 19B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2}) a + a}}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{a^3 (4A - 9B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) - \frac{a^2 (4A - B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{d \sqrt{\cos(c + dx)}}$$

↓ 3253

$$\frac{1}{4} \left(\frac{a^2 (20A + 19B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx) a + a}}} d \left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx) a + a}} \right)}{d} - \frac{a^3 (4A - 9B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) - \frac{a^2 (4A - B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{d \sqrt{\cos(c + dx)}}$$

↓ 223

$$- \frac{a^2 (4A - B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} + \frac{1}{4} \left(\frac{a^{5/2} (20A + 19B) \arcsin \left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}} \right)}{d} - \frac{a^3 (4A - 9B) \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{d \sqrt{\cos(c + dx)}}$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `-1/2*(a^2*(4*A - B)*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/d + (2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) + ((a^(5/2)*(20*A + 19*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (a^3*(4*A - 9*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4`

3.185.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.185.4 Maple [A] (verified)

Time = 18.33 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.48

method	result
default	$\frac{a^2 \left(2B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 11B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 20A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{4d(1+\cos(dx+c)) \sqrt{\cos(dx+c)}}$
parts	$\frac{A \left(5 \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + \cos(dx+c) \sin(dx+c) + 5 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{d(1+\cos(dx+c)) \sqrt{\cos(dx+c)}}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/4*a^2/d*(2*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4
*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+11*B*cos(d*x+c)
*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+20*A*cos(d*x+c)*arctan(tan(d
*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+8*A*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)*sin(d*x+c)+19*B*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)
)))^(1/2))*((a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c)))/(cos(d*x+c)/(1+cos(d*x
+c)))^(1/2)/cos(d*x+c)^(1/2)
```

3.185.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.92

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \frac{(2Ba^2 \cos(dx + c)^2 + (4A + 11B)a^2 \cos(dx + c) + 8A)}{\cos^{3/2}(c + dx)}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algo
rithm="fricas")
```

3.185. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

output $1/4*((2*B*a^2*\cos(d*x + c)^2 + (4*A + 11*B)*a^2*\cos(d*x + c) + 8*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c) - ((20*A + 19*B)*a^2*\cos(d*x + c)^2 + (20*A + 19*B)*a^2*\cos(d*x + c))*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c)))/(d*\cos(d*x + c)^2 + d*\cos(d*x + c))$

3.185.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output Timed out

3.185.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2080 vs. $2(154) = 308$.

Time = 0.54 (sec) , antiderivative size = 2080, normalized size of antiderivative = 11.69

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/16*((2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*((a^2*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(2*d*x + 2*c) + a^2*sin(2*d*x + 2*c) - (a^2*cos(2*d*x + 2*c) - 10*a^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + (a^2*sin(2*d*x + 2*c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) - a^2*cos(2*d*x + 2*c) + 10*a^2 + (a^2*cos(2*d*x + 2*c) - 10*a^2)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(a) + 19*(a^2*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - a^2*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arcta...`

3.185.8 Giac [**F(-1)**]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorith="giac")`

output `Timed out`

3.185.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(3/2),x)`

3.186
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$$

3.186.1 Optimal result	1770
3.186.2 Mathematica [A] (verified)	1770
3.186.3 Rubi [A] (verified)	1771
3.186.4 Maple [A] (verified)	1774
3.186.5 Fricas [A] (verification not implemented)	1775
3.186.6 Sympy [F(-1)]	1775
3.186.7 Maxima [B] (verification not implemented)	1776
3.186.8 Giac [F(-1)]	1776
3.186.9 Mupad [F(-1)]	1777

3.186.1 Optimal result

Integrand size = 35, antiderivative size = 173

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx = \frac{a^{5/2}(2A+5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d} - \frac{a^3(14A+3B)\sqrt{\cos(c+dx)} \sin(c+dx)}{3d\sqrt{a+a \cos(c+dx)}} + \frac{2a^2(2A+B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}} + \frac{2aA(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{3d \cos^{3/2}(c+dx)}$$

output `a^(5/2)*(2*A+5*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+2/3*a*A*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(3/2)-1/3*a^3*(14*A+3*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)+2*a^2*(2*A+B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)`

3.186.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.75

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx = \frac{a^2 \sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(3\sqrt{2}(2A+5B)\right)}{\cos^{5/2}(c+dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(3*sqrt[2]*(2*A + 5*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^(3/2) + (4*A + 3*B + 4*(8*A + 3*B)*Cos[c + d*x] + 3*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(6*d*Cos[c + d*x]^(3/2))`

3.186.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3454

$$\frac{2}{3} \int \frac{(\cos(c + dx)a + a)^{3/2} (3a(2A + B) - a(2A - 3B) \cos(c + dx))}{2 \cos^{3/2}(c + dx)} dx + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{3d \cos^{3/2}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{(\cos(c + dx)a + a)^{3/2} (3a(2A + B) - a(2A - 3B) \cos(c + dx))}{\cos^{3/2}(c + dx)} dx + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{3d \cos^{3/2}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (3a(2A+B) - a(2A-3B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3454

$$\frac{1}{3} \left(2 \int \frac{\sqrt{\cos(c+dx)a+a} (a^2(10A+9B) - a^2(14A+3B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx + \frac{6a^2(2A+B)\sin(c+dx)\sqrt{a \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{3} \left(\int \frac{\sqrt{\cos(c+dx)a+a} (a^2(10A+9B) - a^2(14A+3B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx + \frac{6a^2(2A+B)\sin(c+dx)\sqrt{a \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a} (a^2(10A+9B) - a^2(14A+3B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^2(2A+B)\sin(c+dx)\sqrt{a \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3460

$$\frac{1}{3} \left(\frac{3}{2} a^2(2A+5B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx - \frac{a^3(14A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{6a^2(2A+B)\sin(c+dx)\sqrt{a \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(\frac{3}{2} a^2(2A+5B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{a^3(14A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{6a^2(2A+B)\sin(c+dx)\sqrt{a \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

3.186. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

↓ 3253

$$\frac{1}{3} \left(\frac{3a^2(2A + 5B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} - \frac{a^3(14A + 3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx) + a}} + \frac{6a^2(2A + B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) - \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 223

$$\frac{1}{3} \left(\frac{3a^{5/2}(2A + 5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx) + a}}\right)}{d} - \frac{a^3(14A + 3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx) + a}} + \frac{6a^2(2A + B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) - \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2), x]`

output `(2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((3*a^(5/2)*(2*A + 5*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (a^3*(14*A + 3*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]) + (6*a^2*(2*A + B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/3`

3.186.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.186.4 Maple [A] (verified)

Time = 18.65 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.54

method	result
default	$a^2 \left(3B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 6A \cos^2(dx+c) \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + 16A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) + 3d(1+\cos(dx+c))$
parts	$\frac{2A \left(3 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \cos^2(dx+c) \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + 3 \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{3d(1+\cos(dx+c)) \cos(dx+c)^{\frac{3}{2}}}$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2), x, method=_RET URNVERBOSE)`

$$3.186. \int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

output $\frac{1}{3}a^2/d*(3B*\cos(dx+c)^2*\sin(dx+c)*(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)}+6*A*\cos(dx+c)^2*\arctan(\tan(dx+c)*(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)})+16*A*\cos(dx+c)*\sin(dx+c)*(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)}+15*B*\cos(dx+c)^2*\arctan(\tan(dx+c)*(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)})+6*B*\cos(dx+c)*\sin(dx+c)*(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)}+2*A*(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)}*\sin(dx+c))*(a*(1+\cos(dx+c)))^{(1/2)}/(1+\cos(dx+c))/(\cos(dx+c)/(1+\cos(dx+c)))^{(1/2)}/\cos(dx+c)^{(3/2)}$

3.186.5 Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.98

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \frac{(3 B a^2 \cos(dx + c)^2 + 2 (8 A + 3 B) a^2 \cos(dx + c) + 2 A a^2) \sqrt{\cos(dx + c)} \sin(dx + c) - 3 ((2 A + 5 B) a^2 \cos(dx + c)^3 + (2 A + 5 B) a^2 \cos(dx + c)^2) \sqrt{a} \arctan(\sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}) / (\sqrt{a} \sin(dx + c))}{d \cos(dx + c)^3 + d \cos(dx + c)^2}$$

input `integrate((a+a*cos(dx+c))^(5/2)*(A+B*cos(dx+c))/cos(dx+c)^(5/2),x, algorith="fricas")`

output $\frac{1}{3}*((3*B*a^2*\cos(dx + c)^2 + 2*(8*A + 3*B)*a^2*\cos(dx + c) + 2*A*a^2)*\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)}*\sin(dx + c) - 3*((2*A + 5*B)*a^2*\cos(dx + c)^3 + (2*A + 5*B)*a^2*\cos(dx + c)^2)*\sqrt{a}*\arctan(\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)})/(\sqrt{a}*\sin(dx + c)))/d*\cos(dx + c)^3 + d*\cos(dx + c)^2)$

3.186.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(dx+c))**(5/2)*(A+B*cos(dx+c))/cos(dx+c)**(5/2),x)`

output Timed out

3.186.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2370 vs. $2(151) = 302$.

Time = 0.56 (sec) , antiderivative size = 2370, normalized size of antiderivative = 13.70

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/12*(3*(2*(a^2*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (a^2*cos(d*x + c) - a^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*sqrt(a) + 5*(a^2*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4))*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) + 1 - a^2*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - a^2*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + 1) + a^2*arctan2((cos(2*d*x + 2*c)^2 ...`

3.186.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`

3.186. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$

output Timed out

3.186.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(5/2),x)`

$$3.187 \quad \int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

3.187.1 Optimal result	1778
3.187.2 Mathematica [A] (verified)	1778
3.187.3 Rubi [A] (verified)	1779
3.187.4 Maple [A] (verified)	1782
3.187.5 Fricas [A] (verification not implemented)	1783
3.187.6 Sympy [F(-1)]	1783
3.187.7 Maxima [B] (verification not implemented)	1784
3.187.8 Giac [F(-1)]	1784
3.187.9 Mupad [F(-1)]	1785

3.187.1 Optimal result

Integrand size = 35, antiderivative size = 172

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx = \frac{2a^{5/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{d}$$

$$+ \frac{2a^3(32A+35B) \sin(c+dx)}{15d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} + \frac{2a^2(8A+5B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{15d \cos^{3/2}(c+dx)}$$

$$+ \frac{2aA(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{5d \cos^{5/2}(c+dx)}$$

```
output 2*a^(5/2)*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d+2/5*a*A*(a
+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(5/2)+2/15*a^3*(32*A+35*B)*si
n(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)+2/15*a^2*(8*A+5*B)*sin(
d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)
```

3.187.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.76

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx = \frac{a^2 \sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \left(30\sqrt{2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)\right)}{d}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*(30*sqrt[2]*B*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^(5/2) + 2*(49*A + 40*B + 2*(14*A + 5*B)*Cos[c + d*x] + (43*A + 40*B)*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(30*d*Cos[c + d*x]^(5/2))`

3.187.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

↓ 3454

$$\frac{2}{5} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(8A + 5B) + 5aB \cos(c + dx))}{2 \cos^{5/2}(c + dx)} dx + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d \cos^{5/2}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(8A + 5B) + 5aB \cos(c + dx))}{\cos^{5/2}(c + dx)} dx + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d \cos^{5/2}(c + dx)}$$

↓ 3042

3.187. $\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$

$$\frac{1}{5} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2} (a(8A+5B)+5aB \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

↓ 3454

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{\sqrt{\cos(c+dx)a+a}((32A+35B)a^2+15B \cos(c+dx)a^2)}{2 \cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a^2(8A+5B) \sin(c+dx) \sqrt{a \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{\sqrt{\cos(c+dx)a+a}((32A+35B)a^2+15B \cos(c+dx)a^2)}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a^2(8A+5B) \sin(c+dx) \sqrt{a \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((32A+35B)a^2+15B \sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2a^2(8A+5B) \sin(c+dx) \sqrt{a \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

↓ 3459

$$\frac{1}{5} \left(\frac{1}{3} \left(15a^2B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx + \frac{2a^3(32A+35B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} \right) + \frac{2a^2(8A+5B) \sin(c+dx) \sqrt{a \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(15a^2B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a^3(32A+35B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} \right) + \frac{2a^2(8A+5B) \sin(c+dx) \sqrt{a \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

3.187. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3253} \\
 & \frac{1}{5} \left(\frac{1}{3} \left(\frac{2a^3(32A + 35B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}\sqrt{a \cos(c + dx) + a}} - \frac{30a^2 B \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d} \right) + \frac{2a^2(8A + 5B) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) \\
 & \quad + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \\
 & \quad \downarrow \text{223} \\
 & \frac{1}{5} \left(\frac{2a^2(8A + 5B) \sin(c + dx)\sqrt{a \cos(c + dx) + a}}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{1}{3} \left(\frac{30a^{5/2} B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{2a^3(32A + 35B) \sin(c + dx)}{d\sqrt{\cos(c + dx)}\sqrt{a \cos(c + dx) + a}} \right) \right) \\
 & \quad + \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)}
 \end{aligned}$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*a^2*(8*A + 5*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2))) + ((30*a^(5/2)*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (2*a^3*(32*A + 35*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/3)/5`

3.187.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.187. $\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.187.4 Maple [A] (verified)

Time = 7.94 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.32

method	result
default	$\frac{2a^2 \left(15B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) (\cos^3(dx+c)) + 15B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) (\cos^3(dx+c)) \right)}{\dots}$
parts	$\frac{2A \sin(dx+c) (43 (\cos^2(dx+c)) + 14 \cos(dx+c) + 3) \sqrt{a(1+\cos(dx+c))} a^2}{15d(1+\cos(dx+c)) \cos(dx+c)^{\frac{5}{2}}} + \frac{2B \left(3 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} (\cos^2(dx+c)) \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{\dots}$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2), x, method=_RET URNVERBOSE)`

3.187.
$$\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

output $2/15*a^2/d*(15*B*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})*\cos(d*x+c)^3+15*B*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})*\cos(d*x+c)^2+43*A*\sin(d*x+c)*\cos(d*x+c)^2+40*B*\sin(d*x+c)*\cos(d*x+c)^2+14*A*\sin(d*x+c)*\cos(d*x+c)+5*B*\sin(d*x+c)*\cos(d*x+c)+3*A*\sin(d*x+c)*(a*(1+\cos(d*x+c)))^{(1/2)}/(1+\cos(d*x+c))/\cos(d*x+c)^{(5/2)}$

3.187.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.94

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \frac{2 \left(((43A + 40B)a^2 \cos(dx + c)^2 + (14A + 5B)a^2 \cos(dx + c) + 3Aa^2) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c) - 15(Ba^2 \cos(dx + c)^4 + Ba^2 \cos(dx + c)^3) \sqrt{a} \arctan(\sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}) / (\sqrt{a} \sin(dx + c)) \right)}{d \cos(dx + c)^4 + d \cos(dx + c)^3}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo rithm="fracas")`

output $2/15*(((43*A + 40*B)*a^2*\cos(d*x + c)^2 + (14*A + 5*B)*a^2*\cos(d*x + c) + 3*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c) - 15*(B*a^2*\cos(d*x + c)^4 + B*a^2*\cos(d*x + c)^3)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c)))/(d*\cos(d*x + c)^4 + d*\cos(d*x + c)^3)$

3.187.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output Timed out

3.187.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1548 vs. $2(148) = 296$.

Time = 0.47 (sec) , antiderivative size = 1548, normalized size of antiderivative = 9.00

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorith="maxima")`

output

```
1/30*(5*(30*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(3/4)*a^(5/2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*((12*a^2*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(2*d*x + 2*c) - 3*a^2*sin(2*d*x + 2*c) - 4*(3*a^2*cos(2*d*x + 2*c) + 4*a^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + (12*a^2*sin(2*d*x + 2*c)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 3*a^2*cos(2*d*x + 2*c) - a^2 + 4*(3*a^2*cos(2*d*x + 2*c) + 4*a^2)*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(a) + 3*((a^2*cos(2*d*x + 2*c)^2 + a^2*sin(2*d*x + 2*c)^2 + 2*a^2*cos(2*d*x + 2*c) + a^2)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) + 1) - (a^2*cos(2*d*x + 2*c)^2 + a^2*sin(...
```

3.187.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorith="giac")`

output Timed out

3.187.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\cos(c + dx)^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(7/2),x)`

$$3.188 \quad \int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.188.1 Optimal result 1786
 3.188.2 Mathematica [A] (verified) 1786
 3.188.3 Rubi [A] (verified) 1787
 3.188.4 Maple [A] (verified) 1790
 3.188.5 Fricas [A] (verification not implemented) 1790
 3.188.6 Sympy [F(-1)] 1791
 3.188.7 Maxima [B] (verification not implemented) 1791
 3.188.8 Giac [C] (verification not implemented) 1792
 3.188.9 Mupad [B] (verification not implemented) 1793

3.188.1 Optimal result

Integrand size = 35, antiderivative size = 181

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{2a^3(10A+11B) \sin(c+dx)}{15d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a^3(230A+301B) \sin(c+dx)}{105d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} + \frac{2a^2(10A+7B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{35d \cos^{\frac{5}{2}}(c+dx)} + \frac{2aA(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)}$$

```
output 2/7*a*A*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(7/2)+2/15*a^3*(10*
A+11*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+2/105*a^3*(23
0*A+301*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)+2/35*a^2*(
10*A+7*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)
```

3.188.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.57

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{a^2 \sqrt{a(1+\cos(c+dx))}(290A+196B+(930A+987B))}{\cos^{\frac{9}{2}}(c+dx)}$$

```
input Integrate[((a+a*Cos[c+d*x])^(5/2)*(A+B*Cos[c+d*x]))/Cos[c+d*x]^(
9/2),x]
```

3.188. $\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

output $(a^2 \sqrt{a(1 + \cos[c + dx])} (290A + 196B + (930A + 987B) \cos[c + dx] + 2(115A + 98B) \cos[2(c + dx)] + 230A \cos[3(c + dx)] + 301B \cos[3(c + dx)]) \tan[(c + dx)/2]) / (210d \cos[c + dx]^{7/2})$

3.188.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.06, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx$$

↓ 3454

$$\frac{2}{7} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(10A + 7B) + a(2A + 7B) \cos(c + dx))}{2 \cos^{7/2}(c + dx)} dx + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{7d \cos^{7/2}(c + dx)}$$

↓ 27

$$\frac{1}{7} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(10A + 7B) + a(2A + 7B) \cos(c + dx))}{\cos^{7/2}(c + dx)} dx + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{7d \cos^{7/2}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (a(10A + 7B) + a(2A + 7B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx + \frac{2aA \sin(c + dx) (a \cos(c + dx) + a)^{3/2}}{7d \cos^{7/2}(c + dx)}$$

↓ 3454

3.188. $\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx$

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{\sqrt{\cos(c+dx)a+a}(7(10A+11B)a^2+(30A+49B)\cos(c+dx)a^2)}{2\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^2(10A+7B)\sin(c+dx)\sqrt{a}}{5d\cos^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \downarrow 27$$

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{\sqrt{\cos(c+dx)a+a}(7(10A+11B)a^2+(30A+49B)\cos(c+dx)a^2)}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^2(10A+7B)\sin(c+dx)\sqrt{a}}{5d\cos^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \downarrow 3042$$

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(7(10A+11B)a^2+(30A+49B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2a^2(10A+7B)\sin(c+dx)\sqrt{a}}{5d\cos^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \downarrow 3459$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} a^2(230A+301B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{14a^3(10A+11B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2a^2(10A+7B)\sin(c+dx)\sqrt{a}}{5d\cos^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \downarrow 3042$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} a^2(230A+301B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{14a^3(10A+11B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) + \frac{2a^2(10A+7B)\sin(c+dx)\sqrt{a}}{5d\cos^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \downarrow 3250$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{14a^3(10A+11B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} + \frac{2a^3(230A+301B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right) + \frac{2a^2(10A+7B)\sin(c+dx)\sqrt{a}}{5d\cos^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2aA\sin(c+dx)(a\cos(c+dx)+a)^{3/2}}{7d\cos^{\frac{7}{2}}(c+dx)} \right)$$

input `Int[((a + a*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]))/cos[c + d*x]^(9/2),x]`

output `(2*a*A*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)) + ((2*a^2*(10*A + 7*B)*sqrt[a + a*cos[c + d*x]]*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)) + ((14*a^3*(10*A + 11*B)*sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)*sqrt[a + a*cos[c + d*x]]) + (2*a^3*(230*A + 301*B)*sin[c + d*x])/(3*d*sqrt[cos[c + d*x]]*sqrt[a + a*cos[c + d*x]]))/5/7`

3.188.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*sqrt[a + b*sin[e + f*x]]*sqrt[c + d*sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*sin[e + f*x])^(m - 1)*((c + d*sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*sin[e + f*x])^(m - 1)*(c + d*sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.188.4 Maple [A] (verified)

Time = 8.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.61

method	result
default	$\frac{2a^2 \sin(dx+c)(230A(\cos^3(dx+c))+301B(\cos^3(dx+c))+115A(\cos^2(dx+c))+98B(\cos^2(dx+c))+60A \cos(dx+c)+21B \cos(dx+c)+15A)}{105d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}}$
parts	$\frac{2A \sin(dx+c)(46(\cos^3(dx+c))+23(\cos^2(dx+c))+12 \cos(dx+c)+3) \sqrt{a(1+\cos(dx+c))} a^2}{21d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}} + \frac{2B \sin(dx+c)(43(\cos^2(dx+c))+14 \cos(dx+c)+3) a^2}{15d(1+\cos(dx+c)) \cos(dx+c)^{\frac{7}{2}}}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RET
URNVERBOSE)
```

```
output 2/105*a^2/d*sin(d*x+c)*(230*A*cos(d*x+c)^3+301*B*cos(d*x+c)^3+115*A*cos(d*
x+c)^2+98*B*cos(d*x+c)^2+60*A*cos(d*x+c)+21*B*cos(d*x+c)+15*A)*(a*(1+cos(d
*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(7/2)
```

3.188.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.63

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \frac{2((230A + 301B)a^2 \cos(dx + c)^3 + (115A + 98B)a^2 \cos(dx + c)^2 + (60A + 21B)a^2 \cos(dx + c) + 15A)a^2}{105d(1 + \cos(dx + c)) \cos(dx + c)^{\frac{7}{2}}} + \frac{2B \sin(dx + c)(43 \cos^2(dx + c) + 14 \cos(dx + c) + 3)a^2}{15d(1 + \cos(dx + c)) \cos(dx + c)^{\frac{7}{2}}}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="fracas")
```

output $2/105*((230*A + 301*B)*a^2*\cos(d*x + c)^3 + (115*A + 98*B)*a^2*\cos(d*x + c)^2 + 3*(20*A + 7*B)*a^2*\cos(d*x + c) + 15*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^5 + d*\cos(d*x + c)^4)$

3.188.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output Timed out

3.188.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 396 vs. $2(157) = 314$.

Time = 0.34 (sec) , antiderivative size = 396, normalized size of antiderivative = 2.19

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \frac{8 \left(\frac{7 \left(\frac{15 \sqrt{2} a^{5/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{35 \sqrt{2} a^{5/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{28 \sqrt{2} a^{5/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{8 \sqrt{2} a^{5/2}}{\cos(dx+c)+1} \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{7/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{7/2}} \right)}{1}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo rithm="maxima")`

```
output 8/105*(7*(15*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 35*sqrt(2)*
a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 28*sqrt(2)*a^(5/2)*sin(d*x +
c)^5/(cos(d*x + c) + 1)^5 - 8*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x + c
) + 1)^7)*B/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(-sin(d*x + c)/(c
os(d*x + c) + 1) + 1)^(7/2)) + 5*(21*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x
+ c) + 1) - 56*sqrt(2)*a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 63*s
qrt(2)*a^(5/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 36*sqrt(2)*a^(5/2)*si
n(d*x + c)^7/(cos(d*x + c) + 1)^7 + 8*sqrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(
d*x + c) + 1)^9)*A*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^2/((sin(d*x +
c)/(cos(d*x + c) + 1) + 1)^(9/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(
9/2)*(2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + sin(d*x + c)^4/(cos(d*x + c)
+ 1)^4 + 1))) / d
```

3.188.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 244.27 (sec) , antiderivative size = 169051, normalized size of antiderivative = 933.98

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="giac")
```

output

```
-33554432/105*sqrt(2)*sqrt(-tan(1/4*d*x + c)^4*tan(1/2*c)^8 + 14*tan(1/4*d*x + c)^4*tan(1/2*c)^6 - 24*tan(1/4*d*x + c)^3*tan(1/2*c)^7 + 6*tan(1/4*d*x + c)^2*tan(1/2*c)^8 + 56*tan(1/4*d*x + c)^3*tan(1/2*c)^5 - 84*tan(1/4*d*x + c)^2*tan(1/2*c)^6 + 24*tan(1/4*d*x + c)*tan(1/2*c)^7 - tan(1/2*c)^8 - 14*tan(1/4*d*x + c)^4*tan(1/2*c)^2 + 56*tan(1/4*d*x + c)^3*tan(1/2*c)^3 - 56*tan(1/4*d*x + c)*tan(1/2*c)^5 + 14*tan(1/2*c)^6 + tan(1/4*d*x + c)^4 - 24*tan(1/4*d*x + c)^3*tan(1/2*c) + 84*tan(1/4*d*x + c)^2*tan(1/2*c)^2 - 56*tan(1/4*d*x + c)*tan(1/2*c)^3 - 6*tan(1/4*d*x + c)^2 + 24*tan(1/4*d*x + c)*tan(1/2*c) - 14*tan(1/2*c)^2 + 1)*((((((((((((((((((230*I*A*a^2*e^(1027/2*I*c) + 301*I*B*a^2*e^(1027/2*I*c) + 89700*I*A*a^2*e^(1025/2*I*c) + 117390*I*B*a^2*e^(1025/2*I*c) + 17446650*I*A*a^2*e^(1023/2*I*c) + 22832355*I*B*a^2*e^(1023/2*I*c) + 2256433400*I*A*a^2*e^(1021/2*I*c) + 2952984475*I*B*a^2*e^(1021/2*I*c) + 218309931450*I*A*a^2*e^(1019/2*I*c) + 285701217165*I*B*a^2*e^(1019/2*I*c) + 16853526707940*I*A*a^2*e^(1017/2*I*c) + 22056129161703*I*B*a^2*e^(1017/2*I*c) + 1081434630427540*I*A*a^2*e^(1015/2*I*c) + 1415267768839799*I*B*a^2*e^(1015/2*I*c) + 59324414012490900*I*A*a^2*e^(1013/2*I*c) + 77637503022633645*I*B*a^2*e^(1013/2*I*c) + 2840156320927487250*I*A*a^2*e^(1011/2*I*c) + 3716892534681247740*I*B*a^2*e^(1011/2*I*c) + 120548857186305888250*I*A*a^2*e^(1009/2*I*c) + 157761271574607074465*I*B*a^2*e^(1009/2*I*c) + 4592911459598047037790*I*A*a^2*e^(1007/2*I*c) + 601069617499231...
```

3.188.9 Mupad [B] (verification not implemented)

Time = 7.19 (sec) , antiderivative size = 551, normalized size of antiderivative = 3.04

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \frac{\sqrt{a + a \left(\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2} \right)}}{\sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}} + e^{c \operatorname{li} + dx \operatorname{li}} \sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}}}$$

input

```
int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(9/2),x)
```


output $((a + a*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2))^{(1/2)}*((a^2*(230*A + 301*B)*2i)/(105*d) - (a^2*\exp(c*3i + d*x*3i)*(10*A + 17*B)*2i)/(3*d) + (a^2*\exp(c*4i + d*x*4i)*(10*A + 17*B)*2i)/(3*d) + (a^2*\exp(c*2i + d*x*2i)*(100*A + 113*B)*2i)/(15*d) - (a^2*\exp(c*5i + d*x*5i)*(100*A + 113*B)*2i)/(15*d) - (a^2*\exp(c*7i + d*x*7i)*(230*A + 301*B)*2i)/(105*d) - (B*a^2*\exp(c*1i + d*x*1i)*2i)/d + (B*a^2*\exp(c*6i + d*x*6i)*2i)/d)/((\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + \exp(c*1i + d*x*1i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*2i + d*x*2i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*3i + d*x*3i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*4i + d*x*4i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + 3*\exp(c*5i + d*x*5i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + \exp(c*6i + d*x*6i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2)} + \exp(c*7i + d*x*7i)*(\exp(-c*1i - d*x*1i)/2 + \exp(c*1i + d*x*1i)/2)^{(1/2}))$

3.188. $\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

3.189
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

3.189.1 Optimal result 1795
 3.189.2 Mathematica [A] (verified) 1796
 3.189.3 Rubi [A] (verified) 1796
 3.189.4 Maple [A] (verified) 1800
 3.189.5 Fricas [A] (verification not implemented) 1800
 3.189.6 Sympy [F(-1)] 1801
 3.189.7 Maxima [B] (verification not implemented) 1801
 3.189.8 Giac [F(-1)] 1802
 3.189.9 Mupad [B] (verification not implemented) 1802

3.189.1 Optimal result

Integrand size = 35, antiderivative size = 228

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{2a^3(124A+135B) \sin(c+dx)}{315d \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a^3(292A+345B) \sin(c+dx)}{315d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{4a^3(292A+345B) \sin(c+dx)}{315d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} + \frac{2a^2(4A+3B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{21d \cos^{\frac{7}{2}}(c+dx)} + \frac{2aA(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)}$$

output

```
2/9*a*A*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(9/2)+2/315*a^3*(12
4*A+135*B)*sin(d*x+c)/d/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+2/315*a^3*
(292*A+345*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+4/315*a
^3*(292*A+345*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)+2/21
*a^2*(4*A+3*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(7/2)
```

3.189.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.55

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (1454A + 1395B + (1396A + 1215B) \cos(c + dx) + 2(803A + 870B) \cos(2(c + dx)) + 292A \cos(3(c + dx)) + 345B \cos(3(c + dx)) + 292A \cos(4(c + dx)) + 345B \cos(4(c + dx))) \operatorname{Tan}[(c + dx)/2]}{630d \cos^{9/2}(c + dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*(1454*A + 1395*B + (1396*A + 1215*B)*Cos[c + d*x] + 2*(803*A + 870*B)*Cos[2*(c + d*x)] + 292*A*Cos[3*(c + d*x)] + 345*B*Cos[3*(c + d*x)] + 292*A*Cos[4*(c + d*x)] + 345*B*Cos[4*(c + d*x)])*Tan[(c + d*x)/2])/(630*d*cos[c + d*x]^(9/2))`

3.189.3 Rubi [A] (verified)Time = 1.24 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx \\ & \quad \downarrow \text{3454} \\ & \frac{2}{9} \int \frac{(\cos(c + dx)a + a)^{3/2} (3a(4A + 3B) + a(4A + 9B) \cos(c + dx))}{2 \cos^{9/2}(c + dx)} dx + \\ & \quad \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{9d \cos^{9/2}(c + dx)} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.189. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

$$\frac{1}{9} \int \frac{(\cos(c+dx)a+a)^{3/2}(3a(4A+3B)+a(4A+9B)\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(3a(4A+3B)+a(4A+9B)\sin(c+dx+\frac{\pi}{2}))}{\sin^{\frac{9}{2}}(c+dx+\frac{\pi}{2})} dx + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3454

$$\frac{1}{9} \left(\frac{2}{7} \int \frac{\sqrt{\cos(c+dx)a+a}((124A+135B)a^2+(76A+99B)\cos(c+dx)a^2)}{2 \cos^{\frac{7}{2}}(c+dx)} dx + \frac{6a^2(4A+3B)\sin(c+dx)\sqrt{a \cos(c+dx)+a}}{7d \cos^{\frac{7}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{1}{7} \int \frac{\sqrt{\cos(c+dx)a+a}((124A+135B)a^2+(76A+99B)\cos(c+dx)a^2)}{\cos^{\frac{7}{2}}(c+dx)} dx + \frac{6a^2(4A+3B)\sin(c+dx)\sqrt{a \cos(c+dx)+a}}{7d \cos^{\frac{7}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((124A+135B)a^2+(76A+99B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin^{\frac{7}{2}}(c+dx+\frac{\pi}{2})} dx + \frac{6a^2(4A+3B)\sin(c+dx)\sqrt{a \cos(c+dx)+a}}{7d \cos^{\frac{7}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3459

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} a^2(292A+345B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a^3(124A+135B)\sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} \right) + \frac{6a^2(4A+3B)\sin(c+dx)\sqrt{a \cos(c+dx)+a}}{7d \cos^{\frac{7}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

3.189. $\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} a^2 (292A + 345B) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{5/2}} dx + \frac{2a^3(124A + 135B) \sin(c+dx)}{5d \cos^{5/2}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{6a^2(4A + 3B) \sin(c+dx) \sqrt{a \cos(c+dx) + a}}{7d \cos^{7/2}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{9d \cos^{9/2}(c+dx)}$$

↓ 3251

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} a^2 (292A + 345B) \left(\frac{2}{3} \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{3/2}(c+dx)} dx + \frac{2a \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^3(124A + 135B) \sin(c+dx)}{5d \cos^{5/2}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{6a^2(4A + 3B) \sin(c+dx) \sqrt{a \cos(c+dx) + a}}{7d \cos^{7/2}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{9d \cos^{9/2}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} a^2 (292A + 345B) \left(\frac{2}{3} \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^3(124A + 135B) \sin(c+dx)}{5d \cos^{5/2}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{6a^2(4A + 3B) \sin(c+dx) \sqrt{a \cos(c+dx) + a}}{7d \cos^{7/2}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{9d \cos^{9/2}(c+dx)}$$

↓ 3250

$$\frac{1}{9} \left(\frac{6a^2(4A + 3B) \sin(c+dx) \sqrt{a \cos(c+dx) + a}}{7d \cos^{7/2}(c+dx)} + \frac{1}{7} \left(\frac{2a^3(124A + 135B) \sin(c+dx)}{5d \cos^{5/2}(c+dx) \sqrt{a \cos(c+dx) + a}} + \frac{3}{5} a^2 (292A + 345B) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{9d \cos^{9/2}(c+dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2), x]`

output `(2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((6*a^2*(4*A + 3*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2))) + ((2*a^3*(124*A + 135*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x])) + (3*a^2*(292*A + 345*B)*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x])) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/5)/7)/9`

3.189.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`
- rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.189.4 Maple [A] (verified)

Time = 7.87 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.58

method	result
default	$\frac{2a^2 \sin(dx+c)(584A(\cos^4(dx+c))+690B(\cos^4(dx+c))+292A(\cos^3(dx+c))+345B(\cos^3(dx+c))+219A(\cos^2(dx+c))+180B(\cos^2(dx+c)))}{315d(1+\cos(dx+c)) \cos(dx+c)^{\frac{9}{2}}}$
parts	$\frac{2A \sin(dx+c)(584(\cos^4(dx+c))+292(\cos^3(dx+c))+219(\cos^2(dx+c))+130 \cos(dx+c)+35)\sqrt{a(1+\cos(dx+c))} a^2}{315d(1+\cos(dx+c)) \cos(dx+c)^{\frac{9}{2}}} + \frac{2B \sin(dx+c)}{\cos(dx+c)}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x,method=_RE
TURNVERBOSE)
```

```
output 2/315*a^2/d*sin(d*x+c)*(584*A*cos(d*x+c)^4+690*B*cos(d*x+c)^4+292*A*cos(d*
x+c)^3+345*B*cos(d*x+c)^3+219*A*cos(d*x+c)^2+180*B*cos(d*x+c)^2+130*A*cos(
d*x+c)+45*B*cos(d*x+c)+35*A)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d
*x+c)^(9/2)
```

3.189.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.59

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx = \frac{2(2(292A + 345B)a^2 \cos(dx + c)^4 + (292A + 345B)a^2 \cos(dx + c)^3 + \dots)}{\cos^{\frac{11}{2}}(c + dx)}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, alg
orithm="fracas")
```

3.189. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

output $\frac{2}{315}*(2*(292*A + 345*B)*a^2*\cos(d*x + c)^4 + (292*A + 345*B)*a^2*\cos(d*x + c)^3 + 3*(73*A + 60*B)*a^2*\cos(d*x + c)^2 + 5*(26*A + 9*B)*a^2*\cos(d*x + c) + 35*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c)^6 + d*\cos(d*x + c)^5)$

3.189.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(11/2),x)`

output `Timed out`

3.189.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 533 vs. 2(198) = 396.

Time = 0.35 (sec) , antiderivative size = 533, normalized size of antiderivative = 2.34

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \frac{8 \left(15 \left(\frac{21 \sqrt{2} a^{5/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{56 \sqrt{2} a^{5/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{63 \sqrt{2} a^{5/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{36 \sqrt{2} a^{5/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(\frac{2}{\cos(dx+c)+1} \right)^{9/2}}{\dots}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="maxima")`

output
$$\frac{8/315*(15*(21*\sqrt{2})*a^{(5/2)}*\sin(dx + c)/(\cos(dx + c) + 1) - 56*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 63*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 36*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^7/(\cos(dx + c) + 1)^7 + 8*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^9/(\cos(dx + c) + 1)^9)*B*(\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 1)^2/((\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(9/2)}*(-\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(9/2)}*(2*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + \sin(dx + c)^4/(\cos(dx + c) + 1)^4 + 1)) + (315*\sqrt{2})*a^{(5/2)}*\sin(dx + c)/(\cos(dx + c) + 1) - 945*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 1449*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 1287*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^7/(\cos(dx + c) + 1)^7 + 572*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^9/(\cos(dx + c) + 1)^9 - 104*\sqrt{2})*a^{(5/2)}*\sin(dx + c)^{11}/(\cos(dx + c) + 1)^{11})*A*(\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 1)^3/((\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(11/2)}*(-\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{(11/2)}*(3*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 3*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + \sin(dx + c)^6/(\cos(dx + c) + 1)^6 + 1)))}/d$$

3.189.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(dx+c))^(5/2)*(A+B*cos(dx+c))/cos(dx+c)^(11/2),x, algorithm="giac")`

output Timed out

3.189.9 Mupad [B] (verification not implemented)

Time = 8.70 (sec) , antiderivative size = 647, normalized size of antiderivative = 2.84

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \frac{\sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}} + e^{c \operatorname{li} + dx \operatorname{li}} \sqrt{\frac{e^{-c \operatorname{li} - dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li} + dx \operatorname{li}}}{2}}}{\cos^{11/2}(c + dx)}$$

3.189.
$$\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$$

input `int((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(11/2),
x)`

output `((a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a^2*(292*
A + 345*B)*4i)/(315*d) - (a^2*exp(c*3i + d*x*3i)*(2*A + 5*B)*4i)/(3*d) + (
a^2*exp(c*6i + d*x*6i)*(2*A + 5*B)*4i)/(3*d) + (a^2*exp(c*4i + d*x*4i)*(24
*A + 25*B)*4i)/(5*d) - (a^2*exp(c*5i + d*x*5i)*(24*A + 25*B)*4i)/(5*d) + (
a^2*exp(c*2i + d*x*2i)*(146*A + 155*B)*4i)/(35*d) - (a^2*exp(c*7i + d*x*7i
)*(146*A + 155*B)*4i)/(35*d) - (a^2*exp(c*9i + d*x*9i)*(292*A + 345*B)*4i
/(315*d)))/((exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + exp(c*
1i + d*x*1i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + 4*exp
(c*2i + d*x*2i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + 4*
exp(c*3i + d*x*3i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) +
6*exp(c*4i + d*x*4i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2
) + 6*exp(c*5i + d*x*5i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(
1/2) + 4*exp(c*6i + d*x*6i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2
)^(1/2) + 4*exp(c*7i + d*x*7i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i
) /2)^(1/2) + exp(c*8i + d*x*8i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1
i)/2)^(1/2) + exp(c*9i + d*x*9i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*
1i)/2)^(1/2))`

3.189.
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$$

3.190
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$$

3.190.1 Optimal result 1804
 3.190.2 Mathematica [A] (verified) 1805
 3.190.3 Rubi [A] (verified) 1805
 3.190.4 Maple [A] (verified) 1809
 3.190.5 Fricas [A] (verification not implemented) 1810
 3.190.6 Sympy [F(-1)] 1810
 3.190.7 Maxima [B] (verification not implemented) 1810
 3.190.8 Giac [F(-1)] 1811
 3.190.9 Mupad [B] (verification not implemented) 1812

3.190.1 Optimal result

Integrand size = 35, antiderivative size = 275

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx = \frac{2a^3(194A+209B) \sin(c+dx)}{693d \cos^{\frac{7}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{2a^3(710A+803B) \sin(c+dx)}{1155d \cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{8a^3(710A+803B) \sin(c+dx)}{3465d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} + \frac{16a^3(710A+803B) \sin(c+dx)}{3465d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} + \frac{2a^2(14A+11B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{99d \cos^{\frac{9}{2}}(c+dx)} + \frac{2aA(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{11d \cos^{\frac{11}{2}}(c+dx)}$$

output

```
2/11*a*A*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(11/2)+2/693*a^3*(194*A+209*B)*sin(d*x+c)/d/cos(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2)+2/1155*a^3*(710*A+803*B)*sin(d*x+c)/d/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+8/3465*a^3*(710*A+803*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+16/3465*a^3*(710*A+803*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)+2/99*a^2*(14*A+11*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(9/2)
```

3.190.2 Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.53

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (9070A + 7678B + (25070A + 24827B) \cos(c + dx) + (9230A + 9284B) \cos(2(c + dx)) + 9230A \cos(3(c + dx)) + 10439B \cos(3(c + dx)) + 1420A \cos(4(c + dx)) + 1606B \cos(4(c + dx)) + 1420A \cos(5(c + dx)) + 1606B \cos(5(c + dx))) \tan((c + dx)/2)}{(6930d \cos(c + dx))^{11/2}}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(13/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x]])*(9070*A + 7678*B + (25070*A + 24827*B)*Cos[c + d*x] + (9230*A + 9284*B)*Cos[2*(c + d*x)] + 9230*A*cos[3*(c + d*x)] + 10439*B*cos[3*(c + d*x)] + 1420*A*cos[4*(c + d*x)] + 1606*B*cos[4*(c + d*x)] + 1420*A*cos[5*(c + d*x)] + 1606*B*cos[5*(c + d*x)])*Tan[(c + d*x)/2])/(6930*d*cos[c + d*x]^(11/2))`

3.190.3 Rubi [A] (verified)Time = 1.48 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{13/2}} dx \\ & \quad \downarrow \text{3454} \\ & \frac{2}{11} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(14A + 11B) + a(6A + 11B) \cos(c + dx))}{2 \cos^{11/2}(c + dx)} dx + \\ & \quad \frac{2aA \sin(c + dx)(a \cos(c + dx) + a)^{3/2}}{11d \cos^{11/2}(c + dx)} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.190. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{13/2}(c+dx)} dx$

$$\begin{aligned}
& \frac{1}{11} \int \frac{(\cos(c+dx)a+a)^{3/2}(a(14A+11B)+a(6A+11B)\cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{11} \int \frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(14A+11B)+a(6A+11B)\sin(c+dx+\frac{\pi}{2}))}{\sin^{\frac{11}{2}}(c+dx+\frac{\pi}{2})} dx + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \\
& \quad \downarrow \text{3454} \\
& \frac{1}{11} \left(\frac{2}{9} \int \frac{\sqrt{\cos(c+dx)a+a}((194A+209B)a^2+3(46A+55B)\cos(c+dx)a^2)}{2 \cos^{\frac{9}{2}}(c+dx)} dx + \frac{2a^2(14A+11B)\sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{11} \left(\frac{1}{9} \int \frac{\sqrt{\cos(c+dx)a+a}((194A+209B)a^2+3(46A+55B)\cos(c+dx)a^2)}{\cos^{\frac{9}{2}}(c+dx)} dx + \frac{2a^2(14A+11B)\sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{11} \left(\frac{1}{9} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((194A+209B)a^2+3(46A+55B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sin^{\frac{9}{2}}(c+dx+\frac{\pi}{2})} dx + \frac{2a^2(14A+11B)\sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \\
& \quad \downarrow \text{3459} \\
& \frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2(710A+803B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{7}{2}}(c+dx)} dx + \frac{2a^3(194A+209B)\sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} \right) + \frac{2a^2(14A+11B)\sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \\
& \quad \frac{2aA \sin(c+dx)(a \cos(c+dx)+a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.190. $\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (710A + 803B) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{7/2}} dx + \frac{2a^3(194A + 209B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^2(14A + 11B) \sin(c+dx) \sqrt{a \cos(c+dx) + a}}{9d \cos^{\frac{9}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3251

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (710A + 803B) \left(\frac{4}{5} \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)} dx + \frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^3(194A + 209B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (710A + 803B) \left(\frac{4}{5} \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{5/2}} dx + \frac{2a \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^3(194A + 209B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \right)$$

↓ 3251

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (710A + 803B) \left(\frac{4}{5} \left(\frac{2}{3} \int \frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^3(194A + 209B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{3}{7} a^2 (710A + 803B) \left(\frac{4}{5} \left(\frac{2}{3} \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2a^3(194A + 209B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \right)$$

↓ 3250

$$\frac{1}{11} \left(\frac{2a^2(14A + 11B) \sin(c+dx) \sqrt{a \cos(c+dx) + a}}{9d \cos^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \left(\frac{2a^3(194A + 209B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} + \frac{3}{7} a^2 (710A + 803B) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}}{\sin(c+dx + \frac{\pi}{2})^{7/2}} dx \right) + \frac{2aA \sin(c+dx)(a \cos(c+dx) + a)^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)} \right)$$

3.190. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$

input `Int[((a + a*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]))/cos[c + d*x]^(13/2), x]`

output `(2*a*A*(a + a*cos[c + d*x])^(3/2)*sin[c + d*x])/(11*d*cos[c + d*x]^(11/2)) + ((2*a^2*(14*A + 11*B)*sqrt[a + a*cos[c + d*x]]*sin[c + d*x])/(9*d*cos[c + d*x]^(9/2))) + ((2*a^3*(194*A + 209*B)*sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)*sqrt[a + a*cos[c + d*x]])) + (3*a^2*(710*A + 803*B)*((2*a*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)*sqrt[a + a*cos[c + d*x]])) + (4*((2*a*sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)*sqrt[a + a*cos[c + d*x]])) + (4*a*sin[c + d*x])/(3*d*sqrt[cos[c + d*x]]*sqrt[a + a*cos[c + d*x]])))/5)/7)/9)/11`

3.190.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*sqrt[a + b*sin[e + f*x]]*sqrt[c + d*sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*sqrt[a + b*sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*sin[e + f*x]]*(c + d*sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.190.4 Maple [A] (verified)

Time = 7.42 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.56

method	result
default	$\frac{2a^2 \sin(dx+c)(5680A(\cos^5(dx+c))+6424B(\cos^5(dx+c))+2840A(\cos^4(dx+c))+3212B(\cos^4(dx+c))+2130A(\cos^3(dx+c))+2409B(\cos^3(dx+c))+3465d(1+\cos(dx+c))c}{693d(1+\cos(dx+c)) \cos(dx+c)^{\frac{11}{2}}}$
parts	$\frac{2A \sin(dx+c)(1136(\cos^5(dx+c))+568(\cos^4(dx+c))+426(\cos^3(dx+c))+355(\cos^2(dx+c))+224 \cos(dx+c)+63) \sqrt{a(1+\cos(dx+c))} a}{693d(1+\cos(dx+c)) \cos(dx+c)^{\frac{11}{2}}}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x,method=_RE
TURNVERBOSE)
```

```
output 2/3465*a^2/d*sin(d*x+c)*(5680*A*cos(d*x+c)^5+6424*B*cos(d*x+c)^5+2840*A*co
s(d*x+c)^4+3212*B*cos(d*x+c)^4+2130*A*cos(d*x+c)^3+2409*B*cos(d*x+c)^3+177
5*A*cos(d*x+c)^2+1430*B*cos(d*x+c)^2+1120*A*cos(d*x+c)+385*B*cos(d*x+c)+31
5*A)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(11/2)
```

$$3.190. \int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$$

3.190.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.57

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \frac{2(8(710A + 803B)a^2 \cos(dx + c)^5 + 4(710A + 803B) \dots}{\dots}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x, algorithm="fricas")`

output `2/3465*(8*(710*A + 803*B)*a^2*cos(d*x + c)^5 + 4*(710*A + 803*B)*a^2*cos(d*x + c)^4 + 3*(710*A + 803*B)*a^2*cos(d*x + c)^3 + 5*(355*A + 286*B)*a^2*cos(d*x + c)^2 + 35*(32*A + 11*B)*a^2*cos(d*x + c) + 315*A*a^2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^7 + d*cos(d*x + c)^6)`

3.190.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(13/2),x)`

output `Timed out`

3.190.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 626 vs. 2(239) = 478.

Time = 0.36 (sec) , antiderivative size = 626, normalized size of antiderivative = 2.28

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \frac{8 \left(\frac{11 \left(\frac{315 \sqrt{2} a^2 \sin(dx+c)}{\cos(dx+c)+1} - \frac{945 \sqrt{2} a^2 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{1449 \sqrt{2} a^2 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{12 \dots}{\dots} \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)} \right)}{\dots}$$

3.190. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{13/2}(c+dx)} dx$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x, algorithm="maxima")`

output `8/3465*(11*(315*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 945*sqrt(2)*a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 1449*sqrt(2)*a^(5/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 1287*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 572*sqrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 104*sqrt(2)*a^(5/2)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11)*B*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^3/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(11/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(11/2)*(3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1)) + 5*(693*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 2310*sqrt(2)*a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 4620*sqrt(2)*a^(5/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 5478*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 3575*sqrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 1300*sqrt(2)*a^(5/2)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11 + 200*sqrt(2)*a^(5/2)*sin(d*x + c)^13/(cos(d*x + c) + 1)^13)*A*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^4/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(13/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(13/2)*(4*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 6*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 4*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + sin(d*x + c)^8/(cos(d*x + c) + 1)^8 + 1)))/d`

3.190.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x, algorithm="giac")`

output Timed out

3.190.9 Mupad [B] (verification not implemented)

Time = 7.57 (sec) , antiderivative size = 773, normalized size of antiderivative = 2.81

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \text{Too large to display}$$

```
input int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/cos(c + d*x)^(13/2),
x)
```

```
output ((a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a^2*(710*
A + 803*B)*16i)/(3465*d) - (a^2*exp(c*5i + d*x*5i)*(30*A + 41*B)*8i)/(15*d
) + (a^2*exp(c*6i + d*x*6i)*(30*A + 41*B)*8i)/(15*d) + (a^2*exp(c*4i + d*x
*4i)*(160*A + 157*B)*8i)/(35*d) - (a^2*exp(c*7i + d*x*7i)*(160*A + 157*B)*
8i)/(35*d) + (a^2*exp(c*2i + d*x*2i)*(710*A + 803*B)*8i)/(315*d) - (a^2*ex
p(c*9i + d*x*9i)*(710*A + 803*B)*8i)/(315*d) - (a^2*exp(c*11i + d*x*11i)*(
710*A + 803*B)*16i)/(3465*d) - (B*a^2*exp(c*3i + d*x*3i)*8i)/(3*d) + (B*a^
2*exp(c*8i + d*x*8i)*8i)/(3*d)))/((exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x
*1i)/2)^(1/2) + exp(c*1i + d*x*1i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*
x*1i)/2)^(1/2) + 5*exp(c*2i + d*x*2i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1i +
d*x*1i)/2)^(1/2) + 5*exp(c*3i + d*x*3i)*(exp(- c*1i - d*x*1i)/2 + exp(c*1
i + d*x*1i)/2)^(1/2) + 10*exp(c*4i + d*x*4i)*(exp(- c*1i - d*x*1i)/2 + exp
(c*1i + d*x*1i)/2)^(1/2) + 10*exp(c*5i + d*x*5i)*(exp(- c*1i - d*x*1i)/2 +
exp(c*1i + d*x*1i)/2)^(1/2) + 10*exp(c*6i + d*x*6i)*(exp(- c*1i - d*x*1i)
/2 + exp(c*1i + d*x*1i)/2)^(1/2) + 10*exp(c*7i + d*x*7i)*(exp(- c*1i - d*x
*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + 5*exp(c*8i + d*x*8i)*(exp(- c*1i -
d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + 5*exp(c*9i + d*x*9i)*(exp(- c*1i
- d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + exp(c*10i + d*x*10i)*(exp(- c
*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2) + exp(c*11i + d*x*11i)*(exp(
- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2)^(1/2))
```

$$3.191 \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$$

3.191.1 Optimal result	1813
3.191.2 Mathematica [A] (verified)	1814
3.191.3 Rubi [A] (verified)	1814
3.191.4 Maple [A] (verified)	1818
3.191.5 Fricas [A] (verification not implemented)	1819
3.191.6 Sympy [F(-1)]	1819
3.191.7 Maxima [F]	1820
3.191.8 Giac [F(-1)]	1820
3.191.9 Mupad [F(-1)]	1820

3.191.1 Optimal result

Integrand size = 35, antiderivative size = 190

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx = -\frac{(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{4\sqrt{ad}} + \frac{\sqrt{2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{(4A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{4d\sqrt{a+a \cos(c+dx)}} + \frac{B \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2d\sqrt{a+a \cos(c+dx)}}$$

output
$$\begin{aligned} & -1/4*(4*A-7*B)*\arcsin(\sin(d*x+c)*a^{(1/2)}/(a+a*\cos(d*x+c))^{(1/2)})/d/a^{(1/2)} \\ & + (A-B)*\arctan(1/2*\sin(d*x+c)*a^{(1/2)}*2^{(1/2)}/\cos(d*x+c)^{(1/2)}/(a+a*\cos(d*x+c))^{(1/2)})*2^{(1/2)}/d/a^{(1/2)} + 1/2*B*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)} \\ & + 1/4*(4*A-B)*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(a+a*\cos(d*x+c))^{(1/2)} \end{aligned}$$

$$3.191. \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$$

3.191.2 Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.20

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{\left((4A-B) \arcsin\left(\sqrt{1-\cos(c+dx)}\right) + 8(A-B) \arcsin\left(\sqrt{\cos(c+dx)}\right) - 4\sqrt{2}A \arctan\left(\frac{\sqrt{\cos(c+dx)}}{\sqrt{\sin^2\left(\frac{1}{2}(c+dx)\right)}}\right) \right)}{1}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `((4*A - B)*ArcSin[Sqrt[1 - Cos[c + d*x]]] + 8*(A - B)*ArcSin[Sqrt[Cos[c + d*x]]] - 4*Sqrt[2]*A*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]] + 4*Sqrt[2]*B*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]] + 2*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) + 4*A*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])] - B*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])]*Sin[c + d*x])/(4*d*Sqrt[1 - Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])])]`

3.191.3 Rubi [A] (verified)Time = 1.18 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3462, 27, 3042, 3462, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}(A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\sqrt{a\sin\left(c+dx+\frac{\pi}{2}\right)+a}} dx$$

$$\downarrow \text{3462}$$

$$\frac{\int \frac{\sqrt{\cos(c+dx)}(3aB+a(4A-B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a} + \frac{B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}$$

3.191. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3aB+a(4A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 3042 \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3aB+a(4A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 3462 \\
& \frac{\int \frac{a^2(4A-B)-a^2(4A-7B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 27 \\
& \frac{\int \frac{a^2(4A-B)-a^2(4A-7B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 3042 \\
& \frac{\int \frac{a^2(4A-B)-a^2(4A-7B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 3461 \\
& \frac{8a^2(A-B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - a(4A-7B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{4a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \\
& \quad \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 3042 \\
& \frac{8a^2(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - a(4A-7B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \\
& \quad \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}} \\
& \downarrow 3253
\end{aligned}$$

3.191. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

$$\frac{8a^2(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})^{a+a}}} dx + \frac{2a(4A-7B) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{2a} + \frac{a(4A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}}{2a} + \frac{4a}{2d\sqrt{a \cos(c+dx)+a}} \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}$$

223

$$\frac{8a^2(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})^{a+a}}} dx - \frac{2a^{3/2}(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a(4A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}}{2a} + \frac{4a}{2d\sqrt{a \cos(c+dx)+a}} \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}$$

3261

$$\frac{16a^3(A-B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}}\right) - \frac{2a^{3/2}(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a(4A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}}{2a} + \frac{4a}{2d\sqrt{a \cos(c+dx)+a}} \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}$$

218

$$\frac{8\sqrt{2}a^{3/2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right) - \frac{2a^{3/2}(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a(4A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}}{2a} + \frac{4a}{2d\sqrt{a \cos(c+dx)+a}} \frac{B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(B*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (((-2*a^(3/2)*(4*A - 7*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d + (8*Sqrt[2]*a^(3/2)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/(2*a) + (a*(4*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a)`

3.191. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

3.191.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`


```
rule 3462 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.191.4 Maple [A] (verified)

Time = 16.23 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.33

method	result
default	$\frac{(2B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) - 4A\sqrt{2} \arcsin(\cot(dx+c) - \csc(dx+c)) - B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})}{4d}$
parts	$\frac{A \left(\sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - \sqrt{2} \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) - 2 \arcsin(\cot(dx+c) - \csc(dx+c)) \right) \left(\sqrt{\cos(dx+c)} \sqrt{a(1+\cos(dx+c))} \right)}{2d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a}$

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 1/4/d*(2*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*A*(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)-4*A*2^(1/2)*arcsin(cot(d*x+c)-cs
c(d*x+c))-B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*B*2^(1/2)*arcsi
n(cot(d*x+c)-csc(d*x+c))-4*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))
^(1/2))+7*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*(a*(1+co
s(d*x+c)))^(1/2)*cos(d*x+c)^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c
)))^(1/2)/a
```

3.191.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$$

3.191.5 Fricas [A] (verification not implemented)

Time = 2.44 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.97

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{(2B\cos(dx+c) + 4A - B)\sqrt{a\cos(dx+c) + a}\sqrt{\cos(dx+c)}\sin(dx+c) + ((4A - 7B)\cos(dx+c) + 4A - 7B)\sqrt{a}\arctan(\sqrt{a\cos(dx+c) + a}\sqrt{\cos(dx+c)}) - 4\sqrt{2}\sqrt{(A-B)a\cos(dx+c) + (A-B)a}\arctan(\sqrt{2}\sqrt{a\cos(dx+c) + a}\sqrt{\cos(dx+c)})}{4(ad\cos(dx+c) + a^2)}$$

```
input integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output 1/4*((2*B*cos(d*x + c) + 4*A - B)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) + ((4*A - 7*B)*cos(d*x + c) + 4*A - 7*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - 4*sqrt(2)*((A - B)*a*cos(d*x + c) + (A - B)*a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a)/(a*d*cos(d*x + c) + a*d)
```

3.191.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(1/2),x)
```

```
output Timed out
```

3.191.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{\sqrt{a\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/sqrt(a*cos(d*x + c) + a), x)`

3.191.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.191.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(1/2),x)`

3.191. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

3.192
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$$

3.192.1 Optimal result 1821
 3.192.2 Mathematica [A] (verified) 1822
 3.192.3 Rubi [A] (verified) 1822
 3.192.4 Maple [A] (verified) 1825
 3.192.5 Fricas [A] (verification not implemented) 1826
 3.192.6 Sympy [F] 1826
 3.192.7 Maxima [F(-2)] 1827
 3.192.8 Giac [F(-1)] 1827
 3.192.9 Mupad [F(-1)] 1827

3.192.1 Optimal result

Integrand size = 35, antiderivative size = 141

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx = \frac{(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} - \frac{\sqrt{2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{B \sqrt{\cos(c+dx)} \sin(c+dx)}{d \sqrt{a+a \cos(c+dx)}}$$

output `(2*A-B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d/a^(1/2)-(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+B*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.192.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.18

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{\left(B \arcsin\left(\sqrt{1-\cos(c+dx)}\right) - 2(A-B) \arcsin\left(\sqrt{\cos(c+dx)}\right) + \sqrt{2}A \arctan\left(\frac{\sqrt{\cos(c+dx)}}{\sqrt{\sin^2\left(\frac{1}{2}(c+dx)\right)}}\right) - \sqrt{2}B \arctan\left(\frac{\sqrt{\cos(c+dx)}}{\sqrt{\sin^2\left(\frac{1}{2}(c+dx)\right)}}\right) \right)}{d\sqrt{1-\cos(c+dx)}\sqrt{a(1+\cos(c+dx))}}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `((B*ArcSin[Sqrt[1 - Cos[c + d*x]]] - 2*(A - B)*ArcSin[Sqrt[Cos[c + d*x]]] + Sqrt[2]*A*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]] - Sqrt[2]*B*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]] + B*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])])*Sin[c + d*x])/(d*Sqrt[1 - Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])])`

3.192.3 Rubi [A] (verified)Time = 0.81 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3462, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}(A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\sqrt{a\sin\left(c+dx+\frac{\pi}{2}\right)+a}} dx$$

$$\downarrow \text{3462}$$

$$\frac{\int \frac{aB+a(2A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}$$

3.192. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{aB+a(2A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{27} \\
& \int \frac{aB+a(2A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{(2A-B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx - 2a(A-B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a} + \\
& \quad \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3042} \\
& \frac{(2A-B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 2a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{2a} + \\
& \quad \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3253} \\
& \frac{-2a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2(2A-B) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}}{2a} + \\
& \quad \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{223} \\
& \frac{2\sqrt{a}(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - 2a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \\
& \quad \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} \\
& \quad \downarrow \text{3261}
\end{aligned}$$

3.192. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$

$$\frac{4a^2(A-B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2\sqrt{a}(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} +$$

$$\frac{2a}{d\sqrt{a \cos(c+dx)+a}} \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}$$

↓ 218

$$\frac{2\sqrt{a}(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{2\sqrt{2}\sqrt{a}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{d} +$$

$$\frac{2a}{d\sqrt{a \cos(c+dx)+a}} \frac{B \sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[a + a*Cos[c + d*x]],x]`

output `((2*Sqrt[a]*(2*A - B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (2*Sqrt[2]*Sqrt[a]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d)/(2*a) + (B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])`

3.192.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] :> Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] :> Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] :> Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3462 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m + n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.192.4 Maple [A] (verified)

Time = 16.63 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.33

method	result
default	$\frac{(A\sqrt{2} \arcsin(\cot(dx+c)) - \csc(dx+c)) - B\sqrt{2} \arcsin(\cot(dx+c)) + B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 2A \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a}$
parts	$\frac{A(\sqrt{\cos(dx+c)} \sqrt{a(1+\cos(dx+c))}) \left(\sqrt{2} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + \arcsin(\cot(dx+c) - \csc(dx+c))\right) \sqrt{2}}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a} + \frac{B(\sin(dx+c) \sqrt{a(1+\cos(dx+c))})}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a}$

3.192.
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+a \cos(c+dx)}} dx$$

input `int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*(A^2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))-B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/a`

3.192.5 Fracas [A] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.19

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

$$= \frac{\sqrt{a\cos(dx+c)+a}B\sqrt{\cos(dx+c)}\sin(dx+c) - ((2A-B)\cos(dx+c)+2A-B)\sqrt{a}\arctan\left(\frac{\sqrt{a\cos(dx+c)+a}}{\sin(dx+c)}\right)}{ad\cos(dx+c)+ad}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x,algorithm="fricas")`

output `(sqrt(a*cos(d*x + c) + a)*B*sqrt(cos(d*x + c))*sin(d*x + c) - ((2*A - B)*cos(d*x + c) + 2*A - B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + sqrt(2)*((A - B)*a*cos(d*x + c) + (A - B)*a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a))/(a*d*cos(d*x + c) + a*d)`

3.192.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \int \frac{(A+B\cos(c+dx))\sqrt{\cos(c+dx)}}{\sqrt{a(\cos(c+dx)+1)}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(cos(c + d*x))/sqrt(a*(cos(c + d*x) + 1)), x)`

3.192. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$

3.192.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: sign: argument cannot be imaginary; found %i`

3.192.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.192.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+a\cos(c+dx)}} dx$$

input `int((cos(c+d*x)^(1/2)*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(1/2),x)`

output `int((cos(c+d*x)^(1/2)*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(1/2),x)`

3.193
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}} dx$$

3.193.1 Optimal result 1828
 3.193.2 Mathematica [A] (verified) 1828
 3.193.3 Rubi [A] (verified) 1829
 3.193.4 Maple [A] (verified) 1831
 3.193.5 Fricas [A] (verification not implemented) 1832
 3.193.6 Sympy [F] 1832
 3.193.7 Maxima [C] (verification not implemented) 1832
 3.193.8 Giac [F(-1)] 1833
 3.193.9 Mupad [F(-1)] 1834

3.193.1 Optimal result

Integrand size = 35, antiderivative size = 100

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}}$$

output `2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/d/a^(1/2)+(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)`

3.193.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.82

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}} dx = \frac{2\left(\sqrt{2}B \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + (A - B) \arctan\left(\frac{\sin\left(\frac{1}{2}(c + dx)\right)}{\sqrt{\cos(c + dx)}}\right)\right) \cos\left(\frac{1}{2}(c + dx)\right)}{d\sqrt{a(1 + \cos(c + dx))}}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]),x]`

3.193.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}} dx$$

```
output (2*(Sqrt[2]*B*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + (A - B)*ArcTan[Sin[(c + d
*x)/2]/Sqrt[Cos[c + d*x]])*Cos[(c + d*x)/2])/(d*Sqrt[a*(1 + Cos[c + d*x])
])
```

3.193.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx$$

↓ 3461

$$(A - B) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{\cos(c + dx) a + a}} dx + \frac{B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{a}$$

↓ 3042

$$(A - B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a}} dx + \frac{B \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{a}$$

↓ 3253

$$(A - B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a}} dx - \frac{2B \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{ad}$$

↓ 223

$$(A - B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right) a + a}} dx + \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}}$$

3.193. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned} & \downarrow \text{3261} \\ & \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} - \frac{2a(A-B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} dx \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}}\right)}{d} \\ & \downarrow \text{218} \\ & \frac{\sqrt{2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} + \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]),x]`

output `(2*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/(Sqrt[a]*d) + (Sqrt[2]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]])*Sqrt[a + a*Cos[c + d*x]])/(Sqrt[a]*d)`

3.193.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] :> Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] :> Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.193.4 Maple [A] (verified)

Time = 8.21 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.32

method	result
default	$-\frac{\sqrt{a(1+\cos(dx+c))}(\sqrt{\cos(dx+c)})(A\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c))-B\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c))-2B\arctan(\tan(dx+c)))}{d(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a}$
parts	$-\frac{A\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{a(1+\cos(dx+c))}\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{2}}{d\sqrt{\cos(dx+c)}a} + \frac{B(\sqrt{\cos(dx+c)})\sqrt{a(1+\cos(dx+c))}(\sqrt{2}\arctan(\tan(dx+c))))}{d(1+\cos(dx+c))}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/d*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^(1/2)*(A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-2*B*arctan(tan(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/a
```

3.193.5 Fricas [A] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.96

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} dx = \frac{\sqrt{2}(A - B)\sqrt{a} \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) + 2B\sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right)}{ad}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `-(sqrt(2)*(A - B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos
(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*B*sqrt(a)*arctan(sqrt(a*cos(d*x + c)
+ a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/a*d`

3.193.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a(\cos(c + dx) + 1)} \sqrt{\cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a*(cos(c + d*x) + 1))*sqrt(cos(c + d*x
))), x)`

3.193.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.97 (sec) , antiderivative size = 1221, normalized size of antiderivative = 12.21

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `(sqrt(2)*A*arctan2(((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sin(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sin(d*x + c))/abs(e^(I*d*x + I*c) + 1), ((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sqrt(a)*cos(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sqrt(a)*cos(d*x + c) - sqrt(a))/(sqrt(a)*abs(e^(I*d*x + I*c) + 1))/sqrt(a) - (sqrt(2)*sqrt(a)*arctan2(((abs(2*e^(I*d*x + I*c) + 2)^4 + 16*cos(d*x + c)^4 + 16*sin(d*x + c)^4 + 8*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(2*e^(I*d*x + I*c) + 2)^2 - 64*cos(d*x + c)^3 + 32*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 96*cos(d*x + c)^2 - 64*cos(d*x + c) + 16)^(1/4)*sin(1/2*arctan2(8*(cos(d*x + c) - 1)*sin(d*x + c)/abs(2*e^(I*d*x + I*c) + 2)^2, (abs(2*e^(...`

3.193.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `Timed out`

3.193.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(1/2)),x)`

3.194
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$$

3.194.1 Optimal result 1835
 3.194.2 Mathematica [C] (verified) 1835
 3.194.3 Rubi [A] (verified) 1836
 3.194.4 Maple [B] (verified) 1838
 3.194.5 Fricas [A] (verification not implemented) 1839
 3.194.6 Sympy [F] 1839
 3.194.7 Maxima [C] (verification not implemented) 1839
 3.194.8 Giac [F] 1840
 3.194.9 Mupad [F(-1)] 1841

3.194.1 Optimal result

Integrand size = 35, antiderivative size = 99

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = -\frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2} \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}}\right)}{\sqrt{ad}} + \frac{2A \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a + a \cos(c + dx)}}$$

output `-(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+2*A*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)`

3.194.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 2.05

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{1}{2}(c + dx)\right) \left(10B \cos(c + dx) - (A - B) \left(-\frac{5}{4}(1 + 4 \cos(c + dx) + \cos(2(c + dx)))\right)\right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]),x]`

output `(2*Cos[(c + d*x)/2]*Sin[(c + d*x)/2]*(10*B*Cos[c + d*x] - (A - B)*((-5*(1 + 4*Cos[c + d*x] + Cos[2*(c + d*x)])*Csc[(c + d*x)/2]^4*(1 - Cos[c + d*x] + ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)])*Cos[c + d*x]*Sqrt[2 - 2*Sec[c + d*x]]))/4 + (Hypergeometric2F1[2, 5/2, 7/2, -(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]*Sin[c + d*x]*Tan[c + d*x])/2))/(5*d*Cos[c + d*x]^(3/2)*Sqrt[a*(1 + Cos[c + d*x])])`

3.194.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx \\
 & \quad \downarrow \text{3463} \\
 & \frac{2 \int -\frac{a(A-B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2A \sin(c + dx)}{d\sqrt{\cos(c + dx)}\sqrt{a \cos(c + dx) + a}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2A \sin(c + dx)}{d\sqrt{\cos(c + dx)}\sqrt{a \cos(c + dx) + a}} - (A - B) \int \frac{1}{\sqrt{\cos(c + dx)}\sqrt{\cos(c + dx)a + a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2A \sin(c + dx)}{d\sqrt{\cos(c + dx)}\sqrt{a \cos(c + dx) + a}} - (A - B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}} dx \\
 & \quad \downarrow \text{3261}
 \end{aligned}$$

3.194. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+a \cos(c+dx)}} dx$

$$\frac{2a(A - B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2A \sin(c+dx)} + \frac{d}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx) + a}}$$

↓ 218

$$\frac{2A \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx) + a}} - \frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]),x]`

output `-((Sqrt[2]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]])*Sqrt[a + a*Cos[c + d*x]])]/(Sqrt[a]*d)) + (2*A*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])`

3.194.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.194.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. $2(84) = 168$.

Time = 7.98 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.01

method	result
parts	$\frac{A \left(\cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c)) + \sqrt{2} \sin(dx+c) + \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c)) \right) \sqrt{a(1+\cos(dx+c))}}{d(1+\cos(dx+c)) \sqrt{\cos(dx+c)} a}$
default	$\frac{\left(A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \cos(dx+c) \arcsin(\cot(dx+c)-\csc(dx+c)) \sqrt{2} - B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \cos(dx+c) \arcsin(\cot(dx+c)-\csc(dx+c)) \sqrt{2} + \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \sqrt{a(1+\cos(dx+c))}}{da(1+\cos(dx+c)) \sqrt{\cos(dx+c)} a}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output A/d*(cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*
x+c))+2^(1/2)*sin(d*x+c)+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+
c)-csc(d*x+c)))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/cos(d*x+c)^(1/2)*2
^(1/2)/a-B/d/cos(d*x+c)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(a*(1+cos(
d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)/a
```

3.194.5 Fricas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.44

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{2 \sqrt{a \cos(dx + c)} + aA \sqrt{\cos(dx + c)} \sin(dx + c) - \frac{\sqrt{2}((A-B)a \cos(dx+c)^2 + (A-B)a \cos(dx+c)) \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(dx+c)}}{2(\cos(dx+c))}\right)}{\sqrt{a}}}{ad \cos(dx + c)^2 + ad \cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `(2*sqrt(a*cos(d*x + c) + a)*A*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(2)*((
A - B)*a*cos(d*x + c)^2 + (A - B)*a*cos(d*x + c))*arctan(1/2*sqrt(2)*sqrt(
a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/((cos(d*x + c)^2 + cos
(d*x + c))*sqrt(a)))/sqrt(a))/(a*d*cos(d*x + c)^2 + a*d*cos(d*x + c))`

3.194.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a} (\cos(c + dx) + 1) \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a*(cos(c + d*x) + 1))*cos(c + d*x)**(3
/2)), x)`

3.194.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 1188, normalized size of antiderivative = 12.00

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

3.194. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `(sqrt(2)*B*arctan2(((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sin(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sin(d*x + c))/abs(e^(I*d*x + I*c) + 1), ((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sqrt(a)*cos(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sqrt(a)*cos(d*x + c) - sqrt(a))/(sqrt(a)*abs(e^(I*d*x + I*c) + 1))/sqrt(a) + (2*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - 2*(cos(d*x + c) - 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - sqrt(2)*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*arctan2(((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d...`

3.194.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{a \cos(dx + c) + a \cos(dx + c)^{\frac{3}{2}}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(a*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.194.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(1/2)),x)`

3.195
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$$

3.195.1 Optimal result 1842
 3.195.2 Mathematica [C] (warning: unable to verify) 1842
 3.195.3 Rubi [A] (verified) 1843
 3.195.4 Maple [B] (verified) 1846
 3.195.5 Fricas [A] (verification not implemented) 1847
 3.195.6 Sympy [F] 1847
 3.195.7 Maxima [C] (verification not implemented) 1848
 3.195.8 Giac [F] 1848
 3.195.9 Mupad [F(-1)] 1849

3.195.1 Optimal result

Integrand size = 35, antiderivative size = 142

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx = \frac{\sqrt{2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{\sqrt{ad}} + \frac{2A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} - \frac{2(A-3B) \sin(c+dx)}{3d \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}$$

output

```
(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+2/3*A*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)-2/3*(A-3*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)
```

3.195.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

3.195.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$$

Time = 6.67 (sec) , antiderivative size = 627, normalized size of antiderivative = 4.42

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \frac{4B \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{3d \sqrt{a(1 + \cos(c + dx))} \left(1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)\right)^{3/2}}$$

$$+ \frac{8B \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{3d \sqrt{a(1 + \cos(c + dx))} \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}}$$

$$2(A - B) \cot\left(\frac{c}{2} + \frac{dx}{2}\right) \csc^4\left(\frac{c}{2} + \frac{dx}{2}\right) \left(-12 \cos^4\left(\frac{1}{2}(c + dx)\right) {}_3F_2\left(2, 2, \frac{7}{2}; 1, \frac{9}{2}; -\frac{\sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \sin^8\left(\frac{c}{2} + \frac{dx}{2}\right) + \dots$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]),x]`

output `(4*B*Cos[c/2 + (d*x)/2]*Sin[c/2 + (d*x)/2])/(3*d*Sqrt[a*(1 + Cos[c + d*x])] * (1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (8*B*Cos[c/2 + (d*x)/2]*Sin[c/2 + (d*x)/2])/(3*d*Sqrt[a*(1 + Cos[c + d*x])] * Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (2*(A - B)*Cot[c/2 + (d*x)/2]*Csc[c/2 + (d*x)/2]^4*(-12*Cos[(c + d*x)/2]^4*HypergeometricPFQ[{2, 2, 7/2}, {1, 9/2}, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))] * Sin[c/2 + (d*x)/2]^8 - 12*Hypergeometric2F1[2, 7/2, 9/2, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))] * Sin[c/2 + (d*x)/2]^8*(4 - 7*Sin[c/2 + (d*x)/2]^2 + 3*Sin[c/2 + (d*x)/2]^4) + 7*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))] * (1 - 2*Sin[c/2 + (d*x)/2]^2)^3*(15 - 20*Sin[c/2 + (d*x)/2]^2 + 8*Sin[c/2 + (d*x)/2]^4)*((3 - 7*Sin[c/2 + (d*x)/2]^2)*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))] - 3*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]) * (1 - 2*Sin[c/2 + (d*x)/2]^2)))/(63*d*Sqrt[a*(1 + Cos[c + d*x])] * (1 - 2*Sin[c/2 + (d*x)/2]^2)^(7/2))`

3.195.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.06, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.195. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx \\
& \quad \downarrow \text{3463} \\
& \frac{2 \int -\frac{a(A-3B) - 2aA \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \\
& \quad \downarrow \text{27} \\
& \frac{2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{a(A-3B) - 2aA \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{3a} \\
& \quad \downarrow \text{3042} \\
& \frac{2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} - \frac{\int \frac{a(A-3B) - 2aA \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{3a} \\
& \quad \downarrow \text{3463} \\
& \frac{2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} - \frac{2 \int -\frac{3a^2(A-B)}{2 \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a(A-3B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx) + a}} \\
& \quad \downarrow \text{27} \\
& \frac{2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} - \frac{2a(A-3B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx) + a}} - \frac{3a(A-B) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx}{3a} \\
& \quad \downarrow \text{3042} \\
& \frac{2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} - \frac{2a(A-3B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx) + a}} - \frac{3a(A-B) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{3a} \\
& \quad \downarrow \text{3261}
\end{aligned}$$

3.195. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$

$$\frac{6a^2(A-B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2a(A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}$$

3a

↓ 218

$$\frac{2A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}\sqrt{a}(A-B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d}$$

3a

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]),x]`

output `(2*A*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*Sqrt[a]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a*(A - 3*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/ (3*a)`

3.195.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.195.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 270 vs. 2(119) = 238.

Time = 7.80 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.91

method	result
default	$-\frac{\left(3A\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\left(\cos^2(dx+c)\arcsin(\cot(dx+c)-\csc(dx+c))-3B\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\left(\cos^2(dx+c)\arcsin(\cot(dx+c)-\csc(dx+c))\right)\right)\right)}{3d(1+\cos(dx+c))\cos(dx+c)^{\frac{3}{2}}a}$
parts	$-\frac{A\left(3\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\left(\cos^2(dx+c)\arcsin(\cot(dx+c)-\csc(dx+c))+\sin(dx+c)\cos(dx+c)\sqrt{2}+3\cos(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))\right)\right)}{3d(1+\cos(dx+c))\cos(dx+c)^{\frac{3}{2}}a}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -1/3/d*(3*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(
cot(d*x+c)-csc(d*x+c))-3*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d
*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+3*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2
)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-3*B*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+2*A*sin(d*
x+c)*cos(d*x+c)-6*B*sin(d*x+c)*cos(d*x+c)-2*A*sin(d*x+c))*(a*(1+cos(d*x+c
)))^(1/2)/cos(d*x+c)^(3/2)/a/(1+cos(d*x+c))
```

3.195.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+a \cos(c+dx)}} dx$$

3.195.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.15

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{2((A - 3B) \cos(dx + c) - A) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c) - \frac{3\sqrt{2}((A-B)a \cos(dx+c)^3 + (A-B)a \cos(dx+c)^2)}{3(ad \cos(dx + c)^3 + ad \cos(dx + c)^2)}}{3(ad \cos(dx + c)^3 + ad \cos(dx + c)^2)}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="fricas")
```

```
output -1/3*(2*((A - 3*B)*cos(d*x + c) - A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x
+ c))*sin(d*x + c) - 3*sqrt(2)*((A - B)*a*cos(d*x + c)^3 + (A - B)*a*cos(
d*x + c)^2)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))
*sin(d*x + c)/((cos(d*x + c)^2 + cos(d*x + c))*sqrt(a))/sqrt(a))/(a*d*cos
(d*x + c)^3 + a*d*cos(d*x + c)^2)
```

3.195.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a(\cos(c + dx) + 1)} \cos^{\frac{5}{2}}(c + dx)} dx$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(1/2),x)
```

```
output Integral((A + B*cos(c + d*x))/(sqrt(a*(cos(c + d*x) + 1))*cos(c + d*x)**(5
/2)), x)
```

3.195.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.94 (sec) , antiderivative size = 1485, normalized size of antiderivative = 10.46

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `1/3*(3*(2*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x
+ c) - 2*(cos(d*x + c) - 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c) + 1)) - sqrt(2)*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d
*x + 2*c) + 1)^(1/4)*arctan2(((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4
+ sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) +
1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*c
os(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(
1/4)*sin(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c)
+ 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2
*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sin(d*x + c))/abs(e^(I*d
*x + I*c) + 1), ((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x +
c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d
*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) +
1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sqrt(a)*
cos(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)
^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(
d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sqrt(a)*cos(d*x + c) - sqrt(a
))/sqrt(a)*abs(e^(I*d*x + I*c) + 1))) * B / ((cos(2*d*x + 2*c)^2 + sin(2*d*x
+ 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sqrt(a)) + (3*(sqrt(2)*cos(2*d*x
+ 2*c)^2 + sqrt(2)*sin(2*d*x + 2*c)^2 + 2*sqrt(2)*cos(2*d*x + 2*c) + s...`

3.195.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{a \cos(dx + c) + a \cos(dx + c)}^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

3.195. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$

output `integrate((B*cos(d*x + c) + A)/(sqrt(a*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.195.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{5}{2}} \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(1/2)), x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(1/2)), x)`

3.196
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+a \cos(c+dx)}} dx$$

3.196.1 Optimal result 1850
 3.196.2 Mathematica [C] (warning: unable to verify) 1851
 3.196.3 Rubi [A] (verified) 1851
 3.196.4 Maple [A] (verified) 1855
 3.196.5 Fricas [A] (verification not implemented) 1855
 3.196.6 Sympy [F(-1)] 1856
 3.196.7 Maxima [C] (verification not implemented) 1856
 3.196.8 Giac [F] 1857
 3.196.9 Mupad [F(-1)] 1858

3.196.1 Optimal result

Integrand size = 35, antiderivative size = 187

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx)\sqrt{a + a \cos(c + dx)}} dx = -\frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{2}\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}\right)}{\sqrt{ad}} + \frac{2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)\sqrt{a + a \cos(c + dx)}} - \frac{2(A - 5B) \sin(c + dx)}{15d \cos^{\frac{3}{2}}(c + dx)\sqrt{a + a \cos(c + dx)}} + \frac{2(13A - 5B) \sin(c + dx)}{15d\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}$$

```
output -(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)/d/a^(1/2)+2/5*A*sin(d*x+c)/d/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)-2/15*(A-5*B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+2/15*(13*A-5*B)*sin(d*x+c)/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)
```

3.196.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.24 (sec) , antiderivative size = 1759, normalized size of antiderivative = 9.41

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]),x]`

output `(4*B*Cos[c/2 + (d*x)/2]*Sin[c/2 + (d*x)/2])/(5*d*Sqrt[a*(1 + Cos[c + d*x])]*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) + (16*B*Cos[c/2 + (d*x)/2]*Sin[c/2 + (d*x)/2])/(15*d*Sqrt[a*(1 + Cos[c + d*x])]*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (32*B*Cos[c/2 + (d*x)/2]*Sin[c/2 + (d*x)/2])/(15*d*Sqrt[a*(1 + Cos[c + d*x])]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) - (2*(A - B)*Cot[c/2 + (d*x)/2]*Csc[c/2 + (d*x)/2]^6*(4725*Sin[c/2 + (d*x)/2]^2 - 48825*Sin[c/2 + (d*x)/2]^4 + 210105*Sin[c/2 + (d*x)/2]^6 - 486630*Sin[c/2 + (d*x)/2]^8 + 655812*Sin[c/2 + (d*x)/2]^10 - 710*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^10 - 40*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 9/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^10 - 518760*Sin[c/2 + (d*x)/2]^12 + 1770*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12 + 226656*Sin[c/2 + (d*x)/2]^14 - 1500*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14 - 42048*Sin[c/2 + (d*x)/2]^16 + 440*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^16 + 4725*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)] - 56700*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Sin[c/2 + (d*x)/2]^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]`

3.196.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.10, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.196. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$

$$\begin{array}{c}
 \downarrow \text{3463} \\
 \frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \\
 \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2 \int -\frac{15a^3(A-B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a^2(13A-5B) \sin(c+dx)}{3a \sqrt{a \cos(c+dx)+a}} \\
 \hline
 5a \\
 \downarrow \text{27} \\
 \frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \\
 \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(13A-5B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} - 15a^2(A-B) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx \\
 \hline
 5a \\
 \downarrow \text{3042} \\
 \frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \\
 \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(13A-5B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} - 15a^2(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
 \hline
 5a \\
 \downarrow \text{3261} \\
 \frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \\
 \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{30a^3(A-B) \int \frac{1}{\sin(c+dx) \tan(c+dx)a^3 + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2a^2(13A-5B) \sin(c+dx)}{3a \sqrt{a \cos(c+dx)+a}} \\
 \hline
 5a \\
 \downarrow \text{218} \\
 \frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \\
 \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(13A-5B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} - \frac{15\sqrt{2}a^{3/2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right)}{d} \\
 \hline
 5a
 \end{array}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]),x]`

3.196. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} dx$

```
output (2*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2
*a*(A - 5*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]
]) - ((-15*Sqrt[2]*a^(3/2)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*
Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^2*(13*A - 5*B)*Sin
[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])/(3*a))/(5*a)
```

3.196.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e
_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Ssin[e + f*x])^m*(c + d*Ssin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.196.4 Maple [A] (verified)

Time = 7.20 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.65

method	result
default	$(15A(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c))-15B(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c)))$
parts	$A(15(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))+13\sqrt{2}(\cos^2(dx+c))\sin(dx+c)+15\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\sin(dx+c))+15d(1+\cos(dx+c))\cos(dx+c)^{\frac{5}{2}}a$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/15/d*(15*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-15*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+15*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))-15*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+26*A*sin(d*x+c)*cos(d*x+c)^2-10*B*sin(d*x+c)*cos(d*x+c)^2-2*A*sin(d*x+c)*cos(d*x+c)+10*B*sin(d*x+c)*cos(d*x+c)+6*A*sin(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)/a/(1+cos(d*x+c))/cos(d*x+c)^(5/2)
```

3.196.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.96

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{2((13A - 5B) \cos(dx + c)^2 - (A - 5B) \cos(dx + c) + 3A) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c) + 15(ad \cos(dx + c))^4 + ad \cos(dx + c)}{15(ad \cos(dx + c))^4 + ad \cos(dx + c)}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="fracas")
```

output `1/15*(2*((13*A - 5*B)*cos(d*x + c)^2 - (A - 5*B)*cos(d*x + c) + 3*A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - 15*sqrt(2)*((A - B)*a*cos(d*x + c)^4 + (A - B)*a*cos(d*x + c)^3)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/((cos(d*x + c)^2 + cos(d*x + c))*sqrt(a)))/sqrt(a))/(a*d*cos(d*x + c)^4 + a*d*cos(d*x + c)^3)`

3.196.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(7/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.196.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.97 (sec) , antiderivative size = 1826, normalized size of antiderivative = 9.76

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2),x, algo rithm="maxima")`

output `1/15*(5*(3*(sqrt(2)*cos(2*d*x + 2*c)^2 + sqrt(2)*sin(2*d*x + 2*c)^2 + 2*sqrt(2)*cos(2*d*x + 2*c) + sqrt(2))*arctan2(((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sin(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sin(d*x + c))/abs(e^(I*d*x + I*c) + 1), ((abs(e^(I*d*x + I*c) + 1)^4 + cos(d*x + c)^4 + sin(d*x + c)^4 + 2*(cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)*abs(e^(I*d*x + I*c) + 1)^2 - 4*cos(d*x + c)^3 + 2*(cos(d*x + c)^2 - 2*cos(d*x + c) + 1)*sin(d*x + c)^2 + 6*cos(d*x + c)^2 - 4*cos(d*x + c) + 1)^(1/4)*sqrt(a)*cos(1/2*arctan2(2*(cos(d*x + c) - 1)*sin(d*x + c)/abs(e^(I*d*x + I*c) + 1)^2, (abs(e^(I*d*x + I*c) + 1)^2 + cos(d*x + c)^2 - sin(d*x + c)^2 - 2*cos(d*x + c) + 1)/abs(e^(I*d*x + I*c) + 1)^2)) + sqrt(a)*cos(d*x + c) - sqrt(a))/(sqrt(a)*abs(e^(I*d*x + I*c) + 1))) - 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(3/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (cos(d*x + c) - 3)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) - 4*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(3/2*ar...`

3.196.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{a \cos(dx + c) + a} \cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(a*cos(d*x + c) + a)*cos(d*x + c)^(7/2)), x)`

3.196.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + a \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{7/2} \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(7/2)*(a + a*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(7/2)*(a + a*cos(c + d*x))^(1/2)),x)`

3.197 $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$

3.197.1 Optimal result 1859
 3.197.2 Mathematica [A] (verified) 1860
 3.197.3 Rubi [A] (verified) 1860
 3.197.4 Maple [B] (verified) 1864
 3.197.5 Fricas [A] (verification not implemented) 1865
 3.197.6 Sympy [F(-1)] 1866
 3.197.7 Maxima [F] 1866
 3.197.8 Giac [F(-1)] 1866
 3.197.9 Mupad [F(-1)] 1867

3.197.1 Optimal result

Integrand size = 35, antiderivative size = 197

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx = \frac{(2A-3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{\frac{3}{2}}d} - \frac{(5A-9B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{\frac{3}{2}}d} + \frac{(A-B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2d(a+a \cos(c+dx))^{\frac{3}{2}}} - \frac{(A-3B) \sqrt{\cos(c+dx)} \sin(c+dx)}{2ad\sqrt{a+a \cos(c+dx)}}$$

output

```
(2*A-3*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d+1/2*(A-B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)-1/4*(5*A-9*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)-1/2*(A-3*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.197.2 Mathematica [A] (verified)

Time = 0.88 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.29

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \frac{\left(-4(A-3B)\arcsin\left(\sqrt{1-\cos(c+dx)}\right)\cos^2\left(\frac{1}{2}(c+dx)\right) - 20A\arcsin\left(\sqrt{\cos(c+dx)}\right)\cos\left(\frac{c+dx}{2}\right)^2 + 36B\arcsin\left(\sqrt{\cos(c+dx)}\right)\cos\left(\frac{c+dx}{2}\right)^2 + 2\sqrt{2}\left(5A-9B\right)\arctan\left(\sqrt{\cos(c+dx)}\right)/\sqrt{\sin\left(\frac{c+dx}{2}\right)^2}\cos\left(\frac{c+dx}{2}\right)^2 + 4B\sqrt{1-\cos(c+dx)}\cos\left(\frac{c+dx}{2}\right)^2 - 2A\sqrt{-((-1+\cos(c+dx))\cos(c+dx))} + 6B\sqrt{-((-1+\cos(c+dx))\cos(c+dx))}\sin(c+dx)\right)}{4d\sqrt{1-\cos(c+dx)}(a(1+\cos(c+dx)))^{3/2}}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output `((-4*(A - 3*B)*ArcSin[Sqrt[1 - Cos[c + d*x]]]*Cos[(c + d*x)/2]^2 - 20*A*ArcSin[Sqrt[Cos[c + d*x]]]*Cos[(c + d*x)/2]^2 + 36*B*ArcSin[Sqrt[Cos[c + d*x]]]*Cos[(c + d*x)/2]^2 + 2*Sqrt[2]*(5*A - 9*B)*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]]*Cos[(c + d*x)/2]^2 + 4*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) - 2*A*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])] + 6*B*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])]*Sin[c + d*x])/(4*d*Sqrt[1 - Cos[c + d*x]]*(a*(1 + Cos[c + d*x]))^(3/2))`

3.197.3 Rubi [A] (verified)Time = 1.18 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3456, 27, 3042, 3462, 25, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx$$

↓ 3456

$$\frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)-2a(A-3B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}a+a} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}}$$

↓ 27

3.197. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)-2a(A-3B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)-2a(A-3B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3462} \\
& \frac{\int -\frac{a^2(A-3B)-2a^2(2A-3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{25} \\
& -\frac{\int \frac{a^2(A-3B)-2a^2(2A-3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& -\frac{\int \frac{a^2(A-3B)-2a^2(2A-3B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3461} \\
& -\frac{a^2(5A-9B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - 2a(2A-3B)\int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{4a^2} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& -\frac{a^2(5A-9B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - 2a(2A-3B)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a^2} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \\
& \quad \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3253}
\end{aligned}$$

3.197. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{a^2(5A-9B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{4a(2A-3B) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{a} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}}{\frac{4a^2}{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}} \\
 & \quad \downarrow \text{223} \\
 & \frac{a^2(5A-9B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^{3/2}(2A-3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}}{\frac{4a^2}{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}} + \\
 & \quad \downarrow \text{3261} \\
 & \frac{2a^3(5A-9B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx) \sqrt{\cos(c+dx)a+a}}}\right) - \frac{4a^{3/2}(2A-3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}}{\frac{4a^2}{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}} \\
 & \quad \downarrow \text{218} \\
 & \frac{\sqrt{2}a^{3/2}(5A-9B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx) \sqrt{a \cos(c+dx)+a}}}\right) - \frac{4a^{3/2}(2A-3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}}{\frac{4a^2}{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)} - \frac{2a(A-3B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}}} +
 \end{aligned}$$

```
input Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2), x
]
```

3.197. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

```
output ((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2))
+ (-((( -4*a^(3/2)*(2*A - 3*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])/d
+ (Sqrt[2]*a^(3/2)*(5*A - 9*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/a) - (2*a*(A - 3*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x]/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2)
```

3.197.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3253 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3462 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.197.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 380 vs. 2(166) = 332.

Time = 16.22 (sec) , antiderivative size = 381, normalized size of antiderivative = 1.93

method	result
default	$\frac{(5A\sqrt{2} \arcsin(\cot(dx+c)) - \csc(dx+c)) \cos(dx+c) - 9B\sqrt{2} \arcsin(\cot(dx+c)) - \csc(dx+c)) \cos(dx+c) + 4B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}{4d(1+\cos(dx+c))^2 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$-\frac{A \left(\sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 4\sqrt{2} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \cos(dx+c) - 4\sqrt{2} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) - 5A \sqrt{2} \arcsin(\cot(dx+c)) - \csc(dx+c) \right) \cos(dx+c) + 4B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}{4d(1+\cos(dx+c))^2 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

$$3.197. \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{3}{2}}} dx$$

input `int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVERBOSE)`

output `1/4/a^2/d*(5*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)-9*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+4*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+5*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+8*A*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))-2*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)-9*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-12*B*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))+6*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+8*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))-12*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))*a*(1+cos(d*x+c))^(1/2)*cos(d*x+c)^(1/2)/(1+cos(d*x+c))^2/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)`

3.197.5 Fracas [A] (verification not implemented)

Time = 4.05 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.20

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{3}{2}}} dx = \frac{\sqrt{2}((5A-9B)\cos(dx+c)^2 + 2(5A-9B)\cos(dx+c) + 5A - 9B)\sqrt{a}\arctan(\sqrt{2}\sqrt{a\cos(dx+c)+a})\sqrt{\cos(dx+c)}}{(a+a\cos(c+dx))^{\frac{3}{2}}} + \frac{2(2B\cos(dx+c) - A + 3B)\sqrt{a\cos(dx+c)+a}\sqrt{\cos(dx+c)}\sin(dx+c) - 4((2A-3B)\cos(dx+c)^2 + 2(2A-3B)\cos(dx+c) + 2A-3B)\sqrt{a}\arctan(\sqrt{a\cos(dx+c)+a})\sqrt{\cos(dx+c)}}{(a+a\cos(c+dx))^{\frac{3}{2}}}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x,algorithm="fracas")`

output `1/4*(sqrt(2)*((5*A - 9*B)*cos(d*x + c)^2 + 2*(5*A - 9*B)*cos(d*x + c) + 5*A - 9*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c)) + 2*(2*B*cos(d*x + c) - A + 3*B)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) - 4*((2*A - 3*B)*cos(d*x + c)^2 + 2*(2*A - 3*B)*cos(d*x + c) + 2*A - 3*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c)))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.197. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{3}{2}}} dx$

3.197.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.197.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{3}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(a\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(3/
2), x)`

3.197.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.197.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)`

$$3.198 \quad \int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$$

3.198.1 Optimal result	1868
3.198.2 Mathematica [A] (verified)	1868
3.198.3 Rubi [A] (verified)	1869
3.198.4 Maple [B] (verified)	1872
3.198.5 Fracas [A] (verification not implemented)	1873
3.198.6 Sympy [F]	1873
3.198.7 Maxima [F]	1874
3.198.8 Giac [F(-1)]	1874
3.198.9 Mupad [F(-1)]	1874

3.198.1 Optimal result

Integrand size = 35, antiderivative size = 145

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{3/2}d} + \frac{(A-5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A-B) \sqrt{\cos(c+dx)} \sin(c+dx)}{2d(a+a \cos(c+dx))^{3/2}}$$

output

```
2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d+1/4*(A-5*B)
)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(
1/2))/a^(3/2)/d*2^(1/2)+1/2*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(
d*x+c))^(3/2)
```

3.198.2 Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.18

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx = \frac{\left(B \arcsin\left(\sqrt{1-\cos(c+dx)}\right) (1+\cos(c+dx)) + 5B \arcsin\left(\sqrt{\cos(c+dx)}\right) (1+\cos(c+dx)) + \sqrt{2}\left(\right) \right)}{2d\sqrt{1-\cos(c+dx)}}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output `-1/2*((B*ArcSin[Sqrt[1 - Cos[c + d*x]]]*(1 + Cos[c + d*x]) + 5*B*ArcSin[Sqrt[Cos[c + d*x]]]*(1 + Cos[c + d*x]) + Sqrt[2]*((A - 5*B)*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]]*Cos[(c + d*x)/2]^2 + (-A + B)*Sqrt[Cos[c + d*x]*Sin[(c + d*x)/2]^2]))*Sin[c + d*x])/(d*Sqrt[1 - Cos[c + d*x]]*(a*(1 + Cos[c + d*x]))^(3/2))`

3.198.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3456, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\
 & \quad \downarrow \text{3456} \\
 & \frac{\int \frac{a(A-B)+4aB\cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(A-B)+4aB\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(A-B)+4aB\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \\
 & \quad \downarrow \text{3461}
 \end{aligned}$$

3.198. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{a(A-5B) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx + 4B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{4a^2} + \\
& \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a(A-5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + 4B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a^2} + \\
& \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{3253} \\
& \frac{a(A-5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{8B \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{4a^2}}{+} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{223} \\
& \frac{a(A-5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{8\sqrt{a}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} + \\
& \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{3261} \\
& \frac{8\sqrt{a}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right) - \frac{2a^2(A-5B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} \right)}{4a^2}}{+} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}} \\
& \quad \downarrow \text{218} \\
& \frac{\sqrt{2}\sqrt{a}(A-5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right) + \frac{8\sqrt{a}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} + \\
& \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}}
\end{aligned}$$

3.198. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{3/2}} dx$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(3/2),x]`

output `((8*Sqrt[a]*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (Sqrt[2]*Sqrt[a]*(A - 5*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/(4*a^2) + ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2))`

3.198.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.198.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs. 2(120) = 240.

Time = 5.46 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.92

method	result
default	$-\frac{(A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)-5B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)-2A\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c)-\dots}{4d(1+\cos(dx+c))^2 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a^2}$
parts	$-\frac{A\left(-\sin(dx+c)\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c) + \arcsin(\cot(dx+c)-\csc(dx+c))\right) (\sqrt{\cos(dx+c)}) \sqrt{a}}{4d(1+\cos(dx+c))^2 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a^2}$

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2), x, method=_RET
URNVERBOSE)
```

output
$$\begin{aligned} & -1/4/a^2/d*(A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)-5*B*2^{(1/2)} \\ & * \arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)-2*A*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)} \\ & * \sin(d*x+c)+A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-8*B*\cos(d*x+c)*\ar \\ & \text{ctan}(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})+2*B*\sin(d*x+c)*(\cos(d*x \\ & +c)/(1+\cos(d*x+c)))^{(1/2)}-5*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-8*B*\ar \\ & \text{ctan}(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}))*(\text{a}*(1+\cos(d*x+c)))^{(1/2)} \\ & * \cos(d*x+c)^{(1/2)}/(1+\cos(d*x+c))^{(1/2)}/(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)} \end{aligned}$$

3.198.5 Fracas [A] (verification not implemented)

Time = 3.12 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.40

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \frac{\sqrt{2}((A-5B)\cos(dx+c)^2+2(A-5B)\cos(dx+c)+A-5B)\sqrt{a}\arctan\left(\frac{\sqrt{2}\sqrt{a\cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a}\sin(dx+c)}\right)}{4(a^2d)}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="fricas")`

output
$$\begin{aligned} & -1/4*(\text{sqrt}(2)*((A-5*B)*\cos(d*x+c)^2+2*(A-5*B)*\cos(d*x+c)+A-5 \\ & *B)*\text{sqrt}(a)*\arctan(\text{sqrt}(2)*\text{sqrt}(a*\cos(d*x+c)+a)*\text{sqrt}(\cos(d*x+c))/(\text{sq} \\ & \text{rt}(a)*\sin(d*x+c))) - 2*\text{sqrt}(a*\cos(d*x+c)+a)*(A-B)*\text{sqrt}(\cos(d*x+c) \\ &)*\sin(d*x+c) + 8*(B*\cos(d*x+c)^2+2*B*\cos(d*x+c)+B)*\text{sqrt}(a)*\arct \\ & \text{an}(\text{sqrt}(a*\cos(d*x+c)+a)*\text{sqrt}(\cos(d*x+c))/(\text{sqrt}(a)*\sin(d*x+c))))/(a \\ & ^2*d*\cos(d*x+c)^2+2*a^2*d*\cos(d*x+c)+a^2*d) \end{aligned}$$

3.198.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \int \frac{(A+B\cos(c+dx))\sqrt{\cos(c+dx)}}{(a(\cos(c+dx)+1))^{3/2}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A+B*cos(c+d*x))*sqrt(cos(c+d*x))/(a*(cos(c+d*x)+1))**(3/2),x)`

3.198.
$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$$

3.198.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(a\cos(dx+c)+a)^{3/2}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(a*cos(d*x + c) + a)^(3/2), x)`

3.198.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.198.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(3/2),x)`

3.199 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{3/2}} dx$

3.199.1 Optimal result	1875
3.199.2 Mathematica [A] (verified)	1875
3.199.3 Rubi [A] (verified)	1876
3.199.4 Maple [B] (verified)	1878
3.199.5 Fricas [A] (verification not implemented)	1879
3.199.6 Sympy [F]	1879
3.199.7 Maxima [F]	1879
3.199.8 Giac [F(-1)]	1880
3.199.9 Mupad [F(-1)]	1880

3.199.1 Optimal result

Integrand size = 35, antiderivative size = 107

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \frac{(3A + B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{(A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}}$$

output `1/4*(3*A+B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)-1/2*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(3/2)`

3.199.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.72

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)}\left(2B \arcsin\left(\frac{\sin\left(\frac{1}{2}(c+dx)\right)}{\sqrt{\cos^2\left(\frac{1}{2}(c+dx)\right)}}\right)\right) \cos^3\left(\frac{1}{2}(c + dx)\right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)),x]`

output $(\text{Sqrt}[\text{Cos}[(c + d*x)/2]^2]*(2*B*\text{ArcSin}[\text{Sin}[(c + d*x)/2]/\text{Sqrt}[\text{Cos}[(c + d*x)/2]^2]]*\text{Cos}[(c + d*x)/2]^3 - 3*\text{Sqrt}[2]*A*\text{ArcTanh}[\text{Sqrt}[-(\text{Sec}[c + d*x]*\text{Sin}[(c + d*x)/2]^2)]]*\text{Cos}[(c + d*x)/2]^2*\text{Sqrt}[-1 + \text{Cos}[c + d*x]]*\text{Cot}[(c + d*x)/2] + (-A + B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])]/(\text{Sqrt}[2]*d*\text{Sqrt}[1 + \text{Cos}[c + d*x]]*(a*(1 + \text{Cos}[c + d*x]))^(3/2))$

3.199.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2}} dx$$

↓ 3457

$$\frac{\int \frac{a(3A+B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 27

$$\frac{(3A+B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 3042

$$\frac{(3A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}}$$

↓ 3261

$$-\frac{(3A+B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2d} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}}$$

3.199. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{3/2}} dx$

$$\frac{(3A + B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{(A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{2d(a \cos(c + dx) + a)^{3/2}}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)),x]`

output `((3*A + B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) - ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2))`

3.199.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.199.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(88) = 176.

Time = 7.90 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.99

method	result
default	$-\frac{(3A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)+B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)+3A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)+4a^2 d \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})}{4d \sqrt{-\frac{(\csc^2(dx+c)(1-\cos(dx+c))^2-1}{(\csc^2(dx+c)(1-\cos(dx+c))^2+1)} a^2}}$
parts	$-\frac{A \sqrt{-(\csc^2(dx+c)(1-\cos(dx+c))^2+1)} \sqrt{\frac{a}{(\csc^2(dx+c)(1-\cos(dx+c))^2+1)}} \left(\sqrt{-(\csc^2(dx+c)(1-\cos(dx+c))^2+1} (\csc(dx+c)-\cot(dx+c)) \right)}{4d \sqrt{-\frac{(\csc^2(dx+c)(1-\cos(dx+c))^2-1}{(\csc^2(dx+c)(1-\cos(dx+c))^2+1)} a^2}}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/4/a^2/d*(3*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+B*2^(1/2)
*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+3*A*2^(1/2)*arcsin(cot(d*x+c)-cs
c(d*x+c))+2*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+B*2^(1/2)*arcsi
n(cot(d*x+c)-csc(d*x+c))-2*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))
*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2)/(1+cos(d*x+c))^2
```

3.199.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+a \cos(c+dx))^{3/2}}} dx$$

3.199.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.53

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2}((3A + B) \cos(dx + c)^2 + 2(3A + B) \cos(dx + c) + 3A + B) \arctan\left(\frac{1}{2}\sqrt{2}\sqrt{a \cos(dx + c) + a}\sqrt{\cos(dx + c)}\right) - 2\sqrt{a \cos(dx + c) + a}(A - B)\sqrt{\cos(dx + c)}\sin(dx + c)}{(a^2 d \cos(dx + c) + a^2 d)^{3/2}}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(3/2),x, algo rithm="fricas")`

output `1/4*(sqrt(2)*((3*A + B)*cos(d*x + c)^2 + 2*(3*A + B)*cos(d*x + c) + 3*A + B)*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) - 2*sqrt(a*cos(d*x + c) + a)*(A - B)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.199.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a(\cos(c + dx) + 1))^{3/2} \sqrt{\cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))/((a*(cos(c + d*x) + 1))**(3/2)*sqrt(cos(c + d*x))), x)`

3.199.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{3/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(3/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*sqrt(cos(d*x + c))), x)`

3.199.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.199.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(3/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(3/2)),
x)`

$$3.200 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$$

3.200.1 Optimal result 1881
 3.200.2 Mathematica [C] (warning: unable to verify) 1881
 3.200.3 Rubi [A] (verified) 1882
 3.200.4 Maple [B] (verified) 1885
 3.200.5 Fracas [A] (verification not implemented) 1886
 3.200.6 Sympy [F] 1886
 3.200.7 Maxima [F] 1887
 3.200.8 Giac [F] 1887
 3.200.9 Mupad [F(-1)] 1887

3.200.1 Optimal result

Integrand size = 35, antiderivative size = 156

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{(7A - 3B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d}$$

$$- \frac{(A - B) \sin(c + dx)}{2d\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}} + \frac{(5A - B) \sin(c + dx)}{2ad\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}$$

```
output -1/4*(7*A-3*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a
*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)-1/2*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+
c))^(3/2)/cos(d*x+c)^(1/2)+1/2*(5*A-B)*sin(d*x+c)/a/d/cos(d*x+c)^(1/2)/(a+
a*cos(d*x+c))^(1/2)
```

3.200.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.59 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.71

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \frac{\cos^3\left(\frac{1}{2}(c + dx)\right) \left(30(A - B) \arctan\left(\frac{1 - 2 \sin\left(\frac{1}{2}(c + dx)\right)}{\sqrt{\cos(c + dx)}}\right) - 30(A - B)\right)}{\dots}$$

3.200. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)),x]`

output `(Cos[(c + d*x)/2]^3*(30*(A - B)*ArcTan[(1 - 2*Sin[(c + d*x)/2])/Sqrt[Cos[c + d*x]]] - 30*(A - B)*ArcTan[(1 + 2*Sin[(c + d*x)/2])/Sqrt[Cos[c + d*x]]] - (20*(A - B)*Sqrt[Cos[c + d*x]]/(-1 + Sin[(c + d*x)/2]) - (20*(A - B)*Sqrt[Cos[c + d*x]]/(1 + Sin[(c + d*x)/2]) + (5*(A - B)*(-1 + 2*Sin[(c + d*x)/2]))/(Sqrt[Cos[c + d*x]]*(Cos[(c + d*x)/4] + Sin[(c + d*x)/4])^2) - (5*(A - B)*(1 + 2*Sin[(c + d*x)/2]))/(Sqrt[Cos[c + d*x]]*(-1 + Sin[(c + d*x)/2])) + ((A + 3*B)*Csc[(c + d*x)/2]^3*(5*(1 + 4*Cos[c + d*x] + Cos[2*(c + d*x)]))*(1 - Cos[c + d*x] + ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]*Cos[c + d*x]*Sqrt[2 - 2*Sec[c + d*x]]) - 2*Hypergeometric2F1[2, 5/2, 7/2, -(Sec[c + d*x]*Sin[(c + d*x)/2]^2)*Sin[(c + d*x)/2]^4*Sin[c + d*x]*Tan[c + d*x]))/(2*Cos[c + d*x]^(3/2)))/(10*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.200.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3457, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{3/2}} dx$$

↓ 3457

$$\frac{\int \frac{a(5A-B) - 2a(A-B) \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B) \sin(c+dx)}{2d \sqrt{\cos(c+dx)} (a \cos(c+dx) + a)^{3/2}}$$

↓ 27

$$\frac{\int \frac{a(5A-B) - 2a(A-B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d \sqrt{\cos(c+dx)} (a \cos(c+dx) + a)^{3/2}}$$

↓ 3042

3.200. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{a(5A-B) - 2a(A-B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B) \sin(c+dx)}{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3463} \\
 & \frac{2 \int -\frac{a^2(7A-3B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{2a(5A-B) \sin(c+dx)}{d \sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - \frac{(A-B) \sin(c+dx)}{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2a(5A-B) \sin(c+dx)}{d \sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - a(7A-3B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \\
 & \quad \frac{4a^2}{(A-B) \sin(c+dx)} \\
 & \quad \frac{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2a(5A-B) \sin(c+dx)}{d \sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - a(7A-3B) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}\sqrt{\sin(c+dx + \frac{\pi}{2})a+a}} dx - \\
 & \quad \frac{4a^2}{(A-B) \sin(c+dx)} \\
 & \quad \frac{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}} \\
 & \quad \downarrow \text{3261} \\
 & \frac{2a^2(7A-3B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3 + 2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2a(5A-B) \sin(c+dx)}{d \sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - \\
 & \quad \frac{4a^2}{(A-B) \sin(c+dx)} \\
 & \quad \frac{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}} \\
 & \quad \downarrow \text{218} \\
 & \frac{2a(5A-B) \sin(c+dx)}{d \sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - \frac{\sqrt{2}\sqrt{a}(7A-3B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \\
 & \quad \frac{4a^2}{(A-B) \sin(c+dx)} \\
 & \quad \frac{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)),x]`

3.200. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$

```
output -1/2*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^(3/2)) + (-(Sqrt[2]*Sqrt[a]*(7*A - 3*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c + d*x]])])/d) + (2*a*(5*A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c + d*x]])/(4*a^2)
```

3.200.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x], x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x])/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3457 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.200.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 352 vs. $2(131) = 262$.

Time = 7.91 (sec) , antiderivative size = 353, normalized size of antiderivative = 2.26

method	result
default	$\left(7A\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}(\cos^2(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))-3B\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}(\cos^2(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\right)$
parts	$\frac{A\left(7\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))+5\sin(dx+c)\cos(dx+c)\sqrt{2}+14\cos(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))\right)}{4d(1+\cos(dx+c))^2\sqrt{\cos(dx+c)}}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/4/a^2/d*(7*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcs
in(cot(d*x+c)-csc(d*x+c))-3*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*co
s(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+14*A*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-6*B*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+7*(cos
(d*x+c)/(1+cos(d*x+c)))^(1/2)*A*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-3*(c
os(d*x+c)/(1+cos(d*x+c)))^(1/2)*B*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+10
*A*sin(d*x+c)*cos(d*x+c)-2*B*sin(d*x+c)*cos(d*x+c)+8*A*sin(d*x+c))*(a*(1+c
os(d*x+c)))^(1/2)/(1+cos(d*x+c))^2/cos(d*x+c)^(1/2)
```

3.200.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.29

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{\sqrt{2}((7A - 3B) \cos(dx + c)^3 + 2(7A - 3B) \cos(dx + c)^2 + (7A - 3B) \cos(dx + c))\sqrt{a} \arctan\left(\frac{\sqrt{2}\sqrt{a}\cos(dx + c)}{2}\right)}{4(a^2d \cos(dx + c))^3 + 2a^2d \cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/4*(sqrt(2)*((7*A - 3*B)*cos(d*x + c)^3 + 2*(7*A - 3*B)*cos(d*x + c)^2 + (7*A - 3*B)*cos(d*x + c))*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) - 2*((5*A - B)*cos(d*x + c) + 4*A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^2*d*cos(d*x + c)^3 + 2*a^2*d*cos(d*x + c)^2 + a^2*d*cos(d*x + c))`

3.200.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a(\cos(c + dx) + 1))^{\frac{3}{2}} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))/((a*(cos(c + d*x) + 1))**(3/2)*cos(c + d*x)**(3/2)), x)`

3.200.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3
/2)), x)`

3.200.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3
/2)), x)`

3.200.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(3/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(3/2)),
x)`

3.200. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$

$$3.201 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$$

3.201.1 Optimal result	1888
3.201.2 Mathematica [C] (warning: unable to verify)	1889
3.201.3 Rubi [A] (verified)	1889
3.201.4 Maple [B] (verified)	1893
3.201.5 Fricas [A] (verification not implemented)	1894
3.201.6 Sympy [F(-1)]	1894
3.201.7 Maxima [F]	1894
3.201.8 Giac [F]	1895
3.201.9 Mupad [F(-1)]	1895

3.201.1 Optimal result

Integrand size = 35, antiderivative size = 203

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx = \frac{(11A-7B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{(A-B) \sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} + \frac{(7A-3B) \sin(c+dx)}{6ad \cos^{\frac{3}{2}}(c+dx) \sqrt{a+a \cos(c+dx)}} - \frac{(19A-15B) \sin(c+dx)}{6ad \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}$$

output

```
-1/2*(A-B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2)+1/4*(11*A-7*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(3/2)/d*2^(1/2)+1/6*(7*A-3*B)*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)-1/6*(19*A-15*B)*sin(d*x+c)/a/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)
```

3.201.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.58 (sec) , antiderivative size = 1054, normalized size of antiderivative = 5.19

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(3/2)),x]`

output `-1/6*((A - B)*Cos[c/2 + (d*x)/2]^3*(1 - 2*Sin[c/2 + (d*x)/2]))/(d*(a*(1 + Cos[c + d*x]))^(3/2)*(1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^(2/3)) + ((A - B)*Cos[c/2 + (d*x)/2]^3*(1 + 2*Sin[c/2 + (d*x)/2]))/(6*d*(a*(1 + Cos[c + d*x]))^(3/2)*(1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^(2/3)) - ((A - B)*Cos[c/2 + (d*x)/2]^3*(5*ArcTan[(1 - 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]] + (1 + Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (3*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])/(1 - Sin[c/2 + (d*x)/2]))/(d*(a*(1 + Cos[c + d*x]))^(3/2)) + ((A - B)*Cos[c/2 + (d*x)/2]^3*(5*ArcTan[(1 + 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]] + (1 - Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (3*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])/(1 + Sin[c/2 + (d*x)/2]))/(d*(a*(1 + Cos[c + d*x]))^(3/2)) + ((A + 3*B)*Cot[c/2 + (d*x)/2]^3*Csc[c/2 + (d*x)/2]^2*(-12*Cos[(c + d*x)/2]^4*HypergeometricPFQ[{2, 2, 7/2}, {1, 9/2}, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*Sin[c/2 + (d*x)/2]^8 - 12*Hypergeometric2F1[2, 7/2, 9/2, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^8*(4 - 7*Sin[c/2 + (d*x)/2]^2 + 3*Sin[c/2 + (d*x)/2]^4) + 7*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2)]*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*(15 - 20*Sin[c/2 + (d*x)/2]^2 + 8*Sin[c/2 + (d*x)/2]^4)*((3 - 7*Sin[c/2 + (d*x)/2]^2)*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*...`

3.201.3 Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.201. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2}} dx \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a(7A-3B)-4a(A-B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(7A-3B)-4a(A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(7A-3B)-4a(A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3463} \\
& \frac{2\int -\frac{a^2(19A-15B)-2a^2(7A-3B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \\
& \quad \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \quad \frac{4a^2}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(19A-15B)-2a^2(7A-3B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} - \\
& \quad \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \quad \frac{4a^2}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(19A-15B)-2a^2(7A-3B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{3a} - \\
& \quad \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \quad \frac{4a^2}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}
\end{aligned}$$

3.201. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow \text{3463} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{3a^3(11A-7B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \\
& \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \frac{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{27}} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3a^2(11A-7B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{3a} \\
& \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \frac{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{3042}} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3a^2(11A-7B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} \\
& \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \frac{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{3261}} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{6a^3(11A-7B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \\
& \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \frac{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}{\downarrow \text{218}} \\
& \frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}a^{3/2}(11A-7B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d} \\
& \frac{4a^2}{(A-B)\sin(c+dx)} \\
& \frac{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}}{\downarrow}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(3/2)), x]`

$$3.201. \int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))^{3/2}} dx$$

```
output -1/2*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a*(7*A - 3*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*a^(3/2)*(11*A - 7*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^2*(19*A - 15*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])/(3*a))/(4*a^2)
```

3.201.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3457 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.201.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 402 vs. 2(172) = 344.

Time = 8.15 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.99

method	result
default	$-\frac{\left(33A(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c))-21B(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c))\right)}{\dots}$
parts	$-\frac{A\left(33(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))+19\sqrt{2}(\cos^2(dx+c))\sin(dx+c)+66\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\right)}{\dots}$

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/12/a^2/d*(33*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-21*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+66*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))-42*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+33*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-21*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+38*A*sin(d*x+c)*cos(d*x+c)^2-30*B*sin(d*x+c)*cos(d*x+c)^2+24*A*sin(d*x+c)*cos(d*x+c)-24*B*sin(d*x+c)*cos(d*x+c)-8*A*sin(d*x+c)*(a*(1+cos(d*x+c)))^(1/2)/cos(d*x+c)^(3/2)/(1+cos(d*x+c))^2`

3.201.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$$

3.201.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.09

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \frac{3\sqrt{2}((11A - 7B) \cos(dx + c)^4 + 2(11A - 7B) \cos(dx + c)^3 - \dots}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="fricas")
```

```
output 1/12*(3*sqrt(2)*((11*A - 7*B)*cos(d*x + c)^4 + 2*(11*A - 7*B)*cos(d*x + c)
^3 + (11*A - 7*B)*cos(d*x + c)^2)*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*
x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*
cos(d*x + c))) - 2*((19*A - 15*B)*cos(d*x + c)^2 + 12*(A - B)*cos(d*x + c)
- 4*A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^2*d*c
os(d*x + c)^4 + 2*a^2*d*cos(d*x + c)^3 + a^2*d*cos(d*x + c)^2)
```

3.201.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(3/2),x)
```

```
output Timed out
```

3.201.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="maxima")
```

3.201. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{3/2}} dx$

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(5/2)), x)`

3.201.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(5/2)), x)`

3.201.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(3/2)),x)`

3.202 $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

3.202.1 Optimal result 1896
 3.202.2 Mathematica [A] (verified) 1897
 3.202.3 Rubi [A] (verified) 1897
 3.202.4 Maple [B] (verified) 1902
 3.202.5 Fricas [A] (verification not implemented) 1903
 3.202.6 Sympy [F(-1)] 1904
 3.202.7 Maxima [F] 1904
 3.202.8 Giac [F(-1)] 1904
 3.202.9 Mupad [F(-1)] 1905

3.202.1 Optimal result

Integrand size = 35, antiderivative size = 246

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{5/2}d} - \frac{(43A-115B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A-B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{(7A-15B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}} - \frac{(11A-35B) \sqrt{\cos(c+dx)} \sin(c+dx)}{16a^2d\sqrt{a+a \cos(c+dx)}}$$

output

```
(2*A-5*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d+1/4*(A-B)*cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+1/16*(7*A-15*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)-1/32*(43*A-115*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)-1/16*(11*A-35*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(1/2)
```

3.202.2 Mathematica [A] (verified)

Time = 2.19 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.26

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \frac{\left(-8(11A-35B)\arcsin\left(\sqrt{1-\cos(c+dx)}\right)\cos^4\left(\frac{1}{2}(c+dx)\right) - \dots}{\dots}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `((-8*(11*A - 35*B)*ArcSin[Sqrt[1 - Cos[c + d*x]]]*Cos[(c + d*x)/2]^4 - 344*A*ArcSin[Sqrt[Cos[c + d*x]]]*Cos[(c + d*x)/2]^4 + 920*B*ArcSin[Sqrt[Cos[c + d*x]]]*Cos[(c + d*x)/2]^4 + 4*Sqrt[2]*(43*A - 115*B)*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]]*Cos[(c + d*x)/2]^4 - 30*A*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) + 110*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) + 32*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(5/2) - 22*A*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])] + 70*B*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])])*Sin[c + d*x])/(32*d*Sqrt[1 - Cos[c + d*x]]*(a*(1 + Cos[c + d*x]))^(5/2))`

3.202.3 Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3462, 25, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{\frac{5}{2}}} dx$$

↓ 3042

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{5}{2}}(A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a\sin\left(c+dx+\frac{\pi}{2}\right)+a)^{\frac{5}{2}}} dx$$

↓ 3456

3.202. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx$

$$\begin{aligned}
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)-2a(A-5B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)-2a(A-5B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(5a(A-B)-2a(A-5B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 3456 \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a^2(7A-15B)-2a^2(11A-35B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow 27} \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a^2(7A-15B)-2a^2(11A-35B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow 3042} \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a^2(7A-15B)-2a^2(11A-35B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow 3462} \\
& \frac{\int -\frac{a^3(11A-35B)-16a^3(2A-5B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{4d(a\cos(c+dx)+a)^{5/2}}{\downarrow 25}
\end{aligned}$$

3.202. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{\int \frac{a^3(11A-35B)-16a^3(2A-5B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{a^3(11A-35B)-16a^3(2A-5B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3461

$$\frac{a^3(43A-115B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - 16a^2(2A-5B)\int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{a^3(43A-115B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - 16a^2(2A-5B)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} +$$

$$\frac{8a^2}{(A-B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)} \frac{1}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3253

$$\frac{a^3(43A-115B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{32a^2(2A-5B)\int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2}$$

↓ 223

3.202. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{a^3(43A-115B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})} a+a} dx - \frac{32a^{5/2}(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{2a^2(11A-35B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{a(7A-15B)}{2d} \\
 & \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{3261} \\
 & \frac{2a^4(43A-115B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx) a^3 + 2a^2}{\cos(c+dx) a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx) a+a}}\right) - \frac{32a^{5/2}(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{2a^2(11A-35B) \sin(c+dx)}{d\sqrt{a \cos(c+dx)+a}} \\
 & \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \\
 & \quad \downarrow \text{218} \\
 & \frac{\sqrt{2} a^{5/2} (43A-115B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right) - \frac{32a^{5/2}(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{2a^2(11A-35B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{a(7A-15B)}{2d} \\
 & \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `((A - B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + ((a*(7*A - 15*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2))) + (-(((-32*a^(5/2)*(2*A - 5*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (Sqrt[2]*a^(5/2)*(43*A - 115*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/a) - (2*a^2*(11*A - 35*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2))/(8*a^2)`

3.202. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

3.202.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]))]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.202.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$$

```
rule 3461 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3462 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m + n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.202.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 586 vs. 2(209) = 418.

Time = 17.80 (sec) , antiderivative size = 587, normalized size of antiderivative = 2.39

method	result
default	$(43A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))-115B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))+32B(\cos^2(dx+c)) \sin(dx+c)) \sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}}$
parts	$A \left(-\frac{(\csc^2(dx+c))(1-\cos(dx+c))^2-1}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \right)^{\frac{5}{2}} \left((\csc^2(dx+c))(1-\cos(dx+c))^2+1 \right)^3 \sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}} \left(-2(\csc^3(dx+c)) \sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}} \right)$

```
input int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2), x, method=_RETURNVERBOSE)
```

$$3.202. \int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$$

output $1/32/a^3/d*(43*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2-115*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2+32*B*\cos(d*x+c)^2*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+86*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)+64*A*\cos(d*x+c)^2*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})-30*A*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}-230*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)-160*B*\cos(d*x+c)^2*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})+110*B*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+43*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))+128*A*\cos(d*x+c)*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})-22*A*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\sin(d*x+c)-115*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-320*B*\cos(d*x+c)*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})+70*B*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+64*A*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})-160*B*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}))*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\cos(d*x+c)^{(1/2)}/(1+\cos(d*x+c))^{3/2}/(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}$

3.202.5 Fracas [A] (verification not implemented)

Time = 8.57 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.23

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \frac{\sqrt{2}((43A-115B)\cos(dx+c)^3+3(43A-115B)\cos(dx+c)^2)}{(a+a\cos(c+dx))^{5/2}}$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="fracas")`

output $1/32*(\sqrt{2}*((43*A-115*B)*\cos(d*x+c)^3+3*(43*A-115*B)*\cos(d*x+c)^2+3*(43*A-115*B)*\cos(d*x+c)+43*A-115*B)*\sqrt{a}*\arctan(\sqrt{2)*\sqrt{a*\cos(d*x+c)+a}*\sqrt{\cos(d*x+c)}/(\sqrt{a}*\sin(d*x+c))))+2*(16*B*\cos(d*x+c)^2-5*(3*A-11*B)*\cos(d*x+c)-11*A+35*B)*\sqrt{a*\cos(d*x+c)+a}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-32*((2*A-5*B)*\cos(d*x+c)^3+3*(2*A-5*B)*\cos(d*x+c)^2+3*(2*A-5*B)*\cos(d*x+c)+2*A-5*B)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x+c)+a}*\sqrt{\cos(d*x+c)}/(\sqrt{a}*\sin(d*x+c))))/(a^3*d*\cos(d*x+c)^3+3*a^3*d*\cos(d*x+c)^2+3*a^3*d*\cos(d*x+c)+a^3*d)$

3.202.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.202.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(a\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^(5/
2), x)`

3.202.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.202.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \int \frac{\cos(c+dx)^{\frac{5}{2}}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`

3.203 $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

3.203.1 Optimal result 1906
 3.203.2 Mathematica [A] (verified) 1907
 3.203.3 Rubi [A] (verified) 1907
 3.203.4 Maple [B] (verified) 1911
 3.203.5 Fricas [A] (verification not implemented) 1912
 3.203.6 Sympy [F(-1)] 1913
 3.203.7 Maxima [F] 1913
 3.203.8 Giac [F(-1)] 1913
 3.203.9 Mupad [F(-1)] 1914

3.203.1 Optimal result

Integrand size = 35, antiderivative size = 194

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{5/2}d} + \frac{(3A-43B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A-B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{(3A-11B)\sqrt{\cos(c+dx)} \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}}$$

```
output 2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d+1/4*(A-B)*
cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)+1/32*(3*A-43*B)*arctan(
1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/
a^(5/2)/d*2^(1/2)+1/16*(3*A-11*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos
(d*x+c))^(3/2)
```

3.203.2 Mathematica [A] (verified)

Time = 1.17 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.30

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx =$$

$$\left(88B \arcsin\left(\sqrt{1-\cos(c+dx)}\right) \cos^4\left(\frac{1}{2}(c+dx)\right) + 344B \arcsin\left(\sqrt{\cos(c+dx)}\right) \cos^4\left(\frac{1}{2}(c+dx)\right) + 4\sqrt{\cos(c+dx)}\right)$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `-1/32*((88*B*ArcSin[Sqrt[1 - Cos[c + d*x]]]*Cos[(c + d*x)/2]^4 + 344*B*ArcSin[Sqrt[Cos[c + d*x]]]*Cos[(c + d*x)/2]^4 + 4*Sqrt[2]*(3*A - 43*B)*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]]*Cos[(c + d*x)/2]^4 - 14*A*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) + 30*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) - 6*A*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])] + 22*B*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])]*Sin[c + d*x])/(d*Sqrt[1 - Cos[c + d*x]])*(a*(1 + Cos[c + d*x]))^(5/2))`

3.203.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^{5/2}} dx$$

↓ 3456

3.203. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+8aB \cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+8aB \cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)+8aB \sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3456} \\
& \frac{\int \frac{(3A-11B)a^2+32B \cos(c+dx)a^2}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{a(3A-11B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{(3A-11B)a^2+32B \cos(c+dx)a^2}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{a(3A-11B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(3A-11B)a^2+32B \sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{a(3A-11B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3461} \\
& \frac{a^2(3A-43B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + 32aB \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{8a^2} + \frac{a(3A-11B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a^2(3A-43B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + 32aB \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{8a^2} + \frac{a(3A-11B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \\
& \quad \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx) + a)^{5/2}}
\end{aligned}$$

3.203. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

↓ 3253

$$\frac{a^2(3A-43B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{64aB \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} + \frac{a(3A-11B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

↓ 223

$$\frac{a^2(3A-43B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{64a^{3/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{4a^2} + \frac{a(3A-11B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

↓ 3261

$$\frac{\frac{64a^{3/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{2a^3(3A-43B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}}\right)}{4a^2} + \frac{a(3A-11B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

↓ 218

$$\frac{\frac{\sqrt{2}a^{3/2}(3A-43B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{64a^{3/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{a(3A-11B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}$$

```
input Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2), x]
```

3.203. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$

```
output ((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x]/(4*d*(a + a*Cos[c + d*x])^(5/2))
+ (((64*a^(3/2)*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]]
)/d + (Sqrt[2]*a^(3/2)*(3*A - 43*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]
*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/(4*a^2) + (a*(3*A - 11*
B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x]/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a
^2)
```

3.203.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3253 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Co
s[e + f*x]/Sqrt[a + b*Ssin[e + f*x]])], x] /; FreeQ[{a, b, d, e, f}, x] && E
qQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*S
sin[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.203.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 446 vs. 2(163) = 326.

Time = 6.06 (sec) , antiderivative size = 447, normalized size of antiderivative = 2.30

method	result
default	$-\frac{(3A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))-43B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))+6A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))-6 \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c)))}{32d(1+\cos(dx+c))^3 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A(7\sqrt{2} \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3 \sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 3 \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c)) - 6 \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c)))}{32d(1+\cos(dx+c))^3 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2), x, method=_RET
URNVERBOSE)
```

3.203.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

output
$$\begin{aligned} & -1/32/a^3/d*(3*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2-43*B*2 \\ & ^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2+6*A*2^{(1/2)}*\arcsin(\cot(d \\ & *x+c)-\csc(d*x+c))*\cos(d*x+c)-14*A*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos \\ & (d*x+c)))^{(1/2)}-86*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)-64*B \\ & *\cos(d*x+c)^2*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})+30*B*\cos \\ & (d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+3*A*2^{(1/2)}*\arcsin(\cot \\ & (d*x+c)-\csc(d*x+c))-6*A*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\sin(d*x+c)-43* \\ & B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-128*B*\cos(d*x+c)*\arctan(\tan(d*x+c) \\ & *(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)})+22*B*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x \\ & +c)))^{(1/2)}-64*B*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}))* \\ & (a*(1+\cos(d*x+c)))^{(1/2)}*\cos(d*x+c)^{(1/2)}/(1+\cos(d*x+c))^{3/2}/(\cos(d*x+c)/(1+\cos \\ & (d*x+c)))^{(1/2)} \end{aligned}$$

3.203.5 Fracas [A] (verification not implemented)

Time = 6.87 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.38

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \sqrt{2}((3A-43B)\cos(dx+c)^3 + 3(3A-43B)\cos(dx+c)^2 + 3(3A-43B)\cos(dx+c) + 3A-43B)$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="fracas")`

output
$$\begin{aligned} & -1/32*(\sqrt{2})*((3*A - 43*B)*\cos(d*x + c)^3 + 3*(3*A - 43*B)*\cos(d*x + c)^2 + 3*(3*A - 43*B)*\cos(d*x + c) + 3*A - 43*B)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{ \\ & a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c))) - 2*((7*A - 15*B)*\cos(d*x + c) + 3*A - 11*B)*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + \\ & c)}*\sin(d*x + c) + 64*(B*\cos(d*x + c)^3 + 3*B*\cos(d*x + c)^2 + 3*B*\cos(d*x \\ & + c) + B)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{ \\ & a}*\sin(d*x + c)))/((a^3*d*\cos(d*x + c)^3 + 3*a^3*d*\cos(d*x + c)^2 + 3*a^3 \\ & *d*\cos(d*x + c) + a^3*d) \end{aligned}$$

3.203.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$$

3.203.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.203.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(a\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(5/
2), x)`

3.203.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.203.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(5/2),x)`

3.204
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

3.204.1 Optimal result 1915
 3.204.2 Mathematica [A] (verified) 1915
 3.204.3 Rubi [A] (verified) 1916
 3.204.4 Maple [B] (verified) 1919
 3.204.5 Fracas [A] (verification not implemented) 1919
 3.204.6 Sympy [F] 1920
 3.204.7 Maxima [F] 1920
 3.204.8 Giac [F(-1)] 1921
 3.204.9 Mupad [F(-1)] 1921

3.204.1 Optimal result

Integrand size = 35, antiderivative size = 154

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{(5A+3B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{(A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{4d(a+a \cos(c+dx))^{5/2}} + \frac{(A+7B)\sqrt{\cos(c+dx)} \sin(c+dx)}{16ad(a+a \cos(c+dx))^{3/2}}$$

output `1/32*(5*A+3*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)+1/4*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(5/2)+1/16*(A+7*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(3/2)`

3.204.2 Mathematica [A] (verified)

Time = 4.63 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{12B \arcsin\left(\frac{\sin(\frac{1}{2}(c+dx))}{\sqrt{\cos^2(\frac{1}{2}(c+dx))}}\right) \cos^5\left(\frac{1}{2}(c+dx)\right) - 10\sqrt{2}A \operatorname{arctanh}\left(\frac{\sin(\frac{1}{2}(c+dx))}{\sqrt{\cos^2(\frac{1}{2}(c+dx))}}\right)}{\dots}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

3.204.
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx$$

output $(12*B*ArcSin[\sin[(c + dx)/2]/\sqrt{\cos[(c + dx)/2]^2}]*\cos[(c + dx)/2]^5 - 10*\sqrt{2}*A*ArcTanh[\sqrt{-(\sec[c + dx]*\sin[(c + dx)/2]^2)}]*\cos[(c + dx)/2]^4*\sqrt{-1 + \cos[c + dx]}*Cot[(c + dx)/2] + \sqrt{\cos[c + dx]}*(5*A + 3*B + (A + 7*B)*\cos[c + dx])*sin[c + dx])/(16*d*(a*(1 + \cos[c + dx]))^{(5/2)})$

3.204.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3456, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 3456

$$\frac{\int \frac{a(A-B)+2a(A+3B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 27

$$\frac{\int \frac{a(A-B)+2a(A+3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{a(A-B)+2a(A+3B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 3457

$$\frac{\int \frac{a^2(5A+3B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a(A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}$$

3.204. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\frac{1}{4}(5A + 3B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{a(A+7B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \\
& \quad \frac{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}}{4d(a \cos(c + dx) + a)^{5/2}} \\
& \downarrow 3042 \\
& \frac{\frac{1}{4}(5A + 3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{a(A+7B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \\
& \quad \frac{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}}{4d(a \cos(c + dx) + a)^{5/2}} \\
& \downarrow 3261 \\
& \frac{\frac{a(A+7B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{a(5A+3B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} dx \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} \right)}{8a^2} + \\
& \quad \frac{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}}{4d(a \cos(c + dx) + a)^{5/2}} \\
& \downarrow 218 \\
& \frac{(5A+3B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}\sqrt{ad}} + \frac{a(A+7B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \\
& \quad \frac{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}}{4d(a \cos(c + dx) + a)^{5/2}}
\end{aligned}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(5/2),x]`

output `((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (((5*A + 3*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*Sqrt[a]*d) + (a*(A + 7*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2)`

3.204.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.204.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. 2(129) = 258.

Time = 5.96 (sec) , antiderivative size = 343, normalized size of antiderivative = 2.23

method	result
default	$\frac{(-5A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))-3B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))+2A \cos(dx+c) \sin(dx+c))}{(a+\cos(dx+c))^5}$
parts	$\frac{A(\sqrt{2} \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+5 \sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-5 \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))-10 \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c))}{32d(1+\cos(dx+c))^3 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

input `int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)`

output `1/32/a^3/d*(-5*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2-3*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2+2*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-10*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+14*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-6*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+10*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)-5*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+6*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)/(1+cos(d*x+c))^(1/2)*cos(d*x+c)^(1/2)/(1+cos(d*x+c))^(1/2)+3/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)`

3.204.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.40

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{5/2}} dx = \frac{\sqrt{2}((5A+3B) \cos(dx+c)^3+3(5A+3B) \cos(dx+c)^2+3(5A+3B) \cos(dx+c)+3A)}{(a+a \cos(dx+c))^5}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x,algorithm="fracas")`

output `1/32*(sqrt(2)*((5*A + 3*B)*cos(d*x + c)^3 + 3*(5*A + 3*B)*cos(d*x + c)^2 + 3*(5*A + 3*B)*cos(d*x + c) + 5*A + 3*B)*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) + 2*((A + 7*B)*cos(d*x + c) + 5*A + 3*B)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.204.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{(A+B\cos(c+dx))\sqrt{\cos(c+dx)}}{(a(\cos(c+dx)+1))^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(cos(c + d*x))/(a*(cos(c + d*x) + 1))**(5/2), x)`

3.204.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(a\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(a*cos(d*x + c) + a)^(5/2), x)`

3.204.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2),x, algorith="giac")`

output `Timed out`

3.204.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{5/2}} dx$$

input `int((cos(c+d*x)^(1/2)*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(5/2),x)`

output `int((cos(c+d*x)^(1/2)*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(5/2),x)`

3.205 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{5/2}} dx$

3.205.1 Optimal result 1922
 3.205.2 Mathematica [A] (verified) 1922
 3.205.3 Rubi [A] (verified) 1923
 3.205.4 Maple [B] (verified) 1926
 3.205.5 Fricas [A] (verification not implemented) 1926
 3.205.6 Sympy [F] 1927
 3.205.7 Maxima [F] 1927
 3.205.8 Giac [F(-1)] 1928
 3.205.9 Mupad [F(-1)] 1928

3.205.1 Optimal result

Integrand size = 35, antiderivative size = 156

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \frac{(19A + 5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} - \frac{(A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}} - \frac{(9A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}}$$

output `1/32*(19*A+5*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)-1/4*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(5/2)-1/16*(9*A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(3/2)`

3.205.2 Mathematica [A] (verified)

Time = 1.61 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.96

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \frac{\sec^2\left(\frac{1}{2}(c + dx)\right) \left(4(19A + 5B)\operatorname{arctanh}\left(\sqrt{-\sec(c + dx)} \sin^2\left(\frac{1}{2}(c + dx)\right)\right)\right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(5/2)),x]`

output $(\text{Sec}[(c + d*x)/2]^2*(4*(19*A + 5*B)*\text{ArcTanh}[\text{Sqrt}[-(\text{Sec}[c + d*x]*\text{Sin}[(c + d*x)/2]^2)]]*\text{Cos}[(c + d*x)/2]^4 + \text{Cos}[c + d*x]*(-13*A + 5*B + (-9*A + B)*\text{Cos}[c + d*x])*\text{Sqrt}[2 - 2*\text{Sec}[c + d*x]])*\text{Tan}[(c + d*x)/2])/(32*\text{Sqrt}[2]*a^2*d*\text{Sqrt}[-1 + \text{Cos}[c + d*x]]*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])])$

3.205.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2}} dx$$

↓ 3457

$$\frac{\int \frac{a(7A+B) - 2a(A-B) \cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d(a \cos(c+dx) + a)^{5/2}}$$

↓ 27

$$\frac{\int \frac{a(7A+B) - 2a(A-B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d(a \cos(c+dx) + a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{a(7A+B) - 2a(A-B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(\sin(c+dx + \frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d(a \cos(c+dx) + a)^{5/2}}$$

↓ 3457

$$\frac{\int \frac{a^2(19A+5B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a(9A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx) + a)^{3/2}} - \frac{(A-B) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d(a \cos(c+dx) + a)^{5/2}}$$

↓ 27

3.205. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{5/2}} dx$

$$\frac{\frac{1}{4}(19A + 5B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a(9A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{\frac{8a^2}{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}} \frac{1}{4d(a \cos(c + dx) + a)^{5/2}}} -$$

↓ 3042

$$\frac{\frac{1}{4}(19A + 5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{a(9A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{\frac{8a^2}{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}} \frac{1}{4d(a \cos(c + dx) + a)^{5/2}}} -$$

↓ 3261

$$\frac{a(19A+5B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right) - \frac{a(9A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{\frac{8a^2}{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}} \frac{1}{4d(a \cos(c + dx) + a)^{5/2}}} -$$

↓ 218

$$\frac{(19A+5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right) - \frac{a(9A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{\frac{8a^2}{(A - B) \sin(c + dx)\sqrt{\cos(c + dx)}} \frac{1}{4d(a \cos(c + dx) + a)^{5/2}}} -$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(5/2)),x]`

output `-1/4*((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^(5/2)) + (((19*A + 5*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*Sqrt[a]*d) - (a*(9*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2)`

3.205.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.205.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. 2(131) = 262.

Time = 8.03 (sec) , antiderivative size = 343, normalized size of antiderivative = 2.20

method	result
default	$-\frac{(19A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))+5B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))+38A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c)))+18A\cos(dx+c)\sin(dx+c)(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}+10B2^{1/2}\arcsin(\cot(dx+c)-\csc(dx+c))\cos(dx+c)-2B\cos(dx+c)\sin(dx+c)(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}+19A2^{1/2}\arcsin(\cot(dx+c)-\csc(dx+c))+26A(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}\sin(dx+c)+5B2^{1/2}\arcsin(\cot(dx+c)-\csc(dx+c))-10B\sin(dx+c)(\cos(dx+c)/(1+\cos(dx+c)))^{1/2})\cos(dx+c)^{1/2}(a(1+\cos(dx+c)))^{1/2}/(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}/(1+\cos(dx+c))^{3/2}}{A\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}}\left(-2(\csc^3(dx+c))\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}(1-\cos(dx+c))\right)+32d\sqrt{-\frac{(\csc^2(dx+c))(1-\cos(dx+c))}{(\csc^2(dx+c))(1-\cos(dx+c))}}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/32/a^3/d*(19*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2+5*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2+38*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+18*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+10*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)-2*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+19*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+26*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+5*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-10*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^3
```

3.205.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.39

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2}((19A + 5B) \cos(dx + c)^3 + 3(19A + 5B) \cos(dx + c)^2 + \dots)}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(5/2),x, algorith="fracas")
```

output `1/32*(sqrt(2)*((19*A + 5*B)*cos(d*x + c)^3 + 3*(19*A + 5*B)*cos(d*x + c)^2 + 3*(19*A + 5*B)*cos(d*x + c) + 19*A + 5*B)*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) - 2*((9*A - B)*cos(d*x + c) + 13*A - 5*B)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.205.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{(a(\cos(c + dx) + 1))^{5/2} \sqrt{\cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))/((a*(cos(c + d*x) + 1))**(5/2)*sqrt(cos(c + d*x))), x)`

3.205.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{5/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*sqrt(cos(d*x + c))), x)`

3.205.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.205.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(5/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(5/2)),
x)`

$$3.206 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}} dx$$

3.206.1 Optimal result 1929
 3.206.2 Mathematica [C] (warning: unable to verify) 1930
 3.206.3 Rubi [A] (verified) 1931
 3.206.4 Maple [B] (verified) 1934
 3.206.5 Fricas [A] (verification not implemented) 1935
 3.206.6 Sympy [F(-1)] 1936
 3.206.7 Maxima [F(-1)] 1936
 3.206.8 Giac [F] 1936
 3.206.9 Mupad [F(-1)] 1937

3.206.1 Optimal result

Integrand size = 35, antiderivative size = 203

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{(75A - 19B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d}$$

$$- \frac{(A - B) \sin(c + dx)}{4d\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}} - \frac{(13A - 5B) \sin(c + dx)}{16ad\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}}$$

$$+ \frac{(49A - 9B) \sin(c + dx)}{16a^2d\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}$$

```
output -1/32*(75*A-19*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/
(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)-1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d
*x+c))^(5/2)/cos(d*x+c)^(1/2)-1/16*(13*A-5*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+
c))^(3/2)/cos(d*x+c)^(1/2)+1/16*(49*A-9*B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(1/
2)/(a+a*cos(d*x+c))^(1/2)
```


3.206.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.66 (sec) , antiderivative size = 728, normalized size of antiderivative = 3.59

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{B \cos^5\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^4\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \left(11 - 31 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 18 \sin^4\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{19 \operatorname{arctanh}\left(\sqrt{\frac{-1 - \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{1 - \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}}\right)}{4d(a(1 + \cos(c + dx)))^{5/2} \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}}\right)}{2A \cos^5\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^4\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{8 \cos^6\left(\frac{1}{2}(c + dx)\right) {}_4F_3\left(2, 2, 2, \frac{5}{2}; 1, 1, \frac{11}{2}; \frac{\sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{315(-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right))} + \frac{1}{12}\right)}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(5/2)),x]`

output `-1/4*(B*Cos[c/2 + (d*x)/2]^5*Sec[(c + d*x)/2]^4*Sin[c/2 + (d*x)/2]*(11 - 31*Sin[c/2 + (d*x)/2]^2 + 18*Sin[c/2 + (d*x)/2]^4 - (19*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])*Cos[(c + d*x)/2]^4/Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]))/(d*(a*(1 + Cos[c + d*x]))^(5/2)*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (2*A*Cos[c/2 + (d*x)/2]^5*Sec[(c + d*x)/2]^4*Sin[c/2 + (d*x)/2]*((8*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2])*Sin[c/2 + (d*x)/2]^2/(315*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) + (Csc[c/2 + (d*x)/2]^8*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-15*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)])*Cos[(c + d*x)/2]^4*(-343 + 1465*Sin[c/2 + (d*x)/2]^2 - 2021*Sin[c/2 + (d*x)/2]^4 + 824*Sin[c/2 + (d*x)/2]^6) + Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2]*(-5145 + 33980*Sin[c/2 + (d*x)/2]^2 - 87764*Sin[c/2 + (d*x)/2]^4 + 109737*Sin[c/2 + (d*x)/2]^6 - 66122*Sin[c/2 + (d*x)/2]^8 + 15344*Sin[c/2 + (d*x)/2]^10))/120))/(d*(a*(1 + Cos[c + d*x]))^(5/2)*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2))`

3.206.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} (a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2}} dx$$

↓ 3457

$$\frac{\int \frac{a(9A-B) - 4a(A-B) \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B) \sin(c+dx)}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}}$$

↓ 27

$$\frac{\int \frac{a(9A-B) - 4a(A-B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \sin(c+dx)}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}}$$

↓ 3042

$$\frac{\int \frac{a(9A-B) - 4a(A-B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{3/2} (\sin(c+dx + \frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \sin(c+dx)}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}}$$

↓ 3457

$$\frac{\int \frac{a^2(49A-9B) - 2a^2(13A-5B) \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a(13A-5B) \sin(c+dx)}{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}}$$

$$\frac{8a^2}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}} - \frac{(A-B) \sin(c+dx)}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}}$$

↓ 27

$$\frac{\int \frac{a^2(49A-9B) - 2a^2(13A-5B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(13A-5B) \sin(c+dx)}{2d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{3/2}}$$

$$\frac{8a^2}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}} - \frac{(A-B) \sin(c+dx)}{4d \sqrt{\cos(c+dx)}(a \cos(c+dx) + a)^{5/2}}$$

3.206. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{a^2(49A-9B)-2a^2(13A-5B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx \\
& \quad - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \\
& \quad - \frac{8a^2}{4a^2} \frac{(A-B)\sin(c+dx)}{\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 3042 \\
& \quad \frac{2\int -\frac{a^3(75A-19B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \\
& \quad - \frac{8a^2}{4a^2} \frac{(A-B)\sin(c+dx)}{\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 3463 \\
& \quad \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a^2(75A-19B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx \\
& \quad - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \\
& \quad - \frac{8a^2}{4a^2} \frac{(A-B)\sin(c+dx)}{\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 27 \\
& \quad \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a^2(75A-19B)\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx \\
& \quad - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \\
& \quad - \frac{8a^2}{4a^2} \frac{(A-B)\sin(c+dx)}{\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 3042 \\
& \quad \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a^2(75A-19B)\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx \\
& \quad - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \\
& \quad - \frac{8a^2}{4a^2} \frac{(A-B)\sin(c+dx)}{\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 3261 \\
& \quad \frac{2a^3(75A-19B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} + \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \\
& \quad - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \\
& \quad - \frac{8a^2}{4a^2} \frac{(A-B)\sin(c+dx)}{\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
& \quad \downarrow 218
\end{aligned}$$

3.206. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{5/2}} dx$

$$\frac{\frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{\sqrt{2}a^{3/2}(75A-19B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{4a^2}}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}}$$

$$\frac{8a^2}{(A-B)\sin(c+dx)}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(5/2)),x]`

output `-1/4*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(13*A - 5*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)) + (-((Sqrt[2]*a^(3/2)*(75*A - 19*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d) + (2*a^2*(49*A - 9*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/(4*a^2))/(8*a^2)`

3.206.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.206.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 484 vs. 2(172) = 344.

Time = 8.18 (sec) , antiderivative size = 485, normalized size of antiderivative = 2.39

method	result
default	$\frac{(75A(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c))-19B(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2}\arcsin(\cot(dx+c)-\csc(dx+c)))}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{5/2}}$
parts	$A\left(75(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))+49\sqrt{2}(\cos^2(dx+c))\sin(dx+c)+225\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\right)$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RET
URNVERBOSE)
```

$$3.206. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}} dx$$

output `1/32/a^3/d*(75*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-19*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+225*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))-57*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+225*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-57*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+75*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*A*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+98*A*sin(d*x+c)*cos(d*x+c)^2-19*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*B*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-18*B*sin(d*x+c)*cos(d*x+c)^2+170*A*sin(d*x+c)*cos(d*x+c)-26*B*sin(d*x+c)*cos(d*x+c)+64*A*sin(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^3/cos(d*x+c)^(1/2)`

3.206.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.22

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx =$$

$$\sqrt{2}((75A - 19B) \cos(dx + c)^4 + 3(75A - 19B) \cos(dx + c)^3 + 3(75A - 19B) \cos(dx + c)^2 + (75A -$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2),x, algorith="fricas")`

output `-1/32*(sqrt(2))*((75*A - 19*B)*cos(d*x + c)^4 + 3*(75*A - 19*B)*cos(d*x + c)^3 + 3*(75*A - 19*B)*cos(d*x + c)^2 + (75*A - 19*B)*cos(d*x + c))*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) - 2*((49*A - 9*B)*cos(d*x + c)^2 + (85*A - 13*B)*cos(d*x + c) + 32*A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^3*d*cos(d*x + c)^4 + 3*a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + a^3*d*cos(d*x + c))`

3.206.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.206.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.206.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(3/2)), x)`

3.206.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} (a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(5/2)),x)`

$$3.207 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{\frac{5}{2}}} dx$$

3.207.1 Optimal result	1938
3.207.2 Mathematica [C] (warning: unable to verify)	1939
3.207.3 Rubi [A] (verified)	1939
3.207.4 Maple [B] (verified)	1944
3.207.5 Fracas [A] (verification not implemented)	1944
3.207.6 Sympy [F(-1)]	1945
3.207.7 Maxima [F]	1945
3.207.8 Giac [F]	1946
3.207.9 Mupad [F(-1)]	1946

3.207.1 Optimal result

Integrand size = 35, antiderivative size = 250

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{\frac{5}{2}}} dx = \frac{(163A - 75B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{16\sqrt{2}a^{\frac{5}{2}}d} - \frac{(A - B) \sin(c + dx)}{4d \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{\frac{5}{2}}} - \frac{(17A - 9B) \sin(c + dx)}{16ad \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{\frac{3}{2}}} + \frac{(95A - 39B) \sin(c + dx)}{48a^2d \cos^{\frac{3}{2}}(c + dx)\sqrt{a + a \cos(c + dx)}} - \frac{(299A - 147B) \sin(c + dx)}{48a^2d \sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}$$

output `-1/4*(A-B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2)-1/16*(17*A-9*B)*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2)+1/32*(163*A-75*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(5/2)/d*2^(1/2)+1/48*(95*A-39*B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)-1/48*(299*A-147*B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)`

3.207.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.07 (sec) , antiderivative size = 1148, normalized size of antiderivative = 4.59

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(5/2)),x]`

output `(2*B*Cos[c/2 + (d*x)/2]^5*Sec[(c + d*x)/2]^4*Sin[c/2 + (d*x)/2]*((8*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2)/(315*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) + (Csc[c/2 + (d*x)/2]^8*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-15*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Cos[(c + d*x)/2]^4*(-343 + 1465*Sin[c/2 + (d*x)/2]^2 - 2021*Sin[c/2 + (d*x)/2]^4 + 824*Sin[c/2 + (d*x)/2]^6) + Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-5145 + 33980*Sin[c/2 + (d*x)/2]^2 - 87764*Sin[c/2 + (d*x)/2]^4 + 109737*Sin[c/2 + (d*x)/2]^6 - 66122*Sin[c/2 + (d*x)/2]^8 + 15344*Sin[c/2 + (d*x)/2]^10))/120))/(d*(a*(1 + Cos[c + d*x])^(5/2)*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) - (A*Cot[c/2 + (d*x)/2]^5*Csc[c/2 + (d*x)/2]^4*Sec[(c + d*x)/2]^4*(640*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 7/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12 - 1280*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 7/2}, {1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12*(-6 + 5*Sin[c/2 + (d*x)/2]^2) + 33*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-105*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Cos[(c + d*x)/2]^4*(-10935 + 72902*Sin[c/2 + (d*x)/2]^2 - 188110*Sin[c/2 + (d*x)/2]^4 + 2...`

3.207.3 Rubi [A] (verified)

Time = 1.43 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.207. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}} dx$

$$\begin{aligned} & \downarrow 3463 \\ & \frac{2 \int -\frac{a^3(299A-147B)-2a^3(95A-39B)\cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{2a^2(95A-39B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{a(17A-9B)\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \\ & \frac{8a^2}{(A-B)\sin(c+dx)} \\ & \frac{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{5/2}} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2a^2(95A-39B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^3(299A-147B)-2a^3(95A-39B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \\ & \frac{8a^2}{(A-B)\sin(c+dx)} \\ & \frac{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{5/2}} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2a^2(95A-39B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^3(299A-147B)-2a^3(95A-39B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \\ & \frac{8a^2}{(A-B)\sin(c+dx)} \\ & \frac{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{5/2}} \end{aligned}$$

$$\begin{aligned} & \downarrow 3463 \\ & \frac{2a^2(95A-39B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2 \int -\frac{3a^4(163A-75B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{2a^3(299A-147B)\sin(c+dx)}{3a \sqrt{d \cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - \frac{a(17A-9B)\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \\ & \frac{8a^2}{(A-B)\sin(c+dx)} \\ & \frac{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{5/2}} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2a^2(95A-39B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2a^3(299A-147B)\sin(c+dx)}{d \sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - \frac{3a^3(163A-75B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{a(17A-9B)\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \\ & \frac{8a^2}{(A-B)\sin(c+dx)} \\ & \frac{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{5/2}} \end{aligned}$$

3.207. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}} dx$

↓ 3042

$$\frac{\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 3a^3(163A-75B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2}}{8a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

↓ 3261

$$\frac{\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{6a^4(163A-75B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d \left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} \right)}{4a^2} + \frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}}{8a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

↓ 218

$$\frac{\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}a^{5/2}(163A-75B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{3a}}{4a^2}}{8a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(5/2)),x]`

output `-1/4*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(17*A - 9*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a^2*(95*A - 39*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*a^(5/2)*(163*A - 75*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^3*(299*A - 147*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(4*a^2))/(8*a^2)`

3.207. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))^{5/2}} dx$

3.207.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.207.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 534 vs. $2(213) = 426$.

Time = 7.98 (sec) , antiderivative size = 535, normalized size of antiderivative = 2.14

method	result
default	$-\frac{(489A\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^4(dx+c))-225B\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^4(dx+c)))}{(a+\cos(dx+c))^5}$
parts	$-\frac{A(489\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^4(dx+c))+299\sqrt{2}(\cos^3(dx+c))\sin(dx+c)+1467(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})}{(a+\cos(dx+c))^5}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/96/a^3/d*(489*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^4-225*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^4+1467*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-675*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+1467*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))-675*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+489*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+598*A*cos(d*x+c)^3*sin(d*x+c)-225*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-294*B*sin(d*x+c)*cos(d*x+c)^3+1006*A*sin(d*x+c)*cos(d*x+c)^2-510*B*sin(d*x+c)*cos(d*x+c)^2+320*A*sin(d*x+c)*cos(d*x+c)-192*B*sin(d*x+c)*cos(d*x+c)-64*A*sin(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^3/cos(d*x+c)^(3/2)
```

3.207.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.08

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \frac{3\sqrt{2}((163A - 75B)\cos(dx + c)^5 + 3(163A - 75B)\cos(dx + c))}{(a + a \cos(c + dx))^{5/2}}$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(5/2),x,algorithm="fracas")
```

$$3.207. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{5/2}} dx$$

output $1/96*(3*\sqrt{2})*((163*A - 75*B)*\cos(dx + c)^5 + 3*(163*A - 75*B)*\cos(dx + c)^4 + 3*(163*A - 75*B)*\cos(dx + c)^3 + (163*A - 75*B)*\cos(dx + c)^2)*\sqrt{a}*\arctan(1/2*\sqrt{2}*\sqrt{a*\cos(dx + c) + a}*\sqrt{a}*\sqrt{\cos(dx + c)})*\sin(dx + c)/(a*\cos(dx + c)^2 + a*\cos(dx + c))) - 2*((299*A - 147*B)*\cos(dx + c)^3 + (503*A - 255*B)*\cos(dx + c)^2 + 32*(5*A - 3*B)*\cos(dx + c) - 32*A)*\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)}*\sin(dx + c))/(a^3*d*\cos(dx + c)^5 + 3*a^3*d*\cos(dx + c)^4 + 3*a^3*d*\cos(dx + c)^3 + a^3*d*\cos(dx + c)^2)$

3.207.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(5/2),x)`

output Timed out

3.207.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(5/2)), x)`

3.207.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(5
/2)), x)`

3.207.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(5/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(5/2)),
x)`

3.208
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

3.208.1 Optimal result 1947
 3.208.2 Mathematica [A] (warning: unable to verify) 1948
 3.208.3 Rubi [A] (verified) 1948
 3.208.4 Maple [B] (verified) 1954
 3.208.5 Fricas [A] (verification not implemented) 1955
 3.208.6 Sympy [F(-1)] 1956
 3.208.7 Maxima [F] 1956
 3.208.8 Giac [F(-1)] 1957
 3.208.9 Mupad [F(-1)] 1957

3.208.1 Optimal result

Integrand size = 35, antiderivative size = 293

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx = \frac{(2A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{7/2}d} - \frac{(177A-637B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d} + \frac{(A-B) \cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{6d(a+a \cos(c+dx))^{7/2}} + \frac{(3A-7B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{16ad(a+a \cos(c+dx))^{5/2}} + \frac{(79A-259B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{192a^2d(a+a \cos(c+dx))^{3/2}} - \frac{7(7A-27B) \sqrt{\cos(c+dx)} \sin(c+dx)}{64a^3d\sqrt{a+a \cos(c+dx)}}$$

output

```
(2*A-7*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d+1/6*(A-B)*cos(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)+1/16*(3*A-7*B)*cos(d*x+c)^(5/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)+1/192*(79*A-259*B)*cos(d*x+c)^(3/2)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)-1/128*(177*A-637*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d*2^(1/2)-7/64*(7*A-27*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^3/d/(a+a*cos(d*x+c))^(1/2)
```

3.208.2 Mathematica [A] (warning: unable to verify)

Time = 2.94 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.29

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \frac{\sqrt{a(1+\cos(c+dx))} \left(-336(7A-27B) \arcsin\left(\sqrt{1-\cos(c+dx)}\right) \right)}{\dots}$$

input `Integrate[(Cos[c + d*x]^(7/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])] * (-336*(7*A - 27*B)*ArcSin[Sqrt[1 - Cos[c + d*x]]] * Cos[(c + d*x)/2]^6 - 8496*A*ArcSin[Sqrt[Cos[c + d*x]]] * Cos[(c + d*x)/2]^6 + 30576*B*ArcSin[Sqrt[Cos[c + d*x]]] * Cos[(c + d*x)/2]^6 + 24*Sqrt[2] * (177*A - 637*B)*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2]] * Cos[(c + d*x)/2]^6 - 724*A*Sqrt[1 - Cos[c + d*x]] * Cos[c + d*x]^(3/2) + 2884*B*Sqrt[1 - Cos[c + d*x]] * Cos[c + d*x]^(3/2) - 494*A*Sqrt[1 - Cos[c + d*x]] * Cos[c + d*x]^(5/2) + 2198*B*Sqrt[1 - Cos[c + d*x]] * Cos[c + d*x]^(5/2) + 384*B*Sqrt[1 - Cos[c + d*x]] * Cos[c + d*x]^(7/2) - 294*A*Sqrt[-((-1 + Cos[c + d*x]) * Cos[c + d*x])] + 1134*B*Sqrt[-((-1 + Cos[c + d*x]) * Cos[c + d*x])] * Sin[c + d*x]) / (384*a^4*d*Sqrt[1 - Cos[c + d*x]] * (1 + Cos[c + d*x])^4)`

3.208.3 Rubi [A] (verified)

Time = 1.99 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.08, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3456, 27, 3042, 3462, 25, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{7/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{7/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{7/2}} dx$$

↓ 3456

3.208. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{\cos^{\frac{5}{2}}(c+dx)(7a(A-B)-2a(A-7B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\cos^{\frac{5}{2}}(c+dx)(7a(A-B)-2a(A-7B)\cos(c+dx))}{(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}(7a(A-B)-2a(A-7B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow 3456 \\
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(15a^2(3A-7B)-2a^2(17A-77B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} + \\
& \quad \frac{12a^2}{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \frac{6d(a\cos(c+dx)+a)^{7/2}}{} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(15a^2(3A-7B)-2a^2(17A-77B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} + \\
& \quad \frac{12a^2}{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \frac{6d(a\cos(c+dx)+a)^{7/2}}{} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(15a^2(3A-7B)-2a^2(17A-77B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} + \\
& \quad \frac{12a^2}{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \frac{6d(a\cos(c+dx)+a)^{7/2}}{} \\
& \quad \downarrow 3456 \\
& \frac{\int \frac{3\sqrt{\cos(c+dx)}(a^3(79A-259B)-14a^3(7A-27B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a^2(79A-259B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} + \\
& \quad \frac{12a^2}{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)} \\
& \quad \frac{6d(a\cos(c+dx)+a)^{7/2}}{}
\end{aligned}$$

3.208. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

↓ 27

$$\frac{3 \int \frac{\sqrt{\cos(c+dx)} (a^3(79A-259B) - 14a^3(7A-27B) \cos(c+dx))}{4a^2} dx + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}}{8a^2} + \frac{12a^2}{6d(a \cos(c+dx)+a)^{7/2}} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3042

$$\frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} (a^3(79A-259B) - 14a^3(7A-27B) \sin(c+dx+\frac{\pi}{2}))}{4a^2} dx + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}}{8a^2} + \frac{12a^2}{6d(a \cos(c+dx)+a)^{7/2}} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3462

$$\frac{3 \left(\frac{\int -\frac{7a^4(7A-27B) - 64a^4(2A-7B) \cos(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx - \frac{14a^3(7A-27B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right)}{4a^2} + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}}{8a^2} + \frac{12a^2}{6d(a \cos(c+dx)+a)^{7/2}} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 25

$$\frac{3 \left(-\frac{\int \frac{7a^4(7A-27B) - 64a^4(2A-7B) \cos(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx - \frac{14a^3(7A-27B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right)}{4a^2} + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}}}{8a^2} + \frac{12a^2}{6d(a \cos(c+dx)+a)^{7/2}} (A-B) \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)$$

↓ 3042

3.208. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$

$$3 \left(\frac{\int \frac{7a^4(7A-27B)-64a^4(2A-7B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}a+a} dx - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(79A-259B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B)\sin(c+dx)}{4d(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 3461

$$3 \left(-\frac{a^4(177A-637B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx - 64a^3(2A-7B)\int \frac{\sqrt{\cos(c+dx)}a+a}{\sqrt{\cos(c+dx)}} dx - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(79A-259B)\sin(c+dx)}{2d(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 3042

$$3 \left(-\frac{a^4(177A-637B)\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}a+a} dx - 64a^3(2A-7B)\int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}a+a}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(79A-259B)\sin(c+dx)}{2d(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 3253

$$3 \left(-\frac{a^4(177A-637B)\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}a+a} dx + \frac{128a^3(2A-7B)\int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)}a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}a+a}\right)}{d} - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) + \frac{a^2(79A-259B)\sin(c+dx)}{2d(a\cos(c+dx)+a)}$$

$$\frac{(A-B)\sin(c+dx)\cos^{\frac{7}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 223

3.208. $\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

$$3 \left(\frac{a^4(177A-637B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})} a+a} dx - \frac{128a^{7/2}(2A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{14a^3(7A-27B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}$$

$$4a^2 \qquad 8a^2 \qquad 12a^2$$

$$\frac{(A - B) \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}}$$

↓ 3261

$$3 \left(-\frac{2a^5(177A-637B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)+a}}\right) - \frac{128a^{7/2}(2A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{14a^3(7A-27B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}$$

$$4a^2 \qquad 8a^2 \qquad 12a^2$$

$$\frac{(A - B) \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}}$$

↓ 218

$$3 \left(-\frac{\sqrt{2}a^{7/2}(177A-637B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right) - \frac{128a^{7/2}(2A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{14a^3(7A-27B) \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx)+a}} \right) + \frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}}$$

$$8a^2 \qquad 4a^2 \qquad 12a^2$$

$$\frac{(A - B) \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}}$$

```
input Int[(Cos[c + d*x]^(7/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2),x]
```

```
output ((A - B)*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(6*d*(a + a*Cos[c + d*x])^(7/2))
+ ((3*a*(3*A - 7*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(4*d*(a + a*Cos[c +
d*x])^(5/2)) + ((a^2*(79*A - 259*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*
(a + a*Cos[c + d*x])^(3/2)) + (3*(-((( -128*a^(7/2)*(2*A - 7*B)*ArcSin[(Sqr
t[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]))/d + (Sqrt[2]*a^(7/2)*(177*A
- 637*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a
+ a*Cos[c + d*x]]))/d)/a - (14*a^3*(7*A - 27*B)*Sqrt[Cos[c + d*x]]*Sin[c
+ d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/(4*a^2)/(8*a^2))/(12*a^2)
```

3.208.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3253 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Co
s[e + f*x]/Sqrt[a + b*Ssin[e + f*x]])], x] /; FreeQ[{a, b, d, e, f}, x] && E
qQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.208.
$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$$


```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Ssin[e + f*x])^(m
+ 1)*(c + d*Ssin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Ssin[e + f*x]]/Sqrt[c + d*Ssin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3462 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Ssin[e + f*x])^m*(c + d*S
sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.208.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 794 vs. $2(250) = 500$.

Time = 17.28 (sec) , antiderivative size = 795, normalized size of antiderivative = 2.71

method	result	size
default	Expression too large to display	795
parts	Expression too large to display	828

```
input int(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2),x,method=_RET
URNVERBOSE)
```

$$3.208. \quad \int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{\frac{7}{2}}} dx$$

output `1/384/a^4/d*(531*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^3-1911*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^3+384*B*cos(d*x+c)^3*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+1593*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2+768*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))*cos(d*x+c)^3-494*A*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-5733*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2-2688*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))*cos(d*x+c)^3+2198*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+1593*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+2304*A*cos(d*x+c)^2*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))-724*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-5733*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)-8064*B*cos(d*x+c)^2*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))+2884*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+531*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+2304*A*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))-294*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)-1911*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-8064*B*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))+1134*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+768*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))-2688*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))*((a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^(1/2)/(1+cos(d*x+c))^4...`

3.208.5 Fracas [A] (verification not implemented)

Time = 14.79 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.26

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \frac{3\sqrt{2}((177A-637B)\cos(dx+c)^4 + 4(177A-637B)\cos(dx+c))}{(a+a\cos(c+dx))^{7/2}}$$

input `integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo rithm="fracas")`

output $\frac{1}{384}(3\sqrt{2})((177A - 637B)\cos(dx + c)^4 + 4(177A - 637B)\cos(dx + c)^3 + 6(177A - 637B)\cos(dx + c)^2 + 4(177A - 637B)\cos(dx + c) + 177A - 637B)\sqrt{a}\arctan(\sqrt{2}\sqrt{a\cos(dx + c) + a}\sqrt{\cos(dx + c)})/(\sqrt{a}\sin(dx + c))) + 2(192B\cos(dx + c)^3 - (247A - 1099B)\cos(dx + c)^2 - 2(181A - 721B)\cos(dx + c) - 147A + 567B)\sqrt{a\cos(dx + c) + a}\sqrt{\cos(dx + c)}\sin(dx + c) - 384((2A - 7B)\cos(dx + c)^4 + 4(2A - 7B)\cos(dx + c)^3 + 6(2A - 7B)\cos(dx + c)^2 + 4(2A - 7B)\cos(dx + c) + 2A - 7B)\sqrt{a}\arctan(\sqrt{a\cos(dx + c) + a}\sqrt{\cos(dx + c)})/(\sqrt{a}\sin(dx + c))))/(a^4d\cos(dx + c)^4 + 4a^4d\cos(dx + c)^3 + 6a^4d\cos(dx + c)^2 + 4a^4d\cos(dx + c) + a^4d)$

3.208.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{\frac{7}{2}}} dx = \text{Timed out}$$

input `integrate(cos(dx+c)**(7/2)*(A+B*cos(dx+c))/(a+a*cos(dx+c))**(7/2),x)`

output Timed out

3.208.7 Maxima [F]

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{\frac{7}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{7}{2}}}{(a \cos(dx + c) + a)^{\frac{7}{2}}} dx$$

input `integrate(cos(dx+c)^(7/2)*(A+B*cos(dx+c))/(a+a*cos(dx+c))^(7/2),x, algo rithm="maxima")`

output `integrate((B*cos(dx + c) + A)*cos(dx + c)^(7/2)/(a*cos(dx + c) + a)^(7/2), x)`

3.208.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(7/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="giac")`

output `Timed out`

3.208.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \int \frac{\cos(c+dx)^{7/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$$

input `int((cos(c + d*x)^(7/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(7/2),x
)`

output `int((cos(c + d*x)^(7/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(7/2),
x)`

3.209
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

3.209.1 Optimal result 1958
 3.209.2 Mathematica [A] (warning: unable to verify) 1959
 3.209.3 Rubi [A] (verified) 1959
 3.209.4 Maple [B] (warning: unable to verify) 1964
 3.209.5 Fricas [A] (verification not implemented) 1965
 3.209.6 Sympy [F(-1)] 1966
 3.209.7 Maxima [F] 1966
 3.209.8 Giac [F(-1)] 1966
 3.209.9 Mupad [F(-1)] 1967

3.209.1 Optimal result

Integrand size = 35, antiderivative size = 241

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right)}{a^{7/2}d} + \frac{(5A-177B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d} + \frac{(A-B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{6d(a+a \cos(c+dx))^{7/2}} + \frac{(5A-17B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{48ad(a+a \cos(c+dx))^{5/2}} + \frac{(5A-49B) \sqrt{\cos(c+dx)} \sin(c+dx)}{64a^2d(a+a \cos(c+dx))^{3/2}}$$

```
output 2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d+1/6*(A-B)*
cos(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)+1/48*(5*A-17*B)*cos(d
*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)+1/128*(5*A-177*B)*arctan
(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a
^(7/2)/d*2^(1/2)+1/64*(5*A-49*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d/(a+a*co
s(d*x+c))^(3/2)
```

3.209.2 Mathematica [A] (warning: unable to verify)

Time = 2.17 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.32

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx =$$

$$\sqrt{a(1+\cos(c+dx))} \left(1176B \arcsin\left(\sqrt{1-\cos(c+dx)}\right) \cos^6\left(\frac{1}{2}(c+dx)\right) + 4248B \arcsin\left(\sqrt{\cos(c+dx)}\right) \right)$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2),x]`

output `-1/192*(Sqrt[a*(1 + Cos[c + d*x])]*(1176*B*ArcSin[Sqrt[1 - Cos[c + d*x]]]*Cos[(c + d*x)/2]^6 + 4248*B*ArcSin[Sqrt[Cos[c + d*x]]]*Cos[(c + d*x)/2]^6 + 12*Sqrt[2]*(5*A - 177*B)*ArcTan[Sqrt[Cos[c + d*x]]/Sqrt[Sin[(c + d*x)/2]^2])*Cos[(c + d*x)/2]^6 - 50*A*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) + 362*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(3/2) - 67*A*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(5/2) + 247*B*Sqrt[1 - Cos[c + d*x]]*Cos[c + d*x]^(5/2) - 15*A*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])] + 147*B*Sqrt[-((-1 + Cos[c + d*x])*Cos[c + d*x])])*Sin[c + d*x])/(a^4*d*Sqrt[1 - Cos[c + d*x]]*(1 + Cos[c + d*x])^4)`

3.209.3 Rubi [A] (verified)Time = 1.52 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.08, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3456, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{7/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{7/2}} dx$$

3.209. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

$$\begin{aligned}
& \downarrow 3456 \\
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)+12aB \cos(c+dx))}{2(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}} \\
& \downarrow 27 \\
& \frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)+12aB \cos(c+dx))}{(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}} \\
& \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (5a(A-B)+12aB \sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}} \\
& \downarrow 3456 \\
& \frac{\int \frac{3\sqrt{\cos(c+dx)}((5A-17B)a^2+32B \cos(c+dx)a^2)}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}} \\
& \downarrow 27 \\
& \frac{3 \int \frac{\sqrt{\cos(c+dx)}((5A-17B)a^2+32B \cos(c+dx)a^2)}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}} \\
& \downarrow 3042 \\
& \frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}((5A-17B)a^2+32B \sin(c+dx+\frac{\pi}{2})a^2)}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}} \\
& \downarrow 3456 \\
& \frac{3 \left(\frac{\int \frac{(5A-49B)a^3+128B \cos(c+dx)a^3}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx) + a)^{7/2}}
\end{aligned}$$

3.209. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{3 \left(\frac{\int \frac{(5A-49B)a^3 + 128B \cos(c+dx)a^3}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \\
& \downarrow 3042 \\
& \frac{3 \left(\frac{\int \frac{(5A-49B)a^3 + 128B \sin(c+dx+\frac{\pi}{2})a^3}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \\
& \downarrow 3461 \\
& \frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + 128a^2 B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{4a^2} + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \\
& \downarrow 3042 \\
& \frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + 128a^2 B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a^2} + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \\
& \frac{12a^2}{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)} \\
& \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \\
& \downarrow 3253
\end{aligned}$$

3.209. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$

$$\frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})} a+a} dx - \frac{256a^2 B \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}\right)}{4a^2} \right) + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{12a^2}{6d(a \cos(c+dx)+a)^{7/2}} \right) \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}}$$

223

$$\frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{\sin(c+dx+\frac{\pi}{2})} a+a} dx + \frac{256a^{5/2} B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{4a^2} \right) + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{a(5A-17B) \sin(c+dx)}{4d(a \cos(c+dx)+a)^{3/2}}$$

3261

$$\frac{3 \left(\frac{256a^{5/2} B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{2a^4(5A-177B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}}\right)}{4a^2} \right) + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2}$$

218

$$\frac{3 \left(\frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{\sqrt{2}a^{5/2}(5A-177B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}}\right)}{4a^2} + \frac{256a^{5/2} B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} \right) + \frac{a(5A-17B) \sin(c+dx)}{4d(a \cos(c+dx)+a)^{3/2}}}{8a^2}$$

3.209. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*cos[c + d*x]))/(a + a*cos[c + d*x])^(7/2),x]`

output `((A - B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(6*d*(a + a*cos[c + d*x])^(7/2)) + ((a*(5*A - 17*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(4*d*(a + a*cos[c + d*x])^(5/2)) + (3*((256*a^(5/2)*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*cos[c + d*x]])]/d + (Sqrt[2]*a^(5/2)*(5*A - 177*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]*Sqrt[a + a*cos[c + d*x]])])/d)/(4*a^2) + (a^2*(5*A - 49*B)*Sqrt[Cos[c + d*x]*Sin[c + d*x])/(2*d*(a + a*cos[c + d*x])^(3/2))))/(8*a^2))/(12*a^2)`

3.209.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.209.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.209.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 639 vs. 2(204) = 408.

Time = 5.94 (sec) , antiderivative size = 640, normalized size of antiderivative = 2.66

method	result
parts	$A \left(67\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} (\cos^2(dx+c)) \sin(dx+c) + 50\sqrt{2} \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 15 \arcsin(\cot(dx+c) - \csc(dx+c)) (\cos^3(dx+c)) \right)$
default	$A \left(-\frac{(\csc^2(dx+c))(1-\cos(dx+c))^2-1}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \right)^{\frac{5}{2}} ((\csc^2(dx+c))(1-\cos(dx+c))^2+1)^3 \sqrt{2} \sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}} \left(8(\csc^5(dx+c)) \sqrt{-(\csc^2(dx+c))} \right)$

```
input int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2), x, method=_RET
URNVERBOSE)
```

$$3.209. \int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

output `1/384*A/d*(67*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^2*sin(d*x+c)+50*2^(1/2)*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-15*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^3+15*sin(d*x+c)*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-45*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2-45*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)-15*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^4/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)/a^4-1/384*B/d*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(7/2)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(7/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^4*(a/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(8*csc(d*x+c)^5*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^5-50*csc(d*x+c)^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^3+384*2^(1/2)*arctan(2^(1/2)*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(csc(d*x+c)-cot(d*x+c)))+189*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+c))-531*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)/a^4`

3.209.5 Fracas [A] (verification not implemented)

Time = 8.46 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.36

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx =$$

$$3\sqrt{2}((5A-177B)\cos(dx+c)^4 + 4(5A-177B)\cos(dx+c)^3 + 6(5A-177B)\cos(dx+c)^2 + 4(5A$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo rithm="fricas")`

output `-1/384*(3*sqrt(2)*((5*A - 177*B)*cos(d*x + c)^4 + 4*(5*A - 177*B)*cos(d*x + c)^3 + 6*(5*A - 177*B)*cos(d*x + c)^2 + 4*(5*A - 177*B)*cos(d*x + c) + 5*A - 177*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c))) - 2*((67*A - 247*B)*cos(d*x + c)^2 + 2*(25*A - 181*B)*cos(d*x + c) + 15*A - 147*B)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c) + 768*(B*cos(d*x + c)^4 + 4*B*cos(d*x + c)^3 + 6*B*cos(d*x + c)^2 + 4*B*cos(d*x + c) + B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c)))/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)`

3.209. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

3.209.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{7}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(7/2),x)`

output `Timed out`

3.209.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{7}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(a\cos(dx+c)+a)^{\frac{7}{2}}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^(7/
2), x)`

3.209.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{\frac{7}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="giac")`

output `Timed out`

3.209.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(7/2),x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(7/2),x)`

3.210
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

3.210.1 Optimal result 1968
 3.210.2 Mathematica [B] (verified) 1969
 3.210.3 Rubi [A] (verified) 1970
 3.210.4 Maple [B] (verified) 1973
 3.210.5 Fricas [A] (verification not implemented) 1974
 3.210.6 Sympy [F(-1)] 1974
 3.210.7 Maxima [F] 1974
 3.210.8 Giac [F(-1)] 1975
 3.210.9 Mupad [F(-1)] 1975

3.210.1 Optimal result

Integrand size = 35, antiderivative size = 201

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx = \frac{(7A+5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d} + \frac{(A-B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{6d(a+a \cos(c+dx))^{7/2}} + \frac{(A-13B) \sqrt{\cos(c+dx)} \sin(c+dx)}{48ad(a+a \cos(c+dx))^{5/2}} + \frac{(17A+67B) \sqrt{\cos(c+dx)} \sin(c+dx)}{192a^2d(a+a \cos(c+dx))^{3/2}}$$

output `1/6*(A-B)*cos(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)+1/128*(7*A+5*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d*2^(1/2)+1/48*(A-13*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(5/2)+1/192*(17*A+67*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(3/2)`

3.210.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 613 vs. $2(201) = 402$.

Time = 6.32 (sec) , antiderivative size = 613, normalized size of antiderivative = 3.05

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \frac{5B \arcsin\left(\frac{\sin(\frac{c}{2}+\frac{dx}{2})}{\sqrt{\cos^2(\frac{1}{2}(c+dx))}}\right) \cos^7\left(\frac{c}{2}+\frac{dx}{2}\right)}{8d(a(1+\cos(c+dx)))^{7/2}}$$

$$+ \frac{11B \cos^7\left(\frac{c}{2}+\frac{dx}{2}\right) \sin\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{1-\sec^2\left(\frac{1}{2}(c+dx)\right) \sin^2\left(\frac{c}{2}+\frac{dx}{2}\right)}}{8d\sqrt{\cos^2\left(\frac{1}{2}(c+dx)\right)}(a(1+\cos(c+dx)))^{7/2}}$$

$$- \frac{13B \cos^7\left(\frac{c}{2}+\frac{dx}{2}\right) \sin^3\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{1-\sec^2\left(\frac{1}{2}(c+dx)\right) \sin^2\left(\frac{c}{2}+\frac{dx}{2}\right)}}{12d \cos^2\left(\frac{1}{2}(c+dx)\right)^{3/2} (a(1+\cos(c+dx)))^{7/2}}$$

$$+ \frac{B \cos^7\left(\frac{c}{2}+\frac{dx}{2}\right) \sin^5\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{1-\sec^2\left(\frac{1}{2}(c+dx)\right) \sin^2\left(\frac{c}{2}+\frac{dx}{2}\right)}}{3d \cos^2\left(\frac{1}{2}(c+dx)\right)^{5/2} (a(1+\cos(c+dx)))^{7/2}}$$

$$+ \frac{A \cos^7\left(\frac{c}{2}+\frac{dx}{2}\right) \sec^6\left(\frac{1}{2}(c+dx)\right) \sin\left(\frac{c}{2}+\frac{dx}{2}\right) \left(27-106\sin^2\left(\frac{c}{2}+\frac{dx}{2}\right)+121\sin^4\left(\frac{c}{2}+\frac{dx}{2}\right)-34\sin^6\left(\frac{c}{2}+\frac{dx}{2}\right)\right)}{24d(a(1+\cos(c+dx)))^{7/2} \sqrt{1-2\sin^2\left(\frac{c}{2}+\frac{dx}{2}\right)}}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2), x]`

output `(5*B*ArcSin[Sin[c/2 + (d*x)/2]/Sqrt[Cos[(c + d*x)/2]^2])*Cos[c/2 + (d*x)/2]^7/(8*d*(a*(1 + Cos[c + d*x]))^(7/2)) + (11*B*Cos[c/2 + (d*x)/2]^7*Sin[c/2 + (d*x)/2]*Sqrt[1 - Sec[(c + d*x)/2]^2*Sin[c/2 + (d*x)/2]^2])/(8*d*Sqrt[Cos[(c + d*x)/2]^2]*(a*(1 + Cos[c + d*x]))^(7/2)) - (13*B*Cos[c/2 + (d*x)/2]^7*Sin[c/2 + (d*x)/2]^3*Sqrt[1 - Sec[(c + d*x)/2]^2*Sin[c/2 + (d*x)/2]^2])/(12*d*(Cos[(c + d*x)/2]^2)^(3/2)*(a*(1 + Cos[c + d*x]))^(7/2)) + (B*Cos[c/2 + (d*x)/2]^7*Sin[c/2 + (d*x)/2]^5*Sqrt[1 - Sec[(c + d*x)/2]^2*Sin[c/2 + (d*x)/2]^2])/(3*d*(Cos[(c + d*x)/2]^2)^(5/2)*(a*(1 + Cos[c + d*x]))^(7/2)) + (A*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*(27 - 106*Sin[c/2 + (d*x)/2]^2 + 121*Sin[c/2 + (d*x)/2]^4 - 34*Sin[c/2 + (d*x)/2]^6 + (21*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])*Cos[(c + d*x)/2]^6/Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]))/(24*d*(a*(1 + Cos[c + d*x]))^(7/2)*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])`

3.210. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

3.210.3 Rubi [A] (verified)

Time = 1.08 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3456, 27, 3042, 3456, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{7/2}} dx \\
 & \quad \downarrow \text{3456} \\
 & \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+2a(A+5B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+2a(A+5B)\cos(c+dx))}{(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)+2a(A+5B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3456} \\
 & \frac{\int \frac{(A-13B)a^2+18(A+3B)\cos(c+dx)a^2}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(A-13B)a^2+18(A+3B)\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.210. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

$$\frac{\int \frac{(A-13B)a^2 + 18(A+3B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 3457

$$\frac{\int \frac{3a^3(7A+5B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{8a^2} + \frac{12a^2}{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 27

$$\frac{\frac{3}{4}a(7A+5B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 3042

$$\frac{\frac{3}{4}a(7A+5B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

↓ 3261

$$\frac{\frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{3a^2(7A+5B)\int \frac{1}{\sin(c+dx)\tan(c+dx)a^3+2a^2} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{8a^2}}{12a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}$$

↓ 218

$$\frac{\frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3\sqrt{a}(7A+5B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{2\sqrt{2}d}}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}}$$

3.210. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2),x
]`

output `((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(6*d*(a + a*Cos[c + d*x])^(7/2))
+ ((a*(A - 13*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x]
])^(5/2)) + ((3*Sqrt[a]*(7*A + 5*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]
*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*d) + (a^2*(17*A
+ 67*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2))
)/(8*a^2)/(12*a^2)`

3.210.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e
.) + (f.)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Ssin[e + f*x])^(m
+ 1)*(c + d*Ssin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &&
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])`

3.210.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.210.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs. 2(170) = 340.

Time = 5.97 (sec) , antiderivative size = 475, normalized size of antiderivative = 2.36

method	result
default	$(-21A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))-15B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+34A(\cos^2(dx+c)) \sin(dx+c)-15B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+34A(\cos^2(dx+c)) \sin(dx+c))$
parts	$\frac{A(17\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} (\cos^2(dx+c)) \sin(dx+c)+70\sqrt{2} \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} -21 \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+34A(\cos^2(dx+c)) \sin(dx+c))}{(a+\cos(dx+c))^2}$

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2),x,method=_RET
URNVERBOSE)
```

```
output 1/384/a^4/d*(-21*A*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^3-15*B
*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^3+34*A*cos(d*x+c)^2*sin(
d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-63*A*2^(1/2)*arcsin(cot(d*x+c)-cs
c(d*x+c))*cos(d*x+c)^2+134*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*
x+c)))^(1/2)-45*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^2+140*A
*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-63*A*2^(1/2)*arcs
in(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+100*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+
c)/(1+cos(d*x+c)))^(1/2)-45*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*
x+c)+42*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)-21*A*2^(1/2)*arcsin
(cot(d*x+c)-csc(d*x+c))+30*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-
15*B*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)*cos(d
*x+c)^(1/2)/(1+cos(d*x+c))^4/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

$$3.210. \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

3.210.5 Fricas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.32

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \frac{3\sqrt{2}((7A+5B)\cos(dx+c)^4 + 4(7A+5B)\cos(dx+c)^3 + 6(7A+5B)\cos(dx+c)^2 + 4(7A+5B)\cos(dx+c) + 7A+5B)\sqrt{a}\arctan(1/2\sqrt{2}\sqrt{a\cos(dx+c)+a})\sqrt{a}\sqrt{\cos(dx+c)}\sin(dx+c)/(a\cos(dx+c)^2+a\cos(dx+c)) + 2((17A+67B)\cos(dx+c)^2 + 10(7A+5B)\cos(dx+c) + 21A+15B)\sqrt{a\cos(dx+c)+a}\sqrt{\cos(dx+c)}\sin(dx+c)/(a^4d\cos(dx+c)^4 + 4a^4d\cos(dx+c)^3 + 6a^4d\cos(dx+c)^2 + 4a^4d\cos(dx+c) + a^4d}}{}$$

```
input integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="fricas")
```

```
output 1/384*(3*sqrt(2)*((7*A + 5*B)*cos(d*x + c)^4 + 4*(7*A + 5*B)*cos(d*x + c)^
3 + 6*(7*A + 5*B)*cos(d*x + c)^2 + 4*(7*A + 5*B)*cos(d*x + c) + 7*A + 5*B)
*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x
+ c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) + 2*((17*A + 67*B)
*cos(d*x + c)^2 + 10*(7*A + 5*B)*cos(d*x + c) + 21*A + 15*B)*sqrt(a*cos(d*
x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^4*d*cos(d*x + c)^4 + 4*a^4
*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)
```

3.210.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(7/2),x)
```

```
output Timed out
```

3.210.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(a\cos(dx+c)+a)^{\frac{7}{2}}} dx$$

```
input integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="maxima")
```

3.210. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(7/2), x)`

3.210.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algorith="giac")`

output `Timed out`

3.210.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{\cos(c + dx)^{3/2} (A + B \cos(c + dx))}{(a + a \cos(c + dx))^{7/2}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(7/2),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + a*cos(c + d*x))^(7/2),x)`

3.211
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx$$

3.211.1 Optimal result 1976
 3.211.2 Mathematica [A] (verified) 1977
 3.211.3 Rubi [A] (verified) 1977
 3.211.4 Maple [B] (verified) 1981
 3.211.5 Fricas [A] (verification not implemented) 1981
 3.211.6 Sympy [F(-1)] 1982
 3.211.7 Maxima [F] 1982
 3.211.8 Giac [F(-1)] 1983
 3.211.9 Mupad [F(-1)] 1983

3.211.1 Optimal result

Integrand size = 35, antiderivative size = 201

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx = \frac{(13A+7B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d} + \frac{(A-B)\sqrt{\cos(c+dx)} \sin(c+dx)}{6d(a+a \cos(c+dx))^{7/2}} + \frac{(A+3B)\sqrt{\cos(c+dx)} \sin(c+dx)}{16ad(a+a \cos(c+dx))^{5/2}} - \frac{(5A-17B)\sqrt{\cos(c+dx)} \sin(c+dx)}{192a^2d(a+a \cos(c+dx))^{3/2}}$$

```
output 1/128*(13*A+7*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d*2^(1/2)+1/6*(A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(7/2)+1/16*(A+3*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(5/2)-1/192*(5*A-17*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(3/2)
```

3.211.2 Mathematica [A] (verified)

Time = 1.67 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \frac{\sec^4\left(\frac{1}{2}(c+dx)\right) \left(48(13A+7B)\operatorname{arctanh}\left(\sqrt{-\sec(c+dx)\sin^2\left(\frac{1}{2}(c+dx)\right)}\right)\right)}{(a+a\cos(c+dx))^{7/2}}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2),x]`

output `(Sec[(c + d*x)/2]^4*(48*(13*A + 7*B)*ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)])*Cos[(c + d*x)/2]^6 + Cos[c + d*x]*(73*A + 59*B + 4*(A + 35*B)*Cos[c + d*x] + (-5*A + 17*B)*Cos[2*(c + d*x)])*Sqrt[2 - 2*Sec[c + d*x]]*Tan[(c + d*x)/2])/(1536*Sqrt[2]*a^3*d*Sqrt[-1 + Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])])`

3.211.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3456, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{7/2}} dx \\ & \quad \downarrow \text{3456} \\ & \frac{\int \frac{a(A-B)+4a(A+2B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.211. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{a(A-B)+4a(A+2B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(A-B)+4a(A+2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{(11A+B)a^2+6(A+3B)\cos(c+dx)a^2}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(11A+B)a^2+6(A+3B)\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(11A+B)a^2+6(A+3B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \\
 & \quad \frac{12a^2}{6d(a\cos(c+dx)+a)^{7/2}} \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{3a^3(13A+7B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \\
 & \quad \frac{12a^2}{6d(a\cos(c+dx)+a)^{7/2}} \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\frac{3}{4}a(13A+7B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \\
 & \quad \frac{12a^2}{6d(a\cos(c+dx)+a)^{7/2}} \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.211. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$

$$\frac{\frac{3}{4}a(13A+7B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{a^2(5A-17B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B) \sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a \cos(c+dx)+a)^{5/2}} +$$

$$\frac{12a^2}{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{6d(a \cos(c+dx)+a)^{7/2}}$$

↓ 3261

$$- \frac{3a^2(13A+7B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2}}{2d} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)+a}}\right) - \frac{a^2(5A-17B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3a(A+3B) \sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a \cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{6d(a \cos(c+dx)+a)^{7/2}}$$

↓ 218

$$\frac{3\sqrt{a}(13A+7B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}d} - \frac{a^2(5A-17B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3a(A+3B) \sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a \cos(c+dx)+a)^{5/2}} +$$

$$\frac{12a^2}{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{6d(a \cos(c+dx)+a)^{7/2}}$$

```
input Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + a*Cos[c + d*x])^(7/2),x
]
```

```
output ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(6*d*(a + a*Cos[c + d*x])^(7/2))
+ ((3*a*(A + 3*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*
x])^(5/2)) + ((3*Sqrt[a]*(13*A + 7*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[
2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*d) - (a^2*(5*
A - 17*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)
))/(8*a^2))/(12*a^2)
```

3.211.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.211.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs. 2(170) = 340.

Time = 5.88 (sec) , antiderivative size = 475, normalized size of antiderivative = 2.36

method	result
default	$-\frac{(39A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+21B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+117A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c)))}{(a+\cos(dx+c))^7}$
parts	$-\frac{A\left(5\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} (\cos^2(dx+c) \sin(dx+c)-2\sqrt{2} \cos(dx+c) \sin(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} +39 \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))\right)}{(a+\cos(dx+c))^7}$

input `int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2),x,method=_RETURNVERBOSE)`

output
$$-\frac{1}{384} \frac{a^4}{d} (39A^2)^{1/2} \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)^3 + 21B^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)^3 + 117A^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)^2 + 10A \cos(dx+c)^2 \sin(dx+c) (\cos(dx+c)/(1+\cos(dx+c)))^{1/2} + 63B^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c)^2 - 34B \cos(dx+c)^2 \sin(dx+c) (\cos(dx+c)/(1+\cos(dx+c)))^{1/2} + 117A^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c) - 4A \cos(dx+c) \sin(dx+c) (\cos(dx+c)/(1+\cos(dx+c)))^{1/2} + 63B^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) \cos(dx+c) - 140B \cos(dx+c) \sin(dx+c) (\cos(dx+c)/(1+\cos(dx+c)))^{1/2} + 39A^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) - 78A (\cos(dx+c)/(1+\cos(dx+c)))^{1/2} \sin(dx+c) + 21B^2 (1/2) \arcsin(\cot(dx+c)-\csc(dx+c)) - 42B \sin(dx+c) (\cos(dx+c)/(1+\cos(dx+c)))^{1/2} (a(1+\cos(dx+c)))^{1/2} \cos(dx+c)^{1/2} / (1+\cos(dx+c))^4 / (\cos(dx+c)/(1+\cos(dx+c)))^{1/2}$$

3.211.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+a \cos(c+dx))^{7/2}} dx = \frac{3\sqrt{2}((13A+7B) \cos(dx+c)^4 + 4(13A+7B) \cos(dx+c)^3 + \dots)}{(a+a \cos(c+dx))^{7/2}}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x,algorithm="fracas")`

output $1/384*(3*\sqrt{2})*((13*A + 7*B)*\cos(dx + c)^4 + 4*(13*A + 7*B)*\cos(dx + c)^3 + 6*(13*A + 7*B)*\cos(dx + c)^2 + 4*(13*A + 7*B)*\cos(dx + c) + 13*A + 7*B)*\sqrt{a}*\arctan(1/2*\sqrt{2}*\sqrt{a*\cos(dx + c) + a}*\sqrt{a}*\sqrt{\cos(dx + c)}*\sin(dx + c)/(a*\cos(dx + c)^2 + a*\cos(dx + c))) - 2*((5*A - 17*B)*\cos(dx + c)^2 - 2*(A + 35*B)*\cos(dx + c) - 39*A - 21*B)*\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)}*\sin(dx + c)/(a^4*d*\cos(dx + c)^4 + 4*a^4*d*\cos(dx + c)^3 + 6*a^4*d*\cos(dx + c)^2 + 4*a^4*d*\cos(dx + c) + a^4*d)$

3.211.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate(cos(dx+c)**(1/2)*(A+B*cos(dx+c))/(a+a*cos(dx+c))**(7/2),x)`

output Timed out

3.211.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\cos(dx + c)}}{(a \cos(dx + c) + a)^{7/2}} dx$$

input `integrate(cos(dx+c)^(1/2)*(A+B*cos(dx+c))/(a+a*cos(dx+c))^(7/2),x, algo rithm="maxima")`

output `integrate((B*cos(dx + c) + A)*sqrt(cos(dx + c))/(a*cos(dx + c) + a)^(7/2), x)`

3.211.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2),x, algorith="giac")`

output `Timed out`

3.211.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+a\cos(c+dx))^{7/2}} dx$$

input `int((cos(c+d*x)^(1/2)*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(7/2),x)`

output `int((cos(c+d*x)^(1/2)*(A+B*cos(c+d*x)))/(a+a*cos(c+d*x))^(7/2),x)`

$$3.212 \quad \int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{7/2}} dx$$

3.212.1 Optimal result 1984
 3.212.2 Mathematica [A] (verified) 1985
 3.212.3 Rubi [A] (verified) 1985
 3.212.4 Maple [B] (verified) 1989
 3.212.5 Fracas [A] (verification not implemented) 1989
 3.212.6 Sympy [F(-1)] 1990
 3.212.7 Maxima [F] 1990
 3.212.8 Giac [F(-1)] 1991
 3.212.9 Mupad [F(-1)] 1991

3.212.1 Optimal result

Integrand size = 35, antiderivative size = 203

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \frac{(63A + 13B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d} - \frac{(A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2}} - \frac{(5A - B)\sqrt{\cos(c + dx)} \sin(c + dx)}{16ad(a + a \cos(c + dx))^{5/2}} - \frac{(103A + 5B)\sqrt{\cos(c + dx)} \sin(c + dx)}{192a^2d(a + a \cos(c + dx))^{3/2}}$$

```
output 1/128*(63*A+13*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(
a+a*cos(d*x+c))^(1/2))/a^(7/2)/d*2^(1/2)-1/6*(A-B)*sin(d*x+c)*cos(d*x+c)^(
1/2)/d/(a+a*cos(d*x+c))^(7/2)-1/16*(5*A-B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/d
/(a+a*cos(d*x+c))^(5/2)-1/192*(103*A+5*B)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/
d/(a+a*cos(d*x+c))^(3/2)
```

3.212.2 Mathematica [A] (verified)

Time = 1.59 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.83

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \frac{\sec^4\left(\frac{1}{2}(c + dx)\right) \left(-48(63A + 13B)\operatorname{arctanh}\left(\sqrt{-\sec(c + dx) \sin^2\left(\frac{1}{2}(c + dx)\right)}\right) \cos^6\left(\frac{1}{2}(c + dx)\right) + \cos(c + dx)\right)}{1536\sqrt{2}a^3d\sqrt{-1 + \cos(c + dx)}}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(7/2)),x]`

output `-1/1536*(Sec[(c + d*x)/2]^4*(-48*(63*A + 13*B)*ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]]*Cos[(c + d*x)/2]^6 + Cos[c + d*x]*(493*A - 73*B + (532*A - 4*B)*Cos[c + d*x] + (103*A + 5*B)*Cos[2*(c + d*x)])*Sqrt[2 - 2*Sec[c + d*x]])*Tan[(c + d*x)/2])/(Sqrt[2]*a^3*d*Sqrt[-1 + Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])])`

3.212.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{7/2}} dx \\ & \quad \downarrow \text{3457} \\ & \frac{\int \frac{a(11A+B) - 4a(A-B) \cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} - \frac{(A - B) \sin(c + dx) \sqrt{\cos(c + dx)}}{6d(a \cos(c + dx) + a)^{7/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.212. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{7/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{a(11A+B)-4a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(11A+B)-4a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{a^2(73A+11B)-6a^2(5A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a^2(73A+11B)-6a^2(5A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(73A+11B)-6a^2(5A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} - \\
& \quad \frac{12a^2}{6d(a\cos(c+dx)+a)^{7/2}} \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{3457} \\
& \frac{\int \frac{3a^3(63A+13B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \quad \frac{12a^2}{6d(a\cos(c+dx)+a)^{7/2}} \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\frac{3}{4}a(63A+13B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \\
& \quad \frac{12a^2}{6d(a\cos(c+dx)+a)^{7/2}} \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.212. $\int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a\cos(c+dx))^{7/2}} dx$

$$\frac{\frac{3}{4}a(63A+13B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{a^2(103A+5B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a \cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{6d(a \cos(c+dx)+a)^{7/2}}$$

↓ 3261

$$-\frac{3a^2(63A+13B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2d} - \frac{a^2(103A+5B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{3a(5A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a \cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{6d(a \cos(c+dx)+a)^{7/2}}$$

↓ 218

$$\frac{3\sqrt{a}(63A+13B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}d} - \frac{a^2(103A+5B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} - \frac{3a(5A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a \cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}} \frac{1}{6d(a \cos(c+dx)+a)^{7/2}}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(7/2)),x]`

output `-1/6*((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(a + a*Cos[c + d*x])^(7/2)) + ((-3*a*(5*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2))) + ((3*Sqrt[a]*(63*A + 13*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*d) - (a^2*(103*A + 5*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2)/(12*a^2)`

3.212.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.212.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs. $2(172) = 344$.

Time = 8.15 (sec) , antiderivative size = 475, normalized size of antiderivative = 2.34

method	result
default	$-\frac{(189A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+39B\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c))+567A\sqrt{2} \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^3(dx+c)))}{A\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}} \sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}} \left(8(\csc^5(dx+c))\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}(1-\cos(dx+c))\right)$
parts	

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(7/2),x,method=_RETURNVERBOSE)`

output
$$-1/384/a^4/d*(189*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^3+39*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^3+567*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2+206*A*\cos(d*x+c)^2*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+117*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2+10*B*\cos(d*x+c)^2*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+567*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)+532*A*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+117*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)-4*B*\cos(d*x+c)*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+189*A*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))+390*A*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\sin(d*x+c)+39*B*2^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-78*B*\sin(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1+\cos(d*x+c)))^{(1/2)}/(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}/(1+\cos(d*x+c))^4$$

3.212.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.31

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \frac{3\sqrt{2}((63A + 13B) \cos(dx + c))^4 + 4(63A + 13B) \cos(dx + c)}{\dots}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(7/2),x,algorith="fracas")`

3.212.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+a \cos(c+dx))^{7/2}} dx$$

output $1/384*(3*\sqrt{2})*((63*A + 13*B)*\cos(dx + c)^4 + 4*(63*A + 13*B)*\cos(dx + c)^3 + 6*(63*A + 13*B)*\cos(dx + c)^2 + 4*(63*A + 13*B)*\cos(dx + c) + 63*A + 13*B)*\sqrt{a}*\arctan(1/2*\sqrt{2}*\sqrt{a*\cos(dx + c) + a}*\sqrt{a}*\sqrt{t(\cos(dx + c))*\sin(dx + c)/(a*\cos(dx + c)^2 + a*\cos(dx + c))}) - 2*((103*A + 5*B)*\cos(dx + c)^2 + 2*(133*A - B)*\cos(dx + c) + 195*A - 39*B)*\sqrt{t(a*\cos(dx + c) + a)*\sqrt{\cos(dx + c))*\sin(dx + c)}/(a^4*d*\cos(dx + c)^4 + 4*a^4*d*\cos(dx + c)^3 + 6*a^4*d*\cos(dx + c)^2 + 4*a^4*d*\cos(dx + c) + a^4*d)$

3.212.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(7/2),x)`

output Timed out

3.212.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{7/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*sqrt(cos(d*x + c))), x)`

3.212.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="giac")`

output `Timed out`

3.212.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(7/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + a*cos(c + d*x))^(7/2)),
x)`

3.213
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$$

3.213.1 Optimal result	1992
3.213.2 Mathematica [C] (warning: unable to verify)	1993
3.213.3 Rubi [A] (verified)	1994
3.213.4 Maple [B] (verified)	1998
3.213.5 Fracas [A] (verification not implemented)	1999
3.213.6 Sympy [F(-1)]	2000
3.213.7 Maxima [F(-1)]	2000
3.213.8 Giac [F]	2000
3.213.9 Mupad [F(-1)]	2001

3.213.1 Optimal result

Integrand size = 35, antiderivative size = 250

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx =$$

$$\frac{3(121A - 21B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d}$$

$$- \frac{(A - B) \sin(c + dx)}{6d\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{7/2}} - \frac{(19A - 7B) \sin(c + dx)}{48ad\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{5/2}}$$

$$- \frac{(199A - 43B) \sin(c + dx)}{192a^2d\sqrt{\cos(c + dx)}(a + a \cos(c + dx))^{3/2}}$$

$$+ \frac{(691A - 103B) \sin(c + dx)}{192a^3d\sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}$$

output

```
-3/128*(121*A-21*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)
/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d*2^(1/2)-1/6*(A-B)*sin(d*x+c)/d/(a+a*cos
(d*x+c))^(7/2)/cos(d*x+c)^(1/2)-1/48*(19*A-7*B)*sin(d*x+c)/a/d/(a+a*cos(d*
x+c))^(5/2)/cos(d*x+c)^(1/2)-1/192*(199*A-43*B)*sin(d*x+c)/a^2/d/(a+a*cos(
d*x+c))^(3/2)/cos(d*x+c)^(1/2)+1/192*(691*A-103*B)*sin(d*x+c)/a^3/d/cos(d*
x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)
```

3.213.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.91 (sec) , antiderivative size = 798, normalized size of antiderivative = 3.19

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx =$$

$$\frac{B \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^6\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \left(141 - 518 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 575 \sin^4\left(\frac{c}{2} + \frac{dx}{2}\right) - 206 \sin^6\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{24d(a(1 + \cos(c + dx)))^{7/2} \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} +$$

$$\frac{2A \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^6\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{16 \cos^8\left(\frac{1}{2}(c + dx)\right) {}_5F_4\left(2, 2, 2, 2, \frac{5}{2}; 1, 1, 1, \frac{13}{2}; \frac{\sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{3465(-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right))}\right)}{}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(7/2)),x]`

output

```
-1/24*(B*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*(141 -
518*Sin[c/2 + (d*x)/2]^2 + 575*Sin[c/2 + (d*x)/2]^4 - 206*Sin[c/2 + (d*x)
/2]^6 - (189*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2
^2)])]*Cos[(c + d*x)/2]^6/Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d
*x)/2]^2)])))/(d*(a*(1 + Cos[c + d*x]))^(7/2)*Sqrt[1 - 2*Sin[c/2 + (d*x)/2
]^2]) + (2*A*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*((
16*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 5/2}, {1, 1, 1, 13/2}
, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2
)/(3465*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) - (Csc[c/2 + (d*x)/2]^10*(1 - 2*Sin
[c/2 + (d*x)/2]^2)^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^
2)]*(105*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]
*Cos[(c + d*x)/2]^6*(2187 - 12908*Sin[c/2 + (d*x)/2]^2 + 27986*Sin[c/2 + (
d*x)/2]^4 - 26380*Sin[c/2 + (d*x)/2]^6 + 8752*Sin[c/2 + (d*x)/2]^8) + Sqrt
[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-229635 + 2120790*Si
n[c/2 + (d*x)/2]^2 - 8267707*Sin[c/2 + (d*x)/2]^4 + 17646926*Sin[c/2 + (d*
x)/2]^6 - 22251094*Sin[c/2 + (d*x)/2]^8 + 16548816*Sin[c/2 + (d*x)/2]^10 -
6712984*Sin[c/2 + (d*x)/2]^12 + 1144608*Sin[c/2 + (d*x)/2]^14))/1680))/(
d*(a*(1 + Cos[c + d*x]))^(7/2)*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2))
```

3.213. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$

3.213.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{7/2}} dx \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{a(13A - B) - 6a(A - B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)^{5/2}} dx}{6a^2} - \frac{(A - B) \sin(c + dx)}{6d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{7/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(13A - B) - 6a(A - B) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)^{5/2}} dx}{12a^2} - \frac{(A - B) \sin(c + dx)}{6d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{7/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(13A - B) - 6a(A - B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(\sin\left(c + dx + \frac{\pi}{2}\right)a + a\right)^{5/2}} dx}{12a^2} - \frac{(A - B) \sin(c + dx)}{6d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{7/2}} \\
 & \quad \downarrow \text{3457} \\
 & \frac{\int \frac{3a^2(41A - 5B) - 4a^2(19A - 7B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)^{3/2}} dx}{4a^2} - \frac{a(19A - 7B) \sin(c + dx)}{4d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \frac{12a^2}{(A - B) \sin(c + dx)} \\
 & \quad \frac{(A - B) \sin(c + dx)}{6d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{7/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3a^2(41A - 5B) - 4a^2(19A - 7B) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(\cos(c + dx)a + a)^{3/2}} dx}{8a^2} - \frac{a(19A - 7B) \sin(c + dx)}{4d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{5/2}} \\
 & \quad \frac{12a^2}{(A - B) \sin(c + dx)} \\
 & \quad \frac{(A - B) \sin(c + dx)}{6d \sqrt{\cos(c + dx)}(a \cos(c + dx) + a)^{7/2}}
 \end{aligned}$$

3.213. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx$

$$\begin{array}{c}
\downarrow 3042 \\
\frac{\int \frac{3a^2(41A-5B)-4a^2(19A-7B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{3/2}} dx}{8a^2} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
\hline
\frac{12a^2}{(A-B)\sin(c+dx)} \\
\frac{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}{\downarrow 3457} \\
\frac{\int \frac{a^3(691A-103B)-2a^3(199A-43B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)}a+a} dx}{2a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
\hline
\frac{12a^2}{(A-B)\sin(c+dx)} \\
\frac{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}{\downarrow 27} \\
\frac{\int \frac{a^3(691A-103B)-2a^3(199A-43B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)}a+a} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
\hline
\frac{12a^2}{(A-B)\sin(c+dx)} \\
\frac{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}{\downarrow 3042} \\
\frac{\int \frac{a^3(691A-103B)-2a^3(199A-43B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
\hline
\frac{12a^2}{(A-B)\sin(c+dx)} \\
\frac{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}{\downarrow 3463} \\
\frac{2\int -\frac{9a^4(121A-21B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx}{4a^2} + \frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \\
\hline
\frac{12a^2}{(A-B)\sin(c+dx)} \\
\frac{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}{\downarrow 27}
\end{array}$$

3.213. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{7/2}} dx$

$$\frac{\frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 9a^3(121A-21B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^5}$$

$$\frac{(A-B)\sin(c+dx)12a^2}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}$$

↓ 3042

$$\frac{\frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 9a^3(121A-21B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^5}$$

$$\frac{(A-B)\sin(c+dx)12a^2}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}$$

↓ 3261

$$\frac{18a^4(121A-21B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} + \frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^5}$$

$$\frac{(A-B)\sin(c+dx)12a^2}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}$$

↓ 218

$$\frac{\frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{9\sqrt{2}a^{5/2}(121A-21B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{4a^2}}{8a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^5}$$

$$\frac{(A-B)\sin(c+dx)12a^2}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}}$$

```
input Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(7/2)),x
]
```

3.213. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a\cos(c+dx))^{7/2}} dx$

```
output -1/6*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*cos[c + d*x])^(7/
2)) + (-1/4*(a*(19*A - 7*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*cos
[c + d*x])^(5/2)) + (-1/2*(a^2*(199*A - 43*B)*Sin[c + d*x])/(d*Sqrt[Cos[c
+ d*x]]*(a + a*cos[c + d*x])^(3/2)) + ((-9*Sqrt[2]*a^(5/2)*(121*A - 21*B)*
ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c
+ d*x]])])/d + (2*a^3*(691*A - 103*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]
*Sqrt[a + a*cos[c + d*x]])/(4*a^2)/(8*a^2)/(12*a^2)
```

3.213.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e
_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x])/(Sqrt[a + b*Ssin[e + f*x])*Sqrt[c + d*S
in[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d)), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[B*(a*c*m +
b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.213.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 602 vs. 2(213) = 426.

Time = 8.03 (sec) , antiderivative size = 603, normalized size of antiderivative = 2.41

method	result
parts	$\frac{A \left(1089 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c) - \csc(dx+c)) (\cos^4(dx+c) + 691\sqrt{2} (\cos^3(dx+c) \sin(dx+c) + 4356 (\cos^3(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right. \right.$
default	$\frac{A \left(-(\csc^2(dx+c) (1 - \cos(dx+c))^2 + 1) \right)^{\frac{3}{2}} \sqrt{2} \sqrt{\frac{a}{(\csc^2(dx+c) (1 - \cos(dx+c))^2 + 1)}} \left(-8 (\csc^7(dx+c)) \sqrt{-(\csc^2(dx+c) (1 - \cos(dx+c))^2 + 1} (1 - \cos(dx+c)) \right. \right.$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(7/2),x,method=_RET
URNVERBOSE)
```

$$3.213. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$$

output `1/384*A/d*(1089*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^4+691*2^(1/2)*cos(d*x+c)^3*sin(d*x+c)+4356*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+1874*2^(1/2)*cos(d*x+c)^2*sin(d*x+c)+6534*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+1599*sin(d*x+c)*cos(d*x+c)*2^(1/2)+4356*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+384*2^(1/2)*sin(d*x+c)+1089*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c)))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^4/cos(d*x+c)^(1/2)*2^(1/2)/a^4-1/384*B/d/(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(a/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(8*csc(d*x+c)^5*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^5+46*csc(d*x+c)^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^3+141*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+c))+189*arcsin(cot(d*x+c)-csc(d*x+c)))*2^(1/2)/a^4`

3.213.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.19

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx =$$

$$9\sqrt{2}((121A - 21B) \cos(dx + c)^5 + 4(121A - 21B) \cos(dx + c)^4 + 6(121A - 21B) \cos(dx + c)^3 + 4(121A - 21B) \cos(dx + c)^2 + (121A - 21B) \cos(dx + c)) \sqrt{a} \arctan\left(\frac{1}{2}\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{a \cos(dx + c)} \sin(dx + c)\right) - 2((691A - 103B) \cos(dx + c)^3 + 2(937A - 133B) \cos(dx + c)^2 + 39(41A - 5B) \cos(dx + c) + 384A) \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)} \sin(dx + c) / (a^4 d \cos(dx + c)^5 + 4a^4 d \cos(dx + c)^4 + 6a^4 d \cos(dx + c)^3 + 4a^4 d \cos(dx + c)^2 + a^4 d \cos(dx + c))$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="fracas")`

output `-1/384*(9*sqrt(2))*((121*A - 21*B)*cos(d*x + c)^5 + 4*(121*A - 21*B)*cos(d*x + c)^4 + 6*(121*A - 21*B)*cos(d*x + c)^3 + 4*(121*A - 21*B)*cos(d*x + c)^2 + (121*A - 21*B)*cos(d*x + c))*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) - 2*((691*A - 103*B)*cos(d*x + c)^3 + 2*(937*A - 133*B)*cos(d*x + c)^2 + 39*(41*A - 5*B)*cos(d*x + c) + 384*A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a^4*d*cos(d*x + c)^5 + 4*a^4*d*cos(d*x + c)^4 + 6*a^4*d*cos(d*x + c)^3 + 4*a^4*d*cos(d*x + c)^2 + a^4*d*cos(d*x + c))`

3.213. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$

3.213.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(7/2),x)`

output `Timed out`

3.213.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="maxima")`

output `Timed out`

3.213.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{7}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*cos(d*x + c)^(3/2)), x)`

3.213.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2}(a + a \cos(c + dx))^{7/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(7/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + a*cos(c + d*x))^(7/2)),x)`

3.214
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$$

3.214.1 Optimal result	2002
3.214.2 Mathematica [C] (warning: unable to verify)	2003
3.214.3 Rubi [A] (verified)	2003
3.214.4 Maple [B] (verified)	2009
3.214.5 Fracas [A] (verification not implemented)	2010
3.214.6 Sympy [F(-1)]	2011
3.214.7 Maxima [F(-1)]	2011
3.214.8 Giac [F]	2011
3.214.9 Mupad [F(-1)]	2012

3.214.1 Optimal result

Integrand size = 35, antiderivative size = 297

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \frac{(1015A - 363B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right)}{64\sqrt{2}a^{7/2}d} - \frac{(A - B) \sin(c + dx)}{6d \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} - \frac{(23A - 11B) \sin(c + dx)}{48ad \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{5/2}} - \frac{(109A - 41B) \sin(c + dx)}{64a^2d \cos^{\frac{3}{2}}(c + dx)(a + a \cos(c + dx))^{3/2}} + \frac{(579A - 199B) \sin(c + dx)}{192a^3d \cos^{\frac{3}{2}}(c + dx)\sqrt{a + a \cos(c + dx)}} - \frac{(1887A - 691B) \sin(c + dx)}{192a^3d \sqrt{\cos(c + dx)}\sqrt{a + a \cos(c + dx)}}$$

output

```
-1/6*(A-B)*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2)-1/48*(23*A-11*B)*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2)-1/64*(109*A-41*B)*sin(d*x+c)/a^2/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2)+1/128*(1015*A-363*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))/a^(7/2)/d*2^(1/2)+1/192*(579*A-199*B)*sin(d*x+c)/a^3/d/cos(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)-1/192*(1887*A-691*B)*sin(d*x+c)/a^3/d/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2)
```

3.214.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.44 (sec) , antiderivative size = 1256, normalized size of antiderivative = 4.23

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(7/2)),x]`

output `(2*B*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*((16*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 5/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2)/(3465*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) - (Csc[c/2 + (d*x)/2]^10*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(105*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Cos[(c + d*x)/2]^6*(2187 - 12908*Sin[c/2 + (d*x)/2]^2 + 27986*Sin[c/2 + (d*x)/2]^4 - 26380*Sin[c/2 + (d*x)/2]^6 + 8752*Sin[c/2 + (d*x)/2]^8) + Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-229635 + 2120790*Sin[c/2 + (d*x)/2]^2 - 8267707*Sin[c/2 + (d*x)/2]^4 + 17646926*Sin[c/2 + (d*x)/2]^6 - 22251094*Sin[c/2 + (d*x)/2]^8 + 16548816*Sin[c/2 + (d*x)/2]^10 - 6712984*Sin[c/2 + (d*x)/2]^12 + 1144608*Sin[c/2 + (d*x)/2]^14))/1680))/(d*(a*(1 + Cos[c + d*x]))^(7/2)*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (A*Cot[c/2 + (d*x)/2]^7*Csc[c/2 + (d*x)/2]^4*Sec[(c + d*x)/2]^6*(-7680*Cos[(c + d*x)/2]^10*HypergeometricPFQ[{2, 2, 2, 2, 2, 7/2}, {1, 1, 1, 1, 15/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14 + 19200*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 7/2}, {1, 1, 1, 15/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14*(-7 + 6*Sin[c/2 + (d*x)/2]^2) + 143*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(315*ArcTanh[Sqrt[Sin[c/...`

3.214.3 Rubi [A] (verified)

Time = 1.80 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.09, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3457, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.214. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$

$$\downarrow \text{3457}$$

$$3 \left(\frac{\int \frac{a^3(579A-199B)-4a^3(109A-41B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B)\sin(c+dx)} \frac{1}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

$$\downarrow \text{27}$$

$$3 \left(\frac{\int \frac{a^3(579A-199B)-4a^3(109A-41B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B)\sin(c+dx)} \frac{1}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

$$\downarrow \text{3042}$$

$$3 \left(\frac{\int \frac{a^3(579A-199B)-4a^3(109A-41B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B)\sin(c+dx)} \frac{1}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

$$\downarrow \text{3463}$$

$$3 \left(\frac{2 \int -\frac{a^4(1887A-691B)-2a^4(579A-199B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}}$$

$$\frac{12a^2}{(A-B)\sin(c+dx)} \frac{1}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

$$\downarrow \text{27}$$

3.214. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a\cos(c+dx))^{7/2}} dx$

$$\frac{3 \left(\frac{2a^3(579A-199B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^4(1887A-691B)-2a^4(579A-199B) \cos(c+dx) dx}{\cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}}}{4a^2} - \frac{a^2(109A-41B) \sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-11B) \sin(c+dx)}{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

$$\frac{(A-B) \sin(c+dx) 12a^2}{6d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

↓ 3042

$$\frac{3 \left(\frac{2a^3(579A-199B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^4(1887A-691B)-2a^4(579A-199B) \sin(c+dx+\frac{\pi}{2}) dx}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}}{4a^2} - \frac{a^2(109A-41B) \sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-11B) \sin(c+dx)}{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

$$\frac{(A-B) \sin(c+dx) 12a^2}{6d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

↓ 3463

$$\frac{3 \left(\frac{2a^3(579A-199B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2 \int -\frac{3a^5(1015A-363B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{2a^4(1887A-691B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}}{4a^2} - \frac{a^2(109A-41B) \sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-11B) \sin(c+dx)}{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

$$\frac{(A-B) \sin(c+dx) 12a^2}{6d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

↓ 27

$$\frac{3 \left(\frac{2a^3(579A-199B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\frac{2a^4(1887A-691B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - 3a^4(1015A-363B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a^2(109A-41B) \sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-11B) \sin(c+dx)}{4d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

$$\frac{(A-B) \sin(c+dx) 12a^2}{6d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)^{7/2}}$$

3.214. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$

↓ 3042

$$3 \left(\frac{\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 3a^4(1015A-363B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx}{4a^2} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} \right)$$

$$\frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

12a²

↓ 3261

$$3 \left(\frac{\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{6a^5(1015A-363B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d \left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)+a}} \right)}{4a^2} + \frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right)$$

$$\frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

12a²

↓ 218

$$3 \left(\frac{\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}a^{7/2}(1015A-363B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{d\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{3a}}{4a^2} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} \right)$$

$$\frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}}$$

12a²

```
input Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(7/2)),x
]
```

```
output -1/6*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(7/
2)) + (-1/4*(a*(23*A - 11*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Co
s[c + d*x])^(5/2)) + (3*(-1/2*(a^2*(109*A - 41*B)*Sin[c + d*x])/(d*Cos[c +
d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a^3*(579*A - 199*B)*Sin[c +
d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*a^(
7/2)*(1015*A - 363*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c +
d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^4*(1887*A - 691*B)*Sin[c + d*x]
)/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]))/(3*a)/(4*a^2))/(8*a^2
))/(12*a^2)
```

3.214.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e
_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d)), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.214.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$$

```
rule 3463 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.214.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 666 vs. 2(254) = 508.

Time = 7.90 (sec) , antiderivative size = 667, normalized size of antiderivative = 2.25

method	result
default	$-\frac{(3045A\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))-1089B\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c)))}{A(3045\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))+1887\sqrt{2}(\cos^4(dx+c))\sin(dx+c)+12180\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c)))}$
parts	

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(7/2),x,method=_RET
URNVERBOSE)
```

$$3.214. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+a \cos(c+dx))^{7/2}} dx$$

output

```
-1/384/a^4/d*(3045*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5-1089*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5+12180*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^4-4356*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^4+18270*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-6534*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+12180*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))+3774*A*cos(d*x+c)^4*sin(d*x+c)-4356*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))-1382*B*cos(d*x+c)^4*sin(d*x+c)+3045*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)+10164*A*cos(d*x+c)^3*sin(d*x+c)-1089*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*arcsin(cot(d*x+c)-csc(d*x+c))*2^(1/2)-3748*B*sin(d*x+c)*cos(d*x+c)^3+8502*A*sin(d*x+c)*cos(d*x+c)^2-3198*B*sin(d*x+c)*cos(d*x+c)^2+1792*A*sin(d*x+c)*cos(d*x+c)-768*B*sin(d*x+c)*cos(d*x+c)-256*A*sin(d*x+c)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^4/cos(d*x+c)^(3/2)
```

3.214.5 Fracas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.07

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \frac{3\sqrt{2}((1015A - 363B)\cos(dx + c)^6 + 4(1015A - 363B)\cos(dx + c)^5 + 6(1015A - 363B)\cos(dx + c)^4 + 4(1015A - 363B)\cos(dx + c)^3 + (1015A - 363B)\cos(dx + c)^2)\sqrt{a}\arctan(1/2\sqrt{2})\sqrt{a\cos(dx + c) + a}\sqrt{a}\sqrt{\cos(dx + c)}\sin(dx + c)/(a\cos(dx + c)^2 + a\cos(dx + c)) - 2((1887A - 691B)\cos(dx + c)^4 + 2(2541A - 937B)\cos(dx + c)^3 + 39(109A - 41B)\cos(dx + c)^2 + 128(7A - 3B)\cos(dx + c) - 128A)\sqrt{a\cos(dx + c) + a}\sqrt{\cos(dx + c)}\sin(dx + c))/(a^4d\cos(dx + c)^6 + 4a^4d\cos(dx + c)^5 + 6a^4d\cos(dx + c)^4 + 4a^4d\cos(dx + c)^3 + a^4d\cos(dx + c)^2)}{1}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(7/2),x, algo rithm="fricas")`

output

```
1/384*(3*sqrt(2)*((1015*A - 363*B)*cos(d*x + c)^6 + 4*(1015*A - 363*B)*cos(d*x + c)^5 + 6*(1015*A - 363*B)*cos(d*x + c)^4 + 4*(1015*A - 363*B)*cos(d*x + c)^3 + (1015*A - 363*B)*cos(d*x + c)^2)*sqrt(a)*arctan(1/2*sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(a)*sqrt(cos(d*x + c))*sin(d*x + c)/(a*cos(d*x + c)^2 + a*cos(d*x + c))) - 2*((1887*A - 691*B)*cos(d*x + c)^4 + 2*(2541*A - 937*B)*cos(d*x + c)^3 + 39*(109*A - 41*B)*cos(d*x + c)^2 + 128*(7*A - 3*B)*cos(d*x + c) - 128*A)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))*sin(d*x + c))/(a^4*d*cos(d*x + c)^6 + 4*a^4*d*cos(d*x + c)^5 + 6*a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + a^4*d*cos(d*x + c)^2)
```

3.214.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(7/2),x)`

output `Timed out`

3.214.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="maxima")`

output `Timed out`

3.214.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{7}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*cos(d*x + c)^(5/2)), x)`

3.214.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + a \cos(c + dx))^{7/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + a \cos(c + dx))^{7/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(7/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + a*cos(c + d*x))^(7/2)),x)`

3.215 $\int \cos^2(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$

3.215.1 Optimal result	2013
3.215.2 Mathematica [A] (verified)	2013
3.215.3 Rubi [A] (verified)	2014
3.215.4 Maple [A] (verified)	2017
3.215.5 Fricas [A] (verification not implemented)	2017
3.215.6 Sympy [B] (verification not implemented)	2018
3.215.7 Maxima [A] (verification not implemented)	2018
3.215.8 Giac [A] (verification not implemented)	2019
3.215.9 Mupad [B] (verification not implemented)	2019

3.215.1 Optimal result

Integrand size = 29, antiderivative size = 105

$$\int \cos^2(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{1}{8}(4aA+3bB)x + \frac{(Ab+aB) \sin(c+dx)}{d} + \frac{(4aA+3bB) \cos(c+dx) \sin(c+dx)}{8d}$$

$$+ \frac{bB \cos^3(c+dx) \sin(c+dx)}{4d} - \frac{(Ab+aB) \sin^3(c+dx)}{3d}$$

```
output 1/8*(4*A*a+3*B*b)*x+(A*b+B*a)*sin(d*x+c)/d+1/8*(4*A*a+3*B*b)*cos(d*x+c)*sin(d*x+c)/d+1/4*b*B*cos(d*x+c)^3*sin(d*x+c)/d-1/3*(A*b+B*a)*sin(d*x+c)^3/d
```

3.215.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.87

$$\int \cos^2(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{48aAc+36bBc+48aAdx+36bBdx+96(Ab+aB) \sin(c+dx)-32(Ab+aB) \sin^3(c+dx)+24(aA+}{96d}$$

```
input Integrate[Cos[c + d*x]^2*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]
```

output $(48*a*A*c + 36*b*B*c + 48*a*A*d*x + 36*b*B*d*x + 96*(A*b + a*B)*\text{Sin}[c + d*x] - 32*(A*b + a*B)*\text{Sin}[c + d*x]^3 + 24*(a*A + b*B)*\text{Sin}[2*(c + d*x)] + 3*b*B*\text{Sin}[4*(c + d*x)])/(96*d)$

3.215.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.92, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 3447, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\ & \quad \downarrow 3042 \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow 3447 \\ & \int \cos^2(c + dx) \left((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)\right) dx \\ & \quad \downarrow 3042 \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left((aB + Ab) \sin\left(c + dx + \frac{\pi}{2}\right) + aA + bB \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\ & \quad \downarrow 3502 \\ & \frac{1}{4} \int \cos^2(c + dx) (4aA + 3bB + 4(Ab + aB) \cos(c + dx)) dx + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d} \\ & \quad \downarrow 3042 \\ & \frac{1}{4} \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(4aA + 3bB + 4(Ab + aB) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\ & \quad \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d} \\ & \quad \downarrow 3227 \\ & \frac{1}{4} \left(4(aB + Ab) \int \cos^3(c + dx) dx + (4aA + 3bB) \int \cos^2(c + dx) dx\right) + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d} \\ & \quad \downarrow 3042 \end{aligned}$$

3.215. $\int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\frac{1}{4} \left((4aA + 3bB) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + 4(aB + Ab) \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d}$$

↓ 3113

$$\frac{1}{4} \left((4aA + 3bB) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4(aB + Ab) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right) + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d}$$

↓ 2009

$$\frac{1}{4} \left((4aA + 3bB) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4(aB + Ab) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d}$$

↓ 3115

$$\frac{1}{4} \left((4aA + 3bB) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{4(aB + Ab) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d}$$

↓ 24

$$\frac{1}{4} \left((4aA + 3bB) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) - \frac{4(aB + Ab) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right) + \frac{bB \sin(c + dx) \cos^3(c + dx)}{4d}$$

input `Int[Cos[c + d*x]^2*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(b*B*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + ((4*a*A + 3*b*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (4*(A*b + a*B)*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/4`

3.215.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3113 `Int[sin[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.215.4 Maple [A] (verified)

Time = 2.96 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.82

method	result
parallelrisch	$\frac{(24aA+24Bb) \sin(2dx+2c)+(8Ab+8Ba) \sin(3dx+3c)+3Bb \sin(4dx+4c)+(72Ab+72Ba) \sin(dx+c)+48\left(aA+\frac{3Bb}{4}\right)x d}{96d}$
parts	$\frac{(Ab+Ba)(2+\cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{aA\left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{Bb\left(\frac{(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}) \sin(dx+c)}{4}\right)}{d}$
derivativedivides	$\frac{Bb\left(\frac{(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ab(2+\cos^2(dx+c)) \sin(dx+c)}{3} + \frac{Ba(2+\cos^2(dx+c)) \sin(dx+c)}{3} + aA\left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
default	$\frac{Bb\left(\frac{(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ab(2+\cos^2(dx+c)) \sin(dx+c)}{3} + \frac{Ba(2+\cos^2(dx+c)) \sin(dx+c)}{3} + aA\left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
risch	$\frac{xaA}{2} + \frac{3bBx}{8} + \frac{3 \sin(dx+c)Ab}{4d} + \frac{3aB \sin(dx+c)}{4d} + \frac{Bb \sin(4dx+4c)}{32d} + \frac{\sin(3dx+3c)Ab}{12d} + \frac{\sin(3dx+3c)Ba}{12d} + s$
norman	$\frac{\left(\frac{aA}{2} + \frac{3Bb}{8}\right)x + \left(2aA + \frac{3Bb}{2}\right)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(2aA + \frac{3Bb}{2}\right)x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(3aA + \frac{9Bb}{4}\right)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \left(\frac{aA}{2} + \frac{3Bb}{8}\right)x}{24d}$

input `int(cos(d*x+c)^2*(a+cos(d*x+c)*b)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/96*((24*A*a+24*B*b)*sin(2*d*x+2*c)+(8*A*b+8*B*a)*sin(3*d*x+3*c)+3*B*b*sin(4*d*x+4*c)+(72*A*b+72*B*a)*sin(d*x+c)+48*(a*A+3/4*B*b)*x*d)/d`

3.215.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.77

$$\int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3(4Aa + 3Bb)dx + (6Bb \cos(dx + c))^3 + 8(Ba + Ab) \cos(dx + c)^2 + 16Ba + 16Ab + 3(4Aa + 3Bb) \cos(dx + c)}{24d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output $\frac{1}{24}*(3*(4*A*a + 3*B*b)*d*x + (6*B*b*\cos(d*x + c)^3 + 8*(B*a + A*b)*\cos(d*x + c)^2 + 16*B*a + 16*A*b + 3*(4*A*a + 3*B*b)*\cos(d*x + c))*\sin(d*x + c) / d$

3.215.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(97) = 194.

Time = 0.18 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.40

$$\int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \left\{ \begin{array}{l} \frac{Aax \sin^2(c+dx)}{2} + \frac{Aax \cos^2(c+dx)}{2} + \frac{Aa \sin(c+dx) \cos(c+dx)}{2d} + \frac{2Ab \sin^3(c+dx)}{3d} + \frac{Ab \sin(c+dx) \cos^2(c+dx)}{d} + \frac{2Ba \sin^3(c+dx)}{3d} \\ x(A + B \cos(c))(a + b \cos(c)) \cos^2(c) \end{array} \right.$$

input `integrate(cos(d*x+c)**2*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)), x)`

output `Piecewise((A*a*x*sin(c + d*x)**2/2 + A*a*x*cos(c + d*x)**2/2 + A*a*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*A*b*sin(c + d*x)**3/(3*d) + A*b*sin(c + d*x)*cos(c + d*x)**2/d + 2*B*a*sin(c + d*x)**3/(3*d) + B*a*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*b*x*sin(c + d*x)**4/8 + 3*B*b*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 3*B*b*x*cos(c + d*x)**4/8 + 3*B*b*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 5*B*b*sin(c + d*x)*cos(c + d*x)**3/(8*d), Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))*cos(c)**2, True))`

3.215.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.96

$$\int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{24(2dx + 2c + \sin(2dx + 2c))Aa - 32(\sin(dx + c)^3 - 3\sin(dx + c))Ba - 32(\sin(dx + c)^3 - 3\sin(dx + c))}{96d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)), x, algorithm="maxima")`

output $1/96*(24*(2*d*x + 2*c + \sin(2*d*x + 2*c))*A*a - 32*(\sin(d*x + c)^3 - 3*\sin(d*x + c))*B*a - 32*(\sin(d*x + c)^3 - 3*\sin(d*x + c))*A*b + 3*(12*d*x + 12*c + \sin(4*d*x + 4*c) + 8*\sin(2*d*x + 2*c))*B*b)/d$

3.215.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.85

$$\int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{1}{8} (4 A a + 3 B b) x + \frac{B b \sin(4 d x + 4 c)}{32 d} + \frac{(B a + A b) \sin(3 d x + 3 c)}{12 d}$$

$$+ \frac{(A a + B b) \sin(2 d x + 2 c)}{4 d} + \frac{3 (B a + A b) \sin(d x + c)}{4 d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output $1/8*(4*A*a + 3*B*b)*x + 1/32*B*b*\sin(4*d*x + 4*c)/d + 1/12*(B*a + A*b)*\sin(3*d*x + 3*c)/d + 1/4*(A*a + B*b)*\sin(2*d*x + 2*c)/d + 3/4*(B*a + A*b)*\sin(d*x + c)/d$

3.215.9 Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.11

$$\int \cos^2(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{A a x}{2} + \frac{3 B b x}{8} + \frac{3 A b \sin(c + d x)}{4 d} + \frac{3 B a \sin(c + d x)}{4 d} + \frac{A a \sin(2 c + 2 d x)}{4 d}$$

$$+ \frac{A b \sin(3 c + 3 d x)}{12 d} + \frac{B a \sin(3 c + 3 d x)}{12 d} + \frac{B b \sin(2 c + 2 d x)}{4 d} + \frac{B b \sin(4 c + 4 d x)}{32 d}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x)),x)`

output $(A*a*x)/2 + (3*B*b*x)/8 + (3*A*b*\sin(c + d*x))/(4*d) + (3*B*a*\sin(c + d*x))/(4*d) + (A*a*\sin(2*c + 2*d*x))/(4*d) + (A*b*\sin(3*c + 3*d*x))/(12*d) + (B*a*\sin(3*c + 3*d*x))/(12*d) + (B*b*\sin(2*c + 2*d*x))/(4*d) + (B*b*\sin(4*c + 4*d*x))/(32*d)$

3.216 $\int \cos(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$

3.216.1 Optimal result	2020
3.216.2 Mathematica [A] (verified)	2020
3.216.3 Rubi [A] (verified)	2021
3.216.4 Maple [A] (verified)	2023
3.216.5 Fricas [A] (verification not implemented)	2023
3.216.6 Sympy [B] (verification not implemented)	2024
3.216.7 Maxima [A] (verification not implemented)	2024
3.216.8 Giac [A] (verification not implemented)	2025
3.216.9 Mupad [B] (verification not implemented)	2025

3.216.1 Optimal result

Integrand size = 27, antiderivative size = 84

$$\int \cos(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{1}{2}(Ab+aB)x + \frac{(3aA+2bB) \sin(c+dx)}{3d}$$

$$+ \frac{(Ab+aB) \cos(c+dx) \sin(c+dx)}{2d} + \frac{bB \cos^2(c+dx) \sin(c+dx)}{3d}$$

```
output 1/2*(A*b+B*a)*x+1/3*(3*A*a+2*B*b)*sin(d*x+c)/d+1/2*(A*b+B*a)*cos(d*x+c)*sin(d*x+c)/d+1/3*b*B*cos(d*x+c)^2*sin(d*x+c)/d
```

3.216.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.89

$$\int \cos(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{6Abc+6aBc+6Abdx+6aBdx+3(4aA+3bB) \sin(c+dx)+3(Ab+aB) \sin(2(c+dx))+bB \sin(3(c+dx))}{12d}$$

```
input Integrate[Cos[c+d*x]*(a+b*Cos[c+d*x])*(A+B*Cos[c+d*x]),x]
```

output $(6*A*b*c + 6*a*B*c + 6*A*b*d*x + 6*a*B*d*x + 3*(4*a*A + 3*b*B)*\text{Sin}[c + d*x] + 3*(A*b + a*B)*\text{Sin}[2*(c + d*x)] + b*B*\text{Sin}[3*(c + d*x)])/(12*d)$

3.216.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 3447, 3042, 3502, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \cos(c + dx) \left((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)\right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(\left(aB + Ab\right) \sin\left(c + dx + \frac{\pi}{2}\right) + aA + bB \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{1}{3} \int \cos(c + dx) (3aA + 2bB + 3(Ab + aB) \cos(c + dx)) dx + \frac{bB \sin(c + dx) \cos^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(3aA + 2bB + 3(Ab + aB) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\
 & \quad \frac{bB \sin(c + dx) \cos^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3213} \\
 & \frac{1}{3} \left(\frac{(3aA + 2bB) \sin(c + dx)}{d} + \frac{3(aB + Ab) \sin(c + dx) \cos(c + dx)}{2d} + \frac{3}{2} x(aB + Ab) \right) + \\
 & \quad \frac{bB \sin(c + dx) \cos^2(c + dx)}{3d}
 \end{aligned}$$

input `Int[Cos[c + d*x]*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(b*B*Cos[c + d*x]^2*Sin[c + d*x])/(3*d) + ((3*(A*b + a*B)*x)/2 + ((3*a*A + 2*b*B)*Sin[c + d*x])/d + (3*(A*b + a*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d) /3`

3.216.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] :=> Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; Free Q[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] :=> Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] :=> Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.216.4 Maple [A] (verified)

Time = 2.05 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.77

method	result
parallelrisch	$\frac{3(Ab+Ba) \sin(2dx+2c)+Bb \sin(3dx+3c)+3(4aA+3Bb) \sin(dx+c)+6(Ab+Ba)xd}{12d}$
parts	$\frac{(Ab+Ba) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d} + \frac{Bb(2+\cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{\sin(dx+c)aA}{d}$
derivativedivides	$\frac{Bb(2+\cos^2(dx+c)) \sin(dx+c)}{3} + Ab \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + Ba \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + aA \sin(dx+c)}{d}$
default	$\frac{Bb(2+\cos^2(dx+c)) \sin(dx+c)}{3} + Ab \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + Ba \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + aA \sin(dx+c)}{d}$
risch	$\frac{xAb}{2} + \frac{aBx}{2} + \frac{\sin(dx+c)aA}{d} + \frac{3bB \sin(dx+c)}{4d} + \frac{Bb \sin(3dx+3c)}{12d} + \frac{\sin(2dx+2c)Ab}{4d} + \frac{\sin(2dx+2c)Ba}{4d}$
norman	$\frac{\left(\frac{Ab}{2} + \frac{Ba}{2} \right) x + \left(\frac{Ab}{2} + \frac{Ba}{2} \right) x \left(\tan^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \left(\frac{3Ab}{2} + \frac{3Ba}{2} \right) x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \left(\frac{3Ab}{2} + \frac{3Ba}{2} \right) x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \frac{(2aA - Ab - Ba)}{(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right))^3}}{d}$

input `int(cos(d*x+c)*(a+cos(d*x+c)*b)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/12*(3*(A*b+B*a)*sin(2*d*x+2*c)+B*b*sin(3*d*x+3*c)+3*(4*A*a+3*B*b)*sin(d*x+c)+6*(A*b+B*a)*x*d)/d`

3.216.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

$$\int \cos(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3(Ba + Ab)dx + (2Bb \cos(dx + c))^2 + 6Aa + 4Bb + 3(Ba + Ab) \cos(dx + c) \sin(dx + c)}{6d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `1/6*(3*(B*a + A*b)*d*x + (2*B*b*cos(d*x + c)^2 + 6*A*a + 4*B*b + 3*(B*a + A*b)*cos(d*x + c))*sin(d*x + c))/d`

3.216.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(76) = 152$.

Time = 0.14 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.00

$$\int \cos(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aa \sin(c+dx)}{d} + \frac{Abx \sin^2(c+dx)}{2} + \frac{Abx \cos^2(c+dx)}{2} + \frac{Ab \sin(c+dx) \cos(c+dx)}{2d} + \frac{Bax \sin^2(c+dx)}{2} + \frac{Bax \cos^2(c+dx)}{2} + \frac{Ba \sin(c+dx)}{2d} \\ x(A + B \cos(c))(a + b \cos(c)) \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a*sin(c + d*x)/d + A*b*x*sin(c + d*x)**2/2 + A*b*x*cos(c + d*x)**2/2 + A*b*sin(c + d*x)*cos(c + d*x)/(2*d) + B*a*x*sin(c + d*x)**2/2 + B*a*x*cos(c + d*x)**2/2 + B*a*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*B*b*sin(c + d*x)**3/(3*d) + B*b*sin(c + d*x)*cos(c + d*x)**2/d, Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))*cos(c), True))`

3.216.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.94

$$\int \cos(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{3(2dx + 2c + \sin(2dx + 2c))Ba + 3(2dx + 2c + \sin(2dx + 2c))Ab - 4(\sin(dx + c)^3 - 3\sin(dx + c))}{12d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/12*(3*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a + 3*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*b - 4*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*b + 12*A*a*sin(d*x + c))/d`

3.216.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.81

$$\int \cos(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{1}{2} (Ba + Ab)x + \frac{Bb \sin(3dx + 3c)}{12d}$$

$$+ \frac{(Ba + Ab) \sin(2dx + 2c)}{4d} + \frac{(4Aa + 3Bb) \sin(dx + c)}{4d}$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/2*(B*a + A*b)*x + 1/12*B*b*sin(3*d*x + 3*c)/d + 1/4*(B*a + A*b)*sin(2*d*x + 2*c)/d + 1/4*(4*A*a + 3*B*b)*sin(d*x + c)/d
```

3.216.9 Mupad [B] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00

$$\int \cos(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{A b x}{2} + \frac{B a x}{2} + \frac{A a \sin(c + dx)}{d} + \frac{3 B b \sin(c + dx)}{4 d}$$

$$+ \frac{A b \sin(2 c + 2 d x)}{4 d} + \frac{B a \sin(2 c + 2 d x)}{4 d} + \frac{B b \sin(3 c + 3 d x)}{12 d}$$

```
input int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x)),x)
```

```
output (A*b*x)/2 + (B*a*x)/2 + (A*a*sin(c + d*x))/d + (3*B*b*sin(c + d*x))/(4*d) + (A*b*sin(2*c + 2*d*x))/(4*d) + (B*a*sin(2*c + 2*d*x))/(4*d) + (B*b*sin(3*c + 3*d*x))/(12*d)
```


3.217 $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx$

3.217.1 Optimal result	2026
3.217.2 Mathematica [A] (verified)	2026
3.217.3 Rubi [A] (verified)	2027
3.217.4 Maple [A] (verified)	2028
3.217.5 Fricas [A] (verification not implemented)	2028
3.217.6 Sympy [B] (verification not implemented)	2029
3.217.7 Maxima [A] (verification not implemented)	2029
3.217.8 Giac [A] (verification not implemented)	2030
3.217.9 Mupad [B] (verification not implemented)	2030

3.217.1 Optimal result

Integrand size = 21, antiderivative size = 52

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx = \frac{1}{2}(2aA + bB)x + \frac{(Ab + aB) \sin(c + dx)}{d} + \frac{bB \cos(c + dx) \sin(c + dx)}{2d}$$

```
output 1/2*(2*A*a+B*b)*x+(A*b+B*a)*sin(d*x+c)/d+1/2*b*B*cos(d*x+c)*sin(d*x+c)/d
```

3.217.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.98

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx = \frac{2bBc + 4aAdx + 2bBdx + 4(Ab + aB) \sin(c + dx) + bB \sin(2(c + dx))}{4d}$$

```
input Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]
```

```
output (2*b*B*c + 4*a*A*d*x + 2*b*B*d*x + 4*(A*b + a*B)*Sin[c + d*x] + b*B*Sin[2*(c + d*x)])/(4*d)
```

3.217.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right) \left(A + B \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx$$

$$\downarrow \text{3213}$$

$$\frac{(aB + Ab) \sin(c + dx)}{d} + \frac{1}{2}x(2aA + bB) + \frac{bB \sin(c + dx) \cos(c + dx)}{2d}$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `((2*a*A + b*B)*x)/2 + ((A*b + a*B)*Sin[c + d*x])/d + (b*B*Cos[c + d*x]*Sin[c + d*x])/(2*d)`

3.217.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

3.217.4 Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.98

method	result
risch	$xA A + \frac{bBx}{2} + \frac{\sin(dx+c)Ab}{d} + \frac{aB \sin(dx+c)}{d} + \frac{\sin(2dx+2c)Bb}{4d}$
parallelrisc	$\frac{4Aadx+2Bbdx+4A \sin(dx+c)b+4B \sin(dx+c)a+B \sin(2dx+2c)b}{4d}$
parts	$xA A + \frac{(Ab+Ba) \sin(dx+c)}{d} + \frac{Bb \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d}$
derivativedivides	$\frac{Bb \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + A \sin(dx+c)b + B \sin(dx+c)a + aA(dx+c)}{d}$
default	$\frac{Bb \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + A \sin(dx+c)b + B \sin(dx+c)a + aA(dx+c)}{d}$
norman	$\frac{\left(aA + \frac{Bb}{2} \right) x + \left(aA + \frac{Bb}{2} \right) x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (2aA + Bb) x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \frac{(2Ab + 2Ba - Bb) \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} + \frac{(2Ab + 2Ba + Bb)}{d}}{\left(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^2}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `x*a*A+1/2*b*B*x+sin(d*x+c)/d*A*b+a*B*sin(d*x+c)/d+1/4/d*sin(2*d*x+2*c)*B*b`

3.217.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.81

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{(2Aa + Bb)dx + (Bb \cos(dx + c) + 2Ba + 2Ab) \sin(dx + c)}{2d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `1/2*((2*A*a + B*b)*d*x + (B*b*cos(d*x + c) + 2*B*a + 2*A*b)*sin(d*x + c))/d`

3.217.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(44) = 88$.

Time = 0.09 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.81

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \begin{cases} Aax + \frac{Ab \sin(c+dx)}{d} + \frac{Ba \sin(c+dx)}{d} + \frac{Bbx \sin^2(c+dx)}{2} + \frac{Bbx \cos^2(c+dx)}{2} + \frac{Bb \sin(c+dx) \cos(c+dx)}{2d} & \text{for } d \neq 0 \\ x(A + B \cos(c))(a + b \cos(c)) & \text{otherwise} \end{cases}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a*x + A*b*sin(c + d*x)/d + B*a*sin(c + d*x)/d + B*b*x*sin(c + d*x)**2/2 + B*b*x*cos(c + d*x)**2/2 + B*b*sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c)), True))`

3.217.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.06

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{4(dx + c)Aa + (2dx + 2c + \sin(2dx + 2c))Bb + 4Ba \sin(dx + c) + 4Ab \sin(dx + c)}{4d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/4*(4*(d*x + c)*A*a + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*b + 4*B*a*sin(d*x + c) + 4*A*b*sin(d*x + c))/d`

3.217.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.87

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx = \frac{1}{2} (2 A a + B b) x + \frac{B b \sin(2 dx + 2 c)}{4 d} + \frac{(B a + A b) \sin(dx + c)}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/2*(2*A*a + B*b)*x + 1/4*B*b*sin(2*d*x + 2*c)/d + (B*a + A*b)*sin(d*x + c)/d`**3.217.9 Mupad [B] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.96

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) dx = A a x + \frac{B b x}{2} + \frac{A b \sin(c + dx)}{d} + \frac{B a \sin(c + dx)}{d} + \frac{B b \sin(2 c + 2 d x)}{4 d}$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x)),x)`output `A*a*x + (B*b*x)/2 + (A*b*sin(c + d*x))/d + (B*a*sin(c + d*x))/d + (B*b*sin(2*c + 2*d*x))/(4*d)`

3.218 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec(c+dx) dx$

3.218.1 Optimal result	2031
3.218.2 Mathematica [A] (verified)	2031
3.218.3 Rubi [A] (verified)	2032
3.218.4 Maple [A] (verified)	2034
3.218.5 Fricas [A] (verification not implemented)	2034
3.218.6 Sympy [F]	2035
3.218.7 Maxima [A] (verification not implemented)	2035
3.218.8 Giac [B] (verification not implemented)	2035
3.218.9 Mupad [B] (verification not implemented)	2036

3.218.1 Optimal result

Integrand size = 27, antiderivative size = 35

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= (Ab + aB)x + \frac{aA \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{bB \sin(c + dx)}{d}$$

output `(A*b+B*a)*x+a*A*arctanh(sin(d*x+c))/d+b*B*sin(d*x+c)/d`

3.218.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.31

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= Abx + aBx + \frac{aA \operatorname{ArcTanh}(\sin(c + dx))}{d} + \frac{bB \cos(dx) \sin(c)}{d} + \frac{bB \cos(c) \sin(dx)}{d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `A*b*x + a*B*x + (a*A*ArcTanh[Sin[c + d*x]])/d + (b*B*Cos[d*x]*Sin[c])/d + (b*B*Cos[c]*Sin[d*x])/d`

3.218.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {3042, 3447, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec(c+dx)((aB+Ab)\cos(c+dx)+aA+bB\cos^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aB+Ab)\sin(c+dx+\frac{\pi}{2})+aA+bB\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3502} \\
 & \int (aA+(Ab+aB)\cos(c+dx))\sec(c+dx)dx + \frac{bB\sin(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA+(Ab+aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{bB\sin(c+dx)}{d} \\
 & \quad \downarrow \text{3214} \\
 & aA \int \sec(c+dx)dx + x(aB+Ab) + \frac{bB\sin(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & aA \int \csc(c+dx+\frac{\pi}{2}) dx + x(aB+Ab) + \frac{bB\sin(c+dx)}{d} \\
 & \quad \downarrow \text{4257} \\
 & \frac{aA\operatorname{Arctanh}(\sin(c+dx))}{d} + x(aB+Ab) + \frac{bB\sin(c+dx)}{d}
 \end{aligned}$$

input `Int[(a + b*cos[c + d*x])*(A + B*cos[c + d*x])*Sec[c + d*x],x]`

output `(A*b + a*B)*x + (a*A*ArcTanh[Sin[c + d*x]])/d + (b*B*Sin[c + d*x])/d`

3.218.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.218.4 Maple [A] (verified)

Time = 1.62 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.37

method	result
derivativedivides	$\frac{aA \ln(\sec(dx+c)+\tan(dx+c))+Ba(dx+c)+Ab(dx+c)+B \sin(dx+c)b}{d}$
default	$\frac{aA \ln(\sec(dx+c)+\tan(dx+c))+Ba(dx+c)+Ab(dx+c)+B \sin(dx+c)b}{d}$
parts	$\frac{aA \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{(Ab+Ba)(dx+c)}{d} + \frac{bB \sin(dx+c)}{d}$
parallelrisch	$\frac{-aA \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+aA \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)+B \sin(dx+c)b+(Ab+Ba)xd}{d}$
risch	$xAb + aBx - \frac{iBbe^{i(dx+c)}}{2d} + \frac{iBbe^{-i(dx+c)}}{2d} + \frac{aA \ln(e^{i(dx+c)}+i)}{d} - \frac{aA \ln(e^{i(dx+c)}-i)}{d}$
norman	$\frac{(Ab+Ba)x+(Ab+Ba)x\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(2Ab+2Ba)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\frac{2Bb \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}+\frac{2Bb\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d}}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2} + \frac{aA}{d}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `1/d*(a*A*ln(sec(d*x+c)+tan(d*x+c))+B*a*(d*x+c)+A*b*(d*x+c)+B*sin(d*x+c)*b)`

3.218.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2(Ba + Ab)dx + Aa \log(\sin(dx + c) + 1) - Aa \log(-\sin(dx + c) + 1) + 2Bb \sin(dx + c)}{2d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fracas")`

output `1/2*(2*(B*a + A*b)*d*x + A*a*log(sin(d*x + c) + 1) - A*a*log(-sin(d*x + c) + 1) + 2*B*b*sin(d*x + c))/d`

3.218.6 Sympy [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int (A + B \cos(c + dx))(a + b \cos(c + dx)) \sec(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c), x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sec(c + d*x), x)`

3.218.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.34

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{(dx + c)Ba + (dx + c)Ab + Aa \log(\sec(dx + c) + \tan(dx + c)) + Bb \sin(dx + c)}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c), x, algorithm="maxima")`

output `((d*x + c)*B*a + (d*x + c)*A*b + A*a*log(sec(d*x + c) + tan(d*x + c)) + B*b*sin(d*x + c))/d`

3.218.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(35) = 70$.

Time = 0.29 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.26

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{Aa \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - Aa \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|) + (Ba + Ab)(dx + c) + \frac{2Bb \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1}}{d}$$

3.218. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `(A*a*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - A*a*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + (B*a + A*b)*(d*x + c) + 2*B*b*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 + 1))/d`

3.218.9 Mupad [B] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 100, normalized size of antiderivative = 2.86

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{B b \sin(c + dx)}{d} + \frac{2 A a \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

$$+ \frac{2 A b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2 B a \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x),x)`

output `(B*b*sin(c + d*x))/d + (2*A*a*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2))/d + (2*A*b*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2))/d + (2*B*a*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d`

3.219 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.219.1 Optimal result	2037
3.219.2 Mathematica [A] (verified)	2037
3.219.3 Rubi [A] (verified)	2038
3.219.4 Maple [A] (verified)	2040
3.219.5 Fricas [B] (verification not implemented)	2040
3.219.6 Sympy [F]	2041
3.219.7 Maxima [B] (verification not implemented)	2041
3.219.8 Giac [B] (verification not implemented)	2041
3.219.9 Mupad [B] (verification not implemented)	2042

3.219.1 Optimal result

Integrand size = 29, antiderivative size = 35

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= bBx + \frac{(Ab + aB)\operatorname{arctanh}(\sin(c + dx))}{d} + \frac{aA \tan(c + dx)}{d}$$

output `b*B*x+(A*b+B*a)*arctanh(sin(d*x+c))/d+a*A*tan(d*x+c)/d`

3.219.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.23

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= bBx + \frac{A b \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{a B \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{a A \tan(c + dx)}{d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `b*B*x + (A*b*ArcTanh[Sin[c + d*x]])/d + (a*B*ArcTanh[Sin[c + d*x]])/d + (a*A*Tan[c + d*x])/d`

3.219. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.219.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 3447, 3042, 3500, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec^2(c+dx)((aB+Ab)\cos(c+dx)+aA+bB\cos^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aB+Ab)\sin(c+dx+\frac{\pi}{2})+aA+bB\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3500} \\
 & \int (Ab+B\cos(c+dx)b+aB)\sec(c+dx)dx + \frac{aA\tan(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ab+B\sin(c+dx+\frac{\pi}{2})b+aB}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{aA\tan(c+dx)}{d} \\
 & \quad \downarrow \text{3214} \\
 & (aB+Ab) \int \sec(c+dx)dx + \frac{aA\tan(c+dx)}{d} + bBx \\
 & \quad \downarrow \text{3042} \\
 & (aB+Ab) \int \csc(c+dx+\frac{\pi}{2}) dx + \frac{aA\tan(c+dx)}{d} + bBx \\
 & \quad \downarrow \text{4257} \\
 & \frac{(aB+Ab)\operatorname{arctanh}(\sin(c+dx))}{d} + \frac{aA\tan(c+dx)}{d} + bBx
 \end{aligned}$$

input `Int[(a + b*cos[c + d*x])*(A + B*cos[c + d*x])*Sec[c + d*x]^2,x]`

output `b*B*x + ((A*b + a*B)*ArcTanh[Sin[c + d*x]])/d + (a*A*Tan[c + d*x])/d`

3.219.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.219.4 Maple [A] (verified)

Time = 2.59 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.43

method	result
parts	$\frac{aA \tan(dx+c)}{d} + \frac{(Ab+Ba) \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{Bb(dx+c)}{d}$
derivativedivides	$\frac{aA \tan(dx+c)+Ba \ln(\sec(dx+c)+\tan(dx+c))+Ab \ln(\sec(dx+c)+\tan(dx+c))+Bb(dx+c)}{d}$
default	$\frac{aA \tan(dx+c)+Ba \ln(\sec(dx+c)+\tan(dx+c))+Ab \ln(\sec(dx+c)+\tan(dx+c))+Bb(dx+c)}{d}$
parallelrisch	$\frac{-(Ab+Ba) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+(Ab+Ba) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)+Bbdx \cos(dx+c)+aA \sin(dx+c)}{d \cos(dx+c)}$
risch	$bBx + \frac{2iaA}{d(e^{2i(dx+c)}+1)} + \frac{\ln(e^{i(dx+c)}+i)Ab}{d} + \frac{\ln(e^{i(dx+c)}+i)Ba}{d} - \frac{\ln(e^{i(dx+c)}-i)Ab}{d} - \frac{\ln(e^{i(dx+c)}-i)Ba}{d}$
norman	$\frac{bBx\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+bBx\left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-bBx-\frac{2aA \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}-\frac{4aA\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d}-\frac{2aA\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d}}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBOSE)`

output `a*A*tan(d*x+c)/d+(A*b+B*a)/d*ln(sec(d*x+c)+tan(d*x+c))+B*b/d*(d*x+c)`

3.219.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(35) = 70.

Time = 0.31 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.43

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2 Bbdx \cos(dx + c) + (Ba + Ab) \cos(dx + c) \log(\sin(dx + c) + 1) - (Ba + Ab) \cos(dx + c) \log(-\sin(dx + c) + 1) + 2Aa \sin(dx + c)}{2d \cos(dx + c)}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/2*(2*B*b*d*x*cos(d*x + c) + (B*a + A*b)*cos(d*x + c)*log(sin(d*x + c) + 1) - (B*a + A*b)*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*A*a*sin(d*x + c))/(d*cos(d*x + c))`

3.219.6 Sympy [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int (A + B \cos(c + dx))(a + b \cos(c + dx)) \sec^2(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sec(c + d*x)**2, x)`

3.219.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(35) = 70$.

Time = 0.21 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.09

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2(dx + c)Bb + Ba(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + Ab(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1))}{2d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*B*b + B*a*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + A*b*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*A*a*tan(d*x + c))/d`

3.219.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(35) = 70$.

Time = 0.29 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.40

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{(dx + c)Bb + (Ba + Ab) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - (Ba + Ab) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|) - \frac{2Aa \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)}}{d}$$

3.219. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `((d*x + c)*B*b + (B*a + A*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (B*a + A*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*A*a*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 1))/d`

3.219.9 Mupad [B] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.26

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2 B b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{A a \sin(c + dx)}{d \cos(c + dx)}$$

$$- \frac{A b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} - \frac{B a \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^2,x)`

output `(2*B*b*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d - (B*a*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/d - (A*b*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/d + (A*a*sin(c + d*x))/(d*cos(c + d*x))`

3.220 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.220.1 Optimal result	2043
3.220.2 Mathematica [A] (verified)	2043
3.220.3 Rubi [A] (verified)	2044
3.220.4 Maple [A] (verified)	2046
3.220.5 Fricas [A] (verification not implemented)	2047
3.220.6 Sympy [F]	2047
3.220.7 Maxima [A] (verification not implemented)	2048
3.220.8 Giac [B] (verification not implemented)	2048
3.220.9 Mupad [B] (verification not implemented)	2049

3.220.1 Optimal result

Integrand size = 29, antiderivative size = 61

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(aA + 2bB)\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{(Ab + aB) \tan(c + dx)}{d} + \frac{aA \sec(c + dx) \tan(c + dx)}{2d}$$

output `1/2*(A*a+2*B*b)*arctanh(sin(d*x+c))/d+(A*b+B*a)*tan(d*x+c)/d+1/2*a*A*sec(d*x+c)*tan(d*x+c)/d`

3.220.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.23

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{aA \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{bB \operatorname{arctanh}(\sin(c + dx))}{d}$$

$$+ \frac{Ab \tan(c + dx)}{d} + \frac{aB \tan(c + dx)}{d} + \frac{aA \sec(c + dx) \tan(c + dx)}{2d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output $(a*A*ArcTanh[\sin[c + d*x]])/(2*d) + (b*B*ArcTanh[\sin[c + d*x]])/d + (A*b*\tan[c + d*x])/d + (a*B*\tan[c + d*x])/d + (a*A*\sec[c + d*x]*\tan[c + d*x])/(2*d)$

3.220.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow 3447 \\
 & \int \sec^3(c + dx) ((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow 3500 \\
 & \frac{1}{2} \int (2(Ab + aB) + (aA + 2bB) \cos(c + dx)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow 3042 \\
 & \frac{1}{2} \int \frac{2(Ab + aB) + (aA + 2bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow 3227 \\
 & \frac{1}{2} \left(2(aB + Ab) \int \sec^2(c + dx) dx + (aA + 2bB) \int \sec(c + dx) dx \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left((aA + 2bB) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + 2(aB + Ab) \int \csc \left(c + dx + \frac{\pi}{2} \right)^2 dx \right) + \\
& \quad \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
& \quad \downarrow \text{4254} \\
& \frac{1}{2} \left((aA + 2bB) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx - \frac{2(aB + Ab) \int 1d(-\tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
& \quad \downarrow \text{24} \\
& \frac{1}{2} \left((aA + 2bB) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{2(aB + Ab) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d} \\
& \quad \downarrow \text{4257} \\
& \frac{1}{2} \left(\frac{(aA + 2bB) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{2(aB + Ab) \tan(c + dx)}{d} \right) + \frac{aA \tan(c + dx) \sec(c + dx)}{2d}
\end{aligned}$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (((a*A + 2*b*B)*ArcTanh[Sin[c + d*x]])/d + (2*(A*b + a*B)*Tan[c + d*x])/d)/2`

3.220.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.220.4 Maple [A] (verified)

Time = 3.25 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{aA\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) + Ba\tan(dx+c) + Ab\tan(dx+c) + Bb\ln(\sec(dx+c)+\tan(dx+c))}{d}$
default	$\frac{aA\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right) + Ba\tan(dx+c) + Ab\tan(dx+c) + Bb\ln(\sec(dx+c)+\tan(dx+c))}{d}$
parts	$\frac{aA\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d} + \frac{(Ab+Ba)\tan(dx+c)}{d} + \frac{Bb\ln(\sec(dx+c)+\tan(dx+c))}{d}$
parallelrisc	$\frac{-(1+\cos(2dx+2c))(aA+2Bb)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + (1+\cos(2dx+2c))(aA+2Bb)\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + (2Ab+2Ba)\sin(2dx+2c)}{2d(1+\cos(2dx+2c))}$
risc	$-\frac{i(Aae^{3i(dx+c)} - 2Abe^{2i(dx+c)} - 2Ba e^{2i(dx+c)} - aAe^{i(dx+c)} - 2Ab - 2Ba)}{d(e^{2i(dx+c)} + 1)^2} + \frac{aA\ln(e^{i(dx+c)} + i)}{2d} + \frac{\ln(e^{i(dx+c)} + i)E}{d}$
norman	$\frac{(aA - 2Ab - 2Ba)\left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (aA + 2Ab + 2Ba)\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (3aA - 2Ab - 2Ba)\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (3aA + 2Ab + 2Ba)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2 \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2}$

3.220. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$

```
input int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(a*A*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+B*a*tan(d*x+c)+A*b*tan(d*x+c)+B*b*ln(sec(d*x+c)+tan(d*x+c)))
```

3.220.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.57

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(Aa + 2Bb) \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (Aa + 2Bb) \cos(dx + c)^2 \log(-\sin(dx + c) + 1) + 2(Aa + 2Bb) \cos(dx + c) \sin(dx + c)}{4d \cos(dx + c)^2}$$

```
input integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fricas")
```

```
output 1/4*((A*a + 2*B*b)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A*a + 2*B*b)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(A*a + 2*(B*a + A*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)
```

3.220.6 Sympy [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \int (A + B \cos(c + dx))(a + b \cos(c + dx)) \sec^3(c + dx) dx$$

```
input integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)
```

```
output Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sec(c + d*x)**3, x)
```

3.220.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.56

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx =$$

$$\frac{Aa \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c)+1) + \log(\sin(dx+c)-1) \right) - 2Bb(\log(\sin(dx+c)+1) - \log(\sin(dx+c)-1))}{4d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `-1/4*(A*a*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 2*B*b*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) - 4*B*a*tan(d*x + c) - 4*A*b*tan(d*x + c))/d`

3.220.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. 2(57) = 114.

Time = 0.30 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.48

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(Aa + 2Bb) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - (Aa + 2Bb) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + \frac{2 \left(Aa \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right)^3 - 2}{2d}}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output `1/2*((A*a + 2*B*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - (A*a + 2*B*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + 2*(A*a*tan(1/2*d*x + 1/2*c)^3 - 2*B*a*tan(1/2*d*x + 1/2*c)^3 - 2*A*b*tan(1/2*d*x + 1/2*c)^3 + A*a*tan(1/2*d*x + 1/2*c) + 2*B*a*tan(1/2*d*x + 1/2*c) + 2*A*b*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^2)/d`

3.220.9 Mupad [B] (verification not implemented)

Time = 1.41 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.70

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (Aa + 2Ab + 2Ba) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (2Ab - Aa + 2Ba)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)} + \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (Aa + 2Bb)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^3,x)`output `(tan(c/2 + (d*x)/2)*(A*a + 2*A*b + 2*B*a) - tan(c/2 + (d*x)/2)^3*(2*A*b - A*a + 2*B*a))/(d*(tan(c/2 + (d*x)/2)^4 - 2*tan(c/2 + (d*x)/2)^2 + 1)) + (a*tanh(tan(c/2 + (d*x)/2))*(A*a + 2*B*b))/d`

3.221 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.221.1 Optimal result	2050
3.221.2 Mathematica [A] (verified)	2050
3.221.3 Rubi [A] (verified)	2051
3.221.4 Maple [A] (verified)	2054
3.221.5 Fricas [A] (verification not implemented)	2054
3.221.6 Sympy [F]	2055
3.221.7 Maxima [A] (verification not implemented)	2055
3.221.8 Giac [B] (verification not implemented)	2056
3.221.9 Mupad [B] (verification not implemented)	2056

3.221.1 Optimal result

Integrand size = 29, antiderivative size = 93

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{(Ab + aB) \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{(2aA + 3bB) \tan(c + dx)}{3d}$$

$$+ \frac{(Ab + aB) \sec(c + dx) \tan(c + dx)}{2d} + \frac{aA \sec^2(c + dx) \tan(c + dx)}{3d}$$

```
output 1/2*(A*b+B*a)*arctanh(sin(d*x+c))/d+1/3*(2*A*a+3*B*b)*tan(d*x+c)/d+1/2*(A*
b+B*a)*sec(d*x+c)*tan(d*x+c)/d+1/3*a*A*sec(d*x+c)^2*tan(d*x+c)/d
```

3.221.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.72

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(Ab + aB) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (6aA + 6bB + 3(Ab + aB) \sec(c + dx)) + 2aA \tan^2(c + dx)}{6d}$$

```
input Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]
```

output $(3*(A*b + a*B)*ArcTanh[\text{Sin}[c + d*x]] + \text{Tan}[c + d*x]*(6*a*A + 6*b*B + 3*(A*b + a*B)*\text{Sec}[c + d*x] + 2*a*A*\text{Tan}[c + d*x]^2))/(6*d)$

3.221.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.98, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sec^4(c + dx) ((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{1}{3} \int (3(Ab + aB) + (2aA + 3bB) \cos(c + dx)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{3(Ab + aB) + (2aA + 3bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3227} \\
 & \frac{1}{3} \left(3(aB + Ab) \int \sec^3(c + dx) dx + (2aA + 3bB) \int \sec^2(c + dx) dx \right) + \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3} \left((2aA + 3bB) \int \csc \left(c + dx + \frac{\pi}{2} \right)^2 dx + 3(aB + Ab) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow 4254 \\
& \frac{1}{3} \left(3(aB + Ab) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx - \frac{(2aA + 3bB) \int 1d(-\tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow 24 \\
& \frac{1}{3} \left(3(aB + Ab) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx + \frac{(2aA + 3bB) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow 4255 \\
& \frac{1}{3} \left(3(aB + Ab) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{(2aA + 3bB) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left(3(aB + Ab) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{(2aA + 3bB) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d} \\
& \quad \downarrow 4257 \\
& \frac{1}{3} \left(3(aB + Ab) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{(2aA + 3bB) \tan(c + dx)}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^2(c + dx)}{3d}
\end{aligned}$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + (((2*a*A + 3*b*B)*Tan[c + d*x])/d + 3*(A*b + a*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/d))/(2*d))/3`

3.221.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(- (A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`
- rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.221.4 Maple [A] (verified)

Time = 4.18 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.87

method	result
parts	$-\frac{aA\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c)}{d} + \frac{(Ab+Ba)\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d} + \frac{Bb\tan(dx+c)}{d}$
derivativedivides	$-\frac{aA\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c)+Ba\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)+Ab\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d}$
default	$-\frac{aA\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c)+Ba\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)+Ab\left(\frac{\sec(dx+c)\tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2}\right)}{d}$
parallelrisch	$\frac{-9(Ab+Ba)\left(\frac{\cos(3dx+3c)}{3}+\cos(dx+c)\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+9(Ab+Ba)\left(\frac{\cos(3dx+3c)}{3}+\cos(dx+c)\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{6d(\cos(3dx+3c)+3\cos(dx+c))}$
risch	$-\frac{i(3Ab e^{5i(dx+c)}+3Ba e^{5i(dx+c)}-6Bb e^{4i(dx+c)}-12Aa e^{2i(dx+c)}-12Bb e^{2i(dx+c)}-3Ab e^{i(dx+c)}-3Ba e^{i(dx+c)}-4Aa)}{3d(e^{2i(dx+c)}+1)^3}$
norman	$\frac{\frac{4(aA-3Bb)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3d}-\frac{2(4aA-3Ab-3Ba)\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3d}-\frac{2(4aA+3Ab+3Ba)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3d}-\frac{(2aA-Ab-Ba+2Bb)\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d}}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}$

```
input int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBOSE)
```

```
output -a*A/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(A*b+B*a)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+B*b/d*tan(d*x+c)
```

3.221.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.24

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(Ba + Ab) \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3(Ba + Ab) \cos(dx + c)^3 \log(-\sin(dx + c) + 1) + 2(\dots)}{12d \cos(dx + c)^3}$$

```
input integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")
```

output `1/12*(3*(B*a + A*b)*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*(B*a + A*b)*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(2*(2*A*a + 3*B*b)*cos(d*x + c)^2 + 2*A*a + 3*(B*a + A*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.221.6 Sympy [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \int (A + B \cos(c + dx))(a + b \cos(c + dx)) \sec^4(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sec(c + d*x)**4, x)`

3.221.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.37

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{4 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa - 3 Ba \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) + 12 Bb \tan(dx + c)}{12 d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output `1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a - 3*B*a*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 3*A*b*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 12*B*b*tan(d*x + c))/d`

3.221.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(85) = 170.

Time = 0.33 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.26

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(Ba + Ab) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - 3(Ba + Ab) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right) - \frac{2(6Aa \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 6Ab \tan(\frac{1}{2} dx + \frac{1}{2} c)^5)}{\tan^2(\frac{1}{2} dx + \frac{1}{2} c) - 1}}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")`

output `1/6*(3*(B*a + A*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*(B*a + A*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*A*a*tan(1/2*d*x + 1/2*c)^5 - 3*B*a*tan(1/2*d*x + 1/2*c)^5 - 3*A*b*tan(1/2*d*x + 1/2*c)^5 + 6*B*b*tan(1/2*d*x + 1/2*c)^5 - 4*A*a*tan(1/2*d*x + 1/2*c)^3 - 12*B*b*tan(1/2*d*x + 1/2*c)^3 + 6*A*a*tan(1/2*d*x + 1/2*c) + 3*B*a*tan(1/2*d*x + 1/2*c) + 3*A*b*tan(1/2*d*x + 1/2*c) + 6*B*b*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^3)/d`

3.221.9 Mupad [B] (verification not implemented)

Time = 2.69 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.56

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{\operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) (Ab + Ba)}{d} - \frac{(2Aa - Ab - Ba + 2Bb) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{4Aa}{3} - 4Bb\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + (2Aa + Ab + Ba + 2Bb) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^4,x)`

output `(atanh(tan(c/2 + (d*x)/2))*(A*b + B*a))/d - (tan(c/2 + (d*x)/2)*(2*A*a + A*b + B*a + 2*B*b) - tan(c/2 + (d*x)/2)^3*((4*A*a)/3 + 4*B*b) + tan(c/2 + (d*x)/2)^5*(2*A*a - A*b - B*a + 2*B*b))/(d*(3*tan(c/2 + (d*x)/2)^2 - 3*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 - 1))`

3.222 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.222.1 Optimal result	2057
3.222.2 Mathematica [A] (verified)	2058
3.222.3 Rubi [A] (verified)	2058
3.222.4 Maple [A] (verified)	2061
3.222.5 Fricas [A] (verification not implemented)	2062
3.222.6 Sympy [F]	2062
3.222.7 Maxima [A] (verification not implemented)	2063
3.222.8 Giac [B] (verification not implemented)	2063
3.222.9 Mupad [B] (verification not implemented)	2064

3.222.1 Optimal result

Integrand size = 29, antiderivative size = 114

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{(3aA + 4bB) \operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{(Ab + aB) \tan(c + dx)}{d}$$

$$+ \frac{(3aA + 4bB) \sec(c + dx) \tan(c + dx)}{8d}$$

$$+ \frac{aA \sec^3(c + dx) \tan(c + dx)}{4d} + \frac{(Ab + aB) \tan^3(c + dx)}{3d}$$

```
output 1/8*(3*A*a+4*B*b)*arctanh(sin(d*x+c))/d+(A*b+B*a)*tan(d*x+c)/d+1/8*(3*A*a+
4*B*b)*sec(d*x+c)*tan(d*x+c)/d+1/4*a*A*sec(d*x+c)^3*tan(d*x+c)/d+1/3*(A*b+
B*a)*tan(d*x+c)^3/d
```


3.222.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.75

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3aA + 4bB) \operatorname{arctanh}(\sin(c + dx)) + \sec(c + dx) (9aA + 12bB + 8(Ab + aB)(2 + \cos(2(c + dx)))) \sec(c + dx) + 6aA \sec^3(c + dx) \tan(c + dx)}{24d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(3*(3*a*A + 4*b*B)*ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*(9*a*A + 12*b*B + 8*(A*b + a*B)*(2 + Cos[2*(c + d*x)])*Sec[c + d*x] + 6*a*A*Sec[c + d*x]^2)*Tan[c + d*x])/(24*d)`

3.222.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.93, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 3447, 3042, 3500, 3042, 3227, 3042, 4254, 2009, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3447}$$

$$\int \sec^5(c + dx) ((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3500}$$

$$\begin{aligned}
& \frac{1}{4} \int (4(Ab + aB) + (3aA + 4bB) \cos(c + dx)) \sec^4(c + dx) dx + \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \int \frac{4(Ab + aB) + (3aA + 4bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^4} dx + \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{4} \left(4(aB + Ab) \int \sec^4(c + dx) dx + (3aA + 4bB) \int \sec^3(c + dx) dx \right) + \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left((3aA + 4bB) \int \csc(c + dx + \frac{\pi}{2})^3 dx + 4(aB + Ab) \int \csc(c + dx + \frac{\pi}{2})^4 dx \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{4254} \\
& \frac{1}{4} \left((3aA + 4bB) \int \csc(c + dx + \frac{\pi}{2})^3 dx - \frac{4(aB + Ab) \int (\tan^2(c + dx) + 1) d(-\tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{2009} \\
& \frac{1}{4} \left((3aA + 4bB) \int \csc(c + dx + \frac{\pi}{2})^3 dx - \frac{4(aB + Ab) (-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{4255} \\
& \frac{1}{4} \left((3aA + 4bB) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{4(aB + Ab) (-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left((3aA + 4bB) \left(\frac{1}{2} \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{4(aB + Ab) (-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx))}{d} \right) + \\
& \quad \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \\
& \quad \downarrow \text{4257}
\end{aligned}$$

$$\frac{1}{4} \left((3aA + 4bB) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) - \frac{4(aB + Ab) \left(-\frac{1}{3} \tan^3(c + dx) - \tan(c + dx) \right)}{d} + \frac{aA \tan(c + dx) \sec^3(c + dx)}{4d} \right)$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*A*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((3*a*A + 4*b*B)*(ArcTanh[Sin[c + d*x]])/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)) - (4*(A*b + a*B)*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d)/4`

3.222.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.222.4 Maple [A] (verified)

Time = 4.34 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.04

method	result
parts	$\frac{aA \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)}{d} - \frac{(Ab+Ba) \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d}$
derivativedivides	$\frac{aA \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right) - Ba \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d}$
default	$\frac{aA \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right) - Ba \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d}$
parallelrisch	$\frac{-18 \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) (aA + \frac{4Bb}{3}) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 18 \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) (aA + \frac{4Bb}{3})}{12d(\cos(4dx+4c) + 1)}$
risch	$- \frac{i(9Aa e^{7i(dx+c)} + 12Bb e^{7i(dx+c)} + 33Aa e^{5i(dx+c)} + 12Bb e^{5i(dx+c)} - 48Ab e^{4i(dx+c)} - 48Ba e^{4i(dx+c)} - 33Aa e^{3i(dx+c)} - 12Bb e^{3i(dx+c)})}{12d(e^{2i(dx+c)} + 1)^4}$
norman	$\frac{(5aA - 8Ab - 8Ba + 4Bb) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{4d} + \frac{(5aA + 8Ab + 8Ba + 4Bb) \tan \left(\frac{dx}{2} + \frac{c}{2} \right)}{4d} + \frac{(21aA - 8Ab - 8Ba - 12Bb) \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{6d} + \frac{(21aA - 8Ab - 8Ba - 12Bb) \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{6d} + \frac{(1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right))^2}{12d}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBOSE)`

output $a*A/d*(-(-1/4*\sec(d*x+c)^3-3/8*\sec(d*x+c))*\tan(d*x+c)+3/8*\ln(\sec(d*x+c)+\tan(d*x+c)))-(A*b+B*a)/d*(-2/3-1/3*\sec(d*x+c)^2)*\tan(d*x+c)+B*b/d*(1/2*\sec(d*x+c)*\tan(d*x+c)+1/2*\ln(\sec(d*x+c)+\tan(d*x+c)))$

3.222.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.19

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3Aa + 4Bb) \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(3Aa + 4Bb) \cos(dx + c)^4 \log(-\sin(dx + c) + 1)}{48d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")`

output $1/48*(3*(3*A*a + 4*B*b)*\cos(d*x + c)^4*\log(\sin(d*x + c) + 1) - 3*(3*A*a + 4*B*b)*\cos(d*x + c)^4*\log(-\sin(d*x + c) + 1) + 2*(16*(B*a + A*b)*\cos(d*x + c)^3 + 3*(3*A*a + 4*B*b)*\cos(d*x + c)^2 + 6*A*a + 8*(B*a + A*b)*\cos(d*x + c))*\sin(d*x + c))/(d*\cos(d*x + c)^4)$

3.222.6 Sympy [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \int (A + B \cos(c + dx))(a + b \cos(c + dx)) \sec^5(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sec(c + d*x)**5, x)`

3.222.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.43

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ba + 16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ab - 3 Aa \left(\frac{2(3 \sin(dx+c)^3 - 5 \sin(dx+c)}{\sin(dx+c)^4 - 2 \sin(dx+c)} \right)}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output `1/48*(16*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a + 16*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*b - 3*A*a*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 12*B*b*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)))/d`

3.222.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. 2(106) = 212.

Time = 0.30 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.67

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3Aa + 4Bb) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 3(3Aa + 4Bb) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) + \frac{2(15Aa \tan(\frac{1}{2}dx + \frac{1}{2}c) - 5Aa)}{\sin(dx+c)^4 - 2\sin(dx+c)}}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")`

output $\frac{1}{24}*(3*(3*A*a + 4*B*b)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(3*A*a + 4*B*b)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) + 2*(15*A*a*\tan(1/2*d*x + 1/2*c)^7 - 24*B*a*\tan(1/2*d*x + 1/2*c)^7 - 24*A*b*\tan(1/2*d*x + 1/2*c)^7 + 12*B*b*\tan(1/2*d*x + 1/2*c)^7 + 9*A*a*\tan(1/2*d*x + 1/2*c)^5 + 40*B*a*\tan(1/2*d*x + 1/2*c)^5 + 40*A*b*\tan(1/2*d*x + 1/2*c)^5 - 12*B*b*\tan(1/2*d*x + 1/2*c)^5 + 9*A*a*\tan(1/2*d*x + 1/2*c)^3 - 40*B*a*\tan(1/2*d*x + 1/2*c)^3 - 40*A*b*\tan(1/2*d*x + 1/2*c)^3 - 12*B*b*\tan(1/2*d*x + 1/2*c)^3 + 15*A*a*\tan(1/2*d*x + 1/2*c) + 24*B*a*\tan(1/2*d*x + 1/2*c) + 24*A*b*\tan(1/2*d*x + 1/2*c) + 12*B*b*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^4/d$

3.222.9 Mupad [B] (verification not implemented)

Time = 4.12 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.70

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{\left(\frac{5Aa}{4} - 2Ab - 2Ba + Bb\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{3Aa}{4} + \frac{10Ab}{3} + \frac{10Ba}{3} - Bb\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(\frac{3Aa}{4} - \frac{10Ab}{3} - \frac{10Ba}{3} + Bb\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + \left(\frac{3Aa}{4} + Bb\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \frac{\text{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) \left(\frac{3Aa}{4} + Bb\right)}{d}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)}$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^5,x)`

output $(\tan(c/2 + (d*x)/2)*((5*A*a)/4 + 2*A*b + 2*B*a + B*b) + \tan(c/2 + (d*x)/2)^7*((5*A*a)/4 - 2*A*b - 2*B*a + B*b) - \tan(c/2 + (d*x)/2)^3*((10*A*b)/3 - (3*A*a)/4 + (10*B*a)/3 + B*b) + \tan(c/2 + (d*x)/2)^5*((3*A*a)/4 + (10*A*b)/3 + (10*B*a)/3 - B*b))/(d*(6*\tan(c/2 + (d*x)/2)^4 - 4*\tan(c/2 + (d*x)/2)^2 - 4*\tan(c/2 + (d*x)/2)^6 + \tan(c/2 + (d*x)/2)^8 + 1)) + (\text{atanh}(\tan(c/2 + (d*x)/2))*((3*A*a)/4 + B*b))/d$

3.223 $\int \cos^2(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.223.1 Optimal result	2065
3.223.2 Mathematica [A] (verified)	2066
3.223.3 Rubi [A] (verified)	2066
3.223.4 Maple [A] (verified)	2070
3.223.5 Fricas [A] (verification not implemented)	2070
3.223.6 Sympy [B] (verification not implemented)	2071
3.223.7 Maxima [A] (verification not implemented)	2072
3.223.8 Giac [A] (verification not implemented)	2072
3.223.9 Mupad [B] (verification not implemented)	2073

3.223.1 Optimal result

Integrand size = 31, antiderivative size = 189

$$\int \cos^2(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$$

$$= \frac{1}{8}(4a^2A + 3Ab^2 + 6abB) x + \frac{(4b^2B + 5a(2Ab + aB)) \sin(c+dx)}{5d}$$

$$+ \frac{(4a^2A + 3Ab^2 + 6abB) \cos(c+dx) \sin(c+dx)}{8d}$$

$$+ \frac{b(5Ab + 6aB) \cos^3(c+dx) \sin(c+dx)}{20d}$$

$$+ \frac{bB \cos^3(c+dx)(a+b \cos(c+dx)) \sin(c+dx)}{5d} - \frac{(4b^2B + 5a(2Ab + aB)) \sin^3(c+dx)}{15d}$$

output `1/8*(4*A*a^2+3*A*b^2+6*B*a*b)*x+1/5*(4*b^2*B+5*a*(2*A*b+B*a))*sin(d*x+c)/d`
`+1/8*(4*A*a^2+3*A*b^2+6*B*a*b)*cos(d*x+c)*sin(d*x+c)/d+1/20*b*(5*A*b+6*B*a)`
`*cos(d*x+c)^3*sin(d*x+c)/d+1/5*b*B*cos(d*x+c)^3*(a+b*cos(d*x+c))*sin(d*x+`
`c)/d-1/15*(4*b^2*B+5*a*(2*A*b+B*a))*sin(d*x+c)^3/d`

3.223.2 Mathematica [A] (verified)

Time = 1.15 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.77

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{60(4a^2A + 3Ab^2 + 6abB)(c + dx) + 60(12aAb + 6a^2B + 5b^2B) \sin(c + dx) + 120(a^2A + Ab^2 + 2abB) \sin(2(c + dx)) + 10(8a^2Ab + 4a^2B + 5b^2B) \sin(3(c + dx)) + 15b(Ab + 2aB) \sin(4(c + dx)) + 6b^2B \sin(5(c + dx))}{480d}$$

input `Integrate[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output $(60*(4*a^2*A + 3*A*b^2 + 6*a*b*B)*(c + d*x) + 60*(12*a*A*b + 6*a^2*B + 5*b^2*B)*\text{Sin}[c + d*x] + 120*(a^2*A + A*b^2 + 2*a*b*B)*\text{Sin}[2*(c + d*x)] + 10*(8*a^2*A*b + 4*a^2*B + 5*b^2*B)*\text{Sin}[3*(c + d*x)] + 15*b*(A*b + 2*a*B)*\text{Sin}[4*(c + d*x)] + 6*b^2*B*\text{Sin}[5*(c + d*x)])/(480*d)$

3.223.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.87, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3469, 3042, 3502, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3469}$$

$$\frac{1}{5} \int \cos^2(c + dx) (b(5Ab + 6aB) \cos^2(c + dx) + (4Bb^2 + 5a(2Ab + aB)) \cos(c + dx) + a(5aA + 3bB)) dx + \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(\frac{b(5Ab + 6aB) \sin \left(c + dx + \frac{\pi}{2} \right)^2 + (4Bb^2 + 5a(2Ab + aB)) \sin \left(c + dx + \frac{\pi}{2} \right) + a(5aA - bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx)))}{5d} \right) dx$$

↓ 3502

$$\frac{1}{5} \left(\frac{1}{4} \int \cos^2(c + dx) (5(4Aa^2 + 6bBa + 3Ab^2) + 4(4Bb^2 + 5a(2Ab + aB)) \cos(c + dx)) dx + \frac{b(6aB + 5Ab) \sin(c + dx)}{4} \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 (5(4Aa^2 + 6bBa + 3Ab^2) + 4(4Bb^2 + 5a(2Ab + aB)) \sin \left(c + dx + \frac{\pi}{2} \right)) dx + \frac{b(6aB + 5Ab) \sin(c + dx)}{4} \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d}$$

↓ 3227

$$\frac{1}{5} \left(\frac{1}{4} \left(5(4a^2A + 6abB + 3Ab^2) \int \cos^2(c + dx) dx + 4(5a(aB + 2Ab) + 4b^2B) \int \cos^3(c + dx) dx \right) + \frac{b(6aB + 5Ab) \sin(c + dx)}{4} \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(5(4a^2A + 6abB + 3Ab^2) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + 4(5a(aB + 2Ab) + 4b^2B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^3 dx \right) + \frac{b(6aB + 5Ab) \sin(c + dx)}{4} \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d}$$

↓ 3113

$$\frac{1}{5} \left(\frac{1}{4} \left(5(4a^2A + 6abB + 3Ab^2) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4(5a(aB + 2Ab) + 4b^2B) \int (1 - \sin^2(c + dx)) d(-\sin(c + dx))}{d} \right) + \frac{b(6aB + 5Ab) \sin(c + dx)}{4} \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d}$$

↓ 2009

$$\frac{1}{5} \left(\frac{1}{4} \left(5(4a^2A + 6abB + 3Ab^2) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{4(5a(aB + 2Ab) + 4b^2B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right. \right. \\ \left. \left. \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d} \right) \right. \\ \left. \downarrow \text{3115} \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(5(4a^2A + 6abB + 3Ab^2) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{4(5a(aB + 2Ab) + 4b^2B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right. \right. \\ \left. \left. \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d} \right) \right. \\ \left. \downarrow \text{24} \right.$$

$$\frac{1}{5} \left(\frac{1}{4} \left(5(4a^2A + 6abB + 3Ab^2) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} + \frac{x}{2} \right) - \frac{4(5a(aB + 2Ab) + 4b^2B) \left(\frac{1}{3} \sin^3(c + dx) - \sin(c + dx) \right)}{d} \right. \right. \\ \left. \left. \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))}{5d} \right) \right.$$

input `Int[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(b*B*Cos[c + d*x]^3*(a + b*Cos[c + d*x])*Sin[c + d*x])/(5*d) + ((b*(5*A*b + 6*a*B)*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (5*(4*a^2*A + 3*A*b^2 + 6*a*b*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) - (4*(4*b^2*B + 5*a*(2*A*b + a*B))*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)/4)/5`

3.223.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_.)]^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)]*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)]^(n_)), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)] + (C_.)*sin[(e_.) + (f_.)*(x_.)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.223.4 Maple [A] (verified)

Time = 4.00 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.78

method	result
parallelrisch	$\frac{120(Aa^2+Ab^2+2Bab)\sin(2dx+2c)+10(8Aab+4Ba^2+5Bb^2)\sin(3dx+3c)+15(Ab^2+2Bab)\sin(4dx+4c)+6Bb^2\sin(5dx+5c)}{480d}$
parts	$\frac{(Ab^2+2Bab)\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4}+\frac{3dx}{8}+\frac{3c}{8}\right)}{d}+\frac{(2Aab+Ba^2)(2+\cos^2(dx+c))\sin(dx+c)}{3d}+\frac{Aa^2}{d}$
derivativedivides	$\frac{Bb^2\left(\frac{8}{3}+\cos^4(dx+c)+\frac{4(\cos^2(dx+c))}{3}\right)\sin(dx+c)}{5}+Ab^2\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4}+\frac{3dx}{8}+\frac{3c}{8}\right)+2Bab\left(\frac{\cos^3(dx+c)}{4}\right)$
default	$\frac{Bb^2\left(\frac{8}{3}+\cos^4(dx+c)+\frac{4(\cos^2(dx+c))}{3}\right)\sin(dx+c)}{5}+Ab^2\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4}+\frac{3dx}{8}+\frac{3c}{8}\right)+2Bab\left(\frac{\cos^3(dx+c)}{4}\right)$
risch	$\frac{a^2xA}{2}+\frac{3xA b^2}{8}+\frac{3xBab}{4}+\frac{3\sin(dx+c)Aab}{2d}+\frac{3\sin(dx+c)B a^2}{4d}+\frac{5b^2B\sin(dx+c)}{8d}+\frac{B b^2\sin(5dx+5c)}{80d}+\sin(dx+c)$
norman	$\frac{(\frac{1}{2}Aa^2+\frac{3}{8}Ab^2+\frac{3}{4}Bab)x+(5Aa^2+\frac{15}{4}Ab^2+\frac{15}{2}Bab)x\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(5Aa^2+\frac{15}{4}Ab^2+\frac{15}{2}Bab)x\left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin(dx+c)}{d}$

input `int(cos(d*x+c)^2*(a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/480*(120*(A*a^2+A*b^2+2*B*a*b)*sin(2*d*x+2*c)+10*(8*A*a*b+4*B*a^2+5*B*b^2)*sin(3*d*x+3*c)+15*(A*b^2+2*B*a*b)*sin(4*d*x+4*c)+6*B*b^2*sin(5*d*x+5*c)+60*(12*A*a*b+6*B*a^2+5*B*b^2)*sin(d*x+c)+240*x*d*(A*a^2+3/4*A*b^2+3/2*B*a*b))/d`

3.223.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.75

$$\int \cos^2(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{15(4Aa^2+6Bab+3Ab^2)dx+(24Bb^2\cos(dx+c))^4+30(2Bab+Ab^2)\cos(dx+c)^3+80Ba^2+160A}{d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")`

3.223. $\int \cos^2(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$

output $\frac{1}{120}(15(4Aa^2 + 6Bab + 3Ab^2)dx + (24Bb^2\cos(dx + c)^4 + 30(2Bab + Ab^2)\cos(dx + c)^3 + 80Ba^2 + 160Aab + 64Bb^2 + 8(5Ba^2 + 10Aab + 4Bb^2)\cos(dx + c)^2 + 15(4Aa^2 + 6Bab + 3Ab^2)\cos(dx + c))\sin(dx + c))/d$

3.223.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 459 vs. $2(184) = 368$.

Time = 0.29 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.43

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aa^2x \sin^2(c+dx)}{2} + \frac{Aa^2x \cos^2(c+dx)}{2} + \frac{Aa^2 \sin(c+dx) \cos(c+dx)}{2d} + \frac{4Aab \sin^3(c+dx)}{3d} + \frac{2Aab \sin(c+dx) \cos^2(c+dx)}{d} + \frac{3Ab^2x \sin^4}{8} \\ x(A + B \cos(c))(a + b \cos(c))^2 \cos^2(c) \end{cases}$$

input `integrate(cos(dx+c)**2*(a+b*cos(dx+c))**2*(A+B*cos(dx+c)),x)`

output `Piecewise((A**2*x*sin(c + dx)**2/2 + A**2*x*cos(c + dx)**2/2 + A**2*sin(c + dx)*cos(c + dx)/(2*d) + 4*A*a*b*sin(c + dx)**3/(3*d) + 2*A*a*b*sin(c + dx)*cos(c + dx)**2/d + 3*A*b**2*x*sin(c + dx)**4/8 + 3*A*b**2*x*sin(c + dx)**2*cos(c + dx)**2/4 + 3*A*b**2*x*cos(c + dx)**4/8 + 3*A*b**2*sin(c + dx)**3*cos(c + dx)/(8*d) + 5*A*b**2*sin(c + dx)*cos(c + dx)**3/(8*d) + 2*B*a**2*sin(c + dx)**3/(3*d) + B*a**2*sin(c + dx)*cos(c + dx)**2/d + 3*B*a*b*x*sin(c + dx)**4/4 + 3*B*a*b*x*sin(c + dx)**2*cos(c + dx)**2/2 + 3*B*a*b*x*cos(c + dx)**4/4 + 3*B*a*b*sin(c + dx)**3*cos(c + dx)/(4*d) + 5*B*a*b*sin(c + dx)*cos(c + dx)**3/(4*d) + 8*B*b**2*sin(c + dx)**5/(15*d) + 4*B*b**2*sin(c + dx)**3*cos(c + dx)**2/(3*d) + B*b**2*sin(c + dx)*cos(c + dx)**4/d, Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**2*cos(c)**2, True))`

3.223.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.93

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{120(2dx + 2c + \sin(2dx + 2c))Aa^2 - 160(\sin(dx + c)^3 - 3\sin(dx + c))Ba^2 - 320(\sin(dx + c)^3 - 3\sin(dx + c))Aab + 30(12dx + 12c + \sin(4dx + 4c) + 8\sin(2dx + 2c))Bab + 15(12dx + 12c + \sin(4dx + 4c) + 8\sin(2dx + 2c))Aab^2 + 32(3\sin(dx + c)^5 - 10\sin(dx + c)^3 + 15\sin(dx + c))Bb^2}{d}$$

```
input integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output 1/480*(120*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^2 - 160*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^2 - 320*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a*b + 30*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a*b + 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*b^2 + 32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*b^2)/d
```

3.223.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.83

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{Bb^2 \sin(5dx + 5c)}{80d} + \frac{1}{8}(4Aa^2 + 6Bab + 3Ab^2)x$$

$$+ \frac{(2Bab + Ab^2) \sin(4dx + 4c)}{32d} + \frac{(4Ba^2 + 8Aab + 5Bb^2) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(Aa^2 + 2Bab + Ab^2) \sin(2dx + 2c)}{4d} + \frac{(6Ba^2 + 12Aab + 5Bb^2) \sin(dx + c)}{8d}$$

```
input integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")
```

```
output 1/80*B*b^2*sin(5*d*x + 5*c)/d + 1/8*(4*A*a^2 + 6*B*a*b + 3*A*b^2)*x + 1/32*(2*B*a*b + A*b^2)*sin(4*d*x + 4*c)/d + 1/48*(4*B*a^2 + 8*A*a*b + 5*B*b^2)*sin(3*d*x + 3*c)/d + 1/4*(A*a^2 + 2*B*a*b + A*b^2)*sin(2*d*x + 2*c)/d + 1/8*(6*B*a^2 + 12*A*a*b + 5*B*b^2)*sin(d*x + c)/d
```

3.223.9 Mupad [B] (verification not implemented)

Time = 4.09 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.62

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx = \frac{x \left(A a^2 + \frac{3 B a b}{2} + \frac{3 A b^2}{4} \right)}{2} + \frac{\left(2 B a^2 - \frac{5 A b^2}{4} - A a^2 + 2 B b^2 + 4 A a b - \frac{5 B a b}{2} \right) \tan\left(\frac{c}{2} + \frac{d x}{2}\right)^9 + \left(\frac{16 B a^2}{3} - \frac{A b^2}{2} - 2 A a^2 + \frac{8 B b^2}{3} + \frac{32 A a b}{3} \right) \tan\left(\frac{c}{2} + \frac{d x}{2}\right)^7 + \left(\frac{16 B a^2}{3} - \frac{A b^2}{2} - 2 A a^2 + \frac{8 B b^2}{3} + \frac{32 A a b}{3} \right) \tan\left(\frac{c}{2} + \frac{d x}{2}\right)^5 + \left(\frac{16 B a^2}{3} - \frac{A b^2}{2} - 2 A a^2 + \frac{8 B b^2}{3} + \frac{32 A a b}{3} \right) \tan\left(\frac{c}{2} + \frac{d x}{2}\right)^3 + \left(\frac{16 B a^2}{3} - \frac{A b^2}{2} - 2 A a^2 + \frac{8 B b^2}{3} + \frac{32 A a b}{3} \right) \tan\left(\frac{c}{2} + \frac{d x}{2}\right)}{10}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2,x)`

```
output (x*(A*a^2 + (3*A*b^2)/4 + (3*B*a*b)/2))/2 + (tan(c/2 + (d*x)/2)^5*((20*B*a^2)/3 + (116*B*b^2)/15 + (40*A*a*b)/3) - tan(c/2 + (d*x)/2)^9*(A*a^2 + (5*A*b^2)/4 - 2*B*a^2 - 2*B*b^2 - 4*A*a*b + (5*B*a*b)/2) + tan(c/2 + (d*x)/2)^3*(2*A*a^2 + (A*b^2)/2 + (16*B*a^2)/3 + (8*B*b^2)/3 + (32*A*a*b)/3 + B*a*b) - tan(c/2 + (d*x)/2)^7*(2*A*a^2 + (A*b^2)/2 - (16*B*a^2)/3 - (8*B*b^2)/3 - (32*A*a*b)/3 + B*a*b) + tan(c/2 + (d*x)/2)*(A*a^2 + (5*A*b^2)/4 + 2*B*a^2 + 2*B*b^2 + 4*A*a*b + (5*B*a*b)/2))/(d*(5*tan(c/2 + (d*x)/2)^2 + 10*tan(c/2 + (d*x)/2)^4 + 10*tan(c/2 + (d*x)/2)^6 + 5*tan(c/2 + (d*x)/2)^8 + tan(c/2 + (d*x)/2)^10 + 1))
```


3.224 $\int \cos(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.224.1 Optimal result	2074
3.224.2 Mathematica [A] (verified)	2075
3.224.3 Rubi [A] (verified)	2075
3.224.4 Maple [A] (verified)	2078
3.224.5 Fracas [A] (verification not implemented)	2078
3.224.6 Sympy [B] (verification not implemented)	2079
3.224.7 Maxima [A] (verification not implemented)	2079
3.224.8 Giac [A] (verification not implemented)	2080
3.224.9 Mupad [B] (verification not implemented)	2080

3.224.1 Optimal result

Integrand size = 29, antiderivative size = 170

$$\int \cos(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$$

$$= \frac{1}{8}(8aAb + 4a^2B + 3b^2B)x + \frac{(4a^2Ab + 4Ab^3 - a^3B + 8ab^2B) \sin(c+dx)}{6bd}$$

$$+ \frac{(8aAb - 2a^2B + 9b^2B) \cos(c+dx) \sin(c+dx)}{24d}$$

$$+ \frac{(4Ab - aB)(a+b \cos(c+dx))^2 \sin(c+dx)}{12bd} + \frac{B(a+b \cos(c+dx))^3 \sin(c+dx)}{4bd}$$

output

```
1/8*(8*A*a*b+4*B*a^2+3*B*b^2)*x+1/6*(4*A*a^2*b+4*A*b^3-B*a^3+8*B*a*b^2)*sin(d*x+c)/b/d+1/24*(8*A*a*b-2*B*a^2+9*B*b^2)*cos(d*x+c)*sin(d*x+c)/d+1/12*(4*A*b-B*a)*(a+b*cos(d*x+c))^2*sin(d*x+c)/b/d+1/4*B*(a+b*cos(d*x+c))^3*sin(d*x+c)/b/d
```

3.224.2 Mathematica [A] (verified)

Time = 1.64 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.69

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{12(8aAb + 4a^2B + 3b^2B)(c + dx) + 24(4a^2A + 3Ab^2 + 6abB) \sin(c + dx) + 24(2aAb + a^2B + b^2B) \sin(2(c + dx)) + 8b^2B \sin(3(c + dx)) + 3b^2B \sin(4(c + dx))}{96d}$$

input `Integrate[Cos[c + d*x]*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(12*(8*a*A*b + 4*a^2*B + 3*b^2*B)*(c + d*x) + 24*(4*a^2*A + 3*A*b^2 + 6*a*b*B)*Sin[c + d*x] + 24*(2*a*A*b + a^2*B + b^2*B)*Sin[2*(c + d*x)] + 8*b*(A*b + 2*a*B)*Sin[3*(c + d*x)] + 3*b^2*B*Sine[4*(c + d*x)])/(96*d)`

3.224.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 3447, 3042, 3502, 3042, 3232, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int (a + b \cos(c + dx))^2 (A \cos(c + dx) + B \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{3502}$$

$$\frac{\int (a + b \cos(c + dx))^2 (3bB + (4Ab - aB) \cos(c + dx)) dx}{4b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^3}{4bd}$$

$$\begin{aligned}
 & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (3bB + (4Ab - aB) \sin(c + dx + \frac{\pi}{2})) dx}{\frac{4b}{B \sin(c + dx)(a + b \cos(c + dx))^3}} + \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{\frac{1}{3} \int (a + b \cos(c + dx)) (b(8Ab + 7aB) + (-2Ba^2 + 8Aba + 9b^2B) \cos(c + dx)) dx + \frac{(4Ab - aB) \sin(c + dx)(a + b \cos(c + dx))}{3d}}{\frac{4b}{B \sin(c + dx)(a + b \cos(c + dx))^3}} \\
 & \qquad \qquad \qquad \downarrow \text{3232} \\
 & \frac{\frac{1}{3} \int (a + b \sin(c + dx + \frac{\pi}{2})) (b(8Ab + 7aB) + (-2Ba^2 + 8Aba + 9b^2B) \sin(c + dx + \frac{\pi}{2})) dx + \frac{(4Ab - aB) \sin(c + dx)(a + b \cos(c + dx))}{3d}}{\frac{4b}{B \sin(c + dx)(a + b \cos(c + dx))^3}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{\frac{1}{3} \int (a + b \sin(c + dx + \frac{\pi}{2})) (b(8Ab + 7aB) + (-2Ba^2 + 8Aba + 9b^2B) \sin(c + dx + \frac{\pi}{2})) dx + \frac{(4Ab - aB) \sin(c + dx)(a + b \cos(c + dx))}{3d}}{\frac{4b}{B \sin(c + dx)(a + b \cos(c + dx))^3}} \\
 & \qquad \qquad \qquad \downarrow \text{3213} \\
 & \frac{\frac{1}{3} \left(\frac{b(-2a^2B + 8aAb + 9b^2B) \sin(c + dx) \cos(c + dx)}{2d} + \frac{3}{2}bx(4a^2B + 8aAb + 3b^2B) + \frac{2(a^3(-B) + 4a^2Ab + 8ab^2B + 4Ab^3) \sin(c + dx)}{d} \right) + \frac{(4Ab - aB) \sin(c + dx)(a + b \cos(c + dx))}{3d}}{\frac{4b}{B \sin(c + dx)(a + b \cos(c + dx))^3}}
 \end{aligned}$$

```
input Int[Cos[c + d*x]*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]
```

```
output (B*(a + b*Cos[c + d*x])^3*Sin[c + d*x])/(4*b*d) + (((4*A*b - a*B)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + ((3*b*(8*a*A*b + 4*a^2*B + 3*b^2*B)*x)/2 + (2*(4*a^2*A*b + 4*A*b^3 - a^3*B + 8*a*b^2*B)*Sin[c + d*x])/d + (b*(8*a*A*b - 2*a^2*B + 9*b^2*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/3)/(4*b)
```

3.224. $\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$

3.224.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.224.4 Maple [A] (verified)

Time = 2.76 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.69

method	result
parallelrisc	$\frac{24(2Aab+Ba^2+Bb^2)\sin(2dx+2c)+8(Ab^2+2Bab)\sin(3dx+3c)+3Bb^2\sin(4dx+4c)+24(4Aa^2+3Ab^2+6Bab)\sin(dx+c)}{96d}$
parts	$\frac{(Ab^2+2Bab)(2+\cos^2(dx+c))\sin(dx+c)}{3d} + \frac{(2Aab+Ba^2)\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{\sin(dx+c)Aa^2}{d} + \frac{Bb^2}{d}$
derivativedivides	$Bb^2\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ab^2(2+\cos^2(dx+c))\sin(dx+c)}{3} + \frac{2Bab(2+\cos^2(dx+c))\sin(dx+c)}{3} + \frac{\sin(dx+c)Aa^2}{d}$
default	$Bb^2\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ab^2(2+\cos^2(dx+c))\sin(dx+c)}{3} + \frac{2Bab(2+\cos^2(dx+c))\sin(dx+c)}{3} + \frac{\sin(dx+c)Aa^2}{d}$
risc	$xAab + \frac{a^2Bx}{2} + \frac{3b^2Bx}{8} + \frac{\sin(dx+c)Aa^2}{d} + \frac{3\sin(dx+c)Ab^2}{4d} + \frac{3\sin(dx+c)Bab}{2d} + \frac{Bb^2\sin(4dx+4c)}{32d} + \frac{\sin(dx+c)Aa^2}{d}$
norman	$\frac{(Aab+\frac{1}{2}Ba^2+\frac{3}{8}Bb^2)x+(Aab+\frac{1}{2}Ba^2+\frac{3}{8}Bb^2)x\left(\tan^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(4Aab+2Ba^2+\frac{3}{2}Bb^2)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(4Aab+2Ba^2+\frac{3}{2}Bb^2)x\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{24d}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/96*(24*(2*A*a*b+B*a^2+B*b^2)*sin(2*d*x+2*c)+8*(A*b^2+2*B*a*b)*sin(3*d*x+3*c)+3*B*b^2*sin(4*d*x+4*c)+24*(4*A*a^2+3*A*b^2+6*B*a*b)*sin(d*x+c)+96*x*(A*a*b+1/2*B*a^2+3/8*B*b^2)*d)/d
```

3.224.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.67

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{3(4Ba^2 + 8Aab + 3Bb^2)dx + (6Bb^2 \cos(dx + c))^3 + 24Aa^2 + 32Bab + 16Ab^2 + 8(2Bab + Ab^2) \cos(dx + c)}{24d}$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

output $\frac{1}{24} \cdot (3 \cdot (4B^2a^2 + 8A^2ab + 3B^2b^2) \cdot dx + (6B^2b^2 \cos(dx + c)^3 + 24A^2a^2 + 32B^2a^2b + 16A^2b^2 + 8(2B^2ab + A^2b^2) \cos(dx + c)^2 + 3(4B^2a^2 + 8A^2ab + 3B^2b^2) \cos(dx + c)) \cdot \sin(dx + c)) / d$

3.224.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 338 vs. $2(162) = 324$.

Time = 0.20 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.99

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aa^2 \sin(c+dx)}{d} + Aabx \sin^2(c + dx) + Aabx \cos^2(c + dx) + \frac{Aab \sin(c+dx) \cos(c+dx)}{d} + \frac{2Ab^2 \sin^3(c+dx)}{3d} + \frac{Ab^2 \sin(c+dx) \cos^3(c+dx)}{3d} \\ x(A + B \cos(c))(a + b \cos(c))^2 \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Piecewise((A**2*sin(c + d*x)/d + A*a*b*x*sin(c + d*x)**2 + A*a*b*x*cos(c + d*x)**2 + A*a*b*sin(c + d*x)*cos(c + d*x)/d + 2*A*b**2*sin(c + d*x)**3/(3*d) + A*b**2*sin(c + d*x)*cos(c + d*x)**2/d + B*a**2*x*sin(c + d*x)**2/2 + B*a**2*x*cos(c + d*x)**2/2 + B*a**2*sin(c + d*x)*cos(c + d*x)/(2*d) + 4*B*a*b*sin(c + d*x)**3/(3*d) + 2*B*a*b*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*b**2*x*sin(c + d*x)**4/8 + 3*B*b**2*x*cos(c + d*x)**2*cos(c + d*x)**2/4 + 3*B*b**2*x*cos(c + d*x)**4/8 + 3*B*b**2*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 5*B*b**2*sin(c + d*x)*cos(c + d*x)**3/(8*d), Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**2*cos(c), True))`

3.224.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.84

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{24(2dx + 2c + \sin(2dx + 2c))Ba^2 + 48(2dx + 2c + \sin(2dx + 2c))Aab - 64(\sin(dx + c)^3 - 3\sin(dx + c)\cos^2(dx + c))}{3d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")`

3.224. $\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$

output $1/96*(24*(2*d*x + 2*c + \sin(2*d*x + 2*c))*B*a^2 + 48*(2*d*x + 2*c + \sin(2*d*x + 2*c))*A*a*b - 64*(\sin(d*x + c))^3 - 3*\sin(d*x + c))*B*a*b - 32*(\sin(d*x + c))^3 - 3*\sin(d*x + c))*A*b^2 + 3*(12*d*x + 12*c + \sin(4*d*x + 4*c) + 8*\sin(2*d*x + 2*c))*B*b^2 + 96*A*a^2*\sin(d*x + c))/d$

3.224.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.73

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{Bb^2 \sin(4dx + 4c)}{32d} + \frac{1}{8}(4Ba^2 + 8Aab + 3Bb^2)x + \frac{(2Bab + Ab^2) \sin(3dx + 3c)}{12d}$$

$$+ \frac{(Ba^2 + 2Aab + Bb^2) \sin(2dx + 2c)}{4d} + \frac{(4Aa^2 + 6Bab + 3Ab^2) \sin(dx + c)}{4d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")`

output $1/32*B*b^2*\sin(4*d*x + 4*c)/d + 1/8*(4*B*a^2 + 8*A*a*b + 3*B*b^2)*x + 1/12*(2*B*a*b + A*b^2)*\sin(3*d*x + 3*c)/d + 1/4*(B*a^2 + 2*A*a*b + B*b^2)*\sin(2*d*x + 2*c)/d + 1/4*(4*A*a^2 + 6*B*a*b + 3*A*b^2)*\sin(d*x + c)/d$

3.224.9 Mupad [B] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.99

$$\int \cos(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{B a^2 x}{2} + \frac{3 B b^2 x}{8} + \frac{A a^2 \sin(c + dx)}{d} + \frac{3 A b^2 \sin(c + dx)}{4 d}$$

$$+ A a b x + \frac{B a^2 \sin(2 c + 2 d x)}{4 d} + \frac{A b^2 \sin(3 c + 3 d x)}{12 d}$$

$$+ \frac{B b^2 \sin(2 c + 2 d x)}{4 d} + \frac{B b^2 \sin(4 c + 4 d x)}{32 d} + \frac{3 B a b \sin(c + dx)}{2 d}$$

$$+ \frac{A a b \sin(2 c + 2 d x)}{2 d} + \frac{B a b \sin(3 c + 3 d x)}{6 d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2,x)`

output `(B*a^2*x)/2 + (3*B*b^2*x)/8 + (A*a^2*sin(c + d*x))/d + (3*A*b^2*sin(c + d*x))/(4*d) + A*a*b*x + (B*a^2*sin(2*c + 2*d*x))/(4*d) + (A*b^2*sin(3*c + 3*d*x))/(12*d) + (B*b^2*sin(2*c + 2*d*x))/(4*d) + (B*b^2*sin(4*c + 4*d*x))/(32*d) + (3*B*a*b*sin(c + d*x))/(2*d) + (A*a*b*sin(2*c + 2*d*x))/(2*d) + (B*a*b*sin(3*c + 3*d*x))/(6*d)`

3.225 $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$

3.225.1 Optimal result	2082
3.225.2 Mathematica [A] (verified)	2082
3.225.3 Rubi [A] (verified)	2083
3.225.4 Maple [A] (verified)	2084
3.225.5 Fracas [A] (verification not implemented)	2085
3.225.6 Sympy [A] (verification not implemented)	2085
3.225.7 Maxima [A] (verification not implemented)	2086
3.225.8 Giac [A] (verification not implemented)	2086
3.225.9 Mupad [B] (verification not implemented)	2087

3.225.1 Optimal result

Integrand size = 23, antiderivative size = 107

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx = \frac{1}{2} (2a^2 A + Ab^2 + 2abB) x + \frac{2(3aAb + a^2 B + b^2 B) \sin(c + dx)}{3d} + \frac{b(3Ab + 2aB) \cos(c + dx) \sin(c + dx)}{6d} + \frac{B(a + b \cos(c + dx))^2 \sin(c + dx)}{3d}$$

```
output 1/2*(2*A*a^2+A*b^2+2*B*a*b)*x+2/3*(3*A*a*b+B*a^2+B*b^2)*sin(d*x+c)/d+1/6*b*(3*A*b+2*B*a)*cos(d*x+c)*sin(d*x+c)/d+1/3*B*(a+b*cos(d*x+c))^2*sin(d*x+c)/d
```

3.225.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.84

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx = \frac{6(2a^2 A + Ab^2 + 2abB) (c + dx) + 3(8aAb + 4a^2 B + 3b^2 B) \sin(c + dx) + 3b(Ab + 2aB) \sin(2(c + dx))}{12d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(6*(2*a^2*A + A*b^2 + 2*a*b*B)*(c + d*x) + 3*(8*a*A*b + 4*a^2*B + 3*b^2*B)*Sin[c + d*x] + 3*b*(A*b + 2*a*B)*Sin[2*(c + d*x)] + b^2*B*Sin[3*(c + d*x)])/(12*d)`

3.225.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {3042, 3232, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right) \right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx$$

$$\downarrow \text{3232}$$

$$\frac{1}{3} \int (a + b \cos(c + dx))(3aA + 2bB + (3Ab + 2aB) \cos(c + dx)) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^2}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \int \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right) \right) \left(3aA + 2bB + (3Ab + 2aB) \sin\left(c + dx + \frac{\pi}{2}\right) \right) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^2}{3d}$$

$$\downarrow \text{3213}$$

$$\frac{1}{3} \left(\frac{2(a^2B + 3aAb + b^2B) \sin(c + dx)}{d} + \frac{3}{2} x(2a^2A + 2abB + Ab^2) + \frac{b(2aB + 3Ab) \sin(c + dx) \cos(c + dx)}{2d} \right) + \frac{B \sin(c + dx)(a + b \cos(c + dx))^2}{3d}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output $(B*(a + b*\text{Cos}[c + d*x])^2*\text{Sin}[c + d*x])/(3*d) + ((3*(2*a^2*A + A*b^2 + 2*a*b*B)*x)/2 + (2*(3*a*A*b + a^2*B + b^2*B)*\text{Sin}[c + d*x])/d + (b*(3*A*b + 2*a*B)*\text{Cos}[c + d*x]*\text{Sin}[c + d*x])/(2*d))/3$

3.225.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

3.225.4 Maple [A] (verified)

Time = 2.08 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.82

method	result
parallelrisch	$\frac{(3A b^2 + 6Bab) \sin(2dx + 2c) + B \sin(3dx + 3c)b^2 + (24Aab + 12B a^2 + 9B b^2) \sin(dx + c) + 12x(A a^2 + \frac{1}{2}A b^2 + Bab)d}{12d}$
parts	$a^2 x A + \frac{(A b^2 + 2Bab) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d} + \frac{(2Aab + B a^2) \sin(dx+c)}{d} + \frac{B b^2 (2 + \cos^2(dx+c)) \sin(dx+c)}{3d}$
derivativedivides	$\frac{B b^2 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + \frac{A b^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 2Bab \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 2A \sin(dx+c)a}{d}$
default	$\frac{B b^2 (2 + \cos^2(dx+c)) \sin(dx+c)}{3} + \frac{A b^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 2Bab \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + 2A \sin(dx+c)a}{d}$
risch	$a^2 x A + \frac{x A b^2}{2} + x B a b + \frac{2 \sin(dx+c) A a b}{d} + \frac{\sin(dx+c) B a^2}{d} + \frac{3 b^2 B \sin(dx+c)}{4d} + \frac{\sin(3dx+3c) B b^2}{12d} + \sin(dx+c)$
norman	$\frac{(A a^2 + \frac{1}{2}A b^2 + Bab)x + (A a^2 + \frac{1}{2}A b^2 + Bab)x \left(\tan^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (3A a^2 + \frac{3}{2}A b^2 + 3Bab)x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (3A a^2 + \frac{3}{2}A b^2 + 3Bab)x}{12d}$

3.225. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$

input `int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output $\frac{1}{12} * ((3 * A * b^2 + 6 * B * a * b) * \sin(2 * d * x + 2 * c) + B * \sin(3 * d * x + 3 * c) * b^2 + (24 * A * a * b + 12 * B * a^2 + 9 * B * b^2) * \sin(d * x + c) + 12 * x * (A * a^2 + 1/2 * A * b^2 + B * a * b) * d) / d$

3.225.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.79

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{3(2Aa^2 + 2Bab + Ab^2)dx + (2Bb^2 \cos(dx + c)^2 + 6Ba^2 + 12Aab + 4Bb^2 + 3(2Bab + Ab^2) \cos(dx + c))}{6d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output $\frac{1}{6} * (3 * (2 * A * a^2 + 2 * B * a * b + A * b^2) * d * x + (2 * B * b^2 * \cos(d * x + c)^2 + 6 * B * a^2 + 12 * A * a * b + 4 * B * b^2 + 3 * (2 * B * a * b + A * b^2) * \cos(d * x + c)) * \sin(d * x + c)) / d$

3.225.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.86

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \begin{cases} Aa^2x + \frac{2Aab \sin(c+dx)}{d} + \frac{Ab^2x \sin^2(c+dx)}{2} + \frac{Ab^2x \cos^2(c+dx)}{2} + \frac{Ab^2 \sin(c+dx) \cos(c+dx)}{2d} + \frac{Ba^2 \sin(c+dx)}{d} + Babx \sin^2(c+dx) \\ x(A + B \cos(c)) (a + b \cos(c))^2 \end{cases}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Piecewise((A*a**2*x + 2*A*a*b*sin(c + d*x)/d + A*b**2*x*sin(c + d*x)**2/2 + A*b**2*x*cos(c + d*x)**2/2 + A*b**2*sin(c + d*x)*cos(c + d*x)/(2*d) + B*a**2*sin(c + d*x)/d + B*a*b*x*sin(c + d*x)**2 + B*a*b*x*cos(c + d*x)**2 + B*a*b*sin(c + d*x)*cos(c + d*x)/d + 2*B*b**2*sin(c + d*x)**3/(3*d) + B*b**2*sin(c + d*x)*cos(c + d*x)**2/d, Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**2, True))`

3.225.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.01

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= \frac{12(dx + c)Aa^2 + 6(2dx + 2c + \sin(2dx + 2c))Bab + 3(2dx + 2c + \sin(2dx + 2c))Ab^2 - 4(\sin(dx + c))^3 - 3\sin(dx + c)Bb^2 + 12Ba^2\sin(dx + c) + 24Aab\sin(dx + c)}{12d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="maxima")`output `1/12*(12*(d*x + c)*A*a^2 + 6*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a*b + 3*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*b^2 - 4*(sin(d*x + c))^3 - 3*sin(d*x + c))*B*b^2 + 12*B*a^2*sin(d*x + c) + 24*A*a*b*sin(d*x + c))/d`**3.225.8 Giac [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.87

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx = \frac{Bb^2 \sin(3dx + 3c)}{12d}$$

$$+ \frac{1}{2} (2Aa^2 + 2Bab + Ab^2)x$$

$$+ \frac{(2Bab + Ab^2) \sin(2dx + 2c)}{4d}$$

$$+ \frac{(4Ba^2 + 8Aab + 3Bb^2) \sin(dx + c)}{4d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="giac")`output `1/12*B*b^2*sin(3*d*x + 3*c)/d + 1/2*(2*A*a^2 + 2*B*a*b + A*b^2)*x + 1/4*(2*B*a*b + A*b^2)*sin(2*d*x + 2*c)/d + 1/4*(4*B*a^2 + 8*A*a*b + 3*B*b^2)*sin(d*x + c)/d`

3.225.9 Mupad [B] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.07

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$= A a^2 x + \frac{A b^2 x}{2} + \frac{B a^2 \sin(c + dx)}{d} + \frac{3 B b^2 \sin(c + dx)}{4 d} + B a b x + \frac{A b^2 \sin(2c + 2dx)}{4 d}$$

$$+ \frac{B b^2 \sin(3c + 3dx)}{12 d} + \frac{2 A a b \sin(c + dx)}{d} + \frac{B a b \sin(2c + 2dx)}{2 d}$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2,x)`output `A*a^2*x + (A*b^2*x)/2 + (B*a^2*sin(c + d*x))/d + (3*B*b^2*sin(c + d*x))/(4*d) + B*a*b*x + (A*b^2*sin(2*c + 2*d*x))/(4*d) + (B*b^2*sin(3*c + 3*d*x))/(12*d) + (2*A*a*b*sin(c + d*x))/d + (B*a*b*sin(2*c + 2*d*x))/(2*d)`

3.226 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec(c+dx) dx$

3.226.1 Optimal result	2088
3.226.2 Mathematica [A] (verified)	2088
3.226.3 Rubi [A] (verified)	2089
3.226.4 Maple [A] (verified)	2091
3.226.5 Fricas [A] (verification not implemented)	2092
3.226.6 Sympy [F]	2092
3.226.7 Maxima [A] (verification not implemented)	2093
3.226.8 Giac [B] (verification not implemented)	2093
3.226.9 Mupad [B] (verification not implemented)	2094

3.226.1 Optimal result

Integrand size = 29, antiderivative size = 86

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{1}{2} (4aAb + 2a^2B + b^2B) x + \frac{a^2 A \operatorname{arctanh}(\sin(c + dx))}{d} \\ & \quad + \frac{b(2Ab + 3aB) \sin(c + dx)}{2d} + \frac{bB(a + b \cos(c + dx)) \sin(c + dx)}{2d} \end{aligned}$$

output `1/2*(4*A*a*b+2*B*a^2+B*b^2)*x+a^2*A*arctanh(sin(d*x+c))/d+1/2*b*(2*A*b+3*B*a)*sin(d*x+c)/d+1/2*b*B*(a+b*cos(d*x+c))*sin(d*x+c)/d`

3.226.2 Mathematica [A] (verified)

Time = 0.94 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.40

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\ &= \frac{2(4aAb + 2a^2B + b^2B) (c + dx) - 4a^2 A \log \left(\cos \left(\frac{1}{2}(c + dx) \right) - \sin \left(\frac{1}{2}(c + dx) \right) \right) + 4a^2 A \log \left(\cos \left(\frac{1}{2}(c + dx) \right) + \sin \left(\frac{1}{2}(c + dx) \right) \right)}{4d} \end{aligned}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output $(2*(4*a*A*b + 2*a^2*B + b^2*B)*(c + d*x) - 4*a^2*A*\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] + 4*a^2*A*\text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]] + 4*b*(A*b + 2*a*B)*\text{Sin}[c + d*x] + b^2*B*\text{Sin}[2*(c + d*x)])/(4*d)$

3.226.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 3469, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx \\ & \quad \downarrow 3469 \\ & \frac{1}{2} \int (2Aa^2 + b(2Ab + 3aB) \cos^2(c + dx) + (2Ba^2 + 4Aba + b^2B) \cos(c + dx)) \sec(c + dx) dx + \\ & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d} \\ & \quad \downarrow 3042 \\ & \frac{1}{2} \int \frac{2Aa^2 + b(2Ab + 3aB) \sin(c + dx + \frac{\pi}{2})^2 + (2Ba^2 + 4Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \\ & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d} \\ & \quad \downarrow 3502 \\ & \frac{1}{2} \left(\int (2Aa^2 + (2Ba^2 + 4Aba + b^2B) \cos(c + dx)) \sec(c + dx) dx + \frac{b(3aB + 2Ab) \sin(c + dx)}{d} \right) + \\ & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d} \\ & \quad \downarrow 3042 \\ & \frac{1}{2} \left(\int \frac{2Aa^2 + (2Ba^2 + 4Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{b(3aB + 2Ab) \sin(c + dx)}{d} \right) + \\ & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d} \end{aligned}$$

3.226. $\int (a + b \cos(c + dx))^2(A + B \cos(c + dx)) \sec(c + dx) dx$

$$\begin{aligned}
 & \downarrow \text{3214} \\
 & \frac{1}{2} \left(2a^2 A \int \sec(c + dx) dx + x(2a^2 B + 4aAb + b^2 B) + \frac{b(3aB + 2Ab) \sin(c + dx)}{d} \right) + \\
 & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d} \\
 & \downarrow \text{3042} \\
 & \frac{1}{2} \left(2a^2 A \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + x(2a^2 B + 4aAb + b^2 B) + \frac{b(3aB + 2Ab) \sin(c + dx)}{d} \right) + \\
 & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d} \\
 & \downarrow \text{4257} \\
 & \frac{1}{2} \left(\frac{2a^2 A \operatorname{arctanh}(\sin(c + dx))}{d} + x(2a^2 B + 4aAb + b^2 B) + \frac{b(3aB + 2Ab) \sin(c + dx)}{d} \right) + \\
 & \quad \frac{bB \sin(c + dx)(a + b \cos(c + dx))}{2d}
 \end{aligned}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(b*B*(a + b*Cos[c + d*x])*Sin[c + d*x])/((2*d) + ((4*a*A*b + 2*a^2*B + b^2*B)*x + (2*a^2*A*ArcTanh[Sin[c + d*x]])/d + (b*(2*A*b + 3*a*B)*Sin[c + d*x])/d)/2`

3.226.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.226.4 Maple [A] (verified)

Time = 1.53 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{A a^2 \ln(\sec(dx+c)+\tan(dx+c))+B a^2(dx+c)+2Aab(dx+c)+2B \sin(dx+c)ab+A \sin(dx+c)b^2+B b^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2}\right)}{d}$
default	$\frac{A a^2 \ln(\sec(dx+c)+\tan(dx+c))+B a^2(dx+c)+2Aab(dx+c)+2B \sin(dx+c)ab+A \sin(dx+c)b^2+B b^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2}\right)}{d}$
parts	$\frac{A a^2 \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{(A b^2+2Bab) \sin(dx+c)}{d} + \frac{(2Aab+B a^2)(dx+c)}{d} + \frac{B b^2 \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx+c}{2}\right)}{d}$
parallelrisc	$\frac{-4A a^2 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 4A a^2 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + B \sin(2dx+2c)b^2 + (4A b^2 + 8Bab) \sin(dx+c) + 8xd(Aab + \frac{1}{2}B b^2)}{4d}$
risc	$2x Aab + a^2 Bx + \frac{b^2 Bx}{2} - \frac{ie^{i(dx+c)} A b^2}{2d} - \frac{ie^{i(dx+c)} Bab}{d} + \frac{ie^{-i(dx+c)} A b^2}{2d} + \frac{ie^{-i(dx+c)} Bab}{d} + \frac{A a^2 \ln(e^{i(dx+c)} + \tan(dx+c))}{d}$
norman	$\frac{(2Aab+B a^2+\frac{1}{2}B b^2)x+(2Aab+B a^2+\frac{1}{2}B b^2)x\left(\tan^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(6Aab+3B a^2+\frac{3}{2}B b^2)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(6Aab+3B a^2+\frac{3}{2}B b^2)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right))}$

3.226. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)
```

```
output 1/d*(A*a^2*ln(sec(d*x+c)+tan(d*x+c))+B*a^2*(d*x+c)+2*A*a*b*(d*x+c)+2*B*sin(d*x+c)*a*b+A*sin(d*x+c)*b^2+B*b^2*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c))
```

3.226.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.01

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{Aa^2 \log(\sin(dx + c) + 1) - Aa^2 \log(-\sin(dx + c) + 1) + (2Ba^2 + 4Aab + Bb^2)dx + (Bb^2 \cos(dx + c))}{2d}$$

```
input integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")
```

```
output 1/2*(A*a^2*log(sin(d*x + c) + 1) - A*a^2*log(-sin(d*x + c) + 1) + (2*B*a^2 + 4*A*a*b + B*b^2)*d*x + (B*b^2*cos(d*x + c) + 4*B*a*b + 2*A*b^2)*sin(d*x + c))/d
```

3.226.6 Sympy [F]

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^2 \sec(c + dx) dx$$

```
input integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c),x)
```

```
output Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**2*sec(c + d*x), x)
```

3.226.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.07

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{4(dx + c)Ba^2 + 8(dx + c)Aab + (2dx + 2c + \sin(2dx + 2c))Bb^2 + 4Aa^2 \log(\sec(dx + c) + \tan(dx + c)) + 4A^2 \log(\sec(dx + c) + \tan(dx + c))}{4d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `1/4*(4*(d*x + c)*B*a^2 + 8*(d*x + c)*A*a*b + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*b^2 + 4*A*a^2*log(sec(d*x + c) + tan(d*x + c)) + 8*B*a*b*sin(d*x + c) + 4*A*b^2*sin(d*x + c))/d`

3.226.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. 2(80) = 160.

Time = 0.29 (sec) , antiderivative size = 178, normalized size of antiderivative = 2.07

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2Aa^2 \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 2Aa^2 \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right) + (2Ba^2 + 4Aab + Bb^2)(dx + c) + 2A^2 \log\left(\frac{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1}\right) + 2A^2 \log\left(\frac{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1}\right) + 2A^2 \log\left(\frac{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1}\right) + 2A^2 \log\left(\frac{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1}\right)}{2d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `1/2*(2*A*a^2*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 2*A*a^2*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + (2*B*a^2 + 4*A*a*b + B*b^2)*(d*x + c) + 2*(4*B*a*b*tan(1/2*d*x + 1/2*c)^3 + 2*A*b^2*tan(1/2*d*x + 1/2*c)^3 - B*b^2*tan(1/2*d*x + 1/2*c)^3 + 4*B*a*b*tan(1/2*d*x + 1/2*c) + 2*A*b^2*tan(1/2*d*x + 1/2*c) + B*b^2*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^2)/d`

3.226.9 Mupad [B] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.97

$$\begin{aligned}
& \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec(c + dx) dx \\
&= \frac{A b^2 \sin(c + dx)}{d} + \frac{2 A a^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2 B a^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{B b^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{B b^2 \sin(2c + 2dx)}{4d} \\
&+ \frac{2 B a b \sin(c + dx)}{d} + \frac{4 A a b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d}
\end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x),x)`

output `(A*b^2*sin(c + d*x))/d + (2*A*a^2*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (B*b^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (B*b^2*sin(2*c + 2*d*x))/(4*d) + (2*B*a*b*sin(c + d*x))/d + (4*A*a*b*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d`

3.227 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.227.1 Optimal result	2095
3.227.2 Mathematica [A] (verified)	2095
3.227.3 Rubi [A] (verified)	2096
3.227.4 Maple [A] (verified)	2098
3.227.5 Fricas [A] (verification not implemented)	2099
3.227.6 Sympy [F]	2099
3.227.7 Maxima [A] (verification not implemented)	2099
3.227.8 Giac [B] (verification not implemented)	2100
3.227.9 Mupad [B] (verification not implemented)	2100

3.227.1 Optimal result

Integrand size = 31, antiderivative size = 60

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= b(Ab + 2aB)x + \frac{a(2Ab + aB)\operatorname{arctanh}(\sin(c + dx))}{d} + \frac{b^2 B \sin(c + dx)}{d} + \frac{a^2 A \tan(c + dx)}{d}$$

output `b*(A*b+2*B*a)*x+a*(2*A*b+B*a)*arctanh(sin(d*x+c))/d+b^2*B*sin(d*x+c)/d+a^2*A*tan(d*x+c)/d`

3.227.2 Mathematica [A] (verified)

Time = 1.21 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.82

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{b(Ab + 2aB)(c + dx) - a(2Ab + aB) \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + a(2Ab + aB) \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) + b^2 B \sin(c + dx) + a^2 A \tan(c + dx)}{d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `(b*(A*b + 2*a*B)*(c + d*x) - a*(2*A*b + a*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + a*(2*A*b + a*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + b^2*B*Sin[c + d*x] + a^2*A*Tan[c + d*x])/d`

3.227.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3467, 25, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^2(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^2(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3467} \\
 & \frac{a^2A \tan(c+dx)}{d} - \\
 & \int -((b^2B \cos^2(c+dx) + b(Ab+2aB) \cos(c+dx) + a(2Ab+aB)) \sec(c+dx)) dx \\
 & \quad \downarrow \text{25} \\
 & \int (b^2B \cos^2(c+dx) + b(Ab+2aB) \cos(c+dx) + a(2Ab+aB)) \sec(c+dx) dx + \frac{a^2A \tan(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{b^2B \sin(c+dx+\frac{\pi}{2})^2 + b(Ab+2aB) \sin(c+dx+\frac{\pi}{2}) + a(2Ab+aB)}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{a^2A \tan(c+dx)}{d} \\
 & \quad \downarrow \text{3502} \\
 & \int (a(2Ab+aB) + b(Ab+2aB) \cos(c+dx)) \sec(c+dx) dx + \frac{a^2A \tan(c+dx)}{d} + \frac{b^2B \sin(c+dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a(2Ab+aB) + b(Ab+2aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{a^2A \tan(c+dx)}{d} + \frac{b^2B \sin(c+dx)}{d} \\
 & \quad \downarrow \text{3214} \\
 & a(aB+2Ab) \int \sec(c+dx) dx + \frac{a^2A \tan(c+dx)}{d} + bx(2aB+Ab) + \frac{b^2B \sin(c+dx)}{d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.227. $\int (a+b\cos(c+dx))^2(A+B\cos(c+dx)) \sec^2(c+dx) dx$

$$a(aB + 2Ab) \int \csc\left(c + dx + \frac{\pi}{2}\right) dx + \frac{a^2 A \tan(c + dx)}{d} + bx(2aB + Ab) + \frac{b^2 B \sin(c + dx)}{d}$$

↓ 4257

$$\frac{a^2 A \tan(c + dx)}{d} + \frac{a(aB + 2Ab) \operatorname{arctanh}(\sin(c + dx))}{d} + bx(2aB + Ab) + \frac{b^2 B \sin(c + dx)}{d}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `b*(A*b + 2*a*B)*x + (a*(2*A*b + a*B)*ArcTanh[Sin[c + d*x]])/d + (b^2*B*Sin[c + d*x])/d + (a^2*A*Tan[c + d*x])/d`

3.227.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`


```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.227.4 Maple [A] (verified)

Time = 2.72 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.32

method	result
parts	$\frac{a^2 A \tan(dx+c)}{d} + \frac{(A b^2 + 2 B a b)(dx+c)}{d} + \frac{(2 A a b + B a^2) \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{b^2 B \sin(dx+c)}{d}$
derivativedivides	$\frac{A a^2 \tan(dx+c) + B a^2 \ln(\sec(dx+c) + \tan(dx+c)) + 2 A a b \ln(\sec(dx+c) + \tan(dx+c)) + 2 B a b(dx+c) + A b^2(dx+c) + B \sin(dx+c)}{d}$
default	$\frac{A a^2 \tan(dx+c) + B a^2 \ln(\sec(dx+c) + \tan(dx+c)) + 2 A a b \ln(\sec(dx+c) + \tan(dx+c)) + 2 B a b(dx+c) + A b^2(dx+c) + B \sin(dx+c)}{d}$
parallelrisch	$\frac{(-4 A a b - 2 B a^2) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + (4 A a b + 2 B a^2) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + B \sin(2 dx + 2 c) b^2 + 2 A a b \sin(dx+c)}{2 d \cos(dx+c)}$
risch	$x A b^2 + 2 x B a b - \frac{i B b^2 e^{i(dx+c)}}{2 d} + \frac{i B b^2 e^{-i(dx+c)}}{2 d} + \frac{2 i A a^2}{d(e^{2i(dx+c)} + 1)} + \frac{2 a \ln(e^{i(dx+c)} + i) A b}{d} + \frac{a^2 \ln(e^{i(dx+c)} - i)}{d}$
norman	$\frac{(-A b^2 - 2 B a b)x + (-2 A b^2 - 4 B a b)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (A b^2 + 2 B a b)x \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (2 A b^2 + 4 B a b)x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))}$

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBO
SE)
```

```
output a^2*A*tan(d*x+c)/d+(A*b^2+2*B*a*b)/d*(d*x+c)+(2*A*a*b+B*a^2)/d*ln(sec(d*x+
c)+tan(d*x+c))+b^2*B*sin(d*x+c)/d
```

3.227.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.95

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2(2Bab + Ab^2)dx \cos(dx + c) + (Ba^2 + 2Aab) \cos(dx + c) \log(\sin(dx + c) + 1) - (Ba^2 + 2Aab) \cos(dx + c) \log(\sin(dx + c) - 1) + 2(Bb^2 \cos(dx + c) + Aa^2) \sin(dx + c)}{2d \cos(dx + c)}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `1/2*(2*(2*B*a*b + A*b^2)*d*x*cos(d*x + c) + (B*a^2 + 2*A*a*b)*cos(d*x + c)*log(sin(d*x + c) + 1) - (B*a^2 + 2*A*a*b)*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*(B*b^2*cos(d*x + c) + A*a^2)*sin(d*x + c))/(d*cos(d*x + c))`

3.227.6 Sympy [F]

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^2 \sec^2(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**2*sec(c + d*x)**2, x)`

3.227.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.72

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{4(dx + c)Bab + 2(dx + c)Ab^2 + Ba^2(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 2Aab(\log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1))}{2d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output $\frac{1}{2}*(4*(d*x + c)*B*a*b + 2*(d*x + c)*A*b^2 + B*a^2*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 2*A*a*b*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 2*B*b^2*\sin(d*x + c) + 2*A*a^2*\tan(d*x + c))/d$

3.227.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. $2(60) = 120$.

Time = 0.30 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.53

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{(2 Bab + Ab^2)(dx + c) + (Ba^2 + 2 Aab) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - (Ba^2 + 2 Aab) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right)}{d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output $\frac{((2*B*a*b + A*b^2)*(d*x + c) + (B*a^2 + 2*A*a*b)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - (B*a^2 + 2*A*a*b)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(A*a^2*\tan(1/2*d*x + 1/2*c)^3 - B*b^2*\tan(1/2*d*x + 1/2*c)^3 + A*a^2*\tan(1/2*d*x + 1/2*c) + B*b^2*\tan(1/2*d*x + 1/2*c)))/(\tan(1/2*d*x + 1/2*c)^4 - 1)}{d}$

3.227.9 Mupad [B] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.82

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{A a^2 \tan(c + dx)}{d} + \frac{2 A b^2 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} + \frac{B b^2 \sin(2c + 2dx)}{2d \cos(c + dx)}$$

$$+ \frac{4 B a b \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right)}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right)}{d} - \frac{B a^2 \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right) \operatorname{li}}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right) 2i}{d} - \frac{A a b \operatorname{atan} \left(\frac{\sin \left(\frac{c}{2} + \frac{dx}{2} \right) \operatorname{li}}{\cos \left(\frac{c}{2} + \frac{dx}{2} \right)} \right) 4i}{d}$$

3.227. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^2(c + dx) dx$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^2,x)`

output `(A*a^2*tan(c + d*x))/d + (2*A*b^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d - (B*a^2*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/d + (B*b^2*sin(2*c + 2*d*x))/(2*d*cos(c + d*x)) - (A*a*b*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*4i)/d + (4*B*a*b*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d`

3.228 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.228.1 Optimal result	2102
3.228.2 Mathematica [A] (verified)	2102
3.228.3 Rubi [A] (verified)	2103
3.228.4 Maple [A] (verified)	2105
3.228.5 Fricas [A] (verification not implemented)	2106
3.228.6 Sympy [F]	2107
3.228.7 Maxima [A] (verification not implemented)	2107
3.228.8 Giac [B] (verification not implemented)	2107
3.228.9 Mupad [B] (verification not implemented)	2108

3.228.1 Optimal result

Integrand size = 31, antiderivative size = 80

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= b^2 Bx + \frac{(a^2 A + 2Ab^2 + 4abB) \operatorname{arctanh}(\sin(c + dx))}{2d}$$

$$+ \frac{a(2Ab + aB) \tan(c + dx)}{d} + \frac{a^2 A \sec(c + dx) \tan(c + dx)}{2d}$$

```
output b^2*B*x+1/2*(A*a^2+2*A*b^2+4*B*a*b)*arctanh(sin(d*x+c))/d+a*(2*A*b+B*a)*tan(d*x+c)/d+1/2*a^2*A*sec(d*x+c)*tan(d*x+c)/d
```

3.228.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.84

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2b^2 Bdx + (a^2 A + 2Ab^2 + 4abB) \operatorname{arctanh}(\sin(c + dx)) + a(4Ab + 2aB + aA \sec(c + dx)) \tan(c + dx)}{2d}$$

```
input Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]
```

output $(2b^2Bdx + (a^2A + 2Ab^2 + 4abB) \operatorname{ArcTanh}[\sin(c + dx)] + a(4Ab + 2aB + aA \operatorname{Sec}[c + dx]) \operatorname{Tan}[c + dx]) / (2d)$

3.228.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3467, 25, 3042, 3500, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow 3467 \\
 & \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d} - \\
 & \frac{1}{2} \int -((2b^2 B \cos^2(c + dx) + (Aa^2 + 4bBa + 2Ab^2) \cos(c + dx) + 2a(2Ab + aB)) \sec^2(c + dx)) dx \\
 & \quad \downarrow 25 \\
 & \frac{1}{2} \int (2b^2 B \cos^2(c + dx) + (Aa^2 + 4bBa + 2Ab^2) \cos(c + dx) + 2a(2Ab + aB)) \sec^2(c + dx) dx + \\
 & \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow 3042 \\
 & \frac{1}{2} \int \frac{2b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (Aa^2 + 4bBa + 2Ab^2) \sin(c + dx + \frac{\pi}{2}) + 2a(2Ab + aB)}{\sin(c + dx + \frac{\pi}{2})^2} dx + \\
 & \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d} \\
 & \quad \downarrow 3500 \\
 & \frac{1}{2} \left(\int (Aa^2 + 4bBa + 2Ab^2 + 2b^2 B \cos(c + dx)) \sec(c + dx) dx + \frac{2a(aB + 2Ab) \tan(c + dx)}{d} \right) + \\
 & \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{1}{2} \left(\int \frac{Aa^2 + 4bBa + 2Ab^2 + 2b^2B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a(aB + 2Ab) \tan(c + dx)}{d} \right) + \\
& \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d} \\
& \downarrow \text{3214} \\
& \frac{1}{2} \left((a^2 A + 4abB + 2Ab^2) \int \sec(c + dx) dx + \frac{2a(aB + 2Ab) \tan(c + dx)}{d} + 2b^2 Bx \right) + \\
& \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d} \\
& \downarrow \text{3042} \\
& \frac{1}{2} \left((a^2 A + 4abB + 2Ab^2) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{2a(aB + 2Ab) \tan(c + dx)}{d} + 2b^2 Bx \right) + \\
& \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d} \\
& \downarrow \text{4257} \\
& \frac{1}{2} \left(\frac{(a^2 A + 4abB + 2Ab^2) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{2a(aB + 2Ab) \tan(c + dx)}{d} + 2b^2 Bx \right) + \\
& \quad \frac{a^2 A \tan(c + dx) \sec(c + dx)}{2d}
\end{aligned}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a^2*A*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (2*b^2*B*x + ((a^2*A + 2*A*b^2 + 4*a*b*B)*ArcTanh[Sin[c + d*x]])/d + (2*a*(2*A*b + a*B)*Tan[c + d*x])/d)/2`

3.228.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.228.4 Maple [A] (verified)

Time = 3.72 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.30

method	result
parts	$\frac{A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d} + \frac{(A b^2+2Bab) \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{(2Aab+B a^2) \tan(dx+c)}{d}$
derivatividedives	$\frac{A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) + B a^2 \tan(dx+c) + 2Aab \tan(dx+c) + 2Bab \ln(\sec(dx+c)+\tan(dx+c))}{d}$
default	$\frac{A a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) + B a^2 \tan(dx+c) + 2Aab \tan(dx+c) + 2Bab \ln(\sec(dx+c)+\tan(dx+c))}{d}$
parallelrisch	$\frac{-(1+\cos(2dx+2c))(A a^2+2A b^2+4Bab) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right) + (1+\cos(2dx+2c))(A a^2+2A b^2+4Bab) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{2d(1+\cos(2dx+2c))}$
risch	$b^2 Bx - \frac{ia(Aa e^{3i(dx+c)} - 4Ab e^{2i(dx+c)} - 2Ba e^{2i(dx+c)} - aA e^{i(dx+c)} - 4Ab - 2Ba)}{d(e^{2i(dx+c)} + 1)^2} - \frac{A a^2 \ln(e^{i(dx+c)} - i)}{2d} - \frac{\ln(e^{i(dx+c)} + i)}{2d}$
norman	$b^2 Bx + \frac{a(aA - 4Ab - 2Ba) \left(\tan^9\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{d} + \frac{a(aA + 4Ab + 2Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + b^2 Bx \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + b^2 Bx \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + b^2 Bx \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right)$

input `int((a+cos(d*x+c))*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `A*a^2/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^2+2*B*a*b)/d*ln(sec(d*x+c)+tan(d*x+c))+(2*A*a*b+B*a^2)/d*tan(d*x+c)+B*b^2/d*(d*x+c)`

3.228.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.70

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4 B b^2 dx \cos(dx + c)^2 + (A a^2 + 4 B a b + 2 A b^2) \cos(dx + c)^2 \log(\sin(dx + c) + 1) - (A a^2 + 4 B a b + 2 A b^2) \cos(dx + c)^2 \log(\sin(dx + c) - 1) + 2(A a^2 + 2(B a^2 + 2 A a b) \cos(dx + c)) \sin(dx + c)}{4 d \cos(dx + c)}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fricas")`

output `1/4*(4*B*b^2*d*x*cos(d*x + c)^2 + (A*a^2 + 4*B*a*b + 2*A*b^2)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A*a^2 + 4*B*a*b + 2*A*b^2)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(A*a^2 + 2*(B*a^2 + 2*A*a*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.228. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.228.6 Sympy [F]

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^2 \sec^3(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**2*sec(c + d*x)**3, x)`

3.228.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.75

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4(dx + c)Bb^2 - Aa^2 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) + 4Bab(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1))}{1}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `1/4*(4*(d*x + c)*B*b^2 - A*a^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 4*B*a*b*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 2*A*b^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 4*B*a^2*tan(d*x + c) + 8*A*a*b*tan(d*x + c))/d`

3.228.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 190 vs. $2(76) = 152$.

Time = 0.31 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.38

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2(dx + c)Bb^2 + (Aa^2 + 4Bab + 2Ab^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - (Aa^2 + 4Bab + 2Ab^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{1}$$

3.228. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output $\frac{1}{2}*(2*(d*x + c)*B*b^2 + (A*a^2 + 4*B*a*b + 2*A*b^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - (A*a^2 + 4*B*a*b + 2*A*b^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) + 2*(A*a^2*\tan(1/2*d*x + 1/2*c)^3 - 2*B*a^2*\tan(1/2*d*x + 1/2*c)^3 - 4*A*a*b*\tan(1/2*d*x + 1/2*c)^3 + A*a^2*\tan(1/2*d*x + 1/2*c) + 2*B*a^2*\tan(1/2*d*x + 1/2*c) + 4*A*a*b*\tan(1/2*d*x + 1/2*c)))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^2)/d$

3.228.9 Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 176, normalized size of antiderivative = 2.20

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2 \left(\frac{A a^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{2} + A b^2 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + B b^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + 2 B a b \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \right)}{d \left(\frac{\cos(2c + 2dx)}{2} + \frac{1}{2} \right)} + \frac{B a^2 \sin(2c + 2dx)}{2} + \frac{A a^2 \sin(c + dx)}{2} + A a b \sin(2c + 2dx)$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^3,x)`

output $(2*((A*a^2*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 + A*b^2*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + B*b^2*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + 2*B*a*b*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))))/d + ((B*a^2*\sin(2*c + 2*d*x))/2 + (A*a^2*\sin(c + d*x))/2 + A*a*b*\sin(2*c + 2*d*x))/(d*(\cos(2*c + 2*d*x)/2 + 1/2))$

3.229 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.229.1 Optimal result	2109
3.229.2 Mathematica [A] (verified)	2109
3.229.3 Rubi [A] (verified)	2110
3.229.4 Maple [A] (verified)	2113
3.229.5 Fricas [A] (verification not implemented)	2114
3.229.6 Sympy [F]	2114
3.229.7 Maxima [A] (verification not implemented)	2115
3.229.8 Giac [B] (verification not implemented)	2115
3.229.9 Mupad [B] (verification not implemented)	2116

3.229.1 Optimal result

Integrand size = 31, antiderivative size = 116

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{(2aAb + a^2B + 2b^2B) \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{(2a^2A + 3Ab^2 + 6abB) \tan(c + dx)}{3d}$$

$$+ \frac{a(2Ab + aB) \sec(c + dx) \tan(c + dx)}{2d} + \frac{a^2A \sec^2(c + dx) \tan(c + dx)}{3d}$$

```
output 1/2*(2*A*a*b+B*a^2+2*B*b^2)*arctanh(sin(d*x+c))/d+1/3*(2*A*a^2+3*A*b^2+6*B
*a*b)*tan(d*x+c)/d+1/2*a*(2*A*b+B*a)*sec(d*x+c)*tan(d*x+c)/d+1/3*a^2*A*sec
(d*x+c)^2*tan(d*x+c)/d
```

3.229.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.79

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(2aAb + a^2B + 2b^2B) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (3a(2Ab + aB) \sec(c + dx) + 2(3a^2A + 3Ab^2))}{6d}$$

input `Integrate[(a + b*cos[c + d*x])^2*(A + B*cos[c + d*x])*Sec[c + d*x]^4,x]`

output $(3*(2*a*A*b + a^2*B + 2*b^2*B)*ArcTanh[\sin[c + d*x]] + \tan[c + d*x]*(3*a*(2*A*b + a*B)*\sec[c + d*x] + 2*(3*a^2*A + 3*A*b^2 + 6*a*b*B + a^2*A*\tan[c + d*x]^2)))/(6*d)$

3.229.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3467, 25, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3467} \\
 & \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d} - \\
 & \frac{1}{3} \int -((3b^2 B \cos^2(c + dx) + (2Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(2Ab + aB)) \sec^3(c + dx)) dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \int (3b^2 B \cos^2(c + dx) + (2Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(2Ab + aB)) \sec^3(c + dx) dx + \\
 & \quad \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{3b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (2Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + 3a(2Ab + aB)}{\sin(c + dx + \frac{\pi}{2})^3} dx + \\
 & \quad \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d} \\
 & \quad \downarrow \text{3500}
 \end{aligned}$$

$$\frac{1}{3} \left(\frac{1}{2} \int (2(2Aa^2 + 6bBa + 3Ab^2) + 3(Ba^2 + 2Aba + 2b^2B) \cos(c + dx)) \sec^2(c + dx) dx + \frac{3a(aB + 2Ab) \tan(c + dx)}{2d} \right) + \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \int \frac{2(2Aa^2 + 6bBa + 3Ab^2) + 3(Ba^2 + 2Aba + 2b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{3a(aB + 2Ab) \tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d}$$

↓ 3227

$$\frac{1}{3} \left(\frac{1}{2} \left(2(2a^2A + 6abB + 3Ab^2) \int \sec^2(c + dx) dx + 3(a^2B + 2aAb + 2b^2B) \int \sec(c + dx) dx \right) + \frac{3a(aB + 2Ab) \tan(c + dx)}{2d} \right) + \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3(a^2B + 2aAb + 2b^2B) \int \csc(c + dx + \frac{\pi}{2}) dx + 2(2a^2A + 6abB + 3Ab^2) \int \csc(c + dx + \frac{\pi}{2})^2 dx \right) + \frac{3a(aB + 2Ab) \tan(c + dx)}{2d} \right) + \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d}$$

↓ 4254

$$\frac{1}{3} \left(\frac{1}{2} \left(3(a^2B + 2aAb + 2b^2B) \int \csc(c + dx + \frac{\pi}{2}) dx - \frac{2(2a^2A + 6abB + 3Ab^2) \int 1d(-\tan(c + dx))}{d} \right) + \frac{3a(aB + 2Ab) \tan(c + dx)}{2d} \right) + \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d}$$

↓ 24

$$\frac{1}{3} \left(\frac{1}{2} \left(3(a^2B + 2aAb + 2b^2B) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{2(2a^2A + 6abB + 3Ab^2) \tan(c + dx)}{d} \right) + \frac{3a(aB + 2Ab) \tan(c + dx)}{2d} \right) + \frac{a^2 A \tan(c + dx) \sec^2(c + dx)}{3d}$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{3(a^2B + 2aAb + 2b^2B) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{2(2a^2A + 6abB + 3Ab^2) \tan(c + dx)}{d} \right) + \frac{3a(aB + 2Ab) \tan(c + dx)}{3d} \right)$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a^2*A*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((3*a*(2*A*b + a*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((3*(2*a*A*b + a^2*B + 2*b^2*B)*ArcTanh[Sin[c + d*x]])/d + (2*(2*a^2*A + 3*A*b^2 + 6*a*b*B)*Tan[c + d*x])/d)/2)/3`

3.229.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-A*b^2
- a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.229.4 Maple [A] (verified)

Time = 4.52 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.02

method	result
parts	$-\frac{A a^2 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d} + \frac{(A b^2 + 2Bab) \tan(dx+c)}{d} + \frac{(2Aab + B a^2) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c))}{d} \right)}{d}$
derivativedivides	$-\frac{A a^2 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + B a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 2Aab \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
default	$-\frac{A a^2 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + B a^2 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 2Aab \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
parallelrisch	$-\frac{9(Aab + \frac{1}{2} B a^2 + B b^2) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 9(Aab + \frac{1}{2} B a^2 + B b^2) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right)}{3d(\cos(3dx+c))}$
risch	$-\frac{i(6Aab e^{5i(dx+c)} + 3B a^2 e^{5i(dx+c)} - 6A b^2 e^{4i(dx+c)} - 12Bab e^{4i(dx+c)} - 12A a^2 e^{2i(dx+c)} - 12A b^2 e^{2i(dx+c)} - 24Bab e^{2i(dx+c)})}{3d(e^{2i(dx+c)} + 1)^3}$
norman	$-\frac{2(2A a^2 - 2Aab - 2A b^2 - B a^2 - 4Bab) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} - \frac{(2A a^2 - 2Aab + 2A b^2 - B a^2 + 4Bab) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} - \frac{2(2A a^2 + 2Aab - 2A b^2 - B a^2 - 4Bab) \left(\tan^3 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d}$

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBO
SE)
```

$$3.229. \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

output
$$-Aa^2/d*(-2/3-1/3*\sec(dx+c)^2)*\tan(dx+c)+(Ab^2+2Bab)/d*\tan(dx+c)+(2Aab+Ba^2)/d*(1/2*\sec(dx+c)*\tan(dx+c)+1/2*\ln(\sec(dx+c)+\tan(dx+c)))+Bb^2/d*\ln(\sec(dx+c)+\tan(dx+c))$$

3.229.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.29

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3(Ba^2 + 2Aab + 2Bb^2) \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3(Ba^2 + 2Aab + 2Bb^2) \cos(dx + c)^3 \log(-\sin(dx + c) + 1) + 2(2Aa^2 + 2(2Aa^2 + 6Bab + 3Ab^2) \cos(dx + c)^2 + 3(Ba^2 + 2Aab) \cos(dx + c)) \sin(dx + c)}{d \cos(dx + c)^3}$$

input `integrate((a+b*cos(dx+c))^2*(A+B*cos(dx+c))*sec(dx+c)^4,x, algorithm="fricas")`

output
$$1/12*(3*(Ba^2 + 2Aab + 2Bb^2)*\cos(dx + c)^3*\log(\sin(dx + c) + 1) - 3*(Ba^2 + 2Aab + 2Bb^2)*\cos(dx + c)^3*\log(-\sin(dx + c) + 1) + 2*(2Aa^2 + 2*(2Aa^2 + 6Bab + 3Ab^2)*\cos(dx + c)^2 + 3*(Ba^2 + 2Aab)*\cos(dx + c))*\sin(dx + c))/(d*\cos(dx + c)^3)$$

3.229.6 Sympy [F]

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^2 \sec^4(c + dx) dx$$

input `integrate((a+b*cos(dx+c))**2*(A+B*cos(dx+c))*sec(dx+c)**4,x)`

output `Integral((A + B*cos(c + dx))*(a + b*cos(c + dx))**2*sec(c + dx)**4, x)`

3.229.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.48

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{4 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^2 - 3 Ba^2 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) - 6 Aab^2 (2 \sin(dx + c) / (\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) + 6 Bb^2 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 24 B a b \tan(dx + c) + 12 A b^2 \tan(dx + c)}{d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output `1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^2 - 3*B*a^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 6*A*a*b*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 6*B*b^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 24*B*a*b*tan(d*x + c) + 12*A*b^2*tan(d*x + c))/d`

3.229.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. 2(108) = 216.

Time = 0.33 (sec) , antiderivative size = 294, normalized size of antiderivative = 2.53

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{3 (Ba^2 + 2 Aab + 2 Bb^2) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 3 (Ba^2 + 2 Aab + 2 Bb^2) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + 6 Aab^2 (2 \sin(dx + c) / (\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) + 6 Bb^2 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 24 B a b \tan(dx + c) + 12 A b^2 \tan(dx + c)}{d}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")`

output $\frac{1}{6}(3*(B*a^2 + 2*A*a*b + 2*B*b^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(B*a^2 + 2*A*a*b + 2*B*b^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*A*a^2*\tan(1/2*d*x + 1/2*c)^5 - 3*B*a^2*\tan(1/2*d*x + 1/2*c)^5 - 6*A*a*b*\tan(1/2*d*x + 1/2*c)^5 + 12*B*a*b*\tan(1/2*d*x + 1/2*c)^5 + 6*A*b^2*\tan(1/2*d*x + 1/2*c)^5 - 4*A*a^2*\tan(1/2*d*x + 1/2*c)^3 - 24*B*a*b*\tan(1/2*d*x + 1/2*c)^3 - 12*A*b^2*\tan(1/2*d*x + 1/2*c)^3 + 6*A*a^2*\tan(1/2*d*x + 1/2*c) + 3*B*a^2*\tan(1/2*d*x + 1/2*c) + 6*A*a*b*\tan(1/2*d*x + 1/2*c) + 12*B*a*b*\tan(1/2*d*x + 1/2*c) + 6*A*b^2*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^3)/d$

3.229.9 Mupad [B] (verification not implemented)

Time = 3.85 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.96

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{B a^2}{2} + A a b + B b^2\right)}{2 B a^2 + 4 A a b + 4 B b^2}\right) (B a^2 + 2 A a b + 2 B b^2)}{d} - \frac{(2 A a^2 + 2 A b^2 - B a^2 - 2 A a b + 4 B a b) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \left(-\frac{4 A a^2}{3} - 8 B a b - 4 A b^2\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^4,x)`

output $(\operatorname{atanh}((4*\tan(c/2 + (d*x)/2))*((B*a^2)/2 + B*b^2 + A*a*b))/(2*B*a^2 + 4*B*b^2 + 4*A*a*b))*(B*a^2 + 2*B*b^2 + 2*A*a*b))/d - (\tan(c/2 + (d*x)/2)*(2*A*a^2 + 2*A*b^2 + B*a^2 + 2*A*a*b + 4*B*a*b) - \tan(c/2 + (d*x)/2)^3*((4*A*a^2)/3 + 4*A*b^2 + 8*B*a*b) + \tan(c/2 + (d*x)/2)^5*(2*A*a^2 + 2*A*b^2 - B*a^2 - 2*A*a*b + 4*B*a*b))/(d*(3*\tan(c/2 + (d*x)/2)^2 - 3*\tan(c/2 + (d*x)/2)^4 + \tan(c/2 + (d*x)/2)^6 - 1))$

3.230 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.230.1 Optimal result	2117
3.230.2 Mathematica [A] (verified)	2118
3.230.3 Rubi [A] (verified)	2118
3.230.4 Maple [A] (verified)	2122
3.230.5 Fricas [A] (verification not implemented)	2123
3.230.6 Sympy [F(-1)]	2123
3.230.7 Maxima [A] (verification not implemented)	2123
3.230.8 Giac [B] (verification not implemented)	2124
3.230.9 Mupad [B] (verification not implemented)	2125

3.230.1 Optimal result

Integrand size = 31, antiderivative size = 156

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{(3a^2 A + 4Ab^2 + 8abB) \operatorname{arctanh}(\sin(c + dx))}{8d} + \frac{(4aAb + 2a^2 B + 3b^2 B) \tan(c + dx)}{3d}$$

$$+ \frac{(3a^2 A + 4Ab^2 + 8abB) \sec(c + dx) \tan(c + dx)}{8d}$$

$$+ \frac{a(2Ab + aB) \sec^2(c + dx) \tan(c + dx)}{3d} + \frac{a^2 A \sec^3(c + dx) \tan(c + dx)}{4d}$$

output

```
1/8*(3*A*a^2+4*A*b^2+8*B*a*b)*arctanh(sin(d*x+c))/d+1/3*(4*A*a*b+2*B*a^2+3
*B*b^2)*tan(d*x+c)/d+1/8*(3*A*a^2+4*A*b^2+8*B*a*b)*sec(d*x+c)*tan(d*x+c)/d
+1/3*a*(2*A*b+B*a)*sec(d*x+c)^2*tan(d*x+c)/d+1/4*a^2*A*sec(d*x+c)^3*tan(d*
x+c)/d
```

3.230.2 Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.77

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3a^2A + 4Ab^2 + 8abB) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (24(2aAb + a^2B + b^2B) + 3(3a^2A + 4Ab^2 + 8abB))}{24d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(3*(3*a^2*A + 4*A*b^2 + 8*a*b*B)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(24*(2*a*A*b + a^2*B + b^2*B) + 3*(3*a^2*A + 4*A*b^2 + 8*a*b*B))*Sec[c + d*x] + 6*a^2*A*Sec[c + d*x]^3 + 8*a*(2*A*b + a*B)*Tan[c + d*x]^2)/(24*d)`

3.230.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.96, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3467, 25, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3467}$$

$$\frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} - \frac{1}{4} \int -((4b^2 B \cos^2(c + dx) + (3Aa^2 + 8bBa + 4Ab^2) \cos(c + dx) + 4a(2Ab + aB)) \sec^4(c + dx)) dx$$

$$\downarrow \text{25}$$

$$\frac{1}{4} \int (4b^2 B \cos^2(c + dx) + (3Aa^2 + 8bBa + 4Ab^2) \cos(c + dx) + 4a(2Ab + aB)) \sec^4(c + dx) dx + \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{4} \int \frac{4b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (3Aa^2 + 8bBa + 4Ab^2) \sin(c + dx + \frac{\pi}{2}) + 4a(2Ab + aB)}{\sin(c + dx + \frac{\pi}{2})^4} dx + \\ \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3500 \\ \frac{1}{4} \left(\frac{1}{3} \int (3(3Aa^2 + 8bBa + 4Ab^2) + 4(2Ba^2 + 4Aba + 3b^2 B) \cos(c + dx)) \sec^3(c + dx) dx + \frac{4a(aB + 2Ab) \tan(c + dx)}{3d} \right) \\ \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{4} \left(\frac{1}{3} \int \frac{3(3Aa^2 + 8bBa + 4Ab^2) + 4(2Ba^2 + 4Aba + 3b^2 B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{4a(aB + 2Ab) \tan(c + dx) \sec^3(c + dx)}{3d} \right) \\ \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3227 \\ \frac{1}{4} \left(\frac{1}{3} \left(3(3a^2 A + 8abB + 4Ab^2) \int \sec^3(c + dx) dx + 4(2a^2 B + 4aAb + 3b^2 B) \int \sec^2(c + dx) dx \right) + \frac{4a(aB + 2Ab) \tan(c + dx) \sec^3(c + dx)}{3d} \right) \\ \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{4} \left(\frac{1}{3} \left(4(2a^2 B + 4aAb + 3b^2 B) \int \csc(c + dx + \frac{\pi}{2})^2 dx + 3(3a^2 A + 8abB + 4Ab^2) \int \csc(c + dx + \frac{\pi}{2})^3 dx \right) + \frac{4a(aB + 2Ab) \tan(c + dx) \sec^3(c + dx)}{3d} \right) \\ \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 4254 \\ \frac{1}{4} \left(\frac{1}{3} \left(3(3a^2 A + 8abB + 4Ab^2) \int \csc(c + dx + \frac{\pi}{2})^3 dx - \frac{4(2a^2 B + 4aAb + 3b^2 B) \int 1d(-\tan(c + dx))}{d} \right) + \frac{4a(aB + 2Ab) \tan(c + dx) \sec^3(c + dx)}{3d} \right) \\ \frac{a^2 A \tan(c + dx) \sec^3(c + dx)}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 24 \end{array}$$

$$\frac{1}{4} \left(\frac{1}{3} \left(3(3a^2A + 8abB + 4Ab^2) \int \csc \left(c + dx + \frac{\pi}{2} \right)^3 dx + \frac{4(2a^2B + 4aAb + 3b^2B) \tan(c + dx)}{d} \right) + \frac{4a(aB + 2A)}{4d} \right)$$

↓ 4255

$$\frac{1}{4} \left(\frac{1}{3} \left(3(3a^2A + 8abB + 4Ab^2) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4(2a^2B + 4aAb + 3b^2B) \tan(c + dx)}{d} \right) + \frac{4a(aB + 2A)}{4d} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(3(3a^2A + 8abB + 4Ab^2) \left(\frac{1}{2} \int \csc \left(c + dx + \frac{\pi}{2} \right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4(2a^2B + 4aAb + 3b^2B) \tan(c + dx)}{d} \right) + \frac{4a(aB + 2A)}{4d} \right)$$

↓ 4257

$$\frac{1}{4} \left(\frac{1}{3} \left(3(3a^2A + 8abB + 4Ab^2) \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4(2a^2B + 4aAb + 3b^2B) \tan(c + dx)}{d} \right) + \frac{4a(aB + 2A)}{4d} \right)$$

input `Int[(a + b*cos[c + d*x])^2*(A + B*cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a^2*A*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((4*a*(2*A*b + a*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((4*(4*a*A*b + 2*a^2*B + 3*b^2*B)*Tan[c + d*x])/d + 3*(3*a^2*A + 4*A*b^2 + 8*a*b*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/3)/4`

3.230.3.1 Defintions of rubi rules used

- rule 244 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 255 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`
- rule 4254 `Int[csc[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.230.4 Maple [A] (verified)

Time = 5.23 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.95

method	result
parts	$\frac{A a^2 \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)}{d} + \frac{(A b^2 + 2Bab) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)}{d}$
derivativedivides	$\frac{A a^2 \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right) - B a^2 \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d}$
default	$\frac{A a^2 \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right) - B a^2 \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d}$
parallelrisch	$-36 \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) (A a^2 + \frac{4}{3} A b^2 + \frac{8}{3} Bab) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 36 \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) (A a^2 + \frac{4}{3} A b^2 + \frac{8}{3} Bab)$
norman	$\frac{(7A a^2 - 4A b^2 - 8Bab) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} + \frac{(27A a^2 - 32Aab + 12A b^2 - 16B a^2 + 24Bab) \left(\tan^{11} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{6d} + \frac{(27A a^2 + 32Aab + 12A b^2 - 16B a^2 + 24Bab) \left(\tan^{15} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{6d}$
risch	$- \frac{i(9A a^2 e^{7i(dx+c)} + 12A b^2 e^{7i(dx+c)} + 24Bab e^{7i(dx+c)} - 24B b^2 e^{6i(dx+c)} + 33A a^2 e^{5i(dx+c)} + 12A b^2 e^{5i(dx+c)} + 24Bab e^{5i(dx+c)} - 24B b^2 e^{4i(dx+c)} + 33A a^2 e^{3i(dx+c)} + 12A b^2 e^{3i(dx+c)} + 24Bab e^{3i(dx+c)} - 24B b^2 e^{2i(dx+c)} + 33A a^2 e^{i(dx+c)} + 12A b^2 e^{i(dx+c)} + 24Bab e^{i(dx+c)} - 24B b^2 e^{i(dx+c)})}{6d}$

input `int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBOSE)`

output `A*a^2/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^2+2*B*a*b)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-(2*A*a*b+B*a^2)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+B*b^2/d*tan(d*x+c)`

3.230.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.15

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3Aa^2 + 8Bab + 4Ab^2) \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(3Aa^2 + 8Bab + 4Ab^2) \cos(dx + c)^4 \log(\sin(dx + c) - 1)}{d \cos(dx + c)^4}$$

```
input integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")
```

```
output 1/48*(3*(3*A*a^2 + 8*B*a*b + 4*A*b^2)*cos(d*x + c)^4*log(sin(d*x + c) + 1)
- 3*(3*A*a^2 + 8*B*a*b + 4*A*b^2)*cos(d*x + c)^4*log(-sin(d*x + c) + 1) +
2*(8*(2*B*a^2 + 4*A*a*b + 3*B*b^2)*cos(d*x + c)^3 + 6*A*a^2 + 3*(3*A*a^2
+ 8*B*a*b + 4*A*b^2)*cos(d*x + c)^2 + 8*(B*a^2 + 2*A*a*b)*cos(d*x + c))*si
n(d*x + c))/(d*cos(d*x + c)^4)
```

3.230.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)
```

```
output Timed out
```

3.230.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.46

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{16(\tan(dx + c)^3 + 3 \tan(dx + c))Ba^2 + 32(\tan(dx + c)^3 + 3 \tan(dx + c))Aab - 3Aa^2 \left(\frac{2(3 \sin(dx + c)^3 - \sin(dx + c))}{\sin(dx + c)^4 - 2 \sin(dx + c)} \right)}{d}$$

3.230. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output $\frac{1}{48}(16*(\tan(dx + c))^3 + 3*\tan(dx + c))*B*a^2 + 32*(\tan(dx + c))^3 + 3*\tan(dx + c))*A*a*b - 3*A*a^2*(2*(3*\sin(dx + c)^3 - 5*\sin(dx + c)))/(\sin(dx + c)^4 - 2*\sin(dx + c)^2 + 1) - 3*\log(\sin(dx + c) + 1) + 3*\log(\sin(dx + c) - 1) - 24*B*a*b*(2*\sin(dx + c))/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) - 12*A*b^2*(2*\sin(dx + c))/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) + 48*B*b^2*\tan(dx + c))/d$

3.230.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 478 vs. $2(146) = 292$.

Time = 0.32 (sec) , antiderivative size = 478, normalized size of antiderivative = 3.06

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3Aa^2 + 8Bab + 4Ab^2) \log(|\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1|) - 3(3Aa^2 + 8Bab + 4Ab^2) \log(|\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1|)}{(\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1)^4}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")`

output $\frac{1}{24}(3*(3*A*a^2 + 8*B*a*b + 4*A*b^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(3*A*a^2 + 8*B*a*b + 4*A*b^2)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) + 2*(15*A*a^2*\tan(1/2*d*x + 1/2*c)^7 - 24*B*a^2*\tan(1/2*d*x + 1/2*c)^7 - 48*A*a*b*\tan(1/2*d*x + 1/2*c)^7 + 24*B*a*b*\tan(1/2*d*x + 1/2*c)^7 + 12*A*b^2*\tan(1/2*d*x + 1/2*c)^7 - 24*B*b^2*\tan(1/2*d*x + 1/2*c)^7 + 9*A*a^2*\tan(1/2*d*x + 1/2*c)^5 + 40*B*a^2*\tan(1/2*d*x + 1/2*c)^5 + 80*A*a*b*\tan(1/2*d*x + 1/2*c)^5 - 24*B*a*b*\tan(1/2*d*x + 1/2*c)^5 - 12*A*b^2*\tan(1/2*d*x + 1/2*c)^5 + 72*B*b^2*\tan(1/2*d*x + 1/2*c)^5 + 9*A*a^2*\tan(1/2*d*x + 1/2*c)^3 - 40*B*a^2*\tan(1/2*d*x + 1/2*c)^3 - 80*A*a*b*\tan(1/2*d*x + 1/2*c)^3 - 24*B*a*b*\tan(1/2*d*x + 1/2*c)^3 - 12*A*b^2*\tan(1/2*d*x + 1/2*c)^3 - 72*B*b^2*\tan(1/2*d*x + 1/2*c)^3 + 15*A*a^2*\tan(1/2*d*x + 1/2*c) + 24*B*a^2*\tan(1/2*d*x + 1/2*c) + 48*A*a*b*\tan(1/2*d*x + 1/2*c) + 24*B*a*b*\tan(1/2*d*x + 1/2*c) + 12*A*b^2*\tan(1/2*d*x + 1/2*c) + 24*B*b^2*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^4)/d$

$$3.230. \quad \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

3.230.9 Mupad [B] (verification not implemented)

Time = 4.01 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.01

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3Aa^2}{8} + B a b + \frac{Ab^2}{2}\right)}{\frac{3Aa^2}{2} + 4B a b + 2Ab^2}\right) \left(\frac{3Aa^2}{4} + 2B a b + Ab^2\right)}{d} + \frac{\left(\frac{5Aa^2}{4} + Ab^2 - 2Ba^2 - 2Bb^2 - 4A a b + 2B a b\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(\frac{3Aa^2}{4} - Ab^2 + \frac{10Ba^2}{3} + 6Bb^2 + \dots\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^5,x)`output `(atanh(((4*tan(c/2 + (d*x)/2)*((3*A*a^2)/8 + (A*b^2)/2 + B*a*b))/((3*A*a^2)/2 + 2*A*b^2 + 4*B*a*b))*((3*A*a^2)/4 + A*b^2 + 2*B*a*b))/d + (tan(c/2 + (d*x)/2)^7*((5*A*a^2)/4 + A*b^2 - 2*B*a^2 - 2*B*b^2 - 4*A*a*b + 2*B*a*b) - tan(c/2 + (d*x)/2)^3*(A*b^2 - (3*A*a^2)/4 + (10*B*a^2)/3 + 6*B*b^2 + (20*A*a*b)/3 + 2*B*a*b) + tan(c/2 + (d*x)/2)^5*((3*A*a^2)/4 - A*b^2 + (10*B*a^2)/3 + 6*B*b^2 + (20*A*a*b)/3 - 2*B*a*b) + tan(c/2 + (d*x)/2)*((5*A*a^2)/4 + A*b^2 + 2*B*a^2 + 2*B*b^2 + 4*A*a*b + 2*B*a*b))/(d*(6*tan(c/2 + (d*x)/2)^4 - 4*tan(c/2 + (d*x)/2)^2 - 4*tan(c/2 + (d*x)/2)^6 + tan(c/2 + (d*x)/2)^8 + 1))`

3.231 $\int \cos^2(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx$

3.231.1 Optimal result	2126
3.231.2 Mathematica [A] (verified)	2127
3.231.3 Rubi [A] (verified)	2127
3.231.4 Maple [A] (verified)	2131
3.231.5 Fricas [A] (verification not implemented)	2132
3.231.6 Sympy [B] (verification not implemented)	2133
3.231.7 Maxima [A] (verification not implemented)	2134
3.231.8 Giac [A] (verification not implemented)	2134
3.231.9 Mupad [B] (verification not implemented)	2135

3.231.1 Optimal result

Integrand size = 31, antiderivative size = 269

$$\int \cos^2(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx$$

$$= \frac{1}{16}(8a^3A+18aAb^2+18a^2bB+5b^3B)x + \frac{(15a^2Ab+4Ab^3+5a^3B+12ab^2B)\sin(c+dx)}{5d}$$

$$+ \frac{(8a^3A+18aAb^2+18a^2bB+5b^3B)\cos(c+dx)\sin(c+dx)}{16d}$$

$$+ \frac{b(18aAb+14a^2B+5b^2B)\cos^3(c+dx)\sin(c+dx)}{24d}$$

$$+ \frac{b^2(3Ab+4aB)\cos^4(c+dx)\sin(c+dx)}{15d}$$

$$+ \frac{bB\cos^3(c+dx)(a+b\cos(c+dx))^2\sin(c+dx)}{6d}$$

$$- \frac{(15a^2Ab+4Ab^3+5a^3B+12ab^2B)\sin^3(c+dx)}{15d}$$

```
output 1/16*(8*A*a^3+18*A*a*b^2+18*B*a^2*b+5*B*b^3)*x+1/5*(15*A*a^2*b+4*A*b^3+5*B
*a^3+12*B*a*b^2)*sin(d*x+c)/d+1/16*(8*A*a^3+18*A*a*b^2+18*B*a^2*b+5*B*b^3)
*cos(d*x+c)*sin(d*x+c)/d+1/24*b*(18*A*a*b+14*B*a^2+5*B*b^2)*cos(d*x+c)^3*s
in(d*x+c)/d+1/15*b^2*(3*A*b+4*B*a)*cos(d*x+c)^4*sin(d*x+c)/d+1/6*b*B*cos(d
*x+c)^3*(a+b*cos(d*x+c))^2*sin(d*x+c)/d-1/15*(15*A*a^2*b+4*A*b^3+5*B*a^3+1
2*B*a*b^2)*sin(d*x+c)^3/d
```

3.231.2 Mathematica [A] (verified)

Time = 1.67 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{480a^3Ac + 1080aAb^2c + 1080a^2bBc + 300b^3Bc + 480a^3Adx + 1080aAb^2dx + 1080a^2bBdx + 300b^3Bdx + \dots}{6d}$$

input `Integrate[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(480*a^3*A*c + 1080*a*A*b^2*c + 1080*a^2*b*B*c + 300*b^3*B*c + 480*a^3*A*d*x + 1080*a*A*b^2*d*x + 1080*a^2*b*B*d*x + 300*b^3*B*d*x + 120*(18*a^2*A*b + 5*A*b^3 + 6*a^3*B + 15*a*b^2*B)*Sin[c + d*x] + 15*(16*a^3*A + 48*a*A*b^2 + 48*a^2*b*B + 15*b^3*B)*Sin[2*(c + d*x)] + 240*a^2*A*b*Ssin[3*(c + d*x)] + 100*A*b^3*Ssin[3*(c + d*x)] + 80*a^3*B*Ssin[3*(c + d*x)] + 300*a*b^2*B*Ssin[3*(c + d*x)] + 90*a*A*b^2*Ssin[4*(c + d*x)] + 90*a^2*b*B*Ssin[4*(c + d*x)] + 45*b^3*B*Ssin[4*(c + d*x)] + 12*A*b^3*Ssin[5*(c + d*x)] + 36*a*b^2*B*Ssin[5*(c + d*x)] + 5*b^3*B*Ssin[6*(c + d*x)]/(960*d)`

3.231.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.86, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3469, 3042, 3512, 3042, 3502, 27, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^2 \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3469}$$

$$\frac{1}{6} \int \cos^2(c + dx)(a + b \cos(c + dx)) (2b(3Ab + 4aB) \cos^2(c + dx) + (5Bb^2 + 6a(2Ab + aB)) \cos(c + dx) + 3a(2aA + bB)) dx + \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{6} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right) \left(2b(3Ab + 4aB) \sin \left(c + dx + \frac{\pi}{2} \right)^2 + (5Bb^2 + 6a(2Ab + aB)) \sin \left(c + dx + \frac{\pi}{2} \right) + bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2 \right) dx$$

$$\downarrow \text{3512}$$

$$\frac{1}{6} \left(\frac{1}{5} \int \cos^2(c + dx) (15(2aA + bB)a^2 + 5b(14Ba^2 + 18Aba + 5b^2B) \cos^2(c + dx) + 6(5Ba^3 + 15Aba^2 + 12b^2Ba) \cos(c + dx) + bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2 \right) dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{6} \left(\frac{1}{5} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(15(2aA + bB)a^2 + 5b(14Ba^2 + 18Aba + 5b^2B) \sin \left(c + dx + \frac{\pi}{2} \right)^2 + 6(5Ba^3 + 15Aba^2 + 12b^2Ba) \sin \left(c + dx + \frac{\pi}{2} \right) + bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2 \right) dx \right)$$

$$\downarrow \text{3502}$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \int 3 \cos^2(c + dx) (5(8Aa^3 + 18bBa^2 + 18Ab^2a + 5b^3B) + 8(5Ba^3 + 15Aba^2 + 12b^2Ba + 4Ab^3) \cos(c + dx) + bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2 \right) dx \right)$$

$$\downarrow \text{27}$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int \cos^2(c + dx) (5(8Aa^3 + 18bBa^2 + 18Ab^2a + 5b^3B) + 8(5Ba^3 + 15Aba^2 + 12b^2Ba + 4Ab^3) \cos(c + dx) + bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2 \right) dx \right)$$

$$\downarrow \text{3042}$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(5(8Aa^3 + 18bBa^2 + 18Ab^2a + 5b^3B) + 8(5Ba^3 + 15Aba^2 + 12b^2Ba + 4Ab^3) \sin \left(c + dx + \frac{\pi}{2} \right) + bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2 \right) dx \right) \right)$$

$$\downarrow \text{3227}$$

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(8(5a^3B + 15a^2Ab + 12ab^2B + 4Ab^3) \int \cos^3(c + dx) dx + 5(8a^3A + 18a^2bB + 18aAb^2 + 5b^3B) \int \cos^2(c + dx) dx \right) \right) \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(5(8a^3A + 18a^2bB + 18aAb^2 + 5b^3B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 dx + 8(5a^3B + 15a^2Ab + 12ab^2B + 4Ab^3) \int \cos^2(c + dx) dx \right) \right) \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

↓ 3113

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(5(8a^3A + 18a^2bB + 18aAb^2 + 5b^3B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 dx - \frac{8(5a^3B + 15a^2Ab + 12ab^2B + 4Ab^3)}{d} \int \cos^2(c + dx) dx \right) \right) \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

↓ 2009

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(5(8a^3A + 18a^2bB + 18aAb^2 + 5b^3B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^2 dx - \frac{8(5a^3B + 15a^2Ab + 12ab^2B + 4Ab^3)}{d} \int \cos^2(c + dx) dx \right) \right) \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

↓ 3115

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(5(8a^3A + 18a^2bB + 18aAb^2 + 5b^3B) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{8(5a^3B + 15a^2Ab + 12ab^2B + 4Ab^3)}{d} \int \cos^2(c + dx) dx \right) \right) \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

↓ 24

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{5b(14a^2B + 18aAb + 5b^2B) \sin(c + dx) \cos^3(c + dx)}{4d} + \frac{3}{4} \left(5(8a^3A + 18a^2bB + 18aAb^2 + 5b^3B) \left(\frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{8(5a^3B + 15a^2Ab + 12ab^2B + 4Ab^3)}{d} \int \cos^2(c + dx) dx \right) \right) \right) \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^2}{6d}$$

input `Int[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`


```
output (b*B*Cos[c + d*x]^3*(a + b*Cos[c + d*x])^2*Sin[c + d*x]/(6*d) + ((2*b^2*(
3*A*b + 4*a*B)*Cos[c + d*x]^4*Sin[c + d*x])/(5*d) + ((5*b*(18*a*A*b + 14*a
^2*B + 5*b^2*B)*Cos[c + d*x]^3*Sin[c + d*x])/(4*d) + (3*(5*(8*a^3*A + 18*a
*A*b^2 + 18*a^2*b*B + 5*b^3*B)*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d)) -
(8*(15*a^2*A*b + 4*A*b^3 + 5*a^3*B + 12*a*b^2*B)*(-Sin[c + d*x] + Sin[c +
d*x]^3/3))/d))/4)/5)/6
```

3.231.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3113 Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
and[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

```
rule 3115 Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*Sin
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

```
rule 3227 Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x
_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*SIN[e +
f*x])^(m - 2)*(c + d*SIN[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*SIN[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*SIN[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
  (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f
_.)*(x_)^2], x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2
, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

3.231.4 Maple [A] (verified)

Time = 5.09 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.79

method	result
parts	$\frac{(A b^3 + 3 B a b^2) \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5d} + \frac{(3 A a b^2 + 3 B a^2 b) \left(\frac{\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}}{4} \right) \sin(dx+c)}{d}$
parallelrisch	$(240 A a^3 + 720 A a b^2 + 720 B a^2 b + 225 B b^3) \sin(2dx+2c) + (240 A a^2 b + 100 A b^3 + 80 B a^3 + 300 B a b^2) \sin(3dx+3c) + (90 A a^3 + 270 A a b^2 + 270 B a^2 b + 675 B b^3) \sin(dx+c)$
derivativdivides	$B b^3 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{A b^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5}$
default	$B b^3 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{A b^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5}$
risch	$\frac{a^3 A x}{2} + \frac{9 x A a b^2}{8} + \frac{9 x B a^2 b}{8} + \frac{5 b^3 B x}{16} + \frac{9 \sin(dx+c) A a^2 b}{4d} + \frac{5 \sin(dx+c) A b^3}{8d} + \frac{3 a^3 B \sin(dx+c)}{4d} + \frac{15 \sin(dx+c) B b^3}{8d}$
norman	$\frac{\left(\frac{1}{2} A a^3 + \frac{9}{8} A a b^2 + \frac{9}{8} B a^2 b + \frac{5}{16} B b^3 \right) x + \left(3 A a^3 + \frac{27}{4} A a b^2 + \frac{27}{4} B a^2 b + \frac{15}{8} B b^3 \right) x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \left(3 A a^3 + \frac{27}{4} A a b^2 + \frac{27}{4} B a^2 b + \frac{15}{8} B b^3 \right) x}{1}$

input `int(cos(d*x+c)^2*(a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/5*(A*b^3+3*B*a*b^2)/d*(8/3+cos(d*x+c)^4+4/3*cos(d*x+c)^2)*sin(d*x+c)+(3*A*a*b^2+3*B*a^2*b)/d*(1/4*(cos(d*x+c)^3+3/2*cos(d*x+c))*sin(d*x+c)+3/8*d*x+3/8*c)+1/3*(3*A*a^2*b+B*a^3)/d*(2+cos(d*x+c)^2)*sin(d*x+c)+A*a^3/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+B*b^3/d*(1/6*(cos(d*x+c)^5+5/4*cos(d*x+c)^3+15/8*cos(d*x+c))*sin(d*x+c)+5/16*d*x+5/16*c)`

3.231.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.78

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{15(8 A a^3 + 18 B a^2 b + 18 A a b^2 + 5 B b^3)dx + (40 B b^3 \cos(dx + c)^5 + 48(3 B a b^2 + A b^3) \cos(dx + c)^4 + 10(3 A a^3 + 27 A a b^2 + 27 B a^2 b + 15 B b^3) \cos(dx + c)^3 + 15(8 A a^3 + 18 B a^2 b + 18 A a b^2 + 5 B b^3) \cos(dx + c)^2 + 15(3 A a^3 + 27 A a b^2 + 27 B a^2 b + 15 B b^3) \cos(dx + c) + 15(3 A a^3 + 27 A a b^2 + 27 B a^2 b + 15 B b^3)}{1}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="fracas")`

3.231. $\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$

```
output 1/240*(15*(8*A*a^3 + 18*B*a^2*b + 18*A*a*b^2 + 5*B*b^3)*d*x + (40*B*b^3*cos(d*x + c)^5 + 48*(3*B*a*b^2 + A*b^3)*cos(d*x + c)^4 + 160*B*a^3 + 480*A*a^2*b + 384*B*a*b^2 + 128*A*b^3 + 10*(18*B*a^2*b + 18*A*a*b^2 + 5*B*b^3)*cos(d*x + c)^3 + 16*(5*B*a^3 + 15*A*a^2*b + 12*B*a*b^2 + 4*A*b^3)*cos(d*x + c)^2 + 15*(8*A*a^3 + 18*B*a^2*b + 18*A*a*b^2 + 5*B*b^3)*cos(d*x + c))*sin(d*x + c))/d
```

3.231.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 721 vs. $2(277) = 554$.

Time = 0.43 (sec) , antiderivative size = 721, normalized size of antiderivative = 2.68

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \left\{ \begin{array}{l} \frac{Aa^3x \sin^2(c+dx)}{2} + \frac{Aa^3x \cos^2(c+dx)}{2} + \frac{Aa^3 \sin(c+dx) \cos(c+dx)}{2d} + \frac{2Aa^2b \sin^3(c+dx)}{d} + \frac{3Aa^2b \sin(c+dx) \cos^2(c+dx)}{d} + \frac{9Aab^2x \sin^2(c+dx)}{2d} \\ x(A + B \cos(c)) (a + b \cos(c))^3 \cos^2(c) \end{array} \right.$$

```
input integrate(cos(d*x+c)**2*(a+b*cos(d*x+c))**3*(A+B*cos(d*x+c)), x)
```

```
output Piecewise((A*a**3*x*sin(c + d*x)**2/2 + A*a**3*x*cos(c + d*x)**2/2 + A*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*A*a**2*b*sin(c + d*x)**3/d + 3*A*a**2*b*sin(c + d*x)*cos(c + d*x)**2/d + 9*A*a*b**2*x*sin(c + d*x)**4/8 + 9*A*a*b**2*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 9*A*a*b**2*x*cos(c + d*x)**4/8 + 9*A*a*b**2*x*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 15*A*a*b**2*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 8*A*b**3*sin(c + d*x)**5/(15*d) + 4*A*b**3*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + A*b**3*sin(c + d*x)*cos(c + d*x)**4/d + 2*B*a**3*sin(c + d*x)**3/(3*d) + B*a**3*sin(c + d*x)*cos(c + d*x)**2/d + 9*B*a**2*b*x*sin(c + d*x)**4/8 + 9*B*a**2*b*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 9*B*a**2*b*x*cos(c + d*x)**4/8 + 9*B*a**2*b*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 15*B*a**2*b*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 8*B*a*b**2*sin(c + d*x)**5/(5*d) + 4*B*a*b**2*sin(c + d*x)**3*cos(c + d*x)**2/d + 3*B*a*b**2*sin(c + d*x)*cos(c + d*x)**4/d + 5*B*b**3*x*sin(c + d*x)**6/16 + 15*B*b**3*x*sin(c + d*x)**4*cos(c + d*x)**2/16 + 15*B*b**3*x*sin(c + d*x)**2*cos(c + d*x)**4/16 + 5*B*b**3*x*cos(c + d*x)**6/16 + 5*B*b**3*sin(c + d*x)**5*cos(c + d*x)/(16*d) + 5*B*b**3*sin(c + d*x)**3*cos(c + d*x)**3/(6*d) + 11*B*b**3*sin(c + d*x)*cos(c + d*x)**5/(16*d), Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**3*cos(c)**2, True))
```

3.231.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.99

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{240(2dx + 2c + \sin(2dx + 2c))Aa^3 - 320(\sin(dx + c)^3 - 3\sin(dx + c))Ba^3 - 960(\sin(dx + c)^3 - 3\sin(dx + c))Aa^2b + 90(12dx + 12c + \sin(4dx + 4c) + 8\sin(2dx + 2c))B^2a^2b + 90(12dx + 12c + \sin(4dx + 4c) + 8\sin(2dx + 2c))Aa^2b^2 + 192(3\sin(dx + c)^5 - 10\sin(dx + c)^3 + 15\sin(dx + c))B^2a^2b^2 + 64(3\sin(dx + c)^5 - 10\sin(dx + c)^3 + 15\sin(dx + c))Aa^2b^3 - 5(4\sin(2dx + 2c)^3 - 60dx - 60c - 9\sin(4dx + 4c) - 48\sin(2dx + 2c))B^2b^3}{d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/960*(240*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^3 - 320*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^3 - 960*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a^2*b + 90*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B^2*a^2*b + 90*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*a^2*b^2 + 192*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B^2*a^2*b^2 + 64*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*A*a^2*b^3 - 5*(4*sin(2*d*x + 2*c)^3 - 60*d*x - 60*c - 9*sin(4*d*x + 4*c) - 48*sin(2*d*x + 2*c))*B^2*b^3)/d`

3.231.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.86

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{Bb^3 \sin(6dx + 6c)}{192d} + \frac{1}{16} (8Aa^3 + 18Ba^2b + 18Aab^2 + 5Bb^3)x$$

$$+ \frac{(3Bab^2 + Ab^3) \sin(5dx + 5c)}{80d} + \frac{3(2Ba^2b + 2Aab^2 + Bb^3) \sin(4dx + 4c)}{64d}$$

$$+ \frac{(4Ba^3 + 12Aa^2b + 15Bab^2 + 5Ab^3) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(16Aa^3 + 48Ba^2b + 48Aab^2 + 15Bb^3) \sin(2dx + 2c)}{64d}$$

$$+ \frac{(6Ba^3 + 18Aa^2b + 15Bab^2 + 5Ab^3) \sin(dx + c)}{8d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")`

output $1/192*B*b^3*\sin(6*d*x + 6*c)/d + 1/16*(8*A*a^3 + 18*B*a^2*b + 18*A*a*b^2 + 5*B*b^3)*x + 1/80*(3*B*a*b^2 + A*b^3)*\sin(5*d*x + 5*c)/d + 3/64*(2*B*a^2*b + 2*A*a*b^2 + B*b^3)*\sin(4*d*x + 4*c)/d + 1/48*(4*B*a^3 + 12*A*a^2*b + 15*B*a*b^2 + 5*A*b^3)*\sin(3*d*x + 3*c)/d + 1/64*(16*A*a^3 + 48*B*a^2*b + 48*A*a*b^2 + 15*B*b^3)*\sin(2*d*x + 2*c)/d + 1/8*(6*B*a^3 + 18*A*a^2*b + 15*B*a*b^2 + 5*A*b^3)*\sin(d*x + c)/d$

3.231.9 Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.31

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{Aa^3x}{2} + \frac{5Bb^3x}{16} + \frac{9Aab^2x}{8} + \frac{9Ba^2bx}{8} + \frac{5Ab^3 \sin(c + dx)}{8d}$$

$$+ \frac{3Ba^3 \sin(c + dx)}{4d} + \frac{Aa^3 \sin(2c + 2dx)}{4d} + \frac{5Ab^3 \sin(3c + 3dx)}{48d}$$

$$+ \frac{Ba^3 \sin(3c + 3dx)}{12d} + \frac{Ab^3 \sin(5c + 5dx)}{80d} + \frac{15Bb^3 \sin(2c + 2dx)}{64d}$$

$$+ \frac{3Bb^3 \sin(4c + 4dx)}{64d} + \frac{Bb^3 \sin(6c + 6dx)}{192d} + \frac{3Aab^2 \sin(2c + 2dx)}{4d}$$

$$+ \frac{Aa^2b \sin(3c + 3dx)}{4d} + \frac{3Aab^2 \sin(4c + 4dx)}{32d} + \frac{3Ba^2b \sin(2c + 2dx)}{4d}$$

$$+ \frac{5Bab^2 \sin(3c + 3dx)}{16d} + \frac{3Ba^2b \sin(4c + 4dx)}{32d}$$

$$+ \frac{3Bab^2 \sin(5c + 5dx)}{80d} + \frac{9Aa^2b \sin(c + dx)}{4d} + \frac{15Bab^2 \sin(c + dx)}{8d}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3,x)`

output $(A*a^3*x)/2 + (5*B*b^3*x)/16 + (9*A*a*b^2*x)/8 + (9*B*a^2*b*x)/8 + (5*A*b^3*\sin(c + d*x))/(8*d) + (3*B*a^3*\sin(c + d*x))/(4*d) + (A*a^3*\sin(2*c + 2*d*x))/(4*d) + (5*A*b^3*\sin(3*c + 3*d*x))/(48*d) + (B*a^3*\sin(3*c + 3*d*x))/(12*d) + (A*b^3*\sin(5*c + 5*d*x))/(80*d) + (15*B*b^3*\sin(2*c + 2*d*x))/(64*d) + (3*B*b^3*\sin(4*c + 4*d*x))/(64*d) + (B*b^3*\sin(6*c + 6*d*x))/(192*d) + (3*A*a*b^2*\sin(2*c + 2*d*x))/(4*d) + (A*a^2*b*\sin(3*c + 3*d*x))/(4*d) + (3*A*a*b^2*\sin(4*c + 4*d*x))/(32*d) + (3*B*a^2*b*\sin(2*c + 2*d*x))/(4*d) + (5*B*a*b^2*\sin(3*c + 3*d*x))/(16*d) + (3*B*a^2*b*\sin(4*c + 4*d*x))/(32*d) + (3*B*a*b^2*\sin(5*c + 5*d*x))/(80*d) + (9*A*a^2*b*\sin(c + d*x))/(4*d) + (15*B*a*b^2*\sin(c + d*x))/(8*d)$

3.232 $\int \cos(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx$

3.232.1 Optimal result	2136
3.232.2 Mathematica [A] (verified)	2137
3.232.3 Rubi [A] (verified)	2137
3.232.4 Maple [A] (verified)	2140
3.232.5 Fricas [A] (verification not implemented)	2141
3.232.6 Sympy [B] (verification not implemented)	2141
3.232.7 Maxima [A] (verification not implemented)	2142
3.232.8 Giac [A] (verification not implemented)	2143
3.232.9 Mupad [B] (verification not implemented)	2143

3.232.1 Optimal result

Integrand size = 29, antiderivative size = 243

$$\begin{aligned} & \int \cos(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx \\ &= \frac{1}{8}(12a^2Ab + 3Ab^3 + 4a^3B + 9ab^2B) x \\ &+ \frac{(15a^3Ab + 60aAb^3 - 3a^4B + 52a^2b^2B + 16b^4B) \sin(c+dx)}{30bd} \\ &+ \frac{(30a^2Ab + 45Ab^3 - 6a^3B + 71ab^2B) \cos(c+dx) \sin(c+dx)}{120d} \\ &+ \frac{(15aAb - 3a^2B + 16b^2B) (a+b \cos(c+dx))^2 \sin(c+dx)}{60bd} \\ &+ \frac{(5Ab - aB)(a+b \cos(c+dx))^3 \sin(c+dx)}{20bd} + \frac{B(a+b \cos(c+dx))^4 \sin(c+dx)}{5bd} \end{aligned}$$

```
output 1/8*(12*A*a^2*b+3*A*b^3+4*B*a^3+9*B*a*b^2)*x+1/30*(15*A*a^3*b+60*A*a*b^3-3
*B*a^4+52*B*a^2*b^2+16*B*b^4)*sin(d*x+c)/b/d+1/120*(30*A*a^2*b+45*A*b^3-6*
B*a^3+71*B*a*b^2)*cos(d*x+c)*sin(d*x+c)/d+1/60*(15*A*a*b-3*B*a^2+16*B*b^2)
*(a+b*cos(d*x+c))^2*sin(d*x+c)/b/d+1/20*(5*A*b-B*a)*(a+b*cos(d*x+c))^3*sin
(d*x+c)/b/d+1/5*B*(a+b*cos(d*x+c))^4*sin(d*x+c)/b/d
```

3.232.2 Mathematica [A] (verified)

Time = 2.31 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.72

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{60(12a^2Ab + 3Ab^3 + 4a^3B + 9ab^2B)(c + dx) + 60(8a^3A + 18aAb^2 + 18a^2bB + 5b^3B) \sin(c + dx) + 120}{480d}$$

input `Integrate[Cos[c + d*x]*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(60*(12*a^2*A*b + 3*A*b^3 + 4*a^3*B + 9*a*b^2*B)*(c + d*x) + 60*(8*a^3*A + 18*a*A*b^2 + 18*a^2*b*B + 5*b^3*B)*Sin[c + d*x] + 120*(3*a^2*A*b + A*b^3 + a^3*B + 3*a*b^2*B)*Sin[2*(c + d*x)] + 10*b*(12*a*A*b + 12*a^2*B + 5*b^2*B)*Sin[3*(c + d*x)] + 15*b^2*(A*b + 3*a*B)*Sin[4*(c + d*x)] + 6*b^3*B*SIN[5*(c + d*x)])/(480*d)`

3.232.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 3447, 3042, 3502, 3042, 3232, 3042, 3232, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int (a + b \cos(c + dx))^3 (A \cos(c + dx) + B \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{3502}$$

$$\frac{\int (a + b \cos(c + dx))^3 (4bB + (5Ab - aB) \cos(c + dx)) dx}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

↓ 3042

$$\frac{\int (a + b \sin(c + dx + \frac{\pi}{2}))^3 (4bB + (5Ab - aB) \sin(c + dx + \frac{\pi}{2})) dx}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

↓ 3232

$$\frac{\frac{1}{4} \int (a + b \cos(c + dx))^2 (b(15Ab + 13aB) + (-3Ba^2 + 15Aba + 16b^2B) \cos(c + dx)) dx + \frac{(5Ab - aB) \sin(c + dx)(a + b \cos(c + dx))^4}{4d}}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

↓ 3042

$$\frac{\frac{1}{4} \int (a + b \sin(c + dx + \frac{\pi}{2}))^2 (b(15Ab + 13aB) + (-3Ba^2 + 15Aba + 16b^2B) \sin(c + dx + \frac{\pi}{2})) dx + \frac{(5Ab - aB) \sin(c + dx)(a + b \cos(c + dx))^4}{4d}}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

↓ 3232

$$\frac{\frac{1}{4} \left(\frac{1}{3} \int (a + b \cos(c + dx)) (b(33Ba^2 + 75Aba + 32b^2B) + (-6Ba^3 + 30Aba^2 + 71b^2Ba + 45Ab^3) \cos(c + dx)) dx + \frac{(5Ab - aB) \sin(c + dx)(a + b \cos(c + dx))^4}{4d} \right)}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

↓ 3042

$$\frac{\frac{1}{4} \left(\frac{1}{3} \int (a + b \sin(c + dx + \frac{\pi}{2})) (b(33Ba^2 + 75Aba + 32b^2B) + (-6Ba^3 + 30Aba^2 + 71b^2Ba + 45Ab^3) \sin(c + dx + \frac{\pi}{2})) dx + \frac{(5Ab - aB) \sin(c + dx)(a + b \cos(c + dx))^4}{4d} \right)}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

↓ 3213

$$\frac{\frac{1}{4} \left(\frac{(-3a^2B + 15aAb + 16b^2B) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{1}{3} \left(\frac{b(-6a^3B + 30a^2Ab + 71ab^2B + 45Ab^3) \sin(c + dx) \cos(c + dx)}{2d} + \frac{15}{2} bx(4a^3B + 3a^2Ab + 7ab^2B + 4b^3) \sin(c + dx) \right) \right)}{5b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5bd}$$

3.232. $\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$

input `Int[Cos[c + d*x]*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(B*(a + b*Cos[c + d*x])^4*Sin[c + d*x])/(5*b*d) + (((5*A*b - a*B)*(a + b*Cos[c + d*x])^3*Sin[c + d*x])/(4*d) + (((15*a*A*b - 3*a^2*B + 16*b^2*B)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + ((15*b*(12*a^2*A*b + 3*A*b^3 + 4*a^3*B + 9*a*b^2*B)*x)/2 + (2*(15*a^3*A*b + 60*a*A*b^3 - 3*a^4*B + 52*a^2*b^2*B + 16*b^4*B)*Sin[c + d*x])/d + (b*(30*a^2*A*b + 45*A*b^3 - 6*a^3*B + 71*a*b^2*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/3)/4)/(5*b)`

3.232.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.232.4 Maple [A] (verified)

Time = 3.95 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.72

method	result
parts	$\frac{(A b^3 + 3 B a b^2) \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right)}{d} + \frac{(3 A a b^2 + 3 B a^2 b) (2 + \cos^2(dx+c)) \sin(dx+c)}{3d} +$
parallelrisch	$\frac{120(3 A a^2 b + A b^3 + B a^3 + 3 B a b^2) \sin(2dx+2c) + 10(12 A a b^2 + 12 B a^2 b + 5 B b^3) \sin(3dx+3c) + 15(A b^3 + 3 B a b^2) \sin(4dx+4c)}{480c}$
derivativedivides	$\frac{B b^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + A b^3 \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 3 B a b^2 \left(\frac{\cos^3(dx+c)}{3} + \frac{3dx}{8} + \frac{3c}{8} \right)$
default	$\frac{B b^3 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + A b^3 \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 3 B a b^2 \left(\frac{\cos^3(dx+c)}{3} + \frac{3dx}{8} + \frac{3c}{8} \right)$
risch	$\frac{3x A a^2 b}{2} + \frac{3x A b^3}{8} + \frac{a^3 B x}{2} + \frac{9x B a b^2}{8} + \frac{a^3 A \sin(dx+c)}{d} + \frac{9 \sin(dx+c) A a b^2}{4d} + \frac{9 \sin(dx+c) B a^2 b}{4d} + \frac{5 \sin(dx+c) B b^3}{8}$
norman	$\frac{(\frac{3}{2} A a^2 b + \frac{3}{8} A b^3 + \frac{1}{2} B a^3 + \frac{9}{8} B a b^2) x + (15 A a^2 b + \frac{15}{4} A b^3 + 5 B a^3 + \frac{45}{4} B a b^2) x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (15 A a^2 b + \frac{15}{4} A b^3 + 5 B a^3 + \frac{45}{4} B a b^2) x}{480c}$

```
input int(cos(d*x+c)*(a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (A*b^3+3*B*a*b^2)/d*(1/4*(cos(d*x+c)^3+3/2*cos(d*x+c))*sin(d*x+c)+3/8*d*x+
3/8*c)+1/3*(3*A*a*b^2+3*B*a^2*b)/d*(2+cos(d*x+c)^2)*sin(d*x+c)+(3*A*a^2*b+
B*a^3)/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+a^3*A*sin(d*x+c)/d+1/5*
B*b^3/d*(8/3+cos(d*x+c)^4+4/3*cos(d*x+c)^2)*sin(d*x+c)
```

3.232. $\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.232.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.72

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{15(4Ba^3 + 12Aa^2b + 9Bab^2 + 3Ab^3)dx + (24Bb^3 \cos(dx + c)^4 + 120Aa^3 + 240Ba^2b + 240Aab^2 + 64Bb^3 \cos(dx + c)^3 + 120Aa^2b + 240Aab^2 + 64Bb^3 \cos(dx + c)^2 + 120Aa^2b + 240Aab^2 + 64Bb^3 \cos(dx + c) + 120Aa^2b + 240Aab^2 + 64Bb^3) \sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/120*(15*(4*B*a^3 + 12*A*a^2*b + 9*B*a*b^2 + 3*A*b^3)*d*x + (24*B*b^3*cos(d*x + c)^4 + 120*A*a^3 + 240*B*a^2*b + 240*A*a*b^2 + 64*B*b^3 + 30*(3*B*a*b^2 + A*b^3)*cos(d*x + c)^3 + 8*(15*B*a^2*b + 15*A*a*b^2 + 4*B*b^3)*cos(d*x + c)^2 + 15*(4*B*a^3 + 12*A*a^2*b + 9*B*a*b^2 + 3*A*b^3)*cos(d*x + c))*sin(d*x + c))/d`

3.232.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 551 vs. 2(241) = 482.

Time = 0.30 (sec) , antiderivative size = 551, normalized size of antiderivative = 2.27

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aa^3 \sin(c+dx)}{d} + \frac{3Aa^2bx \sin^2(c+dx)}{2} + \frac{3Aa^2bx \cos^2(c+dx)}{2} + \frac{3Aa^2b \sin(c+dx) \cos(c+dx)}{2d} + \frac{2Aab^2 \sin^3(c+dx)}{d} + \frac{3Aab^2 \sin(c+dx)}{d} \\ x(A + B \cos(c))(a + b \cos(c))^3 \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)`

output `Piecewise((A**3*sin(c + d*x)/d + 3*A**2*b*x*sin(c + d*x)**2/2 + 3*A**2*b*x*cos(c + d*x)**2/2 + 3*A**2*b*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*A*a*b**2*sin(c + d*x)**3/d + 3*A*a*b**2*sin(c + d*x)*cos(c + d*x)**2/d + 3*A*b**3*x*sin(c + d*x)**4/8 + 3*A*b**3*x*cos(c + d*x)**2*cos(c + d*x)**2/4 + 3*A*b**3*x*cos(c + d*x)**4/8 + 3*A*b**3*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 5*A*b**3*sin(c + d*x)*cos(c + d*x)**3/(8*d) + B*a**3*x*sin(c + d*x)**2/2 + B*a**3*x*cos(c + d*x)**2/2 + B*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*B*a**2*b*sin(c + d*x)**3/d + 3*B*a**2*b*sin(c + d*x)*cos(c + d*x)**2/d + 9*B*a*b**2*x*sin(c + d*x)**4/8 + 9*B*a*b**2*x*sin(c + d*x)**2*cos(c + d*x)**2/4 + 9*B*a*b**2*x*cos(c + d*x)**4/8 + 9*B*a*b**2*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 15*B*a*b**2*sin(c + d*x)*cos(c + d*x)**3/(8*d) + 8*B*b**3*sin(c + d*x)**5/(15*d) + 4*B*b**3*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + B*b**3*sin(c + d*x)*cos(c + d*x)**4/d, Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**3*cos(c), True))`

3.232.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.89

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{120(2dx + 2c + \sin(2dx + 2c))Ba^3 + 360(2dx + 2c + \sin(2dx + 2c))Aa^2b - 480(\sin(dx + c)^3 - 3\sin(dx + c)\cos(dx + c))Aa^2b^2 + 480(\sin(dx + c)^3 - 3\sin(dx + c)\cos(dx + c))Aab^2 + 480(\sin(dx + c)^3 - 3\sin(dx + c)\cos(dx + c))Ab^2c + 480(\sin(dx + c)^3 - 3\sin(dx + c)\cos(dx + c))Bb^3 + 480Aa^3\sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `1/480*(120*(2*d*x + 2*c + sin(2*d*x + 2*c))*B*a^3 + 360*(2*d*x + 2*c + sin(2*d*x + 2*c))*A*a^2*b - 480*(sin(d*x + c)^3 - 3*sin(d*x + c))*B*a^2*b^2 - 480*(sin(d*x + c)^3 - 3*sin(d*x + c))*A*a*b^2 + 45*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*B*a*b^2 + 15*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(2*d*x + 2*c))*A*b^3 + 32*(3*sin(d*x + c)^5 - 10*sin(d*x + c)^3 + 15*sin(d*x + c))*B*b^3 + 480*A*a^3*sin(d*x + c))/d`

3.232.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.77

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{Bb^3 \sin(5dx + 5c)}{80d} + \frac{1}{8} (4Ba^3 + 12Aa^2b + 9Bab^2 + 3Ab^3)x$$

$$+ \frac{(3Bab^2 + Ab^3) \sin(4dx + 4c)}{32d} + \frac{(12Ba^2b + 12Aab^2 + 5Bb^3) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(Ba^3 + 3Aa^2b + 3Bab^2 + Ab^3) \sin(2dx + 2c)}{4d}$$

$$+ \frac{(8Aa^3 + 18Ba^2b + 18Aab^2 + 5Bb^3) \sin(dx + c)}{8d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `1/80*B*b^3*sin(5*d*x + 5*c)/d + 1/8*(4*B*a^3 + 12*A*a^2*b + 9*B*a*b^2 + 3*A*b^3)*x + 1/32*(3*B*a*b^2 + A*b^3)*sin(4*d*x + 4*c)/d + 1/48*(12*B*a^2*b + 12*A*a*b^2 + 5*B*b^3)*sin(3*d*x + 3*c)/d + 1/4*(B*a^3 + 3*A*a^2*b + 3*B*a*b^2 + A*b^3)*sin(2*d*x + 2*c)/d + 1/8*(8*A*a^3 + 18*B*a^2*b + 18*A*a*b^2 + 5*B*b^3)*sin(d*x + c)/d`

3.232.9 Mupad [B] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.14

$$\int \cos(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{3Ab^3x}{8} + \frac{Ba^3x}{2} + \frac{3Aa^2bx}{2} + \frac{9Bab^2x}{8} + \frac{Aa^3 \sin(c + dx)}{d} + \frac{5Bb^3 \sin(c + dx)}{8d}$$

$$+ \frac{Ab^3 \sin(2c + 2dx)}{4d} + \frac{Ba^3 \sin(2c + 2dx)}{4d} + \frac{Ab^3 \sin(4c + 4dx)}{32d}$$

$$+ \frac{5Bb^3 \sin(3c + 3dx)}{48d} + \frac{Bb^3 \sin(5c + 5dx)}{80d} + \frac{3Aa^2b \sin(2c + 2dx)}{4d}$$

$$+ \frac{Aab^2 \sin(3c + 3dx)}{4d} + \frac{3Bab^2 \sin(2c + 2dx)}{4d} + \frac{Ba^2b \sin(3c + 3dx)}{4d}$$

$$+ \frac{3Bab^2 \sin(4c + 4dx)}{32d} + \frac{9Aab^2 \sin(c + dx)}{4d} + \frac{9Ba^2b \sin(c + dx)}{4d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3,x)`

output $(3A^3b^3x)/8 + (B^3a^3x)/2 + (3A^2a^2b^3x)/2 + (9B^2a^2b^3x)/8 + (A^3a^3 \sin(c + d*x))/d + (5B^2b^3 \sin(c + d*x))/(8d) + (A^2b^3 \sin(2c + 2d*x))/(4d) + (B^2a^3 \sin(2c + 2d*x))/(4d) + (A^2b^3 \sin(4c + 4d*x))/(32d) + (5B^2b^3 \sin(3c + 3d*x))/(48d) + (B^2b^3 \sin(5c + 5d*x))/(80d) + (3A^2a^2b^3 \sin(2c + 2d*x))/(4d) + (A^2a^2b^2 \sin(3c + 3d*x))/(4d) + (3B^2a^2b^2 \sin(2c + 2d*x))/(4d) + (B^2a^2b^2 \sin(3c + 3d*x))/(4d) + (3B^2a^2b^2 \sin(4c + 4d*x))/(32d) + (9A^2a^2b^2 \sin(c + d*x))/(4d) + (9B^2a^2b^2 \sin(c + d*x))/(4d)$

3.233 $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$

3.233.1 Optimal result	2145
3.233.2 Mathematica [A] (verified)	2145
3.233.3 Rubi [A] (verified)	2146
3.233.4 Maple [A] (verified)	2148
3.233.5 Fricas [A] (verification not implemented)	2148
3.233.6 Sympy [B] (verification not implemented)	2149
3.233.7 Maxima [A] (verification not implemented)	2149
3.233.8 Giac [A] (verification not implemented)	2150
3.233.9 Mupad [B] (verification not implemented)	2151

3.233.1 Optimal result

Integrand size = 23, antiderivative size = 171

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{1}{8} (8a^3 A + 12aAb^2 + 12a^2 bB + 3b^3 B) x + \frac{(16a^2 Ab + 4Ab^3 + 3a^3 B + 12ab^2 B) \sin(c + dx)}{6d}$$

$$+ \frac{b(20aAb + 6a^2 B + 9b^2 B) \cos(c + dx) \sin(c + dx)}{24d}$$

$$+ \frac{(4Ab + 3aB)(a + b \cos(c + dx))^2 \sin(c + dx)}{12d} + \frac{B(a + b \cos(c + dx))^3 \sin(c + dx)}{4d}$$

```
output 1/8*(8*A*a^3+12*A*a*b^2+12*B*a^2*b+3*B*b^3)*x+1/6*(16*A*a^2*b+4*A*b^3+3*B*
a^3+12*B*a*b^2)*sin(d*x+c)/d+1/24*b*(20*A*a*b+6*B*a^2+9*B*b^2)*cos(d*x+c)*
sin(d*x+c)/d+1/12*(4*A*b+3*B*a)*(a+b*cos(d*x+c))^2*sin(d*x+c)/d+1/4*B*(a+b
*cos(d*x+c))^3*sin(d*x+c)/d
```

3.233.2 Mathematica [A] (verified)

Time = 1.36 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.82

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{12(8a^3 A + 12aAb^2 + 12a^2 bB + 3b^3 B) (c + dx) + 24(12a^2 Ab + 3Ab^3 + 4a^3 B + 9ab^2 B) \sin(c + dx) + 24b^3 B \cos(c + dx)}{96d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output $(12*(8*a^3*A + 12*a*A*b^2 + 12*a^2*b*B + 3*b^3*B)*(c + d*x) + 24*(12*a^2*A*b + 3*A*b^3 + 4*a^3*B + 9*a*b^2*B)*\text{Sin}[c + d*x] + 24*b*(3*a*A*b + 3*a^2*B + b^2*B)*\text{Sin}[2*(c + d*x)] + 8*b^2*(A*b + 3*a*B)*\text{Sin}[3*(c + d*x)] + 3*b^3*B*\text{Sin}[4*(c + d*x)])/(96*d)$

3.233.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3042, 3232, 3042, 3232, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3232}$$

$$\frac{1}{4} \int (a + b \cos(c + dx))^2 (4aA + 3bB + (4Ab + 3aB) \cos(c + dx)) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^3}{4d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(4aA + 3bB + (4Ab + 3aB) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^3}{4d}$$

$$\downarrow \text{3232}$$

$$\frac{1}{4} \left(\frac{1}{3} \int (a + b \cos(c + dx)) (12Aa^2 + 15bBa + 8Ab^2 + (6Ba^2 + 20Aba + 9b^2B) \cos(c + dx)) dx + \frac{(3aB + 4Ab) \sin(c + dx)(a + b \cos(c + dx))^3}{4d}\right)$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{3} \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right) \left(12Aa^2 + 15bBa + 8Ab^2 + (6Ba^2 + 20Aba + 9b^2B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \right)$$

↓ 3213

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{b(6a^2B + 20aAb + 9b^2B) \sin(c + dx) \cos(c + dx)}{2d} + \frac{2(3a^3B + 16a^2Ab + 12ab^2B + 4Ab^3) \sin(c + dx)}{d} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(B*(a + b*Cos[c + d*x])^3*Sin[c + d*x])/(4*d) + (((4*A*b + 3*a*B)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + ((3*(8*a^3*A + 12*a*A*b^2 + 12*a^2*b*B + 3*b^3*B)*x)/2 + (2*(16*a^2*A*b + 4*A*b^3 + 3*a^3*B + 12*a*b^2*B)*Sin[c + d*x])/d + (b*(20*a*A*b + 6*a^2*B + 9*b^2*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/3)/4`

3.233.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

3.233.4 Maple [A] (verified)

Time = 3.16 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.83

method	result
parallelrisc	$\frac{24(3Aa^2b^2+3Ba^2b+Bb^3)\sin(2dx+2c)+8(Ab^3+3Bab^2)\sin(3dx+3c)+3B\sin(4dx+4c)b^3+24(12Aa^2b+3Ab^3+4Ba^3+9Bab^2)\sin(dx+c)}{96d}$
parts	$a^3Ax + \frac{(Ab^3+3Bab^2)(2+\cos^2(dx+c))\sin(dx+c)}{3d} + \frac{(3Aa^2b+3Ba^2b)\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{(3Aa^2b^2+3Bab^3)\sin(dx+c)}{3d}$
derivativdivides	$Bb^3\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ab^3(2+\cos^2(dx+c))\sin(dx+c)}{3} + Bab^2(2+\cos^2(dx+c))\sin(dx+c)$
default	$Bb^3\left(\frac{\left(\cos^3(dx+c)+\frac{3\cos(dx+c)}{2}\right)\sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8}\right) + \frac{Ab^3(2+\cos^2(dx+c))\sin(dx+c)}{3} + Bab^2(2+\cos^2(dx+c))\sin(dx+c)$
risc	$a^3Ax + \frac{3xAab^2}{2} + \frac{3xBa^2b}{2} + \frac{3b^3Bx}{8} + \frac{3\sin(dx+c)Aa^2b}{d} + \frac{3\sin(dx+c)Ab^3}{4d} + \frac{a^3B\sin(dx+c)}{d} + \frac{9\sin(dx+c)Bab^2}{4d}$
norman	$\frac{(Aa^3+\frac{3}{2}Aa^2b+\frac{3}{2}Ba^2b+\frac{3}{8}Bb^3)x+(Aa^3+\frac{3}{2}Aa^2b+\frac{3}{2}Ba^2b+\frac{3}{8}Bb^3)x\left(\tan^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(4Aa^3+6Aa^2b+6Ba^2b+\frac{3}{2}Bb^3)}{24d}$

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/96*(24*(3*A*a*b^2+3*B*a^2*b+B*b^3)*sin(2*d*x+2*c)+8*(A*b^3+3*B*a*b^2)*sin(3*d*x+3*c)+3*B*sin(4*d*x+4*c)*b^3+24*(12*A*a^2*b+3*A*b^3+4*B*a^3+9*B*a*b^2)*sin(d*x+c)+96*x*(A*a^3+3/2*A*a*b^2+3/2*B*a^2*b+3/8*B*b^3)*d)/d
```

3.233.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.80

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{3(8Aa^3 + 12Ba^2b + 12Aab^2 + 3Bb^3)dx + (6Bb^3 \cos(dx + c))^3 + 24Ba^3 + 72Aa^2b + 48Bab^2 + 16Ab^3}{24d}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

output $\frac{1}{24} \cdot (3 \cdot (8Aa^3 + 12B^2a^2b + 12A^2ab^2 + 3B^3b^3) \cdot dx + (6B^3b^3 \cos(dx + c))^3 + 24B^2a^3 + 72A^2a^2b + 48B^2a^2b^2 + 16A^2b^3 + 8(3B^2a^2b^2 + A^2b^3) \cos(dx + c)^2 + 9(4B^2a^2b + 4A^2ab^2 + B^3b^3) \cos(dx + c)) \cdot \sin(dx + c) / d$

3.233.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(170) = 340$.

Time = 0.20 (sec) , antiderivative size = 386, normalized size of antiderivative = 2.26

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \left\{ \begin{array}{l} Aa^3x + \frac{3Aab^2 \sin(c+dx)}{d} + \frac{3Aab^2x \sin^2(c+dx)}{2} + \frac{3Aab^2x \cos^2(c+dx)}{2} + \frac{3Aab^2 \sin(c+dx) \cos(c+dx)}{2d} + \frac{2Ab^3 \sin^3(c+dx)}{3d} + \frac{Ab^3}{3d} \\ x(A + B \cos(c)) (a + b \cos(c))^3 \end{array} \right.$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c)), x)`

output `Piecewise((A**3*x + 3*A**2*b*sin(c + d*x)/d + 3*A*a*b**2*x*sin(c + d*x)**2/2 + 3*A*a*b**2*x*cos(c + d*x)**2/2 + 3*A*a*b**2*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*A*b**3*sin(c + d*x)**3/(3*d) + A*b**3*sin(c + d*x)*cos(c + d*x)**2/d + B*a**3*sin(c + d*x)/d + 3*B*a**2*b*x*sin(c + d*x)**2/2 + 3*B*a**2*b*x*cos(c + d*x)**2/2 + 3*B*a**2*b*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*B*a*b**2*sin(c + d*x)**3/d + 3*B*a*b**2*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*b**3*x*sin(c + d*x)**4/8 + 3*B*b**3*x*cos(c + d*x)**2*cos(c + d*x)**2/4 + 3*B*b**3*x*cos(c + d*x)**4/8 + 3*B*b**3*sin(c + d*x)**3*cos(c + d*x)/(8*d) + 5*B*b**3*sin(c + d*x)*cos(c + d*x)**3/(8*d), Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**3, True))`

3.233.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$= \frac{96(dx + c)Aa^3 + 72(2dx + 2c + \sin(2dx + 2c))Ba^2b + 72(2dx + 2c + \sin(2dx + 2c))Aab^2 - 96(\sin(2dx + 2c) + 2c + 2dx)A^2ab + 96(dx + c)A^2b^3 + 96(dx + c)B^2a^2b + 96(dx + c)B^2ab^2 + 96(dx + c)B^3b^3}{96(dx + c)}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output $\frac{1}{96}(96(d*x + c)*A*a^3 + 72*(2*d*x + 2*c + \sin(2*d*x + 2*c))*B*a^2*b + 7*2*(2*d*x + 2*c + \sin(2*d*x + 2*c))*A*a*b^2 - 96*(\sin(d*x + c)^3 - 3*\sin(d*x + c))*B*a*b^2 - 32*(\sin(d*x + c)^3 - 3*\sin(d*x + c))*A*b^3 + 3*(12*d*x + 12*c + \sin(4*d*x + 4*c) + 8*\sin(2*d*x + 2*c))*B*b^3 + 96*B*a^3*\sin(d*x + c) + 288*A*a^2*b*\sin(d*x + c))/d$

3.233.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.87

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx \\ &= \frac{Bb^3 \sin(4dx + 4c)}{32d} + \frac{1}{8} (8Aa^3 + 12Ba^2b + 12Aab^2 + 3Bb^3)x \\ &+ \frac{(3Bab^2 + Ab^3) \sin(3dx + 3c)}{12d} + \frac{(3Ba^2b + 3Aab^2 + Bb^3) \sin(2dx + 2c)}{4d} \\ &+ \frac{(4Ba^3 + 12Aa^2b + 9Bab^2 + 3Ab^3) \sin(dx + c)}{4d} \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")`

output $\frac{1}{32}B*b^3*\sin(4*d*x + 4*c)/d + \frac{1}{8}*(8*A*a^3 + 12*B*a^2*b + 12*A*a*b^2 + 3*B*b^3)*x + \frac{1}{12}*(3*B*a*b^2 + A*b^3)*\sin(3*d*x + 3*c)/d + \frac{1}{4}*(3*B*a^2*b + 3*A*a*b^2 + B*b^3)*\sin(2*d*x + 2*c)/d + \frac{1}{4}*(4*B*a^3 + 12*A*a^2*b + 9*B*a*b^2 + 3*A*b^3)*\sin(d*x + c)/d$

3.233.9 Mupad [B] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.18

$$\begin{aligned}
& \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx \\
&= A a^3 x + \frac{3 B b^3 x}{8} + \frac{3 A a b^2 x}{2} + \frac{3 B a^2 b x}{2} + \frac{3 A b^3 \sin(c + dx)}{4 d} \\
&+ \frac{B a^3 \sin(c + dx)}{d} + \frac{A b^3 \sin(3 c + 3 d x)}{12 d} + \frac{B b^3 \sin(2 c + 2 d x)}{4 d} \\
&+ \frac{B b^3 \sin(4 c + 4 d x)}{32 d} + \frac{3 A a b^2 \sin(2 c + 2 d x)}{4 d} + \frac{3 B a^2 b \sin(2 c + 2 d x)}{4 d} \\
&+ \frac{B a b^2 \sin(3 c + 3 d x)}{4 d} + \frac{3 A a^2 b \sin(c + d x)}{d} + \frac{9 B a b^2 \sin(c + d x)}{4 d}
\end{aligned}$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3,x)`output `A*a^3*x + (3*B*b^3*x)/8 + (3*A*a*b^2*x)/2 + (3*B*a^2*b*x)/2 + (3*A*b^3*sin(c + d*x))/(4*d) + (B*a^3*sin(c + d*x))/d + (A*b^3*sin(3*c + 3*d*x))/(12*d) + (B*b^3*sin(2*c + 2*d*x))/(4*d) + (B*b^3*sin(4*c + 4*d*x))/(32*d) + (3*A*a*b^2*sin(2*c + 2*d*x))/(4*d) + (3*B*a^2*b*sin(2*c + 2*d*x))/(4*d) + (B*a*b^2*sin(3*c + 3*d*x))/(4*d) + (3*A*a^2*b*sin(c + d*x))/d + (9*B*a*b^2*sin(c + d*x))/(4*d)`

3.234 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec(c+dx) dx$

3.234.1 Optimal result	2152
3.234.2 Mathematica [A] (verified)	2153
3.234.3 Rubi [A] (verified)	2153
3.234.4 Maple [A] (verified)	2157
3.234.5 Fricas [A] (verification not implemented)	2157
3.234.6 Sympy [F]	2158
3.234.7 Maxima [A] (verification not implemented)	2158
3.234.8 Giac [B] (verification not implemented)	2159
3.234.9 Mupad [B] (verification not implemented)	2159

3.234.1 Optimal result

Integrand size = 29, antiderivative size = 137

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{1}{2} (6a^2 Ab + Ab^3 + 2a^3 B + 3ab^2 B) x + \frac{a^3 A \operatorname{arctanh}(\sin(c + dx))}{d}$$

$$+ \frac{b(9aAb + 8a^2 B + 2b^2 B) \sin(c + dx)}{3d}$$

$$+ \frac{b^2(3Ab + 5aB) \cos(c + dx) \sin(c + dx)}{6d} + \frac{bB(a + b \cos(c + dx))^2 \sin(c + dx)}{3d}$$

output `1/2*(6*A*a^2*b+A*b^3+2*B*a^3+3*B*a*b^2)*x+a^3*A*arctanh(sin(d*x+c))/d+1/3*b*(9*A*a*b+8*B*a^2+2*B*b^2)*sin(d*x+c)/d+1/6*b^2*(3*A*b+5*B*a)*cos(d*x+c)*sin(d*x+c)/d+1/3*b*B*(a+b*cos(d*x+c))^2*sin(d*x+c)/d`

3.234.2 Mathematica [A] (verified)

Time = 1.70 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.16

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{6(6a^2Ab + Ab^3 + 2a^3B + 3ab^2B)(c + dx) - 12a^3A \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 12a^3A \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{12d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(6*(6*a^2*A*b + A*b^3 + 2*a^3*B + 3*a*b^2*B)*(c + d*x) - 12*a^3*A*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 12*a^3*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 9*b*(4*a*A*b + 4*a^2*B + b^2*B)*Sin[c + d*x] + 3*b^2*(A*b + 3*a*B)*Sin[2*(c + d*x)] + b^3*B*Sin[3*(c + d*x)])/(12*d)`

3.234.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 3469, 3042, 3512, 3042, 3502, 27, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(c + dx)(a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx$$

$$\downarrow \text{3469}$$

$$\frac{1}{3} \int (a + b \cos(c + dx)) (3Aa^2 + b(3Ab + 5aB) \cos^2(c + dx) + (3Ba^2 + 6Aba + 2b^2B) \cos(c + dx)) \sec(c + dx) dx + \frac{bB \sin(c + dx)(a + b \cos(c + dx))^2}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) \left(3Aa^2 + b(3Ab + 5aB) \sin(c + dx + \frac{\pi}{2})^2 + (3Ba^2 + 6Aba + 2b^2B) \sin(c + dx + \frac{\pi}{2}) \right)}{bB \sin(c + dx)(a + b \cos(c + dx))^2} dx$$

\downarrow 3512

$$\frac{1}{3} \left(\frac{1}{2} \int (6Aa^3 + 2b(8Ba^2 + 9Aba + 2b^2B) \cos^2(c + dx) + 3(2Ba^3 + 6Aba^2 + 3b^2Ba + Ab^3) \cos(c + dx)) \sec(c + dx) dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx$$

\downarrow 3042

$$\frac{1}{3} \left(\frac{1}{2} \int \frac{6Aa^3 + 2b(8Ba^2 + 9Aba + 2b^2B) \sin(c + dx + \frac{\pi}{2})^2 + 3(2Ba^3 + 6Aba^2 + 3b^2Ba + Ab^3) \sin(c + dx + \frac{\pi}{2})}{bB \sin(c + dx)(a + b \cos(c + dx))^2} dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx$$

\downarrow 3502

$$\frac{1}{3} \left(\frac{1}{2} \left(\int 3(2Aa^3 + (2Ba^3 + 6Aba^2 + 3b^2Ba + Ab^3) \cos(c + dx)) \sec(c + dx) dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx$$

\downarrow 27

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int (2Aa^3 + (2Ba^3 + 6Aba^2 + 3b^2Ba + Ab^3) \cos(c + dx)) \sec(c + dx) dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx$$

\downarrow 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{2Aa^3 + (2Ba^3 + 6Aba^2 + 3b^2Ba + Ab^3) \sin(c + dx + \frac{\pi}{2})}{bB \sin(c + dx)(a + b \cos(c + dx))^2} dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx + \frac{2b(8a^2B + 9aAb + 2b^2B) \sin(c + dx)}{d} \right) dx$$

\downarrow 3214

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(2a^3 A \int \sec(c+dx) dx + x(2a^3 B + 6a^2 Ab + 3ab^2 B + Ab^3) \right) + \frac{2b(8a^2 B + 9aAb + 2b^2 B) \sin(c+dx)}{d} \right) + \frac{bB \sin(c+dx)(a+b \cos(c+dx))^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(2a^3 A \int \csc \left(c+dx + \frac{\pi}{2} \right) dx + x(2a^3 B + 6a^2 Ab + 3ab^2 B + Ab^3) \right) + \frac{2b(8a^2 B + 9aAb + 2b^2 B) \sin(c+dx)}{d} \right) + \frac{bB \sin(c+dx)(a+b \cos(c+dx))^2}{3d} \right)$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{2b(8a^2 B + 9aAb + 2b^2 B) \sin(c+dx)}{d} + 3 \left(\frac{2a^3 A \operatorname{arctanh}(\sin(c+dx))}{d} + x(2a^3 B + 6a^2 Ab + 3ab^2 B + Ab^3) \right) + \frac{bB \sin(c+dx)(a+b \cos(c+dx))^2}{3d} \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x], x]`

output `(b*B*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + ((b^2*(3*A*b + 5*a*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d) + (3*((6*a^2*A*b + A*b^3 + 2*a^3*B + 3*a*b^2*B)*x + (2*a^3*A*ArcTanh[Sin[c + d*x]])/d) + (2*b*(9*a*A*b + 8*a^2*B + 2*b^2*B)*Sin[c + d*x])/d)/2)/3`

3.234.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2
, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.234.4 Maple [A] (verified)

Time = 2.92 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.99

method	result
parts	$\frac{A a^3 \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{(A b^3+3B a b^2) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{(3A a b^2+3B a^2 b) \sin(dx+c)}{d} +$
parallelrisch	$\frac{-12A a^3 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 12A a^3 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 3(A b^3+3B a b^2) \sin(2dx+2c) + B \sin(3dx+3c)b^3 + 9(4A a b^2 + 3A^2 b^2) \sin(dx+c)}{12d}$
derivativedivides	$\frac{A a^3 \ln(\sec(dx+c)+\tan(dx+c)) + B a^3(dx+c) + 3A a^2 b(dx+c) + 3B \sin(dx+c)a^2 b + 3A \sin(dx+c)a b^2 + 3B a b^2 \left(\frac{\cos(dx+c)}{2}\right)}{d}$
default	$\frac{A a^3 \ln(\sec(dx+c)+\tan(dx+c)) + B a^3(dx+c) + 3A a^2 b(dx+c) + 3B \sin(dx+c)a^2 b + 3A \sin(dx+c)a b^2 + 3B a b^2 \left(\frac{\cos(dx+c)}{2}\right)}{d}$
risch	$3x A a^2 b + \frac{x A b^3}{2} + a^3 B x + \frac{3x B a b^2}{2} - \frac{3ie^{i(dx+c)} A a b^2}{2d} - \frac{3ie^{i(dx+c)} B a^2 b}{2d} - \frac{3ie^{i(dx+c)} B b^3}{8d} + \frac{3ie^{-i(dx+c)} A a b^2}{2d} + \frac{3ie^{-i(dx+c)} B a^2 b}{2d} + \frac{3ie^{-i(dx+c)} B b^3}{8d}$
norman	$\frac{(3A a^2 b + \frac{1}{2} A b^3 + B a^3 + \frac{3}{2} B a b^2)x + (3A a^2 b + \frac{1}{2} A b^3 + B a^3 + \frac{3}{2} B a b^2)x \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (12A a^2 b + 2A b^3 + 4B a^3 + 6B a b^2) \sin(dx+c)}{6d}$

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)
```

```
output A*a^3/d*ln(sec(d*x+c)+tan(d*x+c))+(A*b^3+3*B*a*b^2)/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+(3*A*a*b^2+3*B*a^2*b)/d*sin(d*x+c)+(3*A*a^2*b+B*a^3)/d*(d*x+c)+1/3*B*b^3/d*(2+cos(d*x+c)^2)*sin(d*x+c)
```

3.234.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.96

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{3 A a^3 \log(\sin(dx + c) + 1) - 3 A a^3 \log(-\sin(dx + c) + 1) + 3(2 B a^3 + 6 A a^2 b + 3 B a b^2 + A b^3) dx + (2 B a^3 + 6 A a^2 b + 3 B a b^2 + A b^3) \sin(dx + c)}{6 d}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")
```

output $1/6*(3*A*a^3*\log(\sin(d*x + c) + 1) - 3*A*a^3*\log(-\sin(d*x + c) + 1) + 3*(2*B*a^3 + 6*A*a^2*b + 3*B*a*b^2 + A*b^3)*d*x + (2*B*b^3*\cos(d*x + c)^2 + 18*B*a^2*b + 18*A*a*b^2 + 4*B*b^3 + 3*(3*B*a*b^2 + A*b^3)*\cos(d*x + c))*\sin(d*x + c))/d$

3.234.6 Sympy [F]

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^3 \sec(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**3*sec(c + d*x), x)`

3.234.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.06

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{12(dx + c)Ba^3 + 36(dx + c)Aa^2b + 9(2dx + 2c + \sin(2dx + 2c))Bab^2 + 3(2dx + 2c + \sin(2dx + 2c))Aa^3 + 3(2dx + 2c + \sin(2dx + 2c))Bab^2 + 3(2dx + 2c + \sin(2dx + 2c))Aa^3}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output $1/12*(12*(d*x + c)*B*a^3 + 36*(d*x + c)*A*a^2*b + 9*(2*d*x + 2*c + \sin(2*d*x + 2*c))*B*a*b^2 + 3*(2*d*x + 2*c + \sin(2*d*x + 2*c))*A*b^3 - 4*(\sin(d*x + c))^3 - 3*\sin(d*x + c))*B*b^3 + 12*A*a^3*\log(\sec(d*x + c) + \tan(d*x + c)) + 36*B*a^2*b*\sin(d*x + c) + 36*A*a*b^2*\sin(d*x + c))/d$

3.234.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 314 vs. $2(129) = 258$.

Time = 0.33 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.29

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{6 A a^3 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 6 A a^3 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + 3 (2 B a^3 + 6 A a^2 b + 3 B a b^2 + A b^3) (d x + c) + 2 (18 B a^2 b \tan(1/2 dx + 1/2 c)^5 + 18 A a b^2 \tan(1/2 dx + 1/2 c)^5 - 9 B a b^2 \tan(1/2 dx + 1/2 c)^5 - 3 A b^3 \tan(1/2 dx + 1/2 c)^5 + 6 B b^3 \tan(1/2 dx + 1/2 c)^5 + 36 B a^2 b \tan(1/2 dx + 1/2 c)^3 + 36 A a b^2 \tan(1/2 dx + 1/2 c)^3 + 4 B b^3 \tan(1/2 dx + 1/2 c)^3 + 18 B a^2 b \tan(1/2 dx + 1/2 c) + 18 A a b^2 \tan(1/2 dx + 1/2 c) + 9 B a b^2 \tan(1/2 dx + 1/2 c) + 3 A b^3 \tan(1/2 dx + 1/2 c) + 6 B b^3 \tan(1/2 dx + 1/2 c)) / (\tan(1/2 dx + 1/2 c)^2 + 1)^3}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `1/6*(6*A*a^3*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 6*A*a^3*log(abs(tan(1/2*d*x + 1/2*c) - 1))) + 3*(2*B*a^3 + 6*A*a^2*b + 3*B*a*b^2 + A*b^3)*(d*x + c) + 2*(18*B*a^2*b*tan(1/2*d*x + 1/2*c)^5 + 18*A*a*b^2*tan(1/2*d*x + 1/2*c)^5 - 9*B*a*b^2*tan(1/2*d*x + 1/2*c)^5 - 3*A*b^3*tan(1/2*d*x + 1/2*c)^5 + 6*B*b^3*tan(1/2*d*x + 1/2*c)^5 + 36*B*a^2*b*tan(1/2*d*x + 1/2*c)^3 + 36*A*a*b^2*tan(1/2*d*x + 1/2*c)^3 + 4*B*b^3*tan(1/2*d*x + 1/2*c)^3 + 18*B*a^2*b*tan(1/2*d*x + 1/2*c) + 18*A*a*b^2*tan(1/2*d*x + 1/2*c) + 9*B*a*b^2*tan(1/2*d*x + 1/2*c) + 3*A*b^3*tan(1/2*d*x + 1/2*c) + 6*B*b^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^3/d`

3.234.9 Mupad [B] (verification not implemented)

Time = 2.04 (sec) , antiderivative size = 1924, normalized size of antiderivative = 14.04

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec(c + dx) dx = \text{Too large to display}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x),x)`

output

$$\begin{aligned}
& (\tan(c/2 + (dx)/2)*(A*b^3 + 2*B*b^3 + 6*A*a*b^2 + 3*B*a*b^2 + 6*B*a^2*b) \\
& + \tan(c/2 + (dx)/2)^3*((4*B*b^3)/3 + 12*A*a*b^2 + 12*B*a^2*b) + \tan(c/2 + \\
& (dx)/2)^5*(2*B*b^3 - A*b^3 + 6*A*a*b^2 - 3*B*a*b^2 + 6*B*a^2*b))/(d*(3*t \\
& \tan(c/2 + (dx)/2)^2 + 3*\tan(c/2 + (dx)/2)^4 + \tan(c/2 + (dx)/2)^6 + 1)) \\
& + (\operatorname{atan}((((A*b^3*i)/2 + B*a^3*i + A*a^2*b*3i + (B*a*b^2*3i)/2)*(32*A*a^ \\
& 3 + 16*A*b^3 + 32*B*a^3 + 96*A*a^2*b + 48*B*a*b^2) + \tan(c/2 + (dx)/2)*(3 \\
& 2*A^2*a^6 + 8*A^2*b^6 + 32*B^2*a^6 + 96*A^2*a^2*b^4 + 288*A^2*a^4*b^2 + 72 \\
& *B^2*a^2*b^4 + 96*B^2*a^4*b^2 + 48*A*B*a*b^5 + 192*A*B*a^5*b + 320*A*B*a^3 \\
& *b^3))*((A*b^3*i)/2 + B*a^3*i + A*a^2*b*3i + (B*a*b^2*3i)/2)*i - (((A*b \\
& ^3*i)/2 + B*a^3*i + A*a^2*b*3i + (B*a*b^2*3i)/2)*(32*A*a^3 + 16*A*b^3 + \\
& 32*B*a^3 + 96*A*a^2*b + 48*B*a*b^2) - \tan(c/2 + (dx)/2)*(32*A^2*a^6 + 8*A \\
& ^2*b^6 + 32*B^2*a^6 + 96*A^2*a^2*b^4 + 288*A^2*a^4*b^2 + 72*B^2*a^2*b^4 + \\
& 96*B^2*a^4*b^2 + 48*A*B*a*b^5 + 192*A*B*a^5*b + 320*A*B*a^3*b^3))*((A*b^3* \\
& i)/2 + B*a^3*i + A*a^2*b*3i + (B*a*b^2*3i)/2)*i)/((((A*b^3*i)/2 + B*a^ \\
& 3*i + A*a^2*b*3i + (B*a*b^2*3i)/2)*(32*A*a^3 + 16*A*b^3 + 32*B*a^3 + 96*A \\
& *a^2*b + 48*B*a*b^2) + \tan(c/2 + (dx)/2)*(32*A^2*a^6 + 8*A^2*b^6 + 32*B^2 \\
& *a^6 + 96*A^2*a^2*b^4 + 288*A^2*a^4*b^2 + 72*B^2*a^2*b^4 + 96*B^2*a^4*b^2 \\
& + 48*A*B*a*b^5 + 192*A*B*a^5*b + 320*A*B*a^3*b^3))*((A*b^3*i)/2 + B*a^3*i \\
& + A*a^2*b*3i + (B*a*b^2*3i)/2) + (((A*b^3*i)/2 + B*a^3*i + A*a^2*b*3i \\
& + (B*a*b^2*3i)/2)*(32*A*a^3 + 16*A*b^3 + 32*B*a^3 + 96*A*a^2*b + 48*B*a...
\end{aligned}$$

3.235 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.235.1 Optimal result	2161
3.235.2 Mathematica [A] (verified)	2162
3.235.3 Rubi [A] (verified)	2162
3.235.4 Maple [A] (verified)	2166
3.235.5 Fricas [A] (verification not implemented)	2166
3.235.6 Sympy [F]	2167
3.235.7 Maxima [A] (verification not implemented)	2167
3.235.8 Giac [A] (verification not implemented)	2168
3.235.9 Mupad [B] (verification not implemented)	2168

3.235.1 Optimal result

Integrand size = 31, antiderivative size = 131

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{1}{2}b(6aAb + 6a^2B + b^2B) x + \frac{a^2(3Ab + aB)\operatorname{arctanh}(\sin(c + dx))}{d}$$

$$- \frac{b(2a^2A - Ab^2 - 3abB) \sin(c + dx)}{d} - \frac{b^2(2aA - bB) \cos(c + dx) \sin(c + dx)}{2d}$$

$$+ \frac{aA(a + b \cos(c + dx))^2 \tan(c + dx)}{d}$$

```
output 1/2*b*(6*A*a*b+6*B*a^2+B*b^2)*x+a^2*(3*A*b+B*a)*arctanh(sin(d*x+c))/d-b*(2
*A*a^2-A*b^2-3*B*a*b)*sin(d*x+c)/d-1/2*b^2*(2*A*a-B*b)*cos(d*x+c)*sin(d*x+
c)/d+a*A*(a+b*cos(d*x+c))^2*tan(d*x+c)/d
```


3.235.2 Mathematica [A] (verified)

Time = 1.96 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.66

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{2b(6aAb + 6a^2B + b^2B)(c + dx) - 4a^2(3Ab + aB) \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + 4a^2(3Ab + aB) \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{4d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output $(2*b*(6*a*A*b + 6*a^2*B + b^2*B)*(c + d*x) - 4*a^2*(3*A*b + a*B)*\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] + 4*a^2*(3*A*b + a*B)*\text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]] + (4*a^3*A*\text{Sin}[(c + d*x)/2])/(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]) + (4*a^3*A*\text{Sin}[(c + d*x)/2])/(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]) + 4*b^2*(A*b + 3*a*B)*\text{Sin}[c + d*x] + b^3*B*\text{Sin}[2*(c + d*x)])/(4*d)$

3.235.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3468, 3042, 3512, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx$$

$$\downarrow \text{3468}$$

$$\int (a + b \cos(c + dx)) (-b(2aA - bB) \cos^2(c + dx) + b(Ab + 2aB) \cos(c + dx) + a(3Ab + aB)) \sec(c + dx) dx + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) \left(-b(2aA - bB) \sin(c + dx + \frac{\pi}{2})^2 + b(Ab + 2aB) \sin(c + dx + \frac{\pi}{2}) + a(3Ab + aB) \right)}{\frac{\sin(c + dx + \frac{\pi}{2})}{aA \tan(c + dx)(a + b \cos(c + dx))^2}} dx$$

↓ 3512

$$\frac{1}{2} \int (2(3Ab + aB)a^2 - 2b(2Aa^2 - 3bBa - Ab^2) \cos^2(c + dx) + b(6Ba^2 + 6Aba + b^2B) \cos(c + dx)) \sec(c + dx) dx - \frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

↓ 3042

$$\frac{1}{2} \int \frac{2(3Ab + aB)a^2 - 2b(2Aa^2 - 3bBa - Ab^2) \sin(c + dx + \frac{\pi}{2})^2 + b(6Ba^2 + 6Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\frac{\sin(c + dx + \frac{\pi}{2})}{\frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}}} dx -$$

↓ 3502

$$\frac{1}{2} \left(\int (2(3Ab + aB)a^2 + b(6Ba^2 + 6Aba + b^2B) \cos(c + dx)) \sec(c + dx) dx - \frac{2b(2a^2A - 3abB - Ab^2) \sin(c + dx)}{d} \right) - \frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{2(3Ab + aB)a^2 + b(6Ba^2 + 6Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{2b(2a^2A - 3abB - Ab^2) \sin(c + dx)}{d} \right) - \frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

↓ 3214

$$\frac{1}{2} \left(2a^2(aB + 3Ab) \int \sec(c + dx) dx - \frac{2b(2a^2A - 3abB - Ab^2) \sin(c + dx)}{d} + bx(6a^2B + 6aAb + b^2B) \right) - \frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

↓ 3042

$$\frac{1}{2} \left(2a^2(aB + 3Ab) \int \csc \left(c + dx + \frac{\pi}{2} \right) dx - \frac{2b(2a^2A - 3abB - Ab^2) \sin(c + dx)}{d} + bx(6a^2B + 6aAb + b^2B) \right) - \frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

↓ 4257

$$\frac{1}{2} \left(\frac{2a^2(aB + 3Ab) \operatorname{arctanh}(\sin(c + dx))}{d} - \frac{2b(2a^2A - 3abB - Ab^2) \sin(c + dx)}{d} + bx(6a^2B + 6aAb + b^2B) \right) - \frac{b^2(2aA - bB) \sin(c + dx) \cos(c + dx)}{2d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^2}{d}$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `-1/2*(b^2*(2*a*A - b*B)*Cos[c + d*x]*Sin[c + d*x])/d + (b*(6*a*A*b + 6*a^2*B + b^2*B)*x + (2*a^2*(3*A*b + a*B)*ArcTanh[Sin[c + d*x]])/d - (2*b*(2*a^2*A - A*b^2 - 3*a*b*B)*Sin[c + d*x])/d)/2 + (a*A*(a + b*Cos[c + d*x])^2*Tan[c + d*x])/d`

3.235.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3512 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Sin[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) + A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.235.4 Maple [A] (verified)

Time = 3.24 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.95

method	result
parts	$\frac{A a^3 \tan(dx+c)}{d} + \frac{(A b^3+3B a b^2) \sin(dx+c)}{d} + \frac{(3A a b^2+3B a^2 b)(dx+c)}{d} + \frac{(3A a^2 b+B a^3) \ln(\sec(dx+c)+\tan(dx+c))}{d}$
derivativedivides	$\frac{A a^3 \tan(dx+c)+B a^3 \ln(\sec(dx+c)+\tan(dx+c))+3A a^2 b \ln(\sec(dx+c)+\tan(dx+c))+3B a^2 b(dx+c)+3A a b^2(dx+c)+3B b^3}{d}$
default	$\frac{A a^3 \tan(dx+c)+B a^3 \ln(\sec(dx+c)+\tan(dx+c))+3A a^2 b \ln(\sec(dx+c)+\tan(dx+c))+3B a^2 b(dx+c)+3A a b^2(dx+c)+3B b^3}{d}$
parallelrisch	$\frac{8(-3A a^2 b-B a^3) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+8(3A a^2 b+B a^3) \cos(dx+c) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)+4(A b^3+3B a b^2) \sin(dx+c)}{8d \cos(dx+c)}$
risch	$3x A a b^2 + 3x B a^2 b + \frac{b^3 B x}{2} - \frac{i B b^3 e^{2i(dx+c)}}{8d} - \frac{i e^{i(dx+c)} A b^3}{2d} - \frac{3i e^{i(dx+c)} B a b^2}{2d} + \frac{i e^{-i(dx+c)} A b^3}{2d} + \frac{3i e^{-i(dx+c)} B a^2 b}{2d}$
norman	$\frac{(-3A a b^2-3B a^2 b-\frac{1}{2} B b^3)x+(-9A a b^2-9B a^2 b-\frac{3}{2} B b^3)x\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(3A a b^2+3B a^2 b+\frac{1}{2} B b^3)x\left(\tan^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d}$

input `int((a+cos(d*x+c))*b^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBOSE)`

output `A*a^3/d*tan(d*x+c)+(A*b^3+3*B*a*b^2)/d*sin(d*x+c)+(3*A*a*b^2+3*B*a^2*b)/d*(d*x+c)+(3*A*a^2*b+B*a^3)/d*ln(sec(d*x+c)+tan(d*x+c))+B*b^3/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)`

3.235.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.16

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{(6 B a^2 b + 6 A a b^2 + B b^3) dx \cos(dx + c) + (B a^3 + 3 A a^2 b) \cos(dx + c) \log(\sin(dx + c) + 1) - (B a^3 + 3 A a^2 b) \cos(dx + c) \log(\sin(dx + c) - 1) + (6 B a^2 b + 6 A a b^2 + B b^3) dx \sin(dx + c) + (B a^3 + 3 A a^2 b) \sin(dx + c) \log(\sin(dx + c) + 1) - (B a^3 + 3 A a^2 b) \sin(dx + c) \log(\sin(dx + c) - 1)}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fracas")`

output $1/2*((6*B*a^2*b + 6*A*a*b^2 + B*b^3)*d*x*cos(d*x + c) + (B*a^3 + 3*A*a^2*b)*cos(d*x + c)*log(sin(d*x + c) + 1) - (B*a^3 + 3*A*a^2*b)*cos(d*x + c)*log(-sin(d*x + c) + 1) + (B*b^3*cos(d*x + c)^2 + 2*A*a^3 + 2*(3*B*a*b^2 + A*b^3)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c))$

3.235.6 Sympy [F]

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^3 \sec^2(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**3*sec(c + d*x)**2, x)`

3.235.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.10

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{12(dx + c)Ba^2b + 12(dx + c)Aab^2 + (2dx + 2c + \sin(2dx + 2c))Bb^3 + 2Ba^3(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1))}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output $1/4*(12*(d*x + c)*B*a^2*b + 12*(d*x + c)*A*a*b^2 + (2*d*x + 2*c + \sin(2*d*x + 2*c))*B*b^3 + 2*B*a^3*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 6*A*a^2*b*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 12*B*a*b^2*\sin(d*x + c) + 4*A*b^3*\sin(d*x + c) + 4*A*a^3*\tan(d*x + c))/d$

3.235.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.79

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$\frac{4 A a^3 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1} - (6 B a^2 b + 6 A a b^2 + B b^3)(dx + c) - 2 (B a^3 + 3 A a^2 b) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) +$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `-1/2*(4*A*a^3*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 1) - (6*B*a^2*b + 6*A*a*b^2 + B*b^3)*(d*x + c) - 2*(B*a^3 + 3*A*a^2*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) + 2*(B*a^3 + 3*A*a^2*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*B*a*b^2*tan(1/2*d*x + 1/2*c)^3 + 2*A*b^3*tan(1/2*d*x + 1/2*c)^3 - B*b^3*tan(1/2*d*x + 1/2*c)^3 + 6*B*a*b^2*tan(1/2*d*x + 1/2*c) + 2*A*b^3*tan(1/2*d*x + 1/2*c) + B*b^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^2)/d`

3.235.9 Mupad [B] (verification not implemented)

Time = 1.46 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.80

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{B b^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + 6 A a b^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + 6 B a^2 b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) - B a^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + \frac{A b^3 \sin(2c+2dx)}{2} + \frac{B b^3 \sin(3c+3dx)}{8} + A a^3 \sin(c+dx) + \frac{B b^3 \sin(c+dx)}{8} + \frac{3 B a b^2 \sin(2c+2dx)}{2}}{d \cos(c+dx)}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^2,x)`

output $(B*b^3*\text{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) - B*a^3*\text{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*2i + 6*A*a*b^2*\text{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) - A*a^2*b*\text{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*6i + 6*B*a^2*b*\text{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))/d + ((A*b^3*\sin(2*c + 2*d*x))/2 + (B*b^3*\sin(3*c + 3*d*x))/8 + A*a^3*\sin(c + d*x) + (B*b^3*\sin(c + d*x))/8 + (3*B*a*b^2*\sin(2*c + 2*d*x))/2)/(d*\cos(c + d*x))$

3.236 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.236.1 Optimal result	2170
3.236.2 Mathematica [B] (verified)	2171
3.236.3 Rubi [A] (verified)	2171
3.236.4 Maple [A] (verified)	2175
3.236.5 Fricas [A] (verification not implemented)	2175
3.236.6 Sympy [F]	2176
3.236.7 Maxima [A] (verification not implemented)	2176
3.236.8 Giac [B] (verification not implemented)	2177
3.236.9 Mupad [B] (verification not implemented)	2177

3.236.1 Optimal result

Integrand size = 31, antiderivative size = 124

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= b^2(Ab + 3aB)x + \frac{a(a^2A + 6Ab^2 + 6abB) \operatorname{arctanh}(\sin(c + dx))}{2d}$$

$$- \frac{b^2(aA - 2bB) \sin(c + dx)}{2d} + \frac{a^2(2Ab + aB) \tan(c + dx)}{d}$$

$$+ \frac{aA(a + b \cos(c + dx))^2 \sec(c + dx) \tan(c + dx)}{2d}$$

```
output b^2*(A*b+3*B*a)*x+1/2*a*(A*a^2+6*A*b^2+6*B*a*b)*arctanh(sin(d*x+c))/d-1/2*
b^2*(A*a-2*B*b)*sin(d*x+c)/d+a^2*(2*A*b+B*a)*tan(d*x+c)/d+1/2*a*A*(a+b*cos
(d*x+c))^2*sec(d*x+c)*tan(d*x+c)/d
```

3.236.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 277 vs. $2(124) = 248$.

Time = 3.29 (sec) , antiderivative size = 277, normalized size of antiderivative = 2.23

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4b^2(Ab + 3aB)(c + dx) - 2a(a^2A + 6Ab^2 + 6abB) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2a(a^2A + 6Ab^2 + 6abB) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)}{4d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output $(4b^2(Ab + 3aB)(c + dx) - 2a(a^2A + 6Ab^2 + 6abB) \log[\cos((c + dx)/2) - \sin((c + dx)/2)] + 2a(a^2A + 6Ab^2 + 6abB) \log[\cos((c + dx)/2) + \sin((c + dx)/2)] + (a^3A)/(\cos((c + dx)/2) - \sin((c + dx)/2))^2 + (4a^2(3Ab + aB) \sin((c + dx)/2))/(\cos((c + dx)/2) - \sin((c + dx)/2)) - (a^3A)/(\cos((c + dx)/2) + \sin((c + dx)/2))^2 + (4a^2(3Ab + aB) \sin((c + dx)/2))/(\cos((c + dx)/2) + \sin((c + dx)/2)) + 4b^3B \sin[c + d*x])/(4*d)$

3.236.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3468, 3042, 3510, 25, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^3(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{2} \int (a + b \cos(c + dx)) \left(-b(aA - 2bB) \cos^2(c + dx) + (Aa^2 + 4bBa + 2Ab^2) \cos(c + dx) + 2a(2Ab + aB) \right) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^2}{2d}$$

↓ 3042

$$\frac{1}{2} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) \left(-b(aA - 2bB) \sin(c + dx + \frac{\pi}{2})^2 + (Aa^2 + 4bBa + 2Ab^2) \sin(c + dx + \frac{\pi}{2}) + 2a(2Ab + aB) \right) \sec^2(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^2}{2d}$$

↓ 3510

$$\frac{1}{2} \left(\frac{2a^2(aB + 2Ab) \tan(c + dx)}{d} - \int - \left(-((aA - 2bB) \cos^2(c + dx)b^2) + 2(Ab + 3aB) \cos(c + dx)b^2 + a(Aa^2 + 6bBa + 6Ab^2) \right) \sec(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^2}{2d} \right)$$

↓ 25

$$\frac{1}{2} \left(\int - \left((aA - 2bB) \cos^2(c + dx)b^2 + 2(Ab + 3aB) \cos(c + dx)b^2 + a(Aa^2 + 6bBa + 6Ab^2) \right) \sec(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^2}{2d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{- \left((aA - 2bB) \sin(c + dx + \frac{\pi}{2})^2 b^2 + 2(Ab + 3aB) \sin(c + dx + \frac{\pi}{2}) b^2 + a(Aa^2 + 6bBa + 6Ab^2) \right) \sec(c + dx) dx + \frac{2a^2(aB + 2Ab) \tan(c + dx)}{d}}{\sin(c + dx + \frac{\pi}{2})} + \frac{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^2}{2d} \right)$$

↓ 3502

$$\frac{1}{2} \left(\int \left(2(Ab + 3aB) \cos(c + dx)b^2 + a(Aa^2 + 6bBa + 6Ab^2) \right) \sec(c + dx) dx + \frac{2a^2(aB + 2Ab) \tan(c + dx)}{d} - \frac{b^2(aA - 2bB) \cos^2(c + dx)}{2d} + \frac{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^2}{2d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{2(Ab + 3aB) \sin(c + dx + \frac{\pi}{2}) b^2 + a(Aa^2 + 6bBa + 6Ab^2)}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a^2(aB + 2Ab) \tan(c + dx)}{d} - \frac{b^2(aA - 2bB)}{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^2} \right)$$

↓ 3214

$$\frac{1}{2} \left(a(a^2A + 6abB + 6Ab^2) \int \sec(c + dx) dx + \frac{2a^2(aB + 2Ab) \tan(c + dx)}{d} - \frac{b^2(aA - 2bB) \sin(c + dx)}{d} + 2b^2x(3) \right)$$

↓ 3042

$$\frac{1}{2} \left(a(a^2A + 6abB + 6Ab^2) \int \csc(c + dx + \frac{\pi}{2}) dx + \frac{2a^2(aB + 2Ab) \tan(c + dx)}{d} - \frac{b^2(aA - 2bB) \sin(c + dx)}{d} + 2b^2x(3) \right)$$

↓ 4257

$$\frac{1}{2} \left(\frac{a(a^2A + 6abB + 6Ab^2) \operatorname{arctanh}(\sin(c + dx))}{d} + \frac{2a^2(aB + 2Ab) \tan(c + dx)}{d} - \frac{b^2(aA - 2bB) \sin(c + dx)}{d} + 2b^2x(3) \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(a*A*(a + b*Cos[c + d*x])^2*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (2*b^2*(A*b + 3*a*B)*x + (a*(a^2*A + 6*A*b^2 + 6*a*b*B)*ArcTanh[Sin[c + d*x]])/d - (b^2*(a*A - 2*b*B)*Sin[c + d*x])/d + (2*a^2*(2*A*b + a*B)*Tan[c + d*x])/d)/2`

3.236.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2)), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.236.4 Maple [A] (verified)

Time = 3.73 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.07

method	result
parts	$\frac{A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d} + \frac{(A b^3 + 3B a b^2)(dx+c)}{d} + \frac{(3A a b^2 + 3B a^2 b) \ln(\sec(dx+c)+\tan(dx+c))}{d}$
derivativdivides	$\frac{A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) + B a^3 \tan(dx+c) + 3A a^2 b \tan(dx+c) + 3B a^2 b \ln(\sec(dx+c)+\tan(dx+c))}{d}$
default	$\frac{A a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right) + B a^3 \tan(dx+c) + 3A a^2 b \tan(dx+c) + 3B a^2 b \ln(\sec(dx+c)+\tan(dx+c))}{d}$
parallelrisch	$\frac{-a(1+\cos(2dx+2c))(A a^2+6A b^2+6B ab) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)+a(1+\cos(2dx+2c))(A a^2+6A b^2+6B ab) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{2d}$
risch	$x A b^3 + 3x B a b^2 - \frac{i e^{i(dx+c)} B b^3}{2d} + \frac{i e^{-i(dx+c)} B b^3}{2d} - \frac{i a^2 (A a e^{3i(dx+c)} - 6A b e^{2i(dx+c)} - 2B a e^{2i(dx+c)} - a A e^{i(dx+c)})}{d(e^{2i(dx+c)}+1)^2}$
norman	$\frac{(A b^3 + 3B a b^2)x + (-4A b^3 - 12B a b^2)x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-A b^3 - 3B a b^2)x \left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-A b^3 - 3B a b^2)x \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d}$

input `int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `A*a^3/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^3+3*B*a*b^2)/d*(d*x+c)+(3*A*a*b^2+3*B*a^2*b)/d*ln(sec(d*x+c)+tan(d*x+c))+(3*A*a^2*b+B*a^3)/d*tan(d*x+c)+B*b^3/d*sin(d*x+c)`

3.236.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.35

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4(3Bab^2 + Ab^3)dx \cos(dx+c)^2 + (Aa^3 + 6Ba^2b + 6Aab^2) \cos(dx+c)^2 \log(\sin(dx+c)+1) - (Aa^3 + 6Ba^2b + 6Aab^2) \cos(dx+c)^2}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fracas")`

output $1/4*(4*(3*B*a*b^2 + A*b^3)*d*x*cos(d*x + c)^2 + (A*a^3 + 6*B*a^2*b + 6*A*a*b^2)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A*a^3 + 6*B*a^2*b + 6*A*a*b^2)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(2*B*b^3*cos(d*x + c)^2 + A*a^3 + 2*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)$

3.236.6 Sympy [F]

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^3 \sec^3(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**3*sec(c + d*x)**3, x)`

3.236.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.36

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{12(dx + c)Bab^2 + 4(dx + c)Ab^3 - Aa^3 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) + 6Aa^2b \left(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1) \right) + 6Aa^2b^2 \left(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1) \right) + 4Bb^3 \sin(dx + c) + 4Bb^3 \tan(dx + c) + 12Aa^2b \tan(dx + c)}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output $1/4*(12*(d*x + c)*B*a*b^2 + 4*(d*x + c)*A*b^3 - A*a^3*(2*\sin(d*x + c)/(sin(d*x + c)^2 - 1) - \log(sin(d*x + c) + 1) + \log(sin(d*x + c) - 1)) + 6*B*a^2*b*(\log(sin(d*x + c) + 1) - \log(sin(d*x + c) - 1)) + 6*A*a^2*b^2*(\log(sin(d*x + c) + 1) - \log(sin(d*x + c) - 1)) + 4*B*b^3*\sin(d*x + c) + 4*B*b^3*\tan(d*x + c) + 12*A*a^2*b*\tan(d*x + c))/d$

3.236.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(118) = 236$.

Time = 0.32 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.93

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{4Bb^3 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1} + 2(3Bab^2 + Ab^3)(dx + c) + (Aa^3 + 6Ba^2b + 6Aab^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|) - (A$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output $\frac{1}{2} * (4 * B * b^3 * \tan(1/2 * d * x + 1/2 * c) / (\tan(1/2 * d * x + 1/2 * c)^2 + 1) + 2 * (3 * B * a * b^2 + A * b^3) * (d * x + c) + (A * a^3 + 6 * B * a^2 * b + 6 * A * a * b^2) * \log(\text{abs}(\tan(1/2 * d * x + 1/2 * c) + 1)) - (A * a^3 + 6 * B * a^2 * b + 6 * A * a * b^2) * \log(\text{abs}(\tan(1/2 * d * x + 1/2 * c) - 1)) + 2 * (A * a^3 * \tan(1/2 * d * x + 1/2 * c)^3 - 2 * B * a^3 * \tan(1/2 * d * x + 1/2 * c)^3 - 6 * A * a^2 * b * \tan(1/2 * d * x + 1/2 * c)^3 + A * a^3 * \tan(1/2 * d * x + 1/2 * c) + 2 * B * a^3 * \tan(1/2 * d * x + 1/2 * c) + 6 * A * a^2 * b * \tan(1/2 * d * x + 1/2 * c)) / (\tan(1/2 * d * x + 1/2 * c)^2 - 1)^2) / d$

3.236.9 Mupad [B] (verification not implemented)

Time = 1.67 (sec) , antiderivative size = 249, normalized size of antiderivative = 2.01

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{\frac{Ba^3 \sin(2c+2dx)}{2} + \frac{Bb^3 \sin(3c+3dx)}{4} + \frac{Aa^3 \sin(c+dx)}{2} + \frac{Bb^3 \sin(c+dx)}{4} + \frac{3Aa^2b \sin(2c+2dx)}{2}}{d \left(\frac{\cos(2c+2dx)}{2} + \frac{1}{2} \right)}$$

$$2 \left(\frac{Aa^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right) \operatorname{li}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \operatorname{li}}{2} - Ab^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + Aab^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right) \operatorname{li}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \operatorname{li} - 3Bab^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \right)$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^3,x)`

3.236. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

output $((B*a^3*\sin(2*c + 2*d*x))/2 + (B*b^3*\sin(3*c + 3*d*x))/4 + (A*a^3*\sin(c + d*x))/2 + (B*b^3*\sin(c + d*x))/4 + (3*A*a^2*b*\sin(2*c + 2*d*x))/2)/(d*(\cos(2*c + 2*d*x)/2 + 1/2)) - (2*((A*a^3*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*1i)/2 - A*b^3*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + A*a*b^2*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*3i - 3*B*a*b^2*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + B*a^2*b*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*3i))/d$

3.237 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.237.1 Optimal result	2179
3.237.2 Mathematica [A] (verified)	2180
3.237.3 Rubi [A] (verified)	2180
3.237.4 Maple [A] (verified)	2184
3.237.5 Fricas [A] (verification not implemented)	2185
3.237.6 Sympy [F(-1)]	2185
3.237.7 Maxima [A] (verification not implemented)	2186
3.237.8 Giac [B] (verification not implemented)	2186
3.237.9 Mupad [B] (verification not implemented)	2187

3.237.1 Optimal result

Integrand size = 31, antiderivative size = 145

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= b^3 B x + \frac{(3a^2 A b + 2A b^3 + a^3 B + 6a b^2 B) \operatorname{arctanh}(\sin(c + dx))}{2d}$$

$$+ \frac{a(2a^2 A + 8A b^2 + 9a b B) \tan(c + dx)}{3d} + \frac{a^2(5A b + 3a B) \sec(c + dx) \tan(c + dx)}{6d}$$

$$+ \frac{a A (a + b \cos(c + dx))^2 \sec^2(c + dx) \tan(c + dx)}{3d}$$

output `b^3*B*x+1/2*(3*A*a^2*b+2*A*b^3+B*a^3+6*B*a*b^2)*arctanh(sin(d*x+c))/d+1/3*a*(2*A*a^2+8*A*b^2+9*B*a*b)*tan(d*x+c)/d+1/6*a^2*(5*A*b+3*B*a)*sec(d*x+c)*tan(d*x+c)/d+1/3*a*A*(a+b*cos(d*x+c))^2*sec(d*x+c)^2*tan(d*x+c)/d`

3.237.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.74

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{6b^3 B dx + 3(3a^2 Ab + 2Ab^3 + a^3 B + 6ab^2 B) \operatorname{arctanh}(\sin(c + dx)) + 3a(2a^2 A + 6Ab^2 + 6abB + a(3Ab + a^2 B)) \tan(c + dx) + 2a^3 A \tan^3(c + dx)}{6d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(6*b^3*B*d*x + 3*(3*a^2*A*b + 2*A*b^3 + a^3*B + 6*a*b^2*B)*ArcTanh[Sin[c + d*x]] + 3*a*(2*a^2*A + 6*A*b^2 + 6*a*b*B + a*(3*A*b + a*B))*Tan[c + d*x] + 2*a^3*A*Tan[c + d*x]^3)/(6*d)`

3.237.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3468, 3042, 3510, 25, 3042, 3500, 27, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c + dx)(a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx$$

$$\downarrow 3468$$

$$\frac{1}{3} \int (a + b \cos(c + dx)) (3b^2 B \cos^2(c + dx) + (2Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(5Ab + 3aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^2}{3d}$$

$$\downarrow 3042$$

$$\frac{1}{3} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) \left(3b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (2Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(5Ab + 3aB) \right)}{\sin(c + dx + \frac{\pi}{2})^3} \frac{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2}{3d}$$

↓ 3510

$$\frac{1}{3} \left(\frac{a^2(3aB + 5Ab) \tan(c + dx) \sec(c + dx)}{2d} - \frac{1}{2} \int \frac{-((6B \cos^2(c + dx)b^3 + 2a(2Aa^2 + 9bBa + 8Ab^2) + 3(Ba^3 + 3Aba^2 + 6b^2Ba + 2Ab^3)) \cos(c + dx)) \sec^2(c + dx)}{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2} \frac{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2}{3d} \right)$$

↓ 25

$$\frac{1}{3} \left(\frac{1}{2} \int \frac{(6B \cos^2(c + dx)b^3 + 2a(2Aa^2 + 9bBa + 8Ab^2) + 3(Ba^3 + 3Aba^2 + 6b^2Ba + 2Ab^3)) \cos(c + dx) \sec^2(c + dx)}{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2} \frac{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \int \frac{6B \sin(c + dx + \frac{\pi}{2})^2 b^3 + 2a(2Aa^2 + 9bBa + 8Ab^2) + 3(Ba^3 + 3Aba^2 + 6b^2Ba + 2Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} \frac{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2}{3d} \right)$$

↓ 3500

$$\frac{1}{3} \left(\frac{1}{2} \left(\int 3(Ba^3 + 3Aba^2 + 6b^2Ba + 2Ab^3 + 2b^3B \cos(c + dx)) \sec(c + dx) dx + \frac{2a(2a^2A + 9abB + 8Ab^2) \tan(c + dx)}{d} \right) \frac{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2}{3d} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int (Ba^3 + 3Aba^2 + 6b^2Ba + 2Ab^3 + 2b^3B \cos(c + dx)) \sec(c + dx) dx + \frac{2a(2a^2A + 9abB + 8Ab^2) \tan(c + dx)}{d} \right) \frac{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{Ba^3 + 3Aba^2 + 6b^2Ba + 2Ab^3 + 2b^3B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a(2a^2A + 9abB + 8Ab^2) \tan(c + dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^2}{3d} \right) \\ \downarrow \text{3214}$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left((a^3B + 3a^2Ab + 6ab^2B + 2Ab^3) \int \sec(c + dx) dx + 2b^3Bx \right) + \frac{2a(2a^2A + 9abB + 8Ab^2) \tan(c + dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^2}{3d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left((a^3B + 3a^2Ab + 6ab^2B + 2Ab^3) \int \csc(c + dx + \frac{\pi}{2}) dx + 2b^3Bx \right) + \frac{2a(2a^2A + 9abB + 8Ab^2) \tan(c + dx)}{d} \right) \right. \\ \left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^2}{3d} \right) \\ \downarrow \text{4257}$$

$$\frac{1}{3} \left(\frac{a^2(3aB + 5Ab) \tan(c + dx) \sec(c + dx)}{2d} + \frac{1}{2} \left(\frac{2a(2a^2A + 9abB + 8Ab^2) \tan(c + dx)}{d} + 3 \left(\frac{a^3B + 3a^2Ab + 6ab^2B + 2Ab^3}{d} \right) \right) \right. \\ \left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^2}{3d} \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*(a + b*Cos[c + d*x])^2*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((a^2*(5*A*b + 3*a*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (3*(2*b^3*B*x + ((3*a^2*A*b + 2*A*b^3 + a^3*B + 6*a*b^2*B)*ArcTanh[Sin[c + d*x]]))/d) + (2*a*(2*a^2*A + 8*A*b^2 + 9*a*b*B)*Tan[c + d*x])/d)/2)/3`

3.237.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]) / ((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

```
rule 3510 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

3.237.4 Maple [A] (verified)

Time = 4.88 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.01

method	result
parts	$-\frac{A a^3 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d} + \frac{(A b^3 + 3B a b^2) \ln(\sec(dx+c) + \tan(dx+c))}{d} + \frac{(3A a b^2 + 3B a^2 b) \tan(dx+c)}{d}$
derivativedivides	$-A a^3 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + B a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 3A a^2 b \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)$
default	$-A a^3 \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + B a^3 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 3A a^2 b \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)$
parallelrisch	$-\frac{27(A a^2 b + \frac{2}{3} A b^3 + \frac{1}{3} B a^3 + 2B a b^2) \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 27(A a^2 b + \frac{2}{3} A b^3 + \frac{1}{3} B a^3 + 2B a b^2)}{\dots}$
risch	$b^3 B x - \frac{ia(9Aab e^{5i(dx+c)} + 3B a^2 e^{5i(dx+c)} - 18A b^2 e^{4i(dx+c)} - 18B a b e^{4i(dx+c)} - 12A a^2 e^{2i(dx+c)} - 36A b^2 e^{2i(dx+c)})}{3d(e^{2i(dx+c)} + 1)^3}$
norman	$b^3 B x \left(\tan^{12} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + b^3 B x \left(\tan^{14} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - b^3 B x - \frac{8a(A a^2 - 3A b^2 - 3B a b) \left(\tan^7 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{d} - \frac{a(2A a^2 - 3A a b + 6A b^2 - B a^3)}{\dots}$

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBOSE)
```

3.237. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

output $-Aa^3/d*(-2/3-1/3*\sec(dx+c)^2)*\tan(dx+c)+(Ab^3+3Bab^2)/d*\ln(\sec(dx+c)+\tan(dx+c))+(3Aab^2+3Ba^2b)/d*\tan(dx+c)+(3Aa^2b+Ba^3)/d*(1/2*\sec(dx+c)*\tan(dx+c)+1/2*\ln(\sec(dx+c)+\tan(dx+c)))+Bb^3/d*(dx+c)$

3.237.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.30

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12 B b^3 dx \cos(dx + c)^3 + 3 (B a^3 + 3 A a^2 b + 6 B a b^2 + 2 A b^3) \cos(dx + c)^3 \log(\sin(dx + c) + 1) - 3 (B a^3}{$$

input `integrate((a+b*cos(dx+c))^3*(A+B*cos(dx+c))*sec(dx+c)^4,x, algorithm="fricas")`

output $1/12*(12*B*b^3*d*x*\cos(dx + c)^3 + 3*(B*a^3 + 3*A*a^2*b + 6*B*a*b^2 + 2*A*b^3)*\cos(dx + c)^3*\log(\sin(dx + c) + 1) - 3*(B*a^3 + 3*A*a^2*b + 6*B*a*b^2 + 2*A*b^3)*\cos(dx + c)^3*\log(-\sin(dx + c) + 1) + 2*(2*A*a^3 + 2*(2*A*a^3 + 9*B*a^2*b + 9*A*a*b^2)*\cos(dx + c)^2 + 3*(B*a^3 + 3*A*a^2*b)*\cos(dx + c))*\sin(dx + c))/(d*\cos(dx + c)^3)$

3.237.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(dx+c))**3*(A+B*cos(dx+c))*sec(dx+c)**4,x)`

output `Timed out`

3.237.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.49

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{4 (\tan(dx + c)^3 + 3 \tan(dx + c)) A a^3 + 12 (dx + c) B b^3 - 3 B a^3 \left(\frac{2 \sin(dx + c)}{\sin(dx + c)^2 - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) - 9 A a^2 b \left(\frac{2 \sin(dx + c)}{\sin(dx + c)^2 - 1} - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1) \right) + 18 B a b^2 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 6 A a b^3 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 36 B a^2 b \tan(dx + c) + 36 A a b^2 \tan(dx + c)}{d}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")
```

```
output 1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^3 + 12*(d*x + c)*B*b^3 - 3*B*a^3*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 9*A*a^2*b*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 18*B*a*b^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 6*A*a*b^3*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 36*B*a^2*b*tan(d*x + c) + 36*A*a*b^2*tan(d*x + c))/d
```

3.237.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 336 vs. 2(137) = 274.

Time = 0.32 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.32

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{6 (dx + c) B b^3 + 3 (B a^3 + 3 A a^2 b + 6 B a b^2 + 2 A b^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 3 (B a^3 + 3 A a^2 b + 6 B a b^2 + 2 A b^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + 6 A a b^3 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 36 B a^2 b \tan(dx + c) + 36 A a b^2 \tan(dx + c)}{d}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")
```

output $\frac{1}{6}(6*(d*x + c)*B*b^3 + 3*(B*a^3 + 3*A*a^2*b + 6*B*a*b^2 + 2*A*b^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(B*a^3 + 3*A*a^2*b + 6*B*a*b^2 + 2*A*b^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*A*a^3*\tan(1/2*d*x + 1/2*c)^5 - 3*B*a^3*\tan(1/2*d*x + 1/2*c)^5 - 9*A*a^2*b*\tan(1/2*d*x + 1/2*c)^5 + 18*B*a^2*b*\tan(1/2*d*x + 1/2*c)^5 + 18*A*a*b^2*\tan(1/2*d*x + 1/2*c)^5 - 4*A*a^3*\tan(1/2*d*x + 1/2*c)^3 - 36*B*a^2*b*\tan(1/2*d*x + 1/2*c)^3 - 36*A*a*b^2*\tan(1/2*d*x + 1/2*c)^3 + 6*A*a^3*\tan(1/2*d*x + 1/2*c) + 3*B*a^3*\tan(1/2*d*x + 1/2*c) + 9*A*a^2*b*\tan(1/2*d*x + 1/2*c) + 18*B*a^2*b*\tan(1/2*d*x + 1/2*c) + 18*A*a*b^2*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^3/d$

3.237.9 Mupad [B] (verification not implemented)

Time = 2.17 (sec) , antiderivative size = 526, normalized size of antiderivative = 3.63

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{Aa^3 \sin(3c+3dx)}{6} + \frac{Ba^3 \sin(2c+2dx)}{4} + \frac{Aa^3 \sin(c+dx)}{2} + \frac{3Aab^2 \sin(c+dx)}{4} + \frac{3Ba^2b \sin(c+dx)}{4} - \frac{Ab^3 \cos(c+dx) \operatorname{atan}\left(\frac{\sin(\frac{c}{2} + \frac{dx}{2})}{\cos(\frac{c}{2} + \frac{dx}{2})}\right)}{2}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^4,x)`

output $((A*a^3*\sin(3*c + 3*d*x))/6 + (B*a^3*\sin(2*c + 2*d*x))/4 + (A*a^3*\sin(c + d*x))/2 + (3*A*a*b^2*\sin(c + d*x))/4 + (3*B*a^2*b*\sin(c + d*x))/4 - (A*b^3*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*3i)/2 - (B*a^3*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*3i)/4 + (3*B*b^3*\cos(c + d*x)*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 + (3*A*a^2*b*\sin(2*c + 2*d*x))/4 + (3*A*a*b^2*\sin(3*c + 3*d*x))/4 + (3*B*a^2*b*\sin(3*c + 3*d*x))/4 - (A*b^3*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*1i)/2 - (B*a^3*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*1i)/4 + (B*b^3*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x))/2 - (A*a^2*b*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*3i)/4 - (B*a*b^2*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*3i)/2 - (A*a^2*b*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*9i)/4 - (B*a*b^2*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*9i)/2)/(d*((3*\cos(c + d*x))/4 + \cos(3*c + 3*d*x)/4))$

3.238 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.238.1 Optimal result	2188
3.238.2 Mathematica [A] (verified)	2189
3.238.3 Rubi [A] (verified)	2189
3.238.4 Maple [A] (verified)	2193
3.238.5 Fricas [A] (verification not implemented)	2194
3.238.6 Sympy [F(-1)]	2194
3.238.7 Maxima [A] (verification not implemented)	2195
3.238.8 Giac [B] (verification not implemented)	2195
3.238.9 Mupad [B] (verification not implemented)	2196

3.238.1 Optimal result

Integrand size = 31, antiderivative size = 188

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx \\ &= \frac{(3a^3 A + 12aAb^2 + 12a^2bB + 8b^3 B) \operatorname{arctanh}(\sin(c + dx))}{8d} \\ &+ \frac{(6a^2 Ab + 3Ab^3 + 2a^3 B + 9ab^2 B) \tan(c + dx)}{3d} \\ &+ \frac{a(3a^2 A + 10Ab^2 + 12abB) \sec(c + dx) \tan(c + dx)}{8d} \\ &+ \frac{a^2(3Ab + 2aB) \sec^2(c + dx) \tan(c + dx)}{6d} \\ &+ \frac{aA(a + b \cos(c + dx))^2 \sec^3(c + dx) \tan(c + dx)}{4d} \end{aligned}$$

```
output 1/8*(3*A*a^3+12*A*a*b^2+12*B*a^2*b+8*B*b^3)*arctanh(sin(d*x+c))/d+1/3*(6*A
*a^2*b+3*A*b^3+2*B*a^3+9*B*a*b^2)*tan(d*x+c)/d+1/8*a*(3*A*a^2+10*A*b^2+12*
B*a*b)*sec(d*x+c)*tan(d*x+c)/d+1/6*a^2*(3*A*b+2*B*a)*sec(d*x+c)^2*tan(d*x+
c)/d+1/4*a*A*(a+b*cos(d*x+c))^2*sec(d*x+c)^3*tan(d*x+c)/d
```

3.238.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.74

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3a^3A + 12aAb^2 + 12a^2bB + 8b^3B) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (24(3a^2Ab + Ab^3 + a^3B + 3ab^2A) + 9a^2(A + 4Ab^2 + 4a^2bB) \operatorname{Sec}[c + dx] + 6a^3A \operatorname{Sec}[c + dx]^3 + 8a^2(3Ab + aB) \operatorname{Tan}[c + dx]^2)}{24d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(3*(3*a^3*A + 12*a*A*b^2 + 12*a^2*b*B + 8*b^3*B)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(24*(3*a^2*A*b + A*b^3 + a^3*B + 3*a*b^2*B) + 9*a*(a^2*A + 4*A*b^2 + 4*a*b*B)*Sec[c + d*x] + 6*a^3*A*Sec[c + d*x]^3 + 8*a^2*(3*A*b + a*B)*Tan[c + d*x]^2))/(24*d)`

3.238.3 Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3468, 3042, 3510, 25, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{4} \int (a + b \cos(c + dx)) (b(aA + 4bB) \cos^2(c + dx) + (3Aa^2 + 8bBa + 4Ab^2) \cos(c + dx) + 2a(3Ab + 2aB)) \sec^4(c + dx) dx + \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^2}{4d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) \left(b(aA + 4bB) \sin(c + dx + \frac{\pi}{2})^2 + (3Aa^2 + 8bBa + 4Ab^2) \sin(c + dx + \frac{\pi}{2}) + 2a(3Aa^2 + 12bBa + 10Ab^2) \right)}{aA \tan(c + dx) \sec^3(c + dx) (a + b \cos(c + dx))^2} dx + \frac{\sin(c + dx + \frac{\pi}{2})^4}{4d}$$

↓ 3510

$$\frac{1}{4} \left(\frac{2a^2(2aB + 3Ab) \tan(c + dx) \sec^2(c + dx)}{3d} - \frac{1}{3} \int -((3b^2(aA + 4bB) \cos^2(c + dx) + 4(2Ba^3 + 6Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx) + 3a(3Aa^2 + 12bBa + 10Ab^2)) \sec^2(c + dx))}{aA \tan(c + dx) \sec^3(c + dx) (a + b \cos(c + dx))^2} dx \right) + \frac{\sin(c + dx + \frac{\pi}{2})^4}{4d}$$

↓ 25

$$\frac{1}{4} \left(\frac{1}{3} \int (3b^2(aA + 4bB) \cos^2(c + dx) + 4(2Ba^3 + 6Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx) + 3a(3Aa^2 + 12bBa + 10Ab^2)) \sec^2(c + dx)}{aA \tan(c + dx) \sec^3(c + dx) (a + b \cos(c + dx))^2} dx \right) + \frac{\sin(c + dx + \frac{\pi}{2})^4}{4d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \int \frac{3b^2(aA + 4bB) \sin(c + dx + \frac{\pi}{2})^2 + 4(2Ba^3 + 6Aba^2 + 9b^2Ba + 3Ab^3) \sin(c + dx + \frac{\pi}{2}) + 3a(3Aa^2 + 12bBa + 10Ab^2) \sin(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx) (a + b \cos(c + dx))^2} dx \right) + \frac{\sin(c + dx + \frac{\pi}{2})^3}{4d}$$

↓ 3500

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (8(2Ba^3 + 6Aba^2 + 9b^2Ba + 3Ab^3) + 3(3Aa^3 + 12bBa^2 + 12Ab^2a + 8b^3B) \cos(c + dx)) \sec^2(c + dx)}{aA \tan(c + dx) \sec^3(c + dx) (a + b \cos(c + dx))^2} dx \right) \right) + \frac{\sin(c + dx + \frac{\pi}{2})^4}{4d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{8(2Ba^3 + 6Aba^2 + 9b^2Ba + 3Ab^3) + 3(3Aa^3 + 12bBa^2 + 12Ab^2a + 8b^3B) \sin(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx) (a + b \cos(c + dx))^2} dx \right) \right) + \frac{3a(3Aa^2 + 12bBa + 10Ab^2) \sin(c + dx + \frac{\pi}{2})}{4d}$$

↓ 3227

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(8(2a^3B + 6a^2Ab + 9ab^2B + 3Ab^3) \int \sec^2(c + dx) dx + 3(3a^3A + 12a^2bB + 12aAb^2 + 8b^3B) \int \sec(c + dx) dx \right. \right. \right. \\ \left. \left. \left. \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^2}{4d} \right) \right) \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3(3a^3A + 12a^2bB + 12aAb^2 + 8b^3B) \int \csc\left(c + dx + \frac{\pi}{2}\right) dx + 8(2a^3B + 6a^2Ab + 9ab^2B + 3Ab^3) \int \sec(c + dx) dx \right. \right. \right. \\ \left. \left. \left. \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^2}{4d} \right) \right) \right. \\ \left. \downarrow 4254 \right.$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3(3a^3A + 12a^2bB + 12aAb^2 + 8b^3B) \int \csc\left(c + dx + \frac{\pi}{2}\right) dx - \frac{8(2a^3B + 6a^2Ab + 9ab^2B + 3Ab^3) \int \sec(c + dx) dx}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^2}{4d} \right) \right) \right. \\ \left. \downarrow 24 \right.$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3(3a^3A + 12a^2bB + 12aAb^2 + 8b^3B) \int \csc\left(c + dx + \frac{\pi}{2}\right) dx + \frac{8(2a^3B + 6a^2Ab + 9ab^2B + 3Ab^3) \tan(c + dx)}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^2}{4d} \right) \right) \right. \\ \left. \downarrow 4257 \right.$$

$$\frac{1}{4} \left(\frac{2a^2(2aB + 3Ab) \tan(c + dx) \sec^2(c + dx)}{3d} + \frac{1}{3} \left(\frac{3a(3a^2A + 12abB + 10Ab^2) \tan(c + dx) \sec(c + dx)}{2d} + \frac{1}{2} \left(\frac{3(3a^3A + 12a^2bB + 12aAb^2 + 8b^3B) \tan(c + dx)}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^2}{4d} \right) \right) \right.$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(a*A*(a + b*Cos[c + d*x])^2*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((2*a^2*(3*A*b + 2*a*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((3*a*(3*a^2*A + 10*A*b^2 + 12*a*b*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((3*(3*a^3*A + 12*a*A*b^2 + 12*a^2*b*B + 8*b^3*B)*ArcTanh[Sin[c + d*x]])/d + (8*(6*a^2*A*b + 3*A*b^3 + 2*a^3*B + 9*a*b^2*B)*Tan[c + d*x])/d)/2)/3)/4`

3.238.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2)), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

```
rule 3510 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

3.238.4 Maple [A] (verified)

Time = 5.53 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.98

method	result
parts	$\frac{A a^3 \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)}{d} + \frac{(A b^3 + 3 B a b^2) \tan(dx+c)}{d} + \frac{(3 A a^3 - 3 B a^2 b)}{d}$
derivativedivides	$\frac{A a^3 \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)}{d} - B a^3 \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)$
default	$\frac{A a^3 \left(- \left(- \frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{8} \right)}{d} - B a^3 \left(- \frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)$
parallelrisch	$-18(A a^3 + 4 A a b^2 + 4 B a^2 b + \frac{8}{3} B b^3) \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 18(A a^3 + 4 A a b^2 + 4 B a^2 b + \frac{8}{3} B b^3)$
risch	$\frac{i(36 A a b^2 e^{5i(dx+c)} + 36 B a^2 b e^{5i(dx+c)} - 72 B a b^2 e^{6i(dx+c)} - 36 A a b^2 e^{i(dx+c)} - 36 B a^2 b e^{i(dx+c)} + 36 A a b^2 e^{7i(dx+c)} + 36 B a^3)}{4d}$
norman	$\frac{(5 A a^3 - 24 A a^2 b + 12 A a b^2 - 8 A b^3 - 8 B a^3 + 12 B a^2 b - 24 B a b^2) \left(\tan^{15} \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{4d} + \frac{(5 A a^3 + 24 A a^2 b + 12 A a b^2 + 8 A b^3 + 8 B a^3 + 12 B a^2 b + 24 B a b^2 + 8 B b^3)}{4d}$

3.238. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

input `int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBOSE)`

output `A*a^3/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^3+3*B*a*b^2)/d*tan(d*x+c)+(3*A*a*b^2+3*B*a^2*b)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-(3*A*a^2*b+B*a^3)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+B*b^3/d*ln(sec(d*x+c)+tan(d*x+c))`

3.238.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.12

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3(3Aa^3 + 12Ba^2b + 12Aab^2 + 8Bb^3) \cos(dx + c)^4 \log(\sin(dx + c) + 1) - 3(3Aa^3 + 12Ba^2b + 12Aab^2 + 8Bb^3) \cos(dx + c)^4 \log(-\sin(dx + c) + 1) + 2(6Aa^3 + 8(2Ba^3 + 6Aa^2b + 9Bab^2 + 3Aa^2b^3) \cos(dx + c)^3 + 9(Aa^3 + 4Ba^2b + 4Aa^2b^2) \cos(dx + c)^2 + 8(Ba^3 + 3Aa^2b) \cos(dx + c)) \sin(dx + c)}{(d \cos(dx + c))^4}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")`

output `1/48*(3*(3*A*a^3 + 12*B*a^2*b + 12*A*a*b^2 + 8*B*b^3)*cos(d*x + c)^4*log(sin(d*x + c) + 1) - 3*(3*A*a^3 + 12*B*a^2*b + 12*A*a*b^2 + 8*B*b^3)*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*(6*A*a^3 + 8*(2*B*a^3 + 6*A*a^2*b + 9*B*a*b^2 + 3*A*b^3)*cos(d*x + c)^3 + 9*(A*a^3 + 4*B*a^2*b + 4*A*a^2*b^2)*cos(d*x + c)^2 + 8*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^4)`

3.238.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output `Timed out`

3.238. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$

3.238.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.45

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ba^3 + 48 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^2b - 3 Aa^3 \left(\frac{2 (3 \sin(dx+c)^3}{\sin(dx+c)^4 - 2s} \right)}{\sin(dx+c)^4 - 2s}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")
```

```
output 1/48*(16*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^3 + 48*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^2*b - 3*A*a^3*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 36*B*a^2*b*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 36*A*a*b^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 24*B*b^3*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 144*B*a*b^2*tan(d*x + c) + 48*A*b^3*tan(d*x + c))/d
```

3.238.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 586 vs. 2(178) = 356.

Time = 0.33 (sec) , antiderivative size = 586, normalized size of antiderivative = 3.12

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{3 (3 Aa^3 + 12 Ba^2b + 12 Aab^2 + 8 Bb^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 3 (3 Aa^3 + 12 Ba^2b + 12 Aab^2 + 8 Bb^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right)}{\sin(dx+c)^4 - 2s}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="giac")
```

output

$$\frac{1}{24} \cdot (3 \cdot (3Aa^3 + 12Ba^2b + 12Aab^2 + 8Bb^3) \cdot \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1)) - 3 \cdot (3Aa^3 + 12Ba^2b + 12Aab^2 + 8Bb^3) \cdot \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) + 2 \cdot (15Aa^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 24Ba^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 72Aa^2b \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 36Ba^2b \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 36Aab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 72Bab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 24Ab^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 9Aa^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 40Ba^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 120Aa^2b \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 36Ba^2b \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 36Aab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 216Bab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 72Ab^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 9Aa^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 40Ba^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 120Aa^2b \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 36Ba^2b \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 36Aab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 216Bab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 72Ab^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 15Aa^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 24Ba^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 72Aa^2b \tan(\frac{1}{2}dx + \frac{1}{2}c) + 36Ba^2b \tan(\frac{1}{2}dx + \frac{1}{2}c) + 36Aab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 72Bab^2 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 24Ab^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)) / (\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1)^4 / d$$

3.238.9 Mupad [B] (verification not implemented)

Time = 4.13 (sec) , antiderivative size = 395, normalized size of antiderivative = 2.10

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3Aa^3}{8} + \frac{3Ba^2b}{2} + \frac{3Aab^2}{2} + Bb^3\right)}{\frac{3Aa^3}{2} + 6Ba^2b + 6Aab^2 + 4Bb^3}\right) \left(\frac{3Aa^3}{4} + 3Ba^2b + 3Aab^2 + 2Bb^3\right)}{d} \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^7 + \left(3Aab^2 - 6Ab^3 - \left(2Ab^3 - \frac{5Aa^3}{4} + 2Ba^3 - 3Aab^2 + 6Aa^2b + 6Bab^2 - 3Ba^2b\right)\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^5,x)`

output $(\operatorname{atanh}((4*\tan(c/2 + (d*x)/2)*((3*A*a^3)/8 + B*b^3 + (3*A*a*b^2)/2 + (3*B*a^2*b)/2)))/((3*A*a^3)/2 + 4*B*b^3 + 6*A*a*b^2 + 6*B*a^2*b))*((3*A*a^3)/4 + 2*B*b^3 + 3*A*a*b^2 + 3*B*a^2*b))/d - (\tan(c/2 + (d*x)/2)^7*(2*A*b^3 - (5*A*a^3)/4 + 2*B*a^3 - 3*A*a*b^2 + 6*A*a^2*b + 6*B*a*b^2 - 3*B*a^2*b) + \tan(c/2 + (d*x)/2)^3*(6*A*b^3 - (3*A*a^3)/4 + (10*B*a^3)/3 + 3*A*a*b^2 + 10*A*a^2*b + 18*B*a*b^2 + 3*B*a^2*b) - \tan(c/2 + (d*x)/2)^5*((3*A*a^3)/4 + 6*A*b^3 + (10*B*a^3)/3 - 3*A*a*b^2 + 10*A*a^2*b + 18*B*a*b^2 - 3*B*a^2*b) - \tan(c/2 + (d*x)/2)*((5*A*a^3)/4 + 2*A*b^3 + 2*B*a^3 + 3*A*a*b^2 + 6*A*a^2*b + 6*B*a*b^2 + 3*B*a^2*b))/(d*(6*\tan(c/2 + (d*x)/2)^4 - 4*\tan(c/2 + (d*x)/2)^2 - 4*\tan(c/2 + (d*x)/2)^6 + \tan(c/2 + (d*x)/2)^8 + 1))$

3.239 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^6(c+dx) dx$

3.239.1 Optimal result	2198
3.239.2 Mathematica [A] (verified)	2199
3.239.3 Rubi [A] (verified)	2199
3.239.4 Maple [A] (verified)	2204
3.239.5 Fricas [A] (verification not implemented)	2205
3.239.6 Sympy [F(-1)]	2205
3.239.7 Maxima [A] (verification not implemented)	2206
3.239.8 Giac [B] (verification not implemented)	2206
3.239.9 Mupad [B] (verification not implemented)	2207

3.239.1 Optimal result

Integrand size = 31, antiderivative size = 236

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx \\ &= \frac{(9a^2Ab + 4Ab^3 + 3a^3B + 12ab^2B) \operatorname{arctanh}(\sin(c + dx))}{8d} \\ &+ \frac{(8a^3A + 30aAb^2 + 30a^2bB + 15b^3B) \tan(c + dx)}{15d} \\ &+ \frac{(9a^2Ab + 4Ab^3 + 3a^3B + 12ab^2B) \sec(c + dx) \tan(c + dx)}{8d} \\ &+ \frac{a(4a^2A + 12Ab^2 + 15abB) \sec^2(c + dx) \tan(c + dx)}{15d} \\ &+ \frac{a^2(7Ab + 5aB) \sec^3(c + dx) \tan(c + dx)}{20d} \\ &+ \frac{aA(a + b \cos(c + dx))^2 \sec^4(c + dx) \tan(c + dx)}{5d} \end{aligned}$$

```
output 1/8*(9*A*a^2*b+4*A*b^3+3*B*a^3+12*B*a*b^2)*arctanh(sin(d*x+c))/d+1/15*(8*A
*a^3+30*A*a*b^2+30*B*a^2*b+15*B*b^3)*tan(d*x+c)/d+1/8*(9*A*a^2*b+4*A*b^3+
3*B*a^3+12*B*a*b^2)*sec(d*x+c)*tan(d*x+c)/d+1/15*a*(4*A*a^2+12*A*b^2+15*B*a
*b)*sec(d*x+c)^2*tan(d*x+c)/d+1/20*a^2*(7*A*b+5*B*a)*sec(d*x+c)^3*tan(d*x+
c)/d+1/5*a*A*(a+b*cos(d*x+c))^2*sec(d*x+c)^4*tan(d*x+c)/d
```


$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15(3a^3B + 9a^2Ab + 12ab^2B + 4Ab^3) \int \sec^3(c + dx) dx + 4(8a^3A + 30a^2bB + 30aAb^2 + 15b^3B) \int \sec^2(c + dx) dx \right) \right) \right) \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(4(8a^3A + 30a^2bB + 30aAb^2 + 15b^3B) \int \csc\left(c + dx + \frac{\pi}{2}\right)^2 dx + 15(3a^3B + 9a^2Ab + 12ab^2B + 4Ab^3) \int \sec^2(c + dx) dx \right) \right) \right) \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d}$$

↓ 4254

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15(3a^3B + 9a^2Ab + 12ab^2B + 4Ab^3) \int \csc\left(c + dx + \frac{\pi}{2}\right)^3 dx - \frac{4(8a^3A + 30a^2bB + 30aAb^2 + 15b^3B)}{d} \int \sec^2(c + dx) dx \right) \right) \right) \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d}$$

↓ 24

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15(3a^3B + 9a^2Ab + 12ab^2B + 4Ab^3) \int \csc\left(c + dx + \frac{\pi}{2}\right)^3 dx + \frac{4(8a^3A + 30a^2bB + 30aAb^2 + 15b^3B)}{d} \int \sec^2(c + dx) dx \right) \right) \right) \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d}$$

↓ 4255

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15(3a^3B + 9a^2Ab + 12ab^2B + 4Ab^3) \left(\frac{1}{2} \int \sec(c + dx) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4(8a^3A + 30a^2bB + 30aAb^2 + 15b^3B)}{d} \int \sec^2(c + dx) dx \right) \right) \right) \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(15(3a^3B + 9a^2Ab + 12ab^2B + 4Ab^3) \left(\frac{1}{2} \int \csc\left(c + dx + \frac{\pi}{2}\right) dx + \frac{\tan(c + dx) \sec(c + dx)}{2d} \right) + \frac{4(8a^3A + 30a^2bB + 30aAb^2 + 15b^3B)}{d} \int \sec^2(c + dx) dx \right) \right) \right) \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d}$$

↓ 4257

$$\frac{1}{5} \left(\frac{a^2(5aB + 7Ab) \tan(c + dx) \sec^3(c + dx)}{4d} + \frac{1}{4} \left(\frac{4a(4a^2A + 15abB + 12Ab^2) \tan(c + dx) \sec^2(c + dx)}{3d} + \frac{1}{3} \left(15 \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^2}{5d} \right) \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a*A*(a + b*Cos[c + d*x])^2*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) + ((a^2*(7*A*b + 5*a*B)*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((4*a*(4*a^2*A + 12*A*b^2 + 15*a*b*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((4*(8*a^3*A + 30*a*A*b^2 + 30*a^2*b*B + 15*b^3*B)*Tan[c + d*x])/d + 15*(9*a^2*A*b + 4*A*b^3 + 3*a^3*B + 12*a*b^2*B)*(ArcTanh[Sin[c + d*x])/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d)))/3)/4)/5`

3.239.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.239.4 Maple [A] (verified)

Time = 6.20 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.85

method	result
parts	$-\frac{A a^3 \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c)}{d} + \frac{(A b^3 + 3B a b^2) \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right)}{d}$
derivativedivides	$-A a^3 \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c) + B a^3 \left(-\left(-\frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{2} \right)$
default	$-A a^3 \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c) + B a^3 \left(-\left(-\frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c) + \tan(dx+c))}{2} \right)$
parallelrisch	$-135(A a^2 b + \frac{4}{9} A b^3 + \frac{1}{3} B a^3 + \frac{4}{3} B a b^2) (\cos(5dx+5c) + 5 \cos(3dx+3c) + 10 \cos(dx+c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 135(A a^2 b + \frac{4}{9} A b^3 + \frac{1}{3} B a^3 + \frac{4}{3} B a b^2)$
risch	$-\frac{i(-120 B b^3 - 64 A a^3 - 240 A a b^2 - 240 B a^2 b - 135 A a^2 b e^{i(dx+c)} - 180 B a b^2 e^{i(dx+c)} - 720 A a b^2 e^{6i(dx+c)} - 720 B a^2 b e^{6i(dx+c)})}{d}$

input `int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x,method=_RETURNVERBOSE)`

output `-A*a^3/d*(-8/15-1/5*sec(d*x+c)^4-4/15*sec(d*x+c)^2)*tan(d*x+c)+(A*b^3+3*B*a*b^2)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-(3*A*a*b^2+3*B*a^2*b)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(3*A*a^2*b+B*a^3)/d*(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c))+B*b^3/d*tan(d*x+c)`

3.239.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.06

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{15(3Ba^3 + 9Aa^2b + 12Bab^2 + 4Ab^3) \cos(dx + c)^5 \log(\sin(dx + c) + 1) - 15(3Ba^3 + 9Aa^2b + 12Bab^2 + 4Ab^3) \cos(dx + c)^5 \log(-\sin(dx + c) + 1) + 2(8(8Aa^3 + 30B a^2 b + 30A a b^2 + 15B b^3) \cos(dx + c)^4 + 24Aa^3 + 15(3B a^3 + 9A a^2 b + 12B a b^2 + 4A b^3) \cos(dx + c)^3 + 8(4A a^3 + 15B a^2 b + 15A a b^2) \cos(dx + c)^2 + 30(B a^3 + 3A a^2 b) \cos(dx + c)) \sin(dx + c)}{(d \cos(dx + c))^5}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="fricas")
```

```
output 1/240*(15*(3*B*a^3 + 9*A*a^2*b + 12*B*a*b^2 + 4*A*b^3)*cos(d*x + c)^5*log(sin(d*x + c) + 1) - 15*(3*B*a^3 + 9*A*a^2*b + 12*B*a*b^2 + 4*A*b^3)*cos(d*x + c)^5*log(-sin(d*x + c) + 1) + 2*(8*(8*A*a^3 + 30*B*a^2*b + 30*A*a*b^2 + 15*B*b^3)*cos(d*x + c)^4 + 24*A*a^3 + 15*(3*B*a^3 + 9*A*a^2*b + 12*B*a*b^2 + 4*A*b^3)*cos(d*x + c)^3 + 8*(4*A*a^3 + 15*B*a^2*b + 15*A*a*b^2)*cos(d*x + c)^2 + 30*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)
```

3.239.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**6,x)
```

```
output Timed out
```

3.239.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.44

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{16 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c)) A a^3 + 240 (\tan(dx + c)^3 + 3 \tan(dx + c)) B a^2 b}{d}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="maxima")`

output `1/240*(16*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*A*a^3 + 240*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^2*b + 240*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a*b^2 - 15*B*a^3*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 45*A*a^2*b*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 180*B*a*b^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 60*A*b^3*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 240*B*b^3*tan(d*x + c))/d`

3.239.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. 2(224) = 448.

Time = 0.36 (sec) , antiderivative size = 722, normalized size of antiderivative = 3.06

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Too large to display}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="giac")`

output

```

1/120*(15*(3*B*a^3 + 9*A*a^2*b + 12*B*a*b^2 + 4*A*b^3)*log(abs(tan(1/2*d*x
+ 1/2*c) + 1)) - 15*(3*B*a^3 + 9*A*a^2*b + 12*B*a*b^2 + 4*A*b^3)*log(abs(
tan(1/2*d*x + 1/2*c) - 1)) - 2*(120*A*a^3*tan(1/2*d*x + 1/2*c)^9 - 75*B*a^
3*tan(1/2*d*x + 1/2*c)^9 - 225*A*a^2*b*tan(1/2*d*x + 1/2*c)^9 + 360*B*a^2*
b*tan(1/2*d*x + 1/2*c)^9 + 360*A*a*b^2*tan(1/2*d*x + 1/2*c)^9 - 180*B*a*b^
2*tan(1/2*d*x + 1/2*c)^9 - 60*A*b^3*tan(1/2*d*x + 1/2*c)^9 + 120*B*b^3*tan
(1/2*d*x + 1/2*c)^9 - 160*A*a^3*tan(1/2*d*x + 1/2*c)^7 + 30*B*a^3*tan(1/2*
d*x + 1/2*c)^7 + 90*A*a^2*b*tan(1/2*d*x + 1/2*c)^7 - 960*B*a^2*b*tan(1/2*d
*x + 1/2*c)^7 - 960*A*a*b^2*tan(1/2*d*x + 1/2*c)^7 + 360*B*a*b^2*tan(1/2*d
*x + 1/2*c)^7 + 120*A*b^3*tan(1/2*d*x + 1/2*c)^7 - 480*B*b^3*tan(1/2*d*x +
1/2*c)^7 + 464*A*a^3*tan(1/2*d*x + 1/2*c)^5 + 1200*B*a^2*b*tan(1/2*d*x +
1/2*c)^5 + 1200*A*a*b^2*tan(1/2*d*x + 1/2*c)^5 + 720*B*b^3*tan(1/2*d*x + 1
/2*c)^5 - 160*A*a^3*tan(1/2*d*x + 1/2*c)^3 - 30*B*a^3*tan(1/2*d*x + 1/2*c)
^3 - 90*A*a^2*b*tan(1/2*d*x + 1/2*c)^3 - 960*B*a^2*b*tan(1/2*d*x + 1/2*c)^
3 - 960*A*a*b^2*tan(1/2*d*x + 1/2*c)^3 - 360*B*a*b^2*tan(1/2*d*x + 1/2*c)^
3 - 120*A*b^3*tan(1/2*d*x + 1/2*c)^3 - 480*B*b^3*tan(1/2*d*x + 1/2*c)^3 +
120*A*a^3*tan(1/2*d*x + 1/2*c) + 75*B*a^3*tan(1/2*d*x + 1/2*c) + 225*A*a^2
*b*tan(1/2*d*x + 1/2*c) + 360*B*a^2*b*tan(1/2*d*x + 1/2*c) + 360*A*a*b^2*t
an(1/2*d*x + 1/2*c) + 180*B*a*b^2*tan(1/2*d*x + 1/2*c) + 60*A*b^3*tan(1/2*
d*x + 1/2*c) + 120*B*b^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 ...

```

3.239.9 Mupad [B] (verification not implemented)

Time = 4.44 (sec) , antiderivative size = 470, normalized size of antiderivative = 1.99

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3Ba^3}{8} + \frac{9Aa^2b}{8} + \frac{3Ba^2b^2}{2} + \frac{Ab^3}{2}\right)}{\frac{3Ba^3}{2} + \frac{9Aa^2b}{2} + 6Ba^2b^2 + 2Ab^3}\right) \left(\frac{3Ba^3}{4} + \frac{9Aa^2b}{4} + 3Ba^2b^2 + Ab^3\right)}{d}$$

$$- \frac{\left(2Aa^3 - Ab^3 - \frac{5Ba^3}{4} + 2Bb^3 + 6Aa^2b^2 - \frac{15Aa^2b}{4} - 3Ba^2b^2 + 6Ba^2b\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^9 + \left(2Ab^3 - 8Aa^2b^2 + 4Aa^2b^2 - 4Aa^2b^2 + 4Aa^2b^2\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^8}{d}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^6,x)`

output $(\operatorname{atanh}((4*\tan(c/2 + (d*x)/2)*((A*b^3)/2 + (3*B*a^3)/8 + (9*A*a^2*b)/8 + (3*B*a*b^2)/2)))/(2*A*b^3 + (3*B*a^3)/2 + (9*A*a^2*b)/2 + 6*B*a*b^2))*(A*b^3 + (3*B*a^3)/4 + (9*A*a^2*b)/4 + 3*B*a*b^2))/d - (\tan(c/2 + (d*x)/2)*(2*A*a^3 + A*b^3 + (5*B*a^3)/4 + 2*B*b^3 + 6*A*a*b^2 + (15*A*a^2*b)/4 + 3*B*a*b^2 + 6*B*a^2*b) + \tan(c/2 + (d*x)/2)^5*((116*A*a^3)/15 + 12*B*b^3 + 20*A*a*b^2 + 20*B*a^2*b) + \tan(c/2 + (d*x)/2)^9*(2*A*a^3 - A*b^3 - (5*B*a^3)/4 + 2*B*b^3 + 6*A*a*b^2 - (15*A*a^2*b)/4 - 3*B*a*b^2 + 6*B*a^2*b) - \tan(c/2 + (d*x)/2)^3*((8*A*a^3)/3 + 2*A*b^3 + (B*a^3)/2 + 8*B*b^3 + 16*A*a*b^2 + (3*A*a^2*b)/2 + 6*B*a*b^2 + 16*B*a^2*b) - \tan(c/2 + (d*x)/2)^7*((8*A*a^3)/3 - 2*A*b^3 - (B*a^3)/2 + 8*B*b^3 + 16*A*a*b^2 - (3*A*a^2*b)/2 - 6*B*a*b^2 + 16*B*a^2*b))/(d*(5*\tan(c/2 + (d*x)/2)^2 - 10*\tan(c/2 + (d*x)/2)^4 + 10*\tan(c/2 + (d*x)/2)^6 - 5*\tan(c/2 + (d*x)/2)^8 + \tan(c/2 + (d*x)/2)^{10} - 1))$

3.240 $\int \cos^2(c+dx)(a+b \cos(c+dx))^4(A+B \cos(c+dx)) dx$

3.240.1 Optimal result	2209
3.240.2 Mathematica [A] (verified)	2210
3.240.3 Rubi [A] (verified)	2211
3.240.4 Maple [A] (verified)	2216
3.240.5 Fricas [A] (verification not implemented)	2216
3.240.6 Sympy [B] (verification not implemented)	2217
3.240.7 Maxima [A] (verification not implemented)	2218
3.240.8 Giac [A] (verification not implemented)	2219
3.240.9 Mupad [B] (verification not implemented)	2220

3.240.1 Optimal result

Integrand size = 31, antiderivative size = 366

$$\begin{aligned}
 & \int \cos^2(c+dx)(a+b \cos(c+dx))^4(A+B \cos(c+dx)) dx \\
 &= \frac{1}{16}(8a^4A+36a^2Ab^2+5Ab^4+24a^3bB+20ab^3B)x \\
 &+ \frac{(140a^3Ab+112aAb^3+35a^4B+168a^2b^2B+24b^4B)\sin(c+dx)}{35d} \\
 &+ \frac{(8a^4A+36a^2Ab^2+5Ab^4+24a^3bB+20ab^3B)\cos(c+dx)\sin(c+dx)}{16d} \\
 &+ \frac{b(224a^2Ab+35Ab^3+104a^3B+140ab^2B)\cos^3(c+dx)\sin(c+dx)}{168d} \\
 &+ \frac{b^2(49aAb+31a^2B+18b^2B)\cos^4(c+dx)\sin(c+dx)}{105d} \\
 &+ \frac{b(7Ab+10aB)\cos^3(c+dx)(a+b \cos(c+dx))^2\sin(c+dx)}{42d} \\
 &+ \frac{bB \cos^3(c+dx)(a+b \cos(c+dx))^3\sin(c+dx)}{7d} \\
 &- \frac{(140a^3Ab+112aAb^3+35a^4B+168a^2b^2B+24b^4B)\sin^3(c+dx)}{105d}
 \end{aligned}$$

output $\frac{1}{16}(8Aa^4+36Aa^2b^2+5Ab^4+24B^3b+20B^2ab^3)x+\frac{1}{35}(140Aa^3b+112Aa^2b^3+35B^4+168B^2b^2+24B^3b^4)\sin(dx+c)/d+\frac{1}{16}(8Aa^4+36Aa^2b^2+5Ab^4+24B^3b+20B^2ab^3)\cos(dx+c)\sin(dx+c)/d+\frac{1}{16}8b(224Aa^2b+35Ab^3+104B^3+140B^2ab^2)\cos(dx+c)^3\sin(dx+c)/d+\frac{1}{105}b^2(49Aa^2b+31B^2+18B^2b^2)\cos(dx+c)^4\sin(dx+c)/d+\frac{1}{42}b(7Ab+10B^2)\cos(dx+c)^3(a+b\cos(dx+c))^2\sin(dx+c)/d+\frac{1}{7}bB\cos(dx+c)^3(a+b\cos(dx+c))^3\sin(dx+c)/d-\frac{1}{105}(140Aa^3b+112Aa^2b^3+35B^4+168B^2b^2+24B^3b^4)\sin(dx+c)^3/d$

3.240.2 Mathematica [A] (verified)

Time = 1.89 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.11

$$\int \cos^2(c+dx)(a+b\cos(c+dx))^4(A+B\cos(c+dx))dx$$

$$= \frac{3360a^4Ac + 15120a^2Ab^2c + 2100Ab^4c + 10080a^3bBc + 8400ab^3Bc + 3360a^4Adx + 15120a^2Ab^2dx + 2100Ab^4dx + 10080a^3bBdx + 8400ab^3Bdx + 105(192a^3Ab + 160a^2Ab^3 + 48a^4B + 240a^2b^2B + 35b^4B)\sin[c+dx] + 105(16a^4A + 96a^2Ab^2 + 15Ab^4 + 64a^3bB + 60ab^3B)\sin[2(c+dx)] + 2240a^3Ab\sin[3(c+dx)] + 2800a^2Ab^3\sin[3(c+dx)] + 560a^4B\sin[3(c+dx)] + 4200a^2b^2B\sin[3(c+dx)] + 735b^4B\sin[3(c+dx)] + 1260a^2Ab^2\sin[4(c+dx)] + 315Ab^4\sin[4(c+dx)] + 840a^3bB\sin[4(c+dx)] + 1260ab^3B\sin[4(c+dx)] + 336a^2Ab^3\sin[5(c+dx)] + 504a^2b^2B\sin[5(c+dx)] + 147b^4B\sin[5(c+dx)] + 35Ab^4\sin[6(c+dx)] + 140ab^3B\sin[6(c+dx)] + 15b^4B\sin[7(c+dx)]}{6720d}$$

input `Integrate[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output $(3360a^4Ac + 15120a^2Ab^2c + 2100Ab^4c + 10080a^3bBc + 8400a^2b^3Bc + 3360a^4Adx + 15120a^2Ab^2dx + 2100Ab^4dx + 10080a^3bBdx + 8400ab^3Bdx + 105(192a^3Ab + 160a^2Ab^3 + 48a^4B + 240a^2b^2B + 35b^4B)\sin[c+dx] + 105(16a^4A + 96a^2Ab^2 + 15Ab^4 + 64a^3bB + 60ab^3B)\sin[2(c+dx)] + 2240a^3Ab\sin[3(c+dx)] + 2800a^2Ab^3\sin[3(c+dx)] + 560a^4B\sin[3(c+dx)] + 4200a^2b^2B\sin[3(c+dx)] + 735b^4B\sin[3(c+dx)] + 1260a^2Ab^2\sin[4(c+dx)] + 315Ab^4\sin[4(c+dx)] + 840a^3bB\sin[4(c+dx)] + 1260ab^3B\sin[4(c+dx)] + 336a^2Ab^3\sin[5(c+dx)] + 504a^2b^2B\sin[5(c+dx)] + 147b^4B\sin[5(c+dx)] + 35Ab^4\sin[6(c+dx)] + 140ab^3B\sin[6(c+dx)] + 15b^4B\sin[7(c+dx)])/(6720d)$

3.240.3 Rubi [A] (verified)

Time = 1.81 (sec) , antiderivative size = 315, normalized size of antiderivative = 0.86, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3469, 3042, 3528, 3042, 3512, 3042, 3502, 27, 3042, 3227, 3042, 3113, 2009, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(c+dx)(a+b\cos(c+dx))^4(A+B\cos(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c+dx+\frac{\pi}{2}\right)^2\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^4\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx$$

$$\downarrow \text{3469}$$

$$\frac{1}{7}\int \cos^2(c+dx)(a+b\cos(c+dx))^2(b(7Ab+10aB)\cos^2(c+dx)+(6Bb^2+7a(2Ab+aB))\cos(c+dx)+a(7aA+3bB))dx + \frac{bB\sin(c+dx)\cos^3(c+dx)(a+b\cos(c+dx))^3}{7d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{7}\int \sin\left(c+dx+\frac{\pi}{2}\right)^2\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^2\left(b(7Ab+10aB)\sin\left(c+dx+\frac{\pi}{2}\right)^2+(6Bb^2+7a(2Ab+aB))\right)dx + \frac{bB\sin(c+dx)\cos^3(c+dx)(a+b\cos(c+dx))^3}{7d}$$

$$\downarrow \text{3528}$$

$$\frac{1}{7}\left(\frac{1}{6}\int \cos^2(c+dx)(a+b\cos(c+dx))(2b(31Ba^2+49Aba+18b^2B)\cos^2(c+dx)+(42Ba^3+126Aba^2+104b^2))dx + \frac{bB\sin(c+dx)\cos^3(c+dx)(a+b\cos(c+dx))^3}{7d}\right)$$

$$\downarrow \text{3042}$$

$$\frac{1}{7}\left(\frac{1}{6}\int \sin\left(c+dx+\frac{\pi}{2}\right)^2\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)\left(2b(31Ba^2+49Aba+18b^2B)\sin\left(c+dx+\frac{\pi}{2}\right)^2+(42Ba^3+126Aba^2+104b^2)\right)dx + \frac{bB\sin(c+dx)\cos^3(c+dx)(a+b\cos(c+dx))^3}{7d}\right)$$

$$\downarrow \text{3512}$$

3.240. $\int \cos^2(c+dx)(a+b\cos(c+dx))^4(A+B\cos(c+dx))dx$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \int \cos^2(c+dx) (15(14Aa^2 + 16bBa + 7Ab^2) a^2 + 5b(104Ba^3 + 224Aba^2 + 140b^2Ba + 35Ab^3) \cos^2(c+dx) + \frac{bB \sin(c+dx) \cos^3(c+dx)(a+b \cos(c+dx))^3}{7d} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \int \sin \left(c+dx + \frac{\pi}{2} \right)^2 \left(15(14Aa^2 + 16bBa + 7Ab^2) a^2 + 5b(104Ba^3 + 224Aba^2 + 140b^2Ba + 35Ab^3) \sin \left(c+dx + \frac{\pi}{2} \right) + \frac{bB \sin(c+dx) \cos^3(c+dx)(a+b \cos(c+dx))^3}{7d} \right) \right)$$

↓ 3502

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \int 3 \cos^2(c+dx) (35(8Aa^4 + 24bBa^3 + 36Ab^2a^2 + 20b^3Ba + 5Ab^4) + 8(35Ba^4 + 140Aba^3 + 168b^2Ba + 35Ab^4) + \frac{bB \sin(c+dx) \cos^3(c+dx)(a+b \cos(c+dx))^3}{7d} \right) \right) \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int \cos^2(c+dx) (35(8Aa^4 + 24bBa^3 + 36Ab^2a^2 + 20b^3Ba + 5Ab^4) + 8(35Ba^4 + 140Aba^3 + 168b^2Ba + 35Ab^4) + \frac{bB \sin(c+dx) \cos^3(c+dx)(a+b \cos(c+dx))^3}{7d} \right) \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \int \sin \left(c+dx + \frac{\pi}{2} \right)^2 (35(8Aa^4 + 24bBa^3 + 36Ab^2a^2 + 20b^3Ba + 5Ab^4) + 8(35Ba^4 + 140Aba^3 + 168b^2Ba + 35Ab^4) + \frac{bB \sin(c+dx) \cos^3(c+dx)(a+b \cos(c+dx))^3}{7d} \right) \right) \right)$$

↓ 3227

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(8(35a^4B + 140a^3Ab + 168a^2b^2B + 112aAb^3 + 24b^4B) \int \cos^3(c+dx) dx + 35(8a^4A + 24a^3bB + 35a^2b^2A + 20ab^3B + 5b^4A) + \frac{bB \sin(c+dx) \cos^3(c+dx)(a+b \cos(c+dx))^3}{7d} \right) \right) \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(35(8a^4A + 24a^3bB + 36a^2Ab^2 + 20ab^3B + 5Ab^4) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx + 8(35a^4B + 140a^3Ab + \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^3}{7d} \right) \right) \right) \right) \downarrow \text{3113}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(35(8a^4A + 24a^3bB + 36a^2Ab^2 + 20ab^3B + 5Ab^4) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{8(35a^4B + 140a^3Ab + \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^3}{7d} \right) \right) \right) \right) \downarrow \text{2009}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(35(8a^4A + 24a^3bB + 36a^2Ab^2 + 20ab^3B + 5Ab^4) \int \sin \left(c + dx + \frac{\pi}{2} \right)^2 dx - \frac{8(35a^4B + 140a^3Ab + \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^3}{7d} \right) \right) \right) \right) \downarrow \text{3115}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{1}{5} \left(\frac{3}{4} \left(35(8a^4A + 24a^3bB + 36a^2Ab^2 + 20ab^3B + 5Ab^4) \left(\frac{\int 1 dx}{2} + \frac{\sin(c + dx) \cos(c + dx)}{2d} \right) - \frac{8(35a^4B + \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^3}{7d} \right) \right) \right) \right) \downarrow \text{24}$$

$$\frac{1}{7} \left(\frac{1}{6} \left(\frac{2b^2(31a^2B + 49aAb + 18b^2B) \sin(c + dx) \cos^4(c + dx)}{5d} + \frac{1}{5} \left(\frac{5b(104a^3B + 224a^2Ab + 140ab^2B + 35Ab^3)}{4d} - \frac{bB \sin(c + dx) \cos^3(c + dx)(a + b \cos(c + dx))^3}{7d} \right) \right) \right)$$

input `Int[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

```
output (b*B*cos[c + d*x]^3*(a + b*cos[c + d*x])^3*sin[c + d*x]/(7*d) + ((b*(7*A*
b + 10*A*B)*cos[c + d*x]^3*(a + b*cos[c + d*x])^2*sin[c + d*x]/(6*d) + ((
2*b^2*(49*A*A*b + 31*a^2*B + 18*b^2*B)*cos[c + d*x]^4*sin[c + d*x]/(5*d)
+ ((5*b*(224*a^2*A*b + 35*A*b^3 + 104*a^3*B + 140*a*b^2*B)*cos[c + d*x]^3*
sin[c + d*x]/(4*d) + (3*(35*(8*a^4*A + 36*a^2*A*b^2 + 5*A*b^4 + 24*a^3*b*
B + 20*a*b^3*B)*(x/2 + (cos[c + d*x]*sin[c + d*x])/(2*d)) - (8*(140*a^3*A*
b + 112*A*A*b^3 + 35*a^4*B + 168*a^2*b^2*B + 24*b^4*B)*(-sin[c + d*x] + Si
n[c + d*x]^3/3)/d))/4)/5)/6)/7
```

3.240.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 27 Int[(a_)*(F_), x_Symbol] := Simp[a Int[F, x], x] /; FreeQ[a, x] && !Ma
tchQ[F, (b_)*(G_)] /; FreeQ[b, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3113 Int[sin[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
and[(1 - x^2)^(n - 1)/2], x], x], x, Cos[c + d*x], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

```
rule 3115 Int[((b_)*sin[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*sin
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)]^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3512 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Sin[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) + A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

3.240.4 Maple [A] (verified)

Time = 6.26 (sec) , antiderivative size = 273, normalized size of antiderivative = 0.75

method	result
parts	$(Ab^4+4Bab^3) \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15\cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{(4Aab^3+6Ba^2b^2) \left(\frac{8}{3} + \cos^4(dx+c) \right)}{5d}$
parallelrisch	$\frac{(1680a^4A+10080Aa^2b^2+1575Aa^4+6720Ba^3b+6300Ba^2b^3) \sin(2dx+2c) + (2240Aa^3b+2800Aa^2b^2+560Ba^4+4200Ba^3b) \sin(dx+c)}{1680a^4d + 10080Aa^2b^2d + 1575Aa^4d + 6720Ba^3bd + 6300Ba^2b^3d}$
derivativedivides	$\frac{Bb^4 \left(\frac{16}{5} + \cos^6(dx+c) + \frac{6(\cos^4(dx+c))}{5} + \frac{8(\cos^2(dx+c))}{5} \right) \sin(dx+c)}{7} + Ab^4 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15\cos(dx+c)}{8} \right) \sin(dx+c)}{6} \right)$
default	$\frac{Bb^4 \left(\frac{16}{5} + \cos^6(dx+c) + \frac{6(\cos^4(dx+c))}{5} + \frac{8(\cos^2(dx+c))}{5} \right) \sin(dx+c)}{7} + Ab^4 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15\cos(dx+c)}{8} \right) \sin(dx+c)}{6} \right)$
risch	$\frac{7 \sin(5dx+5c)Bb^4}{320d} + \frac{3 \sin(4dx+4c)Ab^4}{64d} + \frac{7 \sin(3dx+3c)Bb^4}{64d} + \frac{15 \sin(2dx+2c)Bab^3}{16d} + \frac{15 \sin(dx+c)Ba^2b^2}{4d} + \dots$
norman	Expression too large to display

input `int(cos(d*x+c)^2*(a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `(A*b^4+4*B*a*b^3)/d*(1/6*(cos(d*x+c)^5+5/4*cos(d*x+c)^3+15/8*cos(d*x+c))*sin(d*x+c)+5/16*d*x+5/16*c)+1/5*(4*A*a*b^3+6*B*a^2*b^2)/d*(8/3+cos(d*x+c)^4+4/3*cos(d*x+c)^2)*sin(d*x+c)+(6*A*a^2*b^2+4*B*a^3*b)/d*(1/4*(cos(d*x+c)^3+3/2*cos(d*x+c))*sin(d*x+c)+3/8*d*x+3/8*c)+1/3*(4*A*a^3*b+B*a^4)/d*(2+cos(d*x+c)^2)*sin(d*x+c)+a^4*A/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+1/7*B*b^4/d*(16/5+cos(d*x+c)^6+6/5*cos(d*x+c)^4+8/5*cos(d*x+c)^2)*sin(d*x+c)`

3.240.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.79

$$\int \cos^2(c+dx)(a+b\cos(c+dx))^4(A+B\cos(c+dx))dx$$

$$= \frac{105(8Aa^4+24Ba^3b+36Aa^2b^2+20Bab^3+5Ab^4)dx + (240Bb^4\cos(dx+c)^6 + 280(4Bab^3+Ab^4)\cos(dx+c)^5 + \dots}{105(8Aa^4+24Ba^3b+36Aa^2b^2+20Bab^3+5Ab^4) + (240Bb^4\cos(dx+c)^6 + 280(4Bab^3+Ab^4)\cos(dx+c)^5 + \dots)}$$

```
input integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="f
ricas")
```

```
output 1/1680*(105*(8*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 20*B*a*b^3 + 5*A*b^4)*d
*x + (240*B*b^4*cos(d*x + c)^6 + 280*(4*B*a*b^3 + A*b^4)*cos(d*x + c)^5 +
1120*B*a^4 + 4480*A*a^3*b + 5376*B*a^2*b^2 + 3584*A*a*b^3 + 768*B*b^4 + 96
*(21*B*a^2*b^2 + 14*A*a*b^3 + 3*B*b^4)*cos(d*x + c)^4 + 70*(24*B*a^3*b + 3
6*A*a^2*b^2 + 20*B*a*b^3 + 5*A*b^4)*cos(d*x + c)^3 + 16*(35*B*a^4 + 140*A*
a^3*b + 168*B*a^2*b^2 + 112*A*a*b^3 + 24*B*b^4)*cos(d*x + c)^2 + 105*(8*A*
a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 20*B*a*b^3 + 5*A*b^4)*cos(d*x + c))*sin(
d*x + c))/d
```

3.240.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1017 vs. $2(391) = 782$.

Time = 0.59 (sec) , antiderivative size = 1017, normalized size of antiderivative = 2.78

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)**2*(a+b*cos(d*x+c))**4*(A+B*cos(d*x+c)),x)
```


output `Piecewise((A**4*x*sin(c + d*x)**2/2 + A**4*x*cos(c + d*x)**2/2 + A**4*sin(c + d*x)*cos(c + d*x)/(2*d) + 8*A**3*b*sin(c + d*x)**3/(3*d) + 4*A**3*b*sin(c + d*x)*cos(c + d*x)**2/d + 9*A**2*b**2*x*sin(c + d*x)**4/4 + 9*A**2*b**2*x*sin(c + d*x)**2*cos(c + d*x)**2/2 + 9*A**2*b**2*x*cos(c + d*x)**4/4 + 9*A**2*b**2*sin(c + d*x)**3*cos(c + d*x)/(4*d) + 15*A**2*b**2*sin(c + d*x)*cos(c + d*x)**3/(4*d) + 32*A*a*b**3*sin(c + d*x)**5/(15*d) + 16*A*a*b**3*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 4*A*a*b**3*sin(c + d*x)*cos(c + d*x)**4/d + 5*A*b**4*x*sin(c + d*x)**6/16 + 15*A*b**4*x*sin(c + d*x)**4*cos(c + d*x)**2/16 + 15*A*b**4*x*sin(c + d*x)**2*cos(c + d*x)**4/16 + 5*A*b**4*x*cos(c + d*x)**6/16 + 5*A*b**4*sin(c + d*x)**5*cos(c + d*x)/(16*d) + 5*A*b**4*sin(c + d*x)**3*cos(c + d*x)**3/(6*d) + 11*A*b**4*sin(c + d*x)*cos(c + d*x)**5/(16*d) + 2*B*a**4*sin(c + d*x)**3/(3*d) + B*a**4*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*a**3*b*x*sin(c + d*x)**4/2 + 3*B*a**3*b*x*sin(c + d*x)**2*cos(c + d*x)**2 + 3*B*a**3*b*x*cos(c + d*x)**4/2 + 3*B*a**3*b*sin(c + d*x)**3*cos(c + d*x)/(2*d) + 5*B*a**3*b*sin(c + d*x)*cos(c + d*x)**3/(2*d) + 16*B*a**2*b**2*sin(c + d*x)**5/(5*d) + 8*B*a**2*b**2*sin(c + d*x)**3*cos(c + d*x)**2/d + 6*B*a**2*b**2*sin(c + d*x)*cos(c + d*x)**4/d + 5*B*a*b**3*x*sin(c + d*x)**6/4 + 15*B*a*b**3*x*sin(c + d*x)**4*cos(c + d*x)**2/4 + 15*B*a*b**3*x*sin(c + d*x)**2*cos(c + d*x)**4/4 + 5*B*a*b**3*x*cos(c + d*x)**6/4 + 5*B*a*b**3*sin(c + d*x)**5*cos(c + d*x)...`

3.240.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 366, normalized size of antiderivative = 1.00

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{1680(2dx + 2c + \sin(2dx + 2c))Aa^4 - 2240(\sin(dx + c)^3 - 3\sin(dx + c))Ba^4 - 8960(\sin(dx + c)^3 -$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output $1/6720*(1680*(2*d*x + 2*c + \sin(2*d*x + 2*c))*A*a^4 - 2240*(\sin(d*x + c))^3 - 3*\sin(d*x + c))*B*a^4 - 8960*(\sin(d*x + c))^3 - 3*\sin(d*x + c))*A*a^3*b + 840*(12*d*x + 12*c + \sin(4*d*x + 4*c) + 8*\sin(2*d*x + 2*c))*B*a^3*b + 1260*(12*d*x + 12*c + \sin(4*d*x + 4*c) + 8*\sin(2*d*x + 2*c))*A*a^2*b^2 + 2688*(3*\sin(d*x + c))^5 - 10*\sin(d*x + c)^3 + 15*\sin(d*x + c))*B*a^2*b^2 + 1792*(3*\sin(d*x + c))^5 - 10*\sin(d*x + c)^3 + 15*\sin(d*x + c))*A*a*b^3 - 140*(4*\sin(2*d*x + 2*c))^3 - 60*d*x - 60*c - 9*\sin(4*d*x + 4*c) - 48*\sin(2*d*x + 2*c))*B*a*b^3 - 35*(4*\sin(2*d*x + 2*c))^3 - 60*d*x - 60*c - 9*\sin(4*d*x + 4*c) - 48*\sin(2*d*x + 2*c))*A*b^4 - 192*(5*\sin(d*x + c))^7 - 21*\sin(d*x + c)^5 + 35*\sin(d*x + c)^3 - 35*\sin(d*x + c))*B*b^4)/d$

3.240.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.86

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{Bb^4 \sin(7dx + 7c)}{448d} + \frac{1}{16} (8Aa^4 + 24Ba^3b + 36Aa^2b^2 + 20Bab^3 + 5Ab^4)x$$

$$+ \frac{(4Bab^3 + Ab^4) \sin(6dx + 6c)}{192d} + \frac{(24Ba^2b^2 + 16Aab^3 + 7Bb^4) \sin(5dx + 5c)}{320d}$$

$$+ \frac{(8Ba^3b + 12Aa^2b^2 + 12Bab^3 + 3Ab^4) \sin(4dx + 4c)}{64d}$$

$$+ \frac{(16Ba^4 + 64Aa^3b + 120Ba^2b^2 + 80Aab^3 + 21Bb^4) \sin(3dx + 3c)}{192d}$$

$$+ \frac{(16Aa^4 + 64Ba^3b + 96Aa^2b^2 + 60Bab^3 + 15Ab^4) \sin(2dx + 2c)}{64d}$$

$$+ \frac{(48Ba^4 + 192Aa^3b + 240Ba^2b^2 + 160Aab^3 + 35Bb^4) \sin(dx + c)}{64d}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="giac")`

output $1/448*B*b^4*\sin(7*d*x + 7*c)/d + 1/16*(8*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 20*B*a*b^3 + 5*A*b^4)*x + 1/192*(4*B*a*b^3 + A*b^4)*\sin(6*d*x + 6*c)/d + 1/320*(24*B*a^2*b^2 + 16*A*a*b^3 + 7*B*b^4)*\sin(5*d*x + 5*c)/d + 1/64*(8*B*a^3*b + 12*A*a^2*b^2 + 12*B*a*b^3 + 3*A*b^4)*\sin(4*d*x + 4*c)/d + 1/192*(16*B*a^4 + 64*A*a^3*b + 120*B*a^2*b^2 + 80*A*a*b^3 + 21*B*b^4)*\sin(3*d*x + 3*c)/d + 1/64*(16*A*a^4 + 64*B*a^3*b + 96*A*a^2*b^2 + 60*B*a*b^3 + 15*A*b^4)*\sin(2*d*x + 2*c)/d + 1/64*(48*B*a^4 + 192*A*a^3*b + 240*B*a^2*b^2 + 160*A*a*b^3 + 35*B*b^4)*\sin(d*x + c)/d$

3.240.9 Mupad [B] (verification not implemented)

Time = 2.91 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.19

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{420 A a^4 \sin(2c + 2dx) + \frac{1575 A b^4 \sin(2c+2dx)}{4} + 140 B a^4 \sin(3c + 3dx) + \frac{315 A b^4 \sin(4c+4dx)}{4} + \frac{35 A b^4 \sin(6c+6dx)}{4}}{1}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4,x)`

output

```
(420*A*a^4*sin(2*c + 2*d*x) + (1575*A*b^4*sin(2*c + 2*d*x))/4 + 140*B*a^4*
sin(3*c + 3*d*x) + (315*A*b^4*sin(4*c + 4*d*x))/4 + (35*A*b^4*sin(6*c + 6*
d*x))/4 + (735*B*b^4*sin(3*c + 3*d*x))/4 + (147*B*b^4*sin(5*c + 5*d*x))/4
+ (15*B*b^4*sin(7*c + 7*d*x))/4 + 1260*B*a^4*sin(c + d*x) + (3675*B*b^4*si
n(c + d*x))/4 + 4200*A*a*b^3*sin(c + d*x) + 5040*A*a^3*b*sin(c + d*x) + 84
0*A*a^4*d*x + 525*A*b^4*d*x + 700*A*a*b^3*sin(3*c + 3*d*x) + 560*A*a^3*b*s
in(3*c + 3*d*x) + 84*A*a*b^3*sin(5*c + 5*d*x) + 1575*B*a*b^3*sin(2*c + 2*d
*x) + 1680*B*a^3*b*sin(2*c + 2*d*x) + 315*B*a*b^3*sin(4*c + 4*d*x) + 210*B
*a^3*b*sin(4*c + 4*d*x) + 35*B*a*b^3*sin(6*c + 6*d*x) + 6300*B*a^2*b^2*sin
(c + d*x) + 2520*A*a^2*b^2*sin(2*c + 2*d*x) + 315*A*a^2*b^2*sin(4*c + 4*d*
x) + 1050*B*a^2*b^2*sin(3*c + 3*d*x) + 126*B*a^2*b^2*sin(5*c + 5*d*x) + 21
00*B*a*b^3*d*x + 2520*B*a^3*b*d*x + 3780*A*a^2*b^2*d*x)/(1680*d)
```

3.241 $\int \cos(c+dx)(a+b \cos(c+dx))^4(A+B \cos(c+dx)) dx$

3.241.1 Optimal result	2221
3.241.2 Mathematica [A] (verified)	2222
3.241.3 Rubi [A] (verified)	2222
3.241.4 Maple [A] (verified)	2226
3.241.5 Fricas [A] (verification not implemented)	2227
3.241.6 Sympy [B] (verification not implemented)	2227
3.241.7 Maxima [A] (verification not implemented)	2228
3.241.8 Giac [A] (verification not implemented)	2229
3.241.9 Mupad [B] (verification not implemented)	2230

3.241.1 Optimal result

Integrand size = 29, antiderivative size = 325

$$\int \cos(c+dx)(a+b \cos(c+dx))^4(A+B \cos(c+dx)) dx$$

$$= \frac{1}{16}(32a^3Ab + 24aAb^3 + 8a^4B + 36a^2b^2B + 5b^4B) x$$

$$+ \frac{(24a^4Ab + 224a^2Ab^3 + 32Ab^5 - 4a^5B + 121a^3b^2B + 128ab^4B) \sin(c+dx)}{60bd}$$

$$+ \frac{(48a^3Ab + 232aAb^3 - 8a^4B + 178a^2b^2B + 75b^4B) \cos(c+dx) \sin(c+dx)}{240d}$$

$$+ \frac{(24a^2Ab + 32Ab^3 - 4a^3B + 53ab^2B) (a+b \cos(c+dx))^2 \sin(c+dx)}{120bd}$$

$$+ \frac{(24aAb - 4a^2B + 25b^2B) (a+b \cos(c+dx))^3 \sin(c+dx)}{120bd}$$

$$+ \frac{(6Ab - aB)(a+b \cos(c+dx))^4 \sin(c+dx)}{30bd} + \frac{B(a+b \cos(c+dx))^5 \sin(c+dx)}{6bd}$$

output

```
1/16*(32*A*a^3*b+24*A*a*b^3+8*B*a^4+36*B*a^2*b^2+5*B*b^4)*x+1/60*(24*A*a^4
*b+224*A*a^2*b^3+32*A*b^5-4*B*a^5+121*B*a^3*b^2+128*B*a*b^4)*sin(d*x+c)/b/
d+1/240*(48*A*a^3*b+232*A*a*b^3-8*B*a^4+178*B*a^2*b^2+75*B*b^4)*cos(d*x+c)
*sin(d*x+c)/d+1/120*(24*A*a^2*b+32*A*b^3-4*B*a^3+53*B*a*b^2)*(a+b*cos(d*x+
c))^2*sin(d*x+c)/b/d+1/120*(24*A*a*b-4*B*a^2+25*B*b^2)*(a+b*cos(d*x+c))^3*
sin(d*x+c)/b/d+1/30*(6*A*b-B*a)*(a+b*cos(d*x+c))^4*sin(d*x+c)/b/d+1/6*B*(a
+b*cos(d*x+c))^5*sin(d*x+c)/b/d
```

3.241.2 Mathematica [A] (verified)

Time = 2.88 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.02

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{1920a^3Abc + 1440aAb^3c + 480a^4Bc + 2160a^2b^2Bc + 300b^4Bc + 1920a^3Abdx + 1440aAb^3dx + 480a^4Bdx}{960d}$$

input `Integrate[Cos[c + d*x]*(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output $(1920*a^3*A*b*c + 1440*a*A*b^3*c + 480*a^4*B*c + 2160*a^2*b^2*B*c + 300*b^4*B*c + 1920*a^3*A*b*d*x + 1440*a*A*b^3*d*x + 480*a^4*B*d*x + 2160*a^2*b^2*B*d*x + 300*b^4*B*d*x + 120*(8*a^4*A + 36*a^2*A*b^2 + 5*A*b^4 + 24*a^3*b*B + 20*a*b^3*B)*\text{Sin}[c + d*x] + 15*(64*a^3*A*b + 64*a*A*b^3 + 16*a^4*B + 96*a^2*b^2*B + 15*b^4*B)*\text{Sin}[2*(c + d*x)] + 480*a^2*A*b^2*\text{Sin}[3*(c + d*x)] + 100*A*b^4*\text{Sin}[3*(c + d*x)] + 320*a^3*b*B*\text{Sin}[3*(c + d*x)] + 400*a*b^3*B*\text{Sin}[3*(c + d*x)] + 120*a*A*b^3*\text{Sin}[4*(c + d*x)] + 180*a^2*b^2*B*\text{Sin}[4*(c + d*x)] + 45*b^4*B*\text{Sin}[4*(c + d*x)] + 12*A*b^4*\text{Sin}[5*(c + d*x)] + 48*a*b^3*B*\text{Sin}[5*(c + d*x)] + 5*b^4*B*\text{Sin}[6*(c + d*x)])/(960*d)$

3.241.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$, Rules used = {3042, 3447, 3042, 3502, 3042, 3232, 3042, 3232, 27, 3042, 3232, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^4 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int (a + b \cos(c + dx))^4 (A \cos(c + dx) + B \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

3.241. $\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^4 \left(A \sin \left(c + dx + \frac{\pi}{2} \right) + B \sin \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx \\
& \quad \downarrow \text{3502} \\
& \frac{\int (a + b \cos(c + dx))^4 (5bB + (6Ab - aB) \cos(c + dx)) dx}{6b} + \frac{B \sin(c + dx)(a + b \cos(c + dx))^5}{6bd} \\
& \quad \downarrow \text{3042} \\
& \frac{\int (a + b \sin(c + dx + \frac{\pi}{2}))^4 (5bB + (6Ab - aB) \sin(c + dx + \frac{\pi}{2})) dx}{6b} + \\
& \quad \frac{B \sin(c + dx)(a + b \cos(c + dx))^5}{6bd} \\
& \quad \downarrow \text{3232} \\
& \frac{\frac{1}{5} \int (a + b \cos(c + dx))^3 (3b(8Ab + 7aB) + (-4Ba^2 + 24Aba + 25b^2B) \cos(c + dx)) dx + \frac{(6Ab - aB) \sin(c + dx)(a + b \cos(c + dx))^5}{5d}}{6b}}{6bd} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{5} \int (a + b \sin(c + dx + \frac{\pi}{2}))^3 (3b(8Ab + 7aB) + (-4Ba^2 + 24Aba + 25b^2B) \sin(c + dx + \frac{\pi}{2})) dx + \frac{(6Ab - aB) \sin(c + dx)(a + b \cos(c + dx))^5}{5d}}{6b}}{6bd} \\
& \quad \downarrow \text{3232} \\
& \frac{\frac{1}{5} \left(\frac{1}{4} \int 3(a + b \cos(c + dx))^2 (b(24Ba^2 + 56Aba + 25b^2B) + (-4Ba^3 + 24Aba^2 + 53b^2Ba + 32Ab^3) \cos(c + dx)) dx \right)}{6b}}{6bd} \\
& \quad \downarrow \text{27} \\
& \frac{\frac{1}{5} \left(\frac{3}{4} \int (a + b \cos(c + dx))^2 (b(24Ba^2 + 56Aba + 25b^2B) + (-4Ba^3 + 24Aba^2 + 53b^2Ba + 32Ab^3) \cos(c + dx)) dx \right)}{6b}}{6bd} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.241. $\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$

$$\frac{\frac{1}{5} \left(\frac{3}{4} \int (a + b \sin(c + dx + \frac{\pi}{2}))^2 (b(24Ba^2 + 56Aba + 25b^2B) + (-4Ba^3 + 24Aba^2 + 53b^2Ba + 32Ab^3) \sin(c + dx)) \right)}{6b}$$

$$\frac{B \sin(c + dx)(a + b \cos(c + dx))^5}{6bd}$$

↓ 3232

$$\frac{\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \int (a + b \cos(c + dx)) (b(64Ba^3 + 216Aba^2 + 181b^2Ba + 64Ab^3) + (-8Ba^4 + 48Aba^3 + 178b^2Ba^2 + 232Ab^3) \cos(c + dx)) \right) \right)}{6b}$$

$$\frac{B \sin(c + dx)(a + b \cos(c + dx))^5}{6bd}$$

↓ 3042

$$\frac{\frac{1}{5} \left(\frac{3}{4} \left(\frac{1}{3} \int (a + b \sin(c + dx + \frac{\pi}{2})) (b(64Ba^3 + 216Aba^2 + 181b^2Ba + 64Ab^3) + (-8Ba^4 + 48Aba^3 + 178b^2Ba^2 + 232Ab^3) \cos(c + dx)) \right) \right)}{6b}$$

$$\frac{B \sin(c + dx)(a + b \cos(c + dx))^5}{6bd}$$

↓ 3213

$$\frac{\frac{1}{5} \left(\frac{(-4a^2B + 24aAb + 25b^2B) \sin(c + dx)(a + b \cos(c + dx))^3}{4d} + \frac{3}{4} \left(\frac{(-4a^3B + 24a^2Ab + 53ab^2B + 32Ab^3) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{1}{3} \left(\frac{b(-8a^4 + 48Aba^3 + 178b^2Ba^2 + 232Ab^3) \cos(c + dx)}{3d} \right) \right) \right)}{6b}$$

$$\frac{B \sin(c + dx)(a + b \cos(c + dx))^5}{6bd}$$

input `Int[Cos[c + d*x]*(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output `(B*(a + b*Cos[c + d*x])^5*Sin[c + d*x])/(6*b*d) + (((6*A*b - a*B)*(a + b*Cos[c + d*x])^4*Sin[c + d*x])/(5*d) + (((24*a*A*b - 4*a^2*B + 25*b^2*B)*(a + b*Cos[c + d*x])^3*Sin[c + d*x])/(4*d) + (3*(((24*a^2*A*b + 32*A*b^3 - 4*a^3*B + 53*a*b^2*B)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + ((15*b*(3*2*a^3*A*b + 24*a*A*b^3 + 8*a^4*B + 36*a^2*b^2*B + 5*b^4*B)*x)/2 + (2*(24*a^4*A*b + 224*a^2*A*b^3 + 32*A*b^5 - 4*a^5*B + 121*a^3*b^2*B + 128*a*b^4*B)*Sin[c + d*x])/d + (b*(48*a^3*A*b + 232*a*A*b^3 - 8*a^4*B + 178*a^2*b^2*B + 75*b^4*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d))/3))/4)/5)/(6*b)`

3.241.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3213 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2*a*c + b*d)*(x/2), x] + (-Simp[(b*c + a*d)*(Cos[e + f*x]/f), x] - Simp[b*d*Cos[e + f*x]*(Sin[e + f*x]/(2*f)), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.241.4 Maple [A] (verified)

Time = 4.76 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.74

method	result
parts	$\frac{(A b^4 + 4 B a b^3) \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5d} + \frac{(4 A a b^3 + 6 B a^2 b^2) \left(\frac{(\cos^3(dx+c) + \frac{3 \cos(\frac{dx+c}{2})) \sin(dx+c)}{4} \right)}{d}$
parallelrisch	$(960 A a^3 b + 960 A a b^3 + 240 B a^4 + 1440 B a^2 b^2 + 225 B b^4) \sin(2dx+2c) + (480 A a^2 b^2 + 100 A b^4 + 320 B a^3 b + 400 B a b^3) \sin(dx+c)$
derivativedivides	$B b^4 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{A b^4 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5}$
default	$B b^4 \left(\frac{\left(\cos^5(dx+c) + \frac{5(\cos^3(dx+c))}{4} + \frac{15 \cos(dx+c)}{8} \right) \sin(dx+c)}{6} + \frac{5dx}{16} + \frac{5c}{16} \right) + \frac{A b^4 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5}$
risch	$2x A a^3 b + \frac{3x A a b^3}{2} + \frac{9x B a^2 b^2}{4} + \frac{\sin(dx+c) a^4 A}{d} + \frac{5 \sin(dx+c) A b^4}{8d} + \frac{\sin(5dx+5c) A b^4}{80d} + \frac{3 \sin(4dx+4c) B}{64d}$
norman	$\frac{(2 A a^3 b + \frac{3}{2} A a b^3 + \frac{1}{2} B a^4 + \frac{9}{4} B a^2 b^2 + \frac{5}{16} B b^4) x + (2 A a^3 b + \frac{3}{2} A a b^3 + \frac{1}{2} B a^4 + \frac{9}{4} B a^2 b^2 + \frac{5}{16} B b^4) x \left(\tan^{12} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (12 A a^3 b + 3 A a b^3 + B a^4 + 9 B a^2 b^2 + 5 B b^4) \tan \left(\frac{dx}{2} + \frac{c}{2} \right)}{1}$

input `int(cos(d*x+c)*(a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/5*(A*b^4+4*B*a*b^3)/d*(8/3+cos(d*x+c)^4+4/3*cos(d*x+c)^2)*sin(d*x+c)+(4*A*a*b^3+6*B*a^2*b^2)/d*(1/4*(cos(d*x+c)^3+3/2*cos(d*x+c))*sin(d*x+c)+3/8*d*x+3/8*c)+1/3*(6*A*a^2*b^2+4*B*a^3*b)/d*(2+cos(d*x+c)^2)*sin(d*x+c)+(4*A*a^3*b+B*a^4)/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+B*b^4/d*(1/6*(cos(d*x+c)^5+5/4*cos(d*x+c)^3+15/8*cos(d*x+c))*sin(d*x+c)+5/16*d*x+5/16*c)+1/d*sin(d*x+c)*a^4*A`

3.241.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.75

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{15(8Ba^4 + 32Aa^3b + 36Ba^2b^2 + 24Aab^3 + 5Bb^4)dx + (40Bb^4 \cos(dx + c)^5 + 240Aa^4 + 640Ba^3b + 960Aa^2b^2 + 512Bab^3 + 128A^2b^4 + 48(4Bab^3 + A^2b^4)\cos(dx + c)^4 + 10(36Bab^2 + 24Aa^2b^3 + 5B^2b^4)\cos(dx + c)^3 + 32(10Bab^3 + 15Aa^2b^2 + 8B^2ab^3 + 2A^2b^4)\cos(dx + c)^2 + 15(8Ba^4 + 32Aa^3b + 36Ba^2b^2 + 24Aa^2b^3 + 5B^2b^4)\cos(dx + c))\sin(dx + c)}{d}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/240*(15*(8*B*a^4 + 32*A*a^3*b + 36*B*a^2*b^2 + 24*A*a*b^3 + 5*B*b^4)*d*x + (40*B*b^4*cos(d*x + c)^5 + 240*A*a^4 + 640*B*a^3*b + 960*A*a^2*b^2 + 512*B*a*b^3 + 128*A*b^4 + 48*(4*B*a*b^3 + A*b^4)*cos(d*x + c)^4 + 10*(36*B*a^2*b^2 + 24*A*a*b^3 + 5*B*b^4)*cos(d*x + c)^3 + 32*(10*B*a^3*b + 15*A*a^2*b^2 + 8*B*a*b^3 + 2*A*b^4)*cos(d*x + c)^2 + 15*(8*B*a^4 + 32*A*a^3*b + 36*B*a^2*b^2 + 24*A*a*b^3 + 5*B*b^4)*cos(d*x + c))*sin(d*x + c))/d`

3.241.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 811 vs. 2(335) = 670.

Time = 0.43 (sec) , antiderivative size = 811, normalized size of antiderivative = 2.50

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \begin{cases} \frac{Aa^4 \sin(c+dx)}{d} + 2Aa^3bx \sin^2(c + dx) + 2Aa^3bx \cos^2(c + dx) + \frac{2Aa^3b \sin(c+dx) \cos(c+dx)}{d} + \frac{4Aa^2b^2 \sin^3(c+dx)}{d} + \\ x(A + B \cos(c))(a + b \cos(c))^4 \cos(c) \end{cases}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))**4*(A+B*cos(d*x+c)),x)`

output

```
Piecewise((A**4*sin(c + d*x)/d + 2*A**3*b*x*sin(c + d*x)**2 + 2*A**3*b*x*cos(c + d*x)**2 + 2*A**3*b*sin(c + d*x)*cos(c + d*x)/d + 4*A**2*b**2*sin(c + d*x)**3/d + 6*A**2*b**2*sin(c + d*x)*cos(c + d*x)**2/d + 3*A**2*b**3*x*sin(c + d*x)**4/2 + 3*A**2*b**3*x*cos(c + d*x)**4/2 + 3*A**2*b**3*sin(c + d*x)**3*cos(c + d*x))/(2*d) + 5*A**2*b**3*sin(c + d*x)*cos(c + d*x)**3/(2*d) + 8*A**2*b**4*sin(c + d*x)**5/(15*d) + 4*A**2*b**4*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + A**4*sin(c + d*x)*cos(c + d*x)**4/d + B**4*x*sin(c + d*x)**2/2 + B**4*x*cos(c + d*x)**2/2 + B**4*sin(c + d*x)*cos(c + d*x)/(2*d) + 8*B**3*b*sin(c + d*x)**3/(3*d) + 4*B**3*b*sin(c + d*x)*cos(c + d*x)**2/d + 9*B**2*b**2*x*sin(c + d*x)**4/4 + 9*B**2*b**2*x*cos(c + d*x)**4/4 + 9*B**2*b**2*sin(c + d*x)**3*cos(c + d*x)/(4*d) + 15*B**2*b**2*sin(c + d*x)*cos(c + d*x)**3/(4*d) + 32*B**2*b**3*sin(c + d*x)**5/(15*d) + 16*B**2*b**3*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + 4*B**2*b**3*sin(c + d*x)*cos(c + d*x)**4/d + 5*B**4*x*sin(c + d*x)**6/16 + 15*B**4*x*cos(c + d*x)**2/16 + 15*B**4*x*sin(c + d*x)**2*cos(c + d*x)**4/16 + 5*B**4*x*cos(c + d*x)**6/16 + 5*B**4*sin(c + d*x)**5*cos(c + d*x)/(16*d) + 5*B**4*sin(c + d*x)**3*cos(c + d*x)**3/(6*d) + 11*B**4*sin(c + d*x)*cos(c + d*x)**5/(16*d), Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**4*cos(c), True))
```

3.241.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 307, normalized size of antiderivative = 0.94

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{240(2dx + 2c + \sin(2dx + 2c))Ba^4 + 960(2dx + 2c + \sin(2dx + 2c))Aa^3b - 1280(\sin(dx + c))^3 - 3$$

input

```
integrate(cos(d*x+c)*(a+b*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

output $\frac{1}{960}(240(2dx + 2c + \sin(2dx + 2c))B^2a^4 + 960(2dx + 2c + \sin(2dx + 2c))A^2a^3b - 1280(\sin(dx + c)^3 - 3\sin(dx + c))B^2a^3b - 1920(\sin(dx + c)^3 - 3\sin(dx + c))A^2a^2b^2 + 180(12dx + 12c + \sin(4dx + 4c) + 8\sin(2dx + 2c))B^2a^2b^2 + 120(12dx + 12c + \sin(4dx + 4c) + 8\sin(2dx + 2c))A^2a^2b^3 + 256(3\sin(dx + c)^5 - 10\sin(dx + c)^3 + 15\sin(dx + c))B^2a^2b^3 + 64(3\sin(dx + c)^5 - 10\sin(dx + c)^3 + 15\sin(dx + c))A^2a^2b^4 - 5(4\sin(2dx + 2c)^3 - 60dx - 60c - 9\sin(4dx + 4c) - 48\sin(2dx + 2c))B^2b^4 + 960A^2a^4\sin(dx + c))/d$

3.241.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.81

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{Bb^4 \sin(6dx + 6c)}{192d} + \frac{1}{16} (8Ba^4 + 32Aa^3b + 36Ba^2b^2 + 24Aab^3 + 5Bb^4)x$$

$$+ \frac{(4Bab^3 + Ab^4) \sin(5dx + 5c)}{80d} + \frac{(12Ba^2b^2 + 8Aab^3 + 3Bb^4) \sin(4dx + 4c)}{64d}$$

$$+ \frac{(16Ba^3b + 24Aa^2b^2 + 20Bab^3 + 5Ab^4) \sin(3dx + 3c)}{48d}$$

$$+ \frac{(16Ba^4 + 64Aa^3b + 96Ba^2b^2 + 64Aab^3 + 15Bb^4) \sin(2dx + 2c)}{64d}$$

$$+ \frac{(8Aa^4 + 24Ba^3b + 36Aa^2b^2 + 20Bab^3 + 5Ab^4) \sin(dx + c)}{8d}$$

input `integrate(cos(dx+c)*(a+b*cos(dx+c))^4*(A+B*cos(dx+c)),x, algorithm="giac")`

output $\frac{1}{192}B^2b^4\sin(6dx + 6c)/d + \frac{1}{16}(8B^2a^4 + 32A^2a^3b + 36B^2a^2b^2 + 24A^2a^2b^3 + 5B^2b^4)x + \frac{1}{80}(4B^2a^2b^3 + A^2b^4)\sin(5dx + 5c)/d + \frac{1}{64}(12B^2a^2b^2 + 8A^2a^2b^3 + 3B^2b^4)\sin(4dx + 4c)/d + \frac{1}{48}(16B^2a^3b + 24A^2a^2b^2 + 20B^2a^2b^3 + 5A^2b^4)\sin(3dx + 3c)/d + \frac{1}{64}(16B^2a^4 + 64A^2a^3b + 96B^2a^2b^2 + 64A^2a^2b^3 + 15B^2b^4)\sin(2dx + 2c)/d + \frac{1}{8}(8A^2a^4 + 24B^2a^3b + 36A^2a^2b^2 + 20B^2a^2b^3 + 5A^2b^4)\sin(dx + c)/d$

3.241.9 Mupad [B] (verification not implemented)

Time = 1.47 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.24

$$\int \cos(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$= \frac{B a^4 x}{2} + \frac{5 B b^4 x}{16} + \frac{3 A a b^3 x}{2} + 2 A a^3 b x + \frac{A a^4 \sin(c + dx)}{d} + \frac{5 A b^4 \sin(c + dx)}{8 d}$$

$$+ \frac{9 B a^2 b^2 x}{4} + \frac{B a^4 \sin(2c + 2dx)}{4d} + \frac{5 A b^4 \sin(3c + 3dx)}{48d} + \frac{A b^4 \sin(5c + 5dx)}{80d}$$

$$+ \frac{15 B b^4 \sin(2c + 2dx)}{64d} + \frac{3 B b^4 \sin(4c + 4dx)}{64d} + \frac{B b^4 \sin(6c + 6dx)}{192d}$$

$$+ \frac{A a b^3 \sin(2c + 2dx)}{d} + \frac{A a^3 b \sin(2c + 2dx)}{d} + \frac{A a b^3 \sin(4c + 4dx)}{8d}$$

$$+ \frac{9 A a^2 b^2 \sin(c + dx)}{2d} + \frac{5 B a b^3 \sin(3c + 3dx)}{12d} + \frac{B a^3 b \sin(3c + 3dx)}{3d}$$

$$+ \frac{B a b^3 \sin(5c + 5dx)}{20d} + \frac{A a^2 b^2 \sin(3c + 3dx)}{2d} + \frac{3 B a^2 b^2 \sin(2c + 2dx)}{2d}$$

$$+ \frac{3 B a^2 b^2 \sin(4c + 4dx)}{16d} + \frac{5 B a b^3 \sin(c + dx)}{2d} + \frac{3 B a^3 b \sin(c + dx)}{d}$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4,x)`output `(B*a^4*x)/2 + (5*B*b^4*x)/16 + (3*A*a*b^3*x)/2 + 2*A*a^3*b*x + (A*a^4*sin(c + d*x))/d + (5*A*b^4*sin(c + d*x))/(8*d) + (9*B*a^2*b^2*x)/4 + (B*a^4*sin(2*c + 2*d*x))/(4*d) + (5*A*b^4*sin(3*c + 3*d*x))/(48*d) + (A*b^4*sin(5*c + 5*d*x))/(80*d) + (15*B*b^4*sin(2*c + 2*d*x))/(64*d) + (3*B*b^4*sin(4*c + 4*d*x))/(64*d) + (B*b^4*sin(6*c + 6*d*x))/(192*d) + (A*a*b^3*sin(2*c + 2*d*x))/d + (A*a^3*b*sin(2*c + 2*d*x))/d + (A*a*b^3*sin(4*c + 4*d*x))/(8*d) + (9*A*a^2*b^2*sin(c + d*x))/(2*d) + (5*B*a*b^3*sin(3*c + 3*d*x))/(12*d) + (B*a^3*b*sin(3*c + 3*d*x))/(3*d) + (B*a*b^3*sin(5*c + 5*d*x))/(20*d) + (A*a^2*b^2*sin(3*c + 3*d*x))/(2*d) + (3*B*a^2*b^2*sin(2*c + 2*d*x))/(2*d) + (3*B*a^2*b^2*sin(4*c + 4*d*x))/(16*d) + (5*B*a*b^3*sin(c + d*x))/(2*d) + (3*B*a^3*b*sin(c + d*x))/d`

3.242 $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$

3.242.1 Optimal result	2231
3.242.2 Mathematica [A] (verified)	2232
3.242.3 Rubi [A] (verified)	2232
3.242.4 Maple [A] (verified)	2234
3.242.5 Fricas [A] (verification not implemented)	2235
3.242.6 Sympy [B] (verification not implemented)	2236
3.242.7 Maxima [A] (verification not implemented)	2237
3.242.8 Giac [A] (verification not implemented)	2237
3.242.9 Mupad [B] (verification not implemented)	2238

3.242.1 Optimal result

Integrand size = 23, antiderivative size = 241

$$\begin{aligned} & \int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx \\ &= \frac{1}{8} (8a^4 A + 24a^2 Ab^2 + 3Ab^4 + 16a^3 b B + 12ab^3 B) x \\ &+ \frac{(95a^3 Ab + 80aAb^3 + 12a^4 B + 112a^2 b^2 B + 16b^4 B) \sin(c + dx)}{30d} \\ &+ \frac{b(130a^2 Ab + 45Ab^3 + 24a^3 B + 116ab^2 B) \cos(c + dx) \sin(c + dx)}{120d} \\ &+ \frac{(35aAb + 12a^2 B + 16b^2 B) (a + b \cos(c + dx))^2 \sin(c + dx)}{60d} \\ &+ \frac{(5Ab + 4aB) (a + b \cos(c + dx))^3 \sin(c + dx)}{20d} + \frac{B(a + b \cos(c + dx))^4 \sin(c + dx)}{5d} \end{aligned}$$

```
output 1/8*(8*A*a^4+24*A*a^2*b^2+3*A*b^4+16*B*a^3*b+12*B*a*b^3)*x+1/30*(95*A*a^3*b+80*A*a*b^3+12*B*a^4+112*B*a^2*b^2+16*B*b^4)*sin(d*x+c)/d+1/120*b*(130*A*a^2*b+45*A*b^3+24*B*a^3+116*B*a*b^2)*cos(d*x+c)*sin(d*x+c)/d+1/60*(35*A*a*b+12*B*a^2+16*B*b^2)*(a+b*cos(d*x+c))^2*sin(d*x+c)/d+1/20*(5*A*b+4*B*a)*(a+b*cos(d*x+c))^3*sin(d*x+c)/d+1/5*B*(a+b*cos(d*x+c))^4*sin(d*x+c)/d
```


$$\frac{1}{5} \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^3 \left(5aA + 4bB + (5Ab + 4aB) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5d}$$

↓ 3232

$$\frac{1}{5} \left(\frac{1}{4} \int (a + b \cos(c + dx))^2 (20Aa^2 + 28bBa + 15Ab^2 + (12Ba^2 + 35Aba + 16b^2B) \cos(c + dx)) dx + \frac{(4aB + 5A)}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^2 \left(20Aa^2 + 28bBa + 15Ab^2 + (12Ba^2 + 35Aba + 16b^2B) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5d} \right)$$

↓ 3232

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int (a + b \cos(c + dx)) (60Aa^3 + 108bBa^2 + 115Ab^2a + 32b^3B + (24Ba^3 + 130Aba^2 + 116b^2Ba + 45Ab^3)) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5d} \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right) \left(60Aa^3 + 108bBa^2 + 115Ab^2a + 32b^3B + (24Ba^3 + 130Aba^2 + 116b^2Ba + 45Ab^3) \right) dx + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5d} \right) \right)$$

↓ 3213

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{(12a^2B + 35aAb + 16b^2B) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{b(24a^3B + 130a^2Ab + 116ab^2B + 45Ab^3)}{2d} \right) + \frac{B \sin(c + dx)(a + b \cos(c + dx))^4}{5d} \right)$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x]),x]`

output $(B(a + b\cos[c + dx])^4\sin[c + dx])/(5d) + ((5A^2b + 4a^2B)(a + b\cos[c + dx])^3\sin[c + dx])/(4d) + (((35a^2Ab + 12a^2B + 16b^2B)(a + b\cos[c + dx])^2\sin[c + dx])/(3d) + ((15(8a^4A + 24a^2Ab^2 + 3Ab^4 + 16a^3bB + 12ab^3B)x)/2 + (2(95a^3Ab + 80a^2Ab^3 + 12a^4B + 112a^2b^2B + 16b^4B)\sin[c + dx])/d + (b(130a^2Ab + 45Ab^3 + 24a^3B + 116ab^2B)\cos[c + dx]\sin[c + dx])/(2d))/3)/4)/5$

3.242.3.1 Defintions of rubi rules used

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear } Q[u, x]$

rule 3213 $\text{Int}[(a + b\sin[e + f(x)])(c + d\sin[e + f(x)])(x), x_Symbol] \rightarrow \text{Simp}[(2ac + b^2d)(x/2), x] + (-\text{Simp}[(b^2c + a^2d)\cos[e + f(x)]/f], x) - \text{Simp}[b^2d\cos[e + f(x)](\sin[e + f(x)]/(2f)), x] \text{ ; Free } Q[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b^2c - a^2d, 0]$

rule 3232 $\text{Int}[(a + b\sin[e + f(x)])^m(c + d\sin[e + f(x)])(x), x_Symbol] \rightarrow \text{Simp}[(-d)\cos[e + f(x)](a + b\sin[e + f(x)])^m/(f(m + 1)), x] + \text{Simp}[1/(m + 1) \text{ Int}[(a + b\sin[e + f(x)])^{m-1}\text{Simp}[b^2d^m + a^2c(m + 1) + (a^2d^m + b^2c(m + 1))\sin[e + f(x)], x], x], x] \text{ ; Free } Q[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b^2c - a^2d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{IntegerQ}[2m]$

3.242.4 Maple [A] (verified)

Time = 3.92 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.82

method	result
parts	$a^4xA + \frac{(Ab^4+4Bab^3) \left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right)}{d} + \frac{(4Aab^3+6Ba^2b^2)(2+\cos^2(dx+c)) \sin(dx+c)}{3d}$
parallelrisch	$\frac{(720Aa^2b^2+120Ab^4+480Ba^3b+480Bab^3) \sin(2dx+2c) + (160Aab^3+240Ba^2b^2+50Bb^4) \sin(3dx+3c) + (15Ab^4+60Bb^3) \sin(4dx+4c)}{15}$
derivativdivides	$\frac{Bb^4 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + Ab^4 \left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 4Bab^3 \left(\frac{\cos^3(dx+c)}{3} \right)$
default	$\frac{Bb^4 \left(\frac{8}{3} + \cos^4(dx+c) + \frac{4(\cos^2(dx+c))}{3} \right) \sin(dx+c)}{5} + Ab^4 \left(\frac{(\cos^3(dx+c) + \frac{3\cos(dx+c)}{2}) \sin(dx+c)}{4} + \frac{3dx}{8} + \frac{3c}{8} \right) + 4Bab^3 \left(\frac{\cos^3(dx+c)}{3} \right)$
risch	$a^4xA + 3xAa^2b^2 + \frac{3xAb^4}{8} + 2xBa^3b + \frac{3xBab^3}{2} + \frac{4\sin(dx+c)Aa^3b}{d} + \frac{3\sin(dx+c)Aab^3}{d} + \frac{\sin(dx+c)Aa^2b^2}{d}$
norman	$\frac{(a^4A+3Aa^2b^2+\frac{3}{8}Ab^4+2Ba^3b+\frac{3}{2}Bab^3)x + (a^4A+3Aa^2b^2+\frac{3}{8}Ab^4+2Ba^3b+\frac{3}{2}Bab^3)x \left(\tan^{10} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (5a^4A+15Aa^2b^2+\frac{15}{8}Ab^4+10Ba^3b+\frac{15}{2}Bab^3)}{15}$

input `int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `a^4*x*A+(A*b^4+4*B*a*b^3)/d*(1/4*(cos(d*x+c)^3+3/2*cos(d*x+c))*sin(d*x+c)+3/8*d*x+3/8*c)+1/3*(4*A*a*b^3+6*B*a^2*b^2)/d*(2+cos(d*x+c)^2)*sin(d*x+c)+(6*A*a^2*b^2+4*B*a^3*b)/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+(4*A*a^3*b+B*a^4)/d*sin(d*x+c)+1/5*B*b^4/d*(8/3+cos(d*x+c)^4+4/3*cos(d*x+c)^2)*sin(d*x+c)`

3.242.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.82

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \frac{15(8Aa^4 + 16Ba^3b + 24Aa^2b^2 + 12Bab^3 + 3Ab^4)dx + (24Bb^4 \cos(dx + c)^4 + 120Ba^4 + 480Aa^3b + 480Aa^2b^2 + 120Bab^3 + 30Ab^4) \sin(dx + c) + (15Aa^4 + 30Aa^2b^2 + 15Ab^4) \cos(dx + c)^5 + (15Ba^4 + 30Bab^3 + 15Bb^4) \cos(dx + c)^4 + (15Aa^3b + 30Aa^2b^2 + 15Ab^4) \cos(dx + c)^3 + (15Ba^3b + 30Bab^3 + 15Bb^4) \cos(dx + c)^2 + (15Aa^2b^2 + 15Ab^4) \cos(dx + c) + 15Ba^2b^2 + 15Bab^3 + 15Bb^4}{15}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c)),x, algorithm="fracas")`

```
output 1/120*(15*(8*A*a^4 + 16*B*a^3*b + 24*A*a^2*b^2 + 12*B*a*b^3 + 3*A*b^4)*d*x
+ (24*B*b^4*cos(d*x + c)^4 + 120*B*a^4 + 480*A*a^3*b + 480*B*a^2*b^2 + 32
0*A*a*b^3 + 64*B*b^4 + 30*(4*B*a*b^3 + A*b^4)*cos(d*x + c)^3 + 16*(15*B*a^
2*b^2 + 10*A*a*b^3 + 2*B*b^4)*cos(d*x + c)^2 + 15*(16*B*a^3*b + 24*A*a^2*b
^2 + 12*B*a*b^3 + 3*A*b^4)*cos(d*x + c))*sin(d*x + c))/d
```

3.242.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 580 vs. $2(248) = 496$.

Time = 0.31 (sec) , antiderivative size = 580, normalized size of antiderivative = 2.41

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$= \begin{cases} Aa^4x + \frac{4Aa^3b \sin(c+dx)}{d} + 3Aa^2b^2x \sin^2(c + dx) + 3Aa^2b^2x \cos^2(c + dx) + \frac{3Aa^2b^2 \sin(c+dx) \cos(c+dx)}{d} + \frac{8Aab^3}{d} \\ x(A + B \cos(c)) (a + b \cos(c))^4 \end{cases}$$

```
input integrate((a+b*cos(d*x+c))**4*(A+B*cos(d*x+c)),x)
```

```
output Piecewise((A*a**4*x + 4*A*a**3*b*sin(c + d*x)/d + 3*A*a**2*b**2*x*sin(c +
d*x)**2 + 3*A*a**2*b**2*x*cos(c + d*x)**2 + 3*A*a**2*b**2*sin(c + d*x)*cos
(c + d*x)/d + 8*A*a*b**3*sin(c + d*x)**3/(3*d) + 4*A*a*b**3*sin(c + d*x)*c
os(c + d*x)**2/d + 3*A*b**4*x*sin(c + d*x)**4/8 + 3*A*b**4*x*sin(c + d*x)*
**2*cos(c + d*x)**2/4 + 3*A*b**4*x*cos(c + d*x)**4/8 + 3*A*b**4*sin(c + d*x)
)**3*cos(c + d*x)/(8*d) + 5*A*b**4*sin(c + d*x)*cos(c + d*x)**3/(8*d) + B*
a**4*sin(c + d*x)/d + 2*B*a**3*b*x*sin(c + d*x)**2 + 2*B*a**3*b*x*cos(c +
d*x)**2 + 2*B*a**3*b*sin(c + d*x)*cos(c + d*x)/d + 4*B*a**2*b**2*sin(c + d
*x)**3/d + 6*B*a**2*b**2*sin(c + d*x)*cos(c + d*x)**2/d + 3*B*a*b**3*x*sin
(c + d*x)**4/2 + 3*B*a*b**3*x*sin(c + d*x)**2*cos(c + d*x)**2 + 3*B*a*b**3
*x*cos(c + d*x)**4/2 + 3*B*a*b**3*sin(c + d*x)**3*cos(c + d*x)/(2*d) + 5*B
*a*b**3*sin(c + d*x)*cos(c + d*x)**3/(2*d) + 8*B*b**4*sin(c + d*x)**5/(15*
d) + 4*B*b**4*sin(c + d*x)**3*cos(c + d*x)**2/(3*d) + B*b**4*sin(c + d*x)*
cos(c + d*x)**4/d, Ne(d, 0)), (x*(A + B*cos(c))*(a + b*cos(c))**4, True))
```


3.242.9 Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.27

$$\begin{aligned}
& \int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx \\
&= Aa^4x + \frac{3Ab^4x}{8} + \frac{3Ba^3b^3x}{2} + 2Ba^3bx + \frac{Ba^4 \sin(c + dx)}{d} \\
&+ \frac{5Bb^4 \sin(c + dx)}{8d} + 3Aa^2b^2x + \frac{Ab^4 \sin(2c + 2dx)}{4d} \\
&+ \frac{Ab^4 \sin(4c + 4dx)}{32d} + \frac{5Bb^4 \sin(3c + 3dx)}{48d} + \frac{Bb^4 \sin(5c + 5dx)}{80d} \\
&+ \frac{Aab^3 \sin(3c + 3dx)}{3d} + \frac{Bab^3 \sin(2c + 2dx)}{d} + \frac{Ba^3b \sin(2c + 2dx)}{d} \\
&+ \frac{Bab^3 \sin(4c + 4dx)}{8d} + \frac{9Ba^2b^2 \sin(c + dx)}{2d} + \frac{3Aa^2b^2 \sin(2c + 2dx)}{2d} \\
&+ \frac{Ba^2b^2 \sin(3c + 3dx)}{2d} + \frac{3Aab^3 \sin(c + dx)}{d} + \frac{4Aa^3b \sin(c + dx)}{d}
\end{aligned}$$

```
input int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4,x)
```

```
output A*a^4*x + (3*A*b^4*x)/8 + (3*B*a*b^3*x)/2 + 2*B*a^3*b*x + (B*a^4*sin(c + d
*x))/d + (5*B*b^4*sin(c + d*x))/(8*d) + 3*A*a^2*b^2*x + (A*b^4*sin(2*c + 2
*d*x))/(4*d) + (A*b^4*sin(4*c + 4*d*x))/(32*d) + (5*B*b^4*sin(3*c + 3*d*x)
)/(48*d) + (B*b^4*sin(5*c + 5*d*x))/(80*d) + (A*a*b^3*sin(3*c + 3*d*x))/(3
*d) + (B*a*b^3*sin(2*c + 2*d*x))/d + (B*a^3*b*sin(2*c + 2*d*x))/d + (B*a*b
^3*sin(4*c + 4*d*x))/(8*d) + (9*B*a^2*b^2*sin(c + d*x))/(2*d) + (3*A*a^2*b
^2*sin(2*c + 2*d*x))/(2*d) + (B*a^2*b^2*sin(3*c + 3*d*x))/(2*d) + (3*A*a*b
^3*sin(c + d*x))/d + (4*A*a^3*b*sin(c + d*x))/d
```

3.243 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec(c+dx) dx$

3.243.1 Optimal result	2239
3.243.2 Mathematica [A] (verified)	2240
3.243.3 Rubi [A] (verified)	2240
3.243.4 Maple [A] (verified)	2244
3.243.5 Fricas [A] (verification not implemented)	2245
3.243.6 Sympy [F]	2246
3.243.7 Maxima [A] (verification not implemented)	2246
3.243.8 Giac [B] (verification not implemented)	2247
3.243.9 Mupad [B] (verification not implemented)	2248

3.243.1 Optimal result

Integrand size = 29, antiderivative size = 200

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{1}{8} (32a^3 Ab + 16aAb^3 + 8a^4 B + 24a^2 b^2 B + 3b^4 B) x + \frac{a^4 A \operatorname{arctanh}(\sin(c + dx))}{d}$$

$$+ \frac{b(34a^2 Ab + 4Ab^3 + 19a^3 B + 16ab^2 B) \sin(c + dx)}{6d}$$

$$+ \frac{b^2(32aAb + 26a^2 B + 9b^2 B) \cos(c + dx) \sin(c + dx)}{24d}$$

$$+ \frac{b(4Ab + 7aB)(a + b \cos(c + dx))^2 \sin(c + dx)}{12d} + \frac{bB(a + b \cos(c + dx))^3 \sin(c + dx)}{4d}$$

output

```
1/8*(32*A*a^3*b+16*A*a*b^3+8*B*a^4+24*B*a^2*b^2+3*B*b^4)*x+a^4*A*arctanh(sin(d*x+c))/d+1/6*b*(34*A*a^2*b+4*A*b^3+19*B*a^3+16*B*a*b^2)*sin(d*x+c)/d+1/24*b^2*(32*A*a*b+26*B*a^2+9*B*b^2)*cos(d*x+c)*sin(d*x+c)/d+1/12*b*(4*A*b+7*B*a)*(a+b*cos(d*x+c))^2*sin(d*x+c)/d+1/4*b*B*(a+b*cos(d*x+c))^3*sin(d*x+c)/d
```

3.243.2 Mathematica [A] (verified)

Time = 2.03 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.05

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{12(32a^3Ab + 16aAb^3 + 8a^4B + 24a^2b^2B + 3b^4B)(c + dx) - 96a^4A \log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{d}$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`output `(12*(32*a^3*A*b + 16*a*A*b^3 + 8*a^4*B + 24*a^2*b^2*B + 3*b^4*B)*(c + d*x) - 96*a^4*A*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 96*a^4*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 24*b*(24*a^2*A*b + 3*A*b^3 + 16*a^3*B + 12*a*b^2*B)*Sin[c + d*x] + 24*b^2*(4*a*A*b + 6*a^2*B + b^2*B)*Sin[2*(c + d*x)] + 8*b^3*(A*b + 4*a*B)*Sin[3*(c + d*x)] + 3*b^4*B*Sin[4*(c + d*x)])/(96*d)`**3.243.3 Rubi [A] (verified)**Time = 1.45 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$, Rules used = {3042, 3469, 3042, 3528, 3042, 3512, 3042, 3502, 27, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(c + dx)(a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx$$

$$\downarrow \text{3469}$$

$$\frac{1}{4} \int (a + b \cos(c + dx))^2 (4Aa^2 + b(4Ab + 7aB) \cos^2(c + dx) + (4Ba^2 + 8Aba + 3b^2B) \cos(c + dx)) \sec(c + dx) dx + \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d}$$

$$\downarrow \text{3042}$$

3.243. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$

$$\frac{1}{4} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (4Aa^2 + b(4Ab + 7aB) \sin(c + dx + \frac{\pi}{2})^2 + (4Ba^2 + 8Aba + 3b^2B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \downarrow 3528$$

$$\frac{1}{4} \left(\frac{1}{3} \int (a + b \cos(c + dx)) (12Aa^3 + b(26Ba^2 + 32Aba + 9b^2B) \cos^2(c + dx) + (12Ba^3 + 36Aba^2 + 23b^2Ba + 8Aa^2b) \cos(c + dx)) \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \downarrow 3042 \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \int (a + b \sin(c + dx + \frac{\pi}{2})) (12Aa^3 + b(26Ba^2 + 32Aba + 9b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (12Ba^3 + 36Aba^2 + 23b^2Ba + 8Aa^2b) \sin(c + dx + \frac{\pi}{2})) \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \downarrow 3512 \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (24Aa^4 + 4b(19Ba^3 + 34Aba^2 + 16b^2Ba + 4Ab^3) \cos^2(c + dx) + 3(8Ba^4 + 32Aba^3 + 24b^2Ba^2 + 16Aa^2b) \cos(c + dx)) \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \downarrow 3042 \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{24Aa^4 + 4b(19Ba^3 + 34Aba^2 + 16b^2Ba + 4Ab^3) \sin(c + dx + \frac{\pi}{2})^2 + 3(8Ba^4 + 32Aba^3 + 24b^2Ba^2 + 16Aa^2b) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \downarrow 3502 \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\int 3(8Aa^4 + (8Ba^4 + 32Aba^3 + 24b^2Ba^2 + 16Ab^3a + 3b^4B) \cos(c + dx)) \sec(c + dx) dx + \frac{4b(19a^3B + 3a^2bB + 2ab^2B + b^3B)}{4d} \right) \frac{bB \sin(c + dx)(a + b \cos(c + dx))^3}{4d} \downarrow 27 \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \int (8Aa^4 + (8Ba^4 + 32Aba^3 + 24b^2Ba^2 + 16Ab^3a + 3b^4B) \cos(c + dx)) \sec(c + dx) dx + \frac{4b(19a^3B + 34a^2Ab + 16ab^2B + 4Ab^3) \sin(c + dx)}{4d} \right) \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{8Aa^4 + (8Ba^4 + 32Aba^3 + 24b^2Ba^2 + 16Ab^3a + 3b^4B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{4b(19a^3B + 34a^2Ab + 16ab^2B + 4Ab^3) \cos(c + dx)}{4d} \right) \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \left(8a^4A \int \sec(c + dx) dx + x(8a^4B + 32a^3Ab + 24a^2b^2B + 16aAb^3 + 3b^4B) \right) + \frac{4b(19a^3B + 34a^2Ab + 16ab^2B + 4Ab^3) \sin(c + dx)}{4d} \right) \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \left(8a^4A \int \csc(c + dx + \frac{\pi}{2}) dx + x(8a^4B + 32a^3Ab + 24a^2b^2B + 16aAb^3 + 3b^4B) \right) + \frac{4b(19a^3B + 34a^2Ab + 16ab^2B + 4Ab^3) \cos(c + dx)}{4d} \right) \right) \right)$$

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{b^2(26a^2B + 32aAb + 9b^2B) \sin(c + dx) \cos(c + dx)}{2d} + \frac{1}{2} \left(\frac{4b(19a^3B + 34a^2Ab + 16ab^2B + 4Ab^3) \sin(c + dx)}{d} + \frac{4b(19a^3B + 34a^2Ab + 16ab^2B + 4Ab^3) \cos(c + dx)}{d} \right) \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(b*B*(a + b*Cos[c + d*x])^3*Sin[c + d*x])/(4*d) + ((b*(4*A*b + 7*a*B)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d) + ((b^2*(32*a*A*b + 26*a^2*B + 9*b^2*B)*Cos[c + d*x]*Sin[c + d*x])/(2*d) + (3*((32*a^3*A*b + 16*a*A*b^3 + 8*a^4*B + 24*a^2*b^2*B + 3*b^4*B)*x + (8*a^4*A*ArcTanh[Sin[c + d*x]])/d) + (4*b*(34*a^2*A*b + 4*A*b^3 + 19*a^3*B + 16*a*b^2*B)*Sin[c + d*x])/d)/2)/3)/4`

3.243.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2
, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x
])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.243.4 Maple [A] (verified)

Time = 3.08 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.96

method	result
parts	$\frac{a^4 A \ln(\sec(dx+c)+\tan(dx+c))}{d} + \frac{(A b^4+4B a b^3)(2+\cos^2(dx+c)) \sin(dx+c)}{3d} + \frac{(4A a b^3+6B a^2 b^2) \left(\frac{\cos(dx+c) \sin(dx+c)}{2}\right)}{d}$
parallelrisch	$\frac{-96a^4 A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 96a^4 A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) + 24(4A a b^3 + 6B a^2 b^2 + B b^4) \sin(2dx+2c) + 8(A b^4 + 4B a b^3)}{d}$
derivativdivides	$\frac{a^4 A \ln(\sec(dx+c)+\tan(dx+c)) + B a^4(dx+c) + 4A a^3 b(dx+c) + 4B \sin(dx+c) a^3 b + 6A \sin(dx+c) a^2 b^2 + 6B a^2 b^2 \left(\frac{\cos(dx+c)}{2}\right)}{d}$
default	$\frac{a^4 A \ln(\sec(dx+c)+\tan(dx+c)) + B a^4(dx+c) + 4A a^3 b(dx+c) + 4B \sin(dx+c) a^3 b + 6A \sin(dx+c) a^2 b^2 + 6B a^2 b^2 \left(\frac{\cos(dx+c)}{2}\right)}{d}$
risch	$4xA a^3 b + 2xA a b^3 + 3xB a^2 b^2 + a^4 B x + \frac{3b^4 B x}{8} + \frac{3ie^{-i(dx+c)} A a^2 b^2}{d} - \frac{3ie^{i(dx+c)} A a^2 b^2}{d} - \frac{3ie^{i(dx+c)} A a^2 b^2}{d}$
norman	$\frac{(4A a^3 b + 2A a b^3 + 3B a^2 b^2 + B a^4 + \frac{3}{8} B b^4) x + (4A a^3 b + 2A a b^3 + 3B a^2 b^2 + B a^4 + \frac{3}{8} B b^4) x \left(\tan^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (20A a^3 b + 12A a b^3 + 12B a^2 b^2 + 8B a^2 b^2)}{d}$

```
input int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)
```

```
output a^4*A/d*ln(sec(d*x+c)+tan(d*x+c))+1/3*(A*b^4+4*B*a*b^3)/d*(2+cos(d*x+c)^2)*sin(d*x+c)+(4*A*a*b^3+6*B*a^2*b^2)/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+(6*A*a^2*b^2+4*B*a^3*b)/d*sin(d*x+c)+(4*A*a^3*b+B*a^4)/d*(d*x+c)+B*b^4/d*(1/4*(cos(d*x+c)^3+3/2*cos(d*x+c))*sin(d*x+c)+3/8*d*x+3/8*c)
```

3.243.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.92

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{12 A a^4 \log(\sin(dx + c) + 1) - 12 A a^4 \log(-\sin(dx + c) + 1) + 3(8 B a^4 + 32 A a^3 b + 24 B a^2 b^2 + 16 A a b^3)}{d}$$

```
input integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")
```


3.243.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 603 vs. $2(190) = 380$.

Time = 0.33 (sec) , antiderivative size = 603, normalized size of antiderivative = 3.02

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{24 Aa^4 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right| \right) - 24 Aa^4 \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right| \right) + 3 (8 Ba^4 + 32 Aa^3b + 24 Ba^2b^2 + 16 Aa^2b^3 + 3 Bb^4) (dx + c) + 2 (96 B^2 a^3 b \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 144 A^2 a^2 b^2 \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 72 B^2 a^2 b^2 \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 48 A^2 a^2 b^3 \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 96 B^2 a^3 b^3 \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 24 A^2 a^4 b^4 \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 15 B^2 b^4 \tan^7 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 288 B^2 a^3 b \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 432 A^2 a^2 b^2 \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 72 B^2 a^2 b^2 \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 48 A^2 a^2 b^3 \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 160 B^2 a^3 b^3 \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 40 A^2 a^4 b^4 \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 9 B^2 b^4 \tan^5 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 288 B^2 a^3 b \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 432 A^2 a^2 b^2 \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 72 B^2 a^2 b^2 \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 48 A^2 a^2 b^3 \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 160 B^2 a^3 b^3 \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 40 A^2 a^4 b^4 \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 9 B^2 b^4 \tan^3 \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 96 B^2 a^3 b \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 144 A^2 a^2 b^2 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 72 B^2 a^2 b^2 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 48 A^2 a^2 b^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 96 B^2 a^3 b^3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 24 A^2 a^4 b^4 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 15 B^2 b^4 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{\left(\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 + 1 \right)^4} dx$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output $\frac{1}{24} (24 A^2 a^4 \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1) - 24 A^2 a^4 \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1)) + 3 (8 B^2 a^4 + 32 A^2 a^3 b + 24 B^2 a^2 b^2 + 16 A^2 a^2 b^3 + 3 B^2 b^4) (dx + c) + 2 (96 B^2 a^3 b \tan^7(\frac{1}{2} dx + \frac{1}{2} c) + 144 A^2 a^2 b^2 \tan^7(\frac{1}{2} dx + \frac{1}{2} c) - 72 B^2 a^2 b^2 \tan^7(\frac{1}{2} dx + \frac{1}{2} c) - 48 A^2 a^2 b^3 \tan^7(\frac{1}{2} dx + \frac{1}{2} c) + 96 B^2 a^3 b^3 \tan^7(\frac{1}{2} dx + \frac{1}{2} c) + 24 A^2 a^4 b^4 \tan^7(\frac{1}{2} dx + \frac{1}{2} c) - 15 B^2 b^4 \tan^7(\frac{1}{2} dx + \frac{1}{2} c) + 288 B^2 a^3 b \tan^5(\frac{1}{2} dx + \frac{1}{2} c) + 432 A^2 a^2 b^2 \tan^5(\frac{1}{2} dx + \frac{1}{2} c) - 72 B^2 a^2 b^2 \tan^5(\frac{1}{2} dx + \frac{1}{2} c) - 48 A^2 a^2 b^3 \tan^5(\frac{1}{2} dx + \frac{1}{2} c) + 160 B^2 a^3 b^3 \tan^5(\frac{1}{2} dx + \frac{1}{2} c) + 40 A^2 a^4 b^4 \tan^5(\frac{1}{2} dx + \frac{1}{2} c) + 9 B^2 b^4 \tan^5(\frac{1}{2} dx + \frac{1}{2} c) + 288 B^2 a^3 b \tan^3(\frac{1}{2} dx + \frac{1}{2} c) + 432 A^2 a^2 b^2 \tan^3(\frac{1}{2} dx + \frac{1}{2} c) + 72 B^2 a^2 b^2 \tan^3(\frac{1}{2} dx + \frac{1}{2} c) + 48 A^2 a^2 b^3 \tan^3(\frac{1}{2} dx + \frac{1}{2} c) + 160 B^2 a^3 b^3 \tan^3(\frac{1}{2} dx + \frac{1}{2} c) + 40 A^2 a^4 b^4 \tan^3(\frac{1}{2} dx + \frac{1}{2} c) - 9 B^2 b^4 \tan^3(\frac{1}{2} dx + \frac{1}{2} c) + 96 B^2 a^3 b \tan(\frac{1}{2} dx + \frac{1}{2} c) + 144 A^2 a^2 b^2 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 72 B^2 a^2 b^2 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 48 A^2 a^2 b^3 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 96 B^2 a^3 b^3 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 24 A^2 a^4 b^4 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 15 B^2 b^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)) / (\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)^4 dx$

3.243.9 Mupad [B] (verification not implemented)

Time = 1.56 (sec) , antiderivative size = 369, normalized size of antiderivative = 1.84

$$\begin{aligned}
& \int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec(c + dx) dx \\
&= \frac{3 A b^4 \sin(c + dx)}{4 d} + \frac{2 A a^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{2 B a^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{3 B b^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{4 d} + \frac{A b^4 \sin(3 c + 3 d x)}{12 d} + \frac{B b^4 \sin(2 c + 2 d x)}{4 d} \\
&+ \frac{B b^4 \sin(4 c + 4 d x)}{32 d} + \frac{4 A a b^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{8 A a^3 b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} + \frac{A a b^3 \sin(2 c + 2 d x)}{d} + \frac{6 A a^2 b^2 \sin(c + d x)}{d} \\
&+ \frac{B a b^3 \sin(3 c + 3 d x)}{3 d} + \frac{6 B a^2 b^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{d} \\
&+ \frac{3 B a^2 b^2 \sin(2 c + 2 d x)}{2 d} + \frac{3 B a b^3 \sin(c + d x)}{d} + \frac{4 B a^3 b \sin(c + d x)}{d}
\end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4)/cos(c + d*x),x)`output `(3*A*b^4*sin(c + d*x))/(4*d) + (2*A*a^4*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (2*B*a^4*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (3*B*b^4*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(4*d) + (A*b^4*sin(3*c + 3*d*x))/(12*d) + (B*b^4*sin(2*c + 2*d*x))/(4*d) + (B*b^4*sin(4*c + 4*d*x))/(32*d) + (4*A*a*b^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (8*A*a^3*b*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (A*a*b^3*sin(2*c + 2*d*x))/d + (6*A*a^2*b^2*sin(c + d*x))/d + (B*a*b^3*sin(3*c + 3*d*x))/(3*d) + (6*B*a^2*b^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/d + (3*B*a^2*b^2*sin(2*c + 2*d*x))/(2*d) + (3*B*a*b^3*sin(c + d*x))/d + (4*B*a^3*b*sin(c + d*x))/d`

3.244 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.244.1 Optimal result	2249
3.244.2 Mathematica [A] (verified)	2250
3.244.3 Rubi [A] (verified)	2250
3.244.4 Maple [A] (verified)	2254
3.244.5 Fricas [A] (verification not implemented)	2255
3.244.6 Sympy [F]	2256
3.244.7 Maxima [A] (verification not implemented)	2256
3.244.8 Giac [A] (verification not implemented)	2257
3.244.9 Mupad [B] (verification not implemented)	2257

3.244.1 Optimal result

Integrand size = 31, antiderivative size = 195

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{1}{2} b (12a^2 Ab + Ab^3 + 8a^3 B + 4ab^2 B) x + \frac{a^3 (4Ab + aB) \operatorname{arctanh}(\sin(c + dx))}{d}$$

$$- \frac{b(6a^3 A - 12aAb^2 - 17a^2 bB - 2b^3 B) \sin(c + dx)}{3d}$$

$$- \frac{b^2(6a^2 A - 3Ab^2 - 8abB) \cos(c + dx) \sin(c + dx)}{6d}$$

$$- \frac{b(3aA - bB)(a + b \cos(c + dx))^2 \sin(c + dx)}{3d} + \frac{aA(a + b \cos(c + dx))^3 \tan(c + dx)}{d}$$

output

```
1/2*b*(12*A*a^2*b+A*b^3+8*B*a^3+4*B*a*b^2)*x+a^3*(4*A*b+B*a)*arctanh(sin(d*x+c))/d-1/3*b*(6*A*a^3-12*A*a*b^2-17*B*a^2*b-2*B*b^3)*sin(d*x+c)/d-1/6*b^2*(6*A*a^2-3*A*b^2-8*B*a*b)*cos(d*x+c)*sin(d*x+c)/d-1/3*b*(3*A*a-B*b)*(a+b*cos(d*x+c))^2*sin(d*x+c)/d+a*A*(a+b*cos(d*x+c))^3*tan(d*x+c)/d
```


3.244.2 Mathematica [A] (verified)

Time = 2.61 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.32

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{6b(12a^2Ab + Ab^3 + 8a^3B + 4ab^2B)(c + dx) - 12a^3(4Ab + aB) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + \dots}{d}$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`output `(6*b*(12*a^2*A*b + A*b^3 + 8*a^3*B + 4*a*b^2*B)*(c + d*x) - 12*a^3*(4*A*b + a*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 12*a^3*(4*A*b + a*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (12*a^4*A*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]) + (12*a^4*A*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) + 3*b^2*(16*a*A*b + 24*a^2*B + 3*b^2*B)*Sin[c + d*x] + 3*b^3*(A*b + 4*a*B)*Sin[2*(c + d*x)] + b^4*B*Sin[3*(c + d*x)])/(12*d)`**3.244.3 Rubi [A] (verified)**Time = 1.42 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3468, 3042, 3528, 3042, 3512, 3042, 3502, 27, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(c + dx)(a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx$$

$$\downarrow \text{3468}$$

$$\int (a + b \cos(c + dx))^2 (-b(3aA - bB) \cos^2(c + dx) + b(Ab + 2aB) \cos(c + dx) + a(4Ab + aB)) \sec(c + dx) dx + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d}$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 \left(-b(3aA - bB) \sin(c + dx + \frac{\pi}{2})^2 + b(Ab + 2aB) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + aB) \right)}{\frac{\sin(c + dx + \frac{\pi}{2})}{d} \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d}}$$

↓ 3528

$$\frac{1}{3} \int (a + b \cos(c + dx)) \left(3(4Ab + aB)a^2 - b(6Aa^2 - 8bBa - 3Ab^2) \cos^2(c + dx) + b(9Ba^2 + 9Aba + 2b^2B) \cos(c + dx) \right) \sec(c + dx) dx - \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d}$$

↓ 3042

$$\frac{1}{3} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) \left(3(4Ab + aB)a^2 - b(6Aa^2 - 8bBa - 3Ab^2) \sin(c + dx + \frac{\pi}{2})^2 + b(9Ba^2 + 9Aba + 2b^2B) \sin(c + dx + \frac{\pi}{2}) \right)}{\frac{\sin(c + dx + \frac{\pi}{2})}{d} \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d}}$$

↓ 3512

$$\frac{1}{3} \left(\frac{1}{2} \int (6(4Ab + aB)a^3 - 2b(6Aa^3 - 17bBa^2 - 12Ab^2a - 2b^3B) \cos^2(c + dx) + 3b(8Ba^3 + 12Aba^2 + 4b^2Ba + Ab^3) \cos(c + dx)) \sec(c + dx) dx - \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \int \frac{6(4Ab + aB)a^3 - 2b(6Aa^3 - 17bBa^2 - 12Ab^2a - 2b^3B) \sin(c + dx + \frac{\pi}{2})^2 + 3b(8Ba^3 + 12Aba^2 + 4b^2Ba + Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right)$$

↓ 3502

$$\frac{1}{3} \left(\frac{1}{2} \left(\int 3(2(4Ab + aB)a^3 + b(8Ba^3 + 12Aba^2 + 4b^2Ba + Ab^3) \cos(c + dx)) \sec(c + dx) dx - \frac{2b(6a^3A - 17a^2bB - 12Ab^2a - 2b^3B)}{3d} \right) \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int (2(4Ab + aB)a^3 + b(8Ba^3 + 12Aba^2 + 4b^2Ba + Ab^3) \cos(c + dx)) \sec(c + dx) dx - \frac{2b(6a^3A - 17a^2bB)}{d} \right. \right. \\ \left. \left. \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{2(4Ab + aB)a^3 + b(8Ba^3 + 12Aba^2 + 4b^2Ba + Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{2b(6a^3A - 17a^2bB - 12aA)}{d} \right. \right. \\ \left. \left. \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right) \right)$$

↓ 3214

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(2a^3(aB + 4Ab) \int \sec(c + dx) dx + bx(8a^3B + 12a^2Ab + 4ab^2B + Ab^3) \right) - \frac{2b(6a^3A - 17a^2bB - 12aA)}{d} \right. \right. \\ \left. \left. \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(2a^3(aB + 4Ab) \int \csc(c + dx + \frac{\pi}{2}) dx + bx(8a^3B + 12a^2Ab + 4ab^2B + Ab^3) \right) - \frac{2b(6a^3A - 17a^2bB - 12aA)}{d} \right. \right. \\ \left. \left. \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right) \right)$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(\frac{2a^3(aB + 4Ab) \operatorname{arctanh}(\sin(c + dx))}{d} + bx(8a^3B + 12a^2Ab + 4ab^2B + Ab^3) \right) - \frac{2b(6a^3A - 17a^2bB - 12aA)}{d} \right. \right. \\ \left. \left. \frac{b(3aA - bB) \sin(c + dx)(a + b \cos(c + dx))^2}{3d} + \frac{aA \tan(c + dx)(a + b \cos(c + dx))^3}{d} \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `-1/3*(b*(3*a*A - b*B)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/d + (-1/2*(b^2*(6*a^2*A - 3*A*b^2 - 8*a*b*B)*Cos[c + d*x]*Sin[c + d*x])/d + (3*(b*(12*a^2*A*b + A*b^3 + 8*a^3*B + 4*a*b^2*B)*x + (2*a^3*(4*A*b + a*B)*ArcTanh[Sin[c + d*x]]))/d) - (2*b*(6*a^3*A - 12*a*A*b^2 - 17*a^2*b*B - 2*b^3*B)*Sin[c + d*x])/d)/2)/3 + (a*A*(a + b*Cos[c + d*x])^3*Tan[c + d*x])/d`

3.244.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3512 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Sin[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) + A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.244.4 Maple [A] (verified)

Time = 4.01 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.84

method	result
parts	$\frac{a^4 A \tan(dx+c)}{d} + \frac{(A b^4 + 4 B a b^3) \left(\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d} + \frac{(4 A a b^3 + 6 B a^2 b^2) \sin(dx+c)}{d} + \frac{(6 A a^2 b^2 + 4 B a^3 b)}{d}$
derivatividevides	$\frac{a^4 A \tan(dx+c) + B a^4 \ln(\sec(dx+c) + \tan(dx+c)) + 4 A a^3 b \ln(\sec(dx+c) + \tan(dx+c)) + 4 B a^3 b(dx+c) + 6 A a^2 b^2(dx+c) + \dots}{\dots}$
default	$a^4 A \tan(dx+c) + B a^4 \ln(\sec(dx+c) + \tan(dx+c)) + 4 A a^3 b \ln(\sec(dx+c) + \tan(dx+c)) + 4 B a^3 b(dx+c) + 6 A a^2 b^2(dx+c) + \dots$
parallelrisch	$\frac{-96 \cos(dx+c) \left(A b + \frac{B a}{4} \right) a^3 \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 96 \cos(dx+c) \left(A b + \frac{B a}{4} \right) a^3 \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) + (48 A a b^3 + 72 B a^2 b^2)}{\dots}$
risch	$6 x A a^2 b^2 + \frac{x A b^4}{2} + 4 x B a^3 b + 2 x B a b^3 - \frac{3 i e^{i(dx+c)} B a^2 b^2}{d} - \frac{i e^{2i(dx+c)} A b^4}{8 d} - \frac{i e^{2i(dx+c)} B a b^3}{2 d} - \dots$
norman	$\frac{(-6 A a^2 b^2 - \frac{1}{2} A b^4 - 4 B a^3 b - 2 B a b^3) x + (-30 A a^2 b^2 - \frac{5}{2} A b^4 - 20 B a^3 b - 10 B a b^3) x \left(\tan^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (6 A a^2 b^2 + \frac{1}{2} A b^4 + \dots)}{\dots}$

```
input int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNVERBOSE)
```

```
output a^4*A/d*tan(d*x+c)+(A*b^4+4*B*a*b^3)/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)+(4*A*a*b^3+6*B*a^2*b^2)/d*sin(d*x+c)+(6*A*a^2*b^2+4*B*a^3*b)/d*(d*x+c)+(4*A*a^3*b+B*a^4)/d*ln(sec(d*x+c)+tan(d*x+c))+1/3*B*b^4/d*(2+cos(d*x+c))^2*sin(d*x+c)
```

3.244.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.01

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{3(8 B a^3 b + 12 A a^2 b^2 + 4 B a b^3 + A b^4) dx \cos(dx + c) + 3(B a^4 + 4 A a^3 b) \cos(dx + c) \log(\sin(dx + c)) + \dots}{\dots}$$

```
input integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")
```

output $1/6*(3*(8*B*a^3*b + 12*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*d*x*cos(d*x + c) + 3*(B*a^4 + 4*A*a^3*b)*cos(d*x + c)*log(sin(d*x + c) + 1) - 3*(B*a^4 + 4*A*a^3*b)*cos(d*x + c)*log(-sin(d*x + c) + 1) + (2*B*b^4*cos(d*x + c)^3 + 6*A*a^4 + 3*(4*B*a*b^3 + A*b^4)*cos(d*x + c)^2 + 4*(9*B*a^2*b^2 + 6*A*a*b^3 + B*b^4)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c))$

3.244.6 Sympy [F]

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^4 \sec^2(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**4*sec(c + d*x)**2, x)`

3.244.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.01

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{48(dx + c)Ba^3b + 72(dx + c)Aa^2b^2 + 12(2dx + 2c + \sin(2dx + 2c))Bab^3 + 3(2dx + 2c + \sin(2dx + 2c))Aa^4 + 4(2dx + 2c + \sin(2dx + 2c))Aa^3b + 4(2dx + 2c + \sin(2dx + 2c))Aa^2b^2 + 4(2dx + 2c + \sin(2dx + 2c))Aab^3 + 4(2dx + 2c + \sin(2dx + 2c))Ab^4 + 6Bb^4(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 24Aa^3b(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 72Bb^4\sin(dx + c) + 48Aa^3b^3\sin(dx + c) + 12Aa^4\tan(dx + c))/d$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output $1/12*(48*(d*x + c)*B*a^3*b + 72*(d*x + c)*A*a^2*b^2 + 12*(2*d*x + 2*c + \sin(2*d*x + 2*c))*B*a*b^3 + 3*(2*d*x + 2*c + \sin(2*d*x + 2*c))*A*b^4 - 4*(\sin(d*x + c)^3 - 3*\sin(d*x + c))*B*b^4 + 6*B*a^4*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 24*A*a^3*b*(\log(\sin(d*x + c) + 1) - \log(\sin(d*x + c) - 1)) + 72*B*a^2*b^2*\sin(d*x + c) + 48*A*a*b^3*\sin(d*x + c) + 12*A*a^4*\tan(d*x + c))/d$

3.244.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.90

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$\frac{12 A a^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1} - 3 (8 B a^3 b + 12 A a^2 b^2 + 4 B a b^3 + A b^4) (dx + c) - 6 (B a^4 + 4 A a^3 b) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c)|)$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="giac")`

output `-1/6*(12*A*a^4*tan(1/2*d*x + 1/2*c)/(tan(1/2*d*x + 1/2*c)^2 - 1) - 3*(8*B*a^3*b + 12*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) - 6*(B*a^4 + 4*A*a^3*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) + 6*(B*a^4 + 4*A*a^3*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(36*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^5 + 24*A*a*b^3*tan(1/2*d*x + 1/2*c)^5 - 12*B*a*b^3*tan(1/2*d*x + 1/2*c)^5 - 3*A*b^4*tan(1/2*d*x + 1/2*c)^5 + 6*B*b^4*tan(1/2*d*x + 1/2*c)^5 + 72*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^3 + 48*A*a*b^3*tan(1/2*d*x + 1/2*c)^3 + 4*B*b^4*tan(1/2*d*x + 1/2*c)^3 + 36*B*a^2*b^2*tan(1/2*d*x + 1/2*c) + 24*A*a*b^3*tan(1/2*d*x + 1/2*c) + 12*B*a*b^3*tan(1/2*d*x + 1/2*c) + 3*A*b^4*tan(1/2*d*x + 1/2*c) + 6*B*b^4*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 + 1)^3/d`

3.244.9 Mupad [B] (verification not implemented)

Time = 2.46 (sec) , antiderivative size = 2522, normalized size of antiderivative = 12.93

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Too large to display}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4)/cos(c + d*x)^2,x)`

output

$$\begin{aligned}
& (\tan(c/2 + (d*x)/2)*(2*A*a^4 + A*b^4 + 2*B*b^4 + 12*B*a^2*b^2 + 8*A*a*b^3 \\
& + 4*B*a*b^3) + \tan(c/2 + (d*x)/2)^7*(2*A*a^4 + A*b^4 - 2*B*b^4 - 12*B*a^2* \\
& b^2 - 8*A*a*b^3 + 4*B*a*b^3) + \tan(c/2 + (d*x)/2)^3*(6*A*a^4 - A*b^4 - (2* \\
& B*b^4)/3 + 12*B*a^2*b^2 + 8*A*a*b^3 - 4*B*a*b^3) - \tan(c/2 + (d*x)/2)^5*(A \\
& *b^4 - 6*A*a^4 - (2*B*b^4)/3 + 12*B*a^2*b^2 + 8*A*a*b^3 + 4*B*a*b^3))/(d*(\\
& 2*\tan(c/2 + (d*x)/2)^2 - 2*\tan(c/2 + (d*x)/2)^6 - \tan(c/2 + (d*x)/2)^8 + 1 \\
&)) - (\operatorname{atan}(((B*a^4 + 4*A*a^3*b)*((B*a^4 + 4*A*a^3*b)*(16*A*b^4 + 32*B*a^4 \\
& + 192*A*a^2*b^2 + 128*A*a^3*b + 64*B*a*b^3 + 128*B*a^3*b) + \tan(c/2 + (d*x) \\
&)/2)*(8*A^2*b^8 + 32*B^2*a^8 + 192*A^2*a^2*b^6 + 1152*A^2*a^4*b^4 + 512*A^ \\
& 2*a^6*b^2 + 128*B^2*a^2*b^6 + 512*B^2*a^4*b^4 + 512*B^2*a^6*b^2 + 64*A*B*a \\
& *b^7 + 256*A*B*a^7*b + 896*A*B*a^3*b^5 + 1536*A*B*a^5*b^3))*1i - (B*a^4 + \\
& 4*A*a^3*b)*((B*a^4 + 4*A*a^3*b)*(16*A*b^4 + 32*B*a^4 + 192*A*a^2*b^2 + 128 \\
& *A*a^3*b + 64*B*a*b^3 + 128*B*a^3*b) - \tan(c/2 + (d*x)/2)*(8*A^2*b^8 + 32* \\
& B^2*a^8 + 192*A^2*a^2*b^6 + 1152*A^2*a^4*b^4 + 512*A^2*a^6*b^2 + 128*B^2*a \\
& ^2*b^6 + 512*B^2*a^4*b^4 + 512*B^2*a^6*b^2 + 64*A*B*a*b^7 + 256*A*B*a^7*b \\
& + 896*A*B*a^3*b^5 + 1536*A*B*a^5*b^3))*1i)/((B*a^4 + 4*A*a^3*b)*((B*a^4 + \\
& 4*A*a^3*b)*(16*A*b^4 + 32*B*a^4 + 192*A*a^2*b^2 + 128*A*a^3*b + 64*B*a*b^3 \\
& + 128*B*a^3*b) + \tan(c/2 + (d*x)/2)*(8*A^2*b^8 + 32*B^2*a^8 + 192*A^2*a^2 \\
& *b^6 + 1152*A^2*a^4*b^4 + 512*A^2*a^6*b^2 + 128*B^2*a^2*b^6 + 512*B^2*a^4* \\
& b^4 + 512*B^2*a^6*b^2 + 64*A*B*a*b^7 + 256*A*B*a^7*b + 896*A*B*a^3*b^5 \dots
\end{aligned}$$

3.245 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.245.1 Optimal result	2259
3.245.2 Mathematica [A] (verified)	2260
3.245.3 Rubi [A] (verified)	2260
3.245.4 Maple [A] (verified)	2264
3.245.5 Fricas [A] (verification not implemented)	2265
3.245.6 Sympy [F(-1)]	2266
3.245.7 Maxima [A] (verification not implemented)	2266
3.245.8 Giac [B] (verification not implemented)	2266
3.245.9 Mupad [B] (verification not implemented)	2267

3.245.1 Optimal result

Integrand size = 31, antiderivative size = 209

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{1}{2} b^2 (8aAb + 12a^2B + b^2B) x + \frac{a^2(a^2A + 12Ab^2 + 8abB) \operatorname{arctanh}(\sin(c + dx))}{2d}$$

$$- \frac{b(13a^2Ab - 2Ab^3 + 4a^3B - 8ab^2B) \sin(c + dx)}{2d}$$

$$- \frac{b^2(6aAb + 2a^2B - b^2B) \cos(c + dx) \sin(c + dx)}{2d}$$

$$+ \frac{a(5Ab + 2aB)(a + b \cos(c + dx))^2 \tan(c + dx)}{2d}$$

$$+ \frac{aA(a + b \cos(c + dx))^3 \sec(c + dx) \tan(c + dx)}{2d}$$

```
output 1/2*b^2*(8*A*a*b+12*B*a^2+B*b^2)*x+1/2*a^2*(A*a^2+12*A*b^2+8*B*a*b)*arctan
h(sin(d*x+c))/d-1/2*b*(13*A*a^2*b-2*A*b^3+4*B*a^3-8*B*a*b^2)*sin(d*x+c)/d-
1/2*b^2*(6*A*a*b+2*B*a^2-B*b^2)*cos(d*x+c)*sin(d*x+c)/d+1/2*a*(5*A*b+2*B*a
)*(a+b*cos(d*x+c))^2*tan(d*x+c)/d+1/2*a*A*(a+b*cos(d*x+c))^3*sec(d*x+c)*ta
n(d*x+c)/d
```

3.245.2 Mathematica [A] (verified)

Time = 3.24 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.48

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2b^2(8aAb + 12a^2B + b^2B)(c + dx) - 2a^2(a^2A + 12Ab^2 + 8abB) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right)}{4d}$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(2*b^2*(8*a*A*b + 12*a^2*B + b^2*B)*(c + d*x) - 2*a^2*(a^2*A + 12*A*b^2 + 8*a*b*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 2*a^2*(a^2*A + 12*A*b^2 + 8*a*b*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (a^4*A)/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2 + (4*a^3*(4*A*b + a*B)*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]) - (a^4*A)/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 + (4*a^3*(4*A*b + a*B)*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) + 4*b^3*(A*b + 4*a*B)*Sin[c + d*x] + b^4*B*Sin[2*(c + d*x)])/(4*d)`

3.245.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 201, normalized size of antiderivative = 0.96, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3468, 3042, 3526, 3042, 3512, 27, 3042, 3502, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^3(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{2} \int (a + b \cos(c + dx))^2 (-2b(aA - bB) \cos^2(c + dx) + (Aa^2 + 4bBa + 2Ab^2) \cos(c + dx) + a(5Ab + 2aB)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

↓ 3042

$$\frac{1}{2} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (-2b(aA - bB) \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 4bBa + 2Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(5Ab + 2aB)) \sec^2(c + dx + \frac{\pi}{2})}{\sin^2(c + dx + \frac{\pi}{2})} dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

↓ 3526

$$\frac{1}{2} \left(\int (a + b \cos(c + dx)) (-2b(2Ba^2 + 6Aba - b^2B) \cos^2(c + dx) - b(Aa^2 - 6bBa - 2Ab^2) \cos(c + dx) + a(Aa^2 - 6bBa - 2Ab^2)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\int (a + b \sin(c + dx + \frac{\pi}{2})) (-2b(2Ba^2 + 6Aba - b^2B) \sin^2(c + dx + \frac{\pi}{2}) - b(Aa^2 - 6bBa - 2Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(Aa^2 - 6bBa - 2Ab^2)) \sec^2(c + dx + \frac{\pi}{2}) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d} \right)$$

↓ 3512

$$\frac{1}{2} \left(\frac{1}{2} \int 2((Aa^2 + 8bBa + 12Ab^2) a^2 - b(4Ba^3 + 13Aba^2 - 8b^2Ba - 2Ab^3) \cos^2(c + dx) + b^2(12Ba^2 + 8Aba + b^2B)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d} \right)$$

↓ 27

$$\frac{1}{2} \left(\int ((Aa^2 + 8bBa + 12Ab^2) a^2 - b(4Ba^3 + 13Aba^2 - 8b^2Ba - 2Ab^3) \cos^2(c + dx) + b^2(12Ba^2 + 8Aba + b^2B)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{(Aa^2 + 8bBa + 12Ab^2) a^2 - b(4Ba^3 + 13Aba^2 - 8b^2Ba - 2Ab^3) \sin(c + dx + \frac{\pi}{2})^2 + b^2(12Ba^2 + 8Aba + b^2B)}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d} \right)$$

↓ 3502

$$\frac{1}{2} \left(\int ((Aa^2 + 8bBa + 12Ab^2) a^2 + b^2(12Ba^2 + 8Aba + b^2B) \cos(c + dx)) \sec(c + dx) dx - \frac{b^2(2a^2B + 6aAb - b^2B)}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{(Aa^2 + 8bBa + 12Ab^2) a^2 + b^2(12Ba^2 + 8Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{b^2(2a^2B + 6aAb - b^2B) \sin(c + dx)}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

↓ 3214

$$\frac{1}{2} \left(a^2(a^2A + 8abB + 12Ab^2) \int \sec(c + dx) dx - \frac{b^2(2a^2B + 6aAb - b^2B) \sin(c + dx) \cos(c + dx)}{d} + b^2x(12a^2B + 8Ab^2) \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

↓ 3042

$$\frac{1}{2} \left(a^2(a^2A + 8abB + 12Ab^2) \int \csc(c + dx + \frac{\pi}{2}) dx - \frac{b^2(2a^2B + 6aAb - b^2B) \sin(c + dx) \cos(c + dx)}{d} + b^2x(12a^2B + 8Ab^2) \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

↓ 4257

$$\frac{1}{2} \left(\frac{a^2(a^2A + 8abB + 12Ab^2) \operatorname{arctanh}(\sin(c + dx))}{d} - \frac{b^2(2a^2B + 6aAb - b^2B) \sin(c + dx) \cos(c + dx)}{d} + b^2x(12a^2B + 8Ab^2) \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^3}{2d}$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

```
output (a*A*(a + b*Cos[c + d*x])^3*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (b^2*(8*a*A
*b + 12*a^2*B + b^2*B)*x + (a^2*(a^2*A + 12*A*b^2 + 8*a*b*B)*ArcTanh[Sin[c
+ d*x]])/d - (b*(13*a^2*A*b - 2*A*b^3 + 4*a^3*B - 8*a*b^2*B)*Sin[c + d*x]
)/d - (b^2*(6*a*A*b + 2*a^2*B - b^2*B)*Cos[c + d*x]*Sin[c + d*x])/d + (a*(
5*A*b + 2*a*B)*(a + b*Cos[c + d*x])^2*Tan[c + d*x])/d)/2
```

3.245.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3214 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_
)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d
*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3468 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c
+ d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n +
1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n
+ 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a
*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1
- a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c -
a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2
, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2
, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

```
rule 3526 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x
]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 -
d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m -
1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*
d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n +
1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x
] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f
*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d
, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.245.4 Maple [A] (verified)

Time = 3.85 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.85

method	result
parts	$a^4 A \left(\frac{\sec(dx+c) \tan(dx+c) + \ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + \frac{(A b^4 + 4B a b^3) \sin(dx+c)}{d} + \frac{(4A a b^3 + 6B a^2 b^2)(dx+c)}{d} + \dots$
derivatividedivides	$a^4 A \left(\frac{\sec(dx+c) \tan(dx+c) + \ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + B a^4 \tan(dx+c) + 4A a^3 b \tan(dx+c) + 4B a^3 b \ln(\sec(dx+c) + \tan(dx+c)) + \dots$
default	$a^4 A \left(\frac{\sec(dx+c) \tan(dx+c) + \ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + B a^4 \tan(dx+c) + 4A a^3 b \tan(dx+c) + 4B a^3 b \ln(\sec(dx+c) + \tan(dx+c)) + \dots$
parallelrisch	$-4a^2(1+\cos(2dx+2c))(A a^2+12A b^2+8Bab) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right) + 4a^2(1+\cos(2dx+2c))(A a^2+12A b^2+8Bab) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right) + \dots$
risch	$4x A a b^3 + 6x B a^2 b^2 + \frac{b^4 B x}{2} - \frac{i e^{i(dx+c)} A b^4}{2d} - \frac{2i e^{i(dx+c)} B a b^3}{d} + \frac{i e^{-i(dx+c)} A b^4}{2d} + \frac{i B b^4 e^{-2i(dx+c)}}{8d} + \dots$
norman	$\frac{(4A a b^3 + 6B a^2 b^2 + \frac{1}{2} B b^4)x + (-20A a b^3 - 30B a^2 b^2 - \frac{5}{2} B b^4)x \left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-20A a b^3 - 30B a^2 b^2 - \frac{5}{2} B b^4)x \left(\tan^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\dots}$

input `int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `a^4*A/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^4+4*B*a*b^3)/d*sin(d*x+c)+(4*A*a*b^3+6*B*a^2*b^2)/d*(d*x+c)+(6*A*a^2*b^2+4*B*a^3*b)/d*ln(sec(d*x+c)+tan(d*x+c))+(4*A*a^3*b+B*a^4)/d*tan(d*x+c)+B*b^4/d*(1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)`

3.245.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 202, normalized size of antiderivative = 0.97

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{2(12 B a^2 b^2 + 8 A a b^3 + B b^4) dx \cos(dx + c)^2 + (A a^4 + 8 B a^3 b + 12 A a^2 b^2) \cos(dx + c)^2 \log(\sin(dx + c)) - \dots}{\dots}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fracas")`

output `1/4*(2*(12*B*a^2*b^2 + 8*A*a*b^3 + B*b^4)*d*x*cos(d*x + c)^2 + (A*a^4 + 8*B*a^3*b + 12*A*a^2*b^2)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A*a^4 + 8*B*a^3*b + 12*A*a^2*b^2)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(B*b^4*cos(d*x + c)^3 + A*a^4 + 2*(4*B*a*b^3 + A*b^4)*cos(d*x + c)^2 + 2*(B*a^4 + 4*A*a^3*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.245. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.245.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Timed out`

3.245.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.00

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{24(dx+c)Ba^2b^2 + 16(dx+c)Aab^3 + (2dx+2c+\sin(2dx+2c))Bb^4 - Aa^4 \left(\frac{2\sin(dx+c)}{\sin(dx+c)^2-1} - \log(\sin(dx+c)) \right)}{1}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `1/4*(24*(d*x + c)*B*a^2*b^2 + 16*(d*x + c)*A*a*b^3 + (2*d*x + 2*c + sin(2*d*x + 2*c))*B*b^4 - A*a^4*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 8*B*a^3*b*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 12*A*a^2*b^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 16*B*a*b^3*sin(d*x + c) + 4*A*b^4*sin(d*x + c) + 4*B*a^4*tan(d*x + c) + 16*A*a^3*b*tan(d*x + c))/d`

3.245.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 526 vs. 2(197) = 394.

Time = 0.38 (sec) , antiderivative size = 526, normalized size of antiderivative = 2.52

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{(12Ba^2b^2 + 8Aab^3 + Bb^4)(dx+c) + (Aa^4 + 8Ba^3b + 12Aa^2b^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - (Aa^4 + 8Ba^3b + 12Aa^2b^2) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 1\right|\right)}{1}$$

3.245. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="giac")`

output
$$\frac{1}{2} \left((12B^2a^2b^2 + 8A^2ab^3 + B^2b^4)(dx + c) + (A^4a^4 + 8B^3a^3b + 12A^2a^2b^2) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1) - (A^4a^4 + 8B^3a^3b + 12A^2a^2b^2) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) + 2(A^4a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 2B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - 8A^3a^3b \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 8B^2a^2b^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 2A^2ab^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 - B^2b^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 3A^4a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 2B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 8A^3a^3b \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 8B^2a^2b^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 2A^2ab^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 3B^2b^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 + 3A^4a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 2B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 8A^3a^3b \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 8B^2a^2b^3 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 2A^2ab^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 3B^2b^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + A^4a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 2B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 8A^3a^3b \tan(\frac{1}{2}dx + \frac{1}{2}c) + 8B^2a^2b^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 2A^2ab^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + B^2b^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) \right) / (\tan(\frac{1}{2}dx + \frac{1}{2}c)^4 - 1)^2 / d$$

3.245.9 Mupad [B] (verification not implemented)

Time = 2.65 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.58

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= 2 \left(\frac{Aa^4 \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{2} + \frac{Bb^4 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{2} + 4Aab^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) + 4Ba^3b \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \right) + \frac{Ba^4 \sin(2c+2dx)}{2} + \frac{Ab^4 \sin(3c+3dx)}{4} + \frac{Bb^4 \sin(2c+2dx)}{8} + \frac{Bb^4 \sin(4c+4dx)}{16} + \frac{Aa^4 \sin(c+dx)}{2} + \frac{Ab^4 \sin(c+dx)}{4} + Ba^3b \frac{d}{d \left(\frac{\cos(2c+2dx)}{2} + \frac{1}{2} \right)}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4)/cos(c + d*x)^3,x)`

output $(2*((A*a^4*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 + (B*b^4*atan(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 + 4*A*a*b^3*atan(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + 4*B*a^3*b*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + 6*A*a^2*b^2*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)) + 6*B*a^2*b^2*atan(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/d + ((B*a^4*\sin(2*c + 2*d*x))/2 + (A*b^4*\sin(3*c + 3*d*x))/4 + (B*b^4*\sin(2*c + 2*d*x))/8 + (B*b^4*\sin(4*c + 4*d*x))/16 + (A*a^4*\sin(c + d*x))/2 + (A*b^4*\sin(c + d*x))/4 + B*a*b^3*\sin(c + d*x) + 2*A*a^3*b*\sin(2*c + 2*d*x) + B*a*b^3*\sin(3*c + 3*d*x))/(d*(\cos(2*c + 2*d*x)/2 + 1/2))$

3.246 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.246.1 Optimal result	2269
3.246.2 Mathematica [B] (verified)	2270
3.246.3 Rubi [A] (verified)	2270
3.246.4 Maple [A] (verified)	2275
3.246.5 Fricas [A] (verification not implemented)	2276
3.246.6 Sympy [F(-1)]	2276
3.246.7 Maxima [A] (verification not implemented)	2277
3.246.8 Giac [B] (verification not implemented)	2277
3.246.9 Mupad [B] (verification not implemented)	2278

3.246.1 Optimal result

Integrand size = 31, antiderivative size = 198

$$\begin{aligned} & \int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= b^3 (Ab + 4aB)x + \frac{a(4a^2 Ab + 8Ab^3 + a^3 B + 12ab^2 B) \operatorname{arctanh}(\sin(c + dx))}{2d} \\ & \quad - \frac{b^2(8aAb + 3a^2 B - 6b^2 B) \sin(c + dx)}{6d} + \frac{a^2(2a^2 A + 9Ab^2 + 9abB) \tan(c + dx)}{3d} \\ & \quad + \frac{a(2Ab + aB)(a + b \cos(c + dx))^2 \sec(c + dx) \tan(c + dx)}{2d} \\ & \quad + \frac{aA(a + b \cos(c + dx))^3 \sec^2(c + dx) \tan(c + dx)}{3d} \end{aligned}$$

output

```
b^3*(A*b+4*B*a)*x+1/2*a*(4*A*a^2*b+8*A*b^3+B*a^3+12*B*a*b^2)*arctanh(sin(d*x+c))/d-1/6*b^2*(8*A*a*b+3*B*a^2-6*B*b^2)*sin(d*x+c)/d+1/3*a^2*(2*A*a^2+9*A*b^2+9*B*a*b)*tan(d*x+c)/d+1/2*a*(2*A*b+B*a)*(a+b*cos(d*x+c))^2*sec(d*x+c)*tan(d*x+c)/d+1/3*a*A*(a+b*cos(d*x+c))^3*sec(d*x+c)^2*tan(d*x+c)/d
```

3.246.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 415 vs. $2(198) = 396$.

Time = 6.80 (sec) , antiderivative size = 415, normalized size of antiderivative = 2.10

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12b^3(Ab + 4aB)(c + dx) - 6a(4a^2Ab + 8Ab^3 + a^3B + 12ab^2B) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + \dots}{\dots}$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output

$$\frac{(12*b^3*(A*b + 4*a*B)*(c + d*x) - 6*a*(4*a^2*A*b + 8*A*b^3 + a^3*B + 12*a*b^2*B)*\text{Log}[\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]] + 6*a*(4*a^2*A*b + 8*A*b^3 + a^3*B + 12*a*b^2*B)*\text{Log}[\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]] + (a^3*(12*A*b + a*(A + 3*B)))/(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2])^2 + (2*a^4*A*\text{Sin}[(c + d*x)/2])/(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2])^3 + (8*a^2*(a^2*A + 9*A*b^2 + 6*a*b*B)*\text{Sin}[(c + d*x)/2])/(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]) + (2*a^4*A*\text{Sin}[(c + d*x)/2])/(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2])^3 - (a^3*(12*A*b + a*(A + 3*B)))/(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2])^2 + (8*a^2*(a^2*A + 9*A*b^2 + 6*a*b*B)*\text{Sin}[(c + d*x)/2])/(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]) + 12*b^4*B*\text{Sin}[c + d*x])/(12*d)}$$

3.246.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3468, 3042, 3526, 3042, 3510, 25, 3042, 3502, 27, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c + dx)(a + b \cos(c + dx))^4(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx$$

$$\downarrow \text{3468}$$

3.246. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

$$\frac{1}{3} \int (a + b \cos(c + dx))^2 (-b(aA - 3bB) \cos^2(c + dx) + (2Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(2Ab + aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d}$$

↓ 3042

$$\frac{1}{3} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (-b(aA - 3bB) \sin(c + dx + \frac{\pi}{2})^2 + (2Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + 3a(2Ab + aB)) \sec^3(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d}$$

↓ 3526

$$\frac{1}{3} \left(\frac{1}{2} \int (a + b \cos(c + dx)) (-b(3Ba^2 + 8Aba - 6b^2B) \cos^2(c + dx) + (3Ba^3 + 8Aba^2 + 18b^2Ba + 6Ab^3) \cos(c + dx) + 3a(2Ab + aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \int (a + b \sin(c + dx + \frac{\pi}{2})) (-b(3Ba^2 + 8Aba - 6b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (3Ba^3 + 8Aba^2 + 18b^2Ba + 6Ab^3) \sin(c + dx + \frac{\pi}{2}) + 3a(2Ab + aB)) \sec^3(c + dx + \frac{\pi}{2}) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d} \right)$$

↓ 3510

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{2a^2(2a^2A + 9abB + 9Ab^2) \tan(c + dx)}{d} - \int -((6(Ab + 4aB) \cos(c + dx)b^3 - (3Ba^2 + 8Aba - 6b^2B) \cos^2(c + dx))b^2 + 3a(Ba^3 + 4Aba^2 + 12b^2Ba + 6Ab^3) \cos(c + dx) + 3a(2Ab + aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d} \right) \right)$$

↓ 25

$$\frac{1}{3} \left(\frac{1}{2} \left(\int (6(Ab + 4aB) \cos(c + dx)b^3 - (3Ba^2 + 8Aba - 6b^2B) \cos^2(c + dx)b^2 + 3a(Ba^3 + 4Aba^2 + 12b^2Ba + 6Ab^3) \cos(c + dx) + 3a(2Ab + aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(\int \frac{6(Ab + 4aB) \sin(c + dx + \frac{\pi}{2}) b^3 - (3Ba^2 + 8Aba - 6b^2B) \sin(c + dx + \frac{\pi}{2})^2 b^2 + 3a(Ba^3 + 4Aba^2 + 12b^2Ba + 8Ab^3)}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d} \right) \right)$$

↓ 3502

$$\frac{1}{3} \left(\frac{1}{2} \left(\int 3(2(Ab + 4aB) \cos(c + dx)b^3 + a(Ba^3 + 4Aba^2 + 12b^2Ba + 8Ab^3)) \sec(c + dx) dx - \frac{b^2(3a^2B + 8aAb - 6b^2B)}{d} \right) \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int (2(Ab + 4aB) \cos(c + dx)b^3 + a(Ba^3 + 4Aba^2 + 12b^2Ba + 8Ab^3)) \sec(c + dx) dx - \frac{b^2(3a^2B + 8aAb - 6b^2B)}{d} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{2(Ab + 4aB) \sin(c + dx + \frac{\pi}{2}) b^3 + a(Ba^3 + 4Aba^2 + 12b^2Ba + 8Ab^3)}{\sin(c + dx + \frac{\pi}{2})} dx - \frac{b^2(3a^2B + 8aAb - 6b^2B)}{d} \right) \right)$$

↓ 3214

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(a(a^3B + 4a^2Ab + 12ab^2B + 8Ab^3) \int \sec(c + dx) dx + 2b^3x(4aB + Ab) \right) - \frac{b^2(3a^2B + 8aAb - 6b^2B)}{d} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(3 \left(a(a^3B + 4a^2Ab + 12ab^2B + 8Ab^3) \int \csc(c + dx + \frac{\pi}{2}) dx + 2b^3x(4aB + Ab) \right) - \frac{b^2(3a^2B + 8aAb - 6b^2B)}{d} \right) \right)$$

↓ 4257

$$\frac{1}{3} \left(\frac{1}{2} \left(-\frac{b^2(3a^2B + 8aAb - 6b^2B) \sin(c + dx)}{d} + \frac{2a^2(2a^2A + 9abB + 9Ab^2) \tan(c + dx)}{d} \right) + 3 \left(\frac{a(a^3B + 4a^2Ab + aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^3}{3d} \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*(a + b*Cos[c + d*x])^3*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((3*a*(2*A*b + a*B)*(a + b*Cos[c + d*x])^2*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (3*(2*b^3*(A*b + 4*a*B)*x + (a*(4*a^2*A*b + 8*A*b^3 + a^3*B + 12*a*b^2*B)*ArcTan[Sin[c + d*x]])/d) - (b^2*(8*a*A*b + 3*a^2*B - 6*b^2*B)*Sin[c + d*x])/d + (2*a^2*(2*a^2*A + 9*A*b^2 + 9*a*b*B)*Tan[c + d*x])/d)/2)/3`

3.246.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(b*c - a*d))*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

```
rule 3526 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 -
d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m -
1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*
d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n +
1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x
] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f
*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d
, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.246.4 Maple [A] (verified)

Time = 5.03 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.88

method	result
parts	$-\frac{a^4 A \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c)}{d} + \frac{(A b^4 + 4 B a b^3)(dx+c)}{d} + \frac{(4 A a b^3 + 6 B a^2 b^2) \ln(\sec(dx+c) + \tan(dx+c))}{d}$
derivativedivides	$-a^4 A \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + B a^4 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 4 A a^3 b \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)$
default	$-a^4 A \left(-\frac{2}{3} - \frac{\sec^2(dx+c)}{3} \right) \tan(dx+c) + B a^4 \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c) + \tan(dx+c))}{2} \right) + 4 A a^3 b \left(\frac{\sec(dx+c) \tan(dx+c)}{2} \right)$
parallelrisch	$-36 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) a (A a^2 b + 2 A b^3 + \frac{1}{4} B a^3 + 3 B a b^2) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 36 \left(\frac{\cos(3dx+3c)}{3} + \cos(dx+c) \right) a (A a^2 b + 2 A b^3 + \frac{1}{4} B a^3 + 3 B a b^2)$
risch	$x A b^4 + 4 x B a b^3 - \frac{i e^{i(dx+c)} B b^4}{2d} + \frac{i e^{-i(dx+c)} B b^4}{2d} - \frac{i a^2 (12 A a b e^{5i(dx+c)} + 3 B a^2 e^{5i(dx+c)} - 36 A b^2 e^{4i(dx+c)})}{2d}$
norman	$\frac{(-A b^4 - 4 B a b^3) x + (-6 A b^4 - 24 B a b^3) x \left(\tan^{10} \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (-2 A b^4 - 8 B a b^3) x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + (-2 A b^4 - 8 B a b^3) x \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{1}$

```
input int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNVERBO
SE)
```

3.246. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$

output
$$-a^4 A/d*(-2/3-1/3*\sec(dx+c)^2)*\tan(dx+c)+(A*b^4+4*B*a*b^3)/d*(dx+c)+(4*A*a*b^3+6*B*a^2*b^2)/d*\ln(\sec(dx+c)+\tan(dx+c))+(6*A*a^2*b^2+4*B*a^3*b)/d*\tan(dx+c)+(4*A*a^3*b+B*a^4)/d*(1/2*\sec(dx+c)*\tan(dx+c)+1/2*\ln(\sec(dx+c)+\tan(dx+c)))+1/d*\sin(dx+c)*B*b^4$$

3.246.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.11

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12(4Bab^3 + Ab^4)dx \cos(dx + c)^3 + 3(Ba^4 + 4Aa^3b + 12Ba^2b^2 + 8Aab^3) \cos(dx + c)^3 \log(\sin(dx + c))}{1}$$

input `integrate((a+b*cos(dx+c))^4*(A+B*cos(dx+c))*sec(dx+c)^4,x, algorithm="fracas")`

output
$$1/12*(12*(4*B*a*b^3 + A*b^4)*d*x*\cos(dx + c)^3 + 3*(B*a^4 + 4*A*a^3*b + 12*B*a^2*b^2 + 8*A*a*b^3)*\cos(dx + c)^3*\log(\sin(dx + c) + 1) - 3*(B*a^4 + 4*A*a^3*b + 12*B*a^2*b^2 + 8*A*a*b^3)*\cos(dx + c)^3*\log(-\sin(dx + c) + 1) + 2*(6*B*b^4*\cos(dx + c)^3 + 2*A*a^4 + 4*(A*a^4 + 6*B*a^3*b + 9*A*a^2*b^2)*\cos(dx + c)^2 + 3*(B*a^4 + 4*A*a^3*b)*\cos(dx + c))*\sin(dx + c))/d*\cos(dx + c)^3$$

3.246.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(dx+c))**4*(A+B*cos(dx+c))*sec(dx+c)**4,x)`

output Timed out

3.246.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.24

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{4 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^4 + 48(dx + c) Bab^3 + 12(dx + c) Ab^4 - 3Ba^4 \left(\frac{2 \sin(dx+c)}{\sin(dx+c)^2 - 1} - \log(\sin(dx+c) + 1) + \log(\sin(dx+c) - 1) \right)}{d}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output `1/12*(4*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^4 + 48*(d*x + c)*B*a*b^3 + 12*(d*x + c)*A*b^4 - 3*B*a^4*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 12*A*a^3*b*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 36*B*a^2*b^2*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 24*A*a*b^3*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 12*B*b^4*sin(d*x + c) + 48*B*a^3*b*tan(d*x + c) + 72*A*a^2*b^2*tan(d*x + c))/d`

3.246.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 387 vs. 2(188) = 376.

Time = 0.35 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.95

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{12 B b^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1} + 6 (4 B a b^3 + A b^4) (dx + c) + 3 (B a^4 + 4 A a^3 b + 12 B a^2 b^2 + 8 A a b^3) \log \left(\left| \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) \right| \right)$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="giac")`

output $\frac{1}{6}*(12*B*b^4*\tan(1/2*d*x + 1/2*c)/(\tan(1/2*d*x + 1/2*c)^2 + 1) + 6*(4*B*a*b^3 + A*b^4)*(d*x + c) + 3*(B*a^4 + 4*A*a^3*b + 12*B*a^2*b^2 + 8*A*a*b^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) + 1)) - 3*(B*a^4 + 4*A*a^3*b + 12*B*a^2*b^2 + 8*A*a*b^3)*\log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 1)) - 2*(6*A*a^4*\tan(1/2*d*x + 1/2*c)^5 - 3*B*a^4*\tan(1/2*d*x + 1/2*c)^5 - 12*A*a^3*b*\tan(1/2*d*x + 1/2*c)^5 + 24*B*a^3*b*\tan(1/2*d*x + 1/2*c)^5 + 36*A*a^2*b^2*\tan(1/2*d*x + 1/2*c)^5 - 4*A*a^4*\tan(1/2*d*x + 1/2*c)^3 - 48*B*a^3*b*\tan(1/2*d*x + 1/2*c)^3 - 72*A*a^2*b^2*\tan(1/2*d*x + 1/2*c)^3 + 6*A*a^4*\tan(1/2*d*x + 1/2*c) + 3*B*a^4*\tan(1/2*d*x + 1/2*c) + 12*A*a^3*b*\tan(1/2*d*x + 1/2*c) + 24*B*a^3*b*\tan(1/2*d*x + 1/2*c) + 36*A*a^2*b^2*\tan(1/2*d*x + 1/2*c))/(\tan(1/2*d*x + 1/2*c)^2 - 1)^3)/d$

3.246.9 Mupad [B] (verification not implemented)

Time = 3.01 (sec) , antiderivative size = 636, normalized size of antiderivative = 3.21

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{A a^4 \sin(3c+3dx)}{6} + \frac{B a^4 \sin(2c+2dx)}{4} + \frac{B b^4 \sin(2c+2dx)}{4} + \frac{B b^4 \sin(4c+4dx)}{8} + \frac{A a^4 \sin(c+dx)}{2} + B a^3 b \sin(c + dx) +$$

input $\text{int}(((A + B*\cos(c + d*x))*(a + b*\cos(c + d*x))^4)/\cos(c + d*x)^4,x)$

output $((A*a^4*\sin(3*c + 3*d*x))/6 + (B*a^4*\sin(2*c + 2*d*x))/4 + (B*b^4*\sin(2*c + 2*d*x))/4 + (B*b^4*\sin(4*c + 4*d*x))/8 + (A*a^4*\sin(c + d*x))/2 + B*a^3*b*\sin(c + d*x) + (3*A*b^4*\cos(c + d*x)*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 - (B*a^4*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*3i)/4 + A*a^3*b*\sin(2*c + 2*d*x) + (3*A*a^2*b^2*\sin(c + d*x))/2 + B*a^3*b*\sin(3*c + 3*d*x) + (A*b^4*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x))/2 - (B*a^4*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*1i)/4 + (3*A*a^2*b^2*\sin(3*c + 3*d*x))/2 - A*a*b^3*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*2i - A*a^3*b*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*1i + 2*B*a*b^3*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x) - B*a^2*b^2*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*9i - B*a^2*b^2*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*\cos(3*c + 3*d*x)*3i - A*a*b^3*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*6i - A*a^3*b*\cos(c + d*x)*\operatorname{atan}((\sin(c/2 + (d*x)/2)*1i)/\cos(c/2 + (d*x)/2))*3i + 6*B*a*b^3*\cos(c + d*x)*\operatorname{atan}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/(d*((3*\cos(c + d*x))/4 + \cos(3*c + 3*d*x)/4))$

3.247 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.247.1 Optimal result	2280
3.247.2 Mathematica [A] (verified)	2281
3.247.3 Rubi [A] (verified)	2281
3.247.4 Maple [A] (verified)	2286
3.247.5 Fricas [A] (verification not implemented)	2286
3.247.6 Sympy [F(-1)]	2287
3.247.7 Maxima [A] (verification not implemented)	2287
3.247.8 Giac [B] (verification not implemented)	2288
3.247.9 Mupad [B] (verification not implemented)	2289

3.247.1 Optimal result

Integrand size = 31, antiderivative size = 216

$$\begin{aligned} & \int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx \\ &= b^4 B x + \frac{(3a^4 A + 24a^2 A b^2 + 8A b^4 + 16a^3 b B + 32a b^3 B) \operatorname{arctanh}(\sin(c + dx))}{8d} \\ & \quad + \frac{a(16a^2 A b + 19A b^3 + 4a^3 B + 34a b^2 B) \tan(c + dx)}{6d} \\ & \quad + \frac{a^2(9a^2 A + 26A b^2 + 32a b B) \sec(c + dx) \tan(c + dx)}{24d} \\ & \quad + \frac{a(7A b + 4a B)(a + b \cos(c + dx))^2 \sec^2(c + dx) \tan(c + dx)}{12d} \\ & \quad + \frac{a A (a + b \cos(c + dx))^3 \sec^3(c + dx) \tan(c + dx)}{4d} \end{aligned}$$

```
output b^4*B*x+1/8*(3*A*a^4+24*A*a^2*b^2+8*A*b^4+16*B*a^3*b+32*B*a*b^3)*arctanh(sin(d*x+c))/d+1/6*a*(16*A*a^2*b+19*A*b^3+4*B*a^3+34*B*a*b^2)*tan(d*x+c)/d+1/24*a^2*(9*A*a^2+26*A*b^2+32*B*a*b)*sec(d*x+c)*tan(d*x+c)/d+1/12*a*(7*A*b+4*B*a)*(a+b*cos(d*x+c))^2*sec(d*x+c)^2*tan(d*x+c)/d+1/4*a*A*(a+b*cos(d*x+c))^3*sec(d*x+c)^3*tan(d*x+c)/d
```

3.247.2 Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.74

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{24b^4 B dx + 3(3a^4 A + 24a^2 Ab^2 + 8Ab^4 + 16a^3 b B + 32ab^3 B) \operatorname{arctanh}(\sin(c + dx)) + 3a(8(4a^2 Ab + 4Ab^3 +$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output `(24*b^4*B*d*x + 3*(3*a^4*A + 24*a^2*A*b^2 + 8*A*b^4 + 16*a^3*b*B + 32*a*b^3*B)*ArcTanh[Sin[c + d*x]] + 3*a*(8*(4*a^2*A*b + 4*A*b^3 + a^3*B + 6*a*b^2*B) + a*(3*a^2*A + 24*A*b^2 + 16*a*b*B)*Sec[c + d*x] + 2*a^3*A*Sec[c + d*x]^3)*Tan[c + d*x] + 8*a^3*(4*A*b + a*B)*Tan[c + d*x]^3)/(24*d)`

3.247.3 Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3468, 3042, 3526, 3042, 3510, 25, 3042, 3500, 27, 3042, 3214, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{4} \int (a + b \cos(c + dx))^2 (4b^2 B \cos^2(c + dx) + (3Aa^2 + 8bBa + 4Ab^2) \cos(c + dx) + a(7Ab + 4aB)) \sec^4(c + dx) dx + \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (4b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (3Aa^2 + 8bBa + 4Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(7Ab + 4aB)) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3} \frac{1}{4d}$$

↓ 3526

$$\frac{1}{4} \left(\frac{1}{3} \int (a + b \cos(c + dx)) (12B \cos^2(c + dx)b^3 + a(9Aa^2 + 32bBa + 26Ab^2) + (8Ba^3 + 23Aba^2 + 36b^2Ba + 12aB)) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3} \frac{1}{4d} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \int (a + b \sin(c + dx + \frac{\pi}{2})) (12B \sin^2(c + dx + \frac{\pi}{2})b^3 + a(9Aa^2 + 32bBa + 26Ab^2) + (8Ba^3 + 23Aba^2 + 36b^2Ba + 12aB)) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3} \frac{1}{4d} \right)$$

↓ 3510

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{a^2(9a^2A + 32abB + 26Ab^2) \tan(c + dx) \sec(c + dx)}{2d} - \frac{1}{2} \int -((24B \cos^2(c + dx)b^4 + 4a(4Ba^3 + 16Aba^2 + 36b^2Ba + 19Ab^3) + 3(3Aa^4 + 16bBa^3 + 24Ab^2a^2 + 32b^3Ba + 12aB))) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3} \frac{1}{4d} \right) \right)$$

↓ 25

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (24B \cos^2(c + dx)b^4 + 4a(4Ba^3 + 16Aba^2 + 34b^2Ba + 19Ab^3) + 3(3Aa^4 + 16bBa^3 + 24Ab^2a^2 + 32b^3Ba + 12aB)) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3} \frac{1}{4d} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (24B \sin^2(c + dx + \frac{\pi}{2})b^4 + 4a(4Ba^3 + 16Aba^2 + 34b^2Ba + 19Ab^3) + 3(3Aa^4 + 16bBa^3 + 24Ab^2a^2 + 32b^3Ba + 12aB)) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3} \frac{1}{4d} \right) \right)$$

↓ 3500

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(\int 3(3Aa^4 + 16bBa^3 + 24Ab^2a^2 + 32b^3Ba + 8Ab^4 + 8b^4B \cos(c + dx)) \sec(c + dx) dx + \frac{4a(4a^3B + 16a^2Ab + 34ab^2B + 19Ab^3)}{d} \right) \right) \right) \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d}$$

↓ 27

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \int (3Aa^4 + 16bBa^3 + 24Ab^2a^2 + 32b^3Ba + 8Ab^4 + 8b^4B \cos(c + dx)) \sec(c + dx) dx + \frac{4a(4a^3B + 16a^2Ab + 34ab^2B + 19Ab^3)}{d} \right) \right) \right) \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \int \frac{3Aa^4 + 16bBa^3 + 24Ab^2a^2 + 32b^3Ba + 8Ab^4 + 8b^4B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{4a(4a^3B + 16a^2Ab + 34ab^2B + 19Ab^3)}{d} \right) \right) \right) \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d}$$

↓ 3214

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \left((3a^4A + 16a^3bB + 24a^2Ab^2 + 32ab^3B + 8Ab^4) \int \sec(c + dx) dx + 8b^4Bx \right) + \frac{4a(4a^3B + 16a^2Ab + 34ab^2B + 19Ab^3)}{d} \right) \right) \right) \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(3 \left((3a^4A + 16a^3bB + 24a^2Ab^2 + 32ab^3B + 8Ab^4) \int \csc(c + dx + \frac{\pi}{2}) dx + 8b^4Bx \right) + \frac{4a(4a^3B + 16a^2Ab + 34ab^2B + 19Ab^3)}{d} \right) \right) \right) \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d}$$

↓ 4257

$$\frac{1}{4} \left(\frac{1}{3} \left(\frac{a^2(9a^2A + 32abB + 26Ab^2) \tan(c + dx) \sec(c + dx)}{2d} + \frac{1}{2} \left(\frac{4a(4a^3B + 16a^2Ab + 34ab^2B + 19Ab^3) \tan(c + dx)}{d} + \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^3}{4d} \right) \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output $(a*A*(a + b*\cos[c + d*x])^3*\sec[c + d*x]^3*\tan[c + d*x])/(4*d) + ((a*(7*A*b + 4*a*B)*(a + b*\cos[c + d*x])^2*\sec[c + d*x]^2*\tan[c + d*x])/(3*d) + ((a^2*(9*a^2*A + 26*A*b^2 + 32*a*b*B)*\sec[c + d*x]*\tan[c + d*x])/(2*d) + (3*(8*b^4*B*x + ((3*a^4*A + 24*a^2*A*b^2 + 8*A*b^4 + 16*a^3*b*B + 32*a*b^3*B)*\operatorname{ArcTanh}[\sin[c + d*x]])/d) + (4*a*(16*a^2*A*b + 19*A*b^3 + 4*a^3*B + 34*a*b^2*B)*\tan[c + d*x])/d)/2)/3)/4$

3.247.3.1 Defintions of rubi rules used

rule 25 $\operatorname{Int}[-(F_x), x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$

rule 27 $\operatorname{Int}[(a_)*(F_x), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[F_x, (b_)*(G_x)] /; \operatorname{FreeQ}[b, x]$

rule 3042 $\operatorname{Int}[u, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3214 $\operatorname{Int}[(a_ + (b_)*\sin[(e_ + (f_)*(x_))]) / ((c_ + (d_)*\sin[(e_ + (f_)*(x_))])), x_Symbol] \rightarrow \operatorname{Simp}[b*(x/d), x] - \operatorname{Simp}[(b*c - a*d)/d \operatorname{Int}[1/(c + d*\sin[e + f*x]), x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0]$

rule 3468 $\operatorname{Int}[(a_ + (b_)*\sin[(e_ + (f_)*(x_))]^{(m_)} * ((A_ + (B_)*\sin[(e_ + (f_)*(x_))])^{(n_)}), x_Symbol] \rightarrow \operatorname{Simp}[(- (b*c - a*d)) * (B*c - A*d) * \cos[e + f*x] * (a + b*\sin[e + f*x])^{(m - 1)} * ((c + d*\sin[e + f*x])^{(n + 1)} / (d*f*(n + 1)*(c^2 - d^2))), x] + \operatorname{Simp}[1/(d*(n + 1)*(c^2 - d^2)) \operatorname{Int}[(a + b*\sin[e + f*x])^{(m - 2)} * (c + d*\sin[e + f*x])^{(n + 1)} * \operatorname{Simp}[b*(b*c - a*d) * (B*c - A*d) * (m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d) * (n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2))) * (n + 1) - a*(b*c - a*d) * (B*c - A*d) * (n + 2)) * \sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d) * (m + n + 1) - b*B*(c^2*m + d^2*(n + 1))) * \sin[e + f*x]^2, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{NeQ}[a^2 - b^2, 0] \ \&\& \ \operatorname{NeQ}[c^2 - d^2, 0] \ \&\& \ \operatorname{GtQ}[m, 1] \ \&\& \ \operatorname{LtQ}[n, -1]$

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]*(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.247.4 Maple [A] (verified)

Time = 5.47 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.99

method	result
parts	$\frac{a^4 A \left(- \left(- \frac{(\sec^3(dx+c))}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c)+\tan(dx+c))}{8} \right)}{d} + \frac{(A b^4 + 4 B a b^3) \ln(\sec(dx+c)+\tan(dx+c))}{d}$
derivativedivides	$\frac{a^4 A \left(- \left(- \frac{(\sec^3(dx+c))}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c)+\tan(dx+c))}{8} \right)}{d} - B a^4 \left(- \frac{2}{3} - \frac{(\sec^2(dx+c))}{3} \right) \tan(dx+c)$
default	$\frac{a^4 A \left(- \left(- \frac{(\sec^3(dx+c))}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c)+\tan(dx+c))}{8} \right)}{d} - B a^4 \left(- \frac{2}{3} - \frac{(\sec^2(dx+c))}{3} \right) \tan(dx+c)$
parallelrisch	$-36(a^4 A + 8 A a^2 b^2 + \frac{8}{3} A b^4 + \frac{16}{3} B a^3 b + \frac{32}{3} B a b^3) \left(\frac{3}{4} + \frac{\cos(4dx+4c)}{4} + \cos(2dx+2c) \right) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) + 36(a^4 A + 8 A a^2 b^2 + \frac{8}{3} A b^4 + \frac{16}{3} B a^3 b + \frac{32}{3} B a b^3)$
risch	$b^4 B x - \frac{ia(72Aa b^2 e^{5i(dx+c)} + 48B a^2 b e^{5i(dx+c)} - 144Ba b^2 e^{6i(dx+c)} - 72Aa b^2 e^{i(dx+c)} - 48B a^2 b e^{i(dx+c)} + 72Aa b^2 e^{i(dx+c)})}{d}$

```
input int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNVERBOSE)
```

```
output a^4*A/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^4+4*B*a*b^3)/d*ln(sec(d*x+c)+tan(d*x+c))+(4*A*a*b^3+6*B*a^2*b^2)/d*tan(d*x+c)+(6*A*a^2*b^2+4*B*a^3*b)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-(4*A*a^3*b+B*a^4)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+B*b^4/d*(d*x+c)
```

3.247.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.16

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{48 B b^4 dx \cos(dx + c)^4 + 3(3 A a^4 + 16 B a^3 b + 24 A a^2 b^2 + 32 B a b^3 + 8 A b^4) \cos(dx + c)^4 \log(\sin(dx + c))}{d}$$

```
input integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fracas")
```

output $1/48*(48*B*b^4*d*x*cos(d*x + c)^4 + 3*(3*A*a^4 + 16*B*a^3*b + 24*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*cos(d*x + c)^4*log(sin(d*x + c) + 1) - 3*(3*A*a^4 + 16*B*a^3*b + 24*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*cos(d*x + c)^4*log(-sin(d*x + c) + 1) + 2*(6*A*a^4 + 16*(B*a^4 + 4*A*a^3*b + 9*B*a^2*b^2 + 6*A*a*b^3)*cos(d*x + c)^3 + 3*(3*A*a^4 + 16*B*a^3*b + 24*A*a^2*b^2)*cos(d*x + c)^2 + 8*(B*a^4 + 4*A*a^3*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^4)$

3.247.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output `Timed out`

3.247.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.47

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{16 (\tan(dx + c)^3 + 3 \tan(dx + c)) Ba^4 + 64 (\tan(dx + c)^3 + 3 \tan(dx + c)) Aa^3b + 48 (dx + c) Bb^4 - 3}{}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output $\frac{1}{48}(16(\tan(dx + c))^3 + 3\tan(dx + c))*B*a^4 + 64(\tan(dx + c)^3 + 3\tan(dx + c))*A*a^3*b + 48(dx + c)*B*b^4 - 3A*a^4*(2*(3*\sin(dx + c))^3 - 5*\sin(dx + c))/(\sin(dx + c)^4 - 2*\sin(dx + c)^2 + 1) - 3*\log(\sin(dx + c) + 1) + 3*\log(\sin(dx + c) - 1)) - 48B*a^3*b*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) - 72A*a^2*b^2*(2*\sin(dx + c)/(\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) + 96B*a*b^3*(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 24A*b^4*(\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 288B*a^2*b^2*\tan(dx + c) + 192A*a*b^3*\tan(dx + c))/d$

3.247.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(206) = 412$.

Time = 0.35 (sec) , antiderivative size = 635, normalized size of antiderivative = 2.94

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx$$

$$= \frac{24(dx + c)Bb^4 + 3(3Aa^4 + 16Ba^3b + 24Aa^2b^2 + 32Bab^3 + 8Ab^4) \log\left(\left|\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right|\right) - 3(3Aa^4 + 16Ba^3b + 24Aa^2b^2 + 32Bab^3 + 8Ab^4)}{d}$$

input `integrate((a+b*cos(dx+c))^4*(A+B*cos(dx+c))*sec(dx+c)^5,x, algorithm="giac")`

output

```

1/24*(24*(d*x + c)*B*b^4 + 3*(3*A*a^4 + 16*B*a^3*b + 24*A*a^2*b^2 + 32*B*a
*b^3 + 8*A*b^4)*log(abs(tan(1/2*d*x + 1/2*c) + 1)) - 3*(3*A*a^4 + 16*B*a^3
*b + 24*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*log(abs(tan(1/2*d*x + 1/2*c) - 1
)) + 2*(15*A*a^4*tan(1/2*d*x + 1/2*c)^7 - 24*B*a^4*tan(1/2*d*x + 1/2*c)^7
- 96*A*a^3*b*tan(1/2*d*x + 1/2*c)^7 + 48*B*a^3*b*tan(1/2*d*x + 1/2*c)^7 +
72*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^7 - 144*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^7
- 96*A*a*b^3*tan(1/2*d*x + 1/2*c)^7 + 9*A*a^4*tan(1/2*d*x + 1/2*c)^5 + 40
*B*a^4*tan(1/2*d*x + 1/2*c)^5 + 160*A*a^3*b*tan(1/2*d*x + 1/2*c)^5 - 48*B*
a^3*b*tan(1/2*d*x + 1/2*c)^5 - 72*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^5 + 432*B
*a^2*b^2*tan(1/2*d*x + 1/2*c)^5 + 288*A*a*b^3*tan(1/2*d*x + 1/2*c)^5 + 9*A
*a^4*tan(1/2*d*x + 1/2*c)^3 - 40*B*a^4*tan(1/2*d*x + 1/2*c)^3 - 160*A*a^3*
b*tan(1/2*d*x + 1/2*c)^3 - 48*B*a^3*b*tan(1/2*d*x + 1/2*c)^3 - 72*A*a^2*b^
2*tan(1/2*d*x + 1/2*c)^3 - 432*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^3 - 288*A*a*
b^3*tan(1/2*d*x + 1/2*c)^3 + 15*A*a^4*tan(1/2*d*x + 1/2*c) + 24*B*a^4*tan(
1/2*d*x + 1/2*c) + 96*A*a^3*b*tan(1/2*d*x + 1/2*c) + 48*B*a^3*b*tan(1/2*d*
x + 1/2*c) + 72*A*a^2*b^2*tan(1/2*d*x + 1/2*c) + 144*B*a^2*b^2*tan(1/2*d*x
+ 1/2*c) + 96*A*a*b^3*tan(1/2*d*x + 1/2*c))/(tan(1/2*d*x + 1/2*c)^2 - 1)^
4)/d

```

3.247.9 Mupad [B] (verification not implemented)

Time = 3.17 (sec) , antiderivative size = 1969, normalized size of antiderivative = 9.12

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Too large to display}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4)/cos(c + d*x)^5,x)`

output

```

((27*A*a^4*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/8 + 9*A*b^4*atanh
(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)) + (9*A*a^4*sin(3*c + 3*d*x))/8 + 4
*B*a^4*sin(2*c + 2*d*x) + B*a^4*sin(4*c + 4*d*x) + 9*B*b^4*atan((9*A^2*a^8
*sin(c/2 + (d*x)/2) + 64*A^2*b^8*sin(c/2 + (d*x)/2) + 64*B^2*b^8*sin(c/2 +
(d*x)/2) + 384*A^2*a^2*b^6*sin(c/2 + (d*x)/2) + 624*A^2*a^4*b^4*sin(c/2 +
(d*x)/2) + 144*A^2*a^6*b^2*sin(c/2 + (d*x)/2) + 1024*B^2*a^2*b^6*sin(c/2
+ (d*x)/2) + 1024*B^2*a^4*b^4*sin(c/2 + (d*x)/2) + 256*B^2*a^6*b^2*sin(c/2
+ (d*x)/2) + 1792*A*B*a^3*b^5*sin(c/2 + (d*x)/2) + 960*A*B*a^5*b^3*sin(c/
2 + (d*x)/2) + 512*A*B*a*b^7*sin(c/2 + (d*x)/2) + 96*A*B*a^7*b*sin(c/2 + (
d*x)/2))/(cos(c/2 + (d*x)/2)*(9*A^2*a^8 + 64*A^2*b^8 + 64*B^2*b^8 + 384*A^
2*a^2*b^6 + 624*A^2*a^4*b^4 + 144*A^2*a^6*b^2 + 1024*B^2*a^2*b^6 + 1024*B^
2*a^4*b^4 + 256*B^2*a^6*b^2 + 512*A*B*a*b^7 + 96*A*B*a^7*b + 1792*A*B*a^3*
b^5 + 960*A*B*a^5*b^3))) + (33*A*a^4*sin(c + d*x))/8 + 12*B*b^4*cos(2*c +
2*d*x)*atan((9*A^2*a^8*sin(c/2 + (d*x)/2) + 64*A^2*b^8*sin(c/2 + (d*x)/2)
+ 64*B^2*b^8*sin(c/2 + (d*x)/2) + 384*A^2*a^2*b^6*sin(c/2 + (d*x)/2) + 624
*A^2*a^4*b^4*sin(c/2 + (d*x)/2) + 144*A^2*a^6*b^2*sin(c/2 + (d*x)/2) + 102
4*B^2*a^2*b^6*sin(c/2 + (d*x)/2) + 1024*B^2*a^4*b^4*sin(c/2 + (d*x)/2) + 2
56*B^2*a^6*b^2*sin(c/2 + (d*x)/2) + 1792*A*B*a^3*b^5*sin(c/2 + (d*x)/2) +
960*A*B*a^5*b^3*sin(c/2 + (d*x)/2) + 512*A*B*a*b^7*sin(c/2 + (d*x)/2) + 96
*A*B*a^7*b*sin(c/2 + (d*x)/2))/(cos(c/2 + (d*x)/2)*(9*A^2*a^8 + 64*A^2*...

```

3.248 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^6(c+dx) dx$

3.248.1 Optimal result	2291
3.248.2 Mathematica [A] (verified)	2292
3.248.3 Rubi [A] (verified)	2292
3.248.4 Maple [A] (verified)	2297
3.248.5 Fricas [A] (verification not implemented)	2298
3.248.6 Sympy [F(-1)]	2298
3.248.7 Maxima [A] (verification not implemented)	2299
3.248.8 Giac [B] (verification not implemented)	2299
3.248.9 Mupad [B] (verification not implemented)	2300

3.248.1 Optimal result

Integrand size = 31, antiderivative size = 267

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{(12a^3 Ab + 16aAb^3 + 3a^4 B + 24a^2 b^2 B + 8b^4 B) \operatorname{arctanh}(\sin(c + dx))}{8d}$$

$$+ \frac{(8a^4 A + 60a^2 Ab^2 + 15Ab^4 + 40a^3 bB + 60ab^3 B) \tan(c + dx)}{15d}$$

$$+ \frac{a(60a^2 Ab + 56Ab^3 + 15a^3 B + 110ab^2 B) \sec(c + dx) \tan(c + dx)}{40d}$$

$$+ \frac{a^2(8a^2 A + 18Ab^2 + 25abB) \sec^2(c + dx) \tan(c + dx)}{30d}$$

$$+ \frac{a(8Ab + 5aB)(a + b \cos(c + dx))^2 \sec^3(c + dx) \tan(c + dx)}{20d}$$

$$+ \frac{aA(a + b \cos(c + dx))^3 \sec^4(c + dx) \tan(c + dx)}{5d}$$

```
output 1/8*(12*A*a^3*b+16*A*a*b^3+3*B*a^4+24*B*a^2*b^2+8*B*b^4)*arctanh(sin(d*x+c
))/d+1/15*(8*A*a^4+60*A*a^2*b^2+15*A*b^4+40*B*a^3*b+60*B*a*b^3)*tan(d*x+c)
/d+1/40*a*(60*A*a^2*b+56*A*b^3+15*B*a^3+110*B*a*b^2)*sec(d*x+c)*tan(d*x+c)
/d+1/30*a^2*(8*A*a^2+18*A*b^2+25*B*a*b)*sec(d*x+c)^2*tan(d*x+c)/d+1/20*a*(
8*A*b+5*B*a)*(a+b*cos(d*x+c))^2*sec(d*x+c)^3*tan(d*x+c)/d+1/5*a*A*(a+b*cos
(d*x+c))^3*sec(d*x+c)^4*tan(d*x+c)/d
```

3.248.2 Mathematica [A] (verified)

Time = 2.68 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.74

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{15(12a^3Ab + 16aAb^3 + 3a^4B + 24a^2b^2B + 8b^4B) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (120(a^4A + 6a^2Ab^2$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`output `(15*(12*a^3*A*b + 16*a*A*b^3 + 3*a^4*B + 24*a^2*b^2*B + 8*b^4*B)*ArcTanh[Sin[c + d*x]] + Tan[c + d*x]*(120*(a^4*A + 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B + 4*a*b^3*B) + 15*a*(12*a^2*A*b + 16*A*b^3 + 3*a^3*B + 24*a*b^2*B)*Sec[c + d*x] + 30*a^3*(4*A*b + a*B)*Sec[c + d*x]^3 + 80*a^2*(a^2*A + 3*A*b^2 + 2*a*b*B)*Tan[c + d*x]^2 + 24*a^4*A*Tan[c + d*x]^4))/(120*d)`**3.248.3 Rubi [A] (verified)**Time = 1.73 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.06, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3468, 3042, 3526, 3042, 3510, 25, 3042, 3500, 3042, 3227, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^6(c + dx)(a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^6} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{5} \int (a + b \cos(c + dx))^2 (b(aA + 5bB) \cos^2(c + dx) + (4Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(8Ab + 5aB)) \sec^5(c + dx) dx + \frac{aA \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^3}{5d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{5} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (b(aA + 5bB) \sin(c + dx + \frac{\pi}{2})^2 + (4Aa^2 + 10bBa + 5Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(8A + 5bB)) \sin^5(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^4(c + dx) (a + b \cos(c + dx))^3} \frac{1}{5d}$$

↓ 3526

$$\frac{1}{5} \left(\frac{1}{4} \int (a + b \cos(c + dx)) (b(5Ba^2 + 12Aba + 20b^2B) \cos^2(c + dx) + (15Ba^3 + 44Aba^2 + 60b^2Ba + 20Ab^3) \cos(c + dx)) \frac{1}{aA \tan(c + dx) \sec^4(c + dx) (a + b \cos(c + dx))^3} \frac{1}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) (b(5Ba^2 + 12Aba + 20b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (15Ba^3 + 44Aba^2 + 60b^2Ba + 20Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(8A + 5bB)) \sin^4(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^4(c + dx) (a + b \cos(c + dx))^3} \frac{1}{5d} \right)$$

↓ 3510

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{2a^2(8a^2A + 25abB + 18Ab^2) \tan(c + dx) \sec^2(c + dx)}{3d} - \frac{1}{3} \int -((3b^2(5Ba^2 + 12Aba + 20b^2B) \cos^2(c + dx) + 4(8Aa^4 + 40bBa^3 + 60Ab^2a^2 + 60b^3Ba + 15Ab^4) \cos(c + dx))) \frac{1}{aA \tan(c + dx) \sec^4(c + dx) (a + b \cos(c + dx))^3} \frac{1}{5d} \right) \right)$$

↓ 25

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int (3b^2(5Ba^2 + 12Aba + 20b^2B) \cos^2(c + dx) + 4(8Aa^4 + 40bBa^3 + 60Ab^2a^2 + 60b^3Ba + 15Ab^4) \cos(c + dx)) \frac{1}{aA \tan(c + dx) \sec^4(c + dx) (a + b \cos(c + dx))^3} \frac{1}{5d} \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int \frac{3b^2(5Ba^2 + 12Aba + 20b^2B) \sin(c + dx + \frac{\pi}{2})^2 + 4(8Aa^4 + 40bBa^3 + 60Ab^2a^2 + 60b^3Ba + 15Ab^4) \sin(c + dx + \frac{\pi}{2})}{aA \tan(c + dx) \sec^4(c + dx) (a + b \cos(c + dx))^3} \frac{1}{5d} \right) \right)$$

↓ 3500

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (8(8Aa^4 + 40bBa^3 + 60Ab^2a^2 + 60b^3Ba + 15Ab^4) + 15(3Ba^4 + 12Aba^3 + 24b^2Ba^2 + 16Ab^3a + 8aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3) \right) \right) \right) \frac{aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int \frac{8(8Aa^4 + 40bBa^3 + 60Ab^2a^2 + 60b^3Ba + 15Ab^4) + 15(3Ba^4 + 12Aba^3 + 24b^2Ba^2 + 16Ab^3a + 8aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{\sin(c+dx + \frac{\pi}{2})^2} \right) \right) \right) \frac{aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{5d}$$

↓ 3227

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(8(8a^4A + 40a^3bB + 60a^2Ab^2 + 60ab^3B + 15Ab^4) \int \sec^2(c+dx)dx + 15(3a^4B + 12a^3Ab + 24a^2b^2B + 16aAb^3 + 8b^4B) \right) \right) \right) \frac{aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15(3a^4B + 12a^3Ab + 24a^2b^2B + 16aAb^3 + 8b^4B) \int \csc(c+dx + \frac{\pi}{2}) dx + 8(8a^4A + 40a^3bB + 60a^2Ab^2 + 60ab^3B + 15Ab^4) \right) \right) \right) \frac{aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{5d}$$

↓ 4254

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15(3a^4B + 12a^3Ab + 24a^2b^2B + 16aAb^3 + 8b^4B) \int \csc(c+dx + \frac{\pi}{2}) dx - \frac{8(8a^4A + 40a^3bB + 60a^2Ab^2 + 60ab^3B + 15Ab^4)}{\sin(c+dx + \frac{\pi}{2})} \right) \right) \right) \frac{aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{5d}$$

↓ 24

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(15(3a^4B + 12a^3Ab + 24a^2b^2B + 16aAb^3 + 8b^4B) \int \csc(c+dx + \frac{\pi}{2}) dx + \frac{8(8a^4A + 40a^3bB + 60a^2Ab^2 + 60ab^3B + 15Ab^4)}{\sin(c+dx + \frac{\pi}{2})} \right) \right) \right) \frac{aA \tan(c+dx) \sec^4(c+dx)(a+b \cos(c+dx))^3}{5d}$$

↓ 4257

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{2a^2(8a^2A + 25abB + 18Ab^2) \tan(c + dx) \sec^2(c + dx)}{3d} + \frac{1}{3} \left(\frac{3a(15a^3B + 60a^2Ab + 110ab^2B + 56Ab^3) \tan(c + dx) \sec^4(c + dx)(a + b \cos(c + dx))^3}{2d} \right) \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^6,x]`

output `(a*A*(a + b*Cos[c + d*x])^3*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) + ((a*(8*A*b + 5*a*B)*(a + b*Cos[c + d*x])^2*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((2*a^2*(8*a^2*A + 18*A*b^2 + 25*a*b*B)*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + ((3*a*(60*a^2*A*b + 56*A*b^3 + 15*a^3*B + 110*a*b^2*B)*Sec[c + d*x]*Tan[c + d*x])/(2*d) + ((15*(12*a^3*A*b + 16*a*A*b^3 + 3*a^4*B + 24*a^2*b^2*B + 8*b^4*B)*ArcTanh[Sin[c + d*x]])/d + (8*(8*a^4*A + 60*a^2*A*b^2 + 15*A*b^4 + 40*a^3*b*B + 60*a*b^3*B)*Tan[c + d*x])/d)/2)/3)/4)/5`

3.248.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-(b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

```
rule 3526 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 -
d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m -
1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*
d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n +
1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x
] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f
*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d
, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.248.4 Maple [A] (verified)

Time = 6.40 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.89

method	result
parts	$\frac{a^4 A \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c)}{d} + \frac{(A b^4 + 4B a b^3) \tan(dx+c)}{d} + \frac{(4A a b^3 + 6B a^2 b^2) \left(\frac{\sec(dx+c)}{d} \right)}{d}$
derivativedivides	$-a^4 A \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c) + B a^4 \left(-\left(-\frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c))}{d} \right)$
default	$-a^4 A \left(-\frac{8}{15} - \frac{\sec^4(dx+c)}{5} - \frac{4(\sec^2(dx+c))}{15} \right) \tan(dx+c) + B a^4 \left(-\left(-\frac{\sec^3(dx+c)}{4} - \frac{3 \sec(dx+c)}{8} \right) \tan(dx+c) + \frac{3 \ln(\sec(dx+c))}{d} \right)$
parallelrisch	$-180(A a^3 b + \frac{4}{3} A a b^3 + \frac{1}{4} B a^4 + 2B a^2 b^2 + \frac{2}{3} B b^4) (\cos(5dx+5c) + 5 \cos(3dx+3c) + 10 \cos(dx+c)) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) + 180$
risch	$\frac{i(480A a^2 b^2 + 320B a^3 b + 480B a b^3 + 64a^4 A - 720B a^2 b^2 e^{7i(dx+c)} + 1440A a^2 b^2 e^{6i(dx+c)} + 960B a^3 b e^{6i(dx+c)} + 1920B a b^5 e^{5i(dx+c)})}{d}$

```
input int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c)^6,x,method=_RETURNVERBO
SE)
```

3.248. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$

output
$$-a^4 A / d * (-8/15 - 1/5 * \sec(dx+c)^4 - 4/15 * \sec(dx+c)^2) * \tan(dx+c) + (A*b^4 + 4*B*a*b^3) / d * \tan(dx+c) + (4*A*a*b^3 + 6*B*a^2*b^2) / d * (1/2 * \sec(dx+c) * \tan(dx+c) + 1/2 * \ln(\sec(dx+c) + \tan(dx+c))) - (6*A*a^2*b^2 + 4*B*a^3*b) / d * (-2/3 - 1/3 * \sec(dx+c)^2) * \tan(dx+c) + (4*A*a^3*b + B*a^4) / d * (-1/4 * \sec(dx+c)^3 - 3/8 * \sec(dx+c) * \tan(dx+c) + 3/8 * \ln(\sec(dx+c) + \tan(dx+c))) + B*b^4 / d * \ln(\sec(dx+c) + \tan(dx+c))$$

3.248.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.05

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{15(3Ba^4 + 12Aa^3b + 24Ba^2b^2 + 16Aab^3 + 8Bb^4) \cos(dx + c)^5 \log(\sin(dx + c) + 1) - 15(3Ba^4 + 12Aa^3b + 24Ba^2b^2 + 16Aab^3 + 8Bb^4) \cos(dx + c)^5 \log(-\sin(dx + c) + 1) + 2(24Aa^4 + 8(8Aa^4 + 40B*a^3*b + 60A*a^2*b^2 + 60B*a*b^3 + 15A*b^4) * \cos(dx + c)^4 + 15(3B*a^4 + 12A*a^3*b + 24B*a^2*b^2 + 16A*a*b^3) * \cos(dx + c)^3 + 16(2A*a^4 + 10B*a^3*b + 15A*a^2*b^2) * \cos(dx + c)^2 + 30(B*a^4 + 4A*a^3*b) * \cos(dx + c)) * \sin(dx + c)}{(d * \cos(dx + c))^5}$$

input `integrate((a+b*cos(dx+c))^4*(A+B*cos(dx+c))*sec(dx+c)^6,x, algorithm="fricas")`

output
$$\frac{1}{240} * (15 * (3B*a^4 + 12*A*a^3*b + 24*B*a^2*b^2 + 16*A*a*b^3 + 8*B*b^4) * \cos(dx + c)^5 * \log(\sin(dx + c) + 1) - 15 * (3B*a^4 + 12*A*a^3*b + 24*B*a^2*b^2 + 16*A*a*b^3 + 8*B*b^4) * \cos(dx + c)^5 * \log(-\sin(dx + c) + 1) + 2 * (24*A*a^4 + 8 * (8*A*a^4 + 40*B*a^3*b + 60*A*a^2*b^2 + 60*B*a*b^3 + 15*A*b^4) * \cos(dx + c)^4 + 15 * (3B*a^4 + 12*A*a^3*b + 24*B*a^2*b^2 + 16*A*a*b^3) * \cos(dx + c)^3 + 16 * (2*A*a^4 + 10*B*a^3*b + 15*A*a^2*b^2) * \cos(dx + c)^2 + 30 * (B*a^4 + 4*A*a^3*b) * \cos(dx + c)) * \sin(dx + c)) / (d * \cos(dx + c)^5)$$

3.248.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(dx+c))**4*(A+B*cos(dx+c))*sec(dx+c)**6,x)`

output `Timed out`

3.248.
$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

3.248.7 Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 386, normalized size of antiderivative = 1.45

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{16 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c)) A a^4 + 320 (\tan(dx + c)^3 + 3 \tan(dx + c)) B a^3 b + 480 (\tan(dx + c)^3 + 3 \tan(dx + c)) A a^2 b^2 - 15 B a^4 (2 (3 \sin(dx + c)^3 - 5 \sin(dx + c)) / (\sin(dx + c)^4 - 2 \sin(dx + c)^2 + 1) - 3 \log(\sin(dx + c) + 1) + 3 \log(\sin(dx + c) - 1)) - 60 A a^3 b (2 (3 \sin(dx + c)^3 - 5 \sin(dx + c)) / (\sin(dx + c)^4 - 2 \sin(dx + c)^2 + 1) - 3 \log(\sin(dx + c) + 1) + 3 \log(\sin(dx + c) - 1)) - 360 B a^2 b^2 (2 \sin(dx + c) / (\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) - 240 A a b^3 (2 \sin(dx + c) / (\sin(dx + c)^2 - 1) - \log(\sin(dx + c) + 1) + \log(\sin(dx + c) - 1)) + 120 B b^4 (\log(\sin(dx + c) + 1) - \log(\sin(dx + c) - 1)) + 960 B a b^3 \tan(dx + c) + 240 A b^4 \tan(dx + c)) / d}{1}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="maxima")`

output `1/240*(16*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*A*a^4 + 320*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^3*b + 480*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a^2*b^2 - 15*B*a^4*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 60*A*a^3*b*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 360*B*a^2*b^2*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 240*A*a*b^3*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 120*B*b^4*(log(sin(d*x + c) + 1) - log(sin(d*x + c) - 1)) + 960*B*a*b^3*tan(d*x + c) + 240*A*b^4*tan(d*x + c))/d`

3.248.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 850 vs. 2(255) = 510.

Time = 0.38 (sec) , antiderivative size = 850, normalized size of antiderivative = 3.18

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx = \text{Too large to display}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^6,x, algorithm="giac")`

```

output 1/120*(15*(3*B*a^4 + 12*A*a^3*b + 24*B*a^2*b^2 + 16*A*a*b^3 + 8*B*b^4)*log
(abs(tan(1/2*d*x + 1/2*c) + 1)) - 15*(3*B*a^4 + 12*A*a^3*b + 24*B*a^2*b^2
+ 16*A*a*b^3 + 8*B*b^4)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) - 2*(120*A*a^4*
tan(1/2*d*x + 1/2*c)^9 - 75*B*a^4*tan(1/2*d*x + 1/2*c)^9 - 300*A*a^3*b*tan
(1/2*d*x + 1/2*c)^9 + 480*B*a^3*b*tan(1/2*d*x + 1/2*c)^9 + 720*A*a^2*b^2*t
an(1/2*d*x + 1/2*c)^9 - 360*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^9 - 240*A*a*b^3
*tan(1/2*d*x + 1/2*c)^9 + 480*B*a*b^3*tan(1/2*d*x + 1/2*c)^9 + 120*A*b^4*t
an(1/2*d*x + 1/2*c)^9 - 160*A*a^4*tan(1/2*d*x + 1/2*c)^7 + 30*B*a^4*tan(1/
2*d*x + 1/2*c)^7 + 120*A*a^3*b*tan(1/2*d*x + 1/2*c)^7 - 1280*B*a^3*b*tan(1
/2*d*x + 1/2*c)^7 - 1920*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^7 + 720*B*a^2*b^2*
tan(1/2*d*x + 1/2*c)^7 + 480*A*a*b^3*tan(1/2*d*x + 1/2*c)^7 - 1920*B*a*b^3
*tan(1/2*d*x + 1/2*c)^7 - 480*A*b^4*tan(1/2*d*x + 1/2*c)^7 + 464*A*a^4*tan
(1/2*d*x + 1/2*c)^5 + 1600*B*a^3*b*tan(1/2*d*x + 1/2*c)^5 + 2400*A*a^2*b^2
*tan(1/2*d*x + 1/2*c)^5 + 2880*B*a*b^3*tan(1/2*d*x + 1/2*c)^5 + 720*A*b^4*
tan(1/2*d*x + 1/2*c)^5 - 160*A*a^4*tan(1/2*d*x + 1/2*c)^3 - 30*B*a^4*tan(1
/2*d*x + 1/2*c)^3 - 120*A*a^3*b*tan(1/2*d*x + 1/2*c)^3 - 1280*B*a^3*b*tan(
1/2*d*x + 1/2*c)^3 - 1920*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^3 - 720*B*a^2*b^2
*tan(1/2*d*x + 1/2*c)^3 - 480*A*a*b^3*tan(1/2*d*x + 1/2*c)^3 - 1920*B*a*b^
3*tan(1/2*d*x + 1/2*c)^3 - 480*A*b^4*tan(1/2*d*x + 1/2*c)^3 + 120*A*a^4*ta
n(1/2*d*x + 1/2*c) + 75*B*a^4*tan(1/2*d*x + 1/2*c) + 300*A*a^3*b*tan(1/...

```

3.248.9 Mupad [B] (verification not implemented)

Time = 4.17 (sec) , antiderivative size = 555, normalized size of antiderivative = 2.08

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^6(c + dx) dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{3Ba^4}{8} + \frac{3Aa^3b}{2} + 3Ba^2b^2 + 2Aab^3 + Bb^4\right)}{\frac{3Ba^4}{2} + 6Aa^3b + 12Ba^2b^2 + 8Aab^3 + 4Bb^4}\right) \left(\frac{3Ba^4}{4} + 3Aa^3b + 6Ba^2b^2 + 4Aab^3 + 2Bb^4\right)}{d}$$

$$- \frac{\left(2Aa^4 + 2Ab^4 - \frac{5Ba^4}{4} + 12Aa^2b^2 - 6Ba^2b^2 - 4Aab^3 - 5Aa^3b + 8Aab^3 + 8Ba^3b\right) \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d}$$

```

input int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4)/cos(c + d*x)^6,x)

```

output $(\operatorname{atanh}((4*\tan(c/2 + (d*x)/2)*((3*B*a^4)/8 + B*b^4 + 3*B*a^2*b^2 + 2*A*a*b^3 + (3*A*a^3*b)/2)))/((3*B*a^4)/2 + 4*B*b^4 + 12*B*a^2*b^2 + 8*A*a*b^3 + 6*A*a^3*b))*((3*B*a^4)/4 + 2*B*b^4 + 6*B*a^2*b^2 + 4*A*a*b^3 + 3*A*a^3*b))/d - (\tan(c/2 + (d*x)/2)*(2*A*a^4 + 2*A*b^4 + (5*B*a^4)/4 + 12*A*a^2*b^2 + 6*B*a^2*b^2 + 4*A*a*b^3 + 5*A*a^3*b + 8*B*a*b^3 + 8*B*a^3*b) + \tan(c/2 + (d*x)/2)^5*((116*A*a^4)/15 + 12*A*b^4 + 40*A*a^2*b^2 + 48*B*a*b^3 + (80*B*a^3*b)/3) + \tan(c/2 + (d*x)/2)^9*(2*A*a^4 + 2*A*b^4 - (5*B*a^4)/4 + 12*A*a^2*b^2 + b^2 - 6*B*a^2*b^2 - 4*A*a*b^3 - 5*A*a^3*b + 8*B*a*b^3 + 8*B*a^3*b) - \tan(c/2 + (d*x)/2)^3*((8*A*a^4)/3 + 8*A*b^4 + (B*a^4)/2 + 32*A*a^2*b^2 + 12*B*a^2*b^2 + 8*A*a*b^3 + 2*A*a^3*b + 32*B*a*b^3 + (64*B*a^3*b)/3) - \tan(c/2 + (d*x)/2)^7*((8*A*a^4)/3 + 8*A*b^4 - (B*a^4)/2 + 32*A*a^2*b^2 - 12*B*a^2*b^2 - 8*A*a*b^3 - 2*A*a^3*b + 32*B*a*b^3 + (64*B*a^3*b)/3))/d*(5*\tan(c/2 + (d*x)/2)^2 - 10*\tan(c/2 + (d*x)/2)^4 + 10*\tan(c/2 + (d*x)/2)^6 - 5*\tan(c/2 + (d*x)/2)^8 + \tan(c/2 + (d*x)/2)^{10} - 1))$

3.249 $\int (a+b \cos(c+dx))^4 (A+B \cos(c+dx)) \sec^7(c+dx) dx$

3.249.1 Optimal result	2302
3.249.2 Mathematica [A] (verified)	2303
3.249.3 Rubi [A] (verified)	2303
3.249.4 Maple [A] (verified)	2309
3.249.5 Fricas [A] (verification not implemented)	2310
3.249.6 Sympy [F(-1)]	2310
3.249.7 Maxima [A] (verification not implemented)	2311
3.249.8 Giac [B] (verification not implemented)	2311
3.249.9 Mupad [B] (verification not implemented)	2312

3.249.1 Optimal result

Integrand size = 31, antiderivative size = 324

$$\begin{aligned} & \int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx \\ &= \frac{(5a^4 A + 36a^2 Ab^2 + 8Ab^4 + 24a^3 bB + 32ab^3 B) \operatorname{arctanh}(\sin(c + dx))}{16d} \\ &+ \frac{(32a^3 Ab + 40aAb^3 + 8a^4 B + 60a^2 b^2 B + 15b^4 B) \tan(c + dx)}{15d} \\ &+ \frac{(5a^4 A + 36a^2 Ab^2 + 8Ab^4 + 24a^3 bB + 32ab^3 B) \sec(c + dx) \tan(c + dx)}{16d} \\ &+ \frac{a(16a^2 Ab + 13Ab^3 + 4a^3 B + 27ab^2 B) \sec^2(c + dx) \tan(c + dx)}{15d} \\ &+ \frac{a^2(25a^2 A + 48Ab^2 + 72abB) \sec^3(c + dx) \tan(c + dx)}{120d} \\ &+ \frac{a(3Ab + 2aB)(a + b \cos(c + dx))^2 \sec^4(c + dx) \tan(c + dx)}{10d} \\ &+ \frac{aA(a + b \cos(c + dx))^3 \sec^5(c + dx) \tan(c + dx)}{6d} \end{aligned}$$

output $\frac{1}{16}(5Aa^4+36Aa^2b^2+8Ab^4+24B^3b+32B^2ab^3)\operatorname{arctanh}(\sin(dx+c)) + \frac{1}{15}(32Aa^3b+40Aa^2b^2+8B^3a^4+60B^2a^2b^2+15B^2b^4)\tan(dx+c) + \frac{1}{16}(5Aa^4+36Aa^2b^2+8Ab^4+24B^3b+32B^2ab^3)\sec(dx+c)\tan(dx+c) + \frac{1}{15}a(16Aa^2b+13Ab^3+4B^3a^3+27B^2ab^2)\sec(dx+c)^2\tan(dx+c) + \frac{1}{120}a^2(25Aa^2+48Ab^2+72B^2ab)\sec(dx+c)^3\tan(dx+c) + \frac{1}{10}a(3Ab+2B^2a)(a+b\cos(dx+c))^2\sec(dx+c)^4\tan(dx+c) + \frac{1}{6}a^2(a+b\cos(dx+c))^3\sec(dx+c)^5\tan(dx+c) + \frac{1}{d}$

3.249.2 Mathematica [A] (verified)

Time = 1.96 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.75

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{15(5a^4A + 36a^2Ab^2 + 8Ab^4 + 24a^3bB + 32ab^3B) \operatorname{arctanh}(\sin(c + dx)) + \tan(c + dx) (240(4a^3Ab + 4aAb^3) + 15(5a^4A + 36a^2Ab^2 + 8Ab^4 + 24a^3bB + 32ab^3B))}{d}$$

input `Integrate[(a + b*Cos[c + d*x])^4*(A + B*Cos[c + d*x])*Sec[c + d*x]^7,x]`

output $(15(5a^4A + 36a^2Ab^2 + 8Ab^4 + 24a^3bB + 32a^2b^3B)\operatorname{ArcTanh}[\sin[c + dx]] + \tan[c + dx](240(4a^3Ab + 4aAb^3) + 15(5a^4A + 36a^2Ab^2 + 8Ab^4 + 24a^3bB + 32a^2b^3B)\sec[c + dx] + 10a^2(5a^2A + 36Ab^2 + 24a^2bB)\sec[c + dx]^3 + 40a^4A\sec[c + dx]^5 + 160a(4a^2Ab + 2Ab^3 + a^3B + 3a^2b^2B)\tan[c + dx]^2 + 48a^3(4Ab + aB)\tan[c + dx]^4))/(240*d)$

3.249.3 Rubi [A] (verified)

Time = 1.97 (sec) , antiderivative size = 302, normalized size of antiderivative = 0.93, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 3468, 3042, 3526, 3042, 3510, 25, 3042, 3500, 27, 3042, 3227, 3042, 4254, 24, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^7(c + dx)(a + b \cos(c + dx))^4 (A + B \cos(c + dx)) dx$$

↓ 3042

3.249. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^4 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^7} dx$$

↓ 3468

$$\frac{1}{6} \int (a + b \cos(c + dx))^2 (2b(aA + 3bB) \cos^2(c + dx) + (5Aa^2 + 12bBa + 6Ab^2) \cos(c + dx) + 3a(3Ab + 2aB)) \sec^6(c + dx) dx + \frac{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3}{6d}$$

↓ 3042

$$\frac{1}{6} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (2b(aA + 3bB) \sin(c + dx + \frac{\pi}{2})^2 + (5Aa^2 + 12bBa + 6Ab^2) \sin(c + dx + \frac{\pi}{2}) + 3a) \sin(c + dx + \frac{\pi}{2})^6}{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3} dx + \frac{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3}{6d}$$

↓ 3526

$$\frac{1}{6} \left(\frac{1}{5} \int (a + b \cos(c + dx)) (2b(6Ba^2 + 14Aba + 15b^2B) \cos^2(c + dx) + (24Ba^3 + 71Aba^2 + 90b^2Ba + 30Ab^3) \cos(c + dx) + aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3) dx + \frac{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3}{6d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) (2b(6Ba^2 + 14Aba + 15b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (24Ba^3 + 71Aba^2 + 90b^2Ba + 30Ab^3) \sin(c + dx + \frac{\pi}{2}) + aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3) \sin(c + dx + \frac{\pi}{2})^5}{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3} dx + \frac{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3}{6d} \right)$$

↓ 3510

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{a^2(25a^2A + 72abB + 48Ab^2) \tan(c + dx) \sec^3(c + dx)}{4d} - \frac{1}{4} \int -((8b^2(6Ba^2 + 14Aba + 15b^2B) \cos^2(c + dx) + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \cos(c + dx) + aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3) dx + \frac{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3}{6d} \right) \right)$$

↓ 25

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \int (8b^2(6Ba^2 + 14Aba + 15b^2B) \cos^2(c + dx) + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \cos(c + dx) + aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3) dx + \frac{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3}{6d} \right) \right)$$

3.249. $\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \int \frac{8b^2(6Ba^2 + 14Aba + 15b^2B) \sin(c + dx + \frac{\pi}{2})^2 + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \sin(c + dx + \frac{\pi}{2})^4}{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3} dx \right) \right)$$

↓ 3500

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int 3(15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) + 8(8Ba^4 + 32Aba^3 + 60b^2Ba^2 + 40Ab^3a + 15b^4B)) \sec^3(c + dx) dx + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \sin(c + dx + \frac{\pi}{2})^4}{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3} \right) \right) \right)$$

↓ 27

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(\int (15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) + 8(8Ba^4 + 32Aba^3 + 60b^2Ba^2 + 40Ab^3a + 15b^4B)) \sec^3(c + dx) dx + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \sin(c + dx + \frac{\pi}{2})^4}{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3} \right) \right) \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(\int \frac{15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) + 8(8Ba^4 + 32Aba^3 + 60b^2Ba^2 + 40Ab^3a + 15b^4B)}{\sin(c + dx + \frac{\pi}{2})^3} dx + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \sin(c + dx + \frac{\pi}{2})^4}{aA \tan(c + dx) \sec^5(c + dx)(a + b \cos(c + dx))^3} \right) \right) \right)$$

↓ 3227

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(15(5a^4A + 24a^3bB + 36a^2Ab^2 + 32ab^3B + 8Ab^4) \int \sec^3(c + dx) dx + 8(8a^4B + 32a^3Ab + 60a^2b^2B + 40aAb^3 + 15b^4B) \sin(c + dx + \frac{\pi}{2})^4 \right) \right) \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(8(8a^4B + 32a^3Ab + 60a^2b^2B + 40aAb^3 + 15b^4B) \int \csc(c + dx + \frac{\pi}{2})^2 dx + 15(5a^4A + 24a^3bB + 36a^2Ab^2 + 32ab^3B + 8Ab^4) \int \sec^3(c + dx) dx + 15(5Aa^4 + 24bBa^3 + 36Ab^2a^2 + 32b^3Ba + 8Ab^4) \sin(c + dx + \frac{\pi}{2})^4 \right) \right) \right)$$

↓ 4254

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(-\frac{8(8a^4B + 32a^3Ab + 60a^2b^2B + 40aAb^3 + 15b^4B) \int 1d(-\tan(c+dx))}{d} + 15(5a^4A + 24a^3bB + 36a^2Ab^2 + 32ab^3B + 8Ab^4) \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx + \frac{8a(4a^3B + 16a^2Ab + 27ab^2B + 13Ab^3) \tan(c+dx)}{d} \right) \right) \right) \frac{aA \tan(c+dx) \sec^5(c+dx)(a+b \cos(c+dx))^3}{6d}$$

↓ 24

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(15(5a^4A + 24a^3bB + 36a^2Ab^2 + 32ab^3B + 8Ab^4) \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx + \frac{8a(4a^3B + 16a^2Ab + 27ab^2B + 13Ab^3) \tan(c+dx)}{d} \right) \right) \right) \frac{aA \tan(c+dx) \sec^5(c+dx)(a+b \cos(c+dx))^3}{6d}$$

↓ 4255

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(15(5a^4A + 24a^3bB + 36a^2Ab^2 + 32ab^3B + 8Ab^4) \left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{8a(4a^3B + 16a^2Ab + 27ab^2B + 13Ab^3) \tan(c+dx)}{d} \right) \right) \right) \frac{aA \tan(c+dx) \sec^5(c+dx)(a+b \cos(c+dx))^3}{6d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{1}{4} \left(15(5a^4A + 24a^3bB + 36a^2Ab^2 + 32ab^3B + 8Ab^4) \left(\frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx) \sec(c+dx)}{2d} \right) + \frac{8a(4a^3B + 16a^2Ab + 27ab^2B + 13Ab^3) \tan(c+dx)}{d} \right) \right) \right) \frac{aA \tan(c+dx) \sec^5(c+dx)(a+b \cos(c+dx))^3}{6d}$$

↓ 4257

$$\frac{1}{6} \left(\frac{1}{5} \left(\frac{a^2(25a^2A + 72abB + 48Ab^2) \tan(c+dx) \sec^3(c+dx)}{4d} + \frac{1}{4} \left(\frac{8a(4a^3B + 16a^2Ab + 27ab^2B + 13Ab^3) \tan(c+dx)}{d} \right) \right) \right) \frac{aA \tan(c+dx) \sec^5(c+dx)(a+b \cos(c+dx))^3}{6d}$$

input `Int[(a + b*cos[c + d*x])^4*(A + B*cos[c + d*x])*Sec[c + d*x]^7,x]`

```
output (a*A*(a + b*Cos[c + d*x])^3*Sec[c + d*x]^5*Tan[c + d*x]/(6*d) + ((3*a*(3*
A*b + 2*a*B)*(a + b*Cos[c + d*x])^2*Sec[c + d*x]^4*Tan[c + d*x])/(5*d) + (
(a^2*(25*a^2*A + 48*A*b^2 + 72*a*b*B)*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) +
((8*(32*a^3*A*b + 40*a*A*b^3 + 8*a^4*B + 60*a^2*b^2*B + 15*b^4*B)*Tan[c +
d*x])/d + (8*a*(16*a^2*A*b + 13*A*b^3 + 4*a^3*B + 27*a*b^2*B)*Sec[c + d*x
]^2*Tan[c + d*x])/d + 15*(5*a^4*A + 36*a^2*A*b^2 + 8*A*b^4 + 24*a^3*b*B +
32*a*b^3*B)*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*
d)))/4)/5)/6
```

3.249.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)])], x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int
[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3468 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c
+ d*SIN[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n +
1)*(c^2 - d^2)) Int[(a + b*SIN[e + f*x])^(m - 2)*(c + d*SIN[e + f*x])^(n
+ 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a
*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1)
- a*(b*c - a*d)*(B*c - A*d)*(n + 2))*SIN[e + f*x] + b*(d*(A*b*c + a*B*c -
a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*SIN[e + f*x]^2, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2
, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]*(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.249.4 Maple [A] (verified)

Time = 7.46 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.85

method	result
parts	$a^4 A \left(- \left(- \frac{(\sec^5(dx+c))}{6} - \frac{5(\sec^3(dx+c))}{24} - \frac{5 \sec(dx+c)}{16} \right) \tan(dx+c) + \frac{5 \ln(\sec(dx+c) + \tan(dx+c))}{16} \right) + \frac{(A b^4 + 4 B a b^3) \left(\frac{\sec^4(dx+c)}{5} \right)}{d}$
derivativedivides	$a^4 A \left(- \left(- \frac{(\sec^5(dx+c))}{6} - \frac{5(\sec^3(dx+c))}{24} - \frac{5 \sec(dx+c)}{16} \right) \tan(dx+c) + \frac{5 \ln(\sec(dx+c) + \tan(dx+c))}{16} \right) - B a^4 \left(- \frac{8}{15} - \frac{(\sec^4(dx+c))}{5} \right)$
default	$a^4 A \left(- \left(- \frac{(\sec^5(dx+c))}{6} - \frac{5(\sec^3(dx+c))}{24} - \frac{5 \sec(dx+c)}{16} \right) \tan(dx+c) + \frac{5 \ln(\sec(dx+c) + \tan(dx+c))}{16} \right) - B a^4 \left(- \frac{8}{15} - \frac{(\sec^4(dx+c))}{5} \right)$
parallelrisch	$-75(a^4 A + \frac{36}{5} A a^2 b^2 + \frac{8}{5} A b^4 + \frac{24}{5} B a^3 b + \frac{32}{5} B a b^3) (\cos(6dx+6c) + 6 \cos(4dx+4c) + 15 \cos(2dx+2c) + 10) \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) \right)$
risch	Expression too large to display

input `int((a+cos(d*x+c)*b)^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x,method=_RETURNVERBOSE)`

output `a^4*A/d*(-(-1/6*sec(d*x+c)^5-5/24*sec(d*x+c)^3-5/16*sec(d*x+c))*tan(d*x+c)+5/16*ln(sec(d*x+c)+tan(d*x+c)))+(A*b^4+4*B*a*b^3)/d*(1/2*sec(d*x+c)*tan(d*x+c)+1/2*ln(sec(d*x+c)+tan(d*x+c)))-(4*A*a*b^3+6*B*a^2*b^2)/d*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)+(6*A*a^2*b^2+4*B*a^3*b)/d*(-(-1/4*sec(d*x+c)^3-3/8*sec(d*x+c))*tan(d*x+c)+3/8*ln(sec(d*x+c)+tan(d*x+c)))-(4*A*a^3*b+B*a^4)/d*(-8/15-1/5*sec(d*x+c)^4-4/15*sec(d*x+c)^2)*tan(d*x+c)+B*b^4/d*tan(d*x+c)`

3.249.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.01

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{15(5Aa^4 + 24Ba^3b + 36Aa^2b^2 + 32Bab^3 + 8Ab^4) \cos(dx + c)^6 \log(\sin(dx + c) + 1) - 15(5Aa^4 + 24Ba^3b + 36Aa^2b^2 + 32Bab^3 + 8Ab^4) \cos(dx + c)^6 \log(-\sin(dx + c) + 1) + 2(16(B^2a^4 + 32Aa^3b + 60B^2a^2b^2 + 40Aa^2b^3 + 15B^2b^4) \cos(dx + c)^5 + 40Aa^4 + 15(5Aa^4 + 24Ba^3b + 36Aa^2b^2 + 32Ba^2b^3 + 8Aa^2b^4) \cos(dx + c)^4 + 32(2Ba^4 + 8Aa^3b + 15Ba^2b^2 + 10Aa^2b^3) \cos(dx + c)^3 + 10(5Aa^4 + 24Ba^3b + 36Aa^2b^2) \cos(dx + c)^2 + 48(Ba^4 + 4Aa^3b) \cos(dx + c)) \sin(dx + c)}{(d \cos(dx + c))^6}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x, algorithm="fricas")`

output `1/480*(15*(5*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*cos(d*x + c)^6*log(sin(d*x + c) + 1) - 15*(5*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*cos(d*x + c)^6*log(-sin(d*x + c) + 1) + 2*(16*(8*B*a^4 + 32*A*a^3*b + 60*B*a^2*b^2 + 40*A*a*b^3 + 15*B*b^4)*cos(d*x + c)^5 + 40*A*a^4 + 15*(5*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*cos(d*x + c)^4 + 32*(2*B*a^4 + 8*A*a^3*b + 15*B*a^2*b^2 + 10*A*a*b^3)*cos(d*x + c)^3 + 10*(5*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2)*cos(d*x + c)^2 + 48*(B*a^4 + 4*A*a^3*b)*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^6)`

3.249.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**4*(A+B*cos(d*x+c))*sec(d*x+c)**7,x)`

output `Timed out`

3.249.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 474, normalized size of antiderivative = 1.46

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{32 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 + 15 \tan(dx + c)) Ba^4 + 128 (3 \tan(dx + c)^5 + 10 \tan(dx + c)^3 -$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x, algorithm="maxima")`

output `1/480*(32*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*B*a^4 + 128*(3*tan(d*x + c)^5 + 10*tan(d*x + c)^3 + 15*tan(d*x + c))*A*a^3*b + 960*(tan(d*x + c)^3 + 3*tan(d*x + c))*B*a^2*b^2 + 640*(tan(d*x + c)^3 + 3*tan(d*x + c))*A*a*b^3 - 5*A*a^4*(2*(15*sin(d*x + c)^5 - 40*sin(d*x + c)^3 + 33*sin(d*x + c))/(sin(d*x + c)^6 - 3*sin(d*x + c)^4 + 3*sin(d*x + c)^2 - 1) - 15*log(sin(d*x + c) + 1) + 15*log(sin(d*x + c) - 1)) - 120*B*a^3*b*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 180*A*a^2*b^2*(2*(3*sin(d*x + c)^3 - 5*sin(d*x + c))/(sin(d*x + c)^4 - 2*sin(d*x + c)^2 + 1) - 3*log(sin(d*x + c) + 1) + 3*log(sin(d*x + c) - 1)) - 480*B*a*b^3*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) - 120*A*b^4*(2*sin(d*x + c)/(sin(d*x + c)^2 - 1) - log(sin(d*x + c) + 1) + log(sin(d*x + c) - 1)) + 480*B*b^4*tan(d*x + c))/d`

3.249.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1186 vs. 2(310) = 620.

Time = 0.37 (sec) , antiderivative size = 1186, normalized size of antiderivative = 3.66

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx = \text{Too large to display}$$

input `integrate((a+b*cos(d*x+c))^4*(A+B*cos(d*x+c))*sec(d*x+c)^7,x, algorithm="giac")`

```

output 1/240*(15*(5*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2 + 32*B*a*b^3 + 8*A*b^4)*log
(abs(tan(1/2*d*x + 1/2*c) + 1)) - 15*(5*A*a^4 + 24*B*a^3*b + 36*A*a^2*b^2
+ 32*B*a*b^3 + 8*A*b^4)*log(abs(tan(1/2*d*x + 1/2*c) - 1)) + 2*(165*A*a^4*
tan(1/2*d*x + 1/2*c)^11 - 240*B*a^4*tan(1/2*d*x + 1/2*c)^11 - 960*A*a^3*b*
tan(1/2*d*x + 1/2*c)^11 + 600*B*a^3*b*tan(1/2*d*x + 1/2*c)^11 + 900*A*a^2*
b^2*tan(1/2*d*x + 1/2*c)^11 - 1440*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^11 - 960
*A*a*b^3*tan(1/2*d*x + 1/2*c)^11 + 480*B*a*b^3*tan(1/2*d*x + 1/2*c)^11 + 1
20*A*b^4*tan(1/2*d*x + 1/2*c)^11 - 240*B*b^4*tan(1/2*d*x + 1/2*c)^11 + 25*
A*a^4*tan(1/2*d*x + 1/2*c)^9 + 560*B*a^4*tan(1/2*d*x + 1/2*c)^9 + 2240*A*a
^3*b*tan(1/2*d*x + 1/2*c)^9 - 840*B*a^3*b*tan(1/2*d*x + 1/2*c)^9 - 1260*A*
a^2*b^2*tan(1/2*d*x + 1/2*c)^9 + 5280*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^9 + 3
520*A*a*b^3*tan(1/2*d*x + 1/2*c)^9 - 1440*B*a*b^3*tan(1/2*d*x + 1/2*c)^9 -
360*A*b^4*tan(1/2*d*x + 1/2*c)^9 + 1200*B*b^4*tan(1/2*d*x + 1/2*c)^9 + 45
0*A*a^4*tan(1/2*d*x + 1/2*c)^7 - 1248*B*a^4*tan(1/2*d*x + 1/2*c)^7 - 4992*
A*a^3*b*tan(1/2*d*x + 1/2*c)^7 + 240*B*a^3*b*tan(1/2*d*x + 1/2*c)^7 + 360*
A*a^2*b^2*tan(1/2*d*x + 1/2*c)^7 - 8640*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^7 -
5760*A*a*b^3*tan(1/2*d*x + 1/2*c)^7 + 960*B*a*b^3*tan(1/2*d*x + 1/2*c)^7
+ 240*A*b^4*tan(1/2*d*x + 1/2*c)^7 - 2400*B*b^4*tan(1/2*d*x + 1/2*c)^7 + 4
50*A*a^4*tan(1/2*d*x + 1/2*c)^5 + 1248*B*a^4*tan(1/2*d*x + 1/2*c)^5 + 4992
*A*a^3*b*tan(1/2*d*x + 1/2*c)^5 + 240*B*a^3*b*tan(1/2*d*x + 1/2*c)^5 + ...

```

3.249.9 Mupad [B] (verification not implemented)

Time = 4.18 (sec) , antiderivative size = 706, normalized size of antiderivative = 2.18

$$\int (a + b \cos(c + dx))^4 (A + B \cos(c + dx)) \sec^7(c + dx) dx$$

$$= \frac{\operatorname{atanh}\left(\frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{5Aa^4}{16} + \frac{3Ba^3b}{2} + \frac{9Aa^2b^2}{4} + 2Bab^3 + \frac{Ab^4}{2}\right)}{\frac{5Aa^4}{4} + 6Ba^3b + 9Aa^2b^2 + 8Bab^3 + 2Ab^4}\right) \left(\frac{5Aa^4}{8} + 3Ba^3b + \frac{9Aa^2b^2}{2} + 4Bab^3 + Ab^4\right)}{d}$$

$$+ \frac{\left(\frac{11Aa^4}{8} + Ab^4 - 2Ba^4 - 2Bb^4 + \frac{15Aa^2b^2}{2} - 12Ba^2b^2 - 8Aab^3 - 8Aa^3b + 4Bab^3 + 5Ba^3b\right) \tan}{\dots}$$

```

input int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^4)/cos(c + d*x)^7,x)

```

output $(\operatorname{atanh}((4*\tan(c/2 + (d*x)/2)*((5*A*a^4)/16 + (A*b^4)/2 + (9*A*a^2*b^2)/4 + 2*B*a*b^3 + (3*B*a^3*b)/2)))/((5*A*a^4)/4 + 2*A*b^4 + 9*A*a^2*b^2 + 8*B*a*b^3 + 6*B*a^3*b))/d + (\tan(c/2 + (d*x)/2)*((5*A*a^4)/8 + A*b^4 + (9*A*a^2*b^2)/2 + 4*B*a*b^3 + 3*B*a^3*b))/d + (\tan(c/2 + (d*x)/2)*((11*A*a^4)/8 + A*b^4 + 2*B*a^4 + 2*B*b^4 + (15*A*a^2*b^2)/2 + 12*B*a^2*b^2 + 8*A*a*b^3 + 8*A*a^3*b + 4*B*a*b^3 + 5*B*a^3*b) + \tan(c/2 + (d*x)/2)^{11}*((11*A*a^4)/8 + A*b^4 - 2*B*a^4 - 2*B*b^4 + (15*A*a^2*b^2)/2 - 12*B*a^2*b^2 - 8*A*a*b^3 - 8*A*a^3*b + 4*B*a*b^3 + 5*B*a^3*b) - \tan(c/2 + (d*x)/2)^3*(3*A*b^4 - (5*A*a^4)/24 + (14*B*a^4)/3 + 10*B*b^4 + (21*A*a^2*b^2)/2 + 44*B*a^2*b^2 + (88*A*a*b^3)/3 + (56*A*a^3*b)/3 + 12*B*a*b^3 + 7*B*a^3*b) + \tan(c/2 + (d*x)/2)^9*((5*A*a^4)/24 - 3*A*b^4 + (14*B*a^4)/3 + 10*B*b^4 - (21*A*a^2*b^2)/2 + 44*B*a^2*b^2 + (88*A*a*b^3)/3 + (56*A*a^3*b)/3 - 12*B*a*b^3 - 7*B*a^3*b) + \tan(c/2 + (d*x)/2)^5*((15*A*a^4)/4 + 2*A*b^4 + (52*B*a^4)/5 + 20*B*b^4 + 3*A*a^2*b^2 + 72*B*a^2*b^2 + 48*A*a*b^3 + (208*A*a^3*b)/5 + 8*B*a*b^3 + 2*B*a^3*b) + \tan(c/2 + (d*x)/2)^7*((15*A*a^4)/4 + 2*A*b^4 - (52*B*a^4)/5 - 20*B*b^4 + 3*A*a^2*b^2 - 72*B*a^2*b^2 - 48*A*a*b^3 - (208*A*a^3*b)/5 + 8*B*a*b^3 + 2*B*a^3*b))/(d*(15*\tan(c/2 + (d*x)/2)^4 - 6*\tan(c/2 + (d*x)/2)^2 - 20*\tan(c/2 + (d*x)/2)^6 + 15*\tan(c/2 + (d*x)/2)^8 - 6*\tan(c/2 + (d*x)/2)^{10} + \tan(c/2 + (d*x)/2)^{12} + 1))$

3.250 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.250.1 Optimal result 2314
 3.250.2 Mathematica [A] (verified) 2315
 3.250.3 Rubi [A] (verified) 2315
 3.250.4 Maple [A] (verified) 2319
 3.250.5 Fricas [A] (verification not implemented) 2320
 3.250.6 Sympy [F(-1)] 2320
 3.250.7 Maxima [F(-2)] 2321
 3.250.8 Giac [B] (verification not implemented) 2321
 3.250.9 Mupad [B] (verification not implemented) 2322

3.250.1 Optimal result

Integrand size = 31, antiderivative size = 178

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{(2a^2+b^2)(Ab-aB)x}{2b^4} - \frac{2a^3(Ab-aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{\sqrt{a-b}b^4\sqrt{a+bd}} - \frac{(3aAb-3a^2B-2b^2B) \sin(c+dx)}{3b^3d} + \frac{(Ab-aB) \cos(c+dx) \sin(c+dx)}{2b^2d} + \frac{B \cos^2(c+dx) \sin(c+dx)}{3bd}$$

output

```
1/2*(2*a^2+b^2)*(A*b-B*a)*x/b^4-1/3*(3*A*a*b-3*B*a^2-2*B*b^2)*sin(d*x+c)/b^3/d+1/2*(A*b-B*a)*cos(d*x+c)*sin(d*x+c)/b^2/d+1/3*B*cos(d*x+c)^2*sin(d*x+c)/b/d-2*a^3*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b^4/d/(a-b)^(1/2)/(a+b)^(1/2)
```

3.250.2 Mathematica [A] (verified)

Time = 0.94 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.85

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \frac{6(2a^2+b^2)(Ab-aB)(c+dx) - \frac{24a^3(-Ab+aB)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + 3b(-4aAb+4a^2B+3b^2B)\sin(c+dx)}{12b^4d}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(6*(2*a^2 + b^2)*(A*b - a*B)*(c + d*x) - (24*a^3*(-(A*b) + a*B)*ArcTanh[(((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2])]/Sqrt[-a^2 + b^2] + 3*b*(-4*a*A*b + 4*a^2*B + 3*b^2*B)*Sin[c + d*x] + 3*b^2*(A*b - a*B)*Sin[2*(c + d*x)] + b^3*B*Sin[3*(c + d*x)])/(12*b^4*d)`

3.250.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.10, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3469, 3042, 3528, 3042, 3502, 27, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^3 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} dx$$

$$\downarrow \text{3469}$$

$$\frac{\int \frac{\cos(c+dx)(3(Ab-aB)\cos^2(c+dx)+2bB\cos(c+dx)+2aB)}{a+b\cos(c+dx)} dx}{3b} + \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd}$$

$$\downarrow \text{3042}$$

$$\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)\left(3(Ab-aB)\sin\left(c+dx+\frac{\pi}{2}\right)^2+2bB\sin\left(c+dx+\frac{\pi}{2}\right)+2aB\right)}{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} dx}{3b} + \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd}$$

3.250. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3528} \\
 & \frac{\int \frac{-2(-3Ba^2+3Aba-2b^2B)\cos^2(c+dx)+b(3Ab+aB)\cos(c+dx)+3a(Ab-aB)}{a+b\cos(c+dx)} dx}{2b} + \frac{3(Ab-aB)\sin(c+dx)\cos(c+dx)}{2bd} + \\
 & \quad \frac{3b}{3bd} \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd} \\
 & \downarrow \text{3042} \\
 & \frac{\int \frac{-2(-3Ba^2+3Aba-2b^2B)\sin(c+dx+\frac{\pi}{2})^2+b(3Ab+aB)\sin(c+dx+\frac{\pi}{2})+3a(Ab-aB)}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{2b} + \frac{3(Ab-aB)\sin(c+dx)\cos(c+dx)}{2bd} + \\
 & \quad \frac{3b}{3bd} \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd} \\
 & \downarrow \text{3502} \\
 & \frac{\int \frac{3(ab(Ab-aB)+(2a^2+b^2)\cos(c+dx)(Ab-aB))}{a+b\cos(c+dx)} dx}{2b} - \frac{2(-3a^2B+3aAb-2b^2B)\sin(c+dx)}{bd} + \frac{3(Ab-aB)\sin(c+dx)\cos(c+dx)}{2bd} + \\
 & \quad \frac{3b}{3bd} \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd} \\
 & \downarrow \text{27} \\
 & \frac{3\int \frac{ab(Ab-aB)+(2a^2+b^2)\cos(c+dx)(Ab-aB)}{a+b\cos(c+dx)} dx}{2b} - \frac{2(-3a^2B+3aAb-2b^2B)\sin(c+dx)}{bd} + \frac{3(Ab-aB)\sin(c+dx)\cos(c+dx)}{2bd} + \\
 & \quad \frac{3b}{3bd} \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd} \\
 & \downarrow \text{3042} \\
 & \frac{3\int \frac{ab(Ab-aB)+(2a^2+b^2)\sin(c+dx+\frac{\pi}{2})(Ab-aB)}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{2b} - \frac{2(-3a^2B+3aAb-2b^2B)\sin(c+dx)}{bd} + \frac{3(Ab-aB)\sin(c+dx)\cos(c+dx)}{2bd} + \\
 & \quad \frac{3b}{3bd} \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd} \\
 & \downarrow \text{3214} \\
 & \frac{3\left(\frac{x(2a^2+b^2)(Ab-aB)}{b} - \frac{2a^3(Ab-aB)\int \frac{1}{a+b\cos(c+dx)} dx}{b}\right)}{2b} - \frac{2(-3a^2B+3aAb-2b^2B)\sin(c+dx)}{bd} + \frac{3(Ab-aB)\sin(c+dx)\cos(c+dx)}{2bd} + \\
 & \quad \frac{3b}{3bd} \frac{B\sin(c+dx)\cos^2(c+dx)}{3bd} \\
 & \downarrow \text{3042}
 \end{aligned}$$

3.250. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

$$\frac{\frac{3 \left(\frac{x(2a^2+b^2)(Ab-aB)}{b} - \frac{2a^3(Ab-aB) \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} \right)}{2b} - \frac{2(-3a^2B+3aAb-2b^2B) \sin(c+dx)}{bd} + \frac{3(Ab-aB) \sin(c+dx) \cos(c+dx)}{2bd}}{\frac{3b}{3bd} \frac{B \sin(c+dx) \cos^2(c+dx)}}{\downarrow 3138}$$

$$\frac{\frac{3 \left(\frac{x(2a^2+b^2)(Ab-aB)}{b} - \frac{4a^3(Ab-aB) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx))+a+b} d \tan(\frac{1}{2}(c+dx))}{bd} \right)}{2b} - \frac{2(-3a^2B+3aAb-2b^2B) \sin(c+dx)}{bd} + \frac{3(Ab-aB) \sin(c+dx) \cos(c+dx)}{2bd}}{\frac{3b}{3bd} \frac{B \sin(c+dx) \cos^2(c+dx)}}{\downarrow 218}$$

$$\frac{\frac{3 \left(\frac{x(2a^2+b^2)(Ab-aB)}{b} - \frac{4a^3(Ab-aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{bd\sqrt{a-b}\sqrt{a+b}} \right)}{2b} - \frac{2(-3a^2B+3aAb-2b^2B) \sin(c+dx)}{bd} + \frac{3(Ab-aB) \sin(c+dx) \cos(c+dx)}{2bd}}{\frac{3b}{3bd} \frac{B \sin(c+dx) \cos^2(c+dx)}}{+}$$

```
input Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]
```

```
output (B*Cos[c + d*x]^2*Sin[c + d*x])/(3*b*d) + ((3*(A*b - a*B)*Cos[c + d*x]*Sin[c + d*x])/(2*b*d) + ((3*((2*a^2 + b^2)*(A*b - a*B)*x)/b - (4*a^3*(A*b - a*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d))/b - (2*(3*a*A*b - 3*a^2*B - 2*b^2*B)*Sin[c + d*x])/(b*d))/(3*b)
```

3.250.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Ssin[e + f*x
])^m*((c + d*Ssin[e + f*x])^(n + 1)/(d*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Ssin[e + f*x])^(m - 1)*(c + d*Ssin[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

3.250.4 Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.35

method	result
derivativedivides	$-\frac{2a^3(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^4 \sqrt{(a-b)(a+b)}} + \frac{2\left((-Aa b^2 - \frac{1}{2}A b^3 + B a^2 b + \frac{1}{2}B a b^2 + B b^3\right) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-2Aa b^2 + 2B a^2 b + \frac{2}{3} B b^3)}{(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))}$
default	$-\frac{2a^3(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^4 \sqrt{(a-b)(a+b)}} + \frac{2\left((-Aa b^2 - \frac{1}{2}A b^3 + B a^2 b + \frac{1}{2}B a b^2 + B b^3\right) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-2Aa b^2 + 2B a^2 b + \frac{2}{3} B b^3)}{(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))}$
risch	$\frac{x A a^2}{b^3} + \frac{x A}{2b} - \frac{x B a^3}{b^4} - \frac{a B x}{2b^2} - \frac{i e^{i(dx+c)} B a^2}{2b^3 d} + \frac{i e^{-i(dx+c)} B a^2}{2b^3 d} + \frac{i e^{i(dx+c)} A a}{2b^2 d} + \frac{3 i e^{-i(dx+c)} B}{8bd} - \frac{3 i e^{i(dx+c)} B}{8bd}$

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b), x, method=_RETURNVERBOSE
)
```

```
output 1/d*(-2*a^3*(A*B-B*a)/b^4/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2
*c)/((a-b)*(a+b))^(1/2))+2/b^4*(((A*a*b^2-1/2*A*b^3+B*a^2*b+1/2*B*a*b^2+B
*b^3)*tan(1/2*d*x+1/2*c)^5+(-2*A*a*b^2+2*B*a^2*b+2/3*B*b^3)*tan(1/2*d*x+1/
2*c)^3+(-A*a*b^2+B*a^2*b+B*b^3+1/2*A*b^3-1/2*B*a*b^2)*tan(1/2*d*x+1/2*c)))/
(1+tan(1/2*d*x+1/2*c)^2)^3+1/2*(2*A*a^2*b+A*b^3-2*B*a^3-B*a*b^2)*arctan(ta
n(1/2*d*x+1/2*c)))
```

$$3.250. \int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.250.5 Fricas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 541, normalized size of antiderivative = 3.04

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \left[\frac{3(2Ba^5 - 2Aa^4b - Ba^3b^2 + Aa^2b^3 - Bab^4 + Ab^5)dx - 3(Ba^4 - Aa^3b)\sqrt{-a^2 + b^2} \log\left(\frac{2ab\cos(dx+c) + (2a^2 - b^2)\cos(dx+c)^2 - 2\sqrt{-a^2 + b^2}(a\cos(dx+c) + b)\sin(dx+c) - a^2 + 2b^2}{(b^2\cos(dx+c)^2 + 2a*b*\cos(dx+c) + a^2)}\right) - (6Ba^4b - 6Aa^3b^2 - 2Ba^2b^3 + 6Aa*b^4 - 4Bb^5 + 2*(Ba^2b^3 - Bb^5)*\cos(dx+c)^2 - 3*(Ba^3b^2 - Aa^2b^3 - Ba*b^4 + Ab^5)*\cos(dx+c))*\sin(dx+c)}{(a^2b^4 - b^6)*d}, \right.$$

$$\left. \frac{3(2Ba^5 - 2Aa^4b - Ba^3b^2 + Aa^2b^3 - Bab^4 + Ab^5)dx - 6(Ba^4 - Aa^3b)\sqrt{a^2 - b^2} \arctan\left(-\frac{a\cos(dx+c)}{\sqrt{a^2 - b^2}\sin(dx+c)}\right) - (6Ba^4b - 6Aa^3b^2 - 2Ba^2b^3 + 6Aa*b^4 - 4Bb^5 + 2*(Ba^2b^3 - Bb^5)*\cos(dx+c)^2 - 3*(Ba^3b^2 - Aa^2b^3 - Ba*b^4 + Ab^5)*\cos(dx+c))*\sin(dx+c)}{(a^2b^4 - b^6)*d} \right]$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")
```

```
output [-1/6*(3*(2*B*a^5 - 2*A*a^4*b - B*a^3*b^2 + A*a^2*b^3 - B*a*b^4 + A*b^5)*d*x - 3*(B*a^4 - A*a^3*b)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - (6*B*a^4*b - 6*A*a^3*b^2 - 2*B*a^2*b^3 + 6*A*a*b^4 - 4*B*b^5 + 2*(B*a^2*b^3 - B*b^5)*cos(d*x + c)^2 - 3*(B*a^3*b^2 - A*a^2*b^3 - B*a*b^4 + A*b^5)*cos(d*x + c))*sin(d*x + c))/((a^2*b^4 - b^6)*d), -1/6*(3*(2*B*a^5 - 2*A*a^4*b - B*a^3*b^2 + A*a^2*b^3 - B*a*b^4 + A*b^5)*d*x - 6*(B*a^4 - A*a^3*b)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (6*B*a^4*b - 6*A*a^3*b^2 - 2*B*a^2*b^3 + 6*A*a*b^4 - 4*B*b^5 + 2*(B*a^2*b^3 - B*b^5)*cos(d*x + c)^2 - 3*(B*a^3*b^2 - A*a^2*b^3 - B*a*b^4 + A*b^5)*cos(d*x + c))*sin(d*x + c))/((a^2*b^4 - b^6)*d)]
```

3.250.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x)
```

```
output Timed out
```

3.250.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Exception raised: ValueError}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="max
ima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

3.250.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 360 vs. 2(161) = 322.

Time = 0.29 (sec) , antiderivative size = 360, normalized size of antiderivative = 2.02

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx =$$

$$\frac{3(2Ba^3-2Aa^2b+Bab^2-Ab^3)(dx+c)}{b^4} + \frac{12(Ba^4-Aa^3b)\left(\pi\left[\frac{dx+c}{2\pi}+\frac{1}{2}\right]\text{sgn}(-2a+2b)+\arctan\left(-\frac{a\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)-b\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{\sqrt{a^2-b^2}}\right)\right)}{\sqrt{a^2-b^2}b^4}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="gia
c")
```

```
output -1/6*(3*(2*B*a^3 - 2*A*a^2*b + B*a*b^2 - A*b^3)*(d*x + c)/b^4 + 12*(B*a^4
- A*a^3*b)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*
tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/(sqrt(a^2
- b^2)*b^4) - 2*(6*B*a^2*tan(1/2*d*x + 1/2*c)^5 - 6*A*a*b*tan(1/2*d*x + 1
/2*c)^5 + 3*B*a*b*tan(1/2*d*x + 1/2*c)^5 - 3*A*b^2*tan(1/2*d*x + 1/2*c)^5
+ 6*B*b^2*tan(1/2*d*x + 1/2*c)^5 + 12*B*a^2*tan(1/2*d*x + 1/2*c)^3 - 12*A*
a*b*tan(1/2*d*x + 1/2*c)^3 + 4*B*b^2*tan(1/2*d*x + 1/2*c)^3 + 6*B*a^2*tan(
1/2*d*x + 1/2*c) - 6*A*a*b*tan(1/2*d*x + 1/2*c) - 3*B*a*b*tan(1/2*d*x + 1/
2*c) + 3*A*b^2*tan(1/2*d*x + 1/2*c) + 6*B*b^2*tan(1/2*d*x + 1/2*c))/((tan(
1/2*d*x + 1/2*c)^2 + 1)^3*b^3))/d
```

$$3.250. \int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

3.250.9 Mupad [B] (verification not implemented)

Time = 5.32 (sec) , antiderivative size = 4568, normalized size of antiderivative = 25.66

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Too large to display}$$

```
input int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)),x)
```

```
output ((tan(c/2 + (d*x)/2)*(A*b^2 + 2*B*a^2 + 2*B*b^2 - 2*A*a*b - B*a*b))/b^3 +
(tan(c/2 + (d*x)/2)^5*(2*B*a^2 - A*b^2 + 2*B*b^2 - 2*A*a*b + B*a*b))/b^3 +
(4*tan(c/2 + (d*x)/2)^3*(3*B*a^2 + B*b^2 - 3*A*a*b))/(3*b^3))/(d*(3*tan(c
/2 + (d*x)/2)^2 + 3*tan(c/2 + (d*x)/2)^4 + tan(c/2 + (d*x)/2)^6 + 1)) + (a
tan(((2*a^2 + b^2)*(A*b - B*a)*((8*tan(c/2 + (d*x)/2)*(A^2*b^9 - 8*B^2*a^
9 - 3*A^2*a*b^8 + 16*B^2*a^8*b + 7*A^2*a^2*b^7 - 13*A^2*a^3*b^6 + 16*A^2*a
^4*b^5 - 16*A^2*a^5*b^4 + 16*A^2*a^6*b^3 - 8*A^2*a^7*b^2 + B^2*a^2*b^7 - 3
*B^2*a^3*b^6 + 7*B^2*a^4*b^5 - 13*B^2*a^5*b^4 + 16*B^2*a^6*b^3 - 16*B^2*a^
7*b^2 - 2*A*B*a*b^8 + 16*A*B*a^8*b + 6*A*B*a^2*b^7 - 14*A*B*a^3*b^6 + 26*A
*B*a^4*b^5 - 32*A*B*a^5*b^4 + 32*A*B*a^6*b^3 - 32*A*B*a^7*b^2))/b^6 + ((2*
a^2 + b^2)*(A*b - B*a)*((8*(2*A*b^13 + 2*A*a^2*b^11 - 6*A*a^3*b^10 + 4*A*a
^4*b^9 + 2*B*a^2*b^11 - 2*B*a^3*b^10 + 6*B*a^4*b^9 - 4*B*a^5*b^8 - 2*A*a*b
^12 - 2*B*a*b^12))/b^9 - (tan(c/2 + (d*x)/2)*(2*a^2 + b^2)*(A*b - B*a)*(8*
a*b^10 - 16*a^2*b^9 + 8*a^3*b^8)*4i)/b^10)*1i)/(2*b^4)))/(2*b^4) + ((2*a^2
+ b^2)*(A*b - B*a)*((8*tan(c/2 + (d*x)/2)*(A^2*b^9 - 8*B^2*a^9 - 3*A^2*a*
b^8 + 16*B^2*a^8*b + 7*A^2*a^2*b^7 - 13*A^2*a^3*b^6 + 16*A^2*a^4*b^5 - 16*
A^2*a^5*b^4 + 16*A^2*a^6*b^3 - 8*A^2*a^7*b^2 + B^2*a^2*b^7 - 3*B^2*a^3*b^6
+ 7*B^2*a^4*b^5 - 13*B^2*a^5*b^4 + 16*B^2*a^6*b^3 - 16*B^2*a^7*b^2 - 2*A*
B*a*b^8 + 16*A*B*a^8*b + 6*A*B*a^2*b^7 - 14*A*B*a^3*b^6 + 26*A*B*a^4*b^5 -
32*A*B*a^5*b^4 + 32*A*B*a^6*b^3 - 32*A*B*a^7*b^2))/b^6 - ((2*a^2 + b^2...
```

3.251 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.251.1 Optimal result 2323
 3.251.2 Mathematica [A] (verified) 2323
 3.251.3 Rubi [A] (verified) 2324
 3.251.4 Maple [A] (verified) 2326
 3.251.5 Fricas [A] (verification not implemented) 2327
 3.251.6 Sympy [B] (verification not implemented) 2328
 3.251.7 Maxima [F(-2)] 2329
 3.251.8 Giac [A] (verification not implemented) 2330
 3.251.9 Mupad [B] (verification not implemented) 2330

3.251.1 Optimal result

Integrand size = 31, antiderivative size = 134

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = -\frac{(2aAb - 2a^2B - b^2B)x}{2b^3} + \frac{2a^2(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{\sqrt{a-b}b^3\sqrt{a+bd}} + \frac{(Ab - aB) \sin(c+dx)}{b^2d} + \frac{B \cos(c+dx) \sin(c+dx)}{2bd}$$

```
output -1/2*(2*A*a*b-2*B*a^2-B*b^2)*x/b^3+(A*b-B*a)*sin(d*x+c)/b^2/d+1/2*B*cos(d*x+c)*sin(d*x+c)/b/d+2*a^2*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b^3/d/(a-b)^(1/2)/(a+b)^(1/2)
```

3.251.2 Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.90

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2(-2aAb + 2a^2B + b^2B)(c+dx) + \frac{8a^2(-Ab+aB) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + 4b(Ab - aB) \sin(c+dx) + b^2}{4b^3d}$$

3.251. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output $(2*(-2*a*A*b + 2*a^2*B + b^2*B)*(c + d*x) + (8*a^2*(-(A*b) + a*B)*ArcTanh[(a - b)*Tan[(c + d*x)/2]]/Sqrt[-a^2 + b^2])/Sqrt[-a^2 + b^2] + 4*b*(A*b - a*B)*Sin[c + d*x] + b^2*B*Sin[2*(c + d*x)]/(4*b^3*d)$

3.251.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3469, 3042, 3502, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{a + b \cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{a + b \sin(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3469} \\
 & \frac{\int \frac{2(Ab - aB) \cos^2(c + dx) + bB \cos(c + dx) + aB}{a + b \cos(c + dx)} dx}{2b} + \frac{B \sin(c + dx) \cos(c + dx)}{2bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2(Ab - aB) \sin(c + dx + \frac{\pi}{2})^2 + bB \sin(c + dx + \frac{\pi}{2}) + aB}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{2b} + \frac{B \sin(c + dx) \cos(c + dx)}{2bd} \\
 & \quad \downarrow \text{3502} \\
 & \frac{\int \frac{abB - (-2Ba^2 + 2Aba - b^2B) \cos(c + dx)}{a + b \cos(c + dx)} dx}{2b} + \frac{2(Ab - aB) \sin(c + dx)}{bd} + \frac{B \sin(c + dx) \cos(c + dx)}{2bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{abB + (2Ba^2 - 2Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{2b} + \frac{2(Ab - aB) \sin(c + dx)}{bd} + \frac{B \sin(c + dx) \cos(c + dx)}{2bd} \\
 & \quad \downarrow \text{3214}
 \end{aligned}$$

3.251. $\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{a + b \cos(c + dx)} dx$

$$\begin{aligned}
 & \frac{2a^2(Ab-aB) \int \frac{1}{a+b \cos(c+dx)} dx - \frac{x(-2a^2B+2aAb-b^2B)}{b}}{2b} + \frac{2(Ab-aB) \sin(c+dx)}{bd} + \frac{B \sin(c+dx) \cos(c+dx)}{2bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2a^2(Ab-aB) \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx - \frac{x(-2a^2B+2aAb-b^2B)}{b}}{2b} + \frac{2(Ab-aB) \sin(c+dx)}{bd} + \frac{B \sin(c+dx) \cos(c+dx)}{2bd} \\
 & \quad \downarrow \text{3138} \\
 & \frac{4a^2(Ab-aB) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx)) - \frac{x(-2a^2B+2aAb-b^2B)}{b}}{2bd} + \frac{2(Ab-aB) \sin(c+dx)}{bd} + \\
 & \quad \frac{B \sin(c+dx) \cos(c+dx)}{2bd} \\
 & \quad \downarrow \text{218} \\
 & \frac{4a^2(Ab-aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - \frac{x(-2a^2B+2aAb-b^2B)}{b}}{bd\sqrt{a-b}\sqrt{a+b}} + \frac{2(Ab-aB) \sin(c+dx)}{bd} + \frac{B \sin(c+dx) \cos(c+dx)}{2bd}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(B*Cos[c + d*x]*Sin[c + d*x])/(2*b*d) + (((-(((2*a*A*b - 2*a^2*B - b^2*B)*x)/b) + (4*a^2*(A*b - a*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d))/b + (2*(A*b - a*B)*Sin[c + d*x])/(b*d))/(2*b)`

3.251.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.251.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.26

method	result
derivativedivides	$\frac{2a^2(Ab-Ba) \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^3\sqrt{(a-b)(a+b)}} - \frac{2\left(\frac{-Ab^2+Bab+\frac{1}{2}Bb^2}{(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right))}\right)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\left(\frac{-Ab^2+Bab-\frac{1}{2}Bb^2}{(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right))}\right)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{b^3} + (2A$
default	$\frac{2a^2(Ab-Ba) \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^3\sqrt{(a-b)(a+b)}} - \frac{2\left(\frac{-Ab^2+Bab+\frac{1}{2}Bb^2}{(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right))}\right)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\left(\frac{-Ab^2+Bab-\frac{1}{2}Bb^2}{(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right))}\right)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{b^3} + (2A$
risch	$-\frac{xAa}{b^2} + \frac{xBa^2}{b^3} + \frac{Bx}{2b} - \frac{ie^{i(dx+c)}A}{2bd} + \frac{ie^{i(dx+c)}Ba}{2b^2d} + \frac{ie^{-i(dx+c)}A}{2bd} - \frac{ie^{-i(dx+c)}Ba}{2b^2d} - \frac{a^2 \ln\left(e^{i(dx+c)} + ia^2 - \sqrt{-a^2 + b^2}\right)}{\sqrt{-a^2 + b^2}}$

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)
```

```
output 1/d*(2*a^2*(A*b-B*a)/b^3/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-2/b^3*(((A*b^2+B*a*b+1/2*B*b^2)*tan(1/2*d*x+1/2*c)^3+(-A*b^2+B*a*b-1/2*B*b^2)*tan(1/2*d*x+1/2*c))/(1+tan(1/2*d*x+1/2*c)^2)^2+1/2*(2*A*a*b-2*B*a^2-B*b^2)*arctan(tan(1/2*d*x+1/2*c))))
```

3.251.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 426, normalized size of antiderivative = 3.18

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \left[\frac{(2Ba^4 - 2Aa^3b - Ba^2b^2 + 2Aab^3 - Bb^4)dx + (Ba^3 - Aa^2b)\sqrt{-a^2 + b^2} \log\left(\frac{2ab\cos(dx+c) + (2a^2 - b^2)\cos(dx+c)}{b^2\cos(dx+c)}\right)}{\sqrt{-a^2 + b^2}} \right]$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")
```

output `[1/2*((2*B*a^4 - 2*A*a^3*b - B*a^2*b^2 + 2*A*a*b^3 - B*b^4)*d*x + (B*a^3 - A*a^2*b)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c))^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - (2*B*a^3*b - 2*A*a^2*b^2 - 2*B*a*b^3 + 2*A*b^4 - (B*a^2*b^2 - B*b^4)*cos(d*x + c))*sin(d*x + c))/((a^2*b^3 - b^5)*d), 1/2*((2*B*a^4 - 2*A*a^3*b - B*a^2*b^2 + 2*A*a*b^3 - B*b^4)*d*x - 2*(B*a^3 - A*a^2*b)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (2*B*a^3*b - 2*A*a^2*b^2 - 2*B*a*b^3 + 2*A*b^4 - (B*a^2*b^2 - B*b^4)*cos(d*x + c))*sin(d*x + c))/((a^2*b^3 - b^5)*d)]`

3.251.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10409 vs. $2(116) = 232$.

Time = 173.25 (sec) , antiderivative size = 10409, normalized size of antiderivative = 77.68

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)), x)`

output `Piecewise((zoo*x*(A + B*cos(c))*cos(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (-2*A*d*x**tan(c/2 + d*x/2)**4/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) - 4*A*d*x**tan(c/2 + d*x/2)**2/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) - 2*A*d*x/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) + 2*A*tan(c/2 + d*x/2)**5/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) + 8*A*tan(c/2 + d*x/2)**3/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) + 6*A*tan(c/2 + d*x/2)/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) + 3*B*d*x**tan(c/2 + d*x/2)**4/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) + 6*B*d*x**tan(c/2 + d*x/2)**2/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) + 3*B*d*x/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) - 2*B*tan(c/2 + d*x/2)**5/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) - 10*B*tan(c/2 + d*x/2)**3/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d) - 4*B*tan(c/2 + d*x/2)/(2*b*d*tan(c/2 + d*x/2)**4 + 4*b*d*tan(c/2 + d*x/2)**2 + 2*b*d), Eq(a, b)), (2*A*d*x**tan(c/2 + d*x/2)**5/(2*b*d*tan(c/2 + d*x/2)**5 + 4*b*d*tan(c/2 + d*x/2)**3 + 2*b*d*tan(c/2 + d*x/2)) + 4*A*d*x**tan(c/2 + d*x/2)**3/(2*b*d*tan(c/2 + d*x/2)**5 + 4*b*d*tan(c/2 + d*x/2)**3 + 2*b*d*tan(c/2 + d*x/2)) + 2*A*d*x**tan(c/2 + d*x/2)/(2*b*d*tan(c/2 + d*x/2)**5 + 4*b*d*tan(c/2 + d*x/2)**3 + 2*b*d*tan(c/2 + d*x/2)) + 6*...`

3.251.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.251.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.69

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\frac{(2Ba^2-2Aab+Bb^2)(dx+c)}{b^3} + \frac{4(Ba^3-Aa^2b)\left(\pi\left[\frac{dx+c}{2\pi}+\frac{1}{2}\right]\operatorname{sgn}(-2a+2b)+\arctan\left(-\frac{a\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)-b\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{\sqrt{a^2-b^2}}\right)\right)}{\sqrt{a^2-b^2}b^3} - \frac{2(2Ba\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right))}{2d}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/2*((2*B*a^2 - 2*A*a*b + B*b^2)*(d*x + c)/b^3 + 4*(B*a^3 - A*a^2*b)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/(sqrt(a^2 - b^2)*b^3) - 2*(2*B*a*tan(1/2*d*x + 1/2*c)^3 - 2*A*b*tan(1/2*d*x + 1/2*c)^3 + B*b*tan(1/2*d*x + 1/2*c)^3 + 2*B*a*tan(1/2*d*x + 1/2*c) - 2*A*b*tan(1/2*d*x + 1/2*c) - B*b*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 + 1)^2*b^2))/d`

3.251.9 Mupad [B] (verification not implemented)

Time = 4.29 (sec) , antiderivative size = 3761, normalized size of antiderivative = 28.07

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Too large to display}$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output

$$\begin{aligned}
& ((\tan(c/2 + (d*x)/2)*(2*A*b - 2*B*a + B*b))/b^2 - (\tan(c/2 + (d*x)/2)^3*(2 \\
& *B*a - 2*A*b + B*b))/b^2)/(d*(2*\tan(c/2 + (d*x)/2)^2 + \tan(c/2 + (d*x)/2)^ \\
& 4 + 1)) - (\operatorname{atan}((((((8*(2*B*b^{10} + 8*A*a^2*b^8 - 4*A*a^3*b^7 + 2*B*a^2*b^ \\
& 8 - 6*B*a^3*b^7 + 4*B*a^4*b^6 - 4*A*a*b^9 - 2*B*a*b^9))/b^6 - (4*\tan(c/2 + \\
& (d*x)/2)*(B*a^2*2i + B*b^2*1i - A*a*b*2i)*(8*a*b^8 - 16*a^2*b^7 + 8*a^3*b \\
& ^6))/b^7)*(B*a^2*2i + B*b^2*1i - A*a*b*2i))/(2*b^3) - (8*\tan(c/2 + (d*x)/2 \\
&)*(8*B^2*a^7 - B^2*b^7 + 3*B^2*a*b^6 - 16*B^2*a^6*b - 4*A^2*a^2*b^5 + 12*A \\
& ^2*a^3*b^4 - 16*A^2*a^4*b^3 + 8*A^2*a^5*b^2 - 7*B^2*a^2*b^5 + 13*B^2*a^3*b \\
& ^4 - 16*B^2*a^4*b^3 + 16*B^2*a^5*b^2 + 4*A*B*a*b^6 - 16*A*B*a^6*b - 12*A*B \\
& *a^2*b^5 + 20*A*B*a^3*b^4 - 28*A*B*a^4*b^3 + 32*A*B*a^5*b^2))/b^4)*(B*a^2* \\
& 2i + B*b^2*1i - A*a*b*2i)*1i)/(2*b^3) - (((((8*(2*B*b^{10} + 8*A*a^2*b^8 - 4 \\
& *A*a^3*b^7 + 2*B*a^2*b^8 - 6*B*a^3*b^7 + 4*B*a^4*b^6 - 4*A*a*b^9 - 2*B*a*b \\
& ^9))/b^6 + (4*\tan(c/2 + (d*x)/2)*(B*a^2*2i + B*b^2*1i - A*a*b*2i)*(8*a*b^8 \\
& - 16*a^2*b^7 + 8*a^3*b^6))/b^7)*(B*a^2*2i + B*b^2*1i - A*a*b*2i))/(2*b^3) \\
& + (8*\tan(c/2 + (d*x)/2)*(8*B^2*a^7 - B^2*b^7 + 3*B^2*a*b^6 - 16*B^2*a^6*b \\
& - 4*A^2*a^2*b^5 + 12*A^2*a^3*b^4 - 16*A^2*a^4*b^3 + 8*A^2*a^5*b^2 - 7*B^2 \\
& *a^2*b^5 + 13*B^2*a^3*b^4 - 16*B^2*a^4*b^3 + 16*B^2*a^5*b^2 + 4*A*B*a*b^6 \\
& - 16*A*B*a^6*b - 12*A*B*a^2*b^5 + 20*A*B*a^3*b^4 - 28*A*B*a^4*b^3 + 32*A*B \\
& *a^5*b^2))/b^4)*(B*a^2*2i + B*b^2*1i - A*a*b*2i)*1i)/(2*b^3))/((16*(4*B^3* \\
& a^8 - 6*B^3*a^7*b + 4*A^3*a^4*b^4 - 4*A^3*a^5*b^3 - B^3*a^3*b^5 + 2*B^3*...
\end{aligned}$$

3.252 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.252.1 Optimal result 2332
 3.252.2 Mathematica [A] (verified) 2332
 3.252.3 Rubi [A] (verified) 2333
 3.252.4 Maple [A] (verified) 2335
 3.252.5 Fricas [A] (verification not implemented) 2336
 3.252.6 Sympy [B] (verification not implemented) 2336
 3.252.7 Maxima [F(-2)] 2337
 3.252.8 Giac [A] (verification not implemented) 2338
 3.252.9 Mupad [B] (verification not implemented) 2339

3.252.1 Optimal result

Integrand size = 29, antiderivative size = 89

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

$$= \frac{(Ab - aB)x}{b^2} - \frac{2a(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b}b^2\sqrt{a+b}} + \frac{B \sin(c+dx)}{bd}$$

output `(A*b-B*a)*x/b^2+B*sin(d*x+c)/b/d-2*a*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b^2/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.252.2 Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.96

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

$$= \frac{(Ab - aB)(c+dx) - \frac{2a(-Ab+aB) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}}{b^2d} + bB \sin(c+dx)$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `((A*b - a*B)*(c + d*x) - (2*a*(-(A*b) + a*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + b*B*Sin[c + d*x])/(b^2*d)`

3.252. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.252.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.97, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{a+b\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{a+b\cos(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{a+b\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{\int \frac{(Ab-aB)\cos(c+dx)}{a+b\cos(c+dx)} dx}{b} + \frac{B\sin(c+dx)}{bd} \\
 & \quad \downarrow \text{27} \\
 & \frac{(Ab-aB) \int \frac{\cos(c+dx)}{a+b\cos(c+dx)} dx}{b} + \frac{B\sin(c+dx)}{bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab-aB) \int \frac{\sin(c+dx+\frac{\pi}{2})}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B\sin(c+dx)}{bd} \\
 & \quad \downarrow \text{3214} \\
 & \frac{(Ab-aB) \left(\frac{x}{b} - \frac{a \int \frac{1}{a+b\cos(c+dx)} dx}{b} \right)}{b} + \frac{B\sin(c+dx)}{bd} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{(Ab - aB) \left(\frac{x}{b} - \frac{a \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} \right)}{b} + \frac{B \sin(c+dx)}{bd}$$

↓ 3138

$$\frac{(Ab - aB) \left(\frac{x}{b} - \frac{2a \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{bd} \right)}{b} + \frac{B \sin(c+dx)}{bd}$$

↓ 218

$$\frac{(Ab - aB) \left(\frac{x}{b} - \frac{2a \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{bd\sqrt{a-b}\sqrt{a+b}} \right)}{b} + \frac{B \sin(c+dx)}{bd}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `((A*b - a*B)*(x/b - (2*a*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d))/b + (B*Sin[c + d*x])/(b*d)`

3.252.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.252.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.24

method	result
derivativedivides	$\frac{\frac{2Bb \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + 2(Ab - Ba) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^2} - \frac{2a(Ab - Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^2 \sqrt{(a-b)(a+b)}}$
default	$\frac{\frac{2Bb \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + 2(Ab - Ba) \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^2} - \frac{2a(Ab - Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^2 \sqrt{(a-b)(a+b)}}$
risch	$\frac{x A}{b} - \frac{a B x}{b^2} - \frac{i e^{i(dx+c)} B}{2bd} + \frac{i e^{-i(dx+c)} B}{2bd} - \frac{a \ln\left(\frac{e^{i(dx+c)} - i a^2 - i b^2 - a \sqrt{-a^2 + b^2}}{b \sqrt{-a^2 + b^2}}\right) A}{\sqrt{-a^2 + b^2} db} + \frac{a^2 \ln\left(\frac{e^{i(dx+c)} - i a^2 - i b^2 - a \sqrt{-a^2 + b^2}}{b \sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2} d}$

input `int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b), x, method=_RETURNVERBOSE)`

output `1/d*(2/b^2*(B*b*tan(1/2*d*x+1/2*c)/(1+tan(1/2*d*x+1/2*c)^2)+(A*b-B*a)*arctan(tan(1/2*d*x+1/2*c)))-2*a*(A*b-B*a)/b^2/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))`

3.252.
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.252.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 322, normalized size of antiderivative = 3.62

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \left[\frac{2(Ba^3 - Aa^2b - Bab^2 + Ab^3)dx - (Ba^2 - Aab)\sqrt{-a^2 + b^2} \log\left(\frac{2ab\cos(dx+c) + (2a^2 - b^2)\cos(dx+c)^2 - 2\sqrt{-a^2 + b^2}\sin(dx+c)}{b^2\cos(dx+c)^2 + 2ab\cos(dx+c) + a^2}\right)}{2(a^2b^2 - b^4)d} \right. \\ \left. - \frac{(Ba^3 - Aa^2b - Bab^2 + Ab^3)dx - (Ba^2 - Aab)\sqrt{a^2 - b^2} \arctan\left(-\frac{a\cos(dx+c)+b}{\sqrt{a^2 - b^2}\sin(dx+c)}\right) - (Ba^2b - Bb^3)\sin(dx+c)}{(a^2b^2 - b^4)d} \right]$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `[-1/2*(2*(B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*d*x - (B*a^2 - A*a*b)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(B*a^2*b - B*b^3)*sin(d*x + c))/((a^2*b^2 - b^4)*d), -((B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*d*x - (B*a^2 - A*a*b)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (B*a^2*b - B*b^3)*sin(d*x + c))/((a^2*b^2 - b^4)*d)]`

3.252.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3225 vs. 2(76) = 152.

Time = 56.39 (sec) , antiderivative size = 3225, normalized size of antiderivative = 36.24

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Piecewise((zoo*x*(A + B*cos(c)), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (A*d*x*tan(c/2 + d*x/2)**2/(b*d*tan(c/2 + d*x/2)**2 + b*d) + A*d*x/(b*d*tan(c/2 + d*x/2)**2 + b*d) - A*tan(c/2 + d*x/2)**3/(b*d*tan(c/2 + d*x/2)**2 + b*d) - A*tan(c/2 + d*x/2)/(b*d*tan(c/2 + d*x/2)**2 + b*d) - B*d*x*tan(c/2 + d*x/2)**2/(b*d*tan(c/2 + d*x/2)**2 + b*d) - B*d*x/(b*d*tan(c/2 + d*x/2)**2 + b*d) + B*tan(c/2 + d*x/2)**3/(b*d*tan(c/2 + d*x/2)**2 + b*d) + 3*B*tan(c/2 + d*x/2)/(b*d*tan(c/2 + d*x/2)**2 + b*d), Eq(a, b)), (A*d*x*tan(c/2 + d*x/2)**3/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + A*d*x*tan(c/2 + d*x/2)/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + A*tan(c/2 + d*x/2)**2/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + A/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + B*d*x*tan(c/2 + d*x/2)**3/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + B*d*x*tan(c/2 + d*x/2)/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + 3*B*tan(c/2 + d*x/2)**2/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)) + B/(b*d*tan(c/2 + d*x/2)**3 + b*d*tan(c/2 + d*x/2)), Eq(a, -b)), ((A*sin(c + d*x)/d + B*x*sin(c + d*x)**2/2 + B*x*cos(c + d*x)**2/2 + B*sin(c + d*x)*cos(c + d*x)/(2*d))/a, Eq(b, 0)), (x*(A + B*cos(c))*cos(c)/(a + b*cos(c)), Eq(d, 0)), (A*a*b*d*x*sqrt(-a/(a - b) - b/(a - b))*tan(c/2 + d*x/2)**2/(a*b**2*d*sqrt(-a/(a - b) - b/(a - b)))*tan(c/2 + d*x/2)**2 + a*b**2*d*sqrt(-a/(a - b) - b/(a - b)) - b**3*d*sqrt(-a/(a - b) - b/(a - b))*tan(c/2 + d*x/2)**2 - b**3*d*sqrt(-a/(a - b) ...`

3.252.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` for more de`

3.252.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.60

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx =$$

$$\frac{(Ba-Ab)(dx+c)}{b^2} - \frac{2B\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{\left(\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)^2+1\right)b} + \frac{2(Ba^2-Aab)\left(\pi\left\lfloor\frac{dx+c}{2\pi}+\frac{1}{2}\right\rfloor\operatorname{sgn}(-2a+2b)+\arctan\left(-\frac{a\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)-b\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{\sqrt{a^2-b^2}}\right)\right)}{\sqrt{a^2-b^2}b^2}$$

d

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `-(B*a - A*b)*(d*x + c)/b^2 - 2*B*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 + 1)*b) + 2*(B*a^2 - A*a*b)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/(sqrt(a^2 - b^2)*b^2)/d`

3.252.9 Mupad [B] (verification not implemented)

Time = 1.23 (sec) , antiderivative size = 541, normalized size of antiderivative = 6.08

$$\begin{aligned}
& \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx \\
&= \frac{2Ba \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2}+\frac{dx}{2}\right)}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{d(a^2-b^2)} - \frac{2Ab \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2}+\frac{dx}{2}\right)}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{d(a^2-b^2)} - \frac{Bb \sin(c+dx)}{d(a^2-b^2)} \\
&+ \frac{2Aa^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2}+\frac{dx}{2}\right)}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{bd(a^2-b^2)} - \frac{2Ba^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2}+\frac{dx}{2}\right)}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{b^2d(a^2-b^2)} \\
&+ \frac{Aa \ln\left(\frac{a \sin\left(\frac{c}{2}+\frac{dx}{2}\right) - b \sin\left(\frac{c}{2}+\frac{dx}{2}\right) + \cos\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{b^2-a^2}}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{bd\sqrt{b^2-a^2}} \\
&- \frac{Aa \ln\left(\frac{b \sin\left(\frac{c}{2}+\frac{dx}{2}\right) - a \sin\left(\frac{c}{2}+\frac{dx}{2}\right) + \cos\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{b^2-a^2}}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{bd\sqrt{b^2-a^2}} \\
&- \frac{Ba^2 \ln\left(\frac{a \sin\left(\frac{c}{2}+\frac{dx}{2}\right) - b \sin\left(\frac{c}{2}+\frac{dx}{2}\right) + \cos\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{b^2-a^2}}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{b^2d\sqrt{b^2-a^2}} \\
&+ \frac{Ba^2 \ln\left(\frac{b \sin\left(\frac{c}{2}+\frac{dx}{2}\right) - a \sin\left(\frac{c}{2}+\frac{dx}{2}\right) + \cos\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{b^2-a^2}}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{b^2d\sqrt{b^2-a^2}} + \frac{Ba^2 \sin(c+dx)}{bd(a^2-b^2)}
\end{aligned}$$

```
input int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)),x)
```

```
output (2*B*a*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(d*(a^2 - b^2)) - (2*A
*b*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(d*(a^2 - b^2)) - (B*b*sin
(c + d*x))/(d*(a^2 - b^2)) + (2*A*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d
*x)/2)))/(b*d*(a^2 - b^2)) - (2*B*a^3*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d
*x)/2)))/(b^2*d*(a^2 - b^2)) + (A*a*log((a*sin(c/2 + (d*x)/2) - b*sin(c/2
+ (d*x)/2) + cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/cos(c/2 + (d*x)/2)))/(b
*d*(b^2 - a^2)^(1/2)) - (A*a*log((b*sin(c/2 + (d*x)/2) - a*sin(c/2 + (d*x)
/2) + cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/cos(c/2 + (d*x)/2)))/(b*d*(b^2
- a^2)^(1/2)) - (B*a^2*log((a*sin(c/2 + (d*x)/2) - b*sin(c/2 + (d*x)/2) +
cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/cos(c/2 + (d*x)/2)))/(b^2*d*(b^2 -
a^2)^(1/2)) + (B*a^2*log((b*sin(c/2 + (d*x)/2) - a*sin(c/2 + (d*x)/2) + co
s(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/cos(c/2 + (d*x)/2)))/(b^2*d*(b^2 - a^2
)^(1/2)) + (B*a^2*sin(c + d*x))/(b*d*(a^2 - b^2))
```

3.253 $\int \frac{A+B \cos(c+dx)}{a+b \cos(c+dx)} dx$

3.253.1 Optimal result	2340
3.253.2 Mathematica [A] (verified)	2340
3.253.3 Rubi [A] (verified)	2341
3.253.4 Maple [A] (verified)	2342
3.253.5 Fricas [A] (verification not implemented)	2343
3.253.6 Sympy [B] (verification not implemented)	2343
3.253.7 Maxima [F(-2)]	2344
3.253.8 Giac [B] (verification not implemented)	2344
3.253.9 Mupad [B] (verification not implemented)	2345

3.253.1 Optimal result

Integrand size = 23, antiderivative size = 67

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx = \frac{Bx}{b} + \frac{2(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b}b\sqrt{a+bd}}$$

output `B*x/b+2*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.253.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.01

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx = \frac{B(c + dx)}{bd} + \frac{2(-Ab + aB) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x]),x]`

output `(B*(c + d*x) + (2*(-A*b) + a*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2]/(b*d)`

3.253.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3214} \\
 & \frac{(Ab - aB)}{b} \int \frac{1}{a + b \cos(c + dx)} dx + \frac{Bx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab - aB)}{b} \int \frac{1}{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{Bx}{b} \\
 & \quad \downarrow \text{3138} \\
 & \frac{2(Ab - aB)}{bd} \int \frac{1}{(a-b) \tan^2\left(\frac{1}{2}(c + dx)\right) + a + b} d \tan\left(\frac{1}{2}(c + dx)\right) + \frac{Bx}{b} \\
 & \quad \downarrow \text{218} \\
 & \frac{2(Ab - aB)}{bd\sqrt{a-b}\sqrt{a+b}} \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c + dx)\right)}{\sqrt{a+b}}\right) + \frac{Bx}{b}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x]),x]`

output `(B*x)/b + (2*(A*b - a*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d)`

3.253.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

3.253.4 Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{2(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b\sqrt{(a-b)(a+b)}} + \frac{2B \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b}$
default	$\frac{2(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b\sqrt{(a-b)(a+b)}} + \frac{2B \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b}$
risch	$\frac{Bx}{b} - \frac{\ln\left(\frac{e^{i(dx+c)} + ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)A}{\sqrt{-a^2+b^2}d} + \frac{\ln\left(\frac{e^{i(dx+c)} + ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)Ba}{\sqrt{-a^2+b^2}db} + \frac{\ln\left(\frac{e^{i(dx+c)} - ia^2 - ib^2 - a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}d}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b), x, method=_RETURNVERBOSE)`

output `1/d*(2*(A*b-B*a)/b/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))+2*B/b*arctan(tan(1/2*d*x+1/2*c)))`

3.253.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 242, normalized size of antiderivative = 3.61

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \left[\frac{2(Ba^2 - Bb^2)dx + (Ba - Ab)\sqrt{-a^2 + b^2} \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 + 2\sqrt{-a^2 + b^2}(a \cos(dx+c) + b) \sin(dx+c)}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{2(a^2b - b^3)d} \right]$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `[1/2*(2*(B*a^2 - B*b^2)*d*x + (B*a - A*b)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)))/(a^2*b - b^3*d), ((B*a^2 - B*b^2)*d*x - (B*a - A*b)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c)))/(a^2*b - b^3*d)]`

3.253.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 524 vs. 2(56) = 112.

Time = 11.24 (sec) , antiderivative size = 524, normalized size of antiderivative = 7.82

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \left\{ \begin{array}{l} \frac{\tilde{\alpha}x(A+B \cos(c))}{\cos(c)} \\ \frac{A \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{bd} + \frac{Bx}{b} - \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{bd} \\ \frac{A}{bd \tan\left(\frac{c}{2} + \frac{dx}{2}\right)} + \frac{Bx}{b} + \frac{B}{bd \tan\left(\frac{c}{2} + \frac{dx}{2}\right)} \\ \frac{Ax + \frac{B \sin(c+dx)}{d}}{a} \\ \frac{x(A+B \cos(c))}{a+b \cos(c)} \\ \frac{Ab \log\left(-\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{abd\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}} - b^2d\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}}} - \frac{Ab \log\left(\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{abd\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}} - b^2d\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}}} + \frac{Badx\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}}}{abd\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}} - b^2d\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}}} - \frac{Ba \log\left(-\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{abd\sqrt{-\frac{a}{a-b}-\frac{b}{a-b}}} \end{array} \right.$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

```
output Piecewise((zoo*x*(A + B*cos(c))/cos(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (
A*tan(c/2 + d*x/2)/(b*d) + B*x/b - B*tan(c/2 + d*x/2)/(b*d), Eq(a, b)), (A
/(b*d*tan(c/2 + d*x/2)) + B*x/b + B/(b*d*tan(c/2 + d*x/2)), Eq(a, -b)), ((
A*x + B*sin(c + d*x)/d)/a, Eq(b, 0)), (x*(A + B*cos(c))/(a + b*cos(c)), Eq
(d, 0)), (A*b*log(-sqrt(-a/(a - b) - b/(a - b)) + tan(c/2 + d*x/2))/(a*b*d
*sqrt(-a/(a - b) - b/(a - b)) - b**2*d*sqrt(-a/(a - b) - b/(a - b))) - A*b
*log(sqrt(-a/(a - b) - b/(a - b)) + tan(c/2 + d*x/2))/(a*b*d*sqrt(-a/(a -
b) - b/(a - b)) - b**2*d*sqrt(-a/(a - b) - b/(a - b))) + B*a*d*x*sqrt(-a/(
a - b) - b/(a - b))/(a*b*d*sqrt(-a/(a - b) - b/(a - b)) - b**2*d*sqrt(-a/(
a - b) - b/(a - b))) - B*a*log(-sqrt(-a/(a - b) - b/(a - b)) + tan(c/2 + d
*x/2))/(a*b*d*sqrt(-a/(a - b) - b/(a - b)) - b**2*d*sqrt(-a/(a - b) - b/(a
- b))) + B*a*log(sqrt(-a/(a - b) - b/(a - b)) + tan(c/2 + d*x/2))/(a*b*d*
sqrt(-a/(a - b) - b/(a - b)) - b**2*d*sqrt(-a/(a - b) - b/(a - b))) - B*b*
d*x*sqrt(-a/(a - b) - b/(a - b))/(a*b*d*sqrt(-a/(a - b) - b/(a - b)) - b**
2*d*sqrt(-a/(a - b) - b/(a - b))), True))
```

3.253.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

3.253.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 296 vs. $2(58) = 116$.

Time = 0.30 (sec) , antiderivative size = 296, normalized size of antiderivative = 4.42

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx = \frac{\left(\sqrt{a^2 - b^2} B (2a - b) |a - b| - \sqrt{a^2 - b^2} A b |a - b| - \sqrt{a^2 - b^2} A |a - b| |b| + \sqrt{a^2 - b^2} B |a - b| |b| \right) \left(\pi \left[\frac{dx + c}{2\pi} + \frac{1}{2} \right] + \arctan \left(\frac{2\sqrt{1} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\sqrt{2a + \sqrt{-4(a+b)(a-b) + 4a^2}} \frac{a-b}{a-b}} \right) \right)}{(a^2 - 2ab + b^2)b^2 + (a^3 - 2a^2b + ab^2)|b|} d$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `-((sqrt(a^2 - b^2)*B*(2*a - b)*abs(a - b) - sqrt(a^2 - b^2)*A*b*abs(a - b) - sqrt(a^2 - b^2)*A*abs(a - b)*abs(b) + sqrt(a^2 - b^2)*B*abs(a - b)*abs(b))*(pi*floor(1/2*(d*x + c)/pi + 1/2) + arctan(2*sqrt(1/2)*tan(1/2*d*x + 1/2*c)/sqrt((2*a + sqrt(-4*(a + b)*(a - b) + 4*a^2))/(a - b))))/(a^2 - 2*a*b + b^2)*b^2 + (a^3 - 2*a^2*b + a*b^2)*abs(b) + (2*B*a - A*b - B*b + A*abs(b) - B*abs(b))*(pi*floor(1/2*(d*x + c)/pi + 1/2) + arctan(2*sqrt(1/2)*tan(1/2*d*x + 1/2*c)/sqrt((2*a - sqrt(-4*(a + b)*(a - b) + 4*a^2))/(a - b))))/(b^2 - a*abs(b))/d`

3.253.9 Mupad [B] (verification not implemented)

Time = 1.87 (sec) , antiderivative size = 344, normalized size of antiderivative = 5.13

$$\int \frac{A + B \cos(c + dx)}{a + b \cos(c + dx)} dx = \frac{a \left(B \ln \left(\frac{b \sin\left(\frac{c}{2} + \frac{dx}{2}\right) - a \sin\left(\frac{c}{2} + \frac{dx}{2}\right) + \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)} \right) \sqrt{-(a + b)(a - b)} - B \ln \left(\frac{a \sin\left(\frac{c}{2} + \frac{dx}{2}\right) - b \sin\left(\frac{c}{2} + \frac{dx}{2}\right) + \cos\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)} \right)}{2 B \operatorname{atan} \left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)} \right) + b d}$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x)),x)`

output

```
(a*(B*log((b*sin(c/2 + (d*x)/2) - a*sin(c/2 + (d*x)/2) + cos(c/2 + (d*x)/2)
)*(b^2 - a^2)^(1/2))/cos(c/2 + (d*x)/2))*(-(a + b)*(a - b))^(1/2) - B*log(
(a*sin(c/2 + (d*x)/2) - b*sin(c/2 + (d*x)/2) + cos(c/2 + (d*x)/2)*(b^2 - a
^2)^(1/2))/cos(c/2 + (d*x)/2))*(b^2 - a^2)^(1/2)) - A*b*log((b*sin(c/2 + (
d*x)/2) - a*sin(c/2 + (d*x)/2) + cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/cos
(c/2 + (d*x)/2))*(-(a + b)*(a - b))^(1/2) + A*b*log((a*sin(c/2 + (d*x)/2)
- b*sin(c/2 + (d*x)/2) + cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/cos(c/2 + (
d*x)/2))*(b^2 - a^2)^(1/2))/(b*d*(a^2 - b^2)) + (2*B*atan(sin(c/2 + (d*x)/
2)/cos(c/2 + (d*x)/2)))/(b*d)
```

3.254 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx$

3.254.1 Optimal result 2347
 3.254.2 Mathematica [A] (verified) 2347
 3.254.3 Rubi [A] (verified) 2348
 3.254.4 Maple [A] (verified) 2350
 3.254.5 Fricas [A] (verification not implemented) 2350
 3.254.6 Sympy [F] 2351
 3.254.7 Maxima [F(-2)] 2351
 3.254.8 Giac [A] (verification not implemented) 2351
 3.254.9 Mupad [B] (verification not implemented) 2352

3.254.1 Optimal result

Integrand size = 29, antiderivative size = 76

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = -\frac{2(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a\sqrt{a-b}\sqrt{a+bd}} + \frac{A \operatorname{arctanh}(\sin(c + dx))}{ad}$$

output `A*arctanh(sin(d*x+c))/a/d-2*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.254.2 Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.47

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \frac{2(Ab - aB) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + A(-\log(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) + \log(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))))$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x]),x]`

output $((2*(A*b - a*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + A*(-Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]))/(a*d)$

3.254.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.207$, Rules used = {3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow 3042$$

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))} dx$$

$$\downarrow 3480$$

$$\frac{A \int \sec(c+dx) dx}{a} - \frac{(Ab-aB) \int \frac{1}{a+b\cos(c+dx)} dx}{a}$$

$$\downarrow 3042$$

$$\frac{A \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{(Ab-aB) \int \frac{1}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{a}$$

$$\downarrow 3138$$

$$\frac{A \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2(Ab-aB) \int \frac{1}{(a-b)\tan^2(\frac{1}{2}(c+dx))+a+b} d \tan(\frac{1}{2}(c+dx))}{ad}$$

$$\downarrow 218$$

$$\frac{A \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2(Ab-aB) \arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}$$

$$\downarrow 4257$$

$$\frac{A \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2(Ab-aB) \arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x]),x]`

output `(-2*(A*b - a*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]]/(a*Sqrt[a - b]*Sqrt[a + b]*d) + (A*ArcTanh[Sin[c + d*x]])/(a*d)`

3.254.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.254.4 Maple [A] (verified)

Time = 1.38 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a} + \frac{2(-Ab+Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a\sqrt{(a-b)(a+b)}} - \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a}$
default	$\frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a} + \frac{2(-Ab+Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a\sqrt{(a-b)(a+b)}} - \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a}$
risch	$-\frac{\ln\left(e^{i(dx+c)} + \frac{-ia^2+ib^2+a\sqrt{-a^2+b^2}}{\sqrt{-a^2+b^2}b}\right)Ab}{\sqrt{-a^2+b^2}da} + \frac{\ln\left(e^{i(dx+c)} + \frac{-ia^2+ib^2+a\sqrt{-a^2+b^2}}{\sqrt{-a^2+b^2}b}\right)B}{\sqrt{-a^2+b^2}d} + \frac{\ln\left(e^{i(dx+c)} + \frac{ia^2-ib^2+a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}da}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `1/d*(A/a*ln(tan(1/2*d*x+1/2*c)+1)+2/a*(-A*b+B*a)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-A/a*ln(tan(1/2*d*x+1/2*c)-1))`

3.254.5 Fracas [A] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 304, normalized size of antiderivative = 4.00

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \left[\frac{(Ba - Ab)\sqrt{-a^2 + b^2} \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 - 2\sqrt{-a^2+b^2}(a \cos(dx+c) + b) \sin(dx+c) - a^2 + 2b^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{2(a^3 - ab^2)d} + (Aa^2 - \dots) \right]$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `[1/2*((B*a - A*b)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + (A*a^2 - A*b^2)*log(sin(d*x + c) + 1) - (A*a^2 - A*b^2)*log(-sin(d*x + c) + 1))/((a^3 - a*b^2)*d), 1/2*(2*(B*a - A*b)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (A*a^2 - A*b^2)*log(sin(d*x + c) + 1) - (A*a^2 - A*b^2)*log(-sin(d*x + c) + 1))/((a^3 - a*b^2)*d)]`

3.254. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx$

3.254.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a + b*cos(c + d*x)), x)`

3.254.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` for more de`

3.254.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.67

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a} - \frac{A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a} + \frac{2 \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{d \sqrt{a^2 - b^2 a}} (Ba -$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x, algorithm="giac")`

3.254. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx$

output $(A \log(\abs{\tan(1/2*d*x + 1/2*c) + 1})/a - A \log(\abs{\tan(1/2*d*x + 1/2*c) - 1})/a + 2*(\pi*\text{floor}(1/2*(d*x + c)/\pi + 1/2)*\text{sgn}(2*a - 2*b) + \arctan((a*\tan(1/2*d*x + 1/2*c) - b*\tan(1/2*d*x + 1/2*c))/\sqrt{a^2 - b^2}))* (B*a - A*b) / (\sqrt{a^2 - b^2}*a))/d$

3.254.9 Mupad [B] (verification not implemented)

Time = 1.82 (sec) , antiderivative size = 342, normalized size of antiderivative = 4.50

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \frac{2 A \operatorname{atanh}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{a d} + \frac{b \left(A \ln\left(\frac{a \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + b \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) \sqrt{-(a + b)(a - b)} - A \ln\left(\frac{a \sin\left(\frac{c}{2} + \frac{dx}{2}\right) - b \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{\right)}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))),x)`

output $(2*A*\operatorname{atanh}(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/(a*d) + (b*(A*\log((a*\cos(c/2 + (d*x)/2) + b*\cos(c/2 + (d*x)/2) + \sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/\cos(c/2 + (d*x)/2))*(-(a + b)*(a - b))^(1/2) - A*\log((a*\sin(c/2 + (d*x)/2) - b*\sin(c/2 + (d*x)/2) + \cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/\cos(c/2 + (d*x)/2))* (b^2 - a^2)^(1/2)) - B*a*\log((a*\cos(c/2 + (d*x)/2) + b*\cos(c/2 + (d*x)/2) + \sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/\cos(c/2 + (d*x)/2))*(-(a + b)*(a - b))^(1/2) + B*a*\log((a*\sin(c/2 + (d*x)/2) - b*\sin(c/2 + (d*x)/2) + \cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/\cos(c/2 + (d*x)/2))* (b^2 - a^2)^(1/2))/(a*d*(a^2 - b^2))$

3.255 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{a+b \cos(c+dx)} dx$

3.255.1 Optimal result	2353
3.255.2 Mathematica [A] (verified)	2353
3.255.3 Rubi [A] (verified)	2354
3.255.4 Maple [A] (verified)	2357
3.255.5 Fricas [B] (verification not implemented)	2357
3.255.6 Sympy [F]	2358
3.255.7 Maxima [F(-2)]	2358
3.255.8 Giac [A] (verification not implemented)	2359
3.255.9 Mupad [B] (verification not implemented)	2359

3.255.1 Optimal result

Integrand size = 31, antiderivative size = 99

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \frac{2b(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^2 \sqrt{a-b} \sqrt{a+b} d} - \frac{(Ab - aB) \operatorname{arctanh}(\sin(c + dx))}{a^2 d} + \frac{A \tan(c + dx)}{ad}$$

```
output -(A*b-B*a)*arctanh(sin(d*x+c))/a^2/d+2*b*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^2/d/(a-b)^(1/2)/(a+b)^(1/2)+A*tan(d*x+c)/a/d
```

3.255.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.30

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \frac{2b(Ab - aB) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + (Ab - aB) \left(\log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) - \log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) - \frac{1}{a^2 d}$$

```
input Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x]),x]
```


output $((-2*b*(A*b - a*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + (A*b - a*B)*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + a*A*Tan[c + d*x]/(a^2*d)$

3.255.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.97, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3479, 25, 27, 3042, 3226, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^2(a+b\sin(c+dx+\frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{\int -\frac{(Ab-aB)\sec(c+dx)}{a+b\cos(c+dx)} dx}{a} + \frac{A\tan(c+dx)}{ad} \\
 & \quad \downarrow \text{25} \\
 & \frac{A\tan(c+dx)}{ad} - \frac{\int \frac{(Ab-aB)\sec(c+dx)}{a+b\cos(c+dx)} dx}{a} \\
 & \quad \downarrow \text{27} \\
 & \frac{A\tan(c+dx)}{ad} - \frac{(Ab-aB)\int \frac{\sec(c+dx)}{a+b\cos(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A\tan(c+dx)}{ad} - \frac{(Ab-aB)\int \frac{1}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{a} \\
 & \quad \downarrow \text{3226} \\
 & \frac{A\tan(c+dx)}{ad} - \frac{(Ab-aB)\left(\frac{\int \sec(c+dx) dx}{a} - \frac{b\int \frac{1}{a+b\cos(c+dx)} dx}{a}\right)}{a} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.255. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{a+b\cos(c+dx)} dx$

$$\frac{A \tan(c+dx)}{ad} - \frac{(Ab - aB) \left(\frac{\int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{a} \right)}{a}$$

↓ 3138

$$\frac{A \tan(c+dx)}{ad} - \frac{(Ab - aB) \left(\frac{\int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{ad} \right)}{a}$$

↓ 218

$$\frac{A \tan(c+dx)}{ad} - \frac{(Ab - aB) \left(\frac{\int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right)}{a}$$

↓ 4257

$$\frac{A \tan(c+dx)}{ad} - \frac{(Ab - aB) \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2b \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right)}{a}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x]),x]
```

```
output -(((A*b - a*B)*((-2*b*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/
(a*Sqrt[a - b]*Sqrt[a + b]*d) + ArcTanh[Sin[c + d*x]]/(a*d)))/a) + (A*Tan[
c + d*x])/(a*d)
```

3.255.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

3.255. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{a+b \cos(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3226 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.255.4 Maple [A] (verified)

Time = 1.64 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.45

method	result
derivativedivides	$\frac{2b(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^2 \sqrt{(a-b)(a+b)}} - \frac{A}{a\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(Ab-Ba) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^2} - \frac{A}{a\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(-Ab+Ba)}{d}$
default	$\frac{2b(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^2 \sqrt{(a-b)(a+b)}} - \frac{A}{a\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(Ab-Ba) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^2} - \frac{A}{a\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(-Ab+Ba)}{d}$
risch	$\frac{2iA}{da(e^{2i(dx+c)}+1)} - \frac{b^2 \ln\left(e^{i(dx+c)} + \frac{ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)A}{\sqrt{-a^2+b^2} da^2} + \frac{b \ln\left(e^{i(dx+c)} + \frac{ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)B}{\sqrt{-a^2+b^2} da} + \frac{b^2 \ln\left(e^{i(dx+c)} + \frac{ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} da}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)
```

```
output 1/d*(2*b*(A*b-B*a)/a^2/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-A/a/(tan(1/2*d*x+1/2*c)-1)+(A*b-B*a)/a^2*ln(tan(1/2*d*x+1/2*c)-1)-A/a/(tan(1/2*d*x+1/2*c)+1)+1/a^2*(-A*b+B*a)*ln(tan(1/2*d*x+1/2*c)+1))
```

3.255.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(89) = 178.

Time = 0.39 (sec) , antiderivative size = 460, normalized size of antiderivative = 4.65

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{\left[(Bab - Ab^2) \sqrt{-a^2 + b^2} \cos(dx + c) \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 + 2\sqrt{-a^2+b^2}(a \cos(dx+c)+b) \sin(dx+c) - a^2 + b^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right) \right.}{2(Bab - Ab^2) \sqrt{a^2 - b^2} \arctan\left(-\frac{a \cos(dx+c)+b}{\sqrt{a^2-b^2} \sin(dx+c)}\right) \cos(dx+c) - (Ba^3 - Aa^2b - Bab^2 + Ab^3) \cos(dx+c) - \dots}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c)),x, algorithm="fracas")
```

```
output [1/2*((B*a*b - A*b^2)*sqrt(-a^2 + b^2)*cos(d*x + c)*log((2*a*b*cos(d*x + c)
) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)
*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^
2)) + (B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*cos(d*x + c)*log(sin(d*x + c) +
1) - (B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*cos(d*x + c)*log(-sin(d*x + c) +
1) + 2*(A*a^3 - A*a*b^2)*sin(d*x + c))/((a^4 - a^2*b^2)*d*cos(d*x + c)), -
1/2*(2*(B*a*b - A*b^2)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(
a^2 - b^2)*sin(d*x + c)))*cos(d*x + c) - (B*a^3 - A*a^2*b - B*a*b^2 + A*b^
3)*cos(d*x + c)*log(sin(d*x + c) + 1) + (B*a^3 - A*a^2*b - B*a*b^2 + A*b^3
)*cos(d*x + c)*log(-sin(d*x + c) + 1) - 2*(A*a^3 - A*a*b^2)*sin(d*x + c))/
((a^4 - a^2*b^2)*d*cos(d*x + c))]
```

3.255.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c)),x)
```

```
output Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a + b*cos(c + d*x)), x)
```

3.255.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c)),x, algorithm="max
ima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

3.255.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.77

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{(Ba - Ab) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right)}{a^2} - \frac{(Ba - Ab) \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right)}{a^2} - \frac{2A \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)^2 - 1} a + \frac{2(Bab - Ab^2) \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(-2a)\right)}{d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `((B*a - A*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^2 - (B*a - A*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^2 - 2*A*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 - 1)*a) + 2*(B*a*b - A*b^2)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/(sqrt(a^2 - b^2)*a^2))/d`

3.255.9 Mupad [B] (verification not implemented)

Time = 2.30 (sec) , antiderivative size = 675, normalized size of antiderivative = 6.82

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{A b \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right) \operatorname{li}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) 2i}{d (a^2 - b^2)} - \frac{B a \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right) \operatorname{li}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) 2i}{d (a^2 - b^2)} + \frac{A a \tan(c + dx)}{d (a^2 - b^2)}$$

$$- \frac{A b^3 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right) \operatorname{li}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) 2i}{a^2 d (a^2 - b^2)} + \frac{B b^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right) \operatorname{li}}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)}\right) 2i}{a d (a^2 - b^2)} - \frac{A b^2 \tan(c + dx)}{a d (a^2 - b^2)}$$

$$- \frac{B b \operatorname{atan}\left(\frac{\left(a^5 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} + 2 b^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) (b^2 - a^2)^{3/2} - 2 b^5 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} + 3 a^2 b^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} - a^3 b^2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right) (a b^2 - a^3)^2}}{a d (a^2 - b^2)}$$

$$+ \frac{A b^2 \operatorname{atan}\left(\frac{\left(a^5 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} + 2 b^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) (b^2 - a^2)^{3/2} - 2 b^5 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} + 3 a^2 b^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} - a^3 b^2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right) (a b^2 - a^3)^2}}{a^2 d (a^2 - b^2)}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))),x)`

output `(A*b*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/(d*(a^2 - b^2)) - (B*a*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/(d*(a^2 - b^2)) + (A*a*tan(c + d*x))/(d*(a^2 - b^2)) - (A*b^3*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/(a^2*d*(a^2 - b^2)) + (B*b^2*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*2i)/(a*d*(a^2 - b^2)) - (A*b^2*tan(c + d*x))/(a*d*(a^2 - b^2)) - (B*b*atan(((a^5*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 2*b^3*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(3/2) - 2*b^5*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 3*a^2*b^3*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - a^3*b^2*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - a^4*b*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))*1i)/(cos(c/2 + (d*x)/2)*(a*b^2 - a^3)^2))*(-(a + b)*(a - b))^(1/2)*2i)/(a*d*(a^2 - b^2)) + (A*b^2*atan(((a^5*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 2*b^3*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(3/2) - 2*b^5*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 3*a^2*b^3*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - a^3*b^2*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - a^4*b*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))*1i)/(cos(c/2 + (d*x)/2)*(a*b^2 - a^3)^2))*(-(a + b)*(a - b))^(1/2)*2i)/(a^2*d*(a^2 - b^2))`

3.256 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$

3.256.1 Optimal result 2361
 3.256.2 Mathematica [B] (verified) 2361
 3.256.3 Rubi [A] (verified) 2362
 3.256.4 Maple [A] (verified) 2366
 3.256.5 Fricas [B] (verification not implemented) 2366
 3.256.6 Sympy [F] 2367
 3.256.7 Maxima [F(-2)] 2367
 3.256.8 Giac [B] (verification not implemented) 2368
 3.256.9 Mupad [B] (verification not implemented) 2368

3.256.1 Optimal result

Integrand size = 31, antiderivative size = 143

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = -\frac{2b^2(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^3\sqrt{a-b}\sqrt{a+bd}} + \frac{(a^2A + 2Ab^2 - 2abB) \operatorname{arctanh}(\sin(c + dx))}{2a^3d} - \frac{(Ab - aB) \tan(c + dx)}{a^2d} + \frac{A \sec(c + dx) \tan(c + dx)}{2ad}$$

output `1/2*(A*a^2+2*A*b^2-2*B*a*b)*arctanh(sin(d*x+c))/a^3/d-2*b^2*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^3/d/(a-b)^(1/2)/(a+b)^(1/2)-(A*b-B*a)*tan(d*x+c)/a^2/d+1/2*A*sec(d*x+c)*tan(d*x+c)/a/d`

3.256.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 300 vs. 2(143) = 286.

Time = 1.67 (sec) , antiderivative size = 300, normalized size of antiderivative = 2.10

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \frac{8b^2(Ab - aB) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} - 2(a^2A + 2Ab^2 - 2abB) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2C$$

3.256. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x]),x]`

output `((8*b^2*(A*b - a*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] - 2*(a^2*A + 2*A*b^2 - 2*a*b*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 2*(a^2*A + 2*A*b^2 - 2*a*b*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (a^2*A)/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2 + (4*a*(-(A*b) + a*B)*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]) - (a^2*A)/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 + (4*a*(-(A*b) + a*B)*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(4*a^3*d)`

3.256.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3479, 25, 3042, 3534, 25, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))} dx$$

↓ 3479

$$\frac{\int -\frac{(-Ab\cos^2(c+dx)-aA\cos(c+dx)+2(Ab-aB))\sec^2(c+dx)}{a+b\cos(c+dx)} dx}{2a} + \frac{A\tan(c+dx)\sec(c+dx)}{2ad}$$

↓ 25

$$\frac{A\tan(c+dx)\sec(c+dx)}{2ad} - \frac{\int \frac{(-Ab\cos^2(c+dx)-aA\cos(c+dx)+2(Ab-aB))\sec^2(c+dx)}{a+b\cos(c+dx)} dx}{2a}$$

↓ 3042

$$\frac{A\tan(c+dx)\sec(c+dx)}{2ad} - \frac{\int \frac{-Ab\sin(c+dx+\frac{\pi}{2})^2-aA\sin(c+dx+\frac{\pi}{2})+2(Ab-aB)}{\sin(c+dx+\frac{\pi}{2})^2(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{2a}$$

↓ 3534

3.256. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{a+b\cos(c+dx)} dx$

$$\begin{aligned}
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int -\frac{(Aa^2 - 2bBa + Ab \cos(c+dx)a + 2Ab^2) \sec(c+dx)}{a+b \cos(c+dx)} dx}{2a} + \frac{2(Ab-aB) \tan(c+dx)}{ad} \\
 & \quad \downarrow \text{25} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(Ab-aB) \tan(c+dx)}{ad} - \frac{\int \frac{(Aa^2 - 2bBa + Ab \cos(c+dx)a + 2Ab^2) \sec(c+dx)}{a+b \cos(c+dx)} dx}{2a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(Ab-aB) \tan(c+dx)}{ad} - \frac{\int \frac{Aa^2 - 2bBa + Ab \sin(c+dx + \frac{\pi}{2})a + 2Ab^2}{\sin(c+dx + \frac{\pi}{2})(a+b \sin(c+dx + \frac{\pi}{2}))} dx}{2a} \\
 & \quad \downarrow \text{3480} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(Ab-aB) \tan(c+dx)}{ad} - \frac{(a^2A - 2abB + 2Ab^2) \int \sec(c+dx) dx}{2a} - \frac{2b^2(Ab-aB) \int \frac{1}{a+b \cos(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{(a^2A - 2abB + 2Ab^2) \int \csc(c+dx + \frac{\pi}{2}) dx}{2a} - \frac{2b^2(Ab-aB) \int \frac{1}{a+b \sin(c+dx + \frac{\pi}{2})} dx}{a} \\
 & \quad \downarrow \text{3138} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(Ab-aB) \tan(c+dx)}{ad} - \frac{(a^2A - 2abB + 2Ab^2) \int \csc(c+dx + \frac{\pi}{2}) dx}{2a} - \frac{4b^2(Ab-aB) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{a} \\
 & \quad \downarrow \text{218} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(Ab-aB) \tan(c+dx)}{ad} - \frac{(a^2A - 2abB + 2Ab^2) \int \csc(c+dx + \frac{\pi}{2}) dx}{2a} - \frac{4b^2(Ab-aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

3.256. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(Ab-aB) \tan(c+dx)}{ad} - \frac{(a^2A-2abB+2Ab^2) \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{4b^2(Ab-aB) \operatorname{arctan}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}$$

$$2a$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x]),x]`

output `(A*Sec[c + d*x]*Tan[c + d*x])/(2*a*d) - (-(((-4*b^2*(A*b - a*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + ((a^2*A + 2*A*b^2 - 2*a*b*B)*ArcTanh[Sin[c + d*x]])/(a*d))/a) + (2*(A*b - a*B)*Tan[c + d*x])/(a*d))/(2*a)`

3.256.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.256.4 Maple [A] (verified)

Time = 1.56 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.60

method	result
derivativedivides	$-\frac{2b^2(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^3 \sqrt{(a-b)(a+b)}} + \frac{A}{2a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{-aA - 2Ab + 2Ba}{2a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(-Aa^2 - 2Ab^2 + 2Bab) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2a^3 d}$
default	$-\frac{2b^2(Ab-Ba) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^3 \sqrt{(a-b)(a+b)}} + \frac{A}{2a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{-aA - 2Ab + 2Ba}{2a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(-Aa^2 - 2Ab^2 + 2Bab) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2a^3 d}$
risch	$-\frac{i(Aa e^{3i(dx+c)} + 2Ab e^{2i(dx+c)} - 2Ba e^{2i(dx+c)} - aA e^{i(dx+c)} + 2Ab - 2Ba)}{a^2 d (e^{2i(dx+c)} + 1)^2} + \frac{A \ln(e^{i(dx+c)} + i)}{2ad} + \frac{\ln(e^{i(dx+c)} + i) A}{a^3 d}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `1/d*(-2*b^2*(A*b-B*a)/a^3/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))+1/2*A/a/(tan(1/2*d*x+1/2*c)-1)^2-1/2*(-A*a-2*A*b+2*B*a)/a^2/(tan(1/2*d*x+1/2*c)-1)+1/2/a^3*(-A*a^2-2*A*b^2+2*B*a*b)*ln(tan(1/2*d*x+1/2*c)-1)-1/2*A/a/(tan(1/2*d*x+1/2*c)+1)^2-1/2*(-A*a-2*A*b+2*B*a)/a^2/(tan(1/2*d*x+1/2*c)+1)+1/2*(A*a^2+2*A*b^2-2*B*a*b)/a^3*ln(tan(1/2*d*x+1/2*c)+1))`

3.256.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 260 vs. 2(129) = 258.

Time = 2.78 (sec) , antiderivative size = 589, normalized size of antiderivative = 4.12

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \left[\frac{2(Bab^2 - Ab^3) \sqrt{-a^2 + b^2} \cos(dx + c)^2 \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 - 2\sqrt{-a^2 + b^2}(a \cos(dx+c) + b) \sin(dx+c) - b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{\dots} \right]$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `[1/4*(2*(B*a*b^2 - A*b^3)*sqrt(-a^2 + b^2)*cos(d*x + c)^2*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + (A*a^4 - 2*B*a^3*b + A*a^2*b^2 + 2*B*a*b^3 - 2*A*b^4)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A*a^4 - 2*B*a^3*b + A*a^2*b^2 + 2*B*a*b^3 - 2*A*b^4)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(A*a^4 - A*a^2*b^2 + 2*(B*a^4 - A*a^3*b - B*a^2*b^2 + A*a*b^3)*cos(d*x + c))*sin(d*x + c))/((a^5 - a^3*b^2)*d*cos(d*x + c)^2), 1/4*(4*(B*a*b^2 - A*b^3)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c)))*cos(d*x + c)^2 + (A*a^4 - 2*B*a^3*b + A*a^2*b^2 + 2*B*a*b^3 - 2*A*b^4)*cos(d*x + c)^2*log(sin(d*x + c) + 1) - (A*a^4 - 2*B*a^3*b + A*a^2*b^2 + 2*B*a*b^3 - 2*A*b^4)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) + 2*(A*a^4 - A*a^2*b^2 + 2*(B*a^4 - A*a^3*b - B*a^2*b^2 + A*a*b^3)*cos(d*x + c))*sin(d*x + c))/((a^5 - a^3*b^2)*d*cos(d*x + c)^2)]`

3.256.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c)),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a + b*cos(c + d*x)), x)`

3.256.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.256.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 269 vs. $2(129) = 258$.

Time = 0.36 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.88

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx$$

$$\frac{(Aa^2 - 2Bab + 2Ab^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^3} - \frac{(Aa^2 - 2Bab + 2Ab^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^3} - \frac{4(Bab^2 - Ab^3) \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(-2a+2b) \right)}{a^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/2*((A*a^2 - 2*B*a*b + 2*A*b^2)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^3 - (A*a^2 - 2*B*a*b + 2*A*b^2)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^3 - 4*(B*a*b^2 - A*b^3)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/sqrt(a^2 - b^2)*a^3 + 2*(A*a*tan(1/2*d*x + 1/2*c)^3 - 2*B*a*tan(1/2*d*x + 1/2*c)^3 + 2*A*b*tan(1/2*d*x + 1/2*c)^3 + A*a*tan(1/2*d*x + 1/2*c) + 2*B*a*tan(1/2*d*x + 1/2*c) - 2*A*b*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 - 1)^2*a^2))/d`

3.256.9 Mupad [B] (verification not implemented)

Time = 4.44 (sec) , antiderivative size = 4051, normalized size of antiderivative = 28.33

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))),x)`

output

$$\begin{aligned}
& (B*a*\sin(2*c + 2*d*x))/(2*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) - (A*b \\
& *sin(2*c + 2*d*x))/(2*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) + (A*a*\sin \\
& (c + d*x))/(2*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) - (A*a*atan((sin(c \\
& /2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*1i)/(2*d*(a^2 - b^2)*(cos(2*c + 2*d* \\
& x)/2 + 1/2)) + (B*b*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*1i)/(\\
& d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) - (A*b^2*atan((sin(c/2 + (d*x)/2 \\
&)*1i)/cos(c/2 + (d*x)/2))*1i)/(2*a*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2 \\
&)) + (A*b^4*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*1i)/(a^3*d*(a \\
& ^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) - (B*b^3*atan((sin(c/2 + (d*x)/2)*1i \\
&)/cos(c/2 + (d*x)/2))*1i)/(a^2*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) + \\
& (A*b^3*sin(2*c + 2*d*x))/(2*a^2*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) \\
& - (B*b^2*sin(2*c + 2*d*x))/(2*a*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) \\
& - (A*a*atan((sin(c/2 + (d*x)/2)*1i)/cos(c/2 + (d*x)/2))*cos(2*c + 2*d*x)* \\
& 1i)/(2*d*(a^2 - b^2)*(cos(2*c + 2*d*x)/2 + 1/2)) + (B*b*atan((sin(c/2 + (d \\
& *x)/2)*1i)/cos(c/2 + (d*x)/2))*cos(2*c + 2*d*x)*1i)/(d*(a^2 - b^2)*(cos(2* \\
& c + 2*d*x)/2 + 1/2)) - (A*b^2*sin(c + d*x))/(2*a*d*(a^2 - b^2)*(cos(2*c + \\
& 2*d*x)/2 + 1/2)) + (A*b^3*atan(((A^2*a^9*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1 \\
& /2) + 8*A^2*b^7*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(3/2) - 8*A^2*b^9*sin(c/2 + \\
& (d*x)/2)*(b^2 - a^2)^(1/2) - A^2*a^8*b*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/ \\
& 2) + 8*A^2*a^2*b^7*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 3*A^2*a^4*b^5...
\end{aligned}$$

3.257 $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+b \cos(c+dx)} dx$

3.257.1 Optimal result 2370
 3.257.2 Mathematica [B] (verified) 2371
 3.257.3 Rubi [A] (verified) 2371
 3.257.4 Maple [A] (verified) 2376
 3.257.5 Fricas [A] (verification not implemented) 2376
 3.257.6 Sympy [F] 2377
 3.257.7 Maxima [F(-2)] 2378
 3.257.8 Giac [B] (verification not implemented) 2378
 3.257.9 Mupad [B] (verification not implemented) 2379

3.257.1 Optimal result

Integrand size = 31, antiderivative size = 187

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \frac{2b^3(Ab - aB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^4 \sqrt{a-b} \sqrt{a+b} d} - \frac{(a^2 + 2b^2)(Ab - aB) \operatorname{arctanh}(\sin(c + dx))}{2a^4 d} + \frac{(2a^2 A + 3Ab^2 - 3abB) \tan(c + dx)}{3a^3 d} - \frac{(Ab - aB) \sec(c + dx) \tan(c + dx)}{2a^2 d} + \frac{A \sec^2(c + dx) \tan(c + dx)}{3ad}$$

output

```
-1/2*(a^2+2*b^2)*(A*b-B*a)*arctanh(sin(d*x+c))/a^4/d+2*b^3*(A*b-B*a)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^4/d/(a-b)^(1/2)/(a+b)^(1/2)+1/3*(2*A*a^2+3*A*b^2-3*B*a*b)*tan(d*x+c)/a^3/d-1/2*(A*b-B*a)*sec(d*x+c)*tan(d*x+c)/a^2/d+1/3*A*sec(d*x+c)^2*tan(d*x+c)/a/d
```

3.257.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 422 vs. $2(187) = 374$.

Time = 1.99 (sec) , antiderivative size = 422, normalized size of antiderivative = 2.26

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{24b^3(-Ab + aB) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} - 6(a^2 + 2b^2)(-Ab + aB) \log\left(\cos\left(\frac{1}{2}(c + dx)\right) - \sin\left(\frac{1}{2}(c + dx)\right)\right) + \dots$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + b*Cos[c + d*x]),x]`

output

```
((24*b^3*(-(A*b) + a*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]]/Sqrt[-a^2 + b^2] - 6*(a^2 + 2*b^2)*(-(A*b) + a*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 6*(a^2 + 2*b^2)*(-(A*b) + a*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (a^2*(-3*A*b + a*(A + 3*B)))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2 + (2*a^3*A*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^3 + (4*a*(2*a^2*A + 3*A*b^2 - 3*a*b*B)*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]) + (2*a^3*A*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^3 - (a^2*(-3*A*b + a*(A + 3*B)))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 + (4*a*(2*a^2*A + 3*A*b^2 - 3*a*b*B)*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(12*a^4*d)
```

3.257.3 Rubi [A] (verified)

Time = 1.40 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.09, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3479, 25, 3042, 3534, 25, 3042, 3534, 27, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^4(c + dx)(A + B \cos(c + dx))}{a + b \cos(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^4 (a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx$$

$$\begin{aligned}
& \int -\frac{(-2Ab \cos^2(c+dx) - 2aA \cos(c+dx) + 3(Ab-aB)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx + \frac{A \tan(c+dx) \sec^2(c+dx)}{3ad} \\
& \quad \downarrow \text{3479} \\
& \frac{A \tan(c+dx) \sec^2(c+dx)}{3ad} - \frac{\int \frac{(-2Ab \cos^2(c+dx) - 2aA \cos(c+dx) + 3(Ab-aB)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx}{3a} \\
& \quad \downarrow \text{25} \\
& \frac{A \tan(c+dx) \sec^2(c+dx)}{3ad} - \frac{\int \frac{-2Ab \sin(c+dx+\frac{\pi}{2})^2 - 2aA \sin(c+dx+\frac{\pi}{2}) + 3(Ab-aB)}{\sin(c+dx+\frac{\pi}{2})^3 (a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3a} \\
& \quad \downarrow \text{3042} \\
& \frac{A \tan(c+dx) \sec^2(c+dx)}{3ad} - \frac{\int \frac{-3b(Ab-aB) \cos^2(c+dx) + a(Ab+3aB) \cos(c+dx) + 2(2Aa^2 - 3bBa + 3Ab^2)}{a+b \cos(c+dx)} \sec^2(c+dx) dx}{2a} + \frac{3(Ab-aB) \tan(c+dx) \sec(c+dx)}{2ad} \\
& \quad \downarrow \text{3534} \\
& \frac{3(Ab-aB) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{(-3b(Ab-aB) \cos^2(c+dx) + a(Ab+3aB) \cos(c+dx) + 2(2Aa^2 - 3bBa + 3Ab^2)) \sec^2(c+dx)}{a+b \cos(c+dx)} dx}{2a} \\
& \quad \downarrow \text{25} \\
& \frac{3(Ab-aB) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{-3b(Ab-aB) \sin(c+dx+\frac{\pi}{2})^2 + a(Ab+3aB) \sin(c+dx+\frac{\pi}{2}) + 2(2Aa^2 - 3bBa + 3Ab^2)}{\sin(c+dx+\frac{\pi}{2})^2 (a+b \sin(c+dx+\frac{\pi}{2}))} dx}{2a} \\
& \quad \downarrow \text{3042} \\
& \frac{3(Ab-aB) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{3((a^2+2b^2)(Ab-aB) + ab \cos(c+dx)(Ab-aB)) \sec(c+dx)}{a+b \cos(c+dx)} dx}{a} + \frac{2(2a^2A - 3abB + 3Ab^2) \tan(c+dx)}{ad} \\
& \quad \downarrow \text{3534} \\
& \frac{3(Ab-aB) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{3((a^2+2b^2)(Ab-aB) + ab \cos(c+dx)(Ab-aB)) \sec(c+dx)}{a+b \cos(c+dx)} dx}{a} + \frac{2(2a^2A - 3abB + 3Ab^2) \tan(c+dx)}{ad} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.257. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+b \cos(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 4257 \\
 & \frac{A \tan(c+dx) \sec^2(c+dx)}{3ad} - \frac{3(Ab-aB) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(2a^2A-3abB+3Ab^2) \tan(c+dx)}{ad} \\
 & \left(\frac{(a^2+2b^2)(Ab-aB) \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{4b^3(Ab-aB) \operatorname{arctan}\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right) \\
 & \frac{}{3a} \frac{}{2a} \frac{}{a}
 \end{aligned}$$

```
input Int[((A + B*cos[c + d*x])*Sec[c + d*x]^4)/(a + b*cos[c + d*x]),x]
```

```
output (A*Sec[c + d*x]^2*Tan[c + d*x])/(3*a*d) - ((3*(A*b - a*B)*Sec[c + d*x]*Tan[c + d*x])/(2*a*d) - ((-3*((-4*b^3*(A*b - a*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + ((a^2 + 2*b^2)*(A*b - a*B)*ArcTanh[Sin[c + d*x]])/(a*d)))/a + (2*(2*a^2*A + 3*A*b^2 - 3*a*b*B)*Tan[c + d*x])/(a*d))/(2*a))/(3*a)
```

3.257.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3138 Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

3.257. $\int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+b \cos(c+dx)} dx$

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.257.4 Maple [A] (verified)

Time = 1.86 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.79

method	result
derivativedivides	$-\frac{A}{3a\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}-\frac{-aA-Ab+Ba}{2a^2\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^2}+\frac{\left(-Aa^2b-2A b^3+Ba^3+2Ba b^2\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{2a^4}-\frac{2Aa^2+Aab+2Ab^2-Ba^3}{2a^3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}$
default	$-\frac{A}{3a\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}-\frac{-aA-Ab+Ba}{2a^2\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^2}+\frac{\left(-Aa^2b-2A b^3+Ba^3+2Ba b^2\right)\ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{2a^4}-\frac{2Aa^2+Aab+2Ab^2-Ba^3}{2a^3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}$
risch	$\frac{i\left(3Aab e^{5i(dx+c)}-3B a^2 e^{5i(dx+c)}+6A b^2 e^{4i(dx+c)}-6Bab e^{4i(dx+c)}+12A a^2 e^{2i(dx+c)}+12A b^2 e^{2i(dx+c)}-12Bab e^{2i(dx+c)}\right)}{3d a^3\left(e^{2i(dx+c)}+1\right)^3}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)
```

```
output 1/d*(-1/3*A/a/(tan(1/2*d*x+1/2*c)+1)^3-1/2*(-A*a-A*b+B*a)/a^2/(tan(1/2*d*x+1/2*c)+1)^2+1/2/a^4*(-A*a^2*b-2*A*b^3+B*a^3+2*B*a*b^2)*ln(tan(1/2*d*x+1/2*c)+1)-1/2*(2*A*a^2+A*a*b+2*A*b^2-B*a^2-2*B*a*b)/a^3/(tan(1/2*d*x+1/2*c)+1)+2*b^3*(A*b-B*a)/a^4/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-1/3*A/a/(tan(1/2*d*x+1/2*c)-1)^3-1/2*(A*a+A*b-B*a)/a^2/(tan(1/2*d*x+1/2*c)-1)^2+1/2*(A*a^2*b+2*A*b^3-B*a^3-2*B*a*b^2)/a^4*ln(tan(1/2*d*x+1/2*c)-1)-1/2*(2*A*a^2+A*a*b+2*A*b^2-B*a^2-2*B*a*b)/a^3/(tan(1/2*d*x+1/2*c)-1))
```

3.257.5 Fracas [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 729, normalized size of antiderivative = 3.90

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \left[\frac{6 (Bab^3 - Ab^4) \sqrt{-a^2 + b^2} \cos(dx + c)^3 \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 + 2\sqrt{-a^2 + b^2}(a \cos(dx+c) + b) \sin(dx+c) - b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{12 (Bab^3 - Ab^4) \sqrt{a^2 - b^2} \arctan\left(-\frac{a \cos(dx+c) + b}{\sqrt{a^2 - b^2} \sin(dx+c)}\right) \cos(dx + c)^3 - 3 (Ba^5 - Aa^4b + Ba^3b^2 - Aa^2b^3 -$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `[1/12*(6*(B*a*b^3 - A*b^4)*sqrt(-a^2 + b^2)*cos(d*x + c)^3*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 3*(B*a^5 - A*a^4*b + B*a^3*b^2 - A*a^2*b^3 - 2*B*a*b^4 + 2*A*b^5)*cos(d*x + c)^3*log(sin(d*x + c) + 1) - 3*(B*a^5 - A*a^4*b + B*a^3*b^2 - A*a^2*b^3 - 2*B*a*b^4 + 2*A*b^5)*cos(d*x + c)^3*log(-sin(d*x + c) + 1) + 2*(2*A*a^5 - 2*A*a^3*b^2 + 2*(2*A*a^5 - 3*B*a^4*b + A*a^3*b^2 + 3*B*a^2*b^3 - 3*A*a*b^4)*cos(d*x + c)^2 + 3*(B*a^5 - A*a^4*b - B*a^3*b^2 + A*a^2*b^3)*cos(d*x + c))*sin(d*x + c))/((a^6 - a^4*b^2)*d*cos(d*x + c)^3), -1/12*(12*(B*a*b^3 - A*b^4)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c)))*cos(d*x + c)^3 - 3*(B*a^5 - A*a^4*b + B*a^3*b^2 - A*a^2*b^3 - 2*B*a*b^4 + 2*A*b^5)*cos(d*x + c)^3*log(sin(d*x + c) + 1) + 3*(B*a^5 - A*a^4*b + B*a^3*b^2 - A*a^2*b^3 - 2*B*a*b^4 + 2*A*b^5)*cos(d*x + c)^3*log(-sin(d*x + c) + 1) - 2*(2*A*a^5 - 2*A*a^3*b^2 + 2*(2*A*a^5 - 3*B*a^4*b + A*a^3*b^2 + 3*B*a^2*b^3 - 3*A*a*b^4)*cos(d*x + c)^2 + 3*(B*a^5 - A*a^4*b - B*a^3*b^2 + A*a^2*b^3)*cos(d*x + c))*sin(d*x + c))/((a^6 - a^4*b^2)*d*cos(d*x + c)^3)]`

3.257.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**4/(a+b*cos(d*x+c)),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**4/(a + b*cos(c + d*x)), x)`

3.257.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.257.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 412 vs. 2(170) = 340.

Time = 0.32 (sec) , antiderivative size = 412, normalized size of antiderivative = 2.20

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{3(Ba^3 - Aa^2b + 2Bab^2 - 2Ab^3) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^4} - \frac{3(Ba^3 - Aa^2b + 2Bab^2 - 2Ab^3) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^4} + \frac{12(Bab^3 - Ab^4) \left(\pi \left\lfloor \frac{dx}{2} \right\rfloor + \frac{1}{2} \pi \right)}{a^4}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^4/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/6*(3*(B*a^3 - A*a^2*b + 2*B*a*b^2 - 2*A*b^3)*log(abs(tan(1/2*d*x + 1/2*c) + 1)))/a^4 - 3*(B*a^3 - A*a^2*b + 2*B*a*b^2 - 2*A*b^3)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^4 + 12*(B*a*b^3 - A*b^4)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/(sqrt(a^2 - b^2)*a^4) - 2*(6*A*a^2*tan(1/2*d*x + 1/2*c)^5 - 3*B*a^2*tan(1/2*d*x + 1/2*c)^5 + 3*A*a*b*tan(1/2*d*x + 1/2*c)^5 - 6*B*a*b*tan(1/2*d*x + 1/2*c)^5 + 6*A*b^2*tan(1/2*d*x + 1/2*c)^5 - 4*A*a^2*tan(1/2*d*x + 1/2*c)^3 + 12*B*a*b*tan(1/2*d*x + 1/2*c)^3 - 12*A*b^2*tan(1/2*d*x + 1/2*c)^3 + 6*A*a^2*tan(1/2*d*x + 1/2*c) + 3*B*a^2*tan(1/2*d*x + 1/2*c) - 3*A*a*b*tan(1/2*d*x + 1/2*c) - 6*B*a*b*tan(1/2*d*x + 1/2*c) + 6*A*b^2*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 - 1)^3*a^3))/d`

$$3.257. \int \frac{(A+B \cos(c+dx)) \sec^4(c+dx)}{a+b \cos(c+dx)} dx$$

3.257.9 Mupad [B] (verification not implemented)

Time = 5.41 (sec) , antiderivative size = 4696, normalized size of antiderivative = 25.11

$$\int \frac{(A + B \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^4*(a + b*cos(c + d*x))),x)`

output `(atan((((8*tan(c/2 + (d*x)/2)*(8*A^2*b^9 - B^2*a^9 - 16*A^2*a*b^8 + 3*B^2*a^8*b + 16*A^2*a^2*b^7 - 16*A^2*a^3*b^6 + 13*A^2*a^4*b^5 - 7*A^2*a^5*b^4 + 3*A^2*a^6*b^3 - A^2*a^7*b^2 + 8*B^2*a^2*b^7 - 16*B^2*a^3*b^6 + 16*B^2*a^4*b^5 - 16*B^2*a^5*b^4 + 13*B^2*a^6*b^3 - 7*B^2*a^7*b^2 - 16*A*B*a*b^8 + 2*A*B*a^8*b + 32*A*B*a^2*b^7 - 32*A*B*a^3*b^6 + 32*A*B*a^4*b^5 - 26*A*B*a^5*b^4 + 14*A*B*a^6*b^3 - 6*A*B*a^7*b^2)))/a^6 + (((8*(2*B*a^13 - 4*A*a^8*b^5 + 6*A*a^9*b^4 - 2*A*a^10*b^3 + 2*A*a^11*b^2 + 4*B*a^9*b^4 - 6*B*a^10*b^3 + 2*B*a^11*b^2 - 2*A*a^12*b - 2*B*a^12*b))/a^9 - (4*tan(c/2 + (d*x)/2)*(8*a^10*b + 8*a^8*b^3 - 16*a^9*b^2)*(2*A*b^3 - B*a^3 + A*a^2*b - 2*B*a*b^2))/a^10)*(2*A*b^3 - B*a^3 + A*a^2*b - 2*B*a*b^2))/(2*a^4))*(2*A*b^3 - B*a^3 + A*a^2*b - 2*B*a*b^2)*1i)/(2*a^4) + (((8*tan(c/2 + (d*x)/2)*(8*A^2*b^9 - B^2*a^9 - 16*A^2*a*b^8 + 3*B^2*a^8*b + 16*A^2*a^2*b^7 - 16*A^2*a^3*b^6 + 13*A^2*a^4*b^5 - 7*A^2*a^5*b^4 + 3*A^2*a^6*b^3 - A^2*a^7*b^2 + 8*B^2*a^2*b^7 - 16*B^2*a^3*b^6 + 16*B^2*a^4*b^5 - 16*B^2*a^5*b^4 + 13*B^2*a^6*b^3 - 7*B^2*a^7*b^2 - 16*A*B*a*b^8 + 2*A*B*a^8*b + 32*A*B*a^2*b^7 - 32*A*B*a^3*b^6 + 32*A*B*a^4*b^5 - 26*A*B*a^5*b^4 + 14*A*B*a^6*b^3 - 6*A*B*a^7*b^2))/a^6 - (((8*(2*B*a^13 - 4*A*a^8*b^5 + 6*A*a^9*b^4 - 2*A*a^10*b^3 + 2*A*a^11*b^2 + 4*B*a^9*b^4 - 6*B*a^10*b^3 + 2*B*a^11*b^2 - 2*A*a^12*b - 2*B*a^12*b))/a^9 + (4*tan(c/2 + (d*x)/2)*(8*a^10*b + 8*a^8*b^3 - 16*a^9*b^2)*(2*A*b^3 - B*a^3 + A*a^2*b - 2*B*a*b^2))/a^10)*(2*A*b^3 - B*a^3 + A*a^2*b - 2*B*a*b^2))/a^10)*(2*A*b^3 - B*a^3 + A*a^2*b - 2*B*a*b^2)...`

3.258 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.258.1 Optimal result 2380
 3.258.2 Mathematica [A] (verified) 2381
 3.258.3 Rubi [A] (verified) 2381
 3.258.4 Maple [A] (verified) 2386
 3.258.5 Fricas [A] (verification not implemented) 2386
 3.258.6 Sympy [F(-1)] 2387
 3.258.7 Maxima [F(-2)] 2388
 3.258.8 Giac [A] (verification not implemented) 2388
 3.258.9 Mupad [B] (verification not implemented) 2389

3.258.1 Optimal result

Integrand size = 31, antiderivative size = 263

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

$$= -\frac{(4aAb - 6a^2B - b^2B)x}{2b^4}$$

$$+ \frac{2a^2(2a^2Ab - 3Ab^3 - 3a^3B + 4ab^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{3/2}b^4(a+b)^{3/2}d}$$

$$+ \frac{(2a^2Ab - Ab^3 - 3a^3B + 2ab^2B) \sin(c+dx)}{b^3(a^2-b^2)d}$$

$$- \frac{(2aAb - 3a^2B + b^2B) \cos(c+dx) \sin(c+dx)}{2b^2(a^2-b^2)d} + \frac{a(Ab - aB) \cos^2(c+dx) \sin(c+dx)}{b(a^2-b^2)d(a+b \cos(c+dx))}$$

output

```
-1/2*(4*A*a*b-6*B*a^2-B*b^2)*x/b^4+2*a^2*(2*A*a^2*b-3*A*b^3-3*B*a^3+4*B*a*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(3/2)/b^4/(a+b)^(3/2)/d+(2*A*a^2*b-A*b^3-3*B*a^3+2*B*a*b^2)*sin(d*x+c)/b^3/(a^2-b^2)/d-1/2*(2*A*a*b-3*B*a^2+B*b^2)*cos(d*x+c)*sin(d*x+c)/b^2/(a^2-b^2)/d+a*(A*b-B*a)*cos(d*x+c)^2*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.258.2 Mathematica [A] (verified)

Time = 1.50 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.70

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{2(-4aAb+6a^2B+b^2B)(c+dx) - \frac{8a^2(-2a^2Ab+3Ab^3+3a^3B-4ab^2B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{3/2}} + 4b(Ab-2aB)}{4b^4d}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `(2*(-4*a*A*b + 6*a^2*B + b^2*B)*(c + d*x) - (8*a^2*(-2*a^2*A*b + 3*A*b^3 + 3*a^3*B - 4*a*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(3/2) + 4*b*(A*b - 2*a*B)*Sin[c + d*x] + (4*a^3*b*(A*b - a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x])) + b^2*B*Ssin[2*(c + d*x)]/(4*b^4*d)`

3.258.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3468, 25, 3042, 3528, 25, 3042, 3502, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^3 (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a+b\sin\left(c+dx+\frac{\pi}{2}\right))^2} dx$$

$$\downarrow \text{3468}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))} - \int \frac{\cos(c+dx)\left(-((-3Ba^2+2Aba+b^2B)\cos^2(c+dx))-b(Ab-aB)\cos(c+dx)+2a(Ab-aB)\right)}{a+b\cos(c+dx)} dx}{b(a^2-b^2)}$$

3.258. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
 & \int \frac{\cos(c+dx) \left(-((-3Ba^2+2Aba+b^2B) \cos^2(c+dx)) - b(Ab-aB) \cos(c+dx) + 2a(Ab-aB) \right)}{a+b \cos(c+dx)} dx + \\
 & \quad \frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} + \\
 & \quad \frac{a(Ab-aB) \sin(c+dx) \cos^2(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow 25 \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2}) \left((3Ba^2-2Aba-b^2B) \sin(c+dx+\frac{\pi}{2})^2 - b(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + 2a(Ab-aB) \right)}{a+b \sin(c+dx+\frac{\pi}{2})} dx + \\
 & \quad \frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} + \\
 & \quad \frac{a(Ab-aB) \sin(c+dx) \cos^2(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow 3042 \\
 & \int -\frac{-2(-3Ba^3+2Aba^2+2b^2Ba-Ab^3) \cos^2(c+dx) - b(-Ba^2+2Aba-b^2B) \cos(c+dx) + a(-3Ba^2+2Aba+b^2B)}{a+b \cos(c+dx)} dx - \frac{(-3a^2B+2aAb+b^2B) \sin(c+dx) \cos(c+dx)}{2b} \\
 & \quad \frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow 25 \\
 & \int -\frac{-2(-3Ba^3+2Aba^2+2b^2Ba-Ab^3) \cos^2(c+dx) - b(-Ba^2+2Aba-b^2B) \cos(c+dx) + a(-3Ba^2+2Aba+b^2B)}{a+b \cos(c+dx)} dx - \frac{(-3a^2B+2aAb+b^2B) \sin(c+dx) \cos(c+dx)}{2bd} \\
 & \quad \frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow 25 \\
 & \int -\frac{-2(-3Ba^3+2Aba^2+2b^2Ba-Ab^3) \sin(c+dx+\frac{\pi}{2})^2 - b(-Ba^2+2Aba-b^2B) \sin(c+dx+\frac{\pi}{2}) + a(-3Ba^2+2Aba+b^2B)}{a+b \sin(c+dx+\frac{\pi}{2})} dx - \frac{(-3a^2B+2aAb+b^2B) \sin(c+dx) \cos(c+dx)}{2b} \\
 & \quad \frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow 3042 \\
 & \int \frac{ab(-3Ba^2+2Aba+b^2B) + (a^2-b^2)(-6Ba^2+4Aba-b^2B) \cos(c+dx)}{a+b \cos(c+dx)} dx - \frac{2(-3a^3B+2a^2Ab+2ab^2B-Ab^3) \sin(c+dx)}{bd} - \frac{(-3a^2B+2aAb+b^2B) \sin(c+dx) \cos(c+dx)}{2bd} \\
 & \quad \frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow 3502 \\
 & \int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx
 \end{aligned}$$

3.258. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

↓ 3042

$$\frac{\int \frac{ab(-3Ba^2+2Aba+b^2B)+(a^2-b^2)(-6Ba^2+4Aba-b^2B)\sin(c+dx+\frac{\pi}{2})}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{2b} - \frac{2(-3a^3B+2a^2Ab+2ab^2B-Ab^3)\sin(c+dx)}{bd} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)}{2bd}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} \frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3214

$$\frac{x(a^2-b^2)(-6a^2B+4aAb-b^2B)}{b} - \frac{2a^2(-3a^3B+2a^2Ab+4ab^2B-3Ab^3)\int \frac{1}{a+b\cos(c+dx)} dx}{2b} - \frac{2(-3a^3B+2a^2Ab+2ab^2B-Ab^3)\sin(c+dx)}{bd} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)}{2bd}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} \frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3042

$$\frac{x(a^2-b^2)(-6a^2B+4aAb-b^2B)}{b} - \frac{2a^2(-3a^3B+2a^2Ab+4ab^2B-3Ab^3)\int \frac{1}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{2b} - \frac{2(-3a^3B+2a^2Ab+2ab^2B-Ab^3)\sin(c+dx)}{bd} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)}{2bd}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} \frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3138

$$\frac{x(a^2-b^2)(-6a^2B+4aAb-b^2B)}{b} - \frac{4a^2(-3a^3B+2a^2Ab+4ab^2B-3Ab^3)\int \frac{1}{(a-b)\tan(\frac{1}{2}(c+dx))+a+b} d\tan(\frac{1}{2}(c+dx))}{2b} - \frac{2(-3a^3B+2a^2Ab+2ab^2B-Ab^3)\sin(c+dx)}{bd} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)}{2bd}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} \frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 218

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))} + \frac{x(a^2-b^2)(-6a^2B+4aAb-b^2B)}{b} - \frac{4a^2(-3a^3B+2a^2Ab+4ab^2B-3Ab^3)\arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{bd\sqrt{a-b}\sqrt{a+b}}$$

$$\frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)\cos(c+dx)}{2bd} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)\cos(c+dx)}{2bd}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

3.258. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

output $(a*(A*b - a*B)*\cos[c + d*x]^2*\sin[c + d*x])/(b*(a^2 - b^2)*d*(a + b*\cos[c + d*x])) + (-1/2*((2*a*A*b - 3*a^2*B + b^2*B)*\cos[c + d*x]*\sin[c + d*x])/(b*d) - (((a^2 - b^2)*(4*a*A*b - 6*a^2*B - b^2*B)*x)/b - (4*a^2*(2*a^2*A*b - 3*A*b^3 - 3*a^3*B + 4*a*b^2*B)*\text{ArcTan}[\text{Sqrt}[a - b]*\text{Tan}[(c + d*x)/2]])/\text{Sqrt}[a + b])]/(\text{Sqrt}[a - b]*b*\text{Sqrt}[a + b]*d)/b - (2*(2*a^2*A*b - A*b^3 - 3*a^3*B + 2*a*b^2*B)*\sin[c + d*x])/(b*d)/(2*b))/(b*(a^2 - b^2))$

3.258.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 218 $\text{Int}[(a + (b_*)*(x_*)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] \text{ ; FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3138 $\text{Int}[(a + (b_*)*\sin[\text{Pi}/2 + (c_*) + (d_*)*(x_*)])^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \quad \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

rule 3214 $\text{Int}[(a + (b_*)*\sin[(e_*) + (f_*)*(x_*)]) / ((c_*) + (d_*)*\sin[(e_*) + (f_*)*(x_*)]), x_Symbol] \rightarrow \text{Simp}[b*(x/d), x] - \text{Simp}[(b*c - a*d)/d \quad \text{Int}[1/(c + d*\sin[e + f*x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0]$

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

3.258.4 Maple [A] (verified)

Time = 1.40 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{2 \left(\frac{(-A b^2 + 2B a b + \frac{1}{2} B b^2) \left(\tan^3 \left(\frac{d x}{2} + \frac{c}{2} \right) \right) + (-A b^2 + 2B a b - \frac{1}{2} B b^2) \tan \left(\frac{d x}{2} + \frac{c}{2} \right) + (4A a b - 6B a^2 - B b^2) \arctan \left(\tan \left(\frac{d x}{2} + \frac{c}{2} \right) \right)}{(1 + \tan^2 \left(\frac{d x}{2} + \frac{c}{2} \right))^2} \right)}{b^4}$
default	$\frac{2 \left(\frac{(-A b^2 + 2B a b + \frac{1}{2} B b^2) \left(\tan^3 \left(\frac{d x}{2} + \frac{c}{2} \right) \right) + (-A b^2 + 2B a b - \frac{1}{2} B b^2) \tan \left(\frac{d x}{2} + \frac{c}{2} \right) + (4A a b - 6B a^2 - B b^2) \arctan \left(\tan \left(\frac{d x}{2} + \frac{c}{2} \right) \right)}{(1 + \tan^2 \left(\frac{d x}{2} + \frac{c}{2} \right))^2} \right)}{b^4}$
risch	$-\frac{2x A a}{b^3} + \frac{3x B a^2}{b^4} + \frac{B x}{2b^2} - \frac{ie^{i(dx+c)} A}{2b^2 d} + \frac{ie^{i(dx+c)} B a}{b^3 d} + \frac{iB e^{-2i(dx+c)}}{8b^2 d} - \frac{iB e^{2i(dx+c)}}{8b^2 d} - \frac{2ia^3(-Ab+B)}{b^4(a^2-b^2)d(b e^{2i(c+dx)} + b)}$

input `int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \cdot \left(-\frac{2}{b^4} \cdot \left(\frac{(-A b^2 + 2B a b + \frac{1}{2} B b^2) \tan^3 \left(\frac{1}{2} d x + \frac{1}{2} c \right) + (-A b^2 + 2B a b - \frac{1}{2} B b^2) \tan \left(\frac{1}{2} d x + \frac{1}{2} c \right)}{(1 + \tan^2 \left(\frac{1}{2} d x + \frac{1}{2} c \right))^2} + \frac{4A a b - 6B a^2 - B b^2}{2} \arctan \left(\tan \left(\frac{1}{2} d x + \frac{1}{2} c \right) \right) \right) + \frac{2a^2}{b^4} \cdot \left(\frac{b a (A b - B a)}{a^2 - b^2} \tan \left(\frac{1}{2} d x + \frac{1}{2} c \right) + \frac{2A a^2 b - 3A a b^3 - 3B a^3 + 4B a b^2}{(a-b)(a+b)} \frac{1}{\left(\frac{a-b}{a+b} \right)^{1/2}} \operatorname{ctan} \left(\frac{a-b}{a+b} \tan \left(\frac{1}{2} d x + \frac{1}{2} c \right) \right) \right) \right)$$

3.258.5 Fracas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 965, normalized size of antiderivative = 3.67

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx$$

$$= \left[\frac{(6 B a^6 b - 4 A a^5 b^2 - 11 B a^4 b^3 + 8 A a^3 b^4 + 4 B a^2 b^5 - 4 A a b^6 + B b^7) dx \cos(dx + c) + (6 B a^7 - 4 A a^6 b - 11 B a^5 b^2 + 8 A a^4 b^3 + 4 B a^3 b^4 - 4 A a^2 b^5 + 4 A a b^6 - B b^7) dx \sin(dx + c)}{(a + b \cos(c + dx))^2} \right]$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output

```
[1/2*((6*B*a^6*b - 4*A*a^5*b^2 - 11*B*a^4*b^3 + 8*A*a^3*b^4 + 4*B*a^2*b^5
- 4*A*a*b^6 + B*b^7)*d*x*cos(d*x + c) + (6*B*a^7 - 4*A*a^6*b - 11*B*a^5*b^
2 + 8*A*a^4*b^3 + 4*B*a^3*b^4 - 4*A*a^2*b^5 + B*a*b^6)*d*x - (3*B*a^6 - 2*
A*a^5*b - 4*B*a^4*b^2 + 3*A*a^3*b^3 + (3*B*a^5*b - 2*A*a^4*b^2 - 4*B*a^3*b
^3 + 3*A*a^2*b^4)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) +
(2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*si
n(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2))
- (6*B*a^6*b - 4*A*a^5*b^2 - 10*B*a^4*b^3 + 6*A*a^3*b^4 + 4*B*a^2*b^5 - 2
*A*a*b^6 - (B*a^4*b^3 - 2*B*a^2*b^5 + B*b^7)*cos(d*x + c)^2 + (3*B*a^5*b^2
- 2*A*a^4*b^3 - 6*B*a^3*b^4 + 4*A*a^2*b^5 + 3*B*a*b^6 - 2*A*b^7)*cos(d*x
+ c))*sin(d*x + c))/((a^4*b^5 - 2*a^2*b^7 + b^9)*d*cos(d*x + c) + (a^5*b^4
- 2*a^3*b^6 + a*b^8)*d), 1/2*((6*B*a^6*b - 4*A*a^5*b^2 - 11*B*a^4*b^3 + 8
*A*a^3*b^4 + 4*B*a^2*b^5 - 4*A*a*b^6 + B*b^7)*d*x*cos(d*x + c) + (6*B*a^7
- 4*A*a^6*b - 11*B*a^5*b^2 + 8*A*a^4*b^3 + 4*B*a^3*b^4 - 4*A*a^2*b^5 + B*a
*b^6)*d*x - 2*(3*B*a^6 - 2*A*a^5*b - 4*B*a^4*b^2 + 3*A*a^3*b^3 + (3*B*a^5*
b - 2*A*a^4*b^2 - 4*B*a^3*b^3 + 3*A*a^2*b^4)*cos(d*x + c))*sqrt(a^2 - b^2)
*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (6*B*a^6*b
- 4*A*a^5*b^2 - 10*B*a^4*b^3 + 6*A*a^3*b^4 + 4*B*a^2*b^5 - 2*A*a*b^6 - (B
*a^4*b^3 - 2*B*a^2*b^5 + B*b^7)*cos(d*x + c)^2 + (3*B*a^5*b^2 - 2*A*a^4*b^
3 - 6*B*a^3*b^4 + 4*A*a^2*b^5 + 3*B*a*b^6 - 2*A*b^7)*cos(d*x + c))*sin(...
```

3.258.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.258.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de
```

3.258.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.29

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{4(3Ba^5 - 2Aa^4b - 4Ba^3b^2 + 3Aa^2b^3) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \text{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^2b^4 - b^6)\sqrt{a^2 - b^2}} - \frac{4(Ba^4 \tan(\frac{1}{2} dx + \frac{1}{2} c) + Aa^3 \tan^2(\frac{1}{2} dx + \frac{1}{2} c) + Ab^2 \tan^3(\frac{1}{2} dx + \frac{1}{2} c))}{(a^2b^3 - b^5) \left(a \tan(\frac{1}{2} dx + \frac{1}{2} c) + b \right)}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")
```

```
output 1/2*(4*(3*B*a^5 - 2*A*a^4*b - 4*B*a^3*b^2 + 3*A*a^2*b^3)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^2*b^4 - b^6)*sqrt(a^2 - b^2)) - 4*(B*a^4*tan(1/2*d*x + 1/2*c) - A*a^3*b*tan(1/2*d*x + 1/2*c))/((a^2*b^3 - b^5)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)) + (6*B*a^2 - 4*A*a*b + B*b^2)*(d*x + c)/b^4 - 2*(4*B*a*tan(1/2*d*x + 1/2*c)^3 - 2*A*b*tan(1/2*d*x + 1/2*c)^3 + B*b*tan(1/2*d*x + 1/2*c)^3 + 4*B*a*tan(1/2*d*x + 1/2*c) - 2*A*b*tan(1/2*d*x + 1/2*c) - B*b*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 + 1)^2*b^3))/d
```

3.258. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

3.259 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.259.1 Optimal result 2390
 3.259.2 Mathematica [A] (verified) 2390
 3.259.3 Rubi [A] (verified) 2391
 3.259.4 Maple [A] (verified) 2394
 3.259.5 Fricas [B] (verification not implemented) 2394
 3.259.6 Sympy [F(-1)] 2395
 3.259.7 Maxima [F(-2)] 2396
 3.259.8 Giac [B] (verification not implemented) 2396
 3.259.9 Mupad [B] (verification not implemented) 2397

3.259.1 Optimal result

Integrand size = 31, antiderivative size = 155

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

$$= \frac{(Ab - 2aB)x}{b^3} - \frac{2a(a^2Ab - 2Ab^3 - 2a^3B + 3ab^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{3/2}b^3(a+b)^{3/2}d}$$

$$+ \frac{B \sin(c+dx)}{b^2d} - \frac{a^2(Ab - aB) \sin(c+dx)}{b^2(a^2 - b^2)d(a+b \cos(c+dx))}$$

```
output (A*b-2*B*a)*x/b^3-2*a*(A*a^2*b-2*A*b^3-2*B*a^3+3*B*a*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(3/2)/b^3/(a+b)^(3/2)/d+B*sin(d*x+c)/b^2/d-a^2*(A*b-B*a)*sin(d*x+c)/b^2/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.259.2 Mathematica [A] (verified)

Time = 1.14 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.95

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

$$= \frac{(Ab - 2aB)(c+dx) + \frac{2a(-a^2Ab+2Ab^3+2a^3B-3ab^2B) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{3/2}} + bB \sin(c+dx) + \frac{a^2b(-Ab+aB)}{(a-b)(a+b)(a+b \cos(c+dx))}}{b^3d}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `((A*b - 2*a*B)*(c + d*x) + (2*a*(-(a^2*A*b) + 2*A*b^3 + 2*a^3*B - 3*a*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(3/2) + b*B*Sin[c + d*x] + (a^2*b*(-(A*b) + a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x]))/(b^3*d)`

3.259.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.19, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3467, 3042, 3502, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^2(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx$$

↓ 3467

$$\int \frac{b(a^2-b^2)B\cos^2(c+dx)+(a^2-b^2)(Ab-aB)\cos(c+dx)+ab(Ab-aB)}{a+b\cos(c+dx)} dx - \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3042

$$\int \frac{b(a^2-b^2)B\sin(c+dx+\frac{\pi}{2})^2+(a^2-b^2)(Ab-aB)\sin(c+dx+\frac{\pi}{2})+ab(Ab-aB)}{a+b\sin(c+dx+\frac{\pi}{2})} dx - \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3502

$$\int \frac{\frac{a(Ab-aB)b^2+(a^2-b^2)(Ab-2aB)\cos(c+dx)b}{a+b\cos(c+dx)} dx + \frac{B(a^2-b^2)\sin(c+dx)}{d}}{b^2(a^2-b^2)} - \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3042

$$\int \frac{\frac{a(Ab-aB)b^2+(a^2-b^2)(Ab-2aB)\sin(c+dx+\frac{\pi}{2})b}{a+b\sin(c+dx+\frac{\pi}{2})} dx + \frac{B(a^2-b^2)\sin(c+dx)}{d}}{b^2(a^2-b^2)} - \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))}$$

3.259. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
 & \downarrow \text{3214} \\
 & \frac{x(a^2-b^2)(Ab-2aB)-a(-2a^3B+a^2Ab+3ab^2B-2Ab^3) \int \frac{1}{a+b\cos(c+dx)} dx + \frac{B(a^2-b^2)\sin(c+dx)}{d}}{b^2(a^2-b^2)} - \\
 & \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))} \\
 & \downarrow \text{3042} \\
 & \frac{x(a^2-b^2)(Ab-2aB)-a(-2a^3B+a^2Ab+3ab^2B-2Ab^3) \int \frac{1}{a+b\sin(c+dx+\frac{\pi}{2})} dx + \frac{B(a^2-b^2)\sin(c+dx)}{d}}{b^2(a^2-b^2)} - \\
 & \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))} \\
 & \downarrow \text{3138} \\
 & \frac{x(a^2-b^2)(Ab-2aB)-\frac{2a(-2a^3B+a^2Ab+3ab^2B-2Ab^3) \int \frac{1}{(a-b)\tan^2(\frac{1}{2}(c+dx))+a+b} d \tan(\frac{1}{2}(c+dx))}{b} + \frac{B(a^2-b^2)\sin(c+dx)}{d}}{b^2(a^2-b^2)} - \\
 & \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))} \\
 & \downarrow \text{218} \\
 & \frac{B(a^2-b^2)\sin(c+dx)}{d} + \frac{x(a^2-b^2)(Ab-2aB)-\frac{2a(-2a^3B+a^2Ab+3ab^2B-2Ab^3) \arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{d\sqrt{a-b}\sqrt{a+b}}}{b^2(a^2-b^2)} - \\
 & \frac{a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)(a+b\cos(c+dx))}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `-((a^2*(A*b - a*B)*Sin[c + d*x])/(b^2*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))) + (((a^2 - b^2)*(A*b - 2*a*B)*x - (2*a*(a^2*A*b - 2*A*b^3 - 2*a^3*B + 3*a*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*d))/b + ((a^2 - b^2)*B*SIN[c + d*x])/d)/(b^2*(a^2 - b^2))`

3.259.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.259.4 Maple [A] (verified)

Time = 1.27 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.32

method	result
derivativedivides	$\frac{2a \left(\frac{ba(Ab-Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2-b^2) \left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)} + \frac{(A a^2 b - 2 A b^3 - 2 B a^3 + 3 B a b^2) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b) \sqrt{(a-b)(a+b)}} \right)}{b^3 d}$
default	$\frac{2a \left(\frac{ba(Ab-Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2-b^2) \left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)} + \frac{(A a^2 b - 2 A b^3 - 2 B a^3 + 3 B a b^2) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b) \sqrt{(a-b)(a+b)}} \right)}{b^3 d}$
risch	$\frac{x A}{b^2} - \frac{2 x B a}{b^3} - \frac{i B e^{i(dx+c)}}{2 b^2 d} + \frac{i B e^{-i(dx+c)}}{2 b^2 d} + \frac{2 i a^2 (-A b + B a) (a e^{i(dx+c)} + b)}{b^3 (a^2 - b^2) d (b e^{2i(dx+c)} + 2 a e^{i(dx+c)} + b)} - \frac{a^3 \ln\left(e^{i(dx+c)} - \frac{i a^2 - i b}{b}\right)}{\sqrt{-a^2 + b^2} (a + b)}$

input `int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

output `1/d*(-2*a/b^3*(b*a*(A*b-B*a)/(a^2-b^2)*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)+(A*a^2*b-2*A*b^3-2*B*a^3+3*B*a*b^2)/(a-b)/(a+b)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))+2/b^3*(B*b*tan(1/2*d*x+1/2*c)/(1+tan(1/2*d*x+1/2*c)^2)+(A*b-2*B*a)*arctan(tan(1/2*d*x+1/2*c))))`

3.259.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. 2(148) = 296.

Time = 0.34 (sec) , antiderivative size = 788, normalized size of antiderivative = 5.08

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx$$

$$= \left[\frac{2(2Ba^5b - Aa^4b^2 - 4Ba^3b^3 + 2Aa^2b^4 + 2Bab^5 - Ab^6)dx \cos(dx + c) + 2(2Ba^6 - Aa^5b - 4Ba^4b^2 + 2Ba^5b - Aa^4b^2 - 4Ba^3b^3 + 2Aa^2b^4 + 2Bab^5 - Ab^6)dx \cos(dx + c) + (2Ba^6 - Aa^5b - 4Ba^4b^2 + 2Ba^5b - Aa^4b^2 - 4Ba^3b^3 + 2Aa^2b^4 + 2Bab^5 - Ab^6)dx \cos(dx + c)}{(2Ba^6 - Aa^5b - 4Ba^4b^2 + 2Ba^5b - Aa^4b^2 - 4Ba^3b^3 + 2Aa^2b^4 + 2Bab^5 - Ab^6)dx \cos(dx + c)} \right]$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `[-1/2*(2*(2*B*a^5*b - A*a^4*b^2 - 4*B*a^3*b^3 + 2*A*a^2*b^4 + 2*B*a*b^5 - A*b^6)*d*x*cos(d*x + c) + 2*(2*B*a^6 - A*a^5*b - 4*B*a^4*b^2 + 2*A*a^3*b^3 + 2*B*a^2*b^4 - A*a*b^5)*d*x + (2*B*a^5 - A*a^4*b - 3*B*a^3*b^2 + 2*A*a^2*b^3 + (2*B*a^4*b - A*a^3*b^2 - 3*B*a^2*b^3 + 2*A*a*b^4)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(2*B*a^5*b - A*a^4*b^2 - 3*B*a^3*b^3 + A*a^2*b^4 + B*a*b^5 + (B*a^4*b^2 - 2*B*a^2*b^4 + B*b^6)*cos(d*x + c))*sin(d*x + c))/((a^4*b^4 - 2*a^2*b^6 + b^8)*d*cos(d*x + c) + (a^5*b^3 - 2*a^3*b^5 + a*b^7)*d), -((2*B*a^5*b - A*a^4*b^2 - 4*B*a^3*b^3 + 2*A*a^2*b^4 + 2*B*a*b^5 - A*b^6)*d*x*cos(d*x + c) + (2*B*a^6 - A*a^5*b - 4*B*a^4*b^2 + 2*A*a^3*b^3 + 2*B*a^2*b^4 - A*a*b^5)*d*x - (2*B*a^5 - A*a^4*b - 3*B*a^3*b^2 + 2*A*a^2*b^3 + (2*B*a^4*b - A*a^3*b^2 - 3*B*a^2*b^3 + 2*A*a*b^4)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (2*B*a^5*b - A*a^4*b^2 - 3*B*a^3*b^3 + A*a^2*b^4 + B*a*b^5 + (B*a^4*b^2 - 2*B*a^2*b^4 + B*b^6)*cos(d*x + c))*sin(d*x + c))/((a^4*b^4 - 2*a^2*b^6 + b^8)*d*cos(d*x + c) + (a^5*b^3 - 2*a^3*b^5 + a*b^7)*d)]`

3.259.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.259.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.259.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1116 vs. 2(148) = 296.

Time = 0.39 (sec) , antiderivative size = 1116, normalized size of antiderivative = 7.20

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output

```
((4*B*a^6*b^2 - 2*A*a^5*b^3 - 2*B*a^5*b^3 + A*a^4*b^4 - 9*B*a^4*b^4 + 5*A*
a^3*b^5 + 4*B*a^3*b^5 - 2*A*a^2*b^6 + 5*B*a^2*b^6 - 3*A*a*b^7 - 2*B*a*b^7
+ A*b^8 + 2*B*a^3*abs(-a^2*b^3 + b^5) - A*a^2*b*abs(-a^2*b^3 + b^5) - B*a^
2*b*abs(-a^2*b^3 + b^5) + A*a*b^2*abs(-a^2*b^3 + b^5) - 2*B*a*b^2*abs(-a^2
*b^3 + b^5) + A*b^3*abs(-a^2*b^3 + b^5))*(pi*floor(1/2*(d*x + c)/pi + 1/2)
+ arctan(2*sqrt(1/2)*tan(1/2*d*x + 1/2*c)/sqrt((2*a^3*b^2 - 2*a*b^4 + sqr
t(-4*(a^3*b^2 + a^2*b^3 - a*b^4 - b^5)*(a^3*b^2 - a^2*b^3 - a*b^4 + b^5) +
4*(a^3*b^2 - a*b^4)^2))/(a^3*b^2 - a^2*b^3 - a*b^4 + b^5))))/(a^3*b^2*abs
(-a^2*b^3 + b^5) - a*b^4*abs(-a^2*b^3 + b^5) + (a^2*b^3 - b^5)^2) + ((a^2*
b - a*b^2 - b^3)*sqrt(a^2 - b^2)*A*abs(-a^2*b^3 + b^5)*abs(-a + b) - (2*a^
3 - a^2*b - 2*a*b^2)*sqrt(a^2 - b^2)*B*abs(-a^2*b^3 + b^5)*abs(-a + b) - (
2*a^5*b^3 - a^4*b^4 - 5*a^3*b^5 + 2*a^2*b^6 + 3*a*b^7 - b^8)*sqrt(a^2 - b^
2)*A*abs(-a + b) + (4*a^6*b^2 - 2*a^5*b^3 - 9*a^4*b^4 + 4*a^3*b^5 + 5*a^2*
b^6 - 2*a*b^7)*sqrt(a^2 - b^2)*B*abs(-a + b))*(pi*floor(1/2*(d*x + c)/pi +
1/2) + arctan(2*sqrt(1/2)*tan(1/2*d*x + 1/2*c)/sqrt((2*a^3*b^2 - 2*a*b^4
- sqrt(-4*(a^3*b^2 + a^2*b^3 - a*b^4 - b^5)*(a^3*b^2 - a^2*b^3 - a*b^4 + b
^5) + 4*(a^3*b^2 - a*b^4)^2))/(a^3*b^2 - a^2*b^3 - a*b^4 + b^5))))/((a^2*b
^3 - b^5)^2*(a^2 - 2*a*b + b^2) - (a^5*b^2 - 2*a^4*b^3 + 2*a^2*b^5 - a*b^6
)*abs(-a^2*b^3 + b^5)) + 2*(2*B*a^3*tan(1/2*d*x + 1/2*c)^3 - A*a^2*b*tan(1
/2*d*x + 1/2*c)^3 - B*a^2*b*tan(1/2*d*x + 1/2*c)^3 - B*a*b^2*tan(1/2*d*...
```

3.259.9 Mupad [B] (verification not implemented)

Time = 5.50 (sec) , antiderivative size = 3276, normalized size of antiderivative = 21.14

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

output

$$\begin{aligned}
& (\log(\tan(c/2 + (d*x)/2) + 1i)*(A*b - 2*B*a)*1i)/(b^3*d) - ((2*\tan(c/2 + (d*x)/2)^3*(A*a^2*b - B*b^3 - 2*B*a^3 + B*a*b^2 + B*a^2*b))/(b^2*(a + b)*(a - b)) + (2*\tan(c/2 + (d*x)/2)*(B*b^3 - 2*B*a^3 + A*a^2*b + B*a*b^2 - B*a^2*b))/(b^2*(a + b)*(a - b)))/(d*(a + b + \tan(c/2 + (d*x)/2)^4*(a - b) + 2*a*\tan(c/2 + (d*x)/2)^2) - (\log(\tan(c/2 + (d*x)/2) - 1i)*(A*b*1i - B*a*2i))/(b^3*d) - (a*\operatorname{atan}(((a*((32*\tan(c/2 + (d*x)/2)*(A^2*b^8 + 8*B^2*a^8 - 2*A^2*a*b^7 - 8*B^2*a^7*b + 3*A^2*a^2*b^6 + 4*A^2*a^3*b^5 - 5*A^2*a^4*b^4 - 2*A^2*a^5*b^3 + 2*A^2*a^6*b^2 + 4*B^2*a^2*b^6 - 8*B^2*a^3*b^5 + 5*B^2*a^4*b^4 + 16*B^2*a^5*b^3 - 16*B^2*a^6*b^2 - 4*A*B*a*b^7 - 8*A*B*a^7*b + 8*A*B*a^2*b^6 - 8*A*B*a^3*b^5 - 16*A*B*a^4*b^4 + 18*A*B*a^5*b^3 + 8*A*B*a^6*b^2)))/(a*b^6 + b^7 - a^2*b^5 - a^3*b^4) + (a*((32*(A*a^2*b^10 - A*b^12 - 3*A*a^3*b^9 + A*a^5*b^7 - 3*B*a^2*b^10 - 3*B*a^3*b^9 + 5*B*a^4*b^8 + B*a^5*b^7 - 2*B*a^6*b^6 + 2*A*a*b^11 + 2*B*a*b^11)))/(a*b^8 + b^9 - a^2*b^7 - a^3*b^6) - (32*a*\tan(c/2 + (d*x)/2)*(-(a + b)^3*(a - b)^3)^{(1/2)}*(2*A*b^3 + 2*B*a^3 - A*a^2*b - 3*B*a*b^2)*(2*a*b^11 - 2*a^2*b^10 - 4*a^3*b^9 + 4*a^4*b^8 + 2*a^5*b^7 - 2*a^6*b^6)))/((a*b^6 + b^7 - a^2*b^5 - a^3*b^4)*(b^9 - 3*a^2*b^7 + 3*a^4*b^5 - a^6*b^3)))*(-(a + b)^3*(a - b)^3)^{(1/2)}*(2*A*b^3 + 2*B*a^3 - A*a^2*b - 3*B*a*b^2))/(b^9 - 3*a^2*b^7 + 3*a^4*b^5 - a^6*b^3))*(-(a + b)^3*(a - b)^3)^{(1/2)}*(2*A*b^3 + 2*B*a^3 - A*a^2*b - 3*B*a*b^2)*1i)/(b^9 - 3*a^2*b^7 + 3*a^4*b^5 - a^6*b^3) + (a*((32*\tan(c/2 + (d*x)/2)*(A^2*b^8 + ...
\end{aligned}$$

3.260 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.260.1 Optimal result 2399
 3.260.2 Mathematica [A] (verified) 2399
 3.260.3 Rubi [A] (verified) 2400
 3.260.4 Maple [A] (verified) 2402
 3.260.5 Fricas [B] (verification not implemented) 2403
 3.260.6 Sympy [F(-1)] 2404
 3.260.7 Maxima [F(-2)] 2404
 3.260.8 Giac [A] (verification not implemented) 2404
 3.260.9 Mupad [B] (verification not implemented) 2405

3.260.1 Optimal result

Integrand size = 29, antiderivative size = 122

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{Bx}{b^2} - \frac{2(Ab^3 + a^3B - 2ab^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{3/2}b^2(a+b)^{3/2}d} + \frac{a(Ab - aB) \sin(c+dx)}{b(a^2 - b^2)d(a+b \cos(c+dx))}$$

```
output B*x/b^2-2*(A*b^3+B*a^3-2*B*a*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(3/2)/b^2/(a+b)^(3/2)/d+a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.260.2 Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.98

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{B(c+dx) - \frac{2(Ab^3+a(a^2-2b^2)B) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{3/2}} + \frac{ab(Ab-aB) \sin(c+dx)}{(a-b)(a+b)(a+b \cos(c+dx))}}{b^2d}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output $(B*(c + d*x) - (2*(A*b^3 + a*(a^2 - 2*b^2)*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^{(3/2)} + (a*b*(A*b - a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x]))/(b^2*d)$

3.260.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.22, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$, Rules used = {3042, 3447, 3042, 3500, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c + dx + \frac{\pi}{2})(A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A \cos(c + dx) + B \cos^2(c + dx)}{(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A \sin(c + dx + \frac{\pi}{2}) + B \sin(c + dx + \frac{\pi}{2})^2}{(a + b \sin(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \frac{\int \frac{b(Ab - aB) - (a^2 - b^2)B \cos(c + dx)}{a + b \cos(c + dx)} dx}{b(a^2 - b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \frac{\int \frac{b(Ab - aB) - (a^2 - b^2)B \sin(c + dx + \frac{\pi}{2})}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{b(a^2 - b^2)} \\
 & \quad \downarrow \text{3214}
 \end{aligned}$$

3.260. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \frac{(aB(a^2 - 2b^2) + Ab^3) \int \frac{1}{a + b \cos(c + dx)} dx - \frac{Bx(a^2 - b^2)}{b}}{b(a^2 - b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \frac{(aB(a^2 - 2b^2) + Ab^3) \int \frac{1}{a + b \sin(c + dx + \frac{\pi}{2})} dx - \frac{Bx(a^2 - b^2)}{b}}{b(a^2 - b^2)} \\
& \quad \downarrow \text{3138} \\
& \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \frac{2(aB(a^2 - 2b^2) + Ab^3) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx)) - \frac{Bx(a^2 - b^2)}{b}}{b(a^2 - b^2)} \\
& \quad \downarrow \text{218} \\
& \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \frac{2(aB(a^2 - 2b^2) + Ab^3) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - \frac{Bx(a^2 - b^2)}{b}}{bd\sqrt{a-b}\sqrt{a+b} b(a^2 - b^2)}
\end{aligned}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `-(((a^2 - b^2)*B*x)/b) + (2*(A*b^3 + a*(a^2 - 2*b^2)*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d)/(b*(a^2 - b^2)) + (a*(A*b - a*B)*Sin[c + d*x])/(b*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))`

3.260.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d *Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.260.4 Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.32

method	result
derivativedivides	$2 \left(\frac{ba(Ab - Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2 - b^2) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + a + b \right) \right)} + \frac{(Ab^3 + Ba^3 - 2Ba^2b) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}} \right) \frac{d}{b^2} + \frac{2B \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{d}$
default	$2 \left(\frac{ba(Ab - Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2 - b^2) \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) + a + b \right) \right)} + \frac{(Ab^3 + Ba^3 - 2Ba^2b) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}} \right) \frac{d}{b^2} + \frac{2B \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{d}$
risch	$\frac{Bx}{b^2} - \frac{2ia(Ab - Ba)(ae^{i(dx+c)} + b)}{b^2(-a^2 + b^2)d(be^{2i(dx+c)} + 2ae^{i(dx+c)} + b)} - \frac{b \ln\left(\frac{e^{i(dx+c)} - ia^2 - ib^2 - a\sqrt{-a^2 + b^2}}{b\sqrt{-a^2 + b^2}}\right)A}{\sqrt{-a^2 + b^2}(a+b)(a-b)d} - \frac{\ln\left(\frac{e^{i(dx+c)} - ia^2 - ib^2 - a\sqrt{-a^2 + b^2}}{b\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2}(a-b)}$

input `int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

$$3.260. \quad \int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.260.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.260.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.260.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.63

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{2(Ba^3-2Bab^2+Ab^3) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\sqrt{a^2-b^2}} \right) \right)}{(a^2b^2-b^4)\sqrt{a^2-b^2}} + \frac{(dx+c)B}{b^2} - \frac{2(Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + \frac{1}{2} dx + \frac{1}{2} c)}{(a^2b-b^3)}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

3.260. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

output $(2*(B*a^3 - 2*B*a*b^2 + A*b^3)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^2*b^2 - b^4)*sqrt(a^2 - b^2)) + (d*x + c)*B/b^2 - 2*(B*a^2*tan(1/2*d*x + 1/2*c) - A*a*b*tan(1/2*d*x + 1/2*c))/((a^2*b - b^3)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b))/d$

3.260.9 Mupad [B] (verification not implemented)

Time = 8.38 (sec) , antiderivative size = 3775, normalized size of antiderivative = 30.94

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

output $(2*B*atan((B*((B*((32*(A*a^2*b^7 - B*b^9 - A*b^9 - A*a^3*b^6 + B*a^2*b^7 - 3*B*a^3*b^6 + B*a^5*b^4 + A*a*b^8 + 2*B*a*b^8)))/(a*b^5 + b^6 - a^2*b^4 - a^3*b^3) - (B*tan(c/2 + (d*x)/2)*(2*a*b^9 - 2*a^2*b^8 - 4*a^3*b^7 + 4*a^4*b^6 + 2*a^5*b^5 - 2*a^6*b^4)*32i)/(b^2*(a*b^4 + b^5 - a^2*b^3 - a^3*b^2)))*1i)/b^2 + (32*tan(c/2 + (d*x)/2)*(A^2*b^6 + 2*B^2*a^6 + B^2*b^6 - 2*B^2*a*b^5 - 2*B^2*a^5*b + 3*B^2*a^2*b^4 + 4*B^2*a^3*b^3 - 5*B^2*a^4*b^2 - 4*A*B*a*b^5 + 2*A*B*a^3*b^3))/(a*b^4 + b^5 - a^2*b^3 - a^3*b^2))/b^2 - (B*((B*((32*(A*a^2*b^7 - B*b^9 - A*b^9 - A*a^3*b^6 + B*a^2*b^7 - 3*B*a^3*b^6 + B*a^5*b^4 + A*a*b^8 + 2*B*a*b^8)))/(a*b^5 + b^6 - a^2*b^4 - a^3*b^3) + (B*tan(c/2 + (d*x)/2)*(2*a*b^9 - 2*a^2*b^8 - 4*a^3*b^7 + 4*a^4*b^6 + 2*a^5*b^5 - 2*a^6*b^4)*32i)/(b^2*(a*b^4 + b^5 - a^2*b^3 - a^3*b^2)))*1i)/b^2 - (32*tan(c/2 + (d*x)/2)*(A^2*b^6 + 2*B^2*a^6 + B^2*b^6 - 2*B^2*a*b^5 - 2*B^2*a^5*b + 3*B^2*a^2*b^4 + 4*B^2*a^3*b^3 - 5*B^2*a^4*b^2 - 4*A*B*a*b^5 + 2*A*B*a^3*b^3))/(a*b^4 + b^5 - a^2*b^3 - a^3*b^2))/b^2)/((64*(B^3*a^5 - A*B^2*b^5 + A^2*B*b^5 + 2*B^3*a*b^4 - B^3*a^4*b + 2*B^3*a^2*b^3 - 3*B^3*a^3*b^2 - 3*A*B^2*a*b^4 + A*B^2*a^2*b^3 + A*B^2*a^3*b^2))/(a*b^5 + b^6 - a^2*b^4 - a^3*b^3) + (B*((B*((32*(A*a^2*b^7 - B*b^9 - A*b^9 - A*a^3*b^6 + B*a^2*b^7 - 3*B*a^3*b^6 + B*a^5*b^4 + A*a*b^8 + 2*B*a*b^8)))/(a*b^5 + b^6 - a^2*b^4 - a^3*b^3) - (B*tan(c/2 + (d*x)/2)*(2*a*b^9 - 2*a^2*b^8 - 4*a^3*b^7 + 4*a^4*b^6 + 2*a^5*b^5 - 2*a^6*b^4)*32i)/(b^2*(a*b^4 + b^5 - a^2*b^3 - a^3*b^2)...$

3.261 $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2} dx$

3.261.1 Optimal result	2406
3.261.2 Mathematica [A] (verified)	2406
3.261.3 Rubi [A] (verified)	2407
3.261.4 Maple [A] (verified)	2409
3.261.5 Fricas [A] (verification not implemented)	2409
3.261.6 Sympy [B] (verification not implemented)	2410
3.261.7 Maxima [F(-2)]	2411
3.261.8 Giac [A] (verification not implemented)	2411
3.261.9 Mupad [B] (verification not implemented)	2412

3.261.1 Optimal result

Integrand size = 23, antiderivative size = 100

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2(aA - bB) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{(a - b)^{3/2}(a + b)^{3/2}d} - \frac{(Ab - aB) \sin(c + dx)}{(a^2 - b^2) d(a + b \cos(c + dx))}$$

```
output 2*(A*a-B*b)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(3/2)
/(a+b)^(3/2)/d-(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.261.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.97

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2(aA - bB) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{3/2}} + \frac{(-Ab+aB) \sin(c+dx)}{(a-b)(a+b)(a+b \cos(c+dx))} d$$

```
input Integrate[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^2,x]
```

```
output ((2*(a*A - b*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^
2 + b^2)^(3/2) + (((-A*b) + a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos
[c + d*x]))/d
```

3.261.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.11, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {3042, 3233, 25, 27, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2} dx \\
 & \quad \downarrow \text{3233} \\
 & -\frac{\int -\frac{aA-bB}{a+b \cos(c+dx)} dx}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{aA-bB}{a+b \cos(c+dx)} dx}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{(aA - bB) \int \frac{1}{a+b \cos(c+dx)} dx}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(aA - bB) \int \frac{1}{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} dx}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
 & \quad \downarrow \text{3138} \\
 & \frac{2(aA - bB) \int \frac{1}{(a-b) \tan^2\left(\frac{1}{2}(c+dx)\right)+a+b} d \tan\left(\frac{1}{2}(c + dx)\right)}{d(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
 & \quad \downarrow \text{218} \\
 & \frac{2(aA - bB) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{d\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

3.261. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2} dx$

```
output (2*(a*A - b*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]]/(Sqrt[a
- b]*Sqrt[a + b]*(a^2 - b^2)*d) - ((A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)
*d*(a + b*Cos[c + d*x]))
```

3.261.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

```
rule 3233 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

3.261.4 Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.28

method	result
derivativedivides	$-\frac{2(Ab-Ba)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{(a^2-b^2)\left(\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a-b\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b}\right)+\frac{2(aA-Bb)\arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}}$
default	$-\frac{2(Ab-Ba)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{(a^2-b^2)\left(\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a-b\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b}\right)+\frac{2(aA-Bb)\arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}}$
risch	$\frac{2i(Ab-Ba)(ae^{i(dx+c)}+b)}{b(-a^2+b^2)d(be^{2i(dx+c)}+2ae^{i(dx+c)}+b)}-\frac{\ln\left(\frac{e^{i(dx+c)}+ia^2-ib^2+a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)aA}{\sqrt{-a^2+b^2}(a+b)(a-b)d}+\frac{\ln\left(\frac{e^{i(dx+c)}+ia^2-ib^2+a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)(a-b)}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

output `1/d*(-2*(A*b-B*a)/(a^2-b^2)*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)+2*(A*a-B*b)/(a-b)/(a+b)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))`

3.261.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 379, normalized size of antiderivative = 3.79

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \left[-\frac{(Aa^2 - Bab + (Aab - Bb^2) \cos(dx + c))\sqrt{-a^2 + b^2} \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 + 2\sqrt{-a^2+b^2}(a \cos(dx+c) + a^2)}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{2((a^4b - 2a^2b^3 + b^5)d \cos(dx + c) + (a^5 - 2a^4b))} \right]$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`


```
output [-1/2*((A*a^2 - B*a*b + (A*a*b - B*b^2)*cos(d*x + c))*sqrt(-a^2 + b^2)*log
((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(
a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*
b*cos(d*x + c) + a^2)) - 2*(B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*sin(d*x + c
))/((a^4*b - 2*a^2*b^3 + b^5)*d*cos(d*x + c) + (a^5 - 2*a^3*b^2 + a*b^4)*d
), ((A*a^2 - B*a*b + (A*a*b - B*b^2)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(
-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (B*a^3 - A*a^2*b -
B*a*b^2 + A*b^3)*sin(d*x + c))/((a^4*b - 2*a^2*b^3 + b^5)*d*cos(d*x + c)
+ (a^5 - 2*a^3*b^2 + a*b^4)*d)]
```

3.261.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4974 vs. $2(82) = 164$.

Time = 157.10 (sec) , antiderivative size = 4974, normalized size of antiderivative = 49.74

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)
```

```
output Piecewise((zoo*x*(A + B*cos(c))/cos(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0))
, (A*tan(c/2 + d*x/2)**3/(6*b**2*d) + A*tan(c/2 + d*x/2)/(2*b**2*d) - B*ta
n(c/2 + d*x/2)**3/(6*b**2*d) + B*tan(c/2 + d*x/2)/(2*b**2*d), Eq(a, b)), (
-A/(2*b**2*d*tan(c/2 + d*x/2)) - A/(6*b**2*d*tan(c/2 + d*x/2)**3) + B/(2*b
**2*d*tan(c/2 + d*x/2)) - B/(6*b**2*d*tan(c/2 + d*x/2)**3), Eq(a, -b)), (x
*(A + B*cos(c))/(a + b*cos(c))**2, Eq(d, 0)), (A*a**2*log(-sqrt(-a/(a - b)
- b/(a - b)) + tan(c/2 + d*x/2))*tan(c/2 + d*x/2)**2/(a**4*d*sqrt(-a/(a -
b) - b/(a - b)) - b/(a - b))*tan(c/2 + d*x/2)**2 + a**4*d*sqrt(-a/(a - b)
- b/(a - b)) - 2*a**3*b*d*sqrt(-a/(a - b) - b/(a - b))*tan(c/2 + d*x/2)**2
- 2*a**2*b*
**2*d*sqrt(-a/(a - b) - b/(a - b)) + 2*a*b**3*d*sqrt(-a/(a - b) - b/(a - b)
)*tan(c/2 + d*x/2)**2 - b**4*d*sqrt(-a/(a - b) - b/(a - b))*tan(c/2 + d*x/
2)**2 + b**4*d*sqrt(-a/(a - b) - b/(a - b))) + A*a**2*log(-sqrt(-a/(a - b)
- b/(a - b)) + tan(c/2 + d*x/2))/(a**4*d*sqrt(-a/(a - b) - b/(a - b))*tan
(c/2 + d*x/2)**2 + a**4*d*sqrt(-a/(a - b) - b/(a - b)) - 2*a**3*b*d*sqrt(-
a/(a - b) - b/(a - b))*tan(c/2 + d*x/2)**2 - 2*a**2*b**2*d*sqrt(-a/(a - b)
- b/(a - b)) + 2*a*b**3*d*sqrt(-a/(a - b) - b/(a - b))*tan(c/2 + d*x/2)**
2 - b**4*d*sqrt(-a/(a - b) - b/(a - b))*tan(c/2 + d*x/2)**2 + b**4*d*sqrt(
-a/(a - b) - b/(a - b))) - A*a**2*log(sqrt(-a/(a - b) - b/(a - b)) + tan(c
/2 + d*x/2))*tan(c/2 + d*x/2)**2/(a**4*d*sqrt(-a/(a - b) - b/(a - b))*tan(
c/2 + d*x/2)**2 + a**4*d*sqrt(-a/(a - b) - b/(a - b)) - 2*a**3*b*d*sqrt...
```

3.261. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2} dx$

3.261.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

3.261.8 Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.59

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx =$$

$$2 \left(\frac{\left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right) (Aa - Bb)}{(a^2 - b^2)^{\frac{3}{2}}} - \frac{Ba \tan(\frac{1}{2} dx + \frac{1}{2} c) - Ab \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(a \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - b \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + a + b)} \right) dx$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")
```

```
output -2*((pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2
*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))*(A*a - B*b)/(a^2
- b^2)^(3/2) - (B*a*tan(1/2*d*x + 1/2*c) - A*b*tan(1/2*d*x + 1/2*c))/((a*
tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)*(a^2 - b^2))/d
```

3.261.9 Mupad [B] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.13

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2 \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)(2a - 2b)}{2\sqrt{a+b}\sqrt{a-b}}\right) (Aa - Bb)}{d(a+b)^{3/2}(a-b)^{3/2}} - \frac{2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) (Ab - Ba)}{d(a+b)(a-b)\left((a-b)\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + a + b\right)}$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^2,x)`output `(2*atan((tan(c/2 + (d*x)/2)*(2*a - 2*b))/(2*(a + b)^(1/2)*(a - b)^(1/2)))*
(A*a - B*b))/(d*(a + b)^(3/2)*(a - b)^(3/2)) - (2*tan(c/2 + (d*x)/2)*(A*b
- B*a))/(d*(a + b)*(a - b)*(a + b + tan(c/2 + (d*x)/2)^2*(a - b)))`

3.262 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx$

3.262.1 Optimal result 2413
 3.262.2 Mathematica [A] (verified) 2413
 3.262.3 Rubi [A] (verified) 2414
 3.262.4 Maple [A] (verified) 2416
 3.262.5 Fricas [B] (verification not implemented) 2417
 3.262.6 Sympy [F] 2418
 3.262.7 Maxima [F(-2)] 2418
 3.262.8 Giac [A] (verification not implemented) 2419
 3.262.9 Mupad [B] (verification not implemented) 2419

3.262.1 Optimal result

Integrand size = 29, antiderivative size = 133

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = -\frac{2(2a^2Ab - Ab^3 - a^3B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^2(a-b)^{3/2}(a+b)^{3/2}d} + \frac{A \operatorname{arctanh}(\sin(c + dx))}{a^2d} + \frac{b(Ab - aB) \sin(c + dx)}{a(a^2 - b^2)d(a + b \cos(c + dx))}$$

```
output -2*(2*A*a^2*b-A*b^3-B*a^3)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^2/(a-b)^(3/2)/(a+b)^(3/2)/d+A*arctanh(sin(d*x+c))/a^2/d+b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.262.2 Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.44

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{\cos(c + dx)(B + A \sec(c + dx)) \left(\frac{2(-2a^2Ab + Ab^3 + a^3B) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{3/2}} - A \log\left(\cos\left(\frac{1}{2}(c + dx)\right)\right) - s \right)}{a^2d(A + B \cos(c + dx))}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

output `(Cos[c + d*x]*(B + A*Sec[c + d*x])*((2*(-2*a^2*A*b + A*b^3 + a^3*B)*ArcTan h[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(3/2) - A*Log [Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (a*b*(A*b - a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x]))) / (a^2*d*(A + B*Cos[c + d*x]))`

3.262.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.19, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 3479, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{\int \frac{(A(a^2-b^2)-a(Ab-aB)\cos(c+dx))\sec(c+dx)}{a+b\cos(c+dx)} dx}{a(a^2-b^2)} + \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{A(a^2-b^2)-a(Ab-aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{a(a^2-b^2)} + \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3480} \\
 & \frac{\frac{A(a^2-b^2)}{a} \int \sec(c+dx) dx}{a(a^2-b^2)} - \frac{(a^3(-B)+2a^2Ab-Ab^3)}{a} \int \frac{1}{a+b\cos(c+dx)} dx}{a(a^2-b^2)} + \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{A(a^2-b^2)}{a} \int \csc(c+dx+\frac{\pi}{2}) dx}{a(a^2-b^2)} - \frac{(a^3(-B)+2a^2Ab-Ab^3)}{a} \int \frac{1}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{a(a^2-b^2)} + \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))}
 \end{aligned}$$

3.262. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow \text{3138} \\
& \frac{\frac{A(a^2-b^2) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2(a^3(-B)+2a^2Ab-Ab^3) \int \frac{1}{(a-b)\tan^2(\frac{1}{2}(c+dx))+a+b} d \tan(\frac{1}{2}(c+dx))}{ad}}{a(a^2-b^2)} + \\
& \quad \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))} \\
& \downarrow \text{218} \\
& \frac{\frac{A(a^2-b^2) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2(a^3(-B)+2a^2Ab-Ab^3) \arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{a(a^2-b^2)} + \\
& \quad \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))} \\
& \downarrow \text{4257} \\
& \frac{\frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))} + \frac{A(a^2-b^2)\operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2(a^3(-B)+2a^2Ab-Ab^3) \arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{a(a^2-b^2)}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

output `((-2*(2*a^2*A*b - A*b^3 - a^3*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + (A*(a^2 - b^2)*ArcTanh[Sin[c + d*x]])/(a*d)/(a*(a^2 - b^2)) + (b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))`

3.262.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.262.4 Maple [A] (verified)

Time = 1.57 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.37

3.262.
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx$$

method	result
derivativedivides	$\frac{-\frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^2} + \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^2}}{d} - \frac{2 \left(-\frac{ba(Ab - Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2 - b^2) \left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)} + \frac{(2Aa^2b - Ab^3 - B)}{(a - b)} \right)}{a^2}$
default	$\frac{-\frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^2} + \frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^2}}{d} - \frac{2 \left(-\frac{ba(Ab - Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2 - b^2) \left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)} + \frac{(2Aa^2b - Ab^3 - B)}{(a - b)} \right)}{a^2}$
risch	$\frac{2i(Ab - Ba)(ae^{i(dx+c)} + b)}{(-a^2 + b^2)da(b e^{2i(dx+c)} + 2a e^{i(dx+c)} + b)} - \frac{2b \ln\left(e^{i(dx+c)} - \frac{ia^2 - ib^2 - a\sqrt{-a^2 + b^2}}{b\sqrt{-a^2 + b^2}} \right) A}{\sqrt{-a^2 + b^2} (a+b)(a-b)d} + \frac{\ln\left(e^{i(dx+c)} - \frac{ia^2 - ib^2 - a\sqrt{-a^2 + b^2}}{b\sqrt{-a^2 + b^2}} \right)}{\sqrt{-a^2 + b^2} (a+b)(a-b)}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-A/a^2*ln(tan(1/2*d*x+1/2*c)-1)+A/a^2*ln(tan(1/2*d*x+1/2*c)+1)-2/a^2*(-b*a*(A*b-B*a)/(a^2-b^2)*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)+(2*A*a^2*b-A*b^3-B*a^3)/(a-b)/(a+b)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))))
```

3.262.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 308 vs. 2(123) = 246.

Time = 2.99 (sec) , antiderivative size = 684, normalized size of antiderivative = 5.14

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \left[\frac{(Ba^4 - 2Aa^3b + Aab^3 + (Ba^3b - 2Aa^2b^2 + Ab^4) \cos(dx + c))\sqrt{-a^2 + b^2} \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)}{b^2 \cos(dx+c)}\right)}{\dots} \right]$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^2,x, algorithm="fracas")
```



```
output [1/2*((B*a^4 - 2*A*a^3*b + A*a*b^3 + (B*a^3*b - 2*A*a^2*b^2 + A*b^4)*cos(d
*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x
+ c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^
2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + (A*a^5 - 2*A*a^3*b^2
+ A*a*b^4 + (A*a^4*b - 2*A*a^2*b^3 + A*b^5)*cos(d*x + c))*log(sin(d*x + c
) + 1) - (A*a^5 - 2*A*a^3*b^2 + A*a*b^4 + (A*a^4*b - 2*A*a^2*b^3 + A*b^5)*
cos(d*x + c))*log(-sin(d*x + c) + 1) - 2*(B*a^4*b - A*a^3*b^2 - B*a^2*b^3
+ A*a*b^4)*sin(d*x + c))/((a^6*b - 2*a^4*b^3 + a^2*b^5)*d*cos(d*x + c) + (
a^7 - 2*a^5*b^2 + a^3*b^4)*d), 1/2*(2*(B*a^4 - 2*A*a^3*b + A*a*b^3 + (B*a^
3*b - 2*A*a^2*b^2 + A*b^4)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*
x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (A*a^5 - 2*A*a^3*b^2 + A*a*b
^4 + (A*a^4*b - 2*A*a^2*b^3 + A*b^5)*cos(d*x + c))*log(sin(d*x + c) + 1) -
(A*a^5 - 2*A*a^3*b^2 + A*a*b^4 + (A*a^4*b - 2*A*a^2*b^3 + A*b^5)*cos(d*x
+ c))*log(-sin(d*x + c) + 1) - 2*(B*a^4*b - A*a^3*b^2 - B*a^2*b^3 + A*a*b^
4)*sin(d*x + c))/((a^6*b - 2*a^4*b^3 + a^2*b^5)*d*cos(d*x + c) + (a^7 - 2*
a^5*b^2 + a^3*b^4)*d)]
```

3.262.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**2,x)
```

```
output Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a + b*cos(c + d*x))**2, x)
```

3.262.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^2,x, algorithm="max
ima")
```

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.262.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.68

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2(Ba^3 - 2Aa^2b + Ab^3) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan\left(\frac{a \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^4 - a^2b^2)\sqrt{a^2 - b^2}} + \frac{A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^2} - \frac{A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `(2*(B*a^3 - 2*A*a^2*b + A*b^3)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^4 - a^2*b^2)*sqrt(a^2 - b^2)) + A*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^2 - A*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^2 - 2*(B*a*b*tan(1/2*d*x + 1/2*c) - A*b^2*tan(1/2*d*x + 1/2*c))/((a^3 - a*b^2)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b))/d`

3.262.9 Mupad [B] (verification not implemented)

Time = 8.39 (sec) , antiderivative size = 3763, normalized size of antiderivative = 28.29

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^2),x)`

output

$$\begin{aligned}
& - (A \operatorname{atan}\left(\frac{A \left(\frac{32(A^4 b^5 - B a^9 - A a^9 - 3A^6 b^3 + A^7 b^2 - B a^6 b^3 + B a^7 b^2 + 2A^8 b + B a^8 b)}{a^5 b + a^6 - a^3 b^3 - a^4 b^2} \right) - (32A \tan(c/2 + (d*x)/2) (2a^9 b - 2a^4 b^6 + 2a^5 b^5 + 4a^6 b^4 - 4a^7 b^3 - 2a^8 b^2))}{a^2 (a^4 b + a^5 - a^2 b^3 - a^3 b^2)}\right) \\
&) / a^2 - (32 \tan(c/2 + (d*x)/2) (A^2 a^6 + 2A^2 b^6 + B^2 a^6 - 2A^2 a b^5 - 2A^2 a^5 b - 5A^2 a^2 b^4 + 4A^2 a^3 b^3 + 3A^2 a^4 b^2 - 4A B a^5 b + 2A B a^3 b^3)) / (a^4 b + a^5 - a^2 b^3 - a^3 b^2) * i) / a^2 - (A \left(\frac{32(A^4 b^5 - B a^9 - A a^9 - 3A^6 b^3 + A^7 b^2 - B a^6 b^3 + B a^7 b^2 + 2A^8 b + B a^8 b)}{a^5 b + a^6 - a^3 b^3 - a^4 b^2} \right) + (32A \tan(c/2 + (d*x)/2) (2a^9 b - 2a^4 b^6 + 2a^5 b^5 + 4a^6 b^4 - 4a^7 b^3 - 2a^8 b^2)) / (a^2 (a^4 b + a^5 - a^2 b^3 - a^3 b^2))) / a^2 + (32 \tan(c/2 + (d*x)/2) (A^2 a^6 + 2A^2 b^6 + B^2 a^6 - 2A^2 a b^5 - 2A^2 a^5 b - 5A^2 a^2 b^4 + 4A^2 a^3 b^3 + 3A^2 a^4 b^2 - 4A B a^5 b + 2A B a^3 b^3)) / (a^4 b + a^5 - a^2 b^3 - a^3 b^2) * i) / a^2 / \left(\frac{A \left(\frac{32(A^4 b^5 - B a^9 - A a^9 - 3A^6 b^3 + A^7 b^2 - B a^6 b^3 + B a^7 b^2 + 2A^8 b + B a^8 b)}{a^5 b + a^6 - a^3 b^3 - a^4 b^2} \right) - (32A \tan(c/2 + (d*x)/2) (2a^9 b - 2a^4 b^6 + 2a^5 b^5 + 4a^6 b^4 - 4a^7 b^3 - 2a^8 b^2))}{a^2 (a^4 b + a^5 - a^2 b^3 - a^3 b^2)} \right) / a^2 - (32 \tan(c/2 + (d*x)/2) (A^2 a^6 + 2A^2 b^6 + B^2 a^6 - 2A^2 a b^5 - 2A^2 a^5 b - 5A^2 a^2 b^4 + 4A^2 a^3 b^3 + 3A^2 a^4 b^2 - 4A B a^5 b + 2A B a^3 b^3)) / (a^4 b + a^5 - a^2 b^3 - a^3 b^2) * i) / a^2 \dots
\end{aligned}$$

3.263
$$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.263.1 Optimal result 2421
 3.263.2 Mathematica [A] (verified) 2422
 3.263.3 Rubi [A] (verified) 2422
 3.263.4 Maple [A] (verified) 2426
 3.263.5 Fricas [B] (verification not implemented) 2427
 3.263.6 Sympy [F] 2428
 3.263.7 Maxima [F(-2)] 2428
 3.263.8 Giac [B] (verification not implemented) 2428
 3.263.9 Mupad [B] (verification not implemented) 2429

3.263.1 Optimal result

Integrand size = 31, antiderivative size = 189

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2b(3a^2Ab - 2Ab^3 - 2a^3B + ab^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^3(a-b)^{3/2}(a+b)^{3/2}d}$$

$$- \frac{(2Ab - aB) \operatorname{arctanh}(\sin(c + dx))}{a^3d}$$

$$+ \frac{(a^2A - 2Ab^2 + abB) \tan(c + dx)}{a^2(a^2 - b^2)d} + \frac{b(Ab - aB) \tan(c + dx)}{a(a^2 - b^2)d(a + b \cos(c + dx))}$$

```
output 2*b*(3*A*a^2*b-2*A*b^3-2*B*a^3+B*a*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2
*c)/(a+b)^(1/2))/a^3/(a-b)^(3/2)/(a+b)^(3/2)/d-(2*A*b-B*a)*arctanh(sin(d*x
+c))/a^3/d+(A*a^2-2*A*b^2+B*a*b)*tan(d*x+c)/a^2/(a^2-b^2)/d+b*(A*b-B*a)*ta
n(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.263.2 Mathematica [A] (verified)

Time = 1.70 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.27

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2b(-3a^2Ab + 2Ab^3 + 2a^3B - ab^2B) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right) + 2Ab \log\left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right) - aB}{(-a^2+b^2)^{3/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^2,x]`

output `((-2*b*(-3*a^2*A*b + 2*A*b^3 + 2*a^3*B - a*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(3/2) + 2*A*b*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - a*B*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - 2*A*b*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + a*B*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (a*b^2*(-(A*b) + a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x])) + a*A*Tan[c + d*x]/(a^3*d)`

3.263.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.10, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3479, 3042, 3534, 25, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^2 (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^2} dx$$

↓ 3479

$$\int \frac{(Aa^2 + bBa - (Ab - aB) \cos(c + dx)a - 2Ab^2 + b(Ab - aB) \cos^2(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx + \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

3.263. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{\int \frac{Aa^2 + bBa - (Ab - aB) \sin(c + dx + \frac{\pi}{2}) a - 2Ab^2 + b(Ab - aB) \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^2 (a + b \sin(c + dx + \frac{\pi}{2}))} dx}{a(a^2 - b^2)} + \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \downarrow \text{3534} \\
& \frac{\int - \frac{((a^2 - b^2)(2Ab - aB) - ab(Ab - aB) \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx}{a} + \frac{(a^2 A + abB - 2Ab^2) \tan(c + dx)}{ad} + \\
& \quad \frac{a(a^2 - b^2)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \downarrow \text{25} \\
& \frac{(a^2 A + abB - 2Ab^2) \tan(c + dx)}{ad} - \frac{\int \frac{((a^2 - b^2)(2Ab - aB) - ab(Ab - aB) \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx}{a} + \\
& \quad \frac{a(a^2 - b^2)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \downarrow \text{3042} \\
& \frac{(a^2 A + abB - 2Ab^2) \tan(c + dx)}{ad} - \frac{\int \frac{(a^2 - b^2)(2Ab - aB) - ab(Ab - aB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) (a + b \sin(c + dx + \frac{\pi}{2}))} dx}{a} + \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \downarrow \text{3480} \\
& \frac{(a^2 A + abB - 2Ab^2) \tan(c + dx)}{ad} - \frac{(a^2 - b^2)(2Ab - aB) \int \sec(c + dx) dx}{a} - \frac{b(-2a^3 B + 3a^2 Ab + ab^2 B - 2Ab^3) \int \frac{1}{a + b \cos(c + dx)} dx}{a} + \\
& \quad \frac{a(a^2 - b^2)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \downarrow \text{3042} \\
& \frac{(a^2 A + abB - 2Ab^2) \tan(c + dx)}{ad} - \frac{(a^2 - b^2)(2Ab - aB) \int \csc(c + dx + \frac{\pi}{2}) dx}{a} - \frac{b(-2a^3 B + 3a^2 Ab + ab^2 B - 2Ab^3) \int \frac{1}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{a} + \\
& \quad \frac{a(a^2 - b^2)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} \\
& \downarrow \text{3138}
\end{aligned}$$

3.263. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$

$$\frac{\frac{(a^2 A + abB - 2Ab^2) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(2Ab - aB) \int \csc(c+dx + \frac{\pi}{2}) dx}{a} - \frac{2b(-2a^3 B + 3a^2 Ab + ab^2 B - 2Ab^3) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{ad}}{a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 218

$$\frac{\frac{(a^2 A + abB - 2Ab^2) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(2Ab - aB) \int \csc(c+dx + \frac{\pi}{2}) dx}{a} - \frac{2b(-2a^3 B + 3a^2 Ab + ab^2 B - 2Ab^3) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{a(a^2 - b^2)} +$$

$$\frac{b(Ab - aB) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 4257

$$\frac{\frac{(a^2 A + abB - 2Ab^2) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(2Ab - aB) \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2b(-2a^3 B + 3a^2 Ab + ab^2 B - 2Ab^3) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{a(a^2 - b^2)} +$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^2,x]
```

```
output (b*(A*b - a*B)*Tan[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + (-((
(-2*b*(3*a^2*A*b - 2*A*b^3 - 2*a^3*B + a*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c
+ d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + ((a^2 - b^2)*(2*
A*b - a*B)*ArcTanh[Sin[c + d*x]])/(a*d))/a) + ((a^2*A - 2*A*b^2 + a*b*B)*T
an[c + d*x])/(a*d))/(a*(a^2 - b^2))
```

3.263.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`


```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.263.4 Maple [A] (verified)

Time = 2.11 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.28

method	result
derivativedivides	$2b \left(-\frac{ba(Ab - Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2 - b^2) \left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)} + \frac{(3A a^2 b - 2A b^3 - 2B a^3 + B a b^2) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}} \right) \frac{d}{a^3}$
default	$2b \left(-\frac{ba(Ab - Ba) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{(a^2 - b^2) \left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)} + \frac{(3A a^2 b - 2A b^3 - 2B a^3 + B a b^2) \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{(a-b)(a+b)\sqrt{(a-b)(a+b)}} \right) \frac{d}{a^3}$
risch	$\frac{2i(-A a b^2 e^{3i(dx+c)} + B a^2 b e^{3i(dx+c)} + A a^2 b e^{2i(dx+c)} - 2A b^3 e^{2i(dx+c)} + B a b^2 e^{2i(dx+c)} + 2A a^3 e^{i(dx+c)} - 3A a b^2 e^{i(dx+c)})}{d a^2 (e^{2i(dx+c)} + 1) (a^2 - b^2) (b e^{2i(dx+c)} + 2a e^{i(dx+c)} + b)}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBO
SE)
```

3.263. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$

output $1/d*(2*b/a^3*(-b*a*(A*b-B*a)/(a^2-b^2)*\tan(1/2*d*x+1/2*c)/(\tan(1/2*d*x+1/2*c)^2*a-b*\tan(1/2*d*x+1/2*c)^2+a+b)+(3*A*a^2*b-2*A*b^3-2*B*a^3+B*a*b^2)/(a-b)/(a+b)/((a-b)*(a+b))^{(1/2)*\arctan((a-b)*\tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^{(1/2))}-A/a^2/(\tan(1/2*d*x+1/2*c)-1)+(2*A*b-B*a)/a^3*\ln(\tan(1/2*d*x+1/2*c)-1)-A/a^2/(\tan(1/2*d*x+1/2*c)+1)+1/a^3*(-2*A*b+B*a)*\ln(\tan(1/2*d*x+1/2*c)+1))$

3.263.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 510 vs. $2(180) = 360$.

Time = 8.04 (sec) , antiderivative size = 1088, normalized size of antiderivative = 5.76

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output $[-1/2*((2*B*a^3*b^2 - 3*A*a^2*b^3 - B*a*b^4 + 2*A*b^5)*\cos(d*x + c)^2 + (2*B*a^4*b - 3*A*a^3*b^2 - B*a^2*b^3 + 2*A*a*b^4)*\cos(d*x + c))*\sqrt{-a^2 + b^2}*\log((2*a*b*\cos(d*x + c) + (2*a^2 - b^2)*\cos(d*x + c)^2 - 2*\sqrt{-a^2 + b^2})*(a*\cos(d*x + c) + b)*\sin(d*x + c) - a^2 + 2*b^2)/(b^2*\cos(d*x + c)^2 + 2*a*b*\cos(d*x + c) + a^2)) - ((B*a^5*b - 2*A*a^4*b^2 - 2*B*a^3*b^3 + 4*A*a^2*b^4 + B*a*b^5 - 2*A*b^6)*\cos(d*x + c)^2 + (B*a^6 - 2*A*a^5*b - 2*B*a^4*b^2 + 4*A*a^3*b^3 + B*a^2*b^4 - 2*A*a*b^5)*\cos(d*x + c))*\log(\sin(d*x + c) + 1) + ((B*a^5*b - 2*A*a^4*b^2 - 2*B*a^3*b^3 + 4*A*a^2*b^4 + B*a*b^5 - 2*A*b^6)*\cos(d*x + c)^2 + (B*a^6 - 2*A*a^5*b - 2*B*a^4*b^2 + 4*A*a^3*b^3 + B*a^2*b^4 - 2*A*a*b^5)*\cos(d*x + c))*\log(-\sin(d*x + c) + 1) - 2*(A*a^6 - 2*A*a^4*b^2 + A*a^2*b^4 + (A*a^5*b + B*a^4*b^2 - 3*A*a^3*b^3 - B*a^2*b^4 + 2*A*a*b^5)*\cos(d*x + c))*\sin(d*x + c))/((a^7*b - 2*a^5*b^3 + a^3*b^5)*d*\cos(d*x + c)^2 + (a^8 - 2*a^6*b^2 + a^4*b^4)*d*\cos(d*x + c)), -1/2*(2*((2*B*a^3*b^2 - 3*A*a^2*b^3 - B*a*b^4 + 2*A*b^5)*\cos(d*x + c)^2 + (2*B*a^4*b - 3*A*a^3*b^2 - B*a^2*b^3 + 2*A*a*b^4)*\cos(d*x + c))*\sqrt{a^2 - b^2}*\arctan(-(a*\cos(d*x + c) + b)/(\sqrt{a^2 - b^2}*\sin(d*x + c))) - ((B*a^5*b - 2*A*a^4*b^2 - 2*B*a^3*b^3 + 4*A*a^2*b^4 + B*a*b^5 - 2*A*b^6)*\cos(d*x + c)^2 + (B*a^6 - 2*A*a^5*b - 2*B*a^4*b^2 + 4*A*a^3*b^3 + B*a^2*b^4 - 2*A*a*b^5)*\cos(d*x + c))*\log(\sin(d*x + c) + 1) + ((B*a^5*b - 2*A*a^4*b^2 - 2*B*a^3*b^3 + 4*A*a^2*b^4 + B*a*b^5 - 2*A*b^6)*\cos(d*x + c)^2 + (B*a^6 - 2*A*a^5*b ...$

3.263.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**2,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a + b*cos(c + d*x))**2, x)`

3.263.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.263.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 404 vs. $2(180) = 360$.

Time = 0.31 (sec) , antiderivative size = 404, normalized size of antiderivative = 2.14

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2(2Ba^3b - 3Aa^2b^2 - Bab^3 + 2Ab^4) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^5 - a^3b^2)\sqrt{a^2 - b^2}} - \frac{2(Aa^3 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - A}{\dots}$$

3.263. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `(2*(2*B*a^3*b - 3*A*a^2*b^2 - B*a*b^3 + 2*A*b^4)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^5 - a^3*b^2)*sqrt(a^2 - b^2)) - 2*(A*a^3*tan(1/2*d*x + 1/2*c)^3 - A*a^2*b*tan(1/2*d*x + 1/2*c)^3 - A*a*b^2*tan(1/2*d*x + 1/2*c)^3 - B*a*b^2*tan(1/2*d*x + 1/2*c)^3 + 2*A*b^3*tan(1/2*d*x + 1/2*c)^3 + A*a^3*tan(1/2*d*x + 1/2*c) + A*a^2*b*tan(1/2*d*x + 1/2*c) - A*a*b^2*tan(1/2*d*x + 1/2*c) + B*a*b^2*tan(1/2*d*x + 1/2*c) - 2*A*b^3*tan(1/2*d*x + 1/2*c))/((a*tan(1/2*d*x + 1/2*c)^4 - b*tan(1/2*d*x + 1/2*c)^4 + 2*b*tan(1/2*d*x + 1/2*c)^2 - a - b)*(a^4 - a^2*b^2)) + (B*a - 2*A*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^3 - (B*a - 2*A*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^3)/d`

3.263.9 Mupad [B] (verification not implemented)

Time = 13.25 (sec) , antiderivative size = 5464, normalized size of antiderivative = 28.91

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^2),x)`

output

```
(atan((((32*tan(c/2 + (d*x)/2)*(8*A^2*b^8 + B^2*a^8 - 8*A^2*a*b^7 - 2*B^2*a^7*b - 16*A^2*a^2*b^6 + 16*A^2*a^3*b^5 + 5*A^2*a^4*b^4 - 8*A^2*a^5*b^3 + 4*A^2*a^6*b^2 + 2*B^2*a^2*b^6 - 2*B^2*a^3*b^5 - 5*B^2*a^4*b^4 + 4*B^2*a^5*b^3 + 3*B^2*a^6*b^2 - 8*A*B*a*b^7 - 4*A*B*a^7*b + 8*A*B*a^2*b^6 + 18*A*B*a^3*b^5 - 16*A*B*a^4*b^4 - 8*A*B*a^5*b^3 + 8*A*B*a^6*b^2)))/(a^6*b + a^7 - a^4*b^3 - a^5*b^2) + (((32*(A*a^7*b^5 - 2*A*a^6*b^6 - B*a^12 + 5*A*a^8*b^4 - 3*A*a^9*b^3 - 3*A*a^10*b^2 + B*a^7*b^5 - 3*B*a^9*b^3 + B*a^10*b^2 + 2*A*a^11*b + 2*B*a^11*b)))/(a^8*b + a^9 - a^6*b^3 - a^7*b^2) + (32*tan(c/2 + (d*x)/2)*(2*A*b - B*a)*(2*a^11*b - 2*a^6*b^6 + 2*a^7*b^5 + 4*a^8*b^4 - 4*a^9*b^3 - 2*a^10*b^2)))/(a^3*(a^6*b + a^7 - a^4*b^3 - a^5*b^2)))*(2*A*b - B*a))/a^3)*(2*A*b - B*a)*1i)/a^3 + (((32*tan(c/2 + (d*x)/2)*(8*A^2*b^8 + B^2*a^8 - 8*A^2*a*b^7 - 2*B^2*a^7*b - 16*A^2*a^2*b^6 + 16*A^2*a^3*b^5 + 5*A^2*a^4*b^4 - 8*A^2*a^5*b^3 + 4*A^2*a^6*b^2 + 2*B^2*a^2*b^6 - 2*B^2*a^3*b^5 - 5*B^2*a^4*b^4 + 4*B^2*a^5*b^3 + 3*B^2*a^6*b^2 - 8*A*B*a*b^7 - 4*A*B*a^7*b + 8*A*B*a^2*b^6 + 18*A*B*a^3*b^5 - 16*A*B*a^4*b^4 - 8*A*B*a^5*b^3 + 8*A*B*a^6*b^2)))/(a^6*b + a^7 - a^4*b^3 - a^5*b^2) - (((32*(A*a^7*b^5 - 2*A*a^6*b^6 - B*a^12 + 5*A*a^8*b^4 - 3*A*a^9*b^3 - 3*A*a^10*b^2 + B*a^7*b^5 - 3*B*a^9*b^3 + B*a^10*b^2 + 2*A*a^11*b + 2*B*a^11*b)))/(a^8*b + a^9 - a^6*b^3 - a^7*b^2) - (32*tan(c/2 + (d*x)/2)*(2*A*b - B*a)*(2*a^11*b - 2*a^6*b^6 + 2*a^7*b^5 + 4*a^8*b^4 - 4*a^9*b^3 - 2*a^10*b^2)))/(a^3*(a^6*b + a^7 - a^4*b...
```

3.264 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$

3.264.1 Optimal result 2431
 3.264.2 Mathematica [A] (verified) 2432
 3.264.3 Rubi [A] (verified) 2433
 3.264.4 Maple [A] (verified) 2437
 3.264.5 Fricas [B] (verification not implemented) 2438
 3.264.6 Sympy [F] 2439
 3.264.7 Maxima [F(-2)] 2439
 3.264.8 Giac [A] (verification not implemented) 2439
 3.264.9 Mupad [B] (verification not implemented) 2440

3.264.1 Optimal result

Integrand size = 31, antiderivative size = 270

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= -\frac{2b^2(4a^2Ab - 3Ab^3 - 3a^3B + 2ab^2B) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{a^4(a-b)^{3/2}(a+b)^{3/2}d}$$

$$+ \frac{(a^2A + 6Ab^2 - 4abB) \operatorname{arctanh}(\sin(c + dx))}{2a^4d}$$

$$- \frac{(2a^2Ab - 3Ab^3 - a^3B + 2ab^2B) \tan(c + dx)}{a^3(a^2 - b^2)d}$$

$$+ \frac{(a^2A - 3Ab^2 + 2abB) \sec(c + dx) \tan(c + dx)}{2a^2(a^2 - b^2)d} + \frac{b(Ab - aB) \sec(c + dx) \tan(c + dx)}{a(a^2 - b^2)d(a + b \cos(c + dx))}$$

output

```
-2*b^2*(4*A*a^2*b-3*A*b^3-3*B*a^3+2*B*a*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^4/(a-b)^(3/2)/(a+b)^(3/2)/d+1/2*(A*a^2+6*A*b^2-4*B*a*b)*arctanh(sin(d*x+c))/a^4/d-(2*A*a^2*b-3*A*b^3-B*a^3+2*B*a*b^2)*tan(d*x+c)/a^3/(a^2-b^2)/d+1/2*(A*a^2-3*A*b^2+2*B*a*b)*sec(d*x+c)*tan(d*x+c)/a^2/(a^2-b^2)/d+b*(A*b-B*a)*sec(d*x+c)*tan(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.264.2 Mathematica [A] (verified)

Time = 6.90 (sec) , antiderivative size = 438, normalized size of antiderivative = 1.62

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= -\frac{2b^2(-4a^2Ab + 3Ab^3 + 3a^3B - 2ab^2B) \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{a^4(a^2-b^2)\sqrt{-a^2+b^2}d}$$

$$+ \frac{(-a^2A - 6Ab^2 + 4abB) \log\left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right)}{2a^4d}$$

$$+ \frac{(a^2A + 6Ab^2 - 4abB) \log\left(\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right)\right)}{2a^4d}$$

$$+ \frac{A}{4a^2d\left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right)^2} - \frac{A}{4a^2d\left(\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right)\right)^2}$$

$$+ \frac{-2Ab\sin\left(\frac{1}{2}(c+dx)\right) + aB\sin\left(\frac{1}{2}(c+dx)\right)}{a^3d\left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right)}$$

$$+ \frac{-2Ab\sin\left(\frac{1}{2}(c+dx)\right) + aB\sin\left(\frac{1}{2}(c+dx)\right)}{a^3d\left(\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right)\right)} + \frac{Ab^4\sin(c+dx) - ab^3B\sin(c+dx)}{a^3(a-b)(a+b)d(a+b\cos(c+dx))}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^2,x]`output `(-2*b^2*(-4*a^2*A*b + 3*A*b^3 + 3*a^3*B - 2*a*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(a^4*(a^2 - b^2)*Sqrt[-a^2 + b^2]*d) + ((-a^2*A) - 6*A*b^2 + 4*a*b*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]]/(2*a^4*d) + ((a^2*A + 6*A*b^2 - 4*a*b*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/(2*a^4*d) + A/(4*a^2*d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2) - A/(4*a^2*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2) + (-2*A*b*Sin[(c + d*x)/2] + a*B*Sin[(c + d*x)/2])/(a^3*d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])) + (-2*A*b*Sin[(c + d*x)/2] + a*B*Sin[(c + d*x)/2])/(a^3*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])) + (A*b^4*Sin[c + d*x] - a*b^3*B*Sin[c + d*x])/(a^3*(a - b)*(a + b)*d*(a + b*Cos[c + d*x]))`

3.264.3 Rubi [A] (verified)

Time = 1.74 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3479, 3042, 3534, 25, 3042, 3534, 25, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx$$

↓ 3479

$$\int \frac{(Aa^2+2bBa-(Ab-aB)\cos(c+dx)a-3Ab^2+2b(Ab-aB)\cos^2(c+dx))\sec^3(c+dx)}{a+b\cos(c+dx)} dx +$$

$$\frac{a(a^2-b^2)}{ad(a^2-b^2)(a+b\cos(c+dx))} \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3042

$$\int \frac{Aa^2+2bBa-(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-3Ab^2+2b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))} dx +$$

$$\frac{a(a^2-b^2)}{ad(a^2-b^2)(a+b\cos(c+dx))} \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3534

$$\int -\frac{(-b(Aa^2+2bBa-3Ab^2)\cos^2(c+dx)-a(Aa^2-2bBa+Ab^2)\cos(c+dx)+2(-Ba^3+2Aba^2+2b^2Ba-3Ab^3))\sec^2(c+dx)}{a+b\cos(c+dx)} dx + \frac{(a^2A+2abB-3Ab^2)\tan(c+dx)}{2ad}$$

$$\frac{a(a^2-b^2)}{ad(a^2-b^2)(a+b\cos(c+dx))} \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 25

$$\frac{(a^2A+2abB-3Ab^2)\tan(c+dx)\sec(c+dx)}{2ad} - \int \frac{(-b(Aa^2+2bBa-3Ab^2)\cos^2(c+dx)-a(Aa^2-2bBa+Ab^2)\cos(c+dx)+2(-Ba^3+2Aba^2+2b^2Ba-3Ab^3))\sec^2(c+dx)}{a+b\cos(c+dx)} dx$$

$$\frac{a(a^2-b^2)}{ad(a^2-b^2)(a+b\cos(c+dx))} \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{ad(a^2-b^2)(a+b\cos(c+dx))}$$

3.264. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+b\cos(c+dx))^2} dx$

↓ 3042

$$\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{-b(Aa^2 + 2bBa - 3Ab^2) \sin(c+dx + \frac{\pi}{2})^2 - a(Aa^2 - 2bBa + Ab^2) \sin(c+dx + \frac{\pi}{2}) + 2(-Ba^3 + 2Aba^2 + 2b^2Ba - 3Ab^3)}{\sin(c+dx + \frac{\pi}{2})^2 (a+b \sin(c+dx + \frac{\pi}{2}))} dx}{2a}$$

$$\frac{b(Ab - aB) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))}$$

↓ 3534

$$\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{((a^2 - b^2)(Aa^2 - 4bBa + 6Ab^2) + ab(Aa^2 + 2bBa - 3Ab^2) \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx}{a} + \frac{2(a^3(-B) + 2a^2Ab + 2ab^2B - 3Ab^3) \tan(c+dx)}{2a}$$

$$\frac{b(Ab - aB) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))}$$

↓ 25

$$\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2Ab + 2ab^2B - 3Ab^3) \tan(c+dx)}{ad} - \frac{\int \frac{((a^2 - b^2)(Aa^2 - 4bBa + 6Ab^2) + ab(Aa^2 + 2bBa - 3Ab^2) \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx}{2a}$$

$$\frac{b(Ab - aB) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))}$$

↓ 3042

$$\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2Ab + 2ab^2B - 3Ab^3) \tan(c+dx)}{ad} - \frac{\int \frac{(a^2 - b^2)(Aa^2 - 4bBa + 6Ab^2) + ab(Aa^2 + 2bBa - 3Ab^2) \cos(c+dx)}{\sin(c+dx + \frac{\pi}{2})(a+b \sin(c+dx + \frac{\pi}{2}))} \sin(c+dx) dx}{2a}$$

$$\frac{b(Ab - aB) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))}$$

↓ 3480

$$\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2Ab + 2ab^2B - 3Ab^3) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(a^2 A - 4abB + 6Ab^2) \int \sec(c+dx) dx}{a} - \frac{2b^2(-3a^3 B + 4a^2 ab + 3ab^2)}{a}$$

$$\frac{b(Ab - aB) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))}$$

↓ 3042

3.264. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$

$$\frac{\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2 Ab + 2ab^2 B - 3Ab^3) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(a^2 A - 4abB + 6Ab^2) \int \csc(c+dx + \frac{\pi}{2}) dx}{a} - \frac{2b^2(-3a^3)}{a}}{2a} = \frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} a(a^2 - b^2)$$

3138

$$\frac{\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2 Ab + 2ab^2 B - 3Ab^3) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(a^2 A - 4abB + 6Ab^2) \int \csc(c+dx + \frac{\pi}{2}) dx}{a} - \frac{4b^2(-3a^3)}{a}}{2a} = \frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} a(a^2 - b^2)$$

218

$$\frac{\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2 Ab + 2ab^2 B - 3Ab^3) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(a^2 A - 4abB + 6Ab^2) \int \csc(c+dx + \frac{\pi}{2}) dx}{a} - \frac{4b^2(-3a^3)}{a}}{2a} = \frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} a(a^2 - b^2)$$

4257

$$\frac{\frac{(a^2 A + 2abB - 3Ab^2) \tan(c+dx) \sec(c+dx)}{2ad} - \frac{2(a^3(-B) + 2a^2 Ab + 2ab^2 B - 3Ab^3) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)(a^2 A - 4abB + 6Ab^2) \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{4b^2(-3a^3)}{a}}{2a} + \frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^2,x]`

output `(b*(A*b - a*B)*Sec[c + d*x]*Tan[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + (((a^2*A - 3*A*b^2 + 2*a*b*B)*Sec[c + d*x]*Tan[c + d*x])/(2*a*d) - (((-4*b^2*(4*a^2*A*b - 3*A*b^3 - 3*a^3*B + 2*a*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + ((a^2 - b^2)*(a^2*A + 6*A*b^2 - 4*a*b*B)*ArcTanh[Sin[c + d*x]]/(a*d))/a) + (2*(2*a^2*A*b - 3*A*b^3 - a^3*B + 2*a*b^2*B)*Tan[c + d*x])/(a*d))/(2*a))/(a*(a^2 - b^2))`

3.264. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$

3.264.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`
- rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.264.4 Maple [A] (verified)

Time = 2.06 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.21

method	result
derivativedivides	$-\frac{A}{2a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{-aA - 4Ab + 2Ba}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(Aa^2 + 6Ab^2 - 4Bab) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a^4} + \frac{A}{2a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{-aA - 4Ab + 2Ba}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$
default	$-\frac{A}{2a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{-aA - 4Ab + 2Ba}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(Aa^2 + 6Ab^2 - 4Bab) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a^4} + \frac{A}{2a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{-aA - 4Ab + 2Ba}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$
risch	Expression too large to display

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBO
SE)
```

$$3.264. \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$$

```
output 1/d*(-1/2*A/a^2/(tan(1/2*d*x+1/2*c)+1)^2-1/2*(-A*a-4*A*b+2*B*a)/a^3/(tan(1/2*d*x+1/2*c)+1)+1/2*(A*a^2+6*A*b^2-4*B*a*b)/a^4*ln(tan(1/2*d*x+1/2*c)+1)+1/2*A/a^2/(tan(1/2*d*x+1/2*c)-1)^2-1/2*(-A*a-4*A*b+2*B*a)/a^3/(tan(1/2*d*x+1/2*c)-1)+1/2/a^4*(-A*a^2-6*A*b^2+4*B*a*b)*ln(tan(1/2*d*x+1/2*c)-1)-2*b^2/a^4*(-b*a*(A*b-B*a)/(a^2-b^2)*tan(1/2*d*x+1/2*c)/(tan(1/2*d*x+1/2*c)^2-a-b*tan(1/2*d*x+1/2*c)^2+a+b)+(4*A*a^2*b-3*A*b^3-3*B*a^3+2*B*a*b^2)/(a-b)/(a+b)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))))
```

3.264.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 630 vs. $2(256) = 512$.

Time = 12.98 (sec) , antiderivative size = 1329, normalized size of antiderivative = 4.92

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^2,x, algorithm="fracas")
```

```
output [-1/4*(2*((3*B*a^3*b^3 - 4*A*a^2*b^4 - 2*B*a*b^5 + 3*A*b^6)*cos(d*x + c)^3 + (3*B*a^4*b^2 - 4*A*a^3*b^3 - 2*B*a^2*b^4 + 3*A*a*b^5)*cos(d*x + c)^2)*sqrt(-a^2 + b^2)*log(((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - ((A*a^6*b - 4*B*a^5*b^2 + 4*A*a^4*b^3 + 8*B*a^3*b^4 - 11*A*a^2*b^5 - 4*B*a*b^6 + 6*A*b^7)*cos(d*x + c)^3 + (A*a^7 - 4*B*a^6*b + 4*A*a^5*b^2 + 8*B*a^4*b^3 - 11*A*a^3*b^4 - 4*B*a^2*b^5 + 6*A*a*b^6)*cos(d*x + c)^2)*log(sin(d*x + c) + 1) + ((A*a^6*b - 4*B*a^5*b^2 + 4*A*a^4*b^3 + 8*B*a^3*b^4 - 11*A*a^2*b^5 - 4*B*a*b^6 + 6*A*b^7)*cos(d*x + c)^3 + (A*a^7 - 4*B*a^6*b + 4*A*a^5*b^2 + 8*B*a^4*b^3 - 11*A*a^3*b^4 - 4*B*a^2*b^5 + 6*A*a*b^6)*cos(d*x + c)^2)*log(-sin(d*x + c) + 1) - 2*(A*a^7 - 2*A*a^5*b^2 + A*a^3*b^4 + 2*(B*a^6*b - 2*A*a^5*b^2 - 3*B*a^4*b^3 + 5*A*a^3*b^4 + 2*B*a^2*b^5 - 3*A*a*b^6)*cos(d*x + c)^2 + (2*B*a^7 - 3*A*a^6*b - 4*B*a^5*b^2 + 6*A*a^4*b^3 + 2*B*a^3*b^4 - 3*A*a^2*b^5)*cos(d*x + c))*sin(d*x + c))/((a^8*b - 2*a^6*b^3 + a^4*b^5)*d*cos(d*x + c)^3 + (a^9 - 2*a^7*b^2 + a^5*b^4)*d*cos(d*x + c)^2), 1/4*(4*((3*B*a^3*b^3 - 4*A*a^2*b^4 - 2*B*a*b^5 + 3*A*b^6)*cos(d*x + c)^3 + (3*B*a^4*b^2 - 4*A*a^3*b^3 - 2*B*a^2*b^4 + 3*A*a*b^5)*cos(d*x + c)^2)*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + ((A*a^6*b - 4*B*a^5*b^2 + 4*A*a^4*b^3 + 8*B*a^3*b^4 - 11*A*a^2*b^5 - 4*B*a*b^6 + 6*A*b^7)*cos(d*x + c)^...
```

3.264.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**2,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a + b*cos(c + d*x))**2, x)`

3.264.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.264.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.40

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{4(3Ba^3b^2 - 4Aa^2b^3 - 2Bab^4 + 3Ab^5) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^6 - a^4b^2)\sqrt{a^2 - b^2}} + \frac{4(Bab^3 \tan(\frac{1}{2} c))}{(a^5 - a^3b^2)(a \tan(\frac{1}{2} c))}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `-1/2*(4*(3*B*a^3*b^2 - 4*A*a^2*b^3 - 2*B*a*b^4 + 3*A*b^5)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^6 - a^4*b^2)*sqrt(a^2 - b^2)) + 4*(B*a*b^3*tan(1/2*d*x + 1/2*c) - A*b^4*tan(1/2*d*x + 1/2*c))/((a^5 - a^3*b^2)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)) - (A*a^2 - 4*B*a*b + 6*A*b^2)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^4 + (A*a^2 - 4*B*a*b + 6*A*b^2)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^4 - 2*(A*a*tan(1/2*d*x + 1/2*c)^3 - 2*B*a*tan(1/2*d*x + 1/2*c)^3 + 4*A*b*tan(1/2*d*x + 1/2*c)^3 + A*a*tan(1/2*d*x + 1/2*c) + 2*B*a*tan(1/2*d*x + 1/2*c) - 4*A*b*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 - 1)^2*a^3)/d`

3.264.9 Mupad [B] (verification not implemented)

Time = 9.65 (sec) , antiderivative size = 6692, normalized size of antiderivative = 24.79

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^2),x)`

output

$$\begin{aligned} & \left(\operatorname{atan}\left(-\left(\left(\left(8 \tan\left(\frac{c}{2} + \frac{d \cdot x}{2}\right) \cdot \left(A^2 a^{10} + 72 A^2 b^{10} - 72 A^2 a b^9 - 2\right.\right.\right.\right.\right.\right. \right. \\ & * A^2 a^9 b - 120 A^2 a^2 b^8 + 120 A^2 a^3 b^7 + 17 A^2 a^4 b^6 - 26 A^2 a \\ & ^5 b^5 + 23 A^2 a^6 b^4 - 20 A^2 a^7 b^3 + 11 A^2 a^8 b^2 + 32 B^2 a^2 b^8 \\ & - 32 B^2 a^3 b^7 - 64 B^2 a^4 b^6 + 64 B^2 a^5 b^5 + 20 B^2 a^6 b^4 - 32 * \\ & B^2 a^7 b^3 + 16 B^2 a^8 b^2 - 96 A B a b^9 - 8 A B a^9 b + 96 A B a^2 b^8 \\ & + 176 A B a^3 b^7 - 176 A B a^4 b^6 - 40 A B a^5 b^5 + 64 A B a^6 b^4 - 4 \\ & 0 A B a^7 b^3 + 16 A B a^8 b^2\right)\right) / \left(a^8 b + a^9 - a^6 b^3 - a^7 b^2\right) - \left(\left(\left(8 * \right.\right.\right. \\ & \left.\left.\left(2 A a^{15} - 12 A a^8 b^7 + 6 A a^9 b^6 + 28 A a^{10} b^5 - 14 A a^{11} b^4 - 1\right.\right.\right. \\ & \left.\left.\left.6 A a^{12} b^3 + 6 A a^{13} b^2 + 8 B a^9 b^6 - 4 B a^{10} b^5 - 20 B a^{11} b^4 + \right.\right.\right. \\ & \left.\left.\left.12 B a^{12} b^3 + 12 B a^{13} b^2 - 8 B a^{14} b\right)\right)\right) / \left(a^{11} b + a^{12} - a^9 b^3 - a \right. \\ & \left.^{10} b^2\right) - \left(4 \tan\left(\frac{c}{2} + \frac{d \cdot x}{2}\right) \cdot \left(A a^2 + 6 A b^2 - 4 B a b\right) \cdot \left(8 a^{13} b - 8 \right.\right. \\ & \left.\left. a^8 b^6 + 8 a^9 b^5 + 16 a^{10} b^4 - 16 a^{11} b^3 - 8 a^{12} b^2\right)\right) / \left(a^4 \cdot \left(a^8 * \right.\right. \\ & \left.\left. b + a^9 - a^6 b^3 - a^7 b^2\right)\right) \cdot \left(A a^2 + 6 A b^2 - 4 B a b\right) / \left(2 a^4\right) \cdot \left(A a^2 \right. \\ & \left.+ 6 A b^2 - 4 B a b\right) \cdot i) / \left(2 a^4\right) + \left(\left(\left(8 \tan\left(\frac{c}{2} + \frac{d \cdot x}{2}\right) \cdot \left(A^2 a^{10} + 7\right.\right.\right.\right. \right. \\ & \left.\left.\left.2 A^2 b^{10} - 72 A^2 a b^9 - 2 A^2 a^9 b - 120 A^2 a^2 b^8 + 120 A^2 a^3 b^7 \right.\right.\right. \\ & \left.\left.\left.7 + 17 A^2 a^4 b^6 - 26 A^2 a^5 b^5 + 23 A^2 a^6 b^4 - 20 A^2 a^7 b^3 + 11 \right.\right.\right. \\ & \left.\left.\left. A^2 a^8 b^2 + 32 B^2 a^2 b^8 - 32 B^2 a^3 b^7 - 64 B^2 a^4 b^6 + 64 B^2 a \right.\right.\right. \\ & \left.\left.\left. ^5 b^5 + 20 B^2 a^6 b^4 - 32 B^2 a^7 b^3 + 16 B^2 a^8 b^2 - 96 A B a b^9 - \right.\right.\right. \\ & \left.\left.\left. 8 A B a^9 b + 96 A B a^2 b^8 + 176 A B a^3 b^7 - 176 A B a^4 b^6 - 40 A B \right.\right.\right. \\ & \left.\left.\left. a^5 b^5 + 64 A B a^6 b^4 - 40 A B a^7 b^3 + 16 A B a^8 b^2\right)\right)\right) / \left(a^8 b + \dots \right. \end{aligned}$$

3.265 $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

3.265.1 Optimal result 2442
 3.265.2 Mathematica [A] (verified) 2443
 3.265.3 Rubi [A] (verified) 2443
 3.265.4 Maple [A] (verified) 2449
 3.265.5 Fricas [B] (verification not implemented) 2450
 3.265.6 Sympy [F(-1)] 2450
 3.265.7 Maxima [F(-2)] 2451
 3.265.8 Giac [B] (verification not implemented) 2451
 3.265.9 Mupad [B] (verification not implemented) 2452

3.265.1 Optimal result

Integrand size = 31, antiderivative size = 398

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx = -\frac{(6aAb - 12a^2B - b^2B)x}{2b^5} + \frac{a^2(6a^4Ab - 15a^2Ab^3 + 12Ab^5 - 12a^5B + 29a^3b^2B - 20ab^4B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{5/2}b^5(a+b)^{5/2}d} + \frac{(6a^4Ab - 11a^2Ab^3 + 2Ab^5 - 12a^5B + 21a^3b^2B - 6ab^4B) \sin(c+dx)}{2b^4(a^2-b^2)^2d} - \frac{(3a^3Ab - 6aAb^3 - 6a^4B + 10a^2b^2B - b^4B) \cos(c+dx) \sin(c+dx)}{2b^3(a^2-b^2)^2d} + \frac{a(Ab - aB) \cos^3(c+dx) \sin(c+dx)}{2b(a^2-b^2)d(a+b \cos(c+dx))^2} + \frac{a(2a^2Ab - 5Ab^3 - 4a^3B + 7ab^2B) \cos^2(c+dx) \sin(c+dx)}{2b^2(a^2-b^2)^2d(a+b \cos(c+dx))}$$

output

```
-1/2*(6*A*a*b-12*B*a^2-B*b^2)*x/b^5+a^2*(6*A*a^4*b-15*A*a^2*b^3+12*A*b^5-12*B*a^5+29*B*a^3*b^2-20*B*a*b^4)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(5/2)/b^5/(a+b)^(5/2)/d+1/2*(6*A*a^4*b-11*A*a^2*b^3+2*A*b^5-12*B*a^5+21*B*a^3*b^2-6*B*a*b^4)*sin(d*x+c)/b^4/(a^2-b^2)^2/d-1/2*(3*A*a^3*b-6*A*a*b^3-6*B*a^4+10*B*a^2*b^2-B*b^4)*cos(d*x+c)*sin(d*x+c)/b^3/(a^2-b^2)^2/d+1/2*a*(A*b-B*a)*cos(d*x+c)^3*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/2*a*(2*A*a^2*b-5*A*b^3-4*B*a^3+7*B*a*b^2)*cos(d*x+c)^2*sin(d*x+c)/b^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

3.265.2 Mathematica [A] (verified)

Time = 3.66 (sec) , antiderivative size = 734, normalized size of antiderivative = 1.84

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \frac{16a^2(-6a^4Ab+15a^2Ab^3-12Ab^5+12a^5B-29a^3b^2B+20ab^4B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + \frac{-48a^7Abc+72a^5Ab^3c-24aAb^7c+96a^8Bc-136a^6b^2Bc-12a^4b^4Bc+48a^2b^6Bc+4b^8Bc-48a^7A*b*d*x+72a^5A*b^3*d*x-24aA*b^7*d*x+96a^8B*d*x-136a^6b^2B*d*x-12a^4b^4B*d*x+48a^2b^6B*d*x+4b^8B*d*x+16a*b*(a^2-b^2)^2*(-6aA*b+12a^2B+b^2B)*(c+dx)*\cos[c+dx]+4*(-a^2*b+b^3)^2*(-6aA*b+12a^2B+b^2B)*(c+dx)*\cos[2*(c+dx)]+48a^6A*b^2*\sin[c+dx]-84a^4A*b^4*\sin[c+dx]+8a^2A*b^6*\sin[c+dx]+4A*b^8*\sin[c+dx]-96a^7*b*B*\sin[c+dx]+160a^5*b^3*B*\sin[c+dx]-32a^3*b^5*B*\sin[c+dx]-8a*b^7*B*\sin[c+dx]+36a^5A*b^3*\sin[2*(c+dx)]-64a^3A*b^5*\sin[2*(c+dx)]+16aA*b^7*\sin[2*(c+dx)]-72a^6b^2*B*\sin[2*(c+dx)]+130a^4b^4*B*\sin[2*(c+dx)]-48a^2b^6*B*\sin[2*(c+dx)]+2b^8*B*\sin[2*(c+dx)]+4a^4A*b^4*\sin[3*(c+dx)]-8a^2A*b^6*\sin[3*(c+dx)]+4A*b^8*\sin[3*(c+dx)]-8a^5*b^3*B*\sin[3*(c+dx)]+16a^3*b^5*B*\sin[3*(c+dx)]-8a*b^7*B*\sin[3*(c+dx)]+a^4*b^4*B*\sin[4*(c+dx)]-2a^2*b^6*B*\sin[4*(c+dx)]+b^8*B*\sin[4*(c+dx)]}{(a+b\cos(c+dx))^3}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output `((16*a^2*(-6*a^4*A*b + 15*a^2*A*b^3 - 12*A*b^5 + 12*a^5*B - 29*a^3*b^2*B + 20*a*b^4*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(5/2) + (-48*a^7*A*b*c + 72*a^5*A*b^3*c - 24*a*A*b^7*c + 96*a^8*B*c - 136*a^6*b^2*B*c - 12*a^4*b^4*B*c + 48*a^2*b^6*B*c + 4*b^8*B*c - 48*a^7*A*b*d*x + 72*a^5*A*b^3*d*x - 24*a*A*b^7*d*x + 96*a^8*B*d*x - 136*a^6*b^2*B*d*x - 12*a^4*b^4*B*d*x + 48*a^2*b^6*B*d*x + 4*b^8*B*d*x + 16*a*b*(a^2 - b^2)^2*(-6*a*A*b + 12*a^2*B + b^2*B)*(c + d*x)*Cos[c + d*x] + 4*(-a^2*b + b^3)^2*(-6*a*A*b + 12*a^2*B + b^2*B)*(c + d*x)*Cos[2*(c + d*x)] + 48*a^6*A*b^2*Sin[c + d*x] - 84*a^4*A*b^4*Sin[c + d*x] + 8*a^2*A*b^6*Sin[c + d*x] + 4*A*b^8*Sin[c + d*x] - 96*a^7*b*B*Sin[c + d*x] + 160*a^5*b^3*B*Sin[c + d*x] - 32*a^3*b^5*B*Sin[c + d*x] - 8*a*b^7*B*Sin[c + d*x] + 36*a^5*A*b^3*Sin[2*(c + d*x)] - 64*a^3*A*b^5*Sin[2*(c + d*x)] + 16*a*A*b^7*Sin[2*(c + d*x)] - 72*a^6*b^2*B*Sin[2*(c + d*x)] + 130*a^4*b^4*B*Sin[2*(c + d*x)] - 48*a^2*b^6*B*Sin[2*(c + d*x)] + 2*b^8*B*Sin[2*(c + d*x)] + 4*a^4*A*b^4*Sin[3*(c + d*x)] - 8*a^2*A*b^6*Sin[3*(c + d*x)] + 4*A*b^8*Sin[3*(c + d*x)] - 8*a^5*b^3*B*Sin[3*(c + d*x)] + 16*a^3*b^5*B*Sin[3*(c + d*x)] - 8*a*b^7*B*Sin[3*(c + d*x)] + a^4*b^4*B*Sin[4*(c + d*x)] - 2*a^2*b^6*B*Sin[4*(c + d*x)] + b^8*B*Sin[4*(c + d*x)]/(a^2 - b^2)^2*(a + b*Cos[c + d*x])^2)/(16*b^5*d)`

3.265.3 Rubi [A] (verified)

Time = 2.13 (sec) , antiderivative size = 424, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3468, 25, 3042, 3526, 25, 3042, 3528, 27, 3042, 3502, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.265. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^4(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx \\
& \quad \downarrow \text{3468} \\
& \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} - \\
& \frac{\int -\frac{\cos^2(c+dx)(-2(-2Ba^2+Abab^2B)\cos^2(c+dx)-2b(Ab-aB)\cos(c+dx)+3a(Ab-aB))}{(a+b\cos(c+dx))^2} dx}{2b(a^2-b^2)} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{\cos^2(c+dx)(-2(-2Ba^2+Abab^2B)\cos^2(c+dx)-2b(Ab-aB)\cos(c+dx)+3a(Ab-aB))}{(a+b\cos(c+dx))^2} dx}{2b(a^2-b^2)} + \\
& \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2(-2(-2Ba^2+Abab^2B)\sin(c+dx+\frac{\pi}{2})^2-2b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+3a(Ab-aB))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx}{2b(a^2-b^2)} + \\
& \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \downarrow \text{3526} \\
& \frac{a(-4a^3B+2a^2Ab+7ab^2B-5Ab^3)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))} - \frac{\int -\frac{\cos(c+dx)(-2(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3a-b^4B)\cos^2(c+dx)+b(Ba^3+Abab^2-4b^2Ba+2Ab^3)\cos(c+dx)+2a(-4Ba^3+2Abab^2+7b^2Ba-5Ab^3))}{a+b\cos(c+dx)} dx}{b(a^2-b^2)} + a \\
& \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{\cos(c+dx)(-2(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3a-b^4B)\cos^2(c+dx)+b(Ba^3+Abab^2-4b^2Ba+2Ab^3)\cos(c+dx)+2a(-4Ba^3+2Abab^2+7b^2Ba-5Ab^3))}{a+b\cos(c+dx)} dx}{b(a^2-b^2)} + a \\
& \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.265. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)\left(-2\left(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3a-b^4B\right)\sin\left(c+dx+\frac{\pi}{2}\right)^2+b\left(Ba^3+Aba^2-4b^2Ba+2Ab^3\right)\sin\left(c+dx+\frac{\pi}{2}\right)+2a\left(-4Ba^3+2Aba^2+7b^2Ba-5Ab^3\right)}{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} dx$$

$$\frac{2b(a^2-b^2)}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3528

$$\int \frac{2\left(-\left(-12Ba^5+6Aba^4+21b^2Ba^3-11Ab^3a^2-6b^4Ba+2Ab^5\right)\cos^2(c+dx)\right)-b\left(-2Ba^4+Aba^3+4b^2Ba^2-4Ab^3a+b^4B\right)\cos(c+dx)+a\left(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3\right)}{a+b\cos(c+dx)} dx$$

$$\frac{2b(a^2-b^2)}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 27

$$\int \frac{-\left(-12Ba^5+6Aba^4+21b^2Ba^3-11Ab^3a^2-6b^4Ba+2Ab^5\right)\cos^2(c+dx)-b\left(-2Ba^4+Aba^3+4b^2Ba^2-4Ab^3a+b^4B\right)\cos(c+dx)+a\left(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3\right)}{a+b\cos(c+dx)} dx$$

$$\frac{2b(a^2-b^2)}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3042

$$\int \frac{\left(12Ba^5-6Aba^4-21b^2Ba^3+11Ab^3a^2+6b^4Ba-2Ab^5\right)\sin\left(c+dx+\frac{\pi}{2}\right)^2-b\left(-2Ba^4+Aba^3+4b^2Ba^2-4Ab^3a+b^4B\right)\sin\left(c+dx+\frac{\pi}{2}\right)+a\left(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3\right)}{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} dx$$

$$\frac{2b(a^2-b^2)}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3502

$$\int \frac{\left(-12Ba^2+6Aba-b^2B\right)\cos(c+dx)\left(a^2-b^2\right)^2+ab\left(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3a-b^4B\right)}{a+b\cos(c+dx)} dx - \frac{\left(-12a^5B+6a^4Ab+21a^3b^2B-11a^2Ab^3-6ab^4B+2Ab^5\right)\sin(c+dx)}{bd}$$

$$\frac{2b(a^2-b^2)}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3042

3.265. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

$$\frac{\int \frac{(-12Ba^2+6Aba-b^2B) \sin(c+dx+\frac{\pi}{2})(a^2-b^2)^2+ab(-6Ba^4+3Aba^3+10b^2Ba^2-6Ab^3a-b^4B)}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{\frac{\frac{\frac{(-12a^5B+6a^4Ab+21a^3b^2B-11a^2Ab^3-6ab^4B+2Ab^5)}{bd}}{b(a^2-b^2)}}{b}} = 2b(a^2-b^2)$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3214

$$\frac{\frac{x(a^2-b^2)^2(-12a^2B+6aAb-b^2B)}{b} - \frac{a^2(-12a^5B+6a^4Ab+29a^3b^2B-15a^2Ab^3-20ab^4B+12Ab^5)}{b} \int \frac{1}{a+b \cos(c+dx)} dx}{\frac{\frac{(-12a^5B+6a^4Ab+21a^3b^2B-11a^2Ab^3-6ab^4B+2Ab^5)}{bd}}{b(a^2-b^2)}} = 2b(a^2-b^2)$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3042

$$\frac{\frac{x(a^2-b^2)^2(-12a^2B+6aAb-b^2B)}{b} - \frac{a^2(-12a^5B+6a^4Ab+29a^3b^2B-15a^2Ab^3-20ab^4B+12Ab^5)}{b} \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{\frac{(-12a^5B+6a^4Ab+21a^3b^2B-11a^2Ab^3-6ab^4B+2Ab^5)}{bd}} = 2b(a^2-b^2)$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3138

$$\frac{\frac{x(a^2-b^2)^2(-12a^2B+6aAb-b^2B)}{b} - \frac{2a^2(-12a^5B+6a^4Ab+29a^3b^2B-15a^2Ab^3-20ab^4B+12Ab^5)}{bd} \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx))+a+b} d \tan(\frac{1}{2}(c+dx))}{\frac{(-12a^5B+6a^4Ab+21a^3b^2B-11a^2Ab^3-6ab^4B+2Ab^5)}{bd}}}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 218

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{2bd(a^2-b^2)(a+b \cos(c+dx))^2} +$$

$$\frac{a(-4a^3B+2a^2Ab+7ab^2B-5Ab^3) \sin(c+dx) \cos^2(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} + \frac{(-6a^4B+3a^3Ab+10a^2b^2B-6aAb^3-b^4B) \sin(c+dx) \cos(c+dx)}{bd} - \frac{x(a^2-b^2)^2(-12a^2B+6aAb-b^2B)}{b}$$

3.265. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output `(a*(A*b - a*B)*Cos[c + d*x]^3*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((a*(2*a^2*A*b - 5*A*b^3 - 4*a^3*B + 7*a*b^2*B)*Cos[c + d*x]^2*Sin[c + d*x])/(b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + (-(((3*a^3*A*b - 6*a*A*b^3 - 6*a^4*B + 10*a^2*b^2*B - b^4*B)*Cos[c + d*x]*Sin[c + d*x])/(b*d)) - (((a^2 - b^2)^2*(6*a*A*b - 12*a^2*B - b^2*B)*x)/b - (2*a^2*(6*a^4*A*b - 15*a^2*A*b^3 + 12*A*b^5 - 12*a^5*B + 29*a^3*b^2*B - 20*a*b^4*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d))/b - ((6*a^4*A*b - 11*a^2*A*b^3 + 2*A*b^5 - 12*a^5*B + 21*a^3*b^2*B - 6*a*b^4*B)*Sin[c + d*x])/(b*d))/b)/(b*(a^2 - b^2)))/(2*b*(a^2 - b^2))`

3.265.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sine[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x
])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

3.265.4 Maple [A] (verified)

Time = 1.93 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.01

method	result
derivativedivides	$2a^2 \left(\frac{(4Aa^2b - Aab^2 - 8Ab^3 - 6Ba^3 + Ba^2b + 10Bab^2)ab \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) + ba(4Aa^2b + Aab^2 - 8Ab^3 - 6Ba^3 - Ba^2b + 10Bab^2) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)} + \frac{ba(4Aa^2b + Aab^2 - 8Ab^3 - 6Ba^3 - Ba^2b + 10Bab^2) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{1}{\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^{a-b} \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^{a+b}} \frac{1}{b^5}$
default	$2a^2 \left(\frac{(4Aa^2b - Aab^2 - 8Ab^3 - 6Ba^3 + Ba^2b + 10Bab^2)ab \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) + ba(4Aa^2b + Aab^2 - 8Ab^3 - 6Ba^3 - Ba^2b + 10Bab^2) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)} + \frac{ba(4Aa^2b + Aab^2 - 8Ab^3 - 6Ba^3 - Ba^2b + 10Bab^2) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{1}{b^5}$
risch	Expression too large to display

```
input int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBO
SE)
```

```
output 1/d*(2*a^2/b^5*((1/2*(4*A*a^2*b-A*a*b^2-8*A*b^3-6*B*a^3+B*a^2*b+10*B*a*b^2
)*a*b/(a-b)/(a^2+2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3+1/2*b*a*(4*A*a^2*b+A*a*b^
2-8*A*b^3-6*B*a^3-B*a^2*b+10*B*a*b^2)/(a+b)/(a-b)^2*tan(1/2*d*x+1/2*c))/(t
an(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^2+1/2*(6*A*a^4*b-15*A*a^
2*b^3+12*A*b^5-12*B*a^5+29*B*a^3*b^2-20*B*a*b^4)/(a^4-2*a^2*b^2+b^4)/((a-b
)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))-2/b^5
*(((A*b^2+3*B*a*b+1/2*B*b^2)*tan(1/2*d*x+1/2*c)^3+(-A*b^2+3*B*a*b-1/2*B*b
^2)*tan(1/2*d*x+1/2*c))/(1+tan(1/2*d*x+1/2*c)^2)^2+1/2*(6*A*a*b-12*B*a^2-B
*b^2)*arctan(tan(1/2*d*x+1/2*c))))
```

3.265. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

3.265.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 871 vs. $2(378) = 756$.

Time = 0.50 (sec) , antiderivative size = 1812, normalized size of antiderivative = 4.55

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="fracas")
```

```
output [1/4*(2*(12*B*a^8*b^2 - 6*A*a^7*b^3 - 35*B*a^6*b^4 + 18*A*a^5*b^5 + 33*B*a^4*b^6 - 18*A*a^3*b^7 - 9*B*a^2*b^8 + 6*A*a*b^9 - B*b^10)*d*x*cos(d*x + c)^2 + 4*(12*B*a^9*b - 6*A*a^8*b^2 - 35*B*a^7*b^3 + 18*A*a^6*b^4 + 33*B*a^5*b^5 - 18*A*a^4*b^6 - 9*B*a^3*b^7 + 6*A*a^2*b^8 - B*a*b^9)*d*x*cos(d*x + c) + 2*(12*B*a^10 - 6*A*a^9*b - 35*B*a^8*b^2 + 18*A*a^7*b^3 + 33*B*a^6*b^4 - 18*A*a^5*b^5 - 9*B*a^4*b^6 + 6*A*a^3*b^7 - B*a^2*b^8)*d*x + (12*B*a^9 - 6*A*a^8*b - 29*B*a^7*b^2 + 15*A*a^6*b^3 + 20*B*a^5*b^4 - 12*A*a^4*b^5 + (12*B*a^7*b^2 - 6*A*a^6*b^3 - 29*B*a^5*b^4 + 15*A*a^4*b^5 + 20*B*a^3*b^6 - 12*A*a^2*b^7)*cos(d*x + c)^2 + 2*(12*B*a^8*b - 6*A*a^7*b^2 - 29*B*a^6*b^3 + 15*A*a^5*b^4 + 20*B*a^4*b^5 - 12*A*a^3*b^6)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2))*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(12*B*a^9*b - 6*A*a^8*b^2 - 33*B*a^7*b^3 + 17*A*a^6*b^4 + 27*B*a^5*b^5 - 13*A*a^4*b^6 - 6*B*a^3*b^7 + 2*A*a^2*b^8 - (B*a^6*b^4 - 3*B*a^4*b^6 + 3*B*a^2*b^8 - B*b^10)*cos(d*x + c)^3 + 2*(2*B*a^7*b^3 - A*a^6*b^4 - 6*B*a^5*b^5 + 3*A*a^4*b^6 + 6*B*a^3*b^7 - 3*A*a^2*b^8 - 2*B*a*b^9 + A*b^10)*cos(d*x + c)^2 + (18*B*a^8*b^2 - 9*A*a^7*b^3 - 50*B*a^6*b^4 + 25*A*a^5*b^5 + 43*B*a^4*b^6 - 20*A*a^3*b^7 - 11*B*a^2*b^8 + 4*A*a*b^9)*cos(d*x + c))*sin(d*x + c))/((a^6*b^7 - 3*a^4*b^9 + 3*a^2*b^11 - b^13)*d*cos(d*x + c)^2 + 2*(a^7*b^6 - 3*a^5*b^8 + 3*a^3*b^10 - a*b^12)*d*...
```

3.265.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)
```

3.265. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

output Timed out

3.265.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.265.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2712 vs. 2(378) = 756.

Time = 0.54 (sec) , antiderivative size = 2712, normalized size of antiderivative = 6.81

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output

```

1/2*((3*(2*a^5*b - a^4*b^2 - 4*a^3*b^3 + 2*a^2*b^4 + 2*a*b^5)*sqrt(a^2 - b
^2)*A*abs(a^4*b^5 - 2*a^2*b^7 + b^9)*abs(-a + b) - (12*a^6 - 6*a^5*b - 23*
a^4*b^2 + 10*a^3*b^3 + 10*a^2*b^4 - a*b^5 + b^6)*sqrt(a^2 - b^2)*B*abs(a^4
*b^5 - 2*a^2*b^7 + b^9)*abs(-a + b) + 3*(4*a^10*b^5 - 2*a^9*b^6 - 17*a^8*b
^7 + 8*a^7*b^8 + 28*a^6*b^9 - 12*a^5*b^10 - 21*a^4*b^11 + 8*a^3*b^12 + 6*a
^2*b^13 - 2*a*b^14)*sqrt(a^2 - b^2)*A*abs(-a + b) - (24*a^11*b^4 - 12*a^10
*b^5 - 100*a^9*b^6 + 47*a^8*b^7 + 158*a^7*b^8 - 68*a^6*b^9 - 111*a^5*b^10
+ 42*a^4*b^11 + 28*a^3*b^12 - 8*a^2*b^13 + a*b^14 - b^15)*sqrt(a^2 - b^2)*
B*abs(-a + b))*(pi*floor(1/2*(d*x + c)/pi + 1/2) + arctan(2*tan(1/2*d*x +
1/2*c)/sqrt((4*a^5*b^4 - 8*a^3*b^6 + 4*a*b^8 + sqrt(-16*(a^5*b^4 + a^4*b^5
- 2*a^3*b^6 - 2*a^2*b^7 + a*b^8 + b^9)*(a^5*b^4 - a^4*b^5 - 2*a^3*b^6 + 2
*a^2*b^7 + a*b^8 - b^9) + 16*(a^5*b^4 - 2*a^3*b^6 + a*b^8)^2)))/(a^5*b^4 -
a^4*b^5 - 2*a^3*b^6 + 2*a^2*b^7 + a*b^8 - b^9)))/((a^4*b^5 - 2*a^2*b^7 +
b^9)^2*(a^2 - 2*a*b + b^2) + (a^7*b^4 - 2*a^6*b^5 - a^5*b^6 + 4*a^4*b^7 -
a^3*b^8 - 2*a^2*b^9 + a*b^10)*abs(a^4*b^5 - 2*a^2*b^7 + b^9)) + (24*B*a^11
*b^4 - 12*A*a^10*b^5 - 12*B*a^10*b^5 + 6*A*a^9*b^6 - 100*B*a^9*b^6 + 51*A*
a^8*b^7 + 47*B*a^8*b^7 - 24*A*a^7*b^8 + 158*B*a^7*b^8 - 84*A*a^6*b^9 - 68*
B*a^6*b^9 + 36*A*a^5*b^10 - 111*B*a^5*b^10 + 63*A*a^4*b^11 + 42*B*a^4*b^11
- 24*A*a^3*b^12 + 28*B*a^3*b^12 - 18*A*a^2*b^13 - 8*B*a^2*b^13 + 6*A*a*b^
14 + B*a*b^14 - B*b^15 - 12*B*a^6*abs(a^4*b^5 - 2*a^2*b^7 + b^9) + 6*A*...

```

3.265.9 Mupad [B] (verification not implemented)

Time = 12.39 (sec) , antiderivative size = 10598, normalized size of antiderivative = 26.63

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Too large to display}$$

input `int((cos(c + d*x))^4*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3,x)`

output

$$\begin{aligned} & ((\tan(c/2 + (d*x)/2)^5*(3*B*b^7 - 36*B*a^7 - 2*A*b^7 + 10*A*a^2*b^5 + 16*A \\ & *a^3*b^4 - 35*A*a^4*b^3 - 9*A*a^5*b^2 + 5*B*a^2*b^5 - 26*B*a^3*b^4 - 29*B* \\ & a^4*b^3 + 67*B*a^5*b^2 - 4*A*a*b^6 + 18*A*a^6*b + 4*B*a*b^6 + 18*B*a^6*b)) \\ & /((a + b)^2*(b^6 - 2*a*b^5 + a^2*b^4)) - (\tan(c/2 + (d*x)/2)^3*(2*A*b^7 + \\ & 36*B*a^7 + 3*B*b^7 - 10*A*a^2*b^5 + 16*A*a^3*b^4 + 35*A*a^4*b^3 - 9*A*a^5* \\ & b^2 + 5*B*a^2*b^5 + 26*B*a^3*b^4 - 29*B*a^4*b^3 - 67*B*a^5*b^2 - 4*A*a*b^6 \\ & - 18*A*a^6*b - 4*B*a*b^6 + 18*B*a^6*b))/((a + b)^2*(b^6 - 2*a*b^5 + a^2*b \\ & ^4)) + (\tan(c/2 + (d*x)/2)^7*(B*b^6 - 12*B*a^6 - 2*A*b^6 + 4*A*a^2*b^4 - 1 \\ & 2*A*a^3*b^3 - 3*A*a^4*b^2 - 8*B*a^2*b^4 - 10*B*a^3*b^3 + 23*B*a^4*b^2 + 2* \\ & A*a*b^5 + 6*A*a^5*b + 5*B*a*b^5 + 6*B*a^5*b))/((a*b^4 - b^5)*(a + b)^2) + \\ & (\tan(c/2 + (d*x)/2)*(2*A*b^6 - 12*B*a^6 + B*b^6 - 4*A*a^2*b^4 - 12*A*a^3*b \\ & ^3 + 3*A*a^4*b^2 - 8*B*a^2*b^4 + 10*B*a^3*b^3 + 23*B*a^4*b^2 + 2*A*a*b^5 + \\ & 6*A*a^5*b - 5*B*a*b^5 - 6*B*a^5*b))/((a + b)*(b^6 - 2*a*b^5 + a^2*b^4)))/ \\ & (d*(2*a*b + \tan(c/2 + (d*x)/2)^4*(6*a^2 - 2*b^2) + \tan(c/2 + (d*x)/2)^2*(4 \\ & *a*b + 4*a^2) - \tan(c/2 + (d*x)/2)^6*(4*a*b - 4*a^2) + \tan(c/2 + (d*x)/2)^ \\ & 8*(a^2 - 2*a*b + b^2) + a^2 + b^2)) + (\operatorname{atan}((((8*\tan(c/2 + (d*x)/2)*(288* \\ & B^2*a^14 + B^2*b^14 - 2*B^2*a*b^13 - 288*B^2*a^13*b + 36*A^2*a^2*b^12 - 72 \\ & *A^2*a^3*b^11 + 36*A^2*a^4*b^10 + 288*A^2*a^5*b^9 - 288*A^2*a^6*b^8 - 432* \\ & A^2*a^7*b^7 + 441*A^2*a^8*b^6 + 288*A^2*a^9*b^5 - 288*A^2*a^10*b^4 - 72*A^ \\ & 2*a^11*b^3 + 72*A^2*a^12*b^2 + 21*B^2*a^2*b^12 - 40*B^2*a^3*b^11 + 74*B... \end{aligned}$$

3.266 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

3.266.1 Optimal result 2454
 3.266.2 Mathematica [A] (verified) 2455
 3.266.3 Rubi [A] (verified) 2455
 3.266.4 Maple [A] (verified) 2460
 3.266.5 Fricas [B] (verification not implemented) 2460
 3.266.6 Sympy [F(-1)] 2461
 3.266.7 Maxima [F(-2)] 2462
 3.266.8 Giac [B] (verification not implemented) 2462
 3.266.9 Mupad [B] (verification not implemented) 2463

3.266.1 Optimal result

Integrand size = 31, antiderivative size = 280

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

$$= \frac{(Ab-3aB)x}{b^4}$$

$$- \frac{a(2a^4Ab-5a^2Ab^3+6Ab^5-6a^5B+15a^3b^2B-12ab^4B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{5/2}b^4(a+b)^{5/2}d}$$

$$- \frac{(aAb-3a^2B+2b^2B) \sin(c+dx)}{2b^3(a^2-b^2)d} + \frac{a(Ab-aB) \cos^2(c+dx) \sin(c+dx)}{2b(a^2-b^2)d(a+b \cos(c+dx))^2}$$

$$- \frac{a^2(a^2Ab-4Ab^3-3a^3B+6ab^2B) \sin(c+dx)}{2b^3(a^2-b^2)^2d(a+b \cos(c+dx))}$$

output

```
(A*b-3*B*a)*x/b^4-a*(2*A*a^4*b-5*A*a^2*b^3+6*A*b^5-6*B*a^5+15*B*a^3*b^2-12
*B*a*b^4)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(5/2)/b
^4/(a+b)^(5/2)/d-1/2*(A*a*b-3*B*a^2+2*B*b^2)*sin(d*x+c)/b^3/(a^2-b^2)/d+1/
2*a*(A*b-B*a)*cos(d*x+c)^2*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^2-1/2
*a^2*(A*a^2*b-4*A*b^3-3*B*a^3+6*B*a*b^2)*sin(d*x+c)/b^3/(a^2-b^2)^2/d/(a+b
*cos(d*x+c))
```

3.266.2 Mathematica [A] (verified)

Time = 2.26 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.83

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \frac{2(Ab-3aB)(c+dx) - \frac{2a(-2a^4Ab+5a^2Ab^3-6Ab^5+6a^5B-15a^3b^2B+12ab^4B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + 2bB\sin(c+dx)}{2b^4d}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output `(2*(A*b - 3*a*B)*(c + d*x) - (2*a*(-2*a^4*A*b + 5*a^2*A*b^3 - 6*A*b^5 + 6*a^5*B - 15*a^3*b^2*B + 12*a*b^4*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(5/2) + 2*b*B*Sin[c + d*x] + (a^3*b*(A*b - a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x])^2) + (a^2*b*(-3*a^2*A*b + 6*A*b^3 + 5*a^3*B - 8*a*b^2*B)*Sin[c + d*x])/((a - b)^2*(a + b)^2*(a + b*Cos[c + d*x]))/(2*b^4*d)`

3.266.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.13, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3468, 25, 3042, 3510, 3042, 3502, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx$$

$$\downarrow \text{3468}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} - \int \frac{\cos(c+dx)(-((-3Ba^2+Ab+2b^2B)\cos^2(c+dx)-2b(Ab-aB)\cos(c+dx)+2a(Ab-aB))}{(a+b\cos(c+dx))^2}}{2b(a^2-b^2)} dx$$

$$\begin{aligned}
 & \downarrow 25 \\
 & \int \frac{\cos(c+dx) \left(- \left((-3Ba^2 + Aba + 2b^2B) \cos^2(c+dx) - 2b(Ab - aB) \cos(c+dx) + 2a(Ab - aB) \right) \right)}{(a+b \cos(c+dx))^2} dx + \\
 & \quad \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3042 \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2}) \left((3Ba^2 - Aba - 2b^2B) \sin(c+dx+\frac{\pi}{2})^2 - 2b(Ab - aB) \sin(c+dx+\frac{\pi}{2}) + 2a(Ab - aB) \right)}{(a+b \sin(c+dx+\frac{\pi}{2}))^2} dx + \\
 & \quad \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3510 \\
 & \int \frac{-b(a^2 - b^2) \left((-3Ba^2 + Aba + 2b^2B) \cos^2(c+dx) + (a^2 - b^2) \left(-3Ba^3 + Aba^2 + 4b^2Ba - 2Ab^3 \right) \cos(c+dx) + ab \left(-3Ba^3 + Aba^2 + 6b^2Ba - 4Ab^3 \right) \right)}{a+b \cos(c+dx)} dx - \frac{a^2(-3a^3B + a^2)}{b^2d(a^2 - b^2)} \\
 & \quad \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3042 \\
 & \int \frac{-b(a^2 - b^2) \left((-3Ba^2 + Aba + 2b^2B) \sin(c+dx+\frac{\pi}{2})^2 + (a^2 - b^2) \left(-3Ba^3 + Aba^2 + 4b^2Ba - 2Ab^3 \right) \sin(c+dx+\frac{\pi}{2}) + ab \left(-3Ba^3 + Aba^2 + 6b^2Ba - 4Ab^3 \right) \right)}{a+b \sin(c+dx+\frac{\pi}{2})} dx - \frac{a^2(-3a^3B + a^2)}{b^2d(a^2 - b^2)} \\
 & \quad \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3502 \\
 & \int \frac{a \left(-3Ba^3 + Aba^2 + 6b^2Ba - 4Ab^3 \right) b^2 + 2(a^2 - b^2)^2 (Ab - 3aB) \cos(c+dx) b}{a+b \cos(c+dx)} dx - \frac{(a^2 - b^2) \left(-3a^2B + aAb + 2b^2B \right) \sin(c+dx)}{d} - \frac{a^2(-3a^3B + a^2Ab + 6ab^2B - 4Ab^3)}{b^2d(a^2 - b^2)(a + b \cos(c + dx))} \\
 & \quad \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3042
 \end{aligned}$$

3.266. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

$$\frac{\int \frac{a(-3Ba^3 + Aba^2 + 6b^2Ba - 4Ab^3)b^2 + 2(a^2 - b^2)^2(Ab - 3aB)\sin(c + dx + \frac{\pi}{2})b}{a + b\sin(c + dx + \frac{\pi}{2})} dx - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d} - \frac{a^2(-3a^3B + a^2Ab + 6ab^2B - 4b^3B)}{b^2d(a^2 - b^2)(a + b\cos(c + dx))}}{b^2(a^2 - b^2)} = \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} \frac{a(Ab - aB)\sin(c + dx)\cos^2(c + dx)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2}$$

↓ 3214

$$\frac{2x(a^2 - b^2)^2(Ab - 3aB) - a(-6a^5B + 2a^4Ab + 15a^3b^2B - 5a^2Ab^3 - 12ab^4B + 6Ab^5) \int \frac{1}{a + b\cos(c + dx)} dx - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d} - \frac{a^2(-3a^3B + a^2Ab + 6ab^2B - 4b^3B)}{b^2d(a^2 - b^2)(a + b\cos(c + dx))}}{b^2(a^2 - b^2)} = \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} \frac{a(Ab - aB)\sin(c + dx)\cos^2(c + dx)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2}$$

↓ 3042

$$\frac{2x(a^2 - b^2)^2(Ab - 3aB) - a(-6a^5B + 2a^4Ab + 15a^3b^2B - 5a^2Ab^3 - 12ab^4B + 6Ab^5) \int \frac{1}{a + b\sin(c + dx + \frac{\pi}{2})} dx - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d} - \frac{a^2(-3a^3B + a^2Ab + 6ab^2B - 4b^3B)}{b^2d(a^2 - b^2)(a + b\cos(c + dx))}}{b^2(a^2 - b^2)} = \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} \frac{a(Ab - aB)\sin(c + dx)\cos^2(c + dx)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2}$$

↓ 3138

$$\frac{2a(-6a^5B + 2a^4Ab + 15a^3b^2B - 5a^2Ab^3 - 12ab^4B + 6Ab^5) \int \frac{1}{(a - b)\tan^2(\frac{1}{2}(c + dx)) + a + b} d \tan(\frac{1}{2}(c + dx)) - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d} - \frac{a^2(-3a^3B + a^2Ab + 6ab^2B - 4b^3B)}{b^2d(a^2 - b^2)(a + b\cos(c + dx))}}{b^2(a^2 - b^2)} = \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} \frac{a(Ab - aB)\sin(c + dx)\cos^2(c + dx)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} + \frac{2a(-6a^5B + 2a^4Ab + 15a^3b^2B - 5a^2Ab^3 - 12ab^4B + 6Ab^5) \arctan\left(\frac{\sqrt{a - b}\tan(\frac{1}{2}(c + dx))}{\sqrt{a + b}}\right)}{d\sqrt{a - b}\sqrt{a + b}} - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d}$$

↓ 218

$$\frac{2x(a^2 - b^2)^2(Ab - 3aB) - \frac{2a(-6a^5B + 2a^4Ab + 15a^3b^2B - 5a^2Ab^3 - 12ab^4B + 6Ab^5) \arctan\left(\frac{\sqrt{a - b}\tan(\frac{1}{2}(c + dx))}{\sqrt{a + b}}\right)}{d\sqrt{a - b}\sqrt{a + b}} - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d} - \frac{a^2(-3a^3B + a^2Ab + 6ab^2B - 4b^3B)}{b^2d(a^2 - b^2)(a + b\cos(c + dx))}}{b^2(a^2 - b^2)} = \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} \frac{a(Ab - aB)\sin(c + dx)\cos^2(c + dx)}{2bd(a^2 - b^2)(a + b\cos(c + dx))^2} + \frac{2a(-6a^5B + 2a^4Ab + 15a^3b^2B - 5a^2Ab^3 - 12ab^4B + 6Ab^5) \arctan\left(\frac{\sqrt{a - b}\tan(\frac{1}{2}(c + dx))}{\sqrt{a + b}}\right)}{d\sqrt{a - b}\sqrt{a + b}} - \frac{(a^2 - b^2)(-3a^2B + aAb + 2b^2B)\sin(c + dx)}{d}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

3.266. $\int \frac{\cos^3(c + dx)(A + B\cos(c + dx))}{(a + b\cos(c + dx))^3} dx$


```
output (a*(A*b - a*B)*Cos[c + d*x]^2*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*(a + b*Cos[
c + d*x])^2) + (-((a^2*(a^2*A*b - 4*A*b^3 - 3*a^3*B + 6*a*b^2*B)*Sin[c + d
*x])/(b^2*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))) + ((2*(a^2 - b^2)^2*(A*b -
3*a*B)*x - (2*a*(2*a^4*A*b - 5*a^2*A*b^3 + 6*A*b^5 - 6*a^5*B + 15*a^3*b^2*
B - 12*a*b^4*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[
a - b]*Sqrt[a + b]*d))/b - ((a^2 - b^2)*(a*A*b - 3*a^2*B + 2*b^2*B)*Sin[c
+ d*x])/d)/(b^2*(a^2 - b^2))/(2*b*(a^2 - b^2))
```

3.266.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

```
rule 3214 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d
*Sine[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(b*c - a*d))*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2)), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.266.4 Maple [A] (verified)

Time = 1.70 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.22

method	result
derivativedivides	$2a \left(\frac{(2Aa^2b - Aab^2 - 6Ab^3 - 4Ba^3 + Ba^2b + 8Bab^2)ab \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{2(a-b)(a^2 + 2ab + b^2)} + \frac{ba(2Aa^2b + Aab^2 - 6Ab^3 - 4Ba^3 - Ba^2b + 8Bab^2) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{1}{\left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)^2}$
default	$2a \left(\frac{(2Aa^2b - Aab^2 - 6Ab^3 - 4Ba^3 + Ba^2b + 8Bab^2)ab \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{2(a-b)(a^2 + 2ab + b^2)} + \frac{ba(2Aa^2b + Aab^2 - 6Ab^3 - 4Ba^3 - Ba^2b + 8Bab^2) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{1}{\left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) a - b \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b \right)^2}$
risch	Expression too large to display

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-2*a/b^4*((1/2*(2*A*a^2*b-A*a*b^2-6*A*b^3-4*B*a^3+B*a^2*b+8*B*a*b^2)*a*b/(a-b)/(a^2+2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3+1/2*b*a*(2*A*a^2*b+A*a*b^2-6*A*b^3-4*B*a^3-B*a^2*b+8*B*a*b^2)/(a+b)/(a-b)^2*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^2+1/2*(2*A*a^4*b-5*A*a^2*b^3+6*A*b^5-6*B*a^5+15*B*a^3*b^2-12*B*a*b^4)/(a^4-2*a^2*b^2+b^4)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))+2/b^4*(B*b*tan(1/2*d*x+1/2*c)/(1+tan(1/2*d*x+1/2*c)^2)+(A*b-3*B*a)*arctan(tan(1/2*d*x+1/2*c))))
```

3.266.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 746 vs. 2(268) = 536.

Time = 0.41 (sec) , antiderivative size = 1561, normalized size of antiderivative = 5.58

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="fracas")
```

output

```

[-1/4*(4*(3*B*a^7*b^2 - A*a^6*b^3 - 9*B*a^5*b^4 + 3*A*a^4*b^5 + 9*B*a^3*b^
6 - 3*A*a^2*b^7 - 3*B*a*b^8 + A*b^9)*d*x*cos(d*x + c)^2 + 8*(3*B*a^8*b - A
*a^7*b^2 - 9*B*a^6*b^3 + 3*A*a^5*b^4 + 9*B*a^4*b^5 - 3*A*a^3*b^6 - 3*B*a^2
*b^7 + A*a*b^8)*d*x*cos(d*x + c) + 4*(3*B*a^9 - A*a^8*b - 9*B*a^7*b^2 + 3*
A*a^6*b^3 + 9*B*a^5*b^4 - 3*A*a^4*b^5 - 3*B*a^3*b^6 + A*a^2*b^7)*d*x - (6*
B*a^8 - 2*A*a^7*b - 15*B*a^6*b^2 + 5*A*a^5*b^3 + 12*B*a^4*b^4 - 6*A*a^3*b^
5 + (6*B*a^6*b^2 - 2*A*a^5*b^3 - 15*B*a^4*b^4 + 5*A*a^3*b^5 + 12*B*a^2*b^6
- 6*A*a*b^7)*cos(d*x + c)^2 + 2*(6*B*a^7*b - 2*A*a^6*b^2 - 15*B*a^5*b^3 +
5*A*a^4*b^4 + 12*B*a^3*b^5 - 6*A*a^2*b^6)*cos(d*x + c))*sqrt(-a^2 + b^2)*
log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2
)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2
*a*b*cos(d*x + c) + a^2)) - 2*(6*B*a^8*b - 2*A*a^7*b^2 - 17*B*a^6*b^3 + 7*
A*a^5*b^4 + 13*B*a^4*b^5 - 5*A*a^3*b^6 - 2*B*a^2*b^7 + 2*(B*a^6*b^3 - 3*B*
a^4*b^5 + 3*B*a^2*b^7 - B*b^9)*cos(d*x + c)^2 + (9*B*a^7*b^2 - 3*A*a^6*b^3
- 25*B*a^5*b^4 + 9*A*a^4*b^5 + 20*B*a^3*b^6 - 6*A*a^2*b^7 - 4*B*a*b^8)*co
s(d*x + c))*sin(d*x + c))/((a^6*b^6 - 3*a^4*b^8 + 3*a^2*b^10 - b^12)*d*cos
(d*x + c)^2 + 2*(a^7*b^5 - 3*a^5*b^7 + 3*a^3*b^9 - a*b^11)*d*cos(d*x + c)
+ (a^8*b^4 - 3*a^6*b^6 + 3*a^4*b^8 - a^2*b^10)*d), -1/2*(2*(3*B*a^7*b^2 -
A*a^6*b^3 - 9*B*a^5*b^4 + 3*A*a^4*b^5 + 9*B*a^3*b^6 - 3*A*a^2*b^7 - 3*B*a*
b^8 + A*b^9)*d*x*cos(d*x + c)^2 + 4*(3*B*a^8*b - A*a^7*b^2 - 9*B*a^6*b^...

```

3.266.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.266.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx = \text{Exception raised: ValueError}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de
```

3.266.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 543 vs. 2(268) = 536.

Time = 0.36 (sec) , antiderivative size = 543, normalized size of antiderivative = 1.94

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx = \frac{(6Ba^6 - 2Aa^5b - 15Ba^4b^2 + 5Aa^3b^3 + 12Ba^2b^4 - 6Aab^5) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^4b^4 - 2a^2b^6 + b^8)\sqrt{a^2 - b^2}}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="giac")
```

output

```

-((6*B*a^6 - 2*A*a^5*b - 15*B*a^4*b^2 + 5*A*a^3*b^3 + 12*B*a^2*b^4 - 6*A*a
*b^5)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1
/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^4*b^4 - 2*
a^2*b^6 + b^8)*sqrt(a^2 - b^2)) - (4*B*a^6*tan(1/2*d*x + 1/2*c)^3 - 2*A*a^
5*b*tan(1/2*d*x + 1/2*c)^3 - 5*B*a^5*b*tan(1/2*d*x + 1/2*c)^3 + 3*A*a^4*b^
2*tan(1/2*d*x + 1/2*c)^3 - 7*B*a^4*b^2*tan(1/2*d*x + 1/2*c)^3 + 5*A*a^3*b^
3*tan(1/2*d*x + 1/2*c)^3 + 8*B*a^3*b^3*tan(1/2*d*x + 1/2*c)^3 - 6*A*a^2*b^
4*tan(1/2*d*x + 1/2*c)^3 + 4*B*a^6*tan(1/2*d*x + 1/2*c) - 2*A*a^5*b*tan(1/
2*d*x + 1/2*c) + 5*B*a^5*b*tan(1/2*d*x + 1/2*c) - 3*A*a^4*b^2*tan(1/2*d*x
+ 1/2*c) - 7*B*a^4*b^2*tan(1/2*d*x + 1/2*c) + 5*A*a^3*b^3*tan(1/2*d*x + 1/
2*c) - 8*B*a^3*b^3*tan(1/2*d*x + 1/2*c) + 6*A*a^2*b^4*tan(1/2*d*x + 1/2*c)
)/((a^4*b^3 - 2*a^2*b^5 + b^7)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x +
1/2*c)^2 + a + b)^2) + (3*B*a - A*b)*(d*x + c)/b^4 - 2*B*tan(1/2*d*x + 1/
2*c)/((tan(1/2*d*x + 1/2*c)^2 + 1)*b^3))/d

```

3.266.9 Mupad [B] (verification not implemented)

Time = 7.99 (sec) , antiderivative size = 5542, normalized size of antiderivative = 19.79

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `int((cos(c + d*x))^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3,x)`

output

$$\begin{aligned} & ((\tan(c/2 + (d*x)/2)^5*(6*B*a^5 - 2*B*b^5 + 6*A*a^2*b^3 + A*a^3*b^2 + 4*B* \\ & a^2*b^3 - 12*B*a^3*b^2 - 2*A*a^4*b + 2*B*a*b^4 - 3*B*a^4*b))/((a*b^3 - b^4 \\ &)*(a + b)^2) + (\tan(c/2 + (d*x)/2)*(6*B*a^5 + 2*B*b^5 + 6*A*a^2*b^3 - A*a^ \\ & 3*b^2 - 4*B*a^2*b^3 - 12*B*a^3*b^2 - 2*A*a^4*b + 2*B*a*b^4 + 3*B*a^4*b))/((\\ & (a + b)*(b^5 - 2*a*b^4 + a^2*b^3)) + (2*\tan(c/2 + (d*x)/2)^3*(6*B*a^6 - 2* \\ & B*b^6 + 5*A*a^3*b^3 + 6*B*a^2*b^4 - 13*B*a^4*b^2 - 2*A*a^5*b))/((b*(a*b^2 - \\ & b^3)*(a + b)^2*(a - b)))/(d*(2*a*b + \tan(c/2 + (d*x)/2)^2*(2*a*b + 3*a^2 \\ & - b^2) + \tan(c/2 + (d*x)/2)^6*(a^2 - 2*a*b + b^2) + a^2 + b^2 - \tan(c/2 + \\ & (d*x)/2)^4*(2*a*b - 3*a^2 + b^2))) + (\log(\tan(c/2 + (d*x)/2) + 1i)*(A*b - \\ & 3*B*a)*1i)/(b^4*d) - (\log(\tan(c/2 + (d*x)/2) - 1i)*(A*b*1i - B*a*3i))/(b^4 \\ & *d) - (a*\operatorname{atan}(((a*((8*\tan(c/2 + (d*x)/2)*(4*A^2*b^12 + 72*B^2*a^12 - 8*A^2 \\ & *a*b^11 - 72*B^2*a^11*b + 24*A^2*a^2*b^10 + 32*A^2*a^3*b^9 - 52*A^2*a^4*b^ \\ & 8 - 48*A^2*a^5*b^7 + 57*A^2*a^6*b^6 + 32*A^2*a^7*b^5 - 32*A^2*a^8*b^4 - 8* \\ & A^2*a^9*b^3 + 8*A^2*a^10*b^2 + 36*B^2*a^2*b^10 - 72*B^2*a^3*b^9 + 36*B^2*a^ \\ & ^4*b^8 + 288*B^2*a^5*b^7 - 288*B^2*a^6*b^6 - 432*B^2*a^7*b^5 + 441*B^2*a^8 \\ & *b^4 + 288*B^2*a^9*b^3 - 288*B^2*a^10*b^2 - 24*A*B*a*b^11 - 48*A*B*a^11*b \\ & + 48*A*B*a^2*b^10 - 72*A*B*a^3*b^9 - 192*A*B*a^4*b^8 + 252*A*B*a^5*b^7 + 2 \\ & 88*A*B*a^6*b^6 - 318*A*B*a^7*b^5 - 192*A*B*a^8*b^4 + 192*A*B*a^9*b^3 + 48* \\ & A*B*a^10*b^2)))/(a*b^12 + b^13 - 3*a^2*b^11 - 3*a^3*b^10 + 3*a^4*b^9 + 3*a^ \\ & 5*b^8 - a^6*b^7 - a^7*b^6) + (a*((8*(4*A*b^18 - 8*A*a^2*b^16 + 34*A*a^3... \end{aligned}$$

3.267
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

3.267.1 Optimal result 2465
 3.267.2 Mathematica [A] (verified) 2465
 3.267.3 Rubi [A] (verified) 2466
 3.267.4 Maple [A] (verified) 2469
 3.267.5 Fricas [B] (verification not implemented) 2470
 3.267.6 Sympy [F(-1)] 2471
 3.267.7 Maxima [F(-2)] 2471
 3.267.8 Giac [B] (verification not implemented) 2471
 3.267.9 Mupad [B] (verification not implemented) 2472

3.267.1 Optimal result

Integrand size = 31, antiderivative size = 211

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

$$= \frac{Bx}{b^3} + \frac{(a^2Ab^3 + 2Ab^5 - 2a^5B + 5a^3b^2B - 6ab^4B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{5/2}b^3(a+b)^{5/2}d}$$

$$- \frac{a^2(Ab - aB) \sin(c+dx)}{2b^2(a^2 - b^2)d(a+b \cos(c+dx))^2} + \frac{a(a^2Ab - 4Ab^3 - 3a^3B + 6ab^2B) \sin(c+dx)}{2b^2(a^2 - b^2)^2d(a+b \cos(c+dx))}$$

```
output B*x/b^3+(A*a^2*b^3+2*A*b^5-2*B*a^5+5*B*a^3*b^2-6*B*a*b^4)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(5/2)/b^3/(a+b)^(5/2)/d-1/2*a^2*(A*b-B*a)*sin(d*x+c)/b^2/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/2*a*(A*a^2*b-4*A*b^3-3*B*a^3+6*B*a*b^2)*sin(d*x+c)/b^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

3.267.2 Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.97

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

$$= \frac{2B(c+dx)}{2b^3d} + \frac{2(-a^2Ab^3 - 2Ab^5 + 2a^5B - 5a^3b^2B + 6ab^4B) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + \frac{a^2b(-Ab+aB) \sin(c+dx)}{(a-b)(a+b)(a+b \cos(c+dx))^2} + \frac{ab(a^2Ab - 4Ab^3 - 3a^3B + 6ab^2B) \sin(c+dx)}{(a-b)(a+b)(a+b \cos(c+dx))^2}$$

3.267.
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output $(2*B*(c + d*x) + (2*(-(a^2*A*b^3) - 2*A*b^5 + 2*a^5*B - 5*a^3*b^2*B + 6*a*b^4*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^{(5/2)} + (a^2*b*(-(A*b) + a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x])^2) + (a*b*(a^2*A*b - 4*A*b^3 - 3*a^3*B + 6*a*b^2*B)*Sin[c + d*x])/((a - b)^2*(a + b)^2*(a + b*Cos[c + d*x]))/(2*b^3*d)$

3.267.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.17, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3467, 3042, 3500, 25, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^3} dx$$

↓ 3467

$$\frac{\int \frac{2b(a^2 - b^2)B \cos^2(c + dx) + (a^2 - 2b^2)(Ab - aB) \cos(c + dx) + 2ab(Ab - aB)}{(a + b \cos(c + dx))^2} dx}{2b^2(a^2 - b^2)} - \frac{a^2(Ab - aB) \sin(c + dx)}{2b^2 d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\frac{\int \frac{2b(a^2 - b^2)B \sin(c + dx + \frac{\pi}{2})^2 + (a^2 - 2b^2)(Ab - aB) \sin(c + dx + \frac{\pi}{2}) + 2ab(Ab - aB)}{(a + b \sin(c + dx + \frac{\pi}{2}))^2} dx}{2b^2(a^2 - b^2)} - \frac{a^2(Ab - aB) \sin(c + dx)}{2b^2 d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3500

$$\frac{\frac{a(-3a^3 B + a^2 Ab + 6ab^2 B - 4Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} - \int \frac{(Ba^3 + Aba^2 - 4b^2 Ba + 2Ab^3)b^2 + 2(a^2 - b^2)^2 B \cos(c + dx)b}{a + b \cos(c + dx)} dx}{2b^2(a^2 - b^2)} - \frac{a^2(Ab - aB) \sin(c + dx)}{2b^2 d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3.267. $\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{\int \frac{(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3)b^2 + 2(a^2 - b^2)^2 B \cos(c+dx)b}{a+b \cos(c+dx)} dx + \frac{a(-3a^3B + a^2Ab + 6ab^2B - 4Ab^3) \sin(c+dx)}{d(a^2 - b^2)(a+b \cos(c+dx))}}{2b^2(a^2 - b^2)} \\
& \frac{a^2(Ab - aB) \sin(c+dx)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3)b^2 + 2(a^2 - b^2)^2 B \sin(c+dx+\frac{\pi}{2})b}{a+b \sin(c+dx+\frac{\pi}{2})} dx + \frac{a(-3a^3B + a^2Ab + 6ab^2B - 4Ab^3) \sin(c+dx)}{d(a^2 - b^2)(a+b \cos(c+dx))}}{2b^2(a^2 - b^2)} \\
& \frac{a^2(Ab - aB) \sin(c+dx)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \downarrow 3214 \\
& \frac{(-2a^5B + 5a^3b^2B + a^2Ab^3 - 6ab^4B + 2Ab^5) \int \frac{1}{a+b \cos(c+dx)} dx + 2Bx(a^2 - b^2)^2}{b(a^2 - b^2)} + \frac{a(-3a^3B + a^2Ab + 6ab^2B - 4Ab^3) \sin(c+dx)}{d(a^2 - b^2)(a+b \cos(c+dx))} \\
& \frac{2b^2(a^2 - b^2)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \frac{a^2(Ab - aB) \sin(c+dx)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \downarrow 3042 \\
& \frac{(-2a^5B + 5a^3b^2B + a^2Ab^3 - 6ab^4B + 2Ab^5) \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx + 2Bx(a^2 - b^2)^2}{b(a^2 - b^2)} + \frac{a(-3a^3B + a^2Ab + 6ab^2B - 4Ab^3) \sin(c+dx)}{d(a^2 - b^2)(a+b \cos(c+dx))} \\
& \frac{2b^2(a^2 - b^2)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \frac{a^2(Ab - aB) \sin(c+dx)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \downarrow 3138 \\
& \frac{2(-2a^5B + 5a^3b^2B + a^2Ab^3 - 6ab^4B + 2Ab^5) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{d} + 2Bx(a^2 - b^2)^2}{b(a^2 - b^2)} + \frac{a(-3a^3B + a^2Ab + 6ab^2B - 4Ab^3) \sin(c+dx)}{d(a^2 - b^2)(a+b \cos(c+dx))} \\
& \frac{2b^2(a^2 - b^2)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \frac{a^2(Ab - aB) \sin(c+dx)}{2b^2d(a^2 - b^2)(a+b \cos(c+dx))^2} \\
& \downarrow 218
\end{aligned}$$

3.267. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

$$\frac{\frac{a(-3a^3B+a^2Ab+6ab^2B-4Ab^3)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))} + \frac{2Bx(a^2-b^2)^2 + \frac{2(-2a^5B+5a^3b^2B+a^2Ab^3-6ab^4B+2Ab^5)\arctan\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{d\sqrt{a-b}\sqrt{a+b}}}{b(a^2-b^2)}}{2b^2(a^2-b^2)} + \frac{a^2(Ab-aB)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output `-1/2*(a^2*(A*b - a*B)*Sin[c + d*x])/(b^2*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((2*(a^2 - b^2)^2*B*x + (2*(a^2*A*b^3 + 2*A*b^5 - 2*a^5*B + 5*a^3*b^2*B - 6*a*b^4*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*d))/(b*(a^2 - b^2)) + (a*(a^2*A*b - 4*A*b^3 - 3*a^3*B + 6*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*b^2*(a^2 - b^2))`

3.267.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sine[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

```
rule 3467 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.267.4 Maple [A] (verified)

Time = 1.30 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.34

method	result
derivativedivides	$2 \left(-\frac{(Aa^2b^2 + 4Ab^3 + 2Ba^3 - Ba^2b - 6Ba^2b^2)ab \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)} + \frac{ba(Aa^2b^2 - 4Ab^3 - 2Ba^3 - Ba^2b + 6Ba^2b^2) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{d}{b^3} + \frac{((\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^{a-b} (\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^{a+b})^2}{b^3}$
default	$2 \left(-\frac{(Aa^2b^2 + 4Ab^3 + 2Ba^3 - Ba^2b - 6Ba^2b^2)ab \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)} + \frac{ba(Aa^2b^2 - 4Ab^3 - 2Ba^3 - Ba^2b + 6Ba^2b^2) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{d}{b^3} + \frac{((\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^{a-b} (\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^{a+b})^2}{b^3}$
risch	Expression too large to display

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBOSE)
```

$$3.267. \int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

output $1/d*(2/b^3*((-1/2*(A*a*b^2+4*A*b^3+2*B*a^3-B*a^2*b-6*B*a*b^2)*a*b/(a-b)/(a^2+2*a*b+b^2)*\tan(1/2*d*x+1/2*c)^3+1/2*b*a*(A*a*b^2-4*A*b^3-2*B*a^3-B*a^2*b+6*B*a*b^2)/(a+b)/(a-b)^2*\tan(1/2*d*x+1/2*c))/(\tan(1/2*d*x+1/2*c)^2*a-b*\tan(1/2*d*x+1/2*c)^2+a+b)^2+1/2*(A*a^2*b^3+2*A*b^5-2*B*a^5+5*B*a^3*b^2-6*B*a*b^4)/(a^4-2*a^2*b^2+b^4)/((a-b)*(a+b))^(1/2)*\arctan((a-b)*\tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))+2*B/b^3*\arctan(\tan(1/2*d*x+1/2*c)))$

3.267.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 542 vs. $2(201) = 402$.

Time = 0.37 (sec) , antiderivative size = 1152, normalized size of antiderivative = 5.46

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="fracas")`

output $[1/4*(4*(B*a^6*b^2 - 3*B*a^4*b^4 + 3*B*a^2*b^6 - B*b^8)*d*x*\cos(d*x + c)^2 + 8*(B*a^7*b - 3*B*a^5*b^3 + 3*B*a^3*b^5 - B*a*b^7)*d*x*\cos(d*x + c) + 4*(B*a^8 - 3*B*a^6*b^2 + 3*B*a^4*b^4 - B*a^2*b^6)*d*x + (2*B*a^7 - 5*B*a^5*b^2 - A*a^4*b^3 + 6*B*a^3*b^4 - 2*A*a^2*b^5 + (2*B*a^5*b^2 - 5*B*a^3*b^4 - A*a^2*b^5 + 6*B*a*b^6 - 2*A*b^7)*\cos(d*x + c)^2 + 2*(2*B*a^6*b - 5*B*a^4*b^3 - A*a^3*b^4 + 6*B*a^2*b^5 - 2*A*a*b^6)*\cos(d*x + c))*\sqrt{-a^2 + b^2}*1\log((2*a*b*\cos(d*x + c) + (2*a^2 - b^2)*\cos(d*x + c)^2 + 2*\sqrt{-a^2 + b^2})*(a*\cos(d*x + c) + b)*\sin(d*x + c) - a^2 + 2*b^2)/(b^2*\cos(d*x + c)^2 + 2*a*b*\cos(d*x + c) + a^2)) - 2*(2*B*a^7*b - 7*B*a^5*b^3 + 3*A*a^4*b^4 + 5*B*a^3*b^5 - 3*A*a^2*b^6 + (3*B*a^6*b^2 - A*a^5*b^3 - 9*B*a^4*b^4 + 5*A*a^3*b^5 + 6*B*a^2*b^6 - 4*A*a*b^7)*\cos(d*x + c))*\sin(d*x + c))/((a^6*b^5 - 3*a^4*b^7 + 3*a^2*b^9 - b^11)*d*\cos(d*x + c)^2 + 2*(a^7*b^4 - 3*a^5*b^6 + 3*a^3*b^8 - a*b^10)*d*\cos(d*x + c) + (a^8*b^3 - 3*a^6*b^5 + 3*a^4*b^7 - a^2*b^9)*d), 1/2*(2*(B*a^6*b^2 - 3*B*a^4*b^4 + 3*B*a^2*b^6 - B*b^8)*d*x*\cos(d*x + c)^2 + 4*(B*a^7*b - 3*B*a^5*b^3 + 3*B*a^3*b^5 - B*a*b^7)*d*x*\cos(d*x + c) + 2*(B*a^8 - 3*B*a^6*b^2 + 3*B*a^4*b^4 - B*a^2*b^6)*d*x - (2*B*a^7 - 5*B*a^5*b^2 - A*a^4*b^3 + 6*B*a^3*b^4 - 2*A*a^2*b^5 + (2*B*a^5*b^2 - 5*B*a^3*b^4 - A*a^2*b^5 + 6*B*a*b^6 - 2*A*b^7)*\cos(d*x + c)^2 + 2*(2*B*a^6*b - 5*B*a^4*b^3 - A*a^3*b^4 + 6*B*a^2*b^5 - 2*A*a*b^6)*\cos(d*x + c))*\sqrt{a^2 - b^2}*\arctan(-(a*\cos(d*x + c) + b)/(\sqrt{a^2 - b^2}*\sin(d*x + c))) - (2*B...$

3.267.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)
```

```
output Timed out
```

3.267.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx = \text{Exception raised: ValueError}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

3.267.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 455 vs. 2(201) = 402.

Time = 0.33 (sec) , antiderivative size = 455, normalized size of antiderivative = 2.16

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx =$$

$$\frac{(2Ba^5 - 5Ba^3b^2 - Aa^2b^3 + 6Bab^4 - 2Ab^5) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - b \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^4b^3 - 2a^2b^5 + b^7)\sqrt{a^2 - b^2}} - \frac{(dx+c)B}{b^3} + \frac{2Ba^5}{b^3}$$

3.267. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output `-((2*B*a^5 - 5*B*a^3*b^2 - A*a^2*b^3 + 6*B*a*b^4 - 2*A*b^5)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^4*b^3 - 2*a^2*b^5 + b^7)*sqrt(a^2 - b^2)) - (d*x + c)*B/b^3 + (2*B*a^5*tan(1/2*d*x + 1/2*c)^3 - 3*B*a^4*b*tan(1/2*d*x + 1/2*c)^3 + A*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 - 5*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 + 3*A*a^2*b^3*tan(1/2*d*x + 1/2*c)^3 + 6*B*a^2*b^3*tan(1/2*d*x + 1/2*c)^3 - 4*A*a*b^4*tan(1/2*d*x + 1/2*c)^3 + 2*B*a^5*tan(1/2*d*x + 1/2*c) + 3*B*a^4*b*tan(1/2*d*x + 1/2*c) - A*a^3*b^2*tan(1/2*d*x + 1/2*c) - 5*B*a^3*b^2*tan(1/2*d*x + 1/2*c) + 3*A*a^2*b^3*tan(1/2*d*x + 1/2*c) - 6*B*a^2*b^3*tan(1/2*d*x + 1/2*c) + 4*A*a*b^4*tan(1/2*d*x + 1/2*c))/((a^4*b^2 - 2*a^2*b^4 + b^6)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^2))/d`

3.267.9 Mupad [B] (verification not implemented)

Time = 10.66 (sec) , antiderivative size = 6923, normalized size of antiderivative = 32.81

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Too large to display}$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3,x)`

output

$$\begin{aligned}
& (2*B*atan(-((B*((B*((8*(4*A*b^{15} + 4*B*b^{15} - 6*A*a^2*b^{13} + 6*A*a^3*b^{12} \\
& + 2*A*a^6*b^9 - 2*A*a^7*b^8 - 8*B*a^2*b^{13} + 34*B*a^3*b^{12} + 6*B*a^4*b^{11} \\
& - 36*B*a^5*b^{10} - 4*B*a^6*b^9 + 18*B*a^7*b^8 + 2*B*a^8*b^7 - 4*B*a^9*b^6 - \\
& 4*A*a*b^{14} - 12*B*a*b^{14}))/((a*b^{12} + b^{13} - 3*a^2*b^{11} - 3*a^3*b^{10} + 3*a \\
& ^4*b^9 + 3*a^5*b^8 - a^6*b^7 - a^7*b^6) - (B*tan(c/2 + (d*x)/2)*(8*a*b^{15} \\
& - 8*a^2*b^{14} - 32*a^3*b^{13} + 32*a^4*b^{12} + 48*a^5*b^{11} - 48*a^6*b^{10} - 32* \\
& a^7*b^9 + 32*a^8*b^8 + 8*a^9*b^7 - 8*a^{10}*b^6)*8i)/(b^3*(a*b^{10} + b^{11} - 3 \\
& *a^2*b^9 - 3*a^3*b^8 + 3*a^4*b^7 + 3*a^5*b^6 - a^6*b^5 - a^7*b^4)))*1i)/b^ \\
& 3 + (8*tan(c/2 + (d*x)/2)*(4*A^2*b^{10} + 8*B^2*a^{10} + 4*B^2*b^{10} - 8*B^2*a \\
& b^9 - 8*B^2*a^9*b + 4*A^2*a^2*b^8 + A^2*a^4*b^6 + 24*B^2*a^2*b^8 + 32*B^2* \\
& a^3*b^7 - 52*B^2*a^4*b^6 - 48*B^2*a^5*b^5 + 57*B^2*a^6*b^4 + 32*B^2*a^7*b^ \\
& 3 - 32*B^2*a^8*b^2 - 24*A*B*a*b^9 + 8*A*B*a^3*b^7 + 2*A*B*a^5*b^5 - 4*A*B* \\
& a^7*b^3))/((a*b^{10} + b^{11} - 3*a^2*b^9 - 3*a^3*b^8 + 3*a^4*b^7 + 3*a^5*b^6 - \\
& a^6*b^5 - a^7*b^4))/b^3 - (B*((B*((8*(4*A*b^{15} + 4*B*b^{15} - 6*A*a^2*b^{13} \\
& + 6*A*a^3*b^{12} + 2*A*a^6*b^9 - 2*A*a^7*b^8 - 8*B*a^2*b^{13} + 34*B*a^3*b^{12} \\
& + 6*B*a^4*b^{11} - 36*B*a^5*b^{10} - 4*B*a^6*b^9 + 18*B*a^7*b^8 + 2*B*a^8*b^7 \\
& - 4*B*a^9*b^6 - 4*A*a*b^{14} - 12*B*a*b^{14}))/((a*b^{12} + b^{13} - 3*a^2*b^{11} - \\
& 3*a^3*b^{10} + 3*a^4*b^9 + 3*a^5*b^8 - a^6*b^7 - a^7*b^6) + (B*tan(c/2 + (d* \\
& x)/2)*(8*a*b^{15} - 8*a^2*b^{14} - 32*a^3*b^{13} + 32*a^4*b^{12} + 48*a^5*b^{11} - 4 \\
& 8*a^6*b^{10} - 32*a^7*b^9 + 32*a^8*b^8 + 8*a^9*b^7 - 8*a^{10}*b^6)*8i)/(b^3...
\end{aligned}$$

3.268 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

3.268.1 Optimal result	2474
3.268.2 Mathematica [A] (verified)	2474
3.268.3 Rubi [A] (verified)	2475
3.268.4 Maple [A] (verified)	2478
3.268.5 Fricas [B] (verification not implemented)	2479
3.268.6 Sympy [F(-1)]	2479
3.268.7 Maxima [F(-2)]	2480
3.268.8 Giac [B] (verification not implemented)	2480
3.268.9 Mupad [B] (verification not implemented)	2481

3.268.1 Optimal result

Integrand size = 29, antiderivative size = 180

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx = -\frac{(3aAb - a^2B - 2b^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{5/2}(a+b)^{5/2}d} + \frac{a(Ab - aB) \sin(c+dx)}{2b(a^2 - b^2)d(a+b \cos(c+dx))^2} + \frac{(a^2Ab + 2Ab^3 + a^3B - 4ab^2B) \sin(c+dx)}{2b(a^2 - b^2)^2d(a+b \cos(c+dx))}$$

```
output - (3*A*a*b-B*a^2-2*B*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2)) / (a-b)^(5/2)/(a+b)^(5/2)/d+1/2*a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/2*(A*a^2*b+2*A*b^3+B*a^3-4*B*a*b^2)*sin(d*x+c)/b/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

3.268.2 Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.96

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx = \frac{2(-3aAb+a^2B+2b^2B) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + \frac{a(Ab-aB) \sin(c+dx)}{(a-b)b(a+b)(a+b \cos(c+dx))^2} + \frac{(a^2Ab+2Ab^3+a^3B-4ab^2B) \sin(c+dx)}{(a-b)^2b(a+b)^2(a+b \cos(c+dx))}$$

$2d$

3.268. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output `((-2*(-3*a*A*b + a^2*B + 2*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(5/2) + (a*(A*b - a*B)*Sin[c + d*x])/((a - b)*b*(a + b)*(a + b*Cos[c + d*x])^2) + ((a^2*A*b + 2*A*b^3 + a^3*B - 4*a*b^2*B)*Sin[c + d*x])/((a - b)^2*b*(a + b)^2*(a + b*Cos[c + d*x]))/(2*d)`

3.268.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.14, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 3447, 3042, 3500, 3042, 3233, 25, 27, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{3447} \\
 & \int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a+b\cos(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{a(Ab-aB)\sin(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} - \int \frac{2b(Ab-aB)-(Ba^2+Aba-2b^2B)\cos(c+dx)}{(a+b\cos(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(Ab-aB)\sin(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} - \int \frac{2b(Ab-aB)+(-Ba^2-Aba+2b^2B)\sin(c+dx+\frac{\pi}{2})}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3233}
 \end{aligned}$$

3.268. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

$$\begin{aligned}
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{\int -\frac{b(-Ba^2 + 3Aba - 2b^2B)}{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow 25 \\
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{\int \frac{b(-Ba^2 + 3Aba - 2b^2B)}{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow 27 \\
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{b(a^2(-B) + 3aAb - 2b^2B) \int \frac{1}{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow 3042 \\
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{b(a^2(-B) + 3aAb - 2b^2B) \int \frac{1}{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx}{a^2 - b^2} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow 3138 \\
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{2b(a^2(-B) + 3aAb - 2b^2B) \int \frac{1}{(a-b) \tan^2\left(\frac{1}{2}(c + dx)\right) + a + b} d \tan\left(\frac{1}{2}(c + dx)\right)}{d(a^2 - b^2)} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow 218 \\
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{2b(a^2(-B) + 3aAb - 2b^2B) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c + dx)\right)}{\sqrt{a+b}}\right)}{d\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \\
& \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{(a^3B + a^2Ab - 4ab^2B + 2Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}
\end{aligned}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

```
output (a*(A*b - a*B)*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) -
((2*b*(3*a*A*b - a^2*B - 2*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*d) - ((a^2*A*b + 2*A*b^3 + a^3*B - 4*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*b*(a^2 - b^2))
```

3.268.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 3233 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Ssin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Ssin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Ssin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[(- (A*b^2
- a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.268.4 Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.30

method	result
derivativedivides	$\frac{2 \left(-\frac{(2Aa^2 + Aab + 2Ab^2 - Ba^2 - 4Bab) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)} - \frac{(2Aa^2 - Aab + 2Ab^2 + Ba^2 - 4Bab) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a^2 - 2ab + b^2)} \right) (3Aab - Ba^2 - 2Bb^2)}{\left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a-b} \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a+b} \right)^2} - \frac{(3Aab - Ba^2 - 2Bb^2)}{(a^4 - 2a^2b^2 + b^4)}$
default	$\frac{2 \left(-\frac{(2Aa^2 + Aab + 2Ab^2 - Ba^2 - 4Bab) \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)} - \frac{(2Aa^2 - Aab + 2Ab^2 + Ba^2 - 4Bab) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a+b)(a^2 - 2ab + b^2)} \right) (3Aab - Ba^2 - 2Bb^2)}{\left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a-b} \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a+b} \right)^2} - \frac{(3Aab - Ba^2 - 2Bb^2)}{(a^4 - 2a^2b^2 + b^4)}$
risch	$\frac{i(3Aab^4e^{3i(dx+c)} + 2Ba^4be^{3i(dx+c)} - 5Ba^2b^3e^{3i(dx+c)} + 2Aa^4be^{2i(dx+c)} + 5Aa^2b^3e^{2i(dx+c)} + 2Ab^5e^{2i(dx+c)} + 2Ba^5e^{2i(dx+c)})}{b^2(a^2 + 2ab + b^2)}$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBOSE
)
```

```
output 1/d*(-2*(-1/2*(2*A*a^2+A*a*b+2*A*b^2-B*a^2-4*B*a*b)/(a-b)/(a^2+2*a*b+b^2)*
tan(1/2*d*x+1/2*c)^3-1/2*(2*A*a^2-A*a*b+2*A*b^2+B*a^2-4*B*a*b)/(a+b)/(a^2-
2*a*b+b^2)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c
)^2+a+b)^2-(3*A*a*b-B*a^2-2*B*b^2)/(a^4-2*a^2*b^2+b^4)/((a-b)*(a+b))^(1/2)
*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))
```

3.268.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 335 vs. $2(165) = 330$.

Time = 0.36 (sec) , antiderivative size = 740, normalized size of antiderivative = 4.11

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \left[-\frac{(Ba^4 - 3Aa^3b + 2Ba^2b^2 + (Ba^2b^2 - 3Aab^3 + 2Bb^4)\cos(dx+c)^2 + 2(Ba^3b - 3Aa^2b^2 + 2Bab^3)\cos(dx+c))\sqrt{a^2-b^2} + (Ba^4 - 3Aa^3b + 2Ba^2b^2 + (Ba^2b^2 - 3Aab^3 + 2Bb^4)\cos(dx+c)^2 + 2(Ba^3b - 3Aa^2b^2 + 2Bab^3)\cos(dx+c))\arctan\left(\frac{a\cos(dx+c)+b}{\sqrt{a^2-b^2}\sin(dx+c)}\right) + (2Aa^5 - 3Ba^4b - Aa^3b^2 + 3Ba^2b^3 - Aa^2b^3 + 4Ba^2b^4 - 2Aab^5)\cos(dx+c)\sin(dx+c)}{(a^6b^2 - 3a^4b^4 + 3a^2b^6 - b^8)d\cos(dx+c)^2 + 2(a^7b - 3a^5b^3 + 3a^3b^5 - ab^7)d\cos(dx+c) + (a^8 - 3a^6b^2 + 3a^4b^4 - a^2b^6)d} \right]$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="fricas")`

output `[-1/4*((B*a^4 - 3*A*a^3*b + 2*B*a^2*b^2 + (B*a^2*b^2 - 3*A*a*b^3 + 2*B*b^4)*cos(d*x + c)^2 + 2*(B*a^3*b - 3*A*a^2*b^2 + 2*B*a*b^3)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(2*A*a^5 - 3*B*a^4*b - A*a^3*b^2 + 3*B*a^2*b^3 - A*a*b^4 + (B*a^5 + A*a^4*b - 5*B*a^3*b^2 + A*a^2*b^3 + 4*B*a*b^4 - 2*A*b^5)*cos(d*x + c))*sin(d*x + c))/((a^6*b^2 - 3*a^4*b^4 + 3*a^2*b^6 - b^8)*d*cos(d*x + c)^2 + 2*(a^7*b - 3*a^5*b^3 + 3*a^3*b^5 - a*b^7)*d*cos(d*x + c) + (a^8 - 3*a^6*b^2 + 3*a^4*b^4 - a^2*b^6)*d), 1/2*((B*a^4 - 3*A*a^3*b + 2*B*a^2*b^2 + (B*a^2*b^2 - 3*A*a*b^3 + 2*B*b^4)*cos(d*x + c)^2 + 2*(B*a^3*b - 3*A*a^2*b^2 + 2*B*a*b^3)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (2*A*a^5 - 3*B*a^4*b - A*a^3*b^2 + 3*B*a^2*b^3 - A*a*b^4 + (B*a^5 + A*a^4*b - 5*B*a^3*b^2 + A*a^2*b^3 + 4*B*a*b^4 - 2*A*b^5)*cos(d*x + c))*sin(d*x + c))/((a^6*b^2 - 3*a^4*b^4 + 3*a^2*b^6 - b^8)*d*cos(d*x + c)^2 + 2*(a^7*b - 3*a^5*b^3 + 3*a^3*b^5 - a*b^7)*d*cos(d*x + c) + (a^8 - 3*a^6*b^2 + 3*a^4*b^4 - a^2*b^6)*d)]`

3.268.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)`

3.268. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

output Timed out

3.268.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.268.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 391 vs. 2(165) = 330.

Time = 0.32 (sec) , antiderivative size = 391, normalized size of antiderivative = 2.17

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$\frac{(Ba^2-3Aab+2Bb^2)\left(\pi\left[\frac{dx+c}{2\pi}+\frac{1}{2}\right]\text{sgn}(2a-2b)+\arctan\left(\frac{a\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)-b\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{\sqrt{a^2-b^2}}\right)\right)}{(a^4-2a^2b^2+b^4)\sqrt{a^2-b^2}} + \frac{2Aa^3\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)^3-Ba^3\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{(a^4-2a^2b^2+b^4)\sqrt{a^2-b^2}}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

```
output ((B*a^2 - 3*A*a*b + 2*B*b^2)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2
*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 -
b^2)))/((a^4 - 2*a^2*b^2 + b^4)*sqrt(a^2 - b^2)) + (2*A*a^3*tan(1/2*d*x +
1/2*c)^3 - B*a^3*tan(1/2*d*x + 1/2*c)^3 - A*a^2*b*tan(1/2*d*x + 1/2*c)^3 -
3*B*a^2*b*tan(1/2*d*x + 1/2*c)^3 + A*a*b^2*tan(1/2*d*x + 1/2*c)^3 + 4*B*a
*b^2*tan(1/2*d*x + 1/2*c)^3 - 2*A*b^3*tan(1/2*d*x + 1/2*c)^3 + 2*A*a^3*tan
(1/2*d*x + 1/2*c) + B*a^3*tan(1/2*d*x + 1/2*c) + A*a^2*b*tan(1/2*d*x + 1/2
*c) - 3*B*a^2*b*tan(1/2*d*x + 1/2*c) + A*a*b^2*tan(1/2*d*x + 1/2*c) - 4*B
a*b^2*tan(1/2*d*x + 1/2*c) + 2*A*b^3*tan(1/2*d*x + 1/2*c))/((a^4 - 2*a^2*b
^2 + b^4)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^2
)/d
```

3.268.9 Mupad [B] (verification not implemented)

Time = 3.76 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.38

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \frac{\tan\left(\frac{c}{2}+\frac{dx}{2}\right)^3 (2Aa^2+2Ab^2-Ba^2+Aab-4Bab)}{(a+b)^2(a-b)} + \frac{\tan\left(\frac{c}{2}+\frac{dx}{2}\right) (2Aa^2+2Ab^2+Ba^2-Aab-4Bab)}{(a+b)(a^2-2ab+b^2)}$$

$$d \left(2ab + \tan\left(\frac{c}{2}+\frac{dx}{2}\right)^2 (2a^2-2b^2) + \tan\left(\frac{c}{2}+\frac{dx}{2}\right)^4 (a^2-2ab+b^2) + a^2+b^2 \right)$$

$$+ \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2}+\frac{dx}{2}\right)(2a-2b)(a^2-2ab+b^2)}{2\sqrt{a+b}(a-b)^{5/2}}\right) (Ba^2-3Aab+2Bb^2)}{d(a+b)^{5/2}(a-b)^{5/2}}$$

```
input int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3,x)
```

```
output ((tan(c/2 + (d*x)/2)^3*(2*A*a^2 + 2*A*b^2 - B*a^2 + A*a*b - 4*B*a*b))/((a
+b)^2*(a - b)) + (tan(c/2 + (d*x)/2)*(2*A*a^2 + 2*A*b^2 + B*a^2 - A*a*b -
4*B*a*b))/((a + b)*(a^2 - 2*a*b + b^2)))/(d*(2*a*b + tan(c/2 + (d*x)/2)^2
*(2*a^2 - 2*b^2) + tan(c/2 + (d*x)/2)^4*(a^2 - 2*a*b + b^2) + a^2 + b^2))
+ (atan((tan(c/2 + (d*x)/2)*(2*a - 2*b)*(a^2 - 2*a*b + b^2))/(2*(a + b)^(1
/2)*(a - b)^(5/2)))*(B*a^2 + 2*B*b^2 - 3*A*a*b))/(d*(a + b)^(5/2)*(a - b)^(
5/2))
```


3.269 $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3} dx$

3.269.1 Optimal result 2482
 3.269.2 Mathematica [A] (verified) 2482
 3.269.3 Rubi [A] (verified) 2483
 3.269.4 Maple [A] (verified) 2486
 3.269.5 Fricas [B] (verification not implemented) 2486
 3.269.6 Sympy [F(-1)] 2487
 3.269.7 Maxima [F(-2)] 2487
 3.269.8 Giac [B] (verification not implemented) 2488
 3.269.9 Mupad [B] (verification not implemented) 2489

3.269.1 Optimal result

Integrand size = 23, antiderivative size = 164

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx = \frac{(2a^2 A + Ab^2 - 3abB) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{5/2}(a+b)^{5/2}d} - \frac{(Ab - aB) \sin(c + dx)}{2(a^2 - b^2) d(a + b \cos(c + dx))^2} - \frac{(3aAb - a^2B - 2b^2B) \sin(c + dx)}{2(a^2 - b^2)^2 d(a + b \cos(c + dx))}$$

```
output (2*A*a^2+A*b^2-3*B*a*b)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))
/(a-b)^(5/2)/(a+b)^(5/2)/d-1/2*(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d
*x+c))^2-1/2*(3*A*a*b-B*a^2-2*B*b^2)*sin(d*x+c)/(a^2-b^2)^2/d/(a+b*cos(d*x
+c))
```

3.269.2 Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.96

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx = \frac{2(2a^2 A + Ab^2 - 3abB) \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + \frac{(-Ab+aB) \sin(c+dx)}{(a-b)(a+b)(a+b \cos(c+dx))^2} + \frac{(-3aAb+a^2B+2b^2B) \sin(c+dx)}{(a-b)^2(a+b)^2(a+b \cos(c+dx))}$$

$2d$

input `Integrate[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^3,x]`

output `((-2*(2*a^2*A + A*b^2 - 3*a*b*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(5/2) + ((-A*b) + a*B)*Sin[c + d*x]/((a - b)*(a + b)*(a + b*Cos[c + d*x])^2) + ((-3*a*A*b + a^2*B + 2*b^2*B)*Sin[c + d*x])/((a - b)^2*(a + b)^2*(a + b*Cos[c + d*x]))/(2*d)`

3.269.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.16, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {3042, 3233, 25, 3042, 3233, 25, 27, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3} dx \\
 & \quad \downarrow \text{3233} \\
 & -\frac{\int -\frac{2(aA-bB)-(Ab-aB)\cos(c+dx)}{(a+b\cos(c+dx))^2} dx}{2(a^2-b^2)} - \frac{(Ab-aB)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{2(aA-bB)-(Ab-aB)\cos(c+dx)}{(a+b\cos(c+dx))^2} dx}{2(a^2-b^2)} - \frac{(Ab-aB)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2(aA-bB)+(aB-Ab)\sin\left(c+dx+\frac{\pi}{2}\right)}{(a+b\sin\left(c+dx+\frac{\pi}{2}\right))^2} dx}{2(a^2-b^2)} - \frac{(Ab-aB)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
 & \quad \downarrow \text{3233} \\
 & -\frac{\int -\frac{2Aa^2-3bBa+Ab^2}{a^2-b^2} dx}{2(a^2-b^2)} - \frac{(a^2(-B)+3aAb-2b^2B)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))} - \frac{(Ab-aB)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2}
 \end{aligned}$$

3.269. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3} dx$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{\int \frac{2Aa^2 - 3bBa + Ab^2}{a + b \cos(c + dx)} dx - \frac{(a^2(-B) + 3aAb - 2b^2B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 27 \\
 & \frac{(2a^2A - 3abB + Ab^2) \int \frac{1}{a + b \cos(c + dx)} dx - \frac{(a^2(-B) + 3aAb - 2b^2B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3042 \\
 & \frac{(2a^2A - 3abB + Ab^2) \int \frac{1}{a + b \sin(c + dx + \frac{\pi}{2})} dx - \frac{(a^2(-B) + 3aAb - 2b^2B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 3138 \\
 & \frac{2(2a^2A - 3abB + Ab^2) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx)) - \frac{(a^2(-B) + 3aAb - 2b^2B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{d(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow 218 \\
 & \frac{2(2a^2A - 3abB + Ab^2) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - \frac{(a^2(-B) + 3aAb - 2b^2B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{d\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^3,x]`

output `-1/2*((A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + (2*(2*a^2*A + A*b^2 - 3*a*b*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]]/(Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*d) - ((3*a*A*b - a^2*B - 2*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*(a^2 - b^2))`

3.269.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

3.269.4 Maple [A] (verified)

Time = 1.26 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.41

method	result
derivativedivides	$\frac{-\frac{(4Aab+Ab^2-2Ba^2-Bab-2Bb^2)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(a-b)(a^2+2ab+b^2)}-\frac{(4Aab-Ab^2-2Ba^2+Bab-2Bb^2)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{(a+b)(a^2-2ab+b^2)}}{\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a-b\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b}^2} + \frac{(2Aa^2+Ab^2-3Bab)\arctan\left(\frac{a-b}{a+b}\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(a^4-2a^2b^2+b^4)}$
default	$\frac{-\frac{(4Aab+Ab^2-2Ba^2-Bab-2Bb^2)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(a-b)(a^2+2ab+b^2)}-\frac{(4Aab-Ab^2-2Ba^2+Bab-2Bb^2)\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{(a+b)(a^2-2ab+b^2)}}{\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a-b\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b}^2} + \frac{(2Aa^2+Ab^2-3Bab)\arctan\left(\frac{a-b}{a+b}\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(a^4-2a^2b^2+b^4)}$
risch	$\frac{i(-2Aa^2b^2e^{3i(dx+c)}-Ab^4e^{3i(dx+c)}+3Ba^3b^3e^{3i(dx+c)}-6Aa^3be^{2i(dx+c)}-3Aab^3e^{2i(dx+c)}+2Ba^4e^{2i(dx+c)}+5Bb^2e^{2i(dx+c)})}{b(a^2-b^2)^2d(b^2e^{2i(dx+c)}+a^2)}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(2*(-1/2*(4*A*a*b+A*b^2-2*B*a^2-B*a*b-2*B*b^2)/(a-b)/(a^2+2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3-1/2*(4*A*a*b-A*b^2-2*B*a^2+B*a*b-2*B*b^2)/(a+b)/(a^2-2*a*b+b^2)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^2+(2*A*a^2+A*b^2-3*B*a*b)/(a^4-2*a^2*b^2+b^4)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))
```

3.269.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 336 vs. 2(150) = 300.

Time = 0.34 (sec) , antiderivative size = 742, normalized size of antiderivative = 4.52

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \left[\frac{(2Aa^4 - 3Ba^3b + Aa^2b^2 + (2Aa^2b^2 - 3Bab^3 + Ab^4) \cos(dx + c))^2 + 2(2Aa^3b - 3Ba^2b^2 + Aab^3) \cos(dx + c)}{4((a + b \cos(dx + c))^3 - (a - b \cos(dx + c))^3)} \right]$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="fracas")
```

output `[-1/4*((2*A*a^4 - 3*B*a^3*b + A*a^2*b^2 + (2*A*a^2*b^2 - 3*B*a*b^3 + A*b^4)*cos(d*x + c)^2 + 2*(2*A*a^3*b - 3*B*a^2*b^2 + A*a*b^3)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(2*B*a^5 - 4*A*a^4*b - B*a^3*b^2 + 5*A*a^2*b^3 - B*a*b^4 - A*b^5 + (B*a^4*b - 3*A*a^3*b^2 + B*a^2*b^3 + 3*A*a*b^4 - 2*B*b^5)*cos(d*x + c))*sin(d*x + c))/((a^6*b^2 - 3*a^4*b^4 + 3*a^2*b^6 - b^8)*d*cos(d*x + c)^2 + 2*(a^7*b - 3*a^5*b^3 + 3*a^3*b^5 - a*b^7)*d*cos(d*x + c) + (a^8 - 3*a^6*b^2 + 3*a^4*b^4 - a^2*b^6)*d), 1/2*((2*A*a^4 - 3*B*a^3*b + A*a^2*b^2 + (2*A*a^2*b^2 - 3*B*a*b^3 + A*b^4)*cos(d*x + c)^2 + 2*(2*A*a^3*b - 3*B*a^2*b^2 + A*a*b^3)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (2*B*a^5 - 4*A*a^4*b - B*a^3*b^2 + 5*A*a^2*b^3 - B*a*b^4 - A*b^5 + (B*a^4*b - 3*A*a^3*b^2 + B*a^2*b^3 + 3*A*a*b^4 - 2*B*b^5)*cos(d*x + c))*sin(d*x + c))/((a^6*b^2 - 3*a^4*b^4 + 3*a^2*b^6 - b^8)*d*cos(d*x + c)^2 + 2*(a^7*b - 3*a^5*b^3 + 3*a^3*b^5 - a*b^7)*d*cos(d*x + c) + (a^8 - 3*a^6*b^2 + 3*a^4*b^4 - a^2*b^6)*d)]`

3.269.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.269.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' f or more de

3.269.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 390 vs. $2(150) = 300$.

Time = 0.31 (sec) , antiderivative size = 390, normalized size of antiderivative = 2.38

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$\frac{(2Aa^2 - 3Bab + Ab^2) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^4 - 2a^2b^2 + b^4)\sqrt{a^2 - b^2}} + \frac{2Ba^3 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 - 4Aa^2b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(a^4 - 2a^2b^2 + b^4)\sqrt{a^2 - b^2}}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output $((2Aa^2 - 3Bab + Ab^2) * (\pi * \text{floor}(1/2 * (d*x + c) / \pi + 1/2) * \operatorname{sgn}(2a - 2b) + \arctan((a * \tan(1/2 * d*x + 1/2 * c) - b * \tan(1/2 * d*x + 1/2 * c)) / \sqrt{a^2 - b^2}))) / ((a^4 - 2a^2b^2 + b^4) * \sqrt{a^2 - b^2}) + (2Ba^3 * \tan(1/2 * d*x + 1/2 * c)^3 - 4Aa^2 * b * \tan(1/2 * d*x + 1/2 * c)^3 - Ba^2 * b * \tan(1/2 * d*x + 1/2 * c)^3 + 3Aa * a * b^2 * \tan(1/2 * d*x + 1/2 * c)^3 + Ba * b^2 * \tan(1/2 * d*x + 1/2 * c)^3 + A * b^3 * \tan(1/2 * d*x + 1/2 * c)^3 - 2B * b^3 * \tan(1/2 * d*x + 1/2 * c)^3 + 2B * a^3 * \tan(1/2 * d*x + 1/2 * c) - 4Aa^2 * b * \tan(1/2 * d*x + 1/2 * c) + Ba^2 * b * \tan(1/2 * d*x + 1/2 * c) - 3Aa * a * b^2 * \tan(1/2 * d*x + 1/2 * c) + Ba * a * b^2 * \tan(1/2 * d*x + 1/2 * c) + Ab * b^3 * \tan(1/2 * d*x + 1/2 * c) + 2B * b^3 * \tan(1/2 * d*x + 1/2 * c)) / ((a^4 - 2a^2b^2 + b^4) * (a * \tan(1/2 * d*x + 1/2 * c)^2 - b * \tan(1/2 * d*x + 1/2 * c)^2 + a + b)^2) / d$

3.269.9 Mupad [B] (verification not implemented)

Time = 3.66 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.51

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (2Ba^2 - Ab^2 + 2Bb^2 - 4Aab + Bab)}{(a+b)^2(a-b)} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (Ab^2 + 2Ba^2 + 2Bb^2 - 4Aab - Bab)}{(a+b)(a^2 - 2ab + b^2)}}{d \left(2ab + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 (2a^2 - 2b^2) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 (a^2 - 2ab + b^2) + a^2 + b^2 \right)}$$

$$+ \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (2a - 2b) (a^2 - 2ab + b^2)}{2\sqrt{a+b} (a-b)^{5/2}}\right) (2Aa^2 - 3Bab + Ab^2)}{d(a+b)^{5/2} (a-b)^{5/2}}$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^3,x)`output `((tan(c/2 + (d*x)/2)^3*(2*B*a^2 - A*b^2 + 2*B*b^2 - 4*A*a*b + B*a*b))/((a + b)^2*(a - b)) + (tan(c/2 + (d*x)/2)*(A*b^2 + 2*B*a^2 + 2*B*b^2 - 4*A*a*b - B*a*b))/((a + b)*(a^2 - 2*a*b + b^2)))/(d*(2*a*b + tan(c/2 + (d*x)/2)^2*(2*a^2 - 2*b^2) + tan(c/2 + (d*x)/2)^4*(a^2 - 2*a*b + b^2) + a^2 + b^2)) + (atan((tan(c/2 + (d*x)/2)*(2*a - 2*b)*(a^2 - 2*a*b + b^2))/(2*(a + b)^(1/2)*(a - b)^(5/2))))*(2*A*a^2 + A*b^2 - 3*B*a*b)/(d*(a + b)^(5/2)*(a - b)^(5/2))`

3.270 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^3} dx$

3.270.1 Optimal result 2490
 3.270.2 Mathematica [A] (verified) 2491
 3.270.3 Rubi [A] (verified) 2491
 3.270.4 Maple [A] (verified) 2495
 3.270.5 Fricas [B] (verification not implemented) 2495
 3.270.6 Sympy [F] 2496
 3.270.7 Maxima [F(-2)] 2497
 3.270.8 Giac [B] (verification not implemented) 2497
 3.270.9 Mupad [B] (verification not implemented) 2498

3.270.1 Optimal result

Integrand size = 29, antiderivative size = 214

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= -\frac{(6a^4Ab - 5a^2Ab^3 + 2Ab^5 - 2a^5B - a^3b^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^3(a-b)^{5/2}(a+b)^{5/2}d}$$

$$+ \frac{A \operatorname{arctanh}(\sin(c + dx))}{a^3d} + \frac{b(Ab - aB) \sin(c + dx)}{2a(a^2 - b^2)d(a + b \cos(c + dx))^2}$$

$$+ \frac{b(5a^2Ab - 2Ab^3 - 3a^3B) \sin(c + dx)}{2a^2(a^2 - b^2)^2d(a + b \cos(c + dx))}$$

output

```
-(6*A*a^4*b-5*A*a^2*b^3+2*A*b^5-2*B*a^5-B*a^3*b^2)*arctan((a-b)^(1/2)*tan(
1/2*d*x+1/2*c)/(a+b)^(1/2))/a^3/(a-b)^(5/2)/(a+b)^(5/2)/d+A*arctanh(sin(d*
x+c))/a^3/d+1/2*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/
2*b*(5*A*a^2*b-2*A*b^3-3*B*a^3)*sin(d*x+c)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+
c))
```

3.270.2 Mathematica [A] (verified)

Time = 1.25 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.26

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{\cos(c + dx)(B + A \sec(c + dx)) \left(-\frac{2(-6a^4Ab + 5a^2Ab^3 - 2Ab^5 + 2a^5B + a^3b^2B) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} - 2A \log(\cos(c + dx)) \right)}{(-a^2+b^2)^{5/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^3,x]`

output `(Cos[c + d*x]*(B + A*Sec[c + d*x])*((-2*(-6*a^4*A*b + 5*a^2*A*b^3 - 2*A*b^5 + 2*a^5*B + a^3*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(5/2) - 2*A*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 2*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (a^2*b*(A*b - a*B)*Sin[c + d*x])/((a - b)*(a + b)*(a + b*Cos[c + d*x])^2) + (a*b*(5*a^2*A*b - 2*A*b^3 - 3*a^3*B)*Sin[c + d*x])/((a - b)^2*(a + b)^2*(a + b*Cos[c + d*x]))) / (2*a^3*d*(A + B*Cos[c + d*x]))`

3.270.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.21, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 3479, 3042, 3534, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3} dx$$

↓ 3479

$$\frac{\int \frac{(b(Ab - aB) \cos^2(c + dx) - 2a(Ab - aB) \cos(c + dx) + 2A(a^2 - b^2)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx}{2a(a^2 - b^2)} + \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3.270. $\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \int \frac{b(Ab - aB) \sin(c + dx + \frac{\pi}{2})^2 - 2a(Ab - aB) \sin(c + dx + \frac{\pi}{2}) + 2A(a^2 - b^2)}{2a(a^2 - b^2) \sin(c + dx + \frac{\pi}{2}) (a + b \sin(c + dx + \frac{\pi}{2}))^2} dx + \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow \text{3534} \\
 & \int \frac{(2A(a^2 - b^2)^2 - a(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx + \frac{b(-3a^3B + 5a^2Ab - 2Ab^3) \sin(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} + \\
 & \quad \frac{2a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow \text{3042} \\
 & \int \frac{2A(a^2 - b^2)^2 - a(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sin(c + dx + \frac{\pi}{2})}{a(a^2 - b^2) \sin(c + dx + \frac{\pi}{2}) (a + b \sin(c + dx + \frac{\pi}{2}))} dx + \frac{b(-3a^3B + 5a^2Ab - 2Ab^3) \sin(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} + \\
 & \quad \frac{2a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow \text{3480} \\
 & \frac{2A(a^2 - b^2)^2 \int \sec(c + dx) dx}{a} - \frac{(-2a^5B + 6a^4Ab - a^3b^2B - 5a^2Ab^3 + 2Ab^5) \int \frac{1}{a + b \cos(c + dx)} dx}{a(a^2 - b^2)} + \frac{b(-3a^3B + 5a^2Ab - 2Ab^3) \sin(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} + \\
 & \quad \frac{2a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow \text{3042} \\
 & \frac{2A(a^2 - b^2)^2 \int \csc(c + dx + \frac{\pi}{2}) dx}{a} - \frac{(-2a^5B + 6a^4Ab - a^3b^2B - 5a^2Ab^3 + 2Ab^5) \int \frac{1}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{a(a^2 - b^2)} + \frac{b(-3a^3B + 5a^2Ab - 2Ab^3) \sin(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} + \\
 & \quad \frac{2a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow \text{3138} \\
 & \frac{2A(a^2 - b^2)^2 \int \csc(c + dx + \frac{\pi}{2}) dx}{a} - \frac{2(-2a^5B + 6a^4Ab - a^3b^2B - 5a^2Ab^3 + 2Ab^5) \int \frac{1}{(a - b) \tan^2(\frac{1}{2}(c + dx)) + a + b} d \tan(\frac{1}{2}(c + dx))}{ad(a^2 - b^2)}}{a(a^2 - b^2)} + \frac{b(-3a^3B + 5a^2Ab - 2Ab^3) \sin(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} + \\
 & \quad \frac{2a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \quad \frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \\
 & \downarrow \text{218}
 \end{aligned}$$

3.270. $\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx$

$$\frac{\frac{2A(a^2-b^2)^2 \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2(-2a^5B+6a^4Ab-a^3b^2B-5a^2Ab^3+2Ab^5) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{a(a^2-b^2)} + \frac{b(-3a^3B+5a^2Ab-2Ab^3) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} +$$

$$\frac{2a(a^2-b^2) b(Ab-aB) \sin(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 4257

$$\frac{b(Ab-aB) \sin(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2} +$$

$$\frac{b(-3a^3B+5a^2Ab-2Ab^3) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} + \frac{\frac{2A(a^2-b^2)^2 \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2(-2a^5B+6a^4Ab-a^3b^2B-5a^2Ab^3+2Ab^5) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{2a(a^2-b^2)}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^3,x]
```

```
output (b*(A*b - a*B)*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) +
((( -2*(6*a^4*A*b - 5*a^2*A*b^3 + 2*A*b^5 - 2*a^5*B - a^3*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) +
(2*A*(a^2 - b^2)^2*ArcTanh[Sin[c + d*x]])/(a*d))/(a*(a^2 - b^2)) + (b*(5*a^2*A*b - 2*A*b^3 - 3*a^3*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*a*(a^2 - b^2))
```

3.270.3.1 Defintions of rubi rules used

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3138 Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.270.4 Maple [A] (verified)

Time = 2.22 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.41

method	result
derivativedivides	$2 \left(\frac{-\frac{(6A^2b+Ab^2-2Ab^3-4Ba^3-Ba^2b)ab \tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a-b)(a^2+2ab+b^2)} - \frac{ba(6A^2b-Ab^2-2Ab^3-4Ba^3+Ba^2b) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{d}{\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^{a-b} + a+b} + \frac{(6A^2b+Ab^2-2Ab^3-4Ba^3-Ba^2b)ab \tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a-b)(a^2+2ab+b^2)} - \frac{ba(6A^2b-Ab^2-2Ab^3-4Ba^3+Ba^2b) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a+b)(a-b)^2} \frac{d}{\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^{a-b} + a+b} + \frac{1}{a^3}$
default	$2 \left(\frac{-\frac{(6A^2b+Ab^2-2Ab^3-4Ba^3-Ba^2b)ab \tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a-b)(a^2+2ab+b^2)} - \frac{ba(6A^2b-Ab^2-2Ab^3-4Ba^3+Ba^2b) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a+b)(a-b)^2} \right) \frac{d}{\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^{a-b} + a+b} + \frac{(6A^2b+Ab^2-2Ab^3-4Ba^3-Ba^2b)ab \tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a-b)(a^2+2ab+b^2)} - \frac{ba(6A^2b-Ab^2-2Ab^3-4Ba^3+Ba^2b) \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{2(a+b)(a-b)^2} \frac{d}{\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^{a-b} + a+b} + \frac{1}{a^3}$
risch	Expression too large to display

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-2/a^3*((-1/2*(6*A*a^2*b+A*a*b^2-2*A*b^3-4*B*a^3-B*a^2*b)*a*b/(a-b)/(a^2+2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3-1/2*b*a*(6*A*a^2*b-A*a*b^2-2*A*b^3-4*B*a^3+B*a^2*b)/(a+b)/(a-b)^2*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^2+1/2*(6*A*a^4*b-5*A*a^2*b^3+2*A*b^5-2*B*a^5-B*a^3*b^2)/(a^4-2*a^2*b^2+b^4)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-A/a^3*ln(tan(1/2*d*x+1/2*c)-1)+A/a^3*ln(tan(1/2*d*x+1/2*c)+1))
```

3.270.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 665 vs. 2(199) = 398.

Time = 13.01 (sec) , antiderivative size = 1400, normalized size of antiderivative = 6.54

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^3,x, algorithm="fracas")
```

output `[1/4*((2*B*a^7 - 6*A*a^6*b + B*a^5*b^2 + 5*A*a^4*b^3 - 2*A*a^2*b^5 + (2*B*a^5*b^2 - 6*A*a^4*b^3 + B*a^3*b^4 + 5*A*a^2*b^5 - 2*A*b^7)*cos(d*x + c)^2 + 2*(2*B*a^6*b - 6*A*a^5*b^2 + B*a^4*b^3 + 5*A*a^3*b^4 - 2*A*a*b^6)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 2*(A*a^8 - 3*A*a^6*b^2 + 3*A*a^4*b^4 - A*a^2*b^6 + (A*a^6*b^2 - 3*A*a^4*b^4 + 3*A*a^2*b^6 - A*b^8)*cos(d*x + c)^2 + 2*(A*a^7*b - 3*A*a^5*b^3 + 3*A*a^3*b^5 - A*a*b^7)*cos(d*x + c))*log(sin(d*x + c) + 1) - 2*(A*a^8 - 3*A*a^6*b^2 + 3*A*a^4*b^4 - A*a^2*b^6 + (A*a^6*b^2 - 3*A*a^4*b^4 + 3*A*a^2*b^6 - A*b^8)*cos(d*x + c)^2 + 2*(A*a^7*b - 3*A*a^5*b^3 + 3*A*a^3*b^5 - A*a*b^7)*cos(d*x + c))*log(-sin(d*x + c) + 1) - 2*(4*B*a^7*b - 6*A*a^6*b^2 - 5*B*a^5*b^3 + 9*A*a^4*b^4 + B*a^3*b^5 - 3*A*a^2*b^6 + (3*B*a^6*b^2 - 5*A*a^5*b^3 - 3*B*a^4*b^4 + 7*A*a^3*b^5 - 2*A*a*b^7)*cos(d*x + c))*sin(d*x + c))/((a^9*b^2 - 3*a^7*b^4 + 3*a^5*b^6 - a^3*b^8)*d*cos(d*x + c)^2 + 2*(a^10*b - 3*a^8*b^3 + 3*a^6*b^5 - a^4*b^7)*d*cos(d*x + c) + (a^11 - 3*a^9*b^2 + 3*a^7*b^4 - a^5*b^6)*d), 1/2*((2*B*a^7 - 6*A*a^6*b + B*a^5*b^2 + 5*A*a^4*b^3 - 2*A*a^2*b^5 + (2*B*a^5*b^2 - 6*A*a^4*b^3 + B*a^3*b^4 + 5*A*a^2*b^5 - 2*A*b^7)*cos(d*x + c)^2 + 2*(2*B*a^6*b - 6*A*a^5*b^2 + B*a^4*b^3 + 5*A*a^3*b^4 - 2*A*a*b^6)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(...`

3.270.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**3,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a + b*cos(c + d*x))**3, x)`

3.270.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.270.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 481 vs. 2(199) = 398.

Time = 0.34 (sec) , antiderivative size = 481, normalized size of antiderivative = 2.25

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{(2Ba^5 - 6Aa^4b + Ba^3b^2 + 5Aa^2b^3 - 2Ab^5) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^7 - 2a^5b^2 + a^3b^4) \sqrt{a^2 - b^2}} + \frac{A \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c)|)}{a^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output $((2*B*a^5 - 6*A*a^4*b + B*a^3*b^2 + 5*A*a^2*b^3 - 2*A*b^5)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^7 - 2*a^5*b^2 + a^3*b^4)*sqrt(a^2 - b^2)) + A*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^3 - A*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^3 - (4*B*a^4*b*tan(1/2*d*x + 1/2*c)^3 - 6*A*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 - 3*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 + 5*A*a^2*b^3*tan(1/2*d*x + 1/2*c)^3 - B*a^2*b^3*tan(1/2*d*x + 1/2*c)^3 + 3*A*a*b^4*tan(1/2*d*x + 1/2*c)^3 - 2*A*b^5*tan(1/2*d*x + 1/2*c)^3 + 4*B*a^4*b*tan(1/2*d*x + 1/2*c) - 6*A*a^3*b^2*tan(1/2*d*x + 1/2*c) + 3*B*a^3*b^2*tan(1/2*d*x + 1/2*c) - 5*A*a^2*b^3*tan(1/2*d*x + 1/2*c) - B*a^2*b^3*tan(1/2*d*x + 1/2*c) + 3*A*a*b^4*tan(1/2*d*x + 1/2*c) + 2*A*b^5*tan(1/2*d*x + 1/2*c))/(a^6 - 2*a^4*b^2 + a^2*b^4)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^2)/d$

3.270.9 Mupad [B] (verification not implemented)

Time = 9.90 (sec) , antiderivative size = 6913, normalized size of antiderivative = 32.30

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^3),x)`

output
$$\frac{((\tan(c/2 + (d*x)/2))^3*(2*A*b^4 - 6*A*a^2*b^2 + B*a^2*b^2 - A*a*b^3 + 4*B*a^3*b))/((a^2*b - a^3)*(a + b)^2) - (\tan(c/2 + (d*x)/2)*(2*A*b^4 - 6*A*a^2*b^2 - B*a^2*b^2 + A*a*b^3 + 4*B*a^3*b))/((a + b)*(a^4 - 2*a^3*b + a^2*b^2)))/((d*(2*a*b + \tan(c/2 + (d*x)/2)^2*(2*a^2 - 2*b^2) + \tan(c/2 + (d*x)/2)^4*(a^2 - 2*a*b + b^2) + a^2 + b^2)) - (A*\operatorname{atan}(((A*((8*\tan(c/2 + (d*x)/2)*(4*A^2*a^{10} + 8*A^2*b^{10} + 4*B^2*a^{10} - 8*A^2*a*b^9 - 8*A^2*a^9*b - 32*A^2*a^2*b^8 + 32*A^2*a^3*b^7 + 57*A^2*a^4*b^6 - 48*A^2*a^5*b^5 - 52*A^2*a^6*b^4 + 32*A^2*a^7*b^3 + 24*A^2*a^8*b^2 + B^2*a^6*b^4 + 4*B^2*a^8*b^2 - 24*A*B*a^9*b - 4*A*B*a^3*b^7 + 2*A*B*a^5*b^5 + 8*A*B*a^7*b^3)))/(a^{10}*b + a^{11} - a^4*b^7 - a^5*b^6 + 3*a^6*b^5 + 3*a^7*b^4 - 3*a^8*b^3 - 3*a^9*b^2)) + (A*((8*(4*A*a^{15} + 4*B*a^{15} - 4*A*a^6*b^9 + 2*A*a^7*b^8 + 18*A*a^8*b^7 - 4*A*a^9*b^6 - 36*A*a^{10}*b^5 + 6*A*a^{11}*b^4 + 34*A*a^{12}*b^3 - 8*A*a^{13}*b^2 - 2*B*a^8*b^7 + 2*B*a^9*b^6 + 6*B*a^{12}*b^3 - 6*B*a^{13}*b^2 - 12*A*a^{14}*b - 4*B*a^{14}*b)))/(a^{12}*b + a^{13} - a^6*b^7 - a^7*b^6 + 3*a^8*b^5 + 3*a^9*b^4 - 3*a^{10}*b^3 - 3*a^{11}*b^2) + (8*A*\tan(c/2 + (d*x)/2)*(8*a^{15}*b - 8*a^6*b^{10} + 8*a^7*b^9 + 32*a^8*b^8 - 32*a^9*b^7 - 48*a^{10}*b^6 + 48*a^{11}*b^5 + 32*a^{12}*b^4 - 32*a^{13}*b^3 - 8*a^{14}*b^2)))/(a^3*(a^{10}*b + a^{11} - a^4*b^7 - a^5*b^6 + 3*a^6*b^5 + 3*a^7*b^4 - 3*a^8*b^3 - 3*a^9*b^2))))/a^3 + (A*((8*\tan(c/2 + (d*x)/2)*(4*A^2*a^{10} + 8*A^2*b^{10} + 4*B^2*a^{10} - 8*A^2*a*b^9 - 8*A^2*a^9*b - 32*A^2*a^2*b^8 + 32*A^2*a^3*b^7 + 57*A^2*a^4*b^6 - 48*A^2*a^5*b^5...$$

3.271 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^3} dx$

3.271.1 Optimal result 2500
 3.271.2 Mathematica [A] (verified) 2501
 3.271.3 Rubi [A] (verified) 2501
 3.271.4 Maple [A] (verified) 2506
 3.271.5 Fricas [B] (verification not implemented) 2506
 3.271.6 Sympy [F] 2507
 3.271.7 Maxima [F(-2)] 2508
 3.271.8 Giac [B] (verification not implemented) 2508
 3.271.9 Mupad [B] (verification not implemented) 2509

3.271.1 Optimal result

Integrand size = 31, antiderivative size = 299

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{b(12a^4Ab - 15a^2Ab^3 + 6Ab^5 - 6a^5B + 5a^3b^2B - 2ab^4B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - (3Ab - aB) \operatorname{arctanh}(\sin(c + dx))}{a^4(a - b)^{5/2}(a + b)^{5/2}d} + \frac{a^4d(2a^4A - 11a^2Ab^2 + 6Ab^4 + 5a^3bB - 2ab^3B) \tan(c + dx)}{2a^3(a^2 - b^2)^2d} + \frac{b(Ab - aB) \tan(c + dx)}{2a(a^2 - b^2)d(a + b \cos(c + dx))^2} + \frac{b(6a^2Ab - 3Ab^3 - 4a^3B + ab^2B) \tan(c + dx)}{2a^2(a^2 - b^2)^2d(a + b \cos(c + dx))}$$

output

```
b*(12*A*a^4*b-15*A*a^2*b^3+6*A*b^5-6*B*a^5+5*B*a^3*b^2-2*B*a*b^4)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^4/(a-b)^(5/2)/(a+b)^(5/2)/d-(3*A*b-B*a)*arctanh(sin(d*x+c))/a^4/d+1/2*(2*A*a^4-11*A*a^2*b^2+6*A*b^4+5*B*a^3*b-2*B*a*b^3)*tan(d*x+c)/a^3/(a^2-b^2)^2/d+1/2*b*(A*b-B*a)*tan(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/2*b*(6*A*a^2*b-3*A*b^3-4*B*a^3+B*a*b^2)*tan(d*x+c)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

3.271.2 Mathematica [A] (verified)

Time = 4.34 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.18

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{2b(12a^4Ab - 15a^2Ab^3 + 6Ab^5 - 6a^5B + 5a^3b^2B - 2ab^4B) \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + 2(3Ab - aB) \log\left(\cos\left(\frac{1}{2}(c+dx)\right)\right) -$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^3,x]`output `((-2*b*(12*a^4*A*b - 15*a^2*A*b^3 + 6*A*b^5 - 6*a^5*B + 5*a^3*b^2*B - 2*a*b^4*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]]/(-a^2 + b^2)^(5/2) + 2*(3*A*b - a*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 2*(-3*A*b + a*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (2*a*A*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]) + (2*a*A*Sin[(c + d*x)/2])/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) + (a^2*b^2*(-(A*b) + a*B)*Sin[c + d*x])/(a - b)*(a + b)*(a + b*Cos[c + d*x])^2 + (a*b^2*(-7*a^2*A*b + 4*A*b^3 + 5*a^3*B - 2*a*b^2*B)*Sin[c + d*x])/((a - b)^2*(a + b)^2*(a + b*Cos[c + d*x])))/(2*a^4*d)`**3.271.3 Rubi [A] (verified)**Time = 2.02 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 3479, 3042, 3534, 3042, 3534, 25, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^2 (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^3} dx$$

$$\downarrow \text{3479}$$

$$\int \frac{(2Aa^2 + bBa - 2(Ab - aB) \cos(c + dx) a - 3Ab^2 + 2b(Ab - aB) \cos^2(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx + \frac{2a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\int \frac{2Aa^2 + bBa - 2(Ab - aB) \sin(c + dx + \frac{\pi}{2}) a - 3Ab^2 + 2b(Ab - aB) \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^2 (a + b \sin(c + dx + \frac{\pi}{2}))^2} dx + \frac{2a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3534

$$\int \frac{(2Aa^4 + 5bBa^3 - 11Ab^2a^2 - 2b^3Ba - (-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \cos(c + dx) a + 6Ab^4 + b(-4Ba^3 + 6Aba^2 + b^2Ba - 3Ab^3) \cos^2(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx + \frac{b(-4a^3B + 6a^2bA - 2ab^2A + b^3B)}{a(a^2 - b^2)} + \frac{2a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\int \frac{2Aa^4 + 5bBa^3 - 11Ab^2a^2 - 2b^3Ba - (-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sin(c + dx + \frac{\pi}{2}) a + 6Ab^4 + b(-4Ba^3 + 6Aba^2 + b^2Ba - 3Ab^3) \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^2 (a + b \sin(c + dx + \frac{\pi}{2}))} dx + \frac{b(-4a^3B + 6a^2bA - 2ab^2A + b^3B)}{a(a^2 - b^2)} + \frac{2a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3534

$$\int - \frac{(2(a^2 - b^2)^2(3Ab - aB) - ab(-4Ba^3 + 6Aba^2 + b^2Ba - 3Ab^3) \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx + \frac{(2a^4A + 5a^3bB - 11a^2Ab^2 - 2ab^3B + 6Ab^4) \tan(c + dx)}{ad} + \frac{b(-4a^3B + 6a^2bA - 2ab^2A + b^3B)}{ad(a^2 - b^2)} + \frac{2a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 25

$$\frac{(2a^4A + 5a^3bB - 11a^2Ab^2 - 2ab^3B + 6Ab^4) \tan(c + dx)}{ad} - \int \frac{(2(a^2 - b^2)^2(3Ab - aB) - ab(-4Ba^3 + 6Aba^2 + b^2Ba - 3Ab^3) \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx + \frac{b(-4a^3B + 6a^2bA - 2ab^2A + b^3B)}{ad(a^2 - b^2)} + \frac{2a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3.271. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx$

↓ 3042

$$\frac{\frac{(2a^4A+5a^3bB-11a^2Ab^2-2ab^3B+6Ab^4) \tan(c+dx)}{ad} - \frac{\int \frac{2(a^2-b^2)^2(3Ab-aB)-ab(-4Ba^3+6Aba^2+b^2Ba-3Ab^3) \sin(c+dx+\frac{\pi}{2}) dx}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}}{a(a^2-b^2)}}{2a(a^2-b^2)} + \frac{b(-4a^3B+6a^2Ab+aB^2)}{ad(a^2-b^2)(a+b \sin(c+dx+\frac{\pi}{2}))}$$

$$\frac{b(Ab-aB) \tan(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3480

$$\frac{\frac{(2a^4A+5a^3bB-11a^2Ab^2-2ab^3B+6Ab^4) \tan(c+dx)}{ad} - \frac{2(a^2-b^2)^2(3Ab-aB) \int \sec(c+dx) dx}{a} - \frac{b(-6a^5B+12a^4Ab+5a^3b^2B-15a^2Ab^3-2ab^4B+6Ab^5) \int \frac{1}{a+b \cos(c+dx)}}{a}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \tan(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3042

$$\frac{\frac{(2a^4A+5a^3bB-11a^2Ab^2-2ab^3B+6Ab^4) \tan(c+dx)}{ad} - \frac{2(a^2-b^2)^2(3Ab-aB) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b(-6a^5B+12a^4Ab+5a^3b^2B-15a^2Ab^3-2ab^4B+6Ab^5) \int \frac{1}{a+b \cos(c+dx)}}{a}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \tan(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3138

$$\frac{\frac{(2a^4A+5a^3bB-11a^2Ab^2-2ab^3B+6Ab^4) \tan(c+dx)}{ad} - \frac{2(a^2-b^2)^2(3Ab-aB) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b(-6a^5B+12a^4Ab+5a^3b^2B-15a^2Ab^3-2ab^4B+6Ab^5) \int \frac{1}{a+b \cos(c+dx)}}{ad}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \tan(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 218

$$\frac{\frac{(2a^4A+5a^3bB-11a^2Ab^2-2ab^3B+6Ab^4) \tan(c+dx)}{ad} - \frac{2(a^2-b^2)^2(3Ab-aB) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b(-6a^5B+12a^4Ab+5a^3b^2B-15a^2Ab^3-2ab^4B+6Ab^5) \arctan(\frac{a+b \cos(c+dx)}{a-b \sin(c+dx)})}{ad\sqrt{a-b}\sqrt{a+b}}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \tan(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 4257

3.271. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^3} dx$

$$\frac{b(Ab - aB) \tan(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{b(-4a^3B + 6a^2Ab + ab^2B - 3Ab^3) \tan(c + dx)}{ad(a^2 - b^2)(a + b \cos(c + dx))} + \frac{(2a^4A + 5a^3bB - 11a^2Ab^2 - 2ab^3B + 6Ab^4) \tan(c + dx)}{ad} - \frac{2(a^2 - b^2)^2(3Ab - aB) \operatorname{arctanh}(\sin(c + dx))}{ad} - \frac{2b(-)}{a(a^2 - b^2)}$$

$$2a(a^2 - b^2)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^3,x]`

output `(b*(A*b - a*B)*Tan[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((b*(6*a^2*A*b - 3*A*b^3 - 4*a^3*B + a*b^2*B)*Tan[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + (-(((-2*b*(12*a^4*A*b - 15*a^2*A*b^3 + 6*A*b^5 - 6*a^5*B + 5*a^3*b^2*B - 2*a*b^4*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + (2*(a^2 - b^2)^2*(3*A*b - a*B)*ArcTanh[Sin[c + d*x]])/(a*d))/a) + ((2*a^4*A - 11*a^2*A*b^2 + 6*A*b^4 + 5*a^3*b*B - 2*a*b^3*B)*Tan[c + d*x])/(a*d))/(a*(a^2 - b^2))/(2*a*(a^2 - b^2))`

3.271.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Ssin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.271.4 Maple [A] (verified)

Time = 2.54 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.26

method	result
derivativedivides	$-\frac{A}{a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(3Ab - Ba) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^4} + \frac{\left(\frac{(8A^2b + Aa^2b^2 - 4Ab^3 - 6Ba^3 - Ba^2b + 2Ba^2b^2)ab \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)}\right)}{\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a - b}$
default	$-\frac{A}{a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)} + \frac{(3Ab - Ba) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{a^4} + \frac{\left(\frac{(8A^2b + Aa^2b^2 - 4Ab^3 - 6Ba^3 - Ba^2b + 2Ba^2b^2)ab \tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)}{2(a-b)(a^2 + 2ab + b^2)}\right)}{\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a - b}$
risch	Expression too large to display

input `int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBOSE)`

output `1/d*(-A/a^3/(tan(1/2*d*x+1/2*c)-1)+(3*A*b-B*a)/a^4*ln(tan(1/2*d*x+1/2*c)-1)+2*b/a^4*((-1/2*(8*A*a^2*b+A*a*b^2-4*A*b^3-6*B*a^3-B*a^2*b+2*B*a*b^2))*a*b/(a-b)/(a^2+2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3-1/2*b*a*(8*A*a^2*b-A*a*b^2-4*A*b^3-6*B*a^3+B*a^2*b+2*B*a*b^2)/(a+b)/(a-b)^2*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^2+1/2*(12*A*a^4*b-15*A*a^2*b^3+6*A*b^5-6*B*a^5+5*B*a^3*b^2-2*B*a*b^4)/(a^4-2*a^2*b^2+b^4)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-A/a^3/(tan(1/2*d*x+1/2*c)+1)+1/a^4*(-3*A*b+B*a)*ln(tan(1/2*d*x+1/2*c)+1))`

3.271.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1015 vs. 2(284) = 568.

Time = 27.86 (sec) , antiderivative size = 2100, normalized size of antiderivative = 7.02

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^3,x, algorithm="fracas")`

output `[1/4*((6*B*a^5*b^3 - 12*A*a^4*b^4 - 5*B*a^3*b^5 + 15*A*a^2*b^6 + 2*B*a*b^7 - 6*A*b^8)*cos(d*x + c)^3 + 2*(6*B*a^6*b^2 - 12*A*a^5*b^3 - 5*B*a^4*b^4 + 15*A*a^3*b^5 + 2*B*a^2*b^6 - 6*A*a*b^7)*cos(d*x + c)^2 + (6*B*a^7*b - 12*A*a^6*b^2 - 5*B*a^5*b^3 + 15*A*a^4*b^4 + 2*B*a^3*b^5 - 6*A*a^2*b^6)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 2*((B*a^7*b^2 - 3*A*a^6*b^3 - 3*B*a^5*b^4 + 9*A*a^4*b^5 + 3*B*a^3*b^6 - 9*A*a^2*b^7 - B*a*b^8 + 3*A*b^9)*cos(d*x + c)^3 + 2*(B*a^8*b - 3*A*a^7*b^2 - 3*B*a^6*b^3 + 9*A*a^5*b^4 + 3*B*a^4*b^5 - 9*A*a^3*b^6 - B*a^2*b^7 + 3*A*a*b^8)*cos(d*x + c)^2 + (B*a^9 - 3*A*a^8*b - 3*B*a^7*b^2 + 9*A*a^6*b^3 + 3*B*a^5*b^4 - 9*A*a^4*b^5 - B*a^3*b^6 + 3*A*a^2*b^7)*cos(d*x + c))*log(sin(d*x + c) + 1) - 2*((B*a^7*b^2 - 3*A*a^6*b^3 - 3*B*a^5*b^4 + 9*A*a^4*b^5 + 3*B*a^3*b^6 - 9*A*a^2*b^7 - B*a*b^8 + 3*A*b^9)*cos(d*x + c)^3 + 2*(B*a^8*b - 3*A*a^7*b^2 - 3*B*a^6*b^3 + 9*A*a^5*b^4 + 3*B*a^4*b^5 - 9*A*a^3*b^6 - B*a^2*b^7 + 3*A*a*b^8)*cos(d*x + c)^2 + (B*a^9 - 3*A*a^8*b - 3*B*a^7*b^2 + 9*A*a^6*b^3 + 3*B*a^5*b^4 - 9*A*a^4*b^5 - B*a^3*b^6 + 3*A*a^2*b^7)*cos(d*x + c))*log(-sin(d*x + c) + 1) + 2*(2*A*a^9 - 6*A*a^7*b^2 + 6*A*a^5*b^4 - 2*A*a^3*b^6 + (2*A*a^7*b^2 + 5*B*a^6*b^3 - 13*A*a^5*b^4 - 7*B*a^4*b^5 + 17*A*a^3*b^6 + 2*B*a^2*b^7 - 6*A*a*b^8)*cos(d*x + c)^2 + (4*A*a^8*b + 6*B*a^7*b^2 - 20*A*a^6*b^...`

3.271.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**3,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a + b*cos(c + d*x))**3, x)`

3.271.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Exception raised: ValueError}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^3,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de
```

3.271.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 574 vs. 2(284) = 568.

Time = 0.36 (sec) , antiderivative size = 574, normalized size of antiderivative = 1.92

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{(6Ba^5b - 12Aa^4b^2 - 5Ba^3b^3 + 15Aa^2b^4 + 2Bab^5 - 6Ab^6) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^8 - 2a^6b^2 + a^4b^4)\sqrt{a^2 - b^2}} + \frac{6Ba^4}{\dots}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^3,x, algorithm="giac")
```

output $((6B*a^5*b - 12A*a^4*b^2 - 5B*a^3*b^3 + 15A*a^2*b^4 + 2B*a*b^5 - 6A*b^6)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^8 - 2*a^6*b^2 + a^4*b^4)*sqrt(a^2 - b^2)) + (6B*a^4*b^2*tan(1/2*d*x + 1/2*c)^3 - 8A*a^3*b^3*tan(1/2*d*x + 1/2*c)^3 - 5B*a^3*b^3*tan(1/2*d*x + 1/2*c)^3 + 7A*a^2*b^4*tan(1/2*d*x + 1/2*c)^3 - 3B*a^2*b^4*tan(1/2*d*x + 1/2*c)^3 + 5A*a*b^5*tan(1/2*d*x + 1/2*c)^3 + 2B*a*b^5*tan(1/2*d*x + 1/2*c)^3 - 4A*b^6*tan(1/2*d*x + 1/2*c)^3 + 6B*a^4*b^2*tan(1/2*d*x + 1/2*c) - 8A*a^3*b^3*tan(1/2*d*x + 1/2*c) + 5B*a^3*b^3*tan(1/2*d*x + 1/2*c) - 7A*a^2*b^4*tan(1/2*d*x + 1/2*c) - 3B*a^2*b^4*tan(1/2*d*x + 1/2*c) + 5A*a*b^5*tan(1/2*d*x + 1/2*c) - 2B*a*b^5*tan(1/2*d*x + 1/2*c) + 4A*b^6*tan(1/2*d*x + 1/2*c))/((a^7 - 2*a^5*b^2 + a^3*b^4)*(a*tan(1/2*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^2) + (B*a - 3A*b)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^4 - (B*a - 3A*b)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^4 - 2A*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 - 1)*a^3))/d$

3.271.9 Mupad [B] (verification not implemented)

Time = 13.45 (sec) , antiderivative size = 9312, normalized size of antiderivative = 31.14

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^3),x)`

output

$$\begin{aligned} & ((\tan(c/2 + (d*x)/2)^5*(6*A*b^5 - 2*A*a^5 - 12*A*a^2*b^3 + 4*A*a^3*b^2 + B \\ & *a^2*b^3 + 6*B*a^3*b^2 - 3*A*a*b^4 + 2*A*a^4*b - 2*B*a*b^4))/((a^3*b - a^4 \\ &)*(a + b)^2) + (\tan(c/2 + (d*x)/2)*(2*A*a^5 + 6*A*b^5 - 12*A*a^2*b^3 - 4*A \\ & *a^3*b^2 - B*a^2*b^3 + 6*B*a^3*b^2 + 3*A*a*b^4 + 2*A*a^4*b - 2*B*a*b^4))/((\\ & (a + b)*(a^5 - 2*a^4*b + a^3*b^2)) - (2*\tan(c/2 + (d*x)/2)^3*(2*A*a^6 - 6* \\ & A*b^6 + 13*A*a^2*b^4 - 6*A*a^4*b^2 - 5*B*a^3*b^3 + 2*B*a*b^5)))/(a*(a^2*b - \\ & a^3)*(a + b)^2*(a - b)))/(d*(2*a*b - \tan(c/2 + (d*x)/2)^2*(2*a*b - a^2 + \\ & 3*b^2) - \tan(c/2 + (d*x)/2)^6*(a^2 - 2*a*b + b^2) + a^2 + b^2 - \tan(c/2 + \\ & (d*x)/2)^4*(2*a*b + a^2 - 3*b^2))) + (\operatorname{atan}((((8*\tan(c/2 + (d*x)/2)*(72*A^ \\ & 2*b^12 + 4*B^2*a^12 - 72*A^2*a*b^11 - 8*B^2*a^11*b - 288*A^2*a^2*b^10 + 28 \\ & 8*A^2*a^3*b^9 + 441*A^2*a^4*b^8 - 432*A^2*a^5*b^7 - 288*A^2*a^6*b^6 + 288* \\ & A^2*a^7*b^5 + 36*A^2*a^8*b^4 - 72*A^2*a^9*b^3 + 36*A^2*a^10*b^2 + 8*B^2*a^ \\ & 2*b^10 - 8*B^2*a^3*b^9 - 32*B^2*a^4*b^8 + 32*B^2*a^5*b^7 + 57*B^2*a^6*b^6 \\ & - 48*B^2*a^7*b^5 - 52*B^2*a^8*b^4 + 32*B^2*a^9*b^3 + 24*B^2*a^10*b^2 - 48* \\ & A*B*a*b^11 - 24*A*B*a^11*b + 48*A*B*a^2*b^10 + 192*A*B*a^3*b^9 - 192*A*B*a \\ & ^4*b^8 - 318*A*B*a^5*b^7 + 288*A*B*a^6*b^6 + 252*A*B*a^7*b^5 - 192*A*B*a^8 \\ & *b^4 - 72*A*B*a^9*b^3 + 48*A*B*a^10*b^2)))/(a^12*b + a^13 - a^6*b^7 - a^7*b \\ & ^6 + 3*a^8*b^5 + 3*a^9*b^4 - 3*a^10*b^3 - 3*a^11*b^2) + (((8*(4*B*a^18 + 1 \\ & 2*A*a^8*b^10 - 6*A*a^9*b^9 - 54*A*a^10*b^8 + 24*A*a^11*b^7 + 96*A*a^12*b^6 \\ & - 42*A*a^13*b^5 - 78*A*a^14*b^4 + 36*A*a^15*b^3 + 24*A*a^16*b^2 - 4*B*... \end{aligned}$$

3.272 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^3} dx$

3.272.1 Optimal result 2511
 3.272.2 Mathematica [A] (verified) 2512
 3.272.3 Rubi [A] (verified) 2513
 3.272.4 Maple [A] (verified) 2518
 3.272.5 Fricas [B] (verification not implemented) 2519
 3.272.6 Sympy [F] 2520
 3.272.7 Maxima [F(-2)] 2521
 3.272.8 Giac [B] (verification not implemented) 2521
 3.272.9 Mupad [B] (verification not implemented) 2522

3.272.1 Optimal result

Integrand size = 31, antiderivative size = 402

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{b^2(20a^4Ab - 29a^2Ab^3 + 12Ab^5 - 12a^5B + 15a^3b^2B - 6ab^4B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^5(a-b)^{5/2}(a+b)^{5/2}d}$$

$$+ \frac{(a^2A + 12Ab^2 - 6abB) \operatorname{arctanh}(\sin(c + dx))}{2a^5d}$$

$$- \frac{(6a^4Ab - 21a^2Ab^3 + 12Ab^5 - 2a^5B + 11a^3b^2B - 6ab^4B) \tan(c + dx)}{2a^4(a^2 - b^2)^2d}$$

$$+ \frac{(a^4A - 10a^2Ab^2 + 6Ab^4 + 6a^3bB - 3ab^3B) \sec(c + dx) \tan(c + dx)}{2a^3(a^2 - b^2)^2d}$$

$$+ \frac{b(Ab - aB) \sec(c + dx) \tan(c + dx)}{2a(a^2 - b^2)d(a + b \cos(c + dx))^2}$$

$$+ \frac{b(7a^2Ab - 4Ab^3 - 5a^3B + 2ab^2B) \sec(c + dx) \tan(c + dx)}{2a^2(a^2 - b^2)^2d(a + b \cos(c + dx))}$$

output
$$-b^2(20Aa^4b-29Aa^2b^3+12Ab^5-12Ba^5+15Ba^3b^2-6Bab^4)\arctan((a-b)^{1/2}\tan(1/2dx+1/2c)/(a+b)^{1/2})/a^5(a-b)^{5/2}/(a+b)^{5/2}/d+1/2(Aa^2+12Ab^2-6Bab)\operatorname{arctanh}(\sin(dx+c))/a^5/d-1/2(6Aa^4b-21Aa^2b^3+12Ab^5-2Ba^5+11Ba^3b^2-6Bab^4)\tan(dx+c)/a^4/(a^2-b^2)^2/d+1/2(Aa^4-10Aa^2b^2+6Ab^4+6Ba^3b-3Bab^3)\sec(dx+c)\tan(dx+c)/a^3/(a^2-b^2)^2/d+1/2b(Ab-Ba)\sec(dx+c)\tan(dx+c)/a/(a^2-b^2)/d/(a+b\cos(dx+c))^2+1/2b(7Aa^2b-4Ab^3-5Ba^3+2Bab^2)\sec(dx+c)\tan(dx+c)/a^2/(a^2-b^2)^2/d/(a+b\cos(dx+c))$$

3.272.2 Mathematica [A] (verified)

Time = 3.00 (sec) , antiderivative size = 507, normalized size of antiderivative = 1.26

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{16b^2(20a^4Ab-29a^2Ab^3+12Ab^5-12a^5B+15a^3b^2B-6ab^4B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} - 8(a^2A + 12Ab^2 - 6abB) \log(\cos$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^3,x]`

output
$$\begin{aligned} & ((16b^2(20a^4Ab-29a^2Ab^3+12Ab^5-12a^5B+15a^3b^2B-6ab^4B)\operatorname{ArcTanh}[\frac{(a-b)\tan[(c+d*x)/2]}{\sqrt{-a^2+b^2}}])/\sqrt{-a^2+b^2})/(-a^2+b^2)^{5/2} \\ & - 8(a^2A+12Ab^2-6abB)\log[\cos[(c+d*x)/2]] - 8(a^2A+12Ab^2-6abB)\log[\cos[(c+d*x)/2]] \\ & + 2a(4a^7A-30a^5Ab^2+68a^3A^3b^4-36a^6b^2B-32a^4b^3B+18a^2b^5B+(-16a^6Ab+14a^4A^3b^3+47a^2A^5b^5-36Ab^7+8a^7B-10a^5b^2B-25a^3b^4B+18ab^6B)\cos[c+d*x] \\ & + 2ab(-11a^4Ab+32a^2A^3b^3-18Ab^5+4a^5B-16a^3b^2B+9ab^4B)\cos[2(c+d*x)] - 6a^4A^3b^3\cos[3(c+d*x)] + 21a^2A^5b^5\cos[3(c+d*x)] \\ & - 12Ab^7\cos[3(c+d*x)] + 2a^5b^2B\cos[3(c+d*x)] - 11a^3b^4B\cos[3(c+d*x)] + 6ab^6B\cos[3(c+d*x)])\sec[c+d*x]\tan[c+d*x]/((a^2-b^2)^2(a+b\cos[c+d*x])^2)/(16a^5d) \end{aligned}$$

3.272.3 Rubi [A] (verified)

Time = 2.80 (sec) , antiderivative size = 426, normalized size of antiderivative = 1.06, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3479, 3042, 3534, 3042, 3534, 27, 3042, 3534, 25, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx$$

↓ 3479

$$\int \frac{(3b(Ab-aB)\cos^2(c+dx)-2a(Ab-aB)\cos(c+dx)+2(Aa^2+bBa-2Ab^2))\sec^3(c+dx)}{(a+b\cos(c+dx))^2} dx +$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB)\tan(c+dx)\sec(c+dx)} + \frac{2ad(a^2-b^2)(a+b\cos(c+dx))^2}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3042

$$\int \frac{3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2-2a(Ab-aB)\sin(c+dx+\frac{\pi}{2})+2(Aa^2+bBa-2Ab^2)}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx +$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB)\tan(c+dx)\sec(c+dx)} + \frac{2ad(a^2-b^2)(a+b\cos(c+dx))^2}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3534

$$\int \frac{(2b(-5Ba^3+7Ab a^2+2b^2Ba-4Ab^3)\cos^2(c+dx)-a(-2Ba^3+4Aba^2-b^2Ba-Ab^3)\cos(c+dx)+2(Aa^4+6bBa^3-10Ab^2a^2-3b^3Ba+6Ab^4))\sec^3(c+dx)}{a+b\cos(c+dx)} dx + b($$

$$\frac{2a(a^2-b^2)}{b(Ab-aB)\tan(c+dx)\sec(c+dx)} + \frac{2ad(a^2-b^2)(a+b\cos(c+dx))^2}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3042

3.272. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+b\cos(c+dx))^3} dx$

$$\int \frac{2b(-5Ba^3+7Aba^2+2b^2Ba-4Ab^3) \sin(c+dx+\frac{\pi}{2})^2 - a(-2Ba^3+4Aba^2-b^2Ba-Ab^3) \sin(c+dx+\frac{\pi}{2}) + 2(Aa^4+6bBa^3-10Ab^2a^2-3b^3Ba+6Ab^4) dx}{\sin(c+dx+\frac{\pi}{2})^3 (a+b \sin(c+dx+\frac{\pi}{2}))} + \frac{b(-5a^3}{a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \quad 2a(a^2 - b^2)$$

↓ 3534

$$\int \frac{2(-2Ba^5+6Aba^4+11b^2Ba^3-21Ab^3a^2-6b^4Ba-(Aa^4-4bBa^3+4Ab^2a^2+b^3Ba-2Ab^4) \cos(c+dx)a+12Ab^5-b(Aa^4+6bBa^3-10Ab^2a^2-3b^3Ba+6Ab^4) \cos^2(c+dx))}{a+b \cos(c+dx)} \frac{1}{2a}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \quad 2a(a^2 - b^2)$$

↓ 27

$$\frac{(a^4A+6a^3bB-10a^2Ab^2-3ab^3B+6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \int \frac{(-2Ba^5+6Aba^4+11b^2Ba^3-21Ab^3a^2-6b^4Ba-(Aa^4-4bBa^3+4Ab^2a^2+b^3Ba-2Ab^4) \cos(c+dx)a+b \cos(c+dx)) \cos(c+dx)}{a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \quad 2a(a^2 - b^2)$$

↓ 3042

$$\frac{(a^4A+6a^3bB-10a^2Ab^2-3ab^3B+6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \int \frac{-2Ba^5+6Aba^4+11b^2Ba^3-21Ab^3a^2-6b^4Ba-(Aa^4-4bBa^3+4Ab^2a^2+b^3Ba-2Ab^4) \sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^2 (a+b \sin(c+dx+\frac{\pi}{2}))} \frac{1}{a}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \quad 2a(a^2 - b^2)$$

↓ 3534

$$\frac{(a^4A+6a^3bB-10a^2Ab^2-3ab^3B+6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \int \frac{((Aa^2-6bBa+12Ab^2)(a^2-b^2)^2+ab(Aa^4+6bBa^3-10Ab^2a^2-3b^3Ba+6Ab^4) \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} \frac{1}{a}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} \quad 2a(a^2 - b^2)$$

↓ 25

3.272. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^3} dx$

$$\frac{\frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx)}{ad} - \int \frac{((Aa^2 - 6bBa + 12Ab^2)(a^2 - b^2) \tan(c+dx) \sec(c+dx))}{a(a^2 - b^2)} dx}{2a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\frac{\frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx)}{ad} - \int \frac{(Aa^2 - 6bBa + 12Ab^2)(a^2 - b^2) \tan(c+dx) \sec(c+dx)}{a(a^2 - b^2)} dx}{2a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3480

$$\frac{\frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)^2 (a^2 A - 6abB + 12Ab^2) \tan(c+dx) \sec(c+dx)}{a(a^2 - b^2)}}{2a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\frac{\frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)^2 (a^2 A - 6abB + 12Ab^2) \tan(c+dx) \sec(c+dx)}{a(a^2 - b^2)}}{2a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3138

$$\frac{\frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)^2 (a^2 A - 6abB + 12Ab^2) \tan(c+dx) \sec(c+dx)}{a(a^2 - b^2)}}{2a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 218

3.272. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^3} dx$

$$\frac{\frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx)}{ad} - \frac{(a^2 - b^2)^2 (a^2 A - 6abB + 12Ab^2)}{a^3}}{a(a^2 - b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 4257

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} +$$

$$\frac{b(-5a^3 B + 7a^2 Ab + 2ab^2 B - 4Ab^3) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))} + \frac{(a^4 A + 6a^3 bB - 10a^2 Ab^2 - 3ab^3 B + 6Ab^4) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-2a^5 B + 6a^4 Ab + 11a^3 b^2 B - 21a^2 Ab^3 - 6ab^4 B + 12Ab^5) \tan(c+dx) \sec(c+dx)}{a^3}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^3,x]
```

```
output (b*(A*b - a*B)*Sec[c + d*x]*Tan[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((b*(7*a^2*A*b - 4*A*b^3 - 5*a^3*B + 2*a*b^2*B)*Sec[c + d*x]*Tan[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + (((a^4*A - 10*a^2*A*b^2 + 6*A*b^4 + 6*a^3*b*B - 3*a*b^3*B)*Sec[c + d*x]*Tan[c + d*x])/(a*d) - (((-2*b^2*(20*a^4*A*b - 29*a^2*A*b^3 + 12*A*b^5 - 12*a^5*B + 15*a^3*b^2*B - 6*a*b^4*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/Sqrt[a - b]*Sqrt[a + b]*d) + ((a^2 - b^2)^2*(a^2*A + 12*A*b^2 - 6*a*b*B)*ArcTanh[Sin[c + d*x]])/(a*d))/a + ((6*a^4*A*b - 21*a^2*A*b^3 + 12*A*b^5 - 2*a^5*B + 11*a^3*b^2*B - 6*a*b^4*B)*Tan[c + d*x])/(a*d))/a/(a*(a^2 - b^2)))/(2*a*(a^2 - b^2))
```

3.272.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

$$3.272. \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^3} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.272.4 Maple [A] (verified)

Time = 3.04 (sec) , antiderivative size = 460, normalized size of antiderivative = 1.14

method	result
derivativedivides	$-\frac{A}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{-aA - 6Ab + 2Ba}{2a^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(Aa^2 + 12Ab^2 - 6Bab) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a^5} + \frac{A}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{-aA}{2a^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$
default	$-\frac{A}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{-aA - 6Ab + 2Ba}{2a^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(Aa^2 + 12Ab^2 - 6Bab) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a^5} + \frac{A}{2a^3 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^2} - \frac{-aA}{2a^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$
risch	Expression too large to display

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^3,x,method=_RETURNVERBO
SE)
```

$$3.272. \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^3} dx$$

output
$$\frac{1}{d} \left(-\frac{1}{2} \frac{A}{a^3} \frac{1}{\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1} - \frac{1}{2} \frac{(-Aa - 6Ab + 2Ba)}{a^4} \frac{1}{\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1} + \frac{1}{2} \frac{(Aa^2 + 12Ab^2 - 6Bab)}{a^5 \ln(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1)} + \frac{1}{2} \frac{A}{a^3} \frac{1}{\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1} - \frac{1}{2} \frac{(-Aa - 6Ab + 2Ba)}{a^4} \frac{1}{\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1} + \frac{1}{2} \frac{1}{a^5} \frac{(-Aa^2 - 12Ab^2 + 6Bab) \ln(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1)}{-2b^2 a^5 \left((-\frac{1}{2} (10Aa^2b + Aab^2 - 6Ab^3 - 8Ba^3 - Ba^2b + 4Bab^2)) \frac{ab}{(a-b)} \right) / (a^2 + 2ab + b^2) \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - \frac{1}{2} b^2 a^2 (10Aa^2b - Aab^2 - 6Ab^3 - 8Ba^3 + Ba^2b + 4Bab^2) / (a+b) / (a-b)^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)}{\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 a - b \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + (a+b)^2} + \frac{1}{2} \frac{(20Aa^4b - 29Aa^2b^3 + 12Ab^5 - 12Ba^5 + 15Bab^3b^2 - 6Bab^4)}{(a^4 - 2a^2b^2 + b^4)} / ((a-b)(a+b))^{1/2} \arctan\left(\frac{(a-b)\tan(\frac{1}{2}dx + \frac{1}{2}c)}{(a-b)(a+b)^{1/2}}\right) \right)$$

3.272.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1174 vs. $2(382) = 764$.

Time = 43.61 (sec) , antiderivative size = 2416, normalized size of antiderivative = 6.01

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^3,x, algorithm="fracas")`

output `[1/4*(((12*B*a^5*b^4 - 20*A*a^4*b^5 - 15*B*a^3*b^6 + 29*A*a^2*b^7 + 6*B*a*b^8 - 12*A*b^9)*cos(d*x + c)^4 + 2*(12*B*a^6*b^3 - 20*A*a^5*b^4 - 15*B*a^4*b^5 + 29*A*a^3*b^6 + 6*B*a^2*b^7 - 12*A*a*b^8)*cos(d*x + c)^3 + (12*B*a^7*b^2 - 20*A*a^6*b^3 - 15*B*a^5*b^4 + 29*A*a^4*b^5 + 6*B*a^3*b^6 - 12*A*a^2*b^7)*cos(d*x + c)^2)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + ((A*a^8*b^2 - 6*B*a^7*b^3 + 9*A*a^6*b^4 + 18*B*a^5*b^5 - 33*A*a^4*b^6 - 18*B*a^3*b^7 + 35*A*a^2*b^8 + 6*B*a*b^9 - 12*A*b^10)*cos(d*x + c)^4 + 2*(A*a^9*b - 6*B*a^8*b^2 + 9*A*a^7*b^3 + 18*B*a^6*b^4 - 33*A*a^5*b^5 - 18*B*a^4*b^6 + 35*A*a^3*b^7 + 6*B*a^2*b^8 - 12*A*a*b^9)*cos(d*x + c)^3 + (A*a^10 - 6*B*a^9*b + 9*A*a^8*b^2 + 18*B*a^7*b^3 - 33*A*a^6*b^4 - 18*B*a^5*b^5 + 35*A*a^4*b^6 + 6*B*a^3*b^7 - 12*A*a^2*b^8)*cos(d*x + c)^2)*log(sin(d*x + c) + 1) - ((A*a^8*b^2 - 6*B*a^7*b^3 + 9*A*a^6*b^4 + 18*B*a^5*b^5 - 33*A*a^4*b^6 - 18*B*a^3*b^7 + 35*A*a^2*b^8 + 6*B*a*b^9 - 12*A*b^10)*cos(d*x + c)^4 + 2*(A*a^9*b - 6*B*a^8*b^2 + 9*A*a^7*b^3 + 18*B*a^6*b^4 - 33*A*a^5*b^5 - 18*B*a^4*b^6 + 35*A*a^3*b^7 + 6*B*a^2*b^8 - 12*A*a*b^9)*cos(d*x + c)^3 + (A*a^10 - 6*B*a^9*b + 9*A*a^8*b^2 + 18*B*a^7*b^3 - 33*A*a^6*b^4 - 18*B*a^5*b^5 + 35*A*a^4*b^6 + 6*B*a^3*b^7 - 12*A*a^2*b^8)*cos(d*x + c)^2)*log(-sin(d*x + c) + 1) + 2*(A*a^10 - 3*A*a^8*b^2 + 3*A*a^6*b^4 - A*a^4*b^6 + (2*B*a^8*b^2 ...`

3.272.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**3,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a + b*cos(c + d*x))**3, x)`

3.272.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.272.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1395 vs. 2(382) = 764.

Time = 0.37 (sec) , antiderivative size = 1395, normalized size of antiderivative = 3.47

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output

```

-1/2*(2*(12*B*a^5*b^2 - 20*A*a^4*b^3 - 15*B*a^3*b^4 + 29*A*a^2*b^5 + 6*B*a
*b^6 - 12*A*b^7)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arcta
n(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a
^9 - 2*a^7*b^2 + a^5*b^4)*sqrt(a^2 - b^2)) - 2*(A*a^7*tan(1/2*d*x + 1/2*c)
^7 - 2*B*a^7*tan(1/2*d*x + 1/2*c)^7 + 4*A*a^6*b*tan(1/2*d*x + 1/2*c)^7 + 4
*B*a^6*b*tan(1/2*d*x + 1/2*c)^7 - 13*A*a^5*b^2*tan(1/2*d*x + 1/2*c)^7 + 2*
B*a^5*b^2*tan(1/2*d*x + 1/2*c)^7 - 2*A*a^4*b^3*tan(1/2*d*x + 1/2*c)^7 - 16
*B*a^4*b^3*tan(1/2*d*x + 1/2*c)^7 + 33*A*a^3*b^4*tan(1/2*d*x + 1/2*c)^7 +
9*B*a^3*b^4*tan(1/2*d*x + 1/2*c)^7 - 17*A*a^2*b^5*tan(1/2*d*x + 1/2*c)^7 +
9*B*a^2*b^5*tan(1/2*d*x + 1/2*c)^7 - 18*A*a*b^6*tan(1/2*d*x + 1/2*c)^7 -
6*B*a*b^6*tan(1/2*d*x + 1/2*c)^7 + 12*A*b^7*tan(1/2*d*x + 1/2*c)^7 + 3*A*a
^7*tan(1/2*d*x + 1/2*c)^5 - 2*B*a^7*tan(1/2*d*x + 1/2*c)^5 + 4*A*a^6*b*tan
(1/2*d*x + 1/2*c)^5 - 4*B*a^6*b*tan(1/2*d*x + 1/2*c)^5 + 5*A*a^5*b^2*tan(1
/2*d*x + 1/2*c)^5 + 10*B*a^5*b^2*tan(1/2*d*x + 1/2*c)^5 - 26*A*a^4*b^3*tan
(1/2*d*x + 1/2*c)^5 + 16*B*a^4*b^3*tan(1/2*d*x + 1/2*c)^5 - 29*A*a^3*b^4*t
an(1/2*d*x + 1/2*c)^5 - 35*B*a^3*b^4*tan(1/2*d*x + 1/2*c)^5 + 67*A*a^2*b^5
*tan(1/2*d*x + 1/2*c)^5 - 9*B*a^2*b^5*tan(1/2*d*x + 1/2*c)^5 + 18*A*a*b^6*
tan(1/2*d*x + 1/2*c)^5 + 18*B*a*b^6*tan(1/2*d*x + 1/2*c)^5 - 36*A*b^7*tan(
1/2*d*x + 1/2*c)^5 + 3*A*a^7*tan(1/2*d*x + 1/2*c)^3 + 2*B*a^7*tan(1/2*d*x
+ 1/2*c)^3 - 4*A*a^6*b*tan(1/2*d*x + 1/2*c)^3 - 4*B*a^6*b*tan(1/2*d*x + ...

```

3.272.9 Mupad [B] (verification not implemented)

Time = 12.72 (sec) , antiderivative size = 10547, normalized size of antiderivative = 26.24

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^3),x)`

output

$$\begin{aligned}
& ((\tan(c/2 + (d*x)/2)^3*(3*A*a^7 + 36*A*b^7 + 2*B*a^7 - 67*A*a^2*b^5 - 29*A \\
& *a^3*b^4 + 26*A*a^4*b^3 + 5*A*a^5*b^2 - 9*B*a^2*b^5 + 35*B*a^3*b^4 + 16*B* \\
& a^4*b^3 - 10*B*a^5*b^2 + 18*A*a*b^6 - 4*A*a^6*b - 18*B*a*b^6 - 4*B*a^6*b)) \\
& /((a + b)^2*(a^6 - 2*a^5*b + a^4*b^2)) + (\tan(c/2 + (d*x)/2)^5*(3*A*a^7 - \\
& 36*A*b^7 - 2*B*a^7 + 67*A*a^2*b^5 - 29*A*a^3*b^4 - 26*A*a^4*b^3 + 5*A*a^5* \\
& b^2 - 9*B*a^2*b^5 - 35*B*a^3*b^4 + 16*B*a^4*b^3 + 10*B*a^5*b^2 + 18*A*a*b^6 \\
& + 4*A*a^6*b + 18*B*a*b^6 - 4*B*a^6*b))/((a + b)^2*(a^6 - 2*a^5*b + a^4*b^2)) - (\tan(c/2 + (d*x)/2)^7*(A*a^6 - 12*A*b^6 - 2*B*a^6 + 23*A*a^2*b^4 - \\
& 10*A*a^3*b^3 - 8*A*a^4*b^2 - 3*B*a^2*b^4 - 12*B*a^3*b^3 + 4*B*a^4*b^2 + 6* \\
& A*a*b^5 + 5*A*a^5*b + 6*B*a*b^5 + 2*B*a^5*b))/((a^4*b - a^5)*(a + b)^2) + \\
& (\tan(c/2 + (d*x)/2)*(A*a^6 - 12*A*b^6 + 2*B*a^6 + 23*A*a^2*b^4 + 10*A*a^3* \\
& b^3 - 8*A*a^4*b^2 + 3*B*a^2*b^4 - 12*B*a^3*b^3 - 4*B*a^4*b^2 - 6*A*a*b^5 - \\
& 5*A*a^5*b + 6*B*a*b^5 + 2*B*a^5*b))/((a + b)*(a^6 - 2*a^5*b + a^4*b^2)))/ \\
& (d*(2*a*b - \tan(c/2 + (d*x)/2)^4*(2*a^2 - 6*b^2) - \tan(c/2 + (d*x)/2)^2*(4 \\
& *a*b + 4*b^2) + \tan(c/2 + (d*x)/2)^6*(4*a*b - 4*b^2) + \tan(c/2 + (d*x)/2)^ \\
& 8*(a^2 - 2*a*b + b^2) + a^2 + b^2)) - (\operatorname{atan}(\frac{((8*\tan(c/2 + (d*x)/2)*(A^2* \\
& a^{14} + 288*A^2*b^{14} - 288*A^2*a*b^{13} - 2*A^2*a^{13}*b - 1104*A^2*a^2*b^{12} + \\
& 1104*A^2*a^3*b^{11} + 1538*A^2*a^4*b^{10} - 1538*A^2*a^5*b^9 - 827*A^2*a^6*b^8 \\
& + 872*A^2*a^7*b^7 + 18*A^2*a^8*b^6 - 108*A^2*a^9*b^5 + 74*A^2*a^{10}*b^4 - \\
& 40*A^2*a^{11}*b^3 + 21*A^2*a^{12}*b^2 + 72*B^2*a^2*b^{12} - 72*B^2*a^3*b^{11} - \dots
\end{aligned}$$

3.273 $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

3.273.1 Optimal result 2524
 3.273.2 Mathematica [B] (verified) 2525
 3.273.3 Rubi [A] (verified) 2526
 3.273.4 Maple [A] (verified) 2531
 3.273.5 Fracas [B] (verification not implemented) 2532
 3.273.6 Sympy [F(-1)] 2533
 3.273.7 Maxima [F(-2)] 2534
 3.273.8 Giac [B] (verification not implemented) 2534
 3.273.9 Mupad [B] (verification not implemented) 2535

3.273.1 Optimal result

Integrand size = 31, antiderivative size = 409

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx = \frac{(Ab-4aB)x}{b^5} - \frac{a(2a^6Ab-7a^4Ab^3+8a^2Ab^5-8Ab^7-8a^7B+28a^5b^2B-35a^3b^4B+20ab^6B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - (3a^3Ab-8aAb^3-12a^4B+23a^2b^2B-6b^4B) \sin(c+dx)}{(a-b)^{7/2}b^5(a+b)^{7/2}d} - \frac{6b^4(a^2-b^2)^2d}{(3a^3Ab-8aAb^3-12a^4B+23a^2b^2B-6b^4B) \sin(c+dx)} + \frac{a(Ab-aB) \cos^3(c+dx) \sin(c+dx)}{3b(a^2-b^2)d(a+b \cos(c+dx))^3} + \frac{a(a^2Ab-6Ab^3-4a^3B+9ab^2B) \cos^2(c+dx) \sin(c+dx)}{6b^2(a^2-b^2)^2d(a+b \cos(c+dx))^2} - \frac{a^2(a^4Ab-2a^2Ab^3+6Ab^5-4a^5B+11a^3b^2B-12ab^4B) \sin(c+dx)}{2b^4(a^2-b^2)^3d(a+b \cos(c+dx))}$$

```
output (A*b-4*B*a)*x/b^5-a*(2*A*a^6*b-7*A*a^4*b^3+8*A*a^2*b^5-8*A*b^7-8*B*a^7+28*
B*a^5*b^2-35*B*a^3*b^4+20*B*a*b^6)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(
a+b)^(1/2))/(a-b)^(7/2)/b^5/(a+b)^(7/2)/d-1/6*(3*A*a^3*b-8*A*a*b^3-12*B*a^
4+23*B*a^2*b^2-6*B*b^4)*sin(d*x+c)/b^4/(a^2-b^2)^2/d+1/3*a*(A*b-B*a)*cos(d
*x+c)^3*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^3+1/6*a*(A*a^2*b-6*A*b^3
-4*B*a^3+9*B*a*b^2)*cos(d*x+c)^2*sin(d*x+c)/b^2/(a^2-b^2)^2/d/(a+b*cos(d*x
+c))^2-1/2*a^2*(A*a^4*b-2*A*a^2*b^3+6*A*b^5-4*B*a^5+11*B*a^3*b^2-12*B*a*b^
4)*sin(d*x+c)/b^4/(a^2-b^2)^3/d/(a+b*cos(d*x+c))
```

3.273.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1155 vs. 2(409) = 818.

Time = 7.67 (sec) , antiderivative size = 1155, normalized size of antiderivative = 2.82

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{24a(-2a^6Ab + 7a^4Ab^3 - 8a^2Ab^5 + 8Ab^7 + 8a^7B - 28a^5b^2B + 35a^3b^4B - 20ab^6B) \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right) - 24a^9Abc + 36a^7Ab^3c + 36a^5Ab^5c - 84a^3Ab^7c + 36a^2Ab^9c + 96a^{10}Bc - 144a^8b^2Bc - 144a^6b^4Bc + 336a^4b^6Bc - 144a^2b^8Bc - 24a^9Ab^2dx + 36a^7Ab^4dx + 36a^5Ab^6dx - 84a^3Ab^8dx + 36a^2Ab^{10}dx + 96a^{10}Bdx - 144a^8b^2Bdx - 144a^6b^4Bdx + 336a^4b^6Bdx - 144a^2b^8Bdx + 18b(-a^2 + b^2)^3(4a^2 + b^2)(Ab - 4aB)(c + dx)\cos[c + dx] + 36a^2b^2(a^2 - b^2)^3(-Ab + 4aB)(c + dx)\cos[2(c + dx)] - 6a^6Ab^4c\cos[3(c + dx)] + 18a^4Ab^6c\cos[3(c + dx)] - 18a^2Ab^8c\cos[3(c + dx)] + 6Ab^{10}c\cos[3(c + dx)] + 24a^7b^3Bc\cos[3(c + dx)] - 72a^5b^5Bc\cos[3(c + dx)] + 72a^3b^7Bc\cos[3(c + dx)] - 24a^2b^9Bc\cos[3(c + dx)] - 6a^6Ab^4dx\cos[3(c + dx)] + 18a^4Ab^6dx\cos[3(c + dx)] - 18a^2Ab^8dx\cos[3(c + dx)] + 6Ab^{10}dx\cos[3(c + dx)] + 24a^7b^3Bdx\cos[3(c + dx)] - 72a^5b^5Bdx\cos[3(c + dx)] + 72a^3b^7Bdx\cos[3(c + dx)] - 24a^2b^9Bdx\cos[3(c + dx)] + 24a^8Ab^2\sin[c + dx] - 57a^6Ab^4\sin[c + dx] + 72a^4Ab^6\sin[c + dx] + 36a^2Ab^8\sin[c + dx] - 96a^9bB\sin[c + dx] + 228a^7b^3B\sin[c + dx] - 135a^5b^5B\sin[c + dx] - 90a^3b^7B\sin[c + dx] + 18a^2b^9B\sin[c + dx] \dots}{(-a^2+b^2)^{7/2}}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output `((24*a*(-2*a^6*A*b + 7*a^4*A*b^3 - 8*a^2*A*b^5 + 8*A*b^7 + 8*a^7*B - 28*a^5*b^2*B + 35*a^3*b^4*B - 20*a*b^6*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(7/2) - (-24*a^9*A*b*c + 36*a^7*A*b^3*c + 36*a^5*A*b^5*c - 84*a^3*A*b^7*c + 36*a^2*A*b^9*c + 96*a^10*B*c - 144*a^8*b^2*B*c - 144*a^6*b^4*B*c + 336*a^4*b^6*B*c - 144*a^2*b^8*B*c - 24*a^9*A*b^2*d*x + 36*a^7*A*b^4*d*x + 36*a^5*A*b^6*d*x - 84*a^3*A*b^8*d*x + 36*a^2*A*b^10*d*x + 96*a^10*B*d*x - 144*a^8*b^2*B*d*x - 144*a^6*b^4*B*d*x + 336*a^4*b^6*B*d*x - 144*a^2*b^8*B*d*x + 18*b*(-a^2 + b^2)^3*(4*a^2 + b^2)*(A*b - 4*a*B)*(c + d*x)*Cos[c + d*x] + 36*a^2*b^2*(a^2 - b^2)^3*(-A*b + 4*a*B)*(c + d*x)*Cos[2*(c + d*x)] - 6*a^6*A*b^4*c*Cos[3*(c + d*x)] + 18*a^4*A*b^6*c*Cos[3*(c + d*x)] - 18*a^2*A*b^8*c*Cos[3*(c + d*x)] + 6*A*b^10*c*Cos[3*(c + d*x)] + 24*a^7*b^3*B*c*Cos[3*(c + d*x)] - 72*a^5*b^5*B*c*Cos[3*(c + d*x)] + 72*a^3*b^7*B*c*Cos[3*(c + d*x)] - 24*a^2*b^9*B*c*Cos[3*(c + d*x)] - 6*a^6*A*b^4*d*x*Cos[3*(c + d*x)] + 18*a^4*A*b^6*d*x*Cos[3*(c + d*x)] - 18*a^2*A*b^8*d*x*Cos[3*(c + d*x)] + 6*A*b^10*d*x*Cos[3*(c + d*x)] + 24*a^7*b^3*B*d*x*Cos[3*(c + d*x)] - 72*a^5*b^5*B*d*x*Cos[3*(c + d*x)] + 72*a^3*b^7*B*d*x*Cos[3*(c + d*x)] - 24*a^2*b^9*B*d*x*Cos[3*(c + d*x)] + 24*a^8*A*b^2*Sin[c + d*x] - 57*a^6*A*b^4*Sin[c + d*x] + 72*a^4*A*b^6*Sin[c + d*x] + 36*a^2*A*b^8*Sin[c + d*x] - 96*a^9*b*B*Sin[c + d*x] + 228*a^7*b^3*B*Sin[c + d*x] - 135*a^5*b^5*B*Sin[c + d*x] - 90*a^3*b^7*B*Sin[c + d*x] + 18*a^2*b^9*B*Sin[c + d*x]...`

3.273.3 Rubi [A] (verified)

Time = 2.18 (sec) , antiderivative size = 466, normalized size of antiderivative = 1.14, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 3468, 25, 3042, 3526, 25, 3042, 3510, 3042, 3502, 27, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^4(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^4} dx$$

$$\downarrow \text{3468}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3} - \int \frac{\cos^2(c+dx)(-((-4Ba^2+Aba+3b^2B)\cos^2(c+dx))-3b(Ab-aB)\cos(c+dx)+3a(Ab-aB))}{(a+b\cos(c+dx))^3} dx}{3b(a^2-b^2)}$$

$$\downarrow \text{25}$$

$$\int \frac{\cos^2(c+dx)(-((-4Ba^2+Aba+3b^2B)\cos^2(c+dx))-3b(Ab-aB)\cos(c+dx)+3a(Ab-aB))}{(a+b\cos(c+dx))^3} dx}{3b(a^2-b^2)} + \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^2((4Ba^2-Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+3a(Ab-aB))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx}{3b(a^2-b^2)} + \frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

$$\downarrow \text{3526}$$

$$\frac{a(-4a^3B+a^2Ab+9ab^2B-6Ab^3)\sin(c+dx)\cos^2(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} - \int \frac{\cos(c+dx)(-((-12Ba^4+3Aba^3+23b^2Ba^2-8Ab^3a-6b^4B)\cos^2(c+dx))+2b(Ba^3+2Aba^2-2Ab^3-3b^4))}{(a+b\cos(c+dx))^2}}{3b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

3.273. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

↓ 25

$$\int \frac{\cos(c+dx) \left(- \left((-12Ba^4 + 3Aba^3 + 23b^2Ba^2 - 8Ab^3a - 6b^4B) \cos^2(c+dx) \right) + 2b \left(Ba^3 + 2Aba^2 - 6b^2Ba + 3Ab^3 \right) \cos(c+dx) + 2a \left(-4Ba^3 + Aba^2 + 9b^2Ba - 6Ab^3 \right) \right)}{\frac{(a+b \cos(c+dx))^2}{2b(a^2-b^2)}} dx$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} \quad 3b(a^2 - b^2)$$

↓ 3042

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right) \left((12Ba^4 - 3Aba^3 - 23b^2Ba^2 + 8Ab^3a + 6b^4B) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + 2b \left(Ba^3 + 2Aba^2 - 6b^2Ba + 3Ab^3 \right) \sin\left(c+dx+\frac{\pi}{2}\right) + 2a \left(-4Ba^3 + Aba^2 + 9b^2Ba - 6Ab^3 \right) \right)}{\frac{(a+b \sin\left(c+dx+\frac{\pi}{2}\right))^2}{2b(a^2-b^2)}} dx$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} \quad 3b(a^2 - b^2)$$

↓ 3510

$$\int \frac{-b(a^2-b^2) \left((-12Ba^4 + 3Aba^3 + 23b^2Ba^2 - 8Ab^3a - 6b^4B) \cos^2(c+dx) \right) + (a^2-b^2) \left(-12Ba^5 + 3Aba^4 + 25b^2Ba^3 - 4Ab^3a^2 - 18b^4Ba + 6Ab^5 \right) \cos(c+dx) + 3ab \left(-4Ba^5 + 3Aba^4 + 25b^2Ba^3 - 4Ab^3a^2 - 18b^4Ba + 6Ab^5 \right) \sin(c+dx)}{\frac{a+b \cos(c+dx)}{b^2(a^2-b^2)}} dx$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} \quad 2b(a^2 - b^2)$$

↓ 3042

$$\int \frac{-b(a^2-b^2) \left((-12Ba^4 + 3Aba^3 + 23b^2Ba^2 - 8Ab^3a - 6b^4B) \sin\left(c+dx+\frac{\pi}{2}\right)^2 \right) + (a^2-b^2) \left(-12Ba^5 + 3Aba^4 + 25b^2Ba^3 - 4Ab^3a^2 - 18b^4Ba + 6Ab^5 \right) \sin\left(c+dx+\frac{\pi}{2}\right) + 3ab \left(-4Ba^5 + 3Aba^4 + 25b^2Ba^3 - 4Ab^3a^2 - 18b^4Ba + 6Ab^5 \right) \cos\left(c+dx+\frac{\pi}{2}\right)}{\frac{a+b \sin\left(c+dx+\frac{\pi}{2}\right)}{b^2(a^2-b^2)}} dx$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} \quad 2b(a^2 - b^2)$$

↓ 3502

$$\int \frac{3 \left(2b(Ab - 4aB) \cos(c+dx) (a^2 - b^2)^3 + ab^2 \left(-4Ba^5 + Aba^4 + 11b^2Ba^3 - 2Ab^3a^2 - 12b^4Ba + 6Ab^5 \right) \right)}{\frac{a+b \cos(c+dx)}{b}} dx - \frac{(a^2 - b^2) \left(-12a^4B + 3a^3Ab + 23a^2b^2B - 8aAb^3 - 6b^4B \right) \sin(c+dx)}{d}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} \quad 2b(a^2 - b^2)$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} \quad 3b(a^2 - b^2)$$

3.273. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

↓ 27

$$\frac{3 \int \frac{2b(Ab-4aB) \cos(c+dx) (a^2-b^2)^3 + ab^2(-4Ba^5+Ab^4+11b^2Ba^3-2Ab^3a^2-12b^4Ba+6Ab^5)}{a+b \cos(c+dx)} dx - \frac{(a^2-b^2)(-12a^4B+3a^3Ab+23a^2b^2B-8aAb^3-6b^4B) \sin(c+dx)}{d}}{b^2(a^2-b^2)} = \frac{2b(a^2-b^2)}{3b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{3bd(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3042

$$\frac{3 \int \frac{2b(Ab-4aB) \sin(c+dx+\frac{\pi}{2}) (a^2-b^2)^3 + ab^2(-4Ba^5+Ab^4+11b^2Ba^3-2Ab^3a^2-12b^4Ba+6Ab^5)}{a+b \sin(c+dx+\frac{\pi}{2})} dx - \frac{(a^2-b^2)(-12a^4B+3a^3Ab+23a^2b^2B-8aAb^3-6b^4B) \sin(c+dx+\frac{\pi}{2})}{d}}{b^2(a^2-b^2)} = \frac{2b(a^2-b^2)}{3b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{3bd(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3214

$$\frac{3 \left(2x(a^2-b^2)^3(Ab-4aB) - a(-8a^7B+2a^6Ab+28a^5b^2B-7a^4Ab^3-35a^3b^4B+8a^2Ab^5+20ab^6B-8Ab^7) \int \frac{1}{a+b \cos(c+dx)} dx \right) - \frac{(a^2-b^2)(-12a^4B+3a^3Ab+23a^2b^2B-8aAb^3-6b^4B) \sin(c+dx)}{d}}{b^2(a^2-b^2)} = \frac{2b(a^2-b^2)}{3b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{3bd(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3042

$$\frac{3 \left(2x(a^2-b^2)^3(Ab-4aB) - a(-8a^7B+2a^6Ab+28a^5b^2B-7a^4Ab^3-35a^3b^4B+8a^2Ab^5+20ab^6B-8Ab^7) \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx \right) - \frac{(a^2-b^2)(-12a^4B+3a^3Ab+23a^2b^2B-8aAb^3-6b^4B) \sin(c+dx+\frac{\pi}{2})}{d}}{b^2(a^2-b^2)} = \frac{2b(a^2-b^2)}{3b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^3(c+dx)}{3bd(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3138

3.273. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

$$3 \left(\frac{2a(-8a^7B+2a^6Ab+28a^5b^2B-7a^4Ab^3-35a^3b^4B+8a^2Ab^5+20ab^6B-8Ab^7)}{2x(a^2-b^2)^3(Ab-4aB)} - \frac{1}{(a-b)\tan^2\left(\frac{1}{2}(c+dx)\right)+a+b} \right) \int \frac{d \tan\left(\frac{1}{2}(c+dx)\right)}{d} - (a^2)$$

$$\frac{a^2}{b^2(a^2-b^2)}$$

$$\frac{2b(a^2-b^2)}{2b(a^2-b^2)}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 218

$$\frac{a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} +$$

$$3 \left(\frac{2a(-8a^7B+2a^6Ab+28a^5b^2B-7a^4Ab^3-35a^3b^4B+8a^2Ab^5+20ab^6)}{2x(a^2-b^2)^3(Ab-4aB)} - \frac{1}{d\sqrt{a-b}\sqrt{a+b}} \right)$$

$$\frac{1}{b}$$

$$\frac{a(-4a^3B+a^2Ab+9ab^2B-6Ab^3) \sin(c+dx) \cos^2(c+dx)}{2bd(a^2-b^2)(a+b \cos(c+dx))^2} +$$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output `(a*(A*b - a*B)*Cos[c + d*x]^3*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^3) + ((a*(a^2*A*b - 6*A*b^3 - 4*a^3*B + 9*a*b^2*B)*Cos[c + d*x]^2*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((-3*a^2*(a^4*A*b - 2*a^2*A*b^3 + 6*A*b^5 - 4*a^5*B + 11*a^3*b^2*B - 12*a*b^4*B)*Sin[c + d*x])/(b^2*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + ((3*(2*(a^2 - b^2)^3*(A*b - 4*a*B)*x - (2*a*(2*a^6*A*b - 7*a^4*A*b^3 + 8*a^2*A*b^5 - 8*A*b^7 - 8*a^7*B + 28*a^5*b^2*B - 35*a^3*b^4*B + 20*a*b^6*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*d))/b - ((a^2 - b^2)*(3*a^3*A*b - 8*a*A*b^3 - 12*a^4*B + 23*a^2*b^2*B - 6*b^4*B)*Sin[c + d*x])/d)/(b^2*(a^2 - b^2)))/(2*b*(a^2 - b^2)))/(3*b*(a^2 - b^2))`

3.273.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

3.273. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^n)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

3.273.4 Maple [A] (verified)

Time = 2.48 (sec) , antiderivative size = 552, normalized size of antiderivative = 1.35

method	result
derivativedivides	$2a \left(\frac{(2Aa^4b - Aa^3b^2 - 6Aa^2b^3 + 4Aab^4 + 12Ab^5 - 6Ba^5 + 2Ba^4b + 18Ba^3b^2 - 5Ba^2b^3 - 20Bab^4)ab \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{2(a-b)(a^3 + 3a^2b + 3ab^2 + b^3)} + \frac{2(3Aa^4b - 12Aa^3b^2 + 12Aa^2b^3 - 4Aab^4 + 12Ab^5 - 6Ba^5 + 2Ba^4b + 18Ba^3b^2 - 5Ba^2b^3 - 20Bab^4)ab \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{2(a-b)(a^3 + 3a^2b + 3ab^2 + b^3)} \right)$
default	$2a \left(\frac{(2Aa^4b - Aa^3b^2 - 6Aa^2b^3 + 4Aab^4 + 12Ab^5 - 6Ba^5 + 2Ba^4b + 18Ba^3b^2 - 5Ba^2b^3 - 20Bab^4)ab \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{2(a-b)(a^3 + 3a^2b + 3ab^2 + b^3)} + \frac{2(3Aa^4b - 12Aa^3b^2 + 12Aa^2b^3 - 4Aab^4 + 12Ab^5 - 6Ba^5 + 2Ba^4b + 18Ba^3b^2 - 5Ba^2b^3 - 20Bab^4)ab \left(\tan^5 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{2(a-b)(a^3 + 3a^2b + 3ab^2 + b^3)} \right)$
risch	Expression too large to display

3.273. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

```
input int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-2*a/b^5*((1/2*(2*A*a^4*b-A*a^3*b^2-6*A*a^2*b^3+4*A*a*b^4+12*A*b^5-6*B*a^5+2*B*a^4*b+18*B*a^3*b^2-5*B*a^2*b^3-20*B*a*b^4)*a*b/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*tan(1/2*d*x+1/2*c)^5+2/3*(3*A*a^4*b-11*A*a^2*b^3+18*A*b^5-9*B*a^5+29*B*a^3*b^2-30*B*a*b^4)*a*b/(a^2+2*a*b+b^2)/(a^2-2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3+1/2*(2*A*a^4*b+A*a^3*b^2-6*A*a^2*b^3-4*A*a*b^4+12*A*b^5-6*B*a^5-2*B*a^4*b+18*B*a^3*b^2+5*B*a^2*b^3-20*B*a*b^4)*a*b/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^3+1/2*(2*A*a^6*b-7*A*a^4*b^3+8*A*a^2*b^5-8*A*b^7-8*B*a^7+28*B*a^5*b^2-35*B*a^3*b^4+20*B*a*b^6)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))+2/b^5*(B*b*tan(1/2*d*x+1/2*c)/(1+tan(1/2*d*x+1/2*c)^2)+(A*b-4*B*a)*arctan(tan(1/2*d*x+1/2*c))))
```

3.273.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1249 vs. $2(395) = 790$.

Time = 0.58 (sec) , antiderivative size = 2567, normalized size of antiderivative = 6.28

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="fracas")
```

output

```

[-1/12*(12*(4*B*a^9*b^3 - A*a^8*b^4 - 16*B*a^7*b^5 + 4*A*a^6*b^6 + 24*B*a^
5*b^7 - 6*A*a^4*b^8 - 16*B*a^3*b^9 + 4*A*a^2*b^10 + 4*B*a*b^11 - A*b^12)*d
*x*cos(d*x + c)^3 + 36*(4*B*a^10*b^2 - A*a^9*b^3 - 16*B*a^8*b^4 + 4*A*a^7*
b^5 + 24*B*a^6*b^6 - 6*A*a^5*b^7 - 16*B*a^4*b^8 + 4*A*a^3*b^9 + 4*B*a^2*b^
10 - A*a*b^11)*d*x*cos(d*x + c)^2 + 36*(4*B*a^11*b - A*a^10*b^2 - 16*B*a^9
*b^3 + 4*A*a^8*b^4 + 24*B*a^7*b^5 - 6*A*a^6*b^6 - 16*B*a^5*b^7 + 4*A*a^4*b
^8 + 4*B*a^3*b^9 - A*a^2*b^10)*d*x*cos(d*x + c) + 12*(4*B*a^12 - A*a^11*b
- 16*B*a^10*b^2 + 4*A*a^9*b^3 + 24*B*a^8*b^4 - 6*A*a^7*b^5 - 16*B*a^6*b^6
+ 4*A*a^5*b^7 + 4*B*a^4*b^8 - A*a^3*b^9)*d*x - 3*(8*B*a^11 - 2*A*a^10*b -
28*B*a^9*b^2 + 7*A*a^8*b^3 + 35*B*a^7*b^4 - 8*A*a^6*b^5 - 20*B*a^5*b^6 + 8
*A*a^4*b^7 + (8*B*a^8*b^3 - 2*A*a^7*b^4 - 28*B*a^6*b^5 + 7*A*a^5*b^6 + 35*
B*a^4*b^7 - 8*A*a^3*b^8 - 20*B*a^2*b^9 + 8*A*a*b^10)*cos(d*x + c)^3 + 3*(8
*B*a^9*b^2 - 2*A*a^8*b^3 - 28*B*a^7*b^4 + 7*A*a^6*b^5 + 35*B*a^5*b^6 - 8*A
*a^4*b^7 - 20*B*a^3*b^8 + 8*A*a^2*b^9)*cos(d*x + c)^2 + 3*(8*B*a^10*b - 2*
A*a^9*b^2 - 28*B*a^8*b^3 + 7*A*a^7*b^4 + 35*B*a^6*b^5 - 8*A*a^5*b^6 - 20*B
*a^4*b^7 + 8*A*a^3*b^8)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x
+ c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) +
b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) +
a^2)) - 2*(24*B*a^11*b - 6*A*a^10*b^2 - 92*B*a^9*b^3 + 23*A*a^8*b^4 + 133
*B*a^7*b^5 - 43*A*a^6*b^6 - 71*B*a^5*b^7 + 26*A*a^4*b^8 + 6*B*a^3*b^9 + ...

```

3.273.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**4,x)`

output `Timed out`

3.273.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.273.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 966 vs. 2(395) = 790.

Time = 0.39 (sec) , antiderivative size = 966, normalized size of antiderivative = 2.36

$$\int \frac{\cos^4(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

```

output -1/3*(3*(8*B*a^8 - 2*A*a^7*b - 28*B*a^6*b^2 + 7*A*a^5*b^3 + 35*B*a^4*b^4 -
      8*A*a^3*b^5 - 20*B*a^2*b^6 + 8*A*a*b^7)*(pi*floor(1/2*(d*x + c)/pi + 1/2)
      *sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c
      ))/sqrt(a^2 - b^2)))/((a^6*b^5 - 3*a^4*b^7 + 3*a^2*b^9 - b^11)*sqrt(a^2 -
      b^2)) - (18*B*a^9*tan(1/2*d*x + 1/2*c)^5 - 6*A*a^8*b*tan(1/2*d*x + 1/2*c)^
      5 - 42*B*a^8*b*tan(1/2*d*x + 1/2*c)^5 + 15*A*a^7*b^2*tan(1/2*d*x + 1/2*c)^
      5 - 24*B*a^7*b^2*tan(1/2*d*x + 1/2*c)^5 + 6*A*a^6*b^3*tan(1/2*d*x + 1/2*c)
      ^5 + 117*B*a^6*b^3*tan(1/2*d*x + 1/2*c)^5 - 45*A*a^5*b^4*tan(1/2*d*x + 1/2
      *c)^5 - 24*B*a^5*b^4*tan(1/2*d*x + 1/2*c)^5 + 6*A*a^4*b^5*tan(1/2*d*x + 1/
      2*c)^5 - 105*B*a^4*b^5*tan(1/2*d*x + 1/2*c)^5 + 60*A*a^3*b^6*tan(1/2*d*x +
      1/2*c)^5 + 60*B*a^3*b^6*tan(1/2*d*x + 1/2*c)^5 - 36*A*a^2*b^7*tan(1/2*d*x
      + 1/2*c)^5 + 36*B*a^9*tan(1/2*d*x + 1/2*c)^3 - 12*A*a^8*b*tan(1/2*d*x + 1
      /2*c)^3 - 152*B*a^7*b^2*tan(1/2*d*x + 1/2*c)^3 + 56*A*a^6*b^3*tan(1/2*d*x
      + 1/2*c)^3 + 236*B*a^5*b^4*tan(1/2*d*x + 1/2*c)^3 - 116*A*a^4*b^5*tan(1/2*
      d*x + 1/2*c)^3 - 120*B*a^3*b^6*tan(1/2*d*x + 1/2*c)^3 + 72*A*a^2*b^7*tan(1
      /2*d*x + 1/2*c)^3 + 18*B*a^9*tan(1/2*d*x + 1/2*c) - 6*A*a^8*b*tan(1/2*d*x
      + 1/2*c) + 42*B*a^8*b*tan(1/2*d*x + 1/2*c) - 15*A*a^7*b^2*tan(1/2*d*x + 1/
      2*c) - 24*B*a^7*b^2*tan(1/2*d*x + 1/2*c) + 6*A*a^6*b^3*tan(1/2*d*x + 1/2*c
      ) - 117*B*a^6*b^3*tan(1/2*d*x + 1/2*c) + 45*A*a^5*b^4*tan(1/2*d*x + 1/2*c)
      - 24*B*a^5*b^4*tan(1/2*d*x + 1/2*c) + 6*A*a^4*b^5*tan(1/2*d*x + 1/2*c)...

```

3.273.9 Mupad [B] (verification not implemented)

Time = 12.38 (sec) , antiderivative size = 7823, normalized size of antiderivative = 19.13

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Too large to display}$$

```

input int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^4,x)

```

output

$$\begin{aligned}
& (\log(\tan(c/2 + (d*x)/2) + 1i)*(A*b - 4*B*a)*1i)/(b^5*d) - ((\tan(c/2 + (d*x)/2))^7*(12*A*a^2*b^5 - 2*B*b^7 - 8*B*a^7 + 4*A*a^3*b^4 - 6*A*a^4*b^3 - A*a^5*b^2 + 6*B*a^2*b^5 - 26*B*a^3*b^4 - 11*B*a^4*b^3 + 24*B*a^5*b^2 + 2*A*a^6*b + 2*B*a*b^6 + 4*B*a^6*b))/(b^4*(a + b)^3*(a - b)) - (\tan(c/2 + (d*x)/2))^3*(72*B*a^8 + 18*B*b^8 + 36*A*a^2*b^6 - 96*A*a^3*b^5 - 14*A*a^4*b^4 + 59*A*a^5*b^3 + 3*A*a^6*b^2 - 72*B*a^2*b^6 - 60*B*a^3*b^5 + 273*B*a^4*b^4 + 47*B*a^5*b^3 - 236*B*a^6*b^2 - 18*A*a^7*b - 12*B*a^7*b))/(3*b^4*(a + b)^2*(a - b)^3) - (\tan(c/2 + (d*x)/2))^5*(72*B*a^8 + 18*B*b^8 - 36*A*a^2*b^6 - 96*A*a^3*b^5 + 14*A*a^4*b^4 + 59*A*a^5*b^3 - 3*A*a^6*b^2 - 72*B*a^2*b^6 + 60*B*a^3*b^5 + 273*B*a^4*b^4 - 47*B*a^5*b^3 - 236*B*a^6*b^2 - 18*A*a^7*b + 12*B*a^7*b))/(3*b^4*(a + b)^3*(a - b)^2) + (\tan(c/2 + (d*x)/2)*(2*B*b^7 - 8*B*a^7 + 12*A*a^2*b^5 - 4*A*a^3*b^4 - 6*A*a^4*b^3 + A*a^5*b^2 - 6*B*a^2*b^5 - 26*B*a^3*b^4 + 11*B*a^4*b^3 + 24*B*a^5*b^2 + 2*A*a^6*b + 2*B*a*b^6 - 4*B*a^6*b))/(b^4*(a + b)*(a - b)^3))/(d*(3*a*b^2 + 3*a^2*b - \tan(c/2 + (d*x)/2)^4*(6*a*b^2 - 6*a^3) + \tan(c/2 + (d*x)/2)^2*(6*a^2*b + 4*a^3 - 2*b^3) + \tan(c/2 + (d*x)/2)^6*(4*a^3 - 6*a^2*b + 2*b^3) + a^3 + b^3 + \tan(c/2 + (d*x)/2)^8*(3*a*b^2 - 3*a^2*b + a^3 - b^3))) - (\log(\tan(c/2 + (d*x)/2) - 1i)*(A*b*1i - B*a*4i))/(b^5*d) - (a*atan(((a*((8*\tan(c/2 + (d*x)/2)*(4*A^2*b^16 + 128*B^2*a^16 - 8*A^2*a*b^15 - 128*B^2*a^15*b + 44*A^2*a^2*b^14 + 48*A^2*a^3*b^13 - 92*A^2*a^4*b^12 - 120*A^2*a^5*b^11 + 156*A^2*a^6*b^10 + \dots
\end{aligned}$$

3.274 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

3.274.1 Optimal result 2537
 3.274.2 Mathematica [B] (verified) 2538
 3.274.3 Rubi [A] (verified) 2539
 3.274.4 Maple [A] (verified) 2544
 3.274.5 Fricas [B] (verification not implemented) 2544
 3.274.6 Sympy [F(-1)] 2545
 3.274.7 Maxima [F(-2)] 2546
 3.274.8 Giac [B] (verification not implemented) 2546
 3.274.9 Mupad [B] (verification not implemented) 2547

3.274.1 Optimal result

Integrand size = 31, antiderivative size = 301

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$$

$$= \frac{Bx}{b^4} - \frac{(3a^2Ab^5 + 2Ab^7 + 2a^7B - 7a^5b^2B + 8a^3b^4B - 8ab^6B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{7/2}b^4(a+b)^{7/2}d}$$

$$+ \frac{a(Ab - aB) \cos^2(c+dx) \sin(c+dx)}{3b(a^2 - b^2)d(a+b \cos(c+dx))^3} + \frac{a^2(5Ab^3 + 3a^3B - 8ab^2B) \sin(c+dx)}{6b^3(a^2 - b^2)^2d(a+b \cos(c+dx))^2}$$

$$- \frac{a(a^2Ab^3 - 16Ab^5 + 9a^5B - 28a^3b^2B + 34ab^4B) \sin(c+dx)}{6b^3(a^2 - b^2)^3d(a+b \cos(c+dx))}$$

```
output B*x/b^4-(3*A*a^2*b^5+2*A*b^7+2*B*a^7-7*B*a^5*b^2+8*B*a^3*b^4-8*B*a*b^6)*ar
ctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(7/2)/b^4/(a+b)^(7/
2)/d+1/3*a*(A*b-B*a)*cos(d*x+c)^2*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c)
)^3+1/6*a^2*(5*A*b^3+3*B*a^3-8*B*a*b^2)*sin(d*x+c)/b^3/(a^2-b^2)^2/d/(a+b*
cos(d*x+c))^2-1/6*a*(A*a^2*b^3-16*A*b^5+9*B*a^5-28*B*a^3*b^2+34*B*a*b^4)*s
in(d*x+c)/b^3/(a^2-b^2)^3/d/(a+b*cos(d*x+c))
```


3.274.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 717 vs. $2(301) = 602$.

Time = 3.63 (sec) , antiderivative size = 717, normalized size of antiderivative = 2.38

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

$$= \frac{24(3a^2Ab^5+2Ab^7+2a^7B-7a^5b^2B+8a^3b^4B-8ab^6B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{7/2}} + \frac{24a^9Bc-36a^7b^2Bc-36a^5b^4Bc+84a^3b^6Bc-36ab^8Bc}{(-a^2+b^2)^{7/2}}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output

```
((-24*(3*a^2*A*b^5 + 2*A*b^7 + 2*a^7*B - 7*a^5*b^2*B + 8*a^3*b^4*B - 8*a*b^6*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]]/(-a^2 + b^2)^(7/2) + (24*a^9*B*c - 36*a^7*b^2*B*c - 36*a^5*b^4*B*c + 84*a^3*b^6*B*c - 36*a*b^8*B*c + 24*a^9*B*d*x - 36*a^7*b^2*B*d*x - 36*a^5*b^4*B*d*x + 84*a^3*b^6*B*d*x - 36*a*b^8*B*d*x + 18*b*(a^2 - b^2)^3*(4*a^2 + b^2)*B*(c + d*x)*Cos[c + d*x] + 36*a*b^2*(a^2 - b^2)^3*B*(c + d*x)*Cos[2*(c + d*x)] + 6*a^6*b^3*B*c*Cos[3*(c + d*x)] - 18*a^4*b^5*B*c*Cos[3*(c + d*x)] + 18*a^2*b^7*B*c*Cos[3*(c + d*x)] - 6*b^9*B*c*Cos[3*(c + d*x)] + 6*a^6*b^3*B*d*x*Cos[3*(c + d*x)] - 18*a^4*b^5*B*d*x*Cos[3*(c + d*x)] + 18*a^2*b^7*B*d*x*Cos[3*(c + d*x)] - 6*b^9*B*d*x*Cos[3*(c + d*x)] + 18*a^5*A*b^4*Sin[c + d*x] + 39*a^3*A*b^6*Sin[c + d*x] + 18*a*A*b^8*Sin[c + d*x] - 24*a^8*b*B*Sin[c + d*x] + 57*a^6*b^3*B*Sin[c + d*x] - 72*a^4*b^5*B*Sin[c + d*x] - 36*a^2*b^7*B*Sin[c + d*x] + 6*a^4*A*b^5*Sin[2*(c + d*x)] + 54*a^2*A*b^7*Sin[2*(c + d*x)] - 30*a^7*b^2*B*Sin[2*(c + d*x)] + 90*a^5*b^4*B*Sin[2*(c + d*x)] - 120*a^3*b^6*B*Sin[2*(c + d*x)] + 2*a^5*A*b^4*Sin[3*(c + d*x)] - 5*a^3*A*b^6*Sin[3*(c + d*x)] + 18*a*A*b^8*Sin[3*(c + d*x)] - 11*a^6*b^3*B*Sin[3*(c + d*x)] + 32*a^4*b^5*B*Sin[3*(c + d*x)] - 36*a^2*b^7*B*Sin[3*(c + d*x)]/((a^2 - b^2)^3*(a + b*Cos[c + d*x])^3))/(24*b^4*d)
```

3.274.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.18, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3468, 25, 3042, 3510, 25, 3042, 3500, 27, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^4} dx$$

$$\downarrow \text{3468}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3} - \int \frac{\cos(c+dx)(3(a^2-b^2)B\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+2a(Ab-aB))}{(a+b\cos(c+dx))^3} dx}{3b(a^2-b^2)}$$

$$\downarrow \text{25}$$

$$\int \frac{\cos(c+dx)(3(a^2-b^2)B\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+2a(Ab-aB))}{(a+b\cos(c+dx))^3} dx}{3b(a^2-b^2)} + \frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})(3(a^2-b^2)B\sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+2a(Ab-aB))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx}{3b(a^2-b^2)} + \frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

$$\downarrow \text{3510}$$

$$\int \frac{-6b(a^2-b^2)^2B\cos^2(c+dx)+(3Ba^5-10b^2Ba^3+Ab^3a^2+12b^4Ba-6Ab^5)\cos(c+dx)+2ab(3Ba^3-8b^2Ba+5Ab^3)}{(a+b\cos(c+dx))^2}{2b^2(a^2-b^2)} dx + \frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

3.274. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

↓ 25

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{\int \frac{-6b(a^2-b^2)^2B\cos^2(c+dx)+(3Ba^5-10b^2Ba^3+Ab^3a^2+12b^4Ba-6Ab^5)\cos(c+dx)+2ab(3Ba^3-8b^2Ba+5Ab^3)}{(a+b\cos(c+dx))^2} dx}{2b^2(a^2-b^2)}$$

$$\frac{3b(a^2-b^2)}{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}$$

$$\frac{3bd(a^2-b^2)(a+b\cos(c+dx))^3}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{\int \frac{-6b(a^2-b^2)^2B\sin(c+dx+\frac{\pi}{2})^2+(3Ba^5-10b^2Ba^3+Ab^3a^2+12b^4Ba-6Ab^5)\sin(c+dx+\frac{\pi}{2})+2ab(3Ba^3-8b^2Ba+5Ab^3)}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx}{2b^2(a^2-b^2)}$$

$$\frac{3b(a^2-b^2)}{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}$$

$$\frac{3bd(a^2-b^2)(a+b\cos(c+dx))^3}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3500

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{\frac{a(9a^5B-28a^3b^2B+a^2Ab^3+34ab^4B-16Ab^5)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))} - \frac{3(b(2Ab^6-6aBb^5+3a^2Ab^4+2a^3Bb^3-a^5Bb)-2b(a^2-b^2)^3)\cos(c+dx)}{a+b\cos(c+dx)}}{2b^2(a^2-b^2)}$$

$$\frac{3b(a^2-b^2)}{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}$$

$$\frac{3bd(a^2-b^2)(a+b\cos(c+dx))^3}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 27

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{3\int \frac{b(2Ab^6-6aBb^5+3a^2Ab^4+2a^3Bb^3-a^5Bb)-2b(a^2-b^2)^3B\cos(c+dx)}{a+b\cos(c+dx)} dx + \frac{a(9a^5B-28a^3b^2B+a^2Ab^3+34ab^4B-16Ab^5)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))}}{2b^2(a^2-b^2)}$$

$$\frac{3b(a^2-b^2)}{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}$$

$$\frac{3bd(a^2-b^2)(a+b\cos(c+dx))^3}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{3\int \frac{b(2Ab^6-6aBb^5+3a^2Ab^4+2a^3Bb^3-a^5Bb)-2b(a^2-b^2)^3B\sin(c+dx+\frac{\pi}{2})}{a+b\sin(c+dx+\frac{\pi}{2})} dx + \frac{a(9a^5B-28a^3b^2B+a^2Ab^3+34ab^4B-16Ab^5)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))}}{2b^2(a^2-b^2)}$$

$$\frac{3b(a^2-b^2)}{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}$$

$$\frac{3bd(a^2-b^2)(a+b\cos(c+dx))^3}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3214

3.274. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{3\left(\frac{2a^7B-7a^5b^2B+8a^3b^4B+3a^2Ab^5-8ab^6B+2Ab^7}{a+b\cos(c+dx)}dx-2Bx(a^2-b^2)^3\right)}{b(a^2-b^2)} + \frac{a(9a^5B-28a^3b^2B+a^2b^3)}{d(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{3\left(\frac{2a^7B-7a^5b^2B+8a^3b^4B+3a^2Ab^5-8ab^6B+2Ab^7}{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}dx-2Bx(a^2-b^2)^3\right)}{b(a^2-b^2)} + \frac{a(9a^5B-28a^3b^2B+a^2b^3)}{d(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3138

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{3\left(\frac{2(2a^7B-7a^5b^2B+8a^3b^4B+3a^2Ab^5-8ab^6B+2Ab^7)}{d}\frac{1}{(a-b)\tan^2\left(\frac{1}{2}(c+dx)\right)+a+b}d\tan\left(\frac{1}{2}(c+dx)\right)-2Bx(a^2-b^2)^3\right)}{b(a^2-b^2)} + \frac{a(9a^5B-28a^3b^2B+a^2b^3)}{d(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 218

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3} + \frac{3\left(\frac{2(2a^7B-7a^5b^2B+8a^3b^4B+3a^2Ab^5-8ab^6B+2Ab^7)}{d\sqrt{a-b}\sqrt{a+b}}\arctan\left(\frac{\sqrt{a-b}\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{a+b}\cos\left(c+dx+\frac{\pi}{2}\right)}\right)-2Bx(a^2-b^2)^3\right)}{b(a^2-b^2)}$$

$$\frac{a^2(3a^3B-8ab^2B+5Ab^3)\sin(c+dx)}{2b^2d(a^2-b^2)(a+b\cos(c+dx))^2} - \frac{a(9a^5B-28a^3b^2B+a^2Ab^3+34ab^4B-16Ab^5)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))} + \frac{3\left(\frac{2(2a^7B-7a^5b^2B+8a^3b^4B+3a^2Ab^5-8ab^6B+2Ab^7)}{d\sqrt{a-b}\sqrt{a+b}}\arctan\left(\frac{\sqrt{a-b}\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{a+b}\cos\left(c+dx+\frac{\pi}{2}\right)}\right)-2Bx(a^2-b^2)^3\right)}{b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^3}$$

input `Int[(Cos[c + d*x])^3*(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^4,x]`

3.274. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

```
output (a*(A*b - a*B)*Cos[c + d*x]^2*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[
c + d*x])^3) + ((a^2*(5*A*b^3 + 3*a^3*B - 8*a*b^2*B)*Sin[c + d*x])/(2*b^2*
(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) - ((3*(-2*(a^2 - b^2)^3*B*x + (2*(3*
a^2*A*b^5 + 2*A*b^7 + 2*a^7*B - 7*a^5*b^2*B + 8*a^3*b^4*B - 8*a*b^6*B)*Arc
Tan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*
d)))/(b*(a^2 - b^2)) + (a*(a^2*A*b^3 - 16*A*b^5 + 9*a^5*B - 28*a^3*b^2*B +
34*a*b^4*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*b^2*(a
^2 - b^2)))/(3*b*(a^2 - b^2))
```

3.274.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

```
rule 3214 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d
*Sine[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-(b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.274.4 Maple [A] (verified)

Time = 1.73 (sec) , antiderivative size = 459, normalized size of antiderivative = 1.52

method	result
derivativedivides	$\frac{2B \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^4} - \frac{\left(\frac{2Aa^2b^3+3Aab^4+6Ab^5-2Ba^5+Ba^4b+6Ba^3b^2-4Ba^2b^3-12Bab^4}{2(a-b)(a^3+3a^2b+3ab^2+b^3)}\right)ab\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2(Aa^2b^3+3Aab^4+6Ab^5-2Ba^5+Ba^4b+6Ba^3b^2-4Ba^2b^3-12Bab^4)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)}$
default	$\frac{2B \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b^4} - \frac{\left(\frac{2Aa^2b^3+3Aab^4+6Ab^5-2Ba^5+Ba^4b+6Ba^3b^2-4Ba^2b^3-12Bab^4}{2(a-b)(a^3+3a^2b+3ab^2+b^3)}\right)ab\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2(Aa^2b^3+3Aab^4+6Ab^5-2Ba^5+Ba^4b+6Ba^3b^2-4Ba^2b^3-12Bab^4)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)}$
risch	Expression too large to display

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(2*B/b^4*arctan(tan(1/2*d*x+1/2*c))-2/b^4*((-1/2*(2*A*a^2*b^3+3*A*a*b^4+6*A*b^5-2*B*a^5+B*a^4*b+6*B*a^3*b^2-4*B*a^2*b^3-12*B*a*b^4)*a*b/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*tan(1/2*d*x+1/2*c)^5-2/3*(A*a^2*b^3+9*A*b^5-3*B*a^5+11*B*a^3*b^2-18*B*a*b^4)*a*b/(a^2+2*a*b+b^2)/(a^2-2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3-1/2*(2*A*a^2*b^3-3*A*a*b^4+6*A*b^5-2*B*a^5-B*a^4*b+6*B*a^3*b^2+4*B*a^2*b^3-12*B*a*b^4)*a*b/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^3+1/2*(3*A*a^2*b^5+2*A*b^7+2*B*a^7-7*B*a^5*b^2+8*B*a^3*b^4-8*B*a*b^6)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))))
```

3.274.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 894 vs. 2(286) = 572.

Time = 0.49 (sec) , antiderivative size = 1857, normalized size of antiderivative = 6.17

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="fricas")
```

output `[1/12*(12*(B*a^8*b^3 - 4*B*a^6*b^5 + 6*B*a^4*b^7 - 4*B*a^2*b^9 + B*b^11)*d*x*cos(d*x + c)^3 + 36*(B*a^9*b^2 - 4*B*a^7*b^4 + 6*B*a^5*b^6 - 4*B*a^3*b^8 + B*a*b^10)*d*x*cos(d*x + c)^2 + 36*(B*a^10*b - 4*B*a^8*b^3 + 6*B*a^6*b^5 - 4*B*a^4*b^7 + B*a^2*b^9)*d*x*cos(d*x + c) + 12*(B*a^11 - 4*B*a^9*b^2 + 6*B*a^7*b^4 - 4*B*a^5*b^6 + B*a^3*b^8)*d*x + 3*(2*B*a^10 - 7*B*a^8*b^2 + 8*B*a^6*b^4 + 3*A*a^5*b^5 - 8*B*a^4*b^6 + 2*A*a^3*b^7 + (2*B*a^7*b^3 - 7*B*a^5*b^5 + 8*B*a^3*b^7 + 3*A*a^2*b^8 - 8*B*a*b^9 + 2*A*b^10)*cos(d*x + c)^3 + 3*(2*B*a^8*b^2 - 7*B*a^6*b^4 + 8*B*a^4*b^6 + 3*A*a^3*b^7 - 8*B*a^2*b^8 + 2*A*a*b^9)*cos(d*x + c)^2 + 3*(2*B*a^9*b - 7*B*a^7*b^3 + 8*B*a^5*b^5 + 3*A*a^4*b^6 - 8*B*a^3*b^7 + 2*A*a^2*b^8)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(6*B*a^10*b - 23*B*a^8*b^3 - 4*A*a^7*b^4 + 43*B*a^6*b^5 - 7*A*a^5*b^6 - 26*B*a^4*b^7 + 11*A*a^3*b^8 + (11*B*a^8*b^3 - 2*A*a^7*b^4 - 43*B*a^6*b^5 + 7*A*a^5*b^6 + 68*B*a^4*b^7 - 23*A*a^3*b^8 - 36*B*a^2*b^9 + 18*A*a*b^10)*cos(d*x + c)^2 + 3*(5*B*a^9*b^2 - 20*B*a^7*b^4 - A*a^6*b^5 + 35*B*a^5*b^6 - 8*A*a^4*b^7 - 20*B*a^3*b^8 + 9*A*a^2*b^9)*cos(d*x + c))*sin(d*x + c))/((a^8*b^7 - 4*a^6*b^9 + 6*a^4*b^11 - 4*a^2*b^13 + b^15)*d*cos(d*x + c)^3 + 3*(a^9*b^6 - 4*a^7*b^8 + 6*a^5*b^10 - 4*a^3*b^12 + a*b^14)*d*cos(d*x + c)^2 + 3*(a^10*b^5 - 4*a^8*b^7 + 6*a^6*b^9 - 4*a^4...`

3.274.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**4,x)`

output `Timed out`

3.274.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.274.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 813 vs. 2(286) = 572.

Time = 0.35 (sec) , antiderivative size = 813, normalized size of antiderivative = 2.70

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

output `1/3*(3*(2*B*a^7 - 7*B*a^5*b^2 + 8*B*a^3*b^4 + 3*A*a^2*b^5 - 8*B*a*b^6 + 2*A*b^7)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^6*b^4 - 3*a^4*b^6 + 3*a^2*b^8 - b^10)*sqrt(a^2 - b^2)) + 3*(d*x + c)*B/b^4 - (6*B*a^8*tan(1/2*d*x + 1/2*c)^5 - 15*B*a^7*b*tan(1/2*d*x + 1/2*c)^5 - 6*B*a^6*b^2*tan(1/2*d*x + 1/2*c)^5 - 6*A*a^5*b^3*tan(1/2*d*x + 1/2*c)^5 + 45*B*a^5*b^3*tan(1/2*d*x + 1/2*c)^5 + 3*A*a^4*b^4*tan(1/2*d*x + 1/2*c)^5 - 6*B*a^4*b^4*tan(1/2*d*x + 1/2*c)^5 - 6*A*a^3*b^5*tan(1/2*d*x + 1/2*c)^5 - 60*B*a^3*b^5*tan(1/2*d*x + 1/2*c)^5 + 27*A*a^2*b^6*tan(1/2*d*x + 1/2*c)^5 + 36*B*a^2*b^6*tan(1/2*d*x + 1/2*c)^5 - 18*A*a*b^7*tan(1/2*d*x + 1/2*c)^5 + 12*B*a^8*tan(1/2*d*x + 1/2*c)^3 - 56*B*a^6*b^2*tan(1/2*d*x + 1/2*c)^3 - 4*A*a^5*b^3*tan(1/2*d*x + 1/2*c)^3 + 116*B*a^4*b^4*tan(1/2*d*x + 1/2*c)^3 - 32*A*a^3*b^5*tan(1/2*d*x + 1/2*c)^3 - 72*B*a^2*b^6*tan(1/2*d*x + 1/2*c)^3 + 36*A*a*b^7*tan(1/2*d*x + 1/2*c)^3 + 6*B*a^8*tan(1/2*d*x + 1/2*c) + 15*B*a^7*b*tan(1/2*d*x + 1/2*c) - 6*B*a^6*b^2*tan(1/2*d*x + 1/2*c) - 6*A*a^5*b^3*tan(1/2*d*x + 1/2*c) - 45*B*a^5*b^3*tan(1/2*d*x + 1/2*c) - 3*A*a^4*b^4*tan(1/2*d*x + 1/2*c) - 6*B*a^4*b^4*tan(1/2*d*x + 1/2*c) - 6*A*a^3*b^5*tan(1/2*d*x + 1/2*c) + 60*B*a^3*b^5*tan(1/2*d*x + 1/2*c) - 27*A*a^2*b^6*tan(1/2*d*x + 1/2*c) + 36*B*a^2*b^6*tan(1/2*d*x + 1/2*c) - 18*A*a*b^7*tan(1/2*d*x + 1/2*c)))/((a^6*b^3 - 3*a^4*b^5 + 3*a^2*b^7 - b^9)*(a*tan(1/2*d*x + 1/2*c))^2 ...`

3.274.9 Mupad [B] (verification not implemented)

Time = 12.76 (sec) , antiderivative size = 9733, normalized size of antiderivative = 32.34

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^4,x)`

output $((\tan(c/2 + (d*x)/2)^5*(3*A*a^2*b^4 - 2*B*a^6 + 2*A*a^3*b^3 - 12*B*a^2*b^4 - 4*B*a^3*b^3 + 6*B*a^4*b^2 + 6*A*a*b^5 + B*a^5*b))/((a*b^3 - b^4)*(a + b)^3) - (\tan(c/2 + (d*x)/2)*(2*B*a^6 + 3*A*a^2*b^4 - 2*A*a^3*b^3 + 12*B*a^2*b^4 - 4*B*a^3*b^3 - 6*B*a^4*b^2 - 6*A*a*b^5 + B*a^5*b))/((a + b)*(3*a*b^5 - b^6 - 3*a^2*b^4 + a^3*b^3)) + (4*\tan(c/2 + (d*x)/2)^3*(A*a^3*b^3 - 3*B*a^6 - 18*B*a^2*b^4 + 11*B*a^4*b^2 + 9*A*a*b^5))/(3*(a + b)^2*(b^5 - 2*a*b^4 + a^2*b^3)))/(d*(3*a*b^2 - \tan(c/2 + (d*x)/2)^4*(3*a*b^2 + 3*a^2*b - 3*a^3 - 3*b^3) - \tan(c/2 + (d*x)/2)^2*(3*a*b^2 - 3*a^2*b - 3*a^3 + 3*b^3) + 3*a^2*b + a^3 + b^3 + \tan(c/2 + (d*x)/2)^6*(3*a*b^2 - 3*a^2*b + a^3 - b^3)) + (2*B*\operatorname{atan}((B*((8*\tan(c/2 + (d*x)/2)*(4*A^2*b^14 + 8*B^2*a^14 + 4*B^2*b^14 - 8*B^2*a*b^13 - 8*B^2*a^13*b + 12*A^2*a^2*b^12 + 9*A^2*a^4*b^10 + 44*B^2*a^2*b^12 + 48*B^2*a^3*b^11 - 92*B^2*a^4*b^10 - 120*B^2*a^5*b^9 + 156*B^2*a^6*b^8 + 160*B^2*a^7*b^7 - 164*B^2*a^8*b^6 - 120*B^2*a^9*b^5 + 117*B^2*a^10*b^4 + 48*B^2*a^11*b^3 - 48*B^2*a^12*b^2 - 32*A*B*a*b^13 - 16*A*B*a^3*b^11 + 20*A*B*a^5*b^9 - 34*A*B*a^7*b^7 + 12*A*B*a^9*b^5)))/(a*b^16 + b^17 - 5*a^2*b^15 - 5*a^3*b^14 + 10*a^4*b^13 + 10*a^5*b^12 - 10*a^6*b^11 - 10*a^7*b^10 + 5*a^8*b^9 + 5*a^9*b^8 - a^10*b^7 - a^11*b^6) + (B*((8*(4*A*b^21 + 4*B*b^21 - 6*A*a^2*b^19 + 6*A*a^3*b^18 - 6*A*a^4*b^17 + 6*A*a^5*b^16 + 14*A*a^6*b^15 - 14*A*a^7*b^14 - 6*A*a^8*b^13 + 6*A*a^9*b^12 - 12*B*a^2*b^19 + 64*B*a^3*b^18 + 20*B*a^4*b^17 - 110*B*a^5*b^16 - 30*B*a^6*b^15 + 11...$

3.275 $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

3.275.1 Optimal result 2549
 3.275.2 Mathematica [A] (verified) 2550
 3.275.3 Rubi [A] (verified) 2550
 3.275.4 Maple [A] (verified) 2554
 3.275.5 Fricas [B] (verification not implemented) 2555
 3.275.6 Sympy [F(-1)] 2555
 3.275.7 Maxima [F(-2)] 2556
 3.275.8 Giac [B] (verification not implemented) 2556
 3.275.9 Mupad [B] (verification not implemented) 2557

3.275.1 Optimal result

Integrand size = 31, antiderivative size = 274

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$$

$$= \frac{(a^3A + 4aAb^2 - 3a^2bB - 2b^3B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{7/2}(a+b)^{7/2}d}$$

$$- \frac{a^2(Ab - aB) \sin(c+dx)}{3b^2(a^2 - b^2)d(a+b \cos(c+dx))^3} + \frac{a(a^2Ab - 6Ab^3 - 4a^3B + 9ab^2B) \sin(c+dx)}{6b^2(a^2 - b^2)^2d(a+b \cos(c+dx))^2}$$

$$+ \frac{(a^4Ab - 10a^2Ab^3 - 6Ab^5 + 2a^5B - 5a^3b^2B + 18ab^4B) \sin(c+dx)}{6b^2(a^2 - b^2)^3d(a+b \cos(c+dx))}$$

```
output (A*a^3+4*A*a*b^2-3*B*a^2*b-2*B*b^3)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/
(a+b)^(1/2))/(a-b)^(7/2)/(a+b)^(7/2)/d-1/3*a^2*(A*b-B*a)*sin(d*x+c)/b^2/(a
^2-b^2)/d/(a+b*cos(d*x+c))^3+1/6*a*(A*a^2*b-6*A*b^3-4*B*a^3+9*B*a*b^2)*sin
(d*x+c)/b^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^2+1/6*(A*a^4*b-10*A*a^2*b^3-6*A
*b^5+2*B*a^5-5*B*a^3*b^2+18*B*a*b^4)*sin(d*x+c)/b^2/(a^2-b^2)^3/d/(a+b*cos
(d*x+c))
```

3.275.2 Mathematica [A] (verified)

Time = 1.64 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.92

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

$$= \frac{24(a^3A+4aAb^2-3a^2bB-2b^3B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + \frac{2(-25a^4Ab-14a^2Ab^3-6Ab^5+10a^5B+17a^3b^2B+18ab^4B+6a(a^4A-9a^2Ab^2-2Ab^4+a^3bB+9ab^3B))\cos(c+dx)+(a^4Ab-10a^2Ab^3-6Ab^5+2a^5B-5a^3b^2B+18a^4bB)\cos(2(c+dx))\sin(c+dx)}{24(a^2-b^2)^3d}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output `((-24*(a^3*A + 4*a*A*b^2 - 3*a^2*b*B - 2*b^3*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + (2*(-25*a^4*A*b - 14*a^2*A*b^3 - 6*A*b^5 + 10*a^5*B + 17*a^3*b^2*B + 18*a*b^4*B + 6*a*(a^4*A - 9*a^2*A*b^2 - 2*A*b^4 + a^3*b*B + 9*a*b^3*B))*Cos[c + d*x] + (a^4*A*b - 10*a^2*A*b^3 - 6*A*b^5 + 2*a^5*B - 5*a^3*b^2*B + 18*a*b^4*B)*Cos[2*(c + d*x)])*Sin[c + d*x])/(a + b*Cos[c + d*x])^3)/(24*(a^2 - b^2)^3*d)`

3.275.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.16, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3467, 3042, 3500, 25, 3042, 3233, 27, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^2(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^4} dx$$

↓ 3467

$$\int \frac{3b(a^2-b^2)B\cos^2(c+dx)+(a^2-3b^2)(Ab-aB)\cos(c+dx)+3ab(Ab-aB)}{(a+b\cos(c+dx))^3} dx - \frac{a^2(Ab-aB)\sin(c+dx)}{3b^2d(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

3.275. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

$$\frac{\int \frac{3b(a^2-b^2)B \sin(c+dx+\frac{\pi}{2})^2 + (a^2-3b^2)(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + 3ab(Ab-aB)}{(a+b \sin(c+dx+\frac{\pi}{2}))^3} dx}{\frac{3b^2(a^2-b^2)}{a^2(Ab-aB) \sin(c+dx)} \frac{1}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^3}} -$$

↓ 3500

$$\frac{\frac{a(-4a^3B+a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2} - \int \frac{2(Ba^3+2Aba^2-6b^2Ba+3Ab^3)b^2 + (2Ba^4+Ab^3-3b^2Ba^2-6Ab^3a+6b^4B) \cos(c+dx)b}{(a+b \cos(c+dx))^2} dx}{\frac{3b^2(a^2-b^2)}{a^2(Ab-aB) \sin(c+dx)} \frac{1}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^3}} -$$

↓ 25

$$\frac{\int \frac{2(Ba^3+2Aba^2-6b^2Ba+3Ab^3)b^2 + (2Ba^4+Ab^3-3b^2Ba^2-6Ab^3a+6b^4B) \cos(c+dx)b}{(a+b \cos(c+dx))^2} dx}{\frac{3b^2(a^2-b^2)}{a^2(Ab-aB) \sin(c+dx)} \frac{1}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^3}} + \frac{a(-4a^3B+a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2} -$$

↓ 3042

$$\frac{\int \frac{2(Ba^3+2Aba^2-6b^2Ba+3Ab^3)b^2 + (2Ba^4+Ab^3-3b^2Ba^2-6Ab^3a+6b^4B) \sin(c+dx+\frac{\pi}{2})b}{(a+b \sin(c+dx+\frac{\pi}{2}))^2} dx}{\frac{3b^2(a^2-b^2)}{a^2(Ab-aB) \sin(c+dx)} \frac{1}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^3}} + \frac{a(-4a^3B+a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2} -$$

↓ 3233

$$\frac{\frac{b(2a^5B+a^4Ab-5a^3b^2B-10a^2Ab^3+18ab^4B-6Ab^5) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))} - \int \frac{3b^3(Aa^3-3bBa^2+4Ab^2a-2b^3B)}{a+b \cos(c+dx)} dx}{\frac{3b^2(a^2-b^2)}{a^2(Ab-aB) \sin(c+dx)} \frac{1}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^3}} + \frac{a(-4a^3B+a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2} -$$

↓ 27

3.275. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

$$\frac{3b^3(a^3A - 3a^2bB + 4aAb^2 - 2b^3B) \int \frac{1}{a+b \cos(c+dx)} dx + \frac{b(2a^5B + a^4Ab - 5a^3b^2B - 10a^2Ab^3 + 18ab^4B - 6Ab^5) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{2b(a^2-b^2)} + \frac{a(-4a^3B + a^2Ab + 9ab^2B - 6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3042

$$\frac{3b^3(a^3A - 3a^2bB + 4aAb^2 - 2b^3B) \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx + \frac{b(2a^5B + a^4Ab - 5a^3b^2B - 10a^2Ab^3 + 18ab^4B - 6Ab^5) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{2b(a^2-b^2)} + \frac{a(-4a^3B + a^2Ab + 9ab^2B - 6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3138

$$\frac{6b^3(a^3A - 3a^2bB + 4aAb^2 - 2b^3B) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx)) + \frac{b(2a^5B + a^4Ab - 5a^3b^2B - 10a^2Ab^3 + 18ab^4B - 6Ab^5) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{2b(a^2-b^2)} + \frac{a(-4a^3B + a^2Ab + 9ab^2B - 6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 218

$$\frac{a(-4a^3B + a^2Ab + 9ab^2B - 6Ab^3) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2} + \frac{6b^3(a^3A - 3a^2bB + 4aAb^2 - 2b^3B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) + \frac{b(2a^5B + a^4Ab - 5a^3b^2B - 10a^2Ab^3 + 18ab^4B - 6Ab^5) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{2b(a^2-b^2)}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^3}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output `-1/3*(a^2*(A*b - a*B)*Sin[c + d*x])/(b^2*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^3) + ((a*(a^2*A*b - 6*A*b^3 - 4*a^3*B + 9*a*b^2*B)*Sin[c + d*x])/(2*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + (((6*b^3*(a^3*A + 4*a*A*b^2 - 3*a^2*b*B - 2*b^3*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*d) + (b*(a^4*A*b - 10*a^2*A*b^3 - 6*A*b^5 + 2*a^5*B - 5*a^3*b^2*B + 18*a*b^4*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))) / (2*b*(a^2 - b^2))) / (3*b^2*(a^2 - b^2))`

3.275. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$

3.275.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`
- rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.275.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 575 vs. $2(260) = 520$.

Time = 0.38 (sec) , antiderivative size = 1220, normalized size of antiderivative = 4.45

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="fracas")
```

```
output [-1/12*(3*(A*a^6 - 3*B*a^5*b + 4*A*a^4*b^2 - 2*B*a^3*b^3 + (A*a^3*b^3 - 3*B*a^2*b^4 + 4*A*a*b^5 - 2*B*b^6)*cos(d*x + c))^3 + 3*(A*a^4*b^2 - 3*B*a^3*b^3 + 4*A*a^2*b^4 - 2*B*a*b^5)*cos(d*x + c)^2 + 3*(A*a^5*b - 3*B*a^4*b^2 + 4*A*a^3*b^3 - 2*B*a^2*b^4)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(4*B*a^7 - 13*A*a^6*b + 7*B*a^5*b^2 + 11*A*a^4*b^3 - 11*B*a^3*b^4 + 2*A*a^2*b^5 + (2*B*a^7 + A*a^6*b - 7*B*a^5*b^2 - 11*A*a^4*b^3 + 23*B*a^3*b^4 + 4*A*a^2*b^5 - 18*B*a*b^6 + 6*A*b^7)*cos(d*x + c)^2 + 3*(A*a^7 + B*a^6*b - 10*A*a^5*b^2 + 8*B*a^4*b^3 + 7*A*a^3*b^4 - 9*B*a^2*b^5 + 2*A*a*b^6)*cos(d*x + c))*sin(d*x + c))/((a^8*b^3 - 4*a^6*b^5 + 6*a^4*b^7 - 4*a^2*b^9 + b^11)*d*cos(d*x + c)^3 + 3*(a^9*b^2 - 4*a^7*b^4 + 6*a^5*b^6 - 4*a^3*b^8 + a*b^10)*d*cos(d*x + c)^2 + 3*(a^10*b - 4*a^8*b^3 + 6*a^6*b^5 - 4*a^4*b^7 + a^2*b^9)*d*cos(d*x + c) + (a^11 - 4*a^9*b^2 + 6*a^7*b^4 - 4*a^5*b^6 + a^3*b^8)*d), 1/6*(3*(A*a^6 - 3*B*a^5*b + 4*A*a^4*b^2 - 2*B*a^3*b^3 + (A*a^3*b^3 - 3*B*a^2*b^4 + 4*A*a*b^5 - 2*B*b^6)*cos(d*x + c))^3 + 3*(A*a^4*b^2 - 3*B*a^3*b^3 + 4*A*a^2*b^4 - 2*B*a*b^5)*cos(d*x + c)^2 + 3*(A*a^5*b - 3*B*a^4*b^2 + 4*A*a^3*b^3 - 2*B*a^2*b^4)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (4*B*a^7 - 13*A*a^6*b + 7*B*a^5*b^2 + 11*A*a^4*b^3 - 11*B*a^3*b^4 + 2*A*a^2*b^5 + (...
```

3.275.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**4,x)
```

3.275. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

output Timed out

3.275.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.275.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 689 vs. 2(260) = 520.

Time = 0.35 (sec) , antiderivative size = 689, normalized size of antiderivative = 2.51

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \frac{3(Aa^3-3Ba^2b+4Aab^2-2Bb^3) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2-b^2}} \right) \right)}{(a^6-3a^4b^2+3a^2b^4-b^6)\sqrt{a^2-b^2}} + \frac{3Aa^5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 6Aa^4 b \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 3Aa^3 b^2 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(a^6-3a^4b^2+3a^2b^4-b^6)\sqrt{a^2-b^2}}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

output

```
-1/3*(3*(A*a^3 - 3*B*a^2*b + 4*A*a*b^2 - 2*B*b^3)*(pi*floor(1/2*(d*x + c)/
pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*
x + 1/2*c))/sqrt(a^2 - b^2)))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*sqrt(a^
2 - b^2)) + (3*A*a^5*tan(1/2*d*x + 1/2*c)^5 - 6*B*a^5*tan(1/2*d*x + 1/2*c)
^5 + 12*A*a^4*b*tan(1/2*d*x + 1/2*c)^5 + 3*B*a^4*b*tan(1/2*d*x + 1/2*c)^5
- 27*A*a^3*b^2*tan(1/2*d*x + 1/2*c)^5 - 6*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^5
+ 12*A*a^2*b^3*tan(1/2*d*x + 1/2*c)^5 + 27*B*a^2*b^3*tan(1/2*d*x + 1/2*c)
^5 - 6*A*a*b^4*tan(1/2*d*x + 1/2*c)^5 - 18*B*a*b^4*tan(1/2*d*x + 1/2*c)^5
+ 6*A*b^5*tan(1/2*d*x + 1/2*c)^5 - 4*B*a^5*tan(1/2*d*x + 1/2*c)^3 + 28*A*a
^4*b*tan(1/2*d*x + 1/2*c)^3 - 32*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 - 16*A*a
^2*b^3*tan(1/2*d*x + 1/2*c)^3 + 36*B*a*b^4*tan(1/2*d*x + 1/2*c)^3 - 12*A*b
^5*tan(1/2*d*x + 1/2*c)^3 - 3*A*a^5*tan(1/2*d*x + 1/2*c) - 6*B*a^5*tan(1/2
*d*x + 1/2*c) + 12*A*a^4*b*tan(1/2*d*x + 1/2*c) - 3*B*a^4*b*tan(1/2*d*x +
1/2*c) + 27*A*a^3*b^2*tan(1/2*d*x + 1/2*c) - 6*B*a^3*b^2*tan(1/2*d*x + 1/2
*c) + 12*A*a^2*b^3*tan(1/2*d*x + 1/2*c) - 27*B*a^2*b^3*tan(1/2*d*x + 1/2*c
) + 6*A*a*b^4*tan(1/2*d*x + 1/2*c) - 18*B*a*b^4*tan(1/2*d*x + 1/2*c) + 6*A
*b^5*tan(1/2*d*x + 1/2*c))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*(a*tan(1/2
*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^3))/d
```

3.275.9 Mupad [B] (verification not implemented)

Time = 6.72 (sec) , antiderivative size = 440, normalized size of antiderivative = 1.61

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)(2a - 2b)(a^3 - 3a^2b + 3ab^2 - b^3)}{2\sqrt{a+b}(a-b)^{7/2}}\right)(Aa^3 - 3Ba^2b + 4Aab^2 - 2Bb^3)}{d(a+b)^{7/2}(a-b)^{7/2}}$$

$$- \frac{\frac{4\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3(-Ba^3 + 7Aa^2b - 9Bab^2 + 3Ab^3)}{3(a+b)^2(a^2 - 2ab + b^2)} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5(Aa^3 + 2Ab^3 - 2Ba^3 + 2Aab^2 + 6Aa^2b - 6Bab^2 - 3Ba^2b)}{(a+b)^3(a-b)}}{d\left(3ab^2 - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4(-3a^3 + 3a^2b + 3ab^2 - 3b^3) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2(-3a^3 - 3a^2b + 3ab^2 + 3b^3)\right)}$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^4,x)`

output $(\operatorname{atan}(\tan(c/2 + (d*x)/2)*(2*a - 2*b)*(3*a*b^2 - 3*a^2*b + a^3 - b^3))/(2*(a + b)^{(1/2)}*(a - b)^{(7/2)}))*(A*a^3 - 2*B*b^3 + 4*A*a*b^2 - 3*B*a^2*b))/(d*(a + b)^{(7/2)}*(a - b)^{(7/2)}) - ((4*\tan(c/2 + (d*x)/2)^3*(3*A*b^3 - B*a^3 + 7*A*a^2*b - 9*B*a*b^2))/(3*(a + b)^2*(a^2 - 2*a*b + b^2)) + (\tan(c/2 + (d*x)/2)^5*(A*a^3 + 2*A*b^3 - 2*B*a^3 + 2*A*a*b^2 + 6*A*a^2*b - 6*B*a*b^2 - 3*B*a^2*b)))/((a + b)^3*(a - b)) - (\tan(c/2 + (d*x)/2)*(A*a^3 - 2*A*b^3 + 2*B*a^3 + 2*A*a*b^2 - 6*A*a^2*b + 6*B*a*b^2 - 3*B*a^2*b))/((a + b)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)))/(d*(3*a*b^2 - \tan(c/2 + (d*x)/2)^4*(3*a*b^2 + 3*a^2*b - 3*a^3 - 3*b^3) - \tan(c/2 + (d*x)/2)^2*(3*a*b^2 - 3*a^2*b - 3*a^3 + 3*b^3) + 3*a^2*b + a^3 + b^3 + \tan(c/2 + (d*x)/2)^6*(3*a*b^2 - 3*a^2*b + a^3 - b^3)))$

3.276
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$$

3.276.1 Optimal result 2559
 3.276.2 Mathematica [A] (verified) 2560
 3.276.3 Rubi [A] (verified) 2560
 3.276.4 Maple [A] (verified) 2564
 3.276.5 Fricas [B] (verification not implemented) 2565
 3.276.6 Sympy [F(-1)] 2565
 3.276.7 Maxima [F(-2)] 2566
 3.276.8 Giac [B] (verification not implemented) 2566
 3.276.9 Mupad [B] (verification not implemented) 2567

3.276.1 Optimal result

Integrand size = 29, antiderivative size = 263

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$$

$$= -\frac{(4a^2Ab + Ab^3 - a^3B - 4ab^2B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{7/2}(a+b)^{7/2}d}$$

$$+ \frac{a(Ab - aB) \sin(c+dx)}{3b(a^2 - b^2)d(a+b \cos(c+dx))^3} + \frac{(2a^2Ab + 3Ab^3 + a^3B - 6ab^2B) \sin(c+dx)}{6b(a^2 - b^2)^2d(a+b \cos(c+dx))^2}$$

$$+ \frac{(2a^3Ab + 13aAb^3 + a^4B - 10a^2b^2B - 6b^4B) \sin(c+dx)}{6b(a^2 - b^2)^3d(a+b \cos(c+dx))}$$

```
output - (4*A*a^2*b+A*b^3-B*a^3-4*B*a*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/(a-b)^(7/2)/(a+b)^(7/2)/d+1/3*a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^3+1/6*(2*A*a^2*b+3*A*b^3+B*a^3-6*B*a*b^2)*sin(d*x+c)/b/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^2+1/6*(2*A*a^3*b+13*A*a*b^3+B*a^4-10*B*a^2*b^2-6*B*b^4)*sin(d*x+c)/b/(a^2-b^2)^3/d/(a+b*cos(d*x+c))
```

3.276.2 Mathematica [A] (verified)

Time = 1.29 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.96

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

$$= \frac{24(-4a^2Ab - Ab^3 + a^3B + 4ab^2B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + \frac{2(12a^5A + 22a^3Ab^2 + 11aAb^4 - 25a^4bB - 14a^2b^3B - 6b^5B + 6(2a^4Ab + 9a^3A^2b - 13a^2A^2b^3 + a^4B - 10a^2b^2B - 6b^4B)\cos[2(c+dx)])\sin[c+dx]}{24(a^2 - b^2)^3 d}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output `((-24*(-4*a^2*A*b - A*b^3 + a^3*B + 4*a*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + (2*(12*a^5*A + 22*a^3*A*b^2 + 11*a*A*b^4 - 25*a^4*b*B - 14*a^2*b^3*B - 6*b^5*B + 6*(2*a^4*A*b + 9*a^2*A*b^3 - A*b^5 + a^5*B - 9*a^3*b^2*B - 2*a*b^4*B)*Cos[c + d*x] + b*(2*a^3*A*b + 13*a*A*b^3 + a^4*B - 10*a^2*b^2*B - 6*b^4*B)*Cos[2*(c + d*x)])*Sin[c + d*x])/(a + b*Cos[c + d*x])^3)/(24*(a^2 - b^2)^3*d)`

3.276.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.15, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$, Rules used = {3042, 3447, 3042, 3500, 3042, 3233, 25, 3042, 3233, 27, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)(A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a+b\sin\left(c+dx+\frac{\pi}{2}\right))^4} dx$$

$$\downarrow \text{3447}$$

$$\int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a+b\cos(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

3.276. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

$$\begin{aligned}
 & \int \frac{A \sin \left(c+dx+\frac{\pi}{2}\right)+B \sin \left(c+dx+\frac{\pi}{2}\right)^2}{\left(a+b \sin \left(c+dx+\frac{\pi}{2}\right)\right)^4} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{a(A b-a B) \sin (c+dx)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^3}-\int \frac{3 b(A b-a B)-\left(B a^2+2 A b a-3 b^2 B\right) \cos (c+dx)}{\left(a+b \cos (c+dx)\right)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(A b-a B) \sin (c+dx)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^3}-\int \frac{3 b(A b-a B)+\left(-B a^2-2 A b a+3 b^2 B\right) \sin \left(c+dx+\frac{\pi}{2}\right)}{\left(a+b \sin \left(c+dx+\frac{\pi}{2}\right)\right)^3} dx \\
 & \quad \downarrow \text{3233} \\
 & \frac{a(A b-a B) \sin (c+dx)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^3}- \\
 & -\frac{\int-\frac{2 b\left(-2 B a^2+5 A b a-3 b^2 B\right)-\left(B a^3+2 A b a^2-6 b^2 B a+3 A b^3\right) \cos (c+dx)}{\left(a+b \cos (c+dx)\right)^2} dx}{2\left(a^2-b^2\right)}-\frac{\left(a^3 B+2 a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+dx)}{2 d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{a(A b-a B) \sin (c+dx)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^3}- \\
 & \frac{\int \frac{2 b\left(-2 B a^2+5 A b a-3 b^2 B\right)-\left(B a^3+2 A b a^2-6 b^2 B a+3 A b^3\right) \cos (c+dx)}{\left(a+b \cos (c+dx)\right)^2} dx}{2\left(a^2-b^2\right)}-\frac{\left(a^3 B+2 a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+dx)}{2 d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(A b-a B) \sin (c+dx)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^3}- \\
 & \frac{\int \frac{2 b\left(-2 B a^2+5 A b a-3 b^2 B\right)+\left(-B a^3-2 A b a^2+6 b^2 B a-3 A b^3\right) \sin \left(c+dx+\frac{\pi}{2}\right)}{\left(a+b \sin \left(c+dx+\frac{\pi}{2}\right)\right)^2} dx}{2\left(a^2-b^2\right)}-\frac{\left(a^3 B+2 a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+dx)}{2 d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{a(A b-a B) \sin (c+dx)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^3}- \\
 & \frac{\int-\frac{3 b\left(-B a^3+4 A b a^2-4 b^2 B a+A b^3\right)}{a+b \cos (c+dx)} dx}{a^2-b^2}-\frac{\left(a^4 B+2 a^3 A b-10 a^2 b^2 B+13 a A b^3-6 b^4 B\right) \sin (c+dx)}{d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)}-\frac{\left(a^3 B+2 a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+dx)}{2 d\left(a^2-b^2\right)\left(a+b \cos (c+dx)\right)^2} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.276. $\int \frac{\cos (c+dx)(A+B \cos (c+dx))}{(a+b \cos (c+dx))^4} dx$

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} - \frac{3b(a^3(-B) + 4a^2Ab - 4ab^2B + Ab^3) \int \frac{1}{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{(a^4B + 2a^3Ab - 10a^2b^2B + 13aAb^3 - 6b^4B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{2(a^2 - b^2)} - \frac{(a^3B + 2a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3042

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} - \frac{3b(a^3(-B) + 4a^2Ab - 4ab^2B + Ab^3) \int \frac{1}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{a^2 - b^2} - \frac{(a^4B + 2a^3Ab - 10a^2b^2B + 13aAb^3 - 6b^4B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{2(a^2 - b^2)} - \frac{(a^3B + 2a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3138

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} - \frac{6b(a^3(-B) + 4a^2Ab - 4ab^2B + Ab^3) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{d(a^2 - b^2)} - \frac{(a^4B + 2a^3Ab - 10a^2b^2B + 13aAb^3 - 6b^4B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{2(a^2 - b^2)} - \frac{(a^3B + 2a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))}$$

218

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^3} - \frac{6b(a^3(-B) + 4a^2Ab - 4ab^2B + Ab^3) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{d\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)} - \frac{(a^4B + 2a^3Ab - 10a^2b^2B + 13aAb^3 - 6b^4B) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))}}{2(a^2 - b^2)} - \frac{(a^3B + 2a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{2d(a^2 - b^2)(a + b \cos(c + dx))}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^4,x]`

output `(a*(A*b - a*B)*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^3) - (-1/2*((2*a^2*A*b + 3*A*b^3 + a^3*B - 6*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((6*b*(4*a^2*A*b + A*b^3 - a^3*B - 4*a*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*d) - ((2*a^3*A*b + 13*a*A*b^3 + a^4*B - 10*a^2*b^2*B - 6*b^4*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*(a^2 - b^2)))/(3*b*(a^2 - b^2))`

3.276.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(- (A*b^2
- a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.276.4 Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.46

method	result
derivativedivides	$\frac{2 \left(-\frac{(2Aa^3+2Aa^2b+6Aab^2+Ab^3-Ba^3-6Ba^2b-2Bab^2-2Bb^3) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)} - \frac{2(3Aa^3+7Aab^2-7Ba^2b-3Bb^3) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3(a^2-2ab+b^2)(a^2+2ab+b^2)} \right)}{\left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a-b} \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a+b} \right)}$
default	$\frac{2 \left(-\frac{(2Aa^3+2Aa^2b+6Aab^2+Ab^3-Ba^3-6Ba^2b-2Bab^2-2Bb^3) \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)} - \frac{2(3Aa^3+7Aab^2-7Ba^2b-3Bb^3) \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{3(a^2-2ab+b^2)(a^2+2ab+b^2)} \right)}{\left(\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a-b} \left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)^{a+b} \right)}$
risch	Expression too large to display

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE
)
```

```
output 1/d*(-2*(-1/2*(2*A*a^3+2*A*a^2*b+6*A*a*b^2+A*b^3-B*a^3-6*B*a^2*b-2*B*a*b^2
-2*B*b^3)/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*tan(1/2*d*x+1/2*c)^5-2/3*(3*A*a^
3+7*A*a*b^2-7*B*a^2*b-3*B*b^3)/(a^2-2*a*b+b^2)/(a^2+2*a*b+b^2)*tan(1/2*d*x
+1/2*c)^3-1/2*(2*A*a^3-2*A*a^2*b+6*A*a*b^2-A*b^3+B*a^3-6*B*a^2*b+2*B*a*b^2
-2*B*b^3)/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x
+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^3-(4*A*a^2*b+A*b^3-B*a^3-4*B*a*b^2
)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d
*x+1/2*c)/((a-b)*(a+b))^(1/2)))
```

$$3.276. \int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^4} dx$$

3.276.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 581 vs. $2(247) = 494$.

Time = 0.38 (sec) , antiderivative size = 1232, normalized size of antiderivative = 4.68

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="fracas")
```

```
output [-1/12*(3*(B*a^6 - 4*A*a^5*b + 4*B*a^4*b^2 - A*a^3*b^3 + (B*a^3*b^3 - 4*A*a^2*b^4 + 4*B*a*b^5 - A*b^6)*cos(d*x + c)^3 + 3*(B*a^4*b^2 - 4*A*a^3*b^3 + 4*B*a^2*b^4 - A*a*b^5)*cos(d*x + c)^2 + 3*(B*a^5*b - 4*A*a^4*b^2 + 4*B*a^3*b^3 - A*a^2*b^4)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(6*A*a^7 - 13*B*a^6*b + 4*A*a^5*b^2 + 11*B*a^4*b^3 - 11*A*a^3*b^4 + 2*B*a^2*b^5 + A*a*b^6 + (B*a^6*b + 2*A*a^5*b^2 - 11*B*a^4*b^3 + 11*A*a^3*b^4 + 4*B*a^2*b^5 - 13*A*a*b^6 + 6*B*b^7)*cos(d*x + c)^2 + 3*(B*a^7 + 2*A*a^6*b - 10*B*a^5*b^2 + 7*A*a^4*b^3 + 7*B*a^3*b^4 - 10*A*a^2*b^5 + 2*B*a*b^6 + A*b^7)*cos(d*x + c))*sin(d*x + c))/(a^8*b^3 - 4*a^6*b^5 + 6*a^4*b^7 - 4*a^2*b^9 + b^11)*d*cos(d*x + c)^3 + 3*(a^9*b^2 - 4*a^7*b^4 + 6*a^5*b^6 - 4*a^3*b^8 + a*b^10)*d*cos(d*x + c)^2 + 3*(a^10*b - 4*a^8*b^3 + 6*a^6*b^5 - 4*a^4*b^7 + a^2*b^9)*d*cos(d*x + c) + (a^11 - 4*a^9*b^2 + 6*a^7*b^4 - 4*a^5*b^6 + a^3*b^8)*d), 1/6*(3*(B*a^6 - 4*A*a^5*b + 4*B*a^4*b^2 - A*a^3*b^3 + (B*a^3*b^3 - 4*A*a^2*b^4 + 4*B*a*b^5 - A*b^6)*cos(d*x + c)^3 + 3*(B*a^4*b^2 - 4*A*a^3*b^3 + 4*B*a^2*b^4 - A*a*b^5)*cos(d*x + c)^2 + 3*(B*a^5*b - 4*A*a^4*b^2 + 4*B*a^3*b^3 - A*a^2*b^4)*cos(d*x + c))*sqrt(a^2 - b^2)*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (6*A*a^7 - 13*B*a^6*b + 4*A*a^5*b^2 + 11*B*a^4*b^3 - 11*A*a^3*b^4 + 2*B*a^2*b^5 + A*a*b^6...
```

3.276.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**4,x)
```

3.276. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

output Timed out

3.276.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.276.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. 2(247) = 494.

Time = 0.33 (sec) , antiderivative size = 722, normalized size of antiderivative = 2.75

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx = \frac{3(Ba^3-4Aa^2b+4Bab^2-Ab^3)\left(\pi\left[\frac{dx+c}{2\pi}+\frac{1}{2}\right]\text{sgn}(-2a+2b)+\arctan\left(-\frac{a\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)-b\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)}{\sqrt{a^2-b^2}}\right)\right)}{(a^6-3a^4b^2+3a^2b^4-b^6)\sqrt{a^2-b^2}} - \frac{6Aa^5\tan\left(\frac{1}{2}dx+\frac{1}{2}c\right)^5-3B}{\dots}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

3.276. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^4} dx$

```

output -1/3*(3*(B*a^3 - 4*A*a^2*b + 4*B*a*b^2 - A*b^3)*(pi*floor(1/2*(d*x + c)/pi
+ 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x
+ 1/2*c))/sqrt(a^2 - b^2)))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*sqrt(a^2
- b^2)) - (6*A*a^5*tan(1/2*d*x + 1/2*c)^5 - 3*B*a^5*tan(1/2*d*x + 1/2*c)^5
- 6*A*a^4*b*tan(1/2*d*x + 1/2*c)^5 - 12*B*a^4*b*tan(1/2*d*x + 1/2*c)^5 +
12*A*a^3*b^2*tan(1/2*d*x + 1/2*c)^5 + 27*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^5
- 27*A*a^2*b^3*tan(1/2*d*x + 1/2*c)^5 - 12*B*a^2*b^3*tan(1/2*d*x + 1/2*c)^
5 + 12*A*a*b^4*tan(1/2*d*x + 1/2*c)^5 + 6*B*a*b^4*tan(1/2*d*x + 1/2*c)^5 +
3*A*b^5*tan(1/2*d*x + 1/2*c)^5 - 6*B*b^5*tan(1/2*d*x + 1/2*c)^5 + 12*A*a^
5*tan(1/2*d*x + 1/2*c)^3 - 28*B*a^4*b*tan(1/2*d*x + 1/2*c)^3 + 16*A*a^3*b^
2*tan(1/2*d*x + 1/2*c)^3 + 16*B*a^2*b^3*tan(1/2*d*x + 1/2*c)^3 - 28*A*a*b^
4*tan(1/2*d*x + 1/2*c)^3 + 12*B*b^5*tan(1/2*d*x + 1/2*c)^3 + 6*A*a^5*tan(1
/2*d*x + 1/2*c) + 3*B*a^5*tan(1/2*d*x + 1/2*c) + 6*A*a^4*b*tan(1/2*d*x + 1
/2*c) - 12*B*a^4*b*tan(1/2*d*x + 1/2*c) + 12*A*a^3*b^2*tan(1/2*d*x + 1/2*c
) - 27*B*a^3*b^2*tan(1/2*d*x + 1/2*c) + 27*A*a^2*b^3*tan(1/2*d*x + 1/2*c)
- 12*B*a^2*b^3*tan(1/2*d*x + 1/2*c) + 12*A*a*b^4*tan(1/2*d*x + 1/2*c) - 6*
B*a*b^4*tan(1/2*d*x + 1/2*c) - 3*A*b^5*tan(1/2*d*x + 1/2*c) - 6*B*b^5*tan(
1/2*d*x + 1/2*c))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*(a*tan(1/2*d*x + 1/
2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^3))/d

```

3.276.9 Mupad [B] (verification not implemented)

Time = 6.63 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.71

$$\begin{aligned}
 & \int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx \\
 &= \frac{4 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 (3Aa^3 - 7Ba^2b + 7Aab^2 - 3Bb^3)}{3(a+b)^2(a^2 - 2ab + b^2)} + \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right) (2Aa^3 - Ab^3 + Ba^3 - 2Bb^3 + 6Aab^2 - 2Aa^2b + 2Bab^2 - 6Ba^2b)}{(a+b)(a^3 - 3a^2b + 3ab^2 - b^3)} + \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)(2a-2b)(a^3 - 3a^2b + 3ab^2 - b^3)}{2\sqrt{a+b}(a-b)^{7/2}}\right) (-Ba^3 + 4Aa^2b - 4Bab^2 + Ab^3)}{d(a+b)^{7/2}(a-b)^{7/2}}
 \end{aligned}$$

```

input int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^4,x)

```

output $((4*\tan(c/2 + (d*x)/2)^3*(3*A*a^3 - 3*B*b^3 + 7*A*a*b^2 - 7*B*a^2*b))/(3*(a + b)^2*(a^2 - 2*a*b + b^2)) + (\tan(c/2 + (d*x)/2)*(2*A*a^3 - A*b^3 + B*a^3 - 2*B*b^3 + 6*A*a*b^2 - 2*A*a^2*b + 2*B*a*b^2 - 6*B*a^2*b))/((a + b)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)) + (\tan(c/2 + (d*x)/2)^5*(2*A*a^3 + A*b^3 - B*a^3 - 2*B*b^3 + 6*A*a*b^2 + 2*A*a^2*b - 2*B*a*b^2 - 6*B*a^2*b))/((a + b)^3*(a - b)))/(d*(3*a*b^2 - \tan(c/2 + (d*x)/2)^4*(3*a*b^2 + 3*a^2*b - 3*a^3 - 3*b^3) - \tan(c/2 + (d*x)/2)^2*(3*a*b^2 - 3*a^2*b - 3*a^3 + 3*b^3) + 3*a^2*b + a^3 + b^3 + \tan(c/2 + (d*x)/2)^6*(3*a*b^2 - 3*a^2*b + a^3 - b^3))) - (\operatorname{atan}((\tan(c/2 + (d*x)/2)*(2*a - 2*b)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)))/(2*(a + b)^{(1/2)}*(a - b)^{(7/2)})))*(A*b^3 - B*a^3 + 4*A*a^2*b - 4*B*a*b^2))/(d*(a + b)^{(7/2)}*(a - b)^{(7/2)})$

3.277 $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^4} dx$

3.277.1 Optimal result 2569
 3.277.2 Mathematica [A] (verified) 2570
 3.277.3 Rubi [A] (verified) 2570
 3.277.4 Maple [A] (verified) 2574
 3.277.5 Fricas [B] (verification not implemented) 2574
 3.277.6 Sympy [F(-1)] 2575
 3.277.7 Maxima [F(-2)] 2576
 3.277.8 Giac [B] (verification not implemented) 2576
 3.277.9 Mupad [B] (verification not implemented) 2577

3.277.1 Optimal result

Integrand size = 23, antiderivative size = 237

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^4} dx = \frac{(2a^3 A + 3aAb^2 - 4a^2bB - b^3B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{(a-b)^{7/2}(a+b)^{7/2}d} - \frac{(Ab - aB) \sin(c + dx)}{3(a^2 - b^2) d(a + b \cos(c + dx))^3} - \frac{(5aAb - 2a^2B - 3b^2B) \sin(c + dx)}{6(a^2 - b^2)^2 d(a + b \cos(c + dx))^2} - \frac{(11a^2Ab + 4Ab^3 - 2a^3B - 13ab^2B) \sin(c + dx)}{6(a^2 - b^2)^3 d(a + b \cos(c + dx))}$$

output

```
(2*A*a^3+3*A*a*b^2-4*B*a^2*b-B*b^3)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/
(a+b)^(1/2))/(a-b)^(7/2)/(a+b)^(7/2)/d-1/3*(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/
d/(a+b*cos(d*x+c))^3-1/6*(5*A*a*b-2*B*a^2-3*B*b^2)*sin(d*x+c)/(a^2-b^2)^2/
d/(a+b*cos(d*x+c))^2-1/6*(11*A*a^2*b+4*A*b^3-2*B*a^3-13*B*a*b^2)*sin(d*x+c
)/(a^2-b^2)^3/d/(a+b*cos(d*x+c))
```


$$\begin{aligned}
& \frac{\int \frac{3(aA-bB)-2(Ab-aB)\cos(c+dx)}{(a+b\cos(c+dx))^3} dx}{3(a^2-b^2)} - \frac{(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3(aA-bB)-2(Ab-aB)\sin(c+dx+\frac{\pi}{2})}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx}{3(a^2-b^2)} - \frac{(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3} \\
& \quad \downarrow \text{3233} \\
& \frac{\int -\frac{2(3Aa^2-5bBa+2Ab^2)-(-2Ba^2+5Aba-3b^2B)\cos(c+dx)}{(a+b\cos(c+dx))^2} dx}{2(a^2-b^2)} - \frac{(-2a^2B+5aAb-3b^2B)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \frac{3(a^2-b^2)(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{2(3Aa^2-5bBa+2Ab^2)-(-2Ba^2+5Aba-3b^2B)\cos(c+dx)}{(a+b\cos(c+dx))^2} dx}{2(a^2-b^2)} - \frac{(-2a^2B+5aAb-3b^2B)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \frac{3(a^2-b^2)(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2(3Aa^2-5bBa+2Ab^2)+(2Ba^2-5Aba+3b^2B)\sin(c+dx+\frac{\pi}{2})}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx}{2(a^2-b^2)} - \frac{(-2a^2B+5aAb-3b^2B)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \frac{3(a^2-b^2)(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3} \\
& \quad \downarrow \text{3233} \\
& \frac{\int -\frac{3(2Aa^3-4bBa^2+3Ab^2a-b^3B)}{a^2-b^2} dx}{2(a^2-b^2)} - \frac{(-2a^3B+11a^2Ab-13ab^2B+4Ab^3)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))} - \frac{(-2a^2B+5aAb-3b^2B)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \frac{3(a^2-b^2)(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3} \\
& \quad \downarrow \text{27} \\
& \frac{3(2a^3A-4a^2bB+3aAb^2-b^3B)}{a^2-b^2} \int \frac{1}{a+b\cos(c+dx)} dx - \frac{(-2a^3B+11a^2Ab-13ab^2B+4Ab^3)\sin(c+dx)}{d(a^2-b^2)(a+b\cos(c+dx))} - \frac{(-2a^2B+5aAb-3b^2B)\sin(c+dx)}{2d(a^2-b^2)(a+b\cos(c+dx))^2} \\
& \quad \frac{3(a^2-b^2)(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^3}
\end{aligned}$$

3.277. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^4} dx$

↓ 3042

$$\frac{3(2a^3A - 4a^2bB + 3aAb^2 - b^3B) \int \frac{1}{a+b \sin(c+dx + \frac{\pi}{2})} dx - \frac{(-2a^3B + 11a^2Ab - 13ab^2B + 4Ab^3) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{a^2-b^2} - \frac{(-2a^2B + 5aAb - 3b^2B) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2}$$

$$\frac{3(a^2-b^2)(Ab-aB) \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3138

$$\frac{6(2a^3A - 4a^2bB + 3aAb^2 - b^3B) \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx)) - \frac{(-2a^3B + 11a^2Ab - 13ab^2B + 4Ab^3) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{d(a^2-b^2)} - \frac{(-2a^2B + 5aAb - 3b^2B) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2}$$

$$\frac{3(a^2-b^2)(Ab-aB) \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 218

$$\frac{6(2a^3A - 4a^2bB + 3aAb^2 - b^3B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - \frac{(-2a^3B + 11a^2Ab - 13ab^2B + 4Ab^3) \sin(c+dx)}{d(a^2-b^2)(a+b \cos(c+dx))}}{d\sqrt{a-b}\sqrt{a+b}(a^2-b^2)} - \frac{(-2a^2B + 5aAb - 3b^2B) \sin(c+dx)}{2d(a^2-b^2)(a+b \cos(c+dx))^2}$$

$$\frac{3(a^2-b^2)(Ab-aB) \sin(c+dx)}{3d(a^2-b^2)(a+b \cos(c+dx))^3}$$

input `Int[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^4,x]`

output `-1/3*((A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x])^3) + (-1/2*((5*a*A*b - 2*a^2*B - 3*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + ((6*(2*a^3*A + 3*a*A*b^2 - 4*a^2*b*B - b^3*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*d) - ((11*a^2*A*b + 4*A*b^3 - 2*a^3*B - 13*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(2*(a^2 - b^2))/(3*(a^2 - b^2))`

3.277.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

3.277.4 Maple [A] (verified)

Time = 1.54 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.57

method	result
derivativedivides	$\frac{(6Aa^2b+3Aab^2+2Ab^3-2Ba^3-2Ba^2b-6Bab^2-Bb^3)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(a-b)(a^3+3a^2b+3ab^2+b^3)} - \frac{4(9Aa^2b+Ab^3-3Ba^3-7Bab^2)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3(a^2-2ab+b^2)(a^2+2ab+b^2)} - \frac{6}{\left(\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a-b\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b\right)^3}$
default	$\frac{(6Aa^2b+3Aab^2+2Ab^3-2Ba^3-2Ba^2b-6Bab^2-Bb^3)\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{(a-b)(a^3+3a^2b+3ab^2+b^3)} - \frac{4(9Aa^2b+Ab^3-3Ba^3-7Bab^2)\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3(a^2-2ab+b^2)(a^2+2ab+b^2)} - \frac{6}{\left(\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a-b\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b\right)^3}$
risch	Expression too large to display

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE)`

output `1/d*(2*(-1/2*(6*A*a^2*b+3*A*a*b^2+2*A*b^3-2*B*a^3-2*B*a^2*b-6*B*a*b^2-B*b^3)/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*tan(1/2*d*x+1/2*c)^5-2/3*(9*A*a^2*b+A*b^3-3*B*a^3-7*B*a*b^2)/(a^2-2*a*b+b^2)/(a^2+2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3-1/2*(6*A*a^2*b-3*A*a*b^2+2*A*b^3-2*B*a^3+2*B*a^2*b-6*B*a*b^2+B*b^3)/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^3+(2*A*a^3+3*A*a*b^2-4*B*a^2*b-B*b^3)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))`

3.277.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 579 vs. 2(222) = 444.

Time = 0.40 (sec) , antiderivative size = 1228, normalized size of antiderivative = 5.18

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="fracas")`

output

```

[-1/12*(3*(2*A*a^6 - 4*B*a^5*b + 3*A*a^4*b^2 - B*a^3*b^3 + (2*A*a^3*b^3 -
4*B*a^2*b^4 + 3*A*a*b^5 - B*b^6)*cos(d*x + c)^3 + 3*(2*A*a^4*b^2 - 4*B*a^3
*b^3 + 3*A*a^2*b^4 - B*a*b^5)*cos(d*x + c)^2 + 3*(2*A*a^5*b - 4*B*a^4*b^2
+ 3*A*a^3*b^3 - B*a^2*b^4)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d
*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c
) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c
) + a^2)) - 2*(6*B*a^7 - 18*A*a^6*b + 4*B*a^5*b^2 + 23*A*a^4*b^3 - 11*B*a^
3*b^4 - 7*A*a^2*b^5 + B*a*b^6 + 2*A*b^7 + (2*B*a^5*b^2 - 11*A*a^4*b^3 + 11
*B*a^3*b^4 + 7*A*a^2*b^5 - 13*B*a*b^6 + 4*A*b^7)*cos(d*x + c)^2 + 3*(2*B*a
^6*b - 9*A*a^5*b^2 + 7*B*a^4*b^3 + 8*A*a^3*b^4 - 10*B*a^2*b^5 + A*a*b^6 +
B*b^7)*cos(d*x + c))*sin(d*x + c))/((a^8*b^3 - 4*a^6*b^5 + 6*a^4*b^7 - 4*a
^2*b^9 + b^11)*d*cos(d*x + c)^3 + 3*(a^9*b^2 - 4*a^7*b^4 + 6*a^5*b^6 - 4*a
^3*b^8 + a*b^10)*d*cos(d*x + c)^2 + 3*(a^10*b - 4*a^8*b^3 + 6*a^6*b^5 - 4*
a^4*b^7 + a^2*b^9)*d*cos(d*x + c) + (a^11 - 4*a^9*b^2 + 6*a^7*b^4 - 4*a^5*
b^6 + a^3*b^8)*d), 1/6*(3*(2*A*a^6 - 4*B*a^5*b + 3*A*a^4*b^2 - B*a^3*b^3 +
(2*A*a^3*b^3 - 4*B*a^2*b^4 + 3*A*a*b^5 - B*b^6)*cos(d*x + c)^3 + 3*(2*A*a
^4*b^2 - 4*B*a^3*b^3 + 3*A*a^2*b^4 - B*a*b^5)*cos(d*x + c)^2 + 3*(2*A*a^5*
b - 4*B*a^4*b^2 + 3*A*a^3*b^3 - B*a^2*b^4)*cos(d*x + c))*sqrt(a^2 - b^2)*a
rctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) + (6*B*a^7 - 1
8*A*a^6*b + 4*B*a^5*b^2 + 23*A*a^4*b^3 - 11*B*a^3*b^4 - 7*A*a^2*b^5 + B...

```

3.277.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**4,x)`

output `Timed out`

3.277.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.277.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 691 vs. $2(222) = 444$.

Time = 0.32 (sec) , antiderivative size = 691, normalized size of antiderivative = 2.92

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^4} dx = \frac{3(2Aa^3 - 4Ba^2b + 3Aab^2 - Bb^3) \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(-2a+2b) + \arctan \left(-\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right)}{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\sqrt{a^2 - b^2}} - \frac{6Ba^5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^5 - 10Ba^4 \tan(\frac{1}{2} dx + \frac{1}{2} c)^3 + 6Ba^3 \tan(\frac{1}{2} dx + \frac{1}{2} c)}{5(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\sqrt{a^2 - b^2}}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

output

```
-1/3*(3*(2*A*a^3 - 4*B*a^2*b + 3*A*a*b^2 - B*b^3)*(pi*floor(1/2*(d*x + c)/
pi + 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*
x + 1/2*c))/sqrt(a^2 - b^2)))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*sqrt(a^
2 - b^2)) - (6*B*a^5*tan(1/2*d*x + 1/2*c)^5 - 18*A*a^4*b*tan(1/2*d*x + 1/2
*c)^5 - 6*B*a^4*b*tan(1/2*d*x + 1/2*c)^5 + 27*A*a^3*b^2*tan(1/2*d*x + 1/2*
c)^5 + 12*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^5 - 6*A*a^2*b^3*tan(1/2*d*x + 1/2
*c)^5 - 27*B*a^2*b^3*tan(1/2*d*x + 1/2*c)^5 + 3*A*a*b^4*tan(1/2*d*x + 1/2*
c)^5 + 12*B*a*b^4*tan(1/2*d*x + 1/2*c)^5 - 6*A*b^5*tan(1/2*d*x + 1/2*c)^5
+ 3*B*b^5*tan(1/2*d*x + 1/2*c)^5 + 12*B*a^5*tan(1/2*d*x + 1/2*c)^3 - 36*A*
a^4*b*tan(1/2*d*x + 1/2*c)^3 + 16*B*a^3*b^2*tan(1/2*d*x + 1/2*c)^3 + 32*A*
a^2*b^3*tan(1/2*d*x + 1/2*c)^3 - 28*B*a*b^4*tan(1/2*d*x + 1/2*c)^3 + 4*A*b
^5*tan(1/2*d*x + 1/2*c)^3 + 6*B*a^5*tan(1/2*d*x + 1/2*c) - 18*A*a^4*b*tan(
1/2*d*x + 1/2*c) + 6*B*a^4*b*tan(1/2*d*x + 1/2*c) - 27*A*a^3*b^2*tan(1/2*d
*x + 1/2*c) + 12*B*a^3*b^2*tan(1/2*d*x + 1/2*c) - 6*A*a^2*b^3*tan(1/2*d*x
+ 1/2*c) + 27*B*a^2*b^3*tan(1/2*d*x + 1/2*c) - 3*A*a*b^4*tan(1/2*d*x + 1/2
*c) + 12*B*a*b^4*tan(1/2*d*x + 1/2*c) - 6*A*b^5*tan(1/2*d*x + 1/2*c) - 3*B
*b^5*tan(1/2*d*x + 1/2*c))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*(a*tan(1/2
*d*x + 1/2*c)^2 - b*tan(1/2*d*x + 1/2*c)^2 + a + b)^3))/d
```

3.277.9 Mupad [B] (verification not implemented)

Time = 4.00 (sec) , antiderivative size = 440, normalized size of antiderivative = 1.86

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{\operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)(2a-2b)(a^3-3a^2b+3ab^2-b^3)}{2\sqrt{a+b}(a-b)^{7/2}}\right)(2Aa^3-4Ba^2b+3Aab^2-Bb^3)}{d(a+b)^{7/2}(a-b)^{7/2}}$$

$$- \frac{\frac{4\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3(-3Ba^3+9Aa^2b-7Bab^2+Ab^3)}{3(a+b)^2(a^2-2ab+b^2)} - \frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5(2Ba^3-2Ab^3+Bb^3-3Aab^2-6Aa^2b+6Bab^2+2Ba^2b)}{(a+b)^3(a-b)}}{d\left(3ab^2 - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4(-3a^3+3a^2b+3ab^2-3b^3) - \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2(-3a^3-3a^2b+3ab^2+3b^3)\right)}$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^4,x)`

output $(\operatorname{atan}((\tan(c/2 + (d*x)/2)*(2*a - 2*b)*(3*a*b^2 - 3*a^2*b + a^3 - b^3))/(2*(a + b)^{(1/2)*(a - b)^{(7/2)})))*(2*A*a^3 - B*b^3 + 3*A*a*b^2 - 4*B*a^2*b))/(d*(a + b)^{(7/2)*(a - b)^{(7/2)}} - ((4*\tan(c/2 + (d*x)/2)^3*(A*b^3 - 3*B*a^3 + 9*A*a^2*b - 7*B*a*b^2))/(3*(a + b)^2*(a^2 - 2*a*b + b^2)) - (\tan(c/2 + (d*x)/2)^5*(2*B*a^3 - 2*A*b^3 + B*b^3 - 3*A*a*b^2 - 6*A*a^2*b + 6*B*a*b^2 + 2*B*a^2*b)))/((a + b)^3*(a - b)) + (\tan(c/2 + (d*x)/2)*(2*A*b^3 - 2*B*a^3 + B*b^3 - 3*A*a*b^2 + 6*A*a^2*b - 6*B*a*b^2 + 2*B*a^2*b))/((a + b)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)))/(d*(3*a*b^2 - \tan(c/2 + (d*x)/2)^4*(3*a*b^2 + 3*a^2*b - 3*a^3 - 3*b^3) - \tan(c/2 + (d*x)/2)^2*(3*a*b^2 - 3*a^2*b - 3*a^3 + 3*b^3) + 3*a^2*b + a^3 + b^3 + \tan(c/2 + (d*x)/2)^6*(3*a*b^2 - 3*a^2*b + a^3 - b^3)))$

3.278 $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$

3.278.1 Optimal result 2579
 3.278.2 Mathematica [A] (verified) 2580
 3.278.3 Rubi [A] (verified) 2580
 3.278.4 Maple [A] (verified) 2585
 3.278.5 Fricas [B] (verification not implemented) 2586
 3.278.6 Sympy [F] 2587
 3.278.7 Maxima [F(-2)] 2588
 3.278.8 Giac [B] (verification not implemented) 2588
 3.278.9 Mupad [B] (verification not implemented) 2589

3.278.1 Optimal result

Integrand size = 29, antiderivative size = 301

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$= - \frac{(8a^6 Ab - 8a^4 Ab^3 + 7a^2 Ab^5 - 2Ab^7 - 2a^7 B - 3a^5 b^2 B) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{a^4(a-b)^{7/2}(a+b)^{7/2}d}$$

$$+ \frac{A \operatorname{arctanh}(\sin(c + dx))}{a^4 d} + \frac{b(Ab - aB) \sin(c + dx)}{3a(a^2 - b^2) d(a + b \cos(c + dx))^3}$$

$$+ \frac{b(8a^2 Ab - 3Ab^3 - 5a^3 B) \sin(c + dx)}{6a^2(a^2 - b^2)^2 d(a + b \cos(c + dx))^2}$$

$$+ \frac{b(26a^4 Ab - 17a^2 Ab^3 + 6Ab^5 - 11a^5 B - 4a^3 b^2 B) \sin(c + dx)}{6a^3(a^2 - b^2)^3 d(a + b \cos(c + dx))}$$

output

```
- (8*A*a^6*b-8*A*a^4*b^3+7*A*a^2*b^5-2*A*b^7-2*B*a^7-3*B*a^5*b^2)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^4/(a-b)^(7/2)/(a+b)^(7/2)/d+A*arctanh(sin(d*x+c))/a^4/d+1/3*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^3+1/6*b*(8*A*a^2*b-3*A*b^3-5*B*a^3)*sin(d*x+c)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^2+1/6*b*(26*A*a^4*b-17*A*a^2*b^3+6*A*b^5-11*B*a^5-4*B*a^3*b^2)*sin(d*x+c)/a^3/(a^2-b^2)^3/d/(a+b*cos(d*x+c))
```

3.278.2 Mathematica [A] (verified)

Time = 1.61 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.22

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{\cos(c + dx)(B + A \sec(c + dx)) \left(\frac{24(-8a^6Ab + 8a^4Ab^3 - 7a^2Ab^5 + 2Ab^7 + 2a^7B + 3a^5b^2B) \operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{7/2}} - 24 \right)}{(-a^2+b^2)^{7/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^4,x]`

output `(Cos[c + d*x]*(B + A*Sec[c + d*x])*((24*(-8*a^6*A*b + 8*a^4*A*b^3 - 7*a^2*A*b^5 + 2*A*b^7 + 2*a^7*B + 3*a^5*b^2*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(7/2) - 24*A*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 24*A*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - (2*a*b*(-7*2*a^6*A*b + 38*a^4*A*b^3 - 5*a^2*A*b^5 - 6*A*b^7 + 36*a^7*B + a^5*b^2*B + 8*a^3*b^4*B + 6*a*b*(-20*a^4*A*b + 15*a^2*A*b^3 - 5*A*b^5 + 9*a^5*B + a^3*b^2*B)*Cos[c + d*x] + b^2*(-26*a^4*A*b + 17*a^2*A*b^3 - 6*A*b^5 + 11*a^5*B + 4*a^3*b^2*B)*Cos[2*(c + d*x)]*Sin[c + d*x]))/(a^2 - b^2)^3*(a + b*Cos[c + d*x])^3)))/(24*a^4*d*(A + B*Cos[c + d*x]))`

3.278.3 Rubi [A] (verified)

Time = 1.86 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.21, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$, Rules used = {3042, 3479, 3042, 3534, 3042, 3534, 27, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^4} dx$$

↓ 3479

3.278. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$

$$\frac{\int \frac{(2b(Ab-aB) \cos^2(c+dx) - 3a(Ab-aB) \cos(c+dx) + 3A(a^2-b^2)) \sec(c+dx)}{(a+b \cos(c+dx))^3} dx}{3a(a^2-b^2)} + \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3042

$$\frac{\int \frac{2b(Ab-aB) \sin(c+dx+\frac{\pi}{2})^2 - 3a(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + 3A(a^2-b^2)}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))^3} dx}{3a(a^2-b^2)} + \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3534

$$\frac{\int \frac{(6A(a^2-b^2)^2 + b(-5Ba^3 + 8Aba^2 - 3Ab^3) \cos^2(c+dx) - 2a(-3Ba^3 + 6Aba^2 - 2b^2Ba - Ab^3) \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx}{2a(a^2-b^2)} + \frac{b(-5a^3B + 8a^2Ab - 3Ab^3) \sin(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

$$\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3} \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3042

$$\frac{\int \frac{6A(a^2-b^2)^2 + b(-5Ba^3 + 8Aba^2 - 3Ab^3) \sin(c+dx+\frac{\pi}{2})^2 - 2a(-3Ba^3 + 6Aba^2 - 2b^2Ba - Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))^2} dx}{2a(a^2-b^2)} + \frac{b(-5a^3B + 8a^2Ab - 3Ab^3) \sin(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2}$$

$$\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3} \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3534

$$\frac{\int \frac{3(2A(a^2-b^2)^3 - a(-2Ba^5 + 6Aba^4 - 3b^2Ba^3 - 2Ab^3a^2 + Ab^5) \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx}{a(a^2-b^2)} + \frac{b(-11a^5B + 26a^4Ab - 4a^3b^2B - 17a^2Ab^3 + 6Ab^5) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))}$$

$$\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3} \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 27

$$3 \int \frac{(2A(a^2-b^2)^3 - a(-2Ba^5 + 6Aba^4 - 3b^2Ba^3 - 2Ab^3a^2 + Ab^5) \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx}{a(a^2-b^2)} + \frac{b(-11a^5B + 26a^4Ab - 4a^3b^2B - 17a^2Ab^3 + 6Ab^5) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))}$$

$$\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3} \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3}$$

↓ 3042

3.278. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$

$$\frac{\frac{\frac{2A(a^2-b^2)^3 \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2(-2a^7B+8a^6Ab-3a^5b^2B-8a^4Ab^3+7a^2Ab^5-2Ab^7) \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{a(a^2-b^2)} + \frac{b(-11a^5B+26a^4Ab-4a^3b^2B-17a^2Ab^3+6Ab^5) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))}}{2a(a^2-b^2)} + \frac{b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^3} + \frac{b(-5a^3B+8a^2Ab-3Ab^3) \sin(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2} + \frac{b(-11a^5B+26a^4Ab-4a^3b^2B-17a^2Ab^3+6Ab^5) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} + \frac{2A(a^2-b^2)^3 \operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2(-2a^7B+8a^6Ab-3a^5b^2B-8a^4Ab^3+7a^2Ab^5-2Ab^7)}{ad\sqrt{a-b}\sqrt{a+b}}}{3a(a^2-b^2)}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^4,x]
```

```
output (b*(A*b - a*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^3) +
((b*(8*a^2*A*b - 3*A*b^3 - 5*a^3*B)*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*(a +
b*Cos[c + d*x])^2) + ((3*((-2*(8*a^6*A*b - 8*a^4*A*b^3 + 7*a^2*A*b^5 - 2*A
*b^7 - 2*a^7*B - 3*a^5*b^2*B)*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a
+ b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + (2*A*(a^2 - b^2)^3*ArcTanh[Sin[c +
d*x]])/(a*d)))/(a*(a^2 - b^2)) + (b*(26*a^4*A*b - 17*a^2*A*b^3 + 6*A*b^5
- 11*a^5*B - 4*a^3*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d
*x])))/(2*a*(a^2 - b^2)))/(3*a*(a^2 - b^2))
```

3.278.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

3.278. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x] *(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int [(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

3.278.4 Maple [A] (verified)

Time = 2.58 (sec) , antiderivative size = 479, normalized size of antiderivative = 1.59

method	result
derivativedivides	$\frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^4} - \frac{\left(\frac{(12A^4b + 4A^3b^2 - 6A^2b^3 - Ab^4 + 2Ab^5 - 6B^4a^5 - 3B^3a^4b - 2B^2a^3b^2)ab\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2(18A^4a^4)}{2(a-b)(a^3 + 3a^2b + 3ab^2 + b^3)}\right)^2}{2}$
default	$\frac{A \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^4} - \frac{\left(\frac{(12A^4b + 4A^3b^2 - 6A^2b^3 - Ab^4 + 2Ab^5 - 6B^4a^5 - 3B^3a^4b - 2B^2a^3b^2)ab\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 2(18A^4a^4)}{2(a-b)(a^3 + 3a^2b + 3ab^2 + b^3)}\right)^2}{2}$
risch	Expression too large to display

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE)
```

$$3.278. \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$$

output $1/d*(A/a^4*\ln(\tan(1/2*d*x+1/2*c)+1)-2/a^4*((-1/2*(12*A*a^4*b+4*A*a^3*b^2-6*A*a^2*b^3-A*a*b^4+2*A*b^5-6*B*a^5-3*B*a^4*b-2*B*a^3*b^2)*a*b/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*\tan(1/2*d*x+1/2*c)^5-2/3*(18*A*a^4*b-11*A*a^2*b^3+3*A*b^5-9*B*a^5-B*a^3*b^2)*a*b/(a^2+2*a*b+b^2)/(a^2-2*a*b+b^2)*\tan(1/2*d*x+1/2*c)^3-1/2*(12*A*a^4*b-4*A*a^3*b^2-6*A*a^2*b^3+A*a*b^4+2*A*b^5-6*B*a^5+3*B*a^4*b-2*B*a^3*b^2)*a*b/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*\tan(1/2*d*x+1/2*c)))/(\tan(1/2*d*x+1/2*c)^2*a-b*\tan(1/2*d*x+1/2*c)^2+a+b)^3+1/2*(8*A*a^6*b-8*A*a^4*b^3+7*A*a^2*b^5-2*A*b^7-2*B*a^7-3*B*a^5*b^2)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*\arctan((a-b)*\tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))-A/a^4*\ln(\tan(1/2*d*x+1/2*c)-1))$

3.278.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1100 vs. $2(285) = 570$.

Time = 37.89 (sec) , antiderivative size = 2269, normalized size of antiderivative = 7.54

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^4,x, algorithm="fricas")`

output `[1/12*(3*(2*B*a^10 - 8*A*a^9*b + 3*B*a^8*b^2 + 8*A*a^7*b^3 - 7*A*a^5*b^5 + 2*A*a^3*b^7 + (2*B*a^7*b^3 - 8*A*a^6*b^4 + 3*B*a^5*b^5 + 8*A*a^4*b^6 - 7*A*a^2*b^8 + 2*A*b^10)*cos(d*x + c)^3 + 3*(2*B*a^8*b^2 - 8*A*a^7*b^3 + 3*B*a^6*b^4 + 8*A*a^5*b^5 - 7*A*a^3*b^7 + 2*A*a*b^9)*cos(d*x + c)^2 + 3*(2*B*a^9*b - 8*A*a^8*b^2 + 3*B*a^7*b^3 + 8*A*a^6*b^4 - 7*A*a^4*b^6 + 2*A*a^2*b^8)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 6*(A*a^11 - 4*A*a^9*b^2 + 6*A*a^7*b^4 - 4*A*a^5*b^6 + A*a^3*b^8 + (A*a^8*b^3 - 4*A*a^6*b^5 + 6*A*a^4*b^7 - 4*A*a^2*b^9 + A*b^11)*cos(d*x + c)^3 + 3*(A*a^9*b^2 - 4*A*a^7*b^4 + 6*A*a^5*b^6 - 4*A*a^3*b^8 + A*a*b^10)*cos(d*x + c)^2 + 3*(A*a^10*b - 4*A*a^8*b^3 + 6*A*a^6*b^5 - 4*A*a^4*b^7 + A*a^2*b^9)*cos(d*x + c)*log(sin(d*x + c) + 1) - 6*(A*a^11 - 4*A*a^9*b^2 + 6*A*a^7*b^4 - 4*A*a^5*b^6 + A*a^3*b^8 + (A*a^8*b^3 - 4*A*a^6*b^5 + 6*A*a^4*b^7 - 4*A*a^2*b^9 + A*b^11)*cos(d*x + c)^3 + 3*(A*a^9*b^2 - 4*A*a^7*b^4 + 6*A*a^5*b^6 - 4*A*a^3*b^8 + A*a*b^10)*cos(d*x + c)^2 + 3*(A*a^10*b - 4*A*a^8*b^3 + 6*A*a^6*b^5 - 4*A*a^4*b^7 + A*a^2*b^9)*cos(d*x + c))*log(-sin(d*x + c) + 1) - 2*(18*B*a^10*b - 36*A*a^9*b^2 - 23*B*a^8*b^3 + 68*A*a^7*b^4 + 7*B*a^6*b^5 - 43*A*a^5*b^6 - 2*B*a^4*b^7 + 11*A*a^3*b^8 + (11*B*a^8*b^3 - 26*A*a^7*b^4 - 7*B*a^6*b^5 + 43*A*a^5*b^6 - 4*B*a^4*b^7 - 23*A*a^3*b^8 + 6*A*a*b^10)*cos(d...`

3.278.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**4,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a + b*cos(c + d*x))**4, x)`

3.278.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.278.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 837 vs. 2(285) = 570.

Time = 0.36 (sec) , antiderivative size = 837, normalized size of antiderivative = 2.78

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

output `1/3*(3*(2*B*a^7 - 8*A*a^6*b + 3*B*a^5*b^2 + 8*A*a^4*b^3 - 7*A*a^2*b^5 + 2*A*b^7)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))/((a^10 - 3*a^8*b^2 + 3*a^6*b^4 - a^4*b^6)*sqrt(a^2 - b^2)) + 3*A*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^4 - 3*A*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^4 - (18*B*a^7*b*tan(1/2*d*x + 1/2*c)^5 - 36*A*a^6*b^2*tan(1/2*d*x + 1/2*c)^5 - 27*B*a^6*b^2*tan(1/2*d*x + 1/2*c)^5 + 60*A*a^5*b^3*tan(1/2*d*x + 1/2*c)^5 + 6*B*a^5*b^3*tan(1/2*d*x + 1/2*c)^5 + 6*A*a^4*b^4*tan(1/2*d*x + 1/2*c)^5 - 3*B*a^4*b^4*tan(1/2*d*x + 1/2*c)^5 - 45*A*a^3*b^5*tan(1/2*d*x + 1/2*c)^5 + 6*B*a^3*b^5*tan(1/2*d*x + 1/2*c)^5 + 6*A*a^2*b^6*tan(1/2*d*x + 1/2*c)^5 + 15*A*a*b^7*tan(1/2*d*x + 1/2*c)^5 - 6*A*b^8*tan(1/2*d*x + 1/2*c)^5 + 36*B*a^7*b*tan(1/2*d*x + 1/2*c)^3 - 72*A*a^6*b^2*tan(1/2*d*x + 1/2*c)^3 - 32*B*a^5*b^3*tan(1/2*d*x + 1/2*c)^3 + 116*A*a^4*b^4*tan(1/2*d*x + 1/2*c)^3 - 4*B*a^3*b^5*tan(1/2*d*x + 1/2*c)^3 - 56*A*a^2*b^6*tan(1/2*d*x + 1/2*c)^3 + 12*A*b^8*tan(1/2*d*x + 1/2*c)^3 + 18*B*a^7*b*tan(1/2*d*x + 1/2*c) - 36*A*a^6*b^2*tan(1/2*d*x + 1/2*c) + 27*B*a^6*b^2*tan(1/2*d*x + 1/2*c) - 60*A*a^5*b^3*tan(1/2*d*x + 1/2*c) + 6*B*a^5*b^3*tan(1/2*d*x + 1/2*c) + 6*A*a^4*b^4*tan(1/2*d*x + 1/2*c) + 3*B*a^4*b^4*tan(1/2*d*x + 1/2*c) + 45*A*a^3*b^5*tan(1/2*d*x + 1/2*c) + 6*B*a^3*b^5*tan(1/2*d*x + 1/2*c) + 6*A*a^2*b^6*tan(1/2*d*x + 1/2*c) - 15*A*a*b^7*tan(1/2*d*x + 1/2*c) - 6*A*b^8*tan(1/2*d*x + 1/2*c))...`

3.278.9 Mupad [B] (verification not implemented)

Time = 19.06 (sec) , antiderivative size = 9727, normalized size of antiderivative = 32.32

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^4),x)`

output

$$\begin{aligned} & (A*\operatorname{atan}(-((A*((8*\tan(c/2 + (d*x)/2)*(4*A^2*a^14 + 8*A^2*b^14 + 4*B^2*a^14 \\ & - 8*A^2*a*b^13 - 8*A^2*a^13*b - 48*A^2*a^2*b^12 + 48*A^2*a^3*b^11 + 117*A^2 \\ & *a^4*b^10 - 120*A^2*a^5*b^9 - 164*A^2*a^6*b^8 + 160*A^2*a^7*b^7 + 156*A^2 \\ & *a^8*b^6 - 120*A^2*a^9*b^5 - 92*A^2*a^10*b^4 + 48*A^2*a^11*b^3 + 44*A^2*a^ \\ & 12*b^2 + 9*B^2*a^10*b^4 + 12*B^2*a^12*b^2 - 32*A*B*a^13*b + 12*A*B*a^5*b^9 \\ & - 34*A*B*a^7*b^7 + 20*A*B*a^9*b^5 - 16*A*B*a^11*b^3)))/(a^16*b + a^17 - a^ \\ & 6*b^11 - a^7*b^10 + 5*a^8*b^9 + 5*a^9*b^8 - 10*a^10*b^7 - 10*a^11*b^6 + 10 \\ & *a^12*b^5 + 10*a^13*b^4 - 5*a^14*b^3 - 5*a^15*b^2) + (A*((8*(4*A*a^21 + 4* \\ & B*a^21 - 4*A*a^8*b^13 + 2*A*a^9*b^12 + 26*A*a^10*b^11 - 14*A*a^11*b^10 - 7 \\ & 0*A*a^12*b^9 + 30*A*a^13*b^8 + 110*A*a^14*b^7 - 30*A*a^15*b^6 - 110*A*a^16 \\ & *b^5 + 20*A*a^17*b^4 + 64*A*a^18*b^3 - 12*A*a^19*b^2 + 6*B*a^12*b^9 - 6*B* \\ & a^13*b^8 - 14*B*a^14*b^7 + 14*B*a^15*b^6 + 6*B*a^16*b^5 - 6*B*a^17*b^4 + 6 \\ & *B*a^18*b^3 - 6*B*a^19*b^2 - 16*A*a^20*b - 4*B*a^20*b)))/(a^19*b + a^20 - a \\ & ^9*b^11 - a^10*b^10 + 5*a^11*b^9 + 5*a^12*b^8 - 10*a^13*b^7 - 10*a^14*b^6 \\ & + 10*a^15*b^5 + 10*a^16*b^4 - 5*a^17*b^3 - 5*a^18*b^2) + (8*A*\tan(c/2 + (d \\ & *x)/2)*(8*a^21*b - 8*a^8*b^14 + 8*a^9*b^13 + 48*a^10*b^12 - 48*a^11*b^11 - \\ & 120*a^12*b^10 + 120*a^13*b^9 + 160*a^14*b^8 - 160*a^15*b^7 - 120*a^16*b^6 \\ & + 120*a^17*b^5 + 48*a^18*b^4 - 48*a^19*b^3 - 8*a^20*b^2)))/(a^4*(a^16*b + \\ & a^17 - a^6*b^11 - a^7*b^10 + 5*a^8*b^9 + 5*a^9*b^8 - 10*a^10*b^7 - 10*a^11 \\ & *b^6 + 10*a^12*b^5 + 10*a^13*b^4 - 5*a^14*b^3 - 5*a^15*b^2))))/a^4)*i)... \end{aligned}$$

3.278. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^4} dx$

3.279 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^4} dx$

3.279.1 Optimal result 2591
 3.279.2 Mathematica [A] (verified) 2592
 3.279.3 Rubi [A] (verified) 2592
 3.279.4 Maple [A] (verified) 2598
 3.279.5 Fricas [B] (verification not implemented) 2598
 3.279.6 Sympy [F] 2599
 3.279.7 Maxima [F(-2)] 2600
 3.279.8 Giac [B] (verification not implemented) 2600
 3.279.9 Mupad [B] (verification not implemented) 2601

3.279.1 Optimal result

Integrand size = 31, antiderivative size = 420

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{b(20a^6 Ab - 35a^4 Ab^3 + 28a^2 Ab^5 - 8Ab^7 - 8a^7 B + 8a^5 b^2 B - 7a^3 b^4 B + 2ab^6 B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) - (4Ab - aB) \operatorname{arctanh}(\sin(c + dx))}{a^5(a - b)^{7/2}(a + b)^{7/2}d}$$

$$+ \frac{(6a^6 A - 65a^4 Ab^2 + 68a^2 Ab^4 - 24Ab^6 + 26a^5 b B - 17a^3 b^3 B + 6ab^5 B) \tan(c + dx)}{6a^4(a^2 - b^2)^3 d}$$

$$+ \frac{b(Ab - aB) \tan(c + dx)}{3a(a^2 - b^2)d(a + b \cos(c + dx))^3} + \frac{b(9a^2 Ab - 4Ab^3 - 6a^3 B + ab^2 B) \tan(c + dx)}{6a^2(a^2 - b^2)^2 d(a + b \cos(c + dx))^2}$$

$$+ \frac{b(12a^4 Ab - 11a^2 Ab^3 + 4Ab^5 - 6a^5 B + 2a^3 b^2 B - ab^4 B) \tan(c + dx)}{2a^3(a^2 - b^2)^3 d(a + b \cos(c + dx))}$$

```
output b*(20*A*a^6*b-35*A*a^4*b^3+28*A*a^2*b^5-8*A*b^7-8*B*a^7+8*B*a^5*b^2-7*B*a^3*b^4+2*B*a*b^6)*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^5/(a-b)^(7/2)/(a+b)^(7/2)/d-(4*A*b-B*a)*arctanh(sin(d*x+c))/a^5/d+1/6*(6*A*a^6-65*A*a^4*b^2+68*A*a^2*b^4-24*A*b^6+26*B*a^5*b-17*B*a^3*b^3+6*B*a*b^5)*tan(d*x+c)/a^4/(a^2-b^2)^3/d+1/3*b*(A*b-B*a)*tan(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^3+1/6*b*(9*A*a^2*b-4*A*b^3-6*B*a^3+B*a*b^2)*tan(d*x+c)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^2+1/2*b*(12*A*a^4*b-11*A*a^2*b^3+4*A*b^5-6*B*a^5+2*B*a^3*b^2-B*a*b^4)*tan(d*x+c)/a^3/(a^2-b^2)^3/d/(a+b*cos(d*x+c))
```

3.279.2 Mathematica [A] (verified)

Time = 2.97 (sec) , antiderivative size = 549, normalized size of antiderivative = 1.31

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{48b(-20a^6Ab + 35a^4Ab^3 - 28a^2Ab^5 + 8Ab^7 + 8a^7B - 8a^5b^2B + 7a^3b^4B - 2ab^6B) \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{7/2}} + 48(4Ab - aB) \log\left(\frac{\cos\left(\frac{c+dx}{2}\right) - \sin\left(\frac{c+dx}{2}\right)}{2} - \sin\left(\frac{c+dx}{2}\right)\right) + 48(-4Ab + aB) \log\left(\frac{\cos\left(\frac{c+dx}{2}\right) + \sin\left(\frac{c+dx}{2}\right)}{2}\right) + (2a(24a^9A - 36a^7Ab^2 - 246a^5A^2b^4 + 318a^3A^3b^6 - 120a^2A^4b^8 + 120a^6b^3B - 90a^4b^5B + 30a^2b^7B + b(72a^8A - 438a^6Ab^2 + 305a^4A^2b^4 + 28a^2A^3b^6 - 72Ab^8 + 144a^7bB - 50a^5b^3B - 7a^3b^5B + 18Ab^7B)) \cos(c + dx) + 6a^2b(6a^6A - 53a^4Ab^2 + 57a^2A^2b^4 - 20Ab^6 + 20a^5bB - 15a^3b^3B + 5a^2b^5B) \cos(2(c + dx)) + 6a^6A^2b^3 \cos(3(c + dx)) - 65a^4A^2b^5 \cos(3(c + dx)) + 68a^2A^2b^7 \cos(3(c + dx)) - 24Ab^9 \cos(3(c + dx)) + 26a^5b^4B \cos(3(c + dx)) - 17a^3b^6B \cos(3(c + dx)) + 6a^2b^8B \cos(3(c + dx))) \tan(c + dx)) / ((a^2 - b^2)^3(a + b \cos(c + dx))^3) / (48a^5d)$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^4,x]`

output `((-48*b*(-20*a^6*A*b + 35*a^4*A*b^3 - 28*a^2*A*b^5 + 8*A*b^7 + 8*a^7*B - 8*a^5*b^2*B + 7*a^3*b^4*B - 2*a*b^6*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/((-a^2 + b^2)^(7/2) + 48*(4*A*b - a*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 48*(-4*A*b + a*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (2*a*(24*a^9*A - 36*a^7*A*b^2 - 246*a^5*A^2*b^4 + 318*a^3*A^3*b^6 - 120*a^2*A^4*b^8 + 120*a^6*b^3*B - 90*a^4*b^5*B + 30*a^2*b^7*B + b*(72*a^8*A - 438*a^6*A*b^2 + 305*a^4*A^2*b^4 + 28*a^2*A^3*b^6 - 72*A*b^8 + 144*a^7*b*B - 50*a^5*b^3*B - 7*a^3*b^5*B + 18*a*b^7*B))*Cos[c + d*x] + 6*a*b^2*(6*a^6*A - 53*a^4*A*b^2 + 57*a^2*A^2*b^4 - 20*A*b^6 + 20*a^5*b*B - 15*a^3*b^3*B + 5*a^2*b^5*B)*Cos[2*(c + d*x)] + 6*a^6*A^2*b^3*Cos[3*(c + d*x)] - 65*a^4*A^2*b^5*Cos[3*(c + d*x)] + 68*a^2*A^2*b^7*Cos[3*(c + d*x)] - 24*A*b^9*Cos[3*(c + d*x)] + 26*a^5*b^4*B*Cos[3*(c + d*x)] - 17*a^3*b^6*B*Cos[3*(c + d*x)] + 6*a^2*b^8*B*Cos[3*(c + d*x)])*Tan[c + d*x])/((a^2 - b^2)^3*(a + b*Cos[c + d*x])^3)/(48*a^5*d)`

3.279.3 Rubi [A] (verified)

Time = 3.05 (sec) , antiderivative size = 476, normalized size of antiderivative = 1.13, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 3479, 3042, 3534, 3042, 3534, 3042, 3534, 27, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx$$

↓ 3042

3.279. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^4} dx$

$$\int \frac{A + B \sin \left(c + dx + \frac{\pi}{2} \right)}{\sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^4} dx$$

↓ 3479

$$\int \frac{(3Aa^2 + bBa - 3(Ab - aB) \cos(c + dx)a - 4Ab^2 + 3b(Ab - aB) \cos^2(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^3} dx + \frac{3a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3042

$$\int \frac{3Aa^2 + bBa - 3(Ab - aB) \sin \left(c + dx + \frac{\pi}{2} \right) a - 4Ab^2 + 3b(Ab - aB) \sin \left(c + dx + \frac{\pi}{2} \right)^2}{\sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^3} dx + \frac{3a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3534

$$\int \frac{(6Aa^4 + 8bBa^3 - 23Ab^2a^2 - 3b^3Ba - 2(-3Ba^3 + 6Aba^2 - 2b^2Ba - Ab^3) \cos(c + dx)a + 12Ab^4 + 2b(-6Ba^3 + 9Aba^2 + b^2Ba - 4Ab^3) \cos^2(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx + \frac{b(-6a^3 + 3a^2b + 2ab^2 - b^3) \tan(c + dx)}{2a(a^2 - b^2)}$$

$$\frac{3a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3042

$$\int \frac{6Aa^4 + 8bBa^3 - 23Ab^2a^2 - 3b^3Ba - 2(-3Ba^3 + 6Aba^2 - 2b^2Ba - Ab^3) \sin \left(c + dx + \frac{\pi}{2} \right) a + 12Ab^4 + 2b(-6Ba^3 + 9Aba^2 + b^2Ba - 4Ab^3) \sin \left(c + dx + \frac{\pi}{2} \right)^2}{\sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^2} dx + \frac{b(-6a^3 + 3a^2b + 2ab^2 - b^3) \tan(c + dx)}{2a(a^2 - b^2)}$$

$$\frac{3a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3534

$$\int \frac{(6Aa^6 + 26bBa^5 - 65Ab^2a^4 - 17b^3Ba^3 + 68Ab^4a^2 + 6b^5Ba - (-6Ba^5 + 18Aba^4 - 8b^2Ba^3 - 7Ab^3a^2 - b^4Ba + 4Ab^5) \cos(c + dx)a - 24Ab^6 + 3b(-6Ba^5 + 12Aba^4 + 2b^2Ba^3 - 2Ab^4a - b^5) \cos^2(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))} dx + \frac{b(-6a^3 + 3a^2b + 2ab^2 - b^3) \tan(c + dx)}{a(a^2 - b^2)}$$

$$\frac{3a(a^2 - b^2) b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3042

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$\int \frac{6Aa^6 + 26bBa^5 - 65Ab^2a^4 - 17b^3Ba^3 + 68Ab^4a^2 + 6b^5Ba - (-6Ba^5 + 18Aba^4 - 8b^2Ba^3 - 7Ab^3a^2 - b^4Ba + 4Ab^5) \sin\left(c + dx + \frac{\pi}{2}\right) a - 24Ab^6 + 3b(-6Ba^5 + 12Aba^4 + 2b^2Ba^3 - 7Ab^3a^2 - b^4Ba + 4Ab^5) \cos\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^2 (a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

$$\frac{2a(a^2 - b^2)}{3a}$$

3534

$$\int -\frac{3(2(a^2 - b^2))^3 (4Ab - aB) - ab(-6Ba^5 + 12Aba^4 + 2b^2Ba^3 - 11Ab^3a^2 - b^4Ba + 4Ab^5) \cos(c + dx)}{a + b \cos(c + dx)} \sec(c + dx) dx + \frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c + dx)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

$$\frac{2a(a^2 - b^2)}{3a(a^2 - b^2)}$$

27

$$\frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c + dx)}{ad} - 3 \int \frac{(2(a^2 - b^2))^3 (4Ab - aB) - ab(-6Ba^5 + 12Aba^4 + 2b^2Ba^3 - 11Ab^3a^2 - b^4Ba + 4Ab^5) \cos(c + dx)}{a + b \cos(c + dx)} dx$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

$$\frac{2a(a^2 - b^2)}{3a(a^2 - b^2)}$$

3042

$$\frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c + dx)}{ad} - 3 \int \frac{2(a^2 - b^2)^3 (4Ab - aB) - ab(-6Ba^5 + 12Aba^4 + 2b^2Ba^3 - 11Ab^3a^2 - b^4Ba + 4Ab^5) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) (a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

$$\frac{2a(a^2 - b^2)}{3a(a^2 - b^2)}$$

3480

$$\frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c + dx)}{ad} - 3 \left(\frac{2(a^2 - b^2)^3 (4Ab - aB) \int \sec(c + dx) dx}{a} - \frac{b(-8a^7B + 20a^6Ab + 8a^5b^2B - 35a^4Ab^3 - 35a^3b^3B + 20a^2b^4B - 8ab^5B + 8b^6B)}{a} \right)$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

$$\frac{2a(a^2 - b^2)}{3a(a^2 - b^2)}$$

3.279. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx$

↓ 3042

$$\frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c+dx)}{ad} - \frac{\left(\frac{2(a^2-b^2)^3(4Ab-aB) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b(-8a^7B + 20a^6Ab + 8a^5b^2B - 35a^4A)}{a} \right)}{a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3138

$$\frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c+dx)}{ad} - \frac{\left(\frac{2(a^2-b^2)^3(4Ab-aB) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b(-8a^7B + 20a^6Ab + 8a^5b^2B - 35a^4A)}{a} \right)}{a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 218

$$\frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c+dx)}{ad} - \frac{\left(\frac{2(a^2-b^2)^3(4Ab-aB) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b(-8a^7B + 20a^6Ab + 8a^5b^2B - 35a^4A)}{a} \right)}{a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 4257

$$\frac{b(Ab - aB) \tan(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3} +$$

$$\frac{b(-6a^3B + 9a^2Ab + ab^2B - 4Ab^3) \tan(c+dx)}{2ad(a^2-b^2)(a+b \cos(c+dx))^2} + \frac{3b(-6a^5B + 12a^4Ab + 2a^3b^2B - 11a^2Ab^3 - ab^4B + 4Ab^5) \tan(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} + \frac{(6a^6A + 26a^5bB - 65a^4Ab^2 - 17a^3b^3B + 68a^2Ab^4 + 6ab^5B - 24Ab^6) \tan(c+dx)}{ad}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^4,x]`

3.279. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^4} dx$

```
output (b*(A*b - a*B)*Tan[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^3) +
((b*(9*a^2*A*b - 4*A*b^3 - 6*a^3*B + a*b^2*B)*Tan[c + d*x])/(2*a*(a^2 - b^
2)*d*(a + b*Cos[c + d*x])^2) + ((3*b*(12*a^4*A*b - 11*a^2*A*b^3 + 4*A*b^5
- 6*a^5*B + 2*a^3*b^2*B - a*b^4*B)*Tan[c + d*x])/(a*(a^2 - b^2)*d*(a + b*C
os[c + d*x])) + ((-3*((-2*b*(20*a^6*A*b - 35*a^4*A*b^3 + 28*a^2*A*b^5 - 8*
A*b^7 - 8*a^7*B + 8*a^5*b^2*B - 7*a^3*b^4*B + 2*a*b^6*B)*ArcTan[(Sqrt[a -
b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + (2*(a^2
- b^2)^3*(4*A*b - a*B)*ArcTanh[Sin[c + d*x]])/(a*d)))/a + ((6*a^6*A - 65*
a^4*A*b^2 + 68*a^2*A*b^4 - 24*A*b^6 + 26*a^5*b*B - 17*a^3*b^3*B + 6*a*b^5*
B)*Tan[c + d*x])/(a*d))/(a*(a^2 - b^2))/(2*a*(a^2 - b^2))/(3*a*(a^2 - b^
2))
```

3.279.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3138 Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b +
(a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.279.4 Maple [A] (verified)

Time = 3.30 (sec) , antiderivative size = 587, normalized size of antiderivative = 1.40

method	result
derivativedivides	$-\frac{A}{a^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(-4Ab+Ba) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^5} + \frac{2b \left(-\frac{(20A a^4 b + 5A a^3 b^2 - 18A a^2 b^3 - 2A a b^4 + 6A b^5 - 12B a^5 - 4B a^4 b + 2(a-b)(a^3 + 3a^2 b + 3a b^2)}{2(a-b)(a^3 + 3a^2 b + 3a b^2)} \right)}{2(a-b)(a^3 + 3a^2 b + 3a b^2)}$
default	$-\frac{A}{a^4 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)} + \frac{(-4Ab+Ba) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{a^5} + \frac{2b \left(-\frac{(20A a^4 b + 5A a^3 b^2 - 18A a^2 b^3 - 2A a b^4 + 6A b^5 - 12B a^5 - 4B a^4 b + 2(a-b)(a^3 + 3a^2 b + 3a b^2)}{2(a-b)(a^3 + 3a^2 b + 3a b^2)} \right)}{2(a-b)(a^3 + 3a^2 b + 3a b^2)}$
risch	Expression too large to display

input `int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE)`

output `1/d*(-A/a^4/(tan(1/2*d*x+1/2*c)+1)+1/a^5*(-4*A*b+B*a)*ln(tan(1/2*d*x+1/2*c)+1)+2*b/a^5*((-1/2*(20*A*a^4*b+5*A*a^3*b^2-18*A*a^2*b^3-2*A*a*b^4+6*A*b^5-12*B*a^5-4*B*a^4*b+6*B*a^3*b^2+B*a^2*b^3-2*B*a*b^4)*a*b/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*tan(1/2*d*x+1/2*c)^5-2/3*(30*A*a^4*b-29*A*a^2*b^3+9*A*b^5-18*B*a^5+11*B*a^3*b^2-3*B*a*b^4)*a*b/(a^2+2*a*b+b^2)/(a^2-2*a*b+b^2)*tan(1/2*d*x+1/2*c)^3-1/2*(20*A*a^4*b-5*A*a^3*b^2-18*A*a^2*b^3+2*A*a*b^4+6*A*b^5-12*B*a^5+4*B*a^4*b+6*B*a^3*b^2-B*a^2*b^3-2*B*a*b^4)*a*b/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^3+1/2*(20*A*a^6*b-35*A*a^4*b^3+28*A*a^2*b^5-8*A*b^7-8*B*a^7+8*B*a^5*b^2-7*B*a^3*b^4+2*B*a*b^6)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))-A/a^4/(tan(1/2*d*x+1/2*c)-1)+(4*A*b-B*a)/a^5*ln(tan(1/2*d*x+1/2*c)-1))`

3.279.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1662 vs. 2(402) = 804.

Time = 76.71 (sec) , antiderivative size = 3393, normalized size of antiderivative = 8.08

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^4,x, algorithm="fricas")`

output `[1/12*(3*((8*B*a^7*b^4 - 20*A*a^6*b^5 - 8*B*a^5*b^6 + 35*A*a^4*b^7 + 7*B*a^3*b^8 - 28*A*a^2*b^9 - 2*B*a*b^10 + 8*A*b^11)*cos(d*x + c)^4 + 3*(8*B*a^8*b^3 - 20*A*a^7*b^4 - 8*B*a^6*b^5 + 35*A*a^5*b^6 + 7*B*a^4*b^7 - 28*A*a^3*b^8 - 2*B*a^2*b^9 + 8*A*a*b^10)*cos(d*x + c)^3 + 3*(8*B*a^9*b^2 - 20*A*a^8*b^3 - 8*B*a^7*b^4 + 35*A*a^6*b^5 + 7*B*a^5*b^6 - 28*A*a^4*b^7 - 2*B*a^3*b^8 + 8*A*a^2*b^9)*cos(d*x + c)^2 + (8*B*a^10*b - 20*A*a^9*b^2 - 8*B*a^8*b^3 + 35*A*a^7*b^4 + 7*B*a^6*b^5 - 28*A*a^5*b^6 - 2*B*a^4*b^7 + 8*A*a^3*b^8)*cos(d*x + c))*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 6*((B*a^9*b^3 - 4*A*a^8*b^4 - 4*B*a^7*b^5 + 16*A*a^6*b^6 + 6*B*a^5*b^7 - 24*A*a^4*b^8 - 4*B*a^3*b^9 + 16*A*a^2*b^10 + B*a*b^11 - 4*A*b^12)*cos(d*x + c)^4 + 3*(B*a^10*b^2 - 4*A*a^9*b^3 - 4*B*a^8*b^4 + 16*A*a^7*b^5 + 6*B*a^6*b^6 - 24*A*a^5*b^7 - 4*B*a^4*b^8 + 16*A*a^3*b^9 + B*a^2*b^10 - 4*A*a*b^11)*cos(d*x + c)^3 + 3*(B*a^11*b - 4*A*a^10*b^2 - 4*B*a^9*b^3 + 16*A*a^8*b^4 + 6*B*a^7*b^5 - 24*A*a^6*b^6 - 4*B*a^5*b^7 + 16*A*a^4*b^8 + B*a^3*b^9 - 4*A*a^2*b^10)*cos(d*x + c)^2 + (B*a^12 - 4*A*a^11*b - 4*B*a^10*b^2 + 16*A*a^9*b^3 + 6*B*a^8*b^4 - 24*A*a^7*b^5 - 4*B*a^6*b^6 + 16*A*a^5*b^7 + B*a^4*b^8 - 4*A*a^3*b^9)*cos(d*x + c))*log(sin(d*x + c) + 1) - 6*((B*a^9*b^3 - 4*A*a^8*b^4 - 4*B*a^7*b^5 + 16*A*a^6*b^6 + 6*B*a^5*b^7 - 24*A*a^4*b^8 - 4*B*a^3*b^9 + 1...`

3.279.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**4,x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a + b*cos(c + d*x))**4, x)`

3.279.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.279.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 996 vs. 2(402) = 804.

Time = 0.37 (sec) , antiderivative size = 996, normalized size of antiderivative = 2.37

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

output

```

1/3*(3*(8*B*a^7*b - 20*A*a^6*b^2 - 8*B*a^5*b^3 + 35*A*a^4*b^4 + 7*B*a^3*b^
5 - 28*A*a^2*b^6 - 2*B*a*b^7 + 8*A*b^8)*(pi*floor(1/2*(d*x + c)/pi + 1/2)*
sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c)
)/sqrt(a^2 - b^2)))/((a^11 - 3*a^9*b^2 + 3*a^7*b^4 - a^5*b^6)*sqrt(a^2 - b
^2)) + (36*B*a^7*b^2*tan(1/2*d*x + 1/2*c)^5 - 60*A*a^6*b^3*tan(1/2*d*x + 1
/2*c)^5 - 60*B*a^6*b^3*tan(1/2*d*x + 1/2*c)^5 + 105*A*a^5*b^4*tan(1/2*d*x
+ 1/2*c)^5 - 6*B*a^5*b^4*tan(1/2*d*x + 1/2*c)^5 + 24*A*a^4*b^5*tan(1/2*d*x
+ 1/2*c)^5 + 45*B*a^4*b^5*tan(1/2*d*x + 1/2*c)^5 - 117*A*a^3*b^6*tan(1/2*
d*x + 1/2*c)^5 - 6*B*a^3*b^6*tan(1/2*d*x + 1/2*c)^5 + 24*A*a^2*b^7*tan(1/2
*d*x + 1/2*c)^5 - 15*B*a^2*b^7*tan(1/2*d*x + 1/2*c)^5 + 42*A*a*b^8*tan(1/2
*d*x + 1/2*c)^5 + 6*B*a*b^8*tan(1/2*d*x + 1/2*c)^5 - 18*A*b^9*tan(1/2*d*x
+ 1/2*c)^5 + 72*B*a^7*b^2*tan(1/2*d*x + 1/2*c)^3 - 120*A*a^6*b^3*tan(1/2*d
*x + 1/2*c)^3 - 116*B*a^5*b^4*tan(1/2*d*x + 1/2*c)^3 + 236*A*a^4*b^5*tan(1
/2*d*x + 1/2*c)^3 + 56*B*a^3*b^6*tan(1/2*d*x + 1/2*c)^3 - 152*A*a^2*b^7*ta
n(1/2*d*x + 1/2*c)^3 - 12*B*a*b^8*tan(1/2*d*x + 1/2*c)^3 + 36*A*b^9*tan(1/
2*d*x + 1/2*c)^3 + 36*B*a^7*b^2*tan(1/2*d*x + 1/2*c) - 60*A*a^6*b^3*tan(1/
2*d*x + 1/2*c) + 60*B*a^6*b^3*tan(1/2*d*x + 1/2*c) - 105*A*a^5*b^4*tan(1/2
*d*x + 1/2*c) - 6*B*a^5*b^4*tan(1/2*d*x + 1/2*c) + 24*A*a^4*b^5*tan(1/2*d*
x + 1/2*c) - 45*B*a^4*b^5*tan(1/2*d*x + 1/2*c) + 117*A*a^3*b^6*tan(1/2*d*x
+ 1/2*c) - 6*B*a^3*b^6*tan(1/2*d*x + 1/2*c) + 24*A*a^2*b^7*tan(1/2*d*x...

```

3.279.9 Mupad [B] (verification not implemented)

Time = 25.79 (sec) , antiderivative size = 13119, normalized size of antiderivative = 31.24

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input

```

int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^4),x)

```


output

$$\begin{aligned}
& ((\tan(c/2 + (d*x)/2)^3*(18*A*a^8 + 72*A*b^8 - 236*A*a^2*b^6 + 47*A*a^3*b^5 \\
& + 273*A*a^4*b^4 - 60*A*a^5*b^3 - 72*A*a^6*b^2 + 3*B*a^2*b^6 + 59*B*a^3*b^5 \\
& - 14*B*a^4*b^4 - 96*B*a^5*b^3 + 36*B*a^6*b^2 - 12*A*a*b^7 - 18*B*a*b^7)) \\
& / (3*a^4*(a + b)^2*(a - b)^3) - (\tan(c/2 + (d*x)/2)^7*(24*A*a^2*b^5 - 8*A*b \\
& ^7 - 2*A*a^7 - 11*A*a^3*b^4 - 26*A*a^4*b^3 + 6*A*a^5*b^2 - B*a^2*b^5 - 6*B \\
& *a^3*b^4 + 4*B*a^4*b^3 + 12*B*a^5*b^2 + 4*A*a*b^6 + 2*A*a^6*b + 2*B*a*b^6) \\
&) / (a^4*(a + b)^3*(a - b)) + (\tan(c/2 + (d*x)/2)^5*(18*A*a^8 + 72*A*b^8 - 2 \\
& 36*A*a^2*b^6 - 47*A*a^3*b^5 + 273*A*a^4*b^4 + 60*A*a^5*b^3 - 72*A*a^6*b^2 \\
& - 3*B*a^2*b^6 + 59*B*a^3*b^5 + 14*B*a^4*b^4 - 96*B*a^5*b^3 - 36*B*a^6*b^2 \\
& + 12*A*a*b^7 - 18*B*a*b^7)) / (3*a^4*(a + b)^3*(a - b)^2) + (\tan(c/2 + (d*x) \\
& /2)*(2*A*a^7 - 8*A*b^7 + 24*A*a^2*b^5 + 11*A*a^3*b^4 - 26*A*a^4*b^3 - 6*A* \\
& a^5*b^2 + B*a^2*b^5 - 6*B*a^3*b^4 - 4*B*a^4*b^3 + 12*B*a^5*b^2 - 4*A*a*b^6 \\
& + 2*A*a^6*b + 2*B*a*b^6)) / (a^4*(a + b)*(a - b)^3)) / (d*(3*a*b^2 + 3*a^2*b \\
& - \tan(c/2 + (d*x)/2)^4*(6*a^2*b - 6*b^3) - \tan(c/2 + (d*x)/2)^2*(6*a*b^2 - \\
& 2*a^3 + 4*b^3) - \tan(c/2 + (d*x)/2)^6*(2*a^3 - 6*a*b^2 + 4*b^3) + a^3 + b \\
& ^3 - \tan(c/2 + (d*x)/2)^8*(3*a*b^2 - 3*a^2*b + a^3 - b^3))) + (\operatorname{atan}((((4 \\
& *A*b - B*a)*((8*(4*B*a^24 + 16*A*a^10*b^14 - 8*A*a^11*b^13 - 104*A*a^12*b^ \\
& 12 + 50*A*a^13*b^11 + 286*A*a^14*b^10 - 126*A*a^15*b^9 - 434*A*a^16*b^8 + \\
& 174*A*a^17*b^7 + 386*A*a^18*b^6 - 146*A*a^19*b^5 - 190*A*a^20*b^4 + 72*A*a \\
& ^21*b^3 + 40*A*a^22*b^2 - 4*B*a^11*b^13 + 2*B*a^12*b^12 + 26*B*a^13*b^11...
\end{aligned}$$

3.280 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$

3.280.1 Optimal result 2603
 3.280.2 Mathematica [A] (verified) 2604
 3.280.3 Rubi [A] (verified) 2605
 3.280.4 Maple [A] (verified) 2612
 3.280.5 Fricas [B] (verification not implemented) 2613
 3.280.6 Sympy [F] 2613
 3.280.7 Maxima [F(-2)] 2614
 3.280.8 Giac [B] (verification not implemented) 2614
 3.280.9 Mupad [B] (verification not implemented) 2615

3.280.1 Optimal result

Integrand size = 31, antiderivative size = 547

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx =$$

$$\frac{b^2(40a^6 Ab - 84a^4 Ab^3 + 69a^2 Ab^5 - 20Ab^7 - 20a^7 B + 35a^5 b^2 B - 28a^3 b^4 B + 8ab^6 B) \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right) + (a^2 A + 20Ab^2 - 8abB) \operatorname{arctanh}(\sin(c + dx))}{a^6(a - b)^{7/2}(a + b)^{7/2}d}$$

$$+ \frac{2a^6 d (24a^6 Ab - 146a^4 Ab^3 + 167a^2 Ab^5 - 60Ab^7 - 6a^7 B + 65a^5 b^2 B - 68a^3 b^4 B + 24ab^6 B) \tan(c + dx)}{6a^5 (a^2 - b^2)^3 d}$$

$$+ \frac{(a^6 A - 23a^4 Ab^2 + 27a^2 Ab^4 - 10Ab^6 + 12a^5 b B - 11a^3 b^3 B + 4ab^5 B) \sec(c + dx) \tan(c + dx)}{2a^4 (a^2 - b^2)^3 d}$$

$$+ \frac{b(Ab - aB) \sec(c + dx) \tan(c + dx)}{3a (a^2 - b^2) d(a + b \cos(c + dx))^3}$$

$$+ \frac{b(10a^2 Ab - 5Ab^3 - 7a^3 B + 2ab^2 B) \sec(c + dx) \tan(c + dx)}{6a^2 (a^2 - b^2)^2 d(a + b \cos(c + dx))^2}$$

$$+ \frac{b(48a^4 Ab - 53a^2 Ab^3 + 20Ab^5 - 27a^5 B + 20a^3 b^2 B - 8ab^4 B) \sec(c + dx) \tan(c + dx)}{6a^3 (a^2 - b^2)^3 d(a + b \cos(c + dx))}$$

output
$$-b^2(40Aa^6b-84Aa^4b^3+69Aa^2b^5-20Ab^7-20B^2a^7+35B^2a^5b^2-28B^2a^3b^4+8B^2a^2b^6)\arctan((a-b)^{1/2}\tan(1/2dx+1/2c)/(a+b)^{1/2})/a^6/(a-b)^{7/2}/(a+b)^{7/2}/d+1/2(Aa^2+20Ab^2-8B^2a^2b)\operatorname{arctanh}(\sin(dx+c))/a^6/d-1/6(24Aa^6b-146Aa^4b^3+167Aa^2b^5-60Ab^7-6B^2a^7+65B^2a^5b^2-68B^2a^3b^4+24B^2a^2b^6)\tan(dx+c)/a^5/(a^2-b^2)^3/d+1/2(Aa^6-23Aa^4b^2+27Aa^2b^4-10Ab^6+12B^2a^5b-11B^2a^3b^3+4B^2a^2b^5)\sec(dx+c)\tan(dx+c)/a^4/(a^2-b^2)^3/d+1/3b(Ab-Ba)\sec(dx+c)\tan(dx+c)/a/(a^2-b^2)/d/(a+b\cos(dx+c))^3+1/6b(10Aa^2b-5Ab^3-7B^2a^3+2B^2a^2b^2)\sec(dx+c)\tan(dx+c)/a^2/(a^2-b^2)^2/d/(a+b\cos(dx+c))^2+1/6b(48Aa^4b-53Aa^2b^3+20Ab^5-27B^2a^5+20B^2a^3b^2-8B^2a^2b^4)\sec(dx+c)\tan(dx+c)/a^3/(a^2-b^2)^3/d/(a+b\cos(dx+c))$$

3.280.2 Mathematica [A] (verified)

Time = 4.64 (sec) , antiderivative size = 781, normalized size of antiderivative = 1.43

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx$$

$$= \frac{96b^2(-40a^6Ab+84a^4Ab^3-69a^2Ab^5+20Ab^7+20a^7B-35a^5b^2B+28a^3b^4B-8ab^6B)\operatorname{arctanh}\left(\frac{(a-b)\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{7/2}} - 48(a^2A + 20Ab^2)$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^4,x]`

output $((96*b^2*(-40*a^6*A*b + 84*a^4*A*b^3 - 69*a^2*A*b^5 + 20*A*b^7 + 20*a^7*B - 35*a^5*b^2*B + 28*a^3*b^4*B - 8*a*b^6*B)*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^{(7/2)} - 48*(a^2*A + 20*A*b^2 - 8*a*b*B)*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 48*(a^2*A + 20*A*b^2 - 8*a*b*B)*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + (2*a*(24*a^{10}*A - 324*a^8*A*b^2 + 1116*a^6*A*b^4 - 830*a^4*A*b^6 - 61*a^2*A*b^8 + 180*A*b^{10} + 72*a^9*b*B - 438*a^7*b^3*B + 305*a^5*b^5*B + 28*a^3*b^7*B - 72*a*b^9*B + 6*a*(-20*a^8*A*b - 9*a^6*A*b^3 + 309*a^4*A*b^5 - 400*a^2*A*b^7 + 150*A*b^9 + 8*a^9*B - 6*a^7*b^2*B - 135*a^5*b^4*B + 163*a^3*b^6*B - 60*a*b^8*B)*Cos[c + d*x] + 12*b*(-21*a^8*A*b + 85*a^6*A*b^3 - 55*a^4*A*b^5 - 19*a^2*A*b^7 + 20*A*b^9 + 6*a^9*B - 36*a^7*b^2*B + 20*a^5*b^4*B + 8*a^3*b^6*B - 8*a*b^8*B)*Cos[2*(c + d*x)] - 138*a^7*A*b^3*Cos[3*(c + d*x)] + 738*a^5*A*b^5*Cos[3*(c + d*x)] - 840*a^3*A*b^7*Cos[3*(c + d*x)] + 300*a*A*b^9*Cos[3*(c + d*x)] + 36*a^8*b^2*B*Cos[3*(c + d*x)] - 318*a^6*b^4*B*Cos[3*(c + d*x)] + 342*a^4*b^6*B*Cos[3*(c + d*x)] - 120*a^2*b^8*B*Cos[3*(c + d*x)] - 24*a^6*A*b^4*Cos[4*(c + d*x)] + 146*a^4*A*b^6*Cos[4*(c + d*x)] - 167*a^2*A*b^8*Cos[4*(c + d*x)] + 60*A*b^{10}*Cos[4*(c + d*x)] + 6*a^7*b^3*B*Cos[4*(c + d*x)] - 65*a^5*b^5*B*Cos[4*(c + d*x)] + 68*a^3*b^7*B*Cos[4*(c + d*x)] - 24*a*b^9*B*Cos[4*(c + d*x)])*Sec[c + d*x]*Tan[c + d*x])/((a^2 - b^2)^3*(a + b*Cos[c + d*x])^3))/(96*a^6*d)$

3.280.3 Rubi [A] (verified)

Time = 3.91 (sec) , antiderivative size = 591, normalized size of antiderivative = 1.08, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 3479, 3042, 3534, 3042, 3534, 3042, 3534, 27, 3042, 3534, 27, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^4} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3 (a + b \sin(c + dx + \frac{\pi}{2}))^4} dx$$

↓ 3479

3.280. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$

$$\frac{\int \frac{(3Aa^2+2bBa-3(Ab-aB)\cos(c+dx)a-5Ab^2+4b(Ab-aB)\cos^2(c+dx))\sec^3(c+dx)}{(a+b\cos(c+dx))^3} dx}{3a(a^2-b^2)} + \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

$$\frac{\int \frac{3Aa^2+2bBa-3(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-5Ab^2+4b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx}{3a(a^2-b^2)} + \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3534

$$\frac{\int \frac{(3b(-7Ba^3+10Aba^2+2b^2Ba-5Ab^3)\cos^2(c+dx)-2a(-3Ba^3+6Aba^2-2b^2Ba-Ab^3)\cos(c+dx)+2(3Aa^4+9bBa^3-18Ab^2a^2-4b^3Ba+10Ab^4))\sec^3(c+dx)}{(a+b\cos(c+dx))^2} dx}{2a(a^2-b^2)}$$

$3a(a^2-b^2)$

$$\frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

$$\frac{\int \frac{3b(-7Ba^3+10Aba^2+2b^2Ba-5Ab^3)\sin(c+dx+\frac{\pi}{2})^2-2a(-3Ba^3+6Aba^2-2b^2Ba-Ab^3)\sin(c+dx+\frac{\pi}{2})+2(3Aa^4+9bBa^3-18Ab^2a^2-4b^3Ba+10Ab^4)}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx}{2a(a^2-b^2)} + \frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^3}$$

$3a(a^2-b^2)$

$$\frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3534

$$\frac{\int \frac{(2b(-27Ba^5+48Aba^4+20b^2Ba^3-53Ab^3a^2-8b^4Ba+20Ab^5)\cos^2(c+dx)-a(-6Ba^5+18Aba^4-7b^2Ba^3-8Ab^3a^2-2b^4Ba+5Ab^5)\cos(c+dx)+6(Aa^6+12bBa^5-2b^2Aa^4+4b^3Ba^3-2b^4Ba^2+2b^5A))\sec^3(c+dx)}{a+b\cos(c+dx)} dx}{a(a^2-b^2)}$$

$2a(a^2-b^2)$

$$\frac{b(Ab-aB)\tan(c+dx)\sec(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^3}$$

↓ 3042

3.280. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+b\cos(c+dx))^4} dx$

$$\int \frac{2b(-27Ba^5 + 48Aba^4 + 20b^2Ba^3 - 53Ab^3a^2 - 8b^4Ba + 20Ab^5) \sin(c+dx + \frac{\pi}{2})^2 - a(-6Ba^5 + 18Aba^4 - 7b^2Ba^3 - 8Ab^3a^2 - 2b^4Ba + 5Ab^5) \sin(c+dx + \frac{\pi}{2}) + 6(Aa^6 + 12a^5bB - 23a^4Ab^2 - 11a^3b^3B + 27a^2Ab^4 + 4ab^5B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{\sin(c+dx + \frac{\pi}{2})^3 \frac{(a+b \sin(c+dx + \frac{\pi}{2}))}{a(a^2-b^2)}} dx$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3534

$$\int -\frac{2(-6Ba^7 + 24Aba^6 + 65b^2Ba^5 - 146Ab^3a^4 - 68b^4Ba^3 + 167Ab^5a^2 + 24b^6Ba - (3Aa^6 - 18bBa^5 + 27Ab^2a^4 + 7b^3Ba^3 - 25Ab^4a^2 - 4b^5Ba + 10Ab^6) \cos(c+dx)a - 60Ab^7)}{2a(a+b \cos(c+dx))^3} dx$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 27

$$\frac{3(a^6A + 12a^5bB - 23a^4Ab^2 - 11a^3b^3B + 27a^2Ab^4 + 4ab^5B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \int \frac{(-6Ba^7 + 24Aba^6 + 65b^2Ba^5 - 146Ab^3a^4 - 68b^4Ba^3 + 167Ab^5a^2 + 24b^6Ba)}{2a(a+b \cos(c+dx))^3} dx$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3042

$$\frac{3(a^6A + 12a^5bB - 23a^4Ab^2 - 11a^3b^3B + 27a^2Ab^4 + 4ab^5B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \int \frac{-6Ba^7 + 24Aba^6 + 65b^2Ba^5 - 146Ab^3a^4 - 68b^4Ba^3 + 167Ab^5a^2 + 24b^6Ba}{2a(a+b \cos(c+dx))^3} dx$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

↓ 3534

$$\frac{3(a^6A + 12a^5bB - 23a^4Ab^2 - 11a^3b^3B + 27a^2Ab^4 + 4ab^5B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \int \frac{3((Aa^2 - 8bBa + 20Ab^2)(a^2 - b^2))^3 + ab(Aa^6 + 12bBa^5 - 23Ab^2a^4 - 11a^3b^3B + 27a^2Ab^4 + 4ab^5B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{a(a+b \cos(c+dx))^3} dx$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^3}$$

3.280. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$

↓ 27

$$\frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-6a^7 B + 24a^6 Ab + 65a^5 b^2 B - 146a^4 Ab^3 - 68a^3 b^4 B + 167a^2 Ab^5 + 24ab^6 B)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3}$$

↓ 3042

$$\frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-6a^7 B + 24a^6 Ab + 65a^5 b^2 B - 146a^4 Ab^3 - 68a^3 b^4 B + 167a^2 Ab^5 + 24ab^6 B)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3}$$

↓ 3480

$$\frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-6a^7 B + 24a^6 Ab + 65a^5 b^2 B - 146a^4 Ab^3 - 68a^3 b^4 B + 167a^2 Ab^5 + 24ab^6 B)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3}$$

↓ 3042

$$\frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-6a^7 B + 24a^6 Ab + 65a^5 b^2 B - 146a^4 Ab^3 - 68a^3 b^4 B + 167a^2 Ab^5 + 24ab^6 B)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3}$$

↓ 3138

3.280. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$

$$\frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-6a^7 B + 24a^6 Ab + 65a^5 b^2 B - 146a^4 Ab^3 - 68a^3 b^4 B + 167a^2 Ab^5 + 24ab^6 B)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3}$$

↓ 218

$$\frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad} - \frac{(-6a^7 B + 24a^6 Ab + 65a^5 b^2 B - 146a^4 Ab^3 - 68a^3 b^4 B + 167a^2 Ab^5 + 24ab^6 B)}{ad}$$

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3}$$

↓ 4257

$$\frac{b(Ab - aB) \tan(c + dx) \sec(c + dx)}{3ad (a^2 - b^2) (a + b \cos(c + dx))^3} +$$

$$\frac{b(-7a^3 B + 10a^2 Ab + 2ab^2 B - 5Ab^3) \tan(c+dx) \sec(c+dx)}{2ad(a^2 - b^2)(a + b \cos(c+dx))^2} + \frac{b(-27a^5 B + 48a^4 Ab + 20a^3 b^2 B - 53a^2 Ab^3 - 8ab^4 B + 20Ab^5) \tan(c+dx) \sec(c+dx)}{ad(a^2 - b^2)(a + b \cos(c+dx))} + \frac{3(a^6 A + 12a^5 bB - 23a^4 Ab^2 - 11a^3 b^3 B + 27a^2 Ab^4 + 4ab^5 B - 10Ab^6) \tan(c+dx) \sec(c+dx)}{ad}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^4,x]`

$$3.280. \quad \int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$$

output $(b*(A*b - a*B)*\text{Sec}[c + d*x]*\text{Tan}[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*\text{Cos}[c + d*x])^3) + ((b*(10*a^2*A*b - 5*A*b^3 - 7*a^3*B + 2*a*b^2*B)*\text{Sec}[c + d*x]*\text{Tan}[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*\text{Cos}[c + d*x])^2) + ((b*(48*a^4*A*b - 53*a^2*A*b^3 + 20*A*b^5 - 27*a^5*B + 20*a^3*b^2*B - 8*a*b^4*B)*\text{Sec}[c + d*x]*\text{Tan}[c + d*x])/(a*(a^2 - b^2)*d*(a + b*\text{Cos}[c + d*x]))) + ((3*(a^6*A - 23*a^4*A*b^2 + 27*a^2*A*b^4 - 10*A*b^6 + 12*a^5*b*B - 11*a^3*b^3*B + 4*a*b^5*B)*\text{Sec}[c + d*x]*\text{Tan}[c + d*x])/(a*d) - ((-3*((-2*b^2*(40*a^6*A*b - 84*a^4*A*b^3 + 69*a^2*A*b^5 - 20*A*b^7 - 20*a^7*B + 35*a^5*b^2*B - 28*a^3*b^4*B + 8*a*b^6*B)*\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Tan}[(c + d*x)/2])/\text{Sqrt}[a + b]])/(a*\text{Sqrt}[a - b]*\text{Sqrt}[a + b]*d) + ((a^2 - b^2)^3*(a^2*A + 20*A*b^2 - 8*a*b*B)*\text{ArcTan}[\text{h}[\text{Sin}[c + d*x]])/(a*d)))/a + ((24*a^6*A*b - 146*a^4*A*b^3 + 167*a^2*A*b^5 - 60*A*b^7 - 6*a^7*B + 65*a^5*b^2*B - 68*a^3*b^4*B + 24*a*b^6*B)*\text{Tan}[c + d*x])/(a*d))/a/(a*(a^2 - b^2))/(2*a*(a^2 - b^2))/(3*a*(a^2 - b^2))$

3.280.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$

rule 218 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3138 $\text{Int}[(a_*) + (b_*)\sin[\text{Pi}/2 + (c_*) + (d_*)(x_)]^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \quad \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.280.4 Maple [A] (verified)

Time = 5.06 (sec) , antiderivative size = 672, normalized size of antiderivative = 1.23

method	result
derivativedivides	$2b^2 \left(-\frac{(30Aa^4b+6Aa^3b^2-34Aa^2b^3-3Aab^4+12Ab^5-20Ba^5-5Ba^4b+18Ba^3b^2+2Ba^2b^3-6Bab^4)ab\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)} - \frac{2(45Aa^4b+6Aa^3b^2-34Aa^2b^3-3Aab^4+12Ab^5-20Ba^5-5Ba^4b+18Ba^3b^2+2Ba^2b^3-6Bab^4)ab\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)} \right)$
default	$2b^2 \left(-\frac{(30Aa^4b+6Aa^3b^2-34Aa^2b^3-3Aab^4+12Ab^5-20Ba^5-5Ba^4b+18Ba^3b^2+2Ba^2b^3-6Bab^4)ab\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)} - \frac{2(45Aa^4b+6Aa^3b^2-34Aa^2b^3-3Aab^4+12Ab^5-20Ba^5-5Ba^4b+18Ba^3b^2+2Ba^2b^3-6Bab^4)ab\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{2(a-b)(a^3+3a^2b+3ab^2+b^3)} \right)$
risch	Expression too large to display

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-2*b^2/a^6*((-1/2*(30*A*a^4*b+6*A*a^3*b^2-34*A*a^2*b^3-3*A*a*b^4+12*A*b^5-20*B*a^5-5*B*a^4*b+18*B*a^3*b^2+2*B*a^2*b^3-6*B*a*b^4)*a*b/(a-b)/(a^3+3*a^2*b+3*a*b^2+b^3)*tan(1/2*d*x+1/2*c)^5-2/3*(45*A*a^4*b-53*A*a^2*b^3+18*A*b^5-30*B*a^5+29*B*a^3*b^2-9*B*a*b^4)*a*b/(a^2+2*a*b+b^2)/(a^2-2*a*b+b^2))*tan(1/2*d*x+1/2*c)^3-1/2*(30*A*a^4*b-6*A*a^3*b^2-34*A*a^2*b^3+3*A*a*b^4+12*A*b^5-20*B*a^5+5*B*a^4*b+18*B*a^3*b^2-2*B*a^2*b^3-6*B*a*b^4)*a*b/(a+b)/(a^3-3*a^2*b+3*a*b^2-b^3)*tan(1/2*d*x+1/2*c))/(tan(1/2*d*x+1/2*c)^2*a-b*tan(1/2*d*x+1/2*c)^2+a+b)^3+1/2*(40*A*a^6*b-84*A*a^4*b^3+69*A*a^2*b^5-20*A*b^7-20*B*a^7+35*B*a^5*b^2-28*B*a^3*b^4+8*B*a*b^6)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2)))-1/2*A/a^4/(tan(1/2*d*x+1/2*c)+1)^2-1/2*(-A*a-8*A*b+2*B*a)/a^5/(tan(1/2*d*x+1/2*c)+1)+1/2*(A*a^2+20*A*b^2-8*B*a*b)/a^6*ln(tan(1/2*d*x+1/2*c)+1)+1/2*A/a^4/(tan(1/2*d*x+1/2*c)-1)^2-1/2*(-A*a-8*A*b+2*B*a)/a^5/(tan(1/2*d*x+1/2*c)-1)+1/2/a^6*(-A*a^2-20*A*b^2+8*B*a*b)*ln(tan(1/2*d*x+1/2*c)-1))
```

3.280. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^4} dx$

3.280.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1875 vs. $2(525) = 1050$.

Time = 113.16 (sec) , antiderivative size = 3819, normalized size of antiderivative = 6.98

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^4,x, algorithm="fricas")
```

```
output [1/12*(3*((20*B*a^7*b^5 - 40*A*a^6*b^6 - 35*B*a^5*b^7 + 84*A*a^4*b^8 + 28*B*a^3*b^9 - 69*A*a^2*b^10 - 8*B*a*b^11 + 20*A*b^12)*cos(d*x + c)^5 + 3*(20*B*a^8*b^4 - 40*A*a^7*b^5 - 35*B*a^6*b^6 + 84*A*a^5*b^7 + 28*B*a^4*b^8 - 69*A*a^3*b^9 - 8*B*a^2*b^10 + 20*A*a*b^11)*cos(d*x + c)^4 + 3*(20*B*a^9*b^3 - 40*A*a^8*b^4 - 35*B*a^7*b^5 + 84*A*a^6*b^6 + 28*B*a^5*b^7 - 69*A*a^4*b^8 - 8*B*a^3*b^9 + 20*A*a^2*b^10)*cos(d*x + c)^3 + (20*B*a^10*b^2 - 40*A*a^9*b^3 - 35*B*a^8*b^4 + 84*A*a^7*b^5 + 28*B*a^6*b^6 - 69*A*a^5*b^7 - 8*B*a^4*b^8 + 20*A*a^3*b^9)*cos(d*x + c)^2)*sqrt(-a^2 + b^2)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 3*((A*a^10*b^3 - 8*B*a^9*b^4 + 16*A*a^8*b^5 + 32*B*a^7*b^6 - 74*A*a^6*b^7 - 48*B*a^5*b^8 + 116*A*a^4*b^9 + 32*B*a^3*b^10 - 79*A*a^2*b^11 - 8*B*a*b^12 + 20*A*b^13)*cos(d*x + c)^5 + 3*(A*a^11*b^2 - 8*B*a^10*b^3 + 16*A*a^9*b^4 + 32*B*a^8*b^5 - 74*A*a^7*b^6 - 48*B*a^6*b^7 + 116*A*a^5*b^8 + 32*B*a^4*b^9 - 79*A*a^3*b^10 - 8*B*a^2*b^11 + 20*A*a*b^12)*cos(d*x + c)^4 + 3*(A*a^12*b - 8*B*a^11*b^2 + 16*A*a^10*b^3 + 32*B*a^9*b^4 - 74*A*a^8*b^5 - 48*B*a^7*b^6 + 116*A*a^6*b^7 + 32*B*a^5*b^8 - 79*A*a^4*b^9 - 8*B*a^3*b^10 + 20*A*a^2*b^11)*cos(d*x + c)^3 + (A*a^13 - 8*B*a^12*b + 16*A*a^11*b^2 + 32*B*a^10*b^3 - 74*A*a^9*b^4 - 48*B*a^8*b^5 + 116*A*a^7*b^6 + 32*B*a^6*b^7 - 79*A*a^5*b^8 - 8*B*a^4*b^9 + 20*A*a^3*b^10)*cos(d*x + c)^2)*log(si...
```

3.280.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**4,x)
```

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a + b*cos(c + d*x))**4, x)`

3.280.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.280.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1090 vs. $2(525) = 1050$.

Time = 0.37 (sec) , antiderivative size = 1090, normalized size of antiderivative = 1.99

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^4,x, algorithm="giac")`

output

```

-1/6*(6*(20*B*a^7*b^2 - 40*A*a^6*b^3 - 35*B*a^5*b^4 + 84*A*a^4*b^5 + 28*B*
a^3*b^6 - 69*A*a^2*b^7 - 8*B*a*b^8 + 20*A*b^9)*(pi*floor(1/2*(d*x + c)/pi
+ 1/2)*sgn(-2*a + 2*b) + arctan(-(a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x +
1/2*c))/sqrt(a^2 - b^2)))/((a^12 - 3*a^10*b^2 + 3*a^8*b^4 - a^6*b^6)*sqrt
(a^2 - b^2)) + 2*(60*B*a^7*b^3*tan(1/2*d*x + 1/2*c)^5 - 90*A*a^6*b^4*tan(1
/2*d*x + 1/2*c)^5 - 105*B*a^6*b^4*tan(1/2*d*x + 1/2*c)^5 + 162*A*a^5*b^5*t
an(1/2*d*x + 1/2*c)^5 - 24*B*a^5*b^5*tan(1/2*d*x + 1/2*c)^5 + 48*A*a^4*b^6
*tan(1/2*d*x + 1/2*c)^5 + 117*B*a^4*b^6*tan(1/2*d*x + 1/2*c)^5 - 213*A*a^3
*b^7*tan(1/2*d*x + 1/2*c)^5 - 24*B*a^3*b^7*tan(1/2*d*x + 1/2*c)^5 + 48*A*a
^2*b^8*tan(1/2*d*x + 1/2*c)^5 - 42*B*a^2*b^8*tan(1/2*d*x + 1/2*c)^5 + 81*A
*a*b^9*tan(1/2*d*x + 1/2*c)^5 + 18*B*a*b^9*tan(1/2*d*x + 1/2*c)^5 - 36*A*b
^10*tan(1/2*d*x + 1/2*c)^5 + 120*B*a^7*b^3*tan(1/2*d*x + 1/2*c)^3 - 180*A*
a^6*b^4*tan(1/2*d*x + 1/2*c)^3 - 236*B*a^5*b^5*tan(1/2*d*x + 1/2*c)^3 + 39
2*A*a^4*b^6*tan(1/2*d*x + 1/2*c)^3 + 152*B*a^3*b^7*tan(1/2*d*x + 1/2*c)^3
- 284*A*a^2*b^8*tan(1/2*d*x + 1/2*c)^3 - 36*B*a*b^9*tan(1/2*d*x + 1/2*c)^3
+ 72*A*b^10*tan(1/2*d*x + 1/2*c)^3 + 60*B*a^7*b^3*tan(1/2*d*x + 1/2*c) -
90*A*a^6*b^4*tan(1/2*d*x + 1/2*c) + 105*B*a^6*b^4*tan(1/2*d*x + 1/2*c) - 1
62*A*a^5*b^5*tan(1/2*d*x + 1/2*c) - 24*B*a^5*b^5*tan(1/2*d*x + 1/2*c) + 48
*A*a^4*b^6*tan(1/2*d*x + 1/2*c) - 117*B*a^4*b^6*tan(1/2*d*x + 1/2*c) + 213
*A*a^3*b^7*tan(1/2*d*x + 1/2*c) - 24*B*a^3*b^7*tan(1/2*d*x + 1/2*c) + 4...

```

3.280.9 Mupad [B] (verification not implemented)

Time = 14.28 (sec) , antiderivative size = 14398, normalized size of antiderivative = 26.32

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^4} dx = \text{Too large to display}$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^4),x)`

output

$$\begin{aligned} & ((\tan(c/2 + (d*x)/2)*(A*a^8 + 20*A*b^8 + 2*B*a^8 - 59*A*a^2*b^6 - 27*A*a^3 \\ & *b^5 + 57*A*a^4*b^4 + 21*A*a^5*b^3 - 11*A*a^6*b^2 - 4*B*a^2*b^6 + 24*B*a^3 \\ & *b^5 + 11*B*a^4*b^4 - 26*B*a^5*b^3 - 6*B*a^6*b^2 + 10*A*a*b^7 - 7*A*a^7*b \\ & - 8*B*a*b^7 + 2*B*a^7*b))/(a^5*(a + b)*(a - b)^3) + (2*\tan(c/2 + (d*x)/2)^ \\ & 5*(9*A*a^10 + 180*A*b^10 - 611*A*a^2*b^8 + 740*A*a^4*b^6 - 324*A*a^6*b^4 + \\ & 36*A*a^8*b^2 + 248*B*a^3*b^7 - 320*B*a^5*b^5 + 132*B*a^7*b^3 - 72*B*a*b^9 \\ & - 18*B*a^9*b))/(3*a^5*(a + b)^3*(a - b)^3) + (\tan(c/2 + (d*x)/2)^9*(A*a^8 \\ & + 20*A*b^8 - 2*B*a^8 - 59*A*a^2*b^6 + 27*A*a^3*b^5 + 57*A*a^4*b^4 - 21*A* \\ & a^5*b^3 - 11*A*a^6*b^2 + 4*B*a^2*b^6 + 24*B*a^3*b^5 - 11*B*a^4*b^4 - 26*B* \\ & a^5*b^3 + 6*B*a^6*b^2 - 10*A*a*b^7 + 7*A*a^7*b - 8*B*a*b^7 + 2*B*a^7*b))/(\\ & a^5*(a + b)^3*(a - b)) + (2*\tan(c/2 + (d*x)/2)^3*(6*A*a^9 - 120*A*b^9 + 6* \\ & B*a^9 + 364*A*a^2*b^7 + 71*A*a^3*b^6 - 369*A*a^4*b^5 - 45*A*a^5*b^4 + 111* \\ & A*a^6*b^3 + 3*A*a^7*b^2 + 12*B*a^2*b^7 - 148*B*a^3*b^6 - 29*B*a^4*b^5 + 15 \\ & 9*B*a^5*b^4 + 18*B*a^6*b^3 - 30*B*a^7*b^2 - 30*A*a*b^8 - 21*A*a^8*b + 48*B \\ & *a*b^8 - 6*B*a^8*b))/(3*a^5*(a + b)^2*(a - b)^3) + (2*\tan(c/2 + (d*x)/2)^7 \\ & *(6*A*a^9 + 120*A*b^9 - 6*B*a^9 - 364*A*a^2*b^7 + 71*A*a^3*b^6 + 369*A*a^4 \\ & *b^5 - 45*A*a^5*b^4 - 111*A*a^6*b^3 + 3*A*a^7*b^2 + 12*B*a^2*b^7 + 148*B*a \\ & ^3*b^6 - 29*B*a^4*b^5 - 159*B*a^5*b^4 + 18*B*a^6*b^3 + 30*B*a^7*b^2 - 30*A \\ & *a*b^8 + 21*A*a^8*b - 48*B*a*b^8 - 6*B*a^8*b))/(3*a^5*(a + b)^3*(a - b)^2) \\ &)/(d*(\tan(c/2 + (d*x)/2)^4*(6*a*b^2 - 6*a^2*b - 2*a^3 + 10*b^3) - \tan(c... \end{aligned}$$

3.281
$$\int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.281.1 Optimal result 2617
 3.281.2 Mathematica [A] (verified) 2617
 3.281.3 Rubi [A] (verified) 2618
 3.281.4 Maple [A] (verified) 2619
 3.281.5 Fricas [A] (verification not implemented) 2619
 3.281.6 Sympy [B] (verification not implemented) 2620
 3.281.7 Maxima [F(-2)] 2620
 3.281.8 Giac [A] (verification not implemented) 2620
 3.281.9 Mupad [B] (verification not implemented) 2621

3.281.1 Optimal result

Integrand size = 34, antiderivative size = 28

$$\int \frac{\cos^3(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{B \sin(c + dx)}{d} - \frac{B \sin^3(c + dx)}{3d}$$

output `B*sin(d*x+c)/d-1/3*B*sin(d*x+c)^3/d`

3.281.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\cos^3(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = B \left(\frac{\sin(c + dx)}{d} - \frac{\sin^3(c + dx)}{3d} \right)$$

input `Integrate[(Cos[c + d*x]^3*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `B*(Sin[c + d*x]/d - Sin[c + d*x]^3/(3*d))`

3.281.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2011, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \cos^3(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 dx$$

$$\downarrow \text{3113}$$

$$\frac{B \int (1-\sin^2(c+dx)) d(-\sin(c+dx))}{d}$$

$$\downarrow \text{2009}$$

$$\frac{B\left(\frac{1}{3}\sin^3(c+dx) - \sin(c+dx)\right)}{d}$$

input `Int[(Cos[c + d*x]^3*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `-((B*(-Sin[c + d*x] + Sin[c + d*x]^3/3))/d)`

3.281.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m+n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

3.281.4 Maple [A] (verified)

Time = 1.73 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{B(2+\cos^2(dx+c)) \sin(dx+c)}{3d}$	23
default	$\frac{B(2+\cos^2(dx+c)) \sin(dx+c)}{3d}$	23
parallelrisc	$\frac{B(9 \sin(dx+c)+\sin(3dx+3c))}{12d}$	25
risc	$\frac{3B \sin(dx+c)}{4d} + \frac{B \sin(3dx+3c)}{12d}$	29
norman	$\frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{10B \left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3d} + \frac{10B \left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3d} + \frac{2B \left(\tan^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d}}{\left(1+\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^4}$	84

input `int(cos(d*x+c)^3*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `1/3/d*B*(2+cos(d*x+c)^2)*sin(d*x+c)`

3.281.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \frac{(B\cos(dx+c))^2 + 2B}{3d} \sin(dx+c)$$

input `integrate(cos(d*x+c)^3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `1/3*(B*cos(d*x + c)^2 + 2*B)*sin(d*x + c)/d`

3.281. $\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$

3.281.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(22) = 44$.

Time = 0.42 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.00

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \begin{cases} \frac{2B\sin^3(c+dx)}{3d} + \frac{B\sin(c+dx)\cos^2(c+dx)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\cos(c))\cos^3(c)}{a+b\cos(c)} & \text{otherwise} \end{cases}$$

input `integrate(cos(d*x+c)**3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Piecewise((2*B*sin(c + d*x)**3/(3*d) + B*sin(c + d*x)*cos(c + d*x)**2/d, N
e(d, 0)), (x*(B*a + B*b*cos(c))*cos(c)**3/(a + b*cos(c)), True))`

3.281.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm=
"maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de`

3.281.8 Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = -\frac{B\sin(dx+c)^3 - 3B\sin(dx+c)}{3d}$$

input `integrate(cos(d*x+c)^3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `-1/3*(B*sin(d*x + c)^3 - 3*B*sin(d*x + c))/d`

3.281.9 Mupad [B] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{\cos^3(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{B(9 \sin(c + dx) + \sin(3c + 3dx))}{12d}$$

input `int((cos(c + d*x)^3*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `(B*(9*sin(c + d*x) + sin(3*c + 3*d*x)))/(12*d)`

3.282
$$\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.282.1 Optimal result 2622
 3.282.2 Mathematica [A] (verified) 2622
 3.282.3 Rubi [A] (verified) 2623
 3.282.4 Maple [A] (verified) 2624
 3.282.5 Fricas [A] (verification not implemented) 2624
 3.282.6 Sympy [B] (verification not implemented) 2625
 3.282.7 Maxima [F(-2)] 2625
 3.282.8 Giac [A] (verification not implemented) 2626
 3.282.9 Mupad [B] (verification not implemented) 2626

3.282.1 Optimal result

Integrand size = 34, antiderivative size = 27

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{Bx}{2} + \frac{B \cos(c + dx) \sin(c + dx)}{2d}$$

output `1/2*B*x+1/2*B*cos(d*x+c)*sin(d*x+c)/d`

3.282.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{B(2(c + dx) + \sin(2(c + dx)))}{4d}$$

input `Integrate[(Cos[c + d*x]^2*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(B*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d)`

3.282.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2011, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \cos^2(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 dx$$

$$\downarrow \text{3115}$$

$$B \left(\frac{\int 1 dx}{2} + \frac{\sin(c+dx)\cos(c+dx)}{2d} \right)$$

$$\downarrow \text{24}$$

$$B \left(\frac{\sin(c+dx)\cos(c+dx)}{2d} + \frac{x}{2} \right)$$

input `Int[(Cos[c + d*x]^2*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `B*(x/2 + (Cos[c + d*x]*Sin[c + d*x])/(2*d))`

3.282.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

3.282. $\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

3.282.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

method	result	size
risch	$\frac{xB}{2} + \frac{B \sin(2dx+2c)}{4d}$	21
parallelrisch	$\frac{B(2dx+\sin(2dx+2c))}{4d}$	21
derivativedivides	$\frac{B\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$	28
default	$\frac{B\left(\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$	28
norman	$\frac{\frac{B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{xB}{2} - \frac{B\left(\tan^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d} + \frac{3xB\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2} + \frac{3xB\left(\tan^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2} + \frac{xB\left(\tan^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^3}$	98

input `int(cos(d*x+c)^2*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `1/2*x*B+1/4*B/d*sin(2*d*x+2*c)`

3.282.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{Bdx + B \cos(dx + c) \sin(dx + c)}{2d}$$

input `integrate(cos(d*x+c)^2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

3.282. $\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$

output $1/2*(B*d*x + B*\cos(d*x + c)*\sin(d*x + c))/d$

3.282.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(22) = 44$.

Time = 0.39 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.52

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx$$

$$= \begin{cases} \frac{Bx \sin^2(c+dx)}{2} + \frac{Bx \cos^2(c+dx)}{2} + \frac{B \sin(c+dx) \cos(c+dx)}{2d} & \text{for } d \neq 0 \\ \frac{x(Ba + Bb \cos(c)) \cos^2(c)}{a + b \cos(c)} & \text{otherwise} \end{cases}$$

input `integrate(cos(d*x+c)**2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Piecewise((B*x*sin(c + d*x)**2/2 + B*x*cos(c + d*x)**2/2 + B*sin(c + d*x)*cos(c + d*x)/(2*d), Ne(d, 0)), (x*(B*a + B*b*cos(c))*cos(c)**2/(a + b*cos(c)), True))`

3.282.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.282.8 Giac [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.22

$$\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \frac{(dx+c)B + \frac{B\tan(dx+c)}{\tan(dx+c)^2+1}}{2d}$$

input `integrate(cos(d*x+c)^2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/2*((d*x + c)*B + B*tan(d*x + c)/(tan(d*x + c)^2 + 1))/d`

3.282.9 Mupad [B] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.85

$$\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \frac{Bx}{2} + \frac{B\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - B\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{d\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)^2}$$

input `int((cos(c + d*x)^2*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `(B*x)/2 + (B*tan(c/2 + (d*x)/2) - B*tan(c/2 + (d*x)/2)^3)/(d*(tan(c/2 + (d*x)/2)^2 + 1)^2)`

3.283 $\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.283.1 Optimal result 2627
 3.283.2 Mathematica [B] (verified) 2627
 3.283.3 Rubi [A] (verified) 2628
 3.283.4 Maple [A] (verified) 2629
 3.283.5 Fricas [A] (verification not implemented) 2629
 3.283.6 Sympy [B] (verification not implemented) 2630
 3.283.7 Maxima [F(-2)] 2630
 3.283.8 Giac [A] (verification not implemented) 2630
 3.283.9 Mupad [B] (verification not implemented) 2631

3.283.1 Optimal result

Integrand size = 32, antiderivative size = 11

$$\int \frac{\cos(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{B \sin(c + dx)}{d}$$

output `B*sin(d*x+c)/d`

3.283.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 23 vs. 2(11) = 22.

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.09

$$\int \frac{\cos(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = B \left(\frac{\cos(dx) \sin(c)}{d} + \frac{\cos(c) \sin(dx)}{d} \right)$$

input `Integrate[(Cos[c + d*x]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `B*((Cos[d*x]*Sin[c])/d + (Cos[c]*Sin[d*x])/d)`

3.283.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {2011, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

↓ 2011

$$B \int \cos(c+dx) dx$$

↓ 3042

$$B \int \sin\left(c+dx+\frac{\pi}{2}\right) dx$$

↓ 3117

$$\frac{B \sin(c+dx)}{d}$$

input `Int[(Cos[c + d*x]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(B*Sin[c + d*x])/d`

3.283.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]`

3.283.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{B \sin(dx+c)}{d}$	12
default	$\frac{B \sin(dx+c)}{d}$	12
risch	$\frac{B \sin(dx+c)}{d}$	12
parallelrisch	$\frac{B \sin(dx+c)}{d}$	12
norman	$\frac{2B \tan\left(\frac{dx+c}{2}\right) + 2B \left(\tan^3\left(\frac{dx+c}{2}\right)\right)}{d \left(1 + \tan^2\left(\frac{dx+c}{2}\right)\right)^2}$	50

input `int(cos(d*x+c)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `B*sin(d*x+c)/d`

3.283.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \frac{B \sin(dx+c)}{d}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `B*sin(d*x + c)/d`

3.283.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. $2(8) = 16$.

Time = 0.34 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.82

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \begin{cases} \frac{B\sin(c+dx)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\cos(c))\cos(c)}{a+b\cos(c)} & \text{otherwise} \end{cases}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Piecewise((B*sin(c + d*x)/d, Ne(d, 0)), (x*(B*a + B*b*cos(c))*cos(c)/(a + b*cos(c)), True))`

3.283.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.283.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \frac{B\sin(dx+c)}{d}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `B*sin(d*x + c)/d`

3.283.9 Mupad [B] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \frac{B \sin(c + dx)}{d}$$

input `int((cos(c + d*x)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `(B*sin(c + d*x))/d`

3.284 $\int \frac{aB+bB \cos(c+dx)}{a+b \cos(c+dx)} dx$

3.284.1 Optimal result 2632
 3.284.2 Mathematica [A] (verified) 2632
 3.284.3 Rubi [A] (verified) 2633
 3.284.4 Maple [A] (verified) 2634
 3.284.5 Fricas [A] (verification not implemented) 2634
 3.284.6 Sympy [A] (verification not implemented) 2634
 3.284.7 Maxima [F(-2)] 2635
 3.284.8 Giac [C] (verification not implemented) 2635
 3.284.9 Mupad [B] (verification not implemented) 2635

3.284.1 Optimal result

Integrand size = 26, antiderivative size = 3

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = Bx$$

output

```
B*x
```

3.284.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = Bx$$

input

```
Integrate[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x]),x]
```

output

```
B*x
```

3.284.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2011, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx$$

↓ 2011

$$B \int 1 dx$$

↓ 24

$$Bx$$

input `Int[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x]),x]`

output `B*x`

3.284.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

3.284.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
default	xB	4
risch	xB	4
derivativedivides	$\frac{B(dx+c)}{d}$	11
norman	$\frac{xB + xB \left(\tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{1 + \tan^2 \left(\frac{dx}{2} + \frac{c}{2} \right)}$	35

input `int((B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`output `x*B`**3.284.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = Bx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fracas")`output `B*x`**3.284.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = Bx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`output `B*x`

3.284. $\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx$

3.284.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.284.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 3.33

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = \frac{(dx + c)B}{d}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `(d*x + c)*B/d`

3.284.9 Mupad [B] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx = Bx$$

input `int((B*a + B*b*cos(c + d*x))/(a + b*cos(c + d*x)),x)`

output `B*x`

3.284. $\int \frac{aB + bB \cos(c + dx)}{a + b \cos(c + dx)} dx$

3.285 $\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx$

3.285.1 Optimal result 2636
 3.285.2 Mathematica [A] (verified) 2636
 3.285.3 Rubi [A] (verified) 2637
 3.285.4 Maple [A] (verified) 2638
 3.285.5 Fricas [B] (verification not implemented) 2638
 3.285.6 Sympy [B] (verification not implemented) 2639
 3.285.7 Maxima [F(-2)] 2639
 3.285.8 Giac [B] (verification not implemented) 2639
 3.285.9 Mupad [B] (verification not implemented) 2640

3.285.1 Optimal result

Integrand size = 32, antiderivative size = 12

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \operatorname{arctanh}(\sin(c + dx))}{d}$$

output `B*arctanh(sin(d*x+c))/d`

3.285.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \operatorname{arctanh}(\sin(c + dx))}{d}$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x]),x]`

output `(B*ArcTanh[Sin[c + d*x]])/d`

3.285.3 Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {2011, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

↓ 2011

$$B \int \sec(c+dx) dx$$

↓ 3042

$$B \int \csc\left(c+dx+\frac{\pi}{2}\right) dx$$

↓ 4257

$$\frac{B \operatorname{arctanh}(\sin(c+dx))}{d}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x]),x]`

output `(B*ArcTanh[Sin[c + d*x]])/d`

3.285.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :> Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.285.4 Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

method	result	size
derivativedivides	$\frac{B \ln(\sec(dx+c)+\tan(dx+c))}{d}$	20
default	$\frac{B \ln(\sec(dx+c)+\tan(dx+c))}{d}$	20
parallelrisc	$\frac{B \left(\ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right) - \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right) \right)}{d}$	33
norman	$\frac{B \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) + 1 \right)}{d} - \frac{B \ln \left(\tan \left(\frac{dx}{2} + \frac{c}{2} \right) - 1 \right)}{d}$	37
risc	$\frac{B \ln(e^{i(dx+c)}+i)}{d} - \frac{B \ln(e^{i(dx+c)}-i)}{d}$	39

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)
```

```
output 1/d*B*ln(sec(d*x+c)+tan(d*x+c))
```

3.285.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(12) = 24.

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.58

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{B \log(\sin(dx + c) + 1) - B \log(-\sin(dx + c) + 1)}{2d}$$

```
input integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x, algorithm="fracas")
```

```
output 1/2*(B*log(sin(d*x + c) + 1) - B*log(-sin(d*x + c) + 1))/d
```

3.285.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(10) = 20$.

Time = 1.84 (sec) , antiderivative size = 39, normalized size of antiderivative = 3.25

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \begin{cases} \frac{B \log(\tan(c + dx) + \sec(c + dx))}{d} & \text{for } d \neq 0 \\ \frac{x(Ba + Bb \cos(c)) \sec(c)}{a + b \cos(c)} & \text{otherwise} \end{cases}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x)`

output `Piecewise((B*log(tan(c + d*x) + sec(c + d*x))/d, Ne(d, 0)), (x*(B*a + B*b*cos(c))*sec(c)/(a + b*cos(c)), True))`

3.285.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.285.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. $2(12) = 24$.

Time = 0.28 (sec) , antiderivative size = 47, normalized size of antiderivative = 3.92

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \log \left(\left| \frac{1}{\sin(dx+c)} + \sin(dx+c) + 2 \right| \right) - B \log \left(\left| \frac{1}{\sin(dx+c)} + \sin(dx+c) - 2 \right| \right)}{4d}$$

3.285. $\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{a+b \cos(c+dx)} dx$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/4*(B*log(abs(1/sin(d*x + c) + sin(d*x + c) + 2)) - B*log(abs(1/sin(d*x + c) + sin(d*x + c) - 2)))/d`

3.285.9 Mupad [B] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{a + b \cos(c + dx)} dx = \frac{2B \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d}$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))),x)`

output `(2*B*atanh(tan(c/2 + (d*x)/2)))/d`

$$3.286 \quad \int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx$$

3.286.1 Optimal result	2641
3.286.2 Mathematica [A] (verified)	2641
3.286.3 Rubi [A] (verified)	2642
3.286.4 Maple [A] (verified)	2643
3.286.5 Fracas [A] (verification not implemented)	2643
3.286.6 Sympy [B] (verification not implemented)	2644
3.286.7 Maxima [F(-2)]	2644
3.286.8 Giac [A] (verification not implemented)	2645
3.286.9 Mupad [B] (verification not implemented)	2645

3.286.1 Optimal result

Integrand size = 34, antiderivative size = 11

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \tan(c + dx)}{d}$$

output `B*tan(d*x+c)/d`

3.286.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \tan(c + dx)}{d}$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x]),x]`

output `(B*Tan[c + d*x])/d`

3.286.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2011, 3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

↓ 2011

$$B \int \sec^2(c+dx) dx$$

↓ 3042

$$B \int \csc\left(c+dx+\frac{\pi}{2}\right)^2 dx$$

↓ 4254

$$\frac{B \int 1d(-\tan(c+dx))}{d}$$

↓ 24

$$\frac{B \tan(c+dx)}{d}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x]),x]`

output `(B*Tan[c + d*x])/d`

3.286.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4254 Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

3.286.4 Maple [A] (verified)

Time = 1.83 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{B \tan(dx+c)}{d}$	12
default	$\frac{B \tan(dx+c)}{d}$	12
risch	$\frac{2iB}{d(e^{2i(dx+c)}+1)}$	21
parallelrisc	$-\frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$	31
norman	$\frac{-\frac{2B \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} - \frac{2B\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{d}}{\left(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(\tan^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$	65

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b),x,method=_RETURNVER
BOSE)
```

```
output B*tan(d*x+c)/d
```

3.286.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \sin(dx + c)}{d \cos(dx + c)}$$

```
input integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c)),x, algorithm=
"fracas")
```

output `B*sin(d*x + c)/(d*cos(d*x + c))`

3.286.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs. 2(8) = 16.

Time = 1.44 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.91

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \begin{cases} \frac{B \tan(c + dx)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba + Bb \cos(c)) \sec^2(c)}{a + b \cos(c)} & \text{otherwise} \end{cases}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c)),x)`

output `Piecewise((B*tan(c + d*x)/d, Ne(d, 0)), (x*(B*a + B*b*cos(c))*sec(c)**2/(a + b*cos(c)), True))`

3.286.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.286.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \tan(dx + c)}{d}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `B*tan(d*x + c)/d`

3.286.9 Mupad [B] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.73

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{a + b \cos(c + dx)} dx = -\frac{2B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1\right)}$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))),x)`

output `-(2*B*tan(c/2 + (d*x)/2))/(d*(tan(c/2 + (d*x)/2)^2 - 1))`

3.287 $\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$

3.287.1 Optimal result 2646
 3.287.2 Mathematica [A] (verified) 2646
 3.287.3 Rubi [A] (verified) 2647
 3.287.4 Maple [A] (verified) 2648
 3.287.5 Fricas [A] (verification not implemented) 2649
 3.287.6 Sympy [F] 2649
 3.287.7 Maxima [F(-2)] 2649
 3.287.8 Giac [A] (verification not implemented) 2650
 3.287.9 Mupad [B] (verification not implemented) 2650

3.287.1 Optimal result

Integrand size = 34, antiderivative size = 36

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{B \sec(c + dx) \tan(c + dx)}{2d}$$

output `1/2*B*arctanh(sin(d*x+c))/d+1/2*B*sec(d*x+c)*tan(d*x+c)/d`

3.287.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = B \left(\frac{\operatorname{arctanh}(\sin(c + dx))}{2d} + \frac{\sec(c + dx) \tan(c + dx)}{2d} \right)$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x]),x]`

output `B*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))`

3.287.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {2011, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \sec^3(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \csc\left(c+dx+\frac{\pi}{2}\right)^3 dx$$

$$\downarrow \text{4255}$$

$$B\left(\frac{1}{2} \int \sec(c+dx) dx + \frac{\tan(c+dx)\sec(c+dx)}{2d}\right)$$

$$\downarrow \text{3042}$$

$$B\left(\frac{1}{2} \int \csc\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\tan(c+dx)\sec(c+dx)}{2d}\right)$$

$$\downarrow \text{4257}$$

$$B\left(\frac{\operatorname{arctanh}(\sin(c+dx))}{2d} + \frac{\tan(c+dx)\sec(c+dx)}{2d}\right)$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x]),x]`

output `B*(ArcTanh[Sin[c + d*x]]/(2*d) + (Sec[c + d*x]*Tan[c + d*x])/(2*d))`

3.287.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4255 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
  && IntegerQ[2*n]
```

```
rule 4257 Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
  /; FreeQ[{c, d}, x]
```

3.287.4 Maple [A] (verified)

Time = 1.58 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$	37
default	$\frac{B \left(\frac{\sec(dx+c) \tan(dx+c)}{2} + \frac{\ln(\sec(dx+c)+\tan(dx+c))}{2} \right)}{d}$	37
parallelrisc	$-\frac{\left((1+\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right) + (-1-\cos(2dx+2c)) \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right) - 2\sin(dx+c) \right) B}{2d(1+\cos(2dx+2c))}$	79
risc	$-\frac{iB(e^{3i(dx+c)}-e^{i(dx+c)})}{d(e^{2i(dx+c)}+1)^2} + \frac{B \ln(e^{i(dx+c)}+i)}{2d} - \frac{B \ln(e^{i(dx+c)}-i)}{2d}$	81
norman	$\frac{B \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d} + \frac{B(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right))}{d} + \frac{2B(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right))}{d} - \frac{B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}{2d} + \frac{B \ln\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}{2d}$	117

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b), x, method=_RETURNVER
BOSE)
```

3.287.
$$\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$$

output $1/d*B*(1/2*\sec(d*x+c)*\tan(d*x+c)+1/2*\ln(\sec(d*x+c)+\tan(d*x+c)))$

3.287.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.78

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{B \cos(dx + c)^2 \log(\sin(dx + c) + 1) - B \cos(dx + c)^2 \log(-\sin(dx + c) + 1) + 2 B \sin(dx + c)}{4 d \cos(dx + c)^2}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output $1/4*(B*\cos(d*x + c)^2*\log(\sin(d*x + c) + 1) - B*\cos(d*x + c)^2*\log(-\sin(d*x + c) + 1) + 2*B*\sin(d*x + c))/(d*\cos(d*x + c)^2)$

3.287.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = B \int \sec^3(c + dx) dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c)),x)`

output `B*Integral(sec(c + d*x)**3, x)`

3.287.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.287.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.44

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{B \log(|\sin(dx + c) + 1|) - B \log(|\sin(dx + c) - 1|) - \frac{2B \sin(dx+c)}{\sin(dx+c)^2-1}}{4d}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/4*(B*log(abs(sin(d*x + c) + 1)) - B*log(abs(sin(d*x + c) - 1)) - 2*B*sin(d*x + c)/(sin(d*x + c)^2 - 1))/d`

3.287.9 Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.03

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)} + \frac{B \operatorname{atanh}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{d}$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))),x)`

output `(B*tan(c/2 + (d*x)/2) + B*tan(c/2 + (d*x)/2)^3)/(d*(tan(c/2 + (d*x)/2)^4 - 2*tan(c/2 + (d*x)/2)^2 + 1)) + (B*atanh(tan(c/2 + (d*x)/2)))/d`

3.287. $\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{a+b \cos(c+dx)} dx$

$$3.288 \quad \int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx$$

3.288.1 Optimal result	2651
3.288.2 Mathematica [A] (verified)	2651
3.288.3 Rubi [A] (verified)	2652
3.288.4 Maple [A] (verified)	2653
3.288.5 Fracas [A] (verification not implemented)	2654
3.288.6 Sympy [A] (verification not implemented)	2654
3.288.7 Maxima [F(-2)]	2654
3.288.8 Giac [A] (verification not implemented)	2655
3.288.9 Mupad [B] (verification not implemented)	2655

3.288.1 Optimal result

Integrand size = 34, antiderivative size = 28

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \tan(c + dx)}{d} + \frac{B \tan^3(c + dx)}{3d}$$

output `B*tan(d*x+c)/d+1/3*B*tan(d*x+c)^3/d`

3.288.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \frac{B(\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d}$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + b*Cos[c + d*x]),x]`

output `(B*(Tan[c + d*x] + Tan[c + d*x]^3/3))/d`

3.288.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2011, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^4(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \sec^4(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \csc\left(c+dx+\frac{\pi}{2}\right)^4 dx$$

$$\downarrow \text{4254}$$

$$\frac{B \int (\tan^2(c+dx)+1) d(-\tan(c+dx))}{d}$$

$$\downarrow \text{2009}$$

$$\frac{B\left(-\frac{1}{3}\tan^3(c+dx)-\tan(c+dx)\right)}{d}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^4)/(a + b*Cos[c + d*x]),x]`

output `-((B*(-Tan[c + d*x] - Tan[c + d*x]^3/3))/d)`

3.288.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x, a + b*x])`

3.288. $\int \frac{(aB+bB\cos(c+dx))\sec^4(c+dx)}{a+b\cos(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

3.288.4 Maple [A] (verified)

Time = 1.77 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\frac{B\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c)}{d}$	25
default	$\frac{B\left(-\frac{2}{3}-\frac{\sec^2(dx+c)}{3}\right)\tan(dx+c)}{d}$	25
risch	$\frac{4iB(3e^{2i(dx+c)}+1)}{3d(e^{2i(dx+c)}+1)^3}$	34
parallelrisch	$\frac{2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\tan^4\left(\frac{dx}{2}+\frac{c}{2}\right)-\frac{2\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3}+1\right)B}{d\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}$	68
norman	$\frac{\frac{2B\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}-\frac{2B\left(\tan^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3d}-\frac{2B\left(\tan^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3d}-\frac{2B\left(\tan^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{d}}{\left(1+\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(\tan^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}$	99

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^4/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `-1/d*B*(-2/3-1/3*sec(d*x+c)^2)*tan(d*x+c)`

3.288.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \frac{(2B \cos(dx + c)^2 + B) \sin(dx + c)}{3d \cos(dx + c)^3}$$

```
input integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^4/(a+b*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/3*(2*B*cos(d*x + c)^2 + B)*sin(d*x + c)/(d*cos(d*x + c)^3)
```

3.288.6 Sympy [A] (verification not implemented)

Time = 8.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.50

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \begin{cases} \frac{B \left(\frac{\tan^3(c+dx)}{3} + \tan(c+dx) \right)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba + Bb \cos(c)) \sec^4(c)}{a + b \cos(c)} & \text{otherwise} \end{cases}$$

```
input integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**4/(a+b*cos(d*x+c)),x)
```

```
output Piecewise((B*(tan(c + d*x)**3/3 + tan(c + d*x))/d, Ne(d, 0)), (x*(B*a + B*b*cos(c))*sec(c)**4/(a + b*cos(c)), True))
```

3.288.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \text{Exception raised: ValueError}$$

```
input integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^4/(a+b*cos(d*x+c)),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de
```

3.288.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \frac{B \tan(dx + c)^3 + 3B \tan(dx + c)}{3d}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^4/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `1/3*(B*tan(d*x + c)^3 + 3*B*tan(d*x + c))/d`

3.288.9 Mupad [B] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.39

$$\int \frac{(aB + bB \cos(c + dx)) \sec^4(c + dx)}{a + b \cos(c + dx)} dx = \frac{2B \sin(c + dx) \cos(c + dx)^2 + B \sin(c + dx)}{3d \cos(c + dx)^3}$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^4*(a + b*cos(c + d*x))),x)`

output `(B*sin(c + d*x) + 2*B*cos(c + d*x)^2*sin(c + d*x))/(3*d*cos(c + d*x)^3)`

$$3.289 \quad \int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.289.1 Optimal result	2656
3.289.2 Mathematica [A] (verified)	2656
3.289.3 Rubi [A] (verified)	2657
3.289.4 Maple [A] (verified)	2660
3.289.5 Fricas [A] (verification not implemented)	2661
3.289.6 Sympy [F(-1)]	2661
3.289.7 Maxima [F(-2)]	2662
3.289.8 Giac [A] (verification not implemented)	2662
3.289.9 Mupad [B] (verification not implemented)	2663

3.289.1 Optimal result

Integrand size = 34, antiderivative size = 114

$$\int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{(2a^2+b^2)Bx}{2b^3} - \frac{2a^3B \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{\sqrt{a-b}b^3\sqrt{a+bd}} - \frac{aB \sin(c+dx)}{b^2d} + \frac{B \cos(c+dx) \sin(c+dx)}{2bd}$$

output `1/2*(2*a^2+b^2)*B*x/b^3-a*B*sin(d*x+c)/b^2/d+1/2*B*cos(d*x+c)*sin(d*x+c)/b/d-2*a^3*B*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b^3/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.289.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.86

$$\int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{B \left(2(2a^2+b^2)(c+dx) + \frac{8a^3 \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} - 4ab \sin(c+dx) + b^2 \sin(2(c+dx)) \right)}{4b^3d}$$

input `Integrate[(Cos[c + d*x]^3*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2, x]`

output `(B*(2*(2*a^2 + b^2)*(c + d*x) + (8*a^3*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] - 4*a*b*Sin[c + d*x] + b^2*Sin[2*(c + d*x)])/(4*b^3*d)`

3.289.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2011, 3042, 3272, 3042, 3502, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^3(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\cos^3(c + dx)}{a + b \cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\sin(c + dx + \frac{\pi}{2})^3}{a + b \sin(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3272} \\
 & B \left(\frac{\int \frac{-2a \cos^2(c + dx) + b \cos(c + dx) + a}{a + b \cos(c + dx)} dx}{2b} + \frac{\sin(c + dx) \cos(c + dx)}{2bd} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{\int \frac{-2a \sin(c + dx + \frac{\pi}{2})^2 + b \sin(c + dx + \frac{\pi}{2}) + a}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{2b} + \frac{\sin(c + dx) \cos(c + dx)}{2bd} \right) \\
 & \quad \downarrow \text{3502}
 \end{aligned}$$

$$\begin{aligned}
& B \left(\frac{\int \frac{ab + (2a^2 + b^2) \cos(c+dx)}{a+b \cos(c+dx)} dx - \frac{2a \sin(c+dx)}{bd}}{2b} + \frac{\sin(c+dx) \cos(c+dx)}{2bd} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{\int \frac{ab + (2a^2 + b^2) \sin(c+dx + \frac{\pi}{2})}{a+b \sin(c+dx + \frac{\pi}{2})} dx - \frac{2a \sin(c+dx)}{bd}}{2b} + \frac{\sin(c+dx) \cos(c+dx)}{2bd} \right) \\
& \quad \downarrow \text{3214} \\
& B \left(\frac{\frac{x(2a^2 + b^2)}{b} - \frac{2a^3 \int \frac{1}{a+b \cos(c+dx)} dx}{b}}{2b} - \frac{2a \sin(c+dx)}{bd} + \frac{\sin(c+dx) \cos(c+dx)}{2bd} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{\frac{x(2a^2 + b^2)}{b} - \frac{2a^3 \int \frac{1}{a+b \sin(c+dx + \frac{\pi}{2})} dx}{b}}{2b} - \frac{2a \sin(c+dx)}{bd} + \frac{\sin(c+dx) \cos(c+dx)}{2bd} \right) \\
& \quad \downarrow \text{3138} \\
& B \left(\frac{\frac{x(2a^2 + b^2)}{b} - \frac{4a^3 \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{b}}{2b} - \frac{2a \sin(c+dx)}{bd} + \frac{\sin(c+dx) \cos(c+dx)}{2bd} \right) \\
& \quad \downarrow \text{218} \\
& B \left(\frac{\frac{x(2a^2 + b^2)}{b} - \frac{4a^3 \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{bd\sqrt{a-b}\sqrt{a+b}}}{2b} - \frac{2a \sin(c+dx)}{bd} + \frac{\sin(c+dx) \cos(c+dx)}{2bd} \right)
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `B*((Cos[c + d*x]*Sin[c + d*x])/(2*b*d) + (((((2*a^2 + b^2)*x)/b - (4*a^3*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d))/b - (2*a*Sin[c + d*x])/(b*d))/(2*b))`

3.289. $\int \frac{\cos^3(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.289.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 3272 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 2)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Sin[e + f*x])^(m - 3)*(c + d*Sin[e + f*x])^n*Simp[a^3*d*(m + n) + b^2*(b*c*(m - 2) + a*d*(n + 1)) - b*(a*b*c - b^2*d*(m + n - 1) - 3*a^2*d*(m + n))*Sin[e + f*x] - b^2*(b*c*(m - 1) - a*d*(3*m + 2*n - 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 2] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.289.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.22

method	result
derivativedivides	$2B \left(-\frac{a^3 \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^3 \sqrt{(a-b)(a+b)}} + \frac{(-ab - \frac{1}{2}b^2)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-ab + \frac{1}{2}b^2)\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (2a^2 + b^2)\arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^2} \right) + \frac{d}{b^3}$
default	$2B \left(-\frac{a^3 \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^3 \sqrt{(a-b)(a+b)}} + \frac{(-ab - \frac{1}{2}b^2)\left(\tan^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-ab + \frac{1}{2}b^2)\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + (2a^2 + b^2)\arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{(1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right))^2} \right) + \frac{d}{b^3}$
risch	$\frac{Bxa^2}{b^3} + \frac{Bx}{2b} + \frac{iBa e^{i(dx+c)}}{2b^2d} - \frac{iBa e^{-i(dx+c)}}{2b^2d} - \frac{a^3B \ln\left(\frac{e^{i(dx+c)} - ia^2 - ib^2 - a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}db^3} + \frac{a^3B \ln\left(e^{i(dx+c)} + \dots\right)}{\sqrt{-a^2+b^2}}$

```
input int(cos(d*x+c)^3*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

```
output 2/d*B*(-a^3/b^3/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)
*(a+b))^(1/2))+1/b^3*(((a*b-1/2*b^2)*tan(1/2*d*x+1/2*c)^3+(-a*b+1/2*b^2)*
tan(1/2*d*x+1/2*c))/(1+tan(1/2*d*x+1/2*c)^2)^2+1/2*(2*a^2+b^2)*arctan(tan(
1/2*d*x+1/2*c))))
```

3.289.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 350, normalized size of antiderivative = 3.07

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \left[\frac{\sqrt{-a^2+b^2}Ba^3 \log\left(\frac{2ab\cos(dx+c)+(2a^2-b^2)\cos(dx+c)^2-2\sqrt{-a^2+b^2}(a\cos(dx+c)+b)\sin(dx+c)-a^2+2b^2}{b^2\cos(dx+c)^2+2ab\cos(dx+c)+a^2}\right) - (2Ba^4 - Bb^4)}{2(a^2b^3 - b^5)d} \right. \\ \left. - \frac{2\sqrt{a^2-b^2}Ba^3 \arctan\left(-\frac{a\cos(dx+c)+b}{\sqrt{a^2-b^2}\sin(dx+c)}\right) - (2Ba^4 - Ba^2b^2 - Bb^4)dx + (2Ba^3b - 2Bab^3 - (Ba^2b^2 - Bb^4)\cos(dx+c))\sin(dx+c)}{2(a^2b^3 - b^5)d} \right]$$

```
input integrate(cos(d*x+c)^3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm
m="fricas")
```

```
output [-1/2*(sqrt(-a^2 + b^2)*B*a^3*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(
d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 +
2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - (2*B*a^4 - B*a^2
*b^2 - B*b^4)*d*x + (2*B*a^3*b - 2*B*a*b^3 - (B*a^2*b^2 - B*b^4)*cos(d*x +
c))*sin(d*x + c))/((a^2*b^3 - b^5)*d), -1/2*(2*sqrt(a^2 - b^2)*B*a^3*arct
an(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (2*B*a^4 - B*a^
2*b^2 - B*b^4)*d*x + (2*B*a^3*b - 2*B*a*b^3 - (B*a^2*b^2 - B*b^4)*cos(d*x
+ c))*sin(d*x + c))/((a^2*b^3 - b^5)*d)]
```

3.289.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)
```

```
output Timed out
```

3.289.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^3(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.289.8 Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.62

$$\int \frac{\cos^3(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx =$$

$$\frac{4 \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right) B a^3}{\sqrt{a^2 - b^2} b^3} - \frac{(2Ba^2 + Bb^2)(dx+c)}{b^3} + \frac{2 \left(2Ba \tan(\frac{1}{2} dx + \frac{1}{2} c) \right)^3 + Bb^3}{2d}$$

input `integrate(cos(d*x+c)^3*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output
$$-1/2*(4*(\pi*\text{floor}(1/2*(d*x + c)/\pi + 1/2)*\operatorname{sgn}(2*a - 2*b) + \arctan((a*\tan(1/2*d*x + 1/2*c) - b*\tan(1/2*d*x + 1/2*c))/\sqrt{a^2 - b^2}))*B*a^3/(\sqrt{a^2 - b^2})*b^3 - (2*B*a^2 + B*b^2)*(d*x + c)/b^3 + 2*(2*B*a*\tan(1/2*d*x + 1/2*c))^3 + B*b*\tan(1/2*d*x + 1/2*c)^3 + 2*B*a*\tan(1/2*d*x + 1/2*c) - B*b*\tan(1/2*d*x + 1/2*c)/((\tan(1/2*d*x + 1/2*c)^2 + 1)^2*b^2))/d$$

3.289.9 Mupad [B] (verification not implemented)

Time = 1.24 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.52

$$\int \frac{\cos^3(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \frac{B \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2}+\frac{dx}{2}\right)}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{bd} + \frac{B \sin(2c+2dx)}{4bd}$$

$$+ \frac{2Ba^2 \operatorname{atan}\left(\frac{\sin\left(\frac{c}{2}+\frac{dx}{2}\right)}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right)}\right)}{b^3d} - \frac{Ba \sin(c+dx)}{b^2d}$$

$$- \frac{Ba^3 \operatorname{atan}\left(\frac{\left(a \sin\left(\frac{c}{2}+\frac{dx}{2}\right) - b \sin\left(\frac{c}{2}+\frac{dx}{2}\right)\right) i}{\cos\left(\frac{c}{2}+\frac{dx}{2}\right) \sqrt{b^2-a^2}}\right)}{b^3d\sqrt{b^2-a^2}} 2i$$

input `int((cos(c + d*x))^3*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`output `(B*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(b*d) + (B*sin(2*c + 2*d*x))/(4*b*d) + (2*B*a^2*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(b^3*d) - (B*a*sin(c + d*x))/(b^2*d) - (B*a^3*atan(((a*sin(c/2 + (d*x)/2) - b*sin(c/2 + (d*x)/2))*i)/(cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2)))*2i)/(b^3*d*(b^2 - a^2)^(1/2))`

3.290
$$\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.290.1 Optimal result 2664
 3.290.2 Mathematica [A] (verified) 2664
 3.290.3 Rubi [A] (verified) 2665
 3.290.4 Maple [A] (verified) 2668
 3.290.5 Fricas [A] (verification not implemented) 2668
 3.290.6 Sympy [F(-1)] 2669
 3.290.7 Maxima [F(-2)] 2669
 3.290.8 Giac [A] (verification not implemented) 2670
 3.290.9 Mupad [B] (verification not implemented) 2670

3.290.1 Optimal result

Integrand size = 34, antiderivative size = 79

$$\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = -\frac{aBx}{b^2} + \frac{2a^2B \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{\sqrt{a-b}b^2\sqrt{a+bd}} + \frac{B \sin(c+dx)}{bd}$$

output `-a*B*x/b^2+B*sin(d*x+c)/b/d+2*a^2*B*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b^2/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.290.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.92

$$\int \frac{\cos^2(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{B \left(-a(c+dx) - \frac{2a^2 \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + b \sin(c+dx) \right)}{b^2d}$$

input `Integrate[(Cos[c + d*x]^2*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2, x]`

output $(B*(-(a*(c + d*x)) - (2*a^2*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + b*Sin[c + d*x]))/(b^2*d)$

3.290.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2011, 3042, 3225, 25, 27, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{\cos^2(c + dx)}{a + b \cos(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{\sin(c + dx + \frac{\pi}{2})^2}{a + b \sin(c + dx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{3225} \\ & B \left(\frac{\int -\frac{a \cos(c+dx)}{a+b \cos(c+dx)} dx}{b} + \frac{\sin(c + dx)}{bd} \right) \\ & \quad \downarrow \text{25} \\ & B \left(\frac{\sin(c + dx)}{bd} - \frac{\int \frac{a \cos(c+dx)}{a+b \cos(c+dx)} dx}{b} \right) \\ & \quad \downarrow \text{27} \\ & B \left(\frac{\sin(c + dx)}{bd} - \frac{a \int \frac{\cos(c+dx)}{a+b \cos(c+dx)} dx}{b} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{\sin(c + dx)}{bd} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} \right) \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3214} \\
 B \left(\frac{\sin(c+dx)}{bd} - \frac{a \left(\frac{x}{b} - \frac{a \int \frac{1}{a+b \cos(c+dx)} dx}{b} \right)}{b} \right) \\
 \\
 \downarrow \text{3042} \\
 B \left(\frac{\sin(c+dx)}{bd} - \frac{a \left(\frac{x}{b} - \frac{a \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} \right)}{b} \right) \\
 \\
 \downarrow \text{3138} \\
 B \left(\frac{\sin(c+dx)}{bd} - \frac{a \left(\frac{x}{b} - \frac{2a \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{bd} \right)}{b} \right) \\
 \\
 \downarrow \text{218} \\
 B \left(\frac{\sin(c+dx)}{bd} - \frac{a \left(\frac{x}{b} - \frac{2a \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{bd\sqrt{a-b}\sqrt{a+b}} \right)}{b} \right)
 \end{array}$$

input `Int[(Cos[c + d*x]^2*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `B*(-((a*(x/b - (2*a*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d)))/b) + Sin[c + d*x]/(b*d))`

3.290.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3214 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`
- rule 3225 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b^2)*(Cos[e + f*x]/(d*f)), x] + Simp[1/d Int[Simp[a^2*d - b*(b*c - 2*a*d)*Sin[e + f*x], x]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

3.290.4 Maple [A] (verified)

Time = 0.98 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.24

method	result
derivativedivides	$2B \left(\frac{a^2 \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^2 \sqrt{(a-b)(a+b)}} - \frac{b \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + a \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right) \frac{d}{b^2}$
default	$2B \left(\frac{a^2 \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b^2 \sqrt{(a-b)(a+b)}} - \frac{b \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1 + \tan^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + a \arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right) \frac{d}{b^2}$
risch	$-\frac{aBx}{b^2} - \frac{iB e^{i(dx+c)}}{2db} + \frac{iB e^{-i(dx+c)}}{2db} - \frac{a^2 B \ln\left(e^{i(dx+c)} + \frac{ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} db^2} + \frac{a^2 B \ln\left(e^{i(dx+c)} + \frac{-ia^2 + ib^2}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} db^2}$

input `int(cos(d*x+c)^2*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNV ERBOSE)`

output `2/d*B*(a^2/b^2/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))-1/b^2*(-b*tan(1/2*d*x+1/2*c)/(1+tan(1/2*d*x+1/2*c)^2)+a*arctan(tan(1/2*d*x+1/2*c))))`

3.290.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 281, normalized size of antiderivative = 3.56

$$\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \left[-\frac{\sqrt{-a^2+b^2}Ba^2 \log\left(\frac{2ab\cos(dx+c)+(2a^2-b^2)\cos(dx+c)^2+2\sqrt{-a^2+b^2}(a\cos(dx+c)+b)\sin(dx+c)-a^2+2b^2}{b^2\cos(dx+c)^2+2ab\cos(dx+c)+a^2}\right) + 2(Ba^3 - Bb^3)}{2(a^2b^2 - b^4)d} \right]$$

input `integrate(cos(d*x+c)^2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output `[-1/2*(sqrt(-a^2 + b^2)*B*a^2*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + 2*(B*a^3 - B*a*b^2)*d*x - 2*(B*a^2*b - B*b^3)*sin(d*x + c))/((a^2*b^2 - b^4)*d), (sqrt(a^2 - b^2)*B*a^2*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (B*a^3 - B*a*b^2)*d*x + (B*a^2*b - B*b^3)*sin(d*x + c))/((a^2*b^2 - b^4)*d)]`

3.290.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.290.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)^2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.290.8 Giac [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.62

$$\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{2 \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2-b^2}} \right) \right) B a^2}{\sqrt{a^2-b^2} b^2} - \frac{(dx+c)Ba}{b^2} + \frac{2B \tan(\frac{1}{2} dx + \frac{1}{2} c)}{(\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 1)b}$$

```
input integrate(cos(d*x+c)^2*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")
```

```
output (2*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))*B*a^2/(sqrt(a^2 - b^2)*b^2) - (d*x + c)*B*a/b^2 + 2*B*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 + 1)*b))/d
```

3.290.9 Mupad [B] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 193, normalized size of antiderivative = 2.44

$$\int \frac{\cos^2(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{B \sin(c+dx)}{bd} - \frac{2Ba \operatorname{atan} \left(\frac{\sin(\frac{c}{2} + \frac{dx}{2})}{\cos(\frac{c}{2} + \frac{dx}{2})} \right)}{b^2 d}$$

$$- \frac{Ba^2 \operatorname{atan} \left(\frac{1i \sin(\frac{c}{2} + \frac{dx}{2}) a^2 b - 2i \sin(\frac{c}{2} + \frac{dx}{2}) a b^2 + 1i \sin(\frac{c}{2} + \frac{dx}{2}) b^3}{\cos(\frac{c}{2} + \frac{dx}{2}) (b^2 - a^2)^{3/2} + a^2 \cos(\frac{c}{2} + \frac{dx}{2}) \sqrt{b^2 - a^2} - a b \cos(\frac{c}{2} + \frac{dx}{2}) \sqrt{b^2 - a^2}} \right) 2i}{b^2 d \sqrt{b^2 - a^2}}$$

```
input int((cos(c + d*x))^2*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)
```

```
output (B*sin(c + d*x))/(b*d) - (2*B*a*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(b^2*d) - (B*a^2*atan((b^3*sin(c/2 + (d*x)/2)*1i - a*b^2*sin(c/2 + (d*x)/2)*2i + a^2*b*sin(c/2 + (d*x)/2)*1i)/(cos(c/2 + (d*x)/2)*(b^2 - a^2)^(3/2) + a^2*cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - a*b*cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2)))*2i)/(b^2*d*(b^2 - a^2)^(1/2))
```

3.291
$$\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.291.1 Optimal result 2671
 3.291.2 Mathematica [A] (verified) 2671
 3.291.3 Rubi [A] (verified) 2672
 3.291.4 Maple [A] (verified) 2674
 3.291.5 Fricas [A] (verification not implemented) 2674
 3.291.6 Sympy [F(-1)] 2675
 3.291.7 Maxima [F(-2)] 2675
 3.291.8 Giac [B] (verification not implemented) 2676
 3.291.9 Mupad [B] (verification not implemented) 2676

3.291.1 Optimal result

Integrand size = 32, antiderivative size = 61

$$\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{Bx}{b} - \frac{2aB \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b}b\sqrt{a+b}}$$

output `B*x/b-2*a*B*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/b/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.291.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.97

$$\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{B \left(c + dx + \frac{2a \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} \right)}{bd}$$

input `Integrate[(Cos[c + d*x]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `(B*(c + d*x + (2*a*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2])/(b*d)`

3.291.
$$\int \frac{\cos(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.291.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {2011, 3042, 3214, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\cos(c+dx)}{a+b\cos(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\sin(c+dx+\frac{\pi}{2})}{a+b\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3214} \\
 & B \left(\frac{x}{b} - \frac{a \int \frac{1}{a+b\cos(c+dx)} dx}{b} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{x}{b} - \frac{a \int \frac{1}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} \right) \\
 & \quad \downarrow \text{3138} \\
 & B \left(\frac{x}{b} - \frac{2a \int \frac{1}{(a-b)\tan^2(\frac{1}{2}(c+dx))+a+b} d \tan(\frac{1}{2}(c+dx))}{bd} \right) \\
 & \quad \downarrow \text{218} \\
 & B \left(\frac{x}{b} - \frac{2a \arctan\left(\frac{\sqrt{a-b}\tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{bd\sqrt{a-b}\sqrt{a+b}} \right)
 \end{aligned}$$

input `Int[(Cos[c + d*x]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output $B*(x/b - (2*a*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*b*Sqrt[a + b]*d))$

3.291.3.1 Defintions of rubi rules used

rule 218 $Int[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[\{a, b\}, x] \&\& PosQ[a/b]$

rule 2011 $Int[(u_)*((a_) + (b_)*(v_))^{(m_)}*((c_) + (d_)*(v_))^{(n_)}, x_Symbol] \rightarrow Simp[(b/d)^m Int[u*(c + d*v)^{(m + n)}, x], x] /; FreeQ[\{a, b, c, d, n\}, x] \&\& EqQ[b*c - a*d, 0] \&\& IntegerQ[m] \&\& (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])$

rule 3042 $Int[u_, x_Symbol] \rightarrow Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]$

rule 3138 $Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^{-1}, x_Symbol] \rightarrow With[\{e = FreeFactors[Tan[(c + d*x)/2], x]\}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[\{a, b, c, d\}, x] \&\& NeQ[a^2 - b^2, 0]$

rule 3214 $Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] \rightarrow Simp[b*(x/d), x] - Simp[(b*c - a*d)/d Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[\{a, b, c, d, e, f\}, x] \&\& NeQ[b*c - a*d, 0]$

3.291.4 Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{2B \left(\frac{\arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b} - \frac{a \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b\sqrt{(a-b)(a+b)}} \right)}{d}$	66
default	$\frac{2B \left(\frac{\arctan\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{b} - \frac{a \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{b\sqrt{(a-b)(a+b)}} \right)}{d}$	66
risch	$\frac{Bx}{b} - \frac{aB \ln\left(\frac{e^{i(dx+c)} + \frac{-ia^2+ib^2+a\sqrt{-a^2+b^2}}{\sqrt{-a^2+b^2}b}}{\sqrt{-a^2+b^2}db}\right) + aB \ln\left(\frac{e^{i(dx+c)} + \frac{ia^2-ib^2+a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}}{\sqrt{-a^2+b^2}db}\right)}$	153

input `int(cos(d*x+c)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

output $\frac{2}{d}B\left(\frac{1}{b}\arctan\left(\tan\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)-\frac{1}{b}a/\left(\left(a-b\right)\left(a+b\right)\right)^{\left(1/2\right)}\arctan\left(\left(a-b\right)\tan\left(\frac{1}{2}d*x+\frac{1}{2}c\right)/\left(\left(a-b\right)\left(a+b\right)\right)^{\left(1/2\right)}\right)\right)$

3.291.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 231, normalized size of antiderivative = 3.79

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \left[-\frac{\sqrt{-a^2+b^2}Ba \log\left(\frac{2ab\cos(dx+c)+(2a^2-b^2)\cos(dx+c)^2-2\sqrt{-a^2+b^2}(a\cos(dx+c)+b)\sin(dx+c)-a^2+2b^2}{b^2\cos(dx+c)^2+2ab\cos(dx+c)+a^2}\right) - 2(Ba^2 - Bb^2)}{2(a^2b - b^3)d} - \frac{\sqrt{a^2 - b^2}Ba \arctan\left(-\frac{a\cos(dx+c)+b}{\sqrt{a^2-b^2}\sin(dx+c)}\right) - (Ba^2 - Bb^2)dx}{(a^2b - b^3)d} \right]$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output `[-1/2*(sqrt(-a^2 + b^2)*B*a*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c))^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - 2*(B*a^2 - B*b^2)*d*x)/((a^2*b - b^3)*d), -(sqrt(a^2 - b^2)*B*a*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (B*a^2 - B*b^2)*d*x)/((a^2*b - b^3)*d)]`

3.291.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.291.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.291.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. 2(52) = 104.

Time = 0.34 (sec) , antiderivative size = 245, normalized size of antiderivative = 4.02

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \frac{(\sqrt{a^2-b^2}B(2a-b)|a-b|+\sqrt{a^2-b^2}B|a-b||b|) \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor + \arctan \left(\frac{2\sqrt{\frac{1}{2}} \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{\frac{2a+\sqrt{-4(a+b)(a-b)+4a^2}}{a-b}}} \right) \right)}{(a^2-2ab+b^2)b^2+(a^3-2a^2b+ab^2)|b|} + \frac{(2Ba-Bb-B|b|) \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor + \right)}{d}$$

input `integrate(cos(d*x+c)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `-((sqrt(a^2 - b^2)*B*(2*a - b)*abs(a - b) + sqrt(a^2 - b^2)*B*abs(a - b)*abs(b))*(pi*floor(1/2*(d*x + c)/pi + 1/2) + arctan(2*sqrt(1/2)*tan(1/2*d*x + 1/2*c)/sqrt((2*a + sqrt(-4*(a + b)*(a - b) + 4*a^2))/(a - b))))/(a^2 - 2*a*b + b^2)*b^2 + (a^3 - 2*a^2*b + a*b^2)*abs(b)) + (2*B*a - B*b - B*abs(b))*(pi*floor(1/2*(d*x + c)/pi + 1/2) + arctan(2*sqrt(1/2)*tan(1/2*d*x + 1/2*c)/sqrt((2*a - sqrt(-4*(a + b)*(a - b) + 4*a^2))/(a - b))))/(b^2 - a*abs(b))/d`

3.291.9 Mupad [B] (verification not implemented)

Time = 0.83 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.66

$$\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \frac{2B \operatorname{atan} \left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)} \right)}{bd} + \frac{2Ba \operatorname{atanh} \left(\frac{a \sin\left(\frac{c}{2} + \frac{dx}{2}\right) - b \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2}} \right)}{bd \sqrt{b^2 - a^2}}$$

input `int((cos(c + d*x)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

output `(2*B*atan(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(b*d) + (2*B*a*atanh((a*sin(c/2 + (d*x)/2) - b*sin(c/2 + (d*x)/2))/(cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))))/(b*d*(b^2 - a^2)^(1/2))`

3.291. $\int \frac{\cos(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$3.292 \quad \int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx$$

3.292.1 Optimal result	2677
3.292.2 Mathematica [A] (verified)	2677
3.292.3 Rubi [A] (verified)	2678
3.292.4 Maple [A] (verified)	2679
3.292.5 Fracas [A] (verification not implemented)	2680
3.292.6 Sympy [B] (verification not implemented)	2680
3.292.7 Maxima [F(-2)]	2681
3.292.8 Giac [A] (verification not implemented)	2681
3.292.9 Mupad [B] (verification not implemented)	2682

3.292.1 Optimal result

Integrand size = 26, antiderivative size = 50

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2B \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{\sqrt{a-b} \sqrt{a+bd}}$$

output `2*B*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.292.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.98

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = -\frac{2B \operatorname{Arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}d}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

output `(-2*B*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/(Sqrt[-a^2 + b^2]*d)`

$$3.292. \quad \int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx$$

3.292.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2011, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{a + b \cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3138} \\
 & \frac{2B \int \frac{1}{(a-b) \tan^2\left(\frac{1}{2}(c+dx)\right) + a+b} d \tan\left(\frac{1}{2}(c + dx)\right)}{d} \\
 & \quad \downarrow \text{218} \\
 & \frac{2B \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{d\sqrt{a-b}\sqrt{a+b}}
 \end{aligned}$$

input `Int[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

output `(2*B*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(Sqrt[a - b]*Sqrt[a + b]*d)`

3.292.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

- rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

3.292.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{2B \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{d\sqrt{(a-b)(a+b)}}$	45
default	$\frac{2B \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{d\sqrt{(a-b)(a+b)}}$	45
risch	$-\frac{\ln\left(e^{i(dx+c)} + \frac{ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)B}{\sqrt{-a^2+b^2}d} + \frac{B \ln\left(e^{i(dx+c)} - \frac{ia^2 - ib^2 - a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}d}$	141

input `int((B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

output `2/d*B/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))`

3.292.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 177, normalized size of antiderivative = 3.54

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \left[-\frac{\sqrt{-a^2 + b^2} B \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 + 2\sqrt{-a^2 + b^2}(a \cos(dx+c) + b) \sin(dx+c) - a^2 + 2b^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{2(a^2 - b^2)d}, B \arctan\left(\frac{-\frac{a}{\sqrt{a^2 - b^2}}}{\sqrt{a^2 - b^2} \sin(dx+c)}\right) \right]$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `[-1/2*sqrt(-a^2 + b^2)*B*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2))/((a^2 - b^2)*d), B*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c)))/(sqrt(a^2 - b^2)*d)]`

3.292.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 190 vs. 2(42) = 84.

Time = 155.04 (sec) , antiderivative size = 190, normalized size of antiderivative = 3.80

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \begin{cases} \frac{\infty Bx}{\cos(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{B \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{bd} & \text{for } a = b \\ \frac{B}{bd \tan\left(\frac{c}{2} + \frac{dx}{2}\right)} & \text{for } a = -b \\ \frac{x(Ba + Bb \cos(c))}{(a + b \cos(c))^2} & \text{for } d = 0 \\ \frac{B \log\left(-\sqrt{-\frac{a}{a-b} - \frac{b}{a-b}} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{ad \sqrt{-\frac{a}{a-b} - \frac{b}{a-b}} - bd \sqrt{-\frac{a}{a-b} - \frac{b}{a-b}}} - \frac{B \log\left(\sqrt{-\frac{a}{a-b} - \frac{b}{a-b}} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{ad \sqrt{-\frac{a}{a-b} - \frac{b}{a-b}} - bd \sqrt{-\frac{a}{a-b} - \frac{b}{a-b}}} & \text{otherwise} \end{cases}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

```
output Piecewise((zoo*B*x/cos(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*tan(c/2 + d
*x/2)/(b*d), Eq(a, b)), (B/(b*d*tan(c/2 + d*x/2)), Eq(a, -b)), (x*(B*a + B
*b*cos(c))/(a + b*cos(c))**2, Eq(d, 0)), (B*log(-sqrt(-a/(a - b) - b/(a -
b)) + tan(c/2 + d*x/2))/(a*d*sqrt(-a/(a - b) - b/(a - b)) - b*d*sqrt(-a/(a
- b) - b/(a - b))) - B*log(sqrt(-a/(a - b) - b/(a - b)) + tan(c/2 + d*x/2
))/(a*d*sqrt(-a/(a - b) - b/(a - b)) - b*d*sqrt(-a/(a - b) - b/(a - b))),
True))
```

3.292.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

```
input integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

3.292.8 Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.56

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2 \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a - 2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right) B}{\sqrt{a^2 - b^2} d}$$

```
input integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")
```

```
output 2*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x
+ 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))*B/(sqrt(a^2 - b^2)*d
)
```


3.292.9 Mupad [B] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.88

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2B \operatorname{atan}\left(\frac{\tan\left(\frac{c}{2} + \frac{dx}{2}\right)(a-b)}{\sqrt{a^2 - b^2}}\right)}{d \sqrt{a^2 - b^2}}$$

input `int((B*a + B*b*cos(c + d*x))/(a + b*cos(c + d*x))^2,x)`output `(2*B*atan((tan(c/2 + (d*x)/2)*(a - b))/(a^2 - b^2)^(1/2)))/(d*(a^2 - b^2)^(1/2))`

3.293 $\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx$

3.293.1 Optimal result 2683
 3.293.2 Mathematica [A] (verified) 2683
 3.293.3 Rubi [A] (verified) 2684
 3.293.4 Maple [A] (verified) 2686
 3.293.5 Fricas [A] (verification not implemented) 2686
 3.293.6 Sympy [F] 2687
 3.293.7 Maxima [F(-2)] 2687
 3.293.8 Giac [A] (verification not implemented) 2688
 3.293.9 Mupad [B] (verification not implemented) 2688

3.293.1 Optimal result

Integrand size = 32, antiderivative size = 70

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = -\frac{2bB \arctan\left(\frac{\sqrt{a-b} \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{a\sqrt{a-b}\sqrt{a+bd}} + \frac{B \operatorname{arctanh}(\sin(c + dx))}{ad}$$

output `B*arctanh(sin(d*x+c))/a/d-2*b*B*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a/d/(a-b)^(1/2)/(a+b)^(1/2)`

3.293.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.47

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{B \left(\frac{2b \operatorname{arctanh}\left(\frac{(a-b) \tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} - \log\left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right) + \log\left(\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right)\right) \right)}{ad}$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

3.293. $\int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^2} dx$

output $(B*((2*b*ArcTanh[((a - b)*Tan[(c + d*x)/2]])/Sqrt[-a^2 + b^2]))/Sqrt[-a^2 + b^2] - Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]])/(a*d)$

3.293.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {2011, 3042, 3226, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx$$

$$\downarrow 2011$$

$$B \int \frac{\sec(c + dx)}{a + b \cos(c + dx)} dx$$

$$\downarrow 3042$$

$$B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})(a + b \sin(c + dx + \frac{\pi}{2}))} dx$$

$$\downarrow 3226$$

$$B \left(\frac{\int \sec(c + dx) dx}{a} - \frac{b \int \frac{1}{a + b \cos(c + dx)} dx}{a} \right)$$

$$\downarrow 3042$$

$$B \left(\frac{\int \csc(c + dx + \frac{\pi}{2}) dx}{a} - \frac{b \int \frac{1}{a + b \sin(c + dx + \frac{\pi}{2})} dx}{a} \right)$$

$$\downarrow 3138$$

$$B \left(\frac{\int \csc(c + dx + \frac{\pi}{2}) dx}{a} - \frac{2b \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c + dx))}{ad} \right)$$

$$\downarrow 218$$

$$B \left(\frac{\int \csc(c + dx + \frac{\pi}{2}) dx}{a} - \frac{2b \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right)$$

$$B \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2b \operatorname{arctan}\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right)$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^2,x]`

output `B*((-2*b*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b] *Sqrt[a + b]*d) + ArcTanh[Sin[c + d*x]]/(a*d))`

3.293.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3226 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.293.4 Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{2B \left(-\frac{b \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a\sqrt{(a-b)(a+b)}} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2a} \right)}{d}$
default	$\frac{2B \left(-\frac{b \arctan\left(\frac{(a-b)\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a\sqrt{(a-b)(a+b)}} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2a} \right)}{d}$
risch	$-\frac{bB \ln\left(\frac{e^{i(dx+c)} - ia^2 - ib^2 - a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} da} + \frac{bB \ln\left(\frac{e^{i(dx+c)} + ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} da} + \frac{\ln(e^{i(dx+c)} + i)B}{ad} - \frac{\ln(e^{i(dx+c)} - i)B}{ad}$

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)`

output `2/d*B*(-1/a*b/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))+1/2/a*ln(tan(1/2*d*x+1/2*c)+1)-1/2/a*ln(tan(1/2*d*x+1/2*c)-1))`

3.293.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 292, normalized size of antiderivative = 4.17

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \left[\frac{\sqrt{-a^2 + b^2} B b \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 - 2\sqrt{-a^2+b^2}(a \cos(dx+c) + b) \sin(dx+c) - a^2 + 2b^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right) - (Ba^2 - Bb^2)}{2(a^3 - ab^2)d} \right. \\ \left. - \frac{2\sqrt{a^2 - b^2} B b \arctan\left(-\frac{a \cos(dx+c) + b}{\sqrt{a^2 - b^2} \sin(dx+c)}\right) - (Ba^2 - Bb^2) \log(\sin(dx+c) + 1) + (Ba^2 - Bb^2) \log(-\sin(dx+c))}{2(a^3 - ab^2)d} \right]$$

3.293. $\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `[-1/2*(sqrt(-a^2 + b^2)*B*b*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c))^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - (B*a^2 - B*b^2)*log(sin(d*x + c) + 1) + (B*a^2 - B*b^2)*log(-sin(d*x + c) + 1))/((a^3 - a*b^2)*d), -1/2*(2*sqrt(a^2 - b^2)*B*b*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c))) - (B*a^2 - B*b^2)*log(sin(d*x + c) + 1) + (B*a^2 - B*b^2)*log(-sin(d*x + c) + 1))/((a^3 - a*b^2)*d)]`

3.293.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = B \int \frac{\sec(c + dx)}{a + b \cos(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**2,x)`

output `B*Integral(sec(c + d*x)/(a + b*cos(c + d*x)), x)`

3.293.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.293.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.74

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx =$$

$$\frac{2 \left(\pi \left\lfloor \frac{dx+c}{2} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right) Bb}{\sqrt{a^2 - b^2} a} - \frac{B \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a} + \frac{B \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `-(2*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))*B*b/(sqrt(a^2 - b^2)*a) - B*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a + B*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a)/d`

3.293.9 Mupad [B] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.44

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2 B \operatorname{atanh} \left(\frac{\sin(\frac{c}{2} + \frac{dx}{2})}{\cos(\frac{c}{2} + \frac{dx}{2})} \right)}{a d} + \frac{2 B b \operatorname{atanh} \left(\frac{a \sin(\frac{c}{2} + \frac{dx}{2}) - b \sin(\frac{c}{2} + \frac{dx}{2})}{\cos(\frac{c}{2} + \frac{dx}{2}) \sqrt{b^2 - a^2}} \right)}{a d \sqrt{b^2 - a^2}}$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^2),x)`

output `(2*B*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)))/(a*d) + (2*B*b*atanh((a*sin(c/2 + (d*x)/2) - b*sin(c/2 + (d*x)/2))/(cos(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2)))/(a*d*(b^2 - a^2)^(1/2))`

3.294
$$\int \frac{(aB+bB \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.294.1 Optimal result 2689
 3.294.2 Mathematica [A] (verified) 2689
 3.294.3 Rubi [A] (verified) 2690
 3.294.4 Maple [A] (verified) 2693
 3.294.5 Fricas [B] (verification not implemented) 2694
 3.294.6 Sympy [F] 2694
 3.294.7 Maxima [F(-2)] 2695
 3.294.8 Giac [A] (verification not implemented) 2695
 3.294.9 Mupad [B] (verification not implemented) 2696

3.294.1 Optimal result

Integrand size = 34, antiderivative size = 88

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2b^2 B \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^2 \sqrt{a-b} \sqrt{a+b} d} - \frac{b B \operatorname{arctanh}(\sin(c + dx))}{a^2 d} + \frac{B \tan(c + dx)}{a d}$$

output `-b*B*arctanh(sin(d*x+c))/a^2/d+2*b^2*B*arctan((a-b)^(1/2)*tan(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^2/d/(a-b)^(1/2)/(a+b)^(1/2)+B*tan(d*x+c)/a/d`

3.294.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.32

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$B \left(-\frac{2b^2 \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + b(\log(\cos(\frac{1}{2}(c+dx))) - \sin(\frac{1}{2}(c+dx))) - \log(\cos(\frac{1}{2}(c+dx))) + \sin(\frac{1}{2}(c+dx))) \right)$$

$a^2 d$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^2, x]`

3.294.
$$\int \frac{(aB+bB \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$$

output $(B*((-2*b^2*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] + b*(Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]]) + a*Tan[c + d*x]))/(a^2*d)$

3.294.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {2011, 3042, 3281, 25, 27, 3042, 3226, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^2(c+dx)(aB + bB \cos(c+dx))}{(a + b \cos(c+dx))^2} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{\sec^2(c+dx)}{a + b \cos(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{1}{\sin(c+dx + \frac{\pi}{2})^2 (a + b \sin(c+dx + \frac{\pi}{2}))} dx \\ & \quad \downarrow \text{3281} \\ & B \left(\frac{\int -\frac{b \sec(c+dx)}{a+b \cos(c+dx)} dx}{a} + \frac{\tan(c+dx)}{ad} \right) \\ & \quad \downarrow \text{25} \\ & B \left(\frac{\tan(c+dx)}{ad} - \frac{\int \frac{b \sec(c+dx)}{a+b \cos(c+dx)} dx}{a} \right) \\ & \quad \downarrow \text{27} \\ & B \left(\frac{\tan(c+dx)}{ad} - \frac{b \int \frac{\sec(c+dx)}{a+b \cos(c+dx)} dx}{a} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{\tan(c+dx)}{ad} - \frac{b \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) (a + b \sin(c+dx + \frac{\pi}{2}))} dx}{a} \right) \end{aligned}$$

$$\begin{array}{c}
\downarrow \text{3226} \\
B \left(\frac{\tan(c+dx)}{ad} - \frac{b \left(\frac{\int \sec(c+dx) dx}{a} - \frac{b \int \frac{1}{a+b \cos(c+dx)} dx}{a} \right)}{a} \right) \\
\downarrow \text{3042} \\
B \left(\frac{\tan(c+dx)}{ad} - \frac{b \left(\frac{\int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{a} \right)}{a} \right) \\
\downarrow \text{3138} \\
B \left(\frac{\tan(c+dx)}{ad} - \frac{b \left(\frac{\int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} d \tan(\frac{1}{2}(c+dx))}{ad} \right)}{a} \right) \\
\downarrow \text{218} \\
B \left(\frac{\tan(c+dx)}{ad} - \frac{b \left(\frac{\int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right)}{a} \right) \\
\downarrow \text{4257} \\
B \left(\frac{\tan(c+dx)}{ad} - \frac{b \left(\frac{\operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{2b \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}} \right)}{a} \right)
\end{array}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^2,x]`

output $B * (-(b * (-2 * b * \text{ArcTan}[\sqrt{a - b} * \text{Tan}[(c + d * x) / 2]] / \sqrt{a + b}]) / (a * \sqrt{a - b} * \sqrt{a + b} * d) + \text{ArcTanh}[\text{Sin}[c + d * x]] / (a * d)) / a + \text{Tan}[c + d * x] / (a * d))$

3.294.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F x_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F x, x], x]$

rule 27 $\text{Int}[(a_)*(F x_), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F x, (b_)*(G x_)] /; \text{FreeQ}[b, x]$

rule 218 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b]$

rule 2011 $\text{Int}[(u_)*((a_) + (b_)*(v_))^{(m_)}*((c_) + (d_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(b/d)^m \text{Int}[u*(c + d*v)^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{EqQ}[b*c - a*d, 0] \&\& \text{IntegerQ}[m] \&\& (\text{!IntegerQ}[n] || \text{SimplerQ}[c + d*x, a + b*x])$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3138 $\text{Int}[(a_) + (b_)*\text{sin}[\text{Pi}/2 + (c_) + (d_)*(x_)]^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 3226 $\text{Int}[1/(((a_) + (b_)*\text{sin}[(e_) + (f_)*(x_)])*((c_) + (d_)*\text{sin}[(e_) + (f_)*(x_)])), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{Int}[1/(a + b*\text{Sin}[e + f*x]), x], x] - \text{Simp}[d/(b*c - a*d) \text{Int}[1/(c + d*\text{Sin}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$

```
rule 3281 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Sin[e + f*
x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2
))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) + b^2*d*(m + n
+ 2) - (b^2*c + b*(b*c - a*d)*(m + 1))*Sin[e + f*x] - b^2*d*(m + n + 3)*Si
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && IntegersQ[
2*m, 2*n] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*
n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))
```

```
rule 4257 Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

3.294.4 Maple [A] (verified)

Time = 1.32 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.42

method	result
derivativedivides	$2B \left(-\frac{1}{2a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} + \frac{b \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2a^2} + \frac{b^2 \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^2 \sqrt{(a-b)(a+b)}} - \frac{1}{2a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)} - \frac{b \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a^2} \right)$
default	$2B \left(-\frac{1}{2a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} + \frac{b \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{2a^2} + \frac{b^2 \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^2 \sqrt{(a-b)(a+b)}} - \frac{1}{2a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)} - \frac{b \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{2a^2} \right)$
risch	$\frac{2iB}{da(e^{2i(dx+c)}+1)} - \frac{b^2 B \ln\left(e^{i(dx+c)} + \frac{ia^2 - ib^2 + a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} da^2} + \frac{b^2 B \ln\left(e^{i(dx+c)} - \frac{ia^2 - ib^2 - a\sqrt{-a^2+b^2}}{b\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} da^2} + \frac{Bb \ln(e^{i(dx+c)} + 1)}{da}$

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

```
output 2/d*B*(-1/2/a/(tan(1/2*d*x+1/2*c)-1)+1/2*b/a^2*ln(tan(1/2*d*x+1/2*c)-1)+b^
2/a^2/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1
/2))-1/2/a/(tan(1/2*d*x+1/2*c)+1)-1/2*b/a^2*ln(tan(1/2*d*x+1/2*c)+1))
```

$$3.294. \int \frac{(aB+bB \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.294.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 165 vs. 2(79) = 158.

Time = 0.37 (sec) , antiderivative size = 398, normalized size of antiderivative = 4.52

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \left[-\frac{\sqrt{-a^2 + b^2} B b^2 \cos(dx + c) \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 + 2\sqrt{-a^2 + b^2}(a \cos(dx+c) + b) \sin(dx+c) - a^2 + 2b^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right) + (B a^2 b - B b^3) \cos(dx + c) \log(\sin(dx + c) + 1) - (B a^2 b - B b^3) \cos(dx + c) \log(-\sin(dx + c) + 1) - 2(B a^3 - B a b^2) \sin(dx + c) / ((a^4 - a^2 b^2) d \cos(dx + c))}{(a + b \cos(c + dx))^2} \right]$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `[-1/2*(sqrt(-a^2 + b^2)*B*b^2*cos(d*x + c)*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 + 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) + (B*a^2*b - B*b^3)*cos(d*x + c)*log(sin(d*x + c) + 1) - (B*a^2*b - B*b^3)*cos(d*x + c)*log(-sin(d*x + c) + 1) - 2*(B*a^3 - B*a*b^2)*sin(d*x + c)/((a^4 - a^2*b^2)*d*cos(d*x + c)), 1/2*(2*sqrt(a^2 - b^2)*B*b^2*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c)))*cos(d*x + c) - (B*a^2*b - B*b^3)*cos(d*x + c)*log(sin(d*x + c) + 1) + (B*a^2*b - B*b^3)*cos(d*x + c)*log(-sin(d*x + c) + 1) + 2*(B*a^3 - B*a*b^2)*sin(d*x + c)/((a^4 - a^2*b^2)*d*cos(d*x + c))]`

3.294.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = B \int \frac{\sec^2(c + dx)}{a + b \cos(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**2,x)`

output `B*Integral(sec(c + d*x)**2/(a + b*cos(c + d*x)), x)`

3.294.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de

3.294.8 Giac [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.76

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2 \left(\pi \left\lfloor \frac{dx+c}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right) Bb^2}{\sqrt{a^2 - b^2} a^2} - \frac{Bb \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^2} + \frac{Bb \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|)}{a^2}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `(2*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))*B*b^2/(sqrt(a^2 - b^2)*a^2) - B*b*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^2 + B*b*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^2 - 2*B*tan(1/2*d*x + 1/2*c)/((tan(1/2*d*x + 1/2*c)^2 - 1)*a))/d`

3.294.9 Mupad [B] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 326, normalized size of antiderivative = 3.70

$$\int \frac{(aB + bB \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^2} dx = \frac{2B \left(\frac{a^3 \sin(c+dx)}{2} - \frac{ab^2 \sin(c+dx)}{2} \right)}{a^2 d \cos(c + dx) (a^2 - b^2)}$$

$$\frac{2B \left(a^2 b \operatorname{atanh} \left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)} \right) - b^3 \operatorname{atanh} \left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\cos\left(\frac{c}{2} + \frac{dx}{2}\right)} \right) + b^2 \operatorname{atanh} \left(\frac{a^5 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{b^2 - a^2} + 2b^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) (b^2 - a^2)}{a^2 d (a^2 - b^2)} \right)}{a^2 d (a^2 - b^2)}$$

```
input int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^2),x)
```

```
output (2*B*((a^3*sin(c + d*x))/2 - (a*b^2*sin(c + d*x))/2))/(a^2*d*cos(c + d*x)*
(a^2 - b^2)) - (2*B*(a^2*b*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)) -
b^3*atanh(sin(c/2 + (d*x)/2)/cos(c/2 + (d*x)/2)) + b^2*atanh((a^5*sin(c/2
+ (d*x)/2)*(b^2 - a^2)^(1/2) + 2*b^3*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(3/2)
- 2*b^5*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 3*a^2*b^3*sin(c/2 + (d*x)/2
)*(b^2 - a^2)^(1/2) - a^3*b^2*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - a^4*b
*sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))/(cos(c/2 + (d*x)/2)*(a*b^2 - a^3)^2
))*(b^2 - a^2)^(1/2)))/(a^2*d*(a^2 - b^2))
```

3.295
$$\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.295.1 Optimal result 2697
 3.295.2 Mathematica [A] (verified) 2697
 3.295.3 Rubi [A] (verified) 2698
 3.295.4 Maple [A] (verified) 2702
 3.295.5 Fricas [A] (verification not implemented) 2702
 3.295.6 Sympy [F] 2703
 3.295.7 Maxima [F(-2)] 2703
 3.295.8 Giac [B] (verification not implemented) 2704
 3.295.9 Mupad [B] (verification not implemented) 2704

3.295.1 Optimal result

Integrand size = 34, antiderivative size = 123

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = -\frac{2b^3 B \arctan\left(\frac{\sqrt{a-b} \tan(\frac{1}{2}(c+dx))}{\sqrt{a+b}}\right)}{a^3 \sqrt{a-b} \sqrt{a+bd}} + \frac{(a^2 + 2b^2) B \operatorname{arctanh}(\sin(c + dx))}{2a^3 d} - \frac{bB \tan(c + dx)}{a^2 d} + \frac{B \sec(c + dx) \tan(c + dx)}{2ad}$$

```
output 1/2*(a^2+2*b^2)*B*arctanh(sin(d*x+c))/a^3/d-2*b^3*B*arctan((a-b)^(1/2)*tan
(1/2*d*x+1/2*c)/(a+b)^(1/2))/a^3/d/(a-b)^(1/2)/(a+b)^(1/2)-b*B*tan(d*x+c)/
a^2/d+1/2*B*sec(d*x+c)*tan(d*x+c)/a/d
```

3.295.2 Mathematica [A] (verified)

Time = 0.76 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.94

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$B \left(\frac{8b^3 \operatorname{arctanh}\left(\frac{(a-b) \tan(\frac{1}{2}(c+dx))}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} - 2a^2 \log\left(\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right)\right) - 4b^2 \log\left(\cos\left(\frac{1}{2}(c+dx)\right)\right) \right)$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^2, x]`

output $(B*((8*b^3*ArcTanh[((a - b)*Tan[(c + d*x)/2])/Sqrt[-a^2 + b^2]])/Sqrt[-a^2 + b^2] - 2*a^2*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - 4*b^2*Log[Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 2*a^2*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 4*b^2*Log[Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + a^2/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^2 - a^2/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 - 4*a*b*Tan[c + d*x]))/(4*a^3*d)$

3.295.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {2011, 3042, 3281, 25, 3042, 3534, 25, 3042, 3480, 3042, 3138, 218, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\sec^3(c + dx)}{a + b \cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^3 (a + b \sin(c + dx + \frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{3281} \\
 & B \left(\frac{\int -\frac{(-b \cos^2(c + dx) - a \cos(c + dx) + 2b) \sec^2(c + dx)}{a + b \cos(c + dx)} dx}{2a} + \frac{\tan(c + dx) \sec(c + dx)}{2ad} \right) \\
 & \quad \downarrow \text{25} \\
 & B \left(\frac{\tan(c + dx) \sec(c + dx)}{2ad} - \frac{\int \frac{(-b \cos^2(c + dx) - a \cos(c + dx) + 2b) \sec^2(c + dx)}{a + b \cos(c + dx)} dx}{2a} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.295. $\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx$

$$\begin{aligned}
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int \frac{-b \sin(c+dx+\frac{\pi}{2})^2 - a \sin(c+dx+\frac{\pi}{2}) + 2b}{\sin(c+dx+\frac{\pi}{2})^2 (a+b \sin(c+dx+\frac{\pi}{2}))} dx}{2a} \right) \\
& \quad \downarrow \text{3534} \\
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\int -\frac{(a^2+b \cos(c+dx)a+2b^2) \sec(c+dx)}{a+b \cos(c+dx)} dx + \frac{2b \tan(c+dx)}{ad}}{2a} \right) \\
& \quad \downarrow \text{25} \\
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\frac{2b \tan(c+dx)}{ad} - \int \frac{(a^2+b \cos(c+dx)a+2b^2) \sec(c+dx)}{a+b \cos(c+dx)} dx}{2a} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\frac{2b \tan(c+dx)}{ad} - \int \frac{a^2+b \sin(c+dx+\frac{\pi}{2})a+2b^2}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{2a} \right) \\
& \quad \downarrow \text{3480} \\
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\frac{2b \tan(c+dx)}{ad} - \frac{(a^2+2b^2) \int \sec(c+dx) dx}{a} - \frac{2b^3 \int \frac{1}{a+b \cos(c+dx)} dx}{a}}{2a} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\frac{2b \tan(c+dx)}{ad} - \frac{(a^2+2b^2) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{2b^3 \int \frac{1}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{a}}{2a} \right) \\
& \quad \downarrow \text{3138} \\
& B \left(\frac{\tan(c+dx) \sec(c+dx)}{2ad} - \frac{\frac{2b \tan(c+dx)}{ad} - \frac{(a^2+2b^2) \int \csc(c+dx+\frac{\pi}{2}) dx}{a} - \frac{4b^3 \int \frac{1}{(a-b) \tan^2(\frac{1}{2}(c+dx)) + a+b} dx}{a}}{2a} \right) \\
& \quad \downarrow \text{218}
\end{aligned}$$

$$\begin{aligned}
 & B \left(\frac{\tan(c+dx)\sec(c+dx)}{2ad} - \frac{2b\tan(c+dx)}{ad} - \frac{\frac{(a^2+2b^2)\int\csc(c+dx+\frac{\pi}{2})dx}{a} - \frac{4b^3\arctan\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{2a} \right) \\
 & \quad \downarrow 4257 \\
 & B \left(\frac{\tan(c+dx)\sec(c+dx)}{2ad} - \frac{2b\tan(c+dx)}{ad} - \frac{\frac{(a^2+2b^2)\operatorname{arctanh}(\sin(c+dx))}{ad} - \frac{4b^3\arctan\left(\frac{\sqrt{a-b}\tan\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b}}\right)}{ad\sqrt{a-b}\sqrt{a+b}}}{2a} \right)
 \end{aligned}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^2,x]`

output `B*((Sec[c + d*x]*Tan[c + d*x])/(2*a*d) - (-(((-4*b^3*ArcTan[(Sqrt[a - b]*Tan[(c + d*x)/2])/Sqrt[a + b]])/(a*Sqrt[a - b]*Sqrt[a + b]*d) + ((a^2 + 2*b^2)*ArcTanh[Sin[c + d*x]])/(a*d))/a) + (2*b*Tan[c + d*x])/(a*d))/(2*a))`

3.295.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3138 `Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3281 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) + b^2*d*(m + n + 2) - (b^2*c + b*(b*c - a*d)*(m + 1))*Sin[e + f*x] - b^2*d*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && IntegersQ[2*m, 2*n] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3480 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b - a*B)/(b*c - a*d) Int[1/(a + b*Sin[e + f*x]), x], x] + Simp[(B*c - A*d)/(b*c - a*d) Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

3.295.4 Maple [A] (verified)

Time = 1.42 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.58

method	result
derivativedivides	$2B \left(-\frac{1}{4a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)^2} - \frac{-a-2b}{4a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)} + \frac{(a^2+2b^2) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4a^3} - \frac{b^3 \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^3 \sqrt{(a-b)(a+b)}} + \frac{1}{4a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)^2} + \frac{-a-2b}{4a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} + \frac{(a^2+2b^2) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4a^3} - \frac{b^3 \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^3 \sqrt{(a-b)(a+b)}} + \frac{1}{4a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} \right)$
default	$2B \left(-\frac{1}{4a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)^2} - \frac{-a-2b}{4a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)} + \frac{(a^2+2b^2) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4a^3} - \frac{b^3 \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^3 \sqrt{(a-b)(a+b)}} + \frac{1}{4a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)^2} + \frac{-a-2b}{4a^2 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} + \frac{(a^2+2b^2) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4a^3} - \frac{b^3 \arctan\left(\frac{(a-b) \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{\sqrt{(a-b)(a+b)}}\right)}{a^3 \sqrt{(a-b)(a+b)}} + \frac{1}{4a \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)} \right)$
risch	$-\frac{iB(ae^{3i(dx+c)}+2be^{2i(dx+c)}-ae^{i(dx+c)}+2b)}{da^2(e^{2i(dx+c)}+1)^2} - \frac{\ln(e^{i(dx+c)}-i)B}{2ad} - \frac{\ln(e^{i(dx+c)}-i)Bb^2}{a^3d} + \frac{\ln(e^{i(dx+c)}+i)B}{2ad} + \frac{\ln(e^{i(dx+c)}+i)Bb^2}{a^3d}$

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^2,x,method=_RETURNV ERBOSE)`

output `2/d*B*(-1/4/a/(tan(1/2*d*x+1/2*c)+1)^2-1/4*(-a-2*b)/a^2/(tan(1/2*d*x+1/2*c)+1)+1/4*(a^2+2*b^2)/a^3*ln(tan(1/2*d*x+1/2*c)+1)-b^3/a^3/((a-b)*(a+b))^(1/2)*arctan((a-b)*tan(1/2*d*x+1/2*c)/((a-b)*(a+b))^(1/2))+1/4/a/(tan(1/2*d*x+1/2*c)-1)^2-1/4*(-a-2*b)/a^2/(tan(1/2*d*x+1/2*c)-1)+1/4/a^3*(-a^2-2*b^2)*ln(tan(1/2*d*x+1/2*c)-1))`

3.295.5 Fracas [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 487, normalized size of antiderivative = 3.96

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \left[-\frac{2\sqrt{-a^2 + b^2} B b^3 \cos(dx + c)^2 \log\left(\frac{2ab \cos(dx+c) + (2a^2 - b^2) \cos(dx+c)^2 - 2\sqrt{-a^2 + b^2}(a \cos(dx+c) + b) \sin(dx+c) - a^2 + 2b^2}{b^2 \cos(dx+c)^2 + 2ab \cos(dx+c) + a^2}\right)}{4\sqrt{a^2 - b^2} B b^3 \arctan\left(-\frac{a \cos(dx+c) + b}{\sqrt{a^2 - b^2} \sin(dx+c)}\right) \cos(dx + c)^2 - (Ba^4 + Ba^2b^2 - 2Bb^4) \cos(dx + c)^2 \log(\sin(dx + c))} \right]$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^2,x, algorithm m="fricas")`

3.295.
$$\int \frac{(aB+bB \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^2} dx$$

output `[-1/4*(2*sqrt(-a^2 + b^2)*B*b^3*cos(d*x + c)^2*log((2*a*b*cos(d*x + c) + (2*a^2 - b^2)*cos(d*x + c)^2 - 2*sqrt(-a^2 + b^2)*(a*cos(d*x + c) + b)*sin(d*x + c) - a^2 + 2*b^2)/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2)) - (B*a^4 + B*a^2*b^2 - 2*B*b^4)*cos(d*x + c)^2*log(sin(d*x + c) + 1) + (B*a^4 + B*a^2*b^2 - 2*B*b^4)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) - 2*(B*a^4 - B*a^2*b^2 - 2*(B*a^3*b - B*a*b^3)*cos(d*x + c))*sin(d*x + c))/((a^5 - a^3*b^2)*d*cos(d*x + c)^2), -1/4*(4*sqrt(a^2 - b^2)*B*b^3*arctan(-(a*cos(d*x + c) + b)/(sqrt(a^2 - b^2)*sin(d*x + c)))*cos(d*x + c)^2 - (B*a^4 + B*a^2*b^2 - 2*B*b^4)*cos(d*x + c)^2*log(sin(d*x + c) + 1) + (B*a^4 + B*a^2*b^2 - 2*B*b^4)*cos(d*x + c)^2*log(-sin(d*x + c) + 1) - 2*(B*a^4 - B*a^2*b^2 - 2*(B*a^3*b - B*a*b^3)*cos(d*x + c))*sin(d*x + c))/((a^5 - a^3*b^2)*d*cos(d*x + c)^2)]`

3.295.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = B \int \frac{\sec^3(c + dx)}{a + b \cos(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**2,x)`

output `B*Integral(sec(c + d*x)**3/(a + b*cos(c + d*x)), x)`

3.295.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

3.295.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(110) = 220$.

Time = 0.31 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.80

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx =$$

$$\frac{4 \left(\pi \left[\frac{dx+c}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(2a-2b) + \arctan \left(\frac{a \tan(\frac{1}{2} dx + \frac{1}{2} c) - b \tan(\frac{1}{2} dx + \frac{1}{2} c)}{\sqrt{a^2 - b^2}} \right) \right) Bb^3}{\sqrt{a^2 - b^2} a^3} - \frac{(Ba^2 + 2Bb^2) \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1|)}{a^3} + \frac{(Ba^2 + 2Bb^2)}{2d}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `-1/2*(4*(pi*floor(1/2*(d*x + c)/pi + 1/2)*sgn(2*a - 2*b) + arctan((a*tan(1/2*d*x + 1/2*c) - b*tan(1/2*d*x + 1/2*c))/sqrt(a^2 - b^2)))*B*b^3/(sqrt(a^2 - b^2)*a^3) - (B*a^2 + 2*B*b^2)*log(abs(tan(1/2*d*x + 1/2*c) + 1))/a^3 + (B*a^2 + 2*B*b^2)*log(abs(tan(1/2*d*x + 1/2*c) - 1))/a^3 - 2*(B*a*tan(1/2*d*x + 1/2*c))^3 + 2*B*b*tan(1/2*d*x + 1/2*c)^3 + B*a*tan(1/2*d*x + 1/2*c) - 2*B*b*tan(1/2*d*x + 1/2*c))/((tan(1/2*d*x + 1/2*c)^2 - 1)^2*a^2)/d`

3.295.9 Mupad [B] (verification not implemented)

Time = 1.90 (sec) , antiderivative size = 1099, normalized size of antiderivative = 8.93

$$\int \frac{(aB + bB \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Too large to display}$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^2),x)`

output

$$\begin{aligned}
& ((B*b^2*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 - (B*b^2*\sin(c + d \\
& *x))/2 + (B*b^2*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))*\cos(2*c + 2*d \\
& *x))/2)/(a*d*(a^2 - b^2)*(\cos(2*c + 2*d*x)/2 + 1/2)) + (a*((B*\sin(c + d*x) \\
&)/2 + (B*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2)))/2 + (B*atanh(\sin(c/ \\
& 2 + (d*x)/2)/\cos(c/2 + (d*x)/2))*\cos(2*c + 2*d*x))/2)/(d*(a^2 - b^2)*(\cos \\
& (2*c + 2*d*x)/2 + 1/2)) - (B*b*\sin(2*c + 2*d*x))/(2*d*(a^2 - b^2)*(\cos(2*c \\
& + 2*d*x)/2 + 1/2)) - (B*b^4*atanh(\sin(c/2 + (d*x)/2)/\cos(c/2 + (d*x)/2))) \\
& /(a^3*d*(a^2 - b^2)*(\cos(2*c + 2*d*x)/2 + 1/2)) + (B*b^3*\sin(2*c + 2*d*x)) \\
& /(2*a^2*d*(a^2 - b^2)*(\cos(2*c + 2*d*x)/2 + 1/2)) - (B*b^3*atan(((a^9*\sin(\\
& c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 8*b^7*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(3 \\
& /2) - 8*b^9*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 8*a^2*b^7*\sin(c/2 + (d* \\
& x)/2)*(b^2 - a^2)^(1/2) + 3*a^4*b^5*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - \\
& 3*a^5*b^4*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - 2*a^6*b^3*\sin(c/2 + (d*x \\
&)/2)*(b^2 - a^2)^(1/2) + 2*a^7*b^2*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - \\
& a^8*b*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2))*i)/(\cos(c/2 + (d*x)/2)*(a*b^2 \\
& - a^3)*(a^7 - 3*a^3*b^4 + 2*a^5*b^2))*i)/(a^3*d*(b^2 - a^2)^(1/2)*(\cos(\\
& 2*c + 2*d*x)/2 + 1/2)) - (B*b^3*atan(((a^9*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(\\
& 1/2) + 8*b^7*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(3/2) - 8*b^9*\sin(c/2 + (d*x) \\
& /2)*(b^2 - a^2)^(1/2) + 8*a^2*b^7*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) + 3 \\
& *a^4*b^5*\sin(c/2 + (d*x)/2)*(b^2 - a^2)^(1/2) - 3*a^5*b^4*\sin(c/2 + (d*...
\end{aligned}$$

3.296 $\int \cos^3(c+dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.296.1 Optimal result	2706
3.296.2 Mathematica [A] (verified)	2707
3.296.3 Rubi [A] (verified)	2708
3.296.4 Maple [B] (verified)	2714
3.296.5 Fricas [C] (verification not implemented)	2715
3.296.6 Sympy [F(-1)]	2716
3.296.7 Maxima [F]	2716
3.296.8 Giac [F]	2717
3.296.9 Mupad [F(-1)]	2717

3.296.1 Optimal result

Integrand size = 33, antiderivative size = 386

$$\begin{aligned}
 & \int \cos^3(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\
 = & \frac{2(24a^3Ab + 57aAb^3 - 16a^4B - 24a^2b^2B + 147b^4B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{315b^4d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & - \frac{2(a^2 - b^2) (24a^2Ab + 75Ab^3 - 16a^3B - 36ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{315b^4d \sqrt{a + b \cos(c + dx)}} \\
 & + \frac{2(24a^2Ab + 75Ab^3 - 16a^3B - 36ab^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{315b^3d} \\
 & - \frac{2(36aAb - 24a^2B - 49b^2B) (a + b \cos(c + dx))^{3/2} \sin(c + dx)}{315b^3d} \\
 & + \frac{2(3Ab - 2aB) \cos(c + dx) (a + b \cos(c + dx))^{3/2} \sin(c + dx)}{21b^2d} \\
 & + \frac{2B \cos^2(c + dx) (a + b \cos(c + dx))^{3/2} \sin(c + dx)}{9bd}
 \end{aligned}$$

output
$$\begin{aligned} & -2/315*(36*A*a*b-24*B*a^2-49*B*b^2)*(a+b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/b^3/ \\ & d+2/21*(3*A*b-2*B*a)*\cos(d*x+c)*(a+b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/b^2/d+2/ \\ & 9*B*\cos(d*x+c)^2*(a+b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/b/d+2/315*(24*A*a^2*b+7 \\ & 5*A*b^3-16*B*a^3-36*B*a*b^2)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{1/2}/b^3/d+2/315 \\ & *(24*A*a^3*b+57*A*a*b^3-16*B*a^4-24*B*a^2*b^2+147*B*b^4)*(\cos(1/2*d*x+1/2* \\ & c))^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{1/2}*(b/(a+ \\ & b))^{1/2})*(a+b*\cos(d*x+c))^{1/2}/b^4/d/((a+b*\cos(d*x+c))/(a+b))^{1/2}-2/3 \\ & 15*(a^2-b^2)*(24*A*a^2*b+75*A*b^3-16*B*a^3-36*B*a*b^2)*(\cos(1/2*d*x+1/2*c) \\ & ^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{1/2}*(b/(a+b) \\ &)^{1/2})*((a+b*\cos(d*x+c))/(a+b))^{1/2}/b^4/d/(a+b*\cos(d*x+c))^{1/2} \end{aligned}$$

3.296.2 Mathematica [A] (verified)

Time = 1.41 (sec) , antiderivative size = 292, normalized size of antiderivative = 0.76

$$\begin{aligned} & \int \cos^3(c+dx)\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))dx \\ & = \frac{8\sqrt{\frac{a+b\cos(c+dx)}{a+b}}(b^2(6a^2Ab+75Ab^3-4a^3B+111ab^2B)\text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)+(24a^3Ab+57aAb^3- \end{aligned}$$

input `Integrate[Cos[c + d*x]^3*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output
$$\begin{aligned} & (8*\text{Sqrt}[(a+b*\text{Cos}[c+d*x])/(a+b)]*(b^2*(6*a^2*A*b+75*A*b^3-4*a^3*B \\ & +111*a*b^2*B)*\text{EllipticF}[(c+d*x)/2,(2*b)/(a+b)]+(24*a^3*A*b+57*a \\ & *A*b^3-16*a^4*B-24*a^2*b^2*B+147*b^4*B)*((a+b)*\text{EllipticE}[(c+d*x) \\ & /2,(2*b)/(a+b)]-a*\text{EllipticF}[(c+d*x)/2,(2*b)/(a+b)]))-b*(a+b* \\ & \text{Cos}[c+d*x])*(-2*(-48*a^2*A*b+345*A*b^3+32*a^3*B+57*a*b^2*B)*\text{Sin}[c \\ & +d*x]-b*((36*a*A*b-24*a^2*B+266*b^2*B)*\text{Sin}[2*(c+d*x)]+5*b*(2*(9 \\ & *A*b+a*B)*\text{Sin}[3*(c+d*x)]+7*b*B*\text{Sin}[4*(c+d*x)])))/((1260*b^4*d*\text{Sqrt} \\ & [a+b*\text{Cos}[c+d*x]])) \end{aligned}$$

3.296.3 Rubi [A] (verified)

Time = 2.18 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.06, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3502, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^3(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c+dx+\frac{\pi}{2}\right)^3 \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} \left(A+B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3469} \\
 & \frac{2 \int \frac{1}{2} \cos(c+dx) \sqrt{a+b \cos(c+dx)} (3(3Ab-2aB) \cos^2(c+dx) + 7bB \cos(c+dx) + 4aB) dx}{\frac{9b}{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}} + \frac{9bd}{9bd}} + \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \cos(c+dx) \sqrt{a+b \cos(c+dx)} (3(3Ab-2aB) \cos^2(c+dx) + 7bB \cos(c+dx) + 4aB) dx}{\frac{9b}{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}} + \frac{9bd}{9bd}} + \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sin\left(c+dx+\frac{\pi}{2}\right) \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} \left(3(3Ab-2aB) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + 7bB \sin\left(c+dx+\frac{\pi}{2}\right) + 4aB\right) dx}{\frac{9b}{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}} + \frac{9bd}{9bd}} + \\
 & \quad \downarrow \text{3528} \\
 & \frac{2 \int \frac{1}{2} \sqrt{a+b \cos(c+dx)} \left(-((-24Ba^2+36Aba-49b^2B) \cos^2(c+dx)) + b(45Ab-2aB) \cos(c+dx) + 6a(3Ab-2aB)\right) dx}{\frac{9b}{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}} + \frac{6(3Ab-2aB) \sin(c+dx) \cos(c+dx)}{7b}} + \frac{6(3Ab-2aB) \sin(c+dx) \cos(c+dx)}{7b} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.296. $\int \cos^3(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\frac{\int \sqrt{a+b \cos(c+dx)} \left(-\left((-24Ba^2+36Aba-49b^2B) \cos^2(c+dx) \right) + b(45Ab-2aB) \cos(c+dx) + 6a(3Ab-2aB) \right) dx}{7b} + \frac{6(3Ab-2aB) \sin(c+dx) \cos(c+dx)}{7bd}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3042

$$\frac{\int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} \left((24Ba^2-36Aba+49b^2B) \sin^2(c+dx+\frac{\pi}{2}) + b(45Ab-2aB) \sin(c+dx+\frac{\pi}{2}) + 6a(3Ab-2aB) \right) dx}{7b} + \frac{6(3Ab-2aB) \sin(c+dx) \cos(c+dx)}{7bd}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3502

$$\frac{2 \int -\frac{3}{2} \sqrt{a+b \cos(c+dx)} \left(b(-4Ba^2+6Aba-49b^2B) - \left(-16Ba^3+24Aba^2-36b^2Ba+75Ab^3 \right) \cos(c+dx) \right) dx}{5b} - \frac{2(-24a^2B+36aAb-49b^2B) \sin(c+dx) (a+b \cos(c+dx))^{3/2}}{5bd}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 27

$$\frac{3 \int \sqrt{a+b \cos(c+dx)} \left(b(-4Ba^2+6Aba-49b^2B) - \left(-16Ba^3+24Aba^2-36b^2Ba+75Ab^3 \right) \cos(c+dx) \right) dx}{5b} - \frac{2(-24a^2B+36aAb-49b^2B) \sin(c+dx) (a+b \cos(c+dx))^{3/2}}{5bd}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3042

$$\frac{3 \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} \left(b(-4Ba^2+6Aba-49b^2B) + \left(16Ba^3-24Aba^2+36b^2Ba-75Ab^3 \right) \sin(c+dx+\frac{\pi}{2}) \right) dx}{5b} - \frac{2(-24a^2B+36aAb-49b^2B) \sin(c+dx) (a+b \cos(c+dx))^{3/2}}{5bd}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3232

$$\frac{\frac{2}{3} \int -\frac{b(-4Ba^3+6Aba^2+111b^2Ba+75Ab^3) + (-16Ba^4+24Aba^3-24b^2Ba^2+57Ab^3a+147b^4B) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-16a^3B+24a^2Ab-36ab^2B+75Ab^3) \sin(c+dx)}{3d}}{5b}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) (a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 27

3.296. $\int \cos^3(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$3 \left(-\frac{1}{3} \int \frac{b(-4Ba^3+6Aba^2+111b^2Ba+75Ab^3)+(-16Ba^4+24Aba^3-24b^2Ba^2+57Ab^3a+147b^4B) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-16a^3B+24a^2Ab-36ab^2B+75Ab^3) \sin(c+dx)}{3d} \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd} \quad 9b$$

↓ 3042

$$3 \left(-\frac{1}{3} \int \frac{b(-4Ba^3+6Aba^2+111b^2Ba+75Ab^3)+(-16Ba^4+24Aba^3-24b^2Ba^2+57Ab^3a+147b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-16a^3B+24a^2Ab-36ab^2B+75Ab^3) \sin(c+dx+\frac{\pi}{2})}{3d} \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd} \quad 9b$$

↓ 3231

$$3 \left(\frac{1}{3} \left(\frac{(a^2-b^2)(-16a^3B+24a^2Ab-36ab^2B+75Ab^3)}{b} \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx - \frac{(-16a^4B+24a^3Ab-24a^2b^2B+57aAb^3+147b^4B)}{b} \int \sqrt{a+b \cos(c+dx)} dx \right) - \frac{2(-16a^3B+24a^2Ab-36ab^2B+75Ab^3) \sin(c+dx)}{3d} \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3042

$$3 \left(\frac{1}{3} \left(\frac{(a^2-b^2)(-16a^3B+24a^2Ab-36ab^2B+75Ab^3)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-16a^4B+24a^3Ab-24a^2b^2B+57aAb^3+147b^4B)}{b} \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx \right) - \frac{2(-16a^3B+24a^2Ab-36ab^2B+75Ab^3) \sin(c+dx+\frac{\pi}{2})}{3d} \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3134

$$3 \left(\frac{1}{3} \left(\frac{(a^2-b^2)(-16a^3B+24a^2Ab-36ab^2B+75Ab^3)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-16a^4B+24a^3Ab-24a^2b^2B+57aAb^3+147b^4B)}{b} \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx \right) - \frac{2(-16a^3B+24a^2Ab-36ab^2B+75Ab^3) \sin(c+dx+\frac{\pi}{2})}{3d} \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

↓ 3042

3.296. $\int \cos^3(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$3 \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-16a^3B + 24a^2Ab - 36ab^2B + 75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{(-16a^4B + 24a^3Ab - 24a^2b^2B + 57aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

$$\downarrow \quad \mathbf{3132}$$

$$3 \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-16a^3B + 24a^2Ab - 36ab^2B + 75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{2(-16a^4B + 24a^3Ab - 24a^2b^2B + 57aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

$$\downarrow \quad \mathbf{3142}$$

$$3 \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-16a^3B + 24a^2Ab - 36ab^2B + 75Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-16a^4B + 24a^3Ab - 24a^2b^2B + 57aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

$$\downarrow \quad \mathbf{3042}$$

$$3 \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-16a^3B + 24a^2Ab - 36ab^2B + 75Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-16a^4B + 24a^3Ab - 24a^2b^2B + 57aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right)$$

5b 7b

$$\frac{2B \sin(c+dx) \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}}{9bd}$$

$$\downarrow \quad \mathbf{3140}$$

3.296. $\int \cos^3(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\frac{2(-24a^2B+36aAb-49b^2B)\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5bd} \frac{3\left(\frac{1}{3}\left(\frac{2(a^2-b^2)(-16a^3B+24a^2Ab-36ab^2B+75Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd\sqrt{a+b\cos(c+dx)}}\right)\right)}{}$$

7b

$$\frac{2B\sin(c+dx)\cos^2(c+dx)(a+b\cos(c+dx))^{3/2}}{9bd}$$

input `Int[Cos[c + d*x]^3*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(2*B*Cos[c + d*x]^2*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(9*b*d) + ((6*(3*A*b - 2*a*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(7*b*d) + ((-2*(36*a*A*b - 24*a^2*B - 49*b^2*B)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*b*d) - (3*(((-2*(24*a^3*A*b + 57*a*A*b^3 - 16*a^4*B - 24*a^2*b^2*B + 147*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(a^2 - b^2)*(24*a^2*A*b + 75*A*b^3 - 16*a^3*B - 36*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]])))/3 - (2*(24*a^2*A*b + 75*A*b^3 - 16*a^3*B - 36*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d)))/(5*b))/(7*b))/(9*b)`

3.296.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b]*Sin[c + d*x]/Sqrt[(a + b)*Sin[c + d*x]]/(a + b) Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`


```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x
])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

3.296.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1634 vs. $2(416) = 832$.

Time = 17.28 (sec) , antiderivative size = 1635, normalized size of antiderivative = 4.24

method	result	size
default	Expression too large to display	1635
parts	Expression too large to display	1824

```
input int(cos(d*x+c)^3*(a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```

-2/315*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*
B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b^5+(720*A*b^5+640*B*a*b^4+2240
*B*b^5)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-432*A*a*b^4-1080*A*b^5+8
*B*a^2*b^3-960*B*a*b^4-2072*B*b^5)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)
+(-12*A*a^2*b^3+432*A*a*b^4+840*A*b^5+8*B*a^3*b^2-8*B*a^2*b^3+728*B*a*b^4+
952*B*b^5)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(24*A*a^3*b^2+6*A*a^2*b
^3-258*A*a*b^4-240*A*b^5-16*B*a^4*b-4*B*a^3*b^2-24*B*a^2*b^3-204*B*a*b^4-1
68*B*b^5)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-24*A*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4*b-51*A*(sin(1/2*d*x+1/2*c)^2)^(1/
2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d
*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b^3+75*A*b^5*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*
x+1/2*c),(-2*b/(a-b))^(1/2))+24*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)
*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*
b/(a-b))^(1/2))*a^4*b-24*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/
2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b)
)^^(1/2))*a^3*b^2+57*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x
+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/
2))*a^2*b^3-57*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1...

```

3.296.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.19 (sec) , antiderivative size = 639, normalized size of antiderivative = 1.66

$$\int \cos^3(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2}(-32i Ba^5 + 48i Aa^4b - 36i Ba^3b^2 + 96i Aa^2b^3 - 39i Bab^4 - 225i Ab^5) \sqrt{b} \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}\right)}{\dots}$$

input `integrate(cos(d*x+c)^3*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output

```

1/945*(sqrt(2)*(-32*I*B*a^5 + 48*I*A*a^4*b - 36*I*B*a^3*b^2 + 96*I*A*a^2*b^3 - 39*I*B*a*b^4 - 225*I*A*b^5)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(32*I*B*a^5 - 48*I*A*a^4*b + 36*I*B*a^3*b^2 - 96*I*A*a^2*b^3 + 39*I*B*a*b^4 + 225*I*A*b^5)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(16*I*B*a^4*b - 24*I*A*a^3*b^2 + 24*I*B*a^2*b^3 - 57*I*A*a*b^4 - 147*I*B*b^5)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(-16*I*B*a^4*b + 24*I*A*a^3*b^2 - 24*I*B*a^2*b^3 + 57*I*A*a*b^4 + 147*I*B*b^5)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(35*B*b^5*cos(d*x + c)^3 + 8*B*a^3*b^2 - 12*A*a^2*b^3 + 13*B*a*b^4 + 75*A*b^5 + 5*(B*a*b^4 + 9*A*b^5)*cos(d*x + c)^2 - (6*B*a^2*b^3 - 9*A*a*b^4 - 49*B*b^5)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^5*d)

```

3.296.6 Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output Timed out

3.296.7 Maxima [F]

$$\begin{aligned} & \int \cos^3(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a \cos(dx + c)}^3 dx \end{aligned}$$

input `integrate(cos(d*x+c)^3*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^3, x)`

3.296.8 Giac [F]

$$\begin{aligned} & \int \cos^3(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c)^3 dx \end{aligned}$$

input `integrate(cos(d*x+c)^3*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^3, x)`

3.296.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cos^3(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int \cos(c + dx)^3 (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} dx \end{aligned}$$

input `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^3*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`

3.297 $\int \cos^2(c+dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.297.1 Optimal result	2718
3.297.2 Mathematica [A] (verified)	2719
3.297.3 Rubi [A] (verified)	2719
3.297.4 Maple [B] (verified)	2725
3.297.5 Fricas [C] (verification not implemented)	2726
3.297.6 Sympy [F]	2726
3.297.7 Maxima [F]	2727
3.297.8 Giac [F]	2727
3.297.9 Mupad [F(-1)]	2727

3.297.1 Optimal result

Integrand size = 33, antiderivative size = 303

$$\int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= - \frac{2(14a^2Ab - 63Ab^3 - 8a^3B - 19ab^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{105b^3d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(a^2 - b^2) (14aAb - 8a^2B - 25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{105b^3d \sqrt{a + b \cos(c + dx)}}$$

$$- \frac{2(14aAb - 8a^2B - 25b^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{105b^2d}$$

$$+ \frac{2(7Ab - 4aB)(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{35b^2d}$$

$$+ \frac{2B \cos(c + dx)(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{7bd}$$

output
$$\frac{2}{35}(7A^2b-4B^2a)(a+b\cos(dx+c))^{3/2}\sin(dx+c)/b^2/d+2/7B\cos(dx+c)(a+b\cos(dx+c))^{3/2}\sin(dx+c)/b/d-2/105(14A^2a^2b-8B^2a^2-25B^2b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^2/d-2/105(14A^2a^2b-63A^2b^3-8B^2a^3-19B^2ab^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticE}(\sin(1/2dx+1/2c), 2^{1/2}(b/(a+b))^{1/2})(a+b\cos(dx+c))^{1/2}/b^3/d/((a+b\cos(dx+c))/(a+b))^{1/2}+2/105(a^2-b^2)(14A^2a^2b-8B^2a^2-25B^2b^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticF}(\sin(1/2dx+1/2c), 2^{1/2}(b/(a+b))^{1/2})((a+b\cos(dx+c))/(a+b))^{1/2}/b^3/d/(a+b\cos(dx+c))^{1/2})$$

3.297.2 Mathematica [A] (verified)

Time = 0.98 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.77

$$\int \cos^2(c+dx)\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))dx$$

$$= \frac{4\sqrt{\frac{a+b\cos(c+dx)}{a+b}}(b^2(49aAb+2a^2B+25b^2B)\text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)+(-14a^2Ab+63Ab^3+8a^3B+19a^2b^2B)(a+b)\text{EllipticE}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)-a\text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right))+b(a+b\cos(c+dx))((28a^2Ab-16a^2B+115b^2B)\sin(c+dx)+3b(2(7A^2b+aB)\sin[2(c+dx)]+5bB\sin[3(c+dx)])))/(210b^3d\sqrt{a+b\cos(c+dx)})}{1}$$

input `Integrate[Cos[c + d*x]^2*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output
$$(4\sqrt{(a+b\cos(c+dx))/(a+b)}(b^2(49a^2Ab+2a^2B+25b^2B)\text{EllipticF}[(c+dx)/2, (2b)/(a+b)]+(-14a^2Ab+63Ab^3+8a^3B+19a^2b^2B)(a+b)\text{EllipticE}[(c+dx)/2, (2b)/(a+b)]-a\text{EllipticF}[(c+dx)/2, (2b)/(a+b)]))+b(a+b\cos(c+dx))((28a^2Ab-16a^2B+115b^2B)\sin(c+dx)+3b(2(7A^2b+aB)\sin[2(c+dx)]+5bB\sin[3(c+dx)])))/(210b^3d\sqrt{a+b\cos(c+dx)})$$

3.297.3 Rubi [A] (verified)

Time = 1.58 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.05, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.297.
$$\int \cos^2(c+dx)\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))dx$$

$$\begin{aligned}
& \int \cos^2(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} \left(A+B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3469} \\
& \frac{2 \int \frac{1}{2} \sqrt{a+b \cos(c+dx)} ((7Ab-4aB) \cos^2(c+dx) + 5bB \cos(c+dx) + 2aB) dx}{7b} + \\
& \quad \frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd} \\
& \quad \downarrow \text{27} \\
& \frac{\int \sqrt{a+b \cos(c+dx)} ((7Ab-4aB) \cos^2(c+dx) + 5bB \cos(c+dx) + 2aB) dx}{7b} + \\
& \quad \frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} \left((7Ab-4aB) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + 5bB \sin\left(c+dx+\frac{\pi}{2}\right) + 2aB\right) dx}{7b} + \\
& \quad \frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd} \\
& \quad \downarrow \text{3502} \\
& \frac{2 \int \frac{1}{2} \sqrt{a+b \cos(c+dx)} (b(21Ab-2aB) - (-8Ba^2+14Aba-25b^2B) \cos(c+dx)) dx}{5b} + \frac{2(7Ab-4aB) \sin(c+dx) (a+b \cos(c+dx))^{3/2}}{5bd} \\
& \quad \frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd} \\
& \quad \downarrow \text{27} \\
& \frac{\int \sqrt{a+b \cos(c+dx)} (b(21Ab-2aB) - (-8Ba^2+14Aba-25b^2B) \cos(c+dx)) dx}{5b} + \frac{2(7Ab-4aB) \sin(c+dx) (a+b \cos(c+dx))^{3/2}}{5bd} \\
& \quad \frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} (b(21Ab-2aB) + (8Ba^2-14Aba+25b^2B) \sin\left(c+dx+\frac{\pi}{2}\right)) dx}{5b} + \frac{2(7Ab-4aB) \sin(c+dx) (a+b \cos(c+dx))^{3/2}}{5bd} \\
& \quad \frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}
\end{aligned}$$

3.297. $\int \cos^2(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

↓ 3232

$$\frac{\frac{2}{3} \int \frac{b(2Ba^2+49Aba+25b^2B) - (-8Ba^3+14Aba^2-19b^2Ba-63Ab^3) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-8a^2B+14aAb-25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}}{5b} + \frac{2(7Ab-4aB) \sin(c+dx)}{7b}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 27

$$\frac{\frac{1}{3} \int \frac{b(2Ba^2+49Aba+25b^2B) - (-8Ba^3+14Aba^2-19b^2Ba-63Ab^3) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-8a^2B+14aAb-25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}}{5b} + \frac{2(7Ab-4aB) \sin(c+dx)}{7b}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3042

$$\frac{\frac{1}{3} \int \frac{b(2Ba^2+49Aba+25b^2B) + (8Ba^3-14Aba^2+19b^2Ba+63Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-8a^2B+14aAb-25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}}{5b} + \frac{2(7Ab-4aB) \sin(c+dx)}{7b}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3231

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B)}{b} \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx - \frac{(-8a^3B+14a^2Ab-19ab^2B-63Ab^3)}{b} \int \sqrt{a+b \cos(c+dx)} dx \right) - \frac{2(-8a^2B+14aAb-25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}}{5b} + \frac{2(7Ab-4aB) \sin(c+dx)}{7b}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3042

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-8a^3B+14a^2Ab-19ab^2B-63Ab^3)}{b} \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx \right) - \frac{2(-8a^2B+14aAb-25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}}{5b} + \frac{2(7Ab-4aB) \sin(c+dx)}{7b}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3134

3.297. $\int \cos^2(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-8a^3B+14a^2Ab-19ab^2B-63Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - 2(-8a^2B+14a^2A-25b^2B)}{5b}}{7b} = \frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3042

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-8a^3B+14a^2Ab-19ab^2B-63Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - 2(-8a^2B+14a^2A-25b^2B)}{5b}}{7b} = \frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3132

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-8a^3B+14a^2Ab-19ab^2B-63Ab^3) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - 2(-8a^2B+14a^2A-25b^2B)}{5b}}{7b} = \frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3142

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - \frac{2(-8a^3B+14a^2Ab-19ab^2B-63Ab^3) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - 2(-8a^2B+14a^2A-25b^2B)}{5b}}{7b} = \frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{3/2}}{7bd}$$

↓ 3042

$$\frac{\frac{1}{3} \left(\frac{(a^2-b^2)(-8a^2B+14aAb-25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx - \frac{2(-8a^3B+14a^2Ab-19ab^2B-63Ab^3) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - 2(-8a^2B+14a^2A-25b^2B)}{5b}}{7b} = \frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{3/2}}{7bd}$$

3.297. $\int \cos^2(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

↓ 3140

$$\frac{\frac{1}{3} \left(\frac{2(a^2 - b^2)(-8a^2B + 14aAb - 25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2(-8a^3B + 14a^2Ab - 19ab^2B - 63Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{a+b \cos(c+dx)}} \right)}{5b} \quad 7b$$

$$\frac{2B \sin(c+dx) \cos(c+dx) (a+b \cos(c+dx))^{3/2}}{7bd}$$

input `Int[Cos[c + d*x]^2*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `(2*B*Cos[c + d*x]*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(7*b*d) + ((2*(7*A*b - 4*a*B)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*b*d) + (((-2*(14*a^2*A*b - 63*A*b^3 - 8*a^3*B - 19*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(a^2 - b^2)*(14*a*A*b - 8*a^2*B - 25*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]])))/3 - (2*(14*a*A*b - 8*a^2*B - 25*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(5*b))/(7*b)`

3.297.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b]*Sin[c + d*x]/Sqrt[(a + b)*Sin[c + d*x]]/(a + b) Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m/(f*(m + 1))))], x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos
[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.297.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1304 vs. $2(337) = 674$.

Time = 15.46 (sec) , antiderivative size = 1305, normalized size of antiderivative = 4.31

method	result	size
default	Expression too large to display	1305
parts	Expression too large to display	1494

```
input int(cos(d*x+c)^2*(a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -2/105*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^4+(-168*A*b^4-144*B*a*b^3-360*B*
b^4)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(112*A*a*b^3+168*A*b^4-4*B*a^
2*b^2+144*B*a*b^3+280*B*b^4)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-14*
A*a^2*b^2-56*A*a*b^3-42*A*b^4+8*B*a^3*b+2*B*a^2*b^2-86*B*a*b^3-80*B*b^4)*s
in(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+14*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1
/2*c),(-2*b/(a-b))^(1/2))*a^3*b-14*a*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/
(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))*b^3-14*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin
(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a
-b))^(1/2))*a^3*b+14*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*
x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1
/2))*a^2*b^2+63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2
*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*
a*b^3-63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(
a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^4-8*B
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b)
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4-17*B*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)...
```

3.297.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 561, normalized size of antiderivative = 1.85

$$\int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2}(16i Ba^4 - 28i Aa^3b + 32i Ba^2b^2 - 21i Aab^3 - 75i Bb^4) \sqrt{b} \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}\right)}{\dots}$$

```
input integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/315*(sqrt(2)*(16*I*B*a^4 - 28*I*A*a^3*b + 32*I*B*a^2*b^2 - 21*I*A*a*b^3 - 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(-16*I*B*a^4 + 28*I*A*a^3*b - 32*I*B*a^2*b^2 + 21*I*A*a*b^3 + 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(-8*I*B*a^3*b + 14*I*A*a^2*b^2 - 19*I*B*a*b^3 - 63*I*A*b^4)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(8*I*B*a^3*b - 14*I*A*a^2*b^2 + 19*I*B*a*b^3 + 63*I*A*b^4)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(15*B*b^4*cos(d*x + c)^2 - 4*B*a^2*b^2 + 7*A*a*b^3 + 25*B*b^4 + 3*(B*a*b^3 + 7*A*b^4)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^4*d)
```

3.297.6 Sympy [F]

$$\int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} \cos^2(c + dx) dx$$

```
input integrate(cos(d*x+c)**2*(a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)
```

3.297. $\int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*cos(c + d*x)**2, x)`

3.297.7 Maxima [F]

$$\begin{aligned} & \int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c)^2 dx \end{aligned}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^2, x)`

3.297.8 Giac [F]

$$\begin{aligned} & \int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c)^2 dx \end{aligned}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^2, x)`

3.297.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cos^2(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int \cos(c + dx)^2 (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} dx \end{aligned}$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`

3.298 $\int \cos(c+dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.298.1 Optimal result	2729
3.298.2 Mathematica [A] (verified)	2730
3.298.3 Rubi [A] (verified)	2730
3.298.4 Maple [B] (verified)	2735
3.298.5 Fricas [C] (verification not implemented)	2736
3.298.6 Sympy [F]	2736
3.298.7 Maxima [F]	2737
3.298.8 Giac [F]	2737
3.298.9 Mupad [F(-1)]	2737

3.298.1 Optimal result

Integrand size = 31, antiderivative size = 231

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{2(5aAb - 2a^2B + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{15b^2d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$- \frac{2(a^2 - b^2) (5Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{15b^2d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{2(5Ab - 2aB) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{15bd} + \frac{2B(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd}$$

output

```
2/5*B*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/b/d+2/15*(5*A*b-2*B*a)*sin(d*x+c)*
(a+b*cos(d*x+c))^(1/2)/b/d+2/15*(5*A*a*b-2*B*a^2+9*B*b^2)*(cos(1/2*d*x+1/2
*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a
+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/
15*(a^2-b^2)*(5*A*b-2*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)
*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(
a+b))^(1/2)/b^2/d/(a+b*cos(d*x+c))^(1/2)
```


3.298.2 Mathematica [A] (verified)

Time = 2.08 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.77

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{2 \sqrt{\frac{a+b \cos(c+dx)}{a+b}} (b^2(5Ab + 7aB) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + (5aAb - 2a^2B + 9b^2B) ((a + b)E\left(\frac{1}{2}(c + dx)\right) - 15b^2d\sqrt{a + b})}{15b^2d\sqrt{a + b}}$$

input `Integrate[Cos[c + d*x]*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`output `(2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*(b^2*(5*A*b + 7*a*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + (5*a*A*b - 2*a^2*B + 9*b^2*B)*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)])) + 2*b*(a + b*Cos[c + d*x])*(5*A*b + a*B + 3*b*B*Cos[c + d*x])*Sin[c + d*x]/(15*b^2*d*Sqrt[a + b*Cos[c + d*x]])`**3.298.3 Rubi [A] (verified)**Time = 1.22 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.03, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int \sqrt{a + b \cos(c + dx)} (A \cos(c + dx) + B \cos^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(A \sin\left(c + dx + \frac{\pi}{2}\right) + B \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$3.298. \quad \int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$\begin{aligned}
 & \downarrow \text{3502} \\
 & \frac{2 \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (3bB + (5Ab - 2aB) \cos(c + dx)) dx}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{27} \\
 & \frac{\int \sqrt{a + b \cos(c + dx)} (3bB + (5Ab - 2aB) \cos(c + dx)) dx}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{3042} \\
 & \frac{\int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (3bB + (5Ab - 2aB) \sin(c + dx + \frac{\pi}{2})) dx}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{3232} \\
 & \frac{\frac{2}{3} \int \frac{b(5Ab + 7aB) + (-2Ba^2 + 5Aba + 9b^2B) \cos(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{2(5Ab - 2aB) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{27} \\
 & \frac{\frac{1}{3} \int \frac{b(5Ab + 7aB) + (-2Ba^2 + 5Aba + 9b^2B) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \frac{2(5Ab - 2aB) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{3042} \\
 & \frac{\frac{1}{3} \int \frac{b(5Ab + 7aB) + (-2Ba^2 + 5Aba + 9b^2B) \sin(c + dx + \frac{\pi}{2})}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2(5Ab - 2aB) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{3231} \\
 & \frac{\frac{1}{3} \left(\frac{(-2a^2B + 5aAb + 9b^2B) \int \sqrt{a + b \cos(c + dx)} dx}{b} - \frac{(a^2 - b^2)(5Ab - 2aB) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) + \frac{2(5Ab - 2aB) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}}{5b} + \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5bd} \\
 & \downarrow \text{3042}
 \end{aligned}$$

3.298. $\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

$$\frac{1}{3} \left(\frac{(-2a^2B+5aAb+9b^2B) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{(a^2-b^2)(5Ab-2aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \frac{2(5Ab-2aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5bd}$$

↓ 3134

$$\frac{1}{3} \left(\frac{(-2a^2B+5aAb+9b^2B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(5Ab-2aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \frac{2(5Ab-2aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5bd}$$

↓ 3042

$$\frac{1}{3} \left(\frac{(-2a^2B+5aAb+9b^2B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(5Ab-2aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \frac{2(5Ab-2aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5bd}$$

↓ 3132

$$\frac{1}{3} \left(\frac{2(-2a^2B+5aAb+9b^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(5Ab-2aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \frac{2(5Ab-2aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5bd}$$

↓ 3142

$$\frac{1}{3} \left(\frac{2(-2a^2B+5aAb+9b^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(5Ab-2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} \right) + \frac{2(5Ab-2aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5bd}$$

↓ 3042

3.298. $\int \cos(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\frac{1}{3} \left(\frac{2(-2a^2B+5aAb+9b^2B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(5Ab-2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin\left(c+dx+\frac{\pi}{2}\right)}{a+b}}} dx}{b\sqrt{a+b\cos(c+dx)}} \right) + \frac{2(5Ab-2a^2B)\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5bd}$$

3140

$$\frac{1}{3} \left(\frac{2(-2a^2B+5aAb+9b^2B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{2(a^2-b^2)(5Ab-2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd\sqrt{a+b\cos(c+dx)}} \right) + \frac{2(5Ab-2a^2B)\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5bd}$$

input `Int[Cos[c + d*x]*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]), x]`

output `(2*B*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*b*d) + (((2*(5*a*A*b - 2*a^2*B + 9*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(5*A*b - 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 + (2*(5*A*b - 2*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(5*b)`

3.298.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos
[e + f*x]*((a + b*Ssin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Ssin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.298.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 992 vs. $2(269) = 538$.

Time = 13.50 (sec) , antiderivative size = 993, normalized size of antiderivative = 4.30

method	result	size
default	Expression too large to display	993
parts	Expression too large to display	1119

```
input int(cos(d*x+c)*(a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output -2/15*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*c
os(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^3+(20*A*b^3+16*B*a*b^2+24*B*b^3)*
sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*a*b^2-10*A*b^3-2*B*a^2*b-8*
B*a*b^2-6*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-5*A*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*Ellipt
icF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+5*A*b^3*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(
cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/
2*c),(-2*b/(a-b))^(1/2))*a^2*b-5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)
)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2
*b/(a-b))^(1/2))*a*b^2+2*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/
2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b)
)^(1/2))*a^3-2*B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/
2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))
*b^2-2*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+
b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3+2*B*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+9*B*(sin(1/2*
d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)...
```

3.298.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.15 (sec) , antiderivative size = 492, normalized size of antiderivative = 2.13

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \frac{\sqrt{2}(-4i Ba^3 + 10i Aa^2b - 3i Bab^2 - 15i Ab^3) \sqrt{b} \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c)}{b^3}\right)}{b^3}$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm=
"fricas")
```

```
output 1/45*(sqrt(2)*(-4*I*B*a^3 + 10*I*A*a^2*b - 3*I*B*a*b^2 - 15*I*A*b^3)*sqrt(
b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^
3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(4*I*B*a
^3 - 10*I*A*a^2*b + 3*I*B*a*b^2 + 15*I*A*b^3)*sqrt(b)*weierstrassPInverse(
4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c
) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(2*I*B*a^2*b - 5*I*A*a*b^2 -
9*I*B*b^3)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 -
9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 -
9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*s
qrt(2)*(-2*I*B*a^2*b + 5*I*A*a*b^2 + 9*I*B*b^3)*sqrt(b)*weierstrassZeta(4/
3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/
3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c)
- 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(3*B*b^3*cos(d*x + c) + B*a*b^2 + 5*A*
b^3)*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^3*d)
```

3.298.6 Sympy [F]

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} \cos(c + dx) dx$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)
```

```
output Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*cos(c + d*x), x)
```

3.298. $\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.298.7 Maxima [F]

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c), x)`

3.298.8 Giac [F]

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c), x)`

3.298.9 Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \cos(c + dx) (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} dx$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`

3.299 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$

3.299.1 Optimal result	2738
3.299.2 Mathematica [A] (verified)	2739
3.299.3 Rubi [A] (verified)	2739
3.299.4 Maple [B] (verified)	2743
3.299.5 Fricas [C] (verification not implemented)	2744
3.299.6 Sympy [F]	2744
3.299.7 Maxima [F]	2745
3.299.8 Giac [F]	2745
3.299.9 Mupad [F(-1)]	2745

3.299.1 Optimal result

Integrand size = 25, antiderivative size = 171

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx \\ &= \frac{2(3Ab + aB)\sqrt{a + b \cos(c + dx)}E\left(\frac{1}{2}(c + dx)\middle|\frac{2b}{a+b}\right)}{3bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \\ & \quad - \frac{2(a^2 - b^2)B\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3bd\sqrt{a + b \cos(c + dx)}} \\ & \quad + \frac{2B\sqrt{a + b \cos(c + dx)}\sin(c + dx)}{3d} \end{aligned}$$

output $2/3*B*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/d+2/3*(3*A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*\operatorname{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*\cos(d*x+c))^(1/2)/b/d/((a+b*\cos(d*x+c))/(a+b))^(1/2)-2/3*(a^2-b^2)*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*\cos(d*x+c))/(a+b))^(1/2)/b/d/(a+b*\cos(d*x+c))^(1/2)$

3.299.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.85

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{2(a + b)(3Ab + aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) - 2(a^2 - b^2) B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3bd \sqrt{a + b \cos(c + dx)}}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`output `(2*(a + b)*(3*A*b + a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - 2*(a^2 - b^2)*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + 2*b*B*(a + b*Cos[c + d*x])*Sin[(c + d*x)]/(3*b*d*Sqrt[a + b*Cos[c + d*x]])`**3.299.3 Rubi [A] (verified)**Time = 0.83 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.480$, Rules used = {3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3232}$$

$$\frac{2}{3} \int \frac{3aA + bB + (3Ab + aB) \cos(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

$$\downarrow \text{27}$$

$$\frac{1}{3} \int \frac{3aA + bB + (3Ab + aB) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3aA + bB + (3Ab + aB) \sin(c + dx + \frac{\pi}{2})}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \downarrow \text{3231} \\
& \frac{1}{3} \left(\frac{(aB + 3Ab) \int \sqrt{a + b \cos(c + dx)} dx}{b} - \frac{B(a^2 - b^2) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) + \\
& \quad \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \downarrow \text{3042} \\
& \frac{1}{3} \left(\frac{(aB + 3Ab) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx}{b} - \frac{B(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \right) + \\
& \quad \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \downarrow \text{3134} \\
& \frac{1}{3} \left(\frac{(aB + 3Ab) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{B(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \right) + \\
& \quad \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \downarrow \text{3042} \\
& \frac{1}{3} \left(\frac{(aB + 3Ab) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{B(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \right) + \\
& \quad \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \downarrow \text{3132} \\
& \frac{1}{3} \left(\frac{2(aB + 3Ab) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{B(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \right) + \\
& \quad \frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \downarrow \text{3142}
\end{aligned}$$

$$\frac{1}{3} \left(\frac{2(aB + 3Ab)\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{B(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b\sqrt{a + b \cos(c + dx)}} \right) +$$

$$\frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{2(aB + 3Ab)\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{B(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{b\sqrt{a + b \cos(c + dx)}} \right) +$$

$$\frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3140

$$\frac{1}{3} \left(\frac{2(aB + 3Ab)\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2B(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{bd\sqrt{a + b \cos(c + dx)}} \right) +$$

$$\frac{2B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

input `Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output `((2*(3*A*b + a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 + (2*B*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d)`

3.299.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

```
rule 3232 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]
```

3.299.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 599 vs. 2(213) = 426.

Time = 10.72 (sec) , antiderivative size = 600, normalized size of antiderivative = 3.51

method	result
default	$2\sqrt{\left(2b\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) + a - b\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(4B\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^2 + 3A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)$
parts	$\frac{2A\sqrt{\left(2b\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right), \sqrt{-\frac{2b}{a - b}}(a - b)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (a + b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a + bd}}$

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output -2/3*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*B*cos(1/2*d*x+1/2*c)^5*b^2+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b-3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^2+2*B*cos(1/2*d*x+1/2*c)^3*a*b-6*B*cos(1/2*d*x+1/2*c)^3*b^2-B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2+B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2-B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b-2*B*cos(1/2*d*x+1/2*c)*a*b+2*B*cos(1/2*d*x+1/2*c)*b^2)/b/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d
```

3.299. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$

3.299.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.54

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$= \frac{6 \sqrt{b \cos(dx + c) + a} B b^2 \sin(dx + c) + \sqrt{2}(2i B a^2 - 3i A a b - 3i B b^2) \sqrt{b} \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}\right)}{b^2}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output `1/9*(6*sqrt(b*cos(d*x + c) + a)*B*b^2*sin(d*x + c) + sqrt(2)*(2*I*B*a^2 - 3*I*A*a*b - 3*I*B*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(-2*I*B*a^2 + 3*I*A*a*b + 3*I*B*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(-I*B*a*b - 3*I*A*b^2)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(I*B*a*b + 3*I*A*b^2)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)))/(b^2*d)`

3.299.6 Sympy [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x)), x)`

3.299.7 Maxima [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a), x)`

3.299.8 Giac [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a), x)`

3.299.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2),x)`

output `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`

3.300 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$

3.300.1 Optimal result	2746
3.300.2 Mathematica [A] (verified)	2747
3.300.3 Rubi [A] (verified)	2747
3.300.4 Maple [A] (verified)	2752
3.300.5 Fricas [F]	2752
3.300.6 Sympy [F]	2753
3.300.7 Maxima [F]	2753
3.300.8 Giac [F]	2753
3.300.9 Mupad [F(-1)]	2754

3.300.1 Optimal result

Integrand size = 31, antiderivative size = 178

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2B \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2Ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{2aA \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

output

```
2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+
1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))
/(a+b))^(1/2)+2*A*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellipt
icF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(
1/2)/d/(a+b*cos(d*x+c))^(1/2)+2*a*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d
*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*c
os(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)
```

3.300.2 Mathematica [A] (verified)

Time = 26.06 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.60

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \frac{2\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \left((a+b)BE\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) + A(b \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + a \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx)\right), \frac{2b}{a+b}) \right)}{d\sqrt{a + b \cos(c + dx)}}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*((a + b)*B*EllipticE[(c + d*x)/2, (2*b)/(a + b)] + A*(b*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + a*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]))/(d*Sqrt[a + b*Cos[c + d*x]])`

3.300.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3481, 3042, 3134, 3042, 3132, 3282, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(c + dx) \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{3481}$$

$$A \int \sqrt{a + b \cos(c + dx)} \sec(c + dx) dx + B \int \sqrt{a + b \cos(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$A \int \frac{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}}{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + B \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{3134}$$

$$\begin{aligned}
& A \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{B \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3042} \\
& A \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{B \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3132} \\
& A \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}}{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2B \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3282} \\
& A \left(b \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx + a \int \frac{\sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx \right) + \\
& \quad \frac{2B \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3042} \\
& A \left(b \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + a \int \frac{1}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2B \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3142} \\
& A \left(a \int \frac{1}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \right) + \\
& \quad \frac{2B \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\begin{aligned}
& A \left(a \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} \right) + \\
& \quad \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3140} \\
& A \left(a \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) + \\
& \quad \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3286} \\
& A \left(\frac{a \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) + \\
& \quad \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3042} \\
& A \left(\frac{a \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) + \\
& \quad \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \quad \downarrow \text{3284}
\end{aligned}$$

$$A \left(\frac{2b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} + \frac{2a \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) + \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

input `Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(2*B*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + A*((2*b*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*a*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))`

3.300.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])]/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3282 `Int[Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]/((a_) + (b_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d/b Int[1/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[(b*c - a*d)/b Int[1/((a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3481 `Int[(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^m)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.300.4 Maple [A] (verified)

Time = 7.42 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.39

method	result
default	$-\frac{2\sqrt{\left(2b\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}{a-b}}\left(AbF\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)-aA\Pi\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+(a+b)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{-2b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+bd}}$
parts	$-\frac{2A\sqrt{\left(2b\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}{a-b}}\left(F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)b-\Pi\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+(a+b)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{-2b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+bd}}$

input `int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `-2*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*(A*B*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-a*A*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))+B*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)))*a-B*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d`

3.300.5 Fracas [F]

$$\int \sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec(c+dx)dx$$

$$= \int (B\cos(dx+c)+A)\sqrt{b\cos(dx+c)+a}\sec(dx+c)dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c), x)`

3.300.6 Sympy [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} \sec(c + dx) dx$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c), x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*sec(c + d*x), x)`

3.300.7 Maxima [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c), x)`

3.300.8 Giac [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c), x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c), x)`

3.300.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x),x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x), x)`

3.301 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

3.301.1 Optimal result	2755
3.301.2 Mathematica [C] (verified)	2756
3.301.3 Rubi [A] (verified)	2756
3.301.4 Maple [B] (verified)	2762
3.301.5 Fricas [F(-1)]	2763
3.301.6 Sympy [F]	2764
3.301.7 Maxima [F]	2764
3.301.8 Giac [F]	2764
3.301.9 Mupad [F(-1)]	2765

3.301.1 Optimal result

Integrand size = 33, antiderivative size = 213

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= -\frac{A\sqrt{a + b \cos(c + dx)}E\left(\frac{1}{2}(c + dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\ &+ \frac{(aA + 2bB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d\sqrt{a + b \cos(c + dx)}} \\ &+ \frac{(Ab + 2aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d\sqrt{a + b \cos(c + dx)}} \\ &+ \frac{A\sqrt{a + b \cos(c + dx)} \tan(c + dx)}{d} \end{aligned}$$

output

```
-A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+(A*a+2*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)+(A*b+2*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2, 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)+A*(a+b*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.301.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 15.82 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.75

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \frac{8bB \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2(Ab+4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} - \frac{2iA \sqrt{-\frac{b(-1+\cos(c+dx))}{a+b}} \sqrt{\frac{b(1+\cos(c+dx))}{a+b}}}{\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `((8*b*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(A*b + 4*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] - ((2*I)*A*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)])))/(a*b*Sqrt[-(a + b)^(-1)] + 4*A*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/(4*d)`

3.301.3 Rubi [A] (verified)

Time = 1.78 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3478, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(c + dx) \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx$$

3.301. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

$$\begin{aligned}
& \int \frac{(-Ab \cos^2(c+dx) + 2bB \cos(c+dx) + Ab + 2aB) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3478} \\
& \frac{1}{2} \int \frac{(-Ab \cos^2(c+dx) + 2bB \cos(c+dx) + Ab + 2aB) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \int \frac{-Ab \sin(c+dx+\frac{\pi}{2})^2 + 2bB \sin(c+dx+\frac{\pi}{2}) + Ab + 2aB}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\frac{\int \frac{(b(Ab+2aB)+b(aA+2bB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} - A \int \sqrt{a+b \cos(c+dx)} dx \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3538} \\
& \frac{1}{2} \left(\frac{\int \frac{(b(Ab+2aB)+b(aA+2bB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} - A \int \sqrt{a+b \cos(c+dx)} dx \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left(\frac{\int \frac{(b(Ab+2aB)+b(aA+2bB) \sin(c+dx+\frac{\pi}{2})) \sec(c+dx)}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - A \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\frac{\int \frac{(b(Ab+2aB)+b(aA+2bB) \sin(c+dx+\frac{\pi}{2})) \sec(c+dx)}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - A \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3134}
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{2} \left(\frac{\int \frac{b(Ab+2aB)+b(aA+2bB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{A\sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\frac{\int \frac{b(Ab+2aB)+b(aA+2bB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{A\sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3132} \\
& \frac{1}{2} \left(\frac{\int \frac{b(Ab+2aB)+b(aA+2bB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2A\sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3481} \\
& \frac{1}{2} \left(\frac{b(aA+2bB) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx + b(2aB+Ab) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} - \frac{2A\sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\frac{b(aA+2bB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + b(2aB+Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2A\sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \\
& \quad \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} \\
& \quad \downarrow \text{3142}
\end{aligned}$$

3.301. $\int \sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx)) \sec^2(c+dx) dx$

$$\frac{1}{2} \left(\frac{b(2aB + Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(aA+2bB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{b} - \frac{2A\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \right) - \frac{A \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{b(2aB + Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(aA+2bB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{b} - \frac{2A\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \right) - \frac{A \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3140

$$\frac{1}{2} \left(\frac{b(2aB + Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(aA+2bB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2A\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \right) - \frac{A \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3286

$$\frac{1}{2} \left(\frac{b(2aB+Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2b(aA+2bB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2A\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \right) - \frac{A \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{b(2aB+Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{2b(aA+2bB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} - \frac{2A\sqrt{a+b\cos(c+dx)}}{d} \right) - \frac{A \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3284

$$\frac{1}{2} \left(\frac{2b(aA+2bB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} + \frac{2b(2aB+Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} - \frac{2A\sqrt{a+b\cos(c+dx)}}{d} \right) - \frac{A \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

```
input Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]
```

```
output ((-2*A*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*b*(a*A + 2*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*b*(A*b + 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/2 + (A*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/d
```

3.301.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.301. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`


```
rule 3478 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*
x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a
+ b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(
m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*
Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

```
rule 3481 Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.301.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 745 vs. $2(286) = 572$.

Time = 10.24 (sec) , antiderivative size = 746, normalized size of antiderivative = 3.50

method	result	size
default	Expression too large to display	746
parts	Expression too large to display	818

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*
sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2
*d*x+1/2*c),(-2*b/(a-b))^(1/2))+2*a*A*(-cos(1/2*d*x+1/2*c)/a*(-2*sin(1/2*d
*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)
+1/2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(
1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*Elliptic
F(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-1/2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a
+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))+1/2/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/
(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*
b*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+1/2/a*b*(sin(1/2*d*x+1/2*c
)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x
+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c
),2,(-2*b/(a-b))^(1/2)))-2*(A*b+B*a)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*co
s(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(
1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2)
))/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d

```

3.301.5 Fracas [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Timed out}$$

input

```

integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorith
m="fracas")

```

output

```

Timed out

```

3.301.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} \sec^2(c + dx) dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*sec(c + d*x)**2, x)`

3.301.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^2 dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^2, x)`

3.301.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^2(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^2 dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^2, x)`

3.301.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^2(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^2} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^2,x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^2, x)`

3.302 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$

3.302.1 Optimal result	2766
3.302.2 Mathematica [C] (verified)	2767
3.302.3 Rubi [A] (verified)	2768
3.302.4 Maple [B] (verified)	2775
3.302.5 Fricas [F(-1)]	2776
3.302.6 Sympy [F]	2777
3.302.7 Maxima [F]	2777
3.302.8 Giac [F]	2777
3.302.9 Mupad [F(-1)]	2778

3.302.1 Optimal result

Integrand size = 33, antiderivative size = 292

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= -\frac{(Ab + 4aB)\sqrt{a + b \cos(c + dx)}E\left(\frac{1}{2}(c + dx)\middle|\frac{2b}{a+b}\right)}{4ad\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\ &+ \frac{(3Ab + 4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4d\sqrt{a + b \cos(c + dx)}} \\ &+ \frac{(4a^2A - Ab^2 + 4abB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4ad\sqrt{a + b \cos(c + dx)}} \\ &+ \frac{(Ab + 4aB)\sqrt{a + b \cos(c + dx)} \tan(c + dx)}{4ad} \\ &+ \frac{A\sqrt{a + b \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{2d} \end{aligned}$$

output
$$-1/4*(A*b+4*B*a)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*(a+b*cos(d*x+c))^{(1/2)}/a/d/((a+b*cos(d*x+c))/(a+b))^{(1/2)}+1/4*(3*A*b+4*B*a)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*cos(d*x+c))^{(1/2)}+1/4*(4*A*a^2-A*b^2+4*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/a/d/(a+b*cos(d*x+c))^{(1/2)}+1/4*(A*b+4*B*a)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/a/d+1/2*A*sec(d*x+c)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d$$

3.302.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.02 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.44

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \frac{8Ab\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(8a^2A-3Ab^2+4abB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{a\sqrt{a+b\cos(c+dx)}} - \frac{2i(Ab+4aB)\sqrt{-b(-1}}{\dots}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output
$$\left(\frac{8A*b*\sqrt{(a+b*\cos[c+d*x])/(a+b)}}{\sqrt{a+b*\cos[c+d*x]}}*\operatorname{EllipticF}\left[\frac{(c+d*x)}{2}, \frac{(2*b)/(a+b)}{2}\right] + \frac{(2*(8*a^2*A-3*A*b^2+4*a*b*B))*\sqrt{(a+b*\cos[c+d*x])/(a+b)}}{\sqrt{a+b*\cos[c+d*x]}}*\operatorname{EllipticPi}\left[2, \frac{(c+d*x)}{2}, \frac{(2*b)/(a+b)}{2}\right] - \frac{(2*I)*(A*b+4*a*B)*\sqrt{-(b*(-1+\cos[c+d*x])/(a+b))}}{\sqrt{a+b*\cos[c+d*x]}}*\operatorname{Csc}[c+d*x]*(-2*a*(a-b)*\operatorname{EllipticE}[I*\operatorname{ArcSinh}[\sqrt{-(a+b)^{-1}}]*\sqrt{a+b*\cos[c+d*x]}], (a+b)/(a-b)] + b*(-2*a*\operatorname{EllipticF}[I*\operatorname{ArcSinh}[\sqrt{-(a+b)^{-1}}]*\sqrt{a+b*\cos[c+d*x]}], (a+b)/(a-b)] + b*\operatorname{EllipticPi}[(a+b)/a, I*\operatorname{ArcSinh}[\sqrt{-(a+b)^{-1}}]*\sqrt{a+b*\cos[c+d*x]}], (a+b)/(a-b)]\right) / (a^2*b*\sqrt{-(a+b)^{-1}}) + (4*\sqrt{a+b*\cos[c+d*x]}*(2*a*A+(A*b+4*a*B)*\cos[c+d*x])*Sec[c+d*x]*Tan[c+d*x])/a/(16*d)$$

3.302.3 Rubi [A] (verified)

Time = 2.50 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.02, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3478, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^3(c+dx)\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^3}dx \\
 & \quad \downarrow \text{3478} \\
 & \frac{1}{2} \int \frac{(Ab\cos^2(c+dx)+2(aA+2bB)\cos(c+dx)+Ab+4aB)\sec^2(c+dx)}{2\sqrt{a+b\cos(c+dx)}}dx + \\
 & \quad \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4} \int \frac{(Ab\cos^2(c+dx)+2(aA+2bB)\cos(c+dx)+Ab+4aB)\sec^2(c+dx)}{\sqrt{a+b\cos(c+dx)}}dx + \\
 & \quad \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{4} \int \frac{Ab\sin(c+dx+\frac{\pi}{2})^2+2(aA+2bB)\sin(c+dx+\frac{\pi}{2})+Ab+4aB}{\sin(c+dx+\frac{\pi}{2})^2\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx + \\
 & \quad \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d} \\
 & \quad \downarrow \text{3534} \\
 & \frac{1}{4} \left(\frac{\int \frac{(4Aa^2+4bBa+2Ab\cos(c+dx)a-Ab^2-b(Ab+4aB)\cos^2(c+dx))\sec(c+dx)}{2\sqrt{a+b\cos(c+dx)}}dx}{a} + \frac{(4aB+Ab)\tan(c+dx)\sqrt{a+b\cos(c+dx)}}{ad} \right. \\
 & \quad \left. \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d} \right)
 \end{aligned}$$

3.302. $\int \sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^3(c+dx)dx$

↓ 27

$$\frac{1}{4} \left(\frac{\int \frac{(4Aa^2 + 4bBa + 2Ab \cos(c+dx)a - Ab^2 - b(Ab + 4aB) \cos^2(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} + \frac{(4aB + Ab) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} \right)$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{\int \frac{4Aa^2 + 4bBa + 2Ab \sin(c+dx + \frac{\pi}{2})a - Ab^2 - b(Ab + 4aB) \sin(c+dx + \frac{\pi}{2})^2}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{2a} + \frac{(4aB + Ab) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} \right) +$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3538

$$\frac{1}{4} \left(\frac{\int -\frac{(b(4Aa^2 + 4bBa - Ab^2) + ab(3Ab + 4aB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - \left((4aB + Ab) \int \sqrt{a+b \cos(c+dx)} dx \right) \right) + \frac{(4aB + Ab) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 25

$$\frac{1}{4} \left(\frac{\int \frac{(b(4Aa^2 + 4bBa - Ab^2) + ab(3Ab + 4aB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - (4aB + Ab) \int \sqrt{a+b \cos(c+dx)} dx \right) + \frac{(4aB + Ab) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{\int \frac{b(4Aa^2 + 4bBa - Ab^2) + ab(3Ab + 4aB) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{2a} - (4aB + Ab) \int \sqrt{a+b \sin(c+dx + \frac{\pi}{2})} dx \right) + \frac{(4aB + Ab) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

$$\begin{aligned} & \downarrow \text{3134} \\ & \frac{1}{4} \left(\frac{\int \frac{b(4Aa^2+4bBa-Ab^2)+ab(3Ab+4aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{(4aB+Ab)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} + \frac{(4aB+Ab)\tan(c+dx)}{a} \right) \\ & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ & \frac{1}{4} \left(\frac{\int \frac{b(4Aa^2+4bBa-Ab^2)+ab(3Ab+4aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{(4aB+Ab)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} + \frac{(4aB+Ab)\tan(c+dx)}{a} \right) \\ & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3132} \\ & \frac{1}{4} \left(\frac{\int \frac{b(4Aa^2+4bBa-Ab^2)+ab(3Ab+4aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} + \frac{(4aB+Ab)\tan(c+dx)}{ad} \right) \\ & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3481} \\ & \frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2) \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx + ab(4aB+3Ab) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx}{2a} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} + \frac{(4aB+Ab)\tan(c+dx)}{a} \right) \\ & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d} \\ & \downarrow \text{3042} \end{aligned}$$

$$\frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + ab(4aB+3Ab) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)} E\left(\frac{1}{2}(c+\frac{a+b\cos(c+dx)}{a+b})\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3142

$$\frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{ab(4aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{2a} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{ab(4aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{2a} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3140

$$\frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2ab(4aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2a}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3286

$$\frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2ab(4aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2a}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{b(4a^2A+4abB-Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2ab(4aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2a}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3284

$$\frac{1}{4} \left(\frac{2b(4a^2A+4abB-Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{2ab(4aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(4aB+Ab)\sqrt{a+b\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2a}$$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

input `Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output `(A*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (((-2*(A*b + 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*a*b*(3*A*b + 4*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*b*(4*a^2*A - A*b^2 + 4*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/(2*a) + ((A*b + 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/(a*d))/4`

3.302.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3478 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

rule 3481 `Int((((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^n), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.302.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1289 vs. $2(353) = 706$.

Time = 12.90 (sec) , antiderivative size = 1290, normalized size of antiderivative = 4.42

method	result	size
default	Expression too large to display	1290
parts	Expression too large to display	1601

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNV
ERBOSE)
```

output
$$\begin{aligned} & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*a*A*(-1/ \\ & 2*\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c) \\ & ^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)^2+3/4*b/a^2*\cos(1/2*d*x+1/2*c)*(-2*\sin \\ & (1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2* \\ & c)^2-1)-1/8*b/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a- \\ & b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/ \\ & 2)}*EllipticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+3/8/a*(\sin(1/2*d*x+1/2* \\ & *c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+ \\ & 1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*EllipticE(\cos(1/2*d*x+1/2*c) \\ &),(-2*b/(a-b))^{(1/2)})-3/8*b^2/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1 \\ & /2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2 \\ & *d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})-1/2* \\ & (\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/ \\ & (-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos \\ & (1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-3/8/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ &)*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+ \\ & (a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a \\ & -b))^{(1/2)})*b^2)-2*B*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2* \\ & c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c) \\ & ^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})+2*(A*b+B*...$$

3.302.5 Fracas [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fricas")`

output `Timed out`

3.302.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} \sec^3(c + dx) dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*sec(c + d*x)**3, x)`

3.302.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^3 dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^3, x)`

3.302.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^3(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^3 dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^3, x)`

3.302.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^3(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^3} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^3,x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^3, x)`

3.303 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$

3.303.1 Optimal result	2779
3.303.2 Mathematica [C] (verified)	2780
3.303.3 Rubi [A] (verified)	2781
3.303.4 Maple [B] (verified)	2789
3.303.5 Fricas [F(-1)]	2790
3.303.6 Sympy [F(-1)]	2791
3.303.7 Maxima [F]	2791
3.303.8 Giac [F]	2791
3.303.9 Mupad [F(-1)]	2792

3.303.1 Optimal result

Integrand size = 33, antiderivative size = 378

$$\begin{aligned}
 & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx \\
 &= -\frac{(16a^2A - 3Ab^2 + 6abB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{24a^2d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 &+ \frac{(16a^2A - Ab^2 + 18abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{24ad \sqrt{a + b \cos(c + dx)}} \\
 &+ \frac{(4a^2Ab + Ab^3 + 8a^3B - 2ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{8a^2d \sqrt{a + b \cos(c + dx)}} \\
 &+ \frac{(16a^2A - 3Ab^2 + 6abB) \sqrt{a + b \cos(c + dx)} \tan(c + dx)}{24a^2d} \\
 &+ \frac{(Ab + 6aB) \sqrt{a + b \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{12ad} \\
 &+ \frac{A \sqrt{a + b \cos(c + dx)} \sec^2(c + dx) \tan(c + dx)}{3d}
 \end{aligned}$$

output
$$\begin{aligned} & -1/24*(16*A*a^2-3*A*b^2+6*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+ \\ & 1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*(a+b*cos(d*x+ \\ & c))^{(1/2)}/a^2/d/((a+b*cos(d*x+c))/(a+b))^{(1/2)}+1/24*(16*A*a^2-A*b^2+18*B*a \\ & *b)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+ \\ & 1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/a/d/(a+b*co \\ & s(d*x+c))^{(1/2)}+1/8*(4*A*a^2*b+A*b^3+8*B*a^3-2*B*a*b^2)*(cos(1/2*d*x+1/2*c \\ &)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^{(1/2)}*(b/(\\ & a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/a^2/d/(a+b*cos(d*x+c))^{(1/2)}+1 \\ & /24*(16*A*a^2-3*A*b^2+6*B*a*b)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/a^2/d+1/1 \\ & 2*(A*b+6*B*a)*sec(d*x+c)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/a/d+1/3*A*sec(d \\ & *x+c)^2*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d \end{aligned}$$

3.303.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.12 (sec) , antiderivative size = 480, normalized size of antiderivative = 1.27

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \frac{8ab(Ab+6aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(8a^2Ab+9Ab^3+48a^3B-18ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} +$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output
$$\begin{aligned} & ((8*a*b*(A*b + 6*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d* \\ & x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(8*a^2*A*b + 9*A*b^3 + \\ & 48*a^3*B - 18*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (\\ & c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + ((2*I)*(-16*a^2*A + \\ & 3*A*b^2 - 6*a*b*B)*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[-((b*(1 \\ & + Cos[c + d*x]))/(a - b))]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[\\ & Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a* \\ & EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b) \\ & / (a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + \\ & b*Cos[c + d*x]]], (a + b)/(a - b))))/(a*b*Sqrt[-(a + b)^(-1)]) + 4*Sqrt[\\ & a + b*Cos[c + d*x]]*Sec[c + d*x]^2*(2*a*(A*b + 6*a*B)*Sin[c + d*x] + (8*a^ \\ & 2*A - (3*A*b^2)/2 + 3*a*b*B)*Sin[2*(c + d*x)] + 8*a^2*A*Tan[c + d*x]))/(96 \\ & *a^2*d) \end{aligned}$$

3.303.3 Rubi [A] (verified)

Time = 3.25 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.03, number of steps used = 24, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 3478, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^4(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (A+B \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^4} dx \\
 & \quad \downarrow \text{3478} \\
 & \frac{1}{3} \int \frac{(3Ab \cos^2(c+dx) + 2(2aA + 3bB) \cos(c+dx) + Ab + 6aB) \sec^3(c+dx)}{2\sqrt{a+b \cos(c+dx)} \frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}} dx + \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{6} \int \frac{(3Ab \cos^2(c+dx) + 2(2aA + 3bB) \cos(c+dx) + Ab + 6aB) \sec^3(c+dx)}{\sqrt{a+b \cos(c+dx)} \frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}} dx + \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \int \frac{3Ab \sin(c+dx+\frac{\pi}{2})^2 + 2(2aA + 3bB) \sin(c+dx+\frac{\pi}{2}) + Ab + 6aB}{\sin(c+dx+\frac{\pi}{2})^3 \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} \frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}} dx + \\
 & \quad \downarrow \text{3534} \\
 & \frac{1}{6} \left(\int \frac{(16Aa^2+6bBa+2(7Ab+6aB) \cos(c+dx)a-3Ab^2+b(Ab+6aB) \cos^2(c+dx)) \sec^2(c+dx)}{2\sqrt{a+b \cos(c+dx)} 2a} dx + \frac{(6aB + Ab) \tan(c+dx) \sec(c+dx)}{2ad} \right. \\
 & \quad \left. \frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{1}{6} \left(\frac{\int \frac{(16Aa^2+6bBa+2(7Ab+6aB)\cos(c+dx)a-3Ab^2+b(Ab+6aB)\cos^2(c+dx))\sec^2(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{4a} + \frac{(6aB+Ab)\tan(c+dx)\sec(c+dx)}{2ad} \right) \\
 & \qquad \qquad \qquad \frac{A \tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{1}{6} \left(\frac{\int \frac{16Aa^2+6bBa+2(7Ab+6aB)\sin(c+dx+\frac{\pi}{2})a-3Ab^2+b(Ab+6aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^2\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{4a} + \frac{(6aB+Ab)\tan(c+dx)\sec(c+dx)\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}{2ad} \right) \\
 & \qquad \qquad \qquad \frac{A \tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d} \\
 & \qquad \qquad \qquad \downarrow 3534 \\
 & \frac{1}{6} \left(\frac{\int \frac{(-b(16Aa^2+6bBa-3Ab^2)\cos^2(c+dx)+2ab(Ab+6aB)\cos(c+dx)+3(8Ba^3+4Aba^2-2b^2Ba+Ab^3))\sec(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx}{a} + \frac{(16a^2A+6abB-3Ab^2)\tan(c+dx)}{ad} \right) \\
 & \qquad \qquad \qquad \frac{A \tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{1}{6} \left(\frac{\int \frac{(-b(16Aa^2+6bBa-3Ab^2)\cos^2(c+dx)+2ab(Ab+6aB)\cos(c+dx)+3(8Ba^3+4Aba^2-2b^2Ba+Ab^3))\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{2a} + \frac{(16a^2A+6abB-3Ab^2)\tan(c+dx)}{ad} \right) \\
 & \qquad \qquad \qquad \frac{A \tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{1}{6} \left(\frac{\int \frac{-b(16Aa^2+6bBa-3Ab^2)\sin(c+dx+\frac{\pi}{2})^2+2ab(Ab+6aB)\sin(c+dx+\frac{\pi}{2})+3(8Ba^3+4Aba^2-2b^2Ba+Ab^3)}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} + \frac{(16a^2A+6abB-3Ab^2)\tan(c+dx)\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}{ad} \right) \\
 & \qquad \qquad \qquad \frac{A \tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}
 \end{aligned}$$

$$\frac{1}{6} \left(\frac{-\left((16a^2A+6abB-3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx\right) - \frac{\int \frac{(3b(8Ba^3+4Aba^2-2b^2Ba+Ab^3)+ab(16Aa^2+18bBa-Ab^2) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a}}{4a} + (16a^2A+6abB-3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 25

$$\frac{1}{6} \left(\frac{\frac{\int \frac{(3b(8Ba^3+4Aba^2-2b^2Ba+Ab^3)+ab(16Aa^2+18bBa-Ab^2) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - (16a^2A+6abB-3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx + (16a^2A+6abB-3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx}{4a} \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\frac{\int \frac{3b(8Ba^3+4Aba^2-2b^2Ba+Ab^3)+ab(16Aa^2+18bBa-Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2a} - (16a^2A+6abB-3Ab^2) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx + (16a^2A+6abB-3Ab^2) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{4a} \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3134

$$\frac{1}{6} \left(\frac{\frac{\int \frac{3b(8Ba^3+4Aba^2-2b^2Ba+Ab^3)+ab(16Aa^2+18bBa-Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{(16a^2A+6abB-3Ab^2) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{4a} + (16a^2A+6abB-3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\int \frac{3b(8Ba^3+4Aba^2-2b^2Ba+Ab^3)+ab(16Aa^2+18bBa-Ab^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(16a^2A+6abB-3Ab^2)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + \frac{2a}{4a}$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3132

$$\frac{1}{6} \left(\frac{\int \frac{3b(8Ba^3+4Aba^2-2b^2Ba+Ab^3)+ab(16Aa^2+18bBa-Ab^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(16a^2A+6abB-3Ab^2)\sqrt{a+b\cos(c+dx)} E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + \frac{(16a^2A+6abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{4a}$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3481

$$\frac{1}{6} \left(\frac{ab(16a^2A+18abB-Ab^2) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx + 3b(8a^3B+4a^2Ab-2ab^2B+Ab^3) \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2(16a^2A+6abB-3Ab^2)\sqrt{a+b\cos(c+dx)} E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + \frac{2a}{4a}$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{ab(16a^2A+18abB-Ab^2) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + 3b(8a^3B+4a^2Ab-2ab^2B+Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(16a^2A+6abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + \frac{2a}{4a}$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3142

3.303. $\int \sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^4(c+dx) dx$

$$\left(\frac{1}{6} \right) \left(\frac{ab(16a^2A+18abB-Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{3b(8a^3B+4a^2Ab-2ab^2B+Ab^3)}{b} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(16a^2A+18abB-Ab^2)}{2a} \right) \frac{1}{4a}$$

$$\frac{A \tan(c + dx) \sec^2(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3042

$$\left(\frac{1}{6} \right) \left(\frac{ab(16a^2A+18abB-Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{3b(8a^3B+4a^2Ab-2ab^2B+Ab^3)}{b} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(16a^2A+18abB-Ab^2)}{2a} \right) \frac{1}{4a}$$

$$\frac{A \tan(c + dx) \sec^2(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3140

$$\left(\frac{1}{6} \right) \left(\frac{3b(8a^3B+4a^2Ab-2ab^2B+Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2ab(16a^2A+18abB-Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(16a^2A+18abB-Ab^2)}{2a} \right) \frac{1}{4a}$$

$$\frac{A \tan(c + dx) \sec^2(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3286

$$\frac{1}{6} \left(\frac{3b(8a^3B+4a^2Ab-2ab^2B+Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2ab(16a^2A+18abB-Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}} - \frac{2(16a^2A+6abB-3Ab^2) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3b(8a^3B+4a^2Ab-2ab^2B+Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin\left(c+dx+\frac{\pi}{2}\right) \sqrt{\frac{a}{a+b} + \frac{b \sin\left(c+dx+\frac{\pi}{2}\right)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2ab(16a^2A+18abB-Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}} - \frac{2(16a^2A+6abB-3Ab^2) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3284

$$\frac{1}{6} \left(\frac{(16a^2A+6abB-3Ab^2) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} + \frac{2ab(16a^2A+18abB-Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}} + \frac{6b(8a^3B+4a^2Ab-2ab^2B+Ab^3) \sqrt{a+b \cos(c+dx)}}{b} \right)$$

$$\frac{A \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

input `Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

```
output (A*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + (((A*b +
6*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*a*d) + (((-2
*(16*a^2*A - 3*A*b^2 + 6*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*
x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*a*b*(16
*a^2*A - A*b^2 + 18*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c
+ d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*b*(4*a^2*A*b
+ A*b^3 + 8*a^3*B - 2*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*Elliptic
Pi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/(2*a)
+ ((16*a^2*A - 3*A*b^2 + 6*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/(
a*d))/(4*a))/6
```

3.303.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3478 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

rule 3481 `Int((((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^n), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
&& NeQ[c^2 - d^2, 0]
```

3.303.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2212 vs. $2(435) = 870$.

Time = 18.68 (sec) , antiderivative size = 2213, normalized size of antiderivative = 5.85

method	result	size
default	Expression too large to display	2213
parts	Expression too large to display	2716

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNV
ERBOSE)
```

output
$$\begin{aligned} & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*a*A*(-1/ \\ & 3*\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c) \\ & ^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)^3+5/12*b/a^2*\cos(1/2*d*x+1/2*c)*(-2*s \\ & \sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2 \\ & *c)^2-1)^2-1/24*(16*a^2+15*b^2)/a^3*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2 \\ & *c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)+5/48* \\ & b^2/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b) \\ &)^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*Ellip \\ & ticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/ \\ & 2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b \\ & +(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b) \\ &))^{(1/2)})-1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b) \\ & /a-b)^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & *EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+1/3/a*(\sin(1/2*d*x+1/2*c) \\ &)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/ \\ & 2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*EllipticE(\cos(1/2*d*x+1/2*c), \\ & (-2*b/(a-b))^{(1/2)})-5/16*b^2/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/ \\ & 2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2* \\ & d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+5/16/ \\ & a^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))\dots \end{aligned}$$

3.303.5 Fracas [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="fracas")`

output `Timed out`

3.303.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.303.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^4 dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^4, x)`

3.303.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^4 dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^4, x)`

3.303.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^4(c + dx) dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^4} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^4,x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^4, x)`

3.304 $\int \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

3.304.1 Optimal result	2793
3.304.2 Mathematica [A] (verified)	2794
3.304.3 Rubi [A] (verified)	2795
3.304.4 Maple [B] (verified)	2801
3.304.5 Fricas [C] (verification not implemented)	2802
3.304.6 Sympy [F(-1)]	2803
3.304.7 Maxima [F]	2803
3.304.8 Giac [F]	2803
3.304.9 Mupad [F(-1)]	2804

3.304.1 Optimal result

Integrand size = 33, antiderivative size = 378

$$\int \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx =$$

$$\frac{2(18a^3Ab - 246aAb^3 - 8a^4B - 33a^2b^2B - 147b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{315b^3d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(a^2 - b^2)(18a^2Ab - 75Ab^3 - 8a^3B - 39ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{315b^3d \sqrt{a+b \cos(c+dx)}}$$

$$- \frac{2(18a^2Ab - 75Ab^3 - 8a^3B - 39ab^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{315b^2d}$$

$$- \frac{2(18aAb - 8a^2B - 49b^2B)(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{315b^2d}$$

$$+ \frac{2(9Ab - 4aB)(a+b \cos(c+dx))^{5/2} \sin(c+dx)}{63b^2d}$$

$$+ \frac{2B \cos(c+dx)(a+b \cos(c+dx))^{5/2} \sin(c+dx)}{9bd}$$

output
$$\begin{aligned} & -2/315*(18*A*a*b-8*B*a^2-49*B*b^2)*(a+b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/b^2/d \\ & +2/63*(9*A*b-4*B*a)*(a+b*\cos(d*x+c))^{5/2}*\sin(d*x+c)/b^2/d+2/9*B*\cos(d*x+c) \\ & *(a+b*\cos(d*x+c))^{5/2}*\sin(d*x+c)/b/d-2/315*(18*A*a^2*b-75*A*b^3-8*B*a^3-39*B*a*b^2) \\ & *\sin(d*x+c)*(a+b*\cos(d*x+c))^{1/2}/b^2/d-2/315*(18*A*a^3*b-24*6*A*a*b^3-8*B*a^4-33*B*a^2*b^2-147*B*b^4) \\ & *(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{1/2}*(b/(a+b))^{1/2})*(a+b*\cos(d*x+c))^{1/2} \\ & /b^3/d/((a+b*\cos(d*x+c))/(a+b))^{1/2}+2/315*(a^2-b^2)*(18*A*a^2*b-75*A*b^3-8*B*a^3-39*B*a*b^2) \\ & *(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{1/2}*(b/(a+b))^{1/2})*((a+b*\cos(d*x+c))/(a+b))^{1/2} \\ & /b^3/d/(a+b*\cos(d*x+c))^{1/2} \end{aligned}$$

3.304.2 Mathematica [A] (verified)

Time = 1.83 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.77

$$\int \cos^2(c+dx)(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \frac{8\sqrt{\frac{a+b\cos(c+dx)}{a+b}}(b^2(153a^2Ab+75Ab^3+2a^3B+186ab^2B)\text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)+B\cos(c+dx))}{(1260b^3d\sqrt{a+b\cos(c+dx)})}$$

input `Integrate[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output
$$\begin{aligned} & (8*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/ (a + b)]*(b^2*(153*a^2*A*b + 75*A*b^3 + 2*a^3*B + 186*a*b^2*B)*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)] + (-18*a^3*A*b + 246*a*A*b^3 + 8*a^4*B + 33*a^2*b^2*B + 147*b^4*B)*((a + b)*\text{EllipticE}[(c + d*x)/2, (2*b)/(a + b)] - a*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)])) + b*(a + b*\text{Cos}[c + d*x])*((72*a^2*A*b + 690*A*b^3 - 32*a^3*B + 804*a*b^2*B)*\text{Sin}[c + d*x] + b*(2*(144*a*A*b + 6*a^2*B + 133*b^2*B)*\text{Sin}[2*(c + d*x)] + 5*b*(2*(9*A*b + 10*a*B)*\text{Sin}[3*(c + d*x)] + 7*b*B*\text{Sin}[4*(c + d*x)]))))/(1260*b^3*d*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) \end{aligned}$$

3.304.3 Rubi [A] (verified)

Time = 2.07 (sec) , antiderivative size = 394, normalized size of antiderivative = 1.04, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx)(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{3/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3469} \\
 & \frac{2 \int \frac{1}{2}(a+b\cos(c+dx))^{3/2} ((9Ab-4aB)\cos^2(c+dx)+7bB\cos(c+dx)+2aB) dx}{9b} + \\
 & \quad \frac{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{5/2}}{9bd} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int (a+b\cos(c+dx))^{3/2} ((9Ab-4aB)\cos^2(c+dx)+7bB\cos(c+dx)+2aB) dx}{9b} + \\
 & \quad \frac{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{5/2}}{9bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (a+b\sin(c+dx+\frac{\pi}{2}))^{3/2} ((9Ab-4aB)\sin(c+dx+\frac{\pi}{2})^2+7bB\sin(c+dx+\frac{\pi}{2})+2aB) dx}{9b} + \\
 & \quad \frac{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{5/2}}{9bd} \\
 & \quad \downarrow \text{3502} \\
 & \frac{2 \int \frac{1}{2}(a+b\cos(c+dx))^{3/2} (3b(15Ab-2aB)-(-8Ba^2+18Aba-49b^2B)\cos(c+dx)) dx}{7b} + \frac{2(9Ab-4aB)\sin(c+dx)(a+b\cos(c+dx))^{5/2}}{7bd} + \\
 & \quad \frac{9b}{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{5/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.304. $\int \cos^2(c+dx)(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx$

$$\frac{\int (a+b \cos(c+dx))^{3/2} (3b(15Ab-2aB) - (-8Ba^2+18Aba-49b^2B) \cos(c+dx)) dx}{7b} + \frac{2(9Ab-4aB) \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd} +$$

$$\frac{9b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{5/2}}$$

$$\frac{9bd}{9bd} \quad \downarrow \quad 3042$$

$$\frac{\int (a+b \sin(c+dx+\frac{\pi}{2}))^{3/2} (3b(15Ab-2aB) + (8Ba^2-18Aba+49b^2B) \sin(c+dx+\frac{\pi}{2})) dx}{7b} + \frac{2(9Ab-4aB) \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd} +$$

$$\frac{9b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{5/2}}$$

$$\frac{9bd}{9bd} \quad \downarrow \quad 3232$$

$$\frac{\frac{2}{5} \int \frac{3}{2} \sqrt{a+b \cos(c+dx)} (b(-2Ba^2+57Aba+49b^2B) - (-8Ba^3+18Aba^2-39b^2Ba-75Ab^3) \cos(c+dx)) dx - \frac{2(-8a^2B+18aAb-49b^2B) \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{5d}}{7b} + \frac{9b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{5/2}}$$

$$\frac{9bd}{9bd} \quad \downarrow \quad 27$$

$$\frac{\frac{3}{5} \int \sqrt{a+b \cos(c+dx)} (b(-2Ba^2+57Aba+49b^2B) - (-8Ba^3+18Aba^2-39b^2Ba-75Ab^3) \cos(c+dx)) dx - \frac{2(-8a^2B+18aAb-49b^2B) \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{5d}}{7b} + \frac{9b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{5/2}}$$

$$\frac{9bd}{9bd} \quad \downarrow \quad 3042$$

$$\frac{\frac{3}{5} \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (b(-2Ba^2+57Aba+49b^2B) + (8Ba^3-18Aba^2+39b^2Ba+75Ab^3) \sin(c+dx+\frac{\pi}{2})) dx - \frac{2(-8a^2B+18aAb-49b^2B) \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{5d}}{7b} + \frac{9b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{5/2}}$$

$$\frac{9bd}{9bd} \quad \downarrow \quad 3232$$

$$\frac{\frac{3}{5} \left(\frac{2}{3} \int \frac{b(2Ba^3+153Aba^2+186b^2Ba+75Ab^3) - (-8Ba^4+18Aba^3-33b^2Ba^2-246Ab^3a-147b^4B) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-8a^3B+18a^2Ab-39ab^2B-75Ab^3) \sin(c+dx)}{3d} \right)}{7b} + \frac{9b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{5/2}}$$

$$\frac{9bd}{9bd} \quad \downarrow \quad 27$$

3.304. $\int \cos^2(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

$$\frac{3}{5} \left(\frac{1}{3} \int \frac{b(2Ba^3 + 153Aba^2 + 186b^2Ba + 75Ab^3) - (-8Ba^4 + 18Aba^3 - 33b^2Ba^2 - 246Ab^3a - 147b^4B) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sin(c+dx)}{3d} \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3042

$$\frac{3}{5} \left(\frac{1}{3} \int \frac{b(2Ba^3 + 153Aba^2 + 186b^2Ba + 75Ab^3) + (8Ba^4 - 18Aba^3 + 33b^2Ba^2 + 246Ab^3a + 147b^4B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{2(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sin(c+dx)}{3d} \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3231

$$\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} - \frac{(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \int \sqrt{a+b \cos(c+dx)} dx}{b} \right) - \frac{2(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sin(c+dx)}{3d} \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3042

$$\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \int \sqrt{a+b \sin(c+dx + \frac{\pi}{2})} dx}{b} \right) - \frac{2(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sin(c+dx)}{3d} \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3134

$$\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{2(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sin(c+dx)}{3d} \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3042

$$\int \frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) dx$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3132

$$\int \frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{2(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) dx$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3142

$$\int \frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) dx$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3042

$$\int \frac{1}{3} \left(\frac{(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) dx$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

↓ 3140

$$\int \frac{1}{3} \left(\frac{2(a^2 - b^2)(-8a^3B + 18a^2Ab - 39ab^2B - 75Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd \sqrt{a+b \cos(c+dx)}} - \frac{2(-8a^4B + 18a^3Ab - 33a^2b^2B - 246aAb^3 - 147b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) dx$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{5/2}}{9bd}$$

3.304. $\int \cos^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

input `Int[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*Cos[c + d*x]*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(9*b*d) + ((2*(9*A*b - 4*a*B)*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*b*d) + ((-2*(18*a*A*b - 8*a^2*B - 49*b^2*B)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (3*((-2*(18*a^3*A*b - 246*a*A*b^3 - 8*a^4*B - 33*a^2*b^2*B - 147*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(a^2 - b^2)*(18*a^2*A*b - 75*A*b^3 - 8*a^3*B - 39*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 - (2*(18*a^2*A*b - 75*A*b^3 - 8*a^3*B - 39*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5)/(7*b))/(9*b)`

3.304.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*SIN[c + d*x]]/Sqrt[(a + b*SIN[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])]/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m - 1))*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B))*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(GtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.304.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1634 vs. $2(408) = 816$.

Time = 17.54 (sec) , antiderivative size = 1635, normalized size of antiderivative = 4.33

method	result	size
default	Expression too large to display	1635
parts	Expression too large to display	1824

```
input int(cos(d*x+c)^2*(a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -2/315*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*
B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b^5+(720*A*b^5+1360*B*a*b^4+224
0*B*b^5)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-936*A*a*b^4-1080*A*b^5-
424*B*a^2*b^3-2040*B*a*b^4-2072*B*b^5)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/
2*c)+(324*A*a^2*b^3+936*A*a*b^4+840*A*b^5-4*B*a^3*b^2+424*B*a^2*b^3+1568*B
*a*b^4+952*B*b^5)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-18*A*a^3*b^2-1
62*A*a^2*b^3-384*A*a*b^4-240*A*b^5+8*B*a^4*b+2*B*a^3*b^2-282*B*a^2*b^3-444
*B*a*b^4-168*B*b^5)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+18*A*(sin(1/2*
d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*El
lipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4*b-93*A*(sin(1/2*d*x+1/2
*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF
(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b^3+75*A*b^5*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF
(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-18*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1
/2*c),(-2*b/(a-b))^(1/2))*a^4*b+18*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a
-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),
(-2*b/(a-b))^(1/2))*a^3*b^2+246*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*
sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b
/(a-b))^(1/2))*a^2*b^3-246*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*s...
```


3.304.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.17 (sec) , antiderivative size = 639, normalized size of antiderivative = 1.69

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(16i Ba^5 - 36i Aa^4b + 60i Ba^3b^2 + 33i Aa^2b^3 - 264i Bab^4 - 225i Ab^5)\sqrt{b}\text{weierstrassPInverse}(4/3*(4a^2 - 3b^2)/b^2, -8/27*(8a^3 - 9ab^2)/b^3, 1/3*(3b\cos(dx + c) + 3Ib\sin(dx + c) + 2a)/b) + \sqrt{2}(-16i Ba^5 + 36i Aa^4b - 60i Bba^3b^2 - 33i Aa^2b^3 + 264i Bab^4 + 225i Ab^5)\sqrt{b}\text{weierstrassPInverse}(4/3*(4a^2 - 3b^2)/b^2, -8/27*(8a^3 - 9ab^2)/b^3, 1/3*(3b\cos(dx + c) - 3Ib\sin(dx + c) + 2a)/b) - 3\sqrt{2}(-8i Ba^4b + 18i Aa^3b^2 - 33i Bba^2b^3 - 246i Aa^2b^4 - 147i Bb^5)\sqrt{b}\text{weierstrassZeta}(4/3*(4a^2 - 3b^2)/b^2, -8/27*(8a^3 - 9ab^2)/b^3, \text{weierstrassPInverse}(4/3*(4a^2 - 3b^2)/b^2, -8/27*(8a^3 - 9ab^2)/b^3, 1/3*(3b\cos(dx + c) + 3Ib\sin(dx + c) + 2a)/b)) - 3\sqrt{2}(8i Ba^4b - 18i Aa^3b^2 + 33i Bba^2b^3 + 246i Aa^2b^4 + 147i Bb^5)\sqrt{b}\text{weierstrassZeta}(4/3*(4a^2 - 3b^2)/b^2, -8/27*(8a^3 - 9ab^2)/b^3, \text{weierstrassPInverse}(4/3*(4a^2 - 3b^2)/b^2, -8/27*(8a^3 - 9ab^2)/b^3, 1/3*(3b\cos(dx + c) - 3Ib\sin(dx + c) + 2a)/b)) + 6*(35Bb^5\cos(dx + c)^3 - 4Ba^3b^2 + 9Aa^2b^3 + 88Bab^4 + 75Aab^5 + 5*(10Bab^4 + 9Ab^5)\cos(dx + c)^2 + (3Ba^2b^3 + 72Aa^2b^4 + 49Bb^5)\cos(dx + c))\sqrt{b\cos(dx + c) + a}\sin(dx + c))/(b^4d)$$

```
input integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/945*(sqrt(2)*(16*I*B*a^5 - 36*I*A*a^4*b + 60*I*B*a^3*b^2 + 33*I*A*a^2*b^3 - 264*I*B*a*b^4 - 225*I*A*b^5)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(-16*I*B*a^5 + 36*I*A*a^4*b - 60*I*B*a^3*b^2 - 33*I*A*a^2*b^3 + 264*I*B*a*b^4 + 225*I*A*b^5)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(-8*I*B*a^4*b + 18*I*A*a^3*b^2 - 33*I*B*a^2*b^3 - 246*I*A*a*b^4 - 147*I*B*b^5)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(8*I*B*a^4*b - 18*I*A*a^3*b^2 + 33*I*B*a^2*b^3 + 246*I*A*a*b^4 + 147*I*B*b^5)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(35*B*b^5*cos(d*x + c)^3 - 4*B*a^3*b^2 + 9*A*a^2*b^3 + 88*B*a*b^4 + 75*A*b^5 + 5*(10*B*a*b^4 + 9*A*b^5)*cos(d*x + c)^2 + (3*B*a^2*b^3 + 72*A*a^2*b^4 + 49*B*b^5)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^4*d)
```

3.304.6 Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.304.7 Maxima [F]

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^2, x)`

3.304.8 Giac [F]

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^2, x)`

3.304.9 Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^2 (A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2), x)`

3.305 $\int \cos(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

3.305.1 Optimal result	2805
3.305.2 Mathematica [A] (verified)	2806
3.305.3 Rubi [A] (verified)	2806
3.305.4 Maple [B] (verified)	2812
3.305.5 Fricas [C] (verification not implemented)	2813
3.305.6 Sympy [F(-1)]	2813
3.305.7 Maxima [F]	2814
3.305.8 Giac [F]	2814
3.305.9 Mupad [F(-1)]	2814

3.305.1 Optimal result

Integrand size = 31, antiderivative size = 297

$$\int \cos(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx = \frac{2(21a^2Ab + 63Ab^3 - 6a^3B + 82ab^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) + 2(a^2 - b^2) (21aAb - 6a^2B + 25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{2(21aAb - 6a^2B + 25b^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{105bd} + \frac{2(7Ab - 2aB)(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{35bd} + \frac{2B(a+b \cos(c+dx))^{5/2} \sin(c+dx)}{7bd}$$

```
output 2/35*(7*A*b-2*B*a)*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/b/d+2/7*B*(a+b*cos(d*x+c))^(5/2)*sin(d*x+c)/b/d+2/105*(21*A*a*b-6*B*a^2+25*B*b^2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/b/d+2/105*(21*A*a^2*b+63*A*b^3-6*B*a^3+82*B*a*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/105*(a^2-b^2)*(21*A*a*b-6*B*a^2+25*B*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b^2/d/(a+b*cos(d*x+c))^(1/2)
```

3.305.2 Mathematica [A] (verified)

Time = 3.09 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.78

$$\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{4\sqrt{\frac{a+b \cos(c+dx)}{a+b}}(b^2(84aAb + 51a^2B + 25b^2B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + (21a^2Ab + 6$$

input `Integrate[Cos[c + d*x]*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(4*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*(b^2*(84*a*A*b + 51*a^2*B + 25*b^2*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + (21*a^2*A*b + 63*A*b^3 - 6*a^3*B + 82*a*b^2*B)*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)])) + b*(a + b*Cos[c + d*x])*((168*a*A*b + 12*a^2*B + 115*b^2*B)*Sin[c + d*x] + 3*b*(2*(7*A*b + 8*a*B)*Sin[2*(c + d*x)] + 5*b*B*Ssin[3*(c + d*x)])))/(210*b^2*d*Sqrt[a + b*Cos[c + d*x]])`

3.305.3 Rubi [A] (verified)

Time = 1.58 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.03, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.645$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3447} \\ & \int (a + b \cos(c + dx))^{3/2} (A \cos(c + dx) + B \cos^2(c + dx)) dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.305. $\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(A \sin \left(c + dx + \frac{\pi}{2} \right) + B \sin \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx \\
& \quad \downarrow \text{3502} \\
& \frac{2 \int \frac{1}{2} (a + b \cos(c + dx))^{3/2} (5bB + (7Ab - 2aB) \cos(c + dx)) dx}{7b} + \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7bd} \\
& \quad \downarrow \text{27} \\
& \frac{\int (a + b \cos(c + dx))^{3/2} (5bB + (7Ab - 2aB) \cos(c + dx)) dx}{7b} + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} (5bB + (7Ab - 2aB) \sin \left(c + dx + \frac{\pi}{2} \right)) dx}{7b} + \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7bd} \\
& \quad \downarrow \text{3232} \\
& \frac{\frac{2}{5} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (b(21Ab + 19aB) + (-6Ba^2 + 21Aba + 25b^2B) \cos(c + dx)) dx + \frac{2(7Ab - 2aB) \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{5d}}{7b} \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7bd} \\
& \quad \downarrow \text{27} \\
& \frac{\frac{1}{5} \int \sqrt{a + b \cos(c + dx)} (b(21Ab + 19aB) + (-6Ba^2 + 21Aba + 25b^2B) \cos(c + dx)) dx + \frac{2(7Ab - 2aB) \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{5d}}{7b} \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{5} \int \sqrt{a + b \sin \left(c + dx + \frac{\pi}{2} \right)} (b(21Ab + 19aB) + (-6Ba^2 + 21Aba + 25b^2B) \sin \left(c + dx + \frac{\pi}{2} \right)) dx + \frac{2(7Ab - 2aB) \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{5d}}{7b} \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7bd} \\
& \quad \downarrow \text{3232}
\end{aligned}$$

3.305. $\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{b(51Ba^2+84Aba+25b^2B)+(-6Ba^3+21Aba^2+82b^2Ba+63Ab^3) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx + \frac{2(-6a^2B+21aAb+25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{b(51Ba^2+84Aba+25b^2B)+(-6Ba^3+21Aba^2+82b^2Ba+63Ab^3) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + \frac{2(-6a^2B+21aAb+25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{b(51Ba^2+84Aba+25b^2B)+(-6Ba^3+21Aba^2+82b^2Ba+63Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(-6a^2B+21aAb+25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3231

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \int \sqrt{a+b \cos(c+dx)} dx}{b} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} \right) + \frac{2(-6a^2B+21aAb+25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \frac{2(-6a^2B+21aAb+25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3134

3.305. $\int \cos(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3132

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3142

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-6a^2B+21aAb+25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

↓ 3140

3.305. $\int \cos(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

$$\frac{\frac{1}{5} \left(\frac{2(-6a^2B+21aAb+25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} + \frac{1}{3} \left(\frac{2(-6a^3B+21a^2Ab+82ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right)}{7b} - \frac{2B \sin(c+dx)(a+b \cos(c+dx))^{5/2}}{7bd}$$

input `Int[Cos[c + d*x]*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*b*d) + ((2*(7*A*b - 2*a*B)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (((2*(21*a^2*A*b + 63*A*b^3 - 6*a^3*B + 82*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(21*a*A*b - 6*a^2*B + 25*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 + (2*(21*a*A*b - 6*a^2*B + 25*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5)/(7*b)`

3.305.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*SIN[c + d*x]]/Sqrt[(a + b*SIN[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.305.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1304 vs. $2(331) = 662$.

Time = 15.98 (sec) , antiderivative size = 1305, normalized size of antiderivative = 4.39

method	result	size
default	Expression too large to display	1305
parts	Expression too large to display	1492

```
input int(cos(d*x+c)*(a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output -2/105*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^4+(-168*A*b^4-312*B*a*b^3-360*B*
b^4)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(252*A*a*b^3+168*A*b^4+108*B*
a^2*b^2+312*B*a*b^3+280*B*b^4)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-8
4*A*a^2*b^2-126*A*a*b^3-42*A*b^4-6*B*a^3*b-54*B*a^2*b^2-128*B*a*b^3-80*B*b
^4)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-21*A*(sin(1/2*d*x+1/2*c)^2)^(1
/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3*b+21*a*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1
/2*c),(-2*b/(a-b))^(1/2))*b^3+21*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b
)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2
*b/(a-b))^(1/2))*a^3*b-21*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1
/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b
))^(1/2))*a^2*b^2+63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*
x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1
/2))*a*b^3-63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c
)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^
4+6*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/
(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4-31*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))...
```

3.305.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.15 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.89

$$\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(-12i Ba^4 + 42i Aa^3b + 11i Ba^2b^2 - 126i Aab^3 - 75i Bb^4)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4}{3}(4a^2 - 3b^2)/b^2, -8/27(8a^3 - 9ab^2)/b^3, 1/3(3b\cos(dx + c) + 3Ib\sin(dx + c) + 2a)/b\right) + \sqrt{2}(12i Ba^4 - 42i Aa^3b - 11i Bb^2a^2 + 126i Aa^3b^3 + 75i Bb^4)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4}{3}(4a^2 - 3b^2)/b^2, -8/27(8a^3 - 9ab^2)/b^3, 1/3(3b\cos(dx + c) - 3Ib\sin(dx + c) + 2a)/b\right) - 3\sqrt{2}(6i Ba^3b - 21i Aa^2b^2 - 82i Bb^2a^2b^3 - 63i Aa^2b^4)\sqrt{b}\text{weierstrassZeta}\left(\frac{4}{3}(4a^2 - 3b^2)/b^2, -8/27(8a^3 - 9ab^2)/b^3, \text{weierstrassPInverse}\left(\frac{4}{3}(4a^2 - 3b^2)/b^2, -8/27(8a^3 - 9ab^2)/b^3, 1/3(3b\cos(dx + c) + 3Ib\sin(dx + c) + 2a)/b\right) - 3\sqrt{2}(-6i Ba^3b + 21i Aa^2b^2 + 82i Bb^2a^2b^3 + 63i Aa^2b^4)\sqrt{b}\text{weierstrassZeta}\left(\frac{4}{3}(4a^2 - 3b^2)/b^2, -8/27(8a^3 - 9ab^2)/b^3, \text{weierstrassPInverse}\left(\frac{4}{3}(4a^2 - 3b^2)/b^2, -8/27(8a^3 - 9ab^2)/b^3, 1/3(3b\cos(dx + c) - 3Ib\sin(dx + c) + 2a)/b\right) + 6(15Bb^4\cos(dx + c)^2 + 3Bb^2a^2b^2 + 42Aa^3b^3 + 25Bb^4 + 3(8Bb^4\cos(dx + c) + 7Aa^2b^4)\cos(dx + c))\sqrt{b}\cos(dx + c) + a)\sin(dx + c)\right)}{(b^3d)}$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/315*(sqrt(2)*(-12*I*B*a^4 + 42*I*A*a^3*b + 11*I*B*a^2*b^2 - 126*I*A*a*b^3 - 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(12*I*B*a^4 - 42*I*A*a^3*b - 11*I*B*a^2*b^2 + 126*I*A*a*b^3 + 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(6*I*B*a^3*b - 21*I*A*a^2*b^2 - 82*I*B*a*b^3 - 63*I*A*b^4)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(-6*I*B*a^3*b + 21*I*A*a^2*b^2 + 82*I*B*a*b^3 + 63*I*A*b^4)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(15*B*b^4*cos(d*x + c)^2 + 3*B*a^2*b^2 + 42*A*a*b^3 + 25*B*b^4 + 3*(8*B*a*b^3 + 7*A*b^4)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^3*d)
```

3.305.6 Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.305.7 Maxima [F]

$$\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*cos(d*x + c), x)`

3.305.8 Giac [F]

$$\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*cos(d*x + c), x)`

3.305.9 Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx) (A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2), x)`

3.306 $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx$

3.306.1 Optimal result	2815
3.306.2 Mathematica [A] (verified)	2816
3.306.3 Rubi [A] (verified)	2816
3.306.4 Maple [B] (verified)	2820
3.306.5 Fracas [C] (verification not implemented)	2821
3.306.6 Sympy [F]	2822
3.306.7 Maxima [F]	2822
3.306.8 Giac [F]	2823
3.306.9 Mupad [F(-1)]	2823

3.306.1 Optimal result

Integrand size = 25, antiderivative size = 225

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{2(20aAb + 3a^2B + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) + 2(a^2 - b^2) (5Ab + 3aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + \frac{2(5Ab + 3aB) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{15d} + \frac{2B(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{5d}}{15bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

output

```
2/5*B*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/15*(5*A*b+3*B*a)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/15*(20*A*a*b+3*B*a^2+9*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b)))^(1/2)*(a+b*cos(d*x+c))^(1/2)/b/d/((a+b*cos(d*x+c))/(a+b))^2-2/15*(a^2-b^2)*(5*A*b+3*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b/d/(a+b*cos(d*x+c))^(1/2)
```

3.306.2 Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.90

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{2 \left(b(15a^2A + 5Ab^2 + 12abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + (20aAb + 3a^2 + B \cos(c+dx)) \right)}{5d}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`output `(2*(b*(15*a^2*A + 5*A*b^2 + 12*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + (20*a*A*b + 3*a^2*B + 9*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)]) + b*(a + b*Cos[c + d*x])*(5*A*b + 6*a*B + 3*b*B*Cos[c + d*x])*Sin[c + d*x))/(15*b*d*Sqrt[a + b*Cos[c + d*x]])`**3.306.3 Rubi [A] (verified)**Time = 1.13 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(A + B \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx \\ & \quad \downarrow \text{3232} \\ & \frac{2}{5} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (5aA + 3bB + (5Ab + 3aB) \cos(c + dx)) dx + \\ & \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\begin{aligned}
& \frac{1}{5} \int \sqrt{a + b \cos(c + dx)} (5aA + 3bB + (5Ab + 3aB) \cos(c + dx)) dx + \\
& \quad \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(5aA + 3bB + (5Ab + 3aB) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\
& \quad \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d} \\
& \quad \downarrow \text{3232} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{15Aa^2 + 12bBa + 5Ab^2 + (3Ba^2 + 20Aba + 9b^2B) \cos(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{2(3aB + 5Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right. \\
& \quad \left. \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \left(\frac{1}{3} \int \frac{15Aa^2 + 12bBa + 5Ab^2 + (3Ba^2 + 20Aba + 9b^2B) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \frac{2(3aB + 5Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right. \\
& \quad \left. \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{1}{3} \int \frac{15Aa^2 + 12bBa + 5Ab^2 + (3Ba^2 + 20Aba + 9b^2B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2(3aB + 5Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right. \\
& \quad \left. \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d} \right) \\
& \quad \downarrow \text{3231} \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{(3a^2B + 20aAb + 9b^2B) \int \sqrt{a + b \cos(c + dx)} dx}{b} - \frac{(a^2 - b^2) (3aB + 5Ab) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) + \frac{2(3aB + 5Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \right. \\
& \quad \left. \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(3a^2B + 20aAb + 9b^2B) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx}{b} - \frac{(a^2 - b^2) (3aB + 5Ab) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3134}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(3a^2B + 20aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (3aB + 5Ab) \int \frac{1}{\sqrt{a+b \sin(c + dx)}} dx}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3042}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(3a^2B + 20aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (3aB + 5Ab) \int \frac{1}{\sqrt{a+b \sin(c + dx)}} dx}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3132}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(3a^2B + 20aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (3aB + 5Ab) \int \frac{1}{\sqrt{a+b \sin(c + dx)}} dx}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3142}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(3a^2B + 20aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (3aB + 5Ab) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b \sqrt{a + b \cos(c + dx)}} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right) \downarrow \text{3042}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(3a^2B + 20aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2)(3aB + 5Ab) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b \sqrt{a + b \cos(c + dx)}} \right) - \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right)$$

↓ 3140

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(3a^2B + 20aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(a^2 - b^2)(3aB + 5Ab) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{bd \sqrt{a + b \cos(c + dx)}} \right) - \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d} \right)$$

input `Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (((2*(20*a*A*b + 3*a^2*B + 9*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(5*A*b + 3*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 + (2*(5*A*b + 3*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

3.306.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a
+ b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b)
+ (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && !GtQ[a + b, 0]
```

```
rule 3231 Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (
f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]
], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b
, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

```
rule 3232 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*
d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ
[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m,
0] && IntegerQ[2*m]
```

3.306.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 992 vs. $2(263) = 526$.

Time = 13.12 (sec) , antiderivative size = 993, normalized size of antiderivative = 4.41

method	result	size
default	Expression too large to display	993
parts	Expression too large to display	1115

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

$$3.306. \quad \int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx$$

output

```

-2/15*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^3+(20*A*b^3+36*B*a*b^2+24*B*b^3)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*a*b^2-10*A*b^3-12*B*a^2*b-18*B*a*b^2-6*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+5*A*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+20*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b-20*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b^2-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3+3*B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^2+3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+9*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(...

```

3.306.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 493, normalized size of antiderivative = 2.19

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \frac{\sqrt{2}(6i Ba^3 - 5i Aa^2b - 18i Bab^2 - 15i Ab^3)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4(4a^2-3b^2)}{3b^2}, -\frac{8(8a^3)}{27}\right)}{\dots}$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

```
output 1/45*(sqrt(2)*(6*I*B*a^3 - 5*I*A*a^2*b - 18*I*B*a*b^2 - 15*I*A*b^3)*sqrt(b)
)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3
, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(-6*I*B*a
^3 + 5*I*A*a^2*b + 18*I*B*a*b^2 + 15*I*A*b^3)*sqrt(b)*weierstrassPInverse(
4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c)
) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(-3*I*B*a^2*b - 20*I*A*a*b^2
- 9*I*B*b^3)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3
- 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3
- 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3
*sqrt(2)*(3*I*B*a^2*b + 20*I*A*a*b^2 + 9*I*B*b^3)*sqrt(b)*weierstrassZeta(
4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(
4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c)
) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(3*B*b^3*cos(d*x + c) + 6*B*a*b^2 +
5*A*b^3)*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^2*d)
```

3.306.6 Sympy [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2} dx$$

```
input integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)
```

```
output Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**(3/2), x)
```

3.306.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} dx$$

```
input integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2), x)
```

3.306.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2), x)`

3.306.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2), x)`

3.307 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec(c+dx) dx$

3.307.1 Optimal result	2824
3.307.2 Mathematica [C] (verified)	2825
3.307.3 Rubi [A] (verified)	2825
3.307.4 Maple [B] (verified)	2831
3.307.5 Fricas [F]	2832
3.307.6 Sympy [F(-1)]	2833
3.307.7 Maxima [F]	2833
3.307.8 Giac [F]	2833
3.307.9 Mupad [F(-1)]	2834

3.307.1 Optimal result

Integrand size = 31, antiderivative size = 236

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{2(3Ab + 4aB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{3d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2(3aAb - a^2B + b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3d \sqrt{a + b \cos(c + dx)}} + \frac{2a^2A \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}} + \frac{2bB \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3d}$$

output

```
2/3*b*B*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/3*(3*A*b+4*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2/3*(3*A*a*b-B*a^2+B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)+2*a^2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)
```

3.307.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.20 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.72

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{4(6aAb + 3a^2B + b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2(6a^2A + 3Ab^2 + 4abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `((4*(6*a*A*b + 3*a^2*B + b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(6*a^2*A + 3*A*b^2 + 4*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + ((2*I)*(3*A*b + 4*a*B)*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)])))/(a*b*Sqrt[-(a + b)^(-1)]) + 4*b*B*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(6*d)`

3.307.3 Rubi [A] (verified)

Time = 1.97 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.03, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 3469, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})} dx \\
& \quad \downarrow \text{3469} \\
& \frac{2}{3} \int \frac{(3Aa^2 + b(3Ab + 4aB) \cos^2(c + dx) + (3Ba^2 + 6Aba + b^2B) \cos(c + dx)) \sec(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{(3Aa^2 + b(3Ab + 4aB) \cos^2(c + dx) + (3Ba^2 + 6Aba + b^2B) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3Aa^2 + b(3Ab + 4aB) \sin(c + dx + \frac{\pi}{2})^2 + (3Ba^2 + 6Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3538} \\
& \frac{1}{3} \left((4aB + 3Ab) \int \sqrt{a + b \cos(c + dx)} dx - \frac{\int -\frac{(3Aba^2 + b(-Ba^2 + 3Aba + b^2B) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{25} \\
& \frac{1}{3} \left(\frac{\int \frac{(3Aba^2 + b(-Ba^2 + 3Aba + b^2B) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} + (4aB + 3Ab) \int \sqrt{a + b \cos(c + dx)} dx \right) + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(\frac{\int \frac{3Aba^2 + b(-Ba^2 + 3Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} + (4aB + 3Ab) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx \right) + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}
\end{aligned}$$

$$\begin{aligned} & \downarrow \text{3134} \\ & \frac{1}{3} \left(\frac{\int \frac{3Aba^2 + b(-Ba^2 + 3Aba + b^2B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{(4aB + 3Ab) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\ & \quad \frac{2bB \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ & \frac{1}{3} \left(\frac{\int \frac{3Aba^2 + b(-Ba^2 + 3Aba + b^2B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{(4aB + 3Ab) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\ & \quad \frac{2bB \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3132} \\ & \frac{1}{3} \left(\frac{\int \frac{3Aba^2 + b(-Ba^2 + 3Aba + b^2B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{2(4aB + 3Ab) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\ & \quad \frac{2bB \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3481} \\ & \frac{1}{3} \left(\frac{b(a^2(-B) + 3aAb + b^2B) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx + 3a^2Ab \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} + \frac{2(4aB + 3Ab) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\ & \quad \frac{2bB \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ & \frac{1}{3} \left(\frac{b(a^2(-B) + 3aAb + b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + 3a^2Ab \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{2(4aB + 3Ab) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + \\ & \quad \frac{2bB \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} \end{aligned}$$

$$\downarrow \text{3142}$$

$$\frac{1}{3} \left(\frac{b(a^2(-B)+3aAb+b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx + 3a^2 Ab \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{2(4aB + 3Ab)}{3} \right)$$

$$\frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{b(a^2(-B)+3aAb+b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + 3a^2 Ab \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{2(4aB + 3Ab)}{3} \right)$$

$$\frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3140

$$\frac{1}{3} \left(\frac{3a^2 Ab \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(a^2(-B)+3aAb+b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} + \frac{2(4aB + 3Ab)}{3} \right)$$

$$\frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3286

$$\frac{1}{3} \left(\frac{3a^2 Ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx + \frac{2b(a^2(-B)+3aAb+b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} + \frac{2(4aB + 3Ab)}{3} \right)$$

$$\frac{2bB \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{3a^2 Ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2b(a^2(-B)+3aAb+b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) + \frac{2bB \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3284

$$\frac{1}{3} \left(\frac{2b(a^2(-B)+3aAb+b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} + \frac{6a^2 Ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) + \frac{2(4aB+3b^2) \sqrt{a+b \cos(c+dx)}}{3d}$$

```
input Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x], x]
```

```
output ((2*(3*A*b + 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*b*(3*a*A*b - a^2*B + b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*a^2*A*b*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]])))/b)/3 + (2*b*B*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d)
```

3.307.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.307. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx$

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3481 Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3538 Int(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.307.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 700 vs. 2(303) = 606.

Time = 10.70 (sec) , antiderivative size = 701, normalized size of antiderivative = 2.97

method	result
parts	$-\frac{2A\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}{a-b}}}\left(bF\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)a+bE\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+(a+b)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{-2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}}$
default	$-\frac{2\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(4B\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2+3Aab\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}{a-b}}\right)F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+(a+b)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{-2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}}$

3.307. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx$

input `int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVERBOSE)`

output `-2*A*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*(b*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a+b*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a-b^2*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)))-EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))*a^2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d-2/3*B*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*cos(1/2*d*x+1/2*c)^5*b^2+2*cos(1/2*d*x+1/2*c)^3*a*b-6*cos(1/2*d*x+1/2*c)^3*b^2-a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2))*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2))*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+4*(sin(1/2*d*x+1/2*c)^2)^(1/2))*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2-4*(sin(1/2*d*x+1/2*c)^2)^(1/2))*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b-2*cos(1/2*d*x+1/2*c)*a*b+2*cos(1/2*d*x+1/2*c)*b^2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d`

3.307.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c), x)`

3.307.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `Timed out`

3.307.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c), x)`

3.307.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c), x)`

3.307.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x),x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x), x)`

3.308 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.308.1 Optimal result	2835
3.308.2 Mathematica [C] (verified)	2836
3.308.3 Rubi [A] (verified)	2836
3.308.4 Maple [B] (verified)	2843
3.308.5 Fricas [F]	2844
3.308.6 Sympy [F(-1)]	2844
3.308.7 Maxima [F]	2844
3.308.8 Giac [F]	2845
3.308.9 Mupad [F(-1)]	2845

3.308.1 Optimal result

Integrand size = 33, antiderivative size = 232

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$-\frac{(aA - 2bB)\sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{(a^2 A + 2Ab^2 + 2abB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d\sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{a(3Ab + 2aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d\sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{aA\sqrt{a + b \cos(c + dx)} \tan(c + dx)}{d}$$

output

```
-(A*a-2*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+(A*a^2+2*A*b^2+2*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)+a*(3*A*b+2*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2, 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)+a*A*(a+b*cos(d*x+c))^(1/2)*tan(d*x+c)/d
```

3.308.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.92 (sec) , antiderivative size = 398, normalized size of antiderivative = 1.72

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{8b(Ab+2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(5aAb+4a^2B+2b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \dots$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `((8*b*(A*b + 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(5*a*A*b + 4*a^2*B + 2*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + ((2*I)*(-(a*A) + 2*b*B)*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)]))/((a*b*Sqrt[-(a + b)^(-1)]) + 4*a*A*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/((4*d))`

3.308.3 Rubi [A] (verified)

Time = 1.98 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.06, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3468, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

↓ 3042

3.308. $\int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^2(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^2} dx \\
& \quad \downarrow \text{3468} \\
& \int \frac{(-b(aA - 2bB) \cos^2(c + dx) + 2b(Ab + 2aB) \cos(c + dx) + a(3Ab + 2aB)) \sec(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \\
& \quad \frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \int \frac{(-b(aA - 2bB) \cos^2(c + dx) + 2b(Ab + 2aB) \cos(c + dx) + a(3Ab + 2aB)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \\
& \quad \frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{-b(aA - 2bB) \sin(c + dx + \frac{\pi}{2})^2 + 2b(Ab + 2aB) \sin(c + dx + \frac{\pi}{2}) + a(3Ab + 2aB)}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d} \\
& \quad \downarrow \text{3538} \\
& \frac{1}{2} \left(-\frac{\int \frac{(ab(3Ab + 2aB) + b(Aa^2 + 2bBa + 2Ab^2) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} - \left((aA - 2bB) \int \sqrt{a + b \cos(c + dx)} dx \right) \right) + \\
& \quad \frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d} \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left(\frac{\int \frac{(ab(3Ab + 2aB) + b(Aa^2 + 2bBa + 2Ab^2) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} - (aA - 2bB) \int \sqrt{a + b \cos(c + dx)} dx \right) + \\
& \quad \frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{2} \left(\frac{\int \frac{ab(3Ab+2aB)+b(Aa^2+2bBa+2Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - (aA - 2bB) \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx \right) +$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3134

$$\frac{1}{2} \left(\frac{\int \frac{ab(3Ab+2aB)+b(Aa^2+2bBa+2Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(aA - 2bB) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) +$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{\int \frac{ab(3Ab+2aB)+b(Aa^2+2bBa+2Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(aA - 2bB) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) +$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3132

$$\frac{1}{2} \left(\frac{\int \frac{ab(3Ab+2aB)+b(Aa^2+2bBa+2Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(aA - 2bB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) +$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3481

$$\frac{1}{2} \left(\frac{b(a^2A + 2abB + 2Ab^2) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx + ab(2aB + 3Ab) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} - \frac{2(aA - 2bB) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) +$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3042

3.308. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx$

$$\frac{1}{2} \left(\frac{b(a^2A + 2abB + 2Ab^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + ab(2aB + 3Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(aA - 2b^2)}{d} \right)$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3142

$$\frac{1}{2} \left(\frac{b(a^2A + 2abB + 2Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx + ab(2aB + 3Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(aA - 2b^2)}{d} \right)$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{b(a^2A + 2abB + 2Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + ab(2aB + 3Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(aA - 2b^2)}{d} \right)$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3140

$$\frac{1}{2} \left(\frac{ab(2aB + 3Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(a^2A + 2abB + 2Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} - \frac{2(aA - 2b^2)}{d} \right)$$

$$\frac{aA \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d}$$

↓ 3286

3.308. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx$

$$\frac{1}{2} \left(\frac{ab(2aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2b(a^2A+2abB+2Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{\sqrt{a+b\cos(c+dx)}} - \frac{2(aA - 2b^2)}{b} \right)$$

$$\frac{aA \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{ab(2aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2b(a^2A+2abB+2Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{\sqrt{a+b\cos(c+dx)}} - \frac{2(aA - 2b^2)}{b} \right)$$

$$\frac{aA \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

↓ 3284

$$\frac{1}{2} \left(\frac{2b(a^2A+2abB+2Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{2ab(2aB+3Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(aA - 2b^2)}{b} \right)$$

$$\frac{aA \tan(c+dx)\sqrt{a+b\cos(c+dx)}}{d}$$

input `Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output `((-2*(a*A - 2*b*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*b*(a^2*A + 2*A*b^2 + 2*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*a*b*(3*A*b + 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])/b)/2 + (a*A*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/d`

3.308.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])]/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[(a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.308.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 990 vs. $2(305) = 610$.

Time = 12.20 (sec) , antiderivative size = 991, normalized size of antiderivative = 4.27

method	result	size
parts	Expression too large to display	991
default	Expression too large to display	1168

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNV
ERBOSE)
```

```
output -A*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*cos(1/2*
d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*a*b+(-2*a^2-2*a*b)*sin(1/2*d*x+1/2*c)^2*co
s(1/2*d*x+1/2*c)-2*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)
)*a^2+2*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^2-EllipticE(cos
(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2+b*EllipticE(cos(1/2*d*x+1/2*c),(-2
*b/(a-b))^(1/2))*a-3*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))*a
*b)*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*
d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))*a^2+2*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*
c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2+(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*b*Ellipt
icE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a-3*a*b*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticPi(cos(1
/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2)))/(2*cos(1/2*d*x+1/2*c)^2-1)/(-2*sin(1/
2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(-2*
b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d-2*B*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*si
n(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*...
```

3.308.5 Fricas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^2 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^2, x)`

3.308.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Timed out`

3.308.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^2 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^2, x)`

3.308.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^2 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^2, x)`

3.308.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^2} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^2,x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^2, x)`

3.309 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.309.1 Optimal result	2846
3.309.2 Mathematica [C] (verified)	2847
3.309.3 Rubi [A] (verified)	2848
3.309.4 Maple [B] (verified)	2855
3.309.5 Fricas [F(-1)]	2856
3.309.6 Sympy [F(-1)]	2857
3.309.7 Maxima [F]	2857
3.309.8 Giac [F]	2857
3.309.9 Mupad [F(-1)]	2858

3.309.1 Optimal result

Integrand size = 33, antiderivative size = 295

$$\begin{aligned} & \int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \\ & - \frac{(5Ab + 4aB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{4d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\ & + \frac{(7aAb + 4a^2B + 8b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4d \sqrt{a + b \cos(c + dx)}} \\ & + \frac{(4a^2A + 3Ab^2 + 12abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4d \sqrt{a + b \cos(c + dx)}} \\ & + \frac{(5Ab + 4aB) \sqrt{a + b \cos(c + dx)} \tan(c + dx)}{4d} \\ & + \frac{aA \sqrt{a + b \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{2d} \end{aligned}$$

3.309.3 Rubi [A] (verified)

Time = 2.60 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.05, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3468, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^3(c+dx)(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^3} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{2} \int \frac{(b(aA+4bB)\cos^2(c+dx)+2(Aa^2+4bBa+2Ab^2)\cos(c+dx)+a(5Ab+4aB))\sec^2(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx +$$

$$\frac{aA\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d}$$

$$\downarrow \text{27}$$

$$\frac{1}{4} \int \frac{(b(aA+4bB)\cos^2(c+dx)+2(Aa^2+4bBa+2Ab^2)\cos(c+dx)+a(5Ab+4aB))\sec^2(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx +$$

$$\frac{aA\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \frac{b(aA+4bB)\sin(c+dx+\frac{\pi}{2})^2+2(Aa^2+4bBa+2Ab^2)\sin(c+dx+\frac{\pi}{2})+a(5Ab+4aB)}{\sin(c+dx+\frac{\pi}{2})^2\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx +$$

$$\frac{aA\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2d}$$

$$\downarrow \text{3534}$$

$$\frac{1}{4} \left(\frac{\int \frac{(-ab(5Ab+4aB) \cos^2(c+dx)+2ab(aA+4bB) \cos(c+dx)+a(4Aa^2+12bBa+3Ab^2)) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{a} + \frac{(4aB + 5Ab) \tan(c + dx) \sqrt{a+b \cos(c+dx)}}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 27

$$\frac{1}{4} \left(\frac{\int \frac{(-ab(5Ab+4aB) \cos^2(c+dx)+2ab(aA+4bB) \cos(c+dx)+a(4Aa^2+12bBa+3Ab^2)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} + \frac{(4aB + 5Ab) \tan(c + dx) \sqrt{a+b \cos(c+dx)}}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{\int \frac{-ab(5Ab+4aB) \sin(c+dx+\frac{\pi}{2})^2+2ab(aA+4bB) \sin(c+dx+\frac{\pi}{2})+a(4Aa^2+12bBa+3Ab^2)}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2a} + \frac{(4aB + 5Ab) \tan(c + dx) \sqrt{a+b \cos(c+dx)}}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3538

$$\frac{1}{4} \left(\frac{\int -\frac{(ab(4Aa^2+12bBa+3Ab^2)+ab(4Ba^2+7Aba+8b^2B) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - a(4aB + 5Ab) \int \sqrt{a + b \cos(c + dx)} dx + \frac{(4aB + 5Ab) \tan(c + dx) \sqrt{a+b \cos(c+dx)}}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 25

$$\frac{1}{4} \left(\frac{\int \frac{(ab(4Aa^2+12bBa+3Ab^2)+ab(4Ba^2+7Aba+8b^2B) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - a(4aB + 5Ab) \int \sqrt{a + b \cos(c + dx)} dx + \frac{(4aB + 5Ab) \tan(c + dx) \sqrt{a+b \cos(c+dx)}}{d} \right)$$

$$\frac{aA \tan(c + dx) \sec(c + dx) \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{\int \frac{ab(4Aa^2+12bBa+3Ab^2)+ab(4Ba^2+7Aba+8b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - a(4aB+5Ab) \int \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} dx \right) + (4aB+5Ab)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3134

$$\frac{1}{4} \left(\frac{\int \frac{ab(4Aa^2+12bBa+3Ab^2)+ab(4Ba^2+7Aba+8b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{a(4aB+5Ab)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + (4aB+5Ab)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{\int \frac{ab(4Aa^2+12bBa+3Ab^2)+ab(4Ba^2+7Aba+8b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{a(4aB+5Ab)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + (4aB+5Ab)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3132

$$\frac{1}{4} \left(\frac{\int \frac{ab(4Aa^2+12bBa+3Ab^2)+ab(4Ba^2+7Aba+8b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2a} - \frac{2a(4aB+5Ab)\sqrt{a+b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) + (4aB+5Ab)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3481

$$\frac{1}{4} \left(\frac{ab(4a^2B+7aAb+8b^2B) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx + ab(4a^2A+12abB+3Ab^2) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2a(4aB+5Ab)\sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{ab(4a^2B+7aAb+8b^2B) \int \frac{1}{\sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)}} dx + ab(4a^2A+12abB+3Ab^2) \int \frac{1}{\sin\left(c+dx+\frac{\pi}{2}\right)\sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)}} dx - \frac{2a(4aB+5Ab)\sqrt{a+b \cos(c+dx)}}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3142

$$\frac{1}{4} \left(\frac{ab(4a^2A+12abB+3Ab^2) \int \frac{1}{\sin\left(c+dx+\frac{\pi}{2}\right)\sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{ab(4a^2B+7aAb+8b^2B)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} - \frac{2a(4aB+5Ab)\sqrt{a+b \cos(c+dx)}}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{ab(4a^2A+12abB+3Ab^2) \int \frac{1}{\sin\left(c+dx+\frac{\pi}{2}\right)\sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{ab(4a^2B+7aAb+8b^2B)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin\left(c+dx+\frac{\pi}{2}\right)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} - \frac{2a(4aB+5Ab)\sqrt{a+b \cos(c+dx)}}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d}$$

↓ 3140

$$\frac{1}{4} \left(\frac{ab(4a^2A+12abB+3Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2ab(4a^2B+7aAb+8b^2B) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2a(4aB+5a^2)}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3286

$$\frac{1}{4} \left(\frac{ab(4a^2A+12abB+3Ab^2) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2ab(4a^2B+7aAb+8b^2B) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2a(4aB+5a^2)}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{ab(4a^2A+12abB+3Ab^2) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2ab(4a^2B+7aAb+8b^2B) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2a(4aB+5a^2)}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) \sqrt{a+b\cos(c+dx)}}{2d}$$

↓ 3284

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

rule 3538 `Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.309.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1402 vs. $2(356) = 712$.

Time = 14.72 (sec) , antiderivative size = 1403, normalized size of antiderivative = 4.76

method	result	size
default	Expression too large to display	1403
parts	Expression too large to display	1722

input `int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNV
ERBOSE)`

output `-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b^2*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+2*A*a^2*(-1/2*cos(1/2*d*x+1/2*c)/a*(-2*s
in(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2
*c)^2-1)^2+3/4*b/a^2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*s
in(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)-1/8*b/a*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*
d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2
*c),(-2*b/(a-b))^(1/2))+3/8/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d
*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x
+1/2*c)^2)^(1/2)*b*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-3/8*b^
2/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(
1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-1/2*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(
a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-
b))^(1/2))-3/8/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2
+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))*b^2)-2*b*(A*b...`

3.309.5 Fricas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorith
m="fricas")`

output `Timed out`

3.309.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Timed out`

3.309.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^3 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^3, x)`

3.309.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^3 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^3, x)`

3.309.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^3} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^3,x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^3, x)`

3.310 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.310.1 Optimal result	2859
3.310.2 Mathematica [C] (verified)	2860
3.310.3 Rubi [A] (verified)	2861
3.310.4 Maple [B] (verified)	2869
3.310.5 Fricas [F(-1)]	2870
3.310.6 Sympy [F(-1)]	2871
3.310.7 Maxima [F]	2871
3.310.8 Giac [F]	2871
3.310.9 Mupad [F(-1)]	2872

3.310.1 Optimal result

Integrand size = 33, antiderivative size = 375

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx =$$

$$\frac{(16a^2A + 3Ab^2 + 30abB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{24ad \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} +$$

$$\frac{(16a^2A + 17Ab^2 + 42abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{24d \sqrt{a + b \cos(c + dx)}} +$$

$$\frac{(12a^2Ab - Ab^3 + 8a^3B + 6ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{8ad \sqrt{a + b \cos(c + dx)}} +$$

$$\frac{(16a^2A + 3Ab^2 + 30abB) \sqrt{a + b \cos(c + dx)} \tan(c + dx)}{24ad} +$$

$$\frac{(7Ab + 6aB) \sqrt{a + b \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{12d} +$$

$$\frac{aA \sqrt{a + b \cos(c + dx)} \sec^2(c + dx) \tan(c + dx)}{3d}$$

output
$$\begin{aligned} & -1/24*(16*A*a^2+3*A*b^2+30*B*a*b)*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x \\ & +1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)}*(b/(a+b))^{(1/2)})*(a+b*\cos(d*x \\ & +c))^{(1/2)}/a/d/((a+b*\cos(d*x+c))/(a+b))^{(1/2)}+1/24*(16*A*a^2+17*A*b^2+42*B \\ & *a*b)*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d* \\ & x+1/2*c), 2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*\cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*co \\ & s(d*x+c))^{(1/2)}+1/8*(12*A*a^2*b-A*b^3+8*B*a^3+6*B*a*b^2)*(\cos(1/2*d*x+1/2* \\ & c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticPi}(\sin(1/2*d*x+1/2*c), 2, 2^{(1/2)}*(b/ \\ & (a+b))^{(1/2)})*((a+b*\cos(d*x+c))/(a+b))^{(1/2)}/a/d/(a+b*\cos(d*x+c))^{(1/2)}+1/ \\ & 24*(16*A*a^2+3*A*b^2+30*B*a*b)*(a+b*\cos(d*x+c))^{(1/2)}*\tan(d*x+c)/a/d+1/12* \\ & (7*A*b+6*B*a)*\sec(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}*\tan(d*x+c)/d+1/3*a*A*\sec(d \\ & *x+c)^2*(a+b*\cos(d*x+c))^{(1/2)}*\tan(d*x+c)/d \end{aligned}$$

3.310.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.12 (sec) , antiderivative size = 483, normalized size of antiderivative = 1.29

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{8b(7Ab+6aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(56a^2Ab-9Ab^3+48a^3B+6ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(c+dx)\right)}{a\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output
$$\begin{aligned} & ((8*b*(7*A*b + 6*a*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticF}[(c + d* \\ & x)/2, (2*b)/(a + b)])/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(56*a^2*A*b - 9*A*b^3 \\ & + 48*a^3*B + 6*a*b^2*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticPi}[2, (\\ & c + d*x)/2, (2*b)/(a + b)])/(a*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) - ((2*I)*(16*a^2* \\ & A + 3*A*b^2 + 30*a*b*B)*\text{Sqrt}[-((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[(b*(\\ & 1 + \text{Cos}[c + d*x]))/(-a + b)]*\text{Csc}[c + d*x]*(-2*a*(a - b)*\text{EllipticE}[I*\text{ArcSin} \\ & h[\text{Sqrt}[-(a + b)^{-1}]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(-2* \\ & a*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + \\ & b)/(a - b)] + b*\text{EllipticPi}[(a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]*\text{Sqrt}[a \\ & + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)])))/(a^2*b*\text{Sqrt}[-(a + b)^{-1}]) + (4* \\ & \text{Sqrt}[a + b*\text{Cos}[c + d*x]]*\text{Sec}[c + d*x]^2*(2*a*(7*A*b + 6*a*B)*\text{Sin}[c + d*x] \\ & + (8*a^2*A + (3*A*b^2)/2 + 15*a*b*B)*\text{Sin}[2*(c + d*x)] + 8*a^2*A*\text{Tan}[c + d* \\ & x]))/a)/(96*d) \end{aligned}$$

3.310. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx$

3.310.3 Rubi [A] (verified)

Time = 3.35 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.04, number of steps used = 24, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 3468, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c+dx)(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^4} dx$$

$$\downarrow \text{3468}$$

$$\frac{1}{3} \int \frac{(3b(aA+2bB)\cos^2(c+dx) + 2(2Aa^2+6bBa+3Ab^2)\cos(c+dx) + a(7Ab+6aB))\sec^3(c+dx)}{2\sqrt{a+b\cos(c+dx)} + \frac{aA\tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}} dx +$$

$$\downarrow \text{27}$$

$$\frac{1}{6} \int \frac{(3b(aA+2bB)\cos^2(c+dx) + 2(2Aa^2+6bBa+3Ab^2)\cos(c+dx) + a(7Ab+6aB))\sec^3(c+dx)}{\sqrt{a+b\cos(c+dx)} + \frac{aA\tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}} dx +$$

$$\downarrow \text{3042}$$

$$\frac{1}{6} \int \frac{3b(aA+2bB)\sin(c+dx+\frac{\pi}{2})^2 + 2(2Aa^2+6bBa+3Ab^2)\sin(c+dx+\frac{\pi}{2}) + a(7Ab+6aB)}{\sin(c+dx+\frac{\pi}{2})^3\sqrt{a+b\sin(c+dx+\frac{\pi}{2})} + \frac{aA\tan(c+dx)\sec^2(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}} dx +$$

$$\downarrow \text{3534}$$

$$\frac{1}{6} \left(\frac{\int \frac{(ab(7Ab+6aB) \cos^2(c+dx) + 2a(6Ba^2+13Aba+12b^2B) \cos(c+dx) + a(16Aa^2+30bBa+3Ab^2)) \sec^2(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{2a} + \frac{(6aB + 7Ab) \tan(c+dx)}{d} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{\int \frac{(ab(7Ab+6aB) \cos^2(c+dx) + 2a(6Ba^2+13Aba+12b^2B) \cos(c+dx) + a(16Aa^2+30bBa+3Ab^2)) \sec^2(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{4a} + \frac{(6aB + 7Ab) \tan(c+dx)}{d} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\int \frac{ab(7Ab+6aB) \sin(c+dx+\frac{\pi}{2})^2 + 2a(6Ba^2+13Aba+12b^2B) \sin(c+dx+\frac{\pi}{2}) + a(16Aa^2+30bBa+3Ab^2)}{\sin(c+dx+\frac{\pi}{2})^2 \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{4a} + \frac{(6aB + 7Ab) \tan(c+dx)}{d} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3534

$$\frac{1}{6} \left(\frac{\int \frac{(2b(7Ab+6aB) \cos(c+dx)a^2 - b(16Aa^2+30bBa+3Ab^2) \cos^2(c+dx)a + 3(8Ba^3+12Aba^2+6b^2Ba-Ab^3)a) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{a} + \frac{(16a^2A+30abB+3Ab^2) \tan(c+dx)}{d} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{\int \frac{(2b(7Ab+6aB) \cos(c+dx)a^2 - b(16Aa^2+30bBa+3Ab^2) \cos^2(c+dx)a + 3(8Ba^3+12Aba^2+6b^2Ba-Ab^3)a) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} + \frac{(16a^2A+30abB+3Ab^2) \tan(c+dx)}{d} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\int \frac{2b(7Ab+6aB) \sin(c+dx+\frac{\pi}{2})a^2 - b(16Aa^2+30bBa+3Ab^2) \sin(c+dx+\frac{\pi}{2})^2 a + 3(8Ba^3+12Aba^2+6b^2Ba-Ab^3)a}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2a} + \frac{(16a^2A+30abB+3Ab^2) \tan(c+dx)}{d} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3538

$$\frac{1}{6} \left(\frac{-a(16a^2A+30abB+3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx - \frac{\int -\frac{(b(16Aa^2+42bBa+17Ab^2) \cos(c+dx)a^2 + 3b(8Ba^3+12Aba^2+6b^2Ba-Ab^3)a) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a}}{4a} \right) +$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 25

$$\frac{1}{6} \left(\frac{\frac{\int \frac{(b(16Aa^2+42bBa+17Ab^2) \cos(c+dx)a^2 + 3b(8Ba^3+12Aba^2+6b^2Ba-Ab^3)a) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - a(16a^2A+30abB+3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx}{4a} \right) + (16a^2A+30abB+3Ab^2) \int \sqrt{a+b \cos(c+dx)} dx$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\frac{\int \frac{b(16Aa^2+42bBa+17Ab^2) \sin(c+dx+\frac{\pi}{2})a^2 + 3b(8Ba^3+12Aba^2+6b^2Ba-Ab^3)a}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2a} - a(16a^2A+30abB+3Ab^2) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{4a} \right) + (16a^2A+30abB+3Ab^2) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3134

$$\frac{1}{6} \left(\frac{\int \frac{b(16Aa^2+42bBa+17Ab^2) \sin(c+dx+\frac{\pi}{2}) a^2+3b(8Ba^3+12Aba^2+6b^2Ba-Ab^3) a}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} } dx}{2a} - \frac{a(16a^2A+30abB+3Ab^2) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\int \frac{b(16Aa^2+42bBa+17Ab^2) \sin(c+dx+\frac{\pi}{2}) a^2+3b(8Ba^3+12Aba^2+6b^2Ba-Ab^3) a}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} } dx}{2a} - \frac{a(16a^2A+30abB+3Ab^2) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3132

$$\frac{1}{6} \left(\frac{\int \frac{b(16Aa^2+42bBa+17Ab^2) \sin(c+dx+\frac{\pi}{2}) a^2+3b(8Ba^3+12Aba^2+6b^2Ba-Ab^3) a}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} } dx}{2a} - \frac{2a(16a^2A+30abB+3Ab^2) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) + (1)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3481

$$\frac{1}{6} \left(\frac{a^2b(16a^2A+42abB+17Ab^2) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx + 3ab(8a^3B+12a^2Ab+6ab^2B-Ab^3) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} - \frac{2a(16a^2A+30abB+3Ab^2) \sqrt{a+b \cos(c+dx)} E}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

3d

3.310. $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^4(c+dx) dx$

$$\begin{array}{c} \downarrow 3042 \\ \left(\frac{1}{6} \right) \left(\frac{a^2 b (16a^2 A + 42abB + 17Ab^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + 3ab(8a^3 B + 12a^2 Ab + 6ab^2 B - Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2a(16a^2 A + 30abB + 30a^2 B)}{2a} - \frac{4a}{4a} \right) \end{array}$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$\downarrow 3142$

$$\left(\frac{1}{6} \right) \left(\frac{a^2 b (16a^2 A + 42abB + 17Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + 3ab(8a^3 B + 12a^2 Ab + 6ab^2 B - Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2a(16a^2 A + 30abB + 30a^2 B)}{2a} - \frac{4a}{4a} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$\downarrow 3042$

$$\left(\frac{1}{6} \right) \left(\frac{a^2 b (16a^2 A + 42abB + 17Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + 3ab(8a^3 B + 12a^2 Ab + 6ab^2 B - Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2a(16a^2 A + 30abB + 30a^2 B)}{2a} - \frac{4a}{4a} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

$\downarrow 3140$

$$\frac{1}{6} \left(\frac{3ab(8a^3B+12a^2Ab+6ab^2B-Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a^2b(16a^2A+42abB+17Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{2a} - \frac{2a}{4a} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3286

$$\frac{1}{6} \left(\frac{3ab(8a^3B+12a^2Ab+6ab^2B-Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2a^2b(16a^2A+42abB+17Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{2a} - \frac{2a}{4a} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3ab(8a^3B+12a^2Ab+6ab^2B-Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2a^2b(16a^2A+42abB+17Ab^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{2a} - \frac{2a}{4a} \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3284

$$\frac{1}{6} \left(\frac{(16a^2A + 30abB + 3Ab^2) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{d} + \frac{2a^2b(16a^2A + 42abB + 17Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{6ab(8a^3B + 12a^2A)}{b}}{d\sqrt{a+b \cos(c+dx)}} \right) + \frac{aA \tan(c+dx) \sec^2(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

input `Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output `(a*A*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]^2*Tan[c + d*x])/(3*d) + (((7*A*b + 6*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*d) + (((-2*a*(16*a^2*A + 3*A*b^2 + 30*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*a^2*b*(16*a^2*A + 17*A*b^2 + 42*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*a*b*(12*a^2*A*b - A*b^3 + 8*a^3*B + 6*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])))/b)/(2*a) + ((16*a^2*A + 3*A*b^2 + 30*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/d)/(4*a))/6`

3.310.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.310.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2326 vs. $2(432) = 864$.

Time = 18.46 (sec) , antiderivative size = 2327, normalized size of antiderivative = 6.21

method	result	size
default	Expression too large to display	2327
parts	Expression too large to display	2727

input `int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNV
ERBOSE)`

output `-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*a^2*(-
1/3*cos(1/2*d*x+1/2*c)/a*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*
c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3+5/12*b/a^2*cos(1/2*d*x+1/2*c)*(-2
*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1
/2*c)^2-1)^2-1/24*(16*a^2+15*b^2)/a^3*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1
/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)+5/4
8*b^2/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-
b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*Ell
ipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+1/3*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4
*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a
-b))^(1/2))-1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-
b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/
2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+1/3/a*(sin(1/2*d*x+1/2
c)^2)^(1/2)((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+
1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*EllipticE(cos(1/2*d*x+1/2*c
,(-2*b/(a-b))^(1/2))-5/16*b^2/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(
1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/
2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+5/1
6/a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b...`

3.310.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorith
m="fracas")`

output `Timed out`

3.310.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.310.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^4 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^4, x)`

3.310.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^4 dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^4, x)`

3.310.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^4} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^4,x)`output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^4, x)`

3.311 $\int \cos^2(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.311.1 Optimal result	2873
3.311.2 Mathematica [A] (verified)	2874
3.311.3 Rubi [A] (verified)	2875
3.311.4 Maple [B] (verified)	2881
3.311.5 Fricas [C] (verification not implemented)	2882
3.311.6 Sympy [F(-1)]	2883
3.311.7 Maxima [F]	2884
3.311.8 Giac [F]	2884
3.311.9 Mupad [F(-1)]	2884

3.311.1 Optimal result

Integrand size = 33, antiderivative size = 462

$$\int \cos^2(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx =$$

$$\frac{2(110a^4Ab - 3069a^2Ab^3 - 1617Ab^5 - 40a^5B - 255a^3b^2B - 3705ab^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) + 3465b^3d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{2(a^2 - b^2)(110a^3Ab - 1254aAb^3 - 40a^4B - 285a^2b^2B - 675b^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + 3465b^3d \sqrt{a+b \cos(c+dx)}} + \frac{2(110a^3Ab - 1254aAb^3 - 40a^4B - 285a^2b^2B - 675b^4B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3465b^2d} - \frac{2(110a^2Ab - 539Ab^3 - 40a^3B - 335ab^2B)(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{3465b^2d} - \frac{2(22aAb - 8a^2B - 81b^2B)(a+b \cos(c+dx))^{5/2} \sin(c+dx)}{693b^2d} + \frac{2(11Ab - 4aB)(a+b \cos(c+dx))^{7/2} \sin(c+dx)}{99b^2d} + \frac{2B \cos(c+dx)(a+b \cos(c+dx))^{7/2} \sin(c+dx)}{11bd}$$

output
$$\begin{aligned} & -2/3465*(110*A*a^2*b-539*A*b^3-40*B*a^3-335*B*a*b^2)*(a+b*\cos(d*x+c))^{3/2} \\ &)*\sin(d*x+c)/b^2/d-2/693*(22*A*a*b-8*B*a^2-81*B*b^2)*(a+b*\cos(d*x+c))^{5/2} \\ &)*\sin(d*x+c)/b^2/d+2/99*(11*A*b-4*B*a)*(a+b*\cos(d*x+c))^{7/2}*\sin(d*x+c)/b \\ & ^2/d+2/11*B*\cos(d*x+c)*(a+b*\cos(d*x+c))^{7/2}*\sin(d*x+c)/b-2/3465*(110*A \\ & *a^3*b-1254*A*a*b^3-40*B*a^4-285*B*a^2*b^2-675*B*b^4)*\sin(d*x+c)*(a+b*\cos(\\ & d*x+c))^{1/2}/b^2/d-2/3465*(110*A*a^4*b-3069*A*a^2*b^3-1617*A*b^5-40*B*a^5 \\ & -255*B*a^3*b^2-3705*B*a*b^4)*(cos(1/2*d*x+1/2*c))^{1/2}/cos(1/2*d*x+1/2* \\ & c)*EllipticE(sin(1/2*d*x+1/2*c),2^{1/2}*(b/(a+b))^{1/2})*(a+b*\cos(d*x+c))^{ \\ & (1/2)}/b^3/d/((a+b*\cos(d*x+c))/(a+b))^{1/2}+2/3465*(a^2-b^2)*(110*A*a^3*b-1 \\ & 254*A*a*b^3-40*B*a^4-285*B*a^2*b^2-675*B*b^4)*(cos(1/2*d*x+1/2*c))^{1/2} \\ & /cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{1/2}*(b/(a+b))^{1/2})* \\ & ((a+b*\cos(d*x+c))/(a+b))^{1/2}/b^3/d/(a+b*\cos(d*x+c))^{1/2} \end{aligned}$$

3.311.2 Mathematica [A] (verified)

Time = 2.63 (sec) , antiderivative size = 357, normalized size of antiderivative = 0.77

$$\int \cos^2(c+dx)(a+b\cos(c+dx))^{5/2}(A + B\cos(c+dx)) dx = \frac{16\sqrt{\frac{a+b\cos(c+dx)}{a+b}}(b^2(1705a^3Ab + 2871aAb^3 + 10a^4B + 3315a^2b^2B + 675b^4B) \text{EllipticF}$$

input `Integrate[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output
$$\begin{aligned} & (16*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*(b^2*(1705*a^3*A*b + 2871*a*A*b^3 + \\ & 10*a^4*B + 3315*a^2*b^2*B + 675*b^4*B)*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + \\ & b)] + (-110*a^4*A*b + 3069*a^2*A*b^3 + 1617*A*b^5 + 40*a^5*B + 255*a^3*b^2 \\ & *B + 3705*a*b^4*B)*((a + b)*\text{EllipticE}[(c + d*x)/2, (2*b)/(a + b)] - a*\text{Elli \\ & pticF}[(c + d*x)/2, (2*b)/(a + b)])) + b*(a + b*\text{Cos}[c + d*x])*((880*a^3*A*b \\ & + 32868*a*A*b^3 - 320*a^4*B + 18660*a^2*b^2*B + 13050*b^4*B)*\text{Sin}[c + d*x] \\ & + b*(4*(1650*a^2*A*b + 1463*A*b^3 + 30*a^3*B + 3095*a*b^2*B)*\text{Sin}[2*(c + d \\ & *x)] + 5*b*((836*a*A*b + 452*a^2*B + 513*b^2*B)*\text{Sin}[3*(c + d*x)] + 7*b*((2 \\ & 2*A*b + 46*a*B)*\text{Sin}[4*(c + d*x)] + 9*b*B*\text{Sin}[5*(c + d*x)])))))/(27720*b^3* \\ & d*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) \end{aligned}$$

3.311.3 Rubi [A] (verified)

Time = 2.57 (sec) , antiderivative size = 480, normalized size of antiderivative = 1.04, number of steps used = 24, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c+dx+\frac{\pi}{2}\right)^2 \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3469} \\
 & \frac{2 \int \frac{1}{2}(a+b\cos(c+dx))^{5/2} ((11Ab-4aB)\cos^2(c+dx)+9bB\cos(c+dx)+2aB) dx}{\frac{11b}{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{7/2}}} + \\
 & \quad \downarrow \text{27} \\
 & \frac{\int (a+b\cos(c+dx))^{5/2} ((11Ab-4aB)\cos^2(c+dx)+9bB\cos(c+dx)+2aB) dx}{\frac{11b}{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{7/2}}} + \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (a+b\sin(c+dx+\frac{\pi}{2}))^{5/2} ((11Ab-4aB)\sin(c+dx+\frac{\pi}{2})^2+9bB\sin(c+dx+\frac{\pi}{2})+2aB) dx}{\frac{11b}{2B\sin(c+dx)\cos(c+dx)(a+b\cos(c+dx))^{7/2}}} + \\
 & \quad \downarrow \text{3502} \\
 & \frac{2 \int \frac{1}{2}(a+b\cos(c+dx))^{5/2} (b(77Ab-10aB)-(-8Ba^2+22Aba-81b^2B)\cos(c+dx)) dx}{9b} + \frac{2(11Ab-4aB)\sin(c+dx)(a+b\cos(c+dx))^{7/2}}{9bd} + \\
 & \quad \downarrow \text{27} \\
 & \frac{2 \int \frac{1}{2}(a+b\cos(c+dx))^{5/2} (b(77Ab-10aB)-(-8Ba^2+22Aba-81b^2B)\cos(c+dx)) dx}{9b} + \frac{2(11Ab-4aB)\sin(c+dx)(a+b\cos(c+dx))^{7/2}}{9bd} + \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.311. $\int \cos^2(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx$

$$\frac{\int (a+b \cos(c+dx))^{5/2} (b(77Ab-10aB) - (-8Ba^2+22Aba-81b^2B) \cos(c+dx)) dx}{9b} + \frac{2(11Ab-4aB) \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd} +$$

$$\frac{11b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}$$

$$\frac{11bd}{11bd} \downarrow 3042$$

$$\frac{\int (a+b \sin(c+dx+\frac{\pi}{2}))^{5/2} (b(77Ab-10aB) + (8Ba^2-22Aba+81b^2B) \sin(c+dx+\frac{\pi}{2})) dx}{9b} + \frac{2(11Ab-4aB) \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd} +$$

$$\frac{11b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}$$

$$\frac{11bd}{11bd} \downarrow 3232$$

$$\frac{\frac{2}{7} \int \frac{1}{2} (a+b \cos(c+dx))^{3/2} (3b(-10Ba^2+143Aba+135b^2B) - (-40Ba^3+110Aba^2-335b^2Ba-539Ab^3) \cos(c+dx)) dx - \frac{2(-8a^2B+22aAb-81b^2B) \sin(c+dx)}{7d}}{9b} + \frac{11b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}$$

$$\frac{11bd}{11bd} \downarrow 27$$

$$\frac{\frac{1}{7} \int (a+b \cos(c+dx))^{3/2} (3b(-10Ba^2+143Aba+135b^2B) - (-40Ba^3+110Aba^2-335b^2Ba-539Ab^3) \cos(c+dx)) dx - \frac{2(-8a^2B+22aAb-81b^2B) \sin(c+dx)}{7d}}{9b} + \frac{11b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}$$

$$\frac{11bd}{11bd} \downarrow 3042$$

$$\frac{\frac{1}{7} \int (a+b \sin(c+dx+\frac{\pi}{2}))^{3/2} (3b(-10Ba^2+143Aba+135b^2B) + (40Ba^3-110Aba^2+335b^2Ba+539Ab^3) \sin(c+dx+\frac{\pi}{2})) dx - \frac{2(-8a^2B+22aAb-81b^2B) \sin(c+dx)}{7d}}{9b} + \frac{11b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}$$

$$\frac{11bd}{11bd} \downarrow 3232$$

$$\frac{\frac{1}{7} \left(\frac{2}{5} \int \frac{3}{2} \sqrt{a+b \cos(c+dx)} (b(-10Ba^3+605Aba^2+1010b^2Ba+539Ab^3) - (-40Ba^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \cos(c+dx)) dx - \frac{2(-8a^2B+22aAb-81b^2B) \sin(c+dx)}{7d} \right)}{9b} + \frac{11b}{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}$$

$$\frac{11bd}{11bd} \downarrow 27$$

3.311. $\int \cos^2(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

$$\frac{1}{7} \left(\frac{3}{5} \int \sqrt{a+b \cos(c+dx)} (b(-10Ba^3+605Aba^2+1010b^2Ba+539Ab^3) - (-40Ba^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \cos(c+dx)) dx - \frac{2(-40a^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \sin(c+dx)}{9b} \right)$$

$$\frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}{11bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (b(-10Ba^3+605Aba^2+1010b^2Ba+539Ab^3) + (40Ba^4-110Aba^3+285b^2Ba^2+1254Ab^3a+675b^4B) \sin(c+dx+\frac{\pi}{2})) dx - \frac{2(-40a^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \cos(c+dx+\frac{\pi}{2})}{9b} \right)$$

$$\frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}{11bd}$$

↓ 3232

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{2}{3} \int \frac{b(10Ba^4+1705Aba^3+3315b^2Ba^2+2871Ab^3a+675b^4B) - (-40Ba^5+110Aba^4-255b^2Ba^3-3069Ab^3a^2-3705b^4Ba-1617Ab^5) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-40a^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \sin(c+dx)}{9b} \right) \right)$$

$$\frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}{11bd}$$

↓ 27

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \int \frac{b(10Ba^4+1705Aba^3+3315b^2Ba^2+2871Ab^3a+675b^4B) - (-40Ba^5+110Aba^4-255b^2Ba^3-3069Ab^3a^2-3705b^4Ba-1617Ab^5) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-40a^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \sin(c+dx)}{9b} \right) \right)$$

$$\frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}{11bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \int \frac{b(10Ba^4+1705Aba^3+3315b^2Ba^2+2871Ab^3a+675b^4B) + (40Ba^5-110Aba^4+255b^2Ba^3+3069Ab^3a^2+3705b^4Ba+1617Ab^5) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-40a^4+110Aba^3-285b^2Ba^2-1254Ab^3a-675b^4B) \cos(c+dx+\frac{\pi}{2})}{9b} \right) \right)$$

$$\frac{2B \sin(c+dx) \cos(c+dx)(a+b \cos(c+dx))^{7/2}}{11bd}$$

↓ 3231

3.311. $\int \cos^2(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B)}{b} \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx - \frac{(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2Ab^3 - 3705ab^4B - 161b^5)}{b} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2Ab^3 - 3705ab^4B - 161b^5)}{b} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3134

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2Ab^3 - 3705ab^4B - 161b^5)}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2Ab^3 - 3705ab^4B - 161b^5)}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3132

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B)}{b} \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2Ab^3 - 3705ab^4B - 161b^5)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3142

3.311. $\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - 2(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2b^3 - 1617Ab^5 - 40a^5B - 255a^3b^2B - 3705a^2b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{a+b \cos(c+dx)}} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx - 2(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2b^3 - 1617Ab^5 - 40a^5B - 255a^3b^2B - 3705a^2b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{a+b \cos(c+dx)}} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

↓ 3140

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{2(a^2 - b^2)(-40a^4B + 110a^3Ab - 285a^2b^2B - 1254aAb^3 - 675b^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2(-40a^5B + 110a^4Ab - 255a^3b^2B - 3069a^2b^3 - 1617Ab^5 - 40a^5B - 255a^3b^2B - 3705a^2b^4B) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{a+b \cos(c+dx)}} \right) \right) \right)$$

$$\frac{2B \sin(c + dx) \cos(c + dx)(a + b \cos(c + dx))^{7/2}}{11bd}$$

input `Int[Cos[c + d*x]^2*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*Cos[c + d*x]*(a + b*Cos[c + d*x])^(7/2)*Sin[c + d*x]/(11*b*d) + ((2*(11*A*b - 4*a*B)*(a + b*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b*d) + ((-2*(22*a*A*b - 8*a^2*B - 81*b^2*B)*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*d) + ((-2*(110*a^2*A*b - 539*A*b^3 - 40*a^3*B - 335*a*b^2*B)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (3*((-2*(110*a^4*A*b - 3069*a^2*A*b^3 - 1617*A*b^5 - 40*a^5*B - 255*a^3*b^2*B - 3705*a*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(a^2 - b^2)*(110*a^3*A*b - 1254*a*A*b^3 - 40*a^4*B - 285*a^2*b^2*B - 675*b^4*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 - (2*(110*a^3*A*b - 1254*a*A*b^3 - 40*a^4*B - 285*a^2*b^2*B - 675*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5)/7)/(9*b))/(11*b)`

3.311. $\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

3.311.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*SIN[c + d*x]]/Sqrt[(a + b*SIN[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*SIN[c + d*x])/(a + b)]/Sqrt[a + b*SIN[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*SIN[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*SIN[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

```
rule 3232 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*
d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x] /; FreeQ
[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m,
0] && IntegerQ[2*m]
```

```
rule 3469 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.311.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1982 vs. $2(488) = 976$.

Time = 27.78 (sec) , antiderivative size = 1983, normalized size of antiderivative = 4.29

method	result	size
default	Expression too large to display	1983
parts	Expression too large to display	2137

```
input int(cos(d*x+c)^2*(a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.311. \quad \int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

output

```
-2/3465*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(675*B
*b^6*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(
a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-40*B*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*E
llipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^6-1617*A*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*Elliptic
E(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^6+40*B*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2
*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^6-255*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2
*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2
*c),(-2*b/(a-b))^(1/2))*a^3*b^3+3705*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/
(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))*a^2*b^4+1254*A*a*b^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2
*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2
*c),(-2*b/(a-b))^(1/2))-110*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin
(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a
-b))^(1/2))*a^5*b-3705*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*
d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))*a*b^5+20160*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^12*b^6+(22880*A
*a*b^5+24640*A*b^6+21920*B*a^2*b^4+71680*B*a*b^5+56880*B*b^6)*sin(1/2*d...
```

3.311.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.18 (sec) , antiderivative size = 726, normalized size of antiderivative = 1.57

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(80i Ba^6 - 220i Aa^5b + 480i Ba^4b^2 + 1023i Aa^3b^3 - 2535i Ba^2b^4 - 5379i Aab^5 - 20160i Ab^6 + 21920i B a^2 b^4 + 71680i B a b^5 + 56880i B b^6) \sin(1/2 d x + 1/2 c) + \dots}{\dots}$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

```

output 1/10395*(sqrt(2)*(80*I*B*a^6 - 220*I*A*a^5*b + 480*I*B*a^4*b^2 + 1023*I*A*
a^3*b^3 - 2535*I*B*a^2*b^4 - 5379*I*A*a*b^5 - 2025*I*B*b^6)*sqrt(b)*weiers
trassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3
*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(-80*I*B*a^6 + 22
0*I*A*a^5*b - 480*I*B*a^4*b^2 - 1023*I*A*a^3*b^3 + 2535*I*B*a^2*b^4 + 5379
*I*A*a*b^5 + 2025*I*B*b^6)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)
/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x +
c) + 2*a)/b) - 3*sqrt(2)*(-40*I*B*a^5*b + 110*I*A*a^4*b^2 - 255*I*B*a^3*b
^3 - 3069*I*A*a^2*b^4 - 3705*I*B*a*b^5 - 1617*I*A*b^6)*sqrt(b)*weierstrass
Zeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInv
erse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*
x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(40*I*B*a^5*b - 110*I*A
*a^4*b^2 + 255*I*B*a^3*b^3 + 3069*I*A*a^2*b^4 + 3705*I*B*a*b^5 + 1617*I*A*
b^6)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b
^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b
^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(315*B*
b^6*cos(d*x + c)^4 - 20*B*a^4*b^2 + 55*A*a^3*b^3 + 1025*B*a^2*b^4 + 1793*A
*a*b^5 + 675*B*b^6 + 35*(23*B*a*b^5 + 11*A*b^6)*cos(d*x + c)^3 + 5*(113*B*
a^2*b^4 + 209*A*a*b^5 + 81*B*b^6)*cos(d*x + c)^2 + (15*B*a^3*b^3 + 825*A*a
^2*b^4 + 1145*B*a*b^5 + 539*A*b^6)*cos(d*x + c))*sqrt(b*cos(d*x + c) + ...

```

3.311.6 Sympy [**F(-1)**]

Timed out.

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**2*(a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.311.7 Maxima [F]

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^2, x)`

3.311.8 Giac [F]

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \cos(dx + c)^2 dx$$

input `integrate(cos(d*x+c)^2*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^2, x)`

3.311.9 Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^2 (A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2),x)`

output `int(cos(c + d*x)^2*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2), x)`

3.311. $\int \cos^2(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

3.312 $\int \cos(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.312.1 Optimal result	2885
3.312.2 Mathematica [A] (verified)	2886
3.312.3 Rubi [A] (verified)	2887
3.312.4 Maple [B] (verified)	2893
3.312.5 Fricas [C] (verification not implemented)	2894
3.312.6 Sympy [F(-1)]	2895
3.312.7 Maxima [F]	2895
3.312.8 Giac [F]	2895
3.312.9 Mupad [F(-1)]	2896

3.312.1 Optimal result

Integrand size = 31, antiderivative size = 372

$$\int \cos(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx = \frac{2(45a^3Ab + 435aAb^3 - 10a^4B + 279a^2b^2B + 147b^4B) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx))}{315b^2d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(a^2 - b^2) (45a^2Ab + 75Ab^3 - 10a^3B + 114ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b})}{315b^2d \sqrt{a+b \cos(c+dx)}} + \frac{2(45a^2Ab + 75Ab^3 - 10a^3B + 114ab^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{315bd} + \frac{2(45aAb - 10a^2B + 49b^2B) (a+b \cos(c+dx))^{3/2} \sin(c+dx)}{315bd} + \frac{2(9Ab - 2aB)(a+b \cos(c+dx))^{5/2} \sin(c+dx)}{63bd} + \frac{2B(a+b \cos(c+dx))^{7/2} \sin(c+dx)}{9bd}$$

output
$$\frac{2}{315}(45A^2b-10B^2a+49B^2b^2)(a+b\cos(dx+c))^{3/2}\sin(dx+c)/b/d+2/63(9Ab-2B^2a)(a+b\cos(dx+c))^{5/2}\sin(dx+c)/b/d+2/9B(a+b\cos(dx+c))^{7/2}\sin(dx+c)/b/d+2/315(45A^2b+75Ab^3-10B^2a^3+114B^2ab^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b/d+2/315(45A^2b+435A^2ab^3-10B^2a^4+279B^2a^2b^2+147B^2b^4)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticE}(\sin(1/2dx+1/2c), 2^{1/2}(b/(a+b))^{1/2})*(a+b\cos(dx+c))^{1/2}/b^2/d/((a+b\cos(dx+c))/(a+b))^{1/2}-2/315(a^2-b^2)(45A^2b+75Ab^3-10B^2a^3+114B^2ab^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticF}(\sin(1/2dx+1/2c), 2^{1/2}(b/(a+b))^{1/2})*((a+b\cos(dx+c))/(a+b))^{1/2}/b^2/d/(a+b\cos(dx+c))^{1/2}$$

3.312.2 Mathematica [A] (verified)

Time = 3.98 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.78

$$\int \cos(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \frac{8\sqrt{\frac{a+b\cos(c+dx)}{a+b}}(b^2(405a^2Ab+75Ab^3+155a^3B+261ab^2B)\text{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b}))}{(1260b^2d\sqrt{a+b\cos(c+dx)})}$$

input `Integrate[Cos[c + d*x]*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output
$$(8*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*(b^2*(405*a^2*A*b + 75*A*b^3 + 155*a^3*B + 261*a*b^2*B)*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)] + (45*a^3*A*b + 435*a*A*b^3 - 10*a^4*B + 279*a^2*b^2*B + 147*b^4*B)*((a + b)*\text{EllipticE}[(c + d*x)/2, (2*b)/(a + b)] - a*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)])) + b*(a + b*\text{Cos}[c + d*x])*(2*(540*a^2*A*b + 345*A*b^3 + 20*a^3*B + 747*a*b^2*B)*\text{Sin}[c + d*x] + b*((540*a*A*b + 300*a^2*B + 266*b^2*B)*\text{Sin}[2*(c + d*x)] + 5*b*(2*(9*A*b + 19*a*B)*\text{Sin}[3*(c + d*x)] + 7*b*B*\text{Sin}[4*(c + d*x)]))))/(1260*b^2*d*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])$$

3.312.3 Rubi [A] (verified)

Time = 1.98 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.03, number of steps used = 23, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.742$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c+dx+\frac{\pi}{2}\right)\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int (a+b\cos(c+dx))^{5/2}(A\cos(c+dx)+B\cos^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2}\left(A\sin\left(c+dx+\frac{\pi}{2}\right)+B\sin\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{2\int\frac{1}{2}(a+b\cos(c+dx))^{5/2}(7bB+(9Ab-2aB)\cos(c+dx))dx}{9b} + \\
 & \quad \frac{2B\sin(c+dx)(a+b\cos(c+dx))^{7/2}}{9bd} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int(a+b\cos(c+dx))^{5/2}(7bB+(9Ab-2aB)\cos(c+dx))dx}{9b} + \frac{2B\sin(c+dx)(a+b\cos(c+dx))^{7/2}}{9bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}(7bB+(9Ab-2aB)\sin(c+dx+\frac{\pi}{2})) dx}{9b} + \\
 & \quad \frac{2B\sin(c+dx)(a+b\cos(c+dx))^{7/2}}{9bd} \\
 & \quad \downarrow \text{3232}
 \end{aligned}$$

$$\frac{\frac{2}{7} \int \frac{1}{2} (a + b \cos(c + dx))^{3/2} (3b(15Ab + 13aB) + (-10Ba^2 + 45Aba + 49b^2B) \cos(c + dx)) dx + \frac{2(9Ab-2aB) \sin(c+dx)}{7}}{\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd}} \downarrow 27$$

$$\frac{\frac{1}{7} \int (a + b \cos(c + dx))^{3/2} (3b(15Ab + 13aB) + (-10Ba^2 + 45Aba + 49b^2B) \cos(c + dx)) dx + \frac{2(9Ab-2aB) \sin(c+dx)}{7d}}{\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd}} \downarrow 3042$$

$$\frac{\frac{1}{7} \int (a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (3b(15Ab + 13aB) + (-10Ba^2 + 45Aba + 49b^2B) \sin(c + dx + \frac{\pi}{2})) dx + \frac{2(9Ab-2aB) \cos(c+dx)}{7}}{\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd}} \downarrow 3232$$

$$\frac{\frac{1}{7} \left(\frac{2}{5} \int \frac{3}{2} \sqrt{a + b \cos(c + dx)} (b(55Ba^2 + 120Aba + 49b^2B) + (-10Ba^3 + 45Aba^2 + 114b^2Ba + 75Ab^3) \cos(c + dx)) dx + \frac{9b}{5} \right)}{\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd}} \downarrow 27$$

$$\frac{\frac{1}{7} \left(\frac{3}{5} \int \sqrt{a + b \cos(c + dx)} (b(55Ba^2 + 120Aba + 49b^2B) + (-10Ba^3 + 45Aba^2 + 114b^2Ba + 75Ab^3) \cos(c + dx)) dx + \frac{9b}{5} \right)}{\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd}} \downarrow 3042$$

$$\frac{\frac{1}{7} \left(\frac{3}{5} \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (b(55Ba^2 + 120Aba + 49b^2B) + (-10Ba^3 + 45Aba^2 + 114b^2Ba + 75Ab^3) \sin(c + dx + \frac{\pi}{2})) dx + \frac{9b}{5} \right)}{\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd}} \downarrow 3232$$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{2}{3} \int \frac{b(155Ba^3+405Aba^2+261b^2Ba+75Ab^3)+(-10Ba^4+45Aba^3+279b^2Ba^2+435Ab^3a+147b^4B) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx + \frac{2(-10a^3B+45a^2Ab+10aAb^2+114ab^2B+75Ab^3)}{b\sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd}$$

↓ 27

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \int \frac{b(155Ba^3+405Aba^2+261b^2Ba+75Ab^3)+(-10Ba^4+45Aba^3+279b^2Ba^2+435Ab^3a+147b^4B) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + \frac{2(-10a^3B+45a^2Ab+10aAb^2+114ab^2B+75Ab^3)}{b\sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \int \frac{b(155Ba^3+405Aba^2+261b^2Ba+75Ab^3)+(-10Ba^4+45Aba^3+279b^2Ba^2+435Ab^3a+147b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(-10a^3B+45a^2Ab+10aAb^2+114ab^2B+75Ab^3)}{b\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd}$$

↓ 3231

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(-10a^4B+45a^3Ab+279a^2b^2B+435aAb^3+147b^4B) \int \sqrt{a+b \cos(c+dx)} dx}{b} - \frac{(a^2-b^2)(-10a^3B+45a^2Ab+114ab^2B+75Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} \right) \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd}$$

↓ 3042

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(-10a^4B+45a^3Ab+279a^2b^2B+435aAb^3+147b^4B) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{(a^2-b^2)(-10a^3B+45a^2Ab+114ab^2B+75Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \right) \right)$$

$$\frac{2B \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd}$$

↓ 3134

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(-10a^4B + 45a^3Ab + 279a^2b^2B + 435aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{(a^2 - b^2)(-10a^3B + 45a^2Ab + 114ab^2B + 147b^3B)}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd} \downarrow 3042$$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{(-10a^4B + 45a^3Ab + 279a^2b^2B + 435aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{(a^2 - b^2)(-10a^3B + 45a^2Ab + 114ab^2B + 147b^3B)}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd} \downarrow 3132$$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{2(-10a^4B + 45a^3Ab + 279a^2b^2B + 435aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{(a^2 - b^2)(-10a^3B + 45a^2Ab + 114ab^2B + 147b^3B)}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd} \downarrow 3142$$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{2(-10a^4B + 45a^3Ab + 279a^2b^2B + 435aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{(a^2 - b^2)(-10a^3B + 45a^2Ab + 114ab^2B + 147b^3B)}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd} \downarrow 3042$$

$$\frac{1}{7} \left(\frac{3}{5} \left(\frac{1}{3} \left(\frac{2(-10a^4B + 45a^3Ab + 279a^2b^2B + 435aAb^3 + 147b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{(a^2 - b^2)(-10a^3B + 45a^2Ab + 114ab^2B + 147b^3B)}{b \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2B \sin(c + dx)(a + b \cos(c + dx))^{7/2}}{9bd} \downarrow 3140$$

3.312. $\int \cos(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

$$\frac{1}{7} \left(\frac{2(-10a^2B+45aAb+49b^2B) \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5d} + \frac{3}{5} \left(\frac{2(-10a^3B+45a^2Ab+114ab^2B+75Ab^3) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3d} + \frac{1}{3} \right) \right) + \frac{2B \sin(c+dx)(a+b \cos(c+dx))^{7/2}}{9bd}$$

input `Int[Cos[c + d*x]*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(2*B*(a + b*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b*d) + ((2*(9*A*b - 2*a*B)*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*d) + ((2*(45*a*A*b - 10*a^2*B + 49*b^2*B)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (3*((2*(45*a^3*A*b + 435*a*A*b^3 - 10*a^4*B + 279*a^2*b^2*B + 147*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(45*a^2*A*b + 75*A*b^3 - 10*a^3*B + 114*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/3 + (2*(45*a^2*A*b + 75*A*b^3 - 10*a^3*B + 114*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d)))/5)/7)/(9*b)`

3.312.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b]*Sin[c + d*x]/Sqrt[(a + b)*Sin[c + d*x]/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.312.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1634 vs. $2(402) = 804$.

Time = 19.43 (sec) , antiderivative size = 1635, normalized size of antiderivative = 4.40

method	result	size
default	Expression too large to display	1635
parts	Expression too large to display	1824

```
input int(cos(d*x+c)*(a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output -2/315*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*
B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b^5+(720*A*b^5+2080*B*a*b^4+224
0*B*b^5)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1440*A*a*b^4-1080*A*b^5
-1360*B*a^2*b^3-3120*B*a*b^4-2072*B*b^5)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+
1/2*c)+(1080*A*a^2*b^3+1440*A*a*b^4+840*A*b^5+320*B*a^3*b^2+1360*B*a^2*b^3
+2408*B*a*b^4+952*B*b^5)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-270*A*a
^3*b^2-540*A*a^2*b^3-510*A*a*b^4-240*A*b^5-10*B*a^4*b-160*B*a^3*b^2-666*B*
a^2*b^3-684*B*a*b^4-168*B*b^5)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+45*
A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b
))^1/2*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2)*a^4*b-45*A*(sin(
1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2
)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2)*a^3*b^2+435*A*(sin(1/2*
d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2*El
lipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2)*a^2*b^3-435*A*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2*Ellipt
icE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2)*a*b^4-45*A*(sin(1/2*d*x+1/2*c)^
2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2*EllipticF(cos
(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2)*a^4*b-30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2*EllipticF(cos(1/2*d*
x+1/2*c),(-2*b/(a-b))^1/2)*a^2*b^3+75*A*b^5*(sin(1/2*d*x+1/2*c)^2)^(1...
```

3.312.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.16 (sec) , antiderivative size = 639, normalized size of antiderivative = 1.72

$$\int \cos(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \frac{\sqrt{2}(-20i Ba^5 + 90i Aa^4b + 93i Ba^3b^2 - 345i Aa^2b^3 - 489i Bab^4 - 225i Ab^5)\sqrt{b}\text{weierst} + B \cos(c + dx))}{}$$

```
input integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")
```

```
output 1/945*(sqrt(2)*(-20*I*B*a^5 + 90*I*A*a^4*b + 93*I*B*a^3*b^2 - 345*I*A*a^2*
b^3 - 489*I*B*a*b^4 - 225*I*A*b^5)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2
- 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*s
in(d*x + c) + 2*a)/b) + sqrt(2)*(20*I*B*a^5 - 90*I*A*a^4*b - 93*I*B*a^3*b^
2 + 345*I*A*a^2*b^3 + 489*I*B*a*b^4 + 225*I*A*b^5)*sqrt(b)*weierstrassPInv
erse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*
x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(10*I*B*a^4*b - 45*I*A*a
^3*b^2 - 279*I*B*a^2*b^3 - 435*I*A*a*b^4 - 147*I*B*b^5)*sqrt(b)*weierstras
sZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPIn
verse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*co
s(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(-10*I*B*a^4*b + 45*I*
A*a^3*b^2 + 279*I*B*a^2*b^3 + 435*I*A*a*b^4 + 147*I*B*b^5)*sqrt(b)*weierst
rassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrass
PInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*co
s(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(35*B*b^5*cos(d*x + c)^3 +
5*B*a^3*b^2 + 135*A*a^2*b^3 + 163*B*a*b^4 + 75*A*b^5 + 5*(19*B*a*b^4 + 9*A
*b^5)*cos(d*x + c)^2 + (75*B*a^2*b^3 + 135*A*a*b^4 + 49*B*b^5)*cos(d*x + c
))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^3*d)
```

3.312.6 Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.312.7 Maxima [F]

$$\int \cos(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*cos(d*x + c), x)`

3.312.8 Giac [F]

$$\int \cos(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \cos(dx + c) dx$$

input `integrate(cos(d*x+c)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*cos(d*x + c), x)`

3.312.9 Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)(A + B \cos(c + dx))(a + b \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2), x)`

3.313 $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$

3.313.1 Optimal result	2897
3.313.2 Mathematica [A] (verified)	2898
3.313.3 Rubi [A] (verified)	2898
3.313.4 Maple [B] (verified)	2903
3.313.5 Fricas [C] (verification not implemented)	2904
3.313.6 Sympy [F(-1)]	2905
3.313.7 Maxima [F]	2905
3.313.8 Giac [F]	2906
3.313.9 Mupad [F(-1)]	2906

3.313.1 Optimal result

Integrand size = 25, antiderivative size = 288

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{2(161a^2Ab + 63Ab^3 + 15a^3B + 145ab^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) + 2(a^2 - b^2) (56aAb + 15a^2B + 25b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) - \frac{105bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{105bd \sqrt{a + b \cos(c + dx)}} + \frac{2(56aAb + 15a^2B + 25b^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{105d} + \frac{2(7Ab + 5aB)(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{35d} + \frac{2B(a + b \cos(c + dx))^{5/2} \sin(c + dx)}{7d}$$

output

```
2/35*(7*A*b+5*B*a)*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/d+2/7*B*(a+b*cos(d*x+c))^(5/2)*sin(d*x+c)/d+2/105*(56*A*a*b+15*B*a^2+25*B*b^2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/105*(161*A*a^2*b+63*A*b^3+15*B*a^3+145*B*a*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b)))^(1/2)*(a+b*cos(d*x+c))^(1/2)/b/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/105*(a^2-b^2)*(56*A*a*b+15*B*a^2+25*B*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b)))^(1/2)*(a+b*cos(d*x+c))/(a+b)^(1/2)/b/d/(a+b*cos(d*x+c))^(1/2)
```


3.313.2 Mathematica [A] (verified)

Time = 1.78 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.88

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{2b(105a^3A + 119aAb^2 + 135a^2bB + 25b^3B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + \dots}{\dots}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(2*b*(105*a^3*A + 119*a*A*b^2 + 135*a^2*b*B + 25*b^3*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + 2*(161*a^2*A*b + 63*A*b^3 + 15*a^3*B + 145*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)]) + b*(a + b*Cos[c + d*x])*(154*a*A*b + 90*a^2*B + 65*b^2*B + 6*b*(7*A*b + 15*a*B)*Cos[c + d*x] + 15*b^2*B*Cos[2*(c + d*x)])*Sin[c + d*x])/(105*b*d*Sqrt[a + b*Cos[c + d*x]])`

3.313.3 Rubi [A] (verified)

Time = 1.54 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.04, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.720$, Rules used = {3042, 3232, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{5/2} \left(A + B \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx \\ & \quad \downarrow \text{3232} \\ & \frac{2}{7} \int \frac{1}{2} (a + b \cos(c + dx))^{3/2} (7aA + 5bB + (7Ab + 5aB) \cos(c + dx)) dx + \\ & \quad \frac{2B \sin(c + dx) (a + b \cos(c + dx))^{5/2}}{7d} \end{aligned}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{7} \int (a + b \cos(c + dx))^{3/2} (7aA + 5bB + (7Ab + 5aB) \cos(c + dx)) dx + \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \\
& \downarrow 3042 \\
& \frac{1}{7} \int \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2} \left(7aA + 5bB + (7Ab + 5aB) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \\
& \downarrow 3232 \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (35Aa^2 + 40bBa + 21Ab^2 + (15Ba^2 + 56Aba + 25b^2B) \cos(c + dx)) dx + \frac{2(5aB + 7Aa^2)}{7d} \right) \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \\
& \downarrow 27 \\
& \frac{1}{7} \left(\frac{1}{5} \int \sqrt{a + b \cos(c + dx)} (35Aa^2 + 40bBa + 21Ab^2 + (15Ba^2 + 56Aba + 25b^2B) \cos(c + dx)) dx + \frac{2(5aB + 7Aa^2)}{7d} \right) \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \\
& \downarrow 3042 \\
& \frac{1}{7} \left(\frac{1}{5} \int \sqrt{a + b \sin \left(c + dx + \frac{\pi}{2} \right)} (35Aa^2 + 40bBa + 21Ab^2 + (15Ba^2 + 56Aba + 25b^2B) \sin \left(c + dx + \frac{\pi}{2} \right)) dx + \frac{2(5aB + 7Aa^2)}{7d} \right) \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \\
& \downarrow 3232 \\
& \frac{1}{7} \left(\frac{1}{5} \left(\frac{2}{3} \int \frac{105Aa^3 + 135bBa^2 + 119Ab^2a + 25b^3B + (15Ba^3 + 161Aba^2 + 145b^2Ba + 63Ab^3) \cos(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{2(5aB + 7Aa^2)}{7d} \right) \right) \\
& \quad \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \\
& \downarrow 27
\end{aligned}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{105Aa^3 + 135bBa^2 + 119Ab^2a + 25b^3B + (15Ba^3 + 161Aba^2 + 145b^2Ba + 63Ab^3) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \right) \right) \downarrow \mathbf{3042}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{105Aa^3 + 135bBa^2 + 119Ab^2a + 25b^3B + (15Ba^3 + 161Aba^2 + 145b^2Ba + 63Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \right) \right) \downarrow \mathbf{3231}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \int \sqrt{a + b \cos(c + dx)} dx}{b} - \frac{(a^2 - b^2)(15a^2B + 56aAb + 25b^2)}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \right) \right) \downarrow \mathbf{3042}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx}{b} - \frac{(a^2 - b^2)(15a^2B + 56aAb + 25b^2)}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \right) \right) \downarrow \mathbf{3134}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2)(15a^2B + 56aAb + 25b^2)}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \right) \right) \downarrow \mathbf{3042}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2)(15a^2B + 56aAb + 25b^2)}{b} \right) + \frac{2B \sin(c + dx)(a + b \cos(c + dx))^{5/2}}{7d} \right) \right)$$

↓ 3132

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (15a^2B - 25a^2Ab + 15ab^2B + 63Ab^3)}{7d} \right) \right) \right)$$

↓ 3142

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (15a^2B - 25a^2Ab + 15ab^2B + 63Ab^3)}{7d} \right) \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (15a^2B - 25a^2Ab + 15ab^2B + 63Ab^3)}{7d} \right) \right) \right)$$

↓ 3140

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2(15a^2B + 56aAb + 25b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d} + \frac{1}{3} \left(\frac{2(15a^3B + 161a^2Ab + 145ab^2B + 63Ab^3) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2) (15a^2B - 25a^2Ab + 15ab^2B + 63Ab^3)}{7d} \right) \right) \right)$$

input `Int[(a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]),x]`

```
output (2*B*(a + b*cos[c + d*x])^(5/2)*sin[c + d*x])/(7*d) + ((2*(7*A*b + 5*a*B)*
(a + b*cos[c + d*x])^(3/2)*sin[c + d*x])/(5*d) + (((2*(161*a^2*A*b + 63*A*
b^3 + 15*a^3*B + 145*a*b^2*B)*sqrt[a + b*cos[c + d*x]]*EllipticE[(c + d*x)
/2, (2*b)/(a + b)])/(b*d*sqrt[(a + b*cos[c + d*x])/(a + b)]) - (2*(a^2 - b
^2)*(56*a*A*b + 15*a^2*B + 25*b^2*B)*sqrt[(a + b*cos[c + d*x])/(a + b)]*El
lipticF[(c + d*x)/2, (2*b)/(a + b)]/(b*d*sqrt[a + b*cos[c + d*x]]))/3 + (
2*(56*a*A*b + 15*a^2*B + 25*b^2*B)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/
(3*d))/5)/7
```

3.313.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*sin[c + d*x]]/sqrt[(a + b*sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a
+ b*sin[c + d*x])/(a + b)]/sqrt[a + b*sin[c + d*x]] Int[1/Sqrt[a/(a + b)
+ (b/(a + b))*sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && !GtQ[a + b, 0]
```

```
rule 3231 Int[((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*sin[(e_.) + (
f_.)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x
]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b
, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

```
rule 3232 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(
f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*
d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ
[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m,
0] && IntegerQ[2*m]
```

3.313.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1304 vs. $2(322) = 644$.

Time = 15.98 (sec) , antiderivative size = 1305, normalized size of antiderivative = 4.53

method	result	size
default	Expression too large to display	1305
parts	Expression too large to display	1491

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```

-2/105*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^4+(-168*A*b^4-480*B*a*b^3-360*B*
b^4)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(392*A*a*b^3+168*A*b^4+360*B*
a^2*b^2+480*B*a*b^3+280*B*b^4)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-1
54*A*a^2*b^2-196*A*a*b^3-42*A*b^4-90*B*a^3*b-180*B*a^2*b^2-170*B*a*b^3-80*
B*b^4)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-56*A*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3*b+56*a*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*
x+1/2*c),(-2*b/(a-b))^(1/2))*b^3+161*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/
(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))*a^3*b-161*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*
sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b
/(a-b))^(1/2))*a^2*b^2+63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1
/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b)
))^(1/2))*a*b^3-63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+
1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)
))*b^4-15*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+
(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4-10
*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/...

```

3.313.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.95

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \frac{\sqrt{2}(30i Ba^4 + 7i Aa^3b - 115i Ba^2b^2 - 231i Aab^3 - 75i Bb^4)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4}{2}\right)}{\dots}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="fracas")`

```
output 1/315*(sqrt(2)*(30*I*B*a^4 + 7*I*A*a^3*b - 115*I*B*a^2*b^2 - 231*I*A*a*b^3
- 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(
(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b
) + sqrt(2)*(-30*I*B*a^4 - 7*I*A*a^3*b + 115*I*B*a^2*b^2 + 231*I*A*a*b^3 +
75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8
*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)
- 3*sqrt(2)*(-15*I*B*a^3*b - 161*I*A*a^2*b^2 - 145*I*B*a*b^3 - 63*I*A*b^4)
*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/
b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/
b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(15
*I*B*a^3*b + 161*I*A*a^2*b^2 + 145*I*B*a*b^3 + 63*I*A*b^4)*sqrt(b)*weierst
rassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrass
PInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*co
s(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(15*B*b^4*cos(d*x + c)^2 +
45*B*a^2*b^2 + 77*A*a*b^3 + 25*B*b^4 + 3*(15*B*a*b^3 + 7*A*b^4)*cos(d*x +
c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^2*d)
```

3.313.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.313.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2), x)
```

3.313. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$

3.313.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2), x)`

3.313.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx = \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2), x)`

3.314 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec(c+dx) dx$

3.314.1 Optimal result	2907
3.314.2 Mathematica [C] (verified)	2908
3.314.3 Rubi [A] (verified)	2909
3.314.4 Maple [B] (verified)	2916
3.314.5 Fricas [F]	2917
3.314.6 Sympy [F(-1)]	2917
3.314.7 Maxima [F]	2917
3.314.8 Giac [F]	2918
3.314.9 Mupad [F(-1)]	2918

3.314.1 Optimal result

Integrand size = 31, antiderivative size = 292

$$\begin{aligned}
 & \int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec(c+dx) dx = \\
 & \frac{2(35aAb + 23a^2B + 9b^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{15d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & + \frac{2(10a^2Ab + 5Ab^3 - 8a^3B + 8ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{15d \sqrt{a+b \cos(c+dx)}} \\
 & + \frac{2a^3A \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \\
 & + \frac{2b(5Ab + 8aB) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15d} \\
 & + \frac{2bB(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{5d}
 \end{aligned}$$

output $\frac{2}{5}bB(a+b\cos(dx+c))^{3/2}\sin(dx+c)/d+2/15b(5A^2b+8B^2a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/15(35A^2ab+23B^2a^2+9B^2b^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticE}(\sin(1/2dx+1/2c),2^{1/2})(b/(a+b))^{1/2})(a+b\cos(dx+c))^{1/2}/d/((a+b\cos(dx+c))/(a+b))^{1/2}+2/15(10A^2a^2b+5A^2b^3-8B^2a^3+8B^2ab^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticF}(\sin(1/2dx+1/2c),2^{1/2})(b/(a+b))^{1/2})((a+b\cos(dx+c))/(a+b))^{1/2}/d/(a+b\cos(dx+c))^{1/2}+2a^3A(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticPi}(\sin(1/2dx+1/2c),2,2^{1/2})(b/(a+b))^{1/2})((a+b\cos(dx+c))/(a+b))^{1/2}/d/(a+b\cos(dx+c))^{1/2}$

3.314.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.53 (sec) , antiderivative size = 453, normalized size of antiderivative = 1.55

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \frac{4(45a^2Ab + 5Ab^3 + 15a^3B + 17ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(30a^3A + 35aAb^2 + 23a^2bB + 9b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output $((4*(45*a^2*A*b + 5*A*b^3 + 15*a^3*B + 17*a*b^2*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(30*a^3*A + 35*a*A*b^2 + 23*a^2*b*B + 9*b^3*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + ((2*I)*(35*a*A*b + 23*a^2*B + 9*b^2*B)*\text{Sqrt}[-(b*(-1 + \text{Cos}[c + d*x]))/(a + b)]*\text{Sqrt}[-(b*(1 + \text{Cos}[c + d*x]))/(a - b)]*\text{Csc}[c + d*x]*(-2*a*(a - b)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*\text{EllipticPi}[(a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b))))/(a*b*\text{Sqrt}[-(a + b)^{-1}]) + 4*b*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]*(5*A*b + 11*a*B + 3*b*B*\text{Cos}[c + d*x])*Sin[c + d*x])/(30*d)$

3.314.3 Rubi [A] (verified)

Time = 2.49 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.04, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.677$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})} dx$$

$$\downarrow \text{3469}$$

$$\frac{2}{5} \int \frac{1}{2} \sqrt{a+b\cos(c+dx)} (5Aa^2 + b(5Ab+8aB)\cos^2(c+dx) + (3Bb^2 + 5a(2Ab+aB))\cos(c+dx)) \sec(c+dx) dx + \frac{2bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d}$$

$$\downarrow \text{27}$$

$$\frac{1}{5} \int \sqrt{a+b\cos(c+dx)} (5Aa^2 + b(5Ab+8aB)\cos^2(c+dx) + (3Bb^2 + 5a(2Ab+aB))\cos(c+dx)) \sec(c+dx) dx + \frac{2bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{5} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})} (5Aa^2 + b(5Ab+8aB)\sin(c+dx+\frac{\pi}{2})^2 + (3Bb^2 + 5a(2Ab+aB))\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{2bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d}$$

$$\downarrow \text{3528}$$

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{(15Aa^3 + b(23Ba^2 + 35Aba + 9b^2B)\cos^2(c+dx) + (15Ba^3 + 45Aba^2 + 17b^2Ba + 5Ab^3)\cos(c+dx)) \sec(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx + \frac{2bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d} \right)$$

3.314. $\int (a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) \sec(c+dx) dx$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{(15Aa^3 + b(23Ba^2 + 35Aba + 9b^2B)) \cos^2(c + dx) + (15Ba^3 + 45Aba^2 + 17b^2Ba + 5Ab^3) \cos(c + dx) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx \right) - \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{15Aa^3 + b(23Ba^2 + 35Aba + 9b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (15Ba^3 + 45Aba^2 + 17b^2Ba + 5Ab^3) \sin(c + dx + \frac{\pi}{2}) \sec(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) - \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3538

$$\frac{1}{5} \left(\frac{1}{3} \left((23a^2B + 35aAb + 9b^2B) \int \sqrt{a + b \cos(c + dx)} dx - \frac{\int \frac{(15Aba^3 + b(-8Ba^3 + 10Aba^2 + 8b^2Ba + 5Ab^3)) \cos(c + dx) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) \right) - \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 25

$$\frac{1}{5} \left(\frac{1}{3} \left((23a^2B + 35aAb + 9b^2B) \int \sqrt{a + b \cos(c + dx)} dx + \frac{\int \frac{(15Aba^3 + b(-8Ba^3 + 10Aba^2 + 8b^2Ba + 5Ab^3)) \cos(c + dx) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) \right) - \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left((23a^2B + 35aAb + 9b^2B) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx + \frac{\int \frac{15Aba^3 + b(-8Ba^3 + 10Aba^2 + 8b^2Ba + 5Ab^3) \sin(c + dx + \frac{\pi}{2}) \sec(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \right) \right) - \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3134

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(23a^2B + 35aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{\int \frac{15Aba^3 + b(-8Ba^3 + 10Aba^2 + 8b^2Ba + 5Ab^3) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} \right) \right) + \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(23a^2B + 35aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{\int \frac{15Aba^3 + b(-8Ba^3 + 10Aba^2 + 8b^2Ba + 5Ab^3) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} \right) \right) + \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3132

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{\int \frac{15Aba^3 + b(-8Ba^3 + 10Aba^2 + 8b^2Ba + 5Ab^3) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{2(23a^2B + 35aAb + 9b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right) + \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3481

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3Ab \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + b(-8a^3B + 10a^2Ab + 8ab^2B + 5Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} + \frac{2(23a^2B + 35aAb + 9b^2B) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right) + \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3Ab \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + b(-8a^3B + 10a^2Ab + 8ab^2B + 5Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{2(23a^2B + 35aAb + 9b^2B) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) \right) + \frac{2bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d}$$

↓ 3142

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3 Ab \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(-8a^3 B+10a^2 Ab+8ab^2 B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} \right) + \frac{2bB \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d} \right) +$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3 Ab \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(-8a^3 B+10a^2 Ab+8ab^2 B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} \right) + \frac{2bB \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d} \right) +$$

↓ 3140

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3 Ab \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(-8a^3 B+10a^2 Ab+8ab^2 B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \right) + \frac{2bB \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d} \right) +$$

↓ 3286

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3 Ab \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}} dx + \frac{2b(-8a^3 B+10a^2 Ab+8ab^2 B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \right) + \frac{2bB \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{5d} \right) +$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{15a^3 Ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2b(-8a^3 B + 10a^2 Ab + 8ab^2 B + 5Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \sqrt{\frac{a+b \cos(c+dx)}{a+b}}\right)}{b d \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2bB \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5d}$$

↓ 3284

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(23a^2 B + 35aAb + 9b^2 B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{30a^3 Ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \sqrt{\frac{a+b \cos(c+dx)}{a+b}}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{2bB \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{5d}$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x],x]`

output `(2*b*B*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + (((2*(35*a*A*b + 23*a^2*B + 9*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*b*(10*a^2*A*b + 5*A*b^3 - 8*a^3*B + 8*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (30*a^3*A*b*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/3 + (2*b*(5*A*b + 8*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

3.314.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[(Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.314.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1066 vs. $2(355) = 710$.

Time = 14.27 (sec) , antiderivative size = 1067, normalized size of antiderivative = 3.65

method	result	size
default	Expression too large to display	1067
parts	Expression too large to display	1192

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x,method=_RETURNVER
BOSE)
```

```
output -2/15*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*c
os(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^3+(20*A*b^3+56*B*a*b^2+24*B*b^3)*
sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*a*b^2-10*A*b^3-22*B*a^2*b-2
8*B*a*b^2-6*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+10*A*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*Ell
ipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+5*A*b^3*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*Ellipti
cF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+35*A*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*
x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b-35*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b
/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c
),(-2*b/(a-b))^(1/2))*a*b^2-15*A*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a
-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),
2,(-2*b/(a-b))^(1/2))-8*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2
*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))
^(1/2))*a^3+8*B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2
*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*
b^2+23*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+
b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3-23*B*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a...
```

3.314.5 Fricas [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c), x)`

3.314.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x)`

output `Timed out`

3.314.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c), x)`

3.314.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c) dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c), x)`

3.314.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x), x)`

3.315 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^2(c+dx) dx$

3.315.1 Optimal result	2919
3.315.2 Mathematica [C] (verified)	2920
3.315.3 Rubi [A] (verified)	2921
3.315.4 Maple [B] (verified)	2928
3.315.5 Fricas [F(-1)]	2929
3.315.6 Sympy [F(-1)]	2929
3.315.7 Maxima [F]	2929
3.315.8 Giac [F]	2930
3.315.9 Mupad [F(-1)]	2930

3.315.1 Optimal result

Integrand size = 33, antiderivative size = 296

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx =$$

$$-\frac{(3a^2A - 6Ab^2 - 14abB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{3d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{(3a^3A + 12aAb^2 + 4a^2bB + 2b^3B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{a^2(5Ab + 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

$$- \frac{b(3aA - 2bB) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3d}$$

$$+ \frac{aA(a + b \cos(c + dx))^{3/2} \tan(c + dx)}{d}$$

output
$$-1/3*b*(3*A*a-2*B*b)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{1/2}/d-1/3*(3*A*a^2-6*A*b^2-14*B*a*b)*(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{1/2}*(b/(a+b))^{1/2})*(a+b*\cos(d*x+c))^{1/2}/d/((a+b*\cos(d*x+c))/(a+b))^{1/2}+1/3*(3*A*a^3+12*A*a*b^2+4*B*a^2*b+2*B*b^3)*(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{1/2}*(b/(a+b))^{1/2})*((a+b*\cos(d*x+c))/(a+b))^{1/2}/d/(a+b*\cos(d*x+c))^{1/2}+a^2*(5*A*b+2*B*a)*(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticPi}(\sin(1/2*d*x+1/2*c),2,2^{1/2}*(b/(a+b))^{1/2})*((a+b*\cos(d*x+c))/(a+b))^{1/2}/d/(a+b*\cos(d*x+c))^{1/2}+a*A*(a+b*\cos(d*x+c))^{3/2}*\tan(d*x+c)/d$$

3.315.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.78 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.49

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{8b(9aAb+9a^2B+b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(27a^2Ab+6Ab^3+12a^3B+14ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticPi}\left(\frac{1}{2}(c+dx), 2, \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]`

output
$$\left((8*b*(9*a*A*b + 9*a^2*B + b^2*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(27*a^2*A*b + 6*A*b^3 + 12*a^3*B + 14*a*b^2*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + ((2*I)*(-3*a^2*A + 6*A*b^2 + 14*a*b*B)*\text{Sqrt}[-((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[(b*(1 + \text{Cos}[c + d*x]))/(-a + b)]*\text{Csc}[c + d*x]*(-2*a*(a - b)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b) + b*(-2*a*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b) + b*\text{EllipticPi}[(a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)))/((a*b*\text{Sqrt}[-(a + b)^{-1}]) + 4*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]*(3*a^2*A + 2*b^2*B*\text{Cos}[c + d*x])*Tan[c + d*x])/(12*d) \right)$$

3.315.3 Rubi [A] (verified)

Time = 2.59 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.05, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3468, 27, 3042, 3528, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^2(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^2} dx$$

$$\downarrow \text{3468}$$

$$\int \frac{1}{2} \sqrt{a+b\cos(c+dx)} (-b(3aA-2bB)\cos^2(c+dx) + 2b(Ab+2aB)\cos(c+dx) + a(5Ab+2aB)) \sec(c+dx) dx + \frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

$$\downarrow \text{27}$$

$$\frac{1}{2} \int \sqrt{a+b\cos(c+dx)} (-b(3aA-2bB)\cos^2(c+dx) + 2b(Ab+2aB)\cos(c+dx) + a(5Ab+2aB)) \sec(c+dx) dx + \frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{2} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})} (-b(3aA-2bB)\sin(c+dx+\frac{\pi}{2})^2 + 2b(Ab+2aB)\sin(c+dx+\frac{\pi}{2}) + a(5Ab+2aB)) \sec(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx + \frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

$$\downarrow \text{3528}$$

$$\frac{1}{2} \left(\frac{2}{3} \int \frac{(3(5Ab+2aB)a^2 - b(3Aa^2 - 14bBa - 6Ab^2)\cos^2(c+dx) + 2b(9Ba^2 + 9Aba + b^2B)\cos(c+dx)) \sec(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx + \frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d} \right)$$

3.315. $\int (a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) \sec^2(c+dx) dx$

↓ 27

$$\frac{1}{2} \left(\frac{1}{3} \int \frac{(3(5Ab + 2aB)a^2 - b(3Aa^2 - 14bBa - 6Ab^2)) \cos^2(c + dx) + 2b(9Ba^2 + 9Aba + b^2B) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} \sec(c + dx) dx \right)$$

$$\frac{aA \tan(c + dx)(a + b \cos(c + dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \int \frac{3(5Ab + 2aB)a^2 - b(3Aa^2 - 14bBa - 6Ab^2) \sin(c + dx + \frac{\pi}{2})^2 + 2b(9Ba^2 + 9Aba + b^2B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \sec(c + dx + \frac{\pi}{2}) dx \right)$$

$$\frac{aA \tan(c + dx)(a + b \cos(c + dx))^{3/2}}{d}$$

↓ 3538

$$\frac{1}{2} \left(\frac{1}{3} \left(- \left((3a^2A - 14abB - 6Ab^2) \int \sqrt{a + b \cos(c + dx)} dx \right) - \frac{\int - \frac{(3b(5Ab + 2aB)a^2 + b(3Aa^3 + 4bBa^2 + 12Ab^2a + 2b^3B)) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) \right)$$

$$\frac{aA \tan(c + dx)(a + b \cos(c + dx))^{3/2}}{d}$$

↓ 25

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{\int \frac{(3b(5Ab + 2aB)a^2 + b(3Aa^3 + 4bBa^2 + 12Ab^2a + 2b^3B)) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} \sec(c + dx) dx}{b} - (3a^2A - 14abB - 6Ab^2) \int \sqrt{a + b \cos(c + dx)} dx \right) \right)$$

$$\frac{aA \tan(c + dx)(a + b \cos(c + dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{\int \frac{3b(5Ab + 2aB)a^2 + b(3Aa^3 + 4bBa^2 + 12Ab^2a + 2b^3B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} - (3a^2A - 14abB - 6Ab^2) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx \right) \right)$$

$$\frac{aA \tan(c + dx)(a + b \cos(c + dx))^{3/2}}{d}$$

↓ 3134

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{\int \frac{3b(5Ab+2aB)a^2+b(3Aa^3+4bBa^2+12Ab^2a+2b^3B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(3a^2A - 14abB - 6Ab^2) \sqrt{a+b\cos(c+dx)}}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{\int \frac{3b(5Ab+2aB)a^2+b(3Aa^3+4bBa^2+12Ab^2a+2b^3B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(3a^2A - 14abB - 6Ab^2) \sqrt{a+b\cos(c+dx)}}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3132

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{\int \frac{3b(5Ab+2aB)a^2+b(3Aa^3+4bBa^2+12Ab^2a+2b^3B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(3a^2A - 14abB - 6Ab^2) \sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3481

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB + 5Ab) \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx + b(3a^3A + 4a^2bB + 12aAb^2 + 2b^3B) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2(3a^2A - 14abB - 6Ab^2) \sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB + 5Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + b(3a^3A + 4a^2bB + 12aAb^2 + 2b^3B) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(3a^2A - 14abB - 6Ab^2) \sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3142

3.315. $\int (a+b\cos(c+dx))^{5/2} (A+B\cos(c+dx)) \sec^2(c+dx) dx$

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB + 5Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(3a^3A+4a^2bB+12aAb^2+2b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}}}{\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB + 5Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(3a^3A+4a^2bB+12aAb^2+2b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx)}{a+b}}}}{\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3140

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB + 5Ab) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(3a^3A+4a^2bB+12aAb^2+2b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3286

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB+5Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2b(3a^3A+4a^2bB+12aAb^2+2b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

$$\frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d}$$

↓ 3042

3.315. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx$

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{3a^2b(2aB+5Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2b(3a^3A+4a^2bB+12aAb^2+2b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \right) + \frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d} \right)$$

↓ 3284

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{6a^2b(2aB+5Ab)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} + \frac{2b(3a^3A+4a^2bB+12aAb^2+2b^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \right) + \frac{aA \tan(c+dx)(a+b\cos(c+dx))^{3/2}}{d} \right)$$

```
input Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^2,x]
```

```
output (((-2*(3*a^2*A - 6*A*b^2 - 14*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*b*(3*a^3*A + 12*a*A*b^2 + 4*a^2*b*B + 2*b^3*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*a^2*b*(5*A*b + 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/3 - (2*b*(3*a*A - 2*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/2 + (a*A*(a + b*Cos[c + d*x])^(3/2)*Tan[c + d*x])/d
```

3.315.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.315. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx$

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.315.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1489 vs. $2(361) = 722$.

Time = 22.72 (sec) , antiderivative size = 1490, normalized size of antiderivative = 5.03

method	result	size
parts	Expression too large to display	1490
default	Expression too large to display	1563

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x,method=_RETURNV
ERBOSE)
```

```
output -A*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*cos(1/2*
d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*a^2*b+(-2*a^3-2*a^2*b)*sin(1/2*d*x+1/2*c)^
2*cos(1/2*d*x+1/2*c)-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*
x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))*a^3+4*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b^2-Ellipti
cE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3+EllipticE(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))*a^2*b+2*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/
2))*a*b^2-2*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^3-5*Ellipti
cPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))*a^2*b)*sin(1/2*d*x+1/2*c)^2+(
sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(
1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3+4*a*b^2*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*
EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-sin(1/2*d*x+1/2*c)^2)^(1/
2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*
d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)
)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2
*b/(a-b))^(1/2))*a^2*b+2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*
d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))*a*b^2-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)
^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*...
```

3.315.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="fricas")`

output `Timed out`

3.315.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**2,x)`

output `Timed out`

3.315.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^2 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^2, x)`

3.315.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^2 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^2, x)`

3.315.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^2} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^2,x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^2, x)`

3.316 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^3(c+dx) dx$

3.316.1 Optimal result	2931
3.316.2 Mathematica [C] (verified)	2932
3.316.3 Rubi [A] (verified)	2933
3.316.4 Maple [B] (verified)	2940
3.316.5 Fricas [F(-1)]	2941
3.316.6 Sympy [F(-1)]	2941
3.316.7 Maxima [F]	2941
3.316.8 Giac [F]	2942
3.316.9 Mupad [F(-1)]	2942

3.316.1 Optimal result

Integrand size = 33, antiderivative size = 315

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx =$$

$$\frac{(9aAb + 4a^2B - 8b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{4d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{(11a^2Ab + 8Ab^3 + 4a^3B + 16ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{a(4a^2A + 15Ab^2 + 20abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{a(7Ab + 4aB) \sqrt{a + b \cos(c + dx)} \tan(c + dx)}{4d}$$

$$+ \frac{aA(a + b \cos(c + dx))^{3/2} \sec(c + dx) \tan(c + dx)}{2d}$$

output
$$-1/4*(9*A*a*b+4*B*a^2-8*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)}*(a+b*cos(d*x+c))^{(1/2)})/d/((a+b*cos(d*x+c))/(a+b))^{(1/2)}+1/4*(11*A*a^2*b+8*A*b^3+4*B*a^3+16*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)}*(a+b*cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*cos(d*x+c))^{(1/2)}+1/4*a*(4*A*a^2+15*A*b^2+20*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^{(1/2)}*(b/(a+b))^{(1/2)}*(a+b*cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*cos(d*x+c))^{(1/2)}+1/2*a*A*(a+b*cos(d*x+c))^{(3/2)}*sec(d*x+c)*tan(d*x+c)/d+1/4*a*(7*A*b+4*B*a)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d$$

3.316.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.16 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.43

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \frac{8b(a^2 A + 4Ab^2 + 12abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{2(8a^3 A + 21aAb^2 + 36a^2 bB + 8b^3 B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(\frac{1}{2}(c+dx), 2, \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]`

output
$$\frac{((8*b*(a^2*A + 4*A*b^2 + 12*a*b*B)*\operatorname{Sqrt}[(a + b*\operatorname{Cos}[c + d*x])/(a + b)]*\operatorname{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]] + (2*(8*a^3*A + 21*a*A*b^2 + 36*a^2*b*B + 8*b^3*B)*\operatorname{Sqrt}[(a + b*\operatorname{Cos}[c + d*x])/(a + b)]*\operatorname{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)]/\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]] + ((2*I)*(-9*a*A*b - 4*a^2*B + 8*b^2*B)*\operatorname{Sqrt}[-((b*(-1 + \operatorname{Cos}[c + d*x]))/(a + b))])*\operatorname{Sqrt}[(b*(1 + \operatorname{Cos}[c + d*x]))/(-a + b)]*\operatorname{Csc}[c + d*x]*(-2*a*(a - b)*\operatorname{EllipticE}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-(a + b)^{-1}]]*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*\operatorname{EllipticF}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-(a + b)^{-1}]]*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]], (a + b)/(a - b)] + b*\operatorname{EllipticPi}[(a + b)/a, I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-(a + b)^{-1}]]*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]], (a + b)/(a - b)))/(a*b*\operatorname{Sqrt}[-(a + b)^{-1}]) + 4*a*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]*(2*a*A + (9*A*b + 4*a*B)*\operatorname{Cos}[c + d*x])*Sec[c + d*x]*\operatorname{Tan}[c + d*x])/(16*d)$$

3.316.3 Rubi [A] (verified)

Time = 2.58 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.03, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^3(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^3} dx$$

↓ 3468

$$\frac{1}{2} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (-b(aA - 4bB) \cos^2(c + dx) + 2(Aa^2 + 4bBa + 2Ab^2) \cos(c + dx) + a(7Ab + 4aB)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^{3/2}}{2d}$$

↓ 27

$$\frac{1}{4} \int \sqrt{a + b \cos(c + dx)} (-b(aA - 4bB) \cos^2(c + dx) + 2(Aa^2 + 4bBa + 2Ab^2) \cos(c + dx) + a(7Ab + 4aB)) \sec^2(c + dx) dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (-b(aA - 4bB) \sin(c + dx + \frac{\pi}{2})^2 + 2(Aa^2 + 4bBa + 2Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(7Ab + 4aB)) \sec^2(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2} dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^{3/2}}{2d}$$

↓ 3526

$$\frac{1}{4} \left(\int \frac{(-b(4Ba^2 + 9Aba - 8b^2B) \cos^2(c + dx) + 2b(Aa^2 + 12bBa + 4Ab^2) \cos(c + dx) + a(4Aa^2 + 20bBa + 15Ab^2)) \sec^2(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{aA \tan(c + dx) \sec(c + dx)(a + b \cos(c + dx))^{3/2}}{2d} \right)$$

3.316. $\int (a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^3(c + dx) dx$

↓ 27

$$\frac{1}{4} \left(\frac{1}{2} \int \frac{(-b(4Ba^2 + 9Aba - 8b^2B) \cos^2(c + dx) + 2b(Aa^2 + 12bBa + 4Ab^2) \cos(c + dx) + a(4Aa^2 + 20bBa + 15Ab^2)) \sqrt{a + b \cos(c + dx)}}{aA \tan(c + dx) \sec(c + dx) (a + b \cos(c + dx))^{3/2}} dx \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \int \frac{-b(4Ba^2 + 9Aba - 8b^2B) \sin(c + dx + \frac{\pi}{2})^2 + 2b(Aa^2 + 12bBa + 4Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Aa^2 + 20bBa + 15Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right)$$

↓ 3538

$$\frac{1}{4} \left(\frac{1}{2} \left(-((4a^2B + 9aAb - 8b^2B) \int \sqrt{a + b \cos(c + dx)} dx) - \frac{\int -\frac{(ab(4Aa^2 + 20bBa + 15Ab^2) + b(4Ba^3 + 11Aba^2 + 16b^2Ba + 8Ab^3)) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} \right) \right)$$

↓ 25

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{\int \frac{(ab(4Aa^2 + 20bBa + 15Ab^2) + b(4Ba^3 + 11Aba^2 + 16b^2Ba + 8Ab^3)) \cos(c + dx) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} - (4a^2B + 9aAb - 8b^2B) \int \sqrt{a + b \cos(c + dx)} dx \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{\int \frac{ab(4Aa^2 + 20bBa + 15Ab^2) + b(4Ba^3 + 11Aba^2 + 16b^2Ba + 8Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} - (4a^2B + 9aAb - 8b^2B) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx \right) \right)$$

↓ 3134

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{\int \frac{ab(4Aa^2+20bBa+15Ab^2)+b(4Ba^3+11Aba^2+16b^2Ba+8Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(4a^2B+9aAb-8b^2B) \sqrt{a+b \cos(c+dx)}}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{aA \tan(c+dx) \sec(c+dx)(a+b \cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{\int \frac{ab(4Aa^2+20bBa+15Ab^2)+b(4Ba^3+11Aba^2+16b^2Ba+8Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(4a^2B+9aAb-8b^2B) \sqrt{a+b \cos(c+dx)}}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{aA \tan(c+dx) \sec(c+dx)(a+b \cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3132}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{\int \frac{ab(4Aa^2+20bBa+15Ab^2)+b(4Ba^3+11Aba^2+16b^2Ba+8Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(4a^2B+9aAb-8b^2B) \sqrt{a+b \cos(c+dx)}}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right) - \frac{aA \tan(c+dx) \sec(c+dx)(a+b \cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3481}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A+20abB+15Ab^2) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} - \frac{aA \tan(c+dx) \sec(c+dx)(a+b \cos(c+dx))^{3/2}}{2d} \right) \right) \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A+20abB+15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} - \frac{aA \tan(c+dx) \sec(c+dx)(a+b \cos(c+dx))^{3/2}}{2d} \right) \right) \downarrow \text{3142}$$

3.316. $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^3(c+dx) dx$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A + 20abB + 15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{a+b\cos(c+dx)}} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a+b\cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A + 20abB + 15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{a+b\cos(c+dx)}} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a+b\cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3140}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A + 20abB + 15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{d\sqrt{a+b\cos(c+dx)}} \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a+b\cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3286}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A+20abB+15Ab^2) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{d\sqrt{a+b\cos(c+dx)}} \text{EllipticF}\left(\frac{1}{2}(\right) \right) \right) \right. \\ \left. \frac{aA \tan(c+dx) \sec(c+dx)(a+b\cos(c+dx))^{3/2}}{2d} \right) \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{ab(4a^2A+20abB+15Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) (a+b \cos(c+dx))^{3/2}}{2d}$$

↓ 3284

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{2ab(4a^2A+20abB+15Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} + \frac{2b(4a^3B+11a^2Ab+16ab^2B+8Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticE}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec(c+dx) (a+b \cos(c+dx))^{3/2}}{2d}$$

```
input Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^3,x]
```

```
output (a*A*(a + b*Cos[c + d*x])^(3/2)*Sec[c + d*x]*Tan[c + d*x]/(2*d) + (((-2*(9*a*A*b + 4*a^2*B - 8*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*b*(11*a^2*A*b + 8*A*b^3 + 4*a^3*B + 16*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*a*b*(4*a^2*A + 15*A*b^2 + 20*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/2 + (a*(7*A*b + 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/d)/4
```

3.316.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.316. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx$

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.316.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1741 vs. $2(376) = 752$.

Time = 64.48 (sec) , antiderivative size = 1742, normalized size of antiderivative = 5.53

method	result	size
default	Expression too large to display	1742
parts	Expression too large to display	2096

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*b^3*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-2*B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2
*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)
*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/
2))+6*B*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)
/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)
*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-2*B*b^2*(a-b)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+
1/2*c),(-2*b/(a-b))^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)
))+2*A*a^3*(-1/2*cos(1/2*d*x+1/2*c)/a*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin
(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^2+3/4*b/a^2*cos(1/2*d*
x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*c
os(1/2*d*x+1/2*c)^2-1)-1/8*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*
d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*
x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+3/8/a*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(
-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*EllipticE...
```

3.316.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="fricas")`

output `Timed out`

3.316.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**3,x)`

output `Timed out`

3.316.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^3 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^3, x)`

3.316.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^3 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^3, x)`

3.316.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^3(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^3} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^3,x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^3, x)`

3.317 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^4(c+dx) dx$

3.317.1 Optimal result	2943
3.317.2 Mathematica [C] (verified)	2944
3.317.3 Rubi [A] (verified)	2945
3.317.4 Maple [B] (verified)	2954
3.317.5 Fricas [F(-1)]	2955
3.317.6 Sympy [F(-1)]	2955
3.317.7 Maxima [F]	2955
3.317.8 Giac [F]	2956
3.317.9 Mupad [F(-1)]	2956

3.317.1 Optimal result

Integrand size = 33, antiderivative size = 376

$$\begin{aligned}
 & \int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^4(c+dx) dx = \\
 & \frac{(16a^2A + 33Ab^2 + 54abB) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{24d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & + \frac{(16a^3A + 59aAb^2 + 66a^2bB + 48b^3B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{24d \sqrt{a+b \cos(c+dx)}} \\
 & + \frac{(20a^2Ab + 5Ab^3 + 8a^3B + 30ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{8d \sqrt{a+b \cos(c+dx)}} \\
 & + \frac{(16a^2A + 33Ab^2 + 54abB) \sqrt{a+b \cos(c+dx)} \tan(c+dx)}{24d} \\
 & + \frac{a(3Ab + 2aB) \sqrt{a+b \cos(c+dx)} \sec(c+dx) \tan(c+dx)}{4d} \\
 & + \frac{aA(a+b \cos(c+dx))^{3/2} \sec^2(c+dx) \tan(c+dx)}{3d}
 \end{aligned}$$

output
$$-1/24*(16*A*a^2+33*A*b^2+54*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*(a+b*cos(d*x+c))^{(1/2)}/d/((a+b*cos(d*x+c))/(a+b))^{(1/2)}+1/24*(16*A*a^3+59*A*a*b^2+66*B*a^2*b+48*B*b^3)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*cos(d*x+c))^{(1/2)}+1/8*(20*A*a^2*b+5*A*b^3+8*B*a^3+30*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*cos(d*x+c))^{(1/2)}+1/3*a*A*(a+b*cos(d*x+c))^{(3/2)}*sec(d*x+c)^2*tan(d*x+c)/d+1/24*(16*A*a^2+33*A*b^2+54*B*a*b)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d+1/4*a*(3*A*b+2*B*a)*sec(d*x+c)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d$$

3.317.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.12 (sec) , antiderivative size = 486, normalized size of antiderivative = 1.29

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \frac{8b(13aAb + 6a^2B + 24b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2(104a^2Ab - 3Ab^3 + 48a^3B + 126ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticE}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

output
$$\begin{aligned} & ((8*b*(13*a*A*b + 6*a^2*B + 24*b^2*B)*\operatorname{Sqrt}[(a + b*\operatorname{Cos}[c + d*x])/(a + b)]* \operatorname{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]] + (2*(104*a^2*A*b - 3*A*b^3 + 48*a^3*B + 126*a*b^2*B)*\operatorname{Sqrt}[(a + b*\operatorname{Cos}[c + d*x])/(a + b)]* \operatorname{EllipticE}[(c + d*x)/2, (2*b)/(a + b)]/\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]] - \\ & ((2*I)*(16*a^2*A + 33*A*b^2 + 54*a*b*B)*\operatorname{Sqrt}[-((b*(-1 + \operatorname{Cos}[c + d*x]))/(a + b))]*\operatorname{Sqrt}[(b*(1 + \operatorname{Cos}[c + d*x]))/(-a + b)]*\operatorname{Csc}[c + d*x]*(-2*a*(a - b)* \operatorname{EllipticE}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-(a + b)^{-1}]]*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*\operatorname{EllipticF}[I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-(a + b)^{-1}]]*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]], (a + b)/(a - b)] + b*\operatorname{EllipticPi}[(a + b)/a, I*\operatorname{ArcSinh}[\operatorname{Sqrt}[-(a + b)^{-1}]]*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]], (a + b)/(a - b)])))/(a*b*\operatorname{Sqrt}[-(a + b)^{-1}]) + 4*\operatorname{Sqrt}[a + b*\operatorname{Cos}[c + d*x]]*\operatorname{Sec}[c + d*x]^2*(2*a*(13*A*b + 6*a*B)*\operatorname{Sin}[c + d*x] + (8*a^2*A + (33*A*b^2)/2 + 27*a*b*B)*\operatorname{Sin}[2*(c + d*x)] + 8*a^2*A*\operatorname{Tan}[c + d*x]))/(96*d) \end{aligned}$$

3.317. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx$

3.317.3 Rubi [A] (verified)

Time = 3.30 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.05, number of steps used = 24, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^4(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^4} dx$$

↓ 3468

$$\frac{1}{3} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)} (b(aA + 6bB) \cos^2(c + dx) + 2(2Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(3Ab + 2aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d}$$

↓ 27

$$\frac{1}{6} \int \sqrt{a + b \cos(c + dx)} (b(aA + 6bB) \cos^2(c + dx) + 2(2Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(3Ab + 2aB)) \sec^3(c + dx) dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{6} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (b(aA + 6bB) \sin^2(c + dx + \frac{\pi}{2}) + 2(2Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + 3a(3Ab + 2aB)) \sec^3(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^3} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d}$$

↓ 3526

$$\frac{1}{6} \left(\frac{1}{2} \int \frac{(b(6Ba^2 + 13Aba + 24b^2B) \cos^2(c + dx) + 2(6Ba^3 + 19Aba^2 + 36b^2Ba + 12Ab^3) \cos(c + dx) + a(16Aa^2 + 12Ab^2)) \sec^3(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx + \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d} \right)$$

3.317. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx$

$$\downarrow 27$$

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{(b(6Ba^2 + 13Aba + 24b^2B) \cos^2(c + dx) + 2(6Ba^3 + 19Aba^2 + 36b^2Ba + 12Ab^3) \cos(c + dx) + a(16Aa^2 + 12Ab^2 + 3b^3)) \sqrt{a + b \cos(c + dx)}}{aA \tan(c + dx) \sec^2(c + dx) (a + b \cos(c + dx))^{3/2}} dx \right)$$

$$\downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{b(6Ba^2 + 13Aba + 24b^2B) \sin(c + dx + \frac{\pi}{2})^2 + 2(6Ba^3 + 19Aba^2 + 36b^2Ba + 12Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(16Aa^2 + 12Ab^2 + 3b^3)}{\sin(c + dx + \frac{\pi}{2})^2 \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right)$$

$$\downarrow 3534$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\int \frac{(-ab(16Aa^2 + 54bBa + 33Ab^2) \cos^2(c + dx) + 2ab(6Ba^2 + 13Aba + 24b^2B) \cos(c + dx) + 3a(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3)) \sec(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx \right) \right)$$

$$\downarrow 27$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\int \frac{(-ab(16Aa^2 + 54bBa + 33Ab^2) \cos^2(c + dx) + 2ab(6Ba^2 + 13Aba + 24b^2B) \cos(c + dx) + 3a(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx \right) \right)$$

$$\downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\int \frac{-ab(16Aa^2 + 54bBa + 33Ab^2) \sin(c + dx + \frac{\pi}{2})^2 + 2ab(6Ba^2 + 13Aba + 24b^2B) \sin(c + dx + \frac{\pi}{2}) + 3a(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3)}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) \right)$$

$$\downarrow 3538$$

3.317. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{-a(16a^2A + 54abB + 33Ab^2) \int \sqrt{a + b \cos(c + dx)} dx - \int \frac{(3ab(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3) + ab(16Aa^3 + 66bBa^2 + 59Aa^2b + 48b^3B)) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{2a} \right. \right.$$

$$\left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d} \right)$$

↓ 25

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int \frac{(3ab(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3) + ab(16Aa^3 + 66bBa^2 + 59Aa^2b + 48b^3B)) \cos(c + dx) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{2a} - a(16a^2A + 54abB + 33Ab^2) \right. \right.$$

$$\left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int \frac{3ab(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3) + ab(16Aa^3 + 66bBa^2 + 59Aa^2b + 48b^3B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2a} - a(16a^2A + 54abB + 33Ab^2) \int \sqrt{a + b \cos(c + dx)} dx \right. \right.$$

$$\left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d} \right)$$

↓ 3134

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int \frac{3ab(8Ba^3 + 20Aba^2 + 30b^2Ba + 5Ab^3) + ab(16Aa^3 + 66bBa^2 + 59Aa^2b + 48b^3B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2a} - \frac{a(16a^2A + 54abB + 33Ab^2) \sqrt{a + b \cos(c + dx)}}{\sqrt{\frac{a + b \cos(c + dx)}{a + b}}} \right. \right.$$

$$\left. \frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int \frac{3ab(8Ba^3+20Aba^2+30b^2Ba+5Ab^3)+ab(16Aa^3+66bBa^2+59Ab^2a+48b^3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{a(16a^2A+54abB+33Ab^2)\sqrt{a+b\cos(c+dx)}}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3132

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int \frac{3ab(8Ba^3+20Aba^2+30b^2Ba+5Ab^3)+ab(16Aa^3+66bBa^2+59Ab^2a+48b^3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2a(16a^2A+54abB+33Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3481

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx + 3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2a(16a^2A+54abB+33Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + 3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2a(16a^2A+54abB+33Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3142

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\frac{a}{a+b} + \dots}{\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

2a

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\frac{a}{a+b} + \dots}{\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

2a

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3140

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}(\dots)}{d\sqrt{a+b\cos(c+dx)}}}{b} \right) \right)$$

2a

$$\frac{aA \tan(c+dx) \sec^2(c+dx)(a+b\cos(c+dx))^{3/2}}{3d}$$

↓ 3286

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx + \frac{2ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}\right)}{d\sqrt{a+b \cos(c+dx)}} \right) \right)$$

2a

$$\frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d}$$

3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{3ab(8a^3B+20a^2Ab+30ab^2B+5Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx + \frac{2ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}} \right) \right)$$

2a

$$\frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d}$$

3284

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(16a^2A + 54abB + 33Ab^2) \tan(c + dx) \sqrt{a + b \cos(c + dx)}}{d} + \frac{2ab(16a^3A+66a^2bB+59aAb^2+48b^3B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}\right)}{d\sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{aA \tan(c + dx) \sec^2(c + dx)(a + b \cos(c + dx))^{3/2}}{3d}$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^4,x]`

```
output (a*A*(a + b*cos[c + d*x])^(3/2)*Sec[c + d*x]^2*Tan[c + d*x]/(3*d) + ((3*a
*(3*A*b + 2*a*B)*Sqrt[a + b*cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*d)
+ (((-2*a*(16*a^2*A + 33*A*b^2 + 54*a*b*B)*Sqrt[a + b*cos[c + d*x]]*Ellip
ticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a + b*cos[c + d*x])/(a + b)]) +
((2*a*b*(16*a^3*A + 59*a*A*b^2 + 66*a^2*b*B + 48*b^3*B)*Sqrt[(a + b*cos[c
+ d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*cos
[c + d*x]]) + (6*a*b*(20*a^2*A*b + 5*A*b^3 + 8*a^3*B + 30*a*b^2*B)*Sqrt[(a
+ b*cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*
Sqrt[a + b*cos[c + d*x]]))/b)/(2*a) + ((16*a^2*A + 33*A*b^2 + 54*a*b*B)*Sq
rt[a + b*cos[c + d*x]]*Tan[c + d*x])/d)/4)/6
```

3.317.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*sin[c + d*x]]/Sqrt[(a + b*sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])]/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])]/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int[(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_)))/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x] *(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x] *(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int [(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.317.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2437 vs. $2(433) = 866$.

Time = 174.98 (sec) , antiderivative size = 2438, normalized size of antiderivative = 6.48

method	result	size
default	Expression too large to display	2438
parts	Expression too large to display	2878

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b^3*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+2*A*a^3*(-1/3*cos(1/2*d*x+1/2*c)/a*(-2*s
in(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2
*c)^2-1)^3+5/12*b/a^2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*
sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^2-1/24*(16*a^2+15*b
^2)/a^3*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/
2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)+5/48*b^2/a^2*(sin(1/2*d*x+1/2*c)^
2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*
c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*
b/(a-b))^(1/2))+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^
2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)
^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-1/3*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*
x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c
),(-2*b/(a-b))^(1/2))+1/3/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x
+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1
/2*c)^2)^(1/2)*b*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-5/16*b^2
/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b)...
```

3.317.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="fricas")`

output `Timed out`

3.317.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**4,x)`

output `Timed out`

3.317.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^4 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^4, x)`

3.317.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^4 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^4,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^4, x)`

3.317.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^4(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^4} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^4,x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^4, x)`

3.318 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^5(c+dx) dx$

3.318.1 Optimal result	2957
3.318.2 Mathematica [C] (verified)	2958
3.318.3 Rubi [A] (verified)	2959
3.318.4 Maple [B] (verified)	2969
3.318.5 Fricas [F(-1)]	2970
3.318.6 Sympy [F(-1)]	2970
3.318.7 Maxima [F]	2970
3.318.8 Giac [F]	2971
3.318.9 Mupad [F(-1)]	2971

3.318.1 Optimal result

Integrand size = 33, antiderivative size = 465

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx =$$

$$\frac{(284a^2 Ab + 15Ab^3 + 128a^3 B + 264ab^2 B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid \frac{2b}{a+b}\right)}{192ad \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{(356a^2 Ab + 133Ab^3 + 128a^3 B + 472ab^2 B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{192d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{(48a^4 A + 120a^2 Ab^2 - 5Ab^4 + 160a^3 bB + 40ab^3 B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{64ad \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{(284a^2 Ab + 15Ab^3 + 128a^3 B + 264ab^2 B) \sqrt{a + b \cos(c + dx)} \tan(c + dx)}{192ad}$$

$$+ \frac{(36a^2 A + 59Ab^2 + 104abB) \sqrt{a + b \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{96d}$$

$$+ \frac{a(11Ab + 8aB) \sqrt{a + b \cos(c + dx)} \sec^2(c + dx) \tan(c + dx)}{24d}$$

$$+ \frac{aA(a + b \cos(c + dx))^{3/2} \sec^3(c + dx) \tan(c + dx)}{4d}$$

output

$$\begin{aligned}
& -1/192*(284*A*a^2*b+15*A*b^3+128*B*a^3+264*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*(a+b*cos(d*x+c))^{(1/2)}/a/d/((a+b*cos(d*x+c))/(a+b))^{(1/2)}+1/192*(356 \\
& *A*a^2*b+133*A*b^3+128*B*a^3+472*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/d/(a+b*cos(d*x+c))^{(1/2)}+1/64*(48*A*a^4+120*A*a^2*b^2-5*A*b^4+160*B*a^3*b+40*B*a*b^3)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*cos(d*x+c))/(a+b))^{(1/2)}/a/d/(a+b*cos(d*x+c))^{(1/2)}+1/4*a*A*(a+b*cos(d*x+c))^{(3/2)}*sec(d*x+c)^3*tan(d*x+c)/d+1/192*(284*A*a^2*b+15*A*b^3+128*B*a^3+264*B*a*b^2)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/a/d+1/96*(36*A*a^2+59*A*b^2+104*B*a*b)*sec(d*x+c)*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d+1/24*a*(11*A*b+8*B*a)*sec(d*x+c)^2*(a+b*cos(d*x+c))^{(1/2)}*tan(d*x+c)/d
\end{aligned}$$

3.318.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.83 (sec) , antiderivative size = 549, normalized size of antiderivative = 1.18

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c$$

$$+ dx) dx = \frac{8b(36a^2A+59Ab^2+104abB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b\cos(c+dx)}} + \frac{2(288a^4A+436a^2Ab^2-45Ab^4+832a^3bB-24ab^3B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{a\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

output $((8*b*(36*a^2*A + 59*A*b^2 + 104*a*b*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)] * \text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(288 * a^4*A + 436*a^2*A*b^2 - 45*A*b^4 + 832*a^3*b*B - 24*a*b^3*B)*\text{Sqrt}[(a + b * \text{Cos}[c + d*x])/(a + b)] * \text{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)]/(a*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) - ((2*I)*(284*a^2*A*b + 15*A*b^3 + 128*a^3*B + 264*a * b^2*B)*\text{Sqrt}[-((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[(b*(1 + \text{Cos}[c + d*x]))/(-a + b)] * \text{Csc}[c + d*x]*(-2*a*(a - b)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^(-1)]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*\text{EllipticF}[I*\text{Arc Sinh}[\text{Sqrt}[-(a + b)^(-1)]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b * \text{EllipticPi}[(a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^(-1)]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b))))/(a^2*b*\text{Sqrt}[-(a + b)^(-1)]) + (4*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]*\text{Sec}[c + d*x]^3*((8*a^2*(17*A*b + 8*a*B) + (284*a^2*A*b + 15*A*b^3 + 128*a^3*B + 264*a*b^2*B)*\text{Cos}[c + d*x]^2)*\text{Sin}[c + d*x] + a*(36*a^2*A + 59 *A*b^2 + 104*a*b*B)*\text{Sin}[2*(c + d*x)] + 48*a^3*A*\text{Tan}[c + d*x]))/a)/(768*d)$

3.318.3 Rubi [A] (verified)

Time = 4.17 (sec) , antiderivative size = 485, normalized size of antiderivative = 1.04, number of steps used = 27, number of rules used = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^5(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^5} dx$$

$$\downarrow 3468$$

$$\frac{1}{4} \int \frac{1}{2} \sqrt{a + b \cos(c + dx)}(b(3aA + 8bB) \cos^2(c + dx) + 2(3Aa^2 + 8bBa + 4Ab^2) \cos(c + dx) + a(11Ab + 8aB)) \sec^3(c + dx)(a + b \cos(c + dx))^{3/2} dx + \frac{aA \tan(c + dx) \sec^3(c + dx)(a + b \cos(c + dx))^{3/2}}{4d}$$

$$\downarrow 27$$

$$\frac{1}{8} \int \frac{\sqrt{a + b \cos(c + dx)} (b(3aA + 8bB) \cos^2(c + dx) + 2(3Aa^2 + 8bBa + 4Ab^2) \cos(c + dx) + a(11Ab + 8aB)) \sec^3(c + dx) (a + b \cos(c + dx))^{3/2}}{4d} dx$$

↓ 3042

$$\frac{1}{8} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (b(3aA + 8bB) \sin^2(c + dx + \frac{\pi}{2}) + 2(3Aa^2 + 8bBa + 4Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(11Ab + 8aB)) \sec^3(c + dx) (a + b \cos(c + dx))^{3/2}}{4d} dx$$

↓ 3526

$$\frac{1}{8} \left(\frac{1}{3} \int \frac{(3b(8Ba^2 + 17Aba + 16b^2B) \cos^2(c + dx) + 2(16Ba^3 + 49Aba^2 + 72b^2Ba + 24Ab^3) \cos(c + dx) + a(36Aa^2 + 104bBa + 59Ab^2)) \sec^3(c + dx) (a + b \cos(c + dx))^{3/2}}{2\sqrt{a + b \cos(c + dx)} 4d} dx \right)$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \int \frac{(3b(8Ba^2 + 17Aba + 16b^2B) \cos^2(c + dx) + 2(16Ba^3 + 49Aba^2 + 72b^2Ba + 24Ab^3) \cos(c + dx) + a(36Aa^2 + 104bBa + 59Ab^2)) \sec^3(c + dx) (a + b \cos(c + dx))^{3/2}}{\sqrt{a + b \cos(c + dx)} 4d} dx \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \int \frac{3b(8Ba^2 + 17Aba + 16b^2B) \sin^2(c + dx + \frac{\pi}{2}) + 2(16Ba^3 + 49Aba^2 + 72b^2Ba + 24Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(36Aa^2 + 104bBa + 59Ab^2)}{\sin^3(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \frac{\sec^3(c + dx) (a + b \cos(c + dx))^{3/2}}{4d} dx \right)$$

↓ 3534

$$\frac{1}{8} \left(\frac{1}{6} \left(\int \frac{(ab(36Aa^2 + 104bBa + 59Ab^2) \cos^2(c + dx) + 2a(36Aa^3 + 152bBa^2 + 161Ab^2a + 96b^3B) \cos(c + dx) + a(128Ba^3 + 284Aba^2 + 264b^2Ba + 15Aa^2 + 104bBa + 59Ab^2)) \sec^3(c + dx) (a + b \cos(c + dx))^{3/2}}{2\sqrt{a + b \cos(c + dx)} 2a} dx \right) \right)$$

↓ 27

3.318. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\int \frac{ab(36Aa^2+104bBa+59Ab^2) \cos^2(c+dx)+2a(36Aa^3+152bBa^2+161Ab^2a+96b^3B) \cos(c+dx)+a(128Ba^3+284Aba^2+264b^2Ba+15Ab^3)}{\sqrt{a+b \cos(c+dx)}}}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\int \frac{ab(36Aa^2+104bBa+59Ab^2) \sin(c+dx+\frac{\pi}{2})^2+2a(36Aa^3+152bBa^2+161Ab^2a+96b^3B) \sin(c+dx+\frac{\pi}{2})+a(128Ba^3+284Aba^2+264b^2Ba+15Ab^3)}{\sin(c+dx+\frac{\pi}{2})^2 \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3534

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\int \frac{(2b(36Aa^2+104bBa+59Ab^2) \cos(c+dx)a^2-b(128Ba^3+284Aba^2+264b^2Ba+15Ab^3) \cos^2(c+dx)a+3(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a)}{2\sqrt{a+b \cos(c+dx)}}}{a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\int \frac{(2b(36Aa^2+104bBa+59Ab^2) \cos(c+dx)a^2-b(128Ba^3+284Aba^2+264b^2Ba+15Ab^3) \cos^2(c+dx)a+3(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a)}{\sqrt{a+b \cos(c+dx)}}}{2a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\int \frac{2b(36Aa^2+104bBa+59Ab^2) \sin(c+dx+\frac{\pi}{2})a^2-b(128Ba^3+284Aba^2+264b^2Ba+15Ab^3) \sin(c+dx+\frac{\pi}{2})^2a+3(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}}{2a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3538

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{-a(128a^3B+284a^2Ab+264ab^2B+15Ab^3) \int \sqrt{a+b \cos(c+dx)} dx - \frac{\int - \frac{(b(128Ba^3+356Aba^2+472b^2Ba+133Ab^3) \cos(c+dx)a^2+3b(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a}{\sqrt{a+b \cos(c+dx)}} dx}{2a}}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 25

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\frac{\int \frac{(b(128Ba^3+356Aba^2+472b^2Ba+133Ab^3) \cos(c+dx)a^2+3b(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a}{\sqrt{a+b \cos(c+dx)}} dx}{b} - a(128a^3B+284a^2Ab+264ab^2B+15Ab^3)}{2a}}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\frac{\int \frac{(b(128Ba^3+356Aba^2+472b^2Ba+133Ab^3) \sin(c+dx+\frac{\pi}{2})a^2+3b(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - a(128a^3B+284a^2Ab+264ab^2B+15Ab^3)}{2a}}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3134

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\frac{\int \frac{(b(128Ba^3+356Aba^2+472b^2Ba+133Ab^3) \sin(c+dx+\frac{\pi}{2})a^2+3b(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4)a}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - a(128a^3B+284a^2Ab+264ab^2B+15Ab^3)}{2a}}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\int \frac{b(128Ba^3+356Aba^2+472b^2Ba+133Ab^3) \sin(c+dx+\frac{\pi}{2}) a^2+3b(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4) a}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{a(128a^3B+284a^2Ab+264ab^2B+2a^4)}{2a} - \frac{4a}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3132

$$\frac{1}{8} \left(\frac{1}{6} \left(\int \frac{b(128Ba^3+356Aba^2+472b^2Ba+133Ab^3) \sin(c+dx+\frac{\pi}{2}) a^2+3b(48Aa^4+160bBa^3+120Ab^2a^2+40b^3Ba-5Ab^4) a}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(128a^3B+284a^2Ab+264ab^2B+2a^4)}{2a} - \frac{4a}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3481

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx+3ab(48a^4A+160a^3bB+120a^2Ab^2+40ab^3B-5Ab^4) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2a(128a^3B+284a^2Ab+264ab^2B+2a^4)}{2a} - \frac{4a}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx+3ab(48a^4A+160a^3bB+120a^2Ab^2+40ab^3B-5Ab^4) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(128a^3B+284a^2Ab+264ab^2B+2a^4)}{2a} - \frac{4a}{4a} \right) \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3142

3.318. $\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^5(c+dx) dx$

$$\left(\frac{1}{8} \right) \left(\frac{1}{6} \right) \left(\frac{a^2 b (128a^3 B + 356a^2 Ab + 472ab^2 B + 133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + 3ab (48a^4 A + 160a^3 bB + 120a^2 Ab^2 + 40ab^3 B - 5Ab^4) \int \frac{1}{\sin(c+dx)} dx \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) (a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\left(\frac{1}{8} \right) \left(\frac{1}{6} \right) \left(\frac{a^2 b (128a^3 B + 356a^2 Ab + 472ab^2 B + 133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + 3ab (48a^4 A + 160a^3 bB + 120a^2 Ab^2 + 40ab^3 B - 5Ab^4) \int \frac{1}{\sin(c+dx)} dx \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) (a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3140

$$\left(\frac{1}{8} \right) \left(\frac{1}{6} \right) \left(\frac{3ab (48a^4 A + 160a^3 bB + 120a^2 Ab^2 + 40ab^3 B - 5Ab^4) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \frac{2a^2 b (128a^3 B + 356a^2 Ab + 472ab^2 B + 133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d \sqrt{a+b \cos(c+dx)}}}{2a} \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx) (a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3286

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3ab(48a^4A+160a^3bB+120a^2Ab^2+40ab^3B-5Ab^4) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}} \right) + \frac{2a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}} \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{3ab(48a^4A+160a^3bB+120a^2Ab^2+40ab^3B-5Ab^4) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}} \right) + \frac{2a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}} \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3284

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(36a^2A + 104abB + 59Ab^2) \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2d} + \frac{(128a^3B+284a^2Ab+264ab^2B+15Ab^3)}{d} \right) + \frac{2a^2b(128a^3B+356a^2Ab+472ab^2B+133Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}} \right)$$

$$\frac{aA \tan(c+dx) \sec^3(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^5,x]`

```
output (a*A*(a + b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^3*Tan[c + d*x])/(4*d) + ((a*(
11*A*b + 8*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]^2*Tan[c + d*x])/(3*d
) + (((36*a^2*A + 59*A*b^2 + 104*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d
*x]*Tan[c + d*x])/(2*d) + (((-2*a*(284*a^2*A*b + 15*A*b^3 + 128*a^3*B + 26
4*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])
/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*a^2*b*(356*a^2*A*b + 133*A*b
^3 + 128*a^3*B + 472*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF
[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*a*b*(48*a^
4*A + 120*a^2*A*b^2 - 5*A*b^4 + 160*a^3*b*B + 40*a*b^3*B)*Sqrt[(a + b*Cos[
c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a +
b*Cos[c + d*x]]))/b)/(2*a) + ((284*a^2*A*b + 15*A*b^3 + 128*a^3*B + 264*a*
b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/d)/(4*a))/6)/8
```

3.318.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])]/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])]/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int((((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_))/((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x] *(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x] *(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int [(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.318.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3547 vs. $2(518) = 1036$.

Time = 458.62 (sec) , antiderivative size = 3548, normalized size of antiderivative = 7.63

method	result	size
default	Expression too large to display	3548
parts	Expression too large to display	4064

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*a^3*(-
1/4*cos(1/2*d*x+1/2*c)/a*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*
c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^4+7/24*b/a^2*cos(1/2*d*x+1/2*c)*(-2
*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1
/2*c)^2-1)^3-1/96*(36*a^2+35*b^2)/a^3*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1
/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^2+5
/192*b*(20*a^2+21*b^2)/a^4*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(
a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)-7/96*b/a*(sin(
1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*si
n(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),(-2*b/(a-b))^(1/2))-35/384*b^3/a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(
a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))
^(1/2))+25/96/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-
b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/
2)*b*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-25/96*b^2/a^2*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*si
n(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d
*x+1/2*c),(-2*b/(a-b))^(1/2))+35/128/a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*
b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a...
```


3.318.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="fricas")`

output `Timed out`

3.318.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**5,x)`

output `Timed out`

3.318.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^5 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^5, x)`

3.318.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^5 dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^5,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^5, x)`

3.318.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^5(c + dx) dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^5} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^5,x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^5, x)`

3.319 $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$

3.319.1 Optimal result 2972
 3.319.2 Mathematica [A] (verified) 2973
 3.319.3 Rubi [A] (verified) 2973
 3.319.4 Maple [B] (verified) 2979
 3.319.5 Fricas [C] (verification not implemented) 2980
 3.319.6 Sympy [F(-1)] 2981
 3.319.7 Maxima [F] 2981
 3.319.8 Giac [F] 2982
 3.319.9 Mupad [F(-1)] 2982

3.319.1 Optimal result

Integrand size = 33, antiderivative size = 320

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

$$= \frac{2(56a^2Ab + 63Ab^3 - 48a^3B - 44ab^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{105b^4d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$- \frac{2(56a^3Ab + 49aAb^3 - 48a^4B - 32a^2b^2B - 25b^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{105b^4d \sqrt{a+b \cos(c+dx)}}$$

$$- \frac{2(28aAb - 24a^2B - 25b^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{105b^3d}$$

$$+ \frac{2(7Ab - 6aB) \cos(c+dx) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{35b^2d}$$

$$+ \frac{2B \cos^2(c+dx) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{7bd}$$

output
$$\begin{aligned} & -2/105*(28*A*a*b-24*B*a^2-25*B*b^2)*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/b^3/ \\ & d+2/35*(7*A*b-6*B*a)*\cos(d*x+c)*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/b^2/d+2/ \\ & 7*B*\cos(d*x+c)^2*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/b/d+2/105*(56*A*a^2*b+6 \\ & 3*A*b^3-48*B*a^3-44*B*a*b^2)*(\cos(1/2*d*x+1/2*c)^(1/2) / \cos(1/2*d*x+1/2* \\ & c) * \text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2)) * (a+b*\cos(d*x+c))^(\\ & (1/2) / b^4/d / ((a+b*\cos(d*x+c)) / (a+b))^(1/2) - 2/105*(56*A*a^3*b+49*A*a*b^3-48 \\ & *B*a^4-32*B*a^2*b^2-25*B*b^4) * (\cos(1/2*d*x+1/2*c)^(1/2) / \cos(1/2*d*x+1/2 \\ & *c) * \text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2)) * ((a+b*\cos(d*x+c) \\ &) / (a+b))^(1/2) / b^4/d / (a+b*\cos(d*x+c))^(1/2) \end{aligned}$$

3.319.2 Mathematica [A] (verified)

Time = 1.24 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.72

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$= \frac{4\sqrt{\frac{a+b\cos(c+dx)}{a+b}}(b^2(14aAb-12a^2B+25b^2B)\text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - (-56a^2Ab-63Ab^3+48a^3B+}}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`

output
$$\begin{aligned} & (4*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*(b^2*(14*a*A*b - 12*a^2*B + 25*b^2*B) \\ &)*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)] - (-56*a^2*A*b - 63*A*b^3 + 48*a^3 \\ & *B + 44*a*b^2*B)*((a + b)*\text{EllipticE}[(c + d*x)/2, (2*b)/(a + b)] - a*\text{Elliptic} \\ & \text{F}[(c + d*x)/2, (2*b)/(a + b)]) + 2*b*(a + b*\text{Cos}[c + d*x])*(-56*a*A*b + \\ & 48*a^2*B + 65*b^2*B + 6*b*(7*A*b - 6*a*B)*\text{Cos}[c + d*x] + 15*b^2*B*\text{Cos}[2*(c \\ & + d*x)])*\text{Sin}[c + d*x])/(210*b^4*d*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) \end{aligned}$$

3.319.3 Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.06, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.319.
$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$\begin{aligned}
& \int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3469} \\
& \frac{2 \int \frac{\cos(c+dx)((7Ab-6aB)\cos^2(c+dx)+5bB\cos(c+dx)+4aB)}{2\sqrt{a+b\cos(c+dx)}} dx}{\frac{7b}{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}} + \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cos(c+dx)((7Ab-6aB)\cos^2(c+dx)+5bB\cos(c+dx)+4aB)}{\sqrt{a+b\cos(c+dx)}} dx}{\frac{7b}{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}} + \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})((7Ab-6aB)\sin(c+dx+\frac{\pi}{2})^2+5bB\sin(c+dx+\frac{\pi}{2})+4aB)}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\frac{7b}{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}} + \\
& \quad \downarrow \text{3528} \\
& \frac{2 \int \frac{-((-24Ba^2+28Aba-25b^2B)\cos^2(c+dx)+b(21Ab+2aB)\cos(c+dx)+2a(7Ab-6aB))}{2\sqrt{a+b\cos(c+dx)}} dx}{\frac{7b}{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}} + \frac{2(7Ab-6aB)\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} + \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{-((-24Ba^2+28Aba-25b^2B)\cos^2(c+dx)+b(21Ab+2aB)\cos(c+dx)+2a(7Ab-6aB))}{\sqrt{a+b\cos(c+dx)}} dx}{\frac{7b}{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}} + \frac{2(7Ab-6aB)\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} + \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.319. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

$$\frac{\int \frac{(24Ba^2 - 28Aba + 25b^2B) \sin(c+dx + \frac{\pi}{2})^2 + b(21Ab + 2aB) \sin(c+dx + \frac{\pi}{2}) + 2a(7Ab - 6aB)}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{5b} + \frac{2(7Ab - 6aB) \sin(c+dx) \cos(c+dx) \sqrt{a+b \cos(c+dx)}}{5bd}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

↓ 3502

$$2 \int \frac{b(-12Ba^2 + 14Aba + 25b^2B) + (-48Ba^3 + 56Aba^2 - 44b^2Ba + 63Ab^3) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-24a^2B + 28aAb - 25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} + \frac{2(7Ab - 6aB)}{3}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

↓ 27

$$\frac{\int \frac{b(-12Ba^2 + 14Aba + 25b^2B) + (-48Ba^3 + 56Aba^2 - 44b^2Ba + 63Ab^3) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-24a^2B + 28aAb - 25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}}{5b} + \frac{2(7Ab - 6aB)}{3}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

↓ 3042

$$\frac{\int \frac{b(-12Ba^2 + 14Aba + 25b^2B) + (-48Ba^3 + 56Aba^2 - 44b^2Ba + 63Ab^3) \sin(c+dx + \frac{\pi}{2})}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{2(-24a^2B + 28aAb - 25b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}}{5b} + \frac{2(7Ab - 6aB)}{3}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

↓ 3231

$$\frac{(-48a^3B + 56a^2Ab - 44ab^2B + 63Ab^3) \int \sqrt{a+b \cos(c+dx)} dx - \frac{(-48a^4B + 56a^3Ab - 32a^2b^2B + 49aAb^3 - 25b^4B)}{b} \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{3b} - \frac{2(-24a^2B + 28aAb - 25b^2B)}{3}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

↓ 3042

3.319. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$

$$\frac{(-48a^3B+56a^2Ab-44ab^2B+63Ab^3) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-24a^2B+28aAb-12b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5b}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

7b

↓ 3134

$$\frac{(-48a^3B+56a^2Ab-44ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-24a^2B+28aAb-12b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5b}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

7b

↓ 3042

$$\frac{(-48a^3B+56a^2Ab-44ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-24a^2B+28aAb-12b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5b}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

7b

↓ 3132

$$\frac{2(-48a^3B+56a^2Ab-44ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-24a^2B+28aAb-12b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5b}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

7b

↓ 3142

$$\frac{2(-48a^3B+56a^2Ab-44ab^2B+63Ab^3) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B) \int \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-24a^2B+28aAb-12b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5b}$$

$$\frac{2B \sin(c+dx) \cos^2(c+dx) \sqrt{a+b \cos(c+dx)}}{7bd}$$

7b

↓ 3042

3.319. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$

$$\frac{2\left(-48a^3B+56a^2Ab-44ab^2B+63Ab^3\right)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)-\frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}}}{b\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}{7bd}$$

↓ 3140

$$\frac{2\left(-48a^3B+56a^2Ab-44ab^2B+63Ab^3\right)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)-\frac{(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{bd\sqrt{a+b\cos(c+dx)}}}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{2\left(-48a^4B+56a^3Ab-32a^2b^2B+49aAb^3-25b^4B\right)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{bd\sqrt{a+b\cos(c+dx)}}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx)\right)}{bd\sqrt{a+b\cos(c+dx)}}-\frac{\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}}}{b\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2B\sin(c+dx)\cos^2(c+dx)\sqrt{a+b\cos(c+dx)}}{7bd}$$

```
input Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]
```

```
output (2*B*Cos[c + d*x]^2*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*b*d) + ((2*(7*A*b - 6*a*B)*Cos[c + d*x]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*b*d) + (((2*(56*a^2*A*b + 63*A*b^3 - 48*a^3*B - 44*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(56*a^3*A*b + 49*a*A*b^3 - 48*a^4*B - 32*a^2*b^2*B - 25*b^4*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(3*b) - (2*(28*a*A*b - 24*a^2*B - 25*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*b*d))/(5*b))/(7*b)
```

3.319.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```


rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*SIN[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*SIN[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*SIN[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*SIN[e + f*x
])^m*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*SIN[e + f*x])^(m - 1)*(c + d*SIN[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1))*SIN[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*SIN[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

3.319.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1304 vs. $2(354) = 708$.

Time = 13.14 (sec) , antiderivative size = 1305, normalized size of antiderivative = 4.08

method	result	size
default	Expression too large to display	1305
parts	Expression too large to display	1494

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNV
ERBOSE)
```

output

```
-2/105*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^4+(-168*A*b^4+24*B*a*b^3-360*B*b
^4)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(-28*A*a*b^3+168*A*b^4+24*B*a^
2*b^2-24*B*a*b^3+280*B*b^4)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(56*A*
a^2*b^2+14*A*a*b^3-42*A*b^4-48*B*a^3*b-12*B*a^2*b^2-44*B*a*b^3-80*B*b^4)*s
in(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-56*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1
/2*c),(-2*b/(a-b))^(1/2))*a^3*b-49*a*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/
(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))*b^3+56*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin
(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a
-b))^(1/2))*a^3*b-56*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*
x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1
/2))*a^2*b^2+63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2
*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*
a*b^3-63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(
a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^4+48*
B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b
))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4+32*B*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)...
```

3.319.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.18 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.76

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$= \frac{\sqrt{2}(-96iBa^4 + 112iAa^3b - 52iBa^2b^2 + 84iAab^3 - 75iBb^4)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4(4a^2-3b^2)}{3b^2}, -\frac{8(8a^3-27b^3)}{27b^3}\right)}{...}$$

input

```
integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm
m="fricas")
```

```
output 1/315*(sqrt(2)*(-96*I*B*a^4 + 112*I*A*a^3*b - 52*I*B*a^2*b^2 + 84*I*A*a*b^3 - 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(96*I*B*a^4 - 112*I*A*a^3*b + 52*I*B*a^2*b^2 - 84*I*A*a*b^3 + 75*I*B*b^4)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(48*I*B*a^3*b - 56*I*A*a^2*b^2 + 44*I*B*a*b^3 - 63*I*A*b^4)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(-48*I*B*a^3*b + 56*I*A*a^2*b^2 - 44*I*B*a*b^3 + 63*I*A*b^4)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(15*B*b^4*cos(d*x + c)^2 + 24*B*a^2*b^2 - 28*A*a*b^3 + 25*B*b^4 - 3*(6*B*a*b^3 - 7*A*b^4)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^5*d)
```

3.319.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(1/2),x)
```

```
output Timed out
```

3.319.7 Maxima [F]

$$\int \frac{\cos^3(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^3}{\sqrt{b \cos(dx + c) + a}} dx$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*cos(d*x + c)^3/sqrt(b*cos(d*x + c) + a), x)
```

3.319. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$

3.319.8 Giac [F]

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^3}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^3/sqrt(b*cos(d*x + c) + a), x)`

3.319.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{\cos(c+dx)^3(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

output `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2), x)`

3.320
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

3.320.1 Optimal result 2983
 3.320.2 Mathematica [A] (verified) 2984
 3.320.3 Rubi [A] (verified) 2984
 3.320.4 Maple [B] (verified) 2989
 3.320.5 Fracas [C] (verification not implemented) 2990
 3.320.6 Sympy [F] 2990
 3.320.7 Maxima [F] 2991
 3.320.8 Giac [F] 2991
 3.320.9 Mupad [F(-1)] 2991

3.320.1 Optimal result

Integrand size = 33, antiderivative size = 246

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

$$= -\frac{2(10aAb - 8a^2B - 9b^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{15b^3d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(10a^2Ab + 5Ab^3 - 8a^3B - 7ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{15b^3d \sqrt{a+b \cos(c+dx)}}$$

$$+ \frac{2(5Ab - 4aB) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15b^2d}$$

$$+ \frac{2B \cos(c+dx) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{5bd}$$

output
$$\frac{2}{15} \cdot (5A \cdot b - 4B \cdot a) \cdot \sin(d \cdot x + c) \cdot (a + b \cdot \cos(d \cdot x + c))^{1/2} / b^2 / d + 2/5 \cdot B \cdot \cos(d \cdot x + c) \cdot \sin(d \cdot x + c) \cdot (a + b \cdot \cos(d \cdot x + c))^{1/2} / b / d - 2/15 \cdot (10A \cdot a \cdot b - 8B \cdot a^2 - 9B \cdot b^2) \cdot (\cos(1/2 \cdot d \cdot x + 1/2 \cdot c))^{1/2} / \cos(1/2 \cdot d \cdot x + 1/2 \cdot c) \cdot \text{EllipticE}(\sin(1/2 \cdot d \cdot x + 1/2 \cdot c), 2^{1/2} \cdot (b / (a + b))^{1/2}) \cdot (a + b \cdot \cos(d \cdot x + c))^{1/2} / b^3 / d / ((a + b \cdot \cos(d \cdot x + c)) / (a + b))^{1/2} + 2/15 \cdot (10A \cdot a^2 \cdot b + 5A \cdot b^3 - 8B \cdot a^3 - 7B \cdot a \cdot b^2) \cdot (\cos(1/2 \cdot d \cdot x + 1/2 \cdot c))^{1/2} / \cos(1/2 \cdot d \cdot x + 1/2 \cdot c) \cdot \text{EllipticF}(\sin(1/2 \cdot d \cdot x + 1/2 \cdot c), 2^{1/2} \cdot (b / (a + b))^{1/2}) \cdot (a + b \cdot \cos(d \cdot x + c)) / (a + b)^{1/2} / b^3 / d / (a + b \cdot \cos(d \cdot x + c))^{1/2}$$

3.320.2 Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.73

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2\sqrt{\frac{a+b\cos(c+dx)}{a+b}} (b^2(5Ab + 2aB) \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + (-10aAb + 8a^2B + 9b^2B) ((a + b)E\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) - a\text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)) + 2*b*(a + b*\cos[c + d*x])*(5*A*b - 4*a*B + 3*b*B*\cos[c + d*x])*Sin[c + d*x])}{15b^3d\sqrt{a + b\cos(c + dx)}}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*(b^2*(5*A*b + 2*a*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + (-10*a*A*b + 8*a^2*B + 9*b^2*B)*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)])) + 2*b*(a + b*Cos[c + d*x])*(5*A*b - 4*a*B + 3*b*B*Cos[c + d*x])*Sin[c + d*x])/(15*b^3*d*Sqrt[a + b*Cos[c + d*x]])`

3.320.3 Rubi [A] (verified)

Time = 1.23 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{\sin\left(c + dx + \frac{\pi}{2}\right)^2 (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx$$

↓ 3469

$$\frac{2 \int \frac{(5Ab - 4aB) \cos^2(c + dx) + 3bB \cos(c + dx) + 2aB}{2\sqrt{a + b \cos(c + dx)}} dx}{5b} + \frac{2B \sin(c + dx) \cos(c + dx) \sqrt{a + b \cos(c + dx)}}{5bd}$$

↓ 27

3.320. $\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$

$$\begin{aligned}
 & \frac{\int \frac{(5Ab-4aB)\cos^2(c+dx)+3bB\cos(c+dx)+2aB}{\sqrt{a+b\cos(c+dx)}} dx}{5b} + \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(5Ab-4aB)\sin(c+dx+\frac{\pi}{2})^2+3bB\sin(c+dx+\frac{\pi}{2})+2aB}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{5b} + \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} \\
 & \quad \downarrow \text{3502} \\
 & \frac{2\int \frac{b(5Ab+2aB)-(-8Ba^2+10Aba-9b^2B)\cos(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx}{3b} + \frac{2(5Ab-4aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd} + \\
 & \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{b(5Ab+2aB)-(-8Ba^2+10Aba-9b^2B)\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{3b} + \frac{2(5Ab-4aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd} + \\
 & \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{b(5Ab+2aB)+(8Ba^2-10Aba+9b^2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2(5Ab-4aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd} + \\
 & \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} \\
 & \quad \downarrow \text{3231} \\
 & \frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3)\int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{(-8a^2B+10aAb-9b^2B)\int \sqrt{a+b\cos(c+dx)} dx}{b} + \frac{2(5Ab-4aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd} + \\
 & \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3)\int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(-8a^2B+10aAb-9b^2B)\int \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{2(5Ab-4aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd} + \\
 & \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}
 \end{aligned}$$

3.320. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

↓ 3134

$$\frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(-8a^2B+10aAb-9b^2B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(5Ab-4aB) \sin(c+dx)}{3bd}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) \sqrt{a+b \cos(c+dx)}}{5bd}$$

↓ 3042

$$\frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(-8a^2B+10aAb-9b^2B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(5Ab-4aB) \sin(c+dx)}{3bd}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) \sqrt{a+b \cos(c+dx)}}{5bd}$$

↓ 3132

$$\frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-8a^2B+10aAb-9b^2B) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(5Ab-4aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) \sqrt{a+b \cos(c+dx)}}{5bd}$$

↓ 3142

$$\frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-8a^2B+10aAb-9b^2B) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(5Ab-4aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) \sqrt{a+b \cos(c+dx)}}{5bd}$$

↓ 3042

$$\frac{(-8a^3B+10a^2Ab-7ab^2B+5Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(-8a^2B+10aAb-9b^2B) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(5Ab-4aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}$$

$$\frac{2B \sin(c+dx) \cos(c+dx) \sqrt{a+b \cos(c+dx)}}{5bd}$$

3.320. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$

↓ 3140

$$\frac{2(-8a^3B+10a^2Ab-7ab^2B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2(-8a^2B+10aAb-9b^2B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{a+b\cos(c+dx)}} + \frac{2(5Ab-4aB)}{3b}$$

$$\frac{2B\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*B*Cos[c + d*x]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*b*d) + (((-2*(10*a*A*b - 8*a^2*B - 9*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(10*a^2*A*b + 5*A*b^3 - 8*a^3*B - 7*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(3*b) + (2*(5*A*b - 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*b*d)/(5*b)`

3.320.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b]*Sin[c + d*x]/Sqrt[(a + b)*Sin[c + d*x])/(a + b) Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B))*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.320.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 992 vs. $2(284) = 568$.

Time = 11.78 (sec) , antiderivative size = 993, normalized size of antiderivative = 4.04

method	result	size
default	Expression too large to display	993
parts	Expression too large to display	1120

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/15*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*c
os(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^3+(20*A*b^3-4*B*a*b^2+24*B*b^3)*s
in(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*a*b^2-10*A*b^3+8*B*a^2*b+2*B
*a*b^2-6*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+10*A*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*Ellipt
icF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+5*A*b^3*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(
cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1
/2*c),(-2*b/(a-b))^(1/2))*a^2*b+10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a
-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-
2*b/(a-b))^(1/2))*a*b^2-8*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(
1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-
b))^(1/2))*a^3-7*B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+
1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2
))*b^2+8*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(
a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3-8*B
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b
))^^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2*b+9*B*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*...
```

3.320.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 493, normalized size of antiderivative = 2.00

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$= \frac{\sqrt{2}(16iBa^3 - 20iAa^2b + 12iBab^2 - 15iAb^3)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4(4a^2-3b^2)}{3b^2}, -\frac{8(8a^3-9ab^2)}{27b^3}, \frac{3b\cos(dx+c)}{b^3}\right)}{b^4}$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/45*(sqrt(2)*(16*I*B*a^3 - 20*I*A*a^2*b + 12*I*B*a*b^2 - 15*I*A*b^3)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(-16*I*B*a^3 + 20*I*A*a^2*b - 12*I*B*a*b^2 + 15*I*A*b^3)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(-8*I*B*a^2*b + 10*I*A*a*b^2 - 9*I*B*b^3)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(8*I*B*a^2*b - 10*I*A*a*b^2 + 9*I*B*b^3)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)) + 6*(3*B*b^3*cos(d*x + c) - 4*B*a*b^2 + 5*A*b^3)*sqrt(b*cos(d*x + c) + a)*sin(d*x + c))/(b^4*d)`

3.320.6 Sympy [F]

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(A+B\cos(c+dx))\cos^2(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*cos(c + d*x)**2/sqrt(a + b*cos(c + d*x)), x)`

3.320. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

3.320.7 Maxima [F]

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^2}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^2/sqrt(b*cos(d*x + c) + a), x)`

3.320.8 Giac [F]

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^2}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^2/sqrt(b*cos(d*x + c) + a), x)`

3.320.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{\cos(c+dx)^2(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

output `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2), x)`

3.321
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

3.321.1 Optimal result 2992
 3.321.2 Mathematica [A] (verified) 2993
 3.321.3 Rubi [A] (verified) 2993
 3.321.4 Maple [B] (verified) 2997
 3.321.5 Fricas [C] (verification not implemented) 2998
 3.321.6 Sympy [F] 2999
 3.321.7 Maxima [F] 2999
 3.321.8 Giac [F] 3000
 3.321.9 Mupad [B] (verification not implemented) 3000

3.321.1 Optimal result

Integrand size = 31, antiderivative size = 183

$$\begin{aligned} & \int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx \\ &= \frac{2(3Ab-2aB)\sqrt{a+b \cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{3b^2d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\ & \quad - \frac{2(3aAb-2a^2B-b^2B)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3b^2d\sqrt{a+b \cos(c+dx)}} \\ & \quad + \frac{2B\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3bd} \end{aligned}$$

```
output 2/3*B*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/b/d+2/3*(3*A*b-2*B*a)*(cos(1/2*d*x
+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(
b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2
)-2/3*(3*A*a*b-2*B*a^2-B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2
*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c)
)/(a+b))^(1/2)/b^2/d/(a+b*cos(d*x+c))^(1/2)
```

3.321.2 Mathematica [A] (verified)

Time = 1.31 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.84

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$= \frac{-2(a+b)(-3Ab+2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c+dx)\left|\frac{2b}{a+b}\right.\right) + 2(-3aAb+2a^2B+b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticE}\left(\frac{c+dx}{2}, \frac{2b}{a+b}\right) + 3b^2d\sqrt{a+b\cos(c+dx)}}{3b^2d\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`output `(-2*(a + b)*(-3*A*b + 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)] + 2*(-3*a*A*b + 2*a^2*B + b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + 2*b*B*(a + b)*Cos[c + d*x]*Sin[c + d*x])/(3*b^2*d*Sqrt[a + b*Cos[c + d*x]])`**3.321.3 Rubi [A] (verified)**Time = 0.95 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx$$

$$\downarrow \text{3447}$$

$$\int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A\sin(c+dx+\frac{\pi}{2})+B\sin(c+dx+\frac{\pi}{2})^2}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx$$

3.321. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow \text{3502} \\
 & \frac{2 \int \frac{bB+(3Ab-2aB) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{3b} + \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{27} \\
 & \frac{\int \frac{bB+(3Ab-2aB) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{3b} + \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{3042} \\
 & \frac{\int \frac{bB+(3Ab-2aB) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{3231} \\
 & \frac{(3Ab-2aB) \int \frac{\sqrt{a+b \cos(c+dx)}}{b} dx}{3b} - \frac{(-2a^2B+3aAb-b^2B) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} + \\
 & \quad \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{3042} \\
 & \frac{(3Ab-2aB) \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{b} dx}{3b} - \frac{(-2a^2B+3aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \\
 & \quad \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{3134} \\
 & \frac{(3Ab-2aB) \sqrt{a+b \cos(c+dx)} \int \frac{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} dx}{3b} - \frac{(-2a^2B+3aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \\
 & \quad \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{3042} \\
 & \frac{(3Ab-2aB) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(-2a^2B+3aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \\
 & \quad \frac{2B \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd} \\
 & \downarrow \text{3132}
 \end{aligned}$$

3.321. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
 & \frac{2(3Ab-2aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(-2a^2B+3aAb-b^2B)\int\frac{1}{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}}dx}{b} \\
 & \qquad \qquad \qquad + \\
 & \qquad \qquad \qquad \frac{3b}{2B\sin(c+dx)\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad \frac{3bd}{3bd} \\
 & \qquad \qquad \qquad \downarrow \text{3142} \\
 & \frac{2(3Ab-2aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(-2a^2B+3aAb-b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}}}{b\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad + \\
 & \qquad \qquad \qquad \frac{3b}{2B\sin(c+dx)\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad \frac{3bd}{3bd} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{2(3Ab-2aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(-2a^2B+3aAb-b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin\left(c+dx+\frac{\pi}{2}\right)}{a+b}}}}{b\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad + \\
 & \qquad \qquad \qquad \frac{3b}{2B\sin(c+dx)\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad \frac{3bd}{3bd} \\
 & \qquad \qquad \qquad \downarrow \text{3140} \\
 & \frac{2(3Ab-2aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{2(-2a^2B+3aAb-b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\text{EllipticF}\left(\frac{1}{2}(c+dx),\frac{2b}{a+b}\right)}{bd\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad + \\
 & \qquad \qquad \qquad \frac{3b}{2B\sin(c+dx)\sqrt{a+b\cos(c+dx)}} \\
 & \qquad \qquad \qquad \frac{3bd}{3bd}
 \end{aligned}$$

```
input Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]
```

```
output ((2*(3*A*b - 2*a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(3*a*A*b - 2*a^2*B - b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(3*b) + (2*B*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*b*d)
```

3.321.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.321.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 670 vs. 2(225) = 450.

Time = 10.13 (sec) , antiderivative size = 671, normalized size of antiderivative = 3.67

method	result
default	$2\sqrt{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(-4B\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^2 + 3Aab\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} F\left(\cos\left(\frac{dx}{2}\right)\right.\right.$
parts	$\frac{2A\sqrt{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} \left(F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right), \sqrt{-\frac{2b}{a - b}}\right) a - E\left(\cos\left(\frac{dx}{2}\right)\right.\right.}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b + (a + b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a +}}$

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2), x, method=_RETURNVER
BOSE)
```

output

```

2/3*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B*cos(
1/2*d*x+1/2*c)^5*b^2+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/
2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*
a*b-3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b)
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b+3*A*(sin(1/2*
d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE
(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^2-2*B*cos(1/2*d*x+1/2*c)^3*a*b+6
*B*cos(1/2*d*x+1/2*c)^3*b^2-2*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2
*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^
(1/2))*a^2-B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a
-b)/(a-b))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+2*B*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*Elli
pticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^2-2*B*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x
+1/2*c),(-2*b/(a-b))^(1/2))*a*b+2*B*cos(1/2*d*x+1/2*c)*a*b-2*B*cos(1/2*d*x
+1/2*c)*b^2)/b^2/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1
/2)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d

```

3.321.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.38

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$= \frac{6\sqrt{b\cos(dx+c)+a}Bb^2\sin(dx+c)+\sqrt{2}(-4iBa^2+6iAab-3iBb^2)\sqrt{b}\text{weierstrassPInverse}\left(\frac{4(4a^2-3b^2)}{3b^2}\right)}{...}$$

input

```

integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm=
"fricas")

```

output `1/9*(6*sqrt(b*cos(d*x + c) + a)*B*b^2*sin(d*x + c) + sqrt(2)*(-4*I*B*a^2 + 6*I*A*a*b - 3*I*B*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + sqrt(2)*(4*I*B*a^2 - 6*I*A*a*b + 3*I*B*b^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*sqrt(2)*(2*I*B*a*b - 3*I*A*b^2)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*sqrt(2)*(-2*I*B*a*b + 3*I*A*b^2)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)))/(b^3*d)`

3.321.6 Sympy [F]

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(1/2), x)`

output `Integral((A + B*cos(c + d*x))*cos(c + d*x)/sqrt(a + b*cos(c + d*x)), x)`

3.321.7 Maxima [F]

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)/sqrt(b*cos(d*x + c) + a), x)`

3.321.8 Giac [F]

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)/sqrt(b*cos(d*x + c) + a), x)`

3.321.9 Mupad [B] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.09

$$\begin{aligned} & \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx \\ &= \frac{2B\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd} \\ &+ \frac{2A\left(E\left(\frac{c}{2} + \frac{dx}{2} \middle| \frac{2b}{a+b}\right)(a+b) - aF\left(\frac{c}{2} + \frac{dx}{2} \middle| \frac{2b}{a+b}\right)\right)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{bd\sqrt{a+b\cos(c+dx)}} \\ &+ \frac{2B\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\left(F\left(\frac{c}{2} + \frac{dx}{2} \middle| \frac{2b}{a+b}\right)(2a^2+b^2) - 2aE\left(\frac{c}{2} + \frac{dx}{2} \middle| \frac{2b}{a+b}\right)(a+b)\right)}{3b^2d\sqrt{a+b\cos(c+dx)}} \end{aligned}$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

output `(2*B*sin(c + d*x)*(a + b*cos(c + d*x))^(1/2))/(3*b*d) + (2*A*(ellipticE(c/2 + (d*x)/2, (2*b)/(a + b))*(a + b) - a*ellipticF(c/2 + (d*x)/2, (2*b)/(a + b)))*((a + b*cos(c + d*x))/(a + b))^(1/2))/(b*d*(a + b*cos(c + d*x))^(1/2)) + (2*B*((a + b*cos(c + d*x))/(a + b))^(1/2)*(ellipticF(c/2 + (d*x)/2, (2*b)/(a + b))*(2*a^2 + b^2) - 2*a*ellipticE(c/2 + (d*x)/2, (2*b)/(a + b))*(a + b)))/(3*b^2*d*(a + b*cos(c + d*x))^(1/2))`

3.322 $\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

3.322.1 Optimal result	3001
3.322.2 Mathematica [A] (verified)	3001
3.322.3 Rubi [A] (verified)	3002
3.322.4 Maple [A] (verified)	3004
3.322.5 Fricas [C] (verification not implemented)	3005
3.322.6 Sympy [F]	3006
3.322.7 Maxima [F]	3006
3.322.8 Giac [F]	3006
3.322.9 Mupad [B] (verification not implemented)	3007

3.322.1 Optimal result

Integrand size = 25, antiderivative size = 130

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \frac{2B\sqrt{a + b \cos(c + dx)}E\left(\frac{1}{2}(c + dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2(Ab - aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{bd\sqrt{a + b \cos(c + dx)}}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b/d/(a+b*cos(d*x+c))^(1/2)`

3.322.2 Mathematica [A] (verified)

Time = 2.99 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.72

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \frac{2\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \left((a + b)BE\left(\frac{1}{2}(c + dx)\middle|\frac{2b}{a+b}\right) + (Ab - aB) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) \right)}{bd\sqrt{a + b \cos(c + dx)}}$$

input `Integrate[(A + B*Cos[c + d*x])/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*((a + b)*B*EllipticE[(c + d*x)/2, (2*b)/(a + b)] + (A*b - a*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)]))/(b*d*Sqrt[a + b*Cos[c + d*x]])`

3.322.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3231} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx}{b} + \frac{B \int \sqrt{a + b \cos(c + dx)} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{b} + \frac{B \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx}{b} \\
 & \quad \downarrow \text{3134} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{b} + \frac{B \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a + b} + \frac{b \cos(c + dx)}{a + b}} dx}{b \sqrt{\frac{a + b \cos(c + dx)}{a + b}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{b} + \frac{B \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a + b} + \frac{b \sin\left(c + dx + \frac{\pi}{2}\right)}{a + b}} dx}{b \sqrt{\frac{a + b \cos(c + dx)}{a + b}}}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3132} \\
& \frac{(Ab - aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \downarrow \text{3142} \\
& \frac{(Ab - aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} + \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \downarrow \text{3042} \\
& \frac{(Ab - aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} + \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
& \downarrow \text{3140} \\
& \frac{2(Ab - aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd \sqrt{a+b \cos(c+dx)}} + \frac{2B \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*B*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(A*b - a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]])`

3.322.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

3.322.4 Maple [A] (verified)

Time = 6.03 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.92

method	result
default	$-\frac{2\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (a+b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} \left(AbF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a-b}}\right) - BF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a-b}}\right) \right)$
parts	$\frac{2A\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - a - b}{a + b}}}{d\sqrt{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}} \operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2} \mid \frac{\sqrt{2}\sqrt{b}}{\sqrt{a+b}}\right) + \frac{2B\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (a+b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}}$
risch	Expression too large to display

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

output
$$-2*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}*(A*b*\text{EllipticF}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})-B*\text{EllipticF}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})*a+B*\text{EllipticE}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})*a-B*\text{EllipticE}(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})*b)/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b/\sin(1/2*d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2+a+b)^{(1/2)}/d$$

3.322.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 371, normalized size of antiderivative = 2.85

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{3i\sqrt{2}Bb^{\frac{3}{2}}\text{weierstrassZeta}\left(\frac{4(4a^2-3b^2)}{3b^2}, -\frac{8(8a^3-9ab^2)}{27b^3}\right), \text{weierstrassPInverse}\left(\frac{4(4a^2-3b^2)}{3b^2}, -\frac{8(8a^3-9ab^2)}{27b^3}, \frac{3b\cos(c+dx)}{a+b\cos(c+dx)}\right)}{1}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output
$$\frac{1}{3}*(3*I*\sqrt{2})*B*b^{(3/2)}*\text{weierstrassZeta}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, \text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) + 3*I*b*\sin(d*x + c) + 2*a)/b)) - 3*I*\sqrt{2})*B*b^{(3/2)}*\text{weierstrassZeta}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, \text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) - 3*I*b*\sin(d*x + c) + 2*a)/b)) + \sqrt{2}*(2*I*B*a - 3*I*A*b)*\sqrt{b}*\text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) + 3*I*b*\sin(d*x + c) + 2*a)/b) + \sqrt{2}*(-2*I*B*a + 3*I*A*b)*\sqrt{b}*\text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) - 3*I*b*\sin(d*x + c) + 2*a)/b))/(b^2*d)$$

3.322.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/sqrt(a + b*cos(c + d*x)), x)`

3.322.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/sqrt(b*cos(d*x + c) + a), x)`

3.322.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/sqrt(b*cos(d*x + c) + a), x)`

3.322.9 Mupad [B] (verification not implemented)

Time = 1.02 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.04

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2 A F\left(\frac{c}{2} + \frac{dx}{2} \mid \frac{2b}{a+b}\right) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d \sqrt{a + b \cos(c + dx)}} + \frac{2 B \left(E\left(\frac{c}{2} + \frac{dx}{2} \mid \frac{2b}{a+b}\right) (a + b) - a F\left(\frac{c}{2} + \frac{dx}{2} \mid \frac{2b}{a+b}\right) \right) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b d \sqrt{a + b \cos(c + dx)}}$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^(1/2),x)`output `(2*A*ellipticF(c/2 + (d*x)/2, (2*b)/(a + b))*((a + b*cos(c + d*x))/(a + b))^(1/2))/(d*(a + b*cos(c + d*x))^(1/2)) + (2*B*(ellipticE(c/2 + (d*x)/2, (2*b)/(a + b))*(a + b) - a*ellipticF(c/2 + (d*x)/2, (2*b)/(a + b)))*((a + b*cos(c + d*x))/(a + b))^(1/2))/(b*d*(a + b*cos(c + d*x))^(1/2))`

3.323
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.323.1 Optimal result 3008
 3.323.2 Mathematica [A] (verified) 3008
 3.323.3 Rubi [A] (verified) 3009
 3.323.4 Maple [A] (verified) 3011
 3.323.5 Fracas [F(-1)] 3012
 3.323.6 Sympy [F] 3012
 3.323.7 Maxima [F] 3012
 3.323.8 Giac [F] 3013
 3.323.9 Mupad [F(-1)] 3013

3.323.1 Optimal result

Integrand size = 31, antiderivative size = 118

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}} + \frac{2A \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+
1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(
d*x+c))^(1/2)+2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
Pi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(
1/2)/d/(a+b*cos(d*x+c))^(1/2)
```

3.323.2 Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.69

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \frac{2 \sqrt{\frac{a+b \cos(c+dx)}{a+b}} (B \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right) + A \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right))}{d \sqrt{a + b \cos(c + dx)}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*(B*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + A*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]))/(d*Sqrt[a + b*Cos[c + d*x]])`

3.323.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx$$

↓ 3481

$$A \int \frac{\sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx + B \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$A \int \frac{1}{\sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx + B \int \frac{1}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx$$

↓ 3142

$$A \int \frac{1}{\sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}}$$

↓ 3042

$$A \int \frac{1}{\sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin\left(c+dx+\frac{\pi}{2}\right)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}}$$

3.323. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3140 \\
& A \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \\
& \downarrow 3286 \\
& \frac{A\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{2B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \\
& \downarrow 3042 \\
& \frac{A\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{2B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} \\
& \downarrow 3284 \\
& \frac{2A\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} + \frac{2B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*A*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]])`

3.323.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3140 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])]/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])]/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3481 `Int[(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.323.4 Maple [A] (verified)

Time = 5.78 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.64

method	result
default	$\frac{2\sqrt{(2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) + a - b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) + a - b}{a - b}} (A\Pi(\cos(\frac{dx}{2} + \frac{c}{2}), 2, \sqrt{-\frac{2b}{a-b}}) - BF(\cos(\frac{dx}{2} + \frac{c}{2}), 2, \sqrt{-\frac{2b}{a-b}}))}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2}))b + (a+b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \sin(\frac{dx}{2} + \frac{c}{2}) \sqrt{-2b(\sin^2(\frac{dx}{2} + \frac{c}{2})) + a + b} d}$
parts	$\frac{2A\sqrt{(2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) + a - b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) + a - b}{a - b}} \Pi(\cos(\frac{dx}{2} + \frac{c}{2}), 2, \sqrt{-\frac{2b}{a-b}}) - 2B\sqrt{(2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) + a - b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2}))b + (a+b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \sin(\frac{dx}{2} + \frac{c}{2}) \sqrt{-2b(\sin^2(\frac{dx}{2} + \frac{c}{2})) + a + b} d}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^(1/2), x, method=_RETURNVERBOSE)`

$$3.323. \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

output $2*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}*(A*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-B*\text{EllipticF}(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)}))/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2+a+b)^{(1/2)}/d$

3.323.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output Timed out

3.323.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/sqrt(a + b*cos(c + d*x)), x)`

3.323.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)/sqrt(b*cos(d*x + c) + a), x)`

3.323. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

3.323.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)/sqrt(b*cos(d*x + c) + a), x)`

3.323.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(1/2)), x)`

3.324
$$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.324.1 Optimal result 3014
 3.324.2 Mathematica [C] (verified) 3015
 3.324.3 Rubi [A] (verified) 3015
 3.324.4 Maple [B] (verified) 3021
 3.324.5 Fracas [F(-1)] 3022
 3.324.6 Sympy [F] 3022
 3.324.7 Maxima [F] 3022
 3.324.8 Giac [F] 3023
 3.324.9 Mupad [F(-1)] 3023

3.324.1 Optimal result

Integrand size = 33, antiderivative size = 216

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= -\frac{A\sqrt{a + b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{ad\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{A\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d\sqrt{a + b \cos(c + dx)}}$$

$$- \frac{(Ab - 2aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{ad\sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{A\sqrt{a + b \cos(c + dx)} \tan(c + dx)}{ad}$$

output

```
-A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/a/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)-(A*b-2*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/a/d/(a+b*cos(d*x+c))^(1/2)+A*(a+b*cos(d*x+c))^(1/2)*tan(d*x+c)/a/d
```

3.324.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 13.55 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.48

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$\frac{2(-3Ab+4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2iA\sqrt{-\frac{b(-1+\cos(c+dx))}{a+b}} \sqrt{\frac{b(1+\cos(c+dx))}{-a+b}} \operatorname{csc}(c+dx) (-2a(a-b)E(i \operatorname{arcsinh}(\sqrt{\frac{a+b \cos(c+dx)}{a+b}})))}{\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/Sqrt[a + b*Cos[c + d*x]],x]`

output `((2*(-3*A*b + 4*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] - ((2*I)*A*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)]))/ (a*b*Sqrt[-(a + b)^(-1)]) + 4*A*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/(4*a*d)`

3.324.3 Rubi [A] (verified)

Time = 1.81 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.05, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3479, 27, 3042, 3539, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

3.324. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^2 \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
& \quad \downarrow \text{3479} \\
& \frac{\int -\frac{(Ab \cos^2(c+dx) + Ab - 2aB) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{a} + \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} \\
& \quad \downarrow \text{27} \\
& \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\int \frac{(Ab \cos^2(c+dx) + Ab - 2aB) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{2a} \\
& \quad \downarrow \text{3042} \\
& \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\int \frac{Ab \sin\left(c+dx + \frac{\pi}{2}\right)^2 + Ab - 2aB}{\sin\left(c+dx + \frac{\pi}{2}\right) \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx}{2a} \\
& \quad \downarrow \text{3539} \\
& \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{A \int \sqrt{a+b \cos(c+dx)} dx - \frac{\int -\frac{(b(Ab-2aB) - aAb \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b}}{2a} \\
& \quad \downarrow \text{25} \\
& \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\frac{\int \frac{(b(Ab-2aB) - aAb \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b}}{2a} + A \int \sqrt{a+b \cos(c+dx)} dx}{2a} \\
& \quad \downarrow \text{3042} \\
& \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\frac{\int \frac{b(Ab-2aB) - aAb \sin\left(c+dx + \frac{\pi}{2}\right)}{\sin\left(c+dx + \frac{\pi}{2}\right) \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx}{b}}{2a} + A \int \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)} dx}{2a} \\
& \quad \downarrow \text{3134} \\
& \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\frac{\int \frac{b(Ab-2aB) - aAb \sin\left(c+dx + \frac{\pi}{2}\right)}{\sin\left(c+dx + \frac{\pi}{2}\right) \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx}{b} + \frac{A \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{2a} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\begin{aligned}
 & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\int \frac{b(Ab-2aB) - aAb \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{A \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}} dx}{\sqrt{a+b \cos(c+dx)} \frac{a+b}}{a+b} \\
 & \frac{2a}{2a} \downarrow \text{3132} \\
 & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{\int \frac{b(Ab-2aB) - aAb \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \frac{2a}{2a} \downarrow \text{3481} \\
 & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{b(Ab-2aB) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - aAb \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \frac{2a}{2a} \downarrow \text{3042} \\
 & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{b(Ab-2aB) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - aAb \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \frac{2a}{2a} \downarrow \text{3142} \\
 & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{b(Ab-2aB) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b}}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \frac{2a}{2a} \downarrow \text{3042} \\
 & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{b(Ab-2aB) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{b}}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\
 & \frac{2a}{2a}
 \end{aligned}$$

3.324. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned} & \downarrow 3140 \\ & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \\ & \frac{b(Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \end{aligned}$$

$$\begin{aligned} & 2a \\ & \downarrow 3286 \\ & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \\ & \frac{b(Ab-2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - \frac{2aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \end{aligned}$$

$$\begin{aligned} & 2a \\ & \downarrow 3042 \\ & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \\ & \frac{b(Ab-2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx - \frac{2aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \end{aligned}$$

$$\begin{aligned} & \downarrow 3284 \\ & \frac{A \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \\ & \frac{2b(Ab-2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - \frac{2aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}}}{b} + \frac{2A \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/Sqrt[a + b*Cos[c + d*x]],x]`

output `-1/2*((2*A*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((-2*a*A*b*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*b*(A*b - 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b/a + (A*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/(a*d)`

3.324. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

3.324.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3539 `Int[((A_.) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.324.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 638 vs. $2(289) = 578$.

Time = 8.26 (sec) , antiderivative size = 639, normalized size of antiderivative = 2.96

method	result
default	$\frac{\sqrt{-(-2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) - a + b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{2A \left(-\frac{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2}))b + (a+b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{a(2(\cos^2(\frac{dx}{2} + \frac{c}{2})) - 1)} + \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}}{2\sqrt{-\dots}} \right)}$
parts	$A\sqrt{-(-2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) - a + b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \left(-\frac{2\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2}))b + (a+b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{a(2(\cos^2(\frac{dx}{2} + \frac{c}{2})) - 1)} + \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}}{\sqrt{-2(s\dots}} \right)$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*(-cos(
1/2*d*x+1/2*c)/a*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1
/2)/(2*cos(1/2*d*x+1/2*c)^2-1)+1/2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(
1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/
2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-1/2
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)
/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(co
s(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+1/2/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((
2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b
)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))+1/2/a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b
)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)
)*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))-2*B*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x
+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c
),2,(-2*b/(a-b))^(1/2))/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b
)^(1/2)/d
```

3.324.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `Timed out`

3.324.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/sqrt(a + b*cos(c + d*x)), x)`

3.324.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^2}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^2/sqrt(b*cos(d*x + c) + a), x)`

3.324.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^2}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^2/sqrt(b*cos(d*x + c) + a), x)`

3.324.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^2 \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^(1/2)), x)`

3.325
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.325.1 Optimal result 3024
 3.325.2 Mathematica [C] (verified) 3025
 3.325.3 Rubi [A] (verified) 3026
 3.325.4 Maple [B] (verified) 3032
 3.325.5 Fricas [F(-1)] 3033
 3.325.6 Sympy [F] 3034
 3.325.7 Maxima [F] 3034
 3.325.8 Giac [F] 3034
 3.325.9 Mupad [F(-1)] 3035

3.325.1 Optimal result

Integrand size = 33, antiderivative size = 299

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx \\ &= \frac{(3Ab - 4aB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{4a^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \\ & \quad - \frac{(Ab - 4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4ad \sqrt{a + b \cos(c + dx)}} \\ & \quad + \frac{(4a^2 A + 3Ab^2 - 4abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4a^2 d \sqrt{a + b \cos(c + dx)}} \\ & \quad - \frac{(3Ab - 4aB) \sqrt{a + b \cos(c + dx)} \tan(c + dx)}{4a^2 d} \\ & \quad + \frac{A \sqrt{a + b \cos(c + dx)} \sec(c + dx) \tan(c + dx)}{2ad} \end{aligned}$$

```
output 1/4*(3*A*b-4*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/a^2/d
/((a+b*cos(d*x+c))/(a+b))^(1/2)-1/4*(A*b-4*B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2)
)*(a+b*cos(d*x+c))/(a+b))^(1/2)/a/d/(a+b*cos(d*x+c))^(1/2)+1/4*(4*A*a^2+3
*A*b^2-4*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi
(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1
/2)/a^2/d/(a+b*cos(d*x+c))^(1/2)-1/4*(3*A*b-4*B*a)*(a+b*cos(d*x+c))^(1/2)*
tan(d*x+c)/a^2/d+1/2*A*sec(d*x+c)*(a+b*cos(d*x+c))^(1/2)*tan(d*x+c)/a/d
```

3.325.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.19 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.40

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{8aAb \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2(8a^2A+9Ab^2-12abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2i(3Ab-4aB) \sqrt{-\frac{b}{a+b}} \operatorname{EllipticE}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2i(3Ab-4aB) \sqrt{-\frac{b}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2i(3Ab-4aB) \sqrt{-\frac{b}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2i(3Ab-4aB) \sqrt{-\frac{b}{a+b}} \operatorname{EllipticE}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}}$$

```
input Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/Sqrt[a + b*Cos[c + d*x]],x
]
```

```
output ((8*a*A*b*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/
(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(8*a^2*A + 9*A*b^2 - 12*a*b*B)*Sqr
t[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])
/Sqrt[a + b*Cos[c + d*x]] + ((2*I)*(3*A*b - 4*a*B)*Sqrt[-((b*(-1 + Cos[c +
d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a
*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]]
, (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[
a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh
[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b))))/(a*b*S
qrt[-(a + b)^(-1)] + 4*Sqrt[a + b*Cos[c + d*x]]*(2*a*A + (-3*A*b + 4*a*B)
*Cos[c + d*x])*Sec[c + d*x]*Tan[c + d*x])/(16*a^2*d)
```


3.325.3 Rubi [A] (verified)

Time = 2.50 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.02, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^3 \sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3479} \\
 & \int -\frac{(-Ab\cos^2(c+dx)-2aA\cos(c+dx)+3Ab-4aB)\sec^2(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx \\
 & \quad + \frac{2a}{2ad} \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2ad} \\
 & \quad \downarrow \text{27} \\
 & \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2ad} - \frac{\int \frac{(-Ab\cos^2(c+dx)-2aA\cos(c+dx)+3Ab-4aB)\sec^2(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{4a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2ad} - \frac{\int \frac{-Ab\sin\left(c+dx+\frac{\pi}{2}\right)^2-2aA\sin\left(c+dx+\frac{\pi}{2}\right)+3Ab-4aB}{\sin\left(c+dx+\frac{\pi}{2}\right)^2 \sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{4a} \\
 & \quad \downarrow \text{3534} \\
 & \frac{A\tan(c+dx)\sec(c+dx)\sqrt{a+b\cos(c+dx)}}{2ad} - \frac{\int -\frac{(4Aa^2-4bBa+2Ab\cos(c+dx)a+3Ab^2+b(3Ab-4aB)\cos^2(c+dx))\sec^2(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx}{a} + \frac{(3Ab-4aB)\tan(c+dx)\sqrt{a+b\cos(c+dx)}}{ad} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.325. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$

$$\begin{aligned}
 & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2ad}{\int \frac{(4Aa^2-4bBa+2Ab \cos(c+dx)a+3Ab^2+b(3Ab-4aB) \cos^2(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2ad}{\int \frac{4Aa^2-4bBa+2Ab \sin(c+dx+\frac{\pi}{2})a+3Ab^2+b(3Ab-4aB) \sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx} \\
 & \qquad \qquad \qquad \downarrow 3538 \\
 & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2ad}{(3Ab-4aB) \int \sqrt{a+b \cos(c+dx)} dx - \frac{\int -\frac{(b(4Aa^2-4bBa+3Ab^2)-ab(Ab-4aB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b}} \\
 & \qquad \qquad \qquad \downarrow 25 \\
 & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2ad}{\int \frac{(b(4Aa^2-4bBa+3Ab^2)-ab(Ab-4aB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + (3Ab-4aB) \int \sqrt{a+b \cos(c+dx)} dx} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2ad}{\int \frac{b(4Aa^2-4bBa+3Ab^2)-ab(Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + (3Ab-4aB) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx} \\
 & \qquad \qquad \qquad \downarrow 3134 \\
 & \frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2ad}{\int \frac{b(4Aa^2-4bBa+3Ab^2)-ab(Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{(3Ab-4aB) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}} \\
 & \qquad \qquad \qquad \downarrow 3042
 \end{aligned}$$

3.325. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{\int \frac{b(4Aa^2-4abB+3Ab^2) - ab(Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{(3Ab-4aB) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}}{\sqrt{a+b \cos(c+dx)}}}{2a}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{4a}{2a}$$

3132

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{\int \frac{b(4Aa^2-4abB+3Ab^2) - ab(Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{2(3Ab-4aB) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{4a}{2a}$$

3481

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - ab(Ab-4aB) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{b} + \frac{2(3Ab-4aB) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{4a}{2a}$$

3042

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - ab(Ab-4aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + 2(3Ab-4aB) \sqrt{a+b \cos(c+dx)}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{4a}{2a}$$

3142

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(Ab-4aB) \sqrt{a+b \cos(c+dx)} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}}}{\sqrt{a+b \cos(c+dx)}}}{b} + 2(3Ab-4aB) \sqrt{a+b \cos(c+dx)}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{4a}{2a}$$

3042

3.325. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(Ab-4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2a}{4a}$$

↓ 3140

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2ab(Ab-4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \sqrt{\frac{a+b \cos(c+dx)}{a+b}})}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2a}{4a}$$

↓ 3286

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - \frac{2ab(Ab-4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \sqrt{\frac{a+b \cos(c+dx)}{a+b}})}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2a}{4a}$$

↓ 3042

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{b(4a^2A-4abB+3Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx - \frac{2ab(Ab-4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2a}{4a}$$

↓ 3284

$$\frac{A \tan(c+dx) \sec(c+dx) \sqrt{a+b \cos(c+dx)}}{2ad} - \frac{2b(4a^2A-4abB+3Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticPi}(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}) - \frac{2ab(Ab-4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \sqrt{\frac{a+b \cos(c+dx)}{a+b}})}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(3Ab-4aB) \tan(c+dx) \sqrt{a+b \cos(c+dx)}}{ad} - \frac{2a}{4a}$$

3.325. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/Sqrt[a + b*Cos[c + d*x]],x]`

output `(A*Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]*Tan[c + d*x])/(2*a*d) - (-1/2*((2*(3*A*b - 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((-2*a*b*(A*b - 4*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*b*(4*a^2*A + 3*A*b^2 - 4*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/a + ((3*A*b - 4*a*B)*Sqrt[a + b*Cos[c + d*x]]*Tan[c + d*x])/(a*d))/(4*a)`

3.325.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^n), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.325.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1181 vs. 2(360) = 720.

Time = 10.81 (sec) , antiderivative size = 1182, normalized size of antiderivative = 3.95

method	result	size
default	Expression too large to display	1182
parts	Expression too large to display	1244

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*(-1/2*
cos(1/2*d*x+1/2*c)/a*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2
)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^2+3/4*b/a^2*cos(1/2*d*x+1/2*c)*(-2*sin(
1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)
^2-1)-1/8*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)
/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)
*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))+3/8/a*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/
2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*EllipticE(cos(1/2*d*x+1/2*c),
(-2*b/(a-b))^(1/2))-3/8*b^2/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2
*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d
*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-1/2*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(
1/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))-3/8/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a
+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),2,(-2*b/(a-b
))^(1/2))*b^2+2*B*(-cos(1/2*d*x+1/2*c)/a*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)
*sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)+1/2*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2...

```

3.325.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `Timed out`

3.325.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/sqrt(a + b*cos(c + d*x)), x)`

3.325.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^3}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^3/sqrt(b*cos(d*x + c) + a), x)`

3.325.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^3}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^3/sqrt(b*cos(d*x + c) + a), x)`

3.325.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^3 \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^(1/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^(1/2)), x)`

3.326
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.326.1 Optimal result	3036
3.326.2 Mathematica [A] (verified)	3037
3.326.3 Rubi [A] (verified)	3038
3.326.4 Maple [B] (verified)	3044
3.326.5 Fricas [C] (verification not implemented)	3045
3.326.6 Sympy [F(-1)]	3045
3.326.7 Maxima [F]	3046
3.326.8 Giac [F]	3046
3.326.9 Mupad [F(-1)]	3046

3.326.1 Optimal result

Integrand size = 33, antiderivative size = 387

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx =$$

$$\frac{2(40a^3Ab - 25aAb^3 - 48a^4B + 24a^2b^2B + 9b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{15b^4(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(40a^2Ab + 5Ab^3 - 48a^3B - 12ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{15b^4 d \sqrt{a+b \cos(c+dx)}}$$

$$+ \frac{2a(Ab - aB) \cos^2(c+dx) \sin(c+dx)}{b(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}}$$

$$+ \frac{2(20a^2Ab - 5Ab^3 - 24a^3B + 9ab^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15b^3(a^2 - b^2) d}$$

$$- \frac{2(5aAb - 6a^2B + b^2B) \cos(c+dx) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{5b^2(a^2 - b^2) d}$$

output $2*a*(A*b-B*a)*\cos(d*x+c)^2*\sin(d*x+c)/b/(a^2-b^2)/d/(a+b*\cos(d*x+c))^{(1/2)}+2/15*(20*A*a^2*b-5*A*b^3-24*B*a^3+9*B*a*b^2)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/b^3/(a^2-b^2)/d-2/5*(5*A*a*b-6*B*a^2+B*b^2)*\cos(d*x+c)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/b^2/(a^2-b^2)/d-2/15*(40*A*a^3*b-25*A*a*b^3-48*B*a^4+24*B*a^2*b^2+9*B*b^4)*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*(a+b*\cos(d*x+c))^{(1/2)}/b^4/(a^2-b^2)/d/((a+b*\cos(d*x+c))/(a+b))^{(1/2)}+2/15*(40*A*a^2*b+5*A*b^3-48*B*a^3-12*B*a*b^2)*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)}*(b/(a+b))^{(1/2)})*((a+b*\cos(d*x+c))/(a+b))^{(1/2)}/b^4/d/(a+b*\cos(d*x+c))^{(1/2)}$

3.326.2 Mathematica [A] (verified)

Time = 1.80 (sec) , antiderivative size = 304, normalized size of antiderivative = 0.79

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \frac{2b^2(-10a^2Ab-5Ab^3+12a^3B+3ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + 2(-10a^2Ab-5Ab^3+12a^3B+3ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{(a-b)(a+b)}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]`

output $((2*b^2*(-10*a^2*A*b - 5*A*b^3 + 12*a^3*B + 3*a*b^2*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)])/((a - b)*(a + b)) + (2*(-40*a^3*A*b + 25*a*A*b^3 + 48*a^4*B - 24*a^2*b^2*B - 9*b^4*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/(a + b)]*((a + b)*\text{EllipticE}[(c + d*x)/2, (2*b)/(a + b)] - a*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]))/((a - b)*(a + b)) + (30*a^3*b*(-(A*b) + a*B)*\text{Sin}[c + d*x])/(-a^2 + b^2) + 2*b*(5*A*b - 9*a*B)*(a + b*\text{Cos}[c + d*x])*\text{Sin}[c + d*x] + 3*b^2*B*(a + b*\text{Cos}[c + d*x])*\text{Sin}[2*(c + d*x)])/(15*b^4*d*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])$

3.326.3 Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 391, normalized size of antiderivative = 1.01, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3468, 27, 3042, 3528, 27, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^3 (A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx$$

$$\downarrow \text{3468}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} -$$

$$\frac{2\int -\frac{\cos(c+dx)(-((-6Ba^2+5Aba+b^2B)\cos^2(c+dx)-b(Ab-aB)\cos(c+dx)+4a(Ab-aB)))}{2\sqrt{a+b\cos(c+dx)}} dx}{b(a^2-b^2)}$$

$$\downarrow \text{27}$$

$$\frac{\int \frac{\cos(c+dx)(-((-6Ba^2+5Aba+b^2B)\cos^2(c+dx)-b(Ab-aB)\cos(c+dx)+4a(Ab-aB)))}{\sqrt{a+b\cos(c+dx)}} dx}{b(a^2-b^2)} +$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\downarrow \text{3042}$$

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})((6Ba^2-5Aba-b^2B)\sin(c+dx+\frac{\pi}{2})^2-b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+4a(Ab-aB))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b(a^2-b^2)} +$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\downarrow \text{3528}$$

$$2 \int \frac{-((-24Ba^3 + 20Aba^2 + 9b^2Ba - 5Ab^3) \cos^2(c+dx)) - b(-2Ba^2 + 5Aba - 3b^2B) \cos(c+dx) + 2a(-6Ba^2 + 5Aba + b^2B)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-6a^2B + 5aAb + b^2B) \sin(c+dx)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \quad b(a^2 - b^2)$$

↓ 27

$$\int \frac{-((-24Ba^3 + 20Aba^2 + 9b^2Ba - 5Ab^3) \cos^2(c+dx)) - b(-2Ba^2 + 5Aba - 3b^2B) \cos(c+dx) + 2a(-6Ba^2 + 5Aba + b^2B)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-6a^2B + 5aAb + b^2B) \sin(c+dx)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \quad b(a^2 - b^2)$$

↓ 3042

$$\int \frac{(24Ba^3 - 20Aba^2 - 9b^2Ba + 5Ab^3) \sin(c+dx + \frac{\pi}{2})^2 - b(-2Ba^2 + 5Aba - 3b^2B) \sin(c+dx + \frac{\pi}{2}) + 2a(-6Ba^2 + 5Aba + b^2B)}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{2(-6a^2B + 5aAb + b^2B) \sin(c+dx)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \quad b(a^2 - b^2)$$

↓ 3502

$$2 \int \frac{b(-12Ba^3 + 10Aba^2 - 3b^2Ba + 5Ab^3) + (-48Ba^4 + 40Aba^3 + 24b^2Ba^2 - 25Ab^3a + 9b^4B) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-24a^3B + 20a^2Ab + 9ab^2B - 5Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \quad b(a^2 - b^2)$$

↓ 27

$$\int \frac{b(-12Ba^3 + 10Aba^2 - 3b^2Ba + 5Ab^3) + (-48Ba^4 + 40Aba^3 + 24b^2Ba^2 - 25Ab^3a + 9b^4B) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - \frac{2(-24a^3B + 20a^2Ab + 9ab^2B - 5Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3bd}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \quad b(a^2 - b^2)$$

↓ 3042

3.326. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{\int \frac{b(-12Ba^3+10Aba^2-3b^2Ba+5Ab^3)+(-48Ba^4+40Aba^3+24b^2Ba^2-25Ab^3a+9b^4B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\frac{(-24a^3B+20a^2Ab+9ab^2B-5Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3bd}} = \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \frac{b(a^2-b^2)}{b(a^2-b^2)}$$

3231

$$\frac{\frac{(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\int\sqrt{a+b\cos(c+dx)}dx}{b} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\int\frac{1}{\sqrt{a+b\cos(c+dx)}}dx}{b}}{\frac{2(-24a^3B+20a^2Ab+9ab^2B-5Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}}} = \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \frac{b(a^2-b^2)}{b(a^2-b^2)}$$

3042

$$\frac{\frac{(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\int\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}dx}{b} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\int\frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{b}}{\frac{2(-24a^3B+20a^2Ab+9ab^2B-5Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}}} = \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \frac{b(a^2-b^2)}{b(a^2-b^2)}$$

3134

$$\frac{\frac{(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}dx}{b\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\int\frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{b}}{\frac{2(-24a^3B+20a^2Ab+9ab^2B-5Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}}} = \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \frac{b(a^2-b^2)}{b(a^2-b^2)}$$

3042

$$\frac{\frac{(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}dx}{b\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\int\frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{b}}{\frac{2(-24a^3B+20a^2Ab+9ab^2B-5Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}}} = \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \frac{b(a^2-b^2)}{b(a^2-b^2)}$$

3132

3.326. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{2(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\int\frac{1}{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}}dx}{b} \\
 & \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{3142} \\
 & \frac{2(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}}}}{b\sqrt{a+b\cos(c+dx)}} \\
 & \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-48a^3B+40a^2Ab-12ab^2B+5Ab^3)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}}}}{b\sqrt{a+b\cos(c+dx)}} \\
 & \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{3140} \\
 & \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \\
 & \frac{2(-6a^2B+5aAb+b^2B)\sin(c+dx)\cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd} - \frac{2(-48a^4B+40a^3Ab+24a^2b^2B-25aAb^3+9b^4B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}
 \end{aligned}$$

```
input Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]
```

3.326. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$


```
output (2*a*(A*b - a*B)*Cos[c + d*x]^2*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[a + b*
Cos[c + d*x]]) + ((-2*(5*a*A*b - 6*a^2*B + b^2*B)*Cos[c + d*x]*Sqrt[a + b*
Cos[c + d*x]]*Sin[c + d*x])/(5*b*d) - (((2*(40*a^3*A*b - 25*a*A*b^3 - 48*a
^4*B + 24*a^2*b^2*B + 9*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x
)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 -
b^2)*(40*a^2*A*b + 5*A*b^3 - 48*a^3*B - 12*a*b^2*B)*Sqrt[(a + b*Cos[c + d*
x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c
+ d*x]])))/(3*b) - (2*(20*a^2*A*b - 5*A*b^3 - 24*a^3*B + 9*a*b^2*B)*Sqrt[a
+ b*Cos[c + d*x]]*Sin[c + d*x])/(3*b*d))/(5*b))/(b*(a^2 - b^2))
```

3.326.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*S
qrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a
+ b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b)
+ (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && !GtQ[a + b, 0]
```

rule 3231 `Int[((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(GtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

3.326.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1311 vs. $2(423) = 846$.

Time = 14.77 (sec) , antiderivative size = 1312, normalized size of antiderivative = 3.39

method	result	size
default	Expression too large to display	1312
parts	Expression too large to display	2030

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(16*B/b*(-1
/10/b*cos(1/2*d*x+1/2*c)^3*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/
2*c)^2)^(1/2)-1/60/b^2*(-4*a+12*b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*
c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+1/60/b^2*(-4*a+12*b)*(a-b)*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*si
n(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d
*x+1/2*c),(-2*b/(a-b))^(1/2))-1/60*(4*a^2-15*a*b+27*b^2)/b^3*(a-b)*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin
(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d
*x+1/2*c),(-2*b/(a-b))^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1
/2))))+8/b^2*(A*b-B*a-3*B*b)*(-1/6/b*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/
2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+1/6*(a-b)/b*(sin(1/2*d*x+1/2*c
)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2
*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2
*b/(a-b))^(1/2))-1/12/b^2*(-2*a+6*b)*(a-b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((
2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b
)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))))+2*a^3*(A*b-B*a)/b
^4/sin(1/2*d*x+1/2*c)^2/(2*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*sin(1
/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*cos(1/2*d*x+1/2*...
```

3.326.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.21 (sec) , antiderivative size = 910, normalized size of antiderivative = 2.35

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm
m="fricas")
```

```
output -1/45*(6*(24*B*a^4*b^2 - 20*A*a^3*b^3 - 9*B*a^2*b^4 + 5*A*a*b^5 - 3*(B*a^2
*b^4 - B*b^6)*cos(d*x + c)^2 + (6*B*a^3*b^3 - 5*A*a^2*b^4 - 6*B*a*b^5 + 5*
A*b^6)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c) + (sqrt(2)*(-96
*I*B*a^5*b + 80*I*A*a^4*b^2 + 84*I*B*a^3*b^3 - 80*I*A*a^2*b^4 + 27*I*B*a*b
^5 - 15*I*A*b^6)*cos(d*x + c) + sqrt(2)*(-96*I*B*a^6 + 80*I*A*a^5*b + 84*I
*B*a^4*b^2 - 80*I*A*a^3*b^3 + 27*I*B*a^2*b^4 - 15*I*A*a*b^5))*sqrt(b)*weie
rstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*
(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (sqrt(2)*(96*I*B*a^5*b
- 80*I*A*a^4*b^2 - 84*I*B*a^3*b^3 + 80*I*A*a^2*b^4 - 27*I*B*a*b^5 + 15*I*A
*b^6)*cos(d*x + c) + sqrt(2)*(96*I*B*a^6 - 80*I*A*a^5*b - 84*I*B*a^4*b^2 +
80*I*A*a^3*b^3 - 27*I*B*a^2*b^4 + 15*I*A*a*b^5))*sqrt(b)*weierstrassPInve
rse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x
+ c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(sqrt(2)*(48*I*B*a^4*b^2 - 40*I*A
*a^3*b^3 - 24*I*B*a^2*b^4 + 25*I*A*a*b^5 - 9*I*B*b^6)*cos(d*x + c) + sqrt(
2)*(48*I*B*a^5*b - 40*I*A*a^4*b^2 - 24*I*B*a^3*b^3 + 25*I*A*a^2*b^4 - 9*I*
B*a*b^5))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 -
9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 -
9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*(s
qrt(2)*(-48*I*B*a^4*b^2 + 40*I*A*a^3*b^3 + 24*I*B*a^2*b^4 - 25*I*A*a*b^5 +
9*I*B*b^6)*cos(d*x + c) + sqrt(2)*(-48*I*B*a^5*b + 40*I*A*a^4*b^2 + 24...
```

3.326.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)
```

3.326. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

output Timed out

3.326.7 Maxima [F]

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^3}{(b\cos(dx+c)+a)^{3/2}} dx$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^3/(b*cos(d*x + c) + a)^(3/2), x)`

3.326.8 Giac [F]

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^3}{(b\cos(dx+c)+a)^{3/2}} dx$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^3/(b*cos(d*x + c) + a)^(3/2), x)`

3.326.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^3(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2), x)`

3.326. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

3.327
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.327.1 Optimal result 3047
 3.327.2 Mathematica [A] (verified) 3048
 3.327.3 Rubi [A] (verified) 3048
 3.327.4 Maple [B] (verified) 3053
 3.327.5 Fracas [C] (verification not implemented) 3054
 3.327.6 Sympy [F(-1)] 3055
 3.327.7 Maxima [F] 3055
 3.327.8 Giac [F] 3055
 3.327.9 Mupad [F(-1)] 3056

3.327.1 Optimal result

Integrand size = 33, antiderivative size = 262

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx = \frac{2(6a^2Ab - 3Ab^3 - 8a^3B + 5ab^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{3b^3(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(6aAb - 8a^2B - b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3b^3 d \sqrt{a+b \cos(c+dx)}} - \frac{2a^2(Ab - aB) \sin(c+dx)}{b^2(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}} + \frac{2B \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3b^2 d}$$

output

```
-2*a^2*(A*b-B*a)*sin(d*x+c)/b^2/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+2/3*B*
sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/b^2/d+2/3*(6*A*a^2*b-3*A*b^3-8*B*a^3+5*B*
a*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d
*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^3/(a^2-b^2)/d/
((a+b*cos(d*x+c))/(a+b))^(1/2)-2/3*(6*A*a*b-8*B*a^2-B*b^2)*(cos(1/2*d*x+1/
2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(
a+b))^(1/2))*(a+b*cos(d*x+c))/(a+b))^(1/2)/b^3/d/(a+b*cos(d*x+c))^(1/2)
```

3.327.2 Mathematica [A] (verified)

Time = 1.47 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.72

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2 \left(\frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \left((6a^2Ab - 3Ab^3 - 8a^3B + 5ab^2B) E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) + (a-b)(-6aAb + 8a^2B + b^2B) \operatorname{EllipticF}\left[\frac{c+dx}{2}, \frac{2b}{a+b}\right] \right)}{a-b} \right)}{3b^3}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2), x]`

output `(2*((Sqrt[(a + b*Cos[c + d*x])/(a + b)]*((6*a^2*A*b - 3*A*b^3 - 8*a^3*B + 5*a*b^2*B)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] + (a - b)*(-6*a*A*b + 8*a^2*B + b^2*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)])))/(a - b) + b*((a*(3*a*A*b - 4*a^2*B + b^2*B))/(-a^2 + b^2) + b*B*Cos[c + d*x])*Sin[c + d*x]))/(3*b^3*d*Sqrt[a + b*Cos[c + d*x]])`

3.327.3 Rubi [A] (verified)

Time = 1.44 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3467, 27, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx \\ & \quad \downarrow \text{3467} \\ & \frac{2 \int \frac{b(a^2 - b^2)B \cos^2(c + dx) + (2a^2 - b^2)(Ab - aB) \cos(c + dx) + ab(Ab - aB)}{2\sqrt{a + b \cos(c + dx)}} dx}{b^2(a^2 - b^2)} - \frac{2a^2(Ab - aB) \sin(c + dx)}{b^2d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.327. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{\int \frac{b(a^2-b^2)B \cos^2(c+dx) + (2a^2-b^2)(Ab-aB) \cos(c+dx) + ab(Ab-aB)}{\sqrt{a+b \cos(c+dx)}} dx}{b^2(a^2-b^2)} - \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3042

$$\frac{\int \frac{b(a^2-b^2)B \sin(c+dx+\frac{\pi}{2})^2 + (2a^2-b^2)(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + ab(Ab-aB)}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b^2(a^2-b^2)} - \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3502

$$\frac{2 \int \frac{(-2Ba^2+3Aba-b^2B)b^2 + (-8Ba^3+6Aba^2+5b^2Ba-3Ab^3) \cos(c+dx)b}{2\sqrt{a+b \cos(c+dx)}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} - \frac{b^2(a^2-b^2)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} - \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 27

$$\frac{\int \frac{(-2Ba^2+3Aba-b^2B)b^2 + (-8Ba^3+6Aba^2+5b^2Ba-3Ab^3) \cos(c+dx)b}{\sqrt{a+b \cos(c+dx)}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} - \frac{b^2(a^2-b^2)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} - \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3042

$$\frac{\int \frac{(-2Ba^2+3Aba-b^2B)b^2 + (-8Ba^3+6Aba^2+5b^2Ba-3Ab^3) \sin(c+dx+\frac{\pi}{2})b}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} - \frac{b^2(a^2-b^2)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} - \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3231

$$\frac{(-8a^3B+6a^2Ab+5ab^2B-3Ab^3) \int \sqrt{a+b \cos(c+dx)} dx - (a^2-b^2)(-8a^2B+6aAb-b^2B) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d} - \frac{b^2(a^2-b^2)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} - \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2 d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3042

3.327. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{(-8a^3B+6a^2Ab+5ab^2B-3Ab^3) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx - (a^2-b^2)(-8a^2B+6aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{3d}$$

$$\frac{b^2(a^2-b^2)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3134

$$\frac{(-8a^3B+6a^2Ab+5ab^2B-3Ab^3) \frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx - (a^2-b^2)(-8a^2B+6aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{3d}$$

$$\frac{b^2(a^2-b^2)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3042

$$\frac{(-8a^3B+6a^2Ab+5ab^2B-3Ab^3) \frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx - (a^2-b^2)(-8a^2B+6aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{3d}$$

$$\frac{b^2(a^2-b^2)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3132

$$\frac{2(-8a^3B+6a^2Ab+5ab^2B-3Ab^3) \frac{\sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - (a^2-b^2)(-8a^2B+6aAb-b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{3d}$$

$$\frac{b^2(a^2-b^2)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3142

$$\frac{2(-8a^3B+6a^2Ab+5ab^2B-3Ab^3) \frac{\sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - (a^2-b^2)(-8a^2B+6aAb-b^2B) \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}}}{3b} + \frac{2B(a^2-b^2) \sin(c+dx) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{3d}$$

$$\frac{b^2(a^2-b^2)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \frac{2a^2(Ab-aB) \sin(c+dx)}{b^2d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

↓ 3042

3.327. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{2(-8a^3B+6a^2Ab+5ab^2B-3Ab^3)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{(a^2-b^2)(-8a^2B+6aAb-b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin\left(c+dx+\frac{\pi}{2}\right)}{a+b}}}dx}{\sqrt{a+b\cos(c+dx)}}}{3b}+\frac{2B(a^2-b^2)\sin(c+dx)}{b^2(a^2-b^2)}+\frac{2a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

↓ 3140

$$\frac{2B(a^2-b^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}+\frac{2(-8a^3B+6a^2Ab+5ab^2B-3Ab^3)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)-2(a^2-b^2)(-8a^2B+6aAb-b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{2(a^2-b^2)\sin(c+dx)}{3b}}{b^2(a^2-b^2)}+\frac{2a^2(Ab-aB)\sin(c+dx)}{b^2d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

```
input Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]
```

```
output (-2*a^2*(A*b - a*B)*Sin[c + d*x])/(b^2*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + (((2*(6*a^2*A*b - 3*A*b^3 - 8*a^3*B + 5*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(6*a*A*b - 8*a^2*B - b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])))/(3*b) + (2*(a^2 - b^2)*B*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d))/(b^2*(a^2 - b^2))
```

3.327.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

3.327. $\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3467 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^2*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.327.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1274 vs. $2(302) = 604$.

Time = 14.82 (sec) , antiderivative size = 1275, normalized size of antiderivative = 4.87

method	result	size
parts	Expression too large to display	1275
default	Expression too large to display	1336

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -2*A*(2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a^2*b-2*(sin(1/2*d*x+1/2*c)
)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(c
os(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^3+2*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1/2
*d*x+1/2*c),(-2*b/(a-b))^(1/2))+2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)
*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*
b/(a-b))^(1/2))*a^3-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x
+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/
2))*a^2*b-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a
+b)/(a-b))^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a*b^2+(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b^3/b^2/(a-b)/(a+b)
/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d-2/3*B*(4*cos(1
/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*a^2*b^2-4*cos(1/2*d*x+1/2*c)*sin(1/2*d*
x+1/2*c)^4*b^4-8*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a^3*b-2*cos(1/2*d
*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a^2*b^2+2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/
2*c)^2*a*b^3+2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b^4+8*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*Ellipt
icF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*a^4-7*(sin(1/2*d*x+1/2*c)^2)^(1
/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*EllipticF(cos(1...
```

3.327.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.17 (sec) , antiderivative size = 789, normalized size of antiderivative = 3.01

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \frac{6(4Ba^3b^2 - 3Aa^2b^3 - Bab^4 + (Ba^2b^3 - Bb^5)\cos(dx+c))\sqrt{b\cos(dx+c)}}{(a+b\cos(c+dx))^{3/2}}$$

```
input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm
m="fricas")
```

```
output 1/9*(6*(4*B*a^3*b^2 - 3*A*a^2*b^3 - B*a*b^4 + (B*a^2*b^3 - B*b^5)*cos(d*x
+ c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c) - (sqrt(2)*(16*I*B*a^4*b - 12*
I*A*a^3*b^2 - 16*I*B*a^2*b^3 + 15*I*A*a*b^4 - 3*I*B*b^5)*cos(d*x + c) + sq
rt(2)*(16*I*B*a^5 - 12*I*A*a^4*b - 16*I*B*a^3*b^2 + 15*I*A*a^2*b^3 - 3*I*B
*a*b^4))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3
- 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) - (s
qrt(2)*(-16*I*B*a^4*b + 12*I*A*a^3*b^2 + 16*I*B*a^2*b^3 - 15*I*A*a*b^4 + 3
*I*B*b^5)*cos(d*x + c) + sqrt(2)*(-16*I*B*a^5 + 12*I*A*a^4*b + 16*I*B*a^3*
b^2 - 15*I*A*a^2*b^3 + 3*I*B*a*b^4))*sqrt(b)*weierstrassPInverse(4/3*(4*a^
2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b
*sin(d*x + c) + 2*a)/b) + 3*(sqrt(2)*(-8*I*B*a^3*b^2 + 6*I*A*a^2*b^3 + 5*I
*B*a*b^4 - 3*I*A*b^5)*cos(d*x + c) + sqrt(2)*(-8*I*B*a^4*b + 6*I*A*a^3*b^2
+ 5*I*B*a^2*b^3 - 3*I*A*a*b^4))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^
2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^
2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x
+ c) + 2*a)/b)) + 3*(sqrt(2)*(8*I*B*a^3*b^2 - 6*I*A*a^2*b^3 - 5*I*B*a*b^4
+ 3*I*A*b^5)*cos(d*x + c) + sqrt(2)*(8*I*B*a^4*b - 6*I*A*a^3*b^2 - 5*I*B*
a^2*b^3 + 3*I*A*a*b^4))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -
8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -
8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) ...
```

3.327.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.327.7 Maxima [F]

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^2}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^2/(b*cos(d*x + c) + a)^(3/2), x)`

3.327.8 Giac [F]

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^2}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^2/(b*cos(d*x + c) + a)^(3/2), x)`

3.327.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^2(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`output `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2), x)`

3.328
$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.328.1 Optimal result 3057
 3.328.2 Mathematica [A] (verified) 3058
 3.328.3 Rubi [A] (verified) 3058
 3.328.4 Maple [B] (verified) 3062
 3.328.5 Fracas [C] (verification not implemented) 3063
 3.328.6 Sympy [F(-1)] 3064
 3.328.7 Maxima [F] 3064
 3.328.8 Giac [F] 3065
 3.328.9 Mupad [F(-1)] 3065

3.328.1 Optimal result

Integrand size = 31, antiderivative size = 204

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx =$$

$$\frac{2(aAb - 2a^2B + b^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) - b^2(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b^2 d \sqrt{a+b \cos(c+dx)}} + \frac{2(Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{b^2 d \sqrt{a+b \cos(c+dx)}} + \frac{2a(Ab - aB) \sin(c+dx)}{b(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}}$$

```
output 2*a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)-2*(A*a*b-2*B
*a^2+B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(
1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^2/(a^2-b^
2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2*(A*b-2*B*a)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/
2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b^2/d/(a+b*cos(d*x+c))^(1/2)
```


3.328.2 Mathematica [A] (verified)

Time = 0.90 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.83

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx =$$

$$\frac{2\left(-\left((a+b)(-aAb+2a^2B-b^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)\right)+(a^2-b^2)(-Ab+2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\right)}{(a-b)b^2(a+b)d\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]`

output `(-2*(-((a + b)*(-(a*A*b) + 2*a^2*B - b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]) + (a^2 - b^2)*(-(A*b) + 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + a*b*(-(A*b) + a*B)*Sin[c + d*x])/((a - b)*b^2*(a + b)*d*Sqrt[a + b*Cos[c + d*x]])`

3.328.3 Rubi [A] (verified)Time = 1.10 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx$$

$$\downarrow \text{3447}$$

$$\int \frac{A\cos(c+dx)+B\cos^2(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

3.328. $\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \int \frac{A \sin \left(c+dx+\frac{\pi}{2}\right)+B \sin \left(c+dx+\frac{\pi}{2}\right)^2}{\left(a+b \sin \left(c+dx+\frac{\pi}{2}\right)\right)^{3 / 2}} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{2 a(A b-a B) \sin (c+dx)}{b d\left(a^2-b^2\right) \sqrt{a+b \cos (c+dx)}}-\frac{2 \int \frac{b(A b-a B)+\left(-2 B a^2+A b a+b^2 B\right) \cos (c+dx)}{2 \sqrt{a+b \cos (c+dx)}} dx}{b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 a(A b-a B) \sin (c+dx)}{b d\left(a^2-b^2\right) \sqrt{a+b \cos (c+dx)}}-\frac{\int \frac{b(A b-a B)+\left(-2 B a^2+A b a+b^2 B\right) \cos (c+dx)}{\sqrt{a+b \cos (c+dx)}} dx}{b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 a(A b-a B) \sin (c+dx)}{b d\left(a^2-b^2\right) \sqrt{a+b \cos (c+dx)}}-\frac{\int \frac{b(A b-a B)+\left(-2 B a^2+A b a+b^2 B\right) \sin \left(c+dx+\frac{\pi}{2}\right)}{\sqrt{a+b \sin \left(c+dx+\frac{\pi}{2}\right)}} dx}{b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3231} \\
 & \frac{2 a(A b-a B) \sin (c+dx)}{b d\left(a^2-b^2\right) \sqrt{a+b \cos (c+dx)}}-\frac{\left(-2 a^2 B+a A b+b^2 B\right) \int \sqrt{a+b \cos (c+dx)} dx}{b}-\frac{\left(a^2-b^2\right)(A b-2 a B) \int \frac{1}{\sqrt{a+b \cos (c+dx)}} dx}{b}{b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 a(A b-a B) \sin (c+dx)}{b d\left(a^2-b^2\right) \sqrt{a+b \cos (c+dx)}}-\frac{\left(-2 a^2 B+a A b+b^2 B\right) \int \sqrt{a+b \sin \left(c+dx+\frac{\pi}{2}\right)} dx}{b}-\frac{\left(a^2-b^2\right)(A b-2 a B) \int \frac{1}{\sqrt{a+b \sin \left(c+dx+\frac{\pi}{2}\right)}} dx}{b}{b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3134} \\
 & \frac{2 a(A b-a B) \sin (c+dx)}{b d\left(a^2-b^2\right) \sqrt{a+b \cos (c+dx)}}-\frac{\left(-2 a^2 B+a A b+b^2 B\right) \sqrt{a+b \cos (c+dx)} \int \sqrt{\frac{a}{a+b}+\frac{b \cos (c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos (c+dx)}{a+b}}}-\frac{\left(a^2-b^2\right)(A b-2 a B) \int \frac{1}{\sqrt{a+b \sin \left(c+dx+\frac{\pi}{2}\right)}} dx}{b}{b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.328. $\int \frac{\cos (c+dx)(A+B \cos (c+dx))}{(a+b \cos (c+dx))^{3 / 2}} dx$

$$\begin{aligned}
 & \frac{2a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} - \frac{(-2a^2B + aAb + b^2B) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c + dx + \frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c + dx)}{a+b}}} - \frac{(a^2 - b^2)(Ab - 2aB) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \\
 & \frac{2a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} - \frac{2(-2a^2B + aAb + b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c + dx)}{a+b}}} - \frac{(a^2 - b^2)(Ab - 2aB) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b} \\
 & \frac{2a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} - \frac{2(-2a^2B + aAb + b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c + dx)}{a+b}}} - \frac{(a^2 - b^2)(Ab - 2aB) \sqrt{\frac{a+b \cos(c + dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c + dx)}{a+b}}} dx}{b \sqrt{a + b \cos(c + dx)}} \\
 & \frac{2a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} - \frac{2(-2a^2B + aAb + b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c + dx)}{a+b}}} - \frac{(a^2 - b^2)(Ab - 2aB) \sqrt{\frac{a+b \cos(c + dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c + dx + \frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a + b \cos(c + dx)}} \\
 & \frac{2a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} - \frac{2(-2a^2B + aAb + b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c + dx)}{a+b}}} - \frac{2(a^2 - b^2)(Ab - 2aB) \sqrt{\frac{a+b \cos(c + dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{bd \sqrt{a + b \cos(c + dx)}}
 \end{aligned}$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]`

3.328. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

```
output -(((2*(a*A*b - 2*a^2*B + b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(A*b - 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(b*(a^2 - b^2))) + (2*a*(A*b - a*B)*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])
```

3.328.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]
```

rule 3231 `Int[((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])/Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.328.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 518 vs. 2(252) = 504.

Time = 12.08 (sec) , antiderivative size = 519, normalized size of antiderivative = 2.54

method	result
default	$\frac{\sqrt{-(-2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - a + b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{2\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a-b} + \frac{a+b}{a-b}}\left(AbF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a-b}}\right) - 2BF\left(\sin\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a-b}}\right)\right)}{b^2\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}$
parts	Expression too large to display

input `int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)`

3.328. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2/b^2/(-2*
sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(A*b*Ellip
ticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-2*B*EllipticF(cos(1/2*d*x+1/2*
c),(-2*b/(a-b))^(1/2))*a+B*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)
)*a-B*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b)+2*a*(A*b-B*a)/b^
2/sin(1/2*d*x+1/2*c)^2/(2*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*sin(1/
2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*cos(1/2*d*x+1/2*c)*s
in(1/2*d*x+1/2*c)^2*b+EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2
*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1
/2)*a-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2
*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*b))/sin(1/2*
d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d

```

3.328.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.15 (sec) , antiderivative size = 683, normalized size of antiderivative = 3.35

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx =$$

$$\frac{6(Ba^2b^2 - Aab^3)\sqrt{b\cos(dx+c) + a\sin(dx+c)} + (\sqrt{2}(-4iBa^3b + 2iAa^2b^2 + 5iBab^3 - 3iAb^4)\cos(dx+c) + \dots)}{\dots}$$

input

```

integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm=
"fricas")

```

output

```
-1/3*(6*(B*a^2*b^2 - A*a*b^3)*sqrt(b*cos(d*x + c) + a)*sin(d*x + c) + (sqrt(2)*(-4*I*B*a^3*b + 2*I*A*a^2*b^2 + 5*I*B*a*b^3 - 3*I*A*b^4)*cos(d*x + c) + sqrt(2)*(-4*I*B*a^4 + 2*I*A*a^3*b + 5*I*B*a^2*b^2 - 3*I*A*a*b^3))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (sqrt(2)*(4*I*B*a^3*b - 2*I*A*a^2*b^2 - 5*I*B*a*b^3 + 3*I*A*b^4)*cos(d*x + c) + sqrt(2)*(4*I*B*a^4 - 2*I*A*a^3*b - 5*I*B*a^2*b^2 + 3*I*A*a*b^3))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(sqrt(2)*(2*I*B*a^2*b^2 - I*A*a*b^3 - I*B*b^4)*cos(d*x + c) + sqrt(2)*(2*I*B*a^3*b - I*A*a^2*b^2 - I*B*a*b^3))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*(sqrt(2)*(-2*I*B*a^2*b^2 + I*A*a*b^3 + I*B*b^4)*cos(d*x + c) + sqrt(2)*(-2*I*B*a^3*b + I*A*a^2*b^2 + I*B*a*b^3))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)))/((a^2*b^4 - b^6)*d*cos(d*x + c) + (a^3*b^3 - a*b^5)*d)
```

3.328.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)`

output Timed out

3.328.7 Maxima [F]

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

3.328. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)/(b*cos(d*x + c) + a)^(3/2), x)`

3.328.8 Giac [F]

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)/(b*cos(d*x + c) + a)^(3/2), x)`

3.328.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx) (A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2), x)`

3.329 $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

3.329.1 Optimal result 3066
 3.329.2 Mathematica [A] (verified) 3066
 3.329.3 Rubi [A] (verified) 3067
 3.329.4 Maple [A] (verified) 3070
 3.329.5 Fricas [C] (verification not implemented) 3071
 3.329.6 Sympy [F(-1)] 3072
 3.329.7 Maxima [F] 3072
 3.329.8 Giac [F] 3072
 3.329.9 Mupad [F(-1)] 3073

3.329.1 Optimal result

Integrand size = 25, antiderivative size = 185

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2(Ab - aB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{b(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{bd \sqrt{a + b \cos(c + dx)}} - \frac{2(Ab - aB) \sin(c + dx)}{(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

output

```
-2*(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b/(a^2-b^2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b/d/(a+b*cos(d*x+c))^(1/2)
```

3.329.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.82

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2 \left(- \left((a + b)(-Ab + aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) \right) + (a^2 - b^2) B \sqrt{a + b \cos(c + dx)} \right)}{(a - b)b(a + b)d \sqrt{a + b \cos(c + dx)}}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `(2*(-((a + b)*(-(A*b) + a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]) + (a^2 - b^2)*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + b*(-(A*b) + a*B)*Sin[c + d*x]))/((a - b)*b*(a + b)*d*Sqrt[a + b*Cos[c + d*x]])`

3.329.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.480$, Rules used = {3042, 3233, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{(a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{3/2}} dx \\
 & \quad \downarrow \text{3233} \\
 & -\frac{2 \int \frac{aA - bB + (Ab - aB) \cos(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{aA - bB + (Ab - aB) \cos(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{aA - bB + (Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3231} \\
 & \frac{B(a^2 - b^2) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} + \frac{(Ab - aB) \int \sqrt{a + b \cos(c + dx)} dx}{b} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}}
 \end{aligned}$$

3.329. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$

$$\frac{B(a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} + \frac{(Ab-aB) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{2(Ab-aB) \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{B(a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{(Ab-aB) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{a^2-b^2} - \frac{2(Ab-aB) \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{B(a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{(Ab-aB) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{a^2-b^2} - \frac{2(Ab-aB) \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{B(a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{2(Ab-aB) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{a^2-b^2} - \frac{2(Ab-aB) \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{B(a^2-b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} + \frac{2(Ab-aB) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{a^2-b^2} - \frac{2(Ab-aB) \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{B(a^2-b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{b \sqrt{a+b \cos(c+dx)}} + \frac{2(Ab-aB) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{a^2-b^2} - \frac{2(Ab-aB) \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

3.329. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow \text{3140} \\ & \frac{2B(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{2(Ab - aB) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{bd \sqrt{a+b \cos(c+dx)}} \\ & \frac{a^2 - b^2}{2(Ab - aB) \sin(c + dx)} \\ & \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `((2*(A*b - a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(a^2 - b^2)*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(a^2 - b^2) - (2*(A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])`

3.329.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])]/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

3.329.4 Maple [A] (verified)

Time = 8.42 (sec) , antiderivative size = 432, normalized size of antiderivative = 2.34

method	result
default	$\frac{\sqrt{-(-2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - a + b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{b\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (a + b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} \left(\frac{2B\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}}{a - b}} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a - b}}\right) - \frac{2(Ab - B^2)}{a - b} \right)$
parts	$\frac{2A\left(2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a - b}}\right)\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a - b}} + \frac{a + b}{a - b}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right)a - E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a - b}}\right)\sqrt{-2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a + b}d}{(a - b)(a + b)\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a + b}d}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

output
$$-(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*B/b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})-2*(A*b-B*a)/b/\sin(1/2*d*x+1/2*c)^2/(2*b*\sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2+b+EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a-EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b))/\sin(1/2*d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2+a+b)^{(1/2)}/d$$

3.329.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 610, normalized size of antiderivative = 3.30

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{6 (Bab^2 - Ab^3) \sqrt{b \cos(dx + c) + a \sin(dx + c)} - (\sqrt{2}(2i Ba^2b + i Aab^2 - 3i))}{(a + b \cos(c + dx))^{3/2}}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output
$$\frac{1}{3}*(6*(B*a*b^2 - A*b^3)*\sqrt{b*\cos(d*x + c) + a*\sin(d*x + c)} - (\sqrt{2}*(2*I*B*a^2*b + I*A*a*b^2 - 3*I*B*b^3)*\cos(d*x + c) + \sqrt{2}*(2*I*B*a^3 + I*A*a^2*b - 3*I*B*a*b^2))*\sqrt{b}*\text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) + 3*I*b*\sin(d*x + c) + 2*a)/b) - (\sqrt{2}*(-2*I*B*a^2*b - I*A*a*b^2 + 3*I*B*b^3)*\cos(d*x + c) + \sqrt{2}*(-2*I*B*a^3 - I*A*a^2*b + 3*I*B*a*b^2))*\sqrt{b}*\text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) - 3*I*b*\sin(d*x + c) + 2*a)/b) + 3*(\sqrt{2}*(-I*B*a*b^2 + I*A*b^3)*\cos(d*x + c) + \sqrt{2}*(-I*B*a^2*b + I*A*a*b^2))*\sqrt{b}*\text{weierstrassZeta}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, \text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) + 3*I*b*\sin(d*x + c) + 2*a)/b)) + 3*(\sqrt{2}*(I*B*a*b^2 - I*A*b^3)*\cos(d*x + c) + \sqrt{2}*(I*B*a^2*b - I*A*a*b^2))*\sqrt{b}*\text{weierstrassZeta}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, \text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) - 3*I*b*\sin(d*x + c) + 2*a)/b)))/((a^2*b^3 - b^5)*d*\cos(d*x + c) + (a^3*b^2 - a*b^4)*d)$$

3.329.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.329.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)`output `Timed out`**3.329.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="maxima")`output `integrate((B*cos(d*x + c) + A)/(b*cos(d*x + c) + a)^(3/2), x)`**3.329.8 Giac [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`output `integrate((B*cos(d*x + c) + A)/(b*cos(d*x + c) + a)^(3/2), x)`

3.329.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^(3/2), x)`output `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^(3/2), x)`

3.330
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.330.1 Optimal result 3074
 3.330.2 Mathematica [C] (verified) 3075
 3.330.3 Rubi [A] (verified) 3075
 3.330.4 Maple [A] (verified) 3079
 3.330.5 Fricas [F(-1)] 3080
 3.330.6 Sympy [F] 3081
 3.330.7 Maxima [F] 3081
 3.330.8 Giac [F] 3081
 3.330.9 Mupad [F(-1)] 3082

3.330.1 Optimal result

Integrand size = 31, antiderivative size = 190

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx =$$

$$-\frac{2(Ab - aB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{a(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2A \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{ad \sqrt{a + b \cos(c + dx)}} + \frac{2b(Ab - aB) \sin(c + dx)}{a(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

```
output 2*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)-2*(A*b-B*a)*
(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*
c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/a/(a^2-b^2)/d/((a+b*cos
(d*x+c))/(a+b))^(1/2)+2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))
/(a+b))^(1/2)/a/d/(a+b*cos(d*x+c))^(1/2)
```

3.330.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.78 (sec) , antiderivative size = 460, normalized size of antiderivative = 2.42

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{\cos(c + dx)(B + A \sec(c + dx)) \left(\frac{4a(-Ab + aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c - \dots)}{\sqrt{a+b \cos(c+dx)}}\right)}{\dots} \right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `(Cos[c + d*x]*(B + A*Sec[c + d*x])*(-(((4*a*(-A*b) + a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(2*a^2*A - 3*A*b^2 + a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] - ((2*I)*(A*b - a*B)*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)])))/(a*b*Sqrt[-(a + b)^(-1)]))/((-a + b)*(a + b)) + (4*b*(A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*Sqrt[a + b*Cos[c + d*x]])))/(2*a*d*(A + B*Cos[c + d*x]))`

3.330.3 Rubi [A] (verified)

Time = 1.26 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3479, 27, 3042, 3538, 25, 27, 3042, 3134, 3042, 3132, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

3.330. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2}} dx \\
& \quad \downarrow \text{3479} \\
& \frac{2 \int \frac{(-b(Ab - aB) \cos^2(c + dx) - a(Ab - aB) \cos(c + dx) + A(a^2 - b^2)) \sec(c + dx)}{2\sqrt{a + b \cos(c + dx)}} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{(-b(Ab - aB) \cos^2(c + dx) - a(Ab - aB) \cos(c + dx) + A(a^2 - b^2)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{-b(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)^2 - a(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right) + A(a^2 - b^2)}{\sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3538} \\
& \frac{-\frac{\int \frac{Ab(a^2 - b^2) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} - \left((Ab - aB) \int \sqrt{a + b \cos(c + dx)} dx \right)}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{25} \\
& \frac{\frac{\int \frac{Ab(a^2 - b^2) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx}{b} - (Ab - aB) \int \sqrt{a + b \cos(c + dx)} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{A(a^2 - b^2) \int \frac{\sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx - (Ab - aB) \int \sqrt{a + b \cos(c + dx)} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{A(a^2 - b^2) \int \frac{1}{\sin\left(c + dx + \frac{\pi}{2}\right) \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx - (Ab - aB) \int \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3134}
\end{aligned}$$

3.330. $\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{A(a^2 - b^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{(Ab-aB)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a(a^2 - b^2)} + \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{A(a^2 - b^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{(Ab-aB)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a(a^2 - b^2)} + \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3132} \\
& \frac{A(a^2 - b^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(Ab-aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a(a^2 - b^2)} + \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3286} \\
& \frac{A(a^2 - b^2) \int \frac{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx - \frac{2(Ab-aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a(a^2 - b^2)} + \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{A(a^2 - b^2) \int \frac{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx - \frac{2(Ab-aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a(a^2 - b^2)} + \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3284} \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} + \\
& \frac{2A(a^2 - b^2) \int \frac{\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}} - \frac{2(Ab-aB)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a(a^2 - b^2)}
\end{aligned}$$

3.330. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `((-2*(A*b - a*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*A*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/(a*(a^2 - b^2)) + (2*b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])`

3.330.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

```
rule 3286 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt
[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c +
d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]
```

```
rule 3479 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m +
1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e
+ f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +
2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)
*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n
}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Rat
ionalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(I
ntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])
))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x])], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.330.4 Maple [A] (verified)

Time = 9.94 (sec) , antiderivative size = 433, normalized size of antiderivative = 2.28

$$3.330. \quad \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

method	result
default	$\frac{\sqrt{-(-2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) - a + b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\dots} \left(\frac{2A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b(\cos^2(\frac{dx}{2} + \frac{c}{2})) + a - b}{a - b}} \Pi(\cos(\frac{dx}{2} + \frac{c}{2}), 2, \sqrt{-\frac{2b}{a - b}})}{a\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2}))b + (a + b)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}} - \dots \right)$
parts	$2A \left(2\cos(\frac{dx}{2} + \frac{c}{2})(\sin^2(\frac{dx}{2} + \frac{c}{2}))b^2 + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-\frac{2b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{a - b}} + \frac{a + b}{a - b} \Pi(\cos(\frac{dx}{2} + \frac{c}{2}), 2, \sqrt{-\frac{2b}{a - b}}) a^2 - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \dots \right)$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVER
BOSE)
```

```
output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*A/a*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2
*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1
/2*d*x+1/2*c),2,(-2*b/(a-b))^(1/2))-2*(-A*b+B*a)/a/sin(1/2*d*x+1/2*c)^2/(2
*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b+Ell
ipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*
c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*a-EllipticE(cos(1/2*d
*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b
))^1/2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*b)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*
d*x+1/2*c)^2+a+b)^(1/2)/d
```

3.330.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(3/2),x, algorithm=
"fracas")
```

```
output Timed out
```

3.330.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**(3/2), x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a + b*cos(c + d*x))**(3/2), x)`

3.330.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(3/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)/(b*cos(d*x + c) + a)^(3/2), x)`

3.330.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(3/2), x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)/(b*cos(d*x + c) + a)^(3/2), x)`

3.330.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx) (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(3/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(3/2)), x)`

3.331 $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

3.331.1 Optimal result 3083
 3.331.2 Mathematica [C] (verified) 3084
 3.331.3 Rubi [A] (verified) 3084
 3.331.4 Maple [B] (verified) 3091
 3.331.5 Fricas [F(-1)] 3092
 3.331.6 Sympy [F] 3093
 3.331.7 Maxima [F] 3093
 3.331.8 Giac [F] 3093
 3.331.9 Mupad [F(-1)] 3094

3.331.1 Optimal result

Integrand size = 33, antiderivative size = 303

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx =$$

$$\frac{(a^2 A - 3Ab^2 + 2abB) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{a^2 (a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{A \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{ad \sqrt{a + b \cos(c + dx)}}$$

$$- \frac{(3Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{a^2 d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{b(a^2 A - 3Ab^2 + 2abB) \sin(c + dx)}{a^2 (a^2 - b^2) d \sqrt{a + b \cos(c + dx)}} + \frac{A \tan(c + dx)}{ad \sqrt{a + b \cos(c + dx)}}$$

```
output b*(A*a^2-3*A*b^2+2*B*a*b)*sin(d*x+c)/a^2/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)
)-(A*a^2-3*A*b^2+2*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)
/a^2/(a^2-b^2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+A*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))
*((a+b*cos(d*x+c))/(a+b))^(1/2)/a/d/(a+b*cos(d*x+c))^(1/2)-(3*A*b-2*B
*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x
+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/a^2/d/(a
+b*cos(d*x+c))^(1/2)+A*tan(d*x+c)/a/d/(a+b*cos(d*x+c))^(1/2)
```

3.331.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.04 (sec) , antiderivative size = 482, normalized size of antiderivative = 1.59

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{8ab(-Ab+aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + \frac{2(-7a^2Ab+9Ab^3+4a^3B-6ab^2B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{a+b\cos(c+dx)}}}{\sqrt{a+b\cos(c+dx)}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^(3/2), x]`

output `(((-8*a*b*(-A*b) + a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(-7*a^2*A*b^3 + 4*a^3*B - 6*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + ((2*I)*(a^2*A - 3*A*b^2 + 2*a*b*B)*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[-((b*(1 + Cos[c + d*x]))/(a - b))]*Csc[c + d*x]*(2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] - b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)])))/(a*b*Sqrt[-(a + b)^(-1)])/(a - b)*(a + b) + (4*(a*A*(a^2 - b^2) + b*(a^2*A - 3*A*b^2 + 2*a*b*B)*Cos[c + d*x])*Tan[c + d*x])/((a^2 - b^2)*Sqrt[a + b*Cos[c + d*x]])/(4*a^2*d)`

3.331.3 Rubi [A] (verified)

Time = 2.70 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.11, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3479, 27, 3042, 3535, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

3.331. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \int \frac{A + B \sin \left(c + dx + \frac{\pi}{2} \right)}{\sin \left(c + dx + \frac{\pi}{2} \right)^2 \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{3/2}} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{\int -\frac{(-Ab \cos^2(c+dx)+3Ab-2aB) \sec(c+dx)}{2(a+b \cos(c+dx))^{3/2}} dx}{a} + \frac{A \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{A \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{(-Ab \cos^2(c+dx)+3Ab-2aB) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx}{2a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{-Ab \sin(c+dx+\frac{\pi}{2})^2+3Ab-2aB}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{2a} \\
 & \quad \downarrow \text{3535} \\
 & \frac{A \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{2 \int \frac{(b(Aa^2+2bBa-3Ab^2) \cos^2(c+dx)-2ab(Ab-aB) \cos(c+dx)+(a^2-b^2)(3Ab-2aB)) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)} a(a^2-b^2)} dx}{2a} - \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b \cos(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{A \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{(b(Aa^2+2bBa-3Ab^2) \cos^2(c+dx)-2ab(Ab-aB) \cos(c+dx)+(a^2-b^2)(3Ab-2aB)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)} a(a^2-b^2)} dx}{2a} - \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b \cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{b(Aa^2+2bBa-3Ab^2) \sin(c+dx+\frac{\pi}{2})^2-2ab(Ab-aB) \sin(c+dx+\frac{\pi}{2})+(a^2-b^2)(3Ab-2aB)}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})} a(a^2-b^2)} dx}{2a} - \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b \cos(c+dx)}} \\
 & \quad \downarrow \text{3538}
 \end{aligned}$$

3.331. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \int \frac{(b(a^2-b^2)(3Ab-2aB)-aAb(a^2-b^2)\cos(c+dx)) \sec(c+dx)}{\sqrt{a+b\cos(c+dx)} b} dx}{(a^2A+2abB-3Ab^2) \int \sqrt{a+b\cos(c+dx)} dx - \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \cdot 2a$$

25

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \int \frac{(b(a^2-b^2)(3Ab-2aB)-aAb(a^2-b^2)\cos(c+dx)) \sec(c+dx)}{\sqrt{a+b\cos(c+dx)} b} dx}{(a^2A+2abB-3Ab^2) \int \sqrt{a+b\cos(c+dx)} dx + \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \cdot 2a$$

3042

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \int \frac{b(a^2-b^2)(3Ab-2aB)-aAb(a^2-b^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} b} dx}{(a^2A+2abB-3Ab^2) \int \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} dx + \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \cdot 2a$$

3134

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \int \frac{b(a^2-b^2)(3Ab-2aB)-aAb(a^2-b^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} b} dx}{(a^2A+2abB-3Ab^2) \sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}} dx + \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \cdot 2a$$

3042

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \int \frac{b(a^2-b^2)(3Ab-2aB)-aAb(a^2-b^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} b} dx}{(a^2A+2abB-3Ab^2) \sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx + \frac{2b(a^2A+2abB-3Ab^2) \sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \cdot 2a$$

3132

3.331. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \int \frac{b(a^2-b^2)(3Ab-2aB) - aAb(a^2-b^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2b(a^2A+2abB-3Ab^2)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

2a

↓ 3481

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{b(a^2-b^2)(3Ab-2aB) \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx - aAb(a^2-b^2) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{2a} - \frac{2b(a^2A+2abB-3Ab^2)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

2a

↓ 3042

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{b(a^2-b^2)(3Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - aAb(a^2-b^2) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{2a}$$

2a

↓ 3142

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{aAb(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{a(a^2-b^2)} - \frac{b(a^2-b^2)(3Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b}}{2a} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

2a

↓ 3042

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{aAb(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{a(a^2-b^2)} - \frac{b(a^2-b^2)(3Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b}}{2a} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

2a

↓ 3140

3.331. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx$

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{b(a^2-b^2)(3Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2aAb(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2a}{a(a^2-b^2)}$$

3286

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{b(a^2-b^2)(3Ab-2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx - \frac{2aAb(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{\sqrt{a+b\cos(c+dx)}} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2a}{a(a^2-b^2)}$$

3042

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{b(a^2-b^2)(3Ab-2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx - \frac{2aAb(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{\sqrt{a+b\cos(c+dx)}} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2a}{a(a^2-b^2)}$$

3284

$$\frac{\frac{A \tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} - \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} + \frac{2b(a^2-b^2)(3Ab-2aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2aAb(a^2-b^2)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{d\sqrt{a+b\cos(c+dx)}}}{b} + \frac{2(a^2A+2abB-3Ab^2)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2a}{a(a^2-b^2)}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^(3/2), x]`

3.331. $\int \frac{(A+B\cos(c+dx))\sec^2(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx$

```
output -1/2*(((2*(a^2*A - 3*A*b^2 + 2*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((-2*a*A*b*(a^2 - b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*b*(a^2 - b^2)*(3*A*b - 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/(a*(a^2 - b^2)) - (2*b*(a^2*A - 3*A*b^2 + 2*a*b*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]))/a + (A*Tan[c + d*x])/(a*d*Sqrt[a + b*Cos[c + d*x]])
```

3.331.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*SIN[c + d*x]]/Sqrt[(a + b*SIN[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```


rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^n), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3535 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] :>
Simp[(-(A*b^2 + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
in[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m
+ 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin
[e + f*x])^n*Simp[a*(m + 1)*(b*c - a*d)*(A + C) + d*(A*b^2 + a^2*C)*(m + n
+ 2) - (c*(A*b^2 + a^2*C) + b*(m + 1)*(b*c - a*d)*(A + C))*Sin[e + f*x] - d
*(A*b^2 + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) ||
!(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a,
0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.331.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 911 vs. $2(374) = 748$.

Time = 14.11 (sec) , antiderivative size = 912, normalized size of antiderivative = 3.01

method	result	size
default	Expression too large to display	912
parts	Expression too large to display	1276

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNV
ERBOSE)
```

output
$$\begin{aligned} & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*A/a*(-\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)+1/2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})-1/2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+1/2/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+1/2/a*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-2*(-A*b+B*a)/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-2*(A*b-B*a)*b/a^2/\sin(1/2*d*x+1/2*c)^2/(2*b*\sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*b+EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a-EllipticE(... \end{aligned}$$

3.331.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output Timed out

3.331.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a + b*cos(c + d*x))**(3/2), x)`

3.331.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^2}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^2/(b*cos(d*x + c) + a)^(3/2), x)`

3.331.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^2}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^2/(b*cos(d*x + c) + a)^(3/2), x)`

3.331.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^2 (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^(3/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^(3/2)), x)`

3.332
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.332.1 Optimal result	3095
3.332.2 Mathematica [C] (verified)	3096
3.332.3 Rubi [A] (verified)	3097
3.332.4 Maple [B] (verified)	3105
3.332.5 Fricas [F(-1)]	3106
3.332.6 Sympy [F]	3107
3.332.7 Maxima [F(-1)]	3107
3.332.8 Giac [F]	3107
3.332.9 Mupad [F(-1)]	3108

3.332.1 Optimal result

Integrand size = 33, antiderivative size = 398

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{(7a^2 Ab - 15Ab^3 - 4a^3 B + 12ab^2 B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4a^3 (a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(5Ab - 4aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4a^2 d \sqrt{a + b \cos(c + dx)}} + \frac{(4a^2 A + 15Ab^2 - 12abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4a^3 d \sqrt{a + b \cos(c + dx)}} - \frac{b(7a^2 Ab - 15Ab^3 - 4a^3 B + 12ab^2 B) \sin(c + dx)}{4a^3 (a^2 - b^2) d \sqrt{a + b \cos(c + dx)}} - \frac{(5Ab - 4aB) \tan(c + dx)}{4a^2 d \sqrt{a + b \cos(c + dx)}} + \frac{A \sec(c + dx) \tan(c + dx)}{2ad \sqrt{a + b \cos(c + dx)}}$$

output
$$\begin{aligned} & -1/4*b*(7*A*a^2*b-15*A*b^3-4*B*a^3+12*B*a*b^2)*\sin(d*x+c)/a^3/(a^2-b^2)/d/ \\ & (a+b*\cos(d*x+c))^(1/2)+1/4*(7*A*a^2*b-15*A*b^3-4*B*a^3+12*B*a*b^2)*(\cos(1/ \\ & 2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^(1 \\ & /2)*(b/(a+b))^(1/2))*(a+b*\cos(d*x+c))^(1/2)/a^3/(a^2-b^2)/d/((a+b*\cos(d*x+ \\ & c))/(a+b))^(1/2)-1/4*(5*A*b-4*B*a)*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d* \\ & x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*\cos(d \\ & *x+c))/(a+b))^(1/2)/a^2/d/(a+b*\cos(d*x+c))^(1/2)+1/4*(4*A*a^2+15*A*b^2-12* \\ & B*a*b)*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticPi}(\sin(1/2* \\ & d*x+1/2*c), 2, 2^(1/2)*(b/(a+b))^(1/2))*((a+b*\cos(d*x+c))/(a+b))^(1/2)/a^3/d \\ & /((a+b*\cos(d*x+c))^(1/2)-1/4*(5*A*b-4*B*a)*\tan(d*x+c)/a^2/d/(a+b*\cos(d*x+c) \\ &)^(1/2)+1/2*A*\sec(d*x+c)*\tan(d*x+c)/a/d/(a+b*\cos(d*x+c))^(1/2) \end{aligned}$$

3.332.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.72 (sec) , antiderivative size = 546, normalized size of antiderivative = 1.37

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{8ab(a^2A - 5Ab^2 + 4abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + 2(8a^4A + 29a^2Ab^2 - 45Ab^4)}{\sqrt{a+b \cos(c+dx)}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^(3/2), x]`

output
$$\begin{aligned} & (-(((8*a*b*(a^2*A - 5*A*b^2 + 4*a*b*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/ (a + b)]* \\ & \text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(8*a^4 \\ & 4*A + 29*a^2*A*b^2 - 45*A*b^4 - 28*a^3*b*B + 36*a*b^3*B)*\text{Sqrt}[(a + b*\text{Cos}[c \\ & + d*x])/ (a + b)]*\text{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*Co \\ & s[c + d*x]] - ((2*I)*(-7*a^2*A*b + 15*A*b^3 + 4*a^3*B - 12*a*b^2*B)*\text{Sqrt}[- \\ & ((b*(-1 + \text{Cos}[c + d*x]))/(a + b))]*\text{Sqrt}[-((b*(1 + \text{Cos}[c + d*x]))/(a - b))] \\ & * \text{Csc}[c + d*x]*(-2*a*(a - b)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^(-1)]*\text{Sqrt}[a \\ & + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[- \\ & (a + b)^(-1)]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*\text{EllipticPi}[(\\ & a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^(-1)]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b) \\ & / (a - b)])))/(a*b*\text{Sqrt}[-(a + b)^(-1)])/((-a + b)*(a + b))) + (4*(b*(-7*a^2 \\ & 2*A*b + 15*A*b^3 + 4*a^3*B - 12*a*b^2*B)*\text{Sin}[c + d*x] + a*(a^2 - b^2)*(2*a \\ & *A + (-5*A*b + 4*a*B)*\text{Cos}[c + d*x])* \text{Sec}[c + d*x]*\text{Tan}[c + d*x]))/((a^2 - b^2) \\ & *\text{Sqrt}[a + b*\text{Cos}[c + d*x]]))/(16*a^3*d) \end{aligned}$$

3.332.
$$\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.332.3 Rubi [A] (verified)

Time = 3.51 (sec) , antiderivative size = 425, normalized size of antiderivative = 1.07, number of steps used = 24, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{\int -\frac{(-3Ab\cos^2(c+dx)-2aA\cos(c+dx)+5Ab-4aB)\sec^2(c+dx)}{2(a+b\cos(c+dx))^{3/2}} dx}{2a} + \frac{A\tan(c+dx)\sec(c+dx)}{2ad\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{A\tan(c+dx)\sec(c+dx)}{2ad\sqrt{a+b\cos(c+dx)}} - \frac{\int \frac{(-3Ab\cos^2(c+dx)-2aA\cos(c+dx)+5Ab-4aB)\sec^2(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx}{4a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A\tan(c+dx)\sec(c+dx)}{2ad\sqrt{a+b\cos(c+dx)}} - \frac{\int \frac{-3Ab\sin(c+dx+\frac{\pi}{2})^2-2aA\sin(c+dx+\frac{\pi}{2})+5Ab-4aB}{\sin(c+dx+\frac{\pi}{2})^2(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{4a} \\
 & \quad \downarrow \text{3534} \\
 & \frac{A\tan(c+dx)\sec(c+dx)}{2ad\sqrt{a+b\cos(c+dx)}} - \frac{\int -\frac{(4Aa^2-12bBa+6Ab\cos(c+dx)a+15Ab^2-b(5Ab-4aB)\cos^2(c+dx))\sec(c+dx)}{2(a+b\cos(c+dx))^{3/2}} dx}{a} + \frac{(5Ab-4aB)\tan(c+dx)}{ad\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{(4Aa^2 - 12bBa + 6Ab \cos(c+dx)a + 15Ab^2 - b(5Ab - 4aB) \cos^2(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx \\
 & \frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{4a}{2a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{4Aa^2 - 12bBa + 6Ab \sin(c+dx + \frac{\pi}{2})a + 15Ab^2 - b(5Ab - 4aB) \sin(c+dx + \frac{\pi}{2})^2}{\sin(c+dx + \frac{\pi}{2})(a+b \sin(c+dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{4a}{2a} \\
 & \quad \downarrow \text{3534} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{(b(-4Ba^3 + 7Aba^2 + 12b^2Ba - 15Ab^3) \cos^2(c+dx) + 2ab(Aa^2 + 4bBa - 5Ab^2) \cos(c+dx) + (a^2 - b^2)(4Aa^2 - 12bBa + 15Ab^2)) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)} a(a^2 - b^2)} dx \\
 & \frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{4a}{2a} \\
 & \quad \downarrow \text{27} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{(b(-4Ba^3 + 7Aba^2 + 12b^2Ba - 15Ab^3) \cos^2(c+dx) + 2ab(Aa^2 + 4bBa - 5Ab^2) \cos(c+dx) + (a^2 - b^2)(4Aa^2 - 12bBa + 15Ab^2)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)} a(a^2 - b^2)} dx \\
 & \frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{4a}{2a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{b(-4Ba^3 + 7Aba^2 + 12b^2Ba - 15Ab^3) \sin(c+dx + \frac{\pi}{2})^2 + 2ab(Aa^2 + 4bBa - 5Ab^2) \sin(c+dx + \frac{\pi}{2}) + (a^2 - b^2)(4Aa^2 - 12bBa + 15Ab^2)}{\sin(c+dx + \frac{\pi}{2})\sqrt{a+b \sin(c+dx + \frac{\pi}{2})} a(a^2 - b^2)} dx \\
 & \frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{4a}{2a} \\
 & \quad \downarrow \text{3538}
 \end{aligned}$$

3.332. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{(-4a^3B+7a^2Ab+12ab^2B-15Ab^3) \int \sqrt{a+b \cos(c+dx)} dx - \frac{(b(a^2-b^2)(4Aa^2-12bBa+15Ab^2)-ab(a^2-b^2)(5Ab-4aB) \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}}}{a(a^2-b^2)}}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} - \frac{2a}{4a}}$$

25

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{(b(a^2-b^2)(4Aa^2-12bBa+15Ab^2)-ab(a^2-b^2)(5Ab-4aB) \cos(c+dx)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx + (-4a^3B+7a^2Ab+12ab^2B-15Ab^3) \int \sqrt{a+b \cos(c+dx)} dx}{a(a^2-b^2)}}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} - \frac{2a}{4a}}$$

3042

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{b(a^2-b^2)(4Aa^2-12bBa+15Ab^2)-ab(a^2-b^2)(5Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + (-4a^3B+7a^2Ab+12ab^2B-15Ab^3) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{a(a^2-b^2)}}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} - \frac{2a}{4a}}$$

3134

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{b(a^2-b^2)(4Aa^2-12bBa+15Ab^2)-ab(a^2-b^2)(5Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{(-4a^3B+7a^2Ab+12ab^2B-15Ab^3) \sqrt{a+b \cos(c+dx)}}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{a(a^2-b^2)}}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} - \frac{2a}{4a}}$$

3042

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{b(a^2-b^2)(4Aa^2-12bBa+15Ab^2)-ab(a^2-b^2)(5Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{(-4a^3B+7a^2Ab+12ab^2B-15Ab^3) \sqrt{a+b \cos(c+dx)}}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{a(a^2-b^2)}}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} - \frac{2a}{4a}}$$

3132

3.332. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{b(a^2-b^2)(4Aa^2-12bBa+15Ab^2) - ab(a^2-b^2)(5Ab-4aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} = \frac{\frac{2(-4a^3B+7a^2Ab+12ab^2B-15Ab^3)\sqrt{a+b \cos(c+dx)}}{d\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{a(a^2-b^2)}{2a}}{4a}$$

↓ 3481

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - ab(a^2-b^2)(5Ab-4aB) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} = \frac{\frac{2(-4a^3B+7a^2Ab+12ab^2)}{2a} + \frac{a(a^2-b^2)}{2a}}{4a}$$

↓ 3042

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - ab(a^2-b^2)(5Ab-4aB) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} = \frac{\frac{a(a^2-b^2)}{2a} + 2}{4a}$$

↓ 3142

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \int \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(a^2-b^2)(5Ab-4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b}{a+b \cos(c+dx)}}}}{\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}}} = \frac{a(a^2-b^2)}{4a}$$

↓ 3042

3.332. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(a^2-b^2)(5Ab-4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}}{\sqrt{a+b \cos(c+dx)}}}{b}$$

$$\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{a(a^2-b^2)}{4a}$$

3140

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2ab(a^2-b^2)(5Ab-4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2})}{d\sqrt{a+b \cos(c+dx)}}$$

$$\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{a(a^2-b^2)}{4a}$$

3286

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - \frac{2ab(a^2-b^2)(5Ab-4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}}{\sqrt{a+b \cos(c+dx)}}}{b}$$

$$\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{a(a^2-b^2)}{4a}$$

3042

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad\sqrt{a+b \cos(c+dx)}} - \frac{b(a^2-b^2)(4a^2A-12abB+15Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx - \frac{2ab(a^2-b^2)(5Ab-4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{d\sqrt{a+b \cos(c+dx)}}$$

$$\frac{(5Ab-4aB) \tan(c+dx)}{ad\sqrt{a+b \cos(c+dx)}} - \frac{a(a^2-b^2)}{4a}$$

3284

3.332. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\frac{A \tan(c + dx) \sec(c + dx)}{2ad\sqrt{a + b \cos(c + dx)}} - \frac{2b(a^2 - b^2)(4a^2A - 12abB + 15Ab^2)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}} - \frac{2ab(a^2 - b^2)(5Ab - 4aB)\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticE}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b \cos(c+dx)}}$$

$$\frac{(5Ab - 4aB) \tan(c + dx)}{ad\sqrt{a + b \cos(c + dx)}} - \frac{a(a^2 - b^2)}{4a}$$

4a

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^(3/2), x]`

output `(A*Sec[c + d*x]*Tan[c + d*x])/(2*a*d*Sqrt[a + b*Cos[c + d*x]]) - (-1/2*(((2*(7*a^2*A*b - 15*A*b^3 - 4*a^3*B + 12*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((-2*a*b*(a^2 - b^2)*(5*A*b - 4*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (2*b*(a^2 - b^2)*(4*a^2*A + 15*A*b^2 - 12*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])))/b)/(a*(a^2 - b^2)) - (2*b*(7*a^2*A*b - 15*A*b^3 - 4*a^3*B + 12*a*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])/a + ((5*A*b - 4*a*B)*Tan[c + d*x])/(a*d*Sqrt[a + b*Cos[c + d*x]])/(4*a)`

3.332.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])]), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
] && NeQ[c^2 - d^2, 0]
```

3.332.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1567 vs. $2(455) = 910$.

Time = 19.36 (sec) , antiderivative size = 1568, normalized size of antiderivative = 3.94

method	result	size
default	Expression too large to display	1568
parts	Expression too large to display	2446

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNV
ERBOSE)
```


output
$$\begin{aligned}
& -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*A/a*(-1/ \\
& 2*\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c) \\
& ^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)^2+3/4*b/a^2*\cos(1/2*d*x+1/2*c)*(-2*\sin \\
& (1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2* \\
& c)^2-1)-1/8*b/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a- \\
& b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/ \\
& 2)}*EllipticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+3/8/a*(\sin(1/2*d*x+1/2* \\
& *c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+ \\
& 1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*EllipticE(\cos(1/2*d*x+1/2*c) \\
&),(-2*b/(a-b))^{(1/2)})-3/8*b^2/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1 \\
& /2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2 \\
& *d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})-1/2* \\
& (\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/ \\
& (-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos \\
& (1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-3/8/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\
&)*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4*b+ \\
& (a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a \\
& -b))^{(1/2)})*b^2)+2*(-A*b+B*a)/a^2*(-\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1 \\
& /2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)+1/2 \\
& *(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1...
\end{aligned}$$

3.332.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output Timed out

3.332.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a + b*cos(c + d*x))**(3/2), x)`

3.332.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(3/2),x, algorithm m="maxima")`

output `Timed out`

3.332.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^3}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^3/(b*cos(d*x + c) + a)^(3/2), x)`

3.332.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^3 (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^(3/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^(3/2)), x)`

3.333
$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$$

3.333.1 Optimal result 3109
 3.333.2 Mathematica [A] (verified) 3110
 3.333.3 Rubi [A] (verified) 3111
 3.333.4 Maple [B] (verified) 3118
 3.333.5 Fricas [C] (verification not implemented) 3119
 3.333.6 Sympy [F(-1)] 3120
 3.333.7 Maxima [F] 3121
 3.333.8 Giac [F] 3121
 3.333.9 Mupad [F(-1)] 3121

3.333.1 Optimal result

Integrand size = 33, antiderivative size = 550

$$\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx =$$

$$\frac{2(80a^5Ab - 140a^3Ab^3 + 40aAb^5 - 128a^6B + 212a^4b^2B - 55a^2b^4B - 9b^6B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right) + 2(80a^4Ab - 80a^2Ab^3 - 5Ab^5 - 128a^5B + 116a^3b^2B + 17ab^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + 2a(Ab - aB) \cos^3(c+dx) \sin(c+dx) + 2a(5a^2Ab - 9Ab^3 - 8a^3B + 12ab^2B) \cos^2(c+dx) \sin(c+dx) + 2(40a^4Ab - 65a^2Ab^3 + 5Ab^5 - 64a^5B + 98a^3b^2B - 14ab^4B) \sqrt{a+b \cos(c+dx)} \sin(c+dx) + 2(30a^3Ab - 50aAb^3 - 48a^4B + 71a^2b^2B - 3b^4B) \cos(c+dx) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15b^5(a^2 - b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}} + 15b^5(a^2 - b^2) d \sqrt{a+b \cos(c+dx)} + 3b(a^2 - b^2) d(a+b \cos(c+dx))^{3/2} + 3b^2(a^2 - b^2)^2 d \sqrt{a+b \cos(c+dx)} + 15b^4(a^2 - b^2)^2 d + 15b^3(a^2 - b^2)^2 d}$$

output $\frac{2}{3}a(Ab-Ba)\cos(dx+c)^3\sin(dx+c)/b/(a^2-b^2)/d/(a+b\cos(dx+c))^{3/2} + \frac{2}{3}a(5Aa^2b-9Ab^3-8B^2a^3+12B^2ab^2)\cos(dx+c)^2\sin(dx+c)/b^2/(a^2-b^2)^2/d/(a+b\cos(dx+c))^{1/2} + \frac{2}{15}(40Aa^4b-65Aa^2b^3+5Ab^5-64B^2a^5+98B^2ab^2-14B^2ab^4)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^4/(a^2-b^2)^2/d - \frac{2}{15}(30Aa^3b-50Aa^2b^3-48B^2a^4+71B^2ab^2-3B^2b^4)\cos(dx+c)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^3/(a^2-b^2)^2/d - \frac{2}{15}(80Aa^5b-140Aa^3b^3+40Aa^2b^5-128B^2a^6+212B^2a^4b^2-55B^2a^2b^4-9B^2b^6)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticE}(\sin(1/2dx+1/2c), 2^{1/2}*(b/(a+b))^{1/2})*(a+b\cos(dx+c))^{1/2}/b^5/(a^2-b^2)^2/d/((a+b\cos(dx+c))/(a+b))^{1/2} + \frac{2}{15}(80Aa^4b-80Aa^2b^3-5Ab^5-128B^2a^5+116B^2a^3b^2+17B^2ab^4)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticF}(\sin(1/2dx+1/2c), 2^{1/2}*(b/(a+b))^{1/2})*((a+b\cos(dx+c))/(a+b))^{1/2}/b^5/(a^2-b^2)/d/(a+b\cos(dx+c))^{1/2}$

3.333.2 Mathematica [A] (verified)

Time = 3.19 (sec) , antiderivative size = 372, normalized size of antiderivative = 0.68

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \frac{2\left(\frac{a+b\cos(c+dx)}{a+b}\right)^{3/2} \left(b^2(20a^4Ab-35a^2Ab^3-5Ab^5-32a^5B+44a^3b^2B+8ab^4B)\text{EllipticE}\left(\frac{c+dx}{2}, \frac{2b}{a+b}\right) - (-80a^5Ab+140a^3Ab^3-40a^2Ab^5+128a^6B-212a^4b^2B+55a^2b^4B+9b^6B)\left((a+b)\text{EllipticE}\left(\frac{c+dx}{2}, \frac{2b}{a+b}\right) - a\text{EllipticF}\left(\frac{c+dx}{2}, \frac{2b}{a+b}\right)\right)\right)}{(a-b)^2(a+b)+b\left((10a^4(-Ab)+aB)\text{Sin}[c+dx]/(a^2-b^2)-(10a^3(-8a^2Ab+12Ab^3+11a^3B-15ab^2B)*(a+b\cos[c+dx])\text{Sin}[c+dx])/(a^2-b^2)^2+2(5Ab-14aB)*(a+b\cos[c+dx])^2\text{Sin}[c+dx]+3bB*(a+b\cos[c+dx])^2\text{Sin}[2(c+dx)])\right)}/(15b^5d*(a+b\cos[c+dx]))^{3/2}}$$

input `Integrate[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2), x]`

output $((-2*((a+b\cos[c+dx])/(a+b))^{3/2}*(b^2*(20a^4Ab-35a^2Ab^3-5Ab^5-32a^5B+44a^3b^2B+8ab^4B)*\text{EllipticF}[(c+dx)/2, (2b)/(a+b)] - (-80a^5Ab+140a^3Ab^3-40a^2Ab^5+128a^6B-212a^4b^2B+55a^2b^4B+9b^6B)*((a+b)\text{EllipticE}[(c+dx)/2, (2b)/(a+b)] - a\text{EllipticF}[(c+dx)/2, (2b)/(a+b)])))/((a-b)^2(a+b)+b\left((10a^4(-Ab)+aB)\text{Sin}[c+dx]/(a^2-b^2)-(10a^3(-8a^2Ab+12Ab^3+11a^3B-15ab^2B)*(a+b\cos[c+dx])\text{Sin}[c+dx])/(a^2-b^2)^2+2(5Ab-14aB)*(a+b\cos[c+dx])^2\text{Sin}[c+dx]+3bB*(a+b\cos[c+dx])^2\text{Sin}[2(c+dx)])\right)}/(15b^5d*(a+b\cos[c+dx]))^{3/2})$

3.333.3 Rubi [A] (verified)

Time = 3.02 (sec) , antiderivative size = 555, normalized size of antiderivative = 1.01, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3528, 27, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^4(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 3468

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} - \frac{2 \int -\frac{\cos^2(c+dx)(-((-8Ba^2+5Aba+3b^2B)\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+6a(Ab-aB)))}{2(a+b\cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)}$$

↓ 27

$$\frac{\int \frac{\cos^2(c+dx)(-((-8Ba^2+5Aba+3b^2B)\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+6a(Ab-aB)))}{(a+b\cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^2((8Ba^2-5Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+6a(Ab-aB))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3526

$$\frac{2a(-8a^3B+5a^2Ab+12ab^2B-9Ab^3)\sin(c+dx)\cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} - \frac{2 \int -\frac{\cos(c+dx)(-((-48Ba^4+30Aba^3+71b^2Ba^2-50Ab^3a-3b^4B)\cos^2(c+dx))+b(2Ba^3+2\sqrt{a+b\cos(c+dx)})b(a^2-b^2))}{2\sqrt{a+b\cos(c+dx)}} dx}{b(a^2-b^2)}$$

↓

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

3.333. $\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

↓ 27

$$\int \frac{\cos(c+dx) \left(- \left((-48Ba^4 + 30Aba^3 + 71b^2Ba^2 - 50Ab^3a - 3b^4B) \cos^2(c+dx) \right) + b(2Ba^3 + Aa^2 - 6b^2Ba + 3Ab^3) \cos(c+dx) + 4a(-8Ba^3 + 5Aba^2 + 12b^2Ba - 9Ab^3) \right)}{\sqrt{a+b \cos(c+dx)} b(a^2-b^2)} dx$$

$3b(a^2 - b^2)$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right) \left((48Ba^4 - 30Aba^3 - 71b^2Ba^2 + 50Ab^3a + 3b^4B) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + b(2Ba^3 + Aa^2 - 6b^2Ba + 3Ab^3) \sin\left(c+dx+\frac{\pi}{2}\right) + 4a(-8Ba^3 + 5Aba^2 + 12b^2Ba - 9Ab^3) \right)}{\sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} b(a^2-b^2)} dx$$

$3b(a^2 - b^2)$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3528

$$2 \int - \frac{-3(-64Ba^5 + 40Aba^4 + 98b^2Ba^3 - 65Ab^3a^2 - 14b^4Ba + 5Ab^5) \cos^2(c+dx) - b(-16Ba^4 + 10Aba^3 + 27b^2Ba^2 - 30Ab^3a + 9b^4B) \cos(c+dx) + 2a(-48Ba^4 + 30Aba^3 + 71b^2Ba^2 - 50Ab^3a - 3b^4B)}{2\sqrt{a+b \cos(c+dx)} 5b} dx$$

$b(a^2 - b^2)$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 27

$$\int - \frac{-3(-64Ba^5 + 40Aba^4 + 98b^2Ba^3 - 65Ab^3a^2 - 14b^4Ba + 5Ab^5) \cos^2(c+dx) - b(-16Ba^4 + 10Aba^3 + 27b^2Ba^2 - 30Ab^3a + 9b^4B) \cos(c+dx) + 2a(-48Ba^4 + 30Aba^3 + 71b^2Ba^2 - 50Ab^3a - 3b^4B)}{\sqrt{a+b \cos(c+dx)} 5b} dx$$

$b(a^2 - b^2)$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$\int - \frac{-3(-64Ba^5 + 40Aba^4 + 98b^2Ba^3 - 65Ab^3a^2 - 14b^4Ba + 5Ab^5) \sin\left(c+dx+\frac{\pi}{2}\right)^2 - b(-16Ba^4 + 10Aba^3 + 27b^2Ba^2 - 30Ab^3a + 9b^4B) \sin\left(c+dx+\frac{\pi}{2}\right) + 2a(-48Ba^4 + 30Aba^3 + 71b^2Ba^2 - 50Ab^3a - 3b^4B)}{\sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} 5b} dx$$

$b(a^2 - b^2)$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3502

3.333. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\int \frac{3(b(-32Ba^5+20Aba^4+44b^2Ba^3-35Ab^3a^2+8b^4Ba-5Ab^5)+(-128Ba^6+80Aba^5+212b^2Ba^4-140Ab^3a^3-55b^4Ba^2+40Ab^5a-9b^6B)\cos(c+dx))}{2\sqrt{a+b\cos(c+dx)}} dx - \frac{2(-64a^5B+17ab^4B-80a^2Ab^3+116a^3b^2B-128a^5B+80a^4Ab+116a^3b^2B-80a^2Ab^3+17ab^4B-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 27

$$\int \frac{b(-32Ba^5+20Aba^4+44b^2Ba^3-35Ab^3a^2+8b^4Ba-5Ab^5)+(-128Ba^6+80Aba^5+212b^2Ba^4-140Ab^3a^3-55b^4Ba^2+40Ab^5a-9b^6B)\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx - \frac{2(-64a^5B+17ab^4B-80a^2Ab^3+116a^3b^2B-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$\int \frac{b(-32Ba^5+20Aba^4+44b^2Ba^3-35Ab^3a^2+8b^4Ba-5Ab^5)+(-128Ba^6+80Aba^5+212b^2Ba^4-140Ab^3a^3-55b^4Ba^2+40Ab^5a-9b^6B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-64a^5B+17ab^4B-80a^2Ab^3+116a^3b^2B-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3231

$$\frac{(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B) \int \sqrt{a+b\cos(c+dx)} dx}{b} - \frac{(a^2-b^2)(-128a^5B+80a^4Ab+116a^3b^2B-80a^2Ab^3+17ab^4B-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$\frac{(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B) \int \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}{b} - \frac{(a^2-b^2)(-128a^5B+80a^4Ab+116a^3b^2B-80a^2Ab^3+17ab^4B-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)}{5b}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3134

3.333. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}dx}{b\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

5b

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$\frac{(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}dx}{b\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

5b

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3132

$$\frac{2(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

5b

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3142

$$\frac{2(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

5b

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

3.333. $\int \frac{\cos^4(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{2(-128a^6B+80a^5Ab+212a^4b^2B-140a^3Ab^3-55a^2b^4B+40aAb^5-9b^6B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3140

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^3(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} +$$

$$\frac{2a(-8a^3B+5a^2Ab+12ab^2B-9Ab^3) \sin(c+dx) \cos^2(c+dx)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} + \frac{2(-48a^4B+30a^3Ab+71a^2b^2B-50aAb^3-3b^4B) \sin(c+dx) \cos(c+dx)\sqrt{a+b\cos(c+dx)}}{5bd}$$

input `Int[(Cos[c + d*x]^4*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Cos[c + d*x]^3*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + ((2*a*(5*a^2*A*b - 9*A*b^3 - 8*a^3*B + 12*a*b^2*B)*Cos[c + d*x]^2*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])) + ((-2*(30*a^3*A*b - 50*a*A*b^3 - 48*a^4*B + 71*a^2*b^2*B - 3*b^4*B)*Cos[c + d*x]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*b*d) - (((2*(80*a^5*A*b - 140*a^3*A*b^3 + 40*a*A*b^5 - 128*a^6*B + 212*a^4*b^2*B - 55*a^2*b^4*B - 9*b^6*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(80*a^4*A*b - 80*a^2*A*b^3 - 5*A*b^5 - 128*a^5*B + 116*a^3*b^2*B + 17*a*b^4*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/b - (2*(40*a^4*A*b - 65*a^2*A*b^3 + 5*A*b^5 - 64*a^5*B + 98*a^3*b^2*B - 14*a*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d))/(5*b))/(b*(a^2 - b^2)))/(3*b*(a^2 - b^2))`

3.333.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2)), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(a + b*Sine[e + f*x
])^m*((c + d*Sine[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Sine[e + f*x])^(m - 1)*(c + d*Sine[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sine[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sine[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

3.333.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1749 vs. $2(580) = 1160$.

Time = 30.72 (sec) , antiderivative size = 1750, normalized size of antiderivative = 3.18

method	result	size
default	Expression too large to display	1750
parts	Expression too large to display	2985

```
input int(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(16*B/b^2*(
-1/10/b*cos(1/2*d*x+1/2*c)^3*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+
1/2*c)^2)^(1/2)-1/60/b^2*(-4*a+12*b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/
2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+1/60/b^2*(-4*a+12*b)*(a-b)*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*
sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2
*d*x+1/2*c),(-2*b/(a-b))^(1/2))-1/60*(4*a^2-15*a*b+27*b^2)/b^3*(a-b)*(sin(
1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*s
in(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2
*d*x+1/2*c),(-2*b/(a-b))^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(
1/2))))+8/b^3*(A*b-2*B*a-3*B*b)*(-1/6/b*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*
x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+1/6*(a-b)/b*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x
+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))-1/12/b^2*(-2*a+6*b)*(a-b)*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+
(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b
))^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))))+2*a^4*(A*b-B*
a)/b^5*(1/6/b/(a-b)/(a+b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a
+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2+1/2*(a-b)/b)^2+8/...

```

3.333.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.35 (sec) , antiderivative size = 1538, normalized size of antiderivative = 2.80

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output

```
-1/45*(6*(64*B*a^7*b^2 - 40*A*a^6*b^3 - 98*B*a^5*b^4 + 65*A*a^4*b^5 + 14*B
*a^3*b^6 - 5*A*a^2*b^7 - 3*(B*a^4*b^5 - 2*B*a^2*b^7 + B*b^9)*cos(d*x + c)^
3 + (8*B*a^5*b^4 - 5*A*a^4*b^5 - 16*B*a^3*b^6 + 10*A*a^2*b^7 + 8*B*a*b^8 -
5*A*b^9)*cos(d*x + c)^2 + 5*(16*B*a^6*b^3 - 10*A*a^5*b^4 - 25*B*a^4*b^5 +
16*A*a^3*b^6 + 5*B*a^2*b^7 - 2*A*a*b^8)*cos(d*x + c))*sqrt(b*cos(d*x + c)
+ a)*sin(d*x + c) - (sqrt(2)*(256*I*B*a^7*b^2 - 160*I*A*a^6*b^3 - 520*I*B
*a^5*b^4 + 340*I*A*a^4*b^5 + 242*I*B*a^3*b^6 - 185*I*A*a^2*b^7 + 42*I*B*a*
b^8 - 15*I*A*b^9)*cos(d*x + c)^2 - 2*sqrt(2)*(-256*I*B*a^8*b + 160*I*A*a^7
*b^2 + 520*I*B*a^6*b^3 - 340*I*A*a^5*b^4 - 242*I*B*a^4*b^5 + 185*I*A*a^3*b
^6 - 42*I*B*a^2*b^7 + 15*I*A*a*b^8)*cos(d*x + c) + sqrt(2)*(256*I*B*a^9 -
160*I*A*a^8*b - 520*I*B*a^7*b^2 + 340*I*A*a^6*b^3 + 242*I*B*a^5*b^4 - 185*
I*A*a^4*b^5 + 42*I*B*a^3*b^6 - 15*I*A*a^2*b^7))*sqrt(b)*weierstrassPInvers
e(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x +
c) + 3*I*b*sin(d*x + c) + 2*a)/b) - (sqrt(2)*(-256*I*B*a^7*b^2 + 160*I*A
a^6*b^3 + 520*I*B*a^5*b^4 - 340*I*A*a^4*b^5 - 242*I*B*a^3*b^6 + 185*I*A*a^
2*b^7 - 42*I*B*a*b^8 + 15*I*A*b^9)*cos(d*x + c)^2 - 2*sqrt(2)*(256*I*B*a^8
*b - 160*I*A*a^7*b^2 - 520*I*B*a^6*b^3 + 340*I*A*a^5*b^4 + 242*I*B*a^4*b^5
- 185*I*A*a^3*b^6 + 42*I*B*a^2*b^7 - 15*I*A*a*b^8)*cos(d*x + c) + sqrt(2)
*(-256*I*B*a^9 + 160*I*A*a^8*b + 520*I*B*a^7*b^2 - 340*I*A*a^6*b^3 - 242*I
*B*a^5*b^4 + 185*I*A*a^4*b^5 - 42*I*B*a^3*b^6 + 15*I*A*a^2*b^7))*sqrt(b...
```

3.333.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.333.7 Maxima [F]

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^4}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^4/(b*cos(d*x + c) + a)^(5/2), x)`

3.333.8 Giac [F]

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^4}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^4*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^4/(b*cos(d*x + c) + a)^(5/2), x)`

3.333.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^4(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^4*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2), x)`

3.334
$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$$

3.334.1 Optimal result 3122
 3.334.2 Mathematica [A] (verified) 3123
 3.334.3 Rubi [A] (verified) 3123
 3.334.4 Maple [B] (verified) 3129
 3.334.5 Fricas [C] (verification not implemented) 3130
 3.334.6 Sympy [F(-1)] 3131
 3.334.7 Maxima [F] 3132
 3.334.8 Giac [F] 3132
 3.334.9 Mupad [F(-1)] 3132

3.334.1 Optimal result

Integrand size = 33, antiderivative size = 413

$$\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx = \frac{2(8a^4Ab - 15a^2Ab^3 + 3Ab^5 - 16a^5B + 28a^3b^2B - 8ab^4B) \sqrt{a+b \cos(c+dx)}}{3b^4(a^2-b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(8a^3Ab - 9aAb^3 - 16a^4B + 16a^2b^2B + b^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3b^4(a^2-b^2) d \sqrt{a+b \cos(c+dx)}} + \frac{2a(Ab - aB) \cos^2(c+dx) \sin(c+dx)}{3b(a^2-b^2) d(a+b \cos(c+dx))^{3/2}} - \frac{2a^2(3a^2Ab - 7Ab^3 - 6a^3B + 10ab^2B) \sin(c+dx)}{3b^3(a^2-b^2)^2 d \sqrt{a+b \cos(c+dx)}} - \frac{2(aAb - 2a^2B + b^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3b^3(a^2-b^2) d}$$

output

```
2/3*a*(A*b-B*a)*cos(d*x+c)^2*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)-2/3*a^2*(3*A*a^2*b-7*A*b^3-6*B*a^3+10*B*a*b^2)*sin(d*x+c)/b^3/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)-2/3*(A*a*b-2*B*a^2+B*b^2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/b^3/(a^2-b^2)/d+2/3*(8*A*a^4*b-15*A*a^2*b^3+3*A*b^5-16*B*a^5+28*B*a^3*b^2-8*B*a*b^4)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^4/(a^2-b^2)^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/3*(8*A*a^3*b-9*A*a*b^3-16*B*a^4+16*B*a^2*b^2+B*b^4)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b^4/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)
```

3.334.2 Mathematica [A] (verified)

Time = 2.48 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.81

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \frac{2 \left(\left(\frac{a+b\cos(c+dx)}{a+b} \right)^{3/2} (b^2(2a^3Ab-6aAb^3-4a^4B+7a^2b^2B+b^4B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) \right)}{\dots}$$

input `Integrate[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*(((a + b*Cos[c + d*x])/(a + b))^(3/2)*(b^2*(2*a^3*A*b - 6*a*A*b^3 - 4*a^4*B + 7*a^2*b^2*B + b^4*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)] - (-8*a^4*A*b + 15*a^2*A*b^3 - 3*A*b^5 + 16*a^5*B - 28*a^3*b^2*B + 8*a*b^4*B)*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)])))/((a - b)^2*(a + b)) + (b*(-8*a^5*A*b + 16*a^3*A*b^3 + 16*a^6*B - 25*a^4*b^2*B + b^6*B + 2*a*b*(-5*a^3*A*b + 9*a*A*b^3 + 10*a^4*B - 16*a^2*b^2*B + 2*b^4*B)*Cos[c + d*x] + (-a^2*b + b^3)^2*B*Cos[2*(c + d*x)]*Sin[c + d*x])/(2*(a^2 - b^2)^2))/((3*b^4*d*(a + b*Cos[c + d*x])^(3/2))`

3.334.3 Rubi [A] (verified)

Time = 2.23 (sec) , antiderivative size = 421, normalized size of antiderivative = 1.02, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3468, 27, 3042, 3510, 27, 3042, 3502, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^3(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 3468

$$\begin{aligned}
& \frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \\
& \frac{2 \int -\frac{\cos(c+dx)(-3(-2Ba^2 + Aba + b^2B) \cos^2(c+dx) - 3b(Ab - aB) \cos(c+dx) + 4a(Ab - aB))}{2(a+b \cos(c+dx))^{3/2}} dx}{3b(a^2 - b^2)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\cos(c+dx)(-3(-2Ba^2 + Aba + b^2B) \cos^2(c+dx) - 3b(Ab - aB) \cos(c+dx) + 4a(Ab - aB))}{(a+b \cos(c+dx))^{3/2}} dx}{3b(a^2 - b^2)} + \\
& \frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sin(c+dx+\frac{\pi}{2})(-3(-2Ba^2 + Aba + b^2B) \sin(c+dx+\frac{\pi}{2})^2 - 3b(Ab - aB) \sin(c+dx+\frac{\pi}{2}) + 4a(Ab - aB))}{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3b(a^2 - b^2)} + \\
& \frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
& \quad \downarrow 3510 \\
& \frac{2 \int \frac{-3b(a^2 - b^2)(-2Ba^2 + Aba + b^2B) \cos^2(c+dx) + (-12Ba^5 + 6Aba^4 + 22b^2Ba^3 - 13Ab^3a^2 - 6b^4Ba + 3Ab^5) \cos(c+dx) + ab(-6Ba^3 + 3Aba^2 + 10b^2Ba - 7Ab^3)}{2\sqrt{a+b \cos(c+dx)} b^2(a^2 - b^2)} dx}{3b(a^2 - b^2)} \\
& \quad \downarrow 27 \\
& \frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{-3b(a^2 - b^2)(-2Ba^2 + Aba + b^2B) \sin(c+dx+\frac{\pi}{2})^2 + (-12Ba^5 + 6Aba^4 + 22b^2Ba^3 - 13Ab^3a^2 - 6b^4Ba + 3Ab^5) \sin(c+dx+\frac{\pi}{2}) + ab(-6Ba^3 + 3Aba^2 + 10b^2Ba - 7Ab^3)}{\sqrt{a+b \cos(c+dx)} b^2(a^2 - b^2)} dx}{3b(a^2 - b^2)} \\
& \quad \downarrow 3502 \\
& \frac{2a(Ab - aB) \sin(c + dx) \cos^2(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
& \quad \downarrow 3502 \\
& \frac{\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx}{3b(a^2 - b^2)}
\end{aligned}$$

3.334. $\int \frac{\cos^3(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{2 \int \frac{3((-4Ba^4+2Aba^3+7b^2Ba^2-6Ab^3a+b^4B)b^2+(-16Ba^5+8Aba^4+28b^2Ba^3-15Ab^3a^2-8b^4Ba+3Ab^5)\cos(c+dx)b}{2\sqrt{a+b\cos(c+dx)}} dx}{3b} - \frac{2(a^2-b^2)(-2a^2B+aAb+b^2B)\sin(c+dx)}{d}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3b(a^2-b^2)$

↓ 27

$$\int \frac{(-4Ba^4+2Aba^3+7b^2Ba^2-6Ab^3a+b^4B)b^2+(-16Ba^5+8Aba^4+28b^2Ba^3-15Ab^3a^2-8b^4Ba+3Ab^5)\cos(c+dx)b}{\sqrt{a+b\cos(c+dx)}} dx - \frac{2(a^2-b^2)(-2a^2B+aAb+b^2B)\sin(c+dx)}{d}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3b(a^2-b^2)$

↓ 3042

$$\int \frac{(-4Ba^4+2Aba^3+7b^2Ba^2-6Ab^3a+b^4B)b^2+(-16Ba^5+8Aba^4+28b^2Ba^3-15Ab^3a^2-8b^4Ba+3Ab^5)\sin(c+dx+\frac{\pi}{2})b}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a^2-b^2)(-2a^2B+aAb+b^2B)\sin(c+dx)}{d}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3b(a^2-b^2)$

↓ 3231

$$\frac{(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5) \int \sqrt{a+b\cos(c+dx)} dx - (a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2(a^2-b^2)(-2a^2B+aAb+b^2B)\sin(c+dx)}{d}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3b(a^2-b^2)$

↓ 3042

$$\frac{(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5) \int \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} dx - (a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(a^2-b^2)(-2a^2B+aAb+b^2B)\sin(c+dx)}{d}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3b(a^2-b^2)$

↓ 3134

3.334. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\frac{(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}\frac{b}{b^2(a^2-b^2)}-(a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B)\int\frac{1}{\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}\frac{b}{b^2(a^2-b^2)}-(a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B)\int\frac{1}{\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3132

$$\frac{2(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}\frac{b}{b^2(a^2-b^2)}-(a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B)\int\frac{1}{\sqrt{a+b\sin(c+dx)}}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3142

$$\frac{2(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}\frac{b}{b^2(a^2-b^2)}-\frac{(a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

3.334. $\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \frac{2(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{(a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{\sqrt{a+b\cos(c+dx)}} \\
& \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3140} \\
& \frac{2a(Ab-aB)\sin(c+dx)\cos^2(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} + \\
& \frac{2(-16a^5B+8a^4Ab+28a^3b^2B-15a^2Ab^3-8ab^4B+3Ab^5)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}} - \frac{2(a^2-b^2)(-16a^4B+8a^3Ab+16a^2b^2B-9aAb^3+b^4B)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}{d\sqrt{a+b\cos(c+dx)}}
\end{aligned}$$

input `Int[(Cos[c + d*x]^3*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Cos[c + d*x]^2*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + ((-2*a^2*(3*a^2*A*b - 7*A*b^3 - 6*a^3*B + 10*a*b^2*B)*Sin[c + d*x])/(b^2*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + (((2*(8*a^4*A*b - 15*a^2*A*b^3 + 3*A*b^5 - 16*a^5*B + 28*a^3*b^2*B - 8*a*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(8*a^3*A*b - 9*a*A*b^3 - 16*a^4*B + 16*a^2*b^2*B + b^4*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b - (2*(a^2 - b^2)*(a*A*b - 2*a^2*B + b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/d)/(b^2*(a^2 - b^2)))/(3*b*(a^2 - b^2))`

3.334.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3510 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]*(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[
e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - S
imp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(
m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m
+ 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)
))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x] /; F
reeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2,
0] && LtQ[m, -1]
```

3.334.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1411 vs. $2(447) = 894$.

Time = 27.65 (sec) , antiderivative size = 1412, normalized size of antiderivative = 3.42

method	result	size
default	Expression too large to display	1412
parts	Expression too large to display	2208

```
input int(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNV
ERBOSE)
```


output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*a^2/b^4
*(3*A*b-4*B*a)/sin(1/2*d*x+1/2*c)^2/(2*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^
2)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*cos(1/2
*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b+EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b
))^1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*a-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2))*(-2*b/
(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^1/2*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*b)-2*a^3*(A*b-B*a)/b^4*(1/6/b/(a-b)/(a+b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*
d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2+1/2
*(a-b)/b)^2+8/3*sin(1/2*d*x+1/2*c)^2*b/(a-b)^2/(a+b)^2*cos(1/2*d*x+1/2*c)*
a/(-(-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+(3*a-b)/(3
*a^3+3*a^2*b-3*a*b^2-3*b^3)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x
+1/2*c)^2+a-b)/(a-b))^1/2/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1
/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2))-4/3*a/(a-b
)/(a+b)^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-
b))^1/2/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(El
lipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^1/2))-EllipticE(cos(1/2*d*x+1/2*c
),(-2*b/(a-b))^1/2))))-2/3/b^4/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-4*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*b^2+9*A*a
*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/...

```

3.334.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.23 (sec) , antiderivative size = 1348, normalized size of antiderivative = 3.26

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output

```

1/9*(6*(8*B*a^6*b^2 - 4*A*a^5*b^3 - 13*B*a^4*b^4 + 8*A*a^3*b^5 + B*a^2*b^6
+ (B*a^4*b^4 - 2*B*a^2*b^6 + B*b^8)*cos(d*x + c)^2 + (10*B*a^5*b^3 - 5*A*
a^4*b^4 - 16*B*a^3*b^5 + 9*A*a^2*b^6 + 2*B*a*b^7)*cos(d*x + c))*sqrt(b*cos
(d*x + c) + a)*sin(d*x + c) + (sqrt(2)*(-32*I*B*a^6*b^2 + 16*I*A*a^5*b^3 +
68*I*B*a^4*b^4 - 36*I*A*a^3*b^5 - 37*I*B*a^2*b^6 + 24*I*A*a*b^7 - 3*I*B*b
^8)*cos(d*x + c)^2 - 2*sqrt(2)*(32*I*B*a^7*b - 16*I*A*a^6*b^2 - 68*I*B*a^5
*b^3 + 36*I*A*a^4*b^4 + 37*I*B*a^3*b^5 - 24*I*A*a^2*b^6 + 3*I*B*a*b^7)*cos
(d*x + c) + sqrt(2)*(-32*I*B*a^8 + 16*I*A*a^7*b + 68*I*B*a^6*b^2 - 36*I*A*
a^5*b^3 - 37*I*B*a^4*b^4 + 24*I*A*a^3*b^5 - 3*I*B*a^2*b^6))*sqrt(b)*weiers
trassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3
*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (sqrt(2)*(32*I*B*a^6*b^2
- 16*I*A*a^5*b^3 - 68*I*B*a^4*b^4 + 36*I*A*a^3*b^5 + 37*I*B*a^2*b^6 - 24*I
*A*a*b^7 + 3*I*B*b^8)*cos(d*x + c)^2 - 2*sqrt(2)*(-32*I*B*a^7*b + 16*I*A*a
^6*b^2 + 68*I*B*a^5*b^3 - 36*I*A*a^4*b^4 - 37*I*B*a^3*b^5 + 24*I*A*a^2*b^6
- 3*I*B*a*b^7)*cos(d*x + c) + sqrt(2)*(32*I*B*a^8 - 16*I*A*a^7*b - 68*I*B
*a^6*b^2 + 36*I*A*a^5*b^3 + 37*I*B*a^4*b^4 - 24*I*A*a^3*b^5 + 3*I*B*a^2*b^
6))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*
a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(sqrt
(2)*(16*I*B*a^5*b^3 - 8*I*A*a^4*b^4 - 28*I*B*a^3*b^5 + 15*I*A*a^2*b^6 + 8*
I*B*a*b^7 - 3*I*A*b^8)*cos(d*x + c)^2 + 2*sqrt(2)*(16*I*B*a^6*b^2 - 8*I...

```

3.334.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.334.7 Maxima [F]

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^3}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^3/(b*cos(d*x + c) + a)^(5/2), x)`

3.334.8 Giac [F]

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^3}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^3*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^3/(b*cos(d*x + c) + a)^(5/2), x)`

3.334.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^3(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^3*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2), x)`

3.335
$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$$

3.335.1 Optimal result 3133
 3.335.2 Mathematica [A] (verified) 3134
 3.335.3 Rubi [A] (verified) 3134
 3.335.4 Maple [B] (verified) 3139
 3.335.5 Fricas [C] (verification not implemented) 3140
 3.335.6 Sympy [F(-1)] 3141
 3.335.7 Maxima [F] 3142
 3.335.8 Giac [F] 3142
 3.335.9 Mupad [F(-1)] 3142

3.335.1 Optimal result

Integrand size = 33, antiderivative size = 331

$$\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx =$$

$$\frac{2(2a^3Ab - 6aAb^3 - 8a^4B + 15a^2b^2B - 3b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) - 3b^3(a^2 - b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{3b^3(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(2a^2Ab - 3Ab^3 - 8a^3B + 9ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3b^3(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}}$$

$$- \frac{2a^2(Ab - aB) \sin(c+dx)}{3b^2(a^2 - b^2) d(a+b \cos(c+dx))^{3/2}} + \frac{2a(2a^2Ab - 6Ab^3 - 5a^3B + 9ab^2B) \sin(c+dx)}{3b^2(a^2 - b^2)^2 d \sqrt{a+b \cos(c+dx)}}$$

output

```
-2/3*a^2*(A*b-B*a)*sin(d*x+c)/b^2/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)+2/3*a
*(2*A*a^2*b-6*A*b^3-5*B*a^3+9*B*a*b^2)*sin(d*x+c)/b^2/(a^2-b^2)^2/d/(a+b*c
os(d*x+c))^(1/2)-2/3*(2*A*a^3*b-6*A*a*b^3-8*B*a^4+15*B*a^2*b^2-3*B*b^4)*(c
os(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c)
,2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b^3/(a^2-b^2)^2/d/((a+b*c
os(d*x+c))/(a+b))^(1/2)+2/3*(2*A*a^2*b-3*A*b^3-8*B*a^3+9*B*a*b^2)*(cos(1/2
*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/
2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b^3/(a^2-b^2)/d/(a+b*co
s(d*x+c))^(1/2)
```

3.335.2 Mathematica [A] (verified)

Time = 2.01 (sec) , antiderivative size = 274, normalized size of antiderivative = 0.83

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2 \left(\left(\frac{a+b \cos(c+dx)}{a+b} \right)^{3/2} \left(b^2 (a^2 Ab + 3Ab^3 + 2a^3 B - 6ab^2 B) \operatorname{EllipticF} \left(\frac{1}{2}(c+dx), \frac{2b}{a+b} \right) + (-2a^3 B + 6a^2 Ab + 3Ab^3 + 2a^3 B - 6ab^2 B) \operatorname{EllipticE} \left(\frac{1}{2}(c+dx), \frac{2b}{a+b} \right) \right)}{(a+b \cos(c+dx))^{5/2}} \right)}{(a+b \cos(c+dx))^{5/2}}$$

input `Integrate[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*(((a + b*Cos[c + d*x])/(a + b))^(3/2)*(b^2*(a^2*A*b + 3*A*b^3 + 2*a^3*B - 6*a*b^2*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)] + (-2*a^3*A*b + 6*a*A*b^3 + 8*a^4*B - 15*a^2*b^2*B + 3*b^4*B)*((a + b)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - a*EllipticF[(c + d*x)/2, (2*b)/(a + b)])))/((a - b)^2*(a + b)) - (a*b*(a*(-a^2*A*b) + 5*A*b^3 + 4*a^3*B - 8*a*b^2*B) + b*(-2*a^2*A*b + 6*A*b^3 + 5*a^3*B - 9*a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/(a^2 - b^2))/((3*b^3*d*(a + b*Cos[c + d*x]))^(3/2))`

3.335.3 Rubi [A] (verified)

Time = 1.62 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3467, 27, 3042, 3500, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c + dx + \frac{\pi}{2})^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow \text{3467} \\ & \frac{2 \int \frac{3b(a^2 - b^2)B \cos^2(c + dx) + (2a^2 - 3b^2)(Ab - aB) \cos(c + dx) + 3ab(Ab - aB)}{2(a + b \cos(c + dx))^{3/2}} dx}{3b^2(a^2 - b^2)} - \\ & \quad \frac{2a^2(Ab - aB) \sin(c + dx)}{3b^2 d (a^2 - b^2) (a + b \cos(c + dx))^{3/2}} \end{aligned}$$

3.335. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{array}{c}
\downarrow 27 \\
\frac{\int \frac{3b(a^2-b^2)B \cos^2(c+dx) + (2a^2-3b^2)(Ab-aB) \cos(c+dx) + 3ab(Ab-aB)}{(a+b \cos(c+dx))^{3/2}} dx}{\frac{3b^2(a^2-b^2)}{2a^2(Ab-aB) \sin(c+dx)}} \\
\frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
\downarrow 3042 \\
\frac{\int \frac{3b(a^2-b^2)B \sin(c+dx+\frac{\pi}{2})^2 + (2a^2-3b^2)(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + 3ab(Ab-aB)}{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{\frac{3b^2(a^2-b^2)}{2a^2(Ab-aB) \sin(c+dx)}} \\
\frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
\downarrow 3500 \\
\frac{2a(-5a^3B+2a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} - \frac{2 \int \frac{b^2(2Ba^3+Ab^2-6b^2Ba+3Ab^3) - b(-8Ba^4+2Aba^3+15b^2Ba^2-6Ab^3a-3b^4B) \cos(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{b(a^2-b^2)} \\
\frac{3b^2(a^2-b^2)}{2a^2(Ab-aB) \sin(c+dx)} \\
\frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
\downarrow 27 \\
\frac{\int \frac{b^2(2Ba^3+Ab^2-6b^2Ba+3Ab^3) - b(-8Ba^4+2Aba^3+15b^2Ba^2-6Ab^3a-3b^4B) \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b(a^2-b^2)} + \frac{2a(-5a^3B+2a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} \\
\frac{3b^2(a^2-b^2)}{2a^2(Ab-aB) \sin(c+dx)} \\
\frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
\downarrow 3042 \\
\frac{\int \frac{b^2(2Ba^3+Ab^2-6b^2Ba+3Ab^3) - b(-8Ba^4+2Aba^3+15b^2Ba^2-6Ab^3a-3b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b(a^2-b^2)} + \frac{2a(-5a^3B+2a^2Ab+9ab^2B-6Ab^3) \sin(c+dx)}{d(a^2-b^2)\sqrt{a+b \cos(c+dx)}} \\
\frac{3b^2(a^2-b^2)}{2a^2(Ab-aB) \sin(c+dx)} \\
\frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2 d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
\downarrow 3231
\end{array}$$

3.335. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{(a^2-b^2)(-8a^3B+2a^2Ab+9ab^2B-3Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx - (-8a^4B+2a^3Ab+15a^2b^2B-6aAb^3-3b^4B) \int \sqrt{a+b \cos(c+dx)} dx}{b(a^2-b^2)} + \frac{2a(-5a^3B+9a^2bB-3ab^2B-3b^3B)}{d(a^2-b^2)}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(a^2-b^2)(-8a^3B+2a^2Ab+9ab^2B-3Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - (-8a^4B+2a^3Ab+15a^2b^2B-6aAb^3-3b^4B) \int \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b(a^2-b^2)} + \frac{2a(-5a^3B+9a^2bB-3ab^2B-3b^3B)}{d(a^2-b^2)}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3134

$$\frac{(a^2-b^2)(-8a^3B+2a^2Ab+9ab^2B-3Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-8a^4B+2a^3Ab+15a^2b^2B-6aAb^3-3b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{b(a^2-b^2)}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(a^2-b^2)(-8a^3B+2a^2Ab+9ab^2B-3Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{(-8a^4B+2a^3Ab+15a^2b^2B-6aAb^3-3b^4B) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{b(a^2-b^2)}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3132

$$\frac{(a^2-b^2)(-8a^3B+2a^2Ab+9ab^2B-3Ab^3) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(-8a^4B+2a^3Ab+15a^2b^2B-6aAb^3-3b^4B) \sqrt{a+b \cos(c+dx)} E(\frac{1}{2}(c+dx) | \frac{2b}{a+b})}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{b(a^2-b^2)} + \frac{2(-8a^4B+2a^3Ab+15a^2b^2B-6aAb^3-3b^4B) \sqrt{a+b \cos(c+dx)}}{d(a^2-b^2)}$$

$$\frac{3b^2(a^2-b^2)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \frac{2a^2(Ab-aB) \sin(c+dx)}{3b^2d(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3142

3.335. $\int \frac{\cos^2(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{(a^2 - b^2)(-8a^3B + 2a^2Ab + 9ab^2B - 3Ab^3) \int \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{\sqrt{a+b \cos(c+dx)}} \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{b(a^2 - b^2)} - \frac{2(-8a^4B + 2a^3Ab + 15a^2b^2B - 6aAb^3 - 3b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{2a^2(Ab - aB) \sin(c + dx)}{3b^2d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$\frac{(a^2 - b^2)(-8a^3B + 2a^2Ab + 9ab^2B - 3Ab^3) \int \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{\sqrt{a+b \cos(c+dx)}} \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin\left(c+dx + \frac{\pi}{2}\right)}{a+b}}} dx}{b(a^2 - b^2)} - \frac{2(-8a^4B + 2a^3Ab + 15a^2b^2B - 6aAb^3 - 3b^4B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{2a^2(Ab - aB) \sin(c + dx)}{3b^2d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3140

$$\frac{2a(-5a^3B + 2a^2Ab + 9ab^2B - 6Ab^3) \sin(c+dx)}{d(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} + \frac{2(a^2 - b^2)(-8a^3B + 2a^2Ab + 9ab^2B - 3Ab^3) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2(-8a^4B + 2a^3Ab + 15a^2b^2B - 6aAb^3 - 3b^4B) \sqrt{a+b \cos(c+dx)}}{b(a^2 - b^2)}$$

$$\frac{2a^2(Ab - aB) \sin(c + dx)}{3b^2d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

input `Int[(Cos[c + d*x]^2*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(-2*a^2*(A*b - a*B)*Sin[c + d*x])/(3*b^2*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((-2*(2*a^3*A*b - 6*a*A*b^3 - 8*a^4*B + 15*a^2*b^2*B - 3*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]/(a + b))) + (2*(a^2 - b^2)*(2*a^2*A*b - 3*A*b^3 - 8*a^3*B + 9*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*Cos[c + d*x]]))/(b*(a^2 - b^2)) + (2*a*(2*a^2*A*b - 6*A*b^3 - 5*a^3*B + 9*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])/(3*b^2*(a^2 - b^2))`

3.335.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*SIN[c + d*x]]/Sqrt[(a + b*SIN[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*SIN[c + d*x])/(a + b)]/Sqrt[a + b*SIN[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*SIN[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*SIN[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

```
rule 3467 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.335.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 953 vs. $2(369) = 738$.

Time = 24.43 (sec) , antiderivative size = 954, normalized size of antiderivative = 2.88

method	result	size
default	Expression too large to display	954
parts	Expression too large to display	1763

```
input int(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNV
ERBOSE)
```

```

output -((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2/b^3/(-2*
sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(A*b*Ellip
ticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-3*B*EllipticF(cos(1/2*d*x+1/2*
c),(-2*b/(a-b))^(1/2))*a+B*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)
)*a-B*EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*b)+2*a^2*(A*b-B*a)/
b^3*(1/6/b/(a-b)/(a+b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)
*sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2+1/2*(a-b)/b)^2+8/3*sin(
1/2*d*x+1/2*c)^2*b/(a-b)^2/(a+b)^2*cos(1/2*d*x+1/2*c)*a/(-(-2*b*cos(1/2*d*
x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+(3*a-b)/(3*a^3+3*a^2*b-3*a*b^2
-3*b^3)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b)
)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellip
ticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-4/3*a/(a-b)/(a+b)^2*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+
1/2*c),(-2*b/(a-b))^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2)
)))+2*a/b^3*(2*A*b-3*B*a)/sin(1/2*d*x+1/2*c)^2/(2*b*sin(1/2*d*x+1/2*c)^2-a
-b)/(a^2-b^2)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)
*(2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b+EllipticE(cos(1/2*d*x+1/2*c)
,(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)...

```

3.335.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.21 (sec) , antiderivative size = 1193, normalized size of antiderivative = 3.60

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

```

input integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorith
m="fricas")

```

output

```
-1/9*(6*(4*B*a^5*b^2 - A*a^4*b^3 - 8*B*a^3*b^4 + 5*A*a^2*b^5 + (5*B*a^4*b^3 - 2*A*a^3*b^4 - 9*B*a^2*b^5 + 6*A*a*b^6)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c) - (sqrt(2)*(16*I*B*a^5*b^2 - 4*I*A*a^4*b^3 - 36*I*B*a^3*b^4 + 9*I*A*a^2*b^5 + 24*I*B*a*b^6 - 9*I*A*b^7)*cos(d*x + c)^2 - 2*sqrt(2)*(-16*I*B*a^6*b + 4*I*A*a^5*b^2 + 36*I*B*a^4*b^3 - 9*I*A*a^3*b^4 - 24*I*B*a^2*b^5 + 9*I*A*a*b^6)*cos(d*x + c) + sqrt(2)*(16*I*B*a^7 - 4*I*A*a^6*b - 36*I*B*a^5*b^2 + 9*I*A*a^4*b^3 + 24*I*B*a^3*b^4 - 9*I*A*a^2*b^5))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) - (sqrt(2)*(-16*I*B*a^5*b^2 + 4*I*A*a^4*b^3 + 36*I*B*a^3*b^4 - 9*I*A*a^2*b^5 - 24*I*B*a*b^6 + 9*I*A*b^7)*cos(d*x + c)^2 - 2*sqrt(2)*(16*I*B*a^6*b - 4*I*A*a^5*b^2 - 36*I*B*a^4*b^3 + 9*I*A*a^3*b^4 + 24*I*B*a^2*b^5 - 9*I*A*a*b^6)*cos(d*x + c) + sqrt(2)*(-16*I*B*a^7 + 4*I*A*a^6*b + 36*I*B*a^5*b^2 - 9*I*A*a^4*b^3 - 24*I*B*a^3*b^4 + 9*I*A*a^2*b^5))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) + 3*(sqrt(2)*(-8*I*B*a^4*b^3 + 2*I*A*a^3*b^4 + 15*I*B*a^2*b^5 - 6*I*A*a*b^6 - 3*I*B*b^7)*cos(d*x + c)^2 + 2*sqrt(2)*(-8*I*B*a^5*b^2 + 2*I*A*a^4*b^3 + 15*I*B*a^3*b^4 - 6*I*A*a^2*b^5 - 3*I*B*a*b^6)*cos(d*x + c) + sqrt(2)*(-8*I*B*a^6*b + 2*I*A*a^5*b^2 + 15*I*B*a^4*b^3 - 6*I*A*a^3*b^4 - 3*I*B*a^2*b^5))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, ...
```

3.335.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.335.7 Maxima [F]

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^2}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^2/(b*cos(d*x + c) + a)^(5/2), x)`

3.335.8 Giac [F]

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^2}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^2*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^2/(b*cos(d*x + c) + a)^(5/2), x)`

3.335.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^2(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^2*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2), x)`

3.336 $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

3.336.1 Optimal result 3143
 3.336.2 Mathematica [A] (verified) 3144
 3.336.3 Rubi [A] (verified) 3144
 3.336.4 Maple [B] (verified) 3149
 3.336.5 Fricas [C] (verification not implemented) 3150
 3.336.6 Sympy [F(-1)] 3151
 3.336.7 Maxima [F] 3152
 3.336.8 Giac [F] 3152
 3.336.9 Mupad [F(-1)] 3152

3.336.1 Optimal result

Integrand size = 31, antiderivative size = 307

$$\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx =$$

$$\frac{2(a^2Ab + 3Ab^3 + 2a^3B - 6ab^2B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{3b^2(a^2 - b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(aAb + 2a^2B - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3b^2(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}}$$

$$+ \frac{2a(Ab - aB) \sin(c+dx)}{3b(a^2 - b^2) d(a+b \cos(c+dx))^{3/2}} + \frac{2(a^2Ab + 3Ab^3 + 2a^3B - 6ab^2B) \sin(c+dx)}{3b(a^2 - b^2)^2 d \sqrt{a+b \cos(c+dx)}}$$

```
output 2/3*a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)+2/3*(A*a^2
*b+3*A*b^3+2*B*a^3-6*B*a*b^2)*sin(d*x+c)/b/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(
(1/2)-2/3*(A*a^2*b+3*A*b^3+2*B*a^3-6*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*
(a+b*cos(d*x+c))^(1/2)/b^2/(a^2-b^2)^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2/
3*(A*a*b+2*B*a^2-3*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a
+b))^(1/2)/b^2/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)
```

3.336.2 Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.73

$$\int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2 \left(-\frac{\left(\frac{a+b \cos(c+dx)}{a+b}\right)^{3/2} \left((a^2 A b + 3 A b^3 + 2 a^3 B - 6 a b^2 B) E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right) - (a-b)(a A b + \dots) \right)}{(a-b)^2} \right)}{\dots}$$

input `Integrate[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*(-(((a + b*Cos[c + d*x])/(a + b))^(3/2)*((a^2*A*b + 3*A*b^3 + 2*a^3*B - 6*a*b^2*B)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - (a - b)*(a*A*b + 2*a^2*B - 3*b^2*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)])))/(a - b)^2) + (b*(a*(2*a^2*A*b + 2*A*b^3 + a^3*B - 5*a*b^2*B) + b*(a^2*A*b + 3*A*b^3 + 2*a^3*B - 6*a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/(a^2 - b^2)^2))/(3*b^2*d*(a + b*Cos[c + d*x])^(3/2))`

3.336.3 Rubi [A] (verified)

Time = 1.53 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.03, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3233, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sin(c + dx + \frac{\pi}{2})(A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{A \cos(c + dx) + B \cos^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.336. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \int \frac{A \sin \left(c+dx+\frac{\pi}{2}\right)+B \sin \left(c+dx+\frac{\pi}{2}\right)^2}{\left(a+b \sin \left(c+dx+\frac{\pi}{2}\right)\right)^{5 / 2}} dx \\
 & \quad \downarrow \text{3500} \\
 & \frac{2 a(A b-a B) \sin (c+d x)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+d x)\right)^{3 / 2}}-\frac{2 \int \frac{3 b(A b-a B)-(2 B a^2+A b a-3 b^2 B) \cos (c+d x)}{2\left(a+b \cos (c+d x)\right)^{3 / 2}} dx}{3 b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 a(A b-a B) \sin (c+d x)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+d x)\right)^{3 / 2}}-\frac{\int \frac{3 b(A b-a B)-(2 B a^2+A b a-3 b^2 B) \cos (c+d x)}{\left(a+b \cos (c+d x)\right)^{3 / 2}} dx}{3 b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 a(A b-a B) \sin (c+d x)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+d x)\right)^{3 / 2}}-\frac{\int \frac{3 b(A b-a B)+\left(-2 B a^2-A b a+3 b^2 B\right) \sin \left(c+d x+\frac{\pi}{2}\right)}{\left(a+b \sin \left(c+d x+\frac{\pi}{2}\right)\right)^{3 / 2}} dx}{3 b\left(a^2-b^2\right)} \\
 & \quad \downarrow \text{3233} \\
 & \frac{2 a(A b-a B) \sin (c+d x)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+d x)\right)^{3 / 2}}-\frac{2 \int-\frac{b\left(-B a^2+4 A b a-3 b^2 B\right)+\left(2 B a^3+A b a^2-6 b^2 B a+3 A b^3\right) \cos (c+d x)}{2 \sqrt{a+b \cos (c+d x)}} dx}{\frac{2\left(2 a^3 B+a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+d x)}{d\left(a^2-b^2\right) \sqrt{a+b \cos (c+d x)}}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 a(A b-a B) \sin (c+d x)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+d x)\right)^{3 / 2}}-\frac{\int \frac{b\left(-B a^2+4 A b a-3 b^2 B\right)+\left(2 B a^3+A b a^2-6 b^2 B a+3 A b^3\right) \cos (c+d x)}{\sqrt{a+b \cos (c+d x)}} dx}{\frac{2\left(2 a^3 B+a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+d x)}{d\left(a^2-b^2\right) \sqrt{a+b \cos (c+d x)}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 a(A b-a B) \sin (c+d x)}{3 b d\left(a^2-b^2\right)\left(a+b \cos (c+d x)\right)^{3 / 2}}-\frac{\int \frac{b\left(-B a^2+4 A b a-3 b^2 B\right)+\left(2 B a^3+A b a^2-6 b^2 B a+3 A b^3\right) \sin \left(c+d x+\frac{\pi}{2}\right)}{\sqrt{a+b \sin \left(c+d x+\frac{\pi}{2}\right)}} dx}{\frac{2\left(2 a^3 B+a^2 A b-6 a b^2 B+3 A b^3\right) \sin (c+d x)}{d\left(a^2-b^2\right) \sqrt{a+b \cos (c+d x)}}} \\
 & \quad \downarrow \text{3231}
 \end{aligned}$$

3.336. $\int \frac{\cos (c+d x)(A+B \cos (c+d x))}{\left(a+b \cos (c+d x)\right)^{5 / 2}} dx$

$$\frac{\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \int \sqrt{a+b \cos(c+dx)} dx}{b(a^2 - b^2)(2a^2B + aAb - 3b^2B) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}}{a^2 - b^2} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{d(a^2 - b^2)\sqrt{a+b \cos(c+dx)}}$$

$3b(a^2 - b^2)$

↓ 3042

$$\frac{\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \int \sqrt{a+b \sin(c+dx + \frac{\pi}{2})} dx}{b(a^2 - b^2)(2a^2B + aAb - 3b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}}{a^2 - b^2} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{d(a^2 - b^2)\sqrt{a+b \cos(c+dx)}}$$

$3b(a^2 - b^2)$

↓ 3134

$$\frac{\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} (a^2 - b^2)(2a^2B + aAb - 3b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}}{a^2 - b^2} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{d(a^2 - b^2)\sqrt{a+b \cos(c+dx)}}$$

$3b(a^2 - b^2)$

↓ 3042

$$\frac{\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}} dx}{b \sqrt{\frac{a+b \cos(c+dx)}{a+b}} (a^2 - b^2)(2a^2B + aAb - 3b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}}{a^2 - b^2} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{d(a^2 - b^2)\sqrt{a+b \cos(c+dx)}}$$

$3b(a^2 - b^2)$

↓ 3132

$$\frac{\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}} (a^2 - b^2)(2a^2B + aAb - 3b^2B) \int \frac{1}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}}{a^2 - b^2} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sin(c + dx)}{d(a^2 - b^2)\sqrt{a+b \cos(c+dx)}}$$

$3b(a^2 - b^2)$

↓ 3142

3.336. $\int \frac{\cos(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{(a^2 - b^2)(2a^2B + aAb - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2)(2a^2B + aAb - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3)}{d(a^2 - b^2)}$$

$$3b(a^2 - b^2)$$

↓ 3042

$$\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{(a^2 - b^2)(2a^2B + aAb - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin\left(c+dx + \frac{\pi}{2}\right)}{a+b}}} dx}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(a^2 - b^2)(2a^2B + aAb - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{b \sqrt{a+b \cos(c+dx)}} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3)}{d(a^2 - b^2)}$$

$$3b(a^2 - b^2)$$

↓ 3140

$$\frac{2a(Ab - aB) \sin(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \frac{2(a^2 - b^2)(2a^2B + aAb - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(a^2 - b^2)(2a^2B + aAb - 3b^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(2a^3B + a^2Ab - 6ab^2B + 3Ab^3)}{d(a^2 - b^2)}$$

$$3b(a^2 - b^2)$$

input `Int[(Cos[c + d*x]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) - (((2*(a^2*A*b + 3*A*b^3 + 2*a^3*B - 6*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(a*A*b + 2*a^2*B - 3*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(a^2 - b^2) - (2*(a^2*A*b + 3*A*b^3 + 2*a^3*B - 6*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])/(3*b*(a^2 - b^2))`

3.336.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

```
rule 3233 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

```
rule 3447 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3500 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2
- a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*
(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A
*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A,
B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.336.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 863 vs. $2(345) = 690$.

Time = 21.46 (sec) , antiderivative size = 864, normalized size of antiderivative = 2.81

method	result	size
default	Expression too large to display	864
parts	Expression too large to display	1598

```
input int(cos(d*x+c)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVER
BOSE)
```

output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B/b^2*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-2/b^2*(A*b-2*B*a)/sin(1/2*d*x+1/2*c)^2/(
2*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*s
in(1/2*d*x+1/2*c)^2)^(1/2)*(2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2+b*El
lipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2
*c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*a-EllipticE(cos(1/2*
d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b
))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*b)-2*a*(A*b-B*a)/b^2*(1/6/b/(a-b)/(a
+b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)
^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2+1/2*(a-b)/b)^2+8/3*sin(1/2*d*x+1/2*c)^2*b/
(a-b)^2/(a+b)^2*cos(1/2*d*x+1/2*c)*a/(-(-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin
(1/2*d*x+1/2*c)^2)^(1/2)+(3*a-b)/(3*a^3+3*a^2*b-3*a*b^2-3*b^3)*(sin(1/2*d*
x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2
*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),(-2*b/(a-b))^(1/2))-4/3*a/(a-b)/(a+b)^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a
+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),(-2*b/(a-b))
^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))))/sin(1/2*d*x...

```

3.336.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.16 (sec) , antiderivative size = 1076, normalized size of antiderivative = 3.50

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/9*(6*(B*a^4*b^2 + 2*A*a^3*b^3 - 5*B*a^2*b^4 + 2*A*a*b^5 + (2*B*a^3*b^3 + A*a^2*b^4 - 6*B*a*b^5 + 3*A*b^6)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c) + (sqrt(2)*(-4*I*B*a^4*b^2 - 2*I*A*a^3*b^3 + 9*I*B*a^2*b^4 + 6*I*A*a*b^5 - 9*I*B*b^6)*cos(d*x + c)^2 - 2*sqrt(2)*(4*I*B*a^5*b + 2*I*A*a^4*b^2 - 9*I*B*a^3*b^3 - 6*I*A*a^2*b^4 + 9*I*B*a*b^5)*cos(d*x + c) + sqrt(2)*(-4*I*B*a^6 - 2*I*A*a^5*b + 9*I*B*a^4*b^2 + 6*I*A*a^3*b^3 - 9*I*B*a^2*b^4))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (sqrt(2)*(4*I*B*a^4*b^2 + 2*I*A*a^3*b^3 - 9*I*B*a^2*b^4 - 6*I*A*a*b^5 + 9*I*B*b^6)*cos(d*x + c)^2 - 2*sqrt(2)*(-4*I*B*a^5*b - 2*I*A*a^4*b^2 + 9*I*B*a^3*b^3 + 6*I*A*a^2*b^4 - 9*I*B*a*b^5)*cos(d*x + c) + sqrt(2)*(4*I*B*a^6 + 2*I*A*a^5*b - 9*I*B*a^4*b^2 - 6*I*A*a^3*b^3 + 9*I*B*a^2*b^4))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(sqrt(2)*(2*I*B*a^3*b^3 + I*A*a^2*b^4 - 6*I*B*a*b^5 + 3*I*A*b^6)*cos(d*x + c)^2 + 2*sqrt(2)*(2*I*B*a^4*b^2 + I*A*a^3*b^3 - 6*I*B*a^2*b^4 + 3*I*A*a*b^5)*cos(d*x + c) + sqrt(2)*(2*I*B*a^5*b + I*A*a^4*b^2 - 6*I*B*a^3*b^3 + 3*I*A*a^2*b^4))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*(sqrt(2)*(-2*I*B*a^3*b...`

3.336.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2), x)`

output `Timed out`

3.336.7 Maxima [F]

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)}{(b\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)/(b*cos(d*x + c) + a)^(5/2), x)`

3.336.8 Giac [F]

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)}{(b\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)/(b*cos(d*x + c) + a)^(5/2), x)`

3.336.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2), x)`

3.337 $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

3.337.1 Optimal result 3153
 3.337.2 Mathematica [A] (verified) 3154
 3.337.3 Rubi [A] (verified) 3154
 3.337.4 Maple [B] (verified) 3158
 3.337.5 Fracas [C] (verification not implemented) 3159
 3.337.6 Sympy [F(-1)] 3160
 3.337.7 Maxima [F] 3161
 3.337.8 Giac [F] 3161
 3.337.9 Mupad [F(-1)] 3161

3.337.1 Optimal result

Integrand size = 25, antiderivative size = 275

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2(4aAb - a^2B - 3b^2B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{3b(a^2 - b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2(Ab - aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3b(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}} - \frac{2(Ab - aB) \sin(c + dx)}{3(a^2 - b^2) d (a + b \cos(c + dx))^{3/2}} - \frac{2(4aAb - a^2B - 3b^2B) \sin(c + dx)}{3(a^2 - b^2)^2 d \sqrt{a + b \cos(c + dx)}}$$

output

```
-2/3*(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)-2/3*(4*A*a*b-
B*a^2-3*B*b^2)*sin(d*x+c)/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)+2/3*(4*A*a*
b-B*a^2-3*B*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE
(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/b/(a^2
-b^2)^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-2/3*(A*b-B*a)*(cos(1/2*d*x+1/2*c)
^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b)
)^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/
2)
```


3.337.2 Mathematica [A] (verified)

Time = 1.16 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.70

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2 \left(-\left(\frac{a+b \cos(c+dx)}{a+b}\right)^{3/2} \left((-4aAb+a^2B+3b^2B) E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right) - (a-b)(-Ab+aB) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx) \right) \right)}{(a-b)^2 b} \right)}{3d(a + b \cos(c + dx))^{3/2}}$$

input `Integrate[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(5/2), x]`

output `(2*(-(((a + b*Cos[c + d*x])/(a + b))^(3/2)*((-4*a*A*b + a^2*B + 3*b^2*B)*EllipticE[(c + d*x)/2, (2*b)/(a + b)] - (a - b)*(-A*b) + a*B)*EllipticF[(c + d*x)/2, (2*b)/(a + b)]))/((a - b)^2*b)) + ((-5*a^2*A*b + A*b^3 + 2*a^3*B + 2*a*b^2*B + b*(-4*a*A*b + a^2*B + 3*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/(a^2 - b^2)^2))/(3*d*(a + b*Cos[c + d*x])^(3/2))`

3.337.3 Rubi [A] (verified)

Time = 1.31 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3233, 27, 3042, 3233, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{(a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{5/2}} dx \\ & \quad \downarrow \text{3233} \\ & \frac{2 \int -\frac{3(aA-bB)-(Ab-aB) \cos(c+dx)}{2(a+b \cos(c+dx))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2(Ab - aB) \sin(c + dx)}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{3(aA-bB)-(Ab-aB) \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2(Ab - aB) \sin(c + dx)}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \end{aligned}$$

3.337. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \int \frac{3(aA-bB)+(aB-Ab)\sin(c+dx+\frac{\pi}{2})}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \quad \downarrow \quad \mathbf{3042} \\
 & \frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \downarrow \quad \mathbf{3233} \\
 & \frac{2 \int \frac{3Aa^2-4bBa+Ab^2+(-Ba^2+4Aba-3b^2B)\cos(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} - \frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \downarrow \quad \mathbf{27} \\
 & \frac{\int \frac{3Aa^2-4bBa+Ab^2+(-Ba^2+4Aba-3b^2B)\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} - \frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \downarrow \quad \mathbf{3042} \\
 & \frac{\int \frac{3Aa^2-4bBa+Ab^2+(-Ba^2+4Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} - \frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \downarrow \quad \mathbf{3231} \\
 & \frac{(a^2(-B)+4aAb-3b^2B) \int \sqrt{a+b\cos(c+dx)} dx}{b} - \frac{(a^2-b^2)(Ab-aB) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \downarrow \quad \mathbf{3042} \\
 & \frac{(a^2(-B)+4aAb-3b^2B) \int \sqrt{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{(a^2-b^2)(Ab-aB) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \\
 & \frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
 & \downarrow \quad \mathbf{3134}
 \end{aligned}$$

3.337. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\frac{(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}dx}{b\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{(a^2-b^2)(Ab-aB)\int\frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{b}$$

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)}$$

$$\frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}dx}{b\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{(a^2-b^2)(Ab-aB)\int\frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{b}$$

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)}$$

$$\frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3132

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{(a^2-b^2)(Ab-aB)\int\frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{b}$$

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)}$$

$$\frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3142

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}}dx}{b\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)}$$

$$\frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}-\frac{(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}}dx}{b\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2(a^2(-B)+4aAb-3b^2B)\sin(c+dx)}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)}$$

$$\frac{2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

3.337. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

↓ 3140

$$\frac{\frac{2(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}E\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - 2(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{bd\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}}{a^2-b^2} - \frac{2(a^2(-B)+4aAb-3b^2B)\sqrt{a+b\cos(c+dx)}}{d(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)2(Ab-aB)\sin(c+dx)}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

input `Int[(A + B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(5/2), x]`

output `(-2*(A*b - a*B)*Sin[c + d*x])/(3*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((2*(4*a*A*b - a^2*B - 3*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*(A*b - a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(b*d*Sqrt[a + b*Cos[c + d*x]]))/(a^2 - b^2) - (2*(4*a*A*b - a^2*B - 3*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]))/(3*(a^2 - b^2))`

3.337.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

3.337.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 753 vs. 2(313) = 626.

Time = 19.20 (sec) , antiderivative size = 754, normalized size of antiderivative = 2.74

method	result
default	$\frac{\sqrt{-2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - a + b} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{2B\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (a+b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + \dots\right)$
parts	Expression too large to display

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

$$3.337. \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

output

```

-((-2*b*cos(1/2*d*x+1/2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*B/b/sin
(1/2*d*x+1/2*c)^2/(2*b*sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*sin(1/2*d*x
+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*cos(1/2*d*x+1/2*c)*sin(1/
2*d*x+1/2*c)^2*b+EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a
-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*a
-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+
1/2*c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*b)+2*(A*b-B*a)/b*
(1/6/b/(a-b)/(a+b)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin
(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2+1/2*(a-b)/b)^2+8/3*sin(1/2*
d*x+1/2*c)^2*b/(a-b)^2/(a+b)^2*cos(1/2*d*x+1/2*c)*a/(-(-2*b*cos(1/2*d*x+1/
2*c)^2-a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)+(3*a-b)/(3*a^3+3*a^2*b-3*a*b^2-3*b
^3)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1
/2)/(-2*sin(1/2*d*x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF
(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))-4/3*a/(a-b)/(a+b)^2*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*((2*b*cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^(1/2)/(-2*sin(1/2*d*
x+1/2*c)^4*b+(a+b)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*
c),(-2*b/(a-b))^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b))^(1/2))))
/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)^2+a+b)^(1/2)/d

```

3.337.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.16 (sec) , antiderivative size = 956, normalized size of antiderivative = 3.48

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output

```

1/9*(6*(2*B*a^3*b^2 - 5*A*a^2*b^3 + 2*B*a*b^4 + A*b^5 + (B*a^2*b^3 - 4*A*a
*b^4 + 3*B*b^5)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sin(d*x + c) + (sqr
t(2)*(-2*I*B*a^3*b^2 - I*A*a^2*b^3 + 6*I*B*a*b^4 - 3*I*A*b^5)*cos(d*x + c)
^2 - 2*sqrt(2)*(2*I*B*a^4*b + I*A*a^3*b^2 - 6*I*B*a^2*b^3 + 3*I*A*a*b^4)*c
os(d*x + c) + sqrt(2)*(-2*I*B*a^5 - I*A*a^4*b + 6*I*B*a^3*b^2 - 3*I*A*a^2*
b^3))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 -
9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (sqrt
(2)*(2*I*B*a^3*b^2 + I*A*a^2*b^3 - 6*I*B*a*b^4 + 3*I*A*b^5)*cos(d*x + c)^2
- 2*sqrt(2)*(-2*I*B*a^4*b - I*A*a^3*b^2 + 6*I*B*a^2*b^3 - 3*I*A*a*b^4)*co
s(d*x + c) + sqrt(2)*(2*I*B*a^5 + I*A*a^4*b - 6*I*B*a^3*b^2 + 3*I*A*a^2*b^
3))*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*
a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(sqrt
(2)*(I*B*a^2*b^3 - 4*I*A*a*b^4 + 3*I*B*b^5)*cos(d*x + c)^2 + 2*sqrt(2)*(I*
B*a^3*b^2 - 4*I*A*a^2*b^3 + 3*I*B*a*b^4)*cos(d*x + c) + sqrt(2)*(I*B*a^4*b
- 4*I*A*a^3*b^2 + 3*I*B*a^2*b^3))*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*
b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*
b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d
*x + c) + 2*a)/b)) - 3*(sqrt(2)*(-I*B*a^2*b^3 + 4*I*A*a*b^4 - 3*I*B*b^5)*c
os(d*x + c)^2 + 2*sqrt(2)*(-I*B*a^3*b^2 + 4*I*A*a^2*b^3 - 3*I*B*a*b^4)*cos
(d*x + c) + sqrt(2)*(-I*B*a^4*b + 4*I*A*a^3*b^2 - 3*I*B*a^2*b^3))*sqrt(...

```

3.337.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.337.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/(b*cos(d*x + c) + a)^(5/2), x)`

3.337.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/(b*cos(d*x + c) + a)^(5/2), x)`

3.337.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))/(a + b*cos(c + d*x))^(5/2), x)`

3.338
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

3.338.1 Optimal result 3162
 3.338.2 Mathematica [C] (verified) 3163
 3.338.3 Rubi [A] (verified) 3163
 3.338.4 Maple [B] (verified) 3171
 3.338.5 Fricas [F(-1)] 3172
 3.338.6 Sympy [F] 3173
 3.338.7 Maxima [F] 3173
 3.338.8 Giac [F] 3173
 3.338.9 Mupad [F(-1)] 3174

3.338.1 Optimal result

Integrand size = 31, antiderivative size = 349

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx =$$

$$\frac{2(7a^2Ab - 3Ab^3 - 4a^3B) \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{3a^2 (a^2 - b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{2(Ab - aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{3a (a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{2A \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{a^2 d \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{2b(Ab - aB) \sin(c + dx)}{3a (a^2 - b^2) d (a + b \cos(c + dx))^{3/2}} + \frac{2b(7a^2Ab - 3Ab^3 - 4a^3B) \sin(c + dx)}{3a^2 (a^2 - b^2)^2 d \sqrt{a + b \cos(c + dx)}}$$

output

```
2/3*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)+2/3*b*(7*A
*a^2*b-3*A*b^3-4*B*a^3)*sin(d*x+c)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2
)-2/3*(7*A*a^2*b-3*A*b^3-4*B*a^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x
+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x
+c))^(1/2)/a^2/(a^2-b^2)^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2/3*(A*b-B*a)*
(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*
c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/a/(a^2-b^2)/d/(
a+b*cos(d*x+c))^(1/2)+2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))
/(a+b))^(1/2)/a^2/d/(a+b*cos(d*x+c))^(1/2)
```

3.338.
$$\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

3.338.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.92 (sec) , antiderivative size = 553, normalized size of antiderivative = 1.58

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{\cos(c + dx)(B + A \sec(c + dx)) \left(\frac{4a(-6a^2Ab + 2Ab^3 + 3a^3B + ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{\sqrt{a+b \cos(c+dx)}} \right)}{(a + b \cos(c + dx))^{5/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(5/2),x]`

output `(Cos[c + d*x]*(B + A*Sec[c + d*x])*(((4*a*(-6*a^2*A*b + 2*A*b^3 + 3*a^3*B + a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(6*a^4*A - 19*a^2*A*b^2 + 9*A*b^4 + 4*a^3*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + ((2*I)*(-7*a^2*A*b + 3*A*b^3 + 4*a^3*B)*Sqrt[-((b*(-1 + Cos[c + d*x]))/(a + b))]*Sqrt[-((b*(1 + Cos[c + d*x]))/(a - b))]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b))))/(a*b*Sqrt[-(a + b)^(-1)])/(a^2*(a - b)^2*(a + b)^2) - (4*b*(a*(-8*a^2*A*b + 4*A*b^3 + 5*a^3*B - a*b^2*B) + b*(-7*a^2*A*b + 3*A*b^3 + 4*a^3*B)*Cos[c + d*x])*Sin[c + d*x])/((a^3 - a*b^2)^2*(a + b*Cos[c + d*x])^(3/2)))/(6*d*(A + B*Cos[c + d*x]))`

3.338.3 Rubi [A] (verified)

Time = 2.89 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.07, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.677$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.338. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\sec(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{3479} \\
& \frac{2 \int \frac{(b(Ab-aB)\cos^2(c+dx)-3a(Ab-aB)\cos(c+dx)+3A(a^2-b^2))\sec(c+dx)}{2(a+b\cos(c+dx))^{3/2}} dx}{\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} + \frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{(b(Ab-aB)\cos^2(c+dx)-3a(Ab-aB)\cos(c+dx)+3A(a^2-b^2))\sec(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx}{\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} + \frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2-3a(Ab-aB)\sin(c+dx+\frac{\pi}{2})+3A(a^2-b^2)}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}} + \frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3534} \\
& \frac{2 \int \frac{(3A(a^2-b^2)^2-b(-4Ba^3+7Aba^2-3Ab^3)\cos^2(c+dx)-a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\cos(c+dx))\sec(c+dx)}{2\sqrt{a+b\cos(c+dx)}a(a^2-b^2)} dx}{\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} + \frac{2b(-4a^3B+7a^2Ab-3Ab^3)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{(3A(a^2-b^2)^2-b(-4Ba^3+7Aba^2-3Ab^3)\cos^2(c+dx)-a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\cos(c+dx))\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}a(a^2-b^2)} dx}{\frac{3a(a^2-b^2)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} + \frac{2b(-4a^3B+7a^2Ab-3Ab^3)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx
\end{aligned}$$

3.338. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\frac{\int \frac{3A(a^2-b^2)^2 - b(-4Ba^3 + 7Aba^2 - 3Ab^3) \sin(c+dx + \frac{\pi}{2})^2 - a(-3Ba^3 + 6Aba^2 - b^2Ba - 2Ab^3) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{a(a^2-b^2)} + \frac{2b(-4a^3B + 7a^2Ab - 3Ab^3) \sin(c+dx)}{ad(a^2-b^2) \sqrt{a+b \cos(c+dx)}} +$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$\frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3538

$$\frac{\int -\frac{(3Ab(a^2-b^2)^2 + ab(Ab-aB) \cos(c+dx)(a^2-b^2)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)} - \frac{((-4a^3B + 7a^2Ab - 3Ab^3) \int \sqrt{a+b \cos(c+dx)} dx)}{a(a^2-b^2)} + \frac{2b(-4a^3B + 7a^2Ab - 3Ab^3)}{ad(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$\frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 25

$$\frac{\int \frac{(3Ab(a^2-b^2)^2 + ab(Ab-aB) \cos(c+dx)(a^2-b^2)) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)} - \frac{(-4a^3B + 7a^2Ab - 3Ab^3) \int \sqrt{a+b \cos(c+dx)} dx}{a(a^2-b^2)} + \frac{2b(-4a^3B + 7a^2Ab - 3Ab^3) \sin(c+dx)}{ad(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$\frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{\int \frac{3Ab(a^2-b^2)^2 + ab(Ab-aB) \sin(c+dx + \frac{\pi}{2})(a^2-b^2)}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{a(a^2-b^2)} - \frac{(-4a^3B + 7a^2Ab - 3Ab^3) \int \sqrt{a+b \sin(c+dx + \frac{\pi}{2})} dx}{a(a^2-b^2)} + \frac{2b(-4a^3B + 7a^2Ab - 3Ab^3) \sin(c+dx)}{ad(a^2-b^2) \sqrt{a+b \cos(c+dx)}}$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$\frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3134

$$\frac{\int \frac{3Ab(a^2-b^2)^2 + ab(Ab-aB) \sin(c+dx + \frac{\pi}{2})(a^2-b^2)}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{a(a^2-b^2)} - \frac{(-4a^3B + 7a^2Ab - 3Ab^3) \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$\frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3042

3.338. $\int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{\int \frac{3Ab(a^2-b^2)^2 + ab(Ab-aB)\sin(c+dx+\frac{\pi}{2})(a^2-b^2)}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{3a(a^2-b^2)}{a(a^2-b^2)} + \frac{2b(-4a^3B+7a^2Ab-3Ab^3)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3132

$$\frac{\int \frac{3Ab(a^2-b^2)^2 + ab(Ab-aB)\sin(c+dx+\frac{\pi}{2})(a^2-b^2)}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{3a(a^2-b^2)}{a(a^2-b^2)} + \frac{2b(-4a^3B+7a^2Ab-3Ab^3)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3481

$$\frac{ab(a^2-b^2)(Ab-aB) \int \frac{1}{\sqrt{a+b\cos(c+dx)}} dx + 3Ab(a^2-b^2)^2 \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{3a(a^2-b^2)}{a(a^2-b^2)} + \frac{2b(-4a^3B+7a^2Ab-3Ab^3)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{ab(a^2-b^2)(Ab-aB) \int \frac{1}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + 3Ab(a^2-b^2)^2 \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}E(\frac{1}{2}(c+dx)|\frac{2b}{a+b})}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{3a(a^2-b^2)}{a(a^2-b^2)} + \frac{2b(-4a^3B+7a^2Ab-3Ab^3)\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3142

3.338. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\frac{ab(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{3Ab(a^2-b^2)^2 \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3a(a^2-b^2)$

3042

$$\frac{ab(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} + \frac{3Ab(a^2-b^2)^2 \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3a(a^2-b^2)$

3140

$$\frac{3Ab(a^2-b^2)^2 \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2ab(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3a(a^2-b^2)$

3286

$$\frac{3Ab(a^2-b^2)^2 \sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b\cos(c+dx)}{a+b}}} dx + \frac{2ab(a^2-b^2)(Ab-aB)\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}}{b} - \frac{2(-4a^3B+7a^2Ab-3Ab^3)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\frac{a+b\cos(c+dx)}{a+b}}}$$

$$\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

$3a(a^2-b^2)$

3042

3.338. $\int \frac{(A+B\cos(c+dx))\sec(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{3Ab(a^2-b^2)^2 \int \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} + \frac{2ab(a^2-b^2)(Ab-aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} - \frac{2(-4a^3B+7a^2A^2)}{a(a^2-b^2)} \\
 & \frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3284} \\
 & \frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} + \frac{2ab(a^2-b^2)(Ab-aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} + \frac{6Ab(a^2-b^2)^2 \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d \sqrt{a+b \cos(c+dx)}} \\
 & \frac{2b(-4a^3B+7a^2A^2) \sin(c+dx)}{ad(a^2-b^2) \sqrt{a+b \cos(c+dx)}} + \frac{2(-4a^3B+7a^2A^2)}{a(a^2-b^2)}
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*b*(A*b - a*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((-2*(7*a^2*A*b - 3*A*b^3 - 4*a^3*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((2*a*b*(a^2 - b^2)*(A*b - a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*A*b*(a^2 - b^2)^2*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b/(a*(a^2 - b^2)) + (2*b*(7*a^2*A*b - 3*A*b^3 - 4*a^3*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]))/((3*a*(a^2 - b^2))`

3.338.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.338. \int \frac{(A+B \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Ssin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Ssin[e + f*x])^m/(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*(c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
] && NeQ[c^2 - d^2, 0]
```

3.338.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 857 vs. $2(412) = 824$.

Time = 19.82 (sec) , antiderivative size = 858, normalized size of antiderivative = 2.46

method	result	size
default	Expression too large to display	858
parts	Expression too large to display	1340

```
input int((A+B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVER
BOSE)
```

output
$$\begin{aligned} & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*A/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)})/ \\ & (-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c), \\ & 2, (-2*b/(a-b))^{(1/2)})+2*A*b/a^2/\sin(1/2*d*x+1/2*c)^2/(2*b*\sin(1/2*d*x+1/2*c)^2-a-b)/ \\ & (a^2-b^2)*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2+ \\ & EllipticE(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)}* \\ & (\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a-EllipticE(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)}))*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)}* \\ & (\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b)+2*(-A*b+B*a)/a*(1/6/b/(a-b)/(a+b)*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/ \\ & (\cos(1/2*d*x+1/2*c)^2+1/2*(a-b)/b)^2+8/3*\sin(1/2*d*x+1/2*c)^2*b/(a-b)^2/(a+b)^2*\cos(1/2*d*x+1/2*c)*a/ \\ & (-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+(3*a-b)/(3*a^3+3*a^2*b-3*a*b^2-3*b^3)*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}* \\ & ((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}* \\ & EllipticF(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})-4/3*a/(a-b)/(a+b)^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/ \\ & (-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(EllipticF(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})- \\ & EllipticE(\cos(1/2*d*x+1/2*c), (-2*b/(a-b))^{(1/2)})))/\sin(1/2*d*x+1/2*c) \dots \end{aligned}$$

3.338.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `Timed out`

3.338.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**(5/2), x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)/(a + b*cos(c + d*x))**(5/2), x)`

3.338.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)/(b*cos(d*x + c) + a)^(5/2), x)`

3.338.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(5/2), x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)/(b*cos(d*x + c) + a)^(5/2), x)`

3.338.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx) (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(5/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(5/2)), x)`

$$3.339 \quad \int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

3.339.1 Optimal result	3175
3.339.2 Mathematica [C] (warning: unable to verify)	3176
3.339.3 Rubi [A] (verified)	3177
3.339.4 Maple [B] (warning: unable to verify)	3185
3.339.5 Fracas [F(-1)]	3186
3.339.6 Sympy [F]	3187
3.339.7 Maxima [F]	3187
3.339.8 Giac [F]	3187
3.339.9 Mupad [F(-1)]	3188

3.339.1 Optimal result

Integrand size = 33, antiderivative size = 437

$$\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx =$$

$$\frac{(3a^4A - 26a^2Ab^2 + 15Ab^4 + 14a^3bB - 6ab^3B) \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{3a^3(a^2-b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}$$

$$+ \frac{(3a^2A - 5Ab^2 + 2abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{3a^2(a^2-b^2) d \sqrt{a+b \cos(c+dx)}}$$

$$- \frac{(5Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{a^3 d \sqrt{a+b \cos(c+dx)}}$$

$$+ \frac{b(3a^2A - 5Ab^2 + 2abB) \sin(c+dx)}{3a^2(a^2-b^2) d(a+b \cos(c+dx))^{3/2}}$$

$$+ \frac{b(3a^4A - 26a^2Ab^2 + 15Ab^4 + 14a^3bB - 6ab^3B) \sin(c+dx)}{3a^3(a^2-b^2)^2 d \sqrt{a+b \cos(c+dx)}} + \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}}$$

```
output 1/3*b*(3*A*a^2-5*A*b^2+2*B*a*b)*sin(d*x+c)/a^2/(a^2-b^2)/d/(a+b*cos(d*x+c)
)^3/2+1/3*b*(3*A*a^4-26*A*a^2*b^2+15*A*b^4+14*B*a^3*b-6*B*a*b^3)*sin(d*x
+c)/a^3/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)-1/3*(3*A*a^4-26*A*a^2*b^2+15*
A*b^4+14*B*a^3*b-6*B*a*b^3)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c
)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(
1/2)/a^3/(a^2-b^2)^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+1/3*(3*A*a^2-5*A*b^2
+2*B*a*b)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/
2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/a^2/(
a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)-(5*A*b-2*B*a)*(cos(1/2*d*x+1/2*c))^2^(1/
2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1
/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/a^3/d/(a+b*cos(d*x+c))^(1/2)+A*tan(d*x
+c)/a/d/(a+b*cos(d*x+c))^(3/2)
```

3.339.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.88 (sec) , antiderivative size = 750, normalized size of antiderivative = 1.72

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2(36a^3 Ab^2 - 20aAb^4 - 24a^4 bB + 8a^2 b^3 B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right) + 2(\dots)}{\sqrt{a+b \cos(c+dx)}} + \frac{\sqrt{a+b \cos(c+dx)} \left(\frac{2(-Ab^3 \sin(c+dx) + ab^2 B \sin(c+dx))}{3a^2(a^2-b^2)(a+b \cos(c+dx))^2} + \frac{2(-10a^2 Ab^3 \sin(c+dx) + 6Ab^5 \sin(c+dx) + 7a^3 b^2 B \sin(c+dx) - 3ab^4 B \sin(c+dx))}{3a^3(a^2-b^2)^2(a+b \cos(c+dx))} \right)}{d}$$

```
input Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^(5/2)
,x]
```

output $((2*(36*a^3*A*b^2 - 20*a*A*b^4 - 24*a^4*b*B + 8*a^2*b^3*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])]/(a + b))*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(-33*a^4*A*b + 86*a^2*A*b^3 - 45*A*b^5 + 12*a^5*B - 38*a^3*b^2*B + 18*a*b^4*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])]/(a + b))*\text{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)]/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] - ((2*I)*(-3*a^4*A*b + 26*a^2*A*b^3 - 15*A*b^5 - 14*a^3*b^2*B + 6*a*b^4*B)*\text{Sqrt}[(b - b*\text{Cos}[c + d*x])]/(a + b))*\text{Sqrt}[-((b + b*\text{Cos}[c + d*x])/(a - b))]*\text{Cos}[2*(c + d*x)]*(2*a*(a - b)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)) + b*(2*a*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)) - b*\text{EllipticPi}[(a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b))*\text{Sin}[c + d*x])/((a*\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[1 - \text{Cos}[c + d*x]^2]*\text{Sqrt}[-((a^2 - b^2 - 2*a*(a + b*\text{Cos}[c + d*x]) + (a + b*\text{Cos}[c + d*x])^2)/b^2)]*(2*a^2 - b^2 - 4*a*(a + b*\text{Cos}[c + d*x]) + 2*(a + b*\text{Cos}[c + d*x])^2)))/(12*a^3*(-a + b)^2*(a + b)^2*d) + (\text{Sqrt}[a + b*\text{Cos}[c + d*x]]*((2*(-(A*b^3*\text{Sin}[c + d*x]) + a*b^2*B*\text{Sin}[c + d*x]))/(3*a^2*(a^2 - b^2)*(a + b*\text{Cos}[c + d*x])^2) + (2*(-10*a^2*A*b^3*\text{Sin}[c + d*x] + 6*A*b^5*\text{Sin}[c + d*x] + 7*a^3*b^2*B*\text{Sin}[c + d*x] - 3*a*b^4*B*\text{Sin}[c + d*x]))/(3*a^3*(a^2 - b^2)^2*(a + b*\text{Cos}[c + d*x])) + (A*\text{Tan}[c + d*x])/a^3))/d$

3.339.3 Rubi [A] (verified)

Time = 3.75 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.08, number of steps used = 24, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 3479, 27, 3042, 3535, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^2(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^2 (a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx$$

↓ 3479

$$\frac{\int -\frac{(-3Ab \cos^2(c+dx)+5Ab-2aB) \sec(c+dx)}{2(a+b \cos(c+dx))^{5/2}} dx}{a} + \frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}}$$

3.339. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{(-3Ab \cos^2(c+dx)+5Ab-2aB) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx}{2a} \\
 & \quad \downarrow \text{27} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{-3Ab \sin(c+dx+\frac{\pi}{2})^2+5Ab-2aB}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))^{5/2}} dx}{2a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{2 \int \frac{(-b(3Aa^2+2bBa-5Ab^2) \cos^2(c+dx)-6ab(Ab-aB) \cos(c+dx)+3(a^2-b^2)(5Ab-2aB)) \sec(c+dx)}{2(a+b \cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} - \frac{2b(3a^2A+2abB-5Ab^2) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3535} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{2 \int \frac{(-b(3Aa^2+2bBa-5Ab^2) \cos^2(c+dx)-6ab(Ab-aB) \cos(c+dx)+3(a^2-b^2)(5Ab-2aB)) \sec(c+dx)}{2(a+b \cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} - \frac{2b(3a^2A+2abB-5Ab^2) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{(-b(3Aa^2+2bBa-5Ab^2) \cos^2(c+dx)-6ab(Ab-aB) \cos(c+dx)+3(a^2-b^2)(5Ab-2aB)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} - \frac{2b(3a^2A+2abB-5Ab^2) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{-b(3Aa^2+2bBa-5Ab^2) \sin(c+dx+\frac{\pi}{2})^2-6ab(Ab-aB) \sin(c+dx+\frac{\pi}{2})+3(a^2-b^2)(5Ab-2aB)}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3a(a^2-b^2)} - \frac{2b(3a^2A+2abB-5Ab^2) \sin(c+dx)}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3534} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{2 \int \frac{(3(5Ab-2aB)(a^2-b^2)^2+b(3Aa^4+14bBa^3-26Ab^2a^2-6b^3Ba+15Ab^4) \cos^2(c+dx)-2ab(-6Ba^3+9Aba^2+2b^2Ba-5Ab^3) \cos(c+dx)) \sec(c+dx)}{2\sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)} - \frac{2b(3a^4A+2ab^3B-5Ab^2a^2)}{3a(a^2-b^2)} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.339. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \int \frac{(3(5Ab - 2aB)(a^2 - b^2)^2 + b(3Aa^4 + 14bBa^3 - 26Ab^2a^2 - 6b^3Ba + 15Ab^4) \cos^2(c + dx) - 2ab(-6Ba^3 + 9Aba^2 + 2b^2Ba - 5Ab^3) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)} a(a^2 - b^2)} dx - \frac{2b(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4)}{3a(a^2 - b^2)}$$

2a

3042

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \int \frac{3(5Ab - 2aB)(a^2 - b^2)^2 + b(3Aa^4 + 14bBa^3 - 26Ab^2a^2 - 6b^3Ba + 15Ab^4) \sin(c + dx + \frac{\pi}{2})^2 - 2ab(-6Ba^3 + 9Aba^2 + 2b^2Ba - 5Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} a(a^2 - b^2)} dx - \frac{2b(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4)}{3a(a^2 - b^2)}$$

2a

3538

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \int \frac{(3b(a^2 - b^2)^2(5Ab - 2aB) - ab(a^2 - b^2)(3Aa^2 + 2bBa - 5Ab^2) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)} a(a^2 - b^2)} dx - \frac{(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \int \sqrt{a + b \cos(c + dx)} dx - \frac{2b(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4)}{b}}{3a(a^2 - b^2)}$$

2a

25

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \int \frac{(3b(a^2 - b^2)^2(5Ab - 2aB) - ab(a^2 - b^2)(3Aa^2 + 2bBa - 5Ab^2) \cos(c + dx)) \sec(c + dx)}{\sqrt{a + b \cos(c + dx)} a(a^2 - b^2)} dx + \frac{(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \int \sqrt{a + b \cos(c + dx)} dx - \frac{2b(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4)}{b}}{3a(a^2 - b^2)}$$

2a

3042

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \int \frac{3b(a^2 - b^2)^2(5Ab - 2aB) - ab(a^2 - b^2)(3Aa^2 + 2bBa - 5Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} a(a^2 - b^2)} dx + \frac{(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx - \frac{2b(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4)}{b}}{3a(a^2 - b^2)}$$

2a

3134

3.339. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{\int \frac{3b(a^2 - b^2)^2(5Ab - 2aB) - ab(a^2 - b^2)(3Aa^2 + 2bBa - 5Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \frac{(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c + dx)}{a+b}}}{\sqrt{\frac{a + b \cos(c + dx)}{a+b}}}$$

$$\frac{3a(a^2 - b^2)}{2a}$$

3042

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{\int \frac{3b(a^2 - b^2)^2(5Ab - 2aB) - ab(a^2 - b^2)(3Aa^2 + 2bBa - 5Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \frac{(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c + dx)}{a+b}}}{\sqrt{\frac{a + b \cos(c + dx)}{a+b}}}$$

$$\frac{3a(a^2 - b^2)}{2a}$$

3132

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{\int \frac{3b(a^2 - b^2)^2(5Ab - 2aB) - ab(a^2 - b^2)(3Aa^2 + 2bBa - 5Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \frac{2(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \sqrt{a + b \cos(c + dx)} E(\frac{1}{2}(c + dx) | \frac{2b}{a+b})}{d \sqrt{\frac{a + b \cos(c + dx)}{a+b}}}$$

$$\frac{3a(a^2 - b^2)}{2a}$$

3481

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{3b(a^2 - b^2)^2(5Ab - 2aB) \int \frac{\sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx - ab(a^2 - b^2)(3a^2A + 2abB - 5Ab^2) \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx}{a(a^2 - b^2)} + \frac{2(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\frac{a + b \cos(c + dx)}{a+b}}}$$

$$\frac{3a(a^2 - b^2)}{2a}$$

3042

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{3b(a^2 - b^2)^2(5Ab - 2aB) \int \frac{1}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - ab(a^2 - b^2)(3a^2A + 2abB - 5Ab^2) \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \frac{2(3a^4A + 14a^3bB - 26a^2Ab^2 - 6ab^3B + 15Ab^4) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\frac{a + b \cos(c + dx)}{a+b}}}$$

$$\frac{3a(a^2 - b^2)}{2a}$$

3.339. $\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow \text{3142} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \\
 & \frac{3b(a^2-b^2)^2(5Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(a^2-b^2)(3a^2A+2abB-5Ab^2) \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}}}{b} + \frac{2(3a^4A+14a^3B)}{a(a^2-b^2)} \\
 & \frac{3a(a^2-b^2)}{3a(a^2-b^2)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \\
 & \frac{3b(a^2-b^2)^2(5Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(a^2-b^2)(3a^2A+2abB-5Ab^2) \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}}}{b} + \frac{2(3a^4A+14a^3B)}{a(a^2-b^2)} \\
 & \frac{3a(a^2-b^2)}{3a(a^2-b^2)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3140} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \\
 & \frac{3b(a^2-b^2)^2(5Ab-2aB) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2ab(a^2-b^2)(3a^2A+2abB-5Ab^2) \int \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b})}{d\sqrt{a+b \cos(c+dx)}}}{b}}{a(a^2-b^2)} + \frac{2(3a^4A+14a^3B)}{3a(a^2-b^2)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3286} \\
 & \frac{A \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \\
 & \frac{3b(a^2-b^2)^2(5Ab-2aB) \int \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{sec}(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - \frac{2ab(a^2-b^2)(3a^2A+2abB-5Ab^2) \int \frac{\sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b})}{d\sqrt{a+b \cos(c+dx)}}}{b}}{a(a^2-b^2)} + \frac{2(3a^4A+14a^3B)}{3a(a^2-b^2)}
 \end{aligned}$$

$$\downarrow \text{3042}$$

3.339. $\int \frac{(A+B \cos(c+dx)) \operatorname{sec}^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{3b(a^2 - b^2)^2 (5Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)} b} - \frac{2ab(a^2 - b^2) (3a^2 A + 2abB - 5Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b})}{d \sqrt{a+b \cos(c+dx)} a(a^2 - b^2)}$$

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{6b(a^2 - b^2)^2 (5Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticPi}(2, \frac{1}{2}(c+dx), \frac{2b}{a+b})}{d \sqrt{a+b \cos(c+dx)} b} - \frac{2ab(a^2 - b^2) (3a^2 A + 2abB - 5Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b})}{d \sqrt{a+b \cos(c+dx)} a(a^2 - b^2)} + \frac{2(3a^4 A - 2(3a^2 A + 2abB - 5Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}})}{3a(a^2 - b^2)}$$

↓ 3284

$$\frac{A \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} - \frac{6b(a^2 - b^2)^2 (5Ab - 2aB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticPi}(2, \frac{1}{2}(c+dx), \frac{2b}{a+b})}{d \sqrt{a+b \cos(c+dx)} b} - \frac{2ab(a^2 - b^2) (3a^2 A + 2abB - 5Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}(\frac{1}{2}(c+dx), \frac{2b}{a+b})}{d \sqrt{a+b \cos(c+dx)} a(a^2 - b^2)} + \frac{2(3a^4 A - 2(3a^2 A + 2abB - 5Ab^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}})}{3a(a^2 - b^2)}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^2)/(a + b*Cos[c + d*x])^(5/2),x]
```

```
output -1/2*((-2*b*(3*a^2*A - 5*A*b^2 + 2*a*b*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((2*(3*a^4*A - 26*a^2*A*b^2 + 15*A*b^4 + 14*a^3*b*B - 6*a*b^3*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((-2*a*b*(a^2 - b^2)*(3*a^2*A - 5*A*b^2 + 2*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*b*(a^2 - b^2)^2*(5*A*b - 2*a*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/(a*(a^2 - b^2)) - (2*b*(3*a^4*A - 26*a^2*A*b^2 + 15*A*b^4 + 14*a^3*b*B - 6*a*b^3*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]))/(3*a*(a^2 - b^2)))/a + (A*Tan[c + d*x])/(a*d*(a + b*Cos[c + d*x])^(3/2))
```

3.339.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.339. $\int \frac{(A+B \cos(c+dx)) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m + 1)*((c + d*SIN[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*SIN[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*SIN[e + f*x])^m/(c + d*SIN[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m + 1)*((c + d*SIN[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3535 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
  (f_.)*(x_)])^(n_)*((A_.) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] :=
Simp[(-(A*b^2 + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
in[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m
+ 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin
[e + f*x])^n*Simp[a*(m + 1)*(b*c - a*d)*(A + C) + d*(A*b^2 + a^2*C)*(m + n
+ 2) - (c*(A*b^2 + a^2*C) + b*(m + 1)*(b*c - a*d)*(A + C))*Sin[e + f*x] - d
*(A*b^2 + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) ||
!(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a,
0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2]/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
  (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.339.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1344 vs. 2(498) = 996.

Time = 29.58 (sec) , antiderivative size = 1345, normalized size of antiderivative = 3.08

method	result	size
default	Expression too large to display	1345
parts	Expression too large to display	2175

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNV
ERBOSE)
```


output
$$\begin{aligned} & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*A/a^2*(- \\ & \cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2 \\ &)^{(1/2)})/(2*\cos(1/2*d*x+1/2*c)^2-1)+1/2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b* \\ & \cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin \\ & (1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)}) \\ & -1/2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)} \\ &)/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*Elliptic \\ & E(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+1/2/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & *((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+ \\ & (a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a- \\ & b))^{(1/2)})+1/2/a*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2 \\ & +a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & *EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-2*(-2*A*b+B*a)/ \\ & a^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)} \\ &)/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticP \\ & i(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-2*b*(2*A*b-B*a)/a^3/\sin(1/2*d*x \\ & +1/2*c)^2/(2*b*\sin(1/2*d*x+1/2*c)^2-a-b)/(a^2-b^2)*(-2*\sin(1/2*d*x+1/2*c)^ \\ & 4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/ \\ & 2*c)^2*b+EllipticE(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})*(-2*b/(a-b)*\sin(\\ & 1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*a-Elli... \end{aligned}$$

3.339.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `Timed out`

3.339.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**2/(a+b*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**2/(a + b*cos(c + d*x))**(5/2), x)`

3.339.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^2}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^2/(b*cos(d*x + c) + a)^(5/2), x)`

3.339.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^2}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^2/(a+b*cos(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^2/(b*cos(d*x + c) + a)^(5/2), x)`

3.339.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^2(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^2 (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^(5/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^2*(a + b*cos(c + d*x))^(5/2)), x)`

3.340 $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

3.340.1 Optimal result 3189
 3.340.2 Mathematica [C] (warning: unable to verify) 3190
 3.340.3 Rubi [A] (verified) 3191
 3.340.4 Maple [B] (warning: unable to verify) 3200
 3.340.5 Fricas [F(-1)] 3201
 3.340.6 Sympy [F] 3202
 3.340.7 Maxima [F(-1)] 3202
 3.340.8 Giac [F] 3202
 3.340.9 Mupad [F(-1)] 3203

3.340.1 Optimal result

Integrand size = 33, antiderivative size = 532

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{(33a^4Ab - 170a^2Ab^3 + 105Ab^5 - 12a^5B + 104a^3b^2B - 60ab^4B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{12a^4(a^2 - b^2)^2 d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{(27a^2Ab - 35Ab^3 - 12a^3B + 20ab^2B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{12a^3(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}} + \frac{(4a^2A + 35Ab^2 - 20abB) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{4a^4 d \sqrt{a + b \cos(c + dx)}} - \frac{b(27a^2Ab - 35Ab^3 - 12a^3B + 20ab^2B) \sin(c + dx)}{12a^3(a^2 - b^2) d(a + b \cos(c + dx))^{3/2}} - \frac{b(33a^4Ab - 170a^2Ab^3 + 105Ab^5 - 12a^5B + 104a^3b^2B - 60ab^4B) \sin(c + dx)}{12a^4(a^2 - b^2)^2 d \sqrt{a + b \cos(c + dx)}} - \frac{(7Ab - 4aB) \tan(c + dx)}{4a^2 d(a + b \cos(c + dx))^{3/2}} + \frac{A \sec(c + dx) \tan(c + dx)}{2ad(a + b \cos(c + dx))^{3/2}}$$

output

```

-1/12*b*(27*A*a^2*b-35*A*b^3-12*B*a^3+20*B*a*b^2)*sin(d*x+c)/a^3/(a^2-b^2)
/d/(a+b*cos(d*x+c))^(3/2)-1/12*b*(33*A*a^4*b-170*A*a^2*b^3+105*A*b^5-12*B*
a^5+104*B*a^3*b^2-60*B*a*b^4)*sin(d*x+c)/a^4/(a^2-b^2)^2/d/(a+b*cos(d*x+c)
)^(1/2)+1/12*(33*A*a^4*b-170*A*a^2*b^3+105*A*b^5-12*B*a^5+104*B*a^3*b^2-60
*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/
2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/a^4/(a^2-b^2)
^2/d/((a+b*cos(d*x+c))/(a+b))^(1/2)-1/12*(27*A*a^2*b-35*A*b^3-12*B*a^3+20*
B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2
*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))/(a+b)^(1/2)/a^3/(a
^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+1/4*(4*A*a^2+35*A*b^2-20*B*a*b)*(cos(1/2*
d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^(
1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))/(a+b)^(1/2)/a^4/d/(a+b*cos(d*x+c)
)^(1/2)-1/4*(7*A*b-4*B*a)*tan(d*x+c)/a^2/d/(a+b*cos(d*x+c))^(3/2)+1/2*A*se
c(d*x+c)*tan(d*x+c)/a/d/(a+b*cos(d*x+c))^(3/2)
    
```

3.340.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.19 (sec) , antiderivative size = 820, normalized size of antiderivative = 1.54

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2(12a^5 Ab - 216a^3 Ab^3 + 140aAb^5 + 144a^4 b^2 B - 80a^2 b^4 B) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx)\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{\sqrt{a+b \cos(c+dx)} \left(\frac{\sec(c+dx)(-11Ab \sin(c+dx) + 4aB \sin(c+dx))}{4a^4} - \frac{2(-Ab^4 \sin(c+dx) + ab^3 B \sin(c+dx))}{3a^3(a^2-b^2)(a+b \cos(c+dx))^2} - \frac{2(-13a^2 Ab^4 \sin(c+dx))}{d} \right)}{d}$$

input

```

Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^(5/2)
,x]
    
```

output $((2*(12*a^5*A*b - 216*a^3*A*b^3 + 140*a*A*b^5 + 144*a^4*b^2*B - 80*a^2*b^4*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])]/(a + b))*\text{EllipticF}[(c + d*x)/2, (2*b)/(a + b)])/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] + (2*(24*a^6*A + 195*a^4*A*b^2 - 566*a^2*A*b^4 + 315*A*b^6 - 132*a^5*b*B + 344*a^3*b^3*B - 180*a*b^5*B)*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])]/(a + b))*\text{EllipticPi}[2, (c + d*x)/2, (2*b)/(a + b)])/\text{Sqrt}[a + b*\text{Cos}[c + d*x]] - ((2*I)*(33*a^4*A*b^2 - 170*a^2*A*b^4 + 105*A*b^6 - 12*a^5*b*B + 104*a^3*b^3*B - 60*a*b^5*B)*\text{Sqrt}[(b - b*\text{Cos}[c + d*x])]/(a + b))*\text{Sqrt}[-((b + b*\text{Cos}[c + d*x])/(a - b))]*\text{Cos}[2*(c + d*x)]*(2*a*(a - b)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] + b*(2*a*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)] - b*\text{EllipticPi}[(a + b)/a, I*\text{ArcSinh}[\text{Sqrt}[-(a + b)^{-1}]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]], (a + b)/(a - b)))*\text{Sin}[c + d*x])/(a*\text{Sqrt}[-(a + b)^{-1}]*\text{Sqrt}[1 - \text{Cos}[c + d*x]^2]*\text{Sqrt}[-((a^2 - b^2 - 2*a*(a + b*\text{Cos}[c + d*x]) + (a + b*\text{Cos}[c + d*x])^2)/b^2)]*(2*a^2 - b^2 - 4*a*(a + b*\text{Cos}[c + d*x]) + 2*(a + b*\text{Cos}[c + d*x])^2)))/(48*a^4*(a - b)^2*(a + b)^2*d) + (\text{Sqrt}[a + b*\text{Cos}[c + d*x]]*(\text{Sec}[c + d*x]*(-11*A*b*\text{Sin}[c + d*x] + 4*a*B*\text{Sin}[c + d*x]))/(4*a^4) - (2*(-(A*b^4*\text{Sin}[c + d*x]) + a*b^3*B*\text{Sin}[c + d*x]))/(3*a^3*(a^2 - b^2)*(a + b*\text{Cos}[c + d*x])^2) - (2*(-13*a^2*A*b^4*\text{Sin}[c + d*x] + 9*A*b^6*\text{Sin}[c + d*x] + 10*a^3*b^3*B*\text{Sin}[c + d*x] - 6*a*b^5*B*\text{Sin}[c + d*x]))/(3*a^4*(a^2 - b^2)^2*(a + b*\text{Cos}[c + d*x])) + (A*\text{Sec}[c + d*x]*\text{Tan}[c + d*x])...$

3.340.3 Rubi [A] (verified)

Time = 4.69 (sec) , antiderivative size = 569, normalized size of antiderivative = 1.07, number of steps used = 27, number of rules used = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3134, 3042, 3132, 3481, 3042, 3142, 3042, 3140, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^3(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 3479

$$\frac{\int \frac{(-5Ab\cos^2(c+dx)-2aA\cos(c+dx)+7Ab-4aB)\sec^2(c+dx)}{2(a+b\cos(c+dx))^{5/2}} dx}{2a} + \frac{A\tan(c+dx)\sec(c+dx)}{2ad(a+b\cos(c+dx))^{3/2}}$$

3.340. $\int \frac{(A+B\cos(c+dx))\sec^3(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{(-5Ab \cos^2(c+dx) - 2aA \cos(c+dx) + 7Ab - 4aB) \sec^2(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx}{4a} \\
& \downarrow 3042 \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{-5Ab \sin(c+dx+\frac{\pi}{2})^2 - 2aA \sin(c+dx+\frac{\pi}{2}) + 7Ab - 4aB}{\sin(c+dx+\frac{\pi}{2})^2 (a+b \sin(c+dx+\frac{\pi}{2}))^{5/2}} dx}{4a} \\
& \downarrow 3534 \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int -\frac{(4Aa^2 - 20bBa + 10Ab \cos(c+dx)a + 35Ab^2 - 3b(7Ab - 4aB) \cos^2(c+dx)) \sec(c+dx)}{2(a+b \cos(c+dx))^{5/2}} dx}{a} + \frac{(7Ab - 4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} \\
& \frac{4a}{4a} \\
& \downarrow 27 \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{(4Aa^2 - 20bBa + 10Ab \cos(c+dx)a + 35Ab^2 - 3b(7Ab - 4aB) \cos^2(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx}{2a} \\
& \frac{(7Ab - 4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{4a}{4a} \\
& \downarrow 3042 \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{4Aa^2 - 20bBa + 10Ab \sin(c+dx+\frac{\pi}{2})a + 35Ab^2 - 3b(7Ab - 4aB) \sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2}) (a+b \sin(c+dx+\frac{\pi}{2}))^{5/2}} dx}{2a} \\
& \frac{(7Ab - 4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{4a}{4a} \\
& \downarrow 3534 \\
& \frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{2 \int \frac{(-b(-12Ba^3 + 27Aba^2 + 20b^2Ba - 35Ab^3) \cos^2(c+dx) + 6ab(3Aa^2 + 4bBa - 7Ab^2) \cos(c+dx) + 3(a^2 - b^2)(4Aa^2 - 20bBa + 35Ab^2))}{2(a+b \cos(c+dx))^{3/2}}}{3a(a^2 - b^2)} \\
& \frac{(7Ab - 4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} - \frac{2a}{4a} \\
& \downarrow 27
\end{aligned}$$

3.340. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}}}{\frac{\int \frac{-b(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \cos^2(c+dx)+6ab(3Aa^2+4bBa-7Ab^2) \cos(c+dx)+3(a^2-b^2)(4Aa^2-20bBa+35Ab^2)}{(a+b \cos(c+dx))^{3/2}}}{\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}}}} = \frac{2a}{4a}$$

3042

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}}}{\frac{\int \frac{-b(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \sin(c+dx+\frac{\pi}{2})^2+6ab(3Aa^2+4bBa-7Ab^2) \sin(c+dx+\frac{\pi}{2})+3(a^2-b^2)(4Aa^2-20bBa+35Ab^2)}{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}}}{\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}}}} = \frac{2a}{4a}$$

3534

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}}}{\frac{2 \int \frac{(3(4Aa^2-20bBa+35Ab^2)(a^2-b^2)^2+b(-12Ba^5+33Aba^4+104b^2Ba^3-170Ab^3a^2-60b^4Ba+105Ab^5) \cos^2(c+dx)+2ab(3Aa^4+35Ab^4)) \sqrt{a+b \cos(c+dx)}}{a(a^2-b^2)}}}{\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}}}} = \frac{3a}{3a}$$

27

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}}}{\frac{\int \frac{(3(4Aa^2-20bBa+35Ab^2)(a^2-b^2)^2+b(-12Ba^5+33Aba^4+104b^2Ba^3-170Ab^3a^2-60b^4Ba+105Ab^5) \cos^2(c+dx)+2ab(3Aa^4+35Ab^4)) \sqrt{a+b \cos(c+dx)}}{a(a^2-b^2)}}}{\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}}}} = \frac{3a}{3a}$$

3042

$$\frac{\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}}}{\frac{\int \frac{3(4Aa^2-20bBa+35Ab^2)(a^2-b^2)^2+b(-12Ba^5+33Aba^4+104b^2Ba^3-170Ab^3a^2-60b^4Ba+105Ab^5) \sin(c+dx+\frac{\pi}{2})^2+2ab(3Aa^4+35Ab^4)}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}}{\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}}}} = \frac{3a}{3a}$$

3538

3.340. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{(-12a^5B+33a^4Ab+104a^3b^2B-170a^2Ab^3-60ab^4B+105Ab^5) \int \sqrt{a+b \cos(c+dx)} dx - \frac{3b(a^2-b^2)^2(4Aa^2-20bBa+35Ab^2)-a}{a(a^2-b^2)}}{a(a^2-b^2)}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 25

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{3b(a^2-b^2)^2(4Aa^2-20bBa+35Ab^2)-ab(a^2-b^2)(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \cos(c+dx) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)} + (-12a^5B+33a^4Ab+104a^3b^2B-170a^2Ab^3-60ab^4B+105Ab^5)$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3042

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{3b(a^2-b^2)^2(4Aa^2-20bBa+35Ab^2)-ab(a^2-b^2)(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} + (-12a^5B+33a^4Ab+104a^3b^2B-170a^2Ab^3-60ab^4B+105Ab^5)$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3134

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{\int \frac{3b(a^2-b^2)^2(4Aa^2-20bBa+35Ab^2)-ab(a^2-b^2)(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} + (-12a^5B+33a^4Ab+104a^3b^2B-170a^2Ab^3-60ab^4B+105Ab^5)$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3042

3.340. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \int \frac{3b(a^2-b^2)^2(4Aa^2-20bBa+35Ab^2)-ab(a^2-b^2)(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{(-12a^5B+33a^4Ab+...)}{a(a^2-b^2)}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3132

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \int \frac{3b(a^2-b^2)^2(4Aa^2-20bBa+35Ab^2)-ab(a^2-b^2)(-12Ba^3+27Aba^2+20b^2Ba-35Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(-12a^5B+33a^4Ab+...)}{a(a^2-b^2)}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3481

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{3b(a^2-b^2)^2(4a^2A-20abB+35Ab^2) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3042

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{3b(a^2-b^2)^2(4a^2A-20abB+35Ab^2) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3142

3.340. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)\sqrt{\sin(c+dx+\frac{\pi}{2})}}{3b(a^2-b^2)^2(4a^2A-20abB+35Ab^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)}{b \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3042

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)\sqrt{\sin(c+dx+\frac{\pi}{2})}}{3b(a^2-b^2)^2(4a^2A-20abB+35Ab^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)}{b \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3140

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{2ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)\sqrt{\sin(c+dx+\frac{\pi}{2})}}{3b(a^2-b^2)^2(4a^2A-20abB+35Ab^2) \int \frac{1}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3286

$$\frac{A \tan(c+dx) \sec(c+dx)}{2ad(a+b \cos(c+dx))^{3/2}} - \frac{2ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}{3b(a^2-b^2)^2(4a^2A-20abB+35Ab^2) \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx - \frac{2ab(a^2-b^2)(-12a^3B+27a^2Ab+20ab^2B-35Ab^3)}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(7Ab-4aB) \tan(c+dx)}{ad(a+b \cos(c+dx))^{3/2}} -$$

↓ 3042

3.340. $\int \frac{(A+B \cos(c+dx)) \sec^3(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\frac{A \tan(c + dx) \sec(c + dx)}{2ad(a + b \cos(c + dx))^{3/2}} - \frac{3b(a^2 - b^2)^2(4a^2A - 20abB + 35Ab^2) \int \frac{1}{\sqrt{a+b \cos(c+dx)}} \frac{dx}{\sin(c+dx+\frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}}} - \frac{2ab(a^2 - b^2)(-12a^3B + 27a^2)}{b \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(7Ab - 4aB) \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} -$$

↓ 3284

$$\frac{A \tan(c + dx) \sec(c + dx)}{2ad(a + b \cos(c + dx))^{3/2}} - \frac{6b(a^2 - b^2)^2(4a^2A - 20abB + 35Ab^2) \int \frac{1}{d \sqrt{a+b \cos(c+dx)}} \text{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right) - \frac{2ab(a^2 - b^2)(-12a^3B + 27a^2Ab + 20ab^2B - 35)}{b d \sqrt{a+b \cos(c+dx)}}$$

$$\frac{(7Ab - 4aB) \tan(c + dx)}{ad(a + b \cos(c + dx))^{3/2}} -$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^3)/(a + b*Cos[c + d*x])^(5/2),x]
```

```
output (A*Sec[c + d*x]*Tan[c + d*x])/(2*a*d*(a + b*Cos[c + d*x])^(3/2)) - (-1/2*(-2*b*(27*a^2*A*b - 35*A*b^3 - 12*a^3*B + 20*a*b^2*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((2*(33*a^4*A*b - 170*a^2*A*b^3 + 105*A*b^5 - 12*a^5*B + 104*a^3*b^2*B - 60*a*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + ((-2*a*b*(a^2 - b^2)*(27*a^2*A*b - 35*A*b^3 - 12*a^3*B + 20*a*b^2*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]) + (6*b*(a^2 - b^2)^2*(4*a^2*A + 35*A*b^2 - 20*a*b*B)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]))/b)/(a*(a^2 - b^2)) - (2*b*(33*a^4*A*b - 170*a^2*A*b^3 + 105*A*b^5 - 12*a^5*B + 104*a^3*b^2*B - 60*a*b^4*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])/(3*a*(a^2 - b^2))/a + ((7*A*b - 4*a*B)*Tan[c + d*x])/(a*d*(a + b*Cos[c + d*x])^(3/2))/(4*a)
```

3.340.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(- (A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])]), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
] && NeQ[c^2 - d^2, 0]
```

3.340.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2003 vs. 2(585) = 1170.

Time = 40.59 (sec) , antiderivative size = 2004, normalized size of antiderivative = 3.77

method	result	size
default	Expression too large to display	2004
parts	Expression too large to display	3298

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNV
ERBOSE)
```

output
$$\begin{aligned} & -(-(-2*b*\cos(1/2*d*x+1/2*c)^2-a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*A/a^2*(- \\ & 1/2*\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2* \\ & c)^2)^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1)^2+3/4*b/a^2*\cos(1/2*d*x+1/2*c)*(-2* \\ & \sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(2*\cos(1/2*d*x+1/ \\ & 2*c)^2-1)-1/8*b/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+ \\ & a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(\\ & 1/2)*\text{EllipticF}(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})+3/8/a*(\sin(1/2*d*x+1 \\ & /2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d* \\ & x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*\text{EllipticE}(\cos(1/2*d*x+1/2 \\ & *c),(-2*b/(a-b))^{(1/2)})-3/8*b^2/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos \\ & (1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1 \\ & /2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})-1/ \\ & 2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2 \\ &)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticPi} \\ & (\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})-3/8/a^2*(\sin(1/2*d*x+1/2*c)^2)^{(1 \\ & /2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4* \\ & b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),2,(-2*b/ \\ & (a-b))^{(1/2)})*b^2+2*(-2*A*b+B*a)/a^3*(-\cos(1/2*d*x+1/2*c)/a*(-2*\sin(1/2*d \\ & *x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1) \\ & +1/2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b)} \dots \end{aligned}$$

3.340.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(5/2),x, algorithm="fracas")`

output `Timed out`

3.340.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**3/(a+b*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sec(c + d*x)**3/(a + b*cos(c + d*x))**(5/2), x)`

3.340.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(5/2),x, algorithm m="maxima")`

output `Timed out`

3.340.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^3}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^3/(a+b*cos(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^3/(b*cos(d*x + c) + a)^(5/2), x)`

3.340.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^3(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^3 (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^(5/2)),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^3*(a + b*cos(c + d*x))^(5/2)), x)`

3.341 $\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

3.341.1 Optimal result	3204
3.341.2 Mathematica [A] (verified)	3204
3.341.3 Rubi [A] (verified)	3205
3.341.4 Maple [C] (verified)	3206
3.341.5 Fricas [C] (verification not implemented)	3207
3.341.6 Sympy [F]	3207
3.341.7 Maxima [F]	3208
3.341.8 Giac [F]	3208
3.341.9 Mupad [F(-1)]	3208

3.341.1 Optimal result

Integrand size = 28, antiderivative size = 58

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+
1/2*c), 2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(
d*x+c))^(1/2)
```

3.341.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

```
input Integrate[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(3/2), x]
```

```
output (2*B*sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a +
b)])/(d*sqrt[a + b*Cos[c + d*x]])
```

3.341.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {2011, 3042, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3142} \\
 & \frac{B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{\sqrt{a + b \cos(c + dx)}} \\
 & \quad \downarrow \text{3140} \\
 & \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}
 \end{aligned}$$

input `Int[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `(2*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])`

3.341.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3140 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*
Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[
{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3142 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a
+ b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b)
+ (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && !GtQ[a + b, 0]
```

3.341.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 6.01 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.36

method	result
default	$\frac{2B\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-a-b}{a+b}}}{d\sqrt{2b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a-b}} \operatorname{am}^{-1}\left(\frac{dx}{2}+\frac{c}{2}, \frac{\sqrt{2}\sqrt{b}}{\sqrt{a+b}}\right)$
parts	$\frac{2Ba\left(2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b}{a-b}}\right)\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a-b}}+\frac{a+b}{a-b}\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)a-E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a-b}}\right)}{(a-b)(a+b)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-2b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+a+b}d$

```
input int((B*a+b*B*cos(d*x+c))/(a*cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)
```

3.341. $\int \frac{aB+bB\cos(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx$

output $2*B/d/(2*b*\cos(1/2*d*x+1/2*c)^2+a-b)^{(1/2)}*(-(2*b*\sin(1/2*d*x+1/2*c)^2-a-b)/(a+b))^{(1/2)}*\text{InverseJacobiAM}(1/2*d*x+1/2*c,2^{(1/2)}/(a+b)^{(1/2)}*b^{(1/2)})$

3.341.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 148, normalized size of antiderivative = 2.55

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{-i \sqrt{2} B \sqrt{b} \text{weierstrassPInverse}\left(\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8a^3 - 9ab^2)}{27b^3}, \frac{3b \cos(dx+c) + 3i b \sin(dx-c)}{3b}\right)}{1}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output $(-I*\text{sqrt}(2)*B*\text{sqrt}(b)*\text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) + 3*I*b*\sin(d*x + c) + 2*a)/b) + I*\text{sqrt}(2)*B*\text{sqrt}(b)*\text{weierstrassPInverse}(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*\cos(d*x + c) - 3*I*b*\sin(d*x + c) + 2*a)/b))/ (b*d)$

3.341.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)`

output `B*Integral(1/sqrt(a + b*cos(c + d*x)), x)`

3.341.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/(b*cos(d*x + c) + a)^(3/2), x)`

3.341.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/(b*cos(d*x + c) + a)^(3/2), x)`

3.341.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Ba + Bb \cos(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/(a + b*cos(c + d*x))^(3/2),x)`

output `int((B*a + B*b*cos(c + d*x))/(a + b*cos(c + d*x))^(3/2), x)`

3.342
$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$$

3.342.1 Optimal result 3209
 3.342.2 Mathematica [A] (verified) 3209
 3.342.3 Rubi [A] (verified) 3210
 3.342.4 Maple [A] (verified) 3211
 3.342.5 Fricas [F(-1)] 3212
 3.342.6 Sympy [F] 3212
 3.342.7 Maxima [F] 3212
 3.342.8 Giac [F] 3213
 3.342.9 Mupad [F(-1)] 3213

3.342.1 Optimal result

Integrand size = 34, antiderivative size = 59

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/d/(a+b*cos(d*x+c))^(1/2)`

3.342.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{d \sqrt{a + b \cos(c + dx)}}$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `(2*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])`

3.342.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {2011, 3042, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\sec(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3286} \\
 & \frac{B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b}+\frac{b\cos(c+dx)}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx+\frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b\cos(c+dx)}} \\
 & \quad \downarrow \text{3284} \\
 & \frac{2B\sqrt{\frac{a+b\cos(c+dx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{d\sqrt{a+b\cos(c+dx)}}
 \end{aligned}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(3/2),x]`

output `(2*B*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]])`

3.342.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c,
d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Sin[e + f*x])/(c + d)]/Sqrt
[c + d*Sin[e + f*x]] Int[1/((a + b*Sin[e + f*x])*Sqrt[c/(c + d) + (d/(c +
d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

3.342.4 Maple [A] (verified)

Time = 7.44 (sec) , antiderivative size = 167, normalized size of antiderivative = 2.83

method	result
default	$\frac{2B\sqrt{\left(2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (a+b)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{\frac{2b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a - b}{a - b}} \Pi\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2, \sqrt{-\frac{2b}{a - b}}\right)}{\sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + a + b} d}$
parts	$\frac{2B\left(2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^2 + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-\frac{2b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a - b}} + \frac{a+b}{a-b} \Pi\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2, \sqrt{-\frac{2b}{a - b}}\right)a^2 - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right)}{\dots}$

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)/(a*cos(d*x+c)*b)^(3/2), x, method=_RETUR
NVERBOSE)`

$$3.342. \int \frac{(aB+bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

output $2*B*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*((2*b*\cos(1/2*d*x+1/2*c)^2+a-b)/(a-b))^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4*b+(a+b)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})/\sin(1/2*d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2+a+b)^{(1/2)}/d$

3.342.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output Timed out

3.342.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = B \int \frac{\sec(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**(3/2),x)`

output `B*Integral(sec(c + d*x)/sqrt(a + b*cos(c + d*x)), x)`

3.342.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)/(b*cos(d*x + c) + a)^(3/2), x)`

3.342.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)/(b*cos(d*x + c) + a)^(3/2), x)`

3.342.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx) (a + b \cos(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(3/2)), x)`

3.343 $\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

3.343.1 Optimal result 3214
 3.343.2 Mathematica [A] (verified) 3214
 3.343.3 Rubi [A] (verified) 3215
 3.343.4 Maple [A] (verified) 3217
 3.343.5 Fricas [C] (verification not implemented) 3218
 3.343.6 Sympy [F] 3218
 3.343.7 Maxima [F] 3219
 3.343.8 Giac [F] 3219
 3.343.9 Mupad [F(-1)] 3219

3.343.1 Optimal result

Integrand size = 28, antiderivative size = 108

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2B \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2bB \sin(c + dx)}{(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

output `-2*b*B*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+2*B*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b)))^(1/2)*(a+b*cos(d*x+c))^(1/2)/(a^2-b^2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)`

3.343.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.78

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{B \left(2(a + b) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right) - 2b \sin(c + dx) \right)}{(a - b)(a + b)d \sqrt{a + b \cos(c + dx)}}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(5/2),x]`

```
output (B*(2*(a + b)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticE[(c + d*x)/2, (2
*b)/(a + b)] - 2*b*Sin[c + d*x])/((a - b)*(a + b)*d*Sqrt[a + b*Cos[c + d*
x]])
```

3.343.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2011, 3042, 3143, 27, 3042, 3134, 3042, 3132}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{(a + b \cos(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{3143} \\
 & B \left(-\frac{2 \int -\frac{1}{2} \sqrt{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \right) \\
 & \quad \downarrow \text{27} \\
 & B \left(\frac{\int \sqrt{a + b \cos(c + dx)} dx}{a^2 - b^2} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{\int \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} dx}{a^2 - b^2} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \right) \\
 & \quad \downarrow \text{3134} \\
 & B \left(\frac{\sqrt{a + b \cos(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c + dx)}{d(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & B \left(\frac{\sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx+\frac{\pi}{2})}{a+b}} dx}{(a^2-b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
 & \downarrow 3132 \\
 & B \left(\frac{2\sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{d(a^2-b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} - \frac{2b \sin(c+dx)}{d(a^2-b^2) \sqrt{a+b \cos(c+dx)}} \right)
 \end{aligned}$$

input `Int[(a*B + b*B*Cos[c + d*x])/(a + b*Cos[c + d*x])^(5/2), x]`

output `B*((2*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)])/((a^2 - b^2)*d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) - (2*b*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]])`

3.343.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

```
rule 3134 Int[Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

```
rule 3143 Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

3.343.4 Maple [A] (verified)

Time = 13.90 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.02

method	result
default	$\frac{2B \left(2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) b + E \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-\frac{2b}{a-b}} \right) \sqrt{-\frac{2b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{a-b} + \frac{a+b}{a-b} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}} a - E \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right) \right)}{(a-b)(a+b) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + a + b d}}$
parts	Expression too large to display

```
input int((B*a+b*B*cos(d*x+c))/(a*cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2*B*(2*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b+EllipticE(cos(1/2*d*x+1/
2*c),(-2*b/(a-b))^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/
2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*a-EllipticE(cos(1/2*d*x+1/2*c),(-2*b/(a-b)
)^(1/2))*(-2*b/(a-b)*sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^(1/2)*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*b)/(a-b)/(a+b)/sin(1/2*d*x+1/2*c)/(-2*b*sin(1/2*d*x+1/2*c)
^2+a+b)^(1/2)/d
```


3.343.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 491, normalized size of antiderivative = 4.55

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx =$$

$$6 \sqrt{b \cos(dx + c) + a} B b^2 \sin(dx + c) + (i \sqrt{2} B a b \cos(dx + c) + i \sqrt{2} B a^2) \sqrt{b} \text{weierstrassPInverse} \left(\frac{4(4a^2 - 3b^2)}{3b^2} \right)$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `-1/3*(6*sqrt(b*cos(d*x + c) + a)*B*b^2*sin(d*x + c) + (I*sqrt(2)*B*a*b*cos(d*x + c) + I*sqrt(2)*B*a^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b) + (-I*sqrt(2)*B*a*b*cos(d*x + c) - I*sqrt(2)*B*a^2)*sqrt(b)*weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b) - 3*(I*sqrt(2)*B*b^2*cos(d*x + c) + I*sqrt(2)*B*a*b)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*a)/b)) - 3*(-I*sqrt(2)*B*b^2*cos(d*x + c) - I*sqrt(2)*B*a*b)*sqrt(b)*weierstrassZeta(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, weierstrassPInverse(4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*a^3 - 9*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) + 2*a)/b)))/((a^2*b^2 - b^4)*d*cos(d*x + c) + (a^3*b - a*b^3)*d)`

3.343.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = B \int \frac{1}{a \sqrt{a + b \cos(c + dx)} + b \sqrt{a + b \cos(c + dx)} \cos(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `B*Integral(1/(a*sqrt(a + b*cos(c + d*x)) + b*sqrt(a + b*cos(c + d*x))*cos(c + d*x)), x)`

3.343. $\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

3.343.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/(b*cos(d*x + c) + a)^(5/2), x)`

3.343.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/(b*cos(d*x + c) + a)^(5/2), x)`

3.343.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{Ba + Bb \cos(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((B*a + B*b*cos(c + d*x))/(a + b*cos(c + d*x))^(5/2), x)`

$$3.344 \quad \int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$$

3.344.1 Optimal result 3220
 3.344.2 Mathematica [C] (verified) 3220
 3.344.3 Rubi [A] (verified) 3221
 3.344.4 Maple [A] (verified) 3226
 3.344.5 Fracas [F(-1)] 3226
 3.344.6 Sympy [F] 3227
 3.344.7 Maxima [F] 3227
 3.344.8 Giac [F] 3227
 3.344.9 Mupad [F(-1)] 3228

3.344.1 Optimal result

Integrand size = 34, antiderivative size = 179

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = -\frac{2bB \sqrt{a + b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| \frac{2b}{a+b}\right)}{a(a^2 - b^2) d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2B \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticPi}\left(2, \frac{1}{2}(c + dx), \frac{2b}{a+b}\right)}{ad \sqrt{a + b \cos(c + dx)}} + \frac{2b^2 B \sin(c + dx)}{a(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

output

```
2*b^2*B*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)-2*b*B*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2)*(b/(a+b))^(1/2))*(a+b*cos(d*x+c))^(1/2)/a/(a^2-b^2)/d/((a+b*cos(d*x+c))/(a+b))^(1/2)+2*B*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2,2^(1/2)*(b/(a+b))^(1/2))*((a+b*cos(d*x+c))/(a+b))^(1/2)/a/d/(a+b*cos(d*x+c))^(1/2)
```

3.344.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.41 (sec) , antiderivative size = 403, normalized size of antiderivative = 2.25

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = B \left(-\frac{4ab \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} + \frac{2(2a^2 - 3b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \text{EllipticE}\left(\frac{1}{2}(c+dx) \middle| \frac{2b}{a+b}\right)}{\sqrt{a+b \cos(c+dx)}} \right)$$

3.344. $\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(5/2),x]`

output `(B*(-(((4*a*b*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticF[(c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] + (2*(2*a^2 - 3*b^2)*Sqrt[(a + b*Cos[c + d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/Sqrt[a + b*Cos[c + d*x]] - ((2*I)*Sqrt[-(b*(-1 + Cos[c + d*x]))/(a + b)]*Sqrt[(b*(1 + Cos[c + d*x]))/(-a + b)]*Csc[c + d*x]*(-2*a*(a - b)*EllipticE[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*(-2*a*EllipticF[I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)] + b*EllipticPi[(a + b)/a, I*ArcSinh[Sqrt[-(a + b)^(-1)]*Sqrt[a + b*Cos[c + d*x]]], (a + b)/(a - b)])))/(a*Sqrt[-(a + b)^(-1)]))/((-a + b)*(a + b)) + (4*b^2*Sin[c + d*x])/((a^2 - b^2)*Sqrt[a + b*Cos[c + d*x]]))/(2*a*d)`

3.344.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {2011, 3042, 3281, 27, 3042, 3538, 25, 27, 3042, 3134, 3042, 3132, 3286, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\sec(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{3281} \\
 & B \left(\frac{2 \int \frac{(a^2-b\cos(c+dx)a-b^2-b^2\cos^2(c+dx))\sec(c+dx)}{2\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b^2\sin(c+dx)}{ad(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
& B \left(\frac{\int \frac{(a^2 - b \cos(c+dx))a - b^2 - b^2 \cos^2(c+dx) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{\int \frac{a^2 - b \sin(c+dx + \frac{\pi}{2})a - b^2 - b^2 \sin^2(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 3538 \\
& B \left(\frac{-\int \frac{b(a^2 - b^2) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} - b \int \sqrt{a+b \cos(c+dx)} dx}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 25 \\
& B \left(\frac{\int \frac{b(a^2 - b^2) \sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx}{b} - b \int \sqrt{a+b \cos(c+dx)} dx}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 27 \\
& B \left(\frac{(a^2 - b^2) \int \frac{\sec(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx - b \int \sqrt{a+b \cos(c+dx)} dx}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{(a^2 - b^2) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - b \int \sqrt{a+b \sin(c+dx + \frac{\pi}{2})} dx}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 3134 \\
& B \left(\frac{(a^2 - b^2) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{b \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}}}{a(a^2 - b^2)} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right) \\
& \quad \downarrow 3042
\end{aligned}$$

3.344. $\int \frac{(aB + bB \cos(c+dx)) \sec(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$B \left(\frac{(a^2 - b^2) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{b \sqrt{a+b \cos(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}} dx}{\sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right)$$

↓ 3132

$$B \left(\frac{(a^2 - b^2) \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{2b \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right)$$

↓ 3286

$$B \left(\frac{(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{\sec(c+dx)}{\sqrt{\frac{a}{a+b} + \frac{b \cos(c+dx)}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} - \frac{2b \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right)$$

↓ 3042

$$B \left(\frac{(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \int \frac{1}{\sin(c+dx + \frac{\pi}{2}) \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx + \frac{\pi}{2})}{a+b}}} dx}{\sqrt{a+b \cos(c+dx)}} - \frac{2b \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} + \frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} \right)$$

↓ 3284

$$B \left(\frac{2b^2 \sin(c+dx)}{ad(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} + \frac{2(a^2 - b^2) \sqrt{\frac{a+b \cos(c+dx)}{a+b}} \operatorname{EllipticPi}\left(2, \frac{1}{2}(c+dx), \frac{2b}{a+b}\right)}{a(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}} - \frac{2b \sqrt{a+b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \cos(c+dx)}{a+b}}} \right)$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x])/(a + b*Cos[c + d*x])^(5/2), x]`

```
output B*((( -2*b*Sqrt[a + b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, (2*b)/(a + b)]/
(d*Sqrt[(a + b*Cos[c + d*x])/(a + b)]) + (2*(a^2 - b^2)*Sqrt[(a + b*Cos[c
+ d*x])/(a + b)]*EllipticPi[2, (c + d*x)/2, (2*b)/(a + b)]/(d*Sqrt[a + b*
Cos[c + d*x]])))/(a*(a^2 - b^2)) + (2*b^2*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sq
rt[a + b*Cos[c + d*x]]))
```

3.344.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && ( !IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3132 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

```
rule 3134 Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

rule 3281 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) + b^2*d*(m + n + 2) - (b^2*c + b*(b*c - a*d)*(m + 1))*Sin[e + f*x] - b^2*d*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && IntegersQ[2*m, 2*n] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3286 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[Sqrt[(c + d*Ssin[e + f*x])/(c + d)]/Sqrt[c + d*Ssin[e + f*x]] Int[1/((a + b*Ssin[e + f*x])*Sqrt[c/(c + d) + (d/(c + d))*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !GtQ[c + d, 0]`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Ssin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Ssin[e + f*x])*(c + d*Ssin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.344.4 Maple [A] (verified)

Time = 16.67 (sec) , antiderivative size = 377, normalized size of antiderivative = 2.11

method	result
default	$2B \left(2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b^2 + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-\frac{2b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a-b} + \frac{a+b}{a-b}} \Pi\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2, \sqrt{-\frac{2b}{a-b}}\right) a^2 - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)$
parts	Expression too large to display

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

output
$$2*B*(2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*b^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})*a^2-(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),2,(-2*b/(a-b))^{(1/2)})*b^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)*b*\text{EllipticE}(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})*a-b^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(-2*b/(a-b)*\sin(1/2*d*x+1/2*c)^2+(a+b)/(a-b))^{(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),(-2*b/(a-b))^{(1/2)})})/a/(a-b)/(a+b)/\sin(1/2*d*x+1/2*c)/(-2*b*\sin(1/2*d*x+1/2*c)^2+a+b)^{(1/2)/d}$$

3.344.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(5/2),x,algorithm="fricas")`

output Timed out

3.344.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = B \int \frac{\sec(c + dx)}{a\sqrt{a + b \cos(c + dx)} + b\sqrt{a + b \cos(c + dx)} \cos(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))**(5/2),x)`

output `B*Integral(sec(c + d*x)/(a*sqrt(a + b*cos(c + d*x)) + b*sqrt(a + b*cos(c + d*x))*cos(c + d*x)), x)`

3.344.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)/(b*cos(d*x + c) + a)^(5/2), x)`

3.344.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)/(a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)/(b*cos(d*x + c) + a)^(5/2), x)`

3.344.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx) (a + b \cos(c + dx))^{5/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(5/2)),x)`output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)*(a + b*cos(c + d*x))^(5/2)), x)`

3.345 $\int \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$

3.345.1 Optimal result	3229
3.345.2 Mathematica [A] (verified)	3230
3.345.3 Rubi [A] (verified)	3230
3.345.4 Maple [B] (verified)	3234
3.345.5 Fricas [C] (verification not implemented)	3234
3.345.6 Sympy [F(-1)]	3235
3.345.7 Maxima [F]	3235
3.345.8 Giac [F]	3236
3.345.9 Mupad [B] (verification not implemented)	3236

3.345.1 Optimal result

Integrand size = 31, antiderivative size = 170

$$\int \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

$$= \frac{2(9aA+7bB)E(\frac{1}{2}(c+dx)|2)}{15d} + \frac{10(Ab+aB) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{21d}$$

$$+ \frac{10(Ab+aB)\sqrt{\cos(c+dx)} \sin(c+dx)}{21d} + \frac{2(9aA+7bB) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{45d}$$

$$+ \frac{2(Ab+aB) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{7d} + \frac{2bB \cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{9d}$$

```
output 2/15*(9*A*a+7*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellipti
cE(sin(1/2*d*x+1/2*c),2^(1/2))/d+10/21*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1
/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/45*(9*A*a
+7*B*b)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/7*(A*b+B*a)*cos(d*x+c)^(5/2)*sin(d
*x+c)/d+2/9*b*B*cos(d*x+c)^(7/2)*sin(d*x+c)/d+10/21*(A*b+B*a)*sin(d*x+c)*c
os(d*x+c)^(1/2)/d
```

3.345.2 Mathematica [A] (verified)

Time = 1.72 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.74

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{84(9aA + 7bB)E\left(\frac{1}{2}(c + dx) \mid 2\right) + 300(Ab + aB) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)}(7(36aA + 43bB) \cos(c + dx) + 5(78A^2b + 78a^2B + 18(Ab + aB)\cos(2(c + dx)) + 7bB\cos(3(c + dx)))) \sin(c + dx)}{630d}$$

input `Integrate[Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(84*(9*a*A + 7*b*B)*EllipticE[(c + d*x)/2, 2] + 300*(A*b + a*B)*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(7*(36*a*A + 43*b*B)*Cos[c + d*x] + 5*(78*A*b + 78*a*B + 18*(A*b + a*B)*Cos[2*(c + d*x)] + 7*b*B*Cos[3*(c + d*x)]))*Sin[c + d*x])/(630*d)`

3.345.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.95, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3447}$$

$$\int \cos^{\frac{5}{2}}(c + dx) \left((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)\right) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \left((aB + Ab) \sin\left(c + dx + \frac{\pi}{2}\right) + aA + bB \sin\left(c + dx + \frac{\pi}{2}\right)^2\right) dx$$

$$\downarrow \text{3502}$$

3.345. $\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\frac{2}{9} \int \frac{1}{2} \cos^{\frac{5}{2}}(c+dx)(9aA+7bB+9(Ab+aB)\cos(c+dx))dx + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 27

$$\frac{1}{9} \int \cos^{\frac{5}{2}}(c+dx)(9aA+7bB+9(Ab+aB)\cos(c+dx))dx + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 3042

$$\frac{1}{9} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(9aA+7bB+9(Ab+aB)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 3227

$$\frac{1}{9} \left((9aA+7bB) \int \cos^{\frac{5}{2}}(c+dx)dx + 9(aB+Ab) \int \cos^{\frac{7}{2}}(c+dx)dx \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 3042

$$\frac{1}{9} \left((9aA+7bB) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} dx + 9(aB+Ab) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{7/2} dx \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 3115

$$\frac{1}{9} \left((9aA+7bB) \left(\frac{3}{5} \int \sqrt{\cos(c+dx)}dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + 9(aB+Ab) \left(\frac{5}{7} \int \cos^{\frac{3}{2}}(c+dx)dx + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 3042

$$\frac{1}{9} \left((9aA+7bB) \left(\frac{3}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + 9(aB+Ab) \left(\frac{5}{7} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}dx + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d}$$

↓ 3115

$$\frac{1}{9} \left(9(aB + Ab) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) + (9aA$$

↓ 3042

$$\frac{1}{9} \left(9(aB + Ab) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) +$$

↓ 3119

$$\frac{1}{9} \left(9(aB + Ab) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) +$$

↓ 3120

$$\frac{1}{9} \left(9(aB + Ab) \left(\frac{2 \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} + \frac{5}{7} \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3d} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{9d} \right) +$$

input `Int[Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(2*b*B*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(9*d) + ((9*a*A + 7*b*B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)) + 9*(A*b + a*B)*((2*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(7*d) + (5*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/7)/9`

3.345.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.345.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(202) = 404$.

Time = 17.60 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.65

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(-1120B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+(720Ab+720Ba+2240Bb)\left(\sin^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+48\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}$
parts	$\frac{2(Ab+Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+48\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+48\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}$

```
input int(cos(d*x+c)^(5/2)*(a+cos(d*x+c)*b)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output -2/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b+(720*A*b+720*B*a+2240*B*b)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-504*A*a-1080*A*b-1080*B*a-2072*B*b)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(504*A*a+840*A*b+840*B*a+952*B*b)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-126*A*a-240*A*b-240*B*a-168*B*b)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+75*A*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-189*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a+75*B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-147*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.345.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.24

$$\int \cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{2(35Bb\cos(dx+c)^3+45(Ba+Ab)\cos(dx+c)^2+75Ba+75Ab+7(9Aa+7Bb)\cos(dx+c))\sqrt{\cos(dx+c)}}{21}$$

input `integrate(cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `1/315*(2*(35*B*b*cos(d*x + c)^3 + 45*(B*a + A*b)*cos(d*x + c)^2 + 75*B*a + 75*A*b + 7*(9*A*a + 7*B*b)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 75*sqrt(2)*(I*B*a + I*A*b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 75*sqrt(2)*(-I*B*a - I*A*b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*sqrt(2)*(-9*I*A*a - 7*I*B*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*sqrt(2)*(9*I*A*a + 7*I*B*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.345.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.345.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*cos(d*x + c)^(5/2), x)`

3.345. $\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$

3.345.8 Giac [F]

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*cos(d*x + c)^(5/2), x)`

3.345.9 Mupad [B] (verification not implemented)

Time = 1.51 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.04

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= -\frac{2 A a \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 A b \cos(c + dx)^{9/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c + dx)^2\right)}{9 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 B a \cos(c + dx)^{9/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c + dx)^2\right)}{9 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 B b \cos(c + dx)^{11/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c + dx)^2\right)}{11 d \sqrt{\sin(c + dx)^2}}$$

input `int(cos(c + d*x)^(5/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x)),x)`

output `- (2*A*a*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*A*b*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) - (2*B*b*cos(c + d*x)^(11/2)*sin(c + d*x)*hypergeom([1/2, 11/4], 15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2))`

3.345. $\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$

$$\mathbf{3.346} \quad \int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

3.346.1 Optimal result	3237
3.346.2 Mathematica [A] (verified)	3238
3.346.3 Rubi [A] (verified)	3238
3.346.4 Maple [B] (verified)	3241
3.346.5 Fricas [C] (verification not implemented)	3242
3.346.6 Sympy [F(-1)]	3243
3.346.7 Maxima [F]	3243
3.346.8 Giac [F]	3243
3.346.9 Mupad [B] (verification not implemented)	3244

3.346.1 Optimal result

Integrand size = 31, antiderivative size = 140

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx \\ &= \frac{6(Ab+aB)E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d} + \frac{2(7aA+5bB) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d} \\ &+ \frac{2(7aA+5bB)\sqrt{\cos(c+dx)} \sin(c+dx)}{21d} \\ &+ \frac{2(Ab+aB) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{5d} + \frac{2bB \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{7d} \end{aligned}$$

output `6/5*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/21*(7*A*a+5*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*(A*b+B*a)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/7*b*B*cos(d*x+c)^(5/2)*sin(d*x+c)/d+2/21*(7*A*a+5*B*b)*sin(d*x+c)*cos(d*x+c)^(1/2)/d`

$$3.346. \quad \int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))(A+B \cos(c+dx)) dx$$

3.346.2 Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.74

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{126(Ab+aB)E\left(\frac{1}{2}(c+dx)\middle|2\right) + 10(7aA+5bB)\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right) + \sqrt{\cos(c+dx)}(70aA+65bB)}{105d}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(126*(A*b + a*B)*EllipticE[(c + d*x)/2, 2] + 10*(7*a*A + 5*b*B)*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(70*a*A + 65*b*B + 42*(A*b + a*B)*Cos[c + d*x] + 15*b*B*Cos[2*(c + d*x)])*Sin[c + d*x]/(105*d)`

3.346.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.96, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx$$

$$\downarrow \text{3447}$$

$$\int \cos^{\frac{3}{2}}(c+dx)\left((aB+Ab)\cos(c+dx)+aA+bB\cos^2(c+dx)\right)dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}\left((aB+Ab)\sin\left(c+dx+\frac{\pi}{2}\right)+aA+bB\sin\left(c+dx+\frac{\pi}{2}\right)^2\right)dx$$

$$\downarrow \text{3502}$$

$$\frac{2}{7}\int\frac{1}{2}\cos^{\frac{3}{2}}(c+dx)(7aA+5bB+7(Ab+aB)\cos(c+dx))dx+\frac{2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{7d}$$

3.346. $\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx))dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{7} \int \cos^{\frac{3}{2}}(c+dx)(7aA+5bB+7(Ab+aB)\cos(c+dx))dx + \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \\
& \downarrow 3042 \\
& \frac{1}{7} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}} \left(7aA+5bB+7(Ab+aB)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx + \\
& \quad \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \\
& \downarrow 3227 \\
& \frac{1}{7} \left((7aA+5bB) \int \cos^{\frac{3}{2}}(c+dx)dx + 7(aB+Ab) \int \cos^{\frac{5}{2}}(c+dx)dx \right) + \\
& \quad \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \\
& \downarrow 3042 \\
& \frac{1}{7} \left((7aA+5bB) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}} dx + 7(aB+Ab) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{5}{2}} dx \right) + \\
& \quad \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \\
& \downarrow 3115 \\
& \frac{1}{7} \left(7(aB+Ab) \left(\frac{3}{5} \int \sqrt{\cos(c+dx)}dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + (7aA+5bB) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}}dx + \frac{2}{3d} \int \frac{\cos^{\frac{3}{2}}(c+dx)}{\cos(c+dx)}dx \right) \right) + \\
& \quad \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \\
& \downarrow 3042 \\
& \frac{1}{7} \left(7(aB+Ab) \left(\frac{3}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + (7aA+5bB) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx)}}dx + \frac{2}{3d} \int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sin(c+dx)}dx \right) \right) + \\
& \quad \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} \\
& \downarrow 3119 \\
& \frac{1}{7} \left((7aA+5bB) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + 7(aB+Ab) \left(\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5d} \right) \right) + \\
& \quad \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d}
\end{aligned}$$

↓ 3120

$$\frac{1}{7} \left(7(aB + Ab) \left(\frac{6E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d} + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + (7aA + 5bB) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2bB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)}{7d} \right) \right)$$

input `Int[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(2*b*B*Cos[c + d*x]^(5/2)*Sin[c + d*x]/(7*d) + ((7*a*A + 5*b*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)) + 7*(A*b + a*B)*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)))/7`

3.346.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.346.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 412 vs. $2(176) = 352$.

Time = 11.27 (sec) , antiderivative size = 413, normalized size of antiderivative = 2.95

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(240B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (-168Ab - 168Ba - 360Bb)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$
parts	$-\frac{2(Ab + Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}}$

input `int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*b)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.346. \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

output
$$\begin{aligned} & -2/105*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(240*B*\cos(\\ & 1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^8*b+(-168*A*b-168*B*a-360*B*b)*\sin(1/2*d \\ & *x+1/2*c)^6*\cos(1/2*d*x+1/2*c)+(140*A*a+168*A*b+168*B*a+280*B*b)*\sin(1/2*d \\ & *x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+(-70*A*a-42*A*b-42*B*a-80*B*b)*\sin(1/2*d*x+ \\ & 1/2*c)^2*\cos(1/2*d*x+1/2*c)+35*a*A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2 \\ & *d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-63*A*(\sin(1/2 \\ & *d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d* \\ & x+1/2*c),2^{(1/2)})*b+25*B*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2 \\ & *c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-63*B*(\sin(1/2*d*x+1/2 \\ & *c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c) \\ & ,2^{(1/2)})*a)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2* \\ & d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.346.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.37

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{2(15Bb\cos(dx+c)^2+35Aa+25Bb+21(Ba+Ab)\cos(dx+c))\sqrt{\cos(dx+c)}\sin(dx+c)-5\sqrt{2}(7Aa+5Bb)\operatorname{erstrassPInverse}(-4,0,\cos(dx+c)+I\sin(dx+c))-5\sqrt{2}(-7Aa-5Bb)\operatorname{erstrassPInverse}(-4,0,\cos(dx+c)-I\sin(dx+c))-63\sqrt{2}(-I*Ba-I*Ab)\operatorname{erstrassZeta}(-4,0,\operatorname{erstrassPInverse}(-4,0,\cos(dx+c)+I\sin(dx+c)))-63\sqrt{2}(I*Ba+I*Ab)\operatorname{erstrassZeta}(-4,0,\operatorname{erstrassPInverse}(-4,0,\cos(dx+c)-I\sin(dx+c)))}{d}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & 1/105*(2*(15*B*b*\cos(d*x+c)^2+35*A*a+25*B*b+21*(B*a+A*b)*\cos(d*x \\ & +c))*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-5*\sqrt{2}*(7*I*A*a+5*I*B*b)*\operatorname{wei} \\ & \operatorname{erstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))-5*\sqrt{2}*(-7*I*A \\ & *a-5*I*B*b)*\operatorname{wei} \\ & \operatorname{erstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))- \\ & 63*\sqrt{2}*(-I*B*a-I*A*b)*\operatorname{wei} \\ & \operatorname{erstrassZeta}(-4,0,\operatorname{wei} \\ & \operatorname{erstrassPInverse}(-4, \\ & 0,\cos(d*x+c)+I*\sin(d*x+c))) \\ & -63*\sqrt{2}*(I*B*a+I*A*b)*\operatorname{wei} \\ & \operatorname{erstrassZeta}(-4,0,\operatorname{wei} \\ & \operatorname{erstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))))/ \\ & d \end{aligned}$$

3.346.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.346.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*cos(d*x + c)^(3/2), x)`

3.346.8 Giac [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*cos(d*x + c)^(3/2), x)`

3.346.9 Mupad [B] (verification not implemented)

Time = 1.27 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.19

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{2 A a \left(\sqrt{\cos(c + dx)} \sin(c + dx) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) \right)}{3 d}$$

$$- \frac{2 A b \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 B a \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 B b \cos(c + dx)^{9/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c + dx)^2\right)}{9 d \sqrt{\sin(c + dx)^2}}$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x)),x)`output `(2*A*a*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) - (2*A*b*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*b*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2))`

3.347 $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$

3.347.1 Optimal result	3245
3.347.2 Mathematica [A] (verified)	3245
3.347.3 Rubi [A] (verified)	3246
3.347.4 Maple [B] (verified)	3249
3.347.5 Fricas [C] (verification not implemented)	3250
3.347.6 Sympy [F(-1)]	3251
3.347.7 Maxima [F]	3251
3.347.8 Giac [F]	3251
3.347.9 Mupad [B] (verification not implemented)	3252

3.347.1 Optimal result

Integrand size = 31, antiderivative size = 108

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{2(5aA + 3bB)E(\frac{1}{2}(c + dx)|2)}{5d} + \frac{2(Ab + aB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2(Ab + aB)\sqrt{\cos(c + dx)} \sin(c + dx)}{3d} + \frac{2bB \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{5d}$$

```
output 2/5*(5*A*a+3*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*b*B*cos(d*
x+c)^(3/2)*sin(d*x+c)/d+2/3*(A*b+B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.347.2 Mathematica [A] (verified)

Time = 0.99 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.80

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$= \frac{2\left(3(5aA + 3bB)E\left(\frac{1}{2}(c + dx)\middle|2\right) + 5(Ab + aB) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)}(5Ab + 5aB + 3bB \cos(c + dx))\right)}{15d}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])*(A + B*cos[c + d*x]),x]`

output `(2*(3*(5*a*A + 3*b*B)*EllipticE[(c + d*x)/2, 2] + 5*(A*b + a*B)*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(5*A*b + 5*a*B + 3*b*B*cos[c + d*x])*Sin[c + d*x))/(15*d)`

3.347.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.99, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))(A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3447} \\
 & \int \sqrt{\cos(c+dx)}\left((aB+Ab)\cos(c+dx)+aA+bB\cos^2(c+dx)\right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left((aB+Ab)\sin\left(c+dx+\frac{\pi}{2}\right)+aA+bB\sin\left(c+dx+\frac{\pi}{2}\right)^2\right) dx \\
 & \quad \downarrow \text{3502} \\
 & \frac{2}{5} \int \frac{1}{2} \sqrt{\cos(c+dx)}(5aA+3bB+5(Ab+aB)\cos(c+dx))dx + \frac{2bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{5} \int \sqrt{\cos(c+dx)}(5aA+3bB+5(Ab+aB)\cos(c+dx))dx + \frac{2bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.347. $\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))(A+B\cos(c+dx)) dx$

$$\begin{aligned}
& \frac{1}{5} \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \left(5aA + 3bB + 5(Ab + aB) \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\
& \quad \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{5} \left(5(aB + Ab) \int \cos^{\frac{3}{2}}(c + dx) dx + (5aA + 3bB) \int \sqrt{\cos(c + dx)} dx\right) + \\
& \quad \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left((5aA + 3bB) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5(aB + Ab) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx\right) + \\
& \quad \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \\
& \quad \downarrow \text{3115} \\
& \frac{1}{5} \left((5aA + 3bB) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5(aB + Ab) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}\right)\right) + \\
& \quad \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left((5aA + 3bB) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5(aB + Ab) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}\right)\right) + \\
& \quad \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \\
& \quad \downarrow \text{3119} \\
& \frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}\right) + \frac{2(5aA + 3bB)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d}\right) + \\
& \quad \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \\
& \quad \downarrow \text{3120}
\end{aligned}$$

$$\frac{1}{5} \left(\frac{2(5aA + 3bB)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + 5(aB + Ab) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) + \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d}$$

input `Int[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]),x]`

output `(2*b*B*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d) + ((2*(5*a*A + 3*b*B)*EllipticE[(c + d*x)/2, 2])/d + 5*(A*b + a*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)))/5`

3.347.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3447 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a
+ b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2),
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.347.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 370 vs. 2(148) = 296.

Time = 9.45 (sec) , antiderivative size = 371, normalized size of antiderivative = 3.44

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(-24B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (20Ab + 20Ba + 24Bb)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}}\right)$
parts	$-\frac{2(Ab + Ba)\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}d + \sqrt{\frac{1}{2} - \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}}\right)$

```
input int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*b)*(A+B*cos(d*x+c)),x,method=_RETURNVER
BOSE)
```


output
$$\begin{aligned} & -2/15*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-24*B*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6*b+(20*A*b+20*B*a+24*B*b)*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+(-10*A*b-10*B*a-6*B*b)*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+5*A*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-15*A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})+5*B*a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-9*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})+b)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.347.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.62

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{2(3Bb\cos(dx+c)+5Ba+5Ab)\sqrt{\cos(dx+c)}\sin(dx+c)-5\sqrt{2}(iBa+iAb)\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))}{d}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="fracas")`

output
$$\begin{aligned} & 1/15*(2*(3*B*b*\cos(d*x+c)+5*B*a+5*A*b)*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-5*\sqrt{2}*(I*B*a+I*A*b)*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))-5*\sqrt{2}*(-I*B*a-I*A*b)*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))-3*\sqrt{2}*(-5*I*A*a-3*I*B*b)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c)))-3*\sqrt{2}*(5*I*A*a+3*I*B*b)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))))/d \end{aligned}$$

3.347.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.347.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)\sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.347.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)\sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.347.9 Mupad [B] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.19

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))(A+B\cos(c+dx))dx$$

$$= \frac{2Ab\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)\right)}{3d}$$

$$+ \frac{2Ba\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)\right)}{3d} + \frac{2AaE\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)}{d}$$

$$- \frac{2Bb\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}}$$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x)),x)`output `(2*A*b*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*B*a*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*A*a*ellipticE(c/2 + (d*x)/2, 2))/d - (2*B*b*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))`

3.348
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.348.1 Optimal result 3253
 3.348.2 Mathematica [A] (verified) 3253
 3.348.3 Rubi [A] (verified) 3254
 3.348.4 Maple [B] (verified) 3257
 3.348.5 Fricas [C] (verification not implemented) 3257
 3.348.6 Sympy [F] 3258
 3.348.7 Maxima [F] 3258
 3.348.8 Giac [F] 3259
 3.348.9 Mupad [B] (verification not implemented) 3259

3.348.1 Optimal result

Integrand size = 31, antiderivative size = 75

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{2(Ab + aB)E(\frac{1}{2}(c + dx)|2)}{d} + \frac{2(3aA + bB) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2bB \sqrt{\cos(c + dx)} \sin(c + dx)}{3d}$$

```
output 2*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(
1/2*d*x+1/2*c),2^(1/2))/d+2/3*(3*A*a+B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos
(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*b*B*sin(d*x+c)
*cos(d*x+c)^(1/2)/d
```

3.348.2 Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.89

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \frac{2 \left(3(Ab + aB)E\left(\frac{1}{2}(c + dx)|2\right) + (3aA + bB) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + bB \sqrt{\cos(c + dx)} \sin(c + dx) \right)}{3d}$$

input `Integrate[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x
]`

output `(2*(3*(A*b + a*B)*EllipticE[(c + d*x)/2, 2] + (3*a*A + b*B)*EllipticF[(c +
d*x)/2, 2] + b*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*d)`

3.348.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3447

$$\int \frac{(aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3502

$$\frac{2}{3} \int \frac{3aA + bB + 3(Ab + aB) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}$$

↓ 27

$$\frac{1}{3} \int \frac{3aA + bB + 3(Ab + aB) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}$$

↓ 3042

3.348. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\begin{aligned}
& \frac{1}{3} \int \frac{3aA + bB + 3(Ab + aB) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{3} \left((3aA + bB) \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{3(aB + Ab) \int \sqrt{\cos(c + dx)} dx}{2bB \sin(c + dx) \sqrt{\cos(c + dx)}} \right) + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left((3aA + bB) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{3(aB + Ab) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{2bB \sin(c + dx) \sqrt{\cos(c + dx)}} \right) + \\
& \quad \downarrow \text{3119} \\
& \frac{1}{3} \left((3aA + bB) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{6(aB + Ab) E(\frac{1}{2}(c + dx) | 2)}{d} \right) + \\
& \quad \downarrow \text{3120} \\
& \frac{1}{3} \left(\frac{2(3aA + bB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} + \frac{6(aB + Ab) E(\frac{1}{2}(c + dx) | 2)}{d} \right) + \\
& \quad \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}
\end{aligned}$$

input `Int[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `((6*(A*b + a*B)*EllipticE[(c + d*x)/2, 2])/d + (2*(3*a*A + b*B)*EllipticF[(c + d*x)/2, 2])/d)/3 + (2*b*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)`

3.348.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`
- rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.348.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 325 vs. 2(121) = 242.

Time = 7.77 (sec) , antiderivative size = 326, normalized size of antiderivative = 4.35

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(4B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+3aA\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d}$
parts	$\frac{2(Ab+Ba)\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d} - \frac{2Bb\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output
$$-2/3*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*B*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^4+b*3*a*A*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^(1/2))-3*A*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2))*b-2*B*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2+b*B*b*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^(1/2))-3*B*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2))*a)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^(1/2)/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d$$

3.348.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 156, normalized size of antiderivative = 2.08

$$\int \frac{(a+b\cos(c+dx))(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

$$= \frac{2Bb\sqrt{\cos(dx+c)}\sin(dx+c)+\sqrt{2}(-3iAa-iBb)\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))}{\sqrt{\cos(dx+c)}} + \dots$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,algorithm="fricas")`

3.348.
$$\int \frac{(a+b\cos(c+dx))(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

output `1/3*(2*B*b*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(2)*(-3*I*A*a - I*B*b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(3*I*A*a + I*B*b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(-I*B*a - I*A*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(I*B*a + I*A*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.348.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + b \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))/sqrt(cos(c + d*x)), x)`

3.348.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.348.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.348.9 Mupad [B] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.13

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2 B b \left(\sqrt{\cos(c + dx)} \sin(c + dx) + F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) \right)}{3 d}$$

$$+ \frac{2 A a F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 A b E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 B a E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^(1/2),x)`

output `(2*B*b*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/(3*d) + (2*A*a*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*b*ellipticE(c/2 + (d*x)/2, 2))/d + (2*B*a*ellipticE(c/2 + (d*x)/2, 2))/d`

3.349
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.349.1 Optimal result 3260
 3.349.2 Mathematica [A] (verified) 3260
 3.349.3 Rubi [A] (verified) 3261
 3.349.4 Maple [B] (verified) 3263
 3.349.5 Fricas [C] (verification not implemented) 3264
 3.349.6 Sympy [F(-1)] 3265
 3.349.7 Maxima [F] 3265
 3.349.8 Giac [F] 3265
 3.349.9 Mupad [B] (verification not implemented) 3266

3.349.1 Optimal result

Integrand size = 31, antiderivative size = 71

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = -\frac{2(aA - bB)E(\frac{1}{2}(c + dx) | 2)}{d} + \frac{2(Ab + aB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}}$$

```
output -2*(A*a-B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*a*A*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.349.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.90

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \frac{2\left((-aA + bB)E(\frac{1}{2}(c + dx) | 2) + (Ab + aB) \text{EllipticF}(\frac{1}{2}(c + dx), 2) + \frac{aA \sin(c + dx)}{\sqrt{\cos(c + dx)}}\right)}{d}$$

input `Integrate[((a + b*cos[c + d*x])*(A + B*cos[c + d*x]))/Cos[c + d*x]^(3/2),x
]`

output `(2*((-(a*A) + b*B)*EllipticE[(c + d*x)/2, 2] + (A*b + a*B)*EllipticF[(c +
d*x)/2, 2] + (a*A*Sin[c + d*x])/Sqrt[Cos[c + d*x]]))/d`

3.349.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3447} \\ & \int \frac{(aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)}{\cos^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3500} \\ & 2 \int \frac{Ab + aB - (aA - bB) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\ & \quad \downarrow \text{27} \\ & \int \frac{Ab + aB - (aA - bB) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.349. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{Ab + aB + (bB - aA) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3227} \\
& (aB + Ab) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - (aA - bB) \int \sqrt{\cos(c + dx)} dx + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& (aB + Ab) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx - (aA - bB) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3119} \\
& (aB + Ab) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx - \frac{2(aA - bB)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3120} \\
& \frac{2(aB + Ab) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{2(aA - bB)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + \frac{2aA \sin(c + dx)}{d\sqrt{\cos(c + dx)}}
\end{aligned}$$

input `Int[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(-2*(a*A - b*B)*EllipticE[(c + d*x)/2, 2])/d + (2*(A*b + a*B)*EllipticF[(c + d*x)/2, 2])/d + (2*a*A*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]])`

3.349.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.349.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. 2(121) = 242.

Time = 7.12 (sec) , antiderivative size = 246, normalized size of antiderivative = 3.46

method	result
default	$\frac{4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a - 2Ab \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - 2A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \frac{2(Ab+Ba) a m^{-1} \left(\frac{dx}{2} + \frac{c}{2} \sqrt{2}\right)}{d} + \frac{2Bb \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$
parts	

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2), x, method=_RETURNVERBOSE)`

$$3.349. \int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

output $2*(2*A*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*a-A*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a-B*a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b)/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$

3.349.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.61

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 A a \sqrt{\cos(dx + c)} \sin(dx + c) + \sqrt{2}(-i B a - i A b) \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c)) - \dots}{\dots}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="fracas")`

output $(2*A*a*\sqrt{\cos(d*x + c)}*\sin(d*x + c) + \sqrt{2)*(-I*B*a - I*A*b)*\cos(d*x + c)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + \sqrt{2)*(I*B*a + I*A*b)*\cos(d*x + c)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + \sqrt{2)*(-I*A*a + I*B*b)*\cos(d*x + c)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + \sqrt{2)*(I*A*a - I*B*b)*\cos(d*x + c)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)))})/(d*\cos(d*x + c))$

3.349.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`output `Timed out`**3.349.7 Maxima [F]**

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\cos(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`**3.349.8 Giac [F]**

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\cos(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="giac")`output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`

3.349.9 Mupad [B] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.35

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 A b F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{2 B a F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{2 B b E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$+ \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^(3/2),x)`output `(2*A*b*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*a*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*b*ellipticE(c/2 + (d*x)/2, 2))/d + (2*A*a*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.350
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.350.1 Optimal result 3267
 3.350.2 Mathematica [A] (verified) 3268
 3.350.3 Rubi [A] (verified) 3268
 3.350.4 Maple [B] (verified) 3271
 3.350.5 Fricas [C] (verification not implemented) 3272
 3.350.6 Sympy [F(-1)] 3273
 3.350.7 Maxima [F] 3273
 3.350.8 Giac [F] 3273
 3.350.9 Mupad [B] (verification not implemented) 3274

3.350.1 Optimal result

Integrand size = 31, antiderivative size = 103

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = -\frac{2(Ab + aB)E(\frac{1}{2}(c + dx) | 2)}{d} + \frac{2(aA + 3bB) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2(Ab + aB) \sin(c + dx)}{d \sqrt{\cos(c + dx)}}$$

```
output -2*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(A*a+3*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*A*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2*(A*b+B*a)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.350.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.04

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 \left(-3(Ab + aB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + (aA + 3bB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 3A \right)}{3d \sqrt{\cos(c + dx)}}$$

input `Integrate[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2), x]`

output `(2*(-3*(A*b + a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + (a*A + 3*b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 3*A*b*Sin[c + d*x] + 3*a*B*Sin[c + d*x] + a*A*Tan[c + d*x]))/(3*d*Sqrt[Cos[c + d*x]])`

3.350.3 Rubi [A] (verified)Time = 0.63 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.99, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{\frac{5}{2}}} dx$$

$$\downarrow \text{3447}$$

$$\int \frac{(aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^{\frac{5}{2}}} dx$$

3.350. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow \text{3500} \\
& \frac{2}{3} \int \frac{3(Ab + aB) + (aA + 3bB) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{27} \\
& \frac{1}{3} \int \frac{3(Ab + aB) + (aA + 3bB) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3(Ab + aB) + (aA + 3bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{3227} \\
& \frac{1}{3} \left(3(aB + Ab) \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx + (aA + 3bB) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \right) + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{3042} \\
& \frac{1}{3} \left(3(aB + Ab) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + (aA + 3bB) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{3116} \\
& \frac{1}{3} \left((aA + 3bB) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(aB + Ab) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right) \right) + \\
& \quad \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{3042} \\
& \frac{1}{3} \left((aA + 3bB) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(aB + Ab) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) \right) + \\
& \quad \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \\
& \downarrow \text{3119}
\end{aligned}$$

$$\frac{1}{3} \left((aA + 3bB) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(aB + Ab) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx)|2)}{d} \right) \right) + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \downarrow \text{3120}$$

$$\frac{1}{3} \left(\frac{2(aA + 3bB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} + 3(aB + Ab) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx)|2)}{d} \right) \right) + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

```
input Int[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]
```

```
output (2*a*A*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((2*(a*A + 3*b*B)*EllipticF[(c + d*x)/2, 2])/d + 3*(A*b + a*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/3
```

3.350.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3116 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(- (A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.350.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 400 vs. 2(147) = 294.

Time = 9.12 (sec) , antiderivative size = 401, normalized size of antiderivative = 3.89

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1}(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} \left(\frac{2Bb\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1} F(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2})}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} + 2aA \left(-\frac{\cos(\frac{dx}{2} + \frac{c}{2})}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} \right) \right)$
parts	$-\frac{2(Ab+Ba)\left(-2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}d}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2), x, method=_RETURNVER BOSE)`

$$3.350. \int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

output
$$\begin{aligned} & -(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*B*b*(\sin(1/2 \\ & *d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2 \\ & *c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+2* \\ & a*A*(-1/6*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2 \\ &)^{(1/2)}/(\cos(1/2*d*x+1/2*c)^2-1)^2+1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2* \\ & \cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2 \\ &)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)}))+2*(A*b+B*a)/\sin(1/2*d*x+1/ \\ & 2*c)^2/(2*\sin(1/2*d*x+1/2*c)^2-1)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2 \\ & *c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-(\sin(1/2*d*x+1/2*c \\ &)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2 \\ & ^{(1/2)}))/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.350.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.07

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{2}(-i A a - 3i B b) \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \sqrt{2}(i A a + 3i B b) \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{\cos^{\frac{5}{2}}(c + dx)}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fracas")`

output
$$\begin{aligned} & 1/3*(\sqrt{2})*(-I*A*a - 3*I*B*b)*\cos(d*x + c)^2*\operatorname{weierstrassPInverse}(-4, 0, \\ & \cos(d*x + c) + I*\sin(d*x + c)) + \sqrt{2}*(I*A*a + 3*I*B*b)*\cos(d*x + c)^2* \\ & \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*\sqrt{2}*(I*B \\ & *a + I*A*b)*\cos(d*x + c)^2*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, \\ & 0, \cos(d*x + c) + I*\sin(d*x + c))) - 3*\sqrt{2}*(-I*B*a - I*A*b)*\cos(d*x + \\ & c)^2*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin \\ & (d*x + c))) + 2*(A*a + 3*(B*a + A*b)*\cos(d*x + c))*\sqrt{\cos(d*x + c)}*\sin \\ & (d*x + c)/(d*\cos(d*x + c)^2) \end{aligned}$$

3.350.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output `Timed out`

3.350.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\cos(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`

3.350.8 Giac [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\cos(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`

3.350.9 Mupad [B] (verification not implemented)

Time = 2.38 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.46

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 B b F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 A b \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^(5/2),x)`output `(2*B*b*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2)) + (2*A*b*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) + (2*B*a*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.351
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.351.1 Optimal result 3275
 3.351.2 Mathematica [A] (verified) 3276
 3.351.3 Rubi [A] (verified) 3276
 3.351.4 Maple [B] (verified) 3279
 3.351.5 Fricas [C] (verification not implemented) 3280
 3.351.6 Sympy [F(-1)] 3281
 3.351.7 Maxima [F] 3281
 3.351.8 Giac [F] 3282
 3.351.9 Mupad [B] (verification not implemented) 3282

3.351.1 Optimal result

Integrand size = 31, antiderivative size = 140

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = -\frac{2(3aA + 5bB)E(\frac{1}{2}(c + dx) | 2)}{5d} + \frac{2(Ab + aB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} + \frac{2(Ab + aB) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2(3aA + 5bB) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

output

```
-2/5*(3*A*a+5*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*a*A*sin(d*x+c)/d/cos(d*x+c)^(5/2)+2/3*(A*b+B*a)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/5*(3*A*a+5*B*b)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.351.2 Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.96

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{-6(3aA + 5bB) \cos^{\frac{3}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(Ab + aB) \cos^{\frac{3}{2}}(c + dx) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 10A}{15d \cos^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(-6*(3*a*A + 5*b*B)*Cos[c + d*x]^(3/2)*EllipticE[(c + d*x)/2, 2] + 10*(A*b + a*B)*Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + 10*A*b*Sin[c + d*x] + 10*a*B*Sin[c + d*x] + 9*a*A*Sin[2*(c + d*x)] + 15*b*B*Sin[2*(c + d*x)] + 6*a*A*Tan[c + d*x])/(15*d*Cos[c + d*x]^(3/2))`

3.351.3 Rubi [A] (verified)Time = 0.66 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.93, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3447, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

$$\downarrow \text{3447}$$

$$\int \frac{(aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx)}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(aB + Ab) \sin(c + dx + \frac{\pi}{2}) + aA + bB \sin(c + dx + \frac{\pi}{2})^2}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

3.351. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow \text{3500} \\
& \frac{2}{5} \int \frac{5(Ab + aB) + (3aA + 5bB) \cos(c + dx)}{2 \cos^{\frac{5}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{27} \\
& \frac{1}{5} \int \frac{5(Ab + aB) + (3aA + 5bB) \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{5(Ab + aB) + (3aA + 5bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{3227} \\
& \frac{1}{5} \left(5(aB + Ab) \int \frac{1}{\cos^{\frac{5}{2}}(c + dx)} dx + (3aA + 5bB) \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx \right) + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \left(5(aB + Ab) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx + (3aA + 5bB) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \\
& \quad \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{3116} \\
& \frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + (3aA + 5bB) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} \right) \right) \\
& \quad \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{3042} \\
& \frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + (3aA + 5bB) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\sin(c + dx + \frac{\pi}{2})} \right) \right) \\
& \quad \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \\
& \downarrow \text{3119}
\end{aligned}$$

3.351. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + (3aA + 5bB) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx))}{d} \right) \right. \\ \left. \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{2 \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + (3aA + 5bB) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx))}{d} \right) \right. \\ \left. \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

input `Int[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (5*(A*b + a*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))) + (3*a*A + 5*b*B)*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])))/5`

3.351.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.351.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. 2(176) = 352.

Time = 12.94 (sec) , antiderivative size = 636, normalized size of antiderivative = 4.54

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\sin(\frac{dx}{2} + \frac{c}{2})^2 (2(\sin^2(\frac{dx}{2} + \frac{c}{2})) - 1)} \left(\frac{2Bb\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})} (2(\sin^2(\frac{dx}{2} + \frac{c}{2})) \cos(\frac{dx}{2} + \frac{c}{2}) - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}})}{2(\sin^2(\frac{dx}{2} + \frac{c}{2})) - 1} \right)$
parts	Expression too large to display

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2), x, method=_RETURNVERBOSE)`

$$3.351. \int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b/sin(1/2*
d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2)))+2/5*a*A/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*s
in(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2
*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2
*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(
1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+
1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+2*(A*b+B*a)*(-1
/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2
*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2
)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*
x+1/2*c)^2-1)^(1/2)/d

```

3.351.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.68

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{5\sqrt{2}(iBa + iAb) \cos(dx + c)^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5\sqrt{2}(-iBa$$

input

```

integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm=
"fracas")

```

output
$$\begin{aligned} & -1/15*(5*\sqrt{2}*(I*B*a + I*A*b)*\cos(d*x + c)^3*\text{weierstrassPInverse}(-4, 0, \\ & \cos(d*x + c) + I*\sin(d*x + c)) + 5*\sqrt{2}*(-I*B*a - I*A*b)*\cos(d*x + c)^3 \\ & *\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + 3*\sqrt{2}*(3 \\ & *I*A*a + 5*I*B*b)*\cos(d*x + c)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse} \\ & (-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + 3*\sqrt{2}*(-3*I*A*a - 5*I*B*b)* \\ & \cos(d*x + c)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + \\ & c) - I*\sin(d*x + c))) - 2*(3*(3*A*a + 5*B*b)*\cos(d*x + c)^2 + 3*A*a + 5*(\\ & B*a + A*b)*\cos(d*x + c))*\sqrt{\cos(d*x + c)*\sin(d*x + c)}/(d*\cos(d*x + c)^3) \end{aligned}$$

3.351.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2), x)`

output Timed out

3.351.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\ & = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\cos(dx + c)^{\frac{7}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/cos(d*x + c)^(7/2), x)`

3.351.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/cos(d*x + c)^(7/2), x)`

3.351.9 Mupad [B] (verification not implemented)

Time = 2.67 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.26

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2 A a \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right)}{5 d \cos(c + dx)^{5/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 A b \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B b \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/cos(c + d*x)^(7/2),x)`

output $(2Aa\sin(c+dx)\operatorname{hypergeom}([-5/4, 1/2], -1/4, \cos(c+dx)^2))/(5d\cos(c+dx)^{5/2}(\sin(c+dx)^2)^{1/2}) + (2Ab\sin(c+dx)\operatorname{hypergeom}([-3/4, 1/2], 1/4, \cos(c+dx)^2))/(3d\cos(c+dx)^{3/2}(\sin(c+dx)^2)^{1/2}) + (2Ba\sin(c+dx)\operatorname{hypergeom}([-3/4, 1/2], 1/4, \cos(c+dx)^2))/(3d\cos(c+dx)^{3/2}(\sin(c+dx)^2)^{1/2}) + (2Bb\sin(c+dx)\operatorname{hypergeom}([-1/4, 1/2], 3/4, \cos(c+dx)^2))/(d\cos(c+dx)^{1/2}(\sin(c+dx)^2)^{1/2})$

3.351. $\int \frac{(a+b\cos(c+dx))(A+B\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

3.352 $\int \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.352.1 Optimal result	3284
3.352.2 Mathematica [A] (verified)	3285
3.352.3 Rubi [A] (verified)	3285
3.352.4 Maple [B] (verified)	3290
3.352.5 Fricas [C] (verification not implemented)	3291
3.352.6 Sympy [F(-1)]	3291
3.352.7 Maxima [F]	3292
3.352.8 Giac [F]	3292
3.352.9 Mupad [B] (verification not implemented)	3293

3.352.1 Optimal result

Integrand size = 33, antiderivative size = 264

$$\begin{aligned} & \int \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx \\ &= \frac{2(9a^2A+7Ab^2+14abB)E(\frac{1}{2}(c+dx)|2)}{15d} \\ &+ \frac{10(9b^2B+11a(2Ab+aB))\text{EllipticF}(\frac{1}{2}(c+dx),2)}{231d} \\ &+ \frac{10(9b^2B+11a(2Ab+aB))\sqrt{\cos(c+dx)}\sin(c+dx)}{231d} \\ &+ \frac{2(9a^2A+7Ab^2+14abB)\cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{45d} \\ &+ \frac{2(9b^2B+11a(2Ab+aB))\cos^{\frac{5}{2}}(c+dx)\sin(c+dx)}{77d} \\ &+ \frac{2b(11Ab+13aB)\cos^{\frac{7}{2}}(c+dx)\sin(c+dx)}{99d} \\ &+ \frac{2bB\cos^{\frac{7}{2}}(c+dx)(a+b \cos(c+dx))\sin(c+dx)}{11d} \end{aligned}$$

output $\frac{2}{15}(9Aa^2+7Ab^2+14Bab)(\cos(\frac{1}{2}dx+\frac{1}{2}c))^{\frac{1}{2}}/\cos(\frac{1}{2}dx+\frac{1}{2}c)*\text{EllipticE}(\sin(\frac{1}{2}dx+\frac{1}{2}c),2^{\frac{1}{2}})/d+10/231(9b^2B+11a(2Ab+Ba))*(\cos(\frac{1}{2}dx+\frac{1}{2}c))^{\frac{1}{2}}/\cos(\frac{1}{2}dx+\frac{1}{2}c)*\text{EllipticF}(\sin(\frac{1}{2}dx+\frac{1}{2}c),2^{\frac{1}{2}})/d+2/45(9Aa^2+7Ab^2+14Bab)\cos(dx+c)^{\frac{3}{2}}\sin(dx+c)/d+2/77(9b^2B+11a(2Ab+Ba))\cos(dx+c)^{\frac{5}{2}}\sin(dx+c)/d+2/99*b*(11Ab+13Ba)\cos(dx+c)^{\frac{7}{2}}\sin(dx+c)/d+2/11*b*B\cos(dx+c)^{\frac{7}{2}}*(a+b\cos(dx+c))*\sin(dx+c)/d+10/231(9b^2B+11a(2Ab+Ba))*\sin(dx+c)*\cos(dx+c)^{\frac{1}{2}}/d$

3.352.2 Mathematica [A] (verified)

Time = 2.33 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.74

$$\int \cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{3696(9a^2A+7Ab^2+14abB)E(\frac{1}{2}(c+dx)|2)+1200(22aAb+11a^2B+9b^2B)\text{EllipticF}(\frac{1}{2}(c+dx),2)+\dots}{\dots}$$

input `Integrate[Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output $(3696(9a^2A+7Ab^2+14abB)*\text{EllipticE}[(c+dx)/2,2]+1200(22aAb+11a^2B+9b^2B)*\text{EllipticF}[(c+dx)/2,2]+2*\text{Sqrt}[\text{Cos}[c+d*x]])*(154(36a^2A+43Ab^2+86abB)*\text{Cos}[c+d*x]+180(22aAb+11a^2B+16b^2B)*\text{Cos}[2*(c+d*x)]+770*b*(Ab+2aB)*\text{Cos}[3*(c+d*x)]+15(1144aAb+572a^2B+531b^2B+21b^2B*\text{Cos}[4*(c+d*x)]))*\text{Sin}[c+d*x]/(27720*d)$

3.352.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.88, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

3.352. $\int \cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$

$$\begin{aligned}
& \int \sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{2}{11} \cos^{\frac{5}{2}}(c + dx) \left((b(11Ab + 13aB) \cos^2(c + dx) + (9Bb^2 + 11a(2Ab + aB)) \cos(c + dx) + a(11aA + 7bB)) \right. \\
& \quad \left. + \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) dx + \\
& \quad \downarrow \text{3469} \\
& \int \frac{1}{11} \cos^{\frac{5}{2}}(c + dx) \left((b(11Ab + 13aB) \cos^2(c + dx) + (9Bb^2 + 11a(2Ab + aB)) \cos(c + dx) + a(11aA + 7bB)) \right. \\
& \quad \left. + \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) dx + \\
& \quad \downarrow \text{27} \\
& \int \frac{1}{11} \cos^{\frac{5}{2}}(c + dx) \left((b(11Ab + 13aB) \cos^2(c + dx) + (9Bb^2 + 11a(2Ab + aB)) \cos(c + dx) + a(11aA + 7bB)) \right. \\
& \quad \left. + \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{11} \int \sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left((b(11Ab + 13aB) \sin\left(c + dx + \frac{\pi}{2}\right)^2 + (9Bb^2 + 11a(2Ab + aB)) \sin\left(c + dx + \frac{\pi}{2}\right) + a(11aA + 7bB)) \right. \\
& \quad \left. + \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) dx + \\
& \quad \downarrow \text{3502} \\
& \frac{1}{11} \left(\frac{2}{9} \int \frac{1}{2} \cos^{\frac{5}{2}}(c + dx) (11(9Aa^2 + 14bBa + 7Ab^2) + 9(9Bb^2 + 11a(2Ab + aB)) \cos(c + dx)) dx + \frac{2b(13aB + 11aA)}{11d} \right. \\
& \quad \left. + \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{11} \left(\frac{1}{9} \int \cos^{\frac{5}{2}}(c + dx) (11(9Aa^2 + 14bBa + 7Ab^2) + 9(9Bb^2 + 11a(2Ab + aB)) \cos(c + dx)) dx + \frac{2b(13aB + 11aA)}{11d} \right. \\
& \quad \left. + \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) dx + \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.352. $\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\frac{1}{11} \left(\frac{1}{9} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} \left(11(9Aa^2 + 14bBa + 7Ab^2) + 9(9Bb^2 + 11a(2Ab + aB)) \sin \left(c + dx + \frac{\pi}{2} \right) \right) dx + \frac{2b(13aB + 9a^2)}{11d} \right)$$

$$\frac{2bB \sin(c + dx) \cos^{7/2}(c + dx)(a + b \cos(c + dx))}{11d}$$

↓ 3227

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \int \cos^{5/2}(c + dx) dx + 9(11a(aB + 2Ab) + 9b^2B) \int \cos^{7/2}(c + dx) dx \right) + \frac{2b(13aB + 9a^2)}{11d} \right)$$

$$\frac{2bB \sin(c + dx) \cos^{7/2}(c + dx)(a + b \cos(c + dx))}{11d}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} dx + 9(11a(aB + 2Ab) + 9b^2B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{7/2} dx \right) + \frac{2b(13aB + 9a^2)}{11d} \right)$$

$$\frac{2bB \sin(c + dx) \cos^{7/2}(c + dx)(a + b \cos(c + dx))}{11d}$$

↓ 3115

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \left(\frac{3}{5} \int \sqrt{\cos(c + dx)} dx + \frac{2 \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + 9(11a(aB + 2Ab) + 9b^2B) \int \cos^{7/2}(c + dx) dx \right) + \frac{2b(13aB + 9a^2)}{11d} \right)$$

$$\frac{2bB \sin(c + dx) \cos^{7/2}(c + dx)(a + b \cos(c + dx))}{11d}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \left(\frac{3}{5} \int \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} dx + \frac{2 \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + 9(11a(aB + 2Ab) + 9b^2B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{7/2} dx \right) + \frac{2b(13aB + 9a^2)}{11d} \right)$$

$$\frac{2bB \sin(c + dx) \cos^{7/2}(c + dx)(a + b \cos(c + dx))}{11d}$$

↓ 3115

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \left(\frac{3}{5} \int \sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} dx + \frac{2 \sin(c + dx) \cos^{3/2}(c + dx)}{5d} \right) + 9(11a(aB + 2Ab) + 9b^2B) \int \cos^{7/2}(c + dx) dx \right) + \frac{2b(13aB + 9a^2)}{11d} \right)$$

$$\frac{2bB \sin(c + dx) \cos^{7/2}(c + dx)(a + b \cos(c + dx))}{11d}$$

↓ 3042

3.352. $\int \cos^{5/2}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \left(\frac{3}{5} \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + 9(11a(aB + 2Ab) + 9b^2B) \right) \right. \\ \left. \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) \downarrow \text{3119}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(9(11a(aB + 2Ab) + 9b^2B) \left(\frac{5}{7} \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) + \frac{2 \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d} \right) \right) \downarrow \text{3120}$$

$$\frac{1}{11} \left(\frac{1}{9} \left(11(9a^2A + 14abB + 7Ab^2) \left(\frac{6E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d} + \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} \right) + 9(11a(aB + 2Ab) + 9b^2B) \right) \right) \frac{2bB \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))}{11d}$$

input `Int[Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(2*b*B*Cos[c + d*x]^(7/2)*(a + b*Cos[c + d*x])*Sin[c + d*x]/(11*d) + ((2*b*(11*A*b + 13*a*B)*Cos[c + d*x]^(7/2)*Sin[c + d*x]/(9*d) + (11*(9*a^2*A + 7*A*b^2 + 14*a*b*B))*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)) + 9*(9*b^2*B + 11*a*(2*A*b + a*B))*((2*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(7*d) + (5*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/7)/9)/11`

3.352.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*SIN[e + f*x])^(m - 2)*(c + d*SIN[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*SIN[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*SIN[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*SIN[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*SIN[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*SIN[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.352.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 665 vs. $2(292) = 584$.

Time = 42.91 (sec) , antiderivative size = 666, normalized size of antiderivative = 2.52

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(20160B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^{12}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2+(-12320Ab^2-24640Bab-50400Bb^2)\left(\sin\right)}\right)}$
parts	Expression too large to display

input `int(cos(d*x+c)^(5/2)*(a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)`

output

```
-2/3465*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(20160*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^12*b^2+(-12320*A*b^2-24640*B*a*b-50400*B*b^2)*sin(1/2*d*x+1/2*c)^10*cos(1/2*d*x+1/2*c)+(15840*A*a*b+24640*A*b^2+7920*B*a^2+49280*B*a*b+56880*B*b^2)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-5544*A*a^2-23760*A*a*b-22792*A*b^2-11880*B*a^2-45584*B*a*b-34920*B*b^2)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(5544*A*a^2+18480*A*a*b+10472*A*b^2+9240*B*a^2+20944*B*a*b+13860*B*b^2)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-1386*A*a^2-5280*A*a*b-1848*A*b^2-2640*B*a^2-3696*B*a*b-2790*B*b^2)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+1650*A*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2079*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-1617*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^2+825*B*a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+675*B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3234*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.352.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.13

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{2(315 B b^2 \cos(dx + c)^4 + 385(2 B a b + A b^2) \cos(dx + c)^3 + 825 B a^2 + 1650 A a b + 675 B b^2 + 45(11 B a^2$$

```
input integrate(cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/3465*(2*(315*B*b^2*cos(d*x + c)^4 + 385*(2*B*a*b + A*b^2)*cos(d*x + c)^3
+ 825*B*a^2 + 1650*A*a*b + 675*B*b^2 + 45*(11*B*a^2 + 22*A*a*b + 9*B*b^2)
*cos(d*x + c)^2 + 77*(9*A*a^2 + 14*B*a*b + 7*A*b^2)*cos(d*x + c))*sqrt(cos
(d*x + c))*sin(d*x + c) - 75*sqrt(2)*(11*I*B*a^2 + 22*I*A*a*b + 9*I*B*b^2)
*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 75*sqrt(2)*(-
11*I*B*a^2 - 22*I*A*a*b - 9*I*B*b^2)*weierstrassPInverse(-4, 0, cos(d*x +
c) - I*sin(d*x + c)) - 231*sqrt(2)*(-9*I*A*a^2 - 14*I*B*a*b - 7*I*A*b^2)*w
eierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x
+ c))) - 231*sqrt(2)*(9*I*A*a^2 + 14*I*B*a*b + 7*I*A*b^2)*weierstrassZeta(-
4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d
```

3.352.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(5/2)*(a+b*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.352.7 Maxima [F]

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2), x)`

3.352.8 Giac [F]

$$\int \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2), x)`

3.352.9 Mupad [B] (verification not implemented)

Time = 1.70 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.04

$$\begin{aligned}
& \int \cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx)) dx \\
&= -\frac{2Aa^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}} \\
&\quad -\frac{2Ba^2\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}} \\
&\quad -\frac{2Ab^2\cos(c+dx)^{11/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)^2}} \\
&\quad -\frac{2Bb^2\cos(c+dx)^{13/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{13}{4}; \frac{17}{4}; \cos(c+dx)^2\right)}{13d\sqrt{\sin(c+dx)^2}} \\
&\quad -\frac{4Aab\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}} \\
&\quad -\frac{4Bab\cos(c+dx)^{11/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)^2}}
\end{aligned}$$

```
input int(cos(c + d*x)^(5/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2,x)
```

```
output - (2*A*a^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos
(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*a^2*cos(c + d*x)^(9/2)*s
in(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x
)^2)^(1/2)) - (2*A*b^2*cos(c + d*x)^(11/2)*sin(c + d*x)*hypergeom([1/2, 11
/4], 15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2)) - (2*B*b^2*cos(c
+ d*x)^(13/2)*sin(c + d*x)*hypergeom([1/2, 13/4], 17/4, cos(c + d*x)^2))/
(13*d*(sin(c + d*x)^2)^(1/2)) - (4*A*a*b*cos(c + d*x)^(9/2)*sin(c + d*x)*h
ypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) -
(4*B*a*b*cos(c + d*x)^(11/2)*sin(c + d*x)*hypergeom([1/2, 11/4], 15/4, co
s(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2))
```

3.353 $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$

3.353.1 Optimal result	3294
3.353.2 Mathematica [A] (verified)	3295
3.353.3 Rubi [A] (verified)	3295
3.353.4 Maple [B] (verified)	3299
3.353.5 Fricas [C] (verification not implemented)	3300
3.353.6 Sympy [F(-1)]	3301
3.353.7 Maxima [F]	3301
3.353.8 Giac [F]	3302
3.353.9 Mupad [B] (verification not implemented)	3302

3.353.1 Optimal result

Integrand size = 33, antiderivative size = 223

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2(A+B \cos(c+dx)) dx$$

$$= \frac{2(7b^2B + 9a(2Ab + aB)) E(\frac{1}{2}(c+dx)|2)}{15d}$$

$$+ \frac{2(7a^2A + 5Ab^2 + 10abB) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{21d}$$

$$+ \frac{2(7a^2A + 5Ab^2 + 10abB) \sqrt{\cos(c+dx)} \sin(c+dx)}{21d}$$

$$+ \frac{2(7b^2B + 9a(2Ab + aB)) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{45d}$$

$$+ \frac{2b(9Ab + 11aB) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{63d}$$

$$+ \frac{2bB \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx)) \sin(c+dx)}{9d}$$

output

```
2/15*(7*b^2*B+9*a*(2*A*b+B*a))*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/21*(7*A*a^2+5*A*b^2+10*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/45*(7*b^2*B+9*a*(2*A*b+B*a))*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/63*b*(9*A*b+11*B*a)*cos(d*x+c)^(5/2)*sin(d*x+c)/d+2/9*b*B*cos(d*x+c)^(5/2)*(a+b*cos(d*x+c))*sin(d*x+c)/d+2/21*(7*A*a^2+5*A*b^2+10*B*a*b)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.353.2 Mathematica [A] (verified)

Time = 2.05 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.75

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{84(18aAb+9a^2B+7b^2B)E\left(\frac{1}{2}(c+dx)\middle|2\right)+60(7a^2A+5Ab^2+10abB)\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)+\sqrt{\cos(c+dx)}(72a^2A+36a^2B+43b^2B)\cos(c+dx)+5(84a^2A+78Ab^2+156abB+18b(Ab+2aB))\cos(2(c+dx))+7b^2B\cos(3(c+dx))\sin(c+dx)}{630d}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(84*(18*a*A*b + 9*a^2*B + 7*b^2*B)*EllipticE[(c + d*x)/2, 2] + 60*(7*a^2*A + 5*A*b^2 + 10*a*b*B)*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(7*(72*a^2*A + 36*a^2*B + 43*b^2*B)*Cos[c + d*x] + 5*(84*a^2*A + 78*A*b^2 + 156*a*b*B + 18*b*(A*b + 2*a*B))*Cos[2*(c + d*x)] + 7*b^2*B*Cos[3*(c + d*x)])) *Sin[c + d*x])/(630*d)`

3.353.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.91, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^2\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx$$

$$\downarrow \text{3469}$$

$$\frac{2}{9}\int\frac{1}{2}\cos^{\frac{3}{2}}(c+dx)(b(9Ab+11aB)\cos^2(c+dx)+(7Bb^2+9a(2Ab+aB))\cos(c+dx)+a(9aA+5bB))dx+2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))$$

$$\downarrow \text{27}$$

3.353. $\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$

$$\frac{\frac{1}{9} \int \cos^{\frac{3}{2}}(c + dx) (b(9Ab + 11aB) \cos^2(c + dx) + (7Bb^2 + 9a(2Ab + aB)) \cos(c + dx) + a(9aA + 5bB)) dx + 2bB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))}{9d}$$

↓ 3042

$$\frac{\frac{1}{9} \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(b(9Ab + 11aB) \sin\left(c + dx + \frac{\pi}{2}\right)^2 + (7Bb^2 + 9a(2Ab + aB)) \sin\left(c + dx + \frac{\pi}{2}\right) + a(9aA + 5bB)\right) dx + 2bB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))}{9d}$$

↓ 3502

$$\frac{\frac{1}{9} \left(\frac{2}{7} \int \frac{1}{2} \cos^{\frac{3}{2}}(c + dx) (9(7Aa^2 + 10bBa + 5Ab^2) + 7(7Bb^2 + 9a(2Ab + aB)) \cos(c + dx)) dx + \frac{2b(11aB + 9Ab)}{9d} \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx)) \right)}{9d}$$

↓ 27

$$\frac{\frac{1}{9} \left(\frac{1}{7} \int \cos^{\frac{3}{2}}(c + dx) (9(7Aa^2 + 10bBa + 5Ab^2) + 7(7Bb^2 + 9a(2Ab + aB)) \cos(c + dx)) dx + \frac{2b(11aB + 9Ab)}{9d} \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx)) \right)}{9d}$$

↓ 3042

$$\frac{\frac{1}{9} \left(\frac{1}{7} \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (9(7Aa^2 + 10bBa + 5Ab^2) + 7(7Bb^2 + 9a(2Ab + aB)) \sin\left(c + dx + \frac{\pi}{2}\right)) dx + \frac{2b(11aB + 9Ab)}{9d} \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx)) \right)}{9d}$$

↓ 3227

$$\frac{\frac{1}{9} \left(\frac{1}{7} \left(9(7a^2A + 10abB + 5Ab^2) \int \cos^{\frac{3}{2}}(c + dx) dx + 7(9a(aB + 2Ab) + 7b^2B) \int \cos^{\frac{5}{2}}(c + dx) dx \right) + \frac{2b(11aB + 9Ab)}{9d} \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx)) \right)}{9d}$$

↓ 3042

3.353. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$

$$\frac{1}{9} \left(\frac{1}{7} \left(9(7a^2A + 10abB + 5Ab^2) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} dx + 7(9a(aB + 2Ab) + 7b^2B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{5/2} dx \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx) \cos^{5/2}(c + dx)(a + b \cos(c + dx))}{9d} \right) \right. \\ \left. \downarrow \text{3115} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(9(7a^2A + 10abB + 5Ab^2) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + 7(9a(aB + 2Ab) + 7b^2B) \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx) \cos^{5/2}(c + dx)(a + b \cos(c + dx))}{9d} \right) \right. \\ \left. \downarrow \text{3042} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(9(7a^2A + 10abB + 5Ab^2) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + 7(9a(aB + 2Ab) + 7b^2B) \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx) \cos^{5/2}(c + dx)(a + b \cos(c + dx))}{9d} \right) \right. \\ \left. \downarrow \text{3119} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(9(7a^2A + 10abB + 5Ab^2) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + 7(9a(aB + 2Ab) + 7b^2B) \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx) \cos^{5/2}(c + dx)(a + b \cos(c + dx))}{9d} \right) \right. \\ \left. \downarrow \text{3120} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(9(7a^2A + 10abB + 5Ab^2) \left(\frac{2 \operatorname{EllipticF} \left(\frac{1}{2}(c + dx), 2 \right)}{3d} + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + 7(9a(aB + 2Ab) + 7b^2B) \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx) \cos^{5/2}(c + dx)(a + b \cos(c + dx))}{9d} \right) \right.$$

input `Int[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`


```
output (2*b*B*Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])*Sin[c + d*x]/(9*d) + ((2*b
*(9*A*b + 11*a*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x]/(7*d) + (9*(7*a^2*A + 5
*A*b^2 + 10*a*b*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c +
d*x]]*Sin[c + d*x]/(3*d)) + 7*(7*b^2*B + 9*a*(2*A*b + a*B))*((6*EllipticE
[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x]/(5*d)))/7)/9
```

3.353.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3115 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*SIN[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*SIN
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*SIN[e + f*x])^m, x], x] + Simp[d/b Int
[(b*SIN[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.353.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 609 vs. 2(255) = 510.

Time = 14.21 (sec) , antiderivative size = 610, normalized size of antiderivative = 2.74

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(-1120B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^{10}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2+(720A b^2+1440Bab+2240B^2b^2)\left(\sin^8\left(\frac{dx}{2}\right.\right.\right.}$
parts	Expression too large to display

```
input int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.353. \quad \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

output

```

-2/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b^2+(720*A*b^2+1440*B*a*b+2240*B*b^2)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1008*A*a*b-1080*A*b^2-504*B*a^2-2160*B*a*b-2072*B*b^2)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(420*A*a^2+1008*A*a*b+840*A*b^2+504*B*a^2+1680*B*a*b+952*B*b^2)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-210*A*a^2-252*A*a*b-240*A*b^2-126*B*a^2-480*B*a*b-168*B*b^2)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+105*A*a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+75*A*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-378*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b+150*B*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-189*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-147*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d

```

3.353.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.22

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{2(35Bb^2\cos(dx+c)^3+105Aa^2+150Bab+75Ab^2+45(2Bab+Ab^2)\cos(dx+c)^2+7(9Ba^2+18Aa^2+18Ab^2)\cos(dx+c)+7Bb^2)}{\sin(dx+c)}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")`

```
output 1/315*(2*(35*B*b^2*cos(d*x + c)^3 + 105*A*a^2 + 150*B*a*b + 75*A*b^2 + 45*
(2*B*a*b + A*b^2)*cos(d*x + c)^2 + 7*(9*B*a^2 + 18*A*a*b + 7*B*b^2)*cos(d*
x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 15*sqrt(2)*(7*I*A*a^2 + 10*I*B*a
*b + 5*I*A*b^2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))
- 15*sqrt(2)*(-7*I*A*a^2 - 10*I*B*a*b - 5*I*A*b^2)*weierstrassPInverse(-4,
0, cos(d*x + c) - I*sin(d*x + c)) - 21*sqrt(2)*(-9*I*B*a^2 - 18*I*A*a*b -
7*I*B*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c)
+ I*sin(d*x + c))) - 21*sqrt(2)*(9*I*B*a^2 + 18*I*A*a*b + 7*I*B*b^2)*weie
rstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c
))))/d
```

3.353.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(3/2)*(a+b*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.353.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

```
input integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm
m="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2),
x)
```

3.353.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^2\cos(dx+c)^{\frac{3}{2}}dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2), x)`

3.353.9 Mupad [B] (verification not implemented)

Time = 1.52 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.18

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{2Aa^2\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\middle|2\right)\right)}{3d}$$

$$- \frac{2Ba^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)}^2}$$

$$- \frac{2Ab^2\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)}^2}$$

$$- \frac{2Bb^2\cos(c+dx)^{11/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11d\sqrt{\sin(c+dx)}^2}$$

$$- \frac{4Aab\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)}^2}$$

$$- \frac{4Bab\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)}^2}$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2,x)`

3.353. $\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$

```

output (2*A*a^2*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2))/
(3*d) - (2*B*a^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/
4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*A*b^2*cos(c + d*x)^(
9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c
+ d*x)^2)^(1/2)) - (2*B*b^2*cos(c + d*x)^(11/2)*sin(c + d*x)*hypergeom([1
/2, 11/4], 15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2)) - (4*A*a*b
*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^
2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (4*B*a*b*cos(c + d*x)^(9/2)*sin(c + d*x
)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)
)

```

3.354 $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$

3.354.1 Optimal result	3304
3.354.2 Mathematica [A] (verified)	3305
3.354.3 Rubi [A] (verified)	3305
3.354.4 Maple [B] (verified)	3309
3.354.5 Fricas [C] (verification not implemented)	3310
3.354.6 Sympy [F(-1)]	3311
3.354.7 Maxima [F]	3311
3.354.8 Giac [F]	3311
3.354.9 Mupad [B] (verification not implemented)	3312

3.354.1 Optimal result

Integrand size = 33, antiderivative size = 182

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx$$

$$= \frac{2(5a^2A + 3Ab^2 + 6abB) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d}$$

$$+ \frac{2(5b^2B + 7a(2Ab + aB)) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d}$$

$$+ \frac{2(5b^2B + 7a(2Ab + aB)) \sqrt{\cos(c + dx)} \sin(c + dx)}{21d}$$

$$+ \frac{2b(7Ab + 9aB) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{35d}$$

$$+ \frac{2bB \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx)) \sin(c + dx)}{7d}$$

output

```
2/5*(5*A*a^2+3*A*b^2+6*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/21*(5*b^2*B+7*a*(2*A*b+B*a))*cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/35*b*(7*A*b+9*B*a)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/7*b*B*cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))*sin(d*x+c)/d+2/21*(5*b^2*B+7*a*(2*A*b+B*a))*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.354.2 Mathematica [A] (verified)

Time = 1.86 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.76

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{42(5a^2A+3Ab^2+6abB)E\left(\frac{1}{2}(c+dx)\middle|2\right)+10(14aAb+7a^2B+5b^2B)\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)+\sqrt{\cos(c+dx)}(42b(Ab+2aB)\cos(c+dx)+5(28aAb+14a^2B+13b^2B+3b^2B\cos(2(c+dx)))\sin(c+dx))}{105d}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output `(42*(5*a^2*A + 3*A*b^2 + 6*a*b*B)*EllipticE[(c + d*x)/2, 2] + 10*(14*a*A*b + 7*a^2*B + 5*b^2*B)*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(42*b*(A*b + 2*a*B)*Cos[c + d*x] + 5*(28*a*A*b + 14*a^2*B + 13*b^2*B + 3*b^2*B*Cos[2*(c + d*x)]))*Sin[c + d*x])/(105*d)`

3.354.3 Rubi [A] (verified)Time = 0.91 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.96, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$\downarrow 3042$$

$$\int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^2\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx$$

$$\downarrow 3469$$

$$\frac{2}{7}\int\frac{1}{2}\sqrt{\cos(c+dx)}(b(7Ab+9aB)\cos^2(c+dx)+(5Bb^2+7a(2Ab+aB))\cos(c+dx)+a(7aA+3bB))dx+\frac{2bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))}{7d}$$

$$\downarrow 27$$

$$\frac{1}{7} \int \sqrt{\cos(c+dx)} (b(7Ab+9aB) \cos^2(c+dx) + (5Bb^2+7a(2Ab+aB)) \cos(c+dx) + a(7aA+3bB)) dx + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}{7d}$$

↓ 3042

$$\frac{1}{7} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(b(7Ab+9aB) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + (5Bb^2+7a(2Ab+aB)) \sin\left(c+dx+\frac{\pi}{2}\right) + a(7aA+3bB) \right) dx + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}{7d}$$

↓ 3502

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{1}{2} \sqrt{\cos(c+dx)} (7(5Aa^2+6bBa+3Ab^2) + 5(5Bb^2+7a(2Ab+aB)) \cos(c+dx)) dx + \frac{2b(9aB+7Ab) \sin(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}{7d}$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \int \sqrt{\cos(c+dx)} (7(5Aa^2+6bBa+3Ab^2) + 5(5Bb^2+7a(2Ab+aB)) \cos(c+dx)) dx + \frac{2b(9aB+7Ab) \sin(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(7(5Aa^2+6bBa+3Ab^2) + 5(5Bb^2+7a(2Ab+aB)) \sin\left(c+dx+\frac{\pi}{2}\right) \right) dx + \frac{2b(9aB+7Ab) \sin(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}{7d}$$

↓ 3227

$$\frac{1}{7} \left(\frac{1}{5} \left(7(5a^2A+6abB+3Ab^2) \int \sqrt{\cos(c+dx)} dx + 5(7a(aB+2Ab)+5b^2B) \int \cos^{\frac{3}{2}}(c+dx) dx \right) + \frac{2b(9aB+7Ab) \sin(c+dx)}{7d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}{7d}$$

↓ 3042

3.354. $\int \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(7(5a^2A + 6abB + 3Ab^2) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5(7a(aB + 2Ab) + 5b^2B) \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx \right) \right. \\ \left. \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}{7d} \right) \\ \downarrow \text{3115}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(7(5a^2A + 6abB + 3Ab^2) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5(7a(aB + 2Ab) + 5b^2B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 2 \right) \right) \right. \\ \left. \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}{7d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(7(5a^2A + 6abB + 3Ab^2) \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx + 5(7a(aB + 2Ab) + 5b^2B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + 2 \right) \right) \right. \\ \left. \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}{7d} \right) \\ \downarrow \text{3119}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(5(7a(aB + 2Ab) + 5b^2B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) + \frac{14(5a^2A + 6abB + 3Ab^2) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} \right) \right. \\ \left. \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}{7d} \right) \\ \downarrow \text{3120}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{14(5a^2A + 6abB + 3Ab^2) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + 5(7a(aB + 2Ab) + 5b^2B) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2 \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \right) \right) \right. \\ \left. \frac{2bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}{7d} \right)$$

input `Int[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]),x]`

output $(2*b*B*\cos[c + d*x]^{(3/2)}*(a + b*\cos[c + d*x])*sin[c + d*x])/(7*d) + ((2*b*(7*A*b + 9*a*B)*\cos[c + d*x]^{(3/2)}*sin[c + d*x])/(5*d) + ((14*(5*a^2*A + 3*A*b^2 + 6*a*b*B)*\text{EllipticE}[(c + d*x)/2, 2])/d + 5*(5*b^2*B + 7*a*(2*A*b + a*B))*((2*\text{EllipticF}[(c + d*x)/2, 2])/(3*d) + (2*\text{Sqrt}[\cos[c + d*x]]*sin[c + d*x])/(3*d)))/5)/7$

3.354.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_) /; \text{FreeQ}[b, x]]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3115 $\text{Int}[(b_*)*\sin[(c_*) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((b*\sin[c + d*x])^{(n-1)})/(d*n), x] + \text{Simp}[b^2*((n-1)/n) \text{ Int}[(b*\sin[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

rule 3227 $\text{Int}[(b_*)*\sin[(e_*) + (f_)*(x_)]^{(m_)}*((c_*) + (d_)*\sin[(e_*) + (f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[c \text{ Int}[(b*\sin[e + f*x])^m, x], x] + \text{Simp}[d/b \text{ Int}[(b*\sin[e + f*x])^{(m+1)}, x], x] /; \text{FreeQ}\{b, c, d, e, f, m\}, x]$

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
  mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
  n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
  f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
  m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
  + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
  [e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
  - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
  Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
  + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
  s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
  + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
  + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
  && !LtQ[m, -1]
```

3.354.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(218) = 436$.

Time = 12.63 (sec) , antiderivative size = 548, normalized size of antiderivative = 3.01

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\left(240B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^2 + (-168Ab^2 - 336Bab - 360Bb^2)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$
parts	Expression too large to display

```
input int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```
-2/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*cos(
1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^2+(-168*A*b^2-336*B*a*b-360*B*b^2)*s
in(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(280*A*a*b+168*A*b^2+140*B*a^2+336*
B*a*b+280*B*b^2)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-140*A*a*b-42*A*
b^2-70*B*a^2-84*B*a*b-80*B*b^2)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+70
*A*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellip
ticF(cos(1/2*d*x+1/2*c),2^(1/2))-105*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin
(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-63*A*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(co
s(1/2*d*x+1/2*c),2^(1/2))*b^2+35*B*a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin
(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+25*B*b^2*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))-126*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*
x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b)/(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*
x+1/2*c)^2-1)^(1/2)/d
```

3.354.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.34

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$$

$$= \frac{2(15Bb^2\cos(dx+c)^2+35Ba^2+70Aab+25Bb^2+21(2Bab+Ab^2)\cos(dx+c))\sqrt{\cos(dx+c)}\sin(dx+c)}{d}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output

```
1/105*(2*(15*B*b^2*cos(d*x + c)^2 + 35*B*a^2 + 70*A*a*b + 25*B*b^2 + 21*(2
*B*a*b + A*b^2)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 5*sqrt(2)*
(7*I*B*a^2 + 14*I*A*a*b + 5*I*B*b^2)*weierstrassPInverse(-4, 0, cos(d*x +
c) + I*sin(d*x + c)) - 5*sqrt(2)*(-7*I*B*a^2 - 14*I*A*a*b - 5*I*B*b^2)*wei
erstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*sqrt(2)*(-5*I*
A*a^2 - 6*I*B*a*b - 3*I*A*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse(
-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*sqrt(2)*(5*I*A*a^2 + 6*I*B*a*b
+ 3*I*A*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x +
c) - I*sin(d*x + c))))/d
```

3.354. $\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2(A+B\cos(c+dx))dx$

3.354.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a+b*cos(d*x+c))**2*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.354.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^2*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sqrt(cos(d*x + c)), x)`

3.354.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))**2*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sqrt(cos(d*x + c)), x)`

3.354.9 Mupad [B] (verification not implemented)

Time = 1.49 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.26

$$\begin{aligned}
& \int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2(A+B\cos(c+dx)) dx \\
&= \frac{2Ba^2\left(\sqrt{\cos(c+dx)}\sin(c+dx)+F\left(\frac{c}{2}+\frac{dx}{2}\mid 2\right)\right)}{3d} \\
&+ \frac{2Aa^2E\left(\frac{c}{2}+\frac{dx}{2}\mid 2\right)}{d} + \frac{2Aab\left(\frac{2\sqrt{\cos(c+dx)}\sin(c+dx)}{3}+\frac{2F\left(\frac{c}{2}+\frac{dx}{2}\mid 2\right)}{3}\right)}{d} \\
&- \frac{2Ab^2\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}} \\
&- \frac{2Bb^2\cos(c+dx)^{9/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{9d\sqrt{\sin(c+dx)^2}} \\
&- \frac{4Bab\cos(c+dx)^{7/2}\sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7d\sqrt{\sin(c+dx)^2}}
\end{aligned}$$

```
input int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2,x)
```

```
output (2*B*a^2*(cos(c + d*x)^(1/2)*sin(c + d*x) + ellipticF(c/2 + (d*x)/2, 2)))/
(3*d) + (2*A*a^2*ellipticE(c/2 + (d*x)/2, 2))/d + (2*A*a*b*((2*cos(c + d*x)
)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d - (2*A*b^2
*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^
2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*b^2*cos(c + d*x)^(9/2)*sin(c + d*x
)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)
) - (4*B*a*b*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, c
os(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))
```

3.355
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.355.1 Optimal result 3313
 3.355.2 Mathematica [A] (verified) 3314
 3.355.3 Rubi [A] (verified) 3314
 3.355.4 Maple [B] (verified) 3317
 3.355.5 Fricas [C] (verification not implemented) 3318
 3.355.6 Sympy [F(-1)] 3319
 3.355.7 Maxima [F] 3319
 3.355.8 Giac [F] 3319
 3.355.9 Mupad [B] (verification not implemented) 3320

3.355.1 Optimal result

Integrand size = 33, antiderivative size = 140

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2(3b^2B + 5a(2Ab + aB)) E(\frac{1}{2}(c + dx) | 2)}{5d} \\ &+ \frac{2(3a^2A + Ab^2 + 2abB) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} \\ &+ \frac{2b(5Ab + 7aB) \sqrt{\cos(c + dx)} \sin(c + dx)}{15d} \\ &+ \frac{2bB \sqrt{\cos(c + dx)}(a + b \cos(c + dx)) \sin(c + dx)}{5d} \end{aligned}$$

```
output 2/5*(3*b^2*B+5*a*(2*A*b+B*a))*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(3*A*a^2+A*b^2+2*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/15*b*(5*A*b+7*B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/d+2/5*b*B*(a+b*cos(d*x+c))*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```


3.355.2 Mathematica [A] (verified)

Time = 1.17 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.76

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2 \left(3(10aAb + 5a^2B + 3b^2B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 5(3a^2A + Ab^2 + 2abB) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + b\sqrt{\cos(c + dx)} \right)}{15d}$$

input `Integrate[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(2*(3*(10*a*A*b + 5*a^2*B + 3*b^2*B)*EllipticE[(c + d*x)/2, 2] + 5*(3*a^2*A + A*b^2 + 2*a*b*B)*EllipticF[(c + d*x)/2, 2] + b*Sqrt[Cos[c + d*x]]*(5*A*b + 10*a*B + 3*b*B*Cos[c + d*x])*Sin[c + d*x]))/(15*d)`

3.355.3 Rubi [A] (verified)Time = 0.79 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3469, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3469}$$

$$\frac{2}{5} \int \frac{b(5Ab + 7aB) \cos^2(c + dx) + (3Bb^2 + 5a(2Ab + aB)) \cos(c + dx) + a(5aA + bB)}{2\sqrt{\cos(c + dx)}} dx +$$

$$\frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d}$$

$$\downarrow \text{27}$$

$$\frac{1}{5} \int \frac{b(5Ab + 7aB) \cos^2(c + dx) + (3Bb^2 + 5a(2Ab + aB)) \cos(c + dx) + a(5aA + bB)}{\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \int \frac{b(5Ab + 7aB) \sin(c + dx + \frac{\pi}{2})^2 + (3Bb^2 + 5a(2Ab + aB)) \sin(c + dx + \frac{\pi}{2}) + a(5aA + bB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d}$$

↓ 3502

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{5(3Aa^2 + 2bBa + Ab^2) + 3(3Bb^2 + 5a(2Ab + aB)) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2b(7aB + 5Ab) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{5(3Aa^2 + 2bBa + Ab^2) + 3(3Bb^2 + 5a(2Ab + aB)) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2b(7aB + 5Ab) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{5(3Aa^2 + 2bBa + Ab^2) + 3(3Bb^2 + 5a(2Ab + aB)) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b(7aB + 5Ab) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

↓ 3227

$$\frac{1}{5} \left(\frac{1}{3} \left(5(3a^2A + 2abB + Ab^2) \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3(5a(aB + 2Ab) + 3b^2B) \int \sqrt{\cos(c + dx)} dx \right) + \frac{2b(7aB + 5Ab) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(5(3a^2A + 2abB + Ab^2) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(5a(aB + 2Ab) + 3b^2B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

↓ 3119

$$\frac{1}{5} \left(\frac{1}{3} \left(5(3a^2A + 2abB + Ab^2) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{6(5a(aB + 2Ab) + 3b^2B) E(\frac{1}{2}(c + dx)|2)}{d} \right) + \frac{2b(7aB + b^2) \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{10(3a^2A + 2abB + Ab^2) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} + \frac{6(5a(aB + 2Ab) + 3b^2B) E(\frac{1}{2}(c + dx)|2)}{d} \right) + \frac{2b(7aB + b^2) \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))}{5d} \right)$$

input `Int[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(2*b*B*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])*Sin[c + d*x]/(5*d) + (((6*(3*b^2*B + 5*a*(2*A*b + a*B))*EllipticE[(c + d*x)/2, 2])/d + (10*(3*a^2*A + A*b^2 + 2*a*b*B))*EllipticF[(c + d*x)/2, 2])/d)/3 + (2*b*(5*A*b + 7*a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

3.355.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.355.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 486 vs. 2(180) = 360.

Time = 9.59 (sec) , antiderivative size = 487, normalized size of antiderivative = 3.48

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(-24B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^2 + (20Ab^2 + 40Bab + 24Bb^2)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}}$
parts	$-\frac{2(Ab^2 + 2Bab)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}}$

3.355.
$$\int \frac{(a+b\cos(c+dx))^2(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

input `int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)`

output `-2/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*cos(1
/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^2+(20*A*b^2+40*B*a*b+24*B*b^2)*sin(1/
2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*b^2-20*B*a*b-6*B*b^2)*sin(1/2*d*x
+1/2*c)^2*cos(1/2*d*x+1/2*c)+15*A*a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(
1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+5*A*b^2*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(
1/2*d*x+1/2*c),2^(1/2))-30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b+10*B*a*b*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),2^(1/2))-15*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c
)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-9*B*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c
) ,2^(1/2))*b^2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1
/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.355.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.54

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2(3Bb^2 \cos(dx + c) + 10Bab + 5Ab^2) \sqrt{\cos(dx + c)} \sin(dx + c) - 5\sqrt{2}(3iAa^2 + 2iBab + iAb^2) \text{weiers}}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm
m="fricas")`

output `1/15*(2*(3*B*b^2*cos(d*x + c) + 10*B*a*b + 5*A*b^2)*sqrt(cos(d*x + c))*sin
(d*x + c) - 5*sqrt(2)*(3*I*A*a^2 + 2*I*B*a*b + I*A*b^2)*weierstrassPInvers
e(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*sqrt(2)*(-3*I*A*a^2 - 2*I*B*a*
b - I*A*b^2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3
sqrt(2)(-5*I*B*a^2 - 10*I*A*a*b - 3*I*B*b^2)*weierstrassZeta(-4, 0, weie
rstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(5*I*B*
a^2 + 10*I*A*a*b + 3*I*B*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-
4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.355. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

3.355.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Timed out`

3.355.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\sqrt{\cos(dx + c)}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/sqrt(cos(d*x + c)), x)`

3.355.8 Giac [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\sqrt{\cos(dx + c)}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/sqrt(cos(d*x + c)), x)`

3.355.9 Mupad [B] (verification not implemented)

Time = 1.67 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.26

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{A b^2 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{3} \right)}{d} + \frac{2 A a^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{2 B a^2 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$+ \frac{2 B a b \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{3} \right)}{d} + \frac{4 A a b E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$- \frac{2 B b^2 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^(1/2),x)`output `(A*b^2*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (2*A*a^2*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*a^2*ellipticE(c/2 + (d*x)/2, 2))/d + (2*B*a*b*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (4*A*a*b*ellipticE(c/2 + (d*x)/2, 2))/d - (2*B*b^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))`

3.356
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.356.1 Optimal result 3321
 3.356.2 Mathematica [A] (verified) 3322
 3.356.3 Rubi [A] (verified) 3322
 3.356.4 Maple [B] (verified) 3325
 3.356.5 Fracas [C] (verification not implemented) 3326
 3.356.6 Sympy [F(-1)] 3327
 3.356.7 Maxima [F] 3327
 3.356.8 Giac [F] 3327
 3.356.9 Mupad [B] (verification not implemented) 3328

3.356.1 Optimal result

Integrand size = 33, antiderivative size = 121

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2(a^2 A - Ab^2 - 2abB) E(\frac{1}{2}(c + dx) | 2)}{d}$$

$$+ \frac{2(6aAb + 3a^2 B + b^2 B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 B \sqrt{\cos(c + dx)} \sin(c + dx)}{3d}$$

```
output -2*(A*a^2-A*b^2-2*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(6*A*a*b+3*B*a^2+B*b^2)*(cos(1/
2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1
/2))/d+2*a^2*A*sin(d*x+c)/d/cos(d*x+c)^(1/2)+2/3*b^2*B*sin(d*x+c)*cos(d*x+
c)^(1/2)/d
```


3.356.2 Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.84

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 \left((-3a^2 A + 3Ab^2 + 6abB) E\left(\frac{1}{2}(c + dx) \mid 2\right) + (6aAb + 3a^2 B + b^2 B) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \frac{(3a^2 A + b^2 B) \sin(c + dx)}{\sqrt{\cos(c + dx)}} \right)}{3d}$$

input `Integrate[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(2*((-3*a^2*A + 3*A*b^2 + 6*a*b*B)*EllipticE[(c + d*x)/2, 2] + (6*a*A*b + 3*a^2*B + b^2*B)*EllipticF[(c + d*x)/2, 2] + ((3*a^2*A + b^2*B*Cos[c + d*x])*Sin[c + d*x])/Sqrt[Cos[c + d*x]])/(3*d)`

3.356.3 Rubi [A] (verified)Time = 0.73 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3467, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3467}$$

$$\frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - 2 \int -\frac{b^2 B \cos^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \cos(c + dx) + a(2Ab + aB)}{2\sqrt{\cos(c + dx)}} dx$$

$$\downarrow \text{27}$$

$$\int \frac{b^2 B \cos^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \cos(c + dx) + a(2Ab + aB)}{\sqrt{\cos(c + dx)}} dx + \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}}$$

3.356. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (-Aa^2 + 2bBa + Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(2Ab + aB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}}} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{2}{3} \int \frac{3Ba^2 + 6Aba + b^2 B - 3(Aa^2 - 2bBa - Ab^2) \cos(c + dx)}{2\sqrt{\cos(c + dx)} \frac{2b^2 B \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}} dx + \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \\
& \quad \downarrow \text{3502} \\
& \frac{1}{3} \int \frac{3Ba^2 + 6Aba + b^2 B - 3(Aa^2 - 2bBa - Ab^2) \cos(c + dx)}{\sqrt{\cos(c + dx)} \frac{2b^2 B \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}} dx + \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{3Ba^2 + 6Aba + b^2 B - 3(Aa^2 - 2bBa - Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \frac{2b^2 B \sin(c + dx) \sqrt{\cos(c + dx)}}{3d}} dx + \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left((3a^2 B + 6aAb + b^2 B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - 3(a^2 A - 2abB - Ab^2) \int \sqrt{\cos(c + dx)} dx \right) + \\
& \quad \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 B \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3227} \\
& \frac{1}{3} \left((3a^2 B + 6aAb + b^2 B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - 3(a^2 A - 2abB - Ab^2) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) + \\
& \quad \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 B \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left((3a^2 B + 6aAb + b^2 B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - 3(a^2 A - 2abB - Ab^2) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) + \\
& \quad \frac{2a^2 A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 B \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} \\
& \quad \downarrow \text{3119}
\end{aligned}$$

3.356. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

$$\frac{1}{3} \left((3a^2B + 6aAb + b^2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{6(a^2A - 2abB - Ab^2) E(\frac{1}{2}(c + dx) | 2)}{d} \right) + \frac{2a^2A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2B \sin(c + dx)\sqrt{\cos(c + dx)}}{3d}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2(3a^2B + 6aAb + b^2B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} - \frac{6(a^2A - 2abB - Ab^2) E(\frac{1}{2}(c + dx) | 2)}{d} \right) + \frac{2a^2A \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2B \sin(c + dx)\sqrt{\cos(c + dx)}}{3d}$$

input `Int[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `((-6*(a^2*A - A*b^2 - 2*a*b*B)*EllipticE[(c + d*x)/2, 2])/d + (2*(6*a*A*b + 3*a^2*B + b^2*B)*EllipticF[(c + d*x)/2, 2])/d)/3 + (2*a^2*A*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) + (2*b^2*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)`

3.356.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

3.356. $\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$

```
rule 3467 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]
```

3.356.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 404 vs. 2(165) = 330.

Time = 9.84 (sec) , antiderivative size = 405, normalized size of antiderivative = 3.35

method	result
default	$-\frac{8B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b^2}{3} + 4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a^2 - 4Aab \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \dots\right)$
parts	$\frac{2(Ab^2 + 2Bab) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d} + \frac{2(2Aab - \dots)}{\dots}$

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNV ERBOSE)
```

$$3.356. \int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

output
$$\frac{2}{3}(-4B\cos(\frac{1}{2}dx+\frac{1}{2}c)\sin(\frac{1}{2}dx+\frac{1}{2}c)^4b^2+6A\cos(\frac{1}{2}dx+\frac{1}{2}c)\sin(\frac{1}{2}dx+\frac{1}{2}c)^2a^2-6Aa*b*(\sin(\frac{1}{2}dx+\frac{1}{2}c)^2)^{(1/2)}*(2*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}*EllipticF(\cos(\frac{1}{2}dx+\frac{1}{2}c),2^{(1/2)})-3A*(\sin(\frac{1}{2}dx+\frac{1}{2}c)^2)^{(1/2)}*(2*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}*EllipticE(\cos(\frac{1}{2}dx+\frac{1}{2}c),2^{(1/2)})a^2+3A*(\sin(\frac{1}{2}dx+\frac{1}{2}c)^2)^{(1/2)}*(2*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}*EllipticE(\cos(\frac{1}{2}dx+\frac{1}{2}c),2^{(1/2)})b^2+2B*\cos(\frac{1}{2}dx+\frac{1}{2}c)*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2b^2-3B*a^2*(\sin(\frac{1}{2}dx+\frac{1}{2}c)^2)^{(1/2)}*(2*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}*EllipticF(\cos(\frac{1}{2}dx+\frac{1}{2}c),2^{(1/2)})-B*b^2*(\sin(\frac{1}{2}dx+\frac{1}{2}c)^2)^{(1/2)}*(2*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}*EllipticF(\cos(\frac{1}{2}dx+\frac{1}{2}c),2^{(1/2)})+6B*(\sin(\frac{1}{2}dx+\frac{1}{2}c)^2)^{(1/2)}*(2*\sin(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}*EllipticE(\cos(\frac{1}{2}dx+\frac{1}{2}c),2^{(1/2)})a*b)/\sin(\frac{1}{2}dx+\frac{1}{2}c)/(2*\cos(\frac{1}{2}dx+\frac{1}{2}c)^2-1)^{(1/2)}/d$$

3.356.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.98

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{2}(-3iBa^2 - 6iAab - iBb^2) \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \sqrt{2}}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="fracas")`

output
$$\frac{1}{3}(\sqrt{2}*(-3I*B*a^2 - 6I*A*a*b - I*B*b^2)*\cos(d*x + c)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + \sqrt{2}*(3I*B*a^2 + 6I*A*a*b + I*B*b^2)*\cos(d*x + c)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*\sqrt{2}*(I*A*a^2 - 2I*B*a*b - I*A*b^2)*\cos(d*x + c)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - 3*\sqrt{2}*(-I*A*a^2 + 2I*B*a*b + I*A*b^2)*\cos(d*x + c)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)))) + 2*(B*b^2*\cos(d*x + c) + 3*A*a^2)*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(d*\cos(d*x + c))$$

3.356.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`output `Timed out`**3.356.7 Maxima [F]**

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="maxima")`output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/cos(d*x + c)^(3/2), x)`**3.356.8 Giac [F]**

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="giac")`output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/cos(d*x + c)^(3/2), x)`

3.356.9 Mupad [B] (verification not implemented)

Time = 1.85 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.31

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{B b^2 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{3} \right)}{d} + \frac{2 A b^2 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$+ \frac{2 B a^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 A a b F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 B a b E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$+ \frac{2 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^(3/2),x)`output `(B*b^2*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (2*A*b^2*ellipticE(c/2 + (d*x)/2, 2))/d + (2*B*a^2*ellipticF(c/2 + (d*x)/2, 2))/d + (4*A*a*b*ellipticF(c/2 + (d*x)/2, 2))/d + (4*B*a*b*ellipticE(c/2 + (d*x)/2, 2))/d + (2*A*a^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.357
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.357.1 Optimal result 3329
 3.357.2 Mathematica [A] (verified) 3330
 3.357.3 Rubi [A] (verified) 3330
 3.357.4 Maple [C] (verified) 3333
 3.357.5 Fracas [C] (verification not implemented) 3334
 3.357.6 Sympy [F(-1)] 3335
 3.357.7 Maxima [F] 3335
 3.357.8 Giac [F] 3336
 3.357.9 Mupad [B] (verification not implemented) 3336

3.357.1 Optimal result

Integrand size = 33, antiderivative size = 126

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{2(2aAb + a^2B - b^2B) E(\frac{1}{2}(c + dx) | 2)}{d}$$

$$+ \frac{2(a^2A + 3Ab^2 + 6abB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2a^2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2a(2Ab + aB) \sin(c + dx)}{d \sqrt{\cos(c + dx)}}$$

output

```
-2*(2*A*a*b+B*a^2-B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(A*a^2+3*A*b^2+6*B*a*b)*(cos(1/
2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1
/2))/d+2/3*a^2*A*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2*a*(2*A*b+B*a)*sin(d*x+c)/
d/cos(d*x+c)^(1/2)
```


3.357.2 Mathematica [A] (verified)

Time = 1.31 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.83

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 \left(-3(2aAb + a^2B - b^2B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + (a^2A + 3Ab^2 + 6abB) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \frac{a(aA + 3(2Ab + b^2))}{3d} \right)}{3d}$$

input `Integrate[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*(-3*(2*a*A*b + a^2*B - b^2*B)*EllipticE[(c + d*x)/2, 2] + (a^2*A + 3*A*b^2 + 6*a*b*B)*EllipticF[(c + d*x)/2, 2] + (a*(a*A + 3*(2*A*b + a*B)*Cos[c + d*x])*Sin[c + d*x])/Cos[c + d*x]^(3/2)))/(3*d)`

3.357.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3467, 27, 3042, 3500, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

$$\downarrow \text{3467}$$

$$\frac{2a^2A \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} - \frac{2}{3} \int \frac{3b^2B \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(2Ab + aB)}{2 \cos^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{27}$$

3.357. $\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \int \frac{3b^2 B \cos^2(c+dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c+dx) + 3a(2Ab + aB)}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{3b^2 B \sin(c+dx + \frac{\pi}{2})^2 + (Aa^2 + 6bBa + 3Ab^2) \sin(c+dx + \frac{\pi}{2}) + 3a(2Ab + aB)}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3500

$$\frac{1}{3} \left(\int \frac{Aa^2 + 6bBa + 3Ab^2 + 3(b^2 B - a(2Ab + aB)) \cos(c+dx)}{2\sqrt{\cos(c+dx)}} dx + \frac{6a(aB + 2Ab) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{3} \left(\int \frac{Aa^2 + 6bBa + 3Ab^2 + 3(b^2 B - a(2Ab + aB)) \cos(c+dx)}{\sqrt{\cos(c+dx)}} dx + \frac{6a(aB + 2Ab) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{Aa^2 + 6bBa + 3Ab^2 + 3(b^2 B - a(2Ab + aB)) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{6a(aB + 2Ab) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3227

$$\frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3(b^2 B - a(aB + 2Ab)) \int \sqrt{\cos(c+dx)} dx + \frac{6a(aB + 2Ab) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

3.357. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + 3(b^2 B - a(aB + 2Ab)) \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx + \frac{6a(aB + 2Ab)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3119

$$\frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{6(b^2 B - a(aB + 2Ab)) E(\frac{1}{2}(c+dx)|2)}{d} + \frac{6a(aB + 2Ab)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2(a^2 A + 6abB + 3Ab^2) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6(b^2 B - a(aB + 2Ab)) E(\frac{1}{2}(c+dx)|2)}{d} + \frac{6a(aB + 2Ab)}{d\sqrt{\cos(c+dx)}} \right) + \frac{2a^2 A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

input `Int[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*a^2*A*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((6*(b^2*B - a*(2*A*b + a*B))*EllipticE[(c + d*x)/2, 2])/d + (2*(a^2*A + 3*A*b^2 + 6*a*b*B)*EllipticF[(c + d*x)/2, 2])/d + (6*a*(2*A*b + a*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/3`

3.357.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3467 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C))*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

3.357.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 10.72 (sec) , antiderivative size = 574, normalized size of antiderivative = 4.56

3.357.
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^2(c+dx)} dx$$

method	result
parts	$\frac{2(Ab^2+2Bab) \operatorname{am}^{-1}\left(\frac{dx}{2}+\frac{c}{2} \sqrt{2}\right)}{d} - \frac{2(2Aab+Ba^2) \left(-2 \cos\left(\frac{dx}{2}+\frac{c}{2}\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\right) \left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right) + \sqrt{\frac{1}{2}-\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}}$
default	$-\frac{\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(\frac{2Ab^2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)-2Bb^2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}}\right)}$

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 2*(A*b^2+2*B*a*b)/d*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2))-2*(2*A*a*b+B*a^
2)*(-2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*
c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d-2/3*A*a^2
*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellipti
cF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*x+1/2*c)^2
*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*((2*cos(1/2*d*x+1/2*c)^2-1)
*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3/2)/sin(1/2*d*x+1/2*c)/d+2*B*b^2*((2*
cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*
c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.357.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 255, normalized size of antiderivative = 2.02

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{2}(-i Aa^2 - 6i Bab - 3i Ab^2) \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \dots}{\dots}$$

3.357. $\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fricas")`

output `1/3*(sqrt(2)*(-I*A*a^2 - 6*I*B*a*b - 3*I*A*b^2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(I*A*a^2 + 6*I*B*a*b + 3*I*A*b^2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(I*B*a^2 + 2*I*A*a*b - I*B*b^2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(-I*B*a^2 - 2*I*A*a*b + I*B*b^2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(A*a^2 + 3*(B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^2)`

3.357.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output Timed out

3.357.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/cos(d*x + c)^(5/2), x)`

3.357. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

3.357.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/cos(d*x + c)^(5/2), x)`

3.357.9 Mupad [B] (verification not implemented)

Time = 2.60 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.54

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 A b^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{2 B b^2 E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} + \frac{4 B a b F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d}$$

$$+ \frac{2 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{4 A a b \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^(5/2),x)`

output $(2A^2b^2 \operatorname{ellipticF}(c/2 + (dx)/2, 2))/d + (2B^2b^2 \operatorname{ellipticE}(c/2 + (dx)/2, 2))/d + (4B^2ab \operatorname{ellipticF}(c/2 + (dx)/2, 2))/d + (2A^2a^2 \sin(c + dx) \operatorname{hypergeom}([-3/4, 1/2], 1/4, \cos(c + dx)^2))/(3d \cos(c + dx)^{3/2} (\sin(c + dx)^2)^{1/2}) + (2B^2a^2 \sin(c + dx) \operatorname{hypergeom}([-1/4, 1/2], 3/4, \cos(c + dx)^2))/(d \cos(c + dx)^{1/2} (\sin(c + dx)^2)^{1/2}) + (4A^2ab \sin(c + dx) \operatorname{hypergeom}([-1/4, 1/2], 3/4, \cos(c + dx)^2))/(d \cos(c + dx)^{1/2} (\sin(c + dx)^2)^{1/2})$

3.358
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.358.1 Optimal result 3338
 3.358.2 Mathematica [A] (verified) 3339
 3.358.3 Rubi [A] (verified) 3339
 3.358.4 Maple [B] (verified) 3343
 3.358.5 Fracas [C] (verification not implemented) 3344
 3.358.6 Sympy [F(-1)] 3345
 3.358.7 Maxima [F] 3345
 3.358.8 Giac [F] 3346
 3.358.9 Mupad [B] (verification not implemented) 3346

3.358.1 Optimal result

Integrand size = 33, antiderivative size = 172

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= -\frac{2(3a^2A + 5Ab^2 + 10abB) E(\frac{1}{2}(c + dx) | 2)}{5d}$$

$$+ \frac{2(2aAb + a^2B + 3b^2B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{2a(2Ab + aB) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{2(3a^2A + 5Ab^2 + 10abB) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

output

```
-2/5*(3*A*a^2+5*A*b^2+10*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(2*A*a*b+B*a^2+3*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*a^2*A*sin(d*x+c)/d/cos(d*x+c)^(5/2)+2/3*a*(2*A*b+B*a)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/5*(3*A*a^2+5*A*b^2+10*B*a*b)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.358.2 Mathematica [A] (verified)

Time = 1.35 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.02

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{-6(3a^2A + 5Ab^2 + 10abB) \cos^{\frac{3}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(2aAb + a^2B + 3b^2B) \cos^{\frac{3}{2}}(c + dx) \text{EllipticF}\left(\frac{c + dx}{2}, 2\right) + 20aAb \sin(c + dx) + 10a^2B \sin(c + dx) + 9a^2A \sin[2(c + dx)] + 15Ab^2 \sin[2(c + dx)] + 30aAbB \sin[2(c + dx)] + 6a^2A \tan(c + dx)}{(15d \cos(c + dx))^{\frac{3}{2}}}$$

input `Integrate[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(-6*(3*a^2*A + 5*A*b^2 + 10*a*b*B)*Cos[c + d*x]^(3/2)*EllipticE[(c + d*x)/2, 2] + 10*(2*a*A*b + a^2*B + 3*b^2*B)*Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + 20*a*A*b*Sin[c + d*x] + 10*a^2*B*Sin[c + d*x] + 9*a^2*A*Sin[2*(c + d*x)] + 15*A*b^2*Sin[2*(c + d*x)] + 30*a*b*B*Sin[2*(c + d*x)] + 6*a^2*A*Tan[c + d*x])/(15*d*Cos[c + d*x]^(3/2))`

3.358.3 Rubi [A] (verified)Time = 0.88 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.94, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.394$, Rules used = {3042, 3467, 27, 3042, 3500, 27, 3042, 3227, 3042, 3116, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{\frac{7}{2}}} dx$$

$$\downarrow \text{3467}$$

$$\frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} - \frac{2}{5} \int -\frac{5b^2B \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + 5a(2Ab + aB)}{2 \cos^{\frac{5}{2}}(c + dx)} dx$$

3.358. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned} & \downarrow 27 \\ & \frac{1}{5} \int \frac{5b^2 B \cos^2(c+dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c+dx) + 5a(2Ab + aB)}{\cos^{\frac{5}{2}}(c+dx)} dx + \\ & \quad \frac{2a^2 A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\ & \downarrow 3042 \\ & \frac{1}{5} \int \frac{5b^2 B \sin(c+dx + \frac{\pi}{2})^2 + (3Aa^2 + 10bBa + 5Ab^2) \sin(c+dx + \frac{\pi}{2}) + 5a(2Ab + aB)}{\sin(c+dx + \frac{\pi}{2})^{5/2}} dx + \\ & \quad \frac{2a^2 A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\ & \downarrow 3500 \\ & \frac{1}{5} \left(\frac{2}{3} \int \frac{3(3Aa^2 + 10bBa + 5Ab^2) + 5(Ba^2 + 2Aba + 3b^2 B) \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)} dx + \frac{10a(aB + 2Ab) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \\ & \quad \frac{2a^2 A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\ & \downarrow 27 \\ & \frac{1}{5} \left(\frac{1}{3} \int \frac{3(3Aa^2 + 10bBa + 5Ab^2) + 5(Ba^2 + 2Aba + 3b^2 B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx + \frac{10a(aB + 2Ab) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \\ & \quad \frac{2a^2 A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\ & \downarrow 3042 \\ & \frac{1}{5} \left(\frac{1}{3} \int \frac{3(3Aa^2 + 10bBa + 5Ab^2) + 5(Ba^2 + 2Aba + 3b^2 B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{3/2}} dx + \frac{10a(aB + 2Ab) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \\ & \quad \frac{2a^2 A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\ & \downarrow 3227 \\ & \frac{1}{5} \left(\frac{1}{3} \left(3(3a^2 A + 10abB + 5Ab^2) \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)} dx + 5(a^2 B + 2aAb + 3b^2 B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx \right) + \frac{10a(aB + 2Ab) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \\ & \quad \frac{2a^2 A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\ & \downarrow 3042 \end{aligned}$$

3.358. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{1}{3} \left(3(3a^2A + 10abB + 5Ab^2) \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx + 5(a^2B + 2aAb + 3b^2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3116}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^2B + 2aAb + 3b^2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(3a^2A + 10abB + 5Ab^2) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos} \right) \right) + \frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3042}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^2B + 2aAb + 3b^2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(3a^2A + 10abB + 5Ab^2) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\sin} \right) \right) + \frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3119}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^2B + 2aAb + 3b^2B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(3a^2A + 10abB + 5Ab^2) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx))}{d} \right) \right) + \frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3120}$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{10(a^2B + 2aAb + 3b^2B) \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} + 3(3a^2A + 10abB + 5Ab^2) \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx))}{d} \right) \right) + \frac{2a^2A \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

input `Int[((a + b*cos[c + d*x])^2*(A + B*cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

```
output (2*a^2*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((10*a*(2*A*b + a*B)*Sin
[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((10*(2*a*A*b + a^2*B + 3*b^2*B)*Ell
ipticF[(c + d*x)/2, 2])/d + 3*(3*a^2*A + 5*A*b^2 + 10*a*b*B)*((-2*Elliptic
E[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/3)/5
```

3.358.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3116 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) I
nt[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3227 Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3467 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*(b*c - a*d)^2*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 - d^2))), x] - Simp[1/(d^2*(n + 1)*(c^2 - d^2)) Int[(c + d*Sin[e + f*x])^(n + 1)*Simp[d*(n + 1)*(B*(b*c - a*d)^2 - A*d*(a^2*c + b^2*c - 2*a*b*d)) - ((B*c - A*d)*(a^2*d^2*(n + 2) + b^2*(c^2 + d^2*(n + 1))) + 2*a*b*d*(A*c*d*(n + 2) - B*(c^2 + d^2*(n + 1)))*Sin[e + f*x] - b^2*B*d*(n + 1)*(c^2 - d^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

```
rule 3500 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]
```

3.358.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. $2(208) = 416$.

Time = 13.07 (sec) , antiderivative size = 723, normalized size of antiderivative = 4.20

method	result	size
default	Expression too large to display	723
parts	Expression too large to display	799

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+2/5*A*a^2/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+2*b*(A*b+2*B*a)/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))+2*a*(2*A*b+B*a)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d

```

3.358.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.66

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{5\sqrt{2}(iBa^2 + 2iAab + 3iBb^2) \cos(dx + c)^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \dots}{\dots}$$

input

```

integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm
m="fricas")

```

output `-1/15*(5*sqrt(2)*(I*B*a^2 + 2*I*A*a*b + 3*I*B*b^2)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-I*B*a^2 - 2*I*A*a*b - 3*I*B*b^2)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(3*I*A*a^2 + 10*I*B*a*b + 5*I*A*b^2)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-3*I*A*a^2 - 10*I*B*a*b - 5*I*A*b^2)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*A*a^2 + 3*(3*A*a^2 + 10*B*a*b + 5*A*b^2)*cos(d*x + c)^2 + 5*(B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.358.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.358.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{7}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/cos(d*x + c)^(7/2), x)`

3.358.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/cos(d*x + c)^(7/2), x)`

3.358.9 Mupad [B] (verification not implemented)

Time = 2.89 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.32

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{6 A a^2 \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right) + 30 A b^2 \cos(c + dx)^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right) + 15 d \cos(c + dx)^{5/2} \sqrt{1 - \cos(c + dx)}}{d} + \frac{2 B b^2 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) + 2 B a^2 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}} + \frac{4 B a b \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/cos(c + d*x)^(7/2),x)`

output `(6*A*a^2*sin(c + d*x)*hypergeom([-5/4, 1/2], -1/4, cos(c + d*x)^2) + 30*A*b^2*cos(c + d*x)^2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2) + 20*A*a*b*cos(c + d*x)*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(15*d*cos(c + d*x)^(5/2)*(1 - cos(c + d*x)^2)^(1/2)) + (2*B*b^2*ellipticF(c/2 + (d*x)/2, 2))/d + (2*B*a^2*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, cos(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2)) + (4*B*a*b*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))`

3.358. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

3.359 $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx$

3.359.1 Optimal result	3347
3.359.2 Mathematica [A] (verified)	3348
3.359.3 Rubi [A] (verified)	3348
3.359.4 Maple [B] (verified)	3353
3.359.5 Fricas [C] (verification not implemented)	3354
3.359.6 Sympy [F(-1)]	3355
3.359.7 Maxima [F]	3355
3.359.8 Giac [F]	3356
3.359.9 Mupad [B] (verification not implemented)	3357

3.359.1 Optimal result

Integrand size = 33, antiderivative size = 305

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3(A+B \cos(c+dx)) dx \\ &= \frac{2(27a^2Ab + 7Ab^3 + 9a^3B + 21ab^2B) E(\frac{1}{2}(c+dx)|2)}{15d} \\ &+ \frac{2(77a^3A + 165aAb^2 + 165a^2bB + 45b^3B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{231d} \\ &+ \frac{2(77a^3A + 165aAb^2 + 165a^2bB + 45b^3B) \sqrt{\cos(c+dx)} \sin(c+dx)}{231d} \\ &+ \frac{2(27a^2Ab + 7Ab^3 + 9a^3B + 21ab^2B) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{45d} \\ &+ \frac{2b(33aAb + 26a^2B + 9b^2B) \cos^{\frac{5}{2}}(c+dx) \sin(c+dx)}{77d} \\ &+ \frac{2b^2(11Ab + 15aB) \cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{99d} \\ &+ \frac{2bB \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2 \sin(c+dx)}{11d} \end{aligned}$$

output
$$\frac{2}{15} \cdot (27Aa^2b + 7Ab^3 + 9B^2a^3 + 21B^2ab^2) \cdot (\cos(\frac{1}{2}dx + \frac{1}{2}c))^2 \cdot (\cos(\frac{1}{2}dx + \frac{1}{2}c)) \cdot \text{EllipticE}(\sin(\frac{1}{2}dx + \frac{1}{2}c), 2^{1/2}) / d + \frac{2}{231} \cdot (77Aa^3 + 165Aa^2b + 165B^2a^2b + 45B^2b^3) \cdot (\cos(\frac{1}{2}dx + \frac{1}{2}c))^2 \cdot (\cos(\frac{1}{2}dx + \frac{1}{2}c)) \cdot \text{EllipticF}(\sin(\frac{1}{2}dx + \frac{1}{2}c), 2^{1/2}) / d + \frac{2}{45} \cdot (27Aa^2b + 7Ab^3 + 9B^2a^3 + 21B^2ab^2) \cdot \cos(dx + c)^{3/2} \cdot \sin(dx + c) / d + \frac{2}{77} \cdot b \cdot (33Aa^2b + 26B^2a^2 + 9B^2b^2) \cdot \cos(dx + c)^{5/2} \cdot \sin(dx + c) / d + \frac{2}{99} \cdot b^2 \cdot (11Ab + 15B^2a) \cdot \cos(dx + c)^{7/2} \cdot \sin(dx + c) / d + \frac{2}{11} \cdot b \cdot B \cdot \cos(dx + c)^{5/2} \cdot (a + b \cdot \cos(dx + c))^2 \cdot \sin(dx + c) / d + \frac{2}{231} \cdot (77Aa^3 + 165Aa^2b + 165B^2a^2b + 45B^2b^3) \cdot \sin(dx + c) \cdot \cos(dx + c)^{1/2} / d$$

3.359.2 Mathematica [A] (verified)

Time = 2.54 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.77

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{3696(27a^2Ab + 7Ab^3 + 9a^3B + 21ab^2B) E(\frac{1}{2}(c + dx) | 2) + 240(77a^3A + 165aAb^2 + 165a^2bB + 45b^3B) E(\frac{1}{2}(c + dx) | 2) + 2(154(108a^2Ab + 43Ab^3 + 36a^3B + 129a^2b^2B) \cos(c + dx) + 180b(33a^2Ab + 33a^2B + 16b^2B) \cos(2(c + dx)) + 770b^2(Ab + 3a^2B) \cos(3(c + dx)) + 15(616a^3A + 1716a^2Ab^2 + 1716a^2b^2B + 531b^3B + 21b^3B \cos(4(c + dx))) \sin(c + dx))}{(27720 \cdot d)}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output
$$(3696 \cdot (27a^2Ab + 7Ab^3 + 9a^3B + 21ab^2B) \cdot \text{EllipticE}[(c + dx)/2, 2] + 240 \cdot (77a^3A + 165aAb^2 + 165a^2bB + 45b^3B) \cdot \text{EllipticF}[(c + dx)/2, 2] + 2 \cdot \text{Sqrt}[\text{Cos}[c + d*x]] \cdot (154 \cdot (108a^2Ab + 43Ab^3 + 36a^3B + 129a^2b^2B) \cdot \text{Cos}[c + d*x] + 180 \cdot b \cdot (33a^2Ab + 33a^2B + 16b^2B) \cdot \text{Cos}[2 \cdot (c + d*x)] + 770 \cdot b^2 \cdot (Ab + 3a^2B) \cdot \text{Cos}[3 \cdot (c + d*x)] + 15 \cdot (616a^3A + 1716a^2Ab^2 + 1716a^2b^2B + 531b^3B + 21b^3B \cdot \text{Cos}[4 \cdot (c + d*x)])) \cdot \text{Sin}[c + d*x]) / (27720 \cdot d)$$

3.359.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 272, normalized size of antiderivative = 0.89, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3469, 27, 3042, 3512, 27, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.359. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx \\
& \quad \downarrow \text{3042} \\
& \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^3\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\
& \quad \downarrow \text{3469} \\
& \frac{2}{11}\int\frac{1}{2}\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))\left(b(11Ab+15aB)\cos^2(c+dx)+(9Bb^2+11a(2Ab+aB))\cos(c+dx)+a(11aA+5bB)\right)dx+ \\
& \quad \frac{2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2}{11d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{11}\int\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))\left(b(11Ab+15aB)\cos^2(c+dx)+(9Bb^2+11a(2Ab+aB))\cos(c+dx)+a(11aA+5bB)\right)dx+ \\
& \quad \frac{2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2}{11d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{11}\int\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)\left(b(11Ab+15aB)\sin\left(c+dx+\frac{\pi}{2}\right)^2+(9Bb^2+11a(2Ab+aB))\sin\left(c+dx+\frac{\pi}{2}\right)+a(11aA+5bB)\right)dx+ \\
& \quad \frac{2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2}{11d} \\
& \quad \downarrow \text{3512} \\
& \frac{1}{11}\left(\frac{2}{9}\int\frac{1}{2}\cos^{\frac{3}{2}}(c+dx)\left(9(11aA+5bB)a^2+9b(26Ba^2+33Aba+9b^2B)\cos^2(c+dx)+11(9Ba^3+27Aba^2+21b^2A)\cos(c+dx)+11aA+5bB\right)dx+ \right. \\
& \quad \left. \frac{2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2}{11d}\right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{11}\left(\frac{1}{9}\int\cos^{\frac{3}{2}}(c+dx)\left(9(11aA+5bB)a^2+9b(26Ba^2+33Aba+9b^2B)\cos^2(c+dx)+11(9Ba^3+27Aba^2+21b^2A)\cos(c+dx)+11aA+5bB\right)dx+ \right. \\
& \quad \left. \frac{2bB\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2}{11d}\right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.359. $\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx$

$$\frac{1}{11} \left(\frac{1}{9} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} \left(9(11aA + 5bB)a^2 + 9b(26Ba^2 + 33Aba + 9b^2B) \sin \left(c + dx + \frac{\pi}{2} \right)^2 + 11(9Ba^3 + 2bB \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2 \right) \right)$$

\downarrow 3502

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2}{7} \int \frac{1}{2} \cos^{\frac{3}{2}}(c + dx) (9(77Aa^3 + 165bBa^2 + 165Ab^2a + 45b^3B) + 77(9Ba^3 + 27Aba^2 + 21b^2Ba + 7Ab^3) \cos(c + dx) \right) \right)$$

\downarrow 27

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \int \cos^{\frac{3}{2}}(c + dx) (9(77Aa^3 + 165bBa^2 + 165Ab^2a + 45b^3B) + 77(9Ba^3 + 27Aba^2 + 21b^2Ba + 7Ab^3) \cos(c + dx) \right) \right)$$

\downarrow 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} \left(9(77Aa^3 + 165bBa^2 + 165Ab^2a + 45b^3B) + 77(9Ba^3 + 27Aba^2 + 21b^2Ba + 7Ab^3) \sin \left(c + dx + \frac{\pi}{2} \right)^2 \right) \right)$$

\downarrow 3227

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \left(9(77a^3A + 165a^2bB + 165aAb^2 + 45b^3B) \int \cos^{\frac{3}{2}}(c + dx) dx + 77(9a^3B + 27a^2Ab + 21ab^2B + 7Ab^3) \cos(c + dx) \right) \right)$$

\downarrow 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \left(9(77a^3A + 165a^2bB + 165aAb^2 + 45b^3B) \int \sin \left(c + dx + \frac{\pi}{2} \right)^{3/2} dx + 77(9a^3B + 27a^2Ab + 21ab^2B + 7Ab^3) \sin \left(c + dx + \frac{\pi}{2} \right)^2 \right) \right)$$

\downarrow 3115

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \left(77(9a^3B + 27a^2Ab + 21ab^2B + 7Ab^3) \left(\frac{3}{5} \int \sqrt{\cos(c+dx)} dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + 9(77a^3A + 165a^2bB + 165aAb^2 + 45b^3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right) \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a + b \cos(c+dx))^2}{11d}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \left(77(9a^3B + 27a^2Ab + 21ab^2B + 7Ab^3) \left(\frac{3}{5} \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx + \frac{2 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} \right) + 9(77a^3A + 165a^2bB + 165aAb^2 + 45b^3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right) \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a + b \cos(c+dx))^2}{11d}$$

↓ 3119

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{1}{7} \left(9(77a^3A + 165a^2bB + 165aAb^2 + 45b^3B) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) \right) \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a + b \cos(c+dx))^2}{11d}$$

↓ 3120

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{18b(26a^2B + 33aAb + 9b^2B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{7d} + \frac{1}{7} \left(77(9a^3B + 27a^2Ab + 21ab^2B + 7Ab^3) \left(\frac{6E}{5d} \right) \right) \right) \right) + \frac{2bB \sin(c+dx) \cos^{\frac{5}{2}}(c+dx) (a + b \cos(c+dx))^2}{11d}$$

input `Int[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(2*b*B*Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(11*d) + ((2*b^2*(11*A*b + 15*a*B)*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(9*d) + ((18*b*(3*3*a*A*b + 26*a^2*B + 9*b^2*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(7*d) + (9*(77*a^3*A + 165*a*A*b^2 + 165*a^2*b*B + 45*b^3*B))*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d)) + 77*(27*a^2*A*b + 7*A*b^3 + 9*a^3*B + 21*a*b^2*B))*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d)))/7)/9)/11`

3.359.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`
- rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]*(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d))*(m + 3))*Sin[e + f*x]^2
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

3.359.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 824 vs. $2(333) = 666$.

Time = 18.44 (sec) , antiderivative size = 825, normalized size of antiderivative = 2.70

method	result	size
default	Expression too large to display	825
parts	Expression too large to display	1063

```
input int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```


output

```

-2/3465*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(20160*B*c
os(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^12*b^3+(-12320*A*b^3-36960*B*a*b^2-50
400*B*b^3)*sin(1/2*d*x+1/2*c)^10*cos(1/2*d*x+1/2*c)+(23760*A*a*b^2+24640*A
*b^3+23760*B*a^2*b+73920*B*a*b^2+56880*B*b^3)*sin(1/2*d*x+1/2*c)^8*cos(1/2
*d*x+1/2*c)+(-16632*A*a^2*b-35640*A*a*b^2-22792*A*b^3-5544*B*a^3-35640*B*a
^2*b-68376*B*a*b^2-34920*B*b^3)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(4
620*A*a^3+16632*A*a^2*b+27720*A*a*b^2+10472*A*b^3+5544*B*a^3+27720*B*a^2*b
+31416*B*a*b^2+13860*B*b^3)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-2310
*A*a^3-4158*A*a^2*b-7920*A*a*b^2-1848*A*b^3-1386*B*a^3-7920*B*a^2*b-5544*B
*a*b^2-2790*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+1155*A*a^3*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/
2*d*x+1/2*c),2^(1/2))+2475*A*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2
*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-6237*A*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*
d*x+1/2*c),2^(1/2))*a^2*b-1617*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d
*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^3+2475*B*a^2*
b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(
cos(1/2*d*x+1/2*c),2^(1/2))+675*B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(
1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2079*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(co...

```

3.359.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.17

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx$$

$$= \frac{2(315Bb^3\cos(dx+c)^4 + 1155Aa^3 + 2475Ba^2b + 2475Aab^2 + 675Bb^3 + 385(3Bab^2 + Ab^3)\cos(dx+c)}{...}$$

input

```

integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm
m="fracas")

```

output `1/3465*(2*(315*B*b^3*cos(d*x + c)^4 + 1155*A*a^3 + 2475*B*a^2*b + 2475*A*a*b^2 + 675*B*b^3 + 385*(3*B*a*b^2 + A*b^3))*cos(d*x + c)^3 + 135*(11*B*a^2*b + 11*A*a*b^2 + 3*B*b^3))*cos(d*x + c)^2 + 77*(9*B*a^3 + 27*A*a^2*b + 21*B*a*b^2 + 7*A*b^3))*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 15*sqrt(2)*(77*I*A*a^3 + 165*I*B*a^2*b + 165*I*A*a*b^2 + 45*I*B*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 15*sqrt(2)*(-77*I*A*a^3 - 165*I*B*a^2*b - 165*I*A*a*b^2 - 45*I*B*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 231*sqrt(2)*(-9*I*B*a^3 - 27*I*A*a^2*b - 21*I*B*a*b^2 - 7*I*A*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 231*sqrt(2)*(9*I*B*a^3 + 27*I*A*a^2*b + 21*I*B*a*b^2 + 7*I*A*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.359.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+b*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.359.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*cos(d*x + c)^(3/2), x)`

3.359. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.359.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*cos(d*x + c)^(3/2), x)`

3.359.9 Mupad [B] (verification not implemented)

Time = 2.08 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.19

$$\begin{aligned}
& \int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx \\
&= \frac{A a^3 \left(\frac{2\sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3} \right)}{d} \\
&\quad - \frac{2 B a^3 \cos(c+dx)^{7/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7 d \sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2 A b^3 \cos(c+dx)^{11/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11 d \sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2 B b^3 \cos(c+dx)^{13/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{13}{4}; \frac{17}{4}; \cos(c+dx)^2\right)}{13 d \sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{6 A a^2 b \cos(c+dx)^{7/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)}{7 d \sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2 A a b^2 \cos(c+dx)^{9/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{3 d \sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{2 B a^2 b \cos(c+dx)^{9/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)}{3 d \sqrt{\sin(c+dx)^2}} \\
&\quad - \frac{6 B a b^2 \cos(c+dx)^{11/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)}{11 d \sqrt{\sin(c+dx)^2}}
\end{aligned}$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3,x)`

output $(Aa^3((2\cos(c+dx)^{1/2}\sin(c+dx))/3 + (2\text{ellipticF}(c/2 + (dx)/2, 2))/3))/d - (2Ba^3\cos(c+dx)^{7/2}\sin(c+dx)\text{hypergeom}([1/2, 7/4], 11/4, \cos(c+dx)^2))/(7d(\sin(c+dx)^2)^{1/2}) - (2Ab^3\cos(c+dx)^{11/2}\sin(c+dx)\text{hypergeom}([1/2, 11/4], 15/4, \cos(c+dx)^2))/(11d(\sin(c+dx)^2)^{1/2}) - (2Bb^3\cos(c+dx)^{13/2}\sin(c+dx)\text{hypergeom}([1/2, 13/4], 17/4, \cos(c+dx)^2))/(13d(\sin(c+dx)^2)^{1/2}) - (6Aa^2b\cos(c+dx)^{7/2}\sin(c+dx)\text{hypergeom}([1/2, 7/4], 11/4, \cos(c+dx)^2))/(7d(\sin(c+dx)^2)^{1/2}) - (2Aa^2b^2\cos(c+dx)^{9/2}\sin(c+dx)\text{hypergeom}([1/2, 9/4], 13/4, \cos(c+dx)^2))/(3d(\sin(c+dx)^2)^{1/2}) - (2Ba^2b\cos(c+dx)^{9/2}\sin(c+dx)\text{hypergeom}([1/2, 9/4], 13/4, \cos(c+dx)^2))/(3d(\sin(c+dx)^2)^{1/2}) - (6Bab^2\cos(c+dx)^{11/2}\sin(c+dx)\text{hypergeom}([1/2, 11/4], 15/4, \cos(c+dx)^2))/(11d(\sin(c+dx)^2)^{1/2})$

3.360 $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$

3.360.1 Optimal result	3359
3.360.2 Mathematica [A] (verified)	3360
3.360.3 Rubi [A] (verified)	3360
3.360.4 Maple [B] (verified)	3365
3.360.5 Fricas [C] (verification not implemented)	3366
3.360.6 Sympy [F(-1)]	3366
3.360.7 Maxima [F]	3367
3.360.8 Giac [F]	3367
3.360.9 Mupad [B] (verification not implemented)	3368

3.360.1 Optimal result

Integrand size = 33, antiderivative size = 255

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$= \frac{2(15a^3A + 27aAb^2 + 27a^2bB + 7b^3B) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d}$$

$$+ \frac{2(21a^2Ab + 5Ab^3 + 7a^3B + 15ab^2B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d}$$

$$+ \frac{2(21a^2Ab + 5Ab^3 + 7a^3B + 15ab^2B) \sqrt{\cos(c + dx)} \sin(c + dx)}{21d}$$

$$+ \frac{2b(27aAb + 22a^2B + 7b^2B) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{45d}$$

$$+ \frac{2b^2(9Ab + 13aB) \cos^{\frac{5}{2}}(c + dx) \sin(c + dx)}{63d}$$

$$+ \frac{2bB \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2 \sin(c + dx)}{9d}$$

```
output 2/15*(15*A*a^3+27*A*a*b^2+27*B*a^2*b+7*B*b^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/21*(21*A*a^2
*b+5*A*b^3+7*B*a^3+15*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/
2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/45*b*(27*A*a*b+22*B*a^2+7*B
*b^2)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/63*b^2*(9*A*b+13*B*a)*cos(d*x+c)^(5/
2)*sin(d*x+c)/d+2/9*b*B*cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^2*sin(d*x+c)/d+2
/21*(21*A*a^2*b+5*A*b^3+7*B*a^3+15*B*a*b^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.360.2 Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.77

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx$$

$$= \frac{84(15a^3A+27aAb^2+27a^2bB+7b^3B)E\left(\frac{1}{2}(c+dx)\middle|2\right)+60(21a^2Ab+5Ab^3+7a^3B+15ab^2B)\text{EllipticF}\left(\frac{c+dx}{2}\middle|2\right)+\sqrt{\cos(c+dx)}(7b(108aAb+108a^2B+43b^2B)\cos(c+dx)+5(252a^2Ab+78Ab^3+84a^3B+234ab^2B+18b^2(Ab+3aB))\cos(2(c+dx))+7b^3B\cos(3(c+dx)))\sin(c+dx)}{(630d)}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(84*(15*a^3*A + 27*a*A*b^2 + 27*a^2*b*B + 7*b^3*B)*EllipticE[(c + d*x)/2, 2] + 60*(21*a^2*A*b + 5*A*b^3 + 7*a^3*B + 15*a*b^2*B)*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(7*b*(108*a*A*b + 108*a^2*B + 43*b^2*B)*Cos[c + d*x] + 5*(252*a^2*A*b + 78*A*b^3 + 84*a^3*B + 234*a*b^2*B + 18*b^2*(A*b + 3*a*B))*Cos[2*(c + d*x)] + 7*b^3*B*Cos[3*(c + d*x)]))*Sin[c + d*x])/(630*d)`

3.360.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.96, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3469, 27, 3042, 3512, 27, 3042, 3502, 27, 3042, 3227, 3042, 3115, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^3\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx$$

$$\downarrow \text{3469}$$

$$\frac{2}{9}\int\frac{1}{2}\sqrt{\cos(c+dx)}(a+b\cos(c+dx))(b(9Ab+13aB)\cos^2(c+dx)+(7Bb^2+9a(2Ab+aB))\cos(c+dx)+3a(3aA+bB))dx+\frac{2bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2}{9d}$$

3.360. $\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{9} \int \sqrt{\cos(c+dx)} (a+b \cos(c+dx)) (b(9Ab+13aB) \cos^2(c+dx) + (7Bb^2+9a(2Ab+aB)) \cos(c+dx) + 3a(3aA+bB)) dx + \\
& \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^2}{9d} \\
& \downarrow 3042 \\
& \frac{1}{9} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(a+b \sin\left(c+dx+\frac{\pi}{2}\right)\right) \left(b(9Ab+13aB) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + (7Bb^2+9a(2Ab+aB))\right) dx + \\
& \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^2}{9d} \\
& \downarrow 3512 \\
& \frac{1}{9} \left(\frac{2}{7} \int \frac{1}{2} \sqrt{\cos(c+dx)} (21(3aA+bB)a^2 + 7b(22Ba^2+27Aba+7b^2B)) \cos^2(c+dx) + 9(7Ba^3+21Aba^2+15b^2Ba) \right) dx + \\
& \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^2}{9d} \\
& \downarrow 27 \\
& \frac{1}{9} \left(\frac{1}{7} \int \sqrt{\cos(c+dx)} (21(3aA+bB)a^2 + 7b(22Ba^2+27Aba+7b^2B)) \cos^2(c+dx) + 9(7Ba^3+21Aba^2+15b^2Ba) \right) dx + \\
& \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^2}{9d} \\
& \downarrow 3042 \\
& \frac{1}{9} \left(\frac{1}{7} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(21(3aA+bB)a^2 + 7b(22Ba^2+27Aba+7b^2B)\right) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + 9(7Ba^3+21Aba^2+15b^2Ba) \right) dx + \\
& \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^2}{9d} \\
& \downarrow 3502 \\
& \frac{1}{9} \left(\frac{1}{7} \left(\frac{2}{5} \int \frac{3}{2} \sqrt{\cos(c+dx)} (7(15Aa^3+27bBa^2+27Ab^2a+7b^3B)) + 15(7Ba^3+21Aba^2+15b^2Ba+5Ab^3) \cos(c+dx) \right) \right) dx + \\
& \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^2}{9d} \\
& \downarrow 27
\end{aligned}$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \int \sqrt{\cos(c+dx)} (7(15Aa^3 + 27bBa^2 + 27Ab^2a + 7b^3B) + 15(7Ba^3 + 21Aba^2 + 15b^2Ba + 5Ab^3)) \cos(c+dx) \right. \right. \\ \left. \left. \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right. \right. \\ \left. \left. \downarrow 3042 \right. \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} (7(15Aa^3 + 27bBa^2 + 27Ab^2a + 7b^3B) + 15(7Ba^3 + 21Aba^2 + 15b^2Ba + 5Ab^3)) \right. \right. \\ \left. \left. \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right. \right. \\ \left. \left. \downarrow 3227 \right. \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \left(15(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \int \cos^{\frac{3}{2}}(c+dx) dx + 7(15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \int \sqrt{\cos(c+dx)} \right. \right. \right. \\ \left. \left. \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right. \right. \\ \left. \left. \downarrow 3042 \right. \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \left(7(15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + 15(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \int \sqrt{\cos(c+dx)} \right. \right. \right. \\ \left. \left. \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right. \right. \\ \left. \left. \downarrow 3115 \right. \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \left(7(15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + 15(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \int \sqrt{\cos(c+dx)} \right. \right. \right. \\ \left. \left. \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right. \right. \\ \left. \left. \downarrow 3042 \right. \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \left(7(15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + 15(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \int \sqrt{\cos(c+dx)} \right. \right. \right. \\ \left. \left. \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right. \right. \\ \left. \left. \downarrow 3119 \right. \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{3}{5} \left(15(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right) + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right) \right) \right)$$

↓ 3120

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{14b(22a^2B + 27aAb + 7b^2B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5d} + \frac{3}{5} \left(\frac{14(15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) E(\frac{1}{2})}{d} + \frac{2bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}{9d} \right) \right) \right)$$

input `Int[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]),x]`

output `(2*b*B*Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(9*d) + ((2*b^2*(9*A*b + 13*a*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(7*d) + ((14*b*(27*a*A*b + 22*a^2*B + 7*b^2*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*d) + (3*((14*(15*a^3*A + 27*a*A*b^2 + 27*a^2*b*B + 7*b^3*B)*EllipticE[(c + d*x)/2, 2])/d + 15*(21*a^2*A*b + 5*A*b^3 + 7*a^3*B + 15*a*b^2*B)*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))))/5)/7)/9`

3.360.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3227 $\text{Int}[(b_.)*\sin[(e_.) + (f_.)(x_)]^{(m)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[c \text{ Int}[(b*\sin[e + f*x])^m, x], x] + \text{Simp}[d/b \text{ Int}[(b*\sin[e + f*x])^{(m+1)}, x], x] /; \text{FreeQ}[\{b, c, d, e, f, m\}, x]$

rule 3469 $\text{Int}[(a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)]^{(m)}*((A_.) + (B_.)*\sin[(e_.) + (f_.)(x_)]^{(n)}), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cos}[e + f*x]*(a + b*\sin[e + f*x])^{(m-1)}*((c + d*\sin[e + f*x])^{(n+1)}/(d*f*(m+n+1))), x] + \text{Simp}[1/(d*(m+n+1)) \text{ Int}[(a + b*\sin[e + f*x])^{(m-2)}*(c + d*\sin[e + f*x])^n*\text{Simp}[a^2*A*d*(m+n+1) + b*B*(b*c*(m-1) + a*d*(n+1)) + (a*d*(2*A*b + a*B)*(m+n+1) - b*B*(a*c - b*d*(m+n)))*\sin[e + f*x] + b*(A*b*d*(m+n+1) - B*(b*c*m - a*d*(2*m+n)))*\sin[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[m, 1] \&\& !(\text{IGtQ}[n, 1] \&\& (!\text{IntegerQ}[m] || (\text{EqQ}[a, 0] \&\& \text{NeQ}[c, 0])))$

rule 3502 $\text{Int}[(a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)]^{(m)}*((A_.) + (B_.)*\sin[(e_.) + (f_.)(x_)] + (C_.)*\sin[(e_.) + (f_.)(x_)]^2), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cos}[e + f*x]*(a + b*\sin[e + f*x])^{(m+1)}/(b*f*(m+2)), x] + \text{Simp}[1/(b*(m+2)) \text{ Int}[(a + b*\sin[e + f*x])^m*\text{Simp}[A*b*(m+2) + b*C*(m+1) + (b*B*(m+2) - a*C)*\sin[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, e, f, A, B, C, m\}, x] \&\& !\text{LtQ}[m, -1]$

rule 3512 $\text{Int}[(a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)]^{(m)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)]^{(A_.) + (B_.)*\sin[(e_.) + (f_.)(x_)] + (C_.)*\sin[(e_.) + (f_.)(x_)]^2), x_Symbol] \rightarrow \text{Simp}[(-C)*d*\text{Cos}[e + f*x]*\sin[e + f*x]*((a + b*\sin[e + f*x])^{(m+1)}/(b*f*(m+3))), x] + \text{Simp}[1/(b*(m+3)) \text{ Int}[(a + b*\sin[e + f*x])^m*\text{Simp}[a*C*d + A*b*c*(m+3) + b*(B*c*(m+3) + d*(C*(m+2) + A*(m+3)))*\sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m+3))*\sin[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& !\text{LtQ}[m, -1]$

3.360.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 744 vs. $2(287) = 574$.

Time = 16.22 (sec) , antiderivative size = 745, normalized size of antiderivative = 2.92

method	result	size
default	Expression too large to display	745
parts	Expression too large to display	971

```
input int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -2/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*B*co
s(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b^3+(720*A*b^3+2160*B*a*b^2+2240*B*
b^3)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1512*A*a*b^2-1080*A*b^3-151
2*B*a^2*b-3240*B*a*b^2-2072*B*b^3)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)
+(1260*A*a^2*b+1512*A*a*b^2+840*A*b^3+420*B*a^3+1512*B*a^2*b+2520*B*a*b^2+
952*B*b^3)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-630*A*a^2*b-378*A*a*b
^2-240*A*b^3-210*B*a^3-378*B*a^2*b-720*B*a*b^2-168*B*b^3)*sin(1/2*d*x+1/2*
c)^2*cos(1/2*d*x+1/2*c)+315*A*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/
2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+75*A*b^3*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))-315*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^3-567*A*(sin(1/2*
d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x
+1/2*c),2^(1/2))*a*b^2+105*B*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d
*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+225*B*a*b^2*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))-567*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b-147*B*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d
*x+1/2*c),2^(1/2))*b^3)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^...
```

3.360.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.26

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx$$

$$= \frac{2(35Bb^3\cos(dx+c)^3 + 105Ba^3 + 315Aa^2b + 225Bab^2 + 75Ab^3 + 45(3Bab^2 + Ab^3)\cos(dx+c)^2 + 75A^2b^2 + 15A^2b\cos(dx+c) + 15A^2b^2\cos^2(dx+c) + 15A^2b^2\cos^3(dx+c) + 15A^2b^2\cos^4(dx+c) + 15A^2b^2\cos^5(dx+c) + 15A^2b^2\cos^6(dx+c) + 15A^2b^2\cos^7(dx+c) + 15A^2b^2\cos^8(dx+c) + 15A^2b^2\cos^9(dx+c))}{d}$$

```
input integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm
m="fricas")
```

```
output 1/315*(2*(35*B*b^3*cos(d*x + c)^3 + 105*B*a^3 + 315*A*a^2*b + 225*B*a*b^2
+ 75*A*b^3 + 45*(3*B*a*b^2 + A*b^3)*cos(d*x + c)^2 + 7*(27*B*a^2*b + 27*A*
a*b^2 + 7*B*b^3)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 15*sqrt(2
)*(7*I*B*a^3 + 21*I*A*a^2*b + 15*I*B*a*b^2 + 5*I*A*b^3)*weierstrassPInvers
e(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 15*sqrt(2)*(-7*I*B*a^3 - 21*I*A*
a^2*b - 15*I*B*a*b^2 - 5*I*A*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c)
- I*sin(d*x + c)) - 21*sqrt(2)*(-15*I*A*a^3 - 27*I*B*a^2*b - 27*I*A*a*b^2
- 7*I*B*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c)
+ I*sin(d*x + c))) - 21*sqrt(2)*(15*I*A*a^3 + 27*I*B*a^2*b + 27*I*A*a*b^
2 + 7*I*B*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x +
c) - I*sin(d*x + c))))/d
```

3.360.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(1/2)*(a+b*cos(d*x+c))**3*(A+B*cos(d*x+c)),x)
```

```
output Timed out
```

3.360.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx \\ &= \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^3\sqrt{\cos(dx+c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sqrt(cos(d*x + c)), x)`

3.360.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx \\ &= \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^3\sqrt{\cos(dx+c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^3*(A+B*cos(d*x+c)),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sqrt(cos(d*x + c)), x)`

3.360.9 Mupad [B] (verification not implemented)

Time = 1.88 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.29

$$\begin{aligned}
& \int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3(A+B\cos(c+dx)) dx \\
&= \frac{2\left(Aa^3 E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + Aa^2 b F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + Aa^2 b \sqrt{\cos(c+dx)} \sin(c+dx)\right)}{Ba^3 \left(\frac{2\sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{{}_2F_1\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3}\right)} \\
&+ \frac{d}{2Ab^3 \cos(c+dx)^{9/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)} \\
&- \frac{9d \sqrt{\sin(c+dx)^2}}{2Bb^3 \cos(c+dx)^{11/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{11}{4}; \frac{15}{4}; \cos(c+dx)^2\right)} \\
&- \frac{11d \sqrt{\sin(c+dx)^2}}{6Aab^2 \cos(c+dx)^{7/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)} \\
&- \frac{7d \sqrt{\sin(c+dx)^2}}{6Ba^2 b \cos(c+dx)^{7/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c+dx)^2\right)} \\
&- \frac{7d \sqrt{\sin(c+dx)^2}}{2Bab^2 \cos(c+dx)^{9/2} \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c+dx)^2\right)} \\
&- \frac{3d \sqrt{\sin(c+dx)^2}}{3d \sqrt{\sin(c+dx)^2}}
\end{aligned}$$

```
input int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3,x)
```

```
output (2*(A*a^3*ellipticE(c/2 + (d*x)/2, 2) + A*a^2*b*ellipticF(c/2 + (d*x)/2, 2)
) + A*a^2*b*cos(c + d*x)^(1/2)*sin(c + d*x))/d + (B*a^3*((2*cos(c + d*x)^(
1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d - (2*A*b^3*c
os(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2)
)/(9*d*(sin(c + d*x)^2)^(1/2)) - (2*B*b^3*cos(c + d*x)^(11/2)*sin(c + d*x)
*hypergeom([1/2, 11/4], 15/4, cos(c + d*x)^2))/(11*d*(sin(c + d*x)^2)^(1/2)
) - (6*A*a*b^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4
, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (6*B*a^2*b*cos(c + d*x)^(
7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(
c + d*x)^2)^(1/2)) - (2*B*a*b^2*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom(
[1/2, 9/4], 13/4, cos(c + d*x)^2))/(3*d*(sin(c + d*x)^2)^(1/2))
```

3.361
$$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.361.1 Optimal result 3369
 3.361.2 Mathematica [A] (verified) 3370
 3.361.3 Rubi [A] (verified) 3370
 3.361.4 Maple [B] (verified) 3374
 3.361.5 Fricas [C] (verification not implemented) 3375
 3.361.6 Sympy [F(-1)] 3376
 3.361.7 Maxima [F] 3376
 3.361.8 Giac [F] 3377
 3.361.9 Mupad [B] (verification not implemented) 3377

3.361.1 Optimal result

Integrand size = 33, antiderivative size = 205

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^3(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2(15a^2Ab + 3Ab^3 + 5a^3B + 9ab^2B) E(\frac{1}{2}(c + dx) | 2)}{5d} \\ &+ \frac{2(21a^3A + 21aAb^2 + 21a^2bB + 5b^3B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{21d} \\ &+ \frac{2b(21aAb + 18a^2B + 5b^2B) \sqrt{\cos(c + dx)} \sin(c + dx)}{21d} \\ &+ \frac{2b^2(7Ab + 11aB) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{35d} \\ &+ \frac{2bB \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2 \sin(c + dx)}{7d} \end{aligned}$$

```
output 2/5*(15*A*a^2*b+3*A*b^3+5*B*a^3+9*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/21*(21*A*a^3+21*A*a*b^2+21*B*a^2*b+5*B*b^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/35*b^2*(7*A*b+11*B*a)*cos(d*x+c)^(3/2)*sin(d*x+c)/d+2/21*b*(21*A*a*b+18*B*a^2+5*B*b^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/d+2/7*b*B*(a+b*cos(d*x+c))^2*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```


3.361.2 Mathematica [A] (verified)

Time = 2.38 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.77

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{42(15a^2Ab + 3Ab^3 + 5a^3B + 9ab^2B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(21a^3A + 21aAb^2 + 21a^2bB + 5b^3B) \text{EllipticF}}$$

input `Integrate[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(42*(15*a^2*A*b + 3*A*b^3 + 5*a^3*B + 9*a*b^2*B)*EllipticE[(c + d*x)/2, 2] + 10*(21*a^3*A + 21*a*A*b^2 + 21*a^2*b*B + 5*b^3*B)*EllipticF[(c + d*x)/2, 2] + b*Sqrt[Cos[c + d*x]]*(42*b*(A*b + 3*a*B)*Cos[c + d*x] + 5*(42*a*A*b + 42*a^2*B + 13*b^2*B + 3*b^2*B*Cos[2*(c + d*x)]))*Sin[c + d*x])/(105*d)`

3.361.3 Rubi [A] (verified)Time = 1.16 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3469, 27, 3042, 3512, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3469}$$

$$\frac{2}{7} \int \frac{(a + b \cos(c + dx)) (b(7Ab + 11aB) \cos^2(c + dx) + (5Bb^2 + 7a(2Ab + aB)) \cos(c + dx) + a(7aA + bB))}{2\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d}$$

$$\downarrow \text{27}$$

3.361. $\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$

$$\frac{1}{7} \int \frac{(a + b \cos(c + dx)) (b(7Ab + 11aB) \cos^2(c + dx) + (5Bb^2 + 7a(2Ab + aB)) \cos(c + dx) + a(7aA + bB))}{\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d}$$

↓ 3042

$$\frac{1}{7} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) (b(7Ab + 11aB) \sin^2(c + dx + \frac{\pi}{2}) + (5Bb^2 + 7a(2Ab + aB)) \sin(c + dx + \frac{\pi}{2}) + a(7aA + bB))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d}$$

↓ 3512

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{5(7aA + bB)a^2 + 5b(18Ba^2 + 21Aba + 5b^2B) \cos^2(c + dx) + 7(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5(7aA + bB)a^2 + 5b(18Ba^2 + 21Aba + 5b^2B) \cos^2(c + dx) + 7(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5(7aA + bB)a^2 + 5b(18Ba^2 + 21Aba + 5b^2B) \sin^2(c + dx + \frac{\pi}{2}) + 7(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \right)$$

↓ 3502

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{2}{3} \int \frac{5(21Aa^3 + 21bBa^2 + 21Ab^2a + 5b^3B) + 21(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{10b(10a^2 + 3b^2) \sin(c + dx)}{7d} \right) + \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \right)$$

↓ 27

3.361. $\int \frac{(a+b \cos(c+dx))^3 (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{5(21Aa^3 + 21bBa^2 + 21Ab^2a + 5b^3B) + 21(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{10b(18a^2B + 21aAb + 5b^2B)}{3d} \right) \right) \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{5(21Aa^3 + 21bBa^2 + 21Ab^2a + 5b^3B) + 21(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{42(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3)}{d} \right) \right) \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \downarrow \text{3227}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(5(21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 21(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \right) \right) \right) \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \downarrow \text{3042}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(5(21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \right) \right) \right) \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \downarrow \text{3119}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(5(21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{42(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3)}{d} \right) \right) \right) \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d} \downarrow \text{3120}$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{10b(18a^2B + 21aAb + 5b^2B) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d} + \frac{1}{3} \left(\frac{10(21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \text{Ellip}}{d} \right) \right) \right) \frac{2bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^2}{7d}$$

input `Int[((a + b*cos[c + d*x])^3*(A + B*cos[c + d*x]))/sqrt[Cos[c + d*x]],x]`

output `(2*b*B*sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])^2*sin[c + d*x])/(7*d) + ((2*b^2*(7*A*b + 11*a*B)*cos[c + d*x]^(3/2)*sin[c + d*x])/(5*d) + (((42*(15*a^2*A*b + 3*A*b^3 + 5*a^3*B + 9*a*b^2*B)*EllipticE[(c + d*x)/2, 2])/d + (10*(21*a^3*A + 21*a*A*b^2 + 21*a^2*b*B + 5*b^3*B)*EllipticF[(c + d*x)/2, 2])/d)/3 + (10*b*(21*a*A*b + 18*a^2*B + 5*b^2*B)*sqrt[Cos[c + d*x]]*sin[c + d*x])/(3*d))/5)/7`

3.361.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c Int[(b*sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]*(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2
, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

3.361.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 663 vs. $2(241) = 482$.

Time = 13.34 (sec) , antiderivative size = 664, normalized size of antiderivative = 3.24

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(240B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3 + (-168Ab^3 - 504Ba^2b - 360Bb^3)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{\sqrt{\cos(c+dx)}}$
parts	Expression too large to display

$$3.361. \quad \int \frac{(a+b\cos(c+dx))^3(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*cos(
1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^3+(-168*A*b^3-504*B*a*b^2-360*B*b^3)
*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(420*A*a*b^2+168*A*b^3+420*B*a^2*
b+504*B*a*b^2+280*B*b^3)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-210*A*a
*b^2-42*A*b^3-210*B*a^2*b-126*B*a*b^2-80*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1
/2*d*x+1/2*c)+105*A*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+105*A*a*b^2*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2))-315*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b-63*A*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),
2^(1/2))*b^3+105*B*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+25*B*b^3*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*
c),2^(1/2))-105*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^3-189*B*(sin(1/2*d*x+1/2*c)^
2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(
1/2))*a*b^2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*
d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.361.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.39

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2 (15 B b^3 \cos(dx + c)^2 + 105 B a^2 b + 105 A a b^2 + 25 B b^3 + 21 (3 B a b^2 + A b^3) \cos(dx + c)) \sqrt{\cos(dx + c)}}{d}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorith
m="fricas")
```

output `1/105*(2*(15*B*b^3*cos(d*x + c)^2 + 105*B*a^2*b + 105*A*a*b^2 + 25*B*b^3 + 21*(3*B*a*b^2 + A*b^3)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 5*sqrt(2)*(21*I*A*a^3 + 21*I*B*a^2*b + 21*I*A*a*b^2 + 5*I*B*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*sqrt(2)*(-21*I*A*a^3 - 21*I*B*a^2*b - 21*I*A*a*b^2 - 5*I*B*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*sqrt(2)*(-5*I*B*a^3 - 15*I*A*a^2*b - 9*I*B*a*b^2 - 3*I*A*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*sqrt(2)*(5*I*B*a^3 + 15*I*A*a^2*b + 9*I*B*a*b^2 + 3*I*A*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.361.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output Timed out

3.361.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\sqrt{\cos(dx + c)}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/sqrt(cos(d*x + c)), x)`

3.361.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/sqrt(cos(d*x + c)), x)`

3.361.9 Mupad [B] (verification not implemented)

Time = 1.59 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.34

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{2 \left(B a^3 E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + B a^2 b F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) + B a^2 b \sqrt{\cos(c + dx)} \sin(c + dx) \right)}{d}$$

$$+ \frac{2 A a^3 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{6 A a^2 b E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d}$$

$$+ \frac{3 A a b^2 \left(\frac{2 \sqrt{\cos(c + dx)} \sin(c + dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3} \right)}{d}$$

$$- \frac{2 A b^3 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{2 B b^3 \cos(c + dx)^{9/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(c + dx)^2\right)}{9 d \sqrt{\sin(c + dx)^2}}$$

$$- \frac{6 B a b^2 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^(1/2),x)`

output $(2*(B*a^3*ellipticE(c/2 + (d*x)/2, 2) + B*a^2*b*ellipticF(c/2 + (d*x)/2, 2) + B*a^2*b*cos(c + d*x)^(1/2)*sin(c + d*x))/d + (2*A*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (6*A*a^2*b*ellipticE(c/2 + (d*x)/2, 2))/d + (3*A*a*b^2*(2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3)/d - (2*A*b^3*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2)) - (2*B*b^3*cos(c + d*x)^(9/2)*sin(c + d*x)*hypergeom([1/2, 9/4], 13/4, cos(c + d*x)^2))/(9*d*(sin(c + d*x)^2)^(1/2)) - (6*B*a*b^2*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))$

3.362
$$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.362.1 Optimal result 3379
 3.362.2 Mathematica [A] (verified) 3380
 3.362.3 Rubi [A] (verified) 3380
 3.362.4 Maple [B] (verified) 3385
 3.362.5 Fracas [C] (verification not implemented) 3386
 3.362.6 Sympy [F(-1)] 3386
 3.362.7 Maxima [F] 3387
 3.362.8 Giac [F] 3387
 3.362.9 Mupad [B] (verification not implemented) 3388

3.362.1 Optimal result

Integrand size = 33, antiderivative size = 202

$$\int \frac{(a + b \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2(5a^3A - 15aAb^2 - 15a^2bB - 3b^3B) E(\frac{1}{2}(c + dx) | 2)}{5d}$$

$$+ \frac{2(9a^2Ab + Ab^3 + 3a^3B + 3ab^2B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$- \frac{2b(6a^2A - Ab^2 - 3abB) \sqrt{\cos(c + dx)} \sin(c + dx)}{3d}$$

$$- \frac{2b^2(5aA - bB) \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{5d} + \frac{2aA(a + b \cos(c + dx))^2 \sin(c + dx)}{d\sqrt{\cos(c + dx)}}$$

output

```
-2/5*(5*A*a^3-15*A*a*b^2-15*B*a^2*b-3*B*b^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(9*A*a^2*b+
A*b^3+3*B*a^3+3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d-2/5*b^2*(5*A*a-B*b)*cos(d*x+c)^(3/2
)*sin(d*x+c)/d+2*a*A*(a+b*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+c)^(1/2)-2/3*
b*(6*A*a^2-A*b^2-3*B*a*b)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.362.2 Mathematica [A] (verified)

Time = 2.13 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.74

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(-30a^3A + 90aAb^2 + 90a^2bB + 18b^3B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(9a^2Ab + Ab^3 + 3a^3B + 3ab^2B) \text{EllipticF}\left(\frac{1}{2}(c + dx) \mid 2\right) + ((10b^2(Ab + 3aB)) \cos(c + dx) + 3(10a^3A + b^3B + b^3B \cos(2(c + dx)))) \sin(c + dx)}{15d}$$

input `Integrate[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `((-30*a^3*A + 90*a*A*b^2 + 90*a^2*b*B + 18*b^3*B)*EllipticE[(c + d*x)/2, 2] + 10*(9*a^2*A*b + A*b^3 + 3*a^3*B + 3*a*b^2*B)*EllipticF[(c + d*x)/2, 2] + ((10*b^2*(A*b + 3*a*B))*Cos[c + d*x] + 3*(10*a^3*A + b^3*B + b^3*B*Cos[2*(c + d*x)]))*Sin[c + d*x])/Sqrt[Cos[c + d*x]]/(15*d)`

3.362.3 Rubi [A] (verified)Time = 1.17 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3468, 27, 3042, 3512, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3468}$$

$$2 \int \frac{(a + b \cos(c + dx)) (-b(5aA - bB) \cos^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \cos(c + dx) + a(5Ab + aB))}{2\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}}$$

↓ 27

$$\int \frac{(a + b \cos(c + dx)) (-b(5aA - bB) \cos^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \cos(c + dx) + a(5Ab + aB))}{\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}}$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) (-b(5aA - bB) \sin(c + dx + \frac{\pi}{2})^2 + (-Aa^2 + 2bBa + Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(5Ab + aB))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}}$$

↓ 3512

$$\frac{2}{5} \int \frac{5(5Ab + aB)a^2 - 5b(6Aa^2 - 3bBa - Ab^2) \cos^2(c + dx) - (5Aa^3 - 15bBa^2 - 15Ab^2a - 3b^3B) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}}$$

↓ 27

$$\frac{1}{5} \int \frac{5(5Ab + aB)a^2 - 5b(6Aa^2 - 3bBa - Ab^2) \cos^2(c + dx) - (5Aa^3 - 15bBa^2 - 15Ab^2a - 3b^3B) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}}$$

↓ 3042

$$\frac{1}{5} \int \frac{5(5Ab + aB)a^2 - 5b(6Aa^2 - 3bBa - Ab^2) \sin(c + dx + \frac{\pi}{2})^2 + (-5Aa^3 + 15bBa^2 + 15Ab^2a + 3b^3B) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}}$$

↓ 3502

3.362. $\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{5(3Ba^3 + 9Aba^2 + 3b^2Ba + Ab^3) - 3(5Aa^3 - 15bBa^2 - 15Ab^2a - 3b^3B) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx - \frac{10b(6a^2A - 3b^3B)}{5d} \right. \\ \left. \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \\ \downarrow 27$$

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{5(3Ba^3 + 9Aba^2 + 3b^2Ba + Ab^3) - 3(5Aa^3 - 15bBa^2 - 15Ab^2a - 3b^3B) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx - \frac{10b(6a^2A - 3b^3B)}{5d} \right. \\ \left. \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \\ \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{5(3Ba^3 + 9Aba^2 + 3b^2Ba + Ab^3) - 3(5Aa^3 - 15bBa^2 - 15Ab^2a - 3b^3B) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{10b(6a^2A - 3b^3B)}{5d} \right. \\ \left. \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \\ \downarrow 3227$$

$$\frac{1}{5} \left(\frac{1}{3} \left(5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - 3(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) \int \sqrt{\cos(c + dx)} dx \right. \right. \\ \left. \left. \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \right) \\ \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{3} \left(5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - 3(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right. \right. \\ \left. \left. \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \right) \\ \downarrow 3119$$

$$\frac{1}{5} \left(\frac{1}{3} \left(5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{6(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) E(\frac{1}{2}(c + dx))}{d} \right. \right. \\ \left. \left. + \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{10(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d} - \frac{6(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) E(\frac{1}{2}(c + dx))}{d} \right. \right. \\ \left. \left. + \frac{2b^2(5aA - bB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{5d} + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{d\sqrt{\cos(c + dx)}} \right) \right)$$

input `Int[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(-2*b^2*(5*a*A - b*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x]/(5*d) + (2*a*A*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(d*sqrt[Cos[c + d*x]]) + (((-6*(5*a^3*A - 15*a*A*b^2 - 15*a^2*b*B - 3*b^3*B)*EllipticE[(c + d*x)/2, 2])/d + (10*(9*a^2*A*b + A*b^3 + 3*a^3*B + 3*a*b^2*B)*EllipticF[(c + d*x)/2, 2])/d)/3 - (10*b*(6*a^2*A - A*b^2 - 3*a*b*B)*sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))/5`

3.362.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3512 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Sin[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) + A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]`

3.362.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 640 vs. $2(240) = 480$.

Time = 11.41 (sec) , antiderivative size = 641, normalized size of antiderivative = 3.17

method	result
default	$\frac{2\left(-24B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3 + 20A \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3 + 60B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)ab^2 + 24B \cos\left(\frac{dx}{2}\right)}{\dots}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/15*(-24*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^3+20*A*cos(1/2*d*x+
1/2*c)*sin(1/2*d*x+1/2*c)^4*b^3+60*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)
^4*a*b^2+24*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*b^3-30*A*cos(1/2*d*x
+1/2*c)*sin(1/2*d*x+1/2*c)^2*a^3-10*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c
)^2*b^3+45*A*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)
^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+5*A*b^3*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1
/2))+15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*El
lipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^3-45*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b
^2-30*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a*b^2-6*B*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^2*b^3+15*B*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1
/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+15*B*a*b^2*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))-45*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x
+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b-9*B*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d
*x+1/2*c),2^(1/2))*b^3/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2
)/d
```


3.362.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.50

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{5\sqrt{2}(3iBa^3 + 9iAa^2b + 3iBab^2 + iAb^3) \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\dots}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="fricas")`

output `-1/15*(5*sqrt(2)*(3*I*B*a^3 + 9*I*A*a^2*b + 3*I*B*a*b^2 + I*A*b^3)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-3*I*B*a^3 - 9*I*A*a^2*b - 3*I*B*a*b^2 - I*A*b^3)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(5*I*A*a^3 - 15*I*B*a^2*b - 15*I*A*a*b^2 - 3*I*B*b^3)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-5*I*A*a^3 + 15*I*B*a^2*b + 15*I*A*a*b^2 + 3*I*B*b^3)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*B*b^3*cos(d*x + c)^2 + 15*A*a^3 + 5*(3*B*a*b^2 + A*b^3)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c))`

3.362.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Timed out`

3.362.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/cos(d*x + c)^(3/2), x)`

3.362.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/cos(d*x + c)^(3/2), x)`

3.362.9 Mupad [B] (verification not implemented)

Time = 1.64 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.23

$$\begin{aligned}
& \int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\
&= \frac{A b^3 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3} \right)}{d} + \frac{2 B a^3 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} \\
&+ \frac{6 A a b^2 E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{6 A a^2 b F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{6 B a^2 b E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} \\
&+ \frac{3 B a b^2 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3} \right)}{d} \\
&+ \frac{2 A a^3 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}} \\
&- \frac{2 B b^3 \cos(c + dx)^{7/2} \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(c + dx)^2\right)}{7 d \sqrt{\sin(c + dx)^2}}
\end{aligned}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^(3/2),x)`

output

```

(A*b^3*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (2*B*a^3*ellipticF(c/2 + (d*x)/2, 2))/d + (6*A*a*b^2*ellipticE(c/2 + (d*x)/2, 2))/d + (6*A*a^2*b*ellipticF(c/2 + (d*x)/2, 2))/d + (6*B*a^2*b*ellipticE(c/2 + (d*x)/2, 2))/d + (3*B*a*b^2*((2*cos(c + d*x)^(1/2)*sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (2*A*a^3*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2)) - (2*B*b^3*cos(c + d*x)^(7/2)*sin(c + d*x)*hypergeom([1/2, 7/4], 11/4, cos(c + d*x)^2))/(7*d*(sin(c + d*x)^2)^(1/2))

```

3.363
$$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.363.1 Optimal result 3389
 3.363.2 Mathematica [A] (verified) 3390
 3.363.3 Rubi [A] (verified) 3390
 3.363.4 Maple [C] (verified) 3394
 3.363.5 Fracas [C] (verification not implemented) 3395
 3.363.6 Sympy [F(-1)] 3396
 3.363.7 Maxima [F] 3396
 3.363.8 Giac [F] 3397
 3.363.9 Mupad [B] (verification not implemented) 3397

3.363.1 Optimal result

Integrand size = 33, antiderivative size = 192

$$\int \frac{(a + b \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{2(3a^2Ab - Ab^3 + a^3B - 3ab^2B) E(\frac{1}{2}(c + dx) | 2)}{d}$$

$$+ \frac{2(a^3A + 9aAb^2 + 9a^2bB + b^3B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2a^2(7Ab + 3aB) \sin(c + dx)}{3d\sqrt{\cos(c + dx)}} - \frac{2b^2(aA - bB)\sqrt{\cos(c + dx)} \sin(c + dx)}{3d}$$

$$+ \frac{2aA(a + b \cos(c + dx))^2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

output

```
-2*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(A*a^3+9*A*a*b^2+9*B*a^2*b+B*b^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*a*A*(a+b*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2/3*a^2*(7*A*b+3*B*a)*sin(d*x+c)/d/cos(d*x+c)^(1/2)-2/3*b^2*(A*a-B*b)*sin(d*x+c)*cos(d*x+c)^(1/2)/d
```

3.363.2 Mathematica [A] (verified)

Time = 2.13 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.86

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{-6(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 2(a^3A + 9aAb^2 + 9a^2bB + b^3B) \sqrt{\cos(c + dx)}}{3d}$$

input `Integrate[((a + bCos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2), x]`

output `(-6*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 2*(a^3*A + 9*a*A*b^2 + 9*a^2*b*B + b^3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 18*a^2*A*b*Sin[c + d*x] + 6*a^3*B*Sin[c + d*x] + b^3*B*Sin[2*(c + d*x)] + 2*a^3*A*Tan[c + d*x])/(3*d*Sqrt[Cos[c + d*x]])`

3.363.3 Rubi [A] (verified)Time = 1.14 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.99, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3468, 27, 3042, 3510, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

$$\downarrow \text{3468}$$

$$\frac{2}{3} \int \frac{(a + b \cos(c + dx)) (-3b(aA - bB) \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(7Ab + 3aB))}{2 \cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)}$$

3.363. $\int \frac{(a+b \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

↓ 27

$$\frac{1}{3} \int \frac{(a + b \cos(c + dx)) (-3b(aA - bB) \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(7Ab + 3aB))}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) (-3b(aA - bB) \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(7Ab + 3aB))}{\sin^{\frac{3}{2}}(c + dx + \frac{\pi}{2})} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3510

$$\frac{1}{3} \left(\frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - 2 \int \frac{-3b^2(aA - bB) \cos^2(c + dx) - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cos(c + dx) + a(Aa^2 + 9bBa + 10Ab^2)}{2\sqrt{\cos(c + dx)}} dx \right) + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \left(\int \frac{-3b^2(aA - bB) \cos^2(c + dx) - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cos(c + dx) + a(Aa^2 + 9bBa + 10Ab^2)}{\sqrt{\cos(c + dx)}} dx \right) + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{-3b^2(aA - bB) \sin^2(c + dx + \frac{\pi}{2}) - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(Aa^2 + 9bBa + 10Ab^2)}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3502

$$\frac{1}{3} \left(\frac{2}{3} \int \frac{3(Aa^3 + 9bBa^2 + 9Ab^2a + b^3B - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cos(c + dx))}{2\sqrt{\cos(c + dx)}} dx + \frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)} \right) \\ \downarrow 27$$

$$\frac{1}{3} \left(\int \frac{Aa^3 + 9bBa^2 + 9Ab^2a + b^3B - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)} \right) \\ \downarrow 3042$$

$$\frac{1}{3} \left(\int \frac{Aa^3 + 9bBa^2 + 9Ab^2a + b^3B - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)} \right) \\ \downarrow 3227$$

$$\frac{1}{3} \left((a^3A + 9a^2bB + 9aAb^2 + b^3B) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - 3(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \sqrt{\cos(c + dx)} dx + \frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)} \right) \\ \downarrow 3042$$

$$\frac{1}{3} \left((a^3A + 9a^2bB + 9aAb^2 + b^3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - 3(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)} \right) \\ \downarrow 3119$$

$$\frac{1}{3} \left((a^3A + 9a^2bB + 9aAb^2 + b^3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \frac{6(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{d} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

3.363. $\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

↓ 3120

$$\frac{1}{3} \left(\frac{2a^2(3aB + 7Ab) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} + \frac{2(a^3A + 9a^2bB + 9aAb^2 + b^3B) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{6(a^3B + 3a^2Ab - 2aA \sin(c + dx)(a + b \cos(c + dx))^2)}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

input `Int[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*a*A*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((-6*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*EllipticE[(c + d*x)/2, 2])/d + (2*(a^3*A + 9*a*A*b^2 + 9*a^2*b*B + b^3*B)*EllipticF[(c + d*x)/2, 2])/d + (2*a^2*(7*A*b + 3*a*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - (2*b^2*(a*A - b*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/d)/3`

3.363.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(b*c - a*d))*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.363.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 12.89 (sec) , antiderivative size = 771, normalized size of antiderivative = 4.02

method	result	size
parts	Expression too large to display	771
default	Expression too large to display	1210

$$3.363. \int \frac{(a+b \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^2(c+dx)} dx$$

input `int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)`

output `2*(A*b^3+3*B*a*b^2)*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)
)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE
(cos(1/2*d*x+1/2*c),2^(1/2))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d+2*(3*A*a*b^2
+3*B*a^2*b)/d*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2))-2*(3*A*a^2*b+B*a^3)*(-
2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2
-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(c
os(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d-2/3*A*a^3*(-2
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(c
os(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*x+1/2*c)^2*cos
(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*((2*cos(1/2*d*x+1/2*c)^2-1)*sin
(1/2*d*x+1/2*c)^2)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1
/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3/2)/sin(1/2*d*x+1/2*c)/d-2/3*B*b^3*((2*co
s(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*sin(1/2*d*x+1/2*c)^4*
cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2
*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(...`

3.363.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.59

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{2}(-i A a^3 - 9i B a^2 b - 9i A a b^2 - i B b^3) \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\cos^{\frac{5}{2}}(c + dx)}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorith
m="fricas")`

output $\frac{1}{3}(\sqrt{2})(-I A a^3 - 9 I B a^2 b - 9 I A a b^2 - I B b^3) \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I \sin(dx + c)) + \sqrt{2}(I A a^3 + 9 I B a^2 b + 9 I A a b^2 + I B b^3) \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I \sin(dx + c)) - 3 \sqrt{2}(I B a^3 + 3 I A a^2 b - 3 I B a b^2 - I A b^3) \cos(dx + c)^2 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I \sin(dx + c))) - 3 \sqrt{2}(-I B a^3 - 3 I A a^2 b + 3 I B a b^2 + I A b^3) \cos(dx + c)^2 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I \sin(dx + c))) + 2(B b^3 \cos(dx + c)^2 + A a^3 + 3(B a^3 + 3 A a^2 b) \cos(dx + c)) \sqrt{\cos(dx + c) \sin(dx + c)} / (d \cos(dx + c)^2)$

3.363.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output Timed out

3.363.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/cos(d*x + c)^(5/2), x)`

3.363.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/cos(d*x + c)^(5/2), x)`

3.363.9 Mupad [B] (verification not implemented)

Time = 2.65 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.33

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 \left(A E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) b^3 + 3 A a F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right) b^2 \right)}{d}$$

$$+ \frac{B b^3 \left(\frac{2 \sqrt{\cos(c+dx)} \sin(c+dx)}{3} + \frac{2 F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{3} \right)}{d} + \frac{6 B a b^2 E\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d}$$

$$+ \frac{6 B a^2 b F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d} + \frac{2 A a^3 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{2 B a^3 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

$$+ \frac{6 A a^2 b \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^(5/2),x)`

output $(2*(A*b^3*ellipticE(c/2 + (d*x)/2, 2) + 3*A*a*b^2*ellipticF(c/2 + (d*x)/2, 2)))/d + (B*b^3*((2*\cos(c + d*x)^(1/2)*\sin(c + d*x))/3 + (2*ellipticF(c/2 + (d*x)/2, 2))/3))/d + (6*B*a*b^2*ellipticE(c/2 + (d*x)/2, 2))/d + (6*B*a^2*b*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a^3*\sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, \cos(c + d*x)^2))/(3*d*\cos(c + d*x)^(3/2)*(\sin(c + d*x)^2)^(1/2)) + (2*B*a^3*\sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, \cos(c + d*x)^2))/(d*\cos(c + d*x)^(1/2)*(\sin(c + d*x)^2)^(1/2)) + (6*A*a^2*b*\sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, \cos(c + d*x)^2))/(d*\cos(c + d*x)^(1/2)*(\sin(c + d*x)^2)^(1/2))$

3.364
$$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.364.1 Optimal result 3399
 3.364.2 Mathematica [A] (verified) 3400
 3.364.3 Rubi [A] (verified) 3400
 3.364.4 Maple [C] (verified) 3405
 3.364.5 Fracas [C] (verification not implemented) 3406
 3.364.6 Sympy [F(-1)] 3406
 3.364.7 Maxima [F] 3407
 3.364.8 Giac [F] 3407
 3.364.9 Mupad [B] (verification not implemented) 3408

3.364.1 Optimal result

Integrand size = 33, antiderivative size = 204

$$\int \frac{(a + b \cos(c + dx))^3(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= -\frac{2(3a^3A + 15aAb^2 + 15a^2bB - 5b^3B) E(\frac{1}{2}(c + dx) | 2)}{5d}$$

$$+ \frac{2(3a^2Ab + 3Ab^3 + a^3B + 9ab^2B) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d}$$

$$+ \frac{2a^2(9Ab + 5aB) \sin(c + dx)}{15d \cos^{\frac{3}{2}}(c + dx)} + \frac{2a(3a^2A + 14Ab^2 + 15abB) \sin(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

$$+ \frac{2aA(a + b \cos(c + dx))^2 \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx)}$$

output

```
-2/5*(3*A*a^3+15*A*a*b^2+15*B*a^2*b-5*B*b^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*(3*A*a^2*b+
3*A*b^3+B*a^3+9*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/15*a^2*(9*A*b+5*B*a)*sin(d*x+c)/d
/cos(d*x+c)^(3/2)+2/5*a*A*(a+b*cos(d*x+c))^2*sin(d*x+c)/d/cos(d*x+c)^(5/2)
+2/5*a*(3*A*a^2+14*A*b^2+15*B*a*b)*sin(d*x+c)/d/cos(d*x+c)^(1/2)
```

3.364.2 Mathematica [A] (verified)

Time = 3.02 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.86

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{-6(3a^3A + 15aAb^2 + 15a^2bB - 5b^3B) \cos^{\frac{3}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(3a^2Ab + 3Ab^3 + a^3B + 9ab^2B)}{15d \cos^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(-6*(3*a^3*A + 15*a*A*b^2 + 15*a^2*b*B - 5*b^3*B)*Cos[c + d*x]^(3/2)*EllipticE[(c + d*x)/2, 2] + 10*(3*a^2*A*b + 3*A*b^3 + a^3*B + 9*a*b^2*B)*Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + 10*a^2*(3*A*b + a*B)*Sin[c + d*x] + 9*a*(a^2*A + 5*A*b^2 + 5*a*b*B)*Sin[2*(c + d*x)] + 6*a^3*A*Tan[c + d*x])/(15*d*Cos[c + d*x]^(3/2))`

3.364.3 Rubi [A] (verified)Time = 1.13 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3468, 27, 3042, 3510, 27, 3042, 3500, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

$$\downarrow \text{3468}$$

$$\frac{2}{5} \int \frac{(a + b \cos(c + dx)) (-b(aA - 5bB) \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(9Ab + 5aB))}{2 \cos^{\frac{5}{2}}(c + dx) + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}} dx$$

3.364. $\int \frac{(a+b \cos(c+dx))^3 (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

↓ 27

$$\frac{1}{5} \int \frac{(a + b \cos(c + dx)) (-b(aA - 5bB) \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(9Ab + 5aB))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2})) (-b(aA - 5bB) \sin^2(c + dx + \frac{\pi}{2}) + (3Aa^2 + 10bBa + 5Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(9Ab + 5aB))}{\sin^{\frac{5}{2}}(c + dx + \frac{\pi}{2})} dx$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3510

$$\frac{1}{5} \left(\frac{2a^2(5aB + 9Ab) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} - \frac{2}{3} \int \frac{-3b^2(aA - 5bB) \cos^2(c + dx) + 5(Ba^3 + 3Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)} dx \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{-3b^2(aA - 5bB) \cos^2(c + dx) + 5(Ba^3 + 3Aba^2 + 9b^2Ba + 3Ab^3) \cos(c + dx) + 3a(3Aa^2 + 15bBa + 14Ab^2)}{\cos^{\frac{3}{2}}(c + dx)} dx \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{-3b^2(aA - 5bB) \sin^2(c + dx + \frac{\pi}{2}) + 5(Ba^3 + 3Aba^2 + 9b^2Ba + 3Ab^3) \sin(c + dx + \frac{\pi}{2}) + 3a(3Aa^2 + 15bBa + 14Ab^2)}{\sin^{\frac{3}{2}}(c + dx + \frac{\pi}{2})} dx \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 3500

$$\frac{1}{5} \left(\frac{1}{3} \left(2 \int \frac{5(Ba^3 + 3Aba^2 + 9b^2Ba + 3Ab^3) - 3(3Aa^3 + 15bBa^2 + 15Ab^2a - 5b^3B) \cos(c + dx)}{2\sqrt{\cos(c + dx)}} dx + \frac{6a(3a^2A + 15bBa + 14Ab^2)}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \left(\int \frac{5(Ba^3 + 3Aba^2 + 9b^2Ba + 3Ab^3) - 3(3Aa^3 + 15bBa^2 + 15Ab^2a - 5b^3B) \cos(c + dx)}{\sqrt{\cos(c + dx)}} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \frac{6a(3a^2A + 15abB + 14Ab^2) \cos(c + dx)}{d\sqrt{\cos(c + dx)}} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\int \frac{5(Ba^3 + 3Aba^2 + 9b^2Ba + 3Ab^3) - 3(3Aa^3 + 15bBa^2 + 15Ab^2a - 5b^3B) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \cos(c + dx + \frac{\pi}{2})}{d\sqrt{\sin(c + dx + \frac{\pi}{2})}} - \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx + \frac{\pi}{2})}{d\sqrt{\sin(c + dx + \frac{\pi}{2})}} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3227

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \int \frac{1}{\sqrt{\cos(c + dx)}} dx - 3(3a^3A + 15a^2bB + 15aAb^2 - 5b^3B) \int \sqrt{\cos(c + dx)} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - 3(3a^3A + 15a^2bB + 15aAb^2 - 5b^3B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3119

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \frac{6a(3a^2A + 15abB + 14Ab^2) \cos(c + dx)}{d\sqrt{\cos(c + dx)}} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{2a^2(5aB + 9Ab) \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} + \frac{1}{3} \left(\frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx)}{d \sqrt{\cos(c + dx)}} + \frac{10(a^3B + 3a^2Ab + 9ab^2B + 3A^2)}{d} \right) \right) + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^2}{5d \cos^{\frac{5}{2}}(c + dx)}$$

input `Int[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*(a + b*Cos[c + d*x])^2*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*a^2*(9*A*b + 5*a*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((-6*(3*a^3*A + 15*a*A*b^2 + 15*a^2*b*B - 5*b^3*B)*EllipticE[(c + d*x)/2, 2])/d + (10*(3*a^2*A*b + 3*A*b^3 + a^3*B + 9*a*b^2*B)*EllipticF[(c + d*x)/2, 2])/d + (6*a*(3*a^2*A + 14*A*b^2 + 15*a*b*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/3)/5`

3.364.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3500 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C + b*(A*b - a*B + b*C)*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && LtQ[m, -1] && NeQ[a^2 - b^2, 0]`

rule 3510 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(b*c - a*d)*(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b^2*f*(m + 1)*(a^2 - b^2))), x] - Simp[1/(b^2*(m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[b*(m + 1)*((b*B - a*C)*(b*c - a*d) - A*b*(a*c - b*d)) + (b*B*(a^2*d + b^2*d*(m + 1) - a*b*c*(m + 2)) + (b*c - a*d)*(A*b^2*(m + 2) + C*(a^2 + b^2*(m + 1)))*Sin[e + f*x] - b*C*d*(m + 1)*(a^2 - b^2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]`

3.364.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 14.86 (sec) , antiderivative size = 950, normalized size of antiderivative = 4.66

method	result	size
parts	Expression too large to display	950
default	Expression too large to display	970

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

```
output 2*(A*b^3+3*B*a*b^2)/d*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2))-2*(3*A*a*b^2+
3*B*a^2*b)*(-2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*
d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d
*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d-2
/3*(3*A*a^2*b+B*a^3)*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c
)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*
sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*si
n(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*((2*cos
(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+
sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3/2)/sin(1/2*d*x+1
/2*c)/d-2/5*A*a^3*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)
)/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1
)/sin(1/2*d*x+1/2*c)^3*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*s
in(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/
2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2
*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/
2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*
x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2...
```

3.364.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.14 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.60

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{5\sqrt{2}(iBa^3 + 3iAa^2b + 9iBab^2 + 3iAb^3) \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\dots}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="fricas")`

output `-1/15*(5*sqrt(2)*(I*B*a^3 + 3*I*A*a^2*b + 9*I*B*a*b^2 + 3*I*A*b^3)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-I*B*a^3 - 3*I*A*a^2*b - 9*I*B*a*b^2 - 3*I*A*b^3)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(3*I*A*a^3 + 15*I*B*a^2*b + 15*I*A*a*b^2 - 5*I*B*b^3)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-3*I*A*a^3 - 15*I*B*a^2*b - 15*I*A*a*b^2 + 5*I*B*b^3)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*A*a^3 + 9*(A*a^3 + 5*B*a^2*b + 5*A*a*b^2)*cos(d*x + c)^2 + 5*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)`

3.364.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.364.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/cos(d*x + c)^(7/2), x)`

3.364.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/cos(d*x + c)^(7/2), x)`

3.364.9 Mupad [B] (verification not implemented)

Time = 4.24 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.43

$$\begin{aligned}
& \int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\
&= \frac{2 \left(B E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) b^3 + 3 B a F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right) b^2 \right)}{d} + \frac{2 A b^3 F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d} \\
&+ \frac{2 A a^3 \sin(c + dx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(c + dx)^2\right)}{5 d \cos(c + dx)^{5/2} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{2 B a^3 \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{3 d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{6 A a b^2 \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{2 A a^2 b \sin(c + dx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(c + dx)^2\right)}{d \cos(c + dx)^{3/2} \sqrt{\sin(c + dx)^2}} \\
&+ \frac{6 B a^2 b \sin(c + dx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(c + dx)^2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sin(c + dx)^2}}
\end{aligned}$$

```
input int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/cos(c + d*x)^(7/2),x)
```

```
output (2*(B*b^3*ellipticE(c/2 + (d*x)/2, 2) + 3*B*a*b^2*ellipticF(c/2 + (d*x)/2,
2)))/d + (2*A*b^3*ellipticF(c/2 + (d*x)/2, 2))/d + (2*A*a^3*sin(c + d*x)*
hypergeom([-5/4, 1/2], -1/4, cos(c + d*x)^2))/(5*d*cos(c + d*x)^(5/2)*(sin
(c + d*x)^2)^(1/2)) + (2*B*a^3*sin(c + d*x)*hypergeom([-3/4, 1/2], 1/4, co
s(c + d*x)^2))/(3*d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2)) + (6*A*a*b^
2*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d*cos(c + d*x)
)^(1/2)*(sin(c + d*x)^2)^(1/2)) + (2*A*a^2*b*sin(c + d*x)*hypergeom([-3/4,
1/2], 1/4, cos(c + d*x)^2))/(d*cos(c + d*x)^(3/2)*(sin(c + d*x)^2)^(1/2))
+ (6*B*a^2*b*sin(c + d*x)*hypergeom([-1/4, 1/2], 3/4, cos(c + d*x)^2))/(d
*cos(c + d*x)^(1/2)*(sin(c + d*x)^2)^(1/2))
```

3.365 $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.365.1 Optimal result 3409
 3.365.2 Mathematica [A] (verified) 3410
 3.365.3 Rubi [A] (verified) 3410
 3.365.4 Maple [B] (verified) 3415
 3.365.5 Fricas [F] 3416
 3.365.6 Sympy [F(-1)] 3416
 3.365.7 Maxima [F] 3416
 3.365.8 Giac [F] 3417
 3.365.9 Mupad [F(-1)] 3417

3.365.1 Optimal result

Integrand size = 33, antiderivative size = 182

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = -\frac{2(5aAb-5a^2B-3b^2B)E(\frac{1}{2}(c+dx)|2)}{5b^3d} + \frac{2(3a^2+b^2)(Ab-aB)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{3b^4d} - \frac{2a^3(Ab-aB)\text{EllipticPi}(\frac{2b}{a+b},\frac{1}{2}(c+dx),2)}{b^4(a+b)d} + \frac{2(Ab-aB)\sqrt{\cos(c+dx)}\sin(c+dx)}{3b^2d} + \frac{2B \cos^{\frac{3}{2}}(c+dx)\sin(c+dx)}{5bd}$$

output

```
-2/5*(5*A*a*b-5*B*a^2-3*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^3/d+2/3*(3*a^2+b^2)*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b^4/d-2*a^3*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/b^4/(a+b)/d+2/5*B*cos(d*x+c)^(3/2)*sin(d*x+c)/b/d+2/3*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/b^2/d
```


3.365.2 Mathematica [A] (verified)

Time = 1.72 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.43

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \frac{2b^2(-5aAb+5a^2B+9b^2B)\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + 2b^2(5Ab+4aB) \left(2\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} \right)$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `((2*b^2*(-5*a*A*b + 5*a^2*B + 9*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + 2*b^2*(5*A*b + 4*a*B)*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)) + 4*b^2*Sqrt[Cos[c + d*x]]*(5*A*b - 5*a*B + 3*b*B*Cos[c + d*x])*Sin[c + d*x] + (6*(-5*a*A*b + 5*a^2*B + 3*b^2*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*Sqrt[Sin[c + d*x]^2]))/(30*b^4*d)`

3.365.3 Rubi [A] (verified)Time = 1.48 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.10, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3538, 27, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}}(A+B\sin(c+dx+\frac{\pi}{2}))}{a+b\sin(c+dx+\frac{\pi}{2})} dx$$

$$\downarrow \text{3469}$$

3.365. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

$$\begin{aligned}
& \frac{2 \int \frac{\sqrt{\cos(c+dx)}(5(Ab-aB)\cos^2(c+dx)+3bB\cos(c+dx)+3aB)}{2(a+b\cos(c+dx))} dx}{5b} + \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\sqrt{\cos(c+dx)}(5(Ab-aB)\cos^2(c+dx)+3bB\cos(c+dx)+3aB)}{a+b\cos(c+dx)} dx}{5b} + \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(5(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2+3bB\sin(c+dx+\frac{\pi}{2})+3aB)}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{5b} + \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 3528 \\
& \frac{2 \int \frac{-3(-5Ba^2+5Aba-3b^2B)\cos^2(c+dx)+b(5Ab+4aB)\cos(c+dx)+5a(Ab-aB)}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{10(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} + \\
& \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{-3(-5Ba^2+5Aba-3b^2B)\cos^2(c+dx)+b(5Ab+4aB)\cos(c+dx)+5a(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{10(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} + \\
& \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{-3(-5Ba^2+5Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})^2+b(5Ab+4aB)\sin(c+dx+\frac{\pi}{2})+5a(Ab-aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{3b} + \frac{10(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} + \\
& \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 3538 \\
& \frac{-\frac{3(-5a^2B+5aAb-3b^2B)}{b} \int \sqrt{\cos(c+dx)} dx - \frac{5(ab(Ab-aB)+(3a^2+b^2)\cos(c+dx)(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{10(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} + \\
& \quad \frac{5b}{5bd} \frac{2B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow 27
\end{aligned}$$

3.365. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

$$\frac{5 \int \frac{ab(Ab-aB) + (3a^2+b^2) \cos(c+dx)(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} - \frac{3(-5a^2B+5aAb-3b^2B) \int \sqrt{\cos(c+dx)} dx}{b} + \frac{10(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} +$$

$$\frac{5b}{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{5bd}{\downarrow} \quad 3042$$

$$\frac{5 \int \frac{ab(Ab-aB) + (3a^2+b^2) \sin(c+dx+\frac{\pi}{2})(Ab-aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{3(-5a^2B+5aAb-3b^2B) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{10(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} +$$

$$\frac{5b}{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{5bd}{\downarrow} \quad 3119$$

$$\frac{5 \int \frac{ab(Ab-aB) + (3a^2+b^2) \sin(c+dx+\frac{\pi}{2})(Ab-aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{6(-5a^2B+5aAb-3b^2B) E(\frac{1}{2}(c+dx)|2)}{bd} + \frac{10(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} +$$

$$\frac{5b}{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{5bd}{\downarrow} \quad 3481$$

$$\frac{5 \left(\frac{(3a^2+b^2)(Ab-aB) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{3a^3(Ab-aB) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} \right)}{3b} - \frac{6(-5a^2B+5aAb-3b^2B) E(\frac{1}{2}(c+dx)|2)}{bd} + \frac{10(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} +$$

$$\frac{5b}{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{5bd}{\downarrow} \quad 3042$$

$$\frac{5 \left(\frac{(3a^2+b^2)(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{3a^3(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} \right)}{3b} - \frac{6(-5a^2B+5aAb-3b^2B) E(\frac{1}{2}(c+dx)|2)}{bd} + \frac{10(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} +$$

$$\frac{5b}{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{5bd}{\downarrow} \quad 3120$$

3.365. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

$$\begin{aligned}
& \frac{\left(\frac{2(3a^2+b^2)(Ab-aB)}{bd} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{3a^3(Ab-aB)}{b} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}(a+b\sin\left(c+dx+\frac{\pi}{2}\right))} dx \right)}{b} \\
& \frac{6(-5a^2B+5aAb-3b^2B)}{bd} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + \frac{10(Ab-aB)}{3b} \\
& \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5bd} \\
& \quad \downarrow \text{3284} \\
& \frac{\left(\frac{2(3a^2+b^2)(Ab-aB)}{bd} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{6a^3(Ab-aB)}{bd(a+b)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right) \right)}{b} \\
& \frac{6(-5a^2B+5aAb-3b^2B)}{bd} E\left(\frac{1}{2}(c+dx) \middle| 2\right) + \frac{10(Ab-aB) \sin(c+dx)}{3b} \\
& \frac{2B \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{5bd}
\end{aligned}$$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(2*B*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(5*b*d) + (((-6*(5*a*A*b - 5*a^2*B - 3*b^2*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + (5*((2*(3*a^2 + b^2)*(A*b - a*B)*EllipticF[(c + d*x)/2, 2])/(b*d) - (6*a^3*(A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d)))/b)/(3*b) + (10*(A*b - a*B)*Sqrt[Cos[c + d*x]*Sin[c + d*x])/(3*b*d))/(5*b)`

3.365.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.365. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m - 1)*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Ssin[e + f*x])^(m - 2)*(c + d*Ssin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3481 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Ssin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Ssin[e + f*x])^m/(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Ssin[e + f*x])^(m - 1)*(c + d*Ssin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)] + (C_.)*sin[(e_.) + (f_.)*(x_.)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_.)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
] && NeQ[c^2 - d^2, 0]
```

3.365.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1073 vs. 2(248) = 496.

Time = 7.51 (sec) , antiderivative size = 1074, normalized size of antiderivative = 5.90

method	result	size
default	Expression too large to display	1074

```
input int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVER
BOSE)
```

```
output -2/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*((-24*B*a*b^
3+24*B*b^4)*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+(20*A*a*b^3-20*A*b^4-2
0*B*a^2*b^2+44*B*a*b^3-24*B*b^4)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(
-10*A*a*b^3+10*A*b^4+10*B*a^2*b^2-16*B*a*b^3+6*B*b^4)*sin(1/2*d*x+1/2*c)^2
*cos(1/2*d*x+1/2*c)+15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a^3*b-15*A*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2))*a^2*b^2+5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*
c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a*b^3-5*A*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2))*b^4+15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b^2-15*A*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2))*a*b^3-15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
)^2-1)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a^3*b-15*B*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))*a^4+15*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2
*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a^3*b-5*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))*a^2*b^2+5*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(...
```

$$3.365. \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.365.5 Fracas [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `integral((B*cos(d*x + c)^3 + A*cos(d*x + c)^2)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a), x)`

3.365.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.365.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.365.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.365.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.366
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.366.1 Optimal result 3418
 3.366.2 Mathematica [A] (verified) 3419
 3.366.3 Rubi [A] (verified) 3419
 3.366.4 Maple [B] (verified) 3423
 3.366.5 Fracas [F(-1)] 3423
 3.366.6 Sympy [F(-1)] 3424
 3.366.7 Maxima [F] 3424
 3.366.8 Giac [F] 3424
 3.366.9 Mupad [F(-1)] 3425

3.366.1 Optimal result

Integrand size = 33, antiderivative size = 137

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2(Ab - aB)E(\frac{1}{2}(c+dx)|2)}{b^2d} - \frac{2(3aAb - 3a^2B - b^2B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{3b^3d} + \frac{2a^2(Ab - aB) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{b^3(a+b)d} + \frac{2B\sqrt{\cos(c+dx)} \sin(c+dx)}{3bd}$$

```
output 2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/b^2/d-2/3*(3*A*a*b-3*B*a^2-B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/b^3/d+2*a^2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))/b^3/(a+b)/d+2/3*B*sin(d*x+c)*cos(d*x+c)^(1/2)/b/d
```

3.366.2 Mathematica [A] (verified)

Time = 1.10 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.51

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \frac{(3Ab-aB)\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + B\left(2\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b}\right) + 2B\sqrt{\cos(c+dx)}$$

```
input Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x
]
```

```
output (((3*A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + B*(2*
EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]
)/(a + b)) + 2*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x] + (3*(A*b - a*B)*(-2*a*b*
EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[S
qrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Co
s[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*Sqrt[Sin[c + d*x]^2]))/(3*b*d)
```

3.366.3 Rubi [A] (verified)Time = 1.03 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3469, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{a+b\sin(c+dx+\frac{\pi}{2})} dx$$

$$\downarrow \text{3469}$$

$$\frac{2\int \frac{3(Ab-aB)\cos^2(c+dx)+bB\cos(c+dx)+aB}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd}$$

3.366. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{3(Ab-aB)\cos^2(c+dx)+bB\cos(c+dx)+aB}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 3042 \\
& \frac{\int \frac{3(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2+bB\sin(c+dx+\frac{\pi}{2})+aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{3b} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 3538 \\
& \frac{3(Ab-aB)\int \frac{\sqrt{\cos(c+dx)} dx}{b} - \int \frac{abB - (-3Ba^2+3Aba-b^2B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 25 \\
& \frac{\int \frac{abB - (-3Ba^2+3Aba-b^2B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{3b} + \frac{3(Ab-aB)\int \frac{\sqrt{\cos(c+dx)} dx}{b}}{3b} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 3042 \\
& \frac{\int \frac{abB + (3Ba^2-3Aba+b^2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{3b} + \frac{3(Ab-aB)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b}}{3b} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 3119 \\
& \frac{\int \frac{abB + (3Ba^2-3Aba+b^2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{3b} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{bd} + \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 3481 \\
& \frac{3a^2(Ab-aB)\int \frac{1}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx - (-3a^2B+3aAb-b^2B)\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{bd} + \\
& \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd} \\
& \downarrow 3042 \\
& \frac{3a^2(Ab-aB)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx - (-3a^2B+3aAb-b^2B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3b} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{bd} + \\
& \frac{2B\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd}
\end{aligned}$$

3.366. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

↓ 3120

$$\frac{3a^2(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{\frac{b}{b}} - \frac{2(-3a^2B+3aAb-b^2B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{bd} +$$

$$\frac{3b}{2B \sin(c+dx) \sqrt{\cos(c+dx)}} \frac{3bd}{3bd}$$

↓ 3284

$$\frac{6a^2(Ab-aB) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx),2)}{bd(a+b)} - \frac{2(-3a^2B+3aAb-b^2B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{bd} +$$

$$\frac{3b}{2B \sin(c+dx) \sqrt{\cos(c+dx)}} \frac{3bd}{3bd}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `((6*(A*b - a*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((-2*(3*a*A*b - 3*a^2*B - b^2*B)*EllipticF[(c + d*x)/2, 2])/(b*d) + (6*a^2*(A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))/b)/(3*b) + (2*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*b*d)`

3.366.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m - 1)*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Ssin[e + f*x])^(m - 2)*(c + d*Ssin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Ssin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Ssin[e + f*x])^m/(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3538 `Int(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Ssin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Ssin[e + f*x])*(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.366.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 821 vs. $2(209) = 418$.

Time = 5.88 (sec) , antiderivative size = 822, normalized size of antiderivative = 6.00

method	result	size
default	Expression too large to display	822

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVER
BOSE)
```

```
output -2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*B*cos(1/2*
d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*a*b^2-4*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1
/2*c)^4*b^3+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a^2*b-3*A*a^2*b*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/
2*d*x+1/2*c),2^(1/2))+3*A*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*
x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*A*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2))*a*b^2+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^3-2*B*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^2*a*b^2+2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*
b^3-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elli
pticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a^3+3*B*a^3*(sin(1/2*d*x+1/2
*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c)
,2^(1/2))-3*B*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*a*b^2*(sin(1/2*d*x+1/2*c)^
2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(
1/2))-B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*
EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*
sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2...
```

3.366.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm=
"fricas")
```

3.366. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

output Timed out

3.366.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output Timed out

3.366.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.366.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.366. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$

3.366.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

$$3.367 \quad \int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.367.1 Optimal result	3426
3.367.2 Mathematica [A] (verified)	3426
3.367.3 Rubi [A] (verified)	3427
3.367.4 Maple [A] (verified)	3429
3.367.5 Fricas [F(-1)]	3430
3.367.6 Sympy [F(-1)]	3430
3.367.7 Maxima [F]	3431
3.367.8 Giac [F]	3431
3.367.9 Mupad [F(-1)]	3431

3.367.1 Optimal result

Integrand size = 33, antiderivative size = 89

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} + \frac{2(Ab-aB) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{b^2d} - \frac{2a(Ab-aB) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{b^2(a+b)d}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/b/d+2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/b^2/d-2*a*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))/b^2/(a+b)/d
```

3.367.2 Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.44

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{Ab \left(2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} \right) - 2B \left(bE\left(\arcsin\left(\sqrt{\cos(c+dx)}\right) \mid -1\right) - (a+b) \operatorname{EllipticF}\left(\arcsin\left(\sqrt{\cos(c+dx)}\right), 2\right) \right)}{b^2d}$$

3.367. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{a+b \cos(c+dx)} dx$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x
]`

output `(A*b*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b) - (2*B*(b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] - (a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + a*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/Sqrt[Sin[c + d*x]^2])/(b^2*d)`

3.367.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3481, 3042, 3119, 3282, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{a+b\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3481} \\
 & \frac{(Ab-aB) \int \frac{\sqrt{\cos(c+dx)}}{a+b\cos(c+dx)} dx}{b} + \frac{B \int \sqrt{\cos(c+dx)} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab-aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} \\
 & \quad \downarrow \text{3119} \\
 & \frac{(Ab-aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{2BE(\frac{1}{2}(c+dx)|2)}{bd} \\
 & \quad \downarrow \text{3282}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{(Ab - aB) \left(\frac{\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx}{b} \right)}{b} + \frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab - aB) \left(\frac{\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b \sin\left(c+dx+\frac{\pi}{2}\right))}} dx}{b} \right)}{b} + \frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} \\
 & \quad \downarrow \text{3120} \\
 & \frac{(Ab - aB) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{a \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b \sin\left(c+dx+\frac{\pi}{2}\right))}} dx}{b} \right)}{b} + \frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} \\
 & \quad \downarrow \text{3284} \\
 & \frac{(Ab - aB) \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{bd(a+b)} \right)}{b} + \frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd}
 \end{aligned}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(2*B*EllipticE[(c + d*x)/2, 2])/(b*d) + ((A*b - a*B)*((2*EllipticF[(c + d*x)/2, 2])/(b*d) - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d)))/b`

3.367.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 3282 Int[Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]/((a_.) + (b_.)*sin[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d/b Int[1/Sqrt[c + d*Sin[e + f*x]], x], x
] + Simp[(b*c - a*d)/b Int[1/((a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2
- b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.367.4 Maple [A] (verified)

Time = 4.99 (sec) , antiderivative size = 295, normalized size of antiderivative = 3.31

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}}\left(AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)ab - AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-2}\right)b^2(a-b)\right)$

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b), x, method=_RETURNVER
BOSE)
```

output $-2*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*(A*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a*b-A*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b^2-A*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})*a*b-B*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a^2+B*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a*b-B*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a*b+B*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b^2+B*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})*a^2)/b^2/(a-b)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$

3.367.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output Timed out

3.367.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output Timed out

3.367.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a), x)`

3.367.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a), x)`

3.367.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.368 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx$

3.368.1 Optimal result 3432
 3.368.2 Mathematica [A] (verified) 3432
 3.368.3 Rubi [A] (verified) 3433
 3.368.4 Maple [A] (verified) 3434
 3.368.5 Fricas [F(-1)] 3435
 3.368.6 Sympy [F(-1)] 3435
 3.368.7 Maxima [F] 3436
 3.368.8 Giac [F] 3436
 3.368.9 Mupad [F(-1)] 3436

3.368.1 Optimal result

Integrand size = 33, antiderivative size = 61

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))}} dx = \frac{2B \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{bd} + \frac{2(Ab - aB) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{b(a + b)d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/b/d+2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))/b/(a+b)/d`

3.368.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.95

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))}} dx = \frac{2((a + b)B \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + (Ab - aB) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right))}{b(a + b)d}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])),x]`

output $(2*((a + b)*B*EllipticF[(c + d*x)/2, 2] + (A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(b*(a + b)*d)$

3.368.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {3042, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx \\
 & \quad \downarrow \text{3481} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} + \frac{B \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} + \frac{B \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \quad \downarrow \text{3120} \\
 & \frac{(Ab - aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} + \frac{2B \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{bd} \\
 & \quad \downarrow \text{3284} \\
 & \frac{2(Ab - aB) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2)}{bd(a + b)} + \frac{2B \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{bd}
 \end{aligned}$$

input $\text{Int}[(A + B*\text{Cos}[c + d*x])]/(\text{Sqrt}[\text{Cos}[c + d*x]]*(a + b*\text{Cos}[c + d*x])),x]$

output $(2*B*EllipticF[(c + d*x)/2, 2])/(b*d) + (2*(A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d)$

3.368.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3481 `Int[(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(c_) + (d_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.368.4 Maple [A] (verified)

Time = 3.30 (sec) , antiderivative size = 217, normalized size of antiderivative = 3.56

method	result
default	$-\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\left(\text{ArcPi}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), -\frac{2b}{a-b}, \sqrt{2}\right)b + BF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), (a-b)b\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\right)}{(a-b)b\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}}$

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b), x, method=_RETURNVERBOSE)`

3.368.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx$$

output $-2*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*(A*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})*b+B*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a-B*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b-B*\text{EllipticPi}(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})*a)/(a-b)/b/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$

3.368.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output Timed out

3.368.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c)),x)`

output Timed out

3.368.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)\sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.368.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)\sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.368.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))), x)`

3.369 $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$

3.369.1 Optimal result 3437
 3.369.2 Mathematica [B] (verified) 3437
 3.369.3 Rubi [A] (verified) 3438
 3.369.4 Maple [B] (verified) 3441
 3.369.5 Fricas [F(-1)] 3442
 3.369.6 Sympy [F(-1)] 3442
 3.369.7 Maxima [F] 3442
 3.369.8 Giac [F] 3443
 3.369.9 Mupad [F(-1)] 3443

3.369.1 Optimal result

Integrand size = 33, antiderivative size = 86

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = -\frac{2AE(\frac{1}{2}(c + dx)|2)}{ad} - \frac{2(Ab - aB) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2)}{a(a + b)d} + \frac{2A \sin(c + dx)}{ad\sqrt{\cos(c + dx)}}$$

```
output -2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d-2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))/a/(a+b)/d+2*A*sin(d*x+c)/a/d/cos(d*x+c)^(1/2)
```

3.369.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 206 vs. 2(86) = 172.

Time = 1.68 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.40

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \frac{2(-3Ab+2aB) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{a+b} - \frac{2aA \left(2 \text{EllipticF}(\frac{1}{2}(c+dx), 2) - \frac{2a \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{a+b} \right)}{b} + \frac{4A \sin(c+dx)}{\sqrt{\cos(c+dx)}} - \frac{2A(-2abE}{2a}$$

3.369. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])),x
]`

output `((2*(-3*A*b + 2*a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) -
(2*a*A*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c +
d*x)/2, 2])/(a + b)))/b + (4*A*Sin[c + d*x])/Sqrt[Cos[c + d*x]] - (2*A*(-2
*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[Arc
Sin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sq
rt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b*Sqrt[Sin[c + d*x]^2]))/(2*a*d`

3.369.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3479, 27, 3042, 3538, 25, 27, 3042, 3119, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{2 \int -\frac{Ab \cos^2(c+dx) + aA \cos(c+dx) + Ab - aB}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} + \frac{2A \sin(c + dx)}{ad\sqrt{\cos(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2A \sin(c + dx)}{ad\sqrt{\cos(c + dx)}} - \frac{\int \frac{Ab \cos^2(c+dx) + aA \cos(c+dx) + Ab - aB}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2A \sin(c + dx)}{ad\sqrt{\cos(c + dx)}} - \frac{\int \frac{Ab \sin\left(c+dx+\frac{\pi}{2}\right)^2 + aA \sin\left(c+dx+\frac{\pi}{2}\right) + Ab - aB}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}(a+b \sin\left(c+dx+\frac{\pi}{2}\right))} dx}{a} \\
 & \quad \downarrow \text{3538}
 \end{aligned}$$

3.369. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$

$$\frac{2A \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{A \int \sqrt{\cos(c+dx)} dx - \frac{\int -\frac{b(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{a}}$$

↓ 25

$$\frac{2A \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\frac{\int \frac{b(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{b} + A \int \sqrt{\cos(c+dx)} dx}{a}$$

↓ 27

$$\frac{2A \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{(Ab-aB) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx + A \int \sqrt{\cos(c+dx)} dx}{a}$$

↓ 3042

$$\frac{2A \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx + A \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{a}$$

↓ 3119

$$\frac{2A \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx + \frac{2AE(\frac{1}{2}(c+dx)|2)}{d}}{a}$$

↓ 3284

$$\frac{2A \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\frac{2(Ab-aB) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{d(a+b)} + \frac{2AE(\frac{1}{2}(c+dx)|2)}{d}}{a}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])),x]`

output `-(((2*A*EllipticE[(c + d*x)/2, 2])/d + (2*(A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b*d))/a + (2*A*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]]))`

3.369.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.369.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(136) = 272.

Time = 4.60 (sec) , antiderivative size = 300, normalized size of antiderivative = 3.49

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{2A\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})} \left(\frac{2(\sin^2(\frac{dx}{2} + \frac{c}{2})) \cos(\frac{dx}{2} + \frac{c}{2}) - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}}{a \sin(\frac{dx}{2} + \frac{c}{2})^2 (2(\sin^2(\frac{dx}{2} + \frac{c}{2})) - 1)} \right) \sin(\frac{dx}{2} + \frac{c}{2}) \sqrt{2(\cos^2(\frac{dx}{2} + \frac{c}{2})) - 1}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b), x, method=_RETURNVERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^(1/2)*(2*A/a/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))-4*(-A*b+B*a)/a/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c), -2*b/(a-b), 2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.369. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$

3.369.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `Timed out`

3.369.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.369.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.369.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.369.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} (a + b \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))), x)`

$$3.370 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$$

3.370.1 Optimal result	3444
3.370.2 Mathematica [A] (verified)	3445
3.370.3 Rubi [A] (verified)	3445
3.370.4 Maple [A] (verified)	3450
3.370.5 Fracas [F(-1)]	3451
3.370.6 Sympy [F(-1)]	3451
3.370.7 Maxima [F]	3451
3.370.8 Giac [F]	3452
3.370.9 Mupad [F(-1)]	3452

3.370.1 Optimal result

Integrand size = 33, antiderivative size = 150

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx = \frac{2(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{a^2d} + \frac{2A \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{3ad} + \frac{2b(Ab-aB) \operatorname{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{a^2(a+b)d} + \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2(Ab-aB) \sin(c+dx)}{a^2d \sqrt{\cos(c+dx)}}$$

output

```
2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(
1/2*d*x+1/2*c), 2^(1/2))/a^2/d+2/3*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d
*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d+2*b*(A*b-B*a)*(cos(1/2
*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/
(a+b), 2^(1/2))/a^2/(a+b)/d+2/3*A*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)-2*(A*b-B*
a)*sin(d*x+c)/a^2/d/cos(d*x+c)^(1/2)
```

3.370.2 Mathematica [A] (verified)

Time = 1.57 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.73

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$= \frac{2a(2a^2A + 9Ab^2 - 9abB) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + \frac{a(8aAb - 6a^2B) \left(2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} \right)}{b} + \frac{4a^2A \sin(c + dx)}{\cos^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])),x]`

output `((2*a*(2*a^2*A + 9*A*b^2 - 9*a*b*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (a*(8*a*A*b - 6*a^2*B)*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)))/b + (4*a^2*A*Sin[c + d*x])/Cos[c + d*x]^(3/2) + (12*a*(-(A*b) + a*B)*Sin[c + d*x])/Sqrt[Cos[c + d*x]] + (6*(A*b - a*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(b*Sqrt[Sin[c + d*x]^2]))/(6*a^3*d)`

3.370.3 Rubi [A] (verified)

Time = 1.44 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.08, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx$$

$$\downarrow \text{3479}$$

3.370. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$

$$\begin{aligned}
 & \frac{2 \int -\frac{Ab \cos^2(c+dx) - aA \cos(c+dx) + 3(Ab-aB)}{2 \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx}{3a} + \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 27 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int -\frac{Ab \cos^2(c+dx) - aA \cos(c+dx) + 3(Ab-aB)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx}{3a} \\
 & \quad \downarrow 3042 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int -\frac{Ab \sin(c+dx+\frac{\pi}{2})^2 - aA \sin(c+dx+\frac{\pi}{2}) + 3(Ab-aB)}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3a} \\
 & \quad \downarrow 3534 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{2 \int -\frac{Aa^2 - 3bBa + (4Ab - 3aB) \cos(c+dx)a + 3Ab^2 + 3b(Ab-aB) \cos^2(c+dx)}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} + \frac{6(Ab-aB) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}}}{3a} \\
 & \quad \downarrow 27 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6(Ab-aB) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{Aa^2 - 3bBa + (4Ab - 3aB) \cos(c+dx)a + 3Ab^2 + 3b(Ab-aB) \cos^2(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a}}{3a} \\
 & \quad \downarrow 3042 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6(Ab-aB) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{Aa^2 - 3bBa + (4Ab - 3aB) \sin(c+dx+\frac{\pi}{2})a + 3Ab^2 + 3b(Ab-aB) \sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{a}}{3a} \\
 & \quad \downarrow 3538 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6(Ab-aB) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(Ab-aB) \int \sqrt{\cos(c+dx)} dx - \frac{\int -\frac{aA \cos(c+dx)b^2 + (Aa^2 - 3bBa + 3Ab^2)b}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a}}{3a}}{3a} \\
 & \quad \downarrow 25 \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6(Ab-aB) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{aA \cos(c+dx)b^2 + (Aa^2 - 3bBa + 3Ab^2)b}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} + 3(Ab-aB) \int \sqrt{\cos(c+dx)} dx}{3a}}{3a} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.370. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$

$$\begin{aligned}
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{aA \sin(c+dx+\frac{\pi}{2})b^2 + (Aa^2 - 3bBa + 3Ab^2)b}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx}{a} \\
 & \frac{6(Ab-aB) \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{3(Ab-aB) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{a} \\
 & \frac{3a}{3a} \quad \downarrow \quad \mathbf{3119} \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(Ab-aB) \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{\int \frac{aA \sin(c+dx+\frac{\pi}{2})b^2 + (Aa^2 - 3bBa + 3Ab^2)b}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx}{a} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d} \\
 & \frac{3a}{3a} \quad \downarrow \quad \mathbf{3481} \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(Ab-aB) \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{3b^2(Ab-aB) \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx + aAb \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d} \\
 & \frac{3a}{3a} \quad \downarrow \quad \mathbf{3042} \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{3b^2(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx + aAb \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d} \\
 & \frac{3a}{3a} \quad \downarrow \quad \mathbf{3120} \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{3b^2(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx + \frac{2aAb \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}}{a} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d} \\
 & \frac{3a}{3a} \quad \downarrow \quad \mathbf{3284} \\
 & \frac{2A \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6b^2(Ab-aB) \operatorname{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} + \frac{2aAb \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d} \\
 & \frac{3a}{3a}
 \end{aligned}$$

3.370. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])),x]`

output `(2*A*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)) - (-(((6*(A*b - a*B)*EllipticE[(c + d*x)/2, 2])/d + ((2*a*A*b*EllipticF[(c + d*x)/2, 2])/d + (6*b^2*(A*b - a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d))/b)/a) + (6*(A*b - a*B)*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]])/(3*a)`

3.370.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`


```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.370.4 Maple [A] (verified)

Time = 6.42 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.94

method	result
default	$\frac{\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{2A\left(\frac{-\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}{6\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^2} + \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}\right) + \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{a}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b), x, method=_RETURNVERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A/a*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2)))+2*(-A*b+B*a)/a^2/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))-4*(A*b-B*a)*b^2/a^2/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c), -2*b/(a-b), 2^(1/2))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.370. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$

3.370.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `Timed out`

3.370.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.370.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.370.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.370.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + b \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))), x)`

3.371
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.371.1 Optimal result 3453
 3.371.2 Mathematica [A] (verified) 3454
 3.371.3 Rubi [A] (verified) 3454
 3.371.4 Maple [B] (verified) 3459
 3.371.5 Fricas [F(-1)] 3460
 3.371.6 Sympy [F(-1)] 3461
 3.371.7 Maxima [F] 3461
 3.371.8 Giac [F] 3461
 3.371.9 Mupad [F(-1)] 3462

3.371.1 Optimal result

Integrand size = 33, antiderivative size = 303

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

$$= \frac{(3a^2Ab - 2Ab^3 - 5a^3B + 4ab^2B) E\left(\frac{1}{2}(c+dx) \mid 2\right)}{b^3(a^2 - b^2)d}$$

$$- \frac{(9a^3Ab - 12aAb^3 - 15a^4B + 16a^2b^2B + 2b^4B) \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3b^4(a^2 - b^2)d}$$

$$+ \frac{a^2(3a^2Ab - 5Ab^3 - 5a^3B + 7ab^2B) \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{(a-b)b^4(a+b)^2d}$$

$$- \frac{(3aAb - 5a^2B + 2b^2B) \sqrt{\cos(c+dx)} \sin(c+dx)}{3b^2(a^2 - b^2)d}$$

$$+ \frac{a(Ab - aB) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{b(a^2 - b^2)d(a+b \cos(c+dx))}$$

output

```
(3*A*a^2*b-2*A*b^3-5*B*a^3+4*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^3/(a^2-b^2)/d-1/3*(9*A*a^3*b-12*A*a*b^3-15*B*a^4+16*B*a^2*b^2+2*B*b^4)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b^4/(a^2-b^2)/d+a^2*(3*A*a^2*b-5*A*b^3-5*B*a^3+7*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/(a-b)/b^4/(a+b)^2/d+a*(A*b-B*a)*cos(d*x+c)^(3/2)*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))-1/3*(3*A*a*b-5*B*a^2+2*B*b^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/b^2/(a^2-b^2)/d
```

3.371.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.371.2 Mathematica [A] (verified)

Time = 2.38 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.05

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{4\sqrt{\cos(c+dx)}\left(2B + \frac{3a^2(-Ab+aB)}{(a^2-b^2)(a+b\cos(c+dx))}\right) \sin(c+dx) - \frac{2(-3a^2Ab+6Ab^3+5a^3B-8ab^2B)}{a+b} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right) + \frac{8(-3a^2Ab+6Ab^3+5a^3B-8ab^2B)}{a+b} \text{EllipticE}\left[\text{ArcSin}\left[\sqrt{\cos(c+dx)}\right], -1\right] + 2a(a+b)\text{EllipticF}\left[\text{ArcSin}\left[\sqrt{\cos(c+dx)}\right], -1\right] + (-2a^2+b^2)\text{EllipticPi}\left[-\frac{b}{a}, \text{ArcSin}\left[\sqrt{\cos(c+dx)}\right], -1\right] \sin(c+dx)}{(a-b)(a+b)(12b^2d)}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2, x]`

output `(4*sqrt[Cos[c + d*x]]*(2*B + (3*a^2*(-A*b) + a*B))/((a^2 - b^2)*(a + b*Cos[c + d*x]))*Sin[c + d*x] - ((2*(-3*a^2*A*b + 6*A*b^3 + 5*a^3*B - 8*a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]]/(a + b) + (8*(-3*a*A*b + 2*a^2*B + b^2*B))*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b) + (6*(-3*a^2*A*b + 2*A*b^3 + 5*a^3*B - 4*a*b^2*B))*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*sqrt[Sin[c + d*x]^2]))/((a - b)*(a + b)))/(12*b^2*d)`

3.371.3 Rubi [A] (verified)

Time = 1.87 (sec) , antiderivative size = 295, normalized size of antiderivative = 0.97, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3468, 27, 3042, 3528, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx$$

↓ 3468

3.371. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \\
& \frac{\int -\sqrt{\cos(c+dx)} \left((-5Ba^2 + 3Aba + 2b^2B) \cos^2(c+dx) - 2b(Ab - aB) \cos(c+dx) + 3a(Ab - aB) \right) dx}{2(a + b \cos(c+dx))} \\
& \frac{dx}{b(a^2 - b^2)} \\
& \quad \downarrow \text{27} \\
& \frac{\int \sqrt{\cos(c+dx)} \left((-5Ba^2 + 3Aba + 2b^2B) \cos^2(c+dx) - 2b(Ab - aB) \cos(c+dx) + 3a(Ab - aB) \right) dx}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \sqrt{\sin(c+dx+\frac{\pi}{2})} \left((5Ba^2 - 3Aba - 2b^2B) \sin^2(c+dx+\frac{\pi}{2}) - 2b(Ab - aB) \sin(c+dx+\frac{\pi}{2}) + 3a(Ab - aB) \right) dx}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3528} \\
& \frac{2 \int -\frac{-3(-5Ba^3 + 3Aba^2 + 4b^2Ba - 2Ab^3) \cos^2(c+dx) - 2b(-2Ba^2 + 3Aba - b^2B) \cos(c+dx) + a(-5Ba^2 + 3Aba + 2b^2B)}{2\sqrt{\cos(c+dx)}(a + b \cos(c+dx))} dx}{3b} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c+dx)}{3bd} \\
& \frac{2b(a^2 - b^2)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{27} \\
& \frac{\int -\frac{-3(-5Ba^3 + 3Aba^2 + 4b^2Ba - 2Ab^3) \cos^2(c+dx) - 2b(-2Ba^2 + 3Aba - b^2B) \cos(c+dx) + a(-5Ba^2 + 3Aba + 2b^2B)}{\sqrt{\cos(c+dx)}(a + b \cos(c+dx))} dx}{3b} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c+dx)}{3bd} \\
& \frac{2b(a^2 - b^2)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{\int -\frac{-3(-5Ba^3 + 3Aba^2 + 4b^2Ba - 2Ab^3) \sin^2(c+dx+\frac{\pi}{2}) - 2b(-2Ba^2 + 3Aba - b^2B) \sin(c+dx+\frac{\pi}{2}) + a(-5Ba^2 + 3Aba + 2b^2B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a + b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c+dx)}{3bd} \\
& \frac{2b(a^2 - b^2)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3538}
\end{aligned}$$

3.371. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{3(-5a^3B+3a^2Ab+4ab^2B-2Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} - \frac{\int -\frac{ab(-5Ba^2+3Aba+2b^2B)+(-15Ba^4+9Aba^3+16b^2Ba^2-12Ab^3a+2b^4B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} - 2(-5a^2B) \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{ab(-5Ba^2+3Aba+2b^2B)+(-15Ba^4+9Aba^3+16b^2Ba^2-12Ab^3a+2b^4B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} - \frac{3(-5a^3B+3a^2Ab+4ab^2B-2Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} - 2(-5a^2B) \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{ab(-5Ba^2+3Aba+2b^2B)+(-15Ba^4+9Aba^3+16b^2Ba^2-12Ab^3a+2b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{3(-5a^3B+3a^2Ab+4ab^2B-2Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} - 2(-5a^2B) \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow \text{3119} \\
 & \frac{\int \frac{ab(-5Ba^2+3Aba+2b^2B)+(-15Ba^4+9Aba^3+16b^2Ba^2-12Ab^3a+2b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{6(-5a^3B+3a^2Ab+4ab^2B-2Ab^3) E(\frac{1}{2}(c+dx)|2)}{bd} - 2(-5a^2B) \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow \text{3481} \\
 & \frac{(-15a^4B+9a^3Ab+16a^2b^2B-12aAb^3+2b^4B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{3a^2(-5a^3B+3a^2Ab+7ab^2B-5Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} - 6(-5a^3B+3a^2Ab+4ab^2B-2Ab^3) \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b \cos(c+dx))} \frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b \cos(c+dx))} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.371. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

$$\frac{(-15a^4B+9a^3Ab+16a^2b^2B-12aAb^3+2b^4B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^2(-5a^3B+3a^2Ab+7ab^2B-5Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{6(-5a^2B+3aAb+2b^2B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3bd} - \frac{2b(a^2-b^2)}{2b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3120

$$\frac{2(-15a^4B+9a^3Ab+16a^2b^2B-12aAb^3+2b^4B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{3a^2(-5a^3B+3a^2Ab+7ab^2B-5Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{6(-5a^2B+3aAb+2b^2B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3bd} - \frac{2b(a^2-b^2)}{2b(a^2-b^2)}$$

$$\frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))}$$

↓ 3284

$$\frac{a(Ab-aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{bd(a^2-b^2)(a+b\cos(c+dx))} + \frac{2(-15a^4B+9a^3Ab+16a^2b^2B-12aAb^3+2b^4B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{6a^2(-5a^3B+3a^2Ab+7ab^2B)}{b} - \frac{2(-5a^2B+3aAb+2b^2B) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} - \frac{2b(a^2-b^2)}{2b(a^2-b^2)}$$

```
input Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]
```

```
output (a*(A*b - a*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])) + (-1/3*((-6*(3*a^2*A*b - 2*A*b^3 - 5*a^3*B + 4*a*b^2*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((2*(9*a^3*A*b - 12*a*A*b^3 - 15*a^4*B + 16*a^2*b^2*B + 2*b^4*B)*EllipticF[(c + d*x)/2, 2])/(b*d) - (6*a^2*(3*a^2*A*b - 5*A*b^3 - 5*a^3*B + 7*a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))/b)/b - (2*(3*a*A*b - 5*a^2*B + 2*b^2*B)*Sqrt[Cos[c + d*x]])*Sin[c + d*x])/(3*b*d))/(2*b*(a^2 - b^2))
```

3.371. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.371.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.371.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

rule 3481 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3528 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3538 `Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.371.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1065 vs. 2(373) = 746.

Time = 19.24 (sec) , antiderivative size = 1066, normalized size of antiderivative = 3.52

method	result	size
default	Expression too large to display	1066

input `int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNV ERBOSE)`

$$3.371. \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*a^2/b^3*(3*
A*b-4*B*a)/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2
*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipt
icPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a^3*(A*b-B*a)/b^4*(-1/a*b^2/
(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d
*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^
(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*
c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^
4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(
1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*co
s(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))-2/3/b^4/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B*cos(1/2*d*x+1/2*c)*sin(
1/2*d*x+1/2*c)^4*b^2+6*A*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*...

```

3.371.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output `Timed out`

3.371. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

3.371.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`output `Timed out`**3.371.7 Maxima [F]**

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^2, x)`**3.371.8 Giac [F]**

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^2, x)`

3.371.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2, x)`output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2, x)`

3.372
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.372.1 Optimal result 3463
 3.372.2 Mathematica [A] (verified) 3464
 3.372.3 Rubi [A] (verified) 3464
 3.372.4 Maple [B] (verified) 3468
 3.372.5 Fricas [F(-1)] 3469
 3.372.6 Sympy [F(-1)] 3470
 3.372.7 Maxima [F] 3470
 3.372.8 Giac [F] 3470
 3.372.9 Mupad [F(-1)] 3471

3.372.1 Optimal result

Integrand size = 33, antiderivative size = 224

$$\begin{aligned} & \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx \\ &= -\frac{(aAb - 3a^2B + 2b^2B) E(\frac{1}{2}(c+dx)|2)}{b^2(a^2 - b^2)d} \\ & \quad + \frac{(a^2Ab - 2Ab^3 - 3a^3B + 4ab^2B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{b^3(a^2 - b^2)d} \\ & \quad - \frac{a(a^2Ab - 3Ab^3 - 3a^3B + 5ab^2B) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{(a-b)b^3(a+b)^2d} \\ & \quad + \frac{a(Ab - aB)\sqrt{\cos(c+dx)} \sin(c+dx)}{b(a^2 - b^2)d(a+b \cos(c+dx))} \end{aligned}$$

output

```
-(A*a*b-3*B*a^2+2*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/(a^2-b^2)/d+(A*a^2*b-2*A*b^3-3*B*
a^3+4*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(s
in(1/2*d*x+1/2*c),2^(1/2))/b^3/(a^2-b^2)/d-a*(A*a^2*b-3*A*b^3-3*B*a^3+5*B*
a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*
d*x+1/2*c),2*b/(a+b),2^(1/2))/(a-b)/b^3/(a+b)^2/d+a*(A*b-B*a)*sin(d*x+c)*c
os(d*x+c)^(1/2)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.372.2 Mathematica [A] (verified)

Time = 1.95 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.25

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{-\frac{4a(-Ab+aB)\sqrt{\cos(c+dx)}\sin(c+dx)}{(a^2-b^2)(a+b\cos(c+dx))} + \frac{2(aAb+a^2B-2b^2B)\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + \frac{8(-Ab+aB)((a+b)\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - a\operatorname{EllipticPi}\left(\frac{2b}{a+b}\right))}{a+b}}{1}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2, x]`

output `((-4*a*(-A*b) + a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*(a + b*Cos[c + d*x])) + ((2*(a*A*b + a^2*B - 2*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (8*(-A*b) + a*B)*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b) + (2*(-(a*A*b) + 3*a^2*B - 2*b^2*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*Sqrt[Sin[c + d*x]^2]))/((a - b)*(a + b)))/(4*b*d)`

3.372.3 Rubi [A] (verified)

Time = 1.31 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.98, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3468, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx$$

↓ 3468

3.372. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2)(a + b \cos(c + dx))} - \\
& \frac{\int - \frac{((-3Ba^2 + Aba + 2b^2B) \cos^2(c + dx)) - 2b(Ab - aB) \cos(c + dx) + a(Ab - aB)}{2\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{b(a^2 - b^2)} \\
& \quad \downarrow \text{27} \\
& \frac{\int - \frac{((-3Ba^2 + Aba + 2b^2B) \cos^2(c + dx)) - 2b(Ab - aB) \cos(c + dx) + a(Ab - aB)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(3Ba^2 - Aba - 2b^2B) \sin(c + dx + \frac{\pi}{2})^2 - 2b(Ab - aB) \sin(c + dx + \frac{\pi}{2}) + a(Ab - aB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3538} \\
& - \frac{(-3a^2B + aAb + 2b^2B) \int \sqrt{\cos(c + dx)} dx}{b} - \frac{\int - \frac{ab(Ab - aB) + (-3Ba^3 + Aba^2 + 4b^2Ba - 2Ab^3) \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{b} + \\
& \frac{2b(a^2 - b^2)}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{ab(Ab - aB) + (-3Ba^3 + Aba^2 + 4b^2Ba - 2Ab^3) \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{b} - \frac{(-3a^2B + aAb + 2b^2B) \int \sqrt{\cos(c + dx)} dx}{b} + \\
& \frac{2b(a^2 - b^2)}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{ab(Ab - aB) + (-3Ba^3 + Aba^2 + 4b^2Ba - 2Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx}{b} - \frac{(-3a^2B + aAb + 2b^2B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx}{b} + \\
& \frac{2b(a^2 - b^2)}{2b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2)(a + b \cos(c + dx))} \\
& \quad \downarrow \text{3119}
\end{aligned}$$

3.372. $\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx$

$$\begin{aligned}
 & \frac{\int \frac{ab(Ab-aB)+(-3Ba^3+Ab^2+4b^2Ba-2Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{2(-3a^2B+aAb+2b^2B)E(\frac{1}{2}(c+dx)|2)}{bd} + \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} + \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3481} \\
 & \frac{(-3a^3B+a^2Ab+4ab^2B-2Ab^3)\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a(-3a^3B+a^2Ab+5ab^2B-3Ab^3)\int \frac{1}{\sqrt{\cos(c+dx)(a+b\cos(c+dx))}} dx}{b} - \frac{2(-3a^2B+aAb+2b^2B)E(\frac{1}{2}(c+dx))}{bd} + \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} + \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-3a^3B+a^2Ab+4ab^2B-2Ab^3)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{a(-3a^3B+a^2Ab+5ab^2B-3Ab^3)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{2(-3a^2B+aAb+2b^2B)E(\frac{1}{2}(c+dx))}{bd} + \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} + \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2(-3a^3B+a^2Ab+4ab^2B-2Ab^3)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{a(-3a^3B+a^2Ab+5ab^2B-3Ab^3)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{2(-3a^2B+aAb+2b^2B)E(\frac{1}{2}(c+dx))}{bd} + \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} + \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b\cos(c+dx))} \\
 & \quad \downarrow \text{3284} \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b\cos(c+dx))} + \\
 & \frac{2(-3a^3B+a^2Ab+4ab^2B-2Ab^3)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{2a(-3a^3B+a^2Ab+5ab^2B-3Ab^3)\text{EllipticPi}(\frac{-2b}{a+b},\frac{1}{2}(c+dx),2)}{bd(a+b)} - \frac{2(-3a^2B+aAb+2b^2B)E(\frac{1}{2}(c+dx))}{bd} + \\
 & \frac{2b(a^2-b^2)}{bd(a^2-b^2)(a+b\cos(c+dx))} +
 \end{aligned}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

3.372. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

```
output ((-2*(a*A*b - 3*a^2*B + 2*b^2*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((2*(a
^2*A*b - 2*A*b^3 - 3*a^3*B + 4*a*b^2*B)*EllipticF[(c + d*x)/2, 2])/(b*d) -
(2*a*(a^2*A*b - 3*A*b^3 - 3*a^3*B + 5*a*b^2*B)*EllipticPi[(2*b)/(a + b),
(c + d*x)/2, 2])/(b*(a + b)*d))/b)/(2*b*(a^2 - b^2)) + (a*(A*b - a*B)*Sqrt
[Cos[c + d*x]]*Sin[c + d*x])/(b*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))
```

3.372.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

```
rule 3468 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c
+ d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n +
1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n
+ 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a
*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1)
- a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c -
a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x]
/; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2
, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 3481 Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.372.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 848 vs. $2(300) = 600$.

Time = 7.79 (sec) , antiderivative size = 849, normalized size of antiderivative = 3.79

method	result	size
default	Expression too large to display	849

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

$$3.372. \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

output
$$\begin{aligned} & -(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2/b^3/(-2*\sin(\\ & 1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)} \\ & *(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(A*b*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)} \\ &)-2*B*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a-B*b*EllipticE(\cos(1/2*d*x+1/ \\ & 2*c),2^{(1/2)}))+2*a^2*(A*b-B*a)/b^3*(-1/a*b^2/(a^2-b^2)*\cos(1/2*d*x+1/2*c)* \\ & (-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(2*b*\cos(1/2*d*x+1/2* \\ & c)^2+a-b)-1/2/a/(a+b)*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^ \\ & 2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\\ & \cos(1/2*d*x+1/2*c),2^{(1/2)})-1/2/(a^2-b^2)*b/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & *(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/ \\ & 2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+1/2/(a^2-b^2)*b/a*(\sin \\ & (1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x \\ & +1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)} \\ &)-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2* \\ & d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ & *EllipticPi(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})+1/a/(a^2-b^2)/(-2*a*b+2 \\ & *b^2)*b^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(\\ & -2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticPi(\cos(1/2*d*x \\ & +1/2*c),-2*b/(a-b),2^{(1/2)}))+4*a/b^2*(2*A*b-3*B*a)/(-2*a*b+2*b^2)*(\sin(1/2 \\ & *d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+... \end{aligned}$$

3.372.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `Timed out`

3.372.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`output `Timed out`**3.372.7 Maxima [F]**

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^2, x)`**3.372.8 Giac [F]**

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^2, x)`

3.372.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2, x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2, x)`

3.373 $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.373.1 Optimal result 3472
 3.373.2 Mathematica [A] (verified) 3473
 3.373.3 Rubi [A] (verified) 3473
 3.373.4 Maple [B] (verified) 3477
 3.373.5 Fracas [F(-1)] 3478
 3.373.6 Sympy [F(-1)] 3479
 3.373.7 Maxima [F] 3479
 3.373.8 Giac [F] 3479
 3.373.9 Mupad [F(-1)] 3480

3.373.1 Optimal result

Integrand size = 33, antiderivative size = 198

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

$$= \frac{(Ab - aB)E(\frac{1}{2}(c+dx)|2)}{b(a^2 - b^2)d} + \frac{(aAb + a^2B - 2b^2B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{b^2(a^2 - b^2)d}$$

$$- \frac{(a^2Ab + Ab^3 + a^3B - 3ab^2B) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{(a-b)b^2(a+b)^2d}$$

$$- \frac{(Ab - aB)\sqrt{\cos(c+dx)} \sin(c+dx)}{(a^2 - b^2)d(a+b \cos(c+dx))}$$

```
output (A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/b/(a^2-b^2)/d+(A*a*b+B*a^2-2*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/b^2/(a^2-b^2)/d-(A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))/(a-b)/b^2/(a+b)^2/d-(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/(a^2-b^2)/d/(a+b*cos(d*x+c))
```

3.373.2 Mathematica [A] (verified)

Time = 1.83 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= \frac{4(-Ab+aB)\sqrt{\cos(c+dx)}\sin(c+dx)}{(a^2-b^2)(a+b\cos(c+dx))} - \frac{2(-Ab+aB)\operatorname{EllipticPi}\left(\frac{-2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + \frac{(4aA-4bB)\left(2\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b}\right)}{b}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2, x]`

output `((4*(-A*b) + a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*(a + b*Cos[c + d*x])) - ((2*(-A*b) + a*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + ((4*a*A - 4*b*B)*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b))/b + (2*(A*b - a*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*Sqrt[Sin[c + d*x]^2])/((-a + b)*(a + b))/(4*d)`

3.373.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.97, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3478, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx$$

↓ 3478

3.373. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
& \frac{\int \frac{-((Ab-aB)\cos^2(c+dx))-2(aA-bB)\cos(c+dx)+Ab-aB}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{a^2-b^2} = \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{-((Ab-aB)\cos^2(c+dx))-2(aA-bB)\cos(c+dx)+Ab-aB}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{2(a^2-b^2)} = \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(aB-Ab)\sin(c+dx+\frac{\pi}{2})-2(aA-bB)\sin(c+dx+\frac{\pi}{2})+Ab-aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{2(a^2-b^2)} = \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 3538 \\
& \frac{\int \frac{b(Ab-aB)-\frac{(Ba^2+Ab-a-2b^2B)\cos(c+dx)}{b}}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{2(a^2-b^2)} = \frac{(Ab-aB)\int \frac{\sqrt{\cos(c+dx)} dx}{b}}{2(a^2-b^2)} \\
& \quad \quad \quad \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 25 \\
& \frac{\int \frac{b(Ab-aB)-\frac{(Ba^2+Ab-a-2b^2B)\cos(c+dx)}{b}}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{2(a^2-b^2)} = \frac{(Ab-aB)\int \frac{\sqrt{\cos(c+dx)} dx}{b}}{2(a^2-b^2)} = \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{b(Ab-aB)+\frac{(-Ba^2-Ab-a+2b^2B)\sin(c+dx+\frac{\pi}{2})}{b}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{2(a^2-b^2)} = \frac{(Ab-aB)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b}}{2(a^2-b^2)} \\
& \quad \quad \quad \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 3119 \\
& \frac{\int \frac{b(Ab-aB)+\frac{(-Ba^2-Ab-a+2b^2B)\sin(c+dx+\frac{\pi}{2})}{b}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{2(a^2-b^2)} = \frac{2(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{bd} \\
& \quad \quad \quad \frac{(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} \\
& \quad \downarrow 3481
\end{aligned}$$

3.373. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\frac{\frac{(a^3 B + a^2 A b - 3 a b^2 B + A b^3) \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx}{b} - \frac{(a^2 B + a A b - 2 b^2 B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b}}{b} - \frac{2(Ab - aB)E\left(\frac{1}{2}(c+dx)|2\right)}{bd}$$

$$\frac{2(a^2 - b^2)}{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}} \frac{1}{d(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3042

$$\frac{\frac{(a^3 B + a^2 A b - 3 a b^2 B + A b^3) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b \sin\left(c+dx+\frac{\pi}{2}\right))}} dx}{b} - \frac{(a^2 B + a A b - 2 b^2 B) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{b}}{b} - \frac{2(Ab - aB)E\left(\frac{1}{2}(c+dx)|2\right)}{bd}$$

$$\frac{2(a^2 - b^2)}{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}} \frac{1}{d(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3120

$$\frac{\frac{(a^3 B + a^2 A b - 3 a b^2 B + A b^3) \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b \sin\left(c+dx+\frac{\pi}{2}\right))}} dx}{b} - \frac{2(a^2 B + a A b - 2 b^2 B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd}}{b} - \frac{2(Ab - aB)E\left(\frac{1}{2}(c+dx)|2\right)}{bd}$$

$$\frac{2(a^2 - b^2)}{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}} \frac{1}{d(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3284

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{d(a^2 - b^2)(a + b \cos(c + dx))} - \frac{2(a^3 B + a^2 A b - 3 a b^2 B + A b^3) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{bd(a+b)} - \frac{2(a^2 B + a A b - 2 b^2 B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{2(Ab - aB)E\left(\frac{1}{2}(c+dx)|2\right)}{bd}$$

$$\frac{1}{2(a^2 - b^2)}$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `-1/2*((-2*(A*b - a*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((-2*(a*A*b + a^2*B - 2*b^2*B)*EllipticF[(c + d*x)/2, 2])/(b*d) + (2*(a^2*A*b + A*b^3 + a^3*B - 3*a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))/b)/(a^2 - b^2) - ((A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))`

3.373. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.373.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3478 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
] && NeQ[c^2 - d^2, 0]
```

3.373.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 807 vs. $2(274) = 548$.

Time = 6.71 (sec) , antiderivative size = 808, normalized size of antiderivative = 4.08

method	result	size
default	Expression too large to display	808

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

output `-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B/b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-4/b*(A*b-2*B*a)/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a*(A*b-B*a)/b^2*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))/sin(1/2*d...`

3.373.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `Timed out`

3.373.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.373.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x+c)+A)*sqrt(cos(d*x+c))/(b*cos(d*x+c)+a)^2,x)`

3.373.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x+c)+A)*sqrt(cos(d*x+c))/(b*cos(d*x+c)+a)^2,x)`

3.373.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2, x)`output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^2, x)`

3.374 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))^2}} dx$

3.374.1 Optimal result 3481
 3.374.2 Mathematica [A] (verified) 3482
 3.374.3 Rubi [A] (verified) 3482
 3.374.4 Maple [B] (verified) 3486
 3.374.5 Fracas [F(-1)] 3487
 3.374.6 Sympy [F(-1)] 3488
 3.374.7 Maxima [F] 3488
 3.374.8 Giac [F] 3488
 3.374.9 Mupad [F(-1)] 3489

3.374.1 Optimal result

Integrand size = 33, antiderivative size = 200

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))^2}} dx$$

$$= -\frac{(Ab - aB)E\left(\frac{1}{2}(c + dx) \mid 2\right)}{a(a^2 - b^2)d} - \frac{(Ab - aB) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{b(a^2 - b^2)d}$$

$$+ \frac{(3a^2Ab - Ab^3 - a^3B - ab^2B) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{a(a - b)b(a + b)^2d}$$

$$+ \frac{b(Ab - aB)\sqrt{\cos(c + dx)} \sin(c + dx)}{a(a^2 - b^2)d(a + b \cos(c + dx))}$$

output

```
-(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a/(a^2-b^2)/d-(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b/(a^2-b^2)/d+(3*A*a^2*b-A*b^3-B*a^3-B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/a/(a-b)/b/(a+b)^2/d+b*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))
```


3.374.2 Mathematica [A] (verified)

Time = 1.92 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.37

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$$

$$= \frac{4b(Ab - aB)\sqrt{\cos(c + dx)} \sin(c + dx)}{(a^2 - b^2)(a + b \cos(c + dx))} + \frac{2(4a^2A - 3Ab^2 - abB) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{a+b} + \frac{4a(-Ab + aB)}{b} \left(2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{a+b} \right)$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2), x]`

output `((4*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*(a + b*Cos[c + d*x])) + ((2*(4*a^2*A - 3*A*b^2 - a*b*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (4*a*(-(A*b) + a*B)*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b))/b + (2*(-(A*b) + a*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b*Sqrt[Sin[c + d*x]^2]))/((a - b)*(a + b)))/(4*a*d)`

3.374.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.92, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3479, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))^2} dx$$

↓ 3479

3.374. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} dx$

$$\frac{\int \frac{2Aa^2 - bBa - 2(Ab - aB) \cos(c + dx)a - Ab^2 - b(Ab - aB) \cos^2(c + dx)}{2\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{a(a^2 - b^2)} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 27

$$\frac{\int \frac{2Aa^2 - bBa - 2(Ab - aB) \cos(c + dx)a - Ab^2 - b(Ab - aB) \cos^2(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{2a(a^2 - b^2)} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3042

$$\frac{\int \frac{2Aa^2 - bBa - 2(Ab - aB) \sin(c + dx + \frac{\pi}{2})a - Ab^2 - b(Ab - aB) \sin^2(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx}{2a(a^2 - b^2)} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3538

$$\frac{-\int \frac{b(2Aa^2 - bBa - Ab^2) - ab(Ab - aB) \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{2a(a^2 - b^2)} - \left((Ab - aB) \int \sqrt{\cos(c + dx)} dx \right) + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 25

$$\frac{\int \frac{b(2Aa^2 - bBa - Ab^2) - ab(Ab - aB) \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{2a(a^2 - b^2)} - (Ab - aB) \int \sqrt{\cos(c + dx)} dx + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3042

$$\frac{\int \frac{b(2Aa^2 - bBa - Ab^2) - ab(Ab - aB) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx}{2a(a^2 - b^2)} - (Ab - aB) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

↓ 3119

$$\frac{\int \frac{b(2Aa^2 - bBa - Ab^2) - ab(Ab - aB) \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx}{2a(a^2 - b^2)} - \frac{2(Ab - aB)E(\frac{1}{2}(c + dx)|2)}{d} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{ad(a^2 - b^2)(a + b \cos(c + dx))}$$

3.374. $\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$

$$\begin{aligned} & \downarrow \text{3481} \\ & \frac{(a^3(-B)+3a^2Ab-ab^2B-Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)(a+b\cos(c+dx))}} dx - a(Ab-aB) \int \frac{1}{\sqrt{\cos(c+dx)}} dx - \frac{2(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d}}{2a(a^2-b^2)} + \\ & \frac{b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ & \frac{(a^3(-B)+3a^2Ab-ab^2B-Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx - a(Ab-aB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d}}{2a(a^2-b^2)} + \\ & \frac{b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3120} \\ & \frac{(a^3(-B)+3a^2Ab-ab^2B-Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx - \frac{2a(Ab-aB) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{d}}{2a(a^2-b^2)} - \frac{2(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d} + \\ & \frac{b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3284} \\ & \frac{b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} + \\ & \frac{\frac{2(a^3(-B)+3a^2Ab-ab^2B-Ab^3) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} - \frac{2a(Ab-aB) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}}{2a(a^2-b^2)} - \frac{2(Ab-aB)E(\frac{1}{2}(c+dx)|2)}{d}}{2a(a^2-b^2)} \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2),x]`

output `((-2*(A*b - a*B)*EllipticE[(c + d*x)/2, 2])/d + ((-2*a*(A*b - a*B)*EllipticF[(c + d*x)/2, 2])/d + (2*(3*a^2*A*b - A*b^3 - a^3*B - a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d))/b/(2*a*(a^2 - b^2)) + (b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))`

3.374.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
] && NeQ[c^2 - d^2, 0]
```

3.374.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 720 vs. 2(276) = 552.

Time = 6.46 (sec) , antiderivative size = 721, normalized size of antiderivative = 3.60

method	result
default	$-\frac{\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(-2ab + 2b^2\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}} + \frac{4B\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \Pi\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), -\frac{2b}{a-b}, \sqrt{2}\right) + \frac{2(Ab - Ba)}{\dots}}{\dots}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

$$3.374. \int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))^2}} dx$$

output
$$\begin{aligned}
& -(-(-2\cos(1/2dx+1/2c)^2+1)\sin(1/2dx+1/2c)^2)^{1/2}*(-4B/(-2ab+2 \\
& *b^2)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin \\
& \sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2dx+1/2 \\
& *c),-2b/(a-b),2^{1/2})+2*(A*b-B*a)/b*(-1/a*b^2/(a^2-b^2)*\cos(1/2dx+1/2 \\
& c)*(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}/(2*b*\cos(1/2dx+1/2 \\
& /2c)^2+a-b)-1/2/a/(a+b)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2 \\
& c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{Ellipti \\
& cF}(\cos(1/2dx+1/2c),2^{1/2})-1/2/(a^2-b^2)*b/a*(\sin(1/2dx+1/2c)^2)^{1/2} \\
& *(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx \\
& +1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})+1/2/(a^2-b^2)*b/a*(\\
& \sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2 \\
& dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticE}(\cos(1/2dx+1/2c),2^{1/2}) \\
&)-3a/(a^2-b^2)/(-2ab+2b^2)*b*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2 \\
& /2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2} \\
& * \text{EllipticPi}(\cos(1/2dx+1/2c),-2b/(a-b),2^{1/2})+1/a/(a^2-b^2)/(-2a \\
& b+2b^2)*b^3*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2} \\
&)/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2 \\
& dx+1/2c),-2b/(a-b),2^{1/2}))/\sin(1/2dx+1/2c)/(2*\cos(1/2dx+1/2c)^ \\
& 2-1)^{1/2}/d
\end{aligned}$$

3.374.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output Timed out

3.374.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.374.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sqrt(cos(d*x + c))), x)`

3.374.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sqrt(cos(d*x + c))), x)`

3.374.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^2),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^2), x)`

3.375
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

3.375.1 Optimal result 3490
 3.375.2 Mathematica [A] (verified) 3491
 3.375.3 Rubi [A] (verified) 3491
 3.375.4 Maple [B] (verified) 3496
 3.375.5 Fracas [F(-1)] 3497
 3.375.6 Sympy [F(-1)] 3498
 3.375.7 Maxima [F(-1)] 3498
 3.375.8 Giac [F] 3498
 3.375.9 Mupad [F(-1)] 3499

3.375.1 Optimal result

Integrand size = 33, antiderivative size = 256

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$= -\frac{(2a^2A - 3Ab^2 + abB) E(\frac{1}{2}(c + dx)|2)}{a^2(a^2 - b^2)d} + \frac{(Ab - aB) \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{a(a^2 - b^2)d}$$

$$- \frac{(5a^2Ab - 3Ab^3 - 3a^3B + ab^2B) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2)}{a^2(a - b)(a + b)^2d}$$

$$+ \frac{(2a^2A - 3Ab^2 + abB) \sin(c + dx)}{a^2(a^2 - b^2)d\sqrt{\cos(c + dx)}} + \frac{b(Ab - aB) \sin(c + dx)}{a(a^2 - b^2)d\sqrt{\cos(c + dx)}(a + b \cos(c + dx))}$$

```
output -(2*A*a^2-3*A*b^2+B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/(a^2-b^2)/d+(A*b-B*a)*(cos(1/2*d*
x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))
/a/(a^2-b^2)/d-(5*A*a^2*b-3*A*b^3-3*B*a^3+B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/
a^2/(a-b)/(a+b)^2/d+(2*A*a^2-3*A*b^2+B*a*b)*sin(d*x+c)/a^2/(a^2-b^2)/d/cos
(d*x+c)^(1/2)+b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))/cos(d*
x+c)^(1/2)
```

3.375.2 Mathematica [A] (verified)

Time = 2.90 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.23

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$\frac{2(-10a^2Ab+9Ab^3+4a^3B-3ab^2B)}{a+b} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right) - \frac{8a(a^2A-2Ab^2+abB)}{a+b} \left(\frac{(a+b) \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - a \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{b(a+b)}\right) - \frac{2(2a^2A-2Ab^2+abB)}{a+b} \text{EllipticE}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2(2a^2A-2Ab^2+abB)}{a+b} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2(2a^2A-2Ab^2+abB)}{a+b} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{b(a+b)}$$

```
input Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2),x]
```

```
output (-(((2*(-10*a^2*A*b + 9*A*b^3 + 4*a^3*B - 3*a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]]/(a + b) - (8*a*(a^2*A - 2*A*b^2 + a*b*B)*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(b*(a + b)) - (2*(2*a^2*A - 3*A*b^2 + a*b*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b*Sqrt[Sin[c + d*x]^2]))/((-a + b)*(a + b))) + 4*Sqrt[Cos[c + d*x]]*((b^2*(A*b - a*B)*Sin[c + d*x])/((-a^2 + b^2)*(a + b*Cos[c + d*x])) + 2*A*Tan[c + d*x]))/(4*a^2*d)
```

3.375.3 Rubi [A] (verified)

Time = 1.89 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.95, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^2} dx$$

↓ 3479

3.375. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{\int \frac{2Aa^2+bBa-2(Ab-aB)\cos(c+dx)a-3Ab^2+b(Ab-aB)\cos^2(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} dx}{\frac{a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)}} + \\
 & \frac{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}{\downarrow 27} \\
 & \frac{\int \frac{2Aa^2+bBa-2(Ab-aB)\cos(c+dx)a-3Ab^2+b(Ab-aB)\cos^2(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} dx}{\frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)}} + \\
 & \frac{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}{\downarrow 3042} \\
 & \frac{\int \frac{2Aa^2+bBa-2(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-3Ab^2+b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{\frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)}} + \\
 & \frac{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}{\downarrow 3534} \\
 & \frac{2\int -\frac{-2Ba^3+4Aba^2+b^2Ba+2(Aa^2+bBa-2Ab^2)\cos(c+dx)a-3Ab^3+b(2Aa^2+bBa-3Ab^2)\cos^2(c+dx)}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{a} + \frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} + \\
 & \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} \\
 & \frac{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}{\downarrow 27} \\
 & \frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{-2Ba^3+4Aba^2+b^2Ba+2(Aa^2+bBa-2Ab^2)\cos(c+dx)a-3Ab^3+b(2Aa^2+bBa-3Ab^2)\cos^2(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx}{a} + \\
 & \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} \\
 & \frac{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}{\downarrow 3042} \\
 & \frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{-2Ba^3+4Aba^2+b^2Ba+2(Aa^2+bBa-2Ab^2)\sin(c+dx+\frac{\pi}{2})a-3Ab^3+b(2Aa^2+bBa-3Ab^2)\sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{a} + \\
 & \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} \\
 & \frac{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}{\downarrow 3042}
 \end{aligned}$$

3.375. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2} dx$

↓ 3538

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{(2a^2A+abB-3Ab^2) \int \sqrt{\cos(c+dx)} dx - \frac{\int \frac{b(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) - ab^2(Ab-aB) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a}}{ad\sqrt{\cos(c+dx)}} +$$

$$\frac{2a(a^2-b^2) b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))}$$

↓ 25

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{(2a^2A+abB-3Ab^2) \int \sqrt{\cos(c+dx)} dx + \frac{\int \frac{b(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) - ab^2(Ab-aB) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a}}{ad\sqrt{\cos(c+dx)}} +$$

$$\frac{2a(a^2-b^2) b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))}$$

↓ 3042

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{(2a^2A+abB-3Ab^2) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + \frac{\int \frac{b(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) - ab^2(Ab-aB) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{a}}{ad\sqrt{\cos(c+dx)}} +$$

$$\frac{2a(a^2-b^2) b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))}$$

↓ 3119

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\frac{\int \frac{b(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) - ab^2(Ab-aB) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{a} + \frac{2(2a^2A+abB-3Ab^2) E(\frac{1}{2}(c+dx)|2)}{d}}{ad\sqrt{\cos(c+dx)}} +$$

$$\frac{2a(a^2-b^2) b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))}$$

↓ 3481

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\frac{b(-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx - ab(Ab-aB) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a} + \frac{2(2a^2A+abB-3Ab^2)}{d}}{ad\sqrt{\cos(c+dx)}} +$$

$$\frac{2a(a^2-b^2) b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))}$$

3.375. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

↓ 3042

$$\frac{\frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b(-3a^3B+5a^2Ab+ab^2B-3Ab^3)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx - ab(Ab-aB)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a} + \frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}}{2a(a^2-b^2)} = \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}$$

↓ 3120

$$\frac{\frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b(-3a^3B+5a^2Ab+ab^2B-3Ab^3)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx - \frac{2ab(Ab-aB)\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d}}{a} + \frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}}{2a(a^2-b^2)} = \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}$$

↓ 3284

$$\frac{\frac{2(2a^2A+abB-3Ab^2)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{2(2a^2A+abB-3Ab^2)E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2b(-3a^3B+5a^2Ab+ab^2B-3Ab^3)\text{EllipticPi}(\frac{2b}{a+b},\frac{1}{2}(c+dx),2)}{d(a+b)} - \frac{2ab(Ab-aB)\text{EllipticE}(\frac{1}{2}(c+dx),2)}{d}}{a} + \frac{b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}}{2a(a^2-b^2)}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2),x]`

output `(b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])) + (-(((2*(2*a^2*A - 3*A*b^2 + a*b*B)*EllipticE[(c + d*x)/2, 2])/d + ((-2*a*b*(A*b - a*B)*EllipticF[(c + d*x)/2, 2])/d + (2*b*(5*a^2*A*b - 3*A*b^3 - 3*a^3*B + a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)*d)/b)/a) + (2*(2*a^2*A - 3*A*b^2 + a*b*B)*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]])/(2*a*(a^2 - b^2))`

3.375. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

3.375.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.375.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 855 vs. $2(330) = 660$.

Time = 7.46 (sec) , antiderivative size = 856, normalized size of antiderivative = 3.34

method	result	size
default	Expression too large to display	856

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

$$3.375. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A/a^2/sin(1/
2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+
1/2*c),2^(1/2)))+4*A*b^2/a^2/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*
c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+2*(-A*b+B*a)
/a*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/
(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/
2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos
(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-
2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1...

```

3.375.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output `Timed out`

3.375.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.375.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`

output `Timed out`

3.375.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2)), x)`

3.375.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{3}{2}}(a + b \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^2),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^2), x)`

3.376
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

3.376.1 Optimal result 3500
 3.376.2 Mathematica [A] (verified) 3501
 3.376.3 Rubi [A] (verified) 3501
 3.376.4 Maple [B] (verified) 3507
 3.376.5 Fricas [F(-1)] 3508
 3.376.6 Sympy [F(-1)] 3509
 3.376.7 Maxima [F] 3509
 3.376.8 Giac [F] 3509
 3.376.9 Mupad [F(-1)] 3510

3.376.1 Optimal result

Integrand size = 33, antiderivative size = 345

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$= \frac{(4a^2 Ab - 5Ab^3 - 2a^3 B + 3ab^2 B) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{a^3 (a^2 - b^2) d}$$

$$+ \frac{(2a^2 A - 5Ab^2 + 3abB) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3a^2 (a^2 - b^2) d}$$

$$+ \frac{b(7a^2 Ab - 5Ab^3 - 5a^3 B + 3ab^2 B) \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{a^3 (a - b)(a + b)^2 d}$$

$$+ \frac{(2a^2 A - 5Ab^2 + 3abB) \sin(c + dx)}{3a^2 (a^2 - b^2) d \cos^{\frac{3}{2}}(c + dx)} - \frac{(4a^2 Ab - 5Ab^3 - 2a^3 B + 3ab^2 B) \sin(c + dx)}{a^3 (a^2 - b^2) d \sqrt{\cos(c + dx)}}$$

$$+ \frac{b(Ab - aB) \sin(c + dx)}{a (a^2 - b^2) d \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}$$

output

```
(4*A*a^2*b-5*A*b^3-2*B*a^3+3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^3/(a^2-b^2)/d+1/3*(2*A*a^2-5*A*b^2+3*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/(a^2-b^2)/d+b*(7*A*a^2*b-5*A*b^3-5*B*a^3+3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/a^3/(a-b)/(a+b)^2/d+1/3*(2*A*a^2-5*A*b^2+3*B*a*b)*sin(d*x+c)/a^2/(a^2-b^2)/d/cos(d*x+c)^(3/2)+b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))-(4*A*a^2*b-5*A*b^3-2*B*a^3+3*B*a*b^2)*sin(d*x+c)/a^3/(a^2-b^2)/d/cos(d*x+c)^(1/2)
```

3.376.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

3.376.2 Mathematica [A] (verified)

Time = 4.62 (sec) , antiderivative size = 363, normalized size of antiderivative = 1.05

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$\frac{2(4a^4A + 44a^2Ab^2 - 45Ab^4 - 30a^3bB + 27ab^3B) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right) - 8a(-7a^2Ab + 10Ab^3 + 3a^3B - 6ab^2B) \left(\frac{(a+b) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - a \operatorname{EllipticPi}\left(\frac{2}{a+b}\right)}{b(a+b)}\right)}{a+b}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2),x]`

output `((2*(4*a^4*A + 44*a^2*A*b^2 - 45*A*b^4 - 30*a^3*b*B + 27*a*b^3*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) - (8*a*(-7*a^2*A*b + 10*A*b^3 + 3*a^3*B - 6*a*b^2*B))*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(b*(a + b)) - (6*(-4*a^2*A*b + 5*A*b^3 + 2*a^3*B - 3*a*b^2*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b*Sqrt[Sin[c + d*x]^2))/((a - b)*(a + b)) + 4*Sqrt[Cos[c + d*x]]*((3*b^3*(A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*(a + b*Cos[c + d*x])) + 2*(-6*A*b + 3*a*B + a*A*Sec[c + d*x])*Tan[c + d*x]))/(12*a^3*d)`

3.376.3 Rubi [A] (verified)

Time = 2.59 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.95, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^2} dx$$

3.376. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{2Aa^2+3bBa-2(Ab-aB)\cos(c+dx)a-5Ab^2+3b(Ab-aB)\cos^2(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))} dx \\
& \frac{a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} + \\
& \frac{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} \\
& \downarrow 3479 \\
& \int \frac{2Aa^2+3bBa-2(Ab-aB)\cos(c+dx)a-5Ab^2+3b(Ab-aB)\cos^2(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))} dx \\
& \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} + \\
& \frac{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} \\
& \downarrow 27 \\
& \int \frac{2Aa^2+3bBa-2(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-5Ab^2+3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{5/2}(a+b\sin(c+dx+\frac{\pi}{2}))} dx \\
& \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} + \\
& \frac{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} \\
& \downarrow 3042 \\
& 2 \int -\frac{-b(2Aa^2+3bBa-5Ab^2)\cos^2(c+dx)-2a(Aa^2-3bBa+2Ab^2)\cos(c+dx)+3(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3)}{2\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} dx \\
& \frac{2(2a^2A+3abB-5Ab^2)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} + \\
& \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} + \\
& \frac{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} \\
& \downarrow 27 \\
& \frac{2(2a^2A+3abB-5Ab^2)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \int \frac{-b(2Aa^2+3bBa-5Ab^2)\cos^2(c+dx)-2a(Aa^2-3bBa+2Ab^2)\cos(c+dx)+3(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} dx \\
& \frac{2a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)} + \\
& \frac{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} \\
& \downarrow 3042
\end{aligned}$$

3.376. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^2} dx$

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-b(2Aa^2+3bBa-5Ab^2) \sin(c+dx+\frac{\pi}{2})^2 - 2a(Aa^2-3bBa+2Ab^2) \sin(c+dx+\frac{\pi}{2}) + 3(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3)}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3a}$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB) \sin(c+dx)} \\ \frac{b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}$$

↓ 3534

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2 \int \frac{2Aa^4-12bBa^3+16Ab^2a^2+9b^3Ba+2(-3Ba^3+7Aba^2+6b^2Ba-10Ab^3) \cos(c+dx)a-15Ab^4+3b(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3)}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} \quad 3a$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB) \sin(c+dx)} \\ \frac{b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}$$

↓ 27

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{2Aa^4-12bBa^3+16Ab^2a^2+9b^3Ba+2(-3Ba^3+7Aba^2+6b^2Ba-10Ab^3)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} \quad 3a$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB) \sin(c+dx)} \\ \frac{b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}$$

↓ 3042

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{2Aa^4-12bBa^3+16Ab^2a^2+9b^3Ba+2(-3Ba^3+7Aba^2+6b^2Ba-10Ab^3)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a} \quad 3a$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB) \sin(c+dx)} \\ \frac{b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}$$

↓ 3538

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \sqrt{\cos(c+dx)} dx}{a} - \frac{\int \frac{a(2Aa^2+3bBa-5Ab^2) \sin(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx}{a} \quad 3a$$

$$\frac{2a(a^2-b^2)}{b(Ab-aB) \sin(c+dx)} \\ \frac{b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}$$

3.376. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

↓ 25

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \sqrt{\cos(c+dx)} dx + \int \frac{a(2Aa^2+3bBa-5Ab^2)}{a} dx}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} \quad 2a(a^2 - b^2)$$

↓ 3042

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + \int \frac{a(2Aa^2+3bBa-5Ab^2)}{a} dx}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} \quad 2a(a^2 - b^2)$$

↓ 3119

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{a(2Aa^2+3bBa-5Ab^2) \sin(c+dx+\frac{\pi}{2})b^2 + (2Aa^4-12bBa^3+16Ab^2a^2+a^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{a(2Aa^2+3bBa-5Ab^2)}{a} dx}{b}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} \quad 2a(a^2 - b^2)$$

↓ 3481

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{ab(2a^2A+3abB-5Ab^2) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3b^2(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \int \frac{a(2Aa^2+3bBa-5Ab^2)}{a} dx}{b}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} \quad 2a(a^2 - b^2)$$

↓ 3042

3.376. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{ab(2a^2A+3abB-5Ab^2) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3b^2(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} \frac{a+b \sin(c+dx)}{b} dx}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}$$

↓ 3120

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3b^2(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} \frac{a+b \sin(c+dx)}{b} dx}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))}$$

↓ 3284

$$\frac{b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} + \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2ab(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3a}$$

$$\frac{2(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2ab(2a^2A+3abB-5Ab^2) \sin(c+dx)}{3a}$$

$$2a(a^2 - b^2)$$

```
input Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2),x]
```

```
output (b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])) + ((2*(2*a^2*A - 5*A*b^2 + 3*a*b*B)*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2))) - (((6*(4*a^2*A*b - 5*A*b^3 - 2*a^3*B + 3*a*b^2*B)*EllipticE[(c + d*x)/2, 2])/d + ((2*a*b*(2*a^2*A - 5*A*b^2 + 3*a*b*B)*EllipticF[(c + d*x)/2, 2])/d + (6*b^2*(7*a^2*A*b - 5*A*b^3 - 5*a^3*B + 3*a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d))/b)/a + (6*(4*a^2*A*b - 5*A*b^3 - 2*a^3*B + 3*a*b^2*B)*Sin[c + d*x])/(a*d*sqrt[Cos[c + d*x]])))/(3*a))/(2*a*(a^2 - b^2))
```

3.376. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

3.376.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.376.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1003 vs. $2(413) = 826$.

Time = 11.49 (sec) , antiderivative size = 1004, normalized size of antiderivative = 2.91

method	result	size
default	Expression too large to display	1004

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

$$3.376. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2/a^2*A*(-1/6*
cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(c
os(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*
x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*E
llipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2*(-2*A*b+B*a)/a^3/sin(1/2*d*x+1/2*c
)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2)))-4*b^2*(2*A*b-B*a)/a^3/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+2*(A*b-B*a)*b
/a^2*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1
/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2
*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/
2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(
1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/
2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*c
os(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b...

```

3.376.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output `Timed out`

3.376.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c))**2,x)`output `Timed out`**3.376.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2)), x)`**3.376.8 Giac [F]**

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2)), x)`

3.376.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{5}{2}}(a + b \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^2),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^2), x)`

3.377
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

3.377.1 Optimal result 3511
 3.377.2 Mathematica [A] (verified) 3512
 3.377.3 Rubi [A] (verified) 3513
 3.377.4 Maple [B] (verified) 3518
 3.377.5 Fricas [F(-1)] 3518
 3.377.6 Sympy [F(-1)] 3519
 3.377.7 Maxima [F] 3519
 3.377.8 Giac [F] 3519
 3.377.9 Mupad [F(-1)] 3520

3.377.1 Optimal result

Integrand size = 33, antiderivative size = 367

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

$$= -\frac{(3a^3Ab - 9aAb^3 - 15a^4B + 29a^2b^2B - 8b^4B) E(\frac{1}{2}(c+dx)|2)}{4b^3(a^2 - b^2)^2 d}$$

$$+ \frac{(3a^4Ab - 5a^2Ab^3 + 8Ab^5 - 15a^5B + 33a^3b^2B - 24ab^4B) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{4b^4(a^2 - b^2)^2 d}$$

$$- \frac{a(3a^4Ab - 6a^2Ab^3 + 15Ab^5 - 15a^5B + 38a^3b^2B - 35ab^4B) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{4(a-b)^2b^4(a+b)^3d}$$

$$+ \frac{a(Ab - aB) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2b(a^2 - b^2) d(a+b \cos(c+dx))^2}$$

$$+ \frac{a(a^2Ab - 7Ab^3 - 5a^3B + 11ab^2B) \sqrt{\cos(c+dx)} \sin(c+dx)}{4b^2(a^2 - b^2)^2 d(a+b \cos(c+dx))}$$

output
$$-1/4*(3*A*a^3*b-9*A*a*b^3-15*B*a^4+29*B*a^2*b^2-8*B*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)})/b^3/(a^2-b^2)^2/d+1/4*(3*A*a^4*b-5*A*a^2*b^3+8*A*b^5-15*B*a^5+33*B*a^3*b^2-24*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2)})/b^4/(a^2-b^2)^2/d-1/4*a*(3*A*a^4*b-6*A*a^2*b^3+15*A*b^5-15*B*a^5+38*B*a^3*b^2-35*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^{(1/2)})/(a-b)^2/b^4/(a+b)^3/d+1/2*a*(A*b-B*a)*cos(d*x+c)^{(3/2)}*sin(d*x+c)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/4*a*(A*a^2*b-7*A*b^3-5*B*a^3+11*B*a*b^2)*sin(d*x+c)*cos(d*x+c)^{(1/2)}/b^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))$$

3.377.2 Mathematica [A] (verified)

Time = 3.43 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.06

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \frac{2a\sqrt{\cos(c+dx)}(a(-a^2Ab+7Ab^3+5a^3B-11ab^2B)+b(-3a^2Ab+9Ab^3+7a^3B-13ab^2B)\cos(c+dx))\sin(c+dx)}{(a^2-b^2)^2(a+b\cos(c+dx))^2} + \frac{(-a^3Ab-5aAb^3+5a^4B-7a^2b^2B)}{(a^2-b^2)^2(a+b\cos(c+dx))^2}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3, x]`

output
$$((-2*a*sqrt[Cos[c + d*x]]*(a*(-a^2*A*b) + 7*A*b^3 + 5*a^3*B - 11*a*b^2*B) + b*(-3*a^2*A*b + 9*A*b^3 + 7*a^3*B - 13*a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/((a^2 - b^2)^2*(a + b*Cos[c + d*x])^2) + (((-a^3*A*b) - 5*a*A*b^3 + 5*a^4*B - 7*a^2*b^2*B + 8*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (8*(a^2*A*b + 2*A*b^3 + a^3*B - 4*a*b^2*B)*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b) + ((-3*a^3*A*b + 9*a*A*b^3 + 15*a^4*B - 29*a^2*b^2*B + 8*b^4*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*sqrt[Sin[c + d*x]^2])/((a - b)^2*(a + b)^2)/(8*b^2*d)$$

3.377.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

3.377.3 Rubi [A] (verified)

Time = 2.17 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx$$

$$\downarrow \text{3468}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} - \int \frac{\sqrt{\cos(c+dx)}(-((-5Ba^2+Aba+4b^2B)\cos^2(c+dx)-4b(Ab-aB)\cos(c+dx)+3a(Ab-aB)))}{2(a+b\cos(c+dx))^2} dx}{2b(a^2-b^2)}$$

$$\downarrow \text{27}$$

$$\int \frac{\sqrt{\cos(c+dx)}(-((-5Ba^2+Aba+4b^2B)\cos^2(c+dx)-4b(Ab-aB)\cos(c+dx)+3a(Ab-aB)))}{(a+b\cos(c+dx))^2} dx}{4b(a^2-b^2)} + \frac{a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}((5Ba^2-Aba-4b^2B)\sin(c+dx+\frac{\pi}{2})^2-4b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+3a(Ab-aB))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx}{4b(a^2-b^2)} + \frac{a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

$$\downarrow \text{3526}$$

$$\frac{a(-5a^3B+a^2Ab+11ab^2B-7Ab^3)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b\cos(c+dx))} - \int \frac{-((-15Ba^4+3Aba^3+29b^2Ba^2-9Ab^3a-8b^4B)\cos^2(c+dx)+4b(Ba^3+Aba^2-4b^2Ba+2b^3))\sqrt{\cos(c+dx)}}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))b(a^2-b^2)} dx}{4b(a^2-b^2)}$$

$$\frac{a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2bd(a^2-b^2)(a+b\cos(c+dx))^2}$$

3.377. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

↓ 27

$$\frac{\int \frac{-((-15Ba^4 + 3Aba^3 + 29b^2Ba^2 - 9Ab^3a - 8b^4B) \cos^2(c+dx) + 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \cos(c+dx) + a(-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3))}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{2b(a^2-b^2)} + \frac{a(-5a^3B}{4b(a^2-b^2)}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\frac{\int \frac{(15Ba^4 - 3Aba^3 - 29b^2Ba^2 + 9Ab^3a + 8b^4B) \sin(c+dx + \frac{\pi}{2})^2 + 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \sin(c+dx + \frac{\pi}{2}) + a(-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3)}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \sin(c+dx + \frac{\pi}{2}))} dx}{2b(a^2-b^2)} + \frac{a(-5a^3B}{4b(a^2-b^2)}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3538

$$\frac{-\frac{(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{b} \int \sqrt{\cos(c+dx)} dx}{2b(a^2-b^2)} - \frac{\int \frac{ab(-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) + (-15Ba^5 + 3Aba^4 + 33b^2Ba^3 - 5Ab^3a^2 - 24b^4Ba + 8Ab^5)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b}}{4b(a^2-b^2)}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 25

$$\frac{\int \frac{ab(-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) + (-15Ba^5 + 3Aba^4 + 33b^2Ba^3 - 5Ab^3a^2 - 24b^4Ba + 8Ab^5) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{2b(a^2-b^2)} - \frac{(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \int \sqrt{\cos(c+dx)} dx}{b}}{4b(a^2-b^2)}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3042

$$\frac{\int \frac{ab(-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) + (-15Ba^5 + 3Aba^4 + 33b^2Ba^3 - 5Ab^3a^2 - 24b^4Ba + 8Ab^5) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \sin(c+dx + \frac{\pi}{2}))} dx}{2b(a^2-b^2)} - \frac{(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \int \sqrt{\cos(c+dx)} dx}{b}}{4b(a^2-b^2)}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3119

3.377. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

$$\frac{\int \frac{ab(-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) + (-15Ba^5 + 3Aba^4 + 33b^2Ba^3 - 5Ab^3a^2 - 24b^4Ba + 8Ab^5) \sin(c+dx + \frac{\pi}{2}) dx}{\sqrt{\sin(c+dx + \frac{\pi}{2})} (a+b \sin(c+dx + \frac{\pi}{2}))}}{\frac{b}{2b(a^2-b^2)} - \frac{2(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{bd}}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3481

$$\frac{\frac{(-15a^5B + 3a^4Ab + 33a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))}}{b}}{\frac{b}{2b(a^2-b^2)} - \frac{2(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{bd}}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3042

$$\frac{\frac{(-15a^5B + 3a^4Ab + 33a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx}{b} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \cos(c+dx))}}{b}}{\frac{b}{2b(a^2-b^2)} - \frac{2(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{bd}}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3120

$$\frac{\frac{2(-15a^5B + 3a^4Ab + 33a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{bd} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \cos(c+dx))}}{b}}{\frac{b}{2b(a^2-b^2)} - \frac{2(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{bd}}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3284

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{2(-15a^5B + 3a^4Ab + 33a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5) \text{EllipticF}(\frac{1}{2}(c+dx), 2) - 2a(-15a^5B + 3a^4Ab + 33a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5) \sin(c+dx) \sqrt{\cos(c+dx)}}{bd(a^2-b^2)(a+b \cos(c+dx))} + \frac{2a(-15a^5B + 3a^4Ab + 33a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5)}{b}$$

$$\frac{a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2}$$

```
input Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]
```

3.377. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

```
output (a*(A*b - a*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*(a + b*
Cos[c + d*x])^2) + (((-2*(3*a^3*A*b - 9*a*A*b^3 - 15*a^4*B + 29*a^2*b^2*B
- 8*b^4*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((2*(3*a^4*A*b - 5*a^2*A*b^3
+ 8*A*b^5 - 15*a^5*B + 33*a^3*b^2*B - 24*a*b^4*B)*EllipticF[(c + d*x)/2,
2])/(b*d) - (2*a*(3*a^4*A*b - 6*a^2*A*b^3 + 15*A*b^5 - 15*a^5*B + 38*a^3*b
^2*B - 35*a*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d
))/b)/(2*b*(a^2 - b^2)) + (a*(a^2*A*b - 7*A*b^3 - 5*a^3*B + 11*a*b^2*B)*Sq
rt[Cos[c + d*x]*Sin[c + d*x])/(b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])))/(4*
b*(a^2 - b^2))
```

3.377.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m - 1)*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Ssin[e + f*x])^(m - 2)*(c + d*Ssin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Ssin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Ssin[e + f*x])^m/(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m)*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Ssin[e + f*x])^(m - 1)*(c + d*Ssin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3538 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Ssin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Ssin[e + f*x]]*(c + d*Ssin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.377.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1976 vs. $2(431) = 862$.

Time = 83.35 (sec) , antiderivative size = 1977, normalized size of antiderivative = 5.39

method	result	size
default	Expression too large to display	1977

```
input int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^(1/2)*(2/b^4/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(A*b*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)
)-3*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a-B*b*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2)))+2*a^2/b^4*(3*A*b-4*B*a)*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2
*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+
1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2
*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipt
icF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*
x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(
1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(
1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a
*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/
2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2
*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a^3*(A*b-B*a)/b^4*(-1/2/a*b^2/(a^2-b^2)
*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)...
```

3.377.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x,algorith
m="fricas")
```

3.377. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

output Timed out

3.377.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)`

output Timed out

3.377.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^3, x)`

3.377.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^3, x)`

3.377. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

3.377.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{\cos(c+dx)^{5/2}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3, x)`output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3, x)`

$$3.378 \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

3.378.1 Optimal result	3521
3.378.2 Mathematica [A] (verified)	3522
3.378.3 Rubi [A] (verified)	3522
3.378.4 Maple [B] (verified)	3527
3.378.5 Fricas [F(-1)]	3528
3.378.6 Sympy [F(-1)]	3529
3.378.7 Maxima [F]	3529
3.378.8 Giac [F]	3529
3.378.9 Mupad [F(-1)]	3530

3.378.1 Optimal result

Integrand size = 33, antiderivative size = 344

$$\begin{aligned} & \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx \\ &= -\frac{(a^2Ab + 5Ab^3 + 3a^3B - 9ab^2B) E\left(\frac{1}{2}(c+dx) \mid 2\right)}{4b^2(a^2 - b^2)^2 d} \\ & \quad + \frac{(a^3Ab - 7aAb^3 + 3a^4B - 5a^2b^2B + 8b^4B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{4b^3(a^2 - b^2)^2 d} \\ & \quad - \frac{(a^4Ab - 10a^2Ab^3 - 3Ab^5 + 3a^5B - 6a^3b^2B + 15ab^4B) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{4(a-b)^2b^3(a+b)^3d} \\ & \quad + \frac{a(Ab - aB) \sqrt{\cos(c+dx)} \sin(c+dx)}{2b(a^2 - b^2) d(a+b \cos(c+dx))^2} \\ & \quad + \frac{(a^2Ab + 5Ab^3 + 3a^3B - 9ab^2B) \sqrt{\cos(c+dx)} \sin(c+dx)}{4b(a^2 - b^2)^2 d(a+b \cos(c+dx))} \end{aligned}$$

output

```
-1/4*(A*a^2*b+5*A*b^3+3*B*a^3-9*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(
1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/(a^2-b^2)^2/d+1/4
*(A*a^3*b-7*A*a*b^3+3*B*a^4-5*B*a^2*b^2+8*B*b^4)*(cos(1/2*d*x+1/2*c)^2)^(1
/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b^3/(a^2-b^2)
^2/d-1/4*(A*a^4*b-10*A*a^2*b^3-3*A*b^5+3*B*a^5-6*B*a^3*b^2+15*B*a*b^4)*(co
s(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c)
,2*b/(a+b),2^(1/2))/(a-b)^2/b^3/(a+b)^3/d+1/2*a*(A*b-B*a)*sin(d*x+c)*cos(d
*x+c)^(1/2)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/4*(A*a^2*b+5*A*b^3+3*B*a^3-
9*B*a*b^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/b/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

$$3.378. \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

3.378.2 Mathematica [A] (verified)

Time = 2.54 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.05

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \frac{2\sqrt{\cos(c+dx)}(a(3a^2Ab+3Ab^3+a^3B-7ab^2B)+b(a^2Ab+5Ab^3+3a^3B-9ab^2B)\cos(c+dx))\sin(c+dx)}{(a^2-b^2)^2(a+b\cos(c+dx))^2} - \frac{(-5a^2Ab-Ab^3+a^3B+5ab^2B)\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{c+dx}{2}, 2\right)}{a+b}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3, x]`

output `((2*sqrt(Cos[c + d*x]))*(a*(3*a^2*A*b + 3*A*b^3 + a^3*B - 7*a*b^2*B) + b*(a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/((a^2 - b^2)^2*(a + b*Cos[c + d*x])^2) - (((-5*a^2*A*b - A*b^3 + a^3*B + 5*a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) - (8*(-3*a*A*b + a^2*B + 2*b^2*B)*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b) + ((a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*sqrt(Sin[c + d*x]^2)))/((a - b)^2*(a + b)^2)/(8*b*d)`

3.378.3 Rubi [A] (verified)Time = 2.06 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3468, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

↓ 3042

$$\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}(A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{(a+b\sin\left(c+dx+\frac{\pi}{2}\right))^3} dx$$

3.378. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

$$\begin{aligned}
& \downarrow 3468 \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} - \\
& \frac{\int -\frac{(3Ba^2 + Aba - 4b^2B) \cos^2(c + dx) - 4b(Ab - aB) \cos(c + dx) + a(Ab - aB)}{2\sqrt{\cos(c + dx)(a + b \cos(c + dx))^2}} dx}{2b(a^2 - b^2)} \\
& \downarrow 27 \\
& \frac{\int \frac{(3Ba^2 + Aba - 4b^2B) \cos^2(c + dx) - 4b(Ab - aB) \cos(c + dx) + a(Ab - aB)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))^2}} dx}{4b(a^2 - b^2)} + \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{(3Ba^2 + Aba - 4b^2B) \sin(c + dx + \frac{\pi}{2})^2 - 4b(Ab - aB) \sin(c + dx + \frac{\pi}{2}) + a(Ab - aB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})(a + b \sin(c + dx + \frac{\pi}{2}))^2}} dx}{4b(a^2 - b^2)} + \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
& \downarrow 3534 \\
& \frac{\int \frac{-a(3Ba^3 + Aba^2 - 9b^2Ba + 5Ab^3) \cos^2(c + dx) - 4ab(-Ba^2 + 3Aba - 2b^2B) \cos(c + dx) + a(Ba^3 + 3Aba^2 - 7b^2Ba + 3Ab^3)}{2\sqrt{\cos(c + dx)(a + b \cos(c + dx))}} dx}{a(a^2 - b^2)} + \frac{(3a^3B + a^2Ab - 9ab^2B + 5Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \frac{4b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
& \downarrow 27 \\
& \frac{\int \frac{-a(3Ba^3 + Aba^2 - 9b^2Ba + 5Ab^3) \cos^2(c + dx) - 4ab(-Ba^2 + 3Aba - 2b^2B) \cos(c + dx) + a(Ba^3 + 3Aba^2 - 7b^2Ba + 3Ab^3)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))}} dx}{2a(a^2 - b^2)} + \frac{(3a^3B + a^2Ab - 9ab^2B + 5Ab^3) \sin(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \frac{4b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{-a(3Ba^3 + Aba^2 - 9b^2Ba + 5Ab^3) \sin(c + dx + \frac{\pi}{2})^2 - 4ab(-Ba^2 + 3Aba - 2b^2B) \sin(c + dx + \frac{\pi}{2}) + a(Ba^3 + 3Aba^2 - 7b^2Ba + 3Ab^3)}{\sqrt{\sin(c + dx + \frac{\pi}{2})(a + b \sin(c + dx + \frac{\pi}{2}))}} dx}{2a(a^2 - b^2)} + \frac{(3a^3B + a^2Ab - 9ab^2B + 5Ab^3) \cos(c + dx)}{d(a^2 - b^2)(a + b \cos(c + dx))} \\
& \frac{4b(a^2 - b^2)}{2bd(a^2 - b^2)(a + b \cos(c + dx))^2} \\
& \downarrow 3538
\end{aligned}$$

3.378. $\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^3} dx$

$$\frac{\frac{a(3a^3B+a^2Ab-9ab^2B+5Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} - \frac{\int -\frac{ab(Ba^3+3Aba^2-7b^2Ba+3Ab^3)+a(3Ba^4+Aba^3-5b^2Ba^2-7Ab^3a+8b^4B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{2a(a^2-b^2)}}{4b(a^2-b^2)} + (3a^3B+a^2Ab)$$

$$\frac{a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 25

$$\frac{\frac{\int \frac{ab(Ba^3+3Aba^2-7b^2Ba+3Ab^3)+a(3Ba^4+Aba^3-5b^2Ba^2-7Ab^3a+8b^4B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} - \frac{a(3a^3B+a^2Ab-9ab^2B+5Ab^3) \int \sqrt{\cos(c+dx)} dx}{b}}{2a(a^2-b^2)}}{4b(a^2-b^2)} + (3a^3B+a^2Ab)$$

$$\frac{a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3042

$$\frac{\frac{\int \frac{ab(Ba^3+3Aba^2-7b^2Ba+3Ab^3)+a(3Ba^4+Aba^3-5b^2Ba^2-7Ab^3a+8b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} - \frac{a(3a^3B+a^2Ab-9ab^2B+5Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b}}{2a(a^2-b^2)}}{4b(a^2-b^2)} + (3a^3B+a^2Ab)$$

$$\frac{a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3119

$$\frac{\frac{\int \frac{ab(Ba^3+3Aba^2-7b^2Ba+3Ab^3)+a(3Ba^4+Aba^3-5b^2Ba^2-7Ab^3a+8b^4B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} - \frac{2a(3a^3B+a^2Ab-9ab^2B+5Ab^3) E(\frac{1}{2}(c+dx)|2)}{bd}}{2a(a^2-b^2)}}{4b(a^2-b^2)} + (3a^3B+a^2Ab)$$

$$\frac{a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3481

$$\frac{\frac{a(3a^4B+a^3Ab-5a^2b^2B-7aAb^3+8b^4B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a(3a^5B+a^4Ab-6a^3b^2B-10a^2Ab^3+15ab^4B-3Ab^5) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b}}{2a(a^2-b^2)}}{4b(a^2-b^2)} + 2a(3a^3B+a^2Ab)$$

$$\frac{a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b \cos(c+dx))^2}$$

↓ 3042

3.378. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{a(3a^4B+a^3Ab-5a^2b^2B-7aAb^3+8b^4B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{a(3a^5B+a^4Ab-6a^3b^2B-10a^2Ab^3+15ab^4B-3Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} \\
 & \frac{2a(a^2-b^2)}{4b(a^2-b^2)} \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2a(3a^4B+a^3Ab-5a^2b^2B-7aAb^3+8b^4B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{a(3a^5B+a^4Ab-6a^3b^2B-10a^2Ab^3+15ab^4B-3Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} \\
 & \frac{2a(a^2-b^2)}{4b(a^2-b^2)} \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} \\
 & \quad \downarrow \text{3284} \\
 & \frac{a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2bd(a^2-b^2)(a+b\cos(c+dx))^2} + \\
 & \frac{(3a^3B+a^2Ab-9ab^2B+5Ab^3)\sin(c+dx)\sqrt{\cos(c+dx)}}{d(a^2-b^2)(a+b\cos(c+dx))} + \frac{2a(3a^4B+a^3Ab-5a^2b^2B-7aAb^3+8b^4B) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{2a(3a^5B+a^4Ab-6a^3b^2B-10a^2Ab^3+15ab^4B-3Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} \\
 & \frac{2a(a^2-b^2)}{4b(a^2-b^2)}
 \end{aligned}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]`

output `(a*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + (((-2*a*(a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((2*a*(a^3*A*b - 7*a*A*b^3 + 3*a^4*B - 5*a^2*b^2*B + 8*b^4*B)*EllipticF[(c + d*x)/2, 2])/(b*d) - (2*a*(a^4*A*b - 10*a^2*A*b^3 - 3*A*b^5 + 3*a^5*B - 6*a^3*b^2*B + 15*a*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))/b)/(2*a*(a^2 - b^2)) + ((a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(4*b*(a^2 - b^2))`

3.378. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

3.378.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.378.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

```
rule 3481 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.378.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1936 vs. $2(408) = 816$.

Time = 11.48 (sec) , antiderivative size = 1937, normalized size of antiderivative = 5.63

method	result	size
default	Expression too large to display	1937

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

$$3.378. \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

output
$$\begin{aligned} & -(-(-2\cos(1/2dx+1/2c)^2+1)\sin(1/2dx+1/2c)^2)^{1/2}*(2B/b^3*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})-4/b^2*(A*b-3*B*a)/(-2*a*b+2*b^2)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2dx+1/2c),-2*b/(a-b),2^{1/2})+2*a^2*(A*b-B*a)/b^3*(-1/2/a*b^2/(a^2-b^2)*\cos(1/2dx+1/2c)*(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}/(2*b*\cos(1/2dx+1/2c)^2+a-b)^2-3/4*b^2*(3*a^2-b^2)/a^2/(a^2-b^2)^2*\cos(1/2dx+1/2c)*(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}/(2*b*\cos(1/2dx+1/2c)^2+a-b)-7/8/(a+b)/(a^2-b^2)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})+1/4/(a+b)/(a^2-b^2)/a*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})*b+3/8/(a+b)/(a^2-b^2)/a^2*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})*b^2-9/8*b/(a^2-b^2)^2*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2}))+3/8*b^3/a^2/(a^2-b^2)^2*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2... \end{aligned}$$

3.378.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\cos^{3/2}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(dx+c)^(3/2)*(A+B*cos(dx+c))/(a+b*cos(dx+c))^3,x, algorithm="fracas")`

output `Timed out`

3.378.
$$\int \frac{\cos^{3/2}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

3.378.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)`output `Timed out`**3.378.7 Maxima [F]**

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^3, x)`**3.378.8 Giac [F]**

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="giac")`output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^3, x)`

3.378.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3, x)`output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3, x)`

3.379
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

3.379.1 Optimal result 3531
 3.379.2 Mathematica [A] (verified) 3532
 3.379.3 Rubi [A] (verified) 3532
 3.379.4 Maple [B] (verified) 3537
 3.379.5 Fracas [F(-1)] 3538
 3.379.6 Sympy [F(-1)] 3539
 3.379.7 Maxima [F] 3539
 3.379.8 Giac [F] 3539
 3.379.9 Mupad [F(-1)] 3540

3.379.1 Optimal result

Integrand size = 33, antiderivative size = 337

$$\begin{aligned} & \int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx \\ &= \frac{(5a^2Ab + Ab^3 - a^3B - 5ab^2B) E\left(\frac{1}{2}(c+dx) \mid 2\right)}{4ab(a^2 - b^2)^2 d} \\ &+ \frac{(3a^2Ab + 3Ab^3 + a^3B - 7ab^2B) \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{4b^2(a^2 - b^2)^2 d} \\ &- \frac{(3a^4Ab + 10a^2Ab^3 - Ab^5 + a^5B - 10a^3b^2B - 3ab^4B) \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{4a(a-b)^2b^2(a+b)^3d} \\ &- \frac{(Ab - aB)\sqrt{\cos(c+dx)} \sin(c+dx)}{2(a^2 - b^2) d(a+b \cos(c+dx))^2} \\ &- \frac{(5a^2Ab + Ab^3 - a^3B - 5ab^2B) \sqrt{\cos(c+dx)} \sin(c+dx)}{4a(a^2 - b^2)^2 d(a+b \cos(c+dx))} \end{aligned}$$

output

```
1/4*(5*A*a^2*b+A*b^3-B*a^3-5*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2
*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a/b/(a^2-b^2)^2/d+1/4*(3
*A*a^2*b+3*A*b^3+B*a^3-7*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x
+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/(a^2-b^2)^2/d-1/4*(3*A*a
^4*b+10*A*a^2*b^3-A*b^5+B*a^5-10*B*a^3*b^2-3*B*a*b^4)*(cos(1/2*d*x+1/2*c)^
2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2
))/a/(a-b)^2/b^2/(a+b)^3/d-1/2*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/(a^2-
b^2)/d/(a+b*cos(d*x+c))^2-1/4*(5*A*a^2*b+A*b^3-B*a^3-5*B*a*b^2)*sin(d*x+c)
*cos(d*x+c)^(1/2)/a/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

3.379.
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$$

3.379.2 Mathematica [A] (verified)

Time = 3.12 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

$$= \frac{4\sqrt{\cos(c+dx)}(a(-7a^2Ab+Ab^3+3a^3B+3ab^2B)+b(-5a^2Ab-Ab^3+a^3B+5ab^2B)\cos(c+dx))\sin(c+dx)}{(a^2-b^2)^2(a+b\cos(c+dx))^2} + \frac{2(-9a^2Ab+3Ab^3+5a^3B+ab^2B)\operatorname{EllipticE}(\arcsin(\frac{b}{a+b}\sqrt{\cos(c+dx)}))}{a+b}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3, x]`

output `((4*Sqrt[Cos[c + d*x]]*(a*(-7*a^2*A*b + A*b^3 + 3*a^3*B + 3*a*b^2*B) + b*(-5*a^2*A*b - A*b^3 + a^3*B + 5*a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/((a^2 - b^2)^2*(a + b*Cos[c + d*x])^2) + ((2*(-9*a^2*A*b + 3*A*b^3 + 5*a^3*B + a*b^2*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (16*a*(2*a^2*A + A*b^2 - 3*a*b*B)*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(b*(a + b)) - (2*(-5*a^2*A*b - A*b^3 + a^3*B + 5*a*b^2*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*Sqrt[Sin[c + d*x]^2]))/((a - b)^2*(a + b)^2))/(16*a*d)`

3.379.3 Rubi [A] (verified)Time = 2.01 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.99, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3478, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

↓ 3042

$$\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx$$

3.379. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$

$$\begin{aligned}
 & \int \frac{(Ab-aB) \cos^2(c+dx) - 4(aA-bB) \cos(c+dx) + Ab-aB}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} dx \quad \downarrow \text{3478} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \int \frac{(Ab-aB) \cos^2(c+dx) - 4(aA-bB) \cos(c+dx) + Ab-aB}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} dx \quad \downarrow \text{27} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \int \frac{(Ab-aB) \sin(c+dx+\frac{\pi}{2})^2 - 4(aA-bB) \sin(c+dx+\frac{\pi}{2}) + Ab-aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))^2} dx \quad \downarrow \text{3042} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \int \frac{-3Ba^3+7Aba^2-3b^2Ba-4(2Aa^2-3bBa+Ab^2) \cos(c+dx)a-Ab^3 - (-Ba^3+5Aba^2-5b^2Ba+Ab^3) \cos^2(c+dx)}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} \quad \downarrow \text{3534} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \int \frac{-3Ba^3+7Aba^2-3b^2Ba-4(2Aa^2-3bBa+Ab^2) \cos(c+dx)a-Ab^3 - (-Ba^3+5Aba^2-5b^2Ba+Ab^3) \cos^2(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} \quad \downarrow \text{27} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \int \frac{-3Ba^3+7Aba^2-3b^2Ba-4(2Aa^2-3bBa+Ab^2) \sin(c+dx+\frac{\pi}{2})a-Ab^3 + (Ba^3-5Aba^2+5b^2Ba-Ab^3) \sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} \quad \downarrow \text{3042} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \int \frac{-3Ba^3+7Aba^2-3b^2Ba-4(2Aa^2-3bBa+Ab^2) \sin(c+dx+\frac{\pi}{2})a-Ab^3 + (Ba^3-5Aba^2+5b^2Ba-Ab^3) \sin(c+dx+\frac{\pi}{2})^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \sin(c+dx)}{ad(a^2-b^2)(a+b \cos(c+dx))} \quad \downarrow \text{3538} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2}
 \end{aligned}$$

3.379. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} - \frac{\int -\frac{b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)-a(Ba^3+3Aba^2-7b^2Ba+3Ab^3) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{2a(a^2-b^2)}}{4(a^2-b^2)} + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{\frac{b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)-a(Ba^3+3Aba^2-7b^2Ba+3Ab^3) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{2a(a^2-b^2)} - \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\cos(c+dx)} dx}{b} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)-a(Ba^3+3Aba^2-7b^2Ba+3Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{2a(a^2-b^2)} - \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \quad \downarrow \text{3119} \\
 & \frac{\frac{b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)-a(Ba^3+3Aba^2-7b^2Ba+3Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{2a(a^2-b^2)} - \frac{2(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) E(\frac{1}{2}(c+dx)|2)}{bd} + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \quad \downarrow \text{3481} \\
 & \frac{\frac{(a^5B+3a^4Ab-10a^3b^2B+10a^2Ab^3-3ab^4B-Ab^5) \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} - \frac{a(a^3B+3a^2Ab-7ab^2B+3Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b}}{2a(a^2-b^2)} - \frac{2(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} + \frac{(a^3(-B)+5a^2Ab-5ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} \\
 & \frac{(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a^2-b^2)(a+b \cos(c+dx))^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.379. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

$$\frac{\frac{(a^5 B + 3a^4 Ab - 10a^3 b^2 B + 10a^2 Ab^3 - 3ab^4 B - Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{a(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b}}{2a(a^2 - b^2)} - 2(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3)$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3120

$$\frac{\frac{(a^5 B + 3a^4 Ab - 10a^3 b^2 B + 10a^2 Ab^3 - 3ab^4 B - Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{2a(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{bd}}{2a(a^2 - b^2)} - 2(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3)$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2d(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3284

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2d(a^2 - b^2)(a + b \cos(c + dx))^2} - \frac{2(a^5 B + 3a^4 Ab - 10a^3 b^2 B + 10a^2 Ab^3 - 3ab^4 B - Ab^5) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{bd(a+b)} - \frac{2a(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3)}{b}$$

$$\frac{(a^3(-B) + 5a^2 Ab - 5ab^2 B + Ab^3) \sin(c+dx) \sqrt{\cos(c+dx)}}{ad(a^2 - b^2)(a + b \cos(c+dx))} + \frac{2(a^5 B + 3a^4 Ab - 10a^3 b^2 B + 10a^2 Ab^3 - 3ab^4 B - Ab^5) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{bd(a+b)} - \frac{2a(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3)}{b}$$

$$4(a^2 - b^2)$$

```
input Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^3,x]
```

```
output -1/2*((A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) - (((-2*(5*a^2*A*b + A*b^3 - a^3*B - 5*a*b^2*B)*EllipticE[(c + d*x)/2, 2])/(b*d) + ((-2*a*(3*a^2*A*b + 3*A*b^3 + a^3*B - 7*a*b^2*B)*EllipticF[(c + d*x)/2, 2])/(b*d) + (2*(3*a^4*A*b + 10*a^2*A*b^3 - A*b^5 + a^5*B - 10*a^3*b^2*B - 3*a*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))/b)/(2*a*(a^2 - b^2)) + ((5*a^2*A*b + A*b^3 - a^3*B - 5*a*b^2*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(4*(a^2 - b^2))
```

3.379. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^3} dx$

3.379.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3478 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2)), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.379.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1849 vs. $2(401) = 802$.

Time = 11.05 (sec) , antiderivative size = 1850, normalized size of antiderivative = 5.49

method	result	size
default	Expression too large to display	1850

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

$$3.379. \quad \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B/b/(-2*a*b
+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1
/2*c),-2*b/(a-b),2^(1/2))+2*(A*b-2*B*a)/b^2*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*
x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2
*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*
x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*E
llipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1
/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)
*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*si
n(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c
),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2
*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)
^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/
(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1
)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(co
s(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a*(A*b-B*a)/b^2*(-1/2/a*b^2/(a^2-b
^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2
)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)^2-3/4*b^2*(3*a^2-b^2)/a^2/(a^2-b^2)^2*...

```

3.379.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm="fracas")`

output `Timed out`

3.379.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.379.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)^3, x)`

3.379.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^3} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)^3, x)`

3.379.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3,x)`output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^3, x)`

3.380 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^3} dx$

3.380.1 Optimal result 3541
 3.380.2 Mathematica [A] (verified) 3542
 3.380.3 Rubi [A] (verified) 3542
 3.380.4 Maple [B] (verified) 3547
 3.380.5 Fricas [F(-1)] 3548
 3.380.6 Sympy [F(-1)] 3549
 3.380.7 Maxima [F(-1)] 3549
 3.380.8 Giac [F] 3549
 3.380.9 Mupad [F(-1)] 3550

3.380.1 Optimal result

Integrand size = 33, antiderivative size = 345

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx$$

$$= -\frac{(9a^2 Ab - 3Ab^3 - 5a^3 B - ab^2 B) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{4a^2 (a^2 - b^2)^2 d}$$

$$- \frac{(7a^2 Ab - Ab^3 - 3a^3 B - 3ab^2 B) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{4ab (a^2 - b^2)^2 d}$$

$$+ \frac{(15a^4 Ab - 6a^2 Ab^3 + 3Ab^5 - 3a^5 B - 10a^3 b^2 B + ab^4 B) \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{4a^2 (a - b)^2 b (a + b)^3 d}$$

$$+ \frac{b(Ab - aB) \sqrt{\cos(c + dx)} \sin(c + dx)}{2a (a^2 - b^2) d (a + b \cos(c + dx))^2}$$

$$+ \frac{b(9a^2 Ab - 3Ab^3 - 5a^3 B - ab^2 B) \sqrt{\cos(c + dx)} \sin(c + dx)}{4a^2 (a^2 - b^2)^2 d (a + b \cos(c + dx))}$$

output

```
-1/4*(9*A*a^2*b-3*A*b^3-5*B*a^3-B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(
1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/a^2/(a^2-b^2)^2/d-1/4
*(7*A*a^2*b-A*b^3-3*B*a^3-3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*
d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/a/b/(a^2-b^2)^2/d+1/4*(15
*A*a^4*b-6*A*a^2*b^3+3*A*b^5-3*B*a^5-10*B*a^3*b^2+B*a*b^4)*(cos(1/2*d*x+1/
2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2
^(1/2))/a^2/(a-b)^2/b/(a+b)^3/d+1/2*b*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2
)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^2+1/4*b*(9*A*a^2*b-3*A*b^3-5*B*a^3-B*a*b^
2)*sin(d*x+c)*cos(d*x+c)^(1/2)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))
```

3.380. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^3} dx$

3.380.2 Mathematica [A] (verified)

Time = 3.34 (sec) , antiderivative size = 383, normalized size of antiderivative = 1.11

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx$$

$$= \frac{-\frac{2b\sqrt{\cos(c+dx)}(a(-11a^2Ab+5Ab^3+7a^3B-ab^2B))+b(-9a^2Ab+3Ab^3+5a^3B+ab^2B)\cos(c+dx)\sin(c+dx)}{(a^2-b^2)^2(a+b\cos(c+dx))^2} + \frac{(16a^4A-19a^2Ab^2+9Ab^4-9a^3bB)}{(a^2-b^2)^2(a+b\cos(c+dx))^2}}{(a^2-b^2)^2(a+b\cos(c+dx))^2}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^3),x]`

output `((-2*b*Sqrt[Cos[c + d*x]]*(a*(-11*a^2*A*b + 5*A*b^3 + 7*a^3*B - a*b^2*B) + b*(-9*a^2*A*b + 3*A*b^3 + 5*a^3*B + a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/((a^2 - b^2)^2*(a + b*Cos[c + d*x])^2) + (((16*a^4*A - 19*a^2*A*b^2 + 9*A*b^4 - 9*a^3*b*B + 3*a*b^3*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (8*a*(-4*a^2*A*b + A*b^3 + 2*a^3*B + a*b^2*B)*((a + b)*EllipticF[(c + d*x)/2, 2] - a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(b*(a + b)) + ((-9*a^2*A*b + 3*A*b^3 + 5*a^3*B + a*b^2*B)*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b*Sqrt[Sin[c + d*x]^2]))/((a - b)^2*(a + b)^2)/(8*a^2*d)`

3.380.3 Rubi [A] (verified)

Time = 2.24 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.97, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))^3} dx$$

$$\int \frac{4Aa^2 - bBa - 4(Ab - aB) \cos(c + dx)a - 3Ab^2 + b(Ab - aB) \cos^2(c + dx)}{2\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx \quad \downarrow \text{3479}$$

$$+ \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

$$\int \frac{4Aa^2 - bBa - 4(Ab - aB) \cos(c + dx)a - 3Ab^2 + b(Ab - aB) \cos^2(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx \quad \downarrow \text{27}$$

$$+ \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

$$\int \frac{4Aa^2 - bBa - 4(Ab - aB) \sin(c + dx + \frac{\pi}{2})a - 3Ab^2 + b(Ab - aB) \sin(c + dx + \frac{\pi}{2})^2}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))^2} dx \quad \downarrow \text{3042}$$

$$+ \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

$$\int \frac{8Aa^4 - 7bBa^3 - 5Ab^2a^2 + b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \cos(c + dx)a + 3Ab^4 - b(-5Ba^3 + 9Aba^2 - b^2Ba - 3Ab^3) \cos^2(c + dx)}{2\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx \quad \downarrow \text{3534}$$

$$+ \frac{b(-5a^3B + 9a^2Ab - a^2B)}{ad(a^2 - b^2)}$$

$$+ \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

$$\int \frac{8Aa^4 - 7bBa^3 - 5Ab^2a^2 + b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \cos(c + dx)a + 3Ab^4 - b(-5Ba^3 + 9Aba^2 - b^2Ba - 3Ab^3) \cos^2(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx \quad \downarrow \text{27}$$

$$+ \frac{b(-5a^3B + 9a^2Ab - a^2B)}{ad(a^2 - b^2)}$$

$$+ \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

$$\int \frac{8Aa^4 - 7bBa^3 - 5Ab^2a^2 + b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sin(c + dx + \frac{\pi}{2})a + 3Ab^4 - b(-5Ba^3 + 9Aba^2 - b^2Ba - 3Ab^3) \sin(c + dx + \frac{\pi}{2})^2}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))} dx \quad \downarrow \text{3042}$$

$$+ \frac{b(-5a^3B + 9a^2Ab - a^2B)}{ad(a^2 - b^2)}$$

$$+ \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

$$\int \frac{8Aa^4 - 7bBa^3 - 5Ab^2a^2 + b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sin(c + dx + \frac{\pi}{2})a + 3Ab^4 - b(-5Ba^3 + 9Aba^2 - b^2Ba - 3Ab^3) \sin(c + dx + \frac{\pi}{2})^2}{2a(a^2 - b^2)} dx \quad \downarrow \text{3538}$$

$$+ \frac{b(-5a^3B + 9a^2Ab - a^2B)}{ad(a^2 - b^2)}$$

$$+ \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

3.380. $\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx$

$$\frac{-\left((-5a^3B+9a^2Ab-ab^2B-3Ab^3)\int\sqrt{\cos(c+dx)}dx\right)-\frac{\int-\frac{b(8Aa^4-7bBa^3-5Ab^2a^2+b^3Ba+3Ab^4)-ab(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}dx}{2a(a^2-b^2)}}{4a(a^2-b^2)} + \frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3)}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 25

$$\frac{\int\frac{b(8Aa^4-7bBa^3-5Ab^2a^2+b^3Ba+3Ab^4)-ab(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}dx}{2a(a^2-b^2)} - \frac{(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\int\sqrt{\cos(c+dx)}dx}{2a(a^2-b^2)} + \frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3)}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3042

$$\frac{\int\frac{b(8Aa^4-7bBa^3-5Ab^2a^2+b^3Ba+3Ab^4)-ab(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))}dx}{2a(a^2-b^2)} - \frac{(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\int\sqrt{\sin(c+dx+\frac{\pi}{2})}dx}{2a(a^2-b^2)} + \frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3)}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3119

$$\frac{\int\frac{b(8Aa^4-7bBa^3-5Ab^2a^2+b^3Ba+3Ab^4)-ab(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))}dx}{2a(a^2-b^2)} - \frac{2(-5a^3B+9a^2Ab-ab^2B-3Ab^3)E(\frac{1}{2}(c+dx)|2)}{d} + \frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3)}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3481

$$\frac{(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5)\int\frac{1}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}dx - a(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\int\frac{1}{\sqrt{\cos(c+dx)}}dx}{2a(a^2-b^2)} - \frac{2(-5a^3B+9a^2Ab-ab^2B-3Ab^3)}{d}$$

$$\frac{b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{2ad(a^2-b^2)(a+b\cos(c+dx))^2}$$

↓ 3042

3.380. $\int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^3} dx$

$$\frac{(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx - a(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a(a^2-b^2)} - 2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} + \frac{2(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} - \frac{2a(-3a^3B+7a^2Ab-3ab^2B-Ab^3)}{b}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3120

$$\frac{(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}} dx - \frac{2a(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}}{2a(a^2-b^2)} - 2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} + \frac{2(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} - \frac{2a(-3a^3B+7a^2Ab-3ab^2B-Ab^3)}{b}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2}$$

↓ 3284

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{2ad(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{2(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} - \frac{2a(-3a^3B+7a^2Ab-3ab^2B-Ab^3)}{b}$$

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx)\sqrt{\cos(c+dx)}}{ad(a^2-b^2)(a+b\cos(c+dx))} + \frac{2(-3a^5B+15a^4Ab-10a^3b^2B-6a^2Ab^3+ab^4B+3Ab^5) \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} - \frac{2a(-3a^3B+7a^2Ab-3ab^2B-Ab^3)}{b}$$

$$4a(a^2 - b^2)$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^3),x]`

output `(b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^2) + (((-2*(9*a^2*A*b - 3*A*b^3 - 5*a^3*B - a*b^2*B)*EllipticE[(c + d*x)/2, 2])/d + ((-2*a*(7*a^2*A*b - A*b^3 - 3*a^3*B - 3*a*b^2*B)*EllipticF[(c + d*x)/2, 2])/d + (2*(15*a^4*A*b - 6*a^2*A*b^3 + 3*A*b^5 - 3*a^5*B - 10*a^3*b^2*B + a*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)*d)/b)/(2*a*(a^2 - b^2)) + (b*(9*a^2*A*b - 3*A*b^3 - 5*a^3*B - a*b^2*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(a*(a^2 - b^2)*d*(a + b*Cos[c + d*x]))/(4*a*(a^2 - b^2))`

3.380.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m + 1)*((c + d*SIN[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.380.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1743 vs. $2(409) = 818$.

Time = 10.66 (sec) , antiderivative size = 1744, normalized size of antiderivative = 5.06

method	result	size
default	Expression too large to display	1744

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

$$3.380. \quad \int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^3} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B/b*(-1/a*b^
2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)
^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b
/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),
2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/
2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellip
ticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2
^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*
cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^
2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))+2*(A*b-B*a)/b*
(-1/2/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*
d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)^2-3/4*b^2*(3*a^2-b^2)/a
^2/(a^2-b^2)^2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-7/8/(a+b)/(a^2-b^2)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/...

```

3.380.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^3,x, algorithm="fricas")`

output `Timed out`

3.380.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.380.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output `Timed out`

3.380.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sqrt(cos(d*x + c))), x)`

3.380.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^3),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^3), x)`

3.381
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$$

3.381.1 Optimal result 3551
 3.381.2 Mathematica [A] (verified) 3552
 3.381.3 Rubi [A] (verified) 3553
 3.381.4 Maple [B] (verified) 3559
 3.381.5 Fricas [F(-1)] 3560
 3.381.6 Sympy [F(-1)] 3560
 3.381.7 Maxima [F(-2)] 3560
 3.381.8 Giac [F] 3561
 3.381.9 Mupad [F(-1)] 3561

3.381.1 Optimal result

Integrand size = 33, antiderivative size = 420

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$$

$$= -\frac{(8a^4A - 29a^2Ab^2 + 15Ab^4 + 9a^3bB - 3ab^3B) E\left(\frac{1}{2}(c + dx) \mid 2\right)}{4a^3(a^2 - b^2)^2 d}$$

$$+ \frac{(11a^2Ab - 5Ab^3 - 7a^3B + ab^2B) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{4a^2(a^2 - b^2)^2 d}$$

$$- \frac{(35a^4Ab - 38a^2Ab^3 + 15Ab^5 - 15a^5B + 6a^3b^2B - 3ab^4B) \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{4a^3(a - b)^2(a + b)^3 d}$$

$$+ \frac{(8a^4A - 29a^2Ab^2 + 15Ab^4 + 9a^3bB - 3ab^3B) \sin(c + dx)}{4a^3(a^2 - b^2)^2 d \sqrt{\cos(c + dx)}}$$

$$+ \frac{b(Ab - aB) \sin(c + dx)}{2a(a^2 - b^2) d \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2}$$

$$+ \frac{b(11a^2Ab - 5Ab^3 - 7a^3B + ab^2B) \sin(c + dx)}{4a^2(a^2 - b^2)^2 d \sqrt{\cos(c + dx)}(a + b \cos(c + dx))}$$

output
$$-1/4*(8*A*a^4-29*A*a^2*b^2+15*A*b^4+9*B*a^3*b-3*B*a*b^3)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1/2)})/a^3/(a^2-b^2)^2/d+1/4*(11*A*a^2*b-5*A*b^3-7*B*a^3+B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2)})/a^2/(a^2-b^2)^2/d-1/4*(35*A*a^4*b-38*A*a^2*b^3+15*A*b^5-15*B*a^5+6*B*a^3*b^2-3*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^{(1/2)})/a^3/(a-b)^2/(a+b)^3/d+1/4*(8*A*a^4-29*A*a^2*b^2+15*A*b^4+9*B*a^3*b-3*B*a*b^3)*sin(d*x+c)/a^3/(a^2-b^2)^2/d/cos(d*x+c)^{(1/2)+1/2*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^2/cos(d*x+c)^{(1/2)+1/4*b*(11*A*a^2*b-5*A*b^3-7*B*a^3+B*a*b^2)*sin(d*x+c)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))/cos(d*x+c)^{(1/2)}$$

3.381.2 Mathematica [A] (verified)

Time = 3.81 (sec) , antiderivative size = 458, normalized size of antiderivative = 1.09

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$$

$$\frac{(56a^4Ab - 95a^2Ab^3 + 45Ab^5 - 16a^5B + 19a^3b^2B - 9ab^4B) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right) + 8a(2a^4A - 10a^2Ab^2 + 5Ab^4 + 4a^3bB - ab^3B) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), \frac{b}{a+b}\right)}{a+b}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^3),x]`

output
$$\begin{aligned} & -\left(\left(\left(\left(56*a^4*A*b - 95*a^2*A*b^3 + 45*A*b^5 - 16*a^5*B + 19*a^3*b^2*B - 9*a*b^4*B\right)*\operatorname{EllipticPi}\left[\frac{2*b}{a+b}, \left(\frac{c+d*x}{2}, 2\right)\right]/(a+b) + \left(8*a*(2*a^4*A - 10*a^2*A*b^2 + 5*A*b^4 + 4*a^3*b*B - a*b^3*B)*((a+b)*\operatorname{EllipticF}\left[\left(\frac{c+d*x}{2}, 2\right) - a*\operatorname{EllipticPi}\left[\frac{2*b}{a+b}, \left(\frac{c+d*x}{2}, 2\right)\right]\right)/(b*(a+b)) + \left(8*a^4*A - 29*a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B\right)*(-2*a*b*\operatorname{EllipticE}\left[\operatorname{ArcSin}\left[\operatorname{Sqrt}\left[\operatorname{Cos}\left[c+d*x\right]\right]\right], -1\right] + 2*a*(a+b)*\operatorname{EllipticF}\left[\operatorname{ArcSin}\left[\operatorname{Sqrt}\left[\operatorname{Cos}\left[c+d*x\right]\right]\right], -1\right] + (-2*a^2 + b^2)*\operatorname{EllipticPi}\left[-\frac{b}{a}, \operatorname{ArcSin}\left[\operatorname{Sqrt}\left[\operatorname{Cos}\left[c+d*x\right]\right]\right], -1\right)*\operatorname{Sin}\left[c+d*x\right]\right)/(a*b*\operatorname{Sqrt}\left[\operatorname{Sin}\left[c+d*x\right]^2\right])\right)/(a-b)^2*(a+b)^2) + \left(\operatorname{Sqrt}\left[\operatorname{Cos}\left[c+d*x\right]\right)*(2*a*b*(16*a^4*A - 47*a^2*A*b^2 + 25*A*b^4 + 11*a^3*b*B - 5*a*b^3*B)*\operatorname{Sin}\left[c+d*x\right] + b^2*(8*a^4*A - 29*a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B)*\operatorname{Sin}\left[2*(c+d*x)\right] + 16*A*(a^3 - a*b^2)^2*\operatorname{Tan}\left[c+d*x\right])\right)/(a^2 - b^2)^2*(a + b*\operatorname{Cos}\left[c+d*x\right])^2)\right)/(8*a^3*d) \end{aligned}$$

3.381.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$$

3.381.3 Rubi [A] (verified)

Time = 3.07 (sec) , antiderivative size = 412, normalized size of antiderivative = 0.98, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^3} dx$$

$$\downarrow \text{3479}$$

$$\frac{\int \frac{4Aa^2 + bBa - 4(Ab - aB) \cos(c + dx)a - 5Ab^2 + 3b(Ab - aB) \cos^2(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx}{\frac{2a(a^2 - b^2)}{b(Ab - aB) \sin(c + dx)}} +$$

$$\frac{2ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2}{\downarrow \text{27}}$$

$$\frac{\int \frac{4Aa^2 + bBa - 4(Ab - aB) \cos(c + dx)a - 5Ab^2 + 3b(Ab - aB) \cos^2(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx}{\frac{4a(a^2 - b^2)}{b(Ab - aB) \sin(c + dx)}} +$$

$$\frac{2ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2}{\downarrow \text{3042}}$$

$$\frac{\int \frac{4Aa^2 + bBa - 4(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)a - 5Ab^2 + 3b(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)^2}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^2} dx}{\frac{4a(a^2 - b^2)}{b(Ab - aB) \sin(c + dx)}} +$$

$$\frac{2ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2}{\downarrow \text{3534}}$$

3.381. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$

$$\int \frac{8Aa^4 + 9bBa^3 - 29Ab^2a^2 - 3b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \cos(c+dx)a + 15Ab^4 + b(-7Ba^3 + 11Aba^2 + b^2Ba - 5Ab^3) \cos^2(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx + \frac{b(-7a^3B + 11a^2)}{ad(a^2 - b^2)\sqrt{\cos(c+dx)}}$$

$$\frac{4a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{b(Ab - aB) \sin(c+dx)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2}$$

↓ 27

$$\int \frac{8Aa^4 + 9bBa^3 - 29Ab^2a^2 - 3b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \cos(c+dx)a + 15Ab^4 + b(-7Ba^3 + 11Aba^2 + b^2Ba - 5Ab^3) \cos^2(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx + \frac{b(-7a^3B + 11a^2)}{ad(a^2 - b^2)\sqrt{\cos(c+dx)}}$$

$$\frac{4a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{b(Ab - aB) \sin(c+dx)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2}$$

↓ 3042

$$\int \frac{8Aa^4 + 9bBa^3 - 29Ab^2a^2 - 3b^3Ba - 4(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sin(c+dx + \frac{\pi}{2})a + 15Ab^4 + b(-7Ba^3 + 11Aba^2 + b^2Ba - 5Ab^3) \sin(c+dx + \frac{\pi}{2})^2}{\sin(c+dx + \frac{\pi}{2})^{\frac{3}{2}}(a+b \sin(c+dx + \frac{\pi}{2}))} dx + \frac{b(-7a^3B + 11a^2)}{ad(a^2 - b^2)\sqrt{\sin(c+dx + \frac{\pi}{2})}}$$

$$\frac{4a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{b(Ab - aB) \sin(c+dx)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2}$$

↓ 3534

$$2 \int \frac{-8Ba^5 + 24Aba^4 + 5b^2Ba^3 - 33Ab^3a^2 - 3b^4Ba + 4(2Aa^4 + 4bBa^3 - 10Ab^2a^2 - b^3Ba + 5Ab^4) \cos(c+dx)a + 15Ab^5 + b(8Aa^4 + 9bBa^3 - 29Ab^2a^2 - 3b^3Ba + 15Ab^4) \cos(c+dx)}{2 \sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx$$

$$\frac{2a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{b(Ab - aB) \sin(c+dx)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2}$$

↓ 27

$$\frac{2(8a^4A + 9a^3bB - 29a^2Ab^2 - 3ab^3B + 15Ab^4) \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \int \frac{-8Ba^5 + 24Aba^4 + 5b^2Ba^3 - 33Ab^3a^2 - 3b^4Ba + 4(2Aa^4 + 4bBa^3 - 10Ab^2a^2 - b^3Ba + 5Ab^4) \cos(c+dx)a + 15Ab^5 + b(8Aa^4 + 9bBa^3 - 29Ab^2a^2 - 3b^3Ba + 15Ab^4) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx$$

$$\frac{2a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{b(Ab - aB) \sin(c+dx)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} \frac{4a(a^2 - b^2)}{2ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2}$$

↓ 3042

3.381. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$

$$\frac{\int \frac{-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2-3b^4Ba+4(2Aa^4+4bBa^3-10Ab^2a^2-b^3Ba+5Ab^4)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}} = \frac{2a(a^2-b^2)}{4a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)}{2ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2}$$

↓ 3538

$$\frac{\int \frac{-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2}{\sqrt{\cos(c+dx)}} dx - \frac{b(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2)}{a}}{\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}} = \frac{2a(a^2-b^2)}{4a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)}{2ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2}$$

↓ 25

$$\frac{\int \frac{b(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2)}{\sqrt{\cos(c+dx)}} dx + \frac{b(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2)}{a}}{\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}} = \frac{2a(a^2-b^2)}{4a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)}{2ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2}$$

↓ 3042

$$\frac{\int \frac{b(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2)}{a}}{\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}} = \frac{2a(a^2-b^2)}{4a(a^2-b^2)}$$

$$\frac{b(Ab-aB)\sin(c+dx)}{2ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^2}$$

↓ 3119

3.381. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^3} dx$

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{b(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2-3b^4Ba+15Ab^5)-ab^2(-7Ba^3+11Aba^2+b^2Ba-5Ab^3)\sin(c+dx)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx - ab(-7a^3B+11a^2bB-5ab^2B+b^3B)}{b(a+b\sin(c+dx+\frac{\pi}{2}))} - \frac{a}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\sqrt{\cos(c + dx)}(a + b\cos(c + dx))^2}$$

↓ 3481

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b(-15a^5B+35a^4Ab+6a^3b^2B-38a^2Ab^3-3ab^4B+15Ab^5)\int \frac{1}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} dx - ab(-7a^3B+11a^2bB-5ab^2B+b^3B)}{b(a+b\cos(c+dx))} - \frac{a}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\sqrt{\cos(c + dx)}(a + b\cos(c + dx))^2}$$

↓ 3042

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b(-15a^5B+35a^4Ab+6a^3b^2B-38a^2Ab^3-3ab^4B+15Ab^5)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx - ab(-7a^3B+11a^2bB-5ab^2B+b^3B)}{b(a+b\sin(c+dx+\frac{\pi}{2}))} - \frac{a}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\sqrt{\cos(c + dx)}(a + b\cos(c + dx))^2}$$

↓ 3120

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b(-15a^5B+35a^4Ab+6a^3b^2B-38a^2Ab^3-3ab^4B+15Ab^5)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))} dx - ab(-7a^3B+11a^2bB-5ab^2B+b^3B)}{b(a+b\sin(c+dx+\frac{\pi}{2}))} - \frac{a}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\sqrt{\cos(c + dx)}(a + b\cos(c + dx))^2}$$

↓ 3284

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\sqrt{\cos(c + dx)}(a + b\cos(c + dx))^2} +$$

$$\frac{b(-7a^3B+11a^2Ab+ab^2B-5Ab^3)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))} + \frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4)E(\frac{1}{2})}{d}$$

3.381. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^3} dx$

4a(a^2 - b^2)

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^3),x]`

output `(b*(A*b - a*B)*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2) + ((b*(11*a^2*A*b - 5*A*b^3 - 7*a^3*B + a*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])) + (-(((2*(8*a^4*A - 29*a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B)*EllipticE[(c + d*x)/2, 2])/d + ((-2*a*b*(11*a^2*A*b - 5*A*b^3 - 7*a^3*B + a*b^2*B)*EllipticF[(c + d*x)/2, 2])/d + (2*b*(35*a^4*A*b - 38*a^2*A*b^3 + 15*A*b^5 - 15*a^5*B + 6*a^3*b^2*B - 3*a*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)*d)/b)/a) + (2*(8*a^4*A - 29*a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B)*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]]))/(2*a*(a^2 - b^2)))/(4*a*(a^2 - b^2))`

3.381.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Ssin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Ssin[e + f*x])^m/(c + d*Ssin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m + 1)*((c + d*Ssin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.381.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1974 vs. 2(480) = 960.

Time = 12.74 (sec) , antiderivative size = 1975, normalized size of antiderivative = 4.70

method	result	size
default	Expression too large to display	1975

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A/a^3/sin(1/
2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+
1/2*c),2^(1/2)))+4*A*b^2/a^3/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(
-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*
c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*A*b/a^2*(-
1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+s
in(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-
b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1
/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*
d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x
+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(
a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)
)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*...
```

$$3.381. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$$

3.381.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^3,x, algorithm m="fricas")`

output `Timed out`

3.381.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.381.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^3,x, algorithm m="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.381.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*cos(d*x + c)^(3/2)), x)`

3.381.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} (a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^3), x)`

3.382
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$$

3.382.1 Optimal result 3562
 3.382.2 Mathematica [A] (warning: unable to verify) 3563
 3.382.3 Rubi [A] (verified) 3564
 3.382.4 Maple [B] (warning: unable to verify) 3571
 3.382.5 Fracas [F(-1)] 3572
 3.382.6 Sympy [F(-1)] 3573
 3.382.7 Maxima [F(-1)] 3573
 3.382.8 Giac [F] 3573
 3.382.9 Mupad [F(-1)] 3574

3.382.1 Optimal result

Integrand size = 33, antiderivative size = 523

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$$

$$= \frac{(24a^4Ab - 65a^2Ab^3 + 35Ab^5 - 8a^5B + 29a^3b^2B - 15ab^4B) E\left(\frac{1}{2}(c+dx) \mid 2\right)}{4a^4(a^2 - b^2)^2 d}$$

$$+ \frac{(8a^4A - 61a^2Ab^2 + 35Ab^4 + 33a^3bB - 15ab^3B) \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{12a^3(a^2 - b^2)^2 d}$$

$$+ \frac{b(63a^4Ab - 86a^2Ab^3 + 35Ab^5 - 35a^5B + 38a^3b^2B - 15ab^4B) \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{4a^4(a-b)^2(a+b)^3 d}$$

$$+ \frac{(8a^4A - 61a^2Ab^2 + 35Ab^4 + 33a^3bB - 15ab^3B) \sin(c+dx)}{12a^3(a^2 - b^2)^2 d \cos^{\frac{3}{2}}(c+dx)}$$

$$- \frac{(24a^4Ab - 65a^2Ab^3 + 35Ab^5 - 8a^5B + 29a^3b^2B - 15ab^4B) \sin(c+dx)}{4a^4(a^2 - b^2)^2 d \sqrt{\cos(c+dx)}}$$

$$+ \frac{b(Ab - aB) \sin(c+dx)}{2a(a^2 - b^2) d \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2}$$

$$+ \frac{b(13a^2Ab - 7Ab^3 - 9a^3B + 3ab^2B) \sin(c+dx)}{4a^2(a^2 - b^2)^2 d \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))}$$

output $\frac{1}{4}*(24*A*a^4*b-65*A*a^2*b^3+35*A*b^5-8*B*a^5+29*B*a^3*b^2-15*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^{(1/2)})/a^4/(a^2-b^2)^2/d+1/12*(8*A*a^4-61*A*a^2*b^2+35*A*b^4+33*B*a^3*b-15*B*a*b^3)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^{(1/2)})/a^3/(a^2-b^2)^2/d+1/4*b*(63*A*a^4*b-86*A*a^2*b^3+35*A*b^5-35*B*a^5+38*B*a^3*b^2-15*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^{(1/2)})/a^4/(a-b)^2/(a+b)^3/d+1/12*(8*A*a^4-61*A*a^2*b^2+35*A*b^4+33*B*a^3*b-15*B*a*b^3)*sin(d*x+c)/a^3/(a^2-b^2)^2/d/cos(d*x+c)^{(3/2)}+1/2*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/cos(d*x+c)^{(3/2)}/(a+b*cos(d*x+c))^2+1/4*b*(13*A*a^2*b-7*A*b^3-9*B*a^3+3*B*a*b^2)*sin(d*x+c)/a^2/(a^2-b^2)^2/d/cos(d*x+c)^{(3/2)}/(a+b*cos(d*x+c))-1/4*(24*A*a^4*b-65*A*a^2*b^3+35*A*b^5-8*B*a^5+29*B*a^3*b^2-15*B*a*b^4)*sin(d*x+c)/a^4/(a^2-b^2)^2/d/cos(d*x+c)^{(1/2)}$

3.382.2 Mathematica [A] (warning: unable to verify)

Time = 7.13 (sec) , antiderivative size = 570, normalized size of antiderivative = 1.09

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$$

$$= \frac{2(16a^6A + 328a^4Ab^2 - 641a^2Ab^4 + 315Ab^6 - 168a^5bB + 285a^3b^3B - 135ab^5B) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + \frac{(160a^5Ab - 512a^3Ab^3 + 280aAb^5 - 48a^4b^2B + 128a^2b^4B - 64b^6B) \operatorname{EllipticE}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + \frac{\sqrt{\cos(c + dx)} \left(\frac{2 \sec(c + dx)(-3Ab \sin(c + dx) + aB \sin(c + dx))}{a^4} + \frac{Ab^4 \sin(c + dx) - ab^3 B \sin(c + dx)}{2a^3(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{17a^2Ab^4 \sin(c + dx) - 11Ab^6 \sin(c + dx)}{4a^4(a^2 - b^2)} \right)}{d}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^3), x]`

output $((2*(16*a^6*A + 328*a^4*A*b^2 - 641*a^2*A*b^4 + 315*A*b^6 - 168*a^5*b*B + 285*a^3*b^3*B - 135*a*b^5*B)*\text{EllipticPi}[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + ((160*a^5*A*b - 512*a^3*A*b^3 + 280*a*A*b^5 - 48*a^6*B + 240*a^4*b^2*B - 120*a^2*b^4*B)*(2*\text{EllipticF}[(c + d*x)/2, 2] - (2*a*\text{EllipticPi}[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)))/b + (2*(72*a^4*A*b^2 - 195*a^2*A*b^4 + 105*A*b^6 - 24*a^5*b*B + 87*a^3*b^3*B - 45*a*b^5*B)*\text{Cos}[2*(c + d*x)]*(-2*a*b*\text{EllipticE}[\text{ArcSin}[\text{Sqrt}[\text{Cos}[c + d*x]]], -1] + 2*a*(a + b)*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Cos}[c + d*x]]], -1] + (-2*a^2 + b^2)*\text{EllipticPi}[-(b/a), \text{ArcSin}[\text{Sqrt}[\text{Cos}[c + d*x]]], -1])*Sin[c + d*x])/(a*b^2*\text{Sqrt}[1 - \text{Cos}[c + d*x]^2]*(-1 + 2*\text{Cos}[c + d*x]^2))/(48*a^4*(a - b)^2*(a + b)^2*d) + (\text{Sqrt}[\text{Cos}[c + d*x]]*((2*\text{Sec}[c + d*x]*(-3*A*b*\text{Sin}[c + d*x] + a*B*\text{Sin}[c + d*x]))/a^4 + (A*b^4*\text{Sin}[c + d*x] - a*b^3*B*\text{Sin}[c + d*x])/(2*a^3*(a^2 - b^2)*(a + b*\text{Cos}[c + d*x])^2) + (17*a^2*A*b^4*\text{Sin}[c + d*x] - 11*A*b^6*\text{Sin}[c + d*x] - 13*a^3*b^3*B*\text{Sin}[c + d*x] + 7*a*b^5*B*\text{Sin}[c + d*x])/(4*a^4*(a^2 - b^2)^2*(a + b*\text{Cos}[c + d*x])) + (2*A*\text{Sec}[c + d*x]*\text{Tan}[c + d*x])/(3*a^3)))/d$

3.382.3 Rubi [A] (verified)

Time = 4.00 (sec) , antiderivative size = 513, normalized size of antiderivative = 0.98, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{5/2} (a + b \sin(c + dx + \frac{\pi}{2}))^3} dx$$

↓ 3479

$$\frac{\int \frac{4Aa^2 + 3bBa - 4(Ab - aB) \cos(c + dx)a - 7Ab^2 + 5b(Ab - aB) \cos^2(c + dx)}{2 \cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx}{\frac{2a(a^2 - b^2)}{b(Ab - aB) \sin(c + dx)}} +$$

$$\frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2}$$

↓ 27

3.382. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$

$$\frac{\int \frac{4Aa^2+3bBa-4(Ab-aB)\cos(c+dx)a-7Ab^2+5b(Ab-aB)\cos^2(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2} dx}{\frac{4a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)}} + \frac{2ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2}{b(Ab-aB)\sin(c+dx)}$$

3042

$$\frac{\int \frac{4Aa^2+3bBa-4(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-7Ab^2+5b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{5/2}(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx}{\frac{4a(a^2-b^2)}{b(Ab-aB)\sin(c+dx)}} + \frac{2ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2}{b(Ab-aB)\sin(c+dx)}$$

3534

$$\frac{\int \frac{8Aa^4+33bBa^3-61Ab^2a^2-15b^3Ba-4(-2Ba^3+4Aba^2-b^2Ba-Ab^3)\cos(c+dx)a+35Ab^4+3b(-9Ba^3+13Aba^2+3b^2Ba-7Ab^3)\cos^2(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))} dx}{\frac{4a(a^2-b^2)}{a(a^2-b^2)}} + \frac{b(-9a^3B+1)}{ad(a^2-b^2)}$$

$$\frac{4a(a^2-b^2)}{2ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2} \frac{b(Ab-aB)\sin(c+dx)}{b(Ab-aB)\sin(c+dx)}$$

27

$$\frac{\int \frac{8Aa^4+33bBa^3-61Ab^2a^2-15b^3Ba-4(-2Ba^3+4Aba^2-b^2Ba-Ab^3)\cos(c+dx)a+35Ab^4+3b(-9Ba^3+13Aba^2+3b^2Ba-7Ab^3)\cos^2(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))} dx}{\frac{4a(a^2-b^2)}{2a(a^2-b^2)}} + \frac{b(-9a^3B+1)}{ad(a^2-b^2)}$$

$$\frac{4a(a^2-b^2)}{2ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2} \frac{b(Ab-aB)\sin(c+dx)}{b(Ab-aB)\sin(c+dx)}$$

3042

$$\frac{\int \frac{8Aa^4+33bBa^3-61Ab^2a^2-15b^3Ba-4(-2Ba^3+4Aba^2-b^2Ba-Ab^3)\sin(c+dx+\frac{\pi}{2})a+35Ab^4+3b(-9Ba^3+13Aba^2+3b^2Ba-7Ab^3)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{5/2}(a+b\sin(c+dx+\frac{\pi}{2}))} dx}{\frac{4a(a^2-b^2)}{2a(a^2-b^2)}} + \frac{b(-9a^3B+1)}{ad(a^2-b^2)}$$

$$\frac{4a(a^2-b^2)}{2ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2} \frac{b(Ab-aB)\sin(c+dx)}{b(Ab-aB)\sin(c+dx)}$$

3534

3.382. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^3} dx$

$$2 \int \frac{-b(8Aa^4 + 33bBa^3 - 61Ab^2a^2 - 15b^3Ba + 35Ab^4) \cos^2(c+dx) - 4a(2Aa^4 - 12bBa^3 + 14Ab^2a^2 + 3b^3Ba - 7Ab^4) \cos(c+dx) + 3(-8Ba^5 + 24Aba^4 + 29b^2Ba^3 - 65Ab^3Ba^2 + 35b^4Ba - 7b^5) \sin(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$$

$$\frac{2a(a^2 - b^2)}{3a}$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2}$$

↓ 27

$$2 \int \frac{(8a^4A + 33a^3bB - 61a^2Ab^2 - 15ab^3B + 35Ab^4) \sin(c+dx) - b(8Aa^4 + 33bBa^3 - 61Ab^2a^2 - 15b^3Ba + 35Ab^4) \cos^2(c+dx) - 4a(2Aa^4 - 12bBa^3 + 14Ab^2a^2 + 3b^3Ba - 7Ab^4) \cos(c+dx) + 3(-8Ba^5 + 24Aba^4 + 29b^2Ba^3 - 65Ab^3Ba^2 + 35b^4Ba - 7b^5) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} dx$$

$$\frac{2a(a^2 - b^2)}{3a}$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2}$$

↓ 3042

$$2 \int \frac{(8a^4A + 33a^3bB - 61a^2Ab^2 - 15ab^3B + 35Ab^4) \sin(c+dx) - b(8Aa^4 + 33bBa^3 - 61Ab^2a^2 - 15b^3Ba + 35Ab^4) \sin(c+dx + \frac{\pi}{2})^2 - 4a(2Aa^4 - 12bBa^3 + 14Ab^2a^2 + 3b^3Ba - 7Ab^4) \cos(c+dx) + 3(-8Ba^5 + 24Aba^4 + 29b^2Ba^3 - 65Ab^3Ba^2 + 35b^4Ba - 7b^5) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} dx$$

$$\frac{2a(a^2 - b^2)}{3a}$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2}$$

↓ 3534

$$2 \int \frac{(8a^4A + 33a^3bB - 61a^2Ab^2 - 15ab^3B + 35Ab^4) \sin(c+dx) - 8Aa^6 - 72bBa^5 + 128Ab^2a^4 + 99b^3Ba^3 - 223Ab^4a^2 - 45b^5Ba + 4(-6Ba^5 + 20Aba^4 + 30b^2Ba^3 - 64Ab^3Ba^2 + 35b^4Ba - 7b^5) \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} dx$$

$$\frac{2a(a^2 - b^2)}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx)}{2ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2}$$

↓ 27

3.382. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \int \frac{8Aa^6-72bBa^5+128Ab^2a^4+}{}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3042

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \int \frac{8Aa^6-72bBa^5+128Ab^2a^4+}{}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3538

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 25

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3042

3.382. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^3} dx$

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3119

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{a(8Aa^4+33bBa^3-61Ab^2a^2-15ab^3B+35Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3481

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{ab(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3042

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{ab(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3120

3.382. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^3} dx$

$$\frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{3b^2(-35a^5B+63a^4Ab+38a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2}$$

↓ 3284

$$\frac{b(Ab - aB)\sin(c + dx)}{2ad(a^2 - b^2)\cos^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^2} +$$

$$\frac{b(-9a^3B+13a^2Ab+3ab^2B-7Ab^3)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))} + \frac{2(8a^4A+33a^3bB-61a^2Ab^2-15ab^3B+35Ab^4)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{6(-8a^5B+24a^4Ab+29a^3b^2B-65a^2Ab^3-15ab^4B+35Ab^5)\sin(c+dx)}{ad\sqrt{\cos(c+dx)}}$$

```
input Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^3),x]
```

```
output (b*(A*b - a*B)*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2) + ((b*(13*a^2*A*b - 7*A*b^3 - 9*a^3*B + 3*a*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])) + ((2*(8*a^4*A - 61*a^2*A*b^2 + 35*A*b^4 + 33*a^3*b*B - 15*a*b^3*B)*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2))) - (((6*(24*a^4*A*b - 65*a^2*A*b^3 + 35*A*b^5 - 8*a^5*B + 29*a^3*b^2*B - 15*a*b^4*B)*EllipticE[(c + d*x)/2, 2])/d + ((2*a*b*(8*a^4*A - 61*a^2*A*b^2 + 35*A*b^4 + 33*a^3*b*B - 15*a*b^3*B)*EllipticF[(c + d*x)/2, 2])/d + (6*b^2*(63*a^4*A*b - 86*a^2*A*b^3 + 35*A*b^5 - 35*a^5*B + 38*a^3*b^2*B - 15*a*b^4*B)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d))/b)/a) + (6*(24*a^4*A*b - 65*a^2*A*b^3 + 35*A*b^5 - 8*a^5*B + 29*a^3*b^2*B - 15*a*b^4*B)*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]]))/(3*a)/(2*a*(a^2 - b^2))/(4*a*(a^2 - b^2))
```


3.382.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)]))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3534 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

```
rule 3538 Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x])*(c + d*Sin[e + f*x]), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.382.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2130 vs. 2(579) = 1158.

Time = 20.43 (sec) , antiderivative size = 2131, normalized size of antiderivative = 4.07

method	result	size
default	Expression too large to display	2131

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

$$3.382. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^3} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A/a^3*(-1/6*
cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(c
os(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*
x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*E
llipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2*(-3*A*b+B*a)/a^4/sin(1/2*d*x+1/2*c
)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2)))+2*(A*b-B*a)*b/a^2*(-1/2/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1
/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)
^2-3/4*b^2*(3*a^2-b^2)/a^2/(a^2-b^2)^2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+
1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-7/8/(a
+b)/(a^2-b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/
2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),2^(1/2))+1/4/(a+b)/(a^2-b^2)/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2
*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*b+3/8/(a+b)/(a^2-b^2)/a^2*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(
1/2))*b^2-9/8*b/(a^2-b^2)^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*...

```

3.382.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^3,x, algorithm="fricas")`

output `Timed out`

3.382.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.382.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^3,x, algorithm m="maxima")`

output `Timed out`

3.382.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*cos(d*x + c)^(5/2)), x)`

3.382.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{5}{2}}(a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^3),x)`output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^3), x)`

3.383
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.383.1 Optimal result	3575
3.383.2 Mathematica [A] (verified)	3575
3.383.3 Rubi [A] (verified)	3576
3.383.4 Maple [B] (verified)	3577
3.383.5 Fricas [C] (verification not implemented)	3578
3.383.6 Sympy [F(-1)]	3578
3.383.7 Maxima [F]	3579
3.383.8 Giac [F]	3579
3.383.9 Mupad [F(-1)]	3579

3.383.1 Optimal result

Integrand size = 36, antiderivative size = 44

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$$

$$= \frac{6BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d} + \frac{2B \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{5d}$$

output `6/5*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/5*B*cos(d*x+c)^(3/2)*sin(d*x+c)/d`

3.383.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.93

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$$

$$= \frac{B\left(6E\left(\frac{1}{2}(c+dx) \mid 2\right) + \sqrt{\cos(c+dx)} \sin(2(c+dx))\right)}{5d}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(B*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*d)`

3.383.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {2011, 3042, 3115, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx \\ & \quad \downarrow \text{2011} \\ & B \int \cos^{\frac{5}{2}}(c+dx) dx \\ & \quad \downarrow \text{3042} \\ & B \int \sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} dx \\ & \quad \downarrow \text{3115} \\ & B \left(\frac{3}{5} \int \sqrt{\cos(c+dx)} dx + \frac{2\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{3}{5} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{2\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \right) \\ & \quad \downarrow \text{3119} \\ & B \left(\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5d} + \frac{2\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{5d} \right) \end{aligned}$$

input `Int[(Cos[c + d*x]^(5/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `B*((6*EllipticE[(c + d*x)/2, 2])/(5*d) + (2*Cos[c + d*x]^(3/2)*Sin[c + d*x]))/(5*d)`

3.383. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$

3.383.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :>
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3115 Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*
x]*((b*Sine[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*Sine
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

3.383.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 202 vs. 2(64) = 128.

Time = 4.68 (sec) , antiderivative size = 203, normalized size of antiderivative = 4.61

method	result
default	$-\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}}$

```
input int(cos(d*x+c)^(5/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)
```

3.383. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$

output
$$\frac{-2/5*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*B*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)}))/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$$

3.383.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.75

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \frac{2B\cos(dx+c)^{\frac{3}{2}}\sin(dx+c)+3i\sqrt{2}B\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c)))}{5}$$

input `integrate(cos(d*x+c)^(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output
$$\frac{1/5*(2*B*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)+3*I*\text{sqrt}(2)*B*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c)))-3*I*\text{sqrt}(2)*B*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c)))}{d}$$

3.383.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output Timed out

3.383.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{5}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.383.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{5}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.383.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{\cos(c+dx)^{5/2}(Ba+Bb\cos(c+dx))}{a+b\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(5/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `int((cos(c + d*x)^(5/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.384 $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.384.1 Optimal result 3580
 3.384.2 Mathematica [A] (verified) 3580
 3.384.3 Rubi [A] (verified) 3581
 3.384.4 Maple [B] (verified) 3582
 3.384.5 Fricas [C] (verification not implemented) 3583
 3.384.6 Sympy [F(-1)] 3583
 3.384.7 Maxima [F] 3583
 3.384.8 Giac [F] 3584
 3.384.9 Mupad [F(-1)] 3584

3.384.1 Optimal result

Integrand size = 36, antiderivative size = 44

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2B \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + \frac{2B \sqrt{\cos(c+dx)} \sin(c+dx)}{3d}$$

output `2/3*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*B*sin(d*x+c)*cos(d*x+c)^(1/2)/d`

3.384.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2B \left(\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sqrt{\cos(c+dx)} \sin(c+dx) \right)}{3d}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(2*B*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*d)`

3.384. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$

3.384.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {2011, 3042, 3115, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \cos^{\frac{3}{2}}(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}} dx$$

$$\downarrow \text{3115}$$

$$B \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right)$$

$$\downarrow \text{3120}$$

$$B \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3d} \right)$$

input `Int[(Cos[c + d*x]^(3/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `B*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*d))`

3.384.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3115 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sine[c + d*x])^(n - 1)/(d*n), x] + Simp[b^2*((n - 1)/n Int[(b*Sine
[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[
2*n]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.384.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(64) = 128.

Time = 3.36 (sec) , antiderivative size = 180, normalized size of antiderivative = 4.09

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} B\left(4\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) - 2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{\frac{1}{2} + \frac{\cos(dx+c)}{2}}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d$

input `int(cos(d*x+c)^(3/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURN
NVERBOSE)`

output `-2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(4*sin(1/2*
d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.384.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

3.384.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.61

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$= \frac{2B\sqrt{\cos(dx+c)}\sin(dx+c) - i\sqrt{2}B\text{weierstrassPInverse}(-4,0,\cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}B\text{weierstrassPInverse}(-4,0,\cos(dx+c) - i\sin(dx+c))}{3d}$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `1/3*(2*B*sqrt(cos(d*x + c))*sin(d*x + c) - I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

3.384.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.384.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(Bb\cos(dx+c) + Ba)\cos(dx+c)^{\frac{3}{2}}}{b\cos(dx+c) + a} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.384. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$

3.384.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{3}{2}}}{b\cos(dx+c)+a} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x+c)+B*a)*cos(d*x+c)^(3/2)/(b*cos(d*x+c)+a),x)`

3.384.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{\cos(c+dx)^{\frac{3}{2}}(Ba+Bb\cos(c+dx))}{a+b\cos(c+dx)} dx$$

input `int((cos(c+d*x)^(3/2)*(B*a+B*b*cos(c+d*x)))/(a+b*cos(c+d*x)),x)`

output `int((cos(c+d*x)^(3/2)*(B*a+B*b*cos(c+d*x)))/(a+b*cos(c+d*x)),x)`

3.385
$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx$$

3.385.1 Optimal result 3585
 3.385.2 Mathematica [A] (verified) 3585
 3.385.3 Rubi [A] (verified) 3586
 3.385.4 Maple [B] (verified) 3587
 3.385.5 Fricas [C] (verification not implemented) 3587
 3.385.6 Sympy [F(-1)] 3588
 3.385.7 Maxima [F] 3588
 3.385.8 Giac [F] 3588
 3.385.9 Mupad [F(-1)] 3589

3.385.1 Optimal result

Integrand size = 36, antiderivative size = 17

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2BE(\frac{1}{2}(c+dx)|2)}{d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d`

3.385.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{a+b \cos(c+dx)} dx = \frac{2BE(\frac{1}{2}(c+dx)|2)}{d}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(2*B*EllipticE[(c + d*x)/2, 2])/d`

3.385.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2011, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(aB + bB \cos(c+dx))}{a + b \cos(c+dx)} dx$$

↓ 2011

$$B \int \sqrt{\cos(c+dx)} dx$$

↓ 3042

$$B \int \sqrt{\sin\left(c+dx + \frac{\pi}{2}\right)} dx$$

↓ 3119

$$\frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{d}$$

input `Int[(Sqrt[Cos[c + d*x]]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x]),x]`

output `(2*B*EllipticE[(c + d*x)/2, 2])/d`

3.385.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.385.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. $2(43) = 86$.

Time = 2.36 (sec) , antiderivative size = 134, normalized size of antiderivative = 7.88

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} B\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$
risch	$-\frac{i\sqrt{2}B\sqrt{(e^{2i(dx+c)}+1)e^{-i(dx+c)}}}{d} - i\left(\frac{2(e^{2i(dx+c)}+1)}{\sqrt{(e^{2i(dx+c)}+1)e^{i(dx+c)}}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}}{\sqrt{e^{3i(dx+c)}+e^{i(dx+c)}}}\right)$

input `int(cos(d*x+c)^(1/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.385.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 3.47

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \frac{i\sqrt{2}B\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))) - i\sqrt{2}B\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c)))}{d}$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x,algorithm="fracas")`

3.385.
$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

output `(I*sqrt(2)*B*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*B*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.385.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c + dx)}(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.385.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c + dx)}(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\cos(dx + c)}}{b \cos(dx + c) + a} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a), x)`

3.385.8 Giac [F]

$$\int \frac{\sqrt{\cos(c + dx)}(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\cos(dx + c)}}{b \cos(dx + c) + a} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a), x)`

3.385.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx = \int \frac{\sqrt{\cos(c+dx)}(Ba+Bb\cos(c+dx))}{a+b\cos(c+dx)} dx$$

input `int((cos(c + d*x)^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`output `int((cos(c + d*x)^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.386 $\int \frac{aB+bB \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx$

3.386.1 Optimal result 3590
 3.386.2 Mathematica [A] (verified) 3590
 3.386.3 Rubi [A] (verified) 3591
 3.386.4 Maple [C] (verified) 3592
 3.386.5 Fricas [C] (verification not implemented) 3592
 3.386.6 Sympy [F(-1)] 3593
 3.386.7 Maxima [F] 3593
 3.386.8 Giac [F] 3593
 3.386.9 Mupad [F(-1)] 3594

3.386.1 Optimal result

Integrand size = 36, antiderivative size = 17

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \frac{2B \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d`

3.386.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \frac{2B \operatorname{EllipticF}(\frac{1}{2}(c + dx), 2)}{d}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])),x]`

output `(2*B*EllipticF[(c + d*x)/2, 2])/d`

3.386.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2011, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx$$

↓ 2011

$$B \int \frac{1}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$B \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3120

$$\frac{2B \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d}$$

input `Int[(a*B + b*B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])),x]`

output `(2*B*EllipticF[(c + d*x)/2, 2])/d`

3.386.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :> Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.386.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.49 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{2B \operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2} \mid \sqrt{2}\right)}{d}$	19

input `int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `2*B/d*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2))`

3.386.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.12

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx$$

$$= \frac{-i \sqrt{2} B \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} B \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `(-I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

3.386.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.386.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)\sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.386.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)\sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.386.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx = \int \frac{B a + B b \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))),x)`output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))), x)`

$$3.387 \quad \int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

3.387.1 Optimal result	3595
3.387.2 Mathematica [A] (verified)	3595
3.387.3 Rubi [A] (verified)	3596
3.387.4 Maple [B] (verified)	3597
3.387.5 Fracas [C] (verification not implemented)	3598
3.387.6 Sympy [F(-1)]	3598
3.387.7 Maxima [F]	3598
3.387.8 Giac [F]	3599
3.387.9 Mupad [F(-1)]	3599

3.387.1 Optimal result

Integrand size = 36, antiderivative size = 40

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = -\frac{2BE(\frac{1}{2}(c + dx)|2)}{d} + \frac{2B \sin(c + dx)}{d\sqrt{\cos(c + dx)}}$$

output `-2*B*(cos(1/2*d*x+1/2*c)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d+2*B*sin(d*x+c)/d/cos(d*x+c)^(1/2)`

3.387.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = B \left(-\frac{2E(\frac{1}{2}(c + dx)|2)}{d} + \frac{2 \sin(c + dx)}{d\sqrt{\cos(c + dx)}} \right)$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])),x]`

output `B*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))`

3.387. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$

3.387.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {2011, 3042, 3116, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3116}$$

$$B \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\cos(c + dx)} dx \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right)$$

$$\downarrow \text{3119}$$

$$B \left(\frac{2 \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2E(\frac{1}{2}(c + dx)|2)}{d} \right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])),x]`

output `B*((-2*EllipticE[(c + d*x)/2, 2])/d + (2*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))`

3.387.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3116 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) I
nt[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

3.387.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(64) = 128.

Time = 2.08 (sec) , antiderivative size = 183, normalized size of antiderivative = 4.58

method	result
default	$-\frac{2B \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1}} \right)}{\sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1}} d$

```
input int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b),x,method=_RETUR
NVERBOSE)
```

```
output -2*B*(-2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)
^(1/2)*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/
2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellip
ticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.387. $\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx$

3.387.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.40

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$= \frac{-i \sqrt{2} B \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i}{\dots}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `(-I*sqrt(2)*B*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*B*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*B*sqrt(cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c))`

3.387.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.387.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

3.387. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.387.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.387.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx)^{3/2} (a + b \cos(c + dx))} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))),x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))), x)`

3.388
$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

3.388.1 Optimal result 3600
 3.388.2 Mathematica [A] (verified) 3600
 3.388.3 Rubi [A] (verified) 3601
 3.388.4 Maple [B] (verified) 3602
 3.388.5 Fricas [C] (verification not implemented) 3603
 3.388.6 Sympy [F(-1)] 3603
 3.388.7 Maxima [F] 3604
 3.388.8 Giac [F] 3604
 3.388.9 Mupad [F(-1)] 3604

3.388.1 Optimal result

Integrand size = 36, antiderivative size = 44

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \frac{2B \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2B \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)}$$

output `2/3*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/d+2/3*B*sin(d*x+c)/d/cos(d*x+c)^(3/2)`

3.388.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \frac{2B \left(\operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \frac{\sin(c + dx)}{\cos^{\frac{3}{2}}(c + dx)} \right)}{3d}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])),x]`

output `(2*B*(EllipticF[(c + d*x)/2, 2] + Sin[c + d*x]/Cos[c + d*x]^(3/2)))/(3*d)`

3.388.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {2011, 3042, 3116, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

$$\downarrow \text{3116}$$

$$B \left(\frac{1}{3} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{1}{3} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

$$\downarrow \text{3120}$$

$$B \left(\frac{2 \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3d} + \frac{2 \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])),x]`

output `B*((2*EllipticF[(c + d*x)/2, 2])/(3*d) + (2*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))`

3.388.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :> Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3116 `Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Simp[(n + 2)/(b^2*(n + 1)) Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.388.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(64) = 128.

Time = 2.04 (sec) , antiderivative size = 214, normalized size of antiderivative = 4.86

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}$

input `int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

3.388. $\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))} dx$

output
$$-2/3*(-2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)}))*B*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(3/2)}/\sin(1/2*d*x+1/2*c)/d$$

3.388.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.16

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{5/2}(c + dx)(a + b \cos(c + dx))} dx$$

$$= \frac{-i \sqrt{2} B \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} B \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2B \sqrt{\cos(dx + c)} \sin(dx + c)}{3 d \cos(dx + c)^2}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output
$$1/3*(-I*\sqrt{2}*B*\cos(d*x + c)^2*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*B*\cos(d*x + c)^2*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + 2*B*\sqrt{\cos(d*x + c)}*\sin(d*x + c))/(d*\cos(d*x + c)^2)$$

3.388.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{5/2}(c + dx)(a + b \cos(c + dx))} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c)),x)`

output Timed out

3.388.
$$\int \frac{aB + bB \cos(c + dx)}{\cos^{5/2}(c + dx)(a + b \cos(c + dx))} dx$$

3.388.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.388.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.388.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx)^{\frac{5}{2}}(a + b \cos(c + dx))} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))),x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))), x)`

3.389
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.389.1 Optimal result 3605
 3.389.2 Mathematica [A] (verified) 3606
 3.389.3 Rubi [A] (verified) 3606
 3.389.4 Maple [B] (verified) 3610
 3.389.5 Fricas [F(-1)] 3611
 3.389.6 Sympy [F(-1)] 3611
 3.389.7 Maxima [F] 3612
 3.389.8 Giac [F] 3612
 3.389.9 Mupad [F(-1)] 3612

3.389.1 Optimal result

Integrand size = 36, antiderivative size = 116

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = -\frac{2aBE(\frac{1}{2}(c+dx)|2)}{b^2d} + \frac{2(3a^2+b^2)B \text{EllipticF}(\frac{1}{2}(c+dx),2)}{3b^3d} - \frac{2a^3B \text{EllipticPi}(\frac{2b}{a+b},\frac{1}{2}(c+dx),2)}{b^3(a+b)d} + \frac{2B\sqrt{\cos(c+dx)} \sin(c+dx)}{3bd}$$

output

```
-2*a*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d+2/3*(3*a^2+b^2)*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b^3/d-2*a^3*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/b^3/(a+b)/d+2/3*B*sin(d*x+c)*cos(d*x+c)^(1/2)/b/d
```

3.389.2 Mathematica [A] (verified)

Time = 0.96 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.37

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$= B \left(4 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{6a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + 4\sqrt{\cos(c+dx)} \sin(c+dx) - \frac{6(-2abE(\arcsin(\sqrt{\cos(c+dx)}))}{a+b} \right)$$

6bd

input `Integrate[(Cos[c + d*x]^(5/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `(B*(4*EllipticF[(c + d*x)/2, 2] - (6*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b) + 4*Sqrt[Cos[c + d*x]]*Sin[c + d*x] - (6*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(b^2*Sqrt[Sin[c + d*x]^2]))/(6*b*d)`

3.389.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {2011, 3042, 3272, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{\cos^{\frac{5}{2}}(c+dx)}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2}}{a+b\sin(c+dx+\frac{\pi}{2})} dx$$

$$\downarrow \text{3272}$$

3.389. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
& B \left(\frac{2 \int \frac{-3a \cos^2(c+dx) + b \cos(c+dx) + a}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 27 \\
& B \left(\frac{\int \frac{-3a \cos^2(c+dx) + b \cos(c+dx) + a}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{\int \frac{-3a \sin(c+dx+\frac{\pi}{2})^2 + b \sin(c+dx+\frac{\pi}{2}) + a}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 3538 \\
& B \left(\frac{\int \frac{-\frac{ab+(3a^2+b^2) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} - \frac{3a \int \frac{\sqrt{\cos(c+dx)} dx}{b}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 25 \\
& B \left(\frac{\int \frac{\frac{ab+(3a^2+b^2) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{3b} - \frac{3a \int \frac{\sqrt{\cos(c+dx)} dx}{b}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{\int \frac{\frac{ab+(3a^2+b^2) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{3a \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 3119 \\
& B \left(\frac{\int \frac{\frac{ab+(3a^2+b^2) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3b} - \frac{6aE(\frac{1}{2}(c+dx)|2)}{bd} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right) \\
& \quad \downarrow 3481
\end{aligned}$$

3.389. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

$$B \left(\frac{\frac{(3a^2+b^2) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{3a^3 \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx}{b} - \frac{6aE(\frac{1}{2}(c+dx)|2)}{bd}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right)$$

↓ 3042

$$B \left(\frac{\frac{(3a^2+b^2) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{3a^3 \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{6aE(\frac{1}{2}(c+dx)|2)}{bd}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right)$$

↓ 3120

$$B \left(\frac{\frac{2(3a^2+b^2) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{3a^3 \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx}{b} - \frac{6aE(\frac{1}{2}(c+dx)|2)}{bd}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right)$$

↓ 3284

$$B \left(\frac{\frac{2(3a^2+b^2) \text{EllipticF}(\frac{1}{2}(c+dx),2)}{bd} - \frac{6a^3 \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx),2)}{bd(a+b)}}{b} - \frac{6aE(\frac{1}{2}(c+dx)|2)}{bd}}{3b} + \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd} \right)$$

input `Int[(Cos[c + d*x]^(5/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `B*(((-6*a*EllipticE[(c + d*x)/2, 2])/(b*d) + ((2*(3*a^2 + b^2)*EllipticF[(c + d*x)/2, 2])/(b*d) - (6*a^3*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))/b)/(3*b) + (2*sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*b*d))`

3.389.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3272 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 2)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Sin[e + f*x])^(m - 3)*(c + d*Sin[e + f*x])^n*Simp[a^3*d*(m + n) + b^2*(b*c*(m - 2) + a*d*(n + 1)) - b*(a*b*c - b^2*d*(m + n - 1) - 3*a^2*d*(m + n))*Sin[e + f*x] - b^2*(b*c*(m - 1) - a*d*(3*m + 2*n - 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 2] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`


```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

```
rule 3481 Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])/(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.389.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs. $2(188) = 376$.

Time = 6.43 (sec) , antiderivative size = 553, normalized size of antiderivative = 4.77

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}B\left(4\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)a^2b^2-4\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^3-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(a+b\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2}$

```
input int(cos(d*x+c)^(5/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RET
URNVERBOSE)
```

$$3.389. \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

output
$$-2/3*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*B*(4*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^4*a*b^2-4*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*a*b^3-2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*a*b^2+2*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2*b^3+3*a^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-3*a^2*b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+a*b^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})-b^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a^2*b-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a*b^2-3*a^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})/b^3/(a-b)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$$

3.389.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\cos^{5/2}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fricas")`

output Timed out

3.389.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{5/2}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output Timed out

3.389.
$$\int \frac{\cos^{5/2}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

3.389.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algo
rithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)
^2, x)`

3.389.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(5/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algo
rithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)
^2, x)`

3.389.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{5/2}(Ba+Bb\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(5/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x
)`

output `int((cos(c + d*x)^(5/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,
x)`

3.389. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$

3.390
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.390.1 Optimal result 3613
 3.390.2 Mathematica [A] (verified) 3613
 3.390.3 Rubi [A] (verified) 3614
 3.390.4 Maple [A] (verified) 3617
 3.390.5 Fricas [F(-1)] 3617
 3.390.6 Sympy [F(-1)] 3618
 3.390.7 Maxima [F] 3618
 3.390.8 Giac [F] 3618
 3.390.9 Mupad [F(-1)] 3619

3.390.1 Optimal result

Integrand size = 36, antiderivative size = 78

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{2BE\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} - \frac{2aB \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{b^2d} + \frac{2a^2B \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{b^2(a+b)d}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+
1/2*c), 2^(1/2))/b/d-2*a*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/b^2/d+2*a^2*B*(cos(1/2*d*x+1/2*c)^2)
^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))
/b^2/(a+b)/d
```

3.390.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.05

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{2B\left(bE\left(\arcsin\left(\sqrt{\cos(c+dx)}\right) \mid -1\right) - (a+b) \operatorname{EllipticF}\left(\arcsin\left(\sqrt{\cos(c+dx)}\right), -1\right) + a \operatorname{EllipticP}\right)}{b^2d\sqrt{\sin^2(c+dx)}}$$

3.390.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

input `Integrate[(Cos[c + d*x]^(3/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output `(-2*B*(b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] - (a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + a*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(b^2*d*Sqrt[Sin[c + d*x]^2])`

3.390.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2011, 3042, 3283, 3042, 3119, 3282, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\cos^{\frac{3}{2}}(c+dx)}{a+b\cos(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}}{a+b\sin(c+dx+\frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3283} \\
 & B \left(\frac{\int \sqrt{\cos(c+dx)} dx}{b} - \frac{a \int \frac{\sqrt{\cos(c+dx)}}{a+b\cos(c+dx)} dx}{b} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{\int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} - \frac{a \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{a+b\sin(c+dx+\frac{\pi}{2})} dx}{b} \right) \\
 & \quad \downarrow \text{3119}
 \end{aligned}$$

$$\begin{aligned}
& B \left(\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} - \frac{a \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} \right) \\
& \quad \downarrow \text{3282} \\
& B \left(\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} - \frac{a \left(\frac{\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} \right)}{b} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} - \frac{a \left(\frac{\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} \right)}{b} \right) \\
& \quad \downarrow \text{3120} \\
& B \left(\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} - \frac{a \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{a \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} \right)}{b} \right) \\
& \quad \downarrow \text{3284} \\
& B \left(\frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd} - \frac{a \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{bd(a+b)} \right)}{b} \right)
\end{aligned}$$

input `Int[(Cos[c + d*x]^(3/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x
]`

output `B*((2*EllipticE[(c + d*x)/2, 2])/(b*d) - (a*((2*EllipticF[(c + d*x)/2, 2])
/(b*d) - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d)))/b
)`

3.390. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$

3.390.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3282 `Int[Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]/((a_.) + (b_.)*sin[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d/b Int[1/Sqrt[c + d*Sin[e + f*x]], x], x
+ Simp[(b*c - a*d)/b Int[1/((a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x
])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2
- b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3283 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)/((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[b/d Int[Sqrt[a + b*Sin[e + f*x]], x], x
- Simp[(b*c - a*d)/d Int[Sqrt[a + b*Sin[e + f*x]]/(c + d*Sin[e + f*x]),
x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b
^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]`

3.390.4 Maple [A] (verified)

Time = 3.87 (sec) , antiderivative size = 228, normalized size of antiderivative = 2.92

method	result
default	$\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}\left(F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)a^2-F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)b^2(a-b)\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\right)}{b^2(a-b)\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}\right)dx$

```
input int(cos(d*x+c)^(3/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)
```

```
output 2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a*b+EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b-EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^2-a^2*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))/b^2/(a-b)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.390.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="fracas")
```

```
output Timed out
```


3.390.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.390.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^2, x)`

3.390.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^2, x)`

3.390.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{\cos(c+dx)^{3/2}(Ba+Bb\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

output `int((cos(c + d*x)^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

3.391
$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

3.391.1 Optimal result	3620
3.391.2 Mathematica [A] (verified)	3620
3.391.3 Rubi [A] (verified)	3621
3.391.4 Maple [A] (verified)	3623
3.391.5 Fricas [F(-1)]	3623
3.391.6 Sympy [F(-1)]	3624
3.391.7 Maxima [F]	3624
3.391.8 Giac [F]	3624
3.391.9 Mupad [F(-1)]	3625

3.391.1 Optimal result

Integrand size = 36, antiderivative size = 55

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{2B \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{2aB \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{b(a+b)d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))/b/d-2*a*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/b/(a+b)/d`

3.391.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx = \frac{B\left(2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b}\right)}{bd}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x]`

output $(B*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]))/(a + b))/(b*d)$

3.391.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2011, 3042, 3282, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(aB + bB \cos(c+dx))}{(a+b \cos(c+dx))^2} dx$$

$$\downarrow 2011$$

$$B \int \frac{\sqrt{\cos(c+dx)}}{a+b \cos(c+dx)} dx$$

$$\downarrow 3042$$

$$B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{a+b \sin(c+dx+\frac{\pi}{2})} dx$$

$$\downarrow 3282$$

$$B \left(\frac{\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} \right)$$

$$\downarrow 3042$$

$$B \left(\frac{\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} \right)$$

$$\downarrow 3120$$

$$B \left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{a \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b} \right)$$

$$\downarrow 3284$$

$$B \left(\frac{2 \operatorname{EllipticF} \left(\frac{1}{2}(c + dx), 2 \right)}{bd} - \frac{2a \operatorname{EllipticPi} \left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2 \right)}{bd(a + b)} \right)$$

input `Int[(Sqrt[Cos[c + d*x]]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^2,x
]`

output `B*((2*EllipticF[(c + d*x)/2, 2])/(b*d) - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))`

3.391.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x
, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3282 `Int[Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]/((a_.) + (b_.)*sin[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d/b Int[1/Sqrt[c + d*Sin[e + f*x]], x], x
] + Simp[(b*c - a*d)/b Int[1/((a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2
- b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]`

3.391.4 Maple [A] (verified)

Time = 3.09 (sec) , antiderivative size = 189, normalized size of antiderivative = 3.44

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}B\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}\left(F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)a-F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)b\right)}{(a-b)b\sqrt{-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d$

```
input int(cos(d*x+c)^(1/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^2,x,method=_RETURNVERBOSE)
```

```
output -2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^(1/2)*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a-EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*b-a*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))/(a-b)/b/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.391.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x,algorithm="fricas")
```

```
output Timed out
```

3.391.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.391.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(Bb\cos(dx+c)+Ba)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algo
rithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)
^2, x)`

3.391.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{(Bb\cos(dx+c)+Ba)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^2} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^2,x, algo
rithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)
^2, x)`

3.391.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^2} dx = \int \frac{\sqrt{\cos(c+dx)}(Ba+Bb\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

input `int((cos(c + d*x)^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

output `int((cos(c + d*x)^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^2,x)`

3.392
$$\int \frac{aB+bB \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^2} dx$$

3.392.1 Optimal result 3626
 3.392.2 Mathematica [A] (verified) 3626
 3.392.3 Rubi [A] (verified) 3627
 3.392.4 Maple [B] (verified) 3628
 3.392.5 Fricas [F(-1)] 3628
 3.392.6 Sympy [F(-1)] 3629
 3.392.7 Maxima [F] 3629
 3.392.8 Giac [F] 3629
 3.392.9 Mupad [F(-1)] 3630

3.392.1 Optimal result

Integrand size = 36, antiderivative size = 30

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \frac{2B \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{(a + b)d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/(a+b)/d`

3.392.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \frac{2B \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{(a + b)d}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2),x]`

output `(2*B*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d)`

3.392.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2011, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$$

↓ 2011

$$B \int \frac{1}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx$$

↓ 3042

$$B \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} (a + b \sin(c + dx + \frac{\pi}{2}))} dx$$

↓ 3284

$$\frac{2B \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{d(a + b)}$$

input `Int[(a*B + b*B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^2),x]`

output `(2*B*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d)`

3.392.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :> Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.392. $\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_.)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

3.392.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 150 vs. 2(56) = 112.

Time = 2.00 (sec) , antiderivative size = 151, normalized size of antiderivative = 5.03

method	result	size
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} B \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \Pi\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), -\frac{2b}{a-b}, \sqrt{2}\right)}{(a-b)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$	151

```
input int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^2,x,method=_RET
URNVERBOSE)
```

```
output -2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticPi(cos(1/2*d*x+1
/2*c),-2*b/(a-b),2^(1/2))/(a-b)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.392.5 Fracas [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algo
rithm="fracas")
```

```
output Timed out
```

3.392. $\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$

3.392.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.392.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algo
rithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^2*sqrt(cos(d*x +
c))), x)`

3.392.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^2 \sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algo
rithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^2*sqrt(cos(d*x +
c))), x)`

3.392.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx = \int \frac{B a + B b \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^2} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^2),x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^2),x)`

3.393
$$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

3.393.1 Optimal result 3631
 3.393.2 Mathematica [B] (verified) 3631
 3.393.3 Rubi [A] (verified) 3632
 3.393.4 Maple [B] (verified) 3635
 3.393.5 Fricas [F(-1)] 3636
 3.393.6 Sympy [F(-1)] 3636
 3.393.7 Maxima [F] 3637
 3.393.8 Giac [F] 3637
 3.393.9 Mupad [F(-1)] 3637

3.393.1 Optimal result

Integrand size = 36, antiderivative size = 80

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = -\frac{2BE(\frac{1}{2}(c + dx)|2)}{ad} - \frac{2bB \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2)}{a(a + b)d} + \frac{2B \sin(c + dx)}{ad\sqrt{\cos(c + dx)}}$$

```
output -2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x
+1/2*c),2^(1/2))/a/d-2*b*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)
*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))/a/(a+b)/d+2*B*sin(d*x+c)
/a/d/cos(d*x+c)^(1/2)
```

3.393.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 196 vs. 2(80) = 160.

Time = 1.77 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.45

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx =$$

$$B \left(\frac{6b \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + \frac{2a \left(2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} \right)}{b} - \frac{4 \sin(c+dx)}{\sqrt{\cos(c+dx)}} + \frac{2(-2abE(\arcsin(\frac{\sin(c+dx)}{\sqrt{\cos(c+dx)}})) - 2a \operatorname{EllipticE}\left(\frac{1}{2}(c+dx), 2\right))}{a+b} \right)$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2),x]`

output `-1/2*(B*((6*b*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + (2*a*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)))/b - (4*Sin[c + d*x])/Sqrt[Cos[c + d*x]] + (2*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1])*Sin[c + d*x])/(a*b*Sqrt[Sin[c + d*x]^2])))/(a*d)`

3.393.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.99, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {2011, 3042, 3281, 27, 3042, 3538, 25, 27, 3042, 3119, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2} (a + b \sin(c + dx + \frac{\pi}{2}))} dx$$

$$\downarrow \text{3281}$$

3.393. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$

$$\begin{aligned}
& B \left(\frac{2 \int -\frac{b \cos^2(c+dx) + a \cos(c+dx) + b}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} + \frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} \right) \\
& \quad \downarrow 27 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{b \cos^2(c+dx) + a \cos(c+dx) + b}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \frac{b \sin(c+dx + \frac{\pi}{2})^2 + a \sin(c+dx + \frac{\pi}{2}) + b}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \sin(c+dx + \frac{\pi}{2}))} dx}{a} \right) \\
& \quad \downarrow 3538 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\int \sqrt{\cos(c+dx)} dx - \frac{\int -\frac{b^2}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b}}{a} \right) \\
& \quad \downarrow 25 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{\frac{\int \frac{b^2}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b} + \int \sqrt{\cos(c+dx)} dx}{a} \right) \\
& \quad \downarrow 27 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx + \int \sqrt{\cos(c+dx)} dx}{a} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \sin(c+dx + \frac{\pi}{2}))} dx + \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx}{a} \right) \\
& \quad \downarrow 3119 \\
& B \left(\frac{2 \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} - \frac{b \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \sin(c+dx + \frac{\pi}{2}))} dx + \frac{2E(\frac{1}{2}(c+dx)|2)}{d}}{a} \right) \\
& \quad \downarrow 3284
\end{aligned}$$

3.393. $\int \frac{aB + bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

$$B \left(\frac{2 \sin(c + dx)}{ad\sqrt{\cos(c + dx)}} - \frac{\frac{2b \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{d(a+b)} + \frac{2E\left(\frac{1}{2}(c+dx)|2\right)}{d}}{a} \right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^2),x]`

output `B*(-((2*EllipticE[(c + d*x)/2, 2])/d + (2*b*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/((a + b)*d))/a) + (2*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]])`

3.393.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

```
rule 3281 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Sin[e + f*
x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2
))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x
])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) + b^2*d*(m + n
+ 2) - (b^2*c + b*(b*c - a*d)*(m + 1))*Sin[e + f*x] - b^2*d*(m + n + 3)*Si
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*
d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && IntegersQ[
2*m, 2*n] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*
n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])), x_Symbol] := Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x]
, x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[
e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0
] && NeQ[c^2 - d^2, 0]
```

3.393.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. $2(130) = 260$.

Time = 4.02 (sec) , antiderivative size = 355, normalized size of antiderivative = 4.44

method	result
default	$-\frac{2B\left(-2\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}(a-b)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{\frac{1}{2}-\frac{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}{2}}\right)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2}$

```
input int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^2,x,method=_RET
URNVERBOSE)
```

$$3.393. \quad \int \frac{aB+bB\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^2} dx$$

output
$$-2*B*(-2*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(a-b)*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^2+(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})*a-(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b-b*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticPi(\cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)})/a/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(a-b)/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$$

3.393.5 Fracas [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algo rithm="fricas")`

output Timed out

3.393.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c))**2,x)`

output Timed out

3.393.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorith="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2)), x)`

3.393.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorith="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(3/2)), x)`

3.393.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx)^{3/2} (a + b \cos(c + dx))^2} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^2), x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^2), x)`

3.393. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$

3.394
$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

3.394.1 Optimal result 3638
 3.394.2 Mathematica [A] (verified) 3639
 3.394.3 Rubi [A] (verified) 3639
 3.394.4 Maple [B] (verified) 3644
 3.394.5 Fracas [F(-1)] 3645
 3.394.6 Sympy [F(-1)] 3645
 3.394.7 Maxima [F] 3646
 3.394.8 Giac [F] 3646
 3.394.9 Mupad [F(-1)] 3646

3.394.1 Optimal result

Integrand size = 36, antiderivative size = 133

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \frac{2bBE(\frac{1}{2}(c + dx) | 2)}{a^2d} + \frac{2B \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{3ad}$$

$$+ \frac{2b^2B \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2)}{a^2(a + b)d}$$

$$+ \frac{2B \sin(c + dx)}{3ad \cos^{\frac{3}{2}}(c + dx)} - \frac{2bB \sin(c + dx)}{a^2d \sqrt{\cos(c + dx)}}$$

```
output 2*b*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))/a^2/d+2/3*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))/a/d+2*b^2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))/a^2/(a+b)/d+2/3*B*sin(d*x+c)/a/d/cos(d*x+c)^(3/2)-2*b*B*sin(d*x+c)/a^2/d/cos(d*x+c)^(1/2)
```

3.394.2 Mathematica [A] (verified)

Time = 2.61 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.59

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$= B \left(\frac{2(2a^2 + 9b^2) \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} + 8a \left(2 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{a+b} \right) + \frac{4(a-3b \cos(c + dx)) \operatorname{Sin}[c + dx]}{\cos^{\frac{3}{2}}(c + dx)} \right)$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2),x]`

output `(B*((2*(2*a^2 + 9*b^2)*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b) + 8*a*(2*EllipticF[(c + d*x)/2, 2] - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)) + (4*(a - 3*b*Cos[c + d*x])*Sin[c + d*x])/Cos[c + d*x]^(3/2) + (6*(-2*a*b*EllipticE[ArcSin[Sqrt[Cos[c + d*x]]], -1] + 2*a*(a + b)*EllipticF[ArcSin[Sqrt[Cos[c + d*x]]], -1] + (-2*a^2 + b^2)*EllipticPi[-(b/a), ArcSin[Sqrt[Cos[c + d*x]]], -1]*Sin[c + d*x])/(a*Sqrt[Sin[c + d*x]^2])))/(6*a^2*d)`

3.394.3 Rubi [A] (verified)

Time = 1.28 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.05, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {2011, 3042, 3281, 27, 3042, 3534, 27, 3042, 3538, 25, 3042, 3119, 3481, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{\frac{5}{2}}(a + b \sin(c + dx + \frac{\pi}{2}))} dx$$

3.394. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$

$$\begin{aligned}
& \downarrow \text{3281} \\
& B \left(\frac{2 \int \frac{-b \cos^2(c+dx) - a \cos(c+dx) + 3b}{2 \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx}{3a} + \frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow \text{27} \\
& B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-b \cos^2(c+dx) - a \cos(c+dx) + 3b}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))} dx}{3a} \right) \\
& \downarrow \text{3042} \\
& B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-b \sin(c+dx+\frac{\pi}{2})^2 - a \sin(c+dx+\frac{\pi}{2}) + 3b}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{3a} \right) \\
& \downarrow \text{3534} \\
& B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2 \int \frac{-a^2+4b \cos(c+dx)a+3b^2+3b^2 \cos^2(c+dx)}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} + \frac{6b \sin(c+dx)}{ad\sqrt{\cos(c+dx)}} \right) \\
& \downarrow \text{27} \\
& B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad\sqrt{\cos(c+dx)}}}{3a} - \frac{\int \frac{a^2+4b \cos(c+dx)a+3b^2+3b^2 \cos^2(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a} \right) \\
& \downarrow \text{3042} \\
& B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad\sqrt{\cos(c+dx)}}}{3a} - \frac{\int \frac{a^2+4b \sin(c+dx+\frac{\pi}{2})a+3b^2+3b^2 \sin^2(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{a} \right) \\
& \downarrow \text{3538} \\
& B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad\sqrt{\cos(c+dx)}}}{3a} - \frac{3b \int \sqrt{\cos(c+dx)} dx - \frac{\int \frac{a \cos(c+dx)b^2+(a^2+3b^2)b}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{a}}{3a} \right)
\end{aligned}$$

3.394. $\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$

$$\downarrow 25$$

$$B \left(\frac{\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{\int \frac{a \cos(c+dx)b^2 + (a^2+3b^2)b}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b}}{a} + 3b \int \sqrt{\cos(c+dx)} dx}{3a} \right)$$

$$\downarrow 3042$$

$$B \left(\frac{\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{\int \frac{a \sin(c+dx+\frac{\pi}{2})b^2 + (a^2+3b^2)b}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b}}{a} + 3b \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{3a} \right)$$

$$\downarrow 3119$$

$$B \left(\frac{\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{\int \frac{a \sin(c+dx+\frac{\pi}{2})b^2 + (a^2+3b^2)b}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx}{b}}{a} + \frac{6bE(\frac{1}{2}(c+dx)|2)}{d}}{3a} \right)$$

$$\downarrow 3481$$

$$B \left(\frac{\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{3b^3 \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx + ab \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b}}{a} + \frac{6bE(\frac{1}{2}(c+dx)|2)}{d}}{3a} \right)$$

$$\downarrow 3042$$

$$B \left(\frac{\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{3b^3 \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx + ab \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b}}{a} + \frac{6bE(\frac{1}{2}(c+dx)|2)}{d}}{3a} \right)$$

$$\downarrow 3120$$

$$B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{3b^3 \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx + \frac{2ab \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}}{b} + \frac{6bE(\frac{1}{2}(c+dx)|2)}{d}}{a} \right)$$

↓ 3284

$$B \left(\frac{2 \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{6b \sin(c+dx)}{ad \sqrt{\cos(c+dx)}} - \frac{\frac{6b^3 \operatorname{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} + \frac{2ab \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}}{b} + \frac{6bE(\frac{1}{2}(c+dx)|2)}{d}}{a} \right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^2), x]`

output `B*((2*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)) - (-(((6*b*EllipticE[(c + d*x)/2, 2])/d + ((2*a*b*EllipticF[(c + d*x)/2, 2])/d + (6*b^3*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(a + b)*d)/b)/a) + (6*b*Sin[c + d*x])/(a*d*Sqrt[Cos[c + d*x]]))/(3*a))`

3.394.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3281 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) + b^2*d*(m + n + 2) - (b^2*c + b*(b*c - a*d)*(m + 1))*Sin[e + f*x] - b^2*d*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && IntegersQ[2*m, 2*n] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3481 `Int((((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_))/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x] *(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int [(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))
```

```
rule 3538 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[C/(b*d) Int[Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[1/(b*d) Int[Simp[a*c*C - A*b*d + (b*c*C - b*B*d + a*C*d)*Sin[e + f*x], x]/(Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.394.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 424 vs. 2(203) = 406.

Time = 6.31 (sec) , antiderivative size = 425, normalized size of antiderivative = 3.20

method	result
default	$2\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) B \left(\frac{-\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}{6\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^2} + \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}} \right) / a$

```
input int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^2,x,method=_RET URNVERBOSE)
```

$$3.394. \int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^2} dx$$

output
$$\begin{aligned} & -2*(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*B*(1/a*(-1/6* \\ & \cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(c \\ & \cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d* \\ & x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*E \\ & \text{llipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))-1/a^2*b/\sin(1/2*d*x+1/2*c)^2/(2*\sin(\\ & 1/2*d*x+1/2*c)^2-1)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\\ & 2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2* \\ & \sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))-2*b^3 \\ & /a^2/(-2*a*b+2*b^2)*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+ \\ & 1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\text{EllipticPi}(c \\ & \cos(1/2*d*x+1/2*c),-2*b/(a-b),2^{(1/2)})/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1 \\ & /2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.394.5 Fricas [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{5/2}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algo
rithm="fricas")`

output Timed out

3.394.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{5/2}(c + dx)(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c))**2,x)`

output Timed out

3.394.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorith="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2)), x)`

3.394.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^2 \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorith="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^2*cos(d*x + c)^(5/2)), x)`

3.394.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx)^{5/2} (a + b \cos(c + dx))^2} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^2), x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^2), x)`

3.394. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$

3.395 $\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.395.1 Optimal result	3647
3.395.2 Mathematica [C] (warning: unable to verify)	3648
3.395.3 Rubi [A] (verified)	3649
3.395.4 Maple [B] (warning: unable to verify)	3656
3.395.5 Fricas [F(-1)]	3657
3.395.6 Sympy [F(-1)]	3657
3.395.7 Maxima [F]	3657
3.395.8 Giac [F]	3658
3.395.9 Mupad [F(-1)]	3658

3.395.1 Optimal result

Integrand size = 35, antiderivative size = 560

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx =$$

$$- \frac{(a - b) \sqrt{a + b} (6aAb - 3a^2B + 16b^2B) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a+b}}{24ab^2d}$$

$$+ \frac{\sqrt{a + b} (a + 2b) (6Ab - 3aB + 8bB) \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a+b}}{24b^2d}$$

$$+ \frac{\sqrt{a + b} (2a^2Ab - 8Ab^3 - a^3B - 4ab^2B) \cot(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{a+b}}{8b^3d}$$

$$+ \frac{(6aAb - 3a^2B + 16b^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{24b^2d \sqrt{\cos(c + dx)}}$$

$$+ \frac{(2Ab - aB) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{4bd}$$

$$+ \frac{B \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2} \sin(c + dx)}{3bd}$$

output $\frac{1}{3}B(a+b\cos(dx+c))^{3/2}\sin(dx+c)\cos(dx+c)^{1/2}/b/d+1/24(6Aab-3Ba^2+16Bb^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^2/d/\cos(dx+c)^{1/2}+1/4(2Ab-Ba)\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/b/d-1/24(a-b)(6Aab-3Ba^2+16Bb^2)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/b^2/d+1/24(a+2b)(6Ab-3Ba+8Bb)\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d+1/8(2Aa^2b-8Ab^3-Ba^3-4Bab^2)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^3/d$

3.395.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.56 (sec) , antiderivative size = 1224, normalized size of antiderivative = 2.19

$$\int \cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))dx = \text{Too large to display}$$

input `Integrate[Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output

```

-1/48*((-4*a*(-18*a*A*b + a^2*B - 16*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]
^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((
a + b*cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[S
qrt[((a + b*cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)
]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]
) - 4*a*(-24*A*b^2 - 28*a*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b
)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[(a + b*cos[c
+ d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b
*cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c +
d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) - (Sqrt[(
(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c
+ d*x)/2]^2)/a)]*Sqrt[((a + b*cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c +
d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*cos[c + d*x])*Csc[(c + d*x)/2
]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x
]]*Sqrt[a + b*cos[c + d*x]])) + 2*(-6*a*A*b + 3*a^2*B - 16*b^2*B)*((I*cos[
(c + d*x)/2]*Sqrt[a + b*cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]
/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)
/2]^2*Sec[c + d*x]]*Sqrt[((a + b*cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (
2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c
+ d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*cos[c + d*x])*Csc[(c + d*x)...

```

3.395.3 Rubi [A] (verified)

Time = 2.78 (sec) , antiderivative size = 563, normalized size of antiderivative = 1.01, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3469}
 \end{aligned}$$

3.395. $\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

$$\begin{aligned}
 & \frac{\int \frac{\sqrt{a+b \cos(c+dx)} (3(2Ab-aB) \cos^2(c+dx)+4bB \cos(c+dx)+aB) dx}{2\sqrt{\cos(c+dx)}}}{\frac{3b}{3bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}} + \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{\sqrt{a+b \cos(c+dx)} (3(2Ab-aB) \cos^2(c+dx)+4bB \cos(c+dx)+aB) dx}{\sqrt{\cos(c+dx)}}}{\frac{6b}{3bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}} + \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (3(2Ab-aB) \sin(c+dx+\frac{\pi}{2})^2+4bB \sin(c+dx+\frac{\pi}{2})+aB) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{\frac{6b}{3bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}} + \\
 & \quad \downarrow 3528 \\
 & \frac{\frac{1}{2} \int \frac{(-3Ba^2+6Aba+16b^2B) \cos^2(c+dx)+2b(6Ab+7aB) \cos(c+dx)+a(6Ab+aB)}{2\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx + \frac{3(2Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d}}{\frac{6b}{3bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}} + \\
 & \quad \downarrow 27 \\
 & \frac{\frac{1}{4} \int \frac{(-3Ba^2+6Aba+16b^2B) \cos^2(c+dx)+2b(6Ab+7aB) \cos(c+dx)+a(6Ab+aB)}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx + \frac{3(2Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d}}{\frac{6b}{3bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}} + \\
 & \quad \downarrow 3042 \\
 & \frac{\frac{1}{4} \int \frac{(-3Ba^2+6Aba+16b^2B) \sin(c+dx+\frac{\pi}{2})^2+2b(6Ab+7aB) \sin(c+dx+\frac{\pi}{2})+a(6Ab+aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{3(2Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d}}{\frac{6b}{3bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}} + \\
 & \quad \downarrow 3540
 \end{aligned}$$

3.395. $\int \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx$

$$\frac{1}{4} \left(\frac{\int -\frac{3(-Ba^3+2Aba^2-4b^2Ba-8Ab^3)\cos^2(c+dx)-2ab(6Ab+aB)\cos(c+dx)+a(-3Ba^2+6Aba+16b^2B)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} + \frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} \right)$$

$$\frac{B \sin(c+dx)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}}{3bd} \quad 6b$$

↓ 25

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{3(-Ba^3+2Aba^2-4b^2Ba-8Ab^3)\cos^2(c+dx)-2ab(6Ab+aB)\cos(c+dx)+a(-3Ba^2+6Aba+16b^2B)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} \right)$$

$$\frac{B \sin(c+dx)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}}{3bd} \quad 6b$$

↓ 3042

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{3(-Ba^3+2Aba^2-4b^2Ba-8Ab^3)\sin(c+dx+\frac{\pi}{2})^2-2ab(6Ab+aB)\sin(c+dx+\frac{\pi}{2})+a(-3Ba^2+6Aba+16b^2B)\cos^2(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} \right)$$

$$\frac{B \sin(c+dx)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}}{3bd} \quad 6b$$

↓ 3532

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-3Ba^2+6Aba+16b^2B)-2ab(6Ab+aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + 3(a^3(-B)+2a^2Ab-4ab^2B-8a^2b^2)}{2b} \right)$$

$$\frac{B \sin(c+dx)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}}{3bd} \quad 6b$$

↓ 3042

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-3Ba^2+6Aba+16b^2B)-2ab(6Ab+aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + 3(a^3(-B)+2a^2Ab-4ab^2B-8a^2b^2)}{2b} \right)$$

$$\frac{B \sin(c+dx)\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}}{3bd} \quad 6b$$

3.395. $\int \cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx)) dx$

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-3Ba^2+6Aba+16b^2B) - 2ab(6Ab+aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{6\sqrt{a+b}(a^3(-B)+2a^2Ab-4ab)}{a^3(-B)+2a^2Ab-4ab} \right)$$

↓ 3288

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}$$

↓ 3477

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-3a^2B+6aAb+16b^2B) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - a(a+2b)(-3aB+6aAb+8b^2)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}$$

↓ 3042

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-3a^2B+6aAb+16b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - a(a+2b)(-3aB+6aAb+8b^2)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}$$

↓ 3295

$$\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-3a^2B+6aAb+16b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{6\sqrt{a+b}(a^3(-B)+2a^2Ab-4ab)}{a^3(-B)+2a^2Ab-4ab} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}$$

↓ 3473

$$\frac{1}{4} \left(\frac{(-3a^2B + 6aAb + 16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{2(a-b)\sqrt{a+b}(-3a^2B + 6aAb + 16b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E(\arcsin(\dots))}{ad} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3bd}$$

```
input Int[Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]
```

```
output (B*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*b*d) + (
(3*(2*A*b - a*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])
/(2*d) + (-1/2*((2*(a - b)*Sqrt[a + b]*(6*a*A*b - 3*a^2*B + 16*b^2*B)*Cot[
c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c
+ d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[
(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*Sqrt[a + b]*(a + 2*b)*(6*A*b -
3*a*B + 8*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sq
rt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d
*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (6*Sqrt[a + b]*(2
*a^2*A*b - 8*A*b^3 - a^3*B - 4*a*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b,
ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a +
b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*
x]))/(a - b))]/(b*d))/b + ((6*a*A*b - 3*a^2*B + 16*b^2*B)*Sqrt[a + b*Cos[c
+ d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/4)/(6*b)
```

3.395.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] :> Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.395.5 Fricas [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `Timed out`

3.395.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.395.7 Maxima [F]

$$\begin{aligned} & \int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(3/2), x)`

3.395.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(3/2)
, x)`

3.395.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$= \int \cos(c + dx)^{3/2} (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)} dx$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2),x)`

output `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`

3.396 $\int \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

3.396.1 Optimal result	3659
3.396.2 Mathematica [C] (verified)	3660
3.396.3 Rubi [A] (verified)	3661
3.396.4 Maple [B] (verified)	3666
3.396.5 Fricas [F]	3667
3.396.6 Sympy [F]	3668
3.396.7 Maxima [F]	3668
3.396.8 Giac [F]	3668
3.396.9 Mupad [F(-1)]	3669

3.396.1 Optimal result

Integrand size = 35, antiderivative size = 473

$$\int \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx =$$

$$\frac{(a - b)\sqrt{a + b}(4Ab + aB) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right) \mid -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{4abd}$$

$$+ \frac{\sqrt{a + b}(4Ab + (a + 2b)B) \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{4bd}$$

$$- \frac{\sqrt{a + b}(4aAb - a^2B + 4b^2B) \cot(c + dx) \operatorname{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}}}{4b^2d}$$

$$+ \frac{(4Ab + aB)\sqrt{a + b \cos(c + dx)} \sin(c + dx)}{4bd\sqrt{\cos(c + dx)}}$$

$$+ \frac{B\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{2d}$$

output $\frac{1}{4}(4A^2b + B^2a) \sin(dx+c) (a+b \cos(dx+c))^{1/2} / b/d \cos(dx+c)^{1/2} + 1/2$
 $B \sin(dx+c) \cos(dx+c)^{1/2} (a+b \cos(dx+c))^{1/2} / d - 1/4(a-b) (4A^2b + B^2a) \cot(dx+c) \text{EllipticE}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2})$
 $, ((-a-b)/(a-b))^{1/2} (a+b)^{1/2} (a(1-\sec(dx+c)) / (a+b))^{1/2} (a(1+\sec(dx+c)) / (a-b))^{1/2} / a/b/d + 1/4(4A^2b + (a+2b)B) \cot(dx+c) \text{EllipticF}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2})$
 $, ((-a-b)/(a-b))^{1/2} (a+b)^{1/2} (a(1-\sec(dx+c)) / (a+b))^{1/2} (a(1+\sec(dx+c)) / (a-b))^{1/2} / b/d - 1/4(4A^2a^2b - B^2a^2 + 4B^2b^2) \cot(dx+c) \text{EllipticPi}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2})$
 $, (a+b)/b, ((-a-b)/(a-b))^{1/2} (a+b)^{1/2} (a(1-\sec(dx+c)) / (a+b))^{1/2} (a(1+\sec(dx+c)) / (a-b))^{1/2} / b^2/d$

3.396.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 19.79 (sec) , antiderivative size = 1175, normalized size of antiderivative = 2.48

$$\int \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} (A+B \cos(c+dx)) dx = \text{Too large to display}$$

input `Integrate[Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]`

output

```
(B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + ((-4*a*(4*A*b + 3*a*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(8*a*A + 4*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(4*A*b + a*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[...
```

3.396.3 Rubi [A] (verified)

Time = 2.06 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3482, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3482}$$

$$\frac{1}{4} \int \frac{(4Ab + aB) \cos^2(c + dx) + 2(2aA + bB) \cos(c + dx) + aB}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

3.396. $\int \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{4} \int \frac{(4Ab + aB) \sin(c + dx + \frac{\pi}{2})^2 + 2(2aA + bB) \sin(c + dx + \frac{\pi}{2}) + aB}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}} dx + \\
& \downarrow 3540 \\
& \frac{1}{4} \left(\frac{\int -\frac{((-Ba^2 + 4Aba + 4b^2B) \cos^2(c + dx)) - 2abB \cos(c + dx) + a(4Ab + aB)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{2b} + \frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} \right) \\
& \quad \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\
& \downarrow 25 \\
& \frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{\int \frac{((-Ba^2 + 4Aba + 4b^2B) \cos^2(c + dx)) - 2abB \cos(c + dx) + a(4Ab + aB)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{2b} \right) + \\
& \quad \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\
& \downarrow 3042 \\
& \frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{\int \frac{(Ba^2 - 4Aba - 4b^2B) \sin(c + dx + \frac{\pi}{2})^2 - 2abB \sin(c + dx + \frac{\pi}{2}) + a(4Ab + aB)}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2b} \right) \\
& \quad \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\
& \downarrow 3532 \\
& \frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{\int \frac{a(4Ab + aB) - 2abB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - (a^2(-B) + 4aAb + 4b^2B) \int \frac{\sqrt{a + b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{2b} \right) \\
& \quad \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\
& \downarrow 3042
\end{aligned}$$

$$\frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{\int \frac{a(4Ab+aB)-2abB \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - (a^2(-B) + 4aAb + 4b^2B)}{2b} \right)$$

$$\frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3288

$$\frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{\int \frac{a(4Ab+aB)-2abB \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(a^2(-B)+4aAb+4b^2B)}{2b} \right)$$

$$\frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3477

$$\frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{a(aB + 4Ab) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - a(B(a + 2b) + 4Ab)}{2b} \right)$$

$$\frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{-a(B(a + 2b) + 4Ab) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + a(aB + 4Ab)}{2b} \right)$$

$$\frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3295

$$\frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{a(aB + 4Ab) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(a^2(-B) + 4aAb + 4b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticE}[\text{ArcSin}[\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}], -\frac{a+b}{a-b}]}{bd} \right)$$

$$\frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3473

$$\frac{1}{4} \left(\frac{(aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} - \frac{2\sqrt{a+b}(a^2(-B) + 4aAb + 4b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticE}[\text{ArcSin}[\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}], -\frac{a+b}{a-b}]}{bd} \right)$$

$$\frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

```
input Int[Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]),x]
```

```
output (B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2 * ((2*(a - b)*Sqrt[a + b]*(4*A*b + a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]) * Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/(a*d) - (2*Sqrt[a + b]*(4*A*b + (a + 2*b)*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)])*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d + (2*Sqrt[a + b]*(4*a*A*b - a^2*B + 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b*d))/b + ((4*A*b + a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]])/4
```

3.396.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

3.396. $\int \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`


```
rule 3482 Int[Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[-2*B*Cos[e + f*x]*Sqrt[a + b*Sin[e + f*x]]*((c + d*Sin[e + f*x])^n/(f*(2*
n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*Sin[e + f*x])^(n - 1)/Sqrt[a
+ b*Sin[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d
)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*Sin[e + f*x] + (A*b*d*(2*n + 3) + B*
(a*d + 2*b*c*n))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A,
B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Eq
Q[n^2, 1/4]
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.396.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2771 vs. $2(431) = 862$.

Time = 9.72 (sec) , antiderivative size = 2772, normalized size of antiderivative = 5.86

method	result	size
parts	Expression too large to display	2772
default	Expression too large to display	2794

```
input int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c)),x,method=_RET
URNVERBOSE)
```

$$3.396. \quad \int \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

output `A/d*(-EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)^2-EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b*cos(d*x+c)^2-2*EllipticPi(cot(d*x+c)-csc(d*x+c), -1, (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)^2+2*EllipticF(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)^2-2*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)-2*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b*cos(d*x+c)-4*EllipticPi(cot(d*x+c)-csc(d*x+c), -1, (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)+4*EllipticF(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)+b*cos(d*x+c)^2*sin(d*x+c)-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b)...`

3.396.5 Fracas [F]

$$\int \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx$$

$$= \int (B\cos(dx+c) + A) \sqrt{b\cos(dx+c) + a} \sqrt{\cos(dx+c)} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)), x, algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.396.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx \\ &= \int (A+B\cos(c+dx)) \sqrt{a+b\cos(c+dx)} \sqrt{\cos(c+dx)} dx \end{aligned}$$

input `integrate(cos(d*x+c)**(1/2)*(a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c)),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*sqrt(cos(c + d*x)), x)`

3.396.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx \\ &= \int (B\cos(dx+c) + A) \sqrt{b\cos(dx+c) + a} \sqrt{\cos(dx+c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.396.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx \\ &= \int (B\cos(dx+c) + A) \sqrt{b\cos(dx+c) + a} \sqrt{\cos(dx+c)} dx \end{aligned}$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c)), x)`

3.396.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx$$

$$= \int \sqrt{\cos(c+dx)} (A+B\cos(c+dx)) \sqrt{a+b\cos(c+dx)} dx$$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`output `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2), x)`

$$3.397 \quad \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.397.1 Optimal result	3670
3.397.2 Mathematica [A] (verified)	3671
3.397.3 Rubi [A] (verified)	3671
3.397.4 Maple [B] (verified)	3676
3.397.5 Fricas [F(-1)]	3677
3.397.6 Sympy [F]	3677
3.397.7 Maxima [F]	3677
3.397.8 Giac [F]	3678
3.397.9 Mupad [F(-1)]	3678

3.397.1 Optimal result

Integrand size = 35, antiderivative size = 385

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}B \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ad}$$

$$+ \frac{\sqrt{a+b}(2A+B) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{d}$$

$$- \frac{\sqrt{a+b}(2Ab+aB) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{bd}$$

$$+ \frac{B\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}}$$

```
output B*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)-(a-b)*B*cot(d*x+c)*
EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b
))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a
-b))^(1/2)/a/d+(2*A+B)*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1
/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(
a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d-(2*A*b+B*a)*cot(d*x+c)*Ellipt
icPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(
a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/
(a-b))^(1/2)/b/d
```

$$3.397. \quad \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.397.2 Mathematica [A] (verified)

Time = 7.63 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \frac{\sqrt{\cos(c + dx)} \left(2(a + b)B \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} E\left(\arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right) \middle| \frac{-a + b}{a + b}\right) - 4(Ab + a(-A + B)) \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} \right)}{(a + b)(1 + \cos(c + dx))}$$

input `Integrate[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(Sqrt[Cos[c + d*x]]*(2*(a + b)*B*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 4*(A*b + a*(-A + B))*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 8*A*b*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 4*a*B*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + b*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sin[(3*(c + d*x))/2] + 2*a*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Tan[(c + d*x)/2] - b*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Tan[(c + d*x)/2]))/(2*d*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[a + b*Cos[c + d*x]])`

3.397.3 Rubi [A] (verified)Time = 1.57 (sec) , antiderivative size = 391, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3482, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

3.397. $\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3482} \\
& \frac{1}{2} \int -\frac{((2Ab + aB) \cos^2(c + dx)) - 2aA \cos(c + dx) + aB}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{25} \\
& \frac{B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{1}{2} \int \frac{((2Ab + aB) \cos^2(c + dx)) - 2aA \cos(c + dx) + aB}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \\
& \frac{1}{2} \int \frac{(-2Ab - aB) \sin(c + dx + \frac{\pi}{2})^2 - 2aA \sin(c + dx + \frac{\pi}{2}) + aB}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3532} \\
& \frac{1}{2} \left((aB + 2Ab) \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx - \int \frac{aB - 2aA \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \right) + \\
& \quad \frac{B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left((aB + 2Ab) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \int \frac{aB - 2aA \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3288} \\
& \frac{1}{2} \left(- \int \frac{aB - 2aA \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{2\sqrt{a + b}(aB + 2Ab) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) - 1)}{a + b}}}{d \sqrt{\cos(c + dx)}} \right) + \\
& \quad \frac{B \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \\
& \quad \downarrow \text{3477}
\end{aligned}$$

3.397. $\int \frac{\sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$

$$\frac{1}{2} \left(a(2A + B) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx - aB \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - \frac{2\sqrt{a + b}(aB)}{d\sqrt{\cos(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{2} \left(a(2A + B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - aB \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin^{\frac{3}{2}}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{2\sqrt{a + b}(aB)}{d\sqrt{\cos(c + dx)}} \right)$$

↓ 3295

$$\frac{1}{2} \left(-aB \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin^{\frac{3}{2}}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2\sqrt{a + b}(2A + B) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}}}{d\sqrt{\cos(c + dx)}} - \frac{2\sqrt{a + b}(aB)}{d\sqrt{\cos(c + dx)}} \right)$$

↓ 3473

$$\frac{1}{2} \left(\frac{2\sqrt{a + b}(2A + B) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} \text{EllipticF} \left(\arcsin \left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}} \right), -\frac{a + b}{a - b} \right) - \frac{2\sqrt{a + b}(aB)}{d\sqrt{\cos(c + dx)}} \right)$$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`


```
output ((-2*(a - b)*Sqrt[a + b]*B*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d) + (2 *Sqrt[a + b]*(2*A + B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d - (2*Sqrt[a + b]*(2*A*b + a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b*d)) /2 + (B*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])
```

3.397.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[-2A(c-d) \tan(e+fx)/(fbc^2) \text{Rt}[(c+d)/b, 2] \sqrt{c(1+\csc(e+fx))/(c-d)} \sqrt{c(1-\csc(e+fx))/(c+d)} \text{EllipticE}[\text{ArcSin}[\sqrt{c+d \sin(e+fx)}/\sqrt{b \sin(e+fx)}] \text{Rt}[(c+d)/b, 2], -(c+d)/(c-d)], x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\& \text{PosQ}[(c+d)/b]$$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[(A-B)/(a-b) \int [1/(\sqrt{a+b \sin(e+fx)} \sqrt{c+d \sin(e+fx)})], x], x] - \text{Simp}[(A*b - a*B)/(a-b) \int [(1 + \sin(e+fx))/(a+b \sin(e+fx))^{3/2} \sqrt{c+d \sin(e+fx)}], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[A, B]$$`

rule 3482 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) +

$$\int \sqrt{a + b \sin(e + fx)} (A + B \sin(e + fx)) (c + d \sin(e + fx))^n dx$$

$$\rightarrow \text{Simp}[-2B \cos(e + fx) \sqrt{a + b \sin(e + fx)} (c + d \sin(e + fx))^n / (f(2n + 3)), x] + \text{Simp}[1/(2n + 3) \int ((c + d \sin(e + fx))^{n-1} / \sqrt{a + b \sin(e + fx)}) * \text{Simp}[aA*(2n + 3) + B(b*c + 2a*d*n) + (B(a*c + b*d)*(2n + 1) + A(b*c + a*d)*(2n + 3))*\sin(e + fx) + (A*b*d*(2n + 3) + B(a*d + 2*b*c*n))*\sin(e + fx)^2], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[n^2, 1/4]$$`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2)/

$$\int \frac{(A + B \sin(e + fx) + C \sin^2(e + fx)) \sqrt{c + d \sin(e + fx)}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[C/b^2 \int [\sqrt{a + b \sin(e + fx)} / \sqrt{c + d \sin(e + fx)}], x], x] + \text{Simp}[1/b^2 \int [(A*b^2 - a^2*C + b*(b*B - 2*a*C)*\sin(e + fx)) / ((a + b \sin(e + fx))^{3/2} \sqrt{c + d \sin(e + fx)})], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$$`

3.397.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1267 vs. $2(357) = 714$.

Time = 11.87 (sec) , antiderivative size = 1268, normalized size of antiderivative = 3.29

method	result	size
parts	Expression too large to display	1268
default	Expression too large to display	1702

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2*A/d*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a-EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+2*b*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))/(a+cos(d*x+c)*b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(1+cos(d*x+c))/cos(d*x+c)^(1/2)+B/d*(-EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2-EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*b*cos(d*x+c)^2-2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2-2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)-2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*b*cos(d*x+c)-4*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)+4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(...
```

3.397.5 Fracas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algo
rithm="fricas")
```

```
output Timed out
```

3.397.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx \end{aligned}$$

```
input integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)
```

```
output Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))/sqrt(cos(c + d*x)),
x)
```

3.397.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ &= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\sqrt{\cos(dx + c)}} dx \end{aligned}$$

```
input integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algo
rithm="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sqrt(cos(d*x + c))
, x)
```

3.397. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

3.397.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.397.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2),x)`

$$3.398 \quad \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.398.1 Optimal result	3679
3.398.2 Mathematica [A] (verified)	3680
3.398.3 Rubi [A] (verified)	3680
3.398.4 Maple [B] (warning: unable to verify)	3683
3.398.5 Fricas [F]	3684
3.398.6 Sympy [F]	3685
3.398.7 Maxima [F]	3685
3.398.8 Giac [F]	3685
3.398.9 Mupad [F(-1)]	3686

3.398.1 Optimal result

Integrand size = 35, antiderivative size = 351

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{2A(a-b)\sqrt{a+b} \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \mid -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ad}$$

$$+ \frac{2\sqrt{a+b}(Ab-a(A-B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ad}$$

$$- \frac{2\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{d}$$

```
output 2*A*(a-b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d+2*(A*b-a*(A-B))*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d-2*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d
```

$$3.398. \quad \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.398.2 Mathematica [A] (verified)

Time = 8.40 (sec) , antiderivative size = 273, normalized size of antiderivative = 0.78

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{-2A(a + b)\sqrt{1 + \cos(c + dx)}\sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}}E\left(\arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right)\left|\frac{-a + b}{a + b}\right.\right) + 2(b(A - B) + a(A - B))\sqrt{1 + \cos(c + dx)}}{d}$$

input `Integrate[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(-2*A*(a + b)*Sqrt[1 + Cos[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*(b*(A - B) + a*(A + B))*Sqrt[1 + Cos[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 4*b*B*Sqrt[1 + Cos[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + (2*A*(a + b*Cos[c + d*x])*Tan[(c + d*x)/2])/Sqrt[Cos[c + d*x]])/(d*Sqrt[a + b*Cos[c + d*x]])`

3.398.3 Rubi [A] (verified)Time = 1.20 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3470, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2}} dx$$

$$\downarrow \text{3470}$$

3.398. $\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
& \int \frac{aA + (Ab + aB) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + bB \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{aA + (Ab + aB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + bB \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3288} \\
& \int \frac{aA + (Ab + aB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \\
& \frac{2B\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d} \\
& \quad \downarrow \text{3477} \\
& (Ab - a(A - B)) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx + aA \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - \\
& \frac{2B\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d} \\
& \quad \downarrow \text{3042} \\
& (Ab - a(A - B)) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \\
& aA \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \\
& \frac{2B\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d} \\
& \quad \downarrow \text{3295} \\
& aA \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{\frac{3}{2}} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \\
& \frac{2\sqrt{a+b}(Ab - a(A - B)) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{ad} \\
& \frac{2B\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d} \\
& \quad \downarrow \text{3473}
\end{aligned}$$

3.398. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\frac{2\sqrt{a+b}(Ab - a(A - B)) \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) + 2A(a-b)\sqrt{a+b} \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) + 2B\sqrt{a+b} \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d}$$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(2*A*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d) + (2*Sqrt[a + b]*(A*b - a*(A - B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d) - (2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d`

3.398.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3470 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])]/((b_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[B*(d/b^2) Int[Sqrt[b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Int[(A*c + (B*c + A*d)*Sin[e + f*x])/((b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]
```

3.398.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 952 vs. $2(327) = 654$.

Time = 14.28 (sec) , antiderivative size = 953, normalized size of antiderivative = 2.72

method	result	size
parts	Expression too large to display	953
default	Expression too large to display	1161

$$3.398. \quad \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

input `int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `-2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+csc(d*x+c)^3*(1-cos(d*x+c))^3*a-csc(d*x+c)^3*(1-cos(d*x+c))^3*b+a*(csc(d*x+c)-cot(d*x+c))+b*(csc(d*x+c)-cot(d*x+c)))*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(3/2)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)+2*B/d*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2))*(-EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a+EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b-2*b*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))/(a+cos(d*x+c)*b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(1+cos(d*x+c))...`

3.398.5 Fracas [F]

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

$$= \int \frac{(B \cos(dx+c)+A)\sqrt{b \cos(dx+c)+a}}{\cos(dx+c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,algorithm="fracas")`

output `integral((B*cos(d*x+c)+A)*sqrt(b*cos(d*x+c)+a)/cos(d*x+c)^(3/2),x)`

3.398. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

3.398.6 Sympy [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))/cos(c + d*x)**(3/2), x)`

3.398.7 Maxima [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`

3.398.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(3/2), x)`

3.398.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^{\frac{3}{2}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(3/2),x)`

3.399
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.399.1 Optimal result 3687
 3.399.2 Mathematica [A] (verified) 3688
 3.399.3 Rubi [A] (verified) 3688
 3.399.4 Maple [B] (verified) 3691
 3.399.5 Fracas [F] 3692
 3.399.6 Sympy [F] 3693
 3.399.7 Maxima [F] 3693
 3.399.8 Giac [F] 3693
 3.399.9 Mupad [F(-1)] 3694

3.399.1 Optimal result

Integrand size = 35, antiderivative size = 284

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{2(a-b)\sqrt{a+b}(Ab+3aB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a^2d}$$

$$+ \frac{2(a-b)\sqrt{a+b}(A-3B) \cot(c+dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3ad}$$

$$+ \frac{2A\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

output

```
2/3*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)+2/3*(a-b)*(A*b+
3*B*a)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(
1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*
(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d+2/3*(a-b)*(A-3*B)*cot(d*x+c)*EllipticF((
a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(
a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a
/d
```

3.399.2 Mathematica [A] (verified)

Time = 10.99 (sec) , antiderivative size = 407, normalized size of antiderivative = 1.43

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{4 \cos^2\left(\frac{1}{2}(c + dx)\right)^{5/2} \left(\frac{\cos(c+dx)}{1+\cos(c+dx)}\right)^{3/2} \sqrt{1 + \cos(c + dx)} \sqrt{\cos(c + dx) \sec^2\left(\frac{1}{2}(c + dx)\right)} \left(-2(a + b)(Ab + 3aB) + \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} \left(\frac{2 \sec(c+dx)(Ab \sin(c+dx) + 3aB \sin(c+dx))}{3a} + \frac{2}{3}A \sec(c + dx) \tan(c + dx)\right)}\right)}{d}$$

input `Integrate[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(4*(Cos[(c + d*x)/2]^2)^(5/2)*(Cos[c + d*x]/(1 + Cos[c + d*x]))^(3/2)*Sqrt[1 + Cos[c + d*x]]*Sqrt[Cos[c + d*x]*Sec[(c + d*x)/2]^2*(-2*(a + b)*(A*b + 3*a*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] * Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*a*(a + b)*(A + 3*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] * Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - (A*b + 3*a*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2]))/(3*a*d*Cos[c + d*x]^(5/2)*Sqrt[a + b*Cos[c + d*x]]) + (Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*((2*Sec[c + d*x]*(A*b*Sin[c + d*x] + 3*a*B*Sin[c + d*x]))/(3*a) + (2*A*Sec[c + d*x]*Tan[c + d*x])/3))/d`

3.399.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3478, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

3.399. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3478

$$\frac{2}{3} \int \frac{Ab + 3aB + (aA + 3bB) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{Ab + 3aB + (aA + 3bB) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{Ab + 3aB + (aA + 3bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3477

$$\frac{1}{3} \left((3aB + Ab) \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + (a - b)(A - 3B) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \right) + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left((a - b)(A - 3B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + (3aB + Ab) \int \frac{\sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3295

$$\frac{1}{3} \left((3aB + Ab) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2(a - b) \sqrt{a + b} (A - 3B) \cot(c + dx) \sqrt{a(1 - \sec^2(c + dx))}}{3d \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3473

$$\frac{1}{3} \left(\frac{2(a-b)\sqrt{a+b}(3aB+Ab)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)\Big|_{-\frac{a+b}{a-b}}}{a^2d} + \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)} \right)$$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `((2*(a - b)*Sqrt[a + b]*(A*b + 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(A - 3*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/3 + (2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2))`

3.399.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)]]*Sqrt[(a_.) + (b_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_
.)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

```
rule 3478 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*
x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a
+ b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*
(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*
Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

3.399.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1954 vs. $2(258) = 516$.

Time = 18.36 (sec) , antiderivative size = 1955, normalized size of antiderivative = 6.88

method	result	size
parts	Expression too large to display	1955
default	Expression too large to display	2195

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

$$3.399. \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

```

output -2/3*A/d*((cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c
)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*
cos(d*x+c)^3+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*
x+c)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a
*b*cos(d*x+c)^3-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos
(d*x+c)))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2)
)*a*b*cos(d*x+c)^3-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+
cos(d*x+c)))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1
/2))*b^2*cos(d*x+c)^3+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b
)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b
))^(1/2))*a^2*cos(d*x+c)^2+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b
))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-2*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/
(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1
+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((
a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c
),(-(a-b)/(a+b))^(1/2))*b^2*cos(d*x+c)^2+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2
))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d
*x+c),(-(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)+EllipticF(cot(d*x+c)-csc(d*x+c)
,(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(c...

```

3.399.5 Fracas [F]

$$\begin{aligned}
 & \int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\
 &= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{5}{2}}} dx
 \end{aligned}$$

```

input integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algo
rithm="fracas")

```

```

output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(5/2),
x)

```

3.399.6 Sympy [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))/cos(c + d*x)**(5/2), x)`

3.399.7 Maxima [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`

3.399.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`

3.399.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^{\frac{5}{2}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(5/2),x)`

3.400
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

3.400.1 Optimal result 3695
 3.400.2 Mathematica [C] (verified) 3696
 3.400.3 Rubi [A] (verified) 3696
 3.400.4 Maple [B] (verified) 3701
 3.400.5 Fracas [F] 3702
 3.400.6 Sympy [F(-1)] 3702
 3.400.7 Maxima [F] 3702
 3.400.8 Giac [F] 3703
 3.400.9 Mupad [F(-1)] 3703

3.400.1 Optimal result

Integrand size = 35, antiderivative size = 350

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{2(a-b)\sqrt{a+b}(9a^2A-2Ab^2+5abB) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a+b}}}{15a^3d} - \frac{2(a-b)\sqrt{a+b}(9aA+2Ab-5aB) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a+b}}}{15a^2d} + \frac{2A\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} + \frac{2(Ab+5aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15ad \cos^{\frac{3}{2}}(c+dx)}$$

output

```
2/5*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+2/15*(A*b+5*B*a)
*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d/cos(d*x+c)^(3/2)+2/15*(a-b)*(9*A*a
^2-2*A*b^2+5*B*a*b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)
)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+
b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d-2/15*(a-b)*(9*A*a+2*A*b-5*B
*a)*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/
2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+
sec(d*x+c))/(a-b))^(1/2)/a^2/d
```

3.400.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.55 (sec) , antiderivative size = 1315, normalized size of antiderivative = 3.76

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output

```
-1/15*((-4*a*(2*a^2*A*b - 2*A*b^3 - 5*a^3*B + 5*a*b^2*B)*Sqrt[((a + b)*Cot
[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2
)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*Ellip
ticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-
2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*
Cos[c + d*x]]) - 4*a*(9*a^3*A - 2*a*A*b^2 + 5*a^2*b*B)*((Sqrt[((a + b)*Cot
[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2
)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*Ellip
ticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-
2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*
Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a +
b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c
+ d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c
+ d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2
]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])) + 2*(9*a^2*A*b - 2*A
*b^3 + 5*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[
I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d
*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*S
ec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a +
b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*C...
```

3.400.3 Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3478, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.400. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}(A+B \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow \text{3478} \\
& \frac{2}{5} \int \frac{2Ab \cos^2(c+dx) + (3aA + 5bB) \cos(c+dx) + Ab + 5aB}{\frac{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}} dx + \\
& \quad \frac{5d \cos^{\frac{5}{2}}(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{2Ab \cos^2(c+dx) + (3aA + 5bB) \cos(c+dx) + Ab + 5aB}{\frac{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}} dx + \\
& \quad \frac{5d \cos^{\frac{5}{2}}(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{2Ab \sin(c+dx+\frac{\pi}{2})^2 + (3aA + 5bB) \sin(c+dx+\frac{\pi}{2}) + Ab + 5aB}{\frac{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}} dx + \\
& \quad \frac{5d \cos^{\frac{5}{2}}(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3534} \\
& \frac{1}{5} \left(\frac{2 \int \frac{9Aa^2+5bBa+(7Ab+5aB) \cos(c+dx)a-2Ab^2}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(5aB + Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} \right) + \\
& \quad \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \left(\frac{\int \frac{9Aa^2+5bBa+(7Ab+5aB) \cos(c+dx)a-2Ab^2}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(5aB + Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} \right) + \\
& \quad \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.400. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{\int \frac{9Aa^2+5bBa+(7Ab+5aB) \sin(c+dx+\frac{\pi}{2})a-2Ab^2}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a} + \frac{2(5aB + Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3ad \cos^{\frac{3}{2}}(c + dx)} \right) + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5d \cos^{\frac{5}{2}}(c + dx)} \downarrow \text{3477}$$

$$\frac{1}{5} \left(\frac{(9a^2A + 5abB - 2Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - (a - b)(9aA - 5aB + 2Ab) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3042}$$

$$\frac{1}{5} \left(\frac{(9a^2A + 5abB - 2Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - (a - b)(9aA - 5aB + 2Ab) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a} + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3295}$$

$$\frac{1}{5} \left(\frac{(9a^2A + 5abB - 2Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(9aA-5aB+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a}}{3a} + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \downarrow \text{3473}$$

$$\frac{1}{5} \left(\frac{2(a-b)\sqrt{a+b}(9a^2A+5abB-2Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right) - \frac{a+b}{a-b}}{a^2 d} - \frac{2(a-b)\sqrt{a+b}(9aA-5aB+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a}}{3a} + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

3.400. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(9*a^2*A - 2*A*b^2 + 5*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(9*a*A + 2*A*b - 5*a*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(3*a) + (2*(A*b + 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/5`

3.400.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3478 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

3.400.5 Fricas [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(7/2),
x)`

3.400.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.400.7 Maxima [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(7/2)
, x)`

3.400. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

3.400.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(7/2), x)`

3.400.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(7/2), x)`

3.401
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.401.1 Optimal result	3704
3.401.2 Mathematica [C] (verified)	3705
3.401.3 Rubi [A] (verified)	3705
3.401.4 Maple [B] (verified)	3711
3.401.5 Fricas [F]	3712
3.401.6 Sympy [F(-1)]	3712
3.401.7 Maxima [F]	3712
3.401.8 Giac [F]	3713
3.401.9 Mupad [F(-1)]	3713

3.401.1 Optimal result

Integrand size = 35, antiderivative size = 433

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2(a-b)\sqrt{a+b}(19a^2Ab+8Ab^3+63a^3B-14ab^2B) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\cos(c+dx))}{a+b}}}{105a^4d}$$

$$+ \frac{2(a-b)\sqrt{a+b}(8Ab^2+a^2(25A-63B)+2ab(3A-7B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{105a^3d}$$

$$+ \frac{2A\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)} + \frac{2(Ab+7aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{35ad \cos^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{2(25a^2A-4Ab^2+7abB) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{105a^2d \cos^{\frac{3}{2}}(c+dx)}$$

```
output 2/7*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(7/2)+2/35*(A*b+7*B*a
)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d/cos(d*x+c)^(5/2)+2/105*(25*A*a^2-4
*A*b^2+7*B*a*b)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a^2/d/cos(d*x+c)^(3/2)+2
/105*(a-b)*(19*A*a^2*b+8*A*b^3+63*B*a^3-14*B*a*b^2)*cot(d*x+c)*EllipticE((
a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(
a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a
^4/d+2/105*(a-b)*(8*A*b^2+a^2*(25*A-63*B)+2*a*b*(3*A-7*B))*cot(d*x+c)*Elli
pticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(
1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(
1/2)/a^3/d
```

3.401.
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.401.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.56 (sec) , antiderivative size = 1408, normalized size of antiderivative = 3.25

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `((-4*a*(25*a^4*A - 17*a^2*A*b^2 - 8*A*b^4 - 14*a^3*b*B + 14*a*b^3*B)*Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-19*a^3*A*b - 8*a*A*b^3 - 63*a^4*B + 14*a^2*b^2*B)*((Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-19*a^2*A*b^2 - 8*A*b^4 - 63*a^3*b*B + 14*a*b^3*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])*Sec[c + d*x]]/(a + b)) + (2*a*((a*Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos...`

3.401.3 Rubi [A] (verified)

Time = 2.01 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3478, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.401. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}(A+B \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{9/2}} dx \\
& \quad \downarrow \text{3478} \\
& \frac{2}{7} \int \frac{4Ab \cos^2(c+dx) + (5aA + 7bB) \cos(c+dx) + Ab + 7aB}{\frac{2 \cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}} dx + \\
& \quad \frac{7d \cos^{\frac{7}{2}}(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \frac{4Ab \cos^2(c+dx) + (5aA + 7bB) \cos(c+dx) + Ab + 7aB}{\frac{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}} dx + \\
& \quad \frac{7d \cos^{\frac{7}{2}}(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \frac{4Ab \sin(c+dx+\frac{\pi}{2})^2 + (5aA + 7bB) \sin(c+dx+\frac{\pi}{2}) + Ab + 7aB}{\frac{\sin(c+dx+\frac{\pi}{2})^{7/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}} dx + \\
& \quad \frac{7d \cos^{\frac{7}{2}}(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{3534} \\
& \frac{1}{7} \left(\frac{2 \int \frac{25Aa^2+7bBa+(23Ab+21aB) \cos(c+dx)a-4Ab^2+2b(Ab+7aB) \cos^2(c+dx)}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a} + \frac{2(7aB + Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} \right. \\
& \quad \left. \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \left(\frac{\int \frac{25Aa^2+7bBa+(23Ab+21aB) \cos(c+dx)a-4Ab^2+2b(Ab+7aB) \cos^2(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a} + \frac{2(7aB + Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} \right. \\
& \quad \left. \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)} \right)
\end{aligned}$$

3.401. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned} & \downarrow 3042 \\ \frac{1}{7} & \left(\frac{\int \frac{25Aa^2+7bBa+(23Ab+21aB)\sin(c+dx+\frac{\pi}{2})a-4Ab^2+2b(Ab+7aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{5a} + \frac{2(7aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} \right. \\ & \left. \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3534 \\ \frac{1}{7} & \left(\frac{2\int \frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\cos(c+dx)a+8Ab^3}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{5a} + \frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} + \frac{2(7aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} \right. \\ & \left. \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ \frac{1}{7} & \left(\frac{\int \frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\cos(c+dx)a+8Ab^3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{5a} + \frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} + \frac{2(7aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} \right. \\ & \left. \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ \frac{1}{7} & \left(\frac{\int \frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\sin(c+dx+\frac{\pi}{2})a+8Ab^3}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{5a} + \frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} + \frac{2(7aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} \right. \\ & \left. \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3477 \\ \frac{1}{7} & \left(\frac{\int \frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\sin(c+dx+\frac{\pi}{2})a+8Ab^3}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{5a} + \frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} + \frac{2(7aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} \right. \\ & \left. \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \end{aligned}$$

3.401. $\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{(a-b)(a^2(25A-63B)+2ab(3A-7B)+8Ab^2) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx + (63a^3B+19a^2Ab-14ab^2B+8Ab^3) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{3a} \right)$$

5a

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{(a-b)(a^2(25A-63B)+2ab(3A-7B)+8Ab^2) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + (63a^3B+19a^2Ab-14ab^2B+8Ab^3) \int \frac{\sin(c+dx)}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a} \right)$$

5a

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3295

$$\frac{1}{7} \left(\frac{(63a^3B+19a^2Ab-14ab^2B+8Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)+2ab(3A-7B)+8Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a}}{5a} \right)$$

5a

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3473

$$\frac{1}{7} \left(\frac{2(25a^2A+7abB-4Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)+2ab(3A-7B)+8Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx))}{a-b}}}{ad} \right)$$

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2), x]`

3.401. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

```
output (2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2
*(A*b + 7*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*a*d*Cos[c + d*x]^
(5/2)) + (((2*(a - b)*Sqrt[a + b]*(19*a^2*A*b + 8*A*b^3 + 63*a^3*B - 14*a*
b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]
*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a
+ b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(a^2*d) + (2*(a - b)*Sqrt[a +
b]*(8*A*b^2 + a^2*(25*A - 63*B) + 2*a*b*(3*A - 7*B))*Cot[c + d*x]*Elliptic
F[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a
+ b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d
*x]))/(a - b)))/(a*d))/(3*a) + (2*(25*a^2*A - 4*A*b^2 + 7*a*b*B)*Sqrt[a +
b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/(5*a))/7
```

3.401.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_.) + (f
_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_.) + (f_)*(x_)])/(((b_)*sin[(e_.) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3478 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

3.401.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4603 vs. $2(395) = 790$.

Time = 26.96 (sec) , antiderivative size = 4604, normalized size of antiderivative = 10.63

method	result	size
parts	Expression too large to display	4604
default	Expression too large to display	4663

```
input int((a+cos(d*x+c)*b)^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RET
URNVERBOSE)
```

```
output -2/105*A/d*(-15*a^4*sin(d*x+c)+a^2*b^2*cos(d*x+c)^2*sin(d*x+c)-8*b^4*cos(d
*x+c)^4*sin(d*x+c)-25*a^4*cos(d*x+c)^2*sin(d*x+c)-19*a^2*b^2*cos(d*x+c)^4*
sin(d*x+c)-8*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos
(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b
^4*cos(d*x+c)^5+25*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((
a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)*a^4*cos(d*x+c)^5-16*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1
/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+
c)))^(1/2)*b^4*cos(d*x+c)^4+50*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+
b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+co
s(d*x+c)))^(1/2)*a^4*cos(d*x+c)^4-8*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)
)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/
(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^3+25*EllipticF(cot(d*x+c)-csc(d*x+c),
(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d
*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^3-16*EllipticE(cot(d*x+c)-csc(d
*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*
(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b^3*cos(d*x+c)^4+38*EllipticF(cot(d*x+
c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)
)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3*b*cos(d*x+c)^4+4*EllipticF(c
ot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x...
```

3.401.5 Fricas [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{9}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(9/2),
x)`

3.401.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output `Timed out`

3.401.7 Maxima [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{9}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(9/2)
, x)`

3.401. $\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

3.401.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\cos(dx + c)^{\frac{9}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))^(1/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(9/2), x)`

3.401.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\cos(c + dx)^{9/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(9/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/cos(c + d*x)^(9/2), x)`

3.402 $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

3.402.1 Optimal result	3714
3.402.2 Mathematica [C] (warning: unable to verify)	3715
3.402.3 Rubi [A] (verified)	3716
3.402.4 Maple [B] (warning: unable to verify)	3723
3.402.5 Fricas [F(-1)]	3724
3.402.6 Sympy [F(-1)]	3724
3.402.7 Maxima [F]	3724
3.402.8 Giac [F]	3725
3.402.9 Mupad [F(-1)]	3725

3.402.1 Optimal result

Integrand size = 35, antiderivative size = 670

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx =$$

$$\frac{(a-b)\sqrt{a+b}(24a^2Ab+128Ab^3-9a^3B+156ab^2B) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\middle|-\frac{a+b}{a-b}\right) \sqrt{\frac{a(1+\cos(c+dx))}{a+b}}}{192ab^2d}$$

$$-\frac{\sqrt{a+b}(9a^3B-6a^2b(4A+B)-8b^3(16A+9B)-4ab^2(28A+39B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{192b^2d}$$

$$+\frac{\sqrt{a+b}(8a^3Ab-96aAb^3-3a^4B-24a^2b^2B-48b^4B) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{64b^3d}$$

$$+\frac{(24a^2Ab+128Ab^3-9a^3B+156ab^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{192b^2d\sqrt{\cos(c+dx)}}$$

$$+\frac{(8aAb-3a^2B+12b^2B) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{32bd}$$

$$+\frac{(8Ab-3aB) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{24bd}$$

$$+\frac{B \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{5/2} \sin(c+dx)}{4bd}$$

output $\frac{1}{24}(8Ab-3B^2a)(a+b\cos(dx+c))^{3/2}\sin(dx+c)\cos(dx+c)^{1/2}/b/d + \frac{1}{4}B(a+b\cos(dx+c))^{5/2}\sin(dx+c)\cos(dx+c)^{1/2}/b/d + \frac{1}{192}(24Aa^2b+128A^2b^3-9B^2a^3+156B^2ab^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^2/d\cos(dx+c)^{1/2} + \frac{1}{32}(8A^2ab-3B^2a^2+12B^2b^2)\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/b/d - \frac{1}{192}(a-b)(24A^2a^2b+128A^2b^3-9B^2a^3+156B^2ab^2)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a*(1-\sec(dx+c))/(a+b))^{1/2}(a*(1+\sec(dx+c))/(a-b))^{1/2}/a/b^2/d - \frac{1}{192}(9B^2a^3-6a^2b(4A+B)-8b^3(16A+9B)-4a^2b^2(28A+39B))\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a*(1-\sec(dx+c))/(a+b))^{1/2}(a*(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d + \frac{1}{64}(8A^3b-96A^2ab^3-3B^2a^4-24B^2a^2b^2-48B^2b^4)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a*(1-\sec(dx+c))/(a+b))^{1/2}(a*(1+\sec(dx+c))/(a-b))^{1/2}/b^3/d$

3.402.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.01 (sec) , antiderivative size = 1284, normalized size of antiderivative = 1.92

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{\frac{3}{2}}(A+B\cos(c+dx)) dx = \text{Too large to display}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output

```
-1/384*((-4*a*(-136*a^2*A*b - 128*A*b^3 + 3*a^3*B - 228*a*b^2*B)*Sqrt[((a
+ b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d
*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*
x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt
[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqr
t[a + b*Cos[c + d*x]]) - 4*a*(-416*a*A*b^2 - 228*a^2*b*B - 144*b^3*B)*((Sq
rt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc
[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc
[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/
a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*
x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b
)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c
+ d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqr
t[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*
Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(
-24*a^2*A*b - 128*A*b^3 + 9*a^3*B - 156*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt
[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x
]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x
]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a
+ b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c ...
```

3.402.3 Rubi [A] (verified)

Time = 3.45 (sec) , antiderivative size = 675, normalized size of antiderivative = 1.01, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3469}$$

3.402. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \frac{\int \frac{(a+b \cos(c+dx))^{3/2} ((8Ab-3aB) \cos^2(c+dx)+6bB \cos(c+dx)+aB) dx}{2\sqrt{\cos(c+dx)}} + \frac{4b}{4bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}}{27} \\
& \frac{\int \frac{(a+b \cos(c+dx))^{3/2} ((8Ab-3aB) \cos^2(c+dx)+6bB \cos(c+dx)+aB) dx}{\sqrt{\cos(c+dx)}} + \frac{8b}{4bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}}{3042} \\
& \frac{\int \frac{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2} ((8Ab-3aB) \sin^2(c+dx+\frac{\pi}{2})+6bB \sin(c+dx+\frac{\pi}{2})+aB) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} + \frac{8b}{4bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}}{3528} \\
& \frac{\frac{1}{3} \int \frac{\sqrt{a+b \cos(c+dx)} (3(-3Ba^2+8Aba+12b^2B) \cos^2(c+dx)+2b(16Ab+15aB) \cos(c+dx)+a(8Ab+3aB)) dx}{2\sqrt{\cos(c+dx)}} + \frac{(8Ab-3aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d}}{\frac{8b}{4bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}}{27} \\
& \frac{\frac{1}{6} \int \frac{\sqrt{a+b \cos(c+dx)} (3(-3Ba^2+8Aba+12b^2B) \cos^2(c+dx)+2b(16Ab+15aB) \cos(c+dx)+a(8Ab+3aB)) dx}{\sqrt{\cos(c+dx)}} + \frac{(8Ab-3aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d}}{\frac{8b}{4bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}}{3042} \\
& \frac{\frac{1}{6} \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (3(-3Ba^2+8Aba+12b^2B) \sin^2(c+dx+\frac{\pi}{2})+2b(16Ab+15aB) \sin(c+dx+\frac{\pi}{2})+a(8Ab+3aB)) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} + \frac{(8Ab-3aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3d}}{\frac{8b}{4bd} \frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}}{3528}
\end{aligned}$$

3.402. $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) dx$

$$\frac{1}{6} \left(\frac{1}{2} \int \frac{(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) \cos^2(c+dx)+2b(57Ba^2+104Aba+36b^2B) \cos(c+dx)+a(3Ba^2+56Aba+36b^2B)}{2\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx + \frac{3(-3a^2B}{8b}$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 27

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) \cos^2(c+dx)+2b(57Ba^2+104Aba+36b^2B) \cos(c+dx)+a(3Ba^2+56Aba+36b^2B)}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx + \frac{3(-3a^2B}{8b}$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) \sin(c+dx+\frac{\pi}{2})^2+2b(57Ba^2+104Aba+36b^2B) \sin(c+dx+\frac{\pi}{2})+a(3Ba^2+56Aba+36b^2B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{3(-3a^2B}{8b}$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3540

$$\frac{1}{6} \left(\frac{1}{4} \left(\int -\frac{3(-3Ba^4+8Aba^3-24b^2Ba^2-96Ab^3a-48b^4B) \cos^2(c+dx)-2ab(3Ba^2+56Aba+36b^2B) \cos(c+dx)+a(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx + \frac{3(-3a^2B}{8b} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 25

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{3(-3Ba^4+8Aba^3-24b^2Ba^2-96Ab^3a-48b^4B) \cos^2(c+dx)-2ab(3Ba^2+56Aba+36b^2B) \cos(c+dx)+a(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx + \frac{3(-3a^2B}{8b} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3042

3.402. $\int \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) dx$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{\int \frac{3(-3Ba^4+8Aba^3-24b^2Ba^2-96Ab^3a-48b^4B) \sin(c+dx+\frac{\pi}{2})^2 - 2ab(3Ba^3+24Aba^2+156ab^2B+128Ab^3) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^3} dx \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3532

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) - 2ab(3Ba^2+56Aba+36b^2B) \sin(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) - 2ab(3Ba^2+56Aba+36b^2B) \sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3288

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) - 2ab(3Ba^2+56Aba+36b^2B) \sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3477

3.402. $\int \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) dx$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(9a^3B-6a^2b(4A+B)-4ab^2(28A+39B)-8b^3(16A+9B)) \int \frac{1}{\sqrt{\sin(c+dx)}} dx}{\sqrt{\sin(c+dx)}}$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3295

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \cos(c+dx)}} dx}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \cos(c+dx)}} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

↓ 3473

$$\frac{1}{6} \left(\frac{3(-3a^2B+8aAb+12b^2B) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2}}{4bd}$$

input `Int[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output $(B\sqrt{\cos[c + dx]}(a + b\cos[c + dx])^{5/2}\sin[c + dx]/(4bd) + ((8Ab - 3a^2B)\sqrt{\cos[c + dx]}(a + b\cos[c + dx])^{3/2}\sin[c + dx])/3d + ((3(8a^2Ab - 3a^3B + 12b^2B)\sqrt{\cos[c + dx]}\sqrt{a + b\cos[c + dx]}\sin[c + dx])/(2d) + (-1/2((2(a - b)\sqrt{a + b}(24a^2Ab + 128a^3B - 9a^3B + 156ab^2B)\cot[c + dx]\text{EllipticE}[\text{ArcSin}[\sqrt{a + b\cos[c + dx]}/(\sqrt{a + b}\sqrt{\cos[c + dx]})], -((a + b)/(a - b)))\sqrt{(a(1 - \sec[c + dx])/(a + b))\sqrt{(a(1 + \sec[c + dx])/(a - b))})/(ad) + (2\sqrt{a + b}(9a^3B - 6a^2b(4A + B) - 8b^3(16A + 9B) - 4ab^2(28A + 39B))\cot[c + dx]\text{EllipticF}[\text{ArcSin}[\sqrt{a + b\cos[c + dx]}/(\sqrt{a + b}\sqrt{\cos[c + dx]})], -((a + b)/(a - b))\sqrt{(a(1 - \sec[c + dx])/(a + b))\sqrt{(a(1 + \sec[c + dx])/(a - b))})/d - (6\sqrt{a + b}(8a^3Ab - 96a^2Ab^3 - 3a^4B - 24a^2b^2B - 48b^4B)\cot[c + dx]\text{EllipticPi}[(a + b)/b, \text{ArcSin}[\sqrt{a + b\cos[c + dx]}/(\sqrt{a + b}\sqrt{\cos[c + dx]})], -((a + b)/(a - b))\sqrt{(a(1 - \sec[c + dx])/(a + b))\sqrt{(a(1 + \sec[c + dx])/(a - b))})/(bd))/b + ((24a^2Ab + 128a^3B - 9a^3B + 156ab^2B)\sqrt{a + b\cos[c + dx]}\sin[c + dx])/(bd\sqrt{\cos[c + dx]}))/4)/6)/(8b)$

3.402.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[Fx, x], x]$

rule 27 $\text{Int}[(a_)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)(Gx_)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3288 $\text{Int}[\sqrt{(b_)\sin[e_]} + (f_)(x_)]/\sqrt{(c_)} + (d_)\sin[e_]} + (f_)(x_)]], x_Symbol] \rightarrow \text{Simp}[2b(\tan[e + fx]/(df))\text{Rt}[(c + d)/b, 2]\sqrt{c((1 + \csc[e + fx])/(c - d))}\sqrt{c((1 - \csc[e + fx])/(c + d))}\text{EllipticPi}[(c + d)/d, \text{ArcSin}[\sqrt{c + d\sin[e + fx]}/\sqrt{b\sin[e + fx]}/\text{Rt}[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{PosQ}[(c + d)/b]$

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x]
)^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2]/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2]/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.402.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 5532 vs. $2(616) = 1232$.

Time = 18.24 (sec) , antiderivative size = 5533, normalized size of antiderivative = 8.26

method	result	size
parts	Expression too large to display	5533
default	Expression too large to display	5595

$$3.402. \quad \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{\frac{3}{2}}(A + B \cos(c + dx)) dx$$

input `int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `result too large to display`

3.402.5 Fricas [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="fricas")`

output `Timed out`

3.402.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.402.7 Maxima [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3/
2), x)`

3.402.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3/
2), x)`

3.402.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^{3/2} (A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2),x)`

output `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2), x)`

3.403 $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

3.403.1 Optimal result	3726
3.403.2 Mathematica [C] (verified)	3727
3.403.3 Rubi [A] (verified)	3728
3.403.4 Maple [B] (warning: unable to verify)	3735
3.403.5 Fricas [F]	3736
3.403.6 Sympy [F(-1)]	3737
3.403.7 Maxima [F]	3737
3.403.8 Giac [F]	3737
3.403.9 Mupad [F(-1)]	3738

3.403.1 Optimal result

Integrand size = 35, antiderivative size = 566

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx =$$

$$\frac{(a - b)\sqrt{a + b}(30aAb + 3a^2B + 16b^2B) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a}}{24abd}$$

$$+ \frac{\sqrt{a + b}(30aAb + 12Ab^2 + 3a^2B + 14abB + 16b^2B) \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{24bd}$$

$$- \frac{\sqrt{a + b}(6a^2Ab + 8Ab^3 - a^3B + 12ab^2B) \cot(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{a}}{8b^2d}$$

$$+ \frac{(30aAb + 3a^2B + 16b^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{24bd \sqrt{\cos(c + dx)}}$$

$$+ \frac{(6Ab + 7aB) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{12d}$$

$$+ \frac{bB \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3d}$$

output $\frac{1}{3}bB\cos(dx+c)^{3/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+1/24*(30Aab+3B^2a^2+16B^2b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b/d/\cos(dx+c)^{1/2}+1/12*(6Ab+7Ba)\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/d-1/24*(a-b)*(30Aab+3B^2a^2+16B^2b^2)\cot(dx+c)*\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a/b/d+1/24*(30Aab+12A^2b^2+3B^2a^2+14B^2ab+16B^2b^2)\cot(dx+c)*\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/b/d-1/8*(6A^2b+8A^2b^3-B^2a^3+12B^2ab^2)\cot(dx+c)*\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d$

3.403.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.90 (sec) , antiderivative size = 1227, normalized size of antiderivative = 2.17

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx = \text{Too large to display}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

output

```
((-4*a*(42*a*A*b + 17*a^2*B + 16*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) - 4*a*(48*a^2*A + 24*A*b^2 + 52*a*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[(a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) + 2*(30*a*A*b + 3*a^2*B + 16*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b)*Cos[c + d*x])*Sec[c + d*x])/(a + b)) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c ...
```

3.403.3 Rubi [A] (verified)

Time = 2.78 (sec) , antiderivative size = 568, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}\left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2}\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3469}$$

3.403. $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$

$$\frac{1}{3} \int \frac{\sqrt{\cos(c+dx)}(b(6Ab+7aB)\cos^2(c+dx)+2(3Ba^2+6Aba+2b^2B)\cos(c+dx)+3a(2aA+bB))}{2\sqrt{a+b\cos(c+dx)}} dx + \frac{bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \int \frac{\sqrt{\cos(c+dx)}(b(6Ab+7aB)\cos^2(c+dx)+2(3Ba^2+6Aba+2b^2B)\cos(c+dx)+3a(2aA+bB))}{\sqrt{a+b\cos(c+dx)}} dx + \frac{bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(b(6Ab+7aB)\sin(c+dx+\frac{\pi}{2})^2+2(3Ba^2+6Aba+2b^2B)\sin(c+dx+\frac{\pi}{2})+3a(2aA+bB))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3528

$$\frac{1}{6} \left(\int \frac{b(3Ba^2+30Aba+16b^2B)\cos^2(c+dx)+2b(12Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab(6Ab+7aB)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + \frac{(7aB+6Ab)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} \right) + \frac{bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\int \frac{b(3Ba^2+30Aba+16b^2B)\cos^2(c+dx)+2b(12Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab(6Ab+7aB)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + \frac{(7aB+6Ab)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} \right) + \frac{bB\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{\int \frac{b(3Ba^2+30Aba+16b^2B) \sin(c+dx+\frac{\pi}{2})^2+2b(12Aa^2+13bBa+6Ab^2) \sin(c+dx+\frac{\pi}{2})+ab(6Ab+7aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{4b} + \frac{(7aB+6Ab) \sin(c+dx)}{4b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3540

$$\frac{1}{6} \left(\frac{\int -\frac{-2a(6Ab+7aB) \cos(c+dx)b^2-3(-Ba^3+6Aba^2+12b^2Ba+8Ab^3) \cos^2(c+dx)b+a(3Ba^2+30Aba+16b^2B)b}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{4b} + \frac{(3a^2B+30aAb+16b^2B) \sin(c+dx)}{d \sqrt{\cos(c+dx)}} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 25

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{-2a(6Ab+7aB) \cos(c+dx)b^2-3(-Ba^3+6Aba^2+12b^2Ba+8Ab^3) \cos^2(c+dx)b+a(3Ba^2+30Aba+16b^2B)b}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{2b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{-2a(6Ab+7aB) \sin(c+dx+\frac{\pi}{2})b^2-3(-Ba^3+6Aba^2+12b^2Ba+8Ab^3) \sin(c+dx+\frac{\pi}{2})^2 b+a(3Ba^2+30Aba+16b^2B)b}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3532

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(3Ba^2+30Aba+16b^2B) - 2ab^2(6Ab+7aB) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - 3b(a^3(-B)+6a^2Ab+12ab^2B)}{2b} \right)$$

4b

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(3Ba^2+30Aba+16b^2B) - 2ab^2(6Ab+7aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - 3b(a^3(-B)+6a^2Ab+12ab^2B)}{2b} \right)$$

4b

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3288

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(3Ba^2+30Aba+16b^2B) - 2ab^2(6Ab+7aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{6\sqrt{a+b}(a^3(-B)+6a^2Ab+12ab^2B)}{2b} \right)$$

4b

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3477

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{ab(3a^2B+30aAb+16b^2B) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - ab(3a^2B+30aAb+14abB)}{2b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{-ab(3a^2B+30aAb+14abB+12Ab^2+16b^2B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{\dots} \right)$$

$$\frac{bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3295

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{ab(3a^2B+30aAb+16b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(3a^2B+30aAb+16b^2B)}{\dots}}{\dots} \right)$$

$$\frac{bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

↓ 3473

$$\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{2b\sqrt{a+b}(3a^2B+30aAb+14abB+12Ab^2+16b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx))}{a-b}}}{d} \right)$$

$$\frac{bB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{3d}$$

input `Int[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]),x]`

```
output (b*B*cos[c + d*x]^(3/2)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x]/(3*d) + (((
6*A*b + 7*a*B)*sqrt[cos[c + d*x]]*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/
(2*d) + (-1/2*((2*(a - b)*b*sqrt[a + b]*(30*a*A*b + 3*a^2*B + 16*b^2*B)*cot
[c + d*x]*ellipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(sqrt[a + b]*sqrt[cos[
c + d*x]])], -((a + b)/(a - b)))*sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*sqrt
[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*sqrt[a + b]*(30*a*A*b + 12*
A*b^2 + 3*a^2*B + 14*a*b*B + 16*b^2*B)*cot[c + d*x]*ellipticF[ArcSin[Sqrt[
a + b*cos[c + d*x]]/(sqrt[a + b]*sqrt[cos[c + d*x]])], -((a + b)/(a - b))
]*sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*sqrt[(a*(1 + Sec[c + d*x]))/(a - b)
])/d + (6*sqrt[a + b]*(6*a^2*A*b + 8*A*b^3 - a^3*B + 12*a*b^2*B)*cot[c + d*
x]*ellipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(sqrt[a + b]*sqrt
[cos[c + d*x]])], -((a + b)/(a - b)))*sqrt[(a*(1 - Sec[c + d*x]))/(a + b)
]*sqrt[(a*(1 + Sec[c + d*x]))/(a - b))/d)/b + ((30*a*A*b + 3*a^2*B + 16*b^
2*B)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/(d*sqrt[cos[c + d*x]]))/(4*b)
/6
```

3.403.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]/sqrt[(c_) + (d_)*sin[(e_) + (f_)*
(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*sqrt[c*((1 - Csc[e + f*x])/(c + d))]*ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*sin[e + f*x]]/sqrt[b*sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x]
)^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]
])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.403.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4258 vs. $2(518) = 1036$.

Time = 16.06 (sec) , antiderivative size = 4259, normalized size of antiderivative = 7.52

method	result	size
parts	Expression too large to display	4259
default	Expression too large to display	4305

input `int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/4*A/d*(-5*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-10*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)-4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)+7*cos(d*x+c)^2*sin(d*x+c)*a*b+2*a*b*cos(d*x+c)*sin(d*x+c)-8*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2*cos(d*x+c)^2+4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2*cos(d*x+c)^2-10*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2*cos(d*x+c)-12*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2*cos(d*x+c)-16*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2*cos(d*x+c)+8*EllipticF(cot(d*x+c)-csc...`

3.403.5 Fracas [F]

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))dx = \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^{3/2}\sqrt{\cos(dx+c)}dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x,algorithm="fracas")`

output `integral((B*b*cos(d*x+c)^2+A*a+(B*a+A*b)*cos(d*x+c))*sqrt(b*cos(d*x+c)+a)*sqrt(cos(d*x+c)),x)`

3.403.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.403.7 Maxima [F]

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^{3/2}\sqrt{\cos(dx+c)} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x+c)+A)*(b*cos(d*x+c)+a)^(3/2)*sqrt(cos(d*x+c)),x)`

3.403.8 Giac [F]

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx)) dx = \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^{3/2}\sqrt{\cos(dx+c)} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x+c)+A)*(b*cos(d*x+c)+a)^(3/2)*sqrt(cos(d*x+c)),x)`

3.403.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx = \int \sqrt{\cos(c + dx)}(A + B \cos(c + dx))(a + b \cos(c + dx))^{3/2} dx$$

input `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2),x)`output `int(cos(c + d*x)^(1/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2), x)`

3.404
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.404.1 Optimal result	3739
3.404.2 Mathematica [C] (verified)	3740
3.404.3 Rubi [A] (verified)	3741
3.404.4 Maple [B] (warning: unable to verify)	3747
3.404.5 Fracas [F]	3748
3.404.6 Sympy [F]	3748
3.404.7 Maxima [F]	3748
3.404.8 Giac [F]	3749
3.404.9 Mupad [F(-1)]	3749

3.404.1 Optimal result

Integrand size = 35, antiderivative size = 472

$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}(4Ab+5aB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{4ad}$$

$$+ \frac{\sqrt{a+b}(8aA+4Ab+5aB+2bB) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4d}$$

$$- \frac{\sqrt{a+b}(12aAb+3a^2B+4b^2B) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4bd}$$

$$+ \frac{(4Ab+5aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{4d\sqrt{\cos(c+dx)}}$$

$$+ \frac{bB\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{2d}$$

output $\frac{1}{4}(4A^2b+5B^2a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\cos(dx+c)^{1/2}+1/2$
 $bB\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/d-1/4(a-b)(4A^2b$
 $+5B^2a)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)$
 $^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a$
 $(1+\sec(dx+c))/(a-b))^{1/2}/a/d+1/4(8A^2a+4A^2b+5B^2a+2B^2b)\cot(dx+c)*$
 $\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b)$
 $)^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-$
 $b))^{1/2}/d-1/4(12A^2ab+3B^2a^2+4B^2b^2)\cot(dx+c)\text{EllipticPi}((a+b\cos($
 $dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})($
 $a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b$
 $/d$

3.404.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.02 (sec) , antiderivative size = 1198, normalized size of antiderivative = 2.54

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d`
`*x]],x]`

output `(b*B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + ((-4*a*(8*a^2*A + 4*A*b^2 + 7*a*b*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(16*a*A*b + 8*a^2*B + 4*b^2*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(4*A*b^2 + 5*a*b*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[...`

3.404.3 Rubi [A] (verified)

Time = 2.18 (sec) , antiderivative size = 475, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3469, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3469

3.404. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\frac{1}{2} \int \frac{b(4Ab + 5aB) \cos^2(c + dx) + 2(2Ba^2 + 4Aba + b^2B) \cos(c + dx) + a(4aA + bB)}{2\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx +$$

$$\frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 27

$$\frac{1}{4} \int \frac{b(4Ab + 5aB) \cos^2(c + dx) + 2(2Ba^2 + 4Aba + b^2B) \cos(c + dx) + a(4aA + bB)}{\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx +$$

$$\frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \int \frac{b(4Ab + 5aB) \sin(c + dx + \frac{\pi}{2})^2 + 2(2Ba^2 + 4Aba + b^2B) \sin(c + dx + \frac{\pi}{2}) + a(4aA + bB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})}\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx +$$

$$\frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3540

$$\frac{1}{4} \left(\frac{\int -\frac{b(3Ba^2 + 12Aba + 4b^2B) \cos^2(c + dx) - 2ab(4aA + bB) \cos(c + dx) + ab(4Ab + 5aB)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{2b} + \frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \right)$$

$$\frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 25

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\int -\frac{b(3Ba^2 + 12Aba + 4b^2B) \cos^2(c + dx) - 2ab(4aA + bB) \cos(c + dx) + ab(4Ab + 5aB)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{2b} \right)$$

$$\frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\int -\frac{b(3Ba^2 + 12Aba + 4b^2B) \sin(c + dx + \frac{\pi}{2})^2 - 2ab(4aA + bB) \sin(c + dx + \frac{\pi}{2}) + ab(4Ab + 5aB)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2b} \right)$$

$$\frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3532

3.404. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\int \frac{ab(4Ab+5aB) - 2ab(4aA+bB) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - b(3a^2B + 12aAb + 4b^2)}{2b} \right) \\ \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\ \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\int \frac{ab(4Ab+5aB) - 2ab(4aA+bB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - b(3a^2B + 12aAb + 4b^2)}{2b} \right) \\ \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\ \downarrow \text{3288}$$

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{\int \frac{ab(4Ab+5aB) - 2ab(4aA+bB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(3a^2B+12aAb+4b^2)}{2b} \right) \\ \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\ \downarrow \text{3477}$$

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{ab(5aB + 4Ab) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - ab(8aA + 5aB + 4Ab)}{2b} \right) \\ \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d} \\ \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{-ab(8aA + 5aB + 4Ab + 2bB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} \right) \\ \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

3.404. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

↓ 3295

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{ab(5aB + 4Ab) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(3a^2 B + 4a^2 B^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d} \right) - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

↓ 3473

$$\frac{1}{4} \left(\frac{(5aB + 4Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{2\sqrt{a+b}(3a^2 B + 12aAb + 4b^2 B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d} \right) - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2d}$$

input `Int[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(b*B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*(4*A*b + 5*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(8*a*A + 4*A*b + 5*a*B + 2*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d + (2*Sqrt[a + b]*(12*a*A*b + 3*a^2*B + 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/b + ((4*A*b + 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])/4`

3.404.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)])/Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3469 `Int[((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((A_)] + (B_)*sin[(e_)] + (f_)*(x_)]*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
 $\wedge(3/2)*\text{Sqrt}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]]), x_Symbol] \rightarrow \text{Simp}[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*\text{Sqrt}[c*((1 + \text{Csc}[e + f*x])
)/(c - d)]]*\text{Sqrt}[c*((1 - \text{Csc}[e + f*x])/(c + d))]*\text{EllipticE}[\text{ArcSin}[\text{Sqrt}[c +
d*\text{Sin}[e + f*x]]/\text{Sqrt}[b*\text{Sin}[e + f*x]]/\text{Rt}[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\&
\text{PosQ}[(c + d)/b]$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^3/2)*\text{Sqrt}[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] \rightarrow \text{S}
imp[(A - B)/(a - b) Int[1/(\text{Sqrt}[a + b*\text{Sin}[e + f*x]]*\text{Sqrt}[c + d*\text{Sin}[e + f*
x]]), x], x] - \text{Simp}[(A*b - a*B)/(a - b) Int[(1 + \text{Sin}[e + f*x])/((a + b*\text{Si}
n[e + f*x])^3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x]]), x], x] /; \text{FreeQ}\{a, b, c, d, e
, f, A, B\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2,
0] \&\& \text{NeQ}[A, B]`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^
2)/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^3/2)*\text{Sqrt}[(c_) + (d_)*sin[(e_)
+ (f_)*(x_)]]), x_Symbol] \rightarrow \text{Simp}[C/b^2 Int[\text{Sqrt}[a + b*\text{Sin}[e + f*x]]
/\text{Sqrt}[c + d*\text{Sin}[e + f*x]], x], x] + \text{Simp}[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*\text{Sin}[e + f*x])/((a + b*\text{Sin}[e + f*x])^3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x
]], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\&
\&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]`

rule 3540 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^
2)/(\text{Sqrt}[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*\text{Sqrt}[(c_) + (d_)*sin[(e_) +
(f_)*(x_)]]), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cos}[e + f*x]*(\text{Sqrt}[c + d*\text{Sin}[e + f
*x]]/(d*f*\text{Sqrt}[a + b*\text{Sin}[e + f*x]])), x] + \text{Simp}[1/(2*d) Int[(1/((a + b*\text{Si}
n[e + f*x])^3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x]])]*\text{Simp}[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*\text{Sin}[e + f*x] + (2*b*B*d - C*(b*c + a*d))*\text{Sin}[e +
f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b*c - a
*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]`

3.404.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 3309 vs. $2(430) = 860$.

Time = 17.60 (sec) , antiderivative size = 3310, normalized size of antiderivative = 7.01

method	result	size
parts	Expression too large to display	3310
default	Expression too large to display	3318

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -A/d*(EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)
*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(
d*x+c)^2+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)
))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b^2*c
os(d*x+c)^2+6*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d
*x+c))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/
2))*a*b*cos(d*x+c)^2+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)
/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b)
)^(1/2))*a^2*cos(d*x+c)^2-4*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b)
)^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d
*x+c)))^(1/2)*a*b*cos(d*x+c)^2+2*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(
a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+
cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),
(-(a-b)/(a+b))^(1/2))*b^2*cos(d*x+c)+12*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*
((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*
x+c),-1,(-(a-b)/(a+b))^(1/2))*a*b*cos(d*x+c)+4*(cos(d*x+c)/(1+cos(d*x+c)))
^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-
csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)-8*EllipticF(cot(d*x+c)-csc
(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(...
```

3.404.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.404.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx))(a + b \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**(3/2)/sqrt(cos(c + d*x)), x)`

3.404.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/sqrt(cos(d*x + c)), x)`

3.404.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/sqrt(cos(d*x + c)), x)`

3.404.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(1/2),x)`

3.405
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.405.1 Optimal result 3750
 3.405.2 Mathematica [C] (verified) 3751
 3.405.3 Rubi [A] (verified) 3751
 3.405.4 Maple [B] (warning: unable to verify) 3756
 3.405.5 Fricas [F] 3757
 3.405.6 Sympy [F] 3758
 3.405.7 Maxima [F] 3758
 3.405.8 Giac [F] 3758
 3.405.9 Mupad [F(-1)] 3759

3.405.1 Optimal result

Integrand size = 35, antiderivative size = 449

$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{(a-b)\sqrt{a+b}(2aA-bB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right)\right)}{ad} - \frac{\sqrt{a+b}(2a(A-B)-b(4A+B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{d} - \frac{\sqrt{a+b}(2Ab+3aB) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{d} + \frac{2aA\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{(2aA-bB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}}$$

output

```
2*a*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)-(2*A*a-B*b)*sin
(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)+(a-b)*(2*A*a-B*b)*cot(d*
x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)
/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)
))/(a+b)^(1/2)/a/d-(2*a*(A-B)-b*(4*A+B))*cot(d*x+c)*EllipticF((a+b*cos(d*x
+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*
(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a+b)^(1/2)/d-(2*A*b+3*B
*a)*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1
/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/
2)*(a*(1+sec(d*x+c)))/(a+b)^(1/2)/d
```

3.405.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.58 (sec) , antiderivative size = 1196, normalized size of antiderivative = 2.66

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`

output `(2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) + ((4*a*(-2*a*A*b - 2*a^2*B - b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 4*a*(2*a^2*A - 2*A*b^2 - 4*a*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 2*(2*a*A*b - b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + ...`

3.405.3 Rubi [A] (verified)

Time = 2.17 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3468, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.405. $\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\
& \quad \downarrow \text{3468} \\
& 2 \int \frac{-b(2aA - bB) \cos^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \cos(c + dx) + a(2Ab + aB)}{2\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)} + \frac{2aA \sin(c + dx)\sqrt{a + b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}} dx + \\
& \quad \downarrow \text{27} \\
& \int \frac{-b(2aA - bB) \cos^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \cos(c + dx) + a(2Ab + aB)}{\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)} + \frac{2aA \sin(c + dx)\sqrt{a + b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}} dx + \\
& \quad \downarrow \text{3042} \\
& \int \frac{-b(2aA - bB) \sin(c + dx + \frac{\pi}{2})^2 + (-Aa^2 + 2bBa + Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(2Ab + aB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})}\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} + \frac{2aA \sin(c + dx)\sqrt{a + b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}} dx + \\
& \quad \downarrow \text{3540} \\
& \frac{\int \frac{b^2(2Ab + 3aB) \cos^2(c + dx) + 2ab(2Ab + aB) \cos(c + dx) + ab(2aA - bB)}{\cos^{3/2}(c + dx)\sqrt{a + b \cos(c + dx)}} dx}{\frac{2b}{(2aA - bB) \sin(c + dx)\sqrt{a + b \cos(c + dx)}} + \frac{2aA \sin(c + dx)\sqrt{a + b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}} - \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{b^2(2Ab + 3aB) \sin(c + dx + \frac{\pi}{2})^2 + 2ab(2Ab + aB) \sin(c + dx + \frac{\pi}{2}) + ab(2aA - bB)}{\sin(c + dx + \frac{\pi}{2})^{3/2}\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{\frac{2b}{(2aA - bB) \sin(c + dx)\sqrt{a + b \cos(c + dx)}} + \frac{2aA \sin(c + dx)\sqrt{a + b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}} - \\
& \quad \downarrow \text{3532}
\end{aligned}$$

3.405. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx$

$$\begin{aligned}
& \frac{b^2(3aB + 2Ab) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx + \int \frac{ab(2aA-bB)+2ab(2Ab+aB) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2b}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}} \\
& \quad \downarrow 3042 \\
& \frac{b^2(3aB + 2Ab) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{ab(2aA-bB)+2ab(2Ab+aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2b}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}} \\
& \quad \downarrow 3288 \\
& \frac{\int \frac{ab(2aA-bB)+2ab(2Ab+aB) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(3aB+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}}\right)\right)}{d}}{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2b}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}} \\
& \quad \downarrow 3477 \\
& \frac{ab(2aA - bB) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - ab(2aA - 2aB - 4Ab - bB) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx - \frac{2b\sqrt{a+b}(3aB+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}}\right)\right)}{2b}}{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2b}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}} \\
& \quad \downarrow 3042 \\
& \frac{-ab(2aA - 2aB - 4Ab - bB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + ab(2aA - bB) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2b}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}} \\
& \quad \downarrow 3295
\end{aligned}$$

3.405. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
 & ab(2aA - bB) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(2aA-2aB-4Ab-bB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d} \\
 & \frac{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \\
 & \quad \downarrow \text{3473} \\
 & - \frac{2b\sqrt{a+b}(2aA-2aB-4Ab-bB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d} + \frac{2b(a-b)\sqrt{a+b}(2aA-2aB-4Ab-bB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d} \\
 & \frac{(2aA - bB) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}
 \end{aligned}$$

input `Int[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2), x]`

output `((2*(a - b)*b*Sqrt[a + b]*(2*a*A - b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/ (a*d) - (2*b*Sqrt[a + b]*(2*a*A - 4*A*b - 2*a*B - b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (2*b*Sqrt[a + b]*(2*A*b + 3*a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/d)/(2*b) + (2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - ((2*a*A - b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])`

3.405.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.405. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

```
rule 3477 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] :> Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.405.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2379 vs. $2(417) = 834$.

Time = 17.90 (sec) , antiderivative size = 2380, normalized size of antiderivative = 5.30

method	result	size
parts	Expression too large to display	2380
default	Expression too large to display	2601

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RETURNVERBOSE)
```

$$3.405. \quad \int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

output

```

-2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(
d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(
d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2-2*(-csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
))*a*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*
x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+
1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2
+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a
^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c)
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)
-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b-2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)
^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a
+b)/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))
)*b^2+csc(d*x+c)^3*a^2*(1-cos(d*x+c))^3-csc(d*x+c)^3*a*b*(1-cos(d*x+c))^3+a
^2*(csc(d*x+c)-cot(d*x+c))+a*b*(csc(d*x+c)-cot(d*x+c))*((csc(d*x+c)^2*a*(
1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(
d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*a*(1-c
os(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(-csc(d*x+c)^2*(1-co...

```

3.405.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{3/2}} dx$$

input

```

integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algo
rithm="fricas")

```

output

```

integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(
d*x + c) + a)/cos(d*x + c)^(3/2), x)

```

3.405.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos^{3/2}(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**(3/2)/cos(c + d*x)**(3/2), x)`

3.405.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(3/2), x)`

3.405.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(3/2), x)`

3.405.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(3/2),x)`

3.406
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.406.1 Optimal result 3760
 3.406.2 Mathematica [C] (verified) 3761
 3.406.3 Rubi [A] (verified) 3761
 3.406.4 Maple [B] (warning: unable to verify) 3766
 3.406.5 Fricas [F] 3767
 3.406.6 Sympy [F] 3767
 3.406.7 Maxima [F] 3767
 3.406.8 Giac [F] 3768
 3.406.9 Mupad [F(-1)] 3768

3.406.1 Optimal result

Integrand size = 35, antiderivative size = 419

$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(4Ab+3aB) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}}{\sqrt{a+b}}\right)\right)}{3ad} + \frac{2\sqrt{a+b}(3Ab^2+a^2(A-3B)-a(4Ab-6bB)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{a}}{3ad} - \frac{2b\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{d} + \frac{2aA\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)}$$

output

```
2/3*a*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)+2/3*(a-b)*(4*
A*b+3*B*a)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x
+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)
*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d+2/3*(3*A*b^2+a^2*(A-3*B)-a*(4*A*b-6*B*
b))*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/
2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+
sec(d*x+c))/(a-b))^(1/2)/a/d-2*b*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(
1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/
2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d
```

3.406.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.54 (sec) , antiderivative size = 1236, normalized size of antiderivative = 2.95

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `((-4*a*(a^2*A - A*b^2 + 3*a*b*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-4*a*A*b - 3*a^2*B + 3*b^2*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-4*A*b^2 - 3*a*b*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)] + (2*a*((a*Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[...`

3.406.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3468, 27, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.406. $\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx \\
& \quad \downarrow \text{3468} \\
& \frac{2}{3} \int \frac{3b^2 B \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(4Ab + 3aB)}{2 \cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{3/2}(c + dx)}} dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{3b^2 B \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(4Ab + 3aB)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{3/2}(c + dx)}} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{3/2}(c + dx)}} dx + \\
& \quad \downarrow \text{3532} \\
& \frac{1}{3} \left(\int \frac{a(4Ab + 3aB) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + 3b^2 B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(\int \frac{a(4Ab + 3aB) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{3/2}(c + dx)}} dx + 3b^2 B \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \downarrow \text{3288}
\end{aligned}$$

$$\frac{1}{3} \left(\int \frac{a(4Ab + 3aB) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{6bB\sqrt{a+b} \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a(\sec(c+dx)+1)}}{\frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}} \right)$$

↓ 3477

$$\frac{1}{3} \left((a^2(A - 3B) - a(4Ab - 6bB) + 3Ab^2) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx + a(3aB + 4Ab) \int \frac{\cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)} dx - \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{3} \left((a^2(A - 3B) - a(4Ab - 6bB) + 3Ab^2) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + a(3aB + 4Ab) \int \frac{\sin(c + dx + \frac{\pi}{2})}{\sin^{\frac{3}{2}}(c + dx + \frac{\pi}{2})} dx - \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3295

$$\frac{1}{3} \left(a(3aB + 4Ab) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(a^2(A - 3B) - a(4Ab - 6bB) + 3Ab^2) \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a(\sec(c+dx)+1)}}{\frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)}} \right)$$

↓ 3473

$$\frac{1}{3} \left(\frac{2\sqrt{a+b}(a^2(A - 3B) - a(4Ab - 6bB) + 3Ab^2) \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a(\sec(c+dx)+1)}}{ad} \text{EllipticF} \left(\arcsin \left(\frac{\sin(c + dx + \frac{\pi}{2})}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \right), \frac{a+b}{2\sqrt{a+b}} \right) - \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

```
input Int[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x
]
```

3.406. $\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

```
output ((2*(a - b)*Sqrt[a + b]*(4*A*b + 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt
[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))
]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b
)]/(a*d) + (2*Sqrt[a + b]*(3*A*b^2 + a^2*(A - 3*B) - a*(4*A*b - 6*b*B))*Co
t[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos
[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqr
t[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d) - (6*b*Sqrt[a + b]*B*Cot[c + d*x]
*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[C
os[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*S
qrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d)/3 + (2*a*A*Sqrt[a + b*Cos[c + d*x]
]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2))
```

3.406.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f
_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.406.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2467 vs. $2(385) = 770$.

Time = 19.03 (sec) , antiderivative size = 2468, normalized size of antiderivative = 5.89

method	result	size
parts	Expression too large to display	2468
default	Expression too large to display	2696

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output -2/3*A/d*((cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)
))/a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*
cos(d*x+c)^3+4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(
d*x+c))/a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))
*a*b*cos(d*x+c)^3+3*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*
((a+cos(d*x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^
(1/2)*b^2*cos(d*x+c)^3-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*
b)/(1+cos(d*x+c))/a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+
b))^(1/2))*a*b*cos(d*x+c)^3-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*
x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)
)/(a+b))^(1/2)*b^2*cos(d*x+c)^3+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+c
os(d*x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-
(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)^2+8*EllipticF(cot(d*x+c)-csc(d*x+c),(-
(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)*(cos(d*x
+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2+6*EllipticF(cot(d*x+c)-csc(d*x+
c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)*(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2*cos(d*x+c)^2-8*EllipticE(cot(d*x+c)-csc
(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)
)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-8*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/a+b)^(1/2)*EllipticE(...
```

3.406.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(5/2), x)`

3.406.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos^{5/2}(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**(3/2)/cos(c + d*x)**(5/2), x)`

3.406.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(5/2), x)`

3.406.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(5/2), x)`

3.406.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(5/2),x)`

3.407
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

3.407.1 Optimal result 3769
 3.407.2 Mathematica [C] (verified) 3770
 3.407.3 Rubi [A] (verified) 3770
 3.407.4 Maple [B] (verified) 3775
 3.407.5 Fracas [F] 3776
 3.407.6 Sympy [F(-1)] 3777
 3.407.7 Maxima [F] 3777
 3.407.8 Giac [F] 3777
 3.407.9 Mupad [F(-1)] 3778

3.407.1 Optimal result

Integrand size = 35, antiderivative size = 353

$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(9a^2A+3Ab^2+20abB) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{a(1-\cos(c+dx))}}{15ad} + \frac{2aA\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{5d \cos^{5/2}(c+dx)} + \frac{2(6Ab+5aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15d \cos^{3/2}(c+dx)}$$

output

```
2/5*a*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+2/15*(6*A*b+5*B*a)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)+2/15*(a-b)*(9*A*a^2+3*A*b^2+20*B*a*b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^2/d-2/15*(a-b)*(9*A*a-3*A*b-5*B*a+15*B*b)*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a/d
```


3.407.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.64 (sec) , antiderivative size = 1314, normalized size of antiderivative = 3.72

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `-1/15*((-4*a*(-3*a^2*A*b + 3*A*b^3 - 5*a^3*B + 5*a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(9*a^3*A + 3*a*A*b^2 + 20*a^2*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(9*a^2*A*b + 3*A*b^3 + 20*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b))) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + ...`

3.407.3 Rubi [A] (verified)

Time = 1.53 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3468, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.407. $\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow \text{3468} \\
& \frac{2}{5} \int \frac{b(2aA + 5bB) \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(6Ab + 5aB)}{2 \cos^{5/2}(c + dx) \sqrt{a + b \cos(c + dx)} + 2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)} + 5d \cos^{5/2}(c + dx)} dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{b(2aA + 5bB) \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(6Ab + 5aB)}{\cos^{5/2}(c + dx) \sqrt{a + b \cos(c + dx)} + 2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)} + 5d \cos^{5/2}(c + dx)} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{b(2aA + 5bB) \sin(c + dx + \frac{\pi}{2})^2 + (3Aa^2 + 10bBa + 5Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(6Ab + 5aB)}{\sin(c + dx + \frac{\pi}{2})^{5/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} + 2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)} + 5d \cos^{5/2}(c + dx)} dx + \\
& \quad \downarrow \text{3534} \\
& \frac{1}{5} \left(\frac{2 \int \frac{a(9Aa^2 + 20bBa + 3Ab^2) + a(5Ba^2 + 12Aba + 15b^2B) \cos(c + dx)}{2 \cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{3a} + \frac{2(5aB + 6Ab) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3d \cos^{3/2}(c + dx)} \right) + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5d \cos^{5/2}(c + dx)} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.407. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$

$$\frac{1}{5} \left(\frac{\int \frac{a(9Aa^2+20bBa+3Ab^2)+a(5Ba^2+12Aba+15b^2B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(5aB+6Ab) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \downarrow 3042$$

$$\frac{1}{5} \left(\frac{\int \frac{a(9Aa^2+20bBa+3Ab^2)+a(5Ba^2+12Aba+15b^2B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a} + \frac{2(5aB+6Ab) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \downarrow 3477$$

$$\frac{1}{5} \left(\frac{a(9a^2A+20abB+3Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx - a(a-b)(9aA-5aB-3Ab+15bB) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(5aB+6Ab) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \downarrow 3042$$

$$\frac{1}{5} \left(\frac{a(9a^2A+20abB+3Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - a(a-b)(9aA-5aB-3Ab+15bB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a} + \frac{2(5aB+6Ab) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \downarrow 3295$$

$$\frac{1}{5} \left(\frac{a(9a^2A+20abB+3Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(9aA-5aB-3Ab+15bB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b \cos(c+dx)}}}{3a}}{3a} + \frac{2(5aB+6Ab) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} \right) + \frac{2aA \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

3.407. $\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

↓ 3473

$$\frac{1}{5} \left(\frac{2(a-b)\sqrt{a+b}(9a^2A+20abB+3Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right)}{ad} - \frac{2(a-b)\sqrt{a+b}(9a^2A+20abB+3Ab^2)}{3a} \right) - \frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)}$$

input `Int[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*(a - b)*Sqrt[a + b]*(9*a^2*A + 3*A*b^2 + 20*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(a*d) - (2*(a - b)*Sqrt[a + b]*(9*a*A - 3*A*b - 5*a*B + 15*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/d)/(3*a) + (2*(6*A*b + 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2))/5`

3.407.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

3.407. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.407.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3549 vs. $2(321) = 642$.

Time = 23.43 (sec) , antiderivative size = 3550, normalized size of antiderivative = 10.06

method	result	size
parts	Expression too large to display	3550
default	Expression too large to display	3614

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RET
URNVERBOSE)
```

output

```

2/5*A/d*(EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)
/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2
*cos(d*x+c)^4-4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos
(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2
)*a^2*b*cos(d*x+c)^4-EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))
*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))
^(1/2)*a*b^2*cos(d*x+c)^4+6*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))
^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))
/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3+2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)
/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(
d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3-8*EllipticF(cot(d*x+c)-csc(d*x+c),
(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/
(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3-2*EllipticF(cot(d*x+c)-csc(
d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x
+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3+3*EllipticE(cot(d*x+
c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+
cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^2-6*EllipticF(c
ot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^3*cos(d*x+c)^3+3*Ellipt
icE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+...

```

3.407.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{7/2}} dx$$

input

```

integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="fricas")

```

output

```

integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(
d*x + c) + a)/cos(d*x + c)^(7/2), x)

```

3.407.
$$\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

3.407.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.407.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{7/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(7/
2), x)`

3.407.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{7/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(7/
2), x)`

3.407. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

3.407.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(7/2),x)`

3.408
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.408.1 Optimal result 3779
 3.408.2 Mathematica [C] (verified) 3780
 3.408.3 Rubi [A] (verified) 3780
 3.408.4 Maple [B] (verified) 3786
 3.408.5 Fracas [F] 3787
 3.408.6 Sympy [F(-1)] 3788
 3.408.7 Maxima [F] 3788
 3.408.8 Giac [F] 3788
 3.408.9 Mupad [F(-1)] 3789

3.408.1 Optimal result

Integrand size = 35, antiderivative size = 433

$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(82a^2Ab-6Ab^3+63a^3B+21ab^2B) \cot(c+dx)}{105a^2d} + \frac{2(a-b)\sqrt{a+b}(6Ab^2-a^2(25A-63B)+3ab(19A-7B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{105a^2d} + \frac{2aA\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)} + \frac{2(8Ab+7aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{35d \cos^{\frac{5}{2}}(c+dx)} + \frac{2(25a^2A+3Ab^2+42abB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{105ad \cos^{\frac{3}{2}}(c+dx)}$$

```
output 2/7*a*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(7/2)+2/35*(8*A*b+7
*B*a)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+2/105*(25*A*a^2
+3*A*b^2+42*B*a*b)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d/cos(d*x+c)^(3/2)+
2/105*(a-b)*(82*A*a^2*b-6*A*b^3+63*B*a^3+21*B*a*b^2)*cot(d*x+c)*EllipticE(
(a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*
(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/
a^3/d-2/105*(a-b)*(6*A*b^2-a^2*(25*A-63*B)+3*a*b*(19*A-7*B))*cot(d*x+c)*El
lipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))
^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b)
)^(1/2)/a^2/d
```

3.408.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.68 (sec) , antiderivative size = 1407, normalized size of antiderivative = 3.25

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `((-4*a*(25*a^4*A - 31*a^2*A*b^2 + 6*A*b^4 + 21*a^3*b*B - 21*a*b^3*B)*Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-82*a^3*A*b + 6*a*A*b^3 - 63*a^4*B - 21*a^2*b^2*B)*((Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]*Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-82*a^2*A*b^2 + 6*A*b^4 - 63*a^3*b*B - 21*a*b^3*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[(a + b)*Cot[(c + d*x)/2]^2]/(-a + b))*Sqrt[-((a + b)*Cos...`

3.408.3 Rubi [A] (verified)

Time = 2.06 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3468, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.408. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{9/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx \\
& \quad \downarrow \text{3468} \\
& \frac{2}{7} \int \frac{b(4aA + 7bB) \cos^2(c + dx) + (5Aa^2 + 14bBa + 7Ab^2) \cos(c + dx) + a(8Ab + 7aB)}{2 \cos^{7/2}(c + dx) \sqrt{a + b \cos(c + dx)} + 2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)} + 7d \cos^{7/2}(c + dx)} dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \frac{b(4aA + 7bB) \cos^2(c + dx) + (5Aa^2 + 14bBa + 7Ab^2) \cos(c + dx) + a(8Ab + 7aB)}{\cos^{7/2}(c + dx) \sqrt{a + b \cos(c + dx)} + 2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)} + 7d \cos^{7/2}(c + dx)} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \frac{b(4aA + 7bB) \sin(c + dx + \frac{\pi}{2})^2 + (5Aa^2 + 14bBa + 7Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(8Ab + 7aB)}{\sin(c + dx + \frac{\pi}{2})^{7/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})} + 2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)} + 7d \cos^{7/2}(c + dx)} dx + \\
& \quad \downarrow \text{3534} \\
& \frac{1}{7} \left(2 \int \frac{2ab(8Ab + 7aB) \cos^2(c + dx) + a(21Ba^2 + 44Aba + 35b^2B) \cos(c + dx) + a(25Aa^2 + 42bBa + 3Ab^2)}{2 \cos^{5/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{2(7aB + 8Ab) \sin(c + dx)}{5d \cos^{5/2}(c + dx)} \right) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.408. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx$

$$\frac{1}{7} \left(\frac{\int \frac{2ab(8Ab+7aB) \cos^2(c+dx) + a(21Ba^2+44Aba+35b^2B) \cos(c+dx) + a(25Aa^2+42bBa+3Ab^2)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a} + \frac{2(7aB+8Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{\int \frac{2ab(8Ab+7aB) \sin(c+dx+\frac{\pi}{2})^2 + a(21Ba^2+44Aba+35b^2B) \sin(c+dx+\frac{\pi}{2}) + a(25Aa^2+42bBa+3Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5a} + \frac{2(7aB+8Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3534

$$\frac{1}{7} \left(\frac{2 \int \frac{(25Aa^2+84bBa+51Ab^2) \cos(c+dx)a^2 + (63Ba^3+82Aba^2+21b^2Ba-6Ab^3)a}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(25a^2A+42abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + \frac{2(7aB+8Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{\int \frac{(25Aa^2+84bBa+51Ab^2) \cos(c+dx)a^2 + (63Ba^3+82Aba^2+21b^2Ba-6Ab^3)a}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(25a^2A+42abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + \frac{2(7aB+8Ab) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3042

3.408. $\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}} (A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{\int \frac{(25Aa^2+84bBa+51Ab^2) \sin(c+dx+\frac{\pi}{2})a^2+(63Ba^3+82Aba^2+21b^2Ba-6Ab^3)a}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a} + \frac{2(25a^2A+42abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + 2 \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3477

$$\frac{1}{7} \left(\frac{a(63a^3B+82a^2Ab+21ab^2B-6Ab^3) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - a(a-b)(-(a^2(25A-63B))+a(57Ab-21bB)+6Ab^2) \int \frac{1}{\sqrt{\cos(c+dx) \sqrt{a+b \cos(c+dx)}}}}{3a} + \frac{2(25a^2A+42abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + 2 \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{a(63a^3B+82a^2Ab+21ab^2B-6Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - a(a-b)(-(a^2(25A-63B))+a(57Ab-21bB)+6Ab^2) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}}}{3a} + \frac{2(25a^2A+42abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + 2 \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3295

$$\frac{1}{7} \left(\frac{a(63a^3B+82a^2Ab+21ab^2B-6Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b) \sqrt{a+b} (-(a^2(25A-63B))+a(57Ab-21bB)+6Ab^2) \cot(c+dx) \sqrt{a+b \cos(c+dx)}}{3a}}{3a} + \frac{2(25a^2A+42abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + 2 \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

↓ 3473

3.408. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{2(25a^2A + 42abB + 3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3d \cos^{\frac{3}{2}}(c+dx)} + \frac{2(a-b)\sqrt{a+b} (63a^3B + 82a^2Ab + 21ab^2B - 6Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{ad} \right) \\ \frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)}$$

input `Int[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `(2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(8*A*b + 7*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(82*a^2*A*b - 6*A*b^3 + 63*a^3*B + 21*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*(a - b)*Sqrt[a + b]*(6*A*b^2 - a^2*(25*A - 63*B) + a*(57*A*b - 21*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(3*a) + (2*(25*a^2*A + 3*A*b^2 + 42*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))/(5*a))/7`

3.408.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)]*Sqrt[(a_) + (b_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

3.408. $\int \frac{(a+b \cos(c+dx))^{\frac{3}{2}}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m - 1)*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Ssin[e + f*x])^(m - 2)*(c + d*Ssin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Ssin[e + f*x]]/Sqrt[b*Ssin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Ssin[e + f*x])^(3/2)*Sqrt[c + d*Ssin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`


```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.408.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4604 vs. $2(395) = 790$.

Time = 27.51 (sec) , antiderivative size = 4605, normalized size of antiderivative = 10.64

method	result	size
parts	Expression too large to display	4605
default	Expression too large to display	4663

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RET
URNVERBOSE)
```

output `-2/105*A/d*(-15*a^4*sin(d*x+c)-27*a^2*b^2*cos(d*x+c)^2*sin(d*x+c)+6*b^4*cos(d*x+c)^4*sin(d*x+c)-25*a^4*cos(d*x+c)^2*sin(d*x+c)-82*a^2*b^2*cos(d*x+c)^4*sin(d*x+c)+6*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^5+25*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^5+12*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^4+50*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^4+6*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^3+25*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^3+12*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b^3*cos(d*x+c)^4+164*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3*b*cos(d*x+c)^4+102*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+c...`

3.408.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{9/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorith="fracas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(9/2), x)`

3.408.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output `Timed out`

3.408.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{9/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(9/2), x)`

3.408.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{9/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(9/2), x)`

3.408.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^{9/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(9/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(9/2),x)`

3.409
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

3.409.1 Optimal result 3790
 3.409.2 Mathematica [C] (verified) 3791
 3.409.3 Rubi [A] (verified) 3792
 3.409.4 Maple [B] (verified) 3799
 3.409.5 Fracas [F] 3799
 3.409.6 Sympy [F(-1)] 3800
 3.409.7 Maxima [F] 3800
 3.409.8 Giac [F] 3800
 3.409.9 Mupad [F(-1)] 3801

3.409.1 Optimal result

Integrand size = 35, antiderivative size = 522

$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(147a^4A+33a^2Ab^2+8Ab^4+246a^3bB - 2(a-b)\sqrt{a+b}(8Ab^3-a^3(147A-75B)+3a^2b(13A-57B)+6ab^2(A-3B)) \cot(c+dx) \text{EllipticF}(\arcsin(\frac{\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{\sqrt{a+b}}), \frac{315a^3d}{63d \cos^{\frac{7}{2}}(c+dx)})}{315a^3d} + \frac{2aA\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)} + \frac{2(10Ab+9aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{63d \cos^{\frac{7}{2}}(c+dx)} + \frac{2(49a^2A+3Ab^2+72abB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{315ad \cos^{\frac{5}{2}}(c+dx)} + \frac{2(88a^2Ab-4Ab^3+75a^3B+9ab^2B)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{315a^2d \cos^{\frac{3}{2}}(c+dx)}$$

output $\frac{2}{9}aA\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\cos(dx+c)^{9/2} + \frac{2}{63}(10Ab+9B^2a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\cos(dx+c)^{7/2} + \frac{2}{315}(49A^2+3A^2b^2+72B^2a^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/a^2d\cos(dx+c)^{5/2} + \frac{2}{315}(88A^2b-4A^2b^3+75B^2a^3+9B^2a^2b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/a^2d\cos(dx+c)^{3/2} + \frac{2}{315}(a-b)(147A^4+33A^2b^2+8A^2b^4+246B^2a^3b-18B^2a^2b^3)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a(1-\sec(dx+c)))/(a+b)^{1/2}*(a(1+\sec(dx+c)))/(a-b)^{1/2})/a^4d + \frac{2}{315}(a-b)(8A^3b^3-a^3(147A-75B)+3a^2b(13A-57B)+6a^2b^2(A-3B))\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a(1-\sec(dx+c)))/(a+b)^{1/2}*(a(1+\sec(dx+c)))/(a-b)^{1/2})/a^3d$

3.409.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.87 (sec) , antiderivative size = 1515, normalized size of antiderivative = 2.90

$$\int \frac{(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\cos^{11/2}(c+dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2), x]`

output
$$-1/315*((-4*a*(-39*a^4*A*b + 31*a^2*A*b^3 + 8*A*b^5 - 75*a^5*B + 93*a^3*b^2*B - 18*a*b^4*B)*\text{Sqrt}[\text{((a + b)*\text{Cot}[(c + d*x)/2]^2)/(-a + b)}]*\text{Sqrt}[-\text{((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a}]*\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a}]*\text{Csc}[c + d*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a}]/\text{Sqrt}[2]], (-2*a)/(-a + b)]*\text{Sin}[(c + d*x)/2]^4)/\text{((a + b)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])} - 4*a*(147*a^5*A + 33*a^3*A*b^2 + 8*a*A*b^4 + 246*a^4*b*B - 18*a^2*b^3*B)*(\text{Sqrt}[\text{((a + b)*\text{Cot}[(c + d*x)/2]^2)/(-a + b)}]*\text{Sqrt}[-\text{((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a}]*\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a}]*\text{Csc}[c + d*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a}]/\text{Sqrt}[2]], (-2*a)/(-a + b)]*\text{Sin}[(c + d*x)/2]^4)/\text{((a + b)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])} - (\text{Sqrt}[\text{((a + b)*\text{Cot}[(c + d*x)/2]^2)/(-a + b)}]*\text{Sqrt}[-\text{((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a}]*\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a}]*\text{Csc}[c + d*x]*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a}]/\text{Sqrt}[2]], (-2*a)/(-a + b)]*\text{Sin}[(c + d*x)/2]^4)/(b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])) + 2*(147*a^4*A*b + 33*a^2*A*b^3 + 8*A*b^5 + 246*a^3*b^2*B - 18*a*b^4*B)*((I*\text{Cos}[(c + d*x)/2]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]*\text{EllipticE}[I*\text{ArcSinh}[\text{Sin}[(c + d*x)/2]/\text{Sqrt}[\text{Cos}[c + d*x]]], (-2*a)/(-a - b)]*\text{Sec}[c + d*x])/(b*\text{Sqrt}[\text{Cos}[(c + d*x)/2]^2*\text{Sec}[c + d*x]]*\text{Sqrt}[\text{((a + b*\text{Cos}[c + d*x])*\text{Sec}[c + d*x])/(a + b)}]) + (2*a*((a*\text{Sqrt}[(a...$$

3.409.3 Rubi [A] (verified)

Time = 2.67 (sec) , antiderivative size = 533, normalized size of antiderivative = 1.02, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3468, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx$$

↓ 3468

3.409.
$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx$$

$$\frac{2}{9} \int \frac{3b(2aA + 3bB) \cos^2(c + dx) + (7Aa^2 + 18bBa + 9Ab^2) \cos(c + dx) + a(10Ab + 9aB)}{2 \cos^{\frac{9}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{9} \int \frac{3b(2aA + 3bB) \cos^2(c + dx) + (7Aa^2 + 18bBa + 9Ab^2) \cos(c + dx) + a(10Ab + 9aB)}{\cos^{\frac{9}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{9} \int \frac{3b(2aA + 3bB) \sin(c + dx + \frac{\pi}{2})^2 + (7Aa^2 + 18bBa + 9Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(10Ab + 9aB)}{\sin(c + dx + \frac{\pi}{2})^{9/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 3534

$$\frac{1}{9} \left(\frac{2 \int \frac{4ab(10Ab + 9aB) \cos^2(c + dx) + a(45Ba^2 + 92Aba + 63b^2B) \cos(c + dx) + a(49Aa^2 + 72bBa + 3Ab^2)}{2 \cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{7a} + \frac{2(9aB + 10Ab) \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{\int \frac{4ab(10Ab + 9aB) \cos^2(c + dx) + a(45Ba^2 + 92Aba + 63b^2B) \cos(c + dx) + a(49Aa^2 + 72bBa + 3Ab^2)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{7a} + \frac{2(9aB + 10Ab) \sin(c + dx)}{7d \cos^{\frac{7}{2}}(c + dx)} \right) + \frac{2aA \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 3042

3.409. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$

$$\frac{1}{9} \left(\frac{\int \frac{4ab(10Ab+9aB) \sin(c+dx+\frac{\pi}{2})^2 + a(45Ba^2+92Aba+63b^2B) \sin(c+dx+\frac{\pi}{2}) + a(49Aa^2+72bBa+3Ab^2) dx}{\sin(c+dx+\frac{\pi}{2})^{7/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{7a} + \frac{2(9aB+10Ab) \sin(c+dx+\frac{\pi}{2})}{7d \cos^2(c+dx+\frac{\pi}{2})} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3534

$$\frac{1}{9} \left(\frac{2 \int \frac{(147Aa^2+396bBa+209Ab^2) \cos(c+dx)a^2 + 2b(49Aa^2+72bBa+3Ab^2) \cos^2(c+dx)a + 3(75Ba^3+88Aba^2+9b^2Ba-4Ab^3)a}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a}}{7a} + \frac{2(49a^2A+72abB+3Ab^2)}{5d \cos^2(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{\int \frac{(147Aa^2+396bBa+209Ab^2) \cos(c+dx)a^2 + 2b(49Aa^2+72bBa+3Ab^2) \cos^2(c+dx)a + 3(75Ba^3+88Aba^2+9b^2Ba-4Ab^3)a}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a}}{7a} + \frac{2(49a^2A+72abB+3Ab^2)}{5d \cos^2(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{\int \frac{(147Aa^2+396bBa+209Ab^2) \sin(c+dx+\frac{\pi}{2})a^2 + 2b(49Aa^2+72bBa+3Ab^2) \sin(c+dx+\frac{\pi}{2})^2 a + 3(75Ba^3+88Aba^2+9b^2Ba-4Ab^3)a}{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5a}}{7a} + \frac{2(49a^2A+72abB+3Ab^2)}{5d \cos^2(c+dx)} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3534

3.409. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{2 \int \frac{3((75Ba^3 + 186Aba^2 + 153b^2Ba + 2Ab^3) \cos(c+dx)a^2 + (147Aa^4 + 246bBa^3 + 33Ab^2a^2 - 18b^3Ba + 8Ab^4)a) dx}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}}{3a} + \frac{2(75a^3B + 88a^2Ab + 9ab^2B - 4Ab^3) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} \right) \frac{5a}{7a}$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\int \frac{(75Ba^3 + 186Aba^2 + 153b^2Ba + 2Ab^3) \cos(c+dx)a^2 + (147Aa^4 + 246bBa^3 + 33Ab^2a^2 - 18b^3Ba + 8Ab^4)a}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx + \frac{2(75a^3B + 88a^2Ab + 9ab^2B - 4Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)} \right) \frac{5a}{7a}$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\int \frac{(75Ba^3 + 186Aba^2 + 153b^2Ba + 2Ab^3) \sin(c+dx + \frac{\pi}{2})a^2 + (147Aa^4 + 246bBa^3 + 33Ab^2a^2 - 18b^3Ba + 8Ab^4)a}{\sin(c+dx + \frac{\pi}{2})^{\frac{3}{2}} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \frac{2(75a^3B + 88a^2Ab + 9ab^2B - 4Ab^3) \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} \right) \frac{5a}{7a}$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3477

$$\left(\frac{1}{9} \int \frac{a(a-b)(-3a^3(49A-25B)+a^2(39Ab-171bB)+6ab^2(A-3B)+8Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + a(147a^4A+246a^3bB+33a^2Ab^2-18ab^3B+8Ab^4) \int \frac{1}{\cos(c+dx)} dx}{5a} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\left(\frac{1}{9} \int \frac{a(a-b)(-3a^3(49A-25B)+a^2(39Ab-171bB)+6ab^2(A-3B)+8Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + a(147a^4A+246a^3bB+33a^2Ab^2-18ab^3B+8Ab^4) \int \frac{1}{\sin(c+dx+\frac{\pi}{2})} dx}{5a} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3295

$$\left(\frac{1}{9} \int \frac{a(147a^4A+246a^3bB+33a^2Ab^2-18ab^3B+8Ab^4) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(-3a^3(49A-25B)+a^2(39Ab-171bB)+6ab^2(A-3B)+8Ab^3)}{a} \int \frac{1}{\sin(c+dx+\frac{\pi}{2})} dx}{a} \right)$$

$$\frac{2aA \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3473

$$\frac{1}{9} \left(\frac{2(49a^2A+72abB+3Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} + \frac{2(75a^3B+88a^2Ab+9ab^2B-4Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)} + \frac{2(a-b) \sqrt{a+b} (-3a^3(49A-25B) + 6a^2b(39A-171B) + 6ab^2(A-3B) + a^2(39Ab-171bB)) \cot(c+dx) \operatorname{EllipticF}[\operatorname{ArcSin}[\sqrt{a+b \cos(c+dx)}] / (\sqrt{a+b} \sqrt{\cos(c+dx)})], -((a+b)/(a-b))] \sqrt{(a(1-\sec(c+dx)))/(a+b)} \sqrt{(a(1+\sec(c+dx)))/(a-b))}}{9d \cos^{\frac{9}{2}}(c+dx)} \right)$$

```
input Int[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2), x]
```

```
output (2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + (2*(10*A*b + 9*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(49*a^2*A + 3*A*b^2 + 72*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(147*a^4*A + 33*a^2*A*b^2 + 8*A*b^4 + 246*a^3*b*B - 18*a*b^3*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*(a - b)*Sqrt[a + b]*(8*A*b^3 - 3*a^3*(49*A - 25*B) + 6*a*b^2*(A - 3*B) + a^2*(39*A*b - 171*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/a + (2*(88*a^2*A*b - 4*A*b^3 + 75*a^3*B + 9*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)))/(5*a))/(7*a))/9
```

3.409.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.409.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5952 vs. $2(478) = 956$.

Time = 30.80 (sec) , antiderivative size = 5953, normalized size of antiderivative = 11.40

method	result	size
parts	Expression too large to display	5953
default	Expression too large to display	6034

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x,method=_RE
TURNVERBOSE)
```

```
output result too large to display
```

3.409.5 Fricas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{11/2}} dx$$

```
input integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, alg
orithm="fricas")
```

3.409. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(11/2), x)`

3.409.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(11/2),x)`

output `Timed out`

3.409.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{11/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(11/2), x)`

3.409.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\cos(dx + c)^{11/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/cos(d*x + c)^(11/2), x)`

3.409. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

3.409.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\cos(c + dx)^{11/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(11/2),
x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/cos(c + d*x)^(11/2),
x)`

3.410 $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.410.1 Optimal result	3802
3.410.2 Mathematica [C] (warning: unable to verify)	3803
3.410.3 Rubi [A] (verified)	3804
3.410.4 Maple [B] (warning: unable to verify)	3812
3.410.5 Fricas [F(-1)]	3812
3.410.6 Sympy [F(-1)]	3813
3.410.7 Maxima [F]	3813
3.410.8 Giac [F]	3813
3.410.9 Mupad [F(-1)]	3814

3.410.1 Optimal result

Integrand size = 35, antiderivative size = 779

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx =$$

$$\frac{(a-b)\sqrt{a+b}(150a^3Ab+2840aAb^3-45a^4B+1692a^2b^2B+1024b^4B) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{1920ab^2d}$$

$$-\frac{\sqrt{a+b}(45a^4B-30a^3b(5A+B)-16b^4(45A+64B)-8ab^3(355A+193B)-4a^2b^2(295A+423B)) \cot(c+dx)}{1920b^2d}$$

$$+\frac{\sqrt{a+b}(10a^4Ab-240a^2Ab^3-96Ab^5-3a^5B-40a^3b^2B-240ab^4B) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{128b^3d}$$

$$+\frac{(150a^3Ab+2840aAb^3-45a^4B+1692a^2b^2B+1024b^4B) \sqrt{a+b}\cos(c+dx) \sin(c+dx)}{1920b^2d\sqrt{\cos(c+dx)}}$$

$$+\frac{(50a^2Ab+120Ab^3-15a^3B+172ab^2B) \sqrt{\cos(c+dx)}\sqrt{a+b}\cos(c+dx) \sin(c+dx)}{320bd}$$

$$+\frac{(50aAb-15a^2B+64b^2B) \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2} \sin(c+dx)}{240bd}$$

$$+\frac{(10Ab-3aB) \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2} \sin(c+dx)}{40bd}$$

$$+\frac{B\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{7/2} \sin(c+dx)}{5bd}$$

3.410. $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

output $\frac{1}{240}(50Aab-15B^2a^2+64B^2b^2)(a+b\cos(dx+c))^{3/2}\sin(dx+c)\cos(dx+c)^{1/2}/b/d+1/40(10Ab-3B^2a)(a+b\cos(dx+c))^{5/2}\sin(dx+c)\cos(dx+c)^{1/2}/b/d+1/5B(a+b\cos(dx+c))^{7/2}\sin(dx+c)\cos(dx+c)^{1/2}/b/d+1/1920(150A^3b+2840A^2ab^3-45B^2a^4+1692B^2a^2b^2+1024B^2b^4)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^2/d/\cos(dx+c)^{1/2}+1/320(50A^2b+120A^2b^3-15B^2a^3+172B^2ab^2)\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/b/d-1/1920(a-b)(150A^3b+2840A^2ab^3-45B^2a^4+1692B^2a^2b^2+1024B^2b^4)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/b^2/d-1/1920(45B^2a^4-30a^3b(5A+B)-16b^4(45A+64B)-8a^2b^3(355A+193B)-4a^2b^2(295A+423B))\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d+1/128(10A^4b-240A^2b^3-96A^2b^5-3B^2a^5-40B^2a^3b^2-240B^2ab^4)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^3/d$

3.410.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.57 (sec) , antiderivative size = 1353, normalized size of antiderivative = 1.74

$$\int \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx = \text{Too large to display}$$

input `Integrate[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output

```

-1/3840*((-4*a*(-1330*a^3*A*b - 3560*a*A*b^3 + 15*a^4*B - 3236*a^2*b^2*B -
1024*b^4*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*C
os[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*
x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(
c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*
Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-6440*a^2*A*b^2 - 1440
*A*b^4 - 2292*a^3*b*B - 4624*a*b^3*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/
(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a +
b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[
((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Si
n[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) -
(Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*
Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*
Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(c +
d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[
c + d*x]]*Sqrt[a + b*Cos[c + d*x]])) + 2*(-150*a^3*A*b - 2840*a*A*b^3 + 45
*a^4*B - 1692*a^2*b^2*B - 1024*b^4*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[
c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)
/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a
 + b)*Cos[c + d*x])*Sec[c + d*x])/(a + b)) + (2*a*((a*Sqrt[((a + b)*Cot...

```

3.410.3 Rubi [A] (verified)

Time = 4.33 (sec) , antiderivative size = 786, normalized size of antiderivative = 1.01, number of steps used = 23, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.657$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3469}
 \end{aligned}$$

3.410. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

$$\begin{aligned}
& \frac{\int \frac{(a+b \cos(c+dx))^{5/2}((10Ab-3aB) \cos^2(c+dx)+8bB \cos(c+dx)+aB) dx}{2\sqrt{\cos(c+dx)}}}{\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{7/2}}{5bd}} + \\
& \quad \downarrow 27 \\
& \frac{\int \frac{(a+b \cos(c+dx))^{5/2}((10Ab-3aB) \cos^2(c+dx)+8bB \cos(c+dx)+aB) dx}{\sqrt{\cos(c+dx)}}}{\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{7/2}}{5bd}} + \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(a+b \sin(c+dx+\frac{\pi}{2}))^{5/2}((10Ab-3aB) \sin^2(c+dx+\frac{\pi}{2})+8bB \sin(c+dx+\frac{\pi}{2})+aB) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{7/2}}{5bd}} + \\
& \quad \downarrow 3528 \\
& \frac{\frac{1}{4} \int \frac{(a+b \cos(c+dx))^{3/2}((-15Ba^2+50Aba+64b^2B) \cos^2(c+dx)+6b(10Ab+9aB) \cos(c+dx)+5a(2Ab+aB)) dx}{2\sqrt{\cos(c+dx)}} + \frac{(10Ab-3aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d}}{\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{7/2}}{5bd}} \\
& \quad \downarrow 27 \\
& \frac{\frac{1}{8} \int \frac{(a+b \cos(c+dx))^{3/2}((-15Ba^2+50Aba+64b^2B) \cos^2(c+dx)+6b(10Ab+9aB) \cos(c+dx)+5a(2Ab+aB)) dx}{\sqrt{\cos(c+dx)}} + \frac{(10Ab-3aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d}}{\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{7/2}}{5bd}} \\
& \quad \downarrow 3042 \\
& \frac{\frac{1}{8} \int \frac{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}((-15Ba^2+50Aba+64b^2B) \sin^2(c+dx+\frac{\pi}{2})+6b(10Ab+9aB) \sin(c+dx+\frac{\pi}{2})+5a(2Ab+aB)) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} + \frac{(10Ab-3aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{4d}}{\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{7/2}}{5bd}} \\
& \quad \downarrow 3528
\end{aligned}$$

3.410. $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

$$\frac{1}{8} \left(\frac{1}{3} \int \frac{\sqrt{a+b \cos(c+dx)} (3(-15Ba^3+50Aba^2+172b^2Ba+120Ab^3) \cos^2(c+dx)+2b(147Ba^2+310Aba+128b^2B) \cos(c+dx)+a(15Ba^2+110Aba+12b^2B))}{2\sqrt{\cos(c+dx)}} dx \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \int \frac{\sqrt{a+b \cos(c+dx)} (3(-15Ba^3+50Aba^2+172b^2Ba+120Ab^3) \cos^2(c+dx)+2b(147Ba^2+310Aba+128b^2B) \cos(c+dx)+a(15Ba^2+110Aba+12b^2B))}{\sqrt{\cos(c+dx)}} dx \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (3(-15Ba^3+50Aba^2+172b^2Ba+120Ab^3) \sin(c+dx+\frac{\pi}{2})^2+2b(147Ba^2+310Aba+128b^2B) \sin(c+dx+\frac{\pi}{2})+a(15Ba^2+110Aba+12b^2B))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3528

$$\frac{1}{8} \left(\frac{1}{2} \int \frac{(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B) \cos^2(c+dx)+2b(573Ba^3+1610Aba^2+1156b^2Ba+360Ab^3) \cos(c+dx)+a(15Ba^2+110Aba+12b^2B)}{2\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{4} \int \frac{(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B) \cos^2(c+dx)+2b(573Ba^3+1610Aba^2+1156b^2Ba+360Ab^3) \cos(c+dx)+a(15Ba^2+110Aba+12b^2B)}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{4} \int \frac{(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B) \sin(c+dx+\frac{\pi}{2})^2+2b(573Ba^3+1610Aba^2+1156b^2Ba+360Ab^3) \sin(c+dx+\frac{\pi}{2})+a(15Ba^2+110Aba+12b^2B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

3.410. $\int \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) dx$

↓ 3540

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\int -\frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)\cos^2(c+dx)-2ab(15Ba^3+590Aba^2+772b^2Ba+360Ab^3)\cos(c+dx)+a(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} - \frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)}{2b} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 25

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)}{2b} \right) \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)}{2b} \right) \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3532

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{a(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B)\cos^{\frac{3}{2}}(c+dx)+a(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \right) \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} \right) - \int \frac{a(-45Ba^4 + 150Aba^3 + 1692b^2Ba^2 + 2840Ab^3a)}{\sin(c+dx)} dx \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3288

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} \right) - \int \frac{a(-45Ba^4 + 150Aba^3 + 1692b^2Ba^2 + 2840Ab^3a)}{\sin(c+dx)} dx \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3477

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} \right) - \frac{a(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B)}{\sin(c+dx)} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} \right) - \frac{a(45a^4B - 30a^3b(5A+B) - 4a^2b^2(295A + 423B))}{\sin(c+dx)} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3295

3.410. $\int \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) dx$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-45a^4B + 150a^3Ab + 1692a^2b^2B + 2840aAb^3 + 1024b^4B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{2d} \right) \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

↓ 3473

$$\frac{1}{8} \left(\frac{(-15a^2B + 50aAb + 64b^2B) \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3d} + \frac{1}{6} \left(\frac{3(-15a^3B + 50a^2Ab + 172ab^2B + 120Ab^3) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d} \right) \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{7/2}}{5bd}$$

input `Int[Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

output `(B*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(5*b*d) + ((10*A*b - 3*a*B)*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(4*d) + (((50*a*A*b - 15*a^2*B + 64*b^2*B)*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + ((3*(50*a^2*A*b + 120*A*b^3 - 15*a^3*B + 172*a*b^2*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*Sqrt[a + b]*(150*a^3*A*b + 2840*a*A*b^3 - 45*a^4*B + 1692*a^2*b^2*B + 1024*b^4*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]), -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/(a*d) + (2*Sqrt[a + b]*(45*a^4*B - 30*a^3*b*(5*A + B) - 16*b^4*(45*A + 64*B) - 8*a*b^3*(355*A + 193*B) - 4*a^2*b^2*(295*A + 423*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/d - (30*Sqrt[a + b]*(10*a^4*A*b - 240*a^2*A*b^3 - 96*A*b^5 - 3*a^5*B - 40*a^3*b^2*B - 240*a*b^4*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(b*d))/b + ((150*a^3*A*b + 2840*a*A*b^3 - 45*a^4*B + 1692*a^2*b^2*B + 1024*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/4)/6)/8)/(10*b)`

3.410. $\int \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) dx$

3.410.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3469 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(b \sin(e + fx) + f x)^{3/2}} dx$$

$$\rightarrow \text{Simp}[-2 A (c - d) (\tan(e + fx) / (f b c^2)) \text{Rt}[(c + d)/b, 2] \sqrt{c((1 + \text{Csc}(e + fx)) / (c - d))} \sqrt{c((1 - \text{Csc}(e + fx)) / (c + d))} \text{EllipticE}[\text{ArcSin}[\sqrt{c + d \sin(e + fx)}] / \sqrt{b \sin(e + fx)}] / \text{Rt}[(c + d)/b, 2], -(c + d)/(c - d)], x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\& \text{PosQ}[(c + d)/b]$$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[(A - B)/(a - b) \int [1 / (\sqrt{a + b \sin(e + fx)} \sqrt{c + d \sin(e + fx)})], x], x] - \text{Simp}[(A b - a B)/(a - b) \int [(1 + \sin(e + fx)) / ((a + b \sin(e + fx))^{3/2} \sqrt{c + d \sin(e + fx)})], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[A, B]$$`

rule 3528 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +

$$\int (a + b \sin(e + fx))^m ((c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin(e + fx) + f x)^2) dx$$

$$\rightarrow \text{Simp}[-C \cos(e + fx) (a + b \sin(e + fx))^m ((c + d \sin(e + fx))^{n+1} / (d f (m + n + 2))), x] + \text{Simp}[1 / (d (m + n + 2)) \int [(a + b \sin(e + fx))^{m-1} (c + d \sin(e + fx))^n \text{Simp}[a A d (m + n + 2) + C (b c m + a d (n + 1)) + (d (A b + a B) (m + n + 2) - C (a c - b d (m + n + 1))) \sin(e + fx) + (C (a d m - b c (m + 1)) + b B d (m + n + 2)) \sin^2(e + fx)], x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[m, 0] \&\& (!\text{IGtQ}[n, 0] \&\& (!\text{IntegerQ}[m] || (\text{EqQ}[a, 0] \&\& \text{NeQ}[c, 0])))$$`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2)/

$$\int \frac{(A + B \sin(e + fx) + C \sin^2(e + fx)) \sqrt{c + d \sin(e + fx)}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[C/b^2 \int [\sqrt{a + b \sin(e + fx)}] / \sqrt{c + d \sin(e + fx)}, x], x] + \text{Simp}[1/b^2 \int [(A b^2 - a^2 C + b (b B - 2 a C) \sin(e + fx)) / ((a + b \sin(e + fx))^{3/2} \sqrt{c + d \sin(e + fx)})], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$$`

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])]*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.410.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 7109 vs. $2(719) = 1438$.

Time = 20.89 (sec) , antiderivative size = 7110, normalized size of antiderivative = 9.13

method	result	size
parts	Expression too large to display	7110
default	Expression too large to display	7197

```
input int(cos(d*x+c)^(3/2)*(a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c)),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.410.5 Fracas [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algo
rithm="fracas")
```

```
output Timed out
```

3.410. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

3.410.6 Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.410.7 Maxima [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(3/2), x)`

3.410.8 Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(3/2), x)`

3.410. $\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

3.410.9 Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx = \int \cos(c + dx)^{3/2} (A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2} dx$$

input `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2),x)`output `int(cos(c + d*x)^(3/2)*(A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2), x)`

3.411 $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

3.411.1 Optimal result	3815
3.411.2 Mathematica [C] (warning: unable to verify)	3816
3.411.3 Rubi [A] (verified)	3817
3.411.4 Maple [B] (warning: unable to verify)	3825
3.411.5 Fricas [F]	3826
3.411.6 Sympy [F(-1)]	3826
3.411.7 Maxima [F]	3827
3.411.8 Giac [F]	3827
3.411.9 Mupad [F(-1)]	3827

3.411.1 Optimal result

Integrand size = 35, antiderivative size = 664

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx =$$

$$\frac{(a - b)\sqrt{a + b}(264a^2Ab + 128Ab^3 + 15a^3B + 284ab^2B) \cot(c + dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\cos(c + dx)}}{192abd}$$

$$+ \frac{\sqrt{a + b}(15a^3B + 8b^3(16A + 9B) + 2a^2b(132A + 59B) + 4ab^2(52A + 71B)) \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{192bd}$$

$$- \frac{\sqrt{a + b}(40a^3Ab + 160aAb^3 - 5a^4B + 120a^2b^2B + 48b^4B) \cot(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{64b^2d}$$

$$+ \frac{(264a^2Ab + 128Ab^3 + 15a^3B + 284ab^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{192bd\sqrt{\cos(c + dx)}}$$

$$+ \frac{(24aAb + 5a^2B + 12b^2B) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{32d}$$

$$+ \frac{(8Ab + 11aB) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{24d}$$

$$+ \frac{bB \cos^{3/2}(c + dx)(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{4d}$$

output $\frac{1}{4}bB\cos(dx+c)^{3/2}(a+b\cos(dx+c))^{3/2}\sin(dx+c)/d+1/24*(8A*b+11*B*a)*(a+b\cos(dx+c))^{3/2}\sin(dx+c)\cos(dx+c)^{1/2}/d+1/192*(264A*a^2*b+128A*b^3+15B*a^3+284B*a*b^2)*\sin(dx+c)*(a+b\cos(dx+c))^{1/2}/b/d/\cos(dx+c)^{1/2}+1/32*(24A*a*b+5B*a^2+12B*b^2)*\sin(dx+c)\cos(dx+c)^{(1/2)*(a+b\cos(dx+c))^{1/2}/d-1/192*(a-b)*(264A*a^2*b+128A*b^3+15B*a^3+284B*a*b^2)*\cot(dx+c)*\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a*(1+\sec(dx+c))/(a+b))^{1/2}/a/b/d+1/192*(15B*a^3+8*b^3*(16A+9B)+2*a^2*b*(132A+59B)+4*a*b^2*(52A+71B))*\cot(dx+c)*\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a+b))^{1/2}/b/d-1/64*(40A*a^3*b+160A*a*b^3-5B*a^4+120B*a^2*b^2+48B*b^4)*\cot(dx+c)*\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b,((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d$

3.411.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.50 (sec) , antiderivative size = 1287, normalized size of antiderivative = 1.94

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \text{Too large to display}$$

input `Integrate[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

```
output ((-4*a*(472*a^2*A*b + 128*A*b^3 + 133*a^3*B + 356*a*b^2*B)*Sqrt[((a + b)*C
ot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]
^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*Ell
ipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]],
(-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a +
b*Cos[c + d*x]]) - 4*a*(384*a^3*A + 608*a*A*b^2 + 644*a^2*b*B + 144*b^3*B)
*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x
]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a
]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]
^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c
+ d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-
a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*
Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSi
n[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a +
b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])
+ 2*(264*a^2*A*b + 128*A*b^3 + 15*a^3*B + 284*a*b^2*B)*((I*Cos[(c + d*x)/2
]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c
+ d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c
+ d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqr
t[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*C...
```

3.411.3 Rubi [A] (verified)

Time = 3.53 (sec) , antiderivative size = 672, normalized size of antiderivative = 1.01, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

↓ 3042

$$\int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}\left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2}\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

↓ 3469

3.411. $\int \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$

$$\frac{1}{4} \int \frac{1}{2} \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} (b(8Ab+11aB) \cos^2(c+dx) + 2(4Ba^2+8Aba+3b^2B) \cos(c+dx) + a(8Ab+11aB)) \frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d} dx$$

↓ 27

$$\frac{1}{8} \int \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} (b(8Ab+11aB) \cos^2(c+dx) + 2(4Ba^2+8Aba+3b^2B) \cos(c+dx) + a(8Ab+11aB)) \frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d} dx$$

↓ 3042

$$\frac{1}{8} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)} \left(b(8Ab+11aB) \sin\left(c+dx+\frac{\pi}{2}\right)^2 + 2(4Ba^2+8Aba+3b^2B) \sin\left(c+dx+\frac{\pi}{2}\right) + a(8Ab+11aB) \right) \frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d} dx$$

↓ 3528

$$\frac{1}{8} \left(\frac{\int \frac{\sqrt{a+b \cos(c+dx)} (3b(5Ba^2+24Aba+12b^2B) \cos^2(c+dx) + 2b(24Aa^2+31bBa+16Ab^2) \cos(c+dx) + ab(8Ab+11aB))}{2\sqrt{\cos(c+dx)}} dx}{3b} + \frac{(11aB+8Ab)}{3b} \right) \frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(\frac{\int \frac{\sqrt{a+b \cos(c+dx)} (3b(5Ba^2+24Aba+12b^2B) \cos^2(c+dx) + 2b(24Aa^2+31bBa+16Ab^2) \cos(c+dx) + ab(8Ab+11aB))}{\sqrt{\cos(c+dx)}} dx}{6b} + \frac{(11aB+8Ab)}{6b} \right) \frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{\int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})} (3b(5Ba^2+24Aba+12b^2B) \sin(c+dx+\frac{\pi}{2})^2 + 2b(24Aa^2+31bBa+16Ab^2) \sin(c+dx+\frac{\pi}{2}) + ab(8Ab+11aB))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{6b} + \frac{(11aB+8Ab)}{6b} \right) \frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

3.411. $\int \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) dx$

↓ 3528

$$\frac{1}{8} \left(\frac{\frac{1}{2} \int \frac{b(15Ba^3+264Aba^2+284b^2Ba+128Ab^3) \cos^2(c+dx)+2b(96Aa^3+161bBa^2+152Ab^2a+36b^3B) \cos(c+dx)+ab(59Ba^2+104Aba+36b^2B)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{6b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(\frac{\frac{1}{4} \int \frac{b(15Ba^3+264Aba^2+284b^2Ba+128Ab^3) \cos^2(c+dx)+2b(96Aa^3+161bBa^2+152Ab^2a+36b^3B) \cos(c+dx)+ab(59Ba^2+104Aba+36b^2B)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{6b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{\frac{1}{4} \int \frac{b(15Ba^3+264Aba^2+284b^2Ba+128Ab^3) \sin(c+dx+\frac{\pi}{2})^2+2b(96Aa^3+161bBa^2+152Ab^2a+36b^3B) \sin(c+dx+\frac{\pi}{2})+ab(59Ba^2+104Aba+36b^2B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{6b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}}{4d}$$

↓ 3540

$$\frac{1}{8} \left(\frac{\frac{1}{4} \int \frac{-2a(59Ba^2+104Aba+36b^2B) \cos(c+dx)b^2-3(-5Ba^4+40Aba^3+120b^2Ba^2+160Ab^3a+48b^4B) \cos^2(c+dx)+ab(15Ba^3+264Aba^2+284b^2Ba+128Ab^3)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{2b} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}}{4d}$$

↓ 25

$$\frac{1}{8} \left(\frac{\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{-2a(59Ba^2+104Aba+36b^2B) \cos(c+dx)b^2 - 3(-5Ba^4+40Aba^3+120Ab^2a^2) \cos^{\frac{3}{2}}(c+dx)}{\cos^{\frac{3}{2}}(c+dx)} dx \right)}{\dots} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{-2a(59Ba^2+104Aba+36b^2B) \sin(c+dx+\frac{\pi}{2})b^2 - 3(-5Ba^4+40Aba^3+120Ab^2a^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx \right)}{\dots} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3532

$$\frac{1}{8} \left(\frac{\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{ab(15Ba^3+264Aba^2+284b^2Ba+128Ab^3) - 2ab^2(59Ba^2+104Aba+36b^2a)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx \right)}{\dots} \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{ab(15Ba^3+264Aba^2+284b^2Ba+128Ab^3)-2ab^2(59Ba^2+104Aba+36b)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} \right) \right) \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3288

$$\left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{ab(15Ba^3+264Aba^2+284b^2Ba+128Ab^3)-2ab^2(59Ba^2+104Aba+36b)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} \right) \right) \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3477

$$\left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{ab(15a^3B+264a^2Ab+284ab^2B+128Ab^3)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} \right) \right) \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{-ab(15a^3B+2a^2b(132A+59B)+4ab^2(52A+71B)+8b^3(16A+9B)}{\dots} \right) \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3295

$$\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{ab(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \cos(c+dx)}} dx}{\dots} \right) \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{3/2}}{4d}$$

↓ 3473

$$\frac{1}{8} \left(\frac{3b(5a^2B+24aAb+12b^2B) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} \right) \right)$$

$$\frac{bB \sin(c+dx) \cos^{\frac{3}{2}}(c+dx) (a+b \cos(c+dx))^{3/2}}{4d}$$

input `Int[Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]),x]`

```
output (b*B*Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x]/(4*d) + (
((8*A*b + 11*a*B)*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*
x])/(3*d) + ((3*b*(24*a*A*b + 5*a^2*B + 12*b^2*B)*Sqrt[Cos[c + d*x]]*Sqrt[
a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*
(264*a^2*A*b + 128*A*b^3 + 15*a^3*B + 284*a*b^2*B)*Cot[c + d*x]*EllipticE[
ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a +
b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x
]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(15*a^3*B + 8*b^3*(16*A + 9*B) + 2*
a^2*b*(132*A + 59*B) + 4*a*b^2*(52*A + 71*B))*Cot[c + d*x]*EllipticF[ArcSi
n[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a
- b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(
a - b))]/d + (6*Sqrt[a + b]*(40*a^3*A*b + 160*a*A*b^3 - 5*a^4*B + 120*a^2*
b^2*B + 48*b^4*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos
[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*
(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/b +
((264*a^2*A*b + 128*A*b^3 + 15*a^3*B + 284*a*b^2*B)*Sqrt[a + b*Cos[c + d*x
]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])/4)/(6*b))/8
```

3.411.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3528 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*SIN[e + f*x
])^m*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m +
n + 2)) Int[(a + b*SIN[e + f*x])^(m - 1)*(c + d*SIN[e + f*x])^n*Simp[a*A*
d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a
*c - b*d*(m + n + 1)))*SIN[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m +
n + 2))*SIN[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}
, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[
m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2
]/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*SIN[e + f*x]]
/Sqrt[c + d*SIN[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*SIN[e + f*x])/((a + b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2
]/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*SIN[e + f
*x]]/(d*f*Sqrt[a + b*SIN[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*SIN[e + f*x] + (2*b*B*d - C*(b*c + a*d))*SIN[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.411.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 5790 vs. $2(610) = 1220$.

Time = 18.36 (sec) , antiderivative size = 5791, normalized size of antiderivative = 8.72

method	result	size
parts	Expression too large to display	5791
default	Expression too large to display	5861

input `int(cos(d*x+c)^(1/2)*(a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c)),x,method=_RETURNVERBOSE)`

output `result too large to display`

3.411.5 Fricas [F]

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^{5/2}\sqrt{\cos(dx+c)}dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x,algorithm="fricas")`

output `integral((B*b^2*cos(d*x+c)^3+A*a^2+(2*B*a*b+A*b^2)*cos(d*x+c)^2+(B*a^2+2*A*a*b)*cos(d*x+c))*sqrt(b*cos(d*x+c)+a)*sqrt(cos(d*x+c)),x)`

3.411.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(1/2)*(a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c)),x)`

output `Timed out`

3.411.7 Maxima [F]

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^{5/2}\sqrt{\cos(dx+c)}dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x+c)+A)*(b*cos(d*x+c)+a)^(5/2)*sqrt(cos(d*x+c)),x)`

3.411.8 Giac [F]

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \int (B\cos(dx+c)+A)(b\cos(dx+c)+a)^{5/2}\sqrt{\cos(dx+c)}dx$$

input `integrate(cos(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x+c)+A)*(b*cos(d*x+c)+a)^(5/2)*sqrt(cos(d*x+c)),x)`

3.411.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx = \int \sqrt{\cos(c+dx)}(A+B\cos(c+dx))(a+b\cos(c+dx))^{5/2}dx$$

input `int(cos(c+d*x)^(1/2)*(A+B*cos(c+d*x))*(a+b*cos(c+d*x))^(5/2),x)`

output `int(cos(c+d*x)^(1/2)*(A+B*cos(c+d*x))*(a+b*cos(c+d*x))^(5/2),x)`

3.411. $\int \sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))dx$

$$3.412 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

3.412.1 Optimal result	3828
3.412.2 Mathematica [C] (warning: unable to verify)	3829
3.412.3 Rubi [A] (verified)	3830
3.412.4 Maple [B] (warning: unable to verify)	3837
3.412.5 Fricas [F]	3838
3.412.6 Sympy [F(-1)]	3838
3.412.7 Maxima [F]	3838
3.412.8 Giac [F]	3839
3.412.9 Mupad [F(-1)]	3839

3.412.1 Optimal result

Integrand size = 35, antiderivative size = 564

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}(54aAb+33a^2B+16b^2B) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\cos(c+dx)}}{24ad}$$

$$+ \frac{\sqrt{a+b}(4b^2(3A+4B)+a^2(48A+33B)+a(54Ab+26bB)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{24d}$$

$$- \frac{\sqrt{a+b}(30a^2Ab+8Ab^3+5a^3B+20ab^2B) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\cos(c+dx)}}{8bd}$$

$$+ \frac{(54aAb+33a^2B+16b^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{24d\sqrt{\cos(c+dx)}}$$

$$+ \frac{b(2Ab+3aB) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{4d}$$

$$+ \frac{bB \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{3d}$$

$$3.412. \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

output $\frac{1}{3}bB(a+b\cos(dx+c))^{3/2}\sin(dx+c)\cos(dx+c)^{1/2}/d + \frac{1}{24}(54Aab+33Bb^2+16Bb^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\cos(dx+c)^{1/2} + \frac{1}{4}b(2Aa+3Ba)\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/d - \frac{1}{24}(a-b)(54Aab+33Ba^2+16Bb^2)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a + \frac{1}{24}(4b^2(3A+4B)+a^2(48A+33B)+a(54Ab+26Bb))\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/d - \frac{1}{8}(30Aa^2b+8Ab^3+5Ba^3+20Bab^2)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}\cos(dx+c)^{1/2}, (a+b)/b, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b/d$

3.412.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.47 (sec) , antiderivative size = 1251, normalized size of antiderivative = 2.22

$$\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sqrt[Cos[c + d*x]], x]`

output $((-4*a*(48*a^3*A + 66*a*A*b^2 + 59*a^2*b*B + 16*b^3*B)*\text{Sqrt}[(a + b)*\text{Cot}[(c + d*x)/2]^2]/(-a + b))*\text{Sqrt}[-((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a))*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a]*\text{Csc}[c + d*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a]/\text{Sqrt}[2]], (-2*a)/(-a + b))*\text{Sin}[(c + d*x)/2]^4)/((a + b)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) - 4*a*(144*a^2*A*b + 24*A*b^3 + 48*a^3*B + 76*a*b^2*B)*((\text{Sqrt}[(a + b)*\text{Cot}[(c + d*x)/2]^2]/(-a + b))*\text{Sqrt}[-((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a))*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a]*\text{Csc}[c + d*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a]/\text{Sqrt}[2]], (-2*a)/(-a + b))*\text{Sin}[(c + d*x)/2]^4)/((a + b)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) - (\text{Sqrt}[(a + b)*\text{Cot}[(c + d*x)/2]^2]/(-a + b))*\text{Sqrt}[-((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a))*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a]*\text{Csc}[c + d*x]*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2)/a]/\text{Sqrt}[2]], (-2*a)/(-a + b))*\text{Sin}[(c + d*x)/2]^4)/(b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]]) + 2*(54*a*A*b^2 + 33*a^2*b*B + 16*b^3*B)*((I*\text{Cos}[(c + d*x)/2]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])*\text{EllipticE}[I*\text{ArcSinh}[\text{Sin}[(c + d*x)/2]/\text{Sqrt}[\text{Cos}[c + d*x]]], (-2*a)/(-a - b))*\text{Sec}[c + d*x])/(b*\text{Sqrt}[\text{Cos}[(c + d*x)/2]^2*\text{Sec}[c + d*x]]*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Sec}[c + d*x])/(a + b)) + (2*a*((a*\text{Sqrt}[(a + b)*\text{Cot}[(c + d*x)/2]^2]/(-a + b))*\text{Sqrt}[-((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2)/a]...$

3.412.3 Rubi [A] (verified)

Time = 2.93 (sec) , antiderivative size = 572, normalized size of antiderivative = 1.01, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3469, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3469}$$

3.412. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$

$$\frac{1}{3} \int \frac{\sqrt{a + b \cos(c + dx)} (3b(2Ab + 3aB) \cos^2(c + dx) + 2(3Ba^2 + 6Aba + 2b^2B) \cos(c + dx) + a(6aA + bB))}{2\sqrt{\cos(c + dx)}} dx - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \downarrow 27$$

$$\frac{1}{6} \int \frac{\sqrt{a + b \cos(c + dx)} (3b(2Ab + 3aB) \cos^2(c + dx) + 2(3Ba^2 + 6Aba + 2b^2B) \cos(c + dx) + a(6aA + bB))}{\sqrt{\cos(c + dx)}} dx - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \downarrow 3042$$

$$\frac{1}{6} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (3b(2Ab + 3aB) \sin^2(c + dx + \frac{\pi}{2}) + 2(3Ba^2 + 6Aba + 2b^2B) \sin(c + dx + \frac{\pi}{2}) + a(6aA + bB))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \downarrow 3528$$

$$\frac{1}{6} \left(\frac{1}{2} \int \frac{b(33Ba^2 + 54Aba + 16b^2B) \cos^2(c + dx) + 2(12Ba^3 + 36Aba^2 + 19b^2Ba + 6Ab^3) \cos(c + dx) + a(24Aa^2 + 12AbA + 4b^2B)}{2\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 27$$

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{b(33Ba^2 + 54Aba + 16b^2B) \cos^2(c + dx) + 2(12Ba^3 + 36Aba^2 + 19b^2Ba + 6Ab^3) \cos(c + dx) + a(24Aa^2 + 12AbA + 4b^2B)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{4} \int \frac{b(33Ba^2 + 54Aba + 16b^2B) \sin^2(c + dx + \frac{\pi}{2}) + 2(12Ba^3 + 36Aba^2 + 19b^2Ba + 6Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(24Aa^2 + 12AbA + 4b^2B)}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 3540$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int -\frac{3b(5Ba^3+30Aba^2+20b^2Ba+8Ab^3) \cos^2(c+dx) - 2ab(24Aa^2+13bBa+6Ab^2) \cos(c+dx) + ab(33Ba^2+54Aba+16b^2B)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{2b} + \right. \right. \quad (33)$$

$$\left. \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3d} \right)$$

↓ 25

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int -\frac{3b(5Ba^3+30Aba^2+20b^2Ba+8Ab^3) \cos^2(c+dx) - 2ab(24Aa^2+13bBa+6Ab^2) \cos(c+dx) + ab(33Ba^2+54Aba+16b^2B)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{\sin(c+dx)} \right. \right.$$

$$\left. \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int -\frac{3b(5Ba^3+30Aba^2+20b^2Ba+8Ab^3) \sin(c+dx+\frac{\pi}{2})^2 - 2ab(24Aa^2+13bBa+6Ab^2) \sin(c+dx+\frac{\pi}{2}) + ab(33Ba^2+54Aba+16b^2B)}{\sin(c+dx+\frac{\pi}{2}) \sqrt{a+b \cos(c+dx)}} dx}{\sin(c+dx)} \right. \right.$$

$$\left. \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3d} \right)$$

↓ 3532

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(33Ba^2+54Aba+16b^2B) - 2ab(24Aa^2+13bBa+6Ab^2) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{\sin(c+dx)} \right. \right.$$

$$\left. \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3d} \right)$$

↓ 3042

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(33Ba^2+54Aba+16b^2B) - 2ab(24Aa^2+13bBa+6Ab^2) \sin(c+dx+\frac{\pi}{2})^2 - 2ab(24Aa^2+13bBa+6Ab^2) \sin(c+dx+\frac{\pi}{2}) + ab(33Ba^2+54Aba+16b^2B)}{\sin(c+dx+\frac{\pi}{2})^2 \sqrt{a+b \cos(c+dx)}} dx}{\sin(c+dx)} \right. \right.$$

$$\left. \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)} (a+b \cos(c+dx))^{3/2}}{3d} \right)$$

↓ 3288

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \int \frac{ab(33Ba^2 + 54Aba + 16b^2B) - 2ab(24Aa^2 + 13bBa + 6Ab^2)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 3477$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{ab(33a^2B + 54aAb + 16b^2B) \int \frac{\cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} \right) - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 3042$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{-ab(a^2(48A + 33B) + a(54Ab + 26bB) + 4b^2A)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} \right) - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 3295$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{ab(33a^2B + 54aAb + 16b^2B) \int \frac{\sin(c + dx)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \right) - \frac{bB \sin(c + dx) \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2}}{3d} \right) \downarrow 3473$$

$$\frac{1}{6} \left(\frac{1}{4} \left(\frac{(33a^2B + 54aAb + 16b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{2b\sqrt{a+b}(a^2(48A+33B)+a(54Ab+26bB)+4b^2(3A+4B))}{3d} \right) \right)$$

input `Int[((a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]))/Sqrt[Cos[c + d*x]],x]`

output `(b*B*Sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + (3*b*(2*A*b + 3*a*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*(54*a*A*b + 33*a^2*B + 16*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(4*b^2*(3*A + 4*B) + a^2*(48*A + 33*B) + a*(54*A*b + 26*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/d + (6*Sqrt[a + b]*(30*a^2*A*b + 8*A*b^3 + 5*a^3*B + 20*a*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/d)/b + ((54*a*A*b + 33*a^2*B + 16*b^2*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/4)/6`

3.412.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] :> Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.412.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/sqrt(cos(d*x + c)), x)`

3.412.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(1/2),x)`

output `Timed out`

3.412.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/sqrt(cos(d*x + c)), x)`

3.412.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sqrt{\cos(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/sqrt(cos(d*x + c)), x)`

3.412.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(1/2),x)`

$$3.413 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

3.413.1 Optimal result	3840
3.413.2 Mathematica [C] (verified)	3841
3.413.3 Rubi [A] (verified)	3842
3.413.4 Maple [B] (warning: unable to verify)	3848
3.413.5 Fricas [F(-1)]	3849
3.413.6 Sympy [F(-1)]	3850
3.413.7 Maxima [F]	3850
3.413.8 Giac [F]	3850
3.413.9 Mupad [F(-1)]	3851

3.413.1 Optimal result

Integrand size = 35, antiderivative size = 547

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{(a-b)\sqrt{a+b}(8a^2A-4Ab^2-9abB) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a-b}{a+b}\right) + \sqrt{a+b}(8a^2(A-B)-2b^2(2A+B)-3ab(8A+3B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a-b}{a+b}\right) - \sqrt{a+b}(20aAb+15a^2B+4b^2B) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4d} - \frac{(8a^2A-4Ab^2-9abB) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{4d \sqrt{\cos(c+dx)}} - \frac{b(4aA-bB) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{2d} + \frac{2aA(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{d \sqrt{\cos(c+dx)}}$$

output $2*a*A*(a+b*\cos(d*x+c))^(3/2)*\sin(d*x+c)/d/\cos(d*x+c)^(1/2)-1/4*(8*A*a^2-4*A*b^2-9*B*a*b)*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)-1/2*b*(4*A*a-B*b)*\sin(d*x+c)*\cos(d*x+c)^(1/2)*(a+b*\cos(d*x+c))^(1/2)/d+1/4*(a-b)*(8*A*a^2-4*A*b^2-9*B*a*b)*\cot(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^(1/2)/(a+b)^(1/2)/\cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-\sec(d*x+c)))/(a+b))^(1/2)*(a*(1+\sec(d*x+c)))/(a-b))^(1/2)/a/d-1/4*(8*a^2*(A-B)-2*b^2*(2*A+B)-3*a*b*(8*A+3*B))*\cot(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^(1/2)/(a+b)^(1/2)/\cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-\sec(d*x+c)))/(a+b))^(1/2)*(a*(1+\sec(d*x+c)))/(a-b))^(1/2)/d-1/4*(20*A*a*b+15*B*a^2+4*B*b^2)*\cot(d*x+c)*\text{EllipticPi}((a+b*\cos(d*x+c))^(1/2)/(a+b)^(1/2)/\cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-\sec(d*x+c)))/(a+b))^(1/2)*(a*(1+\sec(d*x+c)))/(a-b))^(1/2)/d$

3.413.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.67 (sec) , antiderivative size = 1241, normalized size of antiderivative = 2.27

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2),x]`


```
output ((4*a*(-16*a^2*A*b - 4*A*b^3 - 8*a^3*B - 11*a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 4*a*(8*a^3*A - 24*a*A*b^2 - 24*a^2*b*B - 4*b^3*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 2*(8*a^2*A*b - 4*A*b^3 - 9*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b)*Cos[c + d*x])*Sec[c + d*x])/(a + b)) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[...
```

3.413.3 Rubi [A] (verified)

Time = 2.88 (sec) , antiderivative size = 555, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3468, 27, 3042, 3528, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

↓ 3468

3.413. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

$$2 \int \frac{\sqrt{a+b \cos(c+dx)}(-b(4aA-bB) \cos^2(c+dx) - (Aa^2-2bBa-Ab^2) \cos(c+dx) + a(4Ab+aB))}{2\sqrt{\cos(c+dx)}} dx + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 27

$$\int \frac{\sqrt{a+b \cos(c+dx)}(-b(4aA-bB) \cos^2(c+dx) - (Aa^2-2bBa-Ab^2) \cos(c+dx) + a(4Ab+aB))}{\sqrt{\cos(c+dx)}} dx + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3042

$$\int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}(-b(4aA-bB) \sin(c+dx+\frac{\pi}{2})^2 + (-Aa^2+2bBa+Ab^2) \sin(c+dx+\frac{\pi}{2}) + a(4Ab+aB))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3528

$$\frac{1}{2} \int \frac{-b(8Aa^2-9bBa-4Ab^2) \cos^2(c+dx) - 2(2Aa^3-6bBa^2-6Ab^2a-b^3B) \cos(c+dx) + a(4Ba^2+12Aba-b^2B)}{2\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx + \frac{b(4aA-bB) \sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 27

$$\frac{1}{4} \int \frac{-b(8Aa^2-9bBa-4Ab^2) \cos^2(c+dx) - 2(2Aa^3-6bBa^2-6Ab^2a-b^3B) \cos(c+dx) + a(4Ba^2+12Aba-b^2B)}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx + \frac{b(4aA-bB) \sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3042

$$\frac{1}{4} \int \frac{-b(8Aa^2-9bBa-4Ab^2) \sin(c+dx+\frac{\pi}{2})^2 - 2(2Aa^3-6bBa^2-6Ab^2a-b^3B) \sin(c+dx+\frac{\pi}{2}) + a(4Ba^2+12Aba-b^2B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{b(4aA-bB) \sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3540

3.413. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

$$\frac{1}{4} \left(\frac{\int \frac{b^2(15Ba^2+20Aba+4b^2B) \cos^2(c+dx)+2ab(4Ba^2+12Aba+b^2B) \cos(c+dx)+ab(8Aa^2-9bBa-4Ab^2)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{2b} - \frac{(8a^2A - 9abB - 4Ab^2)}{d\sqrt{\cos(c+dx)}} \right) + \frac{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3042

$$\frac{1}{4} \left(\frac{\int \frac{b^2(15Ba^2+20Aba+4b^2B) \sin(c+dx+\frac{\pi}{2})^2+2ab(4Ba^2+12Aba+b^2B) \sin(c+dx+\frac{\pi}{2})+ab(8Aa^2-9bBa-4Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{(8a^2A - 9abB - 4Ab^2)}{d\sqrt{\sin(c+dx+\frac{\pi}{2})}} \right) + \frac{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3532

$$\frac{1}{4} \left(\frac{\int \frac{ab(8Aa^2-9bBa-4Ab^2)+2ab(4Ba^2+12Aba+b^2B) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx + b^2(15a^2B + 20aAb + 4b^2B) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx}{2b} - \frac{(8a^2A - 9abB - 4Ab^2)}{d\sqrt{\cos(c+dx)}} \right) + \frac{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3042

$$\frac{1}{4} \left(\frac{b^2(15a^2B + 20aAb + 4b^2B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{ab(8Aa^2-9bBa-4Ab^2)+2ab(4Ba^2+12Aba+b^2B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{(8a^2A - 9abB - 4Ab^2)}{d\sqrt{\sin(c+dx+\frac{\pi}{2})}} \right) + \frac{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

↓ 3288

$$\frac{1}{4} \left(\frac{\int \frac{ab(8Aa^2-9bBa-4Ab^2)+2ab(4Ba^2+12Aba+b^2B) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(15a^2B+20aAb+4b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a}{a+b}}}{2b}}{2b} - \frac{(8a^2A - 9abB - 4Ab^2)}{d\sqrt{\sin(c+dx+\frac{\pi}{2})}} \right) + \frac{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}}$$

3.413. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

↓ 3477

$$\frac{1}{4} \left(\frac{ab(8a^2A - 9abB - 4Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - ab(8a^2(A-B) - 3ab(8A+3B) - 2b^2(2A+B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)}} + \frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{-ab(8a^2(A-B) - 3ab(8A+3B) - 2b^2(2A+B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + ab(8a^2A - 9abB - 4Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)}} + \frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}} \right)$$

↓ 3295

$$\frac{1}{4} \left(\frac{ab(8a^2A - 9abB - 4Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(8a^2(A-B) - 3ab(8A+3B) - 2b^2(2A+B)) \cot(c+dx)}{d\sqrt{\cos(c+dx)}}}{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)}} + \frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}} \right)$$

↓ 3473

$$\frac{1}{4} \left(\frac{2b\sqrt{a+b}(8a^2(A-B) - 3ab(8A+3B) - 2b^2(2A+B)) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{d\sqrt{\cos(c+dx)}}}{b(4aA - bB) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)}} + \frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\sqrt{\cos(c+dx)}} \right)$$

input `Int[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(3/2), x]`

3.413. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$

```
output -1/2*(b*(4*a*A - b*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c +
d*x])/d + (2*a*A*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sqrt[Cos[c +
d*x]]) + (((2*(a - b)*b*Sqrt[a + b]*(8*a^2*A - 4*A*b^2 - 9*a*b*B)*Cot[c +
d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d
*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(
1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(8*a^2*(A - B) - 2*b
^2*(2*A + B) - 3*a*b*(8*A + 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b
*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt
[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d -
(2*b*Sqrt[a + b]*(20*a*A*b + 15*a^2*B + 4*b^2*B)*Cot[c + d*x]*EllipticPi[
(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]
)], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 +
Sec[c + d*x]))/(a - b)]/d)/(2*b) - ((8*a^2*A - 4*A*b^2 - 9*a*b*B)*Sqrt[a
+ b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/4
```

3.413.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)])/Sqrt[(c_) + (d_)*sin[(e_)] + (f_)
*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_)] + (f
_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)])), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.413.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4452 vs. $2(501) = 1002$.

Time = 17.71 (sec) , antiderivative size = 4453, normalized size of antiderivative = 8.14

method	result	size
default	Expression too large to display	4453
parts	Expression too large to display	4742

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

output `-1/4/d*(-8*A*cos(d*x+c)*sin(d*x+c)*a^2*b-8*A*a^3*sin(d*x+c)+8*A*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3+8*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3+9*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2-24*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*b+2*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2-24*A*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2+24*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)+40*A*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^2-48*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)...`

3.413.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algo rithm="fracas")`

output `Timed out`

3.413.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(3/2),x)`

output `Timed out`

3.413.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(3/2), x)`

3.413.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(3/2), x)`

3.413. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{3/2}(c+dx)} dx$

3.413.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(3/2),x)`

$$3.414 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$$

3.414.1 Optimal result	3852
3.414.2 Mathematica [C] (verified)	3853
3.414.3 Rubi [A] (verified)	3854
3.414.4 Maple [B] (warning: unable to verify)	3861
3.414.5 Fricas [F(-1)]	3862
3.414.6 Sympy [F(-1)]	3862
3.414.7 Maxima [F]	3862
3.414.8 Giac [F]	3863
3.414.9 Mupad [F(-1)]	3863

3.414.1 Optimal result

Integrand size = 35, antiderivative size = 536

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx = \frac{(a-b)\sqrt{a+b}(14aAb+6a^2B-3b^2B) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}}\right)\right) - \sqrt{a+b}(2ab(7A-9B)-2a^2(A-3B)-3b^2(6A+B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a+b}{a-b}\right) - b\sqrt{a+b}(2Ab+5aB) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{d} + \frac{2a(2Ab+aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{(14aAb+6a^2B-3b^2B)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3d\sqrt{\cos(c+dx)}} + \frac{2aA(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{3d \cos^{3/2}(c+dx)}$$

output $\frac{2}{3}aA(a+b\cos(dx+c))^{3/2}\sin(dx+c)/d\cos(dx+c)^{3/2}+2a(2Ab+Ba)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\cos(dx+c)^{1/2}-\frac{1}{3}(14Aab+6Ba^2-3Bb^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\cos(dx+c)^{1/2}+\frac{1}{3}(a-b)(14Aab+6Ba^2-3Bb^2)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b))^{1/2}/ad-\frac{1}{3}(2ab(7A-9B)-2a^2(A-3B)-3b^2(6A+B))\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b))^{1/2}/d-b(2Ab+5Ba)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b))^{1/2}/d$

3.414.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.69 (sec) , antiderivative size = 1269, normalized size of antiderivative = 2.37

$$\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\cos^{5/2}(c+dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2), x]`

```
output ((-4*a*(2*a^3*A + 4*a*A*b^2 + 12*a^2*b*B + 3*b^3*B)*Sqrt[((a + b)*Cot[(c +
d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]
*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[
ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/
(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c
+ d*x]]) - 4*a*(-14*a^2*A*b + 6*A*b^3 - 6*a^3*B + 18*a*b^2*B)*((Sqrt[((a
+ b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d
*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*
x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt
[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqr
t[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt
[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x]
)*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a +
b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c
+ d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-14*a*A
*b^2 - 6*a^2*b*B + 3*b^3*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*
EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]
*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c
+ d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]
^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt...
```

3.414.3 Rubi [A] (verified)

Time = 2.84 (sec) , antiderivative size = 547, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3468

3.414. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{5/2}(c+dx)} dx$

$$\frac{2}{3} \int \frac{\sqrt{a + b \cos(c + dx)}(-b(2aA - 3bB) \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(2Ab + aB))}{2 \cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{\sqrt{a + b \cos(c + dx)}(-b(2aA - 3bB) \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + 3a(2Ab + aB))}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}(-b(2aA - 3bB) \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + 3a(2Ab + aB))}{\sin^{\frac{3}{2}}(c + dx + \frac{\pi}{2})} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3526

$$\frac{1}{3} \left(2 \int \frac{-b(6Ba^2 + 14Aba - 3b^2B) \cos^2(c + dx) - (3Ba^3 + 7Aba^2 - 9b^2Ba - 3Ab^3) \cos(c + dx) + a(Aa^2 + 9bBa)}{2\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

↓ 27

$$\frac{1}{3} \left(\int \frac{-b(6Ba^2 + 14Aba - 3b^2B) \cos^2(c + dx) - (3Ba^3 + 7Aba^2 - 9b^2Ba - 3Ab^3) \cos(c + dx) + a(Aa^2 + 9bBa)}{\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{-b(6Ba^2 + 14Aba - 3b^2B) \sin^2(c + dx + \frac{\pi}{2}) + (-3Ba^3 - 7Aba^2 + 9b^2Ba + 3Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(Aa^2 + 9bBa)}{\sqrt{\sin(c + dx + \frac{\pi}{2})}\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3540

$$\frac{1}{3} \left(\frac{\int \frac{3(2Ab+5aB) \cos^2(c+dx)b^3 + a(6Ba^2+14Aba-3b^2B)b + 2a(Aa^2+9bBa+9Ab^2) \cos(c+dx)b}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{2b} - \frac{(6a^2B + 14aAb - 3b^2B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) - \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(\frac{\int \frac{3(2Ab+5aB) \sin(c+dx+\frac{\pi}{2})^2 b^3 + a(6Ba^2+14Aba-3b^2B)b + 2a(Aa^2+9bBa+9Ab^2) \sin(c+dx+\frac{\pi}{2})b}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{(6a^2B + 14aAb - 3b^2B)}{d\sqrt{\cos(c+dx)}} \right) - \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3532

$$\frac{1}{3} \left(\frac{\int \frac{ab(6Ba^2+14Aba-3b^2B) + 2ab(Aa^2+9bBa+9Ab^2) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx + 3b^3(5aB + 2Ab) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx}{2b} - \frac{(6a^2B + 14aAb)}{d\sqrt{\cos(c+dx)}} \right) - \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(\frac{\int \frac{ab(6Ba^2+14Aba-3b^2B) + 2ab(Aa^2+9bBa+9Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + 3b^3(5aB + 2Ab) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{(6a^2B + 14aAb)}{d\sqrt{\cos(c+dx)}} \right) - \frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

↓ 3288

3.414. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(\int \frac{ab(6Ba^2+14Aba-3b^2B)+2ab(Aa^2+9bBa+9Ab^2) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{6b^2\sqrt{a+b}(5aB+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d} \right)$$

2b

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\downarrow \text{3477}$$

$$\frac{1}{3} \left(ab(6a^2B+14aAb-3b^2B) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - ab(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B)) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \left(-ab(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + ab(6a^2B+14aAb) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\downarrow \text{3295}$$

$$\frac{1}{3} \left(ab(6a^2B+14aAb-3b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B)) \cot(c+dx)}{d} \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{3d \cos^{\frac{3}{2}}(c+dx)}$$

$$\downarrow \text{3473}$$

3.414. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(-\frac{2b\sqrt{a+b}(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B)) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right), -\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{3d \cos^{3/2}(c+dx)} \right)$$

input `Int[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(2*a*A*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + (((2*(a - b)*b*Sqrt[a + b]*(14*a*A*b + 6*a^2*B - 3*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(2*a*b*(7*A - 9*B) - 2*a^2*(A - 3*B) - 3*b^2*(6*A + B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (6*b^2*Sqrt[a + b]*(2*A*b + 5*a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(2*b) + (6*a*(2*A*b + a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - ((14*a*A*b + 6*a^2*B - 3*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/3`

3.414.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)]/(((b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.414.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4718 vs. $2(492) = 984$.

Time = 20.90 (sec) , antiderivative size = 4719, normalized size of antiderivative = 8.80

method	result	size
default	Expression too large to display	4719
parts	Expression too large to display	4721

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 1/d*(2/3*A*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*((csc(d*x+c)^2*a*(1-cos(d*x+c)
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1
)^(1/2)*(-csc(d*x+c)^2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/
2))*a^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d
*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x+c))^
2-7*csc(d*x+c)^2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2
*b*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c)
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x+c))^2-9*c
sc(d*x+c)^2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^2*(-
csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-c
sc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x+c))^2+3*csc(d*
x+c)^2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*
x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^3*(1-cos(d*x+c))^2+7*csc(d*x+c)^2*
EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b*(-csc(d*x+c)^2
*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*
b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x+c))^2+7*csc(d*x+c)^2*Ellip
ticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^2*(-csc(d*x+c)^2*(1-c
os(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-
cos(d*x+c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x+c))^2-6*csc(d*x+c)^2*(-csc(d...
```

3.414.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algo
rithm="fricas")
```

output Timed out

3.414.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)
```

output Timed out

3.414.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{5/2}} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algo
rithm="maxima")
```

```
output integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(5/2), x)
```

3.414.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(5/2), x)`

3.414.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(5/2),x)`

3.415
$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

3.415.1 Optimal result 3864
 3.415.2 Mathematica [C] (verified) 3865
 3.415.3 Rubi [A] (verified) 3865
 3.415.4 Maple [B] (warning: unable to verify) 3871
 3.415.5 Fricas [F] 3872
 3.415.6 Sympy [F(-1)] 3872
 3.415.7 Maxima [F] 3872
 3.415.8 Giac [F] 3873
 3.415.9 Mupad [F(-1)] 3873

3.415.1 Optimal result

Integrand size = 35, antiderivative size = 493

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(9a^2A+23Ab^2+35abB) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}\right)\right)}{\cos^{7/2}(c+dx)} + \frac{2\sqrt{a+b}(15Ab^3-ab^2(23A-45B)+a^2b(17A-35B)-a^3(9A-5B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}\right)\right)}{\cos^{7/2}(c+dx)} - \frac{2b^2\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{\cos^{7/2}(c+dx)} + \frac{2a(8Ab+5aB)\sqrt{a+b}\cos(c+dx) \sin(c+dx)}{15d \cos^{3/2}(c+dx)} + \frac{2aA(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{5d \cos^{5/2}(c+dx)}$$

output

```
2/5*a*A*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(5/2)+2/15*a*(8*A*b
+5*B*a)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2)+2/15*(a-b)*(9
*A*a^2+23*A*b^2+35*B*a*b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b
)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c
)))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d+2/15*(15*A*b^3-a*b^2*(2
3*A-45*B)+a^2*b*(17*A-35*B)-a^3*(9*A-5*B))*cot(d*x+c)*EllipticF((a+b*cos(d
*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2
)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d-2*b^2*
B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2
), (a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2
)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d
```

3.415.
$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

3.415.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.74 (sec) , antiderivative size = 1319, normalized size of antiderivative = 2.68

$$\int \frac{(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `((4*a*(-8*a^2*A*b + 8*A*b^3 - 5*a^3*B - 10*a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 4*a*(9*a^3*A + 23*a*A*b^2 + 35*a^2*b*B - 15*b^3*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 2*(9*a^2*A*b + 23*A*b^3 + 35*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a]]*Sqrt...`

3.415.3 Rubi [A] (verified)

Time = 2.24 (sec) , antiderivative size = 499, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.415. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

↓ 3468

$$\frac{2}{5} \int \frac{\sqrt{a + b \cos(c + dx)} (5b^2 B \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(8Ab + 5aB))}{2 \cos^{5/2}(c + dx)} dx +$$

$$\frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d \cos^{5/2}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{\sqrt{a + b \cos(c + dx)} (5b^2 B \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + a(8Ab + 5aB))}{\cos^{5/2}(c + dx)} dx +$$

$$\frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d \cos^{5/2}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (5b^2 B \sin^2(c + dx + \frac{\pi}{2}) + (3Aa^2 + 10bBa + 5Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(8Ab + 5aB))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx +$$

$$\frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d \cos^{5/2}(c + dx)}$$

↓ 3526

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{15B \cos^2(c + dx) b^3 + a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2Ba + 15Ab^3) \cos(c + dx)}{2 \cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - \right.$$

$$\left. \frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d \cos^{5/2}(c + dx)} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{15B \cos^2(c + dx) b^3 + a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2Ba + 15Ab^3) \cos(c + dx)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - \right.$$

$$\left. \frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{5d \cos^{5/2}(c + dx)} \right)$$

3.415. $\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{15B \sin(c + dx + \frac{\pi}{2})^2 b^3 + a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2Ba + 15Ab^3) \sin(c + dx)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3532

$$\frac{1}{5} \left(\frac{1}{3} \left(\int \frac{a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2Ba + 15Ab^3) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + 15b^3 B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\int \frac{a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2Ba + 15Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + 15b^3 B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3288

$$\frac{1}{5} \left(\frac{1}{3} \left(\int \frac{a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2Ba + 15Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{30b^2 B \sqrt{a + b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3477

$$\frac{1}{5} \left(\frac{1}{3} \left(a(9a^2 A + 35abB + 23Ab^2) \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + (-a^3(9A - 5B) + a^2b(17A - 35B)) \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

3.415. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{1}{3} \left(a(9a^2A + 35abB + 23Ab^2) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + (-(a^3(9A - 5B)) + a^2b(1) \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \right.$$

↓ 3295

$$\frac{1}{5} \left(\frac{1}{3} \left(a(9a^2A + 35abB + 23Ab^2) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(-(a^3(9A - 5B))}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \right.$$

↓ 3473

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(a-b)\sqrt{a+b}(9a^2A + 35abB + 23Ab^2) \cot(c + dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{ad} \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{5d \cos^{\frac{5}{2}}(c + dx)} \right) \right.$$

input `Int[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(7/2),x]`

output `(2*a*A*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*(a - b)*Sqrt[a + b]*(9*a^2*A + 23*A*b^2 + 35*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(15*A*b^3 - a*b^2*(23*A - 45*B) + a^2*b*(17*A - 35*B) - a^3*(9*A - 5*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (30*b^2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/3 + (2*a*(8*A*b + 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2))/5`

3.415.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(b \sin(e + fx) + f x)^{3/2}} dx$$

$$\rightarrow \text{Simp}[-2A(c - d) \frac{\tan(e + fx)}{f b c^2} \text{Rt}[(c + d)/b, 2] \sqrt{c(1 + \csc(e + fx))} / (c - d)] \sqrt{c(1 - \csc(e + fx))} / (c + d)] \text{EllipticE}[\text{ArcSin}[\sqrt{c + d \sin(e + fx)}] / \sqrt{b \sin(e + fx)}] / \text{Rt}[(c + d)/b, 2], -(c + d)/(c - d), x] /;$$` `FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[(A - B)/(a - b) \int [1/(\sqrt{a + b \sin(e + fx)} \sqrt{c + d \sin(e + fx)})], x], x] - \text{Simp}[(A*b - a*B)/(a - b) \int [(1 + \sin(e + fx))/(a + b \sin(e + fx))^{3/2} \sqrt{c + d \sin(e + fx)}], x], x] /;$$` `FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\rightarrow \text{Simp}[-(c^2 C - B*c*d + A*d^2) \cos(e + fx) (a + b \sin(e + fx))^m (c + d \sin(e + fx))^{n+1} / (d*f*(n+1)*(c^2 - d^2))] + \text{Simp}[1/(d*(n+1)*(c^2 - d^2)) \int [(a + b \sin(e + fx))^{m-1} (c + d \sin(e + fx))^{n+1} \text{Simp}[A*d*(b*d*m + a*c*(n+1)) + (c*C - B*d)*(b*c*m + a*d*(n+1)) - (d*(A*(a*d*(n+2) - b*c*(n+1)) + B*(b*d*(n+1) - a*c*(n+2))) - C*(b*c*d*(n+1) - a*(c^2 + d^2*(n+1)))] \sin(e + fx) + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m+1) + d^2*(n+1))] \sin^2(e + fx)^2, x], x], x] /;$$` `FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2)/

$$\int \frac{(A + B \sin(e + fx) + C \sin^2(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[C/b^2 \int [\sqrt{a + b \sin(e + fx)}] / \sqrt{c + d \sin(e + fx)}, x], x] + \text{Simp}[1/b^2 \int [(A*b^2 - a^2*C + b*(b*B - 2*a*C) \sin(e + fx)) / ((a + b \sin(e + fx))^{3/2} \sqrt{c + d \sin(e + fx)})], x], x] /;$$` `FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.415.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4432 vs. $2(453) = 906$.

Time = 22.24 (sec) , antiderivative size = 4433, normalized size of antiderivative = 8.99

method	result	size
default	Expression too large to display	4433
parts	Expression too large to display	4586

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x,method=_RET
URNVERBOSE)
```

```
output 2/15/d*(14*A*cos(d*x+c)*sin(d*x+c)*a^2*b+35*B*cos(d*x+c)^3*sin(d*x+c)*a*b^
2+34*A*cos(d*x+c)^2*sin(d*x+c)*a*b^2+40*B*cos(d*x+c)^2*sin(d*x+c)*a^2*b+3*
A*a^3*sin(d*x+c)+23*A*cos(d*x+c)^3*sin(d*x+c)*b^3+5*B*cos(d*x+c)*sin(d*x+c
)*a^3-15*A*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+co
s(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b^3
*cos(d*x+c)^2+9*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+c
os(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/
2))*a^2*b*cos(d*x+c)^4+23*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+
c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/
(a+b))^(1/2))*a*b^2*cos(d*x+c)^4-17*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((
a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c
),(-(a-b)/(a+b))^(1/2))*a^2*b*cos(d*x+c)^4-23*A*(cos(d*x+c)/(1+cos(d*x+c))
)^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)
-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2*cos(d*x+c)^4+35*B*(cos(d*x+c)/(1+c
os(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(
cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*b*cos(d*x+c)^4+35*B*(cos(d
*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*
EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2*cos(d*x+c)^4-3
5*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+
b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*b*c...
```

3.415.5 Fricas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{7/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(7/2), x)`

3.415.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(7/2),x)`

output `Timed out`

3.415.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{7/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(7/2), x)`

3.415. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{7/2}(c+dx)} dx$

3.415.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{7/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(7/2), x)`

3.415.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(7/2),x)`

3.416
$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.416.1 Optimal result	3874
3.416.2 Mathematica [C] (verified)	3875
3.416.3 Rubi [A] (verified)	3875
3.416.4 Maple [B] (verified)	3881
3.416.5 Fricas [F]	3882
3.416.6 Sympy [F(-1)]	3882
3.416.7 Maxima [F]	3882
3.416.8 Giac [F]	3883
3.416.9 Mupad [F(-1)]	3883

3.416.1 Optimal result

Integrand size = 35, antiderivative size = 434

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(145a^2Ab+15Ab^3+63a^3B+161ab^2B)}{105ad} + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)+15b^2(A-7B)-8ab(15A-7B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{c+dx}}\right)\right)}{105ad} + \frac{2a(10Ab+7aB)\sqrt{a+b\cos(c+dx)}\sin(c+dx)}{35d\cos^{\frac{5}{2}}(c+dx)} + \frac{2(25a^2A+45Ab^2+77abB)\sqrt{a+b\cos(c+dx)}\sin(c+dx)}{105d\cos^{\frac{3}{2}}(c+dx)} + \frac{2aA(a+b\cos(c+dx))^{3/2}\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)}$$

```
output 2/7*a*A*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(7/2)+2/35*a*(10*A*
b+7*B*a)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+2/105*(25*A*
a^2+45*A*b^2+77*B*a*b)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(3/2
)+2/105*(a-b)*(145*A*a^2*b+15*A*b^3+63*B*a^3+161*B*a*b^2)*cot(d*x+c)*Ellip
ticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1
/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(
1/2)/a^2/d+2/105*(a-b)*(a^2*(25*A-63*B)+15*b^2*(A-7*B)-8*a*b*(15*A-7*B))*c
ot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((
-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d
*x+c))/(a-b))^(1/2)/a/d
```

3.416.
$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

3.416.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.87 (sec) , antiderivative size = 1409, normalized size of antiderivative = 3.25

$$\int \frac{(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(9/2),x]`

output `((-4*a*(25*a^4*A - 10*a^2*A*b^2 - 15*A*b^4 + 56*a^3*b*B - 56*a*b^3*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-145*a^3*A*b - 15*a*A*b^3 - 63*a^4*B - 161*a^2*b^2*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-145*a^2*A*b^2 - 15*A*b^4 - 63*a^3*b*B - 161*a*b^3*B)*(I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + ...`

3.416.3 Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.416. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{9/2}(c+dx)} dx$

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx$$

↓ 3468

$$\frac{2}{7} \int \frac{\sqrt{a + b \cos(c + dx)} (b(2aA + 7bB) \cos^2(c + dx) + (5Aa^2 + 14bBa + 7Ab^2) \cos(c + dx) + a(10Ab + 7aB))}{2 \cos^{7/2}(c + dx)} dx$$

$$\frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{7d \cos^{7/2}(c + dx)}$$

↓ 27

$$\frac{1}{7} \int \frac{\sqrt{a + b \cos(c + dx)} (b(2aA + 7bB) \cos^2(c + dx) + (5Aa^2 + 14bBa + 7Ab^2) \cos(c + dx) + a(10Ab + 7aB))}{\cos^{7/2}(c + dx)} dx$$

$$\frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{7d \cos^{7/2}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (b(2aA + 7bB) \sin^2(c + dx + \frac{\pi}{2}) + (5Aa^2 + 14bBa + 7Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(10Ab + 7aB))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx$$

$$\frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{7d \cos^{7/2}(c + dx)}$$

↓ 3526

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{b(14Ba^2 + 30Aba + 35b^2B) \cos^2(c + dx) + (21Ba^3 + 65Aba^2 + 105b^2Ba + 35Ab^3) \cos(c + dx) + a(25Aa^2 + 15Ab^2)}{2 \cos^{5/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \right.$$

$$\left. \frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{7d \cos^{7/2}(c + dx)} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{b(14Ba^2 + 30Aba + 35b^2B) \cos^2(c + dx) + (21Ba^3 + 65Aba^2 + 105b^2Ba + 35Ab^3) \cos(c + dx) + a(25Aa^2 + 15Ab^2)}{\cos^{5/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \right.$$

$$\left. \frac{2aA \sin(c + dx) (a + b \cos(c + dx))^{3/2}}{7d \cos^{7/2}(c + dx)} \right)$$

3.416. $\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{b(14Ba^2 + 30Aba + 35b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (21Ba^3 + 65Aba^2 + 105b^2Ba + 35Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{5/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3534

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2 \int \frac{a(63Ba^3 + 145Aba^2 + 161b^2Ba + 15Ab^3) + a(25Aa^3 + 119bBa^2 + 135Ab^2a + 105b^3B) \cos(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{3a} + \frac{2(25a^2A + 77abB + 45Ab^2)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{\int \frac{a(63Ba^3 + 145Aba^2 + 161b^2Ba + 15Ab^3) + a(25Aa^3 + 119bBa^2 + 135Ab^2a + 105b^3B) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{3a} + \frac{2(25a^2A + 77abB + 45Ab^2)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{\int \frac{a(63Ba^3 + 145Aba^2 + 161b^2Ba + 15Ab^3) + a(25Aa^3 + 119bBa^2 + 135Ab^2a + 105b^3B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{3a} + \frac{2(25a^2A + 77abB + 45Ab^2)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3477

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{a(a - b)(a^2(25A - 63B) - 8ab(15A - 7B) + 15b^2(A - 7B)) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx + a(63a^3B + 145Aab^2)}{3a} \right) \right. \\ \left. \frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{7d \cos^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3042

3.416. $\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{a(a-b)(a^2(25A-63B) - 8ab(15A-7B) + 15b^2(A-7B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + a(63a^3B + 145a^2Ab + 161ab^2B + 15Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)-8ab(15A-7B)+15b^2(A-7B))}{3a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{7d \cos^{7/2}(c+dx)}$$

↓ 3295

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{a(63a^3B + 145a^2Ab + 161ab^2B + 15Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)-8ab(15A-7B)+15b^2(A-7B))}{3a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{7d \cos^{7/2}(c+dx)}$$

↓ 3473

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2(25a^2A + 77abB + 45Ab^2) \sin(c+dx) \sqrt{a+b\cos(c+dx)}}{3d \cos^{3/2}(c+dx)} + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)-8ab(15A-7B)+15b^2(A-7B))}{3a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{7d \cos^{7/2}(c+dx)}$$

```
input Int[((a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]))/cos[c + d*x]^(9/2),x
]
```

```
output (2*a*A*(a + b*cos[c + d*x])^(3/2)*sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)) +
((2*a*(10*A*b + 7*a*B)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/(5*d*cos[c
+ d*x]^(5/2)) + (((2*(a - b)*sqrt[a + b]*(145*a^2*A*b + 15*A*b^3 + 63*a^3*
B + 161*a*b^2*B)*cot[c + d*x]*ellipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]]/(S
qrt[a + b]*sqrt[cos[c + d*x]]]), -((a + b)/(a - b)))*sqrt[(a*(1 - Sec[c +
d*x]))/(a + b)]*sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*(a - b)*S
qrt[a + b]*(a^2*(25*A - 63*B) + 15*b^2*(A - 7*B) - 8*a*b*(15*A - 7*B))*cot
[c + d*x]*ellipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]]/(sqrt[a + b]*sqrt[cos[
c + d*x]])], -((a + b)/(a - b)))*sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*sqrt
[(a*(1 + Sec[c + d*x]))/(a - b)]/d)/(3*a) + (2*(25*a^2*A + 45*A*b^2 + 77*
a*b*B)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)))/5
/7
```

3.416. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{9/2}(c+dx)} dx$

3.416.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

3.416.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4873 vs. $2(396) = 792$.

Time = 26.96 (sec) , antiderivative size = 4874, normalized size of antiderivative = 11.23

method	result	size
parts	Expression too large to display	4874
default	Expression too large to display	4939

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x,method=_RET
URNVERBOSE)
```

```
output 2/21*A/d*(3*a^4*sin(d*x+c)+18*a^2*b^2*cos(d*x+c)^2*sin(d*x+c)+3*b^4*cos(d*
x+c)^4*sin(d*x+c)+5*a^4*cos(d*x+c)^2*sin(d*x+c)+29*a^2*b^2*cos(d*x+c)^4*si
n(d*x+c)+3*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d
*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4
*cos(d*x+c)^5-5*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+
cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2
)*a^4*cos(d*x+c)^5+6*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))
*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))
^(1/2)*b^4*cos(d*x+c)^4-10*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(
1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*
x+c)))^(1/2)*a^4*cos(d*x+c)^4+3*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a
+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+c
os(d*x+c)))^(1/2)*b^4*cos(d*x+c)^3-5*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-
b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)
/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^3+6*EllipticE(cot(d*x+c)-csc(d*x+c),
(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d
*x+c)/(1+cos(d*x+c)))^(1/2)*a*b^3*cos(d*x+c)^4-58*EllipticF(cot(d*x+c)-csc
(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2
)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3*b*cos(d*x+c)^4-54*EllipticF(cot(d*
x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/. . .
```


3.416.5 Fricas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^9(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos^9(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(9/2), x)`

3.416.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^9(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(9/2),x)`

output `Timed out`

3.416.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^9(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos^9(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(9/2), x)`

3.416. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^9(c+dx)} dx$

3.416.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{9/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(9/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(9/2), x)`

3.416.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{9/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(9/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(9/2), x)`

$$3.417 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

3.417.1 Optimal result	3884
3.417.2 Mathematica [C] (verified)	3885
3.417.3 Rubi [A] (verified)	3886
3.417.4 Maple [B] (verified)	3893
3.417.5 Fracas [F]	3893
3.417.6 Sympy [F(-1)]	3893
3.417.7 Maxima [F]	3894
3.417.8 Giac [F]	3894
3.417.9 Mupad [F(-1)]	3894

3.417.1 Optimal result

Integrand size = 35, antiderivative size = 522

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(147a^4A+279a^2Ab^2-10Ab^4+435a^3bB)}{315a^2d}$$

$$- \frac{2(a-b)\sqrt{a+b}(10Ab^3-6a^2b(19A-60B)+3a^3(49A-25B)+15ab^2(11A-3B)) \cot(c+dx) \text{EllipticF}}{315a^2d}$$

$$+ \frac{2a(4Ab+3aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{21d \cos^{\frac{7}{2}}(c+dx)}$$

$$+ \frac{2(49a^2A+75Ab^2+135abB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{315d \cos^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{2(163a^2Ab+5Ab^3+75a^3B+135ab^2B)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{315ad \cos^{\frac{3}{2}}(c+dx)}$$

$$+ \frac{2aA(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)}$$

output $2/9*a*A*(a+b*\cos(d*x+c))^(3/2)*\sin(d*x+c)/d/\cos(d*x+c)^(9/2)+2/21*a*(4*A*b+3*B*a)*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(7/2)+2/315*(49*A*a^2+75*A*b^2+135*B*a*b)*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(5/2)+2/315*(163*A*a^2*b+5*A*b^3+75*B*a^3+135*B*a*b^2)*\sin(d*x+c)*(a+b*\cos(d*x+c))^(1/2)/a/d/\cos(d*x+c)^(3/2)+2/315*(a-b)*(147*A*a^4+279*A*a^2*b^2-10*A*b^4+435*B*a^3*b+45*B*a*b^3)*\cot(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^(1/2)/(a+b)^(1/2)/\cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-\sec(d*x+c)))/(a+b))^(1/2)*(a*(1+\sec(d*x+c)))/(a-b))^(1/2)/a^3/d-2/315*(a-b)*(10*A*b^3-6*a^2*b*(19*A-60*B)+3*a^3*(49*A-25*B)+15*a*b^2*(11*A-3*B))*\cot(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^(1/2)/(a+b)^(1/2)/\cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-\sec(d*x+c)))/(a+b))^(1/2)*(a*(1+\sec(d*x+c)))/(a-b))^(1/2)/a^2/d$

3.417.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.99 (sec) , antiderivative size = 1517, normalized size of antiderivative = 2.91

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(11/2),x]`

output

```

-1/315*((-4*a*(-114*a^4*A*b + 124*a^2*A*b^3 - 10*A*b^5 - 75*a^5*B + 30*a^3
*b^2*B + 45*a*b^4*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((
a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc
[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x
])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((
(a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(147*a^5*A + 27
9*a^3*A*b^2 - 10*a*A*b^4 + 435*a^4*b*B + 45*a^2*b^3*B)*((Sqrt[((a + b)*Cot
[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2
)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*Ellip
ticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-
2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*
Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a +
b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c
+ d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c
+ d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2
]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])) + 2*(147*a^4*A*b + 2
79*a^2*A*b^3 - 10*A*b^5 + 435*a^3*b^2*B + 45*a*b^4*B)*((I*Cos[(c + d*x)/2]
*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c
+ d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c
+ d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*S...

```

3.417.3 Rubi [A] (verified)

Time = 2.66 (sec) , antiderivative size = 533, normalized size of antiderivative = 1.02, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx \\
 & \quad \downarrow \text{3468}
 \end{aligned}$$

3.417. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

$$\frac{2}{9} \int \frac{\sqrt{a + b \cos(c + dx)} (b(4aA + 9bB) \cos^2(c + dx) + (7Aa^2 + 18bBa + 9Ab^2) \cos(c + dx) + 3a(4Ab + 3aB))}{2 \cos^{\frac{9}{2}}(c + dx)} dx$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{9} \int \frac{\sqrt{a + b \cos(c + dx)} (b(4aA + 9bB) \cos^2(c + dx) + (7Aa^2 + 18bBa + 9Ab^2) \cos(c + dx) + 3a(4Ab + 3aB))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{9} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (b(4aA + 9bB) \sin(c + dx + \frac{\pi}{2})^2 + (7Aa^2 + 18bBa + 9Ab^2) \sin(c + dx + \frac{\pi}{2}) + 3a(4Ab + 3aB))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 3526

$$\frac{1}{9} \left(\frac{2}{7} \int \frac{b(36Ba^2 + 76Aba + 63b^2B) \cos^2(c + dx) + (45Ba^3 + 137Aba^2 + 189b^2Ba + 63Ab^3) \cos(c + dx) + a(49A^2 + 42AbB + 9b^2B^2)}{2 \cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{1}{7} \int \frac{b(36Ba^2 + 76Aba + 63b^2B) \cos^2(c + dx) + (45Ba^3 + 137Aba^2 + 189b^2Ba + 63Ab^3) \cos(c + dx) + a(49A^2 + 42AbB + 9b^2B^2)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \int \frac{b(36Ba^2 + 76Aba + 63b^2B) \sin(c + dx + \frac{\pi}{2})^2 + (45Ba^3 + 137Aba^2 + 189b^2Ba + 63Ab^3) \sin(c + dx + \frac{\pi}{2}) + a(49A^2 + 42AbB + 9b^2B^2)}{\sin(c + dx + \frac{\pi}{2})^{7/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right)$$

$$\frac{2aA \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c + dx)}$$

3.417. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

↓ 3534

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{2 \int \frac{2ab(49Aa^2+135bBa+75Ab^2) \cos^2(c+dx)+a(147Aa^3+585bBa^2+605Ab^2a+315b^3B) \cos(c+dx)+3a(75Ba^3+163Aba^2+135b^2Ba+585b^3B)}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{\int \frac{2ab(49Aa^2+135bBa+75Ab^2) \cos^2(c+dx)+a(147Aa^3+585bBa^2+605Ab^2a+315b^3B) \cos(c+dx)+3a(75Ba^3+163Aba^2+135b^2Ba+585b^3B)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{\int \frac{2ab(49Aa^2+135bBa+75Ab^2) \sin(c+dx+\frac{\pi}{2})^2+a(147Aa^3+585bBa^2+605Ab^2a+315b^3B) \sin(c+dx+\frac{\pi}{2})+3a(75Ba^3+163Aba^2+135b^2Ba+585b^3B)}{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}}{5a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3534

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{2 \int \frac{3((75Ba^3+261Aba^2+405b^2Ba+155Ab^3) \cos(c+dx)a^2+(147Aa^4+435bBa^3+279Ab^2a^2+45b^3Ba-10Ab^4)a)}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(75a^3B+163a^2Ab+135ab^2)}{d} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 27

3.417. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{(75Ba^3+261Aba^2+405b^2Ba+155Ab^3) \cos(c+dx)a^2 + (147Aa^4+435bBa^3+279Ab^2a^2+45b^3Ba-10Ab^4)a}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2(75a^3B+163a^2Ab+135ab^2B+135a^3b^2)}{d\cos^{\frac{3}{2}}(c+dx)} \right) \right)$$

5a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{(75Ba^3+261Aba^2+405b^2Ba+155Ab^3) \sin(c+dx+\frac{\pi}{2})a^2 + (147Aa^4+435bBa^3+279Ab^2a^2+45b^3Ba-10Ab^4)a}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(75a^3B+163a^2Ab+135ab^2B+135a^3b^2)}{d\cos^{\frac{3}{2}}(c+dx)} \right) \right)$$

5a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3477

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{a(147a^4A+435a^3bB+279a^2Ab^2+45ab^3B-10Ab^4) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - a(a-b)(3a^3(49A-25B)-6a^2b(19A-60B)+15ab^2(19A-60B))}{a} \right) \right)$$

5a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{a(147a^4A+435a^3bB+279a^2Ab^2+45ab^3B-10Ab^4) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - a(a-b)(3a^3(49A-25B)-6a^2b(19A-60B))}{a} \right) \right)$$

5

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{9d \cos^{\frac{9}{2}}(c+dx)}$$

↓ 3295

3.417. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{a(147a^4A + 435a^3bB + 279a^2Ab^2 + 45ab^3B - 10Ab^4) \int \frac{\sin(c+dx + \frac{\pi}{2}) + 1}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(3a^3(49A-25B) - 6a^2b(19A-60B))}{a}}{a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{9d \cos^{9/2}(c+dx)}$$

↓ 3473

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{2(49a^2A + 135abB + 75Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{5/2}(c+dx)} + \frac{2(75a^3B + 163a^2Ab + 135ab^2B + 5Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \cos^{3/2}(c+dx)} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{9d \cos^{9/2}(c+dx)}$$

```
input Int[((a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]))/cos[c + d*x]^(11/2), x]
```

```
output (2*a*A*(a + b*cos[c + d*x])^(3/2)*sin[c + d*x]/(9*d*cos[c + d*x]^(9/2)) +
((6*a*(4*A*b + 3*a*B)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/(7*d*cos[c +
d*x]^(7/2)) + ((2*(49*a^2*A + 75*A*b^2 + 135*a*b*B)*sqrt[a + b*cos[c + d*
x]]*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)) + (((2*(a - b)*sqrt[a + b]*(147
*a^4*A + 279*a^2*A*b^2 - 10*A*b^4 + 435*a^3*b*B + 45*a*b^3*B)*cot[c + d*x]
*ellipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(sqrt[a + b]*sqrt[cos[c + d*x]]
)], -(a + b)/(a - b))*sqrt[(a*(1 - sec[c + d*x]))/(a + b)]*sqrt[(a*(1 +
sec[c + d*x]))/(a - b))]/(a*d) - (2*(a - b)*sqrt[a + b]*(10*A*b^3 - 6*a^2*
b*(19*A - 60*B) + 3*a^3*(49*A - 25*B) + 15*a*b^2*(11*A - 3*B))*cot[c + d*x]
*ellipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(sqrt[a + b]*sqrt[cos[c + d*x]
]]], -(a + b)/(a - b))*sqrt[(a*(1 - sec[c + d*x]))/(a + b)]*sqrt[(a*(1 +
sec[c + d*x]))/(a - b))]/d)/a + (2*(163*a^2*A*b + 5*A*b^3 + 75*a^3*B + 13
5*a*b^2*B)*sqrt[a + b*cos[c + d*x]]*sin[c + d*x])/(d*cos[c + d*x]^(3/2)))/
(5*a))/7)/9
```

3.417. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

3.417.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

3.417.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5954 vs. 2(478) = 956.

Time = 30.97 (sec) , antiderivative size = 5955, normalized size of antiderivative = 11.41

method	result	size
parts	Expression too large to display	5955
default	Expression too large to display	6034

input `int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x,method=_RE
TURNVERBOSE)`

output `result too large to display`

3.417.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{11/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, alg
orithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2
+ (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(1
1/2), x)`

3.417.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(11/2),x)`

output `Timed out`

3.417. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

3.417.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{11/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(11/2), x)`

3.417.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{11/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(11/2), x)`

3.417.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{11/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{11/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(11/2), x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(11/2), x)`

3.417. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{11/2}(c+dx)} dx$

$$3.418 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$$

3.418.1 Optimal result	3895
3.418.2 Mathematica [C] (warning: unable to verify)	3896
3.418.3 Rubi [A] (verified)	3897
3.418.4 Maple [B] (warning: unable to verify)	3905
3.418.5 Fricas [F]	3905
3.418.6 Sympy [F(-1)]	3906
3.418.7 Maxima [F]	3906
3.418.8 Giac [F]	3906
3.418.9 Mupad [F(-1)]	3907

3.418.1 Optimal result

Integrand size = 35, antiderivative size = 622

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(3705a^4Ab+255a^2Ab^3+40Ab^5+1617a^5b)}{3465a^3d} + \frac{2(a-b)\sqrt{a+b}(40Ab^4+3a^4(225A-539B)-6a^3b(505A-209B)+15a^2b^2(19A-121B)+10ab^3(3A-5B))}{3465a^3d} + \frac{2a(14Ab+11aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{99d \cos^{\frac{9}{2}}(c+dx)} + \frac{2(81a^2A+113Ab^2+209abB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{693d \cos^{\frac{7}{2}}(c+dx)} + \frac{2(1145a^2Ab+15Ab^3+539a^3B+825ab^2B)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3465ad \cos^{\frac{5}{2}}(c+dx)} + \frac{2(675a^4A+1025a^2Ab^2-20Ab^4+1793a^3bB+55ab^3B)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3465a^2d \cos^{\frac{3}{2}}(c+dx)} + \frac{2aA(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{11d \cos^{\frac{11}{2}}(c+dx)}$$

3.418. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$

output
$$\begin{aligned} & \frac{2}{11} a A (a+b \cos(dx+c))^{3/2} \sin(dx+c) / d \cos(dx+c)^{11/2} + \frac{2}{99} a (14 A b + 11 B a) \sin(dx+c) (a+b \cos(dx+c))^{1/2} / d \cos(dx+c)^{9/2} + \frac{2}{693} (81 A^2 a^2 + 113 A b^2 + 209 B a b) \sin(dx+c) (a+b \cos(dx+c))^{1/2} / d \cos(dx+c)^{7/2} \\ & + \frac{2}{3465} (1145 A^2 a^2 b + 15 A b^3 + 539 B a^3 + 825 B a b^2) \sin(dx+c) (a+b \cos(dx+c))^{1/2} / a d \cos(dx+c)^{5/2} + \frac{2}{3465} (675 A^4 a^4 + 1025 A^2 a^2 b^2 - 20 A b^4 + 1793 B a^3 b + 55 B a b^3) \sin(dx+c) (a+b \cos(dx+c))^{1/2} / a^2 d \cos(dx+c)^{3/2} \\ & + \frac{2}{3465} (a-b) (3705 A^4 a^4 b + 255 A^2 a^2 b^3 + 40 A b^5 + 1617 B a^5 + 3069 B a^3 b^2 - 110 B a b^4) \cot(dx+c) \operatorname{EllipticE}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2}) (a+b)^{1/2} (a(1-\sec(dx+c)) / (a+b))^{1/2} (a(1+\sec(dx+c)) / (a-b))^{1/2} / a^4 d \\ & + \frac{2}{3465} (a-b) (40 A b^4 + 3 a^4 (225 A - 539 B) - 6 a^3 b (505 A - 209 B) + 15 a^2 b^2 (19 A - 121 B) + 10 a b^3 (3 A - 11 B)) \cot(dx+c) \operatorname{EllipticF}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2}) (a+b)^{1/2} (a(1-\sec(dx+c)) / (a+b))^{1/2} (a(1+\sec(dx+c)) / (a-b))^{1/2} / a^3 d \end{aligned}$$

3.418.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.13 (sec) , antiderivative size = 1640, normalized size of antiderivative = 2.64

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(13/2),x]`

```
output ((-4*a*(675*a^6*A - 390*a^4*A*b^2 - 245*a^2*A*b^4 - 40*A*b^6 + 1254*a^5*b*B - 1364*a^3*b^3*B + 110*a*b^5*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) - 4*a*(-3705*a^5*A*b - 255*a^3*A*b^3 - 40*a*A*b^5 - 1617*a^6*B - 3069*a^4*b^2*B + 110*a^2*b^4*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]) + 2*(-3705*a^4*A*b^2 - 255*a^2*A*b^4 - 40*A*b^6 - 1617*a^5*b*B - 3069*a^3*b^3*B + 110*a*b^5*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*cos[c + d*x]])*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[(a...
```

3.418.3 Rubi [A] (verified)

Time = 3.39 (sec) , antiderivative size = 638, normalized size of antiderivative = 1.03, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{13/2}} dx$$

↓ 3468

3.418. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{13/2}(c+dx)} dx$

$$\frac{2}{11} \int \frac{\sqrt{a+b \cos(c+dx)}(b(6aA+11bB) \cos^2(c+dx) + (9Aa^2+22bBa+11Ab^2) \cos(c+dx) + a(14Ab+11aB))}{2 \cos^{\frac{11}{2}}(c+dx)} dx$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{11} \int \frac{\sqrt{a+b \cos(c+dx)}(b(6aA+11bB) \cos^2(c+dx) + (9Aa^2+22bBa+11Ab^2) \cos(c+dx) + a(14Ab+11aB))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}(b(6aA+11bB) \sin(c+dx+\frac{\pi}{2})^2 + (9Aa^2+22bBa+11Ab^2) \sin(c+dx+\frac{\pi}{2}) + a(14Ab+11aB))}{\sin(c+dx+\frac{\pi}{2})^{11/2}} dx$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3526

$$\frac{1}{11} \left(\frac{2}{9} \int \frac{3b(22Ba^2+46Aba+33b^2B) \cos^2(c+dx) + (77Ba^3+233Aba^2+297b^2Ba+99Ab^3) \cos(c+dx) + a(8b^3+12b^2A+6bA^2+8A^2b)}{2 \cos^{\frac{9}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \int \frac{3b(22Ba^2+46Aba+33b^2B) \cos^2(c+dx) + (77Ba^3+233Aba^2+297b^2Ba+99Ab^3) \cos(c+dx) + a(8b^3+12b^2A+6bA^2+8A^2b)}{\cos^{\frac{9}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \int \frac{3b(22Ba^2+46Aba+33b^2B) \sin(c+dx+\frac{\pi}{2})^2 + (77Ba^3+233Aba^2+297b^2Ba+99Ab^3) \sin(c+dx+\frac{\pi}{2}) + a(8b^3+12b^2A+6bA^2+8A^2b)}{\sin(c+dx+\frac{\pi}{2})^{9/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3534

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2 \int \frac{4ab(81Aa^2+209bBa+113Ab^2) \cos^2(c+dx)+a(405Aa^3+1507bBa^2+1531Ab^2a+693b^3B) \cos(c+dx)+a(539Ba^3+1145Aba^2+825b^2B)}{2 \cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}}{7a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{\int \frac{4ab(81Aa^2+209bBa+113Ab^2) \cos^2(c+dx)+a(405Aa^3+1507bBa^2+1531Ab^2a+693b^3B) \cos(c+dx)+a(539Ba^3+1145Aba^2+825b^2B)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}}{7a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{\int \frac{4ab(81Aa^2+209bBa+113Ab^2) \sin(c+dx+\frac{\pi}{2})^2+a(405Aa^3+1507bBa^2+1531Ab^2a+693b^3B) \sin(c+dx+\frac{\pi}{2})+a(539Ba^3+1145Aba^2+825b^2B)}{\sin(c+dx+\frac{\pi}{2})^{7/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}}{7a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3534

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2 \int \frac{(1617Ba^3+5055Aba^2+6655b^2Ba+2305Ab^3) \cos(c+dx)a^2+2b(539Ba^3+1145Aba^2+825b^2Ba+15Ab^3) \cos^2(c+dx)a+3(675Aa^4+1793bBa^3+1025b^2Ba^2+113Ab^3)}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 27

3.418. $\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(\int \frac{(1617Ba^3 + 5055Aba^2 + 6655b^2Ba + 2305Ab^3) \cos(c+dx)a^2 + 2b(539Ba^3 + 1145Aba^2 + 825b^2Ba + 15Ab^3) \cos^2(c+dx)a + 3(675Aa^4 + 1793bBa^3 + 1025Ab^2a^2 + 110b^3Ba + 10Ab^4)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

7a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\int \frac{(1617Ba^3 + 5055Aba^2 + 6655b^2Ba + 2305Ab^3) \sin(c+dx+\frac{\pi}{2})a^2 + 2b(539Ba^3 + 1145Aba^2 + 825b^2Ba + 15Ab^3) \sin(c+dx+\frac{\pi}{2})a + 3(675Aa^4 + 1793bBa^3 + 1025Ab^2a^2 + 110b^3Ba + 10Ab^4)}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \right)$$

7a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3534

$$\frac{1}{11} \left(\frac{1}{9} \left(2 \int \frac{3((675Aa^4 + 2871bBa^3 + 3315Ab^2a^2 + 1705b^3Ba + 10Ab^4) \cos(c+dx)a^2 + (1617Ba^5 + 3705Aba^4 + 3069b^2Ba^3 + 255Ab^3a^2 - 110b^4Ba + 40Ab^5)a) dx}{2 \cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \right) \right)$$

5a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \left(\int \frac{(675Aa^4 + 2871bBa^3 + 3315Ab^2a^2 + 1705b^3Ba + 10Ab^4) \cos(c+dx)a^2 + (1617Ba^5 + 3705Aba^4 + 3069b^2Ba^3 + 255Ab^3a^2 - 110b^4Ba + 40Ab^5)a}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

5a

$$\frac{2aA \sin(c+dx)(a+b\cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

3.418. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \int \frac{(675Aa^4 + 2871bBa^3 + 3315Ab^2a^2 + 1705b^3Ba + 10Ab^4) \sin(c+dx + \frac{\pi}{2}) a^2 + (1617Ba^5 + 3705Aba^4 + 3069b^2Ba^3 + 255Ab^3a^2 - 110b^4Ba + 40Ab^5) a}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \dots \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{11/2}(c+dx)}$$

↓ 3477

$$\frac{1}{11} \left(\frac{1}{9} \int \frac{a(a-b)(3a^4(225A-539B) - 6a^3b(505A-209B) + 15a^2b^2(19A-121B) + 10ab^3(3A-11B) + 40Ab^4) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx + a(1617a^5B + 3705Aa^4)}{a} \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{11/2}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \int \frac{a(a-b)(3a^4(225A-539B) - 6a^3b(505A-209B) + 15a^2b^2(19A-121B) + 10ab^3(3A-11B) + 40Ab^4) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + a(1617a^5B + 3705Aa^4)}{a} \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{11/2}(c+dx)}$$

↓ 3295

3.418. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{13/2}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{a(1617a^5B + 3705a^4Ab + 3069a^3b^2B + 255a^2Ab^3 - 110ab^4B + 40Ab^5) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(3a^4(225A-539B)-6a^5)}{\dots} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

↓ 3473

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(81a^2A + 209abB + 113Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{7d \cos^{\frac{7}{2}}(c+dx)} + \frac{2(539a^3B + 1145a^2Ab + 825ab^2B + 15Ab^3) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)} \right) \right)$$

$$\frac{2aA \sin(c+dx)(a+b \cos(c+dx))^{3/2}}{11d \cos^{\frac{11}{2}}(c+dx)}$$

```
input Int[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Cos[c + d*x]^(13/2), x]
```

```

output (2*a*A*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(11*d*Cos[c + d*x]^(11/2))
+ ((2*a*(14*A*b + 11*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(9*d*Cos
[c + d*x]^(9/2)) + ((2*(81*a^2*A + 113*A*b^2 + 209*a*b*B)*Sqrt[a + b*Cos[c
+ d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(1145*a^2*A*b + 15*A
*b^3 + 539*a^3*B + 825*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*
d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(3705*a^4*A*b + 255*a^2*A
*b^3 + 40*A*b^5 + 1617*a^5*B + 3069*a^3*b^2*B - 110*a*b^4*B)*Cot[c + d*x]*
EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])
], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + S
ec[c + d*x]))/(a - b))]/(a*d) + (2*(a - b)*Sqrt[a + b]*(40*A*b^4 + 3*a^4*(
225*A - 539*B) - 6*a^3*b*(505*A - 209*B) + 15*a^2*b^2*(19*A - 121*B) + 10*
a*b^3*(3*A - 11*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]
/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c
+ d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))/d)/a + (2*(675*a^
4*A + 1025*a^2*A*b^2 - 20*A*b^4 + 1793*a^3*b*B + 55*a*b^3*B)*Sqrt[a + b*Co
s[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)))/(5*a)/(7*a))/9)/11

```

3.418.3.1 Defintions of rubi rules used

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_)] + (f
_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]

```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

$$3.418. \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx$$

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.418.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 7346 vs. 2(572) = 1144.

Time = 37.18 (sec) , antiderivative size = 7347, normalized size of antiderivative = 11.81

method	result	size
parts	Expression too large to display	7347
default	Expression too large to display	7451

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x,method=_RE
TURNVERBOSE)
```

```
output result too large to display
```

3.418.5 Fricas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos^{13/2}(dx + c)} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x, alg
orithm="fricas")
```

3.418. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{13/2}(c+dx)} dx$

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/cos(d*x + c)^(13/2), x)`

3.418.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)**(13/2),x)`

output `Timed out`

3.418.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{13/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(13/2), x)`

3.418.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\cos(dx + c)^{13/2}} dx$$

3.418. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\cos^{13/2}(c+dx)} dx$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/cos(d*x+c)^(13/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(13/2), x)`

3.418.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{13/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(13/2), x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(13/2), x)`

3.419
$$\int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx)\right)}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.419.1 Optimal result 3908
 3.419.2 Mathematica [A] (verified) 3909
 3.419.3 Rubi [A] (verified) 3909
 3.419.4 Maple [B] (verified) 3914
 3.419.5 Fracas [F] 3915
 3.419.6 Sympy [F(-1)] 3915
 3.419.7 Maxima [F] 3915
 3.419.8 Giac [F] 3916
 3.419.9 Mupad [F(-1)] 3916

3.419.1 Optimal result

Integrand size = 43, antiderivative size = 418

$$\int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx)\right)}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{2(a-b)\sqrt{a+b}(a^2+3b^2)B \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{ad} - \frac{(a-3b)\sqrt{a+b}(2a^2-ab+3b^2)B \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{d} - \frac{b\sqrt{a+b}\left(5a+\frac{3b^2}{a}\right)B \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a+b}}}{d} + \frac{bB(a+b \cos(c+dx))^{3/2} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)}$$

output

```
b*B*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/d/cos(d*x+c)^(3/2)+2*(a-b)*(a^2+3*b^2)*B*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d-(a-3*b)*(2*a^2-a*b+3*b^2)*B*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d-b*(5*a+3*b^2/a)*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d
```

3.419.
$$\int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx)\right)}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.419.2 Mathematica [A] (verified)

Time = 12.69 (sec) , antiderivative size = 458, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx = \frac{B(a + b \cos(c + dx)) (ab + (2a^2 + 7b^2) \cos(c + dx)) \sin(c + dx)}{\cos^{5/2}(c + dx)}$$

input `Integrate[((a + b*Cos[c + d*x])^(5/2)*((3*b*B)/(2*a) + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2),x]`

output `(B*(a + b*Cos[c + d*x])*(a*b + (2*a^2 + 7*b^2)*Cos[c + d*x])*Sin[c + d*x] + (2*B*(Cos[(c + d*x)/2]^2)^(5/2)*(Cos[c + d*x]/(1 + Cos[c + d*x]))^(3/2)*Sqrt[1 + Cos[c + d*x]]*(-2*a*(a^3 + a^2*b + 3*a*b^2 + 3*b^3)*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sec[(c + d*x)/2]^2*Sqrt[((a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2)/(a + b)] + (2*a^4 + 7*a^3*b + a^2*b^2 + 9*a*b^3 - 3*b^4)*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sec[(c + d*x)/2]^2*Sqrt[((a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2)/(a + b)] + 2*b^2*(5*a^2 + 3*b^2)*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sec[(c + d*x)/2]^2*Sqrt[((a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2)/(a + b)] - 2*a*(a^2 + 3*b^2)*(a + b*Cos[c + d*x])*(Cos[c + d*x]*Sec[(c + d*x)/2]^2)^(3/2)*Sec[c + d*x]*Tan[(c + d*x)/2]))/a)/(d*Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]])`

3.419.3 Rubi [A] (verified)

Time = 1.83 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.256$, Rules used = {3042, 3468, 27, 3042, 3470, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} \left(\frac{3bB}{2a} + B \sin(c + dx + \frac{\pi}{2}) \right)}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

3.419. $\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx$

$$\begin{aligned}
& \downarrow \text{3468} \\
& \frac{2}{3} \int \frac{3\sqrt{a+b\cos(c+dx)} \left(2(a^2+3b^2)B + b\left(\frac{3b^2}{a}+5a\right)\cos(c+dx)B \right)}{4\cos^{\frac{3}{2}}(c+dx) \frac{bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\cos^{\frac{3}{2}}(c+dx)}} dx + \\
& \downarrow \text{27} \\
& \frac{1}{2} \int \frac{\sqrt{a+b\cos(c+dx)} \left(2(a^2+3b^2)B + b\left(\frac{3b^2}{a}+5a\right)\cos(c+dx)B \right)}{\cos^{\frac{3}{2}}(c+dx) \frac{bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\cos^{\frac{3}{2}}(c+dx)}} dx + \\
& \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})} \left(2(a^2+3b^2)B + b\left(\frac{3b^2}{a}+5a\right)\sin(c+dx+\frac{\pi}{2})B \right)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \frac{bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\cos^{\frac{3}{2}}(c+dx)}} dx + \\
& \downarrow \text{3470} \\
& \frac{1}{2} \left(\int \frac{2a(a^2+3b^2)B + \left(2b(a^2+3b^2)B + ab\left(\frac{3b^2}{a}+5a\right)B \right)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + b^2B\left(\frac{3b^2}{a}+5a\right) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx \right. \\
& \quad \left. \frac{bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\cos^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow \text{3042} \\
& \frac{1}{2} \left(\int \frac{2a(a^2+3b^2)B + \left(2b(a^2+3b^2)B + ab\left(\frac{3b^2}{a}+5a\right)B \right)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + b^2B\left(\frac{3b^2}{a}+5a\right) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right. \\
& \quad \left. \frac{bB\sin(c+dx)(a+b\cos(c+dx))^{3/2}}{d\cos^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow \text{3288}
\end{aligned}$$

3.419. $\int \frac{(a+b\cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B\cos(c+dx) \right)}{\cos^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{2} \left(\int \frac{2a(a^2 + 3b^2)B + (2b(a^2 + 3b^2)B + ab\left(\frac{3b^2}{a} + 5a\right)B) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{2bB\sqrt{a+b}\left(\frac{3b^2}{a} + 5a\right) \cot(c + dx)}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \right) - \frac{bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3477

$$\frac{1}{2} \left(2aB(a^2 + 3b^2) \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - B(a - 3b)(2a^2 - ab + 3b^2) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \right) - \frac{bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{2} \left(-B(a - 3b)(2a^2 - ab + 3b^2) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + 2aB(a^2 + 3b^2) \int \frac{1}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) - \frac{bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3295

$$\frac{1}{2} \left(2aB(a^2 + 3b^2) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{2B(a - 3b)\sqrt{a+b}(2a^2 - ab + 3b^2) \cot(c + dx)}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} \right) - \frac{bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{d \cos^{\frac{3}{2}}(c + dx)}$$

↓ 3473

$$\frac{1}{2} \left(-\frac{2B(a - 3b)\sqrt{a+b}(2a^2 - ab + 3b^2) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a+b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a-b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c + dx)}}{\sqrt{a+b} \sqrt{\cos(c + dx)}}\right)\right)}{ad} \right) - \frac{bB \sin(c + dx)(a + b \cos(c + dx))^{3/2}}{d \cos^{\frac{3}{2}}(c + dx)}$$

input `Int[((a + b*Cos[c + d*x])^(5/2)*((3*b*B)/(2*a) + B*Cos[c + d*x]))/Cos[c + d*x]^(5/2), x]`

$$3.419. \int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{\frac{5}{2}}(c + dx)} dx$$

```
output ((4*(a - b)*Sqrt[a + b]*(a^2 + 3*b^2)*B*Cot[c + d*x]*EllipticE[ArcSin[Sqrt
[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))
]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b
)]/(a*d) - (2*(a - 3*b)*Sqrt[a + b]*(2*a^2 - a*b + 3*b^2)*B*Cot[c + d*x]*E
llipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])]
, -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Se
c[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(5*a + (3*b^2)/a)*B*Cot[c
+ d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*
Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a +
b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d)/2 + (b*B*(a + b*Cos[c + d*x]
)^(3/2)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))
```

3.419.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f
_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

$$3.419. \int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx) \right)}{\cos^{5/2}(c+dx)} dx$$

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3470 `Int[(((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])]/((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2), x_Symbol] := Simp[B*(d/b^2) Int[Sqrt[b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Int[(A*c + (B*c + A*d)*Sin[e + f*x])/((b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

$$3.419. \int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx) \right)}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.419.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3178 vs. $2(390) = 780$.

Time = 21.44 (sec) , antiderivative size = 3179, normalized size of antiderivative = 7.61

method	result	size
default	Expression too large to display	3179
parts	Expression too large to display	4724

```
input int((a+cos(d*x+c)*b)^(5/2)*(3/2*b*B/a+B*cos(d*x+c))/cos(d*x+c)^(5/2),x,method=_RETURNVERBOSE)
```

```
output B/a/d*(-14*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*b*cos(d*x+c)^2-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b^2*cos(d*x+c)^2-18*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^3*cos(d*x+c)^2-20*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*a^2*b^2*cos(d*x+c)^2+4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*b*cos(d*x+c)^2+12*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b^2*cos(d*x+c)^2-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*a^4*cos(d*x+c)-6*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*b^4*cos(d*x+c)^3+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*a^4*cos(d*x+c)^3+3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x...
```

3.419.
$$\int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx) \right)}{\cos^{\frac{5}{2}}(c+dx)} dx$$

3.419.5 Fricas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx = \int \frac{(2B \cos(dx + c) + \frac{3Bb}{a})(b \cos(dx + c) + a)^{5/2}}{2 \cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(3/2*b*B/a+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="fricas")`

output `integral(1/2*(2*B*a*b^2*cos(d*x + c)^3 + 3*B*a^2*b + (4*B*a^2*b + 3*B*b^3)*cos(d*x + c)^2 + 2*(B*a^3 + 3*B*a*b^2)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/(a*cos(d*x + c)^(5/2)), x)`

3.419.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(3/2*b*B/a+B*cos(d*x+c))/cos(d*x+c)**(5/2),x)`

output `Timed out`

3.419.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx = \int \frac{(2B \cos(dx + c) + \frac{3Bb}{a})(b \cos(dx + c) + a)^{5/2}}{2 \cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(3/2*b*B/a+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/2*integrate((2*B*cos(d*x + c) + 3*B*b/a)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(5/2), x)`

3.419. $\int \frac{(a+b \cos(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c+dx) \right)}{\cos^{5/2}(c+dx)} dx$

3.419.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx = \int \frac{\left(2B \cos(dx + c) + \frac{3Bb}{a} \right) (b \cos(dx + c) + a)^{5/2}}{2 \cos(dx + c)^{5/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(3/2*b*B/a+B*cos(d*x+c))/cos(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate(1/2*(2*B*cos(d*x + c) + 3*B*b/a)*(b*cos(d*x + c) + a)^(5/2)/cos(d*x + c)^(5/2), x)`

3.419.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \cos(c + dx) \right)}{\cos^{5/2}(c + dx)} dx = \int \frac{\left(B \cos(c + dx) + \frac{3Bb}{2a} \right) (a + b \cos(c + dx))^{5/2}}{\cos(c + dx)^{5/2}} dx$$

input `int(((B*cos(c + d*x) + (3*B*b)/(2*a))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(5/2), x)`

output `int(((B*cos(c + d*x) + (3*B*b)/(2*a))*(a + b*cos(c + d*x))^(5/2))/cos(c + d*x)^(5/2), x)`

3.420
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

3.420.1 Optimal result 3917
 3.420.2 Mathematica [C] (verified) 3918
 3.420.3 Rubi [A] (verified) 3919
 3.420.4 Maple [B] (verified) 3924
 3.420.5 Fricas [F] 3925
 3.420.6 Sympy [F(-1)] 3926
 3.420.7 Maxima [F] 3926
 3.420.8 Giac [F] 3926
 3.420.9 Mupad [F(-1)] 3927

3.420.1 Optimal result

Integrand size = 35, antiderivative size = 479

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}(4Ab-3aB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{4ab^2d}$$

$$+ \frac{\sqrt{a+b}(4Ab-3aB+2bB) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{4b^2d}$$

$$+ \frac{\sqrt{a+b}(4aAb-3a^2B-4b^2B) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4b^3d}$$

$$+ \frac{(4Ab-3aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{4b^2d \sqrt{\cos(c+dx)}}$$

$$+ \frac{B \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{2bd}$$

output $\frac{1}{4}(4A^2b-3B^2a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^2/d/\cos(dx+c)^{1/2} + 1/2B\sin(dx+c)\cos(dx+c)^{1/2}(a+b\cos(dx+c))^{1/2}/b/d - 1/4(a-b)(4A^2b-3B^2a)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/b^2/d + 1/4(4A^2b-3B^2a+2B^2b)\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d + 1/4(4A^2ab-3B^2a^2-4B^2b^2)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, (a+b)/b, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^3/d$

3.420.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 21.26 (sec) , antiderivative size = 1175, normalized size of antiderivative = 2.45

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \text{Too large to display}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]], x]`

output $(B\sqrt{\cos[c + dx]}\sqrt{a + b\cos[c + dx]}\sin[c + dx])/(2bd) + ((-4a(4Ab - aB)\sqrt{((a + b)\cot[(c + dx)/2]^2)/(-a + b)}\sqrt{-((a + b)\cos[c + dx]\csc[(c + dx)/2]^2/a)}\sqrt{((a + b\cos[c + dx])\csc[(c + dx)/2]^2/a)\csc[c + dx]}\text{EllipticF}[\text{ArcSin}[\sqrt{((a + b\cos[c + dx])\csc[(c + dx)/2]^2/a)}/\sqrt{2}], (-2a)/(-a + b)]\sin[(c + dx)/2]^4)/((a + b)\sqrt{\cos[c + dx]}\sqrt{a + b\cos[c + dx]}) - 16a*b*B*((\sqrt{((a + b)\cot[(c + dx)/2]^2)/(-a + b)}\sqrt{-((a + b)\cos[c + dx]\csc[(c + dx)/2]^2/a)}\sqrt{((a + b\cos[c + dx])\csc[(c + dx)/2]^2/a)\csc[c + dx]}\text{EllipticF}[\text{ArcSin}[\sqrt{((a + b\cos[c + dx])\csc[(c + dx)/2]^2/a)}/\sqrt{2}], (-2a)/(-a + b)]\sin[(c + dx)/2]^4)/((a + b)\sqrt{\cos[c + dx]}\sqrt{a + b\cos[c + dx]}) - (\sqrt{((a + b)\cot[(c + dx)/2]^2)/(-a + b)}\sqrt{-((a + b)\cos[c + dx]\csc[(c + dx)/2]^2/a)}\sqrt{((a + b\cos[c + dx])\csc[(c + dx)/2]^2/a)\csc[c + dx]}\text{EllipticPi}[-(a/b), \text{ArcSin}[\sqrt{((a + b\cos[c + dx])\csc[(c + dx)/2]^2/a)}/\sqrt{2}], (-2a)/(-a + b)]\sin[(c + dx)/2]^4)/(b\sqrt{\cos[c + dx]}\sqrt{a + b\cos[c + dx]}) + 2*(4Ab - 3aB)*((I*\cos[(c + dx)/2]\sqrt{a + b\cos[c + dx]}\text{EllipticE}[\text{ArcSinh}[\sin[(c + dx)/2]/\sqrt{\cos[c + dx]}], (-2a)/(-a - b)]\sec[c + dx])/(b\sqrt{\cos[(c + dx)/2]^2*\sec[c + dx]}\sqrt{((a + b\cos[c + dx])\sec[c + dx])/(a + b)}) + (2*a*((a*\sqrt{((a + b)\cot[(c + dx)/2]^2)/(-a + b)}\sqrt{-((a + b)\cos[c + dx]\csc[(c + dx)/2]^2/a)}\sqrt{((a + b\cos[c + dx])...$

3.420.3 Rubi [A] (verified)

Time = 2.07 (sec) , antiderivative size = 480, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3469, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3469

$$\frac{\int \frac{(4Ab - 3aB) \cos^2(c + dx) + 2bB \cos(c + dx) + aB}{2\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx}{2b} + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}{2bd}$$

3.420. $\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$

$$\begin{aligned}
 & \int \frac{(4Ab-3aB)\cos^2(c+dx)+2bB\cos(c+dx)+aB}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \\
 & \quad \downarrow 27 \\
 & \int \frac{(4Ab-3aB)\sin(c+dx+\frac{\pi}{2})^2+2bB\sin(c+dx+\frac{\pi}{2})+aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \\
 & \quad \downarrow 3042 \\
 & \int -\frac{(-3Ba^2+4Aba-4b^2B)\cos^2(c+dx)-2abB\cos(c+dx)+a(4Ab-3aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} + \\
 & \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \\
 & \quad \downarrow 25 \\
 & \frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{(-3Ba^2+4Aba-4b^2B)\cos^2(c+dx)-2abB\cos(c+dx)+a(4Ab-3aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \\
 & \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \\
 & \quad \downarrow 3042 \\
 & \frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{(-3Ba^2+4Aba-4b^2B)\sin(c+dx+\frac{\pi}{2})^2-2abB\sin(c+dx+\frac{\pi}{2})+a(4Ab-3aB)}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \\
 & \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \\
 & \quad \downarrow 3532 \\
 & \frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{(-3a^2B+4aAb-4b^2B)\int\frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}}dx+\int\frac{a(4Ab-3aB)-2abB\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{2b} + \\
 & \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.420. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

$$\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{(-3a^2B+4aAb-4b^2B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx + \int\frac{a(4Ab-3aB)-2abB\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{2b} +$$

$$\frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd}$$

↓ 3288

$$\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int\frac{a(4Ab-3aB)-2abB\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx - \frac{2\sqrt{a+b}(-3a^2B+4aAb-4b^2B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{2b}}{4b}$$

$$\frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd}$$

↓ 3477

$$\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(4Ab-3aB)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx - a(-3aB+4Ab+2bB)\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}dx}{4b}$$

$$\frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd}$$

↓ 3042

$$\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{-a(-3aB+4Ab+2bB)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx + a(4Ab-3aB)\int\frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{4b}$$

$$\frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd}$$

↓ 3295

$$\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(4Ab-3aB)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx - \frac{2\sqrt{a+b}(-3a^2B+4aAb-4b^2B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{2b}}{4b}$$

$$\frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd}$$

↓ 3473

3.420. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

$$\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{2\sqrt{a+b}(-3a^2B+4aAb-4b^2B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}}{\sqrt{a+b\cos(c+dx)}}\right)\right)}{bd}$$

$$\frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*b*d) + (-1/2*((2*(a - b)*Sqrt[a + b]*(4*A*b - 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*Sqrt[a + b]*(4*A*b - 3*a*B + 2*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (2*Sqrt[a + b]*(4*a*A*b - 3*a^2*B - 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/b + ((4*A*b - 3*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/(4*b)`

3.420.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.420. \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

```
rule 3477 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] :> Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.420.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2524 vs. $2(437) = 874$.

Time = 15.45 (sec) , antiderivative size = 2525, normalized size of antiderivative = 5.27

method	result	size
parts	Expression too large to display	2525
default	Expression too large to display	2535

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)
```

$$3.420. \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

output $A/d*(-\text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2})*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*a*\cos(dx+c)^2-\text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2}*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*b*\cos(dx+c)^2+2*\text{EllipticPi}(\cot(dx+c)-\csc(dx+c), -1, (-a-b)/(a+b))^{1/2}*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*a*\cos(dx+c)^2-2*\text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2}*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*a*\cos(dx+c)-2*\text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2}*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*b*\cos(dx+c)+4*\text{EllipticPi}(\cot(dx+c)-\csc(dx+c), -1, (-a-b)/(a+b))^{1/2}*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*a*\cos(dx+c)+b*\cos(dx+c)^2*\sin(dx+c)-(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*\text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*a-(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*\text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*b+2*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}*\text{EllipticPi}(\cot(dx+c)-\csc(dx+c), -1, (-a-b)/(a+b))^{1/2}*((a+\cos(dx+c)*b)/(1+\cos(dx+c)))/(a+b)^{1/2}*a+\sin(dx+c)*\cos(dx+c)*a/(1+\cos(dx+c))/(a+\cos(dx+c)*b)^{1/2}/\cos(dx+c)^{1/2}/b+1/4*B/d*(-2*\text{EllipticF}(\cot(dx+c)-\csc(dx+c), (...$

3.420.5 Fracas [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(dx+c)^(3/2)*(A+B*cos(dx+c))/(a+b*cos(dx+c))^(1/2),x, algorithm="fracas")`

output `integral((B*cos(dx+c)^2 + A*cos(dx+c))*sqrt(cos(dx+c))/sqrt(b*cos(dx+c)+a), x)`

3.420.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.420.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a)
, x)`

3.420.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a)
, x)`

3.420.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

3.421
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

3.421.1 Optimal result 3928
 3.421.2 Mathematica [A] (verified) 3929
 3.421.3 Rubi [A] (verified) 3929
 3.421.4 Maple [B] (verified) 3935
 3.421.5 Fricas [F] 3936
 3.421.6 Sympy [F] 3936
 3.421.7 Maxima [F] 3936
 3.421.8 Giac [F] 3937
 3.421.9 Mupad [F(-1)] 3937

3.421.1 Optimal result

Integrand size = 35, antiderivative size = 427

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}B \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{abd}$$

$$+ \frac{\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{bd}$$

$$- \frac{\sqrt{a+b}(2Ab-aB) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^2d}$$

$$+ \frac{aB \sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} + \frac{B\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{a+b \cos(c+dx)}}$$

output

```
a*B*sin(d*x+c)/b/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)+B*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+b*cos(d*x+c))^(1/2)-(a-b)*B*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/b/d+B*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b/d-(2*A*b-B*a)*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2), (a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b^2/d
```

3.421.
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx$$

3.421.2 Mathematica [A] (verified)

Time = 7.04 (sec) , antiderivative size = 402, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

$$= \frac{\sqrt{\cos(c+dx)} \left(2(a+b)B \sqrt{\frac{a+b\cos(c+dx)}{(a+b)(1+\cos(c+dx))}} E\left(\arcsin\left(\tan\left(\frac{1}{2}(c+dx)\right)\right) \middle| \frac{-a+b}{a+b}\right) - 4Ab \sqrt{\frac{a+b\cos(c+dx)}{(a+b)(1+\cos(c+dx))}} E\left(\arcsin\left(\tan\left(\frac{1}{2}(c+dx)\right)\right) \middle| \frac{-a+b}{a+b}\right) \right)}{1}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(Sqrt[Cos[c + d*x]]*(2*(a + b)*B*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 4*A*b*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 8*A*b*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 4*a*B*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + b*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sin[(3*(c + d*x))/2] + 2*a*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Tan[(c + d*x)/2] - b*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Tan[(c + d*x)/2]))/(2*b*d*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[a + b*Cos[c + d*x]])`

3.421.3 Rubi [A] (verified)Time = 1.79 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3482, 3042, 3530, 3042, 3288, 3472, 25, 27, 3042, 3280, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

↓ 3042

3.421. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3482} \\
& \frac{1}{2} \int \frac{(2Ab-aB)\cos^2(c+dx)+2aA\cos(c+dx)+aB}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{(2Ab-aB)\sin(c+dx+\frac{\pi}{2})^2+2aA\sin(c+dx+\frac{\pi}{2})+aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3530} \\
& \frac{1}{2} \left(\frac{\int \frac{B\cos(c+dx)a^2+bBa}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{b} + \frac{(2Ab-aB) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx}{b} \right) + \\
& \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(\frac{\int \frac{B\sin(c+dx+\frac{\pi}{2})a^2+bBa}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} + \frac{(2Ab-aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \\
& \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3288} \\
& \frac{1}{2} \left(\frac{\int \frac{B\sin(c+dx+\frac{\pi}{2})a^2+bBa}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{b^2d} \text{Elliptic} \right) \\
& \quad \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3472}
\end{aligned}$$

3.421. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$

$$\frac{1}{2} \left(\frac{\int -\frac{a(a^2-b^2)B}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)\sqrt{a+b\cos(c+dx)}}}}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{b} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}$$

↓ 25

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)\sqrt{a+b\cos(c+dx)}}} - \frac{\int -\frac{a(a^2-b^2)B}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{b} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}$$

↓ 27

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)\sqrt{a+b\cos(c+dx)}}} - aB \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{b} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}$$

↓ 3042

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)\sqrt{a+b\cos(c+dx)}}} - aB \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{b} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}$$

↓ 3280

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)\sqrt{a+b\cos(c+dx)}}} - aB \left(\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - \int \frac{1}{\sqrt{\cos(c+dx)\sqrt{a+b\cos(c+dx)}}} dx \right)}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{b} \right)$$

$$\frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}$$

3.421. $\int \frac{\sqrt{\cos(c+dx)(A+B\cos(c+dx))}}{\sqrt{a+b\cos(c+dx)}} dx$

↓ 3042

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \left(\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx \right)}{b} - \frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b \cos(c+dx)}} \right)$$

↓ 3295

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \left(\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{a^2 d} \right)}{b} - \frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b \cos(c+dx)}} \right)$$

↓ 3473

$$\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \left(\frac{2(a-b)\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E \left(\arcsin \left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}} \right) \right) - \frac{a+b}{a-b}}{a^2 d} \right)}{b} - \frac{B \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a+b \cos(c+dx)}} \right)$$

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[a + b*Cos[c + d*x]],x]`

```
output (B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + b*Cos[c + d*x]]) + ((-2*Sq
rt[a + b]*(2*A*b - a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a +
b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sq
rt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(
b^2*d) + (-a*B*((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt
[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)
]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b
)]/(a^2*d) - (2*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c
+ d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1
- Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d))) +
(2*a*B*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))/b/
2
```

3.421.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3280 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin
[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[1/(a - b) Int[1/(Sqrt[a + b*Sin
[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[b/(a - b) Int[(1 + Si
n[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /
; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0]
```

```
rule 3288 Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

rule 3473 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3482 `Int[Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*B*Cos[e + f*x]*Sqrt[a + b*Sin[e + f*x]]*((c + d*Sin[e + f*x])^n/(f*(2*n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*Sin[e + f*x])^(n - 1)/Sqrt[a + b*Sin[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*Sin[e + f*x] + (A*b*d*(2*n + 3) + B*(a*d + 2*b*c*n))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && EqQ[n^2, 1/4]`

rule 3530 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[C/(b*d) Int[Sqrt[d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[1/b Int[(A*b + (b*B - a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]`

3.421.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 980 vs. $2(395) = 790$.

Time = 11.54 (sec) , antiderivative size = 981, normalized size of antiderivative = 2.30

method	result	size
parts	Expression too large to display	981
default	Expression too large to display	1469

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 2*A/d*(EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))-2*EllipticPi(
cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2)))*(cos(d*x+c)/(1+cos(d*x+c))
)^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)/(a+cos(d*x+c)*b)^(1/
2)*(1+cos(d*x+c))/cos(d*x+c)^(1/2)+B/d*(-EllipticE(cot(d*x+c)-csc(d*x+c),
-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(
1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2-EllipticE(cot(d*x+c)-csc(d*x+c),
-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/
(1+cos(d*x+c))/(a+b))^(1/2)*b*cos(d*x+c)^2+2*EllipticPi(cot(d*x+c)-csc(d*x
+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x
+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2-2*EllipticE(cot(d*x+c)-c
sc(d*x+c),-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(
d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)-2*EllipticE(cot(d*x+c)-
csc(d*x+c),-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos
(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*b*cos(d*x+c)+4*EllipticPi(cot(d*x+c
)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((
a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)+b*cos(d*x+c)^2*si
n(d*x+c)-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c)
,(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a-(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),-(a-b)/(a+
b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*b+2*(cos(d*x+c...
```

3.421.5 Fricas [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/sqrt(b*cos(d*x + c) + a),
x)`

3.421.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(A+B\cos(c+dx))\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(cos(c + d*x))/sqrt(a + b*cos(c + d*x)),
x)`

3.421.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/sqrt(b*cos(d*x + c) + a)
, x)`

3.421.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{\sqrt{b\cos(dx+c)+a}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/sqrt(b*cos(d*x + c) + a), x)`

3.421.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(1/2),x)`

$$3.422 \quad \int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx$$

3.422.1 Optimal result	3938
3.422.2 Mathematica [A] (verified)	3939
3.422.3 Rubi [A] (verified)	3939
3.422.4 Maple [A] (verified)	3941
3.422.5 Fricas [F]	3942
3.422.6 Sympy [F]	3942
3.422.7 Maxima [F]	3942
3.422.8 Giac [F]	3943
3.422.9 Mupad [F(-1)]	3943

3.422.1 Optimal result

Integrand size = 35, antiderivative size = 228

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2A\sqrt{a + b} \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ad} - \frac{2\sqrt{a + b} B \cot(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{bd}$$

```
output 2*A*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d-2*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2), (a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b/d
```

3.422.2 Mathematica [A] (verified)

Time = 2.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.63

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2\sqrt{2} \sqrt{\cos(c + dx)} \sqrt{\frac{a+b \cos(c+dx)}{(a+b)(1+\cos(c+dx))}} \left((A - B) \operatorname{EllipticF} \left(\arcsin \left(\tan \left(\frac{1}{2}(c + dx) \right) \right), \frac{-a+b}{a+b} \right) + 2B \operatorname{EllipticPi} \right)}{d \sqrt{a + b \cos(c + dx)} \sqrt{\cos(c + dx) \sec^2 \left(\frac{1}{2}(c + dx) \right)}}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]),x]`

output `(2*Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))])*((A - B)*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)])/(d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Cos[c + d*x]*Sec[(c + d*x)/2]^2])`

3.422.3 Rubi [A] (verified)Time = 0.62 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 3485, 3042, 3288, 3295}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin \left(c + dx + \frac{\pi}{2} \right)}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} \sqrt{a + b \sin \left(c + dx + \frac{\pi}{2} \right)}} dx$$

$$\downarrow \text{3485}$$

$$A \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx + B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

3.422. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& A \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3288} \\
& A \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \\
& \frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{bd} \\
& \quad \downarrow \text{3295} \\
& \frac{2A\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{ad} \\
& \frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{bd}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]),x]`

output `(2*A*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d) - (2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b*d)`

3.422.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

3.422. $\int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx$

```
rule 3295 Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3485 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] :> Simp[B/d Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[(B*c - A*d)/d Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.422.4 Maple [A] (verified)

Time = 14.27 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.79

method	result
default	$-\frac{2\left(AF\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)-BF\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)+2B\Pi\left(\cot(dx+c)-\csc(dx+c),-1,\sqrt{-\frac{a-b}{a+b}}\right)\right)\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}}{d\sqrt{a+\cos(dx+c)b}\sqrt{\cos(dx+c)}}$
parts	$-\frac{2A(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}F\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}}{d\sqrt{a+\cos(dx+c)b}\sqrt{\cos(dx+c)}} + \frac{2B\left(F\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)\right)}{\sqrt{\cos(dx+c)}}$

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -2/d*(A*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))-B*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))+2*B*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2)))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2))/cos(d*x+c)^(1/2)
```

3.422.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algo rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/ (b*cos(d*x + c)^2 + a*cos(d*x + c)), x)`

3.422.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a + b*cos(c + d*x))*sqrt(cos(c + d*x))), x)`

3.422.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))), x)`

3.422.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c)
)), x)`

3.422.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(1/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(1/2)),
x)`

3.423
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

3.423.1 Optimal result 3944
 3.423.2 Mathematica [A] (verified) 3945
 3.423.3 Rubi [A] (verified) 3945
 3.423.4 Maple [B] (verified) 3948
 3.423.5 Fricas [F] 3949
 3.423.6 Sympy [F] 3949
 3.423.7 Maxima [F] 3949
 3.423.8 Giac [F] 3950
 3.423.9 Mupad [F(-1)] 3950

3.423.1 Optimal result

Integrand size = 35, antiderivative size = 230

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

$$= \frac{2A(a-b)\sqrt{a+b} \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{a^2 d}$$

$$= \frac{2\sqrt{a+b}(A-B) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ad}$$

```
output 2*A*(a-b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d-2*(A-B)*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d
```

3.423.2 Mathematica [A] (verified)

Time = 9.68 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.30

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

$$= 2 \left(A(a + b \cos(c + dx)) \sin(c + dx) - \frac{2\sqrt{2} \cos^2(\frac{1}{2}(c+dx))^{3/2} (2A(a+b) \cos^2(\frac{1}{2}(c+dx)) \sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}} \sqrt{\frac{a+b \cos(c+dx)}{(a+b)(1+\cos(c+dx))}} E(\arcsin(\frac{\cos(c+dx)}{1+\cos(c+dx)})) - \dots}{\dots} \right)$$

ad√

```
input Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]),x]
```

```
output (2*(A*(a + b*Cos[c + d*x])*Sin[c + d*x] - (2*Sqrt[2]*(Cos[(c + d*x)/2]^2)^(3/2)*(2*A*(a + b)*Cos[(c + d*x)/2]^2*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] *Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 2*a*(A + B)*Cos[(c + d*x)/2]^2*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] *Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + A *Cos[c + d*x]*(a + b*Cos[c + d*x])*Tan[(c + d*x)/2]))/(1 + Cos[c + d*x])^(3/2)))/(a*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])
```

3.423.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3477

3.423. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& A \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (A-B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \\
& \quad \downarrow \text{3042} \\
& A \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3295} \\
& A \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \\
& \frac{2\sqrt{a+b}(A-B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{ad} \\
& \quad \downarrow \text{3473} \\
& \frac{2A(a-b)\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b\cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\middle|-\frac{a+b}{a-b}\right)}{2\sqrt{a+b}(A-B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b\cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)} \\
& \quad \downarrow \\
& \frac{a^2d}{ad}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]),x]`

output `(2*A*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a^2*d) - (2 *Sqrt[a + b]*(A - B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d)`

3.423.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_ + (b_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3473 `Int[((A_) + (B_)*sin[(e_.) + (f_)*(x_)])/(((b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

3.423.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 770 vs. 2(214) = 428.

Time = 15.74 (sec) , antiderivative size = 771, normalized size of antiderivative = 3.35

method	result
parts	$2A \frac{-\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \sqrt{\frac{(\csc^2(dx+c))^a(1-\cos(dx+c))^2-(\csc^2(dx+c))^b(1-\cos(dx+c))^2+a+b}{a+b}}}{a+b} F\left(\cot(dx+c)-\csc(dx+c), \frac{-a-b}{a+b}\right)$
default	Expression too large to display

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(1/2), x, method=_RETURNVERBOSE)
```

```
output -2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b)))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))*b+csc(d*x+c)^3*(1-cos(d*x+c))^3*a-csc(d*x+c)^3*(1-cos(d*x+c))^3*b+a*(csc(d*x+c)-cot(d*x+c))+b*(csc(d*x+c)-cot(d*x+c)))*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(3/2)/a-2*B/d*(1+cos(d*x+c))*cos(d*x+c)/(1+cos(d*x+c))^(1/2)/(a+cos(d*x+c)*b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)/cos(d*x+c)^(1/2)
```

3.423.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

3.423.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a \cos(dx + c)}^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(b*cos(d*x + c)^3 + a*cos(d*x + c)^2), x)`

3.423.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a + b*cos(c + d*x))*cos(c + d*x)**(3/2)), x)`

3.423.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a \cos(dx + c)}^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(3/2)), x)`

3.423.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(3/2
)), x)`

3.423.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{3}{2}} \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(1/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(1/2)),
x)`

$$3.424 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

3.424.1 Optimal result	3951
3.424.2 Mathematica [A] (verified)	3952
3.424.3 Rubi [A] (verified)	3952
3.424.4 Maple [B] (verified)	3955
3.424.5 Fracas [F]	3956
3.424.6 Sympy [F]	3957
3.424.7 Maxima [F]	3957
3.424.8 Giac [F]	3957
3.424.9 Mupad [F(-1)]	3958

3.424.1 Optimal result

Integrand size = 35, antiderivative size = 290

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx =$$

$$\frac{2(a-b)\sqrt{a+b}(2Ab-3aB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a^3 d}$$

$$+ \frac{2\sqrt{a+b}(2Ab+a(A-3B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a^2 d}$$

$$+ \frac{2A\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3ad \cos^{\frac{3}{2}}(c+dx)}$$

```
output 2/3*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d/cos(d*x+c)^(3/2)-2/3*(a-b)*(2*
A*b-3*B*a)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x
+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)
*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d+2/3*(2*A*b+a*(A-3*B))*cot(d*x+c)*Ell
ipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(
1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b)
^(1/2)/a^2/d
```

3.424.2 Mathematica [A] (verified)

Time = 12.14 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.43

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{8 \cos^2\left(\frac{1}{2}(c + dx)\right)^{7/2} \sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}} \sqrt{\cos(c + dx) \sec^2\left(\frac{1}{2}(c + dx)\right)} \left(-2(a + b)(-2Ab + 3aB)\sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}}\right)}{d \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)} \left(\frac{2 \sec(c+dx)(-2Ab \sin(c+dx) + 3aB \sin(c+dx))}{3a^2} + \frac{2A \sec(c+dx) \tan(c+dx)}{3a}\right)}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*Sqrt[a + b*Cos[c + d*x]]),x]`

output `(8*(Cos[(c + d*x)/2]^2)^(7/2)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[Cos[c + d*x]*Sec[(c + d*x)/2]^2]*(-2*(a + b)*(-2*A*b + 3*a*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/(a + b)*(1 + Cos[c + d*x])])*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*a*(-2*A*b + a*(A + 3*B))*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/(a + b)*(1 + Cos[c + d*x])])*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + (2*A*b - 3*a*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/(3*a^2*d*Cos[c + d*x]^(3/2)*(1 + Cos[c + d*x])^(3/2)*Sqrt[a + b*Cos[c + d*x]]) + (Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])*((2*Sec[c + d*x]*(-2*A*b*Sin[c + d*x] + 3*a*B*Sin[c + d*x]))/(3*a^2) + (2*A*Sec[c + d*x]*Tan[c + d*x])/(3*a)))/d`

3.424.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3479, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

3.424. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{2 \int -\frac{2Ab - 3aB - aA \cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{2Ab - 3aB - aA \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{2Ab - 3aB - aA \sin\left(c+dx + \frac{\pi}{2}\right)}{\sin\left(c+dx + \frac{\pi}{2}\right)^{3/2} \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx}{3a} \\
 & \quad \downarrow \text{3477} \\
 & \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(2Ab - 3aB) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - (a(A - 3B) + 2Ab) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx}{3a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(2Ab - 3aB) \int \frac{\sin\left(c+dx + \frac{\pi}{2}\right)+1}{\sin\left(c+dx + \frac{\pi}{2}\right)^{3/2} \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx - (a(A - 3B) + 2Ab) \int \frac{1}{\sqrt{\sin\left(c+dx + \frac{\pi}{2}\right)} \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx}{3a} \\
 & \quad \downarrow \text{3295} \\
 & \frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(2Ab - 3aB) \int \frac{\sin\left(c+dx + \frac{\pi}{2}\right)+1}{\sin\left(c+dx + \frac{\pi}{2}\right)^{3/2} \sqrt{a+b \sin\left(c+dx + \frac{\pi}{2}\right)}} dx - \frac{2\sqrt{a+b}(a(A-3B)+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticF}}{ad}}{3a} \\
 & \quad \downarrow \text{3473}
 \end{aligned}$$

3.424. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$

$$\frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3ad \cos^{\frac{3}{2}}(c + dx)} - \frac{2(a-b)\sqrt{a+b}(2Ab-3aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right)}{a^2 d} - \frac{2\sqrt{a+b}(a(A-3B)+2Ab) \cot(c+dx)}{3a}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*Sqrt[a + b*Cos[c + d*x]]),x]`

output `-1/3*((2*(a - b)*Sqrt[a + b]*(2*A*b - 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]), -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*Sqrt[a + b]*(2*A*b + a*(A - 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]), -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/a + (2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2))`

3.424.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_
.)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

```
rule 3479 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m +
1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e
+ f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +
2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)
*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n
}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Rat
ionalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(I
ntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0]
))
```

3.424.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1843 vs. $2(264) = 528$.

Time = 19.46 (sec) , antiderivative size = 1844, normalized size of antiderivative = 6.36

method	result	size
parts	Expression too large to display	1844
default	Expression too large to display	2068

$$3.424. \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx$$

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

output `2/3*A/d*(-cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)^3+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b*cos(d*x+c)^3-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b*cos(d*x+c)^3-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b^2*cos(d*x+c)^3-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)^2+4*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)^2-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b^2*cos(d*x+c)^2-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*...`

3.424.5 Fracas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos^{\frac{5}{2}}(dx + c)} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(1/2),x,algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(b*cos(d*x + c)^4 + a*cos(d*x + c)^3), x)`

3.424.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \cos^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a + b*cos(c + d*x))*cos(c + d*x)**(5/2)), x)`

3.424.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.424.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(5/2)), x)`

3.424.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{\frac{5}{2}} \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^(1/2)),x)`

$$3.425 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

3.425.1 Optimal result	3959
3.425.2 Mathematica [C] (verified)	3960
3.425.3 Rubi [A] (verified)	3960
3.425.4 Maple [B] (verified)	3965
3.425.5 Fricas [F]	3966
3.425.6 Sympy [F(-1)]	3966
3.425.7 Maxima [F]	3966
3.425.8 Giac [F]	3967
3.425.9 Mupad [F(-1)]	3967

3.425.1 Optimal result

Integrand size = 35, antiderivative size = 363

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

$$= \frac{2(a-b)\sqrt{a+b}(9a^2A+8Ab^2-10abB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{a}}{15a^4d}$$

$$- \frac{2\sqrt{a+b}(8Ab^2+a^2(9A-5B)-2ab(A+5B)) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{15a^3d}$$

$$+ \frac{2A\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{15a^2d \cos^{\frac{3}{2}}(c+dx)}$$

```
output 2/5*A*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d/cos(d*x+c)^(5/2)-2/15*(4*A*b-5
*B*a)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a^2/d/cos(d*x+c)^(3/2)+2/15*(a-b)*
(9*A*a^2+8*A*b^2-10*B*a*b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a
b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x
c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^4/d-2/15*(8*A*b^2+a^2*(9
*A-5*B)-2*a*b*(A+5*B))*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(
1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))
(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^3/d
```

3.425.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.59 (sec) , antiderivative size = 1319, normalized size of antiderivative = 3.63

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(7/2)*Sqrt[a + b*Cos[c + d*x]]),x]`

output `-1/15*((-4*a*(7*a^2*A*b + 8*A*b^3 - 5*a^3*B - 10*a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(9*a^3*A + 8*a*A*b^2 - 10*a^2*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(9*a^2*A*b + 8*A*b^3 - 10*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + ...`

3.425.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.425. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{7/2} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
& \quad \downarrow \text{3479} \\
& \frac{2 \int -\frac{-2Ab \cos^2(c+dx) - 3aA \cos(c+dx) + 4Ab - 5aB}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a} + \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5ad \cos^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5ad \cos^{\frac{5}{2}}(c + dx)} - \frac{\int -\frac{2Ab \cos^2(c+dx) - 3aA \cos(c+dx) + 4Ab - 5aB}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{5a} \\
& \quad \downarrow \text{3042} \\
& \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5ad \cos^{\frac{5}{2}}(c + dx)} - \frac{\int -\frac{2Ab \sin(c+dx+\frac{\pi}{2})^2 - 3aA \sin(c+dx+\frac{\pi}{2}) + 4Ab - 5aB}{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{5a} \\
& \quad \downarrow \text{3534} \\
& \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5ad \cos^{\frac{5}{2}}(c + dx)} - \\
& \frac{2 \int -\frac{9Aa^2 - 10bBa + (2Ab + 5aB) \cos(c+dx)a + 8Ab^2}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(4Ab - 5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{2A \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{5ad \cos^{\frac{5}{2}}(c + dx)} - \\
& \frac{2(4Ab - 5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{9Aa^2 - 10bBa + (2Ab + 5aB) \cos(c+dx)a + 8Ab^2}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.425. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{9Aa^2-10bBa+(2Ab+5aB) \sin(c+dx+\frac{\pi}{2})a+8Ab^2}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{3a}$$

5a
↓ 3477

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(9a^2A-10abB+8Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx - (a^2(9A-5B)-2ab(A+5B)+8Ab^2) \int \frac{1}{\sqrt{\cos(c+dx)}}$$

5a
↓ 3042

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(9a^2A-10abB+8Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - (a^2(9A-5B)-2ab(A+5B)+8Ab^2) \int \frac{1}{\sqrt{\cos(c+dx)}}$$

5a
↓ 3295

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(9a^2A-10abB+8Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(a^2(9A-5B)-2ab(A+5B)+8Ab^2)}{3a}$$

5a
↓ 3473

$$\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2(a-b)\sqrt{a+b}(9a^2A-10abB+8Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{a^2d}$$

5a

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(7/2)*Sqrt[a + b*Cos[c + d*x]]),x]`

3.425. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$

```
output (2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*a*d*Cos[c + d*x]^(5/2)) - (
-1/3*((2*(a - b)*Sqrt[a + b]*(9*a^2*A + 8*A*b^2 - 10*a*b*B)*Cot[c + d*x]*E
llipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])]
, -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Se
c[c + d*x]))/(a - b))]/(a^2*d) - (2*Sqrt[a + b]*(8*A*b^2 + a^2*(9*A - 5*B)
- 2*a*b*(A + 5*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]
/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c
+ d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/a + (2*(4*
A*b - 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3
/2)))/(5*a)
```

3.425.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)]*Sqrt[(a_) + (b_)*sin[(e_)] + (f
_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_)] + (f_)*(x_)]/(((b_)*sin[(e_)] + (f_)*(x_))
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

3.425.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3289 vs. $2(331) = 662$.

Time = 22.40 (sec) , antiderivative size = 3290, normalized size of antiderivative = 9.06

method	result	size
parts	Expression too large to display	3290
default	Expression too large to display	3338

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -2/15*A/d*(-8*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d
*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*
a*b^2*cos(d*x+c)^4+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))
*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))
^(1/2)*a^2*b*cos(d*x+c)^4+8*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))
^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))
/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4-18*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)
)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos
(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3-16*EllipticE(cot(d*x+c)-csc(d*x+c)
),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b
)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3+4*EllipticF(cot(d*x+c)-cs
c(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d
*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3+16*EllipticF(cot(d
*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((
a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3-9*Elliptic
E(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^2+18*
EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d
*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^3*cos(d*x+c)
^3-9*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/...
```

3.425.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/
(b*cos(d*x + c)^5 + a*cos(d*x + c)^4), x)`

3.425.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(7/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.425.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(7/2
)), x)`

3.425.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \cos(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(7/2)/(a+b*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*cos(d*x + c)^(7/2)), x)`

3.425.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{7/2} \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(7/2)*(a + b*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(7/2)*(a + b*cos(c + d*x))^(1/2)),x)`

3.426 $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$

3.426.1 Optimal result	3968
3.426.2 Mathematica [C] (warning: unable to verify)	3969
3.426.3 Rubi [A] (verified)	3970
3.426.4 Maple [B] (warning: unable to verify)	3976
3.426.5 Fricas [F]	3977
3.426.6 Sympy [F(-1)]	3977
3.426.7 Maxima [F]	3977
3.426.8 Giac [F]	3978
3.426.9 Mupad [F(-1)]	3978

3.426.1 Optimal result

Integrand size = 35, antiderivative size = 500

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx = \frac{(2aAb - 3a^2B + b^2B) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a-b}{a+b}\right) - (2Ab - (3a+b)B) \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ab^2 \sqrt{a+bd}} - \frac{\sqrt{a+b}(2Ab - 3aB) \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^3 d} + \frac{2a(Ab - aB) \sqrt{\cos(c+dx)} \sin(c+dx)}{b(a^2 - b^2) d \sqrt{a+b \cos(c+dx)}} - \frac{(2aAb - 3a^2B + b^2B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{b^2(a^2 - b^2) d \sqrt{\cos(c+dx)}}$$

3.426. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$

output $2*a*(A*b-B*a)*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b/(a^2-b^2)/d/(a+b*\cos(d*x+c))^{(1/2)}-(2*A*a*b-3*B*a^2+B*b^2)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/b^2/(a^2-b^2)/d/\cos(d*x+c)^{(1/2)}+(2*A*a*b-3*B*a^2+B*b^2)*\cot(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a*(1-\sec(d*x+c))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c))/(a-b))^{(1/2)}/a/b^2/d/(a+b)^{(1/2)}-(2*A*b-(3*a+b)*B)*\cot(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a*(1-\sec(d*x+c))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c))/(a-b))^{(1/2)}/b^2/d/(a+b)^{(1/2)}-(2*A*b-3*B*a)*\cot(d*x+c)*\text{EllipticPi}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},(a+b)/b,((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*(a*(1-\sec(d*x+c))/(a+b))^{(1/2)}*(a*(1+\sec(d*x+c))/(a-b))^{(1/2)}/b^3/d$

3.426.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.83 (sec) , antiderivative size = 1234, normalized size of antiderivative = 2.47

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = \text{Too large to display}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]`

output $(2\sqrt{\cos[c + dx]}*(-(aA*b*\sin[c + dx]) + a^2*B*\sin[c + dx]))/(b*(-a^2 + b^2)*d*\sqrt{a + b*\cos[c + dx]}) + ((-4*a*(a^2*B - b^2*B)*\sqrt{((a + b)*\cot[(c + dx)/2]^2)/(-a + b)}*\sqrt{-(((a + b)*\cos[c + dx]*\csc[(c + dx)/2]^2)/a})*\sqrt{((a + b*\cos[c + dx])*\csc[(c + dx)/2]^2)/a}*\csc[c + dx]*\text{EllipticF}[\text{ArcSin}[\sqrt{((a + b*\cos[c + dx])*\csc[(c + dx)/2]^2)/a}/\sqrt{2}], (-2*a)/(-a + b)]*\sin[(c + dx)/2]^4)/((a + b)*\sqrt{\cos[c + dx]}*\sqrt{a + b*\cos[c + dx]}) - 4*a*(-2*A*b^2 + 2*a*b*B)*((\sqrt{((a + b)*\cot[(c + dx)/2]^2)/(-a + b)}*\sqrt{-(((a + b)*\cos[c + dx]*\csc[(c + dx)/2]^2)/a})*\sqrt{((a + b*\cos[c + dx])*\csc[(c + dx)/2]^2)/a}*\csc[c + dx]*\text{EllipticF}[\text{ArcSin}[\sqrt{((a + b*\cos[c + dx])*\csc[(c + dx)/2]^2)/a}/\sqrt{2}], (-2*a)/(-a + b)]*\sin[(c + dx)/2]^4)/((a + b)*\sqrt{\cos[c + dx]}*\sqrt{a + b*\cos[c + dx]}) - (\sqrt{((a + b)*\cot[(c + dx)/2]^2)/(-a + b)}*\sqrt{-(((a + b)*\cos[c + dx]*\csc[(c + dx)/2]^2)/a})*\sqrt{((a + b*\cos[c + dx])*\csc[(c + dx)/2]^2)/a}*\csc[c + dx]*\text{EllipticPi}[-(a/b), \text{ArcSin}[\sqrt{((a + b*\cos[c + dx])*\csc[(c + dx)/2]^2)/a}/\sqrt{2}], (-2*a)/(-a + b)]*\sin[(c + dx)/2]^4)/(b*\sqrt{\cos[c + dx]}*\sqrt{a + b*\cos[c + dx]}) + 2*(-2*a*A*b + 3*a^2*B - b^2*B)*((I*\cos[(c + dx)/2]*\sqrt{a + b*\cos[c + dx]}*\text{EllipticE}[I*\text{ArcSinh}[\sin[(c + dx)/2]/\sqrt{\cos[c + dx]}]], (-2*a)/(-a - b)]*\sec[c + dx])/(b*\sqrt{\cos[(c + dx)/2]^2*\sec[c + dx]}*\sqrt{((a + b*\cos[c + dx])*sec[c + dx])/(a + b)}) + (2*a*((a*\sqrt{((a + b)*\cot[(c + dx)/2]^2)/(-a + b)}*\sqrt{...$

3.426.3 Rubi [A] (verified)

Time = 2.43 (sec) , antiderivative size = 527, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3468, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2})^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx$$

↓ 3468

3.426. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} - \\
& \frac{2 \int - \frac{((-3Ba^2 + 2Aba + b^2B) \cos^2(c + dx)) - b(Ab - aB) \cos(c + dx) + a(Ab - aB)}{2\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx}{b(a^2 - b^2)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{((-3Ba^2 + 2Aba + b^2B) \cos^2(c + dx)) - b(Ab - aB) \cos(c + dx) + a(Ab - aB)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx}{b(a^2 - b^2)} + \\
& \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(3Ba^2 - 2Aba - b^2B) \sin(c + dx + \frac{\pi}{2})^2 - b(Ab - aB) \sin(c + dx + \frac{\pi}{2}) + a(Ab - aB)}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{b(a^2 - b^2)} + \\
& \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow 3540 \\
& \frac{\int \frac{(a^2 - b^2)(2Ab - 3aB) \cos^2(c + dx) + 2ab(Ab - aB) \cos(c + dx) + a(-3Ba^2 + 2Aba + b^2B)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{2b} - \frac{(-3a^2B + 2aAb + b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} + \\
& \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(a^2 - b^2)(2Ab - 3aB) \sin(c + dx + \frac{\pi}{2})^2 + 2ab(Ab - aB) \sin(c + dx + \frac{\pi}{2}) + a(-3Ba^2 + 2Aba + b^2B)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2b} - \frac{(-3a^2B + 2aAb + b^2B) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{bd \sqrt{\cos(c + dx)}} + \\
& \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow 3532
\end{aligned}$$

3.426. $\int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$

$$\frac{\int \frac{a(-3Ba^2+2Aba+b^2B)+2ab(Ab-aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + (a^2-b^2)(2Ab-3aB) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx}{2b} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

↓ 3042

$$\frac{(a^2-b^2)(2Ab-3aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{a(-3Ba^2+2Aba+b^2B)+2ab(Ab-aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{(-3a^2B+2aAb+b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

$$\frac{b(a^2-b^2)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

↓ 3288

$$\frac{\int \frac{a(-3Ba^2+2Aba+b^2B)+2ab(Ab-aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(a^2-b^2)(2Ab-3aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{a+b}{b}\right)\right)}{2b}}{2b} - \frac{b(a^2-b^2)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

↓ 3477

$$\frac{a(-3a^2B+2aAb+b^2B) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - a(a-b)(2Ab-B(3a+b)) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx - \frac{2\sqrt{a+b}(a^2-b^2)(2Ab-3aB)\cot(c+dx)}{2b}}{2b} - \frac{b(a^2-b^2)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

↓ 3042

$$\frac{a(-3a^2B+2aAb+b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - a(a-b)(2Ab-B(3a+b)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(a^2-b^2)}{2b}}{2b} - \frac{b(a^2-b^2)}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}}$$

↓ 3295

3.426. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx$

$$a(-3a^2B+2aAb+b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(a^2-b^2)(2Ab-3aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a}{a-b}\right)}{bd}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd (a^2 - b^2) \sqrt{a + b \cos(c + dx)}}$$

↓ 3473

$$\frac{2(a-b)\sqrt{a+b}(-3a^2B+2aAb+b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) - 2\sqrt{a+b}(a^2-b^2)(2Ab-3aB) \cot(c+dx)}{ad}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd (a^2 - b^2) \sqrt{a + b \cos(c + dx)}}$$

input `Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2), x]`

output `(2*a*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + (((2*(a - b)*Sqrt[a + b]*(2*a*A*b - 3*a^2*B + b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(a*d) - (2*(a - b)*Sqrt[a + b]*(2*A*b - (3*a + b)*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/d - (2*Sqrt[a + b]*(a^2 - b^2)*(2*A*b - 3*a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(b*d))/(2*b) - ((2*a*A*b - 3*a^2*B + b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]])/(b*(a^2 - b^2))`

3.426. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$

3.426.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
 $\wedge(3/2)*\text{Sqrt}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]]), x_Symbol] \text{ :> Simp}[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*\text{Sqrt}[c*((1 + Csc[e + f*x])
)/(c - d)]*\text{Sqrt}[c*((1 - Csc[e + f*x])/(c + d))]*\text{EllipticE}[ArcSin[\text{Sqrt}[c +
d*\text{Sin}[e + f*x]]/\text{Sqrt}[b*\text{Sin}[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\&
\text{PosQ}[(c + d)/b]$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*
(x_)]) $\wedge(3/2)*\text{Sqrt}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]]), x_Symbol] \text{ :> S}$`
`imp[(A - B)/(a - b) Int[1/(\text{Sqrt}[a + b*\text{Sin}[e + f*x]]*\text{Sqrt}[c + d*\text{Sin}[e + f*
x]]), x], x] - \text{Simp}[(A*b - a*B)/(a - b) Int[(1 + \text{Sin}[e + f*x])/((a + b*\text{Sin}[e + f*x])
 $\wedge(3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x]])], x], x] /; \text{FreeQ}\{a, b, c, d, e$`
`, f, A, B\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2,`
`0] \&\& \text{NeQ}[A, B]`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)] \wedge
2)/(((a_) + (b_)*sin[(e_) + (f_)*(x_)]) $\wedge(3/2)*\text{Sqrt}[(c_) + (d_)*\sin[(e_)
+ (f_)*(x_)]]), x_Symbol] \text{ :> Simp}[C/b^2 Int[\text{Sqrt}[a + b*\text{Sin}[e + f*x]]
/\text{Sqrt}[c + d*\text{Sin}[e + f*x]], x], x] + \text{Simp}[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*\text{Sin}[e + f*x])/((a + b*\text{Sin}[e + f*x]) $\wedge(3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x]$
)], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\&
\&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$`

rule 3540 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)] \wedge
2)/(\text{Sqrt}[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*\text{Sqrt}[(c_) + (d_)*sin[(e_)
+ (f_)*(x_)]]), x_Symbol] \text{ :> Simp}[(-C)*\text{Cos}[e + f*x]*(\text{Sqrt}[c + d*\text{Sin}[e + f*
x]]/(d*f*\text{Sqrt}[a + b*\text{Sin}[e + f*x]]), x] + \text{Simp}[1/(2*d) Int[(1/((a + b*\text{Sin}[e + f*x])
 $\wedge(3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x]])]*\text{Simp}[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*\text{Sin}[e + f*x] + (2*b*B*d - C*(b*c + a*d))*\text{Sin}[e +
f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b*c - a$`
`*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]`

3.426.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 3561 vs. $2(468) = 936$.

Time = 16.51 (sec) , antiderivative size = 3562, normalized size of antiderivative = 7.12

method	result	size
default	Expression too large to display	3562
parts	Expression too large to display	3618

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/d*(2*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^
2+1))^(3/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^2*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^
2+1))^(1/2)*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1
-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF
(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b-(-csc(d*x+c)^2*(1-cos(d*x
+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*
x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(
1/2))*b^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos
(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot
(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2+(-csc(d*x+c)^2*(1-cos(d*x+c))
^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
)^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
)*a*b-2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d
*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticPi(cot(
d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*a^2+2*(-csc(d*x+c)^2*(1-cos(d*x
+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*
x+c))^2+a+b)/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b
))^(1/2))*b^2+csc(d*x+c)^3*a^2*(1-cos(d*x+c))^3-csc(d*x+c)^3*a*b*(1-cos(d*
x+c))^3-a^2*(csc(d*x+c)-cot(d*x+c))+a*b*(csc(d*x+c)-cot(d*x+c)))/(csc(d...
```

3.426.5 Fricas [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `integral((B*cos(d*x + c)^2 + A*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2), x)`

3.426.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.426.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(3/2), x)`

3.426.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(3/2), x)`

3.426.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

$$3.427 \quad \int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.427.1 Optimal result 3979
 3.427.2 Mathematica [C] (verified) 3980
 3.427.3 Rubi [A] (verified) 3981
 3.427.4 Maple [B] (warning: unable to verify) 3985
 3.427.5 Fricas [F] 3986
 3.427.6 Sympy [F] 3987
 3.427.7 Maxima [F] 3987
 3.427.8 Giac [F] 3987
 3.427.9 Mupad [F(-1)] 3988

3.427.1 Optimal result

Integrand size = 35, antiderivative size = 416

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx =$$

$$\frac{2(Ab - aB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ab\sqrt{a+bd}}$$

$$+ \frac{2(Ab - aB) \cot(c+dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ab\sqrt{a+bd}}$$

$$- \frac{2\sqrt{a+bd} B \cot(c+dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^2 d}$$

$$+ \frac{2a(Ab - aB) \sin(c+dx)}{b(a^2 - b^2) d \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}$$

```
output 2*a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(
1/2)-2*(A*b-B*a)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/c
os(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1
+sec(d*x+c))/(a-b))^(1/2)/a/b/d/(a+b)^(1/2)+2*(A*b-B*a)*cot(d*x+c)*Ellipti
cF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2
))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/b/d/(a+
b)^(1/2)-2*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(
d*x+c)^(1/2), (a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c)))/(
a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b^2/d
```

$$3.427. \quad \int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.427.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 16.32 (sec) , antiderivative size = 1012, normalized size of antiderivative = 2.43

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)}(-Ab\sin(c+dx)+aB\sin(c+dx))}{(a^2-b^2)d\sqrt{a+b\cos(c+dx)}}$$

$$-4a(aA-bB) \left(\frac{\sqrt{\frac{(a+b)\cot^2\left(\frac{1}{2}(c+dx)\right)}{-a+b}} \sqrt{-\frac{(a+b)\cos(c+dx)\csc^2\left(\frac{1}{2}(c+dx)\right)}{a}} \sqrt{\frac{(a+b\cos(c+dx))\csc^2\left(\frac{1}{2}(c+dx)\right)}{a}} \csc(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}}\right)\right)}{(a+b)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right)$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]`

output `(2*sqrt[Cos[c + d*x]]*(-(A*b*sin[c + d*x]) + a*B*sin[c + d*x]))/((a^2 - b^2)*d*sqrt[a + b*cos[c + d*x]]) - (-4*a*(a*A - b*B)*((sqrt[((a + b)*cot[(c + d*x)/2]^2)/(-a + b)]*sqrt[-(((a + b)*cos[c + d*x]*csc[(c + d*x)/2]^2)/a])*sqrt[((a + b*cos[c + d*x])*csc[(c + d*x)/2]^2)/a]*csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*cos[c + d*x])*csc[(c + d*x)/2]^2)/a]/sqrt[2]], (-2*a)/(-a + b)]*sin[(c + d*x)/2]^4)/((a + b)*sqrt[Cos[c + d*x]]*sqrt[a + b*cos[c + d*x]]) - (sqrt[((a + b)*cot[(c + d*x)/2]^2)/(-a + b)]*sqrt[-(((a + b)*cos[c + d*x]*csc[(c + d*x)/2]^2)/a])*sqrt[((a + b*cos[c + d*x])*csc[(c + d*x)/2]^2)/a]*csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*cos[c + d*x])*csc[(c + d*x)/2]^2)/a]/sqrt[2]], (-2*a)/(-a + b)]*sin[(c + d*x)/2]^4)/(b*sqrt[Cos[c + d*x]]*sqrt[a + b*cos[c + d*x]]) + 2*(A*b - a*B)*((I*cos[(c + d*x)/2]*sqrt[a + b*cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*sec[c + d*x])/(b*sqrt[Cos[(c + d*x)/2]^2*sec[c + d*x]]*sqrt[((a + b*cos[c + d*x])*sec[c + d*x])/(a + b)]) + (2*a*((a*sqrt[((a + b)*cot[(c + d*x)/2]^2)/(-a + b)]*sqrt[-(((a + b)*cos[c + d*x]*csc[(c + d*x)/2]^2)/a])*sqrt[((a + b*cos[c + d*x])*csc[(c + d*x)/2]^2)/a]*csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*cos[c + d*x])*csc[(c + d*x)/2]^2)/a]/sqrt[2]], (-2*a)/(-a + b)]*sin[(c + d*x)/2]^4)/((a + b)*sqrt[Cos[c + d*x]]*sqrt[a + b*cos[c + d*x]]) - (a*sqrt[((a + b)*cot[(c + d*x)/2]^2)/(-a + b)]*sqrt[-(((a + b)*cos[c + d*x]*csc[(c + d*x)/2]^2)/a])*sqrt...`

3.427.3 Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3471, 3042, 3273, 3042, 3274, 3042, 3288, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{3471} \\
 & \frac{(Ab-aB)}{b} \int \frac{\sqrt{\cos(c+dx)}}{(a+b\cos(c+dx))^{3/2}} dx + \frac{B}{b} \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx
 \end{aligned}$$

$$3.427. \quad \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \frac{(Ab - aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \downarrow \text{3273} \\
 & \frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{\sqrt{a+b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx}{a^2-b^2} \right)}{b} + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \downarrow \text{3042} \\
 & \frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx}{a^2-b^2} \right)}{b} + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \downarrow \text{3274} \\
 & \frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - \frac{a \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx - (a-b) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx}{a^2-b^2} \right)}{b} + \\
 & \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \downarrow \text{3042} \\
 & \frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - (a-b) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} \right)}{b} + \\
 & \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
 & \downarrow \text{3288}
 \end{aligned}$$

3.427. $\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

$$\frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{b^2d}$$

↓ 3295

$$(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{a^2-b^2} \right)$$

$$\frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{b^2d}$$

↓ 3473

$$(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{2(a-b)\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{ad} \right)$$

$$\frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{b^2d}$$

```
input Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x
]
```

3.427. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

```
output (-2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos
[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*
(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b^2*d)
+ ((A*b - a*B)*(-(((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[S
qrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a -
b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a -
b)]/(a*d) - (2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a
+ b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*S
qrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/
(a*d))/(a^2 - b^2)) + (2*a*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]
*Sqrt[a + b*Cos[c + d*x]])))/b
```

3.427.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3273 Int[Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]/((a_) + (b_)*sin[(e_) + (f_)*(x_
)])^(3/2), x_Symbol] := Simp[-2*a*d*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b
*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] - Simp[d^2/(a^2 - b^2) Int[Sqrt
[a + b*Sin[e + f*x]]/(d*Sin[e + f*x])^(3/2), x], x] /; FreeQ[{a, b, d, e, f
}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 3274 Int[Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]/((a_) + (b_)*sin[(e_) + (
f_)*(x_)])^(3/2), x_Symbol] := Simp[(c - d)/(a - b) Int[1/(Sqrt[a + b*Si
n[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(b*c - a*d)/(a - b)
Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]
), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_ + (b_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3471 Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)])]/((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(3/2), x_Symbol] :> Simp[B/b Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(A*b - a*B)/b Int[Sqrt[c + d*Sin[e + f*x]]/(a + b*Sin[e + f*x])^(3/2), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3473 Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(((b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] :> Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]
```

3.427.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1818 vs. $2(388) = 776$.

Time = 12.14 (sec) , antiderivative size = 1819, normalized size of antiderivative = 4.37

method	result	size
default	Expression too large to display	1819
parts	Expression too large to display	1850

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)
```



```
output 1/d*(2*A*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^
2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)*((csc(d*x+c)^2*a*(1-cos(d*x+
c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+
1))^(1/2)*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-c
os(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(c
ot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a-(-csc(d*x+c)^2*(1-cos(d*x+c))
^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
)^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
)*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c)
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c
)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1
/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)
/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+csc(
d*x+c)^3*(1-cos(d*x+c))^3*a-csc(d*x+c)^3*(1-cos(d*x+c))^3*b-a*(csc(d*x+c)-
cot(d*x+c))+b*(csc(d*x+c)-cot(d*x+c)))/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(
csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a-b)
/(a+b)+2*B*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1))^(3/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^2*((csc(d*x+c)^2*a*(1-cos(
d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1))^(1/2)*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2...
```

3.427.5 Fracas [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{3/2}} dx$$

```
input integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="fracas")
```

```
output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/
(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2), x)
```

3.427.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(A+B\cos(c+dx))\sqrt{\cos(c+dx)}}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(cos(c + d*x))/(a + b*cos(c + d*x))**(3/2), x)`

3.427.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)^(3/2), x)`

3.427.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)^(3/2), x)`

3.427.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

3.428 $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}^{3/2}} dx$

3.428.1 Optimal result	3989
3.428.2 Mathematica [C] (verified)	3990
3.428.3 Rubi [A] (verified)	3990
3.428.4 Maple [B] (warning: unable to verify)	3993
3.428.5 Fricas [F]	3994
3.428.6 Sympy [F]	3995
3.428.7 Maxima [F]	3995
3.428.8 Giac [F]	3995
3.428.9 Mupad [F(-1)]	3996

3.428.1 Optimal result

Integrand size = 35, antiderivative size = 284

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))}^{3/2}} dx = \frac{2(Ab - aB) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{a}}{a^2 \sqrt{a + bd}}$$

$$+ \frac{2(A + B) \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{a \sqrt{a + bd}}$$

$$- \frac{2(Ab - aB) \sin(c + dx)}{(a^2 - b^2) d \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}$$

output

```
-2*(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)
)+2*(A*b-B*a)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(
d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+se
c(d*x+c))/(a-b))^(1/2)/a^2/d/(a+b)^(1/2)+2*(A+B)*cot(d*x+c)*EllipticF((a+b
*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(
1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/(a+b)^(1/2)
```

3.428.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.67 (sec) , antiderivative size = 1223, normalized size of antiderivative = 4.31

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)),x]`

output `(-2*Sqrt[Cos[c + d*x]]*(-(A*b^2*Sin[c + d*x]) + a*b*B*Sin[c + d*x]))/(a*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + ((-4*a*(a^2*A - A*b^2)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-(a*A*b) + a^2*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-(A*b^2) + a*b*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*C...`

3.428.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.428. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
& \quad \downarrow \text{3472} \\
& \frac{\int \frac{Ab - aB + (aA - bB) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{Ab - aB + (aA - bB) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3477} \\
& \frac{(Ab - aB) \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + (a - b)(A + B) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(a - b)(A + B) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + (Ab - aB) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3295} \\
& \frac{(Ab - aB) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{2(a - b) \sqrt{a + b} (A + B) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} \text{EllipticF}\left(\arcsin\right)}{ad}}{a^2 - b^2} - \frac{2(Ab - aB) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3473}
\end{aligned}$$

3.428. $\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$

$$\frac{2(a-b)\sqrt{a+b}(Ab-aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\middle|-\frac{a+b}{a-b}\right)}{a^2d} + \frac{2(a-b)\sqrt{a+b}(A+B)\cot(c+dx)\sqrt{a+b}}{a^2-b^2}$$

$$\frac{2(Ab-aB)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)),x]`

output `((2*(a - b)*Sqrt[a + b]*(A*b - a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(A + B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/((a*d))/(a^2 - b^2) - (2*(A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])`

3.428.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]]), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_
.)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

3.428.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1532 vs. $2(264) = 528$.

Time = 13.70 (sec) , antiderivative size = 1533, normalized size of antiderivative = 5.40

method	result	size
default	Expression too large to display	1533
parts	Expression too large to display	1537

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RET
URNVERBOSE)
```


output `1/d*(2*A*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^2+csc(d*x+c)^3*a*b*(1-cos(d*x+c))^3-csc(d*x+c)^3*b^2*(1-cos(d*x+c))^3-a*b*(csc(d*x+c)-cot(d*x+c))+b^2*(csc(d*x+c)-cot(d*x+c)))*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)/(a+b)/(a-b)/(csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/a/(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)+2*B*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+...`

3.428.5 Fracas [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{3/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(b^2*cos(d*x + c)^3 + 2*a*b*cos(d*x + c)^2 + a^2*cos(d*x + c)), x)`

3.428.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}} \sqrt{\cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))/((a + b*cos(c + d*x))**(3/2)*sqrt(cos(c + d*x))), x)`

3.428.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*sqrt(cos(d*x + c))), x)`

3.428.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*sqrt(cos(d*x + c))), x)`

3.428.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(3/2)),x)`

$$3.429 \quad \int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$$

3.429.1 Optimal result	3997
3.429.2 Mathematica [C] (verified)	3998
3.429.3 Rubi [A] (verified)	3998
3.429.4 Maple [B] (warning: unable to verify)	4002
3.429.5 Fracas [F]	4003
3.429.6 Sympy [F]	4003
3.429.7 Maxima [F]	4003
3.429.8 Giac [F]	4004
3.429.9 Mupad [F(-1)]	4004

3.429.1 Optimal result

Integrand size = 35, antiderivative size = 305

$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx = \frac{2(a^2 A - 2Ab^2 + abB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right) - 2(2Ab + a(A - B)) \cot(c+dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{a^3 \sqrt{a+bd}} + \frac{2b(Ab - aB) \sin(c+dx)}{a(a^2 - b^2) d \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}$$

output

```
2*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)+2*(A*a^2-2*A*b^2+B*a*b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d/(a+b)^(1/2)-2*(2*A*b+a*(A-B))*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d/(a+b)^(1/2)
```

3.429.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.66 (sec) , antiderivative size = 1281, normalized size of antiderivative = 4.20

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)),x]`

output `((-4*a*(2*a^2*A*b - 2*A*b^3 - a^3*B + a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(a^3*A - 2*a*A*b^2 + a^2*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(a^2*A*b - 2*A*b^3 + a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])`

3.429.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 331, normalized size of antiderivative = 1.09, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3479, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.429. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} (a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
& \quad \downarrow \text{3479} \\
& \frac{2 \int \frac{Aa^2 + bBa - (Ab - aB) \cos(c + dx)a - 2Ab^2}{2 \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{Aa^2 + bBa - (Ab - aB) \cos(c + dx)a - 2Ab^2}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{Aa^2 + bBa - (Ab - aB) \sin(c + dx + \frac{\pi}{2})a - 2Ab^2}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3477} \\
& \frac{(a^2A + abB - 2Ab^2) \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - (a - b)(a(A - B) + 2Ab) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx}{a(a^2 - b^2)} + \\
& \quad \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(a^2A + abB - 2Ab^2) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - (a - b)(a(A - B) + 2Ab) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a(a^2 - b^2)} + \\
& \quad \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3295}
\end{aligned}$$

3.429. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{(a^2A + abB - 2Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(a(A-B)+2Ab) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{ad}}{a(a^2 - b^2)} \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3473} \\
& \frac{2(a-b)\sqrt{a+b}(a^2A+abB-2Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) - \frac{2(a-b)\sqrt{a+b}(a(A-B)+2Ab)}{ad}}{a^2d} - \frac{2(a-b)\sqrt{a+b}(a(A-B)+2Ab)}{a(a^2 - b^2)} \\
& \frac{2b(Ab - aB) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)),x]`

output `((2*(a - b)*Sqrt[a + b]*(a^2*A - 2*A*b^2 + a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(2*A*b + a*(A - B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a*(a^2 - b^2)) + (2*b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])`

3.429.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

3.429.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1977 vs. $2(285) = 570$.

Time = 18.71 (sec) , antiderivative size = 1978, normalized size of antiderivative = 6.49

method	result	size
default	Expression too large to display	1978
parts	Expression too large to display	1980

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/d*(-2*A*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(-csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
))^2+a+b)/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2
))*a^3+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b)^(1/2)*EllipticF(cot(d*
x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b+2*(-csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
))^2+a+b)/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2
))*a*b^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(
d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b)^(1/2)*EllipticE(cot(
d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3+(-csc(d*x+c)^2*(1-cos(d*x+c))^
2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
))^2+a+b)/(a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
)*a^2*b-2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(
d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b)^(1/2)*EllipticE(cot(
d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^2-2*(-csc(d*x+c)^2*(1-cos(d*x+
c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x
+c))^2+a+b)/(a+b)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1
/2))*b^3+csc(d*x+c)^3*a^3*(1-cos(d*x+c))^3-csc(d*x+c)^3*a^2*b*(1-cos(d*x+c)
))^3-2*csc(d*x+c)^3*a*b^2*(1-cos(d*x+c))^3+2*csc(d*x+c)^3*b^3*(1-cos(d...
```

3.429.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/
(b^2*cos(d*x + c)^4 + 2*a*b*cos(d*x + c)^3 + a^2*cos(d*x + c)^2), x)`

3.429.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))/((a + b*cos(c + d*x))**(3/2)*cos(c + d*x)**(
3/2)), x)`

3.429.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3
/2)), x)`

3.429.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algo rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(3/2)), x)`

3.429.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(3/2)),x)`

3.430
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$$

3.430.1 Optimal result	4005
3.430.2 Mathematica [C] (verified)	4006
3.430.3 Rubi [A] (verified)	4006
3.430.4 Maple [B] (warning: unable to verify)	4011
3.430.5 Fricas [F]	4012
3.430.6 Sympy [F(-1)]	4012
3.430.7 Maxima [F]	4012
3.430.8 Giac [F]	4013
3.430.9 Mupad [F(-1)]	4013

3.430.1 Optimal result

Integrand size = 35, antiderivative size = 393

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx =$$

$$\frac{2(5a^2 Ab - 8Ab^3 - 3a^3 B + 6ab^2 B) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a^4 \sqrt{a + bd}}$$

$$+ \frac{2(a + 2b)(4Ab + a(A - 3B)) \cot(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a^3 \sqrt{a + bd}}$$

$$+ \frac{2b(Ab - aB) \sin(c + dx)}{a(a^2 - b^2) d \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{2(a^2 A - 4Ab^2 + 3abB) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3a^2 (a^2 - b^2) d \cos^{\frac{3}{2}}(c + dx)}$$

output

```
2*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(
1/2)+2/3*(A*a^2-4*A*b^2+3*B*a*b)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a^2/(a^
2-b^2)/d/cos(d*x+c)^(3/2)-2/3*(5*A*a^2*b-8*A*b^3-3*B*a^3+6*B*a*b^2)*cot(d*
x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)
/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/
2)/a^4/d/(a+b)^(1/2)+2/3*(a+2*b)*(4*A*b+a*(A-3*B))*cot(d*x+c)*EllipticF((a
+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a
*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d/(a+b)^(1
/2)
```

3.430.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$$

3.430.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.78 (sec) , antiderivative size = 1357, normalized size of antiderivative = 3.45

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^(3/2)),x]`

output `((-4*a*(a^4*A + 7*a^2*A*b^2 - 8*A*b^4 - 6*a^3*b*B + 6*a*b^3*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(5*a^3*A*b - 8*a*A*b^3 - 3*a^4*B + 6*a^2*b^2*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(5*a^2*A*b^2 - 8*A*b^4 - 3*a^3*b*B + 6*a*b^3*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x]/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[(a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[...`

3.430.3 Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.430. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{3/2}} dx \\
& \quad \downarrow \text{3479} \\
& \frac{2 \int \frac{Aa^2 + 3bBa - (Ab - aB) \cos(c + dx)a - 4Ab^2 + 2b(Ab - aB) \cos^2(c + dx)}{2 \cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{\frac{a(a^2 - b^2)}{2b(Ab - aB) \sin(c + dx)}} + \\
& \quad \frac{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{Aa^2 + 3bBa - (Ab - aB) \cos(c + dx)a - 4Ab^2 + 2b(Ab - aB) \cos^2(c + dx)}{\cos^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{\frac{a(a^2 - b^2)}{2b(Ab - aB) \sin(c + dx)}} + \\
& \quad \frac{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{Aa^2 + 3bBa - (Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)a - 4Ab^2 + 2b(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)^2}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{\frac{a(a^2 - b^2)}{2b(Ab - aB) \sin(c + dx)}} + \\
& \quad \frac{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{3534} \\
& \frac{2 \int -\frac{-3Ba^3 + 5Aba^2 + 6b^2Ba - (Aa^2 - 3bBa + 2Ab^2) \cos(c + dx)a - 8Ab^3}{2 \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{\frac{3a}{3ad \cos^{\frac{3}{2}}(c + dx)}} + \frac{2(a^2A + 3abB - 4Ab^2) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3ad \cos^{\frac{3}{2}}(c + dx)} + \\
& \quad \frac{a(a^2 - b^2)}{2b(Ab - aB) \sin(c + dx)} \\
& \quad \frac{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{\frac{2(a^2A + 3abB - 4Ab^2) \sin(c + dx) \sqrt{a + b \cos(c + dx)}}{3ad \cos^{\frac{3}{2}}(c + dx)} - \int \frac{-3Ba^3 + 5Aba^2 + 6b^2Ba - (Aa^2 - 3bBa + 2Ab^2) \cos(c + dx)a - 8Ab^3}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{\frac{3a}{3ad \cos^{\frac{3}{2}}(c + dx)}} + \\
& \quad \frac{a(a^2 - b^2)}{2b(Ab - aB) \sin(c + dx)} \\
& \quad \frac{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}{ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}
\end{aligned}$$

3.430. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
 & \int \frac{-3Ba^3+5Aba^2+6b^2Ba-(Aa^2-3bBa+2Ab^2)\sin(c+dx+\frac{\pi}{2})a-8Ab^3}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{a(a^2-b^2)}{3a} + \\
 & \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \\
 & \downarrow 3477 \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-(a-b)(a+2b)(aA-3aB+4a^2)}{3a} \\
 & \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} + \frac{a(a^2-b^2)}{3a} \\
 & \downarrow 3042 \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-(a-b)(a+2b)(aA-3aB+4a^2)}{3a} \\
 & \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} + \frac{a(a^2-b^2)}{3a} \\
 & \downarrow 3042 \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-(a-b)(a+2b)(aA-3aB+4a^2)}{3a} \\
 & \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} + \frac{a(a^2-b^2)}{3a} \\
 & \downarrow 3295 \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2(a-b)\sqrt{a+b}(a+2b)}{a^2d}}{3a} \\
 & \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} + \frac{a(a^2-b^2)}{3a} \\
 & \downarrow 3473 \\
 & \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} + \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{2(a-b)\sqrt{a+b}(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E(\arcsin\frac{\sqrt{a+b}\sin(c+dx+\frac{\pi}{2})}{a})}{a^2d} \\
 & \frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{2(a-b)\sqrt{a+b}(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E(\arcsin\frac{\sqrt{a+b}\sin(c+dx+\frac{\pi}{2})}{a})}{a^2d}
 \end{aligned}$$

3.430. $\int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} dx$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^(3/2)),x]`

output `(2*b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]) + (-1/3*((2*(a - b)*Sqrt[a + b]*(5*a^2*A*b - 8*A*b^3 - 3*a^3*B + 6*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(a + 2*b)*(a*A + 4*A*b - 3*a*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d))/a + (2*(a^2*A - 4*A*b^2 + 3*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/(a*(a^2 - b^2))`

3.430.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_)] + (f_)*(x_)]], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3473 `Int[((A_) + (B_)*sin[(e_)] + (f_)*(x_)]/(((b_)*sin[(e_)] + (f_)*(x_))]^(3/2)*Sqrt[(c_) + (d_)*sin[(e_)] + (f_)*(x_)]], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

3.430.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 3845 vs. 2(363) = 726.

Time = 22.20 (sec) , antiderivative size = 3846, normalized size of antiderivative = 9.79

method	result	size
default	Expression too large to display	3846
parts	Expression too large to display	3848

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/d*(2/3*A*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(5*csc(d*x+c)^2*EllipticF(cot
(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*b*(-csc(d*x+c)^2*(1-cos(d*x+c)
))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+
c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x+c))^2-5*EllipticF(cot(d*x+c)-csc(d*x+c)
,(-a-b)/(a+b))^(1/2))*a^3*b*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((cs
c(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(
1/2)+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b^2*(-cs
c(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc
(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)+8*EllipticF(cot(d*x+c)-csc(
d*x+c),(-a-b)/(a+b))^(1/2))*a*b^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2
)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(
a+b))^(1/2)+5*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*b*
(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2
-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)+5*EllipticE(cot(d*x+c)-
csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b^2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1
)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+
a+b)/(a+b))^(1/2)-8*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*
a*b^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(dx
+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)-csc(d*x+c)^2*Elli
pticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^4*(-csc(d*x+c)^2*(1...
```

3.430.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/
(b^2*cos(d*x + c)^5 + 2*a*b*cos(d*x + c)^4 + a^2*cos(d*x + c)^3), x)`

3.430.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.430.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(5
/2)), x)`

3.430.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(3/2),x, algo rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)^(5/2)), x)`

3.430.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^(3/2)),x)`

3.431 $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$

3.431.1 Optimal result 4014
 3.431.2 Mathematica [C] (warning: unable to verify) 4015
 3.431.3 Rubi [A] (verified) 4016
 3.431.4 Maple [B] (warning: unable to verify) 4022
 3.431.5 Fricas [F(-1)] 4023
 3.431.6 Sympy [F(-1)] 4023
 3.431.7 Maxima [F] 4023
 3.431.8 Giac [F] 4024
 3.431.9 Mupad [F(-1)] 4024

3.431.1 Optimal result

Integrand size = 35, antiderivative size = 674

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx = \frac{(6a^3Ab - 14aAb^3 - 15a^4B + 26a^2b^2B - 3b^4B) \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right)\right) + (6a^2Ab + 2aAb^2 - 12Ab^3 - 15a^3B - 5a^2bB + 21ab^2B + 3b^3B) \cot(c+dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right)\right) - \frac{3(a-b)b^3(a+b)^{\frac{3}{2}}d}{\sqrt{a+b}(2Ab - 5aB) \cot(c+dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^4d} + \frac{2a(Ab - aB) \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{3b(a^2 - b^2) d(a+b \cos(c+dx))^{\frac{3}{2}}} + \frac{2a(2a^2Ab - 6Ab^3 - 5a^3B + 9ab^2B) \sqrt{\cos(c+dx)} \sin(c+dx)}{3b^2(a^2 - b^2)^2 d \sqrt{a+b \cos(c+dx)}} - \frac{(6a^3Ab - 14aAb^3 - 15a^4B + 26a^2b^2B - 3b^4B) \sqrt{a+b \cos(c+dx)} \sin(c+dx)}{3b^3(a^2 - b^2)^2 d \sqrt{\cos(c+dx)}}$$

3.431. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$

output $\frac{2}{3}a(Ab-Ba)\cos(dx+c)^{3/2}\sin(dx+c)/b(a^2-b^2)/d/(a+b\cos(dx+c))^{3/2} + \frac{2}{3}a(2Aa^2b-6Ab^3-5Ba^3+9Bab^2)\sin(dx+c)\cos(dx+c)^{1/2}/b^2(a^2-b^2)^2/d/(a+b\cos(dx+c))^{1/2} - \frac{1}{3}(6Aa^3b-14Aab^3-15Ba^4+26Ba^2b^2-3Bb^4)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b^3(a^2-b^2)^2/d/\cos(dx+c)^{1/2} + \frac{1}{3}(6Aa^3b-14Aab^3-15Ba^4+26Ba^2b^2-3Bb^4)\cot(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2}) * (a(1-\sec(dx+c))/(a+b))^{1/2} * (a(1+\sec(dx+c))/(a-b))^{1/2} / a(a-b)/b^3(a+b)^{3/2}/d - \frac{1}{3}(6Aa^2b+2Aab^2-12Ab^3-15Ba^3-5Ba^2b+21Bab^2+3Bb^3)\cot(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2}) * (a(1-\sec(dx+c))/(a+b))^{1/2} * (a(1+\sec(dx+c))/(a-b))^{1/2} / (a-b)/b^3(a+b)^{3/2}/d - (2Ab-5Ba)\cot(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b, ((-a-b)/(a-b))^{1/2}) * (a+b)^{1/2} * (a(1-\sec(dx+c))/(a+b))^{1/2} * (a(1+\sec(dx+c))/(a-b))^{1/2} / b^4/d$

3.431.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.19 (sec) , antiderivative size = 1396, normalized size of antiderivative = 2.07

$$\int \frac{\cos^{5/2}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output

```
(Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*((-2*(-a^2*A*b*Sin[c + d*x])
+ a^3*B*Sin[c + d*x]))/(3*b^2*(-a^2 + b^2)*(a + b*Cos[c + d*x])^2) - (2*(
-3*a^3*A*b*Sin[c + d*x] + 7*a*A*b^3*Sin[c + d*x] + 6*a^4*B*Sin[c + d*x] -
10*a^2*b^2*B*Sin[c + d*x]))/(3*b^2*(-a^2 + b^2)^2*(a + b*Cos[c + d*x])))/
d + ((-4*a*(-2*a^3*A*b + 2*a*A*b^3 + 5*a^4*B - 8*a^2*b^2*B + 3*b^4*B)*Sqrt
[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(
c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c
+ d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]
/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]
]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(2*a^2*A*b^2 + 6*A*b^4 + 4*a^3*b*B - 12*
a*b^3*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos
[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)
/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c
+ d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sq
rt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/
2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[
((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b
), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*
a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c +
d*x]])) + 2*(-6*a^3*A*b + 14*a*A*b^3 + 15*a^4*B - 26*a^2*b^2*B + 3*b^4*...
```

3.431.3 Rubi [A] (verified)

Time = 3.43 (sec) , antiderivative size = 698, normalized size of antiderivative = 1.04, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3468, 27, 3042, 3526, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}}(A+B \sin(c+dx+\frac{\pi}{2}))}{(a+b \sin(c+dx+\frac{\pi}{2}))^{\frac{5}{2}}} dx$$

↓ 3468

3.431. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$

$$\begin{aligned}
 & \frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} - \\
 & \frac{2 \int -\frac{\sqrt{\cos(c+dx)}(-((-5Ba^2+2Aba+3b^2B) \cos^2(c+dx))-3b(Ab-aB) \cos(c+dx)+3a(Ab-aB))}{2(a+b \cos(c+dx))^{3/2}} dx}{3b(a^2 - b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{\cos(c+dx)}(-((-5Ba^2+2Aba+3b^2B) \cos^2(c+dx))-3b(Ab-aB) \cos(c+dx)+3a(Ab-aB))}{(a+b \cos(c+dx))^{3/2}} dx}{3b(a^2 - b^2)} + \\
 & \frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}((5Ba^2-2Aba-3b^2B) \sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB) \sin(c+dx+\frac{\pi}{2}))+3a(Ab-aB)}{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3b(a^2 - b^2)} + \\
 & \frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3526} \\
 & \frac{2a(-5a^3B+2a^2Ab+9ab^2B-6Ab^3) \sin(c+dx) \sqrt{\cos(c+dx)}}{bd(a^2 - b^2) \sqrt{a+b \cos(c+dx)}} - \frac{2 \int -\frac{((-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B) \cos^2(c+dx))+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3) \cos(c+dx)+a(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)}{2\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx}{b(a^2 - b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{-((-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B) \cos^2(c+dx))+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3) \cos(c+dx)+a(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx}{b(a^2 - b^2)} + \frac{2a(-5a^3)}{b(a^2 - b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(15Ba^4-6Aba^3-26b^2Ba^2+14Ab^3a+3b^4B) \sin(c+dx+\frac{\pi}{2})^2+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3) \sin(c+dx+\frac{\pi}{2}))+a(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b(a^2 - b^2)} + \frac{2a(-5a^3)}{b(a^2 - b^2)} \\
 & \frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}
 \end{aligned}$$

3.431. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

↓ 3540

$$\frac{\int \frac{3(a^2-b^2)^2(2Ab-5aB)\cos^2(c+dx)+2ab(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)\cos(c+dx)+a(-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)dx}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{2b} - \frac{(-15a^4B+6a^3Ab+26a^2bB-14ab^3-3b^4)}{b(a^2-b^2)}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{\int \frac{3(a^2-b^2)^2(2Ab-5aB)\sin(c+dx+\frac{\pi}{2})^2+2ab(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)\sin(c+dx+\frac{\pi}{2})+a(-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)dx}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{2b} - \frac{(-15a^4B+6a^3Ab+26a^2bB-14ab^3-3b^4)}{b(a^2-b^2)}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3532

$$\frac{3(a^2-b^2)^2(2Ab-5aB)\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}}dx + \int \frac{a(-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)+2ab(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{2b} - \frac{(-15a^4B+6a^3Ab+26a^2bB-14ab^3-3b^4)}{b(a^2-b^2)}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{3(a^2-b^2)^2(2Ab-5aB)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx + \int \frac{a(-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)+2ab(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{2b} - \frac{(-15a^4B+6a^3Ab+26a^2bB-14ab^3-3b^4)}{b(a^2-b^2)}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3288

3.431. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\int \frac{a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{6\sqrt{a+b}(a^2 - b^2)^2(2Ab - 5aB) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a+b}}}{2b} \sqrt{\frac{a(1 - \sec(c + dx))}{a+b}}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3477

$$a(a-b)(15a^3B - a^2(6Ab - 5bB) - ab^2(2A + 21B) + 3b^3(4A - B)) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx + a(-15a^4B + 6a^3Ab + 26a^2b^2B - 14aAb^3 - 3b^4B) \int \frac{\cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3042

$$a(a-b)(15a^3B - a^2(6Ab - 5bB) - ab^2(2A + 21B) + 3b^3(4A - B)) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + a(-15a^4B + 6a^3Ab + 26a^2b^2B - 14aAb^3 - 3b^4B) \int \frac{1}{\sin(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3295

$$a(-15a^4B + 6a^3Ab + 26a^2b^2B - 14aAb^3 - 3b^4B) \int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{6\sqrt{a+b}(a^2 - b^2)^2(2Ab - 5aB) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a+b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a+b}}}{bd}$$

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}$$

↓ 3473

$$\frac{2a(Ab - aB) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} +$$

$$\frac{2a(-5a^3B + 2a^2Ab + 9ab^2B - 6Ab^3) \sin(c + dx) \sqrt{\cos(c + dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c + dx)}} + \frac{6\sqrt{a+b}(a^2 - b^2)^2(2Ab - 5aB) \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a+b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a+b}} \text{EllipticPi}\left(\frac{a}{a-b}\right)}{bd}$$

3.431. $\int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx$

input `Int[(Cos[c + d*x]^(5/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + ((2*a*(2*a^2*A*b - 6*A*b^3 - 5*a^3*B + 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + (((2*(a - b)*Sqrt[a + b]*(6*a^3*A*b - 14*a*A*b^3 - 15*a^4*B + 26*a^2*b^2*B - 3*b^4*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*(a - b)*Sqrt[a + b]*(3*b^3*(4*A - B) + 15*a^3*B - a*b^2*(2*A + 21*B) - a^2*(6*A*b - 5*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (6*Sqrt[a + b]*(a^2 - b^2)^2*(2*A*b - 5*a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/(2*b) - ((6*a^3*A*b - 14*a*A*b^3 - 15*a^4*B + 26*a^2*b^2*B - 3*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/(b*(a^2 - b^2)))/(3*b*(a^2 - b^2))`

3.431.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

3.431.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$$

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_.)]]*Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_.)])/(((b_.)*sin[(e_.) + (f_.)*(x_.)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] :> Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x] *(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])]*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.431.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 10416 vs. $2(628) = 1256$.

Time = 18.25 (sec) , antiderivative size = 10417, normalized size of antiderivative = 15.46

method	result	size
parts	Expression too large to display	10417
default	Expression too large to display	11449

3.431.
$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$$

input `int(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

3.431.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x,algorithm="fricas")`

output `Timed out`

3.431.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cos(d*x+c)**(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.431.7 Maxima [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{5}{2}}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x,algorithm="maxima")`

3.431. $\int \frac{\cos^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{5}{2}}} dx$

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^(5/2), x)`

3.431.8 Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \cos(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(5/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^(5/2), x)`

3.431.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^{5/2} (A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx$$

input `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^(5/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

3.432
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$$

3.432.1 Optimal result 4025
 3.432.2 Mathematica [C] (warning: unable to verify) 4026
 3.432.3 Rubi [A] (verified) 4026
 3.432.4 Maple [B] (warning: unable to verify) 4032
 3.432.5 Fricas [F(-1)] 4032
 3.432.6 Sympy [F(-1)] 4032
 3.432.7 Maxima [F] 4033
 3.432.8 Giac [F] 4033
 3.432.9 Mupad [F(-1)] 4033

3.432.1 Optimal result

Integrand size = 35, antiderivative size = 545

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx = \frac{2(4Ab^3 + 3a^3B - 7ab^2B) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{3a(a-b)b^2(a+b)^{3/2}d}$$

$$+ \frac{2(aAb^2 - 3Ab^3 - 3a^3B - a^2bB + 6ab^2B) \cot(c+dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a(a-b)b^2(a+b)^{3/2}d}$$

$$- \frac{2\sqrt{a+b}B \cot(c+dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^3d}$$

$$+ \frac{2a(Ab - aB) \sqrt{\cos(c+dx)} \sin(c+dx)}{3b(a^2 - b^2)d(a+b \cos(c+dx))^{3/2}} - \frac{2a(4Ab^3 + 3a^3B - 7ab^2B) \sin(c+dx)}{3b^2(a^2 - b^2)^2 d \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}$$

output

```
2/3*a*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/b/(a^2-b^2)/d/(a+b*cos(d*x+c))
^(3/2)-2/3*a*(4*A*b^3+3*B*a^3-7*B*a*b^2)*sin(d*x+c)/b^2/(a^2-b^2)^2/d/cos(
d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)+2/3*(4*A*b^3+3*B*a^3-7*B*a*b^2)*cot(d*
x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)
/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/
2)/a/(a-b)/b^2/(a+b)^(3/2)/d+2/3*(A*a*b^2-3*A*b^3-3*B*a^3-B*a^2*b+6*B*a*b^
2)*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2
),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(
a-b))^(1/2)/a/(a-b)/b^2/(a+b)^(3/2)/d-2*B*cot(d*x+c)*EllipticPi((a+b*cos(d
*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a
+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b^
3/d
```

3.432.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$$

3.432.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.95 (sec) , antiderivative size = 1342, normalized size of antiderivative = 2.46

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*((2*(-(a*A*b*Sin[c + d*x]) + a^2*B*Sin[c + d*x]))/(3*b*(-a^2 + b^2)*(a + b*Cos[c + d*x])^2) + (2*(4*A*b^3*Sin[c + d*x] + 3*a^3*B*Sin[c + d*x] - 7*a*b^2*B*Sin[c + d*x]))/(3*b*(-a^2 + b^2)^2*(a + b*Cos[c + d*x])))/d - ((-4*a*(-(a^2*A*b) + A*b^3 + a^3*B - a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(4*a*A*b^2 - a^2*b*B - 3*b^3*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])) + 2*(4*A*b^3 + 3*a^3*B - 7*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]]...`

3.432.3 Rubi [A] (verified)

Time = 2.39 (sec) , antiderivative size = 569, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3468, 27, 3042, 3530, 3042, 3288, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.432. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{3468} \\
& \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} - \frac{2\int -\frac{3(a^2-b^2)B\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+a(Ab-aB)}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{3(a^2-b^2)B\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+a(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3(a^2-b^2)B\sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+a(Ab-aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3530} \\
& \frac{\int \frac{ab(Ab-aB)-3(Ab^3+a(a^2-2b^2)B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{b} + \frac{3B(a^2-b^2)\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx}{b} + \\
& \quad \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{ab(Ab-aB)-3(Ab^3+a(a^2-2b^2)B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} + \frac{3B(a^2-b^2)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} + \\
& \quad \frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3288}
\end{aligned}$$

3.432. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\int \frac{ab(Ab-aB)-3(Ab^3+a(a^2-2b^2)B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))}^{3/2}} dx - \frac{6B\sqrt{a+b}(a^2-b^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticPi}\left(\frac{a+b}{b},\arcsin\left(\frac{\sqrt{a(1-\sec(c+dx))}}{\sqrt{a+b}}\right)\right)}{b^2d}$$

$$\frac{3b(a^2-b^2)}{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}} - \frac{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3472

$$\int \frac{a(3Ba^3-7b^2Ba+4Ab^3)+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - \frac{2a(3a^3B-7ab^2B+4Ab^3)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{6B\sqrt{a+b}(a^2-b^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{b}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\int \frac{a(3Ba^3-7b^2Ba+4Ab^3)+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(3a^3B-7ab^2B+4Ab^3)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{6B\sqrt{a+b}(a^2-b^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{b}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3477

$$a(3a^3B-7ab^2B+4Ab^3) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b)(3a^3B+a^2bB-ab^2(A+6B)+3Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx - \frac{2a(3a^3B-7ab^2B+4Ab^3)}{d(a^2-b^2)\sqrt{\cos(c+dx)}}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$a(3a^3B-7ab^2B+4Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(3a^3B+a^2bB-ab^2(A+6B)+3Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(3a^3B-7ab^2B+4Ab^3)}{d(a^2-b^2)\sqrt{\sin(c+dx+\frac{\pi}{2})}}$$

$$\frac{2a(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3295

3.432. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{a(3a^3B - 7ab^2B + 4Ab^3) \int \frac{\sin(c+dx + \frac{\pi}{2}) + 1}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(3a^3B + a^2bB - ab^2(A+6B) + 3Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx))}{a-b}}}{a^2 - b^2}}{b} \\
 & \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3473} \\
 & \frac{2a(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3bd(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} + \\
 & \frac{2(a-b)\sqrt{a+b}(3a^3B - 7ab^2B + 4Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) - \frac{2(a-b)\sqrt{a+b}(3a^3B + a^2bB - ab^2(A+6B) + 3Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx))}{a-b}}}{a^2 - b^2}}{b}
 \end{aligned}$$

```
input Int[(Cos[c + d*x]^(3/2)*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x
]
```

```
output (2*a*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a +
b*Cos[c + d*x])^(3/2)) + ((-6*Sqrt[a + b]*(a^2 - b^2)*B*Cot[c + d*x]*Ellip
ticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c +
d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a
*(1 + Sec[c + d*x]))/(a - b)]/(b^2*d) + (((2*(a - b)*Sqrt[a + b]*(4*A*b^3
+ 3*a^3*B - 7*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d
*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - S
ec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d) - (2*(a
- b)*Sqrt[a + b]*(3*A*b^3 + 3*a^3*B + a^2*b*B - a*b^2*(A + 6*B))*Cot[c +
d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d
*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(
1 + Sec[c + d*x]))/(a - b)]/(a*d))/(a^2 - b^2) - (2*a*(4*A*b^3 + 3*a^3*B
- 7*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Co
s[c + d*x]]))/b)/(3*b*(a^2 - b^2))
```

3.432. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$

3.432.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3468 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

rule 3473 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3530 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[C/(b*d) Int[Sqrt[d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[1/b Int[(A*b + (b*B - a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]`

3.432.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 6573 vs. $2(505) = 1010$.

Time = 16.56 (sec) , antiderivative size = 6574, normalized size of antiderivative = 12.06

method	result	size
default	Expression too large to display	6574
parts	Expression too large to display	6730

```
input int(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)
```

```
output result too large to display
```

3.432.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output Timed out
```

3.432.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate(cos(d*x+c)**(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)
```

```
output Timed out
```

3.432. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

3.432.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(5/
2), x)`

3.432.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(5/
2), x)`

3.432.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(c+dx)^{3/2}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x
)`

output `int((cos(c + d*x)^(3/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),
x)`

3.432. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

3.433
$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx$$

3.433.1 Optimal result	4034
3.433.2 Mathematica [C] (verified)	4035
3.433.3 Rubi [A] (verified)	4035
3.433.4 Maple [B] (warning: unable to verify)	4039
3.433.5 Fracas [F]	4040
3.433.6 Sympy [F]	4041
3.433.7 Maxima [F]	4041
3.433.8 Giac [F]	4041
3.433.9 Mupad [F(-1)]	4042

3.433.1 Optimal result

Integrand size = 35, antiderivative size = 391

$$\int \frac{\sqrt{\cos(c+dx)}(A+B \cos(c+dx))}{(a+b \cos(c+dx))^{5/2}} dx =$$

$$\frac{2(3a^2 A + Ab^2 - 4abB) \cot(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a^2(a-b)(a+b)^{3/2}d}$$

$$+ \frac{2(3aA - Ab + aB - 3bB) \cot(c+dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{3a(a-b)(a+b)^{3/2}d}$$

$$- \frac{2(Ab - aB) \sqrt{\cos(c+dx)} \sin(c+dx)}{3(a^2 - b^2) d (a+b \cos(c+dx))^{3/2}} + \frac{2(3a^2 A + Ab^2 - 4abB) \sin(c+dx)}{3(a^2 - b^2)^2 d \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}$$

output

```
-2/3*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)+2/3*(3*A*a^2+A*b^2-4*B*a*b)*sin(d*x+c)/(a^2-b^2)^2/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)-2/3*(3*A*a^2+A*b^2-4*B*a*b)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/(a-b)/(a+b)^(3/2)/d+2/3*(3*A*a-A*b+B*a-3*B*b)*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/(a-b)/(a+b)^(3/2)/d
```

3.433.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.91 (sec) , antiderivative size = 1335, normalized size of antiderivative = 3.41

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*((2*(-(A*b*Sin[c + d*x])) + A*B*Sin[c + d*x]))/(3*(a^2 - b^2)*(a + b*Cos[c + d*x])^2) - (2*(3*a^2*A*b*Sin[c + d*x] + A*b^3*Sin[c + d*x] - 4*a*b^2*B*Sin[c + d*x]))/(3*a*(a^2 - b^2)^2*(a + b*Cos[c + d*x])))/d + ((-4*a*(-(a^2*A*b) + A*b^3 + a^3*B - a*b^2*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a)*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a)/Sqrt[2]]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(3*a^3*A + a*A*b^2 - 4*a^2*b*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a)*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a)/Sqrt[2]]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a)*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2/a)/Sqrt[2]]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(3*a^2*A*b + A*b^3 - 4*a*b^2*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)...`

3.433.3 Rubi [A] (verified)

Time = 1.53 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3478, 27, 3042, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.433. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{3478} \\
& \frac{2 \int \frac{Ab-aB-3(aA-bB)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{3(a^2-b^2)} - \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{Ab-aB-3(aA-bB)\cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{3(a^2-b^2)} - \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{Ab-aB-3(aA-bB)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3(a^2-b^2)} - \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \\
& \quad \downarrow \text{3472} \\
& \frac{\int \frac{3Aa^2-4bBa+Ab^2+(-Ba^2+4Aba-3b^2B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} - \frac{2(3a^2A-4abB+Ab^2)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3Aa^2-4bBa+Ab^2+(-Ba^2+4Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} - \frac{2(3a^2A-4abB+Ab^2)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \\
& \quad \downarrow \text{3477} \\
& \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}
\end{aligned}$$

3.433. $\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$

$$\frac{(3a^2A-4abB+Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b)(a(3A+B)-b(A+3B)) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} - \frac{2(3a^2A-4abB+Ab^2)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}} \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(3a^2A-4abB+Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(a(3A+B)-b(A+3B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} - \frac{2(3a^2A-4abB+Ab^2)}{d(a^2-b^2)\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}} \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3295

$$\frac{(3a^2A-4abB+Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(a(3A+B)-b(A+3B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticE}}{a^2d}}{a^2-b^2} - \frac{2(a-b)\sqrt{a+b}(a(3A+B)-b(A+3B))}{ad}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}} \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

↓ 3473

$$\frac{2(a-b)\sqrt{a+b}(3a^2A-4abB+Ab^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) - \frac{a+b}{a-b}}{a^2d} - \frac{2(a-b)\sqrt{a+b}(a(3A+B)-b(A+3B))}{a^2-b^2}$$

$$\frac{3(a^2-b^2)}{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}} \frac{2(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3d(a^2-b^2)(a+b\cos(c+dx))^{3/2}}$$

3(a² - b²)

input `Int[(Sqrt[Cos[c + d*x]]*(A + B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(5/2),x]`

```
output (-2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) - (((2*(a - b)*Sqrt[a + b]*(3*a^2*A + A*b^2 - 4*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(a*(3*A + B) - b*(A + 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a^2 - b^2) - (2*(3*a^2*A + A*b^2 - 4*a*b*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])/(3*(a^2 - b^2))
```

3.433.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)]*Sqrt[(a_) + (b_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3472 Int[((A_) + (B_)*sin[(e_)] + (f_)*(x_)]/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)]*((a_) + (b_)*sin[(e_)] + (f_)*(x_))^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_
.)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

```
rule 3478 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*
x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a
+ b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*
(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*
Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

3.433.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4742 vs. $2(359) = 718$.

Time = 14.55 (sec) , antiderivative size = 4743, normalized size of antiderivative = 12.13

method	result	size
default	Expression too large to display	4743
parts	Expression too large to display	4874

```
input int(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2),x,method=_RET
URNVERBOSE)
```

$$3.433. \quad \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

```

output 1/d*(2/3*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^
2+1))^(1/2)*(-csc(d*x+c)^2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(
1/2))*a^3*b*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-
cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*(1-cos(d*x
+c))^2-7*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*b*(-csc
(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(
d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)-5*EllipticF(cot(d*x+c)-csc(d
*x+c),(-a-b)/(a+b))^(1/2))*a^2*b^2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/
2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/
(a+b))^(1/2)-EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^3*(
-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-
csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)+6*EllipticE(cot(d*x+c)-c
sc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*b*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(
1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b
)/(a+b))^(1/2)+4*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2
*b^2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+
c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)+2*EllipticE(cot(d*
x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^3*(-csc(d*x+c)^2*(1-cos(d*x+c)...

```

3.433.5 Fracas [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{5/2}} dx$$

```

input integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="fracas")

```

```

output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/
(b^3*cos(d*x + c)^3 + 3*a*b^2*cos(d*x + c)^2 + 3*a^2*b*cos(d*x + c) + a^3)
, x)

```

3.433.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(A+B\cos(c+dx))\sqrt{\cos(c+dx)}}{(a+b\cos(c+dx))^{5/2}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(cos(c + d*x))/(a + b*cos(c + d*x))**(5/2), x)`

3.433.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)^(5/2), x)`

3.433.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{5/2}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2),x, algo rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)^(5/2), x)`

3.433.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx = \int \frac{\sqrt{\cos(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

input `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

output `int((cos(c + d*x)^(1/2)*(A + B*cos(c + d*x)))/(a + b*cos(c + d*x))^(5/2),x)`

3.434
$$\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{5/2}} dx$$

3.434.1 Optimal result 4043
 3.434.2 Mathematica [C] (verified) 4044
 3.434.3 Rubi [A] (verified) 4044
 3.434.4 Maple [B] (warning: unable to verify) 4048
 3.434.5 Fricas [F] 4048
 3.434.6 Sympy [F(-1)] 4049
 3.434.7 Maxima [F] 4049
 3.434.8 Giac [F] 4049
 3.434.9 Mupad [F(-1)] 4050

3.434.1 Optimal result

Integrand size = 35, antiderivative size = 429

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \frac{2(6a^2 Ab - 2Ab^3 - 3a^3 B - ab^2 B) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}}\right)\right)}{3a^3(a-b)(a+b)^3} + \frac{2(2Ab^2 - 3a^2(A+B) + ab(3A+B)) \cot(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a^2 \sqrt{a+b}(a^2 - b^2) d} + \frac{2b(Ab - aB) \sqrt{\cos(c + dx)} \sin(c + dx)}{3a(a^2 - b^2) d(a + b \cos(c + dx))^{3/2}} - \frac{2(6a^2 Ab - 2Ab^3 - 3a^3 B - ab^2 B) \sin(c + dx)}{3a(a^2 - b^2)^2 d \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}$$

output

```
2/3*b*(A*b-B*a)*sin(d*x+c)*cos(d*x+c)^(1/2)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))
^(3/2)-2/3*(6*A*a^2*b-2*A*b^3-3*B*a^3-B*a*b^2)*sin(d*x+c)/a/(a^2-b^2)^2/d/
cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)+2/3*(6*A*a^2*b-2*A*b^3-3*B*a^3-B*a
*b^2)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(
1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c)
))/(a-b)^(1/2)/a^3/(a-b)/(a+b)^(3/2)/d-2/3*(2*A*b^2-3*a^2*(A+B)+a*b*(3*A+B
))*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2
),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(
a-b))^(1/2)/a^2/(a^2-b^2)/d/(a+b)^(1/2)
```

3.434.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.62 (sec) , antiderivative size = 1384, normalized size of antiderivative = 3.23

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(5/2)),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*((-2*(-(A*b^2*Sin[c + d*x]) + a*b*B*Sin[c + d*x]))/(3*a*(a^2 - b^2)*(a + b*Cos[c + d*x])^2) - (2*(-6*a^2*A*b^2*Sin[c + d*x] + 2*A*b^4*Sin[c + d*x] + 3*a^3*b*B*Sin[c + d*x] + a*b^3*B*Sin[c + d*x]))/(3*a^2*(a^2 - b^2)^2*(a + b*Cos[c + d*x])))/d + ((-4*a*(3*a^4*A - 5*a^2*A*b^2 + 2*A*b^4 - a^3*b*B + a*b^3*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(-6*a^3*A*b + 2*a*A*b^3 + 3*a^4*B + a^2*b^2*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[(a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(-6*a^2*A*b^2 + 2*A*b^4 + 3*a^3*b*B + a*b^3*B)*((I*Cos[(c + d*x)/2]*Sqrt[a ...`

3.434.3 Rubi [A] (verified)

Time = 1.71 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3479, 27, 3042, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.434. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\sin(c + dx + \frac{\pi}{2})(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{2 \int \frac{3Aa^2 - bBa - 3(Ab - aB) \cos(c + dx)a - 2Ab^2}{2\sqrt{\cos(c + dx)(a + b \cos(c + dx))^{3/2}} dx}{3a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3Aa^2 - bBa - 3(Ab - aB) \cos(c + dx)a - 2Ab^2}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))^{3/2}} dx}{3a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3Aa^2 - bBa - 3(Ab - aB) \sin(c + dx + \frac{\pi}{2})a - 2Ab^2}{\sqrt{\sin(c + dx + \frac{\pi}{2})(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx}{3a(a^2 - b^2)} + \frac{2b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3472} \\
 & \frac{\int \frac{-3Ba^3 + 6Aba^2 - b^2Ba + (3Aa^2 - 4bBa + Ab^2) \cos(c + dx)a - 2Ab^3}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} - \frac{2(-3a^3B + 6a^2Ab - ab^2B - 2Ab^3) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} + \\
 & \quad \frac{3a(a^2 - b^2)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \frac{2b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{-3Ba^3 + 6Aba^2 - b^2Ba + (3Aa^2 - 4bBa + Ab^2) \sin(c + dx + \frac{\pi}{2})a - 2Ab^3}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{a^2 - b^2} - \frac{2(-3a^3B + 6a^2Ab - ab^2B - 2Ab^3) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} + \\
 & \quad \frac{3a(a^2 - b^2)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \frac{2b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3477} \\
 & \frac{(-3a^3B + 6a^2Ab - ab^2B - 2Ab^3) \int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - (a - b)(-3a^2(A + B) + ab(3A + B) + 2Ab^2) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx}{a^2 - b^2} - \frac{2(-3a^3B + 6a^2Ab - ab^2B - 2Ab^3) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} + \\
 & \quad \frac{3a(a^2 - b^2)}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \frac{2b(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3ad(a^2 - b^2)(a + b \cos(c + dx))^{3/2}}
 \end{aligned}$$

3.434. $\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)(a + b \cos(c + dx))^{5/2}} dx$

↓ 3042

$$\frac{(-3a^3B+6a^2Ab-ab^2B-2Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(-3a^2(A+B)+ab(3A+B)+2Ab^2) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2}$$

$$3a(a^2-b^2)$$

$$\frac{2b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3295

$$\frac{(-3a^3B+6a^2Ab-ab^2B-2Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(-3a^2(A+B)+ab(3A+B)+2Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a}{ad}}}{a^2-b^2}}{a^2-b^2}$$

$$3a(a^2-b^2)$$

$$\frac{2b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}}$$

↓ 3473

$$\frac{2b(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b \cos(c+dx))^{3/2}} + \frac{2(a-b)\sqrt{a+b}(-3a^3B+6a^2Ab-ab^2B-2Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) - \frac{2(a-b)\sqrt{a+b}(-3a^2(A+B))}{a^2-b^2}}{a^2d}$$

$$3a(a^2-b^2)$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(5/2)),x]`

output `(2*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((2*(a - b)*Sqrt[a + b]*(6*a^2*A*b - 2*A*b^3 - 3*a^3*B - a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(2*A*b^2 - 3*a^2*(A + B) + a*b*(3*A + B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a^2 - b^2) - (2*(6*a^2*A*b - 2*A*b^3 - 3*a^3*B - a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))/(3*a*(a^2 - b^2))`

3.434. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))^{5/2}}} dx$

3.434.3.1 Defintions of rubi rules used

- rule 277 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_.) + (b_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3472 `Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`
- rule 3473 `Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(((b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`
- rule 3477 `Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3479 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m +
1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e
+ f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +
2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)
*(m + n + 3)*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n
}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Rat
ionalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(I
ntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0]
))
```

3.434.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 5454 vs. $2(397) = 794$.

Time = 18.12 (sec) , antiderivative size = 5455, normalized size of antiderivative = 12.72

method	result	size
default	Expression too large to display	5455
parts	Expression too large to display	5509

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.434.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sqrt{\cos(dx + c)}} dx$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="fricas")
```

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/
(b^3*cos(d*x + c)^4 + 3*a*b^2*cos(d*x + c)^3 + 3*a^2*b*cos(d*x + c)^2 + a^3*cos(d*x + c)), x)`

3.434.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(5/2), x)`

output `Timed out`

3.434.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(5/2), x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sqrt(cos(d*x +
c))), x)`

3.434.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(5/2), x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sqrt(cos(d*x +
c))), x)`

3.434. $\int \frac{A+B \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{5/2}} dx$

3.434.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(5/2)),x)`

3.435
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$$

3.435.1 Optimal result 4051
 3.435.2 Mathematica [C] (warning: unable to verify) 4052
 3.435.3 Rubi [A] (verified) 4052
 3.435.4 Maple [B] (warning: unable to verify) 4057
 3.435.5 Fricas [F] 4057
 3.435.6 Sympy [F(-1)] 4058
 3.435.7 Maxima [F] 4058
 3.435.8 Giac [F] 4058
 3.435.9 Mupad [F(-1)] 4059

3.435.1 Optimal result

Integrand size = 35, antiderivative size = 456

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \frac{2(3a^4 A - 15a^2 Ab^2 + 8Ab^4 + 6a^3 b B - 2ab^3 B) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right) + 2(8Ab^3 - 3a^3(A - B) + 2ab^2(3A - B) - 3a^2 b(3A + B)) \cot(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right) + \frac{2b(Ab - aB) \sin(c + dx)}{3a(a^2 - b^2) d \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} + \frac{2b(8a^2 Ab - 4Ab^3 - 5a^3 B + ab^2 B) \sin(c + dx)}{3a^2(a^2 - b^2)^2 d \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}}{3a^4(a - b)(a - b) \sqrt{a + b} (a^2 - b^2) d}$$

output

```
2/3*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)/cos(d*x+c)
^(1/2)+2/3*b*(8*A*a^2*b-4*A*b^3-5*B*a^3+B*a*b^2)*sin(d*x+c)/a^2/(a^2-b^2)
^2/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)+2/3*(3*A*a^4-15*A*a^2*b^2+8*A*
b^4+6*B*a^3*b-2*B*a*b^3)*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)
^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d*x+c))/(a+b))^(1/
2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^4/(a-b)/(a+b)^(3/2)/d+2/3*(8*A*b^3-3*a
^3*(A-B)+2*a*b^2*(3*A-B)-3*a^2*b*(3*A+B))*cot(d*x+c)*EllipticF((a+b*cos(d*
x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a*(1-sec(d
*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/(a^2-b^2)/d/(a+b)^(
1/2)
```

3.435.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$$

3.435.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.76 (sec) , antiderivative size = 1431, normalized size of antiderivative = 3.14

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(5/2)),x]`

output

```
-1/3*((-4*a*(9*a^4*A*b - 17*a^2*A*b^3 + 8*A*b^5 - 3*a^5*B + 5*a^3*b^2*B -
2*a*b^4*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos
[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)
/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c
+ d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sq
rt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(3*a^5*A - 15*a^3*A*b^2 +
8*a*A*b^4 + 6*a^4*b*B - 2*a^2*b^3*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/
(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a])*Sqrt[((a +
b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[
((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Si
n[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) -
(Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-((a + b)*Cos[c + d*x]*
Csc[(c + d*x)/2]^2)/a])*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*
Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c +
d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[
c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(3*a^4*A*b - 15*a^2*A*b^3 + 8*A*b
^5 + 6*a^3*b^2*B - 2*a*b^4*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]
]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b
)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos
[c + d*x])*Sec[c + d*x])/(a + b)]) + (2*a*((a*Sqrt[((a + b)*Cot[(c + d*...
```

3.435.3 Rubi [A] (verified)

Time = 1.98 (sec) , antiderivative size = 477, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.435. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{5/2}} dx \\
 & \quad \downarrow \text{3479} \\
 & \frac{2 \int \frac{3Aa^2 + bBa - 3(Ab - aB) \cos(c + dx)a - 4Ab^2 + 2b(Ab - aB) \cos^2(c + dx)}{2 \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx}{\frac{3a(a^2 - b^2)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} + \frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3Aa^2 + bBa - 3(Ab - aB) \cos(c + dx)a - 4Ab^2 + 2b(Ab - aB) \cos^2(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx}{\frac{3a(a^2 - b^2)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} + \frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3Aa^2 + bBa - 3(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)a - 4Ab^2 + 2b(Ab - aB) \sin\left(c + dx + \frac{\pi}{2}\right)^2}{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{3/2}} dx}{\frac{3a(a^2 - b^2)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} + \frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}}} \\
 & \quad \downarrow \text{3534} \\
 & \frac{2 \int \frac{3Aa^4 + 6bBa^3 - 15Ab^2a^2 - 2b^3Ba - (-3Ba^3 + 6Aba^2 - b^2Ba - 2Ab^3) \cos(c + dx)a + 8Ab^4}{2 \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} a(a^2 - b^2)} dx + \frac{2b(-5a^3B + 8a^2Ab + ab^2B - 4Ab^3) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}}{\frac{3a(a^2 - b^2)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} + \frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3Aa^4 + 6bBa^3 - 15Ab^2a^2 - 2b^3Ba - (-3Ba^3 + 6Aba^2 - b^2Ba - 2Ab^3) \cos(c + dx)a + 8Ab^4}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} a(a^2 - b^2)} dx + \frac{2b(-5a^3B + 8a^2Ab + ab^2B - 4Ab^3) \sin(c + dx)}{ad(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}}}{\frac{3a(a^2 - b^2)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} + \frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}}}
 \end{aligned}$$

3.435. $\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx$

↓ 3042

$$\frac{\int \frac{3Aa^4 + 6bBa^3 - 15Ab^2a^2 - 2b^3Ba - (-3Ba^3 + 6Aba^2 - b^2Ba - 2Ab^3) \sin(c+dx + \frac{\pi}{2})a + 8Ab^4}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \frac{2b(-5a^3B + 8a^2Ab + ab^2B - 4Ab^3) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}}{\frac{3a(a^2-b^2)}{3ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} + \frac{2b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}} +$$

↓ 3477

$$\frac{(a-b)(-3a^3(A-B) - 3a^2b(3A+B) + 2ab^2(3A-B) + 8Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} dx + (3a^4A + 6a^3bB - 15a^2Ab^2 - 2ab^3B + 8Ab^4) \int \frac{\cos(c+dx)}{\cos^2(c+dx)}}{a(a^2-b^2)} + \frac{3a(a^2-b^2)}{3ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} + \frac{2b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}}$$

↓ 3042

$$\frac{(a-b)(-3a^3(A-B) - 3a^2b(3A+B) + 2ab^2(3A-B) + 8Ab^3) \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + (3a^4A + 6a^3bB - 15a^2Ab^2 - 2ab^3B + 8Ab^4) \int \frac{1}{\sin(c+dx + \frac{\pi}{2})}}{a(a^2-b^2)} + \frac{3a(a^2-b^2)}{3ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} + \frac{2b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}}$$

↓ 3295

$$\frac{(3a^4A + 6a^3bB - 15a^2Ab^2 - 2ab^3B + 8Ab^4) \int \frac{\sin(c+dx + \frac{\pi}{2}) + 1}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(-3a^3(A-B) - 3a^2b(3A+B) + 2ab^2(3A-B) + 8Ab^3) \cot(c+dx)}{a(a^2-b^2)}}{a(a^2-b^2)} + \frac{3a(a^2-b^2)}{3ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} + \frac{2b(Ab-aB) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}}$$

↓ 3473

$$\frac{2b(Ab-aB) \sin(c+dx)}{3ad(a^2-b^2) \sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B - 4Ab^3) \sin(c+dx)}{ad(a^2-b^2) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}} + \frac{2(a-b)\sqrt{a+b}(-3a^3(A-B) - 3a^2b(3A+B) + 2ab^2(3A-B) + 8Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx))}{a-b}}}{ad}$$

3.435. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(5/2)),x
]`

output `(2*b*(A*b - a*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)) + (((2*(a - b)*Sqrt[a + b]*(3*a^4*A - 15*a^2*A*b^2 + 8*A*b^4 + 6*a^3*b*B - 2*a*b^3*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(8*A*b^3 - 3*a^3*(A - B) + 2*a*b^2*(3*A - B) - 3*a^2*b*(3*A + B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a*(a^2 - b^2)) + (2*b*(8*a^2*A*b - 4*A*b^3 - 5*a^3*B + a*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])/(3*a*(a^2 - b^2))`

3.435.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
 $\wedge(3/2)*\text{Sqrt}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]]), x_Symbol] \rightarrow \text{Simp}[-2*A*$
 $(c - d)*(\text{Tan}[e + f*x]/(f*b*c^2))*\text{Rt}[(c + d)/b, 2]*\text{Sqrt}[c*((1 + \text{Csc}[e + f*x])$
 $)/(c - d)]*\text{Sqrt}[c*((1 - \text{Csc}[e + f*x])/(c + d))]*\text{EllipticE}[\text{ArcSin}[\text{Sqrt}[c +$
 $d*\text{Sin}[e + f*x]]/\text{Sqrt}[b*\text{Sin}[e + f*x]]/\text{Rt}[(c + d)/b, 2]], -(c + d)/(c - d),$
 $x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\&$
 $\text{PosQ}[(c + d)/b]$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*
 $(x_)]\wedge(3/2)*\text{Sqrt}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]]), x_Symbol] \rightarrow \text{S}$
 $\text{imp}[(A - B)/(a - b) \text{ Int}[1/(\text{Sqrt}[a + b*\text{Sin}[e + f*x]]*\text{Sqrt}[c + d*\text{Sin}[e + f*$
 $x]]), x], x] - \text{Simp}[(A*b - a*B)/(a - b) \text{ Int}[(1 + \text{Sin}[e + f*x])/((a + b*\text{Si}$
 $n[e + f*x])\wedge(3/2)*\text{Sqrt}[c + d*\text{Sin}[e + f*x]]), x], x] /; \text{FreeQ}\{a, b, c, d, e$
 $, f, A, B\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2,$
 $0] \&\& \text{NeQ}[A, B]$`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
 $(f_)*(x_)]*((c_) + (d_)*\sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] \rightarrow \text{Si}$
 $\text{mp}[(-(A*b^2 - a*b*B))*\text{Cos}[e + f*x]*(a + b*\text{Sin}[e + f*x])^(m + 1)*((c + d*\text{Sin}$
 $[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + \text{Simp}[1/((m +$
 $1)*(b*c - a*d)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[e + f*x])^(m + 1)*(c + d*\text{Sin}[e$
 $+ f*x])^n*\text{Simp}[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +$
 $2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*\text{Sin}[e + f*x] - b*d*(A*b - a*B)$
 $*(m + n + 3)*\text{Sin}[e + f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, n$
 $\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{Rat}$
 $\text{ionalQ}[m] \&\& m < -1 \&\& ((\text{EqQ}[a, 0] \&\& \text{IntegerQ}[m] \&\& !\text{IntegerQ}[n]) \|\ !(\text{I}$
 $n\text{tegerQ}[2*n] \&\& \text{LtQ}[n, -1] \&\& ((\text{IntegerQ}[n] \&\& !\text{IntegerQ}[m]) \|\ \text{EqQ}[a, 0])$
 $))$`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.435.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 6515 vs. $2(424) = 848$.

Time = 20.07 (sec) , antiderivative size = 6516, normalized size of antiderivative = 14.29

method	result	size
default	Expression too large to display	6516
parts	Expression too large to display	6518

```
input int((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.435.5 Fracas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{5}{2}} \cos^{\frac{3}{2}}(dx + c)} dx$$

```
input integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="fracas")
```

3.435. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(b^3*cos(d*x + c)^5 + 3*a*b^2*cos(d*x + c)^4 + 3*a^2*b*cos(d*x + c)^3 + a^3*cos(d*x + c)^2), x)`

3.435.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(5/2), x)`

output `Timed out`

3.435.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(3/2)), x)`

3.435.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(5/2), x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(3/2)), x)`

3.435. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

3.435.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{3/2}(a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(5/2)),x)`

3.436
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$$

3.436.1 Optimal result 4060
 3.436.2 Mathematica [C] (warning: unable to verify) 4061
 3.436.3 Rubi [A] (verified) 4062
 3.436.4 Maple [B] (warning: unable to verify) 4068
 3.436.5 Fracas [F] 4068
 3.436.6 Sympy [F(-1)] 4068
 3.436.7 Maxima [F] 4069
 3.436.8 Giac [F] 4069
 3.436.9 Mupad [F(-1)] 4069

3.436.1 Optimal result

Integrand size = 35, antiderivative size = 567

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx =$$

$$\frac{2(8a^4 Ab - 28a^2 Ab^3 + 16Ab^5 - 3a^5 B + 15a^3 b^2 B - 8ab^4 B) \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right)}{3a^5(a-b)(a+b)^{3/2}d}$$

$$\frac{2(16Ab^4 - a^4(A - 3B) + 4ab^3(3A - 2B) - 9a^3b(A - B) - 2a^2b^2(8A + 3B)) \cot(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right)}{3a^4 \sqrt{a+b} (a^2 - b^2) d}$$

$$+ \frac{2b(Ab - aB) \sin(c + dx)}{3a(a^2 - b^2) d \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}}$$

$$+ \frac{2b(10a^2 Ab - 6Ab^3 - 7a^3 B + 3ab^2 B) \sin(c + dx)}{3a^2(a^2 - b^2)^2 d \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}}$$

$$+ \frac{2(a^4 A - 13a^2 Ab^2 + 8Ab^4 + 8a^3 b B - 4ab^3 B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3a^3(a^2 - b^2)^2 d \cos^{\frac{3}{2}}(c + dx)}$$

3.436.
$$\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$$

output $\frac{2}{3}b(Ab-Ba)\sin(dx+c)/a/(a^2-b^2)/d/\cos(dx+c)^{3/2}/(a+b\cos(dx+c))^{3/2}+2/3b(10Aa^2b-6Ab^3-7B^2a^3+3B^2ab^2)\sin(dx+c)/a^2/(a^2-b^2)^2/d/\cos(dx+c)^{3/2}/(a+b\cos(dx+c))^{1/2}+2/3(Aa^4-13Aa^2b^2+8Ab^4+8B^2a^3b-4B^2ab^3)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/a^3/(a^2-b^2)^2/d/\cos(dx+c)^{3/2}-2/3(8Aa^4b-28Aa^2b^3+16Ab^5-3B^2a^5+15B^2ab^3*b^2-8B^2ab^4)\cot(dx+c)*\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a^5/(a-b)/(a+b)^{3/2}/d-2/3(16Ab^4-a^4(A-3B)+4ab^3(3A-2B)-9a^3b(A-B)-2a^2b^2(8A+3B))\cot(dx+c)*\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a^4/(a^2-b^2)/d/(a+b)^{1/2}$

3.436.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 7.06 (sec) , antiderivative size = 1499, normalized size of antiderivative = 2.64

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^(5/2)),x]`

```
output ((-4*a*(a^6*A + 15*a^4*A*b^2 - 32*a^2*A*b^4 + 16*A*b^6 - 9*a^5*b*B + 17*a^3*b^3*B - 8*a*b^5*B)*Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x]*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - 4*a*(8*a^5*A*b - 28*a^3*A*b^3 + 16*a*A*b^5 - 3*a^6*B + 15*a^4*b^2*B - 8*a^2*b^4*B)*((Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/((a + b)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) - (Sqrt[((a + b)*Cot[(c + d*x)/2]^2)/(-a + b)]*Sqrt[-(((a + b)*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a)]*Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]*Csc[c + d*x]*EllipticPi[-(a/b), ArcSin[Sqrt[((a + b*Cos[c + d*x])*Csc[(c + d*x)/2]^2)/a]/Sqrt[2]], (-2*a)/(-a + b)]*Sin[(c + d*x)/2]^4)/(b*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + 2*(8*a^4*A*b^2 - 28*a^2*A*b^4 + 16*A*b^6 - 3*a^5*b*B + 15*a^3*b^3*B - 8*a*b^5*B)*((I*Cos[(c + d*x)/2]*Sqrt[a + b*Cos[c + d*x]]*EllipticE[I*ArcSinh[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]], (-2*a)/(-a - b)]*Sec[c + d*x])/(b*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*Sqrt[((a + b*Cos[c + d*x])*Sec[c + d*x])/(a + ...
```

3.436.3 Rubi [A] (verified)

Time = 2.62 (sec) , antiderivative size = 581, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{5/2} (a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx$$

↓ 3479

3.436. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

$$2 \int \frac{4b(Ab-aB) \cos^2(c+dx) - 3a(Ab-aB) \cos(c+dx) + 3(Aa^2+bBa-2Ab^2)}{2 \cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$3ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}$$

↓ 27

$$\int \frac{4b(Ab-aB) \cos^2(c+dx) - 3a(Ab-aB) \cos(c+dx) + 3(Aa^2+bBa-2Ab^2)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$3ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}$$

↓ 3042

$$\int \frac{4b(Ab-aB) \sin(c+dx+\frac{\pi}{2})^2 - 3a(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + 3(Aa^2+bBa-2Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{5/2}(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$3ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}$$

↓ 3534

$$2 \int \frac{2b(-7Ba^3+10Aba^2+3b^2Ba-6Ab^3) \cos^2(c+dx) - a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3) \cos(c+dx) + 3(Aa^4+8bBa^3-13Ab^2a^2-4b^3Ba+8Ab^4)}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

$$\frac{3a(a^2-b^2)}{ad(a^2-b^2)} + \frac{2b(-7a^3B+10a^2bA+3b^2aB-6Ab^3)}{ad(a^2-b^2)}$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$3ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}$$

↓ 27

$$\int \frac{2b(-7Ba^3+10Aba^2+3b^2Ba-6Ab^3) \cos^2(c+dx) - a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3) \cos(c+dx) + 3(Aa^4+8bBa^3-13Ab^2a^2-4b^3Ba+8Ab^4)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

$$\frac{3a(a^2-b^2)}{ad(a^2-b^2)} + \frac{2b(-7a^3B+10a^2bA+3b^2aB-6Ab^3)}{ad(a^2-b^2)}$$

$$\frac{3a(a^2-b^2)}{2b(Ab-aB) \sin(c+dx)}$$

$$3ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}$$

↓ 3042

3.436. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

$$\int \frac{2b(-7Ba^3+10Aba^2+3b^2Ba-6Ab^3) \sin(c+dx+\frac{\pi}{2})^2 - a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3) \sin(c+dx+\frac{\pi}{2}) + 3(Aa^4+8bBa^3-13Ab^2a^2-4b^3Ba+8Ab^4)}{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b(-}{a(a^2-b^2)}$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} \quad 3a(a^2 - b^2)$$

3534

$$2 \int - \frac{3(-3Ba^5+8Aba^4+15b^2Ba^3-28Ab^3a^2-8b^4Ba-(Aa^4-6bBa^3+7Ab^2a^2+2b^3Ba-4Ab^4) \cos(c+dx)a+16Ab^5)}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx + \frac{2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4)}{ad \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} \quad 3a(a^2 - b^2)$$

27

$$2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4) \sin(c+dx) \sqrt{a+b \cos(c+dx)} \int \frac{-3Ba^5+8Aba^4+15b^2Ba^3-28Ab^3a^2-8b^4Ba-(Aa^4-6bBa^3+7Ab^2a^2+2b^3Ba-4Ab^4) \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} \quad 3a(a^2 - b^2)$$

3042

$$2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4) \sin(c+dx) \sqrt{a+b \cos(c+dx)} \int \frac{-3Ba^5+8Aba^4+15b^2Ba^3-28Ab^3a^2-8b^4Ba-(Aa^4-6bBa^3+7Ab^2a^2+2b^3Ba-4Ab^4) \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} \quad 3a(a^2 - b^2)$$

3477

$$2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4) \sin(c+dx) \sqrt{a+b \cos(c+dx)} \int \frac{(a-b)(-a^4(A-3B)-9a^3b(A-B)-2a^2b^2(8A+3B)+4ab^3(3A-2B)+16Ab^4)}{\sqrt{\cos(c+dx)}} dx$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad(a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} \quad 3a(a^2 - b^2)$$

3.436. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

↓ 3042

$$\frac{2(a^4 A + 8a^3 bB - 13a^2 Ab^2 - 4ab^3 B + 8Ab^4) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(a-b)(- (a^4(A-3B)) - 9a^3 b(A-B) - 2a^2 b^2(8A+3B) + 4ab^3(3A-2B) + 16Ab^4) \int \frac{\sqrt{\sin(c+dx)}}{\sqrt{\sin(c+dx)}} dx}{a(a^2-b^2)}$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad (a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}}$$

↓ 3295

$$\frac{2(a^4 A + 8a^3 bB - 13a^2 Ab^2 - 4ab^3 B + 8Ab^4) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^5 B + 8a^4 Ab + 15a^3 b^2 B - 28a^2 Ab^3 - 8ab^4 B + 16Ab^5) \int \frac{\sin(c+dx + \frac{\pi}{2}) + 1}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx)}} dx}{ad \cos^{\frac{3}{2}}(c+dx)}$$

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad (a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}}$$

↓ 3473

$$\frac{2b(Ab - aB) \sin(c + dx)}{3ad (a^2 - b^2) \cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} +$$

$$\frac{2b(-7a^3 B + 10a^2 Ab + 3ab^2 B - 6Ab^3) \sin(c+dx)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} + \frac{2(a^4 A + 8a^3 bB - 13a^2 Ab^2 - 4ab^3 B + 8Ab^4) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2(a-b)\sqrt{a+b}(- (a^4(A-3B)) - 9a^3 b(A-B) - 2a^2 b^2(8A+3B) + 4ab^3(3A-2B) + 16Ab^4) \int \frac{\sqrt{\sin(c+dx)}}{\sqrt{\sin(c+dx)}} dx}{ad \cos^{\frac{3}{2}}(c+dx)}$$

input `Int[(A + B*Cos[c + d*x])/(Cos[c + d*x]^(5/2)*(a + b*Cos[c + d*x])^(5/2)),x]`

3.436. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$


```
output (2*b*(A*b - a*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*(a +
b*Cos[c + d*x])^(3/2)) + ((2*b*(10*a^2*A*b - 6*A*b^3 - 7*a^3*B + 3*a*b^2*B
)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x
]]) + (-(((2*(a - b)*Sqrt[a + b]*(8*a^4*A*b - 28*a^2*A*b^3 + 16*A*b^5 - 3*
a^5*B + 15*a^3*b^2*B - 8*a*b^4*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b
*Cos[c + d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt
[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a^
2*d) + (2*(a - b)*Sqrt[a + b]*(16*A*b^4 - a^4*(A - 3*B) + 4*a*b^3*(3*A - 2
*B) - 9*a^3*b*(A - B) - 2*a^2*b^2*(8*A + 3*B))*Cot[c + d*x]*EllipticF[ArcS
in[Sqrt[a + b*Cos[c + d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(
a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/
(a - b)]/(a*d))/a) + (2*(a^4*A - 13*a^2*A*b^2 + 8*A*b^4 + 8*a^3*b*B - 4*a
*b^3*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(a*d*Cos[c + d*x]^(3/2))/(
a*(a^2 - b^2)))/(3*a*(a^2 - b^2))
```

3.436.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)]])*Sqrt[(a_) + (b_)*sin[(e_.) + (f
_)*(x_)]], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_.) + (f_)*(x_)])/(((b_)*sin[(e_.) + (f_)*(x_)]
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_.) + (f_)*(x_)]], x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

3.436.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 8900 vs. $2(529) = 1058$.

Time = 24.34 (sec) , antiderivative size = 8901, normalized size of antiderivative = 15.70

method	result	size
parts	Expression too large to display	8901
default	Expression too large to display	9524

input `int((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

3.436.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(5/2),x,algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(b^3*cos(d*x + c)^6 + 3*a*b^2*cos(d*x + c)^5 + 3*a^2*b*cos(d*x + c)^4 + a^3*cos(d*x + c)^3), x)`

3.436.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)**(5/2)/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.436. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{\frac{5}{2}}} dx$

3.436.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(5
/2)), x)`

3.436.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{5}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/cos(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*cos(d*x + c)^(5
/2)), x)`

3.436.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx = \int \frac{A + B \cos(c + dx)}{\cos(c + dx)^{5/2} (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^(5/2)),x
)`

output `int((A + B*cos(c + d*x))/(cos(c + d*x)^(5/2)*(a + b*cos(c + d*x))^(5/2)),
x)`

3.436. $\int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(a+b \cos(c+dx))^{5/2}} dx$

3.437
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.437.1 Optimal result 4070
 3.437.2 Mathematica [A] (verified) 4071
 3.437.3 Rubi [A] (verified) 4071
 3.437.4 Maple [B] (verified) 4076
 3.437.5 Fricas [F] 4077
 3.437.6 Sympy [F] 4078
 3.437.7 Maxima [F] 4078
 3.437.8 Giac [F] 4078
 3.437.9 Mupad [F(-1)] 4079

3.437.1 Optimal result

Integrand size = 38, antiderivative size = 419

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx =$$

$$\frac{(a-b)\sqrt{a+b}B \cot(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{abd}$$

$$+ \frac{\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{bd}$$

$$+ \frac{a\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^2d}$$

$$+ \frac{aB \sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} + \frac{B\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{a+b \cos(c+dx)}}$$

```
output a*B*sin(d*x+c)/b/d/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2)+B*sin(d*x+c)*cos(d*x+c)^(1/2)/d/(a+b*cos(d*x+c))^(1/2)-(a-b)*B*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/b/d+B*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b/d+a*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2), (a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b^2/d
```

3.437.
$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.437.2 Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.54

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \frac{B\cos^{\frac{3}{2}}(c+dx)\sec^2\left(\frac{1}{2}(c+dx)\right)\left((a+b)\sqrt{\frac{a+b\cos(c+dx)}{(a+b)(1+\cos(c+dx))}}E\left(a\right)\right)}{(a+b\cos(c+dx))^{3/2}}$$

input `Integrate[(Cos[c + d*x]^(3/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]`

output `(B*Cos[c + d*x]^(3/2)*Sec[(c + d*x)/2]^2*((a + b)*Sqrt[(a + b*Cos[c + d*x])/(a + b)*(1 + Cos[c + d*x])])*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 2*a*Sqrt[(a + b*Cos[c + d*x])/(a + b)*(1 + Cos[c + d*x])]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*(a + b*Cos[c + d*x])*Tan[(c + d*x)/2])/(2*b*d*(Cos[c + d*x]/(1 + Cos[c + d*x]))^(3/2)*Sqrt[a + b*Cos[c + d*x]])`

3.437.3 Rubi [A] (verified)Time = 1.69 (sec) , antiderivative size = 426, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {2011, 3042, 3299, 3042, 3288, 3482, 27, 3042, 3472, 25, 27, 3042, 3280, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}}{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} dx \\ & \quad \downarrow \text{3299} \end{aligned}$$

3.437. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & B \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(a+2b \cos(c+dx))}{\sqrt{a+b \cos(c+dx)}} dx}{2b} - \frac{a \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx}{2b} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+2b \sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{a \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} \right) \\
 & \quad \downarrow \text{3288} \\
 & B \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+2b \sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{2b} + \frac{a\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arccos\left(\frac{a+b \sin(c+dx+\frac{\pi}{2})}{a+b}\right)\right)}{b^2 d} \right) \\
 & \quad \downarrow \text{3482} \\
 & B \left(\frac{\frac{1}{2} \int \frac{2(\cos(c+dx)a^2+ba)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} dx + \frac{2b \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a+b \cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{b^2 d} \right) \\
 & \quad \downarrow \text{27} \\
 & B \left(\frac{\int \frac{\cos(c+dx)a^2+ba}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} dx + \frac{2b \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a+b \cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{b^2 d} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})a^2+ba}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx + \frac{2b \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a+b \cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{b^2 d} \right) \\
 & \quad \downarrow \text{3472}
 \end{aligned}$$

3.437. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

$$B \left(\frac{\int -\frac{a(a^2-b^2)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec)}{a+}}}{2b} \right)$$

↓ 25

$$B \left(\frac{\int -\frac{a(a^2-b^2)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec)}{a+}}}{2b} \right)$$

↓ 27

$$B \left(\frac{-a \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec)}{a+}}}{2b} \right)$$

↓ 3042

$$B \left(\frac{-a \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec)}{a+}}}{2b} \right)$$

↓ 3280

$$B \left(\frac{-a \left(\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right) + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec)}{a+}}}{2b} \right)$$

↓ 3042

$$B \left(\frac{-a \left(\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec)}{a+}}}{2b} \right)$$

↓ 3295

3.437. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$

$$B \left(\frac{-a \left(\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \operatorname{EllipticF} \left(\arcsin \left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}} \right) \right)}{ad}}{2b} \right)$$

↓ 3473

$$B \left(\frac{-a \left(\frac{2(a-b)\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E \left(\arcsin \left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}} \right) \right) \Big|_{-\frac{a+b}{a-b}}}{a^2 d} - \frac{2\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{a+b} \right)}{2b}$$

```
input Int[(Cos[c + d*x]^(3/2)*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2),x]
```

```
output B*((a*Sqrt[a + b]*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b^2*d) + (-(a*((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a^2*d) - (2*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d)) + (2*a*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + (2*b*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + b*Cos[c + d*x]]))/(2*b))
```

3.437.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.437. $\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3280 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin
[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[1/(a - b) Int[1/(Sqrt[a + b*Sin
[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[b/(a - b) Int[(1 + Si
n[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /
; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f
.)*(x)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]`

rule 3299 `Int[((d_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)/Sqrt[(a_.) + (b_.)*sin[(e_.) + (f
.)*(x)]], x_Symbol] := Simp[(-a)*(d/(2*b)) Int[Sqrt[d*Sin[e + f*x]]/Sqr
t[a + b*Sin[e + f*x]], x], x] + Simp[d/(2*b) Int[Sqrt[d*Sin[e + f*x]]*((a
+ 2*b*Sin[e + f*x])/Sqrt[a + b*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d, e
, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 3472 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 3473 Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]
```

```
rule 3482 Int[Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*B*Cos[e + f*x]*Sqrt[a + b*Sin[e + f*x]]*((c + d*Sin[e + f*x])^n/(f*(2*n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*Sin[e + f*x])^(n - 1)/Sqrt[a + b*Sin[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*Sin[e + f*x] + (A*b*d*(2*n + 3) + B*(a*d + 2*b*c*n))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && EqQ[n^2, 1/4]
```

3.437.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 834 vs. $2(387) = 774$.

Time = 14.52 (sec) , antiderivative size = 835, normalized size of antiderivative = 1.99

method	result
default	$\frac{B\left(-E\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}\right)a\left(\cos^2(dx+c)\right)-E\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)}{\dots}$
parts	Expression too large to display

```
input int(cos(d*x+c)^(3/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)
```

$$3.437. \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx$$

output `B/d*(-EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)^2-EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b*cos(d*x+c)^2+2*EllipticPi(cot(d*x+c)-csc(d*x+c), -1, (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)^2-2*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)-2*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b*cos(d*x+c)+4*EllipticPi(cot(d*x+c)-csc(d*x+c), -1, (-a-b)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)+b*cos(d*x+c)^2*sin(d*x+c)-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a-(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c), (-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c), -1, (-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+sin(d*x+c)*cos(d*x+c)*a/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)/b`

3.437.5 Fracas [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2), x, algorithm="fricas")`

output `integral(B*cos(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a), x)`

3.437.6 Sympy [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = B \int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$$

input `integrate(cos(d*x+c)**(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2), x)`

output `B*Integral(cos(c + d*x)**(3/2)/sqrt(a + b*cos(c + d*x)), x)`

3.437.7 Maxima [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2), x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(3/2), x)`

3.437.8 Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{\frac{3}{2}}} dx = \int \frac{(Bb\cos(dx+c)+Ba)\cos(dx+c)^{\frac{3}{2}}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(3/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2), x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*cos(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(3/2), x)`

3.437.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{\cos(c+dx)^{3/2}(Ba+Bb\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2),x)`

output `int((cos(c + d*x)^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/2), x)`

$$3.438 \quad \int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$$

3.438.1 Optimal result	4080
3.438.2 Mathematica [A] (verified)	4080
3.438.3 Rubi [A] (verified)	4081
3.438.4 Maple [A] (verified)	4082
3.438.5 Fricas [F]	4083
3.438.6 Sympy [F]	4083
3.438.7 Maxima [F]	4083
3.438.8 Giac [F]	4084
3.438.9 Mupad [F(-1)]	4084

3.438.1 Optimal result

Integrand size = 38, antiderivative size = 117

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{a+b}B \cot(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{bd}$$

```
output -2*B*cot(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2), (a+b)/b, ((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b/d
```

3.438.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.12

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx = \frac{2B \sqrt{\cos(c+dx)} \sqrt{\frac{a+b \cos(c+dx)}{(a+b)(1+\cos(c+dx))}} \left(\operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(c+dx)\right)\right), \frac{-a+b}{a+b}\right) - 2 \operatorname{EllipticPi}\left(-1, \arcsin\left(d \sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}} \sqrt{a+b \cos(c+dx)}\right)\right)\right)}{d \sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}} \sqrt{a+b \cos(c+dx)}}$$

```
input Integrate[(Sqrt[Cos[c + d*x]]*(a*B + b*B*Cos[c + d*x]))/(a + b*Cos[c + d*x])^(3/2), x]
```

3.438. $\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

output $(-2*B*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/((a + b)*(1 + \text{Cos}[c + d*x]))])*(\text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + d*x)/2]], (-a + b)/(a + b)] - 2*\text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + d*x)/2]], (-a + b)/(a + b)])/(d*\text{Sqrt}[\text{Cos}[c + d*x]/(1 + \text{Cos}[c + d*x])]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])$

3.438.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.079$, Rules used = {2011, 3042, 3288}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cos(c+dx)}(aB + bB \cos(c+dx))}{(a + b \cos(c+dx))^{3/2}} dx$$

↓ 2011

$$B \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a + b \cos(c+dx)}} dx$$

↓ 3042

$$B \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c+dx + \frac{\pi}{2})}} dx$$

↓ 3288

$$\frac{2B\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{bd}$$

input $\text{Int}[(\text{Sqrt}[\text{Cos}[c + d*x]]*(a*B + b*B*\text{Cos}[c + d*x]))/(a + b*\text{Cos}[c + d*x])^(3/2), x]$

output $(-2*\text{Sqrt}[a + b]*B*\text{Cot}[c + d*x]*\text{EllipticPi}[(a + b)/b, \text{ArcSin}[\text{Sqrt}[a + b*\text{Cos}[c + d*x]]/(\text{Sqrt}[a + b]*\text{Sqrt}[\text{Cos}[c + d*x]])], -((a + b)/(a - b))]*\text{Sqrt}[(a*(1 - \text{Sec}[c + d*x]))/(a + b)]*\text{Sqrt}[(a*(1 + \text{Sec}[c + d*x]))/(a - b)]/(b*d)$

3.438. $\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

3.438.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*
(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

3.438.4 Maple [A] (verified)

Time = 11.98 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.25

method	result
default	$\frac{2B \left(F \left(\cot(dx+c) - \csc(dx+c), \sqrt{-\frac{a-b}{a+b}} \right) - 2\Pi \left(\cot(dx+c) - \csc(dx+c), -1, \sqrt{-\frac{a-b}{a+b}} \right) \right) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}} (1+\cos(dx+c))}{d\sqrt{a+\cos(dx+c)b} \sqrt{\cos(dx+c)}}$
parts	Expression too large to display

```
input int(cos(d*x+c)^(1/2)*(B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2),x,method=
_RETURNVERBOSE)
```

```
output 2*B/d*(EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))-2*EllipticPi(
cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2)))*(cos(d*x+c)/(1+cos(d*x+c))
)^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)/(a+cos(d*x+c)*b)^(1/
2)*(1+cos(d*x+c))/cos(d*x+c)^(1/2)
```

3.438. $\int \frac{\sqrt{\cos(c+dx)}(aB+bB \cos(c+dx))}{(a+b \cos(c+dx))^{3/2}} dx$

3.438.5 Fricas [F]

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(Bb\cos(dx+c)+Ba)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x,
algorithm="fricas")`

output `integral(B*sqrt(cos(d*x + c))/sqrt(b*cos(d*x + c) + a), x)`

3.438.6 Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = B \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx$$

input `integrate(cos(d*x+c)**(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2),x
)`

output `B*Integral(sqrt(cos(c + d*x))/sqrt(a + b*cos(c + d*x)), x)`

3.438.7 Maxima [F]

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(Bb\cos(dx+c)+Ba)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)
^(3/2), x)`

3.438.8 Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{(Bb\cos(dx+c)+Ba)\sqrt{\cos(dx+c)}}{(b\cos(dx+c)+a)^{3/2}} dx$$

input `integrate(cos(d*x+c)^(1/2)*(a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2),x,
algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(cos(d*x + c))/(b*cos(d*x + c) + a)
^(3/2), x)`

3.438.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}(aB+bB\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cos(c+dx)}(Ba+Bb\cos(c+dx))}{(a+b\cos(c+dx))^{3/2}} dx$$

input `int((cos(c + d*x)^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/
2),x)`

output `int((cos(c + d*x)^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))^(3/
2), x)`

3.439
$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$$

3.439.1 Optimal result	4085
3.439.2 Mathematica [A] (verified)	4085
3.439.3 Rubi [A] (verified)	4086
3.439.4 Maple [A] (verified)	4087
3.439.5 Fricas [F]	4088
3.439.6 Sympy [F]	4088
3.439.7 Maxima [F]	4088
3.439.8 Giac [F]	4089
3.439.9 Mupad [F(-1)]	4089

3.439.1 Optimal result

Integrand size = 38, antiderivative size = 110

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \frac{2\sqrt{a + b}B \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{ad}$$

output

```
2*B*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2), ((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d
```

3.439.2 Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.55

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \frac{4(a + b)B \cos^{\frac{3}{2}}(c + dx) \sqrt{-\frac{(a+b) \cot^2(\frac{1}{2}(c+dx))}{a-b}} \sqrt{\frac{(a+b \cos(c+dx)) \csc^2(\frac{1}{2}(c+dx))}{a}} \csc(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\sqrt{-\frac{(a+b) \cot^2(\frac{1}{2}(c+dx))}{a-b}}\right)\right)}{ad\sqrt{a + b \cos(c + dx)} \left(-\frac{(a+b) \cos(c+dx) \csc^2(\frac{1}{2}(c+dx))}{a}\right)^{3/2}}$$

input

```
Integrate[(a*B + b*B*Cos[c + d*x])/(Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)), x]
```

3.439.
$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$$

output $(-4*(a + b)*B*\text{Cos}[c + d*x]^{(3/2)}*\text{Sqrt}[-((a + b)*\text{Cot}[(c + d*x)/2]^2)/(a - b)))*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])*\text{Csc}[(c + d*x)/2]^2/a]*\text{Csc}[c + d*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[-(a + b*\text{Cos}[c + d*x])/(a*(-1 + \text{Cos}[c + d*x]))]], (2*a)/(a - b)]/(a*d*\text{Sqrt}[a + b*\text{Cos}[c + d*x])*(-((a + b)*\text{Cos}[c + d*x]*\text{Csc}[(c + d*x)/2]^2/a))^{(3/2)})$

3.439.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.079$, Rules used = {2011, 3042, 3295}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$$

↓ 2011

$$B \int \frac{1}{\sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$B \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

↓ 3295

$$\frac{2B\sqrt{a + b} \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b} \cos(c + dx)}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right)}{ad}$$

input $\text{Int}[(a*B + b*B*\text{Cos}[c + d*x])]/(\text{Sqrt}[\text{Cos}[c + d*x]]*(a + b*\text{Cos}[c + d*x])^{(3/2)}), x]$

output $(2*\text{Sqrt}[a + b]*B*\text{Cot}[c + d*x]*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[a + b*\text{Cos}[c + d*x]]]/(\text{Sqrt}[a + b]*\text{Sqrt}[\text{Cos}[c + d*x]]), -(a + b)/(a - b))*\text{Sqrt}[(a*(1 - \text{Sec}[c + d*x]))/(a + b)]*\text{Sqrt}[(a*(1 + \text{Sec}[c + d*x]))/(a - b)]/(a*d)$

3.439.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*
(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

3.439.4 Maple [A] (verified)

Time = 12.30 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.02

method	result	size
default	$-\frac{2B(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}F\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}}{d\sqrt{a+\cos(dx+c)}b\sqrt{\cos(dx+c)}}$	112
parts	Expression too large to display	1535

```
input int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(3/2),x,method=
_RETURNVERBOSE)
```

```
output -2*B/d*(1+cos(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/(a+cos(d*x+c)*b)^(
1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*
b)/(1+cos(d*x+c))/(a+b))^(1/2)/cos(d*x+c)^(1/2)
```

3.439.5 Fricas [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="fricas")`

output `integral(sqrt(b*cos(d*x + c) + a)*B*sqrt(cos(d*x + c))/(b*cos(d*x + c)^2 +
a*cos(d*x + c)), x)`

3.439.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \cos(c + dx)} \sqrt{\cos(c + dx)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(3/2),x
)`

output `B*Integral(1/(sqrt(a + b*cos(c + d*x))*sqrt(cos(c + d*x))), x)`

3.439.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*sqrt(cos(d*
x + c))), x)`

3.439.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sqrt{\cos(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*sqrt(cos(d*
x + c))), x)`

3.439.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Ba + Bb \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(3/2
)),x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(1/2)*(a + b*cos(c + d*x))^(3/2
)), x)`

$$3.440 \quad \int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$$

3.440.1 Optimal result	4090
3.440.2 Mathematica [A] (verified)	4090
3.440.3 Rubi [A] (verified)	4091
3.440.4 Maple [B] (verified)	4093
3.440.5 Fracas [F]	4094
3.440.6 Sympy [F]	4094
3.440.7 Maxima [F]	4095
3.440.8 Giac [F]	4095
3.440.9 Mupad [F(-1)]	4095

3.440.1 Optimal result

Integrand size = 38, antiderivative size = 226

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \frac{2(a - b)\sqrt{a + b}B \cot(c + dx)E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right)}{a^2 d} - \frac{2\sqrt{a + b}B \cot(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{ad}$$

```
output 2*(a-b)*B*cot(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d-2*B*cot(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d
```

3.440.2 Mathematica [A] (verified)

Time = 1.41 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.94

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \frac{2B \left(- \left((a + b) \sqrt{\cos(c + dx)} \sqrt{1 + \cos(c + dx)} \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} \right) \right)}{\dots}$$

```
input Integrate[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)),x]
```

3.440. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$

output $(2*B*(-((a + b)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[1 + \text{Cos}[c + d*x]]*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/((a + b)*(1 + \text{Cos}[c + d*x]))]*\text{EllipticE}[\text{ArcSin}[\text{Tan}[(c + d*x)/2]], (-a + b)/(a + b)]) + a*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[1 + \text{Cos}[c + d*x]]*\text{Sqrt}[(a + b*\text{Cos}[c + d*x])/((a + b)*(1 + \text{Cos}[c + d*x]))]*\text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + d*x)/2]], (-a + b)/(a + b)] + (a + b*\text{Cos}[c + d*x])* \text{Tan}[(c + d*x)/2]))/(a*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a + b*\text{Cos}[c + d*x]])$

3.440.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2011, 3042, 3280, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$$

$$\downarrow 2011$$

$$B \int \frac{1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

$$\downarrow 3042$$

$$B \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow 3280$$

$$B \left(\int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \right)$$

$$\downarrow 3042$$

$$B \left(\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right)$$

$$\downarrow 3295$$

$$B \left(\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{2\sqrt{a + b} \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} \text{EllipticE}(\arcsin(\frac{\sqrt{a + b} \cos(c + dx)}{\sqrt{a + b} \sqrt{\cos(c + dx)}}))}{a d} \right)$$

↓ 3473

$$B \left(\frac{2(a - b)\sqrt{a + b} \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} E\left(\arcsin\left(\frac{\sqrt{a + b} \cos(c + dx)}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right) \mid -\frac{a + b}{a - b}\right)}{a^2 d} - \frac{2\sqrt{a + b} \cot(c + dx)}{a d} \right)$$

```
input Int[(a*B + b*B*Cos[c + d*x])/(Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)),x]
```

```
output B*((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a^2*d) - (2*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d))
```

3.440.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3280 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[1/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[b/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.440. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] :> Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]
```

3.440.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 658 vs. 2(210) = 420.

Time = 17.06 (sec) , antiderivative size = 659, normalized size of antiderivative = 2.92

method	result
default	$-\frac{2B \left(-\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \sqrt{\frac{(\csc^2(dx+c)a(1-\cos(dx+c))^2 - (\csc^2(dx+c)b(1-\cos(dx+c))^2+a+b)}{a+b}} \right) F\left(\cot(dx+c) - \csc(dx+c), \frac{a+b}{a+b}\right)}{a+b}$
parts	Expression too large to display

```
input int((B*a+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)
```

3.440.
$$\int \frac{aB+bB \cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b \cos(c+dx))^{3/2}} dx$$

output
$$-2*B/d*(-(-\csc(dx+c)^2*(1-\cos(dx+c))^2+1)^{1/2}*((\csc(dx+c)^2*a*(1-\cos(dx+c))^2-\csc(dx+c)^2*b*(1-\cos(dx+c))^2+a+b)/(a+b))^{1/2}*EllipticF(\cot(dx+c)-\csc(dx+c),(-a-b)/(a+b))^{1/2})*a+(-\csc(dx+c)^2*(1-\cos(dx+c))^2+1)^{1/2}*((\csc(dx+c)^2*a*(1-\cos(dx+c))^2-\csc(dx+c)^2*b*(1-\cos(dx+c))^2+a+b)/(a+b))^{1/2}*EllipticE(\cot(dx+c)-\csc(dx+c),(-a-b)/(a+b))^{1/2})*a+(-\csc(dx+c)^2*(1-\cos(dx+c))^2+1)^{1/2}*((\csc(dx+c)^2*a*(1-\cos(dx+c))^2-\csc(dx+c)^2*b*(1-\cos(dx+c))^2+a+b)/(a+b))^{1/2}*EllipticE(\cot(dx+c)-\csc(dx+c),(-a-b)/(a+b))^{1/2})*b+\csc(dx+c)^3*(1-\cos(dx+c))^3*a-\csc(dx+c)^3*(1-\cos(dx+c))^3*b+a*(\csc(dx+c)-\cot(dx+c))+b*(\csc(dx+c)-\cot(dx+c)))*((\csc(dx+c)^2*a*(1-\cos(dx+c))^2-\csc(dx+c)^2*b*(1-\cos(dx+c))^2+a+b)/(\csc(dx+c)^2*(1-\cos(dx+c))^2+1))^{1/2}*(\csc(dx+c)^2*(1-\cos(dx+c))^2-1)/(\csc(dx+c)^2*a*(1-\cos(dx+c))^2-\csc(dx+c)^2*b*(1-\cos(dx+c))^2+a+b)/(\csc(dx+c)^2*(1-\cos(dx+c))^2+1)/(-(\csc(dx+c)^2*(1-\cos(dx+c))^2-1)/(\csc(dx+c)^2*(1-\cos(dx+c))^2+1))^{3/2}/a$$

3.440.5 Fracas [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(dx+c))/cos(dx+c)^(3/2)/(a+b*cos(dx+c))^(3/2),x,
algorithm="fricas")`

output `integral(sqrt(b*cos(dx + c) + a)*B*sqrt(cos(dx + c))/(b*cos(dx + c)^3 +
a*cos(dx + c)^2), x)`

3.440.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a*B+b*B*cos(dx+c))/cos(dx+c)**(3/2)/(a+b*cos(dx+c))**(3/2),x
)`

output `B*Integral(1/(sqrt(a + b*cos(c + dx))*cos(c + dx)**(3/2)), x)`

3.440.
$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$$

3.440.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)
^(3/2)), x)`

3.440.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{3}{2}} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/cos(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*cos(d*x + c)
^(3/2)), x)`

3.440.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx = \int \frac{Ba + Bb \cos(c + dx)}{\cos(c + dx)^{3/2} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(3/2)
)),x)`

output `int((B*a + B*b*cos(c + d*x))/(cos(c + d*x)^(3/2)*(a + b*cos(c + d*x))^(3/2)
)), x)`

3.440. $\int \frac{aB + bB \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$

$$3.441 \quad \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{2 + 3 \cos(c + dx)}} dx$$

3.441.1 Optimal result	4096
3.441.2 Mathematica [F]	4096
3.441.3 Rubi [A] (verified)	4097
3.441.4 Maple [B] (verified)	4098
3.441.5 Fricas [F]	4098
3.441.6 Sympy [F]	4099
3.441.7 Maxima [F]	4099
3.441.8 Giac [F]	4099
3.441.9 Mupad [F(-1)]	4100

3.441.1 Optimal result

Integrand size = 33, antiderivative size = 72

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{2 + 3 \cos(c + dx)}} dx = -\frac{\cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{2 + 3 \cos(c + dx)}}{\sqrt{5} \sqrt{\cos(c + dx)}}\right) \middle| 5\right) \sqrt{-1 - \sec(c + dx)} \sqrt{1 - \sec(c + dx)}}{d}$$

```
output -cot(d*x+c)*EllipticE(1/5*(2+3*cos(d*x+c))^(1/2)*5^(1/2)/cos(d*x+c)^(1/2),
5^(1/2))*(-1-sec(d*x+c))^(1/2)*(1-sec(d*x+c))^(1/2)/d
```

3.441.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{2 + 3 \cos(c + dx)}} dx = \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{2 + 3 \cos(c + dx)}} dx$$

```
input Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[2 + 3*Cos[c + d*x]]),x]
```

```
output Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[2 + 3*Cos[c + d*x]]), x]
```

3.441.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{3\cos(c+dx)+2}} dx$$

↓ 3042

$$\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{3\sin(c+dx+\frac{\pi}{2})+2}} dx$$

↓ 3473

$$\frac{\cot(c+dx)\sqrt{-\sec(c+dx)-1}\sqrt{1-\sec(c+dx)}E\left(\arcsin\left(\frac{\sqrt{3\cos(c+dx)+2}}{\sqrt{5}\sqrt{\cos(c+dx)}}\right)\right)}{d}$$

input `Int[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[2 + 3*Cos[c + d*x]]),x]`

output `-((Cot[c + d*x]*EllipticE[ArcSin[Sqrt[2 + 3*Cos[c + d*x]]/(Sqrt[5]*Sqrt[Cos[c + d*x]])], 5]*Sqrt[-1 - Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]])/d)`

3.441.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

3.441. $\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{2+3\cos(c+dx)}} dx$

3.441.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 474 vs. 2(64) = 128.

Time = 11.35 (sec) , antiderivative size = 475, normalized size of antiderivative = 6.60

method	result
parts	$\frac{\left(-2\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\sqrt{-5(\csc^2(dx+c))(1-\cos(dx+c))^2+25}F\left(\cot(dx+c)-\csc(dx+c),\frac{\sqrt{5}}{5}\right)+5\sqrt{-(\csc^2(dx+c))}\right)}{5d\left(\csc^2(dx+c)\right)}$
default	$\frac{5E\left(\cot(dx+c)-\csc(dx+c),\frac{\sqrt{5}}{5}\right)\sqrt{2}\sqrt{10}\sqrt{\frac{2+3\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))-4\sqrt{2}\sqrt{10}F\left(\cot(dx+c)-\csc(dx+c),\frac{\sqrt{5}}{5}\right)\sqrt{\frac{2}{1+\cos(dx+c)}}}{5d\left(\csc^2(dx+c)\right)}$

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2+3*cos(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/5/d*(-2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(-5*csc(d*x+c)^2*(1-cos(d*x+c))^2+25)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),1/5*5^(1/2))+5*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(-5*csc(d*x+c)^2*(1-cos(d*x+c))^2+25)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),1/5*5^(1/2))-5*csc(d*x+c)^3*(1-cos(d*x+c))^3+25*csc(d*x+c)-25*cot(d*x+c))*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-5)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-5)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(-(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(3/2)-1/5/d*(1+cos(d*x+c))*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/(2+3*cos(d*x+c))^(1/2)*10^(1/2)*((2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),1/5*5^(1/2))/cos(d*x+c)^(1/2)`

3.441.5 Fricas [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{2 + 3 \cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{3 \cos(dx + c) + 2 \cos^2(dx + c)}^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2+3*cos(d*x+c))^(1/2),x,algorithm="fricas")`

output `integral(sqrt(3*cos(d*x + c) + 2)*(cos(d*x + c) + 1)*sqrt(cos(d*x + c))/(3*cos(d*x + c)^3 + 2*cos(d*x + c)^2), x)`

3.441. $\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{2+3\cos(c+dx)}} dx$

3.441.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{2 + 3\cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{3\cos(c + dx) + 2}\cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(2+3*cos(d*x+c))**(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(3*cos(c + d*x) + 2)*cos(c + d*x)**(3/2)), x)`

3.441.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{2 + 3\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{3\cos(dx + c) + 2}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2+3*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(3*cos(d*x + c) + 2)*cos(d*x + c)^(3/2)), x)`

3.441.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{2 + 3\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{3\cos(dx + c) + 2}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2+3*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(3*cos(d*x + c) + 2)*cos(d*x + c)^(3/2)), x)`

3.441.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{2 + 3 \cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{\frac{3}{2}} \sqrt{3 \cos(c + dx) + 2}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(3*cos(c + d*x) + 2)^(1/2)),x)`output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(3*cos(c + d*x) + 2)^(1/2)), x)`

$$3.442 \quad \int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{-2+3\cos(c+dx)}} dx$$

3.442.1 Optimal result	4101
3.442.2 Mathematica [F]	4101
3.442.3 Rubi [A] (verified)	4102
3.442.4 Maple [B] (verified)	4103
3.442.5 Fricas [F]	4103
3.442.6 Sympy [F]	4104
3.442.7 Maxima [F]	4104
3.442.8 Giac [F]	4104
3.442.9 Mupad [F(-1)]	4105

3.442.1 Optimal result

Integrand size = 33, antiderivative size = 70

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx$$

$$= -\frac{\sqrt{5} \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{-2+3\cos(c+dx)}}{\sqrt{\cos(c+dx)}}\right) \middle| \frac{1}{5}\right) \sqrt{-1 + \sec(c + dx)} \sqrt{1 + \sec(c + dx)}}{d}$$

output `-cot(d*x+c)*EllipticE((-2+3*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),1/5*5^(1/2))`
`*5^(1/2)*(-1+sec(d*x+c))^(1/2)*(1+sec(d*x+c))^(1/2)/d`

3.442.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx = \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx$$

input `Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[-2 + 3*Cos[c + d*x])`
`], x]`

output `Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[-2 + 3*Cos[c + d*x])`
`], x]`

$$3.442. \quad \int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{-2+3\cos(c+dx)}} dx$$

3.442.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx)\sqrt{3\cos(c + dx) - 2}} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2}\sqrt{3\sin(c + dx + \frac{\pi}{2}) - 2}} dx$$

↓ 3473

$$\frac{\sqrt{5}\cot(c + dx)\sqrt{\sec(c + dx) - 1}\sqrt{\sec(c + dx) + 1}E\left(\arcsin\left(\frac{\sqrt{3\cos(c + dx) - 2}}{\sqrt{\cos(c + dx)}}\right)\middle|\frac{1}{5}\right)}{d}$$

input `Int[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[-2 + 3*Cos[c + d*x]]),x]`

output `-((Sqrt[5]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[-2 + 3*Cos[c + d*x]]/Sqrt[Cos[c + d*x]]], 1/5]*Sqrt[-1 + Sec[c + d*x]]*Sqrt[1 + Sec[c + d*x]])/d)`

3.442.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

3.442. $\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx$

3.442.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 461 vs. 2(61) = 122.

Time = 9.83 (sec) , antiderivative size = 462, normalized size of antiderivative = 6.60

method	result
parts	$\frac{\left(-2\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\sqrt{-5(\csc^2(dx+c))(1-\cos(dx+c))^2+1}F(\cot(dx+c)-\csc(dx+c),\sqrt{5})-\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\right)}{d\left(5(\csc^2(dx+c))(1-\cos(dx+c))^2+1\right)}$
default	$-\frac{E(\cot(dx+c)-\csc(dx+c),\sqrt{5})\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{-2+3\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))+4\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{-2+3\cos(dx+c)}{1+\cos(dx+c)}}F(\cot(dx+c)-\csc(dx+c),\sqrt{5})}{\cos^2(dx+c)}$

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2+3*cos(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*(-2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(-5*csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),5^(1/2))-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(-5*csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),5^(1/2))+5*csc(d*x+c)^3*(1-cos(d*x+c))^3-csc(d*x+c)+cot(d*x+c))*(-5*csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(5*csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(3/2)-2/d*(1+cos(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/(-2+3*cos(d*x+c))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),5^(1/2))/cos(d*x+c)^(1/2)`

3.442.5 Fracas [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{3\cos(dx + c) - 2}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2+3*cos(d*x+c))^(1/2),x,algorith="fricas")`

output `integral(sqrt(3*cos(d*x + c) - 2)*(cos(d*x + c) + 1)*sqrt(cos(d*x + c))/(3*cos(d*x + c)^3 - 2*cos(d*x + c)^2), x)`

3.442.
$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx$$

3.442.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{3\cos(c + dx) - 2}\cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(-2+3*cos(d*x+c))**(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(3*cos(c + d*x) - 2)*cos(c + d*x)**(3/2)), x)`

3.442.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{3\cos(dx + c) - 2}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2+3*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(3*cos(d*x + c) - 2)*cos(d*x + c)^(3/2)), x)`

3.442.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{-2 + 3\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{3\cos(dx + c) - 2}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2+3*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(3*cos(d*x + c) - 2)*cos(d*x + c)^(3/2)), x)`

3.442.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-2 + 3 \cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{\frac{3}{2}} \sqrt{3 \cos(c + dx) - 2}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(3*cos(c + d*x) - 2)^(1/2)),x)`output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(3*cos(c + d*x) - 2)^(1/2)), x)`

3.443
$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.443.1 Optimal result 4106
 3.443.2 Mathematica [F] 4106
 3.443.3 Rubi [A] (verified) 4107
 3.443.4 Maple [B] (verified) 4108
 3.443.5 Fricas [F] 4109
 3.443.6 Sympy [F] 4109
 3.443.7 Maxima [F] 4110
 3.443.8 Giac [F] 4110
 3.443.9 Mupad [F(-1)] 4110

3.443.1 Optimal result

Integrand size = 33, antiderivative size = 93

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \frac{\sqrt{5} \sqrt{-\cos(c + dx)} \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{2 - 3 \cos(c + dx)}}{\sqrt{-\cos(c + dx)}}\right) \middle| \frac{1}{5}\right) \sqrt{-1 + \sec(c + dx)} \sqrt{1 + \sec(c + dx)}}{d}$$

output `csc(d*x+c)*EllipticE((2-3*cos(d*x+c))^(1/2)/(-cos(d*x+c))^(1/2),1/5*5^(1/2))*5^(1/2)*(-cos(d*x+c))^(1/2)*cos(d*x+c)^(1/2)*(-1+sec(d*x+c))^(1/2)*(1+sec(d*x+c))^(1/2)/d`

3.443.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `Integrate[(1 + Cos[c + d*x])/(Sqrt[2 - 3*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]`

output `Integrate[(1 + Cos[c + d*x])/(Sqrt[2 - 3*Cos[c + d*x]]*Cos[c + d*x]^(3/2)), x]`

3.443.
$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.443.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {3042, 3474, 3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)+1}{\sqrt{2-3\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sqrt{2-3\sin(c+dx+\frac{\pi}{2})}\sin(c+dx+\frac{\pi}{2})^{3/2}} dx \\
 & \quad \downarrow \text{3474} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\cos(c+dx)+1}{\sqrt{2-3\cos(c+dx)}(-\cos(c+dx))^{3/2}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sqrt{2-3\sin(c+dx+\frac{\pi}{2})}(-\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3473} \\
 & \frac{\sqrt{5}\sqrt{-\cos(c+dx)}\sqrt{\cos(c+dx)}\csc(c+dx)\sqrt{\sec(c+dx)-1}\sqrt{\sec(c+dx)+1}E\left(\arcsin\left(\frac{\sqrt{2-3\cos(c+dx)}}{\sqrt{-\cos(c+dx)}}\right)\middle|\frac{1}{5}\right)}{d}
 \end{aligned}$$

input `Int[(1 + Cos[c + d*x])/(Sqrt[2 - 3*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]`

output `(Sqrt[5]*Sqrt[-Cos[c + d*x]]*Sqrt[Cos[c + d*x]]*Csc[c + d*x]*EllipticE[ArcSin[Sqrt[2 - 3*Cos[c + d*x]]/Sqrt[-Cos[c + d*x]]], 1/5]*Sqrt[-1 + Sec[c + d*x]]*Sqrt[1 + Sec[c + d*x]])/d`

3.443. $\int \frac{1+\cos(c+dx)}{\sqrt{2-3\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$

3.443.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3474 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-Sqrt[(-b)*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]] Int[(A + B*Sin[e + f*x])/((-b)*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && NegQ[(c + d)/b]`

3.443.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 473 vs. 2(80) = 160.

Time = 11.34 (sec) , antiderivative size = 474, normalized size of antiderivative = 5.10

method	result
default	$\frac{\left(E\left(\cot(dx+c)-\operatorname{csc}(dx+c),\sqrt{5}\right)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{-2+3\cos(dx+c)}{1+\cos(dx+c)}}\left(\cos^2(dx+c)+4\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{-2+3\cos(dx+c)}{1+\cos(dx+c)}}F\left(\cot(dx+c)-\operatorname{csc}(dx+c),\sqrt{5}\right)\right)\right)}{d(-2+3\cos(dx+c))\sqrt{\cos(dx+c)}}$
parts	$\frac{2(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{2-3\cos(dx+c)}\sqrt{\frac{-2+3\cos(dx+c)}{1+\cos(dx+c)}}F\left(\cot(dx+c)-\operatorname{csc}(dx+c),\sqrt{5}\right)}{d(-2+3\cos(dx+c))\sqrt{\cos(dx+c)}} - \frac{\left(-2\sqrt{-(\operatorname{csc}^2(dx+c))(1-\cos(dx+c))}\right)}{d(-2+3\cos(dx+c))\sqrt{\cos(dx+c)}}$

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2-3*cos(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

3.443.
$$\int \frac{1+\cos(c+dx)}{\sqrt{2-3\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$$

output `1/d*(EllipticE(cot(d*x+c)-csc(d*x+c),5^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)^2+4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),5^(1/2))*cos(d*x+c)^2+2*EllipticE(cot(d*x+c)-csc(d*x+c),5^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)+8*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),5^(1/2))*cos(d*x+c)+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),5^(1/2))+4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((-2+3*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),5^(1/2))+3*cos(d*x+c)*sin(d*x+c)-2*sin(d*x+c))*(2-3*cos(d*x+c))^(1/2)/(-2+3*cos(d*x+c))/cos(d*x+c)^(1/2)/(1+cos(d*x+c))`

3.443.5 Fracas [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-3 \cos(dx + c) + 2 \cos(dx + c)}^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2-3*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `integral(-(cos(d*x + c) + 1)*sqrt(-3*cos(d*x + c) + 2)*sqrt(cos(d*x + c))/(3*cos(d*x + c)^3 - 2*cos(d*x + c)^2), x)`

3.443.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(2-3*cos(d*x+c))**(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(2 - 3*cos(c + d*x))*cos(c + d*x)**(3/2)), x)`

3.443. $\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$

3.443.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-3 \cos(dx + c) + 2} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2-3*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-3*cos(d*x + c) + 2)*cos(d*x + c)^(3/2)), x)`

3.443.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-3 \cos(dx + c) + 2} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(2-3*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-3*cos(d*x + c) + 2)*cos(d*x + c)^(3/2)), x)`

3.443.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\sqrt{2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{3/2} \sqrt{2 - 3 \cos(c + dx)}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(2 - 3*cos(c + d*x))^(1/2)),x)`

output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(2 - 3*cos(c + d*x))^(1/2)), x)`

$$3.444 \quad \int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.444.1 Optimal result	4111
3.444.2 Mathematica [F]	4111
3.444.3 Rubi [A] (verified)	4112
3.444.4 Maple [B] (verified)	4113
3.444.5 Fracas [F]	4114
3.444.6 Sympy [F]	4114
3.444.7 Maxima [F]	4115
3.444.8 Giac [F]	4115
3.444.9 Mupad [F(-1)]	4115

3.444.1 Optimal result

Integrand size = 33, antiderivative size = 95

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \frac{\sqrt{-\cos(c + dx)} \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{-2 - 3 \cos(c + dx)}}{\sqrt{5} \sqrt{-\cos(c + dx)}}\right) \middle| 5\right) \sqrt{-1 - \sec(c + dx)} \sqrt{1 - \sec(c + dx)}}{d}$$

```
output csc(d*x+c)*EllipticE(1/5*(-2-3*cos(d*x+c))^(1/2)*5^(1/2)/(-cos(d*x+c))^(1/2),5^(1/2))*(-cos(d*x+c))^(1/2)*cos(d*x+c)^(1/2)*(-1-sec(d*x+c))^(1/2)*(1-sec(d*x+c))^(1/2)/d
```

3.444.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

```
input Integrate[(1 + Cos[c + d*x])/(Sqrt[-2 - 3*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]
```

```
output Integrate[(1 + Cos[c + d*x])/(Sqrt[-2 - 3*Cos[c + d*x]]*Cos[c + d*x]^(3/2)), x]
```

$$3.444. \quad \int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.444.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {3042, 3474, 3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)+1}{\sqrt{-3\cos(c+dx)-2\cos^{\frac{3}{2}}(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sqrt{-3\sin(c+dx+\frac{\pi}{2})-2\sin(c+dx+\frac{\pi}{2})^{3/2}}} dx \\
 & \quad \downarrow \text{3474} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\cos(c+dx)+1}{\sqrt{-3\cos(c+dx)-2(-\cos(c+dx))^{3/2}}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sqrt{-3\sin(c+dx+\frac{\pi}{2})-2(-\sin(c+dx+\frac{\pi}{2}))^{3/2}}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3473} \\
 & \frac{\sqrt{-\cos(c+dx)}\sqrt{\cos(c+dx)}\csc(c+dx)\sqrt{-\sec(c+dx)-1}\sqrt{1-\sec(c+dx)}E\left(\arcsin\left(\frac{\sqrt{-3\cos(c+dx)-2}}{\sqrt{5}\sqrt{-\cos(c+dx)}}\right)\middle|5\right)}{d}
 \end{aligned}$$

input `Int[(1 + Cos[c + d*x])/(Sqrt[-2 - 3*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]`

output `(Sqrt[-Cos[c + d*x]]*Sqrt[Cos[c + d*x]]*Csc[c + d*x]*EllipticE[ArcSin[Sqrt[-2 - 3*Cos[c + d*x]]/(Sqrt[5]*Sqrt[-Cos[c + d*x]])], 5]*Sqrt[-1 - Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]])/d`

3.444. $\int \frac{1+\cos(c+dx)}{\sqrt{-2-3\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$

3.444.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d),
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3474 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-Sqrt
[(-b)*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]] Int[(A + B*Sin[e + f*x])/((-b)*
Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{b, c, d, e,
f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && NegQ[(c + d)/b]
```

3.444.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 315 vs. 2(83) = 166.

Time = 12.19 (sec) , antiderivative size = 316, normalized size of antiderivative = 3.33

method	result
default	$-\frac{\left(-\sqrt{5}\sqrt{10}E\left(\frac{\csc(dx+c)-\cot(dx+c)\sqrt{5}}{5},\sqrt{5}\right)\sqrt{2}\sqrt{\frac{2+3\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))-2\sqrt{5}\sqrt{10}E\left(\frac{\csc(dx+c)-\cot(dx+c)}{5}\right)\sqrt{5}\right)}{5d(2+3\cos(dx+c))\sqrt{\cos(dx+c)}}$
parts	$-\frac{(1+\cos(dx+c))\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{-2-3\cos(dx+c)}\sqrt{10}\sqrt{\frac{2+3\cos(dx+c)}{1+\cos(dx+c)}}F\left(\frac{\csc(dx+c)-\cot(dx+c)\sqrt{5}}{5},\sqrt{5}\right)\sqrt{5}}{5d(2+3\cos(dx+c))\sqrt{\cos(dx+c)}} + \frac{\left(\sqrt{5}\sqrt{-5(\cos(dx+c)-1)}\right)}{5d(2+3\cos(dx+c))\sqrt{\cos(dx+c)}}$

```
input int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2-3*cos(d*x+c))^(1/2), x, method=_RETU
RNVERBOSE)
```

3.444.
$$\int \frac{1+\cos(c+dx)}{\sqrt{-2-3\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$$

output
$$\begin{aligned} & -1/10/d*(-5^{(1/2)}*10^{(1/2)}*EllipticE(1/5*(csc(d*x+c)-cot(d*x+c))*5^{(1/2)},5 \\ & ^{(1/2)})*2^{(1/2)*((2+3*cos(d*x+c))/(1+cos(d*x+c)))^{(1/2)}*(cos(d*x+c)/(1+cos \\ & (d*x+c)))^{(1/2)}*cos(d*x+c)^2-2*5^{(1/2)}*10^{(1/2)}*EllipticE(1/5*(csc(d*x+c)- \\ & cot(d*x+c))*5^{(1/2)},5^{(1/2)})*2^{(1/2)*((2+3*cos(d*x+c))/(1+cos(d*x+c)))^{(1/2)} \\ & *(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*cos(d*x+c)-2^{(1/2)}*EllipticE(1/5*(csc \\ & (d*x+c)-cot(d*x+c))*5^{(1/2)},5^{(1/2)}*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*((2 \\ & +3*cos(d*x+c))/(1+cos(d*x+c)))^{(1/2)}*10^{(1/2)}*5^{(1/2)}+30*cos(d*x+c)*sin(d* \\ & x+c)+20*sin(d*x+c))*(-2-3*cos(d*x+c))^{(1/2)}/(2+3*cos(d*x+c))/cos(d*x+c)^{(1 \\ & /2)}/(1+cos(d*x+c)) \end{aligned}$$

3.444.5 Fricas [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-3 \cos(dx + c) - 2} \cos^{\frac{3}{2}}(dx + c)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2-3*cos(d*x+c))^(1/2),x, algorith="fricas")`

output `integral(-(cos(d*x + c) + 1)*sqrt(-3*cos(d*x + c) - 2)*sqrt(cos(d*x + c))/(3*cos(d*x + c)^3 + 2*cos(d*x + c)^2), x)`

3.444.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{-3 \cos(c + dx) - 2} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(-2-3*cos(d*x+c))**(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(-3*cos(c + d*x) - 2)*cos(c + d*x)**(3/2)), x)`

3.444.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-3 \cos(dx + c) - 2} \cos^{\frac{3}{2}}(dx + c)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2-3*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-3*cos(d*x + c) - 2)*cos(d*x + c)^(3/2)), x)`

3.444.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-3 \cos(dx + c) - 2} \cos^{\frac{3}{2}}(dx + c)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-2-3*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-3*cos(d*x + c) - 2)*cos(d*x + c)^(3/2)), x)`

3.444.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{3/2} \sqrt{-3 \cos(c + dx) - 2}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(- 3*cos(c + d*x) - 2)^(1/2)),x)`

output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(- 3*cos(c + d*x) - 2)^(1/2)), x)`

3.444. $\int \frac{1 + \cos(c + dx)}{\sqrt{-2 - 3 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$

$$3.445 \quad \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx$$

3.445.1 Optimal result	4116
3.445.2 Mathematica [F]	4116
3.445.3 Rubi [A] (verified)	4117
3.445.4 Maple [B] (verified)	4118
3.445.5 Fricas [F]	4118
3.445.6 Sympy [F]	4119
3.445.7 Maxima [F]	4119
3.445.8 Giac [F]	4119
3.445.9 Mupad [F(-1)]	4120

3.445.1 Optimal result

Integrand size = 33, antiderivative size = 72

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx$$

$$= \frac{2 \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{3 + 2 \cos(c + dx)}}{\sqrt{5} \sqrt{\cos(c + dx)}}\right) \middle| -5\right) \sqrt{1 - \sec(c + dx)} \sqrt{1 + \sec(c + dx)}}{3d}$$

output `2/3*cot(d*x+c)*EllipticE(1/5*(3+2*cos(d*x+c))^(1/2)*5^(1/2)/cos(d*x+c)^(1/2), I*5^(1/2))*(1-sec(d*x+c))^(1/2)*(1+sec(d*x+c))^(1/2)/d`

3.445.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx = \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx$$

input `Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[3 + 2*Cos[c + d*x]]), x]`

output `Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[3 + 2*Cos[c + d*x]]), x]`

3.445. $\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx$

3.445.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c + dx) + 1}{\cos^{\frac{3}{2}}(c + dx)\sqrt{2\cos(c + dx) + 3}} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2}\sqrt{2\sin(c + dx + \frac{\pi}{2}) + 3}} dx$$

↓ 3473

$$\frac{2 \cot(c + dx)\sqrt{1 - \sec(c + dx)}\sqrt{\sec(c + dx) + 1}E\left(\arcsin\left(\frac{\sqrt{2\cos(c+dx)+3}}{\sqrt{5}\sqrt{\cos(c+dx)}}\right)\right) - 5}{3d}$$

input `Int[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[3 + 2*Cos[c + d*x]]),x]`

output `(2*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[3 + 2*Cos[c + d*x]]/(Sqrt[5]*Sqrt[Cos[c + d*x]])], -5]*Sqrt[1 - Sec[c + d*x]]*Sqrt[1 + Sec[c + d*x]])/(3*d)`

3.445.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

3.445. $\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{3+2\cos(c+dx)}} dx$

3.445.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 476 vs. 2(65) = 130.

Time = 12.36 (sec) , antiderivative size = 477, normalized size of antiderivative = 6.62

method	result
parts	$\frac{2\left(-3\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\sqrt{5(\csc^2(dx+c))(1-\cos(dx+c))^2+25}F\left(\cot(dx+c)-\csc(dx+c),\frac{i\sqrt{5}}{5}\right)+5\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\right)}{15d\left(\csc(dx+c)\sqrt{1-\cos(dx+c)}\sqrt{5(\csc^2(dx+c))(1-\cos(dx+c))^2+25}\right)}$
default	$-\frac{6\sqrt{2}F\left(\cot(dx+c)-\csc(dx+c),\frac{i\sqrt{5}}{5}\right)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{3+2\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{10}(\cos^2(dx+c))-5\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{3+2\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{10}}{15d\left(\csc(dx+c)\sqrt{1-\cos(dx+c)}\sqrt{5(\csc^2(dx+c))(1-\cos(dx+c))^2+25}\right)}$

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3+2*cos(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output
$$-\frac{2}{15} \frac{1}{d} \frac{(-3(-\csc(dx+c))^2(1-\cos(dx+c))^2+1)^{1/2} (5\csc(dx+c))^2(1-\cos(dx+c))^2+25)^{1/2} \text{EllipticF}(\cot(dx+c)-\csc(dx+c), 1/5 I \sqrt{5}^{1/2}) + 5(-\csc(dx+c))^2(1-\cos(dx+c))^2+1)^{1/2} (5\csc(dx+c))^2(1-\cos(dx+c))^2+25)^{1/2} \text{EllipticE}(\cot(dx+c)-\csc(dx+c), 1/5 I \sqrt{5}^{1/2}) + 5\csc(dx+c)^3(1-\cos(dx+c))^3+25\csc(dx+c)-25\cot(dx+c)) \left(\frac{\csc(dx+c)^2(1-\cos(dx+c))^2+5}{\csc(dx+c)^2(1-\cos(dx+c))^2+1} \right)^{1/2} \frac{\csc(dx+c)^2(1-\cos(dx+c))^2-1}{\csc(dx+c)^2(1-\cos(dx+c))^2+5} \frac{1}{\csc(dx+c)^2(1-\cos(dx+c))^2+1} \left(-\frac{\csc(dx+c)^2(1-\cos(dx+c))^2-1}{\csc(dx+c)^2(1-\cos(dx+c))^2+1} \right)^{3/2} - \frac{1}{5} \frac{1}{d} (1+\cos(dx+c))^2)^{1/2} \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \frac{1}{(3+2\cos(dx+c))^{1/2}} \frac{10^{1/2} \left(\frac{3+2\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \text{EllipticF}(\cot(dx+c)-\csc(dx+c), 1/5 I \sqrt{5}^{1/2})}{\cos(dx+c)^{1/2}}$$

3.445.5 Fracas [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{2} \cos(dx + c) \sqrt{3 \cos(dx + c) + 2}}^{\frac{3}{2}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3+2*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `integral(sqrt(2*cos(d*x + c) + 3)*(cos(d*x + c) + 1)*sqrt(cos(d*x + c))/(2*cos(d*x + c)^3 + 3*cos(d*x + c)^2), x)`

3.445.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{3 + 2\cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{2\cos(c + dx) + 3}\cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(3+2*cos(d*x+c))**(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(2*cos(c + d*x) + 3)*cos(c + d*x)**(3/2)), x)`

3.445.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{3 + 2\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{2\cos(dx + c) + 3}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3+2*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(2*cos(d*x + c) + 3)*cos(d*x + c)^(3/2)), x)`

3.445.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx)\sqrt{3 + 2\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{2\cos(dx + c) + 3}\cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3+2*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(2*cos(d*x + c) + 3)*cos(d*x + c)^(3/2)), x)`

3.445.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{3 + 2 \cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{\frac{3}{2}} \sqrt{2 \cos(c + dx) + 3}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(2*cos(c + d*x) + 3)^(1/2)),x)`output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(2*cos(c + d*x) + 3)^(1/2)), x)`

3.446
$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.446.1 Optimal result 4121
 3.446.2 Mathematica [F] 4121
 3.446.3 Rubi [A] (verified) 4122
 3.446.4 Maple [B] (verified) 4123
 3.446.5 Fricas [F] 4123
 3.446.6 Sympy [F] 4124
 3.446.7 Maxima [F] 4124
 3.446.8 Giac [F] 4124
 3.446.9 Mupad [F(-1)] 4125

3.446.1 Optimal result

Integrand size = 33, antiderivative size = 74

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \frac{2\sqrt{5} \cot(c + dx) E\left(\arcsin\left(\frac{\sqrt{3 - 2 \cos(c + dx)}}{\sqrt{\cos(c + dx)}}\right) \middle| -\frac{1}{5}\right) \sqrt{1 - \sec(c + dx)} \sqrt{1 + \sec(c + dx)}}{3d}$$

output `2/3*cot(d*x+c)*EllipticE((3-2*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),1/5*I*5^(1/2))*5^(1/2)*(1-sec(d*x+c))^(1/2)*(1+sec(d*x+c))^(1/2)/d`

3.446.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `Integrate[(1 + Cos[c + d*x])/(Sqrt[3 - 2*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]`

output `Integrate[(1 + Cos[c + d*x])/(Sqrt[3 - 2*Cos[c + d*x]]*Cos[c + d*x]^(3/2)), x]`

3.446.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(c + dx) + 1}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sqrt{3 - 2 \sin(c + dx + \frac{\pi}{2})} \sin(c + dx + \frac{\pi}{2})^{3/2}} dx$$

↓ 3473

$$\frac{2\sqrt{5} \cot(c + dx) \sqrt{1 - \sec(c + dx)} \sqrt{\sec(c + dx) + 1} E\left(\arcsin\left(\frac{\sqrt{3 - 2 \cos(c + dx)}}{\sqrt{\cos(c + dx)}}\right) \mid -\frac{1}{5}\right)}{3d}$$

input `Int[(1 + Cos[c + d*x])/(Sqrt[3 - 2*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]`

output `(2*Sqrt[5]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[3 - 2*Cos[c + d*x]]/Sqrt[Cos[c + d*x]]], -1/5]*Sqrt[1 - Sec[c + d*x]]*Sqrt[1 + Sec[c + d*x]])/(3*d)`

3.446.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

3.446. $\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$

3.446.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 484 vs. $2(64) = 128$.

Time = 11.59 (sec) , antiderivative size = 485, normalized size of antiderivative = 6.55

method	result
parts	$\frac{(1+\cos(dx+c))\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{3-2\cos(dx+c)}\sqrt{\frac{-2(-3+2\cos(dx+c))}{1+\cos(dx+c)}}F(\cot(dx+c)-\csc(dx+c),i\sqrt{5})}{d(-3+2\cos(dx+c))\sqrt{\cos(dx+c)}} - \frac{2\left(-3\sqrt{-\csc^2(dx+c)}\right)}{d}$
default	$\frac{\left(6F(\cot(dx+c)-\csc(dx+c),i\sqrt{5})\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{-2(-3+2\cos(dx+c))}{1+\cos(dx+c)}}\sqrt{2}(\cos^2(dx+c))-E(\cot(dx+c)-\csc(dx+c),i\sqrt{5})\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{d}$

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3-2*cos(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*(1+cos(d*x+c))*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(3-2*cos(d*x+c))^(1/2)*(-2*(-3+2*cos(d*x+c))/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),I*5^(1/2))/(-3+2*cos(d*x+c))/cos(d*x+c)^(1/2)-2/3/d*(-3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(5*csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),I*5^(1/2))+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(5*csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),I*5^(1/2))+5*csc(d*x+c)^3*(1-cos(d*x+c))^3+csc(d*x+c)-cot(d*x+c))*((5*csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(5*csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2-1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(3/2)`

3.446.5 Fracas [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-2 \cos(dx + c) + 3 \cos(dx + c)}^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3-2*cos(d*x+c))^(1/2),x,algorithm="fracas")`

output `integral(-(cos(d*x + c) + 1)*sqrt(-2*cos(d*x + c) + 3)*sqrt(cos(d*x + c))/
(2*cos(d*x + c)^3 - 3*cos(d*x + c)^2), x)`

3.446.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(3-2*cos(d*x+c))**(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(3 - 2*cos(c + d*x))*cos(c + d*x)**(3/2))
, x)`

3.446.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-2 \cos(dx + c) + 3 \cos(dx + c)^{\frac{3}{2}}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3-2*cos(d*x+c))^(1/2),x, algorith
m="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-2*cos(d*x + c) + 3)*cos(d*x + c)^(3/2))
, x)`

3.446.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-2 \cos(dx + c) + 3 \cos(dx + c)^{\frac{3}{2}}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(3-2*cos(d*x+c))^(1/2),x, algorith
m="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-2*cos(d*x + c) + 3)*cos(d*x + c)^(3/2))
, x)`

3.446.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\sqrt{3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{3/2} \sqrt{3 - 2 \cos(c + dx)}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(3 - 2*cos(c + d*x))^(1/2)),x)`output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(3 - 2*cos(c + d*x))^(1/2)), x)`

$$3.447 \quad \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx$$

3.447.1 Optimal result	4126
3.447.2 Mathematica [F]	4126
3.447.3 Rubi [A] (verified)	4127
3.447.4 Maple [B] (verified)	4128
3.447.5 Fricas [F]	4129
3.447.6 Sympy [F]	4129
3.447.7 Maxima [F]	4130
3.447.8 Giac [F]	4130
3.447.9 Mupad [F(-1)]	4130

3.447.1 Optimal result

Integrand size = 33, antiderivative size = 98

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx = \frac{2\sqrt{5}\sqrt{-\cos(c + dx)}\sqrt{\cos(c + dx)}\csc(c + dx)E\left(\arcsin\left(\frac{\sqrt{-3 + 2 \cos(c + dx)}}{\sqrt{-\cos(c + dx)}}\right) \middle| -\frac{1}{5}\right)\sqrt{1 - \sec(c + dx)}\sqrt{1 + \sec(c + dx)}}{3d}$$

```
output -2/3*csc(d*x+c)*EllipticE((-3+2*cos(d*x+c))^(1/2)/(-cos(d*x+c))^(1/2),1/5*
I*5^(1/2))*5^(1/2)*(-cos(d*x+c))^(1/2)*cos(d*x+c)^(1/2)*(1-sec(d*x+c))^(1/
2)*(1+sec(d*x+c))^(1/2)/d
```

3.447.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx = \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx$$

```
input Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[-3 + 2*Cos[c + d*x]]
),x]
```

```
output Integrate[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[-3 + 2*Cos[c + d*x]]
), x]
```

$$3.447. \quad \int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx$$

3.447.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {3042, 3474, 3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{2\cos(c+dx)-3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{2\sin(c+dx+\frac{\pi}{2})-3}} dx \\
 & \quad \downarrow \text{3474} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\cos(c+dx)+1}{(-\cos(c+dx))^{3/2}\sqrt{2\cos(c+dx)-3}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{(-\sin(c+dx+\frac{\pi}{2}))^{3/2}\sqrt{2\sin(c+dx+\frac{\pi}{2})-3}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3473} \\
 & \frac{2\sqrt{5}\sqrt{-\cos(c+dx)}\sqrt{\cos(c+dx)}\csc(c+dx)\sqrt{1-\sec(c+dx)}\sqrt{\sec(c+dx)+1}E\left(\arcsin\left(\frac{\sqrt{2\cos(c+dx)-3}}{\sqrt{-\cos(c+dx)}}\right)\right)}{3d}
 \end{aligned}$$

input `Int[(1 + Cos[c + d*x])/(Cos[c + d*x]^(3/2)*Sqrt[-3 + 2*Cos[c + d*x]]),x]`

output `(-2*Sqrt[5]*Sqrt[-Cos[c + d*x]]*Sqrt[Cos[c + d*x]]*Csc[c + d*x]*EllipticE[ArcSin[Sqrt[-3 + 2*Cos[c + d*x]]/Sqrt[-Cos[c + d*x]]], -1/5]*Sqrt[1 - Sec[c + d*x]]*Sqrt[1 + Sec[c + d*x]])/(3*d)`

3.447. $\int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{-3+2\cos(c+dx)}} dx$

3.447.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3474 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-Sqrt[(-b)*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]] Int[(A + B*Sin[e + f*x])/((-b)*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && NegQ[(c + d)/b]`

3.447.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 312 vs. $2(84) = 168$.

Time = 13.00 (sec) , antiderivative size = 313, normalized size of antiderivative = 3.19

method	result
default	$-\frac{i\sqrt{5}\sqrt{2}E\left(i(\csc(dx+c)-\cot(dx+c))\sqrt{5}, \frac{i\sqrt{5}}{5}\right)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{-\frac{2(-3+2\cos(dx+c))}{1+\cos(dx+c)}}(\cos^2(dx+c))+2i\sqrt{5}\sqrt{2}E\left(i(\csc(dx+c)-\cot(dx+c))\sqrt{5}, \frac{i\sqrt{5}}{5}\right)}{2\left(3i\sqrt{5}\sqrt{5(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1}F\left(i(\csc(dx+c)-\cot(dx+c))\sqrt{5}, \frac{i\sqrt{5}}{5}\right)-5i\sqrt{5}\sqrt{5(\csc^2(dx+c))(1-\cos(dx+c))^2+1}\right)}$
parts	

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3+2*cos(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

$$3.447. \int \frac{1+\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{-3+2\cos(c+dx)}} dx$$

output
$$-1/3/d*(I*5^{(1/2)}*2^{(1/2)}*EllipticE(I*(csc(d*x+c)-cot(d*x+c))*5^{(1/2)},1/5*I*5^{(1/2)})*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*(-2*(-3+2*cos(d*x+c))/(1+cos(d*x+c)))^{(1/2)}*cos(d*x+c)^2+2*I*5^{(1/2)}*2^{(1/2)}*EllipticE(I*(csc(d*x+c)-cot(d*x+c))*5^{(1/2)},1/5*I*5^{(1/2)})*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*(-2*(-3+2*cos(d*x+c))/(1+cos(d*x+c)))^{(1/2)}*cos(d*x+c)+I*EllipticE(I*(csc(d*x+c)-cot(d*x+c))*5^{(1/2)},1/5*I*5^{(1/2)})*(-2*(-3+2*cos(d*x+c))/(1+cos(d*x+c)))^{(1/2)}*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*2^{(1/2)}*5^{(1/2)}+4*cos(d*x+c)*sin(d*x+c)-6*sin(d*x+c))/(1+cos(d*x+c))/(-3+2*cos(d*x+c))^{(1/2)}/cos(d*x+c)^{(1/2)}$$

3.447.5 Fracas [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{3/2}(c + dx)\sqrt{-3 + 2\cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{2\cos(dx + c) - 3}\cos(dx + c)^{3/2}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3+2*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `integral(sqrt(2*cos(d*x + c) - 3)*(cos(d*x + c) + 1)*sqrt(cos(d*x + c))/(2*cos(d*x + c)^3 - 3*cos(d*x + c)^2), x)`

3.447.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{3/2}(c + dx)\sqrt{-3 + 2\cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{2\cos(c + dx) - 3}\cos^{3/2}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(-3+2*cos(d*x+c))^(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(2*cos(c + d*x) - 3)*cos(c + d*x)**(3/2)), x)`

3.447.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{2 \cos(dx + c) - 3} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3+2*cos(d*x+c))^(1/2),x, algor
ithm="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(2*cos(d*x + c) - 3)*cos(d*x + c)^(3/2))
, x)`

3.447.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{2 \cos(dx + c) - 3} \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3+2*cos(d*x+c))^(1/2),x, algor
ithm="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(2*cos(d*x + c) - 3)*cos(d*x + c)^(3/2))
, x)`

3.447.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{-3 + 2 \cos(c + dx)}} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{\frac{3}{2}} \sqrt{2 \cos(c + dx) - 3}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(2*cos(c + d*x) - 3)^(1/2)),x)`

output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(2*cos(c + d*x) - 3)^(1/2)), x)`

$$3.448 \quad \int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.448.1 Optimal result	4131
3.448.2 Mathematica [F]	4131
3.448.3 Rubi [A] (verified)	4132
3.448.4 Maple [B] (verified)	4133
3.448.5 Fracas [F]	4134
3.448.6 Sympy [F]	4134
3.448.7 Maxima [F]	4135
3.448.8 Giac [F]	4135
3.448.9 Mupad [F(-1)]	4135

3.448.1 Optimal result

Integrand size = 33, antiderivative size = 96

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \frac{2\sqrt{-\cos(c + dx)}\sqrt{\cos(c + dx)}\csc(c + dx)E\left(\arcsin\left(\frac{\sqrt{-3 - 2 \cos(c + dx)}}{\sqrt{5}\sqrt{-\cos(c + dx)}}\right) \middle| -5\right)\sqrt{1 - \sec(c + dx)}\sqrt{1 + \sec(c + dx)}}{3d}$$

```
output -2/3*csc(d*x+c)*EllipticE(1/5*(-3-2*cos(d*x+c))^(1/2)*5^(1/2)/(-cos(d*x+c))^(1/2),I*5^(1/2))*(-cos(d*x+c))^(1/2)*cos(d*x+c)^(1/2)*(1-sec(d*x+c))^(1/2)*(1+sec(d*x+c))^(1/2)/d
```

3.448.2 Mathematica [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

```
input Integrate[(1 + Cos[c + d*x])/(Sqrt[-3 - 2*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]
```

```
output Integrate[(1 + Cos[c + d*x])/(Sqrt[-3 - 2*Cos[c + d*x]]*Cos[c + d*x]^(3/2)), x]
```

$$3.448. \quad \int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$$

3.448.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {3042, 3474, 3042, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(c+dx)+1}{\sqrt{-2\cos(c+dx)-3\cos^{\frac{3}{2}}(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sqrt{-2\sin(c+dx+\frac{\pi}{2})-3\sin(c+dx+\frac{\pi}{2})^{3/2}}} dx \\
 & \quad \downarrow \text{3474} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\cos(c+dx)+1}{\sqrt{-2\cos(c+dx)-3(-\cos(c+dx))^{3/2}}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\sqrt{-\cos(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sqrt{-2\sin(c+dx+\frac{\pi}{2})-3(-\sin(c+dx+\frac{\pi}{2}))^{3/2}}} dx}{\sqrt{\cos(c+dx)}} \\
 & \quad \downarrow \text{3473} \\
 & -\frac{2\sqrt{-\cos(c+dx)}\sqrt{\cos(c+dx)}\csc(c+dx)\sqrt{1-\sec(c+dx)}\sqrt{\sec(c+dx)+1}E\left(\arcsin\left(\frac{\sqrt{-2\cos(c+dx)-3}}{\sqrt{5}\sqrt{-\cos(c+dx)}}\right)\right)}{3d}
 \end{aligned}$$

input `Int[(1 + Cos[c + d*x])/(Sqrt[-3 - 2*Cos[c + d*x]]*Cos[c + d*x]^(3/2)),x]`

output `(-2*Sqrt[-Cos[c + d*x]]*Sqrt[Cos[c + d*x]]*Csc[c + d*x]*EllipticE[ArcSin[Sqrt[-3 - 2*Cos[c + d*x]]/(Sqrt[5]*Sqrt[-Cos[c + d*x]])], -5]*Sqrt[1 - Sec[c + d*x]]*Sqrt[1 + Sec[c + d*x]])/(3*d)`

3.448. $\int \frac{1+\cos(c+dx)}{\sqrt{-3-2\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$

3.448.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3474 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-Sqrt[(-b)*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]] Int[(A + B*Sin[e + f*x])/((-b)*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && NegQ[(c + d)/b]`

3.448.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 330 vs. $2(85) = 170$.

Time = 12.62 (sec) , antiderivative size = 331, normalized size of antiderivative = 3.45

method	result
default	$\left(i\sqrt{10}\sqrt{2}\sqrt{5}\sqrt{\frac{3+2\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}E\left(\frac{i(\csc(dx+c)-\cot(dx+c))\sqrt{5}}{5}, i\sqrt{5}\right)(\cos^2(dx+c))+2i\sqrt{10}\sqrt{2}\sqrt{5}\sqrt{\frac{3+2\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)$
parts	$\frac{i(1+\cos(dx+c))\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{-3-2\cos(dx+c)}\sqrt{10}\sqrt{\frac{3+2\cos(dx+c)}{1+\cos(dx+c)}}F\left(\frac{i(\csc(dx+c)-\cot(dx+c))\sqrt{5}}{5}, i\sqrt{5}\right)\sqrt{5}}{5d(3+2\cos(dx+c))\sqrt{\cos(dx+c)}} - \frac{2\left(i\sqrt{5}\sqrt{5(\csc(dx+c)-\cot(dx+c))}\sqrt{5}\right)}{5d(3+2\cos(dx+c))\sqrt{\cos(dx+c)}}$

input `int((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3-2*cos(d*x+c))^(1/2), x, method=_RETURNVERBOSE)`

3.448.
$$\int \frac{1+\cos(c+dx)}{\sqrt{-3-2\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$$

output $1/15/d*(I*10^{(1/2)}*2^{(1/2)}*5^{(1/2)}*((3+2*\cos(d*x+c))/(1+\cos(d*x+c)))^{(1/2)} * (\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)} * \text{EllipticE}(1/5*I*(\csc(d*x+c)-\cot(d*x+c)) * 5^{(1/2)}, I*5^{(1/2)})*\cos(d*x+c)^2+2*I*10^{(1/2)}*2^{(1/2)}*5^{(1/2)}*((3+2*\cos(d*x+c))/(1+\cos(d*x+c)))^{(1/2)} * (\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)} * \text{EllipticE}(1/5*I*(\csc(d*x+c)-\cot(d*x+c))*5^{(1/2)}, I*5^{(1/2)})*\cos(d*x+c)+I*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)} * \text{EllipticE}(1/5*I*(\csc(d*x+c)-\cot(d*x+c))*5^{(1/2)}, I*5^{(1/2)})*5^{(1/2)}*2^{(1/2)}*((3+2*\cos(d*x+c))/(1+\cos(d*x+c)))^{(1/2)}*10^{(1/2)}-20*\cos(d*x+c)*\sin(d*x+c)-30*\sin(d*x+c))*(-3-2*\cos(d*x+c))^{(1/2)}/(3+2*\cos(d*x+c))/\cos(d*x+c)^{(1/2)}/(1+\cos(d*x+c))$

3.448.5 Fricas [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-2 \cos(dx + c) - 3} \cos^{\frac{3}{2}}(dx + c)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3-2*cos(d*x+c))^(1/2),x, algorith="fricas")`

output `integral(-(cos(d*x + c) + 1)*sqrt(-2*cos(d*x + c) - 3)*sqrt(cos(d*x + c))/(2*cos(d*x + c)^3 + 3*cos(d*x + c)^2), x)`

3.448.6 Sympy [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\sqrt{-2 \cos(c + dx) - 3} \cos^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)**(3/2)/(-3-2*cos(d*x+c))^(1/2),x)`

output `Integral((cos(c + d*x) + 1)/(sqrt(-2*cos(c + d*x) - 3)*cos(c + d*x)**(3/2)), x)`

3.448.7 Maxima [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-2 \cos(dx + c) - 3 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3-2*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-2*cos(d*x + c) - 3)*cos(d*x + c)^(3/2)), x)`

3.448.8 Giac [F]

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(dx + c) + 1}{\sqrt{-2 \cos(dx + c) - 3 \cos(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((1+cos(d*x+c))/cos(d*x+c)^(3/2)/(-3-2*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((cos(d*x + c) + 1)/(sqrt(-2*cos(d*x + c) - 3)*cos(d*x + c)^(3/2)), x)`

3.448.9 Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\cos(c + dx) + 1}{\cos(c + dx)^{\frac{3}{2}} \sqrt{-2 \cos(c + dx) - 3}} dx$$

input `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(- 2*cos(c + d*x) - 3)^(1/2)),x)`

output `int((cos(c + d*x) + 1)/(cos(c + d*x)^(3/2)*(- 2*cos(c + d*x) - 3)^(1/2)),x)`

3.448. $\int \frac{1 + \cos(c + dx)}{\sqrt{-3 - 2 \cos(c + dx)} \cos^{\frac{3}{2}}(c + dx)} dx$

3.449 $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$

3.449.1 Optimal result	4136
3.449.2 Mathematica [N/A]	4136
3.449.3 Rubi [N/A]	4137
3.449.4 Maple [N/A] (verified)	4138
3.449.5 Fricas [N/A]	4138
3.449.6 Sympy [N/A]	4138
3.449.7 Maxima [N/A]	4139
3.449.8 Giac [N/A]	4139
3.449.9 Mupad [N/A]	4140

3.449.1 Optimal result

Integrand size = 33, antiderivative size = 33

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$$

$$= \text{Int}((c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)), x)$$

output `Unintegrable((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^n*(A+B*cos(f*x+e)),x)`

3.449.2 Mathematica [N/A]

Not integrable

Time = 16.14 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$$

input `Integrate[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^n*(A + B*Cos[e + f*x]),x]`

output `Integrate[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^n*(A + B*Cos[e + f*x]),x]`

3.449.3 Rubi [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3486}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (A + B \cos(e + fx))(c \cos(e + fx))^m (a + b \cos(e + fx))^n dx$$

↓ 3042

$$\int \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right) \right) \left(c \sin\left(e + fx + \frac{\pi}{2}\right) \right)^m \left(a + b \sin\left(e + fx + \frac{\pi}{2}\right) \right)^n dx$$

↓ 3486

$$\int (A + B \cos(e + fx))(c \cos(e + fx))^m (a + b \cos(e + fx))^n dx$$

input `Int[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^n*(A + B*Cos[e + f*x]),x]`

output `$Aborted`

3.449.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3486 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(A + B*Sin[e + f*x])*(c + d*Sin[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.449.4 Maple [N/A] (verified)

Not integrable

Time = 0.87 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int (c \cos(fx + e))^m (a + b \cos(fx + e))^n (A + \cos(fx + e) B) dx$$

input `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^n*(A+cos(f*x+e)*B),x)`

output `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^n*(A+cos(f*x+e)*B),x)`

3.449.5 Fricas [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^n (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^n*(A+B*cos(f*x+e)),x, algorithm="fricas")`

output `integral((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^n*(c*cos(f*x + e))^m, x)`

3.449.6 Sympy [N/A]

Not integrable

Time = 123.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx \\ &= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^n dx \end{aligned}$$

input `integrate((c*cos(f*x+e))**m*(a+b*cos(f*x+e))**n*(A+B*cos(f*x+e)),x)`

output `Integral((c*cos(e + f*x))**m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))**n,
x)`

3.449.7 Maxima [N/A]

Not integrable

Time = 3.95 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^n (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^n*(A+B*cos(f*x+e)),x, algorithm
m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^n*(c*cos(f*x + e))^m,
x)`

3.449.8 Giac [N/A]

Not integrable

Time = 1.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^n (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^n*(A+B*cos(f*x+e)),x, algorithm
m="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^n*(c*cos(f*x + e))^m,
x)`

3.449.9 Mupad [N/A]

Not integrable

Time = 2.73 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^n dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^n,x)`output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^n, x)`

3.450 $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx$

3.450.1 Optimal result	4141
3.450.2 Mathematica [A] (verified)	4142
3.450.3 Rubi [A] (verified)	4143
3.450.4 Maple [F]	4147
3.450.5 Fricas [F]	4147
3.450.6 Sympy [F(-1)]	4148
3.450.7 Maxima [F]	4148
3.450.8 Giac [F]	4149
3.450.9 Mupad [F(-1)]	4149

3.450.1 Optimal result

Integrand size = 33, antiderivative size = 595

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx$$

$$= \frac{b(Ab^3(15 + 8m + m^2) + 4ab^2B(15 + 8m + m^2) + 2a^3B(28 + 10m + m^2) + a^2Ab(110 + 47m + 5m^2)) (c \cos(e + fx))^{m+1} \sin(e + fx)}{cf(2 + m)(4 + m)(5 + m)}$$

$$+ \frac{b^2(b^2B(4 + m)^2 + 2aAb(5 + m)^2 + a^2B(36 + 11m + m^2)) \cos(e + fx)(c \cos(e + fx))^{1+m} \sin(e + fx)}{cf(3 + m)(4 + m)(5 + m)}$$

$$+ \frac{b(Ab(5 + m) + aB(8 + m))(c \cos(e + fx))^{1+m} (a + b \cos(e + fx))^2 \sin(e + fx)}{cf(4 + m)(5 + m)}$$

$$+ \frac{bB(c \cos(e + fx))^{1+m} (a + b \cos(e + fx))^3 \sin(e + fx)}{cf(5 + m)}$$

$$- \frac{(Ab^4(3 + 4m + m^2) + 4ab^3B(3 + 4m + m^2) + 6a^2Ab^2(4 + 5m + m^2) + 4a^3bB(4 + 5m + m^2) + a^4A(8 + 6m + m^2)) (c \cos(e + fx))^{m+1} \sin(e + fx)}{cf(1 + m)(2 + m)(4 + m)}$$

$$- \frac{(b^4B(8 + 6m + m^2) + 4aAb^3(10 + 7m + m^2) + 6a^2b^2B(10 + 7m + m^2) + 4a^3Ab(15 + 8m + m^2) + a^4A(15 + 8m + m^2)) (c \cos(e + fx))^{m+1} \sin(e + fx)}{c^2f(2 + m)(3 + m)(5 + m)}$$

output `b*(A*b^3*(m^2+8*m+15)+4*a*b^2*B*(m^2+8*m+15)+2*a^3*B*(m^2+10*m+28)+a^2*A*b*(5*m^2+47*m+110))*(c*cos(f*x+e))^(1+m)*sin(f*x+e)/c/f/(5+m)/(m^2+6*m+8)+b^2*(b^2*B*(4+m)^2+2*a*A*b*(5+m)^2+a^2*B*(m^2+11*m+36))*cos(f*x+e)*(c*cos(f*x+e))^(1+m)*sin(f*x+e)/c/f/(3+m)/(4+m)/(5+m)+b*(A*b*(5+m)+a*B*(8+m))*(c*cos(f*x+e))^(1+m)*(a+b*cos(f*x+e))^2*sin(f*x+e)/c/f/(4+m)/(5+m)+b*B*(c*cos(f*x+e))^(1+m)*(a+b*cos(f*x+e))^3*sin(f*x+e)/c/f/(5+m)-(A*b^4*(m^2+4*m+3)+4*a*b^3*B*(m^2+4*m+3)+6*a^2*A*b^2*(m^2+5*m+4)+4*a^3*b*B*(m^2+5*m+4)+a^4*A*(m^2+6*m+8))*(c*cos(f*x+e))^(1+m)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], cos(f*x+e)^2)*sin(f*x+e)/c/f/(4+m)/(m^2+3*m+2)/(sin(f*x+e)^2)^(1/2)-(b^4*B*(m^2+6*m+8)+4*a*A*b^3*(m^2+7*m+10)+6*a^2*b^2*B*(m^2+7*m+10)+4*a^3*A*b*(m^2+8*m+15)+a^4*B*(m^2+8*m+15))*(c*cos(f*x+e))^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], cos(f*x+e)^2)*sin(f*x+e)/c^2/f/(2+m)/(3+m)/(5+m)/(sin(f*x+e)^2)^(1/2)`

3.450.2 Mathematica [A] (verified)

Time = 4.73 (sec) , antiderivative size = 319, normalized size of antiderivative = 0.54

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx$$

$$= \frac{(c \cos(e + fx))^m \cot(e + fx) \left(-\frac{a^4 A \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{1+m} + \cos(e + fx) \left(-\frac{a^3 (4Ab + aB) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, \cos^2(e + fx)\right)}{2+m} + b \cos(e + fx) \left(-\frac{2a^2 (3Ab + 2aB) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, \cos^2(e + fx)\right)}{3+m} + b \cos(e + fx) \left(-\frac{2a (2Ab + 3aB) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, \cos^2(e + fx)\right)}{4+m} + b \cos(e + fx) \left(-\frac{(Ab + 4aB) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5+m}{2}, \frac{7+m}{2}, \cos^2(e + fx)\right)}{5+m} - (bB \cos(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{6+m}{2}, \frac{8+m}{2}, \cos^2(e + fx)\right)) \right) \right) \right) \right) \operatorname{Sqrt}[\sin(e + fx)^2]}{f}$$

input `Integrate[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^4*(A + B*Cos[e + f*x]),x]`

output `((c*Cos[e + f*x])^m*Cot[e + f*x]*(-((a^4*A*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2)]/(1 + m)) + Cos[e + f*x]*(-((a^3*(4*A*b + a*B)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2)]/(2 + m)) + b*Cos[e + f*x]*((-2*a^2*(3*A*b + 2*a*B)*Hypergeometric2F1[1/2, (3 + m)/2, (5 + m)/2, Cos[e + f*x]^2)]/(3 + m) + b*Cos[e + f*x]*((-2*a*(2*A*b + 3*a*B)*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, Cos[e + f*x]^2)]/(4 + m) + b*Cos[e + f*x]*(-((A*b + 4*a*B)*Hypergeometric2F1[1/2, (5 + m)/2, (7 + m)/2, Cos[e + f*x]^2)]/(5 + m)) - (b*B*Cos[e + f*x]*Hypergeometric2F1[1/2, (6 + m)/2, (8 + m)/2, Cos[e + f*x]^2)]/(6 + m)))))*Sqrt[Sin[e + f*x]^2]/f`

3.450.3 Rubi [A] (verified)

Time = 2.94 (sec) , antiderivative size = 614, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3469, 3042, 3528, 3042, 3512, 3042, 3502, 3042, 3227, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \cos(e + fx))^m dx$$

$$\downarrow 3042$$

$$\int \left(a + b \sin\left(e + fx + \frac{\pi}{2}\right)\right)^4 \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right)\right) \left(c \sin\left(e + fx + \frac{\pi}{2}\right)\right)^m dx$$

$$\downarrow 3469$$

$$\frac{\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (bc(Ab(m + 5) + aB(m + 8)) \cos^2(e + fx) + c(B(m + 4)b^2 + a(2Ab + aB))) dx}{c(m + 5)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^3 (c \cos(e + fx))^{m+1}}{cf(m + 5)}$$

$$\downarrow 3042$$

$$\frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m (a + b \sin(e + fx + \frac{\pi}{2}))^2 (bc(Ab(m + 5) + aB(m + 8)) \sin(e + fx + \frac{\pi}{2})^2 + c(B(m + 4)b^2 + a(2Ab + aB))) dx}{c(m + 5)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^3 (c \cos(e + fx))^{m+1}}{cf(m + 5)}$$

$$\downarrow 3528$$

$$\frac{\int (c \cos(e + fx))^m (a + b \cos(e + fx)) (b(B(m^2 + 11m + 36)a^2 + 2Ab(m + 5)^2 a + b^2 B(m + 4)^2) \cos^2(e + fx) c^2 + a(a(m + 4)(bB(m + 1) + aA(m + 5)) + b(m + 4)A)) dx}{c(m + 4)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^3 (c \cos(e + fx))^{m+1}}{cf(m + 5)}$$

$$\downarrow 3042$$

$$\frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m (a + b \sin(e + fx + \frac{\pi}{2})) (b(B(m^2 + 11m + 36)a^2 + 2Ab(m + 5)^2 a + b^2 B(m + 4)^2) \sin(e + fx + \frac{\pi}{2})^2 c^2 + a(a(m + 4)(bB(m + 1) + aA(m + 5)) + b(m + 4)A)) dx}{c(m + 4)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^3 (c \cos(e + fx))^{m+1}}{cf(m + 5)}$$

3.450. $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx$

↓ 3512

$$f(c \cos(e+fx))^m (b(m+3)(2B(m^2+10m+28)a^3+Ab(5m^2+47m+110)a^2+4b^2B(m^2+8m+15)a+Ab^3(m^2+8m+15)) \cos^2(e+fx)c^3+a^2(m+3)(a(m+4)(bB(m+1$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^3(c \cos(e+fx))^{m+1}}{cf(m+5)}$$

↓ 3042

$$f(c \sin(e+fx+\frac{\pi}{2}))^m (b(m+3)(2B(m^2+10m+28)a^3+Ab(5m^2+47m+110)a^2+4b^2B(m^2+8m+15)a+Ab^3(m^2+8m+15)) \sin(e+fx+\frac{\pi}{2})^2 c^3+a^2(m+3)(a(m+4$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^3(c \cos(e+fx))^{m+1}}{cf(m+5)}$$

↓ 3502

$$f(c \cos(e+fx))^m ((m+3)(A(m^3+11m^2+38m+40)a^4+4bB(m^3+10m^2+29m+20)a^3+6Ab^2(m^3+10m^2+29m+20)a^2+4b^3B(m^3+9m^2+23m+15)a+Ab^4(m^3+9m^2+23m+15))$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^3(c \cos(e+fx))^{m+1}}{cf(m+5)}$$

↓ 3042

$$f(c \sin(e+fx+\frac{\pi}{2}))^m ((m+3)(A(m^3+11m^2+38m+40)a^4+4bB(m^3+10m^2+29m+20)a^3+6Ab^2(m^3+10m^2+29m+20)a^2+4b^3B(m^3+9m^2+23m+15)a+Ab^4(m^3+9m^2+23m+15))$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^3(c \cos(e+fx))^{m+1}}{cf(m+5)}$$

↓ 3227

$$c^4(m+3)(a^4A(m^3+11m^2+38m+40)+4a^3bB(m^3+10m^2+29m+20)+6a^2Ab^2(m^3+10m^2+29m+20)+4ab^3B(m^3+9m^2+23m+15)+Ab^4(m^3+9m^2+23m+15))$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^3(c \cos(e+fx))^{m+1}}{cf(m+5)}$$

↓ 3042

3.450. $\int (c \cos(e+fx))^m (a+b \cos(e+fx))^4 (A+B \cos(e+fx)) dx$

$$\frac{c^4(m+3)(a^4A(m^3+11m^2+38m+40)+4a^3bB(m^3+10m^2+29m+20)+6a^2Ab^2(m^3+10m^2+29m+20)+4ab^3B(m^3+9m^2+23m+15)+Ab^4(m^3+9m^2+23m+15))}{cf(m+5)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^3(c \cos(e + fx))^{m+1}}{cf(m + 5)}$$

↓ 3122

$$\frac{b^2c \sin(e+fx) \cos(e+fx)(a^2B(m^2+11m+36)+2aAb(m+5)^2+b^2B(m+4)^2)(c \cos(e+fx))^{m+1}}{f(m+3)} + \frac{bc^2(m+3) \sin(e+fx)(2a^3B(m^2+10m+28)+a^2Ab(5m^2+47m+15)+Ab^3(m^2+10m+28)+b^4B(m^2+10m+28))}{f(m+3)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^3(c \cos(e + fx))^{m+1}}{cf(m + 5)}$$

input `Int[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^4*(A + B*Cos[e + f*x]),x]`

output

```
(b*B*(c*Cos[e + f*x])^(1 + m)*(a + b*Cos[e + f*x])^3*Sin[e + f*x])/(c*f*(5 + m)) + ((b*(A*b*(5 + m) + a*B*(8 + m))*(c*Cos[e + f*x])^(1 + m)*(a + b*Cos[e + f*x])^2*Sin[e + f*x])/(f*(4 + m)) + ((b^2*c*(b^2*B*(4 + m)^2 + 2*a*A*b*(5 + m)^2 + a^2*B*(36 + 11*m + m^2))*Cos[e + f*x]*(c*Cos[e + f*x])^(1 + m)*Sin[e + f*x])/(f*(3 + m)) + ((b*c^2*(3 + m)*(A*b^3*(15 + 8*m + m^2) + 4*a*b^2*B*(15 + 8*m + m^2) + 2*a^3*B*(28 + 10*m + m^2) + a^2*A*b*(110 + 4*7*m + 5*m^2))*(c*Cos[e + f*x])^(1 + m)*Sin[e + f*x])/(f*(2 + m)) + (-((c^3*(3 + m)*(A*b^4*(15 + 23*m + 9*m^2 + m^3) + 4*a*b^3*B*(15 + 23*m + 9*m^2 + m^3) + 6*a^2*A*b^2*(20 + 29*m + 10*m^2 + m^3) + 4*a^3*b*B*(20 + 29*m + 10*m^2 + m^3) + a^4*A*(40 + 38*m + 11*m^2 + m^3))*(c*Cos[e + f*x])^(1 + m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*Sin[e + f*x])/(f*(1 + m)*Sqrt[Sin[e + f*x]^2])) - (c^2*(4 + m)*(b^4*B*(8 + 6*m + m^2) + 4*a*A*b^3*(10 + 7*m + m^2) + 6*a^2*b^2*B*(10 + 7*m + m^2) + 4*a^3*A*b*(15 + 8*m + m^2) + a^4*B*(15 + 8*m + m^2))*(c*Cos[e + f*x])^(2 + m)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2]*Sin[e + f*x])/(f*Sqrt[Sin[e + f*x]^2]))/(c*(2 + m))/(c*(3 + m))/(c*(4 + m))/(c*(5 + m))
```


3.450.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

rule 3512 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Sin[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) + A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d)*(m + 3))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

3.450.4 Maple [F]

$$\int (c \cos(fx + e))^m (a + b \cos(fx + e))^4 (A + \cos(fx + e) B) dx$$

input `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^4*(A+cos(f*x+e)*B),x)`

output `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^4*(A+cos(f*x+e)*B),x)`

3.450.5 Fracas [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^4 (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^4*(A+B*cos(f*x+e)),x, algorithm m="fracas")`

output `integral((B*b^4*cos(f*x + e)^5 + A*a^4 + (4*B*a*b^3 + A*b^4)*cos(f*x + e)^4 + 2*(3*B*a^2*b^2 + 2*A*a*b^3)*cos(f*x + e)^3 + 2*(2*B*a^3*b + 3*A*a^2*b^2)*cos(f*x + e)^2 + (B*a^4 + 4*A*a^3*b)*cos(f*x + e))*(c*cos(f*x + e))^m, x)`

3.450.6 Sympy [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx = \text{Timed out}$$

input `integrate((c*cos(f*x+e))**m*(a+b*cos(f*x+e))**4*(A+B*cos(f*x+e)),x)`

output `Timed out`

3.450.7 Maxima [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^4 (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^4*(A+B*cos(f*x+e)),x, algorithm m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^4*(c*cos(f*x + e))^m, x)`

3.450.8 Giac [F]

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^4 (c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^4*(A+B*cos(f*x+e)),x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^4*(c*cos(f*x + e))^m, x)`

3.450.9 Mupad [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^4 dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^4,x)`

output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^4, x)`

3.451 $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx$

3.451.1 Optimal result	4150
3.451.2 Mathematica [A] (verified)	4151
3.451.3 Rubi [A] (verified)	4152
3.451.4 Maple [F]	4155
3.451.5 Fricas [F]	4155
3.451.6 Sympy [F(-1)]	4156
3.451.7 Maxima [F]	4156
3.451.8 Giac [F]	4157
3.451.9 Mupad [F(-1)]	4157

3.451.1 Optimal result

Integrand size = 33, antiderivative size = 406

$$\begin{aligned}
 & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx \\
 = & \frac{b(b^2 B(3 + m) + 3aAb(4 + m) + 2a^2 B(5 + m)) (c \cos(e + fx))^{1+m} \sin(e + fx)}{cf(2 + m)(4 + m)} \\
 & + \frac{b^2 (Ab(4 + m) + aB(6 + m)) \cos(e + fx) (c \cos(e + fx))^{1+m} \sin(e + fx)}{cf(3 + m)(4 + m)} \\
 & + \frac{bB(c \cos(e + fx))^{1+m} (a + b \cos(e + fx))^2 \sin(e + fx)}{cf(4 + m)} \\
 & - \frac{(a^2(2 + m)(bB(1 + m) + aA(4 + m)) + b(1 + m)(b^2 B(3 + m) + 3aAb(4 + m) + 2a^2 B(5 + m))) (c \cos(e + fx))^{1+m} \sin(e + fx)}{cf(1 + m)(2 + m)(4 + m)\sqrt{\sin^2(e + fx)}} \\
 & - \frac{(Ab^3(2 + m) + 3ab^2 B(2 + m) + 3a^2 Ab(3 + m) + a^3 B(3 + m)) (c \cos(e + fx))^{2+m} \operatorname{Hypergeometric2F1}(m, 1, m+1, -\frac{b \cos(e + fx)}{c})}{c^2 f(2 + m)(3 + m)\sqrt{\sin^2(e + fx)}}
 \end{aligned}$$

```

output b*(b^2*B*(3+m)+3*a*A*b*(4+m)+2*a^2*B*(5+m))*(c*cos(f*x+e))^(1+m)*sin(f*x+e)
)/c/f/(2+m)/(4+m)+b^2*(A*b*(4+m)+a*B*(6+m))*cos(f*x+e)*(c*cos(f*x+e))^(1+m)
)*sin(f*x+e)/c/f/(3+m)/(4+m)+b*B*(c*cos(f*x+e))^(1+m)*(a+b*cos(f*x+e))^2*s
in(f*x+e)/c/f/(4+m)-(a^2*(2+m)*(b*B*(1+m)+a*A*(4+m))+b*(1+m)*(b^2*B*(3+m)+
3*a*A*b*(4+m)+2*a^2*B*(5+m))*(c*cos(f*x+e))^(1+m)*hypergeom([1/2, 1/2+1/2
*m],[3/2+1/2*m],cos(f*x+e)^2)*sin(f*x+e)/c/f/(1+m)/(2+m)/(4+m)/(sin(f*x+e)
^2)^(1/2)-(A*b^3*(2+m)+3*a*b^2*B*(2+m)+3*a^2*A*b*(3+m)+a^3*B*(3+m))*(c*cos
(f*x+e))^(2+m)*hypergeom([1/2, 1+1/2*m],[2+1/2*m],cos(f*x+e)^2)*sin(f*x+e)
/c^2/f/(2+m)/(3+m)/(sin(f*x+e)^2)^(1/2)

```

3.451.2 Mathematica [A] (verified)

Time = 1.79 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.65

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx$$

$$= \frac{(c \cos(e + fx))^m \cot(e + fx) \left(-\frac{a^3 A \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{1+m} + \cos(e + fx) \left(-\frac{a^2 (3Ab + aB) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{1+m} \right) \right)}{1}$$

```

input Integrate[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^3*(A + B*Cos[e + f*x]),x
]

```

```

output ((c*Cos[e + f*x])^m*Cot[e + f*x]*(-((a^3*A*Hypergeometric2F1[1/2, (1 + m)/
2, (3 + m)/2, Cos[e + f*x]^2])/(1 + m)) + Cos[e + f*x]*(-((a^2*(3*A*b + a*
B)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2])/(2 + m))
+ b*Cos[e + f*x]*((-3*a*(A*b + a*B)*Hypergeometric2F1[1/2, (3 + m)/2, (5 +
m)/2, Cos[e + f*x]^2])/(3 + m) + b*Cos[e + f*x]*(-((A*b + 3*a*B)*Hyperge
ometric2F1[1/2, (4 + m)/2, (6 + m)/2, Cos[e + f*x]^2])/(4 + m)) - (b*B*Cos
[e + f*x]*Hypergeometric2F1[1/2, (5 + m)/2, (7 + m)/2, Cos[e + f*x]^2])/(5
+ m))))*Sqrt[Sin[e + f*x]^2])/f

```

3.451.3 Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 3469, 3042, 3512, 3042, 3502, 3042, 3227, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \cos(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin\left(e + fx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right)\right) \left(c \sin\left(e + fx + \frac{\pi}{2}\right)\right)^m dx$$

$$\downarrow \text{3469}$$

$$\frac{\int (c \cos(e + fx))^m (a + b \cos(e + fx)) (bc(Ab(m + 4) + aB(m + 6)) \cos^2(e + fx) + c(B(m + 3)b^2 + a(2Ab + aB))) dx}{c(m + 4)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^2 (c \cos(e + fx))^{m+1}}{cf(m + 4)}$$

$$\downarrow \text{3042}$$

$$\frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m (a + b \sin(e + fx + \frac{\pi}{2})) (bc(Ab(m + 4) + aB(m + 6)) \sin^2(e + fx + \frac{\pi}{2}) + c(B(m + 3)b^2 + a(2Ab + aB))) dx}{c(m + 4)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^2 (c \cos(e + fx))^{m+1}}{cf(m + 4)}$$

$$\downarrow \text{3512}$$

$$\frac{\int (c \cos(e + fx))^m (b(m + 3)(2B(m + 5)a^2 + 3Ab(m + 4)a + b^2B(m + 3)) \cos^2(e + fx)c^2 + a^2(m + 3)(bB(m + 1) + aA(m + 4))c^2 + (m + 4)(B(m + 3)a^3 + 3Ab^2a)) dx}{c(m + 3)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^2 (c \cos(e + fx))^{m+1}}{cf(m + 4)}$$

$$\downarrow \text{3042}$$

$$\frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m (b(m + 3)(2B(m + 5)a^2 + 3Ab(m + 4)a + b^2B(m + 3)) \sin^2(e + fx + \frac{\pi}{2})c^2 + a^2(m + 3)(bB(m + 1) + aA(m + 4))c^2 + (m + 4)(B(m + 3)a^3 + 3Ab^2a)) dx}{c(m + 3)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))^2 (c \cos(e + fx))^{m+1}}{cf(m + 4)}$$

3.451. $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx$

↓ 3502

$$\frac{\int (c \cos(e+fx))^m \left((m+3) \left((m+2)(bB(m+1)+aA(m+4))a^2+b(m+1)(2B(m+5)a^2+3Ab(m+4)a+b^2B(m+3)) \right) \right) e^{3+(m+2)(m+4)(B(m+3)a^3+3Ab(m+3)a^2+3b^2B(m+3))}}{c^{m+2}}}{c^{m+3}}$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^2 (c \cos(e+fx))^{m+1}}{cf(m+4)}$$

↓ 3042

$$\frac{\int \left(c \sin\left(e+fx+\frac{\pi}{2}\right) \right)^m \left((m+3) \left((m+2)(bB(m+1)+aA(m+4))a^2+b(m+1)(2B(m+5)a^2+3Ab(m+4)a+b^2B(m+3)) \right) \right) e^{3+(m+2)(m+4)(B(m+3)a^3+3Ab(m+3)a^2+3b^2B(m+3))}}{c^{m+2}}}{c^{m+3}}$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^2 (c \cos(e+fx))^{m+1}}{cf(m+4)}$$

↓ 3227

$$\frac{c^3(m+3)(b(m+1)(2a^2B(m+5)+3aAb(m+4)+b^2B(m+3))+a^2(m+2)(aA(m+4)+bB(m+1))) \int (c \cos(e+fx))^m dx + c^2(m+2)(m+4)(a^3B(m+3)+3a^2Ab(m+3)+3aAb(m+4)+b^2B(m+3))}{c^{m+2}}}{c^{m+3}}$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^2 (c \cos(e+fx))^{m+1}}{cf(m+4)}$$

↓ 3042

$$\frac{c^3(m+3)(b(m+1)(2a^2B(m+5)+3aAb(m+4)+b^2B(m+3))+a^2(m+2)(aA(m+4)+bB(m+1))) \int \left(c \sin\left(e+fx+\frac{\pi}{2}\right) \right)^m dx + c^2(m+2)(m+4)(a^3B(m+3)+3a^2Ab(m+3)+3aAb(m+4)+b^2B(m+3))}{c^{m+2}}}{c^{m+3}}$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^2 (c \cos(e+fx))^{m+1}}{cf(m+4)}$$

↓ 3122

$$\frac{bc(m+3) \sin(e+fx)(2a^2B(m+5)+3aAb(m+4)+b^2B(m+3))(c \cos(e+fx))^{m+1}}{f(m+2)} + \frac{c^2(m+3) \sin(e+fx)(b(m+1)(2a^2B(m+5)+3aAb(m+4)+b^2B(m+3))+a^2(m+2)(aA(m+4)+bB(m+1)))}{f(m)}$$

$$\frac{bB \sin(e+fx)(a+b \cos(e+fx))^2 (c \cos(e+fx))^{m+1}}{cf(m+4)}$$

input `Int[(c*cos[e + f*x])^m*(a + b*cos[e + f*x])^3*(A + B*cos[e + f*x]),x]`

3.451. $\int (c \cos(e+fx))^m (a+b \cos(e+fx))^3 (A+B \cos(e+fx)) dx$


```
output (b*B*(c*cos[e + f*x])^(1 + m)*(a + b*cos[e + f*x])^2*sin[e + f*x])/(c*f*(4
+ m)) + ((b^2*(A*b*(4 + m) + a*B*(6 + m))*cos[e + f*x]*(c*cos[e + f*x])^(
1 + m)*sin[e + f*x])/(f*(3 + m)) + ((b*c*(3 + m)*(b^2*B*(3 + m) + 3*a*A*b*
(4 + m) + 2*a^2*B*(5 + m))*(c*cos[e + f*x])^(1 + m)*sin[e + f*x])/(f*(2 +
m)) + (-((c^2*(3 + m)*(a^2*(2 + m)*(b*B*(1 + m) + a*A*(4 + m)) + b*(1 + m)
*(b^2*B*(3 + m) + 3*a*A*b*(4 + m) + 2*a^2*B*(5 + m)))*(c*cos[e + f*x])^(1
+ m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*sin[e +
f*x])/(f*(1 + m)*Sqrt[Sin[e + f*x]^2])) - (c*(4 + m)*(A*b^3*(2 + m) + 3*a*
b^2*B*(2 + m) + 3*a^2*A*b*(3 + m) + a^3*B*(3 + m))*(c*cos[e + f*x])^(2 + m)
)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2]*sin[e + f*x
])/ (f*Sqrt[Sin[e + f*x]^2]))/(c*(2 + m))/(c*(3 + m))/(c*(4 + m))
```

3.451.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 3227 Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x
_)])], x_Symbol] := Simp[c Int[(b*sin[e + f*x])^m, x], x] + Simp[d/b Int
[(b*sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]
```

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*cos[e + f*x]*(a + b*sin[e + f*x])^(m - 1)*((c + d*sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*sin[e +
f*x])^(m - 2)*(c + d*sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

```
rule 3512 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]*(A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-C)*d*Cos[e + f*x]*Sin[e + f*x]*((a + b*Si
n[e + f*x])^(m + 1)/(b*f*(m + 3))), x] + Simp[1/(b*(m + 3)) Int[(a + b*Si
n[e + f*x])^m*Simp[a*C*d + A*b*c*(m + 3) + b*(B*c*(m + 3) + d*(C*(m + 2) +
A*(m + 3)))*Sin[e + f*x] - (2*a*C*d - b*(c*C + B*d))*(m + 3))*Sin[e + f*x]^2
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d,
0] && NeQ[a^2 - b^2, 0] && !LtQ[m, -1]
```

3.451.4 Maple [F]

$$\int (c \cos(fx + e))^m (a + b \cos(fx + e))^3 (A + \cos(fx + e) B) dx$$

```
input int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^3*(A+cos(f*x+e)*B),x)
```

```
output int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^3*(A+cos(f*x+e)*B),x)
```

3.451.5 Fracas [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^3 (c \cos(fx + e))^m dx \end{aligned}$$

```
input integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^3*(A+B*cos(f*x+e)),x, algorith
m="fracas")
```

output `integral((B*b^3*cos(f*x + e)^4 + A*a^3 + (3*B*a*b^2 + A*b^3)*cos(f*x + e)^3 + 3*(B*a^2*b + A*a*b^2)*cos(f*x + e)^2 + (B*a^3 + 3*A*a^2*b)*cos(f*x + e))*(c*cos(f*x + e))^m, x)`

3.451.6 Sympy [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx = \text{Timed out}$$

input `integrate((c*cos(f*x+e))**m*(a+b*cos(f*x+e))**3*(A+B*cos(f*x+e)),x)`

output Timed out

3.451.7 Maxima [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^3 (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^3*(A+B*cos(f*x+e)),x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^3*(c*cos(f*x + e))^m, x)`

3.451.8 Giac [F]

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^3 (c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^3*(A+B*cos(f*x+e)),x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^3*(c*cos(f*x + e))^m, x)`

3.451.9 Mupad [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^3 dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^3,x)`

output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^3, x)`

3.452 $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx$

3.452.1 Optimal result	4158
3.452.2 Mathematica [A] (verified)	4159
3.452.3 Rubi [A] (verified)	4159
3.452.4 Maple [F]	4162
3.452.5 Fricas [F]	4162
3.452.6 Sympy [F(-1)]	4163
3.452.7 Maxima [F]	4163
3.452.8 Giac [F]	4163
3.452.9 Mupad [F(-1)]	4164

3.452.1 Optimal result

Integrand size = 33, antiderivative size = 287

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx$$

$$= \frac{b(Ab(3 + m) + aB(4 + m))(c \cos(e + fx))^{1+m} \sin(e + fx)}{cf(2 + m)(3 + m)} + \frac{bB(c \cos(e + fx))^{1+m} (a + b \cos(e + fx)) \sin(e + fx)}{cf(3 + m)}$$

$$- \frac{(Ab^2(1 + m) + 2abB(1 + m) + a^2A(2 + m)) (c \cos(e + fx))^{1+m} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{cf(1 + m)(2 + m)\sqrt{\sin^2(e + fx)}}$$

$$- \frac{(b^2B(2 + m) + a(2Ab + aB)(3 + m)) (c \cos(e + fx))^{2+m} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, \cos^2(e + fx)\right)}{c^2 f(2 + m)(3 + m)\sqrt{\sin^2(e + fx)}}$$

```
output b*(A*b*(3+m)+a*B*(4+m))*(c*cos(f*x+e))^(1+m)*sin(f*x+e)/c/f/(2+m)/(3+m)+b*B*(c*cos(f*x+e))^(1+m)*(a+b*cos(f*x+e))*sin(f*x+e)/c/f/(3+m)-(A*b^2*(1+m)+2*a*b*B*(1+m)+a^2*A*(2+m))*(c*cos(f*x+e))^(1+m)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], cos(f*x+e)^2)*sin(f*x+e)/c/f/(1+m)/(2+m)/(sin(f*x+e)^2)^(1/2)-(b^2*B*(2+m)+a*(2*A*b+B*a)*(3+m))*(c*cos(f*x+e))^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], cos(f*x+e)^2)*sin(f*x+e)/c^2/f/(2+m)/(3+m)/(sin(f*x+e)^2)^(1/2)
```

3.452.2 Mathematica [A] (verified)

Time = 1.08 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.74

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx$$

$$= \frac{(c \cos(e + fx))^m \cot(e + fx) \left(-\frac{a^2 A \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e+fx)\right)}{1+m} + \cos(e + fx) \left(-\frac{a(2Ab+aB) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e+fx)\right)}{1+m} \right) \right)}{f}$$

input `Integrate[(c*cos[e + f*x])^m*(a + b*cos[e + f*x])^2*(A + B*cos[e + f*x]),x]`

output `((c*cos[e + f*x])^m*Cot[e + f*x]*(-(a^2*A*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2)]/(1 + m)) + Cos[e + f*x]*(-(a*(2*A*b + a*B)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2)]/(2 + m)) + b*cos[e + f*x]*(-((A*b + 2*a*B)*Hypergeometric2F1[1/2, (3 + m)/2, (5 + m)/2, Cos[e + f*x]^2)]/(3 + m)) - (b*B*cos[e + f*x]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, Cos[e + f*x]^2)]/(4 + m))))*Sqrt[Sin[e + f*x]^2])/f`

3.452.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3469, 3042, 3502, 3042, 3227, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \cos(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^2 \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \sin \left(e + fx + \frac{\pi}{2} \right) \right)^m dx$$

$$\downarrow \text{3469}$$

$$\frac{\int (c \cos(e + fx))^m (bc(Ab(m + 3) + aB(m + 4)) \cos^2(e + fx) + c(B(m + 2)b^2 + a(2Ab + aB)(m + 3)) \cos(e + fx) + c^2(m + 3) \sin^2(e + fx)) dx}{cf(m + 3)}$$

↓ 3042

$$\frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m (bc(Ab(m+3) + aB(m+4)) \sin(e + fx + \frac{\pi}{2})^2 + c(B(m+2)b^2 + a(2Ab + aB)(m+3))) dx}{c(m+3)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))(c \cos(e + fx))^{m+1}}{cf(m+3)}$$

↓ 3502

$$\frac{\int (c \cos(e + fx))^m ((a(m+2)(bB(m+1) + aA(m+3)) + b(m+1)(Ab(m+3) + aB(m+4)))c^2 + (m+2)(B(m+2)b^2 + a(2Ab + aB)(m+3)) \cos(e + fx)c^2) dx}{c(m+2)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))(c \cos(e + fx))^{m+1}}{cf(m+3)}$$

↓ 3042

$$\frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m ((a(m+2)(bB(m+1) + aA(m+3)) + b(m+1)(Ab(m+3) + aB(m+4)))c^2 + (m+2)(B(m+2)b^2 + a(2Ab + aB)(m+3)) \sin(e + fx)c^2) dx}{c(m+2)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))(c \cos(e + fx))^{m+1}}{cf(m+3)}$$

↓ 3227

$$\frac{c(m+2)(a(m+3)(aB + 2Ab) + b^2B(m+2)) \int (c \cos(e + fx))^{m+1} dx + c^2(a(m+2)(aA(m+3) + bB(m+1)) + b(m+1)(aB(m+4) + Ab(m+3))) \int (c \cos(e + fx))^m dx}{c(m+2)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))(c \cos(e + fx))^{m+1}}{cf(m+3)}$$

↓ 3042

$$\frac{c(m+2)(a(m+3)(aB + 2Ab) + b^2B(m+2)) \int (c \sin(e + fx + \frac{\pi}{2}))^{m+1} dx + c^2(a(m+2)(aA(m+3) + bB(m+1)) + b(m+1)(aB(m+4) + Ab(m+3))) \int (c \sin(e + fx + \frac{\pi}{2}))^m dx}{c(m+2)}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))(c \cos(e + fx))^{m+1}}{cf(m+3)}$$

↓ 3122

$$\frac{\sin(e + fx)(a(m+3)(aB + 2Ab) + b^2B(m+2))(c \cos(e + fx))^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{2}, \frac{m+4}{2}, \cos^2(e + fx)\right) - c \sin(e + fx)(a(m+2)(aA(m+3) + bB(m+1)) + b^2B(m+2))}{f \sqrt{\sin^2(e + fx)}}$$

$$\frac{bB \sin(e + fx)(a + b \cos(e + fx))(c \cos(e + fx))^{m+1}}{cf(m+3)}$$

3.452. $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx$

input `Int[(c*cos[e + f*x])^m*(a + b*cos[e + f*x])^2*(A + B*cos[e + f*x]),x]`

output `(b*B*(c*cos[e + f*x])^(1 + m)*(a + b*cos[e + f*x])*Sin[e + f*x])/(c*f*(3 + m)) + ((b*(A*b*(3 + m) + a*B*(4 + m))*(c*cos[e + f*x])^(1 + m)*Sin[e + f*x])/(f*(2 + m)) + (-((c*(a*(2 + m)*(b*B*(1 + m) + a*A*(3 + m)) + b*(1 + m)*(A*b*(3 + m) + a*B*(4 + m)))*(c*cos[e + f*x])^(1 + m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*Sin[e + f*x])/(f*(1 + m)*Sqrt[Sin[e + f*x]^2])) - ((b^2*B*(2 + m) + a*(2*A*b + a*B)*(3 + m))*(c*cos[e + f*x])^(2 + m)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2]*Sin[e + f*x])/(f*Sqrt[Sin[e + f*x]^2]))/(c*(2 + m))/(c*(3 + m))`

3.452.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*cos[e + f*x]*(a + b*sin[e + f*x])^(m - 1)*((c + d*sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*sin[e + f*x])^(m - 2)*(c + d*sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`


```
rule 3502 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Co
s[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m
+ 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m
+ 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x]
&& !LtQ[m, -1]
```

3.452.4 Maple [F]

$$\int (c \cos(fx + e))^m (a + b \cos(fx + e))^2 (A + \cos(fx + e) B) dx$$

```
input int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^2*(A+cos(f*x+e)*B),x)
```

```
output int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^2*(A+cos(f*x+e)*B),x)
```

3.452.5 Fracas [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^2 (c \cos(fx + e))^m dx \end{aligned}$$

```
input integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^2*(A+B*cos(f*x+e)),x, algorithm
m="fracas")
```

```
output integral((B*b^2*cos(f*x + e)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(f*x + e)^2
+ (B*a^2 + 2*A*a*b)*cos(f*x + e))*(c*cos(f*x + e))^m, x)
```

3.452.6 Sympy [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx = \text{Timed out}$$

input `integrate((c*cos(f*x+e))**m*(a+b*cos(f*x+e))**2*(A+B*cos(f*x+e)),x)`

output `Timed out`

3.452.7 Maxima [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^2 (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^2*(A+B*cos(f*x+e)),x, algorithm m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^2*(c*cos(f*x + e))^m, x)`

3.452.8 Giac [F]

$$\begin{aligned} & \int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^2 (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^2*(A+B*cos(f*x+e)),x, algorithm m="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^2*(c*cos(f*x + e))^m, x)`

3.452.9 Mupad [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^2 dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^2,x)`output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^2, x)`

3.453 $\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$

3.453.1 Optimal result	4165
3.453.2 Mathematica [A] (verified)	4166
3.453.3 Rubi [A] (verified)	4166
3.453.4 Maple [F]	4168
3.453.5 Fracas [F]	4169
3.453.6 Sympy [F]	4169
3.453.7 Maxima [F]	4169
3.453.8 Giac [F]	4170
3.453.9 Mupad [F(-1)]	4170

3.453.1 Optimal result

Integrand size = 31, antiderivative size = 196

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$$

$$= \frac{bB(c \cos(e + fx))^{1+m} \sin(e + fx)}{cf(2 + m)}$$

$$- \frac{(bB(1 + m) + aA(2 + m))(c \cos(e + fx))^{1+m} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sin(e + fx)}{cf(1 + m)(2 + m)\sqrt{\sin^2(e + fx)}}$$

$$- \frac{(Ab + aB)(c \cos(e + fx))^{2+m} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, \cos^2(e + fx)\right) \sin(e + fx)}{c^2 f(2 + m)\sqrt{\sin^2(e + fx)}}$$

```
output b*B*(c*cos(f*x+e))^(1+m)*sin(f*x+e)/c/f/(2+m)-(b*B*(1+m)+a*A*(2+m))*(c*cos
(f*x+e))^(1+m)*hypergeom([1/2, 1/2+1/2*m],[3/2+1/2*m],cos(f*x+e)^2)*sin(f*
x+e)/c/f/(1+m)/(2+m)/(sin(f*x+e)^2)^(1/2)-(A*b+B*a)*(c*cos(f*x+e))^(2+m)*h
ypergeom([1/2, 1+1/2*m],[2+1/2*m],cos(f*x+e)^2)*sin(f*x+e)/c^2/f/(2+m)/(si
n(f*x+e)^2)^(1/2)
```

3.453.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.83

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx =$$

$$\frac{(c \cos(e + fx))^m \left((bB(1 + m) + aA(2 + m)) \cot(e + fx) \operatorname{Hypergeometric2F1} \left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx) \right) \right)}{f(1 + m)(2 + m)}$$

input `Integrate[(c*cos[e + f*x])^m*(a + b*cos[e + f*x])*(A + B*cos[e + f*x]),x]`output `-(((c*cos[e + f*x])^m*((b*B*(1 + m) + a*A*(2 + m))*Cot[e + f*x]*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*Sqrt[Sin[e + f*x]^2] + (A*b + a*B)*(1 + m)*Cos[e + f*x]*Cot[e + f*x]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2]*Sqrt[Sin[e + f*x]^2] - (b*B*(1 + m)*Sin[2*(e + f*x)]/2))/(f*(1 + m)*(2 + m)))`**3.453.3 Rubi [A] (verified)**Time = 0.65 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 3447, 3042, 3502, 3042, 3227, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \cos(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \sin \left(e + fx + \frac{\pi}{2} \right) \right)^m dx$$

$$\downarrow \text{3447}$$

$$\int (c \cos(e + fx))^m ((aB + Ab) \cos(e + fx) + aA + bB \cos^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \left(c \sin \left(e + fx + \frac{\pi}{2} \right) \right)^m \left((aB + Ab) \sin \left(e + fx + \frac{\pi}{2} \right) + aA + bB \sin \left(e + fx + \frac{\pi}{2} \right)^2 \right) dx$$

$$\begin{aligned}
 & \downarrow \text{3502} \\
 & \frac{\int (c \cos(e + fx))^m (c(bB(m + 1) + aA(m + 2)) + (Ab + aB)c(m + 2) \cos(e + fx)) dx}{c(m + 2)} + \\
 & \quad \frac{bB \sin(e + fx)(c \cos(e + fx))^{m+1}}{cf(m + 2)} \\
 & \downarrow \text{3042} \\
 & \frac{\int (c \sin(e + fx + \frac{\pi}{2}))^m (c(bB(m + 1) + aA(m + 2)) + (Ab + aB)c(m + 2) \sin(e + fx + \frac{\pi}{2})) dx}{c(m + 2)} + \\
 & \quad \frac{bB \sin(e + fx)(c \cos(e + fx))^{m+1}}{cf(m + 2)} \\
 & \downarrow \text{3227} \\
 & \frac{c(aA(m + 2) + bB(m + 1)) \int (c \cos(e + fx))^m dx + (m + 2)(aB + Ab) \int (c \cos(e + fx))^{m+1} dx}{c(m + 2)} + \\
 & \quad \frac{bB \sin(e + fx)(c \cos(e + fx))^{m+1}}{cf(m + 2)} \\
 & \downarrow \text{3042} \\
 & \frac{c(aA(m + 2) + bB(m + 1)) \int (c \sin(e + fx + \frac{\pi}{2}))^m dx + (m + 2)(aB + Ab) \int (c \sin(e + fx + \frac{\pi}{2}))^{m+1} dx}{c(m + 2)} + \\
 & \quad \frac{bB \sin(e + fx)(c \cos(e + fx))^{m+1}}{cf(m + 2)} \\
 & \downarrow \text{3122} \\
 & \frac{\frac{\sin(e + fx)(aA(m + 2) + bB(m + 1))(c \cos(e + fx))^{m+1} \text{Hypergeometric2F1}(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e + fx))}{f(m + 1)\sqrt{\sin^2(e + fx)}} - \frac{(aB + Ab) \sin(e + fx)(c \cos(e + fx))^{m+2}}{cf(m + 2)}}{c(m + 2)} \\
 & \quad \frac{bB \sin(e + fx)(c \cos(e + fx))^{m+1}}{cf(m + 2)}
 \end{aligned}$$

input `Int[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])*(A + B*Cos[e + f*x]),x]`

output `(b*B*(c*Cos[e + f*x])^(1 + m)*Sin[e + f*x])/(c*f*(2 + m)) + (-(((b*B*(1 + m) + a*A*(2 + m))*(c*Cos[e + f*x])^(1 + m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*Sin[e + f*x])/(f*(1 + m)*Sqrt[Sin[e + f*x]^2])) - ((A*b + a*B)*(c*Cos[e + f*x])^(2 + m)*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, Cos[e + f*x]^2]*Sin[e + f*x])/(c*f*Sqrt[Sin[e + f*x]^2]))/(c*(2 + m))`

3.453. $\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$

3.453.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 3227 `Int[((b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Sin[e + f*x])^m, x], x] + Simp[d/b Int[(b*Sin[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x]`

rule 3447 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Int[(a + b*Sin[e + f*x])^m*(A*c + (B*c + A*d)*Sin[e + f*x] + B*d*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0]`

rule 3502 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(m + 2))), x] + Simp[1/(b*(m + 2)) Int[(a + b*Sin[e + f*x])^m*Simp[A*b*(m + 2) + b*C*(m + 1) + (b*B*(m + 2) - a*C)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && !LtQ[m, -1]`

3.453.4 Maple [F]

$$\int (c \cos(fx + e))^m (a + b \cos(fx + e)) (A + \cos(fx + e) B) dx$$

input `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))*(A+cos(f*x+e)*B),x)`

output `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))*(A+cos(f*x+e)*B),x)`

3.453.5 Fricas [F]

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)(c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))*(A+B*cos(f*x+e)),x, algorithm="fricas")`

output `integral((B*b*cos(f*x + e)^2 + A*a + (B*a + A*b)*cos(f*x + e))*(c*cos(f*x + e))^m, x)`

3.453.6 Sympy [F]

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx))(a + b \cos(e + fx)) dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))*(A+B*cos(f*x+e)),x)`

output `Integral((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x)), x)`

3.453.7 Maxima [F]

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)(c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))*(A+B*cos(f*x+e)),x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)*(c*cos(f*x + e))^m, x)`

3.453.8 Giac [F]

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)(c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))*(A+B*cos(f*x+e)),x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)*(c*cos(f*x + e))^m, x)`

3.453.9 Mupad [F(-1)]

Timed out.

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))(A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx)) dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x)),x)`

output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x)), x)`

3.454 $\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{a+b \cos(e+fx)} dx$

3.454.1 Optimal result 4171
 3.454.2 Mathematica [B] (warning: unable to verify) 4172
 3.454.3 Rubi [A] (verified) 4172
 3.454.4 Maple [F] 4175
 3.454.5 Fricas [F] 4175
 3.454.6 Sympy [F(-1)] 4175
 3.454.7 Maxima [F] 4176
 3.454.8 Giac [F(-2)] 4176
 3.454.9 Mupad [F(-1)] 4176

3.454.1 Optimal result

Integrand size = 33, antiderivative size = 286

$$\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{a+b \cos(e+fx)} dx$$

$$= \frac{a(Ab - aB)c \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1-m}{2}, 1, \frac{3}{2}, \sin^2(e+fx), -\frac{b^2 \sin^2(e+fx)}{a^2-b^2}\right) (c \cos(e+fx))^{-1+m} \cos^2(e+fx)^{\frac{1-m}{2}} \sin(e+fx)}{b(a^2 - b^2) f}$$

$$- \frac{(Ab - aB) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{m}{2}, 1, \frac{3}{2}, \sin^2(e+fx), -\frac{b^2 \sin^2(e+fx)}{a^2-b^2}\right) (c \cos(e+fx))^m \cos^2(e+fx)^{-m/2} \sin(e+fx)}{(a^2 - b^2) f}$$

$$- \frac{B(c \cos(e+fx))^{1+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e+fx)\right) \sin(e+fx)}{bcf(1+m)\sqrt{\sin^2(e+fx)}}$$

output

```
a*(A*b-B*a)*c*AppellF1(1/2,-1/2*m+1/2,1,3/2,sin(f*x+e)^2,-b^2*sin(f*x+e)^2/(a^2-b^2))*(c*cos(f*x+e))^(1+m)*(cos(f*x+e)^2)^(-1/2*m+1/2)*sin(f*x+e)/b/(a^2-b^2)/f-(A*b-B*a)*AppellF1(1/2,-1/2*m,1,3/2,sin(f*x+e)^2,-b^2*sin(f*x+e)^2/(a^2-b^2))*(c*cos(f*x+e))^m*sin(f*x+e)/(a^2-b^2)/f/((cos(f*x+e)^2)^(1/2*m))-B*(c*cos(f*x+e))^(1+m)*hypergeom([1/2, 1/2+1/2*m],[3/2+1/2*m],cos(f*x+e)^2)*sin(f*x+e)/b/c/f/(1+m)/(sin(f*x+e)^2)^(1/2)
```

3.454.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 10482 vs. $2(286) = 572$.

Time = 30.99 (sec) , antiderivative size = 10482, normalized size of antiderivative = 36.65

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx = \text{Result too large to show}$$

input `Integrate[((c*cos[e + f*x])^m*(A + B*cos[e + f*x]))/(a + b*cos[e + f*x]),x]`

output `Result too large to show`

3.454.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.98, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 3481, 3042, 3122, 3302, 3042, 3668, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + B \cos(e + fx))(c \cos(e + fx))^m}{a + b \cos(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(A + B \sin(e + fx + \frac{\pi}{2}))(c \sin(e + fx + \frac{\pi}{2}))^m}{a + b \sin(e + fx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{3481} \\ & \frac{(Ab - aB) \int \frac{(c \cos(e + fx))^m}{a + b \cos(e + fx)} dx}{b} + \frac{B \int (c \cos(e + fx))^m dx}{b} \\ & \quad \downarrow \text{3042} \\ & \frac{(Ab - aB) \int \frac{(c \sin(e + fx + \frac{\pi}{2}))^m}{a + b \sin(e + fx + \frac{\pi}{2})} dx}{b} + \frac{B \int (c \sin(e + fx + \frac{\pi}{2}))^m dx}{b} \\ & \quad \downarrow \text{3122} \end{aligned}$$

$$\frac{(Ab - aB) \int \frac{(c \sin(e+fx+\frac{\pi}{2}))^m}{a+b \sin(e+fx+\frac{\pi}{2})} dx}{b}$$

$$\frac{B \sin(e+fx)(c \cos(e+fx))^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{bcf(m+1)\sqrt{\sin^2(e+fx)}}$$

↓ 3302

$$\frac{(Ab - aB) \left(a \int \frac{(c \cos(e+fx))^m}{a^2-b^2 \cos^2(e+fx)} dx - \frac{b \int \frac{(c \cos(e+fx))^{m+1}}{a^2-b^2 \cos^2(e+fx)} dx}{c} \right)}{b}$$

$$\frac{B \sin(e+fx)(c \cos(e+fx))^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{bcf(m+1)\sqrt{\sin^2(e+fx)}}$$

↓ 3042

$$\frac{(Ab - aB) \left(a \int \frac{(c \sin(e+fx+\frac{\pi}{2}))^m}{a^2-b^2 \sin^2(e+fx+\frac{\pi}{2})} dx - \frac{b \int \frac{(c \sin(e+fx+\frac{\pi}{2}))^{m+1}}{a^2-b^2 \sin^2(e+fx+\frac{\pi}{2})} dx}{c} \right)}{b}$$

$$\frac{B \sin(e+fx)(c \cos(e+fx))^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{bcf(m+1)\sqrt{\sin^2(e+fx)}}$$

↓ 3668

$$(Ab - aB) \left(\frac{ac \cos^2(e+fx)^{\frac{1-m}{2}} (c \cos(e+fx))^{m-1} \int \frac{(1-\sin^2(e+fx))^{\frac{m-1}{2}}}{a^2-b^2+b^2 \sin^2(e+fx)} d \sin(e+fx)}{f} - \frac{b \cos^2(e+fx)^{-m/2} (c \cos(e+fx))^m \int \frac{(1-\sin^2(e+fx))^{\frac{m-1}{2}}}{a^2-b^2+b^2 \sin^2(e+fx)} d \sin(e+fx)}{f} \right)$$

$$\frac{B \sin(e+fx)(c \cos(e+fx))^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{bcf(m+1)\sqrt{\sin^2(e+fx)}}$$

↓ 333

$$(Ab - aB) \left(\frac{ac \sin(e+fx) \cos^2(e+fx)^{\frac{1-m}{2}} (c \cos(e+fx))^{m-1} \text{AppellF1}\left(\frac{1}{2}, \frac{1-m}{2}, 1, \frac{3}{2}, \sin^2(e+fx), -\frac{b^2 \sin^2(e+fx)}{a^2-b^2}\right)}{f(a^2-b^2)} - \frac{b \sin(e+fx) \cos^2(e+fx)}{f} \right)$$

$$\frac{B \sin(e+fx)(c \cos(e+fx))^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{bcf(m+1)\sqrt{\sin^2(e+fx)}}$$

input `Int[((c*cos[e + f*x])^m*(A + B*cos[e + f*x]))/(a + b*cos[e + f*x]),x]`

3.454. $\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{a+b \cos(e+fx)} dx$

```
output -((B*(c*cos[e + f*x])^(1 + m)*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2,
  Cos[e + f*x]^2*Sin[e + f*x])/(b*c*f*(1 + m)*Sqrt[Sin[e + f*x]^2])) + ((A
  *b - a*B)*((a*c*AppellF1[1/2, (1 - m)/2, 1, 3/2, Sin[e + f*x]^2, -((b^2*Si
  n[e + f*x]^2)/(a^2 - b^2))]*(c*cos[e + f*x])^(-1 + m)*(Cos[e + f*x]^2)^((1
  - m)/2)*Sin[e + f*x])/(a^2 - b^2)*f - (b*AppellF1[1/2, -1/2*m, 1, 3/2,
  Sin[e + f*x]^2, -((b^2*Sin[e + f*x]^2)/(a^2 - b^2))]*(c*cos[e + f*x])^m*Si
  n[e + f*x])/(a^2 - b^2)*f*(Cos[e + f*x]^2)^(m/2)))/b
```

3.454.3.1 Defintions of rubi rules used

```
rule 333 Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; F
reeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,
0]) && (IntegerQ[q] || GtQ[c, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3122 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

```
rule 3302 Int[((d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)/((a_) + (b_.)*sin[(e_.) + (f_.)*(
x_)]), x_Symbol] := Simp[a Int[(d*Sin[e + f*x])^n/(a^2 - b^2*Sin[e + f*x]
^2), x], x] - Simp[b/d Int[(d*Sin[e + f*x])^(n + 1)/(a^2 - b^2*Sin[e + f*
x]^2), x], x] /; FreeQ[{a, b, d, e, f, n}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 3481 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.)
+ (f_.)*(x_)])/(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[
B/d Int[(a + b*Sin[e + f*x])^m, x], x] - Simp[(B*c - A*d)/d Int[(a + b*
Sin[e + f*x])^m/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A,
B, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3668 Int[((d_.)*sin[(e_.) + (f_.)*(x_)]^(m_))*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Simp[(-ff)*d^(2*IntPart[(m - 1)/2] + 1)*((d*SIN[e + f*x])^(2*FracPart[(m - 1)/2])/(f*(Sin[e + f*x]^2)^(FracPart[(m - 1)/2])) Subst[Int[(1 - ff^2*x^2)^(m - 1)/2]*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, d, e, f, m, p}, x] && !IntegerQ[m]
```

3.454.4 Maple [F]

$$\int \frac{(c \cos (fx + e))^m (A + \cos (fx + e) B)}{a + b \cos (fx + e)} dx$$

```
input int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)/(a+b*cos(f*x+e)),x)
```

```
output int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)/(a+b*cos(f*x+e)),x)
```

3.454.5 Fricas [F]

$$\int \frac{(c \cos (e + fx))^m (A + B \cos (e + fx))}{a + b \cos (e + fx)} dx = \int \frac{(B \cos (fx + e) + A)(c \cos (fx + e))^m}{b \cos (fx + e) + a} dx$$

```
input integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e)),x, algorithm="fricas")
```

```
output integral((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/(b*cos(f*x + e) + a), x)
```

3.454.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(c \cos (e + fx))^m (A + B \cos (e + fx))}{a + b \cos (e + fx)} dx = \text{Timed out}$$

```
input integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e)),x)
```

```
output Timed out
```

3.454. $\int \frac{(c \cos (e + fx))^m (A + B \cos (e + fx))}{a + b \cos (e + fx)} dx$

3.454.7 Maxima [F]

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{b \cos(fx + e) + a} dx$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e)),x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/(b*cos(f*x + e) + a), x)`

3.454.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx = \text{Exception raised: TypeError}$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{-1,[0,1,0,0]%%} / %%{1,[0,0,1,0]%%}+%%{-1,[0,0,0,1]%%} } Error:`

3.454.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx = \int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx$$

input `int(((c*cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x)),x)`

output `int(((c*cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x)), x)`

3.455 $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx$

3.455.1 Optimal result	4177
3.455.2 Mathematica [N/A]	4178
3.455.3 Rubi [N/A]	4178
3.455.4 Maple [N/A] (verified)	4180
3.455.5 Fricas [N/A]	4180
3.455.6 Sympy [N/A]	4181
3.455.7 Maxima [N/A]	4181
3.455.8 Giac [N/A]	4182
3.455.9 Mupad [N/A]	4182

3.455.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \frac{2bB(c \cos(e + fx))^{1+m} \sqrt{a + b \cos(e + fx)} \sin(e + fx)}{cf(5 + 2m)} + \frac{2 \operatorname{Int} \left(\frac{(c \cos(e + fx))^m \left(\frac{1}{2} ac(2bB(1+m) + 2aA(\frac{5}{2} + m)) + \frac{1}{2} c(b^2 B(3+2m) + a(2Ab + aB)(5+2m)) \cos(e + fx) + \frac{1}{2} bc(2aB(3+m) + Ab(5+2m)) \cos^2(e + fx) \right)}{\sqrt{a + b \cos(e + fx)}} \right)}{c(5 + 2m)}$$

output

```
2*b*B*(c*cos(f*x+e))^(1+m)*sin(f*x+e)*(a+b*cos(f*x+e))^(1/2)/c/f/(5+2*m)+2
*Unintegrable((c*cos(f*x+e))^m*(1/2*a*c*(2*b*B*(1+m)+2*a*A*(5/2+m))+1/2*c*
(b^2*B*(3+2*m)+a*(2*A*b+B*a)*(5+2*m))*cos(f*x+e)+1/2*b*c*(2*a*B*(3+m)+A*b*
(5+2*m))*cos(f*x+e)^2)/(a+b*cos(f*x+e))^(1/2),x)/c/(5+2*m)
```


3.455.2 Mathematica [N/A]

Not integrable

Time = 79.33 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx$$

input `Integrate[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^(3/2)*(A + B*Cos[e + f*x]),x]`

output `Integrate[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^(3/2)*(A + B*Cos[e + f*x]), x]`

3.455.3 Rubi [N/A]

Not integrable

Time = 0.84 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3469, 27, 3042, 3544}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \cos(e + fx))^m dx$$

↓ 3042

$$\int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^{3/2} \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \sin \left(e + fx + \frac{\pi}{2} \right) \right)^m dx$$

↓ 3469

$$2 \int \frac{(c \cos(e+fx))^m (bc(2aB(m+3)+Ab(2m+5)) \cos^2(e+fx)+c(B(2m+3)b^2+a(2Ab+aB)(2m+5)) \cos(e+fx)+ac(2bB(m+1)+aA(2m+5)))}{2\sqrt{a+b \cos(e+fx)}} dx$$

$$\frac{c(2m+5)}{2bB \sin(e+fx) \sqrt{a+b \cos(e+fx)} (c \cos(e+fx))^{m+1}} \frac{cf(2m+5)}{cf(2m+5)}$$

↓ 27

3.455. $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx$

$$\int \frac{(c \cos(e+fx))^m (bc(2aB(m+3)+Ab(2m+5)) \cos^2(e+fx)+c(B(2m+3)b^2+a(2Ab+aB)(2m+5)) \cos(e+fx)+ac(2bB(m+1)+aA(2m+5)))}{\sqrt{a+b \cos(e+fx)}} dx$$

$$\frac{c(2m+5)}{2bB \sin(e+fx) \sqrt{a+b \cos(e+fx)} (c \cos(e+fx))^{m+1}} \frac{1}{cf(2m+5)}$$

↓ 3042

$$\int \frac{(c \sin(e+fx+\frac{\pi}{2}))^m (bc(2aB(m+3)+Ab(2m+5)) \sin^2(e+fx+\frac{\pi}{2})+c(B(2m+3)b^2+a(2Ab+aB)(2m+5)) \sin(e+fx+\frac{\pi}{2})+ac(2bB(m+1)+aA(2m+5)))}{\sqrt{a+b \sin(e+fx+\frac{\pi}{2})}} dx$$

$$\frac{c(2m+5)}{2bB \sin(e+fx) \sqrt{a+b \cos(e+fx)} (c \cos(e+fx))^{m+1}} \frac{1}{cf(2m+5)}$$

↓ 3544

$$\int \frac{(c \cos(e+fx))^m (bc(2aB(m+3)+Ab(2m+5)) \cos^2(e+fx)+c(B(2m+3)b^2+a(2Ab+aB)(2m+5)) \cos(e+fx)+ac(2bB(m+1)+aA(2m+5)))}{\sqrt{a+b \cos(e+fx)}} dx$$

$$\frac{c(2m+5)}{2bB \sin(e+fx) \sqrt{a+b \cos(e+fx)} (c \cos(e+fx))^{m+1}} \frac{1}{cf(2m+5)}$$

input `Int[(c*Cos[e + f*x])^m*(a + b*Cos[e + f*x])^(3/2)*(A + B*Cos[e + f*x]),x]`

output `$Aborted`

3.455.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3544 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.455.4 Maple [N/A] (verified)

Not integrable

Time = 0.79 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int (c \cos(fx + e))^m (a + b \cos(fx + e))^{\frac{3}{2}} (A + \cos(fx + e) B) dx$$

input `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^(3/2)*(A+cos(f*x+e)*B),x)`

output `int((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^(3/2)*(A+cos(f*x+e)*B),x)`

3.455.5 Fricas [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^{\frac{3}{2}} (c \cos(fx + e))^m dx$$

3.455. $\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^(3/2)*(A+B*cos(f*x+e)),x, algo
rithm="fricas")`

output `integral((B*b*cos(f*x + e)^2 + A*a + (B*a + A*b)*cos(f*x + e))*sqrt(b*cos(
f*x + e) + a)*(c*cos(f*x + e))^m, x)`

3.455.6 Sympy [N/A]

Not integrable

Time = 147.50 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^{\frac{3}{2}} dx$$

input `integrate((c*cos(f*x+e))**m*(a+b*cos(f*x+e))**(3/2)*(A+B*cos(f*x+e)),x)`

output `Integral((c*cos(e + f*x))**m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))**(3
/2), x)`

3.455.7 Maxima [N/A]

Not integrable

Time = 2.59 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^{\frac{3}{2}} (c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^(3/2)*(A+B*cos(f*x+e)),x, algo
rithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^(3/2)*(c*cos(f*x + e))
^m, x)`

3.455.8 Giac [N/A]

Not integrable

Time = 16.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^{3/2} (c \cos(fx + e))^m dx$$

input `integrate((c*cos(f*x+e))^m*(a+b*cos(f*x+e))^(3/2)*(A+B*cos(f*x+e)),x, algorith="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^(3/2)*(c*cos(f*x + e))^m, x)`

3.455.9 Mupad [N/A]

Not integrable

Time = 4.40 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (c \cos(e + fx))^m (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx = \int (c \cos(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^{3/2} dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(3/2),x)`

output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(3/2), x)`

3.456 $\int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx$

3.456.1 Optimal result	4183
3.456.2 Mathematica [N/A]	4183
3.456.3 Rubi [N/A]	4184
3.456.4 Maple [N/A] (verified)	4185
3.456.5 Fricas [N/A]	4185
3.456.6 Sympy [N/A]	4185
3.456.7 Maxima [N/A]	4186
3.456.8 Giac [N/A]	4186
3.456.9 Mupad [N/A]	4187

3.456.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx$$

$$= \text{Int}\left((c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)), x\right)$$

output `Unintegrable((c*cos(f*x+e))^m*(A+B*cos(f*x+e))*(a+b*cos(f*x+e))^(1/2),x)`

3.456.2 Mathematica [N/A]

Not integrable

Time = 28.85 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx$$

input `Integrate[(c*Cos[e + f*x])^m*Sqrt[a + b*Cos[e + f*x]]*(A + B*Cos[e + f*x]),x]`

output `Integrate[(c*Cos[e + f*x])^m*Sqrt[a + b*Cos[e + f*x]]*(A + B*Cos[e + f*x]), x]`

3.456.3 Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3486}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \cos(e + fx))^m dx$$

↓ 3042

$$\int \sqrt{a + b \sin\left(e + fx + \frac{\pi}{2}\right)}\left(A + B \sin\left(e + fx + \frac{\pi}{2}\right)\right)\left(c \sin\left(e + fx + \frac{\pi}{2}\right)\right)^m dx$$

↓ 3486

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \cos(e + fx))^m dx$$

input `Int[(c*cos[e + f*x])^m*Sqrt[a + b*cos[e + f*x]]*(A + B*cos[e + f*x]),x]`

output `$Aborted`

3.456.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3486 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(A + B*Sin[e + f*x])*(c + d*Sin[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.456.4 Maple [N/A] (verified)

Not integrable

Time = 0.80 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int (c \cos(fx + e))^m (A + \cos(fx + e) B) \sqrt{a + b \cos(fx + e)} dx$$

input `int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)*(a+b*cos(f*x+e))^(1/2),x)`

output `int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)*(a+b*cos(f*x+e))^(1/2),x)`

3.456.5 Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A) \sqrt{b \cos(fx + e) + a} (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))*(a+b*cos(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*cos(f*x + e))^m, x)`

3.456.6 Sympy [N/A]

Not integrable

Time = 7.32 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\begin{aligned} & \int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx \\ &= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) \sqrt{a + b \cos(e + fx)} dx \end{aligned}$$

input `integrate((c*cos(f*x+e))**m*(A+B*cos(f*x+e))*(a+b*cos(f*x+e))**(1/2),x)`

output `Integral((c*cos(e + f*x))**m*(A + B*cos(e + f*x))*sqrt(a + b*cos(e + f*x)), x)`

3.456.7 Maxima [N/A]

Not integrable

Time = 2.59 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A) \sqrt{b \cos(fx + e) + a} (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))*(a+b*cos(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*cos(f*x + e))^m, x)`

3.456.8 Giac [N/A]

Not integrable

Time = 0.96 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx \\ &= \int (B \cos(fx + e) + A) \sqrt{b \cos(fx + e) + a} (c \cos(fx + e))^m dx \end{aligned}$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))*(a+b*cos(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*cos(f*x + e))^m, x)`

3.456.9 Mupad [N/A]

Not integrable

Time = 2.72 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (c \cos(e + fx))^m \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx$$

$$= \int (c \cos(e + fx))^m (A + B \cos(e + fx)) \sqrt{a + b \cos(e + fx)} dx$$

input `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(1/2),x)`output `int((c*cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(1/2), x)`

$$3.457 \quad \int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx$$

3.457.1 Optimal result	4188
3.457.2 Mathematica [N/A]	4188
3.457.3 Rubi [N/A]	4189
3.457.4 Maple [N/A] (verified)	4190
3.457.5 Fricas [N/A]	4190
3.457.6 Sympy [N/A]	4190
3.457.7 Maxima [N/A]	4191
3.457.8 Giac [N/A]	4191
3.457.9 Mupad [N/A]	4191

3.457.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx = \text{Int} \left(\frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}}, x \right)$$

output `Unintegrable((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(1/2),x)`

3.457.2 Mathematica [N/A]

Not integrable

Time = 34.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx = \int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx$$

input `Integrate[((c*Cos[e + f*x])^m*(A + B*Cos[e + f*x]))/Sqrt[a + b*Cos[e + f*x]],x]`

output `Integrate[((c*Cos[e + f*x])^m*(A + B*Cos[e + f*x]))/Sqrt[a + b*Cos[e + f*x]], x]`

$$3.457. \quad \int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx$$

3.457.3 Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3486}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \cos(e + fx))(c \cos(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx$$

↓ 3042

$$\int \frac{(A + B \sin(e + fx + \frac{\pi}{2}))(c \sin(e + fx + \frac{\pi}{2}))^m}{\sqrt{a + b \sin(e + fx + \frac{\pi}{2})}} dx$$

↓ 3486

$$\int \frac{(A + B \cos(e + fx))(c \cos(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx$$

input `Int[((c*Cos[e + f*x])^m*(A + B*Cos[e + f*x]))/Sqrt[a + b*Cos[e + f*x]],x]`

output `$Aborted`

3.457.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3486 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(A + B*Sin[e + f*x])*(c + d*Sin[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.457.4 Maple [N/A] (verified)

Not integrable

Time = 0.92 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(c \cos(fx + e))^m (A + \cos(fx + e) B)}{\sqrt{a + b \cos(fx + e)}} dx$$

```
input int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)/(a+b*cos(f*x+e))^(1/2),x)
```

```
output int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)/(a+b*cos(f*x+e))^(1/2),x)
```

3.457.5 Fracas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{\sqrt{b \cos(fx + e) + a}} dx$$

```
input integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(1/2),x, algo
rithm="fracas")
```

```
output integral((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/sqrt(b*cos(f*x + e) + a),
x)
```

3.457.6 Sympy [N/A]

Not integrable

Time = 3.76 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$$

```
input integrate((c*cos(f*x+e))**m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))**(1/2),x)
```

```
output Integral((c*cos(e + f*x))**m*(A + B*cos(e + f*x))/sqrt(a + b*cos(e + f*x))
, x)
```

3.457. $\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx$

3.457.7 Maxima [N/A]

Not integrable

Time = 2.42 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{\sqrt{b \cos(fx + e) + a}} dx$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/sqrt(b*cos(f*x + e) + a)
, x)`

3.457.8 Giac [N/A]

Not integrable

Time = 0.74 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{\sqrt{b \cos(fx + e) + a}} dx$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/sqrt(b*cos(f*x + e) + a)
, x)`

3.457.9 Mupad [N/A]

Not integrable

Time = 4.31 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$$

input `int(((c*cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(1/2),x)`

output `int(((c*cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(1/2),x)`

3.457. $\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}} dx$

3.458
$$\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{(a+b \cos(e+fx))^{3/2}} dx$$

3.458.1 Optimal result 4193
 3.458.2 Mathematica [N/A] 4193
 3.458.3 Rubi [N/A] 4194
 3.458.4 Maple [N/A] (verified) 4196
 3.458.5 Fricas [N/A] 4196
 3.458.6 Sympy [N/A] 4197
 3.458.7 Maxima [N/A] 4197
 3.458.8 Giac [N/A] 4197
 3.458.9 Mupad [N/A] 4198

3.458.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{(a+b \cos(e+fx))^{3/2}} dx = \frac{2b(Ab-aB)(c \cos(e+fx))^{1+m} \sin(e+fx)}{a(a^2-b^2)cf\sqrt{a+b \cos(e+fx)}} + \frac{2 \operatorname{Int}\left(\frac{(c \cos(e+fx))^m \left(\frac{1}{2}c(aA-bB)+2b(Ab-aB)\left(\frac{1}{2}+m\right)\right) - \frac{1}{2}a(Ab-aB)c \cos(e+fx) - \frac{1}{2}b(Ab-aB)c(3+2m) \cos^2(e+fx)}{\sqrt{a+b \cos(e+fx)}}, x\right)}{a(a^2-b^2)c}$$

output `2*b*(A*b-B*a)*(c*cos(f*x+e))^(1+m)*sin(f*x+e)/a/(a^2-b^2)/c/f/(a+b*cos(f*x+e))^(1/2)+2*Unintegrable((c*cos(f*x+e))^m*(1/2*c*(a*(A*a-B*b)+2*b*(A*b-B*a)*(1/2+m))-1/2*a*(A*b-B*a)*c*cos(f*x+e)-1/2*b*(A*b-B*a)*c*(3+2*m)*cos(f*x+e)^2)/(a+b*cos(f*x+e))^(1/2),x)/a/(a^2-b^2)/c`

3.458.2 Mathematica [N/A]

Not integrable

Time = 32.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{(a+b \cos(e+fx))^{3/2}} dx = \int \frac{(c \cos(e+fx))^m (A+B \cos(e+fx))}{(a+b \cos(e+fx))^{3/2}} dx$$

input `Integrate[((c*cos[e + f*x])^m*(A + B*cos[e + f*x]))/(a + b*cos[e + f*x])^(3/2),x]`

output `Integrate[((c*cos[e + f*x])^m*(A + B*cos[e + f*x]))/(a + b*cos[e + f*x])^(3/2), x]`

3.458.3 Rubi [N/A]

Not integrable

Time = 0.85 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3479, 27, 3042, 3544}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \cos(e + fx))(c \cos(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(A + B \sin(e + fx + \frac{\pi}{2}))(c \sin(e + fx + \frac{\pi}{2}))^m}{(a + b \sin(e + fx + \frac{\pi}{2}))^{3/2}} dx$$

↓ 3479

$$2 \int \frac{(c \cos(e + fx))^m (-b(Ab - aB)c(2m + 3) \cos^2(e + fx) - a(Ab - aB)c \cos(e + fx) + c(Aa^2 - 2bB(m + 1)a + Ab^2(2m + 1)))}{2\sqrt{a + b \cos(e + fx)}} dx +$$

$$\frac{ac(a^2 - b^2)}{2b(Ab - aB) \sin(e + fx)(c \cos(e + fx))^{m+1}} \frac{2b(Ab - aB) \sin(e + fx)(c \cos(e + fx))^{m+1}}{acf(a^2 - b^2) \sqrt{a + b \cos(e + fx)}}$$

↓ 27

$$\int \frac{(c \cos(e + fx))^m (-b(Ab - aB)c(2m + 3) \cos^2(e + fx) - a(Ab - aB)c \cos(e + fx) + c(Aa^2 - 2bB(m + 1)a + Ab^2(2m + 1)))}{\sqrt{a + b \cos(e + fx)}} dx +$$

$$\frac{ac(a^2 - b^2)}{2b(Ab - aB) \sin(e + fx)(c \cos(e + fx))^{m+1}} \frac{2b(Ab - aB) \sin(e + fx)(c \cos(e + fx))^{m+1}}{acf(a^2 - b^2) \sqrt{a + b \cos(e + fx)}}$$

↓ 3042

$$\int \frac{(c \sin(e + fx + \frac{\pi}{2}))^m (-b(Ab - aB)c(2m + 3) \sin^2(e + fx + \frac{\pi}{2}) - a(Ab - aB)c \sin(e + fx + \frac{\pi}{2}) + c(Aa^2 - 2bB(m + 1)a + Ab^2(2m + 1)))}{\sqrt{a + b \sin(e + fx + \frac{\pi}{2})}} dx +$$

$$\frac{ac(a^2 - b^2)}{2b(Ab - aB) \sin(e + fx)(c \cos(e + fx))^{m+1}} \frac{2b(Ab - aB) \sin(e + fx)(c \cos(e + fx))^{m+1}}{acf(a^2 - b^2) \sqrt{a + b \cos(e + fx)}}$$

3.458. $\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx$

$$\int \frac{(c \cos(e+fx))^m (-b(Ab-aB)c(2m+3) \cos^2(e+fx) - a(Ab-aB)c \cos(e+fx) + c(Aa^2 - 2bB(m+1)a + Ab^2(2m+1)))}{\sqrt{a+b \cos(e+fx)}} dx + \frac{ac(a^2 - b^2)}{2b(Ab - aB) \sin(e+fx)(c \cos(e+fx))^{m+1}} + \frac{2b(Ab - aB) \sin(e+fx)(c \cos(e+fx))^{m+1}}{acf(a^2 - b^2) \sqrt{a+b \cos(e+fx)}}$$

input `Int[((c*Cos[e + f*x])^m*(A + B*Cos[e + f*x]))/(a + b*Cos[e + f*x])^(3/2),x]`

output `$Aborted`

3.458.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3544 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Unintegrable[(a + b*Sin[e + f*x])^m*(c + d*
Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x] /; FreeQ[{a, b,
c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0]
```

3.458.4 Maple [N/A] (verified)

Not integrable

Time = 0.90 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(c \cos(fx + e))^m (A + \cos(fx + e) B)}{(a + b \cos(fx + e))^{\frac{3}{2}}} dx$$

```
input int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)/(a+b*cos(f*x+e))^(3/2),x)
```

```
output int((c*cos(f*x+e))^m*(A+cos(f*x+e)*B)/(a+b*cos(f*x+e))^(3/2),x)
```

3.458.5 Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{(b \cos(fx + e) + a)^{\frac{3}{2}}} dx$$

```
input integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(3/2),x, algo
rithm="fricas")
```

```
output integral((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*cos(f*x + e))^m/
(b^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + a^2), x)
```

3.458.6 Sympy [N/A]

Not integrable

Time = 9.25 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{\frac{3}{2}}} dx$$

input `integrate((c*cos(f*x+e))**m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))**(3/2),x)`output `Integral((c*cos(e + f*x))**m*(A + B*cos(e + f*x))/(a + b*cos(e + f*x))**(3/2), x)`**3.458.7 Maxima [N/A]**

Not integrable

Time = 2.59 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{(b \cos(fx + e) + a)^{\frac{3}{2}}} dx$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(3/2),x, algorith="maxima")`output `integrate((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/(b*cos(f*x + e) + a)^(3/2), x)`**3.458.8 Giac [N/A]**

Not integrable

Time = 1.29 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(B \cos(fx + e) + A)(c \cos(fx + e))^m}{(b \cos(fx + e) + a)^{\frac{3}{2}}} dx$$

input `integrate((c*cos(f*x+e))^m*(A+B*cos(f*x+e))/(a+b*cos(f*x+e))^(3/2),x, algorith="giac")`

output `integrate((B*cos(f*x + e) + A)*(c*cos(f*x + e))^m/(b*cos(f*x + e) + a)^(3/2), x)`

3.458.9 Mupad [N/A]

Not integrable

Time = 6.39 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(c \cos(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx$$

input `int(((c*cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(3/2),x)`

output `int(((c*cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(3/2),x)`

3.459 $\int (a+a \cos(c+dx))(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

3.459.1 Optimal result	4199
3.459.2 Mathematica [C] (verified)	4200
3.459.3 Rubi [A] (verified)	4200
3.459.4 Maple [B] (verified)	4204
3.459.5 Fricas [C] (verification not implemented)	4205
3.459.6 Sympy [F(-1)]	4206
3.459.7 Maxima [F]	4206
3.459.8 Giac [F]	4207
3.459.9 Mupad [F(-1)]	4207

3.459.1 Optimal result

Integrand size = 31, antiderivative size = 172

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= -\frac{2a(3A + 5B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d} \\ & \quad + \frac{2a(A + B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d} \\ & \quad + \frac{2a(3A + 5B)\sqrt{\sec(c + dx)}\sin(c + dx)}{5d} \\ & \quad + \frac{2a(A + B)\sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{3d} + \frac{2aA\sec^{\frac{5}{2}}(c + dx)\sin(c + dx)}{5d} \end{aligned}$$

output

```
2/3*a*(A+B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/5*a*A*sec(d*x+c)^(5/2)*sin(d*x+c)/d+2/5*a*(3*A+5*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2/5*a*(3*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.459.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.73 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.70

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{ae^{-ic}(-1 + e^{2ic})(1 + \cos(c + dx)) \csc(c) \left(5A + 5B - 3Ae^{i(c+dx)} - 15Be^{i(c+dx)} - 24Ae^{3i(c+dx)} - 30Be^{3i(c+dx)}\right)}{}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(a*(-1 + E^((2*I)*c))*(1 + Cos[c + d*x])*Csc[c]*(5*A + 5*B - 3*A*E^(I*(c + d*x)) - 15*B*E^(I*(c + d*x)) - 24*A*E^((3*I)*(c + d*x)) - 30*B*E^((3*I)*(c + d*x)) - 5*A*E^((4*I)*(c + d*x)) - 5*B*E^((4*I)*(c + d*x)) - 9*A*E^((5*I)*(c + d*x)) - 15*B*E^((5*I)*(c + d*x)) - (5*I)*(A + B)*(1 + E^((2*I)*(c + d*x))))^2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (3*A + 5*B)*E^(I*(c + d*x))*(1 + E^((2*I)*(c + d*x)))^(5/2)*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*Sec[(c + d*x)/2]^2*sqrt[Sec[c + d*x]]/(30*d*E^(I*c)*(1 + E^((2*I)*(c + d*x)))^2)`

3.459.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.97, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4485, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{7}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)(A \sec(c + dx) + B) dx$$

3.459. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\begin{aligned} & \downarrow 3042 \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A \csc\left(c + dx + \frac{\pi}{2}\right) + B\right) dx \\ & \downarrow 4485 \\ & \frac{2}{5} \int \frac{1}{2} \sec^{3/2}(c + dx) (a(3A + 5B) + 5a(A + B) \sec(c + dx)) dx + \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 27 \\ & \frac{1}{5} \int \sec^{3/2}(c + dx) (a(3A + 5B) + 5a(A + B) \sec(c + dx)) dx + \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 3042 \\ & \frac{1}{5} \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a(3A + 5B) + 5a(A + B) \csc\left(c + dx + \frac{\pi}{2}\right)\right) dx + \\ & \quad \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 4274 \\ & \frac{1}{5} \left(a(3A + 5B) \int \sec^{3/2}(c + dx) dx + 5a(A + B) \int \sec^{5/2}(c + dx) dx \right) + \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 3042 \\ & \frac{1}{5} \left(a(3A + 5B) \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx + 5a(A + B) \int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} dx \right) + \\ & \quad \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 4255 \\ & \frac{1}{5} \left(5a(A + B) \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx) \sec^{3/2}(c + dx)}{3d} \right) + a(3A + 5B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right) \\ & \quad \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 3042 \\ & \frac{1}{5} \left(5a(A + B) \left(\frac{1}{3} \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2 \sin(c + dx) \sec^{3/2}(c + dx)}{3d} \right) + a(3A + 5B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right) \\ & \quad \frac{2aA \sin(c + dx) \sec^{5/2}(c + dx)}{5d} \\ & \downarrow 4258 \end{aligned}$$

3.459. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

$$\frac{1}{5} \left(5a(A+B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + a(3A+5B) \left(\frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d} \right) \right) \downarrow 3042$$

$$\frac{1}{5} \left(5a(A+B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + a(3A+5B) \left(\frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d} \right) \right) \downarrow 3119$$

$$\frac{1}{5} \left(5a(A+B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + a(3A+5B) \left(\frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d} \right) \right) \downarrow 3120$$

$$\frac{1}{5} \left(5a(A+B) \left(\frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} \right) + a(3A+5B) \left(\frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*a*A*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(5*d) + (a*(3*A + 5*B)*((-2*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d + (2*sqrt[Sec[c + d*x]]*Sin[c + d*x])/d) + 5*a*(A + B)*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (2*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d))/5`

3.459.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_))^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4255 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

3.459.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 633 vs. 2(200) = 400.

Time = 37.11 (sec) , antiderivative size = 634, normalized size of antiderivative = 3.69

method	result
default	$4\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a \left(\frac{B\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}{2\sin\left(\frac{dx}{2} + \frac{c}{2}\right)^2\left(2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)} \left(2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right) \right)$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNVER
BOSE)
```

3.459. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

output

```

-4*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(1/2*B/sin(
1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1
/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*
x+1/2*c),2^(1/2)))+1/10*A/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+
6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(
1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(
1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d
*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2
^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+(1/2*A+1/2*B
)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*co
s(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1
/2*d*x+1/2*c)^2-1)^(1/2)/d

```

3.459.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.27

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{-5i\sqrt{2}(A+B)a\cos(dx+c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + 5i\sqrt{2}(A+B)a}{\dots}$$

input

```

integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm=
"fracas")

```

3.459. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

output `1/15*(-5*I*sqrt(2)*(A + B)*a*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*(A + B)*a*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(3*A + 5*B)*a*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(3*A + 5*B)*a*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*(3*A + 5*B)*a*cos(d*x + c)^2 + 5*(A + B)*a*cos(d*x + c) + 3*A*a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2)`

3.459.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.459.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.459.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.459.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + a \cos(c + dx)) dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x)), x)`

3.460 $\int (a+a \cos(c+dx))(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$

3.460.1 Optimal result	4208
3.460.2 Mathematica [C] (verified)	4209
3.460.3 Rubi [A] (verified)	4209
3.460.4 Maple [B] (verified)	4213
3.460.5 Fricas [C] (verification not implemented)	4214
3.460.6 Sympy [F(-1)]	4214
3.460.7 Maxima [F]	4215
3.460.8 Giac [F]	4215
3.460.9 Mupad [F(-1)]	4215

3.460.1 Optimal result

Integrand size = 31, antiderivative size = 135

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= -\frac{2a(A + B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2a(A + 3B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a(A + B)\sqrt{\sec(c + dx)}\sin(c + dx)}{d} + \frac{2aA \sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{3d}$$

```
output 2/3*a*A*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2*a*(A+B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*a*(A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.460.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.10 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.67

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{a(1 + \cos(c + dx)) \left((A + 3B) (1 + e^{2i(c+dx)}) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + i(A - 3Ae^{i(c+dx)} - \dots \right)}{\dots}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(a*(1 + Cos[c + d*x])*((A + 3*B)*(1 + E^((2*I)*(c + d*x))))*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + I*(A - 3*A*E^(I*(c + d*x)) - 3*B*E^(I*(c + d*x)) - A*E^((2*I)*(c + d*x)) - 3*A*E^((3*I)*(c + d*x)) - 3*B*E^((3*I)*(c + d*x)) + (A + B)*E^(I*(c + d*x))*(1 + E^((2*I)*(c + d*x)))^(3/2)*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]))*Sec[(c + d*x)/2]^2*Sqrt[Sec[c + d*x]]/(3*d*(1 + E^((2*I)*(c + d*x))))`

3.460.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4485, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \sqrt{\sec(c + dx)}(a \sec(c + dx) + a)(A \sec(c + dx) + B) dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+a\right)\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)dx \\
& \quad \downarrow 4485 \\
& \frac{2}{3}\int\frac{1}{2}\sqrt{\sec(c+dx)}(a(A+3B)+3a(A+B)\sec(c+dx))dx+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 27 \\
& \frac{1}{3}\int\sqrt{\sec(c+dx)}(a(A+3B)+3a(A+B)\sec(c+dx))dx+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{3}\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a(A+3B)+3a(A+B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)dx+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 4274 \\
& \frac{1}{3}\left(3a(A+B)\int\sec^{\frac{3}{2}}(c+dx)dx+a(A+3B)\int\sqrt{\sec(c+dx)}dx\right)+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{3}\left(a(A+3B)\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}dx+3a(A+B)\int\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}dx\right)+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 4255 \\
& \frac{1}{3}\left(a(A+3B)\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}dx+3a(A+B)\left(\frac{2\sin(c+dx)\sqrt{\sec(c+dx)}}{d}-\int\frac{1}{\sqrt{\sec(c+dx)}}dx\right)\right)+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{3}\left(a(A+3B)\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}dx+3a(A+B)\left(\frac{2\sin(c+dx)\sqrt{\sec(c+dx)}}{d}-\int\frac{1}{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}}dx\right)\right)+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 4258
\end{aligned}$$

$$\frac{1}{3} \left(a(A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3a(A + B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(a(A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a(A + B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \downarrow \text{3119}$$

$$\frac{1}{3} \left(a(A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a(A + B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \downarrow \text{3120}$$

$$\frac{1}{3} \left(\frac{2a(A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + 3a(A + B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right)$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*a*A*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + ((2*a*(A + 3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 3*a*(A + B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3`

3.460.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_))^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4255 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

3.460.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 398 vs. 2(171) = 342.

Time = 34.90 (sec) , antiderivative size = 399, normalized size of antiderivative = 2.96

method	result
default	$4\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)+1}\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a \left(\frac{B\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right),\sqrt{2}\right)}{2\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}} + A\left(\frac{-\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)+1}}{6}\right) \right)$
parts	$\frac{2aA\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RETURNVER
BOSE)
```

```
output -4*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(1/2*B*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x
+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)
)+1/2*A*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*
c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+(1/2*A+1/2*B)/sin(1/2*
d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2))))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.460.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.39

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{-i \sqrt{2}(A + 3B)a \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2}(A + 3B)a \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - 3i \sqrt{2}(A + B)a \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 3i \sqrt{2}(A + B)a \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2(3(A + B)a \cos(dx + c) + Aa) \sin(dx + c) / \sqrt{\cos(dx + c)}}{d \cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="fricas")`

output `1/3*(-I*sqrt(2)*(A + 3*B)*a*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*(A + 3*B)*a*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(A + B)*a*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(A + B)*a*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*(A + B)*a*cos(d*x + c) + A*a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))`

3.460.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.460.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.460.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.460.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + a \cos(c + dx)) dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x)), x)`

3.460. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.461 $\int (a+a \cos(c+dx))(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$

3.461.1 Optimal result	4216
3.461.2 Mathematica [C] (verified)	4217
3.461.3 Rubi [A] (verified)	4217
3.461.4 Maple [A] (verified)	4220
3.461.5 Fricas [C] (verification not implemented)	4221
3.461.6 Sympy [F(-1)]	4221
3.461.7 Maxima [F]	4222
3.461.8 Giac [F]	4222
3.461.9 Mupad [F(-1)]	4222

3.461.1 Optimal result

Integrand size = 31, antiderivative size = 106

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= -\frac{2a(A - B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2a(A + B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2aA\sqrt{\sec(c + dx)}\sin(c + dx)}{d}$$

```
output 2*a*A*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2*a*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)
*sec(d*x+c)^(1/2)/d+2*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2
*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2
)/d
```

3.461.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.39 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.48

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{2ae^{-idx} \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(-3iA \cos(c + dx) + 3iB \cos(c + dx) + 3(A + B) \sqrt{\cos(c + dx)} \right)}{\dots}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(2*a*Sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*((-3*I)*A*Cos[c + d*x] + (3*I)*B*Cos[c + d*x] + 3*(A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + I*(A - B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + 3*A*Sin[c + d*x]))/(3*d*E^(I*d*x))`

3.461.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3439, 3042, 4485, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)(A \sec(c + dx) + B)}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)(A \csc(c + dx + \frac{\pi}{2}) + B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4485} \\
& 2 \int -\frac{a(A - B) - a(A + B) \sec(c + dx)}{2\sqrt{\sec(c + dx)}} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow \text{27} \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{a(A - B) - a(A + B) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{a(A - B) - a(A + B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4274} \\
& -a(A - B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + a(A + B) \int \sqrt{\sec(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& -a(A - B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + a(A + B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow \text{4258} \\
& a(A + B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx - a(A - \\
& B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& a(A + B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - a(A - \\
& B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow \text{3119}
\end{aligned}$$

3.461. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\begin{aligned}
& a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \\
& \frac{2a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2aA\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \\
& \quad \downarrow \text{3120} \\
& \frac{2a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d} - \\
& \frac{2a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E(\frac{1}{2}(c+dx)|2)}{d} + \frac{2aA\sin(c+dx)\sqrt{\sec(c+dx)}}{d}
\end{aligned}$$

input `Int[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(-2*a*(A - B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*(A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*A*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d`

3.461.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m+n) Int[(g*Csc[e + f*x])^(p-m-n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4485 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

3.461.4 Maple [A] (verified)

Time = 7.66 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.28

method	result
default	$2a \left(2A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1 F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right)$
parts	$- \frac{2aA \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1 \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1 \right) d}{\sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1 d}$

input `int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `2*a*(2*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2-A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.461. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.461.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.33

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{-i \sqrt{2}(A + B) \operatorname{awierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2}(A + B) \operatorname{awierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2A \operatorname{awierstrassZeta}(-4, 0, \operatorname{awierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 2B \operatorname{awierstrassZeta}(-4, 0, \operatorname{awierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `(-I*sqrt(2)*(A + B)*awierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*(A + B)*awierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - I*sqrt(2)*(A - B)*awierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*(A - B)*awierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*A*a*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.461.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.461.7 Maxima [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.461.8 Giac [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.461.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + a \cos(c + dx)) dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x)), x)`

3.461. $\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.462 $\int (a+a \cos(c+dx))(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$

3.462.1 Optimal result	4223
3.462.2 Mathematica [C] (verified)	4223
3.462.3 Rubi [A] (verified)	4224
3.462.4 Maple [B] (verified)	4227
3.462.5 Fricas [C] (verification not implemented)	4228
3.462.6 Sympy [F]	4228
3.462.7 Maxima [F]	4229
3.462.8 Giac [F]	4229
3.462.9 Mupad [F(-1)]	4229

3.462.1 Optimal result

Integrand size = 31, antiderivative size = 110

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{2a(A + B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d} + \frac{2a(3A + B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d} + \frac{2aB \sin(c + dx)}{3d \sqrt{\sec(c + dx)}}$$

output `2/3*a*B*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*a*(3*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d`

3.462.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.50 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.35

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{2ae^{-idx} \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left((3A + B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - i(A + B)e^i \right)}{3d}$$

input `Integrate[(a + a*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(2*a*Sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*((3*A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - I*(A + B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((3*I)*(A + B) + B*Sin[c + d*x]))/(3*d*E^(I*d*x))`

3.462.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3439, 3042, 4484, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c + dx)}(a \cos(c + dx) + a)(A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{(a \sec(c + dx) + a)(A \sec(c + dx) + B)}{\sec^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a \csc\left(c + dx + \frac{\pi}{2}\right) + a)\left(A \csc\left(c + dx + \frac{\pi}{2}\right) + B\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2}} dx \\
 & \quad \downarrow \text{4484} \\
 & \frac{2aB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} - \frac{2}{3} \int -\frac{3a(A + B) + a(3A + B)\sec(c + dx)}{2\sqrt{\sec(c + dx)}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \int \frac{3a(A + B) + a(3A + B)\sec(c + dx)}{\sqrt{\sec(c + dx)}} dx + \frac{2aB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{1}{3} \int \frac{3a(A+B) + a(3A+B) \csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 4274

$$\frac{1}{3} \left(3a(A+B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + a(3A+B) \int \sqrt{\sec(c+dx)} dx \right) + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 3042

$$\frac{1}{3} \left(3a(A+B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + a(3A+B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx \right) + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 4258

$$\frac{1}{3} \left(a(3A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a(A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx \right) + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 3042

$$\frac{1}{3} \left(a(3A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a(A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx \right) + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 3119

$$\frac{1}{3} \left(a(3A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a(A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx), 2)}{d} \right) + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2a(3A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6a(A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx), 2)}{d} \right) + \frac{2aB \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

input `Int[(a + a*cos[c + d*x])*(A + B*cos[c + d*x])*sqrt[sec[c + d*x]],x]`

output `((6*a*(A + B)*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d + (2*a*(3*A + B)*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d)/3 + (2*a*B*sin[c + d*x])/(3*d*sqrt[Sec[c + d*x]])`

3.462.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4484 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e +
f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(
n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]
```

3.462.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. 2(148) = 296.

Time = 8.61 (sec) , antiderivative size = 321, normalized size of antiderivative = 2.92

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} \left(4B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 3A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1\right) F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)$
parts	$-\frac{2aA\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}} \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 1}{\sin\left(\frac{dx}{2} + \frac{c}{2}\right)} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right), \sqrt{2}} + \frac{2(aA + Ba)\sqrt{2}}{\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1} d$

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RETURNVER
BOSE)
```

```
output -2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(4*B*cos(1/
2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(
1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*A*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*
d*x+1/2*c),2^(1/2))-2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2+B*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*
x+1/2*c),2^(1/2))-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2
-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+
sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(
1/2)/d
```

3.462.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.29

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{2Ba\sqrt{\cos(dx + c)} \sin(dx + c) - i\sqrt{2}(3A + B)a\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{d}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `1/3*(2*B*a*sqrt(cos(d*x + c))*sin(d*x + c) - I*sqrt(2)*(3*A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*(3*A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.462.6 Sympy [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= a \left(\int A\sqrt{\sec(c + dx)} dx + \int A \cos(c + dx)\sqrt{\sec(c + dx)} dx \right. \\ \left. + \int B \cos(c + dx)\sqrt{\sec(c + dx)} dx + \int B \cos^2(c + dx)\sqrt{\sec(c + dx)} dx \right)$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `a*(Integral(A*sqrt(sec(c + d*x)), x) + Integral(A*cos(c + d*x)*sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)*sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)**2*sqrt(sec(c + d*x)), x))`

3.462.7 Maxima [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)\sqrt{\sec(dx + c)} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.462.8 Giac [F]

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)\sqrt{\sec(dx + c)} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.462.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + a \cos(c + dx)) dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x)), x)`

3.462. $\int (a + a \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$

3.463 $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

3.463.1 Optimal result 4230
 3.463.2 Mathematica [C] (verified) 4231
 3.463.3 Rubi [A] (verified) 4231
 3.463.4 Maple [B] (verified) 4235
 3.463.5 Fracas [C] (verification not implemented) 4236
 3.463.6 Sympy [F] 4236
 3.463.7 Maxima [F] 4237
 3.463.8 Giac [F] 4237
 3.463.9 Mupad [F(-1)] 4237

3.463.1 Optimal result

Integrand size = 31, antiderivative size = 141

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{2a(5A + 3B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2a(A + B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2a(A + B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}}$$

```
output 2/5*a*B*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/3*a*(A+B)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2/5*a*(5*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.463.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 1.60 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.05

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{a\sqrt{\sec(c + dx)} \left(10(A + B)\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - 2i(5A + 3B)e^{i(c+dx)}\sqrt{1 + e^{2i(c+dx)}} \operatorname{Hy} \right)}{15d}$$

input `Integrate[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(a*Sqrt[Sec[c + d*x]]*(10*(A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (2*I)*(5*A + 3*B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((6*I)*(5*A + 3*B) + 10*(A + B)*Sin[c + d*x] + 3*B*Sin[2*(c + d*x)])))/(15*d)`

3.463.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4484, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)(A \sec(c + dx) + B)}{\sec^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.463. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)(A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx \\
& \quad \downarrow 4484 \\
& \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} - \frac{2}{5} \int -\frac{5a(A + B) + a(5A + 3B) \sec(c + dx)}{2 \sec^{\frac{3}{2}}(c + dx)} dx \\
& \quad \downarrow 27 \\
& \frac{1}{5} \int \frac{5a(A + B) + a(5A + 3B) \sec(c + dx)}{\sec^{\frac{3}{2}}(c + dx)} dx + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{5} \int \frac{5a(A + B) + a(5A + 3B) \csc(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 4274 \\
& \frac{1}{5} \left(5a(A + B) \int \frac{1}{\sec^{\frac{3}{2}}(c + dx)} dx + a(5A + 3B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{5} \left(5a(A + B) \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx + a(5A + 3B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 4256 \\
& \frac{1}{5} \left(a(5A + 3B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 5a(A + B) \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) \right) + \\
& \quad \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{5} \left(a(5A + 3B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 5a(A + B) \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) \right) + \\
& \quad \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 4258
\end{aligned}$$

3.463. $\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$

$$\frac{1}{5} \left(a(5A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + 5a(A + B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(5a(A + B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) + a(5A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3119

$$\frac{1}{5} \left(5a(A + B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) + \frac{2a(5A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \int \sqrt{\cos(c + dx)} dx + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{2a(5A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + 5a(A + B) \left(\frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} + \frac{2 \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \int \sqrt{\cos(c + dx)} dx \right) + \frac{2aB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(2*a*B*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((2*a*(5*A + 3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 5*a*(A + B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]))/5`

3.463.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4256 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)])*(d_.)^(n_)*((csc[(e_.) + (f_)*(x_)])*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4484 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e +
f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(
n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]
```

3.463.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. 2(173) = 346.

Time = 10.19 (sec) , antiderivative size = 355, normalized size of antiderivative = 2.52

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a} \left(-24B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (20A + 44B)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + (-10A - 16B)\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 5A \operatorname{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2^{1/2}\right) - 15A \operatorname{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2^{1/2}\right) + 5B \operatorname{EllipticF}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2^{1/2}\right) - 9B \operatorname{EllipticE}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), 2^{1/2}\right)\right) / (-2\sin\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + \sin\left(\frac{dx}{2} + \frac{c}{2}\right)^2) / \sin\left(\frac{dx}{2} + \frac{c}{2}\right) / (2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 1) / d$
parts	$\frac{2aA\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - 2(aA + Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}d}$

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNVER
BOSE)
```

```
output -2/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(-24*B*cos
(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+(20*A+44*B)*sin(1/2*d*x+1/2*c)^4*cos(
1/2*d*x+1/2*c)+(-10*A-16*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+5*A*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(
1/2*d*x+1/2*c),2^(1/2))-15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*B*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c
),2^(1/2))-9*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/
2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.463.
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.463.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.20

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{-5i \sqrt{2}(A + B)a \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2}(A + B)a \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{\sqrt{\sec(c + dx)}}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `1/15*(-5*I*sqrt(2)*(A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*(A + B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(5*A + 3*B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(5*A + 3*B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*B*a*cos(d*x + c)^2 + 5*(A + B)*a*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.463.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = a \left(\int \frac{A}{\sqrt{\sec(c + dx)}} dx + \int \frac{A \cos(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{B \cos(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{B \cos^2(c + dx)}{\sqrt{\sec(c + dx)}} dx \right)$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `a*(Integral(A/sqrt(sec(c + d*x)), x) + Integral(A*cos(c + d*x)/sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)/sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)**2/sqrt(sec(c + d*x)), x))`

3.463.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.463.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.463.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + a \cos(c + dx))}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/(1/cos(c + d*x))^(1/2), x)`

3.464
$$\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.464.1 Optimal result 4238
 3.464.2 Mathematica [C] (verified) 4239
 3.464.3 Rubi [A] (verified) 4239
 3.464.4 Maple [A] (verified) 4243
 3.464.5 Fricas [C] (verification not implemented) 4244
 3.464.6 Sympy [F] 4244
 3.464.7 Maxima [F] 4245
 3.464.8 Giac [F] 4245
 3.464.9 Mupad [F(-1)] 4245

3.464.1 Optimal result

Integrand size = 31, antiderivative size = 172

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{6a(A + B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2a(7A + 5B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} + \frac{2a(A + B) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2a(7A + 5B) \sin(c + dx)}{21d \sqrt{\sec(c + dx)}}$$

output

```
2/7*a*B*sin(d*x+c)/d/sec(d*x+c)^(5/2)+2/5*a*(A+B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/21*a*(7*A+5*B)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+6/5*a*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/21*a*(7*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.464.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.16 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.06

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{ae^{-idx} \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(20(7A + 5B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - 84i(A + B) \right)}{210dE^{(I*d*x)}}$$

input `Integrate[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2), x]`

output `(a*sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*(20*(7*A + 5*B)*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (84*I)*(A + B)*E^(I*(c + d*x))*sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((252*I)*(A + B) + 5*(28*A + 23*B)*Sin[c + d*x] + 42*(A + B)*Sin[2*(c + d*x)] + 15*B*Sin[3*(c + d*x)])))/(210*d*E^(I*d*x))`

3.464.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.99, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4484, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)(A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)(A \sec(c + dx) + B)}{\sec^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.464. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)(A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow 4484 \\
& \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} - \frac{2}{7} \int -\frac{7a(A + B) + a(7A + 5B) \sec(c + dx)}{2 \sec^{\frac{5}{2}}(c + dx)} dx \\
& \quad \downarrow 27 \\
& \frac{1}{7} \int \frac{7a(A + B) + a(7A + 5B) \sec(c + dx)}{\sec^{\frac{5}{2}}(c + dx)} dx + \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \int \frac{7a(A + B) + a(7A + 5B) \csc(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx + \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 4274 \\
& \frac{1}{7} \left(7a(A + B) \int \frac{1}{\sec^{\frac{5}{2}}(c + dx)} dx + a(7A + 5B) \int \frac{1}{\sec^{\frac{3}{2}}(c + dx)} dx \right) + \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left(7a(A + B) \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx + a(7A + 5B) \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \\
& \quad \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 4256 \\
& \frac{1}{7} \left(7a(A + B) \left(\frac{3}{5} \int \frac{1}{\sqrt{\sec(c + dx)}} dx + \frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) + a(7A + 5B) \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) \right) + \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left(7a(A + B) \left(\frac{3}{5} \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) + a(7A + 5B) \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\csc(c + dx + \frac{\pi}{2})}} \right) \right) + \frac{2aB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 4258
\end{aligned}$$

3.464. $\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$

$$\frac{1}{7} \left(7a(A+B) \left(\frac{3}{5} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + a(7A+5B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right) + \frac{2aB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(7a(A+B) \left(\frac{3}{5} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + a(7A+5B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) + 7a(A+B) \left(\frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + \frac{2aB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3119

$$\frac{1}{7} \left(a(7A+5B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) + 7a(A+B) \left(\frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right) + \frac{2aB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3120

$$\frac{1}{7} \left(a(7A+5B) \left(\frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} + \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} \right) + 7a(A+B) \left(\frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right) + \frac{2aB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

input `Int[((a + a*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `(2*a*B*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + (7*a*(A + B)*((6*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(5*d) + (2*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)))) + a*(7*A + 5*B)*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*sqrt[Sec[c + d*x]])))/7`

3.464.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4484 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e +
f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(
n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]
```

3.464.4 Maple [A] (verified)

Time = 11.72 (sec) , antiderivative size = 383, normalized size of antiderivative = 2.23

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{a}\left(240B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+(-168A-528B)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+$
parts	$-\frac{2aA\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d$

```
input int((a+cos(d*x+c)*a)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x,method=_RETURNVER
BOSE)
```

```
output -2/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a*(240*B*co
s(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8+(-168*A-528*B)*sin(1/2*d*x+1/2*c)^6*
cos(1/2*d*x+1/2*c)+(308*A+448*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(
-112*A-122*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+35*A*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c
),2^(1/2))-63*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+25*B*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-6
3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elliptic
E(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.464.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.09

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{-5i \sqrt{2}(7A + 5B) \operatorname{aweierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2}(7A + 5B) \operatorname{aweierst}}$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `1/105*(-5*I*sqrt(2)*(7*A + 5*B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*(7*A + 5*B)*a*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 63*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 63*I*sqrt(2)*(A + B)*a*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(15*B*a*cos(d*x + c)^3 + 21*(A + B)*a*cos(d*x + c)^2 + 5*(7*A + 5*B)*a*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.464.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx = a \left(\int \frac{A}{\sec^{\frac{3}{2}}(c + dx)} dx + \int \frac{A \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)} dx + \int \frac{B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)} dx + \int \frac{B \cos^2(c + dx)}{\sec^{\frac{3}{2}}(c + dx)} dx \right)$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `a*(Integral(A/sec(c + d*x)**(3/2), x) + Integral(A*cos(c + d*x)/sec(c + d*x)**(3/2), x) + Integral(B*cos(c + d*x)/sec(c + d*x)**(3/2), x) + Integral(B*cos(c + d*x)**2/sec(c + d*x)**(3/2), x))`

3.464. $\int \frac{(a+a \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

3.464.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.464.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)}{\sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.464.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + a \cos(c + dx))}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x)))/(1/cos(c + d*x))^(3/2), x)`

3.465 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

3.465.1 Optimal result	4246
3.465.2 Mathematica [C] (verified)	4247
3.465.3 Rubi [A] (verified)	4247
3.465.4 Maple [B] (verified)	4252
3.465.5 Fricas [C] (verification not implemented)	4253
3.465.6 Sympy [F(-1)]	4253
3.465.7 Maxima [F]	4254
3.465.8 Giac [F]	4254
3.465.9 Mupad [F(-1)]	4254

3.465.1 Optimal result

Integrand size = 33, antiderivative size = 199

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= -\frac{4a^2(4A + 5B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{4a^2(A + 2B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{4a^2(4A + 5B)\sqrt{\sec(c + dx)}\sin(c + dx)}{5d} + \frac{2a^2(7A + 5B)\sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{15d}$$

$$+ \frac{2A\sec^{\frac{3}{2}}(c + dx)(a^2 + a^2\sec(c + dx))\sin(c + dx)}{5d}$$

output $2/15*a^2*(7*A+5*B)*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)/d+2/5*A*\sec(d*x+c)^{(3/2)}*(a^2+a^2*\sec(d*x+c))*\sin(d*x+c)/d+4/5*a^2*(4*A+5*B)*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/d-4/5*a^2*(4*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*\sec(d*x+c)^{(1/2)}/d+4/3*a^2*(A+2*B)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*\sec(d*x+c)^{(1/2)}/d$

3.465.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.55 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.50

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{a^2 e^{-ic} (-1 + e^{2ic}) (1 + \cos(c + dx))^2 \csc(c) \left(10A + 5B - 18Ae^{i(c+dx)} - 30Be^{i(c+dx)} - 54Ae^{3i(c+dx)} - 60B \right)}{}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(a^2*(-1 + E^((2*I)*c))*(1 + Cos[c + d*x])^2*Csc[c]*(10*A + 5*B - 18*A*E^(I*(c + d*x)) - 30*B*E^(I*(c + d*x)) - 54*A*E^((3*I)*(c + d*x)) - 60*B*E^((3*I)*(c + d*x)) - 10*A*E^((4*I)*(c + d*x)) - 5*B*E^((4*I)*(c + d*x)) - 24*A*E^((5*I)*(c + d*x)) - 30*B*E^((5*I)*(c + d*x)) - (10*I)*(A + 2*B)*(1 + E^((2*I)*(c + d*x)))^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(4*A + 5*B)*E^(I*(c + d*x))*(1 + E^((2*I)*(c + d*x)))^(5/2)*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*Sec[(c + d*x)/2]^4*Sqrt[Sec[c + d*x]])/(60*d*E^(I*c)*(1 + E^((2*I)*(c + d*x)))^2)`

3.465.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.98, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3439, 3042, 4506, 27, 3042, 4485, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx) (a \cos(c + dx) + a)^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

3.465. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \sqrt{\sec(c+dx)}(a \sec(c+dx) + a)^2(A \sec(c+dx) + B)dx \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a \csc\left(c+dx+\frac{\pi}{2}\right) + a\right)^2\left(A \csc\left(c+dx+\frac{\pi}{2}\right) + B\right) dx \\
& \quad \downarrow \text{4506} \\
& \frac{2}{5} \int \frac{1}{2} \sqrt{\sec(c+dx)}(\sec(c+dx)a + a)(a(A+5B) + a(7A+5B) \sec(c+dx))dx + \\
& \quad \frac{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a^2 \sec(c+dx) + a^2)}{5d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \sqrt{\sec(c+dx)}(\sec(c+dx)a + a)(a(A+5B) + a(7A+5B) \sec(c+dx))dx + \\
& \quad \frac{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a^2 \sec(c+dx) + a^2)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(\csc\left(c+dx+\frac{\pi}{2}\right) a + a\right)\left(a(A+5B) + a(7A+5B) \csc\left(c+dx+\frac{\pi}{2}\right)\right) dx + \\
& \quad \frac{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a^2 \sec(c+dx) + a^2)}{5d} \\
& \quad \downarrow \text{4485} \\
& \frac{1}{5} \left(\frac{2}{3} \int \sqrt{\sec(c+dx)}(5(A+2B)a^2 + 3(4A+5B) \sec(c+dx)a^2) dx + \frac{2a^2(7A+5B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + \\
& \quad \frac{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a^2 \sec(c+dx) + a^2)}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{2}{3} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(5(A+2B)a^2 + 3(4A+5B) \csc\left(c+dx+\frac{\pi}{2}\right) a^2\right) dx + \frac{2a^2(7A+5B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + \\
& \quad \frac{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a^2 \sec(c+dx) + a^2)}{5d} \\
& \quad \downarrow \text{4274}
\end{aligned}$$

3.465. $\int (a + a \cos(c+dx))^2(A + B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^2(4A + 5B) \int \sec^{\frac{3}{2}}(c + dx) dx + 5a^2(A + 2B) \int \sqrt{\sec(c + dx)} dx \right) + \frac{2a^2(7A + 5B) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right)$$

$$\frac{2A \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^2 \sec(c + dx) + a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx + 3a^2(4A + 5B) \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx \right) + \frac{2a^2(7A + 5B) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right)$$

$$\frac{2A \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^2 \sec(c + dx) + a^2)}{5d}$$

↓ 4255

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) \right) \right)$$

$$\frac{2A \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^2 \sec(c + dx) + a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx \right) \right) \right)$$

$$\frac{2A \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^2 \sec(c + dx) + a^2)}{5d}$$

↓ 4258

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) \right) \right)$$

$$\frac{2A \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^2 \sec(c + dx) + a^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A + 2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + 3a^2(4A + 5B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) \right) \right)$$

$$\frac{2A \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^2 \sec(c + dx) + a^2)}{5d}$$

↓ 3119

3.465. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(A+2B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a^2(4A+5B) \left(\frac{2\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right. \right. \right. \\ \left. \left. \left. \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a^2\sec(c+dx)+a^2)}{5d} \right) \right) \right. \\ \left. \downarrow \text{3120} \right. \\ \frac{1}{5} \left(\frac{2a^2(7A+5B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{2}{3} \left(\frac{10a^2(A+2B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} \right. \right. \\ \left. \left. \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a^2\sec(c+dx)+a^2)}{5d} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2), x]`

output `(2*A*Sec[c + d*x]^(3/2)*(a^2 + a^2*Sec[c + d*x])*Sin[c + d*x])/(5*d) + ((2*a^2*(7*A + 5*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + (2*((10*a^2*(A + 2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 3*a^2*(4*A + 5*B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d))/3)/5`

3.465.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(g_.)^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4255 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Simp}[b^2*((n-2)/(n-1)) \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4485 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))*(\text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*(d*\text{Csc}[e + f*x])^n/(f*(n+1)), x] + \text{Simp}[1/(n+1) \text{Int}[(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*a*(n+1) + B*b*n + (A*b + B*a)*(n+1)*\text{Csc}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& !\text{LeQ}[n, -1]$

rule 4506 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m-1)}*((d*\text{Csc}[e + f*x])^n/(f*(m+n))), x] + \text{Simp}[1/(d*(m+n)) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m-1)}*(d*\text{Csc}[e + f*x])^n*\text{Simp}[a*A*d*(m+n) + B*(b*d*n) + (A*b*d*(m+n) + a*B*d*(2*m+n-1))*\text{Csc}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{GtQ}[m, 1/2] \&\& !\text{LtQ}[n, -1]$

3.465.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 713 vs. $2(227) = 454$.

Time = 63.49 (sec) , antiderivative size = 714, normalized size of antiderivative = 3.59

method	result	size
default	Expression too large to display	714
parts	Expression too large to display	916

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

```
output -8*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(1/4*B*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d
*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/
2))+1/20*A/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1
/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^
6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^
4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*s
in(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*s
in(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*si
n(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+(1/2*A+1/4*B)*(-1/6*cos(1/2
*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*
d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c
)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Elliptic
F(cos(1/2*d*x+1/2*c),2^(1/2)))+(1/4*A+1/2*B)/sin(1/2*d*x+1/2*c)^2/(2*sin(1
/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2
*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*s
in(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))/sin(1
/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.465.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.20

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx =$$

$$2 \left(5i \sqrt{2} (A + 2B) a^2 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (A$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="fricas")`

output `-2/15*(5*I*sqrt(2)*(A + 2*B)*a^2*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(A + 2*B)*a^2*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(4*A + 5*B)*a^2*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(4*A + 5*B)*a^2*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (6*(4*A + 5*B)*a^2*cos(d*x + c)^2 + 5*(2*A + B)*a^2*cos(d*x + c) + 3*A*a^2)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2)`

3.465.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.465.7 Maxima [F]

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sec(d*x + c)^(7/2), x)`

3.465.8 Giac [F]

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sec(d*x + c)^(7/2), x)`

3.465.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + a \cos(c + dx))^2 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^2,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^2, x)`

3.465. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.466 $\int (a+a \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$

3.466.1 Optimal result	4255
3.466.2 Mathematica [C] (verified)	4256
3.466.3 Rubi [A] (verified)	4256
3.466.4 Maple [B] (verified)	4260
3.466.5 Fricas [C] (verification not implemented)	4261
3.466.6 Sympy [F(-1)]	4262
3.466.7 Maxima [F]	4262
3.466.8 Giac [F]	4262
3.466.9 Mupad [F(-1)]	4263

3.466.1 Optimal result

Integrand size = 33, antiderivative size = 160

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= -\frac{4a^2 A \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{4a^2 (2A + 3B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a^2 (5A + 3B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3d}$$

$$+ \frac{2A \sqrt{\sec(c + dx)} (a^2 + a^2 \sec(c + dx)) \sin(c + dx)}{3d}$$

output

```
2/3*a^2*(5*A+3*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d+2/3*A*(a^2+a^2*sec(d*x+c))
*sin(d*x+c)*sec(d*x+c)^(1/2)/d-4*a^2*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/
2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*
x+c)^(1/2)/d+4/3*a^2*(2*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/
2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/
2)/d
```

3.466.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.30 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.18

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx =$$

$$\frac{ia^2 \sec^{\frac{3}{2}}(c + dx) \left(-6A - 6A \cos(2(c + dx)) \right) + 6Ae^{-2i(c+dx)} (1 + e^{2i(c+dx)})^{3/2} \text{Hypergeometric2F1} \left(-\frac{1}{4}, \frac{1}{2}, \frac{3}{4}, -E^{\left((2i)(c + dx) \right)} \right)}{d}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `((-1/3*I)*a^2*Sec[c + d*x]^(3/2)*(-6*A - 6*A*Cos[2*(c + d*x)] + (6*A*(1 + E^((2*I)*(c + d*x))))^(3/2)*Hypergeometric2F1[-1/4, 1/2, 3/4, -E^((2*I)*(c + d*x))])/E^((2*I)*(c + d*x)) + (2*(2*A + 3*B)*(1 + E^((2*I)*(c + d*x))))^(3/2)*Hypergeometric2F1[1/4, 1/2, 5/4, -E^((2*I)*(c + d*x))])/E^(I*(c + d*x)) + (2*I)*A*Sin[c + d*x] + (6*I)*A*Sin[2*(c + d*x)] + (3*I)*B*Sin[2*(c + d*x)]))/d`

3.466.3 Rubi [A] (verified)

Time = 1.00 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4506, 27, 3042, 4485, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)^2 (A \sec(c + dx) + B)}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

3.466. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4506} \\
& \frac{2}{3} \int - \frac{(\sec(c + dx)a + a)(a(A - 3B) - a(5A + 3B) \sec(c + dx))}{2\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)}{3d} \\
& \quad \downarrow \text{27} \\
& \frac{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)}{3d} - \\
& \frac{1}{3} \int \frac{(\sec(c + dx)a + a)(a(A - 3B) - a(5A + 3B) \sec(c + dx))}{\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)}{3d} - \\
& \frac{1}{3} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)(a(A - 3B) - a(5A + 3B) \csc(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4485} \\
& \frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \int \frac{3a^2A - a^2(2A + 3B) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx \right) + \\
& \quad \frac{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \int \frac{3a^2A - a^2(2A + 3B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)}{3d} \\
& \quad \downarrow \text{4274} \\
& \frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \left(3a^2A \int \frac{1}{\sqrt{\sec(c + dx)}} dx - a^2(2A + 3B) \int \sqrt{\sec(c + dx)} dx \right) \right) + \\
& \quad \frac{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)}{3d} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.466. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \left(\frac{3a^2 A \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx - a^2(2A + 3B) \int \sqrt{\csc(c + dx)}}{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)} \right) \right)$$

↓ 4258

$$\frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \left(\frac{3a^2 A \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx - a^2(2A + 3B) \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx - a^2(2A + 3B) \int \sqrt{\csc(c + dx)}}{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \left(\frac{3a^2 A \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx - a^2(2A + 3B) \int \sqrt{\csc(c + dx)}}{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)} \right) \right)$$

↓ 3119

$$\frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \left(\frac{6a^2 A \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} - a^2(2A + 3B) \int \sqrt{\csc(c + dx)}}{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)} \right) \right)$$

↓ 3120

$$\frac{1}{3} \left(\frac{2a^2(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - 2 \left(\frac{6a^2 A \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} - \frac{2a^2(2A + 3B) \int \sqrt{\csc(c + dx)}}{2A \sin(c + dx) \sqrt{\sec(c + dx)} (a^2 \sec(c + dx) + a^2)} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

```
output (2*A*Sqrt[Sec[c + d*x]]*(a^2 + a^2*Sec[c + d*x])*Sin[c + d*x])/(3*d) + (-2
*((6*a^2*A*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]]
)/d - (2*a^2*(2*A + 3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt
[Sec[c + d*x]])/d) + (2*a^2*(5*A + 3*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d
)/3
```

3.466.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^{(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^{(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^{(n_.)}, x_Symbol] := Sim
p[g^{(m + n) Int[(g*Csc[e + f*x])^{(p - m - n)}*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^{(n_.)}, x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

```
rule 4274 Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^{(n_.)*((c_.) + (f_.)*(x_) +
(a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d In
t[(d*Csc[e + f*x])^{(n + 1)}, x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

```
rule 4506 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] :> Simp[(-b)*B*
Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))),
x] + Simp[1/(d*(m + n)) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x]
)^n*Simp[a*A*d*(m + n) + B*(b*d*n) + (A*b*d*(m + n) + a*B*d*(2*m + n - 1))*
Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b -
a*B, 0] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1]
```

3.466.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 512 vs. $2(194) = 388$.

Time = 60.97 (sec) , antiderivative size = 513, normalized size of antiderivative = 3.21

method	result
default	$-\frac{4 \left(6 \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right)} (2A+B) \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \sqrt{-2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + \sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right)} (7A+3B) \right)}{\dots}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

output

```

-4/3*(6*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A+B)*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4-(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(7*A+3*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-2*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*sin(1/2*d*x+1/2*c)^2+2*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+3*A*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2))*a^2/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3/2)/sin(1/2*d*x+1/2*c)/d

```

3.466.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.26

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx =$$

$$\frac{2 \left(i \sqrt{2} (2A + 3B) a^2 \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - i \sqrt{2} (2A + 3B) a^2 \sin(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) \right)}{\dots}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="fricas")`

output

```

-2/3*(I*sqrt(2)*(2*A + 3*B)*a^2*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - I*sqrt(2)*(2*A + 3*B)*a^2*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*A*a^2*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*A*a^2*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (3*(2*A + B)*a^2*cos(d*x + c) + A*a^2)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))

```

3.466. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.466.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.466.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2), x)`

3.466.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2), x)`

3.466.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + a \cos(c + dx))^2 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^2,x)`output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^2, x)`

3.467 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$

3.467.1 Optimal result	4264
3.467.2 Mathematica [C] (verified)	4265
3.467.3 Rubi [A] (verified)	4265
3.467.4 Maple [A] (verified)	4269
3.467.5 Fricas [C] (verification not implemented)	4270
3.467.6 Sympy [F(-1)]	4270
3.467.7 Maxima [F]	4271
3.467.8 Giac [F]	4271
3.467.9 Mupad [F(-1)]	4271

3.467.1 Optimal result

Integrand size = 33, antiderivative size = 160

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{4a^2B\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{4a^2(3A + 2B)\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a^2(3A - B)\sqrt{\sec(c + dx)} \sin(c + dx)}{3d} + \frac{2B(a^2 + a^2 \sec(c + dx)) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}}$$

```
output 2/3*B*(a^2+a^2*sec(d*x+c))*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2/3*a^2*(3*A-B)*s
in(d*x+c)*sec(d*x+c)^(1/2)/d+4*a^2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*
d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+
c)^(1/2)/d+4/3*a^2*(3*A+2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*
c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)
/d
```

3.467.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.33 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.95

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{2a^2 \left(12iB \operatorname{Hypergeometric2F1} \left(-\frac{1}{4}, \frac{1}{2}, \frac{3}{4}, -e^{2i(c+dx)} \right) - 4i(3A + 2B)e^{i(c+dx)} \operatorname{Hypergeometric2F1} \left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -e^{2i(c+dx)} \right) \right)}{3d\sqrt{1 + e^{2i(c+dx)}}\sqrt{\sec(c + dx)}}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(2*a^2*((12*I)*B*Hypergeometric2F1[-1/4, 1/2, 3/4, -E^((2*I)*(c + d*x))] - (4*I)*(3*A + 2*B)*E^(I*(c + d*x))*Hypergeometric2F1[1/4, 1/2, 5/4, -E^((2*I)*(c + d*x))] + Sqrt[1 + E^((2*I)*(c + d*x))]*((-6*I)*B + B*Sin[c + d*x] + 3*A*Tan[c + d*x]))/(3*d*Sqrt[1 + E^((2*I)*(c + d*x))]*Sqrt[Sec[c + d*x]])`

3.467.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4485, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^2(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)^2(A \sec(c + dx) + B)}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.467. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\
& \quad \downarrow \text{4505} \\
& \frac{2}{3} \int \frac{(\sec(c + dx)a + a)(a(3A + 5B) + a(3A - B) \sec(c + dx))}{2\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{(\sec(c + dx)a + a)(a(3A + 5B) + a(3A - B) \sec(c + dx))}{\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)(a(3A + 5B) + a(3A - B) \csc(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{4485} \\
& \frac{1}{3} \left(2 \int \frac{3Ba^2 + (3A + 2B) \sec(c + dx)a^2}{\sqrt{\sec(c + dx)}} dx + \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(2 \int \frac{3Ba^2 + (3A + 2B) \csc(c + dx + \frac{\pi}{2})a^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{4274} \\
& \frac{1}{3} \left(2 \left(a^2(3A + 2B) \int \sqrt{\sec(c + dx)} dx + 3a^2B \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) + \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d\sqrt{\sec(c + dx)}}
\end{aligned}$$

3.467. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

↓ 3042

$$\frac{1}{3} \left(2 \left(a^2(3A + 2B) \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx + 3a^2B \int \frac{1}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx \right) + \frac{2a^2(3A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d \sqrt{\sec(c + dx)}}$$

↓ 4258

$$\frac{1}{3} \left(2 \left(a^2(3A + 2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3a^2B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d \sqrt{\sec(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{3} \left(2 \left(a^2(3A + 2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + 3a^2B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d \sqrt{\sec(c + dx)}} \right)$$

↓ 3119

$$\frac{1}{3} \left(2 \left(a^2(3A + 2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{6a^2B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx)\right)}{d} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d \sqrt{\sec(c + dx)}} \right)$$

↓ 3120

$$\frac{1}{3} \left(\frac{2a^2(3A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} + 2 \left(\frac{2a^2(3A + 2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{3d \sqrt{\sec(c + dx)}} \right) \right)$$

input `Int[(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

```
output (2*B*(a^2 + a^2*Sec[c + d*x])*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]) + (2*
((6*a^2*B*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])
/d + (2*a^2*(3*A + 2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[
Sec[c + d*x]])/d + (2*a^2*(3*A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3
```

3.467.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

```
rule 4274 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d In
t[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

```
rule 4505 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] :> Simp[a*A*Cot
[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Sim
p[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Sim
p[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0
] && GtQ[m, 1/2] && LtQ[n, -1]
```

3.467.4 Maple [A] (verified)

Time = 9.71 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.52

method	result
default	$4a^2 \left(-2B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 3A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 3A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \right)$
parts	$-\frac{2A a^2 \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} d \right)}{\sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} d}$

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 4/3*a^2*(-2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+3*A*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^2-3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/
2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*cos(1/2*d*x+1/2*c)
*sin(1/2*d*x+1/2*c)^2-2*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*
c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2
^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.467. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.467.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.04

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx =$$

$$2 \left(i \sqrt{2} (3A + 2B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - i \sqrt{2} (3A + 2B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) \right) / d$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `-2/3*(I*sqrt(2)*(3*A + 2*B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - I*sqrt(2)*(3*A + 2*B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*B*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*B*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (B*a^2*cos(d*x + c) + 3*A*a^2)*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.467.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.467.7 Maxima [F]

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2), x)`

3.467.8 Giac [F]

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2), x)`

3.467.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + a \cos(c + dx))^2 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^2,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^2, x)`

3.467. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.468 $\int (a+a \cos(c+dx))^2(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$

3.468.1 Optimal result	4272
3.468.2 Mathematica [C] (verified)	4273
3.468.3 Rubi [A] (verified)	4273
3.468.4 Maple [A] (verified)	4277
3.468.5 Fricas [C] (verification not implemented)	4278
3.468.6 Sympy [F]	4279
3.468.7 Maxima [F]	4279
3.468.8 Giac [F]	4280
3.468.9 Mupad [F(-1)]	4280

3.468.1 Optimal result

Integrand size = 33, antiderivative size = 166

$$\int (a + a \cos(c + dx))^2(A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{4a^2(5A + 4B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{4a^2(2A + B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a^2(5A + 7B) \sin(c + dx)}{15d \sqrt{\sec(c + dx)}} + \frac{2B(a^2 + a^2 \sec(c + dx)) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)}$$

```
output 2/5*B*(a^2+a^2*sec(d*x+c))*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/15*a^2*(5*A+7*B)
)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+4/5*a^2*(5*A+4*B)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(
1/2)*sec(d*x+c)^(1/2)/d+4/3*a^2*(2*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1
/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d
*x+c)^(1/2)/d
```

3.468.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.57 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.92

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{a^2 \sqrt{\sec(c + dx)} \left(20(2A + B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - 4i(5A + 4B) e^{i(c+dx)} \sqrt{1 + e^{2i(c+dx)}} \right)}{5d}$$

input `Integrate[(a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(a^2*Sqrt[Sec[c + d*x]]*(20*(2*A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (4*I)*(5*A + 4*B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((60*I)*A + (48*I)*B + 10*(A + 2*B)*Sin[c + d*x] + 3*B*Sin[2*(c + d*x)])))/(15*d)`

3.468.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4484, 25, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c + dx)} (a \cos(c + dx) + a)^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)^2 (A \sec(c + dx) + B)}{\sec^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx \\
& \quad \downarrow \text{4505} \\
& \frac{2}{5} \int \frac{(\sec(c + dx)a + a)(a(5A + 7B) + a(5A + B) \sec(c + dx))}{\frac{2 \sec^{\frac{3}{2}}(c + dx)}{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}} dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{(\sec(c + dx)a + a)(a(5A + 7B) + a(5A + B) \sec(c + dx))}{\frac{\sec^{\frac{3}{2}}(c + dx)}{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)(a(5A + 7B) + a(5A + B) \csc(c + dx + \frac{\pi}{2}))}{\frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}} dx + \\
& \quad \downarrow \text{4484} \\
& \frac{1}{5} \left(\frac{2a^2(5A + 7B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} - \frac{2}{3} \int \frac{3(5A + 4B)a^2 + 5(2A + B) \sec(c + dx)a^2}{\sqrt{\sec(c + dx)}} dx \right) + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{25} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{3(5A + 4B)a^2 + 5(2A + B) \sec(c + dx)a^2}{\sqrt{\sec(c + dx)}} dx + \frac{2a^2(5A + 7B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{3(5A + 4B)a^2 + 5(2A + B) \csc(c + dx + \frac{\pi}{2}) a^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2a^2(5A + 7B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \\
& \quad \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4274}
\end{aligned}$$

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^2(5A + 4B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + 5a^2(2A + B) \int \sqrt{\sec(c + dx)} dx \right) + \frac{2a^2(5A + 7B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^2(5A + 4B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 5a^2(2A + B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right) + \frac{2a^2(5A + 7B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)}$$

↓ 4258

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(2A + B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3a^2(5A + 4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(2A + B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a^2(5A + 4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3119

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^2(2A + B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{6a^2(5A + 4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{2a^2(5A + 7B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} + \frac{2}{3} \left(\frac{10a^2(2A + B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{6a^2(5A + 4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

input `Int[(a + a*cos[c + d*x])^2*(A + B*cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(2*B*(a^2 + a^2*Sec[c + d*x])*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((2*((6*a^2*(5*A + 4*B)*Sqrt[Cos[c + d*x])*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*a^2*(2*A + B)*Sqrt[Cos[c + d*x])*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/3 + (2*a^2*(5*A + 7*B)*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]])/5`

3.468.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4484 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]`

rule 4505 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a*A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Simp[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Simp[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && LtQ[n, -1]`

3.468.4 Maple [A] (verified)

Time = 11.11 (sec) , antiderivative size = 357, normalized size of antiderivative = 2.15

method	result
default	$-4\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} a^2 \left(-12B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (10A + 32B)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-$
parts	Expression too large to display

input `int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2), x, method=_RETURNV ERBOSE)`

3.468. $\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

output `-4/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(-12*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+(10*A+32*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-5*A-13*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-12*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.468.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.13

$$\int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx =$$

$$2 \left(5i \sqrt{2} (2A + B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (2A + B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) \right) / \sqrt{\cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="fracas")`

output `-2/15*(5*I*sqrt(2)*(2*A + B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(2*A + B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(5*A + 4*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(5*A + 4*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (3*B*a^2*cos(d*x + c)^2 + 5*(A + 2*B)*a^2*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.468.6 Sympy [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= a^2 \left(\int A \sqrt{\sec(c + dx)} dx + \int 2A \cos(c + dx) \sqrt{\sec(c + dx)} dx \right. \\ &\quad \left. + \int A \cos^2(c + dx) \sqrt{\sec(c + dx)} dx + \int B \cos(c + dx) \sqrt{\sec(c + dx)} dx \right. \\ &\quad \left. + \int 2B \cos^2(c + dx) \sqrt{\sec(c + dx)} dx + \int B \cos^3(c + dx) \sqrt{\sec(c + dx)} dx \right) \end{aligned}$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `a**2*(Integral(A*sqrt(sec(c + d*x)), x) + Integral(2*A*cos(c + d*x)*sqrt(sec(c + d*x)), x) + Integral(A*cos(c + d*x)**2*sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)*sqrt(sec(c + d*x)), x) + Integral(2*B*cos(c + d*x)**2*sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)**3*sqrt(sec(c + d*x)), x))`

3.468.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sqrt(sec(d*x + c)), x)`

3.468.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2*sqrt(sec(d*x + c)), x)`

3.468.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + a \cos(c + dx))^2 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^2,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^2, x)`

$$3.469 \quad \int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.469.1 Optimal result	4281
3.469.2 Mathematica [C] (verified)	4282
3.469.3 Rubi [A] (verified)	4282
3.469.4 Maple [A] (verified)	4287
3.469.5 Fracas [C] (verification not implemented)	4288
3.469.6 Sympy [F]	4288
3.469.7 Maxima [F]	4289
3.469.8 Giac [F]	4289
3.469.9 Mupad [F(-1)]	4290

3.469.1 Optimal result

Integrand size = 33, antiderivative size = 201

$$\begin{aligned} & \int \frac{(a+a \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx \\ &= \frac{4a^2(4A+3B)\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{\sec(c+dx)}}{5d} \\ &+ \frac{4a^2(7A+6B)\sqrt{\cos(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)\sqrt{\sec(c+dx)}}{21d} \\ &+ \frac{2a^2(7A+9B)\sin(c+dx)}{35d\sec^{\frac{3}{2}}(c+dx)} + \frac{4a^2(7A+6B)\sin(c+dx)}{21d\sqrt{\sec(c+dx)}} \\ &+ \frac{2B(a^2+a^2\sec(c+dx))\sin(c+dx)}{7d\sec^{\frac{5}{2}}(c+dx)} \end{aligned}$$

output

```
2/35*a^2*(7*A+9*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/7*B*(a^2+a^2*sec(d*x+c)
)*sin(d*x+c)/d/sec(d*x+c)^(5/2)+4/21*a^2*(7*A+6*B)*sin(d*x+c)/d/sec(d*x+c)
^(1/2)+4/5*a^2*(4*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+4
/21*a^2*(7*A+6*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
F(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```


3.469.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.77 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.96

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{a^2 e^{-idx} \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(40(7A + 6B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - 56i(4A + 3B) \sqrt{\cos(c + dx)} \right)}{\dots}$$

input `Integrate[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(a^2*Sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*(40*(7*A + 6*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (56*I)*(4*A + 3*B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((672*I)*A + (504*I)*B + 5*(56*A + 51*B)*Sin[c + d*x] + 42*(A + 2*B)*Sin[2*(c + d*x)] + 15*B*Sin[3*(c + d*x)])))/(210*d*E^(I*d*x))`

3.469.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4484, 25, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3439}$$

$$\begin{aligned}
& \int \frac{(a \sec(c+dx) + a)^2 (A \sec(c+dx) + B)}{\sec^{\frac{7}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a \csc(c+dx + \frac{\pi}{2}) + a)^2 (A \csc(c+dx + \frac{\pi}{2}) + B)}{\csc(c+dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow \text{4505} \\
& \frac{2}{7} \int \frac{(\sec(c+dx)a + a)(a(7A + 9B) + a(7A + 3B) \sec(c+dx))}{\frac{2 \sec^{\frac{5}{2}}(c+dx)}{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}} dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \frac{(\sec(c+dx)a + a)(a(7A + 9B) + a(7A + 3B) \sec(c+dx))}{\frac{\sec^{\frac{5}{2}}(c+dx)}{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \frac{(\csc(c+dx + \frac{\pi}{2})a + a)(a(7A + 9B) + a(7A + 3B) \csc(c+dx + \frac{\pi}{2}))}{\frac{\csc(c+dx + \frac{\pi}{2})^{5/2}}{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}} dx + \\
& \quad \downarrow \text{4484} \\
& \frac{1}{7} \left(\frac{2a^2(7A + 9B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} - \frac{2}{5} \int \frac{5(7A + 6B)a^2 + 7(4A + 3B) \sec(c+dx)a^2}{\sec^{\frac{3}{2}}(c+dx)} dx \right) + \\
& \quad \downarrow \text{25} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{5(7A + 6B)a^2 + 7(4A + 3B) \sec(c+dx)a^2}{\sec^{\frac{3}{2}}(c+dx)} dx + \frac{2a^2(7A + 9B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.469. $\int \frac{(a+a \cos(c+dx))^2 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{5(7A+6B)a^2 + 7(4A+3B) \csc(c+dx+\frac{\pi}{2}) a^2}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2a^2(7A+9B) \sin(c+dx)}{5d \sec^{3/2}(c+dx)} \right) + \frac{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}{7d \sec^{5/2}(c+dx)}$$

↓ 4274

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(7A+6B) \int \frac{1}{\sec^{3/2}(c+dx)} dx + 7a^2(4A+3B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right) + \frac{2a^2(7A+9B) \sin(c+dx)}{5d \sec^{3/2}(c+dx)} \right) + \frac{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}{7d \sec^{5/2}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(7A+6B) \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx + 7a^2(4A+3B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2a^2(7A+9B) \sin(c+dx)}{5d \sec^{3/2}(c+dx)} \right) + \frac{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}{7d \sec^{5/2}(c+dx)}$$

↓ 4256

$$\frac{1}{7} \left(\frac{2}{5} \left(7a^2(4A+3B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5a^2(7A+6B) \left(\frac{1}{3} \int \sqrt{\sec(c+dx)} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) \right) + \frac{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}{7d \sec^{5/2}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(7a^2(4A+3B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5a^2(7A+6B) \left(\frac{1}{3} \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) \right) + \frac{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}{7d \sec^{5/2}(c+dx)}$$

↓ 4258

$$\frac{1}{7} \left(\frac{2}{5} \left(7a^2(4A+3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + 5a^2(7A+6B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sec(c+dx)} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) \right) + \frac{2B \sin(c+dx) (a^2 \sec(c+dx) + a^2)}{7d \sec^{5/2}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(7A + 6B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) + 7a^2(4A + 3B) \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3119

$$\frac{1}{7} \left(\frac{2}{5} \left(5a^2(7A + 6B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) + \frac{14a^2(4A + 3B)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{7} \left(\frac{2a^2(7A + 9B) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2}{5} \left(\frac{14a^2(4A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + 5a^2(7A + 6B) \right) + \frac{2B \sin(c + dx) (a^2 \sec(c + dx) + a^2)}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

input `Int[((a + a*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(2*B*(a^2 + a^2*Sec[c + d*x])*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + ((2*a^2*(7*A + 9*B)*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + (2*((14*a^2*(4*A + 3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 5*a^2*(7*A + 6*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]))))/5)/7`

3.469.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4484 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]`

```
rule 4505 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*A*Cot
[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Sim
p[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Sim
p[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0
] && GtQ[m, 1/2] && LtQ[n, -1]
```

3.469.4 Maple [A] (verified)

Time = 12.98 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.92

method	result
default	$-\frac{4\sqrt{\left(2\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^2} \left(120B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-84A - 348B) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots\right)$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^2*(120*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8+(-84*A-348*B)*sin(1/2*d*x+1/2*c)^6
*cos(1/2*d*x+1/2*c)+(224*A+378*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+
(-91*A-117*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+35*A*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c
),2^(1/2))-84*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+30*B*(sin(1/2*d*x+1/2*c)^2)^(1/2
)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-6
3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elliptic
E(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.469.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.05

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$2 \left(5i \sqrt{2} (7A + 6B) a^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (7A + 6B) a^2 w \right)$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm m="fricas")`

output `-2/105*(5*I*sqrt(2)*(7*A + 6*B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(7*A + 6*B)*a^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*I*sqrt(2)*(4*A + 3*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*(4*A + 3*B)*a^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (15*B*a^2*cos(d*x + c)^3 + 21*(A + 2*B)*a^2*cos(d*x + c)^2 + 10*(7*A + 6*B)*a^2*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.469.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= a^2 \left(\int \frac{A}{\sqrt{\sec(c + dx)}} dx + \int \frac{2A \cos(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{A \cos^2(c + dx)}{\sqrt{\sec(c + dx)}} dx \right.$$

$$\left. + \int \frac{B \cos(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{2B \cos^2(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{B \cos^3(c + dx)}{\sqrt{\sec(c + dx)}} dx \right)$$

input `integrate((a+a*cos(d*x+c))**2*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `a**2*(Integral(A/sqrt(sec(c + d*x)), x) + Integral(2*A*cos(c + d*x)/sqrt(sec(c + d*x)), x) + Integral(A*cos(c + d*x)**2/sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)/sqrt(sec(c + d*x)), x) + Integral(2*B*cos(c + d*x)**2/sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)**3/sqrt(sec(c + d*x)), x))`

3.469.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/sqrt(sec(d*x + c)), x)`

3.469.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^2}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^2/sqrt(sec(d*x + c)), x)`

3.469.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^2}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^2)/(1/cos(c + d*x))^(1/2),x)`

3.470 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

3.470.1 Optimal result	4291
3.470.2 Mathematica [C] (verified)	4292
3.470.3 Rubi [A] (verified)	4292
3.470.4 Maple [B] (verified)	4297
3.470.5 Fricas [C] (verification not implemented)	4298
3.470.6 Sympy [F(-1)]	4299
3.470.7 Maxima [F]	4299
3.470.8 Giac [F]	4300
3.470.9 Mupad [F(-1)]	4300

3.470.1 Optimal result

Integrand size = 33, antiderivative size = 244

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= -\frac{4a^3(7A + 9B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{4a^3(13A + 21B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{4a^3(7A + 9B)\sqrt{\sec(c + dx)}\sin(c + dx)}{5d} + \frac{4a^3(41A + 42B)\sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{105d}$$

$$+ \frac{2aA\sec^{\frac{3}{2}}(c + dx)(a + a \sec(c + dx))^2 \sin(c + dx)}{7d}$$

$$+ \frac{2(11A + 7B)\sec^{\frac{3}{2}}(c + dx)(a^3 + a^3 \sec(c + dx))\sin(c + dx)}{35d}$$

```
output 4/105*a^3*(41*A+42*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/7*a*A*sec(d*x+c)^(3/2)*(a+a*sec(d*x+c))^2*sin(d*x+c)/d+2/35*(11*A+7*B)*sec(d*x+c)^(3/2)*(a^3+a^3*sec(d*x+c))*sin(d*x+c)/d+4/5*a^3*(7*A+9*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d-4/5*a^3*(7*A+9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+4/21*a^3*(13*A+21*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.470.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.52 (sec) , antiderivative size = 435, normalized size of antiderivative = 1.78

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{a^3 e^{-idx} (1 + \cos(c + dx))^3 \csc(c) \sec^6\left(\frac{1}{2}(c + dx)\right) \left(7\sqrt{2}(7A + 9B)e^{2idx}(-1 + e^{2ic}) \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}}\sqrt{1+e^{2i(c+dx)}}\right)}{}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `(a^3*(1 + Cos[c + d*x])^3*Csc[c]*Sec[(c + d*x)/2]^6*(7*Sqrt[2]*(7*A + 9*B)*E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] - ((-1 + E^((2*I)*c))*(21*B*(-5 + 16*E^(I*(c + d*x)) - 5*E^((2*I)*(c + d*x)) + 54*E^((3*I)*(c + d*x)) + 5*E^((4*I)*(c + d*x)) + 56*E^((5*I)*(c + d*x)) + 5*E^((6*I)*(c + d*x)) + 18*E^((7*I)*(c + d*x))) + 2*A*(-65 + 84*E^(I*(c + d*x)) - 95*E^((2*I)*(c + d*x)) + 441*E^((3*I)*(c + d*x)) + 95*E^((4*I)*(c + d*x)) + 504*E^((5*I)*(c + d*x)) + 65*E^((6*I)*(c + d*x)) + 147*E^((7*I)*(c + d*x))) + (10*I)*(13*A + 21*B)*(1 + E^((2*I)*(c + d*x)))^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])*Sqrt[Sec[c + d*x]])/(2*E^(I*(c - d*x))*(1 + E^((2*I)*(c + d*x)))^3))/(420*d*E^(I*d*x))`

3.470.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.01, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4506, 27, 3042, 4506, 3042, 4485, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{9}{2}}(c + dx)(a \cos(c + dx) + a)^3(A + B \cos(c + dx)) dx$$

↓ 3042

3.470. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \csc\left(c+dx+\frac{\pi}{2}\right)^{9/2}\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^3\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\
& \quad \downarrow \text{3439} \\
& \int \sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^3(A\sec(c+dx)+B)dx \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+a\right)^3\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)dx \\
& \quad \downarrow \text{4506} \\
& \frac{2}{7}\int\frac{1}{2}\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)^2(a(A+7B)+a(11A+7B)\sec(c+dx))dx+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)^2}{7d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7}\int\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)^2(a(A+7B)+a(11A+7B)\sec(c+dx))dx+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)^2}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7}\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(\csc\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2\left(a(A+7B)+a(11A+7B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)dx+ \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)^2}{7d} \\
& \quad \downarrow \text{4506} \\
& \frac{1}{7}\left(\frac{2}{5}\int\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)\left((8A+21B)a^2+(41A+42B)\sec(c+dx)a^2\right)dx+\frac{2(11A+7B)\sin(c+dx)}{7d}\right) \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)^2}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7}\left(\frac{2}{5}\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(\csc\left(c+dx+\frac{\pi}{2}\right)a+a\right)\left((8A+21B)a^2+(41A+42B)\csc\left(c+dx+\frac{\pi}{2}\right)a^2\right)dx+\frac{2(11A+7B)\sin(c+dx)}{7d}\right) \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)^2}{7d} \\
& \quad \downarrow \text{4485}
\end{aligned}$$

3.470. $\int(a+a\cos(c+dx))^3(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{2}{3} \int \frac{1}{2} \sqrt{\sec(c+dx)} (5(13A+21B)a^3 + 21(7A+9B) \sec(c+dx)a^3) dx + \frac{2a^3(41A+42B) \sin(c+dx) \sec(c+dx)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + a)^2}{7d} \right) \\ \downarrow 27$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \sqrt{\sec(c+dx)} (5(13A+21B)a^3 + 21(7A+9B) \sec(c+dx)a^3) dx + \frac{2a^3(41A+42B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + a)^2}{7d} \right) \\ \downarrow 3042$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} (5(13A+21B)a^3 + 21(7A+9B) \csc\left(c+dx+\frac{\pi}{2}\right)a^3) dx + \frac{2a^3(41A+42B) \sin\left(c+dx+\frac{\pi}{2}\right) \csc^{\frac{3}{2}}\left(c+dx+\frac{\pi}{2}\right)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + a)^2}{7d} \right) \\ \downarrow 4274$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(21a^3(7A+9B) \int \sec^{\frac{3}{2}}(c+dx) dx + 5a^3(13A+21B) \int \sqrt{\sec(c+dx)} dx \right) + \frac{2a^3(41A+42B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + a)^2}{7d} \right) \\ \downarrow 3042$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A+21B) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + 21a^3(7A+9B) \int \csc\left(c+dx+\frac{\pi}{2}\right)^{3/2} dx \right) + \frac{2a^3(41A+42B) \sin\left(c+dx+\frac{\pi}{2}\right) \csc^{\frac{3}{2}}\left(c+dx+\frac{\pi}{2}\right)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + a)^2}{7d} \right) \\ \downarrow 4255$$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A+21B) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + 21a^3(7A+9B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right) \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + a)^2}{7d} \right) \\ \downarrow 3042$$

3.470. $\int (a + a \cos(c+dx))^3 (A + B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc(c + dx)}} dx \right) \right) \right) \right) \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)^2}{7d}$$

↓ 4258

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc(c + dx)}} dx \right) \right) \right) \right) \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)^2}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc(c + dx)}} dx \right) \right) \right) \right) \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)^2}{7d}$$

↓ 3119

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(13A + 21B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(7A + 9B) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc(c + dx)}} dx \right) \right) \right) \right) \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)^2}{7d}$$

↓ 3120

$$\frac{1}{7} \left(\frac{2(11A + 7B) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) (a^3 \sec(c + dx) + a^3)}{5d} + \frac{2}{5} \left(\frac{2a^3(41A + 42B) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} - \int \frac{1}{\sqrt{\csc(c + dx)}} dx \right) \right) \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)^2}{7d}$$

input `Int[(a + a*cos[c + d*x])^3*(A + B*cos[c + d*x])*Sec[c + d*x]^(9/2), x]`

```
output (2*a*A*Sec[c + d*x]^(3/2)*(a + a*Sec[c + d*x])^2*Sin[c + d*x])/(7*d) + ((2
*(11*A + 7*B)*Sec[c + d*x]^(3/2)*(a^3 + a^3*Sec[c + d*x])*Sin[c + d*x])/(5
*d) + (2*((2*a^3*(41*A + 42*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + ((
10*a^3*(13*A + 21*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec
[c + d*x]])/d + 21*a^3*(7*A + 9*B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c +
d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d))
/3))/5)/7
```

3.470.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4485 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

rule 4506 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^n*Simp[a*A*d*(m + n) + B*(b*d*n) + (A*b*d*(m + n) + a*B*d*(2*m + n - 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1]`

3.470.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 901 vs. $2(268) = 536$.

Time = 179.70 (sec) , antiderivative size = 902, normalized size of antiderivative = 3.70

method	result	size
default	Expression too large to display	902
parts	Expression too large to display	1171

input `int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x,method=_RETURNV
ERBOSE)`

output

```
-16*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(1/8*B*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*
d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1
/2))+1/8*A*(-1/56*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^4-5/42*cos(1/2*d*x+1/2*c)*(-2*s
in(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)
^2+5/21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2)))+1/5*(1/8*B+3/8*A)/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/
2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-
24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*si
n(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*
x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1
/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+(3/8*
A+3/8*B)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x...
```

3.470.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.08

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx =$$

$$2 \left(5i \sqrt{2} (13A + 21B) a^3 \cos(dx + c)^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} \right)$$

input

```
integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm
m="fracas")
```

3.470. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

output
$$\begin{aligned} & -2/105*(5*I*\sqrt{2}*(13*A + 21*B)*a^3*\cos(dx + c)^3*\text{weierstrassPInverse}(- \\ & 4, 0, \cos(dx + c) + I*\sin(dx + c)) - 5*I*\sqrt{2}*(13*A + 21*B)*a^3*\cos(d \\ & *x + c)^3*\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I*\sin(dx + c)) + 21*I \\ & *\sqrt{2}*(7*A + 9*B)*a^3*\cos(dx + c)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrass} \\ & \text{PInverse}(-4, 0, \cos(dx + c) + I*\sin(dx + c))) - 21*I*\sqrt{2}*(7*A + 9*B) \\ & *a^3*\cos(dx + c)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(\\ & dx + c) - I*\sin(dx + c))) - (42*(7*A + 9*B)*a^3*\cos(dx + c)^3 + 5*(26*A \\ & + 21*B)*a^3*\cos(dx + c)^2 + 21*(3*A + B)*a^3*\cos(dx + c) + 15*A*a^3)*\text{si} \\ & \text{n}(dx + c)/\sqrt{\cos(dx + c)})/(d*\cos(dx + c)^3) \end{aligned}$$

3.470.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(9/2),x)`

output Timed out

3.470.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ & = \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{9}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))3*(A+B*cos(d*x+c))*sec(d*x+c)(9/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)3*sec(d*x + c)(9/2), x)`

3.470.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{9}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(9/2), x)`

3.470.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{9/2} (a + a \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + a*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + a*cos(c + d*x))^3, x)`

3.471 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

3.471.1 Optimal result	4301
3.471.2 Mathematica [C] (verified)	4302
3.471.3 Rubi [A] (verified)	4302
3.471.4 Maple [B] (verified)	4307
3.471.5 Fricas [C] (verification not implemented)	4308
3.471.6 Sympy [F(-1)]	4308
3.471.7 Maxima [F]	4309
3.471.8 Giac [F]	4309
3.471.9 Mupad [F(-1)]	4309

3.471.1 Optimal result

Integrand size = 33, antiderivative size = 211

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= -\frac{4a^3(9A + 5B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{4a^3(3A + 5B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{4a^3(21A + 20B)\sqrt{\sec(c + dx)}\sin(c + dx)}{15d}$$

$$+ \frac{2aA\sqrt{\sec(c + dx)}(a + a \sec(c + dx))^2 \sin(c + dx)}{5d}$$

$$+ \frac{2(9A + 5B)\sqrt{\sec(c + dx)}(a^3 + a^3 \sec(c + dx)) \sin(c + dx)}{15d}$$

```
output 4/15*a^3*(21*A+20*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d+2/5*a*A*(a+a*sec(d*x+c))^2*sin(d*x+c)*sec(d*x+c)^(1/2)/d+2/15*(9*A+5*B)*(a^3+a^3*sec(d*x+c))*sin(d*x+c)*sec(d*x+c)^(1/2)/d-4/5*a^3*(9*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+4/3*a^3*(3*A+5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.471.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.45 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.27

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{a^3 e^{-idx} \csc(c) \sec(c) \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(2(9A + 5B) e^{-i(c-dx)} (-1 + e^{4ic}) \sqrt{1 + e^{2i(c+dx)}} \operatorname{Hy} \right)}{}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(a^3*Csc[c]*Sec[c]*Sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*((2*(9*A + 5*B)*(-1 + E^((4*I)*c))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*(c - d*x)) + (Sec[c + d*x]^2*Sin[2*c]*((-18*I)*(9*A + 5*B)*Cos[c + d*x] - (54*I)*A*Cos[3*(c + d*x)] - (30*I)*B*Cos[3*(c + d*x)] + 40*(3*A + 5*B)*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 66*A*Sin[c + d*x] + 45*B*Sin[c + d*x] + 30*A*Sin[2*(c + d*x)] + 10*B*Sin[2*(c + d*x)] + 54*A*Sin[3*(c + d*x)] + 45*B*Sin[3*(c + d*x)]))/2)/(30*d*E^(I*d*x))`

3.471.3 Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.02, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4506, 27, 3042, 4506, 3042, 4485, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx) (a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

3.471. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \sec(c + dx) + a)^3 (A \sec(c + dx) + B)}{\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4506} \\
& \frac{2}{5} \int - \frac{(\sec(c + dx)a + a)^2 (a(A - 5B) - a(9A + 5B) \sec(c + dx))}{2\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \\
& \quad \downarrow \text{27} \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} - \\
& \frac{1}{5} \int \frac{(\sec(c + dx)a + a)^2 (a(A - 5B) - a(9A + 5B) \sec(c + dx))}{\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} - \\
& \frac{1}{5} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)^2 (a(A - 5B) - a(9A + 5B) \csc(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4506} \\
& \frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \int \frac{(\sec(c + dx)a + a) (a^2(6A - 5B) - a^2(21A + 5B) \sec(c + dx))}{\sqrt{\sec(c + dx)}} dx \right. \\
& \quad \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a) (a^2(6A - 5B) - a^2(21A + 5B) \csc(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right. \\
& \quad \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \\
& \quad \downarrow \text{4485}
\end{aligned}$$

3.471. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(2 \int \frac{3a^3(9A + 5B) - 5a^3(3A + 5B) \sec(c + dx)}{2\sqrt{\sec(c + dx)}} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \downarrow 27$$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(\int \frac{3a^3(9A + 5B) - 5a^3(3A + 5B) \sec(c + dx)}{\sqrt{\sec(c + dx)}} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(\int \frac{3a^3(9A + 5B) - 5a^3(3A + 5B) \csc(c + dx)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \downarrow 4274$$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(3a^3(9A + 5B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx - 5a^3(3A + 5B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(3a^3(9A + 5B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx - 5a^3(3A + 5B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \downarrow 4258$$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(-5a^3(3A + 5B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d} \right) \downarrow 3042$$

3.471. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(-5a^3(3A + 5B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \right) \right) - \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d}$$

↓ 3119

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(-5a^3(3A + 5B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \right) \right) - \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d}$$

↓ 3120

$$\frac{1}{5} \left(\frac{2(9A + 5B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^3)}{3d} - \frac{2}{3} \left(-\frac{2a^3(21A + 20B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right) - \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)^2}{5d}$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*a*A*Sqrt[Sec[c + d*x]]*(a + a*Sec[c + d*x])^2*Sin[c + d*x])/(5*d) + ((2*(9*A + 5*B)*Sqrt[Sec[c + d*x]]*(a^3 + a^3*Sec[c + d*x])*Sin[c + d*x])/(3*d) - (2*((6*a^3*(9*A + 5*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (10*a^3*(3*A + 5*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*a^3*(21*A + 20*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d))/3)/5`

3.471.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(g_.))^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)]^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4485 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.))*(\text{csc}[(e_.) + (f_.)(x_)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*((d*\text{Csc}[e + f*x])^n/(f*(n+1))), x] + \text{Simp}[1/(n+1) \text{Int}[(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*a*(n+1) + B*b*n + (A*b + B*a)*(n+1)*\text{Csc}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& !\text{LeQ}[n, -1]$

rule 4506 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m-1)}*((d*\text{Csc}[e + f*x])^n/(f*(m+n))), x] + \text{Simp}[1/(d*(m+n)) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m-1)}*(d*\text{Csc}[e + f*x])^n*\text{Simp}[a*A*d*(m+n) + B*(b*d*n) + (A*b*d*(m+n) + a*B*d*(2*m+n-1))*\text{Csc}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{GtQ}[m, 1/2] \&\& !\text{LtQ}[n, -1]$

3.471.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 915 vs. $2(239) = 478$.

Time = 181.49 (sec) , antiderivative size = 916, normalized size of antiderivative = 4.34

method	result	size
default	Expression too large to display	916
parts	Expression too large to display	1061

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/15*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3/(8*sin
(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/
2*d*x+1/2*c)^3*(216*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-60*A*Ellipti
cF(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^4-108*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*si
n(1/2*d*x+1/2*c)^4+180*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-100*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-60*B*EllipticE(cos(1/2*d*x+1/2
*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*sin(1/2*d*x+1/2*c)^4-246*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+60*A*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1
/2*d*x+1/2*c)^2-1)^(1/2)*sin(1/2*d*x+1/2*c)^2+108*A*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/
2))*sin(1/2*d*x+1/2*c)^2-190*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+100
*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF
(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+60*B*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(2*sin(1/2*d*x+1/2*c)^2-1)
^(1/2)*sin(1/2*d*x+1/2*c)^2+72*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2-1
5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellip...
```

3.471.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.15

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx =$$

$$2 \left(5i \sqrt{2} (3A + 5B) a^3 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (3A + 5B) a^3 \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 3i \sqrt{2} (9A + 5B) a^3 \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 3i \sqrt{2} (9A + 5B) a^3 \cos(dx + c)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - (9(6A + 5B) a^3 \cos(dx + c)^2 + 5(3A + B) a^3 \cos(dx + c) + 3A a^3) \sin(dx + c) / \sqrt{\cos(dx + c)} \right) / (d \cos(dx + c)^2)$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="fricas")`

output `-2/15*(5*I*sqrt(2)*(3*A + 5*B)*a^3*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(3*A + 5*B)*a^3*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(9*A + 5*B)*a^3*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(9*A + 5*B)*a^3*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (9*(6*A + 5*B)*a^3*cos(d*x + c)^2 + 5*(3*A + B)*a^3*cos(d*x + c) + 3*A*a^3)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2)`

3.471.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.471.7 Maxima [F]

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2), x)`

3.471.8 Giac [F]

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2), x)`

3.471.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + a \cos(c + dx))^3 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^3, x)`

3.471. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.472 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$

3.472.1 Optimal result	4310
3.472.2 Mathematica [C] (verified)	4311
3.472.3 Rubi [A] (verified)	4311
3.472.4 Maple [B] (verified)	4316
3.472.5 Fricas [C] (verification not implemented)	4317
3.472.6 Sympy [F(-1)]	4318
3.472.7 Maxima [F]	4318
3.472.8 Giac [F]	4319
3.472.9 Mupad [F(-1)]	4319

3.472.1 Optimal result

Integrand size = 33, antiderivative size = 199

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= -\frac{4a^3(A - B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{d} \\ & \quad + \frac{20a^3(A + B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d} \\ & \quad + \frac{4a^3(4A + B)\sqrt{\sec(c + dx)}\sin(c + dx)}{3d} + \frac{2aB(a + a \sec(c + dx))^2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\ & \quad + \frac{2(A - B)\sqrt{\sec(c + dx)}(a^3 + a^3 \sec(c + dx)) \sin(c + dx)}{3d} \end{aligned}$$

```
output 2/3*a*B*(a+a*sec(d*x+c))^2*sin(d*x+c)/d/sec(d*x+c)^(1/2)+4/3*a^3*(4*A+B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d+2/3*(A-B)*(a^3+a^3*sec(d*x+c))*sin(d*x+c)*sec(d*x+c)^(1/2)/d-4*a^3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+20/3*a^3*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.472.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.57 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.02

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{a^3 e^{-idx} \sec^{\frac{3}{2}}(c + dx) (\cos(dx) + i \sin(dx)) \left(-12iA + 12iB - 12iA \cos(2(c + dx)) + 12iB \cos(2(c + dx)) \right) + \dots}{\dots}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(a^3*Sec[c + d*x]^(3/2)*(Cos[d*x] + I*Sin[d*x])*((-12*I)*A + (12*I)*B - (12*I)*A*Cos[2*(c + d*x)] + (12*I)*B*Cos[2*(c + d*x)] + 40*(A + B)*Cos[c + d*x])^(3/2)*EllipticF[(c + d*x)/2, 2] + (4*I)*(A - B)*(1 + E^((2*I)*(c + d*x))))^(3/2)*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + 4*A*Sin[c + d*x] + B*Sin[c + d*x] + 18*A*Sin[2*(c + d*x)] + 6*B*Sin[2*(c + d*x)] + B*Sin[3*(c + d*x)])/(6*d*E^(I*d*x))`

3.472.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.01, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4506, 27, 3042, 4485, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx) (a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)^3 (A \sec(c + dx) + B)}{\sec^{\frac{3}{2}}(c + dx)} dx$$

3.472. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2}{3} \int \frac{(\sec(c + dx)a + a)^2 (a(3A + 7B) + 3a(A - B) \sec(c + dx))}{2\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{4505} \\
& \frac{1}{3} \int \frac{(\sec(c + dx)a + a)^2 (a(3A + 7B) + 3a(A - B) \sec(c + dx))}{\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)^2 (a(3A + 7B) + 3a(A - B) \csc(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(\frac{2}{3} \int \frac{3(\sec(c + dx)a + a) ((A + 4B)a^2 + (4A + B) \sec(c + dx)a^2)}{\sqrt{\sec(c + dx)}} dx + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^2)}{d} \right) \\
& \quad \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{4506} \\
& \frac{1}{3} \left(\frac{2}{3} \int \frac{3(\sec(c + dx)a + a) ((A + 4B)a^2 + (4A + B) \sec(c + dx)a^2)}{\sqrt{\sec(c + dx)}} dx + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^2)}{d} \right) \\
& \quad \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \left(\frac{2}{3} \int \frac{3(\sec(c + dx)a + a) ((A + 4B)a^2 + (4A + B) \sec(c + dx)a^2)}{\sqrt{\sec(c + dx)}} dx + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)} (a^3 \sec(c + dx) + a^2)}{d} \right) \\
& \quad \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.472. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\frac{1}{3} \left(2 \int \frac{(\csc(c + dx + \frac{\pi}{2}) a + a) ((A + 4B)a^2 + (4A + B) \csc(c + dx + \frac{\pi}{2}) a^2)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d \sqrt{\sec(c + dx)}}$$

↓ 4485

$$\frac{1}{3} \left(2 \int -\frac{3a^3(A - B) - 5a^3(A + B) \sec(c + dx)}{2\sqrt{\sec(c + dx)}} dx + \frac{2a^3(4A + B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d \sqrt{\sec(c + dx)}}$$

↓ 27

$$\frac{1}{3} \left(2 \left(\frac{2a^3(4A + B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{3a^3(A - B) - 5a^3(A + B) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx \right) + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d \sqrt{\sec(c + dx)}}$$

↓ 3042

$$\frac{1}{3} \left(2 \left(\frac{2a^3(4A + B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{3a^3(A - B) - 5a^3(A + B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \frac{2(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d \sqrt{\sec(c + dx)}}$$

↓ 4274

$$\frac{1}{3} \left(2 \left(-3a^3(A - B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + 5a^3(A + B) \int \sqrt{\sec(c + dx)} dx + \frac{2a^3(4A + B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d \sqrt{\sec(c + dx)}}$$

↓ 3042

$$\frac{1}{3} \left(2 \left(-3a^3(A - B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 5a^3(A + B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2a^3(4A + B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{3d \sqrt{\sec(c + dx)}}$$

3.472. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

↓ 4258

$$\frac{1}{3} \left(2 \left(5a^3(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^3(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{3d\sqrt{\sec(c+dx)}}} \right) \right)$$

↓ 3042

$$\frac{1}{3} \left(2 \left(5a^3(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^3(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{3d\sqrt{\sec(c+dx)}}} \right) \right)$$

↓ 3119

$$\frac{1}{3} \left(2 \left(5a^3(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2a^3(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right) \right) - \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{3d\sqrt{\sec(c+dx)}}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}(a^3 \sec(c+dx) + a^3)}{d} + 2 \left(\frac{2a^3(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} + \frac{10a^3}{3} \right) \right) - \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{3d\sqrt{\sec(c+dx)}}$$

input `Int[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*a*B*(a + a*Sec[c + d*x])^2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]) + ((2*(A - B)*Sqrt[Sec[c + d*x]]*(a^3 + a^3*Sec[c + d*x])*Sin[c + d*x])/d + 2*((-6*a^3*(A - B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*a^3*(A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a^3*(4*A + B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3`

3.472.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4485 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

```
rule 4505 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*A*Cot
[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Sim
p[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Sim
p[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0
] && GtQ[m, 1/2] && LtQ[n, -1]
```

```
rule 4506 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-b)*B*
Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))),
x] + Simp[1/(d*(m + n)) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x]
)^n*Simp[a*A*d*(m + n) + B*(b*d*n) + (A*b*d*(m + n) + a*B*d*(2*m + n - 1))*
Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b -
a*B, 0] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1]
```

3.472.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 653 vs. 2(229) = 458.

Time = 187.01 (sec) , antiderivative size = 654, normalized size of antiderivative = 3.29

method	result
default	$- \frac{4 \left(-4B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 2 \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} (9A+5B) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right)}{\dots}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

3.472. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

output

```

-4/3*(-4*B*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^
2)^(1/2)*sin(1/2*d*x+1/2*c)^6+2*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*(9*A+5*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(5*A+2*B)*sin(1/2*d*x+1/2*c)^2*c
os(1/2*d*x+1/2*c)-2*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(
2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(5*A*Elliptic
F(cos(1/2*d*x+1/2*c),2^(1/2))+3*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*
B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*B*EllipticE(cos(1/2*d*x+1/2*c),2
^(1/2)))*sin(1/2*d*x+1/2*c)^2+5*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*
d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*(-2*sin(1/2*d*
x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+3*A*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/
2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2
-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+5*B*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/
2))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)-3*B*(-2*sin(1/2*d
*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*si
n(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*a^3/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-
1)^(3/2)/sin(1/2*d*x+1/2*c)/d

```

3.472.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.08

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx =$$

$$2 \left(5i \sqrt{2} (A + B) a^3 \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (A +$$

input

```

integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorith
m="fracas")

```

3.472. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

output `-2/3*(5*I*sqrt(2)*(A + B)*a^3*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(A + B)*a^3*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*I*sqrt(2)*(A - B)*a^3*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*(A - B)*a^3*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (B*a^3*cos(d*x + c)^2 + 3*(3*A + B)*a^3*cos(d*x + c) + A*a^3)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))`

3.472.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.472.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2), x)`

3.472.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2), x)`

3.472.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + a \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^3, x)`

3.473 $\int (a+a \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$

3.473.1 Optimal result	4320
3.473.2 Mathematica [C] (verified)	4321
3.473.3 Rubi [A] (verified)	4321
3.473.4 Maple [A] (verified)	4326
3.473.5 Fricas [C] (verification not implemented)	4327
3.473.6 Sympy [F(-1)]	4327
3.473.7 Maxima [F]	4328
3.473.8 Giac [F]	4328
3.473.9 Mupad [F(-1)]	4328

3.473.1 Optimal result

Integrand size = 33, antiderivative size = 211

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{4a^3(5A + 9B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{4a^3(5A + 3B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{4a^3(5A - 6B) \sqrt{\sec(c + dx)} \sin(c + dx)}{15d} + \frac{2aB(a + a \sec(c + dx))^2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{2(5A + 9B)(a^3 + a^3 \sec(c + dx)) \sin(c + dx)}{15d \sqrt{\sec(c + dx)}}$$

```
output 2/5*a*B*(a+a*sec(d*x+c))^2*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/15*(5*A+9*B)*(a
^3+a^3*sec(d*x+c))*sin(d*x+c)/d/sec(d*x+c)^(1/2)+4/15*a^3*(5*A-6*B)*sin(d*
x+c)*sec(d*x+c)^(1/2)/d+4/5*a^3*(5*A+9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos
(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec
(d*x+c)^(1/2)/d+4/3*a^3*(5*A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x
+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(
1/2)/d
```

3.473.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.14 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.98

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{a^3 e^{-idx} \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(120iA \cos(c + dx) + 216iB \cos(c + dx) + 40(5A + 3B) \sqrt{\cos(c + dx)} \right)}{\dots}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(a^3*Sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*((120*I)*A*Cos[c + d*x] + (216*I)*B*Cos[c + d*x] + 40*(5*A + 3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (8*I)*(5*A + 9*B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + 60*A*Sin[c + d*x] + 3*B*Sin[c + d*x] + 10*A*Sin[2*(c + d*x)] + 30*B*Sin[2*(c + d*x)] + 3*B*Sin[3*(c + d*x)])/(30*d*E^(I*d*x))`

3.473.3 Rubi [A] (verified)

Time = 1.35 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.02, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4505, 3042, 4485, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx) (a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)^3 (A \sec(c + dx) + B)}{\sec^{\frac{5}{2}}(c + dx)} dx$$

3.473. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx && \downarrow \text{3042} \\
& \frac{2}{5} \int \frac{(\sec(c + dx)a + a)^2 (a(5A + 9B) + a(5A - B) \sec(c + dx))}{2 \sec^{\frac{3}{2}}(c + dx) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + && \downarrow \text{4505} \\
& \frac{1}{5} \int \frac{(\sec(c + dx)a + a)^2 (a(5A + 9B) + a(5A - B) \sec(c + dx))}{\sec^{\frac{3}{2}}(c + dx) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + && \downarrow \text{27} \\
& \frac{1}{5} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)^2 (a(5A + 9B) + a(5A - B) \csc(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + && \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{(\sec(c + dx)a + a) ((20A + 21B)a^2 + (5A - 6B) \sec(c + dx)a^2)}{\sqrt{\sec(c + dx)} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + \frac{2(5A + 9B) \sin(c + dx) (a^3 \sec(c + dx))}{3d \sqrt{\sec(c + dx)}} \right) && \downarrow \text{4505} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a) ((20A + 21B)a^2 + (5A - 6B) \csc(c + dx + \frac{\pi}{2})a^2)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + \frac{2(5A + 9B) \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) && \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{2}{3} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a) ((20A + 21B)a^2 + (5A - 6B) \csc(c + dx + \frac{\pi}{2})a^2)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + \frac{2(5A + 9B) \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) && \downarrow \text{4485}
\end{aligned}$$

$$\frac{1}{5} \left(\frac{2}{3} \left(2 \int \frac{3(5A+9B)a^3 + 5(5A+3B)\sec(c+dx)a^3}{2\sqrt{\sec(c+dx)}} dx + \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right) + \frac{2(5A+9B)\sin(c+dx)(a\sec(c+dx)+a)^2}{5d\sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{2}{3} \left(\int \frac{3(5A+9B)a^3 + 5(5A+3B)\sec(c+dx)a^3}{\sqrt{\sec(c+dx)}} dx + \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right) + \frac{2(5A+9B)\sin(c+dx)(a\sec(c+dx)+a)^2}{5d\sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(\int \frac{3(5A+9B)a^3 + 5(5A+3B)\csc(c+dx+\frac{\pi}{2})a^3}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right) + \frac{2(5A+9B)\sin(c+dx)(a\sec(c+dx)+a)^2}{5d\sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 4274

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^3(5A+9B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + 5a^3(5A+3B) \int \sqrt{\sec(c+dx)} dx + \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right) + \frac{2(5A+9B)\sin(c+dx)(a\sec(c+dx)+a)^2}{5d\sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(3a^3(5A+9B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5a^3(5A+3B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + \frac{2a^3(5A-6B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d} \right) + \frac{2(5A+9B)\sin(c+dx)(a\sec(c+dx)+a)^2}{5d\sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 4258

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^3(5A+3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^3(5A+9B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \right) + \frac{2(5A+9B)\sin(c+dx)(a\sec(c+dx)+a)^2}{5d\sec^{\frac{3}{2}}(c+dx)} \right)$$

3.473. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

↓ 3042

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^3(5A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3a^3(5A + 9B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3119

$$\frac{1}{5} \left(\frac{2}{3} \left(5a^3(5A + 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{2(5A + 9B) \sin(c + dx) (a^3 \sec(c + dx) + a^3)}{3d \sqrt{\sec(c + dx)}} + \frac{2}{3} \left(\frac{2a^3(5A - 6B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} + \frac{10a^3(5A + 3B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

input `Int[(a + a*cos[c + d*x])^3*(A + B*cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(2*a*B*(a + a*Sec[c + d*x])^2*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((2*(5*A + 9*B)*(a^3 + a^3*Sec[c + d*x])*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]) + (2*((6*a^3*(5*A + 9*B))*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*a^3*(5*A + 3*B))*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]]/d + (2*a^3*(5*A - 6*B))*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3)/5`

3.473.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_.)^(p_))*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_))*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4485 `Int[(csc[(e_.) + (f_)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_)*(x_)]*(b_.) + (a_))*((csc[(e_.) + (f_)*(x_)]*(B_.) + (A_))), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

rule 4505 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a*A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Simp[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Simp[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && LtQ[n, -1]`

3.473.4 Maple [A] (verified)

Time = 11.50 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.60

method	result
default	$-\frac{4a^3 \left(-12B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 10A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 42B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 20A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \right)}{\dots}$
parts	Expression too large to display

input `int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)`

output
$$\begin{aligned} & -4/15*a^3*(-12*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+10*A*cos(1/2*d*x+ \\ & 1/2*c)*sin(1/2*d*x+1/2*c)^4+42*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4-2 \\ & 0*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2+25*A*(sin(1/2*d*x+1/2*c)^2)^(1 \\ & /2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)) \\ & -15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-18*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2+15*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-27*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d \end{aligned}$$

3.473.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.93

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx =$$

$$2 \left(5i \sqrt{2} (5A + 3B) a^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (5A + 3B) a^3 w \right)$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `-2/15*(5*I*sqrt(2)*(5*A + 3*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(5*A + 3*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*I*sqrt(2)*(5*A + 9*B)*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*(5*A + 9*B)*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (3*B*a^3*cos(d*x + c)^2 + 5*(A + 3*B)*a^3*cos(d*x + c) + 15*A*a^3)*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.473.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.473.7 Maxima [F]

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2), x)`

3.473.8 Giac [F]

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2), x)`

3.473.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + a \cos(c + dx))^3 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^3, x)`

3.473. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.474 $\int (a+a \cos(c+dx))^3(A+B \cos(c+dx))\sqrt{\sec(c+dx)} dx$

3.474.1 Optimal result	4329
3.474.2 Mathematica [C] (verified)	4330
3.474.3 Rubi [A] (verified)	4330
3.474.4 Maple [A] (verified)	4335
3.474.5 Fricas [C] (verification not implemented)	4336
3.474.6 Sympy [F]	4336
3.474.7 Maxima [F]	4337
3.474.8 Giac [F]	4337
3.474.9 Mupad [F(-1)]	4338

3.474.1 Optimal result

Integrand size = 33, antiderivative size = 211

$$\begin{aligned} & \int (a + a \cos(c + dx))^3(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx \\ &= \frac{4a^3(9A + 7B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d} \\ & \quad + \frac{4a^3(21A + 13B)\sqrt{\cos(c + dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{21d} \\ & \quad + \frac{4a^3(42A + 41B)\sin(c + dx)}{105d\sqrt{\sec(c + dx)}} + \frac{2aB(a + a \sec(c + dx))^2 \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\ & \quad + \frac{2(7A + 11B)(a^3 + a^3 \sec(c + dx)) \sin(c + dx)}{35d \sec^{\frac{3}{2}}(c + dx)} \end{aligned}$$

```
output 2/7*a*B*(a+a*sec(d*x+c))^2*sin(d*x+c)/d/sec(d*x+c)^(5/2)+2/35*(7*A+11*B)*(
a^3+a^3*sec(d*x+c))*sin(d*x+c)/d/sec(d*x+c)^(3/2)+4/105*a^3*(42*A+41*B)*si
n(d*x+c)/d/sec(d*x+c)^(1/2)+4/5*a^3*(9*A+7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)
*sec(d*x+c)^(1/2)/d+4/21*a^3*(21*A+13*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(
1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(
d*x+c)^(1/2)/d
```


3.474.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.22 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.92

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{a^3 e^{-idx} \sqrt{\sec(c + dx)} (\cos(dx) + i \sin(dx)) \left(40(21A + 13B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - 56i(9A + 7B) E^{\left(\frac{1}{2}(c + dx)\right)} \operatorname{Sqrt}\left[1 + E^{\left(\frac{1}{2}(c + dx)\right)}\right] \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{\left(\frac{1}{2}(c + dx)\right)}\right] + \cos[c + dx] * ((168*I)*(9*A + 7*B) + 5*(84*A + 107*B)*\sin[c + dx] + 42*(A + 3*B)*\sin[2*(c + dx)] + 15*B*\sin[3*(c + dx)])\right)}{210*d*E^{\left(\frac{1}{2}(c + dx)\right)}} \right)}{d}$$

input `Integrate[(a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(a^3*Sqrt[Sec[c + d*x]]*(Cos[d*x] + I*Sin[d*x])*(40*(21*A + 13*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (56*I)*(9*A + 7*B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((168*I)*(9*A + 7*B) + 5*(84*A + 107*B)*Sin[c + d*x] + 42*(A + 3*B)*Sin[2*(c + d*x)] + 15*B*Sin[3*(c + d*x)])))/(210*d*E^(I*d*x))`

3.474.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.05, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4505, 3042, 4484, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c + dx)} (a \cos(c + dx) + a)^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + a)^3 (A \sec(c + dx) + B)}{\sec^{\frac{7}{2}}(c + dx)} dx$$

3.474. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + a)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2}{7} \int \frac{(\sec(c + dx)a + a)^2 (a(7A + 11B) + a(7A + B) \sec(c + dx))}{2 \sec^{\frac{5}{2}}(c + dx) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \\
& \quad \downarrow \text{4505} \\
& \frac{1}{7} \int \frac{(\sec(c + dx)a + a)^2 (a(7A + 11B) + a(7A + B) \sec(c + dx))}{\sec^{\frac{5}{2}}(c + dx) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a)^2 (a(7A + 11B) + a(7A + B) \csc(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{5/2} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \frac{(\sec(c + dx)a + a) ((42A + 41B)a^2 + (21A + 8B) \sec(c + dx)a^2)}{\sec^{\frac{3}{2}}(c + dx) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \frac{2(7A + 11B) \sin(c + dx) (a^3 \sec(c + dx))}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4505} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a) ((42A + 41B)a^2 + (21A + 8B) \csc(c + dx + \frac{\pi}{2})a^2)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \frac{2(7A + 11B) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{(\csc(c + dx + \frac{\pi}{2})a + a) ((42A + 41B)a^2 + (21A + 8B) \csc(c + dx + \frac{\pi}{2})a^2)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \frac{2(7A + 11B) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \\
& \quad \downarrow \text{4484}
\end{aligned}$$

3.474. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{2a^3(42A + 41B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} - \frac{2}{3} \int -\frac{21(9A + 7B)a^3 + 5(21A + 13B) \sec(c + dx)a^3}{2\sqrt{\sec(c + dx)}} dx \right) + \frac{2(7A + 11B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \frac{21(9A + 7B)a^3 + 5(21A + 13B) \sec(c + dx)a^3}{\sqrt{\sec(c + dx)}} dx + \frac{2a^3(42A + 41B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2(7A + 11B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \int \frac{21(9A + 7B)a^3 + 5(21A + 13B) \csc(c + dx + \frac{\pi}{2}) a^3}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2a^3(42A + 41B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2(7A + 11B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)}$$

↓ 4274

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(21a^3(9A + 7B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + 5a^3(21A + 13B) \int \sqrt{\sec(c + dx)} dx \right) + \frac{2a^3(42A + 41B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(21a^3(9A + 7B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 5a^3(21A + 13B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right) + \frac{2a^3(42A + 41B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

↓ 4258

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(21A + 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 21a^3(9A + 7B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \right) + \frac{2a^3(42A + 41B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(21A + 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 21a^3(9A + 7B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)} dx \right) \right) \right)$$

↓ 3119

$$\frac{1}{7} \left(\frac{2}{5} \left(\frac{1}{3} \left(5a^3(21A + 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{42a^3(9A + 7B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \int \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)} dx \right) \right) \right)$$

↓ 3120

$$\frac{1}{7} \left(\frac{2(7A + 11B) \sin(c + dx) (a^3 \sec(c + dx) + a^3)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2}{5} \left(\frac{2a^3(42A + 41B) \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} + \frac{1}{3} \left(\frac{10a^3(21A + 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \int \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{7d \sec^{\frac{5}{2}}(c + dx)} dx \right) \right) \right)$$

input `Int[(a + a*cos[c + d*x])^3*(A + B*cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(2*a*B*(a + a*Sec[c + d*x])^2*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + ((2*(7*A + 11*B)*(a^3 + a^3*Sec[c + d*x])*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + (2*(((42*a^3*(9*A + 7*B))*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*a^3*(21*A + 13*B))*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]]/d)/3 + (2*a^3*(42*A + 41*B)*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]])))/5)/7`

3.474.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_.)^(p_))*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_))*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4484 `Int[(csc[(e_.) + (f_)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_)*(x_)]*(b_.) + (a_))*((csc[(e_.) + (f_)*(x_)]*(B_.) + (A_))), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]`

```
rule 4505 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*A*Cot
[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Sim
p[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Sim
p[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0
] && GtQ[m, 1/2] && LtQ[n, -1]
```

3.474.4 Maple [A] (verified)

Time = 14.00 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.82

method	result
default	$-\frac{4\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^3} \left(120B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (-84A - 432B)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right) + \dots\right)$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(120*B*
cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8+(-84*A-432*B)*sin(1/2*d*x+1/2*c)^6
*cos(1/2*d*x+1/2*c)+(294*A+602*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+
(-126*A-208*B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+105*A*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2
*c),2^(1/2))-189*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)
^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+65*B*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)
)-147*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Elli
pticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/
2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.474. $\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

3.474.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.00

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx =$$

$$2 \left(5i \sqrt{2} (21A + 13B) a^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 5i \sqrt{2} (21A + 13B) \right)$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `-2/105*(5*I*sqrt(2)*(21*A + 13*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) - 5*I*sqrt(2)*(21*A + 13*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*I*sqrt(2)*(9*A + 7*B)*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*(9*A + 7*B)*a^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (15*B*a^3*cos(d*x + c)^3 + 21*(A + 3*B)*a^3*cos(d*x + c)^2 + 5*(21*A + 26*B)*a^3*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.474.6 Sympy [F]

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= a^3 \left(\int A \sqrt{\sec(c + dx)} dx + \int 3A \cos(c + dx) \sqrt{\sec(c + dx)} dx \right.$$

$$+ \int 3A \cos^2(c + dx) \sqrt{\sec(c + dx)} dx + \int A \cos^3(c + dx) \sqrt{\sec(c + dx)} dx$$

$$+ \int B \cos(c + dx) \sqrt{\sec(c + dx)} dx + \int 3B \cos^2(c + dx) \sqrt{\sec(c + dx)} dx$$

$$\left. + \int 3B \cos^3(c + dx) \sqrt{\sec(c + dx)} dx + \int B \cos^4(c + dx) \sqrt{\sec(c + dx)} dx \right)$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `a**3*(Integral(A*sqrt(sec(c + d*x)), x) + Integral(3*A*cos(c + d*x)*sqrt(sec(c + d*x)), x) + Integral(3*A*cos(c + d*x)**2*sqrt(sec(c + d*x)), x) + Integral(A*cos(c + d*x)**3*sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)*sqrt(sec(c + d*x)), x) + Integral(3*B*cos(c + d*x)**2*sqrt(sec(c + d*x)), x) + Integral(3*B*cos(c + d*x)**3*sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)**4*sqrt(sec(c + d*x)), x))`

3.474.7 Maxima [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A) (a \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sqrt(sec(d*x + c)), x)`

3.474.8 Giac [F]

$$\begin{aligned} & \int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A) (a \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3*sqrt(sec(d*x + c)), x)`

3.474.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + a \cos(c + dx))^3 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^3,x)`output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^3, x)`

3.475
$$\int \frac{(a+a \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.475.1 Optimal result 4339
 3.475.2 Mathematica [C] (verified) 4340
 3.475.3 Rubi [A] (verified) 4340
 3.475.4 Maple [A] (verified) 4345
 3.475.5 Fracas [C] (verification not implemented) 4346
 3.475.6 Sympy [F] 4347
 3.475.7 Maxima [F] 4347
 3.475.8 Giac [F] 4348
 3.475.9 Mupad [F(-1)] 4348

3.475.1 Optimal result

Integrand size = 33, antiderivative size = 244

$$\int \frac{(a + a \cos(c + dx))^3(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{4a^3(21A + 17B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{15d}$$

$$+ \frac{4a^3(13A + 11B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{4a^3(24A + 23B)\sin(c + dx)}{105d \sec^{\frac{3}{2}}(c + dx)} + \frac{4a^3(13A + 11B)\sin(c + dx)}{21d\sqrt{\sec(c + dx)}}$$

$$+ \frac{2aB(a + a \sec(c + dx))^2 \sin(c + dx)}{9d \sec^{\frac{7}{2}}(c + dx)} + \frac{2(9A + 13B)(a^3 + a^3 \sec(c + dx)) \sin(c + dx)}{63d \sec^{\frac{5}{2}}(c + dx)}$$

```
output 4/105*a^3*(24*A+23*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/9*a*B*(a+a*sec(d*x+c))
)^2*sin(d*x+c)/d/sec(d*x+c)^(7/2)+2/63*(9*A+13*B)*(a^3+a^3*sec(d*x+c))*si
n(d*x+c)/d/sec(d*x+c)^(5/2)+4/21*a^3*(13*A+11*B)*sin(d*x+c)/d/sec(d*x+c)^(
1/2)+4/15*a^3*(21*A+17*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+
4/21*a^3*(13*A+11*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellip
ticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.475.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.13 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.80

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{a^3 \sqrt{\sec(c + dx)} \left(240(13A + 11B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) - 112i(21A + 17B) e^{i(c+dx)} \sqrt{1 + \dots} \right)}{\dots}$$

input `Integrate[((a + a*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(a^3*Sqrt[Sec[c + d*x]]*(240*(13*A + 11*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] - (112*I)*(21*A + 17*B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))] + Cos[c + d*x]*((7056*I)*A + (5712*I)*B + 30*(107*A + 97*B)*Sin[c + d*x] + 14*(54*A + 73*B)*Sin[2*(c + d*x)] + 90*A*Ssin[3*(c + d*x)] + 270*B*Ssin[3*(c + d*x)] + 35*B*Ssin[4*(c + d*x)])))/(1260*d)`

3.475.3 Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.02, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4505, 27, 3042, 4505, 27, 3042, 4484, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \cos(c + dx) + a)^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3439}$$

$$\begin{aligned}
& \int \frac{(a \sec(c+dx) + a)^3 (A \sec(c+dx) + B)}{\sec^{\frac{9}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a \csc(c+dx + \frac{\pi}{2}) + a)^3 (A \csc(c+dx + \frac{\pi}{2}) + B)}{\csc(c+dx + \frac{\pi}{2})^{9/2}} dx \\
& \quad \downarrow \text{4505} \\
& \frac{2}{9} \int \frac{(\sec(c+dx)a + a)^2 (a(9A + 13B) + 3a(3A + B) \sec(c+dx))}{2 \sec^{\frac{7}{2}}(c+dx)} dx + \\
& \quad \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{9d \sec^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{9} \int \frac{(\sec(c+dx)a + a)^2 (a(9A + 13B) + 3a(3A + B) \sec(c+dx))}{\sec^{\frac{7}{2}}(c+dx)} dx + \\
& \quad \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{9d \sec^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \int \frac{(\csc(c+dx + \frac{\pi}{2})a + a)^2 (a(9A + 13B) + 3a(3A + B) \csc(c+dx + \frac{\pi}{2}))}{\csc(c+dx + \frac{\pi}{2})^{7/2}} dx + \\
& \quad \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{9d \sec^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{4505} \\
& \frac{1}{9} \left(\frac{2}{7} \int \frac{3(\sec(c+dx)a + a) ((24A + 23B)a^2 + 5(3A + 2B) \sec(c+dx)a^2)}{\sec^{\frac{5}{2}}(c+dx)} dx + \frac{2(9A + 13B) \sin(c+dx) (a^3 \sec(c+dx))}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{9d \sec^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{9} \left(\frac{6}{7} \int \frac{(\sec(c+dx)a + a) ((24A + 23B)a^2 + 5(3A + 2B) \sec(c+dx)a^2)}{\sec^{\frac{5}{2}}(c+dx)} dx + \frac{2(9A + 13B) \sin(c+dx) (a^3 \sec(c+dx))}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \frac{2aB \sin(c+dx)(a \sec(c+dx) + a)^2}{9d \sec^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.475. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\frac{1}{9} \left(\frac{6}{7} \int \frac{(\csc(c+dx+\frac{\pi}{2})a+a) \left((24A+23B)a^2 + 5(3A+2B)\csc(c+dx+\frac{\pi}{2})a^2 \right)}{\csc(c+dx+\frac{\pi}{2})^{5/2} \frac{2aB \sin(c+dx)(a \sec(c+dx)+a)^2}{9d \sec^{\frac{7}{2}}(c+dx)}} dx + \frac{2(9A+13B) \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \right)$$

↓ 4484

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{2a^3(24A+23B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} - \frac{2}{5} \int -\frac{15(13A+11B)a^3 + 7(21A+17B) \sec(c+dx)a^3}{2 \sec^{\frac{3}{2}}(c+dx)} dx \right) + \frac{2(9A+13B) \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \int \frac{15(13A+11B)a^3 + 7(21A+17B) \sec(c+dx)a^3}{\sec^{\frac{3}{2}}(c+dx)} dx + \frac{2a^3(24A+23B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + \frac{2(9A+13B) \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \int \frac{15(13A+11B)a^3 + 7(21A+17B) \csc(c+dx+\frac{\pi}{2})a^3}{\csc(c+dx+\frac{\pi}{2})^{3/2} \frac{2aB \sin(c+dx)(a \sec(c+dx)+a)^2}{9d \sec^{\frac{7}{2}}(c+dx)}} dx + \frac{2a^3(24A+23B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + \frac{2(9A+13B) \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \right)$$

↓ 4274

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(13A+11B) \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)} dx + 7a^3(21A+17B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right) + \frac{2a^3(24A+23B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(13A+11B) \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx + 7a^3(21A+17B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{2a^3(24A+23B) \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 4256

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(7a^3(21A + 17B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 15a^3(13A + 11B) \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \right) \right) \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{9d \sec^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(7a^3(21A + 17B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + 15a^3(13A + 11B) \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\csc(c + dx + \frac{\pi}{2})}} \right) \right) \right) \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{9d \sec^{\frac{7}{2}}(c + dx)}$$

↓ 4258

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(7a^3(21A + 17B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + 15a^3(13A + 11B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \right) \right) \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{9d \sec^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(13A + 11B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \right) \right) \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{9d \sec^{\frac{7}{2}}(c + dx)} + 7a^3$$

↓ 3119

$$\frac{1}{9} \left(\frac{6}{7} \left(\frac{1}{5} \left(15a^3(13A + 11B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \right) \right) \right) \frac{2aB \sin(c + dx)(a \sec(c + dx) + a)^2}{9d \sec^{\frac{7}{2}}(c + dx)} + \frac{14a^3}{9}$$

↓ 3120

$$\frac{1}{9} \left(\frac{(2(9A + 13B) \sin(c + dx) (a^3 \sec(c + dx) + a^3))}{7d \sec^{\frac{5}{2}}(c + dx)} + \frac{6}{7} \left(\frac{2a^3(24A + 23B) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{1}{5} \left(\frac{14a^3(21A + 17B) \sqrt{c + dx}}{9d \sec^{\frac{7}{2}}(c + dx)} \right) \right) \right)$$

input `Int[((a + a*cos[c + d*x])^3*(A + B*cos[c + d*x]))/sqrt[Sec[c + d*x]],x]`

output `(2*a*B*(a + a*Sec[c + d*x])^2*Sin[c + d*x])/(9*d*Sec[c + d*x]^(7/2)) + ((2*(9*A + 13*B)*(a^3 + a^3*Sec[c + d*x])*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + (6*((2*a^3*(24*A + 23*B)*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((14*a^3*(21*A + 17*B)*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d + 15*a^3*(13*A + 11*B)*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*sqrt[Sec[c + d*x]]))))/5)/7)/9`

3.475.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4484 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]`

rule 4505 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^m*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] - Simp[b/(a*d*n) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^(n + 1)*Simp[a*A*(m - n - 1) - b*B*n - (a*B*n + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && GtQ[m, 1/2] && LtQ[n, -1]`

3.475.4 Maple [A] (verified)

Time = 16.79 (sec) , antiderivative size = 413, normalized size of antiderivative = 1.69

method	result
default	$-\frac{4\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{a^3} \left(-560B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + (360A + 2200B)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right)$
parts	Expression too large to display

3.475.
$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$


```
input int((a+cos(d*x+c)*a)^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -4/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*a^3*(-560*B
*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10+(360*A+2200*B)*sin(1/2*d*x+1/2*c
)^8*cos(1/2*d*x+1/2*c)+(-1296*A-3412*B)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1
/2*c)+(1806*A+2702*B)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-624*A-738*
B)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+195*A*(sin(1/2*d*x+1/2*c)^2)^(1
/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))
-441*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellip
ticE(cos(1/2*d*x+1/2*c),2^(1/2))+165*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin
(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-357*B*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1
/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/
2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.475.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.95

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$2 \left(15i \sqrt{2} (13A + 11B) a^3 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) - 15i \sqrt{2} (13A + 11B) \right)$$

```
input integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorith
m="fricas")
```

```
output -2/315*(15*I*sqrt(2)*(13*A + 11*B)*a^3*weierstrassPInverse(-4, 0, cos(d*x
+ c) + I*sin(d*x + c)) - 15*I*sqrt(2)*(13*A + 11*B)*a^3*weierstrassPInvers
e(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 21*I*sqrt(2)*(21*A + 17*B)*a^3*w
eierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x
+ c))) + 21*I*sqrt(2)*(21*A + 17*B)*a^3*weierstrassZeta(-4, 0, weierstrass
PInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - (35*B*a^3*cos(d*x + c)^4
+ 45*(A + 3*B)*a^3*cos(d*x + c)^3 + 7*(27*A + 34*B)*a^3*cos(d*x + c)^2 +
30*(13*A + 11*B)*a^3*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c))/d
```

3.475. $\int \frac{(a+a \cos(c+dx))^3 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

3.475.6 Sympy [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= a^3 \left(\int \frac{A}{\sqrt{\sec(c + dx)}} dx + \int \frac{3A \cos(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{3A \cos^2(c + dx)}{\sqrt{\sec(c + dx)}} dx \right. \\ \left. + \int \frac{A \cos^3(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{B \cos(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{3B \cos^2(c + dx)}{\sqrt{\sec(c + dx)}} dx \right. \\ \left. + \int \frac{3B \cos^3(c + dx)}{\sqrt{\sec(c + dx)}} dx + \int \frac{B \cos^4(c + dx)}{\sqrt{\sec(c + dx)}} dx \right)$$

input `integrate((a+a*cos(d*x+c))**3*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `a**3*(Integral(A/sqrt(sec(c + d*x)), x) + Integral(3*A*cos(c + d*x)/sqrt(sec(c + d*x)), x) + Integral(3*A*cos(c + d*x)**2/sqrt(sec(c + d*x)), x) + Integral(A*cos(c + d*x)**3/sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)/sqrt(sec(c + d*x)), x) + Integral(3*B*cos(c + d*x)**2/sqrt(sec(c + d*x)), x) + Integral(3*B*cos(c + d*x)**3/sqrt(sec(c + d*x)), x) + Integral(B*cos(c + d*x)**4/sqrt(sec(c + d*x)), x))`

3.475.7 Maxima [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/sqrt(sec(d*x + c)), x)`

3.475.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^3}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^3/sqrt(sec(d*x + c)), x)`

3.475.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^3}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^3)/(1/cos(c + d*x))^(1/2), x)`

3.476
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+a \cos(c+dx)} dx$$

3.476.1 Optimal result 4349
 3.476.2 Mathematica [C] (verified) 4350
 3.476.3 Rubi [A] (verified) 4350
 3.476.4 Maple [B] (verified) 4354
 3.476.5 Fracas [C] (verification not implemented) 4355
 3.476.6 Sympy [F(-1)] 4356
 3.476.7 Maxima [F] 4356
 3.476.8 Giac [F] 4356
 3.476.9 Mupad [F(-1)] 4357

3.476.1 Optimal result

Integrand size = 33, antiderivative size = 193

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx \\ &= \frac{3(A - B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{ad} \\ &+ \frac{(5A - 3B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3ad} \\ &- \frac{3(A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{ad} \\ &+ \frac{(5A - 3B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3ad} - \frac{(A - B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{d(a + a \sec(c + dx))} \end{aligned}$$

output

```
1/3*(5*A-3*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d-(A-B)*sec(d*x+c)^(5/2)*sin(d
*x+c)/d/(a+a*sec(d*x+c))-3*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d+3*(A-B)*(
cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c
),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d+1/3*(5*A-3*B)*(cos(1/2*d*
x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))
*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d
```

3.476.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.69 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.12

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{\cos^2\left(\frac{1}{2}(c + dx)\right) \left(-6\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -e^{2i(c+dx)}\right]\right) + E^{i(c+dx)} \sqrt{1+e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -e^{2i(c+dx)}\right]\right)\right)}{6a^2 d \cos^2\left(\frac{1}{2}(c + dx)\right)}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x]),x]`

output `(Cos[(c + d*x)/2]^2*((-6*sqrt[2]*A*sqrt[E^(I*(c + d*x))]/(1 + E^((2*I)*(c + d*x))))*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + (6*sqrt[2]*B*sqrt[E^(I*(c + d*x))]/(1 + E^((2*I)*(c + d*x)))*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + 20*A*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] - 12*B*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] - 2*sqrt[Sec[c + d*x]]*(18*(A - B)*Cos[d*x]*Csc[c] - 2*(5*A - 3*B + 2*A*Sec[c + d*x])*Tan[(c + d*x)/2])))/(6*a*d*(1 + Cos[c + d*x]))`

3.476.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.96, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

3.476. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{a\sin\left(c+dx+\frac{\pi}{2}\right)+a} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{\sec^{5/2}(c+dx)\left(A\sec(c+dx)+B\right)}{a\sec(c+dx)+a} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{a\csc\left(c+dx+\frac{\pi}{2}\right)+a} dx \\
& \quad \downarrow \text{4507} \\
& \frac{\int -\frac{1}{2}\sec^{3/2}(c+dx)\left(3a(A-B)-a(5A-3B)\sec(c+dx)\right)dx}{a^2} - \frac{(A-B)\sin(c+dx)\sec^{5/2}(c+dx)}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{27} \\
& -\frac{\int \sec^{3/2}(c+dx)\left(3a(A-B)-a(5A-3B)\sec(c+dx)\right)dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sec^{5/2}(c+dx)}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& -\frac{\int \csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(3a(A-B)-a(5A-3B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sec^{5/2}(c+dx)}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{4274} \\
& \frac{3a(A-B)\int \sec^{3/2}(c+dx)dx - a(5A-3B)\int \sec^{5/2}(c+dx)dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sec^{5/2}(c+dx)}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& -\frac{3a(A-B)\int \csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}dx - a(5A-3B)\int \csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sec^{5/2}(c+dx)}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{4255} \\
& \frac{3a(A-B)\left(\frac{2\sin(c+dx)\sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\sec(c+dx)}}dx\right) - a(5A-3B)\left(\frac{1}{3}\int \sqrt{\sec(c+dx)}dx + \frac{2\sin(c+dx)\sec^{3/2}(c+dx)}{3d}\right)}{2a^2} - \frac{(A-B)\sin(c+dx)\sec^{5/2}(c+dx)}{d(a\sec(c+dx)+a)}
\end{aligned}$$

3.476. $\int \frac{(A+B\cos(c+dx))\sec^{5/2}(c+dx)}{a+a\cos(c+dx)} dx$

↓ 3042

$$\frac{3a(A - B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx \right) - a(5A - 3B) \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c+dx) \sec(c+dx)}{3d} \right)}{2a^2} \\ \frac{(A - B) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{d(a \sec(c + dx) + a)}$$

↓ 4258

$$\frac{3a(A - B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \right) - a(5A - 3B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \right)}{2a^2} \\ \frac{(A - B) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{d(a \sec(c + dx) + a)}$$

↓ 3042

$$\frac{3a(A - B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) - a(5A - 3B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \right)}{2a^2} \\ \frac{(A - B) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{d(a \sec(c + dx) + a)}$$

↓ 3119

$$\frac{3a(A - B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d} \right) - a(5A - 3B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \right)}{2a^2} \\ \frac{(A - B) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{d(a \sec(c + dx) + a)}$$

↓ 3120

$$\frac{3a(A - B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d} \right) - a(5A - 3B) \left(\frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{2 \sqrt{\cos(c+dx)}}{3} \right)}{2a^2} \\ \frac{(A - B) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{d(a \sec(c + dx) + a)}$$

input `Int[((A + B*cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*cos[c + d*x]),x]`

3.476. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+a \cos(c+dx)} dx$

```
output -(((A - B)*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))) - (3
*a*(A - B)*((-2*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c +
d*x]])/d + (2*sqrt[Sec[c + d*x]]*Sin[c + d*x])/d) - a*(5*A - 3*B)*((2*sqrt
[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (2*Se
c[c + d*x]^(3/2)*Sin[c + d*x])/(3*d))/(2*a^2)
```

3.476.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_) + (f_)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_) + (f_)*
(x_)])^(m_.)*((c_) + (d_.)*sin[(e_) + (f_)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4255 Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```


rule 4274 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4507 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

3.476.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(227) = 454.

Time = 8.75 (sec) , antiderivative size = 466, normalized size of antiderivative = 2.41

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1}(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2})}} \left(\frac{(A-B)\left(\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\right) (F(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2}) - E(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2}))}{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2})}} \right)$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*a), x, method=_RETURNVERBOSE)`

$$3.476. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+a \cos(c+dx)} dx$$

output
$$\begin{aligned} & -(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/a*((A-B)*(\cos(1/2*d*x+1/2*c)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)} \\ &)*(\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})) \\ &)-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)/\cos(1/2*d*x+1/2*c)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}+2*A*(-1/6*\cos(1/2*d*x+1/2*c) \\ &)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(\cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))+(2*B-2*A)/\sin(1/2*d*x+1/2*c)^2/(2*\sin(1/2*d*x+1/2*c)^2-1)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})))/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.476.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.60

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{(\sqrt{2}(-5i A + 3i B) \cos(dx + c)^2 + \sqrt{2}(-5i A + 3i B) \cos(dx + c)) \text{weierstrassPInverse}(-4, 0, \cos(dx + c))}{1}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & 1/6*((\text{sqrt}(2)*(-5*I*A + 3*I*B)*\cos(d*x + c)^2 + \text{sqrt}(2)*(-5*I*A + 3*I*B)*\cos(d*x + c))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + (\text{sqrt}(2)*(5*I*A - 3*I*B)*\cos(d*x + c)^2 + \text{sqrt}(2)*(5*I*A - 3*I*B)*\cos(d*x + c))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 9*(\text{sqrt}(2)*(-I*A + I*B)*\cos(d*x + c)^2 + \text{sqrt}(2)*(-I*A + I*B)*\cos(d*x + c))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - 9*(\text{sqrt}(2)*(I*A - I*B)*\cos(d*x + c)^2 + \text{sqrt}(2)*(I*A - I*B)*\cos(d*x + c))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) - 2*(9*(A - B)*\cos(d*x + c)^2 + 2*(2*A - 3*B)*\cos(d*x + c) - 2*A)*\sin(d*x + c)/\text{sqrt}(\cos(d*x + c)))/(a*d*\cos(d*x + c)^2 + a*d*\cos(d*x + c)) \end{aligned}$$

3.476.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+a \cos(c+dx)} dx$$

3.476.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+a*cos(d*x+c)),x)`

output `Timed out`

3.476.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{a \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(a*cos(d*x + c) + a), x)`

3.476.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{a \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(a*cos(d*x + c) + a), x)`

3.476.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{a + a \cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x)),x)`output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x)), x)`

3.477
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+a \cos(c+dx)} dx$$

3.477.1 Optimal result 4358
 3.477.2 Mathematica [C] (verified) 4359
 3.477.3 Rubi [A] (verified) 4359
 3.477.4 Maple [A] (verified) 4363
 3.477.5 Fricas [C] (verification not implemented) 4364
 3.477.6 Sympy [F(-1)] 4364
 3.477.7 Maxima [F] 4365
 3.477.8 Giac [F] 4365
 3.477.9 Mupad [F(-1)] 4365

3.477.1 Optimal result

Integrand size = 33, antiderivative size = 159

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx \\ &= -\frac{(3A - B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{ad} \\ & \quad - \frac{(A - B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{ad} \\ & \quad + \frac{(3A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{ad} - \frac{(A - B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{d(a + a \sec(c + dx))} \end{aligned}$$

output

```
-(A-B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*sec(d*x+c))+
(3*A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d-
(3*A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)
*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/
a/d-(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)
*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d
```

3.477.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.20 (sec) , antiderivative size = 400, normalized size of antiderivative = 2.52

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{\cos^2\left(\frac{1}{2}(c + dx)\right) \left(6\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1 + e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1 + e^{2i(c+dx)}} + e^{2idx}(-1 + e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -e^{2i(c+dx)}\right]\right)\right)}{6a d (1 + \cos(c + dx))}$$

```
input Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x]),x
]
```

```
output (Cos[(c + d*x)/2]^2*((6*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c +
d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*Sqrt[1 + E^((2*I)*(c + d*
x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -
E^((2*I)*(c + d*x))])]/E^(I*d*x) - (2*Sqrt[2]*B*Sqrt[E^(I*(c + d*x))/(1 +
E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*Sqrt[1 + E^
((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/
2, 3/4, 7/4, -E^((2*I)*(c + d*x))])]/E^(I*d*x) - 12*A*Sqrt[Cos[c + d*x]]*E
llipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]] + 12*B*Sqrt[Cos[c + d*x]]*Elli
pticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]] + 6*Sqrt[Sec[c + d*x]]*(2*(3*A -
B)*Cos[d*x]*Csc[c] + 2*(-A + B)*Tan[(c + d*x)/2])))/(6*a*d*(1 + Cos[c + d*
x]))
```

3.477.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.99,
 number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules
 used = {3042, 3439, 3042, 4507, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

↓ 3042

3.477. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{a \sin(c+dx+\frac{\pi}{2})+a} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{\sec^{\frac{3}{2}}(c+dx)(A \sec(c+dx)+B)}{a \sec(c+dx)+a} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2} (A \csc(c+dx+\frac{\pi}{2})+B)}{a \csc(c+dx+\frac{\pi}{2})+a} dx \\
& \quad \downarrow \text{4507} \\
& \frac{\int -\frac{1}{2} \sqrt{\sec(c+dx)}(a(A-B)-a(3A-B) \sec(c+dx)) dx}{a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} \\
& \quad \downarrow \text{27} \\
& - \frac{\int \sqrt{\sec(c+dx)}(a(A-B)-a(3A-B) \sec(c+dx)) dx}{2a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& - \frac{\int \sqrt{\csc(c+dx+\frac{\pi}{2})}(a(A-B)-a(3A-B) \csc(c+dx+\frac{\pi}{2})) dx}{2a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} \\
& \quad \downarrow \text{4274} \\
& - \frac{a(A-B) \int \sqrt{\sec(c+dx)} dx - a(3A-B) \int \sec^{\frac{3}{2}}(c+dx) dx}{2a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& - \frac{a(A-B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - a(3A-B) \int \csc(c+dx+\frac{\pi}{2})^{3/2} dx}{2a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} \\
& \quad \downarrow \text{4255} \\
& - \frac{a(A-B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - a(3A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right)}{2a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)}
\end{aligned}$$

3.477. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{a(A-B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - a(3A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right)}{2a^2} \\
& \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx) + a)} \\
& \downarrow 4258 \\
& \frac{a(A-B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx - a(3A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \right)}{2a^2} \\
& \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx) + a)} \\
& \downarrow 3042 \\
& \frac{a(A-B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - a(3A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \right)}{2a^2} \\
& \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx) + a)} \\
& \downarrow 3119 \\
& \frac{a(A-B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - a(3A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d} \right)}{2a^2} \\
& \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx) + a)} \\
& \downarrow 3120 \\
& \frac{2a(A-B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - a(3A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d} \right)}{2a^2} \\
& \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx) + a)}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x]),x]`

3.477. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+a \cos(c+dx)} dx$


```
output -(((A - B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))) - ((
2*a*(A - B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]
])/d - a*(3*A - B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[
Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/(2*a^2)
```

3.477.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_) + (f_)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_) + (f_.)*
(x_)])^(m_.)*((c_) + (d_.)*sin[(e_) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4255 Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4507 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

3.477.4 Maple [A] (verified)

Time = 4.78 (sec) , antiderivative size = 319, normalized size of antiderivative = 2.01

method	result
default	$-\frac{\sqrt{-(-2(\cos(\frac{dx}{2} + \frac{c}{2})) + 1)(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \left(\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))}\right)}{\dots}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/a*(\cos(1/2*d*x+ \\ & 1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(\sin(1/2*d*x+ \\ & 1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(A*\text{EllipticF}(\cos(1/2*d*x+ \\ & 1/2*c), 2^{(1/2)})-3*A*\text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})-B*\text{EllipticF}(\cos(1 \\ & /2*d*x+1/2*c), 2^{(1/2)})+B*\text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^{(1/2)}))-2*(-2*\sin(\\ & 1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(3*A-B)*\sin(1/2*d*x+1/2*c)^4+ \\ & (-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(5*A-B)*\sin(1/2*d*x+ \\ & 1/2*c)^2)/\cos(1/2*d*x+1/2*c)/\sin(1/2*d*x+1/2*c)^3/(2*\sin(1/2*d*x+1/2*c)^2-1 \\ &)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d \end{aligned}$$

3.477.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.57

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx$$

$$= \frac{(\sqrt{2}(i A - i B) \cos(dx + c) + \sqrt{2}(i A - i B)) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) +$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c)),x, algorithm="fricas")
```

```
output 1/2*((sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + (sqrt(2)*(-3*I*A + I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + (sqrt(2)*(3*I*A - I*B)*cos(d*x + c) + sqrt(2)*(3*I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*((3*A - B)*cos(d*x + c) + 2*A)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a*d*cos(d*x + c) + a*d)
```

3.477.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c)),x)
```

```
output Timed out
```

3.477.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{a \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a), x)`

3.477.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{a \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a), x)`

3.477.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + a \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{a + a \cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x)),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x)), x)`

3.478
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{a+a \cos(c+dx)} dx$$

3.478.1 Optimal result 4366
 3.478.2 Mathematica [C] (verified) 4367
 3.478.3 Rubi [A] (verified) 4367
 3.478.4 Maple [A] (verified) 4370
 3.478.5 Fricas [C] (verification not implemented) 4371
 3.478.6 Sympy [F] 4372
 3.478.7 Maxima [F] 4372
 3.478.8 Giac [F] 4372
 3.478.9 Mupad [F(-1)] 4373

3.478.1 Optimal result

Integrand size = 33, antiderivative size = 123

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx \\ &= \frac{(A - B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{ad} \\ &+ \frac{(A + B)\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{ad} \\ &- \frac{(A - B)\sqrt{\sec(c + dx)} \sin(c + dx)}{d(a + a \sec(c + dx))} \end{aligned}$$

```
output -(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*sec(d*x+c))+
(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*
sec(d*x+c)^(1/2)/a/d+(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*
cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d
```

3.478.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.07 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.63

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx =$$

$$\frac{e^{-\frac{1}{2}i(4c+dx)}(-1 + e^{2ic}) \left(3i(A + B) (1 + e^{i(c+dx)}) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + (A - B) \left(-3\right) \right)}{-}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x]),x]`

output `-1/24*((-1 + E^((2*I)*c))*((3*I)*(A + B)*(1 + E^(I*(c + d*x)))*Sqrt[Cos[c + d*x]])*EllipticF[(c + d*x)/2, 2] + (A - B)*(-3*(1 + E^((2*I)*(c + d*x))) + E^(I*(c + d*x))*(1 + E^(I*(c + d*x)))*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*(Csc[c/2] + I*Sec[c/2])*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]])/(a*d*E^((I/2)*(4*c + d*x)))`

3.478.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{a \cos(c + dx) + a} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\csc(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{a \sin(c + dx + \frac{\pi}{2}) + a} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{\sqrt{\sec(c + dx)}(A \sec(c + dx) + B)}{a \sec(c + dx) + a} dx$$

3.478. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{a+a \cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})} (A \csc(c+dx+\frac{\pi}{2}) + B)}{a \csc(c+dx+\frac{\pi}{2}) + a} dx \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(A-B)+a(A+B)\sec(c+dx)}{2\sqrt{\sec(c+dx)}} dx}{a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{4507} \\
& \frac{\int \frac{a(A-B)+a(A+B)\sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(A-B)+a(A+B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(A-B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + a(A+B) \int \sqrt{\sec(c+dx)} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{4274} \\
& \frac{a(A-B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + a(A+B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{4258} \\
& \frac{a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)}
\end{aligned}$$

3.478. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{a+a \cos(c+dx)} dx$

↓ 3119

$$\frac{a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2\frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)}}$$

↓ 3120

$$\frac{\frac{2a(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}(\frac{1}{2}(c+dx),2)}{d}+\frac{2a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E(\frac{1}{2}(c+dx)|2)}{d}}{2a^2\frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)}}$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x]),x]`

output `((2*a*(A - B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*(A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/(2*a^2) - ((A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))`

3.478.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`


```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^(m_.))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

```
rule 4274 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

```
rule 4507 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]
```

3.478.4 Maple [A] (verified)

Time = 3.64 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.98

method	result
default	$\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(-\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\left(AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - AE\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right) + a\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)}{a^2}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a), x, method=_RETURNVER BOSE)
```

$$3.478. \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{a+a \cos(c+dx)} dx$$

output $((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-\cos(1/2*d*x+1/2*c))*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(A*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})-A*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+B*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+B*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})))+(-2*B+2*A)*\sin(1/2*d*x+1/2*c)^4+(-A+B)*\sin(1/2*d*x+1/2*c)^2)/a/\cos(1/2*d*x+1/2*c)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$

3.478.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.96

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx =$$

$$\frac{2(A - B)\sqrt{\cos(dx + c)}\sin(dx + c) - (\sqrt{2}(-iA - iB)\cos(dx + c) + \sqrt{2}(-iA - iB))\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i\sin(dx + c))}{(a*d*\cos(dx + c) + a*d)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c)),x, algorithm="fricas")`

output $-1/2*(2*(A - B)*\sqrt{\cos(d*x + c)}*\sin(d*x + c) - (\sqrt{2})*(-I*A - I*B)*\cos(d*x + c) + \sqrt{2})*(-I*A - I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) - (\sqrt{2})*(I*A + I*B)*\cos(d*x + c) + \sqrt{2})*(I*A + I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - (\sqrt{2})*(I*A - I*B)*\cos(d*x + c) + \sqrt{2})*(I*A - I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - (\sqrt{2})*(-I*A + I*B)*\cos(d*x + c) + \sqrt{2})*(-I*A + I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/(a*d*\cos(d*x + c) + a*d)$

3.478.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx = \int \frac{A\sqrt{\sec(c+dx)}}{\cos(c+dx)+1} dx + \int \frac{B \cos(c+dx)\sqrt{\sec(c+dx)}}{\cos(c+dx)+1} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c)), x)`

output `(Integral(A*sqrt(sec(c + d*x))/(cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sqrt(sec(c + d*x))/(cos(c + d*x) + 1), x))/a`

3.478.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{a \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c)), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a), x)`

3.478.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{a \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c)), x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a), x)`

3.478.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + a \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{a + a \cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x)),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x)), x)`

3.479 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))\sqrt{\sec(c+dx)}} dx$

3.479.1 Optimal result 4374
 3.479.2 Mathematica [C] (verified) 4375
 3.479.3 Rubi [A] (verified) 4375
 3.479.4 Maple [A] (verified) 4378
 3.479.5 Fricas [C] (verification not implemented) 4379
 3.479.6 Sympy [F] 4380
 3.479.7 Maxima [F] 4380
 3.479.8 Giac [F] 4380
 3.479.9 Mupad [F(-1)] 4381

3.479.1 Optimal result

Integrand size = 33, antiderivative size = 125

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

$$= -\frac{(A - 3B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{ad}$$

$$+ \frac{(A - B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{ad}$$

$$+ \frac{(A - B)\sqrt{\sec(c + dx)}\sin(c + dx)}{d(a + a \sec(c + dx))}$$

output

```
(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*sec(d*x+c))-
(A-3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*
sec(d*x+c)^(1/2)/a/d+(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*
cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d
```

3.479.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.94 (sec) , antiderivative size = 422, normalized size of antiderivative = 3.38

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

$$= \frac{\cos^2\left(\frac{1}{2}(c + dx)\right) \left(2\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1 + e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1 + e^{2i(c+dx)}} + e^{2idx}(-1 + e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -e^{2i(c+dx)}\right]\right) + E^{i d x}(-1 + E^{(2I)c}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{(2I)(c+dx)}\right]\right)}{E^{i d x} - (6\sqrt{2}B\sqrt{E^{i(c+dx)}}/(1 + E^{(2I)(c+dx)}))\sqrt{1 + E^{(2I)(c+dx)}}\csc(c) \left(-3\sqrt{1 + E^{(2I)(c+dx)}} + E^{(2I)d x}(-1 + E^{(2I)c}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{(2I)(c+dx)}\right]\right) + (6((A - 2B)\cos((c - dx)/2) - B\cos((3c + dx)/2))\csc(c/2)\sec(c/2)\sec((c + dx)/2))/\sqrt{\sec(c + dx)} + 12A\sqrt{\cos(c + dx)}\operatorname{EllipticF}[(c + dx)/2, 2]\sqrt{\sec(c + dx)} - 12B\sqrt{\cos(c + dx)}\operatorname{EllipticF}[(c + dx)/2, 2]\sqrt{\sec(c + dx)}}}{(6a d (1 + \cos(c + dx)))}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])*Sqrt[Sec[c + d*x]]),x]`

output `(Cos[(c + d*x)/2]^2*((2*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))]] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]])/E^(I*d*x) - (6*Sqrt[2]*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))]] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]])/E^(I*d*x) + (6*((A - 2*B)*Cos[(c - d*x)/2] - B*Cos[(3*c + d*x)/2])*Csc[c/2]*Sec[c/2]*Sec[(c + d*x)/2])/Sqrt[Sec[c + d*x]] + 12*A*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]] - 12*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(6*a*d*(1 + Cos[c + d*x]))`

3.479.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a \cos(c + dx) + a)} dx$$

↓ 3042

3.479. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))\sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{A + B \sin \left(c + dx + \frac{\pi}{2} \right)}{\sqrt{\csc \left(c + dx + \frac{\pi}{2} \right) \left(a \sin \left(c + dx + \frac{\pi}{2} \right) + a \right)}} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{A \sec(c + dx) + B}{\sqrt{\sec(c + dx) \left(a \sec(c + dx) + a \right)}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A \csc \left(c + dx + \frac{\pi}{2} \right) + B}{\sqrt{\csc \left(c + dx + \frac{\pi}{2} \right) \left(a \csc \left(c + dx + \frac{\pi}{2} \right) + a \right)}} dx \\
& \quad \downarrow \text{4508} \\
& \frac{\int -\frac{a(A-3B)-a(A-B)\sec(c+dx)}{2\sqrt{\sec(c+dx)}} dx}{a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{27} \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{\int \frac{a(A-3B)-a(A-B)\sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{\int \frac{a(A-3B)-a(A-B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{2a^2} \\
& \quad \downarrow \text{4274} \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{a(A-3B)\int \frac{1}{\sqrt{\sec(c+dx)}} dx - a(A-B)\int \sqrt{\sec(c+dx)} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{a(A-3B)\int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - a(A-B)\int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{2a^2} \\
& \quad \downarrow \text{4258} \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{a(A-3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int \sqrt{\cos(c+dx)} dx - a(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{2a^2}
\end{aligned}$$

3.479. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))\sqrt{\sec(c+dx)}} dx$

$$\begin{array}{c}
\downarrow \text{3042} \\
\frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a \sec(c + dx) + a)} - \\
\frac{a(A - 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx - a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx)}} dx}{2a^2} \\
\downarrow \text{3119} \\
\frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a \sec(c + dx) + a)} - \\
\frac{\frac{2a(A - 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2}(c + dx)|2)}{d} - a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{2a^2} \\
\downarrow \text{3120} \\
\frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a \sec(c + dx) + a)} - \\
\frac{\frac{2a(A - 3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2}(c + dx)|2)}{d} - \frac{2a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticF}(\frac{1}{2}(c + dx), 2)}{d}}{2a^2}
\end{array}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])*Sqrt[Sec[c + d*x]]), x]`

output `-1/2*((2*a*(A - 3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*a*(A - B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d/a^2 + ((A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))`

3.479.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.479. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sqrt{\sec(c+dx)}} dx$

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(- (A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.479.4 Maple [A] (verified)

Time = 4.71 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.95

method	result
default	$-\frac{\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \left(AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + AE\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{a \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)/sec(d*x+c)^(1/2), x, method=_RETURNVERBOSE)`

$$3.479. \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sqrt{\sec(c+dx)}} dx$$

output $-\left((2\cos(1/2dx+1/2c)^2-1)\sin(1/2dx+1/2c)^2\right)^{1/2}\left(\cos(1/2dx+1/2c)\right)\left(2\sin(1/2dx+1/2c)^2-1\right)^{1/2}\left(\sin(1/2dx+1/2c)^2\right)^{1/2}\left(A\operatorname{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})+A\operatorname{EllipticE}(\cos(1/2dx+1/2c),2^{1/2})-B\operatorname{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})-3B\operatorname{EllipticE}(\cos(1/2dx+1/2c),2^{1/2})\right)+(-2B+2A)\sin(1/2dx+1/2c)^4+(-A+B)\sin(1/2dx+1/2c)^2/a/\cos(1/2dx+1/2c)/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}/\sin(1/2dx+1/2c)/(2\cos(1/2dx+1/2c)^2-1)^{1/2}/d$

3.479.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.90

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

$$= \frac{2(A - B)\sqrt{\cos(dx + c)}\sin(dx + c) + (\sqrt{2}(-iA + iB)\cos(dx + c) + \sqrt{2}(-iA + iB))\operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i\sin(dx + c))}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fracas")`

output $1/2*(2*(A - B)*\sqrt{\cos(d*x + c)}*\sin(d*x + c) + (\sqrt{2})*(-I*A + I*B)*\cos(d*x + c) + \sqrt{2})*(-I*A + I*B)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + (\sqrt{2})*(I*A - I*B)*\cos(d*x + c) + \sqrt{2})*(I*A - I*B)*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) + (\sqrt{2})*(-I*A + 3*I*B)*\cos(d*x + c) + \sqrt{2})*(-I*A + 3*I*B)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + (\sqrt{2})*(I*A - 3*I*B)*\cos(d*x + c) + \sqrt{2})*(I*A - 3*I*B)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/(a*d*\cos(d*x + c) + a*d)$

3.479.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

$$= \frac{\int \frac{A}{\cos(c+dx)\sqrt{\sec(c+dx)}+\sqrt{\sec(c+dx)}} dx + \int \frac{B \cos(c+dx)}{\cos(c+dx)\sqrt{\sec(c+dx)}+\sqrt{\sec(c+dx)}} dx}{a}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)**(1/2), x)`

output `(Integral(A/(cos(c + d*x)*sqrt(sec(c + d*x)) + sqrt(sec(c + d*x))), x) + Integral(B*cos(c + d*x)/(cos(c + d*x)*sqrt(sec(c + d*x)) + sqrt(sec(c + d*x))), x))/a`

3.479.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)\sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(1/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.479.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)\sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(1/2), x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.479.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + a \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))), x)`

3.480
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$$

3.480.1 Optimal result 4382
 3.480.2 Mathematica [C] (verified) 4383
 3.480.3 Rubi [A] (verified) 4383
 3.480.4 Maple [A] (verified) 4387
 3.480.5 Fracas [C] (verification not implemented) 4387
 3.480.6 Sympy [F] 4388
 3.480.7 Maxima [F] 4388
 3.480.8 Giac [F] 4389
 3.480.9 Mupad [F(-1)] 4389

3.480.1 Optimal result

Integrand size = 33, antiderivative size = 163

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{3(A - B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{ad} - \frac{(3A - 5B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3ad} - \frac{(3A - 5B) \sin(c + dx)}{3ad \sqrt{\sec(c + dx)}} + \frac{(A - B) \sin(c + dx)}{d \sqrt{\sec(c + dx)}(a + a \sec(c + dx))}$$

output

```
-1/3*(3*A-5*B)*sin(d*x+c)/a/d/sec(d*x+c)^(1/2)+(A-B)*sin(d*x+c)/d/(a+a*sec
(d*x+c))/sec(d*x+c)^(1/2)+3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x
+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(
1/2)/a/d-1/3*(3*A-5*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*El
lipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d
```

3.480.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.74 (sec) , antiderivative size = 444, normalized size of antiderivative = 2.72

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\cos^2\left(\frac{1}{2}(c + dx)\right) \left(-6\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}\right]\right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])*Sec[c + d*x]^(3/2)),x]`

output `(Cos[(c + d*x)/2]^2*((-6*sqrt[2]*A*sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + (6*sqrt[2]*B*sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) - 12*A*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] + 20*B*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] - (Csc[c/2]*Sec[c/2]*Sec[(c + d*x)/2]*((12*A - 13*B)*Cos[(c - d*x)/2] + (6*A - 5*B)*Cos[(3*c + d*x)/2] - 2*B*Sin[c]*Sin[(3*(c + d*x))/2]))/sqrt[Sec[c + d*x]]))/(6*a*d*(1 + Cos[c + d*x]))`

3.480.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.98, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)} dx$$

↓ 3042

3.480. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2} (a \sin(c + dx + \frac{\pi}{2}) + a)} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{A \sec(c + dx) + B}{\sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A \csc(c + dx + \frac{\pi}{2}) + B}{\csc(c + dx + \frac{\pi}{2})^{3/2} (a \csc(c + dx + \frac{\pi}{2}) + a)} dx \\
& \quad \downarrow \text{4508} \\
& \frac{\int -\frac{a(3A-5B)-3a(A-B)\sec(c+dx)}{2\sec^{\frac{3}{2}}(c+dx)} dx}{a^2} + \frac{(A-B)\sin(c+dx)}{d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{27} \\
& \frac{(A-B)\sin(c+dx)}{d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)} - \frac{\int \frac{a(3A-5B)-3a(A-B)\sec(c+dx)}{\sec^{\frac{3}{2}}(c+dx)} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\sin(c+dx)}{d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)} - \frac{\int \frac{a(3A-5B)-3a(A-B)\csc(c+dx+\frac{\pi}{2})}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx}{2a^2} \\
& \quad \downarrow \text{4274} \\
& \frac{(A-B)\sin(c+dx)}{d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)} - \frac{a(3A-5B)\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)} dx - 3a(A-B)\int \frac{1}{\sqrt{\sec(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\sin(c+dx)}{d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)} - \frac{a(3A-5B)\int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx - 3a(A-B)\int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{2a^2} \\
& \quad \downarrow \text{4256} \\
& \frac{(A-B)\sin(c+dx)}{d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)} - \frac{a(3A-5B)\left(\frac{1}{3}\int \sqrt{\sec(c+dx)} dx + \frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}}\right) - 3a(A-B)\int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{2a^2}
\end{aligned}$$

3.480. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{(A - B) \sin(c + dx)}{d \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)} - \\
& \frac{a(3A - 5B) \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) - 3a(A - B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx}{2a^2} \\
& \downarrow 4258 \\
& \frac{(A - B) \sin(c + dx)}{d \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)} - \\
& \frac{a(3A - 5B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) - 3a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2a^2} \\
& \downarrow 3042 \\
& \frac{(A - B) \sin(c + dx)}{d \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)} - \\
& \frac{a(3A - 5B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) - 3a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2a^2} \\
& \downarrow 3119 \\
& \frac{(A - B) \sin(c + dx)}{d \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)} - \\
& \frac{a(3A - 5B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) - \frac{6a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx)\right)}{d}}{2a^2} \\
& \downarrow 3120 \\
& \frac{(A - B) \sin(c + dx)}{d \sqrt{\sec(c + dx)} (a \sec(c + dx) + a)} - \\
& \frac{a(3A - 5B) \left(\frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} + \frac{2 \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} \right) - \frac{6a(A - B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx)\right)}{d}}{2a^2}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])*Sec[c + d*x]^(3/2)),x]`

output `((A - B)*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]]*(a + a*Sec[c + d*x])) - ((-6*a*(A - B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + a*(3*A - 5*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]])))/(2*a^2)`

3.480. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

3.480.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^(n_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4508 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(- (A*b
- a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m +
1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc
c[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[
e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B
, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]
```

3.480.4 Maple [A] (verified)

Time = 5.45 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.61

method	result
default	$\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}\left(3AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + 9AE\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)\right)}{3a\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)/sec(d*x+c)^(3/2), x, method=_RETURNVER
BOSE)
```

```
output 1/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(cos(1/2*d*x+1
/2*c)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(3*A*E
llipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+9*A*EllipticE(cos(1/2*d*x+1/2*c), 2^(1
/2))-5*B*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-9*B*EllipticE(cos(1/2*d*x+1
/2*c), 2^(1/2)))+8*B*sin(1/2*d*x+1/2*c)^6+(6*A-18*B)*sin(1/2*d*x+1/2*c)^4+(
-3*A+7*B)*sin(1/2*d*x+1/2*c)^2)/a/cos(1/2*d*x+1/2*c)/(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2
-1)^(1/2)/d
```

3.480.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.60

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(\sqrt{2}(3i A - 5i B) \cos(dx + c) + \sqrt{2}(3i A - 5i B)) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\dots}$$

3.480. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `1/6*((sqrt(2)*(3*I*A - 5*I*B)*cos(d*x + c) + sqrt(2)*(3*I*A - 5*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(-3*I*A + 5*I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A + 5*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 9*(sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 9*(sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(2*B*cos(d*x + c)^2 - (3*A - 5*B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a*d*cos(d*x + c) + a*d)`

3.480.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\int \frac{A}{\cos(c+dx) \sec^{\frac{3}{2}}(c+dx) + \sec^{\frac{3}{2}}(c+dx)} dx + \int \frac{B \cos(c+dx)}{\cos(c+dx) \sec^{\frac{3}{2}}(c+dx) + \sec^{\frac{3}{2}}(c+dx)} dx}{a}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `(Integral(A/(cos(c + d*x)*sec(c + d*x)**(3/2) + sec(c + d*x)**(3/2)), x) + Integral(B*cos(c + d*x)/(cos(c + d*x)*sec(c + d*x)**(3/2) + sec(c + d*x)**(3/2)), x))/a`

3.480.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.480.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.480.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + a \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))), x)`

$$3.481 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)} dx$$

3.481.1 Optimal result 4390
 3.481.2 Mathematica [C] (verified) 4391
 3.481.3 Rubi [A] (verified) 4391
 3.481.4 Maple [A] (verified) 4395
 3.481.5 Fracas [C] (verification not implemented) 4396
 3.481.6 Sympy [F(-1)] 4396
 3.481.7 Maxima [F] 4397
 3.481.8 Giac [F] 4397
 3.481.9 Mupad [F(-1)] 4397

3.481.1 Optimal result

Integrand size = 33, antiderivative size = 196

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{3(5A - 7B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5ad}$$

$$+ \frac{5(A - B)\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3ad}$$

$$- \frac{(5A - 7B) \sin(c + dx)}{5ad \sec^{\frac{3}{2}}(c + dx)} + \frac{5(A - B) \sin(c + dx)}{3ad \sqrt{\sec(c + dx)}} + \frac{(A - B) \sin(c + dx)}{d \sec^{\frac{3}{2}}(c + dx)(a + a \sec(c + dx))}$$

output

```
-1/5*(5*A-7*B)*sin(d*x+c)/a/d/sec(d*x+c)^(3/2)+(A-B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*sec(d*x+c))+5/3*(A-B)*sin(d*x+c)/a/d/sec(d*x+c)^(1/2)-3/5*(5*A-7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d+5/3*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d
```

3.481.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 3.63 (sec) , antiderivative size = 518, normalized size of antiderivative = 2.64

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{\cos^2\left(\frac{1}{2}(c + dx)\right) \left(60\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1 + e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1 + e^{2i(c+dx)}} + e^{2idx}(-1 + e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{((2I)(c + dx))}\right]\right) + E^{((2I)d*x)}(-1 + E^{((2I)c)}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{((2I)(c + dx))}\right]\right)}{E^{I*d*x} - (84\sqrt{2}B\sqrt{E^{I(c + dx)}}/(1 + E^{((2I)(c + dx))}))\sqrt{1 + E^{((2I)(c + dx))}}\csc[c](-3\sqrt{1 + E^{((2I)(c + dx))}} + E^{((2I)d*x)}(-1 + E^{((2I)c)})\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{((2I)(c + dx))}\right])/E^{I*d*x} + 200A\sqrt{\cos[c + dx]}\operatorname{EllipticF}\left[\frac{c + dx}{2}, 2\right]\sqrt{\sec[c + dx]} - 200B\sqrt{\cos[c + dx]}\operatorname{EllipticF}\left[\frac{c + dx}{2}, 2\right]\sqrt{\sec[c + dx]} + \sqrt{\sec[c + dx]}(3(40A - 51B + (20A - 33B)\cos[2c])\cos[d*x]\csc[c/2]\sec[c/2] + 40(A - B)\cos[2d*x]\sin[2c] + 12B\cos[3d*x]\sin[3c] - 120(A - B)\sec[c/2]\sec[(c + dx)/2]\sin[(d*x)/2] - 12(20A - 33B)\cos[c]\sin[d*x] + 40(A - B)\cos[2c]\sin[2d*x] + 12B\cos[3c]\sin[3d*x] - 120(A - B)\tan[c/2])\right)}/(60a*d*(1 + \cos[c + dx]))$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])*Sec[c + d*x]^(5/2)),x]`

output `(Cos[(c + d*x)/2]^2*((60*sqrt[2]*A*sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) - (84*sqrt[2]*B*sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + 200*A*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] - 200*B*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] + sqrt[Sec[c + d*x]]*(3*(40*A - 51*B + (20*A - 33*B)*Cos[2*c])*Cos[d*x]*Csc[c/2]*Sec[c/2] + 40*(A - B)*Cos[2*d*x]*Sin[2*c] + 12*B*Cos[3*d*x]*Sin[3*c] - 120*(A - B)*Sec[c/2]*Sec[(c + d*x)/2]*Sin[(d*x)/2] - 12*(20*A - 33*B)*Cos[c]*Sin[d*x] + 40*(A - B)*Cos[2*c]*Sin[2*d*x] + 12*B*Cos[3*c]*Sin[3*d*x] - 120*(A - B)*Tan[c/2])))/(60*a*d*(1 + Cos[c + d*x]))`

3.481.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.95, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)} dx$$

3.481. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)} dx \\
& \downarrow 3439 \\
& \int \frac{A \sec(c + dx) + B}{\sec^{5/2}(c + dx) \left(a \sec(c + dx) + a\right)} dx \\
& \downarrow 3042 \\
& \int \frac{A \csc\left(c + dx + \frac{\pi}{2}\right) + B}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + a\right)} dx \\
& \downarrow 4508 \\
& \frac{\int \frac{-a(5A-7B) - 5a(A-B) \sec(c+dx)}{2 \sec^{5/2}(c+dx)} dx}{a^2} + \frac{(A-B) \sin(c+dx)}{d \sec^{3/2}(c+dx) \left(a \sec(c+dx) + a\right)} \\
& \downarrow 27 \\
& \frac{(A-B) \sin(c+dx)}{d \sec^{3/2}(c+dx) \left(a \sec(c+dx) + a\right)} - \frac{\int \frac{a(5A-7B) - 5a(A-B) \sec(c+dx)}{\sec^{5/2}(c+dx)} dx}{2a^2} \\
& \downarrow 3042 \\
& \frac{(A-B) \sin(c+dx)}{d \sec^{3/2}(c+dx) \left(a \sec(c+dx) + a\right)} - \frac{\int \frac{a(5A-7B) - 5a(A-B) \csc\left(c+dx+\frac{\pi}{2}\right)}{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}} dx}{2a^2} \\
& \downarrow 4274 \\
& \frac{(A-B) \sin(c+dx)}{d \sec^{3/2}(c+dx) \left(a \sec(c+dx) + a\right)} - \frac{a(5A-7B) \int \frac{1}{\sec^{5/2}(c+dx)} dx - 5a(A-B) \int \frac{1}{\sec^{3/2}(c+dx)} dx}{2a^2} \\
& \downarrow 3042 \\
& \frac{(A-B) \sin(c+dx)}{d \sec^{3/2}(c+dx) \left(a \sec(c+dx) + a\right)} - \frac{a(5A-7B) \int \frac{1}{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}} dx - 5a(A-B) \int \frac{1}{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}} dx}{2a^2} \\
& \downarrow 4256
\end{aligned}$$

3.481. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{5/2}(c+dx)} dx$

$$\frac{\frac{(A-B)\sin(c+dx)}{d\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)} - a(5A-7B)\left(\frac{3}{5}\int\frac{1}{\sqrt{\sec(c+dx)}}dx + \frac{2\sin(c+dx)}{5d\sec^{\frac{3}{2}}(c+dx)}\right) - 5a(A-B)\left(\frac{1}{3}\int\sqrt{\sec(c+dx)}dx + \frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}}\right)}{2a^2}$$

↓ 3042

$$\frac{\frac{(A-B)\sin(c+dx)}{d\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)} - a(5A-7B)\left(\frac{3}{5}\int\frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}}dx + \frac{2\sin(c+dx)}{5d\sec^{\frac{3}{2}}(c+dx)}\right) - 5a(A-B)\left(\frac{1}{3}\int\sqrt{\csc(c+dx+\frac{\pi}{2})}dx + \frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}}\right)}{2a^2}$$

↓ 4258

$$\frac{\frac{(A-B)\sin(c+dx)}{d\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)} - a(5A-7B)\left(\frac{3}{5}\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\cos(c+dx)}dx + \frac{2\sin(c+dx)}{5d\sec^{\frac{3}{2}}(c+dx)}\right) - 5a(A-B)\left(\frac{1}{3}\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\cos(c+dx)}dx + \frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}}\right)}{2a^2}$$

↓ 3042

$$\frac{\frac{(A-B)\sin(c+dx)}{d\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)} - a(5A-7B)\left(\frac{3}{5}\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\sin(c+dx+\frac{\pi}{2})}dx + \frac{2\sin(c+dx)}{5d\sec^{\frac{3}{2}}(c+dx)}\right) - 5a(A-B)\left(\frac{1}{3}\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\sin(c+dx+\frac{\pi}{2})}dx + \frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}}\right)}{2a^2}$$

↓ 3119

$$\frac{\frac{(A-B)\sin(c+dx)}{d\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)} - a(5A-7B)\left(\frac{2\sin(c+dx)}{5d\sec^{\frac{3}{2}}(c+dx)} + \frac{6\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E(\frac{1}{2}(c+dx)|2)}{5d}\right) - 5a(A-B)\left(\frac{1}{3}\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\sin(c+dx+\frac{\pi}{2})}dx + \frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}}\right)}{2a^2}$$

↓ 3120

$$\frac{\frac{(A-B)\sin(c+dx)}{d\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+a)} - a(5A-7B)\left(\frac{2\sin(c+dx)}{5d\sec^{\frac{3}{2}}(c+dx)} + \frac{6\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E(\frac{1}{2}(c+dx)|2)}{5d}\right) - 5a(A-B)\left(\frac{2\sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}}{3d}\right)}{2a^2}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*cos[c + d*x])*Sec[c + d*x]^(5/2)),x]`

$$3.481. \quad \int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)} dx$$


```
output ((A - B)*Sin[c + d*x])/(d*Sec[c + d*x]^(3/2)*(a + a*Sec[c + d*x])) - (a*(5
*A - 7*B)*((6*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*
x]])/(5*d) + (2*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2))) - 5*a*(A - B)*((2*
sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (
2*Sin[c + d*x])/(3*d*sqrt[Sec[c + d*x]])))/(2*a^2)
```

3.481.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4256 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c
+ d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.481.4 Maple [A] (verified)

Time = 5.90 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.43

method	result
default	$-\frac{\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(25AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + 45AE\right)}{15a \cos\left(\frac{dx}{2} + \frac{c}{2}\right)}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)/sec(d*x+c)^(5/2), x, method=_RETURNVERBOSE)`

output `-1/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(cos(1/2*d*x+1/2*c)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(25*A*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+45*A*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))-25*B*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-63*B*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))+48*B*sin(1/2*d*x+1/2*c)^8+(-40*A-56*B)*sin(1/2*d*x+1/2*c)^6+(90*A-30*B)*sin(1/2*d*x+1/2*c)^4+(-35*A+23*B)*sin(1/2*d*x+1/2*c)^2)/a/cos(1/2*d*x+1/2*c)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.481.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)} dx$$

3.481.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.42

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{25 (\sqrt{2}(i A - i B) \cos(dx + c) + \sqrt{2}(i A - i B)) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(5/2),x, algorithm="fricas")
```

```
output -1/30*(25*(sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 25*(sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 9*(sqrt(2)*(5*I*A - 7*I*B)*cos(d*x + c) + sqrt(2)*(5*I*A - 7*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 9*(sqrt(2)*(-5*I*A + 7*I*B)*cos(d*x + c) + sqrt(2)*(-5*I*A + 7*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(6*B*cos(d*x + c)^3 + 2*(5*A - 2*B)*cos(d*x + c)^2 + 25*(A - B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c))/(a*d*cos(d*x + c) + a*d)
```

3.481.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)**(5/2),x)
```

```
output Timed out
```

3.481.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sec(d*x + c)^(5/2)), x)`

3.481.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))/sec(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)*sec(d*x + c)^(5/2)), x)`

3.481.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (a + a \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))), x)`

3.482
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^2} dx$$

3.482.1 Optimal result 4398
 3.482.2 Mathematica [C] (verified) 4399
 3.482.3 Rubi [A] (verified) 4399
 3.482.4 Maple [B] (verified) 4403
 3.482.5 Fricas [C] (verification not implemented) 4404
 3.482.6 Sympy [F(-1)] 4405
 3.482.7 Maxima [F] 4405
 3.482.8 Giac [F] 4406
 3.482.9 Mupad [F(-1)] 4406

3.482.1 Optimal result

Integrand size = 33, antiderivative size = 208

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx \\ &= -\frac{(4A - B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{a^2 d} \\ & \quad - \frac{(5A - 2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3a^2 d} \\ & \quad + \frac{(4A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{a^2 d} \\ & \quad - \frac{(5A - 2B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3a^2 d(1 + \sec(c + dx))} - \frac{(A - B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{3d(a + a \sec(c + dx))^2} \end{aligned}$$

output

```
-1/3*(5*A-2*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a^2/d/(1+sec(d*x+c))-1/3*(A-B)*
sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*sec(d*x+c))^2+(4*A-B)*sin(d*x+c)*sec(d*
x+c)^(1/2)/a^2/d-(4*A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*E
llipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2
/d-1/3*(5*A-2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF
(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/d
```

3.482.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.80 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.46

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx =$$

$$e^{-idx} \cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(29iA - 5iB + 2i(25A - 7B) \cos(c + dx) + 17iA \cos(2(c + dx)) - \dots\right)$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^2, x]`

output `-1/6*(Cos[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*((29*I)*A - (5*I)*B + (2*I)*(25*A - 7*B)*Cos[c + d*x] + (17*I)*A*Cos[2*(c + d*x)] - (5*I)*B*Cos[2*(c + d*x)]) - (I*(4*A - B)*(1 + E^(I*(c + d*x)))^3*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*(c + d*x)) + 8*(5*A - 2*B)*Cos[(c + d*x)/2]^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*(Cos[(c + d*x)/2] - I*Sin[(c + d*x)/2]) - 12*A*Sin[c + d*x] - 7*A*Sin[2*(c + d*x)] + B*Sin[2*(c + d*x)]*(Cos[(c + 3*d*x)/2] + I*Sin[(c + 3*d*x)/2]))/(a^2*d*E^(I*d*x)*(1 + Cos[c + d*x])^2)`

3.482.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4507, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx$$

↓ 3042

$$\int \frac{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^2} dx$$

3.482. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 3439 \\
& \int \frac{\sec^{\frac{5}{2}}(c+dx)(A \sec(c+dx) + B)}{(a \sec(c+dx) + a)^2} dx \\
& \downarrow 3042 \\
& \int \frac{\csc(c+dx + \frac{\pi}{2})^{5/2} (A \csc(c+dx + \frac{\pi}{2}) + B)}{(a \csc(c+dx + \frac{\pi}{2}) + a)^2} dx \\
& \downarrow 4507 \\
& \frac{\int -\frac{\sec^{\frac{3}{2}}(c+dx)(3a(A-B) - a(7A-B) \sec(c+dx))}{2(\sec(c+dx)a+a)} dx}{3a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 27 \\
& -\frac{\int \frac{\sec^{\frac{3}{2}}(c+dx)(3a(A-B) - a(7A-B) \sec(c+dx))}{\sec(c+dx)a+a} dx}{6a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 3042 \\
& -\frac{\int \frac{\csc(c+dx + \frac{\pi}{2})^{3/2} (3a(A-B) - a(7A-B) \csc(c+dx + \frac{\pi}{2}))}{\csc(c+dx + \frac{\pi}{2})a+a} dx}{6a^2} - \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 4507 \\
& -\frac{\int \frac{\sqrt{\sec(c+dx)}(a^2(5A-2B) - 3a^2(4A-B) \sec(c+dx))}{a^2} dx}{6a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)} - \\
& \quad \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 3042 \\
& -\frac{\int \frac{\sqrt{\csc(c+dx + \frac{\pi}{2})}(a^2(5A-2B) - 3a^2(4A-B) \csc(c+dx + \frac{\pi}{2}))}{a^2} dx}{6a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)} - \\
& \quad \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 4274 \\
& -\frac{a^2(5A-2B) \int \sqrt{\sec(c+dx)} dx - 3a^2(4A-B) \int \sec^{\frac{3}{2}}(c+dx) dx}{a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)} - \\
& \quad \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 3042
\end{aligned}$$

3.482. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^2} dx$

$$\frac{\frac{a^2(5A-2B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - 3a^2(4A-B) \int \csc(c+dx+\frac{\pi}{2})^{3/2} dx}{a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)}}{6a^2} -$$

$$\frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2}$$

↓ 4255

$$\frac{\frac{a^2(5A-2B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - 3a^2(4A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right)}{a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)}}{6a^2} -$$

$$\frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2}$$

↓ 3042

$$\frac{\frac{a^2(5A-2B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - 3a^2(4A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right)}{a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)}}{6a^2} -$$

$$\frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2}$$

↓ 4258

$$\frac{\frac{a^2(5A-2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^2(4A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx \right)}{a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)}}{6a^2} -$$

$$\frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2}$$

↓ 3042

$$\frac{\frac{a^2(5A-2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^2(4A-B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right)}{a^2} + \frac{2(5A-2B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(\sec(c+dx)+1)}}{6a^2} -$$

$$\frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2}$$

↓ 3119

3.482. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^2} dx$

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4507 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

3.482.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. 2(240) = 480.

Time = 5.18 (sec) , antiderivative size = 494, normalized size of antiderivative = 2.38

method	result
default	$-\frac{2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(5AF\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)-12AE\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\right)}{\dots}$

$$3.482. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^2} dx$$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)`

output `-1/6*(2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(5*A*EllipticF(cos(1/2*d*
x+1/2*c),2^(1/2))-12*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-2*B*EllipticF
(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*co
s(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*s
in(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
^(1/2)*(5*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-12*A*EllipticE(cos(1/2*d
*x+1/2*c),2^(1/2))-2*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*EllipticE
(cos(1/2*d*x+1/2*c),2^(1/2)))*cos(1/2*d*x+1/2*c)-12*(-2*sin(1/2*d*x+1/2*c)
^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*A-B)*sin(1/2*d*x+1/2*c)^6+2*(-2*sin(1/2*
d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(43*A-10*B)*sin(1/2*d*x+1/2*c)^4-
(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(37*A-7*B)*sin(1/2*d*
x+1/2*c)^2)/a^2/cos(1/2*d*x+1/2*c)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.482.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.76

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{(\sqrt{2}(5iA - 2iB) \cos(dx + c))^2 - 2\sqrt{2}(-5iA + 2iB) \cos(dx + c) + \sqrt{2}(5iA - 2iB)}{\text{weierstrassPInvers}}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2,x, algorith
m="fracas")`

3.482. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^2} dx$

output `1/6*((sqrt(2)*(5*I*A - 2*I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(-5*I*A + 2*I*B)*cos(d*x + c) + sqrt(2)*(5*I*A - 2*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(-5*I*A + 2*I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(5*I*A - 2*I*B)*cos(d*x + c) + sqrt(2)*(-5*I*A + 2*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*(sqrt(2)*(4*I*A - I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(4*I*A - I*B)*cos(d*x + c) + sqrt(2)*(4*I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*(sqrt(2)*(-4*I*A + I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(-4*I*A + I*B)*cos(d*x + c) + sqrt(2)*(-4*I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*(4*A - B)*cos(d*x + c)^2 + (19*A - 4*B)*cos(d*x + c) + 6*A)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.482.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**2,x)`

output `Timed out`

3.482.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2,x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^2, x)`

3.482.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^2, x)`

3.482.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{(a + a \cos(c + dx))^2} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^2,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^2, x)`

3.483
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^2} dx$$

3.483.1 Optimal result 4407
 3.483.2 Mathematica [C] (verified) 4408
 3.483.3 Rubi [A] (verified) 4408
 3.483.4 Maple [A] (verified) 4412
 3.483.5 Fracas [C] (verification not implemented) 4413
 3.483.6 Sympy [F] 4413
 3.483.7 Maxima [F] 4414
 3.483.8 Giac [F] 4414
 3.483.9 Mupad [F(-1)] 4414

3.483.1 Optimal result

Integrand size = 33, antiderivative size = 161

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{A\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{a^2d}$$

$$+ \frac{(2A + B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3a^2d}$$

$$- \frac{A\sqrt{\sec(c + dx)}\sin(c + dx)}{a^2d(1 + \sec(c + dx))} - \frac{(A - B)\sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{3d(a + a \sec(c + dx))^2}$$

output

```
-1/3*(A-B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*sec(d*x+c))^2-A*sin(d*x+c)*s
ec(d*x+c)^(1/2)/a^2/d/(1+sec(d*x+c))+A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/
2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*
x+c)^(1/2)/a^2/d+1/3*(2*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*
c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)
/a^2/d
```

3.483.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.16 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.59

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{e^{-idx} \cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(-iAe^{-i(c+dx)}(1 + e^{i(c+dx)})^3 \sqrt{1 + e^{2i(c+dx)}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{4}, -E^{i(c+dx)}\right)\right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^2, x]`

output `(Cos[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(((-I)*A*(1 + E^(I*(c + d*x))))^3*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*(c + d*x)) + 8*(2*A + B)*Cos[(c + d*x)/2]^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*(Cos[(c + d*x)/2] - I*Sin[(c + d*x)/2]) + (2*I)*Cos[c + d*x]*(7*A - B + (5*A + B)*Cos[c + d*x] + I*(A - B)*Sin[c + d*x])*(Cos[(c + 3*d*x)/2] + I*Sin[(c + 3*d*x)/2]))/(6*a^2*d*E^(I*d*x)*(1 + Cos[c + d*x])^2)`

3.483.3 Rubi [A] (verified)

Time = 1.08 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4507, 25, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^2} dx$$

↓ 3042

$$\int \frac{\sqrt{\csc(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{(a \sin(c + dx + \frac{\pi}{2}) + a)^2} dx$$

↓ 3439

3.483. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{\sec^{\frac{3}{2}}(c+dx)(A \sec(c+dx) + B)}{(a \sec(c+dx) + a)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc(c+dx + \frac{\pi}{2})^{3/2} (A \csc(c+dx + \frac{\pi}{2}) + B)}{(a \csc(c+dx + \frac{\pi}{2}) + a)^2} dx \\
& \quad \downarrow \text{4507} \\
& \frac{\int -\frac{\sqrt{\sec(c+dx)}(a(A-B)-a(5A+B)\sec(c+dx))}{2(\sec(c+dx)a+a)} dx}{3a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{27} \\
& -\frac{\int \frac{\sqrt{\sec(c+dx)}(a(A-B)-a(5A+B)\sec(c+dx))}{\sec(c+dx)a+a} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& -\frac{\int \frac{\sqrt{\csc(c+dx + \frac{\pi}{2})}(a(A-B)-a(5A+B)\csc(c+dx + \frac{\pi}{2}))}{\csc(c+dx + \frac{\pi}{2})a+a} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{4507} \\
& -\frac{\int \frac{3Aa^2+(2A+B)\sec(c+dx)a^2}{\sqrt{\sec(c+dx)}} dx}{6a^2} + \frac{6A \sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{25} \\
& -\frac{\frac{6A \sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \int \frac{3Aa^2+(2A+B)\sec(c+dx)a^2}{\sqrt{\sec(c+dx)}} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& -\frac{\frac{6A \sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \int \frac{3Aa^2+(2A+B)\csc(c+dx + \frac{\pi}{2})a^2}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{4274} \\
& -\frac{\frac{6A \sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{a^2(2A+B) \int \sqrt{\sec(c+dx)} dx + 3a^2 A \int \frac{1}{\sqrt{\sec(c+dx)}} dx}{a^2}}{6a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2}
\end{aligned}$$

3.483. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^2} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{\frac{6A \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{a^2(2A+B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + 3a^2 A \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{a^2}}{6a^2} \\ & \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 4258 \\ & \frac{\frac{6A \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{a^2(2A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^2 A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{a^2}}{6a^2} \\ & \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{\frac{6A \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{a^2(2A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3a^2 A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{a^2}}{6a^2} \\ & \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 3119 \\ & \frac{\frac{6A \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{a^2(2A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6a^2 A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d}}{a^2}}{6a^2} \\ & \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 3120 \\ & \frac{\frac{6A \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{2a^2(2A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} + \frac{6a^2 A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d}}{a^2}}{6a^2} \\ & \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx) + a)^2} \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^2,x]`

```
output -1/3*((A - B)*Sec[c + d*x]^(3/2)*Sin[c + d*x]/(d*(a + a*Sec[c + d*x])^2)
- (((6*a^2*A*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d
*x]])/d + (2*a^2*(2*A + B)*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d)/a^2) + (6*A*sqrt[Sec[c + d*x]]*Sin[c + d*x]/(d*(1 + Sec[c + d*x]))) / (6*a^2)
```

3.483.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_.), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4507 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

3.483.4 Maple [A] (verified)

Time = 4.71 (sec) , antiderivative size = 350, normalized size of antiderivative = 2.17

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(12A\left(\cos^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 4A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right)$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^2,x,method=_RETURNV
ERBOSE)`

output
$$\frac{1}{6} \left((2 \cos(1/2 d x + 1/2 c))^{-2} - 1 \right) \sin(1/2 d x + 1/2 c)^2)^{1/2} \left(12 A \cos(1/2 d x + 1/2 c)^6 - 4 A \left(\sin(1/2 d x + 1/2 c)^2 \right)^{1/2} \left(-2 \cos(1/2 d x + 1/2 c)^{2+1} \right)^{1/2} \operatorname{EllipticF}\left(\cos(1/2 d x + 1/2 c), 2^{1/2}\right) \cos(1/2 d x + 1/2 c)^3 + 6 A \cos(1/2 d x + 1/2 c)^3 \left(\sin(1/2 d x + 1/2 c)^2 \right)^{1/2} \left(-2 \cos(1/2 d x + 1/2 c)^{2+1} \right)^{1/2} \operatorname{EllipticE}\left(\cos(1/2 d x + 1/2 c), 2^{1/2}\right) - 2 B \left(\sin(1/2 d x + 1/2 c)^2 \right)^{1/2} \left(-2 \cos(1/2 d x + 1/2 c)^{2+1} \right)^{1/2} \operatorname{EllipticF}\left(\cos(1/2 d x + 1/2 c), 2^{1/2}\right) \cos(1/2 d x + 1/2 c)^3 - 16 A \cos(1/2 d x + 1/2 c)^4 - 2 B \cos(1/2 d x + 1/2 c)^4 + 3 A \cos(1/2 d x + 1/2 c)^2 + 3 B \cos(1/2 d x + 1/2 c)^2 + A - B \right) / a^2 \cos(1/2 d x + 1/2 c)^3 / \left(-2 \sin(1/2 d x + 1/2 c)^4 + \sin(1/2 d x + 1/2 c)^2 \right)^{1/2} / \sin(1/2 d x + 1/2 c) / \left(2 \cos(1/2 d x + 1/2 c)^2 - 1 \right)^{1/2} / d$$

3.483.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.02

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{(\sqrt{2}(-2iA - iB) \cos(dx + c))^2 - 2\sqrt{2}(2iA + iB) \cos(dx + c) + \sqrt{2}(-2iA - iB)) \text{weierstrassPInverse}(\dots)}{\dots}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^2,x, algorithm="fricas")`

output `1/6*((sqrt(2)*(-2*I*A - I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(2*I*A + I*B)*cos(d*x + c) + sqrt(2)*(-2*I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + (sqrt(2)*(2*I*A + I*B)*cos(d*x + c)^2 - 2*sqrt(2)*(-2*I*A - I*B)*cos(d*x + c) + sqrt(2)*(2*I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*(-I*sqrt(2)*A*cos(d*x + c)^2 - 2*I*sqrt(2)*A*cos(d*x + c) - I*sqrt(2)*A)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*(I*sqrt(2)*A*cos(d*x + c)^2 + 2*I*sqrt(2)*A*cos(d*x + c) + I*sqrt(2)*A)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*A*cos(d*x + c)^2 + (4*A - B)*cos(d*x + c)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.483.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx$$

$$= \frac{\int \frac{A \sqrt{\sec(c+dx)}}{\cos^2(c+dx)+2\cos(c+dx)+1} dx + \int \frac{B \cos(c+dx) \sqrt{\sec(c+dx)}}{\cos^2(c+dx)+2\cos(c+dx)+1} dx}{a^2}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**2,x)`

output `(Integral(A*sqrt(sec(c + d*x))/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sqrt(sec(c + d*x))/(cos(c + d*x)**2 + 2*cos(c + d*x) + 1), x))/a**2`

3.483. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^2} dx$

3.483.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^2, x)`

3.483.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^2, x)`

3.483.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + a \cos(c + dx))^2} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^2,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^2, x)`

3.484
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$$

3.484.1 Optimal result 4415
 3.484.2 Mathematica [C] (verified) 4416
 3.484.3 Rubi [A] (verified) 4416
 3.484.4 Maple [A] (verified) 4420
 3.484.5 Fracas [C] (verification not implemented) 4421
 3.484.6 Sympy [F] 4422
 3.484.7 Maxima [F] 4422
 3.484.8 Giac [F] 4422
 3.484.9 Mupad [F(-1)] 4423

3.484.1 Optimal result

Integrand size = 33, antiderivative size = 168

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

$$= -\frac{B \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{a^2 d}$$

$$+ \frac{(A + 2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3a^2 d}$$

$$+ \frac{(A + 2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3a^2 d(1 + \sec(c + dx))} - \frac{(A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3d(a + a \sec(c + dx))^2}$$

```
output 1/3*(A+2*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d/(1+sec(d*x+c))-1/3*(A-B)*sin
(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*sec(d*x+c))^2-B*(cos(1/2*d*x+1/2*c)^2)^(1/
2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/
2)*sec(d*x+c)^(1/2)/a^2/d+1/3*(A+2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2
*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x
+c)^(1/2)/a^2/d
```

3.484.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 2.33 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.52

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{e^{-idx} \cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(8(A + 2B) \cos^3\left(\frac{1}{2}(c + dx)\right) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^2*Sqrt[Sec[c + d*x]]),x]`

output `(Cos[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(8*(A + 2*B)*Cos[(c + d*x)/2]^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*(Cos[(c + d*x)/2] - I*Sin[(c + d*x)/2])) + I*((B*(1 + E^(I*(c + d*x)))^3*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*(c + d*x)) + 2*Cos[c + d*x]*(-A - 5*B + (A - 7*B)*Cos[c + d*x] - I*(A - B)*Sin[c + d*x]))*(Cos[(c + 3*d*x)/2] + I*Sin[(c + 3*d*x)/2]))/(6*a^2*d*E^(I*d*x)*(1 + Cos[c + d*x])^2)`

3.484.3 Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4508, 25, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a \cos(c + dx) + a)^2} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2} dx$$

↓ 3439

3.484. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{\sec(c+dx)}(A \sec(c+dx) + B)}{(a \sec(c+dx) + a)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A \csc(c+dx+\frac{\pi}{2}) + B)}{(a \csc(c+dx+\frac{\pi}{2}) + a)^2} dx \\
& \quad \downarrow \text{4507} \\
& \frac{\int \frac{a(A-B)+3a(A+B)\sec(c+dx)}{2\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{3a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(A-B)+3a(A+B)\sec(c+dx)}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(A-B)+3a(A+B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(\csc(c+dx+\frac{\pi}{2})a+a)} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{4508} \\
& \frac{\int -\frac{3a^2B-a^2(A+2B)\sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{a^2} + \frac{2(A+2B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{25} \\
& \frac{2(A+2B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{\int \frac{3a^2B-a^2(A+2B)\sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{2(A+2B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{\int \frac{3a^2B-a^2(A+2B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \quad \downarrow \text{4274} \\
& \frac{2(A+2B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{3a^2B \int \frac{1}{\sqrt{\sec(c+dx)}} dx - a^2(A+2B) \int \sqrt{\sec(c+dx)} dx}{a^2} - \\
& \quad \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2}
\end{aligned}$$

3.484. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{\frac{2(A+2B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{3a^2 B \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - a^2(A+2B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{a^2}}{6a^2} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 4258 \\
& \frac{\frac{2(A+2B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{3a^2 B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - a^2(A+2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a^2}}{6a^2} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 3042 \\
& \frac{\frac{2(A+2B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{3a^2 B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - a^2(A+2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2}}{6a^2} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 3119 \\
& \frac{\frac{2(A+2B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{6a^2 B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)|2\right) - a^2(A+2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2}}{6a^2} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} \\
& \downarrow 3120 \\
& \frac{\frac{2(A+2B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} - \frac{6a^2 B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)|2\right) - 2a^2(A+2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{a^2}}{6a^2} \\
& \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2}
\end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^2*sqrt[Sec[c + d*x]]),x]`

3.484. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

```
output -1/3*((A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x])^2)
+ (-(((6*a^2*B*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d
*x]])/d - (2*a^2*(A + 2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sq
rt[Sec[c + d*x]])/d)/a^2) + (2*(A + 2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/
(d*(1 + Sec[c + d*x]))/(6*a^2)
```

3.484.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_.), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4507 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.484.4 Maple [A] (verified)

Time = 5.22 (sec) , antiderivative size = 350, normalized size of antiderivative = 2.08

method	result
default	$-\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\left(2A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 12\right)}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)`

output
$$-1/6*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*A*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))*\cos(1/2*d*x+1/2*c)^3+12*B*\cos(1/2*d*x+1/2*c)^6+4*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^{(1/2)}))*\cos(1/2*d*x+1/2*c)^3+6*B*\cos(1/2*d*x+1/2*c)^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})+2*A*\cos(1/2*d*x+1/2*c)^4-20*B*\cos(1/2*d*x+1/2*c)^4-3*A*\cos(1/2*d*x+1/2*c)^2+9*B*\cos(1/2*d*x+1/2*c)^2+A-B)/a^2/\cos(1/2*d*x+1/2*c)^3/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$$

3.484.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 324, normalized size of antiderivative = 1.93

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{(\sqrt{2}(-iA - 2iB) \cos(dx + c)^2 - 2\sqrt{2}(iA + 2iB) \cos(dx + c) + \sqrt{2}(-iA - 2iB)) \text{weierstrassPInverse}(\dots)}{\dots}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output
$$1/6*((\text{sqrt}(2)*(-I*A - 2*I*B)*\cos(d*x + c)^2 - 2*\text{sqrt}(2)*(I*A + 2*I*B)*\cos(d*x + c) + \text{sqrt}(2)*(-I*A - 2*I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + (\text{sqrt}(2)*(I*A + 2*I*B)*\cos(d*x + c)^2 - 2*\text{sqrt}(2)*(-I*A - 2*I*B)*\cos(d*x + c) + \text{sqrt}(2)*(I*A + 2*I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*(I*\text{sqrt}(2)*B*\cos(d*x + c)^2 + 2*I*\text{sqrt}(2)*B*\cos(d*x + c) + I*\text{sqrt}(2)*B)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - 3*(-I*\text{sqrt}(2)*B*\cos(d*x + c)^2 - 2*I*\text{sqrt}(2)*B*\cos(d*x + c) - I*\text{sqrt}(2)*B)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*(3*B*\cos(d*x + c)^2 + (A + 2*B)*\cos(d*x + c))*\sin(d*x + c)/\text{sqrt}(\cos(d*x + c)))/(a^2*d*\cos(d*x + c)^2 + 2*a^2*d*\cos(d*x + c) + a^2*d)$$

3.484.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{\int \frac{A}{\cos^2(c+dx)\sqrt{\sec(c+dx)} + 2\cos(c+dx)\sqrt{\sec(c+dx)} + \sqrt{\sec(c+dx)}} dx + \int \frac{B \cos(c+dx)}{\cos^2(c+dx)\sqrt{\sec(c+dx)} + 2\cos(c+dx)\sqrt{\sec(c+dx)} + \sqrt{\sec(c+dx)}} dx}{a^2}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2/sec(d*x+c)**(1/2),x)`

output `(Integral(A/(cos(c + d*x)**2*sqrt(sec(c + d*x)) + 2*cos(c + d*x)*sqrt(sec(c + d*x)) + sqrt(sec(c + d*x))), x) + Integral(B*cos(c + d*x)/(cos(c + d*x)**2*sqrt(sec(c + d*x)) + 2*cos(c + d*x)*sqrt(sec(c + d*x)) + sqrt(sec(c + d*x))), x))/a**2`

3.484.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sqrt(sec(d*x + c))), x)`

3.484.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sqrt(sec(d*x + c))), x)`

3.484. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

3.484.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + a \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^2),x)`

3.485 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{3}{2}}(c+dx)} dx$

3.485.1 Optimal result 4424
 3.485.2 Mathematica [C] (verified) 4425
 3.485.3 Rubi [A] (verified) 4425
 3.485.4 Maple [A] (verified) 4429
 3.485.5 Fracas [C] (verification not implemented) 4429
 3.485.6 Sympy [F(-1)] 4430
 3.485.7 Maxima [F] 4430
 3.485.8 Giac [F] 4431
 3.485.9 Mupad [F(-1)] 4431

3.485.1 Optimal result

Integrand size = 33, antiderivative size = 176

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{(A - 4B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{a^2 d}$$

$$+ \frac{(2A - 5B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3a^2 d}$$

$$+ \frac{(2A - 5B)\sqrt{\sec(c + dx)}\sin(c + dx)}{3a^2 d(1 + \sec(c + dx))} + \frac{(A - B)\sqrt{\sec(c + dx)}\sin(c + dx)}{3d(a + a \sec(c + dx))^2}$$

output

```
1/3*(2*A-5*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d/(1+sec(d*x+c))+1/3*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*sec(d*x+c))^2-(A-4*B)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/d+1/3*(2*A-5*B)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/d
```

3.485.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.48 (sec) , antiderivative size = 475, normalized size of antiderivative = 2.70

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \cos^4\left(\frac{1}{2}(c + dx)\right) \left(2\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1 + e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1 + e^{2i(c+dx)}} + e^{2idx}(-1 + e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -e^{2i(c+dx)}\right]\right) + E^{i(c+dx)} \sqrt{1 + e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1 + e^{2i(c+dx)}} + e^{2idx}(-1 + e^{2ic}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -e^{2i(c+dx)}\right]\right)\right)$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^2*Sec[c + d*x]^(3/2)),x]`

output `(Cos[(c + d*x)/2]^4*((2*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) - (8*Sqrt[2]*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + ((5*(A - 4*B)*Cos[(c - d*x)/2] + 4*(A - 4*B)*Cos[(3*c + d*x)/2] + 3*A*Cos[(c + 3*d*x)/2] - 9*B*Cos[(c + 3*d*x)/2] - 3*B*Cos[(5*c + 3*d*x)/2])*Csc[c/2]*Sec[c/2]*Sec[(c + d*x)/2]^3)/(2*Sqrt[Sec[c + d*x]]) + 8*A*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]] - 20*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*a^2*d*(1 + Cos[c + d*x])^2)`

3.485.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4508, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^2} dx$$

↓ 3042

3.485. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{3d(a \sec(c + dx) + a)^2} - \\
 & \frac{3a^2(A-4B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - a^2(2A-5B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{a^2} - \frac{2(2A-5B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} \\
 & \frac{6a^2}{6a^2} \\
 & \downarrow 4258 \\
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{3d(a \sec(c + dx) + a)^2} - \\
 & \frac{3a^2(A-4B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - a^2(2A-5B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a^2} - \frac{2(2A-5B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} \\
 & \frac{6a^2}{6a^2} \\
 & \downarrow 3042 \\
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{3d(a \sec(c + dx) + a)^2} - \\
 & \frac{3a^2(A-4B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - a^2(2A-5B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2} - \frac{2(2A-5B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} \\
 & \frac{6a^2}{6a^2} \\
 & \downarrow 3119 \\
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{3d(a \sec(c + dx) + a)^2} - \\
 & \frac{6a^2(A-4B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) - a^2(2A-5B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a^2} - \frac{2(2A-5B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} \\
 & \frac{6a^2}{6a^2} \\
 & \downarrow 3120 \\
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{3d(a \sec(c + dx) + a)^2} - \\
 & \frac{6a^2(A-4B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) - 2a^2(2A-5B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{a^2} - \frac{2(2A-5B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(\sec(c+dx)+1)} \\
 & \frac{6a^2}{6a^2}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^2*Sec[c + d*x]^(3/2)),x]`

output `((A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Sec[c + d*x])^2) - ((6*a^2*(A - 4*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*a^2*(2*A - 5*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d/a^2 - (2*(2*A - 5*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(1 + Sec[c + d*x]))/(6*a^2)`

3.485. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{3}{2}}(c+dx)} dx$

3.485.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_.), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^(n_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4508 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^(n_.)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_.)*(csc[(e_.) + (f_.)*(x_)])*(B_.) + (A_.), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.485.4 Maple [A] (verified)

Time = 5.67 (sec) , antiderivative size = 421, normalized size of antiderivative = 2.39

method	result
default	$-\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(12A\left(\cos^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+4A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\right)$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2/sec(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/6*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*A*cos(1/2
*d*x+1/2*c)^6+4*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)
^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3+6*A*cos(
1/2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-24*B*cos(1/2*d*x+1/2*c)^6-10*B
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(
cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^3-24*B*cos(1/2*d*x+1/2*c)^3
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(
cos(1/2*d*x+1/2*c),2^(1/2))-20*A*cos(1/2*d*x+1/2*c)^4+38*B*cos(1/2*d*x+1/2
*c)^4+9*A*cos(1/2*d*x+1/2*c)^2-15*B*cos(1/2*d*x+1/2*c)^2-A/B)/a^2/cos(1/2*
d*x+1/2*c)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*
d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.485.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 362, normalized size of antiderivative = 2.06

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(\sqrt{2}(-2iA + 5iB) \cos(dx + c))^2 - 2\sqrt{2}(2iA - 5iB) \cos(dx + c) + \sqrt{2}(-2iA + 5iB)) \text{weierstrassPInve}}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(3/2),x, algorithm
m="fricas")
```

3.485. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{3}{2}}(c+dx)} dx$

output $1/6*((\sqrt{2})*(-2*I*A + 5*I*B)*\cos(d*x + c)^2 - 2*\sqrt{2}*(2*I*A - 5*I*B)*\cos(d*x + c) + \sqrt{2})*(-2*I*A + 5*I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + (\sqrt{2}*(2*I*A - 5*I*B)*\cos(d*x + c)^2 - 2*\sqrt{2}*(-2*I*A + 5*I*B)*\cos(d*x + c) + \sqrt{2}*(2*I*A - 5*I*B))*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)) - 3*(\sqrt{2}*(I*A - 4*I*B)*\cos(d*x + c)^2 + 2*\sqrt{2}*(I*A - 4*I*B)*\cos(d*x + c) + \sqrt{2}*(I*A - 4*I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - 3*(\sqrt{2}*(-I*A + 4*I*B)*\cos(d*x + c)^2 + 2*\sqrt{2}*(-I*A + 4*I*B)*\cos(d*x + c) + \sqrt{2}*(-I*A + 4*I*B))*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*(3*(A - 2*B)*\cos(d*x + c)^2 + (2*A - 5*B)*\cos(d*x + c))*\sin(d*x + c)/\sqrt{\cos(d*x + c)}/(a^2*d*\cos(d*x + c)^2 + 2*a^2*d*\cos(d*x + c) + a^2*d)$

3.485.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2/sec(d*x+c)**(3/2),x)`

output Timed out

3.485.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2)), x)`

3.485.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2)), x)`

3.485.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + a \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^2),x)`

3.486
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$$

3.486.1 Optimal result 4432
 3.486.2 Mathematica [C] (verified) 4433
 3.486.3 Rubi [A] (verified) 4433
 3.486.4 Maple [A] (verified) 4438
 3.486.5 Fracas [C] (verification not implemented) 4438
 3.486.6 Sympy [F(-1)] 4439
 3.486.7 Maxima [F] 4439
 3.486.8 Giac [F] 4440
 3.486.9 Mupad [F(-1)] 4440

3.486.1 Optimal result

Integrand size = 33, antiderivative size = 206

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx \\ &= \frac{(4A - 7B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{a^2 d} \\ & \quad - \frac{5(A - 2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3a^2 d} \\ & \quad - \frac{5(A - 2B) \sin(c + dx)}{3a^2 d \sqrt{\sec(c + dx)}} + \frac{(4A - 7B) \sin(c + dx)}{3a^2 d \sqrt{\sec(c + dx)}(1 + \sec(c + dx))} \\ & \quad + \frac{(A - B) \sin(c + dx)}{3d \sqrt{\sec(c + dx)}(a + a \sec(c + dx))^2} \end{aligned}$$

output

```
-5/3*(A-2*B)*sin(d*x+c)/a^2/d/sec(d*x+c)^(1/2)+1/3*(4*A-7*B)*sin(d*x+c)/a^
2/d/(1+sec(d*x+c))/sec(d*x+c)^(1/2)+1/3*(A-B)*sin(d*x+c)/d/(a+a*sec(d*x+c)
)^2/sec(d*x+c)^(1/2)+(4*A-7*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/
2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/
2)/a^2/d-5/3*(A-2*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellip
ticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/d
```

3.486.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 6.76 (sec) , antiderivative size = 525, normalized size of antiderivative = 2.55

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{\cos^4\left(\frac{1}{2}(c + dx)\right) \left(-8\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{((2I)(c+dx))}\right] + E^{((2I)d*x)}(-1+E^{((2I)c)}) \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{((2I)(c+dx))}\right]\right)}{E^{(I*d*x)} + (14*\sqrt{2}*B*\sqrt{E^{(I*(c+dx))}}/(1+E^{((2I)*(c+dx))}))*\sqrt{1+E^{((2I)*(c+dx))}}*\csc[c]*(-3*\sqrt{1+E^{((2I)*(c+dx))}} + E^{((2I)d*x)}(-1+E^{((2I)c)})*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -E^{((2I)(c+dx))}\right])/E^{(I*d*x)} - ((8*(5*A - 9*B)*\cos[(c-d*x)/2] + (32*A - 54*B)*\cos[(3*c+d*x)/2] + 18*A*\cos[(c+3*d*x)/2] - 3*3*B*\cos[(c+3*d*x)/2] + 6*A*\cos[(5*c+3*d*x)/2] - 9*B*\cos[(5*c+3*d*x)/2] - B*\cos[(3*c+5*d*x)/2] + B*\cos[(7*c+5*d*x)/2])*Csc[c/2]*Sec[c/2]*Sec[(c+dx)/2]^3)/(4*\sqrt{Sec[c+dx]}) - 20*A*\sqrt{Cos[c+dx]}*EllipticF[(c+dx)/2, 2]*\sqrt{Sec[c+dx]} + 40*B*\sqrt{Cos[c+dx]}*EllipticF[(c+dx)/2, 2]*\sqrt{Sec[c+dx]})/(3*a^2*d*(1+Cos[c+dx])^2)}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^2*Sec[c + d*x]^(5/2)),x]`

output `(Cos[(c + d*x)/2]^4*((-8*sqrt[2]*A*sqrt[E^(I*(c + d*x))]/(1 + E^((2*I)*(c + d*x))))*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + (14*sqrt[2]*B*sqrt[E^(I*(c + d*x))]/(1 + E^((2*I)*(c + d*x)))*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))]) + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) - ((8*(5*A - 9*B)*Cos[(c - d*x)/2] + (32*A - 54*B)*Cos[(3*c + d*x)/2] + 18*A*Cos[(c + 3*d*x)/2] - 3*3*B*Cos[(c + 3*d*x)/2] + 6*A*Cos[(5*c + 3*d*x)/2] - 9*B*Cos[(5*c + 3*d*x)/2] - B*Cos[(3*c + 5*d*x)/2] + B*Cos[(7*c + 5*d*x)/2])*Csc[c/2]*Sec[c/2]*Sec[(c + d*x)/2]^3)/(4*sqrt[Sec[c + d*x]]) - 20*A*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] + 40*B*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*a^2*d*(1 + Cos[c + d*x])^2)`

3.486.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.02, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4508, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^2} dx$$

3.486. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^2} dx && \downarrow \text{3042} \\
& \int \frac{A \sec(c + dx) + B}{\sec^{3/2}(c + dx) (a \sec(c + dx) + a)^2} dx && \downarrow \text{3439} \\
& \int \frac{A \csc\left(c + dx + \frac{\pi}{2}\right) + B}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + a\right)^2} dx && \downarrow \text{3042} \\
& \frac{\int \frac{3a(A-3B) - 5a(A-B) \sec(c+dx)}{2 \sec^{3/2}(c+dx)(\sec(c+dx)a+a)} dx}{3a^2} + \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} && \downarrow \text{4508} \\
& \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} - \frac{\int \frac{3a(A-3B) - 5a(A-B) \sec(c+dx)}{\sec^{3/2}(c+dx)(\sec(c+dx)a+a)} dx}{6a^2} && \downarrow \text{27} \\
& \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} - \frac{\int \frac{3a(A-3B) - 5a(A-B) \csc\left(c+dx+\frac{\pi}{2}\right)}{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(\csc\left(c+dx+\frac{\pi}{2}\right)a+a\right)} dx}{6a^2} && \downarrow \text{3042} \\
& \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} - \frac{\int \frac{3(5a^2(A-2B) - a^2(4A-7B) \sec(c+dx))}{\sec^{3/2}(c+dx)} dx}{6a^2} - \frac{2(4A-7B) \sin(c+dx)}{d \sqrt{\sec(c+dx)} (\sec(c+dx)+1)} && \downarrow \text{4508} \\
& \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} - \frac{3 \int \frac{5a^2(A-2B) - a^2(4A-7B) \sec(c+dx)}{\sec^{3/2}(c+dx)} dx}{6a^2} - \frac{2(4A-7B) \sin(c+dx)}{d \sqrt{\sec(c+dx)} (\sec(c+dx)+1)} && \downarrow \text{27} \\
& \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} - \frac{3 \int \frac{5a^2(A-2B) - a^2(4A-7B) \csc\left(c+dx+\frac{\pi}{2}\right)}{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}} dx}{6a^2} - \frac{2(4A-7B) \sin(c+dx)}{d \sqrt{\sec(c+dx)} (\sec(c+dx)+1)} && \downarrow \text{3042} \\
& \frac{(A-B) \sin(c+dx)}{3d \sqrt{\sec(c+dx)} (a \sec(c+dx) + a)^2} - \frac{3 \int \frac{5a^2(A-2B) - a^2(4A-7B) \csc\left(c+dx+\frac{\pi}{2}\right)}{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}} dx}{6a^2} - \frac{2(4A-7B) \sin(c+dx)}{d \sqrt{\sec(c+dx)} (\sec(c+dx)+1)} && \downarrow \text{3042}
\end{aligned}$$

3.486. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{5/2}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 4274 \\
 & \frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - \\
 & \frac{3 \left(5a^2(A-2B) \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)} dx - a^2(4A-7B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right)}{a^2} - \frac{2(4A-7B) \sin(c+dx)}{d\sqrt{\sec(c+dx)}(\sec(c+dx)+1)} \\
 & \frac{6a^2}{} \\
 & \downarrow 3042 \\
 & \frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - \\
 & \frac{3 \left(5a^2(A-2B) \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx - a^2(4A-7B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right)}{a^2} - \frac{2(4A-7B) \sin(c+dx)}{d\sqrt{\sec(c+dx)}(\sec(c+dx)+1)} \\
 & \frac{6a^2}{} \\
 & \downarrow 4256 \\
 & \frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - \\
 & \frac{3 \left(5a^2(A-2B) \left(\frac{1}{3} \int \sqrt{\sec(c+dx)} dx + \frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \right) - a^2(4A-7B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right)}{a^2} - \frac{2(4A-7B) \sin(c+dx)}{d\sqrt{\sec(c+dx)}(\sec(c+dx)+1)} \\
 & \frac{6a^2}{} \\
 & \downarrow 3042 \\
 & \frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - \\
 & \frac{3 \left(5a^2(A-2B) \left(\frac{1}{3} \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + \frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \right) - a^2(4A-7B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right)}{a^2} - \frac{2(4A-7B) \sin(c+dx)}{d\sqrt{\sec(c+dx)}(\sec(c+dx)+1)} \\
 & \frac{6a^2}{} \\
 & \downarrow 4258 \\
 & \frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - \\
 & \frac{3 \left(5a^2(A-2B) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \right) - a^2(4A-7B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx \right)}{a^2} - \frac{2(4A-7B) \sin(c+dx)}{d\sqrt{\sec(c+dx)}(\sec(c+dx)+1)} \\
 & \frac{6a^2}{} \\
 & \downarrow 3042
 \end{aligned}$$

3.486. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\frac{\frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - 3 \left(5a^2(A - 2B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - a^2(4A - 7B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right)}{a^2}}{6a^2}$$

↓ 3119

$$\frac{\frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - 3 \left(5a^2(A - 2B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - \frac{2a^2(4A - 7B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) | 2\right)}{d} \right)}{a^2}}{6a^2} - \frac{2(4A - 7B)}{d\sqrt{\sec(c + dx)}}$$

↓ 3120

$$\frac{\frac{(A - B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} - 3 \left(5a^2(A - 2B) \left(\frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} + \frac{2\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} \right) - \frac{2a^2(4A - 7B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) | 2\right)}{d} \right)}{a^2}}{6a^2} - \frac{2(4A - 7B)}{d\sqrt{\sec(c + dx)}}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^2*Sec[c + d*x]^(5/2)),x]`

output `((A - B)*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]*(a + a*Sec[c + d*x])^2) - ((-2*(4*A - 7*B)*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]]*(1 + Sec[c + d*x])) + (3*((-2*a^2*(4*A - 7*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 5*a^2*(A - 2*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]))))/a^2)/(6*a^2)`

3.486.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.486. \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx$$

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(g_.))^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)])^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4256 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n+1)}/(b*d^n)), x] + \text{Simp}[(n+1)/(b^2*n) \text{Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4508 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-(A*b - a*B))*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^m*((d*\text{Csc}[e + f*x])^n/(b*f*(2*m + 1))), x] - \text{Simp}[1/(a^2*(2*m + 1)) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m+1)}*(d*\text{Csc}[e + f*x])^n*\text{Simp}[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*\text{Csc}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{EqQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -2^{(-1)}] \&\& !\text{GtQ}[n, 0]$

3.486.4 Maple [A] (verified)

Time = 6.12 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.11

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(-16B\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+24A\left(\cos^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+10A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^2/sec(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/6*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-16*B*cos(1/2
*d*x+1/2*c)^8+24*A*cos(1/2*d*x+1/2*c)^6+10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*co
s(1/2*d*x+1/2*c)^3+24*A*cos(1/2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-12
*B*cos(1/2*d*x+1/2*c)^6-20*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+
1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)
^3-42*B*cos(1/2*d*x+1/2*c)^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+
1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-38*A*cos(1/2*d*x+1
/2*c)^4+48*B*cos(1/2*d*x+1/2*c)^4+15*A*cos(1/2*d*x+1/2*c)^2-21*B*cos(1/2*d
*x+1/2*c)^2-A+B)/a^2/cos(1/2*d*x+1/2*c)^3/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.486.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.83

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$5(\sqrt{2}(-iA + 2iB) \cos(dx + c)^2 + 2\sqrt{2}(-iA + 2iB) \cos(dx + c) + \sqrt{2}(-iA + 2iB)) \text{weierstrassPIn}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(5/2),x, algorith
m="fricas")
```

3.486. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$

output `-1/6*(5*(sqrt(2)*(-I*A + 2*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(-I*A + 2*I*B)*cos(d*x + c) + sqrt(2)*(-I*A + 2*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(I*A - 2*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(I*A - 2*I*B)*cos(d*x + c) + sqrt(2)*(I*A - 2*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(sqrt(2)*(-4*I*A + 7*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(-4*I*A + 7*I*B)*cos(d*x + c) + sqrt(2)*(-4*I*A + 7*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(sqrt(2)*(4*I*A - 7*I*B)*cos(d*x + c)^2 + 2*sqrt(2)*(4*I*A - 7*I*B)*cos(d*x + c) + sqrt(2)*(4*I*A - 7*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(2*B*cos(d*x + c)^3 - (6*A - 13*B)*cos(d*x + c)^2 - 5*(A - 2*B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.486.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**2/sec(d*x+c)**(5/2),x)`

output `Timed out`

3.486.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2)), x)`

3.486.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^2/sec(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2)), x)`

3.486.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (a + a \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^2), x)`

3.487
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$$

3.487.1 Optimal result 4441
 3.487.2 Mathematica [C] (verified) 4442
 3.487.3 Rubi [A] (verified) 4442
 3.487.4 Maple [B] (warning: unable to verify) 4447
 3.487.5 Fricas [C] (verification not implemented) 4448
 3.487.6 Sympy [F(-1)] 4449
 3.487.7 Maxima [F(-1)] 4449
 3.487.8 Giac [F] 4450
 3.487.9 Mupad [F(-1)] 4450

3.487.1 Optimal result

Integrand size = 33, antiderivative size = 261

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx \\ &= -\frac{(49A - 9B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{10a^3d} \\ & \quad - \frac{(13A - 3B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{6a^3d} \\ & \quad + \frac{(49A - 9B) \sqrt{\sec(c + dx)} \sin(c + dx)}{10a^3d} - \frac{(A - B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{5d(a + a \sec(c + dx))^3} \\ & \quad - \frac{(8A - 3B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{15ad(a + a \sec(c + dx))^2} - \frac{(13A - 3B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{6d(a^3 + a^3 \sec(c + dx))} \end{aligned}$$

```
output -1/5*(A-B)*sec(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*sec(d*x+c))^3-1/15*(8*A-3*B)
*sec(d*x+c)^(5/2)*sin(d*x+c)/a/d/(a+a*sec(d*x+c))^2-1/6*(13*A-3*B)*sec(d*x
+c)^(3/2)*sin(d*x+c)/d/(a^3+a^3*sec(d*x+c))+1/10*(49*A-9*B)*sin(d*x+c)*sec
(d*x+c)^(1/2)/a^3/d-1/10*(49*A-9*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d
*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c
)^(1/2)/a^3/d-1/6*(13*A-3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*
c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)
/a^3/d
```


3.487.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 4.35 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.37

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx =$$

$$e^{-idx} \cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(-i(49A - 9B)e^{-2i(c+dx)}(1 + e^{i(c+dx)})^5 \sqrt{1 + e^{2i(c+dx)}} \text{Hypergeomet}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^3, x]`

output `-1/120*(Cos[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(((-I)*(49*A - 9*B)*(1 + E^(I*(c + d*x)))^5*Sqrt[1 + E^((2*I)*(c + d*x))]*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^((2*I)*(c + d*x)) + 160*(13*A - 3*B)*Cos[(c + d*x)/2]^5*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*(Cos[(c + d*x)/2] - I*Sin[(c + d*x)/2]) + (2*I)*(642*A - 102*B + (1082*A - 207*B)*Cos[c + d*x] + 6*(87*A - 17*B)*Cos[2*(c + d*x)] + 106*A*Cos[3*(c + d*x)] - 21*B*Cos[3*(c + d*x)] + (161*I)*A*Sin[c + d*x] - (6*I)*B*Sin[c + d*x] + (148*I)*A*Sin[2*(c + d*x)] - (18*I)*B*Sin[2*(c + d*x)] + (41*I)*A*Sin[3*(c + d*x)] - (6*I)*B*Sin[3*(c + d*x)]))*(Cos[(c + 3*d*x)/2] + I*Sin[(c + 3*d*x)/2]))/(a^3*d*E^(I*d*x)*(1 + Cos[c + d*x])^3)`

3.487.3 Rubi [A] (verified)

Time = 1.73 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.02, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4507, 3042, 4507, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^3} dx$$

↓ 3042

3.487. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a\sin\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{\sec^{7/2}(c+dx)\left(A\sec(c+dx)+B\right)}{\left(a\sec(c+dx)+a\right)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{7/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{4507} \\
& \frac{\int \frac{\sec^{5/2}(c+dx)\left(5a(A-B)-a(11A-B)\sec(c+dx)\right)}{2\left(\sec(c+dx)a+a\right)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)\sec^{7/2}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& \quad \downarrow \text{27} \\
& - \frac{\int \frac{\sec^{5/2}(c+dx)\left(5a(A-B)-a(11A-B)\sec(c+dx)\right)}{\left(\sec(c+dx)a+a\right)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{7/2}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& \quad \downarrow \text{3042} \\
& - \frac{\int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}\left(5a(A-B)-a(11A-B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(\csc\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{7/2}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& \quad \downarrow \text{4507} \\
& - \frac{\int \frac{\sec^{3/2}(c+dx)\left(3a^2(8A-3B)-a^2(41A-6B)\sec(c+dx)\right)}{\sec(c+dx)a+a} dx}{10a^2} + \frac{2a(8A-3B)\sin(c+dx)\sec^{5/2}(c+dx)}{3d\left(a\sec(c+dx)+a\right)^2} - \\
& \quad \frac{(A-B)\sin(c+dx)\sec^{7/2}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& \quad \downarrow \text{3042} \\
& - \frac{\int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(3a^2(8A-3B)-a^2(41A-6B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)}{\csc\left(c+dx+\frac{\pi}{2}\right)a+a} dx}{10a^2} + \frac{2a(8A-3B)\sin(c+dx)\sec^{5/2}(c+dx)}{3d\left(a\sec(c+dx)+a\right)^2} - \\
& \quad \frac{(A-B)\sin(c+dx)\sec^{7/2}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& \quad \downarrow \text{4507}
\end{aligned}$$

3.487. $\int \frac{(A+B\cos(c+dx))\sec^{3/2}(c+dx)}{(a+a\cos(c+dx))^3} dx$

$$\frac{\int \frac{1}{2} \sqrt{\sec(c+dx)} (5a^3(13A-3B) - 3a^3(49A-9B) \sec(c+dx)) dx}{a^2} + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} + \frac{2a(8A-3B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 27

$$\frac{\int \sqrt{\sec(c+dx)} (5a^3(13A-3B) - 3a^3(49A-9B) \sec(c+dx)) dx}{2a^2} + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} + \frac{2a(8A-3B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 3042

$$\frac{\int \sqrt{\csc(c+dx+\frac{\pi}{2})} (5a^3(13A-3B) - 3a^3(49A-9B) \csc(c+dx+\frac{\pi}{2})) dx}{2a^2} + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} + \frac{2a(8A-3B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 4274

$$\frac{5a^3(13A-3B) \int \sqrt{\sec(c+dx)} dx - 3a^3(49A-9B) \int \sec^{\frac{3}{2}}(c+dx) dx}{2a^2} + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} + \frac{2a(8A-3B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 3042

$$\frac{5a^3(13A-3B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - 3a^3(49A-9B) \int \csc(c+dx+\frac{\pi}{2})^{3/2} dx}{2a^2} + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)} + \frac{2a(8A-3B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 4255

3.487. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$

$$\frac{5a^3(13A-3B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - 3a^3(49A-9B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right) + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)}}{2a^2} + \frac{2a(8A-3B) \sin(c+dx)}{3d(a \sec(c+dx)+a)}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 3042

$$\frac{5a^3(13A-3B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx - 3a^3(49A-9B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right) + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)}}{2a^2} + \frac{2a(8A-3B) \sin(c+dx)}{3d(a \sec(c+dx)+a)}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 4258

$$\frac{5a^3(13A-3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx - 3a^3(49A-9B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx \right) + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)}}{2a^2} + \frac{2a(8A-3B) \sin(c+dx)}{3d(a \sec(c+dx)+a)}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 3042

$$\frac{5a^3(13A-3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^3(49A-9B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx \right) + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)}}{2a^2} + \frac{2a(8A-3B) \sin(c+dx)}{3d(a \sec(c+dx)+a)}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 3119

$$\frac{5a^3(13A-3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - 3a^3(49A-9B) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)|2\right)}{d} \right) + \frac{5a^2(13A-3B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(a \sec(c+dx)+a)}}{2a^2} + \frac{2a(8A-3B) \sin(c+dx)}{3d(a \sec(c+dx)+a)}$$

$$\frac{10a^2}{3a^2} \frac{(A-B) \sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3}$$

↓ 3120

3.487. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$

$$\frac{5a^2(13A-3B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{d(a\sec(c+dx)+a)} + \frac{10a^3(13A-3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) - 3a^3(49A-9B)\left(\frac{2\sin(c+dx)\sqrt{\sec(c+dx)}}{d} - \frac{2\sqrt{\cos(c+dx)}}{d}\right)}{3a^2} \Bigg/ \frac{10a^2}{(A-B)\sin(c+dx)\sec^{\frac{7}{2}}(c+dx)} \Bigg/ \frac{5d(a\sec(c+dx)+a)^3}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^3,x]`

output `-1/5*((A - B)*Sec[c + d*x]^(7/2)*Sin[c + d*x])/(d*(a + a*Sec[c + d*x])^3) - ((2*a*(8*A - 3*B)*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*(a + a*Sec[c + d*x])^2) + ((5*a^2*(13*A - 3*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))) + ((10*a^3*(13*A - 3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - 3*a^3*(49*A - 9*B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/(2*a^2)/(3*a^2)/(10*a^2)`

3.487.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

3.487. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$

```
rule 4255 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*(n - 2)/(n - 1)
  Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
  && IntegerQ[2*n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
  EqQ[n^2, 1/4]
```

```
rule 4274 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
  (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]
```

```
rule 4507 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
  a_))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(
  2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*
  (d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m -
  n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f,
  A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]
```

3.487.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 684 vs. $2(285) = 570$.

Time = 6.00 (sec) , antiderivative size = 685, normalized size of antiderivative = 2.62

method	result	size
default	Expression too large to display	685

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

$$3.487. \quad \int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$$

output

```

-1/60*(-2*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/
2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(65*A*EllipticF(cos(1/2
*d*x+1/2*c),2^(1/2))-147*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-15*B*Elli
pticF(cos(1/2*d*x+1/2*c),2^(1/2))+27*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2
))) *cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4+4*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2
*c)^2)^(1/2)*(65*A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-147*A*EllipticE(c
os(1/2*d*x+1/2*c),2^(1/2))-15*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+27*B
*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1
/2*c)-2*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*
d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(65*A*EllipticF(cos(1/2*d
*x+1/2*c),2^(1/2))-147*A*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-15*B*Ellipt
icF(cos(1/2*d*x+1/2*c),2^(1/2))+27*B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))
)*cos(1/2*d*x+1/2*c)+12*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/
2)*(49*A-9*B)*sin(1/2*d*x+1/2*c)^8-2*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+
1/2*c)^2)^(1/2)*(817*A-147*B)*sin(1/2*d*x+1/2*c)^6+6*(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(248*A-43*B)*sin(1/2*d*x+1/2*c)^4-(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(439*A-69*B)*sin(1/2*d*x+1/2*
c)^2/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)
^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d

```

3.487.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 481, normalized size of antiderivative = 1.84

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx =$$

$$\frac{5(\sqrt{2}(-13iA + 3iB) \cos(dx + c)^3 + 3\sqrt{2}(-13iA + 3iB) \cos(dx + c)^2 + 3\sqrt{2}(-13iA + 3iB) \cos(dx + c))}{(a + a \cos(c + dx))^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^3,x, algorithm="fracas")`

3.487. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$

output

```
-1/60*(5*(sqrt(2)*(-13*I*A + 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-13*I*A +
3*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-13*I*A + 3*I*B)*cos(d*x + c) + sqrt(2)
*(-13*I*A + 3*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x +
c)) + 5*(sqrt(2)*(13*I*A - 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(13*I*A - 3*I
*B)*cos(d*x + c)^2 + 3*sqrt(2)*(13*I*A - 3*I*B)*cos(d*x + c) + sqrt(2)*(13
*I*A - 3*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) +
3*(sqrt(2)*(49*I*A - 9*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(49*I*A - 9*I*B)*c
os(d*x + c)^2 + 3*sqrt(2)*(49*I*A - 9*I*B)*cos(d*x + c) + sqrt(2)*(49*I*A
- 9*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) +
I*sin(d*x + c))) + 3*(sqrt(2)*(-49*I*A + 9*I*B)*cos(d*x + c)^3 + 3*sqrt(2)
*(-49*I*A + 9*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-49*I*A + 9*I*B)*cos(d*x +
c) + sqrt(2)*(-49*I*A + 9*I*B))*weierstrassZeta(-4, 0, weierstrassPInvers
e(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*(49*A - 9*B)*cos(d*x + c)^
3 + 2*(188*A - 33*B)*cos(d*x + c)^2 + 5*(59*A - 9*B)*cos(d*x + c) + 60*A)*
sin(d*x + c)/sqrt(cos(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x +
c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)
```

3.487.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**3,x)`

output `Timed out`

3.487.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

output `Timed out`

3.487. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^3} dx$

3.487.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^3, x)`

3.487.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{(a + a \cos(c + dx))^3} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^3,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^3, x)`

3.488
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^3} dx$$

3.488.1 Optimal result 4451
 3.488.2 Mathematica [C] (verified) 4452
 3.488.3 Rubi [A] (verified) 4453
 3.488.4 Maple [A] (verified) 4458
 3.488.5 Fricas [C] (verification not implemented) 4458
 3.488.6 Sympy [F] 4459
 3.488.7 Maxima [F] 4459
 3.488.8 Giac [F] 4460
 3.488.9 Mupad [F(-1)] 4460

3.488.1 Optimal result

Integrand size = 33, antiderivative size = 222

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx \\ &= \frac{(9A + B)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{10a^3d} \\ &+ \frac{(3A + B)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{6a^3d} \\ &- \frac{(A - B)\sec^{\frac{5}{2}}(c + dx)\sin(c + dx)}{5d(a + a \sec(c + dx))^3} - \frac{(6A - B)\sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{15ad(a + a \sec(c + dx))^2} \\ &- \frac{(9A + B)\sqrt{\sec(c + dx)}\sin(c + dx)}{10d(a^3 + a^3 \sec(c + dx))} \end{aligned}$$

output

```
-1/5*(A-B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*sec(d*x+c))^3-1/15*(6*A-B)*s
ec(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*sec(d*x+c))^2-1/10*(9*A+B)*sin(d*x+c)*
sec(d*x+c)^(1/2)/d/(a^3+a^3*sec(d*x+c))+1/10*(9*A+B)*(cos(1/2*d*x+1/2*c)^2
)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c
)^(1/2)*sec(d*x+c)^(1/2)/a^3/d+1/6*(3*A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/co
s(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*se
c(d*x+c)^(1/2)/a^3/d
```

3.488.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.61 (sec) , antiderivative size = 793, normalized size of antiderivative = 3.57

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx =$$

$$\frac{3\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic}) \operatorname{Hypergeometric2F1}\left(\frac{c}{2}, \frac{c}{2}, \frac{c}{2} + dx, -1+e^{2ic}\right)\right)}{5d(a + a \cos(c + dx))^3}$$

$$- \frac{\sqrt{2}Be^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic}) \operatorname{Hypergeometric2F1}\left(\frac{c}{2}, \frac{c}{2}, \frac{c}{2} + dx, -1+e^{2ic}\right)\right)}{15d(a + a \cos(c + dx))^3}$$

$$+ \frac{2A \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \csc\left(\frac{c}{2}\right) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c + dx)} \sin(c)}{d(a + a \cos(c + dx))^3}$$

$$+ \frac{2B \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \csc\left(\frac{c}{2}\right) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c + dx)} \sin(c)}{3d(a + a \cos(c + dx))^3}$$

$$+ \frac{\cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c + dx)} \left(-\frac{2(9A+B) \cos(dx) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right)}{5d} + \frac{2 \sec\left(\frac{c}{2}\right) \sec^5\left(\frac{c}{2} + \frac{dx}{2}\right) \left(A \sin\left(\frac{dx}{2}\right) - B \sin\left(\frac{dx}{2}\right)\right)}{5d} + \frac{4 \sec\left(\frac{c}{2}\right)}{5d}\right)}{5d}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^3, x]`

output

```
(-3*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]*Sec[c/2])/(5*d*E^(I*d*x)*(a + a*Cos[c + d*x])^3) - (Sqrt[2]*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]*Sec[c/2])/(15*d*E^(I*d*x)*(a + a*Cos[c + d*x])^3) + (2*A*Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*Csc[c/2]*EllipticF[(c + d*x)/2, 2]*Sec[c/2]*Sqrt[Sec[c + d*x]]*Sin[c])/(d*(a + a*Cos[c + d*x])^3) + (2*B*Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*Csc[c/2]*EllipticF[(c + d*x)/2, 2]*Sec[c/2]*Sqrt[Sec[c + d*x]]*Sin[c])/(3*d*(a + a*Cos[c + d*x])^3) + (Cos[c/2 + (d*x)/2]^6*Sqrt[Sec[c + d*x]]*((-2*(9*A + B)*Cos[d*x]*Csc[c/2]*Sec[c/2])/(5*d) + (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^5*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(5*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(3*A*Sin[(d*x)/2] + B*Sin[(d*x)/2]))/(3*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(3*A*Sin[(d*x)/2] + 2*B*Sin[(d*x)/2]))/(15*d) + (4*(3*A + B)*Tan[c/2])/(3*d) + (4*(3*A + 2*B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(15*d) + (2*(A - B)*Sec[c/2 + (d*x)/2]^4*Tan[c/2])/(5*d))/(a + a*Cos[c + d*x])^3
```

3.488.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4507, 3042, 4507, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^3} dx$$

↓ 3042

$$\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^3} dx$$

↓ 3439

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(A\sec(c+dx)+B)}{(a\sec(c+dx)+a)^3} dx$$

3.488. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \int -\frac{\sec^{\frac{3}{2}}(c+dx)\left(3a(A-B)-a(9A+B)\sec(c+dx)\right)}{2\left(\sec(c+dx)a+a\right)^2} dx - \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& -\frac{\int \frac{\sec^{\frac{3}{2}}(c+dx)\left(3a(A-B)-a(9A+B)\sec(c+dx)\right)}{\left(\sec(c+dx)a+a\right)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& -\frac{\int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(3a(A-B)-a(9A+B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(\csc\left(c+dx+\frac{\pi}{2}\right)a+a\right)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& -\frac{\int \frac{\sqrt{\sec(c+dx)}\left(a^2(6A-B)-a^2(21A+4B)\sec(c+dx)\right)}{\sec(c+dx)a+a} dx}{3a^2} + \frac{2a(6A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d\left(a\sec(c+dx)+a\right)^2} \\
& \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& -\frac{\int \frac{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a^2(6A-B)-a^2(21A+4B)\csc\left(c+dx+\frac{\pi}{2}\right)\right)}{\csc\left(c+dx+\frac{\pi}{2}\right)a+a} dx}{3a^2} + \frac{2a(6A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d\left(a\sec(c+dx)+a\right)^2} \\
& \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& -\frac{\int -\frac{3(9A+B)a^3+5(3A+B)\sec(c+dx)a^3}{2\sqrt{\sec(c+dx)}} dx}{a^2} + \frac{3a^2(9A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d\left(a\sec(c+dx)+a\right)} + \frac{2a(6A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d\left(a\sec(c+dx)+a\right)^2} \\
& \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d\left(a\sec(c+dx)+a\right)^3} \\
& \int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^3} dx
\end{aligned}$$

3.488. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\frac{3a^2(9A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{\int \frac{3(9A+B)a^3+5(3A+B) \sec(c+dx)a^3}{\sqrt{\sec(c+dx)}} dx}{3a^2}}{10a^2} + \frac{2a(6A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2} \\
 & \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \quad \downarrow \quad \text{3042} \\
 & \frac{\frac{3a^2(9A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{\int \frac{3(9A+B)a^3+5(3A+B) \csc(c+dx+\frac{\pi}{2})a^3}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{3a^2}}{10a^2} + \frac{2a(6A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2} \\
 & \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \quad \downarrow \quad \text{4274} \\
 & \frac{\frac{3a^2(9A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{3a^3(9A+B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + 5a^3(3A+B) \int \sqrt{\sec(c+dx)} dx}{3a^2}}{10a^2} + \frac{2a(6A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2} \\
 & \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \quad \downarrow \quad \text{3042} \\
 & \frac{\frac{3a^2(9A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{3a^3(9A+B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5a^3(3A+B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{3a^2}}{10a^2} + \frac{2a(6A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2} \\
 & \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \quad \downarrow \quad \text{4258} \\
 & \frac{\frac{3a^2(9A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{5a^3(3A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^3(9A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{3a^2}}{10a^2} + \frac{2a(6A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d(a \sec(c+dx)+a)^2} \\
 & \frac{(A-B) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \quad \downarrow \quad \text{3042}
 \end{aligned}$$

3.488. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\frac{3a^2(9A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{5a^3(3A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + 3a^3(9A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx}{3a^2} \frac{10a^2}{2a^2} + \dots \\
 & \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
 & \quad \downarrow \text{3119} \\
 & \frac{\frac{3a^2(9A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{5a^3(3A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{6a^3(9A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d}}{3a^2} \frac{10a^2}{2a^2} + 2a(6) \\
 & \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
 & \quad \downarrow \text{3120} \\
 & \frac{\frac{3a^2(9A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{10a^3(3A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{6a^3(9A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d}}{3a^2} \frac{10a^2}{2a^2} + 2a(6) \\
 & \frac{(A-B)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3}
 \end{aligned}$$

```
input Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^3,x]
```

```
output -1/5*((A - B)*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(d*(a + a*Sec[c + d*x])^3)
- ((2*a*(6*A - B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d*(a + a*Sec[c + d*x])^2)
+ (-1/2*((6*a^3*(9*A + B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d
+ (10*a^3*(3*A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d/a^2
+ (3*a^2*(9*A + B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))/(3*a^2))/(10*a^2)
```

3.488. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^3} dx$

3.488.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_))^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4507 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

3.488.4 Maple [A] (verified)

Time = 5.52 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.03

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(108A\left(\cos^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-30A\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\right)$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(108*A*cos(1/
2*d*x+1/2*c)^8-30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+
1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+54*A*c
os(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+
1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))+12*B*cos(1/2*d*x+1/2*c)^8-1
0*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*Ellipti
cF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+6*B*cos(1/2*d*x+1/2*c)
^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*Elliptic
E(cos(1/2*d*x+1/2*c),2^(1/2))-138*A*cos(1/2*d*x+1/2*c)^6-22*B*cos(1/2*d*x+
1/2*c)^6+24*A*cos(1/2*d*x+1/2*c)^4+6*B*cos(1/2*d*x+1/2*c)^4+3*A*cos(1/2*d*
x+1/2*c)^2+7*B*cos(1/2*d*x+1/2*c)^2+3*A-3*B)/a^3/cos(1/2*d*x+1/2*c)^5/(-2*
sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos
(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.488.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 476, normalized size of antiderivative = 2.14

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx =$$

$$\frac{5(\sqrt{2}(3iA + iB) \cos(dx + c)^3 + 3\sqrt{2}(3iA + iB) \cos(dx + c)^2 + 3\sqrt{2}(3iA + iB) \cos(dx + c) + \sqrt{2})}{(a + a \cos(c + dx))^3}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^3,x, algorith
m="fricas")
```

$$3.488. \int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^3} dx$$

output `-1/60*(5*(sqrt(2)*(3*I*A + I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(3*I*A + I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(3*I*A + I*B)*cos(d*x + c) + sqrt(2)*(3*I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(-3*I*A - I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-3*I*A - I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-3*I*A - I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(sqrt(2)*(-9*I*A - I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-9*I*A - I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-9*I*A - I*B)*cos(d*x + c) + sqrt(2)*(-9*I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(sqrt(2)*(9*I*A + I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(9*I*A + I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(9*I*A + I*B)*cos(d*x + c) + sqrt(2)*(9*I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*(9*A + B)*cos(d*x + c)^3 + 2*(33*A + 2*B)*cos(d*x + c)^2 + 5*(9*A - B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.488.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx$$

$$= \frac{\int \frac{A \sqrt{\sec(c + dx)}}{\cos^3(c + dx) + 3 \cos^2(c + dx) + 3 \cos(c + dx) + 1} dx + \int \frac{B \cos(c + dx) \sqrt{\sec(c + dx)}}{\cos^3(c + dx) + 3 \cos^2(c + dx) + 3 \cos(c + dx) + 1} dx}{a^3}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**3,x)`

output `(Integral(A*sqrt(sec(c + d*x))/(cos(c + d*x)**3 + 3*cos(c + d*x)**2 + 3*cos(c + d*x) + 1), x) + Integral(B*cos(c + d*x)*sqrt(sec(c + d*x))/(cos(c + d*x)**3 + 3*cos(c + d*x)**2 + 3*cos(c + d*x) + 1), x))/a**3`

3.488.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^3,x, algorithm="maxima")`

3.488. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^3} dx$

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^3, x)`

3.488.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^3, x)`

3.488.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + a \cos(c + dx))^3} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^3, x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^3, x)`

3.489 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

3.489.1 Optimal result 4461
 3.489.2 Mathematica [C] (verified) 4462
 3.489.3 Rubi [A] (verified) 4463
 3.489.4 Maple [A] (verified) 4468
 3.489.5 Fricas [C] (verification not implemented) 4469
 3.489.6 Sympy [F(-1)] 4470
 3.489.7 Maxima [F] 4470
 3.489.8 Giac [F] 4471
 3.489.9 Mupad [F(-1)] 4471

3.489.1 Optimal result

Integrand size = 33, antiderivative size = 216

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{(A - B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{10a^3d}$$

$$+ \frac{(A + B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{6a^3d}$$

$$- \frac{(A - B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{5d(a + a \sec(c + dx))^3} - \frac{(4A + B) \sqrt{\sec(c + dx)} \sin(c + dx)}{15ad(a + a \sec(c + dx))^2}$$

$$+ \frac{(A + B) \sqrt{\sec(c + dx)} \sin(c + dx)}{6d(a^3 + a^3 \sec(c + dx))}$$

output

```
-1/5*(A-B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*sec(d*x+c))^3-1/15*(4*A+B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*sec(d*x+c))^2+1/6*(A+B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a^3+a^3*sec(d*x+c))+1/10*(A-B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^3/d+1/6*(A+B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^3/d
```

3.489.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.50 (sec) , antiderivative size = 792, normalized size of antiderivative = 3.67

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx =$$

$$\frac{\sqrt{2} A e^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic}) \operatorname{Hypergeometric2F1}\left(\frac{c}{2}, \frac{c}{2}, \frac{c}{2}+1, -\frac{e^{2i(c+dx)}}{1+e^{2i(c+dx)}}\right)\right)}{15d(a+a\cos(c+dx))^3}$$

$$+ \frac{\sqrt{2} B e^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic}) \operatorname{Hypergeometric2F1}\left(\frac{c}{2}, \frac{c}{2}, \frac{c}{2}+1, -\frac{e^{2i(c+dx)}}{1+e^{2i(c+dx)}}\right)\right)}{15d(a+a\cos(c+dx))^3}$$

$$+ \frac{2A \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \csc\left(\frac{c}{2}\right) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c+dx)} \sin(c)}{3d(a+a\cos(c+dx))^3}$$

$$+ \frac{2B \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c+dx)} \csc\left(\frac{c}{2}\right) \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c+dx)} \sin(c)}{3d(a+a\cos(c+dx))^3}$$

$$+ \frac{\cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c+dx)} \left(-\frac{2(A-B) \cos(dx) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right)}{5d} + \frac{4 \sec\left(\frac{c}{2}\right) \sec^3\left(\frac{c}{2} + \frac{dx}{2}\right) \left(2A \sin\left(\frac{dx}{2}\right) - 7B \sin\left(\frac{dx}{2}\right)\right)}{15d} - \frac{2 \sec\left(\frac{c}{2}\right)}{15d}\right)}{15d}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sqrt[Sec[c + d*x]]), x]`

output

$$\begin{aligned}
& -1/15*(\text{Sqrt}[2]*A*\text{Sqrt}[E^{(I*(c+d*x))/(1+E^{(2*I)*(c+d*x)})}]]*\text{Sqrt}[1+E^{(2*I)*(c+d*x)}]]*\text{Cos}[c/2+(d*x)/2]^6*\text{Csc}[c/2]*(-3*\text{Sqrt}[1+E^{(2*I)*(c+d*x)}]]+E^{(2*I)*d*x})*(-1+E^{(2*I)*c}))*\text{Hypergeometric2F1}[1/2, 3/4, 7/4, -E^{(2*I)*(c+d*x)}]]*\text{Sec}[c/2]/(d*E^{(I*d*x)}*(a+a*\text{Cos}[c+d*x])^3) \\
& + (\text{Sqrt}[2]*B*\text{Sqrt}[E^{(I*(c+d*x))/(1+E^{(2*I)*(c+d*x)})}]]*\text{Sqrt}[1+E^{(2*I)*(c+d*x)}]]*\text{Cos}[c/2+(d*x)/2]^6*\text{Csc}[c/2]*(-3*\text{Sqrt}[1+E^{(2*I)*(c+d*x)}]]+E^{(2*I)*d*x})*(-1+E^{(2*I)*c}))*\text{Hypergeometric2F1}[1/2, 3/4, 7/4, -E^{(2*I)*(c+d*x)}]]*\text{Sec}[c/2]/(15*d*E^{(I*d*x)}*(a+a*\text{Cos}[c+d*x])^3) \\
& + (2*A*\text{Cos}[c/2+(d*x)/2]^6*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Csc}[c/2]*\text{EllipticF}[(c+d*x)/2, 2]*\text{Sec}[c/2]*\text{Sqrt}[\text{Sec}[c+d*x]]*\text{Sin}[c])/(3*d*(a+a*\text{Cos}[c+d*x])^3) \\
& + (2*B*\text{Cos}[c/2+(d*x)/2]^6*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Csc}[c/2]*\text{EllipticF}[(c+d*x)/2, 2]*\text{Sec}[c/2]*\text{Sqrt}[\text{Sec}[c+d*x]]*\text{Sin}[c])/(3*d*(a+a*\text{Cos}[c+d*x])^3) + \\
& (\text{Cos}[c/2+(d*x)/2]^6*\text{Sqrt}[\text{Sec}[c+d*x]]*((-2*(A-B)*\text{Cos}[d*x]*\text{Csc}[c/2]*\text{Sec}[c/2])/(5*d) + (4*\text{Sec}[c/2]*\text{Sec}[c/2+(d*x)/2]^3*(2*A*\text{Sin}[(d*x)/2] - 7*B*\text{Sin}[(d*x)/2]))/(15*d) - (2*\text{Sec}[c/2]*\text{Sec}[c/2+(d*x)/2]^5*(A*\text{Sin}[(d*x)/2] - B*\text{Sin}[(d*x)/2]))/(5*d) + (4*\text{Sec}[c/2]*\text{Sec}[c/2+(d*x)/2]*(A*\text{Sin}[(d*x)/2] + B*\text{Sin}[(d*x)/2]))/(3*d) + (4*(A+B)*\text{Tan}[c/2])/(3*d) + (4*(2*A - 7*B)*\text{Sec}[c/2+(d*x)/2]^2*\text{Tan}[c/2])/(15*d) - (2*(A-B)*\text{Sec}[c/2+(d*x)/2]^4*\text{Tan}[c/2])/(5*d)))/(a+a*\text{Cos}[c+d*x])^3
\end{aligned}$$

3.489.3 Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4507, 25, 3042, 4508, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a \cos(c + dx) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{\sec^{\frac{3}{2}}(c + dx)(A \sec(c + dx) + B)}{(a \sec(c + dx) + a)^3} dx
\end{aligned}$$

3.489. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+a\right)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{-\frac{\sqrt{\sec(c+dx)}(a(A-B)-a(7A+3B)\sec(c+dx))}{2(\sec(c+dx)a+a)^2} dx}{5a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{4507} \\
& -\frac{\int \frac{\sqrt{\sec(c+dx)}(a(A-B)-a(7A+3B)\sec(c+dx))}{(\sec(c+dx)a+a)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& -\frac{\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(a(A-B)-a(7A+3B)\csc(c+dx+\frac{\pi}{2}))}{(\csc(c+dx+\frac{\pi}{2})a+a)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& -\frac{\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(a(A-B)-a(7A+3B)\csc(c+dx+\frac{\pi}{2}))}{(\csc(c+dx+\frac{\pi}{2})a+a)^2} dx}{10a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{4507} \\
& -\frac{\int \frac{(4A+B)a^2+3(3A+2B)\sec(c+dx)a^2}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{3a^2} + \frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{10a}^2 \\
& -\frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{\int \frac{(4A+B)a^2+3(3A+2B)\sec(c+dx)a^2}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{3a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{25} \\
& -\frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{\int \frac{(4A+B)a^2+3(3A+2B)\csc(c+dx+\frac{\pi}{2})a^2}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(\csc(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{3042} \\
& -\frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{\int \frac{(4A+B)a^2+3(3A+2B)\csc(c+dx+\frac{\pi}{2})a^2}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(\csc(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} - \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{4508} \\
& -\frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{\int \frac{3(A-B)a^3+5(A+B)\sec(c+dx)a^3}{2\sqrt{\sec(c+dx)}} dx}{a^2} + \frac{5a^2(A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} \\
& \quad \downarrow \text{10a}^2 \\
& \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3}
\end{aligned}$$

3.489. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^3\sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{2a(4A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2} - \frac{\int \frac{3(A-B)a^3+5(A+B) \sec(c+dx)a^3}{\sqrt{\sec(c+dx)}} dx}{2a^2} + \frac{5a^2(A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} \\
 & \frac{10a^2}{5d(a \sec(c+dx)+a)^3} \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \downarrow 3042 \\
 & \frac{2a(4A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2} - \frac{\int \frac{3(A-B)a^3+5(A+B) \csc(c+dx+\frac{\pi}{2})a^3}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{2a^2} + \frac{5a^2(A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} \\
 & \frac{10a^2}{5d(a \sec(c+dx)+a)^3} \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \downarrow 4274 \\
 & \frac{2a(4A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2} - \frac{3a^3(A-B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + 5a^3(A+B) \int \sqrt{\sec(c+dx)} dx}{2a^2} + \frac{5a^2(A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} \\
 & \frac{10a^2}{5d(a \sec(c+dx)+a)^3} \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \downarrow 3042 \\
 & \frac{2a(4A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2} - \frac{3a^3(A-B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5a^3(A+B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{2a^2} + \frac{5a^2(A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} \\
 & \frac{10a^2}{5d(a \sec(c+dx)+a)^3} \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \downarrow 4258 \\
 & \frac{2a(4A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2} - \frac{5a^3(A+B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3a^3(A-B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{2a^2} + \frac{5a^2(A+B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} \\
 & \frac{10a^2}{5d(a \sec(c+dx)+a)^3} \frac{(A-B) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{5d(a \sec(c+dx)+a)^3} \\
 & \downarrow 3042
 \end{aligned}$$

3.489. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
 & \frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{5a^3(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+3a^3(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\sin(c+dx+\frac{\pi}{2})}dx}{2a^2} \\
 & \frac{10a^2}{3a^2} \\
 & \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
 & \quad \downarrow \text{3119} \\
 & \frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{5a^3(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{6a^3(A-B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx),2\right)}{d}}{2a^2} + \frac{5a^2}{3a^2} \\
 & \frac{10a^2}{3a^2} \\
 & \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3} \\
 & \quad \downarrow \text{3120} \\
 & \frac{2a(4A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} - \frac{5a^2(A+B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} + \frac{10a^3(A+B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{6a^3(A-B)\sqrt{\cos(c+dx)}}{2a^2} \\
 & \frac{10a^2}{3a^2} \\
 & \frac{(A-B)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{5d(a\sec(c+dx)+a)^3}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sqrt[Sec[c + d*x]]),x]`

output `-1/5*((A - B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*(a + a*Sec[c + d*x])^3) - ((2*a*(4*A + B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Sec[c + d*x])^2) - (((6*a^3*(A - B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*a^3*(A + B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/(2*a^2) + (5*a^2*(A + B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))/(3*a^2))/(10*a^2)`

3.489. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

3.489.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4507 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[d*(A*b
- a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(
2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*
(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m
- n + 1) + A*b*(m + n))*Csc[e + f*x], x], x] /; FreeQ[{a, b, d, e, f,
A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && G
tQ[n, 0]
```

```
rule 4508 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-(A*b
- a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m +
1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Cs
c[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[
e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B
, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]
```

3.489.4 Maple [A] (verified)

Time = 5.77 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.09

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(12A\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 10A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)\right)$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

3.489.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$$

output `1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*A*cos(1/2*d*x+1/2*c)^8-10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+6*A*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-12*B*cos(1/2*d*x+1/2*c)^8-10*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5-6*B*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-22*A*cos(1/2*d*x+1/2*c)^6+2*B*cos(1/2*d*x+1/2*c)^6+6*A*cos(1/2*d*x+1/2*c)^4+24*B*cos(1/2*d*x+1/2*c)^4+7*A*cos(1/2*d*x+1/2*c)^2-17*B*cos(1/2*d*x+1/2*c)^2-3*A+3*B)/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.489.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 472, normalized size of antiderivative = 2.19

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx =$$

$$\frac{5(\sqrt{2}(iA + iB) \cos(dx + c))^3 + 3\sqrt{2}(iA + iB) \cos(dx + c)^2 + 3\sqrt{2}(iA + iB) \cos(dx + c) + \sqrt{2}(iA + iB)}{\dots}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(1/2),x, algorithm m="fracas")`

output `-1/60*(5*(sqrt(2)*(I*A + I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(I*A + I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(I*A + I*B)*cos(d*x + c) + sqrt(2)*(I*A + I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(-I*A - I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-I*A - I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-I*A - I*B)*cos(d*x + c) + sqrt(2)*(-I*A - I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(sqrt(2)*(-I*A + I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-I*A + I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-I*A + I*B)*cos(d*x + c) + sqrt(2)*(-I*A + I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(sqrt(2)*(I*A - I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(I*A - I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(I*A - I*B)*cos(d*x + c) + sqrt(2)*(I*A - I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*(A - B)*cos(d*x + c)^3 + 2*(2*A - 7*B)*cos(d*x + c)^2 - 5*(A + B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.489.6 SymPy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.489.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sqrt(sec(d*x + c))), x)`

3.489. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

3.489.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sqrt(sec(d*x + c))), x)`

3.489.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^3), x)`

$$3.490 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$$

3.490.1 Optimal result	4472
3.490.2 Mathematica [C] (verified)	4473
3.490.3 Rubi [A] (verified)	4474
3.490.4 Maple [A] (verified)	4479
3.490.5 Fricas [C] (verification not implemented)	4480
3.490.6 Sympy [F(-1)]	4480
3.490.7 Maxima [F]	4481
3.490.8 Giac [F]	4481
3.490.9 Mupad [F(-1)]	4481

3.490.1 Optimal result

Integrand size = 33, antiderivative size = 222

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx \\ &= -\frac{(A + 9B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{10a^3d} \\ & \quad + \frac{(A + 3B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{6a^3d} \\ & \quad - \frac{(A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{5d(a + a \sec(c + dx))^3} + \frac{(2A + 3B) \sqrt{\sec(c + dx)} \sin(c + dx)}{15ad(a + a \sec(c + dx))^2} \\ & \quad + \frac{(A + 3B) \sqrt{\sec(c + dx)} \sin(c + dx)}{6d(a^3 + a^3 \sec(c + dx))} \end{aligned}$$

output

```
-1/5*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*sec(d*x+c))^3+1/15*(2*A+3*B)
*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*sec(d*x+c))^2+1/6*(A+3*B)*sin(d*x+c)
*sec(d*x+c)^(1/2)/d/(a^3+a^3*sec(d*x+c))-1/10*(A+9*B)*(cos(1/2*d*x+1/2*c)
^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+
c)^(1/2)*sec(d*x+c)^(1/2)/a^3/d+1/6*(A+3*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/c
os(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*s
ec(d*x+c)^(1/2)/a^3/d
```

3.490.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 7.98 (sec) , antiderivative size = 793, normalized size of antiderivative = 3.57

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \text{Hypergeometric}}{15d(a + a \cos(c + dx))^3}$$

$$+ \frac{3\sqrt{2}Be^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \text{Hypergeometric}}{5d(a + a \cos(c + dx))^3}$$

$$+ \frac{2A \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \csc\left(\frac{c}{2}\right) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c + dx)} \sin(c)}{3d(a + a \cos(c + dx))^3}$$

$$+ \frac{2B \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \csc\left(\frac{c}{2}\right) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c + dx)} \sin(c)}{d(a + a \cos(c + dx))^3}$$

$$+ \frac{\cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c + dx)} \left(\frac{2(A+9B) \cos(dx) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right)}{5d} - \frac{4 \sec\left(\frac{c}{2}\right) \sec^3\left(\frac{c}{2} + \frac{dx}{2}\right) \left(7A \sin\left(\frac{dx}{2}\right) - 12B \sin\left(\frac{dx}{2}\right)\right)}{15d}\right)}{4 \sec\left(\frac{c}{2}\right)}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sec[c + d*x]^(3/2)),x]`

output

```
(Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]*Sec[c/2])/(15*d*E^(I*d*x)*(a + a*Cos[c + d*x])^3) + (3*Sqrt[2]*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))]*Sec[c/2])/(5*d*E^(I*d*x)*(a + a*Cos[c + d*x])^3) + (2*A*Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*Csc[c/2]*EllipticF[(c + d*x)/2, 2]*Sec[c/2]*Sqrt[Sec[c + d*x]]*Sin[c])/(3*d*(a + a*Cos[c + d*x])^3) + (2*B*Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*Csc[c/2]*EllipticF[(c + d*x)/2, 2]*Sec[c/2]*Sqrt[Sec[c + d*x]]*Sin[c])/(d*(a + a*Cos[c + d*x])^3) + (Cos[c/2 + (d*x)/2]^6*Sqrt[Sec[c + d*x]]*((2*(A + 9*B)*Cos[d*x]*Csc[c/2]*Sec[c/2])/(5*d) - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(7*A*Sin[(d*x)/2] - 12*B*Sin[(d*x)/2]))/(15*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(A*Sin[(d*x)/2] - 9*B*Sin[(d*x)/2]))/(3*d) + (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^5*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(5*d) + (4*(A - 9*B)*Tan[c/2])/(3*d) - (4*(7*A - 12*B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(15*d) + (2*(A - B)*Sec[c/2 + (d*x)/2]^4*Tan[c/2])/(5*d))/(a + a*Cos[c + d*x])^3
```

3.490.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4507, 27, 3042, 4508, 3042, 4508, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^3} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}}(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^3} dx$$

↓ 3439

$$\int \frac{\sqrt{\sec(c + dx)}(A \sec(c + dx) + B)}{(a \sec(c + dx) + a)^3} dx$$

3.490. $\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})} (A \csc(c+dx+\frac{\pi}{2}) + B)}{(a \csc(c+dx+\frac{\pi}{2}) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{a(A-B) + 5a(A+B) \sec(c+dx)}{2\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)^2} dx - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{4507} \\
& \int \frac{a(A-B) + 5a(A+B) \sec(c+dx)}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)^2} dx - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{27} \\
& \int \frac{a(A-B) + 5a(A+B) \csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(\csc(c+dx+\frac{\pi}{2})a+a)^2} dx - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(A-6B)a^2 + 3(2A+3B) \sec(c+dx)a^2}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{3a^2} + \frac{2a(2A+3B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{4508} \\
& \frac{\int \frac{(A-6B)a^2 + 3(2A+3B) \csc(c+dx+\frac{\pi}{2})a^2}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(\csc(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} + \frac{2a(2A+3B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(A-6B)a^2 + 3(2A+3B) \sec(c+dx)a^2}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{3a^2} + \frac{5a^2(A+3B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx) + a)} + \frac{2a(2A+3B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{4508} \\
& \frac{\int \frac{-3a^3(A+9B) - 5a^3(A+3B) \sec(c+dx)}{2\sqrt{\sec(c+dx)}} dx}{a^2} + \frac{5a^2(A+3B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx) + a)} + \frac{2a(2A+3B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx) + a)^2} - \frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.490. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{\int \frac{3a^3(A+9B)-5a^3(A+3B)\sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{3a^2}}{3a^2} + \frac{2a(2A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} -$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}} \frac{1}{5d(a\sec(c+dx)+a)^3}$$

↓ 3042

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{\int \frac{3a^3(A+9B)-5a^3(A+3B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{3a^2}}{3a^2} + \frac{2a(2A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} -$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}} \frac{1}{5d(a\sec(c+dx)+a)^3}$$

↓ 4274

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{3a^3(A+9B)\int \frac{1}{\sqrt{\sec(c+dx)}} dx - 5a^3(A+3B)\int \sqrt{\sec(c+dx)} dx}{3a^2}}{3a^2} + \frac{2a(2A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} -$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}} \frac{1}{5d(a\sec(c+dx)+a)^3}$$

↓ 3042

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{3a^3(A+9B)\int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - 5a^3(A+3B)\int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{3a^2}}{3a^2} + \frac{2a(2A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} -$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}} \frac{1}{5d(a\sec(c+dx)+a)^3}$$

↓ 4258

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{3a^3(A+9B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int \sqrt{\cos(c+dx)} dx - 5a^3(A+3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3a^2}}{3a^2} + \frac{2a(2A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a\sec(c+dx)+a)^2} -$$

$$\frac{10a^2}{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}} \frac{1}{5d(a\sec(c+dx)+a)^3}$$

↓ 3042

3.490. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{3a^3(A+9B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx - 5a^3(A+3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx}{3a^2} + \frac{2a^2}{3a^2}$$

$$\frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a\sec(c+dx)+a)^3} \cdot 10a^2$$

↓ 3119

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{6a^3(A+9B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d} - 5a^3(A+3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx}{3a^2} + \frac{2a^2}{3a^2}$$

$$\frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a\sec(c+dx)+a)^3} \cdot 10a^2$$

↓ 3120

$$\frac{\frac{5a^2(A+3B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a\sec(c+dx)+a)} - \frac{6a^3(A+9B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d} - \frac{10a^3(A+3B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d}}{3a^2} + \frac{2a^2}{3a^2}$$

$$\frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a\sec(c+dx)+a)^3} \cdot 10a^2$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sec[c + d*x]^(3/2)),x]`

output `-1/5*((A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x])^3) + ((2*a*(2*A + 3*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Sec[c + d*x])^2) + (-1/2*((6*a^3*(A + 9*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (10*a^3*(A + 3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/a^2 + (5*a^2*(A + 3*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))/(3*a^2))/(10*a^2)`

3.490. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$

3.490.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_))^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`
- rule 4507 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_))^(m_)*(csc[(e_.) + (f_)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[d*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^(n - 1)/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[A*(a*d*(n - 1)) - B*(b*d*(n - 1)) - d*(a*B*(m - n + 1) + A*b*(m + n))*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0]`

3.490.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^2(c+dx)} dx$$

```
rule 4508 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-(A*b
- a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m +
1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Cs
c[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[
e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B
, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]
```

3.490.4 Maple [A] (verified)

Time = 6.27 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.03

method	result
default	$-\frac{\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(12A \left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 10A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} + 1 F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{\dots}\right)\right)}{\dots}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3/sec(d*x+c)^(3/2), x, method=_RETURNV
ERBOSE)
```

```
output -1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(12*A*cos(1/
2*d*x+1/2*c)^8+10*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+
1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*cos(1/2*d*x+1/2*c)^5+6*A*co
s(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2))+108*B*cos(1/2*d*x+1/2*c)^8+3
0*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*Ellipti
cF(cos(1/2*d*x+1/2*c), 2^(1/2))*cos(1/2*d*x+1/2*c)^5+54*B*cos(1/2*d*x+1/2*c
)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c), 2^(1/2))-2*A*cos(1/2*d*x+1/2*c)^6-198*B*cos(1/2*d*x+
1/2*c)^6-24*A*cos(1/2*d*x+1/2*c)^4+114*B*cos(1/2*d*x+1/2*c)^4+17*A*cos(1/2
*d*x+1/2*c)^2-27*B*cos(1/2*d*x+1/2*c)^2-3*A+3*B)/a^3/cos(1/2*d*x+1/2*c)^5/
(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2
*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.490.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 474, normalized size of antiderivative = 2.14

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{5(\sqrt{2}(iA + 3iB) \cos(dx + c))^3 + 3\sqrt{2}(iA + 3iB) \cos(dx + c)^2 + 3\sqrt{2}(iA + 3iB) \cos(dx + c) + \sqrt{2}}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(3/2),x, algorithm
m="fricas")
```

```
output -1/60*(5*(sqrt(2)*(I*A + 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(I*A + 3*I*B)*c
os(d*x + c)^2 + 3*sqrt(2)*(I*A + 3*I*B)*cos(d*x + c) + sqrt(2)*(I*A + 3*I*
B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)
*(-I*A - 3*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-I*A - 3*I*B)*cos(d*x + c)^2 +
3*sqrt(2)*(-I*A - 3*I*B)*cos(d*x + c) + sqrt(2)*(-I*A - 3*I*B))*weierstra
ssPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(sqrt(2)*(I*A + 9*I*B
)*cos(d*x + c)^3 + 3*sqrt(2)*(I*A + 9*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(I*A
+ 9*I*B)*cos(d*x + c) + sqrt(2)*(I*A + 9*I*B))*weierstrassZeta(-4, 0, wei
erstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(sqrt(2)*(-I*A
- 9*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-I*A - 9*I*B)*cos(d*x + c)^2 + 3*sqr
t(2)*(-I*A - 9*I*B)*cos(d*x + c) + sqrt(2)*(-I*A - 9*I*B))*weierstrassZeta
(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*
(A + 9*B)*cos(d*x + c)^3 + 2*(7*A + 18*B)*cos(d*x + c)^2 + 5*(A + 3*B)*cos
(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*
d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)
```

3.490.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3/sec(d*x+c)**(3/2),x)
```

```
output Timed out
```

3.490. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$

3.490.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2)), x)`

3.490.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2)), x)`

3.490.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^3), x)`

3.491
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$$

3.491.1 Optimal result 4482
 3.491.2 Mathematica [C] (verified) 4483
 3.491.3 Rubi [A] (verified) 4484
 3.491.4 Maple [A] (verified) 4488
 3.491.5 Fricas [C] (verification not implemented) 4489
 3.491.6 Sympy [F(-1)] 4490
 3.491.7 Maxima [F] 4490
 3.491.8 Giac [F] 4491
 3.491.9 Mupad [F(-1)] 4491

3.491.1 Optimal result

Integrand size = 33, antiderivative size = 228

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{(9A - 49B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{10a^3 d}$$

$$+ \frac{(3A - 13B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{6a^3 d}$$

$$+ \frac{(A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{5d(a + a \sec(c + dx))^3} + \frac{(3A - 8B) \sqrt{\sec(c + dx)} \sin(c + dx)}{15ad(a + a \sec(c + dx))^2}$$

$$+ \frac{(3A - 13B) \sqrt{\sec(c + dx)} \sin(c + dx)}{6d(a^3 + a^3 \sec(c + dx))}$$

output

```
1/5*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*sec(d*x+c))^3+1/15*(3*A-8*B)*
sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*sec(d*x+c))^2+1/6*(3*A-13*B)*sin(d*x+
c)*sec(d*x+c)^(1/2)/d/(a^3+a^3*sec(d*x+c))-1/10*(9*A-49*B)*(cos(1/2*d*x+1/
2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos
(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^3/d+1/6*(3*A-13*B)*(cos(1/2*d*x+1/2*c))^2
^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)
^(1/2)*sec(d*x+c)^(1/2)/a^3/d
```

3.491.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 8.45 (sec) , antiderivative size = 817, normalized size of antiderivative = 3.58

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{3\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{e^{2i(c+dx)}}{1+e^{2i(c+dx)}}\right)}{5d(a + a \cos(c + dx))^3}$$

$$- \frac{49\sqrt{2}Be^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \csc\left(\frac{c}{2}\right) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{e^{2i(c+dx)}}{1+e^{2i(c+dx)}}\right)}{15d(a + a \cos(c + dx))^3}$$

$$+ \frac{2A \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \csc\left(\frac{c}{2}\right) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c + dx)} \sin(c)}{d(a + a \cos(c + dx))^3}$$

$$- \frac{26B \cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\cos(c + dx)} \csc\left(\frac{c}{2}\right) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sec\left(\frac{c}{2}\right) \sqrt{\sec(c + dx)} \sin(c)}{3d(a + a \cos(c + dx))^3}$$

$$+ \frac{\cos^6\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c + dx)} \left(-\frac{2(-9A+39B+10B \cos(2c)) \cos(dx) \csc\left(\frac{c}{2}\right) \sec\left(\frac{c}{2}\right)}{5d} - \frac{4 \sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) \left(9A \sin\left(\frac{dx}{2}\right) - 23B \sin\left(\frac{c}{2}\right)\right)}{3d}\right)}{d(a + a \cos(c + dx))^3}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sec[c + d*x]^(5/2)),x]`

output

```
(3*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*Sec[c/2])/(5*d*E^(I*d*x)*(a + a*Cos[c + d*x])^3) - (49*Sqrt[2]*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*Cos[c/2 + (d*x)/2]^6*Csc[c/2]*(-3*Sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])*Sec[c/2])/(15*d*E^(I*d*x)*(a + a*Cos[c + d*x])^3) + (2*A*Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*Csc[c/2]*EllipticF[(c + d*x)/2, 2]*Sec[c/2]*Sqrt[Sec[c + d*x]]*Sin[c])/(d*(a + a*Cos[c + d*x])^3) - (26*B*Cos[c/2 + (d*x)/2]^6*Sqrt[Cos[c + d*x]]*Csc[c/2]*EllipticF[(c + d*x)/2, 2]*Sec[c/2]*Sqrt[Sec[c + d*x]]*Sin[c])/(3*d*(a + a*Cos[c + d*x])^3) + (Cos[c/2 + (d*x)/2]^6*Sqrt[Sec[c + d*x]]*((-2*(-9*A + 39*B + 10*B*Cos[2*c])*Cos[d*x]*Csc[c/2]*Sec[c/2])/(5*d) - (4*Sec[c/2]*Sec[c/2 + (d*x)/2]*(9*A*Sin[(d*x)/2] - 23*B*Sin[(d*x)/2]))/(3*d) + (4*Sec[c/2]*Sec[c/2 + (d*x)/2]^3*(12*A*Sin[(d*x)/2] - 17*B*Sin[(d*x)/2]))/(15*d) - (2*Sec[c/2]*Sec[c/2 + (d*x)/2]^5*(A*Sin[(d*x)/2] - B*Sin[(d*x)/2]))/(5*d) + (16*B*Cos[c]*Sin[d*x])/d - (4*(9*A - 23*B)*Tan[c/2])/(3*d) + (4*(12*A - 17*B)*Sec[c/2 + (d*x)/2]^2*Tan[c/2])/(15*d) - (2*(A - B)*Sec[c/2 + (d*x)/2]^4*Tan[c/2])/(5*d)))/(a + a*Cos[c + d*x])^3
```

3.491.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4508, 3042, 4508, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^3} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}}(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^3} dx$$

↓ 3439

$$\int \frac{A \sec(c + dx) + B}{\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} dx$$

3.491. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A \csc(c+dx+\frac{\pi}{2}) + B}{\sqrt{\csc(c+dx+\frac{\pi}{2})(a \csc(c+dx+\frac{\pi}{2}) + a)^3}} dx \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a(A-11B)-5a(A-B)\sec(c+dx)}{2\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} \\
& \quad \downarrow 4508 \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} - \frac{\int \frac{a(A-11B)-5a(A-B)\sec(c+dx)}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)^2} dx}{10a^2} \\
& \quad \downarrow 27 \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} - \frac{\int \frac{a(A-11B)-5a(A-B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})(\csc(c+dx+\frac{\pi}{2})a+a)^2}} dx}{10a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} - \frac{\int \frac{a^2(6A-41B)-3a^2(3A-8B)\sec(c+dx)}{\sqrt{\sec(c+dx)}(\sec(c+dx)a+a)} dx}{3a^2} - \frac{2a(3A-8B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2} \\
& \quad \downarrow 4508 \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} - \frac{\int \frac{a^2(6A-41B)-3a^2(3A-8B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})(\csc(c+dx+\frac{\pi}{2})a+a)} dx}{3a^2} - \frac{2a(3A-8B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}}{10a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} - \frac{\int \frac{3a^3(9A-49B)-5a^3(3A-13B)\sec(c+dx)}{2\sqrt{\sec(c+dx)}} dx}{a^2} - \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}}{10a^2} \\
& \quad \downarrow 4508 \\
& \frac{(A-B)\sin(c+dx)\sqrt{\sec(c+dx)}}{5d(a \sec(c+dx) + a)^3} - \frac{\int \frac{3a^3(9A-49B)-5a^3(3A-13B)\csc(c+dx+\frac{\pi}{2})}{2\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{a^2} - \frac{5a^2(3A-13B)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B)\sin(c+dx)\sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}}{10a^2} \\
& \quad \downarrow 27
\end{aligned}$$

3.491. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\frac{\int \frac{3a^3(9A-49B) - 5a^3(3A-13B) \sec(c+dx) dx}{\sqrt{\sec(c+dx)}}}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

↓ 3042

$$\frac{\int \frac{3a^3(9A-49B) - 5a^3(3A-13B) \csc(c+dx+\frac{\pi}{2}) dx}{\sqrt{\csc(c+dx+\frac{\pi}{2})}}}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

↓ 4274

$$\frac{3a^3(9A-49B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx - 5a^3(3A-13B) \int \sqrt{\sec(c+dx)} dx}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

↓ 3042

$$\frac{3a^3(9A-49B) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - 5a^3(3A-13B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

↓ 4258

$$\frac{3a^3(9A-49B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - 5a^3(3A-13B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

↓ 3042

$$\frac{3a^3(9A-49B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^3(3A-13B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

↓ 3042

$$\frac{3a^3(9A-49B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - 5a^3(3A-13B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{5a^2(3A-13B) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a \sec(c+dx)+a)} - \frac{2a(3A-8B) \sin(c+dx) \sqrt{\sec(c+dx)}}{3d(a \sec(c+dx)+a)^2}$$

$$\frac{(A-B) \sin(c+dx) \sqrt{\sec(c+dx)}}{5d(a \sec(c+dx)+a)^3} - \frac{10a^2}{3a^2}$$

3.491. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3119} \\
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{5d(a \sec(c + dx) + a)^3} - \\
 & \frac{6a^3(9A - 49B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} - \frac{5a^3(3A - 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx}{2a^2} - \frac{5a^2(3A - 13B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a \sec(c + dx) + a)} - 2 \\
 & \frac{\phantom{6a^3(9A - 49B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}}{2a^2} \qquad \frac{\phantom{5a^3(3A - 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx}}{3a^2} \qquad \frac{\phantom{5a^2(3A - 13B) \sin(c + dx) \sqrt{\sec(c + dx)}}}{10a^2} \\
 & \downarrow \text{3120} \\
 & \frac{(A - B) \sin(c + dx) \sqrt{\sec(c + dx)}}{5d(a \sec(c + dx) + a)^3} - \\
 & \frac{6a^3(9A - 49B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} - \frac{10a^3(3A - 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{5a^2(3A - 13B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a \sec(c + dx) + a)} - 2 \\
 & \frac{\phantom{6a^3(9A - 49B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}}{2a^2} \qquad \frac{\phantom{10a^3(3A - 13B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}}{3a^2} \qquad \frac{\phantom{5a^2(3A - 13B) \sin(c + dx) \sqrt{\sec(c + dx)}}}{10a^2}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sec[c + d*x]^(5/2)),x]`

output `((A - B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/((5*d*(a + a*Sec[c + d*x])^3) - ((-2*a*(3*A - 8*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(3*d*(a + a*Sec[c + d*x])^2) + (((6*a^3*(9*A - 49*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (10*a^3*(3*A - 13*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/(2*a^2) - (5*a^2*(3*A - 13*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*(a + a*Sec[c + d*x]))) / (3*a^2) / (10*a^2)`

3.491.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

$$3.491. \quad \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx$$

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(- (A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.491.4 Maple [A] (verified)

Time = 6.15 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.98

method	result
default	$-\frac{\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(108A\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 30A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3/sec(d*x+c)^(5/2), x, method=_RETURNV ERBOSE)`

$$3.491. \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$$

output

```
-1/60*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(108*A*cos(1/2*d*x+1/2*c)^8+30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5+54*A*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-348*B*cos(1/2*d*x+1/2*c)^8-130*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*cos(1/2*d*x+1/2*c)^5-294*B*cos(1/2*d*x+1/2*c)^5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-198*A*cos(1/2*d*x+1/2*c)^6+578*B*cos(1/2*d*x+1/2*c)^6+114*A*cos(1/2*d*x+1/2*c)^4-264*B*cos(1/2*d*x+1/2*c)^4-27*A*cos(1/2*d*x+1/2*c)^2+37*B*cos(1/2*d*x+1/2*c)^2+3*A-3*B)/a^3/cos(1/2*d*x+1/2*c)^5/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.491.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 478, normalized size of antiderivative = 2.10

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$5 \left(\sqrt{2}(3i A - 13i B) \cos(dx + c) \right)^3 + 3 \sqrt{2}(3i A - 13i B) \cos(dx + c)^2 + 3 \sqrt{2}(3i A - 13i B) \cos(dx + c)$$

input

```
integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(5/2),x, algorithm="fricas")
```


output `-1/60*(5*(sqrt(2)*(3*I*A - 13*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(3*I*A - 13*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(3*I*A - 13*I*B)*cos(d*x + c) + sqrt(2)*(3*I*A - 13*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(-3*I*A + 13*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-3*I*A + 13*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-3*I*A + 13*I*B)*cos(d*x + c) + sqrt(2)*(-3*I*A + 13*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*(sqrt(2)*(9*I*A - 49*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(9*I*A - 49*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(9*I*A - 49*I*B)*cos(d*x + c) + sqrt(2)*(9*I*A - 49*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*(sqrt(2)*(-9*I*A + 49*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-9*I*A + 49*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-9*I*A + 49*I*B)*cos(d*x + c) + sqrt(2)*(-9*I*A + 49*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*(9*A - 29*B)*cos(d*x + c)^3 + 2*(18*A - 73*B)*cos(d*x + c)^2 + 5*(3*A - 13*B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.491.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3/sec(d*x+c)**(5/2),x)`

output `Timed out`

3.491.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2)), x)`

3.491. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

3.491.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(5/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2)), x)`

3.491.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^3),x)`

$$3.492 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$$

3.492.1 Optimal result	4492
3.492.2 Mathematica [C] (verified)	4493
3.492.3 Rubi [A] (verified)	4493
3.492.4 Maple [A] (verified)	4498
3.492.5 Fracas [C] (verification not implemented)	4499
3.492.6 Sympy [F(-1)]	4500
3.492.7 Maxima [F]	4500
3.492.8 Giac [F]	4501
3.492.9 Mupad [F(-1)]	4501

3.492.1 Optimal result

Integrand size = 33, antiderivative size = 259

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx \\ &= \frac{7(7A - 17B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{10a^3 d} \\ & - \frac{(13A - 33B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{6a^3 d} \\ & - \frac{(13A - 33B) \sin(c + dx)}{6a^3 d \sqrt{\sec(c + dx)}} + \frac{(A - B) \sin(c + dx)}{5d \sqrt{\sec(c + dx)} (a + a \sec(c + dx))^3} \\ & + \frac{(A - 2B) \sin(c + dx)}{3ad \sqrt{\sec(c + dx)} (a + a \sec(c + dx))^2} + \frac{7(7A - 17B) \sin(c + dx)}{30d \sqrt{\sec(c + dx)} (a^3 + a^3 \sec(c + dx))} \end{aligned}$$

output

```
-1/6*(13*A-33*B)*sin(d*x+c)/a^3/d/sec(d*x+c)^(1/2)+1/5*(A-B)*sin(d*x+c)/d/
(a+a*sec(d*x+c))^3/sec(d*x+c)^(1/2)+1/3*(A-2*B)*sin(d*x+c)/a/d/(a+a*sec(d*
x+c))^2/sec(d*x+c)^(1/2)+7/30*(7*A-17*B)*sin(d*x+c)/d/(a^3+a^3*sec(d*x+c)
)/sec(d*x+c)^(1/2)+7/10*(7*A-17*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x
+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(
1/2)/a^3/d-1/6*(13*A-33*B)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c
)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/
a^3/d
```

3.492.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 5.18 (sec) , antiderivative size = 589, normalized size of antiderivative = 2.27

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{\cos^6\left(\frac{1}{2}(c + dx)\right) \left(-98\sqrt{2}Ae^{-idx} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \csc(c) \left(-3\sqrt{1+e^{2i(c+dx)}} + e^{2idx}(-1+e^{2ic})\right) \operatorname{Hy}\right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sec[c + d*x]^(7/2)),x]`

output `(Cos[(c + d*x)/2]^6*((-98*sqrt[2]*A*sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) + (238*sqrt[2]*B*sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*sqrt[1 + E^((2*I)*(c + d*x))]*Csc[c]*(-3*sqrt[1 + E^((2*I)*(c + d*x))] + E^((2*I)*d*x)*(-1 + E^((2*I)*c))*Hypergeometric2F1[1/2, 3/4, 7/4, -E^((2*I)*(c + d*x))])/E^(I*d*x) - ((806*A - 1961*B)*Cos[(c - d*x)/2] + (664*A - 1609*B)*Cos[(3*c + d*x)/2] + 470*A*Cos[(c + 3*d*x)/2] - 1165*B*Cos[(c + 3*d*x)/2] + 265*A*Cos[(5*c + 3*d*x)/2] - 620*B*Cos[(5*c + 3*d*x)/2] + 117*A*Cos[(3*c + 5*d*x)/2] - 292*B*Cos[(3*c + 5*d*x)/2] + 30*A*Cos[(7*c + 5*d*x)/2] - 65*B*Cos[(7*c + 5*d*x)/2] - 5*B*Cos[(5*c + 7*d*x)/2] + 5*B*Cos[(9*c + 7*d*x)/2])*Csc[c/2]*Sec[c/2]*Sec[(c + d*x)/2]^5)/(8*sqrt[Sec[c + d*x]]) - 260*A*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]] + 660*B*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])))/(15*a^3*d*(1 + Cos[c + d*x])^3)`

3.492.3 Rubi [A] (verified)

Time = 1.68 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.04, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4508, 27, 3042, 4508, 3042, 4508, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.492. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\sec^{\frac{7}{2}}(c + dx)(a \cos(c + dx) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{A \sec(c + dx) + B}{\sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + a)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A \csc\left(c + dx + \frac{\pi}{2}\right) + B}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + a\right)^3} dx \\
& \quad \downarrow \text{4508} \\
& \frac{\int -\frac{a(3A-13B)-7a(A-B)\sec(c+dx)}{2\sec^{\frac{3}{2}}(c+dx)(\sec(c+dx)a+a)^2} dx}{5a^2} + \frac{(A-B)\sin(c+dx)}{5d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^3} \\
& \quad \downarrow \text{27} \\
& \frac{(A-B)\sin(c+dx)}{5d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^3} - \frac{\int \frac{a(3A-13B)-7a(A-B)\sec(c+dx)}{\sec^{\frac{3}{2}}(c+dx)(\sec(c+dx)a+a)^2} dx}{10a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\sin(c+dx)}{5d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^3} - \frac{\int \frac{a(3A-13B)-7a(A-B)\csc(c+dx+\frac{\pi}{2})}{\csc(c+dx+\frac{\pi}{2})^{3/2}(\csc(c+dx+\frac{\pi}{2})a+a)^2} dx}{10a^2} \\
& \quad \downarrow \text{4508} \\
& \frac{(A-B)\sin(c+dx)}{5d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^3} - \frac{\int \frac{3a^2(8A-23B)-25a^2(A-2B)\sec(c+dx)}{\sec^{\frac{3}{2}}(c+dx)(\sec(c+dx)a+a)} dx}{10a^2} - \frac{10a(A-2B)\sin(c+dx)}{3d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A-B)\sin(c+dx)}{5d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^3} - \frac{\int \frac{3a^2(8A-23B)-25a^2(A-2B)\csc(c+dx+\frac{\pi}{2})}{\csc(c+dx+\frac{\pi}{2})^{3/2}(\csc(c+dx+\frac{\pi}{2})a+a)} dx}{10a^2} - \frac{10a(A-2B)\sin(c+dx)}{3d\sqrt{\sec(c+dx)}(a\sec(c+dx)+a)^2}
\end{aligned}$$

3.492. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 4508 \\
 & \frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \\
 & \frac{\int \frac{3(5a^3(13A - 33B) - 7a^3(7A - 17B) \sec(c + dx))}{2 \sec^{\frac{3}{2}}(c + dx)} dx}{3a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} \\
 & \frac{10a^2}{10a^2} \\
 & \downarrow 27 \\
 & \frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \\
 & \frac{3 \int \frac{5a^3(13A - 33B) - 7a^3(7A - 17B) \sec(c + dx)}{\sec^{\frac{3}{2}}(c + dx)} dx}{3a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} \\
 & \frac{10a^2}{10a^2} \\
 & \downarrow 3042 \\
 & \frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \\
 & \frac{3 \int \frac{5a^3(13A - 33B) - 7a^3(7A - 17B) \csc(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} \\
 & \frac{10a^2}{10a^2} \\
 & \downarrow 4274 \\
 & \frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \\
 & \frac{3 \left(5a^3(13A - 33B) \int \frac{1}{\sec^{\frac{3}{2}}(c + dx)} dx - 7a^3(7A - 17B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} \\
 & \frac{10a^2}{10a^2} \\
 & \downarrow 3042 \\
 & \frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \\
 & \frac{3 \left(5a^3(13A - 33B) \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx - 7a^3(7A - 17B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^2} \\
 & \frac{10a^2}{10a^2} \\
 & \downarrow 4256
 \end{aligned}$$

3.492. $\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx$

$$\frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \frac{3 \left(5a^3(13A - 33B) \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - 7a^3(7A - 17B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)}$$

$10a^2$

↓ 3042

$$\frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \frac{3 \left(5a^3(13A - 33B) \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - 7a^3(7A - 17B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)} - \frac{10a(A - 2B) \sin(c + dx)}{3d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)}$$

$10a^2$

↓ 4258

$$\frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \frac{3 \left(5a^3(13A - 33B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - 7a^3(7A - 17B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)}$$

$3a^2$

$10a^2$

↓ 3042

$$\frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \frac{3 \left(5a^3(13A - 33B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - 7a^3(7A - 17B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)}$$

$3a^2$

$10a^2$

↓ 3119

$$\frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \frac{3 \left(5a^3(13A - 33B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) - \frac{14a^3(7A - 17B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) | 2\right)}{d} \right)}{2a^2} - \frac{7a^2(7A - 17B) \sin(c + dx)}{d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)}$$

$3a^2$

$10a^2$

↓ 3120

3.492. $\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx$

$$\frac{(A - B) \sin(c + dx)}{5d\sqrt{\sec(c + dx)}(a \sec(c + dx) + a)^3} - \frac{3 \left(5a^3(13A - 33B) \left(\frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} \right) - \frac{14a^3(7A - 17B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx), 2\right)}{d} \right)}{2a^2} - \frac{7a^2(7A - 17B)}{d\sqrt{\sec(c+dx)}}$$

$$\frac{\hspace{10em}}{3a^2}$$

$$\frac{\hspace{10em}}{10a^2}$$

```
input Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^3*Sec[c + d*x]^(7/2)),x]
```

```
output ((A - B)*Sin[c + d*x])/(5*d*Sqrt[Sec[c + d*x]]*(a + a*Sec[c + d*x])^3) - ((-10*a*(A - 2*B)*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]*(a + a*Sec[c + d*x])^2) + ((-7*a^2*(7*A - 17*B)*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]]*(a + a*Sec[c + d*x])) + (3*((-14*a^3*(7*A - 17*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 5*a^3*(13*A - 33*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]))))/(2*a^2))/(3*a^2))/(10*a^2)
```

3.492.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

3.492. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^2(c+dx)} dx$

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4508 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^m*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-(A*b - a*B))*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(b*f*(2*m + 1))), x] - Simp[1/(a^2*(2*m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[b*B*n - a*A*(2*m + n + 1) + (A*b - a*B)*(m + n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && EqQ[a^2 - b^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0]`

3.492.4 Maple [A] (verified)

Time = 6.61 (sec) , antiderivative size = 465, normalized size of antiderivative = 1.80

method	result
default	$\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(-160B\left(\cos^{10}\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 348A\left(\cos^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 130A\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^3/sec(d*x+c)^(7/2),x,method=_RETURNV ERBOSE)`

$$3.492. \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^2(c+dx)} dx$$

output $\frac{1}{60} * ((2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) * \sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (-160 * B * \cos(1/2 * d * x + 1/2 * c) ^ 10 + 348 * A * \cos(1/2 * d * x + 1/2 * c) ^ 8 + 130 * A * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (-2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 1) ^ (1/2) * \text{EllipticF}(\cos(1/2 * d * x + 1/2 * c), 2) ^ (1/2)) * \cos(1/2 * d * x + 1/2 * c) ^ 5 + 294 * A * \cos(1/2 * d * x + 1/2 * c) ^ 5 * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (-2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 1) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2) ^ (1/2)) - 468 * B * \cos(1/2 * d * x + 1/2 * c) ^ 8 - 330 * B * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (-2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 1) ^ (1/2) * \text{EllipticF}(\cos(1/2 * d * x + 1/2 * c), 2) ^ (1/2)) * \cos(1/2 * d * x + 1/2 * c) ^ 5 - 714 * B * \cos(1/2 * d * x + 1/2 * c) ^ 5 * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (-2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 1) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2) ^ (1/2)) - 578 * A * \cos(1/2 * d * x + 1/2 * c) ^ 6 + 1058 * B * \cos(1/2 * d * x + 1/2 * c) ^ 6 + 264 * A * \cos(1/2 * d * x + 1/2 * c) ^ 4 - 474 * B * \cos(1/2 * d * x + 1/2 * c) ^ 4 - 37 * A * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 47 * B * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 3 * A - 3 * B) / a ^ 3 / \cos(1/2 * d * x + 1/2 * c) ^ 5 / (-2 * \sin(1/2 * d * x + 1/2 * c) ^ 4 + \sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) / \sin(1/2 * d * x + 1/2 * c) / (2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) / d$

3.492.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 489, normalized size of antiderivative = 1.89

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{5(\sqrt{2}(-13iA + 33iB) \cos(dx + c))^3 + 3\sqrt{2}(-13iA + 33iB) \cos(dx + c)^2 + 3\sqrt{2}(-13iA + 33iB) \cos(dx + c)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(7/2),x, algorithm m="fracas")`

output

```
-1/60*(5*(sqrt(2)*(-13*I*A + 33*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-13*I*A + 33*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-13*I*A + 33*I*B)*cos(d*x + c) + sqrt(2)*(-13*I*A + 33*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*(sqrt(2)*(13*I*A - 33*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(13*I*A - 33*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(13*I*A - 33*I*B)*cos(d*x + c) + sqrt(2)*(13*I*A - 33*I*B))*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*(sqrt(2)*(-7*I*A + 17*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(-7*I*A + 17*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(-7*I*A + 17*I*B)*cos(d*x + c) + sqrt(2)*(-7*I*A + 17*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*(sqrt(2)*(7*I*A - 17*I*B)*cos(d*x + c)^3 + 3*sqrt(2)*(7*I*A - 17*I*B)*cos(d*x + c)^2 + 3*sqrt(2)*(7*I*A - 17*I*B)*cos(d*x + c) + sqrt(2)*(7*I*A - 17*I*B))*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(20*B*cos(d*x + c)^4 - 3*(29*A - 79*B)*cos(d*x + c)^3 - 2*(73*A - 188*B)*cos(d*x + c)^2 - 5*(13*A - 33*B)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)
```

3.492.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**3/sec(d*x+c)**(7/2),x)`

output `Timed out`

3.492.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2)), x)`

3.492. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$

3.492.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^3/sec(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2)), x)`

3.492.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{7/2} (a + a \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^3), x)`

3.493 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$

3.493.1 Optimal result	4502
3.493.2 Mathematica [A] (verified)	4503
3.493.3 Rubi [A] (verified)	4503
3.493.4 Maple [A] (verified)	4506
3.493.5 Fricas [A] (verification not implemented)	4507
3.493.6 Sympy [F(-1)]	4507
3.493.7 Maxima [B] (verification not implemented)	4507
3.493.8 Giac [F(-1)]	4508
3.493.9 Mupad [B] (verification not implemented)	4509

3.493.1 Optimal result

Integrand size = 35, antiderivative size = 220

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$$

$$= \frac{32a(8A + 9B) \sqrt{\sec(c + dx)} \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}} + \frac{16a(8A + 9B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{4a(8A + 9B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{105d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{2a(8A + 9B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{63d \sqrt{a + a \cos(c + dx)}} + \frac{2aA \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{9d \sqrt{a + a \cos(c + dx)}}$$

output

```
16/315*a*(8*A+9*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+4/
105*a*(8*A+9*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/63*
a*(8*A+9*B)*sec(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/9*a*A*s
ec(d*x+c)^(9/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+32/315*a*(8*A+9*B)*sin
(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.493.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.56

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{a(1 + \cos(c + dx))(107A + 81B + 11(8A + 9B) \cos(c + dx) + 11(8A + 9B) \cos(2(c + dx)) + 16A \cos(3(c + dx)) + 18B \cos(3(c + dx)) + 16A \cos(4(c + dx)) + 18B \cos(4(c + dx))) \sec^{\frac{9}{2}}(c + dx) \tan\left(\frac{c + dx}{2}\right)}{315d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*(107*A + 81*B + 11*(8*A + 9*B)*Cos[c + d*x] + 11*(8*A + 9*B)*Cos[2*(c + d*x)] + 16*A*Cos[3*(c + d*x)] + 18*B*Cos[3*(c + d*x)] + 16*A*Cos[4*(c + d*x)] + 18*B*Cos[4*(c + d*x)])*Sec[c + d*x]^(9/2)*Tan[(c + d*x)/2])/(315*d)`

3.493.3 Rubi [A] (verified)Time = 1.17 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{11}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{11/2} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}(A + B \sin\left(c + dx + \frac{\pi}{2}\right)) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)a + a}(A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)^{11/2}} dx$$

3.493. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{9}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{9/2}}dx+\frac{2aA\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{6}{7}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{7}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{6}{7}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2a\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{6}{7}\left(\frac{4}{5}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{6}{7}\left(\frac{4}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{6}{7}\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)+\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{6}{7}\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)+\right)$$

3.493. $\int\sqrt{a+a\cos(c+dx)}(A+B\cos(c+dx))\sec^{\frac{11}{2}}(c+dx)dx$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}(8A+9B)\left(\frac{2a\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{6}{7}\left(\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)*Sqrt[a + a*Cos[c + d*x]]) + ((8*A + 9*B)*((2*a*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) + (6*((2*a*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]))))/5))/7))/9`

3.493.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`


```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```

```
rule 3459 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.493.4 Maple [A] (verified)

Time = 9.80 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.55

method	result
default	$-\frac{2 \cot(dx+c)(\cos(dx+c)-1)((128(\cos^4(dx+c))+64(\cos^3(dx+c))+48(\cos^2(dx+c))+40 \cos(dx+c)+35)A+\cos(dx+c)(144(\cos^3(dx+c)+315d))}{315d}$
parts	$-\frac{2A \cot(dx+c)\left(\sec^{\frac{11}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}(128(\cos^5(dx+c))-64(\cos^4(dx+c))-16(\cos^3(dx+c))-8(\cos^2(dx+c))-5 \cos(dx+c)+45)B*(a*(1+\cos(dx+c)))^{\frac{1}{2}}*\sec(dx+c)^{\frac{11}{2}}}{315d}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)*(a+cos(d*x+c)*a)^(1/2), x, method=_RE
TURNVERBOSE)
```

```
output -2/315/d*cot(d*x+c)*(cos(d*x+c)-1)*((128*cos(d*x+c)^4+64*cos(d*x+c)^3+48*cos
os(d*x+c)^2+40*cos(d*x+c)+35)*A+cos(d*x+c)*(144*cos(d*x+c)^3+72*cos(d*x+c)
^2+54*cos(d*x+c)+45)*B)*(a*(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(11/2)
```

$$3.493. \int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$$

3.493.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.55

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$$

$$= \frac{2(16(8A + 9B) \cos(dx + c)^4 + 8(8A + 9B) \cos(dx + c)^3 + 6(8A + 9B) \cos(dx + c)^2 + 5(8A + 9B) \cos(dx + c) + 35A) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{315(d \cos(dx + c))^5 + d \cos(dx + c)^4} \sqrt{\cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `2/315*(16*(8*A + 9*B)*cos(d*x + c)^4 + 8*(8*A + 9*B)*cos(d*x + c)^3 + 6*(8*A + 9*B)*cos(d*x + c)^2 + 5*(8*A + 9*B)*cos(d*x + c) + 35*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/((d*cos(d*x + c))^5 + d*cos(d*x + c)^4)*sqrt(cos(d*x + c))`

3.493.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(11/2)*(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.493.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 659 vs. $2(190) = 380$.

Time = 0.37 (sec) , antiderivative size = 659, normalized size of antiderivative = 3.00

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$$

$$= \frac{2 \left(A \left(\frac{315 \sqrt{2} \sqrt{a} \sin(dx+c)}{\cos(dx+c)+1} - \frac{735 \sqrt{2} \sqrt{a} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{1302 \sqrt{2} \sqrt{a} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{1206 \sqrt{2} \sqrt{a} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{431 \sqrt{2} \sqrt{a} \sin(dx+c)^9}{(\cos(dx+c)+1)^9} - \frac{107 \sqrt{2} \sqrt{a} \sin(dx+c)^{11}}{(\cos(dx+c)+1)^{11}} \right) \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{11}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{11}{2}} \left(\frac{5 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{10 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{10 \sin(dx+c)^6}{(\cos(dx+c)+1)^6} + \frac{5 \sin(dx+c)^8}{(\cos(dx+c)+1)^8} + \frac{\sin(dx+c)^{10}}{(\cos(dx+c)+1)^{10}} \right)}{\dots}$$

3.493. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `2/315*(A*(315*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 735*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 1302*sqrt(2)*sqrt(a)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 1206*sqrt(2)*sqrt(a)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 431*sqrt(2)*sqrt(a)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 107*sqrt(2)*sqrt(a)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11)*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^5/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(11/2))*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(11/2)*(5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 10*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 5*sin(d*x + c)^8/(cos(d*x + c) + 1)^8 + sin(d*x + c)^10/(cos(d*x + c) + 1)^10 + 1)) + 9*B*(35*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 105*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 154*sqrt(2)*sqrt(a)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 142*sqrt(2)*sqrt(a)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 67*sqrt(2)*sqrt(a)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 9*sqrt(2)*sqrt(a)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11)*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^5/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(11/2))*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(11/2)*(5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 10*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 5*sin(d*x + c)^8/(cos(d*x + c) + 1)^8 + sin(d*x + c)^10/(cos(d*x + c) + 1)^10 + 1))/d`

3.493.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.493.9 Mupad [B] (verification not implemented)

Time = 6.20 (sec) , antiderivative size = 479, normalized size of antiderivative = 2.18

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$$

$$= \frac{\sqrt{\frac{1}{\frac{e^{-c \cdot 1i - dx \cdot 1i}}{2} + \frac{e^{c \cdot 1i + dx \cdot 1i}}{2}}}}{\left(\frac{\sqrt{a + a \left(\frac{e^{-c \cdot 1i - dx \cdot 1i}}{2} + \frac{e^{c \cdot 1i + dx \cdot 1i}}{2} \right)}}{315d} (256A + 288B) \cdot 1i - \frac{e^{c \cdot 9i + dx \cdot 9i} \sqrt{a + a \left(\frac{e^{-c \cdot 1i - dx \cdot 1i}}{2} + \frac{e^{c \cdot 1i + dx \cdot 1i}}{2} \right)}}{315d} (256A + 288B) \right)}{e^{c \cdot 1i}}$$

```
input int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + a*cos(c + d*x))^(1/2),x)
```

```
output ((1/(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(((a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(256*A + 288*B)*1i)/(315*d) - (exp(c*9i + d*x*9i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(256*A + 288*B)*1i)/(315*d) + (exp(c*2i + d*x*2i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(1152*A + 1296*B)*1i)/(315*d) - (exp(c*7i + d*x*7i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(1152*A + 1296*B)*1i)/(315*d) + (exp(c*4i + d*x*4i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(2016*A + 1008*B)*1i)/(315*d) - (exp(c*5i + d*x*5i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(2016*A + 1008*B)*1i)/(315*d)))/(exp(c*1i + d*x*1i) + 4*exp(c*2i + d*x*2i) + 4*exp(c*3i + d*x*3i) + 6*exp(c*4i + d*x*4i) + 6*exp(c*5i + d*x*5i) + 4*exp(c*6i + d*x*6i) + 4*exp(c*7i + d*x*7i) + exp(c*8i + d*x*8i) + exp(c*9i + d*x*9i) + 1)
```

3.494 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

3.494.1 Optimal result	4510
3.494.2 Mathematica [A] (verified)	4511
3.494.3 Rubi [A] (verified)	4511
3.494.4 Maple [A] (verified)	4514
3.494.5 Fricas [A] (verification not implemented)	4514
3.494.6 Sympy [F(-1)]	4515
3.494.7 Maxima [B] (verification not implemented)	4515
3.494.8 Giac [F(-1)]	4516
3.494.9 Mupad [B] (verification not implemented)	4516

3.494.1 Optimal result

Integrand size = 35, antiderivative size = 175

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{16a(6A + 7B) \sqrt{\sec(c + dx)} \sin(c + dx)}{105d \sqrt{a + a \cos(c + dx)}} + \frac{8a(6A + 7B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{105d \sqrt{a + a \cos(c + dx)}}$$

$$+ \frac{2a(6A + 7B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{35d \sqrt{a + a \cos(c + dx)}} + \frac{2aA \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{7d \sqrt{a + a \cos(c + dx)}}$$

```
output 8/105*a*(6*A+7*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/3
5*a*(6*A+7*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/7*a*A
*sec(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+16/105*a*(6*A+7*B)*s
in(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.494.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.58

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{a(1 + \cos(c + dx))(27A + 14B + 9(6A + 7B) \cos(c + dx) + 2(6A + 7B) \cos(2(c + dx)) + 12A \cos(3(c + dx)))}}{105d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*(27*A + 14*B + 9*(6*A + 7*B)*Cos[c + d*x] + 2*(6*A + 7*B)*Cos[2*(c + d*x)] + 12*A*Cos[3*(c + d*x)] + 14*B*Cos[3*(c + d*x)])*Sec[c + d*x]^(7/2)*Tan[(c + d*x)/2]/(105*d)`

3.494.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{9}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{9/2} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}(A + B \sin\left(c + dx + \frac{\pi}{2}\right)) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)a + a}(A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)^{9/2}} dx$$

$$\downarrow \text{3459}$$

3.494. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{7}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2aA\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\left(\frac{4}{5}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\left(\frac{4}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}(6A+7B)\left(\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{4}{5}\left(\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sin[c + d*x])/(7*d*Cos[c + d
*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) + ((6*A + 7*B)*((2*a*Sin[c + d*x])/(5*
d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x])) + (4*((2*a*Sin[c + d*x])/(3
*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x])) + (4*a*Sin[c + d*x])/(3*d*
Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/5)/7)
```

3.494.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3250 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sq
rt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3251 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e
+ f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Sim
p[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e
+ f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```



```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.494.4 Maple [A] (verified)

Time = 9.62 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.58

method	result
default	$-\frac{2 \cot(dx+c)(\cos(dx+c)-1)((48(\cos^3(dx+c))+24(\cos^2(dx+c))+18 \cos(dx+c)+15)A+\cos(dx+c)(56(\cos^2(dx+c))+28 \cos(dx+c)+21)B)}{105d}$
parts	$-\frac{2A \cot(dx+c)\left(\sec^{\frac{9}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(16(\cos^4(dx+c))-8(\cos^3(dx+c))-2(\cos^2(dx+c))-\cos(dx+c)-5\right)}{35d} + \frac{2B \sin(dx+c)}{d}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+cos(d*x+c)*a)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -2/105/d*cot(d*x+c)*(cos(d*x+c)-1)*((48*cos(d*x+c)^3+24*cos(d*x+c)^2+18*co
s(d*x+c)+15)*A+cos(d*x+c)*(56*cos(d*x+c)^2+28*cos(d*x+c)+21)*B)*(a*(1+cos(
d*x+c)))^(1/2)*sec(d*x+c)^(9/2)
```

3.494.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.59

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{2(8(6A + 7B) \cos(dx + c)^3 + 4(6A + 7B) \cos(dx + c)^2 + 3(6A + 7B) \cos(dx + c) + 15A) \sqrt{a \cos(dx + c)}}{105(d \cos(dx + c)^4 + d \cos(dx + c)^3) \sqrt{\cos(dx + c)}}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+a*cos(d*x+c))^(1/2),x, algo
rithm="fracas")
```

output $2/105*(8*(6*A + 7*B)*\cos(d*x + c)^3 + 4*(6*A + 7*B)*\cos(d*x + c)^2 + 3*(6*A + 7*B)*\cos(d*x + c) + 15*A)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/((d*\cos(d*x + c)^4 + d*\cos(d*x + c)^3)*\sqrt{\cos(d*x + c)})$

3.494.6 Sympy [**F(-1)**]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(9/2)*(a+a*cos(d*x+c))**(1/2),x)`

output Timed out

3.494.7 Maxima [**B**] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 568 vs. $2(151) = 302$.

Time = 0.37 (sec) , antiderivative size = 568, normalized size of antiderivative = 3.25

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{2 \left(3A \left(\frac{35\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{70\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{84\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{58\sqrt{2}\sqrt{a}\sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{9\sqrt{2}\sqrt{a}\sin(dx+c)^9}{(\cos(dx+c)+1)^9} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^4 + 7B \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{9}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{9}{2}} \left(\frac{4\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{6\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{4\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + \frac{\sin(dx+c)^8}{(\cos(dx+c)+1)^8} + 1 \right)}{105d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+a*cos(d*x+c))^(1/2),x, algo rithm="maxima")`

output $2/105*(3*A*(35*\sqrt{2}*\sqrt{a}*\sin(dx + c)/(\cos(dx + c) + 1) - 70*\sqrt{2}*\sqrt{a}*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 84*\sqrt{2}*\sqrt{a}*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 58*\sqrt{2}*\sqrt{a}*\sin(dx + c)^7/(\cos(dx + c) + 1)^7 + 9*\sqrt{2}*\sqrt{a}*\sin(dx + c)^9/(\cos(dx + c) + 1)^9)*(\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 1)^4/((\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{9/2})*(-\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{9/2}*(4*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 6*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + 4*\sin(dx + c)^6/(\cos(dx + c) + 1)^6 + \sin(dx + c)^8/(\cos(dx + c) + 1)^8 + 1)) + 7*B*(15*\sqrt{2}*\sqrt{a}*\sin(dx + c)/(\cos(dx + c) + 1) - 40*\sqrt{2}*\sqrt{a}*\sin(dx + c)^3/(\cos(dx + c) + 1)^3 + 42*\sqrt{2}*\sqrt{a}*\sin(dx + c)^5/(\cos(dx + c) + 1)^5 - 24*\sqrt{2}*\sqrt{a}*\sin(dx + c)^7/(\cos(dx + c) + 1)^7 + 7*\sqrt{2}*\sqrt{a}*\sin(dx + c)^9/(\cos(dx + c) + 1)^9)*(\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 1)^4/((\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{9/2})*(-\sin(dx + c)/(\cos(dx + c) + 1) + 1)^{9/2}*(4*\sin(dx + c)^2/(\cos(dx + c) + 1)^2 + 6*\sin(dx + c)^4/(\cos(dx + c) + 1)^4 + 4*\sin(dx + c)^6/(\cos(dx + c) + 1)^6 + \sin(dx + c)^8/(\cos(dx + c) + 1)^8 + 1)))/d$

3.494.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(dx+c))*sec(dx+c)^(9/2)*(a+a*cos(dx+c))^(1/2),x, algorith="giac")`

output `Timed out`

3.494.9 Mupad [B] (verification not implemented)

Time = 4.82 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.52

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{\sqrt{\frac{1}{\frac{e^{-c li - dx li}}{2} + \frac{e^{c li + dx li}}{2}}}}{\left(\frac{\sqrt{a + a \left(\frac{e^{-c li - dx li}}{2} + \frac{e^{c li + dx li}}{2} \right)} (96 A + 112 B) li}{105 d} - \frac{e^{c 7i + dx 7i} \sqrt{a + a \left(\frac{e^{-c li - dx li}}{2} + \frac{e^{c li + dx li}}{2} \right)} (96 A + 112 B) li}{105 d} \right)}$$

3.494. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + a*cos(c + d*x))^(1/2),x)`

output `((1/(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(96*A + 112*B)*1i)/(105*d) - (exp(c*7i + d*x*7i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(96*A + 112*B)*1i)/(105*d) + (exp(c*2i + d*x*2i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(336*A + 392*B)*1i)/(105*d) - (exp(c*5i + d*x*5i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(336*A + 392*B)*1i)/(105*d) - (B*exp(c*3i + d*x*3i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*8i)/(3*d) + (B*exp(c*4i + d*x*4i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*8i)/(3*d)))/(exp(c*1i + d*x*1i) + 3*exp(c*2i + d*x*2i) + 3*exp(c*3i + d*x*3i) + 3*exp(c*4i + d*x*4i) + 3*exp(c*5i + d*x*5i) + exp(c*6i + d*x*6i) + exp(c*7i + d*x*7i) + 1)`

3.495 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.495.1 Optimal result	4518
3.495.2 Mathematica [A] (verified)	4518
3.495.3 Rubi [A] (verified)	4519
3.495.4 Maple [A] (verified)	4521
3.495.5 Fricas [A] (verification not implemented)	4522
3.495.6 Sympy [F(-1)]	4522
3.495.7 Maxima [B] (verification not implemented)	4523
3.495.8 Giac [F(-1)]	4523
3.495.9 Mupad [B] (verification not implemented)	4524

3.495.1 Optimal result

Integrand size = 35, antiderivative size = 130

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{4a(4A + 5B)\sqrt{\sec(c + dx)} \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{2a(4A + 5B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{2aA \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{5d\sqrt{a + a \cos(c + dx)}}$$

output

```
2/15*a*(4*A+5*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/5*
a*A*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+4/15*a*(4*A+5*B)*
sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.495.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.60

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{a(1 + \cos(c + dx))}(7A + 5B + (4A + 5B) \cos(c + dx) + (4A + 5B) \cos(2(c + dx))) \sec^{\frac{5}{2}}(c + dx) \tan}{15d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*Sqrt[a*(1 + Cos[c + d*x])]*(7*A + 5*B + (4*A + 5*B)*Cos[c + d*x] + (4*A + 5*B)*Cos[2*(c + d*x)])*Sec[c + d*x]^(5/2)*Tan[(c + d*x)/2])/(15*d)`

3.495.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.15, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3440, 3042, 3459, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{7}{2}}(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)a + a} (A + B \cos(c + dx))}{\cos^{\frac{7}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)^{7/2}} dx \\
 & \quad \downarrow \text{3459} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} (4A + 5B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{5}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} (4A + 5B) \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}}{\sin\left(c + dx + \frac{\pi}{2}\right)^{5/2}} dx + \frac{2aA \sin(c + dx)}{5d \cos^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right)
 \end{aligned}$$

3.495. $\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}(4A+5B)\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}(4A+5B)\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)+\right.$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}(4A+5B)\left(\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{4a\sin(c+dx)}{3d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)+\right.$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) + ((4*A + 5*B)*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/5`

3.495.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3251 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*(x_)^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

3.495.4 Maple [A] (verified)

Time = 9.73 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.65

method	result
default	$-\frac{2 \cot(dx+c) \left(\sec^{\frac{7}{2}}(dx+c) \right) (\cos(dx+c)-1) (8A \cos^2(dx+c) + 10B \cos^2(dx+c) + 4A \cos(dx+c) + 5B \cos(dx+c) + 3A) \sqrt{a(1+\cos(dx+c))}}{15d}$
parts	$-\frac{2A \cot(dx+c) \left(\sec^{\frac{7}{2}}(dx+c) \right) \sqrt{a(1+\cos(dx+c))} (8 \cos^3(dx+c) - 4 \cos^2(dx+c) - \cos(dx+c) - 3)}{15d} - \frac{2B \cos(dx+c) \cot(dx+c) \left(\sec^{\frac{7}{2}}(dx+c) \right) \sqrt{a(1+\cos(dx+c))}}{15d}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+cos(d*x+c)*a)^(1/2), x, method=_RETURNVERBOSE)
```

$$3.495. \quad \int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

output
$$-2/15/d*\cot(d*x+c)*\sec(d*x+c)^{(7/2)}*(\cos(d*x+c)-1)*(8*A*\cos(d*x+c)^2+10*B*\cos(d*x+c)^2+4*A*\cos(d*x+c)+5*B*\cos(d*x+c)+3*A)*(a*(1+\cos(d*x+c)))^{(1/2)}$$

3.495.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.66

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{2(2(4A + 5B) \cos(dx + c)^2 + (4A + 5B) \cos(dx + c) + 3A) \sqrt{a \cos(dx + c) + a \sin(dx + c)}}{15(d \cos(dx + c)^3 + d \cos(dx + c)^2) \sqrt{\cos(dx + c)}}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output
$$2/15*(2*(4*A + 5*B)*\cos(d*x + c)^2 + (4*A + 5*B)*\cos(d*x + c) + 3*A)*\sqrt{a*\cos(d*x + c) + a*\sin(d*x + c)}/((d*\cos(d*x + c))^3 + d*\cos(d*x + c)^2)*\sqrt{\cos(d*x + c)}$$

3.495.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(7/2)*(a+a*cos(d*x+c))**(1/2),x)`

output Timed out

3.495.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 475 vs. $2(112) = 224$.

Time = 0.35 (sec) , antiderivative size = 475, normalized size of antiderivative = 3.65

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{2 \left(A \left(\frac{15 \sqrt{2} \sqrt{a} \sin(dx+c)}{\cos(dx+c)+1} - \frac{25 \sqrt{2} \sqrt{a} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{17 \sqrt{2} \sqrt{a} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{7 \sqrt{2} \sqrt{a} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^3 + 5B \left(\frac{3 \sqrt{2} \sqrt{a} \sin(dx+c)}{\cos(dx+c)+1} - \frac{7 \sqrt{2} \sqrt{a} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{17 \sqrt{2} \sqrt{a} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{7 \sqrt{2} \sqrt{a} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^3 \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(\frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + 1 \right)} + \frac{5B \left(\frac{3 \sqrt{2} \sqrt{a} \sin(dx+c)}{\cos(dx+c)+1} - \frac{7 \sqrt{2} \sqrt{a} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{17 \sqrt{2} \sqrt{a} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{7 \sqrt{2} \sqrt{a} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^3}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{7}{2}} \left(\frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + 1 \right)}$$

15 d

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+a*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `2/15*(A*(15*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 25*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 17*sqrt(2)*sqrt(a)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 7*sqrt(2)*sqrt(a)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^3/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1)) + 5*B*(3*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 7*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 5*sqrt(2)*sqrt(a)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - sqrt(2)*sqrt(a)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7)*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^3/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(7/2)*(3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1)))/d`

3.495.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+a*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output Timed out

3.495. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.495.9 Mupad [B] (verification not implemented)

Time = 2.82 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.51

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{4 \sqrt{a (\cos(c + dx) + 1)} \sqrt{\frac{1}{\cos(c + dx)}} (14 A \sin(c + dx) + 10 B \sin(c + dx) + 8 A \sin(2c + 2dx) + 18 A \sin(3c + 3dx) + 4 A \sin(4c + 4dx) + 4 A \sin(5c + 5dx) + 10 B \sin(2c + 2dx) + 15 B \sin(3c + 3dx) + 5 B \sin(4c + 4dx) + 5 B \sin(5c + 5dx))}{15 d (10 \cos(c + dx) + 8 \cos(2c + 2dx) + 5 \cos(3c + 3dx) + 2 \cos(4c + 4dx) + \cos(5c + 5dx) + 6)}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^(1/2),x)`

output `(4*(a*(cos(c + d*x) + 1))^(1/2)*(1/cos(c + d*x))^(1/2)*(14*A*sin(c + d*x) + 10*B*sin(c + d*x) + 8*A*sin(2*c + 2*d*x) + 18*A*sin(3*c + 3*d*x) + 4*A*sin(4*c + 4*d*x) + 4*A*sin(5*c + 5*d*x) + 10*B*sin(2*c + 2*d*x) + 15*B*sin(3*c + 3*d*x) + 5*B*sin(4*c + 4*d*x) + 5*B*sin(5*c + 5*d*x)))/(15*d*(10*cos(c + d*x) + 8*cos(2*c + 2*d*x) + 5*cos(3*c + 3*d*x) + 2*cos(4*c + 4*d*x) + cos(5*c + 5*d*x) + 6))`

3.496 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.496.1 Optimal result	4525
3.496.2 Mathematica [A] (verified)	4525
3.496.3 Rubi [A] (verified)	4526
3.496.4 Maple [A] (verified)	4528
3.496.5 Fricas [A] (verification not implemented)	4528
3.496.6 Sympy [F(-1)]	4529
3.496.7 Maxima [B] (verification not implemented)	4529
3.496.8 Giac [F(-1)]	4530
3.496.9 Mupad [B] (verification not implemented)	4530

3.496.1 Optimal result

Integrand size = 35, antiderivative size = 85

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2a(2A + 3B)\sqrt{\sec(c + dx)} \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}} + \frac{2aA \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}}$$

output `2/3*a*A*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/3*a*(2*A+3*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.496.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{a(1 + \cos(c + dx))}(A + (2A + 3B) \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) \tan\left(\frac{1}{2}(c + dx)\right)}{3d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output $(2\sqrt{a(1 + \cos[c + dx])} * (A + (2A + 3B) * \cos[c + dx]) * \sec[c + dx]^{\frac{5}{2}} * \tan[(c + dx)/2]) / (3d)$

3.496.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.25, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3042, 3440, 3042, 3459, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{\frac{5}{2}}(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)a + a} (A + B \cos(c + dx))}{\cos^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{3459} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{3} (2A + 3B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{\frac{3}{2}}(c + dx)} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{3} (2A + 3B) \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}}} dx + \frac{2aA \sin(c + dx)}{3d \cos^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} \right) \\
 & \quad \downarrow \text{3250}
 \end{aligned}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2a(2A+3B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}+\frac{2aA\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (2*a*(2*A + 3*B)*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])`

3.496.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.496.4 Maple [A] (verified)

Time = 9.58 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

method	result
default	$-\frac{2 \cot(dx+c) \left(\sec^{\frac{5}{2}}(dx+c)\right) (\cos(dx+c)-1)(2A \cos(dx+c)+3B \cos(dx+c)+A) \sqrt{a(1+\cos(dx+c))}}{3d}$
parts	$-\frac{2A \cot(dx+c) \left(\sec^{\frac{5}{2}}(dx+c)\right) \sqrt{a(1+\cos(dx+c))} (2(\cos^2(dx+c))-\cos(dx+c)-1)}{3d} - \frac{2B \cos(dx+c) \cot(dx+c) \left(\sec^{\frac{5}{2}}(dx+c)\right) \sqrt{a(1+\cos(dx+c))}}{d}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output
$$-2/3/d*\cot(d*x+c)*\sec(d*x+c)^(5/2)*(\cos(d*x+c)-1)*(2*A*\cos(d*x+c)+3*B*\cos(d*x+c)+A)*(a*(1+\cos(d*x+c)))^(1/2)$$

3.496.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.76

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2((2A + 3B) \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a \sin(dx + c)}}{3(d \cos(dx + c)^2 + d \cos(dx + c)) \sqrt{\cos(dx + c)}}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output
$$2/3*((2*A + 3*B)*\cos(d*x + c) + A)*\sqrt{a*\cos(d*x + c) + a*\sin(d*x + c)}/((d*\cos(d*x + c)^2 + d*\cos(d*x + c))*\sqrt{\cos(d*x + c)})$$

3.496.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)*(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.496.7 Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 380 vs. $2(73) = 146$.

Time = 0.35 (sec) , antiderivative size = 380, normalized size of antiderivative = 4.47

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2 \left(A \left(\frac{3\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{4\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^2 + 3B \left(\frac{\sqrt{2}\sqrt{a}\sin(dx+c)}{\cos(dx+c)+1} - \frac{2\sqrt{2}\sqrt{a}\sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{\sqrt{2}\sqrt{a}\sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right) \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{5}{2}} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{\frac{5}{2}} \left(\frac{2\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)}{3d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+a*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `2/3*(A*(3*sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 4*sqrt(2)*sqrt
(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + sqrt(2)*sqrt(a)*sin(d*x + c)^5/
cos(d*x + c) + 1)^5*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^2/((sin(d*x
+ c)/(cos(d*x + c) + 1) + 1)^(5/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)
^(5/2)*(2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + sin(d*x + c)^4/(cos(d*x +
c) + 1)^4 + 1)) + 3*B*(sqrt(2)*sqrt(a)*sin(d*x + c)/(cos(d*x + c) + 1) - 2
*sqrt(2)*sqrt(a)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + sqrt(2)*sqrt(a)*sin
(d*x + c)^5/(cos(d*x + c) + 1)^5*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1
)^2/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(5/2)*(-sin(d*x + c)/(cos(d*x +
c) + 1) + 1)^(5/2)*(2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + sin(d*x + c)^
4/(cos(d*x + c) + 1)^4 + 1)))/d`

3.496.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.496.9 Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.34

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2 \sqrt{a (\cos(c + dx) + 1)} \sqrt{\frac{1}{\cos(c + dx)}} (2 A \sin(c + dx) + 3 B \sin(c + dx) + 2 A \sin(2c + 2dx) + 2 A \sin(3c + 3dx))}{3d (3 \cos(c + dx) + 2 \cos(2c + 2dx) + \cos(3c + 3dx) + 2)}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(1/2),x)`

output `(2*(a*(cos(c + d*x) + 1))^(1/2)*(1/cos(c + d*x))^(1/2)*(2*A*sin(c + d*x) + 3*B*sin(c + d*x) + 2*A*sin(2*c + 2*d*x) + 2*A*sin(3*c + 3*d*x) + 3*B*sin(3*c + 3*d*x)))/(3*d*(3*cos(c + d*x) + 2*cos(2*c + 2*d*x) + cos(3*c + 3*d*x) + 2))`

3.497 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.497.1 Optimal result	4531
3.497.2 Mathematica [A] (verified)	4531
3.497.3 Rubi [A] (verified)	4532
3.497.4 Maple [A] (verified)	4534
3.497.5 Fricas [A] (verification not implemented)	4535
3.497.6 Sympy [F(-1)]	4535
3.497.7 Maxima [B] (verification not implemented)	4535
3.497.8 Giac [F]	4536
3.497.9 Mupad [F(-1)]	4537

3.497.1 Optimal result

Integrand size = 35, antiderivative size = 96

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{a}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} + \frac{2aA \sqrt{\sec(c + dx)} \sin(c + dx)}{d \sqrt{a + a \cos(c + dx)}}$$

output `2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2*a*A*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.497.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.90

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(\sqrt{2}B \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) \sqrt{\cos(c + dx)} + 2A}{d}$$

input `Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(Sqrt[2]*B *ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Sqrt[Cos[c + d*x]] + 2*A*Sin[(c + d*x)/2]))/d`

3.497.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3440, 3042, 3459, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)a + a} (A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sin\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}}} dx \\ & \quad \downarrow \text{3459} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(B \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx + \frac{2aA \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} \right) \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(B \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2aA \sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}} \right) \end{aligned}$$

3.497. $\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{2B\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} + \frac{2\sqrt{a}B\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}\right)$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*Sqrt[a]*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (2*a*A*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))`

3.497.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^n*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```

```
rule 3459 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.497.4 Maple [A] (verified)

Time = 9.93 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.25

method	result
parts	$-\frac{2A \cot(dx+c) \left(\sec^{\frac{3}{2}}(dx+c)\right) (\cos(dx+c)-1) \sqrt{a(1+\cos(dx+c))}}{d} + \frac{2B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \cos(dx+c) \left(\sec^{\frac{3}{2}}(dx+c)\right) \sqrt{a(1+\cos(dx+c))}}{d}$
default	$\frac{2 \left(\sec^{\frac{3}{2}}(dx+c)\right) \left(B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \cos(dx+c) + B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)\right)}{d(1+\cos(dx+c))}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+cos(d*x+c)*a)^(1/2), x, method=_RET
URNVERBOSE)
```

```
output -2*A/d*cot(d*x+c)*sec(d*x+c)^(3/2)*(cos(d*x+c)-1)*(a*(1+cos(d*x+c)))^(1/2)
+2*B/d*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*sec(d*x+c)^(3/2)*(a*(1
+cos(d*x+c)))^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))
```

3.497.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.95

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= -\frac{2 \left((B \cos(dx + c) + B) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - \frac{\sqrt{a \cos(dx+c)+a} A \sin(dx+c)}{\sqrt{\cos(dx+c)}} \right)}{d \cos(dx + c) + d}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(1/2),x, algo
rithm="fracas")`

output `-2*((B*cos(d*x + c) + B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(
d*x + c))/(sqrt(a)*sin(d*x + c))) - sqrt(a*cos(d*x + c) + a)*A*sin(d*x + c
)/sqrt(cos(d*x + c)))/(d*cos(d*x + c) + d)`

3.497.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)*(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.497.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 906 vs. $2(82) = 164$.

Time = 0.47 (sec) , antiderivative size = 906, normalized size of antiderivative = 9.44

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

3.497. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

output `1/2*(B*sqrt(a)*(arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) - 1) - arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2...`

3.497.8 Giac [F]

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.497.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} \sqrt{a + a \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(1/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(1/2), x)`

3.498 $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)}$

3.498.1 Optimal result	4538
3.498.2 Mathematica [A] (verified)	4538
3.498.3 Rubi [A] (verified)	4539
3.498.4 Maple [A] (verified)	4541
3.498.5 Fricas [A] (verification not implemented)	4541
3.498.6 Sympy [F]	4542
3.498.7 Maxima [B] (verification not implemented)	4542
3.498.8 Giac [F(-1)]	4543
3.498.9 Mupad [F(-1)]	4544

3.498.1 Optimal result

Integrand size = 35, antiderivative size = 98

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{\sqrt{a}(2A + B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{d} + \frac{aB \sin(c + dx)}{d\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)}}$$

output

```
a*B*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+(2*A+B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.498.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.05

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{\sqrt{\cos(c + dx)}\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)}\left(\sqrt{2}(2A + B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right)\right)}{2d}$$

input

```
Integrate[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]
```

output $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*\text{Sec}[(c + d*x)/2]*\text{Sqrt}[\text{Sec}[c + d*x]]*(\text{Sqrt}[2]*(2*A + B)*\text{ArcSin}[\text{Sqrt}[2]*\text{Sin}[(c + d*x)/2]] + 2*B*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[(c + d*x)/2]))/(2*d)$

3.498.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3440, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c+dx)} \sqrt{a \cos(c+dx) + a} (A + B \cos(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \sqrt{a \sin\left(c+dx+\frac{\pi}{2}\right) + a} \left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

$$\downarrow 3440$$

$$\sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{\cos(c+dx)a + a} (A + B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx$$

$$\downarrow 3042$$

$$\sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a + a} \left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx$$

$$\downarrow 3460$$

$$\sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{1}{2}(2A+B) \int \frac{\sqrt{\cos(c+dx)a + a}}{\sqrt{\cos(c+dx)}} dx + \frac{aB \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx) + a}} \right)$$

$$\downarrow 3042$$

$$\sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{1}{2}(2A+B) \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a + a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{aB \sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{a \cos(c+dx) + a}} \right)$$

$$\downarrow 3253$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{aB\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} - \frac{(2A+B)\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{\sqrt{a}(2A+B)\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} + \frac{aB\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)$$

input `Int[Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((Sqrt[a]*(2*A + B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))`

3.498.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m*Int[(a + b*Sin[e + f*x])^n*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.498.4 Maple [A] (verified)

Time = 22.68 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.50

method	result
default	$\frac{\left(B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 2A \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) + B \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right) (\sqrt{\sec(dx+c)}) \sqrt{a(1+\cos(dx+c))}}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{2A \sqrt{a(1+\cos(dx+c))} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) (\sqrt{\sec(dx+c)}) \cos(dx+c)}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}} + \frac{B \left(\sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right) \sqrt{a(1+\cos(dx+c))}}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((A+B*cos(d*x+c))*(a+cos(d*x+c)*a)^(1/2)*sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 1/d*(B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2*A*arctan(tan(d*x+c)*
(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)))*sec(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)/(1+c
os(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

3.498.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.99

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{\sqrt{a \cos(dx + c) + aB} \sqrt{\cos(dx + c)} \sin(dx + c) - ((2A + B) \cos(dx + c) + 2A + B) \sqrt{a} \arctan\left(\sqrt{\frac{a \cos(dx + c)}{1 + \cos(dx + c)}}\right)}{d \cos(dx + c) + d}$$

```
input integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)*sec(d*x+c)^(1/2),x, algo
rithm="fracas")
```

3.498. $\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

output $(\sqrt{a \cos(dx + c) + a} * B * \sqrt{\cos(dx + c)} * \sin(dx + c) - ((2 * A + B) * \cos(dx + c) + 2 * A + B) * \sqrt{a} * \arctan(\sqrt{a \cos(dx + c) + a} * \sqrt{\cos(dx + c)}) / (\sqrt{a} * \sin(dx + c))) / (d * \cos(dx + c) + d)$

3.498.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int \sqrt{a (\cos(c + dx) + 1)} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))**(1/2)*sec(d*x+c)**(1/2),x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))*sqrt(sec(c + d*x)), x)`

3.498.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 939 vs. $2(84) = 168$.

Time = 0.48 (sec) , antiderivative size = 939, normalized size of antiderivative = 9.58

$$\int \sqrt{a + a \cos(c + dx)} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)*sec(d*x+c)^(1/2),x, algorith="maxima")`

output `1/4*(4*A*sqrt(a)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + cos(d*x + c)) + (2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (cos(d*x + c) - 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))))*sqrt(a) + sqrt(a)*(arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) + 1) - arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - arctan2((co...`

3.498.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)*sec(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.498.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} \sqrt{a + a \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(1/2), x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(1/2), x)`

$$3.499 \quad \int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.499.1 Optimal result	4545
3.499.2 Mathematica [A] (verified)	4545
3.499.3 Rubi [A] (verified)	4546
3.499.4 Maple [A] (verified)	4549
3.499.5 Fricas [A] (verification not implemented)	4549
3.499.6 Sympy [F]	4550
3.499.7 Maxima [B] (verification not implemented)	4550
3.499.8 Giac [F]	4551
3.499.9 Mupad [F(-1)]	4552

3.499.1 Optimal result

Integrand size = 35, antiderivative size = 151

$$\begin{aligned} & \int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx \\ &= \frac{\sqrt{a}(4A+3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{4d} \\ & \quad + \frac{aB \sin(c+dx)}{2d \sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} + \frac{a(4A+3B) \sin(c+dx)}{4d \sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}} \end{aligned}$$

```
output 1/2*a*B*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/4*a*(4*A+3*
B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/4*(4*A+3*B)*arcs
in(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)*cos(d*x+c)^(1/2)*sec
(d*x+c)^(1/2)/d
```

3.499.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.79

$$\begin{aligned} & \int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx \\ &= \frac{\sqrt{\cos(c+dx)} \sqrt{a(1+\cos(c+dx))} \sec\left(\frac{1}{2}(c+dx)\right) \sqrt{\sec(c+dx)} \left(\sqrt{2}(4A+3B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c+dx)\right)\right)\right)}{8d} \end{aligned}$$

3.499. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

input `Integrate[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]]],x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])] * Sec[(c + d*x)/2] * Sqrt[Sec[c + d*x]] * (Sqrt[2]*(4*A + 3*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(4*A + 3*B + 2*B*Cos[c + d*x])*Sin[(c + d*x)/2]))/(8*d)`

3.499.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3440, 3042, 3460, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

↓ 3042

$$\int \frac{\sqrt{a \sin(c + dx + \frac{\pi}{2}) + a}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + a}(A + B \cos(c + dx)) dx$$

↓ 3042

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{\sin(c + dx + \frac{\pi}{2})a + a}(A + B \sin(c + dx + \frac{\pi}{2})) dx$$

↓ 3460

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{4}(4A + 3B) \int \sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + a} dx + \frac{aB \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{2d \sqrt{a \cos(c + dx) + a}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}(4A+3B)\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}a+adx+\frac{aB\sin(c+dx)\cos(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3249

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}(4A+3B)\left(\frac{1}{2}\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)+\frac{aB\sin(c+dx)\cos(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}(4A+3B)\left(\frac{1}{2}\int\frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)+\frac{aB\sin(c+dx)\cos(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}(4A+3B)\left(\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}-\frac{\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}\right)+\frac{aB\sin(c+dx)\cos(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}(4A+3B)\left(\frac{\sqrt{a}\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)+\frac{aB\sin(c+dx)\cos(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)$$

input `Int[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (((4*A + 3*B)*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)`

3.499.3.1 Defintions of rubi rules used

- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3249 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.499.4 Maple [A] (verified)

Time = 20.26 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.36

method	result
default	$\frac{(2B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 3B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \arctan(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})) \sqrt{a(1+\cos(dx+c))}}{4d(1+\cos(dx+c)) \sqrt{\sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \left(\sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right) \sqrt{a(1+\cos(dx+c))}}{d(1+\cos(dx+c)) \sqrt{\sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}} + \frac{B \left(2 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \dots \right)}{4 \dots}$

```
input int((A+B*cos(d*x+c))*(a+cos(d*x+c)*a)^(1/2)/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/4/d*(2*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+3*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+3*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/sec(d*x+c)^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

3.499.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.84

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx = \frac{((4A+3B) \cos(dx+c) + 4A+3B) \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - \frac{(2B \cos(dx+c)^2 + (4A+3B) \cos(dx+c)) \sqrt{\cos(dx+c)}}{4(d \cos(dx+c) + d)}}{4(d \cos(dx+c) + d)}$$

```
input integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="fracas")
```

```
output -1/4*((4*A + 3*B)*cos(d*x + c) + 4*A + 3*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (2*B*cos(d*x + c)^2 + (4*A + 3*B)*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c))/(d*cos(d*x + c) + d)
```

3.499.6 Sympy [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{\sqrt{a(\cos(c + dx) + 1)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))**(1/2)/sec(d*x+c)**(1/2),x)`

output `Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))/sqrt(sec(c + d*x)), x)`

3.499.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1851 vs. $2(127) = 254$.

Time = 0.55 (sec) , antiderivative size = 1851, normalized size of antiderivative = 12.26

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

```
output 1/16*(4*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) +
1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*
x + c) - (cos(d*x + c) - 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c) + 1))))*sqrt(a) + sqrt(a)*(arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x +
2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c),
cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d
*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)
^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) + 1)))) + 1) - arctan2(-(cos(2*d*x + 2*c)^2 + sin(2
*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(
sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x
+ 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(si
n(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(
2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - arctan2((cos(2*d*x + 2*c)^2 +
sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*
d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)
^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2
*d*x + 2*c) + 1)) + 1) + arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)...
```

3.499.8 Giac [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{a \cos(dx + c) + a}}{\sqrt{\sec(dx + c)}} dx$$

```
input integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algo
rithm="giac")
```

```
output integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)/sqrt(sec(d*x + c))
, x)
```

3.499.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2), x)`

3.500
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.500.1 Optimal result 4553
 3.500.2 Mathematica [A] (verified) 4554
 3.500.3 Rubi [A] (verified) 4554
 3.500.4 Maple [A] (verified) 4557
 3.500.5 Fricas [A] (verification not implemented) 4558
 3.500.6 Sympy [F] 4558
 3.500.7 Maxima [B] (verification not implemented) 4559
 3.500.8 Giac [F] 4559
 3.500.9 Mupad [F(-1)] 4560

3.500.1 Optimal result

Integrand size = 35, antiderivative size = 196

$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{\sqrt{a}(6A+5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{8d}$$

$$+ \frac{aB \sin(c+dx)}{3d \sqrt{a+a \cos(c+dx)} \sec^{\frac{5}{2}}(c+dx)} + \frac{a(6A+5B) \sin(c+dx)}{12d \sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)}$$

$$+ \frac{a(6A+5B) \sin(c+dx)}{8d \sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}}$$

```
output 1/3*a*B*sin(d*x+c)/d/sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+1/12*a*(6*A+5
*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/8*a*(6*A+5*B)*s
in(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/8*(6*A+5*B)*arcsin(s
in(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*a^(1/2)*cos(d*x+c)^(1/2)*sec(d*x
+c)^(1/2)/d
```


3.500.2 Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{\cos(c + dx)}\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(3\sqrt{2}(6A + 5B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right)}{48d}$$

input `Integrate[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])] * Sec[(c + d*x)/2] * Sqrt[Sec[c + d*x]] * (3*Sqrt[2]*(6*A + 5*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(18*A + 19*B + 2*(6*A + 5*B)*Cos[c + d*x] + 4*B*Cos[2*(c + d*x)]) * Sin[(c + d*x)/2]))/(48*d)`

3.500.3 Rubi [A] (verified)Time = 0.97 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.95, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3440, 3042, 3460, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a \cos(c + dx) + a}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \cos^{\frac{3}{2}}(c + dx) \sqrt{\cos(c + dx)a + a}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

3.500. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \int \cos^{3/2}(c+dx) \sqrt{\cos(c+dx)a+adx} + \frac{aB \sin(c+dx) \cos^{5/2}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{aB \sin(c+dx) \cos^{5/2}(c+dx)}{3d\sqrt{a \cos(c+dx)+a}}\right)$$

↓ 3249

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \left(\frac{3}{4} \int \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+adx} + \frac{a \sin(c+dx) \cos^{3/2}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \left(\frac{3}{4} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx} + \frac{a \sin(c+dx) \cos^{3/2}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}\right)\right)$$

↓ 3249

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a+ax}}{\sqrt{\cos(c+dx)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}\right)\right) + \frac{a \sin(c+dx) \cos^{3/2}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+ax}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}}\right)\right) + \frac{a \sin(c+dx) \cos^{3/2}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6}(6A+5B) \left(\frac{3}{4} \left(\frac{a \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+ax}}\right)}{d}\right)\right) + \frac{a \sin(c+dx) \cos^{3/2}(c+dx)}{2d\sqrt{a \cos(c+dx)+a}}\right)$$

↓ 223

3.500. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{3/2}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}(6A+5B)\left(\frac{3}{4}\left(\frac{\sqrt{a}\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)+\frac{a\sin(c+dx)}{2}\right)$$

input `Int[(Sqrt[a + a*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + ((6*A + 5*B)*((a*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)/6)`

3.500.3.1 Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3249 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.500.4 Maple [A] (verified)

Time = 20.12 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.40

method	result
default	$\frac{\sqrt{a(1+\cos(dx+c))} \left(8B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 10B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 18 \tan(dx+c) \right)}{24d(1+\cos(dx+c))}$
parts	$\frac{A \sqrt{a(1+\cos(dx+c))} \left(2 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3 \tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3 \sec(dx+c) \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right)}{4d(1+\cos(dx+c)) \sec(dx+c)^{\frac{3}{2}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}} +$

```
input int((A+B*cos(d*x+c))*(a+cos(d*x+c)*a)^(1/2)/sec(d*x+c)^(3/2), x, method=_RET
URNVERBOSE)
```

```
output 1/24/d*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/sec(d*x+c)^(3/2)/(cos(d*x+c
)/(1+cos(d*x+c)))^(1/2)*(8*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+
c)))^(1/2)+12*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+10*B*sin(d*x+
c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+18*tan(d*x+c)*A*(cos(d*x+c)/(1+cos(d*
x+c)))^(1/2)+15*tan(d*x+c)*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+18*sec(d*x+
c)*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+15*sec(d*x+c)*B*
arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))
```

3.500.
$$\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.500.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.74

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{3((6A + 5B) \cos(dx + c) + 6A + 5B)\sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - (8B \cos(dx+c)^3 + 2(6A+5B) \cos(dx+c)^2 + 3(6A+5B)\cos(dx+c) + 6A + 5B)\sqrt{a} \sin(dx+c)}{24(d \cos(dx + c) + d)}$$

```
input integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algo
rithm="fricas")
```

```
output -1/24*(3*((6*A + 5*B)*cos(d*x + c) + 6*A + 5*B)*sqrt(a)*arctan(sqrt(a*cos(
d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (8*B*cos(d*x +
c)^3 + 2*(6*A + 5*B)*cos(d*x + c)^2 + 3*(6*A + 5*B)*cos(d*x + c)*sqrt(a*c
os(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c) + d)
```

3.500.6 Sympy [F]

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{\sqrt{a(\cos(c + dx) + 1)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

```
input integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))**(1/2)/sec(d*x+c)**(3/2),x)
```

```
output Integral(sqrt(a*(cos(c + d*x) + 1))*(A + B*cos(c + d*x))/sec(c + d*x)**(3/
2), x)
```

3.500.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2981 vs. $2(166) = 332$.

Time = 0.68 (sec) , antiderivative size = 2981, normalized size of antiderivative = 15.21

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algo
rithm="maxima")
```

```
output 1/96*(6*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) +
1)^(1/4))*((cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(2*d*x
+ 2*c) - (cos(2*d*x + 2*c) - 2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d
*x + 2*c)))) + sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*
x + 2*c) + 1)) + ((cos(2*d*x + 2*c) - 2)*cos(1/2*arctan2(sin(2*d*x + 2*c),
cos(2*d*x + 2*c)))) + sin(2*d*x + 2*c)*sin(1/2*arctan2(sin(2*d*x + 2*c), c
os(2*d*x + 2*c))) - cos(2*d*x + 2*c) + 2)*sin(1/2*arctan2(sin(2*d*x + 2*c)
, cos(2*d*x + 2*c) + 1))) * sqrt(a) + 3*sqrt(a)*(arctan2((cos(2*d*x + 2*c)^2
+ sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin
(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*
d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)
)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)
^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(s
in(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c),
cos(2*d*x + 2*c))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) +
1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - arctan2((
cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(c
os(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*
d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos
(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))...
```

3.500.8 Giac [F]

$$\begin{aligned} & \int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A) \sqrt{a \cos(dx + c) + a}}{\sec(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

3.500. $\int \frac{\sqrt{a+a \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

input `integrate((A+B*cos(d*x+c))*(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(a*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.500.9 Mupad **[F(-1)]**

Timed out.

$$\int \frac{\sqrt{a + a \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + a \cos(c + dx)}}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(3/2), x)`

3.501 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{13}{2}}(c+dx) dx$

3.501.1 Optimal result	4561
3.501.2 Mathematica [A] (verified)	4562
3.501.3 Rubi [A] (verified)	4562
3.501.4 Maple [A] (verified)	4566
3.501.5 Fricas [A] (verification not implemented)	4567
3.501.6 Sympy [F(-1)]	4567
3.501.7 Maxima [B] (verification not implemented)	4567
3.501.8 Giac [F(-1)]	4568
3.501.9 Mupad [B] (verification not implemented)	4569

3.501.1 Optimal result

Integrand size = 35, antiderivative size = 275

$$\begin{aligned} & \int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx \\ &= \frac{32a^2(168A + 187B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3465d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{16a^2(168A + 187B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3465d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{4a^2(168A + 187B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{1155d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^2(168A + 187B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{693d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^2(12A + 11B) \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{99d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2aA \sqrt{a + a \cos(c + dx)} \sec^{\frac{11}{2}}(c + dx) \sin(c + dx)}{11d} \end{aligned}$$

output $16/3465*a^2*(168*A+187*B)*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+4/1155*a^2*(168*A+187*B)*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/693*a^2*(168*A+187*B)*\sec(d*x+c)^{(7/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/99*a^2*(12*A+11*B)*\sec(d*x+c)^{(9/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/11*a*A*\sec(d*x+c)^{(11/2)}*\sin(d*x+c)*(a+a*\cos(d*x+c))^{(1/2)}/d+32/3465*a^2*(168*A+187*B)*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/d/(a+a*\cos(d*x+c))^{(1/2)}$

3.501.2 Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.53

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))}(2478A + 2057B + (6342A + 6193B) \cos(c + dx) + 13(168A + 187B) \cos^2(c + dx))}{3465d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(13/2),x]`

output $(a*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*(2478*A + 2057*B + (6342*A + 6193*B)*\text{Cos}[c + d*x] + 13*(168*A + 187*B)*\text{Cos}[2*(c + d*x)] + 2184*A*\text{Cos}[3*(c + d*x)] + 2431*B*\text{Cos}[3*(c + d*x)] + 336*A*\text{Cos}[4*(c + d*x)] + 374*B*\text{Cos}[4*(c + d*x)] + 336*A*\text{Cos}[5*(c + d*x)] + 374*B*\text{Cos}[5*(c + d*x)])*\text{Sec}[c + d*x]^{(11/2)}*\text{Tan}[(c + d*x)/2])/(3465*d)$

3.501.3 Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{13}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx$$

↓ 3042

3.501. $\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \csc\left(c + dx + \frac{\pi}{2}\right)^{13/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{3/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{13/2}} dx \\
& \quad \downarrow \text{3454} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2}{11} \int \frac{\sqrt{\cos(c + dx)a + a} (a(12A + 11B) + a(8A + 11B) \cos(c + dx))}{2 \cos^{11/2}(c + dx)} dx + \frac{2aA \sin(c + dx)}{\cos^{11/2}(c + dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \int \frac{\sqrt{\cos(c + dx)a + a} (a(12A + 11B) + a(8A + 11B) \cos(c + dx))}{\cos^{11/2}(c + dx)} dx + \frac{2aA \sin(c + dx)}{\cos^{11/2}(c + dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a} (a(12A + 11B) + a(8A + 11B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx + \frac{2aA \sin(c + dx + \frac{\pi}{2})}{\sin^{11/2}(c + dx + \frac{\pi}{2})} \right) \\
& \quad \downarrow \text{3459} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \left(\frac{1}{9} a(168A + 187B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{9/2}(c + dx)} dx + \frac{2a^2(12A + 11B) \sin(c + dx)}{9d \cos^{9/2}(c + dx) \sqrt{a \cos(c + dx)}} \right) \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \left(\frac{1}{9} a(168A + 187B) \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a}}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx + \frac{2a^2(12A + 11B) \sin(c + dx + \frac{\pi}{2})}{9d \cos^{9/2}(c + dx) \sqrt{a \cos(c + dx)}} \right) \right) \\
& \quad \downarrow \text{3251} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \left(\frac{1}{9} a(168A + 187B) \left(\frac{6}{7} \int \frac{\sqrt{\cos(c + dx)a + a}}{\cos^{7/2}(c + dx)} dx + \frac{2a \sin(c + dx)}{7d \cos^{7/2}(c + dx) \sqrt{a \cos(c + dx)}} \right) \right) \right)
\end{aligned}$$

3.501. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}a(168A+187B)\left(\frac{6}{7}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2a\sin(c+dx)}{7d\cos^{7/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}a(168A+187B)\left(\frac{6}{7}\left(\frac{4}{5}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{5/2}(c+dx)}dx+\frac{2a\sin(c+dx)}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}a(168A+187B)\left(\frac{6}{7}\left(\frac{4}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a\sin(c+dx)}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}a(168A+187B)\left(\frac{6}{7}\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{3/2}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{3/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}a(168A+187B)\left(\frac{6}{7}\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{3/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{2a^2(12A+11B)\sin(c+dx)}{9d\cos^{9/2}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{1}{9}a(168A+187B)\left(\frac{2a\sin(c+dx)}{7d\cos^{7/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(13/2),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + a*cos[c + d*x]]*Sin
[c + d*x])/(11*d*cos[c + d*x]^(11/2)) + ((2*a^2*(12*A + 11*B)*Sin[c + d*x]
)/(9*d*cos[c + d*x]^(9/2)*Sqrt[a + a*cos[c + d*x]]) + (a*(168*A + 187*B)*(
(2*a*sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)*Sqrt[a + a*cos[c + d*x]]) + (6*
((2*a*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)*Sqrt[a + a*cos[c + d*x]]) + (4
*((2*a*sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)*Sqrt[a + a*cos[c + d*x]]) + (
4*a*sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c + d*x]])))/5)/
7))/9)/11)
```

3.501.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3250 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqr
t[a + b*sin[e + f*x]]*Sqrt[c + d*sin[e + f*x]])), x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3251 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*sin[e
+ f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*sin[e + f*x]])), x] + Sim
p[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*sin[e
+ f*x]]*(c + d*sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*sin[e + f*x])^p Int[(a + b*sin[e + f*x])^m*((c +
d*sin[e + f*x])^n/(g*sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c
- 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.501.4 Maple [A] (verified)

Time = 10.15 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.52

method	result
default	$-\frac{2a \cot(dx+c)(\cos(dx+c)-1)((2688(\cos^5(dx+c))+1344(\cos^4(dx+c))+1008(\cos^3(dx+c))+840(\cos^2(dx+c))+735 \cos(dx+c)+315)A + \cos(dx+c)(2992 \cos^2(dx+c)+1496 \cos(dx+c)+935)B)}{165d}$
parts	$-\frac{2A \cot(dx+c)\sqrt{a(1+\cos(dx+c))} \left(\sec^{\frac{13}{2}}(dx+c)\right) (128(\cos^6(dx+c))-64(\cos^5(dx+c))-16(\cos^4(dx+c))-8(\cos^3(dx+c))-5(\cos^2(dx+c))+3)B}{165d}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x,method=_RE
TURNVERBOSE)
```

```
output -2/3465*a/d*cot(d*x+c)*(cos(d*x+c)-1)*((2688*cos(d*x+c)^5+1344*cos(d*x+c)^
4+1008*cos(d*x+c)^3+840*cos(d*x+c)^2+735*cos(d*x+c)+315)*A+cos(d*x+c)*(299
2*cos(d*x+c)^4+1496*cos(d*x+c)^3+1122*cos(d*x+c)^2+935*cos(d*x+c)+385)*B)*
(a*(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(13/2)
```

$$3.501. \int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx$$

3.501.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.52

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \frac{2(16(168A + 187B)a \cos(dx + c)^5 + 8(168A + 187B)a \cos(dx + c)^4 + 6(168A + 187B)a \cos(dx + c)^3 + 4(168A + 187B)a \cos(dx + c)^2 + 2(168A + 187B)a \cos(dx + c) + 168A + 187B)a \cos(dx + c)^{13/2}}{3465(d \cos(dx + c))^{13/2}}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="fricas")`

output `2/3465*(16*(168*A + 187*B)*a*cos(d*x + c)^5 + 8*(168*A + 187*B)*a*cos(d*x + c)^4 + 6*(168*A + 187*B)*a*cos(d*x + c)^3 + 5*(168*A + 187*B)*a*cos(d*x + c)^2 + 35*(21*A + 11*B)*a*cos(d*x + c) + 315*A*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/((d*cos(d*x + c))^6 + d*cos(d*x + c)^5)*sqrt(cos(d*x + c))`

3.501.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(13/2),x)`

output `Timed out`

3.501.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 712 vs. 2(239) = 478.

Time = 0.35 (sec) , antiderivative size = 712, normalized size of antiderivative = 2.59

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="maxima")`

output `4/3465*(21*(165*sqrt(2)*a^(3/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 495*sqrt(2)*a^(3/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 1056*sqrt(2)*a^(3/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 1254*sqrt(2)*a^(3/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 781*sqrt(2)*a^(3/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 299*sqrt(2)*a^(3/2)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11 + 46*sqrt(2)*a^(3/2)*sin(d*x + c)^13/(cos(d*x + c) + 1)^13)*A*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^5/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(13/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(13/2)*(5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 10*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 5*sin(d*x + c)^8/(cos(d*x + c) + 1)^8 + sin(d*x + c)^10/(cos(d*x + c) + 1)^10 + 1)) + 11*(315*sqrt(2)*a^(3/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 1155*sqrt(2)*a^(3/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 2184*sqrt(2)*a^(3/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 2586*sqrt(2)*a^(3/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 1759*sqrt(2)*a^(3/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 611*sqrt(2)*a^(3/2)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11 + 94*sqrt(2)*a^(3/2)*sin(d*x + c)^13/(cos(d*x + c) + 1)^13)*B*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^5/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(13/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(13/2)*(5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 10*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 5*sin(d*x + c)^8/(cos(d*x + c) + 1)^8 + ...`

3.501.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="giac")`

output `Timed out`

3.501.9 Mupad [B] (verification not implemented)

Time = 5.30 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.27

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx = \frac{\sqrt{\frac{1}{\frac{e^{-c1i-dx1i}}{2} + \frac{e^{c1i+dx1i}}{2}}} \left(-\frac{32ae^{\frac{c11i}{2} + \frac{dx11i}{2}} \sin\left(\frac{c}{2} + \frac{dx}{2}\right) (2A+3B) \sqrt{a+a \cos(c+dx)}}{5d} + \frac{64ae^{\frac{c11i}{2} + \frac{dx11i}{2}} \sin\left(\frac{3c}{2} + \frac{3dx}{2}\right)}{35} \right)}{20e^{\frac{c11i}{2} + \frac{dx11i}{2}} \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + 20e^{\frac{c11i}{2} + \frac{dx11i}{2}} \cos\left(\frac{3c}{2} + \frac{3dx}{2}\right) + 10e^{\frac{c11i}{2} + \frac{dx11i}{2}}}$$

```
input int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(13/2)*(a + a*cos(c + d*x))^(3/2),x)
```

```
output ((1/(exp(-c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((64*a*exp((c*11i)/2 + (d*x*11i)/2)*sin((3*c)/2 + (3*d*x)/2)*(21*A + 19*B)*(a + a*cos(c + d*x))^(1/2))/(35*d) - (32*a*exp((c*11i)/2 + (d*x*11i)/2)*sin(c/2 + (d*x)/2)*(2*A + 3*B)*(a + a*cos(c + d*x))^(1/2))/(5*d) + (32*a*exp((c*11i)/2 + (d*x*11i)/2)*sin((7*c)/2 + (7*d*x)/2)*(168*A + 187*B)*(a + a*cos(c + d*x))^(1/2))/(315*d) + (64*a*exp((c*11i)/2 + (d*x*11i)/2)*sin((11*c)/2 + (11*d*x)/2)*(168*A + 187*B)*(a + a*cos(c + d*x))^(1/2))/(3465*d))/(20*exp((c*11i)/2 + (d*x*11i)/2)*cos(c/2 + (d*x)/2) + 20*exp((c*11i)/2 + (d*x*11i)/2)*cos((3*c)/2 + (3*d*x)/2) + 10*exp((c*11i)/2 + (d*x*11i)/2)*cos((5*c)/2 + (5*d*x)/2) + 10*exp((c*11i)/2 + (d*x*11i)/2)*cos((7*c)/2 + (7*d*x)/2) + 2*exp((c*11i)/2 + (d*x*11i)/2)*cos((9*c)/2 + (9*d*x)/2) + 2*exp((c*11i)/2 + (d*x*11i)/2)*cos((11*c)/2 + (11*d*x)/2))
```


3.502 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$

3.502.1 Optimal result	4570
3.502.2 Mathematica [A] (verified)	4571
3.502.3 Rubi [A] (verified)	4571
3.502.4 Maple [A] (verified)	4575
3.502.5 Fricas [A] (verification not implemented)	4575
3.502.6 Sympy [F(-1)]	4576
3.502.7 Maxima [B] (verification not implemented)	4576
3.502.8 Giac [F(-1)]	4577
3.502.9 Mupad [B] (verification not implemented)	4577

3.502.1 Optimal result

Integrand size = 35, antiderivative size = 228

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx = \frac{16a^2(34A + 39B)\sqrt{\sec(c + dx)} \sin(c + dx)}{315d\sqrt{a + a \cos(c + dx)}} + \frac{8a^2(34A + 39B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{315d\sqrt{a + a \cos(c + dx)}} + \frac{2a^2(34A + 39B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{105d\sqrt{a + a \cos(c + dx)}} + \frac{2a^2(10A + 9B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{63d\sqrt{a + a \cos(c + dx)}} + \frac{2aA\sqrt{a + a \cos(c + dx)} \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{9d}$$

```
output 8/315*a^2*(34*A+39*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
+2/105*a^2*(34*A+39*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
)+2/63*a^2*(10*A+9*B)*sec(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
+2/9*a*A*sec(d*x+c)^(9/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+16/315*a^2*(
34*A+39*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.502.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.54

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))}(376A + 351B + (374A + 324B) \cos(c + dx) + 11(34A + 39B) \cos(2(c + dx)) + 68A \cos(3(c + dx)) + 78B \cos(3(c + dx)) + 68A \cos(4(c + dx)) + 78B \cos(4(c + dx))) \sec^{\frac{9}{2}}(c + dx) \tan^{\frac{1}{2}}(c + dx)}{(315*d)}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*(376*A + 351*B + (374*A + 324*B)*Cos[c + d*x] + 11*(34*A + 39*B)*Cos[2*(c + d*x)] + 68*A*Cos[3*(c + d*x)] + 78*B*Cos[3*(c + d*x)] + 68*A*Cos[4*(c + d*x)] + 78*B*Cos[4*(c + d*x)])*Sec[c + d*x]^(9/2)*Tan[(c + d*x)/2])/(315*d)`

3.502.3 Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{\frac{11}{2}}(c + dx) (a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{11/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{3/2} (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin^{\frac{11}{2}}(c + dx + \frac{\pi}{2})} dx \end{aligned}$$

3.502. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{9}\int\frac{\sqrt{\cos(c+dx)a+a(a(10A+9B)+3a(2A+3B)\cos(c+dx))}}{2\cos^{\frac{9}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{9}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\int\frac{\sqrt{\cos(c+dx)a+a(a(10A+9B)+3a(2A+3B)\cos(c+dx))}}{\cos^{\frac{9}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{9}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a(a(10A+9B)+3a(2A+3B)\sin(c+dx+\frac{\pi}{2}))}}{\sin(c+dx+\frac{\pi}{2})^{9/2}}dx+\frac{2aA\sin(c+dx+\frac{\pi}{2})}{9}\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{3}{7}a(34A+39B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{7}{2}}(c+dx)}dx+\frac{2a^2(10A+9B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{3}{7}a(34A+39B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2a^2(10A+9B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{3}{7}a(34A+39B)\left(\frac{4}{5}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{3}{7}a(34A+39B)\left(\frac{4}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{3}{7}a(34A+39B)\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)\right)$$

3.502. $\int(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{\frac{11}{2}}(c+dx)dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{3}{7}a(34A+39B)\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{3/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{2a^2(10A+9B)\sin(c+dx)}{7d\cos^{7/2}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{3}{7}a(34A+39B)\left(\frac{2a\sin(c+dx)}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((2*a^2*(10*A + 9*B)*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) + (3*a*(34*A + 39*B))*((2*a*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/5)/7)/9)`

3.502.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.502.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(11/2),x)`

output `Timed out`

3.502.7 Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 619 vs. $2(198) = 396$.

Time = 0.37 (sec) , antiderivative size = 619, normalized size of antiderivative = 2.71

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \frac{4 \left(\left(\frac{315 \sqrt{2} a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{840 \sqrt{2} a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{1344 \sqrt{2} a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{1242 \sqrt{2} a^{3/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{517 \sqrt{2} a^{3/2} \sin(dx+c)^9}{(\cos(dx+c)+1)^9} - \frac{94 \sqrt{2} a^{3/2} \sin(dx+c)^{11}}{(\cos(dx+c)+1)^{11}} \right) \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{11/2} \left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{6 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{4 \sin(dx+c)^6}{(\cos(dx+c)+1)^6} + \frac{\sin(dx+c)^8}{(\cos(dx+c)+1)^8} \right)}{\dots}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="maxima")`

output
$$\frac{4}{315} \left(\frac{315 \sqrt{2} a^{3/2} \sin(dx+c)}{(\cos(dx+c)+1)} - 840 \sqrt{2} a^{3/2} \sin(dx+c)^3 / (\cos(dx+c)+1)^3 + 1344 \sqrt{2} a^{3/2} \sin(dx+c)^5 / (\cos(dx+c)+1)^5 - 1242 \sqrt{2} a^{3/2} \sin(dx+c)^7 / (\cos(dx+c)+1)^7 + 517 \sqrt{2} a^{3/2} \sin(dx+c)^9 / (\cos(dx+c)+1)^9 - 94 \sqrt{2} a^{3/2} \sin(dx+c)^{11} / (\cos(dx+c)+1)^{11} \right) A \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 1} \right)^4 \left(\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 6 \sin(dx+c)^4 / (\cos(dx+c)+1)^4 + 4 \sin(dx+c)^6 / (\cos(dx+c)+1)^6 + \sin(dx+c)^8 / (\cos(dx+c)+1)^8 + 1} \right) + 3 \left(\frac{105 \sqrt{2} a^{3/2} \sin(dx+c)}{(\cos(dx+c)+1)} - 350 \sqrt{2} a^{3/2} \sin(dx+c)^3 / (\cos(dx+c)+1)^3 + 518 \sqrt{2} a^{3/2} \sin(dx+c)^5 / (\cos(dx+c)+1)^5 - 444 \sqrt{2} a^{3/2} \sin(dx+c)^7 / (\cos(dx+c)+1)^7 + 209 \sqrt{2} a^{3/2} \sin(dx+c)^9 / (\cos(dx+c)+1)^9 - 38 \sqrt{2} a^{3/2} \sin(dx+c)^{11} / (\cos(dx+c)+1)^{11} \right) B \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 1} \right)^4 \left(\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(\frac{4 \sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 6 \sin(dx+c)^4 / (\cos(dx+c)+1)^4 + 4 \sin(dx+c)^6 / (\cos(dx+c)+1)^6 + \sin(dx+c)^8 / (\cos(dx+c)+1)^8 + 1} \right) \right) / d$$

3.502.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="giac")`

output Timed out

3.502.9 Mupad [B] (verification not implemented)

Time = 5.30 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.39

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \frac{\sqrt{\frac{1}{\frac{e^{-c} \operatorname{li}(-dx) \operatorname{li}^2}{2} + \frac{e^{c} \operatorname{li}(dx) \operatorname{li}^2}{2}}}}} \left(\frac{96 a e^{\frac{c}{2} + \frac{dx}{2}} \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{a + a \cos(c + dx)} (A + B)}{5d} - \frac{16 B a e^{\frac{c}{2} + \frac{dx}{2}} \sin\left(\frac{3c}{2} + \frac{3dx}{2}\right) \sqrt{a + a \cos(c + dx)}}{3d} \right) \frac{1}{12 e^{\frac{c}{2} + \frac{dx}{2}} \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + 8 e^{\frac{c}{2} + \frac{dx}{2}} \cos\left(\frac{3c}{2} + \frac{3dx}{2}\right) + 8 e^{\frac{c}{2} + \frac{dx}{2}}}$$

$$3.502. \quad \int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + a*cos(c + d*x))^(3/2),x)`

output `((1/(exp(- c*i - d*x*i)/2 + exp(c*i + d*x*i)/2))^(1/2)*((96*a*exp((c*9i)/2 + (d*x*9i)/2)*sin(c/2 + (d*x)/2)*(a + a*cos(c + d*x))^(1/2)*(A + B))/(5*d) - (16*B*a*exp((c*9i)/2 + (d*x*9i)/2)*sin((3*c)/2 + (3*d*x)/2)*(a + a*cos(c + d*x))^(1/2))/(3*d) + (16*a*exp((c*9i)/2 + (d*x*9i)/2)*sin((5*c)/2 + (5*d*x)/2)*(34*A + 39*B)*(a + a*cos(c + d*x))^(1/2))/(35*d) + (32*a*exp((c*9i)/2 + (d*x*9i)/2)*sin((9*c)/2 + (9*d*x)/2)*(34*A + 39*B)*(a + a*cos(c + d*x))^(1/2))/(315*d)))/(12*exp((c*9i)/2 + (d*x*9i)/2)*cos(c/2 + (d*x)/2) + 8*exp((c*9i)/2 + (d*x*9i)/2)*cos((3*c)/2 + (3*d*x)/2) + 8*exp((c*9i)/2 + (d*x*9i)/2)*cos((5*c)/2 + (5*d*x)/2) + 2*exp((c*9i)/2 + (d*x*9i)/2)*cos((7*c)/2 + (7*d*x)/2) + 2*exp((c*9i)/2 + (d*x*9i)/2)*cos((9*c)/2 + (9*d*x)/2))`

3.503 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

3.503.1 Optimal result	4579
3.503.2 Mathematica [A] (verified)	4580
3.503.3 Rubi [A] (verified)	4580
3.503.4 Maple [A] (verified)	4583
3.503.5 Fricas [A] (verification not implemented)	4584
3.503.6 Sympy [F(-1)]	4584
3.503.7 Maxima [B] (verification not implemented)	4584
3.503.8 Giac [F(-1)]	4585
3.503.9 Mupad [B] (verification not implemented)	4586

3.503.1 Optimal result

Integrand size = 35, antiderivative size = 181

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \frac{4a^2(52A + 63B)\sqrt{\sec(c + dx)} \sin(c + dx)}{105d\sqrt{a + a \cos(c + dx)}} + \frac{2a^2(52A + 63B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{105d\sqrt{a + a \cos(c + dx)}} + \frac{2a^2(8A + 7B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{35d\sqrt{a + a \cos(c + dx)}} + \frac{2aA\sqrt{a + a \cos(c + dx)} \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{7d}$$

output

```
2/105*a^2*(52*A+63*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)
+2/35*a^2*(8*A+7*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2
/7*a*A*sec(d*x+c)^(7/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+4/105*a^2*(52*
A+63*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.503.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.56

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))}(82A + 63B + 3(78A + 77B) \cos(c + dx) + (52A + 63B) \cos(2(c + dx)) + 105d}{105d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*(82*A + 63*B + 3*(78*A + 77*B)*Cos[c + d*x] + (52*A + 63*B)*Cos[2*(c + d*x)] + 52*A*Cos[3*(c + d*x)] + 63*B*Cos[3*(c + d*x)])*Sec[c + d*x]^(7/2)*Tan[(c + d*x)/2])/(105*d)`

3.503.3 Rubi [A] (verified)Time = 1.08 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.12, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{9/2}(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{9/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{3/2}(A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx \\ & \quad \downarrow \text{3454} \end{aligned}$$

3.503. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{7}\int\frac{\sqrt{\cos(c+dx)a+a}(a(8A+7B)+a(4A+7B)\cos(c+dx))}{2\cos^{\frac{7}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{7d}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{\sqrt{\cos(c+dx)a+a}(a(8A+7B)+a(4A+7B)\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{7d}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a(8A+7B)+a(4A+7B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2aA\sin(c+dx)}{7d}\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}a(52A+63B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}a(52A+63B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}a(52A+63B)\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}a(52A+63B)\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2a^2(8A+7B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{1}{5}a(52A+63B)\left(\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

3.503. $\int(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

input `Int[(a + a*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + a*cos[c + d*x]]*Sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)) + ((2*a^2*(8*A + 7*B)*Sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)*Sqrt[a + a*cos[c + d*x]]) + (a*(52*A + 63*B)*((2*a*Sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)*Sqrt[a + a*cos[c + d*x]]) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c + d*x]])))/5)/7)`

3.503.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

3.503.4 Maple [A] (verified)

Time = 10.20 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.56

method	result
default	$-\frac{2a \cot(dx+c)(\cos(dx+c)-1)((104(\cos^3(dx+c))+52(\cos^2(dx+c))+39 \cos(dx+c)+15)A+\cos(dx+c)(126(\cos^2(dx+c))+63 \cos(dx+c)+21)B)}{105d}$
parts	$-\frac{2A \cot(dx+c)\sqrt{a(1+\cos(dx+c))} \left(\sec^{\frac{9}{2}}(dx+c)\right) (104(\cos^4(dx+c))-52(\cos^3(dx+c))-13(\cos^2(dx+c))-24 \cos(dx+c)-15)a}{105d} + \dots$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2), x, method=_RETURNVERBOSE)
```

```
output -2/105*a/d*cot(d*x+c)*(cos(d*x+c)-1)*((104*cos(d*x+c)^3+52*cos(d*x+c)^2+39*cos(d*x+c)+15)*A+cos(d*x+c)*(126*cos(d*x+c)^2+63*cos(d*x+c)+21)*B)*(a*(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(9/2)
```

3.503. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$

3.503.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.59

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{2(2(52A + 63B)a \cos(dx + c)^3 + (52A + 63B)a \cos(dx + c)^2 + 3(13A + 7B)a \cos(dx + c) + 15A^2a) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{105(d \cos(dx + c)^4 + d \cos(dx + c)^3) \sqrt{\cos(dx + c)}}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorith="fricas")`

output `2/105*(2*(52*A + 63*B)*a*cos(d*x + c)^3 + (52*A + 63*B)*a*cos(d*x + c)^2 + 3*(13*A + 7*B)*a*cos(d*x + c) + 15*A*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/((d*cos(d*x + c)^4 + d*cos(d*x + c)^3)*sqrt(cos(d*x + c)))`

3.503.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(9/2),x)`

output `Timed out`

3.503.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 527 vs. $2(157) = 314$.

Time = 0.37 (sec) , antiderivative size = 527, normalized size of antiderivative = 2.91

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{4 \left(\left(\frac{105 \sqrt{2} a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{245 \sqrt{2} a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{273 \sqrt{2} a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{171 \sqrt{2} a^{3/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{38 \sqrt{2} a^{3/2} \sin(dx+c)^9}{(\cos(dx+c)+1)^9} \right) A \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} \right) + \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(\frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + 1 \right)}{105(d \cos(dx + c)^4 + d \cos(dx + c)^3) \sqrt{\cos(dx + c)}}$$

3.503. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algo
rithm="maxima")`

output `4/105*((105*sqrt(2)*a^(3/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 245*sqrt(2)*
a^(3/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 273*sqrt(2)*a^(3/2)*sin(d*x
+ c)^5/(cos(d*x + c) + 1)^5 - 171*sqrt(2)*a^(3/2)*sin(d*x + c)^7/(cos(d*x
+ c) + 1)^7 + 38*sqrt(2)*a^(3/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9)*A*(s
in(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^3/((sin(d*x + c)/(cos(d*x + c) + 1
) + 1)^(9/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(9/2)*(3*sin(d*x + c)^
2/(cos(d*x + c) + 1)^2 + 3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + sin(d*x +
c)^6/(cos(d*x + c) + 1)^6 + 1)) + 21*(5*sqrt(2)*a^(3/2)*sin(d*x + c)/(cos
(d*x + c) + 1) - 15*sqrt(2)*a^(3/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 +
17*sqrt(2)*a^(3/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 9*sqrt(2)*a^(3/2)
*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 2*sqrt(2)*a^(3/2)*sin(d*x + c)^9/(c
os(d*x + c) + 1)^9)*B*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^3/((sin(d*
x + c)/(cos(d*x + c) + 1) + 1)^(9/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1
)^(9/2)*(3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3*sin(d*x + c)^4/(cos(d*x
+ c) + 1)^4 + sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 1)))/d`

3.503.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algo
rithm="giac")`

output `Timed out`

3.503.9 Mupad [B] (verification not implemented)

Time = 5.15 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.43

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \frac{\sqrt{\frac{1}{\frac{e^{-c1i-dx1i}}{2} + \frac{e^{c1i+dx1i}}{2}}} \left(-\frac{8ae^{\frac{c7i}{2} + \frac{dx7i}{2}} \sin\left(\frac{c}{2} + \frac{dx}{2}\right) (2A+3B) \sqrt{a+a \cos(c+dx)}}{3d} + \frac{16ae^{\frac{c7i}{2} + \frac{dx7i}{2}} \sin\left(\frac{3c}{2} + \frac{3dx}{2}\right) (13A+3B)}{15d} \right)}{6e^{\frac{c7i}{2} + \frac{dx7i}{2}} \cos\left(\frac{c}{2} + \frac{dx}{2}\right) + 6e^{\frac{c7i}{2} + \frac{dx7i}{2}} \cos\left(\frac{3c}{2} + \frac{3dx}{2}\right) + 2e^{\frac{c7i}{2} + \frac{dx7i}{2}} \cos\left(\frac{5c}{2} + \frac{5dx}{2}\right)}$$

```
input int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + a*cos(c + d*x))^(3/2),x)
```

```
output ((1/(exp(-c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((16*a*exp((c*7i)/2 + (d*x*7i)/2)*sin((3*c)/2 + (3*d*x)/2)*(13*A + 12*B)*(a + a*cos(c + d*x))^(1/2))/(15*d) - (8*a*exp((c*7i)/2 + (d*x*7i)/2)*sin(c/2 + (d*x)/2)*(2*A + 3*B)*(a + a*cos(c + d*x))^(1/2))/(3*d) + (8*a*exp((c*7i)/2 + (d*x*7i)/2)*sin((7*c)/2 + (7*d*x)/2)*(52*A + 63*B)*(a + a*cos(c + d*x))^(1/2))/(10*5*d))/(6*exp((c*7i)/2 + (d*x*7i)/2)*cos(c/2 + (d*x)/2) + 6*exp((c*7i)/2 + (d*x*7i)/2)*cos((3*c)/2 + (3*d*x)/2) + 2*exp((c*7i)/2 + (d*x*7i)/2)*cos((5*c)/2 + (5*d*x)/2) + 2*exp((c*7i)/2 + (d*x*7i)/2)*cos((7*c)/2 + (7*d*x)/2))
```

3.504 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$

3.504.1 Optimal result	4587
3.504.2 Mathematica [A] (verified)	4588
3.504.3 Rubi [A] (verified)	4588
3.504.4 Maple [A] (verified)	4591
3.504.5 Fricas [A] (verification not implemented)	4591
3.504.6 Sympy [F(-1)]	4592
3.504.7 Maxima [B] (verification not implemented)	4592
3.504.8 Giac [F(-1)]	4593
3.504.9 Mupad [B] (verification not implemented)	4593

3.504.1 Optimal result

Integrand size = 35, antiderivative size = 134

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{2a^2(18A + 25B)\sqrt{\sec(c + dx)} \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{2a^2(6A + 5B) \sec^{3/2}(c + dx) \sin(c + dx)}{15d\sqrt{a + a \cos(c + dx)}} + \frac{2aA\sqrt{a + a \cos(c + dx)} \sec^{5/2}(c + dx) \sin(c + dx)}{5d}$$

```
output 2/15*a^2*(6*A+5*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/5*a*A*sec(d*x+c)^(5/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+2/15*a^2*(18*A+25*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.504.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.60

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{a\sqrt{a(1 + \cos(c + dx))(24A + 25B + 2(9A + 5B)\cos(c + dx) + (18A + 25B)\cos(2(c + dx)))} \sec^{5/2}(c + dx)}{15d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*(24*A + 25*B + 2*(9*A + 5*B)*Cos[c + d*x] + (18*A + 25*B)*Cos[2*(c + d*x)])*Sec[c + d*x]^(5/2)*Tan[(c + d*x)/2])/(15*d)`

3.504.3 Rubi [A] (verified)Time = 0.91 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.19, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3459, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{7/2}(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{3/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx \\ & \quad \downarrow \text{3454} \end{aligned}$$

3.504. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{5}\int\frac{\sqrt{\cos(c+dx)a+a(a(6A+5B)+a(2A+5B)\cos(c+dx))}}{2\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{5d}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\int\frac{\sqrt{\cos(c+dx)a+a(a(6A+5B)+a(2A+5B)\cos(c+dx))}}{\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)}{5d}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a(a(6A+5B)+a(2A+5B)\sin(c+dx+\frac{\pi}{2}))}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2aA\sin(c+dx)}{5d}\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}a(18A+25B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a^2(6A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}a(18A+25B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a^2(6A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{2a^2(6A+5B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{2a^2(18A+25B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)\right)+\frac{2aA\sin(c+dx)}{5d}$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((2*a^2*(6*A + 5*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (2*a^2*(18*A + 25*B)*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/5)`

3.504.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1) * (b*c + a*d)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]
```

3.504.4 Maple [A] (verified)

Time = 10.14 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.63

method	result
default	$-\frac{2a \cot(dx+c) \left(\sec^{\frac{7}{2}}(dx+c)\right) (\cos(dx+c)-1) (18A(\cos^2(dx+c))+25B(\cos^2(dx+c))+9A \cos(dx+c)+5B \cos(dx+c)+3A) \sqrt{a(1+\cos(dx+c))}}{15d}$
parts	$-\frac{2A \cot(dx+c) \sqrt{a(1+\cos(dx+c))} \left(\sec^{\frac{7}{2}}(dx+c)\right) (6(\cos^3(dx+c))-3(\cos^2(dx+c))-2 \cos(dx+c)-1)a}{5d} - \frac{2B \cos(dx+c) \cot(dx+c)}{5d}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNERVERBOSE)
```

```
output -2/15*a/d*cot(d*x+c)*sec(d*x+c)^(7/2)*(cos(d*x+c)-1)*(18*A*cos(d*x+c)^2+25*B*cos(d*x+c)^2+9*A*cos(d*x+c)+5*B*cos(d*x+c)+3*A)*(a*(1+cos(d*x+c)))^(1/2)
```

3.504.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.66

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \frac{2 \left((18A + 25B)a \cos(dx + c)^2 + (9A + 5B)a \cos(dx + c) + 3Aa \right) \sqrt{a \cos(dx + c) + a \sin(dx + c)}}{15 \left(d \cos(dx + c)^3 + d \cos(dx + c)^2 \right) \sqrt{\cos(dx + c)}}$$

```
input integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="fracas")
```

3.504. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

output $2/15*((18*A + 25*B)*a*\cos(d*x + c)^2 + (9*A + 5*B)*a*\cos(d*x + c) + 3*A*a)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/((d*\cos(d*x + c)^3 + d*\cos(d*x + c)^2)*\sqrt{\cos(d*x + c)})$

3.504.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output Timed out

3.504.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 436 vs. $2(116) = 232$.

Time = 0.36 (sec) , antiderivative size = 436, normalized size of antiderivative = 3.25

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{4 \left(3 \left(\frac{5 \sqrt{2} a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{10 \sqrt{2} a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{7 \sqrt{2} a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{2 \sqrt{2} a^{3/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) A \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right)^2 + 5 \left(\frac{3 \sqrt{2} a^{3/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{10 \sqrt{2} a^{3/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{7 \sqrt{2} a^{3/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{2 \sqrt{2} a^{3/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} \right) \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{7/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{7/2} \left(\frac{2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)}{15d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="maxima")`

output
$$\frac{4}{15} \cdot (3 \cdot (5 \sqrt{2}) \cdot a^{3/2} \cdot \sin(dx + c) / (\cos(dx + c) + 1) - 10 \sqrt{2}) \cdot a^{3/2} \cdot \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + 7 \sqrt{2} \cdot a^{3/2} \cdot \sin(dx + c)^5 / (\cos(dx + c) + 1)^5 - 2 \sqrt{2}) \cdot a^{3/2} \cdot \sin(dx + c)^7 / (\cos(dx + c) + 1)^7) \cdot A \cdot (\sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + 1)^2 / ((\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{7/2} \cdot (-\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{7/2} \cdot (2 \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 + 1)) + 5 \cdot (3 \sqrt{2}) \cdot a^{3/2} \cdot \sin(dx + c) / (\cos(dx + c) + 1) - 8 \sqrt{2}) \cdot a^{3/2} \cdot \sin(dx + c)^3 / (\cos(dx + c) + 1)^3 + 7 \sqrt{2} \cdot a^{3/2} \cdot \sin(dx + c)^5 / (\cos(dx + c) + 1)^5 - 2 \sqrt{2}) \cdot a^{3/2} \cdot \sin(dx + c)^7 / (\cos(dx + c) + 1)^7) \cdot B \cdot (\sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + 1)^2 / ((\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{7/2} \cdot (-\sin(dx + c) / (\cos(dx + c) + 1) + 1)^{7/2} \cdot (2 \sin(dx + c)^2 / (\cos(dx + c) + 1)^2 + \sin(dx + c)^4 / (\cos(dx + c) + 1)^4 + 1))) / d$$

3.504.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="giac")`

output Timed out

3.504.9 Mupad [B] (verification not implemented)

Time = 2.65 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.47

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{2a \sqrt{a (\cos(c + dx) + 1)} \sqrt{\frac{1}{\cos(c + dx)}} (48A \sin(c + dx) + 50B \sin(c + dx) + 36A \sin(2c + 2d + dx))}{15d (10 \cos(c + dx) + 5)}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^(3/2),x)`

3.504. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

output $(2*a*(a*(\cos(c + d*x) + 1))^{1/2}*(1/\cos(c + d*x))^{1/2}*(48*A*\sin(c + d*x) + 50*B*\sin(c + d*x) + 36*A*\sin(2*c + 2*d*x) + 66*A*\sin(3*c + 3*d*x) + 18*A*\sin(4*c + 4*d*x) + 18*A*\sin(5*c + 5*d*x) + 20*B*\sin(2*c + 2*d*x) + 75*B*\sin(3*c + 3*d*x) + 10*B*\sin(4*c + 4*d*x) + 25*B*\sin(5*c + 5*d*x)))/(15*d*(10*\cos(c + d*x) + 8*\cos(2*c + 2*d*x) + 5*\cos(3*c + 3*d*x) + 2*\cos(4*c + 4*d*x) + \cos(5*c + 5*d*x) + 6))$

3.505 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$

3.505.1 Optimal result	4595
3.505.2 Mathematica [A] (verified)	4596
3.505.3 Rubi [A] (verified)	4596
3.505.4 Maple [A] (verified)	4599
3.505.5 Fracas [A] (verification not implemented)	4600
3.505.6 Sympy [F(-1)]	4600
3.505.7 Maxima [B] (verification not implemented)	4600
3.505.8 Giac [F(-1)]	4601
3.505.9 Mupad [F(-1)]	4602

3.505.1 Optimal result

Integrand size = 35, antiderivative size = 145

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \frac{2a^{3/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} + \frac{2a^2(4A + 3B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3d \sqrt{a + a \cos(c + dx)}} + \frac{2aA \sqrt{a + a \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{3d}$$

output

```
2/3*a*A*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+2*a^(3/2)*B*a
rcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+
c)^(1/2)/d+2/3*a^2*(4*A+3*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c)
)^(1/2)
```

3.505.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.73

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \frac{a \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^{3/2}(c + dx) \left(3\sqrt{2}B \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right)\right) \cos^{3/2}(c + dx)}{3d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(a*sqrt[a*(1 + Cos[c + d*x]])*Sec[(c + d*x)/2]*Sec[c + d*x]^(3/2)*(3*sqrt[2]*B*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^(3/2) + 2*(A + (5*A + 3*B)*Cos[c + d*x])*Sin[(c + d*x)/2]))/(3*d)`

3.505.3 Rubi [A] (verified)Time = 0.88 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3459, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{5/2}(c + dx)(a \cos(c + dx) + a)^{3/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{3/2}(A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx \\ & \quad \downarrow \text{3454} \end{aligned}$$

3.505. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a(a(4A+3B)+3aB\cos(c+dx))}}{2\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\int\frac{\sqrt{\cos(c+dx)a+a(a(4A+3B)+3aB\cos(c+dx))}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a(a(4A+3B)+3aB\sin(c+dx+\frac{\pi}{2}))}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(3aB\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(3aB\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}-\frac{6aB\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}\right)+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{6a^{3/2}B\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}+\frac{2a^2(4A+3B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)+\frac{2aA\sin(c+dx)\sqrt{a\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}\right)$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + a*cos[c + d*x]]*Sin
[c + d*x])/(3*d*cos[c + d*x]^(3/2)) + ((6*a^(3/2)*B*ArcSin[(Sqrt[a]*Sin[c
+ d*x])/Sqrt[a + a*cos[c + d*x]])]/d + (2*a^2*(4*A + 3*B)*Sin[c + d*x])/(d
*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c + d*x]]))/3)
```

3.505.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3253 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Co
s[e + f*x]/Sqrt[a + b*sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && E
qQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sine[e + f*x])^p Int[(a + b*sin[e + f*x])^m*((c +
d*sin[e + f*x])^n/(g*Sine[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.505.4 Maple [A] (verified)

Time = 10.32 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.38

method	result
default	$\frac{2a \left(\sec^{\frac{5}{2}}(dx+c)\right) \sqrt{a(1+\cos(dx+c))} \left(3B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) (\cos^3(dx+c)) + 3B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)\right)}{3d(1+\cos(dx+c))}$
parts	$-\frac{2A \cot(dx+c) \sqrt{a(1+\cos(dx+c))} \left(\sec^{\frac{5}{2}}(dx+c)\right) (5(\cos^2(dx+c)) - 4 \cos(dx+c) - 1)a}{3d} + \frac{2B \left(\sec^{\frac{5}{2}}(dx+c)\right) (\cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}})}{3d}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 2/3*a/d*sec(d*x+c)^(5/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))*(3*B*(cos
(d*x+c)/(1+cos(d*x+c)))^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))
)^(1/2))*cos(d*x+c)^3+3*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arctan(tan(d*x
+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)^2+5*A*sin(d*x+c)*cos(d*x
+c)^2+3*B*sin(d*x+c)*cos(d*x+c)^2+A*sin(d*x+c)*cos(d*x+c))
```

$$3.505. \int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

3.505.5 Fricas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.90

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx =$$

$$\frac{2 \left(3 (Ba \cos(dx + c))^2 + Ba \cos(dx + c) \right) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - \frac{((5A+3B)a \cos(dx+c)+Aa) \sqrt{a \cos(dx+c)+a}}{\sqrt{\cos(dx+c)}}}{3 (d \cos(dx + c))^2 + d \cos(dx + c)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="fricas")`

output `-2/3*(3*(B*a*cos(d*x + c)^2 + B*a*cos(d*x + c))*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - ((5*A + 3*B)*a*cos(d*x + c) + A*a)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2 + d*cos(d*x + c))`

3.505.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.505.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1462 vs. 2(123) = 246.

Time = 0.50 (sec) , antiderivative size = 1462, normalized size of antiderivative = 10.08

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algo
rithm="maxima")`

output `1/6*(3*(6*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) +
1)^(3/4)*a^(3/2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))
- 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/
4)*((2*a*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))*sin(2*d*x +
2*c) - a*sin(2*d*x + 2*c) - 2*(a*cos(2*d*x + 2*c) + a)*sin(3/2*arctan2(sin
(2*d*x + 2*c), cos(2*d*x + 2*c))))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2
*d*x + 2*c) + 1)) + (2*a*sin(2*d*x + 2*c)*sin(3/2*arctan2(sin(2*d*x + 2*c)
, cos(2*d*x + 2*c))) + a*cos(2*d*x + 2*c) + 2*(a*cos(2*d*x + 2*c) + a)*cos
(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + a)*sin(3/2*arctan2(sin
(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(a) + ((a*cos(2*d*x + 2*c)^2 +
a*sin(2*d*x + 2*c)^2 + 2*a*cos(2*d*x + 2*c) + a)*arctan2((cos(2*d*x + 2*c)
^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(s
in(2*d*x + 2*c), cos(2*d*x + 2*c)))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(
2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) +
1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*
c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2
(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c)
, cos(2*d*x + 2*c))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
+ 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - (a*cos(
2*d*x + 2*c)^2 + a*sin(2*d*x + 2*c)^2 + 2*a*cos(2*d*x + 2*c) + a)*arcta...`

3.505.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.505.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + a \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(3/2), x)`

3.506 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$

3.506.1 Optimal result	4603
3.506.2 Mathematica [A] (verified)	4604
3.506.3 Rubi [A] (verified)	4604
3.506.4 Maple [B] (verified)	4607
3.506.5 Fricas [A] (verification not implemented)	4608
3.506.6 Sympy [F(-1)]	4608
3.506.7 Maxima [B] (verification not implemented)	4609
3.506.8 Giac [F(-1)]	4609
3.506.9 Mupad [F(-1)]	4610

3.506.1 Optimal result

Integrand size = 35, antiderivative size = 146

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{a^{3/2}(2A + 3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} - \frac{a^2(2A - B) \sin(c + dx)}{d \sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} + \frac{2aA \sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

```
output -a^2*(2*A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+a^(3/2)*
(2*A+3*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/
2)*sec(d*x+c)^(1/2)/d+2*a*A*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)*sec(d*x+c)^(
1/2)/d
```

3.506.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.73

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{a \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(\sqrt{2}(2A + 3B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right) \right)}{2d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(a*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(Sqrt[2]*(2*A + 3*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Sqrt[Cos[c + d*x]] + 2*(2*A + B*Cos[c + d*x])*Sin[(c + d*x)/2]))/(2*d)`

3.506.3 Rubi [A] (verified)Time = 0.91 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{3/2}(c + dx) (a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{3/2} (A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3454} \end{aligned}$$

3.506. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(2\int\frac{\sqrt{\cos(c+dx)a+a(a(2A+B)-a(2A-B)\cos(c+dx))}}{2\sqrt{\cos(c+dx)}}dx+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\int\frac{\sqrt{\cos(c+dx)a+a(a(2A+B)-a(2A-B)\cos(c+dx))}}{\sqrt{\cos(c+dx)}}dx+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a(a(2A+B)-a(2A-B)\sin(c+dx+\frac{\pi}{2}))}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}a(2A+3B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx-\frac{a^2(2A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}a(2A+3B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx-\frac{a^2(2A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{a(2A+3B)\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}-\frac{a^2(2A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{a^{3/2}(2A+3B)\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}-\frac{a^2(2A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a^(3/2)*(2*A + 3*B)*ArcSin[(Sqrt[a
]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d - (a^2*(2*A - B)*Sqrt[Cos[c +
d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]) + (2*a*A*Sqrt[a + a*Cos[
c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])
```

3.506.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3253 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*
(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Co
s[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && E
qQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.506.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 273 vs. 2(128) = 256.

Time = 22.79 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.88

method	result
default	$a \left(\sec^{\frac{3}{2}}(dx+c) \right) \left(2A \cos(dx+c) \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right)$
parts	$\frac{2A \left(\sec^{\frac{3}{2}}(dx+c) \right) \left(\cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) + \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right)}{d(1+\cos(dx+c))}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

3.506. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

output $a/d*\sec(d*x+c)^{(3/2)}*(2*A*\cos(d*x+c)*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c))))^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+3*B*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c))))^{(1/2)}*\cos(d*x+c)+2*A*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c))))^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+B*\sin(d*x+c)*\cos(d*x+c)+3*B*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\arctan(\tan(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c))))^{(1/2)}+2*A*\sin(d*x+c)*\cos(d*x+c)*(a*(1+\cos(d*x+c)))^{(1/2)}/(1+\cos(d*x+c))$

3.506.5 Fricas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.82

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{((2A + 3B)a \cos(dx + c) + (2A + 3B)a)\sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - \frac{(Ba \cos(dx+c)+2Aa)\sqrt{a \cos(dx+c)}}{\sqrt{\cos(dx+c)}}}{d \cos(dx + c) + d}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo rithm="fricas")`

output $-(((2*A + 3*B)*a*\cos(d*x + c) + (2*A + 3*B)*a)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(sqrt(a)*\sin(d*x + c))) - (B*a*\cos(d*x + c) + 2*A*a)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/\sqrt{\cos(d*x + c)}}/(d*\cos(d*x + c) + d)$

3.506.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.506. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$

3.506.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1801 vs. $2(128) = 256$.

Time = 0.54 (sec) , antiderivative size = 1801, normalized size of antiderivative = 12.34

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `1/4*((2*(a*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (a*cos(d*x + c) - a)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sqrt(a) + 3*(a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) + 1 - a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1 - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + 1) + a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2...`

3.506.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo
rithm="giac")`

3.506. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$

output Timed out

3.506.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + a \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(3/2), x)`

3.507 $\int (a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))\sqrt{\sec(c+dx)}$

3.507.1 Optimal result	4611
3.507.2 Mathematica [A] (verified)	4611
3.507.3 Rubi [A] (verified)	4612
3.507.4 Maple [A] (verified)	4615
3.507.5 Fricas [A] (verification not implemented)	4615
3.507.6 Sympy [F(-1)]	4616
3.507.7 Maxima [B] (verification not implemented)	4616
3.507.8 Giac [F(-1)]	4617
3.507.9 Mupad [F(-1)]	4618

3.507.1 Optimal result

Integrand size = 35, antiderivative size = 153

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx = \frac{a^{3/2}(12A + 7B) \arcsin\left(\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a + a \cos(c + dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{4d} + \frac{a^2(4A + 5B) \sin(c + dx)}{4d\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)}} + \frac{aB\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{2d\sqrt{\sec(c + dx)}}$$

```
output 1/4*a^2*(4*A+5*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/2
*a*B*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)+1/4*a^(3/2)*(12*
A+7*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*
sec(d*x+c)^(1/2)/d
```

3.507.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.79

$$\int (a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx = \frac{a\sqrt{\cos(c + dx)}\sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)}\left(\sqrt{2}(1 + \cos(c + dx))\right)^{3/2}}{2d}$$

input `Integrate[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(a*Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(Sqrt[2]*(12*A + 7*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(4*A + 7*B + 2*B*Cos[c + d*x])*Sin[(c + d*x)/2]))/(8*d)`

3.507.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3455, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c+dx)}(a \cos(c+dx) + a)^{3/2}(A + B \cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a \sin\left(c+dx+\frac{\pi}{2}\right) + a\right)^{3/2}\left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(\cos(c+dx)a + a)^{3/2}(A + B \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(\sin\left(c+dx+\frac{\pi}{2}\right) a + a)^{3/2}\left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3455} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \int \frac{\sqrt{\cos(c+dx)a + a}(a(4A + B) + a(4A + 5B) \cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx + \frac{aB \sin(c+dx)}{\cos(c+dx)}\right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \int \frac{\sqrt{\cos(c+dx)a + a}(a(4A + B) + a(4A + 5B) \cos(c+dx))}{\sqrt{\cos(c+dx)}} dx + \frac{aB \sin(c+dx)}{\cos(c+dx)}\right)
 \end{aligned}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a(a(4A+B)+a(4A+5B)\sin(c+dx+\frac{\pi}{2}))}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{aB}{4}\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}a(12A+7B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{a^2(4A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}a(12A+7B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{a^2(4A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{a^2(4A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}-\frac{a(12A+7B)\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a}{\sqrt{\cos(c+dx)+a}}\right)}{d}\right)\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{a^{3/2}(12A+7B)\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}+\frac{a^2(4A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + ((a^(3/2)*(12*A + 7*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a^2*(4*A + 5*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4`

3.507.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.507.4 Maple [A] (verified)

Time = 22.88 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.39

method	result
default	$\frac{a \left(2B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4A \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 7B \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{4d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \left(\sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 3 \arctan\left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right) (\sqrt{\sec(dx+c)}) \sqrt{a(1+\cos(dx+c))} \cos(dx+c)a}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}} + \frac{B(2 \sin(dx+c))}{d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2), x, method=_RET
URNVERBOSE)
```

```
output 1/4*a/d*(2*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*A*(
cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+7*B*sin(d*x+c)*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)+12*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2
)))+7*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*sec(d*x+c)^(1
/2)*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)
```

3.507.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.87

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \frac{((12A + 7B)a \cos(dx + c) + (12A + 7B)a) \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - \frac{(2Ba \cos(dx+c)^2 + (4A+7B)a)}{4(d \cos(dx+c) + d)}}{4(d \cos(dx+c) + d)}$$

3.507. $\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algo
rithm="fricas")`

output `-1/4*(((12*A + 7*B)*a*cos(d*x + c) + (12*A + 7*B)*a)*sqrt(a)*arctan(sqrt(a
*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (2*B*a*cos
(d*x + c)^2 + (4*A + 7*B)*a*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x
+ c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c) + d)`

3.507.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Timed out`

3.507.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1884 vs. 2(129) = 258.

Time = 0.57 (sec) , antiderivative size = 1884, normalized size of antiderivative = 12.31

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `1/16*(4*(2*(a*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - (a*cos(d*x + c) - a)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sqrt(a) + 3*(a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) + 1) - a*arctan2(-(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(d*x + c) - cos(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(d*x + c)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + sin(d*x + c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))) - 1) - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + 1) + a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x ...`

3.507.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.507.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + a \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(3/2), x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(3/2), x)`

3.508
$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.508.1 Optimal result 4619
 3.508.2 Mathematica [A] (verified) 4620
 3.508.3 Rubi [A] (verified) 4620
 3.508.4 Maple [A] (verified) 4624
 3.508.5 Fricas [A] (verification not implemented) 4624
 3.508.6 Sympy [F(-1)] 4625
 3.508.7 Maxima [B] (verification not implemented) 4625
 3.508.8 Giac [F] 4626
 3.508.9 Mupad [F(-1)] 4627

3.508.1 Optimal result

Integrand size = 35, antiderivative size = 200

$$\int \frac{(a + a \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \frac{a^{3/2}(14A + 11B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}}{8d}$$

$$+ \frac{a^2(6A + 7B) \sin(c + dx)}{12d\sqrt{a + a \cos(c + dx)} \sec^{3/2}(c + dx)}$$

$$+ \frac{aB\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{3d \sec^{3/2}(c + dx)} + \frac{a^2(14A + 11B) \sin(c + dx)}{8d\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}}$$

output

```
1/12*a^2*(6*A+7*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/
3*a*B*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+1/8*a^2*(14*A+1
1*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/8*a^(3/2)*(14*
A+11*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)
*sec(d*x+c)^(1/2)/d
```

3.508.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.70

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \frac{a \sqrt{\cos(c + dx)} \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec\left(\frac{1}{2}(c + dx)\right)}}{\sqrt{\sec(c + dx)}}$$

input `Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]], x]`

output `(a*Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(3*Sqrt[2]*(14*A + 11*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(42*A + 37*B + 2*(6*A + 11*B)*Cos[c + d*x] + 4*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.508.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.98, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} (\cos(c + dx)a + a)^{3/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} \left(\sin(c + dx + \frac{\pi}{2}) a + a\right)^{3/2} \left(A + B \sin(c + dx + \frac{\pi}{2})\right) dx \end{aligned}$$

3.508. $\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\int\frac{1}{2}\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a(3a(2A+B)+a(6A+7B)\cos(c+dx))}dx+\dots\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\int\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a(3a(2A+B)+a(6A+7B)\cos(c+dx))}dx+\dots\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a(3a(2A+B)+a(6A+7B)\sin(c+dx))}dx+\dots\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{3}{4}a(14A+11B)\int\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+adx}+\frac{a^2(6A+7B)\sin(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)+\dots\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{3}{4}a(14A+11B)\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx}+\frac{a^2(6A+7B)\sin(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}\right)+\dots\right)$$

↓ 3249

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{3}{4}a(14A+11B)\left(\frac{1}{2}\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)+\dots\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{3}{4}a(14A+11B)\left(\frac{1}{2}\int\frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)+\dots\right)\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{3}{4}a(14A+11B)\left(\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}-\frac{\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)+a}}\right)}{d}\right)+\dots\right)\right)$$

3.508. $\int\frac{(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}}dx$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{a^2(6A+7B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}}+\frac{3}{4}a(14A+11B)\right)\left(\frac{\sqrt{a}\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}\right)\right)$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((a^2*(6*A + 7*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*a*(14*A + 11*B)*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)/6)`

3.508.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3249 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.508.4 Maple [A] (verified)

Time = 20.70 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.38

method	result
default	$\frac{a \left(8B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 22B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 42A \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{24d(1+\cos(dx+c))}$
parts	$\frac{A \left(2 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 7 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 7 \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right) \sqrt{a(1+\cos(dx+c))} a}{4d(1+\cos(dx+c)) \sqrt{\sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}} +$

input `int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{24} \frac{a}{d} \left(8B \cos^2(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 12A \cos(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 22B \cos(dx+c) \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 42A \sin(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \right) \sqrt{a(1+\cos(dx+c))} a$$

3.508.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.76

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \frac{3 \left((14A + 11B)a \cos(dx + c) + (14A + 11B)a \right) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx+c)} + a \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - \frac{(8Ba \cos(dx+c)^3 + 2(6A + 11B)a^2 \cos(dx+c) + 3(14A + 11B)a^3)}{24(d \cos(dx + c) + d)}}{24(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fracas")`

output
$$-\frac{1}{24} \frac{3 \left((14A + 11B)a \cos(dx + c) + (14A + 11B)a \right) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx+c)} + a \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - (8Ba \cos(dx+c)^3 + 2(6A + 11B)a^2 \cos(dx+c) + 3(14A + 11B)a^3)}{24(d \cos(dx + c) + d)}$$

3.508.
$$\int \frac{(a+a \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.508.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.508.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3023 vs. 2(170) = 340.

Time = 0.71 (sec) , antiderivative size = 3023, normalized size of antiderivative = 15.12

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/96*(6*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*((a*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(2*d*x + 2*c) + a*sin(2*d*x + 2*c) - (a*cos(2*d*x + 2*c) - 6*a)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) * cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + (a*sin(2*d*x + 2*c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) - a*cos(2*d*x + 2*c) + (a*cos(2*d*x + 2*c) - 6*a)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 6*a)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))) * sqrt(a) + 7*(a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))) + 1 - a*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) * sin(1/2*arctan2(sin(2*d*x + 2*c), cos...`

3.508.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^{3/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(3/2)/sqrt(sec(d*x + c)), x)`

3.508.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(1/2), x)`

3.509
$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.509.1 Optimal result 4628
 3.509.2 Mathematica [A] (verified) 4629
 3.509.3 Rubi [A] (verified) 4629
 3.509.4 Maple [A] (verified) 4633
 3.509.5 Fricas [A] (verification not implemented) 4633
 3.509.6 Sympy [F(-1)] 4634
 3.509.7 Maxima [B] (verification not implemented) 4634
 3.509.8 Giac [F] 4635
 3.509.9 Mupad [F(-1)] 4636

3.509.1 Optimal result

Integrand size = 35, antiderivative size = 247

$$\int \frac{(a+a \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx = \frac{a^{3/2}(88A+75B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)}}{64d}$$

$$+ \frac{a^2(8A+9B) \sin(c+dx)}{24d \sqrt{a+a \cos(c+dx)} \sec^{\frac{5}{2}}(c+dx)} + \frac{aB \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{4d \sec^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{a^2(88A+75B) \sin(c+dx)}{96d \sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} + \frac{a^2(88A+75B) \sin(c+dx)}{64d \sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}}$$

```
output 1/24*a^2*(8*A+9*B)*sin(d*x+c)/d/sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+1/
96*a^2*(88*A+75*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/
4*a*B*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(5/2)+1/64*a^2*(88*A+
75*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/64*a^(3/2)*(8
8*A+75*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/
2)*sec(d*x+c)^(1/2)/d
```

3.509.2 Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.63

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \frac{a \sqrt{\cos(c + dx)} \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec\left(\frac{1}{2}(c + dx)\right)}}{\sec^{3/2}(c + dx)}$$

input `Integrate[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `(a*Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(3*Sqrt[2]*(88*A + 75*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(296*A + 285*B + 2*(88*A + 93*B)*Cos[c + d*x] + 4*(8*A + 15*B)*Cos[2*(c + d*x)] + 12*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2])/(384*d)`

3.509.3 Rubi [A] (verified)Time = 1.31 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.97, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \cos^{3/2}(c + dx) (\cos(c + dx)a + a)^{3/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sin(c + dx + \frac{\pi}{2})^{3/2} \left(\sin(c + dx + \frac{\pi}{2})a + a\right)^{3/2} \left(A + B \sin(c + dx + \frac{\pi}{2})\right) dx \end{aligned}$$

3.509. $\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}a(88A+75B)\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)\right)\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}a(88A+75B)\left(\frac{3}{4}\left(\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}-\frac{\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)+a}}\right)}{d}\right)\right)\right)\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{a^2(8A+9B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{3d\sqrt{a\cos(c+dx)+a}}+\frac{1}{6}a(88A+75B)\left(\frac{3}{4}\left(\frac{\sqrt{a}\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}\right)\right)\right)\right)$$

input `Int[((a + a*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(4*d) + ((a^2*(8*A + 9*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (a*(88*A + 75*B)*((a*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))))/4)/6)/8)`

3.509.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.509. $\int \frac{(a+a\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

rule 3249 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.509.4 Maple [A] (verified)

Time = 20.38 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.40

method	result
default	$\frac{a\sqrt{a(1+\cos(dx+c))}\left(48B(\cos^2(dx+c))\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+64A\cos(dx+c)\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+120B\cos(dx+c)\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{24d(1+\cos(dx+c))\sec(dx+c)^{\frac{3}{2}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))}\left(8\sin(dx+c)\cos(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+22\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+33\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+33\sec(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{24d(1+\cos(dx+c))\sec(dx+c)^{\frac{3}{2}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((a+cos(d*x+c)*a)^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/192*a/d*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/sec(d*x+c)^(3/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(48*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+64*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+120*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+176*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+150*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+264*tan(d*x+c)*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+225*tan(d*x+c)*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+264*sec(d*x+c)*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+225*sec(d*x+c)*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))
```

3.509.5 Fracas [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.69

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx = \frac{3 \left((88A + 75B)a \cos(dx + c) + (88A + 75B)a \right) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx+c) + a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - \frac{(48Ba \cos(dx+c)^4 + 8(88A + 75B)a^2 \cos(dx+c)^3 + 8(88A + 75B)a^2 \cos(dx+c)^2 + 8(88A + 75B)a^2 \cos(dx+c) + 8(88A + 75B)a^2)}{192(d \cos(dx + c) + d)}}{192(d \cos(dx + c) + d)}$$

```
input integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fracas")
```


output
$$-1/192*(3*((88*A + 75*B)*a*\cos(d*x + c) + (88*A + 75*B)*a)*\sqrt{a}*\arctan(\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c))) - (48*B*a*\cos(d*x + c)^4 + 8*(8*A + 15*B)*a*\cos(d*x + c)^3 + 2*(88*A + 75*B)*a*\cos(d*x + c)^2 + 3*(88*A + 75*B)*a*\cos(d*x + c))*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/\sqrt{\cos(d*x + c)})/(d*\cos(d*x + c) + d)$$

3.509.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.509.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8901 vs. $2(211) = 422$.

Time = 0.98 (sec) , antiderivative size = 8901, normalized size of antiderivative = 36.04

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/768*(8*(4*(a*cos(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1))*sin(3*d*x + 3*c) - (a*cos(3*d*x + 3*c) - a)*sin(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)))*(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(3/4)*sqrt(a) + 6*(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(1/4)*((3*a*sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 11*a*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)) - (3*a*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 5*a*cos(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) - 8*a)*sin(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)))*sqrt(a) + 33*(a*arctan2(-(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(1/4)*(cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos...`

3.509.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^{3/2}}{\sec(dx + c)^{3/2}} dx$$

input `integrate((a+a*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(3/2)/sec(d*x + c)^(3/2), x)`

3.509.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{3/2}}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(3/2), x)`

3.510 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{15}{2}}(c+dx) dx$

3.510.1 Optimal result	4637
3.510.2 Mathematica [A] (verified)	4638
3.510.3 Rubi [A] (verified)	4638
3.510.4 Maple [A] (verified)	4643
3.510.5 Fricas [A] (verification not implemented)	4643
3.510.6 Sympy [F(-1)]	4644
3.510.7 Maxima [B] (verification not implemented)	4644
3.510.8 Giac [F(-1)]	4645
3.510.9 Mupad [B] (verification not implemented)	4646

3.510.1 Optimal result

Integrand size = 35, antiderivative size = 322

$$\begin{aligned} & \int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{\frac{15}{2}}(c + dx) dx \\ &= \frac{32a^3(4184A + 4615B)\sqrt{\sec(c + dx)} \sin(c + dx)}{45045d\sqrt{a + a \cos(c + dx)}} \\ &+ \frac{16a^3(4184A + 4615B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{45045d\sqrt{a + a \cos(c + dx)}} \\ &+ \frac{4a^3(4184A + 4615B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{15015d\sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^3(4184A + 4615B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{9009d\sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^3(280A + 299B) \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{1287d\sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^2(16A + 13B)\sqrt{a + a \cos(c + dx)} \sec^{\frac{11}{2}}(c + dx) \sin(c + dx)}{143d} \\ &+ \frac{2aA(a + a \cos(c + dx))^{3/2} \sec^{\frac{13}{2}}(c + dx) \sin(c + dx)}{13d} \end{aligned}$$

output $\frac{2}{13}a^3(a+a\cos(dx+c))^{3/2}\sec(dx+c)^{13/2}\sin(dx+c)/d+16/45045a^3(4184A+4615B)\sec(dx+c)^{3/2}\sin(dx+c)/d/(a+a\cos(dx+c))^{1/2}+4/15015a^3(4184A+4615B)\sec(dx+c)^{5/2}\sin(dx+c)/d/(a+a\cos(dx+c))^{1/2}+2/9009a^3(4184A+4615B)\sec(dx+c)^{7/2}\sin(dx+c)/d/(a+a\cos(dx+c))^{1/2}+2/1287a^3(280A+299B)\sec(dx+c)^{9/2}\sin(dx+c)/d/(a+a\cos(dx+c))^{1/2}+2/143a^2(16A+13B)\sec(dx+c)^{11/2}\sin(dx+c)(a+a\cos(dx+c))^{1/2}/d+32/45045a^3(4184A+4615B)\sin(dx+c)\sec(dx+c)^{1/2}/d/(a+a\cos(dx+c))^{1/2}$

3.510.2 Mathematica [A] (verified)

Time = 0.87 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.53

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (171806A + 162955B + 35(5552A + 5083B) \cos(c + dx) + 14(15167A +$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(15/2),x]`

output $(a^2 \sqrt{a(1 + \cos(c + dx))} (171806A + 162955B + 35(5552A + 5083B) \cos(c + dx) + 14(15167A + 15925B) \cos(2(c + dx)) + 62760A \cos(3(c + dx)) + 69225B \cos(3(c + dx)) + 62760A \cos(4(c + dx)) + 69225B \cos(4(c + dx)) + 8368A \cos(5(c + dx)) + 9230B \cos(5(c + dx)) + 8368A \cos(6(c + dx)) + 9230B \cos(6(c + dx))) \sec^{13/2}(c + dx) \tan[(c + dx)/2]) / (90090d)$

3.510.3 Rubi [A] (verified)

Time = 1.95 (sec) , antiderivative size = 342, normalized size of antiderivative = 1.06, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.510. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx$

$$\begin{aligned}
& \int \sec^{\frac{15}{2}}(c+dx)(a \cos(c+dx) + a)^{5/2}(A + B \cos(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \csc\left(c+dx+\frac{\pi}{2}\right)^{15/2} \left(a \sin\left(c+dx+\frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(\cos(c+dx)a + a)^{5/2}(A + B \cos(c+dx))}{\cos^{\frac{15}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(\sin(c+dx+\frac{\pi}{2})a + a)^{5/2}(A + B \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{15/2}} dx \\
& \quad \downarrow \text{3454} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2}{13} \int \frac{(\cos(c+dx)a + a)^{3/2}(a(16A + 13B) + a(8A + 13B)\cos(c+dx))}{2\cos^{\frac{13}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)}{\cos^{\frac{13}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{13} \int \frac{(\cos(c+dx)a + a)^{3/2}(a(16A + 13B) + a(8A + 13B)\cos(c+dx))}{\cos^{\frac{13}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)}{\cos^{\frac{13}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{13} \int \frac{(\sin(c+dx+\frac{\pi}{2})a + a)^{3/2}(a(16A + 13B) + a(8A + 13B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{13/2}} dx + \frac{2aA \cos(c+dx+\frac{\pi}{2})}{\sin^{\frac{13}{2}}(c+dx+\frac{\pi}{2})} \right) \\
& \quad \downarrow \text{3454} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{13} \left(\frac{2}{11} \int \frac{\sqrt{\cos(c+dx)a + a}((280A + 299B)a^2 + (216A + 247B)\cos(c+dx)a^2)}{2\cos^{\frac{11}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)}{\cos^{\frac{11}{2}}(c+dx)} \right) \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{13} \left(\frac{1}{11} \int \frac{\sqrt{\cos(c+dx)a + a}((280A + 299B)a^2 + (216A + 247B)\cos(c+dx)a^2)}{\cos^{\frac{11}{2}}(c+dx)} dx + \frac{2aA \sin(c+dx)}{\cos^{\frac{11}{2}}(c+dx)} \right) \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.510. $\int (a + a \cos(c+dx))^{5/2}(A + B \cos(c+dx)) \sec^{\frac{15}{2}}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a((280A+299B)a^2+(216A+247B)\sin(c+dx))}}{\sin(c+dx+\frac{\pi}{2})^{11/2}}dx\right.\right.$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{9/2}(c+dx)}dx+\frac{2a^3(280A+299B)\sin(c+dx)}{9d\cos^{9/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{9/2}}dx+\frac{2a^3(280A+299B)\sin(c+dx)}{9d\cos^{9/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\left(\frac{6}{7}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{7/2}(c+dx)}dx+\frac{2a\sin(c+dx)}{7d\cos^{7/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\left(\frac{6}{7}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2a\sin(c+dx)}{7d\cos^{7/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.\right.$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\left(\frac{6}{7}\left(\frac{4}{5}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{5/2}(c+dx)}dx+\frac{2a\sin(c+dx)}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.\right.\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\left(\frac{6}{7}\left(\frac{4}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a\sin(c+dx)}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.\right.\right.$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\left(\frac{6}{7}\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{3/2}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{3/2}(c+dx)\sqrt{a\cos(c+dx)}}\right.\right.\right.\right.\right.\right.$$

↓ 3042

3.510. $\int(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sec^{15/2}(c+dx)dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{1}{11}\left(\frac{1}{9}a^2(4184A+4615B)\left(\frac{6}{7}\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{3}{3d\cos^{\frac{3}{2}}(c+dx)}\right)\right)\right)\right)\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{13}\left(\frac{2a^2(16A+13B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{11d\cos^{\frac{11}{2}}(c+dx)}+\frac{1}{11}\left(\frac{2a^3(280A+299B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{9d\cos^{\frac{9}{2}}(c+dx)}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(15/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(13*d*Cos[c + d*x]^(13/2)) + ((2*a^2*(16*A + 13*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(11*d*Cos[c + d*x]^(11/2)) + ((2*a^3*(280*A + 299*B)*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)*Sqrt[a + a*Cos[c + d*x]]) + (a^2*(4184*A + 4615*B)*((2*a*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) + (6*((2*a*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x])) + (4*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x])) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]))))/5))/7))/9)/11)/13)`

3.510.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3459 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1]`

3.510.4 Maple [A] (verified)

Time = 2.20 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.51

$$\frac{2a^2 \cot(dx + c) (\cos(dx + c) - 1) ((66944(\cos^6(dx + c)) + 33472(\cos^5(dx + c)) + 25104(\cos^4(dx + c)) + \dots$$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(15/2),x)`

output `-2/45045*a^2/d*cot(d*x+c)*(cos(d*x+c)-1)*((66944*cos(d*x+c)^6+33472*cos(d*x+c)^5+25104*cos(d*x+c)^4+20920*cos(d*x+c)^3+18305*cos(d*x+c)^2+11970*cos(d*x+c)+3465)*A+cos(d*x+c)*(73840*cos(d*x+c)^5+36920*cos(d*x+c)^4+27690*cos(d*x+c)^3+23075*cos(d*x+c)^2+14560*cos(d*x+c)+4095)*B)*(a*(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(15/2)`

3.510.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.55

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx = \frac{2(16(4184A + 4615B)a^2 \cos(dx + c)^6 + 8(4184A + 4615B)a^2 \cos(dx + c)^5 + 6(4184A + 4615B)a^2 \cos(dx + c)^4 + \dots}{\dots}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(15/2),x, algorithm="fracas")`

output `2/45045*(16*(4184*A + 4615*B)*a^2*cos(d*x + c)^6 + 8*(4184*A + 4615*B)*a^2*cos(d*x + c)^5 + 6*(4184*A + 4615*B)*a^2*cos(d*x + c)^4 + 5*(4184*A + 4615*B)*a^2*cos(d*x + c)^3 + 35*(523*A + 416*B)*a^2*cos(d*x + c)^2 + 315*(38*A + 13*B)*a^2*cos(d*x + c) + 3465*A*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/((d*cos(d*x + c))^7 + d*cos(d*x + c)^6)*sqrt(cos(d*x + c))`

3.510.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(15/2),x)`

output `Timed out`

3.510.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 763 vs. $2(280) = 560$.

Time = 0.37 (sec) , antiderivative size = 763, normalized size of antiderivative = 2.37

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(15/2),x, algo
rithm="maxima")`

output `8/45045*((45045*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 165165*sqrt(2)*a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 414414*sqrt(2)*a^(5/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 604890*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 522665*sqrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 289185*sqrt(2)*a^(5/2)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11 + 88980*sqrt(2)*a^(5/2)*sin(d*x + c)^13/(cos(d*x + c) + 1)^13 - 11864*sqrt(2)*a^(5/2)*sin(d*x + c)^15/(cos(d*x + c) + 1)^15)*A*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^5/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(15/2))*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(15/2)*(5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 10*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 5*sin(d*x + c)^8/(cos(d*x + c) + 1)^8 + sin(d*x + c)^10/(cos(d*x + c) + 1)^10 + 1)) + 65*(693*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 3003*sqrt(2)*a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 6930*sqrt(2)*a^(5/2)*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 10098*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 9053*sqrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9 - 4875*sqrt(2)*a^(5/2)*sin(d*x + c)^11/(cos(d*x + c) + 1)^11 + 1500*sqrt(2)*a^(5/2)*sin(d*x + c)^13/(cos(d*x + c) + 1)^13 - 200*sqrt(2)*a^(5/2)*sin(d*x + c)^15/(cos(d*x + c) + 1)^15)*B*(sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^5/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(15/2))*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(15/2)*(5*sin(d*x + c)^2/(c...`

3.510.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(15/2),x, algorithm="giac")`

output `Timed out`

3.510.9 Mupad [B] (verification not implemented)

Time = 6.74 (sec) , antiderivative size = 789, normalized size of antiderivative = 2.45

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{15/2}(c + dx) dx = \text{Too large to display}$$

```
input int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(15/2)*(a + a*cos(c + d*x))^(5/2),x)
```

```
output ((1/(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a^2*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(4184*A + 4615*B)*32i)/(45045*d) - (a^2*exp(c*5i + d*x*5i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(2*A + 5*B)*16i)/(5*d) + (a^2*exp(c*8i + d*x*8i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(2*A + 5*B)*16i)/(5*d) + (a^2*exp(c*6i + d*x*6i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(116*A + 115*B)*16i)/(35*d) - (a^2*exp(c*7i + d*x*7i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(116*A + 115*B)*16i)/(35*d) + (a^2*exp(c*4i + d*x*4i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(1046*A + 1075*B)*16i)/(315*d) - (a^2*exp(c*9i + d*x*9i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(1046*A + 1075*B)*16i)/(315*d) + (a^2*exp(c*2i + d*x*2i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(4184*A + 4615*B)*16i)/(3465*d) - (a^2*exp(c*11i + d*x*11i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(4184*A + 4615*B)*16i)/(3465*d) - (a^2*exp(c*13i + d*x*13i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(4184*A + 4615*B)*32i)/(45045*d)))/(exp(c*1i + d*x*1i) + 6*exp(c*2i + d*x*2i) + 6*exp(c*3i + d*x*3i) + 15*exp(c*4i + d*x*4i) + 15*exp(c*5i + d*x*5i) + 20*exp(c*6i + d*x*6i) + 20*exp(c*7i + d*x*7i) + 15*exp(c*8i + d*x*8i) + 15*exp(c*9i + d*x*9i) + 6*exp(c*10i + d*x*10i) + 6*exp(c*11i + d*x*11i...
```

3.511 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{13}{2}}(c+dx) dx$

3.511.1 Optimal result	4647
3.511.2 Mathematica [A] (verified)	4648
3.511.3 Rubi [A] (verified)	4648
3.511.4 Maple [A] (verified)	4652
3.511.5 Fricas [A] (verification not implemented)	4653
3.511.6 Sympy [F(-1)]	4653
3.511.7 Maxima [B] (verification not implemented)	4653
3.511.8 Giac [F(-1)]	4654
3.511.9 Mupad [B] (verification not implemented)	4655

3.511.1 Optimal result

Integrand size = 35, antiderivative size = 275

$$\begin{aligned} & \int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx \\ &= \frac{16a^3(710A + 803B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3465d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{8a^3(710A + 803B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3465d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^3(710A + 803B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{1155d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^3(194A + 209B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{693d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^2(14A + 11B) \sqrt{a + a \cos(c + dx)} \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{99d} \\ &+ \frac{2aA(a + a \cos(c + dx))^{3/2} \sec^{\frac{11}{2}}(c + dx) \sin(c + dx)}{11d} \end{aligned}$$

output $2/11*a*A*(a+a*\cos(d*x+c))^{(3/2)}*\sec(d*x+c)^{(11/2)}*\sin(d*x+c)/d+8/3465*a^3*(710*A+803*B)*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/1155*a^3*(710*A+803*B)*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/693*a^3*(194*A+209*B)*\sec(d*x+c)^{(7/2)}*\sin(d*x+c)/d/(a+a*\cos(d*x+c))^{(1/2)}+2/99*a^2*(14*A+11*B)*\sec(d*x+c)^{(9/2)}*\sin(d*x+c)*(a+a*\cos(d*x+c))^{(1/2)}/d+16/3465*a^3*(710*A+803*B)*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/d/(a+a*\cos(d*x+c))^{(1/2)}$

3.511.2 Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.53

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (9070A + 7678B + (25070A + 24827B) \cos(c + dx) + (9230A + 9284B) \cos(2(c + dx)) + 9230A \cos(3(c + dx)) + 10439B \cos(3(c + dx)) + 1420A \cos(4(c + dx)) + 1606B \cos(4(c + dx)) + 1420A \cos(5(c + dx)) + 1606B \cos(5(c + dx))) \sec^{11/2}(c + dx) \tan((c + dx)/2)}{(6930*d)}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(13/2),x]`

output $(a^2*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*(9070*A + 7678*B + (25070*A + 24827*B)*\text{Cos}[c + d*x] + (9230*A + 9284*B)*\text{Cos}[2*(c + d*x)] + 9230*A*\text{Cos}[3*(c + d*x)] + 10439*B*\text{Cos}[3*(c + d*x)] + 1420*A*\text{Cos}[4*(c + d*x)] + 1606*B*\text{Cos}[4*(c + d*x)] + 1420*A*\text{Cos}[5*(c + d*x)] + 1606*B*\text{Cos}[5*(c + d*x)])*\text{Sec}[c + d*x]^{(11/2)}*\text{Tan}[(c + d*x)/2])/(6930*d)$

3.511.3 Rubi [A] (verified)

Time = 1.69 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.09, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{13/2}(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx$$

↓ 3042

3.511. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx$

$$\begin{aligned}
& \int \csc\left(c + dx + \frac{\pi}{2}\right)^{13/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2} (A + B \cos(c + dx))}{\cos^{13/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{13/2}} dx \\
& \quad \downarrow \text{3454} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2}{11} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(14A + 11B) + a(6A + 11B) \cos(c + dx))}{2 \cos^{11/2}(c + dx)} dx + \frac{2aA \sin(c + dx)}{\cos^{11/2}(c + dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \int \frac{(\cos(c + dx)a + a)^{3/2} (a(14A + 11B) + a(6A + 11B) \cos(c + dx))}{\cos^{11/2}(c + dx)} dx + \frac{2aA \sin(c + dx)}{\cos^{11/2}(c + dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{3/2} (a(14A + 11B) + a(6A + 11B) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx + \frac{2aA \sin(c + dx)}{\sin(c + dx + \frac{\pi}{2})^{11/2}} \right) \\
& \quad \downarrow \text{3454} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \left(\frac{2}{9} \int \frac{\sqrt{\cos(c + dx)a + a} ((194A + 209B)a^2 + 3(46A + 55B) \cos(c + dx)a^2)}{2 \cos^{9/2}(c + dx)} dx + \frac{2aA \sin(c + dx)}{\cos^{9/2}(c + dx)} \right) \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \left(\frac{1}{9} \int \frac{\sqrt{\cos(c + dx)a + a} ((194A + 209B)a^2 + 3(46A + 55B) \cos(c + dx)a^2)}{\cos^{9/2}(c + dx)} dx + \frac{2aA \sin(c + dx)}{\cos^{9/2}(c + dx)} \right) \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{11} \left(\frac{1}{9} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})a + a} ((194A + 209B)a^2 + 3(46A + 55B) \sin(c + dx + \frac{\pi}{2})a^2)}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx + \frac{2aA \sin(c + dx)}{\sin(c + dx + \frac{\pi}{2})^{9/2}} \right) \right)
\end{aligned}$$

3.511. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{3}{7}a^2(710A+803B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{7}{2}}(c+dx)}dx+\frac{2a^3(194A+209B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{3}{7}a^2(710A+803B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2a^3(194A+209B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{3}{7}a^2(710A+803B)\left(\frac{4}{5}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{5}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{3}{7}a^2(710A+803B)\left(\frac{4}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2a\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{3}{7}a^2(710A+803B)\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{3}{7}a^2(710A+803B)\left(\frac{4}{5}\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{2a^2(14A+11B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{9d\cos^{\frac{9}{2}}(c+dx)}+\frac{1}{9}\left(\frac{2a^3(194A+209B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(13/2), x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + a*cos[c + d*x])^(3/2)*Sin[c + d*x])/(11*d*cos[c + d*x]^(11/2)) + ((2*a^2*(14*A + 11*B)*Sqrt[a + a*cos[c + d*x]]*Sin[c + d*x])/(9*d*cos[c + d*x]^(9/2)) + ((2*a^3*(194*A + 209*B)*Sin[c + d*x])/(7*d*cos[c + d*x]^(7/2)*Sqrt[a + a*cos[c + d*x]]) + (3*a^2*(710*A + 803*B)*((2*a*sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)*Sqrt[a + a*cos[c + d*x]]) + (4*((2*a*sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)*Sqrt[a + a*cos[c + d*x])) + (4*a*sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*cos[c + d*x]])))/5)/7)/9)/11)
```

3.511.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3250 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3251 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])
```

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.511.4 Maple [A] (verified)

Time = 2.21 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.52

$$\frac{2a^2 \cot(dx + c) (\cos(dx + c) - 1) ((5680 \cos^5(dx + c)) + 2840 \cos^4(dx + c) + 2130 \cos^3(dx + c) + 1775 \cos^2(dx + c) + 1120 \cos(dx + c) + 315) A + \cos(dx + c) (6424 \cos^4(dx + c) + 3212 \cos^3(dx + c) + 2409 \cos^2(dx + c) + 1430 \cos(dx + c) + 385) B}{(a + a \cos(dx + c))^{5/2} (A + B \cos(dx + c)) \sec^{13/2}(c + dx) dx}$$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x)
```

```
output -2/3465*a^2/d*cot(d*x+c)*(cos(d*x+c)-1)*((5680*cos(d*x+c)^5+2840*cos(d*x+c)
)^4+2130*cos(d*x+c)^3+1775*cos(d*x+c)^2+1120*cos(d*x+c)+315)*A+cos(d*x+c)*
(6424*cos(d*x+c)^4+3212*cos(d*x+c)^3+2409*cos(d*x+c)^2+1430*cos(d*x+c)+385
)*B*(a*(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(13/2)
```

3.511. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx$

3.511.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.57

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \frac{2(8(710A + 803B)a^2 \cos(dx + c)^5 + 4(710A + 803B)a^2 \cos(dx + c)^4 + 3(710A + 803B)a^2 \cos(dx + c)^3 + 2(8(710A + 803B)a^2 \cos(dx + c)^2 + 4(710A + 803B)a^2 \cos(dx + c) + 3(710A + 803B)a^2) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{3465 (d \cos(dx + c))^6 + d \cos(dx + c)^5 \sqrt{\cos(dx + c)}}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="fricas")`

output `2/3465*(8*(710*A + 803*B)*a^2*cos(d*x + c)^5 + 4*(710*A + 803*B)*a^2*cos(d*x + c)^4 + 3*(710*A + 803*B)*a^2*cos(d*x + c)^3 + 5*(355*A + 286*B)*a^2*cos(d*x + c)^2 + 35*(32*A + 11*B)*a^2*cos(d*x + c) + 315*A*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/((d*cos(d*x + c))^6 + d*cos(d*x + c)^5)*sqrt(cos(d*x + c))`

3.511.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(13/2),x)`

output `Timed out`

3.511.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 672 vs. 2(239) = 478.

Time = 0.37 (sec) , antiderivative size = 672, normalized size of antiderivative = 2.44

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="maxima")`

output
$$\frac{8}{3465} \cdot (5 \cdot (693 \sqrt{2}) a^{5/2} \sin(dx+c) / (\cos(dx+c)+1) - 2310 \sqrt{2}) a^{5/2} \sin(dx+c)^3 / (\cos(dx+c)+1)^3 + 4620 \sqrt{2} a^{5/2} \sin(dx+c)^5 / (\cos(dx+c)+1)^5 - 5478 \sqrt{2} a^{5/2} \sin(dx+c)^7 / (\cos(dx+c)+1)^7 + 3575 \sqrt{2} a^{5/2} \sin(dx+c)^9 / (\cos(dx+c)+1)^9 - 1300 \sqrt{2} a^{5/2} \sin(dx+c)^{11} / (\cos(dx+c)+1)^{11} + 200 \sqrt{2} a^{5/2} \sin(dx+c)^{13} / (\cos(dx+c)+1)^{13} \cdot A \cdot (\sin(dx+c))^2 / (\cos(dx+c)+1)^2 + 1)^4 / ((\sin(dx+c) / (\cos(dx+c)+1) + 1)^{13/2} \cdot (-\sin(dx+c) / (\cos(dx+c)+1) + 1)^{13/2} \cdot (4 \sin(dx+c)^2 / (\cos(dx+c)+1)^2 + 6 \sin(dx+c)^4 / (\cos(dx+c)+1)^4 + 4 \sin(dx+c)^6 / (\cos(dx+c)+1)^6 + \sin(dx+c)^8 / (\cos(dx+c)+1)^8 + 1)) + 11 \cdot (315 \sqrt{2}) a^{5/2} \sin(dx+c) / (\cos(dx+c)+1) - 1260 \sqrt{2} a^{5/2} \sin(dx+c)^3 / (\cos(dx+c)+1)^3 + 2394 \sqrt{2} a^{5/2} \sin(dx+c)^5 / (\cos(dx+c)+1)^5 - 2736 \sqrt{2} a^{5/2} \sin(dx+c)^7 / (\cos(dx+c)+1)^7 + 1859 \sqrt{2} a^{5/2} \sin(dx+c)^9 / (\cos(dx+c)+1)^9 - 676 \sqrt{2} a^{5/2} \sin(dx+c)^{11} / (\cos(dx+c)+1)^{11} + 104 \sqrt{2} a^{5/2} \sin(dx+c)^{13} / (\cos(dx+c)+1)^{13} \cdot B \cdot (\sin(dx+c))^2 / (\cos(dx+c)+1)^2 + 1)^4 / ((\sin(dx+c) / (\cos(dx+c)+1) + 1)^{13/2} \cdot (-\sin(dx+c) / (\cos(dx+c)+1) + 1)^{13/2} \cdot (4 \sin(dx+c)^2 / (\cos(dx+c)+1)^2 + 6 \sin(dx+c)^4 / (\cos(dx+c)+1)^4 + 4 \sin(dx+c)^6 / (\cos(dx+c)+1)^6 + \sin(dx+c)^8 / (\cos(dx+c)+1)^8 + 1)) / d$$

3.511.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="giac")`

output Timed out

3.511.9 Mupad [B] (verification not implemented)

Time = 5.44 (sec) , antiderivative size = 751, normalized size of antiderivative = 2.73

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx = \text{Too large to display}$$

```
input int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(13/2)*(a + a*cos(c + d*x))^(5/2),x)
```

```
output ((1/(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a^2*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(710*A + 803*B)*16i)/(3465*d) - (B*a^2*exp(c*3i + d*x*3i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*8i)/(3*d) + (B*a^2*exp(c*8i + d*x*8i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*8i)/(3*d) - (a^2*exp(c*5i + d*x*5i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(30*A + 41*B)*8i)/(15*d) + (a^2*exp(c*6i + d*x*6i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(30*A + 41*B)*8i)/(15*d) + (a^2*exp(c*4i + d*x*4i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(160*A + 157*B)*8i)/(35*d) - (a^2*exp(c*7i + d*x*7i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(160*A + 157*B)*8i)/(35*d) + (a^2*exp(c*2i + d*x*2i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(710*A + 803*B)*8i)/(315*d) - (a^2*exp(c*9i + d*x*9i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(710*A + 803*B)*8i)/(315*d) - (a^2*exp(c*11i + d*x*11i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(710*A + 803*B)*16i)/(3465*d))/((exp(c*1i + d*x*1i) + 5*exp(c*2i + d*x*2i) + 5*exp(c*3i + d*x*3i) + 10*exp(c*4i + d*x*4i) + 10*exp(c*5i + d*x*5i) + 10*exp(c*6i + d*x*6i) + 10*exp(c*7i + d*x*7i) + 5*exp(c*8i + d*x*8i) + 5*exp(c*9i + d*x*9i) + exp(c*10i + d*x*10i) + exp(c*11i + d*x*11i) + 1)
```

3.512 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$

3.512.1 Optimal result	4656
3.512.2 Mathematica [A] (verified)	4657
3.512.3 Rubi [A] (verified)	4657
3.512.4 Maple [A] (verified)	4661
3.512.5 Fricas [A] (verification not implemented)	4661
3.512.6 Sympy [F(-1)]	4662
3.512.7 Maxima [B] (verification not implemented)	4662
3.512.8 Giac [F(-1)]	4663
3.512.9 Mupad [B] (verification not implemented)	4663

3.512.1 Optimal result

Integrand size = 35, antiderivative size = 228

$$\begin{aligned} &\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx \\ &= \frac{4a^3(292A + 345B) \sqrt{\sec(c + dx)} \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^3(292A + 345B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^3(124A + 135B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}} \\ &+ \frac{2a^2(4A + 3B) \sqrt{a + a \cos(c + dx)} \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{21d} \\ &+ \frac{2aA(a + a \cos(c + dx))^{3/2} \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{9d} \end{aligned}$$

output

```
2/9*a*A*(a+a*cos(d*x+c))^(3/2)*sec(d*x+c)^(9/2)*sin(d*x+c)/d+2/315*a^3*(29
2*A+345*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/315*a^3*
(124*A+135*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/21*a^
2*(4*A+3*B)*sec(d*x+c)^(7/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+4/315*a^3
*(292*A+345*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.512.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.55

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (1454A + 1395B + (1396A + 1215B) \cos(c + dx) + 2(803A + 870B) \cos^2(c + dx))}{630d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x]])*(1454*A + 1395*B + (1396*A + 1215*B)*Cos[c + d*x] + 2*(803*A + 870*B)*Cos[2*(c + d*x)] + 292*A*Cos[3*(c + d*x)] + 345*B*Cos[3*(c + d*x)] + 292*A*Cos[4*(c + d*x)] + 345*B*Cos[4*(c + d*x)])*Sec[c + d*x]^(9/2)*Tan[(c + d*x)/2])/(630*d)`

3.512.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3251, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{\frac{11}{2}}(c + dx) (a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{11/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2} (A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx \end{aligned}$$

3.512. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{9}\int\frac{(\cos(c+dx)a+a)^{3/2}(3a(4A+3B)+a(4A+9B)\cos(c+dx))}{2\cos^{9/2}(c+dx)}dx+\frac{2aA\sin(c+dx)}{2\cos^{9/2}(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\int\frac{(\cos(c+dx)a+a)^{3/2}(3a(4A+3B)+a(4A+9B)\cos(c+dx))}{\cos^{9/2}(c+dx)}dx+\frac{2aA\sin(c+dx)}{\cos^{9/2}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\int\frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(3a(4A+3B)+a(4A+9B)\sin(c+dx+\frac{\pi}{2}))}{\sin^{9/2}(c+dx+\frac{\pi}{2})}dx+\frac{2aA\cos(c+dx+\frac{\pi}{2})}{\sin^{9/2}(c+dx+\frac{\pi}{2})}\right)$$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{2}{7}\int\frac{\sqrt{\cos(c+dx)a+a}((124A+135B)a^2+(76A+99B)\cos(c+dx)a^2)}{2\cos^{7/2}(c+dx)}dx+\frac{2a((124A+135B)a^2+(76A+99B)\cos(c+dx)a^2)}{2\cos^{7/2}(c+dx)}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\int\frac{\sqrt{\cos(c+dx)a+a}((124A+135B)a^2+(76A+99B)\cos(c+dx)a^2)}{\cos^{7/2}(c+dx)}dx+\frac{2a((124A+135B)a^2+(76A+99B)\cos(c+dx)a^2)}{\cos^{7/2}(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((124A+135B)a^2+(76A+99B)\sin(c+dx+\frac{\pi}{2}))}{\sin^{7/2}(c+dx+\frac{\pi}{2})}dx+\frac{2a((124A+135B)a^2+(76A+99B)\sin(c+dx+\frac{\pi}{2}))}{\sin^{7/2}(c+dx+\frac{\pi}{2})}\right)\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{3}{5}a^2(292A+345B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{5/2}(c+dx)}dx+\frac{2a^3(124A+135B)\sin(c+dx)}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{3}{5}a^2(292A+345B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin^{5/2}(c+dx+\frac{\pi}{2})}dx+\frac{2a^3(124A+135B)\sin(c+dx+\frac{\pi}{2})}{5d\cos^{5/2}(c+dx)\sqrt{a\cos(c+dx+\frac{\pi}{2})}}\right)\right)\right)$$

↓ 3251

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{3}{5}a^2(292A+345B)\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{\frac{3}{2}}(c+dx)}dx+\frac{2a\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{3}{5}a^2(292A+345B)\left(\frac{2}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})}dx+\frac{2a\sin(c+dx+\frac{\pi}{2})}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx+\frac{\pi}{2})}}\right)\right)\right)\right)$$

↓ 3250

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{6a^2(4A+3B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{7d\cos^{\frac{7}{2}}(c+dx)}+\frac{1}{7}\left(\frac{2a^3(124A+135B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((6*a^2*(4*A + 3*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*a^3*(124*A + 135*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) + (3*a^2*(292*A + 345*B)*((2*a*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (4*a*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/5)/7)/9)`

3.512.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3251 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2)*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[(2*n + 3)*((b*c - a*d)/(2*b*(n + 1)*(c^2 - d^2))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && NeQ[2*n + 3, 0] && IntegerQ[2*n]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3454 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.512.4 Maple [A] (verified)

Time = 2.68 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.54

$$2a^2 \cot(dx + c) (\cos(dx + c) - 1) ((584 \cos^4(dx + c)) + 292 \cos^3(dx + c) + 219 \cos^2(dx + c) + 130 \cos(dx + c) + 35) A + \cos(dx + c) (690 \cos^3(dx + c) + 345 \cos^2(dx + c) + 180 \cos(dx + c) + 45) B) (a(1 + \cos(dx + c)))^{1/2} \sec(dx + c)^{11/2}$$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x)
```

```
output -2/315*a^2/d*cot(d*x+c)*(cos(d*x+c)-1)*((584*cos(d*x+c)^4+292*cos(d*x+c)^3
+219*cos(d*x+c)^2+130*cos(d*x+c)+35)*A+cos(d*x+c)*(690*cos(d*x+c)^3+345*co
s(d*x+c)^2+180*cos(d*x+c)+45)*B)*(a*(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(11/2
)
```

3.512.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.59

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \frac{2(2(292A + 345B)a^2 \cos^4(dx + c) + (292A + 345B)a^2 \cos^3(dx + c) + 3(73A + 60B)a^2 \cos^2(dx + c) + 130a^2 \cos(dx + c) + 35)A + \cos(dx + c)(690 \cos^3(dx + c) + 345 \cos^2(dx + c) + 180 \cos(dx + c) + 45)B}{315(d \cos(dx + c))^5 + d \cos(dx + c)}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, alg
orithm="fracas")
```

output $2/315*(2*(292*A + 345*B)*a^2*\cos(d*x + c)^4 + (292*A + 345*B)*a^2*\cos(d*x + c)^3 + 3*(73*A + 60*B)*a^2*\cos(d*x + c)^2 + 5*(26*A + 9*B)*a^2*\cos(d*x + c) + 35*A*a^2)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/((d*\cos(d*x + c))^5 + d*\cos(d*x + c)^4)*\sqrt{\cos(d*x + c)}$

3.512.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(11/2),x)`

output `Timed out`

3.512.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 579 vs. $2(198) = 396$.

Time = 0.38 (sec) , antiderivative size = 579, normalized size of antiderivative = 2.54

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \frac{8 \left(\frac{315 \sqrt{2} a^{5/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{945 \sqrt{2} a^{5/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{1449 \sqrt{2} a^{5/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{1287 \sqrt{2} a^{5/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{572 \sqrt{2} a^{5/2} \sin(dx+c)^9}{(\cos(dx+c)+1)^9} - \frac{104 \sqrt{2} a^{5/2} \sin(dx+c)^{11}}{(\cos(dx+c)+1)^{11}} \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{11/2} \left(\frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{\sin(dx+c)^6}{(\cos(dx+c)+1)^6} + 1 \right)}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="maxima")`

output $\frac{8}{315} \left(\frac{315 \sqrt{2} a^{5/2} \sin(dx+c)}{(\cos(dx+c)+1)} - 945 \sqrt{2} a^{5/2} \sin(dx+c)^3 / (\cos(dx+c)+1)^3 + 1449 \sqrt{2} a^{5/2} \sin(dx+c)^5 / (\cos(dx+c)+1)^5 - 1287 \sqrt{2} a^{5/2} \sin(dx+c)^7 / (\cos(dx+c)+1)^7 + 572 \sqrt{2} a^{5/2} \sin(dx+c)^9 / (\cos(dx+c)+1)^9 - 104 \sqrt{2} a^{5/2} \sin(dx+c)^{11} / (\cos(dx+c)+1)^{11} \right) A \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 1} \right)^3 / \left(\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(\frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 3 \sin(dx+c)^4 / (\cos(dx+c)+1)^4 + \sin(dx+c)^6 / (\cos(dx+c)+1)^6 + 1} \right) + 15 \left(\frac{21 \sqrt{2} a^{5/2} \sin(dx+c)}{(\cos(dx+c)+1)} - 77 \sqrt{2} a^{5/2} \sin(dx+c)^3 / (\cos(dx+c)+1)^3 + 119 \sqrt{2} a^{5/2} \sin(dx+c)^5 / (\cos(dx+c)+1)^5 - 99 \sqrt{2} a^{5/2} \sin(dx+c)^7 / (\cos(dx+c)+1)^7 + 44 \sqrt{2} a^{5/2} \sin(dx+c)^9 / (\cos(dx+c)+1)^9 - 8 \sqrt{2} a^{5/2} \sin(dx+c)^{11} / (\cos(dx+c)+1)^{11} \right) B \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 1} \right)^3 / \left(\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(-\frac{\sin(dx+c)}{(\cos(dx+c)+1)} + 1 \right)^{11/2} \left(\frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2 + 3 \sin(dx+c)^4 / (\cos(dx+c)+1)^4 + \sin(dx+c)^6 / (\cos(dx+c)+1)^6 + 1} \right) / d$

3.512.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{1/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="giac")`

output Timed out

3.512.9 Mupad [B] (verification not implemented)

Time = 5.60 (sec) , antiderivative size = 617, normalized size of antiderivative = 2.71

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{1/2}(c + dx) dx = \frac{\sqrt{\frac{1}{\frac{e^{-c \operatorname{li}-dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li}+dx \operatorname{li}}}{2}}}}{\frac{a^2 \sqrt{a+a \left(\frac{e^{-c \operatorname{li}-dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li}+dx \operatorname{li}}}{2} \right)}{(292 A+345 B) 4i}}}{315 d} - \frac{a^2 e^{c 3i+dx 3i} \sqrt{a+a \left(\frac{e^{-c \operatorname{li}-dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li}+dx \operatorname{li}}}{2} \right)}}{3 d}$$

3.512. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{1/2}(c + dx) dx$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + a*cos(c + d*x))^(5/2),x)`

output `((1/(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a^2*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(292*A + 345*B)*4i)/(315*d) - (a^2*exp(c*3i + d*x*3i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(2*A + 5*B)*4i)/(3*d) + (a^2*exp(c*6i + d*x*6i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(2*A + 5*B)*4i)/(3*d) + (a^2*exp(c*4i + d*x*4i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(24*A + 25*B)*4i)/(5*d) - (a^2*exp(c*5i + d*x*5i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(24*A + 25*B)*4i)/(5*d) + (a^2*exp(c*2i + d*x*2i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(146*A + 155*B)*4i)/(35*d) - (a^2*exp(c*7i + d*x*7i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(146*A + 155*B)*4i)/(35*d) - (a^2*exp(c*9i + d*x*9i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(292*A + 345*B)*4i)/(315*d)))/(exp(c*1i + d*x*1i) + 4*exp(c*2i + d*x*2i) + 4*exp(c*3i + d*x*3i) + 6*exp(c*4i + d*x*4i) + 6*exp(c*5i + d*x*5i) + 4*exp(c*6i + d*x*6i) + 4*exp(c*7i + d*x*7i) + exp(c*8i + d*x*8i) + exp(c*9i + d*x*9i) + 1)`

3.513 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{9/2}(c+dx) dx$

3.513.1 Optimal result	4665
3.513.2 Mathematica [A] (verified)	4666
3.513.3 Rubi [A] (verified)	4666
3.513.4 Maple [A] (verified)	4669
3.513.5 Fricas [A] (verification not implemented)	4670
3.513.6 Sympy [F(-1)]	4670
3.513.7 Maxima [B] (verification not implemented)	4670
3.513.8 Giac [F(-1)]	4671
3.513.9 Mupad [B] (verification not implemented)	4671

3.513.1 Optimal result

Integrand size = 35, antiderivative size = 181

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{2a^3(230A + 301B) \sqrt{\sec(c + dx)} \sin(c + dx)}{105d \sqrt{a + a \cos(c + dx)}} + \frac{2a^3(10A + 11B) \sec^{3/2}(c + dx) \sin(c + dx)}{15d \sqrt{a + a \cos(c + dx)}} + \frac{2a^2(10A + 7B) \sqrt{a + a \cos(c + dx)} \sec^{5/2}(c + dx) \sin(c + dx)}{35d} + \frac{2aA(a + a \cos(c + dx))^{3/2} \sec^{7/2}(c + dx) \sin(c + dx)}{7d}$$

```
output 2/7*a*A*(a+a*cos(d*x+c))^(3/2)*sec(d*x+c)^(7/2)*sin(d*x+c)/d+2/15*a^3*(10*
A+11*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/35*a^2*(10*
A+7*B)*sec(d*x+c)^(5/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+2/105*a^3*(230
*A+301*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```


3.513.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.57

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} (290A + 196B + (930A + 987B) \cos(c + dx) + 2(115A + 98B) \cos(2(c + dx)))}{210d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x]])*(290*A + 196*B + (930*A + 987*B)*Cos[c + d*x] + 2*(115*A + 98*B)*Cos[2*(c + d*x)] + 230*A*Cos[3*(c + d*x)] + 301*B*Cos[3*(c + d*x)])*Sec[c + d*x]^(7/2)*Tan[(c + d*x)/2])/(210*d)`

3.513.3 Rubi [A] (verified)Time = 1.25 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.17, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3250}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{9/2}(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{9/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2}(A + B \cos(c + dx))}{\cos^{9/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx \\ & \quad \downarrow \text{3454} \end{aligned}$$

3.513. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{7}\int\frac{(\cos(c+dx)a+a)^{3/2}(a(10A+7B)+a(2A+7B)\cos(c+dx))}{2\cos^{7/2}(c+dx)}dx+\frac{2aA\sin(c+dx)}{2\cos^{7/2}(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{(\cos(c+dx)a+a)^{3/2}(a(10A+7B)+a(2A+7B)\cos(c+dx))}{\cos^{7/2}(c+dx)}dx+\frac{2aA\sin(c+dx)}{\cos^{7/2}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(10A+7B)+a(2A+7B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx+\frac{2aA\cos(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}}\right)$$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2}{5}\int\frac{\sqrt{\cos(c+dx)a+a}(7(10A+11B)a^2+(30A+49B)\cos(c+dx)a^2)}{2\cos^{5/2}(c+dx)}dx+\frac{2aA\cos(c+dx)}{2\cos^{5/2}(c+dx)}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\int\frac{\sqrt{\cos(c+dx)a+a}(7(10A+11B)a^2+(30A+49B)\cos(c+dx)a^2)}{\cos^{5/2}(c+dx)}dx+\frac{2aA\cos(c+dx)}{\cos^{5/2}(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(7(10A+11B)a^2+(30A+49B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}}\right)\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}a^2(230A+301B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\cos^{3/2}(c+dx)}dx+\frac{14a^3(10A+11B)\sin(c+dx)}{3d\cos^{3/2}(c+dx)\sqrt{a\cos(c+dx)}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{1}{3}a^2(230A+301B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{14a^3(10A+11B)\sin(c+dx+\frac{\pi}{2})}{3d\cos^{3/2}(c+dx)\sqrt{a\cos(c+dx+\frac{\pi}{2})}}\right)\right)\right)$$

↓ 3250

3.513. $\int(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sec^{9/2}(c+dx)dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{14a^3(10A+11B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}+\frac{2a^3(230A+301B)\sin(c+dx)}{3d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*a^2*(10*A + 7*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + ((14*a^3*(10*A + 11*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) + (2*a^3*(230*A + 301*B)*Sin[c + d*x])/(3*d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/5)/7)`

3.513.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3250 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[-2*b^2*(Cos[e + f*x]/(f*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_.))^(p_.)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_.)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3454 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.513.4 Maple [A] (verified)

Time = 2.09 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.57

$$\frac{2a^2 \cot(dx + c) (\cos(dx + c) - 1) ((230 \cos^3(dx + c)) + 115 \cos^2(dx + c) + 60 \cos(dx + c) + 15) A + c}{105d}$$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x)
```

```
output -2/105*a^2/d*cot(d*x+c)*(cos(d*x+c)-1)*((230*cos(d*x+c)^3+115*cos(d*x+c)^2
+60*cos(d*x+c)+15)*A+cos(d*x+c)*(301*cos(d*x+c)^2+98*cos(d*x+c)+21)*B)*(a*
(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(9/2)
```

$$3.513. \quad \int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$$

3.513.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.63

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{2 \left((230A + 301B)a^2 \cos(dx + c)^3 + (115A + 98B)a^2 \cos(dx + c)^2 + 3(20A + 7B)a^2 \cos(dx + c) + 15Aa^2 \right) \sqrt{\cos(dx + c)}}{105 (d \cos(dx + c)^4 + d \cos(dx + c)^3) \sqrt{\cos(dx + c)}}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorith="fricas")`

output `2/105*((230*A + 301*B)*a^2*cos(d*x + c)^3 + (115*A + 98*B)*a^2*cos(d*x + c)^2 + 3*(20*A + 7*B)*a^2*cos(d*x + c) + 15*A*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/((d*cos(d*x + c)^4 + d*cos(d*x + c)^3)*sqrt(cos(d*x + c)))`

3.513.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(9/2),x)`

output `Timed out`

3.513.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 488 vs. $2(157) = 314$.

Time = 0.36 (sec) , antiderivative size = 488, normalized size of antiderivative = 2.70

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \frac{8 \left(5 \left(\frac{21 \sqrt{2} a^{5/2} \sin(dx+c)}{\cos(dx+c)+1} - \frac{56 \sqrt{2} a^{5/2} \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{63 \sqrt{2} a^{5/2} \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{36 \sqrt{2} a^{5/2} \sin(dx+c)^7}{(\cos(dx+c)+1)^7} + \frac{8 \sqrt{2} a^{5/2} \sin(dx+c)^9}{(\cos(dx+c)+1)^9} \right) A \left(\frac{\sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 1 \right) + \frac{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(\frac{2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)}{\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(-\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)^{9/2} \left(\frac{2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{\sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 1 \right)}$$

3.513. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algo
rithm="maxima")`

output `8/105*(5*(21*sqrt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 56*sqrt(2)*
a^(5/2)*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 63*sqrt(2)*a^(5/2)*sin(d*x +
c)^5/(cos(d*x + c) + 1)^5 - 36*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x +
c) + 1)^7 + 8*sqrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9)*A*(sin(
d*x + c)^2/(cos(d*x + c) + 1)^2 + 1)^2/((sin(d*x + c)/(cos(d*x + c) + 1) +
1)^(9/2)*(-sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(9/2)*(2*sin(d*x + c)^2/(
cos(d*x + c) + 1)^2 + sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 1)) + 7*(15*sq
rt(2)*a^(5/2)*sin(d*x + c)/(cos(d*x + c) + 1) - 50*sqrt(2)*a^(5/2)*sin(d*x
+ c)^3/(cos(d*x + c) + 1)^3 + 63*sqrt(2)*a^(5/2)*sin(d*x + c)^5/(cos(d*x
+ c) + 1)^5 - 36*sqrt(2)*a^(5/2)*sin(d*x + c)^7/(cos(d*x + c) + 1)^7 + 8*s
qrt(2)*a^(5/2)*sin(d*x + c)^9/(cos(d*x + c) + 1)^9)*B*(sin(d*x + c)^2/(cos
(d*x + c) + 1)^2 + 1)^2/((sin(d*x + c)/(cos(d*x + c) + 1) + 1)^(9/2)*(-sin
(d*x + c)/(cos(d*x + c) + 1) + 1)^(9/2)*(2*sin(d*x + c)^2/(cos(d*x + c) +
1)^2 + sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 1)))/d`

3.513.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algo
rithm="giac")`

output Timed out

3.513.9 Mupad [B] (verification not implemented)

Time = 5.25 (sec) , antiderivative size = 579, normalized size of antiderivative = 3.20

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \frac{\sqrt{\frac{1}{\frac{e^{-c \operatorname{li}-dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li}+dx \operatorname{li}}}{2}}}}{\left(\frac{a^2 \sqrt{a+a \left(\frac{e^{-c \operatorname{li}-dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li}+dx \operatorname{li}}}{2} \right)}}{105 d} (230 A+301 B) 2i - \frac{B a^2 e^{c \operatorname{li}+dx \operatorname{li}} \sqrt{a+a \left(\frac{e^{-c \operatorname{li}-dx \operatorname{li}}}{2} + \frac{e^{c \operatorname{li}+dx \operatorname{li}}}{2} \right)}}{d} \right)}$$

3.513. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + a*cos(c + d*x))^(5/2),x)`

output `((1/(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*((a^2*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(230*A + 301*B)*2i)/(105*d) - (B*a^2*exp(c*1i + d*x*1i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*2i)/d + (B*a^2*exp(c*6i + d*x*6i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*2i)/d - (a^2*exp(c*3i + d*x*3i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(10*A + 17*B)*2i)/(3*d) + (a^2*exp(c*4i + d*x*4i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(10*A + 17*B)*2i)/(3*d) + (a^2*exp(c*2i + d*x*2i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(100*A + 113*B)*2i)/(15*d) - (a^2*exp(c*5i + d*x*5i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(100*A + 113*B)*2i)/(15*d) - (a^2*exp(c*7i + d*x*7i)*(a + a*(exp(- c*1i - d*x*1i)/2 + exp(c*1i + d*x*1i)/2))^(1/2)*(230*A + 301*B)*2i)/(105*d)))/(exp(c*1i + d*x*1i) + 3*exp(c*2i + d*x*2i) + 3*exp(c*3i + d*x*3i) + 3*exp(c*4i + d*x*4i) + 3*exp(c*5i + d*x*5i) + exp(c*6i + d*x*6i) + exp(c*7i + d*x*7i) + 1)`

3.514 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$

3.514.1 Optimal result	4673
3.514.2 Mathematica [A] (verified)	4674
3.514.3 Rubi [A] (verified)	4674
3.514.4 Maple [A] (verified)	4677
3.514.5 Fricas [A] (verification not implemented)	4678
3.514.6 Sympy [F(-1)]	4678
3.514.7 Maxima [B] (verification not implemented)	4679
3.514.8 Giac [F(-1)]	4679
3.514.9 Mupad [F(-1)]	4680

3.514.1 Optimal result

Integrand size = 35, antiderivative size = 192

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{2a^{5/2}B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} + \frac{2a^3(32A + 35B) \sqrt{\sec(c + dx)} \sin(c + dx)}{15d \sqrt{a + a \cos(c + dx)}} + \frac{2a^2(8A + 5B) \sqrt{a + a \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{15d} + \frac{2aA(a + a \cos(c + dx))^{3/2} \sec^{5/2}(c + dx) \sin(c + dx)}{5d}$$

```
output 2/5*a*A*(a+a*cos(d*x+c))^(3/2)*sec(d*x+c)^(5/2)*sin(d*x+c)/d+2/15*a^2*(8*A
+5*B)*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d+2*a^(5/2)*B*arc
sin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)
^(1/2)/d+2/15*a^3*(32*A+35*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c
))^(1/2)
```


3.514.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.68

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^{5/2}(c + dx) \left(30\sqrt{2}B \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) \cos^{5/2}(c + dx)\right)}{30d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^(5/2)*(30*sqrt[2]*B*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^(5/2) + 2*(49*A + 40*B + 2*(14*A + 5*B)*Cos[c + d*x] + (43*A + 40*B)*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(30*d)`

3.514.3 Rubi [A] (verified)Time = 1.18 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3454, 27, 3042, 3459, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{7/2}(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2}(A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx \end{aligned}$$

3.514. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{5}\int\frac{(\cos(c+dx)a+a)^{3/2}(a(8A+5B)+5aB\cos(c+dx))}{2\cos^{5/2}(c+dx)}dx+\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)}{5d\cos^{5/2}(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\int\frac{(\cos(c+dx)a+a)^{3/2}(a(8A+5B)+5aB\cos(c+dx))}{\cos^{5/2}(c+dx)}dx+\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)}{5d\cos^{5/2}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\int\frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(8A+5B)+5aB\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{5/2}}dx+\frac{2aA\sin(c+dx)(a\cos(c+dx)+a)}{5d\cos^{5/2}(c+dx)}\right)$$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{2}{3}\int\frac{\sqrt{\cos(c+dx)a+a}((32A+35B)a^2+15B\cos(c+dx)a^2)}{2\cos^{3/2}(c+dx)}dx+\frac{2a^2(8A+5B)\sin(c+dx)}{5d\cos^{3/2}(c+dx)}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{\sqrt{\cos(c+dx)a+a}((32A+35B)a^2+15B\cos(c+dx)a^2)}{\cos^{3/2}(c+dx)}dx+\frac{2a^2(8A+5B)\sin(c+dx)}{5d\cos^{3/2}(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((32A+35B)a^2+15B\sin(c+dx+\frac{\pi}{2})a^2)}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2a^2(8A+5B)\sin(c+dx)}{5d\cos^{3/2}(c+dx)}\right)\right)$$

↓ 3459

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(15a^2B\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{2a^3(32A+35B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)\right)+\frac{2a^2(8A+5B)\sin(c+dx)}{5d\cos^{3/2}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(15a^2B\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2a^3(32A+35B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)\right)+\frac{2a^2(8A+5B)\sin(c+dx)}{5d\cos^{3/2}(c+dx)}\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2a^3(32A+35B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}-\frac{30a^2B\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}\frac{d}{d}\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}}\right)}{d}\right)\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{2a^2(8A+5B)\sin(c+dx)\sqrt{a\cos(c+dx)+a}}{3d\cos^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(\frac{30a^{5/2}B\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x]/(5*d*Cos[c + d*x]^(5/2)) + ((2*a^2*(8*A + 5*B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + ((30*a^(5/2)*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (2*a^3*(32*A + 35*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/3)/5)`

3.514.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3454 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[
e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c +
a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp
[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B
*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f
, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
&& GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0
])
```

```
rule 3459 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[(-b^2)*(B*c - A*d)*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)
*(b*c + a*d)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*
c - 2*a*d*(n + 1)))/(2*d*(n + 1)*(b*c + a*d)) Int[Sqrt[a + b*Sin[e + f*x]
]*(c + d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x
] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n,
-1]
```

3.514.4 Maple [A] (verified)

Time = 2.44 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.23

$$2a^2 \left(\sec^{\frac{7}{2}}(dx + c) \right) \sqrt{a(1 + \cos(dx + c))} \left(15B \cos^4(dx + c) \right) \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \arctan \left(\tan(dx + c) \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \right)$$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x)
```

$$3.514. \quad \int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

output $2/15*a^2/d*sec(d*x+c)^{(7/2)}*(a*(1+cos(d*x+c)))^{(1/2)}/(1+cos(d*x+c))*(15*B*cos(d*x+c)^4*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)})+15*B*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)}*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^{(1/2)})*cos(d*x+c)^3+43*A*cos(d*x+c)^3*sin(d*x+c)+40*B*sin(d*x+c)*cos(d*x+c)^3+14*A*sin(d*x+c)*cos(d*x+c)^2+5*B*sin(d*x+c)*cos(d*x+c)^2+3*A*sin(d*x+c)*cos(d*x+c))$

3.514.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.84

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx =$$

$$\frac{2 \left(15 (Ba^2 \cos(dx + c)^3 + Ba^2 \cos(dx + c)^2) \sqrt{a} \arctan \left(\frac{\sqrt{a} \cos(dx+c) + a \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - \frac{((43A+40B)a^2 \cos(dx+c)^2)}{15 (d \cos(dx + c)^3 + d \cos(dx + c)^2)} \right)}{15 (d \cos(dx + c)^3 + d \cos(dx + c)^2)}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorith="fricas")`

output $-2/15*(15*(B*a^2*cos(d*x + c)^3 + B*a^2*cos(d*x + c)^2)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - ((43*A + 40*B)*a^2*cos(d*x + c)^2 + (14*A + 5*B)*a^2*cos(d*x + c) + 3*A*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^3 + d*cos(d*x + c)^2)$

3.514.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output Timed out

3.514. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

3.514.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1713 vs. $2(164) = 328$.

Time = 0.53 (sec) , antiderivative size = 1713, normalized size of antiderivative = 8.92

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algo
rithm="maxima")`

output `1/30*(5*(10*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2
*c) + 1)*a^(5/2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))
+ 3*((a^2*cos(2*d*x + 2*c)^2 + a^2*sin(2*d*x + 2*c)^2 + 2*a^2*cos(2*d*x +
2*c) + a^2)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x
+ 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*s
in(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(
sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c),
cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d
*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) +
1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + sin(1/2*arctan
2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c
, cos(2*d*x + 2*c)))) + 1) - (a^2*cos(2*d*x + 2*c)^2 + a^2*sin(2*d*x + 2*
c)^2 + 2*a^2*cos(2*d*x + 2*c) + a^2)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d
*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
+ 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*
arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2
*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c)))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin...`

3.514.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algo
rithm="giac")`

output Timed out

3.514.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + a \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^(5/2), x)`

3.515 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$

3.515.1 Optimal result	4681
3.515.2 Mathematica [A] (verified)	4682
3.515.3 Rubi [A] (verified)	4682
3.515.4 Maple [A] (verified)	4685
3.515.5 Fricas [A] (verification not implemented)	4686
3.515.6 Sympy [F(-1)]	4686
3.515.7 Maxima [B] (verification not implemented)	4687
3.515.8 Giac [F(-1)]	4687
3.515.9 Mupad [F(-1)]	4688

3.515.1 Optimal result

Integrand size = 35, antiderivative size = 193

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \frac{a^{5/2}(2A + 5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} - \frac{a^3(14A + 3B) \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)}} + \frac{2a^2(2A + B)\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)} \sin(c + dx)}{d} + \frac{2aA(a + a \cos(c + dx))^{3/2} \sec^{3/2}(c + dx) \sin(c + dx)}{3d}$$

```
output 2/3*a*A*(a+a*cos(d*x+c))^(3/2)*sec(d*x+c)^(3/2)*sin(d*x+c)/d-1/3*a^3*(14*A
+3*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+a^(5/2)*(2*A+5*
B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(
d*x+c)^(1/2)/d+2*a^2*(2*A+B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)*sec(d*x+c)^(
1/2)/d
```


3.515.2 Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.67

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sec^{3/2}(c + dx) \left(3\sqrt{2}(2A + 5B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + dx\right)}{6d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(a^2*sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sec[c + d*x]^(3/2)*(3*sqrt[2]*(2*A + 5*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Cos[c + d*x]^(3/2) + (4*A + 3*B + 4*(8*A + 3*B)*Cos[c + d*x] + 3*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(6*d)`

3.515.3 Rubi [A] (verified)Time = 1.22 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3454, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{5/2}(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2}(A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx \end{aligned}$$

3.515. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{3}\int\frac{(\cos(c+dx)a+a)^{3/2}(3a(2A+B)-a(2A-3B)\cos(c+dx))}{2\cos^{3/2}(c+dx)}dx+\frac{2aA\sin(c+dx)}{\cos(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\int\frac{(\cos(c+dx)a+a)^{3/2}(3a(2A+B)-a(2A-3B)\cos(c+dx))}{\cos^{3/2}(c+dx)}dx+\frac{2aA\sin(c+dx)}{\cos(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\int\frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(3a(2A+B)-a(2A-3B)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}}dx+\frac{2aA\cos(c+dx)}{\sin(c+dx+\frac{\pi}{2})}\right)$$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(2\int\frac{\sqrt{\cos(c+dx)a+a}(a^2(10A+9B)-a^2(14A+3B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}}dx+\frac{6a^2(2A+B)\sin(c+dx)}{\cos(c+dx)}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{\sqrt{\cos(c+dx)a+a}(a^2(10A+9B)-a^2(14A+3B)\cos(c+dx))}{\sqrt{\cos(c+dx)}}dx+\frac{6a^2(2A+B)\sin(c+dx)}{\cos(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a^2(10A+9B)-a^2(14A+3B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{6a^2(2A+B)\cos(c+dx)}{\sin(c+dx+\frac{\pi}{2})}\right)\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{3}{2}a^2(2A+5B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx-\frac{a^3(14A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{3}{2}a^2(2A+5B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx-\frac{a^3(14A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(-\frac{3a^2(2A+5B)\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}-\frac{a^3(14A+3B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{3a^{5/2}(2A+5B)\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}-\frac{a^3(14A+3B)\sin(c+dx)\sqrt{\cos(c+dx)+a}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x]/(3*d*Cos[c + d*x]^(3/2)) + ((3*a^(5/2)*(2*A + 5*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (a^3*(14*A + 3*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x]/(d*Sqrt[a + a*Cos[c + d*x]]) + (6*a^2*(2*A + B)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x]/(d*Sqrt[Cos[c + d*x]]))/3)`

3.515.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.515.4 Maple [A] (verified)

Time = 2.51 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.73

$$a^2 \left(\sec^{\frac{5}{2}}(dx + c) \right) \sqrt{a(1 + \cos(dx + c))} \left(6A \cos^3(dx + c) \arctan \left(\tan(dx + c) \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \right) \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \right)$$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x)`

3.515. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

output $\frac{1}{3}a^2/d\sec(dx+c)^{5/2}(a(1+\cos(dx+c)))^{1/2}/(1+\cos(dx+c))*(6A\cos(dx+c)^3\arctan(\tan(dx+c)(\cos(dx+c)/(1+\cos(dx+c))))^{1/2}+(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}+15B(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}\arctan(\tan(dx+c)(\cos(dx+c)/(1+\cos(dx+c))))^{1/2}+\cos(dx+c)^3+6A\cos(dx+c)^2\arctan(\tan(dx+c)(\cos(dx+c)/(1+\cos(dx+c))))^{1/2}+(\cos(dx+c)/(1+\cos(dx+c))))^{1/2}+3B\sin(dx+c)\cos(dx+c)^3+15B(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}\arctan(\tan(dx+c)(\cos(dx+c)/(1+\cos(dx+c))))^{1/2}+\cos(dx+c)^2+16A\sin(dx+c)\cos(dx+c)^2+6B\sin(dx+c)\cos(dx+c)^2+2A\sin(dx+c)\cos(dx+c))$

3.515.5 Fricas [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.86

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx =$$

$$\frac{3((2A + 5B)a^2 \cos(dx + c)^2 + (2A + 5B)a^2 \cos(dx + c))\sqrt{a} \arctan\left(\frac{\sqrt{a} \cos(dx+c) + a\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - \frac{(3Ba^2 \cos(dx+c) + 3Aa^2 \cos(dx+c))\sqrt{a}}{3(d \cos(dx+c))^2 + d \cos(dx+c)}}{3(d \cos(dx+c))^2 + d \cos(dx+c)}$$

input `integrate((a+a*cos(dx+c))^(5/2)*(A+B*cos(dx+c))*sec(dx+c)^(5/2),x, algorithm="fricas")`

output $-1/3*(3*((2A + 5B)*a^2*\cos(dx + c)^2 + (2A + 5B)*a^2*\cos(dx + c))*\sqrt{a}*\arctan(\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)})/(\sqrt{a}*\sin(dx + c))) - (3*B*a^2*\cos(dx + c)^2 + 2*(8*A + 3*B)*a^2*\cos(dx + c) + 2*A*a^2)*\sqrt{a*\cos(dx + c) + a}*\sin(dx + c)/\sqrt{\cos(dx + c)})/(d*\cos(dx + c)^2 + d*\cos(dx + c))$

3.515.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(dx+c))**(5/2)*(A+B*cos(dx+c))*sec(dx+c)**(5/2),x)`

output Timed out

3.515. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

3.515.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2780 vs. $2(167) = 334$.

Time = 0.62 (sec) , antiderivative size = 2780, normalized size of antiderivative = 14.40

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algo
rithm="maxima")`

output `1/12*(2*(30*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c)
+ 1)^(3/4)*a^(5/2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)
) - 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(
1/4)*((12*a^2*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(2*d
*x + 2*c) - 3*a^2*sin(2*d*x + 2*c) - 4*(3*a^2*cos(2*d*x + 2*c) + 4*a^2)*si
n(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*cos(3/2*arctan2(sin(2*
d*x + 2*c), cos(2*d*x + 2*c) + 1)) + (12*a^2*sin(2*d*x + 2*c)*sin(3/2*arct
an2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 3*a^2*cos(2*d*x + 2*c) - a^2 +
4*(3*a^2*cos(2*d*x + 2*c) + 4*a^2)*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2
*d*x + 2*c))))*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*s
qrt(a) + 3*((a^2*cos(2*d*x + 2*c)^2 + a^2*sin(2*d*x + 2*c)^2 + 2*a^2*cos(2
*d*x + 2*c) + a^2)*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*co
s(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*a
rctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*
cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + sin(1/2
*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c)))) + 1) - (a^2*cos(2*d*x + 2*c)^2 + a^2*sin(...`

3.515.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algo
rithm="giac")`

3.515. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

output Timed out

3.515.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + a \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(5/2), x)`

3.516 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$

3.516.1 Optimal result	4689
3.516.2 Mathematica [A] (verified)	4690
3.516.3 Rubi [A] (verified)	4690
3.516.4 Maple [A] (verified)	4694
3.516.5 Fricas [A] (verification not implemented)	4694
3.516.6 Sympy [F(-1)]	4695
3.516.7 Maxima [F(-1)]	4695
3.516.8 Giac [F(-1)]	4695
3.516.9 Mupad [F(-1)]	4696

3.516.1 Optimal result

Integrand size = 35, antiderivative size = 198

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{a^{5/2}(20A + 19B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{4d} - \frac{a^3(4A - 9B) \sin(c + dx)}{4d \sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} - \frac{a^2(4A - B) \sqrt{a + a \cos(c + dx)} \sin(c + dx)}{2d \sqrt{\sec(c + dx)}} + \frac{2aA(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

output

```
-1/4*a^3*(4*A-9*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)-1/2*a^2*(4*A-B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)+2*a*A*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)*sec(d*x+c)^(1/2)/d+1/4*a^(5/2)*(20*A+19*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```


3.516.2 Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.64

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{a^2 \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(\sqrt{2}(20A + 19B) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 8d\right)}{8d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(a^2*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(Sqrt[2]*(20*A + 19*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]]*Sqrt[Cos[c + d*x]] + 2*(8*A + B + (4*A + 11*B)*Cos[c + d*x] + B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2])/(8*d)`

3.516.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3454, 27, 3042, 3455, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{3/2}(c + dx)(a \cos(c + dx) + a)^{5/2}(A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2}(A + B \cos(c + dx))}{\cos^{3/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \end{aligned}$$

3.516. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$

↓ 3454

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(2\int\frac{(\cos(c+dx)a+a)^{3/2}(a(4A+B)-a(4A-B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}}dx+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\int\frac{(\cos(c+dx)a+a)^{3/2}(a(4A+B)-a(4A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)}}dx+\frac{2aA\sin(c+dx)}{d\sqrt{\cos(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\int\frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(4A+B)-a(4A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2aA\cos(c+dx+\frac{\pi}{2})}{d\sqrt{\sin(c+dx+\frac{\pi}{2})}}\right)$$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\int\frac{\sqrt{\cos(c+dx)a+a}(a^2(12A+5B)-a^2(4A-9B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}}dx-\frac{a^2(4A-9B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\int\frac{\sqrt{\cos(c+dx)a+a}(a^2(12A+5B)-a^2(4A-9B)\cos(c+dx))}{\sqrt{\cos(c+dx)}}dx-\frac{a^2(4A-9B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}(a^2(12A+5B)-a^2(4A-9B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx-\frac{a^2(4A-9B)\cos(c+dx+\frac{\pi}{2})}{4d\sqrt{\sin(c+dx+\frac{\pi}{2})}}\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}a^2(20A+19B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx-\frac{a^3(4A-9B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}a^2(20A+19B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx-\frac{a^3(4A-9B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)$$

↓ 3253

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{1}{4} \left(-\frac{a^2(20A + 19B) \int \frac{1}{\sqrt{1 - \frac{a \sin^2(c + dx)}{\cos(c + dx)a + a}}} d\left(-\frac{a \sin(c + dx)}{\sqrt{\cos(c + dx)a + a}}\right)}{d} - \frac{a^3(4A - 9B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) \right)$$

↓ 223

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(-\frac{a^2(4A - B) \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{a \cos(c + dx) + a}}{2d} + \frac{1}{4} \left(\frac{a^{5/2}(20A + 19B) \operatorname{ArcSin}\left[\frac{\sqrt{a} \sin(c + dx)}{\sqrt{a \cos(c + dx) + a}}\right]}{d} - \frac{a^3(4A - 9B) \sin(c + dx)}{d\sqrt{a \cos(c + dx) + a}} \right) \right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/2*(a^2*(4*A - B)*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/d + (2*a*A*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) + ((a^(5/2)*(20*A + 19*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d - (a^3*(4*A - 9*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4`

3.516.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

3.516. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3454 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b^2)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(b*c + a*d))), x] - Simp[b/(d*(n + 1)*(b*c + a*d)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[a*A*d*(m - n - 2) - B*(a*c*(m - 1) + b*d*(n + 1)) - (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(n + 1)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3455 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*(A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.516.4 Maple [A] (verified)

Time = 22.91 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.57

method	result
default	$a^2 \left(\sec^{\frac{3}{2}}(dx+c) \right) \left(20A \cos(dx+c) \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 2B \sin(dx+c) (\cos^2(dx+c)) + 19B \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)$
parts	$\frac{A \left(\sec^{\frac{3}{2}}(dx+c) \right) \left(5 \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) + \cos(dx+c) \sin(dx+c) + 5 \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right)}{d(1+\cos(dx+c))}$

input `int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `1/4*a^2/d*sec(d*x+c)^(3/2)*(20*A*cos(d*x+c)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2*B*sin(d*x+c)*cos(d*x+c)^2+19*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2)*cos(d*x+c)+4*A*sin(d*x+c)*cos(d*x+c)+20*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2)+11*B*sin(d*x+c)*cos(d*x+c)+19*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2)+8*A*sin(d*x+c)*cos(d*x+c)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))`

3.516.5 Fracas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.74

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \frac{((20A + 19B)a^2 \cos(dx + c) + (20A + 19B)a^2) \sqrt{a} \arctan \left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)} \right) - (2Ba^2 \cos(dx+c)^2 + 4a^2 \cos(dx+c))}{4(d \cos(dx + c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,algorithm="fracas")`

output `-1/4*(((20*A + 19*B)*a^2*cos(d*x + c) + (20*A + 19*B)*a^2)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (2*B*a^2*cos(d*x + c)^2 + (4*A + 11*B)*a^2*cos(d*x + c) + 8*A*a^2)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c) + d)`

3.516. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.516.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.516.7 Maxima [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `Timed out`

3.516.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.516.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + a \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(5/2), x)`

3.517 $\int (a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))\sqrt{\sec(c+dx)}$

3.517.1 Optimal result	4697
3.517.2 Mathematica [A] (verified)	4698
3.517.3 Rubi [A] (verified)	4698
3.517.4 Maple [A] (verified)	4702
3.517.5 Fricas [A] (verification not implemented)	4702
3.517.6 Sympy [F(-1)]	4703
3.517.7 Maxima [B] (verification not implemented)	4703
3.517.8 Giac [F(-1)]	4704
3.517.9 Mupad [F(-1)]	4705

3.517.1 Optimal result

Integrand size = 35, antiderivative size = 200

$$\int (a + a \cos(c + dx))^{5/2}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx = \frac{a^{5/2}(38A + 25B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{8d} + \frac{a^3(54A + 49B) \sin(c + dx)}{24d\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)}} + \frac{a^2(2A + 3B)\sqrt{a + a \cos(c + dx)} \sin(c + dx)}{4d\sqrt{\sec(c + dx)}} + \frac{aB(a + a \cos(c + dx))^{3/2} \sin(c + dx)}{3d\sqrt{\sec(c + dx)}}$$

output

```
1/3*a*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+1/24*a^3*(54*
A+49*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/4*a^2*(2*A+
3*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)+1/8*a^(5/2)*(38*
A+25*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)
*sec(d*x+c)^(1/2)/d
```


3.517.2 Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.70

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \frac{a^2 \sqrt{\cos(c + dx)} \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(3\sqrt{2}\right)}{48d}$$

input `Integrate[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(a^2*Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(3*Sqrt[2]*(38*A + 25*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(66*A + 79*B + 2*(6*A + 17*B)*Cos[c + d*x] + 4*B*Cos[2*(c + d*x)])*Sin[(c + d*x)/2]))/(48*d)`

3.517.3 Rubi [A] (verified)Time = 1.22 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\sec(c + dx)} (a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\cos(c + dx)a + a)^{5/2} (A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx \end{aligned}$$

3.517. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\int\frac{(\cos(c+dx)a+a)^{3/2}(a(6A+B)+3a(2A+3B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}}dx+\frac{aB\sin(c+dx)}{2}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\int\frac{(\cos(c+dx)a+a)^{3/2}(a(6A+B)+3a(2A+3B)\cos(c+dx))}{\sqrt{\cos(c+dx)}}dx+\frac{aB\sin(c+dx)}{2}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\int\frac{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}(a(6A+B)+3a(2A+3B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{aB\cos(c+dx)}{2}\right)$$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{2}\int\frac{\sqrt{\cos(c+dx)a+a}((30A+13B)a^2+(54A+49B)\cos(c+dx)a^2)}{2\sqrt{\cos(c+dx)}}dx+\frac{3a^2(38A+25B)\sin(c+dx)}{2}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\int\frac{\sqrt{\cos(c+dx)a+a}((30A+13B)a^2+(54A+49B)\cos(c+dx)a^2)}{\sqrt{\cos(c+dx)}}dx+\frac{3a^2(38A+25B)\sin(c+dx)}{2}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}((30A+13B)a^2+(54A+49B)\sin(c+dx+\frac{\pi}{2})a^2)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{3a^2(38A+25B)\cos(c+dx)}{2}\right)\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{3}{2}a^2(38A+25B)\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{a^3(54A+49B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{3}{2}a^2(38A+25B)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{a^3(54A+49B)\sin(c+dx)\sqrt{\sin(c+dx+\frac{\pi}{2})}}{d\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{a^3(54A+49B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}-\frac{3a^2(38A+25B)\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+1}}}}{d}\right)\right)\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{3a^2(2A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}{2d}+\frac{1}{4}\left(\frac{3a^{5/2}(38A+25B)\operatorname{ArcSin}\left[\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right]}{d}+\frac{a^3(54A+49B)\sqrt{\cos(c+dx)}\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}}\right)\right)\right)$$

input `Int[(a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + ((3*a^2*(2*A + 3*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + ((3*a^(5/2)*(38*A + 25*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a^3*(54*A + 49*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/4)/6)`

3.517.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3455 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3460 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]`

3.517.4 Maple [A] (verified)

Time = 22.48 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.42

method	result
default	$a^2 \left(8B \cos^2(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 12A \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 34B \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \dots \right)$
parts	$\frac{A \left(2 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 11 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 19 \arctan \left(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right) \right) (\sqrt{\sec(dx+c)}) \sqrt{a(1+\cos(dx+c))}}{4d(1+\cos(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 1/24*a^2/d*(8*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+
12*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+34*B*cos(d*x+
c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+66*A*(cos(d*x+c)/(1+cos(d*
x+c)))^(1/2)*sin(d*x+c)+75*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+
114*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+75*B*arctan(tan
(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*sec(d*x+c)^(1/2)*(a*(1+cos(d*x
+c)))^(1/2)*cos(d*x+c)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

3.517.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.82

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \frac{3((38A + 25B)a^2 \cos(dx + c) + (38A + 25B)a^2) \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c) + a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - (8Ba^2 \cos(dx+c)^3 + \dots)}{24(d \cos(dx + c) + d)}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algo
rithm="fricas")
```

```
output -1/24*(3*((38*A + 25*B)*a^2*cos(d*x + c) + (38*A + 25*B)*a^2)*sqrt(a)*arct
an(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (
8*B*a^2*cos(d*x + c)^3 + 2*(6*A + 17*B)*a^2*cos(d*x + c)^2 + 3*(22*A + 25*
B)*a^2*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x +
c)))/(d*cos(d*x + c) + d)
```

3.517. $\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

3.517.6 Sympy [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Timed out`

3.517.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3071 vs. $2(170) = 340$.

Time = 0.70 (sec) , antiderivative size = 3071, normalized size of antiderivative = 15.36

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `1/96*(6*(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*((a^2*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(2*d*x + 2*c) + a^2*sin(2*d*x + 2*c) - (a^2*cos(2*d*x + 2*c) - 10*a^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) + (a^2*sin(2*d*x + 2*c)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) - a^2*cos(2*d*x + 2*c) + 10*a^2 + (a^2*cos(2*d*x + 2*c) - 10*a^2)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)))*sqrt(a) + 19*(a^2*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))), (cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))))) + 1) - a^2*arctan2((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)^(1/4)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1)) - cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) + 1))*sin(1/2*arc...`

3.517.8 Giac [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.517.9 Mupad [F(-1)]

Timed out.

$$\int (a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + a \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(5/2), x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(5/2), x)`

3.518
$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.518.1 Optimal result 4706
 3.518.2 Mathematica [A] (verified) 4707
 3.518.3 Rubi [A] (verified) 4707
 3.518.4 Maple [A] (verified) 4711
 3.518.5 Fricas [A] (verification not implemented) 4712
 3.518.6 Sympy [F(-1)] 4712
 3.518.7 Maxima [B] (verification not implemented) 4713
 3.518.8 Giac [F] 4713
 3.518.9 Mupad [F(-1)] 4714

3.518.1 Optimal result

Integrand size = 35, antiderivative size = 247

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx = \frac{a^{5/2}(200A+163B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)}}{64d}$$

$$+ \frac{a^3(104A+95B) \sin(c+dx)}{96d \sqrt{a+a \cos(c+dx)} \sec^{3/2}(c+dx)} + \frac{a^2(8A+11B) \sqrt{a+a \cos(c+dx)} \sin(c+dx)}{24d \sec^{3/2}(c+dx)}$$

$$+ \frac{aB(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{4d \sec^{3/2}(c+dx)} + \frac{a^3(200A+163B) \sin(c+dx)}{64d \sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}}$$

output

```
1/4*a*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+1/96*a^3*(104
*A+95*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/24*a^2*(8*
A+11*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+1/64*a^3*(200
*A+163*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/64*a^(5/2
)*(200*A+163*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+
c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.518.2 Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.64

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \frac{a^2 \sqrt{\cos(c + dx)} \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{s}}{\dots}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(a^2*Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(3*Sqrt[2]*(200*A + 163*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(632*A + 581*B + (272*A + 362*B)*Cos[c + d*x] + 4*(8*A + 23*B)*Cos[2*(c + d*x)] + 12*B*Cos[3*(c + d*x)])*Sin[(c + d*x)/2]))/(384*d)`

3.518.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} (\cos(c + dx) a + a)^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} (\sin(c + dx + \frac{\pi}{2}) a + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2})) dx \end{aligned}$$

3.518. $\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\int\frac{1}{2}\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}(a(8A+3B)+a(8A+11B)\cos(c+dx))dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\int\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}(a(8A+3B)+a(8A+11B)\cos(c+dx))dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{3/2}\left(a(8A+3B)+a(8A+11B)\cos(c+dx)\right)dx\right)$$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{3}\int\frac{1}{2}\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}(3(24A+17B)a^2+(104A+95B)\cos(c+dx))dx\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\int\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}(3(24A+17B)a^2+(104A+95B)\cos(c+dx))dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}(3(24A+17B)a^2+(104A+95B)\cos(c+dx))dx\right)\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\left(\frac{3}{4}a^2(200A+163B)\int\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+adx}+\frac{a^3(104A+95B)}{2d\sqrt{a}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\left(\frac{3}{4}a^2(200A+163B)\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx}+\frac{a^3(104A+95B)}{2d\sqrt{a}}\right)\right)\right)$$

↓ 3249

3.518. $\int \frac{(a+a\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\left(\frac{3}{4}a^2(200A+163B)\left(\frac{1}{2}\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right.\right.\right.\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\left(\frac{3}{4}a^2(200A+163B)\left(\frac{1}{2}\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}\right.\right.\right.\right.$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{1}{6}\left(\frac{3}{4}a^2(200A+163B)\left(\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}-\frac{\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}d\left(-\frac{a}{\sqrt{c}}\right)}{d}\right.\right.\right.\right.$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{a^2(8A+11B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}{3d}+\frac{1}{6}\left(\frac{a^3(104A+95B)\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}{2d\sqrt{a\cos(c+dx)+a}}\right.\right.\right.$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(4*d) + ((a^2*(8*A + 11*B)*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(3*d) + ((a^3*(104*A + 95*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (3*a^2*(200*A + 163*B)*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/4)/6)/8)`

3.518.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3249 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[2*n*((b*c + a*d)/(b*(2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3455 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e +
f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3460 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) + (
f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.518.4 Maple [A] (verified)

Time = 21.16 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.42

method	result
default	$\frac{a^2 \left(48B (\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 64A (\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 184B (\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{24d(1+\cos(dx+c)) \sqrt{\sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \left(8 \sin(dx+c) (\cos^2(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 34 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 75 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 75 \arctan \left(\tan(dx+c) \right) \right)}{24d(1+\cos(dx+c)) \sqrt{\sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2), x, method=_RET
URNVERBOSE)
```

3.518.
$$\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

output $1/192*a^2/d*(48*B*cos(d*x+c)^3*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+64*A*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+184*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+272*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+326*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+600*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+489*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+600*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+489*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/sec(d*x+c)^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)$

3.518.5 Fracas [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.74

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$\frac{3((200A + 163B)a^2 \cos(dx + c) + (200A + 163B)a^2)\sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - \frac{(48Ba^2 \cos(dx+c) + 8(8A + 23B)a^2 \cos(dx+c)^3 + 2(136A + 163B)a^2 \cos(dx+c)^2 + 3(200A + 163B)a^2 \cos(dx+c)) \sqrt{a} \cos(dx+c) + a \sin(dx+c)}{192(d \cos(dx+c) + d)}}{192(d \cos(dx+c) + d)}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algo rithm="fracas")`

output $-1/192*(3*((200*A + 163*B)*a^2*cos(d*x + c) + (200*A + 163*B)*a^2)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (48*B*a^2*cos(d*x + c)^4 + 8*(8*A + 23*B)*a^2*cos(d*x + c)^3 + 2*(136*A + 163*B)*a^2*cos(d*x + c)^2 + 3*(200*A + 163*B)*a^2*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c) + d)$

3.518.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output Timed out

3.518. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

3.518.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9390 vs. $2(211) = 422$.

Time = 0.96 (sec) , antiderivative size = 9390, normalized size of antiderivative = 38.02

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algo
rithm="maxima")
```

```
output 1/768*(8*(4*(a^2*cos(3/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d
*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1))*si
n(3*d*x + 3*c) - (a^2*cos(3*d*x + 3*c) - a^2)*sin(3/2*arctan2(sin(2/3*arct
an2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c)
, cos(3*d*x + 3*c))) + 1)))*(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x +
3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos
(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)^(3/4)*sqrt(a) + 30*
(cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(
sin(3*d*x + 3*c), cos(3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c)
, cos(3*d*x + 3*c))) + 1)^(1/4))*((a^2*sin(2/3*arctan2(sin(3*d*x + 3*c), c
os(3*d*x + 3*c))) + 5*a^2*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*
c))))*cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))
, cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 1)) - (a^2*cos(2/
3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))) + 3*a^2*cos(1/3*arctan2(sin
(3*d*x + 3*c), cos(3*d*x + 3*c))) - 4*a^2*sin(1/2*arctan2(sin(2/3*arctan2
(sin(3*d*x + 3*c), cos(3*d*x + 3*c))), cos(2/3*arctan2(sin(3*d*x + 3*c), c
os(3*d*x + 3*c))) + 1)))*sqrt(a) + 75*(a^2*arctan2(-(cos(2/3*arctan2(sin(3
*d*x + 3*c), cos(3*d*x + 3*c)))^2 + sin(2/3*arctan2(sin(3*d*x + 3*c), cos(
3*d*x + 3*c)))^2 + 2*cos(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c)))
+ 1)^(1/4)*(cos(1/2*arctan2(sin(2/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x...
```

3.518.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^{5/2}}{\sqrt{\sec(dx + c)}} dx$$

```
input integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algo
rithm="giac")
```

3.518. $\int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(5/2)/sqrt(sec(d*x + c)), x)`

3.518.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(1/2), x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(1/2), x)`

$$3.519 \quad \int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.519.1 Optimal result 4715
 3.519.2 Mathematica [A] (verified) 4716
 3.519.3 Rubi [A] (verified) 4716
 3.519.4 Maple [A] (verified) 4720
 3.519.5 Fricas [A] (verification not implemented) 4721
 3.519.6 Sympy [F(-1)] 4722
 3.519.7 Maxima [B] (verification not implemented) 4722
 3.519.8 Giac [F] 4723
 3.519.9 Mupad [F(-1)] 4724

3.519.1 Optimal result

Integrand size = 35, antiderivative size = 294

$$\int \frac{(a+a \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx = \frac{a^{5/2}(326A+283B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)}}{128d}$$

$$+ \frac{a^3(170A+157B) \sin(c+dx)}{240d\sqrt{a+a \cos(c+dx)} \sec^{\frac{5}{2}}(c+dx)} + \frac{a^2(10A+13B)\sqrt{a+a \cos(c+dx)} \sin(c+dx)}{40d \sec^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{aB(a+a \cos(c+dx))^{3/2} \sin(c+dx)}{5d \sec^{\frac{5}{2}}(c+dx)} + \frac{a^3(326A+283B) \sin(c+dx)}{192d\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)}$$

$$+ \frac{a^3(326A+283B) \sin(c+dx)}{128d\sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}}$$

output

```
1/5*a*B*(a+a*cos(d*x+c))^(3/2)*sin(d*x+c)/d/sec(d*x+c)^(5/2)+1/240*a^3*(17
0*A+157*B)*sin(d*x+c)/d/sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2)+1/192*a^3*
(326*A+283*B)*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/40*a^
2*(10*A+13*B)*sin(d*x+c)*(a+a*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(5/2)+1/128*a
^3*(326*A+283*B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+1/12
8*a^(5/2)*(326*A+283*B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*
cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.519.2 Mathematica [A] (verified)

Time = 1.89 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.61

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \frac{a^2 \sqrt{\cos(c + dx)} \sqrt{a(1 + \cos(c + dx))} \sec\left(\frac{1}{2}(c + dx)\right) \sqrt{s}}$$

input `Integrate[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `(a^2*Sqrt[Cos[c + d*x]]*Sqrt[a*(1 + Cos[c + d*x])]*Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(15*Sqrt[2]*(326*A + 283*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*Sqrt[Cos[c + d*x]]*(5810*A + 5521*B + (3620*A + 3874*B)*Cos[c + d*x] + 4*(230*A + 331*B)*Cos[2*(c + d*x)] + 120*A*Cos[3*(c + d*x)] + 348*B*Cos[3*(c + d*x)] + 48*B*Cos[4*(c + d*x)])*Sin[(c + d*x)/2]))/(3840*d)`

3.519.3 Rubi [A] (verified)Time = 1.71 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3440, 3042, 3455, 27, 3042, 3455, 27, 3042, 3460, 3042, 3249, 3042, 3249, 3042, 3253, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a \cos(c + dx) + a)^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \cos^{3/2}(c + dx) (\cos(c + dx)a + a)^{5/2} (A + B \cos(c + dx)) dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.519. $\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{5/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{5} \int \frac{1}{2} \cos^{3/2}(c+dx)(\cos(c+dx)a+a)^{3/2}(5a(2A+B)+a(10A+13B)\cos(c+dx)) dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{10} \int \cos^{3/2}(c+dx)(\cos(c+dx)a+a)^{3/2}(5a(2A+B)+a(10A+13B)\cos(c+dx)) dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{10} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{3/2} (5a(2A+B)+a(10A+13B)\cos(c+dx)) dx\right)$$

↓ 3455

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{10} \left(\frac{1}{4} \int \frac{1}{2} \cos^{3/2}(c+dx)\sqrt{\cos(c+dx)a+a}(5(26A+21B)a^2+(170A+157B)\cos(c+dx)) dx\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{10} \left(\frac{1}{8} \int \cos^{3/2}(c+dx)\sqrt{\cos(c+dx)a+a}(5(26A+21B)a^2+(170A+157B)\cos(c+dx)) dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{10} \left(\frac{1}{8} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}(5(26A+21B)a^2+(170A+157B)\cos(c+dx)) dx\right)\right)$$

↓ 3460

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{10} \left(\frac{1}{8} \left(\frac{5}{6}a^2(326A+283B) \int \cos^{3/2}(c+dx)\sqrt{\cos(c+dx)a+adx} + \frac{a^3(170A+157B)}{3d\sqrt{a}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{1}{8}\left(\frac{5}{6}a^2(326A+283B)\int\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx}+a^2\right)\right)\right)$$

↓ 3249

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{1}{8}\left(\frac{5}{6}a^2(326A+283B)\left(\frac{3}{4}\int\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+adx}+\frac{a\sin(c+dx)}{2d\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{1}{8}\left(\frac{5}{6}a^2(326A+283B)\left(\frac{3}{4}\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+adx}+\right.\right.\right.\right)$$

↓ 3249

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{1}{8}\left(\frac{5}{6}a^2(326A+283B)\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{1}{8}\left(\frac{5}{6}a^2(326A+283B)\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx+\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)}}\right)\right)\right)\right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{1}{8}\left(\frac{5}{6}a^2(326A+283B)\left(\frac{3}{4}\left(\frac{a\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)}+a}-\frac{\int\frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}}}{d}\right)\right)\right)\right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{10}\left(\frac{a^2(10A+13B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}{4d}+\frac{1}{8}\left(\frac{a^3(170A+13B)}{3d}\right)\right)\right)$$

input `Int[((a + a*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*B*Cos[c + d*x]^(5/2)*(a + a*Cos[
c + d*x])^(3/2)*Sin[c + d*x])/(5*d) + ((a^2*(10*A + 13*B)*Cos[c + d*x]^(5/
2)*Sqrt[a + a*Cos[c + d*x]]*Sin[c + d*x])/(4*d) + ((a^3*(170*A + 157*B)*Co
s[c + d*x]^(5/2)*Sin[c + d*x])/(3*d*Sqrt[a + a*Cos[c + d*x]]) + (5*a^2*(32
6*A + 283*B)*((a*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c +
d*x]]) + (3*((Sqrt[a]*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x
]]))/d + (a*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]))
)/4)/6)/8)/10)
```

3.519.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3249 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((c_.) + (d_.)*sin[(e_.) + (
f_.)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])
^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]))], x] + Simp[2*n*((b*c + a*d)/(b*(
2*n + 1))) Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^(n - 1), x],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2,
0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && IntegerQ[2*n]
```

```
rule 3253 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Co
s[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && E
qQ[a^2 - b^2, 0] && EqQ[d, a/b]
```

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^m Int[(a + b*SIN[e + f*x])^m*((c +
d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3455 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-b)*B*COS[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n
+ 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*SIN[e +
f*x])^(m - 1)*(c + d*SIN[e + f*x])^n*Simp[a*A*d*(m + n + 1) + B*(a*c*(m - 1
) + b*d*(n + 1)) + (A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*SIN[e +
f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d,
0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1/2] && !LtQ[n, -1
] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3460 Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (
f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp
[-2*b*B*COS[e + f*x]*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*SIN[e + f*x]]), x] + Simp[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)) Int[Sqrt[a + b*SIN[e + f*x]]*(c + d*SIN[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

3.519.4 Maple [A] (verified)

Time = 20.83 (sec) , antiderivative size = 419, normalized size of antiderivative = 1.43

method	result
default	$\frac{a^2 \sqrt{a(1+\cos(dx+c))} \left(384B(\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 480A(\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 1392B(\cos^2(dx+c)) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{192d(1+\cos(dx+c)) \sec(dx+c)^{\frac{3}{2}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A \sqrt{a(1+\cos(dx+c))} \left(48 \sin(dx+c) (\cos^2(dx+c)) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 184 \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 326 \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{192d(1+\cos(dx+c)) \sec(dx+c)^{\frac{3}{2}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((a+cos(d*x+c)*a)^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

$$3.519. \int \frac{(a+a \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

output `1/1920*a^2/d*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/sec(d*x+c)^(3/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*(384*B*cos(d*x+c)^3*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+480*A*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+1392*B*cos(d*x+c)^2*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+1840*A*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2264*B*cos(d*x+c)*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+3260*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+2830*B*sin(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4890*tan(d*x+c)*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4245*tan(d*x+c)*B*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4890*sec(d*x+c)*A*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+4245*sec(d*x+c)*B*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)))`

3.519.5 Fracas [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.69

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx =$$

$$\frac{15((326 A + 283 B)a^2 \cos(dx + c) + (326 A + 283 B)a^2)\sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - (384 B a^2 \cos(dx+c)^5 + 48(10 A + 29 B)a^2 \cos(dx+c)^4 + 8(230 A + 283 B)a^2 \cos(dx+c)^3 + 10(326 A + 283 B)a^2 \cos(dx+c)^2 + 15(326 A + 283 B)a^2 \cos(dx+c)) \sqrt{a \cos(dx+c) + a} \sin(dx+c)}{d \cos(dx+c) + d}$$

192

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorith="fricas")`

output `-1/1920*(15*((326*A + 283*B)*a^2*cos(d*x + c) + (326*A + 283*B)*a^2)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - (384*B*a^2*cos(d*x + c)^5 + 48*(10*A + 29*B)*a^2*cos(d*x + c)^4 + 8*(230*A + 283*B)*a^2*cos(d*x + c)^3 + 10*(326*A + 283*B)*a^2*cos(d*x + c)^2 + 15*(326*A + 283*B)*a^2*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c) + d)`

3.519.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+a*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.519.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10042 vs. $2(252) = 504$.

Time = 0.95 (sec) , antiderivative size = 10042, normalized size of antiderivative = 34.16

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/7680*((10*(cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + 2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)^(3/4))*((75*a^2*sin(4/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 88*a^2*sin(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 75*a^2*sin(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))))*cos(3/2*arctan2(sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))), cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)) - (75*a^2*cos(4/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 88*a^2*cos(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) - 75*a^2*cos(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) - 88*a^2)*sin(3/2*arctan2(sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))), cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)))*sqrt(a) + 6*(cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2 + 2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)^(1/4))*((8*(a^2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))^2*sin(5*d*x + 5*c) + a^2*sin(5*d*x + 5*c)*sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))))^2 + 2*a^2*cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c)))*sin(5*d*x + 5*c) + a^2*sin(5*d*x + 5*c))*cos(5/2*arctan2(sin(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))), cos(2/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 1)) - 5*(15*a^2*sin(4/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5...`

3.519.8 Giac [F]

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(a \cos(dx + c) + a)^{5/2}}{\sec(dx + c)^{3/2}} dx$$

input `integrate((a+a*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(a*cos(d*x + c) + a)^(5/2)/sec(d*x + c)^(3/
2), x)`

3.519.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + a \cos(c + dx))^{5/2}}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + a*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(3/2), x)`

3.520
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.520.1 Optimal result 4725
 3.520.2 Mathematica [C] (verified) 4726
 3.520.3 Rubi [A] (verified) 4726
 3.520.4 Maple [A] (verified) 4733
 3.520.5 Fricas [A] (verification not implemented) 4733
 3.520.6 Sympy [F(-1)] 4734
 3.520.7 Maxima [F] 4734
 3.520.8 Giac [F] 4735
 3.520.9 Mupad [F(-1)] 4735

3.520.1 Optimal result

Integrand size = 35, antiderivative size = 295

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= -\frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$+ \frac{2(257A - 129B) \sqrt{\sec(c + dx)} \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}}$$

$$- \frac{2(29A - 93B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{315d \sqrt{a + a \cos(c + dx)}} + \frac{2(19A - 3B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{105d \sqrt{a + a \cos(c + dx)}}$$

$$- \frac{2(A - 9B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{63d \sqrt{a + a \cos(c + dx)}} + \frac{2A \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{9d \sqrt{a + a \cos(c + dx)}}$$

output `-2/315*(29*A-93*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/105*(19*A-3*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-2/63*(A-9*B)*sec(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/9*A*sec(d*x+c)^(9/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)+2/315*(257*A-129*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.520.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.520.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.75 (sec) , antiderivative size = 272, normalized size of antiderivative = 0.92

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{2e^{-\frac{1}{2}i(c+dx)} \cos\left(\frac{1}{2}(c+dx)\right) \left(-315i(A-B) \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) - \frac{1}{4}(-1279\right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2))/Sqrt[a + a*Cos[c + d*x]], x]`

output `(2*Cos[(c + d*x)/2]*((-315*I)*(A - B)*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]])] - ((-1279*A + 423*B + (214*A - 918*B)*Cos[c + d*x] - 8*(157*A - 69*B)*Cos[2*(c + d*x)] + 58*A*Cos[3*(c + d*x)] - 186*B*Cos[3*(c + d*x)] - 257*A*Cos[4*(c + d*x)] + 129*B*Cos[4*(c + d*x)])*Sec[c + d*x]^(9/2)*(Cos[(c + d*x)/2] + I*Sin[(c + d*x)/2])*Sin[(c + d*x)/2])/4)/(315*d*E^((I/2)*(c + d*x))*Sqrt[a*(1 + Cos[c + d*x])])]`

3.520.3 Rubi [A] (verified)

Time = 1.97 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.14, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3440, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{11}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a \cos(c + dx) + a}} dx$$

↓ 3042

$$\int \frac{\csc\left(c + dx + \frac{\pi}{2}\right)^{11/2} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx$$

3.520. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow \text{3440} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{11}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{11/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
& \downarrow \text{3463} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{a(A-9B)-8aA\cos(c+dx)}{2\cos^{\frac{9}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{9a} + \frac{2A\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) \\
& \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-9B)-8aA\cos(c+dx)}{\cos^{\frac{9}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{9a} \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-9B)-8aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{9/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{9a} \right) \\
& \downarrow \text{3463} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2 \int -\frac{3(a^2(19A-3B)-2a^2(A-9B)\cos(c+dx))}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{9a} + \frac{2a}{7d\cos^{\frac{7}{2}}(c+dx)} \right) \\
& \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{9d\cos^{\frac{9}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a(A-9B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{3 \int \frac{a^2(19A-3B)-2a^2}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} dx}{7a}}{9a} \right)
\end{aligned}$$

3.520. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a\cos(c+dx)}} dx$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{3 \int \frac{a^2(19A-3B)-2a^2(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{7/2}} dx}{9a} \right) \end{array}$$

$$\begin{array}{c} \downarrow 3463 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{3 \left(2 \int \frac{a^3(29A-93B)}{2 \cos^{\frac{5}{2}}(c+dx)} dx \right)}{9a} \right) \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{3 \left(\frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right)}{9a} \right) \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{3 \left(\frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right)}{9a} \right) \end{array}$$

$$\begin{array}{c} \downarrow 3463 \\ \int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx \end{array}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}}{1} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}}{1} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}}{1} \right)$$

↓ 3463

3.520. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}}{1} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}}{1} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}}}{1} \right)$$

↓ 3261

3.520. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\left(\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - 3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} \right)$$

↓ 218

$$\left(\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \frac{2A \sin(c+dx)}{9d \cos^{\frac{9}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2a(A-9B) \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - 3 \frac{2a^2(19A-3B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sin[c + d*x])/(9*d*Cos[c + d*x]^ (9/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*a*(A - 9*B)*Sin[c + d*x])/(7*d*Cos [c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) - (3*((2*a^2*(19*A - 3*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x])) - ((2*a^3*(29*A - 93*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]])) - ((-315*Sqrt[2]*a^(7/2)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt [Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/d + (2*a^4*(257*A - 129*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(5*a)))/(7*a))/(9*a))`

3.520. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.520.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.520.4 Maple [A] (verified)

Time = 9.42 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.37

method	result
default	$\left(\sec^{\frac{11}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(315A\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^6(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-315B\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)$
parts	$A\left(\sec^{\frac{11}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(315(\cos^6(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+315\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))\right)$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output $\frac{1}{315}d\sec(d*x+c)^{(11/2)}*(a*(1+\cos(d*x+c)))^{(1/2)}/(1+\cos(d*x+c))*(315*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^6*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}-315*B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^6*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+315*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^5*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+257*A*\sin(d*x+c)*\cos(d*x+c)^5*2^{(1/2)}-315*B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^5*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}-129*B*\sin(d*x+c)*\cos(d*x+c)^5*2^{(1/2)}-29*A*\sin(d*x+c)*\cos(d*x+c)^4*2^{(1/2)}+93*B*\sin(d*x+c)*\cos(d*x+c)^4*2^{(1/2)}+57*A*\sin(d*x+c)*\cos(d*x+c)^3*2^{(1/2)}-9*B*\sin(d*x+c)*\cos(d*x+c)^3*2^{(1/2)}-5*A*\sin(d*x+c)*\cos(d*x+c)^2*2^{(1/2)}+45*B*\sin(d*x+c)*\cos(d*x+c)^2*2^{(1/2)}+35*A*\sin(d*x+c)*\cos(d*x+c)*2^{(1/2)})*2^{(1/2)}/a$

3.520.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.67

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{315\sqrt{2}\left((A-B)a\cos(dx+c)^5+(A-B)a\cos(dx+c)^4\right)\arctan\left(\frac{\sqrt{2}\sqrt{a\cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a}\sin(dx+c)}\right)+2\left((257A-129B)\cos(dx+c)^4-(29A-93B)\cos(dx+c)^3\right)\sqrt{a}}{315(ad\cos(dx+c))^5+ad\cos(dx+c)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)/(a+a*cos(d*x+c))^(1/2),x,algorithm="fricas")`

output $1/315*(315*\sqrt{2})*((A - B)*a*\cos(d*x + c)^5 + (A - B)*a*\cos(d*x + c)^4)*a$
 $rctan(\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x$
 $+ c)))/\sqrt{a} + 2*((257*A - 129*B)*\cos(d*x + c)^4 - (29*A - 93*B)*\cos(d*$
 $x + c)^3 + 3*(19*A - 3*B)*\cos(d*x + c)^2 - 5*(A - 9*B)*\cos(d*x + c) + 35*A$
 $)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/\sqrt{\cos(d*x + c)})/(a*d*\cos(d*x +$
 $c)^5 + a*d*\cos(d*x + c)^4)$

3.520.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(11/2)/(a+a*cos(d*x+c))**(1/2),x)`

output Timed out

3.520.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{11}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)/(a+a*cos(d*x+c))^(1/2),x, alg`
`orithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(11/2)/sqrt(a*cos(d*x + c) + a`
`), x)`

3.520.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{11}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(11/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(11/2)/sqrt(a*cos(d*x + c) + a), x)`

3.520.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{11/2}}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2))/(a + a*cos(c + d*x))^(1/2), x)`

3.521
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.521.1 Optimal result 4736
 3.521.2 Mathematica [C] (warning: unable to verify) 4737
 3.521.3 Rubi [A] (verified) 4737
 3.521.4 Maple [A] (verified) 4742
 3.521.5 Fracas [A] (verification not implemented) 4743
 3.521.6 Sympy [F(-1)] 4743
 3.521.7 Maxima [F] 4744
 3.521.8 Giac [F] 4744
 3.521.9 Mupad [F(-1)] 4744

3.521.1 Optimal result

Integrand size = 35, antiderivative size = 250

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$- \frac{2(43A - 91B)\sqrt{\sec(c + dx)} \sin(c + dx)}{105d\sqrt{a + a \cos(c + dx)}} + \frac{2(31A - 7B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{105d\sqrt{a + a \cos(c + dx)}}$$

$$- \frac{2(A - 7B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{35d\sqrt{a + a \cos(c + dx)}} + \frac{2A \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{7d\sqrt{a + a \cos(c + dx)}}$$

output

```
2/105*(31*A-7*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-2/35
*(A-7*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/7*A*sec(d*
x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+(A-B)*arctan(1/2*sin(d*x+c)
*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+
c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)-2/105*(43*A-91*B)*sin(d*x+c)*sec(d*x+c
)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```

3.521.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.521.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 10.90 (sec) , antiderivative size = 2442, normalized size of antiderivative = 9.77

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Result too large to show}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Cos[c/2 + (d*x)/2]*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*((2*B*Sin[c/2 + (d*x)/2])/(7*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(7/2)) + (12*B*Sin[c/2 + (d*x)/2])/(35*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) + (16*B*Sin[c/2 + (d*x)/2])/(35*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (32*B*Sin[c/2 + (d*x)/2])/(35*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + ((A - B)*Csc[c/2 + (d*x)/2]^9*(363825*Sin[c/2 + (d*x)/2]^2 - 4729725*Sin[c/2 + (d*x)/2]^4 + 26785605*Sin[c/2 + (d*x)/2]^6 - 86790165*Sin[c/2 + (d*x)/2]^8 + 177677808*Sin[c/2 + (d*x)/2]^10 - 239283044*Sin[c/2 + (d*x)/2]^12 + 52080*Hypergeometric2F1[2, 11/2, 13/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12 + 560*HypergeometricPFQ[{2, 2, 2, 2, 11/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12 + 213120160*Sin[c/2 + (d*x)/2]^14 - 168280*Hypergeometric2F1[2, 11/2, 13/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14 - 2240*HypergeometricPFQ[{2, 2, 2, 2, 11/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14 - 121497024*Sin[c/2 + (d*x)/2]^16 + 212520*Hypergeometric2F1[2, 11/2, 13/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^16 + 3360*HypergeometricPFQ[{2, 2, 2, 2, 11/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^16 + 40125184*Sin[c/2 + (d*x)/2]^18 - 124320*Hypergeometric2F1[2, 11/2, ...`

3.521.3 Rubi [A] (verified)

Time = 1.56 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.12, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3440, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.521. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
 & \int \frac{\sec^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{9/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{9}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{9/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
 & \quad \downarrow \text{3463} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{a(A-7B)-6aA\cos(c+dx)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{7a} + \frac{2A\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-7B)-6aA\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{7a} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-7B)-6aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{7a} \right) \\
 & \quad \downarrow \text{3463} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2 \int -\frac{a^2(31A-7B)-4a^2(A-7B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2a(A-7B)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.521. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a\cos(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^2(31A-7B)-4a^2(A-7B) \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)}}{\cos^{\frac{5}{2}}(c+dx) \sqrt{\cos(c+dx)}} dx}{5a} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^2(31A-7B)-4a^2(A-7B) \sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{\cos(c+dx)}}{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{\cos(c+dx)}} dx}{5} \right)$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2 \int \frac{a^3(43A-91B)-2a^3(A-7B) \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)}}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)}} dx}{7a} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}}{7a} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(31A-7B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right)$$

3.521. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{array}{c} \downarrow \text{3463} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(31A-7B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \end{array} \right. \end{array}$$

$$\begin{array}{c} \downarrow \text{27} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(31A-7B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \end{array} \right. \end{array}$$

$$\begin{array}{c} \downarrow \text{3042} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(31A-7B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \end{array} \right. \end{array}$$

$$\begin{array}{c} \downarrow \text{3261} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(31A-7B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \end{array} \right. \end{array}$$

$$\downarrow \text{218}$$

3.521. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{7d \cos^{\frac{7}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2a(A-7B) \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2a^2(31A-7B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sin[c + d*x])/(7*d*Cos[c + d*x])^(7/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*a*(A - 7*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*a^2*(31*A - 7*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-105*Sqrt[2]*a^(5/2)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^3*(43*A - 91*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a)/(5*a)/(7*a))`

3.521.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.521. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m*Int[(a + b*Sin[e + f*x])^n*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])
```

```
rule 3463 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.521.4 Maple [A] (verified)

Time = 9.90 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.46

method	result
default	$-\frac{\left(\sec^{\frac{9}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(105A\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-105B\arcsin(\cot(dx+c)-\csc(dx+c))\cos^5(dx+c)\right)}{a^{\frac{1}{2}}}$
parts	$-\frac{A\left(\sec^{\frac{9}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(105\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))+105\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))\cos^5(dx+c)\right)}{a^{\frac{1}{2}}}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/105/d*sec(d*x+c)^(9/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))*(105*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-105*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+43*A*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)+105*A*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-91*B*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)-105*B*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-31*A*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+7*B*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+3*A*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-21*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-15*A*sin(d*x+c)*cos(d*x+c)*2^(1/2))*2^(1/2)/a
```

$$3.521. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.521.5 Fricas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.72

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{105 \sqrt{2} \left((A - B) a \cos(dx + c)^4 + (A - B) a \cos(dx + c)^3 \right) \arctan \left(\frac{\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}}{\sqrt{a} \sin(dx + c)} \right) + 2 \left((43 A - 91 B) \cos(dx + c)^3 - (31 A - 7 B) \cos(dx + c)^2 + 3(A - 7 B) \cos(dx + c) - 15 A \right) \sqrt{a \cos(dx + c) + a} \sin(dx + c)}{105 (ad \cos(dx + c)^4 + ad \cos(dx + c)^3)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `-1/105*(105*sqrt(2)*((A - B)*a*cos(d*x + c)^4 + (A - B)*a*cos(d*x + c)^3)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a) + 2*((43*A - 91*B)*cos(d*x + c)^3 - (31*A - 7*B)*cos(d*x + c)^2 + 3*(A - 7*B)*cos(d*x + c) - 15*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c))/(a*d*cos(d*x + c)^4 + a*d*cos(d*x + c)^3)`

3.521.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(9/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.521.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{9}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(9/2)/sqrt(a*cos(d*x + c) + a), x)`

3.521.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{9}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(9/2)/sqrt(a*cos(d*x + c) + a), x)`

3.521.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{9/2}}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2))/(a + a*cos(c + d*x))^(1/2), x)`

3.521. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.522
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.522.1 Optimal result 4745
 3.522.2 Mathematica [C] (warning: unable to verify) 4746
 3.522.3 Rubi [A] (verified) 4746
 3.522.4 Maple [A] (verified) 4750
 3.522.5 Fricas [A] (verification not implemented) 4751
 3.522.6 Sympy [F(-1)] 4751
 3.522.7 Maxima [F] 4752
 3.522.8 Giac [F] 4752
 3.522.9 Mupad [F(-1)] 4752

3.522.1 Optimal result

Integrand size = 35, antiderivative size = 207

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= -\frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$+ \frac{2(13A - 5B) \sqrt{\sec(c + dx)} \sin(c + dx)}{15d \sqrt{a + a \cos(c + dx)}}$$

$$- \frac{2(A - 5B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{15d \sqrt{a + a \cos(c + dx)}} + \frac{2A \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{5d \sqrt{a + a \cos(c + dx)}}$$

output

```
-2/15*(A-5*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+2/5*A*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)-(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)+2/15*(13*A-5*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)
```


3.522.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.21 (sec) , antiderivative size = 1719, normalized size of antiderivative = 8.30

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Cos[c/2 + (d*x)/2]*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*((2*B*Sin[c/2 + (d*x)/2])/(5*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) + (8*B*Sin[c/2 + (d*x)/2])/(15*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (16*B*Sin[c/2 + (d*x)/2])/(15*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])) - ((A - B)*Csc[c/2 + (d*x)/2]^7*(4725*Sin[c/2 + (d*x)/2]^2 - 48825*Sin[c/2 + (d*x)/2]^4 + 210105*Sin[c/2 + (d*x)/2]^6 - 486630*Sin[c/2 + (d*x)/2]^8 + 655812*Sin[c/2 + (d*x)/2]^10 - 710*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^10 - 40*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 9/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^10 - 518760*Sin[c/2 + (d*x)/2]^12 + 1770*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12 + 226656*Sin[c/2 + (d*x)/2]^14 - 1500*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14 - 42048*Sin[c/2 + (d*x)/2]^16 + 440*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^16 + 4725*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)] - 56700*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Sin[c/2 + (d*x)/2]^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)] + 291060*ArcTanh[Sqrt[Sin[c/2 + (d*...`

3.522.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.09, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.522. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{\sec^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc(c+dx+\frac{\pi}{2})^{7/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
& \quad \downarrow \text{3463} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{a(A-5B)-4aA\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2A\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-5B)-4aA\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a(A-5B)-4aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a} \right) \\
& \quad \downarrow \text{3463} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2 \int -\frac{a^2(13A-5B)-2a^2(A-5B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a(A-5B)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} \right) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.522. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a\cos(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{\int \frac{a^2(13A-5B) - 2a^2(A-5B) \cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx) \sqrt{\cos(c+dx)}} dx}{3a} \right) \frac{1}{5a}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{\int \frac{a^2(13A-5B) - 2a^2(A-5B) \sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{\cos(c+dx)}}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{\cos(c+dx)}} dx}{3} \right) \frac{1}{5a}$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2 \int \frac{15a^3(A-5B) \cos^2(c+dx) \sqrt{\cos(c+dx)}}{2\sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)}} dx}{a} \right) \frac{1}{5a}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a^2(13A-5B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx) + a}} \right) \frac{1}{5a}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx) + a}} - \frac{2a^2(13A-5B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx) + a}} \right) \frac{1}{5a}$$

↓ 3261

3.522. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{30a^3(A-B) \int \frac{\sin(c+dx)}{\cos(c+dx)} dx}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{5d \cos^{\frac{5}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-5B) \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a^2(13A-5B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*a*(A - 5*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-15*Sqrt[2]*a^(3/2)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^2*(13*A - 5*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(5*a)`

3.522.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.522. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])
```

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.522.4 Maple [A] (verified)

Time = 9.84 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.57

method	result
default	$\frac{\left(\sec^{\frac{7}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(15A(\cos^4(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-15B(\cos^4(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{15d(1+\cos(dx+c))}$
parts	$\frac{A\left(\sec^{\frac{7}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(15\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^4(dx+c))+15(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{15d(1+\cos(dx+c))}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)
```

$$3.522. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

output $1/15/d*\sec(d*x+c)^{(7/2)}*(a*(1+\cos(d*x+c)))^{(1/2)}/(1+\cos(d*x+c))*(15*A*\cos(d*x+c)^4*\arcsin(\cot(d*x+c)-\csc(d*x+c))*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}-15*B*\cos(d*x+c)^4*\arcsin(\cot(d*x+c)-\csc(d*x+c))*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+13*A*\sin(d*x+c)*\cos(d*x+c)^3*2^{(1/2)}+15*A*\cos(d*x+c)^3*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-5*B*\sin(d*x+c)*\cos(d*x+c)^3*2^{(1/2)}-15*B*\cos(d*x+c)^3*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\arcsin(\cot(d*x+c)-\csc(d*x+c))-A*\sin(d*x+c)*\cos(d*x+c)^2*2^{(1/2)}+5*B*\sin(d*x+c)*\cos(d*x+c)^2*2^{(1/2)}+3*A*\sin(d*x+c)*\cos(d*x+c)*2^{(1/2)})*2^{(1/2)}/a$

3.522.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.79

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{15 \sqrt{2} \left((A - B) a \cos(dx + c)^3 + (A - B) a \cos(dx + c)^2 \right) \arctan \left(\frac{\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}}{\sqrt{a} \sin(dx + c)} \right) + \frac{2 \left((13 A - 5 B) \cos(dx + c)^2 - (A - 5 B) \cos(dx + c) \right)}{\sqrt{\cos(dx + c)}}}{15 \left(ad \cos(dx + c)^3 + ad \cos(dx + c)^2 \right)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2),x, algo rithm="fricas")`

output $1/15*(15*\sqrt{2})*((A - B)*a*\cos(d*x + c)^3 + (A - B)*a*\cos(d*x + c)^2)*\arctan(\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c)))/\sqrt{a} + 2*((13*A - 5*B)*\cos(d*x + c)^2 - (A - 5*B)*\cos(d*x + c) + 3*A)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/\sqrt{\cos(d*x + c)})/(a*d*\cos(d*x + c)^3 + a*d*\cos(d*x + c)^2)$

3.522.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(7/2)/(a+a*cos(d*x+c))**(1/2),x)`

output Timed out

3.522. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.522.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/sqrt(a*cos(d*x + c) + a), x)`

3.522.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/sqrt(a*cos(d*x + c) + a), x)`

3.522.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{7/2}}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + a*cos(c + d*x))^(1/2), x)`

3.522. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.523
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.523.1 Optimal result 4753
 3.523.2 Mathematica [C] (warning: unable to verify) 4753
 3.523.3 Rubi [A] (verified) 4754
 3.523.4 Maple [B] (verified) 4757
 3.523.5 Fricas [A] (verification not implemented) 4758
 3.523.6 Sympy [F(-1)] 4758
 3.523.7 Maxima [F] 4759
 3.523.8 Giac [F] 4759
 3.523.9 Mupad [F(-1)] 4759

3.523.1 Optimal result

Integrand size = 35, antiderivative size = 162

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$- \frac{2(A - 3B)\sqrt{\sec(c + dx)} \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}} + \frac{2A \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3d\sqrt{a + a \cos(c + dx)}}$$

output `2/3*A*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)+(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)-2/3*(A-3*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.523.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

3.523.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

Time = 6.59 (sec) , antiderivative size = 617, normalized size of antiderivative = 3.81

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)} \left(\frac{2B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{3 \left(1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)\right)^{3/2}} + \frac{4B \sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{3 \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} + \frac{(A - B) \csc^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{3 \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} \right)}{3 \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} + \frac{(A - B) \csc^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{3 \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Cos[c/2 + (d*x)/2]*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*((2*B*Sin[c/2 + (d*x)/2])/(3*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (4*B*Sin[c/2 + (d*x)/2])/(3*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + ((A - B)*Csc[c/2 + (d*x)/2]^5*(-12*Cos[(c + d*x)/2]^4*HypergeometricPFQ[{2, 2, 7/2}, {1, 9/2}, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*Sin[c/2 + (d*x)/2]^8 - 12*Hypergeometric2F1[2, 7/2, 9/2, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*Sin[c/2 + (d*x)/2]^8*(4 - 7*Sin[c/2 + (d*x)/2]^2 + 3*Sin[c/2 + (d*x)/2]^4) + 7*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*(15 - 20*Sin[c/2 + (d*x)/2]^2 + 8*Sin[c/2 + (d*x)/2]^4)*((3 - 7*Sin[c/2 + (d*x)/2]^2)*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]) - 3*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]])*(1 - 2*Sin[c/2 + (d*x)/2]^2)))/(63*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(7/2)))/(d*Sqrt[a*(1 + Cos[c + d*x]))]`

3.523.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3440, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a \cos(c + dx) + a}} dx$$

↓ 3042

3.523. $\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$

$$\begin{aligned}
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{\sqrt{a \sin(c+dx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{A+B \cos(c+dx)}{\cos^{5/2}(c+dx) \sqrt{\cos(c+dx)a+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{A+B \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
 & \quad \downarrow \text{3463} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{a(A-3B)-2aA \cos(c+dx)}{2 \cos^{3/2}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2A \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx)+a}} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a(A-3B)-2aA \cos(c+dx)}{\cos^{3/2}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{3a} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a(A-3B)-2aA \sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a} \right) \\
 & \quad \downarrow \text{3463} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2 \int -\frac{3a^2(A-B)}{2 \sqrt{\cos(c+dx)} \sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a(A-3B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{3/2}(c+dx) \sqrt{a \cos(c+dx)+a}} - \frac{2a(A-3B) \sin(c+dx)}{d \sqrt{\cos(c+dx)} \sqrt{a \cos(c+dx)+a}} - \frac{3a(A-B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3a} \right)
 \end{aligned}$$

3.523. $\int \frac{(A+B \cos(c+dx)) \sec^{5/2}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\frac{2a(A-3B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - 3a(A-B) \int \frac{1}{\sqrt{\sin}}}{3a} \right) \\
 & \downarrow 3261 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\frac{6a^2(A-B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} \right)}{d}}{3a} \right) \\
 & \downarrow 218 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\frac{2a(A-3B) \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - \frac{3\sqrt{2}\sqrt{a}(A-B) \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(c+dx)+a}}{\sqrt{\cos(c+dx)}}\right)}{3a}}{3a} \right)
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*Sqrt[a]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a*(A - 3*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a)`

3.523.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

$$3.523. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.523.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. $2(135) = 270$.

Time = 9.94 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.75

method	result
default	$-\frac{\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(3A(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))-3B(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{3d(1+\cos(dx+c))a}$
parts	$-\frac{A\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(3(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))+\sqrt{2}(\cos^2(dx+c))\sin(dx+c)\right)}{3d(1+\cos(dx+c))a}$

3.523.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/3/d*sec(d*x+c)^(5/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))*(3*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-3*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+A*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)+3*A*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-3*B*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-A*sin(d*x+c)*cos(d*x+c)*2^(1/2))*2^(1/2)/a`

3.523.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.88

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{3\sqrt{2}\left((A-B)a \cos(dx+c)^2 + (A-B)a \cos(dx+c)\right) \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) + \frac{2\left((A-3B) \cos(dx+c) - A\right)\sqrt{a \cos(dx+c)+a} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{3\left(ad \cos(dx+c)^2 + ad \cos(dx+c)\right)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2),x, algo="fricas")`

output `-1/3*(3*sqrt(2)*((A - B)*a*cos(d*x + c)^2 + (A - B)*a*cos(d*x + c))*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a) + 2*((A - 3*B)*cos(d*x + c) - A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a*d*cos(d*x + c)^2 + a*d*cos(d*x + c))`

3.523.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(1/2),x)`

3.523. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

output Timed out

3.523.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/sqrt(a*cos(d*x + c) + a), x)`

3.523.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/sqrt(a*cos(d*x + c) + a), x)`

3.523.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{5/2}}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(1/2), x)`

3.523. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.524
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

3.524.1 Optimal result 4761
 3.524.2 Mathematica [C] (warning: unable to verify) 4761
 3.524.3 Rubi [A] (verified) 4762
 3.524.4 Maple [B] (verified) 4764
 3.524.5 Fricas [A] (verification not implemented) 4765
 3.524.6 Sympy [F(-1)] 4765
 3.524.7 Maxima [F] 4766
 3.524.8 Giac [F] 4766
 3.524.9 Mupad [F(-1)] 4766

3.524.1 Optimal result

Integrand size = 35, antiderivative size = 119

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= -\frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$+ \frac{2A\sqrt{\sec(c + dx)} \sin(c + dx)}{d\sqrt{a + a \cos(c + dx)}}$$

output `-(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)+2*A*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(1/2)`

3.524.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.06 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.71

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{2 \cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \sin\left(\frac{1}{2}(c + dx)\right) \left(10B - (A - B) \sec(c + dx)\right) \left(-\frac{5}{4}(1 + 4 \cos(c + dx)) + \cos\right)}{\dots}$$

3.524.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `(2*Cos[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*Sin[(c + d*x)/2]*(10*B - (A - B)*Sec[c + d*x]*((-5*(1 + 4*Cos[c + d*x] + Cos[2*(c + d*x)])*Csc[(c + d*x)/2]^4*(1 - Cos[c + d*x] + ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)])*Cos[c + d*x]*Sqrt[2 - 2*Sec[c + d*x]]))/4 + (Hypergeometric2F1[2, 5/2, 7/2, -(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]*Sin[c + d*x]*Tan[c + d*x])/2))/5*d*Sqrt[a*(1 + Cos[c + d*x]))]`

3.524.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3440, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx$$

↓ 3042

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx$$

↓ 3440

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{a(A-B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2A\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right)$$

3.524. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a\cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2A\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - (A-B)\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}d\right) \\
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2A\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - (A-B)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}}d\right) \\
& \downarrow 3261 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2a(A-B)\int\frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}}d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2A\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right) \\
& \downarrow 218 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2A\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{\sqrt{2}(A-B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{\sqrt{ad}}\right)
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/Sqrt[a + a*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-((Sqrt[2]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/(Sqrt[a]*d)) + (2*A*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))`

3.524.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.524. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a\cos(c+dx)}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.524.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(100) = 200.

Time = 10.18 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.77

method	result
parts	$\frac{A \left(\sec^{\frac{3}{2}}(dx+c)\right) \left(\cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c)) + \sqrt{2} \sin(dx+c) + \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))\right)}{d(1+\cos(dx+c))a}$
default	$\left(\sec^{\frac{3}{2}}(dx+c)\right) \left(A \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c)) - B \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))\right)$

3.524.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNERVERBOSE)`

output `A/d*sec(d*x+c)^(3/2)*(cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+2^(1/2)*sin(d*x+c)+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c)))*cos(d*x+c)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))*2^(1/2)/a-B/d*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)*sec(d*x+c)^(3/2)*2^(1/2)/a`

3.524.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.92

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\sqrt{2}((A-B)a \cos(dx+c) + (A-B)a) \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(dx+c)} + a\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) + \frac{2\sqrt{a \cos(dx+c)} + aA \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{ad \cos(dx+c) + ad}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `(sqrt(2)*((A - B)*a*cos(d*x + c) + (A - B)*a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a) + 2*sqrt(a*cos(d*x + c) + a)*A*sin(d*x + c)/sqrt(cos(d*x + c))/(a*d*cos(d*x + c) + a*d)`

3.524.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.524. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

3.524.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/sqrt(a*cos(d*x + c) + a), x)`

3.524.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/sqrt(a*cos(d*x + c) + a), x)`

3.524.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{\frac{3}{2}}}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(1/2), x)`

3.524. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+a \cos(c+dx)}} dx$

$$3.525 \quad \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+a \cos(c+dx)}} dx$$

3.525.1 Optimal result	4767
3.525.2 Mathematica [A] (verified)	4767
3.525.3 Rubi [A] (verified)	4768
3.525.4 Maple [A] (verified)	4771
3.525.5 Fricas [A] (verification not implemented)	4771
3.525.6 Sympy [F]	4772
3.525.7 Maxima [C] (verification not implemented)	4772
3.525.8 Giac [F(-1)]	4773
3.525.9 Mupad [F(-1)]	4774

3.525.1 Optimal result

Integrand size = 35, antiderivative size = 140

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx \\ &= \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{\sqrt{ad}} \\ & \quad + \frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{\sqrt{ad}} \end{aligned}$$

output

```
2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec
(d*x+c)^(1/2)/d/a^(1/2)+(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*
x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/
2)/d/a^(1/2)
```

3.525.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.73

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx \\ &= \frac{2\left(\sqrt{2}B \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + (A - B) \arctan\left(\frac{\sin\left(\frac{1}{2}(c + dx)\right)}{\sqrt{\cos(c + dx)}}\right)\right) \cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d\sqrt{a(1 + \cos(c + dx))}} \end{aligned}$$

3.525. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+a \cos(c+dx)}} dx$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/Sqrt[a + a*Cos[c + d*x]]],x]`

output `(2*(Sqrt[2]*B*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + (A - B)*ArcTan[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]])*Cos[(c + d*x)/2]*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]/(d*Sqrt[a*(1 + Cos[c + d*x])])]`

3.525.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.86, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3440, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{\sqrt{a\cos(c+dx)+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a\sin(c+dx+\frac{\pi}{2})+a}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx \\
 & \quad \downarrow \text{3461} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left((A-B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{a} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left((A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a} \right) \\
& \quad \downarrow \text{3253} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left((A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2B \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} dx}{ad} \right) \\
& \quad \downarrow \text{223} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left((A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} \right) \\
& \quad \downarrow \text{3261} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} - \frac{2a(A-B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} dx}{d} \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)+a}} \right) \right) \\
& \quad \downarrow \text{218} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\sqrt{2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} + \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{\sqrt{ad}} \right)
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/Sqrt[a + a*Cos[c + d*x]],x]`

output `((2*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/(Sqrt[a]*d) + (Sqrt[2]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]])*Sqrt[a + a*Cos[c + d*x]])]/(Sqrt[a]*d))*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]`

3.525.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3461 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.525.4 Maple [A] (verified)

Time = 10.31 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.99

method	result
default	$-\frac{(\sqrt{\sec(dx+c)})(-B\sqrt{2} \arctan(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}) + A \arcsin(\cot(dx+c) - \csc(dx+c)) - B \arcsin(\cot(dx+c) - \csc(dx+c)))}{d(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a}$
parts	$-\frac{A(\sqrt{\sec(dx+c)} \arcsin(\cot(dx+c) - \csc(dx+c))\sqrt{a(1+\cos(dx+c))} \cos(dx+c)\sqrt{2}}{d(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a} + \frac{B(\sqrt{\sec(dx+c)})(\sqrt{2} \arctan(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}))}{d(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/d*sec(d*x+c)^(1/2)*(-B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))+A*arcsin(cot(d*x+c)-csc(d*x+c))-B*arcsin(cot(d*x+c)-csc(d*x+c)))*cos(d*x+c)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)/a`

3.525.5 Fracas [A] (verification not implemented)

Time = 1.05 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.69

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx =$$

$$\frac{\sqrt{2}(A - B)\sqrt{a} \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) + 2 B \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right)}{ad}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `-(sqrt(2)*(A - B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*B*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))))/(a*d)`

3.525.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a (\cos(c + dx) + 1)}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/sqrt(a*(cos(c + d*x) + 1)), x)`

3.525.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.08 (sec) , antiderivative size = 1221, normalized size of antiderivative = 8.72

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output $(\sqrt{2})A \arctan 2\left(\frac{(\operatorname{abs}(e^{I dx + I c}) + 1)^4 + \cos(dx + c)^4 + \sin(dx + c)^4 + 2(\cos(dx + c)^2 - \sin(dx + c)^2 - 2\cos(dx + c) + 1)\operatorname{abs}(e^{I dx + I c}) + 1)^2 - 4\cos(dx + c)^3 + 2(\cos(dx + c)^2 - 2\cos(dx + c) + 1)\sin(dx + c)^2 + 6\cos(dx + c)^2 - 4\cos(dx + c) + 1)^{1/4} \sin(1/2 \arctan 2(2(\cos(dx + c) - 1)\sin(dx + c)/\operatorname{abs}(e^{I dx + I c}) + 1)^2, (\operatorname{abs}(e^{I dx + I c}) + 1)^2 + \cos(dx + c)^2 - \sin(dx + c)^2 - 2\cos(dx + c) + 1)/\operatorname{abs}(e^{I dx + I c}) + 1)^2\right) + \sin(dx + c)/\operatorname{abs}(e^{I dx + I c}) + 1, \left(\frac{(\operatorname{abs}(e^{I dx + I c}) + 1)^4 + \cos(dx + c)^4 + \sin(dx + c)^4 + 2(\cos(dx + c)^2 - \sin(dx + c)^2 - 2\cos(dx + c) + 1)\operatorname{abs}(e^{I dx + I c}) + 1)^2 - 4\cos(dx + c)^3 + 2(\cos(dx + c)^2 - 2\cos(dx + c) + 1)\sin(dx + c)^2 + 6\cos(dx + c)^2 - 4\cos(dx + c) + 1)^{1/4} \sqrt{a} \cos(1/2 \arctan 2(2(\cos(dx + c) - 1)\sin(dx + c)/\operatorname{abs}(e^{I dx + I c}) + 1)^2, (\operatorname{abs}(e^{I dx + I c}) + 1)^2 + \cos(dx + c)^2 - \sin(dx + c)^2 - 2\cos(dx + c) + 1)/\operatorname{abs}(e^{I dx + I c}) + 1)^2\right) + \sqrt{a} \cos(dx + c) - \sqrt{a})/(\sqrt{a}) \operatorname{abs}(e^{I dx + I c}) + 1)/\sqrt{a} - (\sqrt{2})\sqrt{a} \arctan 2\left(\frac{(\operatorname{abs}(2e^{I dx + I c}) + 2)^4 + 16\cos(dx + c)^4 + 16\sin(dx + c)^4 + 8(\cos(dx + c)^2 - \sin(dx + c)^2 - 2\cos(dx + c) + 1)\operatorname{abs}(2e^{I dx + I c}) + 2)^2 - 64\cos(dx + c)^3 + 32(\cos(dx + c)^2 - 2\cos(dx + c) + 1)\sin(dx + c)^2 + 96\cos(dx + c)^2 - 64\cos(dx + c) + 16)^{1/4} \sin(1/2 \arctan 2(8(\cos(dx + c) - 1)\sin(dx + c)/\operatorname{abs}(2e^{I dx + I c}) + 2)^2, (\operatorname{abs}(2e^{I dx + I c}) + 2)^2 + \cos(dx + c)^2 - \sin(dx + c)^2 - 2\cos(dx + c) + 1)/\operatorname{abs}(2e^{I dx + I c}) + 2)^2\right)$

3.525.8 Giac [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="giac")`

output Timed out

3.525.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{\sqrt{a + a \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(1/2), x)`

3.526 $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)}\sqrt{\sec(c+dx)}} dx$

3.526.1 Optimal result 4775
 3.526.2 Mathematica [C] (verified) 4776
 3.526.3 Rubi [A] (verified) 4776
 3.526.4 Maple [A] (verified) 4780
 3.526.5 Fricas [A] (verification not implemented) 4781
 3.526.6 Sympy [F] 4781
 3.526.7 Maxima [F(-2)] 4782
 3.526.8 Giac [F(-1)] 4782
 3.526.9 Mupad [F(-1)] 4782

3.526.1 Optimal result

Integrand size = 35, antiderivative size = 181

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)}} dx$$

$$= \frac{(2A - B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{\sqrt{ad}} - \frac{\sqrt{2}(A - B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{\sqrt{ad}} + \frac{B \sin(c + dx)}{d\sqrt{a + a \cos(c + dx)}\sqrt{\sec(c + dx)}}$$

```
output B*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+(2*A-B)*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)-(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)
```

3.526.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.56 (sec) , antiderivative size = 467, normalized size of antiderivative = 2.58

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$$

$$= \frac{ie^{-2i(c+dx)}(1 + e^{i(c+dx)}) \left(B - Be^{i(c+dx)} + Be^{2i(c+dx)} - Be^{3i(c+dx)} - (2A - B)e^{i(c+dx)} \sqrt{1 + e^{2i(c+dx)}} \operatorname{arcsinh} \right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[a + a*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]),x]`

output `((I/4)*(1 + E^(I*(c + d*x)))*(B - B*E^(I*(c + d*x)) + B*E^((2*I)*(c + d*x)) - B*E^((3*I)*(c + d*x)) - (2*A - B)*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*ArcSinh[E^(I*(c + d*x))] + Sqrt[2]*B*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))])] + 2*Sqrt[2]*A*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*ArcTanh[(-1 + E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))])] - Sqrt[2]*B*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*ArcTanh[(-1 + E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))])] + 2*A*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]] - B*E^(I*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))])*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]])*Sqrt[Sec[c + d*x]]/(d*E^((2*I)*(c + d*x))*Sqrt[a*(1 + Cos[c + d*x])])]`

3.526.3 Rubi [A] (verified)

Time = 1.00 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.94, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3462, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)} \sqrt{a \cos(c + dx) + a}} dx$$

↓ 3042

3.526. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)a + a}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}} dx \\
& \quad \downarrow \text{3462} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{aB + a(2A - B) \cos(c + dx)}{2\sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + a}} dx}{a} + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{aB + a(2A - B) \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + a}} dx}{2a} + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{aB + a(2A - B) \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}} dx}{2a} + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{3461} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{(2A - B) \int \frac{\sqrt{\cos(c + dx)a + a}}{\sqrt{\cos(c + dx)}} dx - 2a(A - B) \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)a + a}} dx}{2a} + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{(2A - B) \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx - 2a(A - B) \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}} dx}{2a} + \frac{B \sin(c + dx) \sqrt{\cos(c + dx)}}{d \sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{3253}
\end{aligned}$$

3.526. $\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{-2a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2(2A-B) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}}{2a} \right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\sqrt{a}(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right) - 2a(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{2a} + \frac{B \sin(c+dx)}{d} \right)$$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{4a^2(A-B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3 + 2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2a} + \frac{2\sqrt{a}(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d} + \frac{B \sin(c+dx)}{d} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\sqrt{a}(2A-B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right) - \frac{2\sqrt{2}\sqrt{a}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a} + \frac{B \sin(c+dx)}{d} \right)$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[a + a*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*Sqrt[a]*(2*A - B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d - (2*Sqrt[2]*Sqrt[a]*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d)/(2*a) + (B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))`

3.526.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3461 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3462 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m + n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.526.4 Maple [A] (verified)

Time = 20.34 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.08

method	result
default	$\frac{(B\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sin(dx+c)+2A\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)-B\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)+2A\arcsin(\cot(dx+c)-\csc(dx+c))}{2d(1+\cos(dx+c))\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))}\left(\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)+\arcsin(\cot(dx+c)-\csc(dx+c))\right)\sqrt{2}}{d(1+\cos(dx+c))\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a} - \frac{B\left(-\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{a}$

```
input int((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/2/d*(B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+2*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))-B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+2*A*arcsin(cot(d*x+c)-csc(d*x+c))-2*B*arcsin(cot(d*x+c)-csc(d*x+c)))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/sec(d*x+c)^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)/a
```

3.526. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)}\sqrt{\sec(c+dx)}} dx$

3.526.5 Fricas [A] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.93

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$$

$$= \frac{\sqrt{a \cos(dx + c) + a} B \sqrt{\cos(dx + c)} \sin(dx + c) - ((2A - B) \cos(dx + c) + 2A - B) \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx + c) + a}}{\sin(dx + c)}\right)}{ad \cos(dx + c) + ad}$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `(sqrt(a*cos(d*x + c) + a)*B*sqrt(cos(d*x + c))*sin(d*x + c) - ((2*A - B)*c
os(d*x + c) + 2*A - B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*
x + c))/(sqrt(a)*sin(d*x + c))) + sqrt(2)*((A - B)*a*cos(d*x + c) + (A - B
)a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*s
in(d*x + c)))/sqrt(a))/(a*d*cos(d*x + c) + a*d)`

3.526.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a (\cos(c + dx) + 1)} \sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a*(cos(c + d*x) + 1))*sqrt(sec(c + d*x
))), x)`

3.526.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: sign: argument cannot be imagi
nary; found %i
```

3.526.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algo
rithm="giac")
```

```
output Timed out
```

3.526.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} \sqrt{a + a \cos(c + dx)}} dx$$

```
input int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(1/2
)),x)
```

```
output int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(1/2
)), x)
```

3.526. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}} dx$

$$3.527 \quad \int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.527.1 Optimal result	4783
3.527.2 Mathematica [C] (verified)	4784
3.527.3 Rubi [A] (verified)	4784
3.527.4 Maple [A] (verified)	4789
3.527.5 Fricas [A] (verification not implemented)	4789
3.527.6 Sympy [F]	4790
3.527.7 Maxima [F]	4790
3.527.8 Giac [F(-1)]	4791
3.527.9 Mupad [F(-1)]	4791

3.527.1 Optimal result

Integrand size = 35, antiderivative size = 230

$$\begin{aligned} & \int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx \\ &= -\frac{(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{4\sqrt{ad}} \\ & \quad + \frac{\sqrt{2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{\sqrt{ad}} \\ & \quad + \frac{B \sin(c+dx)}{2d\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} + \frac{(4A-B) \sin(c+dx)}{4d\sqrt{a+a \cos(c+dx)} \sqrt{\sec(c+dx)}} \end{aligned}$$

```
output 1/2*B*sin(d*x+c)/d/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2)+1/4*(4*A-B)*sin
(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)-1/4*(4*A-7*B)*arcsin(sin
(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/
d/a^(1/2)+(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+
a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d/a^(1/2)
```

3.527.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.65 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.79

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{ie^{-3i(c+dx)}(1 + e^{i(c+dx)}) \left(-B - 4Ae^{i(c+dx)} + 2Be^{i(c+dx)} + 4Ae^{2i(c+dx)} - 3Be^{2i(c+dx)} - 4Ae^{3i(c+dx)} + 3B \right)}{-}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[a + a*Cos[c + d*x]]*Sec[c + d*x]^(3/2)),x]`

output `((-1/16*I)*(1 + E^(I*(c + d*x)))*(-B - 4*A*E^(I*(c + d*x)) + 2*B*E^(I*(c + d*x)) + 4*A*E^((2*I)*(c + d*x)) - 3*B*E^((2*I)*(c + d*x)) - 4*A*E^((3*I)*(c + d*x)) + 3*B*E^((3*I)*(c + d*x)) + 4*A*E^((4*I)*(c + d*x)) - 2*B*E^((4*I)*(c + d*x)) + B*E^((5*I)*(c + d*x)) - (4*A - 7*B)*E^((2*I)*(c + d*x)))*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcSinh[E^(I*(c + d*x))] - 8*Sqrt[2]*(A - B)*E^((2*I)*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]]) + 4*A*E^((2*I)*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]] - 7*B*E^((2*I)*(c + d*x))*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]])*Sqrt[Sec[c + d*x]]/(d*E^((3*I)*(c + d*x))*Sqrt[a*(1 + Cos[c + d*x])])`

3.527.3 Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.97, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3462, 27, 3042, 3462, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx) \sqrt{a \cos(c + dx) + a}} dx$$

↓ 3042

3.527. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \sqrt{a \sin\left(c + dx + \frac{\pi}{2}\right) + a}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{3/2}(c + dx)(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)a + a}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sin\left(c + dx + \frac{\pi}{2}\right)^{3/2} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)a + a}} dx \\
& \quad \downarrow \text{3462} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3aB+a(4A-B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a} + \frac{B \sin(c + dx) \cos^{3/2}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3aB+a(4A-B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a} + \frac{B \sin(c + dx) \cos^{3/2}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3aB+a(4A-B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a} + \frac{B \sin(c + dx) \cos^{3/2}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{3462} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a^2(4A-B) - a^2(4A-7B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{B \sin(c + dx) \cos^{3/2}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a^2(4A-B) - a^2(4A-7B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx)+a}} + \frac{B \sin(c + dx) \cos^{3/2}(c + dx)}{2d\sqrt{a \cos(c + dx) + a}} \right)
\end{aligned}$$

3.527. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{3/2}(c+dx)} dx$

$$\begin{aligned} & \downarrow \text{3042} \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{\int \frac{a^2(4A-B)-a^2(4A-7B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx}{2a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{B\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{a\cos(c+dx)+a}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3461} \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{8a^2(A-B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx - a(4A-7B)\int \frac{\sqrt{\cos(c+dx)}a+a}{\sqrt{\cos(c+dx)}} dx}{2a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{8a^2(A-B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx - a(4A-7B)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3253} \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{8a^2(A-B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx + \frac{2a(4A-7B)\int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)}a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}a+a}\right)}{d}}{2a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow \text{223} \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{8a^2(A-B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx - \frac{2a^{3/2}(4A-7B)\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{2a} + \frac{a(4A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) \end{aligned}$$

3.527. $\int \frac{A+B\cos(c+dx)}{\sqrt{a+a\cos(c+dx)}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{16a^3(A-B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right) - \frac{2a^{3/2}(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a} - \frac{2a^{3/2}(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{4a} \right)$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{8\sqrt{2}a^{3/2}(A-B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right) - \frac{2a^{3/2}(4A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{2a} + \frac{a(4A-B)}{d\sqrt{a \cos(c+dx)+a}} \right)$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[a + a*Cos[c + d*x]]*Sec[c + d*x]^(3/2)),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*Sqrt[a + a*Cos[c + d*x]]) + (((-2*a^(3/2)*(4*A - 7*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (8*Sqrt[2]*a^(3/2)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d)/(2*a) + (a*(4*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a))`

3.527.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

$$3.527. \int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3462 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m + n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.527.
$$\int \frac{A+B \cos(c+dx)}{\sqrt{a+a \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.527.4 Maple [A] (verified)

Time = 19.93 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.15

method	result
default	$-\frac{(-2B \cos(dx+c) \sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 4A \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + B \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 4A \sqrt{2} \arctan(\frac{\sin(dx+c)}{\sqrt{1+\cos(dx+c)}}))}{8d \sqrt{a}}$
parts	$-\frac{A(-\sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + \sqrt{2} \arctan(\tan(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}) + 2 \arcsin(\cot(dx+c) - \csc(dx+c))) \sqrt{a(1+\cos(dx+c))}}{2d(1+\cos(dx+c)) \sqrt{\sec(dx+c)} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} a}$

```
input int((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(1/2), x, method=_RETURNVERBOSE)
```

```
output -1/8/d/sec(d*x+c)^(1/2)*(-2*B*cos(d*x+c)*sin(d*x+c)*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-4*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+4*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))-7*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2))+8*A*arcsin(cot(d*x+c)-csc(d*x+c))-8*B*arcsin(cot(d*x+c)-csc(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)/a
```

3.527.5 Fracas [A] (verification not implemented)

Time = 2.26 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.84

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{((4A - 7B) \cos(dx + c) + 4A - 7B) \sqrt{a} \arctan\left(\frac{\sqrt{a \cos(dx+c)+a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right) - \frac{4\sqrt{2}((A-B)a \cos(dx+c)+(A-B)a)}{\sqrt{a}}}{4(ad \cos(dx + c) + ad)}$$

```
input integrate((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2), x, algo rithm="fracas")
```

output `1/4*((4*A - 7*B)*cos(d*x + c) + 4*A - 7*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - 4*sqrt(2)*((A - B)*a*cos(d*x + c) + (A - B)*a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a) + (2*B*cos(d*x + c)^2 + (4*A - B)*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a*d*cos(d*x + c) + a*d)`

3.527.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a (\cos(c + dx) + 1)} \sec^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a*(cos(c + d*x) + 1))*sec(c + d*x)**(3/2)), x)`

3.527.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{a \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(a*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.527.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(1/2),x, algorith="giac")`

output `Timed out`

3.527.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + a \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} \sqrt{a + a \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(1/2)), x)`

3.528
$$\int \frac{(aA+(Ab+aB)\cos(c+dx)+bB\cos^2(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+a\cos(c+dx)}} dx$$

3.528.1 Optimal result	4792
3.528.2 Mathematica [A] (verified)	4793
3.528.3 Rubi [A] (verified)	4793
3.528.4 Maple [A] (verified)	4797
3.528.5 Fricas [A] (verification not implemented)	4797
3.528.6 Sympy [F]	4798
3.528.7 Maxima [F]	4798
3.528.8 Giac [F(-1)]	4799
3.528.9 Mupad [F(-1)]	4799

3.528.1 Optimal result

Integrand size = 54, antiderivative size = 192

$$\int \frac{(aA + (Ab + aB)\cos(c + dx) + bB\cos^2(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + a\cos(c + dx)}} dx$$

$$= \frac{(2Ab + 2aB - bB)\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a+a\cos(c+dx)}}\right)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$+ \frac{\sqrt{2}(a - b)(A - B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a\cos(c+dx)}}\right)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{\sqrt{ad}}$$

$$+ \frac{bB\sin(c + dx)}{d\sqrt{a + a\cos(c + dx)}\sqrt{\sec(c + dx)}}$$

output

```
b*B*sin(d*x+c)/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+(2*A*b+2*B*a-B*b)
*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*
x+c)^(1/2)/d/a^(1/2)+(a-b)*(A-B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos
(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*2^(1/2)*cos(d*x+c)^(1/2)*sec(d*x+c)^(
1/2)/d/a^(1/2)
```

3.528.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.74

$$\int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \frac{\cos\left(\frac{1}{2}(c + dx)\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\sqrt{2}(2Ab + 2aB - bB) \arcsin\left(\sqrt{2} \sin\left(\frac{1}{2}(c + dx)\right)\right) + 2(a - b) \right)}{d \sqrt{a(1 + \cos(c + dx))}}$$

input `Integrate[((a*A + (A*b + a*B)*Cos[c + d*x] + b*B*Cos[c + d*x]^2)*Sqrt[Sec[c + d*x]])/Sqrt[a + a*Cos[c + d*x]],x]`

output `(Cos[(c + d*x)/2]*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(Sqrt[2]*(2*A*b + 2*a*B - b*B)*ArcSin[Sqrt[2]*Sin[(c + d*x)/2]] + 2*(a - b)*(A - B)*ArcTan[Sin[(c + d*x)/2]/Sqrt[Cos[c + d*x]]] + 2*b*B*Sqrt[Cos[c + d*x]]*Sin[(c + d*x)/2]))/(d*Sqrt[a*(1 + Cos[c + d*x])])`

3.528.3 Rubi [A] (verified)Time = 1.19 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.95, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4709, 3042, 3524, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c + dx)}((aB + Ab) \cos(c + dx) + aA + bB \cos^2(c + dx))}{\sqrt{a \cos(c + dx) + a}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\sec(c + dx)}((aB + Ab) \cos(c + dx) + aA + bB \cos(c + dx)^2)}{\sqrt{a \cos(c + dx) + a}} dx$$

$$\downarrow \text{4709}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{bB \cos^2(c + dx) + (Ab + aB) \cos(c + dx) + aA}{\sqrt{\cos(c + dx)} \sqrt{\cos(c + dx)} a + a} dx$$

$$\downarrow \text{3042}$$

3.528. $\int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{bB \sin(c+dx+\frac{\pi}{2})^2 + (Ab+aB) \sin(c+dx+\frac{\pi}{2}) + aA}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx$$

↓ 3524

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(2aA+bB)+a(2Ab-Bb+2aB) \cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx) + a}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(2aA+bB)+a(2Ab-Bb+2aB) \cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a} + \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx) + a}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(2aA+bB)+a(2Ab-Bb+2aB) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{2a} + \frac{bB \sin(c+dx) \sqrt{\cos(c+dx)}}{d\sqrt{a \cos(c+dx) + a}} \right)$$

↓ 3461

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(a-b)(A-B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + (2aB+2Ab-bB) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}}}{2a} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(a-b)(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + (2aB+2Ab-bB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{2a} \right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(a-b)(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2(2aB+2Ab-bB) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}}}{2a} \right)$$

3.528. $\int \frac{(aA+(Ab+aB) \cos(c+dx)+bB \cos^2(c+dx)) \sqrt{\sec(c+dx)}}{\sqrt{a+a \cos(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 223 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(a-b)(A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx + \frac{2\sqrt{a}(2aB+2Ab-bB) \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{2a} \right) \\
 & \downarrow 3261 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2\sqrt{a}(2aB+2Ab-bB) \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{4a^2(a-b)(A-B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d}{d}}{2a} \right) \\
 & \downarrow 218 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2\sqrt{a}(2aB+2Ab-bB) \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} + \frac{2\sqrt{2}\sqrt{a}(a-b)(A-B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d}}{2a} \right)
 \end{aligned}$$

input `Int[((a*A + (A*b + a*B)*Cos[c + d*x] + b*B*Cos[c + d*x]^2)*Sqrt[Sec[c + d*x]])/Sqrt[a + a*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*Sqrt[a]*(2*A*b + 2*a*B - b*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (2*Sqrt[2]*Sqrt[a]*(a - b)*(A - B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]])*Sqrt[a + a*Cos[c + d*x]])]/d)/(2*a) + (b*B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))`

3.528.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.528. $\int \frac{(aA+(Ab+aB)\cos(c+dx)+bB\cos^2(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+a\cos(c+dx)}} dx$

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3524 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

rule 4709 `Int[(u_)*((c_)*sec[(a_) + (b_)*(x_)])^(m_), x_Symbol] := Simp[(c*Sec[a + b*x])^m*(c*cos[a + b*x])^m Int[ActivateTrig[u]/(c*cos[a + b*x])^m, x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownSineIntegrandQ[u, x]`

3.528.4 Maple [A] (verified)

Time = 24.58 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.46

method	result
default	$\frac{(B\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}b\sin(dx+c)+2A\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)b+2B\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)a-B\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)a}{d(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a}$
parts	$\frac{(Ab+Ba)(\sqrt{\sec(dx+c)})\left(\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)+\arcsin(\cot(dx+c)-\csc(dx+c))\right)\cos(dx+c)\sqrt{a(1+\cos(dx+c))}\sqrt{2}}{d(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a}$

input `int((a*A+(A*b+B*a)*cos(d*x+c)+b*B*cos(d*x+c)^2)*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2/d*(B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b*sin(d*x+c)+2*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*b+2*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*a-B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*b-2*A*arcsin(cot(d*x+c)-csc(d*x+c))*a+2*A*arcsin(cot(d*x+c)-csc(d*x+c))*b+2*B*arcsin(cot(d*x+c)-csc(d*x+c))*a-2*B*arcsin(cot(d*x+c)-csc(d*x+c))*b)*sec(d*x+c)^(1/2)*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)/(1+cos(d*x+c))/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)/a`

3.528.5 Fracas [A] (verification not implemented)

Time = 13.77 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.08

$$\int \frac{(aA + (Ab + aB)\cos(c + dx) + bB\cos^2(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + a\cos(c + dx)}} dx$$

$$= \frac{\sqrt{a\cos(dx + c) + aBb}\sqrt{\cos(dx + c)}\sin(dx + c) - (2Ba + (2A - B)b + (2Ba + (2A - B)b)\cos(dx + c))}{ad c}$$

3.528.
$$\int \frac{(aA+(Ab+aB)\cos(c+dx)+bB\cos^2(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+a\cos(c+dx)}} dx$$

input `integrate((A*a+(A*b+B*a)*cos(d*x+c)+b*B*cos(d*x+c)^2)*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `(sqrt(a*cos(d*x + c) + a)*B*b*sqrt(cos(d*x + c))*sin(d*x + c) - (2*B*a + (2*A - B)*b + (2*B*a + (2*A - B)*b)*cos(d*x + c))*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - sqrt(2)*((A - B)*a^2 - (A - B)*a*b + ((A - B)*a^2 - (A - B)*a*b)*cos(d*x + c))*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/sqrt(a))/(a*d*cos(d*x + c) + a*d)`

3.528.6 Sympy [F]

$$\int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a (\cos(c + dx) + 1)}} dx$$

input `integrate((A*a+(A*b+B*a)*cos(d*x+c)+b*B*cos(d*x+c)**2)*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sqrt(sec(c + d*x))/sqrt(a*(cos(c + d*x) + 1)), x)`

3.528.7 Maxima [F]

$$\int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx$$

$$= \int \frac{(Bb \cos(dx + c)^2 + Aa + (Ba + Ab) \cos(dx + c)) \sqrt{\sec(dx + c)}}{\sqrt{a \cos(dx + c) + a}} dx$$

input `integrate((A*a+(A*b+B*a)*cos(d*x+c)+b*B*cos(d*x+c)^2)*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(sec(d*x + c))/sqrt(a*cos(d*x + c) + a), x)`

3.528. $\int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx$

3.528.8 Giac [F(-1)]

Timed out.

$$\int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A*a+(A*b+B*a)*cos(d*x+c)+b*B*cos(d*x+c)^2)*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.528.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(aA + (Ab + aB) \cos(c + dx) + bB \cos^2(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + a \cos(c + dx)}} dx \\ &= \int \frac{\sqrt{\frac{1}{\cos(c+dx)}} (B b \cos(c + dx)^2 + (Ab + Ba) \cos(c + dx) + Aa)}{\sqrt{a + a \cos(c + dx)}} dx \end{aligned}$$

input `int(((1/cos(c + d*x))^(1/2)*(A*a + cos(c + d*x)*(A*b + B*a) + B*b*cos(c + d*x)^2))/(a + a*cos(c + d*x))^(1/2),x)`

output `int(((1/cos(c + d*x))^(1/2)*(A*a + cos(c + d*x)*(A*b + B*a) + B*b*cos(c + d*x)^2))/(a + a*cos(c + d*x))^(1/2), x)`

3.529
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.529.1 Optimal result	4800
3.529.2 Mathematica [C] (warning: unable to verify)	4801
3.529.3 Rubi [A] (verified)	4801
3.529.4 Maple [A] (verified)	4808
3.529.5 Fricas [A] (verification not implemented)	4809
3.529.6 Sympy [F(-1)]	4809
3.529.7 Maxima [F]	4810
3.529.8 Giac [F]	4810
3.529.9 Mupad [F(-1)]	4810

3.529.1 Optimal result

Integrand size = 35, antiderivative size = 317

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(19A - 15B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{a+a \cos(c+dx)}}{2\sqrt{2}a^{3/2}d} - \frac{(1201A - 1029B)\sqrt{\sec(c + dx)} \sin(c + dx)}{210ad\sqrt{a + a \cos(c + dx)}} + \frac{(397A - 273B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{210ad\sqrt{a + a \cos(c + dx)}} - \frac{(67A - 63B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{70ad\sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}} + \frac{(11A - 7B) \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{14ad\sqrt{a + a \cos(c + dx)}}$$

output

```
-1/2*(A-B)*sec(d*x+c)^(7/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)+1/210*(397
*A-273*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)-1/70*(67*
A-63*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/14*(11*A-
7*B)*sec(d*x+c)^(7/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/4*(19*A-15*B
)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(
1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d*2^(1/2)-1/210*(1201*A-1
029*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.529.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.38 (sec) , antiderivative size = 2966, normalized size of antiderivative = 9.36

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Result too large to show}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2))/(a + a*Cos[c + d*x])^(3/2),x]`

output `(2*Cos[c/2 + (d*x)/2]^3*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*(-1/28*((A - B)*(1 - 2*Sin[c/2 + (d*x)/2]))/((1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(7/2)) + ((A - B)*(1 + 2*Sin[c/2 + (d*x)/2]))/(28*(1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(7/2)) - ((A - B)*(315*ArcTan[(1 - 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]] + (5 + 3*Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) - (11 + 17*Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (61 + 71*Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (193*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]/(1 - Sin[c/2 + (d*x)/2]))/70 + ((A - B)*(315*ArcTan[(1 + 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]] + (5 - 3*Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) - (11 - 17*Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) + (61 - 71*Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (193*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]/(1 + Sin[c/2 + (d*x)/2]))/70 - ((-A - 3*B)*Csc[c/2 + (d*x)/2]^9*(363825*Sin[c/2 + (d*x)/2]^2 - 4729725*Sin[c/2 + (d*x)/2]^4 + 26785605*Sin[c/2 + (d*x)/2]^6 - 86790165*Sin[c/2 + (d*x)/2]^8 + 177677808*Sin[c/2 + (d*x)/2]^10 - 239283044*Sin[c/2 + (d*x)/2]^12 + 52080*Hypergeometric2F1[2, 11/2, 13/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2))`

3.529.3 Rubi [A] (verified)

Time = 2.03 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.09, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.529. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{\sec^{\frac{9}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc(c+dx+\frac{\pi}{2})^{9/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{9}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{9/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx \\
& \quad \downarrow \text{3457} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A-7B)-8a(A-B)\cos(c+dx)}{2\cos^{\frac{9}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{7}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A-7B)-8a(A-B)\cos(c+dx)}{\cos^{\frac{9}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{7}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A-7B)-8a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{9/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{7}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3463} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int \frac{-a^2(67A-63B)-6a^2(11A-7B)\cos(c+dx)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{7a} + \frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{(A-B)}{2d\cos^{\frac{7}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.529. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(67A-63B)-6a^2(11A-7B)\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{7a}}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{7}{2}}(c+dx)(a\cos(c+dx)+a)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(67A-63B)-6a^2(11A-7B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{7a}}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{7}{2}}(c+dx)(a\cos(c+dx)+a)} \right)$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{a^3(397A-273B)-4a^3(67A-63B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{7a}}{4a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(397A-273B)-4a^3(67A-63B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a}}{7a}}{4a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(397A-273B)-4a^3(67A-63B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a}}{7a}}{4a^2}$$

3.529. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{a^4(1201A-1029B)-2a^4(397A-273B)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)+a}}}{3a}}{7a}}{4a^2} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int a^4(1201A-1029B)-2a^4(397A-273B)}{5}}{7a}}{4a^2} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int a^4(1201A-1029B)-2a^4(397A-273B)}{5}}{7a}}{4a^2} \right)$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int - \frac{2\sqrt{c}}{2\sqrt{c}}}{7a}}{4a^2} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^4(1201)}{d\sqrt{\cos(c+dx)}}}{7a}}{4a^2} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^4(1201)}{d\sqrt{\cos(c+dx)}}}{7a}}{4a^2} \right)$$

↓ 3261

3.529. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\left(\frac{\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}}{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{210a^5(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}}} \right) \frac{1}{4a^2}$$

↓ 218

$$\left(\frac{\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}}{\frac{2a(11A-7B)\sin(c+dx)}{7d\cos^{\frac{7}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(67A-63B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(397A-273B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(1201A-1029B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}}} \right) \frac{1}{4a^2}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2))/(a + a*Cos[c + d*x])^(3/2), x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/2*((A - B)*Sin[c + d*x])/(d*Cos[
c + d*x]^(7/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a*(11*A - 7*B)*Sin[c + d*
x])/(7*d*Cos[c + d*x]^(7/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*a^2*(67*A - 63
*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*
a^3*(397*A - 273*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c
+ d*x]]) - ((-105*Sqrt[2]*a^(7/2)*(19*A - 15*B)*ArcTan[(Sqrt[a]*Sin[c + d
*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^4*(1
201*A - 1029*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x
]]))/(3*a))/(5*a))/(7*a))/(4*a^2))
```

3.529.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.529.4 Maple [A] (verified)

Time = 9.58 (sec) , antiderivative size = 497, normalized size of antiderivative = 1.57

method	result
default	$-\frac{\left(\sec^{\frac{9}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(1995A \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^6(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-1575B \arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{A\sqrt{a(1+\cos(dx+c))}\left(\sec^{\frac{9}{2}}(dx+c)\right)\left(1995(\cos^6(dx+c)) \arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+1201(\cos^5(dx+c))\sqrt{2} \sin(dx+c)\right)}$
parts	

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/420/a^2/d*sec(d*x+c)^(9/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^2*(1
995*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^6*(cos(d*x+c)/(1+cos(d*x+c)
))^1/2-1575*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^6*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)+3990*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(
d*x+c)/(1+cos(d*x+c)))^(1/2)+1201*A*sin(d*x+c)*cos(d*x+c)^5*2^(1/2)-3150*B
*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1
/2)-1029*B*sin(d*x+c)*cos(d*x+c)^5*2^(1/2)+1995*A*cos(d*x+c)^4*arcsin(cot(
d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+804*A*sin(d*x+c)*cos(
d*x+c)^4*2^(1/2)-1575*B*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*
x+c)/(1+cos(d*x+c)))^(1/2)-756*B*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)-196*A*sin
(d*x+c)*cos(d*x+c)^3*2^(1/2)+84*B*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+36*A*sin
(d*x+c)*cos(d*x+c)^2*2^(1/2)-84*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-60*A*sin
(d*x+c)*cos(d*x+c)*2^(1/2)*2^(1/2)
```

$$3.529. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.529.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.75

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{105 \sqrt{2} ((19A - 15B) \cos(dx + c)^5 + 2(19A - 15B) \cos(dx + c)^4 + (19A - 15B) \cos(dx + c)^3) \sqrt{a} \arctan\left(\frac{\sqrt{2} \sqrt{a \cos(dx + c)}}{\sqrt{a \sin(dx + c)}}\right) + 2((1201A - 1029B) \cos(dx + c)^4 + 12(67A - 63B) \cos(dx + c)^3 - 28(7A - 3B) \cos(dx + c)^2 + 12(3A - 7B) \cos(dx + c) - 60A) \sqrt{a \cos(dx + c)} + a \sin(dx + c) / \sqrt{\cos(dx + c)}}{420 (a^2 d \cos(dx + c)^5 + 2a^2 d \cos(dx + c)^4 + a^2 d \cos(dx + c)^3)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+a*cos(d*x+c))^(3/2),x, algo rithm="fricas")`

output `-1/420*(105*sqrt(2)*((19*A - 15*B)*cos(d*x + c)^5 + 2*(19*A - 15*B)*cos(d*x + c)^4 + (19*A - 15*B)*cos(d*x + c)^3)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((1201*A - 1029*B)*cos(d*x + c)^4 + 12*(67*A - 63*B)*cos(d*x + c)^3 - 28*(7*A - 3*B)*cos(d*x + c)^2 + 12*(3*A - 7*B)*cos(d*x + c) - 60*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*d*cos(d*x + c)^5 + 2*a^2*d*cos(d*x + c)^4 + a^2*d*cos(d*x + c)^3)`

3.529.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(9/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.529.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{9}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+a*cos(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(9/2)/(a*cos(d*x + c) + a)^(3/2), x)`

3.529.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{9}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)/(a+a*cos(d*x+c))^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(9/2)/(a*cos(d*x + c) + a)^(3/2), x)`

3.529.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{9/2}}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2))/(a + a*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2))/(a + a*cos(c + d*x))^(3/2), x)`

3.529. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.530
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.530.1 Optimal result 4811
 3.530.2 Mathematica [C] (warning: unable to verify) 4812
 3.530.3 Rubi [A] (verified) 4812
 3.530.4 Maple [A] (verified) 4817
 3.530.5 Fricas [A] (verification not implemented) 4818
 3.530.6 Sympy [F(-1)] 4819
 3.530.7 Maxima [F] 4819
 3.530.8 Giac [F] 4819
 3.530.9 Mupad [F(-1)] 4820

3.530.1 Optimal result

Integrand size = 35, antiderivative size = 270

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$-\frac{(15A - 11B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2\sqrt{2}a^{3/2}d}$$

$$+ \frac{(147A - 95B) \sqrt{\sec(c + dx)} \sin(c + dx)}{30ad \sqrt{a + a \cos(c + dx)}} - \frac{(39A - 35B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{30ad \sqrt{a + a \cos(c + dx)}}$$

$$- \frac{(A - B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}} + \frac{(9A - 5B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{10ad \sqrt{a + a \cos(c + dx)}}$$

output

```
-1/2*(A-B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)-1/30*(39*A-35*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/10*(9*A-5*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)-1/4*(15*A-11*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d*2^(1/2)+1/30*(147*A-95*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.530.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.23 (sec) , antiderivative size = 2166, normalized size of antiderivative = 8.02

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Result too large to show}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/(a + a*Cos[c + d*x])^(3/2),x]`

output `(2*Cos[c/2 + (d*x)/2]^3*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*(-1/20*((A - B)*(1 - 2*Sin[c/2 + (d*x)/2]))/((1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) + ((A - B)*(1 + 2*Sin[c/2 + (d*x)/2]))/(20*(1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(5/2)) - ((A - B)*(-105*ArcTan[(1 - 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]] + (4 + 3*Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) - (19 + 29*Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) - (67*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])/((1 - Sin[c/2 + (d*x)/2])))/30 + ((A - B)*(-105*ArcTan[(1 + 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]] + (4 - 3*Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^2)^(3/2)) - (19 - 29*Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) - (67*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])/((1 + Sin[c/2 + (d*x)/2])))/30 + ((-A - 3*B)*Csc[c/2 + (d*x)/2]^7*(4725*Sin[c/2 + (d*x)/2]^2 - 48825*Sin[c/2 + (d*x)/2]^4 + 210105*Sin[c/2 + (d*x)/2]^6 - 486630*Sin[c/2 + (d*x)/2]^8 + 655812*Sin[c/2 + (d*x)/2]^10 - 710*Hypergeometric2F1[2, 9/2, 11/2, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^10 - 40*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 9/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^10 - 518760*Sin[c/2 + (d*x)/2]^12 + 1770*Hypergeometric2F1[...`

3.530.3 Rubi [A] (verified)

Time = 1.63 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.530. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{\sec^{\frac{7}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\csc(c+dx+\frac{\pi}{2})^{7/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx \\
& \quad \downarrow \text{3457} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(9A-5B)-6a(A-B)\cos(c+dx)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(9A-5B)-6a(A-B)\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(9A-5B)-6a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3463} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int \frac{-a^2(39A-35B)-4a^2(9A-5B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a} + \frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{(A-B)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.530. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(39A-35B)-4a^2(9A-5B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{5a}}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^2(39A-35B)-4a^2(9A-5B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{5a}}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)} \right)$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{a^3(147A-95B)-2a^3(39A-35B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{5a}}{4a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(147A-95B)-2a^3(39A-35B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a}}{5a}}{4a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(147A-95B)-2a^3(39A-35B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a}}{5a}}{4a^2}$$

3.530. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{array}{c} \downarrow \text{3463} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{15a^4(15A-11B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2}{d\sqrt{\cos(c+dx)}} \right) \\ \hline 4a^2 \end{array} \right) \end{array}$$

$$\begin{array}{c} \downarrow \text{27} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(147A-95B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^3(15A-11B)}{3a} \right) \\ \hline 4a^2 \end{array} \right) \end{array}$$

$$\begin{array}{c} \downarrow \text{3042} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(147A-95B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^3(15A-11B)}{3a} \right) \\ \hline 4a^2 \end{array} \right) \end{array}$$

$$\begin{array}{c} \downarrow \text{3261} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{30a^4(15A-11B)\int \frac{1}{\sin(c+dx)\tan(c+dx)a^3} dx}{d} \right) \\ \hline 4a^2 \end{array} \right) \end{array}$$

$$\downarrow \text{218}$$

3.530. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(9A-5B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(39A-35B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(147A-95B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15\sqrt{2}a^{5/2}}{5a}}{4a^2} \right)$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/(a + a*Cos[c + d*x])^(3/2),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/2*((A - B)*Sin[c + d*x])/(d*Cos[
c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a*(9*A - 5*B)*Sin[c + d*x
])/((5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) - ((2*a^2*(39*A - 35*
B)*Sin[c + d*x])/((3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-15
*Sqrt[2]*a^(5/2)*(15*A - 11*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt
[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^3*(147*A - 95*B)*Sin[c
+ d*x])/((d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])/(3*a))/(5*a))/(4
*a^2))
```

3.530.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.530. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3457 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3463 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.530.4 Maple [A] (verified)

Time = 10.38 (sec) , antiderivative size = 457, normalized size of antiderivative = 1.69

method	result
default	$\left(\sec^{\frac{7}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(225A\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-165B\arcsin(\cot(dx+c)-\csc(dx+c))\right)$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))}\left(\sec^{\frac{7}{2}}(dx+c)\right)\left(75\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))+49\sqrt{2}(\cos^4(dx+c))\sin(dx+c)\right)}{1}$

3.530.
$$\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$$


```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/60/a^2/d*sec(d*x+c)^(7/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^2*(225
*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)-165*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)+147*A*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)+450*A*cos(d*x+c)^4*ar
csin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-95*B*sin(d*x
+c)*cos(d*x+c)^4*2^(1/2)-330*B*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*
(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+108*A*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+22
5*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d
*x+c))-60*B*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)-165*B*cos(d*x+c)^3*(cos(d*x+c)
/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-12*A*sin(d*x+c)*cos(d
*x+c)^2*2^(1/2)+20*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)+12*A*sin(d*x+c)*cos(d
*x+c)*2^(1/2))*2^(1/2)
```

3.530.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.81

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{15 \sqrt{2} ((15A - 11B) \cos(dx + c)^4 + 2(15A - 11B) \cos(dx + c))}{(a + a \cos(c + dx))^{3/2}}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="fricas")
```

```
output 1/60*(15*sqrt(2))*((15*A - 11*B)*cos(d*x + c)^4 + 2*(15*A - 11*B)*cos(d*x +
c)^3 + (15*A - 11*B)*cos(d*x + c)^2)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*
x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((147*A - 95*B)
*cos(d*x + c)^3 + 12*(9*A - 5*B)*cos(d*x + c)^2 - 4*(3*A - 5*B)*cos(d*x +
c) + 12*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*
d*cos(d*x + c)^4 + 2*a^2*d*cos(d*x + c)^3 + a^2*d*cos(d*x + c)^2)
```

3.530. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.530.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(7/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.530.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^(3/
2), x)`

3.530.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^(3/
2), x)`

3.530.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{7/2}}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + a*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + a*cos(c + d*x))^(3/2), x)`

3.531
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.531.1 Optimal result 4821
 3.531.2 Mathematica [C] (warning: unable to verify) 4822
 3.531.3 Rubi [A] (verified) 4823
 3.531.4 Maple [B] (verified) 4827
 3.531.5 Fricas [A] (verification not implemented) 4828
 3.531.6 Sympy [F(-1)] 4828
 3.531.7 Maxima [F] 4829
 3.531.8 Giac [F] 4829
 3.531.9 Mupad [F(-1)] 4829

3.531.1 Optimal result

Integrand size = 35, antiderivative size = 223

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(11A - 7B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2\sqrt{2}a^{3/2}d} - \frac{(19A - 15B)\sqrt{\sec(c + dx)} \sin(c + dx)}{6ad\sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}} + \frac{(7A - 3B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{6ad\sqrt{a + a \cos(c + dx)}}$$

output

```
-1/2*(A-B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)+1/6*(7*A-3
*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)+1/4*(11*A-7*B)*
arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1
/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d*2^(1/2)-1/6*(19*A-15*B)*s
in(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.531.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.59 (sec) , antiderivative size = 981, normalized size of antiderivative = 4.40

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{2 \cos^3\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)} \left(-\frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{12(1 + \sin(c + dx))} \right)$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x])^(3/2),x]`

output

```
(2*Cos[c/2 + (d*x)/2]^3*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*(-1/12*((A - B)*(1 - 2*Sin[c/2 + (d*x)/2]))/((1 + Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^(3/2)) + ((A - B)*(1 + 2*Sin[c/2 + (d*x)/2]))/(12*(1 - Sin[c/2 + (d*x)/2])*(1 - 2*Sin[c/2 + (d*x)/2]^(3/2)) - ((A - B)*(5*ArcTan[(1 - 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (1 + Sin[c/2 + (d*x)/2])/((1 - Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (3*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])/(1 - Sin[c/2 + (d*x)/2]))/2 + ((A - B)*(5*ArcTan[(1 + 2*Sin[c/2 + (d*x)/2])/Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (1 - Sin[c/2 + (d*x)/2])/((1 + Sin[c/2 + (d*x)/2])*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]) + (3*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2])/(1 + Sin[c/2 + (d*x)/2]))/2 + ((A + 3*B)*Csc[c/2 + (d*x)/2]^5*(-12*Cos[(c + d*x)/2]^4*HypergeometricPFQ[{2, 2, 7/2}, {1, 9/2}, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*Sin[c/2 + (d*x)/2]^8 - 12*Hypergeometric2F1[2, 7/2, 9/2, -(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*Sin[c/2 + (d*x)/2]^8*(4 - 7*Sin[c/2 + (d*x)/2]^2 + 3*Sin[c/2 + (d*x)/2]^4) + 7*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*(15 - 20*Sin[c/2 + (d*x)/2]^2 + 8*Sin[c/2 + (d*x)/2]^4)*((3 - 7*Sin[c/2 + (d*x)/2]^2)*Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))] - 3*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])*(1 - 2*Sin[c/2 + (d*x)/2]^2)))/(126*(1 - 2*Si...
```

3.531.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx$$

$$\downarrow \text{3457}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(7A-3B)-4a(A-B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(7A-3B)-4a(A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(7A-3B)-4a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

3.531. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow 3463 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{2 \int \frac{a^2(19A-15B)-2a^2(7A-3B)\cos(c+dx)}{2 \cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx + \frac{2a(7A-3B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}}}{4a^2} - \frac{(A-B)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^2(19A-15B)-2a^2(7A-3B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\int \frac{a^2(19A-15B)-2a^2(7A-3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3463 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{2 \int \frac{3a^3(11A-7B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}}{4a^2} - \frac{(A-B)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} & \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d \cos^{\frac{3}{2}}(c+dx)\sqrt{a \cos(c+dx)+a}} - \frac{\frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}} - 3a^2(11A-7B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)}{2d \cos^{\frac{3}{2}}(c+dx)(a \cos(c+dx)+a)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \end{aligned}$$

3.531. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 3a^2(11A-7B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx)}}}{3a}}{4a^2} \right)$$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{6a^3(11A-7B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d \left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)+a}} \right)}{3a}}{4a^2} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(7A-3B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^2(19A-15B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}a^{3/2}(11A-7B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{3a}}{4a^2} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x])^(3/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/2*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a*(7*A - 3*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*a^(3/2)*(11*A - 7*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^2*(19*A - 15*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(4*a^2))`

3.531. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

3.531.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.531.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 416 vs. 2(188) = 376.

Time = 10.33 (sec) , antiderivative size = 417, normalized size of antiderivative = 1.87

method	result
default	$-\frac{\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(33A(\cos^4(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-21B(\cos^4(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\right)}{a^{\frac{5}{2}}(1+\cos(dx+c))}$
parts	$-\frac{A\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(33\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^4(dx+c))+19\sqrt{2}(\cos^3(dx+c))\sin(dx+c)\right)}{a^{\frac{5}{2}}(1+\cos(dx+c))}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/12/a^2/d*sec(d*x+c)^(5/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^2*(33
*A*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)-21*B*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d
*x+c)))^(1/2)+19*A*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+66*A*cos(d*x+c)^3*(cos(
d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-15*B*sin(d*x+c)
*cos(d*x+c)^3*2^(1/2)-42*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*
arcsin(cot(d*x+c)-csc(d*x+c))+12*A*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)+33*A*co
s(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
-12*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-21*B*cos(d*x+c)^2*arcsin(cot(d*x+c)-
csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-4*A*sin(d*x+c)*cos(d*x+c)*2^(
1/2))*2^(1/2)
```

3.531. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.531.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.88

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{3\sqrt{2}((11A - 7B) \cos(dx + c)^3 + 2(11A - 7B) \cos(dx + c)^2 + (11A - 7B) \cos(dx + c))\sqrt{a} \arctan\left(\frac{\sqrt{2} \cos(dx + c)}{\sqrt{a \cos(dx + c) + a}}\right) + 2(19A - 15B) \cos(dx + c)^2 + 12(A - B) \cos(dx + c) - 4A \sqrt{a \cos(dx + c) + a} \sin(dx + c) / \sqrt{\cos(dx + c)}}{12(a^2 d \cos(dx + c)^3 + 2a^2 d \cos(dx + c)^2 + a^2 d \cos(dx + c))}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(3/2),x, algo rithm="fricas")`

output `-1/12*(3*sqrt(2)*((11*A - 7*B)*cos(d*x + c)^3 + 2*(11*A - 7*B)*cos(d*x + c)^2 + (11*A - 7*B)*cos(d*x + c))*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((19*A - 15*B)*cos(d*x + c)^2 + 12*(A - B)*cos(d*x + c) - 4*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*d*cos(d*x + c)^3 + 2*a^2*d*cos(d*x + c)^2 + a^2*d*cos(d*x + c))`

3.531.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.531.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^(3/2), x)`

3.531.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^(3/2), x)`

3.531.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{5/2}}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(3/2), x)`

3.531. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$

3.532
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.532.1 Optimal result 4830
 3.532.2 Mathematica [C] (warning: unable to verify) 4830
 3.532.3 Rubi [A] (verified) 4831
 3.532.4 Maple [B] (verified) 4835
 3.532.5 Fricas [A] (verification not implemented) 4835
 3.532.6 Sympy [F(-1)] 4836
 3.532.7 Maxima [F] 4836
 3.532.8 Giac [F] 4836
 3.532.9 Mupad [F(-1)] 4837

3.532.1 Optimal result

Integrand size = 35, antiderivative size = 176

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx =$$

$$\frac{(7A - 3B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2\sqrt{2}a^{3/2}d}$$

$$- \frac{(A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2}} + \frac{(5A - B) \sqrt{\sec(c + dx)} \sin(c + dx)}{2ad \sqrt{a + a \cos(c + dx)}}$$

output

```
-1/2*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(3/2)-1/4*(7*A-3
*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c)
)^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d*2^(1/2)+1/2*(5*A-B)*s
in(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(1/2)
```

3.532.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.90 (sec) , antiderivative size = 443, normalized size of antiderivative = 2.52

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\cos^3\left(\frac{1}{2}(c + dx)\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{\dots} \left(30(A - B) \arctan \dots\right)$$

3.532.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^(3/2),x]`

output `(Cos[(c + d*x)/2]^3*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(30*(A - B)*ArcTan[(1 - 2*Sin[(c + d*x)/2])/Sqrt[Cos[c + d*x]]] - 30*(A - B)*ArcTan[(1 + 2*Sin[(c + d*x)/2])/Sqrt[Cos[c + d*x]]] - (20*(A - B)*Sqrt[Cos[c + d*x]])/(-1 + Sin[(c + d*x)/2]) - (20*(A - B)*Sqrt[Cos[c + d*x]])/(1 + Sin[(c + d*x)/2]) + (5*(A - B)*(-1 + 2*Sin[(c + d*x)/2]))/(Sqrt[Cos[c + d*x]]*(Cos[(c + d*x)/4] + Sin[(c + d*x)/4])^2) - (5*(A - B)*(1 + 2*Sin[(c + d*x)/2]))/(Sqrt[Cos[c + d*x]]*(-1 + Sin[(c + d*x)/2])) + ((A + 3*B)*Csc[(c + d*x)/2]^3*(5*(1 + 4*Cos[c + d*x] + Cos[2*(c + d*x)]))*(1 - Cos[c + d*x] + ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]*Cos[c + d*x]*Sqrt[2 - 2*Sec[c + d*x]]) - 2*Hypergeometric2F1[2, 5/2, 7/2, -(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]*Sin[(c + d*x)/2]^4*Sin[c + d*x]*Tan[c + d*x])/(2*Cos[c + d*x]^(3/2)))/(10*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.532.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx$$

↓ 3440

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx$$

3.532. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow \text{3457} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(5A-B)-2a(A-B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{(A-B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \\ & \downarrow \text{27} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(5A-B)-2a(A-B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \\ & \downarrow \text{3042} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(5A-B)-2a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \\ & \downarrow \text{3463} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int -\frac{a^2(7A-3B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a(5A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{(A-B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \\ & \downarrow \text{27} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(5A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a(7A-3B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \\ & \downarrow \text{3042} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(5A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a(7A-3B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \\ & \downarrow \text{3261} \end{aligned}$$

3.532. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^2(7A-3B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2a(5A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)}} \right) \frac{1}{4a^2}$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a(5A-B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{\sqrt{2}\sqrt{a}(7A-3B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{1}{2d\sqrt{\cos(c+dx)}} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^(3/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/2*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)) + (-((Sqrt[2]*Sqrt[a]*(7*A - 3*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d) + (2*a*(5*A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])))/(4*a^2))`

3.532.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.532.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 352 vs. 2(147) = 294.

Time = 10.23 (sec) , antiderivative size = 353, normalized size of antiderivative = 2.01

method	result
default	$\frac{\left(7A(\cos^2(dx+c)) \arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-3B(\cos^2(dx+c)) \arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+5A\right)}{4d(1+\cos(dx+c))}$
parts	$\frac{A\left(7\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c)) \arcsin(\cot(dx+c)-\csc(dx+c))+5\sin(dx+c)\cos(dx+c)\sqrt{2+14\cos(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))\right)}{4d(1+\cos(dx+c))}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{4} \frac{1}{a^2 d} \left(7A \cos^2(dx+c) \arcsin(\cot(dx+c)-\csc(dx+c)) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} - 3B \cos^2(dx+c) \arcsin(\cot(dx+c)-\csc(dx+c)) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} + 5A \sin(dx+c) \cos(dx+c) \sqrt{2+14\cos(dx+c)} \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \arcsin(\cot(dx+c)-\csc(dx+c)) - B \sqrt{2+14\cos(dx+c)} \cos(dx+c) \sin(dx+c) - 6B \cos(dx+c) \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} \arcsin(\cot(dx+c)-\csc(dx+c)) + 4A \sqrt{2+14\cos(dx+c)} \sin(dx+c) + 7 \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} A \arcsin(\cot(dx+c)-\csc(dx+c)) - 3 \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{1/2} B \arcsin(\cot(dx+c)-\csc(dx+c)) \right) \sec(dx+c)^{3/2} (a + \cos(dx+c))^{1/2} \cos(dx+c) / (1 + \cos(dx+c))^2$$

3.532.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.93

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2}((7A - 3B) \cos(dx + c)^2 + 2(7A - 3B) \cos(dx + c) + 7A - 3B) \arctan\left(\frac{\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}}{\sqrt{a} \sin(dx + c)}\right) + 2((5A - B) \cos(dx + c) + 4A) \sqrt{a \cos(dx + c) + a} \sin(dx + c) / \sqrt{\cos(dx + c)}}{4(a^2 d \cos(dx + c) + a^2 d)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output
$$\frac{1}{4} \frac{\sqrt{2} \left((7A - 3B) \cos^2(dx + c) + 2(7A - 3B) \cos(dx + c) + 7A - 3B \right) \arctan\left(\frac{\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}}{\sqrt{a} \sin(dx + c)}\right) + 2 \left((5A - B) \cos(dx + c) + 4A \right) \sqrt{a \cos(dx + c) + a} \sin(dx + c) / \sqrt{\cos(dx + c)}}{4(a^2 d \cos(dx + c) + a^2 d)}$$

3.532.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{3/2}} dx$$

3.532.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.532.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(3/
2), x)`

3.532.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(3/
2), x)`

3.532.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(3/2), x)`

3.533
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{3/2}} dx$$

3.533.1 Optimal result 4838
 3.533.2 Mathematica [A] (verified) 4838
 3.533.3 Rubi [A] (verified) 4839
 3.533.4 Maple [B] (verified) 4841
 3.533.5 Fricas [A] (verification not implemented) 4842
 3.533.6 Sympy [F] 4842
 3.533.7 Maxima [F] 4843
 3.533.8 Giac [F(-1)] 4843
 3.533.9 Mupad [F(-1)] 4843

3.533.1 Optimal result

Integrand size = 35, antiderivative size = 127

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \frac{(3A + B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} + (A - B) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

output

```
-1/2*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+1/4*(3*A+B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d*2^(1/2)
```

3.533.2 Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.50

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \cot\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(3A \operatorname{arctanh}\left(\sqrt{-\sec(c + dx) \sin^2\left(\frac{1}{2}(c + dx)\right)}\right) \cos^2\left(\frac{1}{2}(c + dx)\right) \cos(c + dx) + B \sin(c + dx)\right)$$

input

```
Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^(3/2), x]
```

3.533.
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{3/2}} dx$$

output
$$\frac{-1/2*(\text{Cot}[(c + d*x)/2]*\text{Sqrt}[\text{Sec}[c + d*x]]*(3*A*\text{ArcTanh}[\text{Sqrt}[-(\text{Sec}[c + d*x]*\text{Sin}[(c + d*x)/2]^2)])*\text{Cos}[(c + d*x)/2]^2*\text{Cos}[c + d*x]*\text{Sqrt}[2 - 2*\text{Sec}[c + d*x]] - 2*(B*\text{ArcSin}[\text{Sin}[(c + d*x)/2]/\text{Sqrt}[\text{Cos}[(c + d*x)/2]^2])* \text{Cos}[(c + d*x)/2]^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[(c + d*x)/2] + (-A + B)*\text{Cos}[c + d*x]*\text{Sin}[(c + d*x)/2]^2)))/(d*(a*(1 + \text{Cos}[c + d*x]))^{3/2})}$$

3.533.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{3/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx \\ & \quad \downarrow \text{3457} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{a(3A+B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\ & \quad \downarrow \text{27} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3A+B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \right) \end{aligned}$$

3.533. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx}{4a} - \frac{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right) \\
 & \downarrow \text{3261} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(-\frac{(3A+B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2d} - \frac{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right) \\
 & \downarrow \text{218} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3A+B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{(A-B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((3*A + B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) - ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))`

3.533.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.533.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 214 vs. 2(104) = 208.

Time = 10.54 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.69

method	result
default	$-\frac{\left(A\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sin(dx+c)-B\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sin(dx+c)+3A\arcsin(\cot(dx+c)-\csc(dx+c))\cos(dx+c)+B\arcsin(\cot(dx+c)-\csc(dx+c))\right)\sqrt{\sec(dx+c)}}{4a^2d(1+\cos(dx+c))}$
parts	$-\frac{A\left(\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+3\arcsin(\cot(dx+c)-\csc(dx+c))\cos(dx+c)+3\arcsin(\cot(dx+c)-\csc(dx+c))\right)\left(\sqrt{\sec(dx+c)}\right)\sqrt{a^2}}{4d(1+\cos(dx+c))^2\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a^2}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(3/2), x, method=_RETURNVERBOSE)
```

$$3.533. \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{3/2}} dx$$

output
$$-1/4/a^2/d*(A*2^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\sin(d*x+c)-B*2^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\sin(d*x+c)+3*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)+B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)+3*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))+B*\arcsin(\cot(d*x+c)-\csc(d*x+c)))*\sec(d*x+c)^{(1/2)}*(a*(1+\cos(d*x+c)))^{(1/2)}*\cos(d*x+c)/(1+\cos(d*x+c))^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*2^{(1/2)}$$

3.533.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.13

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \frac{\sqrt{2}((3A + B) \cos(dx + c)^2 + 2(3A + B) \cos(dx + c) + 3A + B) \sqrt{a} \arctan\left(\frac{\sqrt{2} \sqrt{a \cos(dx+c) + a} \sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right)}{4(a^2 d \cos(dx + c)^2 + 2a^2 d \cos(dx + c) + a^2 d)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output
$$-1/4*(\sqrt{2})*((3*A + B)*\cos(d*x + c)^2 + 2*(3*A + B)*\cos(d*x + c) + 3*A + B)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c)) + 2*\sqrt{a*\cos(d*x + c) + a}*(A - B)*\sqrt{\cos(d*x + c)}*\sin(d*x + c)/(a^2*d*\cos(d*x + c)^2 + 2*a^2*d*\cos(d*x + c) + a^2*d)$$

3.533.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a(\cos(c + dx) + 1))^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/(a*(cos(c + d*x) + 1))**(3/2), x)`

3.533.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^(3/2), x)`

3.533.8 Giac [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.533.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + a \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(3/2), x)`

3.534
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$$

3.534.1 Optimal result 4844
 3.534.2 Mathematica [C] (verified) 4845
 3.534.3 Rubi [A] (verified) 4845
 3.534.4 Maple [A] (verified) 4849
 3.534.5 Fricas [A] (verification not implemented) 4849
 3.534.6 Sympy [F] 4850
 3.534.7 Maxima [F] 4850
 3.534.8 Giac [F(-1)] 4851
 3.534.9 Mupad [F(-1)] 4851

3.534.1 Optimal result

Integrand size = 35, antiderivative size = 185

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{a^{3/2} d}$$

$$+ \frac{(A - 5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2\sqrt{2} a^{3/2} d}$$

$$+ \frac{(A - B) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

output

```
1/2*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+2*B*arcsin(
sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/
2)/a^(3/2)/d+1/4*(A-5*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(
1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d*
2^(1/2)
```

3.534.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.16 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.31

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \frac{\cos^3\left(\frac{1}{2}(c + dx)\right) \left(-i\sqrt{2}e^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}}\sqrt{1+e^{2i(c+dx)}}\right)}{4}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(3/2)*Sqrt[Sec[c + d*x]]),x]`

output `(Cos[(c + d*x)/2]^3*(((-I)*Sqrt[2]*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*(4*B*ArcSinh[E^(I*(c + d*x))] - Sqrt[2]*(A - 5*B)*ArcTanh[(1 - E^(I*(c + d*x))]/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]]) - 4*B*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]])/E^((I/2)*(c + d*x)) + (A - B)*Sec[(c + d*x)/2]^2*Sqrt[Sec[c + d*x]]*(-Sin[(c + d*x)/2] + Sin[(3*(c + d*x)/2])))/(2*d*(a*(1 + Cos[c + d*x]))^(3/2))`

3.534.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.92, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a \cos(c + dx) + a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{3/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.534. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx \\
& \quad \downarrow \text{3456} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+4aB\cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+4aB\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+4aB\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3461} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(A-5B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + 4B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(A-5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + 4B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{3253} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(A-5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{8B \int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}} d \left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{4a^2} + \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \quad \downarrow \text{223}
\end{aligned}$$

3.534. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{3/2}\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(A-5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{8\sqrt{a}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{4a^2} + \frac{(A-B)}{2d} \right)$$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{8\sqrt{a}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{2a^2(A-5B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d \left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\sqrt{2}\sqrt{a}(A-5B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d} + \frac{8\sqrt{a}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} + \frac{(A-B)}{2d} \right)$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(3/2)*Sqrt[Sec[c + d*x]]),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((8*Sqrt[a]*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (Sqrt[2]*Sqrt[a]*(A - 5*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/(4*a^2) + ((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))`

3.534.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.534. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3456 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3461 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.534.4 Maple [A] (verified)

Time = 7.44 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.53

method	result
default	$\frac{(A\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sin(dx+c)+4B\sqrt{2}\arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)\cos(dx+c)-B\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sin(dx+c)-A\arcsin(\cot(dx+c)-\csc(dx+c)))\sqrt{a(1+\cos(dx+c))}\sqrt{2}}{4d(1+\cos(dx+c))^2\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a^2}$
parts	

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2)/sec(d*x+c)^(1/2),x,method=_RETURNERVERBOSE)
```

```
output 1/4/a^2/d*(A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)+4*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*cos(d*x+c)-B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sin(d*x+c)-A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)+4*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+5*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)-A*arcsin(cot(d*x+c)-csc(d*x+c))+5*B*arcsin(cot(d*x+c)-csc(d*x+c))*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^2/sec(d*x+c)^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*2^(1/2)
```

3.534.5 Fracas [A] (verification not implemented)

Time = 2.89 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.10

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \frac{\sqrt{2}((A - 5B) \cos(dx + c)^2 + 2(A - 5B) \cos(dx + c) + A - 5B) \sqrt{a} \arctan\left(\frac{\sqrt{2}\sqrt{a \cos(dx+c)+a}\sqrt{\cos(dx+c)}}{\sqrt{a} \sin(dx+c)}\right)}{4(a^2d}$$

3.534. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/4*(sqrt(2)*((A - 5*B)*cos(d*x + c)^2 + 2*(A - 5*B)*cos(d*x + c) + A - 5*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - 2*sqrt(a*cos(d*x + c) + a)*(A - B)*sqrt(cos(d*x + c))*sin(d*x + c) + 8*(B*cos(d*x + c)^2 + 2*B*cos(d*x + c) + B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c)))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.534.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{(a (\cos(c + dx) + 1))^{\frac{3}{2}} \sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2)/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/((a*(cos(c + d*x) + 1))**(3/2)*sqrt(sec(c + d*x))), x)`

3.534.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{\frac{3}{2}} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*x + c))), x)`

3.534.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.534.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(3/2)), x)`

3.535 $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$

3.535.1 Optimal result	4852
3.535.2 Mathematica [C] (verified)	4853
3.535.3 Rubi [A] (verified)	4854
3.535.4 Maple [A] (verified)	4859
3.535.5 Fricas [A] (verification not implemented)	4860
3.535.6 Sympy [F(-1)]	4860
3.535.7 Maxima [F]	4860
3.535.8 Giac [F(-1)]	4861
3.535.9 Mupad [F(-1)]	4861

3.535.1 Optimal result

Integrand size = 35, antiderivative size = 237

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{(2A - 3B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{a^{3/2}d} - \frac{(5A - 9B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{2\sqrt{2}a^{3/2}d} + \frac{(A - B) \sin(c + dx)}{2d(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} - \frac{(A - 3B) \sin(c + dx)}{2ad \sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}}$$

```
output 1/2*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2)-1/2*(A-3*B)
*sine(d*x+c)/a/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+(2*A-3*B)*arcsin(s
in(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2
)/a^(3/2)/d-1/4*(5*A-9*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)
^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(3/2)/d
*2^(1/2)
```

3.535.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.44 (sec) , antiderivative size = 836, normalized size of antiderivative = 3.53

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \\
& \frac{iAe^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) \cos^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{d(a(1+\cos(c+dx)))^{3/2}} \\
& + \frac{3iBe^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) \cos^3\left(\frac{c}{2} + \frac{dx}{2}\right)}{d(a(1+\cos(c+dx)))^{3/2}} \\
& + \frac{2i\sqrt{2}Ae^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \left(-\operatorname{arcsinh}(e^{i(c+dx)}) + \sqrt{2}\operatorname{arctanh}\left(\frac{-1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) + \operatorname{arctanh}\left(\frac{1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right)\right)}{d(a(1+\cos(c+dx)))^{3/2}} \\
& - \frac{3i\sqrt{2}Be^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \left(-\operatorname{arcsinh}(e^{i(c+dx)}) + \sqrt{2}\operatorname{arctanh}\left(\frac{-1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) + \operatorname{arctanh}\left(\frac{1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right)\right)}{d(a(1+\cos(c+dx)))^{3/2}} \\
& + \frac{\cos^3\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c+dx)} \left(-\frac{2A \cos\left(\frac{dx}{2}\right) \sin\left(\frac{c}{2}\right)}{d} + \frac{\sec\left(\frac{c}{2}\right) \sec\left(\frac{c}{2} + \frac{dx}{2}\right) (A \sin\left(\frac{c}{2}\right) - B \sin\left(\frac{c}{2}\right))}{d} + \frac{2B \cos\left(\frac{3dx}{2}\right) \sin\left(\frac{3c}{2}\right)}{d} - \frac{2A \cos\left(\frac{3dx}{2}\right) \sin\left(\frac{3c}{2}\right)}{d}\right)}{(a(1+\cos(c+dx)))^{3/2}}
\end{aligned}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(3/2)*Sec[c + d*x]^(3/2)), x]`

output

```

((-I)*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*
(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c +
d*x))])] * Cos[c/2 + (d*x)/2]^3 / (d * E^((I/2)*(c + d*x)) * (a * (1 + Cos[c + d*x]
))^ (3/2)) + ((3*I)*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[
1 + E^((2*I)*(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E
^((2*I)*(c + d*x))])] * Cos[c/2 + (d*x)/2]^3 / (d * E^((I/2)*(c + d*x)) * (a * (1 +
Cos[c + d*x]))^ (3/2)) + ((2*I)*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*
I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*(-ArcSinh[E^(I*(c + d*x))] +
Sqrt[2]*ArcTanh[-1 + E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*
x))]))] + ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]] * Cos[c/2 + (d*x)/2]^3 / (d *
E^((I/2)*(c + d*x)) * (a * (1 + Cos[c + d*x]))^ (3/2)) - ((3*I)*Sqrt[2]*B*Sqrt[
E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*(-
ArcSinh[E^(I*(c + d*x))] + Sqrt[2]*ArcTanh[-1 + E^(I*(c + d*x)))/(Sqrt[2
]*Sqrt[1 + E^((2*I)*(c + d*x))]))] + ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]]
)*Cos[c/2 + (d*x)/2]^3 / (d * E^((I/2)*(c + d*x)) * (a * (1 + Cos[c + d*x]))^ (3/2
)) + (Cos[c/2 + (d*x)/2]^3 * Sqrt[Sec[c + d*x]] * ((-2*A * Cos[(d*x)/2] * Sin[c/2]
)/d + (Sec[c/2] * Sec[c/2 + (d*x)/2] * (A * Sin[c/2] - B * Sin[c/2]))/d + (2*B * Cos
[(3*d*x)/2] * Sin[(3*c)/2])/d - (2*A * Cos[c/2] * Sin[(d*x)/2])/d + (Sec[c/2] * Se
c[c/2 + (d*x)/2]^2 * (A * Sin[(d*x)/2] - B * Sin[(d*x)/2]))/d + (2*B * Cos[(3*c)/2
]*Sin[(3*d*x)/2])/d) / (a * (1 + Cos[c + d*x]))^ (3/2)

```

3.535.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.95, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3462, 25, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{3/2}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{3/2}} dx
 \end{aligned}$$

3.535. $\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx \\
& \downarrow \text{3456} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)-2a(A-3B)\cos(c+dx))}{2\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)-2a(A-3B)\cos(c+dx))}{\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)-2a(A-3B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \downarrow \text{3462} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{-\frac{a^2(A-3B)-2a^2(2A-3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \downarrow \text{25} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{-\frac{\int \frac{a^2(A-3B)-2a^2(2A-3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \\
& \downarrow \text{3042}
\end{aligned}$$

3.535. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(A-3B)-2a^2(2A-3B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} + \frac{(A-B)\sin(c+dx)}{2d(a\cos(c+dx)+a)} \right)$$

↓ 3461

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(5A-9B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - 2a(2A-3B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(5A-9B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - 2a(2A-3B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} \right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(5A-9B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{4a(2A-3B) \int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{d}}{4a^2} \right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(5A-9B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{4a^{3/2}(2A-3B) \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{2a(A-3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}}}{4a^2} \right)$$

↓ 3261

3.535. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^3(5A-9B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right) - \frac{4a^{3/2}(2A-3B) \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{a} \right) \frac{1}{4a^2}$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{\sqrt{2}a^{3/2}(5A-9B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right) - \frac{4a^{3/2}(2A-3B) \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{a} - \frac{2a(A-B)}{4a^2} \right)$$

```
input Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(3/2)*Sec[c + d*x]^(3/2)),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)) + (-(((4*a^(3/2)*(2*A - 3*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (Sqrt[2]*a^(3/2)*(5*A - 9*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d)/a) - (2*a*(A - 3*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2))
```

3.535.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

3.535. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sec^2(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3462 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.535.4 Maple [A] (verified)

Time = 20.86 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.63

method	result
default	$\frac{\sqrt{a(1+\cos(dx+c))} \left(2B\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 4A\sqrt{2} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) - \tan(dx+c)A\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 6 \right)}{4d(1+\cos(dx+c))^2 \sec(dx+c)^{\frac{3}{2}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))} \left(4\sqrt{2} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) - \tan(dx+c)\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 4\sec(dx+c)\sqrt{2} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{4d(1+\cos(dx+c))^2 \sec(dx+c)^{\frac{3}{2}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(3/2)/sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output 1/4/a^2/d*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^2/sec(d*x+c)^(3/2)/(cos(
d*x+c)/(1+cos(d*x+c)))^(1/2)*(2*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2
)*sin(d*x+c)+4*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2))-tan(d*x+c)*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-6*B*2^(1/2)*arc
tan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+3*tan(d*x+c)*B*2^(1/2)*(
cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4*sec(d*x+c)*A*2^(1/2)*arctan(tan(d*x+c)*
(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+5*A*arcsin(cot(d*x+c)-csc(d*x+c))-6*sec
(d*x+c)*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))-9*B
*arcsin(cot(d*x+c)-csc(d*x+c))+5*sec(d*x+c)*A*arcsin(cot(d*x+c)-csc(d*x+c)
)-9*sec(d*x+c)*B*arcsin(cot(d*x+c)-csc(d*x+c)))^2^(1/2)
```

$$3.535. \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.535.5 Fricas [A] (verification not implemented)

Time = 3.75 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.04

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \frac{\sqrt{2}((5A - 9B) \cos(dx + c))^2 + 2(5A - 9B) \cos(dx + c) + 5A}{(a + a \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algo
rithm="fricas")`

output `1/4*(sqrt(2)*((5*A - 9*B)*cos(d*x + c)^2 + 2*(5*A - 9*B)*cos(d*x + c) + 5*A - 9*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - 4*((2*A - 3*B)*cos(d*x + c)^2 + 2*(2*A - 3*B)*cos(d*x + c) + 2*A - 3*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*(2*B*cos(d*x + c)^2 - (A - 3*B)*cos(d*x + c)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^2*d*cos(d*x + c)^2 + 2*a^2*d*cos(d*x + c) + a^2*d)`

3.535.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(3/2)/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.535.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algo
rithm="maxima")`

3.535. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{3/2} \sec^{3/2}(c+dx)} dx$

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2)), x)`

3.535.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `Timed out`

3.535.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} (a + a \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(3/2)), x)`

3.536
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

3.536.1 Optimal result 4862
 3.536.2 Mathematica [C] (verified) 4863
 3.536.3 Rubi [A] (verified) 4863
 3.536.4 Maple [B] (verified) 4870
 3.536.5 Fricas [A] (verification not implemented) 4870
 3.536.6 Sympy [F(-1)] 4871
 3.536.7 Maxima [F(-1)] 4871
 3.536.8 Giac [F] 4872
 3.536.9 Mupad [F(-1)] 4872

3.536.1 Optimal result

Integrand size = 35, antiderivative size = 317

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{(283A - 163B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{16\sqrt{2}a^{5/2}d}$$

$$+ \frac{(2671A - 1495B) \sqrt{\sec(c + dx)} \sin(c + dx)}{240a^2d \sqrt{a + a \cos(c + dx)}}$$

$$- \frac{(787A - 475B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{240a^2d \sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}}$$

$$- \frac{(21A - 13B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}} + \frac{(157A - 85B) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{80a^2d \sqrt{a + a \cos(c + dx)}}$$

output

```
-1/4*(A-B)*sec(d*x+c)^(5/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)-1/16*(21*A
-13*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)-1/240*(787*A
-475*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)+1/80*(157
*A-85*B)*sec(d*x+c)^(5/2)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)-1/32*(28
3*A-163*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos
(d*x+c)^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(5/2)/d*2^(1/2)+1/240*
(2671*A-1495*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(1/2)
```

3.536.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 10.24 (sec) , antiderivative size = 261, normalized size of antiderivative = 0.82

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\cos^5\left(\frac{1}{2}(c + dx)\right) \left(-240i(283A - 163B)e^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}}\sqrt{1}\right)}{(a + a \cos(c + dx))^{5/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/(a + a*Cos[c + d*x])^(5/2),x]`

output `(Cos[(c + d*x)/2]^5*(((240*I)*(283*A - 163*B)*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]])]/E^((I/2)*(c + d*x)) + (15053*A - 7685*B + 10*(2605*A - 1381*B)*Cos[c + d*x] + 108*(157*A - 85*B)*Cos[2*(c + d*x)] + 9110*A*Cos[3*(c + d*x)] - 5030*B*Cos[3*(c + d*x)] + 2671*A*Cos[4*(c + d*x)] - 1495*B*Cos[4*(c + d*x)])*Sec[(c + d*x)/2]^3*Sec[c + d*x]^(5/2)*Tan[(c + d*x)/2]))/(960*d*(a*(1 + Cos[c + d*x]))^(5/2))`

3.536.3 Rubi [A] (verified)

Time = 2.00 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.09, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{(a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{5/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{A + B \cos(c + dx)}{\cos^{\frac{7}{2}}(c + dx)(\cos(c + dx)a + a)^{5/2}} dx \end{aligned}$$

3.536. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{7/2} (\sin(c + dx + \frac{\pi}{2})a + a)^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sin(c + dx + \frac{\pi}{2})^{7/2} (\sin(c + dx + \frac{\pi}{2})a + a)^{5/2}} dx \\
& \quad \downarrow \text{3457} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a(13A-5B) - 8a(A-B) \cos(c+dx)}{2 \cos^{\frac{7}{2}}(c+dx) (\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B) \sin(c+dx)}{4d \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^{5/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a(13A-5B) - 8a(A-B) \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) (\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \sin(c+dx)}{4d \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^{5/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a(13A-5B) - 8a(A-B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{7/2} (\sin(c+dx + \frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B) \sin(c+dx)}{4d \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^{5/2}} \right) \\
& \quad \downarrow \text{3457} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a^2(157A-85B) - 6a^2(21A-13B) \cos(c+dx)}{2 \cos^{\frac{7}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{8a^2} - \frac{a(21A-13B) \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^{3/2}} - \frac{(A-B) \sin(c+dx)}{4d \cos^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{a^2(157A-85B) - 6a^2(21A-13B) \cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx) \sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(21A-13B) \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx) (a \cos(c+dx) + a)^{3/2}} - \frac{(A-B) \sin(c+dx)}{4d \cos^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.536. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(157A-85B)-6a^2(21A-13B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{7/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2} - \frac{a(21A-13B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{4d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int \frac{a^3(787A-475B)-4a^3(157A-85B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{a(21A-13B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(787A-475B)-4a^3(157A-85B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2}}{8a^2} - \frac{a(21A-13B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(787A-475B)-4a^3(157A-85B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2}}{8a^2} - \frac{a(21A-13B)\sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)$$

↓ 3463

3.536. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{2\int -\frac{a^4(2671A-1495B)-2a^4(787A-475B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{5a} \right) \frac{1}{8a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{\int \frac{a^4(2671A-1495B)-2a^4(787A-475B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}}}{3a}}{5a} \right) \frac{1}{8a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{\int \frac{a^4(2671A-1495B)-2a^4(787A-475B)\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}}{3a}}{5a} \right) \frac{1}{8a^2}$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{2\int -\frac{15a^5(283A-163B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a^4(2671A-1495B)\cos(c+dx)}{3a}}{5a} \right) \frac{1}{8a^2}$$

3.536. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{array}{c}
 \downarrow 27 \\
 \left(\begin{array}{l}
 \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^4(283A-163B)}{3a} \\
 \hline
 \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^4(283A-163B)}{3a} \\
 \hline
 4a^2 \\
 \hline
 8a^2
 \end{array} \right) \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 3042 \\
 \left(\begin{array}{l}
 \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^4(283A-163B)}{3a} \\
 \hline
 \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^4(283A-163B)}{3a} \\
 \hline
 4a^2 \\
 \hline
 8a^2
 \end{array} \right) \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 3261 \\
 \left(\begin{array}{l}
 \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^4(283A-163B)}{3a} \\
 \hline
 \frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15a^4(283A-163B)}{3a} \\
 \hline
 4a^2 \\
 \hline
 8a^2
 \end{array} \right) \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}
 \end{array}$$

\downarrow 218

3.536. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(157A-85B)\sin(c+dx)}{5d\cos^{\frac{5}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(787A-475B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(2671A-1495B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{15\sqrt{2}a^{7/2}(283A-163B)}{5a}}{4a^2} \right) \frac{1}{8a^2}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/(a + a*Cos[c + d*x])^(5/2),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/4*((A - B)*Sin[c + d*x])/(d*Cos[
c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(21*A - 13*B)*Sin[c
+ d*x])/(d*Cos[c + d*x]^(5/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a^2*(157*A
- 85*B)*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)*Sqrt[a + a*Cos[c + d*x]]) -
((2*a^3*(787*A - 475*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*
Cos[c + d*x]]) - ((-15*Sqrt[2]*a^(7/2)*(283*A - 163*B)*ArcTan[(Sqrt[a]*Sin
[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*
a^4*(2671*A - 1495*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c
+ d*x]]))/(3*a))/(5*a))/(4*a^2))/(8*a^2))
```

3.536.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

3.536. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.536.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 588 vs. $2(270) = 540$.

Time = 10.84 (sec) , antiderivative size = 589, normalized size of antiderivative = 1.86

method	result
default	$\left(\sec^{\frac{7}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(4245A\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^6(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-2445B\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)$
parts	$A\sqrt{a(1+\cos(dx+c))}\left(\sec^{\frac{7}{2}}(dx+c)\right)\left(4245(\cos^6(dx+c))\arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+2671(\cos^5(dx+c))\sqrt{2}\sin(dx+c)\right)$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 1/480/a^3/d*sec(d*x+c)^(7/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^3*(42
45*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^6*(cos(d*x+c)/(1+cos(d*x+c))
)^(1/2)-2445*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^6*(cos(d*x+c)/(1+c
os(d*x+c)))^(1/2)+2671*A*sin(d*x+c)*cos(d*x+c)^5*2^(1/2)+12735*A*arcsin(co
t(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-1495*B
*sin(d*x+c)*cos(d*x+c)^5*2^(1/2)-7335*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(
d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4555*A*sin(d*x+c)*cos(d*x+c)^4*
2^(1/2)+12735*A*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)-2515*B*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)-7335*B*cos(d*x+c
)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+1568*A
*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+4245*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*
x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-800*B*sin(d*x+c)*cos(d*x+c)^3*2
^(1/2)-2445*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*
x+c)-csc(d*x+c))-160*A*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)+160*B*sin(d*x+c)*co
s(d*x+c)^2*2^(1/2)+96*A*sin(d*x+c)*cos(d*x+c)*2^(1/2)*2^(1/2)
```

3.536.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.84

$$\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx = \frac{15\sqrt{2}((283A-163B)\cos(dx+c)^5+3(283A-163B)\cos(dx+c))}{(a+a\cos(c+dx))^{5/2}}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(5/2),x, algo
rithm="fracas")
```

3.536.
$$\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$$

output $1/480*(15*\sqrt{2}*((283*A - 163*B)*\cos(d*x + c)^5 + 3*(283*A - 163*B)*\cos(d*x + c)^4 + 3*(283*A - 163*B)*\cos(d*x + c)^3 + (283*A - 163*B)*\cos(d*x + c)^2)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{a*\cos(d*x + c) + a}*\sqrt{\cos(d*x + c)})/(\sqrt{a}*\sin(d*x + c))) + 2*((2671*A - 1495*B)*\cos(d*x + c)^4 + 5*(911*A - 503*B)*\cos(d*x + c)^3 + 32*(49*A - 25*B)*\cos(d*x + c)^2 - 160*(A - B)*\cos(d*x + c) + 96*A)*\sqrt{a*\cos(d*x + c) + a}*\sin(d*x + c)/\sqrt{\cos(d*x + c)})/(a^3*d*\cos(d*x + c)^5 + 3*a^3*d*\cos(d*x + c)^4 + 3*a^3*d*\cos(d*x + c)^3 + a^3*d*\cos(d*x + c)^2)$

3.536.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(7/2)/(a+a*cos(d*x+c))**(5/2),x)`

output Timed out

3.536.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(5/2),x, algorith="maxima")`

output Timed out

3.536.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{(a \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/(a*cos(d*x + c) + a)^(5/2), x)`

3.536.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{7/2}}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + a*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + a*cos(c + d*x))^(5/2), x)`

3.537
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

3.537.1 Optimal result	4873
3.537.2 Mathematica [C] (warning: unable to verify)	4874
3.537.3 Rubi [A] (verified)	4874
3.537.4 Maple [B] (verified)	4879
3.537.5 Fracas [A] (verification not implemented)	4880
3.537.6 Sympy [F(-1)]	4881
3.537.7 Maxima [F(-1)]	4881
3.537.8 Giac [F]	4881
3.537.9 Mupad [F(-1)]	4882

3.537.1 Optimal result

Integrand size = 35, antiderivative size = 270

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{(163A - 75B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}}{16\sqrt{2}a^{5/2}d} - \frac{(299A - 147B)\sqrt{\sec(c + dx) \sin(c + dx)}}{48a^2d\sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}} - \frac{(17A - 9B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}} + \frac{(95A - 39B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{48a^2d\sqrt{a + a \cos(c + dx)}}$$

output

```
-1/4*(A-B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)-1/16*(17*A-9*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)+1/48*(95*A-39*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)+1/32*(163*A-75*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(5/2)/d*2^(1/2)-1/48*(299*A-147*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(1/2)
```


3.537.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.07 (sec) , antiderivative size = 1152, normalized size of antiderivative = 4.27

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{5}{2}}} dx = \text{Too large to display}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x])^(5/2),x]`

output `(2*B*Cos[c/2 + (d*x)/2]^5*Sec[(c + d*x)/2]^4*Sin[c/2 + (d*x)/2]*((1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1))^3/2*((8*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2)/(315*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) + (Csc[c/2 + (d*x)/2]^8*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-15*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Cos[(c + d*x)/2]^4*(-343 + 1465*Sin[c/2 + (d*x)/2]^2 - 2021*Sin[c/2 + (d*x)/2]^4 + 824*Sin[c/2 + (d*x)/2]^6) + sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-5145 + 33980*Sin[c/2 + (d*x)/2]^2 - 87764*Sin[c/2 + (d*x)/2]^4 + 109737*Sin[c/2 + (d*x)/2]^6 - 66122*Sin[c/2 + (d*x)/2]^8 + 15344*Sin[c/2 + (d*x)/2]^10))/120))/(d*(a*(1 + Cos[c + d*x]))^(5/2)) - (A*Cot[c/2 + (d*x)/2]^5*Csc[c/2 + (d*x)/2]^4*Sec[(c + d*x)/2]^4*((1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1))^7/2*(640*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 7/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12 - 1280*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 7/2}, {1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^12*(-6 + 5*Sin[c/2 + (d*x)/2]^2) + 33*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-105*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Cos[(c + d*x)/2]^4*(-10935 + 72902*Sin[c...`

3.537.3 Rubi [A] (verified)

Time = 1.64 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.537. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{\frac{5}{2}}} dx$

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 3440

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{5/2}} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A-3B)-6a(A-B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A-3B)-6a(A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A-3B)-6a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(95A-39B)-4a^2(17A-9B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

3.537. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(95A-39B)-4a^2(17A-9B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(95A-39B)-4a^2(17A-9B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int -\frac{a^3(299A-147B)-2a^3(95A-39B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(299A-147B)-2a^3(95A-39B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a}}{4a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^3(299A-147B)-2a^3(95A-39B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{3a}}{4a^2} - \frac{a(17A-9B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\cos^{\frac{3}{2}}(c+dx)} \right)$$

3.537. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{array}{c} \downarrow \text{3463} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{3a^4(163A-75B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right) \\ \hline 4a^2 \\ \hline 8a^2 \\ \hline 2d\cos^{\frac{3}{2}} \end{array} \right) \end{array}$$

$$\begin{array}{c} \downarrow \text{27} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3a^3(163A-75B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{3a} \right) \\ \hline 4a^2 \\ \hline 8a^2 \end{array} \right) \end{array}$$

$$\begin{array}{c} \downarrow \text{3042} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3a^3(163A-75B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{3a} \right) \\ \hline 4a^2 \\ \hline 8a^2 \end{array} \right) \end{array}$$

$$\begin{array}{c} \downarrow \text{3261} \\ \left(\begin{array}{l} \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{6a^4(163A-75B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{3a} \right) \\ \hline 4a^2 \\ \hline 8a^2 \end{array} \right) \end{array}$$

$$\downarrow \text{218}$$

3.537. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(95A-39B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^3(299A-147B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}a^{5/2}(163A-75B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{3a}}{4a^2} \right) \frac{1}{8a^2}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x])^(5/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/4*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(17*A - 9*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a^2*(95*A - 39*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*a^(5/2)*(163*A - 75*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^3*(299*A - 147*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(3*a))/(4*a^2))/(8*a^2)`

3.537.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

$$3.537. \quad \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3457 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3463 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.537.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 548 vs. 2(229) = 458.

Time = 11.25 (sec) , antiderivative size = 549, normalized size of antiderivative = 2.03

method	result
default	$-\frac{\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(489A\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-225B\arcsin(\cot(dx+c)-\csc(dx+c))\right)}{1}$
parts	$-\frac{A\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(489\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^5(dx+c))+1467\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))\right)}{1}$

$$3.537. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/96/a^3/d*sec(d*x+c)^(5/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^3*(48
9*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))
^(1/2)-225*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos
(d*x+c)))^(1/2)+299*A*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)+1467*A*cos(d*x+c)^4*
arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-147*B*sin(
d*x+c)*cos(d*x+c)^4*2^(1/2)-675*B*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c
))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+503*A*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)
+1467*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-c
sc(d*x+c))-255*B*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)-675*B*cos(d*x+c)^3*(cos(d
*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+160*A*sin(d*x+c)
*cos(d*x+c)^2*2^(1/2)+489*A*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))*(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2)-96*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-225*B*
cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2)-32*A*sin(d*x+c)*cos(d*x+c)*2^(1/2))*2^(1/2)
```

3.537.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.91

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{3\sqrt{2}((163A - 75B) \cos(dx + c)^4 + 3(163A - 75B) \cos(dx + c)^3 + 3(163A - 75B) \cos(dx + c)^2 + (163A - 75B) \cos(dx + c) + 96(a^3 d \cos(dx + c) + a^3 d))}{(a + a \cos(c + dx))^{5/2}}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output -1/96*(3*sqrt(2)*((163*A - 75*B)*cos(d*x + c)^4 + 3*(163*A - 75*B)*cos(d*x
+ c)^3 + 3*(163*A - 75*B)*cos(d*x + c)^2 + (163*A - 75*B)*cos(d*x + c))*s
qrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)
*sin(d*x + c))) + 2*((299*A - 147*B)*cos(d*x + c)^3 + (503*A - 255*B)*cos(
d*x + c)^2 + 32*(5*A - 3*B)*cos(d*x + c) - 32*A)*sqrt(a*cos(d*x + c) + a)*
sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^4 + 3*a^3*d*cos(d*x +
c)^3 + 3*a^3*d*cos(d*x + c)^2 + a^3*d*cos(d*x + c))
```

3.537. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.537.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.537.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.537.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(a \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^(5/2), x)`

3.537.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(5/2), x)`

3.538 $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.538.1 Optimal result	4883
3.538.2 Mathematica [C] (warning: unable to verify)	4884
3.538.3 Rubi [A] (verified)	4885
3.538.4 Maple [B] (verified)	4889
3.538.5 Fricas [A] (verification not implemented)	4890
3.538.6 Sympy [F(-1)]	4890
3.538.7 Maxima [F(-1)]	4890
3.538.8 Giac [F]	4891
3.538.9 Mupad [F(-1)]	4891

3.538.1 Optimal result

Integrand size = 35, antiderivative size = 223

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{(75A - 19B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{16\sqrt{2}a^{5/2}d}$$

$$- \frac{(A - B)\sqrt{\sec(c + dx)} \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}} - \frac{(13A - 5B)\sqrt{\sec(c + dx)} \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}}$$

$$+ \frac{(49A - 9B)\sqrt{\sec(c + dx)} \sin(c + dx)}{16a^2d\sqrt{a + a \cos(c + dx)}}$$

output `-1/4*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(5/2)-1/16*(13*A-5*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(3/2)-1/32*(75*A-19*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(5/2)/d*2^(1/2)+1/16*(49*A-9*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(1/2)`

3.538.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.78 (sec) , antiderivative size = 732, normalized size of antiderivative = 3.28

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{B \cos^5\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^4\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} \left(11 - 31 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 18 \sin^4\left(\frac{c}{2} + \frac{dx}{2}\right) - \dots\right)}{4d(a(1 + \cos(c + dx)))^{5/2}}$$

$$+ \frac{2A \cos^5\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^4\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)^{3/2} \left(\frac{8 \cos^6\left(\frac{1}{2}(c + dx)\right) {}_4F_3\left(2, 2, 2, \frac{5}{2}; 1, 1, \frac{11}{2}; \frac{\sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{315(-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right))}\right)}{}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^(5/2), x]`

output `-1/4*(B*Cos[c/2 + (d*x)/2]^5*Sec[(c + d*x)/2]^4*Sin[c/2 + (d*x)/2]*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*(11 - 31*Sin[c/2 + (d*x)/2]^2 + 18*Sin[c/2 + (d*x)/2]^4 - (19*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])*Cos[(c + d*x)/2]^4)/Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])/(d*(a*(1 + Cos[c + d*x]))^(5/2)) + (2*A*Cos[c/2 + (d*x)/2]^5*Sec[(c + d*x)/2]^4*Sin[c/2 + (d*x)/2]*((1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1))^(3/2)*((8*Cos[(c + d*x)/2]^6*HypergeometricPFQ[{2, 2, 2, 5/2}, {1, 1, 11/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2)/(315*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) + (Csc[c/2 + (d*x)/2]^8*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-15*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)])*Cos[(c + d*x)/2]^4*(-343 + 1465*Sin[c/2 + (d*x)/2]^2 - 2021*Sin[c/2 + (d*x)/2]^4 + 824*Sin[c/2 + (d*x)/2]^6) + Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-5145 + 33980*Sin[c/2 + (d*x)/2]^2 - 87764*Sin[c/2 + (d*x)/2]^4 + 109737*Sin[c/2 + (d*x)/2]^6 - 66122*Sin[c/2 + (d*x)/2]^8 + 15344*Sin[c/2 + (d*x)/2]^10))/120))/(d*(a*(1 + Cos[c + d*x]))^(5/2))`

3.538. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.538.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{5/2}} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx$$

$$\downarrow \text{3457}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\frac{a(9A-B)-4a(A-B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\frac{a(9A-B)-4a(A-B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\frac{a(9A-B)-4a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \right)$$

$$\downarrow \text{3457}$$

3.538. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(49A-9B)-2a^2(13A-5B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(49A-9B)-2a^2(13A-5B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(49A-9B)-2a^2(13A-5B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int \frac{a^3(75A-19B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} + \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a^2(75A-19B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 3042

3.538. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - a^2(75A-19B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \right)$$

3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^3(75A-19B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{4a^2} + \frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}}{8a^2}$$

218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^2(49A-9B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{\sqrt{2}a^{3/2}(75A-19B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{4a^2}}{8a^2} - \frac{a(13A-5B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^(5/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/4*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a*(13*A - 5*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)) + (-((Sqrt[2]*a^(3/2)*(75*A - 19*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d) + (2*a^2*(49*A - 9*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2))/(8*a^2)`

3.538. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.538.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.538.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 484 vs. 2(188) = 376.

Time = 11.08 (sec) , antiderivative size = 485, normalized size of antiderivative = 2.17

method	result
default	$\frac{(75A(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))-19B(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))+49\sqrt{2}(\cos^2(dx+c))\sin(dx+c)+225\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\sin(dx+c))}{A(75(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))+49\sqrt{2}(\cos^2(dx+c))\sin(dx+c)+225\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\sin(dx+c))}$
parts	

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 1/32/a^3/d*(75*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot
(d*x+c)-csc(d*x+c))-19*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*ar
csin(cot(d*x+c)-csc(d*x+c))+49*A*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)+225*A*cos
(d*x+c)^2*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-
9*B*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)-57*B*cos(d*x+c)^2*arcsin(cot(d*x+c)-cs
c(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+85*A*sin(d*x+c)*cos(d*x+c)*2^(
1/2)+225*A*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-
csc(d*x+c))-13*B*2^(1/2)*cos(d*x+c)*sin(d*x+c)-57*B*cos(d*x+c)*(cos(d*x+c)
/(1+cos(d*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))+32*A*2^(1/2)*sin(d*x+
c)+75*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*A*arcsin(cot(d*x+c)-csc(d*x+c))-19
*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*B*arcsin(cot(d*x+c)-csc(d*x+c))*sec(d*
x+c)^(3/2)*(a*(1+cos(d*x+c)))^(1/2)*cos(d*x+c)/(1+cos(d*x+c))^3*2^(1/2)
```

3.538. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.538.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.94

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sqrt{2}((75A - 19B) \cos(dx + c)^3 + 3(75A - 19B) \cos(dx + c)^2 +$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/32*(sqrt(2)*((75*A - 19*B)*cos(d*x + c)^3 + 3*(75*A - 19*B)*cos(d*x + c)^2 + 3*(75*A - 19*B)*cos(d*x + c) + 75*A - 19*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((49*A - 9*B)*cos(d*x + c)^2 + (85*A - 13*B)*cos(d*x + c) + 32*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.538.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.538.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.538. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{5/2}} dx$

3.538.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(5/2), x)`

3.538.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(5/2), x)`

3.539
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{5/2}} dx$$

3.539.1 Optimal result	4892
3.539.2 Mathematica [A] (verified)	4892
3.539.3 Rubi [A] (verified)	4893
3.539.4 Maple [B] (verified)	4896
3.539.5 Fracas [A] (verification not implemented)	4897
3.539.6 Sympy [F(-1)]	4897
3.539.7 Maxima [F]	4898
3.539.8 Giac [F(-1)]	4898
3.539.9 Mupad [F(-1)]	4898

3.539.1 Optimal result

Integrand size = 35, antiderivative size = 176

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx = \frac{(19A + 5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{16\sqrt{2}a^{5/2}d} - \frac{(A - B) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2}\sqrt{\sec(c + dx)}} - \frac{(9A - B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2}\sqrt{\sec(c + dx)}}$$

output

```
-1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2)-1/16*(9*A-B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+1/32*(19*A+5*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(5/2)/d*2^(1/2)
```

3.539.2 Mathematica [A] (verified)

Time = 1.89 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.92

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx = \frac{\sec^2\left(\frac{1}{2}(c + dx)\right) \left(4(19A + 5B)\operatorname{arctanh}\left(\sqrt{-\sec(c + dx)} \sin^2\left(\frac{1}{2}(c + dx)\right)\right)\right)}{(a + a \cos(c + dx))^{5/2}}$$

input

```
Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^(5/2),x]
```

output $(\text{Sec}[(c + d*x)/2]^2*(4*(19*A + 5*B)*\text{ArcTanh}[\text{Sqrt}[-(\text{Sec}[c + d*x]*\text{Sin}[(c + d*x)/2]^2)])*\text{Cos}[(c + d*x)/2]^4 + \text{Cos}[c + d*x]*(-13*A + 5*B + (-9*A + B)*\text{Cos}[c + d*x])*\text{Sqrt}[2 - 2*\text{Sec}[c + d*x]])*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Tan}[(c + d*x)/2]) / (32*\text{Sqrt}[2]*a^2*d*\text{Sqrt}[a*(1 + \text{Cos}[c + d*x])]*\text{Sqrt}[1 - \text{Sec}[c + d*x]])$

3.539.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{5/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{5/2}} dx$$

↓ 3440

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\frac{a(7A+B)-2a(A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\frac{a(7A+B)-2a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

3.539. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(7A+B)-2a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \\
& \downarrow 3457 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(19A+5B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a(9A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 27 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4}(19A+5B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{8a^2} - \frac{a(9A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4}(19A+5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{8a^2} - \frac{a(9A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 3261 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(19A+5B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2d} - \frac{a(9A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 218 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(19A+5B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{2\sqrt{2}\sqrt{a}d} - \frac{a(9A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right)
\end{aligned}$$

3.539. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^{5/2}} dx$

```
input Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^(5/2),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/4*((A - B)*Sqrt[Cos[c + d*x]]*Si
n[c + d*x])/(d*(a + a*Cos[c + d*x])^(5/2)) + (((19*A + 5*B)*ArcTan[(Sqrt[a
]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/(2
*Sqrt[2]*Sqrt[a]*d) - (a*(9*A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(
a + a*Cos[c + d*x])^(3/2)))/(8*a^2))
```

3.539.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e
_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.539.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. 2(147) = 294.

Time = 11.46 (sec) , antiderivative size = 367, normalized size of antiderivative = 2.09

method	result
default	$\sqrt{-\frac{\csc^2(dx+c)(1-\cos(dx+c))^2+1}{\csc^2(dx+c)(1-\cos(dx+c))^2-1}} \left((\csc^2(dx+c)(1-\cos(dx+c))^2-1) \sqrt{\frac{a}{\csc^2(dx+c)(1-\cos(dx+c))^2+1}} \left(2(\csc^3(dx+c))A\sqrt{-(\csc^2(dx+c)(1-\cos(dx+c))^2-1)} \right) \right)$
parts	$-\frac{A\left(9\sqrt{2}\cos(dx+c)\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+19\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))+13\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+38\right)}{32d(1+\cos(dx+c))^3}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output 1/32/a^3/d*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c)
)^2-1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(a/(csc(d*x+c)^2*(1-cos(d*
x+c))^2+1))^(1/2)*(2*csc(d*x+c)^3*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/
2)*(1-cos(d*x+c))^3-2*csc(d*x+c)^3*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1
/2)*(1-cos(d*x+c))^3+11*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*
x+c)-cot(d*x+c))-3*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-
cot(d*x+c))+19*A*arcsin(cot(d*x+c)-csc(d*x+c))+5*B*arcsin(cot(d*x+c)-csc(d
*x+c)))/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*2^(1/2)
```

$$3.539. \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{5/2}} dx$$

3.539.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.18

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx =$$

$$\frac{\sqrt{2}((19A + 5B) \cos(dx + c)^3 + 3(19A + 5B) \cos(dx + c)^2 + 3(19A + 5B) \cos(dx + c) + 19A + 5B)}{32(a^3 d \cos(dx + c))^3 + 3a^3 d \cos(dx + c)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(5/2),x, algo rithm="fricas")`

output `-1/32*(sqrt(2)*((19*A + 5*B)*cos(d*x + c)^3 + 3*(19*A + 5*B)*cos(d*x + c)^2 + 3*(19*A + 5*B)*cos(d*x + c) + 19*A + 5*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((9*A - B)*cos(d*x + c)^2 + (13*A - 5*B)*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)`

3.539.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.539.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^(5/2), x)`

3.539.8 Giac [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.539.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + a \cos(c + dx))^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(5/2), x)`

3.539. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{5/2}} dx$

$$3.540 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$$

3.540.1 Optimal result	4899
3.540.2 Mathematica [A] (verified)	4899
3.540.3 Rubi [A] (verified)	4900
3.540.4 Maple [B] (verified)	4903
3.540.5 Fricas [A] (verification not implemented)	4904
3.540.6 Sympy [F(-1)]	4904
3.540.7 Maxima [F]	4905
3.540.8 Giac [F(-1)]	4905
3.540.9 Mupad [F(-1)]	4905

3.540.1 Optimal result

Integrand size = 35, antiderivative size = 174

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \frac{(5A + 3B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{a}}{16\sqrt{2}a^{5/2}d}$$

$$+ \frac{(A - B) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} + \frac{(A + 7B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

```
output 1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2)+1/16*(A+7*B
)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+1/32*(5*A+3*B)*ar
ctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2
))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(5/2)/d*2^(1/2)
```

3.540.2 Mathematica [A] (verified)

Time = 3.68 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.05

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \frac{6B \arcsin\left(\frac{\sin(\frac{1}{2}(c+dx))}{\sqrt{\cos^2(\frac{1}{2}(c+dx))}}\right) \cos^3(\frac{1}{2}(c+dx))}{\sqrt{\cos(c+dx)}} - 5A \operatorname{Arctanh}\left(\sqrt{-\sec(c + dx)}\right)$$

```
input Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(5/2)*Sqrt[Sec[c + d*
x]]),x]
```

3.540. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$

```
output ((6*B*ArcSin[Sin[(c + d*x)/2]/Sqrt[Cos[(c + d*x)/2]^2]]*Cos[(c + d*x)/2]^3
)/Sqrt[Cos[c + d*x]] - 5*A*ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2
)]*Cos[(c + d*x)/2]^2*Cot[(c + d*x)/2]*Sqrt[2 - 2*Sec[c + d*x]] + (5*A + 3
*B + (A + 7*B)*Cos[c + d*x])*Tan[(c + d*x)/2]/(16*a*d*(a*(1 + Cos[c + d*x
]))^(3/2)*Sqrt[Sec[c + d*x]])
```

3.540.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a \cos(c + dx) + a)^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})}(a \sin(c + dx + \frac{\pi}{2}) + a)^{5/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{5/2}} dx$$

↓ 3042

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2}} dx$$

↓ 3456

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{\int \frac{a(A-B)+2a(A+3B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{\int \frac{a(A-B)+2a(A+3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

3.540. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+2a(A+3B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \\
& \downarrow 3457 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(5A+3B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a(A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 27 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4}(5A+3B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{8a^2} + \frac{a(A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4}(5A+3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{8a^2} + \frac{a(A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 3261 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{a(A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{a(5A+3B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2d}}{8a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right) \\
& \downarrow 218 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(5A+3B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{2\sqrt{2}\sqrt{ad}} + \frac{a(A+7B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)} \right)
\end{aligned}$$

3.540. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{5/2}\sqrt{\sec(c+dx)}} dx$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(5/2)*Sqrt[Sec[c + d*x]]),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (((5*A + 3*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*Sqrt[a]*d) + (a*(A + 7*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2)`

3.540.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.540.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. 2(145) = 290.

Time = 8.02 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.95

method	result
default	$-\left(-A\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\cos(dx+c)\sin(dx+c)-7B\cos(dx+c)\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+5A\arcsin(\cot(dx+c)-\csc(dx+c))\right)(\cos^2(dx+c)-10\arcsin(\cot(dx+c)-\csc(dx+c)))\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+32d(1+\cos(dx+c))^3\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}$
parts	$\frac{A\left(\sqrt{2}\cos(dx+c)\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+5\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-5\arcsin(\cot(dx+c)-\csc(dx+c))\right)(\cos^2(dx+c)-10\arcsin(\cot(dx+c)-\csc(dx+c)))\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}{32d(1+\cos(dx+c))^3\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2)/sec(d*x+c)^(1/2), x, method=_RET
URNVERBOSE)
```

$$3.540. \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$$

output
$$\begin{aligned} & -1/32/a^3/d*(-A*2^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\cos(d*x+c)*\sin(d \\ & *x+c)-7*B*\cos(d*x+c)*\sin(d*x+c)*2^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}+ \\ & 5*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c)^2-5*A*2^{(1/2)}*(\cos(d*x+c)/(1+ \\ & \cos(d*x+c)))^{(1/2)}*\sin(d*x+c)+3*B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+c) \\ & ^2-3*B*2^{(1/2)}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{(1/2)}*\sin(d*x+c)+10*A*\arcsin(\cot \\ & (d*x+c)-\csc(d*x+c))*\cos(d*x+c)+6*B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*\cos(d*x+ \\ & c)+5*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))+3*B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*(a \\ & *(1+\cos(d*x+c)))^{(1/2)}/(1+\cos(d*x+c))^{-3}/\sec(d*x+c)^{(1/2)}/(\cos(d*x+c)/(1+\cos \\ & (d*x+c)))^{(1/2)}*2^{(1/2)} \end{aligned}$$

3.540.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.18

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \frac{\sqrt{2}((5A + 3B) \cos(dx + c)^3 + 3(5A + 3B) \cos(dx + c)^2 + 3(5A + 3B) \cos(dx + c) + 5A + 3B) \sqrt{a}}{32(a^3 d \cos(dx + c)^3 + 3a^3 d \cos(dx + c)^2}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algo
rithm="fricas")`

output
$$\begin{aligned} & -1/32*(\text{sqrt}(2))*((5*A + 3*B)*\cos(d*x + c)^3 + 3*(5*A + 3*B)*\cos(d*x + c)^2 \\ & + 3*(5*A + 3*B)*\cos(d*x + c) + 5*A + 3*B)*\text{sqrt}(a)*\arctan(\text{sqrt}(2)*\text{sqrt}(a*\cos \\ & (d*x + c) + a)*\text{sqrt}(\cos(d*x + c))/(\text{sqrt}(a)*\sin(d*x + c))) - 2*((A + 7*B)* \\ & \cos(d*x + c)^2 + (5*A + 3*B)*\cos(d*x + c))*\text{sqrt}(a*\cos(d*x + c) + a)*\sin(d* \\ & x + c)/\text{sqrt}(\cos(d*x + c)))/(a^3*d*\cos(d*x + c)^3 + 3*a^3*d*\cos(d*x + c)^2 \\ & + 3*a^3*d*\cos(d*x + c) + a^3*d) \end{aligned}$$

3.540.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2)/sec(d*x+c)**(1/2),x)`

3.540.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$$

output Timed out

3.540.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{5/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*sqrt(sec(d*x + c))), x)`

3.540.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="giac")`

output Timed out

3.540.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(5/2)), x)`

3.540. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$

3.541
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.541.1 Optimal result 4906
 3.541.2 Mathematica [C] (verified) 4907
 3.541.3 Rubi [A] (verified) 4907
 3.541.4 Maple [B] (warning: unable to verify) 4912
 3.541.5 Fricas [A] (verification not implemented) 4912
 3.541.6 Sympy [F(-1)] 4913
 3.541.7 Maxima [F] 4913
 3.541.8 Giac [F(-1)] 4914
 3.541.9 Mupad [F(-1)] 4914

3.541.1 Optimal result

Integrand size = 35, antiderivative size = 234

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{a^{5/2}d}$$

$$+ \frac{(3A - 43B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{16\sqrt{2}a^{5/2}d}$$

$$+ \frac{(A - B) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} + \frac{(3A - 11B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

```
output 1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2)+1/16*(3*A-1
1*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+2*B*arcsin(sin
(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/
a^(5/2)/d+1/32*(3*A-43*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)
^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(5/2)/d
*2^(1/2)
```

3.541.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.85 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.13

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{3/2}(c + dx)} dx = \frac{\cos^5\left(\frac{1}{2}(c + dx)\right) \left(-i\sqrt{2}e^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}}\sqrt{1+e^{2i(c+dx)}}\right)}{(32)}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(3/2)),x]`

output `(Cos[(c + d*x)/2]^5*(((-I)*Sqrt[2]*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*(32*B*ArcSinh[E^(I*(c + d*x))]] - Sqrt[2]*(3*A - 43*B)*ArcTanh[(1 - E^(I*(c + d*x))]/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]]]) - 32*B*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]]))/E^((I/2)*(c + d*x)) + ((3*A - 11*B + (7*A - 15*B)*Cos[c + d*x])*Sec[(c + d*x)/2]^4*Sqrt[Sec[c + d*x]]*(-Sin[(c + d*x)/2] + Sin[(3*(c + d*x))/2]))/(2))/(8*d*(a*(1 + Cos[c + d*x]))^(5/2))`

3.541.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.97, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3456, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{\sec^{3/2}(c + dx)(a \cos(c + dx) + a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a \sin\left(c + dx + \frac{\pi}{2}\right) + a)^{5/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{3/2}(c + dx)(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{5/2}} dx \end{aligned}$$

3.541. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{3/2}(c+dx)} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx$$

↓ 3456

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+8aB \cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3a(A-B)+8aB \cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(3a(A-B)+8aB \sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 3456

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(3A-11B)a^2+32B \cos(c+dx)a^2}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a(3A-11B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(3A-11B)a^2+32B \cos(c+dx)a^2}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a(3A-11B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{(A-B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

3.541. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(3A-11B)a^2+32B\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a(3A-11B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)} \right)$$

↓ 3461

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(3A-43B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + 32aB\int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{4a^2} + \frac{a(3A-11B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(3A-43B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + 32aB\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a^2} + \frac{a(3A-11B)\sin(c+dx)}{2d(a\cos(c+dx)+a)} \right)$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(3A-43B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{64aB\int \frac{1}{\sqrt{1-\frac{a\sin^2(c+dx)}{\cos(c+dx)a+a}}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)a+a}}\right)}{4a^2}}{8a^2} + \frac{a(3A-11B)\sin(c+dx)}{2d(a\cos(c+dx)+a)} \right)$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(3A-43B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{64a^{3/2}B\arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{4a^2} + \frac{a(3A-11B)\sin(c+dx)}{2d(a\cos(c+dx)+a)} \right)$$

↓ 3261

3.541. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{5/2}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{64a^{3/2}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{2a^3(3A-43B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} dx \left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{4a^2}}{8a^2} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{\sqrt{2}a^{3/2}(3A-43B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d} + \frac{64a^{3/2}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d}}{8a^2} + \frac{a(3A-11B)}{2d(a\cos(c+dx)+a)} \right)$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(3/2)),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + (((64*a^(3/2)*B*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (Sqrt[2]*a^(3/2)*(3*A - 43*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d)/(4*a^2) + (a*(3*A - 11*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2))`

3.541.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

3.541. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{3}{2}}(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.541.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^2(c+dx)} dx$$

3.541.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 468 vs. $2(195) = 390$.

Time = 8.39 (sec) , antiderivative size = 469, normalized size of antiderivative = 2.00

method	result
default	$\frac{\left(\left(\csc^2(dx+c)\right)\left(1-\cos(dx+c)\right)^2+1\right)^2\left(\left(\csc^2(dx+c)\right)\left(1-\cos(dx+c)\right)^2-1\right)\sqrt{\frac{a}{\left(\csc^2(dx+c)\right)\left(1-\cos(dx+c)\right)^2+1}}\left(-2\left(\csc^3(dx+c)\right)A\sqrt{\frac{a}{\left(\csc^2(dx+c)\right)\left(1-\cos(dx+c)\right)^2+1}}\right)}{\dots}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))}\left(7\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+3\tan(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-3\arcsin(\cot(dx+c)-\csc(dx+c))\cos(dx+c)-6\right)}{32d(1+\cos(dx+c))^3\sec(dx+c)^{\frac{3}{2}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}a^3}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2)/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/32/a^3/d/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)^(3/2)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(5/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^2*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(a/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(-2*csc(d*x+c)^3*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^3+2*csc(d*x+c)^3*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^3+5*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+c))-32*B^2^(1/2)*arctan(2^(1/2)*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(csc(d*x+c)-cot(d*x+c))))-13*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+c))-3*A*arcsin(cot(d*x+c)-csc(d*x+c))+43*B*arcsin(cot(d*x+c)-csc(d*x+c)))*2^(1/2)`

3.541.5 Fracas [A] (verification not implemented)

Time = 6.03 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.18

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{\sqrt{2}((3A - 43B) \cos(dx + c)^3 + 3(3A - 43B) \cos(dx + c)^2 + 3(3A - 43B) \cos(dx + c) + 3A - 43B)}{\dots}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x,algorithm="fricas")`

3.541.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

output
$$\begin{aligned} & -1/32*(\text{sqrt}(2)*((3*A - 43*B)*\cos(d*x + c)^3 + 3*(3*A - 43*B)*\cos(d*x + c)^2 \\ & + 3*(3*A - 43*B)*\cos(d*x + c) + 3*A - 43*B)*\text{sqrt}(a)*\arctan(\text{sqrt}(2)*\text{sqrt}(\\ & a*\cos(d*x + c) + a)*\text{sqrt}(\cos(d*x + c))/(\text{sqrt}(a)*\sin(d*x + c))) + 64*(B*\cos \\ & (d*x + c)^3 + 3*B*\cos(d*x + c)^2 + 3*B*\cos(d*x + c) + B)*\text{sqrt}(a)*\arctan(\text{sq} \\ & \text{rt}(a*\cos(d*x + c) + a)*\text{sqrt}(\cos(d*x + c))/(\text{sqrt}(a)*\sin(d*x + c))) - 2*((7* \\ & A - 15*B)*\cos(d*x + c)^2 + (3*A - 11*B)*\cos(d*x + c))*\text{sqrt}(a*\cos(d*x + c) \\ & + a)*\sin(d*x + c)/\text{sqrt}(\cos(d*x + c)))/(a^3*d*\cos(d*x + c)^3 + 3*a^3*d*\cos(\\ & d*x + c)^2 + 3*a^3*d*\cos(d*x + c) + a^3*d) \end{aligned}$$

3.541.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2)/sec(d*x+c)**(3/2),x)`

output Timed out

3.541.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{5/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(3
/2)), x)`

3.541.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^3(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algo
rithm="giac")
```

```
output Timed out
```

3.541.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^3(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + a \cos(c + dx))^{5/2}} dx$$

```
input int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(5/2
)),x)
```

```
output int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(5/2
)), x)
```

$$3.542 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$$

3.542.1 Optimal result 4915
 3.542.2 Mathematica [C] (verified) 4916
 3.542.3 Rubi [A] (verified) 4917
 3.542.4 Maple [B] (verified) 4923
 3.542.5 Fricas [A] (verification not implemented) 4924
 3.542.6 Sympy [F(-1)] 4925
 3.542.7 Maxima [F] 4925
 3.542.8 Giac [F(-1)] 4925
 3.542.9 Mupad [F(-1)] 4926

3.542.1 Optimal result

Integrand size = 35, antiderivative size = 286

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx = \frac{(2A - 5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{a^{5/2} d}$$

$$- \frac{(43A - 115B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{16\sqrt{2} a^{5/2} d}$$

$$+ \frac{(A - B) \sin(c + dx)}{4d(a + a \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} + \frac{(7A - 15B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)}$$

$$- \frac{(11A - 35B) \sin(c + dx)}{16a^2 d \sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}}$$

output

```
1/4*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2)+1/16*(7*A-1
5*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2)-1/16*(11*A-35*
B)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+(2*A-5*B)*arcs
in(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(
1/2)/a^(5/2)/d-1/32*(43*A-115*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/co
s(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a
^(5/2)/d*2^(1/2)
```

3.542.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.90 (sec) , antiderivative size = 929, normalized size of antiderivative = 3.25

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{11iAe^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) \cos^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{4d(a(1+\cos(c+dx)))^{5/2}}$$

$$+ \frac{35iBe^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) \cos^5\left(\frac{c}{2} + \frac{dx}{2}\right)}{4d(a(1+\cos(c+dx)))^{5/2}}$$

$$+ \frac{4i\sqrt{2}Ae^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \left(-\operatorname{arcsinh}(e^{i(c+dx)}) + \sqrt{2}\operatorname{arctanh}\left(\frac{-1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) + \operatorname{arctanh}\left(\frac{1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right)\right)}{d(a(1+\cos(c+dx)))^{5/2}}$$

$$- \frac{10i\sqrt{2}Be^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \left(-\operatorname{arcsinh}(e^{i(c+dx)}) + \sqrt{2}\operatorname{arctanh}\left(\frac{-1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) + \operatorname{arctanh}\left(\frac{1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right)\right)}{d(a(1+\cos(c+dx)))^{5/2}}$$

$$+ \frac{\cos^5\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c+dx)} \left(\frac{15(-A+B) \cos\left(\frac{dx}{2}\right) \sin\left(\frac{c}{2}\right)}{2d} + \frac{4B \cos\left(\frac{3dx}{2}\right) \sin\left(\frac{3c}{2}\right)}{d} - \frac{15(A-B) \cos\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right)}{2d} + \frac{\sec\left(\frac{c}{2}\right) \sec^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{d}\right)}{d}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(5/2)),x]`

output

```

(((−11*I)/4)*A*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^
((2*I)*(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)
)*(c + d*x)])]*Cos[c/2 + (d*x)/2]^5)/(d*E^((I/2)*(c + d*x))*(a*(1 + Cos[c
+ d*x]))^(5/2)) + (((35*I)/4)*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d
*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*ArcTanh[(1 - E^(I*(c + d*x)))/(Sqrt[2
]*Sqrt[1 + E^((2*I)*(c + d*x)))]]*Cos[c/2 + (d*x)/2]^5)/(d*E^((I/2)*(c + d
*x))*(a*(1 + Cos[c + d*x]))^(5/2)) + ((4*I)*Sqrt[2]*A*Sqrt[E^(I*(c + d*x))
/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*(-ArcSinh[E^(I*(
c + d*x))] + Sqrt[2]*ArcTanh[(-1 + E^(I*(c + d*x)))/(Sqrt[2]*Sqrt[1 + E^((
2*I)*(c + d*x))]]) + ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]])*Cos[c/2 + (d*
x)/2]^5)/(d*E^((I/2)*(c + d*x))*(a*(1 + Cos[c + d*x]))^(5/2)) - ((10*I)*Sq
rt[2]*B*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*
(c + d*x))]*(-ArcSinh[E^(I*(c + d*x))] + Sqrt[2]*ArcTanh[(-1 + E^(I*(c + d
*x)))/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]]) + ArcTanh[Sqrt[1 + E^((2*I)
*(c + d*x))]])*Cos[c/2 + (d*x)/2]^5)/(d*E^((I/2)*(c + d*x))*(a*(1 + Cos[c
+ d*x]))^(5/2)) + (Cos[c/2 + (d*x)/2]^5*Sqrt[Sec[c + d*x]]*((15*(-A + B)*C
os[(d*x)/2]*Sin[c/2])/(2*d) + (4*B*Cos[(3*d*x)/2]*Sin[(3*c)/2])/d - (15*(A
- B)*Cos[c/2]*Sin[(d*x)/2])/(2*d) + (Sec[c/2]*Sec[c/2 + (d*x)/2]^2*(19*A*
Sin[(d*x)/2] - 27*B*Sin[(d*x)/2]))/(4*d) + (Sec[c/2]*Sec[c/2 + (d*x)/2]^4*
(-A*Sin[(d*x)/2] + B*Sin[(d*x)/2]))/(2*d) + (4*B*Cos[(3*c)/2]*Sin[(3*...

```

3.542.3 Rubi [A] (verified)

Time = 1.83 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.99, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3456, 27, 3042, 3462, 25, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{5/2}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{5/2}} dx
 \end{aligned}$$

3.542. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} (A+B \sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right) a+a)^{5/2}} dx$$

↓ 3456

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{3/2}(c+dx)(5a(A-B)-2a(A-5B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{3/2}(c+dx)(5a(A-B)-2a(A-5B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} (5a(A-B)-2a(A-5B)\sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right) a+a)^{3/2}} dx}{8a^2} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3456

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3a^2(7A-15B)-2a^2(11A-35B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}a+a} dx}{8a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{3/2}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(3a^2(7A-15B)-2a^2(11A-35B)\cos(c+dx))}{\sqrt{\cos(c+dx)}a+a} dx}{8a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{3/2}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{(A-B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

3.542. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{5/2}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} (3a^2(7A-15B)-2a^2(11A-35B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx}{4a^2} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 3462

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int -\frac{a^3(11A-35B)-16a^3(2A-5B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx}{4a^2} - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(-\frac{\int \frac{a^3(11A-35B)-16a^3(2A-5B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx}{4a^2} - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^3(11A-35B)-16a^3(2A-5B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx}{4a^2} - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} + \frac{a(7A-15B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2}$$

↓ 3461

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^3(43A-115B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx - 16a^2(2A-5B)\int \frac{\sqrt{\cos(c+dx)}a+a}{\sqrt{\cos(c+dx)}} dx}{4a^2} - \frac{2a^2(11A-35B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right) \frac{1}{8a^2}$$

↓ 3042

3.542. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{5/2}\sec^{\frac{5}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^3(43A-115B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx - 16a^2(2A-5B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{a} \right) \frac{dx}{4a^2} - \frac{2a^2(11A-35B)}{d\sqrt{a}} \frac{dx}{8a^2}$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^3(43A-115B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx + \frac{32a^2(2A-5B) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)}a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}a+a}\right)}{d}}{a} \right) \frac{dx}{4a^2} - \frac{2a^2(11A-35B)}{d\sqrt{a}} \frac{dx}{8a^2}$$

↓ 223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^3(43A-115B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx - \frac{32a^{5/2}(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a} \cos(c+dx)+a}\right)}{d}}{a} \right) \frac{dx}{4a^2} - \frac{2a^2(11A-35B)}{d\sqrt{a}} \frac{dx}{8a^2}$$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^4(43A-115B) \int \frac{1}{\frac{\sin(c+dx) \tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a}\right) + \frac{32a^{5/2}(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a} \cos(c+dx)+a}\right)}{d}}{a} \right) \frac{dx}{4a^2} - \frac{2a^2(11A-35B)}{d\sqrt{a}} \frac{dx}{8a^2}$$

↓ 218

3.542. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^2(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{-\frac{\sqrt{2}a^{5/2}(43A-115B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{32a^{5/2}(2A-5B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{a} - \frac{2a^2(1}{4a^2} - \frac{8a^2}{8a^2} \right)$$

```
input Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(5/2)),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Cos[c + d*x]^(5/2)*Sin[c +
d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + ((a*(7*A - 15*B)*Cos[c + d*x]^(3
/2)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)) + (-((( -32*a^(5/2)*(2*A
- 5*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]])]/d + (Sqrt
[2]*a^(5/2)*(43*A - 115*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos
[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/a) - (2*a^2*(11*A - 35*B)*Sqrt[C
os[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2)/(8*a^2))
```

3.542.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 223 Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

3.542. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^2(c+dx)} dx$

rule 3253 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3461 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3462 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*S
in[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.542.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 596 vs. 2(241) = 482.

Time = 20.99 (sec) , antiderivative size = 597, normalized size of antiderivative = 2.09

method	result
default	$\frac{\sqrt{a(1+\cos(dx+c))} \left(16B\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c) + 32A\sqrt{2} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) - 15 \tan(dx+c)A\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))} \left(32\sqrt{2} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) - 15 \tan(dx+c)\sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 64 \sec(dx+c)\sqrt{2} \arctan\left(\tan(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right) \right)}{\dots}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(5/2)/sec(d*x+c)^(5/2), x, method=_RET
URNVERBOSE)
```

3.542. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$

```
output 1/32/a^3/d*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^3/sec(d*x+c)^(5/2)/(cos
(d*x+c)/(1+cos(d*x+c)))^(1/2)*(16*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1
/2)*sin(d*x+c)+32*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^
(1/2))-15*tan(d*x+c)*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-80*B*2^(1
/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+55*tan(d*x+c)*B*2
^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+64*sec(d*x+c)*A*2^(1/2)*arctan(ta
n(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))-11*tan(d*x+c)*sec(d*x+c)*A*2^(
1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+43*A*arcsin(cot(d*x+c)-csc(d*x+c))-
160*sec(d*x+c)*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2))+35*tan(d*x+c)*sec(d*x+c)*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-1
15*B*arcsin(cot(d*x+c)-csc(d*x+c))+32*sec(d*x+c)^2*A*2^(1/2)*arctan(tan(d*
x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+86*sec(d*x+c)*A*arcsin(cot(d*x+c)-
csc(d*x+c))-80*sec(d*x+c)^2*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos
(d*x+c)))^(1/2))-230*sec(d*x+c)*B*arcsin(cot(d*x+c)-csc(d*x+c))+43*sec(d*x
+c)^2*A*arcsin(cot(d*x+c)-csc(d*x+c))-115*sec(d*x+c)^2*B*arcsin(cot(d*x+c)
-csc(d*x+c))*2^(1/2)
```

3.542.5 Fracas [A] (verification not implemented)

Time = 7.92 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.09

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx = \frac{\sqrt{2}((43A - 115B) \cos(dx + c))^3 + 3(43A - 115B) \cos(dx + c)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algo
rithm="fricas")
```

```
output 1/32*(sqrt(2))*((43*A - 115*B)*cos(d*x + c)^3 + 3*(43*A - 115*B)*cos(d*x +
c)^2 + 3*(43*A - 115*B)*cos(d*x + c) + 43*A - 115*B)*sqrt(a)*arctan(sqrt(2
)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) - 32
*((2*A - 5*B)*cos(d*x + c)^3 + 3*(2*A - 5*B)*cos(d*x + c)^2 + 3*(2*A - 5*B
)*cos(d*x + c) + 2*A - 5*B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(c
os(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*(16*B*cos(d*x + c)^3 - 5*(3*A - 1
1*B)*cos(d*x + c)^2 - (11*A - 35*B)*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)
*sin(d*x + c)/sqrt(cos(d*x + c))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x
+ c)^2 + 3*a^3*d*cos(d*x + c) + a^3*d)
```

3.542. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$

3.542.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(5/2)/sec(d*x+c)**(5/2),x)`output `Timed out`**3.542.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{5/2} \sec(dx + c)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algo
rithm="maxima")`output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(5
/2)), x)`**3.542.8 Giac [F(-1)]**

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algo
rithm="giac")`output `Timed out`

3.542.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (a + a \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(5/2)), x)`

3.543
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

3.543.1 Optimal result 4927
 3.543.2 Mathematica [C] (warning: unable to verify) 4928
 3.543.3 Rubi [A] (verified) 4928
 3.543.4 Maple [B] (verified) 4935
 3.543.5 Fricas [A] (verification not implemented) 4936
 3.543.6 Sympy [F(-1)] 4937
 3.543.7 Maxima [F(-1)] 4937
 3.543.8 Giac [F] 4938
 3.543.9 Mupad [F(-1)] 4938

3.543.1 Optimal result

Integrand size = 35, antiderivative size = 317

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \frac{(1015A - 363B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}}{64\sqrt{2}a^{7/2}d} - \frac{(1887A - 691B)\sqrt{\sec(c + dx)} \sin(c + dx)}{192a^3d\sqrt{a + a \cos(c + dx)}} - \frac{(A - B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2}} - \frac{(23A - 11B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{48ad(a + a \cos(c + dx))^{5/2}} - \frac{(109A - 41B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{64a^2d(a + a \cos(c + dx))^{3/2}} + \frac{(579A - 199B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{192a^3d\sqrt{a + a \cos(c + dx)}}$$

```
output -1/6*(A-B)*sec(d*x+c)^(3/2)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)-1/48*(23*A
-11*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)-1/64*(109*A-
41*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)+1/192*(579*
A-199*B)*sec(d*x+c)^(3/2)*sin(d*x+c)/a^3/d/(a+a*cos(d*x+c))^(1/2)+1/128*(1
015*A-363*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*c
os(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d*2^(1/2)-1/19
2*(1887*A-691*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^3/d/(a+a*cos(d*x+c))^(1/2)
```

3.543.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

3.543.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.88 (sec) , antiderivative size = 1260, normalized size of antiderivative = 3.97

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \text{Too large to display}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x])^(7/2),x]`

output `(2*B*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*((1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1))^3/2*((16*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 5/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2)/(3465*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) - (Csc[c/2 + (d*x)/2]^10*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(105*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Cos[(c + d*x)/2]^6*(2187 - 12908*Sin[c/2 + (d*x)/2]^2 + 27986*Sin[c/2 + (d*x)/2]^4 - 26380*Sin[c/2 + (d*x)/2]^6 + 8752*Sin[c/2 + (d*x)/2]^8) + Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-229635 + 2120790*Sin[c/2 + (d*x)/2]^2 - 8267707*Sin[c/2 + (d*x)/2]^4 + 17646926*Sin[c/2 + (d*x)/2]^6 - 22251094*Sin[c/2 + (d*x)/2]^8 + 16548816*Sin[c/2 + (d*x)/2]^10 - 6712984*Sin[c/2 + (d*x)/2]^12 + 1144608*Sin[c/2 + (d*x)/2]^14))/1680))/(d*(a*(1 + Cos[c + d*x]))^(7/2)) + (A*Cot[c/2 + (d*x)/2]^7*Csc[c/2 + (d*x)/2]^4*Sec[(c + d*x)/2]^6*((1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1))^7/2*(-7680*Cos[(c + d*x)/2]^10*HypergeometricPFQ[{2, 2, 2, 2, 7/2}, {1, 1, 1, 1, 15/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14 + 19200*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 7/2}, {1, 1, 1, 1, 15/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^14*(-7 + 6*Sin[c/2 + (d*x)/2]^2) + 143*(1 - 2*Sin[c/2 + (d*x)/2]^2)^3*sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + ...`

3.543.3 Rubi [A] (verified)

Time = 2.03 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.09, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.543. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{7/2}} dx$$

↓ 3042

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{7/2}} dx$$

↓ 3440

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{7/2}} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{7/2}} dx$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a(5A-B)-8a(A-B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} - \frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a(5A-B)-8a(A-B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a(5A-B)-8a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}} \right)$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3(a^2(63A-19B)-2a^2(23A-11B)\cos(c+dx))}{2\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{7/2}} \right)$$

↓ 27

3.543. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \int \frac{a^2(63A-19B)-2a^2(23A-11B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)}{6d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \int \frac{a^2(63A-19B)-2a^2(23A-11B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{a(23A-11B)\sin(c+dx)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} - \frac{(A+B)}{6d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\int \frac{a^3(579A-199B)-4a^3(109A-41B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-11B)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\int \frac{a^3(579A-199B)-4a^3(109A-41B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-11B)}{4d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

3.543. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\int \frac{a^3(579A-199B)-4a^3(109A-41B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)}{8a^2} - \frac{a(23A-1)}{4d\cos^{\frac{3}{2}}(c+dx)} \right) \frac{12a^2}{12a^2}$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{2 \int -\frac{a^4(1887A-691B)-2a^4(579A-199B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} + \frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)}{4a^2} \right) \frac{8a^2}{12a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^4(1887A-691B)-2a^4(579A-199B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{3a} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)^{3/2}} \right)}{4a^2} \right) \frac{8a^2}{12a^2}$$

↓ 3042

3.543. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\int \frac{a^4(1887A-691B)-2a^4(579A-199B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} - \frac{a^2(109A-41B)\sin(c+dx)}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} \right)}{8a^2} \right) \frac{1}{12a^2}$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2\int -\frac{3a^5(1015A-363B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}}{4a^2} - \frac{a^2}{2d\cos^{\frac{3}{2}}(c+dx)(a\cos(c+dx)+a)} \right)}{8a^2} \right) \frac{1}{12a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3a^4(1015A-363B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}}{3a} \right)}{4a^2} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3042

3.543. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 3a^4(1015A-363B)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx)}}}{3a}}{4a^2} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{6a^5(1015A-363B)\int \frac{1}{\sin(c+dx)\tan(c+dx)a^3+2a^2} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)+a}}\right)}{3a}}{4a^2} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^3(579A-199B)\sin(c+dx)}{3d\cos^{\frac{3}{2}}(c+dx)\sqrt{a\cos(c+dx)+a}} - \frac{\frac{2a^4(1887A-691B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{3\sqrt{2}a^{7/2}(1015A-363B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)+a}}\right)}{d}}{3a}}{4a^2} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

3.543. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + a*Cos[c + d*x])^(7/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/6*((A - B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(7/2)) + (-1/4*(a*(23*A - 11*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(5/2)) + (3*(-1/2*(a^2*(109*A - 41*B)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)*(a + a*Cos[c + d*x])^(3/2)) + ((2*a^3*(579*A - 199*B)*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)*Sqrt[a + a*Cos[c + d*x]]) - ((-3*Sqrt[2]*a^(7/2)*(1015*A - 363*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^4*(1887*A - 691*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x])))/(3*a))/(4*a^2))/(8*a^2))/(12*a^2))`

3.543.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

3.543.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

```
rule 3463 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n
+ 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a
+ b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n +
1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && Eq
Q[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m
+ 1/2, 0])
```

3.543.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 680 vs. 2(270) = 540.

Time = 10.72 (sec) , antiderivative size = 681, normalized size of antiderivative = 2.15

method	result
default	$-\frac{\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(3045A \arcsin(\cot(dx+c)-\csc(dx+c))(\cos^6(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-1089B \arcsin(\cot(dx+c)-\csc(dx+c))\right)}{a^{\frac{7}{2}}(a+\cos(dx+c))}$
parts	$-\frac{A\left(\sec^{\frac{5}{2}}(dx+c)\right)\sqrt{a(1+\cos(dx+c))}\left(3045(\cos^6(dx+c)) \arcsin(\cot(dx+c)-\csc(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+12180\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \arcsin(\cot(dx+c)-\csc(dx+c))\right)}{a^{\frac{7}{2}}(a+\cos(dx+c))}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*a)^(7/2), x, method=_RET
URNVERBOSE)
```

$$3.543. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

output

```
-1/384/a^4/d*sec(d*x+c)^(5/2)*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^4*(3
045*A*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^6*(cos(d*x+c)/(1+cos(d*x+c)
))^1/2)-1089*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(d*x+c)^6*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)+1887*A*sin(d*x+c)*cos(d*x+c)^5*2^(1/2)+12180*A*arcsin(c
ot(d*x+c)-csc(d*x+c))*cos(d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-691*B
*sin(d*x+c)*cos(d*x+c)^5*2^(1/2)-4356*B*arcsin(cot(d*x+c)-csc(d*x+c))*cos(
d*x+c)^5*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+5082*A*sin(d*x+c)*cos(d*x+c)^4*
2^(1/2)+18270*A*cos(d*x+c)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)-1874*B*sin(d*x+c)*cos(d*x+c)^4*2^(1/2)-6534*B*cos(d*x+c
)^4*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+4251*A
*sin(d*x+c)*cos(d*x+c)^3*2^(1/2)+12180*A*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d
*x+c)))^(1/2)*arcsin(cot(d*x+c)-csc(d*x+c))-1599*B*sin(d*x+c)*cos(d*x+c)^3
*2^(1/2)-4356*B*cos(d*x+c)^3*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*arcsin(cot(
d*x+c)-csc(d*x+c))+896*A*sin(d*x+c)*cos(d*x+c)^2*2^(1/2)+3045*A*cos(d*x+c)
^2*arcsin(cot(d*x+c)-csc(d*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-384*B*s
in(d*x+c)*cos(d*x+c)^2*2^(1/2)-1089*B*cos(d*x+c)^2*arcsin(cot(d*x+c)-csc(d
*x+c))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-128*A*sin(d*x+c)*cos(d*x+c)*2^(1/
2))*2^(1/2)
```

3.543.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 295, normalized size of antiderivative = 0.93

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx =$$

$$3\sqrt{2}((1015A - 363B) \cos(dx + c)^5 + 4(1015A - 363B) \cos(dx + c)^4 + 6(1015A - 363B) \cos(dx + c)^3 + \dots)$$

input

```
integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="fricas")
```

output `-1/384*(3*sqrt(2)*((1015*A - 363*B)*cos(d*x + c)^5 + 4*(1015*A - 363*B)*cos(d*x + c)^4 + 6*(1015*A - 363*B)*cos(d*x + c)^3 + 4*(1015*A - 363*B)*cos(d*x + c)^2 + (1015*A - 363*B)*cos(d*x + c))*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((1887*A - 691*B)*cos(d*x + c)^4 + 2*(2541*A - 937*B)*cos(d*x + c)^3 + 39*(109*A - 41*B)*cos(d*x + c)^2 + 128*(7*A - 3*B)*cos(d*x + c) - 128*A)*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^4*d*cos(d*x + c)^5 + 4*a^4*d*cos(d*x + c)^4 + 6*a^4*d*cos(d*x + c)^3 + 4*a^4*d*cos(d*x + c)^2 + a^4*d*cos(d*x + c))`

3.543.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+a*cos(d*x+c))**(7/2),x)`

output `Timed out`

3.543.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="maxima")`

output `Timed out`

3.543.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(a \cos(dx + c) + a)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(a*cos(d*x + c) + a)^(7/2), x)`

3.543.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{(a + a \cos(c + dx))^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + a*cos(c + d*x))^(7/2), x)`

3.544
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

3.544.1 Optimal result 4939
 3.544.2 Mathematica [C] (warning: unable to verify) 4940
 3.544.3 Rubi [A] (verified) 4941
 3.544.4 Maple [B] (verified) 4946
 3.544.5 Fricas [A] (verification not implemented) 4946
 3.544.6 Sympy [F(-1)] 4947
 3.544.7 Maxima [F(-1)] 4947
 3.544.8 Giac [F] 4948
 3.544.9 Mupad [F(-1)] 4948

3.544.1 Optimal result

Integrand size = 35, antiderivative size = 270

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx =$$

$$\frac{3(121A - 21B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{64\sqrt{2}a^{7/2}d}$$

$$- \frac{(A - B)\sqrt{\sec(c + dx)} \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2}} - \frac{(19A - 7B)\sqrt{\sec(c + dx)} \sin(c + dx)}{48ad(a + a \cos(c + dx))^{5/2}}$$

$$- \frac{(199A - 43B)\sqrt{\sec(c + dx)} \sin(c + dx)}{192a^2d(a + a \cos(c + dx))^{3/2}}$$

$$+ \frac{(691A - 103B)\sqrt{\sec(c + dx)} \sin(c + dx)}{192a^3d\sqrt{a + a \cos(c + dx)}}$$

```
output -1/6*(A-B)*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(a+a*cos(d*x+c))^(7/2)-1/48*(19*A
-7*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d/(a+a*cos(d*x+c))^(5/2)-1/192*(199*A-
43*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d/(a+a*cos(d*x+c))^(3/2)-3/128*(121*
A-21*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*
x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d*2^(1/2)+1/192*(69
1*A-103*B)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^3/d/(a+a*cos(d*x+c))^(1/2)
```

3.544.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

3.544.2 Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.02 (sec) , antiderivative size = 802, normalized size of antiderivative = 2.97

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx =$$

$$\frac{B \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^6\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} \left(141 - 518 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 575 \sin^4\left(\frac{c}{2} + \frac{dx}{2}\right)\right)}{24d(a(1 + \cos(c + dx)))^{7/2}}$$

$$+ \frac{2A \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^6\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)^{3/2} \left(\frac{16 \cos^8\left(\frac{1}{2}(c + dx)\right) {}_5F_4\left(2, 2, 2, 2, \frac{5}{2}; 1, 1, 1, \frac{13}{2}; \frac{\sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}\right)}{3465(-1 + 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right))}\right)}{d(a(1 + \cos(c + dx)))^{7/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^(7/2),x]`

output `-1/24*(B*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*(141 - 518*Sin[c/2 + (d*x)/2]^2 + 575*Sin[c/2 + (d*x)/2]^4 - 206*Sin[c/2 + (d*x)/2]^6 - (189*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]]*Cos[(c + d*x)/2]^6/Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))]))/(d*(a*(1 + Cos[c + d*x]))^(7/2)) + (2*A*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*((1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1))^(3/2)*((16*Cos[(c + d*x)/2]^8*HypergeometricPFQ[{2, 2, 2, 2, 5/2}, {1, 1, 1, 13/2}, Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*Sin[c/2 + (d*x)/2]^2)/(3465*(-1 + 2*Sin[c/2 + (d*x)/2]^2)) - (Csc[c/2 + (d*x)/2]^10*(1 - 2*Sin[c/2 + (d*x)/2]^2)^2*Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(105*ArcTanh[Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]]*Cos[(c + d*x)/2]^6*(2187 - 12908*Sin[c/2 + (d*x)/2]^2 + 27986*Sin[c/2 + (d*x)/2]^4 - 26380*Sin[c/2 + (d*x)/2]^6 + 8752*Sin[c/2 + (d*x)/2]^8) + Sqrt[Sin[c/2 + (d*x)/2]^2/(-1 + 2*Sin[c/2 + (d*x)/2]^2)]*(-229635 + 2120790*Sin[c/2 + (d*x)/2]^2 - 8267707*Sin[c/2 + (d*x)/2]^4 + 17646926*Sin[c/2 + (d*x)/2]^6 - 22251094*Sin[c/2 + (d*x)/2]^8 + 16548816*Sin[c/2 + (d*x)/2]^10 - 6712984*Sin[c/2 + (d*x)/2]^12 + 1144608*Sin[c/2 + (d*x)/2]^14))/(1680))/(d*(a*(1 + Cos[c + d*x]))^(7/2))`

3.544. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$

3.544.3 Rubi [A] (verified)

Time = 1.66 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3463, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a\cos(c+dx)+a)^{7/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a\sin(c+dx+\frac{\pi}{2})+a)^{7/2}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{7/2}} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{7/2}} dx$$

$$\downarrow \text{3457}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(13A-B)-6a(A-B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} - \frac{(A-B)\sin(c+dx)}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}} \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(13A-B)-6a(A-B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}} \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(13A-B)-6a(A-B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)}{6d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{7/2}} \right)$$

3.544. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\begin{array}{c}
 \downarrow 3457 \\
 \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a^2(41A-5B)-4a^2(19A-7B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)}{6d\sqrt{\cos(c+dx)}} \right) \\
 \downarrow 27 \\
 \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a^2(41A-5B)-4a^2(19A-7B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)}{6d\sqrt{\cos(c+dx)}} \right) \\
 \downarrow 3042 \\
 \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a^2(41A-5B)-4a^2(19A-7B)\sin(c+dx+\frac{\pi}{2})}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)}{6d\sqrt{\cos(c+dx)}} \right) \\
 \downarrow 3457 \\
 \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^3(691A-103B)-2a^3(199A-43B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{2a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \right) \\
 \downarrow 27 \\
 \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^3(691A-103B)-2a^3(199A-43B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{5/2}} \right) \\
 \downarrow 3042
 \end{array}$$

3.544. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^3(691A-103B)-2a^3(199A-43B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3463

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\int -\frac{9a^4(121A-21B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 9a^3(121A-21B)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - 9a^3(121A-21B)\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)a+a}} dx}{4a^2} - \frac{a^2(199A-43B)\sin(c+dx)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} - \frac{a(19A-7B)\sin(c+dx)}{4d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3261

3.544. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{18a^4(121A-21B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{d} + \frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} \right) \frac{4a^2}{8a^2} \frac{8a^2}{12a^2}$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^3(691A-103B)\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}} - \frac{9\sqrt{2}a^{5/2}(121A-21B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{4a^2} - \frac{a^2(199A-43B)}{2d\sqrt{\cos(c+dx)}(a\cos(c+dx)+a)} \right) \frac{8a^2}{12a^2}$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + a*Cos[c + d*x])^(7/2),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/6*((A - B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(7/2)) + (-1/4*(a*(19*A - 7*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(5/2)) + (-1/2*(a^2*(199*A - 43*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*(a + a*Cos[c + d*x])^(3/2)) + ((-9*Sqrt[2]*a^(5/2)*(121*A - 21*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/d + (2*a^3*(691*A - 103*B)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]]))/(4*a^2))/(8*a^2))/(12*a^2)
```

3.544.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

3.544. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3463 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*(a*d*m + b*c*(n + 1)) - B*(a*c*m + b*d*(n + 1)) + b*(B*c - A*d)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[n, -1] && (IntegerQ[n] || EqQ[m + 1/2, 0])`

3.544.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 642 vs. $2(229) = 458$.

Time = 10.80 (sec) , antiderivative size = 643, normalized size of antiderivative = 2.38

method	result
default	$-\frac{\left(-\frac{\csc^2(dx+c)(1-\cos(dx+c))^2+1}{\csc^2(dx+c)(1-\cos(dx+c))^2-1}\right)^{\frac{3}{2}}\sqrt{-(\csc^2(dx+c)(1-\cos(dx+c))^2+1)}\sqrt{\frac{a}{\csc^2(dx+c)(1-\cos(dx+c))^2+1}}\left(8(\csc^7(dx+c))A\sqrt{\dots}\right)$
parts	$\frac{A\left(1089\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^4(dx+c))+691\sqrt{2}(\cos^3(dx+c))\sin(dx+c)+4356(\cos^3(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\right)}{\dots}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*a)^(7/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/384/a^4/d*(-(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)/(\csc(d*x+c)^2*(1-\cos(d*x+c))^2-1))^{\frac{3}{2}}*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(a/(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1))^{\frac{1}{2}}*(8*\csc(d*x+c)^7*A*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(1-\cos(d*x+c))^7-8*\csc(d*x+c)^7*B*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(1-\cos(d*x+c))^7+62*\csc(d*x+c)^5*A*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(1-\cos(d*x+c))^5-38*\csc(d*x+c)^5*B*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(1-\cos(d*x+c))^5+299*\csc(d*x+c)^3*A*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(1-\cos(d*x+c))^3-95*\csc(d*x+c)^3*B*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(1-\cos(d*x+c))^3+1089*\csc(d*x+c)^2*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))*(1-\cos(d*x+c))^2-189*\csc(d*x+c)^2*B*\arcsin(\cot(d*x+c)-\csc(d*x+c))*(1-\cos(d*x+c))^2-1137*A*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(\csc(d*x+c)-\cot(d*x+c))+141*B*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*(\csc(d*x+c)-\cot(d*x+c))-1089*A*\arcsin(\cot(d*x+c)-\csc(d*x+c))+189*B*\arcsin(\cot(d*x+c)-\csc(d*x+c)))/(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)*2^{\frac{1}{2}} \end{aligned}$$

3.544.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.96

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \frac{9\sqrt{2}((121A - 21B) \cos(dx + c)^4 + 4(121A - 21B) \cos(dx + c) \dots)}{\dots}$$

3.544.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+a \cos(c+dx))^{7/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="fricas")`

output `1/384*(9*sqrt(2)*((121*A - 21*B)*cos(d*x + c)^4 + 4*(121*A - 21*B)*cos(d*x
+ c)^3 + 6*(121*A - 21*B)*cos(d*x + c)^2 + 4*(121*A - 21*B)*cos(d*x + c)
+ 121*A - 21*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d
*x + c))/(sqrt(a)*sin(d*x + c))) + 2*((691*A - 103*B)*cos(d*x + c)^3 + 2*(
937*A - 133*B)*cos(d*x + c)^2 + 39*(41*A - 5*B)*cos(d*x + c) + 384*A)*sqrt
(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^4*d*cos(d*x + c)^
4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c)
+ a^4*d)`

3.544.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+a*cos(d*x+c))**(7/2),x)`

output Timed out

3.544.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2),x, algo
rithm="maxima")`

output Timed out

3.544.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(a \cos(dx + c) + a)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(a*cos(d*x + c) + a)^(7/2), x)`

3.544.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{(a + a \cos(c + dx))^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + a*cos(c + d*x))^(7/2), x)`

$$3.545 \quad \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{7/2}} dx$$

3.545.1 Optimal result	4949
3.545.2 Mathematica [A] (verified)	4950
3.545.3 Rubi [A] (verified)	4950
3.545.4 Maple [B] (verified)	4954
3.545.5 Fricas [A] (verification not implemented)	4955
3.545.6 Sympy [F(-1)]	4955
3.545.7 Maxima [F]	4956
3.545.8 Giac [F(-1)]	4956
3.545.9 Mupad [F(-1)]	4956

3.545.1 Optimal result

Integrand size = 35, antiderivative size = 223

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx = \frac{(63A + 13B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{64\sqrt{2}a^{7/2}d} - \frac{(A - B) \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2}\sqrt{\sec(c + dx)}} - \frac{(5A - B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{5/2}\sqrt{\sec(c + dx)}} - \frac{(103A + 5B) \sin(c + dx)}{192a^2d(a + a \cos(c + dx))^{3/2}\sqrt{\sec(c + dx)}}$$

```
output -1/6*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(1/2)-1/16*(5*A-
B)*5sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2)-1/192*(103*A+5*B
)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+1/128*(63*A+13*
B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))
^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d*2^(1/2)
```

3.545.2 Mathematica [A] (verified)

Time = 1.86 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.81

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx = \frac{\sec^4\left(\frac{1}{2}(c + dx)\right) \left(-48(63A + 13B) \operatorname{arctanh}\left(\sqrt{-\sec(c + dx) \sin^2\left(\frac{1}{2}(c + dx)\right)}\right) \cos^6\left(\frac{1}{2}(c + dx)\right) + \cos(c + dx)\right)}{1536\sqrt{2}a^3d\sqrt{\sec(c + dx)}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^(7/2), x]`

output `-1/1536*(Sec[(c + d*x)/2]^4*(-48*(63*A + 13*B)*ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]]*Cos[(c + d*x)/2]^6 + Cos[c + d*x]*(493*A - 73*B + (532*A - 4*B)*Cos[c + d*x] + (103*A + 5*B)*Cos[2*(c + d*x)])*Sqrt[2 - 2*Sec[c + d*x]])*Sqrt[Sec[c + d*x]]*Tan[(c + d*x)/2])/(Sqrt[2]*a^3*d*Sqrt[a*(1 + Cos[c + d*x])])*Sqrt[1 - Sec[c + d*x]])`

3.545.3 Rubi [A] (verified)Time = 1.30 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3457, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{(a \cos(c + dx) + a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\csc(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{(a \sin(c + dx + \frac{\pi}{2}) + a)^{7/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(\cos(c + dx)a + a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.545. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{7/2}} dx$

$$\begin{aligned}
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)\left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{7/2}} dx \\
& \quad \downarrow \text{3457} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A+B)-4a(A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A+B)-4a(A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(11A+B)-4a(A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)\left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{5/2}} dx}{12a^2} - \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \quad \downarrow \text{3457} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(73A+11B)-6a^2(5A-B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)\sin(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(73A+11B)-6a^2(5A-B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)\sin(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a^2(73A+11B)-6a^2(5A-B)\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)\left(\sin\left(c+dx+\frac{\pi}{2}\right)a+a\right)^{3/2}} dx}{8a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} - \frac{(A-B)\sin(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right)
\end{aligned}$$

3.545. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a\cos(c+dx))^{7/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a^3(63A+13B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} \right)$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3}{4}a(63A+13B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3}{4}a(63A+13B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3a^2(63A+13B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} dx - \frac{d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{2d} - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} \right)$$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3\sqrt{a}(63A+13B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right) - \frac{a^2(103A+5B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} - \frac{3a(5A-B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} \right)$$

↓ 218

```
input Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + a*Cos[c + d*x])^(7/2), x
]
```

3.545. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{7/2}} dx$

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/6*((A - B)*Sqrt[Cos[c + d*x]]*Si
n[c + d*x])/(d*(a + a*cos[c + d*x])^(7/2)) + ((-3*a*(5*A - B)*Sqrt[Cos[c +
d*x]]*Sin[c + d*x])/(4*d*(a + a*cos[c + d*x])^(5/2)) + ((3*Sqrt[a]*(63*A
+ 13*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a +
a*cos[c + d*x]])])/(2*Sqrt[2]*d) - (a^2*(103*A + 5*B)*Sqrt[Cos[c + d*x]]*
Sin[c + d*x])/(2*d*(a + a*cos[c + d*x])^(3/2)))/(8*a^2)/(12*a^2))
```

3.545.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e
_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*S
in[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c +
d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```



```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.545.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 464 vs. 2(188) = 376.

Time = 11.72 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.09

method	result
default	$-\frac{\sqrt{-\frac{(\csc^2(dx+c)(1-\cos(dx+c))^2+1}{(\csc^2(dx+c)(1-\cos(dx+c))^2-1)} \left((\csc^2(dx+c)(1-\cos(dx+c))^2-1 \right) \sqrt{\frac{a}{(\csc^2(dx+c)(1-\cos(dx+c))^2+1}} \left(-8(\csc^5(dx+c))A\sqrt{-(c} \right.}{}}$
parts	$-\frac{A\left(103\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\left(\cos^2(dx+c)\sin(dx+c)+266\sqrt{2}\cos(dx+c)\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+189\arcsin(\cot(dx+c)-\csc(dx+c))\right)}{}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*a)^(7/2),x,method=_RET
URNVERBOSE)
```

```
output -1/384/a^4/d*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+
c))^2-1))^1/2*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(a/(csc(d*x+c)^2*(1-cos(
d*x+c))^2+1))^1/2*(-8*csc(d*x+c)^5*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^
1/2*(1-cos(d*x+c))^5+8*csc(d*x+c)^5*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)
^1/2*(1-cos(d*x+c))^5-46*csc(d*x+c)^3*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+
1)^1/2*(1-cos(d*x+c))^3+22*csc(d*x+c)^3*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^
2+1)^1/2*(1-cos(d*x+c))^3-141*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^1/2)
*(csc(d*x+c)-cot(d*x+c))+9*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^1/2*(csc
(d*x+c)-cot(d*x+c))-189*A*arcsin(cot(d*x+c)-csc(d*x+c))-39*B*arcsin(cot(d*
x+c)-csc(d*x+c)))/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^1/2)*2^1/2
```

$$3.545. \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{7/2}} dx$$

3.545.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.15

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx =$$

$$\frac{3\sqrt{2}((63A + 13B) \cos(dx + c)^4 + 4(63A + 13B) \cos(dx + c)^3 + 6(63A + 13B) \cos(dx + c)^2 + 4(63A + 13B) \cos(dx + c) + 3A + 13B) \sqrt{a} \arctan(\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}) / (\sqrt{a} \sin(dx + c)) + 2((103A + 5B) \cos(dx + c)^3 + 2(133A - B) \cos(dx + c)^2 + 39(5A - B) \cos(dx + c)) \sqrt{a \cos(dx + c) + a} \sin(dx + c) / \sqrt{\cos(dx + c)}}{384(a^4 d \cos(dx + c)^4 + 4a^4 d \cos(dx + c)^3 + 6a^4 d \cos(dx + c)^2 + 4a^4 d \cos(dx + c) + a^4 d)}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(7/2),x, algo rithm="fricas")`

output `-1/384*(3*sqrt(2)*((63*A + 13*B)*cos(d*x + c)^4 + 4*(63*A + 13*B)*cos(d*x + c)^3 + 6*(63*A + 13*B)*cos(d*x + c)^2 + 4*(63*A + 13*B)*cos(d*x + c) + 3*A + 13*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c))) + 2*((103*A + 5*B)*cos(d*x + c)^3 + 2*(133*A - B)*cos(d*x + c)^2 + 39*(5*A - B)*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)`

3.545.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+a*cos(d*x+c))**(7/2),x)`

output `Timed out`

3.545.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(a \cos(dx + c) + a)^{7/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(a*cos(d*x + c) + a)^(7/2), x)`

3.545.8 Giac [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(7/2),x, algorithm="giac")`

output `Timed out`

3.545.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + a \cos(c + dx))^{7/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + a \cos(c + dx))^{7/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(7/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + a*cos(c + d*x))^(7/2), x)`

3.545. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+a \cos(c+dx))^{7/2}} dx$

3.546
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sqrt{\sec(c+dx)}} dx$$

3.546.1 Optimal result 4957
 3.546.2 Mathematica [A] (verified) 4958
 3.546.3 Rubi [A] (verified) 4958
 3.546.4 Maple [B] (verified) 4962
 3.546.5 Fricas [A] (verification not implemented) 4963
 3.546.6 Sympy [F(-1)] 4964
 3.546.7 Maxima [F] 4964
 3.546.8 Giac [F(-1)] 4964
 3.546.9 Mupad [F(-1)] 4965

3.546.1 Optimal result

Integrand size = 35, antiderivative size = 221

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx = \frac{(13A + 7B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}}{64\sqrt{2}a^{7/2}d}$$

$$+ \frac{(A - B) \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} + \frac{(A + 3B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}}$$

$$- \frac{(5A - 17B) \sin(c + dx)}{192a^2d(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

output

```
1/6*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(1/2)+1/16*(A+3*B
)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2)-1/192*(5*A-17*B)*
sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+1/128*(13*A+7*B)*
arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1
/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d*2^(1/2)
```

3.546.2 Mathematica [A] (verified)

Time = 1.80 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.81

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx = \frac{\sec^4\left(\frac{1}{2}(c + dx)\right) \left(48(13A + 7B) \operatorname{arctanh}\left(\sqrt{-\sec(c + dx) \sin^2}\right)\right)}{\dots}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sqrt[Sec[c + d*x]]),x]`

output `(Sec[(c + d*x)/2]^4*(48*(13*A + 7*B)*ArcTanh[Sqrt[-(Sec[c + d*x]*Sin[(c + d*x)/2]^2)]]*Cos[(c + d*x)/2]^6 + Cos[c + d*x]*(73*A + 59*B + 4*(A + 35*B)*Cos[c + d*x] + (-5*A + 17*B)*Cos[2*(c + d*x)])*Sqrt[2 - 2*Sec[c + d*x]])*Sqrt[Sec[c + d*x]]*Tan[(c + d*x)/2])/(1536*Sqrt[2]*a^3*d*Sqrt[a*(1 + Cos[c + d*x])]*Sqrt[1 - Sec[c + d*x]])`

3.546.3 Rubi [A] (verified)Time = 1.29 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3457, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a \cos(c + dx) + a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{7/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}(A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\left(\sin\left(c + dx + \frac{\pi}{2}\right)a + a\right)^{7/2}} dx \end{aligned}$$

3.546. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \downarrow \text{3456} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+4a(A+2B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+4a(A+2B)\cos(c+dx)}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(A-B)+4a(A+2B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \downarrow \text{3457} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(11A+B)a^2+6(A+3B)\cos(c+dx)a^2}{2\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(11A+B)a^2+6(A+3B)\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(11A+B)a^2+6(A+3B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}}}{12a^2} + \frac{(A-B)\sin(c+dx)\sqrt{\cos(c+dx)}}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
& \downarrow \text{3457}
\end{aligned}$$

3.546. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{7/2}\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a^3(13A+7B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2} +$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3}{4}a(13A+7B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2} +$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3}{4}a(13A+7B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2} +$$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3a^2(13A+7B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3+2a^2}{\cos(c+dx)a+a}} dx - \frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}}{2d} - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2} +$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3\sqrt{a}(13A+7B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{2\sqrt{2}d} - \frac{a^2(5A-17B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{3a(A+3B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2} +$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sqrt[Sec[c + d*x]]),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Sqrt[Cos[c + d*x]]*Sin[c +
d*x])/(6*d*(a + a*cos[c + d*x])^(7/2)) + ((3*a*(A + 3*B)*Sqrt[Cos[c + d*x
]]*Sin[c + d*x])/(4*d*(a + a*cos[c + d*x])^(5/2)) + ((3*Sqrt[a]*(13*A + 7*
B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Co
s[c + d*x]])])/(2*Sqrt[2]*d) - (a^2*(5*A - 17*B)*Sqrt[Cos[c + d*x]]*Sin[c
+ d*x])/(2*d*(a + a*cos[c + d*x])^(3/2)))/(8*a^2)/(12*a^2))
```

3.546.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3261 Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e
_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c
- b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*S
in[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3440 Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Ssin[e + f*x])^p Int[(a + b*Ssin[e + f*x])^m*((c +
d*Ssin[e + f*x])^n/(g*Ssin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```



```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Ssin[e + f*x])^(m
+ 1)*(c + d*Ssin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3457 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(
n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d))
Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n*Simp[B*(a*c*m + b
*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n +
2)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ
[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)]
&& !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])
```

3.546.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs. 2(186) = 372.

Time = 8.29 (sec) , antiderivative size = 488, normalized size of antiderivative = 2.21

method	result
default	$\frac{\left(\csc^2(dx+c)(1-\cos(dx+c))^2+1\right)\left(\csc^2(dx+c)(1-\cos(dx+c))^2-1\right)\sqrt{\frac{a}{\csc^2(dx+c)(1-\cos(dx+c))^2+1}}\left(8(\csc^5(dx+c))A\sqrt{-(\csc^2(dx+c)(1-\cos(dx+c))^2-1}\right)}{\dots}$
parts	$-\frac{A\left(5\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}(\cos^2(dx+c))\sin(dx+c)-2\sqrt{2}\cos(dx+c)\sin(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+39\arcsin(\cot(dx+c)-\csc(dx+c))(\cos^2(dx+c))\right)}{\dots}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2)/sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

$$3.546. \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sqrt{\sec(c+dx)}} dx$$

output
$$-1/384/a^4/d/(-(\csc(dx+c)^2(1-\cos(dx+c))^{2+1}/(\csc(dx+c)^2(1-\cos(dx+c))^{2-1}))^{1/2}/(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{3/2}*(\csc(dx+c)^2(1-\cos(dx+c))^{2+1}*(\csc(dx+c)^2(1-\cos(dx+c))^{2-1}*(a/(\csc(dx+c)^2(1-\cos(dx+c))^{2+1}))^{1/2}*(8*\csc(dx+c)^5*A*(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{1/2}*(1-\cos(dx+c))^{5-8*\csc(dx+c)^5*B*(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{1/2}*(1-\cos(dx+c))^{5+22*\csc(dx+c)^3*A*(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{1/2}*(1-\cos(dx+c))^{3+2*\csc(dx+c)^3*B*(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{1/2}*(1-\cos(dx+c))^{3+9*A*(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{1/2}*(\csc(dx+c)-\cot(dx+c))+27*B*(-\csc(dx+c)^2(1-\cos(dx+c))^{2+1})^{1/2}*(\csc(dx+c)-\cot(dx+c))-39*A*\arcsin(\cot(dx+c)-\csc(dx+c))-21*B*\arcsin(\cot(dx+c)-\csc(dx+c)))^{1/2}$$

3.546.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.15

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx =$$

$$\frac{3\sqrt{2}((13A + 7B) \cos(dx + c)^4 + 4(13A + 7B) \cos(dx + c)^3 + 6(13A + 7B) \cos(dx + c)^2 + 4(13A + 7B) \cos(dx + c) + 13A + 7B) \sqrt{a} \arctan(\sqrt{2} \sqrt{a \cos(dx + c) + a} \sqrt{\cos(dx + c)}) / (\sqrt{a} \sin(dx + c)) + 2*((5A - 17B) \cos(dx + c)^3 - 2*(A + 35B) \cos(dx + c)^2 - 3*(13A + 7B) \cos(dx + c)) \sqrt{a \cos(dx + c) + a} \sin(dx + c) / \sqrt{\cos(dx + c)}}{(a^4 d \cos(dx + c)^4 + 4a^4 d \cos(dx + c)^3 + 6a^4 d \cos(dx + c)^2 + 4a^4 d \cos(dx + c) + a^4 d)}$$

input `integrate((A+B*cos(dx+c))/(a+a*cos(dx+c))^(7/2)/sec(dx+c)^(1/2),x, algorith="fracas")`

output
$$-1/384*(3*\sqrt{2})*((13*A + 7*B)*\cos(dx + c)^4 + 4*(13*A + 7*B)*\cos(dx + c)^3 + 6*(13*A + 7*B)*\cos(dx + c)^2 + 4*(13*A + 7*B)*\cos(dx + c) + 13*A + 7*B)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)})/(\sqrt{a}*\sin(dx + c)) + 2*((5*A - 17*B)*\cos(dx + c)^3 - 2*(A + 35*B)*\cos(dx + c)^2 - 3*(13*A + 7*B)*\cos(dx + c))*\sqrt{a*\cos(dx + c) + a}*\sin(dx + c)/\sqrt{\cos(dx + c)}}/(a^4*d*\cos(dx + c)^4 + 4*a^4*d*\cos(dx + c)^3 + 6*a^4*d*\cos(dx + c)^2 + 4*a^4*d*\cos(dx + c) + a^4*d)$$

3.546.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sqrt{\sec(c+dx)}} dx$$

3.546.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(7/2)/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.546.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{7/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*sqrt(sec(d*x +
c))), x)`

3.546.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(1/2),x, algo
rithm="giac")`

output `Timed out`

3.546.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + a \cos(c + dx))^{7/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(7/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + a*cos(c + d*x))^(7/2)), x)`

$$3.547 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.547.1 Optimal result	4966
3.547.2 Mathematica [B] (verified)	4967
3.547.3 Rubi [A] (verified)	4967
3.547.4 Maple [B] (verified)	4972
3.547.5 Fricas [A] (verification not implemented)	4972
3.547.6 Sympy [F(-1)]	4973
3.547.7 Maxima [F]	4973
3.547.8 Giac [F(-1)]	4974
3.547.9 Mupad [F(-1)]	4974

3.547.1 Optimal result

Integrand size = 35, antiderivative size = 221

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{(7A + 5B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}}{64\sqrt{2}a^{7/2}d}$$

$$+ \frac{(A - B) \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2} \sec^{\frac{3}{2}}(c + dx)} + \frac{(A - 13B) \sin(c + dx)}{48ad(a + a \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}}$$

$$+ \frac{(17A + 67B) \sin(c + dx)}{192a^2d(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

```
output 1/6*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(3/2)+1/48*(A-13*
B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2)+1/192*(17*A+67*B
)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+1/128*(7*A+5*B
)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(
1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d*2^(1/2)
```

3.547.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 561 vs. $2(221) = 442$.

Time = 6.43 (sec) , antiderivative size = 561, normalized size of antiderivative = 2.54

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{A \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sec^6\left(\frac{1}{2}(c + dx)\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}}}{24d(a(1 + \cos(c + dx)))^{7/2}} \left(15 \arcsin\left(\frac{\sin\left(\frac{c}{2} + \frac{dx}{2}\right)}{\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)}}\right) + \frac{33 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{1 - \sec^2\left(\frac{1}{2}(c + dx)\right)}}{\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)}} \right) + \frac{B \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\frac{1}{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}} \sqrt{1 - 2 \sin^2\left(\frac{c}{2} + \frac{dx}{2}\right)}}{24d(a(1 + \cos(c + dx)))^{7/2}}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sec[c + d*x]^(3/2)),x]`

output `(A*Cos[c/2 + (d*x)/2]^7*Sec[(c + d*x)/2]^6*Sin[c/2 + (d*x)/2]*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*(27 - 106*Sin[c/2 + (d*x)/2]^2 + 121*Sin[c/2 + (d*x)/2]^4 - 34*Sin[c/2 + (d*x)/2]^6 + (21*ArcTanh[Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])*Cos[(c + d*x)/2]^6/Sqrt[-(Sin[c/2 + (d*x)/2]^2/(1 - 2*Sin[c/2 + (d*x)/2]^2))])/(24*d*(a*(1 + Cos[c + d*x]))^(7/2)) + (B*Cos[c/2 + (d*x)/2]^7*Sqrt[(1 - 2*Sin[c/2 + (d*x)/2]^2)^(-1)]*Sqrt[1 - 2*Sin[c/2 + (d*x)/2]^2]*(15*ArcSin[Sin[c/2 + (d*x)/2]/Sqrt[Cos[(c + d*x)/2]^2]] + (33*Sin[c/2 + (d*x)/2]*Sqrt[1 - Sec[(c + d*x)/2]^2*Sin[c/2 + (d*x)/2]^2])/Sqrt[Cos[(c + d*x)/2]^2] - (26*Sin[c/2 + (d*x)/2]^3*Sqrt[1 - Sec[(c + d*x)/2]^2*Sin[c/2 + (d*x)/2]^2])/(Cos[(c + d*x)/2]^2)^(3/2) + (8*Sin[c/2 + (d*x)/2]^5*Sqrt[1 - Sec[(c + d*x)/2]^2*Sin[c/2 + (d*x)/2]^2])/(Cos[(c + d*x)/2]^2)^(5/2)))/(24*d*(a*(1 + Cos[c + d*x]))^(7/2))`

3.547.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3456, 27, 3042, 3457, 27, 3042, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.547. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a \cos(c + dx) + a)^{7/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2}(a \sin(c + dx + \frac{\pi}{2}) + a)^{7/2}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{7/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sin(c + dx + \frac{\pi}{2})^{3/2}(A + B \sin(c + dx + \frac{\pi}{2}))}{(\sin(c + dx + \frac{\pi}{2})a + a)^{7/2}} dx \\
& \quad \downarrow \text{3456} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\cos(c + dx)}(3a(A - B) + 2a(A + 5B)\cos(c + dx))}{2(\cos(c + dx)a + a)^{5/2}} dx}{6a^2} + \frac{(A - B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\cos(c + dx)}(3a(A - B) + 2a(A + 5B)\cos(c + dx))}{(\cos(c + dx)a + a)^{5/2}} dx}{12a^2} + \frac{(A - B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}(3a(A - B) + 2a(A + 5B)\sin(c + dx + \frac{\pi}{2}))}{(\sin(c + dx + \frac{\pi}{2})a + a)^{5/2}} dx}{12a^2} + \frac{(A - B) \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}} \right) \\
& \quad \downarrow \text{3456} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{(A - 13B)a^2 + 18(A + 3B)\cos(c + dx)a^2}{2\sqrt{\cos(c + dx)}(\cos(c + dx)a + a)^{3/2}} dx}{4a^2} + \frac{a(A - 13B) \sin(c + dx) \sqrt{\cos(c + dx)}}{4d(a \cos(c + dx) + a)^{5/2}} + \frac{(A - B) \sin(c + dx)}{6d(a \cos(c + dx) + a)^{7/2}} \right) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.547. $\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{3}{2}}(c + dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(A-13B)a^2+18(A+3B)\cos(c+dx)a^2}{\sqrt{\cos(c+dx)}(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)}{6d(a\cos(c+dx)+a)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(A-13B)a^2+18(A+3B)\sin(c+dx+\frac{\pi}{2})a^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)}{6d(a\cos(c+dx)+a)} \right)$$

↓ 3457

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3a^3(7A+5B)}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3}{4}a(7A+5B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + \frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{3}{4}a(7A+5B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + \frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}}}{8a^2} + \frac{a(A-13B)\sin(c+dx)\sqrt{\cos(c+dx)}}{4d(a\cos(c+dx)+a)^{5/2}} \right)$$

↓ 3261

3.547. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{7/2}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} - \frac{3a^2(7A+5B)\int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d\left(-\frac{a\sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}}\right)}{8a^2}}{12a^2} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{a^2(17A+67B)\sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3\sqrt{a}(7A+5B)\arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{8a^2}}{12a^2} + \frac{a(A-13B)\sin(c+dx)}{4d(a\cos(c+dx)+a)^{3/2}} \right)$$

```
input Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sec[c + d*x]^(3/2)),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(6*d*(a + a*Cos[c + d*x])^(7/2)) + ((a*(A - 13*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + ((3*Sqrt[a]*(7*A + 5*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])])/(2*Sqrt[2]*d) + (a^2*(17*A + 67*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)))/(8*a^2))/(12*a^2))
```

3.547.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.547. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{7/2}\sec^{\frac{3}{2}}(c+dx)} dx$

rule 3261 `Int[1/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3456 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m + b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

rule 3457 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(a*f*(2*m + 1)*(b*c - a*d))), x] + Simp[1/(a*(2*m + 1)*(b*c - a*d)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[B*(a*c*m + b*d*(n + 1)) + A*(b*c*(m + 1) - a*d*(2*m + n + 2)) + d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && !GtQ[n, 0] && IntegerQ[2*m] && (IntegerQ[2*n] || EqQ[c, 0])`

3.547.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.547.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 489 vs. 2(186) = 372.

Time = 8.27 (sec) , antiderivative size = 490, normalized size of antiderivative = 2.22

method	result
default	$\frac{\left((\csc^2(dx+c))(1-\cos(dx+c))^2+1 \right)^2 \left((\csc^2(dx+c))(1-\cos(dx+c))^2-1 \right) \sqrt{\frac{a}{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}} \left(-8(\csc^5(dx+c))A\sqrt{\dots} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))} \left(17\sqrt{2} \cos(dx+c) \sin(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} + 70 \sin(dx+c) \sqrt{2} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 21 \arcsin(\cot(dx+c) - \csc(dx+c)) \right)}{\dots}$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2)/sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output -1/384/a^4/d/(-(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1))^(3/2)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(5/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^2*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(a/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(-8*csc(d*x+c)^5*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^5+8*csc(d*x+c)^5*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^5+2*csc(d*x+c)^3*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^3-26*csc(d*x+c)^3*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(1-cos(d*x+c))^3+27*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+c))+33*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+c))-21*A*arcsin(cot(d*x+c)-csc(d*x+c))-15*B*arcsin(cot(d*x+c)-csc(d*x+c)))^2^(1/2)
```

3.547.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.16

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{3/2}(c + dx)} dx = \frac{3\sqrt{2}((7A + 5B) \cos(dx + c)^4 + 4(7A + 5B) \cos(dx + c)^3 + 6(7A + 5B) \cos(dx + c)^2 + 4(7A + 5B) \cos(dx + c) + 3A) + 384(a^4 d \cos(dx + c)^4 + \dots)}{\dots}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(3/2),x, algo
rithm="fracas")
```

3.547. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{3/2}(c+dx)} dx$

output
$$-1/384*(3*\sqrt{2})*((7*A + 5*B)*\cos(dx + c)^4 + 4*(7*A + 5*B)*\cos(dx + c)^3 + 6*(7*A + 5*B)*\cos(dx + c)^2 + 4*(7*A + 5*B)*\cos(dx + c) + 7*A + 5*B)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)})/(\sqrt{a}*\sin(dx + c))) - 2*((17*A + 67*B)*\cos(dx + c)^3 + 10*(7*A + 5*B)*\cos(dx + c)^2 + 3*(7*A + 5*B)*\cos(dx + c))*\sqrt{a*\cos(dx + c) + a}*\sin(dx + c)/\sqrt{\cos(dx + c)})/(a^4*d*\cos(dx + c)^4 + 4*a^4*d*\cos(dx + c)^3 + 6*a^4*d*\cos(dx + c)^2 + 4*a^4*d*\cos(dx + c) + a^4*d)$$

3.547.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(7/2)/sec(d*x+c)**(3/2),x)`

output Timed out

3.547.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{7/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(3/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*sec(d*x + c)^(3/2)), x)`

3.547.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(3/2),x, algo
rithm="giac")
```

```
output Timed out
```

3.547.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + a \cos(c + dx))^{7/2}} dx$$

```
input int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(7/2
)),x)
```

```
output int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + a*cos(c + d*x))^(7/2
)), x)
```

$$3.548 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$$

3.548.1 Optimal result	4975
3.548.2 Mathematica [C] (verified)	4976
3.548.3 Rubi [A] (verified)	4976
3.548.4 Maple [B] (warning: unable to verify)	4982
3.548.5 Fricas [A] (verification not implemented)	4983
3.548.6 Sympy [F(-1)]	4984
3.548.7 Maxima [F]	4984
3.548.8 Giac [F(-1)]	4985
3.548.9 Mupad [F(-1)]	4985

3.548.1 Optimal result

Integrand size = 35, antiderivative size = 281

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{5}{2}}(c + dx)} dx = \frac{2B \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{a^{7/2}d}$$

$$+ \frac{(5A - 177B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{64\sqrt{2}a^{7/2}d}$$

$$+ \frac{(A - B) \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2} \sec^{\frac{5}{2}}(c + dx)} + \frac{(5A - 17B) \sin(c + dx)}{48ad(a + a \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{(5A - 49B) \sin(c + dx)}{64a^2d(a + a \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}$$

output

```
1/6*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(5/2)+1/48*(5*A-17*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2)+1/64*(5*A-49*B)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+2*B*arcsin(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d+1/128*(5*A-177*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^(7/2)/d*2^(1/2)
```

3.548.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.07 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.00

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{5}{2}}(c + dx)} dx = \frac{\cos^7\left(\frac{1}{2}(c + dx)\right) \left(-3i\sqrt{2}e^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}}\sqrt{1+e^{2i(c+dx)}}\right)}{(1 + \dots)}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sec[c + d*x]^(5/2)),x]`

output `(Cos[(c + d*x)/2]^7*(((3*I)*Sqrt[2]*Sqrt[E^(I*(c + d*x))/(1 + E^((2*I)*(c + d*x)))]*Sqrt[1 + E^((2*I)*(c + d*x))]*(128*B*ArcSinh[E^(I*(c + d*x))] - Sqrt[2]*(5*A - 177*B)*ArcTanh[(1 - E^(I*(c + d*x))]/(Sqrt[2]*Sqrt[1 + E^((2*I)*(c + d*x))]]) - 128*B*ArcTanh[Sqrt[1 + E^((2*I)*(c + d*x))]])/E^((I/2)*(c + d*x)) + ((97*A - 541*B + 4*(25*A - 181*B)*Cos[c + d*x] + (67*A - 247*B)*Cos[2*(c + d*x)])*Sec[(c + d*x)/2]^6*Sqrt[Sec[c + d*x]]*(-Sin[(c + d*x)/2] + Sin[(3*(c + d*x))/2]))/8)/(48*d*(a*(1 + Cos[c + d*x]))^(7/2))`

3.548.3 Rubi [A] (verified)

Time = 1.77 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3456, 27, 3042, 3456, 27, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a \cos(c + dx) + a)^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a \sin\left(c + dx + \frac{\pi}{2}\right) + a\right)^{7/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{7/2}} dx \end{aligned}$$

3.548. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} (A+B \sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right) a+a)^{7/2}} dx \\ & \downarrow 3456 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)+12aB \cos(c+dx))}{2(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \right) \\ & \downarrow 27 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{\frac{3}{2}}(c+dx)(5a(A-B)+12aB \cos(c+dx))}{(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \right) \\ & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} (5a(A-B)+12aB \sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right) a+a)^{5/2}} dx}{12a^2} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \right) \\ & \downarrow 3456 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3\sqrt{\cos(c+dx)}((5A-17B)a^2+32B \cos(c+dx)a^2)}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \right) \\ & \downarrow 27 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \int \frac{\sqrt{\cos(c+dx)}((5A-17B)a^2+32B \cos(c+dx)a^2)}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \frac{(A-B) \sin(c+dx) \cos^{\frac{5}{2}}(c+dx)}{6d(a \cos(c+dx)+a)^{7/2}} \right) \\ & \downarrow 3042 \end{aligned}$$

3.548. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} ((5A-17B)a^2+32B \sin(c+dx+\frac{\pi}{2})a^2)}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} + \frac{(A-17B)}{6} \right)$$

↓ 3456

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{\int \frac{(5A-49B)a^3+128B \cos(c+dx)a^3}{2\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{2a^2} + \frac{a^2(5A-49B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{\int \frac{(5A-49B)a^3+128B \cos(c+dx)a^3}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{4a^2} + \frac{a^2(5A-49B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{\int \frac{(5A-49B)a^3+128B \sin(c+dx+\frac{\pi}{2})a^3}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{4a^2} + \frac{a^2(5A-49B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} + \frac{a(5A-17B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{4d(a \cos(c+dx)+a)^{5/2}} \right)$$

↓ 3461

3.548. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx + 128a^2 B \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx + \frac{a^2(5A-49B) \sin(c+dx) \sqrt{\cos(c+dx)}}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} \right)}{12a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx + 128a^2 B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{a^2(5A-49B) \sin(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} \right)}{8a^2} \right)}{12a^2}$$

↓ 3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx - \frac{256a^2 B \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)a+a}}} dx \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)a+a}} \right)}{4a^2} \right)}{8a^2} \right)}{12a^2}$$

↓ 223

3.548. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{a^3(5A-177B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx + \frac{256a^{5/2}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} \right)}{8a^2} + \frac{a^2(5A-49B) \sin(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} \right) \frac{1}{12a^2}$$

3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{256a^{5/2}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{d} - \frac{2a^4(5A-177B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} dx}{4a^2} \right)}{8a^2} - \frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)+a}} \right) \frac{1}{12a^2}$$

218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{a^2(5A-49B) \sin(c+dx)\sqrt{\cos(c+dx)}}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{\sqrt{2}a^{5/2}(5A-177B) \arctan\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a\cos(c+dx)+a}}\right)}{d} \right)}{8a^2} + \frac{256a^{5/2}B \arcsin\left(\frac{\sqrt{a}\sin(c+dx)}{\sqrt{a\cos(c+dx)+a}}\right)}{4a^2} \right) \frac{1}{12a^2}$$

```
input Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sec[c + d*x]^(5/2)),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Cos[c + d*x]^(5/2)*Sin[c +
d*x])/(6*d*(a + a*cos[c + d*x])^(7/2)) + ((a*(5*A - 17*B)*Cos[c + d*x]^(3
/2)*Sin[c + d*x])/(4*d*(a + a*cos[c + d*x])^(5/2)) + (3*(((256*a^(5/2)*B*A
rcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*cos[c + d*x]]])/d + (Sqrt[2]*a^(5/
2)*(5*A - 17*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]
*Sqrt[a + a*cos[c + d*x]]])/d)/(4*a^2) + (a^2*(5*A - 49*B)*Sqrt[Cos[c + d
*x]]*Sin[c + d*x])/(2*d*(a + a*cos[c + d*x])^(3/2))))/(8*a^2))/(12*a^2))
```

3.548.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3456 Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m
+ 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(a_) + (b_.)*sin[(e_.) +
(f_.)*(x_)])]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.548.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 566 vs. 2(236) = 472.

Time = 8.18 (sec) , antiderivative size = 567, normalized size of antiderivative = 2.02

method	result
default	$\frac{\left(\csc^2(dx+c)(1-\cos(dx+c))^2+1\right)^3\left(\csc^2(dx+c)(1-\cos(dx+c))^2-1\right)\sqrt{\frac{a}{\left(\csc^2(dx+c)(1-\cos(dx+c))^2+1\right)}}\left(8(\csc^5(dx+c))A\sqrt{-\right)}{\dots}$
parts	$\frac{A\sqrt{a(1+\cos(dx+c))}\left(67\sin(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}+50\tan(dx+c)\sqrt{2}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}-15\arcsin(\cot(dx+c)-\csc(dx+c))\cos(dx+c)}{\dots}$

3.548.
$$\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$$

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2)/sec(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output -1/384/a^4/d/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+
c))^2-1))^(5/2)/(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(7/2)*(csc(d*x+c)^2*(1-
cos(d*x+c))^2+1)^3*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(a/(csc(d*x+c)^2*(1-
cos(d*x+c))^2+1))^(1/2)*(8*csc(d*x+c)^5*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+
1)^(1/2)*(1-cos(d*x+c))^5-8*csc(d*x+c)^5*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+
1)^(1/2)*(1-cos(d*x+c))^5-26*csc(d*x+c)^3*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^
2+1)^(1/2)*(1-cos(d*x+c))^3+50*csc(d*x+c)^3*B*(-csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1)^(1/2)*(1-cos(d*x+c))^3+33*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2
)*(csc(d*x+c)-cot(d*x+c))-384*B*2^(1/2)*arctan(2^(1/2)*(-csc(d*x+c)^2*(1-c
os(d*x+c))^2+1)^(1/2)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(csc(d*x+c)-cot(d*
x+c)))-189*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*(csc(d*x+c)-cot(d*x+
c))-15*A*arcsin(cot(d*x+c)-csc(d*x+c))+531*B*arcsin(cot(d*x+c)-csc(d*x+c)
)*2^(1/2)
```

3.548.5 Fracas [A] (verification not implemented)

Time = 7.95 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.20

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{3\sqrt{2}((5A - 177B) \cos(dx + c)^4 + 4(5A - 177B) \cos(dx + c)^3 + 6(5A - 177B) \cos(dx + c)^2 + 4(5A - 177B) \cos(dx + c) + 3(5A - 177B))}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{5}{2}}(c + dx)}$$

```
input integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(5/2),x, algo
rithm="fricas")
```

3.548. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{5}{2}}(c+dx)} dx$

output `-1/384*(3*sqrt(2)*((5*A - 177*B)*cos(d*x + c)^4 + 4*(5*A - 177*B)*cos(d*x + c)^3 + 6*(5*A - 177*B)*cos(d*x + c)^2 + 4*(5*A - 177*B)*cos(d*x + c) + 5*A - 177*B)*sqrt(a)*arctan(sqrt(2)*sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c))) + 768*(B*cos(d*x + c)^4 + 4*B*cos(d*x + c)^3 + 6*B*cos(d*x + c)^2 + 4*B*cos(d*x + c) + B)*sqrt(a)*arctan(sqrt(a*cos(d*x + c) + a)*sqrt(cos(d*x + c)))/(sqrt(a)*sin(d*x + c))) - 2*((67*A - 247*B)*cos(d*x + c)^3 + 2*(25*A - 181*B)*cos(d*x + c)^2 + 3*(5*A - 49*B)*cos(d*x + c))*sqrt(a*cos(d*x + c) + a)*sin(d*x + c)/sqrt(cos(d*x + c)))/(a^4*d*cos(d*x + c)^4 + 4*a^4*d*cos(d*x + c)^3 + 6*a^4*d*cos(d*x + c)^2 + 4*a^4*d*cos(d*x + c) + a^4*d)`

3.548.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(7/2)/sec(d*x+c)**(5/2),x)`

output `Timed out`

3.548.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{5/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{7/2} \sec(dx + c)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*sec(d*x + c)^(5/2)), x)`

3.548.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(5/2),x, algorith="giac")`

output `Timed out`

3.548.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{5/2}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (a + a \cos(c + dx))^{7/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(7/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + a*cos(c + d*x))^(7/2)), x)`

$$3.549 \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{7/2}(c+dx)} dx$$

3.549.1 Optimal result	4986
3.549.2 Mathematica [C] (verified)	4987
3.549.3 Rubi [A] (verified)	4988
3.549.4 Maple [B] (verified)	4995
3.549.5 Fricas [A] (verification not implemented)	4996
3.549.6 Sympy [F(-1)]	4997
3.549.7 Maxima [F]	4997
3.549.8 Giac [F(-1)]	4998
3.549.9 Mupad [F(-1)]	4998

3.549.1 Optimal result

Integrand size = 35, antiderivative size = 333

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} dx = \frac{(2A - 7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{a^{7/2} d}$$

$$- \frac{(177A - 637B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2} \sqrt{\cos(c+dx)} \sqrt{a+a \cos(c+dx)}}\right) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{64\sqrt{2}a^{7/2}d}$$

$$+ \frac{(A - B) \sin(c + dx)}{6d(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} + \frac{(3A - 7B) \sin(c + dx)}{16ad(a + a \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)}$$

$$+ \frac{(79A - 259B) \sin(c + dx)}{192a^2d(a + a \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} - \frac{7(7A - 27B) \sin(c + dx)}{64a^3d\sqrt{a + a \cos(c + dx)} \sqrt{\sec(c + dx)}}$$

output

```

1/6*(A-B)*sin(d*x+c)/d/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(7/2)+1/16*(3*A-7
*B)*sin(d*x+c)/a/d/(a+a*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2)+1/192*(79*A-259
*B)*sin(d*x+c)/a^2/d/(a+a*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2)-7/64*(7*A-27*
B)*sin(d*x+c)/a^3/d/(a+a*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+(2*A-7*B)*arcs
in(sin(d*x+c)*a^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(
1/2)/a^(7/2)/d-1/128*(177*A-637*B)*arctan(1/2*sin(d*x+c)*a^(1/2)*2^(1/2)/
cos(d*x+c)^(1/2)/(a+a*cos(d*x+c))^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)
/a^(7/2)/d*2^(1/2)
    
```

3.549. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{7/2}(c+dx)} dx$

3.549.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 8.35 (sec) , antiderivative size = 1017, normalized size of antiderivative = 3.05

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} dx = \\
& \frac{49iAe^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{8d(a(1+\cos(c+dx)))^{7/2}} \\
& + \frac{189iBe^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) \cos^7\left(\frac{c}{2} + \frac{dx}{2}\right)}{8d(a(1+\cos(c+dx)))^{7/2}} \\
& + \frac{8i\sqrt{2}Ae^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \left(-\operatorname{arcsinh}(e^{i(c+dx)}) + \sqrt{2}\operatorname{arctanh}\left(\frac{-1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) + \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right)\right)}{d(a(1+\cos(c+dx)))^{7/2}} \\
& - \frac{28i\sqrt{2}Be^{-\frac{1}{2}i(c+dx)} \sqrt{\frac{e^{i(c+dx)}}{1+e^{2i(c+dx)}}} \sqrt{1+e^{2i(c+dx)}} \left(-\operatorname{arcsinh}(e^{i(c+dx)}) + \sqrt{2}\operatorname{arctanh}\left(\frac{-1+e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right) + \operatorname{arctanh}\left(\frac{1-e^{i(c+dx)}}{\sqrt{2}\sqrt{1+e^{2i(c+dx)}}}\right)\right)}{d(a(1+\cos(c+dx)))^{7/2}} \\
& + \frac{\cos^7\left(\frac{c}{2} + \frac{dx}{2}\right) \sqrt{\sec(c+dx)} \left(\frac{(-247A+427B) \cos\left(\frac{dx}{2}\right) \sin\left(\frac{c}{2}\right)}{12d} + \frac{8B \cos\left(\frac{3dx}{2}\right) \sin\left(\frac{3c}{2}\right)}{d} - \frac{(247A-427B) \cos\left(\frac{c}{2}\right) \sin\left(\frac{dx}{2}\right)}{12d} + \frac{\sec\left(\frac{c}{2}\right)}{d}\right)}{d}
\end{aligned}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sec[c + d*x]^(7/2)),x]`

output $(((-49*I)/8)*A*\text{Sqrt}[E^{(I*(c + d*x))}/(1 + E^{((2*I)*(c + d*x))})*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]]*\text{ArcTanh}[(1 - E^{(I*(c + d*x))})/(\text{Sqrt}[2]*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}])]*\text{Cos}[c/2 + (d*x)/2]^7)/(d*E^{((I/2)*(c + d*x))}*(a*(1 + \text{Cos}[c + d*x]))^{(7/2)} + (((189*I)/8)*B*\text{Sqrt}[E^{(I*(c + d*x))}/(1 + E^{((2*I)*(c + d*x))})*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]]*\text{ArcTanh}[(1 - E^{(I*(c + d*x))})/(\text{Sqrt}[2]*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}])]*\text{Cos}[c/2 + (d*x)/2]^7)/(d*E^{((I/2)*(c + d*x))}*(a*(1 + \text{Cos}[c + d*x]))^{(7/2)} + ((8*I)*\text{Sqrt}[2]*A*\text{Sqrt}[E^{(I*(c + d*x))}/(1 + E^{((2*I)*(c + d*x))})*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]]*(-\text{ArcSinh}[E^{(I*(c + d*x))}] + \text{Sqrt}[2]*\text{ArcTanh}[-1 + E^{(I*(c + d*x))})/(\text{Sqrt}[2]*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]]]) + \text{ArcTanh}[\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]])*\text{Cos}[c/2 + (d*x)/2]^7)/(d*E^{((I/2)*(c + d*x))}*(a*(1 + \text{Cos}[c + d*x]))^{(7/2)} - ((28*I)*\text{Sqrt}[2]*B*\text{Sqrt}[E^{(I*(c + d*x))}/(1 + E^{((2*I)*(c + d*x))})*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]]*(-\text{ArcSinh}[E^{(I*(c + d*x))}] + \text{Sqrt}[2]*\text{ArcTanh}[-1 + E^{(I*(c + d*x))})/(\text{Sqrt}[2]*\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]]]) + \text{ArcTanh}[\text{Sqrt}[1 + E^{((2*I)*(c + d*x))}]])*\text{Cos}[c/2 + (d*x)/2]^7)/(d*E^{((I/2)*(c + d*x))}*(a*(1 + \text{Cos}[c + d*x]))^{(7/2)} + (\text{Cos}[c/2 + (d*x)/2]^7*\text{Sqrt}[\text{Sec}[c + d*x]]*(((-247*A + 427*B)*\text{Cos}[(d*x)/2]*\text{Sin}[c/2])/(12*d) + (8*B*\text{Cos}[(3*d*x)/2]*\text{Sin}[(3*c)/2])/d - ((247*A - 427*B)*\text{Cos}[c/2]*\text{Sin}[(d*x)/2])/(12*d) + (\text{Sec}[c/2]*\text{Sec}[c/2 + (d*x)/2]^2*(379*A*\text{Sin}[(d*x)/2] - 703*B*\text{Sin}[(d*x)/2]))/(24*d) + (\text{Sec}[c/2]*\text{Sec}[c/2 + (d*x)/2]^6*(A*\text{Sin}[(d*x)/2] - B*\text{Sin}[(d*x)/2]))/(3*d) + (\text{Sec}[c/2]*\text{Se...$

3.549.3 Rubi [A] (verified)

Time = 2.24 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.02, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3440, 3042, 3456, 27, 3042, 3456, 27, 3042, 3456, 27, 3042, 3462, 25, 3042, 3461, 3042, 3253, 223, 3261, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{7}{2}}(c + dx)(a \cos(c + dx) + a)^{7/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{7/2} (a \sin(c + dx + \frac{\pi}{2}) + a)^{7/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{(\cos(c + dx)a + a)^{7/2}} dx$$

3.549. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{7/2} (A+B \sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right) a+a)^{7/2}} dx \\
 & \downarrow \text{3456} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{5/2}(c+dx)(7a(A-B)-2a(A-7B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{5/2}} dx}{6a^2} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{5/2}(c+dx)(7a(A-B)-2a(A-7B)\cos(c+dx))}{(\cos(c+dx)a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} (7a(A-B)-2a(A-7B)\sin\left(c+dx+\frac{\pi}{2}\right))}{(\sin\left(c+dx+\frac{\pi}{2}\right) a+a)^{5/2}} dx}{12a^2} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
 & \downarrow \text{3456} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{3/2}(c+dx)(15a^2(3A-7B)-2a^2(17A-77B)\cos(c+dx))}{2(\cos(c+dx)a+a)^{3/2}} dx}{4a^2} + \frac{3a(3A-7B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos^{3/2}(c+dx)(15a^2(3A-7B)-2a^2(17A-77B)\cos(c+dx))}{(\cos(c+dx)a+a)^{3/2}} dx}{8a^2} + \frac{3a(3A-7B)\sin(c+dx)\cos^{5/2}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} + \frac{(A-B)\sin(c+dx)\cos^{7/2}(c+dx)}{6d(a\cos(c+dx)+a)^{7/2}} \right) \\
 & \downarrow \text{3042}
 \end{aligned}$$

3.549. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^{7/2}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (15a^2(3A-7B)-2a^2(17A-77B)\sin(c+dx+\frac{\pi}{2}))}{(\sin(c+dx+\frac{\pi}{2})a+a)^{3/2}} dx}{8a^2} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2}$$

↓ 3456

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3\sqrt{\cos(c+dx)}(a^3(79A-259B)-14a^3(7A-27B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}a+a} dx}{2a^2} + \frac{a^2(79A-259B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \int \frac{\sqrt{\cos(c+dx)}(a^3(79A-259B)-14a^3(7A-27B)\cos(c+dx))}{\sqrt{\cos(c+dx)}a+a} dx}{4a^2} + \frac{a^2(79A-259B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a^3(79A-259B)-14a^3(7A-27B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx}{4a^2} + \frac{a^2(79A-259B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2}$$

↓ 3462

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{\int -\frac{7a^4(7A-27B)-64a^4(2A-7B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)}a+a} dx}{a} - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right)}{4a^2} + \frac{a^2(79A-259B)\sin(c+dx)\cos^{\frac{3}{2}}(c+dx)}{2d(a\cos(c+dx)+a)^{3/2}} + \frac{3a(3A-7B)\sin(c+dx)\cos^{\frac{5}{2}}(c+dx)}{4d(a\cos(c+dx)+a)^{5/2}} \right) \frac{1}{12a^2}$$

3.549. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{7/2}\sec^2(c+dx)} dx$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{\int \frac{7a^4(7A-27B)-64a^4(2A-7B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx}{a} - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right)}{4a^2} + \frac{a^2(79A-259B)\sin(c+dx)}{2d(a\cos(c+dx)+a)} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{\int \frac{7a^4(7A-27B)-64a^4(2A-7B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})a+a}} dx}{a} - \frac{14a^3(7A-27B)\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a\cos(c+dx)+a}} \right)}{4a^2} + \frac{a^2(79A-259B)\sin(c+dx)}{2d(a\cos(c+dx)+a)} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3461

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3 \left(\frac{a^4(177A-637B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} dx - 64a^3(2A-7B) \int \frac{\sqrt{\cos(c+dx)a+a}}{\sqrt{\cos(c+dx)}} dx}{a} - \frac{14a^3(7A-27B)\sin(c+dx)}{d\sqrt{a\cos(c+dx)+a}} \right)}{4a^2} \right) \frac{1}{8a^2} \frac{1}{12a^2}$$

↓ 3042

3.549. $\int \frac{A+B\cos(c+dx)}{(a+a\cos(c+dx))^{7/2} \sec^2(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^4(177A-637B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx - 64a^3(2A-7B) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{4a^2} - \frac{14a^3(7A-3B)}{8a^2} - \frac{12a^3}{12a^2} \right)$$

3253

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^4(177A-637B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx + \frac{128a^3(2A-7B) \int \frac{1}{\sqrt{1-\frac{a \sin^2(c+dx)}{\cos(c+dx)}a+a}} d\left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}}\right)}{4a^2}}{8a^2} - \frac{14a^3}{12a^2} \right)$$

223

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^4(177A-637B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{\sin(c+dx+\frac{\pi}{2})}a+a} dx - \frac{128a^{7/2}(2A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{a \cos(c+dx)+a}}\right)}{d}}{4a^2} - \frac{14a^3}{8a^2} - \frac{12a^3}{12a^2} \right)$$

3.549. $\int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^2(c+dx)} dx$

↓ 3261

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a^5(177A-637B) \int \frac{1}{\frac{\sin(c+dx)\tan(c+dx)a^3}{\cos(c+dx)a+a} + 2a^2} d \left(-\frac{a \sin(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{\cos(c+dx)a+a}} \right) - \frac{128a^{7/2}(2A-7B) \arcsin\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right)}{d} - \frac{128a^{7/2}}{8a^2}}{3} \right)$$

↓ 218

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a^2(79A-259B) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d(a \cos(c+dx)+a)^{3/2}} + \frac{3 \left(\frac{\sqrt{2}a^{7/2}(177A-637B) \arctan\left(\frac{\sqrt{a} \sin(c+dx)}{\sqrt{2}\sqrt{\cos(c+dx)}\sqrt{a \cos(c+dx)+a}}\right) - \frac{128a^{7/2}}{8a^2}}{d} \right)}{8a^2} \right)$$

input `Int[(A + B*Cos[c + d*x])/((a + a*Cos[c + d*x])^(7/2)*Sec[c + d*x]^(7/2)),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((A - B)*Cos[c + d*x]^(7/2)*Sin[c + d*x])/(6*d*(a + a*Cos[c + d*x])^(7/2)) + ((3*a*(3*A - 7*B)*Cos[c + d*x]^(5/2)*Sin[c + d*x])/(4*d*(a + a*Cos[c + d*x])^(5/2)) + ((a^2*(79*A - 259*B)*Cos[c + d*x]^(3/2)*Sin[c + d*x])/(2*d*(a + a*Cos[c + d*x])^(3/2)) + (3*(-((-128*a^(7/2)*(2*A - 7*B)*ArcSin[(Sqrt[a]*Sin[c + d*x])/Sqrt[a + a*Cos[c + d*x]]])/d + (Sqrt[2]*a^(7/2)*(177*A - 637*B)*ArcTan[(Sqrt[a]*Sin[c + d*x])/(Sqrt[2]*Sqrt[Cos[c + d*x]]*Sqrt[a + a*Cos[c + d*x]])]/d)/a) - (14*a^3*(7*A - 27*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + a*Cos[c + d*x]])))/(4*a^2))/(8*a^2))/(12*a^2))`

3.549.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3253 `Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2/f Subst[Int[1/Sqrt[1 - x^2/a], x], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])]], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 - b^2, 0] && EqQ[d, a/b]`
- rule 3261 `Int[1/(Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(a/f) Subst[Int[1/(2*b^2 - (a*c - b*d)*x^2), x], x, b*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`
- rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3456 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(
a*f*(2*m + 1))), x] - Simp[1/(a*b*(2*m + 1)) Int[(a + b*Ssin[e + f*x])^(m
+ 1)*(c + d*Ssin[e + f*x])^(n - 1)*Simp[A*(a*d*n - b*c*(m + 1)) - B*(a*c*m +
b*d*n) - d*(a*B*(m - n) + A*b*(m + n + 1))*Sin[e + f*x], x], x], x] /; Fre
eQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0] && LtQ[m, -2^(-1)] && GtQ[n, 0] && IntegerQ[2*m] && (In
tegerQ[2*n] || EqQ[c, 0])
```

```
rule 3461 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Sim
p[(A*b - a*B)/b Int[1/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]])
, x], x] + Simp[B/b Int[Sqrt[a + b*Ssin[e + f*x]]/Sqrt[c + d*Ssin[e + f*x]]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3462 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^n/(f*(m +
n + 1))), x] + Simp[1/(b*(m + n + 1)) Int[(a + b*Ssin[e + f*x])^m*(c + d*S
sin[e + f*x])^(n - 1)*Simp[A*b*c*(m + n + 1) + B*(a*c*m + b*d*n) + (A*b*d*(m
+ n + 1) + B*(a*d*m + b*c*n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && GtQ[n, 0] && (IntegerQ[n] || EqQ[m + 1/2, 0])
```

3.549.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 810 vs. $2(282) = 564$.

Time = 20.98 (sec) , antiderivative size = 811, normalized size of antiderivative = 2.44

method	result	size
default	Expression too large to display	811
parts	Expression too large to display	843

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*a)^(7/2)/sec(d*x+c)^(7/2),x,method=_RET
URNVERBOSE)
```

$$3.549. \quad \int \frac{A+B \cos(c+dx)}{(a+a \cos(c+dx))^{7/2} \sec^2(c+dx)} dx$$

output

```

-1/384/a^4/d*(a*(1+cos(d*x+c)))^(1/2)/(1+cos(d*x+c))^4/sec(d*x+c)^(7/2)/(c
os(d*x+c)/(1+cos(d*x+c)))^(1/2)*(-192*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c))
)^(1/2)*sin(d*x+c)+247*tan(d*x+c)*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1
/2)-384*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))-109
9*tan(d*x+c)*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+1344*B*2^(1/2)*ar
ctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+362*tan(d*x+c)*sec(d*x+
c)*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-531*A*arcsin(cot(d*x+c)-csc
(d*x+c))-1152*sec(d*x+c)*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*
x+c)))^(1/2))-1442*tan(d*x+c)*sec(d*x+c)*B*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+
c)))^(1/2)+1911*B*arcsin(cot(d*x+c)-csc(d*x+c))+4032*sec(d*x+c)*B*2^(1/2)*
arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+147*tan(d*x+c)*sec(d*
x+c)^2*A*2^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-1593*sec(d*x+c)*A*arcsi
n(cot(d*x+c)-csc(d*x+c))-1152*sec(d*x+c)^2*A*2^(1/2)*arctan(tan(d*x+c)*(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2))-567*tan(d*x+c)*sec(d*x+c)^2*B*2^(1/2)*(cos
(d*x+c)/(1+cos(d*x+c)))^(1/2)+5733*sec(d*x+c)*B*arcsin(cot(d*x+c)-csc(d*x+
c))+4032*sec(d*x+c)^2*B*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c
)))^(1/2))-1593*sec(d*x+c)^2*A*arcsin(cot(d*x+c)-csc(d*x+c))-384*sec(d*x+c
)^3*A*2^(1/2)*arctan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))+5733*se
c(d*x+c)^2*B*arcsin(cot(d*x+c)-csc(d*x+c))+1344*sec(d*x+c)^3*B*2^(1/2)*arc
tan(tan(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))-531*sec(d*x+c)^3*A*ar...

```

3.549.5 Fracas [A] (verification not implemented)

Time = 13.73 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.14

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{7}{2}}(c + dx)} dx = \frac{3\sqrt{2}((177A - 637B) \cos(dx + c)^4 + 4(177A - 637B) \cos(dx + c))}{(a + a \cos(c + dx))^{7/2} \sec^{\frac{7}{2}}(c + dx)}$$

input

```

integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(7/2),x, algo
rithm="fracas")

```

output $1/384*(3*\sqrt{2})*((177*A - 637*B)*\cos(dx + c)^4 + 4*(177*A - 637*B)*\cos(dx + c)^3 + 6*(177*A - 637*B)*\cos(dx + c)^2 + 4*(177*A - 637*B)*\cos(dx + c) + 177*A - 637*B)*\sqrt{a}*\arctan(\sqrt{2}*\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)})/(\sqrt{a}*\sin(dx + c))) - 384*((2*A - 7*B)*\cos(dx + c)^4 + 4*(2*A - 7*B)*\cos(dx + c)^3 + 6*(2*A - 7*B)*\cos(dx + c)^2 + 4*(2*A - 7*B)*\cos(dx + c) + 2*A - 7*B)*\sqrt{a}*\arctan(\sqrt{a*\cos(dx + c) + a}*\sqrt{\cos(dx + c)})/(\sqrt{a}*\sin(dx + c))) + 2*(192*B*\cos(dx + c)^4 - (247*A - 1099*B)*\cos(dx + c)^3 - 2*(181*A - 721*B)*\cos(dx + c)^2 - 21*(7*A - 27*B)*\cos(dx + c))*\sqrt{a*\cos(dx + c) + a}*\sin(dx + c)/\sqrt{\cos(dx + c)})/(a^4*d*\cos(dx + c)^4 + 4*a^4*d*\cos(dx + c)^3 + 6*a^4*d*\cos(dx + c)^2 + 4*a^4*d*\cos(dx + c) + a^4*d)$

3.549.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))**(7/2)/sec(d*x+c)**(7/2),x)`

output Timed out

3.549.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(a \cos(dx + c) + a)^{7/2} \sec(dx + c)^{7/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(7/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((a*cos(d*x + c) + a)^(7/2)*sec(d*x + c)^(7/2)), x)`

3.549.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+a*cos(d*x+c))^(7/2)/sec(d*x+c)^(7/2),x, algorith="giac")`

output `Timed out`

3.549.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + a \cos(c + dx))^{7/2} \sec^{7/2}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{7/2} (a + a \cos(c + dx))^{7/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^(7/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(7/2)*(a + a*cos(c + d*x))^(7/2)), x)`

3.550 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

3.550.1 Optimal result	4999
3.550.2 Mathematica [A] (verified)	5000
3.550.3 Rubi [A] (verified)	5000
3.550.4 Maple [B] (verified)	5004
3.550.5 Fricas [C] (verification not implemented)	5005
3.550.6 Sympy [F(-1)]	5006
3.550.7 Maxima [F]	5006
3.550.8 Giac [F]	5007
3.550.9 Mupad [F(-1)]	5007

3.550.1 Optimal result

Integrand size = 31, antiderivative size = 180

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= -\frac{2(3aA + 5bB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d} \\ & \quad + \frac{2(Ab + aB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d} \\ & \quad + \frac{2(3aA + 5bB) \sqrt{\sec(c + dx)} \sin(c + dx)}{5d} \\ & \quad + \frac{2(Ab + aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3d} + \frac{2aA \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{5d} \end{aligned}$$

```
output 2/3*(A*b+B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/5*a*A*sec(d*x+c)^(5/2)*sin(d
*x+c)/d+2/5*(3*A*a+5*B*b)*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2/5*(3*A*a+5*B*b)*
(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*
c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(A*b+B*a)*(cos(1/2*d*x
+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*
cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.550.2 Mathematica [A] (verified)

Time = 1.75 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.73

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{\sec^{\frac{5}{2}}(c + dx) \left(-12(3aA + 5bB) \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20(Ab + aB) \cos^{\frac{5}{2}}(c + dx) \text{EllipticF}\left(\frac{1}{2}(c + dx) \mid 2\right) \right)}{30d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`output `(Sec[c + d*x]^(5/2)*(-12*(3*a*A + 5*b*B)*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 20*(A*b + a*B)*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 2*(15*(a*A + b*B) + 10*(A*b + a*B)*Cos[c + d*x] + 3*(3*a*A + 5*b*B)*Cos[2*(c + d*x)])*Sin[c + d*x])/(30*d)`**3.550.3 Rubi [A] (verified)**Time = 0.85 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.94, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4485, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)(A \sec(c + dx) + B) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + b\right) \left(A \csc\left(c + dx + \frac{\pi}{2}\right) + B\right) dx$$

$$\downarrow \text{4485}$$

3.550. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\frac{2}{5} \int \frac{1}{2} \sec^{\frac{3}{2}}(c+dx)(3aA+5bB+5(Ab+aB)\sec(c+dx))dx + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 27

$$\frac{1}{5} \int \sec^{\frac{3}{2}}(c+dx)(3aA+5bB+5(Ab+aB)\sec(c+dx))dx + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 3042

$$\frac{1}{5} \int \csc\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(3aA+5bB+5(Ab+aB)\csc\left(c+dx+\frac{\pi}{2}\right)\right) dx + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 4274

$$\frac{1}{5} \left((3aA+5bB) \int \sec^{\frac{3}{2}}(c+dx)dx + 5(aB+Ab) \int \sec^{\frac{5}{2}}(c+dx)dx \right) + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 3042

$$\frac{1}{5} \left((3aA+5bB) \int \csc\left(c+dx+\frac{\pi}{2}\right)^{3/2} dx + 5(aB+Ab) \int \csc\left(c+dx+\frac{\pi}{2}\right)^{5/2} dx \right) + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 4255

$$\frac{1}{5} \left(5(aB+Ab) \left(\frac{1}{3} \int \sqrt{\sec(c+dx)}dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + (3aA+5bB) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} \right) \right) + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(5(aB+Ab) \left(\frac{1}{3} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right) + (3aA+5bB) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} \right) \right) + \frac{2aA \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d}$$

↓ 4258

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right) + (3aA + 5bB) \frac{2aA \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right) + (3aA + 5bB) \frac{2aA \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{5d} \right)$$

↓ 3119

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} \right) + (3aA + 5bB) \frac{2aA \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{5d} \right)$$

↓ 3120

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} + \frac{2 \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} \right) + (3aA + 5bB) \frac{2aA \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{5d} \right)$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*a*A*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(5*d) + ((3*a*A + 5*b*B)*((-2*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d + (2*sqrt[Sec[c + d*x]]*Sin[c + d*x])/d) + 5*(A*b + a*B)*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (2*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d))/5`

3.550.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

3.550.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 635 vs. 2(208) = 416.

Time = 102.97 (sec) , antiderivative size = 636, normalized size of antiderivative = 3.53

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1}(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{\sin(\frac{dx}{2} + \frac{c}{2})^2(2(\sin^2(\frac{dx}{2} + \frac{c}{2})) - 1)} \left(\frac{2Bb\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})} (2(\sin^2(\frac{dx}{2} + \frac{c}{2})) \cos(\frac{dx}{2} + \frac{c}{2}) - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}})}{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \right)$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2), x, method=_RETURNVER
BOSE)
```

3.550. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b/sin(1/2*
d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2)))+2/5*a*A/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*s
in(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2
*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2
*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(
1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+
1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+2*(A*b+B*a)*(-1
/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2
*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2
)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*
x+1/2*c)^2-1)^(1/2)/d

```

3.550.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.31

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx =$$

$$5\sqrt{2}(iBa + iAb) \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5\sqrt{2}(-iBa$$

input

```

integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm=
"fricas")

```

output `-1/15*(5*sqrt(2)*(I*B*a + I*A*b)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-I*B*a - I*A*b)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(3*I*A*a + 5*I*B*b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-3*I*A*a - 5*I*B*b)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*(3*A*a + 5*B*b)*cos(d*x + c)^2 + 3*A*a + 5*(B*a + A*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2)`

3.550.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.550.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.550.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.550.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + b \cos(c + dx)) dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x)), x)`

3.551 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$

3.551.1 Optimal result	5008
3.551.2 Mathematica [A] (verified)	5009
3.551.3 Rubi [A] (verified)	5009
3.551.4 Maple [B] (verified)	5013
3.551.5 Fricas [C] (verification not implemented)	5014
3.551.6 Sympy [F(-1)]	5014
3.551.7 Maxima [F]	5015
3.551.8 Giac [F]	5015
3.551.9 Mupad [F(-1)]	5015

3.551.1 Optimal result

Integrand size = 31, antiderivative size = 143

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= -\frac{2(Ab + aB)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2(aA + 3bB)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2(Ab + aB)\sqrt{\sec(c + dx)}\sin(c + dx)}{d} + \frac{2aA \sec^{\frac{3}{2}}(c + dx)\sin(c + dx)}{3d}$$

```
output 2/3*a*A*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2*(A*b+B*a)*sin(d*x+c)*sec(d*x+c)^(1
/2)/d-2*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Elliptic
E(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(A*a
+3*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*
d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.551.2 Mathematica [A] (verified)

Time = 0.97 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.73

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}\left(-3(Ab + aB)E\left(\frac{1}{2}(c + dx) \mid 2\right) + (aA + 3bB)\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \frac{(aA - 3bB)\text{EllipticE}\left(\frac{1}{2}(c + dx) \mid 2\right)}{2}\right)}{3d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*sqrt[Cos[c + d*x]]*sqrt[Sec[c + d*x]]*(-3*(A*b + a*B)*EllipticE[(c + d*x)/2, 2] + (a*A + 3*b*B)*EllipticF[(c + d*x)/2, 2] + ((a*A + 3*(A*b + a*B)*Cos[c + d*x])*Sin[c + d*x])/Cos[c + d*x]^(3/2)))/(3*d)`

3.551.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.99, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4485, 27, 3042, 4274, 3042, 4255, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)(A \sec(c + dx) + B) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + b\right) \left(A \csc\left(c + dx + \frac{\pi}{2}\right) + B\right) dx$$

$$\downarrow \text{4485}$$

3.551. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\frac{2}{3} \int \frac{1}{2} \sqrt{\sec(c+dx)} (aA + 3bB + 3(Ab + aB) \sec(c+dx)) dx + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 27

$$\frac{1}{3} \int \sqrt{\sec(c+dx)} (aA + 3bB + 3(Ab + aB) \sec(c+dx)) dx + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \left(aA + 3bB + 3(Ab + aB) \csc\left(c+dx+\frac{\pi}{2}\right) \right) dx + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 4274

$$\frac{1}{3} \left(3(aB + Ab) \int \sec^{\frac{3}{2}}(c+dx) dx + (aA + 3bB) \int \sqrt{\sec(c+dx)} dx \right) + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \left((aA + 3bB) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + 3(aB + Ab) \int \csc\left(c+dx+\frac{\pi}{2}\right)^{3/2} dx \right) + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 4255

$$\frac{1}{3} \left((aA + 3bB) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + 3(aB + Ab) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right) \right) + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \left((aA + 3bB) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + 3(aB + Ab) \left(\frac{2 \sin(c+dx) \sqrt{\sec(c+dx)}}{d} - \int \frac{1}{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}} dx \right) \right) + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 4258

$$\frac{1}{3} \left((aA + 3bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3(aB + Ab) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \right) \right) - \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 3042

$$\frac{1}{3} \left((aA + 3bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(aB + Ab) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \right) \right) - \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 3119

$$\frac{1}{3} \left((aA + 3bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + 3(aB + Ab) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \right) \right) - \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2(aA + 3bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + 3(aB + Ab) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \right) \right) - \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d}$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*a*A*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + ((2*(a*A + 3*b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 3*(A*b + a*B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3`

3.551.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.)^(p_))*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4485 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) +
(a_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[
e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc
[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x
], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[
n, -1]
```

3.551.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 400 vs. 2(179) = 358.

Time = 100.54 (sec) , antiderivative size = 401, normalized size of antiderivative = 2.80

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1}(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} \left(\frac{2Bb\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))} + 1 F(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2})}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} + 2aA \left(-\frac{\cos(\frac{dx}{2} + \frac{c}{2})}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} \right) \right)$
parts	$-\frac{2aA \left(-2\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1 F(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2}) (\sin^2(\frac{dx}{2} + \frac{c}{2})) - 2(\sin^2(\frac{dx}{2} + \frac{c}{2})) \cos(\frac{dx}{2} + \frac{c}{2}) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)}{3\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})} (2(\cos^2(\frac{dx}{2} + \frac{c}{2})))$

```
input int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RETURNVER
BOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2
*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+2*
a*A*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*
cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^
2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2*(A*b+B*a)/sin(1/2*d*x+1/
2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2
^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.551. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.551.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.43

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{\sqrt{2}(-i Aa - 3i Bb) \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \sqrt{2}(i Aa + 3i Bb) \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) - 3\sqrt{2}(i B^2 a + i A^2 b) \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 3\sqrt{2}(i B^2 a + i A^2 b) \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2(A^2 a + 3(B^2 a + A^2 b) \cos(dx + c)) \sin(dx + c) / \sqrt{\cos(dx + c)}}{d \cos(dx + c)}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="fricas")`

output `1/3*(sqrt(2)*(-I*A*a - 3*I*B*b)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(I*A*a + 3*I*B*b)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(I*B*a + I*A*b)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(-I*B*a - I*A*b)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(A*a + 3*(B*a + A*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))`

3.551.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.551.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.551.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.551.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + b \cos(c + dx)) dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x)), x)`

3.551. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.552 $\int (a+b \cos(c+dx))(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$

3.552.1 Optimal result	5016
3.552.2 Mathematica [A] (verified)	5017
3.552.3 Rubi [A] (verified)	5017
3.552.4 Maple [A] (verified)	5020
3.552.5 Fricas [C] (verification not implemented)	5021
3.552.6 Sympy [F(-1)]	5021
3.552.7 Maxima [F]	5022
3.552.8 Giac [F]	5022
3.552.9 Mupad [F(-1)]	5022

3.552.1 Optimal result

Integrand size = 31, antiderivative size = 111

$$\begin{aligned} & \int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= -\frac{2(aA - bB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d} \\ & \quad + \frac{2(Ab + aB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{d} \\ & \quad + \frac{2aA \sqrt{\sec(c + dx)} \sin(c + dx)}{d} \end{aligned}$$

output

```
2*a*A*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2*(A*a-B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.552.2 Mathematica [A] (verified)

Time = 1.38 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.77

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{\sec(c + dx)} \left(- \left((aA - bB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \right) + (Ab + aB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx) \mid 2\right) \right) + aA \sin(c + dx)}{d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(2*Sqrt[Sec[c + d*x]]*(-((a*A - b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + (A*b + a*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + a*A*Sin[c + d*x]))/d`

3.552.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3439, 3042, 4485, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right) \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)(A \sec(c + dx) + B)}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \csc\left(c + dx + \frac{\pi}{2}\right) + b)(A \csc\left(c + dx + \frac{\pi}{2}\right) + B)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx$$

$$\downarrow \text{4485}$$

3.552. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\begin{aligned}
& 2 \int -\frac{aA - bB - (Ab + aB) \sec(c + dx)}{2\sqrt{\sec(c + dx)}} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow 27 \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{aA - bB - (Ab + aB) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow 3042 \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{aA - bB + (-Ab - aB) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow 4274 \\
& -(aA - bB) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + (aB + Ab) \int \sqrt{\sec(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow 3042 \\
& -(aA - bB) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + (aB + Ab) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow 4258 \\
& (aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx - (aA - \\
& bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow 3042 \\
& (aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - (aA - \\
& bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow 3119 \\
& (aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx - \\
& \frac{2(aA - bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2}(c + dx) | 2)}{d} + \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \\
& \quad \downarrow 3120
\end{aligned}$$

3.552. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\frac{2(aB + Ab)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{2(aA - bB)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} + \frac{2aA \sin(c + dx)\sqrt{\sec(c + dx)}}{d}$$

input `Int[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(-2*(a*A - b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*(A*b + a*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*A*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d`

3.552.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4485 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

3.552.4 Maple [A] (verified)

Time = 9.32 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.22

method	result
default	$4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a - 2Ab \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - 2A \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}$
parts	$- \frac{2aA \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `2*(2*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a-A*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a-B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.552.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.38

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{2Aa \sin(dx+c)}{\sqrt{\cos(dx+c)}} + \sqrt{2}(-iBa - iAb) \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c)) + \sqrt{2}(iBa + iAb) \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i \sin(dx+c))$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `(2*A*a*sin(d*x + c)/sqrt(cos(d*x + c)) + sqrt(2)*(-I*B*a - I*A*b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(I*B*a + I*A*b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + sqrt(2)*(-I*A*a + I*B*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + sqrt(2)*(I*A*a - I*B*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.552.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.552.7 Maxima [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.552.8 Giac [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.552.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + b \cos(c + dx)) dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x)), x)`

3.552. $\int (a + b \cos(c + dx))(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.553 $\int (a+b \cos(c+dx))(A+B \cos(c+dx))\sqrt{\sec(c+dx)} dx$

3.553.1 Optimal result	5023
3.553.2 Mathematica [A] (verified)	5023
3.553.3 Rubi [A] (verified)	5024
3.553.4 Maple [B] (verified)	5027
3.553.5 Fricas [C] (verification not implemented)	5028
3.553.6 Sympy [F]	5028
3.553.7 Maxima [F]	5029
3.553.8 Giac [F]	5029
3.553.9 Mupad [F(-1)]	5029

3.553.1 Optimal result

Integrand size = 31, antiderivative size = 115

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{2(Ab + aB)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{d} + \frac{2(3aA + bB)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d} + \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}}$$

```
output 2/3*b*B*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(3*A*a+B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.553.2 Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.78

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{\sqrt{\sec(c + dx)}\left(6(Ab + aB)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) + 2(3aA + bB)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\right)}{3d}$$

input `Integrate[(a + b*Cos[c + d*x])*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(6*(A*b + a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 2*(3*a*A + b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + b*B*Sin[2*(c + d*x)]))/(3*d)`

3.553.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 3439, 3042, 4484, 27, 3042, 4274, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c+dx)}(a+b\cos(c+dx))(A+B\cos(c+dx))dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{(a\sec(c+dx)+b)(A\sec(c+dx)+B)}{\sec^{\frac{3}{2}}(c+dx)}dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a\csc\left(c+dx+\frac{\pi}{2}\right)+b)(A\csc\left(c+dx+\frac{\pi}{2}\right)+B)}{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}}dx \\
 & \quad \downarrow \text{4484} \\
 & \frac{2bB\sin(c+dx)}{3d\sqrt{\sec(c+dx)}} - \frac{2}{3} \int \frac{3(Ab+aB) + (3aA+bB)\sec(c+dx)}{2\sqrt{\sec(c+dx)}}dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \int \frac{3(Ab+aB) + (3aA+bB)\sec(c+dx)}{\sqrt{\sec(c+dx)}}dx + \frac{2bB\sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3} \int \frac{3(Ab + aB) + (3aA + bB) \csc\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 4274 \\
& \frac{1}{3} \left(3(aB + Ab) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + (3aA + bB) \int \sqrt{\sec(c + dx)} dx \right) + \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left(3(aB + Ab) \int \frac{1}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx + (3aA + bB) \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx \right) + \\
& \quad \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 4258 \\
& \frac{1}{3} \left((3aA + bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + 3(aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos} \right. \\
& \quad \left. \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left((3aA + bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + 3(aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos} \right. \\
& \quad \left. \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\
& \quad \downarrow 3119 \\
& \frac{1}{3} \left((3aA + bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{6(aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E}{d} \right. \\
& \quad \left. \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\
& \quad \downarrow 3120 \\
& \frac{1}{3} \left(\frac{2(3aA + bB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{6(aB + Ab) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E}{d} \right. \\
& \quad \left. \frac{2bB \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right)
\end{aligned}$$

input `Int[(a + b*cos[c + d*x])*(A + B*cos[c + d*x])*sqrt[sec[c + d*x]],x]`

output `((6*(A*b + a*B)*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d + (2*(3*a*A + b*B)*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/d)/3 + (2*b*B*sin[c + d*x])/(3*d*sqrt[Sec[c + d*x]])`

3.553.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_) + (f_)*(x_)]*(b_) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4484 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e +
f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(
n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]
```

3.553.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 325 vs. 2(153) = 306.

Time = 9.84 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.83

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(4B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + 3aA\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1\right)F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{d}$
parts	$-\frac{2aA\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) + \frac{2(Ab+Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1$

```
input int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RETURNVER
BOSE)
```

```
output -2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*B*cos(1/2*
d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*b+3*a*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*si
n(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*A*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/
2*d*x+1/2*c),2^(1/2))*b-2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b+B*b*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+
1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a)/(-2*sin(1/2*d*x
+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/
2*c)^2-1)^(1/2)/d
```

3.553. $\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$

3.553.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.36

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{2 B b \sqrt{\cos(dx + c)} \sin(dx + c) + \sqrt{2}(-3i A a - i B b) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `1/3*(2*B*b*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(2)*(-3*I*A*a - I*B*b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(3*I*A*a + I*B*b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(-I*B*a - I*A*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(I*B*a + I*A*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d`

3.553.6 Sympy [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx))(a + b \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))*sqrt(sec(c + d*x)), x)`

3.553.7 Maxima [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)\sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.553.8 Giac [F]

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)\sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.553.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx)) dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x)),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x)), x)`

3.553. $\int (a + b \cos(c + dx))(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$

3.554
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.554.1 Optimal result 5030
 3.554.2 Mathematica [A] (verified) 5031
 3.554.3 Rubi [A] (verified) 5031
 3.554.4 Maple [B] (verified) 5035
 3.554.5 Fricas [C] (verification not implemented) 5036
 3.554.6 Sympy [F] 5036
 3.554.7 Maxima [F] 5037
 3.554.8 Giac [F] 5037
 3.554.9 Mupad [F(-1)] 5037

3.554.1 Optimal result

Integrand size = 31, antiderivative size = 148

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{2(5aA + 3bB)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(Ab + aB)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2bB \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2(Ab + aB) \sin(c + dx)}{3d \sqrt{\sec(c + dx)}}$$

```
output 2/5*b*B*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/3*(A*b+B*a)*sin(d*x+c)/d/sec(d*x+c)
)^(1/2)+2/5*(5*A*a+3*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*
EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+
2/3*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(si
n(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.554.2 Mathematica [A] (verified)

Time = 1.00 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.73

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(6(5aA + 3bB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(Ab + aB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx) \mid 2\right) + (5A^2b + 5a^2B + 3b^2B \cos(c + dx)) \sin[2(c + dx)] \right)}{15d}$$

input `Integrate[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(6*(5*a*A + 3*b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 10*(A*b + a*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (5*A*b + 5*a*B + 3*b*B*Cos[c + d*x])*Sin[2*(c + d*x)])/(15*d)`

3.554.3 Rubi [A] (verified)Time = 0.83 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.99, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4484, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)(A \sec(c + dx) + B)}{\sec^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)(A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx$$

$$\begin{aligned}
& \downarrow 4484 \\
& \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} - \frac{2}{5} \int \frac{5(Ab+aB) + (5aA+3bB) \sec(c+dx)}{2 \sec^{\frac{3}{2}}(c+dx)} dx \\
& \downarrow 27 \\
& \frac{1}{5} \int \frac{5(Ab+aB) + (5aA+3bB) \sec(c+dx)}{\sec^{\frac{3}{2}}(c+dx)} dx + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \int \frac{5(Ab+aB) + (5aA+3bB) \csc(c+dx+\frac{\pi}{2})}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \\
& \downarrow 4274 \\
& \frac{1}{5} \left(5(aB+Ab) \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)} dx + (5aA+3bB) \int \frac{1}{\sqrt{\sec(c+dx)}} dx \right) + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(5(aB+Ab) \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx + (5aA+3bB) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \\
& \downarrow 4256 \\
& \frac{1}{5} \left((5aA+3bB) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5(aB+Ab) \left(\frac{1}{3} \int \sqrt{\sec(c+dx)} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) + \\
& \quad \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left((5aA+3bB) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + 5(aB+Ab) \left(\frac{1}{3} \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) + \\
& \quad \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \\
& \downarrow 4258
\end{aligned}$$

3.554. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\frac{1}{5} \left((5aA + 3bB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + 5(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + (5aA + 3bB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx$$

↓ 3119

$$\frac{1}{5} \left(5(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) + \frac{2(5aA + 3bB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d} + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{2(5aA + 3bB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{d} + 5(aB + Ab) \left(\frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} + \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d} \right) + \frac{2bB \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

input `Int[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(2*b*B*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((2*(5*a*A + 3*b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + 5*(A*b + a*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]))/5`

3.554.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.)^(p_))*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4484 Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (
a_))*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e +
f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(
n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x]
/; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]
```

3.554.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 370 vs. 2(180) = 360.

Time = 10.47 (sec) , antiderivative size = 371, normalized size of antiderivative = 2.51

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(-24B\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b + (20Ab + 20Ba + 24Bb)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\cos\left(\frac{dx}{2} + \frac{c}{2}\right)}\right)$
parts	$\frac{2aA\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right) - \frac{2(Ab + Ba)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}}d$

```
input int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNVER
BOSE)
```

```
output -2/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*cos(1
/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b+(20*A*b+20*B*a+24*B*b)*sin(1/2*d*x+1/
2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*b-10*B*a-6*B*b)*sin(1/2*d*x+1/2*c)^2*cos(
1/2*d*x+1/2*c)+5*A*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-
1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-15*A*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/
2))*a+5*B*a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*
EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-9*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*
sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b)/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*co
s(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

$$3.554. \int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.554.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.24

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$\frac{5 \sqrt{2}(i Ba + i Ab)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5 \sqrt{2}(-i Ba - i Ab)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 3 \sqrt{2}(-5 I A^2 a - 3 I B^2 b)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 3 \sqrt{2}(5 I A^2 a + 3 I B^2 b)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - 2(3 B^2 b \cos^2(dx + c) + 5(B^2 a + A^2 b) \cos(dx + c)) \sin(dx + c) / \sqrt{\cos(dx + c)}}}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/15*(5*sqrt(2)*(I*B*a + I*A*b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-I*B*a - I*A*b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(-5*I*A*a - 3*I*B*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(5*I*A*a + 3*I*B*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*B*b*cos(d*x + c)^2 + 5*(B*a + A*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.554.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + b \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))/sqrt(sec(c + d*x)), x)`

3.554.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.554.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.554.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + b \cos(c + dx))}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/(1/cos(c + d*x))^(1/2), x)`

3.555
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.555.1 Optimal result 5038
 3.555.2 Mathematica [A] (verified) 5039
 3.555.3 Rubi [A] (verified) 5039
 3.555.4 Maple [A] (verified) 5043
 3.555.5 Fricas [C] (verification not implemented) 5044
 3.555.6 Sympy [F] 5044
 3.555.7 Maxima [F] 5045
 3.555.8 Giac [F] 5045
 3.555.9 Mupad [F(-1)] 5045

3.555.1 Optimal result

Integrand size = 31, antiderivative size = 180

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{6(Ab + aB)\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(7aA + 5bB)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{2bB \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} + \frac{2(Ab + aB) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2(7aA + 5bB) \sin(c + dx)}{21d \sqrt{\sec(c + dx)}}$$

output

```
2/7*b*B*sin(d*x+c)/d/sec(d*x+c)^(5/2)+2/5*(A*b+B*a)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/21*(7*A*a+5*B*b)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+6/5*(A*b+B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/21*(7*A*a+5*B*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.555.2 Mathematica [A] (verified)

Time = 1.39 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.69

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(252(Ab + aB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20(7aA + 5bB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx) \mid 2\right) \right)}{210d}$$

input `Integrate[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `(Sqrt[Sec[c + d*x]]*(252*(A*b + a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 20*(7*a*A + 5*b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (70*a*A + 65*b*B + 42*(A*b + a*B)*Cos[c + d*x] + 15*b*B*Cos[2*(c + d*x)])*Sin[2*(c + d*x)])/(210*d)`

3.555.3 Rubi [A] (verified)Time = 0.91 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.97, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 3439, 3042, 4484, 27, 3042, 4274, 3042, 4256, 3042, 4258, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))(A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)(A \sec(c + dx) + B)}{\sec^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)(A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx$$

3.555. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow 4484 \\
& \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} - \frac{2}{7} \int -\frac{7(Ab+aB) + (7aA+5bB) \sec(c+dx)}{2 \sec^{\frac{5}{2}}(c+dx)} dx \\
& \downarrow 27 \\
& \frac{1}{7} \int \frac{7(Ab+aB) + (7aA+5bB) \sec(c+dx)}{\sec^{\frac{5}{2}}(c+dx)} dx + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{7} \int \frac{7(Ab+aB) + (7aA+5bB) \csc(c+dx+\frac{\pi}{2})}{\csc(c+dx+\frac{\pi}{2})^{5/2}} dx + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \\
& \downarrow 4274 \\
& \frac{1}{7} \left(7(aB+Ab) \int \frac{1}{\sec^{\frac{5}{2}}(c+dx)} dx + (7aA+5bB) \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)} dx \right) + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{7} \left(7(aB+Ab) \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{5/2}} dx + (7aA+5bB) \int \frac{1}{\csc(c+dx+\frac{\pi}{2})^{3/2}} dx \right) + \\
& \quad \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \\
& \downarrow 4256 \\
& \frac{1}{7} \left(7(aB+Ab) \left(\frac{3}{5} \int \frac{1}{\sqrt{\sec(c+dx)}} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + (7aA+5bB) \left(\frac{1}{3} \int \sqrt{\sec(c+dx)} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) + \\
& \quad \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{7} \left(7(aB+Ab) \left(\frac{3}{5} \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + (7aA+5bB) \left(\frac{1}{3} \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\csc(c+dx+\frac{\pi}{2})}} \right) \right) + \\
& \quad \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)} \\
& \downarrow 4258
\end{aligned}$$

3.555. $\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\frac{1}{7} \left((7aB + Ab) \left(\frac{3}{5} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + (7aA + 5bB) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right) + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left((7aB + Ab) \left(\frac{3}{5} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) + (7aA + 5bB) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right) \right) + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3119

$$\frac{1}{7} \left((7aA + 5bB) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) + 7(aB + Ab) \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} \right) \right) + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3120

$$\frac{1}{7} \left((7aA + 5bB) \left(\frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} + \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} \right) + 7(aB + Ab) \left(\frac{2 \sin(c+dx)}{3d \sqrt{\sec(c+dx)}} + \frac{2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} \right) \right) + \frac{2bB \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

input `Int[((a + b*Cos[c + d*x])*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `(2*b*B*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + (7*(A*b + a*B)*((6*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(5*d) + (2*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)))) + (7*a*A + 5*b*B)*((2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*sqrt[Sec[c + d*x]])))/7`

3.555.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4256 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)])*(d_.)^(n_)*((csc[(e_.) + (f_)*(x_)])*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4484 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^n]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (B*b*n + A*a*(n + 1))*Csc[e + f*x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && LeQ[n, -1]`

3.555.4 Maple [A] (verified)

Time = 12.43 (sec) , antiderivative size = 413, normalized size of antiderivative = 2.29

method	result
default	$-\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(240B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+(-168Ab-168Ba-360Bb)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d}$
parts	$-\frac{2aA\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d}$

input `int((a+cos(d*x+c)*b)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output
$$-2/105*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^8*b+(-168*A*b-168*B*a-360*B*b)*\sin(1/2*d*x+1/2*c)^6*\cos(1/2*d*x+1/2*c)+(140*A*a+168*A*b+168*B*a+280*B*b)*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+(-70*A*a-42*A*b-42*B*a-80*B*b)*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+35*a*A*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^(1/2))-63*A*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2))*b+25*B*b*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^(1/2))-63*B*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2))*a)/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^(1/2)/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d$$

3.555.
$$\int \frac{(a+b \cos(c+dx))(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.555.5 Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.13

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{5\sqrt{2}(7iAa + 5iBb)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5\sqrt{2}(-7iAa - 5iBb)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 63\sqrt{2}(-iB^2a - iA^2b)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 63\sqrt{2}(iB^2a + iA^2b)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) - 2(15B^2b \cos(dx + c)^3 + 21(B^2a + A^2b) \cos(dx + c)^2 + 5(7A^2a + 5B^2b) \cos(dx + c)) \sin(dx + c) / \sqrt{\cos(dx + c)}}}{d}$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `-1/105*(5*sqrt(2)*(7*I*A*a + 5*I*B*b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-7*I*A*a - 5*I*B*b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 63*sqrt(2)*(-I*B*a - I*A*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 63*sqrt(2)*(I*B*a + I*A*b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(15*B*b*cos(d*x + c)^3 + 21*(B*a + A*b)*cos(d*x + c)^2 + 5*(7*A*a + 5*B*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c))/d`

3.555.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + b \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))/sec(c + d*x)**(3/2), x)`

3.555.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.555.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)}{\sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*cos(d*x+c))*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.555.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx))(a + b \cos(c + dx))}{\left(\frac{1}{\cos(c + dx)}\right)^{\frac{3}{2}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x)))/(1/cos(c + d*x))^(3/2), x)`

3.556 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

3.556.1 Optimal result	5046
3.556.2 Mathematica [A] (verified)	5047
3.556.3 Rubi [A] (verified)	5047
3.556.4 Maple [B] (verified)	5052
3.556.5 Fricas [C] (verification not implemented)	5053
3.556.6 Sympy [F(-1)]	5053
3.556.7 Maxima [F]	5054
3.556.8 Giac [F]	5054
3.556.9 Mupad [F(-1)]	5054

3.556.1 Optimal result

Integrand size = 33, antiderivative size = 221

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= -\frac{2(3a^2 A + 5b(Ab + 2aB)) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(2aAb + a^2 B + 3b^2 B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2(3a^2 A + 5b(Ab + 2aB)) \sqrt{\sec(c + dx)} \sin(c + dx)}{5d}$$

$$+ \frac{2a(7Ab + 5aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{15d}$$

$$+ \frac{2aA \sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx)) \sin(c + dx)}{5d}$$

output

```
2/15*a*(7*A*b+5*B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/5*a*A*sec(d*x+c)^(3/2)
)*(b+a*sec(d*x+c))*sin(d*x+c)/d+2/5*(3*A*a^2+5*b*(A*b+2*B*a))*sin(d*x+c)*s
ec(d*x+c)^(1/2)/d-2/5*(3*A*a^2+5*b*(A*b+2*B*a))*(cos(1/2*d*x+1/2*c)^2)^(1/
2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/
2)*sec(d*x+c)^(1/2)/d+2/3*(2*A*a*b+B*a^2+3*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(
1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(
1/2)*sec(d*x+c)^(1/2)/d
```

3.556.2 Mathematica [A] (verified)

Time = 2.94 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.77

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{\sec^{\frac{5}{2}}(c + dx) \left(-12(3a^2A + 5Ab^2 + 10abB) \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20(2aAb + a^2B + 3b^2B) \cos^{\frac{5}{2}}(c + dx) \right)}{30d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(Sec[c + d*x]^(5/2)*(-12*(3*a^2*A + 5*A*b^2 + 10*a*b*B)*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 20*(2*a*A*b + a^2*B + 3*b^2*B)*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 2*(15*(a^2*A + A*b^2 + 2*a*b*B) + 10*a*(2*A*b + a*B)*Cos[c + d*x] + 3*(3*a^2*A + 5*A*b^2 + 10*a*b*B)*Cos[2*(c + d*x)])*Sin[c + d*x])/(30*d)`

3.556.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.93, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3439, 3042, 4514, 27, 3042, 4535, 3042, 4255, 3042, 4258, 3042, 3119, 4534, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx) (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2 (A \sec(c + dx) + B) dx$$

$$\downarrow \text{3042}$$

3.556. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+b\right)^2\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)dx$$

↓ 4514

$$\frac{2}{5}\int\frac{1}{2}\sqrt{\sec(c+dx)}\left(a(7Ab+5aB)\sec^2(c+dx)+(3Aa^2+5b(Ab+2aB))\sec(c+dx)+b(aA+5bB)\right)dx+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)}{5d}$$

↓ 27

$$\frac{1}{5}\int\sqrt{\sec(c+dx)}\left(a(7Ab+5aB)\sec^2(c+dx)+(3Aa^2+5b(Ab+2aB))\sec(c+dx)+b(aA+5bB)\right)dx+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)}{5d}$$

↓ 3042

$$\frac{1}{5}\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a(7Ab+5aB)\csc\left(c+dx+\frac{\pi}{2}\right)^2+(3Aa^2+5b(Ab+2aB))\csc\left(c+dx+\frac{\pi}{2}\right)+b(aA+5bB)\right)dx+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)}{5d}$$

↓ 4535

$$\frac{1}{5}\left(\left(3a^2A+5b(2aB+Ab)\right)\int\sec^{\frac{3}{2}}(c+dx)dx+\int\sqrt{\sec(c+dx)}\left(a(7Ab+5aB)\sec^2(c+dx)+b(aA+5bB)\right)dx\right)+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)}{5d}$$

↓ 3042

$$\frac{1}{5}\left(\left(3a^2A+5b(2aB+Ab)\right)\int\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}dx+\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a(7Ab+5aB)\csc\left(c+dx+\frac{\pi}{2}\right)+b(aA+5bB)\right)dx\right)+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)}{5d}$$

↓ 4255

$$\frac{1}{5}\left(\left(3a^2A+5b(2aB+Ab)\right)\left(\frac{2\sin(c+dx)\sqrt{\sec(c+dx)}}{d}-\int\frac{1}{\sqrt{\sec(c+dx)}}dx\right)+\int\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a(7Ab+5aB)\csc\left(c+dx+\frac{\pi}{2}\right)+b(aA+5bB)\right)dx\right)+\frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)}{5d}$$

↓ 3042

3.556. $\int(a+b\cos(c+dx))^2(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)dx$

$$\frac{1}{5} \left((3a^2A + 5b(2aB + Ab)) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right) \\ \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)}{5d} \\ \downarrow 4258$$

$$\frac{1}{5} \left((3a^2A + 5b(2aB + Ab)) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \right) + \int \sqrt{\cos(c + dx)} dx \right) \\ \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)}{5d} \\ \downarrow 3042$$

$$\frac{1}{5} \left((3a^2A + 5b(2aB + Ab)) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) + \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \right) \\ \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)}{5d} \\ \downarrow 3119$$

$$\frac{1}{5} \left(\int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \left(a(7Ab + 5aB) \csc(c + dx + \frac{\pi}{2})^2 + b(aA + 5bB) \right) + (3a^2A + 5b(2aB + Ab)) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) \right) \\ \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)}{5d} \\ \downarrow 4534$$

$$\frac{1}{5} \left(\frac{5}{3} (a^2B + 2aAb + 3b^2B) \int \sqrt{\sec(c + dx)} dx + (3a^2A + 5b(2aB + Ab)) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \frac{2\sqrt{\cos(c + dx)}}{d} \int \sqrt{\cos(c + dx)} dx \right) \right) \\ \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)}{5d} \\ \downarrow 3042$$

$$\frac{1}{5} \left(\frac{5}{3} (a^2B + 2aAb + 3b^2B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + (3a^2A + 5b(2aB + Ab)) \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \frac{2\sqrt{\cos(c + dx)}}{d} \int \sqrt{\cos(c + dx)} dx \right) \right) \\ \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)}{5d} \\ \downarrow 4258$$

3.556. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\frac{1}{5} \left(\frac{5}{3} (a^2 B + 2aAb + 3b^2 B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + (3a^2 A + 5b(2aB + Ab)) \left(\frac{2 \sin(c+dx)}{5d} \right) \right)$$

$$\frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a \sec(c+dx) + b)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{5}{3} (a^2 B + 2aAb + 3b^2 B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + (3a^2 A + 5b(2aB + Ab)) \left(\frac{2 \sin(c+dx)}{5d} \right) \right)$$

$$\frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a \sec(c+dx) + b)}{5d}$$

↓ 3120

$$\frac{1}{5} \left(\frac{10(a^2 B + 2aAb + 3b^2 B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d} + (3a^2 A + 5b(2aB + Ab)) \left(\frac{2 \sin(c+dx)}{5d} \right) \right)$$

$$\frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a \sec(c+dx) + b)}{5d}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*a*A*Sec[c + d*x]^(3/2)*(b + a*Sec[c + d*x])*Sin[c + d*x])/(5*d) + ((10*(2*a*A*b + a^2*B + 3*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*a*(7*A*b + 5*a*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + (3*a^2*A + 5*b*(A*b + 2*a*B))*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/5`

3.556.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(g_.))^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)])^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4255 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Simp}[b^2*((n-2)/(n-1)) \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4514 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)(x_)]*(B_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m-1)}*((d*\text{Csc}[e + f*x])^n/(f*(m+n))), x] + \text{Simp}[1/(m+n) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m-2)}*(d*\text{Csc}[e + f*x])^n*\text{Simp}[a^2*A*(m+n) + a*b*B*n + (a*(2*A*b + a*B)*(m+n) + b^2*B*(m+n-1))*\text{Csc}[e + f*x] + b*(A*b*(m+n) + a*B*(2*m+n-1))*\text{Csc}[e + f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{GtQ}[m, 1] \&\& !(\text{IGtQ}[n, 1] \&\& !\text{IntegerQ}[m])$

rule 4534 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(b_.))^{(m_.)}*(\text{csc}[(e_.) + (f_.)(x_)]^2*(C_.) + (A_.)), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cot}[e + f*x]*(b*\text{Csc}[e + f*x])^m/(f*(m+1)), x] + \text{Simp}[(C*m + A*(m+1))/(m+1) \text{Int}[(b*\text{Csc}[e + f*x])^m, x], x] /; \text{FreeQ}\{b, e, f, A, C, m\}, x] \&\& \text{NeQ}[C*m + A*(m+1), 0] \&\& !\text{LeQ}[m, -1]$

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.556.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. $2(249) = 498$.

Time = 490.58 (sec) , antiderivative size = 723, normalized size of antiderivative = 3.27

method	result	size
default	Expression too large to display	723
parts	Expression too large to display	914

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

```
output -(-(-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+2/5*A*a^2/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+2*b*(A*b+2*B*a)/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))+2*a*(2*A*b+B*a)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.556.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.29

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx =$$

$$\frac{5\sqrt{2}(iBa^2 + 2iAab + 3iBb^2) \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \dots}{\dots}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="fricas")`

output `-1/15*(5*sqrt(2)*(I*B*a^2 + 2*I*A*a*b + 3*I*B*b^2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-I*B*a^2 - 2*I*A*a*b - 3*I*B*b^2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(3*I*A*a^2 + 10*I*B*a*b + 5*I*A*b^2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-3*I*A*a^2 - 10*I*B*a*b - 5*I*A*b^2)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*A*a^2 + 3*(3*A*a^2 + 10*B*a*b + 5*A*b^2)*cos(d*x + c)^2 + 5*(B*a^2 + 2*A*a*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2)`

3.556.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.556.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sec(d*x + c)^(7/2), x)`

3.556.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sec(d*x + c)^(7/2), x)`

3.556.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + b \cos(c + dx))^2 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^2,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^2, x)`

3.556. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.557 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$

3.557.1 Optimal result	5055
3.557.2 Mathematica [A] (verified)	5056
3.557.3 Rubi [A] (verified)	5056
3.557.4 Maple [B] (verified)	5060
3.557.5 Fracas [C] (verification not implemented)	5061
3.557.6 Sympy [F(-1)]	5062
3.557.7 Maxima [F]	5062
3.557.8 Giac [F]	5063
3.557.9 Mupad [F(-1)]	5063

3.557.1 Optimal result

Integrand size = 33, antiderivative size = 177

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= -\frac{2(2aAb + a^2B - b^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2(a^2A + 3Ab^2 + 6abB) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a(5Ab + 3aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{3d}$$

$$+ \frac{2aA \sqrt{\sec(c + dx)} (b + a \sec(c + dx)) \sin(c + dx)}{3d}$$

```
output 2/3*a*(5*A*b+3*B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/d+2/3*a*A*(b+a*sec(d*x+c))
*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2*(2*A*a*b+B*a^2-B*b^2)*(cos(1/2*d*x+1/2*c)
^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x
+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(A*a^2+3*A*b^2+6*B*a*b)*(cos(1/2*d*x+1/2*
c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d
*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.557.2 Mathematica [A] (verified)

Time = 4.40 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.71

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(-3(2aAb + a^2B - b^2B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + (a^2A + 3Ab^2 + 6abB) \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \right)}{3d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*sqrt[Cos[c + d*x]]*sqrt[Sec[c + d*x]]*(-3*(2*a*A*b + a^2*B - b^2*B)*EllipticE[(c + d*x)/2, 2] + (a^2*A + 3*A*b^2 + 6*a*b*B)*EllipticF[(c + d*x)/2, 2] + (a*(a*A + 3*(2*A*b + a*B)*Cos[c + d*x])*Sin[c + d*x])/Cos[c + d*x]^(3/2)))/(3*d)`

3.557.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3439, 3042, 4514, 27, 3042, 4535, 3042, 4258, 3042, 3120, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx) (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^2 (A \sec(c + dx) + B)}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

3.557. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4514} \\
& \frac{2}{3} \int \frac{-a(5Ab + 3aB) \sec^2(c + dx) - (Aa^2 + 6bBa + 3Ab^2) \sec(c + dx) + b(aA - 3bB)}{2\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)}{3d} \\
& \quad \downarrow \text{27} \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)}{3d} - \\
& \frac{1}{3} \int \frac{-a(5Ab + 3aB) \sec^2(c + dx) - (Aa^2 + 6bBa + 3Ab^2) \sec(c + dx) + b(aA - 3bB)}{\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)}{3d} - \\
& \frac{1}{3} \int \frac{-a(5Ab + 3aB) \csc(c + dx + \frac{\pi}{2})^2 + (-Aa^2 - 6bBa - 3Ab^2) \csc(c + dx + \frac{\pi}{2}) + b(aA - 3bB)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4535} \\
& \frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \int \sqrt{\sec(c + dx)} dx - \int \frac{b(aA - 3bB) - a(5Ab + 3aB) \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx \right) + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx - \int \frac{b(aA - 3bB) - a(5Ab + 3aB) \csc(c + dx + \frac{\pi}{2})^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)}{3d} \\
& \quad \downarrow \text{4258} \\
& \frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx - \int \frac{b(aA - 3bB) - a(5Ab + 3aB) \csc}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} \right) + \\
& \quad \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)}{3d}
\end{aligned}$$

3.557. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

↓ 3042

$$\frac{1}{3} \left((a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx - \int \frac{b(aA - 3bB) - a(5Ab + 3aB)}{\sqrt{\csc(c+dx)}} dx \right) \\ \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2(a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} - \int \frac{b(aA - 3bB) - a(5Ab + 3aB)}{\sqrt{\csc(c+dx)}} dx \right) \\ \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

↓ 4534

$$\frac{1}{3} \left(-3(a^2 B + 2aAb - b^2 B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + \frac{2(a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} \right) \\ \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-3(a^2 B + 2aAb - b^2 B) \int \frac{1}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx + \frac{2(a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} \right) \\ \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

↓ 4258

$$\frac{1}{3} \left(-3(a^2 B + 2aAb - b^2 B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + \frac{2(a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} \right) \\ \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-3(a^2 B + 2aAb - b^2 B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx + \frac{\pi}{2}\right)} dx + \frac{2(a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} \right) \\ \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

↓ 3119

$$\frac{1}{3} \left(\frac{2(a^2 A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} - \frac{6(a^2 B + 2aAb - b^2 B) \sqrt{\cos(c+dx)}}{3d} \right) - \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)}{3d}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*a*A*Sqrt[Sec[c + d*x]]*(b + a*Sec[c + d*x])*Sin[c + d*x])/(3*d) + ((-6*(2*a*A*b + a^2*B - b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*(a^2*A + 3*A*b^2 + 6*a*b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*(5*A*b + 3*a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3`

3.557.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_.)], x_Symbol] := Simp[g^(m+n) Int[(g*Csc[e + f*x])^(p-m-n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4514 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))), x] + Simp[1/(m + n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^n*Simp[a^2*A*(m + n) + a*b*B*n + (a*(2*A*b + a*B)*(m + n) + b^2*B*(m + n - 1))*Csc[e + f*x] + b*(A*b*(m + n) + a*B*(2*m + n - 1))*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && !IntegerQ[m])`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.557.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 649 vs. 2(211) = 422.

Time = 495.72 (sec) , antiderivative size = 650, normalized size of antiderivative = 3.67

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1}(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} \left(\frac{2Ab^2\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1}F(\cos(\frac{dx}{2} + \frac{c}{2}), \sqrt{2})}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} - \frac{2Bb^2\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}}{\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})}} \right)$
parts	Expression too large to display

input `int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2), x, method=_RETURNV ERBOSE)`

$$3.557. \quad \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2*B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+4*B*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+2*B*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))+2*A*a^2*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2*a*(2*A*b+B*a)/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d

```

3.557.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.40

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{\sqrt{2}(-i Aa^2 - 6i Bab - 3i Ab^2) \cos(dx + c) \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \sqrt{2}}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="fracas")`

output `1/3*(sqrt(2)*(-I*A*a^2 - 6*I*B*a*b - 3*I*A*b^2)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(I*A*a^2 + 6*I*B*a*b + 3*I*A*b^2)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(I*B*a^2 + 2*I*A*a*b - I*B*b^2)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(-I*B*a^2 - 2*I*A*a*b + I*B*b^2)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(A*a^2 + 3*(B*a^2 + 2*A*a*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))`

3.557.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.557.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2), x)`

3.557.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2), x)`

3.557.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + b \cos(c + dx))^2 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^2,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^2, x)`

3.558 $\int (a+b \cos(c+dx))^2 (A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$

3.558.1 Optimal result	5064
3.558.2 Mathematica [A] (verified)	5065
3.558.3 Rubi [A] (verified)	5065
3.558.4 Maple [B] (verified)	5069
3.558.5 Fricas [C] (verification not implemented)	5070
3.558.6 Sympy [F(-1)]	5071
3.558.7 Maxima [F]	5071
3.558.8 Giac [F]	5071
3.558.9 Mupad [F(-1)]	5072

3.558.1 Optimal result

Integrand size = 33, antiderivative size = 161

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= -\frac{2(a^2 A - b(Ab + 2aB)) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2(6aAb + 3a^2 B + b^2 B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2b^2 B \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} + \frac{2a^2 A \sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

```
output 2/3*b^2*B*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2*a^2*A*sin(d*x+c)*sec(d*x+c)^(1/2)
)/d-2*(A*a^2-b*(A*b+2*B*a))*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)
)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/
d+2/3*(6*A*a*b+3*B*a^2+B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2
*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2
)/d
```

3.558.2 Mathematica [A] (verified)

Time = 4.13 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.77

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left((-6a^2 A + 6Ab^2 + 12abB) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 2(6aAb + 3a^2 B + b^2 B) \operatorname{EllipticF}\left(\frac{c + dx}{2}, 2\right) + (2(3a^2 A + b^2 B \cos(c + dx)) \sin(c + dx)) / \sqrt{\cos(c + dx)} \right)}{3d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((-6*a^2*A + 6*A*b^2 + 12*a*b*B)*EllipticE[(c + d*x)/2, 2] + 2*(6*a*A*b + 3*a^2*B + b^2*B)*EllipticF[(c + d*x)/2, 2] + (2*(3*a^2*A + b^2*B*Cos[c + d*x])*Sin[c + d*x])/Sqrt[Cos[c + d*x]]))/(3*d)`

3.558.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3439, 3042, 4512, 27, 3042, 4535, 3042, 4258, 3042, 3120, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx) (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^2 (A \sec(c + dx) + B)}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.558. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\
& \quad \downarrow 4512 \\
& \frac{2b^2 B \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} - \\
& \frac{2}{3} \int - \frac{3a^2 A \sec^2(c + dx) + (3Ba^2 + 6Aba + b^2 B) \sec(c + dx) + 3b(Ab + 2aB)}{2\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow 27 \\
& \frac{1}{3} \int \frac{3a^2 A \sec^2(c + dx) + (3Ba^2 + 6Aba + b^2 B) \sec(c + dx) + 3b(Ab + 2aB)}{\sqrt{\sec(c + dx)}} dx + \frac{2b^2 B \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \int \frac{3a^2 A \csc(c + dx + \frac{\pi}{2})^2 + (3Ba^2 + 6Aba + b^2 B) \csc(c + dx + \frac{\pi}{2}) + 3b(Ab + 2aB)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2b^2 B \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 4535 \\
& \frac{1}{3} \left((3a^2 B + 6aAb + b^2 B) \int \sqrt{\sec(c + dx)} dx + \int \frac{3a^2 A \sec^2(c + dx) + 3b(Ab + 2aB)}{\sqrt{\sec(c + dx)}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left((3a^2 B + 6aAb + b^2 B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \int \frac{3a^2 A \csc(c + dx + \frac{\pi}{2})^2 + 3b(Ab + 2aB)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 4258 \\
& \frac{1}{3} \left((3a^2 B + 6aAb + b^2 B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \int \frac{3a^2 A \csc(c + dx + \frac{\pi}{2})^2 + 3b(Ab + 2aB)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow 3042
\end{aligned}$$

3.558. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\frac{1}{3} \left((3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \int \frac{3a^2A \csc(c+dx + \frac{\pi}{2})^2 + 3b(Ab + 2aB)}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx \right) \\ \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \\ \downarrow \text{3120}$$

$$\frac{1}{3} \left(\int \frac{3a^2A \csc(c+dx + \frac{\pi}{2})^2 + 3b(Ab + 2aB)}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx + \frac{2(3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{\pi}{2}, \sqrt{\sec(c+dx)})}{d} \right) \\ \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \\ \downarrow \text{4534}$$

$$\frac{1}{3} \left(-3(a^2A - 2abB - Ab^2) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + \frac{2(3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{\pi}{2}, \sqrt{\sec(c+dx)})}{d} \right) \\ \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(-3(a^2A - 2abB - Ab^2) \int \frac{1}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx + \frac{2(3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{\pi}{2}, \sqrt{\sec(c+dx)})}{d} \right) \\ \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \\ \downarrow \text{4258}$$

$$\frac{1}{3} \left(-3(a^2A - 2abB - Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + \frac{2(3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{\pi}{2}, \sqrt{\sec(c+dx)})}{d} \right) \\ \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} \\ \downarrow \text{3042}$$

$$\frac{1}{3} \left(-3(a^2A - 2abB - Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx + \frac{2(3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{\pi}{2}, \sqrt{\sec(c+dx)})}{d} \right) \\ \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

↓ 3119

$$\frac{1}{3} \left(\frac{2(3a^2B + 6aAb + b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} - \frac{6(a^2A - 2abB - Ab^2) \sqrt{\cos(c+dx)}}{3d\sqrt{\sec(c+dx)}} \right) + \frac{2b^2B \sin(c+dx)}{3d\sqrt{\sec(c+dx)}}$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(2*b^2*B*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]) + ((-6*(a^2*A - A*b^2 - 2*a*b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*(6*a*A*b + 3*a^2*B + b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (6*a^2*A*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3`

3.558.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m+n) Int[(g*Csc[e + f*x])^(p-m-n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4512 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^2*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a^2*A*Cos[e + f*x]*((d*Csc[e + f*x])^(n + 1)/(d*f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*(a*(2*A*b + a*B)*n + (2*a*b*B*n + A*(b^2*n + a^2*(n + 1)))*Csc[e + f*x] + b^2*B*n*Csc[e + f*x]^2), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LeQ[n, -1]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.558.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 404 vs. 2(197) = 394.

Time = 11.53 (sec) , antiderivative size = 405, normalized size of antiderivative = 2.52

method	result
default	$-\frac{8B \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b^2}{3} + 4A \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) a^2 - 4Aab \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \dots\right)$
parts	$-\frac{2A a^2 \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d$

input `int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2), x, method=_RETURNV ERBOSE)`

3.558. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

output
$$\frac{2}{3}(-4B\cos(1/2dx+1/2c)\sin(1/2dx+1/2c)^4b^2+6A\cos(1/2dx+1/2c)\sin(1/2dx+1/2c)^2a^2-6Aa*b*(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2*\sin(1/2dx+1/2c)^2-1)^{(1/2)}*EllipticF(\cos(1/2dx+1/2c),2^{(1/2)})-3A*(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2*\sin(1/2dx+1/2c)^2-1)^{(1/2)}*EllipticE(\cos(1/2dx+1/2c),2^{(1/2)})a^2+3A*(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2*\sin(1/2dx+1/2c)^2-1)^{(1/2)}*EllipticE(\cos(1/2dx+1/2c),2^{(1/2)})b^2+2B*\cos(1/2dx+1/2c)*\sin(1/2dx+1/2c)^2b^2-3B*a^2*(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2*\sin(1/2dx+1/2c)^2-1)^{(1/2)}*EllipticF(\cos(1/2dx+1/2c),2^{(1/2)})-B*b^2*(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2*\sin(1/2dx+1/2c)^2-1)^{(1/2)}*EllipticF(\cos(1/2dx+1/2c),2^{(1/2)})+6B*(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2*\sin(1/2dx+1/2c)^2-1)^{(1/2)}*EllipticE(\cos(1/2dx+1/2c),2^{(1/2)})a*b)/\sin(1/2dx+1/2c)/(2*\cos(1/2dx+1/2c)^2-1)^{(1/2)}/d$$

3.558.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.29

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{\sqrt{2}(-3iBa^2 - 6iAab - iBb^2)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + \sqrt{2}(3iBa^2 + 6iAab + iBb^2)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{(2*\cos(1/2dx+1/2c)^2-1)^{(1/2)}}$$

input `integrate((a+b*cos(dx+c))^2*(A+B*cos(dx+c))*sec(dx+c)^(3/2),x, algorithm="fracas")`

output
$$\frac{1}{3}(\sqrt{2}*(-3I*B*a^2 - 6I*A*a*b - I*B*b^2)*\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I*\sin(dx + c)) + \sqrt{2}*(3I*B*a^2 + 6I*A*a*b + I*B*b^2)*\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I*\sin(dx + c)) - 3*\sqrt{2}*(I*A*a^2 - 2I*B*a*b - I*A*b^2)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I*\sin(dx + c))) - 3*\sqrt{2}*(-I*A*a^2 + 2I*B*a*b + I*A*b^2)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I*\sin(dx + c))) + 2*(B*b^2*\cos(dx + c) + 3*A*a^2)*\sin(dx + c)/\sqrt{\cos(dx + c)})/d$$

3.558.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.558.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2), x)`

3.558.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2), x)`

3.558. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.558.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{\frac{3}{2}} (a + b \cos(c + dx))^2 dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^2,x)`output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^2, x)`

3.559 $\int (a+b \cos(c+dx))^2(A+B \cos(c+dx))\sqrt{\sec(c+dx)} dx$

3.559.1 Optimal result	5073
3.559.2 Mathematica [A] (verified)	5074
3.559.3 Rubi [A] (verified)	5074
3.559.4 Maple [B] (verified)	5079
3.559.5 Fricas [C] (verification not implemented)	5079
3.559.6 Sympy [F]	5080
3.559.7 Maxima [F]	5080
3.559.8 Giac [F]	5081
3.559.9 Mupad [F(-1)]	5081

3.559.1 Optimal result

Integrand size = 33, antiderivative size = 171

$$\int (a+b \cos(c+dx))^2(A+B \cos(c+dx))\sqrt{\sec(c+dx)} dx$$

$$= \frac{2(10aAb + 5a^2B + 3b^2B) \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{\sec(c+dx)}}{5d}$$

$$+ \frac{2(3a^2A + Ab^2 + 2abB) \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) \sqrt{\sec(c+dx)}}{3d}$$

$$+ \frac{2b^2B \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} + \frac{2b(Ab + 2aB) \sin(c+dx)}{3d \sqrt{\sec(c+dx)}}$$

```
output 2/5*b^2*B*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/3*b*(A*b+2*B*a)*sin(d*x+c)/d/sec
(d*x+c)^(1/2)+2/5*(10*A*a*b+5*B*a^2+3*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*
sec(d*x+c)^(1/2)/d+2/3*(3*A*a^2+A*b^2+2*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)
/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)
)*sec(d*x+c)^(1/2)/d
```


3.559.2 Mathematica [A] (verified)

Time = 4.19 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.75

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(6(10aAb + 5a^2B + 3b^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 10(3a^2A + Ab^2 + 2abB) \sqrt{\cos(c + dx)} \right)}{15d}$$

input `Integrate[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(6*(10*a*A*b + 5*a^2*B + 3*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 10*(3*a^2*A + A*b^2 + 2*a*b*B)*Sqrt[Cos[c + d*x]])*EllipticF[(c + d*x)/2, 2] + b*(5*A*b + 10*a*B + 3*b*B*Cos[c + d*x])*Sin[2*(c + d*x)])/(15*d)`

3.559.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3439, 3042, 4512, 27, 3042, 4535, 3042, 4258, 3042, 3119, 4533, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c + dx)} (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^2 (A \sec(c + dx) + B)}{\sec^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx \\
& \quad \downarrow \text{4512} \\
& \frac{2b^2 B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} - \\
& \frac{2}{5} \int - \frac{5a^2 A \sec^2(c + dx) + (5Ba^2 + 10Aba + 3b^2 B) \sec(c + dx) + 5b(Ab + 2aB)}{2 \sec^{\frac{3}{2}}(c + dx)} dx \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{5a^2 A \sec^2(c + dx) + (5Ba^2 + 10Aba + 3b^2 B) \sec(c + dx) + 5b(Ab + 2aB)}{\sec^{\frac{3}{2}}(c + dx)} dx + \\
& \quad \frac{2b^2 B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{5a^2 A \csc(c + dx + \frac{\pi}{2})^2 + (5Ba^2 + 10Aba + 3b^2 B) \csc(c + dx + \frac{\pi}{2}) + 5b(Ab + 2aB)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx + \\
& \quad \frac{2b^2 B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4535} \\
& \frac{1}{5} \left((5a^2 B + 10aAb + 3b^2 B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + \int \frac{5a^2 A \sec^2(c + dx) + 5b(Ab + 2aB)}{\sec^{\frac{3}{2}}(c + dx)} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left((5a^2 B + 10aAb + 3b^2 B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \int \frac{5a^2 A \csc(c + dx + \frac{\pi}{2})^2 + 5b(Ab + 2aB)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4258} \\
& \frac{1}{5} \left((5a^2 B + 10aAb + 3b^2 B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \int \frac{5a^2 A \csc(c + dx + \frac{\pi}{2})^2 + 5b(Ab + 2aB)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)}
\end{aligned}$$

3.559. $\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

↓ 3042

$$\frac{1}{5} \left((5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} dx + \int \frac{5a^2A \csc\left(c+dx+\frac{\pi}{2}\right)^2 + \frac{2b^2B \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)}}{\csc\left(c+dx+\frac{\pi}{2}\right)} dx \right)$$

↓ 3119

$$\frac{1}{5} \left(\int \frac{5a^2A \csc\left(c+dx+\frac{\pi}{2}\right)^2 + 5b(Ab+2aB)}{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}} dx + \frac{2(5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d} + \frac{2b^2B \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 4533

$$\frac{1}{5} \left(\frac{5}{3} (3a^2A + 2abB + Ab^2) \int \sqrt{\sec(c+dx)} dx + \frac{2(5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d} + \frac{2b^2B \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{5}{3} (3a^2A + 2abB + Ab^2) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{2(5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d} + \frac{2b^2B \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 4258

$$\frac{1}{5} \left(\frac{5}{3} (3a^2A + 2abB + Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2(5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx)\right)}{d} + \frac{2b^2B \sin(c+dx)}{5d \sec^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{5}{3} (3a^2A + 2abB + Ab^2) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2(5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c + dx)}}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

$\frac{2b^2B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)}$
 \downarrow 3120

$$\frac{1}{5} \left(\frac{10(3a^2A + 2abB + Ab^2) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} + \frac{2(5a^2B + 10aAb + 3b^2B) \sqrt{\cos(c + dx)}}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

input `Int[(a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(2*b^2*B*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((2*(10*a*A*b + 5*a^2*B + 3*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*(3*a^2*A + A*b^2 + 2*a*b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (10*b*(A*b + 2*a*B)*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]])/5`

3.559.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4512 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^2*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a^2*A*Cos[e + f*x]*((d*Csc[e + f*x])^(n + 1)/(d*f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*(a*(2*A*b + a*B)*n + (2*a*b*B*n + A*(b^2*n + a^2*(n + 1)))*Csc[e + f*x] + b^2*B*n*Csc[e + f*x]^2), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LeQ[n, -1]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

3.559.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 486 vs. 2(203) = 406.

Time = 12.23 (sec) , antiderivative size = 487, normalized size of antiderivative = 2.85

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(-24B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2+(20Ab^2+40Bab+24Bb^2)\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/15*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-24*B*cos(1
/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^2+(20*A*b^2+40*B*a*b+24*B*b^2)*sin(1/
2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-10*A*b^2-20*B*a*b-6*B*b^2)*sin(1/2*d*x
+1/2*c)^2*cos(1/2*d*x+1/2*c)+15*A*a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(
1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+5*A*b^2*(s
in(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(
1/2*d*x+1/2*c),2^(1/2))-30*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b+10*B*a*b*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),2^(1/2))-15*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c
)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-9*B*(sin(1/2*d*x+1/
2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c
),2^(1/2))*b^2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1
/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.559.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.32

$$\int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx =$$

$$\frac{5\sqrt{2}(3iAa^2 + 2iBab + iAb^2)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5\sqrt{2}(-3iAa^2 + 2iBab + iAb^2)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{\dots}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/15*(5*sqrt(2)*(3*I*A*a^2 + 2*I*B*a*b + I*A*b^2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-3*I*A*a^2 - 2*I*B*a*b - I*A*b^2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(-5*I*B*a^2 - 10*I*A*a*b - 3*I*B*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(5*I*B*a^2 + 10*I*A*a*b + 3*I*B*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*B*b^2*cos(d*x + c)^2 + 5*(2*B*a*b + A*b^2)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.559.6 Sympy [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2*sqrt(sec(c + d*x)), x)`

3.559.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A) (b \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sqrt(sec(d*x + c)), x)`

3.559.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2*sqrt(sec(d*x + c)), x)`

3.559.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^2 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx))^2 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^2,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^2, x)`

3.560
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.560.1 Optimal result 5082
 3.560.2 Mathematica [A] (verified) 5083
 3.560.3 Rubi [A] (verified) 5083
 3.560.4 Maple [B] (verified) 5088
 3.560.5 Fricas [C] (verification not implemented) 5089
 3.560.6 Sympy [F] 5090
 3.560.7 Maxima [F] 5090
 3.560.8 Giac [F] 5090
 3.560.9 Mupad [F(-1)] 5091

3.560.1 Optimal result

Integrand size = 33, antiderivative size = 213

$$\int \frac{(a + b \cos(c + dx))^2(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{2(5a^2A + 3Ab^2 + 6abB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(14aAb + 7a^2B + 5b^2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{2b^2B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} + \frac{2b(Ab + 2aB) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{2(14aAb + 7a^2B + 5b^2B) \sin(c + dx)}{21d \sqrt{\sec(c + dx)}}$$

output

```
2/7*b^2*B*sin(d*x+c)/d/sec(d*x+c)^(5/2)+2/5*b*(A*b+2*B*a)*sin(d*x+c)/d/sec
(d*x+c)^(3/2)+2/21*(14*A*a*b+7*B*a^2+5*B*b^2)*sin(d*x+c)/d/sec(d*x+c)^(1/2
)+2/5*(5*A*a^2+3*A*b^2+6*B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1
/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1
/2)/d+2/21*(14*A*a*b+7*B*a^2+5*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2
*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x
+c)^(1/2)/d
```

3.560.2 Mathematica [A] (verified)

Time = 4.47 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.76

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(84(5a^2 A + 3Ab^2 + 6abB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20(14aAb + 7a^2 B + 5b^2 B) \sqrt{\cos(c + dx)} \right)}{210d}$$

input `Integrate[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(84*(5*a^2*A + 3*A*b^2 + 6*a*b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 20*(14*a*A*b + 7*a^2*B + 5*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (42*b*(A*b + 2*a*B)*Cos[c + d*x] + 5*(28*a*A*b + 14*a^2*B + 13*b^2*B + 3*b^2*B*Cos[2*(c + d*x)]))*Sin[2*(c + d*x)])/(210*d)`

3.560.3 Rubi [A] (verified)Time = 1.41 (sec) , antiderivative size = 202, normalized size of antiderivative = 0.95, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 3439, 3042, 4512, 27, 3042, 4535, 3042, 4256, 3042, 4258, 3042, 3120, 4533, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^2 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^2 (A \sec(c + dx) + B)}{\sec^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.560. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^2 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow 4512 \\
& \frac{2b^2 B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} - \\
& \frac{2}{7} \int - \frac{7a^2 A \sec^2(c + dx) + (7Ba^2 + 14Aba + 5b^2 B) \sec(c + dx) + 7b(Ab + 2aB)}{2 \sec^{\frac{5}{2}}(c + dx)} dx \\
& \quad \downarrow 27 \\
& \frac{1}{7} \int \frac{7a^2 A \sec^2(c + dx) + (7Ba^2 + 14Aba + 5b^2 B) \sec(c + dx) + 7b(Ab + 2aB)}{\sec^{\frac{5}{2}}(c + dx)} dx + \\
& \quad \frac{2b^2 B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \int \frac{7a^2 A \csc(c + dx + \frac{\pi}{2})^2 + (7Ba^2 + 14Aba + 5b^2 B) \csc(c + dx + \frac{\pi}{2}) + 7b(Ab + 2aB)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx + \\
& \quad \frac{2b^2 B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 4535 \\
& \frac{1}{7} \left((7a^2 B + 14aAb + 5b^2 B) \int \frac{1}{\sec^{\frac{3}{2}}(c + dx)} dx + \int \frac{7a^2 A \sec^2(c + dx) + 7b(Ab + 2aB)}{\sec^{\frac{5}{2}}(c + dx)} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left((7a^2 B + 14aAb + 5b^2 B) \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx + \int \frac{7a^2 A \csc(c + dx + \frac{\pi}{2})^2 + 7b(Ab + 2aB)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow 4256 \\
& \frac{1}{7} \left((7a^2 B + 14aAb + 5b^2 B) \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} \right) + \int \frac{7a^2 A \csc(c + dx + \frac{\pi}{2})^2 + 7b(Ab + 2aB)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx \right) + \\
& \quad \frac{2b^2 B \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)}
\end{aligned}$$

3.560. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

↓ 3042

$$\frac{1}{7} \left((7a^2B + 14aAb + 5b^2B) \left(\frac{1}{3} \int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \int \frac{7a^2A \csc\left(c + dx + \frac{\pi}{2}\right)^2 + 7b(Ab + 2aB)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2}} dx \right) + \frac{2b^2B \sin(c + dx)}{7d \sec^{5/2}(c + dx)}$$

↓ 4258

$$\frac{1}{7} \left((7a^2B + 14aAb + 5b^2B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \int \frac{7a^2A \csc\left(c + dx + \frac{\pi}{2}\right)^2 + 7b(Ab + 2aB)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2}} dx \right) + \frac{2b^2B \sin(c + dx)}{7d \sec^{5/2}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left((7a^2B + 14aAb + 5b^2B) \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) + \int \frac{7a^2A \csc\left(c + dx + \frac{\pi}{2}\right)^2 + 7b(Ab + 2aB)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2}} dx \right) + \frac{2b^2B \sin(c + dx)}{7d \sec^{5/2}(c + dx)}$$

↓ 3120

$$\frac{1}{7} \left(\int \frac{7a^2A \csc\left(c + dx + \frac{\pi}{2}\right)^2 + 7b(Ab + 2aB)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2}} dx + (7a^2B + 14aAb + 5b^2B) \left(\frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} + \frac{2\sqrt{\cos(c + dx)}}{\sqrt{\sec(c + dx)}} \right) \right) + \frac{2b^2B \sin(c + dx)}{7d \sec^{5/2}(c + dx)}$$

↓ 4533

$$\frac{1}{7} \left(\frac{7}{5} (5a^2A + 6abB + 3Ab^2) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + (7a^2B + 14aAb + 5b^2B) \left(\frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} + \frac{2\sqrt{\cos(c + dx)}}{\sqrt{\sec(c + dx)}} \right) \right) + \frac{2b^2B \sin(c + dx)}{7d \sec^{5/2}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{7}{5} (5a^2A + 6abB + 3Ab^2) \int \frac{1}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx + (7a^2B + 14aAb + 5b^2B) \left(\frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)}}{3d\sqrt{\sec(c+dx)}} \right) \right) \frac{2b^2B \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 4258

$$\frac{1}{7} \left(\frac{7}{5} (5a^2A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + (7a^2B + 14aAb + 5b^2B) \left(\frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)}}{3d\sqrt{\sec(c+dx)}} \right) \right) \frac{2b^2B \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{7}{5} (5a^2A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx + (7a^2B + 14aAb + 5b^2B) \left(\frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)}}{3d\sqrt{\sec(c+dx)}} \right) \right) \frac{2b^2B \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

↓ 3119

$$\frac{1}{7} \left(\frac{14(5a^2A + 6abB + 3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d} + (7a^2B + 14aAb + 5b^2B) \left(\frac{2 \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)}}{3d\sqrt{\sec(c+dx)}} \right) \right) \frac{2b^2B \sin(c+dx)}{7d \sec^{\frac{5}{2}}(c+dx)}$$

input `Int[((a + b*Cos[c + d*x])^2*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(2*b^2*B*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + ((14*(5*a^2*A + 3*A*b^2 + 6*a*b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(5*d) + (14*b*(A*b + 2*a*B)*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + (14*a*A*b + 7*a^2*B + 5*b^2*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]])))/7`

3.560.3.1 Defintions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_*)(x_)]*(g_.)^{(p_.)}*((a_.) + (b_*)\sin[(e_.) + (f_*)(x_)]))^{(m_.)}*((c_.) + (d_*)\sin[(e_.) + (f_*)(x_)]))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{IntegerQ}[n]$
- rule 4256 $\text{Int}[(\text{csc}[(c_.) + (d_*)(x_)]*(b_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n+1)}/(b*d^n)), x] + \text{Simp}[(n+1)/(b^2*n) \text{Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_*)(x_)]*(b_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{EqQ}[n^2, 1/4]$
- rule 4512 $\text{Int}[(\text{csc}[(e_.) + (f_*)(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_*)(x_)]*(b_.) + (a_.)^2*(\text{csc}[(e_.) + (f_*)(x_)]*(B_.) + (A_))), x_Symbol] \rightarrow \text{Simp}[a^2*A*\text{Cos}[e + f*x]*((d*\text{Csc}[e + f*x])^{(n+1)}/(d*f^n)), x] + \text{Simp}[1/(d*n) \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}*(a*(2*A*b + a*B)*n + (2*a*b*B*n + A*(b^2*n + a^2*(n+1)))*\text{Csc}[e + f*x] + b^2*B*n*\text{Csc}[e + f*x]^2), x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LeQ}[n, -1]$

```
rule 4533 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
+ (A_.)), x_Symbol] :> Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] +
Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; Fr
eeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Cs
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.560.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs. 2(241) = 482.

Time = 15.02 (sec) , antiderivative size = 548, normalized size of antiderivative = 2.57

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}}{\left(240B\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2+(-168Ab^2-336Bab-360Bb^2)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}$
parts	Expression too large to display

```
input int((a+cos(d*x+c)*b)^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

3.560.
$$\int \frac{(a+b \cos(c+dx))^2(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

output

```
-2/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*cos(
1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^2+(-168*A*b^2-336*B*a*b-360*B*b^2)*s
in(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(280*A*a*b+168*A*b^2+140*B*a^2+336*
B*a*b+280*B*b^2)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-140*A*a*b-42*A*
b^2-70*B*a^2-84*B*a*b-80*B*b^2)*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+70
*A*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*Ellip
ticF(cos(1/2*d*x+1/2*c),2^(1/2))-105*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin
(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2-63*A*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(co
s(1/2*d*x+1/2*c),2^(1/2))*b^2+35*B*a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin
(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+25*B*b^2*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))-126*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*
x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b)/(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*
x+1/2*c)^2-1)^(1/2)/d
```

3.560.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.19

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$\frac{5\sqrt{2}(7iBa^2 + 14iAab + 5iBb^2)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5\sqrt{2}(-7iBb^2 - 14iAab - 5iBa^2)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{-2\sin(dx + c)^4 + \sin(dx + c)^2} dx$$

input

```
integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorith
m="fracas")
```

output

```
-1/105*(5*sqrt(2)*(7*I*B*a^2 + 14*I*A*a*b + 5*I*B*b^2)*weierstrassPInverse
(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-7*I*B*a^2 - 14*I*A*a*
b - 5*I*B*b^2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) +
21*sqrt(2)*(-5*I*A*a^2 - 6*I*B*a*b - 3*I*A*b^2)*weierstrassZeta(-4, 0, we
ierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*sqrt(2)*(5*I
*A*a^2 + 6*I*B*a*b + 3*I*A*b^2)*weierstrassZeta(-4, 0, weierstrassPInverse
(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(15*B*b^2*cos(d*x + c)^3 + 21*
(2*B*a*b + A*b^2)*cos(d*x + c)^2 + 5*(7*B*a^2 + 14*A*a*b + 5*B*b^2)*cos(d*
x + c))*sin(d*x + c)/sqrt(cos(d*x + c))/d
```

3.560. $\int \frac{(a+b \cos(c+dx))^2 (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

3.560.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^2}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((a+b*cos(d*x+c))**2*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**2/sqrt(sec(c + d*x)), x)`

3.560.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/sqrt(sec(d*x + c)), x)`

3.560.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^2}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^2*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^2/sqrt(sec(d*x + c)), x)`

3.560.9 Mupad **[F(-1)]**

Timed out.

$$\int \frac{(a + b \cos(c + dx))^2 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^2}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^2)/(1/cos(c + d*x))^(1/2),x)`

3.561 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

3.561.1 Optimal result	5092
3.561.2 Mathematica [A] (verified)	5093
3.561.3 Rubi [A] (verified)	5093
3.561.4 Maple [B] (verified)	5099
3.561.5 Fricas [C] (verification not implemented)	5100
3.561.6 Sympy [F(-1)]	5101
3.561.7 Maxima [F]	5101
3.561.8 Giac [F]	5102
3.561.9 Mupad [F(-1)]	5102

3.561.1 Optimal result

Integrand size = 33, antiderivative size = 295

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= -\frac{2(9a^2 Ab + 5Ab^3 + 3a^3 B + 15ab^2 B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(5a^3 A + 21aAb^2 + 21a^2 bB + 21b^3 B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{2(9a^2 Ab + 5Ab^3 + 3a^3 B + 15ab^2 B) \sqrt{\sec(c + dx)} \sin(c + dx)}{21d}$$

$$+ \frac{2a(5a^2 A + 18Ab^2 + 21abB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{21d}$$

$$+ \frac{2a^2(11Ab + 7aB) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{35d}$$

$$+ \frac{2aA \sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))^2 \sin(c + dx)}{7d}$$

output
$$\frac{2}{21}a(5Aa^2+18Ab^2+21Bab)\sec(dx+c)^{3/2}\sin(dx+c)/d+2/35a^2(11Ab+7Ba)\sec(dx+c)^{5/2}\sin(dx+c)/d+2/7aA\sec(dx+c)^{3/2}(b+a\sec(dx+c))^2\sin(dx+c)/d+2/5(9Aa^2b+5Ab^3+3Ba^3+15Bab^2)\sin(dx+c)\sec(dx+c)^{1/2}/d-2/5(9Aa^2b+5Ab^3+3Ba^3+15Bab^2)(\cos(1/2dx+1/2c)^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticE}(\sin(1/2dx+1/2c),2^{1/2})*\cos(dx+c)^{1/2}\sec(dx+c)^{1/2}/d+2/21(5Aa^3+21Aa^2b+21Ba^2b+21Bb^3)(\cos(1/2dx+1/2c)^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticF}(\sin(1/2dx+1/2c),2^{1/2})*\cos(dx+c)^{1/2}\sec(dx+c)^{1/2}/d$$

3.561.2 Mathematica [A] (verified)

Time = 3.82 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.76

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{\sec(c + dx)} \left(-21(9a^2Ab + 5Ab^3 + 3a^3B + 15ab^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 5(5a^3A + 21aAb^2 + 21Ba^2b + 21Bb^3) \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \mid 2\right) + 21(9Aa^2b + 5Ab^3 + 3Ba^3 + 15Bab^2) \sin(c + dx) + 5a(5a^2A + 21Aab + 21Ba^2) \tan(c + dx) + 21a^2(3Ab + aB) \sec(c + dx) \tan(c + dx) + 15a^3A \sec(c + dx)^2 \tan(c + dx) \right)}{(105d)}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output
$$(2\sqrt{\text{Sec}[c + d*x]}*(-21*(9a^2A*b + 5A*b^3 + 3a^3*B + 15*a*b^2*B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2] + 5*(5a^3*A + 21*a*A*b^2 + 21*a^2*b*B + 21*b^3*B)*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2] + 21*(9a^2A*b + 5A*b^3 + 3a^3*B + 15*a*b^2*B)*\text{Sin}[c + d*x] + 5*a*(5a^2*A + 21A*b^2 + 21*a*b*B)*\text{Tan}[c + d*x] + 21*a^2*(3A*b + a*B)*\text{Sec}[c + d*x]*\text{Tan}[c + d*x] + 15*a^3*A*\text{Sec}[c + d*x]^2*\text{Tan}[c + d*x]))/(105*d)$$

3.561.3 Rubi [A] (verified)

Time = 1.94 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.94, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3439, 3042, 4514, 27, 3042, 4564, 27, 3042, 4535, 3042, 4255, 3042, 4258, 3042, 3119, 4534, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.561.
$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$\begin{aligned}
& \int \sec^{\frac{9}{2}}(c+dx)(a+b\cos(c+dx))^3(A+B\cos(c+dx))dx \\
& \quad \downarrow \text{3042} \\
& \int \csc\left(c+dx+\frac{\pi}{2}\right)^{9/2}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^3\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right)dx \\
& \quad \downarrow \text{3439} \\
& \int \sqrt{\sec(c+dx)}(a\sec(c+dx)+b)^3(A\sec(c+dx)+B)dx \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+b\right)^3\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)dx \\
& \quad \downarrow \text{4514} \\
& \frac{2}{7} \int \frac{1}{2} \sqrt{\sec(c+dx)}(b+a\sec(c+dx))\left(a(11Ab+7aB)\sec^2(c+dx)+(5Aa^2+7b(Ab+2aB))\sec(c+dx)+b(aA+7bB)\right)dx + \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)^2}{7d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \sqrt{\sec(c+dx)}(b+a\sec(c+dx))\left(a(11Ab+7aB)\sec^2(c+dx)+(5Aa^2+7b(Ab+2aB))\sec(c+dx)+b(aA+7bB)\right)dx + \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)^2}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(b+a\csc\left(c+dx+\frac{\pi}{2}\right)\right)\left(a(11Ab+7aB)\csc\left(c+dx+\frac{\pi}{2}\right)^2+(5Aa^2+7b(Ab+2aB))\right)dx + \\
& \quad \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)^2}{7d} \\
& \quad \downarrow \text{4564} \\
& \frac{1}{7} \left(\frac{2}{5} \int \frac{1}{2} \sqrt{\sec(c+dx)}(5(aA+7bB)b^2+5a(5Aa^2+21bBa+18Ab^2))\sec^2(c+dx)+7(3Ba^3+9Aba^2+15b^2Ba) \right. \\
& \quad \left. \frac{2aA\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)(a\sec(c+dx)+b)^2}{7d} \right) \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.561. $\int (a+b\cos(c+dx))^3(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

$$\frac{1}{7} \left(\frac{1}{5} \int \sqrt{\sec(c+dx)} (5(aA+7bB)b^2 + 5a(5Aa^2+21bBa+18Ab^2)) \sec^2(c+dx) + 7(3Ba^3+9Aba^2+15b^2Ba + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx)+b)^2}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \left(5(aA+7bB)b^2 + 5a(5Aa^2+21bBa+18Ab^2) \csc\left(c+dx+\frac{\pi}{2}\right)^2 + 7(3Ba^3+9Aba^2+15b^2Ba + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx)+b)^2}{7d} \right) \right)$$

↓ 4535

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\sec(c+dx)} (5(aA+7bB)b^2 + 5a(5Aa^2+21bBa+18Ab^2)) \sec^2(c+dx) \right) dx + 7(3a^3B+9a^2Ab+15a^2bA + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx)+b)^2}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \left(5(aA+7bB)b^2 + 5a(5Aa^2+21bBa+18Ab^2) \csc\left(c+dx+\frac{\pi}{2}\right)^2 \right) dx + 7(3a^3B+9a^2Ab+15a^2bA + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx)+b)^2}{7d} \right) \right)$$

↓ 4255

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \left(5(aA+7bB)b^2 + 5a(5Aa^2+21bBa+18Ab^2) \csc\left(c+dx+\frac{\pi}{2}\right)^2 \right) dx + 7(3a^3B+9a^2Ab+15a^2bA + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx)+b)^2}{7d} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \left(5(aA+7bB)b^2 + 5a(5Aa^2+21bBa+18Ab^2) \csc\left(c+dx+\frac{\pi}{2}\right)^2 \right) dx + 7(3a^3B+9a^2Ab+15a^2bA + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx)+b)^2}{7d} \right) \right)$$

↓ 4258

3.561. $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\csc \left(c + dx + \frac{\pi}{2} \right)} \left(5(aA + 7bB)b^2 + 5a(5Aa^2 + 21bBa + 18Ab^2) \csc \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx + 7(3a^3B + \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2}{7d} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\csc \left(c + dx + \frac{\pi}{2} \right)} \left(5(aA + 7bB)b^2 + 5a(5Aa^2 + 21bBa + 18Ab^2) \csc \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx + 7(3a^3B + \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2}{7d} \right) \right)$$

↓ 3119

$$\frac{1}{7} \left(\frac{1}{5} \left(\int \sqrt{\csc \left(c + dx + \frac{\pi}{2} \right)} \left(5(aA + 7bB)b^2 + 5a(5Aa^2 + 21bBa + 18Ab^2) \csc \left(c + dx + \frac{\pi}{2} \right)^2 \right) dx + 7(3a^3B + \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2}{7d} \right) \right)$$

↓ 4534

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (5a^3A + 21a^2bB + 21aAb^2 + 21b^3B) \int \sqrt{\sec(c + dx)} dx + \frac{10a(5a^2A + 21abB + 18Ab^2) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} + \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2}{7d} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (5a^3A + 21a^2bB + 21aAb^2 + 21b^3B) \int \sqrt{\csc \left(c + dx + \frac{\pi}{2} \right)} dx + \frac{10a(5a^2A + 21abB + 18Ab^2) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} + \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2}{7d} \right) \right)$$

↓ 4258

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (5a^3A + 21a^2bB + 21aAb^2 + 21b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{10a(5a^2A + 21abB + 18Ab^2) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} + \frac{2aA \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2}{7d} \right) \right)$$

↓ 3042

3.561. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (5a^3A + 21a^2bB + 21aAb^2 + 21b^3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}} dx + \frac{10a(5a^2A + 21abB + 18Ab^2) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a \sec(c+dx) + b)^2}{7d} \right) \right)$$

↓ 3120

$$\frac{1}{7} \left(\frac{2a^2(7aB + 11Ab) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{5d} + \frac{1}{5} \left(\frac{10a(5a^2A + 21abB + 18Ab^2) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{10a(5a^2A + 21abB + 18Ab^2) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{2aA \sin(c+dx) \sec^{\frac{3}{2}}(c+dx) (a \sec(c+dx) + b)^2}{7d} \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `(2*a*A*Sec[c + d*x]^(3/2)*(b + a*Sec[c + d*x])^2*Sin[c + d*x])/(7*d) + ((2*a^2*(11*A*b + 7*a*B)*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(5*d) + ((10*(5*a^3*A + 21*a*A*b^2 + 21*a^2*b*B + 21*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (10*a*(5*a^2*A + 18*A*b^2 + 21*a*b*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + 7*(9*a^2*A*b + 5*A*b^3 + 3*a^3*B + 15*a*b^2*B)*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/5)/7`

3.561.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.561. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4514 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))), x] + Simp[1/(m + n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^n*Simp[a^2*A*(m + n) + a*b*B*n + (a*(2*A*b + a*B)*(m + n) + b^2*B*(m + n - 1))*Csc[e + f*x] + b*(A*b*(m + n) + a*B*(2*m + n - 1))*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && !IntegerQ[m])`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

```
rule 4564 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (
a_)), x_Symbol] :> Simp[(-b)*C*Csc[e + f*x]*Cot[e + f*x]*((d*Csc[e + f*x])^
n/(f*(n + 2))), x] + Simp[1/(n + 2) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n +
2) + (B*a*(n + 2) + b*(C*(n + 1) + A*(n + 2)))*Csc[e + f*x] + (a*C + B*b)*(
n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x]
&& !LtQ[n, -1]
```

3.561.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 916 vs. $2(319) = 638$.

Time = 1876.70 (sec) , antiderivative size = 917, normalized size of antiderivative = 3.11

method	result	size
default	Expression too large to display	917
parts	Expression too large to display	1175

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+2*A*a^3*(-1/56*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^4-5/42*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+5/21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2/5*a^2*(3*A*b+B*a)/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+2*b^2*(A*b+3*B*a)/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*E...

```

3.561.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.23

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx =$$

$$\frac{5\sqrt{2}(5i Aa^3 + 21i Ba^2b + 21i Aab^2 + 21i Bb^3) \cos(dx + c)^3 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i)}{...}$$

input

```

integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm="fricas")

```

output `-1/105*(5*sqrt(2)*(5*I*A*a^3 + 21*I*B*a^2*b + 21*I*A*a*b^2 + 21*I*B*b^3)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-5*I*A*a^3 - 21*I*B*a^2*b - 21*I*A*a*b^2 - 21*I*B*b^3)*cos(d*x + c)^3*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*sqrt(2)*(3*I*B*a^3 + 9*I*A*a^2*b + 15*I*B*a*b^2 + 5*I*A*b^3)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*sqrt(2)*(-3*I*B*a^3 - 9*I*A*a^2*b - 15*I*B*a*b^2 - 5*I*A*b^3)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(15*A*a^3 + 21*(3*B*a^3 + 9*A*a^2*b + 15*B*a*b^2 + 5*A*b^3)*cos(d*x + c)^3 + 5*(5*A*a^3 + 21*B*a^2*b + 21*A*a*b^2)*cos(d*x + c)^2 + 21*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^3)`

3.561.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(9/2),x)`

output `Timed out`

3.561.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{9}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(9/2), x)`

3.561. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

3.561.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{9}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(9/2), x)`

3.561.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{9/2} (a + b \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^3, x)`

3.562 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx) dx$

3.562.1 Optimal result	5103
3.562.2 Mathematica [A] (verified)	5104
3.562.3 Rubi [A] (verified)	5104
3.562.4 Maple [B] (verified)	5109
3.562.5 Fricas [C] (verification not implemented)	5110
3.562.6 Sympy [F(-1)]	5111
3.562.7 Maxima [F]	5111
3.562.8 Giac [F]	5112
3.562.9 Mupad [F(-1)]	5112

3.562.1 Optimal result

Integrand size = 33, antiderivative size = 244

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= -\frac{2(3a^3A + 15aAb^2 + 15a^2bB - 5b^3B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(3a^2Ab + 3Ab^3 + a^3B + 9ab^2B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a(3a^2A + 14Ab^2 + 15abB) \sqrt{\sec(c + dx)} \sin(c + dx)}{5d}$$

$$+ \frac{2a^2(9Ab + 5aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{15d}$$

$$+ \frac{2aA \sqrt{\sec(c + dx)} (b + a \sec(c + dx))^2 \sin(c + dx)}{5d}$$

output

```
2/15*a^2*(9*A*b+5*B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/5*a*(3*A*a^2+14*A*b^2+15*B*a*b)*sin(d*x+c)*sec(d*x+c)^(1/2)/d+2/5*a*A*(b+a*sec(d*x+c))^2*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2/5*(3*A*a^3+15*A*a*b^2+15*B*a^2*b-5*B*b^3)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(3*A*a^2*b+3*A*b^3+B*a^3+9*B*a*b^2)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.562.2 Mathematica [A] (verified)

Time = 10.30 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.79

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(-3(3a^3A + 15aAb^2 + 15a^2bB - 5b^3B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 5(3a^2Ab + 3Ab^3) \right)}{15d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*sqrt[Cos[c + d*x]]*sqrt[Sec[c + d*x]]*(-3*(3*a^3*A + 15*a*A*b^2 + 15*a^2*b*B - 5*b^3*B)*EllipticE[(c + d*x)/2, 2] + 5*(3*a^2*A*b + 3*A*b^3 + a^3*B + 9*a*b^2*B)*EllipticF[(c + d*x)/2, 2] + (a*(15*(a^2*A + 3*A*b^2 + 3*a*b*B) + 10*a*(3*A*b + a*B)*Cos[c + d*x] + 9*(a^2*A + 5*A*b^2 + 5*a*b*B)*Cos[2*(c + d*x)])*Sin[c + d*x])/(2*Cos[c + d*x]^(5/2)))/(15*d)`

3.562.3 Rubi [A] (verified)

Time = 1.75 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.02, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4514, 27, 3042, 4564, 27, 3042, 4535, 3042, 4258, 3042, 3120, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))^3(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^3 (A \sec(c + dx) + B)}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

3.562. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

↓ 4514

$$\frac{2}{5} \int - \frac{(b + a \sec(c + dx)) (-a(9Ab + 5aB) \sec^2(c + dx) - (3Aa^2 + 5b(Ab + 2aB)) \sec(c + dx) + b(aA - 5bB))}{2\sqrt{\sec(c + dx)}} dx$$

$$\frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d}$$

↓ 27

$$\frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d} -$$

$$\frac{1}{5} \int \frac{(b + a \sec(c + dx)) (-a(9Ab + 5aB) \sec^2(c + dx) - (3Aa^2 + 5b(Ab + 2aB)) \sec(c + dx) + b(aA - 5bB))}{\sqrt{\sec(c + dx)}} dx$$

↓ 3042

$$\frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d} -$$

$$\frac{1}{5} \int \frac{(b + a \csc(c + dx + \frac{\pi}{2})) (-a(9Ab + 5aB) \csc^2(c + dx + \frac{\pi}{2}) + (-3Aa^2 - 5b(Ab + 2aB)) \csc(c + dx + \frac{\pi}{2}) + b(aA - 5bB))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

↓ 4564

$$\frac{1}{5} \left(\frac{2a^2(5aB + 9Ab) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} - \frac{2}{3} \int \frac{3(aA - 5bB)b^2 - 3a(3Aa^2 + 15bBa + 14Ab^2) \sec^2(c + dx) - b^3}{2\sqrt{\sec(c + dx)}} dx \right)$$

$$\frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d}$$

↓ 27

$$\frac{1}{5} \left(\frac{2a^2(5aB + 9Ab) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} - \frac{1}{3} \int \frac{3(aA - 5bB)b^2 - 3a(3Aa^2 + 15bBa + 14Ab^2) \sec^2(c + dx) - b^3}{\sqrt{\sec(c + dx)}} dx \right)$$

$$\frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{2a^2(5aB + 9Ab) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} - \frac{1}{3} \int \frac{3(aA - 5bB)b^2 - 3a(3Aa^2 + 15bBa + 14Ab^2) \csc(c + dx + \frac{\pi}{2}) + b^3}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \right)$$

$$\frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d}$$

3.562. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

↓ 4535

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \int \sqrt{\sec(c+dx)} dx - \int \frac{3b^2(aA - 5bB) - 3a(3Aa^2 + 15bBa + 14Ab^2)}{\sqrt{\sec(c+dx)}} dx \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx - \int \frac{3b^2(aA - 5bB) - 3a(3Aa^2 + 15bBa + 14Ab^2)}{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}} dx \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)^2}{5d} \right)$$

↓ 4258

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx - \int \frac{3b^2(aA - 5bB) - 3a(3Aa^2 + 15bBa + 14Ab^2)}{\sqrt{\cos(c+dx)}} dx \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(5(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx - \int \frac{3b^2(aA - 5bB) - 3a(3Aa^2 + 15bBa + 14Ab^2)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)^2}{5d} \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{10(a^3B + 3a^2Ab + 9ab^2B + 3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d} - \int \frac{3b^2(aA - 5bB) - 3a(3Aa^2 + 15bBa + 14Ab^2)}{\sqrt{\cos(c+dx)}} dx \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)^2}{5d} \right)$$

↓ 4534

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(3a^3A + 15a^2bB + 15aAb^2 - 5b^3B) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{d} \right) \right. \\ \left. \frac{2aA \sin(c+dx) \sqrt{\sec(c+dx)} (a \sec(c+dx) + b)^2}{5d} \right)$$

3.562. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(3a^3A + 15a^2bB + 15aAb^2 - 5b^3B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx)}{d} \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d} \right) \right.$$

↓ 4258

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(3a^3A + 15a^2bB + 15aAb^2 - 5b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx)}{d} \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d} \right) \right.$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(3a^3A + 15a^2bB + 15aAb^2 - 5b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx)}{d} \right. \right. \\ \left. \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d} \right) \right.$$

↓ 3119

$$\frac{1}{5} \left(\frac{2a^2(5aB + 9Ab) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{3d} + \frac{1}{3} \left(\frac{6a(3a^2A + 15abB + 14Ab^2) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} + \frac{10(a^3B + 9a^2bB + 6aAb^2 + 2b^3B) \sin(c + dx)}{3d} \right) \right. \\ \left. \frac{2aA \sin(c + dx) \sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2}{5d} \right)$$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*a*A*Sqrt[Sec[c + d*x]]*(b + a*Sec[c + d*x])^2*Sin[c + d*x])/(5*d) + ((2*a^2*(9*A*b + 5*a*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d) + ((-6*(3*a^3*A + 15*a*A*b^2 + 15*a^2*b*B - 5*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*(3*a^2*A*b + 3*A*b^3 + a^3*B + 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (6*a*(3*a^2*A + 14*A*b^2 + 15*a*b*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3)/5`

3.562.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.)^(p_))*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4514 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.)^(n_))*((csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_))*((csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))), x] + Simp[1/(m + n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^n*Simp[a^2*A*(m + n) + a*b*B*n + (a*(2*A*b + a*B))*(m + n) + b^2*B*(m + n - 1)]*Csc[e + f*x] + b*(A*b*(m + n) + a*B*(2*m + n - 1))*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && !IntegerQ[m])`

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.
+ (A_.)), x_Symbol] :> Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
)), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] :> Simp[B/b Int[(b*Cs
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

```
rule 4564 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_)), x_Symbol] :> Simp[(-b)*C*Csc[e + f*x]*Cot[e + f*x]*((d*Csc[e + f*x])^
n/(f*(n + 2))), x] + Simp[1/(n + 2) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n +
2) + (B*a*(n + 2) + b*(C*(n + 1) + A*(n + 2)))*Csc[e + f*x] + (a*C + B*b)*(
n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x]
&& !LtQ[n, -1]
```

3.562.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 969 vs. $2(272) = 544$.

Time = 1955.72 (sec) , antiderivative size = 970, normalized size of antiderivative = 3.98

method	result	size
default	Expression too large to display	970
parts	Expression too large to display	1065

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-2*B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+6*B*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+2*B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))+2/5*A*a^3/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)^2*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)+2*a^2*(3*A*b+B*a)*(-1/6*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)...

```

3.562.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.34

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx =$$

$$5\sqrt{2}(iBa^3 + 3iAa^2b + 9iBab^2 + 3iAb^3) \cos(dx + c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))$$

input

```

integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="fricas")

```

output `-1/15*(5*sqrt(2)*(I*B*a^3 + 3*I*A*a^2*b + 9*I*B*a*b^2 + 3*I*A*b^3)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-I*B*a^3 - 3*I*A*a^2*b - 9*I*B*a*b^2 - 3*I*A*b^3)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(3*I*A*a^3 + 15*I*B*a^2*b + 15*I*A*a*b^2 - 5*I*B*b^3)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-3*I*A*a^3 - 15*I*B*a^2*b - 15*I*A*a*b^2 + 5*I*B*b^3)*cos(d*x + c)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*A*a^3 + 9*(A*a^3 + 5*B*a^2*b + 5*A*a*b^2)*cos(d*x + c)^2 + 5*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^2)`

3.562.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.562.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2), x)`

3.562.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2), x)`

3.562.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + b \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^3, x)`

3.563 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx) dx$

3.563.1 Optimal result	5113
3.563.2 Mathematica [A] (verified)	5114
3.563.3 Rubi [A] (verified)	5114
3.563.4 Maple [B] (verified)	5119
3.563.5 Fricas [C] (verification not implemented)	5120
3.563.6 Sympy [F(-1)]	5121
3.563.7 Maxima [F]	5121
3.563.8 Giac [F]	5122
3.563.9 Mupad [F(-1)]	5122

3.563.1 Optimal result

Integrand size = 33, antiderivative size = 239

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= -\frac{2(3a^2 Ab - Ab^3 + a^3 B - 3ab^2 B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

$$+ \frac{2(a^3 A + 9aAb^2 + 9a^2 bB + b^3 B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2a(9aAb + 3a^2 B - 2b^2 B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3d}$$

$$+ \frac{2a^2(aA - bB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3d} + \frac{2bB(b + a \sec(c + dx))^2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}}$$

output

```
2/3*a^2*(A*a-B*b)*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/3*b*B*(b+a*sec(d*x+c))^2
*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2/3*a*(9*A*a*b+3*B*a^2-2*B*b^2)*sin(d*x+c)*
sec(d*x+c)^(1/2)/d-2*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*(cos(1/2*d*x+1/2*c)
^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x
+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(A*a^3+9*A*a*b^2+9*B*a^2*b+B*b^3)*(cos(1/
2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1
/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```


3.563.2 Mathematica [A] (verified)

Time = 7.28 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.69

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(-6(3a^2 Ab - Ab^3 + a^3 B - 3ab^2 B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 2(a^3 A + 9aAb^2 + 9a^2 B) \right)}{3d}$$

3d

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-6*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*EllipticE[(c + d*x)/2, 2] + 2*(a^3*A + 9*a*A*b^2 + 9*a^2*b*B + b^3*B)*EllipticF[(c + d*x)/2, 2] + ((2*a^3*A + b^3*B + 6*a^2*(3*A*b + a*B)*Cos[c + d*x] + b^3*B*Cos[2*(c + d*x)])*Sin[c + d*x])/Cos[c + d*x]^(3/2)))/(3*d)`

3.563.3 Rubi [A] (verified)

Time = 1.66 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4513, 27, 3042, 4564, 27, 3042, 4535, 3042, 4258, 3042, 3120, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^3 (A \sec(c + dx) + B)}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.563. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\
& \quad \downarrow \text{4513} \\
& \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} - \\
& \frac{2}{3} \int - \frac{(b + a \sec(c + dx)) (3a(aA - bB) \sec^2(c + dx) + (3Ba^2 + 6Aba + b^2B) \sec(c + dx) + b(3Ab + 7aB))}{2\sqrt{\sec(c + dx)}} dx \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{(b + a \sec(c + dx)) (3a(aA - bB) \sec^2(c + dx) + (3Ba^2 + 6Aba + b^2B) \sec(c + dx) + b(3Ab + 7aB))}{\sqrt{\sec(c + dx)}} dx + \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{(b + a \csc(c + dx + \frac{\pi}{2})) (3a(aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + (3Ba^2 + 6Aba + b^2B) \csc(c + dx + \frac{\pi}{2}) + b(3Ab + 7aB))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \\
& \quad \downarrow \text{4564} \\
& \frac{1}{3} \left(\frac{2}{3} \int \frac{3((3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \sec^2(c + dx) + (Aa^3 + 9bBa^2 + 9Ab^2a + b^3B) \sec(c + dx))}{2\sqrt{\sec(c + dx)}} dx + \right. \\
& \quad \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \sec^2(c + dx) + (Aa^3 + 9bBa^2 + 9Ab^2a + b^3B) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx + \right. \\
& \quad \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.563. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \csc(c + dx + \frac{\pi}{2})^2 + (Aa^3 + 9bBa^2 + 9Ab^2a + b^3B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 4535

$$\frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \sec^2(c + dx)}{\sqrt{\sec(c + dx)}} dx + (a^3A + 9a^2bB + 9aAb^2 + b^3B) \int \sqrt{\sec(c + dx)} dx + \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \csc(c + dx + \frac{\pi}{2})^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + (a^3A + 9a^2bB + 9aAb^2 + b^3B) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 4258

$$\frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \csc(c + dx + \frac{\pi}{2})^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + (a^3A + 9a^2bB + 9aAb^2 + b^3B) \sqrt{\cos(c + dx)} + \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \csc(c + dx + \frac{\pi}{2})^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + (a^3A + 9a^2bB + 9aAb^2 + b^3B) \sqrt{\cos(c + dx)} + \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 3120

$$\frac{1}{3} \left(\int \frac{(3Ab + 7aB)b^2 + a(3Ba^2 + 9Aba - 2b^2B) \csc(c + dx + \frac{\pi}{2})^2}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2a^2(aA - bB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{d} \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 4534

$$\frac{1}{3} \left(-3(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + \frac{2a(3a^2B + 9aAb - 2b^2B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{3} \left(-3(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2a(3a^2B + 9aAb - 2b^2B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 4258

$$\frac{1}{3} \left(-3(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{2a(3a^2B + 9aAb - 2b^2B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{3} \left(-3(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2a(3a^2B + 9aAb - 2b^2B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

↓ 3119

$$\frac{1}{3} \left(\frac{2a(3a^2B + 9aAb - 2b^2B) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} + \frac{2a^2(aA - bB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{d} + \frac{2(a^3A + 9a^2Ab - 6ab^2B - Ab^3) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{d} \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{3d\sqrt{\sec(c + dx)}} \right)$$

3.563. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*b*B*(b + a*Sec[c + d*x])^2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]) + ((-6*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*(a^3*A + 9*a*A*b^2 + 9*a^2*b*B + b^3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*(9*a*A*b + 3*a^2*B - 2*b^2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d + (2*a^2*(a*A - b*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/d)/3`

3.563.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_.), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

```
rule 4513 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*A*Cot
[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] + Sim
p[1/(d*n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^(n + 1)*Simp[
a*(a*B*n - A*b*(m - n - 1)) + (2*a*b*B*n + A*(b^2*n + a^2*(1 + n)))*Csc[e +
f*x] + b*(b*B*n + a*A*(m + n))*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b,
d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] &
& LeQ[n, -1]
```

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
+ (A_)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
)), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Cs
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

```
rule 4564 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)
)*(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_)), x_Symbol] := Simp[(-b)*C*Csc[e + f*x]*Cot[e + f*x]*((d*Csc[e + f*x])^
n/(f*(n + 2))), x] + Simp[1/(n + 2) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n +
2) + (B*a*(n + 2) + b*(C*(n + 1) + A*(n + 2)))*Csc[e + f*x] + (a*C + B*b)*
(n + 2)*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x]
&& !LtQ[n, -1]
```

3.563.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 885 vs. $2(269) = 538$.

Time = 2002.35 (sec) , antiderivative size = 886, normalized size of antiderivative = 3.71

method	result	size
parts	Expression too large to display	886
default	Expression too large to display	1210

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/3*A*a^3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)
*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*
x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+
1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*((2*cos(1/2*d*x+1
/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*
x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3/2)/sin(1/2*d*x+1/2*c)/d+2*
(A*b^3+3*B*a*b^2)*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(c
os(1/2*d*x+1/2*c),2^(1/2))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d-2*(3*A*a*b^2+3
*B*a^2*b)*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2
*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))/si
n(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d-2*(3*A*a^2*b+B*a^3)*(-
2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*
sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-
1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(co
s(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d-2/3*B*b^3*((2*
cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*sin(1/2*d*x+1/2*...
```

3.563.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.25

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{\sqrt{2}(-i Aa^3 - 9i Ba^2b - 9i Aab^2 - i Bb^3) \cos(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\dots}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,algorith
m="fricas")
```

$$3.563. \quad \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

output `1/3*(sqrt(2)*(-I*A*a^3 - 9*I*B*a^2*b - 9*I*A*a*b^2 - I*B*b^3)*cos(d*x + c) *weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + sqrt(2)*(I*A*a^3 + 9*I*B*a^2*b + 9*I*A*a*b^2 + I*B*b^3)*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) - 3*sqrt(2)*(I*B*a^3 + 3*I*A*a^2*b - 3*I*B*a*b^2 - I*A*b^3)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*sqrt(2)*(-I*B*a^3 - 3*I*A*a^2*b + 3*I*B*a*b^2 + I*A*b^3)*cos(d*x + c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(B*b^3*cos(d*x + c)^2 + A*a^3 + 3*(B*a^3 + 3*A*a^2*b)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))`

3.563.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.563.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2), x)`

3.563.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2), x)`

3.563.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + b \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^3, x)`

3.564 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx) dx$

3.564.1 Optimal result	5123
3.564.2 Mathematica [A] (verified)	5124
3.564.3 Rubi [A] (verified)	5124
3.564.4 Maple [B] (verified)	5129
3.564.5 Fricas [C] (verification not implemented)	5130
3.564.6 Sympy [F(-1)]	5131
3.564.7 Maxima [F]	5131
3.564.8 Giac [F]	5132
3.564.9 Mupad [F(-1)]	5132

3.564.1 Optimal result

Integrand size = 33, antiderivative size = 237

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= -\frac{2(5a^3A - 15aAb^2 - 15a^2bB - 3b^3B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(9a^2Ab + Ab^3 + 3a^3B + 3ab^2B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d}$$

$$+ \frac{2b^2(5Ab + 9aB) \sin(c + dx)}{15d\sqrt{\sec(c + dx)}} + \frac{2a^2(5aA - bB) \sqrt{\sec(c + dx)} \sin(c + dx)}{5d}$$

$$+ \frac{2bB(b + a \sec(c + dx))^2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)}$$

```
output 2/5*b*B*(b+a*sec(d*x+c))^2*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/15*b^2*(5*A*b+9
*B*a)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2/5*a^2*(5*A*a-B*b)*sin(d*x+c)*sec(d*x
+c)^(1/2)/d-2/5*(5*A*a^3-15*A*a*b^2-15*B*a^2*b-3*B*b^3)*(cos(1/2*d*x+1/2*c
)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*
x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/3*(9*A*a^2*b+A*b^3+3*B*a^3+3*B*a*b^2)*(cos
(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2
^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.564.2 Mathematica [A] (verified)

Time = 6.70 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.73

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(12(-5a^3 A + 15aAb^2 + 15a^2 bB + 3b^3 B) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20(9a^2 Ab + Ab^3 \right)}{30d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(12*(-5*a^3*A + 15*a*A*b^2 + 15*a^2*b*B + 3*b^3*B)*EllipticE[(c + d*x)/2, 2] + 20*(9*a^2*A*b + A*b^3 + 3*a^3*B + 3*a*b^2*B)*EllipticF[(c + d*x)/2, 2] + (2*(10*b^2*(A*b + 3*a*B))*Cos[c + d*x] + 3*(10*a^3*A + b^3*B + b^3*B*Cos[2*(c + d*x)]))*Sin[c + d*x])/Sqrt[Cos[c + d*x]])/(30*d)`

3.564.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.02, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4513, 27, 3042, 4562, 27, 3042, 4535, 3042, 4258, 3042, 3120, 4534, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx) (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^3 (A \sec(c + dx) + B)}{\sec^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.564. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 4513

$$\frac{2}{5} \int - \frac{\frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)} - (b + a \sec(c + dx)) (a(5aA - bB) \sec^2(c + dx) + (5Ba^2 + 10Aba + 3b^2B) \sec(c + dx) + b(5Ab + 9aB))}{2 \sec^{\frac{3}{2}}(c + dx)} dx$$

↓ 27

$$\frac{1}{5} \int \frac{(b + a \sec(c + dx)) (a(5aA - bB) \sec^2(c + dx) + (5Ba^2 + 10Aba + 3b^2B) \sec(c + dx) + b(5Ab + 9aB))}{\sec^{\frac{3}{2}}(c + dx)} dx + \frac{\frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}}{2 \sec^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{(b + a \csc(c + dx + \frac{\pi}{2})) (a(5aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + (5Ba^2 + 10Aba + 3b^2B) \csc(c + dx + \frac{\pi}{2}) + b(5Ab + 9aB))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx + \frac{\frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}}{2 \sec^{\frac{3}{2}}(c + dx)}$$

↓ 4562

$$\frac{1}{5} \left(\frac{2b^2(9aB + 5Ab) \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} - \frac{2}{3} \int - \frac{3a^2(5aA - bB) \sec^2(c + dx) + 5(3Ba^3 + 9Aba^2 + 3b^2Ba + Ab^3) \sec(c + dx) + 3b(14Ba^2 + 15Aba + 3b^2B)}{2 \sqrt{\sec(c + dx)}} dx \right) + \frac{\frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}}{2 \sqrt{\sec(c + dx)}}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{3a^2(5aA - bB) \sec^2(c + dx) + 5(3Ba^3 + 9Aba^2 + 3b^2Ba + Ab^3) \sec(c + dx) + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\sec(c + dx)}} dx \right) + \frac{\frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}}{\sqrt{\sec(c + dx)}}$$

↓ 3042

3.564. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{3a^2(5aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + 5(3Ba^3 + 9Aba^2 + 3b^2Ba + Ab^3) \csc(c + dx + \frac{\pi}{2}) + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + 5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right)$$

↓ 4535

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{3a^2(5aA - bB) \sec^2(c + dx) + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\sec(c + dx)} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + 5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \sqrt{\sec(c + dx)} dx \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{3a^2(5aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + 5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right)$$

↓ 4258

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{3a^2(5aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + 5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{3a^2(5aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)}} dx + 5(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx \right)$$

↓ 3120

$$\frac{1}{5} \left(\frac{1}{3} \left(\int \frac{3a^2(5aA - bB) \csc(c + dx + \frac{\pi}{2})^2 + 3b(14Ba^2 + 15Aba + 3b^2B)}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{10(3a^3B + 9a^2Ab + 3ab^2B + Ab^3)}{2bB \sin(c + dx)(a \sec(c + dx) + b)^2} \right) \right. \\ \left. \frac{5d \sec^{\frac{3}{2}}(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \downarrow 4534$$

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + \frac{6a^2(5aA - bB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{6a^2(5aA - bB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \downarrow 4258$$

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{6a^2(5aA - bB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{3} \left(-3(5a^3A - 15a^2bB - 15aAb^2 - 3b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{6a^2(5aA - bB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} \right) \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \downarrow 3119$$

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{6a^2(5aA - bB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d} + \frac{10(3a^3B + 9a^2Ab + 3ab^2B + Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \right) \right. \\ \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{5d \sec^{\frac{3}{2}}(c + dx)} \right)$$

3.564. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

input `Int[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(2*b*B*(b + a*Sec[c + d*x])^2*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((2*b^2*(5*A*b + 9*a*B)*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]) + ((-6*(5*a^3*A - 15*a*A*b^2 - 15*a^2*b*B - 3*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*(9*a^2*A*b + A*b^3 + 3*a^3*B + 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (6*a^2*(5*a*A - b*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d)/3)/5`

3.564.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4513 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a*A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^(n + 1)*Simp[a*(a*B*n - A*b*(m - n - 1)) + (2*a*b*B*n + A*(b^2*n + a^2*(1 + n)))*Csc[e + f*x] + b*(b*B*n + a*A*(m + n))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && LeQ[n, -1]`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

rule 4562 `Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (n*(a*C + B*b) + A*a*(n + 1))*Csc[e + f*x] + b*C*n*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && LtQ[n, -1]`

3.564.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 640 vs. $2(265) = 530$.

Time = 15.57 (sec) , antiderivative size = 641, normalized size of antiderivative = 2.70

method	result
default	$\frac{2\left(-24B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3 + 20A \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3 + 60B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)a b^2 + 24B \cos\left(\frac{dx}{2}\right)}{\dots}$
parts	Expression too large to display

3.564. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$


```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/15*(-24*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6*b^3+20*A*cos(1/2*d*x+
1/2*c)*sin(1/2*d*x+1/2*c)^4*b^3+60*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)
^4*a*b^2+24*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*b^3-30*A*cos(1/2*d*x
+1/2*c)*sin(1/2*d*x+1/2*c)^2*a^3-10*A*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c
)^2*b^3+45*A*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)
^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+5*A*b^3*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1
/2))+15*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*El
lipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^3-45*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*
(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b
^2-30*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a*b^2-6*B*cos(1/2*d*x+1/2*
c)*sin(1/2*d*x+1/2*c)^2*b^3+15*B*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1
/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+15*B*a*b^2*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(co
s(1/2*d*x+1/2*c),2^(1/2))-45*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x
+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b-9*B*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d
*x+1/2*c),2^(1/2))*b^3/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2
)/d
```

3.564.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.14

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx =$$

$$\frac{5\sqrt{2}(3iBa^3 + 9iAa^2b + 3iBab^2 + iAb^3)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorith
m="fracas")
```

output `-1/15*(5*sqrt(2)*(3*I*B*a^3 + 9*I*A*a^2*b + 3*I*B*a*b^2 + I*A*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-3*I*B*a^3 - 9*I*A*a^2*b - 3*I*B*a*b^2 - I*A*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 3*sqrt(2)*(5*I*A*a^3 - 15*I*B*a^2*b - 15*I*A*a*b^2 - 3*I*B*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*sqrt(2)*(-5*I*A*a^3 + 15*I*B*a^2*b + 15*I*A*a*b^2 + 3*I*B*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(3*B*b^3*cos(d*x + c)^2 + 15*A*a^3 + 5*(3*B*a*b^2 + A*b^3)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.564.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.564.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2), x)`

3.564.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2), x)`

3.564.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + b \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^3, x)`

3.565 $\int (a+b \cos(c+dx))^3 (A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$

3.565.1 Optimal result	5133
3.565.2 Mathematica [A] (verified)	5134
3.565.3 Rubi [A] (verified)	5134
3.565.4 Maple [B] (verified)	5139
3.565.5 Fricas [C] (verification not implemented)	5140
3.565.6 Sympy [F]	5141
3.565.7 Maxima [F]	5141
3.565.8 Giac [F]	5142
3.565.9 Mupad [F(-1)]	5142

3.565.1 Optimal result

Integrand size = 33, antiderivative size = 245

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{2(15a^2Ab + 3Ab^3 + 5a^3B + 9ab^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d}$$

$$+ \frac{2(21a^3A + 21aAb^2 + 21a^2bB + 5b^3B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{2b^2(7Ab + 11aB) \sin(c + dx)}{35d \sec^{\frac{3}{2}}(c + dx)} + \frac{2b(21aAb + 18a^2B + 5b^2B) \sin(c + dx)}{21d \sqrt{\sec(c + dx)}}$$

$$+ \frac{2bB(b + a \sec(c + dx))^2 \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)}$$

output

```
2/35*b^2*(7*A*b+11*B*a)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/7*b*B*(b+a*sec(d*x+c))^2*sin(d*x+c)/d/sec(d*x+c)^(5/2)+2/21*b*(21*A*a*b+18*B*a^2+5*B*b^2)*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2/5*(15*A*a^2*b+3*A*b^3+5*B*a^3+9*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/21*(21*A*a^3+21*A*a*b^2+21*B*a^2*b+5*B*b^3)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.565.2 Mathematica [A] (verified)

Time = 7.10 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.73

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(84(15a^2 Ab + 3Ab^3 + 5a^3 B + 9ab^2 B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 20(21a^3 A + 21aAb^2 \right)}{210d}$$

input `Integrate[(a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(84*(15*a^2*A*b + 3*A*b^3 + 5*a^3*B + 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 20*(21*a^3*A + 21*a*A*b^2 + 21*a^2*b*B + 5*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + b*(42*b*(A*b + 3*a*B)*Cos[c + d*x] + 5*(42*a*A*b + 42*a^2*B + 13*b^2*B + 3*b^2*B*Cos[2*(c + d*x)]))*Sin[2*(c + d*x)])/(210*d)`

3.565.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.03, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4513, 27, 3042, 4562, 27, 3042, 4535, 3042, 4258, 3042, 3119, 4533, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c + dx)} (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3 \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^3 (A \sec(c + dx) + B)}{\sec^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.565. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow \text{4513} \\
& \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)} - \\
& \frac{2}{7} \int - \frac{(b + a \sec(c + dx))(a(7aA + bB) \sec^2(c + dx) + (7Ba^2 + 14Aba + 5b^2B) \sec(c + dx) + b(7Ab + 11aB))}{2 \sec^{\frac{5}{2}}(c + dx)} dx \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \frac{(b + a \sec(c + dx))(a(7aA + bB) \sec^2(c + dx) + (7Ba^2 + 14Aba + 5b^2B) \sec(c + dx) + b(7Ab + 11aB))}{\sec^{\frac{5}{2}}(c + dx)} dx + \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \frac{(b + a \csc(c + dx + \frac{\pi}{2})) (a(7aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + (7Ba^2 + 14Aba + 5b^2B) \csc(c + dx + \frac{\pi}{2}) + b(7Ab + 11aB))}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx + \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4562} \\
& \frac{1}{7} \left(\frac{2b^2(11aB + 7Ab) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} - \frac{2}{5} \int - \frac{5a^2(7aA + bB) \sec^2(c + dx) + 7(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \sec(c + dx) + 5b(18Ba^2 + 21Aba + 5b^2B)}{2 \sec^{\frac{3}{2}}(c + dx)} dx \right) \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \sec^2(c + dx) + 7(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \sec(c + dx) + 5b(18Ba^2 + 21Aba + 5b^2B)}{\sec^{\frac{3}{2}}(c + dx)} dx \right) \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.565. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7(5Ba^3 + 15Aba^2 + 9b^2Ba + 3Ab^3) \csc(c + dx + \frac{\pi}{2}) + 5b(18Ba^2 + 21Aba + 5b^2B)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + 7(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \right)$$

↓ 4535

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \sec^2(c + dx) + 5b(18Ba^2 + 21Aba + 5b^2B)}{\sec^{\frac{3}{2}}(c + dx) \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + 7(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 5b(18Ba^2 + 21Aba + 5b^2B)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + 7(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \right)$$

↓ 4258

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 5b(18Ba^2 + 21Aba + 5b^2B)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + 7(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 5b(18Ba^2 + 21Aba + 5b^2B)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + 7(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3) \right)$$

↓ 3119

$$\frac{1}{7} \left(\frac{1}{5} \int \frac{5a^2(7aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 5b(18Ba^2 + 21Aba + 5b^2B)}{\csc(c + dx + \frac{\pi}{2})^{3/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{7d \sec^{\frac{5}{2}}(c + dx)}} dx + \frac{14(5a^3B + 15a^2Ab + 9ab^2B + 3Ab^3)}{7d \sec^{\frac{5}{2}}(c + dx)} \right)$$

3.565. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

↓ 4533

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \int \sqrt{\sec(c+dx)} dx + \frac{10b(18a^2B + 21aAb + 5b^2B) \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2bB \sin(c+dx)(a \sec(c+dx) + b)^2}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{10b(18a^2B + 21aAb + 5b^2B) \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2bB \sin(c+dx)(a \sec(c+dx) + b)^2}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \right)$$

↓ 4258

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{10b(18a^2B + 21aAb + 5b^2B) \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2bB \sin(c+dx)(a \sec(c+dx) + b)^2}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{5}{3} (21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{10b(18a^2B + 21aAb + 5b^2B) \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{2bB \sin(c+dx)(a \sec(c+dx) + b)^2}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \right)$$

↓ 3120

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{10b(18a^2B + 21aAb + 5b^2B) \sin(c+dx)}{3d\sqrt{\sec(c+dx)}} + \frac{10(21a^3A + 21a^2bB + 21aAb^2 + 5b^3B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{3d} + \frac{2bB \sin(c+dx)(a \sec(c+dx) + b)^2}{7d \sec^{\frac{5}{2}}(c+dx)} \right) \right)$$

input `Int[(a + b*cos[c + d*x])^3*(A + B*cos[c + d*x])*sqrt[Sec[c + d*x]],x]`


```
output (2*b*B*(b + a*Sec[c + d*x])^2*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + ((2
*b^2*(7*A*b + 11*a*B)*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + ((14*(15*a^
2*A*b + 3*A*b^3 + 5*a^3*B + 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d
*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (10*(21*a^3*A + 21*a*A*b^2 + 21*a^2*b*B
+ 5*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]]
)/(3*d) + (10*b*(21*a*A*b + 18*a^2*B + 5*b^2*B)*Sin[c + d*x])/(3*d*Sqrt[Se
c[c + d*x]]))/5)/7
```

3.565.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4258 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

rule 4513 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a*A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^(n + 1)*Simp[a*(a*B*n - A*b*(m - n - 1)) + (2*a*b*B*n + A*(b^2*n + a^2*(1 + n)))*Csc[e + f*x] + b*(b*B*n + a*A*(m + n))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && LeQ[n, -1]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

rule 4562 `Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (n*(a*C + B*b) + A*a*(n + 1))*Csc[e + f*x] + b*C*n*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && LtQ[n, -1]`

3.565.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 663 vs. $2(273) = 546$.

Time = 16.21 (sec) , antiderivative size = 664, normalized size of antiderivative = 2.71

method	result
default	$-2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \left(240B \cos\left(\frac{dx}{2} + \frac{c}{2}\right)\left(\sin^8\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3 + (-168A b^3 - 504B a b^2 - 360B b^3)\left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)$
parts	Expression too large to display

3.565. $\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/105*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(240*B*cos(
1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^8*b^3+(-168*A*b^3-504*B*a*b^2-360*B*b^3)
*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)+(420*A*a*b^2+168*A*b^3+420*B*a^2*
b+504*B*a*b^2+280*B*b^3)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-210*A*a
*b^2-42*A*b^3-210*B*a^2*b-126*B*a*b^2-80*B*b^3)*sin(1/2*d*x+1/2*c)^2*cos(1
/2*d*x+1/2*c)+105*A*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+105*A*a*b^2*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/
2*c),2^(1/2))-315*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b-63*A*(sin(1/2*d*x+1/2*
c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),
2^(1/2))*b^3+105*B*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)
)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+25*B*b^3*(sin(1/2*d*x+1
/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*
c),2^(1/2))-105*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^3-189*B*(sin(1/2*d*x+1/2*c)^
2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(
1/2))*a*b^2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*
d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.565.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.20

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx =$$

$$\frac{5\sqrt{2}(21i Aa^3 + 21i Ba^2b + 21i Aab^2 + 5i Bb^3)\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{\dots}$$

```
input integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorith
m="fricas")
```

output `-1/105*(5*sqrt(2)*(21*I*A*a^3 + 21*I*B*a^2*b + 21*I*A*a*b^2 + 5*I*B*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*sqrt(2)*(-21*I*A*a^3 - 21*I*B*a^2*b - 21*I*A*a*b^2 - 5*I*B*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*sqrt(2)*(-5*I*B*a^3 - 15*I*A*a^2*b - 9*I*B*a*b^2 - 3*I*A*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*sqrt(2)*(5*I*B*a^3 + 15*I*A*a^2*b + 9*I*B*a*b^2 + 3*I*A*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(15*B*b^3*cos(d*x + c)^3 + 21*(3*B*a*b^2 + A*b^3)*cos(d*x + c)^2 + 5*(21*B*a^2*b + 21*A*a*b^2 + 5*B*b^3)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.565.6 Sympy [F]

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx)) (a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)} dx$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**3*sqrt(sec(c + d*x)), x)`

3.565.7 Maxima [F]

$$\int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sqrt(sec(d*x + c)), x)`

3.565.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3*sqrt(sec(d*x + c)), x)`

3.565.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \cos(c + dx))^3 (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx \\ &= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx))^3 dx \end{aligned}$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^3,x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^3, x)`

3.566
$$\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.566.1 Optimal result 5143
 3.566.2 Mathematica [A] (verified) 5144
 3.566.3 Rubi [A] (verified) 5144
 3.566.4 Maple [B] (verified) 5150
 3.566.5 Fricas [C] (verification not implemented) 5151
 3.566.6 Sympy [F] 5151
 3.566.7 Maxima [F] 5152
 3.566.8 Giac [F] 5152
 3.566.9 Mupad [F(-1)] 5152

3.566.1 Optimal result

Integrand size = 33, antiderivative size = 295

$$\int \frac{(a + b \cos(c + dx))^3(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{2(15a^3A + 27aAb^2 + 27a^2bB + 7b^3B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{15d}$$

$$+ \frac{2(21a^2Ab + 5Ab^3 + 7a^3B + 15ab^2B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{21d}$$

$$+ \frac{2b^2(9Ab + 13aB) \sin(c + dx)}{63d \sec^{\frac{5}{2}}(c + dx)} + \frac{2b(27aAb + 22a^2B + 7b^2B) \sin(c + dx)}{45d \sec^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{2(21a^2Ab + 5Ab^3 + 7a^3B + 15ab^2B) \sin(c + dx)}{21d \sqrt{\sec(c + dx)}} + \frac{2bB(b + a \sec(c + dx))^2 \sin(c + dx)}{9d \sec^{\frac{7}{2}}(c + dx)}$$

output

```
2/63*b^2*(9*A*b+13*B*a)*sin(d*x+c)/d/sec(d*x+c)^(5/2)+2/45*b*(27*A*a*b+22*
B*a^2+7*B*b^2)*sin(d*x+c)/d/sec(d*x+c)^(3/2)+2/9*b*B*(b+a*sec(d*x+c))^2*si
n(d*x+c)/d/sec(d*x+c)^(7/2)+2/21*(21*A*a^2*b+5*A*b^3+7*B*a^3+15*B*a*b^2)*s
in(d*x+c)/d/sec(d*x+c)^(1/2)+2/15*(15*A*a^3+27*A*a*b^2+27*B*a^2*b+7*B*b^3)
*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2
*c), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d+2/21*(21*A*a^2*b+5*A*b^3+
7*B*a^3+15*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellipt
icF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.566.2 Mathematica [A] (verified)

Time = 7.45 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.74

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(168(15a^3 A + 27aAb^2 + 27a^2 bB + 7b^3 B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 120(21a^2 Ab + 5A^2 b^2 + 7a^3 B + 15a^2 b^2 B) \operatorname{Sqrt}[\cos(c + dx)] \operatorname{EllipticF}\left[\frac{c + dx}{2}, 2\right] + (7b^3 (108a^2 A^2 b + 108a^2 b^2 B + 43b^2 B^2) \cos(c + dx) + 5(252a^2 A^2 b + 78A^2 b^3 + 84a^3 B + 234a^2 b^2 B + 18b^2 (A^2 b + 3a^2 B)) \cos(2(c + dx)) + 7b^3 B \cos(3(c + dx))) \sin(2(c + dx)) \right)}{1260d}$$

input `Integrate[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(Sqrt[Sec[c + d*x]]*(168*(15*a^3*A + 27*a*A*b^2 + 27*a^2*b*B + 7*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 120*(21*a^2*A*b + 5*A*b^3 + 7*a^3*B + 15*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + (7*b*(108*a*A*b + 108*a^2*B + 43*b^2*B)*Cos[c + d*x] + 5*(252*a^2*A*b + 78*A*b^3 + 84*a^3*B + 234*a*b^2*B + 18*b^2*(A*b + 3*a*B))*Cos[2*(c + d*x)] + 7*b^3*B*Cos[3*(c + d*x)]))*Sin[2*(c + d*x)])/(1260*d)`

3.566.3 Rubi [A] (verified)Time = 1.93 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.95, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3439, 3042, 4513, 27, 3042, 4562, 27, 3042, 4535, 3042, 4256, 3042, 4258, 3042, 3120, 4533, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^3 (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{(a \sec(c + dx) + b)^3 (A \sec(c + dx) + B)}{\sec^{\frac{9}{2}}(c + dx)} dx$$

$$\begin{aligned}
& \int \frac{(a \csc(c + dx + \frac{\pi}{2}) + b)^3 (A \csc(c + dx + \frac{\pi}{2}) + B)}{\csc(c + dx + \frac{\pi}{2})^{9/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} - \\
& \frac{2}{9} \int \frac{(b + a \sec(c + dx)) (3a(3aA + bB) \sec^2(c + dx) + (9Ba^2 + 18Aba + 7b^2B) \sec(c + dx) + b(9Ab + 13aB))}{2 \sec^{\frac{7}{2}}(c + dx)} dx \\
& \quad \downarrow \text{4513} \\
& \frac{1}{9} \int \frac{(b + a \sec(c + dx)) (3a(3aA + bB) \sec^2(c + dx) + (9Ba^2 + 18Aba + 7b^2B) \sec(c + dx) + b(9Ab + 13aB))}{\sec^{\frac{7}{2}}(c + dx)} dx \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{9} \int \frac{(b + a \csc(c + dx + \frac{\pi}{2})) (3a(3aA + bB) \csc^2(c + dx + \frac{\pi}{2}) + (9Ba^2 + 18Aba + 7b^2B) \csc(c + dx + \frac{\pi}{2}) + b(9Ab + 13aB))}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \int \frac{(b + a \csc(c + dx + \frac{\pi}{2})) (3a(3aA + bB) \csc^2(c + dx + \frac{\pi}{2}) + (9Ba^2 + 18Aba + 7b^2B) \csc(c + dx + \frac{\pi}{2}) + b(9Ab + 13aB))}{\csc(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow \text{4562} \\
& \frac{1}{9} \left(\frac{2b^2(13aB + 9Ab) \sin(c + dx)}{7d \sec^{\frac{5}{2}}(c + dx)} - \frac{2}{7} \int \frac{21a^2(3aA + bB) \sec^2(c + dx) + 9(7Ba^3 + 21Aba^2 + 15b^2Ba + 5Ab^3) \sec(c + dx) + 7b(22Ba^2 + 27Aba + 5Ab^2)}{2 \sec^{\frac{5}{2}}(c + dx)} dx \right) \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{9} \left(\frac{1}{7} \int \frac{21a^2(3aA + bB) \sec^2(c + dx) + 9(7Ba^3 + 21Aba^2 + 15b^2Ba + 5Ab^3) \sec(c + dx) + 7b(22Ba^2 + 27Aba + 5Ab^2)}{\sec^{\frac{5}{2}}(c + dx)} dx \right) \\
& \quad \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.566. $\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\frac{1}{9} \left(\frac{1}{7} \int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 9(7Ba^3 + 21Aba^2 + 15b^2Ba + 5Ab^3) \csc(c + dx + \frac{\pi}{2}) + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \right)$$

↓ 4535

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \sec^2(c + dx) + 7b(22Ba^2 + 27Aba + 7b^2B)}{\sec^{\frac{5}{2}}(c + dx) \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \right) \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \right) \right)$$

↓ 4256

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \right) \right)$$

↓ 3042

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \right) \right)$$

↓ 4258

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2} \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5Ab^3) \right) \right)$$

↓ 3042

3.566. $\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5) \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{7/2}(c + dx)} \right) \right. \\ \left. \downarrow \text{3120} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\int \frac{21a^2(3aA + bB) \csc(c + dx + \frac{\pi}{2})^2 + 7b(22Ba^2 + 27Aba + 7b^2B)}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx + 9(7a^3B + 21a^2Ab + 15ab^2B + 5) \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{7/2}(c + dx)} \right) \right. \\ \left. \downarrow \text{4533} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{21}{5} (15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + \frac{14b(22a^2B + 27aAb + 7b^2B) \sin(c + dx)}{5d \sec^{3/2}(c + dx)} \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{7/2}(c + dx)} \right) \right. \\ \left. \downarrow \text{3042} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{21}{5} (15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{14b(22a^2B + 27aAb + 7b^2B) \sin(c + dx)}{5d \sec^{3/2}(c + dx)} \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{7/2}(c + dx)} \right) \right. \\ \left. \downarrow \text{4258} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{21}{5} (15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{14b(22a^2B + 27aAb + 7b^2B) \sin(c + dx)}{5d \sec^{3/2}(c + dx)} \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{7/2}(c + dx)} \right) \right. \\ \left. \downarrow \text{3042} \right.$$

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{21}{5} (15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{14b(22a^2B + 27aAb + 7b^2B) \sin(c + dx)}{5d \sec^{3/2}(c + dx)} \right. \right. \\ \left. \left. \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{7/2}(c + dx)} \right) \right.$$

↓ 3119

$$\frac{1}{9} \left(\frac{1}{7} \left(\frac{14b(22a^2B + 27aAb + 7b^2B) \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{42(15a^3A + 27a^2bB + 27aAb^2 + 7b^3B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{5d} \right) + \frac{2bB \sin(c + dx)(a \sec(c + dx) + b)^2}{9d \sec^{\frac{7}{2}}(c + dx)} \right)$$

input `Int[((a + b*Cos[c + d*x])^3*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `(2*b*B*(b + a*Sec[c + d*x])^2*Sin[c + d*x])/(9*d*Sec[c + d*x]^(7/2)) + ((2*b^2*(9*A*b + 13*a*B)*Sin[c + d*x])/(7*d*Sec[c + d*x]^(5/2)) + ((42*(15*a^3*A + 27*a*A*b^2 + 27*a^2*b*B + 7*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(5*d) + (14*b*(27*a*A*b + 22*a^2*B + 7*b^2*B)*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)) + 9*(21*a^2*A*b + 5*A*b^3 + 7*a^3*B + 15*a*b^2*B)*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]]))/7)/9`

3.566.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4513 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a*A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*n)), x] + Simp[1/(d*n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^(n + 1)*Simp[a*(a*B*n - A*b*(m - n - 1)) + (2*a*b*B*n + A*(b^2*n + a^2*(1 + n)))*Csc[e + f*x] + b*(b*B*n + a*A*(m + n))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && LeQ[n, -1]`

rule 4533 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_.)), x_Symbol] := Simp[A*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*m)), x] + Simp[(C*m + A*(m + 1))/(b^2*m) Int[(b*Csc[e + f*x])^(m + 2), x], x] /; FreeQ[{b, e, f, A, C}, x] && NeQ[C*m + A*(m + 1), 0] && LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

```
rule 4562 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a
_)), x_Symbol] :> Simp[A*a*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*n)), x] + Si
mp[1/(d*n) Int[(d*Csc[e + f*x])^(n + 1)*Simp[n*(B*a + A*b) + (n*(a*C + B*
b) + A*a*(n + 1))*Csc[e + f*x] + b*C*n*Csc[e + f*x]^2, x], x], x] /; FreeQ[
{a, b, d, e, f, A, B, C}, x] && LtQ[n, -1]
```

3.566.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 744 vs. $2(319) = 638$.

Time = 18.82 (sec) , antiderivative size = 745, normalized size of antiderivative = 2.53

method	result	size
default	Expression too large to display	745
parts	Expression too large to display	971

```
input int((a+cos(d*x+c)*b)^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -2/315*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1120*B*co
s(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^10*b^3+(720*A*b^3+2160*B*a*b^2+2240*B*
b^3)*sin(1/2*d*x+1/2*c)^8*cos(1/2*d*x+1/2*c)+(-1512*A*a*b^2-1080*A*b^3-151
2*B*a^2*b-3240*B*a*b^2-2072*B*b^3)*sin(1/2*d*x+1/2*c)^6*cos(1/2*d*x+1/2*c)
+(1260*A*a^2*b+1512*A*a*b^2+840*A*b^3+420*B*a^3+1512*B*a^2*b+2520*B*a*b^2+
952*B*b^3)*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+(-630*A*a^2*b-378*A*a*b
^2-240*A*b^3-210*B*a^3-378*B*a^2*b-720*B*a*b^2-168*B*b^3)*sin(1/2*d*x+1/2*
c)^2*cos(1/2*d*x+1/2*c)+315*A*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/
2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+75*A*b^3*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))-315*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^3-567*A*(sin(1/2*
d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x
+1/2*c),2^(1/2))*a*b^2+105*B*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d
*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+225*B*a*b^2*(si
n(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1
/2*d*x+1/2*c),2^(1/2))-567*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1
/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2*b-147*B*(sin(1/
2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d
*x+1/2*c),2^(1/2))*b^3)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^...
```

3.566.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.13

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$15 \sqrt{2} (7i Ba^3 + 21i Aa^2b + 15i Bab^2 + 5i Ab^3) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/315*(15*sqrt(2)*(7*I*B*a^3 + 21*I*A*a^2*b + 15*I*B*a*b^2 + 5*I*A*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 15*sqrt(2)*(-7*I*B*a^3 - 21*I*A*a^2*b - 15*I*B*a*b^2 - 5*I*A*b^3)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 21*sqrt(2)*(-15*I*A*a^3 - 27*I*B*a^2*b - 27*I*A*a*b^2 - 7*I*B*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*sqrt(2)*(15*I*A*a^3 + 27*I*B*a^2*b + 27*I*A*a*b^2 + 7*I*B*b^3)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) - 2*(35*B*b^3*cos(d*x + c)^4 + 45*(3*B*a*b^2 + A*b^3)*cos(d*x + c)^3 + 7*(27*B*a^2*b + 27*A*a*b^2 + 7*B*b^3)*cos(d*x + c)^2 + 15*(7*B*a^3 + 21*A*a^2*b + 15*B*a*b^2 + 5*A*b^3)*cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.566.6 Sympy [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^3}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((a+b*cos(d*x+c))**3*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*(a + b*cos(c + d*x))**3/sqrt(sec(c + d*x)), x)`

3.566.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/sqrt(sec(d*x + c)), x)`

3.566.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^3}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^3*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^3/sqrt(sec(d*x + c)), x)`

3.566.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^3 (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^3}{\sqrt{\frac{1}{\cos(c+dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^3)/(1/cos(c + d*x))^(1/2),x)`

3.566. $\int \frac{(a+b \cos(c+dx))^3(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

3.567
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$$

3.567.1 Optimal result 5154
 3.567.2 Mathematica [A] (verified) 5155
 3.567.3 Rubi [A] (verified) 5155
 3.567.4 Maple [A] (verified) 5161
 3.567.5 Fricas [F(-1)] 5162
 3.567.6 Sympy [F(-1)] 5162
 3.567.7 Maxima [F] 5162
 3.567.8 Giac [F] 5163
 3.567.9 Mupad [F(-1)] 5163

3.567.1 Optimal result

Integrand size = 33, antiderivative size = 210

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx \\ &= \frac{2(Ab - aB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{a^2 d} \\ &+ \frac{2A \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3ad} \\ &+ \frac{2b(Ab - aB) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{a^2(a + b)d} \\ &- \frac{2(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{a^2 d} + \frac{2A \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3ad} \end{aligned}$$

```
output 2/3*A*sec(d*x+c)^(3/2)*sin(d*x+c)/a/d-2*(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/d+2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/d+2/3*A*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d+2*b*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/(a+b)/d
```

3.567.2 Mathematica [A] (verified)

Time = 9.76 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.07

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx =$$

$$\frac{\cot(c + dx) \left(-a^2 A \sec^{\frac{5}{2}}(c + dx) + a^2 A \cos(2(c + dx)) \sec^{\frac{5}{2}}(c + dx) - 6a(-Ab + aB) E \left(\arcsin \left(\sqrt{\sec(c + dx)} \right) \right) \right)}{a^3 d}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x]),x]`

output `-1/3*(Cot[c + d*x]*(-(a^2*A*Sec[c + d*x]^(5/2)) + a^2*A*Cos[2*(c + d*x)]*Sec[c + d*x]^(5/2) - 6*a*(-(A*b) + a*B)*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2] - 2*(3*A*b^2 + a^2*(A - 3*B) + 3*a*b*(A - B))*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2] + 6*A*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2] - 6*a*b*B*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2]))/(a^3*d)`

3.567.3 Rubi [A] (verified)Time = 1.97 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.07, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4521, 27, 3042, 4590, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{a + b \cos(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(c + dx + \frac{\pi}{2})^{\frac{5}{2}}(A + B \sin(c + dx + \frac{\pi}{2}))}{a + b \sin(c + dx + \frac{\pi}{2})} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A \sec(c + dx) + B)}{a \sec(c + dx) + b} dx$$

3.567. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{a\csc\left(c+dx+\frac{\pi}{2}\right)+b} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\sec(c+dx)}(-3(Ab-aB)\sec^2(c+dx)+aA\sec(c+dx)+Ab)}{2(b+a\sec(c+dx))} dx + \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{4521} \\
& \int \frac{\sqrt{\sec(c+dx)}(-3(Ab-aB)\sec^2(c+dx)+aA\sec(c+dx)+Ab)}{b+a\sec(c+dx)} dx + \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{27} \\
& \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(-3(Ab-aB)\csc(c+dx+\frac{\pi}{2})^2+aA\csc(c+dx+\frac{\pi}{2})+Ab)}{b+a\csc(c+dx+\frac{\pi}{2})} dx + \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(-3(Ab-aB)\csc(c+dx+\frac{\pi}{2})^2+aA\csc(c+dx+\frac{\pi}{2})+Ab)}{b+a\csc(c+dx+\frac{\pi}{2})} dx + \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{4590} \\
& \frac{2\int \frac{(Aa^2-3bBa+3Ab^2)\sec^2(c+dx)+a(4Ab-3aB)\sec(c+dx)+3b(Ab-aB)}{2\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx - \frac{6(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad}}{3a} + \\
& \quad \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{27} \\
& \frac{2\int \frac{(Aa^2-3bBa+3Ab^2)\sec^2(c+dx)+a(4Ab-3aB)\sec(c+dx)+3b(Ab-aB)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx - \frac{6(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad}}{3a} + \\
& \quad \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(Aa^2-3bBa+3Ab^2)\csc(c+dx+\frac{\pi}{2})^2+a(4Ab-3aB)\csc(c+dx+\frac{\pi}{2})+3b(Ab-aB)}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(b+a\csc(c+dx+\frac{\pi}{2}))} dx - \frac{6(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad}}{3a} + \\
& \quad \frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow \text{4594}
\end{aligned}$$

3.567. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{a+b\cos(c+dx)} dx$

$$\begin{aligned}
 & \frac{\int \frac{3(Ab-aB)b^2+aA\sec(c+dx)b^2}{\sqrt{\sec(c+dx)}} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a\sec(c+dx)} dx}{a} - \frac{6(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} \\
 & \frac{3a}{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)} \\
 & \frac{3ad}{3042} \\
 & \frac{\int \frac{3(Ab-aB)b^2+aA\csc(c+dx+\frac{\pi}{2})b^2}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a\csc(c+dx+\frac{\pi}{2})} dx}{a} - \frac{6(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} \\
 & \frac{3a}{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)} \\
 & \frac{3ad}{4274} \\
 & \frac{3b^2(Ab-aB) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + aAb^2 \int \sqrt{\sec(c+dx)} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a\csc(c+dx+\frac{\pi}{2})} dx}{a} - \frac{6(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} \\
 & \frac{3a}{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)} \\
 & \frac{3ad}{3042} \\
 & \frac{3b^2(Ab-aB) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + aAb^2 \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a\csc(c+dx+\frac{\pi}{2})} dx}{a} - \frac{6(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} \\
 & \frac{3a}{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)} \\
 & \frac{3ad}{4258} \\
 & \frac{3b^2(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + aAb^2 \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a\csc(c+dx+\frac{\pi}{2})} dx}{a} - \frac{6(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} \\
 & \frac{3a}{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)} \\
 & \frac{3ad}{3042}
 \end{aligned}$$

3.567. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

$$\frac{3b^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}dx+aAb^2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx}{b^2}+3b(Ab-aB)\int\frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}}{b+a\csc\left(c+dx+\frac{\pi}{2}\right)}dx$$

$$\frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad}$$

3119

$$\frac{aAb^2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}dx+\frac{6b^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d}}{b^2}+3b(Ab-aB)\int\frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}}{b+a\csc\left(c+dx+\frac{\pi}{2}\right)}dx$$

$$\frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad}$$

3120

$$3b(Ab-aB)\int\frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}}{b+a\csc\left(c+dx+\frac{\pi}{2}\right)}dx+\frac{6b^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d}+\frac{2aAb^2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d}$$

$$\frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad}$$

4336

$$3b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\cos(c+dx)(a+b\cos(c+dx))}}dx+\frac{6b^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d}+\frac{2aAb^2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}}{b^2}$$

$$\frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad}$$

3042

$$3b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b\sin\left(c+dx+\frac{\pi}{2}\right))}}dx+\frac{6b^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d}+\frac{2aAb^2\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2A\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3ad}$$

3284

3.567. $\int\frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{a+b\cos(c+dx)}dx$

$$\frac{\frac{6b^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2aAb^2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d}}{b^2} + \frac{6b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticPi}\left(\frac{2b}{a+b}\right)}{d(a+b)}}{a} = \frac{2A \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3ad}$$

3a

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x]),x]`

output `(2*A*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*a*d) + (((6*b^2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*A*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (6*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/((a + b)*d))/a - (6*(A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*d))/(3*a)`

3.567.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

3.567. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^(m_.))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4521 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[(-B)*d^2*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 2)/(b*f*(m + n))), x] + Simp[d^2/(b*(m + n)) Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n - 2)*Simp[a*B*(n - 2) + B*b*(m + n - 1)*Csc[e + f*x] + (A*b*(m + n) - a*B*(n - 1))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, m}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[n, 1] && NeQ[m + n, 0] && !IGtQ[m, 1]`

rule 4590 `Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Simp[(-C)*d*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 1)/(b*f*(m + n + 1))), x] + Simp[d/(b*(m + n + 1)) Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n - 1)*Simp[a*C*(n - 1) + (A*b*(m + n + 1) + b*C*(m + n))*Csc[e + f*x] + (b*B*(m + n + 1) - a*C*n)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, m}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 0]`

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))/ (Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.567.4 Maple [A] (verified)

Time = 20.24 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.10

method	result
default	$\frac{\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{2A\left(\frac{-\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}}{6\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^2} + \frac{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}\right)} + \frac{\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}}{a}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*b),x,method=_RETURNVER
BOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A/a*(-1/6*co
s(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(cos
(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+
1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ell
ipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2*(-A*b+B*a)/a^2/sin(1/2*d*x+1/2*c)^2/
(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1
/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))
)-4*(A*b-B*a)*b^2/a^2/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(
1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))/sin(1/2*d*x+1/2*c)
/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

$$3.567. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$$

3.567.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `Timed out`

3.567.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.567.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.567.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.567.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{a + b \cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x)),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x)), x)`

3.568
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$$

3.568.1 Optimal result 5164
 3.568.2 Mathematica [A] (verified) 5165
 3.568.3 Rubi [A] (verified) 5165
 3.568.4 Maple [A] (verified) 5169
 3.568.5 Fricas [F(-1)] 5169
 3.568.6 Sympy [F(-1)] 5170
 3.568.7 Maxima [F] 5170
 3.568.8 Giac [F] 5170
 3.568.9 Mupad [F(-1)] 5171

3.568.1 Optimal result

Integrand size = 33, antiderivative size = 126

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= -\frac{2A\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{ad}$$

$$- \frac{2(Ab - aB)\sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{a(a + b)d}$$

$$+ \frac{2A\sqrt{\sec(c + dx)} \sin(c + dx)}{ad}$$

```
output 2*A*sin(d*x+c)*sec(d*x+c)^(1/2)/a/d-2*A*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/d-2*(A*b-B*a)*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/(a+b)/d
```

3.568.2 Mathematica [A] (verified)

Time = 2.02 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.99

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \frac{2 \cos(2(c + dx)) \csc(c + dx) \left(aAE \left(\arcsin \left(\sqrt{\sec(c + dx)} \right) \middle| -1 \right) - (aA + Ab - aB) \text{EllipticF} \left(\arcsin \left(\sqrt{\sec(c + dx)} \right) \middle| -1 \right) \right)}{a^2 d (-1)}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x]),x]`

output `(-2*Cos[2*(c + d*x)]*Csc[c + d*x]*(a*A*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1] - (a*A + A*b - a*B)*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] + (A*b - a*B)*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sec[c + d*x]*Sqrt[-Tan[c + d*x]^2])/(a^2*d*(-2 + Sec[c + d*x]^2))`

3.568.3 Rubi [A] (verified)Time = 1.11 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 3439, 3042, 4521, 27, 3042, 4594, 27, 3042, 4258, 3042, 3119, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{a + b \cos(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{a + b \sin(c + dx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{3439} \\ & \int \frac{\sec^{\frac{3}{2}}(c + dx)(A \sec(c + dx) + B)}{a \sec(c + dx) + b} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.568. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2} (A \csc(c+dx+\frac{\pi}{2}) + B)}{a \csc(c+dx+\frac{\pi}{2}) + b} dx \\
& \quad \downarrow 4521 \\
& \frac{2 \int -\frac{(Ab-aB) \sec^2(c+dx) + aA \sec(c+dx) + Ab}{2\sqrt{\sec(c+dx)(b+a \sec(c+dx))}} dx}{a} + \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} \\
& \quad \downarrow 27 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{(Ab-aB) \sec^2(c+dx) + aA \sec(c+dx) + Ab}{\sqrt{\sec(c+dx)(b+a \sec(c+dx))}} dx}{a} \\
& \quad \downarrow 3042 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{(Ab-aB) \csc(c+dx+\frac{\pi}{2})^2 + aA \csc(c+dx+\frac{\pi}{2}) + Ab}{\sqrt{\csc(c+dx+\frac{\pi}{2})(b+a \csc(c+dx+\frac{\pi}{2}))}} dx}{a} \\
& \quad \downarrow 4594 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a \sec(c+dx)} dx + \int \frac{Ab^2}{\sqrt{\sec(c+dx)}} dx}{a} \\
& \quad \downarrow 27 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a \sec(c+dx)} dx + A \int \frac{1}{\sqrt{\sec(c+dx)}} dx}{a} \\
& \quad \downarrow 3042 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + A \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{a} \\
& \quad \downarrow 4258 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{a} \\
& \quad \downarrow 3042 \\
& \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{a}
\end{aligned}$$

3.568. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3119} \\
 & \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{2A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d}}{a} \\
 & \downarrow \text{4336} \\
 & \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx + \frac{2A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d}}{a} \\
 & \downarrow \text{3042} \\
 & \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))} dx + \frac{2A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d}}{a} \\
 & \downarrow \text{3284} \\
 & \frac{2A \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{2(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{d(a+b)} + \frac{2A \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d}}{a}
 \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x]),x]`

output `-(((2*A*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/((a + b)*d))/a + (2*A*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*d)`

3.568.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.568. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4521 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-B)*d^2*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 2)/(b*f*(m + n))), x] + Simp[d^2/(b*(m + n)) Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n - 2)*Simp[a*B*(n - 2) + B*b*(m + n - 1)*Csc[e + f*x] + (A*b*(m + n) - a*B*(n - 1))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, m}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[n, 1] && NeQ[m + n, 0] && !IGtQ[m, 1]`

rule 4594 `Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e + f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]`

3.568.4 Maple [A] (verified)

Time = 5.45 (sec) , antiderivative size = 300, normalized size of antiderivative = 2.38

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1}(\sin^2(\frac{dx}{2} + \frac{c}{2}))}{\sin(\frac{dx}{2} + \frac{c}{2})} \left(\frac{2A\sqrt{-2(\sin^4(\frac{dx}{2} + \frac{c}{2})) + \sin^2(\frac{dx}{2} + \frac{c}{2})} (2(\sin^2(\frac{dx}{2} + \frac{c}{2})) \cos(\frac{dx}{2} + \frac{c}{2}) - \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}})}{a \sin(\frac{dx}{2} + \frac{c}{2})^2 (2(\sin^2(\frac{dx}{2} + \frac{c}{2})) - 1)} \right)$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^(1/2)*(2*A/a/sin(1/2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))-4*(-A*b+B*a)/a/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.568.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

3.568. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

output Timed out

3.568.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c)),x)`

output Timed out

3.568.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.568.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.568. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

3.568.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{3/2}}{a + b \cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x)),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x)), x)`

3.569
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{a+b \cos(c+dx)} dx$$

3.569.1 Optimal result 5172
 3.569.2 Mathematica [A] (verified) 5172
 3.569.3 Rubi [A] (verified) 5173
 3.569.4 Maple [A] (verified) 5176
 3.569.5 Fracas [F(-1)] 5176
 3.569.6 Sympy [F] 5177
 3.569.7 Maxima [F] 5177
 3.569.8 Giac [F] 5177
 3.569.9 Mupad [F(-1)] 5178

3.569.1 Optimal result

Integrand size = 33, antiderivative size = 101

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

$$= \frac{2B\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{bd} + \frac{2(Ab - aB)\sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{b(a + b)d}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+
1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b/d+2*(A*b-B*a)*(cos(1/2
*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/
(a+b),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b/(a+b)/d
```

3.569.2 Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.75

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

$$= \frac{2 \cot(c + dx) \left(Ab \operatorname{EllipticF}\left(\arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) + (-Ab + aB) \operatorname{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c + dx)}\right)\right) \right)}{abd}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x]),x]`

output `(2*Cot[c + d*x]*(A*b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] + -(A*b) + a*B)*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2])/(a*b*d)`

3.569.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3439, 3042, 4526, 3042, 4258, 3042, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{a+b\cos(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{a+b\sin(c+dx+\frac{\pi}{2})} dx \\ & \quad \downarrow \text{3439} \\ & \int \frac{\sqrt{\sec(c+dx)}(A\sec(c+dx)+B)}{a\sec(c+dx)+b} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A\csc(c+dx+\frac{\pi}{2})+B)}{a\csc(c+dx+\frac{\pi}{2})+b} dx \\ & \quad \downarrow \text{4526} \\ & \frac{(Ab-aB) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a\sec(c+dx)} dx}{b} + \frac{B \int \sqrt{\sec(c+dx)} dx}{b} \\ & \quad \downarrow \text{3042} \\ & \frac{(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a\csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{b} \end{aligned}$$

$$\begin{aligned}
& \downarrow 4258 \\
& \frac{(Ab - aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b} \\
& \downarrow 3120 \\
& \frac{(Ab - aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{bd} \\
& \downarrow 4336 \\
& \frac{(Ab - aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx}{b} + \\
& \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{bd} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx}{b} + \\
& \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{bd} \\
& \downarrow 3284 \\
& \frac{2(Ab - aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{bd(a+b)} + \\
& \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{bd}
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x]),x]`

output `(2*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b*d) + (2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b*(a + b)*d)`

3.569.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4526 `Int[((csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[A/a Int[(d*Csc[e + f*x])^n, x], x] - Simp[(A*b - a*B)/(a*d) Int[(d*Csc[e + f*x])^(n + 1)/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0]`

3.569.4 Maple [A] (verified)

Time = 3.76 (sec) , antiderivative size = 217, normalized size of antiderivative = 2.15

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}\left(\text{A}\Pi\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),-\frac{2b}{a-b},\sqrt{2}\right)b+BF\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),2^{\frac{1}{2}}\right)a-B*E\right)}{(a-b)b\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `-2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*(A*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*b+B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a-B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*b-B*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a)/(a-b)/b/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.569.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="fricas")`

output `Timed out`

3.569.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c)),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/(a + b*cos(c + d*x)), x)`

3.569.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{b \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a), x)`

3.569.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{b \cos(dx + c) + a} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a), x)`

3.569.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{a + b \cos(c + dx)} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x)),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x)), x)`

3.570
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))\sqrt{\sec(c+dx)}} dx$$

3.570.1 Optimal result 5179
 3.570.2 Mathematica [A] (verified) 5180
 3.570.3 Rubi [A] (verified) 5180
 3.570.4 Maple [A] (verified) 5184
 3.570.5 Fricas [F(-1)] 5185
 3.570.6 Sympy [F] 5185
 3.570.7 Maxima [F] 5185
 3.570.8 Giac [F] 5186
 3.570.9 Mupad [F(-1)] 5186

3.570.1 Optimal result

Integrand size = 33, antiderivative size = 149

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx \\ &= \frac{2B\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{\sec(c + dx)}}{bd} \\ & \quad + \frac{2(Ab - aB)\sqrt{\cos(c + dx)}\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{b^2d} \\ & \quad - \frac{2a(Ab - aB)\sqrt{\cos(c + dx)}\text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)\sqrt{\sec(c + dx)}}{b^2(a + b)d} \end{aligned}$$

```
output 2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+
1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b/d+2*(A*b-B*a)*(cos(1/2
*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/
2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^2/d-2*a*(A*b-B*a)*(cos(1/2*d*x+1/2
*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^
(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^2/(a+b)/d
```

3.570.2 Mathematica [A] (verified)

Time = 19.91 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.48

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

$$= \frac{\cot(c + dx) \left(-bB \sec^{\frac{3}{2}}(c + dx) - bB \cos(2(c + dx)) \sec^{\frac{3}{2}}(c + dx) + bB \sec^{\frac{7}{2}}(c + dx) + bB \cos(2(c + dx)) \right)}{b^2 d}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]),x]`

output `(Cot[c + d*x]*(-(b*B*Sec[c + d*x]^(3/2)) - b*B*Cos[2*(c + d*x)]*Sec[c + d*x]^(3/2) + b*B*Sec[c + d*x]^(7/2) + b*B*Cos[2*(c + d*x)]*Sec[c + d*x]^(7/2)) - 2*b*B*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2] + 2*b*B*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2] + 2*A*b*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2] - 2*a*B*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[-Tan[c + d*x]^2]))/(b^2*d)`

3.570.3 Rubi [A] (verified)Time = 1.00 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.85, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4526, 3042, 4258, 3042, 3119, 4335, 3042, 3282, 3042, 3120, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx$$

$$\downarrow \text{3439}$$

$$\int \frac{A \sec(c + dx) + B}{\sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{A \csc(c + dx + \frac{\pi}{2}) + B}{\sqrt{\csc(c + dx + \frac{\pi}{2})} (a \csc(c + dx + \frac{\pi}{2}) + b)} dx \\
& \downarrow 4526 \\
& \frac{(Ab - aB) \int \frac{\sqrt{\sec(c+dx)}}{b+a \sec(c+dx)} dx}{b} + \frac{B \int \frac{1}{\sqrt{\sec(c+dx)}} dx}{b} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b} \\
& \downarrow 4258 \\
& \frac{(Ab - aB) \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b} \\
& \downarrow 3119 \\
& \frac{(Ab - aB) \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b} + \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{bd} \\
& \downarrow 4335 \\
& \frac{(Ab - aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{\cos(c+dx)}}{a+b \cos(c+dx)} dx}{b} + \\
& \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{bd} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{a+b \sin(c+dx+\frac{\pi}{2})} dx}{b} + \\
& \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{bd} \\
& \downarrow 3282
\end{aligned}$$

3.570. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
 & \frac{(Ab - aB)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}\left(\frac{\int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx}{b}\right)}{2B\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)} + \\
 & \qquad \qquad \qquad \frac{b}{bd} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{(Ab - aB)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}\left(\frac{\int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{b} - \frac{a \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b \sin\left(c+dx+\frac{\pi}{2}\right))}} dx}{b}\right)}{2B\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)} + \\
 & \qquad \qquad \qquad \frac{b}{bd} \\
 & \qquad \qquad \qquad \downarrow \text{3120} \\
 & \frac{(Ab - aB)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}\left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{a \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)(a+b \sin\left(c+dx+\frac{\pi}{2}\right))}} dx}{b}\right)}{2B\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)} + \\
 & \qquad \qquad \qquad \frac{b}{bd} \\
 & \qquad \qquad \qquad \downarrow \text{3284} \\
 & \frac{(Ab - aB)\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}\left(\frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd} - \frac{2a \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)}{bd(a+b)}\right)}{2B\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)} + \\
 & \qquad \qquad \qquad \frac{b}{bd}
 \end{aligned}$$

```
input Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]),x]
```

```
output (2*B*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]]/(b*d) + ((A*b - a*B)*Sqrt[Cos[c + d*x]]*((2*EllipticF[(c + d*x)/2, 2])/(b*d) - (2*a*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2])/(b*(a + b)*d))*Sqrt[Sec[c + d*x]])/b
```

3.570. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))\sqrt{\sec(c+dx)}} dx$

3.570.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3282 `Int[Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d/b Int[1/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[(b*c - a*d)/b Int[1/((a + b*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^((p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^((n_.), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4335 `Int[Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[Sqrt[d*Sin[e + f*x]]*(Sqrt[d*Csc[e + f*x]]/d) Int[Sqrt[d*Sin[e + f*x]]/(b + a*Sin[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4526 `Int[((csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[A/a Int[(d*Csc[e + f*x])^n, x], x] - Simp[(A*b - a*B)/(a*d) Int[(d*Csc[e + f*x])^(n + 1)/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0]`

3.570.4 Maple [A] (verified)

Time = 5.42 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.98

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}}\left(AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)ab - AF\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{-2}\right)b^2(a-b)\right)\sqrt{-2}$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*(A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a*b-A*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*b^2-A*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a*b-B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a^2+B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*a*b-B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b+B*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^2+B*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a^2)/b^2/(a-b)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.570.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))\sqrt{\sec(c+dx)}} dx$$

3.570.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `Timed out`

3.570.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/((a + b*cos(c + d*x))*sqrt(sec(c + d*x))), x)`

3.570.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)\sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.570.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)\sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.570.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + b \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))), x)`

$$3.571 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$$

3.571.1 Optimal result	5187
3.571.2 Mathematica [A] (verified)	5188
3.571.3 Rubi [A] (verified)	5188
3.571.4 Maple [B] (verified)	5193
3.571.5 Fracas [F(-1)]	5194
3.571.6 Sympy [F]	5195
3.571.7 Maxima [F]	5195
3.571.8 Giac [F]	5195
3.571.9 Mupad [F(-1)]	5196

3.571.1 Optimal result

Integrand size = 33, antiderivative size = 197

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx \\ &= \frac{2(Ab - aB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{b^2 d} \\ & \quad - \frac{2(3aAb - 3a^2 B - b^2 B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3b^3 d} \\ & \quad + \frac{2a^2 (Ab - aB) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{b^3 (a + b) d} \\ & \quad + \frac{2B \sin(c + dx)}{3bd \sqrt{\sec(c + dx)}} \end{aligned}$$

```
output 2/3*B*sin(d*x+c)/b/d/sec(d*x+c)^(1/2)+2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^2/d-2/3*(3*A*a*b-3*B*a^2-B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^3/d+2*a^2*(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^3/(a+b)/d
```

3.571.2 Mathematica [A] (verified)

Time = 5.01 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.41

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2 \csc(c + dx) \left(-3Ab^2 + 3abB + 3Ab^2 \sec^2(c + dx) - 3abB \sec^2(c + dx) + b^2 B \sin(c + dx) \tan(c + dx) - \dots \right)}{\dots}$$

```
input Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sec[c + d*x]^(3/2)),x
]
```

```
output (2*Csc[c + d*x]*(-3*A*b^2 + 3*a*b*B + 3*A*b^2*Sec[c + d*x]^2 - 3*a*b*B*Sec
[c + d*x]^2 + b^2*B*Sin[c + d*x]*Tan[c + d*x] - 3*b*(A*b - a*B)*EllipticE[
ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] +
b*(3*A*b - 3*a*B + b*B)*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Se
c[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 3*a*A*b*EllipticPi[-(a/b), ArcSin[Sqrt
[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + 3*a^2*B*El
lipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-
Tan[c + d*x]^2]))/(3*b^3*d*Sec[c + d*x]^(3/2))
```

3.571.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.03, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4522, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx$$

↓ 3439

3.571. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A \sec(c+dx) + B}{\sec^{\frac{3}{2}}(c+dx)(a \sec(c+dx) + b)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A \csc(c+dx + \frac{\pi}{2}) + B}{\csc(c+dx + \frac{\pi}{2})^{\frac{3}{2}}(a \csc(c+dx + \frac{\pi}{2}) + b)} dx \\
& \quad \downarrow \text{4522} \\
& \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}} - \frac{2 \int -\frac{aB \sec^2(c+dx) + bB \sec(c+dx) + 3(Ab-aB)}{2\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx}{3b} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{aB \sec^2(c+dx) + bB \sec(c+dx) + 3(Ab-aB)}{\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx}{3b} + \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{aB \csc(c+dx + \frac{\pi}{2})^2 + bB \csc(c+dx + \frac{\pi}{2}) + 3(Ab-aB)}{\sqrt{\csc(c+dx + \frac{\pi}{2})}(b+a \csc(c+dx + \frac{\pi}{2}))} dx}{3b} + \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}} \\
& \quad \downarrow \text{4594} \\
& \frac{3a^2(Ab-aB) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a \sec(c+dx)} dx}{b^2} + \frac{\int \frac{3b(Ab-aB) - (-3Ba^2 + 3Aba - b^2B) \sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{b^2} + \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{3a^2(Ab-aB) \int \frac{\csc(c+dx + \frac{\pi}{2})^{\frac{3}{2}}}{b+a \csc(c+dx + \frac{\pi}{2})} dx}{b^2} + \frac{\int \frac{3b(Ab-aB) + (3Ba^2 - 3Aba + b^2B) \csc(c+dx + \frac{\pi}{2})}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx}{b^2} + \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}} \\
& \quad \downarrow \text{4274} \\
& \frac{3a^2(Ab-aB) \int \frac{\csc(c+dx + \frac{\pi}{2})^{\frac{3}{2}}}{b+a \csc(c+dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{1}{\sqrt{\sec(c+dx)}} dx - (-3a^2B + 3aAb - b^2B) \int \sqrt{\sec(c+dx)} dx}{b^2} + \\
& \quad \frac{3b}{3bd\sqrt{\sec(c+dx)}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.571. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

$$\frac{3a^2(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{3b(Ab-aB) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - (-3a^2B+3aAb-b^2B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{b^2} +$$

$$\frac{3b}{3bd\sqrt{\sec(c+dx)}} \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 4258

$$\frac{3a^2(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{3b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - (-3a^2B+3aAb-b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b^2} +$$

$$\frac{3b}{3bd\sqrt{\sec(c+dx)}} \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 3042

$$\frac{3a^2(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{3b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - (-3a^2B+3aAb-b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^2} +$$

$$\frac{3b}{3bd\sqrt{\sec(c+dx)}} \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 3119

$$\frac{3a^2(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{6b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d} - (-3a^2B+3aAb-b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{b^2} +$$

$$\frac{3b}{3bd\sqrt{\sec(c+dx)}} \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 3120

$$\frac{3a^2(Ab-aB) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2} + \frac{6b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d} - \frac{2(-3a^2B+3aAb-b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{1}{2}(c+dx+\frac{\pi}{2})|2)}{b^2 d} +$$

$$\frac{3b}{3bd\sqrt{\sec(c+dx)}} \frac{2B \sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 4336

3.571. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)} dx$

$$\frac{3a^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))}dx}{b^2} + \frac{6b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d} - \frac{2(-3a^2B+3aAb-b^2B)\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2B\sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 3042

$$\frac{3a^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}(a+b\sin\left(c+dx+\frac{\pi}{2}\right))}dx}{b^2} + \frac{6b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d} - \frac{2(-3a^2B+3aAb-b^2B)\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2B\sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

↓ 3284

$$\frac{6a^2(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticPi}\left(\frac{2b}{a+b},\frac{1}{2}(c+dx),2\right)}{b^2d(a+b)} + \frac{6b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)|2\right)}{d} - \frac{2(-3a^2B+3aAb-b^2B)\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2B\sin(c+dx)}{3bd\sqrt{\sec(c+dx)}}$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sec[c + d*x]^(3/2)),x]`

output `((6*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*(3*a*A*b - 3*a^2*B - b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (6*a^2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d)/(3*b) + (2*B*Sin[c + d*x])/(3*b*d*Sqrt[Sec[c + d*x]])`

3.571.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.571. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)} dx$

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3284 $\text{Int}[1/(((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)])*\text{Sqrt}[(c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)])), x_Symbol] \rightarrow \text{Simp}[(2/(f*(a + b)*\text{Sqrt}[c + d]))*\text{EllipticPi}[2*(b/(a + b)), (1/2)*(e - \text{Pi}/2 + f*x), 2*(d/(c + d))], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[c + d, 0]$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(g_.))^{\text{p}_.}*((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)])^{\text{m}_.}*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)])^{\text{n}_.}, x_Symbol] \rightarrow \text{Simp}[g^{\text{m} + \text{n}} \text{Int}[(g*\text{Csc}[e + f*x])^{\text{p} - \text{m} - \text{n}}*(b + a*\text{Csc}[e + f*x])^{\text{m}}*(d + c*\text{Csc}[e + f*x])^{\text{n}}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{!IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{\text{n}_.}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{\text{n}}*\text{Sin}[c + d*x]^{\text{n}} \text{Int}[1/\text{Sin}[c + d*x]^{\text{n}}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{\text{n}_.}*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^{\text{n}}, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{\text{n} + 1}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4336 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{\text{3}/2}/(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[d*\text{Sqrt}[d*\text{Sin}[e + f*x]]*\text{Sqrt}[d*\text{Csc}[e + f*x]] \text{Int}[1/(\text{Sqrt}[d*\text{Sin}[e + f*x]]*(b + a*\text{Sin}[e + f*x])), x], x] /; \text{FreeQ}[\{a, b, d, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

```
rule 4522 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[A*Cot[e
+ f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*n)), x] + Sim
p[1/(a*d*n) Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n + 1)*Simp[a*B*
n - A*b*(m + n + 1) + A*a*(n + 1)*Csc[e + f*x] + A*b*(m + n + 2)*Csc[e + f*
x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, m}, x] && NeQ[A*b - a*B, 0]
&& NeQ[a^2 - b^2, 0] && LeQ[n, -1]
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/ (Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.571.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 821 vs. $2(257) = 514$.

Time = 6.48 (sec) , antiderivative size = 822, normalized size of antiderivative = 4.17

method	result	size
default	Expression too large to display	822

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)/sec(d*x+c)^(3/2),x,method=_RETURNVER
BOSE)
```


output

```
-2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*a*b^2-4*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^4*b^3+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a^2*b-3*A*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*A*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a*b^2+3*A*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*b^3-2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*a*b^2+2*B*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^2*b^3-3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))*a^3+3*B*a^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-3*B*a^2*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+B*a*b^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-B*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+3*B*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*a^2...
```

3.571.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fracas")`

output `Timed out`

3.571.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `Integral((A + B*cos(c + d*x))/((a + b*cos(c + d*x))*sec(c + d*x)**(3/2)),
x)`

3.571.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm=
"maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)),
x)`

3.571.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm=
"giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)),
x)`

3.571.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} (a + b \cos(c + dx))} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))),x)`output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))), x)`

3.572
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.572.1 Optimal result 5197
 3.572.2 Mathematica [A] (warning: unable to verify) 5198
 3.572.3 Rubi [A] (verified) 5199
 3.572.4 Maple [B] (verified) 5206
 3.572.5 Fricas [F(-1)] 5207
 3.572.6 Sympy [F(-1)] 5207
 3.572.7 Maxima [F(-1)] 5207
 3.572.8 Giac [F] 5208
 3.572.9 Mupad [F(-1)] 5208

3.572.1 Optimal result

Integrand size = 33, antiderivative size = 405

$$\begin{aligned} & \int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx \\ &= \frac{(4a^2Ab - 5Ab^3 - 2a^3B + 3ab^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{a^3(a^2 - b^2)d} \\ &+ \frac{(2a^2A - 5Ab^2 + 3abB) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3a^2(a^2 - b^2)d} \\ &+ \frac{b(7a^2Ab - 5Ab^3 - 5a^3B + 3ab^2B) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{a^3(a - b)(a + b)^2d} \\ &- \frac{(4a^2Ab - 5Ab^3 - 2a^3B + 3ab^2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{a^3(a^2 - b^2)d} \\ &+ \frac{(2a^2A - 5Ab^2 + 3abB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3a^2(a^2 - b^2)d} + \frac{b(Ab - aB) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{a(a^2 - b^2)d(b + a \sec(c + dx))} \end{aligned}$$

output $\frac{1}{3}(2Aa^2-5Ab^2+3Bab)\sec(dx+c)^{3/2}\sin(dx+c)/a^2/(a^2-b^2)/d+b(Ab-Ba)\sec(dx+c)^{5/2}\sin(dx+c)/a/(a^2-b^2)/d/(b+a\sec(dx+c))-(4Aa^2b-5Ab^3-2Ba^3+3Bab^2)\sin(dx+c)\sec(dx+c)^{1/2}/a^3/(a^2-b^2)/d+(4Aa^2b-5Ab^3-2Ba^3+3Bab^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticE}(\sin(1/2dx+1/2c),2^{1/2})*\cos(dx+c)^{1/2}\sec(dx+c)^{1/2}/a^3/(a^2-b^2)/d+1/3(2Aa^2-5Ab^2+3Bab)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticF}(\sin(1/2dx+1/2c),2^{1/2})*\cos(dx+c)^{1/2}\sec(dx+c)^{1/2}/a^2/(a^2-b^2)/d+b(7Aa^2b-5Ab^3-5Ba^3+3Bab^2)(\cos(1/2dx+1/2c))^2)^{1/2}/\cos(1/2dx+1/2c)*\text{EllipticPi}(\sin(1/2dx+1/2c),2b/(a+b),2^{1/2})*\cos(dx+c)^{1/2}\sec(dx+c)^{1/2}/a^3/(a-b)/(a+b)^2/d$

3.572.2 Mathematica [A] (warning: unable to verify)

Time = 6.98 (sec) , antiderivative size = 735, normalized size of antiderivative = 1.81

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{2(-4a^4A - 44a^2Ab^2 + 45Ab^4 + 30a^3bB - 27ab^3B) \cos^2(c + dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) \right)}{a(a + b \cos(c + dx))(1 - \cos^2(c + dx))} + \frac{\sqrt{\sec(c + dx)} \left(\frac{(-4a^2Ab + 5Ab^3 + 2a^3B - 3ab^2B) \sin(c + dx)}{a^3(a^2 - b^2)} + \frac{-Ab^3 \sin(c + dx) + ab^2B \sin(c + dx)}{a^2(a^2 - b^2)(a + b \cos(c + dx))} + \frac{2A \tan(c + dx)}{3a^2} \right)}{d}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x])^2, x]`

output $((2*(-4*a^4*A - 44*a^2*A*b^2 + 45*A*b^4 + 30*a^3*b*B - 27*a*b^3*B)*\text{Cos}[c + d*x]^2*(\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1] - \text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1])*(b + a*\text{Sec}[c + d*x])* \text{Sqrt}[1 - \text{Sec}[c + d*x]^2]*\text{Sin}[c + d*x])/(a*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)) + (2*(-28*a^3*A*b + 40*a*A*b^3 + 12*a^4*B - 24*a^2*b^2*B)*\text{Cos}[c + d*x]^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*(b + a*\text{Sec}[c + d*x])* \text{Sqrt}[1 - \text{Sec}[c + d*x]^2]*\text{Sin}[c + d*x])/(b*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)) + ((-12*a^2*A*b^2 + 15*A*b^4 + 6*a^3*b*B - 9*a*b^3*B)*\text{Cos}[2*(c + d*x)]*(b + a*\text{Sec}[c + d*x])*(-4*a*b + 4*a*b*\text{Sec}[c + d*x]^2 - 4*a*b*\text{EllipticE}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] + 2*(2*a - b)*b*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] - 4*a^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] + 2*b^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2])* \text{Sin}[c + d*x])/(a*b^2*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)*\text{Sqrt}[\text{Sec}[c + d*x]]*(2 - \text{Sec}[c + d*x]^2)))/(12*a^3*(-a + b)*(a + b)*d + (\text{Sqrt}[\text{Sec}[c + d*x]]*((-4*a^2*A*b + 5*A*b^3 + 2*a^3*B - 3*a*b^2*B)*\text{Sin}[c + d*x])/(a^3*(a^2 - b^2)) + -(A*b^3*\text{Sin}[c + d*x]) + a*b^2*B*\text{Sin}[c + d*x])/(a^2*(a^2 - b^2))*(a + b*\text{Cos}[c + d*x])) + (2*A*\text{Tan}[c + d*x])/(3*a^2))/d$

3.572.3 Rubi [A] (verified)

Time = 3.15 (sec) , antiderivative size = 389, normalized size of antiderivative = 0.96, number of steps used = 23, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 3439, 3042, 4517, 27, 3042, 4590, 27, 3042, 4590, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx$$

↓ 3439

$$\int \frac{\sec^{\frac{7}{2}}(c+dx)(A\sec(c+dx)+B)}{(a\sec(c+dx)+b)^2} dx$$

3.572. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+b\cos(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{7/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+b\right)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \\
& \quad \downarrow \text{4517} \\
& \int \frac{-\frac{\sec^{\frac{3}{2}}(c+dx)\left((2Aa^2+3bBa-5Ab^2)\sec^2(c+dx)-2a(Ab-aB)\sec(c+dx)+3b(Ab-aB)\right)}{2(b+a\sec(c+dx))}}{a(a^2-b^2)} dx \\
& \quad \downarrow \text{27} \\
& \int \frac{\frac{\sec^{\frac{3}{2}}(c+dx)\left((2Aa^2+3bBa-5Ab^2)\sec^2(c+dx)-2a(Ab-aB)\sec(c+dx)+3b(Ab-aB)\right)}{b+a\sec(c+dx)}}{2a(a^2-b^2)} dx + \\
& \quad \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a\sec(c+dx)+b)} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left((2Aa^2+3bBa-5Ab^2)\csc\left(c+dx+\frac{\pi}{2}\right)^2-2a(Ab-aB)\csc\left(c+dx+\frac{\pi}{2}\right)+3b(Ab-aB)\right)}{b+a\csc\left(c+dx+\frac{\pi}{2}\right)}}{2a(a^2-b^2)} dx + \\
& \quad \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a\sec(c+dx)+b)} \\
& \quad \downarrow \text{4590} \\
& \frac{2\int \frac{\sqrt{\sec(c+dx)}\left(-3\left(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3\right)\sec^2(c+dx)+2a\left(Aa^2-3bBa+2Ab^2\right)\sec(c+dx)+b\left(2Aa^2+3bBa-5Ab^2\right)\right)}{2(b+a\sec(c+dx))}}{3a} dx + \frac{2(2a^2A+3abB-5Ab^2)\sin(c+dx)}{3ad} \\
& \quad \frac{2a(a^2-b^2)}{ad(a^2-b^2)(a\sec(c+dx)+b)} \\
& \quad \downarrow \text{27} \\
& \int \frac{\frac{\sqrt{\sec(c+dx)}\left(-3\left(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3\right)\sec^2(c+dx)+2a\left(Aa^2-3bBa+2Ab^2\right)\sec(c+dx)+b\left(2Aa^2+3bBa-5Ab^2\right)\right)}{b+a\sec(c+dx)}}{3a} dx + \frac{2(2a^2A+3abB-5Ab^2)\sin(c+dx)}{3ad} \\
& \quad \frac{2a(a^2-b^2)}{ad(a^2-b^2)(a\sec(c+dx)+b)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.572. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+b\cos(c+dx))^2} dx$

$$\int \frac{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(-3\left(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3\right)\csc\left(c+dx+\frac{\pi}{2}\right)^2+2a\left(Aa^2-3bBa+2Ab^2\right)\csc\left(c+dx+\frac{\pi}{2}\right)+b\left(2Aa^2+3bBa-5Ab^2\right)\right)dx}{b+a\csc\left(c+dx+\frac{\pi}{2}\right)} + \frac{2(2a^2A+3abB)}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)} \quad 2a(a^2 - b^2)$$

↓ 4590

$$2 \int \frac{\left(2Aa^4 - 12bBa^3 + 16Ab^2a^2 + 9b^3Ba - 15Ab^4\right) \sec^2(c+dx) + 2a\left(-3Ba^3 + 7Aba^2 + 6b^2Ba - 10Ab^3\right) \sec(c+dx) + 3b\left(-2Ba^3 + 4Aba^2 + 3b^2Ba - 5Ab^3\right)}{2\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx - \frac{6(-2a^3B + 4a^2Ab + 3ab^2B - 5Ab^3)}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)} \quad 2a(a^2 - b^2)$$

↓ 27

$$\int \frac{\left(2Aa^4 - 12bBa^3 + 16Ab^2a^2 + 9b^3Ba - 15Ab^4\right) \sec^2(c+dx) + 2a\left(-3Ba^3 + 7Aba^2 + 6b^2Ba - 10Ab^3\right) \sec(c+dx) + 3b\left(-2Ba^3 + 4Aba^2 + 3b^2Ba - 5Ab^3\right)}{\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx - \frac{6(-2a^3B + 4a^2Ab + 3ab^2B - 5Ab^3)}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)} \quad 2a(a^2 - b^2)$$

↓ 3042

$$\int \frac{\left(2Aa^4 - 12bBa^3 + 16Ab^2a^2 + 9b^3Ba - 15Ab^4\right) \csc\left(c+dx+\frac{\pi}{2}\right)^2 + 2a\left(-3Ba^3 + 7Aba^2 + 6b^2Ba - 10Ab^3\right) \csc\left(c+dx+\frac{\pi}{2}\right) + 3b\left(-2Ba^3 + 4Aba^2 + 3b^2Ba - 5Ab^3\right)}{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(b+a \csc\left(c+dx+\frac{\pi}{2}\right)\right)} dx - \frac{6(-2a^3B + 4a^2Ab + 3ab^2B - 5Ab^3)}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)} \quad 2a(a^2 - b^2)$$

↓ 4594

$$3b\left(-5a^3B + 7a^2Ab + 3ab^2B - 5Ab^3\right) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a \sec(c+dx)} dx + \int \frac{3\left(-2Ba^3 + 4Aba^2 + 3b^2Ba - 5Ab^3\right)b^2 + a\left(2Aa^2 + 3bBa - 5Ab^2\right) \sec(c+dx)b^2}{\sqrt{\sec(c+dx)}b^2} dx - \frac{6(-2a^3B + 4a^2Ab + 3ab^2B - 5Ab^3)}{3a}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)} \quad 2a(a^2 - b^2)$$

↓ 3042

3.572. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

$$\frac{3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \int \frac{3(-2Ba^3+4Aba^2+3b^2Ba-5Ab^3)b^2+a(2Aa^2+3bBa-5Ab^2) \csc(c+dx+\frac{\pi}{2})b^2 dx}{\sqrt{\csc(c+dx+\frac{\pi}{2})} b^2}}{a} \quad \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3)}{3a} \quad 2a(a^2-b^2)$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4274

$$\frac{3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{ab^2(2a^2A+3abB-5Ab^2) \int \sqrt{\sec(c+dx)} dx + 3b^2(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \frac{1}{\sqrt{\sec(c+dx)}} dx}{b^2}}{a} \quad \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3)}{3a} \quad 2a(a^2-b^2)$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

$$\frac{ab^2(2a^2A+3abB-5Ab^2) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx + 3b^2(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b^2} + \frac{3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{a} \quad \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3)}{3a} \quad 2a(a^2-b^2)$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4258

$$\frac{3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{ab^2(2a^2A+3abB-5Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + 3b^2(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2}}{a} \quad \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3)}{3a} \quad 2a(a^2-b^2)$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

$$\frac{3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{ab^2(2a^2A+3abB-5Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + 3b^2(-2a^3B+4a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2}}{a} \quad \frac{6(-2a^3B+4a^2Ab+3ab^2B-5Ab^3)}{3a} \quad 2a(a^2-b^2)$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

3.572. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

↓ 3119

$$3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{ab^2(2a^2A+3abB-5Ab^2)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{6b^2(-2a^3B+4a^2Ab+3ab^2)}{b^2}$$

a $3a$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 3120

$$3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{2ab^2(2a^2A+3abB-5Ab^2)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{6b^2(-2a^3B+4a^2Ab+3ab^2)}{b^2}$$

a $3a$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 4336

$$3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)(a+b \cos(c+dx))}} dx + \frac{2ab^2(2a^2A+3abB-5Ab^2)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}$$

a $3a$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 3042

$$3b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx + \frac{2ab^2(2a^2A+3abB-5Ab^2)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}$$

a $3a$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 3284

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{ad(a^2 - b^2)(a \sec(c + dx) + b)} + \frac{6b(-5a^3B+7a^2Ab+3ab^2B-5Ab^3)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \operatorname{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{d(a+b)} + \frac{2ab^2(2a^2A+3abB-5Ab^2)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \operatorname{EllipticF}(\frac{1}{2}(c+dx), 2)}{d}$$

$3ad$

3.572. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

input `Int[((A + B*cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*cos[c + d*x])^2,x]`

output `(b*(A*b - a*B)*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(a*(a^2 - b^2)*d*(b + a*Sec[c + d*x])) + ((2*(2*a^2*A - 5*A*b^2 + 3*a*b*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*a*d) + (((6*b^2*(4*a^2*A*b - 5*A*b^3 - 2*a^3*B + 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*b^2*(2*a^2*A - 5*A*b^2 + 3*a*b*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (6*b*(7*a^2*A*b - 5*A*b^3 - 5*a^3*B + 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/((a + b)*d)/a - (6*(4*a^2*A*b - 5*A*b^3 - 2*a^3*B + 3*a*b^2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*d)/(3*a)/(2*a*(a^2 - b^2))`

3.572.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(g_.)^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\sin[c + d*x]^n \text{Int}[1/\sin[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4336 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(3/2)}/(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[d*\text{Sqrt}[d*\sin[e + f*x]]*\text{Sqrt}[d*\text{Csc}[e + f*x]] \text{Int}[1/(\text{Sqrt}[d*\sin[e + f*x]]*(b + a*\sin[e + f*x])), x], x] /; \text{FreeQ}[\{a, b, d, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 4517 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[a*d^2*(A*b - a*B)*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m+1)}*((d*\text{Csc}[e + f*x])^{(n-2)})/(b*f*(m+1)*(a^2 - b^2)), x] - \text{Simp}[d/(b*(m+1)*(a^2 - b^2)) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m+1)}*(d*\text{Csc}[e + f*x])^{(n-2)}*\text{Simp}[a*d*(A*b - a*B)*(n-2) + b*d*(A*b - a*B)*(m+1)*\text{Csc}[e + f*x] - (a*A*b*d*(m+n) - d*B*(a^2*(n-1) + b^2*(m+1)))*\text{Csc}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -1] \&\& \text{GtQ}[n, 1]$

rule 4590 $\text{Int}[(A_.) + \text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + \text{csc}[(e_.) + (f_.)*(x_)]^2*(C_.)*(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(-C)*d*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m+1)}*((d*\text{Csc}[e + f*x])^{(n-1)})/(b*f*(m+n+1)), x] + \text{Simp}[d/(b*(m+n+1)) \text{Int}[(a + b*\text{Csc}[e + f*x])^m*(d*\text{Csc}[e + f*x])^{(n-1)}*\text{Simp}[a*C*(n-1) + (A*b*(m+n+1) + b*C*(m+n))*\text{Csc}[e + f*x] + (b*B*(m+n+1) - a*C*n)*\text{Csc}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B, C, m\}, x] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{GtQ}[n, 0]$

3.572. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))/ (Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.572.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1003 vs. $2(461) = 922$.

Time = 44.15 (sec) , antiderivative size = 1004, normalized size of antiderivative = 2.48

method	result	size
default	Expression too large to display	1004

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2/a^2*A*(-1/6*
cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(c
os(1/2*d*x+1/2*c)^2-1/2)^2+1/3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*
x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*E
llipticF(cos(1/2*d*x+1/2*c),2^(1/2)))+2*(-2*A*b+B*a)/a^3/sin(1/2*d*x+1/2*c
)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2)))-4*b^2*(2*A*b-B*a)/a^3/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c
)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+2*(A*b-B*a)*b
/a^2*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1
/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2
*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/
2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(
1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/
2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*c
os(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b...
```

$$3.572. \quad \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.572.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorithm
m="fricas")
```

```
output Timed out
```

3.572.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+b*cos(d*x+c))**2,x)
```

```
output Timed out
```

3.572.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorithm
m="maxima")
```

```
output Timed out
```

3.572.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^2, x)`

3.572.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{(a + b \cos(c + dx))^2} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^2,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^2, x)`

3.573
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.573.1 Optimal result 5209
 3.573.2 Mathematica [B] (warning: unable to verify) 5210
 3.573.3 Rubi [A] (verified) 5211
 3.573.4 Maple [B] (verified) 5217
 3.573.5 Fricas [F(-1)] 5218
 3.573.6 Sympy [F(-1)] 5218
 3.573.7 Maxima [F(-1)] 5218
 3.573.8 Giac [F] 5219
 3.573.9 Mupad [F(-1)] 5219

3.573.1 Optimal result

Integrand size = 33, antiderivative size = 316

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx$$

$$= - \frac{(2a^2 A - 3Ab^2 + abB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{a^2 (a^2 - b^2) d}$$

$$+ \frac{(Ab - aB) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{a (a^2 - b^2) d}$$

$$- \frac{(5a^2 Ab - 3Ab^3 - 3a^3 B + ab^2 B) \sqrt{\cos(c + dx)} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{a^2 (a - b)(a + b)^2 d}$$

$$+ \frac{(2a^2 A - 3Ab^2 + abB) \sqrt{\sec(c + dx)} \sin(c + dx)}{a^2 (a^2 - b^2) d} + \frac{b(Ab - aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{a (a^2 - b^2) d (b + a \sec(c + dx))}$$

```
output b*(A*b-B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/(a^2-b^2)/d/(b+a*sec(d*x+c))+(2*
A*a^2-3*A*b^2+B*a*b)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/(a^2-b^2)/d-(2*A*a^2-
3*A*b^2+B*a*b)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(s
in(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a^2/(a^2-b^2)
/d+(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin
(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/(a^2-b^2)/d-(
5*A*a^2*b-3*A*b^3-3*B*a^3+B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*
x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/2)
*sec(d*x+c)^(1/2)/a^2/(a-b)/(a+b)^2/d
```

3.573.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.573.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 681 vs. $2(316) = 632$.

Time = 6.87 (sec) , antiderivative size = 681, normalized size of antiderivative = 2.16

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx =$$

$$\frac{2(10a^2Ab - 9Ab^3 - 4a^3B + 3ab^2B) \cos^2(c + dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) \right) (b + a \sec(c + dx))}{a(a + b \cos(c + dx))(1 - \cos^2(c + dx))}$$

$$+ \frac{\sqrt{\sec(c + dx)} \left(\frac{(2a^2A - 3Ab^2 + abB) \sin(c + dx)}{a^2(a^2 - b^2)} + \frac{Ab^2 \sin(c + dx) - abB \sin(c + dx)}{a(a^2 - b^2)(a + b \cos(c + dx))} \right)}{d}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^2, x]`

output

```
-1/4*((2*(10*a^2*A*b - 9*A*b^3 - 4*a^3*B + 3*a*b^2*B)*Cos[c + d*x]^2*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] - EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(a*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)) + (2*(4*a^3*A - 8*a*A*b^2 + 4*a^2*b*B)*Cos[c + d*x]^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(b*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)) + ((2*a^2*A*b - 3*A*b^3 + a*b^2*B)*Cos[2*(c + d*x)]*(b + a*Sec[c + d*x])*(-4*a*b + 4*a*b*Sec[c + d*x]^2 - 4*a*b*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*(2*a - b)*b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] - 4*a^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2])*Sin[c + d*x])/(a*b^2*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)*Sqrt[Sec[c + d*x]]*(2 - Sec[c + d*x]^2)))/(a^2*(a - b)*(a + b)*d) + (Sqrt[Sec[c + d*x]]*(((2*a^2*A - 3*A*b^2 + a*b*B)*Sin[c + d*x])/(a^2*(a^2 - b^2)) + (A*b^2*Sin[c + d*x] - a*b*B*Sin[c + d*x])/(a*(a^2 - b^2)*(a + b*Cos[c + d*x]))))/d
```

3.573.3 Rubi [A] (verified)

Time = 2.40 (sec) , antiderivative size = 306, normalized size of antiderivative = 0.97, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4517, 27, 3042, 4590, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{\sec^{\frac{5}{2}}(c+dx)(A\sec(c+dx)+B)}{(a\sec(c+dx)+b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2}(A\csc(c+dx+\frac{\pi}{2})+B)}{(a\csc(c+dx+\frac{\pi}{2})+b)^2} dx \\
 & \quad \downarrow \text{4517} \\
 & \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \\
 & \frac{\int -\frac{\sqrt{\sec(c+dx)}((2Aa^2+bBa-3Ab^2)\sec^2(c+dx)-2a(Ab-aB)\sec(c+dx)+b(Ab-aB))}{2(b+a\sec(c+dx))} dx}{a(a^2-b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{\sec(c+dx)}((2Aa^2+bBa-3Ab^2)\sec^2(c+dx)-2a(Ab-aB)\sec(c+dx)+b(Ab-aB))}{b+a\sec(c+dx)} dx}{2a(a^2-b^2)} + \\
 & \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a\sec(c+dx)+b)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.573. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+b\cos(c+dx))^2} dx$

$$\frac{\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})} \left((2Aa^2+bBa-3Ab^2) \csc(c+dx+\frac{\pi}{2})^2 - 2a(Ab-aB) \csc(c+dx+\frac{\pi}{2}) + b(Ab-aB) \right)}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{2a(a^2-b^2)} +$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4590

$$\frac{2 \int -\frac{(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) \sec^2(c+dx)+2a(Aa^2+bBa-2Ab^2) \sec(c+dx)+b(2Aa^2+bBa-3Ab^2)}{2\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx}{a} + \frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad}$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 27

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) \sec^2(c+dx)+2a(Aa^2+bBa-2Ab^2) \sec(c+dx)+b(2Aa^2+bBa-3Ab^2)}{\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx}{a}$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{(-2Ba^3+4Aba^2+b^2Ba-3Ab^3) \csc(c+dx+\frac{\pi}{2})^2+2a(Aa^2+bBa-2Ab^2) \csc(c+dx+\frac{\pi}{2})+b(2Aa^2+bBa-3Ab^2)}{\sqrt{\csc(c+dx+\frac{\pi}{2})} (b+a \csc(c+dx+\frac{\pi}{2}))} dx}{a}$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4594

$$\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{b^2(2Aa^2+bBa-3Ab^2)-ab^2(Ab-aB) \sec(c+dx)}{\sqrt{\sec(c+dx)}} dx}{b^2} + \frac{(-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a \sec(c+dx)}}{a}$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

3.573. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{b^2(2Aa^2+bBa-3Ab^2) - ab^2(Ab-aB) \csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b^2}}{a} + (-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\csc(c+dx)}{b+a \csc(c+dx)} dx$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4274

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(2a^2A+abB-3Ab^2) \int \frac{1}{\sqrt{\sec(c+dx)}} dx - ab^2(Ab-aB) \int \sqrt{\sec(c+dx)} dx}{b^2}}{a} + (-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\csc(c+dx)}{b+a \csc(c+dx)} dx$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(2a^2A+abB-3Ab^2) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - ab^2(Ab-aB) \int \sqrt{\csc(c+dx+\frac{\pi}{2})} dx}{b^2}}{a} + (-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\csc(c+dx)}{b+a \csc(c+dx)} dx$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4258

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - ab^2(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sec(c+dx)} dx}{b^2}}{a} + (-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\csc(c+dx)}{b+a \csc(c+dx)} dx$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - ab^2(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sec(c+dx)} dx}{b^2}}{a} + (-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\csc(c+dx)}{b+a \csc(c+dx)} dx$$

$$\frac{2a(a^2-b^2)}{ad(a^2-b^2)(a \sec(c+dx)+b)} \frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3119

3.573. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{2b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) - ab^2(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} f}{d}}{b^2} \frac{a}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3120

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(-3a^3B+5a^2Ab+ab^2B-3Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}}{a}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 4336

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(-3a^3B+5a^2Ab+ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \cos(c+dx))}} dx + \frac{2b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}}{a}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3042

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(-3a^3B+5a^2Ab+ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b \sin(c+dx+\frac{\pi}{2}))}} dx + \frac{2b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}}{a}}{2a(a^2-b^2)}$$

$$\frac{b(Ab-aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)}$$

↓ 3284

$$\frac{\frac{2(2a^2A+abB-3Ab^2) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{2b^2(2a^2A+abB-3Ab^2) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) - \frac{2ab^2(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d}}{b^2}}{2a(a^2-b^2)}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^2,x]`

3.573. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$

```
output (b*(A*b - a*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(a*(a^2 - b^2)*d*(b + a*Sec[c + d*x])) + (-((((2*b^2*(2*a^2*A - 3*A*b^2 + a*b*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*a*b^2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (2*(5*a^2*A*b - 3*A*b^3 - 3*a^3*B + a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/((a + b)*d))/a + (2*(2*a^2*A - 3*A*b^2 + a*b*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*d)/(2*a*(a^2 - b^2))
```

3.573.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3284 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]
```

```
rule 3439 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

rule 4258 $\text{Int}[(\text{csc}[c] + (d)(x))(b)^n, x_Symbol] \rightarrow \text{Simp}[(b \text{Csc}[c + dx])^n \text{Sin}[c + dx]^n \text{Int}[1/\text{Sin}[c + dx]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[e] + (f)(x))(d)^n (\text{csc}[e] + (f)(x))(b) + (a)], x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d \text{Csc}[e + fx])^n, x], x] + \text{Simp}[b/d \text{Int}[(d \text{Csc}[e + fx])^{n+1}, x], x] /; \text{FreeQ}\{a, b, d, e, f, n\}, x]$

rule 4336 $\text{Int}[(\text{csc}[e] + (f)(x))(d)^{3/2} / (\text{csc}[e] + (f)(x))(b) + (a)], x_Symbol] \rightarrow \text{Simp}[d \text{Sqrt}[d \text{Sin}[e + fx]] \text{Sqrt}[d \text{Csc}[e + fx]] \text{Int}[1/(\text{Sqrt}[d \text{Sin}[e + fx]](b + a \text{Sin}[e + fx])), x], x] /; \text{FreeQ}\{a, b, d, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 4517 $\text{Int}[(\text{csc}[e] + (f)(x))(d)^n (\text{csc}[e] + (f)(x))(b) + (a))^m (\text{csc}[e] + (f)(x))(B) + (A)], x_Symbol] \rightarrow \text{Simp}[a d^2 (A b - a B) \text{Cot}[e + fx] (a + b \text{Csc}[e + fx])^{m+1} ((d \text{Csc}[e + fx])^{n-2} / (b f (m+1) (a^2 - b^2))), x] - \text{Simp}[d / (b (m+1) (a^2 - b^2)) \text{Int}[(a + b \text{Csc}[e + fx])^{m+1} (d \text{Csc}[e + fx])^{n-2} \text{Simp}[a d (A b - a B) (n-2) + b d (A b - a B) (m+1) \text{Csc}[e + fx] - (a A b d (m+n) - d B (a^2 (n-1) + b^2 (m+1))) \text{Csc}[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, d, e, f, A, B\}, x] \&\& \text{NeQ}[A b - a B, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -1] \&\& \text{GtQ}[n, 1]$

rule 4590 $\text{Int}[(A) + \text{csc}[e] + (f)(x))(B) + \text{csc}[e] + (f)(x)]^2 (C) (\text{csc}[e] + (f)(x))(d)^n (\text{csc}[e] + (f)(x))(b) + (a))^m, x_Symbol] \rightarrow \text{Simp}[(-C) d \text{Cot}[e + fx] (a + b \text{Csc}[e + fx])^{m+1} ((d \text{Csc}[e + fx])^{n-1} / (b f (m+n+1))), x] + \text{Simp}[d / (b (m+n+1)) \text{Int}[(a + b \text{Csc}[e + fx])^m (d \text{Csc}[e + fx])^{n-1} \text{Simp}[a C (n-1) + (A b (m+n+1) + b C (m+n)) \text{Csc}[e + fx] + (b B (m+n+1) - a C n) \text{Csc}[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, d, e, f, A, B, C, m\}, x] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{GtQ}[n, 0]$

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))/ (Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.573.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 855 vs. $2(378) = 756$.

Time = 8.25 (sec) , antiderivative size = 856, normalized size of antiderivative = 2.71

method	result	size
default	Expression too large to display	856

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^2,x,method=_RETURNV
ERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*A/a^2/sin(1/
2*d*x+1/2*c)^2/(2*sin(1/2*d*x+1/2*c)^2-1)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+
1/2*c),2^(1/2)))+4*A*b^2/a^2/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-
2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*
c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+2*(-A*b+B*a)
/a*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/
(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/
2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*
d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos
(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-
2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)
^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1...
```

3.573.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^2} dx$$

3.573.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorithm m="fricas")`

output `Timed out`

3.573.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**2,x)`

output `Timed out`

3.573.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`

output `Timed out`

3.573.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^2, x)`

3.573.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{(a + b \cos(c + dx))^2} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^2,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^2, x)`

$$3.574 \quad \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx$$

3.574.1 Optimal result	5220
3.574.2 Mathematica [A] (verified)	5221
3.574.3 Rubi [A] (verified)	5221
3.574.4 Maple [B] (verified)	5226
3.574.5 Fracas [F(-1)]	5227
3.574.6 Sympy [F]	5228
3.574.7 Maxima [F]	5228
3.574.8 Giac [F]	5228
3.574.9 Mupad [F(-1)]	5229

3.574.1 Optimal result

Integrand size = 33, antiderivative size = 260

$$\begin{aligned} & \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx \\ &= -\frac{(Ab-aB)\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\mid 2\right)\sqrt{\sec(c+dx)}}{a(a^2-b^2)d} \\ & \quad -\frac{(Ab-aB)\sqrt{\cos(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)\sqrt{\sec(c+dx)}}{b(a^2-b^2)d} \\ & \quad +\frac{(3a^2Ab-Ab^3-a^3B-ab^2B)\sqrt{\cos(c+dx)}\operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2\right)\sqrt{\sec(c+dx)}}{a(a-b)b(a+b)^2d} \\ & \quad +\frac{b(Ab-aB)\sqrt{\sec(c+dx)}\sin(c+dx)}{a(a^2-b^2)d(b+a \sec(c+dx))} \end{aligned}$$

output

```

b*(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/(a^2-b^2)/d/(b+a*sec(d*x+c))-
(A*b-B*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d
*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/a/(a^2-b^2)/d-(A*b-B*
a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1
/2*c), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b/(a^2-b^2)/d+(3*A*a^2*b-
A*b^3-B*a^3-B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellip
ticPi(sin(1/2*d*x+1/2*c), 2*b/(a+b), 2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1
/2)/a/(a-b)/b/(a+b)^2/d

```

$$3.574. \quad \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx$$

3.574.2 Mathematica [A] (verified)

Time = 6.01 (sec) , antiderivative size = 443, normalized size of antiderivative = 1.70

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx$$

$$= \frac{-\frac{4ab(-Ab+aB) \sin(c+dx)}{a^2-b^2} + \frac{4 \cos(c+dx) \cot(c+dx)(b+a \sec(c+dx))(aAb^2-a^2bB-aAb^2 \sec^2(c+dx)+a^2bB \sec^2(c+dx)-ab(-Ab+aB)E(\ar$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^2, x]`

output `((-4*a*b*(-(A*b) + a*B)*Sin[c + d*x])/(a^2 - b^2) + (4*Cos[c + d*x]*Cot[c + d*x]*(b + a*Sec[c + d*x])*(a*A*b^2 - a^2*b*B - a*A*b^2*Sec[c + d*x]^2 + a^2*b*B*Sec[c + d*x]^2 - a*b*(-(A*b) + a*B)*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + (a - b)*b*(2*a*A + A*b + a*B)*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 3*a^2*A*b*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + A*b^3*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + a^3*B*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + a*b^2*B*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2]))/(4*a^2*d*(a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]])`

3.574.3 Rubi [A] (verified)

Time = 1.83 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.97, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4517, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{(a + b \cos(c + dx))^2} dx$$

↓ 3042

3.574. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx$

$$\begin{aligned}
 & \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{\sec^{\frac{3}{2}}(c+dx)(A\sec(c+dx)+B)}{(a\sec(c+dx)+b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A\csc(c+dx+\frac{\pi}{2})+B)}{(a\csc(c+dx+\frac{\pi}{2})+b)^2} dx \\
 & \quad \downarrow \text{4517} \\
 & \frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \\
 & \frac{\int \frac{-((2Aa^2-bBa-Ab^2)\sec^2(c+dx))+2a(Ab-aB)\sec(c+dx)+b(Ab-aB)}{2\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx}{a(a^2-b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \\
 & \frac{\int \frac{-((2Aa^2-bBa-Ab^2)\sec^2(c+dx))+2a(Ab-aB)\sec(c+dx)+b(Ab-aB)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx}{2a(a^2-b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \\
 & \frac{\int \frac{(-2Aa^2+bBa+Ab^2)\csc(c+dx+\frac{\pi}{2})^2+2a(Ab-aB)\csc(c+dx+\frac{\pi}{2})+b(Ab-aB)}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(b+a\csc(c+dx+\frac{\pi}{2}))} dx}{2a(a^2-b^2)} \\
 & \quad \downarrow \text{4594} \\
 & \frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \\
 & \frac{\int \frac{(Ab-aB)b^2+a(Ab-aB)\sec(c+dx)b}{\sqrt{\sec(c+dx)}} dx}{b^2} - \frac{(a^3(-B)+3a^2Ab-ab^2B-Ab^3)\int \frac{\sec^{\frac{3}{2}}(c+dx)}{b+a\sec(c+dx)} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b\cos(c+dx))^2} dx}{2a(a^2-b^2)}
 \end{aligned}$$

3.574. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b\cos(c+dx))^2} dx$

$$\frac{\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{\int \frac{(Ab - aB)b^2 + a(Ab - aB) \csc(c + dx + \frac{\pi}{2})b}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx}{b^2} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b}}{2a(a^2 - b^2)}$$

↓ 4274

$$\frac{\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{b^2(Ab - aB) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + ab(Ab - aB) \int \sqrt{\sec(c + dx)} dx}{b^2} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b}}{2a(a^2 - b^2)}$$

↓ 3042

$$\frac{\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{b^2(Ab - aB) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + ab(Ab - aB) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx}{b^2} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b}}{2a(a^2 - b^2)}$$

↓ 4258

$$\frac{\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{b^2} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3)}{b}}{2a(a^2 - b^2)}$$

↓ 3042

$$\frac{\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{b^2} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3)}{b}}{2a(a^2 - b^2)}$$

↓ 3119

$$\frac{\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2}(c + dx) | 2)}{d}}{b^2} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx}{b}}{2a(a^2 - b^2)}$$

↓ 3120

3.574. $\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx$

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} + \frac{2ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \int \frac{\csc(c + dx)}{b + a \cos(c + dx)} dx}{b}$$

$2a(a^2 - b^2)$

↓ 4336

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} + \frac{2ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \sqrt{\cos(c + dx)}}{b}$$

$2a(a^2 - b^2)$

↓ 3042

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} + \frac{2ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \sqrt{\cos(c + dx)}}{b}$$

$2a(a^2 - b^2)$

↓ 3284

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{ad(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2b^2(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d} + \frac{2ab(Ab - aB) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} - \frac{2(a^3(-B) + 3a^2Ab - ab^2B - Ab^3) \sqrt{\cos(c + dx)}}{b}$$

$2a(a^2 - b^2)$

```
input Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^2,x]
```

```
output -1/2*(((2*b^2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 - (2*(3*a^2*A*b - A*b^3 - a^3*B - a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b*(a + b)*d)/(a*(a^2 - b^2)) + (b*(A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*(a^2 - b^2)*d*(b + a*Sec[c + d*x]))
```

3.574. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx$

3.574.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)])*(d_.)^(n_)*(csc[(e_.) + (f_)*(x_)])*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4517 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[a*d^2*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 2)/(b*f*(m + 1)*(a^2 - b^2))), x] - Simp[d/(b*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 2)*Simp[a*d*(A*b - a*B)*(n - 2) + b*d*(A*b - a*B)*(m + 1)*Csc[e + f*x] - (a*A*b*d*(m + n) - d*B*(a^2*(n - 1) + b^2*(m + 1)))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && GtQ[n, 1]`

rule 4594 `Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e + f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]`

3.574.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 720 vs. 2(324) = 648.

Time = 7.46 (sec) , antiderivative size = 721, normalized size of antiderivative = 2.77

method	result
default	$-\frac{\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(-2ab + 2b^2\right) \sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)}} \left(\frac{4B \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} \Pi\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), -\frac{2b}{a-b}, \sqrt{2}\right)}{2(Ab - Ba)} \right) + \dots$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^2,x,method=_RETURNV ERBOSE)`

$$3.574. \int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^2} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B/(-2*a*b+2
*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*s
in(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2
*c),-2*b/(a-b),2^(1/2))+2*(A*b-B*a)/b*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*
c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1
/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*
c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1
/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x
+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(
sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*
d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1
/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1
/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1
/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a*
b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2
)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*
d*x+1/2*c),-2*b/(a-b),2^(1/2)))/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^
2-1)^(1/2)/d

```

3.574.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algorithm="fracas")`

output `Timed out`

3.574.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**2,x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/(a + b*cos(c + d*x))**2, x)`

3.574.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^2, x)`

3.574.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^2} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^2, x)`

3.574.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^2} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + b \cos(c + dx))^2} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^2,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^2,x)`

3.575
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$$

3.575.1 Optimal result 5230
 3.575.2 Mathematica [A] (verified) 5231
 3.575.3 Rubi [A] (verified) 5231
 3.575.4 Maple [B] (verified) 5236
 3.575.5 Fricas [F(-1)] 5237
 3.575.6 Sympy [F] 5238
 3.575.7 Maxima [F] 5238
 3.575.8 Giac [F] 5238
 3.575.9 Mupad [F(-1)] 5239

3.575.1 Optimal result

Integrand size = 33, antiderivative size = 258

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{(Ab - aB) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{b(a^2 - b^2)d}$$

$$+ \frac{(aAb + a^2B - 2b^2B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{b^2(a^2 - b^2)d}$$

$$- \frac{(a^2Ab + Ab^3 + a^3B - 3ab^2B) \sqrt{\cos(c + dx)} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{(a - b)b^2(a + b)^2d}$$

$$- \frac{(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{(a^2 - b^2)d(b + a \sec(c + dx))}$$

output

```
-(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/(a^2-b^2)/d/(b+a*sec(d*x+c))+(A*b-B
*a)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+
1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b/(a^2-b^2)/d+(A*a*b+B*a
^2-2*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(
1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^2/(a^2-b^2)/d-
(A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1
/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/2)*se
c(d*x+c)^(1/2)/(a-b)/b^2/(a+b)^2/d
```

3.575.2 Mathematica [A] (verified)

Time = 5.29 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.71

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{\frac{4b^2(Ab - aB) \sin(c + dx)}{-a^2 + b^2} + \frac{4 \cos(c + dx) \cot(c + dx)(b + a \sec(c + dx))(-aAb^2 + a^2bB + aAb^2 \sec^2(c + dx) - a^2bB \sec^2(c + dx) + ab(-Ab + aB)E(\arcsin(\frac{b + a \sec(c + dx)}{\sqrt{a^2 + b^2}}))}{-a^2 + b^2}}{-a^2 + b^2}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^2*Sqrt[Sec[c + d*x]]),x]`

output `((4*b^2*(A*b - a*B)*Sin[c + d*x])/(-a^2 + b^2) + (4*Cos[c + d*x]*Cot[c + d*x]*(b + a*Sec[c + d*x])*(-(a*A*b^2) + a^2*b*B + a*A*b^2*Sec[c + d*x]^2 - a^2*b*B*Sec[c + d*x]^2 + a*b*(-(A*b) + a*B)*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - (a - b)*b*(-(A*b) + a*B)*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + a^2*A*b*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + A*b^3*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + a^3*B*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 3*a*b^2*B*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2]))/(a*(a - b)*(a + b)))/(4*b^2*d*(a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]])`

3.575.3 Rubi [A] (verified)Time = 1.73 (sec) , antiderivative size = 247, normalized size of antiderivative = 0.96, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4515, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))^2} dx$$

↓ 3042

3.575. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right) \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{\sqrt{\sec(c + dx)}(A \sec(c + dx) + B)}{(a \sec(c + dx) + b)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(A \csc\left(c + dx + \frac{\pi}{2}\right) + B)}{\left(a \csc\left(c + dx + \frac{\pi}{2}\right) + b\right)^2} dx \\
& \quad \downarrow \text{4515} \\
& \frac{\int \frac{-((Ab - aB) \sec^2(c + dx)) + 2(aA - bB) \sec(c + dx) + Ab - aB}{2\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{a^2 - b^2} - \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{-((Ab - aB) \sec^2(c + dx)) + 2(aA - bB) \sec(c + dx) + Ab - aB}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{2(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(aB - Ab) \csc\left(c + dx + \frac{\pi}{2}\right)^2 + 2(aA - bB) \csc\left(c + dx + \frac{\pi}{2}\right) + Ab - aB}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(b + a \csc\left(c + dx + \frac{\pi}{2}\right))} dx}{2(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} \\
& \quad \downarrow \text{4594} \\
& \frac{\int \frac{b(Ab - aB) + (Ba^2 + Aba - 2b^2B) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx}{b^2} - \frac{(a^3B + a^2Ab - 3ab^2B + Ab^3) \int \frac{\sec^{\frac{3}{2}}(c + dx)}{b + a \sec(c + dx)} dx}{b^2} \\
& \quad \frac{2(a^2 - b^2)}{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}} \\
& \quad \frac{d(a^2 - b^2)(a \sec(c + dx) + b)}{\quad} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{b(Ab - aB) + (Ba^2 + Aba - 2b^2B) \csc\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx}{b^2} - \frac{(a^3B + a^2Ab - 3ab^2B + Ab^3) \int \frac{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2}}{b + a \csc\left(c + dx + \frac{\pi}{2}\right)} dx}{b^2} \\
& \quad \frac{2(a^2 - b^2)}{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}} \\
& \quad \frac{d(a^2 - b^2)(a \sec(c + dx) + b)}{\quad} \\
& \quad \downarrow \text{4274}
\end{aligned}$$

3.575. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$

$$\frac{(a^2B+aAb-2b^2B) \int \sqrt{\sec(c+dx)}dx+b(Ab-aB) \int \frac{1}{\sqrt{\sec(c+dx)}} dx}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2}$$

$$\frac{2(a^2-b^2)}{d(a^2-b^2)(a \sec(c+dx)+b)} \frac{(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a^2-b^2)(a \sec(c+dx)+b)}$$

3042

$$\frac{(a^2B+aAb-2b^2B) \int \sqrt{\csc(c+dx+\frac{\pi}{2})}dx+b(Ab-aB) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx}{b^2}$$

$$\frac{2(a^2-b^2)}{d(a^2-b^2)(a \sec(c+dx)+b)} \frac{(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a^2-b^2)(a \sec(c+dx)+b)}$$

4258

$$\frac{(a^2B+aAb-2b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx+b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3) \int \sqrt{\cos(c+dx)} dx}{b^2}$$

$$\frac{2(a^2-b^2)}{d(a^2-b^2)(a \sec(c+dx)+b)} \frac{(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a^2-b^2)(a \sec(c+dx)+b)}$$

3042

$$\frac{(a^2B+aAb-2b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx+b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b^2}$$

$$\frac{2(a^2-b^2)}{d(a^2-b^2)(a \sec(c+dx)+b)} \frac{(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a^2-b^2)(a \sec(c+dx)+b)}$$

3119

$$\frac{(a^2B+aAb-2b^2B) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx+\frac{2b(Ab-aB) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E(\frac{1}{2}(c+dx)|2)}{d}}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3) \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx}{b^2}$$

$$\frac{2(a^2-b^2)}{d(a^2-b^2)(a \sec(c+dx)+b)} \frac{(Ab-aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{d(a^2-b^2)(a \sec(c+dx)+b)}$$

3120

3.575. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

$$\frac{\frac{2(a^2B+aAb-2b^2B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)\right)}{d}}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3)\int \frac{\cos(c+dx)}{b-a}}{b^2}$$

$$\frac{2(a^2 - b^2)}{d(a^2 - b^2)(a \sec(c + dx) + b)} \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 4336

$$\frac{\frac{2(a^2B+aAb-2b^2B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)\right)}{d}}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3)\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2(a^2 - b^2)}{d(a^2 - b^2)(a \sec(c + dx) + b)} \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 3042

$$\frac{\frac{2(a^2B+aAb-2b^2B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)\right)}{d}}{b^2} - \frac{(a^3B+a^2Ab-3ab^2B+Ab^3)\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2(a^2 - b^2)}{d(a^2 - b^2)(a \sec(c + dx) + b)} \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)}$$

↓ 3284

$$\frac{\frac{2(a^2B+aAb-2b^2B)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b(Ab-aB)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}E\left(\frac{1}{2}(c+dx)\right)}{d}}{b^2} - \frac{2(a^3B+a^2Ab-3ab^2B+Ab^3)\sqrt{\cos(c+dx)}}{b^2}$$

$$\frac{2(a^2 - b^2)}{d(a^2 - b^2)(a \sec(c + dx) + b)} \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)}$$

```
input Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^2*Sqrt[Sec[c + d*x]]),x]
```

```
output (((2*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*(a*A*b + a^2*B - 2*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 - (2*(a^2*A*b + A*b^3 + a^3*B - 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d)/(2*(a^2 - b^2)) - ((A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*d*(b + a*Sec[c + d*x]))
```

3.575. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sqrt{\sec(c+dx)}} dx$

3.575.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_.)^(p_))*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

```
rule 4336 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[
1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e,
f}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 4515 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) +
(a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-d)*(A
*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n -
1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b
*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[d*(n - 1)*(A*b - a*B)
+ d*(a*A - b*B)*(m + 1)*Csc[e + f*x] - d*(A*b - a*B)*(m + n + 1)*Csc[e + f
*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] &&
NeQ[a^2 - b^2, 0] && LtQ[m, -1] && LtQ[0, n, 1]
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/ (Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.575.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 807 vs. $2(322) = 644$.

Time = 7.94 (sec) , antiderivative size = 808, normalized size of antiderivative = 3.13

method	result	size
default	Expression too large to display	808

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

output
$$\begin{aligned} & -(-(-2\cos(1/2dx+1/2c)^2+1)\sin(1/2dx+1/2c)^2)^{1/2}*(2B/b^2*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})- \\ & 4/b*(A*b-2*B*a)/(-2*a*b+2*b^2)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2dx+1/2c),-2*b/(a-b),2^{1/2})-2*a*(A*b-B*a)/b^2*(-1/a*b \\ & ^2/(a^2-b^2)*\cos(1/2dx+1/2c)*(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}/(2*b*\cos(1/2dx+1/2c)^2+a-b)-1/2/a/(a+b)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2})-1/2/(a^2-b^2)* \\ & b/a*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2}))+1/2/(a^2-b^2)*b/a*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticE}(\cos(1/2dx+1/2c),2^{1/2})-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2dx+1/2c),-2*b/(a-b),2^{1/2}))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2dx+1/2c),-2*b/(a-b),2^{1/2}))/\sin(1/2d\dots \end{aligned}$$

3.575.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `Timed out`

3.575.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/((a + b*cos(c + d*x))**2*sqrt(sec(c + d*x))), x)`

3.575.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sqrt(sec(d*x + c))), x)`

3.575.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sqrt(sec(d*x + c))), x)`

3.575.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^2),x)`

$$3.576 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{3}{2}}(c+dx)} dx$$

3.576.1 Optimal result	5240
3.576.2 Mathematica [A] (verified)	5241
3.576.3 Rubi [A] (verified)	5241
3.576.4 Maple [B] (verified)	5246
3.576.5 Fricas [F(-1)]	5247
3.576.6 Sympy [F(-1)]	5248
3.576.7 Maxima [F]	5248
3.576.8 Giac [F]	5248
3.576.9 Mupad [F(-1)]	5249

3.576.1 Optimal result

Integrand size = 33, antiderivative size = 284

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx \\ &= - \frac{(aAb - 3a^2B + 2b^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{b^2 (a^2 - b^2) d} \\ &+ \frac{(a^2Ab - 2Ab^3 - 3a^3B + 4ab^2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{b^3 (a^2 - b^2) d} \\ &- \frac{a(a^2Ab - 3Ab^3 - 3a^3B + 5ab^2B) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{(a - b)b^3(a + b)^2d} \\ &+ \frac{a(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{b(a^2 - b^2) d(b + a \sec(c + dx))} \end{aligned}$$

output

```
a*(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/b/(a^2-b^2)/d/(b+a*sec(d*x+c))-
(A*a*b-3*B*a^2+2*B*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*Ellip
ticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^2/(a^
2-b^2)/d+(A*a^2*b-2*A*b^3-3*B*a^3+4*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/
cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*
sec(d*x+c)^(1/2)/b^3/(a^2-b^2)/d-a*(A*a^2*b-3*A*b^3-3*B*a^3+5*B*a*b^2)*(co
s(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c)
,2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/(a-b)/b^3/(a+b)^2/d
```

3.576.2 Mathematica [A] (verified)

Time = 6.52 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.62

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{4ab^2(-Ab+aB)\sin(c+dx)}{-a^2+b^2} + \frac{\cos(c+dx)\cot(c+dx)(b+a\sec(c+dx))(-2b^2(-aAb-a^2B+2b^2B))\left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c+dx)}\right),-1\right)-\text{EllipticE}\left(\arcsin\left(\sqrt{\sec(c+dx)}\right),-1\right)\right)}{-a^2+b^2}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^2*Sec[c + d*x]^(3/2)),x]`

output `((4*a*b^2*(-(A*b) + a*B)*Sin[c + d*x])/(-a^2 + b^2) + (Cos[c + d*x]*Cot[c + d*x]*(b + a*Sec[c + d*x])*(-2*b^2*(-(a*A*b) - a^2*B + 2*b^2*B)*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] - EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + 8*a*b^2*(-(A*b) + a*B)*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + (a*A*b - 3*a^2*B + 2*b^2*B)*(4*a*b - 4*a*b*Sec[c + d*x]^2 + 4*a*b*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 2*(2*a - b)*b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + 4*a^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 2*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2]))/(a*(a - b)*(a + b)))/(4*b^3*d*(a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]])`

3.576.3 Rubi [A] (verified)

Time = 1.91 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.97, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 3439, 3042, 4518, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^2} dx$$

↓ 3042

3.576. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2} dx \\
& \quad \downarrow \text{3439} \\
& \int \frac{A \sec(c + dx) + B}{\sqrt{\sec(c + dx)} (a \sec(c + dx) + b)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A \csc\left(c + dx + \frac{\pi}{2}\right) + B}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + b\right)^2} dx \\
& \quad \downarrow \text{4518} \\
& \frac{\int -\frac{-3Ba^2 - (Ab - aB) \sec^2(c + dx)a + Aba + 2b^2 B + 2b(Ab - aB) \sec(c + dx)}{2\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{b(a^2 - b^2)} + \\
& \quad \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} \\
& \quad \downarrow \text{27} \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{\int -\frac{-3Ba^2 - (Ab - aB) \sec^2(c + dx)a + Aba + 2b^2 B + 2b(Ab - aB) \sec(c + dx)}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{2b(a^2 - b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\
& \frac{\int -\frac{-3Ba^2 - (Ab - aB) \csc\left(c + dx + \frac{\pi}{2}\right)^2 a + Aba + 2b^2 B + 2b(Ab - aB) \csc\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(b + a \csc\left(c + dx + \frac{\pi}{2}\right))} dx}{2b(a^2 - b^2)} \\
& \quad \downarrow \text{4594} \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\
& \frac{a(-3a^3 B + a^2 Ab + 5ab^2 B - 3Ab^3) \int \frac{\sec^{\frac{3}{2}}(c + dx)}{b + a \sec(c + dx)} dx + \int \frac{b(-3Ba^2 + Aba + 2b^2 B) - (-3Ba^3 + Aba^2 + 4b^2 Ba - 2Ab^3) \sec(c + dx)}{\sqrt{\sec(c + dx)}} dx}{b^2}{2b(a^2 - b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\
& \frac{a(-3a^3 B + a^2 Ab + 5ab^2 B - 3Ab^3) \int \frac{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2}}{b + a \csc\left(c + dx + \frac{\pi}{2}\right)} dx + \int \frac{b(-3Ba^2 + Aba + 2b^2 B) + (3Ba^3 - Aba^2 - 4b^2 Ba + 2Ab^3) \csc\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx}{b^2}{2b(a^2 - b^2)}
\end{aligned}$$

3.576. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned} & \downarrow 4274 \\ & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\ & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{b(-3a^2B + aAb + 2b^2B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx - (-3a^3B + a^2Ab + 4ab^2B - 2Ab^3) \int \sqrt{\sec(c + dx)}}{b^2} \\ & \frac{\phantom{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}}{2b(a^2 - b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\ & \frac{b(-3a^2B + aAb + 2b^2B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx - (-3a^3B + a^2Ab + 4ab^2B - 2Ab^3) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} \\ & \frac{\phantom{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}}{2b(a^2 - b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4258 \\ & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\ & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx - (-3a^3B + a^2Ab + 4ab^2B - 2Ab^3) \int \sqrt{\sec(c + dx)}}{b^2} \\ & \frac{\phantom{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}}{2b(a^2 - b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\ & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx - (-3a^3B + a^2Ab + 4ab^2B - 2Ab^3) \int \sqrt{\sec(c + dx)}}{b^2} \\ & \frac{\phantom{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}}{2b(a^2 - b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3119 \\ & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\ & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{2b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) - (-3a^3B + a^2Ab + 4ab^2B - 2Ab^3) \int \sqrt{\sec(c + dx)}}{b^2} \\ & \frac{\phantom{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}}{2b(a^2 - b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3120 \\ & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\ & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{2b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) - 2(-3a^3B + a^2Ab + 4ab^2B - 2Ab^3) \int \sqrt{\sec(c + dx)}}{b^2} \\ & \frac{\phantom{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}}{2b(a^2 - b^2)} \end{aligned}$$

3.576. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow 4336 \\
 & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\
 & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{b^2} + \frac{2b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx)\right)}{d} \\
 & \frac{\hspace{10em}}{2b(a^2 - b^2)} \\
 & \downarrow 3042 \\
 & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\
 & \frac{a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))} dx}{b^2} + \frac{2b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \\
 & \frac{\hspace{10em}}{2b(a^2 - b^2)} \\
 & \downarrow 3284 \\
 & \frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{bd(a^2 - b^2)(a \sec(c + dx) + b)} - \\
 & \frac{2a(-3a^3B + a^2Ab + 5ab^2B - 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a + b}, \frac{1}{2}(c + dx), 2\right)}{b^2d(a + b)} + \frac{2b(-3a^2B + aAb + 2b^2B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx)\right)}{d} \\
 & \frac{\hspace{10em}}{2b(a^2 - b^2)}
 \end{aligned}$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^2*Sec[c + d*x]^(3/2)),x]`

output `-1/2*(((2*b*(a*A*b - 3*a^2*B + 2*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*(a^2*A*b - 2*A*b^3 - 3*a^3*B + 4*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (2*a*(a^2*A*b - 3*A*b^3 - 3*a^3*B + 5*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d))/b*(a^2 - b^2) + (a*(A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b*(a^2 - b^2)*d*(b + a*Sec[c + d*x]))`

3.576.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.576. \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx$$

rule 3119 $\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3284 $\text{Int}[1/(((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)])*\text{Sqrt}[(c_.) + (d_.)*\sin[(e_.) + (f_.)*(x_)]]), x_Symbol] \rightarrow \text{Simp}[(2/(f*(a + b)*\text{Sqrt}[c + d]))*\text{EllipticPi}[2*(b/(a + b)), (1/2)*(e - \text{Pi}/2 + f*x), 2*(d/(c + d))], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[c + d, 0]$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(g_.)^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m + n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p - m - n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n + 1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4336 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(3/2)}/(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[d*\text{Sqrt}[d*\text{Sin}[e + f*x]]*\text{Sqrt}[d*\text{Csc}[e + f*x]] \text{Int}[1/(\text{Sqrt}[d*\text{Sin}[e + f*x]]*(b + a*\text{Sin}[e + f*x])), x], x] /; \text{FreeQ}[\{a, b, d, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

```
rule 4518 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[b*(A*b
- a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(
m + 1)*(a^2 - b^2))), x] + Simp[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[
e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[A*(a^2*(m + 1) - b^2*(m + n + 1))
+ a*b*B*n - a*(A*b - a*B)*(m + 1)*Csc[e + f*x] + b*(A*b - a*B)*(m + n + 2)
*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*
b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && IL
tQ[n, 0])
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.576.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 848 vs. $2(348) = 696$.

Time = 9.53 (sec) , antiderivative size = 849, normalized size of antiderivative = 2.99

method	result	size
default	Expression too large to display	849

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2/sec(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2/b^3/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(A*b*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)
)-2*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*a-B*b*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2)))+2*a^2*(A*b-B*a)/b^3*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*
(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*
c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^
2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(
cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/
2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*(sin
(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x
+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)
)-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*
d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2
*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-
2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x
+1/2*c),-2*b/(a-b),2^(1/2))+4*a/b^2*(2*A*b-3*B*a)/(-2*a*b+2*b^2)*(sin(1/2
*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+...

```

3.576.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `Timed out`

3.576.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2/sec(d*x+c)**(3/2),x)`output `Timed out`**3.576.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(3/2),x, algorithm m="maxima")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2)), x)`**3.576.8 Giac [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(3/2),x, algorithm m="giac")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sec(d*x + c)^(3/2)), x)`

3.576.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} (a + b \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^2),x)`

$$3.577 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$$

3.577.1 Optimal result	5250
3.577.2 Mathematica [A] (warning: unable to verify)	5251
3.577.3 Rubi [A] (verified)	5252
3.577.4 Maple [B] (verified)	5257
3.577.5 Fricas [F(-1)]	5258
3.577.6 Sympy [F(-1)]	5259
3.577.7 Maxima [F]	5259
3.577.8 Giac [F]	5259
3.577.9 Mupad [F(-1)]	5260

3.577.1 Optimal result

Integrand size = 33, antiderivative size = 363

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx \\ &= \frac{(3a^2Ab - 2Ab^3 - 5a^3B + 4ab^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{b^3(a^2 - b^2)d} \\ & \quad - \frac{(9a^3Ab - 12aAb^3 - 15a^4B + 16a^2b^2B + 2b^4B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3b^4(a^2 - b^2)d} \\ & \quad + \frac{a^2(3a^2Ab - 5Ab^3 - 5a^3B + 7ab^2B) \sqrt{\cos(c + dx)} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{(a - b)b^4(a + b)^2d} \\ & \quad - \frac{(3aAb - 5a^2B + 2b^2B) \sin(c + dx)}{3b^2(a^2 - b^2)d\sqrt{\sec(c + dx)}} + \frac{a(Ab - aB) \sin(c + dx)}{b(a^2 - b^2)d\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} \end{aligned}$$

output

```
-1/3*(3*A*a*b-5*B*a^2+2*B*b^2)*sin(d*x+c)/b^2/(a^2-b^2)/d/sec(d*x+c)^(1/2)
+a*(A*b-B*a)*sin(d*x+c)/b/(a^2-b^2)/d/(b+a*sec(d*x+c))/sec(d*x+c)^(1/2)+(3
*A*a^2*b-2*A*b^3-5*B*a^3+4*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d
*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c
)^(1/2)/b^3/(a^2-b^2)/d-1/3*(9*A*a^3*b-12*A*a*b^3-15*B*a^4+16*B*a^2*b^2+2*
B*b^4)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d
*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/b^4/(a^2-b^2)/d+a^2*(
3*A*a^2*b-5*A*b^3-5*B*a^3+7*B*a*b^2)*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*
d*x+1/2*c)*EllipticPi(sin(1/2*d*x+1/2*c),2*b/(a+b),2^(1/2))*cos(d*x+c)^(1/
2)*sec(d*x+c)^(1/2)/(a-b)/b^4/(a+b)^2/d
```

$$3.577. \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$$

3.577.2 Mathematica [A] (warning: unable to verify)

Time = 6.84 (sec) , antiderivative size = 701, normalized size of antiderivative = 1.93

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{2(-3a^2Ab + 6Ab^3 + 5a^3B - 8ab^2B) \cos^2(c + dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) \right) (b + a \sec(c + dx))}{a(a + b \cos(c + dx))(1 - \cos^2(c + dx))}$$

$$+ \frac{\sqrt{\sec(c + dx)} \left(\frac{a^2(-Ab + aB) \sin(c + dx)}{b^3(a^2 - b^2)} - \frac{a^3Ab \sin(c + dx) - a^4B \sin(c + dx)}{b^3(-a^2 + b^2)(a + b \cos(c + dx))} + \frac{B \sin(2(c + dx))}{3b^2} \right)}{d}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^2*Sec[c + d*x]^(5/2)),x]`

output

```
-1/12*((2*(-3*a^2*A*b + 6*A*b^3 + 5*a^3*B - 8*a*b^2*B)*Cos[c + d*x]^2*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] - EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(a*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)) + (2*(-12*a*A*b^2 + 8*a^2*b*B + 4*b^3*B)*Cos[c + d*x]^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(b*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)) + ((-9*a^2*A*b + 6*A*b^3 + 15*a^3*B - 12*a*b^2*B)*Cos[2*(c + d*x)]*(b + a*Sec[c + d*x])*(-4*a*b + 4*a*b*Sec[c + d*x]^2 - 4*a*b*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*(2*a - b)*b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] - 4*a^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2])*Sin[c + d*x])/(a*b^2*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)*Sqrt[Sec[c + d*x]]*(2 - Sec[c + d*x]^2)))/((a - b)*b^2*(a + b)*d) + (Sqrt[Sec[c + d*x]]*((a^2*(-(A*b) + a*B)*Sin[c + d*x])/(b^3*(a^2 - b^2)) - (a^3*A*b*Sin[c + d*x] - a^4*B*Sin[c + d*x])/(b^3*(-a^2 + b^2)*(a + b*Cos[c + d*x])) + (B*Sin[2*(c + d*x)])/(3*b^2)))/d
```

3.577.3 Rubi [A] (verified)

Time = 2.58 (sec) , antiderivative size = 350, normalized size of antiderivative = 0.96, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4518, 27, 3042, 4592, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^2} dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{A \sec(c + dx) + B}{\sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A \csc\left(c + dx + \frac{\pi}{2}\right) + B}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a \csc\left(c + dx + \frac{\pi}{2}\right) + b\right)^2} dx \\
 & \quad \downarrow \text{4518} \\
 & \frac{\int -\frac{5Ba^2 - 3(Ab - aB) \sec^2(c + dx)a + 3Aba + 2b^2B + 2b(Ab - aB) \sec(c + dx)}{2 \sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))} dx}{b(a^2 - b^2)} + \\
 & \quad \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} \\
 & \quad \downarrow \text{27} \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{\int -\frac{5Ba^2 - 3(Ab - aB) \sec^2(c + dx)a + 3Aba + 2b^2B + 2b(Ab - aB) \sec(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))} dx}{2b(a^2 - b^2)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.577. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx$

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \int \frac{-5Ba^2 - 3(Ab - aB) \csc(c + dx + \frac{\pi}{2})^2 a + 3Aba + 2b^2 B + 2b(Ab - aB) \csc(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2} (b + a \csc(c + dx + \frac{\pi}{2}))} dx}{2b(a^2 - b^2)}$$

↓ 4592

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{2 \int \frac{-a(-5Ba^2 + 3Aba + 2b^2 B) \sec^2(c + dx) + 2b(-2Ba^2 + 3Aba - b^2 B) \sec(c + dx) + 3(-5Ba^3 + 3Aba^2 + 4b^2 Ba - 2Ab^3)}{2\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{3b}}{\frac{2(-5a^2 B + 3aAb + 2b^2 B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{2b(a^2 - b^2)}{3b}}$$

↓ 27

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \int \frac{-a(-5Ba^2 + 3Aba + 2b^2 B) \sec^2(c + dx) + 2b(-2Ba^2 + 3Aba - b^2 B) \sec(c + dx) + 3(-5Ba^3 + 3Aba^2 + 4b^2 Ba - 2Ab^3)}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{3b}}{\frac{2(-5a^2 B + 3aAb + 2b^2 B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{2b(a^2 - b^2)}{3b}}$$

↓ 3042

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \int \frac{-a(-5Ba^2 + 3Aba + 2b^2 B) \csc(c + dx + \frac{\pi}{2})^2 + 2b(-2Ba^2 + 3Aba - b^2 B) \csc(c + dx + \frac{\pi}{2}) + 3(-5Ba^3 + 3Aba^2 + 4b^2 Ba - 2Ab^3)}{\sqrt{\csc(c + dx + \frac{\pi}{2})} (b + a \csc(c + dx + \frac{\pi}{2}))} dx}{3b}}{\frac{2(-5a^2 B + 3aAb + 2b^2 B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{2b(a^2 - b^2)}{3b}}$$

↓ 4594

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{3a^2(-5a^3 B + 3a^2 Ab + 7ab^2 B - 5Ab^3) \int \frac{\sec^{\frac{3}{2}}(c + dx)}{b + a \sec(c + dx)} dx + \int \frac{3b(-5Ba^3 + 3Aba^2 + 4b^2 Ba - 2Ab^3) - (-15Ba^4 + 9Aba^3 + \sqrt{\sec(c + dx)})}{b^2} dx}{b^2}}{\frac{2(-5a^2 B + 3aAb + 2b^2 B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{2b(a^2 - b^2)}{3b}}$$

↓ 3042

$$\frac{\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{3a^2(-5a^3 B + 3a^2 Ab + 7ab^2 B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + \int \frac{3b(-5Ba^3 + 3Aba^2 + 4b^2 Ba - 2Ab^3) + (15Ba^4 - 9Aba^3 + \sqrt{\csc(c + dx)})}{b^2} dx}{b^2}}{\frac{2(-5a^2 B + 3aAb + 2b^2 B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{2b(a^2 - b^2)}{3b}}$$

3.577. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow 4274 \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \int \frac{1}{\sqrt{\sec(c + dx)}} dx}{3b} \\
 & \hline
 & 2b(a^2 - b^2)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx}{3b} \\
 & \hline
 & 2b(a^2 - b^2)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 4258 \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{3b} \\
 & \hline
 & 2b(a^2 - b^2)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{3b} \\
 & \hline
 & 2b(a^2 - b^2)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3119 \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{6b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \\
 & \hline
 & 2b(a^2 - b^2)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3120 \\
 & \frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \\
 & \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{6b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d} \\
 & \hline
 & 2b(a^2 - b^2)
 \end{aligned}$$

3.577. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{6b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d}$$

$2b(a^2 - b^2)$

↓ 4336

$$\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{b^2} + \frac{6b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d}$$

$2b(a^2 - b^2)$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} (a + b \sin(c + dx + \frac{\pi}{2}))} dx}{b^2} + \frac{6b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d}$$

$2b(a^2 - b^2)$

↓ 3284

$$\frac{a(Ab - aB) \sin(c + dx)}{bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)} - \frac{2(-5a^2B + 3aAb + 2b^2B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{6a^2(-5a^3B + 3a^2Ab + 7ab^2B - 5Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2)}{b^2 d(a+b)} + \frac{6b(-5a^3B + 3a^2Ab + 4ab^2B - 2Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{d}$$

$2b(a^2 - b^2)$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^2*Sec[c + d*x]^(5/2)),x]`

output `(a*(A*b - a*B)*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[Sec[c + d*x]]*(b + a*Sec[c + d*x])) - (-1/3*(((6*b*(3*a^2*A*b - 2*A*b^3 - 5*a^3*B + 4*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*(9*a^3*A*b - 12*a*A*b^3 - 15*a^4*B + 16*a^2*b^2*B + 2*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (6*a^2*(3*a^2*A*b - 5*A*b^3 - 5*a^3*B + 7*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d)/b + (2*(3*a*A*b - 5*a^2*B + 2*b^2*B)*Sin[c + d*x])/(3*b*d*Sqrt[Sec[c + d*x]])/(2*b*(a^2 - b^2))`

3.577. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$

3.577.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3119 `Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`
- rule 3284 `Int[1/(((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`
- rule 3439 `Int[(csc[(e_.) + (f_)*(x_)]*(g_))^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4258 `Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`
- rule 4274 `Int[(csc[(e_.) + (f_)*(x_)]*(d_))^(n_)*(csc[(e_.) + (f_)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4518 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)), x_Symbol] := Simp[b*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[A*(a^2*(m + 1) - b^2*(m + n + 1)) + a*b*B*n - a*(A*b - a*B)*(m + 1)*Csc[e + f*x] + b*(A*b - a*B)*(m + n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && ILtQ[n, 0])`

rule 4592 `Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_), x_Symbol] := Simp[A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*n)), x] + Simp[1/(a*d*n) Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n + 1)*Simp[a*B*n - A*b*(m + n + 1) + a*(A + A*n + C*n)*Csc[e + f*x] + A*b*(m + n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, m}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1]`

rule 4594 `Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))/(Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e + f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]`

3.577.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1065 vs. 2(421) = 842.

Time = 11.30 (sec) , antiderivative size = 1066, normalized size of antiderivative = 2.94

method	result	size
default	Expression too large to display	1066

$$3.577. \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$$

input `int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^2/sec(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)`

output `-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*a^2/b^3*(3*
A*b-4*B*a)/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*
c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipt
icPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a^3*(A*b-B*a)/b^4*(-1/a*b^2/
a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2
)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d
*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(
1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*
c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipti
cE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^
4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(
1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*co
s(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)
)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2/3/b^4/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B*cos(1/2*d*x+1/2*c)*sin(
1/2*d*x+1/2*c)^4*b^2+6*A*a*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*...`

3.577.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(5/2),x, algorith
m="fracas")`

output `Timed out`

3.577. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^2 \sec^{\frac{5}{2}}(c+dx)} dx$

3.577.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**2/sec(d*x+c)**(5/2),x)`output `Timed out`**3.577.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(5/2),x, algorithm="maxima")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2)), x)`**3.577.8 Giac [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^2 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^2/sec(d*x+c)^(5/2),x, algorithm="giac")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^2*sec(d*x + c)^(5/2)), x)`

3.577.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^2 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{5/2} (a + b \cos(c + dx))^2} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^2),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^2),x)`

$$3.578 \quad \int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$$

3.578.1 Optimal result	5261
3.578.2 Mathematica [A] (warning: unable to verify)	5262
3.578.3 Rubi [A] (verified)	5263
3.578.4 Maple [B] (verified)	5270
3.578.5 Fricas [F(-1)]	5271
3.578.6 Sympy [F(-1)]	5272
3.578.7 Maxima [F(-1)]	5272
3.578.8 Giac [F]	5272
3.578.9 Mupad [F(-1)]	5273

3.578.1 Optimal result

Integrand size = 33, antiderivative size = 480

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx =$$

$$\frac{(8a^4A - 29a^2Ab^2 + 15Ab^4 + 9a^3bB - 3ab^3B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{4a^3(a^2 - b^2)^2 d}$$

$$+ \frac{(11a^2Ab - 5Ab^3 - 7a^3B + ab^2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4a^2(a^2 - b^2)^2 d}$$

$$- \frac{(35a^4Ab - 38a^2Ab^3 + 15Ab^5 - 15a^5B + 6a^3b^2B - 3ab^4B) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{4a^3(a - b)^2(a + b)^3 d}$$

$$+ \frac{(8a^4A - 29a^2Ab^2 + 15Ab^4 + 9a^3bB - 3ab^3B) \sqrt{\sec(c + dx)} \sin(c + dx)}{4a^3(a^2 - b^2)^2 d}$$

$$+ \frac{b(Ab - aB) \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{2a(a^2 - b^2) d(b + a \sec(c + dx))^2}$$

$$+ \frac{b(11a^2Ab - 5Ab^3 - 7a^3B + ab^2B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{4a^2(a^2 - b^2)^2 d(b + a \sec(c + dx))}$$

3.578. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$

output $\frac{1}{2} * b * (A * b - B * a) * \sec(d * x + c)^{(5/2)} * \sin(d * x + c) / a / (a^2 - b^2) / d / (b + a * \sec(d * x + c))^{2 + 1/4 * b * (11 * A * a^2 * b - 5 * A * b^3 - 7 * B * a^3 + B * a * b^2) * \sec(d * x + c)^{(3/2)} * \sin(d * x + c) / a^2 / (a^2 - b^2)^2 / d / (b + a * \sec(d * x + c)) + 1/4 * (8 * A * a^4 - 29 * A * a^2 * b^2 + 15 * A * b^4 + 9 * B * a^3 * b - 3 * B * a * b^3) * \sin(d * x + c) * \sec(d * x + c)^{(1/2)} / a^3 / (a^2 - b^2)^2 / d - 1/4 * (8 * A * a^4 - 29 * A * a^2 * b^2 + 15 * A * b^4 + 9 * B * a^3 * b - 3 * B * a * b^3) * (\cos(1/2 * d * x + 1/2 * c))^2)^{(1/2)} / \cos(1/2 * d * x + 1/2 * c) * \text{EllipticE}(\sin(1/2 * d * x + 1/2 * c), 2^{(1/2)}) * \cos(d * x + c)^{(1/2)} * \sec(d * x + c)^{(1/2)} / a^3 / (a^2 - b^2)^2 / d + 1/4 * (11 * A * a^2 * b - 5 * A * b^3 - 7 * B * a^3 + B * a * b^2) * (\cos(1/2 * d * x + 1/2 * c))^2)^{(1/2)} / \cos(1/2 * d * x + 1/2 * c) * \text{EllipticF}(\sin(1/2 * d * x + 1/2 * c), 2^{(1/2)}) * \cos(d * x + c)^{(1/2)} * \sec(d * x + c)^{(1/2)} / a^2 / (a^2 - b^2)^2 / d - 1/4 * (35 * A * a^4 * b - 38 * A * a^2 * b^3 + 15 * A * b^5 - 15 * B * a^5 + 6 * B * a^3 * b^2 - 3 * B * a * b^4) * (\cos(1/2 * d * x + 1/2 * c))^2)^{(1/2)} / \cos(1/2 * d * x + 1/2 * c) * \text{EllipticPi}(\sin(1/2 * d * x + 1/2 * c), 2 * b / (a + b), 2^{(1/2)}) * \cos(d * x + c)^{(1/2)} * \sec(d * x + c)^{(1/2)} / a^3 / (a - b)^2 / (a + b)^3 / d$

3.578.2 Mathematica [A] (warning: unable to verify)

Time = 7.14 (sec) , antiderivative size = 844, normalized size of antiderivative = 1.76

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx =$$

$$\frac{2(56a^4Ab - 95a^2Ab^3 + 45Ab^5 - 16a^5B + 19a^3b^2B - 9ab^4B) \cos^2(c + dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c + dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c + dx)}\right)\right) \right)}{a(a + b \cos(c + dx))(1 - \cos^2(c + dx))} +$$

$$\frac{\sqrt{\sec(c + dx)} \left(\frac{(8a^4A - 29a^2Ab^2 + 15Ab^4 + 9a^3bB - 3ab^3B) \sin(c + dx)}{4a^3(a^2 - b^2)^2} + \frac{Ab^2 \sin(c + dx) - abB \sin(c + dx)}{2a(a^2 - b^2)(a + b \cos(c + dx))^2} + \frac{11a^2Ab^2 \sin(c + dx) - 5Ab^4}{4a^2(c + dx)} \right)}{d}$$

input `Integrate[((A + B * Cos[c + d * x]) * Sec[c + d * x]^(3/2)) / (a + b * Cos[c + d * x])^3, x]`

output

```

-1/16*((2*(56*a^4*A*b - 95*a^2*A*b^3 + 45*A*b^5 - 16*a^5*B + 19*a^3*b^2*B
- 9*a*b^4*B)*Cos[c + d*x]^2*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] - E
llipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*(b + a*Sec[c + d*x])*Sq
rt[1 - Sec[c + d*x]^2]*Sin[c + d*x))/(a*(a + b*Cos[c + d*x])*(1 - Cos[c +
d*x]^2)) + (2*(16*a^5*A - 80*a^3*A*b^2 + 40*a*A*b^4 + 32*a^4*b*B - 8*a^2*b
^3*B)*Cos[c + d*x]^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*(b
+ a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x))/(b*(a + b*Cos[c
+ d*x])*(1 - Cos[c + d*x]^2)) + ((8*a^4*A*b - 29*a^2*A*b^3 + 15*A*b^5 + 9*
a^3*b^2*B - 3*a*b^4*B)*Cos[2*(c + d*x)]*(b + a*Sec[c + d*x])*(-4*a*b + 4*a
*b*Sec[c + d*x]^2 - 4*a*b*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[S
ec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*(2*a - b)*b*EllipticF[ArcSin[Sqr
t[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] - 4*a^2*
EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt
[1 - Sec[c + d*x]^2] + 2*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]]
, -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2])*Sin[c + d*x))/(a*b^2*(a
+ b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)*Sqrt[Sec[c + d*x]]*(2 - Sec[c + d*
x]^2)))/(a^3*(a - b)^2*(a + b)^2*d) + (Sqrt[Sec[c + d*x]]*(((8*a^4*A - 29*
a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B)*Sin[c + d*x]))/(4*a^3*(a^2 -
b^2)^2) + (A*b^2*Sin[c + d*x] - a*b*B*Sin[c + d*x]))/(2*a*(a^2 - b^2)*(a +
b*Cos[c + d*x])^2) + (11*a^2*A*b^2*Sin[c + d*x] - 5*A*b^4*Sin[c + d*x] ...

```

3.578.3 Rubi [A] (verified)

Time = 3.67 (sec) , antiderivative size = 476, normalized size of antiderivative = 0.99, number of steps used = 23, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 3439, 3042, 4517, 27, 3042, 4586, 27, 3042, 4590, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{\sec^{\frac{7}{2}}(c+dx)(A\sec(c+dx)+B)}{(a\sec(c+dx)+b)^3} dx
 \end{aligned}$$

3.578. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+b\cos(c+dx))^3} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{7/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+b\right)^3} dx \\
& \downarrow \text{4517} \\
& \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} - \\
& \frac{\int -\frac{\sec^{\frac{3}{2}}(c+dx)\left((4Aa^2+bBa-5Ab^2)\sec^2(c+dx)-4a(Ab-aB)\sec(c+dx)+3b(Ab-aB)\right)}{2(b+a\sec(c+dx))^2} dx}{2a(a^2-b^2)} \\
& \downarrow \text{27} \\
& \frac{\int \frac{\sec^{\frac{3}{2}}(c+dx)\left((4Aa^2+bBa-5Ab^2)\sec^2(c+dx)-4a(Ab-aB)\sec(c+dx)+3b(Ab-aB)\right)}{(b+a\sec(c+dx))^2} dx}{4a(a^2-b^2)} + \\
& \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} \\
& \downarrow \text{3042} \\
& \frac{\int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left((4Aa^2+bBa-5Ab^2)\csc\left(c+dx+\frac{\pi}{2}\right)^2-4a(Ab-aB)\csc\left(c+dx+\frac{\pi}{2}\right)+3b(Ab-aB)\right)}{(b+a\csc\left(c+dx+\frac{\pi}{2}\right))^2} dx}{4a(a^2-b^2)} + \\
& \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} \\
& \downarrow \text{4586} \\
& \frac{\int \frac{\sqrt{\sec(c+dx)}\left((8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4)\sec^2(c+dx)-4a(-2Ba^3+4Aba^2-b^2Ba-Ab^3)\sec(c+dx)+b(-7Ba^3+11Aba^2+b^2Ba-5Ab^3)\right)}{2(b+a\sec(c+dx))}}{a(a^2-b^2)} dx}{4a(a^2-b^2)} + b \\
& \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} \\
& \downarrow \text{27} \\
& \frac{\int \frac{\sqrt{\sec(c+dx)}\left((8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4)\sec^2(c+dx)-4a(-2Ba^3+4Aba^2-b^2Ba-Ab^3)\sec(c+dx)+b(-7Ba^3+11Aba^2+b^2Ba-5Ab^3)\right)}{b+a\sec(c+dx)}}{2a(a^2-b^2)} dx}{4a(a^2-b^2)} + b \\
& \frac{b(Ab-aB)\sin(c+dx)\sec^{\frac{5}{2}}(c+dx)}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} \\
& \downarrow \text{3042}
\end{aligned}$$

3.578. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+b\cos(c+dx))^3} dx$

$$\int \frac{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(\left(8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4\right)\csc\left(c+dx+\frac{\pi}{2}\right)^2-4a\left(-2Ba^3+4Aba^2-b^2Ba-Ab^3\right)\csc\left(c+dx+\frac{\pi}{2}\right)+b\left(-7Ba^3+11Aba^2+b^2Ba-5Ab^3\right)}{b+a\csc\left(c+dx+\frac{\pi}{2}\right)} dx$$

$$\frac{2a\left(a^2-b^2\right)}{2a\left(a^2-b^2\right)}$$

$$4a\left(a^2-b^2\right)$$

$$\frac{b\left(Ab-aB\right)\sin\left(c+dx\right)\sec^{\frac{5}{2}}\left(c+dx\right)}{2ad\left(a^2-b^2\right)\left(a\sec\left(c+dx\right)+b\right)^2}$$

↓ 4590

$$2\int -\frac{\left(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2-3b^4Ba+15Ab^5\right)\sec^2\left(c+dx\right)+4a\left(2Aa^4+4bBa^3-10Ab^2a^2-b^3Ba+5Ab^4\right)\sec\left(c+dx\right)+b\left(8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4\right)}{2\sqrt{\sec\left(c+dx\right)}\left(b+a\sec\left(c+dx\right)\right)} dx$$

$$\frac{2a\left(a^2-b^2\right)}{2a\left(a^2-b^2\right)}$$

$$2a\left(a^2-b^2\right)$$

$$4a\left(a^2-b^2\right)$$

$$\frac{b\left(Ab-aB\right)\sin\left(c+dx\right)\sec^{\frac{5}{2}}\left(c+dx\right)}{2ad\left(a^2-b^2\right)\left(a\sec\left(c+dx\right)+b\right)^2}$$

↓ 27

$$2\left(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4\right)\sin\left(c+dx\right)\sqrt{\sec\left(c+dx\right)} - \int \frac{\left(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2-3b^4Ba+15Ab^5\right)\sec^2\left(c+dx\right)+4a\left(2Aa^4+4bBa^3-10Ab^2a^2-b^3Ba+5Ab^4\right)\sec\left(c+dx\right)+b\left(8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4\right)}{\sqrt{\sec\left(c+dx\right)}\left(b+a\sec\left(c+dx\right)\right)} dx$$

$$\frac{2a\left(a^2-b^2\right)}{2a\left(a^2-b^2\right)}$$

$$4a\left(a^2-b^2\right)$$

$$\frac{b\left(Ab-aB\right)\sin\left(c+dx\right)\sec^{\frac{5}{2}}\left(c+dx\right)}{2ad\left(a^2-b^2\right)\left(a\sec\left(c+dx\right)+b\right)^2}$$

↓ 3042

$$2\left(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4\right)\sin\left(c+dx\right)\sqrt{\sec\left(c+dx\right)} - \int \frac{\left(-8Ba^5+24Aba^4+5b^2Ba^3-33Ab^3a^2-3b^4Ba+15Ab^5\right)\csc\left(c+dx+\frac{\pi}{2}\right)^2+4a\left(2Aa^4+4bBa^3-10Ab^2a^2-b^3Ba+5Ab^4\right)\csc\left(c+dx+\frac{\pi}{2}\right)+b\left(8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4\right)}{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}\left(b+a\csc\left(c+dx+\frac{\pi}{2}\right)\right)} dx$$

$$\frac{2a\left(a^2-b^2\right)}{2a\left(a^2-b^2\right)}$$

$$2a\left(a^2-b^2\right)$$

$$4a\left(a^2-b^2\right)$$

$$\frac{b\left(Ab-aB\right)\sin\left(c+dx\right)\sec^{\frac{5}{2}}\left(c+dx\right)}{2ad\left(a^2-b^2\right)\left(a\sec\left(c+dx\right)+b\right)^2}$$

↓ 4594

$$2\left(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4\right)\sin\left(c+dx\right)\sqrt{\sec\left(c+dx\right)} - \int \frac{b^2\left(8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4\right)-ab^2\left(-7Ba^3+11Aba^2+b^2Ba-5Ab^3\right)\sec\left(c+dx\right)}{\sqrt{\sec\left(c+dx\right)}b^2}$$

$$\frac{2a\left(a^2-b^2\right)}{2a\left(a^2-b^2\right)}$$

$$2a\left(a^2-b^2\right)$$

$$4a\left(a^2-b^2\right)$$

$$\frac{b\left(Ab-aB\right)\sin\left(c+dx\right)\sec^{\frac{5}{2}}\left(c+dx\right)}{2ad\left(a^2-b^2\right)\left(a\sec\left(c+dx\right)+b\right)^2}$$

↓ 3042

3.578. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{\int \frac{b^2(8Aa^4+9bBa^3-29Ab^2a^2-3b^3Ba+15Ab^4) - ab^2(-7Ba^3+11Aba^2+b^2Ba-5Ab^3) \csc(c+dx)}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b^2}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

$$2a(a^2 - b^2)$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4274

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \int \frac{1}{\sqrt{\sec(c+dx)}} dx - ab^2(-7a^3B+11a^2Ab+ab^2B)}{b^2}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

$$2a(a^2 - b^2)$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx - ab^2(-7a^3B+11a^2Ab+ab^2B)}{b^2}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

$$2a(a^2 - b^2)$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4258

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - ab^2(-7a^3B+11a^2Ab+ab^2B)}{b^2}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

$$2a(a^2 - b^2)$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx - ab^2(-7a^3B+11a^2Ab+ab^2B)}{b^2}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

$$2a(a^2 - b^2)$$

$$4a(a^2 - b^2)$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

3.578. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$

↓ 3119

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{2b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d} - \frac{ab^2}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3120

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(-15a^5B+35a^4Ab+6a^3b^2B-38a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}}{b+a \csc(c+dx+\frac{\pi}{2})} dx + \frac{2b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4336

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(-15a^5B+35a^4Ab+6a^3b^2B-38a^2Ab^3-3ab^4B+15Ab^5) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{(-15a^5B+35a^4Ab+6a^3b^2B-38a^2Ab^3-3ab^4B+15Ab^5) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx)}} dx + \frac{2b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3284

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{5}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} + \frac{b(-7a^3B+11a^2Ab+ab^2B-5Ab^3) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{ad(a^2-b^2)(a \sec(c+dx)+b)} + \frac{2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad} - \frac{2b^2(8a^4A+9a^3bB-29a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{d}$$

3.578. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^3,x]`

output `(b*(A*b - a*B)*Sec[c + d*x]^(5/2)*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*(b + a*Sec[c + d*x])^2) + ((b*(11*a^2*A*b - 5*A*b^3 - 7*a^3*B + a*b^2*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(a*(a^2 - b^2)*d*(b + a*Sec[c + d*x])) + (-((((2*b^2*(8*a^4*A - 29*a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*a*b^2*(11*a^2*A*b - 5*A*b^3 - 7*a^3*B + a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (2*(35*a^4*A*b - 38*a^2*A*b^3 + 15*A*b^5 - 15*a^5*B + 6*a^3*b^2*B - 3*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/((a + b)*d))/a + (2*(8*a^4*A - 29*a^2*A*b^2 + 15*A*b^4 + 9*a^3*b*B - 3*a*b^3*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*d))/(2*a*(a^2 - b^2))/(4*a*(a^2 - b^2))`

3.578.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

3.578.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(g_.)^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\sin[c + d*x]^n \text{Int}[1/\sin[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4336 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(3/2)}/(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[d*\text{Sqrt}[d*\sin[e + f*x]]*\text{Sqrt}[d*\text{Csc}[e + f*x]] \text{Int}[1/(\text{Sqrt}[d*\sin[e + f*x]]*(b + a*\sin[e + f*x])), x], x] /; \text{FreeQ}[\{a, b, d, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 4517 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[a*d^2*(A*b - a*B)*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m+1)}*((d*\text{Csc}[e + f*x])^{(n-2)}/(b*f*(m+1)*(a^2 - b^2))), x] - \text{Simp}[d/(b*(m+1)*(a^2 - b^2)) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m+1)}*(d*\text{Csc}[e + f*x])^{(n-2)}*\text{Simp}[a*d*(A*b - a*B)*(n-2) + b*d*(A*b - a*B)*(m+1)*\text{Csc}[e + f*x] - (a*A*b*d*(m+n) - d*B*(a^2*(n-1) + b^2*(m+1)))*\text{Csc}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, d, e, f, A, B\}, x] \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -1] \&\& \text{GtQ}[n, 1]$

```
rule 4586 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(-d)*(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a +
b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 1)/(b*f*(a^2 - b^2)*(m + 1)
), x] + Simp[d/(b*(a^2 - b^2)*(m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(
d*Csc[e + f*x])^(n - 1)*Simp[A*b^2*(n - 1) - a*(b*B - a*C)*(n - 1) + b*(a*A
- b*B + a*C)*(m + 1)*Csc[e + f*x] - (b*(A*b - a*B)*(m + n + 1) + C*(a^2*n
+ b^2*(m + 1)))*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, C
}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

```
rule 4590 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(-C)*d*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1
)*((d*Csc[e + f*x])^(n - 1)/(b*f*(m + n + 1))), x] + Simp[d/(b*(m + n + 1)
) Int[(a + b*Csc[e + f*x])^m*(d*Csc[e + f*x])^(n - 1)*Simp[a*C*(n - 1) + (
A*b*(m + n + 1) + b*C*(m + n))*Csc[e + f*x] + (b*B*(m + n + 1) - a*C*n)*Csc
[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, C, m}, x] && NeQ[a^2
- b^2, 0] && GtQ[n, 0]
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.578.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1974 vs. $2(528) = 1056$.

Time = 15.52 (sec) , antiderivative size = 1975, normalized size of antiderivative = 4.11

method	result	size
default	Expression too large to display	1975

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

$$3.578. \quad \int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^3} dx$$

output
$$\begin{aligned} & -(-(-2\cos(1/2dx+1/2c)^2+1)\sin(1/2dx+1/2c)^2)^{1/2}*(2A/a^3/\sin(1/2dx+1/2c)^2/(2\sin(1/2dx+1/2c)^2-1)*(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*(2\sin(1/2dx+1/2c)^2\cos(1/2dx+1/2c)-(\sin(1/2dx+1/2c)^2)^{1/2}*(2\sin(1/2dx+1/2c)^2-1)^{1/2}*\text{EllipticE}(\cos(1/2dx+1/2c),2^{1/2})) \\ & +4Ab^2/a^3/(-2ab+2b^2)*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticPi}(\cos(1/2dx+1/2c),-2b/(a-b),2^{1/2}) \\ & -2Ab/a^2*(-1/ab^2/(a^2-b^2)\cos(1/2dx+1/2c)*(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}/(2b\cos(1/2dx+1/2c)^2+a-b)-1/2a/(a+b)*(\sin(1/2dx+1/2c)^2)^{1/2} \\ & *(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2}*\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2}) \\ & -1/2/(a^2-b^2)*b/a*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2} \\ & *\text{EllipticF}(\cos(1/2dx+1/2c),2^{1/2}) \\ & +1/2/(a^2-b^2)*b/a*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2} \\ & *\text{EllipticE}(\cos(1/2dx+1/2c),2^{1/2}) \\ & -3a/(a^2-b^2)/(-2ab+2b^2)*b*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2} \\ & *\text{EllipticPi}(\cos(1/2dx+1/2c),-2b/(a-b),2^{1/2}) \\ & +1/a/(a^2-b^2)/(-2ab+2b^2)*b^3*(\sin(1/2dx+1/2c)^2)^{1/2}*(-2\cos(1/2dx+1/2c)^2+1)^{1/2}/(-2\sin(1/2dx+1/2c)^4+\sin(1/2dx+1/2c)^2)^{1/2} \end{aligned}$$

3.578.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^3,x, algorithm="fracas")`

output `Timed out`

3.578.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**3,x)`

output `Timed out`

3.578.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^3,x, algorithm="maxima")`

output `Timed out`

3.578.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^3, x)`

3.578.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{3/2}}{(a + b \cos(c + dx))^3} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^3,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^3,x)`

$$3.579 \quad \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$$

3.579.1 Optimal result	5274
3.579.2 Mathematica [A] (warning: unable to verify)	5275
3.579.3 Rubi [A] (verified)	5276
3.579.4 Maple [B] (verified)	5282
3.579.5 Fracas [F(-1)]	5283
3.579.6 Sympy [F]	5284
3.579.7 Maxima [F(-1)]	5284
3.579.8 Giac [F]	5284
3.579.9 Mupad [F(-1)]	5285

3.579.1 Optimal result

Integrand size = 33, antiderivative size = 405

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx$$

$$= - \frac{(9a^2 Ab - 3Ab^3 - 5a^3 B - ab^2 B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{4a^2 (a^2 - b^2)^2 d}$$

$$- \frac{(7a^2 Ab - Ab^3 - 3a^3 B - 3ab^2 B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4ab (a^2 - b^2)^2 d}$$

$$+ \frac{(15a^4 Ab - 6a^2 Ab^3 + 3Ab^5 - 3a^5 B - 10a^3 b^2 B + ab^4 B) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4a^2 (a - b)^2 b (a + b)^3 d}$$

$$+ \frac{b(Ab - aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2a (a^2 - b^2) d (b + a \sec(c + dx))^2}$$

$$+ \frac{b(9a^2 Ab - 3Ab^3 - 5a^3 B - ab^2 B) \sqrt{\sec(c + dx)} \sin(c + dx)}{4a^2 (a^2 - b^2)^2 d (b + a \sec(c + dx))}$$

output $\frac{1}{2}b(Ab - Ba) \sec(dx+c)^{3/2} \sin(dx+c) / (a^2 - b^2) / d / (b + a \sec(dx+c))^{2+1/4} b^4 (9Aa^2b - 3A^2b^3 - 5B^2a^3 - B^2ab^2) \sin(dx+c) \sec(dx+c)^{1/2} / a^2 / (a^2 - b^2)^2 / d / (b + a \sec(dx+c)) - 1/4 (9Aa^2b - 3A^2b^3 - 5B^2a^3 - B^2ab^2) (\cos(1/2 dx + 1/2 c))^2)^{1/2} / \cos(1/2 dx + 1/2 c) \text{EllipticE}(\sin(1/2 dx + 1/2 c), 2^{1/2}) \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / a^2 / (a^2 - b^2)^2 / d - 1/4 (7Aa^2b - Ab^3 - 3B^2a^3 - 3B^2ab^2) (\cos(1/2 dx + 1/2 c))^2)^{1/2} / \cos(1/2 dx + 1/2 c) \text{EllipticF}(\sin(1/2 dx + 1/2 c), 2^{1/2}) \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / a/b / (a^2 - b^2)^2 / d + 1/4 (15Aa^4b - 6Aa^2b^3 + 3A^2b^5 - 3B^2a^5 - 10B^2a^3b^2 + B^2ab^4) (\cos(1/2 dx + 1/2 c))^2)^{1/2} / \cos(1/2 dx + 1/2 c) \text{EllipticPi}(\sin(1/2 dx + 1/2 c), 2b/(a+b), 2^{1/2}) \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / a^2 / (a-b)^2 / b / (a+b)^3 / d$

3.579.2 Mathematica [A] (warning: unable to verify)

Time = 6.98 (sec) , antiderivative size = 797, normalized size of antiderivative = 1.97

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx$$

$$= \frac{2(16a^4A - 19a^2Ab^2 + 9Ab^4 - 9a^3bB + 3ab^3B) \cos^2(c+dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c+dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c+dx)}\right), -1\right) \right) (b+a)}{a(a+b \cos(c+dx))(1 - \cos^2(c+dx))} + \frac{\sqrt{\sec(c + dx)} \left(-\frac{(-9a^2Ab + 3Ab^3 + 5a^3B + ab^2B) \sin(c+dx)}{4a^2(a^2 - b^2)^2} + \frac{-Ab \sin(c+dx) + aB \sin(c+dx)}{2(a^2 - b^2)(a + b \cos(c+dx))^2} + \frac{-7a^2Ab \sin(c+dx) + Ab^3 \sin(c+dx)}{4a(a^2 - b^2)^2(a} \right)}{d}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^3, x]`

output $((2*(16*a^4*A - 19*a^2*A*b^2 + 9*A*b^4 - 9*a^3*b*B + 3*a*b^3*B)*\text{Cos}[c + d*x]^2*(\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1] - \text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1])*(b + a*\text{Sec}[c + d*x])* \text{Sqrt}[1 - \text{Sec}[c + d*x]^2]*\text{Sin}[c + d*x])/(a*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)) + (2*(-32*a^3*A*b + 8*a*A*b^3 + 16*a^4*B + 8*a^2*b^2*B)*\text{Cos}[c + d*x]^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*(b + a*\text{Sec}[c + d*x])* \text{Sqrt}[1 - \text{Sec}[c + d*x]^2]*\text{Sin}[c + d*x])/(b*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)) + ((-9*a^2*A*b^2 + 3*A*b^4 + 5*a^3*b*B + a*b^3*B)*\text{Cos}[2*(c + d*x)]*(b + a*\text{Sec}[c + d*x])*(-4*a*b + 4*a*b*\text{Sec}[c + d*x]^2 - 4*a*b*\text{EllipticE}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] + 2*(2*a - b)*b*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] - 4*a^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] + 2*b^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2])* \text{Sin}[c + d*x])/(a*b^2*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)*\text{Sqrt}[\text{Sec}[c + d*x]]*(2 - \text{Sec}[c + d*x]^2)))/(16*a^2*(a - b)^2*(a + b)^2*d + (\text{Sqrt}[\text{Sec}[c + d*x]]*(-1/4*((-9*a^2*A*b + 3*A*b^3 + 5*a^3*B + a*b^2*B)*\text{Sin}[c + d*x])/(a^2*(a^2 - b^2)^2) + (-(A*b*\text{Sin}[c + d*x]) + a*B*\text{Sin}[c + d*x])/(2*(a^2 - b^2)*(a + b*\text{Cos}[c + d*x])^2) + (-7*a^2*A*b*\text{Sin}[c + d*x] + A*b^3*\text{Sin}[c + d*x] + 3*a^3*B*\text{Sin}[c + d*x] + 3*a*b^2*B*\text{Sin}[c + d*x]))/(4*a*(a^2 - b^2)^2*(a + ...$

3.579.3 Rubi [A] (verified)

Time = 2.83 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4517, 27, 3042, 4586, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{(a+b\cos(c+dx))^3} dx$$

↓ 3042

$$\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^3} dx$$

↓ 3439

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(A\sec(c+dx)+B)}{(a\sec(c+dx)+b)^3} dx$$

3.579. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b\cos(c+dx))^3} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \int \frac{\csc(c+dx+\frac{\pi}{2})^{5/2} (A \csc(c+dx+\frac{\pi}{2}) + B)}{(a \csc(c+dx+\frac{\pi}{2}) + b)^3} dx \\
 & \downarrow \text{4517} \\
 & \frac{b(Ab - aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{2ad(a^2 - b^2)(a \sec(c+dx) + b)^2} - \\
 & \frac{\int -\frac{\sqrt{\sec(c+dx)}((4Aa^2 - bBa - 3Ab^2) \sec^2(c+dx) - 4a(Ab - aB) \sec(c+dx) + b(Ab - aB))}{2(b+a \sec(c+dx))^2} dx}{2a(a^2 - b^2)} \\
 & \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{\sec(c+dx)}((4Aa^2 - bBa - 3Ab^2) \sec^2(c+dx) - 4a(Ab - aB) \sec(c+dx) + b(Ab - aB))}{(b+a \sec(c+dx))^2} dx}{4a(a^2 - b^2)} + \\
 & \frac{b(Ab - aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{2ad(a^2 - b^2)(a \sec(c+dx) + b)^2} \\
 & \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})}((4Aa^2 - bBa - 3Ab^2) \csc(c+dx+\frac{\pi}{2})^2 - 4a(Ab - aB) \csc(c+dx+\frac{\pi}{2}) + b(Ab - aB))}{(b+a \csc(c+dx+\frac{\pi}{2}))^2} dx}{4a(a^2 - b^2)} + \\
 & \frac{b(Ab - aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{2ad(a^2 - b^2)(a \sec(c+dx) + b)^2} \\
 & \downarrow \text{4586} \\
 & \frac{\int -\frac{((8Aa^4 - 7bBa^3 - 5Ab^2a^2 + b^3Ba + 3Ab^4) \sec^2(c+dx) + 4a(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sec(c+dx) + b(-5Ba^3 + 9Aba^2 - b^2Ba - 3Ab^3))}{2\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx}{a(a^2 - b^2)} + \frac{b(-5a^3B + 9a^2Ab - ab^2B - 3Ab^3)}{4a(a^2 - b^2)} \\
 & \frac{b(Ab - aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{2ad(a^2 - b^2)(a \sec(c+dx) + b)^2} \\
 & \downarrow \text{27} \\
 & \frac{b(-5a^3B + 9a^2Ab - ab^2B - 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{\int -\frac{((8Aa^4 - 7bBa^3 - 5Ab^2a^2 + b^3Ba + 3Ab^4) \sec^2(c+dx) + 4a(-2Ba^3 + 4Aba^2 - b^2Ba - Ab^3) \sec(c+dx) + b(-5Ba^3 + 9Aba^2 - b^2Ba - 3Ab^3))}{\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx}{2a(a^2 - b^2)} \\
 & \frac{b(Ab - aB) \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{2ad(a^2 - b^2)(a \sec(c+dx) + b)^2} \\
 & \downarrow \text{3042}
 \end{aligned}$$

3.579. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{\int \frac{(-8Aa^4+7bBa^3+5Ab^2a^2-b^3Ba-3Ab^4) \csc(c+dx+\frac{\pi}{2})^2+4a(-2Ba^3+4Aba^2-b^2Ba-Ab^3)}{\sqrt{\csc(c+dx+\frac{\pi}{2})(b+a \csc(c+dx+\frac{\pi}{2}))}} dx}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4594

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{\int \frac{(-5Ba^3+9Aba^2-b^2Ba-3Ab^3)b^2+a(-3Ba^3+7Aba^2-3b^2Ba-Ab^3) \sec(c+dx)b}{\sqrt{\sec(c+dx)}} dx}{b^2} - \frac{(-3a^5)}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{\int \frac{(-5Ba^3+9Aba^2-b^2Ba-3Ab^3)b^2+a(-3Ba^3+7Aba^2-3b^2Ba-Ab^3) \csc(c+dx+\frac{\pi}{2})b}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx}{b^2} - \frac{(-3a^5)}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4274

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \int \frac{1}{\sqrt{\sec(c+dx)}} dx + ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \int \sqrt{\sec(c+dx)}}{b^2} - \frac{2a(a^2-b^2)}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \int \frac{1}{\sqrt{\csc(c+dx+\frac{\pi}{2})}} dx + ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \int \sqrt{\csc(c+dx+\frac{\pi}{2})}}{b^2} - \frac{2a(a^2-b^2)}{2a(a^2-b^2)}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4258

3.579. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx + ab(-3a^3B+7a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{b^2}$$

$4a(a^2 - b^2)$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx+\frac{\pi}{2})} dx + ab(-3a^3B+7a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{b^2}$$

$4a(a^2 - b^2)$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3119

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{b^2}}{b^2}$$

$4a(a^2 - b^2)$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3120

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{2ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{2b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{b^2}}$$

$4a(a^2 - b^2)$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4336

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a \sec(c+dx)+b)} - \frac{2ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticF}(\frac{1}{2}(c+dx), 2)}{d} + \frac{2b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)}}{b^2}}$$

$$\frac{b(Ab - aB) \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

3.579. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \frac{2ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2}$$

$$\frac{b(Ab - aB)\sin(c + dx)\sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a\sec(c + dx) + b)^2}$$

↓ 3284

$$\frac{b(Ab - aB)\sin(c + dx)\sec^{\frac{3}{2}}(c + dx)}{2ad(a^2 - b^2)(a\sec(c + dx) + b)^2} + \frac{2ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2}$$

$$\frac{b(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{ad(a^2-b^2)(a\sec(c+dx)+b)} - \frac{2ab(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\operatorname{EllipticF}\left(\frac{1}{2}(c+dx),2\right)}{d} + \frac{2b^2(-5a^3B+9a^2Ab-ab^2B-3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2}$$

4

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^3,x]`

output `(b*(A*b - a*B)*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*(b + a*Sec[c + d*x])^2) + (-1/2*((2*b^2*(9*a^2*A*b - 3*A*b^3 - 5*a^3*B - a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*b*(7*a^2*A*b - A*b^3 - 3*a^3*B - 3*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 - (2*(15*a^4*A*b - 6*a^2*A*b^3 + 3*A*b^5 - 3*a^5*B - 10*a^3*b^2*B + a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b*(a + b)*d)/(a*(a^2 - b^2)) + (b*(9*a^2*A*b - 3*A*b^3 - 5*a^3*B - a*b^2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*(a^2 - b^2)*d*(b + a*Sec[c + d*x]))/(4*a*(a^2 - b^2))`

3.579.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.579. $\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$

rule 3120 $\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3284 $\text{Int}[1/((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)])*\text{Sqrt}[(c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/(f*(a + b)*\text{Sqrt}[c + d]))*\text{EllipticPi}[2*(b/(a + b)), (1/2)*(e - \text{Pi}/2 + f*x), 2*(d/(c + d))], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[c + d, 0]$

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(g_.))^{(p_.)*((a_.) + (b_.)*\sin[(e_.) + (f_.)(x_)]^{(m_.)*((c_.) + (d_.)*\sin[(e_.) + (f_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m + n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p - m - n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(n_.)*(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n + 1)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x]$

rule 4336 $\text{Int}[(\text{csc}[(e_.) + (f_.)(x_)]*(d_.))^{(3/2)}/(\text{csc}[(e_.) + (f_.)(x_)]*(b_.) + (a_.)), x_Symbol] \rightarrow \text{Simp}[d*\text{Sqrt}[d*\text{Sin}[e + f*x]]*\text{Sqrt}[d*\text{Csc}[e + f*x]] \text{Int}[1/(\text{Sqrt}[d*\text{Sin}[e + f*x]]*(b + a*\text{Sin}[e + f*x])), x], x] /; \text{FreeQ}[\{a, b, d, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$


```
rule 4517 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*d^2*(
A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n -
2)/(b*f*(m + 1)*(a^2 - b^2))), x] - Simp[d/(b*(m + 1)*(a^2 - b^2)) Int[(
a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 2)*Simp[a*d*(A*b - a*B)*(
n - 2) + b*d*(A*b - a*B)*(m + 1)*Csc[e + f*x] - (a*A*b*d*(m + n) - d*B*(a^2
*(n - 1) + b^2*(m + 1)))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f
, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && GtQ[
n, 1]
```

```
rule 4586 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(-d)*(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a +
b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 1)/(b*f*(a^2 - b^2)*(m + 1)
), x] + Simp[d/(b*(a^2 - b^2)*(m + 1)) Int[(a + b*Csc[e + f*x])^(m + 1)*(
d*Csc[e + f*x])^(n - 1)*Simp[A*b^2*(n - 1) - a*(b*B - a*C)*(n - 1) + b*(a*A
- b*B + a*C)*(m + 1)*Csc[e + f*x] - (b*(A*b - a*B)*(m + n + 1) + C*(a^2*n
+ b^2*(m + 1)))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C
}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.579.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1743 vs. 2(457) = 914.

Time = 12.60 (sec) , antiderivative size = 1744, normalized size of antiderivative = 4.31

method	result	size
default	Expression too large to display	1744

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^3,x,method=_RETURNV
ERBOSE)
```

$$3.579. \quad \int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^3} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B/b*(-1/a*b^
2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)
^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2
)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2
*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b
/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),
2^(1/2))+1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/
2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellip
ticE(cos(1/2*d*x+1/2*c),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c
)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2
^(1/2))+1/a/(a^2-b^2)/(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*
cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^
2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2)))+2*(A*b-B*a)/b*
(-1/2/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*
d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)^2-3/4*b^2*(3*a^2-b^2)/a
^2/(a^2-b^2)^2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2
*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-7/8/(a+b)/(a^2-b^2)*(sin(1/2*d
*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/...

```

3.579.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^3,x, algorithm="fricas")`

output `Timed out`

3.579.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**3,x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/(a + b*cos(c + d*x))**3, x)`

3.579.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^3,x, algorithm m="maxima")`

output `Timed out`

3.579.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^3} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^3, x)`

3.579.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^3} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + b \cos(c + dx))^3} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^3,x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^3,x)`

3.580
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$$

3.580.1 Optimal result 5286
 3.580.2 Mathematica [A] (warning: unable to verify) 5287
 3.580.3 Rubi [A] (verified) 5288
 3.580.4 Maple [B] (verified) 5294
 3.580.5 Fricas [F(-1)] 5295
 3.580.6 Sympy [F(-1)] 5296
 3.580.7 Maxima [F] 5296
 3.580.8 Giac [F] 5296
 3.580.9 Mupad [F(-1)] 5297

3.580.1 Optimal result

Integrand size = 33, antiderivative size = 402

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{(5a^2Ab + Ab^3 - a^3B - 5ab^2B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{4ab(a^2 - b^2)^2 d}$$

$$+ \frac{(3a^2Ab + 3Ab^3 + a^3B - 7ab^2B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4b^2(a^2 - b^2)^2 d}$$

$$- \frac{(3a^4Ab + 10a^2Ab^3 - Ab^5 + a^5B - 10a^3b^2B - 3ab^4B) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4a(a - b)^2 b^2 (a + b)^3 d}$$

$$+ \frac{b(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{2a(a^2 - b^2) d (b + a \sec(c + dx))^2}$$

$$- \frac{(7a^2Ab - Ab^3 - 3a^3B - 3ab^2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{4a(a^2 - b^2)^2 d (b + a \sec(c + dx))}$$

output $\frac{1}{2}b(Ab - B^2a) \sin(dx+c) \sec(dx+c)^{1/2} / a(a^2 - b^2) / d + (b + a \sec(dx+c))^{-2} - \frac{1}{4}(7Aa^2b - Ab^3 - 3B^2a^3 - 3B^2ab^2) \sin(dx+c) \sec(dx+c)^{1/2} / a(a^2 - b^2)^2 / d + (b + a \sec(dx+c)) + \frac{1}{4}(5Aa^2b + Ab^3 - B^2a^3 - 5B^2ab^2) (\cos(1/2 dx + 1/2 c))^2)^{1/2} / \cos(1/2 dx + 1/2 c) * \text{EllipticE}(\sin(1/2 dx + 1/2 c), 2^{1/2}) * \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / a/b(a^2 - b^2)^2 / d + \frac{1}{4}(3Aa^2b + 3Ab^3 + B^2a^3 - 7B^2ab^2) (\cos(1/2 dx + 1/2 c))^2)^{1/2} / \cos(1/2 dx + 1/2 c) * \text{EllipticF}(\sin(1/2 dx + 1/2 c), 2^{1/2}) * \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / b^2 / (a^2 - b^2)^2 / d - \frac{1}{4}(3Aa^4b + 10Aa^2b^3 - Ab^5 + B^2a^5 - 10B^2a^3b^2 - 3B^2ab^4) (\cos(1/2 dx + 1/2 c))^2)^{1/2} / \cos(1/2 dx + 1/2 c) * \text{EllipticPi}(\sin(1/2 dx + 1/2 c), 2b/(a+b), 2^{1/2}) * \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / a(a-b)^2 / b^2 / (a+b)^3 / d$

3.580.2 Mathematica [A] (warning: unable to verify)

Time = 6.95 (sec) , antiderivative size = 784, normalized size of antiderivative = 1.95

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx$$

$$= \frac{2(-9a^2Ab + 3Ab^3 + 5a^3B + ab^2B) \cos^2(c+dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c+dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c+dx)}\right), -1\right) \right) (b+a \sec(c+dx))}{a(a+b \cos(c+dx))(1-\cos^2(c+dx))} + \frac{\sqrt{\sec(c+dx)} \left(\frac{(-5a^2Ab - Ab^3 + a^3B + 5ab^2B) \sin(c+dx)}{4ab(a^2 - b^2)^2} - \frac{aAb \sin(c+dx) - a^2B \sin(c+dx)}{2b(-a^2 + b^2)(a+b \cos(c+dx))^2} + \frac{3a^2Ab \sin(c+dx) + 3Ab^3 \sin(c+dx) + a^3}{4b(-a^2 + b^2)^2(a+b \cos(c+dx))} \right)}{d}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sqrt[Sec[c + d*x]]),x]`

output

```
((2*(-9*a^2*A*b + 3*A*b^3 + 5*a^3*B + a*b^2*B)*Cos[c + d*x]^2*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] - EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(a*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)) + (2*(16*a^3*A + 8*a*A*b^2 - 24*a^2*b*B)*Cos[c + d*x]^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(b*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)) + ((5*a^2*A*b + A*b^3 - a^3*B - 5*a*b^2*B)*Cos[2*(c + d*x)]*(b + a*Sec[c + d*x])*(-4*a*b + 4*a*b*Sec[c + d*x]^2 - 4*a*b*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*(2*a - b)*b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] - 4*a^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2])*Sin[c + d*x])/(a*b^2*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)*Sqrt[Sec[c + d*x]]*(2 - Sec[c + d*x]^2))/(16*a*(a - b)^2*(a + b)^2*d) + (Sqrt[Sec[c + d*x]]*(((5*a^2*A*b - A*b^3 + a^3*B + 5*a*b^2*B)*Sin[c + d*x])/(4*a*b*(a^2 - b^2)^2) - (a*A*b*SIN[c + d*x] - a^2*B*SIN[c + d*x])/(2*b*(-a^2 + b^2)*(a + b*Cos[c + d*x])^2) + (3*a^2*A*b*SIN[c + d*x] + 3*A*b^3*SIN[c + d*x] + a^3*B*SIN[c + d*x] - 7*a*b^2*B*SIN[c + d*x])/(4*b*(-a^2 + b^2)^2*(a + b*Cos[c + d*x]))))/d
```

3.580.3 Rubi [A] (verified)

Time = 2.74 (sec) , antiderivative size = 397, normalized size of antiderivative = 0.99, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4517, 27, 3042, 4588, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))^3} dx$$

↓ 3439

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(A \sec(c + dx) + B)}{(a \sec(c + dx) + b)^3} dx$$

3.580. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{3/2}\left(A\csc\left(c+dx+\frac{\pi}{2}\right)+B\right)}{\left(a\csc\left(c+dx+\frac{\pi}{2}\right)+b\right)^3} dx \\
& \downarrow 4517 \\
& \frac{\frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} - \int \frac{-((4Aa^2-3bBa-Ab^2)\sec^2(c+dx))+4a(Ab-aB)\sec(c+dx)+b(Ab-aB)}{2\sqrt{\sec(c+dx)}(b+a\sec(c+dx))^2} dx}{2a(a^2-b^2)} \\
& \downarrow 27 \\
& \frac{\frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} - \int \frac{-((4Aa^2-3bBa-Ab^2)\sec^2(c+dx))+4a(Ab-aB)\sec(c+dx)+b(Ab-aB)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))^2} dx}{4a(a^2-b^2)} \\
& \downarrow 3042 \\
& \frac{\frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} - \int \frac{(-4Aa^2+3bBa+Ab^2)\csc\left(c+dx+\frac{\pi}{2}\right)^2+4a(Ab-aB)\csc\left(c+dx+\frac{\pi}{2}\right)+b(Ab-aB)}{\sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)}(b+a\csc\left(c+dx+\frac{\pi}{2}\right))^2} dx}{4a(a^2-b^2)} \\
& \downarrow 4588 \\
& \frac{\frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} - \int \frac{-b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\sec^2(c+dx)+4ab(2Aa^2-3bBa+Ab^2)\sec(c+dx)+b(-Ba^3+5Aba^2-5b^2Ba+Ab^3)}{2\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx}{b(a^2-b^2)} + \frac{(-3a^3B+7a^2Ab-3ab^2B-Ab^3)}{d(a^2-b^2)(a\sec(c+dx)+b)} \\
& \downarrow 27 \\
& \frac{\frac{b(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2ad(a^2-b^2)(a\sec(c+dx)+b)^2} - \int \frac{-b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\sec^2(c+dx)+4ab(2Aa^2-3bBa+Ab^2)\sec(c+dx)+b(-Ba^3+5Aba^2-5b^2Ba+Ab^3)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx}{d(a^2-b^2)(a\sec(c+dx)+b)} - \frac{(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a^2-b^2)(a\sec(c+dx)+b)} \\
& \downarrow 3042 \\
& \frac{(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a^2-b^2)(a\sec(c+dx)+b)} - \int \frac{-b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\sec^2(c+dx)+4ab(2Aa^2-3bBa+Ab^2)\sec(c+dx)+b(-Ba^3+5Aba^2-5b^2Ba+Ab^3)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx}{2b(a^2-b^2)} \\
& \downarrow 3042 \\
& \frac{(-3a^3B+7a^2Ab-3ab^2B-Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{d(a^2-b^2)(a\sec(c+dx)+b)} - \int \frac{-b(-3Ba^3+7Aba^2-3b^2Ba-Ab^3)\sec^2(c+dx)+4ab(2Aa^2-3bBa+Ab^2)\sec(c+dx)+b(-Ba^3+5Aba^2-5b^2Ba+Ab^3)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))} dx}{2b(a^2-b^2)}
\end{aligned}$$

3.580. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^3\sqrt{\sec(c+dx)}} dx$

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{\int \frac{-b(-3Ba^3 + 7Aba^2 - 3b^2Ba - Ab^3) \csc(c + dx + \frac{\pi}{2})^2 + 4ab(2Aa^2 - 3bBa + Ab^2) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\sec(c + dx + \frac{\pi}{2})} (b + a \csc(c + dx + \frac{\pi}{2}))} dx}{2b(a^2 - b^2)}$$

$$\frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{4a(a^2 - b^2)}{2b(a^2 - b^2)}$$

↓ 4594

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{\int \frac{(-Ba^3 + 5Aba^2 - 5b^2Ba + Ab^3)b^2 + a(Ba^3 + 3Aba^2 - 7b^2Ba + 3Ab^3) \sec(c + dx)b}{\sqrt{\sec(c + dx)}} dx}{b^2} - \frac{(a^5B + 3a^4B)}{2b(a^2 - b^2)}$$

$$\frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{4a(a^2 - b^2)}{2b(a^2 - b^2)}$$

↓ 3042

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{\int \frac{(-Ba^3 + 5Aba^2 - 5b^2Ba + Ab^3)b^2 + a(Ba^3 + 3Aba^2 - 7b^2Ba + 3Ab^3) \csc(c + dx + \frac{\pi}{2})b}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx}{b^2} - \frac{(a^5B + 3a^4B)}{2b(a^2 - b^2)}$$

$$\frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{4a(a^2 - b^2)}{2b(a^2 - b^2)}$$

↓ 4274

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \int \frac{1}{\sqrt{\sec(c + dx)}} dx + ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \int \sqrt{\sec(c + dx)}}{b^2} - \frac{(a^5B + 3a^4B)}{2b(a^2 - b^2)}$$

$$\frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{4a(a^2 - b^2)}{2b(a^2 - b^2)}$$

↓ 3042

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \int \sqrt{\csc(c + dx + \frac{\pi}{2})}}{b^2} - \frac{(a^5B + 3a^4B)}{2b(a^2 - b^2)}$$

$$\frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{4a(a^2 - b^2)}{2b(a^2 - b^2)}$$

↓ 4258

3.580. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx$

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + ab(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{b^2}$$

$4a(a^2 - b^2)$

↓ 3042

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + ab(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{b^2}$$

$4a(a^2 - b^2)$

↓ 3119

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{b^2}}$$

$4a(a^2 - b^2)$

↓ 3120

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticF}(\frac{1}{2}(c + dx), 2) + \frac{2b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{b^2}}$$

$4a(a^2 - b^2)$

↓ 4336

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticF}(\frac{1}{2}(c + dx), 2) + \frac{2b^2(a^3(-B) + 5a^2Ab - 5ab^2B + Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}{b^2}}$$

$4a(a^2 - b^2)$

↓ 3042

3.580. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx$

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{2b^2(a^3(-B))}{b^2}$$

↓ 3284

$$\frac{b(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2ad(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{(-3a^3B + 7a^2Ab - 3ab^2B - Ab^3) \sin(c + dx) \sqrt{\sec(c + dx)}}{d(a^2 - b^2)(a \sec(c + dx) + b)} - \frac{2ab(a^3B + 3a^2Ab - 7ab^2B + 3Ab^3) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d} + \frac{2b^2(a^3(-B))}{b^2}$$

4a

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sqrt[Sec[c + d*x]]),x]`

output `(b*(A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(2*a*(a^2 - b^2)*d*(b + a*Sec[c + d*x])^2) - (-1/2*((2*b^2*(5*a^2*A*b + A*b^3 - a^3*B - 5*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*a*b*(3*a^2*A*b + 3*A*b^3 + a^3*B - 7*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 - (2*(3*a^4*A*b + 10*a^2*A*b^3 - A*b^5 + a^5*B - 10*a^3*b^2*B - 3*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b*(a + b)*d)/(b*(a^2 - b^2)) + ((7*a^2*A*b - A*b^3 - 3*a^3*B - 3*a*b^2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*d*(b + a*Sec[c + d*x]))/(4*a*(a^2 - b^2))`

3.580.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.580. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 4517 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[a*d^2*(
A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n -
2)/(b*f*(m + 1)*(a^2 - b^2))), x] - Simp[d/(b*(m + 1)*(a^2 - b^2)) Int[(
a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 2)*Simp[a*d*(A*b - a*B)*(
n - 2) + b*d*(A*b - a*B)*(m + 1)*Csc[e + f*x] - (a*A*b*d*(m + n) - d*B*(a^2
*(n - 1) + b^2*(m + 1)))*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f
, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && GtQ[
n, 1]
```

```
rule 4588 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a + b*Csc
[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(m + 1)*(a^2 - b^2))), x] + Sim
p[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f
*x])^n*Simp[a*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C)*(m + n +
1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m
+ n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x
] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && ILtQ[n, 0])
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.580.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1849 vs. $2(454) = 908$.

Time = 13.38 (sec) , antiderivative size = 1850, normalized size of antiderivative = 4.60

method	result	size
default	Expression too large to display	1850

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3/sec(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

$$3.580. \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sqrt{\sec(c+dx)}} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-4*B/b/(-2*a*b
+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2
*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2*d*x+1
/2*c),-2*b/(a-b),2^(1/2))+2*(A*b-2*B*a)/b^2*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*
x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2
*d*x+1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*
x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*E
llipticF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)
^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1
/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)
*b/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*si
n(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c
),2^(1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2
*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)
^2)^(1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/
(-2*a*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1
)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(co
s(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a*(A*b-B*a)/b^2*(-1/2/a*b^2/(a^2-b
^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2
)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)^2-3/4*b^2*(3*a^2-b^2)/a^2/(a^2-b^2)^2*...

```

3.580.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `Timed out`

3.580.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.580.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sqrt(sec(d*x + c))), x)`

3.580.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sqrt(sec(d*x + c))), x)`

3.580.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^3),x)`

3.581
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$$

3.581.1 Optimal result 5298
 3.581.2 Mathematica [A] (verified) 5299
 3.581.3 Rubi [A] (verified) 5300
 3.581.4 Maple [B] (verified) 5306
 3.581.5 Fricas [F(-1)] 5307
 3.581.6 Sympy [F(-1)] 5307
 3.581.7 Maxima [F] 5307
 3.581.8 Giac [F] 5308
 3.581.9 Mupad [F(-1)] 5308

3.581.1 Optimal result

Integrand size = 33, antiderivative size = 400

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= - \frac{(a^2 Ab + 5Ab^3 + 3a^3 B - 9ab^2 B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{4b^2 (a^2 - b^2)^2 d}$$

$$+ \frac{(a^3 Ab - 7aAb^3 + 3a^4 B - 5a^2 b^2 B + 8b^4 B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4b^3 (a^2 - b^2)^2 d}$$

$$- \frac{(a^4 Ab - 10a^2 Ab^3 - 3Ab^5 + 3a^5 B - 6a^3 b^2 B + 15ab^4 B) \sqrt{\cos(c + dx)} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4(a - b)^2 b^3 (a + b)^3 d}$$

$$- \frac{(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{2(a^2 - b^2) d (b + a \sec(c + dx))^2}$$

$$+ \frac{(3a^2 Ab + 3Ab^3 + a^3 B - 7ab^2 B) \sqrt{\sec(c + dx)} \sin(c + dx)}{4b(a^2 - b^2)^2 d (b + a \sec(c + dx))}$$

output
$$-1/2*(A*b-B*a)*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/(a^2-b^2)/d/(b+a*\sec(d*x+c))^{2+}$$

$$1/4*(3*A*a^2*b+3*A*b^3+B*a^3-7*B*a*b^2)*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/b/(a^2$$

$$-b^2)^2/d/(b+a*\sec(d*x+c))-1/4*(A*a^2*b+5*A*b^3+3*B*a^3-9*B*a*b^2)*(cos(1/$$

$$2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^{(1$$

$$/2))*cos(d*x+c)^{(1/2)*\sec(d*x+c)^{(1/2)}/b^2/(a^2-b^2)^2/d+1/4*(A*a^3*b-7*A*$$

$$a*b^3+3*B*a^4-5*B*a^2*b^2+8*B*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*$$

$$x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^{(1/2))*cos(d*x+c)^{(1/2)*\sec(d*x+c)$$

$$^{(1/2)}/b^3/(a^2-b^2)^2/d-1/4*(A*a^4*b-10*A*a^2*b^3-3*A*b^5+3*B*a^5-6*B*a^3$$

$$*b^2+15*B*a*b^4)*(cos(1/2*d*x+1/2*c)^2)^{(1/2)}/cos(1/2*d*x+1/2*c)*EllipticP$$

$$i(sin(1/2*d*x+1/2*c),2*b/(a+b),2^{(1/2))*cos(d*x+c)^{(1/2)*\sec(d*x+c)^{(1/2)}/$$

$$(a-b)^2/b^3/(a+b)^3/d$$

3.581.2 Mathematica [A] (verified)

Time = 6.01 (sec) , antiderivative size = 551, normalized size of antiderivative = 1.38

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{4b^2(a(3a^2Ab+3Ab^3+a^3B-7ab^2B)+b(a^2Ab+5Ab^3+3a^3B-9ab^2B)\cos(c+dx))\sin(c+dx)}{(a^2-b^2)^2} + \frac{\cos(c+dx)(a+b\cos(c+dx))\cot(c+dx)(b+a\sec(c+dx))}{(a-b)^2(b^3/(a+b)^3)}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sec[c + d*x]^(3/2)),x]`

output
$$((4*b^2*(a*(3*a^2*A*b + 3*A*b^3 + a^3*B - 7*a*b^2*B) + b*(a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*Cos[c + d*x])*Sin[c + d*x])/((a^2 - b^2)^2 + (Cos[c + d*x]*(a + b*Cos[c + d*x])*Cot[c + d*x]*(b + a*Sec[c + d*x]))*(2*b^2*(5*a^2*A*b + A*b^3 - a^3*B - 5*a*b^2*B)*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] - EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + 16*a*b^2*(-3*a*A*b + a^2*B + 2*b^2*B)*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + (a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*(4*a*b - 4*a*b*Sec[c + d*x]^2 + 4*a*b*EllipticE[ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 2*(2*a - b)*b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] + 4*a^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2] - 2*b^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*Sqrt[Sec[c + d*x]]*Sqrt[-Tan[c + d*x]^2]))/(a*(a - b)^2*(a + b)^2)/(16*b^3*d*(a + b*Cos[c + d*x])^2*Sqrt[Sec[c + d*x]])$$

3.581.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$$

3.581.3 Rubi [A] (verified)

Time = 2.71 (sec) , antiderivative size = 397, normalized size of antiderivative = 0.99, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4515, 27, 3042, 4588, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^3} dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{\sqrt{\sec(c + dx)}(A \sec(c + dx) + B)}{(a \sec(c + dx) + b)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(A \csc\left(c + dx + \frac{\pi}{2}\right) + B\right)}{\left(a \csc\left(c + dx + \frac{\pi}{2}\right) + b\right)^3} dx \\
 & \quad \downarrow \text{4515} \\
 & \frac{\int \frac{-3(Ab - aB) \sec^2(c + dx) + 4(aA - bB) \sec(c + dx) + Ab - aB}{2\sqrt{\sec(c + dx)}(b + a \sec(c + dx))^2} dx}{2(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{-3(Ab - aB) \sec^2(c + dx) + 4(aA - bB) \sec(c + dx) + Ab - aB}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))^2} dx}{4(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{-3(Ab - aB) \csc\left(c + dx + \frac{\pi}{2}\right)^2 + 4(aA - bB) \csc\left(c + dx + \frac{\pi}{2}\right) + Ab - aB}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(b + a \csc\left(c + dx + \frac{\pi}{2}\right))^2} dx}{4(a^2 - b^2)} - \frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2} \\
 & \quad \downarrow \text{4588}
 \end{aligned}$$

3.581. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx$

$$\int -\frac{3Ba^3+Ab^2-9b^2Ba+5Ab^3-(Ba^3+3Aba^2-7b^2Ba+3Ab^3)\sec^2(c+dx)+4b(-Ba^2+3Aba-2b^2B)\sec(c+dx)}{2\sqrt{\sec(c+dx)}(b+a\sec(c+dx))}dx + \frac{(a^3B+3a^2Ab-7ab^2B+3Ab^3)\sin(c+dx)}{bd(a^2-b^2)(a\sec(c+dx)+b)}$$

$$\frac{4(a^2-b^2)}{2d(a^2-b^2)(a\sec(c+dx)+b)^2} \frac{(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2d(a^2-b^2)(a\sec(c+dx)+b)^2}$$

↓ 27

$$\frac{(a^3B+3a^2Ab-7ab^2B+3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{bd(a^2-b^2)(a\sec(c+dx)+b)} - \int \frac{3Ba^3+Ab^2-9b^2Ba+5Ab^3-(Ba^3+3Aba^2-7b^2Ba+3Ab^3)\sec^2(c+dx)+4b(-Ba^2+3Aba-2b^2B)\sec(c+dx)}{\sqrt{\sec(c+dx)}(b+a\sec(c+dx))}dx$$

$$\frac{4(a^2-b^2)}{2d(a^2-b^2)(a\sec(c+dx)+b)^2} \frac{(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2d(a^2-b^2)(a\sec(c+dx)+b)^2}$$

↓ 3042

$$\frac{(a^3B+3a^2Ab-7ab^2B+3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{bd(a^2-b^2)(a\sec(c+dx)+b)} - \int \frac{3Ba^3+Ab^2-9b^2Ba+5Ab^3+(Ba^3-3Aba^2+7b^2Ba-3Ab^3)\csc(c+dx+\frac{\pi}{2})+4b(-Ba^2+3Aba-2b^2B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}(b+a\csc(c+dx+\frac{\pi}{2}))}dx$$

$$\frac{4(a^2-b^2)}{2d(a^2-b^2)(a\sec(c+dx)+b)^2} \frac{(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2d(a^2-b^2)(a\sec(c+dx)+b)^2}$$

↓ 4594

$$\frac{(a^3B+3a^2Ab-7ab^2B+3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{bd(a^2-b^2)(a\sec(c+dx)+b)} - \int \frac{b(3Ba^3+Ab^2-9b^2Ba+5Ab^3)-(3Ba^4+Ab^3-5b^2Ba^2-7Ab^3a+8b^4B)\sec(c+dx)}{\sqrt{\sec(c+dx)}b^2}dx + \frac{(3a^5B-3a^4B+3a^3B-3a^2B+3aB-B)\csc(c+dx+\frac{\pi}{2})}{2b(a^2-b^2)}$$

$$\frac{4(a^2-b^2)}{2d(a^2-b^2)(a\sec(c+dx)+b)^2} \frac{(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2d(a^2-b^2)(a\sec(c+dx)+b)^2}$$

↓ 3042

$$\frac{(a^3B+3a^2Ab-7ab^2B+3Ab^3)\sin(c+dx)\sqrt{\sec(c+dx)}}{bd(a^2-b^2)(a\sec(c+dx)+b)} - \int \frac{b(3Ba^3+Ab^2-9b^2Ba+5Ab^3)+(-3Ba^4-Ab^3+5b^2Ba^2+7Ab^3a-8b^4B)\csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})}b^2}dx + \frac{(3a^5B-3a^4B+3a^3B-3a^2B+3aB-B)\csc(c+dx+\frac{\pi}{2})}{2b(a^2-b^2)}$$

$$\frac{4(a^2-b^2)}{2d(a^2-b^2)(a\sec(c+dx)+b)^2} \frac{(Ab-aB)\sin(c+dx)\sqrt{\sec(c+dx)}}{2d(a^2-b^2)(a\sec(c+dx)+b)^2}$$

↓ 4274

3.581. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^3\sec^{\frac{3}{2}}(c+dx)}dx$

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{b(3a^3 B + a^2 Ab - 9ab^2 B + 5Ab^3) \int \frac{1}{\sqrt{\sec(c+dx)}} dx - (3a^4 B + a^3 Ab - 5a^2 b^2 B - 7aAb^3 + 8b^4 B) \int \sqrt{\sec(c+dx)}}{b^2} - \frac{2b(a^2 - b^2)}{4(a^2 - b^2)}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{b(3a^3 B + a^2 Ab - 9ab^2 B + 5Ab^3) \int \frac{1}{\sqrt{\csc(c+dx + \frac{\pi}{2})}} dx - (3a^4 B + a^3 Ab - 5a^2 b^2 B - 7aAb^3 + 8b^4 B) \int \sqrt{\csc(c+dx + \frac{\pi}{2})}}{b^2} - \frac{2b(a^2 - b^2)}{4(a^2 - b^2)}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4258

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{b(3a^3 B + a^2 Ab - 9ab^2 B + 5Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\cos(c+dx)} dx - (3a^4 B + a^3 Ab - 5a^2 b^2 B - 7aAb^3 + 8b^4 B) \int \sqrt{\sec(c+dx)}}{b^2} - \frac{2b(a^2 - b^2)}{4(a^2 - b^2)}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{b(3a^3 B + a^2 Ab - 9ab^2 B + 5Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \sqrt{\sin(c+dx + \frac{\pi}{2})} dx - (3a^4 B + a^3 Ab - 5a^2 b^2 B - 7aAb^3 + 8b^4 B) \int \sqrt{\sec(c+dx)}}{b^2} - \frac{2b(a^2 - b^2)}{4(a^2 - b^2)}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3119

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{2b(3a^3 B + a^2 Ab - 9ab^2 B + 5Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) - (3a^4 B + a^3 Ab - 5a^2 b^2 B - 7aAb^3 + 8b^4 B) \int \sqrt{\sec(c+dx)}}{d} - \frac{2b(a^2 - b^2)}{4(a^2 - b^2)}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3120

3.581. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{(3a^5 B + a^4 Ab - 6a^3 b^2 B - 10a^2 Ab^3 + 15ab^4 B - 3Ab^5) \int \frac{\csc(c+dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c+dx + \frac{\pi}{2})} dx}{b^2} + \frac{2b(3a^3 B + a^2 Ab - 9a^2 b^2 B + 3ab^3)}{4(a^2 - b^2)}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 4336

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{(3a^5 B + a^4 Ab - 6a^3 b^2 B - 10a^2 Ab^3 + 15ab^4 B - 3Ab^5) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))} dx}{b^2}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3042

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{(3a^5 B + a^4 Ab - 6a^3 b^2 B - 10a^2 Ab^3 + 15ab^4 B - 3Ab^5) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})}(a+b \cos(c+dx))} dx}{b^2}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

↓ 3284

$$\frac{(a^3 B + 3a^2 Ab - 7ab^2 B + 3Ab^3) \sin(c+dx) \sqrt{\sec(c+dx)}}{bd(a^2 - b^2)(a \sec(c+dx) + b)} - \frac{2b(3a^3 B + a^2 Ab - 9ab^2 B + 5Ab^3) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d} - \frac{2(3a^4 B + a^3 Ab - 5a^2 b^2 B + 3ab^3)}{b^2}$$

$$\frac{(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2d(a^2 - b^2)(a \sec(c + dx) + b)^2}$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sec[c + d*x]^(3/2)),x]`

```
output -1/2*((A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/((a^2 - b^2)*d*(b + a*Sec[c + d*x])^2) + (-1/2*(((2*b*(a^2*A*b + 5*A*b^3 + 3*a^3*B - 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*(a^3*A*b - 7*a*A*b^3 + 3*a^4*B - 5*a^2*b^2*B + 8*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (2*(a^4*A*b - 10*a^2*A*b^3 - 3*A*b^5 + 3*a^5*B - 6*a^3*b^2*B + 15*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d))/(b*(a^2 - b^2)) + ((3*a^2*A*b + 3*A*b^3 + a^3*B - 7*a*b^2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b*(a^2 - b^2)*d*(b + a*Sec[c + d*x]))/(4*(a^2 - b^2))
```

3.581.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3284 Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]
```

```
rule 3439 Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

3.581.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^2(c+dx)} dx$$

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 4515 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] := Simp[(-d)*(A*b - a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^(n - 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^(n - 1)*Simp[d*(n - 1)*(A*b - a*B) + d*(a*A - b*B)*(m + 1)*Csc[e + f*x] - d*(A*b - a*B)*(m + n + 1)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && LtQ[0, n, 1]`

rule 4588 `Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(m_.), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(m + 1)*(a^2 - b^2))), x] + Simp[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[a*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C)*(m + n + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m + n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && ILtQ[n, 0])`

3.581.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{3}{2}}(c+dx)} dx$$


```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))/ (Sqrt[csc[(e_.) + (f_.)*(x_.)]*(d_.)]*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.581.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1936 vs. $2(452) = 904$.

Time = 13.43 (sec) , antiderivative size = 1937, normalized size of antiderivative = 4.84

method	result	size
default	Expression too large to display	1937

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3/sec(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output -((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*B/b^3*(sin(1
/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1
/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-
4/b^2*(A*b-3*B*a)/(-2*a*b+2*b^2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*
d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)
*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+2*a^2*(A*b-B*a)/b^3*(-1
/2/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x
+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+1/2*c)^2+a-b)-7/8*(a+b)/(a^2-b^2)*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4
+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/4/(a+
b)/(a^2-b^2)/a*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1
/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2
*d*x+1/2*c),2^(1/2))*b+3/8/(a+b)/(a^2-b^2)/a^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)
)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1
/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*b^2-9/8*b/(a^2-b^2)^2
*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/
2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^
(1/2))+3/8*b^3/a^2/(a^2-b^2)^2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2...
```

3.581.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(3/2),x, algorithm m="fricas")`

output `Timed out`

3.581.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.581.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2)), x)`

3.581.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sec(d*x + c)^(3/2)), x)`

3.581.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^3), x)`

$$3.582 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$$

3.582.1 Optimal result	5309
3.582.2 Mathematica [A] (warning: unable to verify)	5310
3.582.3 Rubi [A] (verified)	5311
3.582.4 Maple [B] (verified)	5317
3.582.5 Fracas [F(-1)]	5318
3.582.6 Sympy [F(-1)]	5319
3.582.7 Maxima [F]	5319
3.582.8 Giac [F]	5319
3.582.9 Mupad [F(-1)]	5320

3.582.1 Optimal result

Integrand size = 33, antiderivative size = 427

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{(3a^3Ab - 9aAb^3 - 15a^4B + 29a^2b^2B - 8b^4B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{4b^3(a^2 - b^2)^2 d}$$

$$+ \frac{(3a^4Ab - 5a^2Ab^3 + 8Ab^5 - 15a^5B + 33a^3b^2B - 24ab^4B) \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{4b^4(a^2 - b^2)^2 d}$$

$$- \frac{a(3a^4Ab - 6a^2Ab^3 + 15Ab^5 - 15a^5B + 38a^3b^2B - 35ab^4B) \sqrt{\cos(c + dx)} \text{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx), 2\right)}{4(a - b)^2 b^4 (a + b)^3 d}$$

$$+ \frac{a(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{2b(a^2 - b^2) d(b + a \sec(c + dx))^2}$$

$$+ \frac{a(a^2Ab - 7Ab^3 - 5a^3B + 11ab^2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{4b^2(a^2 - b^2)^2 d(b + a \sec(c + dx))}$$

output $\frac{1}{2}a(Ab - B^2a) \sin(dx+c) \sec(dx+c)^{1/2} / b(a^2 - b^2) / d(b + a \sec(dx+c))^{2+1/4} a(A^2b - 7A^2b^3 - 5B^2a^3 + 11B^2a^2b) \sin(dx+c) \sec(dx+c)^{1/2} / b^2(a^2 - b^2)^2 / d(b + a \sec(dx+c)) - 1/4(3A^2a^3b - 9A^2a^2b^3 - 15B^2a^4 + 29B^2a^2b^2 - 8B^2b^4) (\cos(1/2dx + 1/2c))^2)^{1/2} / \cos(1/2dx + 1/2c) \text{EllipticE}(\sin(1/2dx + 1/2c), 2^{1/2}) \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / b^3(a^2 - b^2)^2 / d + 1/4(3A^2a^4b - 5A^2a^2b^3 + 8A^2b^5 - 15B^2a^5 + 33B^2a^3b^2 - 24B^2a^2b^4) (\cos(1/2dx + 1/2c))^2)^{1/2} / \cos(1/2dx + 1/2c) \text{EllipticF}(\sin(1/2dx + 1/2c), 2^{1/2}) \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / b^4(a^2 - b^2)^2 / d - 1/4a(3A^2a^4b - 6A^2a^2b^3 + 15A^2b^5 - 15B^2a^5 + 38B^2a^3b^2 - 35B^2a^2b^4) (\cos(1/2dx + 1/2c))^2)^{1/2} / \cos(1/2dx + 1/2c) \text{EllipticPi}(\sin(1/2dx + 1/2c), 2b/(a+b), 2^{1/2}) \cos(dx+c)^{1/2} \sec(dx+c)^{1/2} / (a-b)^2 / b^4 / (a+b)^3 / d$

3.582.2 Mathematica [A] (warning: unable to verify)

Time = 7.14 (sec) , antiderivative size = 820, normalized size of antiderivative = 1.92

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2(-a^3Ab - 5aAb^3 + 5a^4B - 7a^2b^2B + 8b^4B) \cos^2(c+dx) \left(\text{EllipticF}\left(\arcsin\left(\sqrt{\sec(c+dx)}\right), -1\right) - \text{EllipticPi}\left(-\frac{a}{b}, \arcsin\left(\sqrt{\sec(c+dx)}\right), -1\right) \right) (b+a \sec(c+dx))}{a(a+b \cos(c+dx))(1-\cos^2(c+dx))} + \frac{\sqrt{\sec(c+dx)} \left(-\frac{a(-3a^2Ab + 9Ab^3 + 7a^3B - 13ab^2B) \sin(c+dx)}{4b^3(a^2 - b^2)^2} - \frac{a^3Ab \sin(c+dx) - a^4B \sin(c+dx)}{2b^3(-a^2 + b^2)(a+b \cos(c+dx))^2} + \frac{-5a^4Ab \sin(c+dx) + 11a^2Ab^3 \sin(c+dx)}{4b^3(-a^2 + b^2)} \right)}{d}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sec[c + d*x]^(5/2)),x]`

output $((2*(-a^3A*b) - 5*a*A*b^3 + 5*a^4*B - 7*a^2*b^2*B + 8*b^4*B)*\text{Cos}[c + d*x]^{2*(\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1] - \text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1])*(b + a*\text{Sec}[c + d*x])* \text{Sqrt}[1 - \text{Sec}[c + d*x]^2]*\text{Sin}[c + d*x]/(a*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)) + (2*(8*a^2*A*b^2 + 16*A*b^4 + 8*a^3*b*B - 32*a*b^3*B)*\text{Cos}[c + d*x]^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*(b + a*\text{Sec}[c + d*x])* \text{Sqrt}[1 - \text{Sec}[c + d*x]^2]*\text{Sin}[c + d*x]/(b*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)) + ((-3*a^3*A*b + 9*a*A*b^3 + 15*a^4*B - 29*a^2*b^2*B + 8*b^4*B)*\text{Cos}[2*(c + d*x)]*(b + a*\text{Sec}[c + d*x])*(-4*a*b + 4*a*b*\text{Sec}[c + d*x]^2 - 4*a*b*\text{EllipticE}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] + 2*(2*a - b)*b*\text{EllipticF}[\text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] - 4*a^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2] + 2*b^2*\text{EllipticPi}[-(a/b), \text{ArcSin}[\text{Sqrt}[\text{Sec}[c + d*x]]], -1]*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[1 - \text{Sec}[c + d*x]^2])* \text{Sin}[c + d*x]/(a*b^2*(a + b*\text{Cos}[c + d*x])*(1 - \text{Cos}[c + d*x]^2)*\text{Sqrt}[\text{Sec}[c + d*x]]*(2 - \text{Sec}[c + d*x]^2)))/(16*(a - b)^2*b^2*(a + b)^2*d) + (\text{Sqrt}[\text{Sec}[c + d*x]]*(-1/4*(a*(-3*a^2*A*b + 9*A*b^3 + 7*a^3*B - 13*a*b^2*B)*\text{Sin}[c + d*x])/(b^3*(a^2 - b^2)^2) - (a^3*A*b*\text{Sin}[c + d*x] - a^4*B*\text{Sin}[c + d*x])/(2*b^3*(-a^2 + b^2)*(a + b*\text{Cos}[c + d*x])^2) + (-5*a^4*A*b*\text{Sin}[c + d*x] + 11*a^2*A*b^3*\text{Sin}[c + d*x] + 9*a^5*B*\text{Sin}[c + d*x] - 15*a^3*b^2*B*\text{Sin}...$

3.582.3 Rubi [A] (verified)

Time = 3.03 (sec) , antiderivative size = 428, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 3439, 3042, 4518, 27, 3042, 4588, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{5/2} (a + b \sin(c + dx + \frac{\pi}{2}))^3} dx$$

↓ 3439

$$\int \frac{A \sec(c + dx) + B}{\sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^3} dx$$

3.582. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{A \csc(c+dx+\frac{\pi}{2}) + B}{\sqrt{\csc(c+dx+\frac{\pi}{2})} (a \csc(c+dx+\frac{\pi}{2}) + b)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{-5Ba^2 - 3(Ab-aB) \sec^2(c+dx)a + Aba + 4b^2B + 4b(Ab-aB) \sec(c+dx)}{2\sqrt{\sec(c+dx)}(b+a \sec(c+dx))^2} dx \\
& \quad \downarrow \text{4518} \\
& \frac{2b(a^2 - b^2)}{a(Ab - aB) \sin(c+dx) \sqrt{\sec(c+dx)}} + \frac{a(Ab - aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{2bd(a^2 - b^2)(a \sec(c+dx) + b)^2} \\
& \quad \downarrow \text{27} \\
& \frac{a(Ab - aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{2bd(a^2 - b^2)(a \sec(c+dx) + b)^2} - \int \frac{-5Ba^2 - 3(Ab-aB) \sec^2(c+dx)a + Aba + 4b^2B + 4b(Ab-aB) \sec(c+dx)}{\sqrt{\sec(c+dx)}(b+a \sec(c+dx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \frac{a(Ab - aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{2bd(a^2 - b^2)(a \sec(c+dx) + b)^2} - \int \frac{-5Ba^2 - 3(Ab-aB) \csc(c+dx+\frac{\pi}{2})^2 a + Aba + 4b^2B + 4b(Ab-aB) \csc(c+dx+\frac{\pi}{2})}{\sqrt{\csc(c+dx+\frac{\pi}{2})} (b+a \csc(c+dx+\frac{\pi}{2}))^2} dx \\
& \quad \downarrow \text{4588} \\
& \frac{a(Ab - aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{2bd(a^2 - b^2)(a \sec(c+dx) + b)^2} - \int \frac{-15Ba^4 + 3Aba^3 + 29b^2Ba^2 - 9Ab^3a - (-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) \sec^2(c+dx)a - 8b^4B - 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \sec(c+dx)}{2\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx - \frac{a(-5a^3B + a^2Ab)}{b} \\
& \quad \downarrow \text{27} \\
& \frac{a(Ab - aB) \sin(c+dx) \sqrt{\sec(c+dx)}}{2bd(a^2 - b^2)(a \sec(c+dx) + b)^2} - \int \frac{-15Ba^4 + 3Aba^3 + 29b^2Ba^2 - 9Ab^3a - (-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) \sec^2(c+dx)a - 8b^4B - 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \sec(c+dx)}{\sqrt{\sec(c+dx)}(b+a \sec(c+dx))} dx - \frac{a(-5a^3B + a^2Ab)}{b} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.582. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{\int \frac{-15Ba^4 + 3Aba^3 + 29b^2Ba^2 - 9Ab^3a - (-5Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) \csc(c + dx + \frac{\pi}{2})^2 a - 8b^4B - 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\csc(c + dx + \frac{\pi}{2})(b + a \csc(c + dx + \frac{\pi}{2}))}} dx}{2b(a^2 - b^2)} - \frac{a(-5a^3)}{4b(a^2 - b^2)}$$

↓ 4594

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\sec^3(c + dx)}{b + a \sec(c + dx)} dx + \int \frac{b(-15Ba^4 + 3Aba^3 + 29b^2Ba^2 - 9Ab^3a - 8b^4B) - (-15Ba^5 + 3Aba^4 + 33b^2Ba^3 - 5a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \sqrt{\sec(c + dx)}}{b^2}}{2b(a^2 - b^2)} - \frac{4b(a^2 - b^2)}{4b(a^2 - b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + \int \frac{b(-15Ba^4 + 3Aba^3 + 29b^2Ba^2 - 9Ab^3a - 8b^4B) + (15Ba^5 - 3Aba^4 - 33b^2Ba^3 - 5a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \sqrt{\csc(c + dx + \frac{\pi}{2})}}{b^2}}{2b(a^2 - b^2)} - \frac{4b(a^2 - b^2)}{4b(a^2 - b^2)}$$

↓ 4274

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \int \frac{1}{\sqrt{\sec(c + dx)}} dx - (-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5)}{2b(a^2 - b^2)} - \frac{4b(a^2 - b^2)}{4b(a^2 - b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx - (-15a^5B + 3a^4Ab + 38a^3b^2B - 5a^2Ab^3 - 24ab^4B + 8Ab^5) \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5)}{2b(a^2 - b^2)} - \frac{4b(a^2 - b^2)}{4b(a^2 - b^2)}$$

↓ 4258

3.582. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)}}{2b(a^2 - b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx)}}{2b(a^2 - b^2)}$$

↓ 3119

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{2b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2})}{d \cdot 2b(a^2 - b^2)}$$

↓ 3120

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{2b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2})}{d \cdot 2b(a^2 - b^2)}$$

↓ 4336

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))} dx}{b^2} + \frac{2b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{2b(a^2 - b^2)}$$

↓ 3042

3.582. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})(a+b \sin(c+dx + \frac{\pi}{2}))}} dx}{b^2} + \frac{2b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{2b(a^2 - b^2)}$$

↓ 3284

$$\frac{a(Ab - aB) \sin(c + dx) \sqrt{\sec(c + dx)}}{2bd(a^2 - b^2)(a \sec(c + dx) + b)^2} - \frac{2a(-15a^5B + 3a^4Ab + 38a^3b^2B - 6a^2Ab^3 - 35ab^4B + 15Ab^5) \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \text{EllipticPi}(\frac{2b}{a+b}, \frac{1}{2}(c+dx), 2)}{b^2d(a+b)} + \frac{2b(-15a^4B + 3a^3Ab + 29a^2b^2B - 9aAb^3 - 8b^4B)}{2b(a^2 - b^2)}$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sec[c + d*x]^(5/2)),x]`

output `(a*(A*b - a*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/((2*b*(a^2 - b^2)*d*(b + a*Sec[c + d*x])^2) - (((2*b*(3*a^3*A*b - 9*a*A*b^3 - 15*a^4*B + 29*a^2*b^2*B - 8*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d - (2*(3*a^4*A*b - 5*a^2*A*b^3 + 8*A*b^5 - 15*a^5*B + 33*a^3*b^2*B - 24*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (2*a*(3*a^4*A*b - 6*a^2*A*b^3 + 15*A*b^5 - 15*a^5*B + 38*a^3*b^2*B - 35*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d))/(2*b*(a^2 - b^2)) - (a*(a^2*A*b - 7*A*b^3 - 5*a^3*B + 11*a*b^2*B)*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b*(a^2 - b^2)*d*(b + a*Sec[c + d*x]))/(4*b*(a^2 - b^2))`

3.582.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

3.582. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3284 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[c + d, 0]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4336 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(3/2)/((csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[d*Sqrt[d*Sin[e + f*x]]*Sqrt[d*Csc[e + f*x]] Int[1/(Sqrt[d*Sin[e + f*x]]*(b + a*Sin[e + f*x])), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

```
rule 4518 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[b*(A*b
- a*B)*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(
m + 1)*(a^2 - b^2))), x] + Simp[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[
e + f*x])^(m + 1)*(d*Csc[e + f*x])^n*Simp[A*(a^2*(m + 1) - b^2*(m + n + 1))
+ a*b*B*n - a*(A*b - a*B)*(m + 1)*Csc[e + f*x] + b*(A*b - a*B)*(m + n + 2)
*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*
b - a*B, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && IL
tQ[n, 0])
```

```
rule 4588 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a + b*Csc
[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(m + 1)*(a^2 - b^2))), x] + Sim
p[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f
*x])^n*Simp[a*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C)*(m + n +
1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m
+ n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x
] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && ILtQ[n, 0])
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.582.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1976 vs. 2(479) = 958.

Time = 15.44 (sec) , antiderivative size = 1977, normalized size of antiderivative = 4.63

method	result	size
default	Expression too large to display	1977

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3/sec(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

$$3.582. \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{5}{2}}(c+dx)} dx$$

output

```

-((-2*cos(1/2*d*x+1/2*c)^2+1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*(2/b^4/(-2*sin(
1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1
/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(A*b*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)
)-3*B*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*a-B*b*EllipticE(cos(1/2*d*x+1/
2*c),2^(1/2)))+2*a^2/b^4*(3*A*b-4*B*a)*(-1/a*b^2/(a^2-b^2)*cos(1/2*d*x+1/2
*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*b*cos(1/2*d*x+
1/2*c)^2+a-b)-1/2/a/(a+b)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2
*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*Ellipt
icF(cos(1/2*d*x+1/2*c),2^(1/2))-1/2/(a^2-b^2)*b/a*(sin(1/2*d*x+1/2*c)^2)^(
1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*
x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+1/2/(a^2-b^2)*b/a*
(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2
*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(
1/2))-3*a/(a^2-b^2)/(-2*a*b+2*b^2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(
1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(
1/2)*EllipticPi(cos(1/2*d*x+1/2*c),-2*b/(a-b),2^(1/2))+1/a/(a^2-b^2)/(-2*a
*b+2*b^2)*b^3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/
2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticPi(cos(1/2
*d*x+1/2*c),-2*b/(a-b),2^(1/2))-2*a^3*(A*b-B*a)/b^4*(-1/2/a*b^2/(a^2-b^2)
*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)...

```

3.582.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(5/2),x, algorithm="fracas")`

output `Timed out`

3.582.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3/sec(d*x+c)**(5/2),x)`output `Timed out`**3.582.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(5/2),x, algorithm="maxima")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2)), x)`**3.582.8 Giac [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(5/2),x, algorithm="giac")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sec(d*x + c)^(5/2)), x)`

3.582.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{5/2} (a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^3),x)`

$$3.583 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$$

3.583.1 Optimal result	5321
3.583.2 Mathematica [A] (warning: unable to verify)	5322
3.583.3 Rubi [A] (verified)	5323
3.583.4 Maple [B] (warning: unable to verify)	5330
3.583.5 Fricas [F(-1)]	5331
3.583.6 Sympy [F(-1)]	5332
3.583.7 Maxima [F]	5332
3.583.8 Giac [F]	5332
3.583.9 Mupad [F(-1)]	5333

3.583.1 Optimal result

Integrand size = 33, antiderivative size = 521

$$\begin{aligned} & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx \\ &= \frac{(15a^4Ab - 29a^2Ab^3 + 8Ab^5 - 35a^5B + 65a^3b^2B - 24ab^4B) \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{4b^4 (a^2 - b^2)^2 d} \\ & \quad - \frac{(45a^5Ab - 99a^3Ab^3 + 72aAb^5 - 105a^6B + 223a^4b^2B - 128a^2b^4B - 8b^6B) \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx) \mid 2\right)}{12b^5 (a^2 - b^2)^2 d} \\ & \quad + \frac{a^2(15a^4Ab - 38a^2Ab^3 + 35Ab^5 - 35a^5B + 86a^3b^2B - 63ab^4B) \sqrt{\cos(c + dx)} \operatorname{EllipticPi}\left(\frac{2b}{a+b}, \frac{1}{2}(c + dx)\right)}{4(a - b)^2 b^5 (a + b)^3 d} \\ & \quad - \frac{(15a^3Ab - 33aAb^3 - 35a^4B + 61a^2b^2B - 8b^4B) \sin(c + dx)}{12b^3 (a^2 - b^2)^2 d \sqrt{\sec(c + dx)}} \\ & \quad + \frac{a(Ab - aB) \sin(c + dx)}{2b (a^2 - b^2) d \sqrt{\sec(c + dx)} (b + a \sec(c + dx))^2} \\ & \quad + \frac{a(3a^2Ab - 9Ab^3 - 7a^3B + 13ab^2B) \sin(c + dx)}{4b^2 (a^2 - b^2)^2 d \sqrt{\sec(c + dx)} (b + a \sec(c + dx))} \end{aligned}$$

3.583. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$

output

$$\begin{aligned}
& -1/12*(15*A*a^3*b-33*A*a*b^3-35*B*a^4+61*B*a^2*b^2-8*B*b^4)*\sin(d*x+c)/b^3 \\
& / (a^2-b^2)^2/d/\sec(d*x+c)^{(1/2)}+1/2*a*(A*b-B*a)*\sin(d*x+c)/b/(a^2-b^2)/d/ \\
& (b+a*\sec(d*x+c))^2/\sec(d*x+c)^{(1/2)}+1/4*a*(3*A*a^2*b-9*A*b^3-7*B*a^3+13*B*a \\
& *b^2)*\sin(d*x+c)/b^2/(a^2-b^2)^2/d/(b+a*\sec(d*x+c))/\sec(d*x+c)^{(1/2)}+1/4*(\\
& 15*A*a^4*b-29*A*a^2*b^3+8*A*b^5-35*B*a^5+65*B*a^3*b^2-24*B*a*b^4)*(\cos(1/2 \\
& *d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/ \\
& 2)})*\cos(d*x+c)^{(1/2)}*\sec(d*x+c)^{(1/2)}/b^4/(a^2-b^2)^2/d-1/12*(45*A*a^5*b-9 \\
& 9*A*a^3*b^3+72*A*a*b^5-105*B*a^6+223*B*a^4*b^2-128*B*a^2*b^4-8*B*b^6)*(\cos \\
& (1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2 \\
& ^{(1/2)})*\cos(d*x+c)^{(1/2)}*\sec(d*x+c)^{(1/2)}/b^5/(a^2-b^2)^2/d+1/4*a^2*(15*A \\
& a^4*b-38*A*a^2*b^3+35*A*b^5-35*B*a^5+86*B*a^3*b^2-63*B*a*b^4)*(\cos(1/2*d*x \\
& +1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticPi}(\sin(1/2*d*x+1/2*c), 2*b/(a+b \\
&), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*\sec(d*x+c)^{(1/2)}/(a-b)^2/b^5/(a+b)^3/d
\end{aligned}$$

3.583.2 Mathematica [A] (warning: unable to verify)

Time = 7.35 (sec) , antiderivative size = 865, normalized size of antiderivative = 1.66

$$\begin{aligned}
& \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \\
& \frac{2(-15a^4Ab+21a^2Ab^3-24Ab^5+35a^5B-73a^3b^2B+56ab^4B) \cos^2(c+dx) \left(\text{EllipticF} \left(\arcsin \left(\sqrt{\sec(c+dx)} \right), -1 \right) - \text{EllipticPi} \left(-\frac{a}{b}, \arcsin \left(\sqrt{\sec(c+dx)} \right) \right) \right)}{a(a+b \cos(c+dx))(1-\cos^2(c+dx))} \\
& + \frac{\sqrt{\sec(c+dx)} \left(\frac{a^2(-7a^2Ab+13Ab^3+11a^3B-17ab^2B) \sin(c+dx)}{4b^4(a^2-b^2)^2} - \frac{-a^4Ab \sin(c+dx)+a^5B \sin(c+dx)}{2b^4(-a^2+b^2)(a+b \cos(c+dx))^2} + \frac{9a^5Ab \sin(c+dx)-15a^3Ab^3}{4b^4} \right)}{d}
\end{aligned}$$

input

```
Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sec[c + d*x]^(7/2)),x]
```

3.583. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$

output

```

-1/48*((2*(-15*a^4*A*b + 21*a^2*A*b^3 - 24*A*b^5 + 35*a^5*B - 73*a^3*b^2*B
+ 56*a*b^4*B)*Cos[c + d*x]^2*(EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1] -
EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1])*(b + a*Sec[c + d*x])*
Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(a*(a + b*Cos[c + d*x])*(1 - Cos[c
+ d*x]^2)) + (2*(-24*a^3*A*b^2 + 96*a*A*b^4 + 56*a^4*b*B - 112*a^2*b^3*B -
16*b^5*B)*Cos[c + d*x]^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -
1])*(b + a*Sec[c + d*x])*Sqrt[1 - Sec[c + d*x]^2]*Sin[c + d*x])/(b*(a + b*C
os[c + d*x])*(1 - Cos[c + d*x]^2)) + ((-45*a^4*A*b + 87*a^2*A*b^3 - 24*A*b
^5 + 105*a^5*B - 195*a^3*b^2*B + 72*a*b^4*B)*Cos[2*(c + d*x)]*(b + a*Sec[c
+ d*x])*(-4*a*b + 4*a*b*Sec[c + d*x]^2 - 4*a*b*EllipticE[ArcSin[Sqrt[Sec[
c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*(2*a - b)*
b*EllipticF[ArcSin[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Se
c[c + d*x]^2] - 4*a^2*EllipticPi[-(a/b), ArcSin[Sqrt[Sec[c + d*x]]], -1]*S
qrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2] + 2*b^2*EllipticPi[-(a/b), ArcS
in[Sqrt[Sec[c + d*x]]], -1]*Sqrt[Sec[c + d*x]]*Sqrt[1 - Sec[c + d*x]^2])*S
in[c + d*x])/(a*b^2*(a + b*Cos[c + d*x])*(1 - Cos[c + d*x]^2)*Sqrt[Sec[c +
d*x])*(2 - Sec[c + d*x]^2))/((a - b)^2*b^3*(a + b)^2*d) + (Sqrt[Sec[c +
d*x]]*((a^2*(-7*a^2*A*b + 13*A*b^3 + 11*a^3*B - 17*a*b^2*B)*Sin[c + d*x])/
(4*b^4*(a^2 - b^2)^2) - (-a^4*A*b*Sin[c + d*x]) + a^5*B*Sin[c + d*x])/(2*
b^4*(-a^2 + b^2)*(a + b*Cos[c + d*x])^2) + (9*a^5*A*b*Sin[c + d*x] - 15...

```

3.583.3 Rubi [A] (verified)

Time = 3.86 (sec) , antiderivative size = 519, normalized size of antiderivative = 1.00, number of steps used = 23, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 3439, 3042, 4518, 27, 3042, 4588, 27, 3042, 4592, 27, 3042, 4594, 3042, 4274, 3042, 4258, 3042, 3119, 3120, 4336, 3042, 3284}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sec^{\frac{7}{2}}(c + dx)(a + b \cos(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^3} dx \\
 & \quad \downarrow \text{3439} \\
 & \int \frac{A \sec(c + dx) + B}{\sec^{\frac{3}{2}}(c + dx)(a \sec(c + dx) + b)^3} dx
 \end{aligned}$$

3.583. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \int \frac{A \csc(c + dx + \frac{\pi}{2}) + B}{\csc(c + dx + \frac{\pi}{2})^{3/2} (a \csc(c + dx + \frac{\pi}{2}) + b)^3} dx \\
 & \int \frac{-7Ba^2 - 5(Ab - aB) \sec^2(c + dx)a + 3Aba + 4b^2B + 4b(Ab - aB) \sec(c + dx)}{2 \sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))^2} dx \\
 & \frac{2b(a^2 - b^2)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} + \frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} \\
 & \int \frac{-7Ba^2 - 5(Ab - aB) \sec^2(c + dx)a + 3Aba + 4b^2B + 4b(Ab - aB) \sec(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))^2} dx \\
 & \frac{a(Ab - aB) \sin(c + dx)}{4b(a^2 - b^2)} \\
 & \int \frac{-7Ba^2 - 5(Ab - aB) \csc(c + dx + \frac{\pi}{2})^2 a + 3Aba + 4b^2B + 4b(Ab - aB) \csc(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2} (b + a \csc(c + dx + \frac{\pi}{2}))^2} dx \\
 & \frac{a(Ab - aB) \sin(c + dx)}{4b(a^2 - b^2)} \\
 & \int \frac{-35Ba^4 + 15Aba^3 + 61b^2Ba^2 - 33Ab^3a - 3(-7Ba^3 + 3Aba^2 + 13b^2Ba - 9Ab^3) \sec^2(c + dx)a - 8b^4B - 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \sec(c + dx)}{2 \sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))} dx \\
 & \frac{a(Ab - aB) \sin(c + dx)}{4b(a^2 - b^2)} - \frac{a(-7a^3B + 3b^4B)}{bd(a^2 - b^2)} \\
 & \int \frac{-35Ba^4 + 15Aba^3 + 61b^2Ba^2 - 33Ab^3a - 3(-7Ba^3 + 3Aba^2 + 13b^2Ba - 9Ab^3) \sec^2(c + dx)a - 8b^4B - 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \sec(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(b + a \sec(c + dx))} dx \\
 & \frac{a(Ab - aB) \sin(c + dx)}{2b(a^2 - b^2)} - \frac{a(-7a^3B + 3b^4B)}{bd(a^2 - b^2)} \\
 & \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx
 \end{aligned}$$

3.583. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \int \frac{-35Ba^4 + 15Aba^3 + 61b^2Ba^2 - 33Ab^3a - 3(-7Ba^3 + 3Aba^2 + 13b^2Ba - 9Ab^3) \csc(c + dx + \frac{\pi}{2})^2 a - 8b^4B - 4b(Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \csc(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2} (b + a \csc(c + dx + \frac{\pi}{2}))} dx$$

$$4b(a^2 - b^2)$$

↓ 4592

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{2 \int \frac{-a(-35Ba^4 + 15Aba^3 + 61b^2Ba^2 - 33Ab^3a - 8b^4B) \sec^2(c + dx) + 4b(-7Ba^4 + 3Aba^3 + 14b^2Ba^2 - 12Ab^3a - 8b^4B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{2b(a^2 - b^2)}$$

$$4b(a^2 - b^2)$$

↓ 27

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{\int \frac{-a(-35Ba^4 + 15Aba^3 + 61b^2Ba^2 - 33Ab^3a - 8b^4B) \sec^2(c + dx) + 4b(-7Ba^4 + 3Aba^3 + 14b^2Ba^2 - 12Ab^3a - 8b^4B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{2b(a^2 - b^2)}$$

$$4b(a^2 - b^2)$$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{\int \frac{-a(-35Ba^4 + 15Aba^3 + 61b^2Ba^2 - 33Ab^3a - 8b^4B) \csc(c + dx + \frac{\pi}{2})^2 + 4b(-7Ba^4 + 3Aba^3 + 14b^2Ba^2 - 12Ab^3a - 8b^4B) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{2b(a^2 - b^2)}$$

$$4b(a^2 - b^2)$$

↓ 4594

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\sec^{\frac{3}{2}}(c + dx)}{b + a \sec(c + dx)} dx}{b^2} + \frac{\int \frac{3b(-35Ba^5 + 15Aba^4 + 86b^2Ba^3 - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \csc(c + dx + \frac{\pi}{2})}{\sqrt{\sec(c + dx)}(b + a \sec(c + dx))} dx}{2b(a^2 - b^2)}$$

$$2b(a^2 - b^2)$$

↓ 3042

3.583. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{2b(a^2 - b^2)}$$

↓ 4274

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{2b(a^2 - b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{2b(a^2 - b^2)}$$

↓ 4258

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{2b(a^2 - b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{b^2} + \frac{3b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx}{2b(a^2 - b^2)}$$

↓ 3119

3.583. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^2(c + dx)} dx$

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}}{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx) - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}} b^2}$$

3120

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}}{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx) - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{b + a \csc(c + dx + \frac{\pi}{2})} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}} b^2}$$

4336

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}}{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx) - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}} b^2}$$

3042

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}}{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx) - \frac{3a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}} b^2}$$

3284

$$\frac{a(Ab - aB) \sin(c + dx)}{2bd(a^2 - b^2) \sqrt{\sec(c + dx)}(a \sec(c + dx) + b)^2} - \frac{6a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticPi}(\frac{2}{a}, \frac{c + dx}{2}) + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}}{2(-35a^4B + 15a^3Ab + 61a^2b^2B - 33aAb^3 - 8b^4B) \sin(c + dx) - \frac{6a^2(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \text{EllipticPi}(\frac{2}{a}, \frac{c + dx}{2}) + \frac{6b(-35a^5B + 15a^4Ab + 86a^3b^2B - 38a^2Ab^3 - 63ab^4B + 35Ab^5) \sin(c + dx)}{3bd\sqrt{\sec(c + dx)}}} b^2 d(a + b)}$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^3*Sec[c + d*x]^(7/2)),x]`

3.583. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^2(c + dx)} dx$

```
output (a*(A*b - a*B)*Sin[c + d*x])/(2*b*(a^2 - b^2)*d*Sqrt[Sec[c + d*x]]*(b + a*
Sec[c + d*x])^2) - (-((a*(3*a^2*A*b - 9*A*b^3 - 7*a^3*B + 13*a*b^2*B)*Sin[
c + d*x])/(b*(a^2 - b^2)*d*Sqrt[Sec[c + d*x]]*(b + a*Sec[c + d*x]))) + (-1
/3*(((6*b*(15*a^4*A*b - 29*a^2*A*b^3 + 8*A*b^5 - 35*a^5*B + 65*a^3*b^2*B -
24*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x
]])/d - (2*(45*a^5*A*b - 99*a^3*A*b^3 + 72*a*A*b^5 - 105*a^6*B + 223*a^4*b
^2*B - 128*a^2*b^4*B - 8*b^6*B)*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2,
2]*Sqrt[Sec[c + d*x]])/d)/b^2 + (6*a^2*(15*a^4*A*b - 38*a^2*A*b^3 + 35*A*b
^5 - 35*a^5*B + 86*a^3*b^2*B - 63*a*b^4*B)*Sqrt[Cos[c + d*x]]*EllipticPi[(
2*b)/(a + b), (c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(b^2*(a + b)*d))/b + (2*
(15*a^3*A*b - 33*a*A*b^3 - 35*a^4*B + 61*a^2*b^2*B - 8*b^4*B)*Sin[c + d*x]
)/(3*b*d*Sqrt[Sec[c + d*x]]))/(2*b*(a^2 - b^2)))/(4*b*(a^2 - b^2))
```

3.583.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 3284 Int[1/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_)
+ (f_)*(x_)])], x_Symbol] := Simp[(2/(f*(a + b)*Sqrt[c + d]))*EllipticPi[
2*(b/(a + b)), (1/2)*(e - Pi/2 + f*x), 2*(d/(c + d))], x] /; FreeQ[{a, b, c
, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && GtQ[c + d, 0]
```

rule 3439 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(g_.)^{(p_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_.)}*((c_.) + (d_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\text{Csc}[e + f*x])^{(p-m-n)}*(b + a*\text{Csc}[e + f*x])^m*(d + c*\text{Csc}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{!IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4258 $\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^n*\sin[c + d*x]^n \text{Int}[1/\sin[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x\} \&\& \text{EqQ}[n^2, 1/4]$

rule 4274 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[(d*\text{Csc}[e + f*x])^n, x], x] + \text{Simp}[b/d \text{Int}[(d*\text{Csc}[e + f*x])^{(n+1)}, x], x] /; \text{FreeQ}\{a, b, d, e, f, n\}, x\}$

rule 4336 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(3/2)} / (\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[d*\text{Sqrt}[d*\sin[e + f*x]]*\text{Sqrt}[d*\text{Csc}[e + f*x]] \text{Int}[1/(\text{Sqrt}[d*\sin[e + f*x]]*(b + a*\sin[e + f*x])), x], x] /; \text{FreeQ}\{a, b, d, e, f\}, x\} \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 4518 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.)^{(n_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^{(m_.)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m+1)}*((d*\text{Csc}[e + f*x])^n/(a*f*(m+1)*(a^2 - b^2))), x] + \text{Simp}[1/(a*(m+1)*(a^2 - b^2)) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m+1)}*(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*(a^2*(m+1) - b^2*(m+n+1)) + a*b*B*n - a*(A*b - a*B)*(m+1)*\text{Csc}[e + f*x] + b*(A*b - a*B)*(m+n+2)*\text{Csc}[e + f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, d, e, f, A, B, n\}, x\} \&\& \text{NeQ}[A*b - a*B, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -1] \&\& \text{!(IntegerQ}[m + 1/2, 0] \&\& \text{IntegerQ}[n, 0])$


```
rule 4588 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*(a + b*Csc
[e + f*x])^(m + 1)*((d*Csc[e + f*x])^n/(a*f*(m + 1)*(a^2 - b^2))), x] + Sim
p[1/(a*(m + 1)*(a^2 - b^2)) Int[(a + b*Csc[e + f*x])^(m + 1)*(d*Csc[e + f
*x])^n*Simp[a*(a*A - b*B + a*C)*(m + 1) - (A*b^2 - a*b*B + a^2*C)*(m + n +
1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m
+ n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x
] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && !(ILtQ[m + 1/2, 0] && ILtQ[n, 0])
```

```
rule 4592 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))^(m_), x_Symbol] := Simp[A*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m + 1)*((d
*Csc[e + f*x])^n/(a*f*n)), x] + Simp[1/(a*d*n) Int[(a + b*Csc[e + f*x])^m
*(d*Csc[e + f*x])^(n + 1)*Simp[a*B*n - A*b*(m + n + 1) + a*(A + A*n + C*n)*
Csc[e + f*x] + A*b*(m + n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d
, e, f, A, B, C, m}, x] && NeQ[a^2 - b^2, 0] && LeQ[n, -1]
```

```
rule 4594 Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.
))/(Sqrt[csc[(e_.) + (f_.)*(x_)]*(d_.)]*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a
_))), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)/(a^2*d^2) Int[(d*Csc[e +
f*x])^(3/2)/(a + b*Csc[e + f*x]), x], x] + Simp[1/a^2 Int[(a*A - (A*b - a
*B)*Csc[e + f*x])/Sqrt[d*Csc[e + f*x]], x], x] /; FreeQ[{a, b, d, e, f, A,
B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.583.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2194 vs. 2(569) = 1138.

Time = 17.79 (sec) , antiderivative size = 2195, normalized size of antiderivative = 4.21

method	result	size
default	Expression too large to display	2195

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^3/sec(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)
```

$$3.583. \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^3 \sec^2(c+dx)} dx$$

output
$$\begin{aligned} & -(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*a^4*(A*b-B*a) \\ &)/b^5*(-1/2/a*b^2/(a^2-b^2)*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin \\ & (1/2*d*x+1/2*c)^2)^{(1/2)}/(2*b*\cos(1/2*d*x+1/2*c)^2+a-b)^2-3/4*b^2*(3*a^2- \\ & b^2)/a^2/(a^2-b^2)^2*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d \\ & *x+1/2*c)^2)^{(1/2)}/(2*b*\cos(1/2*d*x+1/2*c)^2+a-b)-7/8/(a+b)/(a^2-b^2)*(\sin \\ & (1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x \\ & +1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)} \\ &)+1/4/(a+b)/(a^2-b^2)/a*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c \\ &)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*Elliptic \\ & F(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b+3/8/(a+b)/(a^2-b^2)/a^2*(\sin(1/2*d*x+1/2*c \\ &)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(\\ & 1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*b^2-9/8*b/(a \\ & ^2-b^2)^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(\\ & -2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticF(\cos(1/2*d*x+ \\ & 1/2*c),2^{(1/2)})+3/8*b^3/a^2/(a^2-b^2)^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*c \\ & \cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2 \\ &)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})+9/8*b/(a^2-b^2)^2*(\sin(1/2*d \\ & *x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-2*\sin(1/2*d*x+1/2*c \\ &)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})-3/8* \\ & b^3/a^2/(a^2-b^2)^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)... \end{aligned}$$

3.583.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(7/2),x, algorithm="fracas")`

output `Timed out`

3.583.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**3/sec(d*x+c)**(7/2),x)`output `Timed out`**3.583.7 Maxima [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(7/2),x, algorithm m="maxima")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2)), x)`**3.583.8 Giac [F]**

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^3 \sec(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^3/sec(d*x+c)^(7/2),x, algorithm m="giac")`output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^3*sec(d*x + c)^(7/2)), x)`

3.583.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^3 \sec^{\frac{7}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{7/2} (a + b \cos(c + dx))^3} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^3),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^3),x)`

3.584
$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

3.584.1 Optimal result	5334
3.584.2 Mathematica [A] (verified)	5334
3.584.3 Rubi [A] (verified)	5335
3.584.4 Maple [B] (verified)	5337
3.584.5 Fricas [C] (verification not implemented)	5337
3.584.6 Sympy [F(-1)]	5338
3.584.7 Maxima [F]	5338
3.584.8 Giac [F]	5338
3.584.9 Mupad [F(-1)]	5339

3.584.1 Optimal result

Integrand size = 36, antiderivative size = 64

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d} + \frac{2B \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3d}$$

output

```
2/3*B*sec(d*x+c)^(3/2)*sin(d*x+c)/d+2/3*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d
```

3.584.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.73

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{2B \sec^{\frac{3}{2}}(c + dx) \left(\cos^{\frac{3}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(c + dx) \right)}{3d}$$

input

```
Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x]),x]
```

output $(2*B*\text{Sec}[c + d*x]^{(3/2)}*(\text{Cos}[c + d*x]^{(3/2)}*\text{EllipticF}[(c + d*x)/2, 2] + \text{Sin}[c + d*x]))/(3*d)$

3.584.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {2011, 3042, 4255, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c+dx)(aB+bB\cos(c+dx))}{a+b\cos(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \sec^{\frac{5}{2}}(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \csc\left(c+dx+\frac{\pi}{2}\right)^{5/2} dx$$

$$\downarrow \text{4255}$$

$$B \left(\frac{1}{3} \int \sqrt{\sec(c+dx)} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{1}{3} \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right)$$

$$\downarrow \text{4258}$$

$$B \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{1}{3} \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{2 \sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{3d} \right)$$

$$\downarrow \text{3120}$$

3.584. $\int \frac{(aB+bB\cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b\cos(c+dx)} dx$

$$B\left(\frac{2\sin(c+dx)\sec^{\frac{3}{2}}(c+dx)}{3d} + \frac{2\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d}\right)$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x]),x]`

output `B*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(3*d)`

3.584.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_.))^(m_.)*((c_) + (d_.)*(v_.))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]`

3.584.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(80) = 160.

Time = 3.66 (sec) , antiderivative size = 214, normalized size of antiderivative = 3.34

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}$

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `-2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*B*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^(3/2)/sin(1/2*d*x+1/2*c)/d`

3.584.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.42

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{-i \sqrt{2} B \cos(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} B \cos(dx + c) \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 * B * \sin(dx + c) / \sqrt{\cos(dx + c)}}{3 d \cos(dx + c)}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `1/3*(-I*sqrt(2)*B*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*cos(d*x + c)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*B*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))`

3.584. $\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{a+b \cos(c+dx)} dx$

3.584.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+b*cos(d*x+c)),x)`

output `Timed out`

3.584.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{5}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.584.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{5}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a), x)`

3.584.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (Ba + Bb \cos(c + dx))}{a + b \cos(c + dx)} dx$$

input `int(((1/cos(c + d*x))^(5/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `int(((1/cos(c + d*x))^(5/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.585
$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

3.585.1 Optimal result	5340
3.585.2 Mathematica [A] (verified)	5340
3.585.3 Rubi [A] (verified)	5341
3.585.4 Maple [B] (verified)	5343
3.585.5 Fricas [C] (verification not implemented)	5343
3.585.6 Sympy [F]	5344
3.585.7 Maxima [F]	5344
3.585.8 Giac [F]	5344
3.585.9 Mupad [F(-1)]	5345

3.585.1 Optimal result

Integrand size = 36, antiderivative size = 60

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= -\frac{2B \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d} + \frac{2B \sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

output

```
2*B*sin(d*x+c)*sec(d*x+c)^(1/2)/d-2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2
*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x
+c)^(1/2)/d
```

3.585.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.77

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{2B \sqrt{\sec(c + dx)} \left(-\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d}$$

input

```
Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x
]),x]
```

output $(2*B*\text{Sqrt}[\text{Sec}[c + d*x]]*(-(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2]) + \text{Sin}[c + d*x]))/d$

3.585.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {2011, 3042, 4255, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(aB + bB \cos(c + dx))}{a + b \cos(c + dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \sec^{\frac{3}{2}}(c + dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} dx$$

$$\downarrow \text{4255}$$

$$B \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\sec(c + dx)}} dx \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \int \frac{1}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}} dx \right)$$

$$\downarrow \text{4258}$$

$$B \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin\left(c + dx + \frac{\pi}{2}\right)} dx \right)$$

$$\downarrow \text{3119}$$

3.585. $\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx$

$$B \left(\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d} - \frac{2 \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d} \right)$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x]),x]`

output `B*((-2*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d + (2*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/d`

3.585.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1))
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
&& IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]`

3.585.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(80) = 160.

Time = 2.95 (sec) , antiderivative size = 183, normalized size of antiderivative = 3.05

method	result
default	$-\frac{2B \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1} \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) - 1}} \right) - 1}{d}$

input `int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b),x,method=_RETURNVERBOSE)`

output `-2*B*(-2*cos(1/2*d*x+1/2*c)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*sin(1/2*d*x+1/2*c)^2+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.585.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.27

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx$$

$$= \frac{-i \sqrt{2} B \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2} B \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 * B * \sin(dx + c) / \sqrt{\cos(dx + c)}}{d}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `(-I*sqrt(2)*B*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + I*sqrt(2)*B*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*B*sin(d*x + c)/sqrt(cos(d*x + c)))/d`

3.585.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = B \int \sec^{\frac{3}{2}}(c + dx) dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c)),x)`

output `B*Integral(sec(c + d*x)**(3/2), x)`

3.585.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{3}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.585.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{3}{2}}}{b \cos(dx + c) + a} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a), x)`

3.585.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{a + b \cos(c + dx)} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (Ba + Bb \cos(c + dx))}{a + b \cos(c + dx)} dx$$

input `int(((1/cos(c + d*x))^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `int(((1/cos(c + d*x))^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.586
$$\int \frac{(aB + bB \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

3.586.1 Optimal result	5346
3.586.2 Mathematica [A] (verified)	5346
3.586.3 Rubi [A] (verified)	5347
3.586.4 Maple [B] (verified)	5348
3.586.5 Fricas [C] (verification not implemented)	5349
3.586.6 Sympy [F]	5349
3.586.7 Maxima [F]	5349
3.586.8 Giac [F]	5350
3.586.9 Mupad [F(-1)]	5350

3.586.1 Optimal result

Integrand size = 36, antiderivative size = 37

$$\int \frac{(aB + bB \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

$$= \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d`

3.586.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{(aB + bB \cos(c + dx)) \sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

$$= \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{d}$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x]),x]`

output `(2*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d`

3.586.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {2011, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\sec(c+dx)}(aB + bB \cos(c+dx))}{a + b \cos(c+dx)} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \sqrt{\sec(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \sqrt{\csc\left(c+dx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{4258} \\
 & B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin\left(c+dx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3120} \\
 & \frac{2B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d}
 \end{aligned}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x]),x]`

output `(2*B*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d`

3.586.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :>
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && ( !IntegerQ[n] || SimplrQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3120 Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 4258 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

3.586.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. 2(59) = 118.

Time = 2.07 (sec) , antiderivative size = 134, normalized size of antiderivative = 3.62

method	result	size
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}B\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d}$	134

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b),x,method=_RETUR
NVERBOSE)
```

```
output -2*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(sin(1/2*d*x+
1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-2*sin(1/2*d*x+1/2*c)^4
+sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))/sin(1/2
*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.586. $\int \frac{(aB+bB \cos(c+dx))\sqrt{\sec(c+dx)}}{a+b \cos(c+dx)} dx$

3.586.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.43

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx$$

$$= \frac{-i\sqrt{2}B\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i\sqrt{2}B\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="fracas")`

output `(-I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

3.586.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = B \int \sqrt{\sec(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c)),x)`

output `B*Integral(sqrt(sec(c + d*x)), x)`

3.586.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\sec(dx + c)}}{b \cos(dx + c) + a} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a), x)`

3.586. $\int \frac{(aB+bB \cos(c+dx))\sqrt{\sec(c+dx)}}{a+b \cos(c+dx)} dx$

3.586.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\sec(dx + c)}}{b \cos(dx + c) + a} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a), x)`

3.586.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{a + b \cos(c + dx)} dx = \int \frac{\sqrt{\frac{1}{\cos(c+dx)}} (Ba + Bb \cos(c + dx))}{a + b \cos(c + dx)} dx$$

input `int(((1/cos(c + d*x))^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)),x)`

output `int(((1/cos(c + d*x))^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x)), x)`

3.587
$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sqrt{\sec(c + dx)}} dx$$

3.587.1 Optimal result 5351
 3.587.2 Mathematica [A] (verified) 5351
 3.587.3 Rubi [A] (verified) 5352
 3.587.4 Maple [B] (verified) 5353
 3.587.5 Fricas [C] (verification not implemented) 5354
 3.587.6 Sympy [F] 5354
 3.587.7 Maxima [F] 5355
 3.587.8 Giac [F] 5355
 3.587.9 Mupad [F(-1)] 5355

3.587.1 Optimal result

Integrand size = 36, antiderivative size = 37

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sqrt{\sec(c + dx)}} dx = \frac{2B \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{d}$$

output `2*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d`

3.587.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sqrt{\sec(c + dx)}} dx = \frac{2BE\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)}}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]),x]`

output `(2*B*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]])`

3.587.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {2011, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sqrt{\sec(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4258} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3119} \\
 & \frac{2B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E(\frac{1}{2}(c + dx) | 2)}{d}
 \end{aligned}$$

input `Int[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]),x]`

output `(2*B*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/d`

3.587.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x],
  a + b*x)
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3119 Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

```
rule 4258 Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

3.587.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. 2(59) = 118.

Time = 3.26 (sec) , antiderivative size = 134, normalized size of antiderivative = 3.62

method	result
default	$\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} B\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} d$
risch	$-\frac{i\sqrt{2} B}{d\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} - \frac{i\left(-\frac{2(e^{2i(dx+c)+1})}{\sqrt{(e^{2i(dx+c)+1})e^{i(dx+c)}}} + \frac{i\sqrt{-i(e^{i(dx+c)+i})}\sqrt{2}\sqrt{i(e^{i(dx+c)-i})}\sqrt{ie^{i(dx+c)}}(-2iE\left(\sqrt{-i(e^{i(dx+c)+i})}, \frac{\sqrt{2}}{2}\right)}{\sqrt{e^{3i(dx+c)+e^{i(dx+c)}}}}}\right)}{d\sqrt{\frac{e^{i(dx+c)}}{e^{2i(dx+c)+1}}}} (e^{2i(dx+c)+1})$

```
input int((B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)/sec(d*x+c)^(1/2), x, method=_RETUR
NVERBOSE)
```

3.587.
$$\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))\sqrt{\sec(c+dx)}} dx$$

output $2*((2*\cos(1/2*d*x+1/2*c)^2-1)*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*B*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})/(-2*\sin(1/2*d*x+1/2*c)^4+\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)^{(1/2)}/d$

3.587.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx$$

$$= \frac{i\sqrt{2}B\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i\sqrt{2}B\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output $(I*\text{sqrt}(2)*B*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - I*\text{sqrt}(2)*B*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/d$

3.587.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = B \int \frac{1}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `B*Integral(1/sqrt(sec(c + d*x)), x)`

3.587.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)\sqrt{\sec(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.587.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)\sqrt{\sec(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.587.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))\sqrt{\sec(c + dx)}} dx = \int \frac{Ba + Bb \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + b \cos(c + dx))} dx$$

input `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))),x)`

output `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))), x)`

3.588
$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

3.588.1 Optimal result	5356
3.588.2 Mathematica [A] (verified)	5356
3.588.3 Rubi [A] (verified)	5357
3.588.4 Maple [B] (verified)	5359
3.588.5 Fracas [C] (verification not implemented)	5359
3.588.6 Sympy [F]	5360
3.588.7 Maxima [F]	5360
3.588.8 Giac [F]	5360
3.588.9 Mupad [F(-1)]	5361

3.588.1 Optimal result

Integrand size = 36, antiderivative size = 64

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \frac{2B \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) \sqrt{\sec(c + dx)}}{3d} + \frac{2B \sin(c + dx)}{3d \sqrt{\sec(c + dx)}}$$

output `2/3*B*sin(d*x+c)/d/sec(d*x+c)^(1/2)+2/3*B*(cos(1/2*d*x+1/2*c)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d`

3.588.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.78

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \frac{B \sqrt{\sec(c + dx)} \left(2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx)) \right)}{3d}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sec[c + d*x]^(3/2)),x]`

3.588.
$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

output $(B\sqrt{\sec(c + dx)}*(2\sqrt{\cos(c + dx)}*\text{EllipticF}[(c + dx)/2, 2] + \text{Sin}[2*(c + dx)]))/(3*d)$

3.588.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {2011, 3042, 4256, 3042, 4258, 3042, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{aB + bB \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{1}{\sec^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx \\ & \quad \downarrow \text{4256} \\ & B \left(\frac{1}{3} \int \sqrt{\sec(c + dx)} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{1}{3} \int \sqrt{\csc(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\ & \quad \downarrow \text{4258} \\ & B \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{1}{3} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{3d\sqrt{\sec(c + dx)}} \right) \\ & \quad \downarrow \text{3120} \end{aligned}$$

3.588. $\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$

$$B \left(\frac{2 \sin(c + dx)}{3d \sqrt{\sec(c + dx)}} + \frac{2 \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d} \right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sec[c + d*x]^(3/2)),x]`

output `B*((2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(3*d) + (2*Sin[c + d*x])/(3*d*Sqrt[Sec[c + d*x]])`

3.588.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c
+ d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]`

3.588.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(80) = 160.

Time = 4.02 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.81

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}}{3\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}d$

input `int((B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `-2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d`

3.588.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.11

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2B\sqrt{\cos(dx+c)}\sin(dx+c) - i\sqrt{2}B\text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}B\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{3d}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fracas")`

output `1/3*(2*B*sqrt(cos(d*x + c))*sin(d*x + c) - I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*B*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d`

3.588.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = B \int \frac{1}{\sec^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `B*Integral(sec(c + d*x)**(-3/2), x)`

3.588.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.588.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.588.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{Ba + Bb \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (a + b \cos(c + dx))} dx$$

input `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))),x)`

output `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))), x)`

3.589
$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx$$

3.589.1 Optimal result	5362
3.589.2 Mathematica [A] (verified)	5362
3.589.3 Rubi [A] (verified)	5363
3.589.4 Maple [B] (verified)	5365
3.589.5 Fracas [C] (verification not implemented)	5365
3.589.6 Sympy [F(-1)]	5366
3.589.7 Maxima [F]	5366
3.589.8 Giac [F]	5366
3.589.9 Mupad [F(-1)]	5367

3.589.1 Optimal result

Integrand size = 36, antiderivative size = 64

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \frac{6B \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{\sec(c + dx)}}{5d} + \frac{2B \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)}$$

output `2/5*B*sin(d*x+c)/d/sec(d*x+c)^(3/2)+6/5*B*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*sec(d*x+c)^(1/2)/d`

3.589.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.88

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \frac{B \sqrt{\sec(c + dx)} \left(12 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) + \sin(3(c + dx)) \right)}{10d}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sec[c + d*x]^(5/2)),x]`

```
output (B*Sqrt[Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*d)
```

3.589.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {2011, 3042, 4256, 3042, 4258, 3042, 3119}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sec^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{5/2}} dx \\
 & \quad \downarrow \text{4256} \\
 & B \left(\frac{3}{5} \int \frac{1}{\sqrt{\sec(c + dx)}} dx + \frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{3}{5} \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx + \frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \\
 & \quad \downarrow \text{4258} \\
 & B \left(\frac{3}{5} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} dx + \frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{3}{5} \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\sin(c + dx + \frac{\pi}{2})} dx + \frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} \right) \\
 & \quad \downarrow \text{3119}
 \end{aligned}$$

3.589. $\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx$

$$B \left(\frac{2 \sin(c + dx)}{5d \sec^{\frac{3}{2}}(c + dx)} + \frac{6 \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d} \right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])*Sec[c + d*x]^(5/2)),x]`

output `B*((6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]*Sqrt[Sec[c + d*x]])/(5*d) + (2*Sin[c + d*x])/(5*d*Sec[c + d*x]^(3/2)))`

3.589.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 4256 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d*n)), x] + Simp[(n + 1)/(b^2*n) Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

rule 4258 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^n*Sin[c + d*x]^n Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

3.589.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 202 vs. 2(80) = 160.

Time = 5.03 (sec) , antiderivative size = 203, normalized size of antiderivative = 3.17

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)B\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}{d}$

```
input int((B*a+b*B*cos(d*x+c))/(a+cos(d*x+c)*b)/sec(d*x+c)^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/5*((2*cos(1/2*d*x+1/2*c)^2-1)*sin(1/2*d*x+1/2*c)^2)^(1/2)*B*(-8*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+8*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-2*sin(1/2*d*x+1/2*c)^4+sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)/d
```

3.589.5 Fracas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.20

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2 B \cos(dx + c)^{\frac{3}{2}} \sin(dx + c) + 3i \sqrt{2} B \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 3i \sqrt{2} B \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{5}$$

```
input integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(5/2),x, algorithm="fracas")
```

```
output 1/5*(2*B*cos(d*x + c)^(3/2)*sin(d*x + c) + 3*I*sqrt(2)*B*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 3*I*sqrt(2)*B*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d
```

3.589.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)**(5/2),x)`output `Timed out`**3.589.7 Maxima [F]**

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(5/2),x, algorithm="maxima")`output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sec(d*x + c)^(5/2)), x)`**3.589.8 Giac [F]**

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a) \sec(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))/sec(d*x+c)^(5/2),x, algorithm="giac")`output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)*sec(d*x + c)^(5/2)), x)`

3.589.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{Ba + Bb \cos(c + dx)}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2} (a + b \cos(c + dx))} dx$$

input `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))),x)`

output `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))), x)`

3.590 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

3.590.1 Optimal result	5368
3.590.2 Mathematica [A] (warning: unable to verify)	5369
3.590.3 Rubi [A] (verified)	5370
3.590.4 Maple [B] (verified)	5375
3.590.5 Fracas [F]	5376
3.590.6 Sympy [F(-1)]	5377
3.590.7 Maxima [F]	5377
3.590.8 Giac [F]	5377
3.590.9 Mupad [F(-1)]	5378

3.590.1 Optimal result

Integrand size = 35, antiderivative size = 473

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{2(a - b)\sqrt{a + b}(19a^2Ab + 8Ab^3 + 63a^3B - 14ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right)\right)}{105a^4d\sqrt{\sec(c + dx)}} + \frac{2(a - b)\sqrt{a + b}(8Ab^2 + a^2(25A - 63B) + 2ab(3A - 7B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right)\right)}{105a^3d\sqrt{\sec(c + dx)}} + \frac{2(25a^2A - 4Ab^2 + 7abB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{105a^2d} + \frac{2(Ab + 7aB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{35ad} + \frac{2A\sqrt{a + b \cos(c + dx)} \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{7d}$$

output $\frac{2}{105}*(25*A*a^2-4*A*b^2+7*B*a*b)*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/a^2/d+2/35*(A*b+7*B*a)*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/a/d+2/7*A*\sec(d*x+c)^{(7/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/d+2/105*(a-b)*(19*A*a^2*b+8*A*b^3+63*B*a^3-14*B*a*b^2)*\csc(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b))^{(1/2)}/a^4/d/\sec(d*x+c)^{(1/2)}+2/105*(a-b)*(8*A*b^2+a^2*(25*A-63*B)+2*a*b*(3*A-7*B))*\csc(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b))^{(1/2)}/a^3/d/\sec(d*x+c)^{(1/2)}$

3.590.2 Mathematica [A] (warning: unable to verify)

Time = 15.83 (sec) , antiderivative size = 519, normalized size of antiderivative = 1.10

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \frac{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)} \sec(c + dx) \left(-2(a + b)(19a^2Ab + 8Ab^3 + 63a^3B - 14ab^2B)\sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}}\sqrt{\frac{a+b\cos(c+dx)}{(a+b)(1+\cos(c+dx))}}\right)}{105a^3} + \frac{2\sec^2(c+dx)(Ab\sin(c+dx)+7aB\sin(c+dx))}{35a}$$

d

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output $(2\sqrt{\cos[(c+dx)/2]^2 \sec[c+dx]} \cdot (-2(a+b)(19a^2Ab + 8Ab^3 + 63a^3B - 14ab^2B)) \sqrt{\cos[c+dx]/(1+\cos[c+dx])} \sqrt{(a+b\cos[c+dx])/((a+b)(1+\cos[c+dx]))} \text{EllipticE}[\text{ArcSin}[\text{Tan}[(c+dx)/2]]], (-a+b)/(a+b) + 2a(a+b)(8Ab^2 - 2ab(3A+7B) + a^2(25A+63B)) \sqrt{\cos[c+dx]/(1+\cos[c+dx])} \sqrt{(a+b\cos[c+dx])/((a+b)(1+\cos[c+dx]))} \text{EllipticF}[\text{ArcSin}[\text{Tan}[(c+dx)/2]]], (-a+b)/(a+b) - (19a^2Ab + 8Ab^3 + 63a^3B - 14ab^2B) \cos[c+dx] \cdot (a+b\cos[c+dx]) \sec[(c+dx)/2]^2 \text{Tan}[(c+dx)/2]) / (105a^3 d \sqrt{a+b\cos[c+dx]} \sqrt{\sec[(c+dx)/2]^2} + (\sqrt{a+b\cos[c+dx]} \sqrt{\sec[c+dx]} \cdot ((2(19a^2Ab + 8Ab^3 + 63a^3B - 14ab^2B) \sin[c+dx]) / (105a^3) + (2\sec[c+dx]^2 (Ab\sin[c+dx] + 7aB\sin[c+dx])) / (35a) + (2\sec[c+dx] (25a^2A\sin[c+dx] - 4Ab^2\sin[c+dx] + 7abB\sin[c+dx])) / (105a^2) + (2A\sec[c+dx]^2 \text{Tan}[c+dx]) / 7)) / d$

3.590.3 Rubi [A] (verified)

Time = 2.34 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3478, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{9}{2}}(c+dx) \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c+dx+\frac{\pi}{2}\right)^{\frac{9}{2}} \sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} (A+B\sin\left(c+dx+\frac{\pi}{2}\right)) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{9}{2}}} dx$$

$$\downarrow \text{3478}$$

3.590. $\int \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{7}\int\frac{4Ab\cos^2(c+dx)+(5aA+7bB)\cos(c+dx)+Ab+7aB}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{4Ab\cos^2(c+dx)+(5aA+7bB)\cos(c+dx)+Ab+7aB}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{4Ab\sin(c+dx+\frac{\pi}{2})^2+(5aA+7bB)\sin(c+dx+\frac{\pi}{2})+Ab+7aB}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)}\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2\int\frac{25Aa^2+7bBa+(23Ab+21aB)\cos(c+dx)a-4Ab^2+2b(Ab+7aB)\cos^2(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{5a}+\frac{2(7aB+Ab)}{7d\cos^{\frac{7}{2}}(c+dx)}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{25Aa^2+7bBa+(23Ab+21aB)\cos(c+dx)a-4Ab^2+2b(Ab+7aB)\cos^2(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{5a}+\frac{2(7aB+Ab)}{7d\cos^{\frac{7}{2}}(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{25Aa^2+7bBa+(23Ab+21aB)\sin(c+dx+\frac{\pi}{2})a-4Ab^2+2b(Ab+7aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{5a}+\frac{2(7aB+Ab)}{7d\cos^{\frac{7}{2}}(c+dx)}\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2\int\frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\cos(c+dx)a+8Ab^3}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{3a}+\frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)}\right)\right)$$

↓ 27

3.590. $\int\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\cos(c+dx)a+8Ab^3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{3a}+\frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)}{3ad\cos^{\frac{3}{2}}(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{63Ba^3+19Aba^2-14b^2Ba+(25Aa^2+49bBa+2Ab^2)\sin(c+dx+\frac{\pi}{2})a+8Ab^3}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{3a}+\frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx+\frac{\pi}{2})}{3ad\cos^{\frac{3}{2}}(c+dx+\frac{\pi}{2})}\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{(a-b)(a^2(25A-63B)+2ab(3A-7B)+8Ab^2)\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}dx+(63a^3B+19a^2Ab-14ab^2B)}{3a}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{(a-b)(a^2(25A-63B)+2ab(3A-7B)+8Ab^2)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+(63a^3B+19a^2Ab-14ab^2B)}{3a}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{(63a^3B+19a^2Ab-14ab^2B+8Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)+2ab(3A-7B)+8Ab^2)}{3a}\right)\right)$$

↓ 3473

3.590. $\int\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2(25a^2A+7abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} + \frac{2(a-b)\sqrt{a+b}(a^2(25A-63B)+2ab(3A-7B)+8Ab^2)}{\dots}\right)\right)$$

```
input Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(A*b + 7*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*a*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(19*a^2*A*b + 8*A*b^3 + 63*a^3*B - 14*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(8*A*b^2 + a^2*(25*A - 63*B) + 2*a*b*(3*A - 7*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(3*a) + (2*(25*a^2*A - 4*A*b^2 + 7*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/(5*a))/7)
```

3.590.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*SIN[e + f*x]]/Sqrt[b*SIN[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*SIN[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + SIN[e + f*x])/((a + b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3478 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m + 1)*((c + d*SIN[e + f*x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*SIN[e + f*x] - d*(A*b - a*B)*(m + n + 2)*SIN[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.590.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4627 vs. $2(427) = 854$.

Time = 29.53 (sec) , antiderivative size = 4628, normalized size of antiderivative = 9.78

method	result	size
parts	Expression too large to display	4628
default	Expression too large to display	4673

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

```

output 2/105*A/d*sec(d*x+c)^(9/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(25*a^4*cos
os(d*x+c)^4*sin(d*x+c)+15*a^4*cos(d*x+c)^2*sin(d*x+c)-a^2*b^2*cos(d*x+c)^4
*sin(d*x+c)+8*cos(d*x+c)^5*b^4*sin(d*x+c)+16*EllipticE(cot(d*x+c)-csc(d*x+
c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(co
s(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^5-50*EllipticF(cot(d*x+c)-cs
c(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/
2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^5+8*EllipticE(cot(d*x+
c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b
))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^4-25*EllipticF(co
t(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))
)/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^4+8*Ellipti
cE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*
x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b^3*cos(d*x+c)^4-19
*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(
1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3*b*cos(d*x
+c)^4-2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+
c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2*b^
2*cos(d*x+c)^4-8*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a
+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/
2)*a*b^3*cos(d*x+c)^4+38*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))...

```

3.590.5 Fracas [F]

$$\begin{aligned}
 & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\
 &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{9}{2}} dx
 \end{aligned}$$

```

input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fricas")

```

```

output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(9/2),
x)

```

3.590.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(9/2)*(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.590.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{9}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(9/2), x)`

3.590.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{9}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(9/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(9/2), x)`

3.590. $\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

3.590.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{9/2} \sqrt{a + b \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^(1/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^(1/2), x)`

3.591 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.591.1 Optimal result	5379
3.591.2 Mathematica [A] (warning: unable to verify)	5380
3.591.3 Rubi [A] (verified)	5380
3.591.4 Maple [B] (verified)	5384
3.591.5 Fricas [F]	5385
3.591.6 Sympy [F(-1)]	5386
3.591.7 Maxima [F]	5386
3.591.8 Giac [F]	5386
3.591.9 Mupad [F(-1)]	5387

3.591.1 Optimal result

Integrand size = 35, antiderivative size = 390

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \frac{2(a - b)\sqrt{a + b}(9a^2 A - 2Ab^2 + 5abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right) \mid -\frac{a + b}{a - b}\right) \sqrt{\frac{a + b}{a - b}}}{15a^3 d \sqrt{\sec(c + dx)}} - \frac{2(a - b)\sqrt{a + b}(9aA + 2Ab - 5aB) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), -\frac{a + b}{a - b}\right)}{15a^2 d \sqrt{\sec(c + dx)}} + \frac{2(Ab + 5aB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{15ad} + \frac{2A \sqrt{a + b \cos(c + dx)} \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{5d}$$

output

```
2/15*(A*b+5*B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d+2/5*A*sec(d*x+c)^(5/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/15*(a-b)*(9*A*a^2-2*A*b^2+5*B*a*b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2))/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d/sec(d*x+c)^(1/2)-2/15*(a-b)*(9*A*a+2*A*b-5*B*a)*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d/sec(d*x+c)^(1/2)
```

3.591.2 Mathematica [A] (warning: unable to verify)

Time = 11.88 (sec) , antiderivative size = 373, normalized size of antiderivative = 0.96

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= 2 \left(\sqrt{\cos^2\left(\frac{1}{2}(c+dx)\right) \sec(c+dx)} \left(-2(a+b)(9a^2A - 2Ab^2 + 5abB) E\left(\arcsin\left(\tan\left(\frac{1}{2}(c+dx)\right)\right) \middle| \frac{-a+b}{a+b}\right) \sqrt{\frac{1}{1+\sec(c+dx)}} \sqrt{\frac{b+a \sec(c+dx)}{(a+b)(1+\sec(c+dx))}} + 2a(a+b) \right) \right)$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*((Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(9*a^2*A - 2*A*b^2 + 5*a*b*B)*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x])]) + 2*a*(a + b)*(9*a*A - 2*A*b + 5*a*B)*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x])]) - (9*a^2*A - 2*A*b^2 + 5*a*b*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2 + (a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]*((9*a^2*A - 2*A*b^2 + 5*a*b*B)*Sin[c + d*x] + a*(A*b + 5*a*B + 3*a*A*Sec[c + d*x])*Tan[c + d*x])))/(15*a^2*d*Sqrt[a + b*Cos[c + d*x]])`

3.591.3 Rubi [A] (verified)Time = 1.61 (sec) , antiderivative size = 380, normalized size of antiderivative = 0.97, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3478, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{7}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}(A + B \sin\left(c + dx + \frac{\pi}{2}\right)) dx$$

$$\downarrow \text{3440}$$

3.591. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)}dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int\frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx$$

↓ 3478

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{5}\int\frac{2Ab\cos^2(c+dx)+(3aA+5bB)\cos(c+dx)+Ab+5aB}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5d\cos^{\frac{5}{2}}(c+dx)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\int\frac{2Ab\cos^2(c+dx)+(3aA+5bB)\cos(c+dx)+Ab+5aB}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5d\cos^{\frac{5}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\int\frac{2Ab\sin(c+dx+\frac{\pi}{2})^2+(3aA+5bB)\sin(c+dx+\frac{\pi}{2})+Ab+5aB}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2A\cos(c+dx)\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}{5d\sin^{\frac{5}{2}}(c+dx+\frac{\pi}{2})}\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{2\int\frac{9Aa^2+5bBa+(7Ab+5aB)\cos(c+dx)a-2Ab^2}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{3a}+\frac{2(5aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{\int\frac{9Aa^2+5bBa+(7Ab+5aB)\cos(c+dx)a-2Ab^2}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{3a}+\frac{2(5aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{\int\frac{9Aa^2+5bBa+(7Ab+5aB)\sin(c+dx+\frac{\pi}{2})a-2Ab^2}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{3a}+\frac{2(5aB+Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)}\right)\right)$$

↓ 3477

3.591. $\int\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{(9a^2A+5abB-2Ab^2)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-(a-b)(9aA-5aB+2Ab^2)}{3a}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{(9a^2A+5abB-2Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-(a-b)(9aA-5aB+2Ab^2)}{3a}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{(9a^2A+5abB-2Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2(a-b)\sqrt{a+b}(9aA-5aB+2Ab^2)}{3a}}{3a}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{2(a-b)\sqrt{a+b}(9a^2A+5abB-2Ab^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{a^2d}\right)\right)$$

```
input Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(9*a^2*A - 2*A*b^2 + 5*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(9*a*A + 2*A*b - 5*a*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(3*a) + (2*(A*b + 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/5)
```

3.591.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_)] + (f_)*(x_))*(g_)]^(p_)*((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3473 `Int[((A_)] + (B_)*sin[(e_)] + (f_)*(x_)]/(((b_)*sin[(e_)] + (f_)*(x_)]^(3/2)*Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`
- rule 3477 `Int[((A_)] + (B_)*sin[(e_)] + (f_)*(x_)]/(((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(3/2)*Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3478 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(B*a - A*b)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m + 1)*((c + d*SIN[e + f*
x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a
+ b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(
m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*
SIN[e + f*x] - d*(A*b - a*B)*(m + n + 2)*SIN[e + f*x]^2, x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*SIN[e + f*x])^(m + 1)*((c + d*SIN[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*SIN[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*SIN[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.591.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3319 vs. 2(350) = 700.

Time = 25.14 (sec) , antiderivative size = 3320, normalized size of antiderivative = 8.51

method	result	size
parts	Expression too large to display	3320
default	Expression too large to display	3349

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

```

output -2/15*A/d*sec(d*x+c)^(7/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(4*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4+14*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^4-4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4-9*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3+2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3+7*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3-2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3+9*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^3*cos(d*x+c)^3-3*sin(d*x+c)*cos(d*x+c)*a^3-4*sin(d*x+c)*cos(d*x+c)^2*a^2*b-18*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+...

```

3.591.5 Fracas [F]

$$\begin{aligned}
 & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\
 &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{7}{2}} dx
 \end{aligned}$$

```

input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")

```

```

output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)

```


3.591.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(7/2)*(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.591.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.591.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{7}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.591. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$

3.591.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} \sqrt{a + b \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^(1/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^(1/2), x)`

3.592 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.592.1 Optimal result	5388
3.592.2 Mathematica [A] (verified)	5389
3.592.3 Rubi [A] (verified)	5389
3.592.4 Maple [B] (verified)	5392
3.592.5 Fracas [F]	5393
3.592.6 Sympy [F(-1)]	5394
3.592.7 Maxima [F]	5394
3.592.8 Giac [F]	5394
3.592.9 Mupad [F(-1)]	5395

3.592.1 Optimal result

Integrand size = 35, antiderivative size = 324

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2(a - b)\sqrt{a + b}(Ab + 3aB)\sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{3a^2 d \sqrt{\sec(c + dx)}} + \frac{2(a - b)\sqrt{a + b}(A - 3B)\sqrt{\cos(c + dx)} \csc(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{3ad \sqrt{\sec(c + dx)}} + \frac{2A\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3d}$$

output

```
2/3*A*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/3*(a-b)*(A*b+
3*B*a)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(
1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))
/(a+b))^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^2/d/sec(d*x+c)^(1/2)+2/3*(a
-b)*(A-3*B)*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*
x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*
x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a/d/sec(d*x+c)^(1/2)
```

3.592.2 Mathematica [A] (verified)

Time = 11.00 (sec) , antiderivative size = 318, normalized size of antiderivative = 0.98

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \frac{2 \left((a + b \cos(c + dx))(aA + (Ab + 3aB) \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx) + \sqrt{\cos^2\left(\frac{1}{2}(c+dx)\right) \sec(c+dx)} \left(-2 \right) \right)}{}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `(2*((a + b*Cos[c + d*x])*(a*A + (A*b + 3*a*B)*Cos[c + d*x])*Sec[c + d*x]^(3/2)*Sin[c + d*x] + (Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(A*b + 3*a*B)*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x])]) + 2*a*(a + b)*(A + 3*B)*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x])])]) - (A*b + 3*a*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2])/(3*a*d*Sqrt[a + b*Cos[c + d*x]])`

3.592.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 306, normalized size of antiderivative = 0.94, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3478, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{5}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{5}{2}} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

3.592. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\cos^{\frac{5}{2}}(c+dx)} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{5/2}} dx$$

↓ 3478

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2}{3} \int \frac{Ab+3aB+(aA+3bB)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \int \frac{Ab+3aB+(aA+3bB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \int \frac{Ab+3aB+(aA+3bB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \left((3aB+Ab) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + (a-b)(A-3B) \int \frac{1}{\sqrt{\cos(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \left((a-b)(A-3B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + (3aB+Ab) \int \frac{1}{\sqrt{\cos(c+dx)}} dx \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \left((3aB+Ab) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 3473

3.592. $\int \sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{2(a-b)\sqrt{a+b}(3aB+Ab)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E(\arcsin}{a^2d}\right.\right.$$

input `Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*(a - b)*Sqrt[a + b]*(A*b + 3*A*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(A - 3*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/3 + (2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))`

3.592.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)]*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_)] + (f_)*(x_)]*(g_))^(p_)*((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

3.592. $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_
.)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

```
rule 3478 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*
x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a
+ b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*
(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*
Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]
```

3.592.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1974 vs. $2(290) = 580$.

Time = 18.41 (sec) , antiderivative size = 1975, normalized size of antiderivative = 6.10

method	result	size
parts	Expression too large to display	1975
default	Expression too large to display	2578

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

$$3.592. \quad \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

output

```

-2/3*A/d*sec(d*x+c)^(5/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(cos(d*x+c)
)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b
)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2+cos(d*
x+c)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)
)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b-cos
(d*x+c)^4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos
(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*a*b-
cos(d*x+c)^4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*b
^2+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a
+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*a^2*cos(d
*x+c)^3+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c
)))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*a*b*
cos(d*x+c)^3-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(
d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
*a*b*cos(d*x+c)^3-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1
+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(
1/2))*b^2*cos(d*x+c)^3+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b
)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)
)^(1/2))*a^2*cos(d*x+c)^2+EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)...

```

3.592.5 Fracas [F]

$$\begin{aligned}
 & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\
 &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{5}{2}} dx
 \end{aligned}$$

input

```

integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fricas")

```

output

```

integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2),
x)

```


3.592.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)*(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.592.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.592.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.592. $\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$

3.592.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} \sqrt{a + b \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(1/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(1/2), x)`

3.593 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

3.593.1 Optimal result	5396
3.593.2 Mathematica [A] (verified)	5397
3.593.3 Rubi [A] (verified)	5397
3.593.4 Maple [B] (warning: unable to verify)	5400
3.593.5 Fricas [F]	5401
3.593.6 Sympy [F(-1)]	5402
3.593.7 Maxima [F]	5402
3.593.8 Giac [F]	5402
3.593.9 Mupad [F(-1)]	5403

3.593.1 Optimal result

Integrand size = 35, antiderivative size = 411

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{2A(a - b)\sqrt{a + b}\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{ad\sqrt{\sec(c + dx)}} + \frac{2\sqrt{a + b}(Ab - a(A - B))\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}}}{ad\sqrt{\sec(c + dx)}} - \frac{2\sqrt{a + b}B\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{d\sqrt{\sec(c + dx)}}$$

```
output 2*A*(a-b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/sec(d*x+c)^(1/2)+2*(A*b-a*(A-B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/sec(d*x+c)^(1/2)-2*B*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d/sec(d*x+c)^(1/2)
```

3.593.2 Mathematica [A] (verified)

Time = 7.17 (sec) , antiderivative size = 358, normalized size of antiderivative = 0.87

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \frac{\sqrt{\sec(c + dx)} \left(2A(a + b \cos(c + dx)) \sin(c + dx) - \cos^2\left(\frac{1}{2}(c + dx)\right) \left(4A(a + b) \sqrt{\frac{\cos(c + dx)}{1 + \cos(c + dx)}} \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} \right) \right)}{2(a + b)}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `(Sqrt[Sec[c + d*x]]*(2*A*(a + b*Cos[c + d*x])*Sin[c + d*x] - Cos[(c + d*x)/2]^2*(4*A*(a + b)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 4*(b*(A - B) + a*(A + B))*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 8*b*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + A*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^3*(-Sin[(c + d*x)/2] + Sin[(3*(c + d*x))/2])))/(d*Sqrt[a + b*Cos[c + d*x]])`

3.593.3 Rubi [A] (verified)Time = 1.38 (sec) , antiderivative size = 372, normalized size of antiderivative = 0.91, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3470, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

3.593. $\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\cos^{\frac{3}{2}}(c+dx)} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx$$

↓ 3470

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{aA+(Ab+aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + bB \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{aA+(Ab+aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + bB \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{aA+(Ab+aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{\sin(c+dx+\frac{\pi}{2})} \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left((Ab-a(A-B)) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + aA \int \frac{\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left((Ab-a(A-B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + aA \int \frac{\cos(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(aA \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2\sqrt{a+b}(Ab-a(A-B))\cot(c+dx)}{\sin(c+dx+\frac{\pi}{2})} \right)$$

3.593. $\int \sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx) dx$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2\sqrt{a+b}(Ab-a(A-B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{ad} \text{EllipticF}(\arcsin(\dots)) \right)$$

```
input Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(A*b - a*(A - B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)
```

3.593.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)])/Sqrt[(c_) + (d_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)]^(m_.))*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^m Int[(a + b*SIN[e + f*x])^m*((c +
d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3470 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)]]/((b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2), x_Symbol] := Simp[B*(d
/b^2) Int[Sqrt[b*SIN[e + f*x]]/Sqrt[c + d*SIN[e + f*x]], x], x] + Int[(A*
c + (B*c + A*d)*SIN[e + f*x])/((b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f
x]]), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0]
```

```
rule 3473 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)]
^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*SIN[e + f*x]]/Sqrt[b*SIN[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_
.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*SIN[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + SIN[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

3.593.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 959 vs. $2(375) = 750$.

Time = 18.25 (sec) , antiderivative size = 960, normalized size of antiderivative = 2.34

method	result	size
parts	Expression too large to display	960
default	Expression too large to display	1126

3.593. $\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+csc(d*x+c)^3*(1-cos(d*x+c))^3*a-csc(d*x+c)^3*(1-cos(d*x+c))^3*b+a*(csc(d*x+c)-cot(d*x+c))+b*(csc(d*x+c)-cot(d*x+c)))*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1))^(3/2)/(csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)-2*B/d*sec(d*x+c)^(3/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/(a+cos(d*x+c)*b)^(1/2)*(EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a-EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b+2*b*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b))^(1/2)...`

3.593.5 Fracas [F]

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(1/2),x,algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2),x)`

3.593.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)*(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.593.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.593.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx \\ &= \int (B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2), x)`

3.593.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$= \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} \sqrt{a + b \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(1/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(1/2), x)`

3.594 $\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)}$

3.594.1 Optimal result	5404
3.594.2 Mathematica [A] (verified)	5405
3.594.3 Rubi [A] (verified)	5406
3.594.4 Maple [B] (verified)	5410
3.594.5 Fricas [F]	5411
3.594.6 Sympy [F]	5412
3.594.7 Maxima [F]	5412
3.594.8 Giac [F]	5412
3.594.9 Mupad [F(-1)]	5413

3.594.1 Optimal result

Integrand size = 35, antiderivative size = 445

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx =$$

$$\frac{(a - b)\sqrt{a + b}B\sqrt{\cos(c + dx)}\csc(c + dx)E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{ad\sqrt{\sec(c + dx)}} +$$

$$\frac{\sqrt{a + b}(2A + B)\sqrt{\cos(c + dx)}\csc(c + dx)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{d\sqrt{\sec(c + dx)}} +$$

$$\frac{\sqrt{a + b}(2Ab + aB)\sqrt{\cos(c + dx)}\csc(c + dx)\text{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{bd\sqrt{\sec(c + dx)}} +$$

$$\frac{B\sqrt{a + b \cos(c + dx)}\sqrt{\sec(c + dx)}\sin(c + dx)}{d}$$

```
output B*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)*sec(d*x+c)^(1/2)/d-(a-b)*B*csc(d*x+c)*
EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*
(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/
a/d/sec(d*x+c)^(1/2)+(2*A+B)*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*
(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d/
sec(d*x+c)^(1/2)-(2*A*b+B*a)*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),
(a+b)/b,((-a-b)/(a-b))^(1/2))*
(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b/d/
sec(d*x+c)^(1/2)
```

3.594.2 Mathematica [A] (verified)

Time = 16.63 (sec) , antiderivative size = 787, normalized size of antiderivative = 1.77

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \frac{-aB \tan\left(\frac{1}{2}(c + dx)\right) - bB \tan\left(\frac{1}{2}(c + dx)\right) + 2bB \tan^3\left(\frac{1}{2}(c + dx)\right) + aB \tan^5\left(\frac{1}{2}(c + dx)\right) - bB \tan^5\left(\frac{1}{2}(c + dx)\right)}{d}$$

input `Integrate[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `(-(a*B*Tan[(c + d*x)/2]) - b*B*Tan[(c + d*x)/2] + 2*b*B*Tan[(c + d*x)/2]^3 + a*B*Tan[(c + d*x)/2]^5 - b*B*Tan[(c + d*x)/2]^5 - 4*A*b*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 2*a*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 4*A*b*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 2*a*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - (a + b)*B*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*(1 + Tan[(c + d*x)/2]^2)*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + 2*(A*b + a*(-A + B))*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*(1 + Tan[(c + d*x)/2]^2)*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)))/(d*Sqrt[(1 + Tan[(c + d*x)/2]^2)/(1 - Tan[(c + d*x)/2]^2)]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(1 + Tan[(c + d*x)/2]^2)]*(-1 + Tan[(c + d*x)/2]^4))`

3.594.3 Rubi [A] (verified)

Time = 1.73 (sec) , antiderivative size = 412, normalized size of antiderivative = 0.93, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3482, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\sec(c+dx)} \sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\csc\left(c+dx+\frac{\pi}{2}\right)} \sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\cos(c+dx)} (A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} (A+B\sin\left(c+dx+\frac{\pi}{2}\right))}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3482} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{1}{2} \int -\frac{((2Ab+aB)\cos^2(c+dx)) - 2aA\cos(c+dx) + aB}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + \frac{B\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \right) \\
 & \quad \downarrow \text{25} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{B\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{1}{2} \int \frac{((2Ab+aB)\cos^2(c+dx)) - 2aA\cos(c+dx) + aB}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{B\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{1}{2} \int \frac{(-2Ab-aB)\sin\left(c+dx+\frac{\pi}{2}\right)^2 - 2aA\sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} dx \right) \\
 & \quad \downarrow \text{3532}
 \end{aligned}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left((aB+2Ab)\int\frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}}dx-\int\frac{aB-2aA\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left((aB+2Ab)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\int\frac{aB-2aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left(-\int\frac{aB-2aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2\sqrt{a+b}(aB+2Ab)\cot(c+dx)}{\sin(c+dx+\frac{\pi}{2})}\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left(a(2A+B)\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}dx-aB\int\frac{\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left(a(2A+B)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-aB\int\frac{\cos(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left(-aB\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2\sqrt{a+b}(2A+B)\cot(c+dx)}{\sin(c+dx+\frac{\pi}{2})}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{2}\left(\frac{2\sqrt{a+b}(2A+B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}\right)\right)}{d}\right)\right)$$

input `Int[Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((( -2*(a - b)*Sqrt[a + b]*B*Cot[c +
d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d
*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(
1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(2*A + B)*Cot[c + d*x]
*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]
)], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 +
Sec[c + d*x]))/(a - b))]/d - (2*Sqrt[a + b]*(2*A*b + a*B)*Cot[c + d*x]*Ell
ipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c
+ d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[
(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/2 + (B*Sqrt[a + b*Cos[c + d*x]]*Si
n[c + d*x])/(d*Sqrt[Cos[c + d*x]))
```

3.594.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3288 Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)])/Sqrt[(c_) + (d_)*sin[(e_)] + (f_)*
(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_)] + (f
_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m*Int[(a + b*Sin[e + f*x])^n*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3482 `Int[Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[-2*B*Cos[e + f*x]*Sqrt[a + b*Sin[e + f*x]]*((c + d*Sin[e + f*x])^n/(f*(2*n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*Sin[e + f*x])^(n - 1)/Sqrt[a + b*Sin[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*Sin[e + f*x] + (A*b*d*(2*n + 3) + B*(a*d + 2*b*c*n))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && EqQ[n^2, 1/4]`


```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)])), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.594.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1267 vs. 2(405) = 810.

Time = 15.05 (sec) , antiderivative size = 1268, normalized size of antiderivative = 2.85

method	result	size
parts	Expression too large to display	1268
default	Expression too large to display	2084

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)*(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 2*A/d/(a*cos(d*x+c)*b)^(1/2)*sec(d*x+c)^(1/2)*(-EllipticF(cot(d*x+c)-csc(d
*x+c),(-a-b)/(a+b))^(1/2))*a+EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b
))^(1/2))*b-2*b*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))
*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))
^(1/2)*(1+cos(d*x+c))+B/d*(-EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)
)^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+c)*b)/(1+cos(d*x+c)
)/(a+b))^(1/2)*a*cos(d*x+c)^2-EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)
)^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+c)*b)/(1+cos(d*x+c)
)/(a+b))^(1/2)*b*cos(d*x+c)^2-2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b
)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+c)*b)/(1+cos
(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-
a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+c)*b)/(1+
cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)^2-2*EllipticE(cot(d*x+c)-csc(d*x+c),
(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+c)*b)/
(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)-2*EllipticE(cot(d*x+c)-csc(d*x+c)
,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+c)*b)
/(1+cos(d*x+c))/(a+b))^(1/2)*b*cos(d*x+c)-4*EllipticPi(cot(d*x+c)-csc(d*x+
c),-1,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(d*x+
c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)+4*EllipticF(cot(d*x+c)-csc(
d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a*cos(...
```

3.594.5 Fracas [F]

$$\int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$= \int (B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}\sqrt{\sec(dx + c)} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fracas")
```

```
output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c)),
x)
```

3.594.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx \\ &= \int (A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}\sqrt{\sec(c + dx)} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)*(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))*sqrt(sec(c + d*x)), x)`

3.594.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}\sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.594.8 Giac [F]

$$\begin{aligned} & \int \sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx \\ &= \int (B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}\sqrt{\sec(dx + c)} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)*(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.594.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$$

$$= \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} \sqrt{a + b \cos(c + dx)} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(1/2), x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(1/2), x)`

$$3.595 \quad \int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.595.1 Optimal result	5414
3.595.2 Mathematica [B] (verified)	5415
3.595.3 Rubi [A] (verified)	5416
3.595.4 Maple [B] (verified)	5421
3.595.5 Fracas [F]	5422
3.595.6 Sympy [F]	5423
3.595.7 Maxima [F]	5423
3.595.8 Giac [F]	5423
3.595.9 Mupad [F(-1)]	5424

3.595.1 Optimal result

Integrand size = 35, antiderivative size = 533

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}(4Ab+aB)\sqrt{\cos(c+dx)}\csc(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\middle|-\frac{a+b}{a-b}\right)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4abd\sqrt{\sec(c+dx)}} -$$

$$+ \frac{\sqrt{a+b}(4Ab+(a+2b)B)\sqrt{\cos(c+dx)}\csc(c+dx)\operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right),-\frac{a+b}{a-b}\right)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4bd\sqrt{\sec(c+dx)}} -$$

$$\frac{\sqrt{a+b}(4aAb-a^2B+4b^2B)\sqrt{\cos(c+dx)}\csc(c+dx)\operatorname{EllipticPi}\left(\frac{a+b}{b},\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right),-\frac{a+b}{a-b}\right)}{4b^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{B\sqrt{a+b \cos(c+dx)}\sin(c+dx)}{2d\sqrt{\sec(c+dx)}} +$$

$$\frac{(4Ab+aB)\sqrt{a+b \cos(c+dx)}\sqrt{\sec(c+dx)}\sin(c+dx)}{4bd}$$

output $\frac{1}{2}B\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\sec(dx+c)^{1/2}+1/4*(4A*b+B*a)*\sin(dx+c)*(a+b\cos(dx+c))^{1/2}*\sec(dx+c)^{1/2}/b/d-1/4*(a-b)*(4A*b+B*a)*\csc(dx+c)*\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c)))/(a+b)^{1/2}*(a*(1+\sec(dx+c)))/(a-b)^{1/2}/a/b/d\sec(dx+c)^{1/2}+1/4*(4A*b+(a+2*b)*B)*\csc(dx+c)*\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c)))/(a+b)^{1/2}*(a*(1+\sec(dx+c)))/(a-b)^{1/2}/b/d\sec(dx+c)^{1/2}-1/4*(4A*a*b-B*a^2+4*B*b^2)*\csc(dx+c)*\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b,((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c)))/(a+b)^{1/2}*(a*(1+\sec(dx+c)))/(a-b)^{1/2}/b^2/d\sec(dx+c)^{1/2}$

3.595.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1121 vs. $2(533) = 1066$.

Time = 16.84 (sec) , antiderivative size = 1121, normalized size of antiderivative = 2.10

$$\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx = \text{Too large to display}$$

input `Integrate[(Sqrt[a + bCos[c + d*x]]*(A + B*cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output $(B\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}\sin[2(c + dx)]/(4d) + (\sqrt{(1 - \tan[(c + dx)/2]^2})^{-1}*(4*a*A*b*\tan[(c + dx)/2] + 4*A*b^2*\tan[(c + dx)/2] + a^2*B*\tan[(c + dx)/2] + a*b*B*\tan[(c + dx)/2] - 8*A*b^2*\tan[(c + dx)/2]^3 - 2*a*b*B*\tan[(c + dx)/2]^3 - 4*a*A*b*\tan[(c + dx)/2]^5 + 4*A*b^2*\tan[(c + dx)/2]^5 - a^2*B*\tan[(c + dx)/2]^5 + a*b*B*\tan[(c + dx)/2]^5 + 8*a*A*b*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + dx)/2]^2}*\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} - 2*a^2*B*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + dx)/2]^2}*\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + 8*b^2*B*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + dx)/2]^2}*\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + 8*a*A*b*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]*\tan[(c + dx)/2]^2*\sqrt{1 - \tan[(c + dx)/2]^2}*\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} - 2*a^2*B*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]*\tan[(c + dx)/2]^2*\sqrt{1 - \tan[(c + dx)/2]^2}*\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + 8*b^2*B*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]*\tan[(c + dx)/2]^2*\sqrt{1 - \tan[(c + dx)/2]^2}*\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + (a + b)*(4*A*b + a*B)*\text{EllipticE}[\text{ArcSin}[\dots$

3.595.3 Rubi [A] (verified)

Time = 2.28 (sec) , antiderivative size = 492, normalized size of antiderivative = 0.92, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3482, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

↓ 3042

$$\int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)}\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx)) dx$$

3.595. $\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3482

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\int\frac{(4Ab+aB)\cos^2(c+dx)+2(2aA+bB)\cos(c+dx)+aB}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}dx+\frac{B\sin(c+dx)}{\sqrt{\cos(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\int\frac{(4Ab+aB)\sin\left(c+dx+\frac{\pi}{2}\right)^2+2(2aA+bB)\sin\left(c+dx+\frac{\pi}{2}\right)+aB}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}}dx+\frac{B\sin\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}\right)$$

↓ 3540

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{\int\frac{-((-Ba^2+4Aba+4b^2B)\cos^2(c+dx))-2abB\cos(c+dx)+a(4Ab+aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{2b}+\frac{(aB+4Ab)\sin(c+dx)}{bd}\right)\right)$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}-\frac{\int\frac{-((-Ba^2+4Aba+4b^2B)\cos^2(c+dx))-2abB\cos(c+dx)+a(4Ab+aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{2b}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}-\frac{\int\frac{(Ba^2-4Aba-4b^2B)\sin\left(c+dx+\frac{\pi}{2}\right)^2-2abB\sin\left(c+dx+\frac{\pi}{2}\right)+a(4Ab+aB)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{3/2}\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}}dx}{2b}\right)\right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}-\frac{\int\frac{a(4Ab+aB)-2abB\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-(a^2)}{2b}\right)\right)$$

↓ 3042

3.595. $\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{a(4Ab+aB)-2abB\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} \right) \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{a(4Ab+aB)-2abB\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} \right) \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(aB+4Ab) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{-a(B(a+2b)+4Ab) \int \frac{1}{\sqrt{\sin(c+dx)}} dx}{\sqrt{\sin(c+dx)}} \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(aB+4Ab) \int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{2\sqrt{a+b}(a^2(-B)+4aAb+4b^2B) \cot(c+dx)}{\sqrt{\sec(c+dx)}} \right) \right)$$

3.595. $\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*Sqrt[a + b]*(4*A*b + a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*Sqrt[a + b]*(4*A*b + (a + 2*b)*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d + (2*Sqrt[a + b]*(4*a*A*b - a^2*B + 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/b + ((4*A*b + a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/4)`

3.595.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

3.595.
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^m*Int[(a + b*SIN[e + f*x])^n*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*SIN[e + f*x]]/Sqrt[b*SIN[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*SIN[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + SIN[e + f*x])/((a + b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3482 `Int[Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[-2*B*COS[e + f*x]*Sqrt[a + b*SIN[e + f*x]]*((c + d*SIN[e + f*x])^n/(f*(2*n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*SIN[e + f*x])^(n - 1)/Sqrt[a + b*SIN[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*SIN[e + f*x] + (A*b*d*(2*n + 3) + B*(a*d + 2*b*c*n))*SIN[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && EqQ[n^2, 1/4]`

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)])), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.595.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2727 vs. 2(479) = 958.

Time = 14.56 (sec) , antiderivative size = 2728, normalized size of antiderivative = 5.12

method	result	size
parts	Expression too large to display	2728
default	Expression too large to display	2750

```
input int((A+B*cos(d*x+c))*(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output A/d/(1+cos(d*x+c))/sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(1/2)*(2*EllipticF(cot
(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2
)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)-EllipticE(cot
(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)-EllipticE(cot(
d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*
((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*b*cos(d*x+c)-2*EllipticPi(cot
(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*cos(d*x+c)+4*(cos(d*x
+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(
1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a-2*(cos(d*x+c)/(1+cos
(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+
cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a-2*(cos(d*x+c)/(1+cos(d*x+c)))^(
1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)
*b)/(1+cos(d*x+c))/(a+b))^(1/2)*b-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*Elli
pticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1
+cos(d*x+c))/(a+b))^(1/2)*a+2*sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(
1+cos(d*x+c))/(a+b))^(1/2)*a-sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*
EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)...
```

3.595.5 Fracas [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\sqrt{\sec(dx + c)}} dx$$

```
input integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algo
rithm="fracas")
```

```
output integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sqrt(sec(d*x + c)),
x)
```

3.595.6 Sympy [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))**(1/2)/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))/sqrt(sec(c + d*x)), x)`

3.595.7 Maxima [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.595.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.595.9 Mupad **[F(-1)]**

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2), x)`

3.596
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.596.1 Optimal result	5425
3.596.2 Mathematica [B] (verified)	5426
3.596.3 Rubi [A] (verified)	5427
3.596.4 Maple [B] (verified)	5435
3.596.5 Fracas [F(-1)]	5436
3.596.6 Sympy [F]	5436
3.596.7 Maxima [F]	5436
3.596.8 Giac [F]	5437
3.596.9 Mupad [F(-1)]	5437

3.596.1 Optimal result

Integrand size = 35, antiderivative size = 620

$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx =$$

$$\frac{(a-b)\sqrt{a+b}(6Ab-3a^2B+16b^2B)\sqrt{\cos(c+dx)}\csc(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\middle|-\frac{a+b}{a-b}\right)\sqrt{\sec(c+dx)}}{24ab^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(a+2b)(6Ab-3aB+8bB)\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right),-\frac{a+b}{a-b}\right)\sqrt{\sec(c+dx)}}{24b^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(2a^2Ab-8Ab^3-a^3B-4ab^2B)\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticPi}\left(\frac{a+b}{b},\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)\sqrt{\sec(c+dx)}}{8b^3d\sqrt{\sec(c+dx)}} +$$

$$\frac{(2Ab-aB)\sqrt{a+b \cos(c+dx)}\sin(c+dx)}{4bd\sqrt{\sec(c+dx)}} + \frac{B(a+b \cos(c+dx))^{3/2}\sin(c+dx)}{3bd\sqrt{\sec(c+dx)}} +$$

$$\frac{(6Ab-3a^2B+16b^2B)\sqrt{a+b \cos(c+dx)}\sqrt{\sec(c+dx)}\sin(c+dx)}{24b^2d}$$

3.596.
$$\int \frac{\sqrt{a+b \cos(c+dx)}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

output $\frac{1}{3}B(a+b\cos(dx+c))^{3/2}\sin(dx+c)/b/d/\sec(dx+c)^{1/2} + \frac{1}{4}(2A*b - B*a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b/d/\sec(dx+c)^{1/2} + \frac{1}{24}(6A*a*b - 3B*a^2 + 16B*b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/b^2/d - \frac{1}{24}(a-b)(6A*a*b - 3B*a^2 + 16B*b^2)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2}/a/b^2/d/\sec(dx+c)^{1/2} + \frac{1}{24}(a+2b)*(6A*b - 3B*a + 8B*b)\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2}/b^2/d/\sec(dx+c)^{1/2} + \frac{1}{8}(2A*a^2*b - 8A*b^3 - B*a^3 - 4B*a*b^2)\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, (a+b)/b, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2}/b^3/d/\sec(dx+c)^{1/2}$

3.596.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1533 vs. $2(620) = 1240$.

Time = 13.94 (sec) , antiderivative size = 1533, normalized size of antiderivative = 2.47

$$\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx = \text{Too large to display}$$

input `Integrate[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2), x]`

output $(\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} ((B \sin(c + dx))/12 + ((6A b + aB) \sin[2(c + dx)])/(24b) + (B \sin[3(c + dx)])/12))/d + (\sqrt{(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2)/(1 + \tan[(c + dx)/2]^2)} (6a^2 A b \tan[(c + dx)/2] + 6a A b^2 \tan[(c + dx)/2] - 3a^3 B \tan[(c + dx)/2] - 3a^2 b B \tan[(c + dx)/2] + 16a b^2 B \tan[(c + dx)/2] + 16b^3 B \tan[(c + dx)/2] - 12a A b^2 \tan[(c + dx)/2]^3 + 6a^2 b B \tan[(c + dx)/2]^3 - 32b^3 B \tan[(c + dx)/2]^3 - 6a^2 A b \tan[(c + dx)/2]^5 + 6a A b^2 \tan[(c + dx)/2]^5 + 3a^3 B \tan[(c + dx)/2]^5 - 3a^2 b B \tan[(c + dx)/2]^5 - 16a b^2 B \tan[(c + dx)/2]^5 + 16b^3 B \tan[(c + dx)/2]^5 - 12a^2 A b \operatorname{EllipticPi}[-1, \operatorname{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \sqrt{1 - \tan[(c + dx)/2]^2} \sqrt{(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2)/(a + b)} + 48A b^3 \operatorname{EllipticPi}[-1, \operatorname{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \sqrt{1 - \tan[(c + dx)/2]^2} \sqrt{(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2)/(a + b)} + 6a^3 B \operatorname{EllipticPi}[-1, \operatorname{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \sqrt{1 - \tan[(c + dx)/2]^2} \sqrt{(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2)/(a + b)} + 24a b^2 B \operatorname{EllipticPi}[-1, \operatorname{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \sqrt{1 - \tan[(c + dx)/2]^2} \sqrt{(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2)/(a + b)} - 12a^2 A b \operatorname{EllipticPi}[-1, \operatorname{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \tan[(c + dx)/2]^2 \sqrt{1 - \tan[(c + dx)/2]^2} \sqrt{(a + b \dots$

3.596.3 Rubi [A] (verified)

Time = 2.91 (sec) , antiderivative size = 584, normalized size of antiderivative = 0.94, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx)) dx$$

3.596. $\int \frac{\sqrt{a + b \cos(c + dx)} (A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3469

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{a+b\cos(c+dx)}(3(2Ab-aB)\cos^2(c+dx)+4bB\cos(c+dx)+aB)}{2\sqrt{\cos(c+dx)}} dx}{3b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{3b} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{a+b\cos(c+dx)}(3(2Ab-aB)\cos^2(c+dx)+4bB\cos(c+dx)+aB)}{\sqrt{\cos(c+dx)}} dx}{6b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{3b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})} \left(3(2Ab-aB)\sin(c+dx+\frac{\pi}{2})^2+4bB\sin(c+dx+\frac{\pi}{2})+aB\right)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{6b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{3b} \right)$$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{2} \int \frac{(-3Ba^2+6Aba+16b^2B)\cos^2(c+dx)+2b(6Ab+7aB)\cos(c+dx)+a(6Ab+aB)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{6b} + \frac{3(2Ab-aB)\sin(c+dx)}{3b} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4} \int \frac{(-3Ba^2+6Aba+16b^2B)\cos^2(c+dx)+2b(6Ab+7aB)\cos(c+dx)+a(6Ab+aB)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{6b} + \frac{3(2Ab-aB)\sin(c+dx)}{3b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4} \int \frac{(-3Ba^2+6Aba+16b^2B)\sin(c+dx+\frac{\pi}{2})^2+2b(6Ab+7aB)\sin(c+dx+\frac{\pi}{2})+a(6Ab+aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{6b} + \frac{3(2Ab-aB)\sin(c+dx)}{3b} \right)$$

3.596. $\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

↓ 3540

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{\int -\frac{3(-Ba^3+2Aba^2-4b^2Ba-8Ab^3)\cos^2(c+dx)-2ab(6Ab+aB)\cos(c+dx)+a(-3Ba^2+6Aba+16b^2B)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} + \dots \right) \right)$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{3(-Ba^3+2Aba^2-4b^2Ba-8Ab^3)\cos^2(c+dx)-2a}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{3(-Ba^3+2Aba^2-4b^2Ba-8Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})} dx}{2} \right) \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{a(-3Ba^2+6Aba+16b^2B)-2ab(6Ab+aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2} \right) \right)$$

↓ 3042

3.596. $\int \frac{\sqrt{a+b\cos(c+dx)}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{a(-3Ba^2+6Aba+16b^2B)-2ab(6Ab+aB)\sin(c+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{a(-3Ba^2+6Aba+16b^2B)-2ab(6Ab+aB)\sin(c+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(-3a^2B+6aAb+16b^2B)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \int \frac{\cos(c+dx)+\sin(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(-3a^2B+6aAb+16b^2B)\int \frac{\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}} \sqrt{a+b\cos(c+dx)}}{3/2} \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(-3a^2B+6aAb+16b^2B)\int \frac{\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}} \sqrt{a+b\cos(c+dx)}}{3/2} \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(-3a^2B+6aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{2(a-b)\sqrt{a+b}(-3a^2B+6aAb+16b^2B)\cot(c+dx)\sqrt{a}}{3/2} \right) \right)$$

input `Int[(Sqrt[a + b*Cos[c + d*x]]*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

```

output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Sqrt[Cos[c + d*x]]*(a + b*Cos[c
+ d*x])^(3/2)*Sin[c + d*x])/(3*b*d) + ((3*(2*A*b - a*B)*Sqrt[Cos[c + d*x]]
*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*Sqrt[a +
b]*(6*a*A*b - 3*a^2*B + 16*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a +
b*Cos[c + d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqr
t[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a
*d) - (2*Sqrt[a + b]*(a + 2*b)*(6*A*b - 3*a*B + 8*b*B)*Cot[c + d*x]*Ellipt
icF[ArcSin[Sqrt[a + b*Cos[c + d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((
a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c +
d*x]))/(a - b)]/d - (6*Sqrt[a + b]*(2*a^2*A*b - 8*A*b^3 - a^3*B - 4*a*b^
2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]])/(S
qrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c +
d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b*d))/b + ((6*a*A*b
- 3*a^2*B + 16*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Co
s[c + d*x]))/4)/(6*b))

```

3.596.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3288 Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)]/Sqrt[(c_) + (d_)*sin[(e_)] + (f_
)*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]

```

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.596.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3921 vs. $2(560) = 1120$.

Time = 17.61 (sec) , antiderivative size = 3922, normalized size of antiderivative = 6.33

method	result	size
parts	Expression too large to display	3922
default	Expression too large to display	3963

```
input int((A+B*cos(d*x+c))*(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*A/d/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(3/2)*(2*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b-4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2+((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2+((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b-2*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2+8*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2+4*sec(d*x+c)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b-8*sec(d*x+c)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2+2*sec(d*x+c)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2+2*sec(d*x+c)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(c...
```

3.596.5 Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="fricas")`

output Timed out

3.596.6 Sympy [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\sec^{\frac{3}{2}}(c + dx)} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))**(1/2)/sec(d*x+c)**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(a + b*cos(c + d*x))/sec(c + d*x)**(3/2), x)`

3.596.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \cos(dx + c) + A) \sqrt{b \cos(dx + c) + a}}{\sec(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.596.8 Giac [F]

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \cos(dx + c) + A)\sqrt{b \cos(dx + c) + a}}{\sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/sec(d*x + c)^(3/2), x)`

3.596.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \cos(c + dx)}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \cos(c + dx)) \sqrt{a + b \cos(c + dx)}}{\left(\frac{1}{\cos(c + dx)}\right)^{\frac{3}{2}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(3/2), x)`

3.597 $\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$

3.597.1 Optimal result	5438
3.597.2 Mathematica [A] (warning: unable to verify)	5439
3.597.3 Rubi [A] (verified)	5440
3.597.4 Maple [B] (warning: unable to verify)	5446
3.597.5 Fricas [F]	5447
3.597.6 Sympy [F(-1)]	5447
3.597.7 Maxima [F]	5447
3.597.8 Giac [F]	5448
3.597.9 Mupad [F(-1)]	5448

3.597.1 Optimal result

Integrand size = 35, antiderivative size = 562

$$\int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx =$$

$$\frac{2(a - b)\sqrt{a + b}(147a^4A + 33a^2Ab^2 + 8Ab^4 + 246a^3bB - 18ab^3B) \sqrt{\cos(c + dx)} \csc(c + dx) E(\arcsin(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}}))}{315a^4d\sqrt{\sec(c + dx)}} +$$

$$\frac{2(a - b)\sqrt{a + b}(8Ab^3 - a^3(147A - 75B) + 3a^2b(13A - 57B) + 6ab^2(A - 3B)) \sqrt{\cos(c + dx)} \csc(c + dx)}{315a^3d\sqrt{\sec(c + dx)}} +$$

$$\frac{2(88a^2Ab - 4Ab^3 + 75a^3B + 9ab^2B) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{315a^2d} +$$

$$\frac{2(49a^2A + 3Ab^2 + 72abB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{315ad} +$$

$$\frac{2(10Ab + 9aB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{63d} +$$

$$\frac{2aA \sqrt{a + b \cos(c + dx)} \sec^{\frac{9}{2}}(c + dx) \sin(c + dx)}{9d}$$

output $2/315*(88*A*a^2*b-4*A*b^3+75*B*a^3+9*B*a*b^2)*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/a^2/d+2/315*(49*A*a^2+3*A*b^2+72*B*a*b)*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/a/d+2/63*(10*A*b+9*B*a)*\sec(d*x+c)^{(7/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/d+2/9*a*A*\sec(d*x+c)^{(9/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/d+2/315*(a-b)*(147*A*a^4+33*A*a^2*b^2+8*A*b^4+246*B*a^3*b-18*B*a*b^3)*\csc(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b))^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b))^{(1/2)}/a^4/d/\sec(d*x+c)^{(1/2)}+2/315*(a-b)*(8*A*b^3-a^3*(147*A-75*B)+3*a^2*b*(13*A-57*B)+6*a*b^2*(A-3*B))*\csc(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b))^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b))^{(1/2)}/a^3/d/\sec(d*x+c)^{(1/2)}$

3.597.2 Mathematica [A] (warning: unable to verify)

Time = 18.97 (sec) , antiderivative size = 625, normalized size of antiderivative = 1.11

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \frac{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)} \sec(c + dx) \left(-2(a + b)(147a^4A + 33a^2Ab^2 + 8Ab^4 + 246a^3bB - 18ab^3B)\right) \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2(147a^4A + 33a^2Ab^2 + 8Ab^4 + 246a^3bB - 18ab^3B) \sin(c + dx)}{315a^3} + \frac{2}{63} \sec^3(c + dx)(10Ab \sin(c + dx) - 10aB)\right)}{315a^3}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output $(2\sqrt{\cos[(c + dx)/2]}^2 \sec[c + dx]) \cdot (-2(a + b) \cdot (147a^4A + 33a^2Ab^2 + 8A^2b^4 + 246a^3bB - 18ab^3B)) \sqrt{\cos[c + dx]/(1 + \cos[c + dx])} \cdot \sqrt{(a + b\cos[c + dx])/((a + b)(1 + \cos[c + dx]))} \cdot \text{EllipticE}[\text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] + 2a(a + b)(8A^2b^3 - 6a^2b^2(A + 3B) + 3a^3(49A + 25B) + 3a^2b(13A + 57B)) \sqrt{\cos[c + dx]/(1 + \cos[c + dx])} \cdot \sqrt{(a + b\cos[c + dx])/((a + b)(1 + \cos[c + dx]))} \cdot \text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] - (147a^4A + 33a^2Ab^2 + 8A^2b^4 + 246a^3bB - 18ab^3B) \cos[c + dx] \cdot (a + b\cos[c + dx]) \cdot \sec[(c + dx)/2]^2 \cdot \text{Tan}[(c + dx)/2]) / (315a^3d \sqrt{a + b\cos[c + dx]} \cdot \sqrt{\sec[(c + dx)/2]^2} + (\sqrt{a + b\cos[c + dx]} \cdot \sqrt{\sec[c + dx]}) \cdot ((2(147a^4A + 33a^2Ab^2 + 8A^2b^4 + 246a^3bB - 18ab^3B) \sin[c + dx]) / (315a^3) + (2\sec[c + dx]^3(10Ab \sin[c + dx] + 9a^2B \sin[c + dx])) / 63 + (2\sec[c + dx]^2(49a^2A \sin[c + dx] + 3A^2b^2 \sin[c + dx] + 72abB \sin[c + dx])) / (315a) + (2\sec[c + dx](88a^2Ab \sin[c + dx] - 4A^2b^3 \sin[c + dx] + 75a^3B \sin[c + dx] + 9a^2b^2B \sin[c + dx])) / (315a^2) + (2aA \sec[c + dx]^3 \text{Tan}[c + dx]) / 9) / d$

3.597.3 Rubi [A] (verified)

Time = 2.86 (sec) , antiderivative size = 554, normalized size of antiderivative = 0.99, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{11}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{11/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{11/2}} dx$$

3.597. $\int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx$

↓ 3468

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{9}\int\frac{3b(2aA+3bB)\cos^2(c+dx)+(7Aa^2+18bBa+9Ab^2)\cos(c+dx)+a(10Ab+9A^2)}{2\cos^{\frac{9}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\int\frac{3b(2aA+3bB)\cos^2(c+dx)+(7Aa^2+18bBa+9Ab^2)\cos(c+dx)+a(10Ab+9A^2)}{\cos^{\frac{9}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\int\frac{3b(2aA+3bB)\sin(c+dx+\frac{\pi}{2})^2+(7Aa^2+18bBa+9Ab^2)\sin(c+dx+\frac{\pi}{2})+a(10Ab+9A^2)}{\sin(c+dx+\frac{\pi}{2})^{9/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{2\int\frac{4ab(10Ab+9aB)\cos^2(c+dx)+a(45Ba^2+92Aba+63b^2B)\cos(c+dx)+a(49Aa^2+72bBa+3Ab^2)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{7a}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{\int\frac{4ab(10Ab+9aB)\cos^2(c+dx)+a(45Ba^2+92Aba+63b^2B)\cos(c+dx)+a(49Aa^2+72bBa+3Ab^2)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{7a}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{\int\frac{4ab(10Ab+9aB)\sin(c+dx+\frac{\pi}{2})^2+a(45Ba^2+92Aba+63b^2B)\sin(c+dx+\frac{\pi}{2})+a(49Aa^2+72bBa+3Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{7a}\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{2\int\frac{(147Aa^2+396bBa+209Ab^2)\cos(c+dx)a^2+2b(49Aa^2+72bBa+3Ab^2)\cos^2(c+dx)a+3(75Ba^3+88Aba^2+9b^2Ba^2)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{5a}\right)\right)$$

3.597. $\int(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{\frac{11}{2}}(c+dx)dx$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{\int \frac{(147Aa^2+396bBa+209Ab^2)\cos(c+dx)a^2+2b(49Aa^2+72bBa+3Ab^2)\cos^2(c+dx)a+3(75Ba^3+88Aba^2+9b^2Ba-}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{5a} \right) \right) \quad 7a$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{\int \frac{(147Aa^2+396bBa+209Ab^2)\sin(c+dx+\frac{\pi}{2})a^2+2b(49Aa^2+72bBa+3Ab^2)\sin(c+dx+\frac{\pi}{2})^2a+3(75Ba^3+88Aba^2-}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{5a} \right) \right) \quad 7a$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{2 \int \frac{3((75Ba^3+186Aba^2+153b^2Ba+2Ab^3)\cos(c+dx)a^2+(147Aa^4+246bBa^3+33Ab^2a^2-18b^3Ba+8Ab^4)a) dx}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{3a} \right) \right) \quad 5a$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{\int \frac{(75Ba^3+186Aba^2+153b^2Ba+2Ab^3)\cos(c+dx)a^2+(147Aa^4+246bBa^3+33Ab^2a^2-18b^3Ba+8Ab^4)a}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a} + \frac{2(75a}{5a} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\int \frac{(75Ba^3+186Aba^2+153b^2Ba+2Ab^3)\sin(c+dx+\frac{\pi}{2})a^2+(147Aa^4+246bBa^3+33Ab^2a^2-18b^3Ba+8Ab^4)a}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}\sqrt{-3a^2}}{a} \right) \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\int \frac{a(a-b)(-3a^3(49A-25B)+a^2(39Ab-171bB)+6ab^2(A-3B)+8Ab^3)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + a(147a^4A+246a^3bB+33a^2Ab^2-18ab^3Ba+8Ab^4) \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\int \frac{a(a-b)(-3a^3(49A-25B)+a^2(39Ab-171bB)+6ab^2(A-3B)+8Ab^3)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + a(147a^4A+246a^3bB+33a^2Ab^2-18ab^3Ba+8Ab^4) \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\int \frac{a(147a^4A+246a^3bB+33a^2Ab^2-18ab^3Ba+8Ab^4)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}\sqrt{-3a^2}}{a} \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{2(49a^2A+72abB+3Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5d\cos^{\frac{5}{2}}(c+dx)} + \frac{2(75a^3B+88a^2Ab+9ab^2B-4Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\cos^{\frac{3}{2}}(c+dx)} \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((2*(10*A*b + 9*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(49*a^2*A + 3*A*b^2 + 72*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2))) + (((2*(a - b)*Sqrt[a + b]*(147*a^4*A + 33*a^2*A*b^2 + 8*A*b^4 + 246*a^3*b*B - 18*a*b^3*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*(a - b)*Sqrt[a + b]*(8*A*b^3 - 3*a^3*(49*A - 25*B) + 6*a*b^2*(A - 3*B) + a^2*(39*A*b - 171*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/a + (2*(88*a^2*A*b - 4*A*b^3 + 75*a^3*B + 9*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2)))/(5*a))/(7*a))/9`

3.597.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_.)]]*Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_.)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_.)])/(((b_.)*sin[(e_.) + (f_.)*(x_.)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

```
rule 3477 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]
```

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))
```

3.597.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 5976 vs. $2(510) = 1020$.

Time = 32.82 (sec) , antiderivative size = 5977, normalized size of antiderivative = 10.64

method	result	size
parts	Expression too large to display	5977
default	Expression too large to display	6046

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x,method=_RE
TURNVERBOSE)
```

```
output result too large to display
```

3.597.5 Fricas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{11/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(11/2), x)`

3.597.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(11/2),x)`

output `Timed out`

3.597.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{11/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(11/2), x)`

3.597.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{11/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(11/2), x)`

3.597.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{11/2} (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + b*cos(c + d*x))^(3/2), x)`

3.598 $\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sec^{\frac{9}{2}}(c+dx) dx$

3.598.1 Optimal result	5449
3.598.2 Mathematica [A] (warning: unable to verify)	5450
3.598.3 Rubi [A] (verified)	5451
3.598.4 Maple [B] (verified)	5456
3.598.5 Fricas [F]	5457
3.598.6 Sympy [F(-1)]	5458
3.598.7 Maxima [F]	5458
3.598.8 Giac [F]	5458
3.598.9 Mupad [F(-1)]	5459

3.598.1 Optimal result

Integrand size = 35, antiderivative size = 473

$$\int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \frac{2(a - b)\sqrt{a + b}(82a^2Ab - 6Ab^3 + 63a^3B + 21ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{105a^3d\sqrt{\sec(c + dx)}} - \frac{2(a - b)\sqrt{a + b}(6Ab^2 - a^2(25A - 63B) + 3ab(19A - 7B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{105a^2d\sqrt{\sec(c + dx)}} + \frac{2(25a^2A + 3Ab^2 + 42abB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{105ad} + \frac{2(8Ab + 7aB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{35d} + \frac{2aA \sqrt{a + b \cos(c + dx)} \sec^{\frac{7}{2}}(c + dx) \sin(c + dx)}{7d}$$

output $2/105*(25*A*a^2+3*A*b^2+42*B*a*b)*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/a/d+2/35*(8*A*b+7*B*a)*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/d+2/7*a*A*\sec(d*x+c)^{(7/2)}*\sin(d*x+c)*(a+b*\cos(d*x+c))^{(1/2)}/d+2/105*(a-b)*(82*A*a^2*b-6*A*b^3+63*B*a^3+21*B*a*b^2)*\csc(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b)^{(1/2)}/a^3/d/\sec(d*x+c)^{(1/2)}-2/105*(a-b)*(6*A*b^2-a^2*(25*A-63*B)+3*a*b*(19*A-7*B))*\csc(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b)^{(1/2)}/a^2/d/\sec(d*x+c)^{(1/2)}$

3.598.2 Mathematica [A] (warning: unable to verify)

Time = 16.28 (sec) , antiderivative size = 518, normalized size of antiderivative = 1.10

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \frac{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right) \sec(c + dx)} \left(-2(a + b) (82a^2 Ab - 6Ab^3 + 63a^3 B + 21ab^2 B) \sqrt{\frac{\cos(c + dx)}{1 + \cos(c + dx)}} \sqrt{\frac{\cos(c + dx)}{1 - \cos(c + dx)}}\right)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} \left(-\frac{2(-82a^2 Ab + 6Ab^3 - 63a^3 B - 21ab^2 B) \sin(c + dx)}{105a^2} + \frac{2}{35} \sec^2(c + dx) (8Ab \sin(c + dx) + \dots)\right)$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output $(2\sqrt{\cos[(c + dx)/2]^2 \sec[c + dx]} * (-2*(a + b)*(82*a^2*A*b - 6*A*b^3 + 63*a^3*B + 21*a*b^2*B)) * \sqrt{\cos[c + dx]/(1 + \cos[c + dx])} * \sqrt{(a + b*\cos[c + dx])/((a + b)*(1 + \cos[c + dx]))} * \text{EllipticE}[\text{ArcSin}[\text{Tan}[(c + dx)/2]]], (-a + b)/(a + b)] + 2*a*(a + b)*(-6*A*b^2 + 3*a*b*(19*A + 7*B) + a^2*(25*A + 63*B)) * \sqrt{\cos[c + dx]/(1 + \cos[c + dx])} * \sqrt{(a + b*\cos[c + dx])/((a + b)*(1 + \cos[c + dx]))} * \text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + dx)/2]]], (-a + b)/(a + b)] - (82*a^2*A*b - 6*A*b^3 + 63*a^3*B + 21*a*b^2*B) * \cos[c + dx] * (a + b*\cos[c + dx]) * \sec[(c + dx)/2]^2 * \text{Tan}[(c + dx)/2]) / (105*a^2*d*\sqrt{a + b*\cos[c + dx]} * \sqrt{\sec[(c + dx)/2]^2}) + (\sqrt{a + b*\cos[c + dx]} * \sqrt{\sec[c + dx]} * ((-2*(-82*a^2*A*b + 6*A*b^3 - 63*a^3*B - 21*a*b^2*B) * \sin[c + dx]) / (105*a^2) + (2*\sec[c + dx]^2 * (8*A*b*\sin[c + dx] + 7*a*B*\sin[c + dx])) / 35 + (2*\sec[c + dx] * (25*a^2*A*\sin[c + dx] + 3*A*b^2*\sin[c + dx] + 42*a*b*B*\sin[c + dx])) / (105*a) + (2*a*A*\sec[c + dx]^2 * \text{Tan}[c + dx]) / 7)) / d$

3.598.3 Rubi [A] (verified)

Time = 2.25 (sec) , antiderivative size = 465, normalized size of antiderivative = 0.98, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{9}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{9/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow 3440$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$\downarrow 3042$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx$$

$$\downarrow 3468$$

3.598. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{7}\int\frac{b(4aA+7bB)\cos^2(c+dx)+(5Aa^2+14bBa+7Ab^2)\cos(c+dx)+a(8Ab+7A^2)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{b(4aA+7bB)\cos^2(c+dx)+(5Aa^2+14bBa+7Ab^2)\cos(c+dx)+a(8Ab+7A^2)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{b(4aA+7bB)\sin(c+dx+\frac{\pi}{2})^2+(5Aa^2+14bBa+7Ab^2)\sin(c+dx+\frac{\pi}{2})+a(8Ab+7A^2)}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2\int\frac{2ab(8Ab+7aB)\cos^2(c+dx)+a(21Ba^2+44Aba+35b^2B)\cos(c+dx)+a(25Aa^2+42bBa+3Ab^2)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{5a}\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{2ab(8Ab+7aB)\cos^2(c+dx)+a(21Ba^2+44Aba+35b^2B)\cos(c+dx)+a(25Aa^2+42bBa+3Ab^2)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{5a}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{2ab(8Ab+7aB)\sin(c+dx+\frac{\pi}{2})^2+a(21Ba^2+44Aba+35b^2B)\sin(c+dx+\frac{\pi}{2})+a(25Aa^2+42bBa+3Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{5a}\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2\int\frac{(25Aa^2+84bBa+51Ab^2)\cos(c+dx)a^2+(63Ba^3+82Aba^2+21b^2Ba-6Ab^3)a}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{3a}\right)+\frac{2(25a^2A+42abB+3Ab^2)}{3d\cos(c+dx)}\right)$$

↓ 27

3.598. $\int(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{(25Aa^2+84bBa+51Ab^2)\cos(c+dx)a^2+(63Ba^3+82Aba^2+21b^2Ba-6Ab^3)a}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{3a}+\frac{2(25a^2A+42abB+3Ab^2)}{3d\cos(c+dx)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{\int\frac{(25Aa^2+84bBa+51Ab^2)\sin(c+dx+\frac{\pi}{2})a^2+(63Ba^3+82Aba^2+21b^2Ba-6Ab^3)a}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{3a}+\frac{2(25a^2A+42abB+3Ab^2)}{3d\cos(c+dx)}\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{a(63a^3B+82a^2Ab+21ab^2B-6Ab^3)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-a(a-b)(-a^2(25A-63B))+a(57A-63B)}{3a}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{a(63a^3B+82a^2Ab+21ab^2B-6Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-a(a-b)(-a^2(25A-63B))+a(57A-63B)}{3a}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{a(63a^3B+82a^2Ab+21ab^2B-6Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2(a-b)\sqrt{a+b}}{3d\cos(c+dx)}(-a^2(25A-63B))+a(57A-63B)}{3a}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2(25a^2A+42abB+3Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)} + \frac{2(a-b)\sqrt{a+b}(63a^3B+82a^2Ab+21ab^2B-6Ab^3)\cot(c+dx)}{\dots}\right)\right)$$

input `Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(8*A*b + 7*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(82*a^2*A*b - 6*A*b^3 + 63*a^3*B + 21*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*(a - b)*Sqrt[a + b]*(6*A*b^2 - a^2*(25*A - 63*B) + a*(57*A*b - 21*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(3*a) + (2*(25*a^2*A + 3*A*b^2 + 42*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))/(5*a))/7)`

3.598.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)]])*Sqrt[(a_) + (b_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.598.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4626 vs. $2(427) = 854$.

Time = 28.85 (sec) , antiderivative size = 4627, normalized size of antiderivative = 9.78

method	result	size
parts	Expression too large to display	4627
default	Expression too large to display	4675

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x,method=_RET
URNVERBOSE)
```

output

```
-2/105*A/d*sec(d*x+c)^(9/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(-25*a^4
*cos(d*x+c)^4*sin(d*x+c)-15*a^4*cos(d*x+c)^2*sin(d*x+c)-27*a^2*b^2*cos(d*x
+c)^4*sin(d*x+c)+6*cos(d*x+c)^5*b^4*sin(d*x+c)+12*EllipticE(cot(d*x+c)-csc
(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2
)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^5+50*EllipticF(cot(d*x+
c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b
)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^5+6*EllipticE(cot
(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/
(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^4*cos(d*x+c)^4+25*Ellipti
cF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*
x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^4*cos(d*x+c)^4+6*El
lipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+c
os(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b^3*cos(d*x+c)
^4+82*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)
*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3*b*co
s(d*x+c)^4+51*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+co
s(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*
a^2*b^2*cos(d*x+c)^4-6*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2
))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)
))^(1/2)*a*b^3*cos(d*x+c)^4-164*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)...
```

3.598.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{9/2} dx$$

input

```
integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algo
rithm="fracas")
```

output

```
integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(
d*x + c) + a)*sec(d*x + c)^(9/2), x)
```


3.598.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(9/2),x)`

output `Timed out`

3.598.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{9/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(9/2), x)`

3.598.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{9/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(9/2), x)`

3.598. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx$

3.598.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{9/2} (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^(3/2), x)`

3.599 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$

3.599.1 Optimal result	5460
3.599.2 Mathematica [A] (warning: unable to verify)	5461
3.599.3 Rubi [A] (verified)	5461
3.599.4 Maple [B] (verified)	5465
3.599.5 Fricas [F]	5466
3.599.6 Sympy [F(-1)]	5467
3.599.7 Maxima [F]	5467
3.599.8 Giac [F]	5467
3.599.9 Mupad [F(-1)]	5468

3.599.1 Optimal result

Integrand size = 35, antiderivative size = 393

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{2(a - b)\sqrt{a + b}(9a^2A + 3Ab^2 + 20abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right) \mid -\frac{a - b}{a + b}\right) + 2(a - b)\sqrt{a + b}(9aA - 3Ab - 5aB + 15bB) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), \frac{a - b}{a + b}\right) + \frac{2(6Ab + 5aB)\sqrt{a + b \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{15d} + \frac{2aA\sqrt{a + b \cos(c + dx)} \sec^{5/2}(c + dx) \sin(c + dx)}{5d}$$

```
output 2/15*(6*A*b+5*B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/5*a*A*sec(d*x+c)^(5/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d+2/15*(a-b)*(9*A*a^2+3*A*b^2+20*B*a*b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^2/d/sec(d*x+c)^(1/2)-2/15*(a-b)*(9*A*a-3*A*b-5*B*a+15*B*b)*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a/d/sec(d*x+c)^(1/2))
```

3.599.2 Mathematica [A] (warning: unable to verify)

Time = 12.71 (sec) , antiderivative size = 379, normalized size of antiderivative = 0.96

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{2 \left(\sqrt{\cos^2(\frac{1}{2}(c+dx)) \sec(c+dx)} (-2(a+b)(9a^2A+3Ab^2+20abB)) E(\arcsin(\tan(\frac{1}{2}(c+dx))) | \frac{-a+b}{a+b}) \sqrt{\frac{1}{1+\sec(c+dx)}} \sqrt{\frac{b+a \sec(c+dx)}{(a+b)(1+\sec(c+dx))}} \right)}{\dots}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `(2*((Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(9*a^2*A + 3*A*b^2 + 20*a*b*B))*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))]) + 2*a*(a + b)*(3*b*(A + 5*B) + a*(9*A + 5*B))*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))]) - (9*a^2*A + 3*A*b^2 + 20*a*b*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2 + (a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]*((9*a^2*A + 3*A*b^2 + 20*a*b*B)*Sin[c + d*x] + a*(6*A*b + 5*a*B + 3*a*A*Sec[c + d*x])*Tan[c + d*x]))/(15*a*d*Sqrt[a + b*Cos[c + d*x]])`

3.599.3 Rubi [A] (verified)Time = 1.71 (sec) , antiderivative size = 380, normalized size of antiderivative = 0.97, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{7/2}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

3.599. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\cos^{7/2}(c+dx)} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}} dx$$

↓ 3468

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2}{5} \int \frac{b(2aA+5bB)\cos^2(c+dx) + (3Aa^2+10bBa+5Ab^2)\cos(c+dx) + a(6Ab+5A^2)}{2\cos^{5/2}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{5} \int \frac{b(2aA+5bB)\cos^2(c+dx) + (3Aa^2+10bBa+5Ab^2)\cos(c+dx) + a(6Ab+5A^2)}{\cos^{5/2}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{5} \int \frac{b(2aA+5bB)\sin(c+dx+\frac{\pi}{2})^2 + (3Aa^2+10bBa+5Ab^2)\sin(c+dx+\frac{\pi}{2}) + a(6Ab+5A^2)}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{5} \left(\frac{2 \int \frac{a(9Aa^2+20bBa+3Ab^2)+a(5Ba^2+12Aba+15b^2B)\cos(c+dx)}{2\cos^{3/2}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a} + \frac{2(5aB+6Ab)\sin(c+dx)}{3d\cos^{3/2}(c+dx)} \right) \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{5} \left(\frac{\int \frac{a(9Aa^2+20bBa+3Ab^2)+a(5Ba^2+12Aba+15b^2B)\cos(c+dx)}{\cos^{3/2}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a} + \frac{2(5aB+6Ab)\sin(c+dx)}{3d\cos^{3/2}(c+dx)} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{5} \left(\frac{\int \frac{a(9Aa^2+20bBa+3Ab^2)+a(5Ba^2+12Aba+15b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{3a} + \frac{2(5aB+6Ab)\sin(c+dx+\frac{\pi}{2})}{3d\cos^{3/2}(c+dx)} \right) \right)$$

↓ 3477

3.599. $\int (a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{7/2}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{a(9a^2A+20abB+3Ab^2)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-a(a-b)(9aA-5aB-3a)}{3a}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{a(9a^2A+20abB+3Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-a(a-b)(9aA-5aB-3a)}{3a}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{a(9a^2A+20abB+3Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2(a-b)\sqrt{a+b}(9aA-5aB-3a)}{3a}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{2(a-b)\sqrt{a+b}(9a^2A+20abB+3Ab^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{ad}\right)\right)$$

input `Int[(a + b*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(9*a^2*A + 3*A*b^2 + 20*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]), -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*(a - b)*Sqrt[a + b]*(9*a*A - 3*A*b - 5*a*B + 15*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]), -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(3*a) + (2*(6*A*b + 5*a*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)))/5)`

3.599.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_)] + (f_)*(x_))*(g_))^(p_)*((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`
- rule 3468 `Int[((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((A_)] + (B_)*sin[(e_)] + (f_)*(x_)]*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_
.)*(x_)])^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := S
imp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*
x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e
, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2,
0] && NeQ[A, B]
```

```
rule 3534 Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_)
+ (f_)*(x_)^2), x_Symbol] := Simp[-(A*b^2 - a*b*B + a^2*C)*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.599.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3581 vs. $2(353) = 706$.

Time = 24.06 (sec) , antiderivative size = 3582, normalized size of antiderivative = 9.11

method	result	size
parts	Expression too large to display	3582
default	Expression too large to display	3626

input `int((a*cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,method=_RETURNVERBOSE)`

output `2/5*A/d*sec(d*x+c)^(7/2)/(1+cos(d*x+c))/(a*cos(d*x+c)*b)^(1/2)*(2*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4-8*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^4-2*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4+3*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3+EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3-4*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3-EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3-3*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^3*cos(d*x+c)^3+sin(d*x+c)*cos(d*x+c)*a^3+3*sin(d*x+c)*cos(d*x+c)^2*a^2*b+6*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a*cos(d*x+c)*b)/(1+cos(d*x+c))...`

3.599.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{7/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x,algorithm="fracas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.599.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.599.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{7/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(7/2), x)`

3.599.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{7/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(7/2), x)`

3.599. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

3.599.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^(3/2), x)`

3.600 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$

3.600.1 Optimal result	5469
3.600.2 Mathematica [B] (warning: unable to verify)	5470
3.600.3 Rubi [A] (verified)	5470
3.600.4 Maple [B] (warning: unable to verify)	5475
3.600.5 Fricas [F]	5476
3.600.6 Sympy [F(-1)]	5476
3.600.7 Maxima [F]	5476
3.600.8 Giac [F]	5477
3.600.9 Mupad [F(-1)]	5477

3.600.1 Optimal result

Integrand size = 35, antiderivative size = 479

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \frac{2(a - b)\sqrt{a + b}(4Ab + 3aB)\sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}}}{3ad\sqrt{\sec(c + dx)}} + \frac{2\sqrt{a + b}(3Ab^2 + a^2(A - 3B) - a(4Ab - 6bB))\sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right)\right)}{3ad\sqrt{\sec(c + dx)}} - \frac{2b\sqrt{a + b}B\sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a + b}}}{d\sqrt{\sec(c + dx)}} + \frac{2aA\sqrt{a + b \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{3d}$$

output $\frac{2}{3}aA\sec(dx+c)^{3/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/3(a-b)(4A^2b+3B^2a)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/a/d/\sec(dx+c)^{1/2}+2/3(3A^2b^2+a^2(A-3B)-a(4Ab-6B^2b))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2})/a/d/\sec(dx+c)^{1/2}-2bB\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2})/d/\sec(dx+c)^{1/2}$

3.600.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 5981 vs. $2(479) = 958$.

Time = 22.34 (sec) , antiderivative size = 5981, normalized size of antiderivative = 12.49

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Result too large to show}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `Result too large to show`

3.600.3 Rubi [A] (verified)

Time = 1.90 (sec) , antiderivative size = 441, normalized size of antiderivative = 0.92, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{5/2}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

↓ 3042

3.600. $\int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

↓ 3440

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx$$

↓ 3042

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx$$

↓ 3468

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{2}{3} \int \frac{3b^2 B \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(4Ab + 3aB)}{2 \cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{2}{3} \int \frac{3b^2 B \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{2 \sin^{3/2}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right)$$

↓ 27

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{1}{3} \int \frac{3b^2 B \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(4Ab + 3aB)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + \frac{1}{3} \int \frac{3b^2 B \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{\sin^{3/2}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right)$$

↓ 3042

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{1}{3} \int \frac{3b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + \frac{1}{3} \int \frac{3b^2 B \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx) + a(4Ab + 3aB)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \right)$$

↓ 3532

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{1}{3} \left(\int \frac{a(4Ab + 3aB) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + 3b^2 B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) + \frac{1}{3} \left(\int \frac{3b^2 B \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{\sin^{3/2}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + 3b^2 B \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) \right)$$

↓ 3042

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{1}{3} \left(\int \frac{a(4Ab + 3aB) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + 3b^2 B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) + \frac{1}{3} \left(\int \frac{3b^2 B \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{\sin^{3/2}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + 3b^2 B \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) \right)$$

↓ 3288

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \left(\frac{1}{3} \left(\int \frac{a(4Ab + 3aB) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx + 3b^2 B \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \right) + \frac{1}{3} \left(\int \frac{3b^2 B \sin^2(c + dx + \frac{\pi}{2}) + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}) + a(4Ab + 3aB)}{\sin^{3/2}(c + dx + \frac{\pi}{2}) \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx + 3b^2 B \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) \right) - \frac{6bB\sqrt{a + b} \cot(c + dx)}{3}$$

3.600. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left((a^2(A-3B)-a(4Ab-6bB)+3Ab^2)\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}dx+\right.\right.$$

↓ 3042

$$\left.\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left((a^2(A-3B)-a(4Ab-6bB)+3Ab^2)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx)}}dx+\right.\right.$$

↓ 3295

$$\left.\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(a(3aB+4Ab)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2\sqrt{a+b}(a^2(A-3B)-a(4Ab-6bB)+3Ab^2)\cot(c+dx)}{ad}\right.\right.$$

↓ 3473

$$\left.\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{2\sqrt{a+b}(a^2(A-3B)-a(4Ab-6bB)+3Ab^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a+b}}}{ad}\right.\right.\right.$$

input `Int[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*(a - b)*Sqrt[a + b]*(4*A*b + 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(3*A*b^2 + a^2*(A - 3*B) - a*(4*A*b - 6*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (6*b*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/d)/3 + (2*a*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))`

3.600.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m*Int[(a + b*Sin[e + f*x])^n*(c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.600.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2488 vs. $2(433) = 866$.

Time = 17.87 (sec) , antiderivative size = 2489, normalized size of antiderivative = 5.20

method	result	size
parts	Expression too large to display	2489
default	Expression too large to display	3349

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2/3*A/d*sec(d*x+c)^(5/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(4*cos(d*x+c)^4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b+4*cos(d*x+c)^4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b^2-cos(d*x+c)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2-4*cos(d*x+c)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*a*b-3*cos(d*x+c)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*b^2+8*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b*cos(d*x+c)^3+8*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b^2*cos(d*x+c)^3-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)^3-8*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b*cos(d*x+c)^3-6*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a...
```

3.600.5 Fricas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{5/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorith="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2), x)`

3.600.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.600.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{5/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(5/2), x)`

3.600. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

3.600.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{5/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(5/2), x)`

3.600.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(3/2), x)`

3.601 $\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$

3.601.1 Optimal result	5478
3.601.2 Mathematica [A] (verified)	5479
3.601.3 Rubi [A] (verified)	5480
3.601.4 Maple [B] (warning: unable to verify)	5485
3.601.5 Fricas [F]	5486
3.601.6 Sympy [F(-1)]	5487
3.601.7 Maxima [F]	5487
3.601.8 Giac [F]	5487
3.601.9 Mupad [F(-1)]	5488

3.601.1 Optimal result

Integrand size = 35, antiderivative size = 509

$$\int (a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx = \frac{(a-b)\sqrt{a+b}(2aA-bB)\sqrt{\cos(c+dx)} \csc(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{ad\sqrt{\sec(c+dx)}} - \frac{\sqrt{a+b}(2a(A-B)-b(4A+B))\sqrt{\cos(c+dx)} \csc(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{d\sqrt{\sec(c+dx)}} - \frac{\sqrt{a+b}(2Ab+3aB)\sqrt{\cos(c+dx)} \csc(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{d\sqrt{\sec(c+dx)}} + \frac{2aA\sqrt{a+b \cos(c+dx)}\sqrt{\sec(c+dx)} \sin(c+dx)}{d} - \frac{(2aA-bB)\sqrt{a+b \cos(c+dx)}\sqrt{\sec(c+dx)} \sin(c+dx)}{d}$$

output $2*a*A*\sin(d*x+c)*(a+b*\cos(d*x+c))^{1/2}*sec(d*x+c)^{1/2}/d-(2*A*a-B*b)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{1/2}*sec(d*x+c)^{1/2}/d+(a-b)*(2*A*a-B*b)*csc(d*x+c)*EllipticE((a+b*\cos(d*x+c))^{1/2}/(a+b)^{1/2}/\cos(d*x+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(d*x+c)^{1/2}*(a*(1-sec(d*x+c)))/(a+b)^{1/2}*(a*(1+sec(d*x+c)))/(a-b)^{1/2}/a/d/sec(d*x+c)^{1/2}-(2*a*(A-B)-b*(4*A+B))*csc(d*x+c)*EllipticF((a+b*\cos(d*x+c))^{1/2}/(a+b)^{1/2}/\cos(d*x+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(d*x+c)^{1/2}*(a*(1-sec(d*x+c)))/(a+b)^{1/2}*(a*(1+sec(d*x+c)))/(a-b)^{1/2}/d/sec(d*x+c)^{1/2}-(2*A*b+3*B*a)*csc(d*x+c)*EllipticPi((a+b*\cos(d*x+c))^{1/2}/(a+b)^{1/2}/\cos(d*x+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(d*x+c)^{1/2}*(a*(1-sec(d*x+c)))/(a+b)^{1/2}*(a*(1+sec(d*x+c)))/(a-b)^{1/2}/d/sec(d*x+c)^{1/2}$

3.601.2 Mathematica [A] (verified)

Time = 14.77 (sec) , antiderivative size = 927, normalized size of antiderivative = 1.82

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{2aA\sqrt{a + b \cos(c + dx)}\sqrt{\sec(c + dx)} \sin(c + dx)}{d} + \frac{\sqrt{\frac{1}{1 - \tan^2(\frac{1}{2}(c + dx))}} \left(-2a^2A \tan\left(\frac{1}{2}(c + dx)\right) - 2aAb \tan\left(\frac{1}{2}(c + dx)\right) + abB \tan\left(\frac{1}{2}(c + dx)\right) + b^2B \tan\left(\frac{1}{2}(c + dx)\right) \right)}{d}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output $(2*a*A*\sqrt{a + b*\cos[c + d*x]}\sqrt{\sec[c + d*x]}\sin[c + d*x])/d + (\sqrt{[(1 - \tan[(c + d*x)/2]^2)^{-1}]*(-2*a^2*A*\tan[(c + d*x)/2] - 2*a*A*b*\tan[(c + d*x)/2] + a*b*B*\tan[(c + d*x)/2] + b^2*B*\tan[(c + d*x)/2] + 4*a*A*b*\tan[(c + d*x)/2]^3 - 2*b^2*B*\tan[(c + d*x)/2]^3 + 2*a^2*A*\tan[(c + d*x)/2]^5 - 2*a*A*b*\tan[(c + d*x)/2]^5 - a*b*B*\tan[(c + d*x)/2]^5 + b^2*B*\tan[(c + d*x)/2]^5 + 4*A*b^2*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + d*x)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + d*x)/2]^2}\sqrt{(a + b + a*\tan[(c + d*x)/2]^2 - b*\tan[(c + d*x)/2]^2)/(a + b)} + 6*a*b*B*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + d*x)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + d*x)/2]^2}\sqrt{(a + b + a*\tan[(c + d*x)/2]^2 - b*\tan[(c + d*x)/2]^2)/(a + b)} + 4*A*b^2*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + d*x)/2]], (-a + b)/(a + b)]*\tan[(c + d*x)/2]^2*\sqrt{1 - \tan[(c + d*x)/2]^2}\sqrt{(a + b + a*\tan[(c + d*x)/2]^2 - b*\tan[(c + d*x)/2]^2)/(a + b)} + 6*a*b*B*\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + d*x)/2]], (-a + b)/(a + b)]*\tan[(c + d*x)/2]^2*\sqrt{1 - \tan[(c + d*x)/2]^2}\sqrt{(a + b + a*\tan[(c + d*x)/2]^2 - b*\tan[(c + d*x)/2]^2)/(a + b)} - (a + b)*(2*a*A - b*B)*\text{EllipticE}[\text{ArcSin}[\tan[(c + d*x)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + d*x)/2]^2}*(1 + \tan[(c + d*x)/2]^2)\sqrt{(a + b + a*\tan[(c + d*x)/2]^2 - b*\tan[(c + d*x)/2]^2)/(a + b)} + 2*(-(A*b^2) + 2*a*b*(A - B) + a^2*(A + B))*\text{EllipticF}[\text{ArcSin}[\tan[(c + d*x)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + d*x)/2]^2}*(1 + \tan[(c + d*x)/2]^2)\sqrt{(a + b + a*\tan[(c + d*x)/2]^2 - b*\tan...$

3.601.3 Rubi [A] (verified)

Time = 2.38 (sec) , antiderivative size = 482, normalized size of antiderivative = 0.95, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$3.601. \quad \int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx \\ & \downarrow 3468 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(2 \int \frac{-b(2aA-bB)\cos^2(c+dx) - (Aa^2-2bBa-Ab^2)\cos(c+dx) + a(2Ab+aB)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right) \\ & \downarrow 27 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{-b(2aA-bB)\cos^2(c+dx) - (Aa^2-2bBa-Ab^2)\cos(c+dx) + a(2Ab+aB)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right) \\ & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{-b(2aA-bB)\sin(c+dx+\frac{\pi}{2})^2 + (-Aa^2+2bBa+Ab^2)\sin(c+dx+\frac{\pi}{2}) + a(2Ab+aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \\ & \downarrow 3540 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{b^2(2Ab+3aB)\cos^2(c+dx)+2ab(2Ab+aB)\cos(c+dx)+ab(2aA-bB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} - \frac{(2aA-bB)\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) \\ & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{b^2(2Ab+3aB)\sin(c+dx+\frac{\pi}{2})^2+2ab(2Ab+aB)\sin(c+dx+\frac{\pi}{2})+ab(2aA-bB)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} - \frac{(2aA-bB)\sin(c+dx+\frac{\pi}{2})}{d\sqrt{\sin(c+dx+\frac{\pi}{2})}} \right) \\ & \downarrow 3532 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{b^2(3aB+2Ab) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx + \int \frac{ab(2aA-bB)+2ab(2Ab+aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} - \frac{(2aA-bB)\sin(c+dx)}{d\sqrt{\cos(c+dx)}} \right) \\ & \downarrow 3042 \end{aligned}$$

3.601. $\int (a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{b^2(3aB+2Ab) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{ab(2aA-bB)+2ab(2Ab+aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} - (2) \right.$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{ab(2aA-bB)+2ab(2Ab+aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(3aB+2Ab)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a+b}}}{2b}}{2b} \right.$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{ab(2aA-bB) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - ab(2aA-2aB-4Ab-bB) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{2b} \right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{-ab(2aA-2aB-4Ab-bB) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + ab(2aA-bB) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{2b} \right.$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{ab(2aA-bB) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2b\sqrt{a+b}(2aA-2aB-4Ab-bB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a+b}}}{2b}}{2b} \right.$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2b\sqrt{a+b}(2aA-2aB-4Ab-bB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a+b}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{d} \right.$$

3.601. $\int (a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx) dx$

input `Int[(a + b*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*(a - b)*b*Sqrt[a + b]*(2*a*A - b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(2*a*A - 4*A*b - 2*a*B - b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (2*b*Sqrt[a + b]*(2*A*b + 3*a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(2*b) + (2*a*A*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - ((2*a*A - b*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))`

3.601.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)]/Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*sin[e + f*x]]/Sqrt[b*sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)]*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*sin[e + f*x]]/Sqrt[d*sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

$$3.601. \quad \int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$$

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*SIN[e + f*x])^(m - 2)*(c + d*SIN[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*SIN[e + f*x]]/Sqrt[b*SIN[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*SIN[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + SIN[e + f*x])/((a + b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)])), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.601.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 2385 vs. $2(465) = 930$.

Time = 19.43 (sec) , antiderivative size = 2386, normalized size of antiderivative = 4.69

method	result	size
parts	Expression too large to display	2386
default	Expression too large to display	3276

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

output

```

-2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(
d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(
d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2-2*(-csc(d*x+c)^2*(1-cos(d*x+c)
)^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c)
))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
))*a*b+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*
x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+
1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2
+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a
^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c)
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)
-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b-2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)
^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a
+b)/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))
)*b^2+csc(d*x+c)^3*a^2*(1-cos(d*x+c))^3-csc(d*x+c)^3*a*b*(1-cos(d*x+c))^3+a
^2*(csc(d*x+c)-cot(d*x+c))+a*b*(csc(d*x+c)-cot(d*x+c))*((csc(d*x+c)^2*a*(
1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(
d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(-csc(d*x+c)^2*(1-c
os(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1))^(3/2)/(csc(d*x+c)^2*...

```

3.601.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2} dx$$

input

```

integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo
rithm="fricas")

```

output

```

integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(
d*x + c) + a)*sec(d*x + c)^(3/2), x)

```

3.601.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.601.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2), x)`

3.601.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2), x)`

3.601.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(3/2), x)`

3.602 $\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}$

3.602.1 Optimal result	5489
3.602.2 Mathematica [B] (verified)	5490
3.602.3 Rubi [A] (verified)	5491
3.602.4 Maple [B] (verified)	5497
3.602.5 Fricas [F]	5498
3.602.6 Sympy [F(-1)]	5498
3.602.7 Maxima [F]	5498
3.602.8 Giac [F]	5499
3.602.9 Mupad [F(-1)]	5499

3.602.1 Optimal result

Integrand size = 35, antiderivative size = 532

$$\int (a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx =$$

$$\frac{(a-b)\sqrt{a+b}(4Ab+5aB)\sqrt{\cos(c+dx)} \operatorname{csc}(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\sec(c+dx)}}{4ad\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(8aA+4Ab+5aB+2bB)\sqrt{\cos(c+dx)} \operatorname{csc}(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\sec(c+dx)}}{4d\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(12aAb+3a^2B+4b^2B)\sqrt{\cos(c+dx)} \operatorname{csc}(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\sec(c+dx)}}{4bd\sqrt{\sec(c+dx)}} +$$

$$\frac{bB\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{2d\sqrt{\sec(c+dx)}} +$$

$$\frac{(4Ab+5aB)\sqrt{a+b \cos(c+dx)} \sqrt{\sec(c+dx)} \sin(c+dx)}{4d}$$

output $\frac{1}{2}bB\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\sec(dx+c)^{1/2}+1/4(4Ab+5B^2a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/d-1/4(a-b)(4A^2b+5B^2a)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/a/d\sec(dx+c)^{1/2}+1/4(8A^2a+4A^2b+5B^2a+2B^2b)\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/d\sec(dx+c)^{1/2}-1/4(12A^2ab+3B^2a^2+4B^2b^2)\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b))^{1/2}/b/d\sec(dx+c)^{1/2}$

3.602.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1134 vs. $2(532) = 1064$.

Time = 15.12 (sec) , antiderivative size = 1134, normalized size of antiderivative = 2.13

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output

```
(b*B*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*Sin[2*(c + d*x)]/(4*d) +
(Sqrt[(1 - Tan[(c + d*x)/2]^2)^(-1)]*(4*a*A*b*Tan[(c + d*x)/2] + 4*A*b^2*
Tan[(c + d*x)/2] + 5*a^2*B*Tan[(c + d*x)/2] + 5*a*b*B*Tan[(c + d*x)/2] - 8
*A*b^2*Tan[(c + d*x)/2]^3 - 10*a*b*B*Tan[(c + d*x)/2]^3 - 4*a*A*b*Tan[(c +
d*x)/2]^5 + 4*A*b^2*Tan[(c + d*x)/2]^5 - 5*a^2*B*Tan[(c + d*x)/2]^5 + 5*a
*b*B*Tan[(c + d*x)/2]^5 + 24*a*A*b*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]]
, (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c +
d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + 6*a^2*B*EllipticPi[-1, ArcSin
[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a
+ b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + 8*b^2*B*Ell
ipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c +
d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a +
b)] + 24*a*A*b*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]
*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c +
d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + 6*a^2*B*EllipticPi[-1, ArcSin
[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c +
d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a
+ b)] + 8*b^2*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]
*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c +
d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + (a + b)*(4*A*b + 5*a*B)*El...
```

3.602.3 Rubi [A] (verified)

Time = 2.41 (sec) , antiderivative size = 496, normalized size of antiderivative = 0.93, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c + dx)}(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2}\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

3.602. $\int (a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx$$

$$\downarrow \text{3469}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \int \frac{b(4Ab+5aB)\cos^2(c+dx)+2(2Ba^2+4Aba+b^2B)\cos(c+dx)+a(4aA+bB)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \int \frac{b(4Ab+5aB)\cos^2(c+dx)+2(2Ba^2+4Aba+b^2B)\cos(c+dx)+a(4aA+bB)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \int \frac{b(4Ab+5aB)\sin(c+dx+\frac{\pi}{2})^2+2(2Ba^2+4Aba+b^2B)\sin(c+dx+\frac{\pi}{2})+a(4aA+bB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

$$\downarrow \text{3540}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{\int -\frac{b(3Ba^2+12Aba+4b^2B)\cos^2(c+dx)-2ab(4aA+bB)\cos(c+dx)+ab(4Ab+5aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} + \frac{(5aB+b^2)}{2b} \right) \right)$$

$$\downarrow \text{25}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{-b(3Ba^2+12Aba+4b^2B)\cos^2(c+dx)-2ab(4aA+bB)\cos(c+dx)+ab(4Ab+5aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} \right) \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{-b(3Ba^2+12Aba+4b^2B)\sin(c+dx+\frac{\pi}{2})-2ab(4aA+bB)\sin(c+dx+\frac{\pi}{2})+ab(4Ab+5aB)}{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} \right) \right)$$

$$\downarrow \text{3532}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{\int\frac{ab(4Ab+5aB)-2ab(4aA+bB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{\int\frac{ab(4Ab+5aB)-2ab(4aA+bB)\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{\int\frac{ab(4Ab+5aB)-2ab(4aA+bB)\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{ab(5aB+4Ab)\int\frac{\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{1}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{-ab(8aA+5aB+4Ab+2bB)\int\frac{\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx}{1}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{ab(5aB+4Ab)\int\frac{\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{1}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{2\sqrt{a+b}(3a^2B+12aAb+4b^2B)\cot(c+dx)}{d\sqrt{\cos(c+dx)}}\right)\right)$$

input `Int[(a + b*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x])*Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((b*B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*(4*A*b + 5*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(8*a*A + 4*A*b + 5*a*B + 2*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d + (2*Sqrt[a + b]*(12*a*A*b + 3*a^2*B + 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/b + ((4*A*b + 5*a*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/4`

3.602.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]], x_Symbol] :> Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B))*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])3/2*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])3/2*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])3/2*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]2)/(((a_) + (b_)*sin[(e_) + (f_)*(x_)])3/2*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])3/2*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]2)/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])3/2*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.602.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3309 vs. $2(478) = 956$.

Time = 18.06 (sec) , antiderivative size = 3310, normalized size of antiderivative = 6.22

method	result	size
parts	Expression too large to display	3310
default	Expression too large to display	3318

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -A/d*(EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)
*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(
d*x+c)^2+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)
))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^2*c
os(d*x+c)^2+6*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d
*x+c))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/
2))*a*b*cos(d*x+c)^2+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)
/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)
)^(1/2))*a^2*cos(d*x+c)^2-4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)
)^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d
*x+c)))^(1/2)*a*b*cos(d*x+c)^2+2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(
a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+
cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+
cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),
(-a-b)/(a+b))^(1/2))*b^2*cos(d*x+c)+12*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*
((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*
x+c),-1,(-a-b)/(a+b))^(1/2))*a*b*cos(d*x+c)+4*(cos(d*x+c)/(1+cos(d*x+c)))
^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-
csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*cos(d*x+c)-8*EllipticF(cot(d*x+c)-csc
(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(...
```


3.602.5 Fricas [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algo rithm="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c)), x)`

3.602.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Timed out`

3.602.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*x + c)), x)`

3.602. $\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

3.602.8 Giac [F]

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*x + c)), x)`

3.602.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx))^{3/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(3/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(3/2), x)`

3.603
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.603.1 Optimal result 5500
 3.603.2 Mathematica [B] (verified) 5501
 3.603.3 Rubi [A] (verified) 5502
 3.603.4 Maple [B] (verified) 5509
 3.603.5 Fricas [F] 5510
 3.603.6 Sympy [F(-1)] 5510
 3.603.7 Maxima [F] 5510
 3.603.8 Giac [F] 5511
 3.603.9 Mupad [F(-1)] 5511

3.603.1 Optimal result

Integrand size = 35, antiderivative size = 626

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx =$$

$$\frac{(a - b)\sqrt{a + b}(30aAb + 3a^2B + 16b^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{a + b}}{24abd \sqrt{\sec(c + dx)}} +$$

$$\frac{\sqrt{a + b}(30aAb + 12Ab^2 + 3a^2B + 14abB + 16b^2B) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right)\right)}{24bd \sqrt{\sec(c + dx)}} +$$

$$\frac{\sqrt{a + b}(6a^2Ab + 8Ab^3 - a^3B + 12ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right)\right)}{8b^2d \sqrt{\sec(c + dx)}} +$$

$$\frac{bB \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{3d \sec^{\frac{3}{2}}(c + dx)} + \frac{(6Ab + 7aB) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{12d \sqrt{\sec(c + dx)}} +$$

$$\frac{(30aAb + 3a^2B + 16b^2B) \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{24bd}$$

output $\frac{1}{3}bB\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d/\sec(dx+c)^{3/2}+1/12(6A^2b+7B^2a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d/\sec(dx+c)^{1/2}+1/24(30A^2ab+3B^2a^2+16B^2b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/bd-1/24(a-b)(30A^2ab+3B^2a^2+16B^2b^2)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2})/a/b/d/\sec(dx+c)^{1/2}+1/24(30A^2ab+12A^2b^2+3B^2a^2+14B^2ab+16B^2b^2)\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2})/b/d/\sec(dx+c)^{1/2}-1/8(6A^2a^2b+8A^2ab^3-B^2a^3+12B^2ab^2)\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2})/b^2/d/\sec(dx+c)^{1/2}$

3.603.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1489 vs. $2(626) = 1252$.

Time = 15.36 (sec) , antiderivative size = 1489, normalized size of antiderivative = 2.38

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Too large to display}$$

input `Integrate[((a + bCos[c + d*x])^(3/2)*(A + BCos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output $(\text{Sqrt}[a + b \cos[c + dx]] \cdot \text{Sqrt}[\text{Sec}[c + dx]] \cdot ((b \cdot B \cdot \sin[c + dx])/12 + ((6 \cdot A \cdot b + 7 \cdot a \cdot B) \cdot \sin[2 \cdot (c + dx)])/24 + (b \cdot B \cdot \sin[3 \cdot (c + dx)])/12))/d + (\text{Sqrt}[(1 - \tan[(c + dx)/2]^2)^{-1}] \cdot (30 \cdot a^2 \cdot A \cdot b \cdot \tan[(c + dx)/2] + 30 \cdot a \cdot A \cdot b^2 \cdot \tan[(c + dx)/2] + 3 \cdot a^3 \cdot B \cdot \tan[(c + dx)/2] + 3 \cdot a^2 \cdot b \cdot B \cdot \tan[(c + dx)/2] + 16 \cdot a \cdot b^2 \cdot B \cdot \tan[(c + dx)/2] + 16 \cdot b^3 \cdot B \cdot \tan[(c + dx)/2] - 60 \cdot a \cdot A \cdot b^2 \cdot \tan[(c + dx)/2]^3 - 6 \cdot a^2 \cdot b \cdot B \cdot \tan[(c + dx)/2]^3 - 32 \cdot b^3 \cdot B \cdot \tan[(c + dx)/2]^3 - 30 \cdot a^2 \cdot A \cdot b \cdot \tan[(c + dx)/2]^5 + 30 \cdot a \cdot A \cdot b^2 \cdot \tan[(c + dx)/2]^5 - 3 \cdot a^3 \cdot B \cdot \tan[(c + dx)/2]^5 + 3 \cdot a^2 \cdot b \cdot B \cdot \tan[(c + dx)/2]^5 - 16 \cdot a \cdot b^2 \cdot B \cdot \tan[(c + dx)/2]^5 + 16 \cdot b^3 \cdot B \cdot \tan[(c + dx)/2]^5 + 36 \cdot a^2 \cdot A \cdot b \cdot \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \cdot \tan[(c + dx)/2]^2 - b \cdot \tan[(c + dx)/2]^2)/(a + b)] + 48 \cdot A \cdot b^3 \cdot \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \cdot \tan[(c + dx)/2]^2 - b \cdot \tan[(c + dx)/2]^2)/(a + b)] - 6 \cdot a^3 \cdot B \cdot \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \cdot \tan[(c + dx)/2]^2 - b \cdot \tan[(c + dx)/2]^2)/(a + b)] + 72 \cdot a \cdot b^2 \cdot B \cdot \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \cdot \tan[(c + dx)/2]^2 - b \cdot \tan[(c + dx)/2]^2)/(a + b)] + 36 \cdot a^2 \cdot A \cdot b \cdot \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \tan[(c + dx)/2]^2 \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \cdot \tan[(c + dx)/2]^2 - b \cdot \tan[(c + dx)/2]^2) \dots$

3.603.3 Rubi [A] (verified)

Time = 3.05 (sec) , antiderivative size = 589, normalized size of antiderivative = 0.94, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx$$

3.603. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{3/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3469

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \int \frac{\sqrt{\cos(c+dx)}(b(6Ab+7aB)\cos^2(c+dx)+2(3Ba^2+6Aba+2b^2B)\cos(c+dx))}{2\sqrt{a+b\cos(c+dx)}} dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \int \frac{\sqrt{\cos(c+dx)}(b(6Ab+7aB)\cos^2(c+dx)+2(3Ba^2+6Aba+2b^2B)\cos(c+dx))}{\sqrt{a+b\cos(c+dx)}} dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \int \frac{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}(b(6Ab+7aB)\sin\left(c+dx+\frac{\pi}{2}\right)^2+2(3Ba^2+6Aba+2b^2B)\sin\left(c+dx+\frac{\pi}{2}\right))}{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} dx\right)$$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{\int \frac{b(3Ba^2+30Aba+16b^2B)\cos^2(c+dx)+2b(12Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab(6Ab+7aB)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{2b} + \dots\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{\int \frac{b(3Ba^2+30Aba+16b^2B)\cos^2(c+dx)+2b(12Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab(6Ab+7aB)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{4b} + \dots\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{\int \frac{b(3Ba^2+30Aba+16b^2B)\sin\left(c+dx+\frac{\pi}{2}\right)^2+2b(12Aa^2+13bBa+6Ab^2)\sin\left(c+dx+\frac{\pi}{2}\right)+ab(6Ab+7aB)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{4b} + \dots\right)\right)$$

↓ 3540

3.603. $\int \frac{(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{\int \frac{-2a(6Ab+7aB)\cos(c+dx)b^2-3(-Ba^3+6Aba^2+12b^2Ba+8Ab^3)\cos^2(c+dx)b+a(3Ba^2+30Aba+16b^2B)b}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} \right) \right)$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{-2a(6Ab+7aB)\cos(c+dx)b^2-3(-Ba^3+6Aba^2+12b^2Ba+8Ab^3)\cos^2(c+dx)b+a(3Ba^2+30Aba+16b^2B)b}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{4b} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{-2a(6Ab+7aB)\sin(c+dx+\frac{\pi}{2})b^2-3(-Ba^3+6Aba^2+12b^2Ba+8Ab^3)\sin^2(c+dx+\frac{\pi}{2})b+a(3Ba^2+30Aba+16b^2B)b}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{4b} \right) \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(3Ba^2+30Aba+16b^2B)-2ab^2(6Ab+7aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{4b} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(3Ba^2+30Aba+16b^2B)-2ab^2(6Ab+7aB)\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{4b} \right) \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{\int \frac{ab(3Ba^2+30Aba+16b^2B)-2ab^2(6Ab+7aB)\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{ab(3a^2B+30aAb+16b^2B)\int \frac{\cos(c+dx)+\frac{3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{-ab(3a^2B+30aAb+14abB+12Ab^2+16b^2B)\int \frac{\cos(c+dx)+\frac{3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{ab(3a^2B+30aAb+16b^2B)\int \frac{\sin(c+dx)+\frac{3}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{(3a^2B+30aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} - \frac{2b\sqrt{a+b}(3a^2B+30aAb+14abB+12Ab^2+16b^2B)\int \frac{\cos(c+dx)+\frac{3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \right) \right)$$

input `Int[((a + b*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((b*B*cos[c + d*x]^(3/2)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(3*d) + (((6*A*b + 7*a*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*(30*a*A*b + 3*a^2*B + 16*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(30*a*A*b + 12*A*b^2 + 3*a^2*B + 14*a*b*B + 16*b^2*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d + (6*Sqrt[a + b]*(6*a^2*A*b + 8*A*b^3 - a^3*B + 12*a*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/b + ((30*a*A*b + 3*a^2*B + 16*b^2*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])/(4*b))/6`

3.603.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.603.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4188 vs. $2(566) = 1132$.

Time = 17.61 (sec) , antiderivative size = 4189, normalized size of antiderivative = 6.69

method	result	size
parts	Expression too large to display	4189
default	Expression too large to display	4235

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*A/d/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(1/2)*(5*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b*cos(d*x+c)-2*a*b*sin(d*x+c)-4*sec(d*x+c)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2+6*sec(d*x+c)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2+8*sec(d*x+c)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2-7*a*b*cos(d*x+c)*sin(d*x+c)+5*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2*cos(d*x+c)+6*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2*cos(d*x+c)+8*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*b^2*cos...
```

3.603.5 Fricas [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral((B*b*cos(d*x + c)^2 + A*a + (B*a + A*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/sqrt(sec(d*x + c)), x)`

3.603.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.603.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/sqrt(sec(d*x + c)), x)`

3.603.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/sqrt(sec(d*x + c)), x)`

3.603.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\sqrt{\frac{1}{\cos(c+dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(1/2), x)`

3.604
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.604.1 Optimal result 5512
 3.604.2 Mathematica [B] (verified) 5513
 3.604.3 Rubi [A] (verified) 5514
 3.604.4 Maple [B] (verified) 5522
 3.604.5 Fracas [F(-1)] 5522
 3.604.6 Sympy [F(-1)] 5523
 3.604.7 Maxima [F] 5523
 3.604.8 Giac [F] 5523
 3.604.9 Mupad [F(-1)] 5524

3.604.1 Optimal result

Integrand size = 35, antiderivative size = 730

$$\int \frac{(a + b \cos(c + dx))^{3/2}(A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{(a - b)\sqrt{a + b}(24a^2Ab + 128Ab^3 - 9a^3B + 156ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{192ab^2d\sqrt{\sec(c + dx)}} -$$

$$\frac{\sqrt{a + b}(9a^3B - 6a^2b(4A + B) - 8b^3(16A + 9B) - 4ab^2(28A + 39B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{192b^2d\sqrt{\sec(c + dx)}} +$$

$$\frac{\sqrt{a + b}(8a^3Ab - 96aAb^3 - 3a^4B - 24a^2b^2B - 48b^4B) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{64b^3d\sqrt{\sec(c + dx)}} +$$

$$\frac{(8aAb - 3a^2B + 12b^2B) \sqrt{a + b \cos(c + dx)} \sin(c + dx)}{32bd\sqrt{\sec(c + dx)}} +$$

$$\frac{(8Ab - 3aB)(a + b \cos(c + dx))^{3/2} \sin(c + dx)}{24bd\sqrt{\sec(c + dx)}} + \frac{B(a + b \cos(c + dx))^{5/2} \sin(c + dx)}{4bd\sqrt{\sec(c + dx)}} +$$

$$\frac{(24a^2Ab + 128Ab^3 - 9a^3B + 156ab^2B) \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{192b^2d}$$

3.604.
$$\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

output $\frac{1}{24}(8Ab-3B^2a)(a+b\cos(dx+c))^{3/2}\sin(dx+c)/b/d/\sec(dx+c)^{1/2} + \frac{1}{4}B(a+b\cos(dx+c))^{5/2}\sin(dx+c)/b/d/\sec(dx+c)^{1/2} + \frac{1}{32}(8A^2ab-3B^2a^2+12B^2b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b/d/\sec(dx+c)^{1/2} + \frac{1}{192}(24A^2a^2b+128A^2b^3-9B^2a^3+156B^2ab^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/b^2/d - \frac{1}{192}(a-b)(24A^2a^2b+128A^2b^3-9B^2a^3+156B^2ab^2)\operatorname{csc}(dx+c)\operatorname{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2})/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/a/b^2/d/\sec(dx+c)^{1/2} - \frac{1}{192}(9B^2a^3-6a^2b(4A+B)-8b^3(16A+9B)-4ab^2(28A+39B))\operatorname{csc}(dx+c)\operatorname{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/b^2/d/\sec(dx+c)^{1/2} + \frac{1}{64}(8A^3b-96A^2ab^3-3B^2a^4-24B^2a^2b^2-48B^2b^4)\operatorname{csc}(dx+c)\operatorname{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/b^3/d/\sec(dx+c)^{1/2}$

3.604.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1888 vs. $2(730) = 1460$.

Time = 19.45 (sec) , antiderivative size = 1888, normalized size of antiderivative = 2.59

$$\int \frac{(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx = \text{Too large to display}$$

input `Integrate[((a + b*Cos[c + d*x])^(3/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2), x]`

output $(\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}*((8A*b + 9a*B)\sin[c + dx])/96 + ((56*a*A*b + 3*a^2*B + 48*b^2*B)\sin[2*(c + dx)]/(192*b) + ((8A*b + 9a*B)\sin[3*(c + dx)]/96 + (b*B\sin[4*(c + dx)]/32))/d - (\sqrt{(1 - \tan[(c + dx)/2]^2)^{-1}}\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(1 + \tan[(c + dx)/2]^2)}*(24*a^3*A*b*\tan[(c + dx)/2] + 24*a^2*A*b^2*\tan[(c + dx)/2] + 128*a*A*b^3*\tan[(c + dx)/2] + 128*A*b^4*\tan[(c + dx)/2] - 9*a^4*B*\tan[(c + dx)/2] - 9*a^3*b*B*\tan[(c + dx)/2] + 156*a^2*b^2*B*\tan[(c + dx)/2] + 156*a*b^3*B*\tan[(c + dx)/2] - 48*a^2*A*b^2*\tan[(c + dx)/2]^3 - 256*A*b^4*\tan[(c + dx)/2]^3 + 18*a^3*b*B*\tan[(c + dx)/2]^3 - 312*a*b^3*B*\tan[(c + dx)/2]^3 - 24*a^3*A*b*\tan[(c + dx)/2]^5 + 24*a^2*A*b^2*\tan[(c + dx)/2]^5 - 128*a*A*b^3*\tan[(c + dx)/2]^5 + 128*A*b^4*\tan[(c + dx)/2]^5 + 9*a^4*B*\tan[(c + dx)/2]^5 - 9*a^3*b*B*\tan[(c + dx)/2]^5 - 156*a^2*b^2*B*\tan[(c + dx)/2]^5 + 156*a*b^3*B*\tan[(c + dx)/2]^5 - 48*a^3*A*b*EllipticPi[-1, ArcSin[Tan[(c + dx)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + 576*a*A*b^3*EllipticPi[-1, ArcSin[Tan[(c + dx)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + 18*a^4*B*EllipticPi[-1, ArcSin[Tan[(c + dx)/2]], (-a + b)/(a + b)]*\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a*\tan[(c + dx)/2]^2 - b*\tan[(c + dx)/2]^2)/(a + b)} + 144...$

3.604.3 Rubi [A] (verified)

Time = 3.66 (sec) , antiderivative size = 696, normalized size of antiderivative = 0.95, number of steps used = 22, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.629$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \cos^{\frac{3}{2}}(c + dx) (a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx)) dx$$

3.604. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{3/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3469

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a+b\cos(c+dx))^{3/2}((8Ab-3aB)\cos^2(c+dx)+6bB\cos(c+dx)+aB) dx}{2\sqrt{\cos(c+dx)}}}{4b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{8b} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a+b\cos(c+dx))^{3/2}((8Ab-3aB)\cos^2(c+dx)+6bB\cos(c+dx)+aB) dx}{\sqrt{\cos(c+dx)}}}{8b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{8b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}((8Ab-3aB)\sin^2(c+dx+\frac{\pi}{2})+6bB\sin(c+dx+\frac{\pi}{2})+aB) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{8b} + \frac{B\sin(c+dx)}{8b} \right)$$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{3} \int \frac{\sqrt{a+b\cos(c+dx)}(3(-3Ba^2+8Aba+12b^2B)\cos^2(c+dx)+2b(16Ab+15aB)\cos(c+dx)+a(8Ab+3aB)) dx}{2\sqrt{\cos(c+dx)}}}{8b} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{6} \int \frac{\sqrt{a+b\cos(c+dx)}(3(-3Ba^2+8Aba+12b^2B)\cos^2(c+dx)+2b(16Ab+15aB)\cos(c+dx)+a(8Ab+3aB)) dx}{\sqrt{\cos(c+dx)}}}{8b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{6} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(3(-3Ba^2+8Aba+12b^2B)\sin^2(c+dx+\frac{\pi}{2})+2b(16Ab+15aB)\sin(c+dx+\frac{\pi}{2})+a(8Ab+3aB)) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{8b} \right)$$

3.604. $\int \frac{(a+b\cos(c+dx))^{3/2}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) \cos^2(c+dx)+2b(57Ba^2+104Aba+36b^2B) \cos(c+dx)+a(3B}{2\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} \right) \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) \cos^2(c+dx)+2b(57Ba^2+104Aba+36b^2B) \cos(c+dx)+a(3B}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{(-9Ba^3+24Aba^2+156b^2Ba+128Ab^3) \sin(c+dx+\frac{\pi}{2})^2+2b(57Ba^2+104Aba+36b^2B) \sin(c+dx+\frac{\pi}{2}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} \right) \right)$$

↓ 3540

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\int -\frac{3(-3Ba^4+8Aba^3-24b^2Ba^2-96Ab^3a-48b^4B) \cos^2(c+dx)-2ab(3Ba^2+56Aba+36b^2B) \cos(c+dx)+a(-9}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} \right) \right) \right)$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{3(-3Ba^4+8Aba^3-24b^2Ba^2}{\sec^{\frac{3}{2}}(c+dx)} \right) \right) \right)$$

↓ 3042

3.604. $\int \frac{(a+b \cos(c+dx))^{3/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{3(-3Ba^4+8Aba^3-24b^2Ba^2}{\cos(c+dx)} dx \right) \right) \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{a(-9Ba^3+24Aba^2+156b^2Ba)}{\cos(c+dx)} dx \right) \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{a(-9Ba^3+24Aba^2+156b^2Ba)}{\sin(c+dx)} dx \right) \right) \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \int \frac{a(-9Ba^3+24Aba^2+156b^2Ba)}{\sin(c+dx)} dx \right) \right) \right)$$

↓ 3477

3.604. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-9a^3B+24a^2Ab+156ab^2)}{\dots} \right) \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(9a^3B-6a^2b(4A+B)-4ab)}{\dots} \right) \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{bd \sqrt{\cos(c+dx)}} - \frac{a(-9a^3B+24a^2Ab+156ab^2)}{\dots} \right) \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{3(-3a^2B+8aAb+12b^2B) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(-9a^3B+24a^2Ab+156ab^2B)}{bd} \right) \right) \right)$$

input `Int[((a + b*cos[c + d*x])^(3/2)*(A + B*cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])^(5/2)*Sin[c + d*x])/(4*b*d) + (((8*A*b - 3*a*B)*Sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + ((3*(8*a*A*b - 3*a^2*B + 12*b^2*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*Sqrt[a + b]*(24*a^2*A*b + 128*A*b^3 - 9*a^3*B + 156*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(9*a^3*B - 6*a^2*b*(4*A + B) - 8*b^3*(16*A + 9*B) - 4*a*b^2*(28*A + 39*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (6*Sqrt[a + b]*(8*a^3*A*b - 96*a*A*b^3 - 3*a^4*B - 24*a^2*b^2*B - 48*b^4*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/b + ((24*a^2*A*b + 128*A*b^3 - 9*a^3*B + 156*a*b^2*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/4)/6)/(8*b))`

3.604.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]], x_Symbol] :> Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B))*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}^{3/2}}{(b \sin(e + fx) + f x)} dx$$

$$\rightarrow \text{Simp}[-2 A (c - d) (\tan[e + f x] / (f b c^2)) \text{Rt}[(c + d) / b, 2] \sqrt{c ((1 + \text{Csc}[e + f x]) / (c - d))} \sqrt{c ((1 - \text{Csc}[e + f x]) / (c + d))} \text{EllipticE}[\text{ArcSin}[\sqrt{c + d \sin[e + f x]}] / \sqrt{b \sin[e + f x]}] / \text{Rt}[(c + d) / b, 2], -(c + d) / (c - d)], x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\& \text{PosQ}[(c + d) / b]$$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}^{3/2}}{(a + b \sin(e + fx))} dx$$

$$\rightarrow \text{Simp}[(A - B) / (a - b) \text{Int}[1 / (\sqrt{a + b \sin[e + f x]} \sqrt{c + d \sin[e + f x]}), x], x] - \text{Simp}[(A b - a B) / (a - b) \text{Int}[(1 + \sin[e + f x]) / ((a + b \sin[e + f x])^2) \sqrt{c + d \sin[e + f x]}], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[A, B]$$`

rule 3528 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +

$$\int (a + b \sin(e + fx))^m ((c + d \sin(e + fx))^n ((A + B \sin(e + fx)) + (C \sin(e + fx) + f x)^2) dx$$

$$\rightarrow \text{Simp}[(-C) \cos[e + f x] (a + b \sin[e + f x])^m ((c + d \sin[e + f x])^{n+1} / (d f (m + n + 2))), x] + \text{Simp}[1 / (d (m + n + 2)) \text{Int}[(a + b \sin[e + f x])^{m-1} (c + d \sin[e + f x])^n \text{Simp}[a A d (m + n + 2) + C (b c m + a d (n + 1)) + (d (A b + a B) (m + n + 2) - C (a c - b d (m + n + 1))) \sin[e + f x] + (C (a d m - b c (m + 1)) + b B d (m + n + 2)) \sin^2[e + f x], x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[m, 0] \&\& (!\text{IGtQ}[n, 0] \&\& (!\text{IntegerQ}[m] || (\text{EqQ}[a, 0] \&\& \text{NeQ}[c, 0])))$$`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2)/

$$\int \frac{(A + B \sin(e + fx) + C \sin^2(e + fx)) \sqrt{(c + d \sin(e + fx))}^{3/2}}{(a + b \sin(e + fx))} dx$$

$$\rightarrow \text{Simp}[C / b^2 \text{Int}[\sqrt{a + b \sin[e + f x]} / \sqrt{c + d \sin[e + f x]}, x], x] + \text{Simp}[1 / b^2 \text{Int}[(A b^2 - a^2 C + b (b B - 2 a C) \sin[e + f x]) / ((a + b \sin[e + f x])^2 \sqrt{c + d \sin[e + f x]}), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$$`


```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])]*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.604.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5424 vs. $2(664) = 1328$.

Time = 21.10 (sec) , antiderivative size = 5425, normalized size of antiderivative = 7.43

method	result	size
parts	Expression too large to display	5425
default	Expression too large to display	5487

```
input int((a+cos(d*x+c)*b)^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.604.5 Fracas [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algo
rithm="fracas")
```

```
output Timed out
```

3.604. $\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx$

3.604.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.604.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sec(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/sec(d*x + c)^(3/2), x)`

3.604.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{3/2}}{\sec(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(3/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algo rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(3/2)/sec(d*x + c)^(3/2), x)`

3.604. $\int \frac{(a+b \cos(c+dx))^{3/2} (A+B \cos(c+dx))}{\sec^{3/2}(c+dx)} dx$

3.604.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{3/2} (A + B \cos(c + dx))}{\sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{3/2}}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(3/2))/(1/cos(c + d*x))^(3/2), x)`

3.605 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{13/2} (c+dx) dx$

3.605.1 Optimal result	5525
3.605.2 Mathematica [B] (warning: unable to verify)	5526
3.605.3 Rubi [A] (verified)	5527
3.605.4 Maple [F(-1)]	5534
3.605.5 Fricas [F]	5535
3.605.6 Sympy [F(-1)]	5535
3.605.7 Maxima [F]	5535
3.605.8 Giac [F]	5536
3.605.9 Mupad [F(-1)]	5536

3.605.1 Optimal result

Integrand size = 35, antiderivative size = 662

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2} (c + dx) dx = \frac{2(a - b)\sqrt{a + b}(3705a^4Ab + 255a^2Ab^3 + 40Ab^5 + 1617a^5B + 3069a^3b^2B - 110ab^4B) \sqrt{\cos(c + dx)}}{3465a^4d\sqrt{\sec(c + dx)}} + \frac{2(a - b)\sqrt{a + b}(40Ab^4 + 3a^4(225A - 539B) - 6a^3b(505A - 209B) + 15a^2b^2(19A - 121B) + 10ab^3(3A - 5B)) \sqrt{\cos(c + dx)}}{3465a^3d\sqrt{\sec(c + dx)}} + \frac{2(675a^4A + 1025a^2Ab^2 - 20Ab^4 + 1793a^3bB + 55ab^3B) \sqrt{a + b \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{3465a^2d} + \frac{2(1145a^2Ab + 15Ab^3 + 539a^3B + 825ab^2B) \sqrt{a + b \cos(c + dx)} \sec^{5/2}(c + dx) \sin(c + dx)}{3465ad} + \frac{2(81a^2A + 113Ab^2 + 209abB) \sqrt{a + b \cos(c + dx)} \sec^{7/2}(c + dx) \sin(c + dx)}{693d} + \frac{2a(14Ab + 11aB) \sqrt{a + b \cos(c + dx)} \sec^{9/2}(c + dx) \sin(c + dx)}{99d} + \frac{2aA(a + b \cos(c + dx))^{3/2} \sec^{11/2}(c + dx) \sin(c + dx)}{11d}$$

output $\frac{2}{11} a A (a + b \cos(dx + c))^{3/2} \sec(dx + c)^{11/2} \sin(dx + c) / d + \frac{2}{3465} (675 A a^4 + 1025 A a^2 b^2 - 20 A b^4 + 1793 B a^3 b + 55 B a b^3) \sec(dx + c)^{3/2} \sin(dx + c) (a + b \cos(dx + c))^{1/2} / a^2 + \frac{2}{3465} (1145 A a^2 b + 15 A b^3 + 539 B a^3 + 825 B a b^2) \sec(dx + c)^{5/2} \sin(dx + c) (a + b \cos(dx + c))^{1/2} / a + \frac{2}{693} (81 A a^2 + 113 A b^2 + 209 B a b) \sec(dx + c)^{7/2} \sin(dx + c) (a + b \cos(dx + c))^{1/2} / d + \frac{2}{99} a (14 A b + 11 B a) \sec(dx + c)^{9/2} \sin(dx + c) (a + b \cos(dx + c))^{1/2} / d + \frac{2}{3465} (a - b) (3705 A a^4 b + 255 A a^2 b^3 + 40 A b^5 + 1617 B a^5 + 3069 B a^3 b^2 - 110 B a b^4) \operatorname{csc}(dx + c) \operatorname{EllipticE}((a + b \cos(dx + c))^{1/2} / (a + b)^{1/2} / \cos(dx + c)^{1/2}, ((-a - b) / (a - b))^{1/2}) (a + b)^{1/2} \cos(dx + c)^{1/2} (a (1 - \sec(dx + c)) / (a + b))^{1/2} (a (1 + \sec(dx + c)) / (a - b))^{1/2} / a^4 / d / \sec(dx + c)^{1/2} + \frac{2}{3465} (a - b) (40 A b^4 + 3 a^4 (225 A - 539 B) - 6 a^3 b (505 A - 209 B) + 15 a^2 b^2 (19 A - 121 B) + 10 a b^3 (3 A - 11 B)) \operatorname{csc}(dx + c) \operatorname{EllipticF}((a + b \cos(dx + c))^{1/2} / (a + b)^{1/2} / \cos(dx + c)^{1/2}, ((-a - b) / (a - b))^{1/2}) (a + b)^{1/2} \cos(dx + c)^{1/2} (a (1 - \sec(dx + c)) / (a + b))^{1/2} (a (1 + \sec(dx + c)) / (a - b))^{1/2} / a^3 / d / \sec(dx + c)^{1/2}$

3.605.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 4198 vs. $2(662) = 1324$.

Time = 24.74 (sec) , antiderivative size = 4198, normalized size of antiderivative = 6.34

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Result too large to show}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(13/2),x]`

output $(\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}*((2*(3705*a^4*A*b + 255*a^2*A*b^3 + 40*A*b^5 + 1617*a^5*B + 3069*a^3*b^2*B - 110*a*b^4*B)*\sin[c + dx]) / (3465*a^3) + (2*\sec[c + dx]^4*(23*a*A*b*\sin[c + dx] + 11*a^2*B*\sin[c + dx]))/99 + (2*\sec[c + dx]^3*(81*a^2*A*\sin[c + dx] + 113*A*b^2*\sin[c + dx] + 209*a*b*B*\sin[c + dx]))/693 + (2*\sec[c + dx]^2*(1145*a^2*A*b*\sin[c + dx] + 15*A*b^3*\sin[c + dx] + 539*a^3*B*\sin[c + dx] + 825*a*b^2*B*\sin[c + dx]))/(3465*a) + (2*\sec[c + dx]*(675*a^4*A*\sin[c + dx] + 1025*a^2*A*b^2*\sin[c + dx] - 20*A*b^4*\sin[c + dx] + 1793*a^3*b*B*\sin[c + dx] + 55*a*b^3*B*\sin[c + dx]))/(3465*a^2) + (2*a^2*A*\sec[c + dx]^4*\tan[c + dx])/11)/d + (2*((-247*a^2*A*b)/(231*\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}) - (17*A*b^3)/(231*\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}) - (8*A*b^5)/(693*a^2*\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}) - (7*a^3*B)/(15*\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}) - (31*a*b^2*B)/(35*\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}) + (2*b^4*B)/(63*a*\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}) + (15*a^3*A*\sqrt{\sec[c + dx]})/(77*\sqrt{a + b\cos[c + dx]}) - (26*a*A*b^2*\sqrt{\sec[c + dx]})/(231*\sqrt{a + b\cos[c + dx]}) - (7*A*b^4*\sqrt{\sec[c + dx]})/(99*a*\sqrt{a + b\cos[c + dx]}) - (8*A*b^6*\sqrt{\sec[c + dx]})/(693*a^3*\sqrt{a + b\cos[c + dx]}) + (38*a^2*b*B*\sqrt{\sec[c + dx]})/(105*\sqrt{a + b\cos[c + dx]}) - (124*b^3*B*\sqrt{\sec[c + dx]})/(315*\sqrt{a + b\cos[c + dx]}) + (2*b^5*B*\sqrt{\sec[c + dx]})...$

3.605.3 Rubi [A] (verified)

Time = 3.67 (sec) , antiderivative size = 659, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.629$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{13}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{13/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx))}{\cos^{\frac{13}{2}}(c + dx)} dx$$

3.605. $\int (a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{\frac{13}{2}}(c + dx) dx$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{13/2}} dx \\ & \downarrow 3468 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2}{11} \int \frac{\sqrt{a+b\cos(c+dx)}(b(6aA+11bB)\cos^2(c+dx)+(9Aa^2+22bBa+11Ab^2)\cos(c+dx))}{2\cos^{\frac{11}{2}}(c+dx)} dx \right. \\ & \downarrow 27 \\ & \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \int \frac{\sqrt{a+b\cos(c+dx)}(b(6aA+11bB)\cos^2(c+dx)+(9Aa^2+22bBa+11Ab^2)\cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx \right) \right. \\ & \downarrow 3042 \\ & \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(b(6aA+11bB)\sin(c+dx+\frac{\pi}{2})^2+(9Aa^2+22bBa+11Ab^2)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{11/2}} dx \right) \right. \\ & \downarrow 3526 \\ & \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \left(\frac{2}{9} \int \frac{3b(22Ba^2+46Aba+33b^2B)\cos^2(c+dx)+(77Ba^3+233Aba^2+297b^2B)\cos(c+dx)}{2\cos^{\frac{9}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right. \right. \\ & \downarrow 27 \\ & \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \int \frac{3b(22Ba^2+46Aba+33b^2B)\cos^2(c+dx)+(77Ba^3+233Aba^2+297b^2B)\cos(c+dx)}{\cos^{\frac{9}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right) \right. \right. \\ & \downarrow 3042 \\ & \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \int \frac{3b(22Ba^2+46Aba+33b^2B)\sin(c+dx+\frac{\pi}{2})^2+(77Ba^3+233Aba^2+297b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{9/2}\sqrt{a+b\cos(c+dx)}} dx \right) \right) \right. \right. \\ & \downarrow 3534 \\ & \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \left(\frac{2 \int \frac{4ab(81Aa^2+209bBa+113Ab^2)\cos^2(c+dx)+a(405Aa^3+1507bBa^2+1531Ab^2a+693b^3B)\cos(c+dx)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{7a} \right) \right) \right) \right. \right. \end{aligned}$$

3.605. $\int (a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sec^{\frac{13}{2}}(c+dx) dx$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{\int \frac{4ab(81Aa^2+209bBa+113Ab^2)\cos^2(c+dx)+a(405Aa^3+1507bBa^2+1531Ab^2a+693b^3B)\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{7a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{\int \frac{4ab(81Aa^2+209bBa+113Ab^2)\sin(c+dx+\frac{\pi}{2})^2+a(405Aa^3+1507bBa^2+1531Ab^2a+693b^3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{7a}\right)\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{2\int \frac{(1617Ba^3+5055Aba^2+6655b^2Ba+2305Ab^3)\cos(c+dx)a^2+2b(539Ba^3+1145Aba^2+825b^2Ba+15Ab^3)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{5a}\right)\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{\int \frac{(1617Ba^3+5055Aba^2+6655b^2Ba+2305Ab^3)\cos(c+dx)a^2+2b(539Ba^3+1145Aba^2+825b^2Ba+15Ab^3)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{5a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{\int \frac{(1617Ba^3+5055Aba^2+6655b^2Ba+2305Ab^3)\sin(c+dx+\frac{\pi}{2})a^2+2b(539Ba^3+1145Aba^2+825b^2Ba+15Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{5a}\right)\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\int \frac{3\left(\left(675Aa^4+2871bBa^3+3315Ab^2a^2+1705b^3Ba+10Ab^4\right)\cos(c+dx)a^2+\left(1617Ba^5+3705Aba^4+3069b^2Ba^3+\right.\right.}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \frac{1}{3a} dx \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\int \frac{\left(675Aa^4+2871bBa^3+3315Ab^2a^2+1705b^3Ba+10Ab^4\right)\cos(c+dx)a^2+\left(1617Ba^5+3705Aba^4+3069b^2Ba^3+\right.}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \frac{1}{a} dx \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\int \frac{\left(675Aa^4+2871bBa^3+3315Ab^2a^2+1705b^3Ba+10Ab^4\right)\sin\left(c+dx+\frac{\pi}{2}\right)a^2+\left(1617Ba^5+3705Aba^4+3069b^2Ba^3+\right.}{\sin\left(c+dx+\frac{\pi}{2}\right)^{\frac{3}{2}}\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}} \frac{1}{a} dx \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\int \frac{a(a-b)\left(3a^4(225A-539B)-6a^3b(505A-209B)+15a^2b^2(19A-121B)+10ab^3(3A-11B)+40Ab^4\right)}{\sqrt{\cos(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\frac{a(a-b)(3a^4(225A-539B)-6a^3b(505A-209B)+15a^2b^2(19A-121B)+10ab^3(3A-11B)+40Ab^4)}{\sqrt{\sin(c+dx)}} \int \frac{1}{\sqrt{\sin(c+dx)}} dx \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\frac{a(1617a^5B+3705a^4Ab+3069a^3b^2B+255a^2Ab^3-110ab^4B+40Ab^5)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \left(\frac{2(81a^2A+209abB+113Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{7d\cos^{\frac{7}{2}}(c+dx)} + \frac{2(539a^3B+113a^2bB+113ab^2)}{7d\cos^{\frac{7}{2}}(c+dx)} \right)$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(13/2),x]`

```

output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + b*Cos[c + d*x])^(3/2)*S
in[c + d*x])/(11*d*Cos[c + d*x]^(11/2)) + ((2*a*(14*A*b + 11*a*B)*Sqrt[a +
b*Cos[c + d*x]]*Sin[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((2*(81*a^2*A +
113*A*b^2 + 209*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c +
d*x]^(7/2)) + ((2*(1145*a^2*A*b + 15*A*b^3 + 539*a^3*B + 825*a*b^2*B)*Sqr
t[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b
)*Sqrt[a + b]*(3705*a^4*A*b + 255*a^2*A*b^3 + 40*A*b^5 + 1617*a^5*B + 3069
*a^3*b^2*B - 110*a*b^4*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c +
d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 -
Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*
(a - b)*Sqrt[a + b]*(40*A*b^4 + 3*a^4*(225*A - 539*B) - 6*a^3*b*(505*A - 2
09*B) + 15*a^2*b^2*(19*A - 121*B) + 10*a*b^3*(3*A - 11*B))*Cot[c + d*x]*El
lipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])],
-((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec
[c + d*x]))/(a - b))]/d)/a + (2*(675*a^4*A + 1025*a^2*A*b^2 - 20*A*b^4 + 1
793*a^3*b*B + 55*a*b^3*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c
+ d*x]^(3/2)))/(5*a)/(7*a))/9)/11)

```

3.605.3.1 Defintions of rubi rules used

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_.) + (f
_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Ssin[e + f*x]]/Sqrt[d*Ssin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]

```

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*SIN[e + f*x])^(m - 2)*(c + d*SIN[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*SIN[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*SIN[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*SIN[e + f*x]]/Sqrt[b*SIN[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*SIN[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + SIN[e + f*x])/((a + b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] :> Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3534 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] :> Simp[-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 - a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) | EqQ[a, 0])))`

3.605.4 Maple [F(-1)]

Timed out.

hanged

input `int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x)`

output `int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x)`

3.605.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{13/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(13/2), x)`

3.605.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(13/2),x)`

output `Timed out`

3.605.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{13/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(13/2), x)`

3.605. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx$

3.605.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{13/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(13/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(13/2), x)`

3.605.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{13/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{13/2} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(13/2)*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(13/2)*(a + b*cos(c + d*x))^(5/2), x)`

3.606 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$

3.606.1 Optimal result	5537
3.606.2 Mathematica [A] (warning: unable to verify)	5538
3.606.3 Rubi [A] (verified)	5539
3.606.4 Maple [F(-1)]	5545
3.606.5 Fricas [F]	5545
3.606.6 Sympy [F(-1)]	5546
3.606.7 Maxima [F]	5546
3.606.8 Giac [F]	5546
3.606.9 Mupad [F(-1)]	5547

3.606.1 Optimal result

Integrand size = 35, antiderivative size = 562

$$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx =$$

$$\frac{2(a-b)\sqrt{a+b}(147a^4A+279a^2Ab^2-10Ab^4+435a^3bB+45ab^3B)\sqrt{\cos(c+dx)}\csc(c+dx)E(\frac{c+dx}{2}, \sqrt{\frac{a+b\cos(c+dx)}{a+b}})}{315a^3d\sqrt{\sec(c+dx)}} +$$

$$\frac{2(a-b)\sqrt{a+b}(10Ab^3-6a^2b(19A-60B)+3a^3(49A-25B)+15ab^2(11A-3B))\sqrt{\cos(c+dx)}\csc(c+dx)}{315a^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{2(163a^2Ab+5Ab^3+75a^3B+135ab^2B)\sqrt{a+b\cos(c+dx)}\sec^{\frac{3}{2}}(c+dx)\sin(c+dx)}{315ad} +$$

$$\frac{2(49a^2A+75Ab^2+135abB)\sqrt{a+b\cos(c+dx)}\sec^{\frac{5}{2}}(c+dx)\sin(c+dx)}{315d} +$$

$$\frac{2a(4Ab+3aB)\sqrt{a+b\cos(c+dx)}\sec^{\frac{7}{2}}(c+dx)\sin(c+dx)}{21d} +$$

$$\frac{2aA(a+b\cos(c+dx))^{3/2}\sec^{\frac{9}{2}}(c+dx)\sin(c+dx)}{9d}$$

output $\frac{2}{9}aA(a+b\cos(dx+c))^{3/2}\sec(dx+c)^{9/2}\sin(dx+c)/d+2/315(163Aa^2b+5Aab^3+75B^2a^3+135B^2a^2b)\sec(dx+c)^{3/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/a/d+2/315(49Aa^2+75Aab^2+135B^2a^2b)\sec(dx+c)^{5/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/21a(4Aab+3B^2a)\sec(dx+c)^{7/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/315(a-b)(147Aa^4+279Aa^2b^2-10Aab^4+435B^2a^3b+45B^2a^2b^3)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2}/a^3/d/\sec(dx+c)^{1/2}-2/315(a-b)(10Aab^3-6a^2b(19A-60B)+3a^3(49A-25B)+15a^2b^2(11A-3B))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2}/a^2/d/\sec(dx+c)^{1/2}$

3.606.2 Mathematica [A] (warning: unable to verify)

Time = 19.36 (sec) , antiderivative size = 629, normalized size of antiderivative = 1.12

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \frac{2\sqrt{\cos^2(\frac{1}{2}(c + dx)) \sec(c + dx)} \left(-2(a + b) (147a^4A + 279a^2Ab^2 - 10Ab^4 + 435a^3bB + 45ab^3B) \right)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} \left(\frac{2(147a^4A + 279a^2Ab^2 - 10Ab^4 + 435a^3bB + 45ab^3B) \sin(c + dx)}{315a^2} + \frac{2}{63} \sec^3(c + dx) (19aAb + 15ab^2B) \right)$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

output $(2\sqrt{\cos[(c+dx)/2]^2 \sec[c+dx]}(-2(a+b)(147a^4A+279a^2Ab^2-10Ab^4+435a^3bB+45ab^3B))\sqrt{\cos[c+dx]/(1+\cos[c+dx])}\sqrt{(a+b\cos[c+dx])/((a+b)(1+\cos[c+dx]))}\text{EllipticE}[\text{ArcSin}[\text{Tan}[(c+dx)/2]], (-a+b)/(a+b)]+2a(a+b)(-10Ab^3+15a^2b(11A+3B)+3a^3(49A+25B)+6a^2b(19A+60B))\sqrt{\cos[c+dx]/(1+\cos[c+dx])}\sqrt{(a+b\cos[c+dx])/((a+b)(1+\cos[c+dx]))}\text{EllipticF}[\text{ArcSin}[\text{Tan}[(c+dx)/2]], (-a+b)/(a+b)]-(147a^4A+279a^2Ab^2-10Ab^4+435a^3bB+45ab^3B)\cos[c+dx](a+b\cos[c+dx])\sec[(c+dx)/2]^2\tan[(c+dx)/2])/(315a^2d\sqrt{(a+b\cos[c+dx])}\sqrt{\sec[(c+dx)/2]^2})+(\sqrt{a+b\cos[c+dx]}\sqrt{\sec[c+dx]}((2(147a^4A+279a^2Ab^2-10Ab^4+435a^3bB+45ab^3B)\sin[c+dx])/(315a^2)+(2\sec[c+dx]^3(19aAb\sin[c+dx]+9a^2B\sin[c+dx]))/63+(2\sec[c+dx]^2(49a^2A\sin[c+dx]+75Ab^2\sin[c+dx]+135abB\sin[c+dx]))/315+(2\sec[c+dx](163a^2Ab\sin[c+dx]+5Ab^3\sin[c+dx]+75a^3B\sin[c+dx]+135ab^2B\sin[c+dx]))/(315a)+(2a^2A\sec[c+dx]^3\tan[c+dx])/9))/d$

3.606.3 Rubi [A] (verified)

Time = 2.87 (sec) , antiderivative size = 554, normalized size of antiderivative = 0.99, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{11}{2}}(c+dx)(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c+dx+\frac{\pi}{2}\right)^{11/2}\left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2}\left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\cos^{\frac{11}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

3.606. $\int (a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sec^{\frac{11}{2}}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2} (A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{11/2}} dx$$

↓ 3468

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2}{9} \int \frac{\sqrt{a+b\cos(c+dx)}(b(4aA+9bB)\cos^2(c+dx) + (7Aa^2+18bBa+9Ab^2)\cos(c+dx))}{2\cos^{\frac{9}{2}}(c+dx)} dx \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \int \frac{\sqrt{a+b\cos(c+dx)}(b(4aA+9bB)\cos^2(c+dx) + (7Aa^2+18bBa+9Ab^2)\cos(c+dx))}{\cos^{\frac{9}{2}}(c+dx)} dx \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})} (b(4aA+9bB)\sin(c+dx+\frac{\pi}{2})^2 + (7Aa^2+18bBa+9Ab^2)\cos(c+dx))}{\sin(c+dx+\frac{\pi}{2})^{9/2}} dx \right)$$

↓ 3526

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{2}{7} \int \frac{b(36Ba^2+76Aba+63b^2B)\cos^2(c+dx) + (45Ba^3+137Aba^2+189b^2Ba-189b^3B)\cos(c+dx)}{2\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{1}{7} \int \frac{b(36Ba^2+76Aba+63b^2B)\cos^2(c+dx) + (45Ba^3+137Aba^2+189b^2Ba-189b^3B)\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{1}{7} \int \frac{b(36Ba^2+76Aba+63b^2B)\sin(c+dx+\frac{\pi}{2})^2 + (45Ba^3+137Aba^2+189b^2Ba-189b^3B)\cos(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{a+b\cos(c+dx)}} dx \right) \right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{1}{7} \left(2 \int \frac{2ab(49Aa^2+135bBa+75Ab^2)\cos^2(c+dx) + a(147Aa^3+585bBa^2+605Ab^2a+315b^3B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \right) \right) \right)$$

↓ 27

3.606. $\int (a+b\cos(c+dx))^{5/2} (A+B\cos(c+dx)) \sec^{\frac{11}{2}}(c+dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{\int \frac{2ab(49Aa^2+135bBa+75Ab^2)\cos^2(c+dx)+a(147Aa^3+585bBa^2+605Ab^2a+315b^3B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{5a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{\int \frac{2ab(49Aa^2+135bBa+75Ab^2)\sin(c+dx+\frac{\pi}{2})^2+a(147Aa^3+585bBa^2+605Ab^2a+315b^3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{5a}\right)\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{2\int \frac{3((75Ba^3+261Aba^2+405b^2Ba+155Ab^3)\cos(c+dx)a^2+(147Aa^4+435bBa^3+279Ab^2a^2+45b^3Ba-10Ab^4)a}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{3a}}}{5a}\right)\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{\int \frac{(75Ba^3+261Aba^2+405b^2Ba+155Ab^3)\cos(c+dx)a^2+(147Aa^4+435bBa^3+279Ab^2a^2+45b^3Ba-10Ab^4)a}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{a}}}{5a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{\int \frac{(75Ba^3+261Aba^2+405b^2Ba+155Ab^3)\sin(c+dx+\frac{\pi}{2})a^2+(147Aa^4+435bBa^3+279Ab^2a^2+45b^3Ba-10Ab^4)a}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{a}}}{5a}\right)\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{9}\left(\frac{1}{7}\left(\frac{a(147a^4A+435a^3bB+279a^2Ab^2+45ab^3B-10Ab^4)\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-a(a-b)(3a^3(4a^2+3ab+2b^2))}{a}}}{5a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{1}{7} \left(\frac{a(147a^4A+435a^3bB+279a^2Ab^2+45ab^3B-10Ab^4) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx - a(a-2b \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx - \frac{2(a-2b \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx} \right) \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{1}{7} \left(\frac{a(147a^4A+435a^3bB+279a^2Ab^2+45ab^3B-10Ab^4) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx - a(a-2b \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx - \frac{2(a-2b \sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})} dx} \right) \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{9} \left(\frac{1}{7} \left(\frac{2(49a^2A + 135abB + 75Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5d \cos^{\frac{5}{2}}(c+dx)} + \frac{2(75a^3B+163a^2)}{\dots} \right) \right) \right)$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(11/2),x]`

```

output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + b*Cos[c + d*x])^(3/2)*S
in[c + d*x])/(9*d*Cos[c + d*x]^(9/2)) + ((6*a*(4*A*b + 3*a*B)*Sqrt[a + b*C
os[c + d*x]]*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*(49*a^2*A + 75*A
*b^2 + 135*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]
^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(147*a^4*A + 279*a^2*A*b^2 - 10*A*b^4 +
435*a^3*b*B + 45*a*b^3*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c
+ d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1
- Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2
*(a - b)*Sqrt[a + b]*(10*A*b^3 - 6*a^2*b*(19*A - 60*B) + 3*a^3*(49*A - 25*
B) + 15*a*b^2*(11*A - 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c
+ d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1
- Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/a + (2
*(163*a^2*A*b + 5*A*b^3 + 75*a^3*B + 135*a*b^2*B)*Sqrt[a + b*Cos[c + d*x]]
*Ssin[c + d*x])/(d*Cos[c + d*x]^(3/2)))/(5*a))/7)/9)

```

3.606.3.1 Defintions of rubi rules used

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f
_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]

```

```

rule 3440 Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*
(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])

```

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.606.4 Maple [F(-1)]

Timed out.

hanged

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x)
```

```
output int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x)
```

3.606.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{11/2} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, alg
orithm="fricas")
```

```
output integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2
+ (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(1
1/2), x)
```

3.606. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx$

3.606.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(11/2),x)`

output `Timed out`

3.606.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{11/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(11/2), x)`

3.606.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{11/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(11/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(11/2), x)`

3.606. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{11/2}(c + dx) dx$

3.606.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{11}{2}}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{11/2} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(11/2)*(a + b*cos(c + d*x))^(5/2), x)`

3.607 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{9/2}(c+dx) dx$

3.607.1 Optimal result	5548
3.607.2 Mathematica [A] (warning: unable to verify)	5549
3.607.3 Rubi [A] (verified)	5550
3.607.4 Maple [F(-1)]	5555
3.607.5 Fricas [F]	5555
3.607.6 Sympy [F(-1)]	5556
3.607.7 Maxima [F]	5556
3.607.8 Giac [F]	5556
3.607.9 Mupad [F(-1)]	5557

3.607.1 Optimal result

Integrand size = 35, antiderivative size = 474

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx =$$

$$\frac{2(a - b)\sqrt{a + b}(145a^2Ab + 15Ab^3 + 63a^3B + 161ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}}\right)\right)}{105a^2d\sqrt{\sec(c + dx)}} +$$

$$\frac{2(a - b)\sqrt{a + b}(a^2(25A - 63B) + 15b^2(A - 7B) - 8ab(15A - 7B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}}\right)\right)}{105ad\sqrt{\sec(c + dx)}} +$$

$$\frac{2(25a^2A + 45Ab^2 + 77abB) \sqrt{a + b \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{105d} +$$

$$\frac{2a(10Ab + 7aB) \sqrt{a + b \cos(c + dx)} \sec^{5/2}(c + dx) \sin(c + dx)}{35d} +$$

$$\frac{2aA(a + b \cos(c + dx))^{3/2} \sec^{7/2}(c + dx) \sin(c + dx)}{7d}$$

output $\frac{2}{7}aA(a+b\cos(dx+c))^{3/2}\sec(dx+c)^{7/2}\sin(dx+c)/d+2/105(25Aa^2+45Ab^2+77Bab)\sec(dx+c)^{3/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/35a(10Ab+7Ba)\sec(dx+c)^{5/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/105(a-b)(145Aa^2b+15Ab^3+63Ba^3+161Bab^2)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a^2d/\sec(dx+c)^{1/2}+2/105(a-b)(a^2(25A-63B)+15b^2(A-7B)-8ab(15A-7B))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a/d/\sec(dx+c)^{1/2}$

3.607.2 Mathematica [A] (warning: unable to verify)

Time = 17.20 (sec) , antiderivative size = 524, normalized size of antiderivative = 1.11

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^2(c + dx) dx = \frac{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right) \sec(c + dx)} \left(-2(a + b)(145a^2Ab + 15Ab^3 + 63a^3B + 161ab^2B)\sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}}\right)}{105a} + \frac{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2(145a^2Ab + 15Ab^3 + 63a^3B + 161ab^2B) \sin(c + dx)}{105a} + \frac{2}{35} \sec^2(c + dx) (15aAb \sin(c + dx) + \dots)\right)}{105a}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

output $(2\sqrt{\cos[(c + dx)/2]^2 \sec[c + dx]} * (-2(a + b)(145a^2Ab + 15A^3b^3 + 63a^3B + 161ab^2B)) * \sqrt{\cos[c + dx]/(1 + \cos[c + dx])} * \sqrt{(a + b\cos[c + dx])/((a + b)(1 + \cos[c + dx]))} * \text{EllipticE}[\text{ArcSin}[\text{Tan}[(c + dx)/2]]], (-a + b)/(a + b) + 2a(a + b)(15b^2(A + 7B) + 8ab(15A + 7B) + a^2(25A + 63B)) * \sqrt{\cos[c + dx]/(1 + \cos[c + dx])} * \sqrt{(a + b\cos[c + dx])/((a + b)(1 + \cos[c + dx]))} * \text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + dx)/2]]], (-a + b)/(a + b) - (145a^2Ab + 15A^3b^3 + 63a^3B + 161ab^2B) * \cos[c + dx] * (a + b\cos[c + dx]) * \sec[(c + dx)/2]^2 * \text{Tan}[(c + dx)/2]) / (105ad\sqrt{a + b\cos[c + dx]} * \sqrt{\sec[(c + dx)/2]^2} + (\sqrt{a + b\cos[c + dx]} * \sqrt{\sec[c + dx]} * ((2(145a^2Ab + 15A^3b^3 + 63a^3B + 161ab^2B) * \sin[c + dx]) / (105a) + (2\sec[c + dx]^2(15aAb\sin[c + dx] + 7a^2B\sin[c + dx])) / 35 + (2\sec[c + dx] * (25a^2A\sin[c + dx] + 45Ab^2\sin[c + dx] + 77abB\sin[c + dx])) / 105 + (2a^2A\sec[c + dx]^2 * \text{Tan}[c + dx]) / 7) / d$

3.607.3 Rubi [A] (verified)

Time = 2.30 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.98, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3526, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{9}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{9/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{9/2}} dx$$

$$\downarrow \text{3468}$$

3.607. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2}{7}\int\frac{\sqrt{a+b\cos(c+dx)}(b(2aA+7bB)\cos^2(c+dx)+(5Aa^2+14bBa+7Ab^2)\cos(c+dx))}{2\cos^{\frac{7}{2}}(c+dx)}dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{\sqrt{a+b\cos(c+dx)}(b(2aA+7bB)\cos^2(c+dx)+(5Aa^2+14bBa+7Ab^2)\cos(c+dx))}{\cos^{\frac{7}{2}}(c+dx)}dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\int\frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(b(2aA+7bB)\sin(c+dx+\frac{\pi}{2})^2+(5Aa^2+14bBa+7Ab^2)\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{7/2}}dx\right)$$

↓ 3526

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{2}{5}\int\frac{b(14Ba^2+30Aba+35b^2B)\cos^2(c+dx)+(21Ba^3+65Aba^2+105b^2Ba+105b^3B)\cos(c+dx)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\int\frac{b(14Ba^2+30Aba+35b^2B)\cos^2(c+dx)+(21Ba^3+65Aba^2+105b^2Ba+105b^3B)\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\int\frac{b(14Ba^2+30Aba+35b^2B)\sin(c+dx+\frac{\pi}{2})^2+(21Ba^3+65Aba^2+105b^2Ba+105b^3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(2\int\frac{a(63Ba^3+145Aba^2+161b^2Ba+15Ab^3)+a(25Aa^3+119bBa^2+135Ab^2a+105b^3B)\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\int\frac{a(63Ba^3+145Aba^2+161b^2Ba+15Ab^3)+a(25Aa^3+119bBa^2+135Ab^2a+105b^3B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)\right)$$

3.607. $\int(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sec^{\frac{9}{2}}(c+dx)dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\int\frac{a(63Ba^3+145Aba^2+161b^2Ba+15Ab^3)+a(25Aa^3+119bBa^2+135Ab^2a+105b^3B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}\right)\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{a(a-b)(a^2(25A-63B)-8ab(15A-7B)+15b^2(A-7B))\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}\right)}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{a(a-b)(a^2(25A-63B)-8ab(15A-7B)+15b^2(A-7B))\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}\right)}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{a(63a^3B+145a^2Ab+161ab^2B+15Ab^3)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}\right)}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2(25a^2A+77abB+45Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3d\cos^{\frac{3}{2}}(c+dx)}+\frac{2(a-b)\sqrt{a+b}(a^2)}{\sin(c+dx+\frac{\pi}{2})}\right)\right)\right)$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(9/2),x]`

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(7*d*Cos[c + d*x]^(7/2)) + ((2*a*(10*A*b + 7*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*d*Cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(145*a^2*A*b + 15*A*b^3 + 63*a^3*B + 161*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*(a - b)*Sqrt[a + b]*(a^2*(25*A - 63*B) + 15*b^2*(A - 7*B) - 8*a*b*(15*A - 7*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(3*a) + (2*(25*a^2*A + 45*A*b^2 + 77*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*d*Cos[c + d*x]^(3/2)))/5/7)
```

3.607.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]
```

```
rule 3440 Int[(csc[(e_)] + (f_)*(x_))*(g_.)^(p_)*((a_.) + (b_)*sin[(e_)] + (f_)*(x_))^(m_)*((c_) + (d_)*sin[(e_)] + (f_)*(x_))^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])
```


rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[-(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.607.4 Maple [F(-1)]

Timed out.

hanged

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x)
```

```
output int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x)
```

3.607.5 Fracas [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{\frac{5}{2}} \sec(dx + c)^{\frac{9}{2}} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algo
rithm="fricas")
```

```
output integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2
+ (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(9
/2), x)
```

$$3.607. \quad \int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{9}{2}}(c + dx) dx$$

3.607.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(9/2),x)`

output `Timed out`

3.607.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{9/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(9/2), x)`

3.607.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{9/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(9/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(9/2), x)`

3.607.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{9/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{9/2} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(9/2)*(a + b*cos(c + d*x))^(5/2), x)`

3.608 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{7/2}(c+dx) dx$

3.608.1 Optimal result	5558
3.608.2 Mathematica [B] (warning: unable to verify)	5559
3.608.3 Rubi [A] (verified)	5559
3.608.4 Maple [F(-1)]	5564
3.608.5 Fracas [F]	5565
3.608.6 Sympy [F(-1)]	5565
3.608.7 Maxima [F]	5565
3.608.8 Giac [F]	5566
3.608.9 Mupad [F(-1)]	5566

3.608.1 Optimal result

Integrand size = 35, antiderivative size = 553

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \frac{2(a - b)\sqrt{a + b}(9a^2A + 23Ab^2 + 35abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right) \mid -\right)}{15ad\sqrt{\sec(c + dx)}} + \frac{2\sqrt{a + b}(15Ab^3 - ab^2(23A - 45B) + a^2b(17A - 35B) - a^3(9A - 5B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), \sqrt{\frac{a + b}{a + b \sqrt{\cos(c + dx)}}}\right)}{15ad\sqrt{\sec(c + dx)}} - \frac{2b^2\sqrt{a + b}B\sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a}{a + b}}}{d\sqrt{\sec(c + dx)}} + \frac{2a(8Ab + 5aB)\sqrt{a + b \cos(c + dx)} \sec^{3/2}(c + dx) \sin(c + dx)}{15d} + \frac{2aA(a + b \cos(c + dx))^{3/2} \sec^{5/2}(c + dx) \sin(c + dx)}{5d}$$

output $\frac{2}{5}aA(a+b\cos(dx+c))^{3/2}\sec(dx+c)^{5/2}\sin(dx+c)/d+2/15a(8Ab+5Ba)\sec(dx+c)^{3/2}\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d+2/15(a-b)(9Aa^2+23Ab^2+35Bab)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2})*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a/d/\sec(dx+c)^{1/2}+2/15(15Ab^3-ab^2(23A-45B)+a^2b(17A-35B)-a^3(9A-5B))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/a/d/\sec(dx+c)^{1/2}-2b^2B\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a(1-\sec(dx+c))/(a+b))^{1/2}*(a(1+\sec(dx+c))/(a-b))^{1/2}/d/\sec(dx+c)^{1/2}$

3.608.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 7032 vs. $2(553) = 1106$.

Time = 24.85 (sec) , antiderivative size = 7032, normalized size of antiderivative = 12.72

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Result too large to show}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `Result too large to show`

3.608.3 Rubi [A] (verified)

Time = 2.48 (sec) , antiderivative size = 520, normalized size of antiderivative = 0.94, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3526, 27, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{7/2}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

3.608. $\int (a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

$$\begin{aligned}
& \int \csc\left(c + dx + \frac{\pi}{2}\right)^{7/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{7/2}(c + dx)} dx \\
& \quad \downarrow \text{3440} \\
& \int \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{7/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2}{5} \int \frac{\sqrt{a + b \cos(c + dx)} (5b^2 B \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + \dots)}{2 \cos^{5/2}(c + dx)} \right) \\
& \quad \downarrow \text{3468} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} \int \frac{\sqrt{a + b \cos(c + dx)} (5b^2 B \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + \dots)}{\cos^{5/2}(c + dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} \int \frac{\sqrt{a + b \cos(c + dx)} (5b^2 B \cos^2(c + dx) + (3Aa^2 + 10bBa + 5Ab^2) \cos(c + dx) + \dots)}{\cos^{5/2}(c + dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (5b^2 B \sin(c + dx + \frac{\pi}{2})^2 + (3Aa^2 + 10bBa + 5Ab^2) \sin(c + dx + \frac{\pi}{2})^{5/2}}{\sin(c + dx + \frac{\pi}{2})^{5/2}} \right) \\
& \quad \downarrow \text{3526} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} \left(\frac{2}{3} \int \frac{15B \cos^2(c + dx) b^3 + a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2)}{2 \cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} \right) \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{15B \cos^2(c + dx) b^3 + a(9Aa^2 + 35bBa + 23Ab^2) + (5Ba^3 + 17Aba^2 + 45b^2)}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} \right) \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.608. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{15B\sin(c+dx+\frac{\pi}{2})^2b^3+a(9Aa^2+35bBa+23Ab^2)+(5Ba^3+17Aba^2-}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+}}$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{a(9Aa^2+35bBa+23Ab^2)+(5Ba^3+17Aba^2+45b^2Ba+15Ab^3)\cos(c+}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx}}\right.\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{a(9Aa^2+35bBa+23Ab^2)+(5Ba^3+17Aba^2+45b^2Ba+15Ab^3)\sin(c+}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2}}}\right.\right.$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\int\frac{a(9Aa^2+35bBa+23Ab^2)+(5Ba^3+17Aba^2+45b^2Ba+15Ab^3)\sin(c+}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2}}}\right.\right.$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(a(9a^2A+35abB+23Ab^2)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+(-a^3(9\right.\right.\right.$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(a(9a^2A+35abB+23Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2}}}\right.\right.\right.$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(a(9a^2A+35abB+23Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2}}}\right.\right.\right.$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2(a-b)\sqrt{a+b}(9a^2A+35abB+23Ab^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a}}}{ad}\right)\right)\right)$$

input `Int[(a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x])*Sec[c + d*x]^(7/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(5*d*cos[c + d*x]^(5/2)) + (((2*(a - b)*Sqrt[a + b]*(9*a^2*A + 23*A*b^2 + 35*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(15*A*b^3 - a*b^2*(23*A - 45*B) + a^2*b*(17*A - 35*B) - a^3*(9*A - 5*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (30*b^2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/3 + (2*a*(8*A*b + 5*a*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(3*d*cos[c + d*x]^(3/2)))/5)`

3.608.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_)*sin[(e_.) + (f_)*(x_)]/Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*sin[e + f*x]]/Sqrt[b*sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

```
rule 3477 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]
```

```
rule 3526 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.608.4 Maple [F(-1)]

Timed out.

hanged

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x)
```

```
output int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x)
```

$$3.608. \quad \int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$$

3.608.5 Fricas [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{7/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="fricas")`

output `integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2 + (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(7/2), x)`

3.608.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(7/2),x)`

output `Timed out`

3.608.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{7/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(7/2), x)`

3.608. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx$

3.608.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{7/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(7/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(7/2), x)`

3.608.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{7/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{7/2} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2)*(a + b*cos(c + d*x))^(5/2), x)`

3.609 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx$

3.609.1 Optimal result	5567
3.609.2 Mathematica [B] (warning: unable to verify)	5568
3.609.3 Rubi [A] (verified)	5568
3.609.4 Maple [F(-1)]	5574
3.609.5 Fricas [F(-1)]	5575
3.609.6 Sympy [F(-1)]	5575
3.609.7 Maxima [F]	5575
3.609.8 Giac [F]	5576
3.609.9 Mupad [F(-1)]	5576

3.609.1 Optimal result

Integrand size = 35, antiderivative size = 596

$$\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{5/2}(c+dx) dx = \frac{(a-b)\sqrt{a+b}(14aAb+6a^2B-3b^2B)\sqrt{\cos(c+dx)}\csc(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) - \frac{a+b}{a-b}\sqrt{a+b}(2ab(7A-9B)-2a^2(A-3B)-3b^2(6A+B))\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) - b\sqrt{a+b}(2Ab+5aB)\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3ad\sqrt{\sec(c+dx)}} + \frac{2a(2Ab+aB)\sqrt{a+b}\cos(c+dx)\sqrt{\sec(c+dx)}\sin(c+dx)}{d} - \frac{(14aAb+6a^2B-3b^2B)\sqrt{a+b}\cos(c+dx)\sqrt{\sec(c+dx)}\sin(c+dx)}{3d} + \frac{2aA(a+b\cos(c+dx))^{3/2}\sec^{3/2}(c+dx)\sin(c+dx)}{3d}$$

output $\frac{2}{3}aA(a+b\cos(dx+c))^{3/2}\sec(dx+c)^{3/2}\sin(dx+c)/d+2a(2Ab+Ba)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/d-1/3(14Aab+6Ba^2-3Bb^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/d+1/3(a-b)(14Aab+6Ba^2-3Bb^2)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/d/\sec(dx+c)^{1/2}-1/3(2ab(7A-9B)-2a^2(A-3B)-3b^2(6A+B))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/d/\sec(dx+c)^{1/2}-b(2Ab+5Ba)\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/d/\sec(dx+c)^{1/2}$

3.609.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 7700 vs. $2(596) = 1192$.

Time = 23.26 (sec) , antiderivative size = 7700, normalized size of antiderivative = 12.92

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Result too large to show}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

output `Result too large to show`

3.609.3 Rubi [A] (verified)

Time = 3.08 (sec) , antiderivative size = 568, normalized size of antiderivative = 0.95, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3526, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{5/2}(c + dx)(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

3.609. $\int (a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \int \csc\left(c + dx + \frac{\pi}{2}\right)^{5/2} \left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2} \left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
& \downarrow \text{3440} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\cos^{5/2}(c + dx)} dx \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{5/2}} dx \\
& \downarrow \text{3468} \\
& \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2}{3} \int \frac{\sqrt{a + b \cos(c + dx)} (-b(2aA - 3bB) \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx))}{2 \cos^{3/2}(c + dx)} dx \right. \\
& \downarrow \text{27} \\
& \left. \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{3} \int \frac{\sqrt{a + b \cos(c + dx)} (-b(2aA - 3bB) \cos^2(c + dx) + (Aa^2 + 6bBa + 3Ab^2) \cos(c + dx))}{\cos^{3/2}(c + dx)} dx \right) \right. \\
& \downarrow \text{3042} \\
& \left. \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{3} \int \frac{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})} (-b(2aA - 3bB) \sin(c + dx + \frac{\pi}{2})^2 + (Aa^2 + 6bBa + 3Ab^2) \sin(c + dx + \frac{\pi}{2}))}{\sin(c + dx + \frac{\pi}{2})^{3/2}} dx \right) \right. \\
& \downarrow \text{3526} \\
& \left. \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{3} \left(2 \int \frac{-b(6Ba^2 + 14Aba - 3b^2B) \cos^2(c + dx) - (3Ba^3 + 7Aba^2 - 9b^2Ba - 3Ab^3)}{2 \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \right) \right. \right. \\
& \downarrow \text{27} \\
& \left. \left. \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{3} \left(\int \frac{-b(6Ba^2 + 14Aba - 3b^2B) \cos^2(c + dx) - (3Ba^3 + 7Aba^2 - 9b^2Ba - 3Ab^3)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \right) \right) \right) \right. \\
& \downarrow \text{3042}
\end{aligned}$$

3.609. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{-b(6Ba^2+14Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})^2+(-3Ba^3-7Aba^2+9b^2Ba+9Ab^2)\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx)}}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx)}}dx\right)\right)$$

↓ 3540

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{3(2Ab+5aB)\cos^2(c+dx)b^3+a(6Ba^2+14Aba-3b^2B)b+2a(Aa^2+9bBa+9Ab^2)\cos(c+dx)b}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{3(2Ab+5aB)\sin(c+dx+\frac{\pi}{2})^2b^3+a(6Ba^2+14Aba-3b^2B)b+2a(Aa^2+9bBa+9Ab^2)\sin(c+dx+\frac{\pi}{2})b}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{ab(6Ba^2+14Aba-3b^2B)+2ab(Aa^2+9bBa+9Ab^2)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+3b^3(5aB+2Ab)\int\frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{ab(6Ba^2+14Aba-3b^2B)+2ab(Aa^2+9bBa+9Ab^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+3b^3(5aB+2Ab)\int\frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)\right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\int\frac{ab(6Ba^2+14Aba-3b^2B)+2ab(Aa^2+9bBa+9Ab^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{6b^2\sqrt{a+b}(5aB+2Ab)\cot(c+dx+\frac{\pi}{2})}{2b}\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{ab(6a^2B+14aAb-3b^2B)\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-ab(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B))\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2b\sqrt{a+b}(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{-ab(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B))\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2b\sqrt{a+b}(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{ab(6a^2B+14aAb-3b^2B)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2b\sqrt{a+b}(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d}\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{3}\left(\frac{2b\sqrt{a+b}(-2a^2(A-3B)+2ab(7A-9B)-3b^2(6A+B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{d}\right)\right)$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2),x]`

```

output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*A*(a + b*Cos[c + d*x])^(3/2)*S
in[c + d*x])/(3*d*Cos[c + d*x]^(3/2)) + (((2*(a - b)*b*Sqrt[a + b]*(14*a*A
*b + 6*a^2*B - 3*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d
*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])), -((a + b)/(a - b))*Sqrt[(a*(1 - S
ec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*
Sqrt[a + b]*(2*a*b*(7*A - 9*B) - 2*a^2*(A - 3*B) - 3*b^2*(6*A + B))*Cot[c
+ d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c +
d*x]])), -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a
*(1 + Sec[c + d*x]))/(a - b)]/d - (6*b^2*Sqrt[a + b]*(2*A*b + 5*a*B)*Cot[
c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b
]*Sqrt[Cos[c + d*x]])), -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a
+ b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d)/(2*b) + (6*a*(2*A*b + a*B)
*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) - ((14*a*A*
b + 6*a^2*B - 3*b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[
c + d*x]]))/3

```

3.609.3.1 Defintions of rubi rules used

```

rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3288 Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]

```

```

rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f
_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]

```

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*SIN[e + f*x])^(m - 1)*((c + d*SIN[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*SIN[e + f*x])^(m - 2)*(c + d*SIN[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*SIN[e + f*x]]/Sqrt[b*SIN[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*SIN[e + f*x]]*Sqrt[c + d*SIN[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + SIN[e + f*x])/((a + b*SIN[e + f*x])^(3/2)*Sqrt[c + d*SIN[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3526 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] :> Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x] *(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(b*d*m + a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - (d*(A*(a*d*(n + 2) - b*c*(n + 1)) + B*(b*d*(n + 1) - a*c*(n + 2))) - C*(b*c*d*(n + 1) - a*(c^2 + d^2*(n + 1)))]*Sin[e + f*x] + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])]*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.609.4 Maple [F(-1)]

Timed out.

hanged

input `int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x)`

output `int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x)`

$$3.609. \quad \int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx) dx$$

3.609.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="fricas")`

output `Timed out`

3.609.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(5/2),x)`

output `Timed out`

3.609.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{5/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(5/2), x)`

3.609.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{5/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(5/2),x, algo rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(5/2), x)`

3.609.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{5/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{5/2} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(5/2), x)`

3.610 $\int (a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx)) \sec^{3/2}(c+dx) dx$

3.610.1 Optimal result	5577
3.610.2 Mathematica [B] (verified)	5578
3.610.3 Rubi [A] (verified)	5579
3.610.4 Maple [B] (verified)	5585
3.610.5 Fricas [F(-1)]	5586
3.610.6 Sympy [F(-1)]	5587
3.610.7 Maxima [F]	5587
3.610.8 Giac [F]	5587
3.610.9 Mupad [F(-1)]	5588

3.610.1 Optimal result

Integrand size = 35, antiderivative size = 607

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \frac{(a - b)\sqrt{a + b}(8a^2A - 4Ab^2 - 9abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \mid -\frac{a+b}{a-b}\right) + \sqrt{a + b}(8a^2(A - B) - 2b^2(2A + B) - 3ab(8A + 3B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) + \sqrt{a + b}(20aAb + 15a^2B + 4b^2B) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) + \frac{b(4aA - bB)\sqrt{a + b \cos(c + dx)} \sin(c + dx)}{2d\sqrt{\sec(c + dx)}} + \frac{(8a^2A - 4Ab^2 - 9abB) \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{4d} + \frac{2aA(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)} \sin(c + dx)}{d}$$

output

```

-1/2*b*(4*A*a-B*b)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)+2*
a*A*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)*sec(d*x+c)^(1/2)/d-1/4*(8*A*a^2-4*A*
b^2-9*B*a*b)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)*sec(d*x+c)^(1/2)/d+1/4*(a-b
)*(8*A*a^2-4*A*b^2-9*B*a*b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a
+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1
/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/sec(
d*x+c)^(1/2)-1/4*(8*a^2*(A-B)-2*b^2*(2*A+B)-3*a*b*(8*A+3*B))*csc(d*x+c)*El
lipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))
^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+
sec(d*x+c))/(a-b))^(1/2)/d/sec(d*x+c)^(1/2)-1/4*(20*A*a*b+15*B*a^2+4*B*b^2
)*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2
),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x
+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/d/sec(d*x+c)^(1/2)

```

3.610.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1278 vs. $2(607) = 1214$.

Time = 16.12 (sec) , antiderivative size = 1278, normalized size of antiderivative = 2.11

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Too large to display}$$

input

```

Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/
2),x]

```

output $(\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}(2a^2A\sin[c + dx] + (b^2B\sin[2(c + dx)]/4))/d + (\sqrt{(1 - \tan[(c + dx)/2]^2)^{-1}}(-8a^3A\tan[(c + dx)/2] - 8a^2Ab\tan[(c + dx)/2] + 4aAb^2\tan[(c + dx)/2] + 4Ab^3\tan[(c + dx)/2] + 9a^2bB\tan[(c + dx)/2] + 9ab^2B\tan[(c + dx)/2] + 16a^2A\tan[(c + dx)/2]^3 - 8Ab^3\tan[(c + dx)/2]^3 - 18ab^2B\tan[(c + dx)/2]^3 + 8a^3A\tan[(c + dx)/2]^5 - 8a^2Ab\tan[(c + dx)/2]^5 - 4aAb^2\tan[(c + dx)/2]^5 + 4Ab^3\tan[(c + dx)/2]^5 - 9a^2bB\tan[(c + dx)/2]^5 + 9ab^2B\tan[(c + dx)/2]^5 + 40aAb^2\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 30a^2bB\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 8b^3B\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 40aAb^2\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\tan[(c + dx)/2]^2\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 30a^2bB\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\tan[(c + dx)/2]^2\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 8b^3B\text{Ell}...$

3.610.3 Rubi [A] (verified)

Time = 3.05 (sec) , antiderivative size = 576, normalized size of antiderivative = 0.95, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3528, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{\frac{3}{2}}(c + dx)(a + b\cos(c + dx))^{5/2}(A + B\cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} \left(a + b\sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2} \left(A + B\sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b\cos(c + dx))^{5/2}(A + B\cos(c + dx))}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$3.610. \quad \int (a + b\cos(c + dx))^{5/2}(A + B\cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) dx$$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sin(c+dx+\frac{\pi}{2})^{3/2}} dx \\ & \downarrow 3468 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(2 \int \frac{\sqrt{a+b\cos(c+dx)}(-b(4aA-bB)\cos^2(c+dx)-(Aa^2-2bBa-Ab^2)\cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx \right. \\ & \downarrow 27 \\ & \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\sqrt{a+b\cos(c+dx)}(-b(4aA-bB)\cos^2(c+dx)-(Aa^2-2bBa-Ab^2)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \right. \right. \\ & \downarrow 3042 \\ & \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(-b(4aA-bB)\sin(c+dx+\frac{\pi}{2})^2+(-Aa^2+2bBa+Ab^2)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \right. \right. \right. \\ & \downarrow 3528 \\ & \left. \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \int \frac{-b(8Aa^2-9bBa-4Ab^2)\cos^2(c+dx)-2(2Aa^3-6bBa^2-6Ab^2a-b^3B)\cos(c+dx)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right. \right. \right. \\ & \downarrow 27 \\ & \left. \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \int \frac{-b(8Aa^2-9bBa-4Ab^2)\cos^2(c+dx)-2(2Aa^3-6bBa^2-6Ab^2a-b^3B)\cos(c+dx)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right. \right. \right. \\ & \downarrow 3042 \\ & \left. \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \int \frac{-b(8Aa^2-9bBa-4Ab^2)\sin(c+dx+\frac{\pi}{2})^2-2(2Aa^3-6bBa^2-6Ab^2a-b^3B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right. \right. \right. \\ & \downarrow 3540 \\ & \left. \left. \left. \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \left(\frac{\int \frac{b^2(15Ba^2+20Aba+4b^2B)\cos^2(c+dx)+2ab(4Ba^2+12Aba+b^2B)\cos(c+dx)+ab(8Aa^2-9bBa-4Ab^2)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b} \right) \right. \right. \right. \end{aligned}$$

3.610. $\int (a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx) dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{\int \frac{b^2(15Ba^2+20Aba+4b^2B)\sin(c+dx+\frac{\pi}{2})^2+2ab(4Ba^2+12Aba+b^2B)\sin(c+dx+\frac{\pi}{2})+ab(8Aa^2-9bBa-4Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{2b}\right)\right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{\int \frac{ab(8Aa^2-9bBa-4Ab^2)+2ab(4Ba^2+12Aba+b^2B)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+b^2(15a^2B+20aAb+4b^2B)}{2b}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{b^2(15a^2B+20aAb+4b^2B)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\int \frac{ab(8Aa^2-9bBa-4Ab^2)+2ab(4Ba^2+12Aba+b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{2b}\right)\right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{\int \frac{ab(8Aa^2-9bBa-4Ab^2)+2ab(4Ba^2+12Aba+b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2b\sqrt{a+b}(15a^2B+20aAb+4b^2B)}{2b}\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{ab(8a^2A-9abB-4Ab^2)\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-ab(8a^2(A-B)-3ab(8A+3B)-2b^2(2A+B))}{2b}\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{-ab(8a^2(A-B)-3ab(8A+3B)-2b^2(2A+B))\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx}{2b}\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{ab(8a^2A-9abB-4Ab^2)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2b\sqrt{a+b}(8a^2(A-B))}{d}\right)}\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{4}\left(\frac{2b\sqrt{a+b}(8a^2(A-B)-3ab(8A+3B)-2b^2(2A+B))\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{Ellip}}{d}\right)}\right)$$

input `Int[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/2*(b*(4*a*A - b*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/d + (2*a*A*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]) + (((2*(a - b)*b*Sqrt[a + b]*(8*a^2*A - 4*A*b^2 - 9*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(8*a^2*(A - B) - 2*b^2*(2*A + B) - 3*a*b*(8*A + 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (2*b*Sqrt[a + b]*(20*a*A*b + 15*a^2*B + 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/(2*b) - ((8*a^2*A - 4*A*b^2 - 9*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/4)`

3.610.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^(m - 1)*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Ssin[e + f*x])^(m - 2)*(c + d*Ssin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Ssin[e + f*x]]/Sqrt[b*Ssin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Ssin[e + f*x])^(3/2)*Sqrt[c + d*Ssin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Ssin[e + f*x])^(m - 1)*(c + d*Ssin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)])), x_Symbol] :> Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)])), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.610.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4458 vs. $2(549) = 1098$.

Time = 21.31 (sec) , antiderivative size = 4459, normalized size of antiderivative = 7.35

method	result	size
default	Expression too large to display	4459
parts	Expression too large to display	4748

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```


output `1/4/d*(8*A*cos(d*x+c)*sin(d*x+c)*a^2*b+8*A*a^3*sin(d*x+c)-8*A*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3-8*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^3-9*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2+24*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*b-2*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2+24*A*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^2-24*A*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^2*b*cos(d*x+c)^2-80*A*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)-40*A*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^2+48*B*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))...`

3.610.5 Fracas [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algo rithm="fricas")`

output `Timed out`

3.610.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(3/2),x)`

output `Timed out`

3.610.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{3/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(3/2), x)`

3.610.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{3/2} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(3/2), x)`

3.610. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx$

3.610.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sec^{3/2}(c + dx) dx = \int (A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)} \right)^{3/2} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(5/2),x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(5/2), x)`

3.611 $\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx$

3.611.1 Optimal result	5589
3.611.2 Mathematica [B] (verified)	5590
3.611.3 Rubi [A] (verified)	5591
3.611.4 Maple [B] (verified)	5597
3.611.5 Fricas [F]	5598
3.611.6 Sympy [F(-1)]	5599
3.611.7 Maxima [F]	5599
3.611.8 Giac [F]	5599
3.611.9 Mupad [F(-1)]	5600

3.611.1 Optimal result

Integrand size = 35, antiderivative size = 624

$$\int (a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx)) \sqrt{\sec(c+dx)} dx =$$

$$\frac{(a-b)\sqrt{a+b}(54aAb+33a^2B+16b^2B)\sqrt{\cos(c+dx)}\csc(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\middle|-\frac{a+b}{a-b}\right)\sqrt{a+b}}{24ad\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(4b^2(3A+4B)+a^2(48A+33B)+a(54Ab+26bB))\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{24d\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(30a^2Ab+8Ab^3+5a^3B+20ab^2B)\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticPi}\left(\frac{a+b}{b},\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{8bd\sqrt{\sec(c+dx)}} +$$

$$\frac{b(2Ab+3aB)\sqrt{a+b}\cos(c+dx)\sin(c+dx)}{4d\sqrt{\sec(c+dx)}} + \frac{bB(a+b\cos(c+dx))^{3/2}\sin(c+dx)}{3d\sqrt{\sec(c+dx)}} +$$

$$\frac{(54aAb+33a^2B+16b^2B)\sqrt{a+b}\cos(c+dx)\sqrt{\sec(c+dx)}\sin(c+dx)}{24d}$$

output $\frac{1}{3}bB(a+b\cos(dx+c))^{3/2}\sin(dx+c)/d\sec(dx+c)^{1/2} + \frac{1}{4}b(2A*b+3B*a)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\sec(dx+c)^{1/2} + \frac{1}{24}(54A*a*b+33B*a^2+16B*b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/d - \frac{1}{24}(a-b)(54A*a*b+33B*a^2+16B*b^2)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2})/a/d\sec(dx+c)^{1/2} + \frac{1}{24}(4*b^2*(3A+4B)+a^2*(48A+33B)+a*(54A*b+26B*b))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2})/d\sec(dx+c)^{1/2} - \frac{1}{8}(30A*a^2*b+8A*b^3+5B*a^3+20B*a*b^2)\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, (a+b)/b, ((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c))/(a+b))^{1/2}*(a*(1+\sec(dx+c))/(a-b))^{1/2})/b/d\sec(dx+c)^{1/2}$

3.611.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1504 vs. $2(624) = 1248$.

Time = 16.97 (sec) , antiderivative size = 1504, normalized size of antiderivative = 2.41

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]], x]`

output $(\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}((b^2B\sin[c + dx])/12 + (b(6Ab + 13aB)\sin[2(c + dx)])/24 + (b^2B\sin[3(c + dx)])/12))/d + (\sqrt{(1 - \tan[(c + dx)/2]^{-1})}(54a^2Ab\tan[(c + dx)/2] + 54aAb^2\tan[(c + dx)/2] + 33a^3B\tan[(c + dx)/2] + 33a^2bB\tan[(c + dx)/2] + 16ab^2B\tan[(c + dx)/2] + 16b^3B\tan[(c + dx)/2] - 108aAb^2\tan[(c + dx)/2]^3 - 66a^2bB\tan[(c + dx)/2]^3 - 32b^3B\tan[(c + dx)/2]^3 - 54a^2Ab\tan[(c + dx)/2]^5 + 54aAb^2\tan[(c + dx)/2]^5 - 33a^3B\tan[(c + dx)/2]^5 + 33a^2bB\tan[(c + dx)/2]^5 - 16ab^2B\tan[(c + dx)/2]^5 + 16b^3B\tan[(c + dx)/2]^5 + 180a^2Ab\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 48Ab^3\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 30a^3B\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 120ab^2B\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2)/(a + b)} + 180a^2Ab\text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)]\tan[(c + dx)/2]^2\sqrt{1 - \tan[(c + dx)/2]^2}\sqrt{(a + b + a\tan[(c + dx)/2]^2 - b\tan[(c + dx)/2]^2 - b\dots$

3.611.3 Rubi [A] (verified)

Time = 3.09 (sec) , antiderivative size = 593, normalized size of antiderivative = 0.95, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\sec(c + dx)}(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}\left(a + b \sin\left(c + dx + \frac{\pi}{2}\right)\right)^{5/2}\left(A + B \sin\left(c + dx + \frac{\pi}{2}\right)\right) dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{(a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx))}{\sqrt{\cos(c + dx)}} dx$$

$$3.611. \quad \int (a + b \cos(c + dx))^{5/2}(A + B \cos(c + dx))\sqrt{\sec(c + dx)} dx$$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \\ & \downarrow 3469 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{3} \int \frac{\sqrt{a+b\cos(c+dx)}(3b(2Ab+3aB)\cos^2(c+dx)+2(3Ba^2+6Aba+2b^2B)\cos(c+dx))}{2\sqrt{\cos(c+dx)}} dx \right. \\ & \downarrow 27 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \int \frac{\sqrt{a+b\cos(c+dx)}(3b(2Ab+3aB)\cos^2(c+dx)+2(3Ba^2+6Aba+2b^2B)\cos(c+dx))}{\sqrt{\cos(c+dx)}} dx \right. \\ & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(3b(2Ab+3aB)\sin(c+dx+\frac{\pi}{2})^2+2(3Ba^2+6Aba+2b^2B)\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \right. \\ & \downarrow 3528 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{b(33Ba^2+54Aba+16b^2B)\cos^2(c+dx)+2(12Ba^3+36Aba^2+19b^2Ba+16b^3)\cos(c+dx)+2ab(24Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab^3}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right. \right. \\ & \downarrow 27 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{b(33Ba^2+54Aba+16b^2B)\cos^2(c+dx)+2(12Ba^3+36Aba^2+19b^2Ba+16b^3)\cos(c+dx)+2ab(24Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab^3}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right. \right. \\ & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{b(33Ba^2+54Aba+16b^2B)\sin(c+dx+\frac{\pi}{2})^2+2(12Ba^3+36Aba^2+19b^2Ba+16b^3)\sin(c+dx+\frac{\pi}{2})+2ab(24Aa^2+13bBa+6Ab^2)\sin(c+dx+\frac{\pi}{2})+ab^3}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right. \right. \\ & \downarrow 3540 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{\int -\frac{3b(5Ba^3+30Aba^2+20b^2Ba+8Ab^3)\cos^2(c+dx)-2ab(24Aa^2+13bBa+6Ab^2)\cos(c+dx)+ab^3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{2b} dx \right. \right. \right. \end{aligned}$$

3.611. $\int (a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))\sqrt{\sec(c+dx)} dx$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{-3b(5Ba^3+30Aa^2)}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{-3b(5Ba^3+30Aa^2)}{\dots}\right)\right)\right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{ab(33Ba^2+54Aa)}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{ab(33Ba^2+54Aa)}{\dots}\right)\right)\right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{ab(33Ba^2+54Aa)}{\dots}\right)\right)\right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{ab(33a^2B+54Aa)}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{-ab(a^2(48A+54aB+16b^2)+2b^2(a^2+2ab+2b^2))}{2d\sqrt{a+b\cos(c+dx)}}\right)\right)\right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{ab(33a^2B+54a^2+16b^2)}{2d\sqrt{a+b\cos(c+dx)}}\right)\right)\right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{6}\left(\frac{1}{4}\left(\frac{(33a^2B+54aAb+16b^2B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\frac{2b\sqrt{a+b}(a^2(48A+54aB+16b^2)+2b^2(a^2+2ab+2b^2))}{2d\sqrt{a+b\cos(c+dx)}}\right)\right)\right)$$

input `Int[(a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x])*sqrt[sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((b*B*Sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + ((3*b*(2*A*b + 3*a*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*(54*a*A*b + 33*a^2*B + 16*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(4*b^2*(3*A + 4*B) + a^2*(48*A + 33*B) + a*(54*A*b + 26*b*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d + (6*Sqrt[a + b]*(30*a^2*A*b + 8*A*b^3 + 5*a^3*B + 20*a*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/b + ((54*a*A*b + 33*a^2*B + 16*b^2*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])/4)/6)`

3.611.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

```
rule 3532 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e
_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]
/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*
B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x
]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] &
& NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.611.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4801 vs. $2(564) = 1128$.

Time = 20.15 (sec) , antiderivative size = 4802, normalized size of antiderivative = 7.70

method	result	size
parts	Expression too large to display	4802
default	Expression too large to display	4828

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```


3.611.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)**(1/2),x)`

output `Timed out`

3.611.7 Maxima [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sqrt(sec(d*x + c)), x)`

3.611.8 Giac [F]

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2} \sqrt{\sec(dx + c)} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))*sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)*sqrt(sec(d*x + c)), x)`

3.611. $\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx$

3.611.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) \sqrt{\sec(c + dx)} dx = \int (A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}} (a + b \cos(c + dx))^{5/2} dx$$

input `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(5/2), x)`

output `int((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(5/2), x)`

$$3.612 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

3.612.1 Optimal result	5601
3.612.2 Mathematica [B] (warning: unable to verify)	5602
3.612.3 Rubi [A] (verified)	5603
3.612.4 Maple [B] (verified)	5611
3.612.5 Fracas [F]	5611
3.612.6 Sympy [F(-1)]	5612
3.612.7 Maxima [F]	5612
3.612.8 Giac [F]	5612
3.612.9 Mupad [F(-1)]	5613

3.612.1 Optimal result

Integrand size = 35, antiderivative size = 724

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx =$$

$$\frac{(a-b)\sqrt{a+b}(264a^2Ab+128Ab^3+15a^3B+284ab^2B)\sqrt{\cos(c+dx)}\csc(c+dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{192abd\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(15a^3B+8b^3(16A+9B)+2a^2b(132A+59B)+4ab^2(52A+71B))\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticE}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{192bd\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(40a^3Ab+160aAb^3-5a^4B+120a^2b^2B+48b^4B)\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticPi}\left(\frac{a+b}{b},\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{64b^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{bB(a+b \cos(c+dx))^{3/2}\sin(c+dx)}{4d \sec^{\frac{3}{2}}(c+dx)} +$$

$$\frac{(24aAb+5a^2B+12b^2B)\sqrt{a+b \cos(c+dx)}\sin(c+dx)}{32d\sqrt{\sec(c+dx)}} +$$

$$\frac{(8Ab+11aB)(a+b \cos(c+dx))^{3/2}\sin(c+dx)}{24d\sqrt{\sec(c+dx)}} +$$

$$\frac{(264a^2Ab+128Ab^3+15a^3B+284ab^2B)\sqrt{a+b \cos(c+dx)}\sqrt{\sec(c+dx)}\sin(c+dx)}{192bd}$$

$$3.612. \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$$

output $\frac{1}{4}bB(a+b\cos(dx+c))^{3/2}\sin(dx+c)/d\sec(dx+c)^{3/2} + \frac{1}{24}(8A^2b+11B^2a)(a+b\cos(dx+c))^{3/2}\sin(dx+c)/d\sec(dx+c)^{1/2} + \frac{1}{32}(24A^2ab+5B^2a^2+12B^2b^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}/d\sec(dx+c)^{1/2} + \frac{1}{192}(264A^2a^2b+128A^2b^3+15B^2a^3+284B^2ab^2)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/b/d - \frac{1}{192}(a-b)(264A^2a^2b+128A^2b^3+15B^2a^3+284B^2ab^2)\operatorname{csc}(dx+c)\operatorname{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2})/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/a/b/d\sec(dx+c)^{1/2} + \frac{1}{192}(15B^2a^3+8b^3(16A+9B)+2a^2b(132A+59B)+4ab^2(52A+71B))\operatorname{csc}(dx+c)\operatorname{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/b/d\sec(dx+c)^{1/2} - \frac{1}{64}(40A^2a^3b+160A^2ab^3-5B^2a^4+120B^2a^2b^2+48B^2b^4)\operatorname{csc}(dx+c)\operatorname{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b, ((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c)))/(a+b)^{1/2}(a(1+\sec(dx+c)))/(a-b)^{1/2}/b^2/d\sec(dx+c)^{1/2}$

3.612.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 1857 vs. $2(724) = 1448$.

Time = 16.90 (sec) , antiderivative size = 1857, normalized size of antiderivative = 2.56

$$\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx = \text{Too large to display}$$

input `Integrate[((a + bCos[c + d*x])^(5/2)*(A + BCos[c + d*x]))/Sqrt[Sec[c + d*x]], x]`

output $(\text{Sqrt}[a + b \cos[c + dx]] \cdot \text{Sqrt}[\text{Sec}[c + dx]] \cdot ((b(8Ab + 17aB) \sin[c + dx])/96 + ((104a^2A + 59a^2B + 48b^2B) \sin[2(c + dx)])/192 + (b(8Ab + 17aB) \sin[3(c + dx)])/96 + (b^2B \sin[4(c + dx)])/32))/d + (\text{Sqrt}[(1 - \text{Tan}[(c + dx)/2]^2)^{-1}] \cdot (264a^3Ab \text{Tan}[(c + dx)/2] + 264a^2Ab^2 \text{Tan}[(c + dx)/2] + 128a^2A^2 \text{Tan}[(c + dx)/2] + 128Ab^4 \text{Tan}[(c + dx)/2] + 15a^4B \text{Tan}[(c + dx)/2] + 15a^3bB \text{Tan}[(c + dx)/2] + 284a^2b^2B \text{Tan}[(c + dx)/2] + 284ab^3B \text{Tan}[(c + dx)/2] - 528a^2Ab^2 \text{Tan}[(c + dx)/2]^3 - 256Ab^4 \text{Tan}[(c + dx)/2]^3 - 30a^3bB \text{Tan}[(c + dx)/2]^3 - 568ab^3B \text{Tan}[(c + dx)/2]^3 - 264a^3Ab \text{Tan}[(c + dx)/2]^5 + 264a^2Ab^2 \text{Tan}[(c + dx)/2]^5 - 128a^2A^2 \text{Tan}[(c + dx)/2]^5 + 128Ab^4 \text{Tan}[(c + dx)/2]^5 - 15a^4B \text{Tan}[(c + dx)/2]^5 + 15a^3bB \text{Tan}[(c + dx)/2]^5 - 284a^2b^2B \text{Tan}[(c + dx)/2]^5 + 284ab^3B \text{Tan}[(c + dx)/2]^5 + 240a^3Ab \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] + 960a^2Ab^3 \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] - 30a^4B \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] + 720a^2b^2B \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \dots$

3.612.3 Rubi [A] (verified)

Time = 3.77 (sec) , antiderivative size = 693, normalized size of antiderivative = 0.96, number of steps used = 22, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.629$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{\csc(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \sqrt{\cos(c + dx)} (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$$

3.612. $\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3469

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{4} \int \frac{1}{2} \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (b(8Ab+11aB)\cos^2(c+dx) + 2(4Ba^2+8Ab$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \int \sqrt{\cos(c+dx)} \sqrt{a+b\cos(c+dx)} (b(8Ab+11aB)\cos^2(c+dx) + 2(4Ba^2+8Ab$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \int \sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)} \sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)} \left(b(8Ab+11aB)\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{\int \frac{\sqrt{a+b\cos(c+dx)}(3b(5Ba^2+24Aba+12b^2B)\cos^2(c+dx)+2b(24Aa^2+31bBa+16Ab^2)\cos(c+dx)+ab}{2\sqrt{\cos(c+dx)}}}{3b}\right) dx\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{\int \frac{\sqrt{a+b\cos(c+dx)}(3b(5Ba^2+24Aba+12b^2B)\cos^2(c+dx)+2b(24Aa^2+31bBa+16Ab^2)\cos(c+dx)+ab}{\sqrt{\cos(c+dx)}}}{6b}\right) dx\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{\int \frac{\sqrt{a+b\sin\left(c+dx+\frac{\pi}{2}\right)}\left(3b(5Ba^2+24Aba+12b^2B)\sin\left(c+dx+\frac{\pi}{2}\right)^2+2b(24Aa^2+31bBa+16Ab^2)\sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\sqrt{\sin\left(c+dx+\frac{\pi}{2}\right)}}}{6b}\right) dx\right)$$

↓ 3528

3.612. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{\frac{1}{2}\int\frac{b(15Ba^3+264Aba^2+284b^2Ba+128Ab^3)\cos^2(c+dx)+2b(96Aa^3+161bBa^2+152Ab^2a+36b^3B)\cos}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{\right)}\right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{\frac{1}{4}\int\frac{b(15Ba^3+264Aba^2+284b^2Ba+128Ab^3)\cos^2(c+dx)+2b(96Aa^3+161bBa^2+152Ab^2a+36b^3B)\cos}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{\right)}\right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{\frac{1}{4}\int\frac{b(15Ba^3+264Aba^2+284b^2Ba+128Ab^3)\sin(c+dx+\frac{\pi}{2})^2+2b(96Aa^3+161bBa^2+152Ab^2a+36b^3B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{\right)}\right)$$

↓ 3540

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{\frac{1}{4}\left(\int\frac{-2a(59Ba^2+104Aba+36b^2B)\cos(c+dx)b^2-3(-5Ba^4+40Aba^3+120b^2Ba^2+160Ab^3a+48b^4B)\cos^2(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}\right)}{2b}\right)}$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{1}{8}\left(\frac{\frac{1}{4}\left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}}-\int\frac{-2a(59Ba^2+104Aba+36b^2B)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{\right)}{2b}\right)}$$

↓ 3042

3.612. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{-2a(59Ba^2+104Aba+36b^2)}{\dots} \right) \right) \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{ab(15Ba^3+264Aba^2+284b^2)}{\dots} \right) \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{ab(15Ba^3+264Aba^2+284b^2)}{\dots} \right) \right) \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \int \frac{ab(15Ba^3+264Aba^2+284b^2)}{\dots} \right) \right) \right)$$

↓ 3477

3.612. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{ab(15a^3B+264a^2Ab+284a^2b^2B+128Ab^3)}{d} \right) \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{-ab(15a^3B+2a^2b(132A+5b^2))}{d} \right) \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{ab(15a^3B+264a^2Ab+284a^2b^2B+128Ab^3)}{d} \right) \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{3b(5a^2B+24aAb+12b^2B) \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{a+b \cos(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(15a^3B+264a^2Ab+284ab^2B+128Ab^3) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{ab(15a^3B+264a^2Ab+284a^2b^2B+128Ab^3)}{d} \right) \right) \right)$$

input `Int[((a + b*cos[c + d*x])^(5/2)*(A + B*cos[c + d*x]))/Sqrt[Sec[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((b*B*cos[c + d*x]^(3/2)*(a + b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(4*d) + (((8*A*b + 11*a*B)*Sqrt[Cos[c + d*x]]*(a + b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(3*d) + ((3*b*(24*a*A*b + 5*a^2*B + 12*b^2*B)*Sqrt[Cos[c + d*x]]*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*b*Sqrt[a + b]*(264*a^2*A*b + 128*A*b^3 + 15*a^3*B + 284*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*b*Sqrt[a + b]*(15*a^3*B + 8*b^3*(16*A + 9*B) + 2*a^2*b*(132*A + 59*B) + 4*a*b^2*(52*A + 71*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d + (6*Sqrt[a + b]*(40*a^3*A*b + 160*a*A*b^3 - 5*a^4*B + 120*a^2*b^2*B + 48*b^4*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d)/b + ((264*a^2*A*b + 128*A*b^3 + 15*a^3*B + 284*a*b^2*B)*Sqrt[a + b*cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]))/4)/(6*b))/8`

3.612.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B))*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{((b \sin(e + fx) + f x) \sqrt{c + d \sin(e + fx)})^{3/2}} dx$$

$$:= \text{Simp}[-2 A (c - d) (\tan[e + f x] / (f b c^2)) \text{Rt}[(c + d) / b, 2] \sqrt{c ((1 + \text{Csc}[e + f x]) / (c - d))} \sqrt{c ((1 - \text{Csc}[e + f x]) / (c + d))} \text{EllipticE}[\text{ArcSin}[\sqrt{c + d \sin[e + f x]}] / \sqrt{b \sin[e + f x]}] / \text{Rt}[(c + d) / b, 2], -(c + d) / (c - d)], x] /;$$` `FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d) / b]`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$:= \text{Simp}[(A - B) / (a - b) \text{Int}[1 / (\sqrt{a + b \sin[e + f x]} \sqrt{c + d \sin[e + f x]}), x], x] - \text{Simp}[(A b - a B) / (a - b) \text{Int}[(1 + \sin[e + f x]) / ((a + b \sin[e + f x])^{3/2} \sqrt{c + d \sin[e + f x]}), x], x] /;$$` `FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +

$$\int ((a + b \sin(e + fx))^m ((c + d \sin(e + fx))^n ((A + B \sin(e + fx) + C \sin(e + fx) + f x)^2)) dx$$

$$:= \text{Simp}[(-C) \cos[e + f x] (a + b \sin[e + f x])^m ((c + d \sin[e + f x])^{n+1} / (d f (m + n + 2))), x] + \text{Simp}[1 / (d (m + n + 2)) \text{Int}[(a + b \sin[e + f x])^{m-1} (c + d \sin[e + f x])^n \text{Simp}[a A d (m + n + 2) + C (b c m + a d (n + 1)) + (d (A b + a B) (m + n + 2) - C (a c - b d (m + n + 1))) \sin[e + f x] + (C (a d m - b c (m + 1)) + b B d (m + n + 2)) \sin[e + f x]^2, x], x], x] /;$$` `FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2)/

$$\int \frac{(A + B \sin(e + fx) + C \sin(e + fx) + f x)^2 ((a + b \sin(e + fx)) \sqrt{c + d \sin(e + fx)})^{3/2}}{((a + b \sin(e + fx)) \sqrt{c + d \sin(e + fx)})^{3/2}} dx$$

$$:= \text{Simp}[C / b^2 \text{Int}[\sqrt{a + b \sin[e + f x]} / \sqrt{c + d \sin[e + f x]}, x], x] + \text{Simp}[1 / b^2 \text{Int}[(A b^2 - a^2 C + b (b B - 2 a C) \sin[e + f x]) / ((a + b \sin[e + f x])^{3/2} \sqrt{c + d \sin[e + f x]}), x], x] /;$$` `FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]]), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])]*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.612.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5694 vs. $2(658) = 1316$.

Time = 21.54 (sec) , antiderivative size = 5695, normalized size of antiderivative = 7.87

method	result	size
parts	Expression too large to display	5695
default	Expression too large to display	5765

```
input int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.612.5 Fracas [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sqrt{\sec(dx + c)}} dx$$

```
input integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algo
rithm="fricas")
```

```
output integral((B*b^2*cos(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(d*x + c)^2
+ (B*a^2 + 2*A*a*b)*cos(d*x + c))*sqrt(b*cos(d*x + c) + a)/sqrt(sec(d*x +
c)), x)
```

3.612. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sqrt{\sec(c+dx)}} dx$

3.612.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.612.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/sqrt(sec(d*x + c)), x)`

3.612.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sqrt{\sec(dx + c)}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/sqrt(sec(d*x + c)), x)`

3.612.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sqrt{\sec(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\sqrt{\frac{1}{\cos(c + dx)}}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(1/2), x)`

$$3.613 \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

3.613.1 Optimal result	5614
3.613.2 Mathematica [A] (verified)	5615
3.613.3 Rubi [A] (verified)	5616
3.613.4 Maple [B] (verified)	5625
3.613.5 Fracas [F(-1)]	5625
3.613.6 Sympy [F(-1)]	5625
3.613.7 Maxima [F]	5626
3.613.8 Giac [F]	5626
3.613.9 Mupad [F(-1)]	5626

3.613.1 Optimal result

Integrand size = 35, antiderivative size = 839

$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx =$$

$$\frac{(a-b)\sqrt{a+b}(150a^3Ab+2840aAb^3-45a^4B+1692a^2b^2B+1024b^4B)\sqrt{\cos(c+dx)}\csc(c+dx)E(\arcsin(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}))}{1920ab^2d\sqrt{\sec(c+dx)}} -$$

$$\frac{\sqrt{a+b}(45a^4B-30a^3b(5A+B)-16b^4(45A+64B)-8ab^3(355A+193B)-4a^2b^2(295A+423B))\sqrt{\cos(c+dx)}\csc(c+dx)}{1920b^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(10a^4Ab-240a^2Ab^3-96Ab^5-3a^5B-40a^3b^2B-240ab^4B)\sqrt{\cos(c+dx)}\csc(c+dx)\text{EllipticPi}(\arcsin(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}}), \frac{a+b\cos(c+dx)}{a+b})}{128b^3d\sqrt{\sec(c+dx)}} +$$

$$\frac{(50a^2Ab+120Ab^3-15a^3B+172ab^2B)\sqrt{a+b\cos(c+dx)}\sin(c+dx)}{320bd\sqrt{\sec(c+dx)}} +$$

$$\frac{(50aAb-15a^2B+64b^2B)(a+b\cos(c+dx))^{3/2}\sin(c+dx)}{240bd\sqrt{\sec(c+dx)}} +$$

$$\frac{(10Ab-3aB)(a+b\cos(c+dx))^{5/2}\sin(c+dx)}{40bd\sqrt{\sec(c+dx)}} + \frac{B(a+b\cos(c+dx))^{7/2}\sin(c+dx)}{5bd\sqrt{\sec(c+dx)}} +$$

$$\frac{(150a^3Ab+2840aAb^3-45a^4B+1692a^2b^2B+1024b^4B)\sqrt{a+b\cos(c+dx)}\sqrt{\sec(c+dx)}\sin(c+dx)}{1920b^2d}$$

$$3.613. \quad \int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

output

```

1/240*(50*A*a*b-15*B*a^2+64*B*b^2)*(a+b*cos(d*x+c))^(3/2)*sin(d*x+c)/b/d/
sec(d*x+c)^(1/2)+1/40*(10*A*b-3*B*a)*(a+b*cos(d*x+c))^(5/2)*sin(d*x+c)/b/d/
sec(d*x+c)^(1/2)+1/5*B*(a+b*cos(d*x+c))^(7/2)*sin(d*x+c)/b/d/sec(d*x+c)^(1
/2)+1/320*(50*A*a^2*b+120*A*b^3-15*B*a^3+172*B*a*b^2)*sin(d*x+c)*(a+b*cos(
d*x+c))^(1/2)/b/d/sec(d*x+c)^(1/2)+1/1920*(150*A*a^3*b+2840*A*a*b^3-45*B*a
^4+1692*B*a^2*b^2+1024*B*b^4)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)*sec(d*x+c)
^(1/2)/b^2/d-1/1920*(a-b)*(150*A*a^3*b+2840*A*a*b^3-45*B*a^4+1692*B*a^2*b^
2+1024*B*b^4)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(
d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(
d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/b^2/d/sec(d*x+c)^(1/
2)-1/1920*(45*B*a^4-30*a^3*b*(5*A+B)-16*b^4*(45*A+64*B)-8*a*b^3*(355*A+193
*B)-4*a^2*b^2*(295*A+423*B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(
a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(
1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b^2/d/s
ec(d*x+c)^(1/2)+1/128*(10*A*a^4*b-240*A*a^2*b^3-96*A*b^5-3*B*a^5-40*B*a^3*
b^2-240*B*a*b^4)*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/
cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2
)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b^3/d/sec(
d*x+c)^(1/2)

```

3.613.2 Mathematica [A] (verified)

Time = 13.17 (sec) , antiderivative size = 703, normalized size of antiderivative = 0.84

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^3(c + dx)} dx = \frac{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{1}{960} (170aAb + 93a^2B - \dots) \right)}{-b(a + b) (150a^3Ab + 2840aAb^3 - 45a^4B + 1692a^2b^2B + 1024b^4B) E(\arcsin(\tan(\frac{1}{2}(c + dx))) | \frac{-a+b}{a+b}) \sec}$$

input

```

Integrate[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(
3/2),x]

```

output $(\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}*((((170*a*A*b + 93*a^2*B + 88*b^2*B)*\sin[c + dx])/960 + ((590*a^2*A*b + 480*A*b^3 + 15*a^3*B + 1024*a*b^2*B)*\sin[2*(c + dx)])/(1920*b) + ((170*a*A*b + 93*a^2*B + 100*b^2*B)*\sin[3*(c + dx)])/960 + (b*(10*A*b + 21*a*B)*\sin[4*(c + dx)])/320 + (b^2*B*\sin[5*(c + dx)]/80))/d - ((-b*(a + b)*(150*a^3*A*b + 2840*a*A*b^3 - 45*a^4*B + 1692*a^2*b^2*B + 1024*b^4*B)*\text{EllipticE}[\text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)]*\sec[(c + dx)/2]^2*\sqrt{((a + b*\cos[c + dx])*\sec[(c + dx)/2]^2)/(a + b)} + a*(a + b)*(45*a^4*B - 30*a^3*b*(5*A + 3*B) + 60*a^2*b^2*(5*A + 11*B) + 16*b^4*(45*A + 64*B) + 8*a*b^3*(265*A + 129*B))*\text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)]*\sec[(c + dx)/2]^2*\sqrt{((a + b*\cos[c + dx])*\sec[(c + dx)/2]^2)/(a + b)} + 15*(10*a^4*A*b - 240*a^2*A*b^3 - 96*A*b^5 - 3*a^5*B - 40*a^3*b^2*B - 240*a*b^4*B)*(a - b)*\text{EllipticF}[\text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] + 2*b*\text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)]*\sec[(c + dx)/2]^2*\sqrt{((a + b*\cos[c + dx])*\sec[(c + dx)/2]^2)/(a + b)} - b*(150*a^3*A*b + 2840*a*A*b^3 - 45*a^4*B + 1692*a^2*b^2*B + 1024*b^4*B)*(a + b*\cos[c + dx])*(\cos[c + dx]*\sec[(c + dx)/2]^2)^{(3/2)}*\sec[c + dx]*\text{Tan}[(c + dx)/2])/(1920*b^3*d*\sqrt{a + b*\cos[c + dx]}*(\cos[c + dx]*\sec[(c + dx)/2]^2)^{(3/2)}*\sec[c + dx]^{(3/2)}))$

3.613.3 Rubi [A] (verified)

Time = 4.58 (sec) , antiderivative size = 807, normalized size of antiderivative = 0.96, number of steps used = 25, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3528, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\csc(c + dx + \frac{\pi}{2})^{3/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \cos^{3/2}(c + dx) (a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx)) dx$$

3.613. $\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \sin\left(c+dx+\frac{\pi}{2}\right)^{3/2} \left(a+b\sin\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2} \left(A+B\sin\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

↓ 3469

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a+b\cos(c+dx))^{5/2}((10Ab-3aB)\cos^2(c+dx)+8bB\cos(c+dx)+aB) dx}{2\sqrt{\cos(c+dx)}}}{5b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{10b} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a+b\cos(c+dx))^{5/2}((10Ab-3aB)\cos^2(c+dx)+8bB\cos(c+dx)+aB) dx}{\sqrt{\cos(c+dx)}}}{10b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{10b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}((10Ab-3aB)\sin^2(c+dx+\frac{\pi}{2})+8bB\sin(c+dx+\frac{\pi}{2})+aB) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{10b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{10b} \right)$$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{4} \int \frac{(a+b\cos(c+dx))^{3/2}((-15Ba^2+50Aba+64b^2B)\cos^2(c+dx)+6b(10Ab+9aB)\cos(c+dx)+5a(2Ab+aB)) dx}{2\sqrt{\cos(c+dx)}}}{10b} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \int \frac{(a+b\cos(c+dx))^{3/2}((-15Ba^2+50Aba+64b^2B)\cos^2(c+dx)+6b(10Ab+9aB)\cos(c+dx)+5a(2Ab+aB)) dx}{\sqrt{\cos(c+dx)}}}{10b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \int \frac{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}((-15Ba^2+50Aba+64b^2B)\sin^2(c+dx+\frac{\pi}{2})+6b(10Ab+9aB)\sin(c+dx+\frac{\pi}{2})+5a(2Ab+aB)) dx}{\sqrt{\sin(c+dx+\frac{\pi}{2})}}}{10b} \right)$$

3.613. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \left(\frac{1}{3} \int \frac{\sqrt{a+b\cos(c+dx)}(3(-15Ba^3+50Aba^2+172b^2Ba+120Ab^3)\cos^2(c+dx)+2b(147Ba^2+310Aba+128Ab^2))}{2\sqrt{\cos(c+dx)}} dx \right)}{\sqrt{\cos(c+dx)}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \left(\frac{1}{6} \int \frac{\sqrt{a+b\cos(c+dx)}(3(-15Ba^3+50Aba^2+172b^2Ba+120Ab^3)\cos^2(c+dx)+2b(147Ba^2+310Aba+128Ab^2))}{\sqrt{\cos(c+dx)}} dx \right)}{\sqrt{\cos(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \left(\frac{1}{6} \int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}(3(-15Ba^3+50Aba^2+172b^2Ba+120Ab^3)\sin^2(c+dx+\frac{\pi}{2})+2b(147Ba^2+310Aba+128Ab^2))}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx \right)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} \right)$$

↓ 3528

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{2} \int \frac{(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B)\cos^2(c+dx)+2b(573Ba^3+1610Aba^2+1280Ab^2a+1024b^3B)}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right) \right)}{\sqrt{\cos(c+dx)}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B)\cos^2(c+dx)+2b(573Ba^3+1610Aba^2+1280Ab^2a+1024b^3B)}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right) \right)}{\sqrt{\cos(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \int \frac{(-45Ba^4+150Aba^3+1692b^2Ba^2+2840Ab^3a+1024b^4B)\sin^2(c+dx+\frac{\pi}{2})+2b(573Ba^3+1610Aba^2+1280Ab^2a+1024b^3B)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \right)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} \right)$$

↓ 3540

3.613. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\int \frac{-15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)\cos^2(c+dx)-2ab(15Ba^3+590Aba^2+7\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)})}{2b} dx \right) \right) \right) \right)$$

↓ 25

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)\cos^2(c+dx)-2ab(15Ba^3+590Aba^2+7\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)})}{2b} dx \right) \right) \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)\cos^2(c+dx)-2ab(15Ba^3+590Aba^2+7\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)})}{2b} dx \right) \right) \right) \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \int \frac{15(-3Ba^5+10Aba^4-40b^2Ba^3-240Ab^3a^2-240b^4Ba-96Ab^5)\cos^2(c+dx)-2ab(15Ba^3+590Aba^2+7\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)})}{2b} dx \right) \right) \right) \right)$$

↓ 3042

3.613. $\int \frac{(a+b\cos(c+dx))^{5/2}(A+B\cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} \right) \right) \right) \right) - \frac{f(a(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b \cos(c+dx)})}{bd\sqrt{\cos(c+dx)}}$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} \right) \right) \right) \right) - \frac{f(a(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b \cos(c+dx)})}{bd\sqrt{\cos(c+dx)}}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} \right) \right) \right) \right) - \frac{a(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B)}{bd\sqrt{\cos(c+dx)}} \right) \right) \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{8} \left(\frac{1}{6} \left(\frac{1}{4} \left(\frac{(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B) \sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(-45a^4B+150a^3Ab+1692a^2b^2B+2840aAb^3+1024b^4B)}{bd\sqrt{\cos(c+dx)}} \right) \right) \right) \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{B\sqrt{\cos(c+dx)}\sin(c+dx)(a+b\cos(c+dx))^{7/2}}{5bd} + \frac{(10Ab-3aB)\sqrt{\cos(c+dx)}\sin(c+dx)(a+b\cos(c+dx))^{5/2}}{4d} \right)$$

input `Int[((a + b*Cos[c + d*x])^(5/2)*(A + B*Cos[c + d*x]))/Sec[c + d*x]^(3/2),x]`

```

output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Sqrt[Cos[c + d*x]]*(a + b*Cos[c
+ d*x])^(7/2)*Sin[c + d*x])/(5*b*d) + (((10*A*b - 3*a*B)*Sqrt[Cos[c + d*x]
]*(a + b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(4*d) + ((50*a*A*b - 15*a^2*B
+ 64*b^2*B)*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(3
*d) + ((3*(50*a^2*A*b + 120*A*b^3 - 15*a^3*B + 172*a*b^2*B)*Sqrt[Cos[c + d
*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*d) + (-1/2*((2*(a - b)*Sqrt
[a + b]*(150*a^3*A*b + 2840*a*A*b^3 - 45*a^4*B + 1692*a^2*b^2*B + 1024*b^4
*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sq
rt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b
)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) + (2*Sqrt[a + b]*(45*a^4*B
- 30*a^3*b*(5*A + B) - 16*b^4*(45*A + 64*B) - 8*a*b^3*(355*A + 193*B) - 4*
a^2*b^2*(295*A + 423*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c +
d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 -
Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/d - (30*Sqrt
[a + b]*(10*a^4*A*b - 240*a^2*A*b^3 - 96*A*b^5 - 3*a^5*B - 40*a^3*b^2*B -
240*a*b^4*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c +
d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 -
Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(b*d))/b + (
(150*a^3*A*b + 2840*a*A*b^3 - 45*a^4*B + 1692*a^2*b^2*B + 1024*b^4*B)*Sqrt
[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]))/4)/6)/8)/(...

```

3.613.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3288 Int[Sqrt[(b_)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)
*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c
*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*Ellipti
cPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c +
d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 -
d^2, 0] && PosQ[(c + d)/b]

```

3.613.
$$\int \frac{(a+b \cos(c+dx))^{5/2}(A+B \cos(c+dx))}{\sec^{\frac{3}{2}}(c+dx)} dx$$

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGTQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3528 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n*Simp[a*A*d*(m + n + 2) + C*(b*c*m + a*d*(n + 1)) + (d*(A*b + a*B)*(m + n + 2) - C*(a*c - b*d*(m + n + 1)))*Sin[e + f*x] + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3540 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]))*Simp[2*a*A*d - C*(b*c - a*d) - 2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.613.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6969 vs. 2(767) = 1534.

Time = 23.12 (sec) , antiderivative size = 6970, normalized size of antiderivative = 8.31

method	result	size
parts	Expression too large to display	6970
default	Expression too large to display	7057

input `int((a+cos(d*x+c)*b)^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

3.613.5 Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorithm="fricas")`

output `Timed out`

3.613.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*cos(d*x+c))**(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.613. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sec^{3/2}(c+dx)} dx$

3.613.7 Maxima [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sec(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/sec(d*x + c)^(3/2), x)`

3.613.8 Giac [F]

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(B \cos(dx + c) + A)(b \cos(dx + c) + a)^{5/2}}{\sec(dx + c)^{3/2}} dx$$

input `integrate((a+b*cos(d*x+c))^(5/2)*(A+B*cos(d*x+c))/sec(d*x+c)^(3/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*(b*cos(d*x + c) + a)^(5/2)/sec(d*x + c)^(3/2), x)`

3.613.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \cos(c + dx))^{5/2} (A + B \cos(c + dx))}{\sec^{3/2}(c + dx)} dx = \int \frac{(A + B \cos(c + dx)) (a + b \cos(c + dx))^{5/2}}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(a + b*cos(c + d*x))^(5/2))/(1/cos(c + d*x))^(3/2), x)`

3.613. $\int \frac{(a+b \cos(c+dx))^{5/2} (A+B \cos(c+dx))}{\sec^{3/2}(c+dx)} dx$

3.614
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.614.1 Optimal result 5627
 3.614.2 Mathematica [A] (warning: unable to verify) 5628
 3.614.3 Rubi [A] (verified) 5628
 3.614.4 Maple [B] (verified) 5633
 3.614.5 Fricas [F] 5634
 3.614.6 Sympy [F(-1)] 5635
 3.614.7 Maxima [F] 5635
 3.614.8 Giac [F] 5635
 3.614.9 Mupad [F(-1)] 5636

3.614.1 Optimal result

Integrand size = 35, antiderivative size = 403

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2(a - b)\sqrt{a + b}(9a^2 A + 8Ab^2 - 10abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{a}}{15a^4 d \sqrt{\sec(c + dx)}} - \frac{2\sqrt{a + b}(8Ab^2 + a^2(9A - 5B) - 2ab(A + 5B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right)\right)}{15a^3 d \sqrt{\sec(c + dx)}} - \frac{2(4Ab - 5aB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{15a^2 d} + \frac{2A \sqrt{a + b \cos(c + dx)} \sec^{\frac{5}{2}}(c + dx) \sin(c + dx)}{5ad}$$

output

```
-2/15*(4*A*b-5*B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a^2
/d+2/5*A*sec(d*x+c)^(5/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d+2/15*(a-b)
*(9*A*a^2+8*A*b^2-10*B*a*b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a
+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1
/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^4/d/se
c(d*x+c)^(1/2)-2/15*(8*A*b^2+a^2*(9*A-5*B)-2*a*b*(A+5*B))*csc(d*x+c)*Ellip
ticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1
/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec
(d*x+c))/(a-b))^(1/2)/a^3/d/sec(d*x+c)^(1/2)
```

3.614.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.614.2 Mathematica [A] (warning: unable to verify)

Time = 13.68 (sec) , antiderivative size = 385, normalized size of antiderivative = 0.96

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= 2 \left(-\frac{\sqrt{\cos^2(\frac{1}{2}(c+dx)) \sec(c+dx)} (2(a+b)(9a^2A+8Ab^2-10abB) E(\arcsin(\tan(\frac{1}{2}(c+dx))) | \frac{-a+b}{a+b}) \sqrt{\frac{1}{1+\sec(c+dx)}} \sqrt{\frac{b+a \sec(c+dx)}{(a+b)(1+\sec(c+dx))}} - 2a(8A$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*(-((Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(2*(a + b)*(9*a^2*A + 8*A*b^2 - 10*a*b*B)*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x])])]) - 2*a*(8*A*b^2 + 2*a*b*(A - 5*B) + a^2*(9*A + 5*B))*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x])])]) + (9*a^2*A + 8*A*b^2 - 10*a*b*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2] + (a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]*((9*a^2*A + 8*A*b^2 - 10*a*b*B)*Sin[c + d*x] + a*(-4*A*b + 5*a*B + 3*a*A*Sec[c + d*x])*Tan[c + d*x]))/(15*a^3*d*Sqrt[a + b*Cos[c + d*x]])`

3.614.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 396, normalized size of antiderivative = 0.98, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{7}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{\csc(c + dx + \frac{\pi}{2})^{7/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

3.614. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3440 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{7}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \\
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{7/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \downarrow 3479 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{-2Ab\cos^2(c+dx)-3A\cos(c+dx)+4Ab-5aB}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{5a} + \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} \right) \\
& \downarrow 27 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} - \frac{\int \frac{-2Ab\cos^2(c+dx)-3A\cos(c+dx)+4Ab-5aB}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{5a} \right) \\
& \downarrow 3042 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} - \frac{\int \frac{-2Ab\sin(c+dx+\frac{\pi}{2})^2-3A\sin(c+dx+\frac{\pi}{2})+4Ab-5aB}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{5a} \right) \\
& \downarrow 3534 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} - \frac{2 \int \frac{-9Aa^2-10bBa+(2Ab+5aB)\cos(c+dx)a+8Ab^2}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a} + \frac{2(4Ab-5aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5a} \right) \\
& \downarrow 27 \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{5ad\cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{9Aa^2-10bBa}{\cos^{\frac{3}{2}}}}{5a} \right)
\end{aligned}$$

3.614. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{9Aa^2-10bBa}{\sin(c+dx)} dx}{5a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3477 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(9a^2A-10abB)}{\sin(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(9a^2A-10abB)}{\sin(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3295 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(9a^2A-10abB)}{\sin(c+dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3473 \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{5ad \cos^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2(a-b)\sqrt{a+b}(9a^2A-10abB)}{\sin(c+dx)} \right) \end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(7/2))/Sqrt[a + b*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(5*a*d*Cos[c + d*x]^(5/2)) - (-1/3*((2*(a - b)*Sqrt[a + b]*(9*a^2 *A + 8*A*b^2 - 10*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2 *Sqrt[a + b]*(8*A*b^2 + a^2*(9*A - 5*B) - 2*a*b*(A + 5*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]])/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/a + (2*(4*A*b - 5*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/(5*a)`

3.614.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_.) + (b_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

3.614.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[-2A(c-d) \frac{\tan(e+fx)}{f \sqrt{c^2+d^2}} \text{Rt}[(c+d)/b, 2] \sqrt{c(1+\csc(e+fx))/(c-d)} \sqrt{c(1-\csc(e+fx))/(c+d)} \text{EllipticE}[\text{ArcSin}[\sqrt{c+d \sin(e+fx)}/\sqrt{b \sin(e+fx)}] \text{Rt}[(c+d)/b, 2], -(c+d)/(c-d)], x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\& \text{PosQ}[(c+d)/b]$$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{c + d \sin(e + fx)}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[(A-B)/(a-b) \int 1/(\sqrt{a+b \sin(e+fx)} \sqrt{c+d \sin(e+fx)}), x], x] - \text{Simp}[(A*b - a*B)/(a-b) \int (1 + \sin(e+fx))/(a+b \sin(e+fx))^{3/2} \sqrt{c+d \sin(e+fx)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[A, B]$$`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx))^{n+1} (c + d \sin(e + fx))^{m+n} dx$$

$$\rightarrow \text{Simp}[(-A*b^2 - a*b*B) \cos(e+fx) (a + b \sin(e+fx))^{m+1} (c + d \sin(e+fx))^{m+n} / (f*(m+1)*(b*c - a*d)*(a^2 - b^2)), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 - b^2)) \int (a + b \sin(e+fx))^{m+1} (c + d \sin(e+fx))^{m+n} \text{Simp}[(a*A - b*B)*(b*c - a*d)*(m+1) + b*d*(A*b - a*B)*(m+n+2) + (A*b - a*B)*(a*d*(m+1) - b*c*(m+2))*\sin(e+fx) - b*d*(A*b - a*B)*(m+n+3)*\sin(e+fx)^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{RationalQ}[m] \&\& m < -1 \&\& ((\text{EqQ}[a, 0] \&\& \text{IntegerQ}[m] \&\& !\text{IntegerQ}[n]) || !(\text{IntegerQ}[2*n] \&\& \text{LtQ}[n, -1] \&\& ((\text{IntegerQ}[n] \&\& !\text{IntegerQ}[m]) || \text{EqQ}[a, 0])))$$`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.614.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3319 vs. $2(363) = 726$.

Time = 25.69 (sec) , antiderivative size = 3320, normalized size of antiderivative = 8.24

method	result	size
parts	Expression too large to display	3320
default	Expression too large to display	3350

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

3.614.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

output `-2/15*A/d*sec(d*x+c)^(7/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(-16*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4+4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^4+16*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^4-9*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3-8*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3+2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^2*b*cos(d*x+c)^3+8*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a*b^2*cos(d*x+c)^3+9*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*a^3*cos(d*x+c)^3-3*sin(d*x+c)*cos(d*x+c)*a^3+sin(d*x+c)*cos(d*x+c)^2*a^2*b-18*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+...`

3.614.5 Fracas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/sqrt(b*cos(d*x + c) + a), x)`

3.614. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{7}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

3.614.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(7/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.614.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/sqrt(b*cos(d*x + c) + a)
, x)`

3.614.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{7}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(7/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(7/2)/sqrt(b*cos(d*x + c) + a)
, x)`

3.614.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{7}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{7/2}}{\sqrt{a + b \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + b*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(7/2))/(a + b*cos(c + d*x))^(1/2), x)`

3.615
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.615.1 Optimal result 5637
 3.615.2 Mathematica [A] (verified) 5638
 3.615.3 Rubi [A] (verified) 5638
 3.615.4 Maple [B] (verified) 5642
 3.615.5 Fracas [F] 5643
 3.615.6 Sympy [F(-1)] 5644
 3.615.7 Maxima [F] 5644
 3.615.8 Giac [F] 5644
 3.615.9 Mupad [F(-1)] 5645

3.615.1 Optimal result

Integrand size = 35, antiderivative size = 330

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx =$$

$$\frac{2(a - b)\sqrt{a + b}(2Ab - 3aB)\sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}}}{3a^3 d \sqrt{\sec(c + dx)}} +$$

$$\frac{2\sqrt{a + b}(2Ab + a(A - 3B))\sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}}}{3a^2 d \sqrt{\sec(c + dx)}} +$$

$$\frac{2A\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3ad}$$

output

```
2/3*A*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a/d-2/3*(a-b)*(2*
A*b-3*B*a)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x
+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x
+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d/sec(d*x+c)^(1/2)+2/
3*(2*A*b+a*(A-3*B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2
)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(
1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d/sec(d*x+c)
^(1/2)
```

3.615.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.615.2 Mathematica [A] (verified)

Time = 11.15 (sec) , antiderivative size = 324, normalized size of antiderivative = 0.98

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2 \left((a + b \cos(c + dx))(aA + (-2Ab + 3aB) \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx) + \sqrt{\cos^2(\frac{1}{2}(c + dx)) \sec(c + dx)} \right)}{\dots}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*((a + b*Cos[c + d*x])*(a*A + (-2*A*b + 3*a*B)*Cos[c + d*x])*Sec[c + d*x]^(3/2)*Sin[c + d*x] + (Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(-2*A*b + 3*a*B)*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))] + 2*a*(-2*A*b + a*(A + 3*B))*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))] + (2*A*b - 3*a*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2]))/(3*a^2*d*Sqrt[a + b*Cos[c + d*x]])`

3.615.3 Rubi [A] (verified)Time = 1.17 (sec) , antiderivative size = 315, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc(c + dx + \frac{\pi}{2})^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

3.615. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow \text{3440} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{5}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
 & \downarrow \text{3479} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{2Ab-3aB-aA\cos(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a} + \frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{2Ab-3aB-aA\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{2Ab-3aB-aA\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{3a} \right) \\
 & \downarrow \text{3477} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(2Ab-3aB) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a+b)}{3a} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(2Ab-3aB) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a+b)}{3a} \right) \\
 & \downarrow \text{3295}
 \end{aligned}$$

3.615. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{(2Ab-3aB) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{a^2 d} \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A \sin(c+dx)\sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} - \frac{2(a-b)\sqrt{a+b}(2Ab-3aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a+b}}}{a^2 d} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/Sqrt[a + b*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(-1/3*((2*(a - b)*Sqrt[a + b]*(2*A*b - 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*Sqrt[a + b]*(2*A*b + a*(A - 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/a + (2*A*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))`

3.615.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.615. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

3.615.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$


```
rule 3479 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m +
1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e
+ f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +
2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)
*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n
}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Rat
ionalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(I
ntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0]
))
```

3.615.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1863 vs. $2(296) = 592$.

Time = 21.36 (sec) , antiderivative size = 1864, normalized size of antiderivative = 5.65

method	result	size
parts	Expression too large to display	1864
default	Expression too large to display	2308

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

3.615.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

```
output -2/3*A/d*sec(d*x+c)^(5/2)/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)*(cos(d*x+c)
)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*((a+cos(d*x+c)*b
)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a^2-2*cos(
d*x+c)^4*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*((a+cos(d*x
+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*a*b+2
*cos(d*x+c)^4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+c)/(1
+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*
a*b+2*cos(d*x+c)^4*((a+cos(d*x+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*(cos(d*x+
c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1
/2))*b^2+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+
c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*a^2
*cos(d*x+c)^3-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos
(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)
)*a*b*cos(d*x+c)^3+4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(
1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(
1/2))*a*b*cos(d*x+c)^3+4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)
*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a
+b))^(1/2))*b^2*cos(d*x+c)^3+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x
+c)*b)/(1+cos(d*x+c))/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)
/(a+b))^(1/2))*a^2*cos(d*x+c)^2-2*EllipticF(cot(d*x+c)-csc(d*x+c),(-a...
```

3.615.5 Fracas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fracas")
```

```
output integral((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/sqrt(b*cos(d*x + c) + a),
x)
```

3.615.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.615.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/sqrt(b*cos(d*x + c) + a)
, x)`

3.615.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/sqrt(b*cos(d*x + c) + a)
, x)`

3.615.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{\sqrt{a + b \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^(1/2), x)`

3.616
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.616.1 Optimal result 5646
 3.616.2 Mathematica [A] (verified) 5647
 3.616.3 Rubi [A] (verified) 5647
 3.616.4 Maple [B] (verified) 5650
 3.616.5 Fricas [F] 5651
 3.616.6 Sympy [F(-1)] 5651
 3.616.7 Maxima [F] 5651
 3.616.8 Giac [F] 5652
 3.616.9 Mupad [F(-1)] 5652

3.616.1 Optimal result

Integrand size = 35, antiderivative size = 270

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2A(a - b)\sqrt{a + b}\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{a^2 d \sqrt{\sec(c + dx)}} - \frac{2\sqrt{a + b}(A - B)\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{ad \sqrt{\sec(c + dx)}}$$

```
output 2*A*(a-b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d/sec(d*x+c)^(1/2)-2*(A-B)*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/sec(d*x+c)^(1/2)
```

3.616.2 Mathematica [A] (verified)

Time = 9.70 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.03

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= 2 \left(A(a + b \cos(c + dx)) \sqrt{\sec(c + dx)} \sin(c + dx) - \sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right) \sec(c + dx)} \left(2A(a + b) \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} E\left(\arcsin\left(\tan\left(\frac{c + dx}{2}\right)\right)\right) \right) \right)$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*(A*(a + b*Cos[c + d*x])*Sqrt[Sec[c + d*x]]*Sin[c + d*x] - (Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(2*A*(a + b)*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x])])]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)] - 2*a*(A + B)*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x])])]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)] + A*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2])/(a*d*Sqrt[a + b*Cos[c + d*x]])`

3.616.3 Rubi [A] (verified)Time = 0.87 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3440, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} (A + B \sin\left(c + dx + \frac{\pi}{2}\right))}{\sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx$$

$$\downarrow \text{3440}$$

3.616. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3477} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(A \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (A-B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(A \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (A-B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \\
& \quad \downarrow \text{3295} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(A \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(A-B)\cot(c+dx)}{a^2d} \right) \\
& \quad \downarrow \text{3473} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A(a-b)\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{a^2d} \right)
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/Sqrt[a + b*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*Sqrt[a + b]*(A - B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))`

3.616. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\cos(c+dx)}} dx$

3.616.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_ + (b_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(((b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_)*sin[(e_.) + (f_)*(x_)])/(((a_.) + (b_)*sin[(e_.) + (f_)*(x_)])^(3/2)*Sqrt[(c_.) + (d_)*sin[(e_.) + (f_)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

3.616.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 777 vs. 2(246) = 492.

Time = 18.68 (sec) , antiderivative size = 778, normalized size of antiderivative = 2.88

method	result
parts	$\frac{2A \left(-\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \sqrt{\frac{(\csc^2(dx+c))^a(1-\cos(dx+c))^2 - (\csc^2(dx+c))^b(1-\cos(dx+c))^2+a+b}{a+b}} F(\cot(dx+c) - \csc(dx+c)) \right)}{\dots}$
default	$\frac{2 \left(-\frac{(\csc^2(dx+c))(1-\cos(dx+c))^2+1}{(\csc^2(dx+c))(1-\cos(dx+c))^2-1} \right)^{\frac{3}{2}} \left((\csc^2(dx+c))(1-\cos(dx+c))^2-1 \right) \left(-A \sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \sqrt{\frac{(\csc^2(dx+c))^a(1-\cos(dx+c))^2 - (\csc^2(dx+c))^b(1-\cos(dx+c))^2+a+b}{a+b}} F(\cot(dx+c) - \csc(dx+c)) \right)}{\dots}$

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*b+csc(d*x+c)^3*(1-cos(d*x+c))^3*a-csc(d*x+c)^3*(1-cos(d*x+c))^3*b+a*(csc(d*x+c)-cot(d*x+c))+b*(csc(d*x+c)-cot(d*x+c)))*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1))^(3/2)/(csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/a-2*B/d*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))/(a+cos(d*x+c)*b)^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*sec(d*x+c)^(3/2)*(cos(d*x+c)^2+cos(d*x+c))`

$$3.616. \int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{\sqrt{a+b \cos(c+dx)}} dx$$

3.616.5 Fricas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a), x)`

3.616.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Timed out`

3.616.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a), x)`

3.616.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a), x)`

3.616.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{\sqrt{a + b \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^(1/2), x)`

$$3.617 \quad \int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx$$

3.617.1 Optimal result	5653
3.617.2 Mathematica [A] (verified)	5654
3.617.3 Rubi [A] (verified)	5654
3.617.4 Maple [A] (verified)	5657
3.617.5 Fricas [F]	5657
3.617.6 Sympy [F]	5658
3.617.7 Maxima [F]	5658
3.617.8 Giac [F]	5658
3.617.9 Mupad [F(-1)]	5659

3.617.1 Optimal result

Integrand size = 35, antiderivative size = 268

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2A\sqrt{a + b}\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{ad\sqrt{\sec(c + dx)}} - \frac{2\sqrt{a + b}B\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b}\sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{bd\sqrt{\sec(c + dx)}}$$

```
output 2*A*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/sec(d*x+c)^(1/2)-2*B*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/b/d/sec(d*x+c)^(1/2)
```

3.617.2 Mathematica [A] (verified)

Time = 3.02 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.59

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx$$

$$= \frac{2 \sqrt{\frac{a+b \cos(c+dx)}{(a+b)(1+\cos(c+dx))}} ((A - B) \text{EllipticF}(\arcsin(\tan(\frac{1}{2}(c + dx))), \frac{-a+b}{a+b}) + 2B \text{EllipticPi}(-1, \arcsin(\tan(\frac{1}{2}(c + dx))))}{d \sqrt{a + b \cos(c + dx)} \sqrt{\sec^2(\frac{1}{2}(c + dx))}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/Sqrt[a + b*Cos[c + d*x]],x]`

output `(2*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*((A - B)*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)])*Sqrt[Cos[c + d*x]*Sec[(c + d*x)/2]^2]*Sqrt[1 + Sec[c + d*x]])/(d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2])`

3.617.3 Rubi [A] (verified)Time = 0.79 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3440, 3042, 3485, 3042, 3288, 3295}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\csc(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3440}$$

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx$$

3.617. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
& \downarrow \text{3485} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(A \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + B \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(A \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \right) \\
& \downarrow \text{3288} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(A \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2B\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{ad} \right) \\
& \downarrow \text{3295} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2A\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{ad} \right)
\end{aligned}$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/Sqrt[a + b*Cos[c + d*x]],x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*A*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))`

3.617.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)])/Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_)] + (f_)*(x_)]*(g_))^(p_)*((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3485 `Int[((A_)] + (B_)*sin[(e_)] + (f_)*(x_)]/(Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)]*Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)]], x_Symbol] := Simp[B/d Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] - Simp[(B*c - A*d)/d Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.617.4 Maple [A] (verified)

Time = 16.18 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.68

method	result
default	$-\frac{2\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}}{d\sqrt{a+\cos(dx+c)b}}\left(AF\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)-BF\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)+2B\Pi\left(\cot(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)\right)$
parts	$-\frac{2A(1+\cos(dx+c))\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}F\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)(\sqrt{\sec(dx+c)})}{d\sqrt{a+\cos(dx+c)b}}+\frac{2B(\sqrt{\sec(dx+c)})}{d\sqrt{a+\cos(dx+c)b}}$

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output -2/d*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a
+b)^(1/2)*(A*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))-B*Elli
pticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))+2*B*EllipticPi(cot(d*x+c
)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))/(a+cos(d*x+c)*b)^(1/2)*sec(d*x+c)^(
1/2)*(1+cos(d*x+c))
```

3.617.5 Fricas [F]

$$\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx = \int \frac{(B\cos(dx+c)+A)\sqrt{\sec(dx+c)}}{\sqrt{b\cos(dx+c)+a}} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="fricas")
```

```
output integral((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/sqrt(b*cos(d*x + c) + a),
x)
```


3.617.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/sqrt(a + b*cos(c + d*x)), x)`

3.617.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/sqrt(b*cos(d*x + c) + a), x)`

3.617.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(B \cos(dx + c) + A)\sqrt{\sec(dx + c)}}{\sqrt{b \cos(dx + c) + a}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/sqrt(b*cos(d*x + c) + a), x)`

3.617.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{\sqrt{a + b \cos(c + dx)}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^(1/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^(1/2), x)`

$$3.618 \quad \int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sqrt{\sec(c+dx)}} dx$$

3.618.1 Optimal result	5660
3.618.2 Mathematica [A] (verified)	5661
3.618.3 Rubi [A] (verified)	5662
3.618.4 Maple [B] (verified)	5667
3.618.5 Fricas [F]	5668
3.618.6 Sympy [F]	5669
3.618.7 Maxima [F]	5669
3.618.8 Giac [F]	5669
3.618.9 Mupad [F(-1)]	5670

3.618.1 Optimal result

Integrand size = 35, antiderivative size = 487

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx =$$

$$\frac{(a - b)\sqrt{a + b}B\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right) \middle| -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{abd\sqrt{\sec(c + dx)}} -$$

$$+ \frac{\sqrt{a + b}B\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{bd\sqrt{\sec(c + dx)}} -$$

$$\frac{\sqrt{a + b}(2Ab - aB)\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b \sqrt{\cos(c + dx)}}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{b^2d\sqrt{\sec(c + dx)}} +$$

$$+ \frac{B \sin(c + dx)}{d\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} + \frac{aB\sqrt{\sec(c + dx)} \sin(c + dx)}{bd\sqrt{a + b \cos(c + dx)}}$$

output `B*sin(d*x+c)/d/(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+a*B*sin(d*x+c)*sec(d*x+c)^(1/2)/b/d/(a+b*cos(d*x+c))^(1/2)-(a-b)*B*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a/b/d/sec(d*x+c)^(1/2)+B*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/b/d/sec(d*x+c)^(1/2)-(2*A*b-B*a)*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/b^2/d/sec(d*x+c)^(1/2)`

3.618.2 Mathematica [A] (verified)

Time = 9.36 (sec) , antiderivative size = 610, normalized size of antiderivative = 1.25

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$$

$$= \frac{\sec\left(\frac{1}{2}(c + dx)\right) \sqrt{\sec(c + dx)} \left(8(a + b)B \cos^3\left(\frac{1}{2}(c + dx)\right) \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} E\left(\arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right)\right)\right)}{1}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]),x]`

output

```
(Sec[(c + d*x)/2]*Sqrt[Sec[c + d*x]]*(8*(a + b)*B*Cos[(c + d*x)/2]^3*Sqrt[
(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c
+ d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)] - 16*A*b*Cos[
(c + d*x)/2]^3*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[
(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d
*x]))] + 24*A*b*Cos[(c + d*x)/2]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]],
(-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/
((a + b)*(1 + Sec[c + d*x]))] - 12*a*B*Cos[(c + d*x)/2]*EllipticPi[-1, Arc
Sin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqr
t[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))] + 8*A*b*Cos[(3*(c + d
*x))/2]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1
+ Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x
]))] - 4*a*B*Cos[(3*(c + d*x))/2]*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]],
(-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])
/((a + b)*(1 + Sec[c + d*x]))] - 2*a*B*Sin[(c + d*x)/2] + 2*b*B*Sin[(c + d
*x)/2] + 2*a*B*Sin[(3*(c + d*x))/2] - b*B*Sin[(3*(c + d*x))/2] + b*B*Sin[(
5*(c + d*x))/2]))/(4*b*d*Sqrt[a + b*Cos[c + d*x]])
```

3.618.3 Rubi [A] (verified)

Time = 1.96 (sec) , antiderivative size = 457, normalized size of antiderivative = 0.94, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3482, 3042, 3530, 3042, 3288, 3472, 25, 27, 3042, 3280, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)} (A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.618. $\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx$$

↓ 3482

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \int \frac{(2Ab-aB)\cos^2(c+dx)+2aA\cos(c+dx)+aB}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \int \frac{(2Ab-aB)\sin(c+dx+\frac{\pi}{2})^2+2aA\sin(c+dx+\frac{\pi}{2})+aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx + \frac{B\sin(c+dx)}{d\sqrt{a+b\cos(c+dx)}} \right)$$

↓ 3530

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\int \frac{B\cos(c+dx)a^2+bBa}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{b} + \frac{(2Ab-aB) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx}{b} \right) + \frac{B\sin(c+dx)}{d\sqrt{a+b\cos(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\int \frac{B\sin(c+dx+\frac{\pi}{2})a^2+bBa}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} + \frac{(2Ab-aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) + \frac{B\sin(c+dx)}{d\sqrt{a+b\cos(c+dx)}} \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\int \frac{B\sin(c+dx+\frac{\pi}{2})a^2+bBa}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{d\sqrt{a+b\cos(c+dx)}} \right) + \frac{B\sin(c+dx)}{d\sqrt{a+b\cos(c+dx)}} \right)$$

↓ 3472

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\int \frac{a(a^2-b^2)B}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} + \frac{2aB\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{2\sqrt{a+b}(2Ab-aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{d\sqrt{a+b\cos(c+dx)}} \right) + \frac{B\sin(c+dx)}{d\sqrt{a+b\cos(c+dx)}} \right)$$

↓ 25

3.618. $\int \frac{A+B\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - \frac{\int \frac{a(a^2-b^2)B}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{a^2-b^2}}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx)}{b} \right) \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx)}{b} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \int \frac{1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx)}{b} \right) \right)$$

↓ 3280

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \left(\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx - \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx \right)}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx)}{b} \right) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \left(\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx \right)}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx)}{b} \right) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - aB \left(\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b} \cot(c+dx)}{b} \right)}{b} - \frac{2\sqrt{a+b}(2Ab-aB) \cot(c+dx)}{b} \right) \right)$$

↓ 3473

3.618. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)}\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{1}{2} \left(\frac{2aB \sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - aB \left(\frac{2(a-b)\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{a^2d} \right) \right) \right)$$

input `Int[(A + B*Cos[c + d*x])/(Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + b*Cos[c + d*x]]) + ((-2*Sqrt[a + b]*(2*A*b - a*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b^2*d) + (-a*B*((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d)) + (2*a*B*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))/b)/2)`

3.618.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3280 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[1/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[b/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] / ; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] / ; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] / ; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^m Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] / ; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]]), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] / ; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3482 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[-2*B*Cos[e + f*x]*Sqrt[a + b*Sin[e + f*x]]*((c + d*Sin[e + f*x])^n/(f*(2*
n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*Sin[e + f*x])^(n - 1)/Sqrt[a
+ b*Sin[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d
)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*Sin[e + f*x] + (A*b*d*(2*n + 3) + B*
(a*d + 2*b*c*n))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A,
B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Eq
Q[n^2, 1/4]
```

```
rule 3530 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)]^
2)/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*((a_) + (b_)*sin[(e_) + (f_)*(x_
)]))^(3/2), x_Symbol] := Simp[C/(b*d) Int[Sqrt[d*Sin[e + f*x]]/Sqrt[a + b
*Sin[e + f*x]], x], x] + Simp[1/b Int[(A*b + (b*B - a*C)*Sin[e + f*x])/((
a + b*Sin[e + f*x])^(3/2)*Sqrt[d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, d,
e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.618.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 957 vs. $2(443) = 886$.

Time = 16.82 (sec) , antiderivative size = 958, normalized size of antiderivative = 1.97

method	result	size
parts	Expression too large to display	958
default	Expression too large to display	1344

```
input int((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

$$3.618. \quad \int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sqrt{\sec(c+dx)}} dx$$

output `2*A/d/(a+cos(d*x+c)*b)^(1/2)*(EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b)))^(1/2))-2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)/sec(d*x+c)^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-B/d/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(1/2)*(EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)+EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b*cos(d*x+c)-2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a*cos(d*x+c)+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-4*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-2*sec(d*x+c)*(co...`

3.618.5 Fracas [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.618.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a + b*cos(c + d*x))*sqrt(sec(c + d*x))), x)`

3.618.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.618.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)`

3.618.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c + dx)}} \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(1/2)), x)`

$$3.619 \quad \int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.619.1 Optimal result	5671
3.619.2 Mathematica [B] (verified)	5672
3.619.3 Rubi [A] (verified)	5673
3.619.4 Maple [B] (verified)	5679
3.619.5 Fricas [F]	5680
3.619.6 Sympy [F]	5681
3.619.7 Maxima [F]	5681
3.619.8 Giac [F]	5681
3.619.9 Mupad [F(-1)]	5682

3.619.1 Optimal result

Integrand size = 35, antiderivative size = 539

$$\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx =$$

$$\frac{(a-b)\sqrt{a+b}(4Ab-3aB)\sqrt{\cos(c+dx)} \csc(c+dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4ab^2d\sqrt{\sec(c+dx)}} -$$

$$\frac{\sqrt{a+b}(4Ab-3aB+2bB)\sqrt{\cos(c+dx)} \csc(c+dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{4b^2d\sqrt{\sec(c+dx)}} +$$

$$\frac{\sqrt{a+b}(4aAb-3a^2B-4b^2B)\sqrt{\cos(c+dx)} \csc(c+dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{4b^3d\sqrt{\sec(c+dx)}} +$$

$$\frac{B\sqrt{a+b \cos(c+dx)} \sin(c+dx)}{2bd\sqrt{\sec(c+dx)}} +$$

$$\frac{(4Ab-3aB)\sqrt{a+b \cos(c+dx)} \sqrt{\sec(c+dx)} \sin(c+dx)}{4b^2d}$$

3.619. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$

output $\frac{1}{2}B\sin(dx+c)(a+b\cos(dx+c))^{1/2}/b/d/\sec(dx+c)^{1/2}+1/4*(4A*b-3B*a)*\sin(dx+c)*(a+b\cos(dx+c))^{1/2}*\sec(dx+c)^{1/2}/b^2/d-1/4*(a-b)*(4A*b-3B*a)*\csc(dx+c)*\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c)))/(a+b)^{1/2}*(a*(1+\sec(dx+c)))/(a-b)^{1/2}/a/b^2/d/\sec(dx+c)^{1/2}+1/4*(4A*b-3B*a+2*B*b)*\csc(dx+c)*\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c)))/(a+b)^{1/2}*(a*(1+\sec(dx+c)))/(a-b)^{1/2}/b^2/d/\sec(dx+c)^{1/2}+1/4*(4A*a*b-3B*a^2-4B*b^2)*\csc(dx+c)*\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2},(a+b)/b,((-a-b)/(a-b))^{1/2})*(a+b)^{1/2}*\cos(dx+c)^{1/2}*(a*(1-\sec(dx+c)))/(a+b)^{1/2}*(a*(1+\sec(dx+c)))/(a-b)^{1/2}/b^3/d/\sec(dx+c)^{1/2}$

3.619.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1157 vs. $2(539) = 1078$.

Time = 15.45 (sec) , antiderivative size = 1157, normalized size of antiderivative = 2.15

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/(Sqrt[a + b*Cos[c + d*x]]*Sec[c + d*x]^(3/2)),x]`

output

```
(B*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*Sin[2*(c + d*x)]/(4*b*d) +
(Sqrt[(1 - Tan[(c + d*x)/2]^2)^(-1)]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 -
b*Tan[(c + d*x)/2]^2)/(1 + Tan[(c + d*x)/2]^2)]*(-4*a*A*b*Tan[(c + d*x)/2
] - 4*A*b^2*Tan[(c + d*x)/2] + 3*a^2*B*Tan[(c + d*x)/2] + 3*a*b*B*Tan[(c +
d*x)/2] + 8*A*b^2*Tan[(c + d*x)/2]^3 - 6*a*b*B*Tan[(c + d*x)/2]^3 + 4*a*A
*b*Tan[(c + d*x)/2]^5 - 4*A*b^2*Tan[(c + d*x)/2]^5 - 3*a^2*B*Tan[(c + d*x)
/2]^5 + 3*a*b*B*Tan[(c + d*x)/2]^5 + 8*a*A*b*EllipticPi[-1, ArcSin[Tan[(c
+ d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a
*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 6*a^2*B*EllipticPi[
-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^
2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 8
*b^2*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 -
Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/
2]^2)/(a + b)] + 8*a*A*b*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)
/(a + b)]*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*
Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 6*a^2*B*EllipticPi[-
1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Tan[(c + d*x)/2]^2*Sqrt[1 -
Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/
2]^2)/(a + b)] - 8*b^2*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)
/(a + b)]*Tan[(c + d*x)/2]^2*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + ...
```

3.619.3 Rubi [A] (verified)

Time = 2.25 (sec) , antiderivative size = 501, normalized size of antiderivative = 0.93, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3540, 25, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{\frac{3}{2}} \sqrt{a + b \sin\left(c + dx + \frac{\pi}{2}\right)}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{3}{2}}(c + dx) (A + B \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx$$

3.619. $\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx \\
 & \downarrow \text{3469} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(4Ab-3aB)\cos^2(c+dx)+2bB\cos(c+dx)+aB}{2\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{2b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(4Ab-3aB)\cos^2(c+dx)+2bB\cos(c+dx)+aB}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{4b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(4Ab-3aB)\sin(c+dx+\frac{\pi}{2})^2+2bB\sin(c+dx+\frac{\pi}{2})+aB}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{4b} + \frac{B\sin(c+dx)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}{2bd} \right) \\
 & \downarrow \text{3540} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(-3Ba^2+4Aba-4b^2B)\cos^2(c+dx)-2abB\cos(c+dx)+a(4Ab-3aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{4b} + \frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} \right) \\
 & \downarrow \text{25} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{(-3Ba^2+4Aba-4b^2B)\cos^2(c+dx)-2abB\cos(c+dx)+a(4Ab-3aB)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{4b} \right) \\
 & \downarrow \text{3042}
 \end{aligned}$$

3.619. $\int \frac{A+B\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{(-3Ba^2+4Aba-4b^2B)\sin(c+dx+\frac{\pi}{2})^2-2abB\sin(c+dx+\frac{\pi}{2})+a(4Aa-3Ba^2)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b}}{4b} \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{(-3a^2B+4aAb-4b^2B)\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx + \int \frac{a(4Ab-3aB)-2abB}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{2b}}{4b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{(-3a^2B+4aAb-4b^2B)\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{a(4Ab-3aB)-2abB}{\sin(c+dx+\frac{\pi}{2})} dx}{2b}}{4b} \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{\int \frac{a(4Ab-3aB)-2abB\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(-3a^2B+4aAb-4b^2B)}{\sin(c+dx+\frac{\pi}{2})}}{2b}}{4b} \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(4Ab-3aB)\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - a(-3aB+4Ab+2a^2)}{2b}}{4b} \right)$$

↓ 3042

3.619. $\int \frac{A+B\cos(c+dx)}{\sqrt{a+b\cos(c+dx)}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{-a(-3aB+4Ab+2bB)\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+}{\dots} \right)$$

3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{a(4Ab-3aB)\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2\sqrt{a+b}}{\dots}}{\dots} \right)$$

3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(4Ab-3aB)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{bd\sqrt{\cos(c+dx)}} - \frac{2\sqrt{a+b}(-3a^2B+4aAb-4b^2B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{bd} \right)$$

```
input Int[(A + B*cos[c + d*x])/(sqrt[a + b*cos[c + d*x]]*sec[c + d*x]^(3/2)),x]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((B*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(2*b*d) + (-1/2*((2*(a - b)*Sqrt[a + b]*(4*A*b - 3*a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*Sqrt[a + b]*(4*A*b - 3*a*B + 2*b*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/d - (2*Sqrt[a + b]*(4*A*A*b - 3*a^2*B - 4*b^2*B)*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/b + ((4*A*b - 3*a*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]]))/(4*b))
```

3.619. $\int \frac{A+B \cos(c+dx)}{\sqrt{a+b \cos(c+dx)} \sec^{\frac{3}{2}}(c+dx)} dx$

3.619.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_)] + (f_)*(x_)])/Sqrt[(c_)] + (d_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]]/Rt[(c + d)/b, 2], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_)] + (f_)*(x_)])*Sqrt[(a_)] + (b_)*sin[(e_)] + (f_)*(x_)], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_)] + (f_)*(x_))*(g_))^(p_)*((a_)] + (b_)*sin[(e_)] + (f_)*(x_)]^(m_)*((c_)] + (d_)*sin[(e_)] + (f_)*(x_)]^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3469 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m + n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2)/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))]*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.619.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2479 vs. $2(485) = 970$.

Time = 17.49 (sec) , antiderivative size = 2480, normalized size of antiderivative = 4.60

method	result	size
parts	Expression too large to display	2480
default	Expression too large to display	2491

```
input int((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(1/2),x,method=_RET
URNVERBOSE)
```

output `-A/d/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(3/2)*(sec(d*x+c)^2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+sec(d*x+c)^2*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-2*sec(d*x+c)^2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+2*sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+2*sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-4*sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-2*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a-tan(d*x+c)*a-b*sin(d*x+c))/b+1/4*B/d/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(3/2)*(3*((a...`

3.619.5 Fracas [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.619.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/(sqrt(a + b*cos(c + d*x))*sec(c + d*x)**(3/2)), x)`

3.619.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.619.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{\sqrt{b \cos(dx + c) + a} \sec(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))/sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/(sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)`

3.619.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} \sqrt{a + b \cos(c + dx)}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(1/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(1/2)), x)`

3.620
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.620.1 Optimal result	5683
3.620.2 Mathematica [A] (warning: unable to verify)	5684
3.620.3 Rubi [A] (verified)	5684
3.620.4 Maple [B] (warning: unable to verify)	5689
3.620.5 Fricas [F]	5690
3.620.6 Sympy [F(-1)]	5691
3.620.7 Maxima [F]	5691
3.620.8 Giac [F]	5691
3.620.9 Mupad [F(-1)]	5692

3.620.1 Optimal result

Integrand size = 35, antiderivative size = 433

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx =$$

$$\frac{2(5a^2 Ab - 8Ab^3 - 3a^3 B + 6ab^2 B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a^4 \sqrt{a + bd} \sqrt{\sec(c + dx)}} +$$

$$\frac{2(a + 2b)(4Ab + a(A - 3B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a^3 \sqrt{a + bd} \sqrt{\sec(c + dx)}} +$$

$$\frac{2b(Ab - aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{a(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}} +$$

$$\frac{2(a^2 A - 4Ab^2 + 3abB) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3a^2 (a^2 - b^2) d}$$

output

```
2*b*(A*b-B*a)*sec(d*x+c)^(3/2)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+2/3*(A*a^2-4*A*b^2+3*B*a*b)*sec(d*x+c)^(3/2)*sin(d*x+c)*(a+b*cos(d*x+c))^(1/2)/a^2/(a^2-b^2)/d-2/3*(5*A*a^2*b-8*A*b^3-3*B*a^3+6*B*a*b^2)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^4/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)+2/3*(a+2*b)*(4*A*b+a*(A-3*B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^3/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)
```

3.620.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.620.2 Mathematica [A] (warning: unable to verify)

Time = 16.32 (sec) , antiderivative size = 498, normalized size of antiderivative = 1.15

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right) \sec(c + dx)} \left(-2(a + b)(-5a^2Ab + 8Ab^3 + 3a^3B - 6ab^2B)\right)}{3a^3(a^2 - b^2)} + \frac{2(-Ab^3 \sin(c + dx) + ab^2B \sin(c + dx))}{a^2(a^2 - b^2)(a + b \cos(c + dx))} + \frac{2A \tan\left(\frac{c + dx}{2}\right)}{3a^2} + \frac{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2(-5a^2Ab + 8Ab^3 + 3a^3B - 6ab^2B) \sin(c + dx)}{3a^3(a^2 - b^2)} + \frac{2(-Ab^3 \sin(c + dx) + ab^2B \sin(c + dx))}{a^2(a^2 - b^2)(a + b \cos(c + dx))} + \frac{2A \tan\left(\frac{c + dx}{2}\right)}{3a^2}\right)}{d}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x])^(3/2), x]`

output `(2*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(-5*a^2*A*b + 8*A*b^3 + 3*a^3*B - 6*a*b^2*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*a*(a^2 - a*b - 2*b^2)*(-4*A*b + a*(A + 3*B))*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - (-5*a^2*A*b + 8*A*b^3 + 3*a^3*B - 6*a*b^2*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2]))/(3*a^3*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2]) + (Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*(-5*a^2*A*b + 8*A*b^3 + 3*a^3*B - 6*a*b^2*B)*Sin[c + d*x])/(3*a^3*(a^2 - b^2)) + (2*(-(A*b^3*Sin[c + d*x]) + a*b^2*B*Sin[c + d*x]))/(a^2*(a^2 - b^2)*(a + b*Cos[c + d*x])) + (2*A*Tan[c + d*x])/(3*a^2)))/d`

3.620.3 Rubi [A] (verified)

Time = 1.93 (sec) , antiderivative size = 433, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

3.620. $\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
& \int \frac{\csc\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(A+B \sin\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(a+b \sin\left(c+dx+\frac{\pi}{2}\right)\right)^{3/2}} dx \\
& \quad \downarrow \text{3440} \\
& \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{A+B \cos(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \left(a+b \cos(c+dx)\right)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{A+B \sin\left(c+dx+\frac{\pi}{2}\right)}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \left(a+b \sin\left(c+dx+\frac{\pi}{2}\right)\right)^{3/2}} dx \\
& \quad \downarrow \text{3479} \\
& \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{Aa^2+3bBa-(Ab-aB) \cos(c+dx)a-4Ab^2+2b(Ab-aB) \cos^2(c+dx)}{2 \cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{\int \frac{Aa^2+3bBa-(Ab-aB) \cos(c+dx)a-4Ab^2+2b(Ab-aB) \cos^2(c+dx)}{\cos^{\frac{5}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{\int \frac{Aa^2+3bBa-(Ab-aB) \sin\left(c+dx+\frac{\pi}{2}\right)a-4Ab^2+2b(Ab-aB) \sin\left(c+dx+\frac{\pi}{2}\right)^2}{\sin\left(c+dx+\frac{\pi}{2}\right)^{5/2} \sqrt{a+b \sin\left(c+dx+\frac{\pi}{2}\right)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)}{ad(a^2-b^2) \cos^{\frac{3}{2}}(c+dx)} \right) \\
& \quad \downarrow \text{3534} \\
& \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \left(\frac{2 \int -\frac{-3Ba^3+5Aba^2+6b^2Ba-(Aa^2-3bBa+2Ab^2) \cos(c+dx)a-8Ab^3}{2 \cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{3a} + \frac{2(a^2A+3abB-4Ab^2) \sin(c+dx) \sqrt{a+b \cos(c+dx)}}{3ad \cos^{\frac{3}{2}}(c+dx)} \right)}{a(a^2-b^2)} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.620. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-3Ba^3+5Aba^2+6b^2Ba-(Aa^2-3bBa+2Ab^2)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{3a} \right) \frac{1}{a(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-3Ba^3+5Aba^2+6b^2Ba-(Aa^2-3bBa+2Ab^2)\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{3a} \right) \frac{1}{a(a^2-b^2)}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}}{3a} \right) \frac{1}{a(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int \frac{\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{3a} \right) \frac{1}{a(a^2-b^2)}$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^3B+5a^2Ab+6ab^2B-8Ab^3)\int \frac{\sin(c+dx)}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}{3a} \right) \frac{1}{a(a^2-b^2)}$$

↓ 3473

3.620. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}+\frac{2(a^2A+3abB-4Ab^2)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad\cos^{\frac{3}{2}}(c+dx)}\right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x])^(3/2), x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]) + (-1/3*((2*(a - b)*Sqrt[a + b]*(5*a^2*A*b - 8*A*b^3 - 3*a^3*B + 6*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)])*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(a + 2*b)*(a*A + 4*A*b - 3*a*B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)])*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/a + (2*(a^2*A - 4*A*b^2 + 3*a*b*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(3*a*d*Cos[c + d*x]^(3/2)))/(a*(a^2 - b^2))`

3.620.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)])*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

3.620. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.620.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 3849 vs. $2(395) = 790$.

Time = 25.29 (sec) , antiderivative size = 3850, normalized size of antiderivative = 8.89

method	result	size
parts	Expression too large to display	3850
default	Expression too large to display	4714

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RET
URNVERBOSE)
```

3.620.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

output $\frac{2}{3}A/d * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)/(\csc(dx+c)^2(1-\cos(dx+c))^2-1)^{5/2} * (\csc(dx+c)^2(1-\cos(dx+c))^2-1)^{5/2} * \text{EllipticF}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a^3 * b * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} * (1-\cos(dx+c))^2 - 5 * \text{EllipticF}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a^3 * b * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} + 2 * \text{EllipticF}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a^2 * b^2 * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} + 8 * \text{EllipticF}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a * b^3 * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} + 5 * \text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a^3 * b * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} + 5 * \text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a^2 * b^2 * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} - 8 * \text{EllipticE}(\cot(dx+c)-\csc(dx+c), (-a-b)/(a+b))^{1/2} * a * b^3 * (-\csc(dx+c)^2(1-\cos(dx+c))^2+1)^{1/2} * ((\csc(dx+c)^2 * a * (1-\cos(dx+c))^2 - \csc(dx+c)^2 * b * (1-\cos(dx+c))^2 + a+b)/(a+b))^{1/2} - \csc(dx+c)^2 * E...$

3.620.5 Fracas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(dx+c))*sec(dx+c)^(5/2)/(a+b*cos(dx+c))^(3/2),x, algo
rithm="fracas")`

output `integral((B*cos(dx + c) + A)*sqrt(b*cos(dx + c) + a)*sec(dx + c)^(5/2)/
(b^2*cos(dx + c)^2 + 2*a*b*cos(dx + c) + a^2), x)`

3.620.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.620.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^(3/
2), x)`

3.620.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^(3/
2), x)`

3.620.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^(3/2), x)`

3.621
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.621.1 Optimal result 5693
 3.621.2 Mathematica [A] (warning: unable to verify) 5694
 3.621.3 Rubi [A] (verified) 5694
 3.621.4 Maple [B] (warning: unable to verify) 5698
 3.621.5 Fricas [F] 5699
 3.621.6 Sympy [F(-1)] 5700
 3.621.7 Maxima [F] 5700
 3.621.8 Giac [F] 5700
 3.621.9 Mupad [F(-1)] 5701

3.621.1 Optimal result

Integrand size = 35, antiderivative size = 345

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2(a^2 A - 2Ab^2 + abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}}{\sqrt{a+b \cos(c+dx)}}\right)\right) + 2(2Ab + a(A - B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{a^2 \sqrt{a + b d} \sqrt{\sec(c + dx)}} + \frac{2b(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{a(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

output

```
2*b*(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)+2*(A*a^2-2*A*b^2+B*a*b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)-2*(2*A*b+a*(A-B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)
```

3.621.2 Mathematica [A] (warning: unable to verify)

Time = 13.78 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.09

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2 \left(\sqrt{2} \sqrt{\frac{1}{1 + \cos(c + dx)}} (aA(a^2 - b^2) + b(a^2A - 2Ab^2 + abB)) \cos(c + dx) \right)}{(a + b \cos(c + dx))^{3/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^(3/2), x]`

output `(2*(Sqrt[2]*Sqrt[(1 + Cos[c + d*x])^(-1)]*(a*A*(a^2 - b^2) + b*(a^2*A - 2*A*b^2 + a*b*B)*Cos[c + d*x])*Sqrt[Sec[c + d*x]]*Sin[c + d*x] - Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(2*(a + b)*(a^2*A - 2*A*b^2 + a*b*B)*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))] - 2*a*(a + b)*(-2*A*b + a*(A + B))*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))]) + (a^2*A - 2*A*b^2 + a*b*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2]))/(a^2*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2])`

3.621.3 Rubi [A] (verified)Time = 1.38 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx$$

↓ 3440

3.621. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$

$$\begin{aligned} & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\ & \quad \downarrow \text{3479} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{Aa^2+bBa-(Ab-aB)\cos(c+dx)a-2Ab^2}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\ & \quad \downarrow \text{27} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{Aa^2+bBa-(Ab-aB)\cos(c+dx)a-2Ab^2}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{Aa^2+bBa-(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-2Ab^2}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)\sin(c+dx)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\ & \quad \downarrow \text{3477} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a^2A+abB-2Ab^2) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b)(a(A-B)+2Ab) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a(a^2-b^2)} \right) \\ & \quad \downarrow \text{3042} \\ & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a^2A+abB-2Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(a(A-B)+2Ab) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a(a^2-b^2)} \right) \\ & \quad \downarrow \text{3295} \end{aligned}$$

3.621. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+b\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a^2A + abB - 2Ab^2) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(a(A-B)+2Ab) \cot(c+dx)}{a(a^2-b^2)}}{a^2d} \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a-b)\sqrt{a+b}(a^2A+abB-2Ab^2) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{a^2d} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*(a - b)*Sqrt[a + b]*(a^2*A - 2*A*b^2 + a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(2*A*b + a*(A - B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a*(a^2 - b^2)) + (2*b*(A*b - a*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))`

3.621.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`


```
rule 3479 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin
[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2)), x] + Simp[1/((m +
1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e
+ f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +
2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)
*(m + n + 3)*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n
}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Rat
ionalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(I
ntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0]
))
```

3.621.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1810 vs. $2(317) = 634$.

Time = 22.10 (sec) , antiderivative size = 1811, normalized size of antiderivative = 5.25

method	result	size
default	Expression too large to display	1811
parts	Expression too large to display	2028

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RET
URNVERBOSE)
```

output

```
-2/d*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)
)^(3/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2-1)*(-A*EllipticF(cot(d*x+c)-csc(d*x
+c),(-a-b)/(a+b))^(1/2))*a^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((c
sc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))
^(1/2)+A*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b*(-csc
(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(
d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)+2*A*EllipticF(cot(d*x+c)-csc
(d*x+c),(-a-b)/(a+b))^(1/2))*a*b^2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/
2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/
(a+b))^(1/2)+A*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a^3*(
-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-
csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)+A*EllipticE(cot(d*x+c)-c
sc(d*x+c),(-a-b)/(a+b))^(1/2))*a^2*b*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(
1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b
)/(a+b))^(1/2)-2*A*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a
*b^2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+
c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)-2*A*EllipticE(cot(
d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^3*(-csc(d*x+c)^2*(1-cos(d*x+c))^
2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))
^2+a+b)/(a+b))^(1/2)+csc(d*x+c)^3*A*a^3*(1-cos(d*x+c))^3-csc(d*x+c)^3*A...
```

3.621.5 Fracas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)/
(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2), x)`

3.621.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.621.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(3/
2), x)`

3.621.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(3/
2), x)`

3.621.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{3/2}}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^(3/2), x)`

3.622
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{3/2}} dx$$

3.622.1 Optimal result 5702
 3.622.2 Mathematica [A] (verified) 5703
 3.622.3 Rubi [A] (verified) 5703
 3.622.4 Maple [B] (warning: unable to verify) 5706
 3.622.5 Fricas [F] 5707
 3.622.6 Sympy [F] 5708
 3.622.7 Maxima [F] 5708
 3.622.8 Giac [F] 5708
 3.622.9 Mupad [F(-1)] 5709

3.622.1 Optimal result

Integrand size = 35, antiderivative size = 324

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2(Ab - aB)\sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right)}{a^2 \sqrt{a + bd} \sqrt{\sec(c + dx)}} + \frac{2(A + B)\sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{a \sqrt{a + bd} \sqrt{\sec(c + dx)}} - \frac{2(Ab - aB)\sqrt{\sec(c + dx)} \sin(c + dx)}{(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

output

```
-2*(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(1/2)
)+2*(A*b-B*a)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(
d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b
))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)
+2*(A+B)*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c
))^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1
/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)
```

3.622.2 Mathematica [A] (verified)

Time = 9.27 (sec) , antiderivative size = 305, normalized size of antiderivative = 0.94

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2 \left(\frac{b(Ab - aB) \sin(c + dx)}{\sqrt{\sec(c + dx)}} + \frac{\sqrt{\cos^2(\frac{1}{2}(c + dx)) \sec(c + dx)} (2(a + b)(-Ab + aB)E(\arcsin(\frac{\sqrt{\sec(c + dx)} \sin(\frac{1}{2}(c + dx))}{\sqrt{a + b \cos(c + dx)}}))}{\sqrt{\sec(c + dx)}} \right)}{(a + b \cos(c + dx))^{3/2}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^(3/2),x]`

output `(2*((b*(A*b - a*B)*Sin[c + d*x])/Sqrt[Sec[c + d*x]] + (Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(2*(a + b)*(-A*b) + a*B)*EllipticE[ArcSin[Tan[(c + d*x)/2]]], (-a + b)/(a + b))*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))] + 2*a*(a + b)*(A - B)*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[(1 + Sec[c + d*x])^(-1)]*Sqrt[(b + a*Sec[c + d*x])/((a + b)*(1 + Sec[c + d*x]))] - (A*b - a*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/Sqrt[Sec[(c + d*x)/2]^2])/((a^3 - a*b^2)*d*Sqrt[a + b*Cos[c + d*x]])`

3.622.3 Rubi [A] (verified)Time = 1.24 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 3440, 3042, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\csc(c + dx + \frac{\pi}{2})}(A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx \\ & \quad \downarrow \text{3440} \\ & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{A + B \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx \end{aligned}$$

3.622. $\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
& \downarrow \text{3472} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{Ab-aB+(aA-bB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} - \frac{2(Ab-aB)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{Ab-aB+(aA-bB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} - \frac{2(Ab-aB)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\
& \downarrow \text{3477} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab-aB) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + (a-b)(A+B) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} \right) \\
& \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a-b)(A+B) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + (Ab-aB) \int \frac{\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} \right) \\
& \downarrow \text{3295} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab-aB) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(A+B)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{a^2-b^2}}{a^2-b^2} \right) \\
& \downarrow \text{3473}
\end{aligned}$$

3.622. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b\cos(c+dx))^{3/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a-b)\sqrt{a+b}(Ab-aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)-\frac{a+b}{a-b}}{a^2d} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^(3/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*(((2*(a - b)*Sqrt[a + b]*(A*b - a*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]], -(a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(A + B)*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]], -(a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a^2 - b^2) - (2*(A*b - a*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))`

3.622.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^p*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^m*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`


```
rule 3472 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] :> Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Ssin[e + f*x]]*Sqrt[d*Ssin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Ssin[e + f*x]]*(d*Ssin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]
```

```
rule 3473 Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Ssin[e + f*x]]/Sqrt[b*Ssin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]
```

```
rule 3477 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Ssin[e + f*x]]*Sqrt[c + d*Ssin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Ssin[e + f*x])^(3/2)*Sqrt[c + d*Ssin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]
```

3.622.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1315 vs. $2(296) = 592$.

Time = 18.91 (sec) , antiderivative size = 1316, normalized size of antiderivative = 4.06

method	result	size
default	Expression too large to display	1316
parts	Expression too large to display	1487

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)
```

output `2/d*(-(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)/(csc(d*x+c)^2*(1-cos(d*x+c))^2-1))^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2)*(-A*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*a^2-A*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*a*b+A*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*a*b+A*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*b^2+csc(d*x+c)^3*A*a*b*(1-cos(d*x+c))^3-csc(d*x+c)^3*A*b^2*(1-cos(d*x+c))^3+B*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*a^2+B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b-B*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((c...`

3.622.5 Fracas [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x, algorithm="fracas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c))/(b^2*cos(d*x + c)^2 + 2*a*b*cos(d*x + c) + a^2), x)`

3.622.6 Sympy [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Integral((A + B*cos(c + d*x))*sqrt(sec(c + d*x))/(a + b*cos(c + d*x))**(3/2), x)`

3.622.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x, algo rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^(3/2), x)`

3.622.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x, algo rithm="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^(3/2), x)`

3.622.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^(3/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^(3/2), x)`

$$3.623 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$$

3.623.1 Optimal result	5710
3.623.2 Mathematica [B] (verified)	5711
3.623.3 Rubi [A] (verified)	5712
3.623.4 Maple [B] (warning: unable to verify)	5717
3.623.5 Fricas [F]	5718
3.623.6 Sympy [F]	5719
3.623.7 Maxima [F]	5719
3.623.8 Giac [F]	5719
3.623.9 Mupad [F(-1)]	5720

3.623.1 Optimal result

Integrand size = 35, antiderivative size = 476

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx =$$

$$\frac{2(Ab - aB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \middle| -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ab\sqrt{a + bd} \sqrt{\sec(c + dx)}} -$$

$$+ \frac{2(Ab - aB) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ab\sqrt{a + bd} \sqrt{\sec(c + dx)}} -$$

$$\frac{2\sqrt{a + b} B \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{b^2 d \sqrt{\sec(c + dx)}} +$$

$$\frac{2a(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{b(a^2 - b^2) d \sqrt{a + b \cos(c + dx)}}$$

output $2*a*(A*b-B*a)*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/b/(a^2-b^2)/d/(a+b*\cos(d*x+c))^{(1/2)}-2*(A*b-B*a)*\csc(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c))/(a+b))^{(1/2)}*(a*(1+\sec(d*x+c))/(a-b))^{(1/2)}/a/b/d/(a+b)^{(1/2)}/\sec(d*x+c)^{(1/2)}+2*(A*b-B*a)*\csc(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{(1/2)})*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c))/(a+b))^{(1/2)}*(a*(1+\sec(d*x+c))/(a-b))^{(1/2)}/a/b/d/(a+b)^{(1/2)}/\sec(d*x+c)^{(1/2)}-2*B*\csc(d*x+c)*\text{EllipticPi}((a+b*\cos(d*x+c))^{(1/2)}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},(a+b)/b,((-a-b)/(a-b))^{(1/2)})*(a+b)^{(1/2)}*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c))/(a+b))^{(1/2)}*(a*(1+\sec(d*x+c))/(a-b))^{(1/2)}/b^2/d/\sec(d*x+c)^{(1/2)}$

3.623.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1050 vs. $2(476) = 952$.

Time = 12.80 (sec) , antiderivative size = 1050, normalized size of antiderivative = 2.21

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sqrt[Sec[c + d*x]]),x]`

output $(\text{Sqrt}[a + b \cos[c + dx]] \cdot \text{Sqrt}[\text{Sec}[c + dx]] \cdot ((2(Ab - aB) \sin[c + dx]) / (b(-a^2 + b^2)) - (2(aAb \sin[c + dx] - a^2B \sin[c + dx])) / (b(-a^2 + b^2)(a + b \cos[c + dx]))) / d + (2 \text{Sqrt}[(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2) / (1 + \tan[(c + dx)/2]^2)] \cdot (aAb \tan[(c + dx)/2] + Ab^2 \tan[(c + dx)/2] - a^2B \tan[(c + dx)/2] - abB \tan[(c + dx)/2] - 2Ab^2 \tan[(c + dx)/2]^3 + 2aAbB \tan[(c + dx)/2]^3 - aAb \tan[(c + dx)/2]^5 + Ab^2 \tan[(c + dx)/2]^5 + a^2B \tan[(c + dx)/2]^5 - abB \tan[(c + dx)/2]^5 + 2a^2B \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2) / (a + b)] - 2b^2B \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2) / (a + b)] + 2a^2B \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \tan[(c + dx)/2]^2 \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2) / (a + b)] - 2b^2B \text{EllipticPi}[-1, \text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \tan[(c + dx)/2]^2 \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2) / (a + b)] - (a + b) \cdot (-Ab + aB) \cdot \text{EllipticE}[\text{ArcSin}[\tan[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \tan[(c + dx)/2]^2] \cdot (1 + \tan[(c + dx)/2]^2) \cdot \text{Sqrt}[(a + b + a \tan[(c + dx)/2]^2 - b \tan[(c + dx)/2]^2) / (a + b)] - b(a + b)(A - B) \cdot \text{EllipticF}[\text{ArcSin}[\dots$

3.623.3 Rubi [A] (verified)

Time = 1.74 (sec) , antiderivative size = 441, normalized size of antiderivative = 0.93, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3471, 3042, 3273, 3042, 3274, 3042, 3288, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{\csc\left(c + dx + \frac{\pi}{2}\right)}(a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{3/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

3.623. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})} (A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \downarrow \text{3471} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab-aB) \int \frac{\sqrt{\cos(c+dx)}}{(a+b\cos(c+dx))^{3/2}} dx}{b} + \frac{B \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx}{b} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab-aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \\
 & \downarrow \text{3273} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab-aB) \left(\frac{2a\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{\int \frac{\sqrt{a+b\cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx}{a^2-b^2} \right)}{b} + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab-aB) \left(\frac{2a\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{\int \frac{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}{\sin^{\frac{3}{2}}(c+dx+\frac{\pi}{2})} dx}{a^2-b^2} \right)}{b} + \frac{B \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} \right) \\
 & \downarrow \text{3274}
 \end{aligned}$$

3.623. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{3/2}\sqrt{\sec(c+dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{a \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a^2-b^2} \right)}{b} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a^2-b^2} \right)}{b} \right)$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a^2-b^2} \right)}{b} \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} - \frac{a \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{a^2-b^2} \right)}{b} \right)$$

↓ 3473

3.623. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

$$\frac{\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}}{\left((Ab - aB) \left(\frac{2a \sin(c+dx)}{d(a^2 - b^2)\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} - \frac{2(a-b)\sqrt{a+b} \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)-1)}{a-b}}}{ad} \right) \right)}$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sqrt[Sec[c + d*x]]),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((-2*Sqrt[a + b]*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b^2*d) + ((A*b - a*B)*(-((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a^2 - b^2) + (2*a*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))/b)`

3.623.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3273 `Int[Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2), x_Symbol] := Simp[-2*a*d*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] - Simp[d^2/(a^2 - b^2) Int[Sqrt[a + b*Sin[e + f*x]]/(d*Sin[e + f*x])^(3/2), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 3274 `Int[Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]/((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[(c - d)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(b*c - a*d)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x])], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3471 `Int[(((A_) + (B_)*sin[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]])/((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(3/2), x_Symbol] := Simp[B/b Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[(A*b - a*B)/b Int[Sqrt[c + d*Sin[e + f*x]]/(a + b*Sin[e + f*x])^(3/2), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

3.623.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1601 vs. 2(436) = 872.

Time = 17.63 (sec) , antiderivative size = 1602, normalized size of antiderivative = 3.37

method	result	size
default	Expression too large to display	1602
parts	Expression too large to display	1823

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2)/sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

output

```

2/d*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-cs
c(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1/2
)*(-A*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2
*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*
b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*a*b-A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2
+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^
2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*
b^2+A*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2
*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*
b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*a*b+A*EllipticE(cot(d*x+c)-csc(d*x+c)
,(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+
c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*
b^2+csc(d*x+c)^3*A*a*b*(1-cos(d*x+c))^3-csc(d*x+c)^3*A*b^2*(1-cos(d*x+c))^
3+B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c)
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)
)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b+B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1
)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+
a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*b^
2-B*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(-csc(d*x+c)^2*(
1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2...

```

3.623.5 Fracas [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)}} dx$$

input

```

integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algo
rithm="fracas")

```

output

```

integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/((b^2*cos(d*x + c))^
2 + 2*a*b*cos(d*x + c) + a^2)*sqrt(sec(d*x + c))), x)

```

3.623.6 Sympy [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}} \sqrt{\sec(c + dx)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2)/sec(d*x+c)**(1/2),x)`

output `Integral((A + B*cos(c + d*x))/((a + b*cos(c + d*x))**(3/2)*sqrt(sec(c + d*x))), x)`

3.623.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*x + c))), x)`

3.623.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*x + c))), x)`

3.623.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(3/2)), x)`

$$3.624 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.624.1 Optimal result	5721
3.624.2 Mathematica [B] (warning: unable to verify)	5722
3.624.3 Rubi [A] (verified)	5723
3.624.4 Maple [B] (warning: unable to verify)	5729
3.624.5 Fracas [F]	5730
3.624.6 Sympy [F(-1)]	5731
3.624.7 Maxima [F]	5731
3.624.8 Giac [F]	5731
3.624.9 Mupad [F(-1)]	5732

3.624.1 Optimal result

Integrand size = 35, antiderivative size = 560

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{(2aAb - 3a^2B + b^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b}}\right)\right) - (2Ab - (3a + b)B) \sqrt{\cos(c + dx)} \csc(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a+b \cos(c+dx)}{\cos(c+dx)}}}{b^2 \sqrt{a + b} \sqrt{\sec(c + dx)}} - \frac{\sqrt{a + b} (2Ab - 3aB) \sqrt{\cos(c + dx)} \csc(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{b^3 d \sqrt{\sec(c + dx)}} + \frac{2a(Ab - aB) \sin(c + dx)}{b(a^2 - b^2) d \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} - \frac{(2aAb - 3a^2B + b^2B) \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{b^2 (a^2 - b^2) d}$$

3.624. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$

output $2*a*(A*b-B*a)*\sin(d*x+c)/b/(a^2-b^2)/d/(a+b*\cos(d*x+c))^{1/2}/\sec(d*x+c)^{(1/2)}-(2*A*a*b-3*B*a^2+B*b^2)*\sin(d*x+c)*(a+b*\cos(d*x+c))^{1/2}*\sec(d*x+c)^{(1/2)}/b^2/(a^2-b^2)/d+(2*A*a*b-3*B*a^2+B*b^2)*\csc(d*x+c)*\text{EllipticE}((a+b*\cos(d*x+c))^{1/2}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{1/2})*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b)^{(1/2)}/a/b^2/d/(a+b)^{(1/2)}/\sec(d*x+c)^{(1/2)}-(2*A*b-(3*a+b)*B)*\csc(d*x+c)*\text{EllipticF}((a+b*\cos(d*x+c))^{1/2}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},((-a-b)/(a-b))^{1/2})*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b)^{(1/2)}/b^2/d/(a+b)^{(1/2)}/\sec(d*x+c)^{(1/2)}-(2*A*b-3*B*a)*\csc(d*x+c)*\text{EllipticPi}((a+b*\cos(d*x+c))^{1/2}/(a+b)^{(1/2)}/\cos(d*x+c)^{(1/2)},(a+b)/b,((-a-b)/(a-b))^{1/2})*\cos(d*x+c)^{(1/2)}*(a*(1-\sec(d*x+c)))/(a+b)^{(1/2)}*(a*(1+\sec(d*x+c)))/(a-b)^{(1/2)}/b^3/d/\sec(d*x+c)^{(1/2)}$

3.624.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 1551 vs. $2(560) = 1120$.

Time = 15.81 (sec) , antiderivative size = 1551, normalized size of antiderivative = 2.77

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^(3/2)),x]`

output $(\text{Sqrt}[a + b \cos[c + dx]] \text{Sqrt}[\text{Sec}[c + dx]] * ((-2*a*(-A*b) + a*B) \text{Sin}[c + dx]) / (b^2*(a^2 - b^2)) + (2*(a^2*A*b \text{Sin}[c + dx] - a^3*B \text{Sin}[c + dx])) / (b^2*(-a^2 + b^2)*(a + b \cos[c + dx])) / d - (\text{Sqrt}[(1 - \text{Tan}[(c + dx)/2]^2)^{-1}] \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2) / (1 + \text{Tan}[(c + dx)/2]^2)] * (2*a^2*A*b \text{Tan}[(c + dx)/2] + 2*a*A*b^2 \text{Tan}[(c + dx)/2] - 3*a^3*B \text{Tan}[(c + dx)/2] - 3*a^2*b*B \text{Tan}[(c + dx)/2] + a*b^2*B \text{Tan}[(c + dx)/2] + b^3*B \text{Tan}[(c + dx)/2] - 4*a*A*b^2 \text{Tan}[(c + dx)/2]^3 + 6*a^2*b*B \text{Tan}[(c + dx)/2]^3 - 2*b^3*B \text{Tan}[(c + dx)/2]^3 - 2*a^2*A*b \text{Tan}[(c + dx)/2]^5 + 2*a*A*b^2 \text{Tan}[(c + dx)/2]^5 + 3*a^3*B \text{Tan}[(c + dx)/2]^5 - 3*a^2*b*B \text{Tan}[(c + dx)/2]^5 - a*b^2*B \text{Tan}[(c + dx)/2]^5 + b^3*B \text{Tan}[(c + dx)/2]^5 - 4*a^2*A*b \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] + 4*A*b^3 \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] + 6*a^3*B \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] - 6*a*b^2*B \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] - 4*a^2*A*b \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + ...$

3.624.3 Rubi [A] (verified)

Time = 2.60 (sec) , antiderivative size = 548, normalized size of antiderivative = 0.98, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{3/2} (a + b \sin(c + dx + \frac{\pi}{2}))^{3/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

3.624. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \downarrow \text{3468} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{bd(a^2-b^2) \sqrt{a+b \cos(c+dx)}} - \frac{2 \int -\frac{((-3Ba^2+2Aba+b^2B) \cos^2(c+dx))-b(Ab-aB)}{2\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx}{b(a^2-b^2)} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{-((-3Ba^2+2Aba+b^2B) \cos^2(c+dx))-b(Ab-aB) \cos(c+dx)+a(Ab-aB)}{\sqrt{\cos(c+dx)}\sqrt{a+b \cos(c+dx)}} dx}{b(a^2-b^2)} + \frac{2a(Ab-aB) \sin(c+dx)}{bd(a^2-b^2) \sqrt{\cos(c+dx)}} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(3Ba^2-2Aba-b^2B) \sin(c+dx+\frac{\pi}{2})^2 - b(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + a(Ab-aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b(a^2-b^2)} + \frac{2a(Ab-aB) \sin(c+dx)}{bd(a^2-b^2) \sqrt{\cos(c+dx)}} \right) \\
 & \downarrow \text{3540} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a^2-b^2)(2Ab-3aB) \cos^2(c+dx)+2ab(Ab-aB) \cos(c+dx)+a(-3Ba^2+2Aba+b^2B)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b \cos(c+dx)}} dx}{b(a^2-b^2)} - \frac{(-3a^2B+2aAb+b^2B) \sin(c+dx)}{bd \sqrt{\cos(c+dx)}} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(a^2-b^2)(2Ab-3aB) \sin(c+dx+\frac{\pi}{2})^2 + 2ab(Ab-aB) \sin(c+dx+\frac{\pi}{2}) + a(-3Ba^2+2Aba+b^2B)}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b(a^2-b^2)} - \frac{(-3a^2B+2aAb+b^2B) \cos(c+dx)}{bd \sqrt{\cos(c+dx)}} \right) \\
 & \downarrow \text{3532}
 \end{aligned}$$

3.624. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(-3Ba^2+2Aba+b^2B)+2ab(Ab-aB)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx + (a^2-b^2)(2Ab-3aB) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b\cos(c+dx)}} dx}{2b} - (-3a^2B+2aAb+b^2B) \right) \frac{1}{b(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a^2-b^2)(2Ab-3aB) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \int \frac{a(-3Ba^2+2Aba+b^2B)+2ab(Ab-aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} - (-3a^2B+2aAb+b^2B) \right) \frac{1}{b(a^2-b^2)}$$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(-3Ba^2+2Aba+b^2B)+2ab(Ab-aB)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(a^2-b^2)(2Ab-3aB)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{2b}}{2b} - (-3a^2B+2aAb+b^2B) \right) \frac{1}{b(a^2-b^2)}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(-3a^2B+2aAb+b^2B) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - a(a-b)(2Ab-B(3a+b)) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{2b} - (-3a^2B+2aAb+b^2B) \right) \frac{1}{b(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(-3a^2B+2aAb+b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - a(a-b)(2Ab-B(3a+b)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}} dx}{2b} - (-3a^2B+2aAb+b^2B) \right) \frac{1}{b(a^2-b^2)}$$

↓ 3295

3.624. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{3/2}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(-3a^2B+2aAb+b^2B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{a+b}(a^2-b^2)(2Ab-3aB) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{ad}}{\dots} \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a-b)\sqrt{a+b}(-3a^2B+2aAb+b^2B) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(\sec(c+dx)+1)}{a-b}} E\left(\arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right)\right) - \frac{a}{a-b}}{ad}}{\dots} \right)$$

```
input Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^(3/2)),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*(A*b - a*B)*Sqrt[Cos[c + d*x]]
*Sin[c + d*x])/(b*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + (((2*(a - b)*S
qrt[a + b]*(2*a*A*b - 3*a^2*B + b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[
a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]
)*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)
)/(a*d) - (2*(a - b)*Sqrt[a + b]*(2*A*b - (3*a + b)*B)*Cot[c + d*x]*Ellipt
icF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((
a + b)/(a - b))] *Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c +
d*x]))/(a - b)))/d - (2*Sqrt[a + b]*(a^2 - b^2)*(2*A*b - 3*a*B)*Cot[c + d
*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqr
t[Cos[c + d*x]])], -((a + b)/(a - b))] *Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)
]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(b*d))/(2*b) - ((2*a*A*b - 3*a^2*B
+ b^2*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x])/(b*d*Sqrt[Cos[c + d*x]])
/(b*(a^2 - b^2)))
```

3.624.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d))*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3532 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[C/b^2 Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[1/b^2 Int[(A*b^2 - a^2*C + b*(b*B - 2*a*C)*Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]))]*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.624.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 3269 vs. $2(516) = 1032$.

Time = 18.01 (sec) , antiderivative size = 3270, normalized size of antiderivative = 5.84

method	result	size
parts	Expression too large to display	3270
default	Expression too large to display	4195

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(3/2)/sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```


output

```

2*A/d*(-(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d
*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticF(cot(d
*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*a*b-(-csc(d*x+c)^2*(1-cos(d*x+c))^2
+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^
2+a+b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*
b^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c
))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c
)-csc(d*x+c),(-a-b)/(a+b))^(1/2)*a^2+(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^
(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+
b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*a*b-
2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))
^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(a+b))^(1/2)*EllipticPi(cot(d*x+c
)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2)*a^2+2*(-csc(d*x+c)^2*(1-cos(d*x+c))^2
+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^
2+a+b)/(a+b))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/
2))*b^2+csc(d*x+c)^3*a^2*(1-cos(d*x+c))^3-csc(d*x+c)^3*a*b*(1-cos(d*x+c))^
3-a^2*(csc(d*x+c)-cot(d*x+c))+a*b*(csc(d*x+c)-cot(d*x+c))*((csc(d*x+c)^2*
a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a+b)/(csc(d*x+c)^2*(1-c
os(d*x+c))^2+1)^(1/2)*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^2/(csc(d*x+c)^4*a
*(1-cos(d*x+c))^4-csc(d*x+c)^4*b*(1-cos(d*x+c))^4+2*csc(d*x+c)^2*b*(1-c...

```

3.624.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{\frac{3}{2}} \sec^{\frac{3}{2}}(dx + c)} dx$$

input

```

integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algo
rithm="fricas")

```

output

```

integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/((b^2*cos(d*x + c)^
2 + 2*a*b*cos(d*x + c) + a^2)*sec(d*x + c)^(3/2)), x)

```

3.624.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2)/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.624.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2)), x)`

3.624.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2)), x)`

3.624.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(3/2)), x)`

3.625
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

3.625.1 Optimal result	5733
3.625.2 Mathematica [B] (warning: unable to verify)	5734
3.625.3 Rubi [A] (verified)	5735
3.625.4 Maple [B] (warning: unable to verify)	5741
3.625.5 Fricas [F]	5741
3.625.6 Sympy [F(-1)]	5742
3.625.7 Maxima [F]	5742
3.625.8 Giac [F]	5742
3.625.9 Mupad [F(-1)]	5743

3.625.1 Optimal result

Integrand size = 35, antiderivative size = 607

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx =$$

$$\frac{2(8a^4Ab - 28a^2Ab^3 + 16Ab^5 - 3a^5B + 15a^3b^2B - 8ab^4B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) + 2(16Ab^4 - a^4(A - 3B) + 4ab^3(3A - 2B) - 9a^3b(A - B) - 2a^2b^2(8A + 3B)) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) + 2b(Ab - aB) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx) + 2b(10a^2Ab - 6Ab^3 - 7a^3B + 3ab^2B) \sec^{\frac{3}{2}}(c + dx) \sin(c + dx) + 2(a^4A - 13a^2Ab^2 + 8Ab^4 + 8a^3bB - 4ab^3B) \sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx) \sin(c + dx)}{3a^5(a - b)(a + b)^{3/2}d\sqrt{\sec(c + dx)} + 3a^4\sqrt{a + b}(a^2 - b^2)d\sqrt{\sec(c + dx)} + 3a(a^2 - b^2)d(a + b \cos(c + dx))^{3/2} + 3a^2(a^2 - b^2)^2d\sqrt{a + b \cos(c + dx)} + 3a^3(a^2 - b^2)^2d}$$

3.625.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

output $\frac{2}{3}b(Ab - Ba) \sec(dx+c)^{3/2} \sin(dx+c) / a(a^2 - b^2) / d(a+b \cos(dx+c))^{3/2} + \frac{2}{3}b(10Aa^2b - 6Ab^3 - 7B^2a^3 + 3B^2ab^2) \sec(dx+c)^{3/2} \sin(dx+c) / a^2(a^2 - b^2)^2 / d(a+b \cos(dx+c))^{1/2} + \frac{2}{3}(Aa^4 - 13Aa^2b^2 + 8A^2b^4 + 8B^2a^3b - 4B^2ab^3) \sec(dx+c)^{3/2} \sin(dx+c) (a+b \cos(dx+c))^{1/2} / a^3(a^2 - b^2)^2 / d - \frac{2}{3}(8Aa^4b - 28Aa^2b^3 + 16A^2b^5 - 3B^2a^5 + 15B^2ab^3 - 8B^2ab^4) \csc(dx+c) \operatorname{EllipticE}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2}) \cos(dx+c)^{1/2} (a(1 - \sec(dx+c)) / (a+b))^{1/2} (a(1 + \sec(dx+c)) / (a-b))^{1/2} / a^5(a-b) / (a+b)^{3/2} / d / \sec(dx+c)^{1/2} - \frac{2}{3}(16A^2b^4 - a^4(A - 3B) + 4a^2b^3(3A - 2B) - 9a^3b(A - B) - 2a^2b^2(8A + 3B)) \csc(dx+c) \operatorname{EllipticF}((a+b \cos(dx+c))^{1/2} / (a+b)^{1/2} / \cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2}) \cos(dx+c)^{1/2} (a(1 - \sec(dx+c)) / (a+b))^{1/2} (a(1 + \sec(dx+c)) / (a-b))^{1/2} / a^4(a^2 - b^2) / d / (a+b)^{1/2} / \sec(dx+c)^{1/2}$

3.625.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 4316 vs. $2(607) = 1214$.

Time = 24.30 (sec) , antiderivative size = 4316, normalized size of antiderivative = 7.11

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \text{Result too large to show}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x])^(5/2), x]`

output $(\sqrt{a + b\cos[c + dx]}\sqrt{\sec[c + dx]}*((2*(-8*a^4*A*b + 28*a^2*A*b^3 - 16*A*b^5 + 3*a^5*B - 15*a^3*b^2*B + 8*a*b^4*B)*\sin[c + dx])/(3*a^4*(a^2 - b^2)^2) + (2*(-(A*b^3*\sin[c + dx]) + a*b^2*B*\sin[c + dx]))/(3*a^2*(a^2 - b^2)*(a + b*\cos[c + dx])^2) + (2*(-11*a^2*A*b^3*\sin[c + dx] + 7*A*b^5*\sin[c + dx] + 8*a^3*b^2*B*\sin[c + dx] - 4*a*b^4*B*\sin[c + dx]))/(3*a^3*(a^2 - b^2)^2*(a + b*\cos[c + dx])) + (2*A*\tan[c + dx])/(3*a^3)))/d + (2*((8*A*A*b)/(3*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}\sqrt{\sec[c + dx]}) - (28*A*b^3)/(3*a*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}\sqrt{\sec[c + dx]}) + (16*A*b^5)/(3*a^3*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}\sqrt{\sec[c + dx]}) - (a^2*B)/((a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}\sqrt{\sec[c + dx]}) + (5*b^2*B)/((a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}\sqrt{\sec[c + dx]}) - (8*b^4*B)/(3*a^2*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}\sqrt{\sec[c + dx]}) + (a^2*A*\sqrt{\sec[c + dx]})/(3*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) + (5*A*b^2*\sqrt{\sec[c + dx]})/((a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) - (32*A*b^4*\sqrt{\sec[c + dx]})/(3*a^2*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) + (16*A*b^6*\sqrt{\sec[c + dx]})/(3*a^4*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) - (3*a*b*B*\sqrt{\sec[c + dx]})/((a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) + (17*b^3*B*\sqrt{\sec[c + dx]})/(3*a*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) - (8*b^5*B*\sqrt{\sec[c + dx]})/(3*a^3*(a^2 - b^2)^2*\sqrt{a + b*\cos[c + dx]}) + (8*A*b^2*\cos[2*(c + dx)]*\sqrt{\sec[c + dx]})/(3*...$

3.625.3 Rubi [A] (verified)

Time = 2.90 (sec) , antiderivative size = 602, normalized size of antiderivative = 0.99, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3534, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(c + dx + \frac{\pi}{2})^{5/2} (A + B \sin(c + dx + \frac{\pi}{2}))}{(a + b \sin(c + dx + \frac{\pi}{2}))^{5/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)}\sqrt{\sec(c + dx)} \int \frac{A + B \cos(c + dx)}{\cos^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx$$

3.625. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{5/2}(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
 & \downarrow 3479 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{4b(Ab-aB)\cos^2(c+dx)-3a(Ab-aB)\cos(c+dx)+3(Aa^2+bBa-2Ab^2)}{2\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-aB)}{3ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)} \right) \\
 & \downarrow 27 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{4b(Ab-aB)\cos^2(c+dx)-3a(Ab-aB)\cos(c+dx)+3(Aa^2+bBa-2Ab^2)}{\cos^{\frac{5}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-aB)}{3ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)} \right) \\
 & \downarrow 3042 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{4b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2-3a(Ab-aB)\sin(c+dx+\frac{\pi}{2})+3(Aa^2+bBa-2Ab^2)}{\sin(c+dx+\frac{\pi}{2})^{5/2}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-aB)}{3ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)} \right) \\
 & \downarrow 3534 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{2b(-7Ba^3+10Aba^2+3b^2Ba-6Ab^3)\cos^2(c+dx)-a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\cos(c+dx)+3(Aa^4+8bBa^3-13Ab^2a^2)}{2\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)}{3a(a^2-b^2)} \right) \\
 & \downarrow 27 \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{2b(-7Ba^3+10Aba^2+3b^2Ba-6Ab^3)\cos^2(c+dx)-a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\cos(c+dx)+3(Aa^4+8bBa^3-13Ab^2a^2)}{\cos^{\frac{5}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(Ab-aB)}{3a(a^2-b^2)} \right) \\
 & \downarrow 3042
 \end{aligned}$$

3.625. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{2b(-7Ba^3+10Aba^2+3b^2Ba-6Ab^3)\sin(c+dx+\frac{\pi}{2})^2 - a(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\sin(c+dx+\frac{\pi}{2}) + 3(Aa^4+8bBa^3)}{\sin(c+dx+\frac{\pi}{2})^{5/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{3(-3Ba^5+8Aba^4+15b^2Ba^3-28Ab^3a^2-8b^4Ba - (Aa^4-6bBa^3+7Ab^2a^2+2b^3Ba-4Ab^4)\cos(c+dx)a+16Ab^5)}{2 \cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a} \right) \frac{1}{a(a^2-b^2)} \frac{1}{3a(a^2-b^2)}$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-3Ba^5+8Aba^4+15b^2Ba^3-28Ab^3a^2-8b^4Ba - (Aa^4-6bBa^3+7Ab^2a^2+2b^3Ba-4Ab^4)\cos(c+dx)a+16Ab^5}{\cos^{\frac{3}{2}}(c+dx)} dx}{a(a^2-b^2)} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad \cos^{\frac{3}{2}}(c+dx)} - \frac{\int \frac{-3Ba^5+8Aba^4+15b^2Ba^3-28Ab^3a^2-8b^4Ba - (Aa^4-6bBa^3+7Ab^2a^2+2b^3Ba-4Ab^4)\cos(c+dx)a+16Ab^5}{\sin(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3477

3.625. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(a-b)(-a^4(A-3B)-9a^3b(A-B)-2a^2b^2(8A+3b^2))\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(a-b)(-a^4(A-3B)-9a^3b(A-B)-2a^2b^2(8A+3b^2))\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a^4A+8a^3bB-13a^2Ab^2-4ab^3B+8Ab^4)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad\cos^{\frac{3}{2}}(c+dx)} - \frac{(-3a^5B+8a^4Ab+15a^3b^2B-28a^2Ab^3-8ab^4B+3b^5)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad\cos^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2b(Ab-aB)\sin(c+dx)}{3ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} + \frac{2b(-7a^3B+10a^2Ab+3ab^2B-6Ab^3)\sin(c+dx)\sqrt{a+b\cos(c+dx)}}{ad(a^2-b^2)\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} \right)$$

```
input Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(5/2))/(a + b*Cos[c + d*x])^(5/2), x]
```

3.625. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{5}{2}}(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*b*(A*b - a*B)*Sin[c + d*x])/(3*a
*(a^2 - b^2)*d*Cos[c + d*x]^(3/2)*(a + b*Cos[c + d*x])^(3/2)) + ((2*b*(10*
a^2*A*b - 6*A*b^3 - 7*a^3*B + 3*a*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Co
s[c + d*x]^(3/2)*Sqrt[a + b*Cos[c + d*x]]) + (-(((2*(a - b)*Sqrt[a + b]*(8
*a^4*A*b - 28*a^2*A*b^3 + 16*A*b^5 - 3*a^5*B + 15*a^3*b^2*B - 8*a*b^4*B)*C
ot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Co
s[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sq
rt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(16*A
*b^4 - a^4*(A - 3*B) + 4*a*b^3*(3*A - 2*B) - 9*a^3*b*(A - B) - 2*a^2*b^2*(
8*A + 3*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a
+ b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x])
)/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)]/(a*d))/a + (2*(a^4*A - 1
3*a^2*A*b^2 + 8*A*b^4 + 8*a^3*b*B - 4*a*b^3*B)*Sqrt[a + b*Cos[c + d*x]]*Si
n[c + d*x])/(a*d*Cos[c + d*x]^(3/2)))/(a*(a^2 - b^2))/(3*a*(a^2 - b^2))
```

3.625.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_.) + (b_)*sin[(e_.) + (f
_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 3440 Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*
(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

3.625.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$$

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(b \sin(e + fx))^{3/2}} dx$$

$$:= \text{Simp}[-2A(c - d)(\tan[e + fx]/(fbc^2)) \text{Rt}[(c + d)/b, 2] \sqrt{c((1 + \text{Csc}[e + fx])/(c - d))} \sqrt{c((1 - \text{Csc}[e + fx])/(c + d))} \text{EllipticE}[\text{ArcSin}[\sqrt{c + d \sin[e + fx]}/\sqrt{b \sin[e + fx]}/\text{Rt}[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\& \text{PosQ}[(c + d)/b]$$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$:= \text{Simp}[(A - B)/(a - b) \int[1/(\sqrt{a + b \sin[e + fx]}\sqrt{c + d \sin[e + fx]}), x], x] - \text{Simp}[(A*b - a*B)/(a - b) \int[(1 + \sin[e + fx])/((a + b \sin[e + fx])^{3/2}\sqrt{c + d \sin[e + fx]}), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[A, B]$$`

rule 3479 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +

$$\int ((a + b \sin(e + fx))^m ((A + B \sin(e + fx))^n) dx$$

$$:= \text{Simp}[(-(A*b^2 - a*b*B))*\text{Cos}[e + fx]*(a + b*\text{Sin}[e + fx])^{m+1}*((c + d*\text{Sin}[e + fx])^{1+n}/(f*(m+1)*(b*c - a*d)*(a^2 - b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 - b^2)) \int[(a + b*\text{Sin}[e + fx])^{m+1}*(c + d*\text{Sin}[e + fx])^n \text{Simp}[(a*A - b*B)*(b*c - a*d)*(m+1) + b*d*(A*b - a*B)*(m+n+2) + (A*b - a*B)*(a*d*(m+1) - b*c*(m+2))*\text{Sin}[e + fx] - b*d*(A*b - a*B)*(m+n+3)*\text{Sin}[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{RationalQ}[m] \&\& m < -1 \&\& ((\text{EqQ}[a, 0] \&\& \text{IntegerQ}[m] \&\& !\text{IntegerQ}[n]) || !(\text{IntegerQ}[2*n] \&\& \text{LtQ}[n, -1] \&\& ((\text{IntegerQ}[n] \&\& !\text{IntegerQ}[m]) || \text{EqQ}[a, 0])))$$`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.625.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 9527 vs. 2(561) = 1122.

Time = 25.71 (sec) , antiderivative size = 9528, normalized size of antiderivative = 15.70

method	result	size
parts	Expression too large to display	9528
default	Expression too large to display	9947

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+cos(d*x+c)*b)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.625.5 Fricas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="fricas")
```

3.625. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(5/2)/(b^3*cos(d*x + c)^3 + 3*a*b^2*cos(d*x + c)^2 + 3*a^2*b*cos(d*x + c) + a^3), x)`

3.625.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(5/2)/(a+b*cos(d*x+c))**(5/2), x)`

output `Timed out`

3.625.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^(5/2), x)`

3.625.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(5/2)/(a+b*cos(d*x+c))^(5/2), x, algorithm="giac")`

3.625. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{5}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(5/2)/(b*cos(d*x + c) + a)^(5/2), x)`

3.625.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{5}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{\frac{5}{2}}}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(5/2))/(a + b*cos(c + d*x))^(5/2), x)`

3.626
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

3.626.1 Optimal result	5744
3.626.2 Mathematica [A] (warning: unable to verify)	5745
3.626.3 Rubi [A] (verified)	5745
3.626.4 Maple [B] (warning: unable to verify)	5750
3.626.5 Fricas [F]	5750
3.626.6 Sympy [F(-1)]	5751
3.626.7 Maxima [F]	5751
3.626.8 Giac [F]	5751
3.626.9 Mupad [F(-1)]	5752

3.626.1 Optimal result

Integrand size = 35, antiderivative size = 496

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2(3a^4A - 15a^2Ab^2 + 8Ab^4 + 6a^3bB - 2ab^3B) \sqrt{\cos(c + dx)} \csc(c + dx)}{3a^4(a - b)(a + b)} + \frac{2(8Ab^3 - 3a^3(A - B) + 2ab^2(3A - B) - 3a^2b(3A + B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a-b}}{\sqrt{a+b}}\right), \frac{a-b}{a+b}\right)}{3a^3\sqrt{a+b}(a^2 - b^2) d\sqrt{\sec(c + dx)}} + \frac{2b(Ab - aB) \sqrt{\sec(c + dx)} \sin(c + dx)}{3a(a^2 - b^2) d(a + b \cos(c + dx))^{3/2}} + \frac{2b(8a^2Ab - 4Ab^3 - 5a^3B + ab^2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3a^2(a^2 - b^2)^2 d\sqrt{a + b \cos(c + dx)}}$$

output

```
2/3*b*(A*b-B*a)*sin(d*x+c)*sec(d*x+c)^(1/2)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))
^(3/2)+2/3*b*(8*A*a^2*b-4*A*b^3-5*B*a^3+B*a*b^2)*sin(d*x+c)*sec(d*x+c)^(1/2)/a^2/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)+2/3*(3*A*a^4-15*A*a^2*b^2+8*A*b^4+6*B*a^3*b-2*B*a*b^3)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^4/(a-b)/(a+b)^(3/2)/d/sec(d*x+c)^(1/2)+2/3*(8*A*b^3-3*a^3*(A-B)+2*a*b^2*(3*A-B)-3*a^2*b*(3*A+B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/(a^2-b^2)/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)
```

3.626.
$$\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$$

3.626.2 Mathematica [A] (warning: unable to verify)

Time = 18.73 (sec) , antiderivative size = 610, normalized size of antiderivative = 1.23

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2(3a^4A - 15a^2Ab^2 + 8Ab^4 + 6a^3bB - 2ab^3B)}{3a^3(a^2 - b^2)^2} \right)}{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)} \sec(c + dx) \left(-2(a + b)(3a^4A - 15a^2Ab^2 + 8Ab^4 + 6a^3bB - 2ab^3B) \sqrt{\frac{\cos(c + dx)}{1 + \cos(c + dx)}} \sqrt{\frac{1}{(a + b \cos(c + dx))}} \right)}$$

input `Integrate[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^(5/2),x]`

output `(Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*(3*a^4*A - 15*a^2*A*b^2 + 8*A*b^4 + 6*a^3*b*B - 2*a*b^3*B)*Sin[c + d*x])/(3*a^3*(a^2 - b^2)^2) - (2*(-(A*b^2*Sin[c + d*x]) + a*b*B*Sin[c + d*x]))/(3*a*(a^2 - b^2)*(a + b*Cos[c + d*x])^2) - (2*(-8*a^2*A*b^2*Sin[c + d*x] + 4*A*b^4*Sin[c + d*x] + 5*a^3*b*B*Sin[c + d*x] - a*b^3*B*Sin[c + d*x]))/(3*a^2*(a^2 - b^2)^2*(a + b*Cos[c + d*x])))/d + (2*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(3*a^4*A - 15*a^2*A*b^2 + 8*A*b^4 + 6*a^3*b*B - 2*a*b^3*B)*Sqrt[Cos[c + d*x]]/(1 + Cos[c + d*x]))*Sqrt[(a + b*Cos[c + d*x])/(a + b)*(1 + Cos[c + d*x])])*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*a*(a + b)*(8*A*b^3 + 3*a^2*b*(-3*A + B) + 3*a^3*(A + B) - 2*a*b^2*(3*A + B))*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/(a + b)*(1 + Cos[c + d*x])])*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - (3*a^4*A - 15*a^2*A*b^2 + 8*A*b^4 + 6*a^3*b*B - 2*a*b^3*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/(3*a^3*(a^2 - b^2)^2*d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2])`

3.626.3 Rubi [A] (verified)Time = 2.16 (sec) , antiderivative size = 498, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3534, 27, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.626. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)(A+B\cos(c+dx))}{(a+b\cos(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\csc(c+dx+\frac{\pi}{2})^{3/2}(A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 3440

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{5/2}} dx$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{A+B\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx$$

↓ 3479

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{3Aa^2+bBa-3(Ab-aB)\cos(c+dx)a-4Ab^2+2b(Ab-aB)\cos^2(c+dx)}{2\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-a)}{3ad(a^2-b^2)\sqrt{\cos(c+dx)}} \right)$$

↓ 27

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^2+bBa-3(Ab-aB)\cos(c+dx)a-4Ab^2+2b(Ab-aB)\cos^2(c+dx)}{\cos^{\frac{3}{2}}(c+dx)(a+b\cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-a)}{3ad(a^2-b^2)\sqrt{\cos(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^2+bBa-3(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-4Ab^2+2b(Ab-aB)\sin(c+dx+\frac{\pi}{2})^2}{\sin(c+dx+\frac{\pi}{2})^{3/2}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-a)}{3ad(a^2-b^2)\sqrt{\cos(c+dx)}} \right)$$

↓ 3534

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{3Aa^4+6bBa^3-15Ab^2a^2-2b^3Ba-(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\cos(c+dx)a+8Ab^4}{2\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a(a^2-b^2)} + \frac{2b(-5a^3B+8a^2Ab+ad^2)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}} \right)$$

↓ 27

3.626. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+b\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^4+6bBa^3-15Ab^2a^2-2b^3Ba-(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\cos(c+dx)a+8Ab^4}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a(a^2-b^2)} + \frac{2b(-5a^3B+8a^2Ab+a^2B)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^4+6bBa^3-15Ab^2a^2-2b^3Ba-(-3Ba^3+6Aba^2-b^2Ba-2Ab^3)\sin(c+dx+\frac{\pi}{2})a+8Ab^4}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a(a^2-b^2)} + \frac{2b(-5a^3B+8a^2Ab+a^2B)}{ad(a^2-b^2)\sqrt{\cos(c+dx)}} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a-b)(-3a^3(A-B)-3a^2b(3A+B)+2ab^2(3A-B)+8Ab^3) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + (3a^4A+6a^3bB-15a^2Ab^2-2ab^3B+8Ab^4)}{a(a^2-b^2)} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(a-b)(-3a^3(A-B)-3a^2b(3A+B)+2ab^2(3A-B)+8Ab^3) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + (3a^4A+6a^3bB-15a^2Ab^2-2ab^3B+8Ab^4)}{a(a^2-b^2)} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3a^4A+6a^3bB-15a^2Ab^2-2ab^3B+8Ab^4) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{\frac{3}{2}}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + \frac{2(a-b)\sqrt{a+b}(-3a^3(A-B)-3a^2b(3A+B)+2ab^2(3A-B)+8Ab^3)}{a(a^2-b^2)} \right) \frac{1}{3a(a^2-b^2)}$$

↓ 3473

3.626. $\int \frac{(A+B\cos(c+dx))\sec^{\frac{3}{2}}(c+dx)}{(a+b\cos(c+dx))^{\frac{5}{2}}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2b(Ab - aB) \sin(c+dx)}{3ad(a^2 - b^2) \sqrt{\cos(c+dx)}(a + b \cos(c+dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B - 4Ab^3) \sin(c+dx)}{ad(a^2 - b^2) \sqrt{\cos(c+dx)} \sqrt{a + b \cos(c+dx)}} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^(5/2),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*b*(A*b - a*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*(a + b*Cos[c + d*x])^(3/2)) + (((2*(a - b)*Sqrt[a + b]*(3*a^4*A - 15*a^2*A*b^2 + 8*A*b^4 + 6*a^3*b*B - 2*a*b^3*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) + (2*(a - b)*Sqrt[a + b]*(8*A*b^3 - 3*a^3*(A - B) + 2*a*b^2*(3*A - B) - 3*a^2*b*(3*A + B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)))/(a*d))/(a*(a^2 - b^2)) + (2*b*(8*a^2*A*b - 4*A*b^3 - 5*a^3*B + a*b^2*B)*Sin[c + d*x])/(a*(a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))/(3*a*(a^2 - b^2)))`

3.626.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_.) + (b_)*sin[(e_.) + (f_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]]/Rt[(a + b)/d, 2], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

3.626. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

```
rule 3534 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] :> Simp[(-(A*b^2 - a*b*B + a^2*C))*Cos[e + f*x
]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(n + 1)/(f*(m + 1)*(b*
c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int
[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(m + 1)*(b*c - a*
d)*(a*A - b*B + a*C) + d*(A*b^2 - a*b*B + a^2*C)*(m + n + 2) - (c*(A*b^2 -
a*b*B + a^2*C) + (m + 1)*(b*c - a*d)*(A*b - a*B + b*C))*Sin[e + f*x] - d*(A
*b^2 - a*b*B + a^2*C)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0] && LtQ[m, -1] && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ
[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) |
| EqQ[a, 0])))
```

3.626.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 6407 vs. $2(456) = 912$.

Time = 22.02 (sec) , antiderivative size = 6408, normalized size of antiderivative = 12.92

method	result	size
default	Expression too large to display	6408
parts	Expression too large to display	6566

```
input int((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.626.5 Fracas [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

```
input integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(5/2),x, algo
rithm="fracas")
```

3.626. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{5/2}} dx$

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)/(b^3*cos(d*x + c)^3 + 3*a*b^2*cos(d*x + c)^2 + 3*a^2*b*cos(d*x + c) + a^3), x)`

3.626.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(5/2), x)`

output `Timed out`

3.626.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(5/2), x)`

3.626.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{(B \cos(dx + c) + A) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(5/2), x, algorithm="giac")`

3.626. $\int \frac{(A+B \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{5}{2}}} dx$

output `integrate((B*cos(d*x + c) + A)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)^(5/2), x)`

3.626.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \left(\frac{1}{\cos(c + dx)}\right)^{3/2}}{(a + b \cos(c + dx))^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(3/2))/(a + b*cos(c + d*x))^(5/2), x)`

3.627
$$\int \frac{(A+B \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{5/2}} dx$$

3.627.1 Optimal result	5753
3.627.2 Mathematica [A] (warning: unable to verify)	5754
3.627.3 Rubi [A] (verified)	5754
3.627.4 Maple [B] (warning: unable to verify)	5759
3.627.5 Fracas [F]	5759
3.627.6 Sympy [F(-1)]	5759
3.627.7 Maxima [F]	5760
3.627.8 Giac [F]	5760
3.627.9 Mupad [F(-1)]	5760

3.627.1 Optimal result

Integrand size = 35, antiderivative size = 469

$$\int \frac{(A + B \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \frac{2(6a^2Ab - 2Ab^3 - 3a^3B - ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) + 2(2Ab^2 - 3a^2(A + B) + ab(3A + B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) - \frac{2b(Ab - aB) \sin(c + dx)}{3a(a^2 - b^2) d(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}}}{3a^2\sqrt{a + b}(a^2 - b^2) d\sqrt{\sec(c + dx)}} + \frac{2(6a^2Ab - 2Ab^3 - 3a^3B - ab^2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3a(a^2 - b^2)^2 d\sqrt{a + b \cos(c + dx)}}$$

output

```
2/3*b*(A*b-B*a)*sin(d*x+c)/a/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)
^(1/2)-2/3*(6*A*a^2*b-2*A*b^3-3*B*a^3-B*a*b^2)*sin(d*x+c)*sec(d*x+c)^(1/2)
/a/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)+2/3*(6*A*a^2*b-2*A*b^3-3*B*a^3-B*a
*b^2)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(
1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)
*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^3/(a-b)/(a+b)^(3/2)/d/sec(d*x+c)^(1/2)-2
/3*(2*A*b^2-3*a^2*(A+B)+a*b*(3*A+B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))
^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)
*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/(a^2-b^
2)/d/(a+b)^(1/2)/sec(d*x+c)^(1/2)
```


3.627.2 Mathematica [A] (warning: unable to verify)

Time = 17.15 (sec) , antiderivative size = 557, normalized size of antiderivative = 1.19

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(-\frac{2(-6a^2Ab + 2Ab^3 + 3a^3B + ab^2B) \sin(c + dx)}{3a^2(a^2 - b^2)^2} \right)}{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)} \sec(c + dx) (2(a + b) (-6a^2Ab + 2Ab^3 + 3a^3B + ab^2B)) \sqrt{\frac{\cos(c + dx)}{1 + \cos(c + dx)}} \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}}}$$

input `Integrate[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^(5/2), x]`

output `(Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((-2*(-6*a^2*A*b + 2*A*b^3 + 3*a^3*B + a*b^2*B)*Sin[c + d*x])/(3*a^2*(a^2 - b^2)^2) + (2*(-(A*b*SIN[c + d*x]) + a*B*SIN[c + d*x]))/(3*(a^2 - b^2)*(a + b*Cos[c + d*x])^2) + (2*(-5*a^2*A*b*SIN[c + d*x] + A*b^3*SIN[c + d*x] + 2*a^3*B*SIN[c + d*x] + 2*a*b^2*B*SIN[c + d*x]))/(3*a*(a^2 - b^2)^2*(a + b*Cos[c + d*x])))/d + (2*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(2*(a + b)*(-6*a^2*A*b + 2*A*b^3 + 3*a^3*B + a*b^2*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] *Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*a*(a + b)*(-2*A*b^2 + 3*a^2*(A - B) + a*b*(3*A - B))*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] *Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + (-6*a^2*A*b + 2*A*b^3 + 3*a^3*B + a*b^2*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2]))/(3*(a^3 - a*b^2)^2*d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2])`

3.627.3 Rubi [A] (verified)Time = 1.96 (sec) , antiderivative size = 465, normalized size of antiderivative = 0.99, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c + dx)}(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx$$

3.627. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{\csc(c+dx+\frac{\pi}{2})(A+B\sin(c+dx+\frac{\pi}{2}))}}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}(A+B\cos(c+dx))}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{5/2}} dx \\
& \quad \downarrow \text{3440} \\
& \int \frac{\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}(A+B\sin(c+dx+\frac{\pi}{2}))}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2 \int \frac{3Aa^2-bBa-3(Ab-aB)\cos(c+dx)a-2Ab^2}{2\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{3479} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^2-bBa-3(Ab-aB)\cos(c+dx)a-2Ab^2}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^2-bBa-3(Ab-aB)\sin(c+dx+\frac{\pi}{2})a-2Ab^2}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3a(a^2-b^2)} + \frac{2b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{-3Ba^3+6Aba^2-b^2Ba+(3Aa^2-4bBa+Ab^2)\cos(c+dx)a-2Ab^3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a(a^2-b^2)} - \frac{2(-3a^3B+6a^2Ab-ab^2B-2Ab^3)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\
& \quad \downarrow \text{3472} \\
& \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{-3Ba^3+6Aba^2-b^2Ba+(3Aa^2-4bBa+Ab^2)\cos(c+dx)a-2Ab^3}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{3a(a^2-b^2)} - \frac{2(-3a^3B+6a^2Ab-ab^2B-2Ab^3)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.627. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b\cos(c+dx))^{5/2}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{-3Ba^3+6Aba^2-b^2Ba+(3Aa^2-4bBa+Ab^2)\sin(c+dx+\frac{\pi}{2})a-2Ab^3}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} - \frac{2(-3a^3B+6a^2Ab-ab^2B-2Ab^3)\sin(c+dx+\frac{\pi}{2})}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{3a(a^2-b^2)}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(-3a^3B+6a^2Ab-ab^2B-2Ab^3) \int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b)(-3a^2(A+B)+ab(3A+B)+2Ab^2)}{a^2-b^2}}{3a(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(-3a^3B+6a^2Ab-ab^2B-2Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(-3a^2(A+B)+ab(3A+B)+2Ab^2)}{a^2-b^2}}{3a(a^2-b^2)}$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(-3a^3B+6a^2Ab-ab^2B-2Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(-3a^2(A+B)+ab(3A+B)+2Ab^2)}{a^2-b^2}}{a^2-b^2}}{3a(a^2-b^2)}$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2b(Ab-aB)\sin(c+dx)\sqrt{\cos(c+dx)}}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} + \frac{2(a-b)\sqrt{a+b}(-3a^3B+6a^2Ab-ab^2B-2Ab^3)\cot(c+dx)\sqrt{a+b\cos(c+dx)}}{3ad(a^2-b^2)(a+b\cos(c+dx))^{3/2}} \right)$$

input `Int[((A + B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^(5/2),x]`

3.627. $\int \frac{(A+B\cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b\cos(c+dx))^{5/2}} dx$

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*b*(A*b - a*B)*Sqrt[Cos[c + d*x]]
*Sin[c + d*x])/(3*a*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + (((2*(a -
b)*Sqrt[a + b]*(6*a^2*A*b - 2*A*b^3 - 3*a^3*B - a*b^2*B)*Cot[c + d*x]*Elli
pticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -
((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c
+ d*x]))/(a - b))]/(a^2*d) - (2*(a - b)*Sqrt[a + b]*(2*A*b^2 - 3*a^2*(A +
B) + a*b*(3*A + B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]
]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[
c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a^2 - b^
2) - (2*(6*a^2*A*b - 2*A*b^3 - 3*a^3*B - a*b^2*B)*Sin[c + d*x])/((a^2 - b^
2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]])/(3*a*(a^2 - b^2)))
```

3.627.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)])*Sqrt[(a_.) + (b_)*sin[(e_.) + (f
_)*(x_)])], x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 3440 Int[(csc[(e_.) + (f_)*(x_)])*(g_.)^(p_)*((a_.) + (b_)*sin[(e_.) + (f_)*
(x_)])^(m_)*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3479 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n + 2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)*(m + n + 3)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && RationalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(IntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0])))`

3.627.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 5286 vs. $2(429) = 858$.

Time = 18.88 (sec) , antiderivative size = 5287, normalized size of antiderivative = 11.27

method	result	size
default	Expression too large to display	5287
parts	Expression too large to display	5409

input `int((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

3.627.5 Fricas [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c))/(b^3*cos(d*x + c)^3 + 3*a*b^2*cos(d*x + c)^2 + 3*a^2*b*cos(d*x + c) + a^3), x)`

3.627.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(5/2),x)`

output `Timed out`

3.627. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{5/2}} dx$

3.627.7 Maxima [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(5/2),x, algorith="maxima")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^(5/2), x)`

3.627.8 Giac [F]

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(B \cos(dx + c) + A) \sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(5/2),x, algorith="giac")`

output `integrate((B*cos(d*x + c) + A)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)^(5/2), x)`

3.627.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(c + dx)) \sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{5/2}} dx = \int \frac{(A + B \cos(c + dx)) \sqrt{\frac{1}{\cos(c + dx)}}}{(a + b \cos(c + dx))^{5/2}} dx$$

input `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^(5/2),x)`

output `int(((A + B*cos(c + d*x))*(1/cos(c + d*x))^(1/2))/(a + b*cos(c + d*x))^(5/2), x)`

3.627. $\int \frac{(A+B \cos(c+dx)) \sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{5/2}} dx$

3.628
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$$

3.628.1 Optimal result 5761
 3.628.2 Mathematica [A] (warning: unable to verify) 5762
 3.628.3 Rubi [A] (verified) 5762
 3.628.4 Maple [B] (warning: unable to verify) 5767
 3.628.5 Fricas [F] 5768
 3.628.6 Sympy [F(-1)] 5768
 3.628.7 Maxima [F] 5768
 3.628.8 Giac [F] 5769
 3.628.9 Mupad [F(-1)] 5769

3.628.1 Optimal result

Integrand size = 35, antiderivative size = 431

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx =$$

$$\frac{2(3a^2 A + Ab^2 - 4abB) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right) \mid -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a+b}}}{3a^2(a-b)(a+b)^{3/2} d \sqrt{\sec(c+dx)}} +$$

$$\frac{2(a(3A + B) - b(A + 3B)) \sqrt{\cos(c + dx)} \csc(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{3a(a-b)(a+b)^{3/2} d \sqrt{\sec(c+dx)}} -$$

$$\frac{2(Ab - aB) \sin(c + dx)}{3(a^2 - b^2) d(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} +$$

$$\frac{2(3a^2 A + Ab^2 - 4abB) \sqrt{\sec(c + dx)} \sin(c + dx)}{3(a^2 - b^2)^2 d \sqrt{a + b \cos(c + dx)}}$$

output

```
-2/3*(A*b-B*a)*sin(d*x+c)/(a^2-b^2)/d/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2)+2/3*(3*A*a^2+A*b^2-4*B*a*b)*sin(d*x+c)*sec(d*x+c)^(1/2)/(a^2-b^2)^2/d/(a+b*cos(d*x+c))^(1/2)-2/3*(3*A*a^2+A*b^2-4*B*a*b)*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a^2/(a-b)/(a+b)^(3/2)/d/sec(d*x+c)^(1/2)+2/3*(a*(3*A+B)-b*(A+3*B))*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/(a-b)/(a+b)^(3/2)/d/sec(d*x+c)^(1/2)
```

3.628.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sqrt{\sec(c+dx)}} dx$$

3.628.2 Mathematica [A] (warning: unable to verify)

Time = 14.53 (sec) , antiderivative size = 528, normalized size of antiderivative = 1.23

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \frac{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \left(-\frac{2(3a^2A + Ab^2 - 4abB) \sin(c + dx)}{3a(a^2 - b^2)^2} + \dots \right)}{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)} \sec(c + dx) \left(-2(a + b)(3a^2A + Ab^2 - 4abB) \sqrt{\frac{\cos(c + dx)}{1 + \cos(c + dx)}} \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} E(\arcsin \dots) \right)}$$

```
input Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(5/2)*Sqrt[Sec[c + d*x]]),x]
```

```
output (Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((-2*(3*a^2*A + A*b^2 - 4*a*b*B)*Sin[c + d*x])/(3*a*(a^2 - b^2)^2) + (2*(-(a*A*b*Sin[c + d*x]) + a^2*B*Sin[c + d*x]))/(3*b*(-a^2 + b^2)*(a + b*Cos[c + d*x])^2) + (2*(2*a^2*A*b*Sin[c + d*x] + 2*A*b^3*Sin[c + d*x] + a^3*B*Sin[c + d*x] - 5*a*b^2*B*Sin[c + d*x]))/(3*b*(-a^2 + b^2)^2*(a + b*Cos[c + d*x])))/d - (2*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(-2*(a + b)*(3*a^2*A + A*b^2 - 4*a*b*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] *Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + 2*a*(a + b)*(3*a*A + A*b - a*B - 3*b*B)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] *Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - (3*a^2*A + A*b^2 - 4*a*b*B)*Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/(3*a*(a^2 - b^2)^2*d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2])
```

3.628.3 Rubi [A] (verified)

Time = 1.79 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 3440, 3042, 3478, 27, 3042, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))^{5/2}} dx \quad \downarrow \quad 3042$$

3.628. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx$

$$\begin{aligned}
 & \int \frac{A + B \sin \left(c + dx + \frac{\pi}{2} \right)}{\sqrt{\csc \left(c + dx + \frac{\pi}{2} \right)} \left(a + b \sin \left(c + dx + \frac{\pi}{2} \right) \right)^{5/2}} dx \\
 & \quad \downarrow \text{3440} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)} (A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} (A + B \sin \left(c + dx + \frac{\pi}{2} \right))}{(a + b \sin \left(c + dx + \frac{\pi}{2} \right))^{5/2}} dx \\
 & \quad \downarrow \text{3478} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(-\frac{2 \int \frac{Ab - aB - 3(aA - bB) \cos(c + dx)}{2\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(-\frac{\int \frac{Ab - aB - 3(aA - bB) \cos(c + dx)}{\sqrt{\cos(c + dx)}(a + b \cos(c + dx))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(-\frac{\int \frac{Ab - aB - 3(aA - bB) \sin \left(c + dx + \frac{\pi}{2} \right)}{\sqrt{\sin \left(c + dx + \frac{\pi}{2} \right)} (a + b \sin \left(c + dx + \frac{\pi}{2} \right))^{3/2}} dx}{3(a^2 - b^2)} - \frac{2(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3472} \\
 & \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(-\frac{\int \frac{3Aa^2 - 4bBa + Ab^2 + (-Ba^2 + 4Aba - 3b^2B) \cos(c + dx)}{\cos^{\frac{3}{2}}(c + dx) \sqrt{a + b \cos(c + dx)} (a^2 - b^2)} dx}{3(a^2 - b^2)} - \frac{2(3a^2A - 4abB + Ab^2) \sin(c + dx)}{d(a^2 - b^2) \sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} - \frac{2(Ab - aB) \sin(c + dx) \sqrt{\cos(c + dx)}}{3d(a^2 - b^2)(a + b \cos(c + dx))^{3/2}} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.628. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3Aa^2-4bBa+Ab^2+(-Ba^2+4Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2} - \frac{2(3a^2A-4abB+Ab^2)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} \right) \frac{1}{3(a^2-b^2)}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3a^2A-4abB+Ab^2)\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b)(a(3A+B)-b(A+3B))\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{a^2-b^2} \right) \frac{1}{3(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3a^2A-4abB+Ab^2)\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(a(3A+B)-b(A+3B))\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}}}{a^2-b^2} \right) \frac{1}{3(a^2-b^2)}$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{(3a^2A-4abB+Ab^2)\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(a(3A+B)-b(A+3B))\cot(c+dx)}{a^2-b^2}}{a^2-b^2} \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2(a-b)\sqrt{a+b}(3a^2A-4abB+Ab^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right) - \frac{a}{a-b}}{a^2d} \right)$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(5/2)*Sqrt[Sec[c + d*x]]),x]`

3.628. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{5/2}\sqrt{\sec(c+dx)}} dx$

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((-2*(A*b - a*B)*Sqrt[Cos[c + d*x]]*
Sin[c + d*x])/(3*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) - (((2*(a - b)*
Sqrt[a + b]*(3*a^2*A + A*b^2 - 4*a*b*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt
[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))
]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)
])/ (a^2*d) - (2*(a - b)*Sqrt[a + b]*(a*(3*A + B) - b*(A + 3*B))*Cot[c + d*
x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]
])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1
+ Sec[c + d*x]))/(a - b)]/(a*d))/(a^2 - b^2) - (2*(3*a^2*A + A*b^2 - 4*a*
b*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*
x]]))/(3*(a^2 - b^2)))
```

3.628.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_.) + (f_)*(x_)]]*Sqrt[(a_.) + (b_)*sin[(e_.) + (f
_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 3440 Int[(csc[(e_.) + (f_)*(x_)]*(g_.)^(p_))*((a_.) + (b_)*sin[(e_.) + (f_)*
(x_)]^(m_))*((c_.) + (d_)*sin[(e_.) + (f_)*(x_)]^(n_), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
negerQ[n])
```

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

rule 3473 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`

rule 3478 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(B*a - A*b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin[e + f*x])^n/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^(n - 1)*Simp[c*(a*A - b*B)*(m + 1) + d*n*(A*b - a*B) + (d*(a*A - b*B)*(m + 1) - c*(A*b - a*B)*(m + 2))*Sin[e + f*x] - d*(A*b - a*B)*(m + n + 2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && LtQ[m, -1] && GtQ[n, 0]`

3.628.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4566 vs. $2(391) = 782$.

Time = 17.01 (sec) , antiderivative size = 4567, normalized size of antiderivative = 10.60

method	result	size
default	Expression too large to display	4567
parts	Expression too large to display	4847

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2)/sec(d*x+c)^(1/2),x,method=_RET
URNVERBOSE)
```

```
output 2/3/d*(csc(d*x+c)^2*(1-cos(d*x+c))^2+1)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-
csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a*b)/(csc(d*x+c)^2*(1-cos(d*x+c))^2+1))^(1
/2)*(-2*B*a^4*(csc(d*x+c)-cot(d*x+c))-3*A*a^4*(csc(d*x+c)-cot(d*x+c))+A*b^
4*(csc(d*x+c)-cot(d*x+c))+6*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((c
sc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a*b)/(a+b))
^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^3*b+4*A*(-c
sc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-cs
c(d*x+c)^2*b*(1-cos(d*x+c))^2+a*b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d
*x+c),(-(a-b)/(a+b))^(1/2))*a^2*b^2+2*A*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)
^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a
+b)/(a+b))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b
^3+5*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*
x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a*b)/(a+b))^(1/2)*EllipticF(cot(d*
x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a^3*b+7*B*(-csc(d*x+c)^2*(1-cos(d*x+
c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x
+c))^2+a*b)/(a+b))^(1/2)*EllipticF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1
/2))*a^2*b^2+3*B*(-csc(d*x+c)^2*(1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a
*(1-cos(d*x+c))^2-csc(d*x+c)^2*b*(1-cos(d*x+c))^2+a*b)/(a+b))^(1/2)*Ellipt
icF(cot(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*a*b^3-4*B*(-csc(d*x+c)^2*(
1-cos(d*x+c))^2+1)^(1/2)*((csc(d*x+c)^2*a*(1-cos(d*x+c))^2-csc(d*x+c)^2...
```

3.628.5 Fricas [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral((B*cos(d*x + c) + A)*sqrt(b*cos(d*x + c) + a)/((b^3*cos(d*x + c)^3 + 3*a*b^2*cos(d*x + c)^2 + 3*a^2*b*cos(d*x + c) + a^3)*sqrt(sec(d*x + c))), x)`

3.628.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2)/sec(d*x+c)**(1/2),x)`

output `Timed out`

3.628.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sqrt(sec(d*x + c))), x)`

3.628.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sqrt(sec(d*x +
c))), x)`

3.628.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sqrt{\sec(c + dx)}} dx = \int \frac{A + B \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(5/2
)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(5/2
)), x)`

$$3.629 \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

3.629.1 Optimal result	5770
3.629.2 Mathematica [B] (verified)	5771
3.629.3 Rubi [A] (verified)	5772
3.629.4 Maple [B] (warning: unable to verify)	5778
3.629.5 Fricas [F(-1)]	5778
3.629.6 Sympy [F(-1)]	5779
3.629.7 Maxima [F]	5779
3.629.8 Giac [F]	5779
3.629.9 Mupad [F(-1)]	5780

3.629.1 Optimal result

Integrand size = 35, antiderivative size = 602

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{2(4Ab^3 + 3a^3B - 7ab^2B) \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a}{a-b}\right) + 2(3Ab^3 + 3a^3B + a^2bB - ab^2(A + 6B)) \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a}{a-b}\right) - 3a(a-b)b^2(a+b)^{3/2}d\sqrt{\sec(c+dx)} + 2\sqrt{a+b}B\sqrt{\cos(c+dx)} \csc(c+dx) \text{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b \cos(c+dx)}}{\sqrt{a+b \sqrt{\cos(c+dx)}}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a+b}}}{b^3d\sqrt{\sec(c+dx)}} + \frac{2a(Ab - aB) \sin(c + dx)}{3b(a^2 - b^2) d(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} - \frac{2a(4Ab^3 + 3a^3B - 7ab^2B) \sqrt{\sec(c + dx)} \sin(c + dx)}{3b^2(a^2 - b^2)^2 d\sqrt{a + b \cos(c + dx)}}$$

$$3.629. \quad \int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^{\frac{3}{2}}(c+dx)} dx$$

output $\frac{2}{3}a(Ab - Ba)\sin(dx+c)/b/(a^2-b^2)/d/(a+b\cos(dx+c))^{3/2}/\sec(dx+c)^{1/2} - \frac{2}{3}a(4Ab^3+3Ba^3-7Bab^2)\sin(dx+c)\sec(dx+c)^{1/2}/b^2/(a^2-b^2)^2/d/(a+b\cos(dx+c))^{1/2} + \frac{2}{3}(4Ab^3+3Ba^3-7Bab^2)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/(a-b)/b^2/(a+b)^{3/2}/d/\sec(dx+c)^{1/2} - \frac{2}{3}(3Aab^3+3Ba^3+Bab^2-ab^2(A+6B))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, ((-a-b)/(a-b))^{1/2})\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/(a-b)/b^2/(a+b)^{3/2}/d/\sec(dx+c)^{1/2} - 2B\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}, (a+b)/b, ((-a-b)/(a-b))^{1/2})\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^3/d/\sec(dx+c)^{1/2}$

3.629.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1496 vs. $2(602) = 1204$.

Time = 14.20 (sec) , antiderivative size = 1496, normalized size of antiderivative = 2.49

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(3/2)),x]`

output

```
(Sqrt[a + b*cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*(4*A*b^3 + 3*a^3*B - 7*a*b^2*B)*Sin[c + d*x])/(3*b^2*(-a^2 + b^2)^2) - (2*(-(a^2*A*b*SIN[c + d*x]) + a^3*B*SIN[c + d*x]))/(3*b^2*(-a^2 + b^2)*(a + b*cos[c + d*x])^2) - (2*(-(a^3*A*b*SIN[c + d*x]) + 5*a*A*b^3*SIN[c + d*x] + 4*a^4*B*SIN[c + d*x] - 8*a^2*b^2*B*SIN[c + d*x]))/(3*b^2*(-a^2 + b^2)^2*(a + b*cos[c + d*x])))/d + (2*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(1 + Tan[(c + d*x)/2]^2)]*(-4*a*A*b^3*Tan[(c + d*x)/2] - 4*A*b^4*Tan[(c + d*x)/2] - 3*a^4*B*Tan[(c + d*x)/2] - 3*a^3*b*B*Tan[(c + d*x)/2] + 7*a^2*b^2*B*Tan[(c + d*x)/2] + 7*a*b^3*B*Tan[(c + d*x)/2] + 8*A*b^4*Tan[(c + d*x)/2]^3 + 6*a^3*b*B*Tan[(c + d*x)/2]^3 - 14*a*b^3*B*Tan[(c + d*x)/2]^3 + 4*a*A*b^3*Tan[(c + d*x)/2]^5 - 4*A*b^4*Tan[(c + d*x)/2]^5 + 3*a^4*B*Tan[(c + d*x)/2]^5 - 3*a^3*b*B*Tan[(c + d*x)/2]^5 - 7*a^2*b^2*B*Tan[(c + d*x)/2]^5 + 7*a*b^3*B*Tan[(c + d*x)/2]^5 + 6*a^4*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] - 12*a^2*b^2*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + 6*b^4*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]*Sqrt[1 - Tan[(c + d*x)/2]^2]*Sqrt[(a + b + a*Tan[(c + d*x)/2]^2 - b*Tan[(c + d*x)/2]^2)/(a + b)] + 6*a^4*B*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)...
```

3.629.3 Rubi [A] (verified)

Time = 2.71 (sec) , antiderivative size = 590, normalized size of antiderivative = 0.98, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3530, 3042, 3288, 3472, 3042, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin\left(c + dx + \frac{\pi}{2}\right)}{\csc\left(c + dx + \frac{\pi}{2}\right)^{3/2} (a + b \sin\left(c + dx + \frac{\pi}{2}\right))^{5/2}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{3}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{5/2}} dx$$

3.629. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})^{3/2} (A+B \sin(c+dx+\frac{\pi}{2}))}{(a+b \sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
 & \downarrow \text{3468} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b \cos(c+dx))^{3/2}} - \frac{2 \int -\frac{3(a^2-b^2)B \cos^2(c+dx)-3b(Ab-aB) \cos(c+dx)}{2\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3(a^2-b^2)B \cos^2(c+dx)-3b(Ab-aB) \cos(c+dx)+a(Ab-aB)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3(a^2-b^2)B \sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB) \sin(c+dx+\frac{\pi}{2})+a(Ab-aB)}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \right) \\
 & \downarrow \text{3530} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{ab(Ab-aB)-3(Ab^3+a(a^2-2b^2)B) \cos(c+dx)}{\sqrt{\cos(c+dx)}(a+b \cos(c+dx))^{3/2}} dx}{b} + \frac{3B(a^2-b^2) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx}{b} + \frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{ab(Ab-aB)-3(Ab^3+a(a^2-2b^2)B) \sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b \sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{b} + \frac{3B(a^2-b^2) \int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx}{b} + \frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b \cos(c+dx))^{3/2}} \right) \\
 & \downarrow \text{3288}
 \end{aligned}$$

3.629. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^3(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{ab(Ab-aB)-3(Ab^3+a(a^2-2b^2)B)\sin(c+dx+\frac{\pi}{2})}{\sqrt{\sin(c+dx+\frac{\pi}{2})(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}}} dx}{b} - \frac{6B\sqrt{a+b}(a^2-b^2)\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)-1)}{a+b}}}{3b(a^2-b^2)} \right)$$

↓ 3472

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(3Ba^3-7b^2Ba+4Ab^3)+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3)\cos(c+dx)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{b} - \frac{2a(3a^3B-7ab^2B+4Ab^3)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{3b(a^2-b^2)}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{a(3Ba^3-7b^2Ba+4Ab^3)+b(2Ba^3+Ab^2-6b^2Ba+3Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{b} - \frac{2a(3a^3B-7ab^2B+4Ab^3)\sin(c+dx)}{d(a^2-b^2)\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{3b(a^2-b^2)}$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(3a^3B-7ab^2B+4Ab^3)\int \frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx - (a-b)(3a^3B+a^2bB-ab^2(A+6B)+3Ab^3)\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2}}{b}$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(3a^3B-7ab^2B+4Ab^3)\int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - (a-b)(3a^3B+a^2bB-ab^2(A+6B)+3Ab^3)\int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{a^2-b^2}}{b}$$

↓ 3295

3.629. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{5/2}\sec^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{a(3a^3B-7ab^2B+4Ab^3) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx+\frac{\pi}{2})}} dx - \frac{2(a-b)\sqrt{a+b}(3a^3B+a^2bB-ab^2(A+6B)+3Ab^3)}{a^2-b^2}}{a^2-b^2} \right)$$

↓ 3473

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(Ab-aB) \sin(c+dx) \sqrt{\cos(c+dx)}}{3bd(a^2-b^2)(a+b \cos(c+dx))^{3/2}} + \frac{2(a-b)\sqrt{a+b}(3a^3B-7ab^2B+4Ab^3) \cot(c+dx) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{a^2-b^2} \right)$$

input `Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(3/2)),x]`

output `Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*(A*b - a*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + ((-6*Sqrt[a + b]*(a^2 - b^2)*B*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b^2*d) + (((2*(a - b)*Sqrt[a + b]*(4*A*b^3 + 3*a^3*B - 7*a*b^2*B)*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d) - (2*(a - b)*Sqrt[a + b]*(3*A*b^3 + 3*a^3*B + a^2*b*B - a*b^2*(A + 6*B))*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))/(a^2 - b^2) - (2*a*(4*A*b^3 + 3*a^3*B - 7*a*b^2*B)*Sin[c + d*x])/((a^2 - b^2)*d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]))/b)/(3*b*(a^2 - b^2))`

3.629.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3288 `Int[Sqrt[(b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d), x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`
- rule 3295 `Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b), x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`
- rule 3440 `Int[(csc[(e_) + (f_)*(x_)])*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)])*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]])), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

rule 3473 `Int[((A_) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*A*(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] && PosQ[(c + d)/b]`

rule 3477 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[(A - B)/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[(A*b - a*B)/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A, B]`


```
rule 3530 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_
)])^(3/2)), x_Symbol] := Simp[C/(b*d) Int[Sqrt[d*Sin[e + f*x]]/Sqrt[a + b
*Sin[e + f*x]], x], x] + Simp[1/b Int[(A*b + (b*B - a*C)*Sin[e + f*x])/((
a + b*Sin[e + f*x])^(3/2)*Sqrt[d*Sin[e + f*x])], x], x] /; FreeQ[{a, b, d,
e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0]
```

3.629.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 6417 vs. $2(550) = 1100$.

Time = 19.33 (sec) , antiderivative size = 6418, normalized size of antiderivative = 10.66

method	result	size
default	Expression too large to display	6418
parts	Expression too large to display	6730

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2)/sec(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.629.5 Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algo
rithm="fricas")
```

```
output Timed out
```

3.629.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2)/sec(d*x+c)**(3/2),x)`

output `Timed out`

3.629.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(3/2)), x)`

3.629.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{3/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(3/2)), x)`

3.629.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(5/2)), x)`

3.630
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$$

3.630.1 Optimal result	5781
3.630.2 Mathematica [B] (warning: unable to verify)	5782
3.630.3 Rubi [A] (verified)	5783
3.630.4 Maple [B] (warning: unable to verify)	5790
3.630.5 Fricas [F(-1)]	5790
3.630.6 Sympy [F(-1)]	5791
3.630.7 Maxima [F]	5791
3.630.8 Giac [F]	5791
3.630.9 Mupad [F(-1)]	5792

3.630.1 Optimal result

Integrand size = 35, antiderivative size = 733

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx = \frac{(6a^3Ab - 14aAb^3 - 15a^4B + 26a^2b^2B - 3b^4B) \sqrt{\cos(c + dx)} \operatorname{csc}(c + dx)}{3a(a - b)b^3} + \frac{(3b^3(4A - B) + 15a^3B - ab^2(2A + 21B) - a^2(6Ab - 5bB)) \sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\sqrt{\cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{3(a - b)b^3(a + b)^{3/2}d\sqrt{\sec(c + dx)}} - \frac{\sqrt{a + b}(2Ab - 5aB) \sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b}\sqrt{\cos(c+dx)}}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}}}{b^4d\sqrt{\sec(c + dx)}} + \frac{2a(Ab - aB) \sin(c + dx)}{3b(a^2 - b^2) d(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} + \frac{2a(2a^2Ab - 6Ab^3 - 5a^3B + 9ab^2B) \sin(c + dx)}{3b^2(a^2 - b^2)^2 d \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} - \frac{(6a^3Ab - 14aAb^3 - 15a^4B + 26a^2b^2B - 3b^4B) \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{3b^3(a^2 - b^2)^2 d}$$

3.630.
$$\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$$

output $\frac{2}{3}a(Ab-Ba)\sin(dx+c)/b/(a^2-b^2)/d/(a+b\cos(dx+c))^{3/2}/\sec(dx+c)^{3/2}+2/3a(2Aa^2b-6Ab^3-5Ba^3+9Bab^2)\sin(dx+c)/b^2/(a^2-b^2)^2/d/(a+b\cos(dx+c))^{1/2}/\sec(dx+c)^{1/2}-1/3(6Aa^3b-14Aab^3-15Ba^4+26Ba^2b^2-3Bb^4)\sin(dx+c)(a+b\cos(dx+c))^{1/2}\sec(dx+c)^{1/2}/b^3/(a^2-b^2)^2/d+1/3(6Aa^3b-14Aab^3-15Ba^4+26Ba^2b^2-3Bb^4)\csc(dx+c)\text{EllipticE}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/a/(a-b)/b^3/(a+b)^{3/2}/d/\sec(dx+c)^{1/2}+1/3(3b^3(4A-B)+15Ba^3-ab^2(2A+21B)-a^2(6Ab-5Bb))\csc(dx+c)\text{EllipticF}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}),((-a-b)/(a-b))^{1/2})\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/(a-b)/b^3/(a+b)^{3/2}/d/\sec(dx+c)^{1/2}-(2Ab-5Ba)\csc(dx+c)\text{EllipticPi}((a+b\cos(dx+c))^{1/2}/(a+b)^{1/2}/\cos(dx+c)^{1/2}), (a+b)/b,((-a-b)/(a-b))^{1/2})(a+b)^{1/2}\cos(dx+c)^{1/2}(a(1-\sec(dx+c))/(a+b))^{1/2}(a(1+\sec(dx+c))/(a-b))^{1/2}/b^4/d/\sec(dx+c)^{1/2}$

3.630.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 2318 vs. $2(733) = 1466$.

Time = 20.14 (sec) , antiderivative size = 2318, normalized size of antiderivative = 3.16

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^2(c + dx)} dx = \text{Result too large to show}$$

input `Integrate[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(5/2)),x]`

output $(\text{Sqrt}[a + b \cos[c + dx]] \cdot \text{Sqrt}[\text{Sec}[c + dx]] \cdot ((-2a(-3a^2Ab + 7A^3b^3 + 6a^3B - 10ab^2B) \sin[c + dx]) / (3b^3(a^2 - b^2)^2) + (2(-a^3Ab \sin[c + dx]) + a^4B \sin[c + dx])) / (3b^3(-a^2 + b^2)(a + b \cos[c + dx])^2) + (2(-4a^4Ab \sin[c + dx] + 8a^2Ab^3 \sin[c + dx] + 7a^5B \sin[c + dx] - 11a^3b^2B \sin[c + dx])) / (3b^3(-a^2 + b^2)^2(a + b \cos[c + dx])))) / d + (\text{Sqrt}[(1 - \text{Tan}[(c + dx)/2])^{-1}] \cdot \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2) / (1 + \text{Tan}[(c + dx)/2]^2)] \cdot (6a^4Ab \text{Tan}[(c + dx)/2] + 6a^3Ab^2 \text{Tan}[(c + dx)/2] - 14a^2Ab^3 \text{Tan}[(c + dx)/2] - 14aAb^4 \text{Tan}[(c + dx)/2] - 15a^5B \text{Tan}[(c + dx)/2] - 15a^4bB \text{Tan}[(c + dx)/2] + 26a^3b^2B \text{Tan}[(c + dx)/2] + 26a^2b^3B \text{Tan}[(c + dx)/2] - 3ab^4B \text{Tan}[(c + dx)/2] - 3b^5B \text{Tan}[(c + dx)/2] - 12a^3Ab^2 \text{Tan}[(c + dx)/2]^3 + 28aAb^4 \text{Tan}[(c + dx)/2]^3 + 30a^4bB \text{Tan}[(c + dx)/2]^3 - 52a^2b^3B \text{Tan}[(c + dx)/2]^3 + 6b^5B \text{Tan}[(c + dx)/2]^3 - 6a^4Ab \text{Tan}[(c + dx)/2]^5 + 6a^3Ab^2 \text{Tan}[(c + dx)/2]^5 + 14a^2Ab^3 \text{Tan}[(c + dx)/2]^5 - 14aAb^4 \text{Tan}[(c + dx)/2]^5 + 15a^5B \text{Tan}[(c + dx)/2]^5 - 15a^4bB \text{Tan}[(c + dx)/2]^5 - 26a^3b^2B \text{Tan}[(c + dx)/2]^5 + 26a^2b^3B \text{Tan}[(c + dx)/2]^5 + 3ab^4B \text{Tan}[(c + dx)/2]^5 - 3b^5B \text{Tan}[(c + dx)/2]^5 - 12a^4Ab \text{EllipticPi}[-1, \text{ArcSin}[\text{Tan}[(c + dx)/2]], (-a + b)/(a + b)] \cdot \text{Sqrt}[1 - \text{Tan}[(c + dx)/2]^2] \cdot \text{Sqrt}[(a + b + a \text{Tan}[(c + dx)/2]^2 - b \text{Tan}[(c + dx)/2]^2)/(a + b)] + 24a^2Ab^3 \dots$

3.630.3 Rubi [A] (verified)

Time = 3.65 (sec) , antiderivative size = 719, normalized size of antiderivative = 0.98, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 3440, 3042, 3468, 27, 3042, 3526, 27, 3042, 3540, 3042, 3532, 3042, 3288, 3477, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \cos(c + dx)}{\sec^{\frac{5}{2}}(c + dx)(a + b \cos(c + dx))^{\frac{5}{2}}} dx$$

↓ 3042

$$\int \frac{A + B \sin(c + dx + \frac{\pi}{2})}{\csc(c + dx + \frac{\pi}{2})^{\frac{5}{2}}(a + b \sin(c + dx + \frac{\pi}{2}))^{\frac{5}{2}}} dx$$

↓ 3440

$$\sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{5}{2}}(c + dx)(A + B \cos(c + dx))}{(a + b \cos(c + dx))^{\frac{5}{2}}} dx$$

3.630. $\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{\frac{5}{2}} \sec^{\frac{5}{2}}(c + dx)} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \int \frac{\sin(c+dx+\frac{\pi}{2})^{5/2} (A+B\sin(c+dx+\frac{\pi}{2}))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
 & \downarrow \text{3468} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(Ab-aB)\sin(c+dx)\cos^3(c+dx)}{3bd(a^2-b^2)(a+b\cos(c+dx))^{3/2}} - \frac{2 \int -\frac{\sqrt{\cos(c+dx)}(-((-5Ba^2+2Aba+3b^2B)\cos^2(c+dx)-2(a+b\cos(c+dx)))}{2(a+b\cos(c+dx))}}{3b(a^2-b^2)} dx \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\cos(c+dx)}(-((-5Ba^2+2Aba+3b^2B)\cos^2(c+dx)-3b(Ab-aB)\cos(c+dx)+3a(Ab-aB)))}{(a+b\cos(c+dx))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB)}{3b(a^2-b^2)} \right) \\
 & \downarrow \text{3042} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}((5Ba^2-2Aba-3b^2B)\sin(c+dx+\frac{\pi}{2})^2-3b(Ab-aB)\sin(c+dx+\frac{\pi}{2})+3a(Ab-aB))}{(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx}{3b(a^2-b^2)} + \frac{2a(Ab-aB)}{3b(a^2-b^2)} \right) \\
 & \downarrow \text{3526} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(-5a^3B+2a^2Ab+9ab^2B-6Ab^3)\sin(c+dx)\sqrt{\cos(c+dx)}}{bd(a^2-b^2)\sqrt{a+b\cos(c+dx)}} - \frac{2 \int -\frac{((-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)\cos^2(c+dx)+b(2Ba^3+Aba^2-6b^2Ba+3Ab^3)\cos(c+dx)+a(-5Ba^3-3b^4B))}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{b(a^2-b^2)} dx}{3b(a^2-b^2)} \right) \\
 & \downarrow \text{27} \\
 & \sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{-((-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)\cos^2(c+dx)+b(2Ba^3+Aba^2-6b^2Ba+3Ab^3)\cos(c+dx)+a(-5Ba^3-3b^4B))}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{b(a^2-b^2)} dx}{3b(a^2-b^2)} \right) \\
 & \downarrow \text{3042}
 \end{aligned}$$

3.630. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{5/2}\sec^2(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{(15Ba^4 - 6Aba^3 - 26b^2Ba^2 + 14Ab^3a + 3b^4B) \sin(c+dx + \frac{\pi}{2})^2 + b(2Ba^3 + Aba^2 - 6b^2Ba + 3Ab^3) \sin(c+dx + \frac{\pi}{2}) + a(-5Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{\sqrt{\sin(c+dx + \frac{\pi}{2})} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \frac{a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{b(a^2 - b^2)} \right) \quad 3b(a^2 - b^2)$$

↓ 3540

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3(a^2 - b^2)^2(2Ab - 5aB) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx + \frac{a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{b(a^2 - b^2)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{3(a^2 - b^2)^2(2Ab - 5aB) \sin(c+dx + \frac{\pi}{2})^2 + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \sin(c+dx + \frac{\pi}{2}) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \frac{a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{b(a^2 - b^2)} \right)$$

↓ 3532

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3(a^2 - b^2)^2(2Ab - 5aB) \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{a+b \cos(c+dx)}} dx + \int \frac{a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{\cos^{\frac{3}{2}}(c+dx) \sqrt{a+b \cos(c+dx)}} dx}{b(a^2 - b^2)} \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{3(a^2 - b^2)^2(2Ab - 5aB) \int \frac{\sqrt{\sin(c+dx + \frac{\pi}{2})}}{\sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx + \int \frac{a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos^2(c+dx) + 2ab(-5Ba^3 + 2Aba^2 + 9b^2Ba - 6Ab^3) \cos(c+dx) + a(-15Ba^4 + 6Aba^3 + 26b^2Ba^2 - 14Ab^3a - 3b^4B) \sin(c+dx + \frac{\pi}{2})}{\sin(c+dx + \frac{\pi}{2})^{3/2} \sqrt{a+b \sin(c+dx + \frac{\pi}{2})}} dx}{b(a^2 - b^2)} \right)$$

3.630. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^2(c+dx)} dx$

↓ 3288

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\int \frac{a(-15Ba^4+6Aba^3+26b^2Ba^2-14Ab^3a-3b^4B)+2ab(-5Ba^3+2Aba^2+9b^2Ba-6Ab^3)\sin(c+dx+\frac{\pi}{2})}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6\sqrt{a+b}(a^2-b^2)}{2b} \right)$$

↓ 3477

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(a(a-b)(15a^3B-a^2(6Ab-5bB)-ab^2(2A+21B)+3b^3(4A-B)) \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}} dx + a(-15a^4B+6a^3Ab+26a^2b^2B-14aAb^3-3b^4B) \right)$$

↓ 3042

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(a(a-b)(15a^3B-a^2(6Ab-5bB)-ab^2(2A+21B)+3b^3(4A-B)) \int \frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx + a(-15a^4B+6a^3Ab+26a^2b^2B-14aAb^3-3b^4B) \right)$$

↓ 3295

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(a(-15a^4B+6a^3Ab+26a^2b^2B-14aAb^3-3b^4B) \int \frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx - \frac{6\sqrt{a+b}(a^2-b^2)^2(2Ab-5b^2)}{2b^3} \right)$$

↓ 3473

3.630. $\int \frac{A+B\cos(c+dx)}{(a+b\cos(c+dx))^{5/2}\sec^2(c+dx)} dx$

$$\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{2a(Ab - aB) \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{3bd(a^2 - b^2)(a + b \cos(c+dx))^{3/2}} + \frac{2a(-5a^3B + 2a^2Ab + 9ab^2B - 6Ab^3) \sin(c+dx) \sqrt{\cos(c+dx)}}{bd(a^2 - b^2) \sqrt{a + b \cos(c+dx)}} \right)$$

```
input Int[(A + B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^(5/2)),x
]
```

```
output Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*a*(A*b - a*B)*Cos[c + d*x]^(3/2)
*Sin[c + d*x])/(3*b*(a^2 - b^2)*d*(a + b*Cos[c + d*x])^(3/2)) + ((2*a*(2*a
^2*A*b - 6*A*b^3 - 5*a^3*B + 9*a*b^2*B)*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(
b*(a^2 - b^2)*d*Sqrt[a + b*Cos[c + d*x]]) + (((2*(a - b)*Sqrt[a + b]*(6*a^
3*A*b - 14*a*A*b^3 - 15*a^4*B + 26*a^2*b^2*B - 3*b^4*B)*Cot[c + d*x]*Ellip
ticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(
(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c
+ d*x]))/(a - b))]/(a*d) + (2*(a - b)*Sqrt[a + b]*(3*b^3*(4*A - B) + 15*a^
3*B - a*b^2*(2*A + 21*B) - a^2*(6*A*b - 5*b*B))*Cot[c + d*x]*EllipticF[Arc
Sin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/
(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))
/(a - b))]/d - (6*Sqrt[a + b]*(a^2 - b^2)^2*(2*A*b - 5*a*B)*Cot[c + d*x]*E
llipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos
[c + d*x]])], -((a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqr
t[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d))/(2*b) - ((6*a^3*A*b - 14*a*A*b^3
- 15*a^4*B + 26*a^2*b^2*B - 3*b^4*B)*Sqrt[a + b*Cos[c + d*x]]*Sin[c + d*x
])/(b*d*Sqrt[Cos[c + d*x]])/(b*(a^2 - b^2)))/(3*b*(a^2 - b^2)))
```

3.630.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]]], x_Symbol] :> Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^p*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^m*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3468 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^m*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^n*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Simp[(-b*c - a*d)*(B*c - A*d)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^(m - 2)*(c + d*Sin[e + f*x])^(n + 1)*Simp[b*(b*c - a*d)*(B*c - A*d)*(m - 1) + a*d*(a*A*c + b*B*c - (A*b + a*B)*d)*(n + 1) + (b*(b*d*(B*c - A*d) + a*(A*c*d + B*(c^2 - 2*d^2)))*(n + 1) - a*(b*c - a*d)*(B*c - A*d)*(n + 2))*Sin[e + f*x] + b*(d*(A*b*c + a*B*c - a*A*d)*(m + n + 1) - b*B*(c^2*m + d^2*(n + 1)))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 3473 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[-2A(c - d)(\tan[e + fx]/(fbc^2)) \text{Rt}[(c + d)/b, 2] \sqrt{c((1 + \text{Csc}[e + fx])/(c - d))} \sqrt{c((1 - \text{Csc}[e + fx])/(c + d))} \text{EllipticE}[\text{ArcSin}[\sqrt{c + d \sin[e + fx]}/\sqrt{b \sin[e + fx]}/\text{Rt}[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; \text{FreeQ}\{b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{EqQ}[A, B] \&\& \text{PosQ}[(c + d)/b]$$`

rule 3477 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/((a_) + (b_)*sin[(e_) + (f_)*

$$\int \frac{(A + B \sin(e + fx)) \sqrt{(c + d \sin(e + fx))}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[(A - B)/(a - b) \int[1/(\sqrt{a + b \sin[e + fx]}\sqrt{c + d \sin[e + fx]}), x], x] - \text{Simp}[(A*b - a*B)/(a - b) \int[(1 + \sin[e + fx])/((a + b \sin[e + fx])^{3/2}\sqrt{c + d \sin[e + fx]}), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[A, B]$$`

rule 3526 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +

$$\int ((a + b \sin(e + fx))^m ((c + d \sin(e + fx))^n ((A + B \sin(e + fx)) + (C + f(x))^2))^{n+1} dx$$

$$\rightarrow \text{Simp}[-(c^2C - Bcd + Ad^2) \cos[e + fx] (a + b \sin[e + fx])^m ((c + d \sin[e + fx])^{n+1}) / (d f (n+1) (c^2 - d^2)), x] + \text{Simp}[1 / (d (n+1) (c^2 - d^2)) \int [(a + b \sin[e + fx])^{m-1} (c + d \sin[e + fx])^{n+1} \text{Simp}[Ad(bd^m + ac(n+1)) + (cC - Bd)(b^m + ad(n+1)) - (d(A(ad(n+2) - b^m(n+1)) + B(bd^m(n+1) - ac(n+2))) - C(bcd(n+1) - a(c^2 + d^2(n+1))) \sin[e + fx] + b(d(Bc - Ad)(m + n + 2) - C(c^2(m+1) + d^2(n+1))) \sin[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{GtQ}[m, 0] \&\& \text{LtQ}[n, -1]$$`

rule 3532 `Int[((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)]^2)/

$$\int \frac{(A + B \sin(e + fx) + (C + f(x))^2) \sqrt{(c + d \sin(e + fx))}}{(a + b \sin(e + fx))^{3/2}} dx$$

$$\rightarrow \text{Simp}[C/b^2 \int[\sqrt{a + b \sin[e + fx]}/\sqrt{c + d \sin[e + fx]}, x], x] + \text{Simp}[1/b^2 \int[(A*b^2 - a^2*C + b*(b*B - 2*a*C) \sin[e + fx]) / ((a + b \sin[e + fx])^{3/2} \sqrt{c + d \sin[e + fx]}), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{NeQ}[c^2 - d^2, 0]$$`

```
rule 3540 Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(c_.) + (d_.)*sin[(e_.)
+ (f_.)*(x_)]]), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(Sqrt[c + d*Sin[e + f
*x]]/(d*f*Sqrt[a + b*Sin[e + f*x]])), x] + Simp[1/(2*d) Int[(1/((a + b*Si
n[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]])]*Simp[2*a*A*d - C*(b*c - a*d) -
2*(a*c*C - d*(A*b + a*B))*Sin[e + f*x] + (2*b*B*d - C*(b*c + a*d))*Sin[e +
f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

3.630.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 10352 vs. $2(675) = 1350$.

Time = 21.29 (sec) , antiderivative size = 10353, normalized size of antiderivative = 14.12

method	result	size
parts	Expression too large to display	10353
default	Expression too large to display	10874

```
input int((A+B*cos(d*x+c))/(a+cos(d*x+c)*b)^(5/2)/sec(d*x+c)^(5/2),x,method=_RET
URNVERBOSE)
```

```
output result too large to display
```

3.630.5 Fracas [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algo
rithm="fracas")
```

```
output Timed out
```

3.630. $\int \frac{A+B \cos(c+dx)}{(a+b \cos(c+dx))^{5/2} \sec^{\frac{5}{2}}(c+dx)} dx$

3.630.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))**(5/2)/sec(d*x+c)**(5/2),x)`

output `Timed out`

3.630.7 Maxima [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algo
rithm="maxima")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(5
/2)), x)`

3.630.8 Giac [F]

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{5/2}(c + dx)} dx = \int \frac{B \cos(dx + c) + A}{(b \cos(dx + c) + a)^{5/2} \sec(dx + c)^{5/2}} dx$$

input `integrate((A+B*cos(d*x+c))/(a+b*cos(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algo
rithm="giac")`

output `integrate((B*cos(d*x + c) + A)/((b*cos(d*x + c) + a)^(5/2)*sec(d*x + c)^(5
/2)), x)`

3.630.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \cos(c + dx)}{(a + b \cos(c + dx))^{5/2} \sec^{\frac{5}{2}}(c + dx)} dx = \int \frac{A + B \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{5/2} (a + b \cos(c + dx))^{5/2}} dx$$

input `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(5/2)),x)`

output `int((A + B*cos(c + d*x))/((1/cos(c + d*x))^(5/2)*(a + b*cos(c + d*x))^(5/2)), x)`

3.631
$$\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

3.631.1 Optimal result 5793
 3.631.2 Mathematica [A] (verified) 5794
 3.631.3 Rubi [A] (verified) 5794
 3.631.4 Maple [B] (warning: unable to verify) 5797
 3.631.5 Fricas [F] 5798
 3.631.6 Sympy [F(-1)] 5798
 3.631.7 Maxima [F] 5799
 3.631.8 Giac [F] 5799
 3.631.9 Mupad [F(-1)] 5799

3.631.1 Optimal result

Integrand size = 38, antiderivative size = 266

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2(a - b)\sqrt{a + b}B\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx)E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)}{a^2 d \sqrt{\sec(c + dx)}} - \frac{2\sqrt{a + b}B\sqrt{\cos(c + dx)} \operatorname{csc}(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a-b}}}{ad\sqrt{\sec(c + dx)}}$$

output

```
2*(a-b)*B*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a^2/d/sec(d*x+c)^(1/2)-2*B*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/a/d/sec(d*x+c)^(1/2)
```


3.631.2 Mathematica [A] (verified)

Time = 6.06 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.12

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = B \left(\frac{2\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)} \sin(c + dx)}{ad} \right. \\ \left. - \frac{2\sqrt{\cos^2\left(\frac{1}{2}(c + dx)\right)} \sec(c + dx) \left(2(a + b) \sqrt{\frac{\cos(c + dx)}{1 + \cos(c + dx)}} \sqrt{\frac{a + b \cos(c + dx)}{(a + b)(1 + \cos(c + dx))}} E\left(\arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right) \mid \frac{-a + b}{a + b}\right)\right)}{ad} \right)$$

input `Integrate[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^(3/2),x]`

output `B*((2*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(a*d) - (2*Sqrt[Cos[(c + d*x)/2]^2*Sec[c + d*x]]*(2*(a + b)*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticE[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] - 2*a*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)] + Cos[c + d*x]*(a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2])/(a*d*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[(c + d*x)/2]^2]))`

3.631.3 Rubi [A] (verified)Time = 0.81 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.92, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {2011, 3042, 4710, 3042, 3280, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{\frac{3}{2}}(c + dx)(aB + bB \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

↓ 2011

$$B \int \frac{\sec^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx$$

↓ 3042

3.631. $\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$

$$\begin{aligned}
& B \int \frac{\csc(c + dx + \frac{\pi}{2})^{3/2}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4710} \\
& B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{3280} \\
& B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\int \frac{\cos(c + dx) + 1}{\cos^{3/2}(c + dx) \sqrt{a + b \cos(c + dx)}} dx - \int \frac{1}{\sqrt{\cos(c + dx)} \sqrt{a + b \cos(c + dx)}} dx \right) \\
& \quad \downarrow \text{3042} \\
& B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \int \frac{1}{\sqrt{\sin(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \right) \\
& \quad \downarrow \text{3295} \\
& B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\int \frac{\sin(c + dx + \frac{\pi}{2}) + 1}{\sin(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx - \frac{2\sqrt{a + b} \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}}}{a^2 d} \right) \\
& \quad \downarrow \text{3473} \\
& B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{2(a - b) \sqrt{a + b} \cot(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} E\left(\arcsin\left(\frac{\sqrt{a + b} \cos(c + dx)}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right)\right)}{a^2 d} \right)
\end{aligned}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sec[c + d*x]^(3/2))/(a + b*Cos[c + d*x])^(3/2),x]`

```
output B*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((2*(a - b)*Sqrt[a + b]*Cot[c + d*
x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x
]]]), -(a + b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1
+ Sec[c + d*x]))/(a - b)]/(a^2*d) - (2*Sqrt[a + b]*Cot[c + d*x]*EllipticF
[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]]]), -(a +
b)/(a - b))*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*
x]))/(a - b)]/(a*d))
```

3.631.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :>
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x
] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x
, a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3280 Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin
[(e_.) + (f_.)*(x_)]]), x_Symbol] :> Simp[1/(a - b) Int[1/(Sqrt[a + b*Sin
[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[b/(a - b) Int[(1 + Si
n[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] /
; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &&
NeQ[c^2 - d^2, 0]
```

```
rule 3295 Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f
_.)*(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

3.631.
$$\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{3/2}} dx$$

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)])
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 4710 Int[(csc[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Csc[a
+ b*x])^m*(c*Sin[a + b*x])^m Int[ActivateTrig[u]/(c*Sin[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownSineIntegrandQ[u, x]
```

3.631.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 658 vs. $2(242) = 484$.

Time = 20.43 (sec) , antiderivative size = 659, normalized size of antiderivative = 2.48

method	result
default	$- \frac{2B \left(- \frac{\csc^2(dx+c)(1-\cos(dx+c))^2+1}{\csc^2(dx+c)(1-\cos(dx+c))^2-1} \right)^{\frac{3}{2}} \left((\csc^2(dx+c))(1-\cos(dx+c))^2-1 \right) \left(-\sqrt{-(\csc^2(dx+c))(1-\cos(dx+c))^2+1} \sqrt{\csc^2(dx+c)} \right)}{\dots}$
parts	Expression too large to display

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+cos(d*x+c)*b)^(3/2),x,method=
_RETURNVERBOSE)
```

output
$$\begin{aligned} & -2*B/d*(-(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)/(\csc(d*x+c)^2*(1-\cos(d*x+c))^2-1))^{\frac{3}{2}}*(\csc(d*x+c)^2*(1-\cos(d*x+c))^2-1)*(-(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*((\csc(d*x+c)^2*a*(1-\cos(d*x+c))^2-\csc(d*x+c)^2*b*(1-\cos(d*x+c))^2+a+b)/(a+b))^{\frac{1}{2}}*\text{EllipticF}(\cot(d*x+c)-\csc(d*x+c),(-a-b)/(a+b))^{\frac{1}{2}}) \\ & +a*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*((\csc(d*x+c)^2*a*(1-\cos(d*x+c))^2-\csc(d*x+c)^2*b*(1-\cos(d*x+c))^2+a+b)/(a+b))^{\frac{1}{2}}*\text{EllipticE}(\cot(d*x+c)-\csc(d*x+c),(-a-b)/(a+b))^{\frac{1}{2}}) \\ & +a*(-\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)^{\frac{1}{2}}*((\csc(d*x+c)^2*a*(1-\cos(d*x+c))^2-\csc(d*x+c)^2*b*(1-\cos(d*x+c))^2+a+b)/(a+b))^{\frac{1}{2}}*\text{EllipticE}(\cot(d*x+c)-\csc(d*x+c),(-a-b)/(a+b))^{\frac{1}{2}})*b+\csc(d*x+c)^3*(1-\cos(d*x+c))^3*a-\csc(d*x+c)^3*(1-\cos(d*x+c))^3*b+a*(\csc(d*x+c)-\cot(d*x+c))+b*(\csc(d*x+c)-\cot(d*x+c))) \\ & *((\csc(d*x+c)^2*a*(1-\cos(d*x+c))^2-\csc(d*x+c)^2*b*(1-\cos(d*x+c))^2+a+b)/(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1))^{\frac{1}{2}}/(\csc(d*x+c)^2*(1-\cos(d*x+c))^2+1)/(\csc(d*x+c)^2*a*(1-\cos(d*x+c))^2-\csc(d*x+c)^2*b*(1-\cos(d*x+c))^2+a+b)/a \end{aligned}$$

3.631.5 Fracas [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

output `integral(B*sec(d*x + c)^(3/2)/sqrt(b*cos(d*x + c) + a), x)`

3.631.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**(3/2)/(a+b*cos(d*x+c))**(3/2),x)`

output `Timed out`

3.631.
$$\int \frac{(aB+bB \cos(c+dx)) \sec^{\frac{3}{2}}(c+dx)}{(a+b \cos(c+dx))^{\frac{3}{2}}} dx$$

3.631.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)
^(3/2), x)`

3.631.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba) \sec(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(3/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sec(d*x + c)^(3/2)/(b*cos(d*x + c) + a)
^(3/2), x)`

3.631.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{3/2} (Ba + Bb \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int(((1/cos(c + d*x))^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))
^(3/2),x)`

output `int(((1/cos(c + d*x))^(3/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))
^(3/2), x)`

3.631. $\int \frac{(aB + bB \cos(c + dx)) \sec^{\frac{3}{2}}(c + dx)}{(a + b \cos(c + dx))^{3/2}} dx$

$$3.632 \quad \int \frac{(aB+bB \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{3/2}} dx$$

3.632.1 Optimal result	5800
3.632.2 Mathematica [A] (verified)	5800
3.632.3 Rubi [A] (verified)	5801
3.632.4 Maple [A] (verified)	5802
3.632.5 Fracas [F]	5803
3.632.6 Sympy [F]	5803
3.632.7 Maxima [F]	5803
3.632.8 Giac [F]	5804
3.632.9 Mupad [F(-1)]	5804

3.632.1 Optimal result

Integrand size = 38, antiderivative size = 130

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2\sqrt{a + b}B\sqrt{\cos(c + dx)} \csc(c + dx) \operatorname{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\sec(c+dx)}}\right)\right)}{ad\sqrt{\sec(c + dx)}}$$

```
output 2*B*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c))/(a+b))^(1/2)*(a*(1+sec(d*x+c))/(a-b))^(1/2)/a/d/sec(d*x+c)^(1/2)
```

3.632.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.80

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \frac{2B\sqrt{\frac{a+b \cos(c+dx)}{(a+b)(1+\cos(c+dx))}} \operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right)\right), \frac{-a+b \cos(c+dx)}{a+b}}{d\sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}}\sqrt{a + b \cos(c + dx)}\sqrt{\sec(c + dx)}}$$

```
input Integrate[((a*B + b*B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^(3/2),x]
```

```
output (2*B*Sqrt[(a + b*Cos[c + d*x])/((a + b)*(1 + Cos[c + d*x]))]*EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b)]/(d*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])]*Sqrt[a + b*Cos[c + d*x]]*Sqrt[Sec[c + d*x]])
```

3.632. $\int \frac{(aB+bB \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{3/2}} dx$

3.632.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.132$, Rules used = {2011, 3042, 4710, 3042, 3295}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sec(c+dx)}(aB + bB \cos(c+dx))}{(a + b \cos(c+dx))^{3/2}} dx$$

↓ 2011

$$B \int \frac{\sqrt{\sec(c+dx)}}{\sqrt{a + b \cos(c+dx)}} dx$$

↓ 3042

$$B \int \frac{\sqrt{\csc(c+dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c+dx + \frac{\pi}{2})}} dx$$

↓ 4710

$$B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{a + b \cos(c+dx)}} dx$$

↓ 3042

$$B \sqrt{\cos(c+dx)} \sqrt{\sec(c+dx)} \int \frac{1}{\sqrt{\sin(c+dx + \frac{\pi}{2})} \sqrt{a + b \sin(c+dx + \frac{\pi}{2})}} dx$$

↓ 3295

$$\frac{2B\sqrt{a+b}\sqrt{\cos(c+dx)}\csc(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}\text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right)}{ad\sqrt{\sec(c+dx)}}$$

input `Int[((a*B + b*B*Cos[c + d*x])*Sqrt[Sec[c + d*x]])/(a + b*Cos[c + d*x])^(3/2), x]`

output `(2*Sqrt[a + b]*B*Sqrt[Cos[c + d*x]]*Csc[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b)])/(a*d*Sqrt[Sec[c + d*x]])`

3.632. $\int \frac{(aB+bB \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{3/2}} dx$

3.632.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :>
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3295 Int[1/(Sqrt[(d_)*sin[(e_) + (f_)*(x_)]*Sqrt[(a_) + (b_)*sin[(e_) + (f_)*
(x_)]]), x_Symbol] :> Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqr
t[a*((1 - Csc[e + f*x])/(a + b))]*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*Elli
pticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]
], -(a + b)/(a - b)], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]
&& PosQ[(a + b)/d]
```

```
rule 4710 Int[(csc[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] :> Simp[(c*Csc[a
+ b*x])^m*(c*Sin[a + b*x])^m Int[ActivateTrig[u]/(c*Sin[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownSineIntegrandQ[u, x]
```

3.632.4 Maple [A] (verified)

Time = 15.87 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.86

method	result	size
default	$-\frac{2B(1+\cos(dx+c))(\sqrt{\sec(dx+c)})F\left(\cot(dx+c)-\csc(dx+c),\sqrt{-\frac{a-b}{a+b}}\right)\sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}{d\sqrt{a+\cos(dx+c)}b}$	112
parts	Expression too large to display	1485

```
input int((B*a+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(3/2),x,method=
_RETURNVERBOSE)
```

```
output -2*B/d*(1+cos(d*x+c))*sec(d*x+c)^(1/2)/(a+cos(d*x+c)*b)^(1/2)*EllipticF(co
t(d*x+c)-csc(d*x+c),(-(a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c))
/(a+b))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)
```

$$3.632. \quad \int \frac{(aB+bB \cos(c+dx))\sqrt{\sec(c+dx)}}{(a+b \cos(c+dx))^{3/2}} dx$$

3.632.5 Fricas [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="fricas")`

output `integral(B*sqrt(sec(d*x + c))/sqrt(b*cos(d*x + c) + a), x)`

3.632.6 Sympy [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = B \int \frac{\sqrt{\sec(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)**(1/2)/(a+b*cos(d*x+c))**(3/2),x
)`

output `B*Integral(sqrt(sec(c + d*x))/sqrt(a + b*cos(c + d*x)), x)`

3.632.7 Maxima [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)
^(3/2), x)`

3.632.8 Giac [F]

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{(Bb \cos(dx + c) + Ba)\sqrt{\sec(dx + c)}}{(b \cos(dx + c) + a)^{3/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))*sec(d*x+c)^(1/2)/(a+b*cos(d*x+c))^(3/2),x,
algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)*sqrt(sec(d*x + c))/(b*cos(d*x + c) + a)
^(3/2), x)`

3.632.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(aB + bB \cos(c + dx))\sqrt{\sec(c + dx)}}{(a + b \cos(c + dx))^{3/2}} dx = \int \frac{\sqrt{\frac{1}{\cos(c+dx)}} (Ba + Bb \cos(c + dx))}{(a + b \cos(c + dx))^{3/2}} dx$$

input `int(((1/cos(c + d*x))^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))
^(3/2),x)`

output `int(((1/cos(c + d*x))^(1/2)*(B*a + B*b*cos(c + d*x)))/(a + b*cos(c + d*x))
^(3/2), x)`

3.633 $\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

3.633.1 Optimal result	5805
3.633.2 Mathematica [A] (verified)	5805
3.633.3 Rubi [A] (verified)	5806
3.633.4 Maple [A] (verified)	5808
3.633.5 Fricas [F]	5808
3.633.6 Sympy [F]	5808
3.633.7 Maxima [F]	5809
3.633.8 Giac [F]	5809
3.633.9 Mupad [F(-1)]	5809

3.633.1 Optimal result

Integrand size = 38, antiderivative size = 137

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \frac{2\sqrt{a + b}B \sqrt{\cos(c + dx)} \csc(c + dx) \operatorname{EllipticPi}\left(\frac{a+b}{b}, \arcsin\left(\frac{\sqrt{a+b} \cos(c+dx)}{\sqrt{a+b} \sqrt{\cos(c+dx)}}\right), -\frac{a+b}{a-b}\right) \sqrt{\frac{a(1-\sec(c+dx))}{a+b}} \sqrt{\frac{a(1+\sec(c+dx))}{a+b}}}{bd \sqrt{\sec(c + dx)}}$$

output

```
-2*B*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2), (a+b)/b, ((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b)^(1/2)/b/d/sec(d*x+c)^(1/2)
```

3.633.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.07

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \frac{2B \sqrt{\frac{\cos(c+dx)}{1+\cos(c+dx)}} \sqrt{\frac{a+b \cos(c+dx)}{(a+b)(1+\cos(c+dx))}} \left(\operatorname{EllipticF}\left(\arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right), \frac{-a+b}{a+b}\right) - 2 \operatorname{EllipticPi}\left(-1, \arcsin\left(\tan\left(\frac{1}{2}(c + dx)\right)\right)\right) \right)}{d \sqrt{\frac{1}{1+\cos(c+dx)}} \sqrt{a + b \cos(c + dx)}}$$

input

```
Integrate[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sqrt[Sec[c + d*x]]), x]
```

3.633. $\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

```
output (-2*B*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])] * Sqrt[(a + b * Cos[c + d*x]) / ((a + b) * (1 + Cos[c + d*x]))] * (EllipticF[ArcSin[Tan[(c + d*x)/2]], (-a + b) / (a + b)] - 2 * EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b) / (a + b)]) * Sqrt[1 + Sec[c + d*x]]) / (d * Sqrt[(1 + Cos[c + d*x])^(-1)] * Sqrt[a + b * Cos[c + d * x]])
```

3.633.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.132$, Rules used = {2011, 3042, 4710, 3042, 3288}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{\sqrt{\sec(c + dx)}(a + b \cos(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sqrt{\csc(c + dx + \frac{\pi}{2})} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4710} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3288} \\
 & \frac{2B \sqrt{a + b} \sqrt{\cos(c + dx)} \csc(c + dx) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(\sec(c + dx) + 1)}{a - b}} \text{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b} \cos(c + dx)}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right)\right) - \dots}{bd \sqrt{\sec(c + dx)}}
 \end{aligned}$$

3.633. $\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx$

input `Int[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sqrt[Sec[c + d*x]]),x]`

output `(-2*Sqrt[a + b]*B*Sqrt[Cos[c + d*x]]*Csc[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -(a + b)/(a - b)]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b*d*Sqrt[Sec[c + d*x]])`

3.633.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 4710 `Int[(csc[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Csc[a + b*x])^m*(c*Sin[a + b*x])^m Int[ActivateTrig[u]/(c*Sin[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownSineIntegrandQ[u, x]`

3.633.4 Maple [A] (verified)

Time = 15.26 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.01

method	result	size
default	$\frac{2B \left(F \left(\cot(dx+c) - \csc(dx+c), \sqrt{-\frac{a-b}{a+b}} \right) - 2\Pi \left(\cot(dx+c) - \csc(dx+c), -1, \sqrt{-\frac{a-b}{a+b}} \right) \right) \sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}}}{d\sqrt{a+\cos(dx+c)}b\sqrt{\sec(dx+c)}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}}$	138
parts	Expression too large to display	1821

```
input int((B*a+b*B*cos(d*x+c))/(a*cos(d*x+c)*b)^(3/2)/sec(d*x+c)^(1/2),x,method=
_RETURNVERBOSE)
```

```
output 2*B/d/(a*cos(d*x+c)*b)^(1/2)*(EllipticF(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b)
)^(1/2))-2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b)^(1/2))*((a
+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)/sec(d*x+c)^(1/2)/(cos(d*x+c)/(1
+cos(d*x+c)))^(1/2)
```

3.633.5 Fracas [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)}} dx$$

```
input integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x,
algorithm="fricas")
```

```
output integral(B/(sqrt(b*cos(d*x + c) + a)*sqrt(sec(d*x + c))), x)
```

3.633.6 Sympy [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = B \int \frac{1}{\sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} dx$$

```
input integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2)/sec(d*x+c)**(1/2),x
)
```

```
output B*Integral(1/(sqrt(a + b*cos(c + d*x))*sqrt(sec(c + d*x))), x)
```

3.633. $\int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sqrt{\sec(c+dx)}} dx$

3.633.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x,
algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*
x + c))), x)`

3.633.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sqrt{\sec(dx + c)}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x,
algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*sqrt(sec(d*
x + c))), x)`

3.633.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx = \int \frac{Ba + Bb \cos(c + dx)}{\sqrt{\frac{1}{\cos(c+dx)}} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(
3/2)),x)`

output `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(1/2)*(a + b*cos(c + d*x))^(
3/2)), x)`

3.633. $\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sqrt{\sec(c + dx)}} dx$

3.634
$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx$$

3.634.1 Optimal result	5810
3.634.2 Mathematica [A] (warning: unable to verify)	5811
3.634.3 Rubi [A] (verified)	5812
3.634.4 Maple [A] (verified)	5817
3.634.5 Fracas [F]	5818
3.634.6 Sympy [F(-1)]	5819
3.634.7 Maxima [F]	5819
3.634.8 Giac [F]	5819
3.634.9 Mupad [F(-1)]	5820

3.634.1 Optimal result

Integrand size = 38, antiderivative size = 479

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{(a - b)\sqrt{a + b} \sqrt{\cos(c + dx)} \csc(c + dx) E\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right) \mid -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{abd \sqrt{\sec(c + dx)}} +$$

$$\frac{\sqrt{a + b} B \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticF}\left(\arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{bd \sqrt{\sec(c + dx)}} +$$

$$\frac{a\sqrt{a + b} B \sqrt{\cos(c + dx)} \csc(c + dx) \text{EllipticPi}\left(\frac{a + b}{b}, \arcsin\left(\frac{\sqrt{a + b \cos(c + dx)}}{\sqrt{a + b} \sqrt{\cos(c + dx)}}\right), -\frac{a + b}{a - b}\right) \sqrt{\frac{a(1 - \sec(c + dx))}{a + b}} \sqrt{\frac{a(1 + \sec(c + dx))}{a - b}}}{b^2 d \sqrt{\sec(c + dx)}} +$$

$$\frac{B \sin(c + dx)}{d \sqrt{a + b \cos(c + dx)} \sqrt{\sec(c + dx)}} + \frac{aB \sqrt{\sec(c + dx)} \sin(c + dx)}{bd \sqrt{a + b \cos(c + dx)}}$$

3.634.
$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx$$

output `B*sin(d*x+c)/d/(a+b*cos(d*x+c))^(1/2)/sec(d*x+c)^(1/2)+a*B*sin(d*x+c)*sec(d*x+c)^(1/2)/b/d/(a+b*cos(d*x+c))^(1/2)-(a-b)*B*csc(d*x+c)*EllipticE((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b))^(1/2)/a/b/d/sec(d*x+c)^(1/2)+B*csc(d*x+c)*EllipticF((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b))^(1/2)/b/d/sec(d*x+c)^(1/2)+a*B*csc(d*x+c)*EllipticPi((a+b*cos(d*x+c))^(1/2)/(a+b)^(1/2)/cos(d*x+c)^(1/2),(a+b)/b,((-a-b)/(a-b))^(1/2))*(a+b)^(1/2)*cos(d*x+c)^(1/2)*(a*(1-sec(d*x+c)))/(a+b)^(1/2)*(a*(1+sec(d*x+c)))/(a-b))^(1/2)/b^2/d/sec(d*x+c)^(1/2)`

3.634.2 Mathematica [A] (warning: unable to verify)

Time = 0.98 (sec) , antiderivative size = 236, normalized size of antiderivative = 0.49

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \frac{2B \cos(c + dx) \sqrt{\frac{\cos(c+dx)}{(1+\cos(c+dx))^2}} (\cos^2(\frac{1}{2}(c + dx)) \sec(c + dx))^{3/2}}{}$$

input `Integrate[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^(3/2)),x]`

output `(2*B*Cos[c + d*x]*Sqrt[Cos[c + d*x]/(1 + Cos[c + d*x])^2]*(Cos[(c + d*x)/2]^2*Sec[c + d*x])^(3/2)*((a + b)*EllipticE[ArcSin[Tan[(c + d*x)/2]]], (-a + b)/(a + b))*Sqrt[((a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2)/(a + b)] - 2*a*EllipticPi[-1, ArcSin[Tan[(c + d*x)/2]], (-a + b)/(a + b))*Sqrt[((a + b*Cos[c + d*x])*Sec[(c + d*x)/2]^2)/(a + b)] + (a + b*Cos[c + d*x])*Sqrt[Cos[c + d*x]*Sec[(c + d*x)/2]^2*Tan[(c + d*x)/2]]/(b*d*Sqrt[a + b*Cos[c + d*x]])]`

3.634.3 Rubi [A] (verified)

Time = 1.86 (sec) , antiderivative size = 446, normalized size of antiderivative = 0.93, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {2011, 3042, 4710, 3042, 3299, 3042, 3288, 3482, 27, 3042, 3472, 25, 27, 3042, 3280, 3042, 3295, 3473}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \cos(c + dx)}{\sec^{\frac{3}{2}}(c + dx)(a + b \cos(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sqrt{a + b \cos(c + dx)} \sec^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\csc(c + dx + \frac{\pi}{2})^{3/2} \sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4710} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\cos^{\frac{3}{2}}(c + dx)}{\sqrt{a + b \cos(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \int \frac{\sin(c + dx + \frac{\pi}{2})^{3/2}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3299} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\cos(c + dx)}(a + 2b \cos(c + dx))}{\sqrt{a + b \cos(c + dx)}} dx}{2b} - \frac{a \int \frac{\sqrt{\cos(c + dx)}}{\sqrt{a + b \cos(c + dx)}} dx}{2b} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\cos(c + dx)} \sqrt{\sec(c + dx)} \left(\frac{\int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}(a + 2b \sin(c + dx + \frac{\pi}{2}))}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2b} - \frac{a \int \frac{\sqrt{\sin(c + dx + \frac{\pi}{2})}}{\sqrt{a + b \sin(c + dx + \frac{\pi}{2})}} dx}{2b} \right) \\
 & \quad \downarrow \text{3288}
 \end{aligned}$$

3.634. $\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx$

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+2b\sin(c+dx+\frac{\pi}{2}))}{\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}} dx}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{2b} \right)$$

↓ 3482

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\frac{1}{2} \int \frac{2(\cos(c+dx)a^2+ba)}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{2b} \right)$$

↓ 27

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\cos(c+dx)a^2+ba}{\sqrt{\cos(c+dx)}(a+b\cos(c+dx))^{3/2}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{2b} \right)$$

↓ 3042

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int \frac{\sin(c+dx+\frac{\pi}{2})a^2+ba}{\sqrt{\sin(c+dx+\frac{\pi}{2})}(a+b\sin(c+dx+\frac{\pi}{2}))^{3/2}} dx + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{2b} \right)$$

↓ 3472

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(\frac{\int -\frac{a(a^2-b^2)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{2b} \right)$$

↓ 25

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)} \left(-\frac{\int \frac{a(a^2-b^2)}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}} dx}{a^2-b^2} + \frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}} + \frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b} + \frac{a\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx))}{a+b}}}{2b} \right)$$

↓ 27

3.634. $\int \frac{aB+bB\cos(c+dx)}{(a+b\cos(c+dx))^{3/2}\sec^{\frac{3}{2}}(c+dx)} dx$

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{-a\int\frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx+\frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}+\frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b}\right)$$

↓ 3042

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{-a\int\frac{1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx+\frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}+\frac{2a\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}}{2b}\right)$$

↓ 3280

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{-a\left(\int\frac{\cos(c+dx)+1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{a+b\cos(c+dx)}}dx-\int\frac{1}{\sqrt{\cos(c+dx)}\sqrt{a+b\cos(c+dx)}}dx\right)+\frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b}\right)$$

↓ 3042

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{-a\left(\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\int\frac{1}{\sqrt{\sin(c+dx+\frac{\pi}{2})}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx\right)+\frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b}\right)$$

↓ 3295

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{-a\left(\int\frac{\sin(c+dx+\frac{\pi}{2})+1}{\sin(c+dx+\frac{\pi}{2})^{3/2}\sqrt{a+b\sin(c+dx+\frac{\pi}{2})}}dx-\frac{2\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}}{a^2d}\right)+\frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b}\right)$$

↓ 3473

$$B\sqrt{\cos(c+dx)}\sqrt{\sec(c+dx)}\left(\frac{-a\left(\frac{2(a-b)\sqrt{a+b}\cot(c+dx)\sqrt{\frac{a(1-\sec(c+dx))}{a+b}}\sqrt{\frac{a(\sec(c+dx)+1)}{a-b}}E\left(\arcsin\left(\frac{\sqrt{a+b}\cos(c+dx)}{\sqrt{a+b}\sqrt{\cos(c+dx)}}\right)\right)-\frac{a+b}{a-b}}{a^2d}\right)+\frac{2b\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{a+b\cos(c+dx)}}}{2b}\right)$$

input `Int[(a*B + b*B*Cos[c + d*x])/((a + b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^(3/2)),x]`

output `B*Sqrt[Cos[c + d*x]]*Sqrt[Sec[c + d*x]]*((a*Sqrt[a + b]*Cot[c + d*x]*EllipticPi[(a + b)/b, ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(b^2*d) + (-a*((2*(a - b)*Sqrt[a + b]*Cot[c + d*x]*EllipticE[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a^2*d) - (2*Sqrt[a + b]*Cot[c + d*x]*EllipticF[ArcSin[Sqrt[a + b*Cos[c + d*x]]/(Sqrt[a + b]*Sqrt[Cos[c + d*x]])], -((a + b)/(a - b))]*Sqrt[(a*(1 - Sec[c + d*x]))/(a + b)]*Sqrt[(a*(1 + Sec[c + d*x]))/(a - b))]/(a*d))) + (2*a*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]]*Sqrt[a + b*Cos[c + d*x]]) + (2*b*Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[a + b*Cos[c + d*x]])/(2*b))`

3.634.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3280 `Int[1/(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)*Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[1/(a - b) Int[1/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]]), x], x] - Simp[b/(a - b) Int[(1 + Sin[e + f*x])/((a + b*Sin[e + f*x])^(3/2)*Sqrt[c + d*Sin[e + f*x]]), x], x] / ; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

rule 3288 `Int[Sqrt[(b_.)*sin[(e_.) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*b*(Tan[e + f*x]/(d*f))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x])/(c - d))*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticPi[(c + d)/d, ArcSin[Sqrt[c + d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)], x] / ; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && PosQ[(c + d)/b]`

rule 3295 `Int[1/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Simp[-2*(Tan[e + f*x]/(a*f))*Rt[(a + b)/d, 2]*Sqrt[a*((1 - Csc[e + f*x])/(a + b))*Sqrt[a*((1 + Csc[e + f*x])/(a - b))]*EllipticF[ArcSin[Sqrt[a + b*Sin[e + f*x]]/Sqrt[d*Sin[e + f*x]]/Rt[(a + b)/d, 2]], -(a + b)/(a - b)], x] / ; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && PosQ[(a + b)/d]`

rule 3299 `Int[((d_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)/Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(-a)*(d/(2*b)) Int[Sqrt[d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/(2*b) Int[Sqrt[d*Sin[e + f*x]]*((a + 2*b*Sin[e + f*x])/Sqrt[a + b*Sin[e + f*x]]), x], x] / ; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0]`

rule 3472 `Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])/(Sqrt[(d_.)*sin[(e_.) + (f_.)*(x_)]]*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(3/2)), x_Symbol] := Simp[2*(A*b - a*B)*(Cos[e + f*x]/(f*(a^2 - b^2)*Sqrt[a + b*Sin[e + f*x]]*Sqrt[d*Sin[e + f*x]]), x] + Simp[d/(a^2 - b^2) Int[(A*b - a*B + (a*A - b*B)*Sin[e + f*x])/(Sqrt[a + b*Sin[e + f*x]]*(d*Sin[e + f*x])^(3/2)), x], x] / ; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[a^2 - b^2, 0]`

```
rule 3473 Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(((b_)*sin[(e_) + (f_)*(x_)]
^(3/2)*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] := Simp[-2*A*
(c - d)*(Tan[e + f*x]/(f*b*c^2))*Rt[(c + d)/b, 2]*Sqrt[c*((1 + Csc[e + f*x]
)/(c - d))]*Sqrt[c*((1 - Csc[e + f*x])/(c + d))]*EllipticE[ArcSin[Sqrt[c +
d*Sin[e + f*x]]/Sqrt[b*Sin[e + f*x]]/Rt[(c + d)/b, 2]], -(c + d)/(c - d)],
x] /; FreeQ[{b, c, d, e, f, A, B}, x] && NeQ[c^2 - d^2, 0] && EqQ[A, B] &&
PosQ[(c + d)/b]
```

```
rule 3482 Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[-2*B*Cos[e + f*x]*Sqrt[a + b*Sin[e + f*x]]*((c + d*Sin[e + f*x])^n/(f*(2*
n + 3))), x] + Simp[1/(2*n + 3) Int[((c + d*Sin[e + f*x])^(n - 1)/Sqrt[a
+ b*Sin[e + f*x]])*Simp[a*A*c*(2*n + 3) + B*(b*c + 2*a*d*n) + (B*(a*c + b*d
)*(2*n + 1) + A*(b*c + a*d)*(2*n + 3))*Sin[e + f*x] + (A*b*d*(2*n + 3) + B*
(a*d + 2*b*c*n))*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A,
B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Eq
Q[n^2, 1/4]
```

```
rule 4710 Int[(csc[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Csc[a
+ b*x])^m*(c*Sin[a + b*x])^m Int[ActivateTrig[u]/(c*Sin[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownSineIntegrandQ[u, x]
```

3.634.4 Maple [A] (verified)

Time = 16.90 (sec) , antiderivative size = 820, normalized size of antiderivative = 1.71

method	result
default	$-\frac{B \left(\sec^2(dx+c) E \left(\cot(dx+c) - \csc(dx+c), \sqrt{-\frac{a-b}{a+b}} \right) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sqrt{\frac{a+\cos(dx+c)b}{(1+\cos(dx+c))(a+b)}} \right)}{a + (\sec^2(dx+c) E \left(\cot(dx+c) - \csc(dx+c) \right))}$
parts	Expression too large to display

```
input int((B*a+b*B*cos(d*x+c))/(a*cos(d*x+c)*b)^(3/2)/sec(d*x+c)^(3/2),x,method=
_RETURNVERBOSE)
```

$$3.634. \int \frac{aB+bB \cos(c+dx)}{(a+b \cos(c+dx))^{3/2} \sec^2(c+dx)} dx$$

output

```

-B/d/(1+cos(d*x+c))/(a+cos(d*x+c)*b)^(1/2)/sec(d*x+c)^(3/2)*(sec(d*x+c)^2*
EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(d
*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+sec(d*x+c)^2
*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-2*sec(d*x+c
)^2*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*(cos(d*x+c)/
(1+cos(d*x+c)))^(1/2)*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+2*se
c(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c)
,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+2*s
ec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c)
,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-4*
sec(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x
+c),-1,(-a-b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)
*a+(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-
b)/(a+b))^(1/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a+(cos(d*x+
c)/(1+cos(d*x+c)))^(1/2)*EllipticE(cot(d*x+c)-csc(d*x+c),(-a-b)/(a+b))^(1
/2))*((a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*b-2*(cos(d*x+c)/(1+cos(
d*x+c)))^(1/2)*EllipticPi(cot(d*x+c)-csc(d*x+c),-1,(-a-b)/(a+b))^(1/2))*((
a+cos(d*x+c)*b)/(1+cos(d*x+c)))/(a+b)^(1/2)*a-tan(d*x+c)*a-b*sin(d*x+c))/
b

```

3.634.5 Fracas [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{\frac{3}{2}} \sec^{\frac{3}{2}}(dx + c)} dx$$

input

```

integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x,
algorithm="fricas")

```

output

```

integral(B/(sqrt(b*cos(d*x + c) + a)*sec(d*x + c)^(3/2)), x)

```

3.634.6 Sympy [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))**(3/2)/sec(d*x+c)**(3/2), x)`

output `Timed out`

3.634.7 Maxima [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2), x, algorithm="maxima")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2)), x)`

3.634.8 Giac [F]

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{3/2}(c + dx)} dx = \int \frac{Bb \cos(dx + c) + Ba}{(b \cos(dx + c) + a)^{3/2} \sec(dx + c)^{3/2}} dx$$

input `integrate((a*B+b*B*cos(d*x+c))/(a+b*cos(d*x+c))^(3/2)/sec(d*x+c)^(3/2), x, algorithm="giac")`

output `integrate((B*b*cos(d*x + c) + B*a)/((b*cos(d*x + c) + a)^(3/2)*sec(d*x + c)^(3/2)), x)`

3.634.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \cos(c + dx)}{(a + b \cos(c + dx))^{3/2} \sec^{\frac{3}{2}}(c + dx)} dx = \int \frac{Ba + Bb \cos(c + dx)}{\left(\frac{1}{\cos(c + dx)}\right)^{3/2} (a + b \cos(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(3/2)),x)`

output `int((B*a + B*b*cos(c + d*x))/((1/cos(c + d*x))^(3/2)*(a + b*cos(c + d*x))^(3/2)), x)`

3.635 $\int (a+b \cos(e+fx))^n (A+B \cos(e+fx))(c \sec(e+fx))^m dx$

3.635.1 Optimal result	5821
3.635.2 Mathematica [N/A]	5821
3.635.3 Rubi [N/A]	5822
3.635.4 Maple [N/A] (verified)	5823
3.635.5 Fracas [N/A]	5824
3.635.6 Sympy [N/A]	5824
3.635.7 Maxima [N/A]	5825
3.635.8 Giac [N/A]	5825
3.635.9 Mupad [N/A]	5826

3.635.1 Optimal result

Integrand size = 33, antiderivative size = 33

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= (c \cos(e + fx))^m (c \sec(e + fx))^m \text{Int}((c \cos(e + fx))^{-m} (a + b \cos(e + fx))^n (A + B \cos(e + fx)), x)$$

output `(c*cos(f*x+e))^m*(c*sec(f*x+e))^m*Unintegrable((a+b*cos(f*x+e))^n*(A+B*cos(f*x+e))/((c*cos(f*x+e))^m),x)`

3.635.2 Mathematica [N/A]

Not integrable

Time = 20.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (a + b \cos(e + fx))^n (A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

input `Integrate[(a + b*Cos[e + f*x])^n*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `Integrate[(a + b*Cos[e + f*x])^n*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m, x]`

3.635.3 Rubi [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3440, 3042, 3486}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (A + B \cos(e + fx))(c \sec(e + fx))^m (a + b \cos(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right) \right) \left(c \csc\left(e + fx + \frac{\pi}{2}\right) \right)^m \left(a + b \sin\left(e + fx + \frac{\pi}{2}\right) \right)^n dx$$

$$\downarrow \text{3440}$$

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int (c \cos(e + fx))^{-m} (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int \left(c \sin\left(e + fx + \frac{\pi}{2}\right) \right)^{-m} \left(a + b \sin\left(e + fx + \frac{\pi}{2}\right) \right)^n \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right) \right) dx$$

$$\downarrow \text{3486}$$

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int (c \cos(e + fx))^{-m} (a + b \cos(e + fx))^n (A + B \cos(e + fx)) dx$$

input `Int[(a + b*Cos[e + f*x])^n*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `$Aborted`

3.635.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3486 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(A + B*Sin[e + f*x])*(c + d*Sin[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.635.4 Maple [N/A] (verified)

Not integrable

Time = 1.88 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int (a + b \cos(fx + e))^n (A + \cos(fx + e) B) (c \sec(fx + e))^m dx$$

input `int((a+b*cos(f*x+e))^n*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

output `int((a+b*cos(f*x+e))^n*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

3.635.5 Fricas [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^n (c \sec(fx + e))^m dx$$

```
input integrate((a+b*cos(f*x+e))^n*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm
m="fricas")
```

```
output integral((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^n*(c*sec(f*x + e))^m, x
)
```

3.635.6 Sympy [N/A]

Not integrable

Time = 165.51 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (c \sec(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^n dx$$

```
input integrate((a+b*cos(f*x+e))^n*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x)
```

```
output Integral((c*sec(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^n,
x)
```

3.635.7 Maxima [N/A]

Not integrable

Time = 5.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^n (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^n*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^n*(c*sec(f*x + e))^m, x)`

3.635.8 Giac [N/A]

Not integrable

Time = 0.66 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.06

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^n (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^n*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^n*(c*sec(f*x + e))^m, x)`

3.635.9 Mupad [N/A]

Not integrable

Time = 2.85 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.12

$$\int (a + b \cos(e + fx))^n (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^n dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^n,x)`output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^n, x)`

3.636 $\int (a+b \cos(e+fx))^4(A+B \cos(e+fx))(c \sec(e+fx))^m dx$

3.636.1 Optimal result	5827
3.636.2 Mathematica [A] (verified)	5828
3.636.3 Rubi [A] (verified)	5829
3.636.4 Maple [F]	5835
3.636.5 Fricas [F]	5835
3.636.6 Sympy [F(-1)]	5836
3.636.7 Maxima [F]	5836
3.636.8 Giac [F]	5836
3.636.9 Mupad [F(-1)]	5837

3.636.1 Optimal result

Integrand size = 33, antiderivative size = 644

$$\int (a + b \cos(e + fx))^4(A + B \cos(e + fx))(c \sec(e + fx))^m dx =$$

$$\frac{c^6(4a^3Ab(15 - 8m + m^2) + a^4B(15 - 8m + m^2) + 4aAb^3(10 - 7m + m^2) + 6a^2b^2B(10 - 7m + m^2) + f(2 - m)(4 - m)(6 - m))}{f(2 - m)(4 - m)(6 - m)}$$

$$\frac{c^5(a^4A(8 - 6m + m^2) + 6a^2Ab^2(4 - 5m + m^2) + 4a^3bB(4 - 5m + m^2) + Ab^4(3 - 4m + m^2) + 4ab^3B(3 - 4m + m^2))}{f(1 - m)(3 - m)(5 - m)}$$

$$\frac{ac^5(4a^2Ab(3 - 4m + m^2) + a^3B(3 - 4m + m^2) + 2Ab^3(4 - 2m + m^2) + ab^2B(8 - 13m + 5m^2))(c \sec(e + fx))^m}{f(1 - m)(2 - m)(4 - m)}$$

$$\frac{a^2c^5(2abB(1 - m)^2 + a^2A(2 - m)^2 + Ab^2(6 - m + m^2)) \sec(e + fx)(c \sec(e + fx))^{-5+m} \tan(e + fx)}{f(1 - m)(2 - m)(3 - m)}$$

$$\frac{ac^5(aB(1 - m) - Ab(2 + m))(c \sec(e + fx))^{-5+m}(b + a \sec(e + fx))^2 \tan(e + fx)}{f(1 - m)(2 - m)}$$

$$\frac{aAc^5(c \sec(e + fx))^{-5+m}(b + a \sec(e + fx))^3 \tan(e + fx)}{f(1 - m)}$$

output
$$-c^6(4a^3Ab(m^2-8m+15)+a^4B(m^2-8m+15)+4aAb^3(m^2-7m+10)+6a^2b^2B(m^2-7m+10)+b^4B(m^2-6m+8))*\text{hypergeom}([1/2, 3-1/2m], [4-1/2m], \cos(fx+e)^2*(c*\sec(fx+e))^{(-6+m)}*\sin(fx+e)/f/(-m^3+12m^2-44m+48)/(\sin(fx+e)^2)^{(1/2)}-c^5(a^4A(m^2-6m+8)+6a^2Ab^2(m^2-5m+4)+4a^3bB(m^2-5m+4)+Ab^4(m^2-4m+3)+4aAb^3B(m^2-4m+3))*\text{hypergeom}([1/2, 5/2-1/2m], [7/2-1/2m], \cos(fx+e)^2*(c*\sec(fx+e))^{(-5+m)}*\sin(fx+e)/f/(1-m)/(m^2-8m+15)/(\sin(fx+e)^2)^{(1/2)}-ac^5(4a^2Ab(m^2-4m+3)+a^3B(m^2-4m+3)+2Ab^3(m^2-2m+4)+Ab^2B(5m^2-13m+8))*(c*\sec(fx+e))^{(-5+m)}*\tan(fx+e)/f/(1-m)/(m^2-6m+8)-a^2c^5(2AbB(1-m)^2+a^2A(2-m)^2+Ab^2(m^2-m+6))*\sec(fx+e)*(c*\sec(fx+e))^{(-5+m)}*\tan(fx+e)/f/(-m^3+6m^2-11m+6)-ac^5(Ab(1-m)-Ab(2+m))*(c*\sec(fx+e))^{(-5+m)}*(b+a*\sec(fx+e))^2*\tan(fx+e)/f/(m^2-3m+2)-aAc^5(c*\sec(fx+e))^{(-5+m)}*(b+a*\sec(fx+e))^3*\tan(fx+e)/f/(1-m)$$

3.636.2 Mathematica [A] (verified)

Time = 2.66 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.49

$$\int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \frac{\cot(e + fx) \left(\frac{b^4 B \cos^5(e + fx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-5+m), \frac{1}{2}(-3+m), \sec^2(e + fx)\right)}{-5+m} + \frac{b^3 (Ab + 4aB) \cos^4(e + fx) \text{Hypergeometric2F1}}{-4+m} \right)}{1}$$

input `Integrate[(a + b*Cos[e + f*x])^4*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output
$$(\text{Cot}[e + f*x]*((b^4*B*\text{Cos}[e + f*x]^5*\text{Hypergeometric2F1}[1/2, (-5 + m)/2, (-3 + m)/2, \text{Sec}[e + f*x]^2])/(-5 + m) + (b^3*(A*b + 4*a*B)*\text{Cos}[e + f*x]^4*\text{Hypergeometric2F1}[1/2, (-4 + m)/2, (-2 + m)/2, \text{Sec}[e + f*x]^2])/(-4 + m) + a*((2*b^2*(2*A*b + 3*a*B)*\text{Cos}[e + f*x]^3*\text{Hypergeometric2F1}[1/2, (-3 + m)/2, (-1 + m)/2, \text{Sec}[e + f*x]^2])/(-3 + m) + a*((2*b*(3*A*b + 2*a*B)*\text{Cos}[e + f*x]^2*\text{Hypergeometric2F1}[1/2, (-2 + m)/2, m/2, \text{Sec}[e + f*x]^2])/(-2 + m) + a*(((4*A*b + a*B)*\text{Cos}[e + f*x]*\text{Hypergeometric2F1}[1/2, (-1 + m)/2, (1 + m)/2, \text{Sec}[e + f*x]^2])/(-1 + m) + (a*A*\text{Hypergeometric2F1}[1/2, m/2, (2 + m)/2, \text{Sec}[e + f*x]^2])/m)))*c*\text{Sec}[e + f*x]^m*\text{Sqrt}[-\text{Tan}[e + f*x]^2])/f$$

3.636.3 Rubi [A] (verified)

Time = 4.27 (sec) , antiderivative size = 617, normalized size of antiderivative = 0.96, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 3439, 3042, 4514, 25, 3042, 4584, 3042, 4564, 25, 3042, 4535, 3042, 4259, 3042, 3122, 4534, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^4 \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^m dx$$

$$\downarrow \text{3439}$$

$$c^5 \int (c \sec(e + fx))^{m-5} (b + a \sec(e + fx))^4 (B + A \sec(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$c^5 \int \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^{m-5} \left(b + a \csc \left(e + fx + \frac{\pi}{2} \right) \right)^4 \left(B + A \csc \left(e + fx + \frac{\pi}{2} \right) \right) dx$$

$$\downarrow \text{4514}$$

$$c^5 \left(- \frac{\int -(c \sec(e + fx))^{m-5} (b + a \sec(e + fx))^2 (a(aB(1 - m) - Ab(m + 2)) \sec^2(e + fx) + (A(2 - m)a^2 + b(A(2 - m) + Ab))) dx}{1 - m} \right)$$

$$\downarrow \text{25}$$

$$c^5 \left(\frac{\int (c \sec(e + fx))^{m-5} (b + a \sec(e + fx))^2 (a(aB(1 - m) - Ab(m + 2)) \sec^2(e + fx) + (A(2 - m)a^2 + b(Ab + A(2 - m)))) dx}{1 - m} \right)$$

$$\downarrow \text{3042}$$

$$c^5 \left(\frac{\int (c \csc \left(e + fx + \frac{\pi}{2} \right))^{m-5} (b + a \csc \left(e + fx + \frac{\pi}{2} \right))^2 \left(a(aB(1 - m) - Ab(m + 2)) \csc \left(e + fx + \frac{\pi}{2} \right)^2 + (A(2 - m)a^2 + b(A(2 - m) + Ab)) \right) dx}{1 - m} \right)$$

$$\downarrow \text{4584}$$

$$c^5 \left(\frac{-\int (c \sec(e+fx))^{m-5} (b+a \sec(e+fx)) (-a(A(m^2-m+6)b^2+2aB(1-m)^2b+a^2A(2-m)^2) \sec^2(e+fx) - (B(m^2-4m+3)a^3+Ab(3m^2-12m-2-m))}{2-m} \right)$$

↓ 3042

$$c^5 \left(\frac{-\int (c \csc(e+fx+\frac{\pi}{2}))^{m-5} (b+a \csc(e+fx+\frac{\pi}{2})) (-a(A(m^2-m+6)b^2+2aB(1-m)^2b+a^2A(2-m)^2) \csc(e+fx+\frac{\pi}{2})^2 + (-B(m^2-4m+3)a^3+Ab(3m^2-12m-2-m))}{2-m} \right)$$

↓ 4564

$$c^5 \left(\frac{a^2 \tan(e+fx) \sec(e+fx) (a^2 A(2-m)^2 + 2abB(1-m)^2 + Ab^2(m^2-m+6)) (c \sec(e+fx))^{m-5} - \int (c \sec(e+fx))^{m-5} ((3-m)(-B(m^2-6m+5)a^2+2Ab(5-m))}{f(3-m)} \right)$$

↓ 25

$$c^5 \left(\frac{\int (c \sec(e+fx))^{m-5} ((3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))b^2-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m-2-m)))}{f(3-m)} \right)$$

↓ 3042

$$c^5 \left(\frac{\int (c \csc(e+fx+\frac{\pi}{2}))^{m-5} ((3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))b^2-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m-2-m)))}{f(3-m)} \right)$$

↓ 4535

$$c^5 \left(\frac{\int (c \sec(e+fx))^{m-5} (b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m-2-m)))}{f(3-m)} \right)$$

↓ 3042

$$c^5 \left(\frac{f(c \csc(e+fx+\frac{\pi}{2}))^{m-5} (b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))}{(b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))} \right)$$

↓ 4259

$$c^5 \left(\frac{f(c \csc(e+fx+\frac{\pi}{2}))^{m-5} (b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))}{(b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))} \right)$$

↓ 3042

$$c^5 \left(\frac{f(c \csc(e+fx+\frac{\pi}{2}))^{m-5} (b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))}{(b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))} \right)$$

↓ 3122

$$c^5 \left(\frac{f(c \csc(e+fx+\frac{\pi}{2}))^{m-5} (b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))}{(b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))} \right)$$

↓ 4534

$$c^5 \left(\frac{-(m^2-4m+3)(a^4B(m^2-8m+15)+4a^3Ab(m^2-8m+15)+6a^2b^2B(m^2-7m+10)+4aAb^3(m^2-7m+10)+b^4B(m^2-6m+8))f(c \sec(e+fx))^{m-5} dx}{(b^2(3-m)(-B(m^2-6m+5)a^2+2Ab(5-m)ma-b^2B(m^2-3m+2))-a(3-m)(B(m^2-4m+3)a^3+4Ab(m^2-4m+3)a^2+b^2B(5m^2-12m+8)ab-b^3B(m^2-3m+2)))} \right)$$

↓ 3042

$$c^5 \left(\frac{(m^2-4m+3)(a^4B(m^2-8m+15)+4a^3Ab(m^2-8m+15)+6a^2b^2B(m^2-7m+10)+4aAb^3(m^2-7m+10)+b^4B(m^2-6m+8))}{4-m} f(c \csc(e+fx+\frac{\pi}{2}))^{m-5} \right)$$

↓ 4259

$$c^5 \left(\frac{(m^2-4m+3)(a^4B(m^2-8m+15)+4a^3Ab(m^2-8m+15)+6a^2b^2B(m^2-7m+10)+4aAb^3(m^2-7m+10)+b^4B(m^2-6m+8))}{4-m} \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^{m-5} \right)$$

↓ 3042

$$c^5 \left(\frac{(m^2-4m+3)(a^4B(m^2-8m+15)+4a^3Ab(m^2-8m+15)+6a^2b^2B(m^2-7m+10)+4aAb^3(m^2-7m+10)+b^4B(m^2-6m+8))}{4-m} \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^{m-5} \right)$$

↓ 3122

$$c^5 \left(\frac{a^2 \tan(e+fx) \sec(e+fx) (a^2 A(2-m)^2 + 2abB(1-m)^2 + Ab^2(m^2-m+6)) (c \sec(e+fx))^{m-5}}{f(3-m)} + \frac{a(3-m) \tan(e+fx) (a^3 B(m^2-4m+3) + 4a^2 Ab(m^2-4m+3))}{f(3-m)} \right)$$

input `Int[(a + b*Cos[e + f*x])^4*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

```

output c^5*(-((a*A*(c*Sec[e + f*x])^(-5 + m)*(b + a*Sec[e + f*x])^3*Tan[e + f*x])
/(f*(1 - m))) + (-((a*(a*B*(1 - m) - A*b*(2 + m))*(c*Sec[e + f*x])^(-5 + m)
)*(b + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(2 - m))) - ((a^2*(2*a*b*B*(1 -
m)^2 + a^2*A*(2 - m)^2 + A*b^2*(6 - m + m^2))*Sec[e + f*x]*(c*Sec[e + f*x]
)^(-5 + m)*Tan[e + f*x])/(f*(3 - m)) + ((c*(3 - 4*m + m^2)*(4*a^3*A*b*(15
- 8*m + m^2) + a^4*B*(15 - 8*m + m^2) + 4*a*A*b^3*(10 - 7*m + m^2) + 6*a^2
*b^2*B*(10 - 7*m + m^2) + b^4*B*(8 - 6*m + m^2))*Hypergeometric2F1[1/2, (6
- m)/2, (8 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-6 + m)*Sin[e + f*x]
)/(f*(4 - m)*(6 - m)*Sqrt[Sin[e + f*x]^2]) + ((2 - m)*(a^4*A*(8 - 6*m + m^
2) + 6*a^2*A*b^2*(4 - 5*m + m^2) + 4*a^3*b*B*(4 - 5*m + m^2) + A*b^4*(3 -
4*m + m^2) + 4*a*b^3*B*(3 - 4*m + m^2))*Hypergeometric2F1[1/2, (5 - m)/2,
(7 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-5 + m)*Sin[e + f*x])/(f*(5 -
m)*Sqrt[Sin[e + f*x]^2]) + (a*(3 - m)*(4*a^2*A*b*(3 - 4*m + m^2) + a^3*B*
(3 - 4*m + m^2) + 2*A*b^3*(4 - 2*m + m^2) + a*b^2*B*(8 - 13*m + 5*m^2))*(c
*Sec[e + f*x])^(-5 + m)*Tan[e + f*x])/(f*(4 - m))/(3 - m)/(2 - m)/(1 -
m))

```

3.636.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3122 Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]

```

```

rule 3439 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]

```


rule 4259 $\text{Int}[(\text{csc}[(e_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)} \text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /;$
 $\text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[n]$

rule 4514 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.))^{(n_)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^{(m_)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[(-b)*B*\text{Cot}[e + f*x]*(a + b*\text{Csc}[e + f*x])^{(m-1)}*((d*\text{Csc}[e + f*x])^n/(f*(m+n))), x] + \text{Simp}[1/(m+n) \text{Int}[(a + b*\text{Csc}[e + f*x])^{(m-2)}*(d*\text{Csc}[e + f*x])^n*\text{Simp}[a^2*A*(m+n) + a*b*B*n + (a*(2*A*b + a*B))*(m+n) + b^2*B*(m+n-1)]*\text{Csc}[e + f*x] + b*(A*b*(m+n) + a*B*(2*m+n-1))*\text{Csc}[e + f*x]^2, x], x] /;$
 $\text{FreeQ}\{a, b, d, e, f, A, B, n\}, x \ \&\& \ \text{NeQ}[A*b - a*B, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ !(\text{IGtQ}[n, 1] \ \&\& \ !\text{IntegerQ}[m])$

rule 4534 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.))^{(m_)}*(\text{csc}[(e_.) + (f_.)*(x_)]^2*(C_.) + (A_)), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cot}[e + f*x]*((b*\text{Csc}[e + f*x])^m/(f*(m+1))), x] + \text{Simp}[(C*m + A*(m+1))/(m+1) \text{Int}[(b*\text{Csc}[e + f*x])^m, x], x] /;$
 $\text{FreeQ}\{b, e, f, A, C, m\}, x \ \&\& \ \text{NeQ}[C*m + A*(m+1), 0] \ \&\& \ !\text{LeQ}[m, -1]$

rule 4535 $\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.))^{(m_)}*((A_.) + \text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + \text{csc}[(e_.) + (f_.)*(x_)]^2*(C_)), x_Symbol] \rightarrow \text{Simp}[B/b \text{Int}[(b*\text{Csc}[e + f*x])^{(m+1)}, x], x] + \text{Int}[(b*\text{Csc}[e + f*x])^m*(A + C*\text{Csc}[e + f*x]^2), x] /;$
 $\text{FreeQ}\{b, e, f, A, B, C, m\}, x]$

rule 4564 $\text{Int}[(A_.) + \text{csc}[(e_.) + (f_.)*(x_)]*(B_.) + \text{csc}[(e_.) + (f_.)*(x_)]^2*(C_.))*(\text{csc}[(e_.) + (f_.)*(x_)]*(d_.))^{(n_)}*(\text{csc}[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] \rightarrow \text{Simp}[(-b)*C*\text{Csc}[e + f*x]*\text{Cot}[e + f*x]*((d*\text{Csc}[e + f*x])^n/(f*(n+2))), x] + \text{Simp}[1/(n+2) \text{Int}[(d*\text{Csc}[e + f*x])^n*\text{Simp}[A*a*(n+2) + (B*a*(n+2) + b*(C*(n+1) + A*(n+2))]*\text{Csc}[e + f*x] + (a*C + B*b)*(n+2)*\text{Csc}[e + f*x]^2, x], x], x] /;$
 $\text{FreeQ}\{a, b, d, e, f, A, B, C, n\}, x \ \&\& \ !\text{LtQ}[n, -1]$

rule 4584 `Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^n*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^m, x_Symbol] :> Simp[(-C)*Cot[e + f*x]*(a + b*Csc[e + f*x])^m*((d*Csc[e + f*x])^n/(f*(m + n + 1))), x] + Simp[1/(m + n + 1) Int[(a + b*Csc[e + f*x])^(m - 1)*(d*Csc[e + f*x])^n*Simp[a*A*(m + n + 1) + a*C*n + ((A*b + a*B)*(m + n + 1) + b*C*(m + n))*Csc[e + f*x] + (b*B*(m + n + 1) + a*C*m)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && !LeQ[n, -1]`

3.636.4 Maple [F]

$$\int (a + b \cos(fx + e))^4 (A + \cos(fx + e) B) (c \sec(fx + e))^m dx$$

input `int((a+b*cos(f*x+e))^4*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

output `int((a+b*cos(f*x+e))^4*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

3.636.5 Fracas [F]

$$\begin{aligned} & \int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx \\ &= \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^4 (c \sec(fx + e))^m dx \end{aligned}$$

input `integrate((a+b*cos(f*x+e))^4*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm="fricas")`

output `integral((B*b^4*cos(f*x + e)^5 + A*a^4 + (4*B*a*b^3 + A*b^4)*cos(f*x + e)^4 + 2*(3*B*a^2*b^2 + 2*A*a*b^3)*cos(f*x + e)^3 + 2*(2*B*a^3*b + 3*A*a^2*b^2)*cos(f*x + e)^2 + (B*a^4 + 4*A*a^3*b)*cos(f*x + e))*(c*sec(f*x + e))^m, x)`

3.636.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \text{Timed out}$$

input `integrate((a+b*cos(f*x+e))**4*(A+B*cos(f*x+e))*(c*sec(f*x+e))**m,x)`

output `Timed out`

3.636.7 Maxima [F]

$$\begin{aligned} & \int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^4 (c \sec(fx + e))^m dx \end{aligned}$$

input `integrate((a+b*cos(f*x+e))^4*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^4*(c*sec(f*x + e))^m, x)`

3.636.8 Giac [F]

$$\begin{aligned} & \int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^4 (c \sec(fx + e))^m dx \end{aligned}$$

input `integrate((a+b*cos(f*x+e))^4*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^4*(c*sec(f*x + e))^m, x)`

3.636.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))^4 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^4 dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^4,x)`output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^4, x)`

3.637 $\int (a+b \cos(e+fx))^3 (A+B \cos(e+fx))(c \sec(e+fx))^m dx$

3.637.1 Optimal result	5838
3.637.2 Mathematica [A] (verified)	5839
3.637.3 Rubi [A] (verified)	5840
3.637.4 Maple [F]	5845
3.637.5 Fracas [F]	5845
3.637.6 Sympy [F]	5846
3.637.7 Maxima [F]	5846
3.637.8 Giac [F]	5846
3.637.9 Mupad [F(-1)]	5847

3.637.1 Optimal result

Integrand size = 33, antiderivative size = 455

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx))(c \sec(e + fx))^m dx =$$

$$\frac{c^5(a^3 A(8 - 6m + m^2) + 3aAb^2(4 - 5m + m^2) + 3a^2bB(4 - 5m + m^2) + b^3B(3 - 4m + m^2)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4-m}{2}, \frac{6-m}{2}, \cos^2(e + fx)\right)}{f(1 - m)(3 - m)(5 - m)\sqrt{\sin^2(e + fx)}} +$$

$$\frac{c^4(Ab^3(2 - m) + 3ab^2B(2 - m) + 3a^2Ab(3 - m) + a^3B(3 - m)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4-m}{2}, \frac{6-m}{2}, \cos^2(e + fx)\right)}{f(2 - m)(4 - m)\sqrt{\sin^2(e + fx)}} +$$

$$\frac{ac^4(3abB(1 - m) + a^2A(2 - m) - 2Ab^2m) (c \sec(e + fx))^{-4+m} \tan(e + fx)}{f(1 - m)(3 - m)} +$$

$$\frac{a^2c^4(aB(1 - m) - Ab(1 + m)) \sec(e + fx)(c \sec(e + fx))^{-4+m} \tan(e + fx)}{f(1 - m)(2 - m)} +$$

$$\frac{aAc^4(c \sec(e + fx))^{-4+m}(b + a \sec(e + fx))^2 \tan(e + fx)}{f(1 - m)}$$

output $-c^5(a^3A(m^2-6m+8)+3aAb^2(m^2-5m+4)+3a^2bB(m^2-5m+4)+b^3B(m^2-4m+3))*\text{hypergeom}([1/2, 5/2-1/2m], [7/2-1/2m], \cos(f*x+e)^2)*(c*\sec(f*x+e))^{(-5+m)}*\sin(f*x+e)/f/(1-m)/(m^2-8m+15)/(\sin(f*x+e)^2)^{(1/2)}-c^4(A*b^3(2-m)+3a*b^2B(2-m)+3a^2A*b(3-m)+a^3B(3-m))*\text{hypergeom}([1/2, 2-1/2m], [3-1/2m], \cos(f*x+e)^2)*(c*\sec(f*x+e))^{(-4+m)}*\sin(f*x+e)/f/(m^2-6m+8)/(\sin(f*x+e)^2)^{(1/2)}-a*c^4(3a*b*B(1-m)+a^2A(2-m)-2A*b^2m)*(c*\sec(f*x+e))^{(-4+m)}*\tan(f*x+e)/f/(m^2-4m+3)-a^2*c^4(a*B(1-m)-A*b(1+m))*\sec(f*x+e)*(c*\sec(f*x+e))^{(-4+m)}*\tan(f*x+e)/f/(m^2-3m+2)-aA*c^4(c*\sec(f*x+e))^{(-4+m)}*(b+a*\sec(f*x+e))^2*\tan(f*x+e)/f/(1-m)$

3.637.2 Mathematica [A] (verified)

Time = 1.42 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.57

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \frac{\cot(e + fx) \left(\frac{b^3 B \cos^4(e + fx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-4 + m), \frac{1}{2}(-2 + m), \sec^2(e + fx)\right)}{-4 + m} + \frac{b^2 (Ab + 3aB) \cos^3(e + fx) \text{Hypergeometric2F1}}{-3 + m} \right)}{1}$$

input `Integrate[(a + b*Cos[e + f*x])^3*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output $(\text{Cot}[e + f*x]*((b^3*B*\text{Cos}[e + f*x]^4*\text{Hypergeometric2F1}[1/2, (-4 + m)/2, (-2 + m)/2, \text{Sec}[e + f*x]^2)]/(-4 + m) + (b^2*(A*b + 3*a*B)*\text{Cos}[e + f*x]^3*\text{Hypergeometric2F1}[1/2, (-3 + m)/2, (-1 + m)/2, \text{Sec}[e + f*x]^2)]/(-3 + m) + a*((3*b*(A*b + a*B)*\text{Cos}[e + f*x]^2*\text{Hypergeometric2F1}[1/2, (-2 + m)/2, m/2, \text{Sec}[e + f*x]^2)]/(-2 + m) + a*((3*A*b + a*B)*\text{Cos}[e + f*x]*\text{Hypergeometric2F1}[1/2, (-1 + m)/2, (1 + m)/2, \text{Sec}[e + f*x]^2)]/(-1 + m) + (a*A*\text{Hypergeometric2F1}[1/2, m/2, (2 + m)/2, \text{Sec}[e + f*x]^2)]/m))*c*\text{Sec}[e + f*x]^m*\text{Sqrt}[-\text{Tan}[e + f*x]^2])/f$

3.637.3 Rubi [A] (verified)

Time = 2.83 (sec) , antiderivative size = 450, normalized size of antiderivative = 0.99, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 3439, 3042, 4514, 25, 3042, 4564, 25, 3042, 4535, 3042, 4259, 3042, 3122, 4534, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^3 \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^m dx$$

$$\downarrow \text{3439}$$

$$c^4 \int (c \sec(e + fx))^{m-4} (b + a \sec(e + fx))^3 (B + A \sec(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$c^4 \int \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^{m-4} \left(b + a \csc \left(e + fx + \frac{\pi}{2} \right) \right)^3 \left(B + A \csc \left(e + fx + \frac{\pi}{2} \right) \right) dx$$

$$\downarrow \text{4514}$$

$$c^4 \left(- \frac{\int - (c \sec(e + fx))^{m-4} (b + a \sec(e + fx)) (a(aB(1 - m) - Ab(m + 1)) \sec^2(e + fx) + (A(2 - m)a^2 + b(Ab + 2Aa))) dx}{1 - m} \right)$$

$$\downarrow \text{25}$$

$$c^4 \left(\frac{\int (c \sec(e + fx))^{m-4} (b + a \sec(e + fx)) (a(aB(1 - m) - Ab(m + 1)) \sec^2(e + fx) + (A(2 - m)a^2 + b(Ab + 2Aa))) dx}{1 - m} \right)$$

$$\downarrow \text{3042}$$

$$c^4 \left(\frac{\int (c \csc \left(e + fx + \frac{\pi}{2} \right))^{m-4} (b + a \csc \left(e + fx + \frac{\pi}{2} \right)) \left(a(aB(1 - m) - Ab(m + 1)) \csc \left(e + fx + \frac{\pi}{2} \right)^2 + (A(2 - m)a^2 + b(Ab + 2Aa)) \right) dx}{1 - m} \right)$$

$$\downarrow \text{4564}$$

$$c^4 \left(\frac{\int -(\csc(e+fx))^{m-4} ((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \sec^2(e+fx)+(B(3-m)a^3+3Ab(3-m)a^2))}{2-m} dx + \frac{(1-m)(a^3B(3-m)+3a^2A(3-m)a)}{1-m} \right)$$

↓ 25

$$c^4 \left(\frac{\int (\csc(e+fx))^{m-4} ((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \sec^2(e+fx)+(B(3-m)a^3+3Ab(3-m)a^2))}{2-m} dx + \frac{(1-m)(a^3B(3-m)+3a^2A(3-m)a)}{1-m} \right)$$

↓ 3042

$$c^4 \left(\frac{\int (\csc(e+fx+\frac{\pi}{2}))^{m-4} ((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \csc(e+fx+\frac{\pi}{2})^2+(B(3-m)a^3+3Ab(3-m)a^2))}{2-m} dx + \frac{(1-m)(a^3B(3-m)+3a^2A(3-m)a)}{1-m} \right)$$

↓ 4535

$$c^4 \left(\frac{\int (\csc(e+fx))^{m-4} ((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \sec^2(e+fx)) dx + \frac{(1-m)(a^3B(3-m)+3a^2A(3-m)a)}{1-m}}{1-m} \right)$$

↓ 3042

$$c^4 \left(\frac{\int (\csc(e+fx+\frac{\pi}{2}))^{m-4} ((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \csc(e+fx+\frac{\pi}{2})^2) dx + \frac{(1-m)(a^3B(3-m)+3a^2A(3-m)a)}{1-m}}{1-m} \right)$$

↓ 4259

$$c^4 \left(\frac{\int (\csc(e+fx+\frac{\pi}{2}))^{m-4} ((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \csc(e+fx+\frac{\pi}{2})^2) dx + \frac{(1-m)(a^3B(3-m)+3a^2A(3-m)a)}{1-m}}{1-m} \right)$$

↓ 3042

$$c^4 \left(\frac{\int (c \csc(e+fx+\frac{\pi}{2}))^{m-4} \left((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \csc(e+fx+\frac{\pi}{2})^2 \right) dx + \frac{(1-m)(a^3B(3-m) + \dots)}{2-m}}{2-m} \right)$$

↓ 3122

$$c^4 \left(\frac{\int (c \csc(e+fx+\frac{\pi}{2}))^{m-4} \left((bB(1-m)+aA(4-m))(2-m)b^2+a(2-m)(A(2-m)a^2+3bB(1-m)a-2Ab^2m) \csc(e+fx+\frac{\pi}{2})^2 \right) dx - \frac{(1-m) \sin(e+fx)}{2-m}}{2-m} \right)$$

↓ 4534

$$c^4 \left(\frac{\frac{(2-m)(a^3A(m^2-6m+8)+3a^2bB(m^2-5m+4)+3aAb^2(m^2-5m+4)+b^3B(m^2-4m+3)) \int (c \sec(e+fx))^{m-4} dx}{3-m} - \frac{a(2-m) \tan(e+fx)(a^2A(2-m)+3abB(1-m))}{f(3-m)}}{2-m} \right)$$

↓ 3042

$$c^4 \left(\frac{\frac{(2-m)(a^3A(m^2-6m+8)+3a^2bB(m^2-5m+4)+3aAb^2(m^2-5m+4)+b^3B(m^2-4m+3)) \int (c \csc(e+fx+\frac{\pi}{2}))^{m-4} dx}{3-m} - \frac{a(2-m) \tan(e+fx)(a^2A(2-m)+3abB(1-m))}{f(3-m)}}{2-m} \right)$$

↓ 4259

$$c^4 \left(\frac{\frac{(2-m)(a^3A(m^2-6m+8)+3a^2bB(m^2-5m+4)+3aAb^2(m^2-5m+4)+b^3B(m^2-4m+3)) \left(\frac{\cos(e+fx)}{c} \right)^m (c \sec(e+fx))^m \int \left(\frac{\cos(e+fx)}{c} \right)^{4-m} dx}{3-m} - \frac{a(2-m)}{f(3-m)}}{2-m} \right)$$

↓ 3042

$$c^4 \left(\frac{(2-m)(a^3 A(m^2-6m+8)+3a^2 b B(m^2-5m+4)+3a A b^2(m^2-5m+4)+b^3 B(m^2-4m+3)) \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^m \int \left(\frac{\sin(e+fx+\frac{\pi}{2})}{c}\right)^{4-m} dx}{3-m} \right)$$

↓ 3122

$$c^4 \left(-\frac{a(2-m) \tan(e+fx)(a^2 A(2-m)+3ab B(1-m)-2Ab^2 m)(c \sec(e+fx))^{m-4}}{f(3-m)} - \frac{c(2-m) \sin(e+fx)(a^3 A(m^2-6m+8)+3a^2 b B(m^2-5m+4)+3a A b^2(m^2-5m+4)+b^3 B(m^2-4m+3))}{f(3-m)} \right)$$

```
input Int[(a + b*Cos[e + f*x])^3*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]
```

```
output c^4*((-(a*A*(c*Sec[e + f*x])^(-4 + m)*(b + a*Sec[e + f*x])^2*Tan[e + f*x])/(f*(1 - m))) + (-(a^2*(a*B*(1 - m) - A*b*(1 + m))*Sec[e + f*x]*(c*Sec[e + f*x])^(-4 + m)*Tan[e + f*x])/(f*(2 - m))) + (-(c*(2 - m)*(a^3*A*(8 - 6*m + m^2) + 3*a*A*b^2*(4 - 5*m + m^2) + 3*a^2*b*B*(4 - 5*m + m^2) + b^3*B*(3 - 4*m + m^2))*Hypergeometric2F1[1/2, (5 - m)/2, (7 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-5 + m)*Sin[e + f*x])/(f*(3 - m)*(5 - m)*Sqrt[Sin[e + f*x]^2])) - ((A*b^3*(2 - m) + 3*a*b^2*B*(2 - m) + 3*a^2*A*b*(3 - m) + a^3*B*(3 - m))*(1 - m)*Hypergeometric2F1[1/2, (4 - m)/2, (6 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-4 + m)*Sin[e + f*x])/(f*(4 - m)*Sqrt[Sin[e + f*x]^2]) - (a*(2 - m)*(3*a*b*B*(1 - m) + a^2*A*(2 - m) - 2*A*b^2*m)*(c*Sec[e + f*x])^(-4 + m)*Tan[e + f*x])/(f*(3 - m)))/(2 - m)/(1 - m))
```

3.637.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

3.637. $\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4514 `Int[(csc[(e_.) + (f_.)*(x_)])*(d_.)^(n_)*(csc[(e_.) + (f_.)*(x_)])*(b_.) + (a_.)^(m_)*(csc[(e_.) + (f_.)*(x_)])*(B_.) + (A_.)), x_Symbol] := Simp[(-b)*B*Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))), x] + Simp[1/(m + n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^n*Simp[a^2*A*(m + n) + a*b*B*n + (a*(2*A*b + a*B)*(m + n) + b^2*B*(m + n - 1))*Csc[e + f*x] + b*(A*b*(m + n) + a*B*(2*m + n - 1))*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && !IntegerQ[m])`

rule 4534 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.)^(m_.)*(csc[(e_.) + (f_.)*(x_)])^2*(C_.) + (A_.)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1))), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /; FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]`

rule 4535 `Int[(csc[(e_.) + (f_.)*(x_)])*(b_.)^(m_.)*((A_.) + csc[(e_.) + (f_.)*(x_)])*(B_.) + csc[(e_.) + (f_.)*(x_)])^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Csc[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2), x] /; FreeQ[{b, e, f, A, B, C, m}, x]`

```
rule 4564 Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))*(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (
a_)), x_Symbol] :> Simp[(-b)*C*Csc[e + f*x]*Cot[e + f*x]*((d*Csc[e + f*x])^
n/(f*(n + 2))), x] + Simp[1/(n + 2) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n +
2) + (B*a*(n + 2) + b*(C*(n + 1) + A*(n + 2)))*Csc[e + f*x] + (a*C + B*b)*(
n + 2)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, d, e, f, A, B, C, n}, x]
&& !LtQ[n, -1]
```

3.637.4 Maple [F]

$$\int (a + b \cos(fx + e))^3 (A + \cos(fx + e) B) (c \sec(fx + e))^m dx$$

```
input int((a+b*cos(f*x+e))^3*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)
```

```
output int((a+b*cos(f*x+e))^3*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)
```

3.637.5 Fracas [F]

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^3 (c \sec(fx + e))^m dx$$

```
input integrate((a+b*cos(f*x+e))^3*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorith
m="fricas")
```

```
output integral((B*b^3*cos(f*x + e)^4 + A*a^3 + (3*B*a*b^2 + A*b^3)*cos(f*x + e)^
3 + 3*(B*a^2*b + A*a*b^2)*cos(f*x + e)^2 + (B*a^3 + 3*A*a^2*b)*cos(f*x + e
))*c*sec(f*x + e))^m, x)
```

3.637.6 Sympy [F]

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (c \sec(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^3 dx$$

input `integrate((a+b*cos(f*x+e))**3*(A+B*cos(f*x+e))*(c*sec(f*x+e))**m,x)`

output `Integral((c*sec(e + f*x))**m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))**3, x)`

3.637.7 Maxima [F]

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^3 (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^3*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^3*(c*sec(f*x + e))^m, x)`

3.637.8 Giac [F]

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^3 (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^3*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^3*(c*sec(f*x + e))^m, x)`

3.637.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))^3 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^3 dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^3,x)`output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^3, x)`

3.638 $\int (a+b \cos(e+fx))^2(A+B \cos(e+fx))(c \sec(e+fx))^m dx$

3.638.1 Optimal result	5848
3.638.2 Mathematica [A] (verified)	5849
3.638.3 Rubi [A] (verified)	5849
3.638.4 Maple [F]	5853
3.638.5 Fracas [F]	5854
3.638.6 Sympy [F]	5854
3.638.7 Maxima [F]	5854
3.638.8 Giac [F]	5855
3.638.9 Mupad [F(-1)]	5855

3.638.1 Optimal result

Integrand size = 33, antiderivative size = 327

$$\int (a + b \cos(e + fx))^2(A + B \cos(e + fx))(c \sec(e + fx))^m dx =$$

$$\frac{c^4(b^2B(2 - m) + 2aAb(3 - m) + a^2B(3 - m)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4-m}{2}, \frac{6-m}{2}, \cos^2(e + fx)\right) (c \sec(e + fx))^{m-4}}{f(2 - m)(4 - m)\sqrt{\sin^2(e + fx)}}$$

$$\frac{c^3(Ab^2(1 - m) + 2abB(1 - m) + a^2A(2 - m)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3-m}{2}, \frac{5-m}{2}, \cos^2(e + fx)\right) (c \sec(e + fx))^{m-3}}{f(1 - m)(3 - m)\sqrt{\sin^2(e + fx)}}$$

$$\frac{ac^3(aB(1 - m) - Abm)(c \sec(e + fx))^{-3+m} \tan(e + fx)}{f(1 - m)(2 - m)}$$

$$\frac{aAc^3(c \sec(e + fx))^{-3+m}(b + a \sec(e + fx)) \tan(e + fx)}{f(1 - m)}$$

```
output -c^4*(b^2*B*(2-m)+2*a*A*b*(3-m)+a^2*B*(3-m))*hypergeom([1/2, 2-1/2*m], [3-1/2*m], cos(f*x+e)^2)*(c*sec(f*x+e))^(m-4)*sin(f*x+e)/f/(m^2-6*m+8)/(sin(f*x+e)^2)^(1/2)-c^3*(A*b^2*(1-m)+2*a*b*B*(1-m)+a^2*A*(2-m))*hypergeom([1/2, 3/2-1/2*m], [5/2-1/2*m], cos(f*x+e)^2)*(c*sec(f*x+e))^(m-3)*sin(f*x+e)/f/(m^2-4*m+3)/(sin(f*x+e)^2)^(1/2)-a*c^3*(a*B*(1-m)-A*b*m)*(c*sec(f*x+e))^(m-3)*tan(f*x+e)/f/(m^2-3*m+2)-a*A*c^3*(c*sec(f*x+e))^(m-3)*(b+a*sec(f*x+e))*tan(f*x+e)/f/(1-m)
```

3.638.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.63

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \frac{\cot(e + fx) \left(\frac{b^2 B \cos^3(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-3 + m), \frac{1}{2}(-1 + m), \sec^2(e + fx)\right)}{-3 + m} + \frac{b(Ab + 2aB) \cos^2(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2 + m), \frac{1}{2}(-1 + m), \sec^2(e + fx)\right)}{-2 + m} \right)}{c^m}$$

input `Integrate[(a + b*Cos[e + f*x])^2*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `(Cot[e + f*x]*((b^2*B*Cos[e + f*x]^3*Hypergeometric2F1[1/2, (-3 + m)/2, (-1 + m)/2, Sec[e + f*x]^2)]/(-3 + m) + (b*(A*b + 2*a*B)*Cos[e + f*x]^2*Hypergeometric2F1[1/2, (-2 + m)/2, m/2, Sec[e + f*x]^2)]/(-2 + m) + a*((2*A*b + a*B)*Cos[e + f*x]*Hypergeometric2F1[1/2, (-1 + m)/2, (1 + m)/2, Sec[e + f*x]^2)]/(-1 + m) + (a*A*Hypergeometric2F1[1/2, m/2, (2 + m)/2, Sec[e + f*x]^2])/m)*(c*Sec[e + f*x])^m*sqrt[-Tan[e + f*x]^2])/f`

3.638.3 Rubi [A] (verified)

Time = 1.80 (sec) , antiderivative size = 315, normalized size of antiderivative = 0.96, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 3439, 3042, 4514, 25, 3042, 4535, 3042, 4259, 3042, 3122, 4534, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^2 \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^m dx$$

$$\downarrow \text{3439}$$

$$c^3 \int (c \sec(e + fx))^{m-3} (b + a \sec(e + fx))^2 (B + A \sec(e + fx)) dx$$

$$\downarrow \text{3042}$$

3.638. $\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$

$$\begin{aligned}
& c^3 \int \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^{m-3} \left(b + a \csc \left(e + fx + \frac{\pi}{2} \right) \right)^2 \left(B + A \csc \left(e + fx + \frac{\pi}{2} \right) \right) dx \\
& \quad \downarrow \text{4514} \\
& c^3 \left(\frac{\int - (c \sec(e + fx))^{m-3} (a(aB(1-m) - Abm) \sec^2(e + fx) + (A(2-m)a^2 + b(Ab + 2aB)(1-m)) \sec(e + fx)) dx}{1-m} \right) \\
& \quad \downarrow \text{25} \\
& c^3 \left(\frac{\int (c \sec(e + fx))^{m-3} (a(aB(1-m) - Abm) \sec^2(e + fx) + (A(2-m)a^2 + b(Ab + 2aB)(1-m)) \sec(e + fx)) dx}{1-m} \right) \\
& \quad \downarrow \text{3042} \\
& c^3 \left(\frac{\int (c \csc(e + fx + \frac{\pi}{2}))^{m-3} (a(aB(1-m) - Abm) \csc(e + fx + \frac{\pi}{2})^2 + (A(2-m)a^2 + b(Ab + 2aB)(1-m)) \csc(e + fx + \frac{\pi}{2})) dx}{1-m} \right) \\
& \quad \downarrow \text{4535} \\
& c^3 \left(\frac{\frac{(a^2A(2-m)+b(1-m)(2aB+Ab))}{c} \int (c \sec(e + fx))^{m-2} dx}{1-m} + \int (c \sec(e + fx))^{m-3} (a(aB(1-m) - Abm) \sec^2(e + fx) + b(Ab + 2aB)(1-m)) dx}{1-m} \right) \\
& \quad \downarrow \text{3042} \\
& c^3 \left(\frac{\frac{(a^2A(2-m)+b(1-m)(2aB+Ab))}{c} \int (c \csc(e + fx + \frac{\pi}{2}))^{m-2} dx}{1-m} + \int (c \csc(e + fx + \frac{\pi}{2}))^{m-3} (a(aB(1-m) - Abm) \csc(e + fx + \frac{\pi}{2})^2 + b(Ab + 2aB)(1-m)) dx}{1-m} \right) \\
& \quad \downarrow \text{4259} \\
& c^3 \left(\frac{\frac{(a^2A(2-m)+b(1-m)(2aB+Ab)) \left(\frac{\cos(e+fx)}{c} \right)^m (c \sec(e+fx))^m \int \left(\frac{\cos(e+fx)}{c} \right)^{2-m} dx}{c} + \int (c \csc(e + fx + \frac{\pi}{2}))^{m-3} (a(aB(1-m) - Abm) \csc(e + fx + \frac{\pi}{2})^2 + b(Ab + 2aB)(1-m)) dx}{1-m} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.638. $\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$

$$c^3 \left(\frac{(a^2A(2-m)+b(1-m)(2aB+Ab)) \left(\frac{\cos(e+fx)}{c}\right)^m (c \operatorname{csc}(e+fx))^m \int \left(\frac{\sin(e+fx+\frac{\pi}{2})}{c}\right)^{2-m} dx}{c} + \int (c \operatorname{csc}(e+fx+\frac{\pi}{2}))^{m-3} (a(aB(1-m) - Abm) \operatorname{csc}(e+fx+\frac{\pi}{2})^2 + b(bB(1-m) + aA(3-m))) dx - \frac{\sin(e+fx)(a^2A(2-m)+b(1-m)(2aB+Ab))(c \operatorname{csc}(e+fx))^{m-3}}{f(3-m)\sqrt{\sin^2(e+fx)}}}{1-m} \right)$$

↓ 3122

$$c^3 \left(\frac{\int (c \operatorname{csc}(e+fx+\frac{\pi}{2}))^{m-3} (a(aB(1-m) - Abm) \operatorname{csc}(e+fx+\frac{\pi}{2})^2 + b(bB(1-m) + aA(3-m))) dx - \frac{\sin(e+fx)(a^2A(2-m)+b(1-m)(2aB+Ab))(c \operatorname{csc}(e+fx))^{m-3}}{f(3-m)\sqrt{\sin^2(e+fx)}}}{1-m} \right)$$

↓ 4534

$$c^3 \left(\frac{(1-m)(a^2B(3-m)+2aAb(3-m)+b^2B(2-m)) \int (c \operatorname{csc}(e+fx))^{m-3} dx - \frac{\sin(e+fx)(a^2A(2-m)+b(1-m)(2aB+Ab))(c \operatorname{csc}(e+fx))^{m-3}}{f(3-m)\sqrt{\sin^2(e+fx)}}}{2-m} \right)$$

↓ 3042

$$c^3 \left(\frac{(1-m)(a^2B(3-m)+2aAb(3-m)+b^2B(2-m)) \int (c \operatorname{csc}(e+fx+\frac{\pi}{2}))^{m-3} dx - \frac{\sin(e+fx)(a^2A(2-m)+b(1-m)(2aB+Ab))(c \operatorname{csc}(e+fx))^{m-3}}{f(3-m)\sqrt{\sin^2(e+fx+\frac{\pi}{2})}}}{2-m} \right)$$

↓ 4259

$$c^3 \left(\frac{(1-m)(a^2B(3-m)+2aAb(3-m)+b^2B(2-m)) \left(\frac{\cos(e+fx)}{c}\right)^m (c \operatorname{csc}(e+fx))^m \int \left(\frac{\cos(e+fx)}{c}\right)^{3-m} dx - \frac{\sin(e+fx)(a^2A(2-m)+b(1-m)(2aB+Ab))(c \operatorname{csc}(e+fx))^{m-3}}{f(3-m)\sqrt{\sin^2(e+fx)}}}{2-m} \right)$$

↓ 3042

$$c^3 \left(\frac{(1-m)(a^2B(3-m)+2aAb(3-m)+b^2B(2-m)) \left(\frac{\cos(e+fx)}{c}\right)^m (c \operatorname{csc}(e+fx))^m \int \left(\frac{\sin(e+fx+\frac{\pi}{2})}{c}\right)^{3-m} dx - \frac{\sin(e+fx)(a^2A(2-m)+b(1-m)(2aB+Ab))(c \operatorname{csc}(e+fx))^{m-3}}{f(3-m)\sqrt{\sin^2(e+fx+\frac{\pi}{2})}}}{2-m} \right)$$

↓ 3122

$$c^3 \left(\frac{-c(1-m)\sin(e+fx)(a^2B(3-m)+2aAb(3-m)+b^2B(2-m))(c\sec(e+fx))^{m-4} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4-m}{2}, \frac{6-m}{2}, \cos^2(e+fx)\right) - \sin(e+fx)}{f(2-m)(4-m)\sqrt{\sin^2(e+fx)}} \right)$$

input `Int[(a + b*Cos[e + f*x])^2*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `c^3*(-((a*A*(c*Sec[e + f*x])^(-3 + m)*(b + a*Sec[e + f*x])*Tan[e + f*x])/(f*(1 - m))) + (-((c*(b^2*B*(2 - m) + 2*a*A*b*(3 - m) + a^2*B*(3 - m))*(1 - m)*Hypergeometric2F1[1/2, (4 - m)/2, (6 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-4 + m)*Sin[e + f*x])/(f*(2 - m)*(4 - m)*Sqrt[Sin[e + f*x]^2])) - ((b*(A*b + 2*a*B)*(1 - m) + a^2*A*(2 - m))*Hypergeometric2F1[1/2, (3 - m)/2, (5 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-3 + m)*Sin[e + f*x])/(f*(3 - m)*Sqrt[Sin[e + f*x]^2]) - (a*(a*B*(1 - m) - A*b*m)*(c*Sec[e + f*x])^(-3 + m)*Tan[e + f*x])/(f*(2 - m)))/(1 - m))`

3.638.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4259 Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

```
rule 4514 Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (
a_))^(m_)*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_)), x_Symbol] := Simp[(-b)*B*
Cot[e + f*x]*(a + b*Csc[e + f*x])^(m - 1)*((d*Csc[e + f*x])^n/(f*(m + n))),
x] + Simp[1/(m + n) Int[(a + b*Csc[e + f*x])^(m - 2)*(d*Csc[e + f*x])^n*
Simp[a^2*A*(m + n) + a*b*B*n + (a*(2*A*b + a*B)*(m + n) + b^2*B*(m + n - 1)
)*Csc[e + f*x] + b*(A*b*(m + n) + a*B*(2*m + n - 1))*Csc[e + f*x]^2, x], x]
/; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2
- b^2, 0] && GtQ[m, 1] && !(IGtQ[n, 1] && !IntegerQ[m])
```

```
rule 4534 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_)*(csc[(e_.) + (f_.)*(x_)]^2*(C_.)
+ (A_)), x_Symbol] := Simp[(-C)*Cot[e + f*x]*((b*Csc[e + f*x])^m/(f*(m + 1)
)), x] + Simp[(C*m + A*(m + 1))/(m + 1) Int[(b*Csc[e + f*x])^m, x], x] /;
FreeQ[{b, e, f, A, C, m}, x] && NeQ[C*m + A*(m + 1), 0] && !LeQ[m, -1]
```

```
rule 4535 Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(m_)*((A_.) + csc[(e_.) + (f_.)*(x_)]*
(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.)), x_Symbol] := Simp[B/b Int[(b*Cs
c[e + f*x])^(m + 1), x], x] + Int[(b*Csc[e + f*x])^m*(A + C*Csc[e + f*x]^2)
, x] /; FreeQ[{b, e, f, A, B, C, m}, x]
```

3.638.4 Maple [F]

$$\int (a + b \cos(fx + e))^2 (A + \cos(fx + e) B) (c \sec(fx + e))^m dx$$

```
input int((a+b*cos(f*x+e))^2*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)
```

```
output int((a+b*cos(f*x+e))^2*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)
```

3.638.5 Fracas [F]

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^2 (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^2*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="fricas")`

output `integral((B*b^2*cos(f*x + e)^3 + A*a^2 + (2*B*a*b + A*b^2)*cos(f*x + e)^2 + (B*a^2 + 2*A*a*b)*cos(f*x + e))*(c*sec(f*x + e))^m, x)`

3.638.6 Sympy [F]

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (c \sec(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^2 dx$$

input `integrate((a+b*cos(f*x+e))**2*(A+B*cos(f*x+e))*(c*sec(f*x+e))**m,x)`

output `Integral((c*sec(e + f*x))**m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))**2, x)`

3.638.7 Maxima [F]

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^2 (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^2*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^2*(c*sec(f*x + e))^m, x)`

3.638.8 Giac [F]

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)^2 (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^2*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm m="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^2*(c*sec(f*x + e))^m, x)`

3.638.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))^2 (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^2 dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^2,x)`

output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^2, x)`

3.639 $\int (a+b \cos(e+fx))(A+B \cos(e+fx))(c \sec(e+fx))^m dx$

3.639.1 Optimal result	5856
3.639.2 Mathematica [A] (verified)	5857
3.639.3 Rubi [A] (verified)	5857
3.639.4 Maple [F]	5860
3.639.5 Fricas [F]	5860
3.639.6 Sympy [F]	5861
3.639.7 Maxima [F]	5861
3.639.8 Giac [F]	5861
3.639.9 Mupad [F(-1)]	5862

3.639.1 Optimal result

Integrand size = 31, antiderivative size = 217

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx =$$

$$\frac{c^3(bB(1 - m) + aA(2 - m)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3-m}{2}, \frac{5-m}{2}, \cos^2(e + fx)\right) (c \sec(e + fx))^{-3+m} \sin(e + fx)}{f(1 - m)(3 - m)\sqrt{\sin^2(e + fx)}} -$$

$$\frac{(Ab + aB)c^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-m}{2}, \frac{4-m}{2}, \cos^2(e + fx)\right) (c \sec(e + fx))^{-2+m} \sin(e + fx)}{f(2 - m)\sqrt{\sin^2(e + fx)}} -$$

$$\frac{aAc^2(c \sec(e + fx))^{-2+m} \tan(e + fx)}{f(1 - m)}$$

```
output -c^3*(b*B*(1-m)+a*A*(2-m))*hypergeom([1/2, 3/2-1/2*m],[5/2-1/2*m],cos(f*x+
e)^2)*(c*sec(f*x+e))^(3+m)*sin(f*x+e)/f/(m^2-4*m+3)/(sin(f*x+e)^2)^(1/2)-
(A*b+B*a)*c^2*hypergeom([1/2, 1-1/2*m],[2-1/2*m],cos(f*x+e)^2)*(c*sec(f*x+
e))^(2+m)*sin(f*x+e)/f/(2-m)/(sin(f*x+e)^2)^(1/2)-a*A*c^2*(c*sec(f*x+e))^(
2+m)*tan(f*x+e)/f/(1-m)
```

3.639.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.75

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \frac{\cot(e + fx) (bB(-1 + m)m \cos^2(e + fx) \text{Hypergeometric2F1}(\frac{1}{2}, \frac{1}{2}(-2 + m), \frac{m}{2}, \sec^2(e + fx)) + (-2 + m) \dots)}{\dots}$$

input `Integrate[(a + b*Cos[e + f*x])*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`output `(Cot[e + f*x]*(b*B*(-1 + m)*m*Cos[e + f*x]^2*Hypergeometric2F1[1/2, (-2 + m)/2, m/2, Sec[e + f*x]^2] + (-2 + m)*((A*b + a*B)*m*Cos[e + f*x]*Hypergeometric2F1[1/2, (-1 + m)/2, (1 + m)/2, Sec[e + f*x]^2] + a*A*(-1 + m)*Hypergeometric2F1[1/2, m/2, (2 + m)/2, Sec[e + f*x]^2]))*(c*Sec[e + f*x])^m*Sqrt[-Tan[e + f*x]^2])/(f*(-2 + m)*(-1 + m)*m)`**3.639.3 Rubi [A] (verified)**Time = 0.91 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 3439, 3042, 4485, 25, 3042, 4274, 3042, 4259, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$\downarrow \text{3042}$$

$$\int \left(a + b \sin\left(e + fx + \frac{\pi}{2}\right) \right) \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right) \right) \left(c \csc\left(e + fx + \frac{\pi}{2}\right) \right)^m dx$$

$$\downarrow \text{3439}$$

$$c^2 \int (c \sec(e + fx))^{m-2} (b + a \sec(e + fx))(B + A \sec(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$c^2 \int \left(c \csc\left(e + fx + \frac{\pi}{2}\right) \right)^{m-2} \left(b + a \csc\left(e + fx + \frac{\pi}{2}\right) \right) \left(B + A \csc\left(e + fx + \frac{\pi}{2}\right) \right) dx$$

$$\downarrow \text{4485}$$

$$c^2 \left(\frac{\int -(c \sec(e + fx))^{m-2} (bB(1-m) + (Ab + aB) \sec(e + fx)(1-m) + aA(2-m)) dx}{1-m} - \frac{aA \tan(e + fx)(c \sec(e + fx))}{f(1-m)} \right)$$

↓ 25

$$c^2 \left(\frac{\int (c \sec(e + fx))^{m-2} (bB(1-m) + (Ab + aB) \sec(e + fx)(1-m) + aA(2-m)) dx}{1-m} - \frac{aA \tan(e + fx)(c \sec(e + fx))}{f(1-m)} \right)$$

↓ 3042

$$c^2 \left(\frac{\int (c \csc(e + fx + \frac{\pi}{2}))^{m-2} (bB(1-m) + (Ab + aB) \csc(e + fx + \frac{\pi}{2})(1-m) + aA(2-m)) dx}{1-m} - \frac{aA \tan(e + fx + \frac{\pi}{2})(c \csc(e + fx + \frac{\pi}{2}))}{f(1-m)} \right)$$

↓ 4274

$$c^2 \left(\frac{(aA(2-m) + bB(1-m)) \int (c \sec(e + fx))^{m-2} dx + \frac{(1-m)(aB+Ab) \int (c \sec(e+fx))^{m-1} dx}{c}}{1-m} - \frac{aA \tan(e + fx)(c \sec(e + fx))}{f(1-m)} \right)$$

↓ 3042

$$c^2 \left(\frac{(aA(2-m) + bB(1-m)) \int (c \csc(e + fx + \frac{\pi}{2}))^{m-2} dx + \frac{(1-m)(aB+Ab) \int (c \csc(e+fx+\frac{\pi}{2}))^{m-1} dx}{c}}{1-m} - \frac{aA \tan(e + fx + \frac{\pi}{2})(c \csc(e + fx + \frac{\pi}{2}))}{f(1-m)} \right)$$

↓ 4259

$$c^2 \left(\frac{\frac{(1-m)(aB+Ab) \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^m \int \left(\frac{\cos(e+fx)}{c}\right)^{1-m} dx}{c} + (aA(2-m) + bB(1-m)) \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e + fx))^{m-2}}{1-m} - \frac{aA \tan(e + fx)(c \sec(e + fx))}{f(1-m)} \right)$$

↓ 3042

$$c^2 \left(\frac{\frac{(1-m)(aB+Ab) \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^m \int \left(\frac{\sin(e+fx+\frac{\pi}{2})}{c}\right)^{1-m} dx}{c} + (aA(2-m) + bB(1-m)) \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e + fx))^{m-2}}{1-m} - \frac{aA \tan(e + fx)(c \sec(e + fx))}{f(1-m)} \right)$$

↓ 3122

$$c^2 \left(\frac{-\frac{c \sin(e+fx)(aA(2-m)+bB(1-m))(c \sec(e+fx))^{m-3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3-m}{2}, \frac{5-m}{2}, \cos^2(e+fx)\right)}{f(3-m)\sqrt{\sin^2(e+fx)}} - \frac{(1-m)(aB+Ab) \sin(e+fx)(c \sec(e+fx))^{m-2}}{1-m} \right)$$

input `Int[(a + b*Cos[e + f*x])*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `c^2*((-((c*(b*B*(1 - m) + a*A*(2 - m))*Hypergeometric2F1[1/2, (3 - m)/2, (5 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-3 + m)*Sin[e + f*x])/(f*(3 - m)*Sqrt[Sin[e + f*x]^2])) - ((A*b + a*B)*(1 - m)*Hypergeometric2F1[1/2, (2 - m)/2, (4 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-2 + m)*Sin[e + f*x])/(f*(2 - m)*Sqrt[Sin[e + f*x]^2]))/(1 - m) - (a*A*(c*Sec[e + f*x])^(-2 + m)*Tan[e + f*x])/(f*(1 - m)))`

3.639.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d + c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4274 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> Simp[a Int[(d*Csc[e + f*x])^n, x], x] + Simp[b/d Int[(d*Csc[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, n}, x]`

rule 4485 `Int[(csc[(e_.) + (f_.)*(x_)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))*(csc[(e_.) + (f_.)*(x_)]*(B_.) + (A_.)), x_Symbol] :> Simp[(-b)*B*Cot[e + f*x]*((d*Csc[e + f*x])^n/(f*(n + 1))), x] + Simp[1/(n + 1) Int[(d*Csc[e + f*x])^n*Simp[A*a*(n + 1) + B*b*n + (A*b + B*a)*(n + 1)*Csc[e + f*x], x], x], x] /; FreeQ[{a, b, d, e, f, A, B}, x] && NeQ[A*b - a*B, 0] && !LeQ[n, -1]`

3.639.4 Maple [F]

$$\int (a + b \cos(fx + e))(A + \cos(fx + e)B)(c \sec(fx + e))^m dx$$

input `int((a+b*cos(f*x+e))*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

output `int((a+b*cos(f*x+e))*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

3.639.5 Fracas [F]

$$\begin{aligned} & \int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx \\ &= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)(c \sec(fx + e))^m dx \end{aligned}$$

input `integrate((a+b*cos(f*x+e))*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm="fracas")`

output `integral((B*b*cos(f*x + e)^2 + A*a + (B*a + A*b)*cos(f*x + e))*(c*sec(f*x + e))^m, x)`

3.639.6 Sympy [F]

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (c \sec(e + fx))^m (A + B \cos(e + fx)) (a + b \cos(e + fx)) dx$$

input `integrate((a+b*cos(f*x+e))*(A+B*cos(f*x+e))*(c*sec(f*x+e))**m,x)`

output `Integral((c*sec(e + f*x))**m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x)), x)`

3.639.7 Maxima [F]

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)(c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m, x)`

3.639.8 Giac [F]

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A)(b \cos(fx + e) + a)(c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m, x)`

3.639.9 Mupad [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) (a + b \cos(e + fx)) dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x)),x)`output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x)), x)`

$$3.640 \quad \int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{a+b \cos(e+fx)} dx$$

3.640.1 Optimal result	5863
3.640.2 Mathematica [B] (warning: unable to verify)	5864
3.640.3 Rubi [A] (verified)	5864
3.640.4 Maple [F]	5868
3.640.5 Fracas [F]	5868
3.640.6 Sympy [F]	5869
3.640.7 Maxima [F]	5869
3.640.8 Giac [F(-2)]	5869
3.640.9 Mupad [F(-1)]	5870

3.640.1 Optimal result

Integrand size = 33, antiderivative size = 299

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx =$$

$$\frac{(Ab - aB) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{m}{2}, 1, \frac{3}{2}, \sin^2(e + fx), -\frac{b^2 \sin^2(e + fx)}{a^2 - b^2}\right) \cos(e + fx) \cos^2(e + fx)^{m/2} (c \sec(e + fx))}{(a^2 - b^2) cf}$$

$$+ \frac{a(Ab - aB) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1+m}{2}, 1, \frac{3}{2}, \sin^2(e + fx), -\frac{b^2 \sin^2(e + fx)}{a^2 - b^2}\right) \cos^2(e + fx)^{\frac{1+m}{2}} (c \sec(e + fx))^{1+m} \sin(e + fx)}{b(a^2 - b^2) cf}$$

$$- \frac{Bc \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e + fx)\right) (c \sec(e + fx))^{-1+m} \sin(e + fx)}{bf(1 - m) \sqrt{\sin^2(e + fx)}}$$

output

```

-(A*b-B*a)*AppellF1(1/2,1/2*m,1,3/2,sin(f*x+e)^2,-b^2*sin(f*x+e)^2/(a^2-b^2))*cos(f*x+e)*(cos(f*x+e)^2)^(1/2*m)*(c*sec(f*x+e))^(1+m)*sin(f*x+e)/(a^2-b^2)/c/f+a*(A*b-B*a)*AppellF1(1/2,1/2+1/2*m,1,3/2,sin(f*x+e)^2,-b^2*sin(f*x+e)^2/(a^2-b^2))*(cos(f*x+e)^2)^(1/2+1/2*m)*(c*sec(f*x+e))^(1+m)*sin(f*x+e)/b/(a^2-b^2)/c/f-B*c*hypergeom([1/2,-1/2*m+1/2],[3/2-1/2*m],cos(f*x+e)^2)*(c*sec(f*x+e))^(-1+m)*sin(f*x+e)/b/f/(1-m)/(sin(f*x+e)^2)^(1/2)
    
```

3.640.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 10630 vs. $2(299) = 598$.

Time = 32.02 (sec) , antiderivative size = 10630, normalized size of antiderivative = 35.55

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx = \text{Result too large to show}$$

input `Integrate[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/(a + b*Cos[e + f*x]),x]`

output `Result too large to show`

3.640.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 3439, 3042, 4526, 3042, 4259, 3042, 3122, 4356, 3042, 3302, 3042, 3668, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(A + B \sin(e + fx + \frac{\pi}{2}))(c \csc(e + fx + \frac{\pi}{2}))^m}{a + b \sin(e + fx + \frac{\pi}{2})} dx \\ & \quad \downarrow \text{3439} \\ & \int \frac{(A \sec(e + fx) + B)(c \sec(e + fx))^m}{a \sec(e + fx) + b} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(A \csc(e + fx + \frac{\pi}{2}) + B)(c \csc(e + fx + \frac{\pi}{2}))^m}{a \csc(e + fx + \frac{\pi}{2}) + b} dx \\ & \quad \downarrow \text{4526} \\ & \frac{(Ab - aB) \int \frac{(c \sec(e + fx))^{m+1}}{b + a \sec(e + fx)} dx}{bc} + \frac{B \int (c \sec(e + fx))^m dx}{b} \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{(Ab - aB) \int \frac{(c \csc(e+fx+\frac{\pi}{2}))^{m+1}}{b+a \csc(e+fx+\frac{\pi}{2})} dx}{bc} + \frac{B \int (c \csc(e+fx+\frac{\pi}{2}))^m dx}{b} \\
& \downarrow 4259 \\
& \frac{(Ab - aB) \int \frac{(c \csc(e+fx+\frac{\pi}{2}))^{m+1}}{b+a \csc(e+fx+\frac{\pi}{2})} dx}{bc} + \frac{B \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^m \int \left(\frac{\cos(e+fx)}{c}\right)^{-m} dx}{b} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \int \frac{(c \csc(e+fx+\frac{\pi}{2}))^{m+1}}{b+a \csc(e+fx+\frac{\pi}{2})} dx}{bc} + \frac{B \left(\frac{\cos(e+fx)}{c}\right)^m (c \sec(e+fx))^m \int \left(\frac{\sin(e+fx+\frac{\pi}{2})}{c}\right)^{-m} dx}{b} \\
& \downarrow 3122 \\
& \frac{(Ab - aB) \int \frac{(c \csc(e+fx+\frac{\pi}{2}))^{m+1}}{b+a \csc(e+fx+\frac{\pi}{2})} dx}{bc} - \\
& \frac{Bc \sin(e+fx)(c \sec(e+fx))^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e+fx)\right)}{bf(1-m)\sqrt{\sin^2(e+fx)}} \\
& \downarrow 4356 \\
& \frac{(Ab - aB) \cos^{m+1}(e+fx)(c \sec(e+fx))^{m+1} \int \frac{\cos^{-m}(e+fx)}{a+b \cos(e+fx)} dx}{bc} - \\
& \frac{Bc \sin(e+fx)(c \sec(e+fx))^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e+fx)\right)}{bf(1-m)\sqrt{\sin^2(e+fx)}} \\
& \downarrow 3042 \\
& \frac{(Ab - aB) \cos^{m+1}(e+fx)(c \sec(e+fx))^{m+1} \int \frac{\sin(e+fx+\frac{\pi}{2})^{-m}}{a+b \sin(e+fx+\frac{\pi}{2})} dx}{bc} - \\
& \frac{Bc \sin(e+fx)(c \sec(e+fx))^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e+fx)\right)}{bf(1-m)\sqrt{\sin^2(e+fx)}} \\
& \downarrow 3302 \\
& \frac{(Ab - aB) \cos^{m+1}(e+fx)(c \sec(e+fx))^{m+1} \left(a \int \frac{\cos^{-m}(e+fx)}{a^2-b^2 \cos^2(e+fx)} dx - b \int \frac{\cos^{1-m}(e+fx)}{a^2-b^2 \cos^2(e+fx)} dx \right)}{bc} - \\
& \frac{Bc \sin(e+fx)(c \sec(e+fx))^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e+fx)\right)}{bf(1-m)\sqrt{\sin^2(e+fx)}} \\
& \downarrow 3042
\end{aligned}$$

3.640. $\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{a+b \cos(e+fx)} dx$

$$\begin{aligned}
& \frac{(Ab - aB) \cos^{m+1}(e + fx)(c \sec(e + fx))^{m+1} \left(a \int \frac{\sin(e+fx+\frac{\pi}{2})^{-m}}{a^2-b^2 \sin(e+fx+\frac{\pi}{2})^2} dx - b \int \frac{\sin(e+fx+\frac{\pi}{2})^{1-m}}{a^2-b^2 \sin(e+fx+\frac{\pi}{2})^2} dx \right)}{bc} \\
& \frac{Bc \sin(e + fx)(c \sec(e + fx))^{m-1} \operatorname{Hypergeometric2F1} \left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e + fx) \right)}{bf(1-m)\sqrt{\sin^2(e + fx)}} \\
& \quad \downarrow \text{3668} \\
& \frac{(Ab - aB) \cos^{m+1}(e + fx)(c \sec(e + fx))^{m+1} \left(\frac{a \cos^{-m-1}(e+fx) \cos^2(e+fx)^{\frac{m+1}{2}} \int \frac{(1-\sin^2(e+fx))^{\frac{1}{2}(-m-1)}}{a^2-b^2+b^2 \sin^2(e+fx)} d \sin(e+fx)}{f} - b \cos \right)}{bc} \\
& \frac{Bc \sin(e + fx)(c \sec(e + fx))^{m-1} \operatorname{Hypergeometric2F1} \left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e + fx) \right)}{bf(1-m)\sqrt{\sin^2(e + fx)}} \\
& \quad \downarrow \text{333} \\
& \frac{(Ab - aB) \cos^{m+1}(e + fx)(c \sec(e + fx))^{m+1} \left(\frac{a \sin(e+fx) \cos^{-m-1}(e+fx) \cos^2(e+fx)^{\frac{m+1}{2}} \operatorname{AppellF1} \left(\frac{1}{2}, \frac{m+1}{2}, 1, \frac{3}{2}, \sin^2(e+fx), - \right)}{f(a^2-b^2)} \right)}{bc} \\
& \frac{Bc \sin(e + fx)(c \sec(e + fx))^{m-1} \operatorname{Hypergeometric2F1} \left(\frac{1}{2}, \frac{1-m}{2}, \frac{3-m}{2}, \cos^2(e + fx) \right)}{bf(1-m)\sqrt{\sin^2(e + fx)}}
\end{aligned}$$

input `Int[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/(a + b*Cos[e + f*x]),x]`

output `-((B*c*Hypergeometric2F1[1/2, (1 - m)/2, (3 - m)/2, Cos[e + f*x]^2]*(c*Sec[e + f*x])^(-1 + m)*Sin[e + f*x])/(b*f*(1 - m)*Sqrt[Sin[e + f*x]^2])) + ((A*b - a*B)*Cos[e + f*x]^(1 + m)*(c*Sec[e + f*x])^(1 + m)*(-((b*AppellF1[1/2, m/2, 1, 3/2, Sin[e + f*x]^2, -((b^2*Sin[e + f*x]^2)/(a^2 - b^2))]*(Cos[e + f*x]^2)^(m/2)*Sin[e + f*x])/((a^2 - b^2)*f*Cos[e + f*x]^m)) + (a*AppellF1[1/2, (1 + m)/2, 1, 3/2, Sin[e + f*x]^2, -((b^2*Sin[e + f*x]^2)/(a^2 - b^2))]*Cos[e + f*x]^(-1 - m)*(Cos[e + f*x]^2)^((1 + m)/2)*Sin[e + f*x])/((a^2 - b^2)*f)))/(b*c)`

3.640.3.1 Defintions of rubi rules used

- rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp
p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; F
reeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,
0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 3122 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]`
- rule 3302 `Int[((d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)/((a_) + (b_.)*sin[(e_.) + (f_.)*(
x_)]), x_Symbol] := Simp[a Int[(d*Sin[e + f*x])^n/(a^2 - b^2*Sin[e + f*x]
^2), x], x] - Simp[b/d Int[(d*Sin[e + f*x])^(n + 1)/(a^2 - b^2*Sin[e + f*
x]^2), x], x] /; FreeQ[{a, b, d, e, f, n}, x] && NeQ[a^2 - b^2, 0]`
- rule 3439 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[g^(m + n) Int[(g*Csc[e + f*x])^(p - m - n)*(b + a*Csc[e + f*x])^m*(d +
c*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b*c
- a*d, 0] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 3668 `Int[((d_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x
)])^(p.), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Simp[(
-ff)*d^(2*IntPart[(m - 1)/2] + 1)*((d*Sin[e + f*x])^(2*FracPart[(m - 1)/2])
/(f*(Sin[e + f*x]^2)^FracPart[(m - 1)/2])) Subst[Int[(1 - ff^2*x^2)^((m -
1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b,
d, e, f, m, p}, x] && !IntegerQ[m]`
- rule 4259 `Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1) Int[1/(Sin[c + d*x]/b)^n, x]), x] /;
FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4356 `Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] :> Simp[Sin[e + f*x]^n*(d*Csc[e + f*x])^n Int[(b + a*Sin[e + f*x])^m/Sin[e + f*x]^(m + n), x], x] /; FreeQ[{a, b, d, e, f, n}, x] && NeQ[a^2 - b^2, 0] && IntegerQ[m]`

rule 4526 `Int[((csc[(e_.) + (f_.)*(x_.)]*(d_.))^(n_.)*(csc[(e_.) + (f_.)*(x_.)]*(B_.) + (A_.)))/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[A/a Int[(d*Csc[e + f*x])^n, x], x] - Simp[(A*b - a*B)/(a*d) Int[(d*Csc[e + f*x])^(n + 1)/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f, A, B, n}, x] && NeQ[A*b - a*B, 0] && NeQ[a^2 - b^2, 0]`

3.640.4 Maple [F]

$$\int \frac{(A + \cos(fx + e)B)(c \sec(fx + e))^m}{a + b \cos(fx + e)} dx$$

input `int((A+cos(f*x+e)*B)*(c*sec(f*x+e))^m/(a+b*cos(f*x+e)),x)`

output `int((A+cos(f*x+e)*B)*(c*sec(f*x+e))^m/(a+b*cos(f*x+e)),x)`

3.640.5 Fracas [F]

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{b \cos(fx + e) + a} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e)),x, algorithm="fricas")`

output `integral((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/(b*cos(f*x + e) + a), x)`

3.640.6 Sympy [F]

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx = \int \frac{(c \sec(e + fx))^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))**m/(a+b*cos(f*x+e)),x)`

output `Integral((c*sec(e + f*x))**m*(A + B*cos(e + f*x))/(a + b*cos(e + f*x)), x)`

3.640.7 Maxima [F]

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{b \cos(fx + e) + a} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e)),x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/(b*cos(f*x + e) + a), x)`

3.640.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{-1, [0, 1, 0, 0]%%} / %%{1, [0, 0, 1, 0]%%}+%%{-1, [0, 0, 0, 1]%%} Error:`

3.640.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{a + b \cos(e + fx)} dx = \int \frac{\left(\frac{c}{\cos(e + fx)}\right)^m (A + B \cos(e + fx))}{a + b \cos(e + fx)} dx$$

input `int(((c/cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x)),x)`output `int(((c/cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x)), x)`

3.641 $\int (a+b \cos(e+fx))^{3/2}(A+B \cos(e+fx))(c \sec(e+fx))^m dx$

3.641.1 Optimal result	5871
3.641.2 Mathematica [F(-1)]	5872
3.641.3 Rubi [N/A]	5872
3.641.4 Maple [N/A] (verified)	5874
3.641.5 Fricas [N/A]	5875
3.641.6 Sympy [F(-1)]	5875
3.641.7 Maxima [N/A]	5875
3.641.8 Giac [N/A]	5876
3.641.9 Mupad [N/A]	5876

3.641.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int (a + b \cos(e + fx))^{3/2}(A + B \cos(e + fx))(c \sec(e + fx))^m dx = \frac{2bB \cos(e + fx) \sqrt{a + b \cos(e + fx)}(c \sec(e + fx))^m \sin(e + fx)}{f(5 - 2m)} + \frac{2(c \cos(e + fx))^m (c \sec(e + fx))^m \text{Int}\left(\frac{(c \cos(e + fx))^{-m} (\frac{1}{2}ac(2bB(1-m) + 2aA(\frac{5}{2}-m)) + \frac{1}{2}c(b^2B(3-2m) + a(2Ab+aB)(5-2m))}{\sqrt{a+b \cos(e+fx)}})\right)}{c(5 - 2m)}$$

output

```
2*b*B*cos(f*x+e)*(c*sec(f*x+e))^m*sin(f*x+e)*(a+b*cos(f*x+e))^(1/2)/f/(5-2
*m)+2*(c*cos(f*x+e))^m*(c*sec(f*x+e))^m*Unintegrable((1/2*a*c*(2*b*B*(1-m)
+2*a*A*(5/2-m))+1/2*c*(b^2*B*(3-2*m)+a*(2*A*b+B*a)*(5-2*m))*cos(f*x+e)+1/2
*b*c*(A*b*(5-2*m)+2*a*B*(3-m))*cos(f*x+e)^2)/((c*cos(f*x+e))^m)/(a+b*cos(f
*x+e))^(1/2),x)/c/(5-2*m)
```

3.641.2 Mathematica [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \$Aborted$$

input `Integrate[(a + b*Cos[e + f*x])^(3/2)*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])
^m,x]`

output `$Aborted`

3.641.3 Rubi [N/A]

Not integrable

Time = 1.06 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3440, 3042, 3469, 27, 3042, 3544}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^{3/2} \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) \left(c \csc \left(e + fx + \frac{\pi}{2} \right) \right)^m dx \\ & \quad \downarrow \text{3440} \\ & (c \cos(e + fx))^m (c \sec(e + fx))^m \int (c \cos(e + fx))^{-m} (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & (c \cos(e + fx))^m (c \sec(e + fx))^m \int \left(c \sin \left(e + fx + \frac{\pi}{2} \right) \right)^{-m} \left(a + b \sin \left(e + fx + \frac{\pi}{2} \right) \right)^{3/2} \left(A + B \sin \left(e + fx + \frac{\pi}{2} \right) \right) dx \\ & \quad \downarrow \text{3469} \end{aligned}$$

$$\begin{aligned}
 & (c \cos(e + fx))^m (c \sec(e + fx))^m \left(\frac{2 \int \frac{(c \cos(e+fx))^{-m} (bc(Ab(5-2m)+2aB(3-m)) \cos^2(e+fx) + c(B(3-2m)b^2 + a(2Ab+aB)(5-2m)) \cos(e+fx) + ac(aA(5-2m)+2bB(3-m)))}{2\sqrt{a+b \cos(e+fx)}}}{c(5-2m)} \right) \\
 & \quad \downarrow \text{27} \\
 & (c \cos(e + fx))^m (c \sec(e + fx))^m \left(\frac{\int \frac{(c \cos(e+fx))^{-m} (bc(Ab(5-2m)+2aB(3-m)) \cos^2(e+fx) + c(B(3-2m)b^2 + a(2Ab+aB)(5-2m)) \cos(e+fx) + ac(aA(5-2m)+2bB(3-m)))}{\sqrt{a+b \cos(e+fx)}}}{c(5-2m)} \right) \\
 & \quad \downarrow \text{3042} \\
 & (c \cos(e + fx))^m (c \sec(e + fx))^m \left(\frac{\int \frac{(c \sin(e+fx+\frac{\pi}{2}))^{-m} (bc(Ab(5-2m)+2aB(3-m)) \sin^2(e+fx+\frac{\pi}{2}) + c(B(3-2m)b^2 + a(2Ab+aB)(5-2m)) \sin(e+fx+\frac{\pi}{2}) + ac(aA(5-2m)+2bB(3-m)))}{\sqrt{a+b \sin(e+fx+\frac{\pi}{2})}}}{c(5-2m)} \right) \\
 & \quad \downarrow \text{3544} \\
 & (c \cos(e + fx))^m (c \sec(e + fx))^m \left(\frac{\int \frac{(c \cos(e+fx))^{-m} (bc(Ab(5-2m)+2aB(3-m)) \cos^2(e+fx) + c(B(3-2m)b^2 + a(2Ab+aB)(5-2m)) \cos(e+fx) + ac(aA(5-2m)+2bB(3-m)))}{\sqrt{a+b \cos(e+fx)}}}{c(5-2m)} \right)
 \end{aligned}$$

input `Int[(a + b*Cos[e + f*x])^(3/2)*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `$Aborted`

3.641.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 3440 Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*
(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Sim
p[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c +
d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g
, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && I
ntegerQ[n])
```

```
rule 3469 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-b)*B*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^(
n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Sin[e +
f*x])^(m - 2)*(c + d*Sin[e + f*x])^n*Simp[a^2*A*d*(m + n + 1) + b*B*(b*c*(
m - 1) + a*d*(n + 1)) + (a*d*(2*A*b + a*B)*(m + n + 1) - b*B*(a*c - b*d*(m
+ n)))*Sin[e + f*x] + b*(A*b*d*(m + n + 1) - B*(b*c*m - a*d*(2*m + n)))*Sin
[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && GtQ[m, 1] && !(IGt
Q[n, 1] && (!IntegerQ[m] || (EqQ[a, 0] && NeQ[c, 0])))
```

```
rule 3544 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(c + d*
Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x] /; FreeQ[{a, b,
c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0]
```

3.641.4 Maple [N/A] (verified)

Not integrable

Time = 2.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int (a + b \cos(fx + e))^{\frac{3}{2}} (A + \cos(fx + e) B) (c \sec(fx + e))^m dx$$

```
input int((a+b*cos(f*x+e))^(3/2)*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)
```

```
output int((a+b*cos(f*x+e))^(3/2)*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)
```

3.641.5 Fricas [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^{3/2} (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^(3/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algo rithm="fricas")`

output `integral((B*b*cos(f*x + e)^2 + A*a + (B*a + A*b)*cos(f*x + e))*sqrt(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m, x)`

3.641.6 Sympy [F(-1)]

Timed out.

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \text{Timed out}$$

input `integrate((a+b*cos(f*x+e))**(3/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))**m,x)`

output `Timed out`

3.641.7 Maxima [N/A]

Not integrable

Time = 2.69 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^{3/2} (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^(3/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algo
rithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^(3/2)*(c*sec(f*x + e))
^m, x)`

3.641.8 Giac [N/A]

Not integrable

Time = 1.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \int (B \cos(fx + e) + A) (b \cos(fx + e) + a)^{3/2} (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^(3/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algo
rithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(b*cos(f*x + e) + a)^(3/2)*(c*sec(f*x + e))
^m, x)`

3.641.9 Mupad [N/A]

Not integrable

Time = 5.76 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int (a + b \cos(e + fx))^{3/2} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx = \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) (a + b \cos(e + fx))^{3/2} dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(3/2),x)`

output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(3/2), x)`

3.642 $\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$

3.642.1 Optimal result	5877
3.642.2 Mathematica [N/A]	5877
3.642.3 Rubi [N/A]	5878
3.642.4 Maple [N/A] (verified)	5879
3.642.5 Fricas [N/A]	5880
3.642.6 Sympy [N/A]	5880
3.642.7 Maxima [N/A]	5881
3.642.8 Giac [N/A]	5881
3.642.9 Mupad [N/A]	5882

3.642.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= (c \cos(e + fx))^m (c \sec(e + fx))^m \text{Int}\left((c \cos(e + fx))^{-m} \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx)), x\right)$$

output $(c*\cos(f*x+e))^m*(c*\sec(f*x+e))^m*\text{Unintegrable}((A+B*\cos(f*x+e))*(a+b*\cos(f*x+e))^{1/2}/((c*\cos(f*x+e))^m),x)$

3.642.2 Mathematica [N/A]

Not integrable

Time = 33.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

input `Integrate[Sqrt[a + b*Cos[e + f*x]]*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m ,x]`

output `Integrate[Sqrt[a + b*Cos[e + f*x]]*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m , x]`

3.642.3 Rubi [N/A]

Not integrable

Time = 0.48 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3440, 3042, 3486}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{a + b \sin\left(e + fx + \frac{\pi}{2}\right)} \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right)\right) \left(c \csc\left(e + fx + \frac{\pi}{2}\right)\right)^m dx \\ & \quad \downarrow \text{3440} \\ & (c \cos(e + fx))^m (c \sec(e + fx))^m \int (c \cos(e + fx))^{-m} \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & (c \cos(e + fx))^m (c \sec(e + fx))^m \int \left(c \sin\left(e + fx + \frac{\pi}{2}\right)\right)^{-m} \sqrt{a + b \sin\left(e + fx + \frac{\pi}{2}\right)} \left(A + B \sin\left(e + fx + \frac{\pi}{2}\right)\right) dx \\ & \quad \downarrow \text{3486} \\ & (c \cos(e + fx))^m (c \sec(e + fx))^m \int (c \cos(e + fx))^{-m} \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) dx \end{aligned}$$

input `Int[Sqrt[a + b*Cos[e + f*x]]*(A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m,x]`

output `$Aborted`

3.642.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*SIN[e + f*x])^p Int[(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^n/(g*SIN[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3486 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(a + b*SIN[e + f*x])^m*(A + B*SIN[e + f*x])*(c + d*SIN[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.642.4 Maple [N/A] (verified)

Not integrable

Time = 1.77 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \sqrt{a + b \cos(fx + e)} (A + \cos(fx + e) B) (c \sec(fx + e))^m dx$$

input `int((a+b*cos(f*x+e))^(1/2)*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

output `int((a+b*cos(f*x+e))^(1/2)*(A+cos(f*x+e)*B)*(c*sec(f*x+e))^m,x)`

3.642.5 Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) \sqrt{b \cos(fx + e) + a} (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^(1/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algo rithm="fricas")`

output `integral((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m, x)`

3.642.6 Sympy [N/A]

Not integrable

Time = 8.46 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (c \sec(e + fx))^m (A + B \cos(e + fx)) \sqrt{a + b \cos(e + fx)} dx$$

input `integrate((a+b*cos(f*x+e))**(1/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))**m,x)`

output `Integral((c*sec(e + f*x))**m*(A + B*cos(e + f*x))*sqrt(a + b*cos(e + f*x)), x)`

3.642.7 Maxima [N/A]

Not integrable

Time = 2.71 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) \sqrt{b \cos(fx + e) + a} (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^(1/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m, x)`

3.642.8 Giac [N/A]

Not integrable

Time = 0.69 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos(e + fx)}(A + B \cos(e + fx))(c \sec(e + fx))^m dx$$

$$= \int (B \cos(fx + e) + A) \sqrt{b \cos(fx + e) + a} (c \sec(fx + e))^m dx$$

input `integrate((a+b*cos(f*x+e))^(1/2)*(A+B*cos(f*x+e))*(c*sec(f*x+e))^m,x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m, x)`

3.642.9 Mupad [N/A]

Not integrable

Time = 3.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \sqrt{a + b \cos(e + fx)} (A + B \cos(e + fx)) (c \sec(e + fx))^m dx$$

$$= \int \left(\frac{c}{\cos(e + fx)} \right)^m (A + B \cos(e + fx)) \sqrt{a + b \cos(e + fx)} dx$$

input `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(1/2),x)`output `int((c/cos(e + f*x))^m*(A + B*cos(e + f*x))*(a + b*cos(e + f*x))^(1/2), x)`

$$3.643 \quad \int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{\sqrt{a+b \cos(e+fx)}} dx$$

3.643.1 Optimal result	5883
3.643.2 Mathematica [N/A]	5883
3.643.3 Rubi [N/A]	5884
3.643.4 Maple [N/A] (verified)	5885
3.643.5 Fricas [N/A]	5886
3.643.6 Sympy [N/A]	5886
3.643.7 Maxima [N/A]	5886
3.643.8 Giac [N/A]	5887
3.643.9 Mupad [N/A]	5887

3.643.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{\sqrt{a+b \cos(e+fx)}} dx$$

$$= (c \cos(e+fx))^m (c \sec(e+fx))^m \text{Int} \left(\frac{(c \cos(e+fx))^{-m} (A+B \cos(e+fx))}{\sqrt{a+b \cos(e+fx)}}, x \right)$$

output `(c*cos(f*x+e))^m*(c*sec(f*x+e))^m*Unintegrable((A+B*cos(f*x+e))/((c*cos(f*x+e))^m)/(a+b*cos(f*x+e))^(1/2),x)`

3.643.2 Mathematica [N/A]

Not integrable

Time = 39.77 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{\sqrt{a+b \cos(e+fx)}} dx = \int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{\sqrt{a+b \cos(e+fx)}} dx$$

input `Integrate[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/Sqrt[a + b*Cos[e + f*x]],x]`

output `Integrate[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/Sqrt[a + b*Cos[e + f*x]], x]`

$$3.643. \quad \int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{\sqrt{a+b \cos(e+fx)}} dx$$

3.643.3 Rubi [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3440, 3042, 3486}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx$$

↓ 3042

$$\int \frac{(A + B \sin(e + fx + \frac{\pi}{2}))(c \csc(e + fx + \frac{\pi}{2}))^m}{\sqrt{a + b \sin(e + fx + \frac{\pi}{2})}} dx$$

↓ 3440

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int \frac{(c \cos(e + fx))^{-m} (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$$

↓ 3042

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int \frac{(c \sin(e + fx + \frac{\pi}{2}))^{-m} (A + B \sin(e + fx + \frac{\pi}{2}))}{\sqrt{a + b \sin(e + fx + \frac{\pi}{2})}} dx$$

↓ 3486

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int \frac{(c \cos(e + fx))^{-m} (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$$

input `Int[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/Sqrt[a + b*Cos[e + f*x]],x]`

output `$Aborted`

3.643.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3440 `Int[(csc[(e_.) + (f_.)*(x_)])*(g_.)^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

rule 3486 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(A + B*Sin[e + f*x])*(c + d*Sin[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]`

3.643.4 Maple [N/A] (verified)

Not integrable

Time = 1.58 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(A + \cos(fx + e)B)(c \sec(fx + e))^m}{\sqrt{a + b \cos(fx + e)}} dx$$

input `int((A+cos(f*x+e)*B)*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(1/2),x)`

output `int((A+cos(f*x+e)*B)*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(1/2),x)`

3.643.5 Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{\sqrt{b \cos(fx + e) + a}} dx$$

```
input integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(1/2),x, algo
rithm="fricas")
```

```
output integral((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/sqrt(b*cos(f*x + e) + a),
x)
```

3.643.6 Sympy [N/A]

Not integrable

Time = 3.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(c \sec(e + fx))^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$$

```
input integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(1/2),x)
```

```
output Integral((c*sec(e + f*x))^m*(A + B*cos(e + f*x))/sqrt(a + b*cos(e + f*x))
, x)
```

3.643.7 Maxima [N/A]

Not integrable

Time = 2.61 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{\sqrt{b \cos(fx + e) + a}} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(1/2),x, algorith="maxima")`

output `integrate((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/sqrt(b*cos(f*x + e) + a), x)`

3.643.8 Giac [N/A]

Not integrable

Time = 0.73 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{\sqrt{b \cos(fx + e) + a}} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(1/2),x, algorith="giac")`

output `integrate((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/sqrt(b*cos(f*x + e) + a), x)`

3.643.9 Mupad [N/A]

Not integrable

Time = 4.94 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{\sqrt{a + b \cos(e + fx)}} dx = \int \frac{\left(\frac{c}{\cos(e + fx)}\right)^m (A + B \cos(e + fx))}{\sqrt{a + b \cos(e + fx)}} dx$$

input `int(((c/cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(1/2),x)`

output `int(((c/cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(1/2),x)`

3.644 $\int \frac{(A+B \cos(e+fx))(c \sec(e+fx))^m}{(a+b \cos(e+fx))^{3/2}} dx$

3.644.1 Optimal result	5888
3.644.2 Mathematica [N/A]	5888
3.644.3 Rubi [N/A]	5889
3.644.4 Maple [N/A] (verified)	5891
3.644.5 Fricas [N/A]	5892
3.644.6 Sympy [N/A]	5892
3.644.7 Maxima [N/A]	5892
3.644.8 Giac [N/A]	5893
3.644.9 Mupad [N/A]	5893

3.644.1 Optimal result

Integrand size = 35, antiderivative size = 35

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \frac{2b(Ab - aB) \cos(e + fx)(c \sec(e + fx))^m \sin(e + fx)}{a(a^2 - b^2) f \sqrt{a + b \cos(e + fx)}} + \frac{2(c \cos(e + fx))^m (c \sec(e + fx))^m \operatorname{Int}\left(\frac{(c \cos(e + fx))^{-m} (\frac{1}{2}c(a^2 A + Ab^2(1-2m) - 2abB(1-m)) - \frac{1}{2}a(Ab - aB)c \cos(e + fx) - \frac{1}{2}b(Ab - aB)c \cos(e + fx)^2)}{\sqrt{a + b \cos(e + fx)}}\right)}{a(a^2 - b^2) c}$$

output

```
2*b*(A*b-B*a)*cos(f*x+e)*(c*sec(f*x+e))^m*sin(f*x+e)/a/(a^2-b^2)/f/(a+b*cos(f*x+e))^(1/2)+2*(c*cos(f*x+e))^m*(c*sec(f*x+e))^m*Unintegrable((1/2*c*(A*a^2+A*b^2*(1-2*m))-2*a*b*B*(1-m))-1/2*a*(A*b-B*a)*c*cos(f*x+e)-1/2*b*(A*b-B*a)*c*(3-2*m)*cos(f*x+e)^2)/((c*cos(f*x+e))^m)/(a+b*cos(f*x+e))^(1/2),x)/a/(a^2-b^2)/c
```

3.644.2 Mathematica [N/A]

Not integrable

Time = 31.59 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx$$

input

```
Integrate[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/(a + b*Cos[e + f*x])^(3/2),x]
```

output `Integrate[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/(a + b*Cos[e + f*x])^(3/2), x]`

3.644.3 Rubi [N/A]

Not integrable

Time = 1.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3440, 3042, 3479, 27, 3042, 3544}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(A + B \sin(e + fx + \frac{\pi}{2}))(c \csc(e + fx + \frac{\pi}{2}))^m}{(a + b \sin(e + fx + \frac{\pi}{2}))^{3/2}} dx$$

↓ 3440

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int \frac{(c \cos(e + fx))^{-m} (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx$$

↓ 3042

$$(c \cos(e + fx))^m (c \sec(e + fx))^m \int \frac{(c \sin(e + fx + \frac{\pi}{2}))^{-m} (A + B \sin(e + fx + \frac{\pi}{2}))}{(a + b \sin(e + fx + \frac{\pi}{2}))^{3/2}} dx$$

↓ 3479

$$fx))^m \left(\frac{2 \int \frac{(c \cos(e + fx))^{-m} (-b(Ab - aB)c(3 - 2m) \cos^2(e + fx) - a(Ab - aB)c \cos(e + fx) + c(Aa^2 - 2bB(1 - m)a + Ab^2(1 - 2m)))}{2\sqrt{a + b \cos(e + fx)}} dx}{ac(a^2 - b^2)} + \frac{2b(Ab - aB)}{ac(a^2 - b^2)} \right)$$

↓ 27

$$fx))^m \left(\frac{\int \frac{(c \cos(e + fx))^{-m} (-b(Ab - aB)c(3 - 2m) \cos^2(e + fx) - a(Ab - aB)c \cos(e + fx) + c(Aa^2 - 2bB(1 - m)a + Ab^2(1 - 2m)))}{\sqrt{a + b \cos(e + fx)}} dx}{ac(a^2 - b^2)} + \frac{2b(Ab - aB)}{ac(a^2 - b^2)} \right)$$

↓ 3042

3.644. $\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx$

$$\begin{aligned}
 & \int \frac{(c \cos(e + fx))^m (c \sec(e + fx))^m (c \sin(e + fx + \frac{\pi}{2}))^{-m} (-b(Ab - aB)c(3 - 2m) \sin(e + fx + \frac{\pi}{2})^2 - a(Ab - aB)c \sin(e + fx + \frac{\pi}{2}) + c(Aa^2 - 2bB(1 - m)a + Ab^2(1 - 2m)))}{\sqrt{a + b \sin(e + fx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3544} \\
 & \int \frac{(c \cos(e + fx))^m (c \sec(e + fx))^m (c \cos(e + fx))^{-m} (-b(Ab - aB)c(3 - 2m) \cos^2(e + fx) - a(Ab - aB)c \cos(e + fx) + c(Aa^2 - 2bB(1 - m)a + Ab^2(1 - 2m)))}{\sqrt{a + b \cos(e + fx)}} dx + \frac{2b(Ab - aB)}{ac(a^2 - b^2)}
 \end{aligned}$$

input `Int[((A + B*Cos[e + f*x])*(c*Sec[e + f*x])^m)/(a + b*Cos[e + f*x])^(3/2),x]`

output `$Aborted`

3.644.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3440 `Int[(csc[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(g*Csc[e + f*x])^p*(g*Sin[e + f*x])^p Int[(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(g*Sin[e + f*x])^p), x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[p] && !(IntegerQ[m] && IntegerQ[n])`

```
rule 3479 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(-(A*b^2 - a*b*B))*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*((c + d*Sin
[e + f*x])^(1 + n)/(f*(m + 1)*(b*c - a*d)*(a^2 - b^2))), x] + Simp[1/((m +
1)*(b*c - a*d)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e
+ f*x])^n*Simp[(a*A - b*B)*(b*c - a*d)*(m + 1) + b*d*(A*b - a*B)*(m + n +
2) + (A*b - a*B)*(a*d*(m + 1) - b*c*(m + 2))*Sin[e + f*x] - b*d*(A*b - a*B)
*(m + n + 3)*Sin[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n
}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && Rat
ionalQ[m] && m < -1 && ((EqQ[a, 0] && IntegerQ[m] && !IntegerQ[n]) || !(I
ntegerQ[2*n] && LtQ[n, -1] && ((IntegerQ[n] && !IntegerQ[m]) || EqQ[a, 0]
))
```

```
rule 3544 Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(c + d*
Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x] /; FreeQ[{a, b,
c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] &
& NeQ[c^2 - d^2, 0]
```

3.644.4 Maple [N/A] (verified)

Not integrable

Time = 1.71 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(A + \cos(fx + e)B)(c \sec(fx + e))^m}{(a + b \cos(fx + e))^{\frac{3}{2}}} dx$$

```
input int((A+cos(f*x+e)*B)*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(3/2),x)
```

```
output int((A+cos(f*x+e)*B)*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(3/2),x)
```

3.644.5 Fricas [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.80

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{(b \cos(fx + e) + a)^{3/2}} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(3/2),x, algo
rithm="fricas")`

output `integral((B*cos(f*x + e) + A)*sqrt(b*cos(f*x + e) + a)*(c*sec(f*x + e))^m/
(b^2*cos(f*x + e)^2 + 2*a*b*cos(f*x + e) + a^2), x)`

3.644.6 Sympy [N/A]

Not integrable

Time = 11.40 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(c \sec(e + fx))^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(3/2),x)`

output `Integral((c*sec(e + f*x))^m*(A + B*cos(e + f*x))/(a + b*cos(e + f*x))^(3
/2), x)`

3.644.7 Maxima [N/A]

Not integrable

Time = 2.32 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{(b \cos(fx + e) + a)^{3/2}} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/(b*cos(f*x + e) + a)^(3/2), x)`

3.644.8 Giac [N/A]

Not integrable

Time = 1.26 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{(B \cos(fx + e) + A)(c \sec(fx + e))^m}{(b \cos(fx + e) + a)^{3/2}} dx$$

input `integrate((A+B*cos(f*x+e))*(c*sec(f*x+e))^m/(a+b*cos(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*cos(f*x + e) + A)*(c*sec(f*x + e))^m/(b*cos(f*x + e) + a)^(3/2), x)`

3.644.9 Mupad [N/A]

Not integrable

Time = 8.63 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{(A + B \cos(e + fx))(c \sec(e + fx))^m}{(a + b \cos(e + fx))^{3/2}} dx = \int \frac{\left(\frac{c}{\cos(e + fx)}\right)^m (A + B \cos(e + fx))}{(a + b \cos(e + fx))^{3/2}} dx$$

input `int(((c/cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(3/2),x)`

output `int(((c/cos(e + f*x))^m*(A + B*cos(e + f*x)))/(a + b*cos(e + f*x))^(3/2), x)`

APPENDIX

4.1 Listing of Grading functions	5894
--	------

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3,ExpnType[expn[[1]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
          If[HypergeometricFunctionQ[Head[expn]],
            Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
          If[AppellFunctionQ[Head[expn]],
            Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
          If[Head[expn]===RootSum,
            Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
          If[Head[expn]===Integrate || Head[expn]===Int,
            Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
          9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```



```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```

if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):

```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```



```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```